


TRUMPETER SWAN SURVEYS ON THE
CHUGACH NATIONAL FOREST
2007



Deborah J. Groves¹, Karen Bollinger², and Rob MacDonald¹

¹U.S. Fish and Wildlife Service, Migratory Bird Management, Waterfowl Management Branch, 3000 Vintage Blvd., Suite 240, Juneau, Alaska 99801

²U.S. Fish and Wildlife Service, Migratory Bird Management, Waterfowl Management Branch, 1412 Airport Way, Fairbanks, Alaska 99701

January 2008

ABSTRACT

Trumpeter swan (*Cygnus buccinator*) aerial surveys were conducted during May and August 2007 on the Copper River Delta and surrounding areas of the Chugach National Forest in southcentral Alaska. The surveys were accomplished through cooperation between the U.S. Fish and Wildlife Service and the U.S. Forest Service. In May 906 white (adult and subadult) swans and 137 nests were counted. In August 776 white swans and 125 cygnets in 43 broods were recorded. Production was below average. Nest success was 0.31, mean brood size was 2.9, and young made up 14% of the early fall population. The total fall swan population decreased 22% from 2006 and was equal to the previous 31-year mean. There are now 61 comparable swan surveys in 32 different years for this area. A continued standardized trumpeter swan survey program is recommended.

INTRODUCTION

The Copper River Delta and surrounding coastal wetlands in the Chugach National Forest support a large and dense nesting and summering population of trumpeter swans (*Cygnus buccinator*). Aerial surveys were conducted in this area as part of U.S. Fish and Wildlife Service (USFWS) statewide trumpeter censuses in 1968, 1975, 1980, 1985, 1990, 1995, 2000, and 2005 (Hansen et al. 1971, King 1976, King and Conant 1981, Conant et al. 1985, Conant et al. 1991, Conant et al. 1996, Conant et al. 2001, USFWS unpubl. data). The U.S. Forest Service (USFS) initiated aerial surveys in 1978, and annual swan surveys have been jointly accomplished under a formal agreement between the USFWS and USFS since 1981.

SURVEY AREA

The survey area was comprised of portions of the Copper River Delta and Controller Bay drainages within 11 U.S. Geological Survey 1:63,360 scale topographic maps (Figure 1). The area actually surveyed consisted of all potential swan nesting and summering habitat delineated on these maps (1787 km²). In general, potential swan habitat included most lakes, rivers, streams and all wetland areas under 760 m (2500 ft) elevation.

METHODS

Aerial surveys were conducted 19-20 May 2007 with a Cessna 206 and 20-24 August 2007 with the USFWS Turbo-Beaver (N754). The aerial survey method used for both surveys was described by King (1973), with modifications that allowed swan observations and the survey flight path to be recorded digitally. Generally, a system of parallel tracks was flown 150-180 m above ground level over all known and potential swan habitat within the survey area. The pilot was responsible for navigation, searching for swans, and ensuring that all swan habitat was adequately surveyed, considering factors such as visibility and observer experience. The primary observer searched for swans and recorded the type and location of each observation onto a digital 1:63,360 topographic map, using a custom program run on a notebook computer linked to the

aircraft's GPS unit. The program produced two text files: A swan data file containing the geographic coordinates and attributes of each swan observation, and a flight path file containing point locations of the aircraft during the survey flight in five-second intervals. The program, developed by John Hodges, USFWS, Migratory Bird Management, Juneau, Alaska, has been used for the spring survey since 2002 and for the fall survey since 2000. Prior to that, data were recorded on paper maps as per King 1973 and afterwards transcribed into a computer and digitized to produce the final data files.

RESULTS

Population Trend

In May 2007, 906 white swans (adults and subadults) were counted, down 9% from spring 2006 and 35% above the previous 28-year mean (Table 1). The numbers of single, paired, and flocked birds all decreased from last year, but the number of single/paired birds was still the second-highest count on record for the May survey.

In August 2007, 776 white swans were counted, down 17% from August 2006 and 10% above the previous 31-year mean (Figures 1 and 2, Table 2). The numbers of single, paired, and flocked birds all decreased from last year. The total number of swans, including cygnets, was 901 in 2007, down 22% from 2006 and equal to the previous mean (Figure 2, Table 2).

Productivity

During the May survey 137 nests were observed, 12% above the previous 28-year mean (Figures 3 and 4, Table 3). Thirty-nine percent of pairs had a nest, 24% below the previous mean (Table 3). In August 125 cygnets in 43 broods were recorded, resulting in a mean brood size of 2.9 (Figures 3 and 4, Tables 2 and 3). The number of cygnets, number of broods, and mean brood size were 35%, 25%, and 13% below their previous means, respectively. Nest success, which we defined as the proportion of occupied nests in May that produced at least one cygnet still alive in August, was 0.31, 37% below the previous mean (Table 3). The number of young per occupied nest, a productivity statistic based on the number of known territorial pairs (as evidenced by the presence of a nest or brood), was 0.9, 46% below the previous mean. Young made up 14% of the early fall population, 35% below the previous mean (Tables 2 and 3).

DISCUSSION

The size of the fall swan population in the survey area has fluctuated between about 500 and 1300 birds since surveys began in 1968. The population increased in the early survey years until it peaked at 1118 swans in 1984, then decreased to 662 in 1991. It reached a second peak of 1325 swans in 2005, then declined successively during the past two years to its 2007 level of 901 swans. Annual variation in weather conditions on the breeding and wintering grounds were likely the main factors causing these fluctuations, through its influence on breeding propensity, nest success, cygnet survival, and overwinter survival of adults and juveniles. The below-

average productivity experienced in 2007 should result in reduced recruitment into the population this year.

BIAS

Possible sources of bias in these data come from using different pilots and observers with variable levels of experience and training, using more than one type of aircraft, and surveying in variable weather conditions. However, by using a standardized system, comparable sets of data were collected as evidenced by comparable recorded flight paths and mileages flown.

RECOMMENDATIONS

We now have 61 comparable trumpeter swan surveys (29 spring and 32 fall) on the Copper River Delta, one of the most complete records for any swan population in Alaska. We recommend continuing a cooperative program of two surveys per year. Information acquired from both the early and late phases of the breeding season has greatly enhanced our ability to understand the factors influencing the population's reproductive success. Long-term, standardized data sets such as these are an invaluable tool for evaluating population dynamics and properly managing trumpeter swan breeding populations.

SURVEY PERSONNEL

Swan surveys summarized in this report were conducted by the following individuals and agencies:

Year	Spring Survey Personnel	Fall Survey Personnel
1968		J. King, J. Bartonek - USFWS
1975		J. King, T. Schoenfelder - USFWS
1978	G. Bucaria, S. Watson – USFS	G. Bucaria, S. Watson – USFS
1979		C. Moitoret – USFS
1980	G. Bucaria, D. Logan – USFS	J. King, B. Conant – USFWS
1981	G. Bucaria – USFS	J. King, B. Conant – USFWS
1982	R. King, G. Bucaria – USFWS, USFS	J. King, B. Conant – USFWS
1983	R. King, Zimmerman – USFWS	B. Conant, D. Derksen, J. Baker, M. Jacobson, G. Covell, Broekema – USFWS, USFS
1984	R. King, R. Leedy – USFWS	B. Conant, J. Hodges – USFWS
1985	R. King, T. Simon-Jackson – USFWS	B. Conant, S. Cain – USFWS
1986	R. King, K. Bollinger – USFWS	B. Conant, J. Hodges – USFWS
1987	R. King, M. North – USFWS	B. Conant, J. Hodges – USFWS
1988	R. King, R. Pospahala – USFWS	B. Conant, J. Hodges – USFWS
1989	R. King, F. Gerhardt – USFWS	B. Conant, R. Oates, M. Jacobson – USFWS
1990	R. King, A. Brackney – USFWS	B. Conant, D. Groves, J. King – USFWS
1991	R. King, A. Brackney – USFWS	B. Conant, D. Groves – USFWS
1992	R. King, D. Youkey – USFWS, USFS	J. Hodges, J. King – USFWS
1993	R. King, D. Youkey – USFWS, USFS	J. Hodges, D. Groves, D. Youkey – USFWS, USFS
1994	R. King, P. Greene – USFWS, USFS	B. Conant, D. Groves – USFWS
1995	R. King, R. Leedy – USFWS	B. Conant, E. Lucas – USFWS
1996	R. King, S. Hill – USFWS	J. Hodges, D. Groves – USFWS
1997	R. King, T. Tiplady – USFWS	B. Conant, G. Fowler – USFWS, Ducks Unlimited Canada
1998	W. Larned, J. King – USFWS	B. Conant, J. King – USFWS
1999	W. Larned, T. Tiplady – USFWS	B. Conant, J. King – USFWS
2000	J. Sarvis, R. Oates – USFWS	B. Conant, D. Petersen – USFWS
2001	E. Mallek, P. Anderson – USFWS	B. Conant, J. King – USFWS
2002	E. Mallek, L. Lysne – USFWS	B. Conant, L. Lysne, D. Groves – USFWS
2003	E. Mallek, L. Lysne - USFWS	B. Conant, H. Wilson – USFWS
2004	E. Mallek, J. King - USFWS	J. Hodges, H. Wilson - USFWS
2005	E. Mallek, J. King - USFWS	B. Conant, J. King - USFWS
2006	W. Larned, J. Seyfried - USFWS, USGS	P. Anderson, E. Mallek - USFWS
2007	K. Bollinger, C. Dau - USFWS	R. MacDonald, P. Anderson - USFWS

LITERATURE CITED

- Conant, B., J. I. Hodges, J. G. King, and S. L. Cain. 1985. Alaska trumpeter swan status report - 1985. U.S. Fish and Wildlife Service Report, Juneau, Alaska. 10pp.
- _____, _____, D. J. Groves, and J. G. King. 1991. Alaska trumpeter swan status report - 1990. U.S. Fish and Wildlife Service Report, Juneau, Alaska. 31pp.
- _____, _____, _____, and _____. 1996. Alaska trumpeter swan status report - 1995. U.S. Fish and Wildlife Service Report, Juneau, Alaska.
- _____, _____, _____, and _____. 2001. Alaska trumpeter swan status report - 2000. U.S. Fish and Wildlife Service Report, Juneau, Alaska. 37pp.
- Hansen, H. A., P. E. K. Shepherd, J. G. King, and W. A. Troyer. 1971. The trumpeter swan in Alaska. Wildl. Mono. No. 26: 1-83.
- King, J. G. 1973. The use of small airplanes to gather swan data in Alaska. Wildfowl 24:15-20.
- _____. 1976. Trumpeter swan survey in Alaska. The 5th Trumpeter Swan Society Conference.
- _____ and B. Conant. 1981. The 1980 census of trumpeter swans in Alaskan nesting habitats. American Birds, vol. 35, no. 5, pp. 789-93.




Figure 1. Copper River Delta survey area showing fall 2007 trumpeter swan distribution.




Figure 2. Population trend of trumpeter swans on the Copper River Delta survey area from fall surveys, 1978-2007.




Figure 3. Copper River Delta study area showing spring 2007 trumpeter swan nest and fall 2007 brood locations.




Figure 4. Number of trumpeter swan nests recorded during spring surveys, and number of broods and cygnets recorded during fall surveys, 1978-2007. A spring survey was not conducted in 1979.

Table 1. Spring survey swan observations - Chugach National Forest (1978-2007).

Year	Dates Flown	Observations	White Swans				Cygnets	Total Swans
			Paired	Single	Flocked	Subtotal		
1978	5/15-5/18	192	278	20	362	660	--	660
1980	May ^a	222	320	45	169	534	--	534
1981	5/21-5/23	244	350	37	235	622	--	622
1982	6/2-6/4	247	356	44	200	600	--	600
1983	6/6-6/8	297	448	47	144	639	23	662
1984	5/21-5/23	324	502	43	190	735	--	735
1985	6/8-6/10	309	452	50	235	737	--	737
1986	5/22-5/24	304	508	35	123	666	--	666
1987	5/22-5/26	291	462	39	101	602	--	602
1988	5/25-5/26	263	418	42	116	576	5	581
1989	5/17-5/19	241	400	28	174	602	--	602
1990	5/23-5/25	226	374	25	121	520	--	520
1991	5/20-5/22	250	394	34	152	580	--	580
1992	5/19-5/20	249	412	25	195	632	--	632
1993	5/15-5/17	248	394	25	159	578	--	578
1994	5/24-5/25	278	436	31	204	671	--	671
1995	5/25-5/26	246	402	24	157	583	--	583
1996	5/21-5/22	267	442	27	125	594	--	594
1997	5/26-5/27	246	406	27	84	517	--	517
1998	5/26-5/29	285	462	40	110	612	--	612
1999	6/3-6/5	229	358	33	79	470	--	470
2000	5/25-5/26	256	408	28	255	691	--	691
2001	5/29-5/31	261	410	39	117	566	31	597
2002	5/29-5/31	400	536	88	391	1015	--	1015
2003	5/27-5/29	417	632	76	205	913	--	913
2004	5/26-5/28	389	606	52	282	940	--	940
2005	5/23-5/26	425	664	66	213	943	4	947
2006	5/24-5/25	484	714	100	177	991	--	991
2007	5/19-5/20	433	670	84	152	906	--	906
Mean ^b		289	448	42	181	671	--	673
% Change from:								
2006		-11%	-6%	-16%	-14%	-9%	--	-9%
Mean ^b		50%	50%	101%	-16%	35%	--	35%

^a Exact dates unknown.

^b Mean excludes 2007.

Table 2. Fall survey swan observations - Chugach National Forest (1968-2007).

Year	Dates Flown	Observations	White Swans				Cygnets	Young in Pop.(%)	Total Swans
			Paired	Single	Flocked	Subtotal			
1968	8/14-8/16	199	326	24	181	531	267	33	798
1975	8/14-8/20	196	312	24	142	478	131	22	609
1978	8/9-8/12	186	248	36	127	411	103	20	514
1979	8/18-8/23	160	234	20	217	471	143	23	614
1980	8/5-8/12	262	410	33	187	630	216	26	846
1981	8/4-8/22	234	374	16	287	677	266	28	943
1982	8/11-8/14	271	436	27	227	690	152	18	842
1983	8/4-8/15	314	512	32	219	763	259	25	1022
1984	8/6-8/10	303	448	42	400	890	228	20	1118
1985	8/11-8/16	348	534	45	319	898	111	11	1009
1986	8/6-8/8	298	490	25	200	715	188	22	903
1987	8/1-8/3	318	510	46	175	731	64	8	795
1988	8/3-8/5	281	472	29	145	646	217	25	863
1989	8/3-8/6	278	460	41	98	599	117	16	716
1990	8/5-8/12	267	424	35	169	628	245	28	873
1991	8/2-8/4	253	400	36	90	526	136	21	662
1992	8/29-9/1	197	314	19	231	564	250	31	814
1993	8/22-8/24	237	368	29	218	615	201	25	816
1994	8/24-8/28	260	382	24	404	810	131	14	941
1995	8/2-8/6	280	408	51	185	644	97	13	741
1996	8/25-8/28	259	430	23	176	629	151	19	780
1997	8/14-8/17	259	416	25	157	598	175	23	773
1998	8/2-8/4	273	428	37	185	650	217	25	867
1999	8/10-8/14	258	408	27	199	634	182	22	816
2000	8/2-8/7	325	512	41	202	755	209	22	964
2001	8/22-8/24	314	520	26	324	870	352	29	1222
2002	8/4-8/6	313	472	50	376	898	228	20	1126
2003	8/6-8/8	349	518	50	360	928	259	22	1187
2004	8/23-8/25	284	446	27	497	970	189	16	1159
2005	8/2-8/5	361	552	56	423	1031	294	22	1325
2006	8/5-8/8	356	544	59	337	940	216	19	1156
2007	8/20-8/24	301	486	30	260	776	125	14	901
Mean ^a		274	429	34	241	704	193	22	897
% Change from:									
2006		-15%	-11%	-49%	-23%	-17%	-42%	-26%	-22%
Mean ^a		10%	13%	-12%	8%	10%	-35%	-35%	0%

^aMean excludes 2007.

Table 3. Swan productivity - Chugach National Forest (1968-2007).

Year	% Pairs with Nest in Spring	% Pairs with Brood in Fall	Nests	Broods	Nest Success ^a	Mean Brood Size	Young Per Occupied Nest ^b
1968	-- ^c	40	-- ^c	67	-- ^c	4.0	-- ^c
1975	-- ^c	24	-- ^c	39	-- ^c	3.4	-- ^c
1978	51	26	78	32	0.41	3.2	1.3
1979	-- ^c	34	-- ^c	41	-- ^c	3.5	-- ^c
1980	59	30	94	62	0.66	3.5	2.3
1981	58	34	120	67	0.56	4.0	2.2
1982	40	23	83	51	0.61	3.0	1.8
1983	27	27	68	71	1.04	3.6	3.8
1984	53	27	143	61	0.43	3.7	1.6
1985	42	13	103	37	0.36	3.0	1.1
1986	52	24	140	60	0.43	3.1	1.3
1987	43	10	115	25	0.22	2.6	0.6
1988	59	29	133	68	0.51	3.2	1.6
1989	63	17	130	38	0.29	3.1	0.9
1990	67	33	130	70	0.54	3.5	1.9
1991	64	25	129	49	0.38	2.8	1.1
1992	65	46	134	73	0.54	3.4	1.9
1993	59	32	118	61	0.52	3.3	1.7
1994	57	23	130	44	0.34	3.0	1.0
1995	56	17	115	35	0.30	2.8	0.8
1996	48	19	106	40	0.38	3.8	1.4
1997	55	25	117	54	0.46	3.2	1.5
1998	57	30	137	66	0.48	3.3	1.6
1999	54	26	105	55	0.52	3.3	1.7
2000	52	25	115	66	0.57	3.2	1.8
2001	39	35	90	93	1.03	3.8	3.9
2002	43	29	129	69	0.53	3.3	1.8
2003	41	26	144	70	0.49	3.7	1.8
2004	47	26	161	59	0.37	3.2	1.2
2005	48	33	179	91	0.51	3.2	1.6
2006	45	23	189	68	0.36	3.2	1.1
2007	39	16	137	43	0.31	2.9	0.9
Mean ^d	52	27	123	57	0.49	3.3	1.7
% Change from:							
2006	-13%	-30%	-28%	-37%	-14%	-9%	-18%
Mean ^d	-24%	-40%	12%	-25%	-37%	-13%	-46%

^a Proportion of total nests that produced 1 or more young to near fledging age.

^b Total number of young in fall divided by total number of nests and broods (known territorial pairs) in spring.

^c Spring survey not conducted.

^d Mean excludes 2007.

Appendix. Results of 2007 spring and fall trumpeter swan surveys by 1:63,360 topographic map.

**TRUMPETER SWAN SURVEYS
COPPER RIVER DELTA**

SPRING 2007

MAP	QUAD	DATE	NO. OF OBS	AS SNG	IN PRS	IN FKS	NO. OF PRS	NO. OF FKS	PRS w/ NEST	SNG w/ NEST	0 w/ NEST	PRS w/ BRD	SNG w/ BRD	0 w/ BRD	NO. OF NESTS	NO. OF BRDS	TOT ADU	TOT YNG	TOT SWANS
CORDOVA	A1	05/19/07	47	6	80	5	40	1	16	1	0	0	0	0	17	0	91	0	91
CORDOVA	A2	05/19/07	11	5	12	0	6	0	4	0	0	0	0	0	4	0	17	0	17
CORDOVA	B1	05/19/07	41	11	52	73	26	4	7	1	0	0	0	0	8	0	136	0	136
CORDOVA	B2	05/20/07	49	12	74	0	37	0	14	0	0	0	0	0	14	0	86	0	86
CORDOVA	B3	05/20/07	85	8	148	38	74	3	34	1	0	0	0	0	35	0	194	0	194
CORDOVA	B4	05/19/07	125	21	198	25	99	4	40	0	1	0	0	0	41	0	244	0	244
CORDOVA	B5	05/20/07	44	14	60	0	30	0	9	2	0	0	0	0	11	0	74	0	74
CORDOVA	C2	05/20/07	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
CORDOVA	C3	05/20/07	3	2	2	0	1	0	0	0	0	0	0	0	0	0	4	0	4
CORDOVA	C4	05/20/07	4	2	4	0	2	0	1	0	0	0	0	0	1	0	6	0	6
CORDOVA	C5	05/20/07	24	3	40	11	20	1	6	0	0	0	0	0	6	0	54	0	54
TOTAL			433	84	670	152	335	13	131	5	1	0	0	0	137	0	906	0	906

FALL 2007

MAP	QUAD	DATE	NO. OF OBS	AS SNG	IN PRS	IN FKS	NO. OF PRS	NO. OF FKS	PRS w/ NEST	SNG w/ NEST	0 w/ NEST	PRS w/ BRD	SNG w/ BRD	0 w/ BRD	NO. OF NESTS	NO. OF BRDS	TOT ADU	TOT YNG	TOT SWANS
CORDOVA	A1	08/22/07	41	4	72	4	36	1	--	--	--	5	1	0	--	6	80	15	95
CORDOVA	A2	08/22/07	5	0	10	0	5	0	--	--	--	0	0	0	--	0	10	0	10
CORDOVA	B1	08/22/07	42	1	62	151	31	10	--	--	--	4	0	0	--	4	214	11	225
CORDOVA	B2	08/22/07	33	4	54	37	27	2	--	--	--	5	0	0	--	5	95	16	111
CORDOVA	B3	08/22/07	61	2	108	24	54	5	--	--	--	9	0	0	--	9	134	27	161
CORDOVA	B4	08/22/07	79	16	116	28	58	5	--	--	--	15	3	0	--	18	160	53	213
CORDOVA	B5	08/23/07	17	3	22	10	11	3	--	--	--	0	1	0	--	1	35	3	38
CORDOVA	C2	08/23/07	3	0	6	0	3	0	--	--	--	0	0	0	--	0	6	0	6
CORDOVA	C3	08/23/07	4	0	8	0	4	0	--	--	--	0	0	0	--	0	8	0	8
CORDOVA	C4	08/20/07	4	0	6	3	3	1	--	--	--	0	0	0	--	0	9	0	9
CORDOVA	C5	08/24/07	12	0	22	3	11	1	--	--	--	0	0	0	--	0	25	0	25
TOTAL			301	30	486	260	243	28	--	--	--	38	5	0	--	43	776	125	901