

Methadone Dosing Recommendations for Treatment of Chronic Pain

Prepared by Francine Goodman, PharmD, BCPS; William N. Jones, BSc, MSc; and Peter Glassman, MBBS, MSc

Summary

- Methadone is a safe and effective long-acting opioid analgesic that is useful in managing chronic pain.
- Although it has unique pharmacokinetic and pharmacodynamic properties, the general principles of dosing methadone are similar to those of other opioids.
- In general, as with other opioids, methadone should be used as one aspect of a comprehensive pain management plan, as agreed upon by the practitioner and the patient.
- Methadone is most easily titrated by using small initial doses or adjusting the initial dose according to the previous opioid dose.
- A number of methods are available for titrating methadone using conversion ratios, as detailed below. However, titration should be based on patient response and not solely based on equianalgesic dosing tables.
- Consultation with a pain specialist, clinical pharmacist, or other practitioner who has experience with using methadone for chronic pain is recommended if questions arise about dosing or titrating methadone.^a

Background

Methadone should be used when a strong opioid is needed and the patient has not achieved adequate pain relief on escalating doses of controlled-release morphine or has experienced intolerable adverse effects on controlled-release morphine. Commonly, nonsteroidal anti-inflammatory drugs and adjuvant agents (e.g., tricyclic antidepressants) should be used in combination with methadone. Methadone's duration of effect is not dependent upon a specialized delivery system, as is the case with transdermal fentanyl or sustained release formulations of morphine or oxycodone. It is the only long-duration opioid available as an oral solution.

While methadone has gained increasing acceptance as an alternative to morphine for treatment of moderate to severe pain, a number of authors have cautioned clinicians about the complexities of dosing methadone or have suggested the drug be prescribed by practitioners with relevant experience in an adequately monitored setting.¹⁻⁷ Significant toxicity has occurred particularly when dosage increments were made too frequently, conversion doses were too high, or dosing intervals were too close.^{5,8-10} Accruing experience, however, suggests that methadone can be safely used when initial doses are small, conversion ratios are adjusted to the previous opioid dose, and dosage is slowly titrated to patient response.^{2,3,5,6,9,11-15} The general principles of dosing methadone are similar to those of other opioids.

The pharmacokinetic and pharmacodynamic properties of methadone are complex and incompletely documented.^{16,17} Although methadone may have a long elimination half-life (up to a mean/median of 128 h),¹⁸⁻³¹ the elimination half-life does not necessarily reflect duration of analgesia.^{28,32} Patients may require dosing intervals of 6 hours to achieve adequate pain relief, although repeated oral administration of methadone for cancer pain may lead to progressively longer dosing intervals.^{33,34} As a result of the dissociation between half-life and analgesic duration, tissue accumulation of methadone can occur. Patients need to be reassessed more frequently (e.g., every few days) when methadone is initiated and when the dose is increased. However, once a stable dosing is established, follow-up can be as clinically indicated. With a 3-day phased conversion from morphine to methadone, the analgesic effects have taken a median of 5 days (range: 4 to 13 days) to stabilize.³

It is important to note that the equianalgesic conversion ratios between methadone and other opioids are imprecise (Table 1).

Table 1 Points to consider about equianalgesic conversion ratios

A number of equianalgesic dosing tables underestimate the potency of methadone. [†]	† Management of Cancer Pain, Clinical Practice Guidelines, AHCPR (1994) ³⁵ ; Cancer pain: a monograph on the management of cancer pain, Health & Welfare Canada (1984) ³⁶ ; Twycross (1990) ³⁷ ; Levy (1985) ³⁸
Conversion ratios in many equianalgesic dosing tables do not apply to repeated doses of opioids.	‡ The oral morphine to oral methadone conversion ratio may be unexpectedly much higher in patients who previously received very high doses of morphine. ^{24,39}
The morphine- or hydromorphone-to-methadone conversion ratio increases (i.e., the potency of methadone increases) as the previous dose of morphine or hydromorphone increases. [‡]	§ Bruera (1999) ⁴⁰
Conversion ratios may not be bi-directional (i.e., the morphine-to-methadone conversion ratio may not be the same as the methadone-to-morphine ratio). [§]	Estimated ratio based on single-dose, double-blind, double-dummy, cross-over studies in patients with moderate to severe cancer pain. ¹
There may be large interpatient variability in the equianalgesic conversion ratio; a single ratio may not be applicable to all patients. [§]	
The use of high but ineffective doses of previous opioid may result in overestimation of the equivalent dose of methadone.	
The relative analgesic potency ratio of oral to parenteral methadone is 2:1; however, confidence intervals are wide.	

a For more information on identifying patients who should be referred to a pain specialist or pain clinic and on dosing methadone, see the Web-based educational program for VA employees entitled *Opioids in the Management of Acute and Chronic Pain*; available at: <http://vaww.sites.lrn.va.gov/pain/opioids/>.

Certain drug interactions may also potentially affect methadone dosage requirements (Table 2).

Table 2 Potential clinically relevant drug interactions with methadone

Agents that may DECREASE methadone concentrations	Agents that may INCREASE methadone concentrations	Agents that may increase the adverse effects of methadone
<i>Antiepileptics:</i> carbamazepine, phenobarbital, phenytoin (no interaction with valproic acid and gabapentin)	<i>Antidepressants:</i> selective serotonin reuptake inhibitors (venlafaxine least likely to interact); amitriptyline	Benzodiazepines St. John's Wort
<i>Antipsychotics:</i> risperidone	<i>Antifungals:</i> fluconazole, ketoconazole	
<i>Antiretrovirals:</i> nevirapine, ritonavir		
<i>Antitubercular:</i> rifampin (no interaction with rifabutin)		

Sources: Davis (2001)⁴¹; Natural Medicines Comprehensive Database⁴²; Plummer (1988)^{30,43}

The present dosing recommendations are provided to offer guidance on dosing methadone in the treatment of patients with chronic nonmalignant pain (CNMP) or chronic cancer pain, particularly when converting from another opioid to methadone. If in doubt, a practitioner should consult a pain management specialist, clinical pharmacist, or another practitioner who has the relevant knowledge.^b

Dosing Strategies

The best titration strategy has not been determined. Any methadone dosing strategy could be used for treating either CNMP or chronic cancer pain. The rapidity of conversion may be more important than type of pain in determining which method is useful in a given clinical situation. Therefore, the dosing strategies have been categorized by previous opioid exposure and rapidity of titration or conversion (Table 3 and Table 4). The dosing recommendations shown here represent a conservative approach to titrating methadone.

Table 3 Dosing recommendations for patients receiving codeine preparations or no previous opioids

Dosing strategy	Initial MET dose	Increments	Comments
Gradual titration (For CNMP and situations necessitating less frequent monitoring) ⁴⁴	2.5 mg q 8 h	2.5 mg q 8 h every 5 to 7 d	As a general rule, <i>start low and go slow</i> .
Faster titration (For cancer pain and situations where frequent monitoring is possible)	2.5 mg q 6 or 8 h	2.5 mg q 6 or 8 h as often as every day over about 4 d	

The dosing recommendations for gradual titration were modified with permission from *Evidence-Based Recommendations for Medical Management of Chronic Non-Malignant Pain*, College of Physicians and Surgeons of Ontario, November 2000. **All doses refer to oral administration.** CNMP = Chronic nonmalignant pain; MET = Methadone

Table 4 Dosing recommendations for patients previously receiving other opioids

Dosing strategy	Initial MET dose		Increments	Comments
Gradual conversion (For CNMP and situations necessitating less frequent monitoring) ⁴⁴	<u>MOR-E Dose</u> < 200 mg/d	<u>MET Dose</u> 5 mg q 8 h	Increase by the calculated MET dose every 5 to 7 d as needed. <i>For patients switched from greater than 500 mg/d MOR-E:</i> add 1/3 rd of MET dose (2.3% of daily MOR-E dose) every 5 d (total conversion dose [period]: ~7% of MOR-E dose [15 d]). Reduce dose of previous opioid by 1/3 rd every 5 d.	Previous MOR-E dose < 200 mg/d includes patients already on a major opioid analgesic like oxycodone with or without acetaminophen. In patients with CNMP, look for a graded analgesic response to dosage increments; if absent, the patient may have opioid-nonresponsive pain.
	200 to 500 mg/d	~7% of MOR-E dose, given in divided doses q 8 h (see below)		
	> 500 mg/d	~7% of MOR-E dose; give 1/3 rd of MET dose (2.3% of MOR-E dose) in divided doses q 8 h (see below)		
	For patients with CNMP who have received repeated doses of > 200 mg/d MOR-E, calculate MET dose using the table below.			
	<u>Drug</u>	<u>Oral</u>	<u>Drug/MOR</u>	
	MET	2 mg	7%	
	MOR	30 mg	100%	
	HMO	8 mg	27%	
	OXY	15 mg	50%	
Faster conversion (For cancer pain and situations where frequent monitoring is possible) ^{2,3,5,11,12,45,46}	<u>MOR-E Dose</u> < 200 mg/d	<u>MET/MOR-E</u> 10% to 30%	Two methods may be used: (1) <i>Phased conversion.</i> Replace 1/3 of MOR-E dose with the calculated equivalent dose of MET daily for 3 days. (2) <i>Rapid ("stop-and-go") conversion.</i> Discontinue MOR-E and start MET on day 1.	Titrate MET day by day according to patient's symptoms and the number of BTP doses administered. Smaller MET-to-MOR-E conversion proportions (%) should be used the larger the previous MOR-E dose.
	200 to 500 mg/d	10% to 20%		
	501 to 1000 mg/d	5% to 10%		
	> 1000 mg/d	5% or less		
	For the most conservative approach, use 5% MET/MOR-E (or less with very high MOR-E doses) to calculate the initial MET dose irrespective of the previous MOR-E dose.			
	Give starting dose of MET in divided doses q 8 h (up to a maximum 50 mg q 8 h).			

The dosing recommendations for gradual conversion were modified with permission from *Evidence-Based Recommendations for Medical Management of Chronic Non-Malignant Pain*, College of Physicians and Surgeons of Ontario, November 2000. The dosing strategy for faster conversion is based on a synthesis of the most recent versions of the more notable dosing strategies used in opioid-tolerant patients with mostly cancer-related pain.

All doses refer to oral administration.

BTP = Breakthrough pain; CNMP = Chronic nonmalignant pain; HMO = Hydromorphone; MET = Methadone; MOR = Morphine; MOR-E = Morphine-equivalent; OXY = Oxycodone

^b See footnote "a" on page 1.

It is important to note that various dosing methods have been used (including a patient-controlled regimen^{6,47}) and are still evolving. Two dosing strategies^{2,11} have been prospectively studied, but no clinical trials comparing systematic dosing methods have been performed. A literature search (PubMed 1966 to 2001) identified only a small case series that discussed methadone dosing during the treatment of CNMP.⁴⁸ The lack of prospective and comparative studies highlights the need to carefully individualize the dosing regimen of methadone, as is done with other opioids.

As a general rule, smaller methadone-to-morphine conversion proportions (%) should be used the larger the previous morphine-equivalent dose, remembering that precise conversions from another opioid to methadone are impossible. Disproportionately smaller methadone doses may be required with the larger morphine doses. However, it is important to remember that the equianalgesic conversion ratio is only one part of the process of properly dosing methadone and other opioids.

For breakthrough pain (BTP), a short-acting opioid preparation (such as acetaminophen with codeine, oxycodone with or without acetaminophen, or immediate-release morphine) may be used as necessary. Keep in mind that the use of BTP medications in patients with CNMP is controversial. If opioid medications for BTP are indicated following titration to a stable methadone dose in a patient with CNMP, they should be used sparingly.⁴⁴ Methadone has also been used (in doses 10% to 30% of the calculated daily methadone dose up to 3 to 8 doses per day as needed)^{6,11,46,47}; however, the short-acting opioids are generally preferred to avoid drug accumulation.

Special patient populations

Patients 65 years and older may have a decreased clearance of methadone.³⁰ In patients with stable chronic liver disease, no dosage adjustments appear to be necessary.⁴⁹ Methadone and its metabolites do not accumulate in patients with renal failure.⁵⁰ The two prospective studies on methadone dosing strategies excluded patients with liver or renal disease.^{2,11} Use extra caution when dosing any opioid in all of these patient populations.^c

General principles for dosing methadone

- Use methadone for treatment of patients with chronic pain.
- Individualize doses and slowly titrate to response.
- An acceptable balance between analgesic effects and tolerable and manageable adverse effects generally indicates a favorable response to pain medication. In the treatment of CNMP, the main goals are to improve the patient's ability to function and to increase the patient's quality of life.
- Once the daily dosage for adequate analgesia has been determined, a trial of longer dosing intervals may be attempted. Many patients can take the same total daily dose divided every 8 hours. Intervals of 12 hours may be attempted when patients are stable at 8-hour intervals.
- If a patient develops sedation (which may be a precursor to respiratory depression), hold or decrease the following dose of previous opioid or methadone (depending on the dosing strategy) and decrease subsequent doses and/or make dosage increments less frequently. Do not increase the dose of methadone.
- Short-acting opioids may be used for treatment of BTP, at least initially and when pain is severe and escalating.
- The use of medications for BTP in the treatment of CNMP is controversial. If medications for BTP are indicated after titration to a stable methadone dose, they should be used sparingly.⁴⁴
- Reassess patients at appropriate intervals; at least once weekly during titration and at least once monthly after the daily dosage is stabilized.
- Use additional caution with elderly patients (≥ 65 years), patients with liver, renal, or pulmonary disease, debilitated patients, and patients previously receiving high doses of opioid. Patients who cannot be adequately monitored at home may be considered for inpatient titration of methadone.

Patient education

- Explain to patients that the initial dose will often be inadequate for pain relief. BTP medication should be used during the dose titration period. A pain and pain medicine diary should be kept.
- Reassure patients that the dose will be titrated to achieve adequate analgesia.
- Advise patients that the effects of methadone will increase over at least one week following a dosage increment. Pain relief during the last few days of that week will be greater than at the first few days of the week.
- Remind patients about the need for and the frequency of monitoring during the titration and maintenance periods. Provide patients with instructions on what to do if they develop increasing or intolerable adverse effects.
- Advise patients to avoid abrupt discontinuation of their opioid medication without first consulting their physician. Educate patients about withdrawal symptoms.
- Since patients may become concerned about the social stigma associated with the use of methadone for treatment of opioid dependence, reassure them that methadone is also an accepted pain medication and that they are not "addicts" because they are taking methadone for pain control. Explain the difference between addiction and dependence.^d

^c For patients with liver or renal disease, special consideration can be given locally to use an alternative opioid at the discretion of the care team or provider.

^d For more information on the definitions of addiction and dependence, see the Web-based educational program for VA employees entitled *Opioids in the Management of Acute and Chronic Pain*; available at: <http://vaww.sites.lrn.va.gov/pain/opioids/> or reference 51.

References

1. Foley KM, Houde RW. Methadone in cancer pain management: individualize dose and titrate to effect. *J Clin Oncol* 1998;16:3213-5.
2. Ripamonti C, Groff L, Brunelli C, Polastri D, Stavakis A, De Conno F. Switching from morphine to oral methadone in treating cancer pain: what is the equianalgesic dose ratio? *J Clin Oncol* 1998;16:3216-21.
3. Lawlor PG, Turner KS, Hanson J, Bruera ED. Dose ratio between morphine and methadone in patients with cancer pain: a retrospective study. *Cancer* 1998;82:1167-73.
4. Bruera E, Pereira J, Watanabe S, Belzile M, Kuehn N, Hanson J. Opioid rotation in patients with cancer pain. A retrospective comparison of dose ratios between methadone, hydromorphone, and morphine. *Cancer* 1996;78:852-7.
5. Ayonrinde OT, Bridge DT. The rediscovery of methadone for cancer pain management. *Med J Aust* 2000;173:536-40.
6. Morley JS, Makin MK. Comments on Ripamonti et al., Pain, 70 (1997) 109-115. *Pain* 1997;73:114-5.
7. Hanks GW, Conno F, Cherny N et al. Morphine and alternative opioids in cancer pain: the EAPC recommendations. *Br J Cancer* 2001;84:587-93.
8. Symonds P. Methadone and the elderly (letter). *Br Med J* 1977;1:512.
9. Bruera E, Watanabe S, Fainsinger RL, Spachynski K, Suarez-Almazor M, Inturrisi C. Custom-made capsules and suppositories of methadone for patients on high-dose opioids for cancer pain. *Pain* 1995;62:141-6.
10. Ettinger DS, Vitale PJ, Trump DL. Important clinical pharmacologic considerations in the use of methadone in cancer patients. *Cancer Treat Rep* 1979;63:457-9.
11. Mercadante S, Casuccio A, Fulfaro F et al. Switching from morphine to methadone to improve analgesia and tolerability in cancer patients: a prospective study. *J Clin Oncol* 2001;19:2898-904.
12. Gagnon B, Bruera E. Differences in the ratios of morphine to methadone in patients with neuropathic pain versus non-neuropathic pain. *J Pain Symptom Manage* 1999;18:120-5.
13. Mercadante S, Casuccio A, Calderone L. Rapid switching from morphine to methadone in cancer patients with poor response to morphine. *J Clin Oncol* 1999;17:3307-12.
14. Hagen NA, Wasylenko E. Methadone: outpatient titration and monitoring strategies in cancer patients. *J Pain Symptom Manage* 1999;18:369-75.
15. Krames E. The Bruera/Neumann article reviewed. Discussion of Bruera E, Neumann CM. Role of methadone in the management of pain in cancer patients. *Oncology* 1999;13:1275-1282. *Oncology* 1999;13:1288-1289.
16. Ripamonti C, Zecca E, Bruera E. An update on the clinical use of methadone for cancer pain. *Pain* 1997;70:109-15.
17. Garrido MJ, Troconiz IF. Methadone: a review of its pharmacokinetic/pharmacodynamic properties. *J Pharmacol Toxicol Methods* 1999;42:61-6.
18. Wolff K, Rostami-Hodjegan A, Shires S et al. The pharmacokinetics of methadone in healthy subjects and opiate users. *Br J Clin Pharmacol* 1997;44:325-34.
19. Olsen GD, Wendel HA, Livermore JD, Leger RM, Lynn RK, Gerber N. Clinical effects and pharmacokinetics of racemic methadone and its optical isomers. *Clin Pharmacol Ther* 1977;21:147-57.
20. Verebely K, Volavka J, Mule S, Resnick R. Methadone in man: pharmacokinetic and excretion studies in acute and chronic treatment. *Clin Pharmacol Ther* 1975;18:180-90.
21. Inturrisi CE, Verebely K. Disposition of methadone in man after a single oral dose. *Clin Pharmacol Ther* 1972;13:923-30.
22. Wolff K, Rostami-Hodjegan A, Hay AW, Raistrick D, Tucker G. Population-based pharmacokinetic approach for methadone monitoring of opiate addicts: potential clinical utility. *Addiction* 2000;95:1771-83.
23. de Vos JW, Geerlings PJ, van den Brink W, Ufkes JG, van Wilgenburg H. Pharmacokinetics of methadone and its primary metabolite in 20 opiate addicts. *Eur J Clin Pharmacol* 1995;48:361-6.
24. Wolff K, Hay AW, Raistrick D, Calvert R. Steady-state pharmacokinetics of methadone in opioid addicts. *Eur J Clin Pharmacol* 1993;44:189-94.
25. Nilsson MI, Gronbladh L, Widerlov E, Anggard E. Pharmacokinetics of methadone in methadone maintenance treatment: characterization of therapeutic failures. *Eur J Clin Pharmacol* 1983;25:497-501.
26. Anggard E, Nilsson MI, Holmstrand J, Gunne LM. Pharmacokinetics of methadone during maintenance therapy: pulse labeling with deuterated methadone in the steady state. *Eur J Clin Pharmacol* 1979;16:53-7.
27. Nilsson MI, Anggard E, Holmstrand J, Gunne LM. Pharmacokinetics of methadone during maintenance treatment: adaptive changes during the induction phase. *Eur J Clin Pharmacol* 1982;22:343-9.
28. Inturrisi CE, Colburn WA, Kaiko RF, Houde RW, Foley KM. Pharmacokinetics and pharmacodynamics of methadone in patients with chronic pain. *Clin Pharmacol Ther* 1987;41:392-401.
29. Gourlay GK, Cherry DA, Cousins MJ. A comparative study of the efficacy and pharmacokinetics of oral methadone and morphine in the treatment of severe pain in patients with cancer. *Pain* 1986;25:297-312.
30. Plummer JL, Gourlay GK, Cherry DA, Cousins MJ. Estimation of methadone clearance: application in the management of cancer pain. *Pain* 1988;33:313-22.
31. Denson DD, Concilus RR, Warden G, Raj PP. Pharmacokinetics of continuous intravenous infusion of methadone in the early post-burn period. *J Clin Pharmacol* 1990;30:70-5.
32. Grochow L, Sheidler V, Grossman S, Green L, Enterline J. Does intravenous methadone provide longer lasting analgesia than intravenous morphine? A randomized, double-blind study. *Pain* 1989;38:151-7.
33. Hanson J, Gimman C, Hartvig P, et al. Clinical evaluation of oral methadone in treatment of cancer pain. *Acta Anaesthesiol Scand* 1982;74:124-127.
34. Sawe J, Hansen J, Gimman C et al. Patient-controlled dose regimen of methadone for chronic cancer pain. *Br Med J (Clin Res Ed)* 1981;282:771-3.
35. AHCPR. Management of Cancer Pain, Clinical Practice Guidelines. Rockville, MD: Agency for Health Care Policy and Research; U.S. Department of Health and Human Services; 1994. AHCPR Pub. No. 94-0592.
36. Health & Welfare Canada. Cancer pain: a monograph on the management of cancer pain. Ottawa, Canada: Health & Welfare Canada, Minister of Supply and Services; 1984. H42-2/5.
37. Twycross R, Lack S. Pain relief. In: Twycross R, Lack S, eds. *Therapeutics in terminal cancer, 2nd edition*. Edinburgh: Churchill Livingstone; 1990;2:11-39.
38. Levy MH. Pain management in advanced cancer. *Semin Oncol* 1985;12:394-410.
39. Ripamonti C, De Conno F, Groff L et al. Equianalgesic dose/ratio between methadone and other opioid agonists in cancer pain: comparison of two clinical experiences. *Ann Oncol* 1998;9:79-83.
40. Bruera E, Neumann CM. Role of methadone in the management of pain in cancer patients. *Oncology (Huntingt)* 1999;13:1275-82; discussion 1285-8, 1291.
41. Davis MP, Walsh D. Methadone for relief of cancer pain: a review of pharmacokinetics, pharmacodynamics, drug interactions and protocols of administration. *Support Care Cancer* 2001;9:73-83.
42. Jellin JM, Gregory P, Batz F, Hitchens K, et al. Pharmacist's Letter/Prescriber's Letter Natural Medicines Comprehensive Database, 3rd ed. Stockton, CA: Therapeutic Research Faculty; 2000.
43. Brown LS, Sawyer RC, Li R, Cobb MN, Colborn DC, Narang PK. Lack of a pharmacologic interaction between rifabutin and methadone in HIV-infected former injecting drug users. *Drug Alcohol Depend* 1996;43:71-7.
44. CPSO Task Force on CNMP. Evidence-based recommendations for medical management of chronic non-malignant pain: College of Physicians and Surgeons of Ontario (CPSO); Nov 2000.
45. De Conno F, Groff L, Brunelli C, Zecca E, Ventafredda V, Ripamonti C. Clinical experience with oral methadone administration in the treatment of pain in 196 advanced cancer patients. *J Clin Oncol* 1996;14:2836-42.
46. Friedman LL. Using Methadone. Lecture presented at: American Academy of Hospice and Palliative Medicine, 13th Annual Assembly; 22 June 2001; Phoenix, AZ.
47. Morley J, Makin M. The use of methadone in cancer pain poorly responsive to other opioids. *Pain Rev* 1998;5:51-58.
48. Gardner-Nix JS. Oral methadone for managing chronic nonmalignant pain. *J Pain Symptom Manage* 1996;11:321-8.
49. Novick DM, Kreek MJ, Fanizza AM, Yancovitz SR, Gelb AM, Stenger RJ. Methadone disposition in patients with chronic liver disease. *Clin Pharmacol Ther* 1981;30:353-62.
50. Kreek MJ, Schechter AJ, Gutjahr CL, Hecht M. Methadone use in patients with chronic renal disease. *Drug Alcohol Depend* 1980;5:197-205.
51. Portenoy RK. Pain specialists and addiction medicine specialists unite to address critical issues. American Pain Society Web site. APS bulletin (online). 1999;9(2). Available at: <http://www.ampainoc.org/pub/bulletin/mar99/president.htm>. Accessed 5 October 2001.