

# TABLES

TABLE 1. Sample Sizes and Response Rates, Selected U.S. Sites: School Health Profiles, Principals' and Teachers' Surveys, 2002

**STATE SURVEYS**

Site	Principals' surveys		Teachers' surveys	
	Sample size	Response rate (%)	Sample size	Response rate (%)
<b>Weighted Data</b>				
Alabama	285	77	269	73
Alaska	231	72	225	71
Arizona	345	91	329	87
Arkansas	261	73	NA*	NA
Connecticut	236	72	237	72
Delaware	52	80	49	75
Georgia	266	71	260	70
Hawaii	66	74	62	70
Idaho	192	87	194	87
Illinois**	380	78	382	78
Iowa	263	76	262	76
Kentucky	235	72	236	72
Maine	197	86	204	90
Massachusetts	608	88	591	86
Michigan	331	82	322	80
Minnesota	282	75	269	72
Missouri	354	84	351	84
Montana	255	77	255	77
Nebraska	320	79	309	76
New Hampshire	171	81	154	73
New Jersey	333	79	NA	NA
New Mexico	179	76	174	74
New York	319	72	321	72
North Carolina	319	74	NA	NA
North Dakota	175	77	164	73
Oklahoma	322	73	NA	NA
Tennessee	316	84	314	83
Texas**	358	71	NA	NA
Utah	215	87	189	76
Vermont	116	76	107	70
Virginia	267	73	255	70
Wisconsin	361	79	352	77
<b>Unweighted Data</b>				
Arkansas	NA	NA	236	66
California	351	68	314	61
North Carolina	NA	NA	251	59
Oklahoma	NA	NA	300	68
Oregon	183	50	NA	NA
Pennsylvania**	302	66	268	58
Rhode Island	65	60	65	60
South Carolina	265	62	281	65
Texas**	NA	NA	266	53
Washington	248	60	NA	NA

(continued)

TABLE 1. Sample Sizes and Response Rates, Selected U.S. Sites: School Health Profiles, Principals' and Teachers' Surveys, 2002  
(continued)

### LOCAL SURVEYS

Site	Principals' surveys		Teachers' surveys	
	Sample size	Response rate (%)	Sample size	Response rate (%)
<b>Weighted Data</b>				
Chicago	223	80	217	78
Dallas	51	98	51	98
Fort Lauderdale	43	73	44	75
Houston	46	77	52	87
Los Angeles	104	87	104	87
Miami	94	92	93	91
Orange County	41	84	40	82
Palm Beach	39	85	41	89
San Bernardino	14	93	NA	NA
San Diego	49	100	49	100
San Francisco	38	84	34	76
<b>Unweighted Data</b>				
District of Columbia	31	69	31	69
Philadelphia	87	65	85	64
San Bernardino	NA	NA	8	53

\* NA = not available.

\*\* Survey did not include students from one of the state's large school districts.

TABLE 2. Percentage of Schools That Required Health Education in Grades 6-12 and, Among Those Schools, Percentage That Taught One or More Separate Required Health Education Courses, Selected U.S. Sites: School Health Profiles, Principals' Surveys, 2002

**STATE SURVEYS**

Site	Required health education	Taught one or more separate required health education courses
<b>Weighted Data</b>		
Alabama	89.8	84.5
Alaska	91.2	93.2
Arizona	68.3	83.4
Arkansas	96.7	99.2
Connecticut	92.8	93.2
Delaware	98.0	91.8
Georgia	92.5	94.0
Hawaii	100.0	98.6
Idaho	95.9	97.4
Illinois*	96.6	97.4
Iowa	80.3	91.2
Kentucky	91.3	93.8
Maine	96.5	95.3
Massachusetts	93.6	96.0
Michigan	83.3	91.6
Minnesota	96.6	98.1
Missouri	91.8	93.7
Montana	95.2	95.3
Nebraska	89.7	93.0
New Hampshire	88.5	91.7
New Jersey	99.0	95.7
New Mexico	74.8	92.2
New York	99.1	99.7
North Carolina	97.1	91.6
North Dakota	92.0	98.9
Oklahoma	32.7	77.6
Tennessee	81.9	88.4
Texas*	86.6	93.7
Utah	97.2	100.0
Vermont	90.5	82.0
Virginia	91.2	90.5
Wisconsin	94.6	97.8
<b>Unweighted Data</b>		
California	83.5	82.6
Oregon	100.0	97.1
Pennsylvania*	98.3	99.0
Rhode Island	95.3	100.0
South Carolina	87.0	88.1
Washington	88.1	94.2
<b>State Median</b>	<b>92.3</b>	<b>93.7</b>

(continued)

TABLE 2. Percentage of Schools That Required Health Education in Grades 6-12 and, Among Those Schools, Percentage That Taught One or More Separate Required Health Education Courses, Selected U.S. Sites: School Health Profiles, Principals' Surveys, 2002 (continued)

### LOCAL SURVEYS

Site	Required health education	Taught one or more separate required health education courses
<b>Weighted Data</b>		
Chicago	88.3	77.1
Dallas	69.7	96.9
Fort Lauderdale	89.9	79.8
Houston	93.3	97.5
Los Angeles	97.1	97.0
Miami	68.8	79.5
Orange County	81.1	82.2
Palm Beach	80.2	87.3
San Bernardino	71.4	80.0
San Diego	87.5	64.3
San Francisco	100.0	89.5
<b>Unweighted Data</b>		
District of Columbia	96.7	100.0
Philadelphia	97.7	90.1
<b>Local Median</b>	<b>88.3</b>	<b>87.3</b>

\* Survey did not include students from one of the state's large school districts.

TABLE 3. Percentage of Schools That Required Teachers to Use Standards, a Specific Curriculum, Guidelines, Framework, or Other Selected Materials,\* Selected U.S. Sites: School Health Profiles, Teachers' Surveys, 2002

**STATE SURVEYS**

Site	National Health Education Standards	State, district, or school curriculum, guidelines, or framework	Materials from health organizations	Commercial teacher's guide
<b>Weighted Data</b>				
Alabama	47.5	100.0	61.7	70.8
Alaska	28.9	91.1	18.8	44.9
Arizona	36.9	95.0	49.8	49.7
Connecticut	50.8	98.4	34.4	25.6
Delaware	68.9	95.6	29.4	34.5
Georgia	41.9	100.0	46.7	80.2
Hawaii	62.1	96.8	27.5	27.6
Idaho	26.9	89.9	22.5	55.6
Illinois**	38.9	94.6	32.6	60.1
Iowa	36.0	88.7	26.8	39.5
Kentucky	42.9	98.9	33.2	57.0
Maine	26.5	95.6	17.7	25.8
Massachusetts	45.1	97.5	45.2	37.2
Michigan	30.3	94.9	36.4	35.7
Minnesota	36.7	93.2	39.0	39.1
Missouri	46.5	98.1	36.9	55.8
Montana	40.3	93.7	33.5	42.8
Nebraska	22.1	87.4	36.1	49.4
New Hampshire	44.7	85.7	28.3	33.2
New Mexico	64.6	97.1	49.6	63.4
New York	55.2	97.6	36.9	34.7
North Dakota	31.5	82.9	28.6	50.9
Tennessee	38.1	98.8	42.8	65.5
Utah	19.9	98.8	23.7	42.9
Vermont	34.3	87.3	33.2	34.4
Virginia	37.3	98.5	42.9	60.8
Wisconsin	44.0	96.7	34.9	36.5
<b>Unweighted Data</b>				
Arkansas	37.2	96.7	29.9	67.6
California	34.0	92.8	38.6	45.7
North Carolina	40.1	99.5	55.5	63.0
Oklahoma	37.5	90.8	43.8	56.0
Pennsylvania**	56.0	95.9	46.4	52.4
Rhode Island	68.3	91.8	43.3	30.5
South Carolina	55.9	97.9	33.5	72.1
Texas**	30.8	99.5	43.8	67.8
<b>State Median</b>	<b>38.9</b>	<b>95.9</b>	<b>36.1</b>	<b>49.4</b>

(continued)

TABLE 3. Percentage of Schools That Required Teachers to Use Standards, a Specific Curriculum, Guidelines, Framework, or Other Selected Materials,\* Selected U.S. Sites: School Health Profiles, Teachers' Surveys, 2002 (continued)

### LOCAL SURVEYS

Site	National Health Education Standards	State, district, or school curriculum, guidelines, or framework	Materials from health organizations	Commercial teacher's guide
<b>Weighted Data</b>				
Chicago	47.2	98.0	58.0	63.3
Dallas	57.0	100.0	89.6	75.8
Fort Lauderdale	52.1	100.0	58.5	69.4
Houston	58.5	100.0	71.1	76.6
Los Angeles	36.4	100.0	32.8	47.6
Miami	47.5	100.0	67.7	62.5
Orange County	51.6	100.0	40.7	62.5
Palm Beach	32.1	100.0	58.0	68.6
San Diego	24.4	100.0	92.9	35.7
San Francisco	45.3	90.7	31.1	33.1
<b>Unweighted Data</b>				
District of Columbia	72.0	96.4	64.3	44.4
Philadelphia	40.0	97.2	31.3	34.9
San Bernardino	66.7	80.0	40.0	80.0
<b>Local Median</b>	<b>47.5</b>	<b>100.0</b>	<b>58.0</b>	<b>62.5</b>

\* In a required health education course.

\*\* Survey did not include students from one of the state's large school districts.

TABLE 4a. Percentage of Schools That Tried to Increase Student Knowledge, by Specific Health-Related Topic,\* Selected U.S. Sites: School Health Profiles, Teachers' Surveys, 2002

**STATE SURVEYS**

Site	Accident or injury prevention	Alcohol or other drug-use prevention	Consumer health	CPR**	Death and dying	Dental and oral health	Emotional and mental health
<b>Weighted Data</b>							
Alabama	97.3	99.5	85.0	87.6	75.2	81.0	96.8
Alaska	89.5	96.4	75.7	46.9	56.3	69.7	89.0
Arizona	82.2	97.9	77.7	45.7	43.9	66.9	85.1
Connecticut	86.4	99.5	70.6	48.9	53.2	40.2	97.8
Delaware	91.1	100.0	82.9	52.3	54.4	47.1	97.6
Georgia	94.3	99.6	81.6	72.0	62.6	71.4	98.0
Hawaii	85.8	100.0	88.4	50.2	38.9	54.8	98.3
Idaho	94.3	98.9	83.7	75.8	74.1	66.6	97.1
Illinois <sup>§</sup>	91.4	98.8	77.5	65.6	58.7	72.3	94.7
Iowa	83.7	98.7	80.4	63.2	57.9	67.6	91.9
Kentucky	96.4	98.7	87.4	73.3	62.8	71.1	98.2
Maine	87.6	99.4	86.3	58.1	47.3	45.6	97.5
Massachusetts	86.9	98.9	79.2	54.3	50.9	52.0	95.4
Michigan	77.0	99.2	73.6	47.7	41.7	56.4	89.7
Minnesota	94.6	99.3	74.9	76.7	57.3	53.0	95.8
Missouri	93.9	98.6	87.9	72.6	61.7	77.0	93.9
Montana	88.0	98.5	79.7	66.6	54.4	63.3	89.8
Nebraska	84.7	98.4	74.4	66.2	61.7	65.8	91.8
New Hampshire	86.3	99.1	81.3	56.7	44.4	53.1	92.7
New Mexico	87.0	100.0	77.9	58.3	67.6	61.8	91.5
New York	91.4	100.0	85.1	62.3	67.2	60.1	98.9
North Dakota	93.8	98.6	85.2	65.9	59.9	76.5	97.9
Tennessee	94.7	98.4	78.1	70.8	58.9	72.7	90.1
Utah	94.7	99.3	88.4	72.0	68.9	59.6	97.9
Vermont	88.6	97.6	74.8	44.1	34.5	54.3	96.0
Virginia	93.8	98.2	84.3	73.0	51.3	64.6	93.5
Wisconsin	88.6	98.8	78.9	63.4	61.2	58.5	96.7
<b>Unweighted Data</b>							
Arkansas	96.2	100.0	81.0	76.7	68.7	78.8	96.2
California	75.2	97.1	73.2	55.4	45.3	54.9	84.4
North Carolina	92.6	99.5	77.8	76.9	55.7	55.8	95.2
Oklahoma	90.9	97.4	77.9	68.8	46.8	82.1	84.4
Pennsylvania <sup>§</sup>	91.8	100.0	77.6	74.2	49.2	59.5	97.2
Rhode Island	88.5	100.0	80.3	59.0	44.3	49.2	93.4
South Carolina	90.3	97.9	81.4	55.9	46.8	73.0	90.9
Texas <sup>§</sup>	93.0	99.5	80.8	77.7	61.0	73.6	90.7
<b>State Median</b>	<b>90.9</b>	<b>98.9</b>	<b>80.3</b>	<b>65.6</b>	<b>56.3</b>	<b>63.3</b>	<b>95.2</b>

(continued)



TABLE 4a. Percentage of Schools That Tried to Increase Student Knowledge, by Specific Health-Related Topic,\*  
Selected U.S. Sites: School Health Profiles, Teachers' Surveys, 2002 (continued)

### LOCAL SURVEYS

Site	Accident or injury prevention	Alcohol or other drug-use prevention	Consumer health	CPR**	Death and dying	Dental and oral health	Emotional and mental health
<b>Weighted Data</b>							
Chicago	90.7	97.4	75.2	55.7	59.1	84.9	84.2
Dallas	100.0	100.0	93.4	93.4	69.9	60.1	100.0
Fort Lauderdale	90.5	100.0	81.3	76.1	63.6	54.8	100.0
Houston	97.8	100.0	82.2	77.1	68.7	81.8	100.0
Los Angeles	94.3	100.0	92.5	63.7	64.5	88.5	92.7
Miami	91.9	100.0	88.2	70.9	58.8	57.8	88.0
Orange County	84.2	100.0	72.9	78.4	69.7	55.4	93.0
Palm Beach	84.5	100.0	89.0	88.7	57.1	68.0	100.0
San Diego	46.3	100.0	66.7	17.1	19.5	19.5	65.0
San Francisco	86.3	100.0	76.4	44.7	67.1	64.0	90.7
<b>Unweighted Data</b>							
District of Columbia	82.1	96.4	71.4	78.6	46.4	82.1	89.3
Philadelphia	91.8	97.3	83.3	52.8	57.7	77.5	91.8
San Bernardino	83.3	100.0	66.7	33.3	50.0	66.7	83.3
<b>Local Median</b>	<b>90.5</b>	<b>100.0</b>	<b>81.3</b>	<b>70.9</b>	<b>59.1</b>	<b>66.7</b>	<b>91.8</b>

\* In a required health education course.

\*\* CPR = cardiopulmonary resuscitation.

§ Survey did not include students from one of the state's large school districts.

TABLE 4b. Percentage of Schools That Tried to Increase Student Knowledge, by Specific Health-Related Topic,\* Selected U.S. Sites: School Health Profiles, Teachers' Surveys, 2002

**STATE SURVEYS**

Site	Environmental health	First aid	Growth and development	HIV** prevention	Human sexuality	Immunization and vaccinations	Nutrition and dietary behavior
<b>Weighted Data</b>							
Alabama	86.1	92.6	90.4	100.0	88.1	78.8	99.0
Alaska	72.8	61.6	80.7	84.4	72.3	68.8	94.2
Arizona	76.9	68.1	88.0	90.6	74.4	62.4	95.5
Connecticut	57.7	65.1	97.1	99.4	94.2	49.6	98.3
Delaware	72.8	72.5	95.6	100.0	97.6	51.9	100.0
Georgia	75.7	87.7	93.9	97.2	93.7	75.0	98.1
Hawaii	59.8	56.7	90.0	100.0	92.9	36.2	98.3
Idaho	72.5	82.8	86.2	96.5	87.3	73.3	98.8
Illinois <sup>§</sup>	75.1	80.1	93.6	97.0	87.7	71.8	98.1
Iowa	73.8	76.0	91.0	98.2	92.2	64.7	98.3
Kentucky	81.7	87.9	90.8	99.0	88.4	72.0	99.5
Maine	65.8	71.5	88.3	97.8	92.6	57.1	98.9
Massachusetts	63.7	68.6	92.8	96.1	91.1	54.0	97.0
Michigan	61.5	57.1	86.7	93.9	85.5	46.4	96.9
Minnesota	60.4	85.6	88.4	99.1	96.0	57.7	98.7
Missouri	78.4	86.5	91.9	96.4	84.4	76.5	99.7
Montana	75.4	77.5	87.5	92.9	80.1	60.2	98.0
Nebraska	72.1	73.8	84.8	95.1	84.6	63.6	97.4
New Hampshire	53.9	65.2	89.1	99.1	93.7	62.4	99.2
New Mexico	70.0	75.2	95.8	99.2	91.2	61.4	98.0
New York	75.6	72.3	94.4	100.0	93.2	71.2	99.6
North Dakota	76.2	80.5	89.4	91.2	82.4	75.1	99.2
Tennessee	75.0	86.5	91.3	95.4	87.9	73.7	97.3
Utah	79.2	73.6	91.5	98.6	89.3	65.1	99.3
Vermont	60.3	61.0	91.6	98.5	96.3	56.5	96.0
Virginia	75.8	85.4	90.3	93.4	86.0	67.3	97.6
Wisconsin	67.0	75.3	93.3	96.8	93.3	55.6	95.9
<b>Unweighted Data</b>							
Arkansas	75.0	87.7	85.8	95.6	79.2	67.8	100.0
California	66.5	65.2	87.0	97.1	88.9	57.8	92.9
North Carolina	68.9	85.6	82.4	98.9	82.3	51.9	99.4
Oklahoma	80.5	88.3	89.6	88.3	62.3	71.4	97.2
Pennsylvania <sup>§</sup>	65.4	83.1	92.7	99.6	90.2	62.4	96.9
Rhode Island	63.3	73.8	90.2	98.3	91.8	56.7	94.9
South Carolina	77.3	79.7	94.7	97.8	94.1	61.3	97.8
Texas <sup>§</sup>	78.0	85.4	90.2	96.6	83.6	67.3	97.5
<b>State Median</b>	<b>72.8</b>	<b>76.0</b>	<b>90.3</b>	<b>97.2</b>	<b>88.9</b>	<b>62.4</b>	<b>98.1</b>

(continued)

TABLE 4b. Percentage of Schools That Tried to Increase Student Knowledge, by Specific Health-Related Topic,\*  
Selected U.S. Sites: School Health Profiles, Teachers' Surveys, 2002 (continued)

### LOCAL SURVEYS

Site	Environmental health	First aid	Growth and development	HIV** prevention	Human sexuality	Immunization and vaccinations	Nutrition and dietary behavior
<b>Weighted Data</b>							
Chicago	78.0	79.9	93.4	90.2	86.3	75.8	98.6
Dallas	89.7	100.0	100.0	100.0	96.6	93.4	100.0
Fort Lauderdale	84.5	87.8	100.0	100.0	100.0	80.6	100.0
Houston	81.9	90.9	100.0	97.7	86.3	77.2	100.0
Los Angeles	88.2	80.5	97.9	100.0	95.8	85.9	100.0
Miami	75.4	72.8	96.4	100.0	98.2	69.6	100.0
Orange County	63.5	67.7	100.0	100.0	100.0	58.1	100.0
Palm Beach	88.3	80.8	96.1	100.0	100.0	73.5	100.0
San Diego	27.5	24.4	63.4	100.0	100.0	34.1	67.6
San Francisco	82.0	59.0	91.3	95.0	95.7	59.0	100.0
<b>Unweighted Data</b>							
District of Columbia	75.0	78.6	92.6	100.0	92.9	88.9	96.2
Philadelphia	76.4	75.0	95.9	98.6	95.9	64.3	100.0
San Bernardino	83.3	50.0	100.0	100.0	100.0	83.3	100.0
<b>Local Median</b>	<b>81.9</b>	<b>78.6</b>	<b>96.4</b>	<b>100.0</b>	<b>96.6</b>	<b>75.8</b>	<b>100.0</b>

\* In a required health education course.

\*\* HIV = human immunodeficiency virus.

§ Survey did not include students from one of the state's large school districts.

TABLE 4c. Percentage of Schools That Tried to Increase Student Knowledge, by Specific Health-Related Topic,\* Selected U.S. Sites: School Health Profiles, Teachers' Surveys, 2002

**STATE SURVEYS**

Site	Personal hygiene	Physical activity and fitness	Pregnancy prevention	STD** prevention	Suicide prevention	Sun safety	Tobacco-use prevention	Violence prevention
<b>Weighted Data</b>								
Alabama	94.0	100.0	89.1	96.8	83.8	82.8	98.4	95.1
Alaska	80.4	96.5	67.1	75.4	67.0	42.7	96.5	85.2
Arizona	87.9	97.8	69.0	75.3	58.2	67.8	99.5	84.8
Connecticut	72.6	95.7	85.1	97.3	80.4	60.8	98.5	91.4
Delaware	80.9	100.0	90.8	95.2	79.5	63.4	100.0	83.6
Georgia	89.3	100.0	91.8	96.4	80.0	78.0	100.0	90.1
Hawaii	80.2	100.0	91.0	94.7	69.3	66.1	100.0	91.6
Idaho	83.9	98.9	78.8	92.1	85.2	78.2	99.4	83.0
Illinois <sup>§</sup>	88.9	99.0	83.7	93.5	77.3	78.0	98.8	84.5
Iowa	86.2	98.3	87.7	94.5	67.4	76.7	98.1	78.7
Kentucky	85.7	99.1	91.4	96.4	77.7	81.5	99.4	90.3
Maine	77.5	98.2	90.0	94.7	73.4	68.0	100.0	87.8
Massachusetts	84.6	97.3	84.5	91.3	74.3	71.6	98.6	96.0
Michigan	79.7	98.2	72.9	87.4	55.3	63.8	99.0	83.7
Minnesota	77.6	97.7	95.3	97.7	83.4	68.4	100.0	83.6
Missouri	90.9	100.0	82.3	90.0	74.1	81.1	97.6	84.6
Montana	88.0	99.5	72.5	82.0	65.0	73.0	98.1	82.6
Nebraska	88.1	99.1	73.2	87.0	67.2	70.3	98.8	80.5
New Hampshire	75.1	98.1	84.4	91.1	72.0	61.9	98.2	87.1
New Mexico	83.5	100.0	94.0	96.1	76.4	73.0	100.0	95.5
New York	84.8	98.5	89.9	95.8	82.7	79.9	100.0	90.9
North Dakota	91.5	98.7	70.1	81.7	78.7	81.4	99.3	87.5
Tennessee	91.3	98.6	84.0	94.3	72.5	72.8	98.8	83.3
Utah	79.7	98.6	79.6	92.3	92.0	73.9	99.3	94.4
Vermont	81.9	94.9	90.7	94.6	63.4	55.0	96.5	82.6
Virginia	89.1	100.0	81.3	85.5	71.5	76.1	100.0	91.1
Wisconsin	87.6	97.8	89.5	92.6	80.0	70.2	99.1	81.9
<b>Unweighted Data</b>								
Arkansas	93.8	99.5	87.1	91.5	77.3	75.5	99.5	89.2
California	75.1	95.8	84.1	92.3	65.2	66.3	98.0	80.4
North Carolina	77.7	100.0	80.7	89.4	69.6	67.9	100.0	87.2
Oklahoma	90.9	98.6	61.0	74.0	59.7	67.5	96.1	87.0
Pennsylvania <sup>§</sup>	84.4	99.2	85.7	94.3	67.9	71.9	99.6	83.7
Rhode Island	81.7	98.3	86.9	91.8	70.5	75.0	96.7	93.4
South Carolina	88.3	97.7	88.2	94.7	59.2	68.1	96.2	81.7
Texas <sup>§</sup>	87.1	98.6	83.8	88.9	76.1	80.0	99.5	87.4
<b>State Median</b>	<b>84.8</b>	<b>98.6</b>	<b>84.5</b>	<b>92.3</b>	<b>73.4</b>	<b>71.9</b>	<b>99.1</b>	<b>87.0</b>

(continued)

TABLE 4c. Percentage of Schools That Tried to Increase Student Knowledge, by Specific Health-Related Topic,\*  
Selected U.S. Sites: School Health Profiles, Teachers' Surveys, 2002 (continued)

### LOCAL SURVEYS

Site	Personal hygiene	Physical activity and fitness	Pregnancy prevention	STD** prevention	Suicide prevention	Sun safety	Tobacco-use prevention	Violence prevention
<b>Weighted Data</b>								
Chicago	95.4	98.7	78.2	81.0	52.8	51.5	95.9	98.0
Dallas	83.5	100.0	93.3	96.7	96.6	76.6	100.0	100.0
Fort Lauderdale	96.6	100.0	88.0	100.0	81.8	81.3	100.0	97.0
Houston	93.4	100.0	88.6	97.7	84.4	84.0	100.0	97.8
Los Angeles	97.9	100.0	93.7	98.0	77.6	88.6	100.0	88.1
Miami	88.2	94.3	91.9	98.0	84.4	80.7	100.0	82.7
Orange County	80.0	100.0	100.0	100.0	89.8	83.6	100.0	85.0
Palm Beach	80.1	100.0	100.0	96.3	84.8	88.7	100.0	81.4
San Diego	36.6	76.5	95.2	97.6	51.2	22.0	100.0	83.3
San Francisco	82.0	100.0	95.7	95.0	90.7	71.4	100.0	95.7
<b>Unweighted Data</b>								
District of Columbia	96.4	100.0	92.9	89.3	59.3	52.0	100.0	92.9
Philadelphia	97.2	98.6	83.6	91.8	62.0	65.3	97.2	89.0
San Bernardino	83.3	100.0	100.0	100.0	66.7	80.0	100.0	100.0
<b>Local Median</b>	<b>88.2</b>	<b>100.0</b>	<b>93.3</b>	<b>97.6</b>	<b>81.8</b>	<b>80.0</b>	<b>100.0</b>	<b>92.9</b>

\* In a required health education course.

\*\* STD = sexually transmitted disease.

§ Survey did not include students from one of the state's large school districts.

TABLE 5. Percentage of Schools That Tried to Improve Specific Student Skills,\* Selected U.S. Sites: School Health Profiles, Teachers' Surveys, 2002

**STATE SURVEYS**

Site	Accessing health information	Advocating for health	Analysis of media messages	Communication	Decision making	Goal setting	Conflict resolution	Resisting peer pressure	Stress management
<b>Weighted Data</b>									
Alabama	90.0	88.2	80.6	95.2	97.9	97.2	94.2	98.5	93.5
Alaska	76.3	74.8	66.6	84.0	89.7	83.9	82.6	90.0	70.5
Arizona	77.2	79.9	71.4	89.2	94.1	91.4	86.7	93.0	79.1
Connecticut	84.8	76.9	90.2	94.8	97.0	89.7	88.0	97.7	89.5
Delaware	95.6	90.4	93.5	93.5	100.0	100.0	95.2	100.0	97.6
Georgia	86.7	85.2	85.9	96.9	98.0	98.1	92.6	99.2	94.4
Hawaii	93.4	91.6	98.5	98.1	100.0	93.2	93.1	100.0	89.7
Idaho	87.0	80.2	82.3	92.2	97.7	94.2	85.6	96.6	92.9
Illinois**	83.0	79.3	83.4	91.1	97.6	93.4	85.8	96.9	92.1
Iowa	80.7	81.8	79.5	89.6	96.5	89.8	78.3	95.5	85.8
Kentucky	90.2	87.8	87.9	95.5	99.2	97.1	94.9	98.3	93.5
Maine	92.1	79.4	91.1	95.1	98.9	96.1	88.2	96.3	94.7
Massachusetts	89.6	80.8	91.9	96.2	99.1	93.1	93.2	98.7	88.7
Michigan	76.7	75.9	83.2	94.1	96.0	90.3	83.2	95.7	83.5
Minnesota	87.4	81.7	89.6	94.0	99.7	92.6	82.1	96.2	89.7
Missouri	89.9	84.8	84.6	92.5	97.0	94.3	89.7	97.7	92.1
Montana	87.0	82.7	82.1	91.4	95.6	91.1	89.5	95.6	86.1
Nebraska	76.8	77.4	74.4	86.0	94.8	91.6	84.9	96.0	89.5
New Hampshire	82.2	77.5	91.4	90.3	95.9	89.6	79.7	94.4	93.4
New Mexico	86.1	85.2	84.9	94.3	98.0	93.9	89.0	99.1	87.1
New York	92.4	89.5	94.2	96.2	98.6	96.2	91.0	98.0	95.3
North Dakota	82.9	79.4	87.4	95.0	96.4	92.8	87.8	97.9	93.1
Tennessee	82.9	81.6	70.8	91.9	97.4	92.5	86.2	95.0	84.8
Utah	88.5	86.5	91.4	96.1	99.3	98.3	95.8	99.3	98.6
Vermont	84.3	81.8	90.8	96.2	100.0	93.2	81.1	97.9	86.8
Virginia	89.0	87.9	87.6	94.2	97.8	95.2	88.9	97.7	94.3
Wisconsin	87.0	81.0	89.1	96.0	98.1	90.7	87.1	97.4	91.2
<b>Unweighted Data</b>									
Arkansas	82.1	79.7	70.6	90.5	97.2	92.5	88.2	97.2	88.2
California	73.3	75.4	81.0	87.3	93.2	83.4	81.0	94.7	78.3
North Carolina	86.6	80.0	76.2	93.0	98.4	94.1	95.2	96.3	94.1
Oklahoma	77.6	75.6	61.0	88.3	93.6	93.5	89.2	94.7	87.2
Pennsylvania**	87.7	82.6	87.2	93.4	98.4	95.1	85.7	98.0	88.1
Rhode Island	96.7	88.5	91.8	100.0	100.0	95.1	95.1	98.4	88.5
South Carolina	85.4	83.0	74.9	89.8	97.3	95.2	89.9	96.3	86.1
Texas**	84.6	78.1	72.9	91.2	96.7	94.4	89.3	96.3	88.8
<b>State Median</b>	<b>86.6</b>	<b>81.6</b>	<b>84.9</b>	<b>93.5</b>	<b>97.7</b>	<b>93.4</b>	<b>88.2</b>	<b>97.2</b>	<b>89.5</b>

(continued)

TABLE 5. Percentage of Schools That Tried to Improve Specific Student Skills,\* Selected U.S. Sites: School Health Profiles, Teachers' Surveys, 2002 (continued)

**LOCAL SURVEYS**

Site	Accessing health information	Advocating for health	Analysis of media messages	Communication	Decision making	Goal setting	Conflict resolution	Resisting peer pressure	Stress management
<b>Weighted Data</b>									
Chicago	78.4	78.0	66.7	87.5	96.7	94.1	94.6	97.3	78.3
Dallas	96.8	93.3	73.2	96.7	100.0	93.4	93.4	96.7	96.6
Fort Lauderdale	96.9	97.0	91.0	97.0	97.0	97.0	90.8	100.0	97.0
Houston	88.5	93.2	77.2	97.7	100.0	100.0	97.8	100.0	97.8
Los Angeles	94.8	88.1	92.5	92.5	99.0	95.7	96.9	97.9	91.4
Miami	88.6	80.6	84.7	94.3	98.1	96.1	92.4	94.4	84.0
Orange County	87.4	91.2	84.8	100.0	100.0	96.2	96.8	100.0	100.0
Palm Beach	92.7	96.3	77.4	96.3	96.3	100.0	92.7	100.0	96.3
San Diego	66.7	64.3	95.2	97.6	100.0	97.6	97.6	97.6	97.6
San Francisco	86.3	82.0	82.0	95.7	95.7	100.0	90.7	100.0	95.7
<b>Unweighted Data</b>									
District of Columbia	82.1	82.1	67.9	77.8	89.3	92.9	89.3	85.7	75.0
Philadelphia	84.9	81.9	85.9	91.7	98.6	91.7	91.7	100.0	77.8
San Bernardino	83.3	100.0	83.3	100.0	100.0	100.0	100.0	100.0	100.0
<b>Local Median</b>	<b>87.4</b>	<b>88.1</b>	<b>83.3</b>	<b>96.3</b>	<b>98.6</b>	<b>96.2</b>	<b>93.4</b>	<b>100.0</b>	<b>96.3</b>

\* In a required health education course.

\*\* Survey did not include students from one of the state's large school districts.

TABLE 6a. Percentage of Schools That Taught Specific Topics Related to Tobacco-Use Prevention,\* Selected U.S. Sites: School Health Profiles, Teachers' Surveys, 2002

**STATE SURVEYS**

Site	Addictive effects of nicotine	Benefits of not smoking cigarettes	Benefits of not using smokeless tobacco	Health consequences of cigarette smoking	Health consequences of smokeless tobacco use	Health effects of ETS**	Number of illnesses and deaths related to tobacco use	Risks of cigar or pipe smoking
<b>Weighted Data</b>								
Alabama	97.8	97.7	96.6	97.7	94.8	96.0	96.6	93.0
Alaska	90.3	90.0	88.3	93.3	87.7	85.9	84.8	76.7
Arizona	94.1	96.7	93.5	96.2	94.1	90.4	90.9	87.5
Connecticut	96.5	96.7	86.0	97.3	91.5	95.1	94.6	84.1
Delaware	100.0	100.0	95.4	100.0	100.0	100.0	100.0	90.0
Georgia	99.5	100.0	99.5	100.0	99.5	99.1	97.6	96.1
Hawaii	98.1	98.5	89.9	100.0	94.7	98.1	92.6	87.9
Idaho	98.2	98.8	98.1	98.8	99.4	95.1	93.9	88.2
Illinois <sup>§</sup>	97.4	98.0	94.2	98.5	95.6	97.3	94.7	89.0
Iowa	95.2	95.8	90.1	96.8	93.9	94.7	91.4	83.6
Kentucky	97.0	97.0	93.8	98.9	97.7	97.2	95.4	89.2
Maine	97.4	97.8	90.9	100.0	95.7	95.8	92.0	80.9
Massachusetts	97.4	97.6	91.2	98.4	95.2	97.0	95.7	88.5
Michigan	96.3	97.4	90.9	98.2	94.9	97.0	93.1	86.9
Minnesota	99.3	99.6	95.3	99.6	97.3	98.1	96.6	82.8
Missouri	95.3	95.5	91.1	96.9	94.6	93.8	93.5	88.9
Montana	94.4	95.7	93.0	96.8	94.9	94.8	94.4	83.1
Nebraska	95.2	95.7	92.1	96.9	93.7	93.3	91.7	76.8
New Hampshire	97.3	94.0	90.0	97.3	94.2	93.2	92.5	81.9
New Mexico	99.0	99.0	90.9	99.0	94.7	95.9	96.8	91.1
New York	99.2	98.9	97.6	99.3	98.6	98.2	97.2	95.3
North Dakota	96.3	97.2	93.5	98.6	94.8	95.6	94.1	84.7
Tennessee	95.1	97.1	92.5	97.5	95.6	96.2	93.3	87.3
Utah	99.3	98.7	95.9	99.3	98.7	98.2	96.9	90.9
Vermont	96.5	92.9	92.9	95.1	92.6	93.0	86.5	78.0
Virginia	97.5	99.0	91.4	99.0	94.5	96.1	93.4	89.5
Wisconsin	97.8	97.8	94.1	98.5	96.1	95.4	94.8	86.3
<b>Unweighted Data</b>								
Arkansas	98.6	98.1	95.7	99.0	98.1	98.6	96.2	89.9
California	95.5	95.5	87.1	96.5	91.0	91.0	93.1	84.7
North Carolina	97.3	99.4	96.7	99.5	97.8	98.9	94.5	90.1
Oklahoma	94.6	96.1	93.2	93.4	93.4	92.2	94.6	87.0
Pennsylvania <sup>§</sup>	98.3	98.8	95.5	98.8	97.1	96.7	96.7	92.1
Rhode Island	95.0	96.7	90.0	96.7	89.8	95.0	95.0	90.0
South Carolina	92.9	92.9	89.8	95.1	92.9	91.3	86.3	84.2
Texas <sup>§</sup>	99.1	99.1	96.2	98.6	97.2	97.6	96.7	92.0
<b>State Median</b>	<b>97.3</b>	<b>97.6</b>	<b>93.0</b>	<b>98.5</b>	<b>94.9</b>	<b>95.9</b>	<b>94.5</b>	<b>87.9</b>

(continued)



TABLE 6a. Percentage of Schools That Taught Specific Topics Related to Tobacco-Use Prevention,\* Selected U.S. Sites:  
School Health Profiles, Teachers' Surveys, 2002 (continued)

### LOCAL SURVEYS

Site	Addictive effects of nicotine	Benefits of not smoking cigarettes	Benefits of not using smokeless tobacco	Health consequences of cigarette smoking	Health consequences of smokeless tobacco use	Health effects of ETS**	Number of illnesses and deaths related to tobacco use	Risks of cigar or pipe smoking
<b>Weighted Data</b>								
Chicago	93.9	91.3	76.9	93.9	77.9	91.3	92.6	83.8
Dallas	96.5	96.5	93.0	96.6	93.0	100.0	100.0	96.3
Fort Lauderdale	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Houston	100.0	97.7	100.0	100.0	100.0	100.0	100.0	100.0
Los Angeles	98.8	100.0	95.7	100.0	97.9	98.0	98.8	94.9
Miami	97.9	100.0	90.0	100.0	92.2	98.1	96.3	94.2
Orange County	96.7	92.8	89.4	100.0	89.4	96.7	96.7	92.8
Palm Beach	100.0	100.0	100.0	100.0	96.3	100.0	100.0	100.0
San Diego	97.5	97.5	95.0	97.5	95.0	97.5	95.0	97.5
San Francisco	94.8	90.3	63.0	94.8	81.2	81.2	80.5	76.6
<b>Unweighted Data</b>								
District of Columbia	84.0	91.7	64.0	88.0	61.5	80.0	80.0	75.0
Philadelphia	97.2	98.6	81.7	98.6	87.1	97.2	94.4	91.4
San Bernardino	100.0	100.0	100.0	100.0	100.0	100.0	83.3	83.3
<b>Local Median</b>	<b>97.5</b>	<b>97.7</b>	<b>93.0</b>	<b>100.0</b>	<b>93.0</b>	<b>98.0</b>	<b>96.3</b>	<b>94.2</b>

\* In a required health education course.

\*\* ETS = environmental tobacco smoke.

§ Survey did not include students from one of the state's large school districts.

TABLE 6b. Percentage of Schools That Taught Specific Topics Related to Tobacco-Use Prevention,\* Selected U.S. Sites: School Health Profiles, Teachers' Surveys, 2002

**STATE SURVEYS**

Site	Influence of families**	Influence of media**	Social or cultural influences**	How students can influence others to prevent tobacco use	How students can influence others to quit using tobacco	How many young people use tobacco
<b>Weighted Data</b>						
Alabama	94.9	96.6	94.8	93.9	92.8	92.8
Alaska	81.2	84.2	82.8	80.1	76.2	80.9
Arizona	88.0	91.0	88.2	86.8	85.8	86.5
Connecticut	89.6	94.7	88.6	85.8	78.7	90.6
Delaware	93.2	100.0	92.5	97.5	95.0	100.0
Georgia	97.7	98.3	94.2	92.3	92.6	96.2
Hawaii	95.2	98.1	82.7	78.9	77.2	91.9
Idaho	90.2	97.6	93.1	88.6	85.3	94.8
Illinois <sup>§</sup>	91.8	95.4	90.7	86.0	83.1	92.7
Iowa	88.6	93.6	88.9	79.1	79.7	90.4
Kentucky	93.8	95.8	94.8	93.0	92.1	93.9
Maine	87.7	95.7	85.6	82.4	80.5	88.1
Massachusetts	92.2	95.5	90.7	90.3	88.7	93.5
Michigan	91.4	96.6	91.1	88.3	82.4	90.9
Minnesota	91.7	98.0	90.0	90.0	85.7	95.6
Missouri	91.2	92.5	92.2	88.1	87.8	89.8
Montana	87.7	90.7	88.4	84.2	82.7	93.0
Nebraska	89.2	90.2	87.1	85.3	84.2	88.5
New Hampshire	91.0	91.8	89.8	77.6	78.2	90.2
New Mexico	87.3	93.8	86.1	86.9	83.7	91.6
New York	97.6	99.3	95.9	90.3	88.7	96.2
North Dakota	89.4	95.6	91.2	87.8	82.9	93.4
Tennessee	91.4	93.0	90.1	89.4	88.2	93.0
Utah	95.6	97.4	93.5	92.3	90.0	95.7
Vermont	81.8	94.0	86.3	82.4	75.6	87.5
Virginia	93.6	96.4	91.4	91.6	91.1	92.0
Wisconsin	89.8	96.5	88.8	85.5	82.6	93.0
<b>Unweighted Data</b>						
Arkansas	92.3	93.8	88.0	92.8	90.9	94.8
California	87.2	92.5	85.7	78.7	74.3	88.1
North Carolina	94.0	96.2	92.8	90.4	90.0	95.0
Oklahoma	90.9	92.2	86.7	87.0	82.7	92.0
Pennsylvania <sup>§</sup>	92.6	95.9	91.7	88.3	89.6	94.7
Rhode Island	95.0	93.3	86.7	90.0	85.0	93.3
South Carolina	84.2	87.0	83.6	81.5	79.3	83.8
Texas <sup>§</sup>	95.3	96.7	92.4	89.0	89.0	94.8
<b>State Median</b>	<b>91.4</b>	<b>95.5</b>	<b>90.0</b>	<b>88.1</b>	<b>85.0</b>	<b>92.8</b>

(continued)

TABLE 6b. Percentage of Schools That Taught Specific Topics Related to Tobacco-Use Prevention,\* Selected U.S. Sites: School Health Profiles, Teachers' Surveys, 2002 (continued)

### LOCAL SURVEYS

Site	Influence of families**	Influence of media**	Social or cultural influences**	How students can influence others to prevent tobacco use	How students can influence others to quit using tobacco	How many young people use tobacco
<b>Weighted Data</b>						
Chicago	86.6	87.1	86.6	84.8	85.4	87.9
Dallas	89.6	93.0	86.2	89.6	93.0	93.0
Fort Lauderdale	93.8	96.7	100.0	100.0	100.0	100.0
Houston	97.8	95.6	97.7	100.0	97.7	97.8
Los Angeles	95.9	100.0	95.9	89.5	86.4	93.5
Miami	90.4	94.3	90.2	90.2	84.2	96.0
Orange County	90.9	96.7	87.6	92.8	77.8	92.8
Palm Beach	96.3	92.7	81.4	89.0	92.7	96.3
San Diego	90.0	97.5	90.0	92.5	87.5	95.0
San Francisco	94.8	90.3	90.3	80.5	75.3	80.5
<b>Unweighted Data</b>						
District of Columbia	76.0	83.3	75.0	76.0	76.0	84.0
Philadelphia	93.0	95.7	91.5	87.3	87.1	93.0
San Bernardino	100.0	100.0	83.3	100.0	100.0	83.3
<b>Local Median</b>	<b>93.0</b>	<b>95.6</b>	<b>90.0</b>	<b>89.6</b>	<b>87.1</b>	<b>93.0</b>

\* In a required health education course.

\*\* On tobacco-use.

§ Survey did not include students from one of the state's large school districts.

TABLE 6c. Percentage of Schools That Taught Specific Topics Related to Tobacco-Use Prevention,\* Selected U.S. Sites: School Health Profiles, Teachers' Surveys, 2002

**STATE SURVEYS**

Site	How to find information on tobacco-use cessation	Making a personal commitment not to use tobacco	How to say no to tobacco use
<b>Weighted Data</b>			
Alabama	77.2	80.3	98.4
Alaska	64.4	72.7	90.4
Arizona	77.8	78.2	92.9
Connecticut	68.9	68.0	93.8
Delaware	85.0	77.1	100.0
Georgia	84.1	80.0	98.8
Hawaii	74.2	65.4	95.0
Idaho	72.8	69.7	97.0
Illinois**	68.8	66.3	95.3
Iowa	62.1	65.2	91.4
Kentucky	86.5	83.0	97.0
Maine	74.3	65.9	89.5
Massachusetts	79.4	79.8	95.7
Michigan	68.8	68.2	93.4
Minnesota	70.7	73.3	95.4
Missouri	67.3	72.6	94.5
Montana	74.8	74.4	92.8
Nebraska	69.3	66.8	92.0
New Hampshire	65.4	65.2	87.2
New Mexico	72.6	72.8	93.0
New York	81.4	80.8	95.9
North Dakota	65.1	70.9	95.7
Tennessee	70.5	77.7	95.0
Utah	73.2	78.9	98.1
Vermont	66.0	60.9	93.1
Virginia	74.7	80.7	96.1
Wisconsin	71.7	75.1	93.6
<b>Unweighted Data</b>			
Arkansas	69.7	75.7	97.6
California	70.6	67.2	90.6
North Carolina	76.1	76.1	98.3
Oklahoma	65.3	78.7	93.3
Pennsylvania**	82.6	73.3	98.3
Rhode Island	83.1	76.3	95.0
South Carolina	68.9	72.8	88.6
Texas**	78.0	73.1	97.6
<b>State Median</b>	<b>72.6</b>	<b>73.3</b>	<b>95.0</b>

(continued)

TABLE 6c. Percentage of Schools That Taught Specific Topics Related to Tobacco-Use Prevention,\* Selected U.S. Sites: School Health Profiles, Teachers' Surveys, 2002 (continued)

### LOCAL SURVEYS

Site	How to find information on tobacco-use cessation	Making a personal commitment not to use tobacco	How to say no to tobacco use
<b>Weighted Data</b>			
Chicago	68.5	81.2	91.3
Dallas	75.8	79.4	96.6
Fort Lauderdale	87.8	100.0	100.0
Houston	84.5	89.0	100.0
Los Angeles	91.5	84.4	97.9
Miami	82.6	84.4	94.2
Orange County	70.0	76.4	93.4
Palm Beach	92.7	81.1	100.0
San Diego	90.0	75.0	97.5
San Francisco	80.5	71.4	94.8
<b>Unweighted Data</b>			
District of Columbia	68.0	80.0	84.0
Philadelphia	78.9	78.9	97.2
San Bernardino	83.3	66.7	100.0
<b>Local Median</b>	<b>82.6</b>	<b>80.0</b>	<b>97.2</b>

\* In a required health education course.

\*\* Survey did not include students from one of the state's large school districts.