TABLE 7a. Percentage of Schools That Taught Specific Topics Related to HIV\*/AIDS\*\* Prevention,§ Selected U.S. Sites: School Health Profiles, Teachers' Surveys, 2002

	Abstinence to avoid HIV	How HIV is	How HIV affects	How to correctly use	Condom	Number of young people
Site	infection	transmitted	the body	a condom	efficacy	who get HIV
<b>Weighted Data</b>						
Alabama	98.4	97.8	97.2	39.2	75.8	92.3
Alaska	70.5	75.7	74.3	28.7	48.2	66.5
Arizona	83.2	83.3	81.7	29.3	48.6	74.8
Connecticut	99.4	99.4	98.6	50.6	79.9	91.3
Delaware	95.6	97.6	95.2	49.1	70.0	92.8
Georgia	96.7	96.3	95.6	23.6	74.5	91.7
Hawaii	98.5	98.5	96.5	58.4	83.2	84.7
Idaho	94.7	93.5	92.3	20.2	59.8	87.8
Illinois <sup>§§</sup>	95.0	95.5	94.7	35.8	71.2	83.7
lowa	97.0	97.0	95.7	44.2	76.2	86.2
Kentucky	97.0	96.3	94.8	41.7	70.4	87.8
Maine	97.4	97.8	95.7	65.5	88.1	88.3
Massachusetts	94.2	94.4	93.8	51.2	75.3	86.4
Michigan	90.5	91.3	89.5	28.9	60.3	79.3
Minnesota	99.1	99.1	98.1	46.5	78.5	89.6
Missouri	93.5	94.2	92.3	27.8	68.4	80.0
Montana	85.9	87.4	86.9	32.7	64.6	82.6
Nebraska	86.5	87.7	86.4	21.8	53.3	76.9
New Hampshire	95.9	95.8	95.0	52.2	73.3	81.2
New Mexico	97.0	97.0	97.2	53.2	79.4	92.0
New York	98.6	99.3	99.3	61.3	84.3	94.9
North Dakota	88.3	87.6	86.8	22.5	51.5	76.2
Tennessee	92.0	91.2	92.6	37.6	68.5	82.8
Utah	97.3	97.3	96.1	7.8	40.6	89.6
Vermont	97.3	94.5	93.3	56.8	86.3	80.1
Virginia	88.8	90.1	88.7	NA <sup>+</sup>	NA <sup>+</sup>	79.3
Wisconsin	94.9	95.9	94.8	46.9	79.1	86.5
Unweighted Data						
Arkansas	92.7	91.3	89.4	25.6	63.9	80.6
California	95.7	96.1	95.7	54.9	79.2	88.0
North Carolina	97.1	91.4	91.3	18.3	60.8	80.6
Oklahoma	86.8	88.3	88.2	25.7	50.0	81.6
Pennsylvania <sup>§§</sup>	99.2	99.2	99.2	47.3	78.6	93.5
Rhode Island	95.0	95.1	95.1	58.3	71.7	85.2
South Carolina	94.6	93.0	92.5	44.1	73.0	82.9
Texas <sup>§§</sup>	94.2	94.2	92.3	22.2	59.7	85.3
State Median	95.0	95.1	94.7	40.4	71.4	<b>85.2</b>

TABLE 7a. Percentage of Schools That Taught Specific Topics Related to HIV\*/AIDS\*\* Prevention,§ Selected U.S. Sites: School Health Profiles, Teachers' Surveys, 2002 (continued)

Site	Abstinence to avoid HIV infection	How HIV is transmitted	How HIV affects the body	How to correctly use a condom	Condom efficacy	Number of young people who get HIV
<b>Weighted Data</b>						
Chicago	85.0	84.7	85.1	39.0	50.0	76.8
Dallas	100.0	100.0	100.0	40.0	86.4	93.3
Fort Lauderdale	100.0	97.0	97.0	52.2	85.0	94.0
Houston	95.5	95.5	95.5	46.3	70.9	91.1
Los Angeles	98.0	99.0	97.9	69.5	92.5	90.4
Miami	100.0	100.0	100.0	86.1	89.9	100.0
Orange County	100.0	100.0	96.8	42.9	89.8	96.8
Palm Beach	100.0	100.0	100.0	NA <sup>+</sup>	92.1	92.4
San Diego	100.0	100.0	100.0	100.0	95.2	90.5
San Francisco	90.1	90.1	90.1	85.7	85.7	81.4
<b>Unweighted Data</b>						
District of Columbia	100.0	100.0	100.0	89.3	96.4	92.9
Philadelphia	97.2	97.2	97.2	59.7	77.8	92.9
San Bernardino	100.0	100.0	100.0	66.7	100.0	83.3
Local Median	100.0	100.0	97.9	63.2	89.8	92.4

<sup>\*</sup> HIV = human immunodeficiency virus.

<sup>\*\*</sup> AIDS = acquired immunodeficiency syndrome.

 $<sup>\</sup>S$  In a required health education course.

 $<sup>\</sup>S\S$  Survey did not include students from one of the state's large school districts.

<sup>+</sup> NA = not available.

TABLE 7b. Percentage of Schools That Taught Specific Topics Related to HIV\*/AIDS\*\* Prevention,§ Selected U.S. Sites: School Health Profiles, Teachers' Surveys, 2002

Site	Influence of alcohol or other drugs <sup>§§</sup>	Social or cultural influences <sup>§§</sup>	How to find valid information on HIV	Compassion for persons with HIV or AIDS
Weighted Data	other drugs	imachees	OHTHV	1117 01 71123
Alabama	95.0	91.2	87.7	85.6
Alaska	69.2	62.5	60.5	59.6
Arizona	79.0	72.0	64.6	69.5
Connecticut	96.7	86.7	84.5	91.7
Delaware	97.9	91.1	88.0	86.3
Georgia	95.1	90.3	81.6	84.8
Hawaii	91.6	78.4	84.1	83.3
daho	90.0	83.8	75.1	84.5
Illinois <sup>+</sup>	89.1	79.3	74.8	77.3
		79.3 83.4	74.8 76.1	
lowa Kontucky	89.9 91.2	83.4 87.2	76.1 82.2	80.4 82.8
Kentucky				
Maine	94.0	82.9	85.2	83.8
Massachusetts	92.1	82.0	83.1	86.4
Michigan	88.1	77.5	77.7	78.4
Minnesota	96.2	84.9	78.2	81.6
Missouri	88.0	83.4	77.6	76.2
Montana	85.9	76.9	74.9	78.3
Nebraska	81.3	72.4	63.7	71.7
New Hampshire	89.0	75.2	75.8	82.8
New Mexico	92.2	87.9	85.7	87.4
New York	98.2	89.0	92.0	94.5
North Dakota	87.3	75.5	66.2	75.5
Tennessee	84.4	83.2	76.0	77.1
Utah	93.1	89.0	76.8	89.7
Vermont	93.1	76.6	72.1	82.6
Virginia	83.2	74.6	70.1	73.8
Wisconsin	93.8	83.6	77.6	82.2
Unweighted Data				
Arkansas	90.6	83.7	72.2	71.5
California	89.9	83.1	84.7	86.0
North Carolina	87.8	80.0	69.0	77.5
Oklahoma	82.9	77.3	68.0	75.3
Pennsylvania <sup>+</sup>	98.4	87.8	87.3	88.9
Rhode Island	89.8	78.7	85.2	88.5
South Carolina	87.5	81.7	73.0	76.6
Texas <sup>+</sup>	85.4	81.7	73.6	75.4
State Median	89.9	82.9	76.8	82.2

TABLE 7b. Percentage of Schools That Taught Specific Topics Related to HIV\*/AIDS\*\* Prevention,§ Selected U.S. Sites: School Health Profiles, Teachers' Surveys, 2002 (continued)

Site	Influence of alcohol or other drugs <sup>§§</sup>	Social or cultural influences <sup>§§</sup>	How to find valid information on HIV	Compassion for persons with HIV or AIDS
Weighted Data				
Chicago	77.3	71.6	66.6	73.3
Dallas	93.3	90.0	96.6	90.0
Fort Lauderdale	97.0	91.0	94.0	97.0
Houston	95.5	95.5	93.1	95.5
Los Angeles	95.0	92.8	90.5	93.7
Miami	97.9	94.4	98.1	94.4
Orange County	100.0	100.0	89.2	96.8
Palm Beach	92.4	85.3	89.0	96.1
San Diego	95.2	90.5	100.0	92.9
San Francisco	90.1	85.1	85.7	81.4
Unweighted Data				
District of Columbia	100.0	92.9	92.9	89.3
Philadelphia	91.7	86.1	81.9	86.1
San Bernardino	100.0	100.0	83.3	100.0
Local Median	95.2	91.0	90.5	93.7

<sup>\*</sup> HIV = human immunodeficiency virus.

<sup>\*\*</sup> AIDS = acquired immunodeficiency syndrome.

<sup>§</sup> In a required health education course.

 $<sup>\</sup>S\S$  On HIV-related risk behaviors.

<sup>+</sup> Survey did not include students from one of the state's large school districts.

TABLE 8a. Percentage of Schools That Taught Specific Topics Related to Nutrition and Dietary Intake,\* Selected U.S. Sites: School Health Profiles, Teachers' Surveys, 2002

Site	Benefits of healthy eating	Using food labels	Food guide pyramid	Choosing a variety of grains**	Choosing a variety of fruits and vegetables	Choosing a low-fat diet <sup>§</sup>	Using less salt	Moderating intake of sugars	Eating more calcium-rich foods
Weighted D	ata								
Alabama	98.3	92.8	91.8	90.1	95.0	93.5	87.4	91.1	85.8
Alaska	90.4	78.1	83.5	79.9	83.5	77.9	68.5	81.9	74.5
Arizona	93.8	84.8	91.9	83.5	85.9	83.1	76.6	83.2	78.8
Connecticut	94.1	78.3	80.7	78.2	83.7	83.3	63.3	79.5	66.2
Delaware	93.2	88.1	84.3	81.5	83.6	88.0	72.3	86.0	75.9
Georgia	97.3	88.9	91.4	86.8	90.1	93.2	86.9	90.5	88.3
Hawaii	96.4	84.2	91.3	80.1	89.0	84.2	77.8	91.1	73.3
Idaho	98.0	87.6	92.5	86.0	88.4	88.8	81.6	89.4	86.1
Illinois <sup>§§</sup>	95.4	85.3	89.1	81.5	88.6	89.0	73.9	85.3	78.3
lowa	96.6	87.6	92.7	86.1	89.9	90.4	75.8	87.7	84.1
Kentucky	99.5	96.5	97.4	92.5	96.3	97.4	88.1	95.3	86.8
Maine	96.8	88.9	92.1	87.6	92.5	92.9	81.3	90.7	82.2
Massachusetts	s 94.5	84.0	87.2	84.5	87.8	86.2	74.1	83.9	81.8
Michigan	93.1	83.8	88.0	84.7	87.4	85.3	74.5	85.3	77.4
Minnesota	95.0	80.0	82.4	78.4	82.8	84.4	68.7	80.7	73.1
Missouri	98.2	95.8	97.6	93.0	95.4	94.3	83.3	93.9	88.3
Montana	96.5	85.8	91.6	86.3	90.8	88.6	75.6	92.4	83.5
Nebraska	93.4	78.5	83.7	80.8	81.6	83.6	71.7	79.6	77.1
New Hampshire	e 94.2	86.4	89.2	84.8	87.1	88.4	80.3	88.2	80.7
New Mexico	96.9	90.8	91.4	87.9	92.0	89.1	80.5	92.3	85.4
New York	96.4	86.3	85.4	81.8	87.3	92.5	78.1	85.8	79.2
North Dakota	96.6	86.2	90.8	85.9	88.6	89.7	76.9	87.5	85.9
Tennessee	94.3	87.9	93.3	85.0	89.9	88.2	82.8	87.2	78.8
Utah	98.0	93.1	96.3	92.8	94.5	96.9	83.9	95.4	85.9
Vermont	93.2	84.8	90.8	81.0	85.7	83.9	73.6	81.6	77.8
Virginia	95.7	89.5	90.7	87.6	90.3	90.9	80.7	90.5	85.2
Wisconsin	94.7	86.8	90.0	85.7	88.3	86.4	78.6	87.5	78.2
Unweighted	d Data								
Arkansas	99.0	88.3	90.8	84.1	91.7	92.2	85.4	88.3	82.5
California	88.7	81.9	84.2	77.3	84.7	81.4	66.7	79.7	72.9
North Carolina	a 97.8	94.1	95.7	91.3	92.9	95.1	87.5	92.9	84.7
Oklahoma	90.5	77.3	88.2	81.3	88.0	87.8	73.7	81.6	76.0
Pennsylvania <sup>§§</sup>	93.6	85.8	85.8	78.0	84.2	87.3	78.4	83.8	78.7
Rhode Island	91.8	80.3	83.3	81.7	88.3	84.7	67.2	81.7	75.4
South Carolina	a 95.6	91.6	90.6	86.2	90.1	89.5	83.9	86.7	82.8
Texas <sup>§§</sup>	95.2	87.6	91.9	84.5	89.3	91.8	84.5	90.9	81.6
<b>State Median</b>	95.4	86.4	90.8	84.7	88.6	88.6	78.1	87.5	80.7

TABLE 8a. Percentage of Schools That Taught Specific Topics Related to Nutrition and Dietary Intake,\* Selected U.S. Sites: School Health Profiles, Teachers' Surveys, 2002 (continued)

Site	Benefits of healthy eating	Using food labels	Food guide pyramid	Choosing a variety of grains**	Choosing a variety of fruits and vegetables	Choosing a low-fat diet <sup>§</sup>	Using less salt	Moderating intake of sugars	Eating more calcium-rich foods	
Weighted Da	Weighted Data									
Chicago	96.6	84.0	93.8	85.6	93.7	82.0	78.1	84.9	81.3	
Dallas	100.0	93.2	100.0	86.3	96.6	96.6	89.7	89.7	82.8	
Fort Lauderdale	97.0	97.0	100.0	97.0	97.0	97.0	97.0	97.0	97.0	
Houston	100.0	95.4	95.5	97.8	100.0	97.7	95.6	100.0	93.4	
Los Angeles	100.0	99.0	98.0	94.5	98.8	96.7	93.6	98.8	94.6	
Miami	98.1	96.1	96.2	86.3	92.1	91.9	80.3	86.2	90.1	
<b>Orange County</b>	100.0	86.8	96.2	96.2	96.2	90.6	81.8	96.2	74.6	
Palm Beach	100.0	89.0	96.3	100.0	100.0	100.0	92.4	100.0	92.4	
San Diego	54.1	44.4	51.4	40.0	47.5	47.5	37.5	45.0	45.0	
San Francisco	95.0	90.1	95.7	90.7	95.0	90.7	81.4	95.0	85.7	
Unweighted	Data									
District of Columbia	96.3	74.1	88.9	88.9	88.9	70.4	66.7	74.1	70.4	
Philadelphia	95.8	91.7	94.4	84.7	91.7	94.4	83.1	84.7	82.2	
San Bernardino	100.0	83.3	83.3	83.3	83.3	83.3	83.3	83.3	66.7	
<b>Local Median</b>	98.1	90.1	95.7	88.9	95.0	91.9	83.1	89.7	82.8	

<sup>\*</sup> In a required health education course.

<sup>\*\*</sup> Especially whole grains.

<sup>§</sup> Low in saturated fat and cholesterol and moderate in total fat.

<sup>§§</sup> Survey did not include students from one of the state's large school districts.

TABLE 8b. Percentage of Schools That Taught Specific Topics Related to Nutrition and Dietary Intake,\* Selected U.S. Sites: School Health Profiles, Teachers' Surveys, 2002

Site	Keeping food safe to eat	Preparing healthy meals and snacks	Aiming for a healthy weight**	Risks of unhealthy weight-control practices	Accepting body size differences	Eating disorders
Weighted Data	10 041	and shacks	g.nc	practices	uniter entrees	uiso. uc. s
Alabama	81.7	89.5	97.2	93.8	88.3	93.9
Alaska	69.4	75.5	84.1	76.7	76.2	74.9
Arizona	78.2	78.8	85.3	83.3	82.2	82.0
Connecticut	55.7	67.8	89.8	88.2	88.1	92.6
Delaware	74.6	81.5	92.7	85.7	83.3	88.4
Georgia	80.4	86.3	96.0	93.2	90.8	92.4
Hawaii	65.7	77.9	92.8	84.2	74.1	89.8
Idaho	75.3	81.9	94.0	94.7	92.8	96.4
Illinois§	72.4	83.0	89.9	89.7	86.8	90.3
lowa	75.4	80.1	90.9	91.5	84.3	88.4
Kentucky	85.5	87.7	97.5	95.9	93.8	98.4
Maine	72.7	80.2	93.1	92.9	87.9	93.9
Massachusetts	68.7	80.2	89.4	89.7	90.7	91.0
Michigan	66.1	78.5	89.6	85.7	84.2	87.6
Minnesota	64.6	71.5	90.7	89.4	86.1	92.8
Missouri	83.7	90.4	95.9	94.4	89.8	95.4
Montana	73.9	86.4	93.8	92.2	90.4	90.0
Nebraska	71.5	74.7	88.1	87.0	83.6	86.8
New Hampshire	64.6	78.4	91.0	89.6	90.9	90.7
New Mexico	75.9	83.0	91.7	91.4	86.9	89.7
New York	70.5	75.5	95.4	95.1	91.0	97.1
North Dakota	81.7	81.5	91.5	89.0	84.8	90.4
Tennessee	79.3	83.7	92.3	89.4	83.1	85.6
Utah	76.7	85.8	98.1	97.5	94.1	99.3
Vermont	67.4	83.3	89.7	93.3	87.5	85.5
Virginia	81.5	86.3	93.7	92.2	90.3	89.7
Wisconsin	69.0	79.0	90.8	87.8	87.3	88.8
Unweighted Data						
Arkansas	74.1	84.2	97.1	96.1	87.8	94.6
California	62.4	71.3	83.2	81.9	79.8	81.4
North Carolina	79.3	90.3	96.7	95.7	89.1	96.7
Oklahoma	74.7	78.7	83.8	81.3	77.0	86.3
Pennsylvania <sup>§</sup>	67.2	77.8	91.5	89.7	83.1	90.8
Rhode Island	71.7	76.7	83.3	88.3	88.1	80.3
South Carolina	77.3	85.7	90.0	90.5	86.7	88.3
Texas§	81.5	86.1	91.8	91.3	87.9	89.9
State Median	74.1	81.5	91.5	89.7	87.5	90.0

TABLE 8b. Percentage of Schools That Taught Specific Topics Related to Nutrition and Dietary Intake,\* Selected U.S. Sites: School Health Profiles, Teachers' Surveys, 2002 (continued)

	Keeping food safe	Preparing healthy meals	Aiming for a healthy	Risks of unhealthy weight-control	Accepting body size	Eating
Site	to eat	and snacks	weight**	practices	differences	disorders
Weighted Data						
Chicago	81.7	83.6	86.2	84.9	84.3	75.5
Dallas	93.0	93.0	93.0	100.0	86.2	93.1
Fort Lauderdale	94.0	97.0	97.0	97.0	97.0	97.0
Houston	88.7	97.8	100.0	100.0	97.8	97.7
Los Angeles	89.2	91.3	96.7	94.7	92.6	93.9
Miami	79.7	87.3	86.0	87.4	82.1	85.8
Orange County	62.9	75.6	94.4	93.0	88.0	93.0
Palm Beach	89.0	92.4	100.0	96.3	92.4	100.0
San Diego	37.5	40.5	51.4	51.4	54.1	57.6
San Francisco	76.4	76.4	90.1	90.1	85.7	85.1
<b>Unweighted Data</b>						
District of Columbia	59.3	63.0	92.6	70.4	70.4	70.4
Philadelphia	84.7	83.3	90.3	83.6	84.9	80.6
San Bernardino	80.0	100.0	100.0	100.0	100.0	100.0
Local Median	81.7	87.3	93.0	93.0	86.2	93.0

<sup>\*</sup> In a required health education course.

<sup>\*\*</sup> Balancing food intake and physical activity.

 $<sup>\</sup>S$  Survey did not include students from one of the state's large school districts.

TABLE 9a. Percentage of Schools That Taught Specific Topics Related to Physical Activity,\* Selected U.S. Sites: School Health Profiles, Teachers' Surveys, 2002

Site	Physical, psychological, or social benefits	Health-related fitness	Phases of a workout	How much physical activity is enough	Decreasing sedentary activities
Weighted Data					
Alabama	97.8	97.3	95.7	92.3	90.8
Alaska	87.6	83.4	74.7	69.4	76.5
Arizona	92.0	86.9	83.6	78.6	83.2
Connecticut	84.9	74.0	60.1	59.5	71.8
Delaware	100.0	89.1	84.6	84.6	88.7
Georgia	93.1	88.6	85.7	84.8	86.7
Hawaii	93.2	84.5	71.0	63.6	67.8
daho	94.9	91.7	86.4	86.7	84.2
Illinois**	92.4	87.8	76.6	73.4	83.8
lowa	94.4	89.7	80.1	79.0	79.0
Kentucky	97.7	96.5	92.4	92.7	92.2
Maine	90.4	82.3	75.9	74.3	79.1
Massachusetts	93.0	86.4	77.0	77.5	83.1
Michigan	90.1	86.7	75.8	75.4	82.2
Minnesota	88.3	81.7	71.0	73.0	75.7
Missouri	96.3	92.9	91.0	88.4	90.0
Montana	96.6	95.2	96.2	87.4	88.1
Nebraska	92.1	91.3	86.4	81.5	82.6
New Hampshire	90.8	80.0	69.8	70.1	81.1
New Mexico	94.9	91.5	84.6	84.5	91.3
New York	90.7	76.2	63.4	67.7	82.8
North Dakota	92.8	88.4	86.9	79.5	83.1
Tennessee	95.8	95.7	94.2	84.7	90.7
Utah	95.6	87.3	80.1	81.8	90.8
Vermont	83.3	84.0	64.8	70.2	72.6
Virginia	96.1	97.6	95.7	89.8	92.7
Wisconsin	90.2	84.3	75.3	77.9	79.6
Unweighted Data					
Arkansas	97.6	97.1	94.7	92.3	92.8
California	86.9	83.3	69.8	69.8	74.1
North Carolina	97.8	97.8	94.1	85.5	90.8
Oklahoma	90.4	93.2	88.9	85.1	77.8
Pennsylvania**	96.3	92.1	88.1	85.9	86.7
Rhode Island	85.0	85.0	78.0	78.3	78.3
South Carolina	94.9	89.8	84.5	84.9	91.0
Texas**	96.6	95.2	82.9	86.7	89.5
State Median	93.1	88.6	83.6	81.5	83.2

TABLE 9a. Percentage of Schools That Taught Specific Topics Related to Physical Activity,\* Selected U.S. Sites: School Health Profiles, Teachers' Surveys, 2002 (continued)

Site	Physical, psychological, or social benefits	Health-related fitness	Phases of a workout	How much physical activity is enough	Decreasing sedentary activities
Weighted Data					
Chicago	94.6	94.6	92.4	85.5	86.4
Dallas	100.0	93.6	83.4	80.2	90.0
Fort Lauderdale	86.7	83.7	71.1	74.4	84.0
Houston	100.0	95.5	93.4	93.4	93.2
Los Angeles	96.7	92.4	78.9	77.8	90.1
Miami	86.5	75.0	67.9	65.8	74.4
Orange County	87.7	70.8	60.2	59.6	100.0
Palm Beach	100.0	80.3	62.0	69.6	88.3
San Diego	58.3	55.6	47.4	36.1	35.9
San Francisco	86.3	90.7	73.3	77.6	82.0
Unweighted Data					
District of Columbia	100.0	96.3	88.9	77.8	77.8
Philadelphia	90.4	88.9	86.3	82.2	90.1
San Bernardino	100.0	66.7	50.0	66.7	83.3
<b>Local Median</b>	94.6	88.9	73.3	77.6	86.4

<sup>\*</sup> In a required health education course.

<sup>\*\*</sup> Survey did not include students from one of the state's large school districts.

TABLE 9b. Percentage of Schools That Taught Specific Topics Related to Physical Activity,\* Selected U.S. Sites: School Health Profiles, Teachers' Surveys, 2002

Site	Overcoming barriers to physical activity	Developing an individualized physical activity plan	Monitoring progress toward reaching goals	Opportunities for physical activity in the community	Preventing injury during physical activity	Weather-related safety	Dangers of performance- enhancing drugs**
<b>Weighted Data</b>							
Alabama	79.1	80.1	79.1	87.5	94.6	93.9	91.8
Alaska	59.8	59.3	54.0	74.3	81.2	73.6	63.3
Arizona	72.7	67.2	64.5	74.5	85.6	83.3	81.6
Connecticut	50.2	45.7	41.2	56.2	63.6	60.1	82.7
Delaware	67.8	65.9	51.9	68.9	82.6	77.8	93.2
Georgia	75.2	74.1	73.1	81.0	84.6	88.8	95.0
Hawaii	57.1	44.2	47.3	66.1	76.9	70.0	83.9
Idaho	73.3	69.2	62.0	70.7	85.6	84.0	91.8
Illinois <sup>§</sup>	58.7	58.4	50.7	66.4	78.9	78.1	89.9
lowa	64.9	64.8	57.2	70.0	80.4	80.1	88.0
Kentucky	81.2	76.6	72.1	85.3	93.7	87.4	93.9
Maine	63.8	61.6	56.2	67.7	75.5	64.3	78.8
Massachusetts	67.6	62.9	60.8	71.0	80.3	74.9	88.2
Michigan	64.1	60.9	54.1	67.7	75.4	68.8	83.8
Minnesota	59.9	62.8	60.8	63.9	73.2	74.1	84.0
Missouri	74.4	75.4	72.0	79.8	89.2	89.7	89.7
Montana	78.7	69.4	70.9	79.8	94.3	84.3	87.1
Nebraska	65.3	70.3	63.5	72.6	84.4	78.4	88.8
New Hampshire	61.4	54.9	49.1	59.2	76.0	71.1	87.7
New Mexico	80.2	72.1	67.9	77.1	89.5	79.9	89.7
New York	61.5	49.6	49.3	63.5	69.3	76.4	95.9
North Dakota	66.5	63.4	58.2	70.7	82.2	81.3	86.7
Tennessee	75.9	79.2	74.5	85.2	94.0	85.5	91.2
Utah	68.5	68.6	63.2	69.7	82.0	81.3	93.6
Vermont	50.2	46.6	44.5	51.0	60.5	57.2	88.0
Virginia	81.8	77.7	75.1	82.8	93.2	89.5	88.9
Wisconsin	62.5	60.8	51.7	66.9	74.7	75.9	86.0
Unweighted Da	ta						
Arkansas	76.9	76.3	69.2	84.2	93.3	89.5	91.8
California	58.2	54.0	54.7	61.7	73.5	69.4	82.6
North Carolina	77.2	77.7	71.2	84.2	93.5	90.3	89.7
Oklahoma	68.5	65.8	69.9	76.1	87.7	87.5	84.7
Pennsylvania <sup>§</sup>	67.8	65.1	60.7	71.7	87.2	83.9	91.7
Rhode Island	65.0	66.1	61.0	66.1	78.3	71.7	85.0
South Carolina	76.8	74.7	71.5	84.3	88.3	85.5	87.3
Texas§	76.2	72.6	71.0	73.8	87.5	92.9	96.2
State Median	67.8	65.9	61.0	71.0	82.6	80.1	88.2

TABLE 9b. Percentage of Schools That Taught Specific Topics Related to Physical Activity,\* Selected U.S. Sites: School Health Profiles, Teachers' Surveys, 2002 (continued)

Site	Overcoming barriers to physical activity	Developing an individualized physical activity plan	Monitoring progress toward reaching goals	Opportunities for physical activity in the community	Preventing injury during physical activity	Weather-related safety	Dangers of performance- enhancing drugs**
<b>Weighted Data</b>							
Chicago	75.2	63.2	61.3	85.6	90.4	76.3	76.1
Dallas	72.4	70.1	56.8	70.1	93.2	93.3	96.6
Fort Lauderdale	77.5	66.0	61.8	80.6	80.9	90.4	93.6
Houston	93.0	84.4	75.1	92.9	95.5	97.7	97.6
Los Angeles	65.5	60.0	59.8	66.3	74.8	80.7	88.8
Miami	65.1	63.6	60.2	66.5	70.5	80.7	80.1
Orange County	67.5	61.7	60.2	68.9	74.7	81.2	89.7
Palm Beach	65.9	54.9	54.9	80.8	69.8	84.8	92.0
San Diego	30.8	28.9	28.9	33.3	48.6	33.3	50.0
San Francisco	73.3	68.9	68.0	73.3	73.3	77.6	95.4
<b>Unweighted Dat</b>	a						
District of Columbia	77.8	70.4	74.1	81.5	88.9	74.1	92.6
Philadelphia	76.1	72.6	65.8	80.6	83.6	73.6	77.8
San Bernardino	66.7	33.3	33.3	50.0	66.7	50.0	100.0
<b>Local Median</b>	72.4	63.6	60.2	73.3	74.8	80.7	92.0

<sup>\*</sup> In a required health education course.

<sup>\*\*</sup> Such as steroids.

<sup>§</sup> Survey did not include students from one of the state's large school districts.

TABLE 10. Percentage of Schools That Used Specific Teaching Methods,\* Selected U.S. Sites: School Health Profiles, Teachers' Surveys, 2002

	Group	Cooperative group		Language, performing,	Pledges or	Adult guest	Peer		Computer- assisted
Site	discussions	activities	Role play	or visual arts	contracts	speakers	educators	The Internet	instruction
<b>Weighted Da</b>	ita								
Alabama	98.5	94.4	79.0	69.3	49.7	88.0	63.0	73.7	58.8
Alaska	92.1	88.0	65.8	62.2	40.7	83.5	58.0	78.8	42.5
Arizona	94.5	90.1	74.4	64.2	55.4	78.8	56.0	69.2	43.9
Connecticut	99.4	98.7	88.0	65.9	40.9	90.5	53.2	81.7	55.8
Delaware	100.0	100.0	92.5	64.3	51.1	95.0	55.4	90.0	50.0
Georgia	98.6	97.0	87.0	74.7	58.6	87.7	64.2	77.9	65.5
Hawaii	100.0	98.3	93.0	82.5	44.6	83.5	68.1	78.2	39.9
Idaho	99.4	96.3	81.0	74.9	39.9	86.8	61.3	82.7	55.1
Illinois**	97.7	93.6	74.6	63.6	37.4	79.8	44.4	74.5	52.8
Iowa	98.9	95.4	79.2	61.6	39.2	83.2	50.9	84.1	52.6
Kentucky	99.4	97.2	88.0	66.8	54.3	91.2	62.2	89.6	69.0
Maine	100.0	99.5	90.9	67.1	38.9	81.2	45.8	81.9	46.4
Massachusetts	99.6	99.3	89.9	76.7	47.8	86.8	61.0	81.8	56.8
Michigan	98.1	95.4	83.4	67.6	40.6	82.3	51.0	79.3	43.0
Minnesota	97.2	95.4	84.4	63.9	43.4	91.5	59.1	93.3	57.2
Missouri	98.9	95.6	79.0	67.7	35.5	83.8	61.1	79.7	54.8
Montana	98.0	95.1	77.0	65.6	39.7	78.8	56.5	73.8	57.5
Nebraska	99.2	96.3	74.8	51.9	41.2	75.0	50.0	80.6	58.7
New Hampshire	99.1	99.3	88.7	67.6	26.3	81.1	54.0	82.8	45.5
New Mexico	98.9	95.8	79.3	64.1	57.7	91.3	64.8	79.5	58.9
New York	99.0	98.0	91.8	73.8	53.7	86.5	66.8	83.1	61.4
North Dakota	97.9	95.0	80.0	60.1	40.2	78.2	50.8	90.7	58.6
Tennessee	97.6	92.3	76.2	67.1	50.7	87.8	63.6	70.3	54.6
Utah	97.7	97.1	87.5	69.1	49.3	91.1	50.8	71.1	50.1
Vermont	100.0	96.1	93.7	60.9	42.4	84.1	56.1	80.6	50.4
Virginia	98.6	93.2	85.2	65.2	55.7	83.2	61.1	85.1	71.6
Wisconsin	99.4	98.0	85.7	72.2	50.0	82.3	55.5	84.6	57.1
Unweighted	Data								
Arkansas	99.1	92.9	68.2	47.2	29.9	71.0	48.6	66.7	42.4
California	96.6	97.1	83.3	69.3	50.7	81.0	62.8	72.0	55.0
North Carolina	98.9	96.8	89.8	67.0	58.9	89.2	64.3	79.8	67.2
Oklahoma	98.7	93.5	62.8	58.4	43.2	69.3	46.8	63.5	42.9
Pennsylvania**	98.4	97.2	83.3	69.4	50.0	87.4	59.5	84.4	59.2
Rhode Island	100.0	100.0	86.7	64.4	50.0	78.7	49.2	80.0	67.8
South Carolina	98.4	95.7	78.2	63.8	51.6	78.0	50.3	79.6	61.0
Texas**	98.1	94.4	73.3	62.1	38.6	81.8	63.0	82.2	59.5
State Median	98.9	96.1	83.3	65.9	44.6	83.5	56.5	80.0	55.8

TABLE 10. Percentage of Schools That Used Specific Teaching Methods,\* Selected U.S. Sites: School Health Profiles, Teachers' Surveys, 2002 (continued)

Site	Group discussions	Cooperative group activities	Role play	Language, performing, or visual arts	Pledges or contracts	Adult guest speakers	Peer educators	The Internet	Computer- assisted instruction	
Weighted Dat	Weighted Data									
Chicago	98.0	94.7	87.2	74.2	67.1	85.5	58.1	58.8	54.4	
Dallas	100.0	100.0	86.2	72.2	51.8	93.0	69.0	62.0	48.2	
Fort Lauderdale	100.0	100.0	93.8	93.5	67.4	88.0	78.5	88.0	59.8	
Houston	100.0	95.5	90.4	83.2	76.9	80.8	65.0	66.6	57.1	
Los Angeles	97.8	98.8	88.4	83.7	70.5	89.3	61.1	80.1	72.7	
Miami	100.0	98.2	86.9	86.5	60.6	90.6	75.1	83.7	62.5	
Orange County	100.0	94.0	100.0	69.4	55.8	89.0	57.3	76.9	63.4	
Palm Beach	100.0	100.0	91.8	95.9	60.5	96.2	87.5	71.6	48.5	
San Diego	100.0	100.0	100.0	53.7	46.3	68.3	43.9	39.0	29.3	
San Francisco	95.7	86.3	82.0	87.0	65.2	95.7	77.0	71.4	72.7	
Unweighted I	Unweighted Data									
District of										
Columbia	89.3	89.3	82.1	71.4	71.4	89.3	64.3	82.1	64.3	
Philadelphia	98.6	97.2	91.7	80.3	62.0	68.5	60.6	74.6	48.6	
San Bernardino	100.0	83.3	100.0	83.3	33.3	50.0	50.0	50.0	33.3	
<b>Local Median</b>	100.0	97.2	90.4	83.2	62.0	89.0	64.3	71.6	57.1	

<sup>\*</sup> In a required health education course.

<sup>\*\*</sup> Survey did not include students from one of the state's large school districts.

Table 11. Percentage of Schools in Which a Specific Person Was Responsible for Coordinating Health Education, by Type of Position, Selected U.S. Sites: School Health Profiles, Principals' Surveys, 2002

Site	School district administrator*	School administrator	Health education teacher	School nurse	Other	No coordinator
Weighted Data**		darimistrator	tederier	Harse	Other	110 coordinator
Alabama	16.2	33.9	34.2	3.4	6.0	6.4
Alaska	25.3	23.5	29.5	4.4	6.5	11.0
Arizona	27.0	19.2	21.6	8.4	8.3	15.7
Arkansas	18.0	26.0	49.0	0.4	1.3	5.3
Connecticut	42.0	15.0	30.4	2.0	3.6	7.0
Delaware	34.5	9.5	52.0	0.0	2.0	2.0
Georgia	31.2	19.6	44.6	0.0	2.3	2.3
Hawaii	3.1	17.4	70.4	0.0	6.3	2.9
ldaho	23.2	15.8	55.8	0.0	3.2	2.0
Illinois <sup>§</sup>	20.3	20.6	49.2	0.8	3.2	5.9
lowa	28.0	13.3	45.4	3.2	4.7	5.3
Kentucky	7.1	24.1	60.4	0.0	3.0	5.3
Maine	28.8	11.4	44.5	6.2	2.4	6.6
Massachusetts	61.5	10.2	22.0	1.5	2.6	2.2
Michigan	31.4	19.5	34.9	4.1	5.1	5.0
Minnesota	25.3	14.2	53.1	0.7	3.7	3.0
Missouri	39.5	15.4	35.2	4.2	2.7	3.1
Montana	16.1	15.6	62.0	1.2	2.1	3.0
Nebraska	20.0	21.3	47.7	2.9	4.5	3.6
New Hampshire	13.0	21.6	44.5	9.6	6.1	5.2
New Jersey	50.3	26.3	15.1	2.4	3.4	2.6
New Mexico	16.7	16.0	46.0	7.9	5.5	7.8
New York	40.6	22.4	31.2	0.0	3.0	2.9
North Carolina	25.7	17.9	46.7	1.0	6.2	2.6
North Dakota	20.6	24.7	42.6	0.0	3.7	8.4
Oklahoma	19.0	26.6	18.0	14.0	6.3	16.1
Tennessee	19.3	20.5	40.1	3.4	5.9	10.8
Texas <sup>§</sup>	22.7	27.4	34.0	5.6	3.9	6.4
Utah	23.6	17.8	54.2	1.0	1.0	2.4
Vermont	7.1	19.9	44.6	9.2	9.8	9.5
Virginia	35.9	16.1	39.9	0.8	2.1	5.1
Wisconsin	28.8	8.4	49.7	1.8	7.1	4.2
Unweighted Data	a**					
California	23.2	20.5	32.7	5.4	6.5	11.6
Oregon	21.1	17.5	55.6	0.6	2.3	2.9
Pennsylvania <sup>§</sup>	40.2	21.3	34.5	0.3	1.7	2.0
Rhode Island	37.5	9.4	43.8	1.6	6.3	1.6
South Carolina	26.7	17.4	45.3	1.9	4.3	4.3
Washington	24.8	17.1	37.6	6.0	7.3	7.3
State Median	25.0	18.5	44.5	1.8	3.8	5.1

Table 11. Percentage of Schools in Which a Specific Person Was Responsible for Coordinating Health Education, by Type of Position, Selected U.S. Sites: School Health Profiles, Principals' Surveys, 2002 (continued)

	Calca al aliatoriat	Cabaal	Health	Calanal		
Site	School district administrator*	School administrator	education teacher	School nurse	Other	No coordinator
Weighted Data**						
Chicago	3.6	13.3	37.1	5.1	22.5	18.4
Dallas	40.7	12.2	28.3	0.0	4.2	14.5
Fort Lauderdale	25.5	11.7	30.6	0.0	27.5	4.7
Houston	27.1	15.6	50.3	2.3	2.3	2.3
Los Angeles	19.7	17.5	57.8	0.0	2.0	3.0
Miami	19.4	28.5	25.7	2.3	11.5	12.5
Orange County	17.8	27.3	35.4	2.5	11.7	5.2
Palm Beach	13.0	18.2	38.8	4.9	15.0	10.1
San Bernardino	42.9	35.7	7.1	0.0	0.0	14.3
San Diego	44.7	17.0	21.3	6.4	8.5	2.1
San Francisco	19.5	28.2	27.7	5.1	19.5	0.0
Unweighted Data*	*					
District of Columbia	7.1	17.9	75.0	0.0	0.0	0.0
Philadelphia	23.2	9.8	59.8	2.4	1.2	3.7
Local Median	19.7	17.5	35.4	2.3	8.5	4.7

<sup>\*</sup> District administrator or district health education or curriculum coordinator.

<sup>\*\*</sup> Percentages for each row might not add up to 100.0 due to rounding.

<sup>§</sup> Survey did not include students from one of the state's large school districts.

TABLE 12. Percentage of Schools in Which Health Education Staff Planned or Coordinated Health-Related Projects or Activities with Other Groups, Selected U.S. Sites: School Health Profiles, Teachers' Surveys, 2002

Cit-	Physical	School health	School mental	Food	Community
Site	education staff	services staff	health staff	service staff	members
Weighted Data	747	74.0	<b></b>	20.0	52.4
Alabama	74.7	74.0	60.2	29.9	53.1
Alaska	56.2	50.1	56.0	19.6	58.0
Arizona	54.9	57.6	50.3	21.1	48.4
Connecticut	69.0	70.4	67.2	13.1	60.6
Delaware	91.2	83.4	73.5	20.9	68.0
Georgia	83.0	64.1	55.3	29.4	62.6
Hawaii	70.8	52.3	43.7	18.6	65.3
daho	70.9	60.4	51.9	20.7	54.8
llinois*	70.0	51.9	55.6	13.9	53.2
owa	64.2	72.9	48.1	21.4	54.4
Kentucky	81.6	65.7	55.3	25.3	66.7
Maine	72.8	79.4	64.4	25.6	56.6
Massachusetts	81.9	83.8	79.4	29.8	73.0
Michigan	61.6	38.8	44.9	13.7	46.8
Minnesota	74.4	66.9	66.3	21.0	61.9
Missouri	87.6	82.0	59.0	26.5	50.0
Montana	86.1	55.1	52.9	26.1	52.0
Nebraska	64.7	60.8	40.7	18.6	44.0
New Hampshire	62.0	77.1	63.8	21.1	58.4
New Mexico	66.8	78.6	67.1	27.1	61.7
New York	70.2	67.0	68.3	22.1	68.5
North Dakota	65.8	36.4	57.1	31.5	50.4
Tennessee	72.9	66.0	61.2	35.0	57.5
Utah	69.9	47.1	60.5	9.8	64.5
Vermont	64.2	77.2	74.2	32.5	60.1
Virginia	87.9	73.9	53.5	20.4	50.0
Wisconsin	68.7	70.4	64.1	17.0	56.2
Unweighted Data					
Arkansas	76.1	65.5	48.0	15.9	46.5
California	51.6	50.3	43.2	14.8	44.1
North Carolina	84.6	69.8	63.1	23.6	58.3
Oklahoma	52.8	44.1	47.1	23.9	30.2
Pennsylvania*	89.7	78.3	56.5	15.8	59.1
Rhode Island	80.0	81.3	65.1	15.9	55.6
South Carolina	78.8	66.7	45.5	25.7	48.7
Texas*	65.5	65.6	47.7	20.3	46.3
State Median	70.8	66.7	56.5	21.1	56.2

TABLE 12. Percentage of Schools in Which Health Education Staff Planned or Coordinated Health-Related Projects or Activities with Other Groups, Selected U.S. Sites: School Health Profiles, Teachers' Surveys, 2002 (continued)

Site	Physical education staff	School health services staff	School mental health staff	Food service staff	Community members
Weighted Data					
Chicago	77.0	71.2	71.4	39.5	40.8
Dallas	52.9	75.9	50.0	22.5	36.4
Fort Lauderdale	56.6	28.2	51.4	29.9	49.7
Houston	92.3	90.4	66.7	24.5	51.1
Los Angeles	47.2	60.0	52.7	28.0	56.2
Miami	40.8	32.1	58.8	12.9	45.1
Orange County	52.4	50.9	51.3	14.5	62.1
Palm Beach	42.1	64.0	61.7	12.8	61.2
San Diego	53.1	79.6	65.3	8.3	59.2
San Francisco	67.7	71.4	65.1	15.3	69.4
<b>Unweighted Data</b>					
District of Columbia	96.7	90.3	71.0	19.4	64.5
Philadelphia	85.7	70.6	54.2	32.5	42.9
San Bernardino	28.6	71.4	57.1	0.0	28.6
<b>Local Median</b>	53.1	71.2	58.8	19.4	51.1

<sup>\*</sup> Survey did not include students from one of the state's large school districts.

TABLE 13. Percentage of Schools in Which the Lead Health Education Teacher Had Professional Preparation, by Specific Discipline, Selected U.S. Sites: School Health Profiles, Teachers' Surveys, 2002

Site	Health and physical education	Health education only	Physical education only	Science or other education degree	Nursing or counseling	Another discipline*
		education only	education only	education degree	or counseling	discipline"
Weighted Data** Alabama	51.1	8.2	20.2	13.2	3.9	3.3
Alaska	13.3	1.9	8.5	48.3	6.4	21.6
Arizona	19.3	5.3	10.5	26.4	21.1	17.4
Connecticut	44.5	18.3	13.2	6.8	6.3	10.8
Delaware	83.2	4.6	7.1	2.5	0.0	2.5
Georgia	72.0	5.6	3.5	11.6	3.6	3.8
Hawaii	41.9	9.2	21.3	18.3	1.8	7.5
Idaho	57.9	5.3	18.2	9.1	1.8	7.7
Illinois§	42.2	11.0	17.9	19.6	3.1	6.1
lowa	29.2	6.0	13.5	14.8	6.8	29.6
Kentucky	59.0	12.5	10.9	8.9	3.0	5.7
Maine	37.8	12.5	9.5	13.6	3.0 8.1	12.1
Massachusetts	46.8	23.9	5.3	5.5	8.7	9.8
	34.8	9.4	14.2	17.6	4.6	19.5
Michigan Minnesota	71.2	11.2	9.4	5.7	1.1	1.3
Missouri	46.8	3.0	25.4	6.7	2.5	15.7
Montana	63.8	1.8	10.7	16.9	2.5	4.4
Nebraska	37.2	2.0	20.9	20.8	4.8	14.5
New Hampshire	24.1	20.3	13.4	6.0	23.2	13.0
New Mexico	34.2	8.8	15.4	15.2	13.1	13.5
New York	47.1	37.7	6.5	3.7	1.3	3.7
North Dakota	37.2	4.5	14.1	15.6	1.9	26.7
Tennessee	55.8	4.0	5.7	19.1	9.0	6.3
Utah	48.7	14.8	14.7	12.9	0.7	8.3
Vermont	27.8	11.5	4.5	9.6	23.2	23.3
Virginia	84.5	1.3	6.6	4.2	1.0	23.3
Wisconsin	55.6	7.1	15.5	10.0	3.0	8.9
WISCOTISITI	33.0	7.1	15.5	10.0	3.0	0.9
<b>Unweighted Data</b>	**					
Arkansas	68.1	2.9	20.0	2.9	1.9	4.3
California	16.0	12.2	19.8	30.8	4.6	16.7
North Carolina	55.8	4.3	25.8	4.3	5.6	4.3
Oklahoma	39.0	3.4	6.0	20.6	21.0	10.1
Pennsylvania§	87.3	5.0	1.9	1.5	1.2	3.1
Rhode Island	55.7	16.4	11.5	1.6	13.1	1.6
South Carolina	46.2	5.4	31.2	8.5	5.0	3.8
Texas <sup>§</sup>	51.1	8.4	7.9	11.5	4.8	16.3
State Median	46.8	7.1	13.2	11.5	4.6	8.3

TABLE 13. Percentage of Schools in Which the Lead Health Education Teacher Had Professional Preparation, by Specific Discipline, Selected U.S. Sites: School Health Profiles, Teachers' Surveys, 2002 (continued)

Site	Health and physical education	Health education only	Physical education only	Science or other education degree	Nursing or counseling	Another discipline*
Weighted Data**						
Chicago	35.1	2.3	25.1	24.0	11.2	2.3
Dallas	34.4	17.3	2.1	41.8	0.0	4.3
Fort Lauderdale	28.1	10.9	8.0	39.0	5.7	8.3
Houston	81.8	0.0	11.4	0.0	0.0	6.8
Los Angeles	20.5	31.1	1.1	36.6	2.1	8.5
Miami	17.2	11.4	8.3	38.4	10.5	14.2
Orange County	19.4	26.2	2.7	38.4	2.3	10.9
Palm Beach	44.1	15.9	2.6	26.2	5.4	5.6
San Diego	0.0	25.6	0.0	30.8	35.9	7.7
San Francisco	12.9	3.1	0.0	57.3	6.7	20.0
<b>Unweighted Data</b> *	<b>**</b>					
District of Columbia	88.9	0.0	3.7	3.7	3.7	0.0
Philadelphia	77.8	3.7	4.9	3.7	4.9	4.9
San Bernardino	20.0	40.0	0.0	40.0	0.0	0.0
Local Median	28.1	11.4	2.7	36.6	4.9	6.8

<sup>\*</sup> Includes kinesiology, exercise science or exercise physiology, public health, home economics or family and consumer science, and other.

<sup>\*\*</sup> Percentages for each row might not add up to 100.0 because of rounding.

<sup>§</sup> Survey did not include students from one of the state's large school districts.