

TABLE 14a. Percentage of Schools in Which the Lead Health Education Teacher Had Received Staff Development During the Preceding 2 Years, by Specific Health Education Topic, Selected U.S. Sites: School Health Profiles, Teachers' Surveys, 2002

**STATE SURVEYS**

Site	Accident or injury prevention	Alcohol or other drug-use prevention	Consumer health	CPR*	Death and dying	Dental and oral health	Emotional and mental health
<b>Weighted Data</b>							
Alabama	57.6	56.8	18.9	76.0	13.6	15.4	33.7
Alaska	50.5	46.4	15.6	48.3	13.1	19.1	33.5
Arizona	55.9	49.3	15.6	61.3	16.6	19.6	38.1
Connecticut	30.5	49.6	10.8	55.0	12.5	5.3	36.0
Delaware	23.3	55.6	12.5	58.7	10.5	2.3	32.9
Georgia	40.2	50.9	10.3	65.0	8.6	8.1	23.5
Hawaii	30.4	57.0	22.6	37.0	6.4	5.4	43.7
Idaho	25.9	46.7	10.5	40.6	9.7	6.4	26.1
Illinois**	30.9	45.8	12.0	54.1	10.4	7.1	30.2
Iowa	31.9	38.9	15.4	50.8	10.6	8.6	27.1
Kentucky	47.3	36.8	14.0	69.4	7.1	6.2	21.8
Maine	28.0	40.8	10.4	48.3	10.1	6.6	34.0
Massachusetts	35.2	65.5	16.9	66.6	16.3	10.8	52.1
Michigan	23.7	50.5	15.7	48.0	8.7	8.6	30.4
Minnesota	41.0	57.2	14.8	54.0	13.7	9.8	40.6
Missouri	44.1	53.3	18.5	68.0	14.6	12.0	31.2
Montana	48.2	57.4	17.0	66.8	13.9	12.2	30.5
Nebraska	34.7	34.8	11.4	52.6	18.6	9.6	22.2
New Hampshire	40.0	60.9	19.4	64.9	12.6	4.4	49.2
New Mexico	53.1	52.4	26.9	67.7	15.2	19.4	42.3
New York	35.2	52.6	14.3	56.2	17.6	9.4	37.5
North Dakota	31.3	53.0	21.6	57.8	12.8	7.9	30.1
Tennessee	55.8	52.8	21.0	75.0	16.8	20.0	38.4
Utah	37.4	59.9	27.0	61.6	11.5	8.1	44.5
Vermont	31.3	68.0	17.5	59.0	17.1	17.1	48.0
Virginia	46.8	50.2	17.2	69.4	12.2	10.6	25.9
Wisconsin	42.9	50.8	13.9	60.7	13.1	5.8	38.2
<b>Unweighted Data</b>							
Arkansas	47.4	43.7	11.3	60.9	8.3	7.9	21.0
California	38.7	45.9	11.2	55.0	9.6	9.9	26.3
North Carolina	42.7	34.9	11.8	70.2	7.5	7.2	28.3
Oklahoma	61.0	57.1	19.6	63.4	15.2	15.9	33.0
Pennsylvania**	41.1	62.3	12.7	81.3	9.7	5.4	34.2
Rhode Island	34.4	50.8	12.7	57.8	9.7	7.9	28.6
South Carolina	40.8	37.0	17.2	58.5	8.9	15.9	22.6
Texas**	52.5	50.2	12.7	80.1	10.1	12.0	25.1
<b>State Median</b>	<b>40.2</b>	<b>50.8</b>	<b>15.4</b>	<b>60.7</b>	<b>12.2</b>	<b>8.6</b>	<b>32.9</b>

(continued)

TABLE 14a. Percentage of Schools in Which the Lead Health Education Teacher Had Received Staff Development During the Preceding 2 Years, by Specific Health Education Topic, Selected U.S. Sites: School Health Profiles, Teachers' Surveys, 2002 (continued)

### LOCAL SURVEYS

Site	Accident or injury prevention	Alcohol or other drug-use prevention	Consumer health	CPR*	Death and dying	Dental and oral health	Emotional and mental health
<b>Weighted Data</b>							
Chicago	47.8	50.9	28.5	45.5	26.7	31.7	45.7
Dallas	59.5	54.0	27.0	65.4	12.4	14.7	43.7
Fort Lauderdale	40.5	68.1	17.2	71.4	19.3	12.2	45.9
Houston	82.9	80.9	43.1	98.1	37.3	38.6	55.7
Los Angeles	46.2	52.2	20.2	37.8	13.4	14.5	36.8
Miami	32.1	44.5	19.8	53.9	16.5	13.2	28.5
Orange County	29.2	46.1	7.4	52.3	16.0	4.7	25.6
Palm Beach	74.6	74.8	24.1	85.5	15.3	7.7	40.2
San Diego	20.8	93.8	29.2	27.1	37.5	0.0	58.3
San Francisco	21.9	79.2	32.9	46.7	18.6	15.7	44.3
<b>Unweighted Data</b>							
District of Columbia	29.0	71.0	22.6	63.3	32.3	31.0	53.3
Philadelphia	36.1	34.9	14.6	49.4	7.2	14.5	30.1
San Bernardino	37.5	71.4	14.3	50.0	0.0	0.0	42.9
<b>Local Median</b>	<b>37.5</b>	<b>68.1</b>	<b>22.6</b>	<b>52.3</b>	<b>16.5</b>	<b>14.5</b>	<b>43.7</b>

\* CPR = cardiopulmonary resuscitation.

\*\* Survey did not include students from one of the state's large school districts.

TABLE 14b. Percentage of Schools in Which the Lead Health Education Teacher Had Received Staff Development During the Preceding 2 Years, by Specific Health Education Topic, Selected U.S. Sites: School Health Profiles, Teachers' Surveys, 2002

**STATE SURVEYS**

Site	Environmental health	First aid	Growth and development	HIV* prevention	Human sexuality	Immunization and vaccinations	Nutrition and dietary behavior
<b>Weighted Data</b>							
Alabama	21.9	66.8	23.7	50.7	25.6	14.3	29.5
Alaska	17.9	53.6	19.2	28.1	14.7	32.6	21.3
Arizona	27.1	57.5	26.5	49.1	23.1	37.2	26.7
Connecticut	9.8	50.0	28.7	45.9	37.4	12.3	23.9
Delaware	10.5	40.5	18.9	42.2	39.9	4.2	28.7
Georgia	15.4	60.0	28.6	55.8	36.9	11.5	19.4
Hawaii	19.3	24.6	31.1	46.7	44.4	3.6	41.7
Idaho	9.4	38.1	12.0	53.7	26.1	8.8	24.3
Illinois**	13.8	47.8	16.9	33.7	26.7	12.0	23.3
Iowa	16.2	36.6	18.5	38.9	19.2	20.8	26.0
Kentucky	11.2	64.5	18.0	34.4	17.9	13.4	23.1
Maine	11.0	37.5	21.5	49.1	33.2	16.4	24.6
Massachusetts	15.1	50.4	33.0	44.7	38.0	14.1	45.5
Michigan	10.8	41.3	29.0	57.2	44.9	19.4	33.5
Minnesota	17.1	47.1	21.1	51.6	40.0	16.6	25.4
Missouri	17.9	57.5	23.1	31.0	21.2	19.8	33.3
Montana	12.8	66.4	24.6	54.3	35.0	13.9	31.5
Nebraska	13.8	39.8	19.0	24.8	20.3	17.2	22.9
New Hampshire	20.1	55.5	36.6	54.6	45.8	20.1	48.2
New Mexico	29.6	60.5	29.0	52.4	34.9	34.2	34.0
New York	16.4	48.8	23.8	58.5	33.0	14.7	32.6
North Dakota	14.9	37.3	16.5	38.8	17.4	16.5	31.7
Tennessee	23.8	68.7	28.8	50.7	26.9	29.3	33.7
Utah	16.3	53.2	25.8	71.6	59.0	9.1	43.2
Vermont	15.0	40.3	28.7	48.1	41.3	21.2	23.0
Virginia	14.0	59.1	23.4	38.8	23.3	17.3	23.9
Wisconsin	19.7	54.1	28.5	43.0	25.3	18.6	20.9
<b>Unweighted Data</b>							
Arkansas	13.9	53.9	16.0	33.2	13.9	12.2	17.0
California	14.5	49.8	20.5	50.0	31.3	18.8	24.3
North Carolina	12.2	63.6	22.1	39.7	33.6	14.1	28.3
Oklahoma	26.3	64.7	23.1	76.7	23.7	41.9	28.4
Pennsylvania**	11.6	61.3	19.9	41.6	29.5	8.5	24.0
Rhode Island	15.9	47.7	25.4	49.2	45.3	22.2	26.6
South Carolina	16.6	57.3	24.4	47.8	42.1	22.9	29.3
Texas**	13.6	67.1	20.9	32.2	20.2	18.3	22.4
<b>State Median</b>	<b>15.1</b>	<b>53.6</b>	<b>23.4</b>	<b>47.8</b>	<b>31.3</b>	<b>16.6</b>	<b>26.6</b>

(continued)

TABLE 14b. Percentage of Schools in Which the Lead Health Education Teacher Had Received Staff Development During the Preceding 2 Years, by Specific Health Education Topic, Selected U.S. Sites: School Health Profiles, Teachers' Surveys, 2002 (continued)

### LOCAL SURVEYS

Site	Environmental health	First aid	Growth and development	HIV* prevention	Human sexuality	Immunization and vaccinations	Nutrition and dietary behavior
<b>Weighted Data</b>							
Chicago	29.9	54.4	43.1	54.7	40.4	42.9	42.4
Dallas	40.0	68.4	50.8	56.9	50.7	16.2	26.9
Fort Lauderdale	25.4	63.6	45.3	91.0	75.0	22.1	42.5
Houston	50.1	98.1	54.0	75.9	58.7	34.6	56.9
Los Angeles	29.8	41.0	29.5	71.9	37.7	30.4	36.3
Miami	19.8	35.9	52.2	68.2	51.4	29.6	34.5
Orange County	15.2	50.2	27.4	97.6	80.4	10.9	16.5
Palm Beach	29.6	77.3	33.9	80.1	46.5	26.3	29.3
San Diego	25.0	25.0	39.6	85.4	89.6	14.6	43.8
San Francisco	21.5	41.7	38.4	70.6	60.7	25.1	26.3
<b>Unweighted Data</b>							
District of Columbia	25.8	54.8	66.7	71.0	71.0	45.2	38.7
Philadelphia	19.3	41.2	24.1	35.4	25.6	13.3	30.1
San Bernardino	14.3	42.9	42.9	100.0	75.0	14.3	28.6
<b>Local Median</b>	<b>25.4</b>	<b>50.2</b>	<b>42.9</b>	<b>71.9</b>	<b>58.7</b>	<b>25.1</b>	<b>34.5</b>

\* HIV = human immunodeficiency virus.

\*\* Survey did not include students from one of the state's large school districts.

TABLE 14c. Percentage of Schools in Which the Lead Health Education Teacher Had Received Staff Development During the Preceding 2 Years, by Specific Health Education Topic, Selected U.S. Sites: School Health Profiles, Teachers' Surveys, 2002

**STATE SURVEYS**

Site	Personal hygiene	Physical activity and fitness	Pregnancy prevention	STD* prevention	Suicide prevention	Sun safety	Tobacco-use prevention	Violence prevention
<b>Weighted Data</b>								
Alabama	18.7	53.9	30.6	44.8	24.4	15.4	44.2	57.6
Alaska	13.4	29.7	14.4	19.1	23.5	8.0	33.9	47.3
Arizona	21.9	36.1	24.9	31.9	28.9	22.1	49.7	60.9
Connecticut	9.3	40.2	23.7	36.0	24.1	5.9	28.8	57.6
Delaware	6.4	56.2	33.9	40.3	14.1	14.1	43.1	57.5
Georgia	11.6	39.0	33.8	48.9	22.1	9.7	36.3	42.7
Hawaii	17.2	45.5	37.2	47.5	27.4	13.9	55.5	49.2
Idaho	7.3	31.6	16.3	41.0	21.2	6.5	30.5	45.4
Illinois**	8.2	36.5	22.7	29.5	21.8	7.3	39.3	51.5
Iowa	10.3	30.8	19.1	25.4	18.7	10.4	27.4	40.4
Kentucky	10.1	40.0	18.2	25.7	9.0	8.8	31.0	50.0
Maine	8.1	40.5	31.1	37.4	36.6	6.1	34.3	54.0
Massachusetts	10.6	48.3	28.2	35.8	34.2	18.6	49.3	74.6
Michigan	13.9	39.9	30.8	47.9	17.9	9.4	41.1	48.9
Minnesota	11.1	41.2	35.7	46.6	32.3	10.4	51.0	51.5
Missouri	15.9	41.5	21.9	29.6	22.7	13.8	32.9	50.6
Montana	13.7	46.6	26.2	38.1	21.6	13.7	57.1	51.2
Nebraska	12.9	32.8	15.7	20.7	17.4	9.6	28.8	39.6
New Hampshire	7.4	52.1	32.0	40.1	29.1	11.4	40.7	57.5
New Mexico	23.1	49.7	35.3	47.6	30.5	19.0	45.7	54.2
New York	12.0	34.8	31.1	41.5	26.6	13.6	43.4	61.3
North Dakota	14.8	43.2	15.7	28.9	23.2	11.1	41.2	57.9
Tennessee	21.4	51.9	28.6	41.5	25.0	17.5	39.8	60.1
Utah	9.8	38.2	43.0	64.0	47.9	10.9	52.8	49.1
Vermont	5.3	40.9	27.9	47.1	22.7	9.9	49.7	51.9
Virginia	11.7	67.3	20.1	27.5	19.2	12.7	32.6	52.6
Wisconsin	11.0	32.3	21.4	25.9	21.6	7.3	34.6	46.3
<b>Unweighted Data</b>								
Arkansas	9.6	40.3	18.2	29.4	16.7	12.7	26.6	40.0
California	11.2	30.1	27.5	40.0	19.5	13.1	41.5	44.7
North Carolina	9.2	50.4	31.9	38.9	15.5	12.2	31.1	41.9
Oklahoma	20.9	38.3	26.0	47.6	26.7	11.5	36.9	61.5
Pennsylvania**	8.1	47.7	24.6	35.6	24.1	9.7	34.1	62.3
Rhode Island	14.3	43.8	21.9	33.3	12.7	15.6	38.1	45.3
South Carolina	16.4	53.1	33.1	40.8	12.6	9.3	28.0	34.1
Texas**	12.5	47.1	22.7	31.7	22.0	13.1	34.1	50.8
<b>State Median</b>	<b>11.6</b>	<b>40.9</b>	<b>26.2</b>	<b>38.1</b>	<b>22.7</b>	<b>11.4</b>	<b>38.1</b>	<b>51.2</b>

(continued)

TABLE 14c. Percentage of Schools in Which the Lead Health Education Teacher Had Received Staff Development During the Preceding 2 Years, by Specific Health Education Topic, Selected U.S. Sites: School Health Profiles, Teachers' Surveys, 2002 (continued)

### LOCAL SURVEYS

Site	Personal hygiene	Physical activity and fitness	Pregnancy prevention	STD* prevention	Suicide prevention	Sun safety	Tobacco-use prevention	Violence prevention
<b>Weighted Data</b>								
Chicago	35.5	54.9	31.0	48.2	22.4	13.7	43.1	61.9
Dallas	16.8	39.3	32.3	42.6	36.5	18.7	40.8	59.2
Fort Lauderdale	14.3	51.7	42.3	82.2	35.6	50.3	64.8	59.7
Houston	48.2	90.4	56.0	72.5	63.5	52.0	75.1	86.7
Los Angeles	20.3	31.7	32.7	56.9	24.4	18.6	72.7	54.4
Miami	21.9	32.8	45.0	60.1	27.9	18.6	37.4	49.2
Orange County	15.3	15.3	62.6	84.6	22.7	16.9	26.8	59.7
Palm Beach	16.2	42.4	43.9	56.8	33.9	41.5	57.0	82.7
San Diego	0.0	27.1	81.3	89.6	72.9	20.8	83.3	93.8
San Francisco	20.8	35.7	48.2	66.7	42.9	9.3	79.2	68.2
<b>Unweighted Data</b>								
District of Columbia	32.3	64.5	48.4	58.1	25.8	12.9	35.5	51.6
Philadelphia	16.0	61.7	22.9	32.5	15.7	6.0	28.0	51.8
San Bernardino	14.3	28.6	57.1	75.0	14.3	14.3	42.9	28.6
<b>Local Median</b>	<b>16.8</b>	<b>39.3</b>	<b>45.0</b>	<b>60.1</b>	<b>27.9</b>	<b>18.6</b>	<b>43.1</b>	<b>59.7</b>

\* STD = sexually transmitted disease.

\*\* Survey did not include students from one of the state's large school districts.

TABLE 15a. Percentage of Schools in Which the Lead Health Education Teacher Wanted Staff Development, by Specific Health Education Topic, Selected U.S. Sites: School Health Profiles, Teachers' Surveys, 2002

**STATE SURVEYS**

Site	Accident or injury prevention	Alcohol or other drug-use prevention	Consumer health	CPR*	Death and dying	Dental and oral health	Emotional and mental health
<b>Weighted Data</b>							
Alabama	60.4	76.1	52.0	74.3	57.1	51.1	67.4
Alaska	43.8	60.0	43.4	64.9	55.2	37.2	63.9
Arizona	57.0	72.0	52.7	67.6	53.9	48.1	67.0
Connecticut	41.6	74.0	53.3	54.0	68.3	33.2	77.8
Delaware	51.3	78.9	57.8	61.3	74.4	33.2	70.3
Georgia	54.9	72.7	43.9	66.2	57.6	46.5	63.1
Hawaii	62.1	60.3	70.1	71.4	64.8	50.2	64.7
Idaho	48.2	65.0	48.1	61.8	51.6	33.6	66.4
Illinois**	43.1	65.2	42.0	55.9	46.5	29.6	58.9
Iowa	35.3	60.3	44.6	54.7	48.7	29.2	56.3
Kentucky	56.2	66.8	50.4	61.2	58.4	41.8	59.3
Maine	33.9	62.3	50.5	51.6	51.7	22.1	62.5
Massachusetts	50.7	76.1	53.4	60.2	64.9	36.5	75.1
Michigan	41.6	64.6	46.3	61.3	51.1	33.2	58.8
Minnesota	41.0	71.2	45.2	52.2	55.6	26.4	72.5
Missouri	53.6	67.5	49.3	67.9	47.8	37.3	61.0
Montana	49.8	65.3	47.5	61.3	59.2	42.7	67.8
Nebraska	33.3	52.7	31.4	49.7	35.3	22.9	48.8
New Hampshire	52.1	70.5	60.0	52.9	56.4	41.7	73.9
New Mexico	60.3	75.6	53.4	65.5	59.7	44.1	69.9
New York	44.9	75.0	53.7	56.6	64.9	36.1	70.2
North Dakota	42.8	62.1	38.7	58.7	45.3	26.7	57.6
Tennessee	59.4	73.8	51.0	72.9	59.8	48.3	69.9
Utah	46.5	65.5	57.5	57.8	61.4	35.2	72.5
Vermont	31.7	57.2	39.0	37.8	54.4	20.5	61.6
Virginia	48.7	63.0	44.9	61.9	55.9	35.9	62.7
Wisconsin	39.7	62.3	49.6	54.9	54.2	28.2	63.4
<b>Unweighted Data</b>							
Arkansas	55.9	74.8	47.2	66.5	49.4	42.2	61.6
California	37.7	61.8	41.3	58.4	42.4	29.3	66.1
North Carolina	50.2	71.4	38.2	64.6	52.3	35.9	62.3
Oklahoma	55.5	73.0	42.7	71.3	45.6	41.3	62.6
Pennsylvania**	56.0	75.9	52.5	65.8	66.8	36.3	75.2
Rhode Island	47.5	64.1	59.7	62.9	71.0	30.0	71.9
South Carolina	54.9	70.6	50.0	68.3	54.9	45.6	63.4
Texas**	55.6	73.4	47.0	71.7	56.2	43.0	64.9
<b>State Median</b>	<b>49.8</b>	<b>67.5</b>	<b>49.3</b>	<b>61.3</b>	<b>55.6</b>	<b>36.1</b>	<b>64.7</b>

(continued)

TABLE 15a. Percentage of Schools in Which the Lead Health Education Teacher Wanted Staff Development, by Specific Health Education Topic, Selected U.S. Sites: School Health Profiles, Teachers' Surveys, 2002 (continued)

**LOCAL SURVEYS**

Site	Accident or injury prevention	Alcohol or other drug-use prevention	Consumer health	CPR*	Death and dying	Dental and oral health	Emotional and mental health
<b>Weighted Data</b>							
Chicago	67.4	74.0	64.4	83.4	70.2	61.5	73.1
Dallas	51.9	65.0	47.7	81.7	47.9	46.6	66.6
Fort Lauderdale	63.5	71.3	60.0	75.7	71.0	56.2	70.3
Houston	72.0	88.0	71.5	76.0	85.6	73.5	84.0
Los Angeles	56.7	74.7	59.8	70.9	62.4	46.6	71.6
Miami	63.1	71.9	56.2	80.4	62.9	52.8	71.7
Orange County	33.1	68.9	48.1	50.7	58.5	35.6	63.5
Palm Beach	45.7	63.9	56.2	57.7	53.0	54.1	77.3
San Diego	22.9	79.2	37.5	50.0	45.8	20.8	66.7
San Francisco	65.1	73.7	60.3	63.2	64.3	53.3	76.5
<b>Unweighted Data</b>							
District of Columbia	63.3	80.6	70.0	74.2	64.3	73.3	73.3
Philadelphia	62.5	73.5	63.3	80.2	63.1	46.3	77.8
San Bernardino	71.4	62.5	42.9	71.4	42.9	42.9	75.0
<b>Local Median</b>	<b>63.1</b>	<b>73.5</b>	<b>59.8</b>	<b>74.2</b>	<b>62.9</b>	<b>52.8</b>	<b>73.1</b>

\* CPR = cardiopulmonary resuscitation.

\*\* Survey did not include students from one of the state's large school districts.



TABLE 15b. Percentage of Schools in Which the Lead Health Education Teacher Wanted Staff Development, by Specific Health Education Topic, Selected U.S. Sites: School Health Profiles, Teachers' Surveys, 2002

**STATE SURVEYS**

Site	Environmental health	First aid	Growth and development	HIV* prevention	Human sexuality	Immunization and vaccinations	Nutrition and dietary behavior
<b>Weighted Data</b>							
Alabama	55.8	76.6	53.3	67.7	59.6	47.4	69.1
Alaska	50.9	63.3	44.2	51.6	46.1	44.6	59.1
Arizona	58.0	71.1	54.2	60.4	47.7	47.8	63.7
Connecticut	61.4	54.4	61.9	69.7	73.5	40.1	68.3
Delaware	48.6	53.7	58.1	75.1	72.5	43.8	58.1
Georgia	52.5	67.6	56.0	71.0	63.9	48.4	64.4
Hawaii	62.8	64.3	60.6	70.4	68.3	60.3	73.3
Idaho	44.9	69.6	49.1	60.0	56.5	41.1	59.0
Illinois**	44.6	56.4	39.6	56.9	50.6	34.7	53.8
Iowa	43.4	52.9	41.3	53.8	49.6	36.1	52.3
Kentucky	54.7	63.9	48.0	60.1	56.4	46.3	59.6
Maine	51.5	50.3	39.7	52.6	56.7	33.6	57.6
Massachusetts	58.5	57.8	58.4	68.9	70.8	42.6	65.4
Michigan	48.2	61.2	45.7	57.0	53.0	38.6	59.4
Minnesota	48.8	54.2	44.7	63.0	60.3	42.9	52.3
Missouri	52.1	70.1	48.1	61.7	51.4	42.3	63.4
Montana	53.2	61.2	51.1	64.2	58.4	48.6	66.6
Nebraska	32.7	49.8	32.8	46.9	41.4	31.4	46.3
New Hampshire	66.4	57.0	56.1	62.6	69.6	49.5	69.8
New Mexico	56.5	69.9	56.4	65.5	60.1	50.2	62.9
New York	58.8	53.3	55.6	73.0	73.2	46.4	61.6
North Dakota	37.7	57.2	32.6	53.4	42.0	32.9	50.5
Tennessee	59.0	73.8	58.0	69.4	56.0	49.9	69.2
Utah	54.2	58.8	55.4	62.4	61.9	48.2	62.1
Vermont	47.8	39.1	46.1	56.6	57.4	23.0	57.0
Virginia	48.3	63.3	43.7	52.0	49.3	37.6	61.4
Wisconsin	52.2	53.6	51.8	58.6	58.8	35.0	61.9
<b>Unweighted Data</b>							
Arkansas	52.8	71.4	49.4	71.3	52.8	46.1	63.2
California	45.7	58.0	44.9	56.3	50.5	34.4	57.9
North Carolina	44.3	65.8	46.2	63.1	55.1	37.2	67.1
Oklahoma	49.3	71.3	44.6	61.6	41.0	46.8	57.5
Pennsylvania**	58.9	69.0	59.9	83.4	75.1	48.6	72.7
Rhode Island	67.8	53.3	60.7	72.6	72.6	36.7	61.9
South Carolina	55.1	69.5	52.4	63.6	61.3	45.8	67.8
Texas**	52.4	68.8	48.8	66.4	59.2	45.3	62.8
<b>State Median</b>	<b>52.4</b>	<b>61.2</b>	<b>49.4</b>	<b>62.6</b>	<b>57.4</b>	<b>43.8</b>	<b>61.9</b>

(continued)

TABLE 15b. Percentage of Schools in Which the Lead Health Education Teacher Wanted Staff Development, by Specific Health Education Topic, Selected U.S. Sites: School Health Profiles, Teachers' Surveys, 2002 (continued)

**LOCAL SURVEYS**

Site	Environmental health	First aid	Growth and development	HIV* prevention	Human sexuality	Immunization and vaccinations	Nutrition and dietary behavior
<b>Weighted Data</b>							
Chicago	65.4	79.0	64.2	76.7	68.5	61.1	66.6
Dallas	56.5	72.8	62.1	67.9	58.5	48.8	68.1
Fort Lauderdale	71.5	65.9	56.2	66.1	71.8	58.4	71.5
Houston	77.6	76.0	81.3	88.0	75.4	67.3	82.1
Los Angeles	63.9	63.7	60.8	65.1	65.1	55.4	65.4
Miami	63.7	80.3	61.0	64.1	62.8	59.5	77.4
Orange County	50.3	35.2	41.0	58.3	59.5	34.8	58.9
Palm Beach	65.2	47.1	57.3	53.5	62.3	55.9	62.3
San Diego	43.8	50.0	43.8	66.7	66.7	22.9	52.1
San Francisco	63.9	61.6	61.6	62.0	74.1	47.1	65.1
<b>Unweighted Data</b>							
District of Columbia	72.4	72.4	73.3	86.7	73.3	62.1	74.2
Philadelphia	65.4	79.5	65.9	75.9	70.7	52.4	72.3
San Bernardino	71.4	71.4	71.4	62.5	87.5	57.1	62.5
<b>Local Median</b>	<b>65.2</b>	<b>71.4</b>	<b>61.6</b>	<b>66.1</b>	<b>68.5</b>	<b>55.9</b>	<b>66.6</b>

\* HIV = human immunodeficiency virus.

\*\* Survey did not include students from one of the state's large school districts.

TABLE 15c. Percentage of Schools in Which the Lead Health Education Teacher Wanted Staff Development, by Specific Health Education Topic, Selected U.S. Sites: School Health Profiles, Teachers' Surveys, 2002

**STATE SURVEYS**

Site	Personal hygiene	Physical activity and fitness	Pregnancy prevention	STD* prevention	Suicide prevention	Sun safety	Tobacco-use prevention	Violence prevention
<b>Weighted Data</b>								
Alabama	49.8	68.9	62.7	70.7	75.4	58.7	69.6	80.3
Alaska	38.5	54.1	46.7	52.5	72.3	38.3	60.7	73.2
Arizona	51.6	60.7	53.9	56.9	70.0	59.8	65.8	77.2
Connecticut	36.9	56.9	64.7	73.4	81.8	48.8	63.0	79.9
Delaware	41.5	70.6	64.2	70.9	74.4	45.4	72.5	81.2
Georgia	43.6	65.3	64.5	70.6	70.7	57.1	63.1	77.2
Hawaii	49.0	65.3	65.9	68.2	84.8	66.2	62.5	78.8
Idaho	34.6	51.9	53.3	60.6	72.7	42.0	60.9	80.3
Illinois**	30.3	49.7	47.9	56.5	67.3	39.8	55.3	70.7
Iowa	30.8	50.5	46.9	54.7	63.9	44.5	50.5	67.8
Kentucky	43.8	59.6	56.6	60.3	72.0	47.9	61.4	76.9
Maine	25.3	48.7	50.3	54.4	65.8	40.6	54.9	68.9
Massachusetts	41.2	59.9	62.2	69.7	78.3	56.1	60.2	80.9
Michigan	37.6	50.0	52.2	58.4	65.9	45.0	57.9	68.9
Minnesota	29.5	54.2	60.2	65.8	71.8	43.4	59.7	78.2
Missouri	40.8	57.9	54.8	60.4	66.1	47.7	56.6	72.2
Montana	41.3	66.0	54.6	62.8	74.2	53.6	60.4	74.6
Nebraska	26.5	47.0	39.2	45.9	57.9	34.1	50.4	65.6
New Hampshire	40.4	63.3	56.3	65.7	70.4	53.1	59.0	77.8
New Mexico	48.0	66.5	66.6	66.8	75.9	50.6	66.2	78.3
New York	43.9	55.3	71.0	74.7	76.8	54.4	70.2	78.9
North Dakota	35.2	49.3	41.3	54.7	63.4	42.2	48.7	66.0
Tennessee	50.0	68.3	62.0	67.5	76.9	54.6	67.1	83.5
Utah	39.3	57.3	58.8	65.1	80.6	50.8	62.6	87.7
Vermont	29.5	34.9	54.7	53.7	72.7	34.0	50.0	79.2
Virginia	39.7	72.8	50.7	54.0	67.2	53.5	56.1	78.2
Wisconsin	35.4	54.9	55.9	58.8	67.2	40.4	58.7	70.2
<b>Unweighted Data</b>								
Arkansas	50.2	67.4	60.7	67.1	69.8	50.7	65.5	78.9
California	35.5	49.0	45.7	53.3	65.6	44.1	54.8	74.3
North Carolina	40.8	67.7	53.2	60.8	67.4	51.7	65.4	78.4
Oklahoma	45.7	57.0	55.6	57.1	68.3	52.9	67.6	78.3
Pennsylvania**	42.4	74.4	68.0	79.0	79.1	63.4	67.6	82.2
Rhode Island	37.7	63.9	69.4	66.7	70.8	63.9	56.5	85.5
South Carolina	46.6	64.2	61.4	64.0	68.1	58.4	64.9	75.8
Texas**	44.4	61.4	64.0	67.1	71.5	56.9	68.3	75.0
<b>State Median</b>	<b>40.8</b>	<b>59.6</b>	<b>56.3</b>	<b>62.8</b>	<b>70.8</b>	<b>50.7</b>	<b>60.9</b>	<b>78.2</b>

(continued)

TABLE 15c. Percentage of Schools in Which the Lead Health Education Teacher Wanted Staff Development, by Specific Health Education Topic, Selected U.S. Sites: School Health Profiles, Teachers' Surveys, 2002 (continued)

**LOCAL SURVEYS**

Site	Personal hygiene	Physical activity and fitness	Pregnancy prevention	STD* prevention	Suicide prevention	Sun safety	Tobacco-use prevention	Violence prevention
<b>Weighted Data</b>								
Chicago	64.8	66.6	67.2	73.6	73.6	58.8	70.5	86.3
Dallas	45.5	60.0	71.6	69.3	68.5	59.9	58.5	70.7
Fort Lauderdale	53.9	63.7	71.0	68.7	74.2	66.1	61.2	71.7
Houston	67.3	84.0	87.8	87.8	92.0	73.0	81.9	96.0
Los Angeles	47.9	55.1	60.4	64.8	70.4	55.2	58.9	78.0
Miami	53.8	60.6	64.3	65.2	77.0	64.7	69.7	72.5
Orange County	35.6	39.9	62.0	67.2	81.0	56.4	55.7	66.9
Palm Beach	57.2	55.7	56.2	56.2	65.2	61.1	56.2	74.0
San Diego	25.0	33.3	62.5	56.3	75.0	22.9	39.6	83.3
San Francisco	56.5	62.7	65.1	68.6	76.5	49.8	52.5	76.9
<b>Unweighted Data</b>								
District of Columbia	67.7	74.2	74.2	83.9	77.4	61.3	77.4	93.5
Philadelphia	52.4	70.7	69.5	70.4	85.4	56.6	66.7	83.3
San Bernardino	50.0	62.5	75.0	75.0	62.5	42.9	62.5	75.0
<b>Local Median</b>	<b>53.8</b>	<b>62.5</b>	<b>67.2</b>	<b>68.7</b>	<b>75.0</b>	<b>58.8</b>	<b>61.2</b>	<b>76.9</b>

\* STD = sexually transmitted disease.

\*\* Survey did not include students from one of the state's large school districts.

TABLE 16. Percentage of Schools in Which the Lead Health Education Teacher Received Staff Development During the Preceding 2 Years, by Specific Teaching Method, Selected U.S. Sites: School Health Profiles, Teachers' Surveys, 2002

**STATE SURVEYS**

Site	Teaching students with physical or cognitive disabilities	Teaching students of various cultural backgrounds	Teaching students with limited English proficiency	Using interactive teaching methods	Encouraging family or community involvement	Teaching skills for behavior change
<b>Weighted Data</b>						
Alabama	62.5	49.3	30.2	54.3	43.8	53.6
Alaska	37.9	57.0	33.4	55.4	46.2	38.1
Arizona	54.4	54.5	52.8	57.1	48.0	58.0
Connecticut	44.2	34.9	18.2	54.5	29.2	42.1
Delaware	50.9	48.4	17.2	65.3	34.1	61.3
Georgia	37.1	38.7	26.1	58.0	32.6	42.2
Hawaii	32.7	32.7	26.1	70.0	21.9	48.3
Idaho	22.7	20.8	16.0	43.7	20.1	34.6
Illinois*	37.3	28.2	15.8	50.7	30.1	44.8
Iowa	45.7	36.2	18.6	54.7	40.5	46.5
Kentucky	50.6	44.5	16.3	57.7	41.2	52.8
Maine	37.8	15.8	6.9	44.6	25.2	38.9
Massachusetts	47.1	34.9	18.3	61.3	34.5	55.8
Michigan	31.2	25.6	9.8	48.3	32.2	41.7
Minnesota	40.9	39.1	18.8	47.4	32.9	42.3
Missouri	40.4	35.5	15.4	61.1	43.3	47.4
Montana	31.4	21.4	3.5	50.5	31.6	46.0
Nebraska	38.2	37.1	15.3	40.9	31.0	40.9
New Hampshire	52.8	14.4	11.0	58.3	30.9	52.5
New Mexico	36.1	51.7	45.6	55.2	48.8	45.0
New York	39.1	31.5	18.3	54.7	32.5	54.2
North Dakota	37.7	16.1	8.1	42.9	27.8	42.0
Tennessee	46.0	40.8	25.1	55.9	44.5	49.4
Utah	42.3	52.6	46.0	50.7	30.2	46.3
Vermont	46.6	18.5	10.3	61.0	28.0	55.2
Virginia	44.5	41.2	21.8	60.8	34.9	45.0
Wisconsin	40.4	32.9	14.7	54.3	37.5	53.7
<b>Unweighted Data</b>						
Arkansas	43.7	35.2	24.6	51.1	35.1	46.5
California	43.0	54.5	58.8	52.1	32.7	41.0
North Carolina	44.1	45.7	30.0	55.5	28.8	39.9
Oklahoma	50.7	65.3	25.8	57.8	53.7	55.9
Pennsylvania*	51.1	26.7	11.8	56.6	36.9	45.6
Rhode Island	35.4	26.2	16.9	44.6	23.1	29.2
South Carolina	35.3	39.1	20.1	46.7	37.8	38.8
Texas*	54.7	52.9	47.8	60.8	39.4	49.6
<b>State Median</b>	<b>42.3</b>	<b>36.2</b>	<b>18.3</b>	<b>54.7</b>	<b>32.9</b>	<b>46.0</b>

(continued)

TABLE 16. Percentage of Schools in Which the Lead Health Education Teacher Received Staff Development During the Preceding 2 Years, by Specific Teaching Method, Selected U.S. Sites: School Health Profiles, Teachers' Surveys, 2002 (continued)

### LOCAL SURVEYS

Site	Teaching students with physical or cognitive disabilities	Teaching students of various cultural backgrounds	Teaching students with limited English proficiency	Using interactive teaching methods	Encouraging family or community involvement	Teaching skills for behavior change
<b>Weighted Data</b>						
Chicago	54.5	39.5	25.2	62.5	49.6	54.7
Dallas	78.3	57.1	53.1	69.3	38.7	61.9
Fort Lauderdale	39.7	67.5	62.8	65.0	53.8	57.9
Houston	72.5	80.0	88.0	85.7	69.4	77.6
Los Angeles	56.0	72.6	77.2	71.8	44.5	54.7
Miami	27.9	51.2	60.5	63.3	33.4	31.9
Orange County	47.8	66.8	70.4	55.2	36.5	50.4
Palm Beach	62.2	85.8	67.8	76.9	40.2	62.6
San Diego	29.2	47.9	33.3	85.4	50.0	72.9
San Francisco	59.2	76.9	56.1	74.1	52.5	53.3
<b>Unweighted Data</b>						
District of Columbia	54.8	54.8	32.3	80.6	54.8	64.5
Philadelphia	28.9	43.4	19.3	53.6	28.9	37.3
San Bernardino	85.7	85.7	85.7	100.0	71.4	85.7
<b>Local Median</b>	<b>54.8</b>	<b>66.8</b>	<b>60.5</b>	<b>71.8</b>	<b>49.6</b>	<b>57.9</b>

\* Survey did not include students from one of the state's large school districts.

TABLE 17. Percentage of Schools in Which the Lead Health Education Teacher Wanted Staff Development, by Specific Teaching Method, Selected U.S. Sites: School Health Profiles, Teachers' Surveys, 2002

**STATE SURVEYS**

Site	Teaching students with physical or cognitive disabilities	Teaching students of various cultural backgrounds	Teaching students with limited English proficiency	Using interactive teaching methods	Encouraging family or community involvement	Teaching skills for behavior change
<b>Weighted Data</b>						
Alabama	62.2	59.1	56.5	63.2	65.4	71.5
Alaska	60.7	50.5	45.6	57.5	63.8	72.4
Arizona	65.2	64.1	61.0	61.2	67.2	74.4
Connecticut	64.2	63.1	54.6	71.4	70.4	80.8
Delaware	72.6	61.5	53.2	69.4	70.4	79.7
Georgia	66.1	64.4	64.3	64.2	67.0	73.0
Hawaii	72.7	63.6	66.1	69.0	78.1	86.6
Idaho	60.9	53.6	50.6	56.5	65.1	78.6
Illinois*	52.4	40.8	34.8	52.7	58.9	66.0
Iowa	49.9	42.4	38.2	50.0	55.4	66.3
Kentucky	55.5	54.5	52.6	56.3	63.0	71.4
Maine	50.3	39.9	33.7	58.7	68.9	76.8
Massachusetts	69.1	63.0	55.1	70.9	72.3	84.4
Michigan	58.1	48.6	41.5	57.1	61.6	74.7
Minnesota	53.0	49.3	42.2	58.9	62.0	74.6
Missouri	60.2	47.7	39.7	60.3	66.2	74.1
Montana	57.9	43.6	34.8	63.3	66.2	78.2
Nebraska	46.0	39.5	33.1	41.1	53.6	63.6
New Hampshire	63.4	51.0	43.2	73.4	71.5	82.7
New Mexico	66.5	58.0	54.3	62.7	66.8	78.9
New York	67.7	59.3	53.1	72.9	72.6	81.7
North Dakota	48.8	33.7	27.1	51.1	57.1	68.8
Tennessee	68.1	60.3	59.0	63.9	72.4	78.1
Utah	62.7	58.7	61.7	65.9	63.8	80.3
Vermont	51.0	32.6	27.6	55.1	62.7	73.5
Virginia	66.6	51.1	53.3	59.3	64.3	80.1
Wisconsin	55.3	50.3	47.0	60.0	64.2	74.3
<b>Unweighted Data</b>						
Arkansas	56.3	51.7	45.9	58.3	63.5	73.8
California	51.3	50.7	50.5	52.5	56.5	72.3
North Carolina	69.6	62.3	63.8	58.4	60.3	74.8
Oklahoma	68.5	60.9	53.2	62.7	69.9	77.9
Pennsylvania*	68.1	57.1	46.9	64.8	69.9	82.7
Rhode Island	66.7	64.1	54.0	73.0	75.0	87.7
South Carolina	69.3	63.2	59.3	60.4	68.4	73.6
Texas*	62.5	61.3	58.6	61.4	65.9	76.6
<b>State Median</b>	<b>62.5</b>	<b>54.5</b>	<b>52.6</b>	<b>60.4</b>	<b>65.9</b>	<b>74.8</b>

(continued)

TABLE 17. Percentage of Schools in Which the Lead Health Education Teacher Wanted Staff Development, by Specific Teaching Method, Selected U.S. Sites: School Health Profiles, Teachers' Surveys, 2002 (continued)

**LOCAL SURVEYS**

Site	Teaching students with physical or cognitive disabilities	Teaching students of various cultural backgrounds	Teaching students with limited English proficiency	Using interactive teaching methods	Encouraging family or community involvement	Teaching skills for behavior change
<b>Weighted Data</b>						
Chicago	74.5	64.2	58.7	72.4	78.2	86.3
Dallas	58.8	66.0	74.9	59.5	68.9	79.1
Fort Lauderdale	65.8	63.2	58.7	65.4	74.0	72.4
Houston	85.7	75.4	73.4	75.9	81.5	87.7
Los Angeles	69.4	66.1	64.6	72.3	74.6	81.9
Miami	63.5	67.3	62.9	63.9	74.1	76.3
Orange County	43.0	48.0	52.8	66.6	71.7	70.5
Palm Beach	58.3	46.3	53.0	59.4	77.8	74.7
San Diego	56.3	60.4	60.4	66.0	68.8	75.0
San Francisco	66.8	70.0	64.0	49.0	72.9	78.5
<b>Unweighted Data</b>						
District of Columbia	83.3	77.4	80.0	86.7	83.9	100.0
Philadelphia	72.9	72.0	65.4	75.3	79.5	90.4
San Bernardino	62.5	42.9	57.1	71.4	75.0	62.5
<b>Local Median</b>	<b>65.8</b>	<b>66.0</b>	<b>62.9</b>	<b>66.6</b>	<b>74.6</b>	<b>78.5</b>

\* Survey did not include students from one of the state's large school districts.



TABLE 18. Percentage of Schools That Required Physical Education in Grades 6-12 and, Among Those Schools, Percentage That Allowed Students to Be Exempted from Required Physical Education, Selected U.S. Sites: School Health Profiles, Principals' Surveys, 2002

**STATE SURVEYS**

Site	Required physical education	Reason for exemption from required physical education			
		Enrolled in another course*	Participated in school sports	Participated in other school activities**	Participated in community sports activities
<b>Weighted Data</b>					
Alabama	98.4	3.0	20.3	41.7	1.2
Alaska	95.7	6.0	20.2	10.0	7.3
Arizona	75.6	10.5	11.3	17.3	8.2
Arkansas	96.4	3.4	31.5	14.3	2.1
Connecticut	99.0	2.6	3.0	1.7	1.5
Delaware	100.0	2.0	5.8	6.0	2.0
Georgia	77.8	10.6	0.5	9.6	0.0
Hawaii	100.0	5.7	0.0	6.0	0.0
Idaho	92.3	9.4	5.9	6.5	2.9
Illinois <sup>§</sup>	98.5	24.1	29.1	20.2	2.1
Iowa	99.3	41.2	20.7	5.9	0.8
Kentucky	91.5	6.1	0.6	7.3	0.0
Maine	99.5	3.1	2.7	1.0	1.6
Massachusetts	97.2	8.4	2.9	4.2	2.1
Michigan	91.3	12.1	18.8	23.6	2.7
Minnesota	96.7	3.4	3.8	1.6	2.9
Missouri	99.7	2.6	0.6	1.9	0.3
Montana	100.0	0.4	1.3	0.0	0.8
Nebraska	98.7	4.0	2.6	2.5	0.4
New Hampshire	98.3	4.2	6.6	4.5	2.0
New Jersey	99.7	0.3	1.1	1.5	0.8
New Mexico	98.2	5.1	25.3	21.4	3.8
New York	99.4	2.1	7.8	3.1	3.6
North Carolina	97.5	4.9	0.7	4.9	0.4
North Dakota	99.4	4.0	1.2	0.0	0.0
Oklahoma	47.7	27.7	48.0	31.0	4.9
Tennessee	85.8	4.8	4.4	22.6	0.4
Texas <sup>§</sup>	97.7	8.9	81.6	58.5	15.7
Utah	99.5	7.6	15.9	10.8	8.3
Vermont	100.0	2.7	12.5	1.0	3.8
Virginia	95.1	9.9	1.7	6.9	2.2
Wisconsin	97.8	2.0	1.1	0.7	0.7
<b>Unweighted Data</b>					
California	100.0	2.6	18.2	21.5	13.6
Oregon	97.8	9.4	7.7	3.6	8.9
Pennsylvania <sup>§</sup>	99.3	3.1	3.2	2.1	1.4
Rhode Island	98.5	0.0	1.6	1.6	6.3
South Carolina	97.0	8.4	0.4	43.9	0.0
Washington	96.3	23.0	38.4	20.6	22.7
<b>State Median</b>	<b>98.2</b>	<b>4.9</b>	<b>5.1</b>	<b>6.0</b>	<b>2.0</b>

(continued)

TABLE 18. Percentage of Schools That Required Physical Education in Grades 6-12 and, Among Those Schools, Percentage That Allowed Students to Be Exempted from Required Physical Education, Selected U.S. Sites: School Health Profiles, Principals' Surveys, 2002 (continued)

### LOCAL SURVEYS

Site	Required physical education	Reason for exemption from required physical education			
		Enrolled in another course*	Participated in school sports	Participated in other school activities**	Participated in community sports activities
<b>Weighted Data</b>					
Chicago	98.6	2.8	2.9	6.1	1.0
Dallas	92.2	2.4	49.4	72.3	2.3
Fort Lauderdale	79.3	3.3	45.3	6.5	0.0
Houston	100.0	9.6	26.5	45.5	9.3
Los Angeles	100.0	1.0	36.6	55.9	8.8
Miami	60.8	19.9	35.6	15.1	2.0
Orange County	69.0	15.5	58.0	39.0	0.0
Palm Beach	82.3	12.7	50.1	13.5	0.0
San Bernardino	92.9	0.0	0.0	9.1	0.0
San Diego	89.8	2.3	35.7	40.5	26.2
San Francisco	94.0	0.0	9.6	19.2	9.6
<b>Unweighted Data</b>					
District of Columbia	100.0	3.4	0.0	10.3	0.0
Philadelphia	95.4	3.7	2.4	6.1	3.7
<b>Local Median</b>	<b>92.9</b>	<b>3.3</b>	<b>35.6</b>	<b>15.1</b>	<b>2.0</b>

\* Such as math or science.

\*\* Such as ROTC, marching band, chorus, or cheerleading.

§ Survey did not include students from one of the state's large school districts.

TABLE 19. Percentage of Schools That Offered Intramural Activities or Physical Activity Clubs for Students and, Among Those Schools, Percentage That Provided Transportation Home from Intramural Activities, Selected U.S. Sites: School Health Profiles, Principals' Surveys, 2002

**STATE SURVEYS**

Site	Offered intramural activities or physical activity clubs	Provided transportation home*
<b>Weighted Data</b>		
Alabama	48.8	4.8
Alaska	78.7	11.6
Arizona	76.1	53.4
Arkansas	45.7	19.0
Connecticut	85.3	58.2
Delaware	71.7	58.1
Georgia	55.5	15.1
Hawaii	92.7	13.1
Idaho	66.1	11.2
Illinois**	58.0	43.3
Iowa	48.8	35.1
Kentucky	59.0	33.4
Maine	91.7	50.6
Massachusetts	91.6	54.7
Michigan	78.1	17.9
Minnesota	63.3	50.7
Missouri	59.5	33.2
Montana	58.1	9.7
Nebraska	41.7	14.6
New Hampshire	83.1	28.1
New Jersey	90.4	43.4
New Mexico	64.1	33.3
New York	90.3	66.6
North Carolina	65.3	22.2
North Dakota	51.0	25.9
Oklahoma	37.8	17.6
Tennessee	59.8	7.5
Texas**	39.6	38.3
Utah	76.9	31.4
Vermont	83.6	28.5
Virginia	66.4	59.3
Wisconsin	72.8	34.9
<b>Unweighted Data</b>		
California	81.4	29.0
Oregon	65.2	26.5
Pennsylvania**	86.7	50.2
Rhode Island	75.4	70.8
South Carolina	70.7	11.2
Washington	70.6	50.6
<b>State Median</b>	<b>68.5</b>	<b>32.3</b>

(continued)

TABLE 19. Percentage of Schools That Offered Intramural Activities or Physical Activity Clubs for Students and, Among Those Schools, Percentage That Provided Transportation Home from Intramural Activities, Selected U.S. Sites: School Health Profiles, Principals' Surveys, 2002 (continued)

### LOCAL SURVEYS

Site	Offered intramural activities or physical activity clubs	Provided transportation home*
<b>Weighted Data</b>		
Chicago	89.2	26.8
Dallas	78.6	39.7
Fort Lauderdale	90.9	35.6
Houston	87.2	17.6
Los Angeles	92.4	50.5
Miami	93.9	52.3
Orange County	97.1	4.5
Palm Beach	86.6	96.9
San Bernardino	92.9	61.5
San Diego	74.5	66.7
San Francisco	97.5	20.9
<b>Unweighted Data</b>		
District of Columbia	100.0	10.0
Philadelphia	98.9	19.8
<b>Local Median</b>	<b>92.4</b>	<b>35.6</b>

\* For students who participate in after-school intramural activities or physical activity clubs.

\*\* Survey did not include students from one of the state's large school districts.