

**Developing a geographic exposure profile of methylmercury
availability in salt marshes of New England, 2004-05**

(BRI 2006-01)



BioDiversity Research Institute (BRI) is a nonprofit organization located in Gorham, Maine. Founded in 1994, BRI is dedicated to progressive environmental research and education that furthers global sustainability and conservation policies. BRI's research efforts emphasize conservation biology issues in New England and across North America.

To obtain copies of this report contact:

*BioDiversity Research Institute
19 Flaggy Meadow Road
Gorham, ME 04038
(207) 839-7600*

*staff@briloon.org
www.BRIloon.org*

Developing a geographic exposure profile of methylmercury availability in salt marshes of New England

(BRI 2006-01)

Submitted to:

Tom Hodgman, Maine DIFW, Bangor, Maine
Andrew Major, US FWS, Concord, New Hampshire,
Kate O'Brien and Graham Taylor, US FWS, Rachel Carson National Wildlife Refuge,
Wells, Maine

Submitted By:

Oksana P. Lane and David C. Evers
BioDiversity Research Institute, Gorham Maine

January 10, 2006



Please cite this report as: Lane, O.P. and D.C. Evers. 2006. Developing a geographic exposure profile of methylmercury availability in salt marshes of New England. Report BRI 2006-01. BioDiversity Research Institute, Gorham, Maine.

Table of Contents

ABSTRACT	5
INTRODUCTION	5
NATURAL HISTORY	7
METHODS	7
STUDY AREA	7
2004.....	7
2005.....	8
CAPTURE AND SAMPLING	8
STATISTICAL ANALYSES.....	9
RESULTS AND DISCUSSION	9
SAMPLING EFFORT	9
<i>Saltmarsh Sharp-tailed Sparrows</i>	9
2004	9
2005	9
<i>Tree Swallows</i>	9
MORPHOLOGICAL MEASUREMENTS IN SPARROWS 2004.....	11
COMPARISONS AMONG SALT MARSH BIRDS BLOOD MERCURY CONCENTRATIONS	12
<i>Saltmarsh Sharp-tailed Sparrows</i>	12
2004	12
2005	12
All years	13
<i>Other salt marsh birds</i>	14
COMPARISONS BETWEEN TREE SWALLOWS AND SALTMARSH SHARP-TAILED SPARROWS ..	16
<i>Blood mercury</i>	16
2004-2005	16
<i>Egg mercury</i>	18
CONCLUSIONS	20
FUTURE RECOMMENDATIONS:	20
ACKNOWLEDGMENTS	20
LITERATURE CITED	21

Abstract

In 2004-05 we conducted a mercury (Hg) exposure survey of Saltmarsh Sharp-tailed Sparrows (*Ammodramus caudacutus*) nesting in the estuaries at four National Wildlife Refuges across New England and additional sites in Maine. Our goal was to assess methylmercury (MeHg) availability to insectivorous birds. In 2004 we sampled sparrow blood and collected failed eggs from five estuaries of the Rachel Carson National Wildlife Refuge (NWR) in Maine, one site at Parker River NWR in Massachusetts, two sites at Ninigret NWR in Rhode Island, one site at Stewart B. McKinney NWR and a nearby off-refuge site Hammock River in Connecticut, and in Scarborough Marsh State Wildlife Management Area (SWMA) in Maine. In 2005, in addition to sampling Parker River, McKinney and Ninigret NWRs we sampled two marshes at the Rachel Carson NWR and added two new sites in Scarborough, Maine. Concurrent Tree Swallow (*Tachycineta bicolor*) sampling efforts were conducted in 2004 and 2005 at one of the Rachel Carson NWR sites in Wells and at Scarborough Marsh SWMA, and in 2005 we added Parker River NWR to relate established Hg risks in swallows to the sparrows.

In 2004, mean site-specific sparrow blood Hg levels ranged from 0.23 to 1.09 ppm (wet weight). Sparrow blood Hg levels from Parker River NWR and Ninigret NWR were significantly higher than the concentrations in sparrows sampled in Maine and Connecticut.

In 2005, mean refuge site-specific sparrow blood Hg levels ranged from 0.61 ppm (wet wt.) from Spurwink Marsh (RCNWR in Cape Elizabeth) in Maine and McKinney NWR in Connecticut to 1.24 ppm in Parker River NWR in Massachusetts. Sparrow blood Hg concentrations from Parker River NWR were significantly higher than all other sampled sites. Three of the plausible explanations are: (1) there are higher levels of Hg in the Parker River NWR system originating from waterborne and/or airborne sources, (2) Hg is methylated at a higher rate in these marshes (3) sparrows in Parker River may consume prey at a higher trophic level.

In Maine, sparrows from Libby River site in Scarborough had the lowest mean Hg levels of 0.31 ppm and from Furbish Road Marsh in Wells had the highest concentrations of 0.76 ppm.

Blood Hg concentrations in Tree Swallows at all study sites were significantly lower than in Saltmarsh Sharp-tailed Sparrows, although, egg Hg levels tended to be similar. Swallow blood and egg Hg concentrations were significantly correlated. Additional sampling efforts are needed to further document (1) site-specific exposure profiles, (2) origin of Hg input, (3) Hg relationship between swallows and sparrows and (4) Hg effects in sparrows.

Introduction

Sharp-tailed Sparrows have been recently (1995) split into two species: a northern species, (*Ammodramus nelsoni*) with three subspecies (*A. n. nelsoni*, *A. n. alterus*, and *A. n. subvirgatus*) and the southern species, (*Ammodramus caudacutus*) with two subspecies (*A. c. caudacutus* and *A. c. diverus*). *A. n. subvirgatus* (hereafter Nelson's Sparrow) and *A. c. caudacutus* (hereafter Saltmarsh Sparrow) (Figure 1) are sympatric in coastal Maine, New Hampshire and northern Massachusetts (Hodgman et al. 2002, Shriver et al. 2005). Based on a previous study we concluded that Saltmarsh Sparrows were more numerous in the study area and had significantly higher blood mercury concentrations than closely related Nelson's

Sparrow (Shriver et al. 2002). Consequently we focused our efforts on the Saltmarsh Sharp-tailed Sparrow.

Figure 1. Nelson's (left) and Saltmarsh Sharp-tailed Sparrows, Scarborough Marsh SWMA, Maine, 2004.



Exposure to mercury can impact bird behavior, physiology, and reproductive success (Thompson 1996, Wolfe and Norman 1998). Extensive studies with Common Loons have documented these impacts in Maine (Evers et al. 2004). It is well known that freshwater wetlands generally serve as areas of high Hg methylation, thus making obligate birds especially vulnerable to high levels of Hg contamination (Evers et al. 2005). The role of salt marsh habitats in methylating Hg and enhancing its bioavailability (Marvin-DiPasquale et al. 2003), however, is less documented, but is of increasing concern especially in urban areas.

Sharp-tailed Sparrows are obligate salt marsh passerines with $\approx 95\%$ of their global population breeding within the Northeast. Spending their entire annual cycle in salt marsh habitats makes them excellent indicators of Hg contamination. According to Partners in Flight, Saltmarsh Sharp-tailed Sparrows are among the highest priority species in the northeast, and considered a “bird of conservation concern” by the US Fish and Wildlife Service (USFWS) Region 5. These designations result from their near endemic status in our region, a lack of population trend data, and threats on their breeding and wintering grounds (Hodgman pers. com.). Saltmarsh Sparrows spend their entire lifecycle within the salt marsh ecosystem. An estimated 98 percent of their global population nests in USFWS Region 5.

Between 1998-2000, we found 5 sparrows with beak and/or foot deformities at Scarborough Marsh SWMA, and when we later analyzed the blood of Saltmarsh Sparrows from that site, we found elevated levels of Hg. In 2001, we expanded sampling to include blood from 75 sparrows from five Maine estuaries to determine relative differences in Hg

levels. Sparrow blood samples from estuarine habitat in Rachel Carson NWR in Wells contained some of the highest Hg levels in Maine (Shriver et al 2002).

Concurrently, work in Maine and Massachusetts has shown that Tree Swallows may be appropriate indicators of Hg exposure in wetlands. We included swallows in the study to compare Hg levels in the Saltmarsh Sharp-tailed Sparrows with the established risk concentrations from other studies conducted with Tree Swallows (Heinz, pers. comm). Because recent dosing studies have now identified lethal effects at 0.8 ppm (wet weight=ww), and lowest observed adverse effect levels (LOAEL) for Tree Swallow eggs at 0.4 ppm (ww), linking Hg exposure levels in Tree Swallows with those in Sharp-tailed Sparrows is of great interest. Tree Swallows nest in artificial boxes and can be easily attracted to breed on the targeted salt marshes.

Our objectives include:

1. Determine comparative exposure profile for Saltmarsh Sparrows on four National Wildlife Refuges across New England and the Scarborough marsh area in Maine.
2. Further understand the relationship of (a) Tree Swallow blood and egg levels and (b) swallow-sparrow Hg exposure.
3. Identify areas with lowest and highest exposure for focused efforts in 2006 to measure potential Hg effects on Saltmarsh Sparrows from a high risk population.

Natural history

The Saltmarsh Sharp-tailed Sparrow has a limited range, occupying estuaries along the Atlantic Coast from Florida up to the southern coast of Maine where it overlaps with the Nelson's Sharp-tailed Sparrow (Hodgman et al. 2002). Across its range the species is non-territorial and exhibits a bondless form of polygyny in which males provide no parental care (Greenlaw 1993, Greenlaw and Rising 1994).

The diet during the breeding season consists mainly of animal matter: immature and adult insects and other arthropods (Greenlaw and Rising 1994, Merriam 1979). Tidal flooding causes most of nest failures but usually the birds renest within a short period of time. Young leave the nest 23-27 days after clutch initiation. Nestlings fledge between 8-11 days after hatching (Greenlaw and Rising 1994).

Because Saltmarsh Sparrows require nesting habitat that is becoming increasingly limited due to habitat fragmentation, rise in sea level, pollution and human encroachment into estuaries, the species' conservation calls for immediate attention.

Methods

Study area

2004

The study encompassed six estuaries in Maine (Appendix I and II): 1) Scarborough Marsh State Wildlife Management Area (hereafter Scarborough Marsh SWMA) in Scarborough and 2) five estuaries in the Rachel Carson NWR: Furbish Road Marsh (hereafter Furbish Marsh) in Wells, Little River Marsh (hereafter Little River) in Wells and Kennebunk, Granite Point Marsh (hereafter Granite Point) in Biddeford, Goosefare Brook Marsh (hereafter Goosefare) in Old Orchard Beach and Saco, and Spurwink Marsh on Cape Elizabeth (hereafter Spurwink). We sampled one site in Massachusetts: Parker River NWR

(hereafter Parker River) on Plum Island in Newburyport, two sites in Rhode Island: Ninigret NWR, Chaffee and Sachuest Marshes, and two sites in Connecticut: Stewart B. McKinney NWR-Great Meadows salt marsh (hereafter McKinney) in Stratford and Hammock River Marsh (hereafter Hammock River) in Clinton (this site is a part of a University of Connecticut study area) (Appendix I). Saltmarsh Sparrow blood samples were collected from all of the sites listed above. Tree Swallows were sampled from Scarborough Marsh SWMA and Furbish Marsh in Maine.

2005

The study encompassed four estuaries in Maine: 1) Libby River and Nonesuch River estuaries in Scarborough and 2) two estuaries in Rachel Carson NWR: Furbish Marsh in Wells and Spurwink Marsh on Cape Elizabeth. We sampled the same site as in 2004 in Parker River, one site in Ninigret (Chafee Marsh) and one site in McKinney (Salt Meadow Unit in Westbrook).

We sampled sparrows in Furbish Marsh (Wells) in June and again in August to determine if Hg levels in the sparrows change from early to late summer. Tree Swallows were sampled from Scarborough Marsh SWMA and Furbish Marsh in Maine and Parker River NWR in Massachusetts.

Capture and sampling

All capture and blood sampling occurred in the summers of 2004 and 2005. We used two to six 12-m mist nets with 36 mm mesh size. We positioned the nets perpendicular to drainage ditches and tidal creeks, regularly found in many of the estuaries. A team of 5-6 people “swept” the area of approximately 100-200 m² “rounding up” the sparrows and coaxing them to fly towards the nets. The birds were extracted from the nets and banded with a USFWS band. A beach umbrella was used for shade to prevent birds from overheating. Sex, age and breeding status were determined for each bird. Venipuncture of the cutaneous ulnar vein with a 26 gauge sterile disposable needle allowed collection of 1-2 capillary tubes of blood into heparinized tubes for Hg analysis. The capillary tubes were sealed with critoseal or Critocaps® and stored in 10 cc plastic vacutainer, labeled with date, site, species, age and sex information. All birds were released unharmed within 10-20 minutes of capture.

We placed 14 nest boxes in Scarborough Marsh SWMA and 16 boxes in Furbish Marsh, to attract Tree Swallows. In 2005 we placed additional 20 boxes in Parker River NWR on Plum Island in Massachusetts. We collected one egg from each occupied box (unless the nest failed then we collected all eggs in the nest) and blood samples from nesting adult swallows for Hg analysis.

We opportunistically collected sparrow eggs and dead chicks from nests flooded during the high tide floods. All blood and 2004 egg samples were analyzed for total Hg at Texas A&M University, Trace Element Research Lab. All analyses are for total Hg because 95% of the total Hg is methylmercury in songbird blood (Rimmer et al. 2005). All blood and egg Hg concentrations are in parts per million (ppm) wet weight (ww). Feathers, dead chicks and 2005 eggs were archived at BioDiversity Research Institute (BRI) for future testing if funding is available.

Statistical analyses

All statistical analyses were conducted using JMP 4.0 software with $\alpha=0.05$. We used one-way Analysis of Variance (ANOVA) and Tukey-Kramer HSD pairwise comparisons tests to determine significant differences among sites and between species. All means are reported as arithmetic means unless otherwise stated. When conditions of normal distribution of data were not met and/or variances were unequal a non-parametric Welch ANOVA and Wilcoxon/Kruskal-Wallis Tests were used.

Results and Discussion

Sampling effort

Saltmarsh Sharp-tailed Sparrows

2004

From 11 June to 28 July, we collected blood samples for Hg analysis from 63 Saltmarsh Sharp-tailed Sparrows in Maine and 42 from other states (Table 1, Appendix I). Blood samples were also opportunistically collected from 55 other individual birds representing 10 species in our study sites. Twenty three abandoned sparrow eggs from several marshes in Maine and New Hampshire were also collected and tested for Hg. The highest density of sparrows within our study appeared to be at Scarborough Marsh SWMA in Maine, followed by Furbish Marsh and Goosefare of the Rachel Carson NWR and the lowest appeared to be at the Chafee site at Ninigret NWR, Rhode Island and Parker River NWR in Massachusetts. The last two locations were the most difficult, as they required more effort to catch our target sample size of sparrows.

We analyzed 20 (8 from Scarborough and 12 from Wells) Tree Swallow blood samples and 21 (6 from Scarborough and 15 from Wells) eggs. Nest box occupancy by Tree Swallows was 50% (7/14) in Scarborough Marsh SWMA and 56% (9/16) in Furbish Marsh. Typically, the nest box occupancy rate by Tree Swallows increases in the years following the installation season.

2005

We sampled a total of 90 adult (Table 1) and 16 juvenile (hatch year) Saltmarsh Sparrows from all sites combined. In Maine we collected samples from 50 adult and 12 juvenile Saltmarsh Sparrows. We have captured and sampled significantly greater (ChiSquare analysis, $p<0.0001$) number of male sparrows than female (124 and 70 respectively) in the two years of our investigation. Blood samples were also opportunistically collected from 18 adult and 2 juvenile birds representing three species in our study sites.

Tree Swallows

Tree Swallow nest box occupancy rates in Maine were higher in 2005 than in 2004. In Scarborough, 13/14 boxes contained eggs (93% occupancy); in Furbish Marsh 16/16 boxes were occupied (100% occupancy rate); and on our new site in Parker River 9/20 boxes had eggs (45% occupancy rate).

Table 1. Summary of adult (AHY=after hatch year=adult) bird blood sampling efforts from 13 New England salt marshes, June-August 2004-05.

Location	Site	Species	2004		2005										
			Sex M F	Total AHY	Sex M F	Total AHY									
Rachel Carson NWR, ME	Furbish Marsh	Saltmarsh Sharp-tailed Sparrow	11	4	15	12	9	21							
		Nelson's Sharp-tailed Sparrow	3	3	6				1	1	2				
	Goosefare Brook	Saltmarsh Sharp-tailed Sparrow	11	2	13										
		Nelson's Sharp-tailed Sparrow	5	2	7										
		Hybrid Saltmarsh/Nelson's	1		1										
	Granite Point	Saltmarsh Sharp-tailed Sparrow	1	2	3										
		Nelson's Sharp-tailed Sparrow	1		1										
		Hybrid Saltmarsh/Nelson's	1		1										
	Little River- Wells	Saltmarsh Sharp-tailed Sparrow		6	1							7			
			Nelson's Sharp-tailed Sparrow	2	1							3			
		Hybrid Saltmarsh/Nelson's	2		2										
		Least Sandpiper	2		2										
		Semipalmated Sandpiper	5		5										
		Spurwink Marsh	Saltmarsh Sharp-tailed Sparrow	6	4							10	10	3	13
			Nelson's Sharp-tailed Sparrow	5	1							6	5	2	7
	American Goldfinch		1	1	2										
	Barn Swallow				2										
Bobolink	1			1											
Chipping Sparrow					1		1								
Least Sandpiper					unknown		1								
Total for Rachel Carson NWR, ME	Saltmarsh Sharp-tailed Sparrow	35	13	48	22	12	34								
	<i>All other species</i>			39	6	4	11								
Scarborough, ME	Scarborough Marsh WMA	Saltmarsh Sharp-tailed Sparrow	7	8	15										
		Nelson's Sharp-tailed Sparrow	4	1	5										
	Libby River	Saltmarsh Sharp-tailed Sparrow				5	2	7							
		Nelson's Sharp-tailed Sparrow				6	1	7							
	Nonesuch River	Saltmarsh Sharp-tailed Sparrow					2	2							
Total for Scarborough, ME	Saltmarsh Sharp-tailed Sparrow	7	8	15	11	5	16								
	<i>All other species</i>			5	6	1	7								
Total for Parker River NWR, MA	Plum Island	Saltmarsh Sharp-tailed Sparrow	6	4	10	12	3	15							
Ninigret NWR, RI	Chafee Marsh	Saltmarsh Sharp-tailed Sparrow	3	3	6	11	4	15							
	Sachuest Marsh	Saltmarsh Sharp-tailed Sparrow	5	4	9										
		Song Sparrow	1		1										
Total for Ninigret NWR, RI	Saltmarsh Sharp-tailed Sparrow	8	7	15	11	4	15								
	<i>All other species</i>			1											
McKinney NWR, CT	Great Meadows	Saltmarsh Sharp-tailed Sparrow	9	6	15										
		Seaside Sparrow	2	1	3										
		Marsh Wren	1	1	2										
		Common Yellowthroat	1		1										
		Bobolink	1		1										
	Salt Meadows	Saltmarsh Sharp-tailed Sparrow				6	4	10							
	Total for McKinney NWR, CT	Saltmarsh Sharp-tailed Sparrow	9	6	15	6	4	10							
<i>All other species</i>				7											
Hammock River, CT		Saltmarsh Sharp-tailed Sparrow	2		2										
		Seaside Sparrow	3		3										
		Saltmarsh Sharp-tailed Sparrow	2		2										
		<i>All other species</i>	3		3										
Total for Hammock River, CT	Saltmarsh Sharp-tailed Sparrow	67	38	105	62	28	90								
	<i>All other species</i>			55			18								
Grand Total															

We collected 16 Tree Swallow eggs from Scarborough Marsh, 19 eggs from Furbish Marsh, and 21 eggs from Parker River. Due to the lack of funding we have analyzed only 10 eggs from 2005 (Hg analysis pending) and we only have limited productivity data from 2005: In Scarborough Marsh, a total of 69 eggs were laid (13 nests), in Furbish Marsh 82 eggs were laid (16 nests) and in Parker River 48 eggs were laid (9 nests), making the average clutch size 5.3, 5.1, and 5.3 eggs respectively (Table 2).

In Parker River out of 40 available eggs (8 were collected for Hg analysis) 22 eggs hatched (55%) and 18 chicks fledged.

Table 2. Summary of Tree Swallow reproductive effort in Maine and Massachusetts study sites, 2005.

	Scarborough Marsh, Maine	Rachel Carson NWR-Furbish Marsh, Maine	Parker River NWR Massachusetts
# Nests/ # nest boxes	13/14	16/16	9/20
# Eggs laid	69	82	48 ²
# Eggs (viable) collected	13	10	8
# Eggs available for hatching ¹	56	72	40
Average clutch size	5.3	5.1	5.3
# Eggs hatched	50 ²	61 ²	22
Hatch success	89%	85%	55%
# Nestlings fledged	unknown	59 ²	18

¹ one egg was collected from most of the nests and it was subtracted from total clutch size

² estimated, visits to the sites were infrequent

Morphological measurements in sparrows 2004

We conducted a oneway analysis of variance (ANOVA) of bird weight in Maine, where both Nelson's and Saltmarsh Sparrows are sympatric, and found that Saltmarsh Sparrows' (20.3 +/- 0.2 g, n=62) mean weight was significantly greater than Nelson's (17.3 +/- 0.3 g, n=28) (ANOVA df=89, F=93, P<0.0001), which is consistent with the findings from previous studies (Shriver, et al. 2005).

We also found that culmen length (from end of nares to tip of the bill) (ANOVA df=84, F=44, P<0.0001) and total bill length (df=86, F=33, P<0.003) were significantly longer in Saltmarsh than Nelson's Sparrow. We compared other bill measurements (bill width and depth) and found no statistically significant differences between species (width: df=83, F=0.25, P<0.6; depth: df=83, F=1.06, P<0.3) (Table 3). Shriver (2002) found that Saltmarsh Sparrows' bill was significantly wider than Nelson's, however unlike Shriver's results, our data analysis does not indicate significant bill width differences between species.

Table 3. Morphological variables (bill measurements in mm, weight in g) measured on sympatric Saltmarsh and Nelson's Sharp-tailed Sparrows in Maine, 2004 (n=birds measured).

Species	Length	Bill mean +/-sd			Weight +/-sd
		Culmen	Width	Depth	
Nelson's	13.1 +/-0.1 (27)	8.8 +/-0.4 (26)	4.3 +/-0.05 (26)	5.35 +/-0.2 (26)	17.3 +/-0.3 (28)
Saltmarsh	13.5 +/-0.1 (60)	9.4 +/-0.4 (59)	4.3 +/-0.04 (59)	5.29 +/-0.3 (59)	20.3 +/-0.2 (62)

Male Sharp-tailed Sparrows tend to have longer wing chords than females (Greenlaw 1993). We found that wing chord was significantly longer in male Nelson's and Saltmarsh

Sparrows than in females (Table 4) (ANOVA, df=28, F=19, P<0.0002; df=102, F=92, P<0.0001 respectively).

Table 4. Mean wing chord measurements (mm) in Saltmarsh and Nelson’s Sharp-tailed Sparrows in New England, 2004 (n=birds measured).

Species	Wing chord mean +/-sd	
	Male	Female
Nelson’s	57.8 +/-0.4 (21)	54.6 +/-0.6 (8)
Saltmarsh	58.0 +/-1.4 (65)	55.4 +/-1.1 (38)

Comparisons among salt marsh birds blood mercury concentrations

Saltmarsh Sharp-tailed Sparrows

2004

We found the lowest Hg concentrations in the sparrows from Hammock River in Connecticut and the highest in Parker River NWR, Massachusetts (Table 5, Figure 2). Using oneway ANOVA we detected statistically significant differences in blood Hg among sites (df=103, F=17.6, P<0.0001). Tukey-Kramer HSD pairwise comparisons test indicates significant site differences and showed that Chafee Marsh of Ninigret NWR and Parker River NWR sparrows had significantly higher blood Hg levels than sparrows from other sites. Sparrows from Sachuest site in Ninigret NWR and Little River in Wells have significantly higher Hg concentrations than Hammock River, Scarborough Marsh WMA and Spurwink.

Table 5. Mean blood Hg concentrations in adult Saltmarsh Sharp-tailed Sparrows, sampled across New England, 2004. (Sites arranged in increasing order of Hg).

State	Site	Mean Hg (ppm, ww)	SD	Hg range (ppm, ww)	n
CT	Hammock River	0.23	0.06	0.18-0.24	6
ME	Rachel Carson NWR - Spurwink Marsh	0.45	0.10	0.26-0.60	10
ME	Scarborough Marsh SWMA	0.47	0.16	0.23-0.82	15
ME	Rachel Carson NWR - Goosefare Brook	0.50	0.12	0.32-0.75	13
CT	McKinney NWR	0.54	0.11	0.39-0.73	15
ME	Rachel Carson NWR - Granite Point	0.54	0.11	0.46-0.66	3
ME	Rachel Carson NWR - Furbish Marsh	0.56	0.09	0.33-0.69	14
RI	Ninigret NWR – Sachuest Marsh	0.72	0.11	0.54-0.87	9
ME	Rachel Carson NWR - Little River	0.74	0.08	0.64-0.84	7
RI	Ninigret NWR - Chafee	1.08	0.22	0.86-1.36	6
MA	Parker River NWR	1.09	0.38	0.67-1.68	10

Using oneway ANOVA we detected statistically significant differences in blood Hg among sites (df=103, F=17.6, P<0.0001). Tukey-Kramer HSD pairwise comparisons test indicates significant site differences and showed that Chafee Marsh of Ninigret NWR and Parker River NWR sparrows had significantly higher blood Hg levels than sparrows from other sites. Sparrows from Sachuest site in Ninigret NWR and Little River in Wells have significantly higher Hg concentrations than Hammock River, Scarborough Marsh WMA and Spurwink.

2005

Among the refuges, the lowest Hg concentrations were detected in the sparrows from McKinney NWR in Westbrook, Connecticut and RCNWR-Spurwink in Maine, and the highest, as in 2004, in Parker River NWR, Massachusetts (Table 6).


Table 6. Mean blood Hg concentrations in adult Saltmarsh Sharp-tailed Sparrows, sampled across four NWR systems in New England and additional estuaries in Maine, 2005. (Sites arranged in increasing order of Hg).

State	Site	Mean Hg (ppm, ww)	SD	Hg range (ppm, ww)	n
ME	Libby River-Scarborough	0.31	0.06	0.28-0.42	7
ME	Nonesuch River-Scarborough	0.45	0.09	0.39-0.52	2
CT	McKinney NWR-Salt Meadow Unit	0.61	0.14	0.44-0.96	10
ME	Rachel Carson NWR - Spurwink Marsh	0.61	0.14	0.40-0.87	13
ME	Rachel Carson NWR - Furbish Marsh	0.76	0.20	0.47-1.44	21
RI	Ninigret NWR – Chafee Marsh	0.79	0.19	0.41-1.17	15
MA	Parker River NWR	1.24	0.38	0.81-2.22	15

Both years

We found a general increasing trend in blood Hg concentrations between 2004 and 2005 from the same site (Figure 2) with the exception of Chafee Marsh in Rhode Island. Sparrows from both sites in Maine had significantly higher Hg levels in 2005 than 2004 ($p < 0.005$) and sparrows from Chafee Marsh had significantly lower Hg levels in 2005 than 2004 ($p < 0.005$). Parker River sparrows had higher mean blood Hg levels in 2005 but not significantly ($p = 0.3$). There might be several explanations for the increasing Hg levels. First, we captured and sampled sparrows later in the summer in 2005 than in 2004. Mercury methylation rates increase with temperature (Driscoll et al. in press) and temperatures increase as the summer progresses. Therefore, the prey items in the salt marshes may have bioaccumulated higher Hg concentrations later in the summer than during the first half of the summer. Second, 2005 might have been a hotter year than 2004, therefore contributing to higher Hg levels in the biota, and third, Hg might be increasing in the estuarine environment on an annual basis perhaps from atmospheric and point sources.


Figure 2. Mean blood Hg concentrations in Saltmarsh Sharp-tailed Sparrows sampled on consecutive years in New England.



We conducted statistical analyses on the pooled data that include 2004 and 2005 results.

Saltmarsh Sparrows from Parker River NWR have blood Hg levels that are significantly higher than sparrows from all other sites (Tukey-Kramer HSD test) (Figure 3).

Figure 3. Mean blood Hg levels in Saltmarsh Sparrows in 2004 and 2005. Lines indicate statistically similar Hg levels. Yellow bars are Maine sites.



One likely reason for high Hg levels in sparrows from the Parker River NWR is that sampling efforts were concentrated in the salt marsh situated between the Merrimack and Parker Rivers (Appendix 3). Both potentially carry Hg-polluted waters from interior watersheds to the coast. The Merrimack River, flowing through New Hampshire and Massachusetts is well known as a historical source of Hg. Mercury that has been deposited in the sediment is likely still present and may continue to methylate and enter the aquatic food chain. Parker River NWR is located in the northeastern region of the state that is also a well-known biological hotspot for Hg (Evers et al. 2005).

Other salt marsh birds

We opportunistically sampled Seaside Sparrows (*Ammodramus maritimus*) in Connecticut and found their blood Hg levels to be similar to Saltmarsh Sparrow levels from the same sites (Hammock River: 0.25 +/-0.11 (n=3) and 0.23 +/-0.06 (n=6) respectively; McKinney NWR-Great Meadows: 0.56 +/-0.14 (n=3) and 0.54 +/-0.11 (n=15)).

Nelson's Sparrow blood Hg levels are known to be lower than the sympatric Saltmarsh Sparrows (Shriver et al. 2002) and we documented similar patterns with this study (Table 5, Figure 4). We found (ANOVA F=21, df=155, p<0.001) that Maine mean Saltmarsh Sparrow blood Hg concentrations (0.57 +/-0.19 ppm, wet wt., n=104) were significantly higher than


Nelson's (0.43 +/-0.16 ppm, n=52). However, when we compared means within a site, only Spurwink Saltmarsh Sparrows had significantly higher blood Hg than Nelson's sparrows (p<0.006). Furbish, Little River and Scarborough SWMA Saltmarsh Sparrow Hg levels were marginally significantly higher than Nelson's (p<0.08, p<0.09, and p<0.1 respectively).

We also collected blood samples from other species opportunistically caught during mist-netting efforts. Saltmarsh Sparrow Blood Hg concentrations tended to be higher than the other 10 species tested (Table 7).

It is likely that Saltmarsh Sparrows had significantly higher blood Hg levels (Shriver et al. 2002) because they feed at a higher trophic level or consume different prey base than sympatric Nelson's Sparrows. For example, based on limited stomach analyses data, the diet of Saltmarsh Sparrows (from New England) had 24 % amphipods by volume during breeding season and Nelson's (in New Brunswick) had higher insect percent by volume but no amphipods in their diet (U.S. Fish and Wildlife Service data, in Greenlaw and Rising 1994). George et al. (2001) found that amphipods contained higher concentrations of Hg than other organisms found higher on the food chain, such as odonates and crayfish. Contaminants collect in "modern mud" (i.e., mud buildup over the last century) so bottom-dwelling animals that dwell in mud habitats tend to accumulate contaminants.

Shriver (2002) found that in the areas of overlap the Saltmarsh Sharp-tailed Sparrows had 25 % of their alleles from the Nelson's Sparrow. Shriver also found that "hybrids" resembled Saltmarsh Sparrows closer than the Nelson's. When comparing Maine Sharp-tailed Sparrows to the other states where there is no hybridization with Nelson's the genetic difference should be considered.

Figure 4. Sympatric species of Sharp-tailed Sparrows sampled across several estuaries in Maine, 2001-2005, NSTS=Nelson's Sharp-tailed Sparrow, SSTS=Saltmarsh Sharp-tailed Sparrow, RCNWR=Rachel Carson National Wildlife Refuge.



There are several factors that may explain differences in interspecies Hg levels: As body mass increases so does MeHg concentration in insectivorous birds (Evers et al. 2005). This

finding holds true for several species in this study with the exception of Seaside Sparrow, which has body mass greater (by ~ 1 gram) than the Saltmarsh Sparrow but Hg levels are similar. The blood Hg levels in the sandpipers (also heavier than sparrows) were lower but they likely reflect multiple feeding sites because of migration. Food chain length also affects the amount of MeHg available to insectivorous birds. It is likely that Saltmarsh Sparrows are feeding at a higher trophic level than other birds in the estuary. Species such as the Bobolink and Common Yellowthroat, had comparatively lower blood Hg levels than Saltmarsh Sparrows within a study site, which indicates a diet with a shorter food chain length and less of an ability for MeHg biomagnification.

Table 7. Mean blood mercury concentrations (ppm, wet wt.) in species opportunistically sampled in 2004 and 2005. Sites are arranged in geographical order-North to South.

State	Site	Species	Age	n	mean Hg +/- st dev 2004	2005
ME	Scarborough Marsh SWMA	Nelson's Sparrow	adult	5	0.34 +/-0.07	
ME	RCNWR - Spurwink Marsh	American Goldfinch	adult	1	0.01	
		Bobolink	adult	1	0.02	
		Barn Swallow	juvenile	2	0.14 +/-0.04	
		Chipping Sparrow	adult	1		0.31
		Least Sandpiper	juvenile	1		0.21
		Nelson's Sparrow	adult	6/7	0.29 +/-0.10	0.47+/-0.15
ME	RCNWR - Granite Point	Nelson's Sparrow	adult	1	0.51	
		Hybrid Sharp-tailed Sp.	adult	1	0.44	
ME	RCNWR - Goosefare Brook	Nelson's Sparrow	adult	7	0.51 +/-0.10	
		Hybrid Sharp-tailed Sp.	adult	1	0.44	
ME	RCNWR - Little River	Least Sandpiper	adult	2	0.15 +/-0.00	
		Semipalmated Sandpiper	adult	5	0.38 +/-0.19	
		Nelson's Sparrow	adult	3	0.60 +/-0.17	
ME	RCNWR - Furbish Marsh	Least Sandpiper	juvenile	2		0.39+/-0.36
		Nelson's Sparrow	adult	6/2	0.47 +/-0.18	0.73+/-0.07
RI	Ninigret NWR-Sachuest Marsh	Song Sparrow	adult	1	0.12	
CT	McKinney NWR-Great Meadows Unit	Common Yellowthroat	adult	1	0.05	
		Bobolink	adult	1	0.05	
		Marsh Wren	adult	2	0.24 +/-0.00	
		Seaside Sparrow	adult	3	0.56 +/-0.14	
CT	Hammock River	Seaside Sparrow	adult	3	0.26 +/-0.11	

Comparisons between Tree Swallows and Saltmarsh Sharp-tailed Sparrows

Blood mercury

2004-2005


Tree Swallow blood Hg levels were significantly higher in Furbish Marsh in Wells than in Scarborough Marsh ($p < 0.005$) when we combined 2004 and 2005 data or when we analyzed 2005 only ($p < 0.001$). There was no significant difference in Hg concentrations between the sites in 2004, however, swallows at Furbish site had higher mean blood Hg levels than in Scarborough. There was no significant difference between Parker River swallows and two sites in Maine ($p > 0.1$). We used a mean of two Hg concentrations when including re-trapped (within a year) individuals.

Mercury levels in Saltmarsh Sparrow blood were significantly higher than in Tree Swallows from all three sites (one-way ANOVA: Furbish Marsh: $df=59$, $F=87$, $p<0.0001$; Scarborough SWMA: $df=35$, $F=44$, $p<0.0001$ and Parker River NWR: $df=35$, $F=66$, $p<0.0001$ (Figure 5).

The mean blood Hg levels are approximately 60% higher in Saltmarsh Sparrows versus Tree Swallows in both Maine sites and 80% higher in Parker River. The difference in blood Hg levels between the species is likely a reflection of their foraging habits. Tree Swallows are likely feeding on flying insects, prey that are lower on the food chain therefore have lower Hg concentrations. In other studies flies (Diptera) made up about 70% of adult Tree Swallows' diet; egg-laying females also consumed mayflies (15%) and females during nestling stage preyed on Odonata (Dragonflies and Damselflies 10%) and a variety of small terrestrial prey (Blancher and McNicol 1991). Food of aquatic origin constituted 65% of the nestling diet by mass in Ontario (Blancher and McNicol 1991). During the breeding season in Ontario Tree Swallows consumed small insects (<1 cm): mostly adult flies (order Diptera) and small leafhoppers (order Homoptera) (Quinney and Ankney 1985).

In a study conducted in New York State, McCarty and Winkler (1999) found at least 11 orders of insects in the diet of Tree Swallow nestlings, with insects in the 3-5 mm range making up the largest proportion of the diet. Diptera (Nematocera and Brachycera) were the most frequent items followed by Hemiptera and Odonata (McCarty and Winkler 1999). Adult Tree Swallows with damselflies (Odonata) in their bills were observed feeding the nestlings on the Sudbury River in Massachusetts (Lane, pers. obser. 2003). We collected one bolus of food in Scarborough Marsh while trapping the female and most of the items were small adult dipterans (flies).

Figure 5. Blood mercury concentrations in Tree Swallows and Saltmarsh Sharp-tailed Sparrows sampled in the summers of 2004-05 in Maine and Massachusetts.



*TRES=Tree Swallow, SSTS=Saltmarsh Sharp-tailed Sparrow

There may be a significant difference in Hg levels between larval and adult stages of many insects. Elwood et al. (1976) demonstrated that gut contents of crane flies could be a significant fraction of Hg in the organism. When most insects and other invertebrates emerge, they molt their final exoskeleton, which includes the gut lining and the guts, leaving a significant portion of the Hg behind (Hildebrand et al. 1980). Sarica et al. (2005) showed that blowfly maggots accumulated Hg when feeding on fish carcasses and took that Hg with them

into the pupal stage, but then eliminated it when they emerged as adults. In addition, many adult insects do not feed and therefore lack a functional gut and consequently have lower Hg levels than the larval forms.

We speculate that prey items in Tree Swallow diet likely contain less Hg than the prey consumed by Saltmarsh Sparrows. Tree Swallows' estimated foraging radius is approximately 400 m from the nest box (Quinney and Ankney 1985) and their foraging area might be larger than the Saltmarsh Sparrows'.


Saltmarsh Sparrows forage entirely in the salt marsh (Greenlaw and Rising 1994, Merriam 1979). On Long Island, New York Merriam (1979) found that the two most common insect orders in Saltmarsh Sharp-tailed Sparrow's diet were Diptera, ranging between 13 % in June to 47% of all items in July (predominantly adults and larvae of Stratiomyidae) and Hemiptera, ranging between 4% in June to 37% in July) (nymphs and adults of Miridae). Other invertebrates found in the diet of nestling Sharp-tailed Sparrows were Homoptera (Deplhacidae) ranging between 0 and 34%, the Araneida (spiders), ranging between 3% in July to 56% in June, and amphipods (between 5% in first half of July to 15 % in the second half of the month (Merriam 1979).

Egg mercury

We collected one egg from each box occupied by a nesting pair of Tree Swallows from Scarborough Marsh WMA and Furbish Marsh. We also collected failed Saltmarsh Sharp-tailed Sparrows eggs from various sites in Maine and New Hampshire. A total of 20 Tree Swallow eggs were analyzed for Hg and 17 Saltmarsh Sparrow eggs from seven estuaries were opportunistically collected and tested.

In general, Hg levels in the eggs were low (Figure 6). Based on a Hg dosing study (Heinz pers. comm.) conducted on Tree Swallow eggs the lethal LOAEL in swallow egg =0.8 ppm (ww) and sublethal effects are estimated at 0.4 ppm (ww) (were detected at as low as 0.1 ppm, further experiments are necessary to more accurately identify the sublethal concentration). All eggs tested in this study are well below 0.4 ppm and the lethal LOAEL.

Figure 6. Mean (+/-sd) egg mercury concentrations from flooded out Saltmarsh Sharp-tailed Sparrow nests collected in Maine and New Hampshire in 2004, (n) =egg sample size.



It might be of benefit to test amphipods and other prey groups from the selected sparrow sites for Hg content.

We found a positive correlation between female Tree Swallow blood and egg Hg levels (Figure 7). In a previously funded EPA study we also found a positive relationship between adult female swallow blood and egg Hg levels (n=45, r²=0.49) (BRI unpubl. data). We are unable to correlate sparrow blood and eggs because we do not have Hg values from the eggs and blood from the same bird.

Based on our small sample size of eggs collected from Rachel Carson NWR in Wells-Furbish Road, there was no significant difference in Hg levels between Tree Swallow and Saltmarsh Sparrow eggs (Table 8), (F=0.12; df=12, P=0.8). We found a significantly higher blood Hg levels in the Sparrows vs. Swallows but the egg Hg concentrations were not significantly different between the species. Therefore, contaminant levels measured in eggs might underestimate the total Hg exposure to the species. Interpreting Sharp-tailed Sparrow egg contaminant levels should be done with caution and eggs might not be the best tissue to use in estuarine contaminant studies. It is possible that at the time of egg laying the females' body burden of Hg is not at high enough levels to be depurated into the eggs.

Figure 7. Relationship between egg and female blood mercury in nesting Tree Swallows in two estuaries in Maine, 2004.




Table 8. Mean mercury concentrations (ppm, ww) in Tree Swallow and Saltmarsh Sparrow eggs collected in Maine, 2004.

Site	Mean egg Hg +/- sd (n)	
	Tree Swallows	Saltmarsh Sharp-tailed Sparrows
Furbish Marsh-RCNWR	0.11 +/- 0.06 (9)	0.10 +/- 0.02 (4)
Scarborough Marsh WMA	0.09 +/- 0.03 (6)	--

Conclusions

Based on the results of two years of sampling of Saltmarsh Sharp-tailed Sparrows we conclude that:

1. Bird blood is an appropriate tissue to use to evaluate the Hg exposure to insectivorous birds in salt marshes.
2. Saltmarsh Sharp-tailed Sparrows have elevated blood Hg levels across the sampling sites and are at potential risk in Parker River NWR in Massachusetts and Ninigret NWR in Rhode Island.
3. Mercury levels in sparrow blood from all sites (except RI) were higher in 2005 than in 2004.
4. Mercury levels in sparrow blood sampled in late July-August appear higher than in birds captured in June-early July.
5. It appears that sparrow eggs may not be the best indicator tissue to use in assessing Hg exposure in Saltmarsh Sharp-tailed Sparrows.
6. Tree Swallows have significantly less mercury in blood than Saltmarsh Sharp-tailed Sparrows indicating that aerial feeders such as swallows may not best represent Hg risk in estuaries.

Future Recommendations:

1. Continue sampling Saltmarsh Sparrows at the sites with the highest Hg levels to increase sample size and understanding of risk.
2. Sample other species that could be at risk to mercury contamination in the estuaries, such as willets.
3. Collect sparrow blood samples from additional sites in estuaries near possible point sources of Hg.
4. Conduct a vigorous nest search to determine productivity.
5. Continue to collect Tree Swallow blood and egg samples for Hg analysis at the established in 2004 sites.
6. Collect prey samples in the study estuaries for diet composition, Hg and isotope analysis.

Acknowledgments

Funding for this project was provided by US Fish and Wildlife Service, Maine Department of Inland Fisheries and Wildlife and Maine Department of Environmental Protection. We are grateful to Kate O'Brien, Graham Taylor, and the staff of Rachel Carson NWR for their logistical support and help in the field. Kate O'Brien provided photographs for this report. We thank Tom Hodgman and Greg Shriver for their expert technical advice. Many thanks to Andrew Major and Steve Mierzykowski for help in the field and with the project. We thank the following individuals for their help and support: Kate Laskowski for her assistance in the field; Dr. Kim Babbit and Megan


McElroy and field crew from UNH for collecting failed eggs; biologists: Nancy Pau at Parker River NWR, Sara Williams at Steward B. McKinney NWR and Suzanne Paton at Ninigret NWR; Carina Gjerdrum and Chris Elphick from UCONN for providing sparrow blood from Hammock River site in CT; and all the volunteers who came out to help catch birds, especially Frank Dehler for his many hours of assistance and valuable help. Many thanks to Bob Taylor and his team at Texas A&M University for providing lab analysis. BRI biologists Theresa Daigle, Melissa Duron and Wing Goodale were a tremendous help in the field and in the office.

Literature Cited

- Blancher, P.J. and D.K. McNicol. 1991. Tree swallow diet in relation to wetland acidity. *Can. J. Zool.* 69:2629-2637.
- Driscoll, in pres.
- Elwood, J.W., S.G. Hildebrand, and J.J. Beauchamp. 1976. Contribution of gut contents to the concentration and body burden of elements in *Tipula* spp. from a spring fed stream. *J. Fish. Res. Bd. Can.* 33:1930-1938.
- Evers, D. C., O. P. Lane, L. Savoy and W. Goodale. 2004. Assessing the impacts of methylmercury on piscivorous wildlife using a wildlife criterion value based on the Common Loon, 1998-2003. Report BRI 2004-05 submitted to the Maine Department of Environmental Protection. BioDiversity Research Institute, Gorham, Maine.
- Evers, D.C, N.M. Burgess, L. Champoux, B. Hoskins, A. Major, W. Goodale, R. Taylor, and T. Daigle. 2005. Patterns and interpretation of mercury exposure in freshwater avian communities in Northeastern North America. *Ecotoxicology*, 14:193-222.
- George, B.M., D. P. Batzer, and R. Noblet. 2001. Assessment of mercury concentrations in invertebrates of the Okefenokee Swamp of Southeast Georgia. Presented at the NABS Annual meeting, La Crosse, Wisconsin, 2001
- Greenlaw, J.S. 1993. Behavioral and morphological diversification in Sharp-tailed Sparrows (*Ammodramus caudacutus*) of the Atlantic Coast. *Auk* 110:286-303.
- Greenlaw, J.S. and J.D. Rising. 1994. Sharp-tailed Sparrow (*Ammodramus caudacutus*). In *The Birds of North America*, No. 112 (A. Poole and F. Gill, Eds.). Philadelphia: The Academy of Natural Sciences; Washington, D.C.: The American Ornithologists' Union.
- Hildebrand, S.G., H. Rodney, J. Strand, and W. Huckabee. 1980. Mercury accumulation in fish and invertebrates of the North Fork Holston River, Virginia and Tennessee. *L. Environ. Quality*. 9:393-400.
- Hodgman, T.P., W.G. Shriver, and P.D. Vickery. 2002. Redefining range overlap between the sharp-tailed sparrows of coastal New England. *Willson Bull.*, 114: 38-43.
- Marvin-DiPasquale, M.C., J.L. Agee, R.M. Bouse, and B.E. Jaffe. 2003. Microbial cycling of mercury in contaminated pelagic and wetland sediments of San Pablo Bay, California. *Environmental Geology*, 43:260-267.
- McCarty, J.P. and D.W. Winkler. 1999. Foraging ecology and diet selectivity of tree swallows feeding nestlings. *Condor* 101:246-254.
- Merriam, T.L. 1979. Feeding ecology and food overlap of two sympatric marsh-inhabiting sparrows (Aves: Ammospiza). Master of Science thesis. Long Island University, New York.
- Quinney T.E. and C. D. Ankney. 1985. Prey size selection by tree swallows. *Auk*, 102: 245-250.
- Rimmer, C.C., K.P. McFarland, D.C. Evers, E. K. Miller, Y. Aubry, D. Busby and R J. Taylor. 2005. Mercury Concentrations in Bicknell's Thrush and Other Insectivorous Passerines in Montane Forests of Northeastern North America. *Ecotoxicology*, 14: 223-240.
- Sarica, J., M. Amoyt, J. Bien, and L. Hare. 2005. Fate of mercury accumulated by blowflies feeding on fish carcasses. *Environ. Tox. Chem.* 24:526-529.
- Shriver, W.G. 2002. Conservation Ecology of Salt Marsh Birds in New England. Ph.D. Dissertation. State University of New York, Syracuse, NY.
- Shriver, W.G., D.C. Evers, and T. Hodgman. 2002. Mercury exposure profile for Sharp-tailed Sparrows breeding in coastal Maine salt marshes. Report BRI 2002-11 submitted to the Maine Department of Environmental Protection. BioDiversity Research Institute, Falmouth, Maine.


- Shriver, W.G., J.P. Gibbs, P.D. Vickery, H.L. Gibbs, T.P. Hodgman, P.T. Jones, and C.N. Jacques. 2005. Concordance between morphological and molecular markers in assessing hybridization between sharp-tailed sparrows in New England. *Auk* 122:94-107.
- Thompson, D. R. 1996. Mercury in Birds and Terrestrial Animals. In: W. N. Beyer, Gary H. Heinz, Amy W. Redmon-Norwood (Ed.), *Environmental Contaminants in Wildlife: Interpreting Tissue Concentrations* (pp. 341-355). Clemson, SC: Lewis Publisher.
- Wolfe, M. and D. Norman. 1998. Effects of waterborne mercury on terrestrial wildlife at Clear Lake: evaluation and testing of a predictive model. *Environmental Toxicology and Chemistry* 17, 214-227.

Appendix I. Saltmarsh Sharp-tailed Sparrow sampling locations in New England, 2004.



Appendix II

Saltmarsh Sharp-tailed Sparrow sampling locations in the Rachel Carson NWR and other locations in Maine, 2001 and 2004.



Appendix III. Saltmarsh Sharp-tailed Sparrow sampling location in Parker River NWR, Massachusetts, 2004 (map provided by Nancy Pau).

