

Chemical Assay of Drugs and Drug Metabolites

Sanford P. Markey
Laboratory of Neurotoxicology
NIMH

September 18, 2008


Lecture Outline

- Quantification principles
 - Analytical PK lab tasks
- Chromatography
- Detection - spectroscopies
 - Optical
 - Mass
- Examples
 - Resveratrol
 - Aminoflavone
 - CYP450 Assays
 - Cyclosporin A
- References


Definition of Analytical Terms

- Limits of detection (LOD)
 - Sensitivity is the minimum detectable concentration change that can be observed at a specified concentration
 - LOD is the minimum mass or concentration of analyte that can be detected at an acceptable signal to noise (S/N) ratio
- Limits of quantification (LOQ)
 - Analyte mass or concentration required to give an acceptable level of confidence in the measured analyte quantity
 - Always greater (usually 3x) than the minimum LOD


Accuracy vs. Precision



Good accuracy
Poor precision



Poor accuracy
Good precision



Good accuracy
Good precision

Pharmaceutical Industry PK Lab

Analytical Assays (1)

- Parent drug usually the target analyte for Phase 1 dose response and safety determinations
- Scale of runs: 30-50 samples/patient, plus 10-15 standards, procedural blanks, plus 10-15 QC pools or previously analyzed samples
- Several patients per run - effort to optimize patient/(standards + QC) ratio. Result is >100 samples/run
- Analytical runs require automation & rugged instrumentation, continuous operation for assay cycle time X number of samples
- Develop assays on 96 well or 384 well devices

Pharmaceutical Industry PK Lab Analytical Assays (2)

- Speed of assay development principal determinant of methodology choice
- Avoid derivatization chemistry
- Use solid phase extraction or simple methanol/acetonitrile protein precipitation
- Time is money (5 min LC/MS/MS assay vs. 40 min HPLC)
- Use automated LC/MS/MS methods with high sensitivity and specificity

Assay Issues

- What to assay (what is important?)
 - Species -
 - man, non-human primate, rat, mouse (transgenic)
 - Tissue/Fluid
 - liver, target organ, plasma, excreta
 - Isolated organ/tissue fluids
 - liver slices, human liver microsomes, CYPs, other enzymes

Assay Issues

- Commercial Aides
 - Drug metabolizing preparations
 - Human liver tissue or hepatocytes – all enzymes present in fresh (not frozen) tissue – single use only
 - Microsomes from frozen liver; easily stored
 - Recombinant CYPs and other enzymes - widely available (yeast, baculovirus, bacteria) and some mammalian cells with NADPH CYP reductase
 - CYP substrates, antibodies, inhibitors, inducers
 - Computer software - predict metabolites, pKa, pLogD, logP
 - Contract Research Organizations

Liquid Chromatography


- High Performance (HPLC)
 - Reverse Phase - polarity separation
 - Immunoaffinity
 - Cation & Anion Exchange - charge separation
 - Smaller particle size, higher pressures - higher performance




Liquid Chromatography

stationary phase

column



HPLC Apparatus




Detection Principles (1)


- Ultraviolet or Fluorescence
Spectroscopy
 - chromophore in drug or derivatized drug
 - most useful for known target analytes
- Nuclear Magnetic Resonance
Spectrometry
 - most useful for totally unknown chemical structure characterization
 - least sensitive



UV Absorption Spectrophotometer



Emission Spectrophotometer




Detection Principles (2)


- Mass Spectrometry
 - versatile ionization modes for liquids and gases
 - electron, chemical, electrospray, desorption
 - versatile mass analyzers with varying capabilities
 - magnetic, ion trap, quadrupole, time-of-flight
 - combination analyzers
 - triple quadrupole
 - quadrupole-time-of-flight
 - linear trap-orbitrap, etc, etc
 - very sensitive and structurally informative –
example: air, acetaminophen
 - added specificity through mass chromatography
 - tandem mass chromatography = multiple reaction monitoring



Mass Spectrometer Component Overview




Mass Spectrometer Ionizers



mass analyzer


Electron
Ionization
(in vacuo)




Electrospray
Ionization
(external)



Mass Analyzers




Time-of-flight
(TOF)



Quadrupole
(q)

Quadrupole Ion Trap

C.




Electrospray-Ion Trap Mass Spectrometer


esilcq

Overview of Electrospray Ionization using an Ion Trap Mass Spectrometer

DETECTION



Mass Spectrum





Mass-to-Charge Ratio (m/z)	Relative Intensity (%)
100	~20
200	~20
300	~25
400	100


TRAPPING SCANNING DETECTION

PLAY STOP LABEL


ThermoFinnigan




Mass Spectrum of Air




Mass spectrum of acetaminophen (Electron Ionization)



Mass Chromatography



Multidimensional Analyses



Pharmaceutical Industry PK Lab Analytical Assay Work Load for New Chemical Entities

Method	1990	1998	2000	2008
HPLC	75%	50-60%	20%	2%
GC/MS	12%	3%	2%	0
LC/MS/MS	3%	40-50%	60-75%	98%
RIA	10%	10%	10%	0
Preliminary lead profile time	18 m	4 m	0	0

Conclusion: requirement for speed (not instrumentation cost)
dictates choice of analytical methods

Popular Methods for Qualitative & Quantitative Assays in Clinical Pharmacology

- LC/MS/MS
 - High speed, reduced requirement for sample preparation
- HPLC/UV or Fluorescence
 - Very robust, routine assay technology
- Enzyme Linked Immunoassay (ELISA)
 - Many 96 well formatted colorimetric or radiometric commercial assay kits for specific compounds
- Fluorescence polarization immunoassay (FPIA)
 - Measures difference in fluorescence between bound and free antigen
 - Important in therapeutic drug monitoring - CsA

Examples of Analytical Methods Applied in Drug Analyses

- 1. Resveratrol - bioavailability
- 2. CYP450 Assays - LC/MS/MS
- 3. Cyclosporin - FPIA, HPLC/UV, LC/MS/MS

Example 1 -Where Do Drugs Go?

- Radiochemical tracers (^{14}C , ^3H)
 - requires availability of labeled drug
 - useful for bioavailability, kinetics - **Resveratrol**
 - detection of protein adducts/localization (autoradiography)
- Non-radiochemical methods
 - Unique drug elements (fluorine, etc.) or structural property (fluorescence)
 - Specific atom or isotope detectors
 - Accelerator mass spectrometry (AMS) - detection of ^{14}C at near natural background levels for drug pharmacokinetics
 - Ideal for human studies of toxic mechanisms - DNA
 - Calcium metabolism

Resveratrol

Washington Post, November 2, 2006

A Compound in Red Wine Makes Fat Mice Healthy

By Rob Stein □

A substance found in red wine protected mice from the ill effects of obesity and extended their life spans, raising the tantalizing prospect that the compound could do the same for humans and may also help people live longer, healthier lives, researchers reported yesterday...

"We've been looking for something like this for the last 100,000 years, and maybe it's right around the corner -- a molecule that could be taken in a single pill to delay the diseases of aging and keep you healthier as you grow old," said David A. Sinclair, a Harvard Medical School molecular biologist who led the study.


Resveratrol

JA Baur, et al...DA Sinclair

Nature **444**, 337-342 (16 November 2006)

Resveratrol improves health and survival of mice on a high-calorie diet

- $22.4 \pm 0.4 \text{ mg/kg}^{-1}/\text{day}^{-1}$ in food




- Resveratrol is a polyphenolic SIRT1 activator
 - ameliorates insulin resistance
 - increases mitochondrial content


Lagouge, M. et al. Cell 127, 1109–1122 (2006)

Resveratrol Delays Age-Related Deterioration and Mimics Transcriptional Aspects of Dietary Restriction without Extending Life Span

Pearson KJ, et al. Cell Metabolism 8, 6 August 2008, 157-168




Resveratrol



- Widely sold at health food stores as antioxidant
- Proposed chemopreventive for cardiac diseases, cancer based on *in vitro* evidence
- Absorption?
- Bioavailability?
- Metabolism?

HIGH ABSORPTION BUT VERY LOW BIOAVAILABILITY OF ORAL RESVERATROL IN HUMANS
T.Walle et al., Drug Metab Disp 32:1377-1382 (2004)

Resveratrol plasma concentration-time curves (total radioactivity)



Oral 25 mg ● i.v. 0.2 mg □

T.Walle et al., Drug Metab Disp 32:1377-1382 (2004)

Resveratrol Urine Metabolites (0-12 hr)

LC/MS


(selected ions)

M1,2=RV-Glu

M3= RV-H₂-Glu

M4= RV-SO₄


M5 = RV-H₂-SO₄



LC/UV

305 nm

(note
absence of
M3, M5)




Resveratrol

Recovery of Radioactivity

	25 mg Oral		0.2 mg i.v.	
	Urine	Feces	Urine	Feces
N=6	70.5 ± 4.3	12.7 ± 6.1	64.1 ± 7.7	10.4 ± 3.7

T.Walle et al., Drug Metab Disp 32:1377-1382 (2004)

HPLC Radiochromatogram 0-12 hr urine extract



□ *Urine* ● *After aryl sulfatase*

Glucuronidase shifts M1-M3 to Resveratrol r.t.

T.Walle et al., Drug Metab Disp 32:1377-1382 (2004)

Resveratrol Study

Conclusions

T.Walle et al., Drug Metab Disp 32:1377-1382 (2004)

- Unmetabolized resveratrol not detectable in plasma
- Absorption of resveratrol is at least 70%
- No evidence for further oxidation - only conjugation \pm reduction
- Bioavailability of resveratrol limited
 - Highly accumulated in intestinal epithelial cells
 - Target sites of breast and prostate unlikely unless RV-SO₄ is active species or reservoir of parent
- Small molecule activators of SIRT1 sought as alternative therapeutics
 - Milne JC et al. Nature 450, 712-716, 2007

Example 2: LC/MS/MS CYP GLP Assays


- 12 Semi-automated assays for 10 human CYP450 enzymes described
- Microsomes pooled from 54 human livers
- Microsomes, NADPH, substrate in 96 well plate; stable isotope internal standards added with quenching solvent
- Recombinant CYP450 enzymes (Sf9 cells) from PanVera run in parallel; reference values published
- High speed LC/MS/MS conditions established for each analyte and internal standard (2 min/assay)
- Interassay precision of reaction velocity <10%

Validated Assays for Human Cytochrome P450 Activities, RL Walsky and

RS Obach, *Drug Metab Disp* 32:647-660, 2004




CYP 450 Validated Assay

Bupropion and hydroxy metabolite




multiple reaction monitoring

Hydroxybutpropion - ESI-MS + [D6]-hydroxybutpropion




Hydroxybupropion - CID of MH^+ 256

[D6]-Hydroxybupropion - CID of MH^+ 262




Example: CYP2B6 Assay Bupropion substrate



From RL Walsky & RS Obach

Example: CYP2B6 Results

BUPROPION HYDROXYLASE
HLM-13 0.05 mg/ml
Product Formed vs Time



From RL Walsky & RS Obach


Partial Summary of CYP Activities

RL Walsky and RS Obach, Drug Metab Disp 32:647-660, 2004

Enzyme	Assay	Inhibitor	IC ₅₀ (μM)	
			Human	Recomb
CYP1A2	Phenacetin O-deethylase	Furafylline	1.76±0.28	1.54±0.16
CYP2A6	Coumarin 7- hydroxylase	Tranlylcyp- romine	0.449±.073	0.895±.262
CYP2B6	Bupropriion hydroxylase	PPP	7.74±0.47	2.02±0.19
CYP2C8	Amodiaquine N-deethylase	Quercetin	3.06±0.31	3.33±0.20
CYP2C9	Diclofenac 4'- hydroxylase	Sulfaphen- azole	0.272±.031	0.169±.004

Example 3: Cyclosporin A (CsA)

*Potent immunosuppressive drug
for transplantation; irreversible
kidney damage if dose too high*




- HPLC - UV (210 nm) method first used for clinical analyses
 - LOQ - 20-45 $\mu\text{g/L}$ (therapeutic range 80-300 $\mu\text{g/L}$)
- LC/MS/MS method for fingerprick samples
 - 25 μL ; LOQ 10 $\mu\text{g/L}$
 - Keevil BG, Ther Drug Monitor 24: 757-67 (2002)

Cyclosporin Immuno Assays


- Florescence polarization immunoassay (FPIA)
 - Homogeneous immunoassay
 - Fluorescein tagged drug competes with patient drug for monoclonal Ab
 - Polarized light excites Ab-tagged drug complex most efficiently
 - LOQ 25 µg/L; analysis of 20 samples in 19 min
- Enzyme monitored Immunoassay Technique (EMIT) and Cloned Enzyme Donor Immunoassay (CEDIA)
 - Competitive: enzyme labeled antigen competes with sample antigen; enzyme labeled antigen-Ab complex changes rate
- Multiple cyclosporin metabolites exhibit cross-reactivity in immunoassays

Monoclonal CEDIA


a)




a)



b)



b)



Blood concentrations of cyclosporine (CSA)

Mai I, et al. *Clinical Pharmacology & Therapeutics* (2004) 76, 330–340

WWW Sites (1) HPLC & Drug Metabolism

- Tutorial for HPLC
 - <http://kerouac.pharm.uky.edu/asrg/hplc/HPLCMYTRY.HTML>
- Prediction Software – pK, structure
 - <http://www.acdlabs.com/>
- Human Drug Metabolizing Enzymes:
 - Celsis (<http://www.celsis.com/index.php>)

WWW Sites (2): Mass Spectrometry Information Education

- <http://ull.chemistry.uakron.edu/classroom.html>
 - Excellent introductory tutorials in analytical methods including chromatography and mass spectrometry
- <http://ionsource.com/>
 - Site with very useful links for mass spectrometry including tutorials, freeware