Allotment Management Plan

North Fork-Boulder Allotment Ketchum Ranger District Sawtooth National Forest

Prepared by:	(Title)	Date:	
Reviewed by:	Permittee	Date:	
	Permittee	Date:	
	Permittee	Date:	
2003 Revision Plan and the S Allotment Final	Management Plan imple of the Sawtooth National eptember 30, 2004 Reco Environmental Impact S eart of your Term Grazing ermit.	Forest Land and Re rd of Decision for the Statement. This Alle	source Management North Fork Boulder otment Management
Approved by:	District Ranger	Date:	

Table of Contents

1. 1	PURPOSE	3
A.	Area Map	4
II.	GOALS & OBJECTIVES	5
A	Forest Plan I. Forest-Wide Goals 2. Forest-Wide Objectives 3. Management Area Prescriptions 4. Management Area Objectives North Fork-Boulder and Baker Creek ROD I. Goals Objectives Desired Conditions I. Forest Plan Level 2. Allotment Specific a) Riparian b) Upland Mesic Riparian c) Sagebrush & Non-forested Uplands	
	d) Desired Condition for Aspen Stands	
III.	ADAPTIVE MANAGEMENT ACTIONS	
I	Adaptive Management Decision Tree	
IV.	MANAGEMENT REQUIREMENTS	20
	1. Livestock kind, class, numbers permitted, and season of use	
V. 1	IMPROVEMENTS	22
VI.	MONITORING	22
	1. Implementation Monitoring	24 25
	a) Riparian Effectiveness Monitoringb) Photo Points	
	3. Upland Effectiveness Monitoring	
4	4. Aspen Monitoringa) Aspen Monitoring Location Map	
VII.	REFERENCES	

I. PURPOSE


The Federal Land Policy Management Act (FLPMA), as amended by the Public Rangelands Improvement Act (PRIA) allows for Allotment Management Plans (AMPs) to be included in grazing permits at the discretion of the Secretary of Agriculture (43 U.S.C. (1752(d), as amended by 92 Stat. 1803 (1978)). The Secretary has elected to exercise this discretion, and has delegated his authority to issue regulations in this area to the Chief of the Forest Service (36 CFR 222.1 et.seq.). An AMP is defined in FLPMA as a document prepared in consultation with permittees applying to livestock operations on the public lands (1) prescribing the manner in and extent to which livestock operations will be conducted in order to meet multiple use, sustained yield, economic and other needs and objectives,(2) describing range improvements to be installed and maintained, and (3) containing such other provisions relating to livestock grazing and other objectives found by the Secretary to be consistent with the provisions of the FLPMA (43 USC 1702(k), and 36 CFR 222.1(b)(2), and FSM 1023).

This AMP contains the pertinent livestock management direction specific to implementing the July 2003 revision of the Sawtooth National Forest Land and Resource Management Plan (Forest Plan), the September 30, 2004 North Fork-Boulder and Baker Creek Allotment Management Plan Updates Record of Decision (ROD), and the February 2008 Supplement to the North Fork-Boulder and Baker Creek ROD. This AMP is included as part of any term grazing permit that authorizes livestock grazing on the North Fork Boulder Allotment per Section 8(a) Part 2 of the term grazing permit). The primary triggers for considering adaptive management changes are exceeding annual allowable grazing use criteria and not achieving or trending toward achieving long-term desired conditions

This AMP implements an adaptive management approach to livestock management on the North Fork Boulder Allotment. Adaptive management provides for: 1) identification of site specific desired conditions; 2) definition of appropriate decision criteria (constraints) to guide management; 3) identification of pre-determined optional courses of action, as part of a proposed action, from which to adjust management decisions over time; and 4) establishment of carefully focused project monitoring to be used to make adaptive adjustments in management over time (R4 FSH 2209.13 chapter 90). This AMP is an essential part of implementing this direction. As such, the management requirements, improvement needs, and monitoring plan may be modified as needed to insure that management direction is consistent with achieving the direction established in the Forest Plan and ROD. Additional NEPA analysis and decision processes will not be needed as long as the direction identified in this plan is consistent with the analysis conducted in the September 2004 North Sheep Final Environmental Impact Statement (FEIS). Vegetation improvement projects not specifically addressed in the FEIS will require additional analysis at the appropriate level and scale.

Adaptive management protocols will follow the direction established in the Rangeland Resources Forest Plan Implementation Guide. (USDA, 2005).

A. Area Map



II. GOALS & OBJECTIVES

A. Forest Plan

Forest Plan goals and objectives are identified at two scales. The Forest Plan scale describes goals and objectives that are generally applied to all areas within the Forest. The management area scale describes management direction for specific management areas within the National Forest. Management Area objectives form the basis for development of project-level actions or proposals to help achieve Forest goals. However, not all projects proposed and designed are expected to respond to all objectives in the Forest Plan. This allotment falls within the Big Wood River Management Area (Management Area No. 4).

1. Forest-Wide Goals

RAGO01- Provide for livestock forage within existing open allotments, in a manner that is consistent with other resource management direction and uses.

RAGO02- Manage rangelands using controlled livestock grazing, range structural and non-structural improvements, vegetative and ground rehabilitation, fire, and timber management in various combinations to meet desired conditions.

RAGO03- Manage upland vegetation on suitable rangelands to maintain or restore hydrologic function and soil productivity of watersheds.

RAGO04- Manage herbaceous and shrub vegetation on suitable rangelands to meet resource objectives in an efficient manner.

RAGO05- Manage livestock grazing within riparian areas to accommodate the maintenance or restoration of aquatic and riparian processes and functions.

RAGO06- Coordinate livestock grazing to address conflicts with other resource uses in a manner that is consistent with Forest Plan management direction.

VEOB03- Utilize emerging technologies and science, and implement adaptive management process to provide for increasing the effectiveness of vegetation monitoring. **VEOB06-** Determine high priority areas for vegetation management actions that restore or maintain desired vegetation attributes.

2. Forest-Wide Objectives

RAOB01 - Coordinate the design, update and/or revision of AMPs with adjacent landowners to maximize opportunities and minimize potential management conflicts.

RAOB02 - Coordinate livestock grazing with timber harvest and forest regeneration activities to capitalize on management opportunities, while minimizing activity conflicts to help meet Forest Plan Vegetation and Rangeland Resource goals.

RAOB03 - Identify rangeland facilities that are degrading resource conditions and prioritize opportunities to mitigate their effects or to initiate restoration of resource conditions.

3. Management Area Prescriptions

Management prescriptions are defined as, "Management practices and intensity selected and scheduled for application on a specific area to attain multiple use and other goals and objectives" (36 CFR 219.3). Management prescription categories (MPC) are broad categories of management prescriptions that indicate the general management emphasis prescribed for a given area. They are based on Forest Service definitions developed at the national level, and represent management emphasis themes, ranging from Wilderness (1.0) to Concentrated Development (8.0). The national MPCs have been customized during Forest Plan revision to better fit the needs and issues of the Forest. The allotment includes two MPCs, MPC 4.1c and 4.2 (see map).

<u>4.1c – Undeveloped Recreation</u>: Maintain Unroaded Character with Allowance for Restoration Activities

This prescription applies to lands where dispersed recreation uses are the primary emphasis. Providing dispersed recreation opportunities in an unroaded landscape is the main objective. Both motorized and non-motorized recreation opportunities may be provided. Other resource uses are allowed to the extent that they do not compromise the area's Recreation Opportunity Spectrum (ROS) settings.

4.2 – Roaded Recreation Emphasis

This prescription applies to lands where dispersed and developed recreation uses are the primary emphasis. A wide range of recreational activities and developments occurs. Facilities are maintained, and both motorized and non-motorized recreation opportunities may be provided. Multiple uses such as timber harvest and grazing are allowed to the extent that they do not compromise recreation resource objectives. Generally, a mix of mechanical and fire activities are used to treat vegetation to achieve desired conditions for recreation settings and developments, and to reduce the risk of uncharacteristic vegetative damage or loss from insects, diseases, and fire.

4. Management Area Objectives

- Restore dry meadows by improving specie composition, reducing compaction and increasing plant vigor due to the effects of livestock grazing, dispersed recreation, and road alteration on natural drainage patterns. due to the effects of livestock grazing. (Objective 0447)
- Restore structure and native species composition, as described in Appendix A, in the Alpine Meadows, Dry Meadows, and Mountain Big Sagebrush vegetation groups. (Objective 0448).
- Prevent the spread of noxious weed seeds due to domestic sheep by adjusting or changing management practices, such as trailing route locations and driveway/grazing area seasons of use. (Objective 04111)
- Confine, contain, or reduce the density of noxious weed infestations, particularly spotted knapweed, diffuse knapweed, and dalmation toadflax, within the big wood River drainage.(Objective 0452)

- Continue weed management coordination efforts with local management agencies and private individuals. (Objective 0453)
- Provide high –quality mountain goat forage by minimizing or reducing summer and fall forage competition between domestic sheep in the tributaries to the North Fork Big Wood River. (Objective 0454)
- Maintain and restore habitat for deer, elk, migratory land birds, and sage grouse in lower elevation sagebrush communities. (Objective 0456)
- Within the SNRA, manage federal and private lands to retain a pastoral or natural-appearing landscape consistent with the scenic values for which the SNRA was established. (Objective 0488)
- Adjust management practices to minimize livestock/recreation conflicts within high density recreation areas including North Fork Big Wood River. (Objective 04109)
- Prevent the spread of noxious weed seeds due to domestic sheep by adjusting or changing management practices, such as trailing route locations and driveway/grazing area seasons of use. (Objective 04111)
- Do not bed sheep within designated campgrounds or summer home tracts, or on trails, trailheads, or maintained roads. (Objective 04112)

B. North Fork-Boulder and Baker Creek ROD

The following goals are described in Record of Decision for the North Fork-Boulder & Baker Creek Allotment Management Plan Updates. They are found in the Decision Section (p. 1) and the Adaptive Management Section (p. 2). The objectives were derived from the Record of Decision discussions on "How the Decision Responds to Public Concerns and Needs for Change" (pp. 5-9).

1. Goals

- Achieve and maintain sustainable grazing system on the North Fork Boulder Allotment
- Implement adaptive management approach
- Authorize grazing in a manner that will meet FLRMP standards
- Achieve or make progress towards FLRMP desired conditions affected by grazing

Objectives

- Confine, reduce or eradicate, noxious weed infestations, particularly spotted knapweed in the Phantom hill areas. Prevent the spread of noxious weeds seeds due to domestic sheep by adjusting or changing management practices, such as trailing routes, and grazing area seasons of use. Use sheep driveway to access allotment during early season before seed production. Avoid area trailing off in fall.
- Provide for high –quality mountain goat forage by adhering to closures of summer and fall forage use by domestic sheep in the high basins.
- Restore the protective soil mantle and indigenous plants species that are characteristic of these extreme high altitude environments.

• Increase vegetative cover, density, and structure to uplands sites in the Boulder Front and Dip Creek areas of the allotment through proper herding techniques Implementation monitoring will be emphasized.

C. Desired Conditions

1. Forest Plan Level

Rangeland Resource Desired Condition. A sustainable level of forage, consistent with other resource management direction, is available for use through the Forest Service grazing permit system. Rangeland forage quality is maintained or improved in areas where vegetation management projects and range management actions occur. Riparian areas continue to be a focal point for providing vegetative diversity, landscape capability, soil productivity, wildlife habitat, proper stream channel function, and water quality important to sustaining beneficial uses. Riparian areas are functioning properly and/or have improving trends in vegetative composition, age class structure and vigor. Upland range vegetation is contributing to proper hydrologic function. The composition and densities of shrubs, grasses and forbs are variable and dynamic across the landscape.

2. Allotment Specific

a) Riparian

Streambank vegetation in late seral condition per Winword 2000 Classification Bank Stability at 90 % of potential in areas impacted by grazing. Manage for maximum 45% use or retain a 4-inch stubble height of hydric greenline species.

Grazing management on the allotment will focus on maintaining riparian plant communities:

Salix exigua/mesic graminoid- Characterized by overstory of Salix exigua with moderate to dense cover of graminoid species, including Carex nebrascensis, C. lanuginose, Juncus balticus, Eleocharis palustris, Agrostis stolonifera, Scirpus pungens, and Agropyron repens. Manage these plant communities to maintain plant vigor, composition, and density and prevent conversion bluegrass plant communities.

Salix geyeriana/mesic forb- Characterized by Salix Geyeriana with some S. drummondiana and S. boothii, and understory shrubs of Ribes inerme, Amelanchier alnifolia, and forb component of Veratrum, Senecio, Heracleum, Geum, and graminoids such as Agrostis stolonifera, Carex nebrascensis, and poa pratensis. Manage these plant communities to maintain plant vigor, composition, and density of willow species.

Salix geyeriana/mesic graminoid- Characterized by Salix geyeriana and a diverse mix of graminoids such as Carex lanuginose, Deschampsia cespitosa, C. microptera, C. aquatilis, C. ultriculata, and forbs including Taraxacum officinale, Achillea millefolium, Trifolium, and Potentella gracilis. Manage these plant communities to maintain plant vigor, composition, and density of willow species.

b) Upland Mesic Riparian

There is an adequate cover of key native species (both woody and herbaceous) such as carex, rush, and willow species in a variety of age classes that are adequately reproducing and maintaining good vigor. Desired condition can be further explained as cover of these species to be 70 percent or greater of estimated potential. No active head cutting is occurring.

c) Sagebrush & Non-forested Uplands

Sagebrush meets FLRMP standards of:

30-40% of area in 0-10% canopy cover class

30-40% of area in 11-20% canopy cover class

20-30% of area in 21-30% canopy cover class

<= 5% of area in > 30% canopy cover class

Note: Recovery to meet this condition will require disturbance treatment by fire or other means irrespective of sheep management practices. This will require planning treatment projects with additional environmental analysis as funds and priorities warrant.

Soil Cover on grazed areas will be within the desired range based on information available for existing land type classifications.

Plant composition on grazed areas will include species diversity and presence as described in FLRMP App. B.

Perennial Grass Slopes (10-18 in. precipitation zone). Bluebunch wheatgrass is the dominant bunch grass. Perennial grasses compose 80-90% of production. Sandberg bluegrass is a minor component of the vegetation. Common forbs include Indian wheat, shining chickweed, salsify, yarrow, lupine, balsamroot, biscuit root, hawksbeard, fleabane, milkvetch and phlox. Noxious weeds are absent.

Perennial Grass Montane (14-30 inch precipitation zone). This type is interspersed with sagebrush, forested and mountain brush types. Idaho fescue is the dominant grass. Other species may include intermediate oatgrass, western needlegrass and Richardson needlegrass. Forbs compose 40-65% of production. Common forbs are yarrow, besssaya, geum, Indian paintbrush, lupines, phlox, and balsamroot.

d) Desired Condition for Aspen Stands

Stand Specific Desired Conditions

Aspen dominates the overstory canopy (aspen >8" DBH. Over 2/3rds of the overstory is aspen.

Aspen dominates the mid-level canopy (aspen 1-8" DBH). Over 2/3rds of this level of canopy is composed of aspen.

There is significant aspen regeneration occurring to support a healthy stand.

The stand has over 500 stems per acre <1in. DBH with less than 20% having multiple leaders or are hedged from browsing.

Less than 20% of the stand contains sagebrush.

Note: Recovery to meet this condition in some stands will require disturbance treatment by fire or other means irrespective of sheep management practices. This may require planning treatment projects with additional environmental analysis as funds and priorities warrant.

e) Noxious Weeds

Areas on the Allotment which do not currently have noxious weed infestations will be maintained noxious weed free. Current noxious weed infestations will be contained and weed density reduced.

III. Adaptive Management Actions

Adaptive management is a strategy based on three principles:

- 1. Achievement of realistic, clearly defined objectives;
- 2. Ongoing monitoring to assess progress toward those objectives; and
- 3. Flexibility to alter management when adequate progress is not being achieved.


This management strategy is most appropriate in dynamic situations, where change is the norm. Change can be a characteristic of the management setting, or the result of management activities, or both. In such situations, adaptive management is the most efficient way to achieve desired objectives.

The Sawtooth Forest Plan recognizes that most physical, biological, social, and economic systems are dynamic and that management must be correspondingly flexible in order to be effective.

The adaptive management procedure is based on both annual grazing use and long-term monitoring to determine if management is achieving long-term management objectives. Establishing a relationship between annual grazing use and achievement of long-term objectives necessarily emphasizes use of end-of-season annual grazing use indicators, as well as long-term indicators of rangeland condition. Within-season annual grazing use indicators may also be established through the adaptive management process to determine when livestock should be moved from a grazing unit to achieve appropriate end-of-season grazing use levels and resource management objectives. Annual grazing use indicators

(including Forest Plan Standards and Guidelines), both within-season and end-of-season, along with other required management practices, are a total package that, when implemented and adhered to, will result in a reasonable expectation that long-term desired condition objectives will be achieved.

Adaptive Management Decision Tree



In Block 1, the grazing permittee(s) and/or land manager evaluates whether the annual grazing use indicator or standard was met. This assumes that the correct indicator and value was being used. These annual indicators are initially set in the Forest Plan (Forest Plan pp. III-45 through III-47) and the monitoring sections of the Allotment Management Plans (Supplement. App. C). The adaptive management process provides the opportunity to evaluate and adjust annual grazing indicators. As the adaptive management process is followed, indicators may be modified based on the results of annual and long-term monitoring. This may be subject to re-evaluation later in the process.

- A. <u>Annual Indicator or Standard is Met</u>: If the annual grazing use indicator is met, current management will continue, including short- and long-term monitoring as indicated in Block 2.
- A1. <u>Continue Current Management and Monitoring (Block 2)</u>: Long-term monitoring indicators are used to assess whether management objectives for

resource conditions and values are being achieved. This data will be used over time to determine the effectiveness of management direction and/or annual grazing use indicators in achieving the desired conditions. Note: The adaptive management process may begin with this block when long-term monitoring is completed and evaluated.

A2.Modify the Annual Indicator and/or Management as Appropriate (Block 3): If the desired condition objective is not being achieved, there is a need to change management and/or modify either the type or value of annual grazing use indicators being used. The primary situations that could lead to modifying annual indicators are. 1) When long-term monitoring results indicate that desired conditions are not being achieved. Along with other management changes, it may be necessary to change the indicator to a more restrictive use criterion. For example, if bank stability goals are not being achieved with a 4 inch stubble height annual use criteria, raising the threshold to require leaving a 6 inch stubble height after grazing use may be appropriate. 2) The indicator is not sensitive to achieving desired conditions. For example, using a stubble height use criteria may not adequately address recovery of willow species on a site. It may be more appropriate to add or change the indicator to a limit on browsing on seedling and young willow plants. 3) New resource issues or use conflicts surface. If areas are significantly disturbed by fire, flood, or other disturbances that significantly change resource conditions, new or additional use criteria may be needed. 4) When desired conditions have been achieved, criteria may not need to be as restrictive to provide for maintenance of resource conditions as when managing for recovery of resource conditions.

- B. <u>Annual Grazing Use Indicator or Standard Is Not Met</u>: If the grazing use exceeds the annual grazing use indicator or standard, proceed to the evaluation steps in Block 4.
- B1. Analysis and Determination of the Need for an Adaptive Management Adjustment (Block 4): If the grazing use exceeds the established annual grazing use indicator or standard, the resource manager, in consultation with the permittee(s) and others as appropriate, determine: 1) the potential cause for exceeding the standard, and 2) the significance of the excessive grazing use relative to its impact on the achievement of the desired resource conditions.

The resource manager, in consultation with the permittee(s), should determine whether the failure to meet the annual grazing use indicator is an infrequent occurrence or whether there is routine difficulty in meeting annual grazing use standards. A one-time occurrence due to some unique variable may not be significant and may not require further evaluation or adaptive management adjustments. Routine difficulty in meeting the annual grazing use indicator may indicate further evaluation and the need for adaptive management adjustments.

If further evaluation is warranted, comparison of the current condition with the desired condition should be made. If there is a large departure between current conditions and desired resource conditions, it may be fairly obvious that the need to achieve the annual use indicator is significant and that adaptive management actions are needed to provide for the achievement of the annual use indicator and meet long-term objectives.

While the evaluation of current versus desired conditions should be made with the use of long-term monitoring data, this information may not be available. In that case, utilize the best available information or complete a simple and rapid qualitative analysis to compare current conditions with desired conditions. While long-term trend and condition information is preferred, the lack of such information should not delay the evaluation of the current rangeland condition and needed adaptive management adjustments. Adaptive management adjustments should be temporary modifications until quantitative long-term condition and trend information is available to support permanent changes. If the resource manager's evaluation concludes that current conditions are close to desired resource conditions, then failure to achieve the annual grazing use indicator during that grazing season may not be significant in terms of achieving long-term objectives. In this case, adaptive management adjustments may not be necessary. Existing management and monitoring to achieve desired conditions would continue (Blocks 2 and 3). The exception to this situation may be where available information indicates that the long-term trend is negative, and adaptive management adjustments are needed.

If the resource manager's evaluation concludes that there is a significant gap between current and desired conditions and there is no indication of a positive trend, then the need for adaptive management adjustments are indicated.

Note: Determination of "large departure" may be either qualitative or quantitative depending on available information. Interdisciplinary teams or resource specialists may rely on personal experience, observations, and/or quantitative assessments to make this determination. Where available, quantitative data such as is found in the Natural Conditions Database (Overton et.al. 1995), could be used. For example, a bank stability rating that is greater than the standard error in the Database could be used to define "large departure". Where observational data is used for this determination, specialists should use photographs and/or descriptions of the observed conditions related to desired conditions to support the need for changing management and/or use indicators.

B2. Development and Implementation of Adaptive Management Adjustments (Block 5): If adaptive management adjustments are warranted, the resource manager develops these actions in collaboration with the permittee(s) and others, as appropriate. The adaptive actions are implemented through annual authorizations or operating instructions issued by the resource manager. These actions typically include, implementation of additional or more restrictive annual

use criteria; change in season, timing, or duration of grazing; changes in numbers of livestock; changes in herding or routing practices; changes in grazing rotations; closures or resting areas from grazing; changes in salting and watering practices, and changes in other livestock management practices and requirements.

Once adaptive management adjustments are developed and assigned, the resource manager, in collaboration with permittee(s) and others, as appropriate, must assess whether the adaptive management adjustments were implemented as designed during the following grazing period.

If adaptive management adjustments were implemented by the permittee(s), then a determination as to whether these adjustments achieved the annual grazing use indicator would be made the following grazing period (Block 1). If the adaptive management adjustments were effective in achieving the annual grazing use indicator, then management and monitoring would continue as planned (Blocks 2 and 3). If they were not effective, then the resource manager, in collaboration with permittee(s) and others, as appropriate, must determine what additional adaptive management actions are needed (return to Block 5). Adaptive management actions considered in the proposed action are described below.

- B3. Adaptive Management Adjustment Not Implemented (Block 6): If the adaptive management adjustments were not implemented, the resource manager must determine if the failure results from a design problem or changed condition, outside the control of the permittee(s). If there were problems with the design or ability to implement the adaptive management adjustments outside the control of the permittee(s), the resource manager and/or permittee(s) would revisit the design or selection of the adaptive management adjustment (return to Block 5).
- B4. <u>Determination of Non-compliance (Block 7)</u>: If failure to implement the adaptive management adjustment is not related to the design or inability to implement the adaptive action by the permittee(s), the resource manager would assess the need for an administrative action. If the resource manager determines that an administrative action is not warranted, additional changes or adaptive management direction should be considered (return to Block 5).
- B5. <u>Issue Notice of Non-compliance (Block 8)</u>: If failure to implement adaptive management adjustments is an issue of permittee(s) performance and compliance or is repetitive, then take appropriate action under the grazing regulations (36 CFR Part 222.4), Forest Service Manual direction (FSM 2231.6), and Forest Service Handbook direction (FSH 2209.13 sec. 16 & R4 FSH 2209.13 sec. 16).

This AMP is a component of the grazing permit that authorizes grazing use on National Forest System lands. The AMP implements management direction designed to achieve the goals and directives identified in the Forest Plan and allotment level NEPA analysis and related decisions.

Annually, Agency personnel meet with the grazing permittees to evaluate management activities and accomplishment of the grazing objectives. During these annual meetings, the previous year's grazing use and monitoring is reviewed, and annual operating instructions (AOI) are developed for the following grazing season. The AOI adapts management direction from the AMP, term grazing permit, and NEPA analysis and related decisions to the current conditions and expectations for the grazing season. The AOI sets the stage for the on-the-ground application of management direction for livestock grazing on the allotment. The AOI are used to implement direction within the context of the existing allotment specific decisions and the Agency's administrative authority established by law and regulation. Actions implemented through the AOI must be consistent with the direction evaluated in the existing NEPA analyses and/or the existing administrative authority of the Agency.

Adaptive management actions may be implemented as long as they are consistent with existing NEPA analyses and decisions and/or the administrative authority of the Forest Service. The administrative authority of the Forest Service is described in Title 36 of the Code of Federal Regulations, part 222; and in Forest Service Manuals and Handbooks. Courses of action that would be considered if monitoring did not indicate progress toward desired future conditions, particularly in light of the constraints discussed above are described in the following section. Such changes would generally be determined in advance and documented in the AOI describing authorized management actions for the upcoming grazing season. Additional NEPA analysis would not be required.

Adaptive management actions should be applied where:

- Monitoring shows management objectives are not being achieved or that trend towards achieving desired conditions is not improving or improving at an adequate rate.
- Annual indicators of grazing use or grazing standards are not met.
- Climatic events, fire, flood or uses and activities detrimentally impact resource conditions and a modification of grazing use is needed to provide for recovery of the site.

Implementation of adaptive management actions will be consistent with the direction established in the December 19, 2005, Forest Plan Grazing Implementation Guide (1920/2200 Memo to District Rangers signed by SE Idaho Forest Supervisors on Dec. 19, 2005). Adaptive actions may be needed and applied in both the short-term and long-term. Adaptive management actions may be implemented singly or as a set of management actions. Short-term actions will be implemented through the AOI. Modifications to the AMP and/or term grazing permit should be considered where monitoring shows that these actions need to be continued in the long-term or are implemented repeatedly or consistently over time.

The following table lists and describes the probable actions that would be considered and implemented under adaptive management. However, it is not intended to exclude other actions which may be authorized by the grazing permit or under authority of 36 CFR 222, Forest Service Manuals and Handbooks, and other laws and regulations as they exist or may be enacted.

Table of Potential Adaptive Management Actions	Authority
1. Modify the terms and conditions of a permit to conform to current situations brought about by changes in law, regulation, executive order, development or	36 CFR 222.4
revision of an allotment management plan, or other management needs.	
2. Modify the seasons of use, numbers, kind, and class of livestock allowed or the allotment to be used under the permit, because of resource condition, or permittee request.	36 CFR 222.4 (Change in livestock kind will require additional NEPA
3. Adjustments to sheep numbers and seasons of use.	evaluation.) NSEIS, p. 2-11
4. Implement periods of rest for the allotment or areas within the allotment.	NSEIS, p. 2-11
 5. Closure of grazing areas within the allotment. 6. Implementation of additional grazing restrictions. Includes: annual grazing use indicators (end of season and/or within season), salting practices, herding practices, and other management practices. 	NSEIS, p. 2-11 NSEIS, p. 2-11; FLRMP p. III-44 - 47
7. Alteration of trailing routes (timing and location).	NSEIS, p. 2-11; FLRMP p. III-44 - 47
8. Adjust grazing to address conflicts with other resource uses.	FLRMP, p. III- 44 - 47
9. Adjust grazing to provide for maintenance or restoration of aquatic and riparian processes and functions and beneficial uses.	FLRMP, p. III- 44 - 47
10. Coordinate grazing with timber harvest and forest regeneration activities.	FLRMP, p. III- 44 - 47
11. Temporary corrals.	NSEIS p. 2-12 & associated RODs.
12. Structural range improvements and handling facilities (water developments, fences, permanent corrals, etc.	Will require additional NEPA evaluation.
13. Vegetation treatments (prescribed fire, brush control, seedings, etc.) implemented to achieve management objectives and desired conditions.	Will require additional NEPA evaluation.

- 1. Modification of Terms and Conditions of the Grazing Permit. Term grazing permits may be modified at the request of the permit holder to adjust the permit to his/her ranch operation. It may also be modified to achieve consistency with changes in law and regulation, Forest Plan direction, NEPA analysis and related decisions, AMP direction, monitoring results, etc. Permit modifications are administrative actions and do not require additional NEPA unless they are inconsistent with existing NEPA analyses and decisions. Permit modifications may include the actions described below.
- 2. Modify the seasons of use, numbers, kind, and class of livestock allowed or the allotment to be used under the permit, because of resource condition, or permittee request. This action may include changing the timing, duration and intensity of grazing use, class of livestock grazed (ewes with lambs, dry ewes, and rams), changes in allotment boundaries, etc. without additional NEPA as long as these actions are consistent with current NEPA analysis and related decisions. Changes in kind of livestock such as changing from sheep to cattle use will require additional NEPA analysis. These changes may be implemented at the request of the permittee to adapt grazing to his/her ranch operation or they may be the result monitoring and the need to adapt management to changing conditions using actions such as those described below to achieve resource desired conditions and or resolve conflicts in resource uses.
- **3a. Modify Season of Use.** As appropriate, adjust the season of use for the allotment or areas within an allotment to reduce grazing impacts. These actions include shortening the period of use to reduce or eliminate grazing impacts during periods where plants or other resources are most susceptible to damage, or avoid conflicts with other uses such as during periods of high recreation use. They may include: changing the season of use to avoid grazing impacts or conflicts with critical resource needs of TES species and other wildlife, adjusting the season of use at the request of the permittee to provide a better fit to his/her ranch operation, adjusting the season of use to take advantage of the availability of additional forage through extending the grazing season, and adapting the grazing season in response to seasonal variations in climate and productivity such as during periods of drought. Adjustments to stocking and season of use may be considered jointly or separately as appropriate.
- **3.b Modify Stocking.** As appropriate, adjust authorized or permitted livestock numbers during all or a portion of the grazing season to match grazing use to resource conditions and productivity. Adjustments to stocking and season of use may be considered jointly or separately as appropriate.
- **4. Rest (i.e. closure to grazing for a full year).** Rest the allotment or areas within the allotment for a specific period of years or on a periodic rotation where monitoring shows that trend towards achieving desired conditions are not stable,

improving, or improving at an adequate rate. May also be implemented where fire, flood, etc; detrimentally impact resource conditions or where treatment activities require a period of rest to provide for recovery of the site. Where this occurs, specific recovery criteria for when grazing will be allowed should be specified.

- **5. Closure of Areas.** Close areas within allotments where monitoring shows that desired conditions cannot be met while sustaining grazing use. This may include alteration of allotment boundaries or identification of specific areas within an allotment where livestock grazing will not be allowed. Modify the AMP and term grazing permit to identify the change in the allotment boundary or the area closure.
- **6. Grazing Restrictions Modification of Indicators of Annual Grazing Use.** Annual grazing use indicators generally consist of measures of allowable grazing use including: forage utilization limits, woody species utilization limits, streambank disturbance limits, soil disturbance limits, once-over grazing, open herding, one-time use of bedding areas, one-time use of watering areas, location of nooning areas, location of watering areas, location of bedding areas, camp locations, length of stay at camp locations, corral locations, use limits around corrals, season and duration of use, etc. These indicators of livestock use may be modified or other indicators identified as needed to facilitate achievement of objectives and desired conditions. Levels of acceptable use such as forage utilization are set for some of these practices in the Forest Plan and/or the NSEIS. Where specific allowable use limits are set in the Forest Plan or in the NSEIS and ROD, they may be modified, if needed, to be more restrictive without additional NEPA analysis.
- **6a. Grazing Restrictions Changes in Grazing Use Indicators.** Changes in end-of-year and in-season grazing use indicators may be made based on results of short-term and/or long-term monitoring. Indicators evaluated during monitoring are described in the AMP Monitoring Plan. Modification and/or implementation of these annual use indicators will be consistent with the direction established in the December 19, 2005, Forest Plan Grazing Implementation Guide (1920/2200 Memo, dated 12/19/2005, signed by Boise, Payette, and Sawtooth Forest Supervisors).
- **6b.** Grazing Restrictions Modification of Management Practices. This includes a range of management and herding practices that vary according to conditions and use that are found on individual grazing allotments. These practices may include specification of areas where trailing or open herding techniques are used, location of bedding and nooning areas, use of salt and mineral supplement, location and duration of use of herder camps, etc.
- **7. Alteration of grazing routes**. Alteration of designated trailing routes and route rotations to avoid resource damage, avoid use conflicts, reduce grazing

pressure in specific areas, improve distribution, access unused grazing areas, facilitate shipping, or facilitate rest or deferred rotation grazing.

- **8.** Adjust grazing to address conflicts with other resource uses. Modification of grazing use may be appropriate to prevent or manage conflicts with other uses such as dispersed recreation, coordinate with other management activities such as timber harvest and forest regeneration, or mitigate conflicts or impacts to other resources. Examples include management of impacts to roads and trails, herding and trailing practices around developed recreation sites, use of sheep grazing as a tool for noxious weed management and site preparation for reforestation, management of sheep camps, fire and noxious weed prevention, etc.
- 9. Adjust grazing to provide for maintenance or restoration of aquatic and riparian processes and functions and beneficial uses. This practice may involve use of the adaptive actions described in this section with the specific purpose of reducing grazing impacts or managing grazing use to achieve functioning riparian systems. The focus of these actions will be on ecological conditions or processes that may be impacted by grazing. They include managing for properly functioning riparian vegetation, bank stability, sedimentation, etc.
- **10.** Coordinate grazing with timber harvest and forest regeneration activities. This covers three areas of coordination actions. First, the potential for physical conflict between grazing and timber activities (harvest, thinning, site preparation, etc.) as the timber activities are implemented; second, the potential for physical damage to tree seedlings on new plantations or regeneration sites; and third, the potential for using grazing for vegetation management and site preparation to facilitate timber stand regeneration and reduce competition from other vegetation, (noxious weeds, brush, etc.). Coordination may include changing use routes, closing or resting areas for periods needed for regeneration, adjusting grazing intensity to remove competing vegetation prior to planting, etc.
- 11. Temporary Corrals. The location and use of temporary corrals has been provided for in the North Sheep FEIS. These corrals are composed of panels that may be erected at the time of shipping and removed afterwards. They include some permanent structures required to support the corral panels and associated use. Use of fully portable corrals with no associated permanent structures may be considered in other locations as long as they are consistent with direction for management of heritage and archeological resources, Threatened, Endangered, and Sensitive (TES) species requirements, noxious weed management direction, recreation management direction, etc. identified in the Forest Plan and other appropriate NEPA analysis and related decisions.
- **12. Range Improvements Structural.** Structural range improvements include construction of water developments, fences, corrals and other permanent livestock handling facilities, trails, bridges, etc. These actions may be proposed as adaptive

management actions. Additional NEPA analysis will be required for these activities unless they are currently covered under existing NEPA analyses.

13. Vegetative Treatments – Nonstructural range improvements. Actions include implementing vegetation treatments to achieve desired rangeland conditions including prescribed fire, noxious weed treatment, seedings, aspen stand treatments, sagebrush manipulation, etc. These actions may be proposed as adaptive management actions. Additional NEPA analysis will be required for these activities unless they are currently covered under existing NEPA analyses such as is the case with noxious weed management activities.

IV. MANAGEMENT REQUIREMENTS

	MPC 4.1c Standards and Guideline				
Standard	Management actions—including mechanical vegetation treatments, salvage harvest, wildland fire use, prescribed fire, special use authorizations, and road maintenance—must be designed and implemented in a manner that would be consistent with the unroaded landscape in the temporary, short term, and long term. Exceptions to this standard are actions in the 4.1c road standards, below.				
	MPC 4.2 Standard and Guideline				
	Vegetation management actions—including wildland fire use, prescribed fire, and mechanical treatments—may be used to maintain or restore desired vegetation and fuel conditions provided they do not prevent achievement of recreation resource objectives.				

1. Livestock kind, class, numbers permitted, and season of use

Grazing use will be authorized through a term grazing permit which specifies variable numbers of sheep and a variable season of use which allows for up to 3,518 sheep head months of use within the grazing season of June 1 through September 30. Band size is limited to a maximum size of 1000 head of ewes with lambs (not to exceed 2500 total animals) or 1000 head of dry ewes. At the maximum band size, the season of use would be limited to 105 days within the permitted season of use.

Livestock Management: The following band management practices will be followed to reduce adverse grazing impacts on soil, vegetation, water, and wildlife resources.

- Reduce sheep activity around corrals at shipping time by bringing bands directly into the corral the night before shipping, and only allowing them out for water until handling is completed and they are being removed from the area.
- Avoid sheep bedding and nooning in areas of noxious weed infestation to the extent possible.
- Utilize open herding where possible, and once over grazing (No repeated use of an area previously grazed) except on designated driveways, travel routes, or where

- specifically authorized in the AOI. Criteria specific to the exceptions, as to duration of travel, location, nooning/bedding, and watering instructions will be displayed in the AOI.
- Utilize open herding where possible, and once over grazing (No repeated use of an
 area previously grazed) except on designated driveways, travel routes, or where
 specifically authorized in the AOI. Criteria specific to the exceptions as to duration of
 travel, locations, nooning/bedding, and watering instructions will be displayed in the
 AOI.
- Only one night/one time use of bed grounds is allowed.
- Livestock salting is prohibited in Riparian Conservation Areas(RCA's) Salting will be confined to bed grounds and placed in some form of containers or tarps.
- Sheep shall not be shaded or bedded within 200 yards of any stream or standing water unless topography is restrictive. Only one watering at each site is allowed, excluding troughs.
- Sheep shall be routed to avoid steep slopes, loose soil, rehabilitation sites and areas of active gullies.
- Sheep will be bedded away from known calving or fawning sites, and outside of a 30 acre buffer from known goshawk nest sites. These locations to be determined by the District Biologist.
- Avoid sheep nooning and bedding in areas of noxious weed infestation to the extent possible.
- Coordinate with the permittees to identify and manage noxious weed infestations within the allotment to prevent further expansion or reduce existing weed densities.

2. General Grazing Prescription

The Ewe/lamb band will enter the allotment at Lake Creek on or about 6/16. The band will trail above the Forest boundary at Lake Creek on or near the unnamed drainage below Taylor gulch, and grazing above Dip Creek toward Neal Canyon. The band will then cross Eagle Creek and graze toward North Fork. The band will then cross behind the SNRA headquarters and graze toward Silver Creek, where the band will ship at either the Boulder or Russian John temporary corral locations. The dry band will then return to Silver Creek and graze along the unused portions of the Boulder unit back toward Headquarters creek, and down into the North Fork canyon via the sheep bridge. The band will then graze up to the East Fork canyon, using Murdock on the way out. The sheep will exit the allotment via the sheep driveway on or about 9/14.

Each year the permittee will meet with the District Range Staff to discuss the previous years grazing summary to include both positive results due to management as well as areas needing further attention. Included in this discussion will be management area direction, authorized use, maintenance needs, routing, shipping dates, special instructions, and possible influences that may affect the current seasons grazing. These discussions are designed to manage annual grazing practices and associated impacts within the ability of the range to support and achieve resource management objectives

and values. From these meetings Annual Operating Instructions will be developed and distributed to the permittee and his agents.

Camp locations: Both pack strings and sheep wagons will be utilized to service the herders. Where wagon campsites become unavailable due to conflicts or loss of access, new locations will be approved by the District Ranger.

V. IMPROVEMENTS

Murdock corral #4 – SW1/4 Sec. 34, 5N. 17E. Wood planks/posts (1 acre)

North Fork Sheep Bridge #4107- NW1/4 Sec. 15, 5N. 17E. Wood plank/stringers


Maintenance of the improvements is the responsibility of the permittee.

VI. MONITORING

The use of the monitoring protocols identified in the following sections will be adaptive. Changes in protocols used may occur where new or more effective protocols are developed, identified protocols are found to not be effective in evaluating achievement of objectives, etc. The permittee conducts additional monitoring of grazing patterns and grazing effects as part of their management program. As appropriate, this information will be used along with Forest Service monitoring to evaluate achievement of desired conditions and annual grazing use.

Three general types of monitoring activities will be carried out on the Allotment. These are allotment field inspections or reviews, site-specific grazing use monitoring, and longterm effectiveness monitoring. The allotment reviews and site-specific use monitoring are Implementation Monitoring actions or monitoring of annual management to determine if specified management direction is being implemented and criteria or limits on annual grazing use are being met. Effectiveness monitoring evaluates achievement of long-term desired condition of the allotment resources. Site-specific annual use monitoring and long-term effectiveness monitoring are conducted at specific locations identified in this monitoring plan. These sites are chosen to represent areas of the allotment where grazing impacts are occurring and specific improvement of resource conditions are monitored. These sites are used to link annual grazing use monitoring to the achievement of specified desired conditions for allotment resources associated with grazing use. The allotment reviews are used to relate monitoring results from these sites to observed conditions and grazing use across the allotment in general. The combined use of these three monitoring actions provide adequate information to determine if management is being implemented as directed and that the results of that management are leading to achievement of management goals and objectives. Linking annual field inspections, site-specific use monitoring and long-term desired condition monitoring

allows the Ranger District to keep the number of more labor intensive monitoring sites and protocols to a manageable level.



Baker/North Fork Sheep Allotment Monitoring Sites

D. Kenney, Sawtooth N.F., 8/29/07

1. Implementation Monitoring

Annual grazing use limits listed in Section IV, Management Requirements, and the grazing standards and guidelines found on pages III-45 through III-47 in the Sawtooth National Forest Land and Resource Management Plan will be monitored for compliance within the allotment.

Seasonal monitoring and allotment administration will include field reviews of grazing practices, to include allotment inspections with permittees and their agents, review of annual grazing use information provided by the permittee, and pertinent information provided by Forest specialists working within the allotment. Allotment inspections or reviews are general inspections of grazing use across the allotment or large areas (pastures, watersheds, etc.) within the allotment. These reviews assess the general compliance with directions established in the grazing permit and annual operating instructions. They provide an overview of grazing impacts associated with the areas that have been grazed. Items such as maintenance of improvements, compliance with designated use and rest areas, trailing effects, overall use levels, etc. are evaluated during these field inspections. These field reviews will include evaluation of compliance with the grazing standards and other direction listed in the Management Requirements section of this AMP. This information will be evaluated on a yearly basis to insure management deficiencies are corrected and that successful management practices continue moving the

resource in the desired direction. The results of this monitoring will help determine the need for adaptive changes to livestock management.

Annual inspections will also focus on:

- Presence of livestock in closed or outside the permitted area or season.
- Extent and location of impact areas (e.g. salting, nooning, bedding, and trailing).
- General patterns of utilization and identification of areas of concentrated use or areas underutilized.
- Areas showing recent impacts due to drought or other weather related influences such as drying up of springs and seeps, increased use on vegetation or areas not generally used, and obvious increase in bare ground through lack of production.

a) Riparian Implementation Monitoring

The main purpose of this type of monitoring is to aid in within season grazing management. The monitoring protocol and location of the implementation monitoring sites are intended to measure whether the desired level of use has occurred on the allotment and whether Forest Plan standards and guidelines are met.

Protocols: Stubble height, woody vegetation regeneration, and stream bank alteration, per Burton et. al.2007. Stubble height measurements will be used where appropriate and streambank alteration and /or woody vegetation regeneration measurements at the remainder of sites.

Location/Number of Sites: See following table.

Timing/Frequency: Monitoring would be performed annually within two weeks of date of band routing through/past monitoring site. Implementation monitoring would not be performed at a site if it were determined that sheep were not grazed or routed through that site in that year.

Participants: Range staff would perform stubble height monitoring; either range or aquatics staff would take stream bank alteration measurements. (Note: Insert a table showing the following type of information.)

DMA Description and Location (GPS)	Site Type	Study Type/Protocol	Short Term Monitoring End-of-Season Indicator
DMA: Snow Creek Lat: .43° 48.273' Long: 114° 32.911'	Streambank	MIM (Burton et. Al. 2007)	Stubble Ht. > 4 in. Bank Alteration < 20% Woody Utiliz. < 25%
DMA: Murdock Creek Lat: 43° 48.978' Long: 114° 25.5626'	Streambank	MIM (Burton et. Al. 2007)	Stubble Ht. > 4 in Bank Alteration < 20% Woody Utiliz. < 25%

2. Effectiveness Monitoring

The main purpose of this type of monitoring is to determine condition/trend and progress toward achievement of desired conditions as described in the Sawtooth National Forest Land and Resource Management Plan and the North Sheep Final Environmental Impact Statement. Data from this monitoring will allow the Forest to document successful adaptive management and compliance with Forest Plan goals and objectives.

a) Riparian Effectiveness Monitoring

Protocols: Green line vegetation community composition, woody species regeneration, and stream bank stability; per Burton et al. 2007 and /or Pacfish Infish Biological Opinion monitoring protocols.

Location /Number of sites: Each effectiveness monitoring site will be co-located with an implementation monitoring site. Sites selected are:

- 1. Snow Creek- T.5N. R.16E, SE ¹/₄ Sec. 33- (GPS: N.43 48.273' W. 114 32.901'')
- 2. Murdock Creek- T. 5N. R.17E, SE1/4 Sec.34- (GPS: N.43 48.978' W. 114 25.5626').

Timing/ Frequency: After establishment, data collection at each site would follow at three to five-year intervals. Because this effort is intended to be long term monitoring, the data collection would not occur at any particular time relative to sheep routing or location on the allotment.

Participants: Range staff would perform vegetation community and woody regeneration monitoring in cooperation with the permittee. Range and/or aquatics staff would take stream bank stability measurements.

Resource conditions and trends, relative to desired conditions within the allotments, will be determined. Effectiveness monitoring will be used to determine if the grazing management practices are achieving the desired resource conditions.

Riparian	Last	Location	Protocol	Desired	Current
DMA	Monitored			Condition	Condition
Snow Creek	2007	Lat: .43° 48.273'	MIM	Late Seral	Mid Seral
		Long: 114° 32.911'	(Burton et.	Bank Stability =	Bank Stability =
			Àl. 2007)	78%	100%
Murdock Cr.	2006	Lat: 43° 48.978'	MIM	Late Seral	Early Seral
		Long: 114°	(Burton et.	Bank Stability =	Bank Stability =
		25.5626'	Al. 2007)	78%	88%

b) Photo Points

In order to collect additional information on the long-term effectiveness of the adaptive management strategy, additional photo points will be established at sites that reflect impacts from grazing. Repeat photos will be taken at these sites at a three to five year interval (per Hall 2002 protocol).

Photo Points

	Year Last Monitored	Location	Timing	Frequency	Responsibility
(to be added)					

3. Upland Effectiveness Monitoring

Nested Frequency Plots (USDA et. al. 1996): These are established to provide quantitative measurements of frequency and ground cover. Frequency is the presence or absence of a species on a quadrant by quadrant basis. It is a useful index for monitoring changes in vegetation over time and comparing different plant communities. Ground cover is the percentage of material, other than bare ground, covering the land surface. These measurements will be made along the frequency belts using the sharp pins of the frame.

Line Intercept measurements of sagebrush crown canopy cover will be conducted along each leg of the Nested Frequency Plot. These measurements provide an estimate of the relative cover of the shrub species measured on the study site.

Photo-points will be used to evaluate changes in conditions of key vegetative communities at locations impacted from grazing use including upland mesic sites. Additional photo points may be added if needed. The Process for Assessing Proper Functioning Condition for Lentic Riparian-Wetland Areas, per TR 1737-11 1994 will be utilized as part of the monitoring starting in 2008.

Snow Creek: T.5N. R.16E. NE1/4 Sec.04 (GPS: 43 48.273', 114 32.901') Little Boulder Creek: T.5N. R.16E. NW1/4 Sec.02 (GPS: 43 47.978' 114 31.393')

Upland	Year Last	Location	Protocol	Desired	Current
DMA	Monitored			Condition	Condition
NFB-1	2005	Lat: 43 48.273'	Nested Frequency &	<30%	24%
Snow Cr		Long: 114 32.901'	Line Intercept	sagebrush	sagebrush
				cover	cover
				>70% Soil	77% Soil
				cover	cover
NFB-2	2005	Lat: 43 47.978'	Nested Frequency &	<30%	32%
Little		Long: 114 31.393'	Line Intercept	sagebrush	sagebrush
Boulder Cr.				cover	cover
				>70% Soil	85% Soil
				cover	cover

Additional photo-points or ground cover measurements will be installed in 2008 within the key vegetative communities addressed in Appendix C of the North Sheep FEIS. These will include locations selected to evaluate areas with trailing concerns and upland seeps and springs.

4. Aspen Monitoring


Aspen stands selected for monitoring have a history of sheep use. Monitoring will focus on:

- Aspen regeneration- stems per acre. (Belt transect)
- Aspen browsing- utilization of terminal leader. (Belt transect)
- Domestic vs. wild ungulate use-Fecal count/browsing height. (Ocular estimate)
- Ground cover- Bare ground. (Ocular estimate)

Transects will be read every 3-5 years

Aspen	Year Last	Location	Desired Condition	Current Condition
DMA	Monitored	UTM		
Stand 1	2006	See Map	>500 Stems/Acre < 1" dbh	691 stems/ac < 1" dbh
Silver Cr.			< 20% leader use	0% leader use
Stand 2	2006	See Map	>500 Stems/Acre < 1" dbh	1636 stems/ac < 1" dbh
Snow Cr.			< 20% leader use	15% leader use (wildlife
				only)
Stand 3	2006	See Map	>500 Stems/Acre < 1" dbh	473 stems/ac < 1" dbh
Boulder Cr		_	< 20% leader use	15% leader use (wildlife
				only)
Stand 4	2006	See Map	>500 Stems/Acre < 1" dbh	764 stems/ac < 1" dbh
Boulder Cr			< 20% leader use	33% leader use (wildlife
				only)

a) Aspen Monitoring Location Map



VII. References

- Bartos, Dale L. 2001. Landscape Dynamics of Aspen and Conifer Forests, in Sustaining Aspen in Western Landscapes: Symposium Proceedings. USDA Forest Service, Rocky Mountain Research Station. RMRS-P-18.
- Burton, David; et al. 2002. Aspen Delineation Project Protocol for Recording Aspen Location and Condition. USDA Forest Service, Pacific Southwest Region.
- Burton, Timothy A., Cowley, Ervin R., and Smith Steven J. 2007. Monitoring Stream Channels and Riparian Vegetation-Multiple Indicators. Technical Bulletin 2007-01.USDI Bureau of Land Management, Idaho State Office.
- Mueggler, Walter F. 1988. Aspen Community Types of the Intermountain Region. USDA Forest Service, Intermountain Research Station, GTR INT-250.
- Overton, C. Kerry, McIntyre, John D., Armstrong, Robyn, Whitwell, Shari L., and Duncan, Kelly A. 1995. User's Guide to Fish Habitat: Descriptions that Represent Natural Conditions in the Salmon River Basin, Idaho. USDA Forest Service. Intermountain Research Station, GTR INT-322.
- USDI BLM, 1994. Revised 1998. Prichard, D., C. Bridges, S. Leonard, R. Krapf, and W. Hagenbuck. 1994. Riparian area management: process for assessing proper functioning condition for Lentic riparian-wetland areas. TR 1737-11. Bureau of Land Management, BLM/SC/ST-94/008+1737, Service Center, CO. USDI.
- USDA Forest Service, USDA Natural Resource Conservation Service, USDI Bureau of Land Management. 1996. Sampling Vegetation Attributes Interagency Technical Reference. BLM/RS/ST-96/002+1730.
- USDA Forest Service, USDA Natural Resource Conservation Service, USDI Bureau of Land Management. 1996. Utilization Studies and Residual Measurements Interagency Technical Reference. BLM/RS/ST-96/004+1730.
- USDA Forest Service, Pacific Southwest Region. 2004. Browsed Plant Method for Young Quaking Aspen An Annual Monitoring Method for Determining the Incidence of Use on Sprouts and Young Plants During the Growing Season

USDA Forest Service, 2005. Forest Plan Grazing Implementation Guide. Rangeland Resources Grazing Use Standards and Guidelines - Boise, Sawtooth, and Payette National Forest Land and Resource Management Plans. 8 pages. (Cover Memo dated December 19, 2005). Document dated November 3, 2005.

Winward, Alma H. 2000. Monitoring The Vegetation Resources In Riparian Areas. Gen. Tech. Rep. RMRS-GTR-47. Ogden, UT. USDA Forest Service, Rocky Mountain Research Station.