



Photo credit: Maria Luiza Delgado Assad, Fishbase.

Current Status:

KEY INFORMATION

Areas of Concern

Central and South America.

Year Identified as “Species of Concern”

1988 (removed in 1997, returned to list in 1999).

Factors for Decline

- Bycatch
- Fishing
- Habitat degradation

Conservation Designations

IUCN: Critically Endangered

AFS: Endangered

Species of Greatest Conservation Need:

TX, FL

Demographic and Genetic Diversity Concerns:

Simpfendorfer (2000) estimated a relatively slow intrinsic rate of increase for *Pristis perotteti* was from 0.05 to 0.07 per yr, with population doubling times of 10.3 to 13.6 years. Musick et al. (2000) noted that intrinsic rates of increase less than ten percent (0.1) were low, and make a species particularly vulnerable to excessive mortalities and rapid population declines, after which recovery may take decades.

P. perotteti historically inhabited warm-temperate to tropical marine waters in the Atlantic and eastern Pacific Ocean, possibly in the eastern Mediterranean, and freshwater habitats in Central and South America and Africa. Historical occurrences of largetooth sawfish in North America were much more limited than those of the related smalltooth sawfish and were strictly confined to shallow (< 33 feet), near-shore, warm-temperate and tropical (>64-86 °F), estuarine localities, partly enclosed lagoons, and similar situations. Thorson noted large catches of *P.*

perotteti during visits to Lake Nicaragua in 1963 (Cook *et al.*, 2006). Directed fisheries removed an estimated 60,000 to 100,000 sawfishes between 1970 and 1975 (Thorson 1976), and sawfish are now rare in Nicaragua. In the United States, largetooth sawfish were reported mainly along the Texas coast and east into Florida waters, but now it is considered extirpated in the United States. The last confirmed largetooth sawfish reported in US waters was in 1941 in Florida and 1943 in Texas (Burgess and Curtis 2003).

Existing Protections and Conservation Actions: Largetooth sawfish are a prohibited species in Florida, Louisiana, and Alabama. Texas Parks and Wildlife Division has listed largetooth sawfish as endangered under the Parks and Wildlife Code Chapter 68, due to the extreme difficulty that fishermen have in distinguishing the smalltooth sawfish from the largetooth sawfish.



Species of Concern

NOAA National Marine Fisheries Service

Factors for Decline:

Sawfish are extremely vulnerable to overexploitation due to their exceptional propensity for entanglement in net gear, their restricted habitat, and their low intrinsic rate of increase. Habitat degradation likely impacts the species given their inshore distribution.

Status Reviews/Research Underway:

In 2000, NMFS denied a petition to list them under the Endangered Species Act (65 FR 12959).

Brief Species Description:

Sawfishes are generally a tropical marine and estuarine **elasmobranch**. All modern sawfishes appear in some respects to be more shark-like than ray-like, with only the trunk and especially the head ventrally flattened. The presence of a rostrum having laterally protruding teeth separates sawfishes from skates and rays (Figure 1). The rostrum has a saw-like appearance and hence the name of sawfish. The largetooth sawfish and the smalltooth sawfish (*P. pectinata*) are similar in appearance. The two species can usually be differentiated by noting the number of teeth on one side of the rostrum. *P. perotteti* can have between 14 and 21 rostral teeth on one edge of the saw whereas *P. pectinata* usually has 23 to 34 (Compagno and Last 1999).

These two species can also be distinguished by noting that the first dorsal fin in *P. perotteti* originates anterior to the pelvic fins, while in *P. pectinata* the first dorsal fin originates along the same axis as the pelvic fins. The pectoral fins of *P. perotteti* are proportionally larger than those of *P. pectinata*. Only *P. perotteti* has a distinct lower lobe on its caudal fin (Compagno and Last 1999). Maximum size of *P. perotteti* is 21.3 feet (6.5 m) length with weights between 1100 and 1300 pounds (500 to 600 kg) (Thorson 1976). Studies of *Pristis perotteti* in Lake Nicaragua (Thorson 1976) report litter sizes of 1 to 13 individuals, with a mean of 7.3 individuals. The gestation period for largetooth sawfish is approximately 5 months, and females likely produce litters every second year.




Figure 1. A comparison of smalltooth sawfish rostrum (left) and largetooth sawfish rostrum (right). Photo courtesy George Burgess, Florida Museum of Natural History.

References:

- Burgess, G.H. and T.H. Curtis. 2003. Temporal reductions in the distribution and abundance of U.S. Atlantic sawfishes. Abstract: ASIH Annual Meeting. Manaus, Brazil.
- Compagno, L. and P. Last. 1999. Pristidae. p. 1410. In: Carpenter, K. and V. Niem (eds.), FAO Identification Guide for Fishery Purposes. FAO, Rome.
- Musick, J.A., et al. 2000. Fisheries 25:6-30.
- National Marine Fisheries Service. 2006. Recovery Plan for Smalltooth Sawfish. Smalltooth Sawfish Recovery Team. Silver Spring, Maryland.
- Simpfendorfer, C.A. 2000. *Env'tl Biol Fish* 58:371-377.
- Thorson, T. 1976. The status of the Lake Nicaragua shark. In: Investigations of the ichthyofauna of Nicaraguan lakes. T.B. Thorson ed. Univ NE-Lincoln. p. 561.
- Thorson, T.B. 1982. *Env. Bio. Fish.* 7: 207-228.

Contact Information

For largetooth sawfish, contact:

Shelley Norton
NOAA Fisheries, Southeast Region
Protected Resources Division
263 13th Avenue South
St. Petersburg, FL 33701
(727) 824-5312

Shelley.Norton@noaa.gov

<http://www.nmfs.noaa.gov/pr/species/concern>

For Species of Concern, contact

NOAA Fisheries
Office of Protected Resources
1315 East West Highway
Silver Spring, MD 20910
(301) 713-1401

soc.list@noaa.gov