

Appendix D

Sample indicators

- Proportion of verified TB cases reported to the local health jurisdiction within 1 working day from treatment start date.
- Demonstrated ability (organization, staffing, resources, and facilities) to carry out the core components of a TB control program.
- Number of TB cases identified per 100,000 people.
- Proportion of cases with complete data on key variables (homelessness, injecting drug use, non-injecting drug use, excess alcohol use)
- Proportion of pulmonary or laryngeal TB cases > 12 years of age with sputum culture obtained.
- Proportion of TB cases started on the recommended 4-drug regimen.
- Proportion of sputum smear-positive pulmonary or laryngeal TB cases initiating treatment in < 7 days of specimen collection.
- Proportion of sputum culture-positive TB cases with documented conversion to sputum culture-negative within 90 days of initiation of treatment.
- Proportion of TB cases for whom DOT is recommended who receive DOT throughout the course of treatment.
- Proportion of TB cases for whom DOT is recommended who receive inappropriate SAT throughout the course of treatment.
- Proportion of TB cases who complete treatment in < 12 months.
- Proportion of TB cases who default prior to completing treatment.
- Proportion of sputum smear-positive cases with at least one contact identified.
- Proportion of identified contacts to smear-positive cases who complete evaluation for TB infection or disease.
- Proportion of infected contacts to pulmonary cases who started treatment for LTBI.
- Proportion of infected contacts to pulmonary cases started on treatment for LTBI who complete treatment.
- Number of TB cases in children 0-4 years old.
- Number of persons who die with TB.

*As developed by California's Tuberculosis Indicators Project