

LA-UR-07-2053

*Approved for public release;
distribution is unlimited.*

<i>Title:</i>	MCNP5 + Data + MCNPX Workshop
<i>Author(s):</i>	Forrest B. Brown, Albert C. Kahler, Gregg W. McKinney, Russell D. Mosteller, Morgan C. White
<i>Intended for:</i>	Workshop at M&C+SNA-2007, ANS Mathematics & Computation Topical Meeting Monterey, CA, 15-19 April 2007



Los Alamos National Laboratory, an affirmative action/equal opportunity employer, is operated by the Los Alamos National Security, LLC for the National Nuclear Security Administration of the U.S. Department of Energy under contract DE-AC52-06NA25396. By acceptance of this article, the publisher recognizes that the U.S. Government retains a nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or to allow others to do so, for U.S. Government purposes. Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy. Los Alamos National Laboratory strongly supports academic freedom and a researcher's right to publish; as an institution, however, the Laboratory does not endorse the viewpoint of a publication or guarantee its technical correctness.

MCNP5 + Data + MCNPX

Workshop

**Forrest Brown, Skip Kahler, Gregg McKinney,
Russ Mosteller, Morgan White**

Los Alamos National Laboratory

Abstract



MCNP5 + Data + MCNPX Workshop

F.B. Brown, A.C. Kahler, G.W. McKinney, R.D. Mosteller, M.C. White
Los Alamos National Laboratory

This workshop provides an overview of recent advances in Monte Carlo particle transport codes and data at Los Alamos National Laboratory.

The MCNP5 and MCNPX Monte Carlo codes are used worldwide for particle transport calculations for reactors, radiation shielding, accelerators, medical physics, dosimetry, homeland security, experiment analysis, and many other applications. Updates are provided on the status and recent progress for each code.

Transport codes are only as good as the cross-section data they use. One of the most significant computational events of this year is the introduction of the ENDF/B-VII nuclear datasets. The extensive work in evaluating, preparing, and testing the data over the past few years will be described, along with some remaining challenges. Plans for making the ENDF/B-VII data available to MCNP5 and MCNPX users will also be presented.

MCNP5 + Data + MCNPX Workshop



F.B. Brown, A.C. Kahler, G.W. McKinney, R.D. Mosteller, M.C. White

- **MCNP5 Advances** (Brown, .5 hr)
 - Status & current work
 - Performance: multi-core threading, clusters
- **ENDF/B-VII Data** (Kahler, Mosteller, White, 1.5 hr)
 - Why ENDF/B-VII; What was wrong with ENDF/B-VI?
 - Processing ENDF/B-VII neutron files into an MCNP Library
 - Data Testing with ICSBEP Benchmarks
 - What's improved, what deficiencies remain?
 - Plans for release of MCNP ENDF/B-VII libraries
- **MCNPX Advances** (McKinney, .5 hr)
 - Overview of MCNPX 2006-2007 Features
 - Decay Particle Production from Fission & Activation
 - Coupled Space, Energy, Time Spherical Weight Windows
 - Background Radioactive Sources

MCNP5 Advances

Forrest B. Brown

Monte Carlo Codes (X-3-MCC)
Los Alamos National Laboratory
fbrown@lanl.gov

- **MCNP5 Releases**
 - Status
 - New features
- **MCNP5/6 - Ongoing work**
 - MCNP6 / MCNPX merger
 - Major revisions to classes
 - Acceleration & convergence for criticality problems
- **MCNP5 Performance**
 - Compilers & Threading


MCNP5 Releases



- **Current, RSICC:** **MCNP5 / 1.40** **(Spring 2006)**
 - Lethargy plots for energy dependent tallies
 - Logarithmic interpolation for input data
 - Fission neutron multiplicity
 - Stochastic geometry
 - Source entropy with plots
 - Mesh tally plotter
 - New electron energy-loss straggling logic
 - Source particle type specification
 - Positron sources
- **Nearing completion:** **MCNP5 / 1.50** **(Spring 2007)**
 - Variance reduction for pulse height tallies (F8)
 - MAKXSF - Doppler broadening & temperature interpolation
 - Long filenames
 - Numerous minor bug fixes
- **Planned:** **new data** **(Summer 2007 ?)**
 - ENDF/B-VII.0 data libraries in ACE format

- **Pulse height tallies (F8) depend on collections of particles (e.g., the entire particle history)**
 - Two 1 MeV photons deposited in one cell registered as one 2 MeV count, not two 1 MeV counts
 - Assumes both photons have weight one
 - Not the case when using variance reduction
- **Create “trees” to keep track of relationship between individual particles**
 - Weight assigned to branches of the tree
 - Energy deposited and weight stored for each branch

- (S) Source node
- (V) Variance reduction node
- (P) Physical node
- (T) Terminal node



Pulse height tallies require knowledge of the entire particle history

E_{ij} is the energy deposited on branch_{ij}
 w_{ij} is the weight multiplier for branch_{ij}

- **Variance reduction permitted**

Geometry splitting
Weight Window
DXTRAN

Energy splitting
Exponential Transform
Weight Cutoff

Time splitting
Forced Collision
Implicit Capture

- **Variance reduction NOT permitted**

- Neutrons
- Electron specific variance reduction that controls types and how many specific types of electrons are produced:


Bremsstrahlung


electron-induced x-rays

photon-induced secondary electrons

knock-on electrons

- 100 cm x 100 cm cylinder
- 10 cm thick disks
- Two Aluminum disks followed by one Oxygen disk
- 5 MeV isotropic photon source at the base
- Tally in final Al slab (90-100 cm)





- **Traditional MAKXSF functions**
 - Convert cross-section libraries to/from ASCII & binary
 - Copy entire libraries to new files
 - Copy selected nuclide tablesets to new libraries
 - Create new **xsdir** file for the new libraries
- **Create nuclide table-sets at new temperatures [new]**
 - Doppler broaden resolved resonance data to higher T
 - Interpolate unresolved resonance probability tables to new T
 - Interpolate S(α, β) thermal data to new T

- Taken from "doppler" code by MacFarlane, taken from NJOY
 1. R.E. MacFarlane & P. Talou, "DOPPLER: A Utility Code for Preparing Customized Temperature-Dependent Data Libraries for the MCNP Monte Carlo Transport Code", unpublished (Oct 3, 2003)
 2. R.E. MacFarlane & D.W. Muir, "The NJOY Nuclear Data Processing System, Version 91", LA-12740-M (1994).
- Doppler
 - Doppler broaden the resolved resonance data to new (higher) temperature
 - Temperatures can be specified in degrees-K or in MeV
 - Only need base cross-section, at lower temperature
- Interpolation
 - For **S(α, β) thermal data or unresolved resonance probability table** data
 - Must have existing datasets at BOTH lower & higher temperature

- **Mosteller, MacFarlane, Little, White, "Analysis of Hot and Cold Kritz Benchmarks With MCNP5 and Temperature-specific Nuclear Data Libraries", LA-UR-03-7071 (2003).**
 - Separate nuclear data sets were generated at 245 C using DOPPLER and NJOY
 - Basic data were taken from ENDF/B-VI Release 6 (ENDF66)
 - **MCNP5** calculations were performed for the hot 2-D benchmarks, and the results were compared
 - Each calculation employed 550 generations with 10,000 neutron histories per generation
 - Results from first 50 generations were discarded, giving 5,000,000 active histories for each case

Case	Library	keff	Δk (vs NJOY)
Kritz:2-1	NJOY	0.9914 ± 0.0003	—
	DOPPLER	0.9911 ± 0.0003	-0.0003 ± 0.0004
	new MAKXSF	0.9913 ± 0.0003	-0.0001 ± 0.0004
Kritz:2-13	NJOY	0.9944 ± 0.0003	—
	DOPPLER	0.9942 ± 0.0003	-0.0002 ± 0.0004
	new MAKXSF	0.9940 ± 0.0003	-0.0004 ± 0.0004
Kritz:2-19	NJOY	1.0005 ± 0.0003	—
	DOPPLER	1.0009 ± 0.0003	0.0004 ± 0.0004
	new MAKXSF	1.0004 ± 0.0003	-0.0001 ± 0.0004

- **MCNP6 - MCNPX Merger**
 - Happening now
 - Preserve features of both codes
 - Internal versions available this year
 - Will undergo extensive cleanup, testing, V&V
- **MCNP Classes**
 - Major upgrades to lectures
 - CD format, with extra references
- **Acceleration & convergence testing for criticality problems**
 - Improved tests using Keff & Shannon entropy of source distribution
 - Wielandt acceleration (paper this conference)
 - Fission matrix (& variants)

- MCNP5 performance - both serial & parallel - depends strongly on the Fortran-90 compiler & options used
 - Runtime factors of 2-4x with different compilers on same hardware
 - Runtime factors of 2-4x with different options on same hardware & compiler
- Parallel performance
 - MCNP5 has always supported parallel calculations with **message-passing** (MPI, PVM) & **threading** (OpenMP)
 - Prior to mid-2006, Fortran compilers for Windows/Linux/Mac did a terrible job at threading. We recommended using only MPI.
 - Recently, using OpenMP threading with Intel compilers on Windows/Linux/Mac shows excellent speedups -- nearly 2x on dual-cores.
- Examples & demo for MacBook Pro (Core2 Duo)
 - KCODE: BAWXI2 benchmark, kcode 5000 1 10 50
 - Fixed source: inp12, oil-well log, mode n, nps 105000

MCNP5 - Compilers & Threads

- Mac G5, 2.5 GHz PowerPC, OS X 10.3.9

Fortran-90	Regression tests OK ?	These runs correct ?	Wall-clock time (sec)	
			KCODE	Fixed Source
g95 -O0	Yes	Yes	535	164
g95 -O1	Yes	Yes	363	107
g95 -O2	Yes	Yes	370	113
absoft -O0	Yes	Yes	324	109
absoft -O1	Yes	Yes	352	121
absoft -O2	aborted	-	-	-
ibm xlf -O0	Yes	Yes	237	73
ibm xlf -O1	Yes	Yes	236	73
ibm xlf -O2	Yes	Yes	154	46

g95 - 0.90, absoft - 9.2, xlf - 8.1

MCNP5 - Compilers & Threads

- MacBook Pro, 2.16 GHz Intel Core2 Duo, OS X 10.4.8

Fortran-90	Regression tests OK ?	These runs correct ?	Threads	Wall-clock time (sec)	
				KCODE	Fixed Source
g95 -O1 (on G5), Run using Rosetta	Yes	Yes	1	727	281
g95 -O0	Yes	Yes	1	379	135
g95 -O1	Yes	Yes	1	277	87
g95 -O2	Yes	Yes	1	272	86
intel -O0 -openmp	Yes	Yes	1	256	105
			2	142	57
intel -O1 -openmp	No	Yes	1	122	55
			2	65	29
intel -O2 -openmp	No	Yes	1	113	54
			2	60	28

g95 - 0.91, intel - 9.1

MCNP and the ENDF/B-VII Cross Section Library

A. C. (Skip) Kahler, R. D. Mosteller, M. C. White

Theoretical Division & Applied Physics Division
Los Alamos National Laboratory

akahler@lanl.gov, mosteller@lanl.gov, morgan@lanl.gov

Presented at the MCNP Workshop at MCD + SNA 2007
Monterey, CA April 15, 2007



EST. 1943
This work was carried out under the auspices of the National Nuclear Security Administration of the U.S. Department of Energy at Los Alamos National Laboratory under Contract No. DE-AC52-06NA25396.

LA-UR-07-2053



The ENDF/B-VII Cross Section Library

- Why ENDF/B-VII; What was wrong with ENDF/B-VI?
- Processing ENDF/B-VII Neutron Files into an MCNP Library.
- Data Testing with ICSBEP Benchmarks.
- What's Improved, What Deficiencies Remain.
- Plans for Release of ENDF/B-VII Libraries for MCNP.

Why ENDF/B-VII; What was wrong with ENDF/B-VI?

- Does anyone know what ENDF/B-VI is?
 - ◆ Originally released in 1990.
 - Followed by 8 “mods” over the next 11 years (including substantial changes to ^{235}U in mods 1, 3 and 5).
 - Was frozen in 2001 following release of ENDF/B-VI.8.
 - ◆ Changes to many structural nuclide files in mods 1 & 5.
 - ◆ Changes to thermal scattering kernels in mod 2.
 - ◆ Changes to ^1H and ^{238}U in mod 5.
 - ◆ New photon production data for some nuclides in mod 5 and later.
 - ◆ Some revised fission product evaluations throughout, but most fp data files remain as is (including many from ENDF/B-IV).
 - ◆ Extension from 20 MeV to 150 MeV for some nuclides in mod 6.
 - ◆ Confusing nomenclature – MCNPDATA uses .60c for mod 2, uses .66c for mod 6 and uses .62c for release 8.

Why ENDF/B-VII; What was wrong with ENDF/B-VI?

- ENDF/B-VI was significantly more complex than previous ENDF/B libraries.
 - ◆ “complex” meant physically more complete (good).
 - For example, the resolved resonance region description for ^{235}U expanded from 1 eV to 82 eV in ENDF/B-V to 10^{-5} eV to 2250 eV in ENDF/B-VI.0.
 - ◆ but “complex” also meant that greater resources (i.e., greater cost) are required to process and use this library (bad).
 - There were no dominant customers crying for ENDF/B-VI, coupled with a moribund domestic commercial reactor industry, which led to slow (or no) adoption of the new file.
 - ENDF/B-V (late 1970s/early 1980s);
 - ENDF/B-IV (early 1970s) had fast reactor support;
 - ENDF/B-III (late 1960s) had commercial reactor support.
 - Therefore, organizations that failed to adopt ENDF/B-VI are now using data that are likely to be at least 30 years old.

Why ENDF/B-VII; What was wrong with ENDF/B-VI?

- ENDF/B-VII is more complete than ENDF/B-VI.
 - ◆ 393 versus 328 neutron files;
 - ◆ 20 versus 15 thermal scattering files;
 - ◆ 3838 versus 979 radioactive decay files;
 - ◆ 163 versus 0 photonuclear files.
- Neutron files continue movement to isotopic data.
 - ◆ 390 isotopic files (carbon, vanadium and zinc are elemental).
 - ◆ 219 revised or new fission product evaluations.
 - primarily from international collaboration (WPEC-21 & 23) or developed at BNL (with KAERI collaboration).
 - ◆ have adopted existing files from other sources (e.g., JENDL-3.3) when judged appropriate.
 - ◆ revised evaluations benefit from new experimental data and from more sophisticated nuclear model codes.

Why ENDF/B-VII; What was wrong with ENDF/B-VI?



Available online at www.sciencedirect.com



Nuclear Data Sheets 107 (2006) 2931–3060

**Nuclear Data
Sheets**

www.elsevier.com/locate/nds



ENDF/B-VII.0: Next Generation Evaluated Nuclear Data Library for Nuclear Science and Technology

M.B. Chadwick,¹ P. Obložinský,^{2*} M. Herman,² N.M. Greene,⁶ R.D. McKnight,³ D.L. Smith,³
P.G. Young,¹ R.E. MacFarlane,¹ G.M. Hale,¹ S.C. Frankle,¹ A.C. Kahler,^{1,12} T. Kawano,¹ R.C. Little,¹
D.G. Madland,¹ P. Moller,¹ R.D. Mosteller,¹ P.R. Page,¹ P. Talou,¹ H. Trellue,¹ M.C. White,¹ W.B. Wilson,¹
R. Arcilla,² C.L. Dunford,² S.F. Mughabghab,² B. Pritychenko,² D. Rochman,² A.A. Sonzogni,²
C.R. Lubitz,⁴ T.H. Trumbull,⁴ J.P. Weinman,⁴ D.A. Brown,⁵ D.E. Cullen,⁵ D.P. Heinrichs,⁵ D.P. McNabb,⁵
H. Derrien,⁶ M.E. Dunn,⁶ N.M. Larson,⁶ L.C. Leal,⁶ A.D. Carlson,⁷ R.C. Block,⁸ J.B. Briggs,⁹ E.T. Cheng,¹⁰
H.C. Huria,¹¹ M.L. Zerkle,¹² K.S. Kozier,¹³ A. Courcelle,¹⁴ V. Pronyaev,¹⁵ S.C. van der Marck¹⁶

¹ Los Alamos National Laboratory, Los Alamos, NM 87545

² National Nuclear Data Center, Brookhaven National Laboratory, Upton, NY 11973-5000

³ Argonne National Laboratory, 9700 South Cass Ave, Argonne, IL 60439-4842

⁴ Knolls Atomic Power Laboratory, P.O. Box 1072, Schenectady, NY 12301-1072

⁵ Lawrence Livermore National Laboratory, P.O. Box 808, Livermore, CA 94551-0808

⁶ Oak Ridge National Laboratory, P.O. Box 2008, Oak Ridge, TN 37831-6171

⁷ National Institute of Standards and Technology, Gaithersburg, MD 20899-8463

⁸ Guerttner LINAC Laboratory, Rensselaer Polytechnic Institute, Troy, NY 12180-3590

⁹ Idaho National Laboratory, P.O. Box 1625, Idaho Falls, ID 83415-3860

¹⁰ TSI Research Corp., 312 S. Cedros Ave, Solana Beach, CA 92067

¹¹ Westinghouse Electric Corp., 4350 Northern Pike, Monroeville, PA 15146

¹² Bettis Atomic Power Laboratory, P.O. Box 79, West Mifflin, PA 15122

¹³ Atomic Energy of Canada Ltd., Chalk River, ON K0J 1J0, Canada

¹⁴ Centre d'Etudes Nucléaires de Cadarache, F-13108 Saint-Paul lèz Durance, Cedex, France

¹⁵ Institute for Physics and Power Engineering, Ploschad Bondarenko 1, 249 020 Obninsk, Russia and

¹⁶ Nuclear Research and Consultancy Group, P.O. Box 25, NL-1755 ZG Petten, The Netherlands

- The ENDF/B-VII.0 cross section library may be downloaded from Brookhaven National Laboratory's National Nuclear Data Center (NNDC),
<http://www.nndc.bnl.gov/exfor7/endf00.htm>.

(Received October 2, 2006)

Processing ENDF/B-VII Neutron Files into an MCNP Library

- The Los Alamos NJOY Nuclear Data Processing Program converts an ENDF-formatted file into an MCNP ACE file.
 - ◆ The base version of NJOY (currently NJOY99) is available from ORNL (RSICC) or the Nuclear Energy Agency;
 - ◆ Updates are available at <http://t2.lanl.gov/codes/njoy99/index.html>.
 - includes machine and compiler guidance plus sample problem I/O.
 - ◆ User information is available at <http://www.nea.fr/lists/njoy.html> or at <http://www-rsicc.ornl.gov/ENOTE/enotnjoy.html>.
- For each ENDF evaluation NJOY can create:
 - ◆ a unique, doppler broadened, linearly interpolable, unionized cross section mesh for all cross sections for that material;
 - ◆ angular distribution probability functions;
 - ◆ secondary energy spectra.

Processing ENDF/B-VII Neutron Files into an MCNP Library


- The most commonly used NJOY modules include:
 - ◆ MODER: ascii to binary conversion for efficient I/O (optional).
 - ◆ RECONR: resonance reconstruction, cross section linearization and energy mesh unionization at zero degrees.
 - ◆ BROADR: doppler broadening to User specified temperature, including energy mesh adjustment to maintain user specified linear-linear interpolation accuracy.
 - ◆ UNRESR: unresolved resonance processing (to support HEATR).
 - ◆ HEATR: heating and energy damage calculations (a standard part of MCNP ACE files, but not needed for criticality calculations).
 - ◆ PURR: unresolved resonance processing (probability tables).
 - ◆ THERMR: thermal scattering kernel processing (maybe in conjunction with the LEAPR module).
 - ◆ ACER: create an MCNP continuous energy (.c) and thermal kernel (.t) library files.

Processing ENDF/B-VII Neutron Files into an MCNP Library

- Independently compiled versions of NJOY (99.162 at Brookhaven National Laboratory's National Nuclear Data Center (NNDC) or 99.167 at LANL) have successfully created ACE-formatted files for all 393 ENDF/B-VII.0 neutron files.
- The ENDF/B-VII.0 cross section library may be obtained at <http://www.nndc.bnl.gov/exfor7/endf00.htm>.

Data Testing with ICSBEP Benchmarks

- ICSBEP = International Criticality Safety Benchmark Evaluation Project.
- Responsible for the content of the International Handbook of Evaluated Criticality Safety Benchmark Experiments.
- The Handbook is updated and re-issued annually in the Fall.
 - ◆ For 2006 there are 442 evaluations with 3955 configurations.
- <http://icsbep.inel.gov/>



Data Testing with ICSBEP Benchmarks

- The Handbook has distinct categories for fissile material, physical form of that material and spectrum.
 - ◆ Fissile material categories include
 - HEU: > 60 wt.% ^{235}U ;
 - IEU: 10 wt.% < ^{235}U < 60 wt.%;
 - LEU: < 10 wt.% ^{235}U ;
 - Pu: mostly ^{239}Pu , some configurations with > 40 at.% ^{240}Pu ;
 - U233:
 - Mixed Pu-U:
 - Special Isotopes (e.g. ^{237}Np).
 - ◆ Categories for the physical form of the fissile material include:
 - Metal;
 - Compound (e.g. UO_2);
 - Solution (mostly nitrates and fluorides).

Data Testing with ICSBEP Benchmarks

- The Handbook has distinct categories for fissile material, physical form of that material and spectrum (con't).
 - ◆ Fission spectrum categories include:
 - Fast: > 50% of the fissions occur above 100 keV;
 - Intermediate: > 50% of the fissions occur between 0.625 eV & 100 keV.
 - Thermal: > 50% of the fission occur below 0.625 eV;
 - Mixed: < 50% of the fissions occur in each category.
 - ◆ In recent years the scope of the Handbook has expanded to include radiation transport measurements applicable to Criticality Alarm & Shielding Benchmarks as well as Fundamental Physics Measurements.


Data Testing with ICSBEP Benchmarks

- ICSBEP Nomenclature
 - ◆ A given benchmark is identified by its fissile material, its physical form, the dominant fission spectrum category and a sequence number.
 - Examples
 - HEU-MET-FAST-001: Godiva
 - HEU-MET-FAST-028: Flattop-28
 - PU-MET-FAST-001: Jezebel
 - PU-MET-FAST-006: Thor
 - U233-MET-FAST-001: Jezebel-233
- Although not a formal category, many benchmarks include metallic and/or non-hydrogenous reflectors.

Data Testing with ICSBEP Benchmarks


- LANL has developed MCNP models for hundreds of the ICSBEP benchmarks and performed critical eigenvalue calculations for them with various cross section data sets.
- Comparison of calculated critical eigenvalues illustrate areas of improvement and point to where future efforts to improve fundamental nuclear data are needed.
- Back to the original question –
 - ◆ why ENDF/B-VII, what was wrong with ENDF/B-VI?
- Representative results follow:

Data Testing with ICSBEP Benchmarks




- ENDF/B-VI.8 eigenvalues for Historical LANL "Fast" Benchmarks.
- Eigenvalues are biased low by several hundred pcm for all bare systems.
- A reflector bias is readily observed.

Data Testing with ICSBEP Benchmarks




- An ENDF/B-VII.0 success story.
- Calculated eigenvalues are significantly closer to unity than those obtained with ENDF/B-VI.8 cross sections.
- The reflector bias is virtually eliminated.

Data Testing with ICSBEP Benchmarks



- Calculated eigenvalues for uranium solutions are an ENDF/B-VI success story.
- Eigenvalue biases and trends observed in previous ENDF/B cross sections were eliminated starting with ENDF/B-VI.3.


Data Testing with ICSBEP Benchmarks



- Results with ENDF/B-VII.0 remain excellent, despite changes in the underlying $^{235,238}\text{U}$, ^1H and ^{16}O cross sections.


Data Testing with ICSBEP Benchmarks

Calculated LEU-COMP-THERM Eigenvalues
with ENDF/B-VI.8 Cross Sections




- ENDF/B-VI based LEU-COMP-THERM calculated eigenvalues are low and exhibit trends with ²³⁸U absorption fraction and fundamental lattice parameters.
- These are long-standing, historical deficiencies inherited from ENDF/B-V (and earlier).

Data Testing with ICSBEP Benchmarks




- LEU-COMP-THERM-006 (Japan) provides an excellent example of the ENDF/B-VI.8 calculated eigenvalue trend as a function of ^{238}U absorption.

Data Testing with ICSBEP Benchmarks




- LEU-COMP-THERM-007 (France) provides an excellent example of the ENDF/B-VI.8 calculated eigenvalue trend as a function of H/U ratio (lattice pitch).

Data Testing with ICSBEP Benchmarks




- ENDF/B-VII.0 based LEU-COMP-THERM calculated eigenvalues are significantly closer to unity.

Data Testing with ICSBEP Benchmarks



- ENDF/B-VI.8 Based Pu Solution System Calculated Eigenvalues are Biased High.
- There are no obvious trends in these eigenvalues when correlated against various system parameters (e.g. ²⁴⁰Pu at. %, leakage, ¹H absorption fraction, thermal fission fraction, Pu concentration, ...).

Data Testing with ICSBEP Benchmarks



- There has been little work on thermal Pu cross sections for ENDF/B-VII, hence the calculated eigenvalue deficiency remains.

Data Testing with ICSBEP Benchmarks

- Acknowledgements
- Preliminary ACE formatted cross section files derived from ENDF/B-VII.0 neutron and thermal scattering kernel files used in these calculations were processed with NJOY99.167 by R. E. MacFarlane. Bob has been and continues to be a significant contributor to LANL's ENDF/B-VII data testing effort.
- ENDF/B-VI.8 results were obtained using publicly available MCNPDATA files (.66c and .62c and associated ENDF/B-VI.2 thermal scattering kernels).

Data Testing with ICSBEP Benchmarks

- And now for the rest of the story ...

ENDF70 - ACE Data Based On ENDF/B-VI.0

- Expected Release in Late Summer (On Track!)
- Initial release will include continuous-energy neutron and photon data and S(alpha,beta) tables
 - ◆ Continuous-energy neutron library will include all 393 tables
 - Initial indications are promising that ENDF/B-VII.0 is “complete”
 - Data at 293.6, 600, 900, 1200 & 2500 K
 - Doppler broadening to other temperatures via MAKXSF
 - ◆ All 20 thermal scattering tables
 - Data at all evaluated temperatures
 - ◆ MCPLIB04 includes photoatomic data for 100 elements
 - No changes were made to the photon data between VI.8 and VII.0
- Incident charged particle and photonuclear tables will be released “at a later date”

Verification, Verification, Verification

The ENDF66 Success Story

- Zero (0!) bug reports since ENDF66 was released
- Our team consider themselves to be master data breakers
 - ◆ We have stressed the data, NJOY, MCNP and PARTISN to their limits and helped make their use more robust
 - ◆ Our friendly users add even more stress to the system
 - ◆ Most of our time on this project will be spent on the 80/20 rule
 - And it's actually more like 95/5
- It's All About Verification
 - ◆ A wide variety of tools are employed to check the data “six-ways-from-Sunday”
- ENDF70 is in the process of undergoing similar QA

Benchmark Anomalies and Persistent Problems in Nuclear-Data Testing for ENDF/B-VII.0

Russell D. Mosteller

Applied Physics Division
Los Alamos National Laboratory
mosteller@lanl.gov

Presented at the MCNP Workshop at MCD + SNA 2007
Monterey, CA April 15, 2007

OVERVIEW OF PRESENTATION

Identification of nuclear data libraries and details of calculations

Cases and results (all input specifications are taken from *International Handbook of Evaluated Criticality Safety Benchmark Experiments*)

- Zeus HEU-graphite benchmarks (^{235}U in unresolved resonance range)
- Unmoderated Zeus HEU benchmark (copper in fast range)
- Neptunium sphere benchmark (^{237}Np in fast range)
- Benchmarks of HEU nitrate solutions in heavy water (angular scattering distributions for ^2H)
- MOX lattice benchmarks (^{239}Pu in thermal range)
- Plutonium nitrate solution benchmarks (^{239}Pu in thermal range)
- HEU nitrate solution benchmark with Cd (^{113}Cd in thermal range)

Conclusions

NUCLEAR DATA LIBRARIES

Evaluated Nuclear Data File VII, Release 0 (ENDF/B-VII.0)

Evaluated Nuclear Data File VI, Release 8 (ENDF/B-VI)

Evaluated Nuclear Data File V (ENDF/B-V)

Japanese Evaluated Nuclear Data Library, version 3.3 (JENDL-3.3)

Joint Evaluated Fission and Fusion file, version 3.1 (JEFF-3.1)

MCNP5 CALCULATIONS

Each calculation employed at least 550 generations with 10,000 neutrons per generation

Results from first 50 generations were excluded from the statistics

Results therefore are based on at least 5,000,000 active histories for each case



The World's Greatest Science Protecting America


Applied Physics Division



CRITICALITY BENCHMARKS

Name	Spectrum	Handbook ID	Description
Zeus-1	Intermediate	HEU-MET-INTER-006, case 1	HEU platters moderated by graphite and reflected by copper
Zeus-2	Intermediate	HEU-MET-INTER-006, case 2	HEU platters moderated by graphite and reflected by copper
Zeus-3	Intermediate	HEU-MET-INTER-006, case 3	HEU platters moderated by graphite and reflected by copper
Zeus-4	Intermediate	HEU-MET-INTER-006, case 4	HEU platters moderated by graphite and reflected by copper
Unmod Zeus	Fast	HEU-MET-FAST-073	HEU platters reflected by copper
Np Sphere	Fast	SPEC-MET-FAST-008	Neptunium sphere reflected by HEU shells
HST-4	Intermediate	HEU-SOL-THERM-004, cases 1-2	Reflected spheres of HEU solution in heavy water
HST-4	Thermal	HEU-SOL-THERM-004, cases 3-6	Reflected spheres of HEU solution in heavy water
HST-20	Thermal	HEU-SOL-THERM-020, cases 1-5	Unreflected cylinders of HEU solution in heavy water
PNL-30	Thermal	MIX-COMP-THERM-002, case 30	Lattice of 469 MOX pins (2 wt.% PuO ₂) in water (1.7 PPM)
PNL-31	Thermal	MIX-COMP-THERM-002, case 31	Lattice of 761 MOX pins (2 wt.% PuO ₂) in water (687.9 PPM)
PNL-32	Thermal	MIX-COMP-THERM-002, case 32	Lattice of 195 MOX pins (2 wt.% PuO ₂) in water (0.9 PPM)
PNL-33	Thermal	MIX-COMP-THERM-002, case 33	Lattice of 761 MOX pins (2 wt.% PuO ₂) in water (1090.4 PPM)
PNL-34	Thermal	MIX-COMP-THERM-002, case 34	Lattice of 161 MOX pins (2 wt.% PuO ₂) in water (1.6 PPM)
PNL-35	Thermal	MIX-COMP-THERM-002, case 35	Lattice of 689 MOX pins (2 wt.% PuO ₂) in water (767.2 PPM)
PNL-1	Thermal	PU-SOL-THERM-021, case 7	Unreflected 14-inch sphere of plutonium nitrate solution
PNL-3	Thermal	PU-SOL-THERM-011, case 18-1	Unreflected 18-inch sphere of plutonium nitrate solution
PNL-4	Thermal	PU-SOL-THERM-011, case 18-6	Unreflected 18-inch sphere of plutonium nitrate solution
PNL-5	Thermal	PU-SOL-THERM-011, case 16-5	Unreflected 16-inch sphere of plutonium nitrate solution
PNL-6	Thermal	PU-SOL-THERM-021, case 3	Unreflected 15.2-inch sphere of plutonium nitrate solution
PNL-8	Thermal	PU-SOL-THERM-021, case 2	Unreflected 15.2-inch sphere of plutonium nitrate solution
PST-9	Thermal	PU-SOL-THERM-009, case 3a	Unreflected 48-inch sphere of plutonium nitrate solution
HST-49c20	Thermal	HEU-SOL-THERM-049, case 20	Reflected cylinder of HEU nitrate solution with Cd

ZEUS HEU-GRAphite BENCHMARKS



⇒ Cross sections for ^{235}U in the unresolved resonance region should be re-examined

UNMODERATED ZEUS BENCHMARK

Source	k_{eff}	Δk
Benchmark	1.0004 ± 0.0016	—
ENDF/B-VII.0	1.0116 ± 0.0003	0.0112 ± 0.0016
ENDF/B-VI	1.0082 ± 0.0003	0.0078 ± 0.0016
JENDL-3.3	1.0242 ± 0.0003	0.0238 ± 0.0016

$$|\Delta k| > 4\sigma$$

Benchmark contains no moderator and therefore has a fast spectrum

All three calculated results differ from the benchmark value for k_{eff} by more than 4 standard deviations

Results for k_{eff} with ENDF/B-V Cross Sections for Copper

Source	k_{eff}	Δk
Benchmark	1.0004 ± 0.0016	—
ENDF/B-VII.0	1.0001 ± 0.0003	-0.0003 ± 0.0016
ENDF/B-VI	0.9971 ± 0.0003	-0.0033 ± 0.0016
JENDL-3.3	1.0001 ± 0.0003	-0.0003 ± 0.0016


$$|\Delta k| > 2\sigma$$

Both JENDL-3.3 and ENDF/B-VII.0 now produce results for k_{eff} that are within a single standard deviation of the experimental value

Differences between ENDF/B-V and ENDF/B-VII.0 Cu cross sections have little net reactivity impact on previous Zeus benchmarks, which have intermediate spectra

⇒ Fast cross sections for Cu should be reviewed

NEPTUNIUM SPHERE REFLECTED BY HEU



Benchmark k_{eff}	Calculated k_{eff}		
	ENDF/B-VII.0	ENDF/B-VI	JENDL-3.3
1.0019 ± 0.0036	0.9951 ± 0.0003	0.9889 ± 0.0002	0.9967 ± 0.0002

$$\sigma < |\Delta k| \leq 2\sigma$$

$$|\Delta k| > 2\sigma$$

→ Fast cross sections for ^{237}Np should be reviewed

HEAVY-WATER SOLUTIONS

Case	Benchmark k_{eff}	Calculated k_{eff}		
		ENDF/B-VII.0 + JENDL-3.3 ^2H	ENDF/B-VII.0	JENDL-3.3
Reflected Spheres (HEU-SOL-THERM-004)				
1	1.0000 ± 0.0033	0.9967 ± 0.0004	0.9847 ± 0.0004	0.9918 ± 0.0004
2	1.0000 ± 0.0036	0.9919 ± 0.0004	0.9815 ± 0.0004	0.9873 ± 0.0004
3	1.0000 ± 0.0039	0.9985 ± 0.0004	0.9883 ± 0.0004	0.9944 ± 0.0004
4	1.0000 ± 0.0046	1.0012 ± 0.0004	0.9904 ± 0.0004	0.9971 ± 0.0004
5	1.0000 ± 0.0052	0.9994 ± 0.0005	0.9891 ± 0.0004	0.9956 ± 0.0004
6	1.0000 ± 0.0059	0.9959 ± 0.0004	0.9854 ± 0.0004	0.9913 ± 0.0004
Unreflected Cylinders (HEU-SOL-THERM-020)				
1	0.9966 ± 0.0116	1.0039 ± 0.0005	0.9932 ± 0.0005	1.0006 ± 0.0005
2	0.9956 ± 0.0093	1.0105 ± 0.0005	0.9983 ± 0.0005	1.0066 ± 0.0005
3	0.9957 ± 0.0079	1.0185 ± 0.0005	1.0070 ± 0.0005	1.0149 ± 0.0005
4	0.9955 ± 0.0078	1.0162 ± 0.0005	1.0048 ± 0.0005	1.0160 ± 0.0005
5	0.9959 ± 0.0077	1.0234 ± 0.0005	1.0131 ± 0.0005	1.0167 ± 0.0005

$$\sigma < |\Delta k| \leq 2\sigma \quad |\Delta k| > 2\sigma$$

PNL MOX Lattices

Case	Pitch (cm)	Soluble Boron (PPM)	Benchmark k_{eff}	Calculated k_{eff}		
				ENDF/B-VII.0	ENDF/B-VI	JENDL-3.3
PNL-30	1.77800	1.7	1.0010 ± 0.0059	1.0019 ± 0.0003	0.9933 ± 0.0003	0.9987 ± 0.0003
PNL-31	1.77800	687.9	1.0009 ± 0.0045	1.0028 ± 0.0003	0.9960 ± 0.0004	1.0008 ± 0.0004
PNL-32	2.20914	0.9	1.0024 ± 0.0029	1.0035 ± 0.0003	0.9965 ± 0.0004	1.0018 ± 0.0003
PNL-33	2.20914	1090.4	1.0024 ± 0.0021	1.0072 ± 0.0003	1.0029 ± 0.0003	1.0069 ± 0.0003
PNL-34	2.51447	1.6	1.0038 ± 0.0022	1.0041 ± 0.0003	0.9989 ± 0.0003	1.0033 ± 0.0003
PNL-35	2.51447	767.2	1.0029 ± 0.0024	1.0069 ± 0.0003	1.0031 ± 0.0003	1.0062 ± 0.0004

$$\sigma < |\Delta k| \leq 2\sigma$$


$$|\Delta k| > 2\sigma$$

Overall, ENDF/B-VII.0 produces best results

None of the ENDF/B-VII.0 values for k_{eff} are lower than the corresponding mean benchmark value, but two are higher by more than 1 standard deviation


All 3 libraries produce different shapes for slightly and highly borated cases

Slightly Borated PNL MOX Lattices



ENDF/B-VII.0 produces significant improvement relative to ENDF/B-VI

Highly Borated PNL MOX Lattices



⇒ Thermal cross sections for ^{239}Pu probably should be reviewed

Unreflected Spheres of Plutonium-Nitrate Solutions

Case	H / ^{239}Pu Atom Radio	Benchmark k_{eff}	Calculated k_{eff}		
			ENDF/B-VII.0	ENDF/B-VI	JENDL-3.3
PNL-6	131.83	1.0000 ± 0.0065	1.0040 ± 0.0005	1.0029 ± 0.0005	1.0070 ± 0.0004
PNL-5	574.52	1.0000 ± 0.0052	1.0066 ± 0.0004	1.0058 ± 0.0004	1.0098 ± 0.0004
PNL-1	701.70	1.0000 ± 0.0032	1.0066 ± 0.0004	1.0066 ± 0.0004	1.0101 ± 0.0004
PNL-8	797.62	1.0000 ± 0.0032	1.0052 ± 0.0004	1.0060 ± 0.0004	1.0101 ± 0.0004
PNL-4	907.13	1.0000 ± 0.0052	1.0002 ± 0.0004	1.0012 ± 0.0004	1.0043 ± 0.0004
PNL-3	1207.8	1.0000 ± 0.0052	0.9950 ± 0.0004	0.9954 ± 0.0004	0.9985 ± 0.0004
PST-9	2806.8	1.0003 ± 0.0033	1.0189 ± 0.0002	1.0190 ± 0.0002	1.0226 ± 0.0002

$$\sigma < |\Delta k| \leq 2\sigma \quad |\Delta k| > 2\sigma$$

ENDF/B-VII.0 results are very similar to those from ENDF/B-VI

All differences greater than 1σ are high, except for PNL-3

Is PST-9 (case 3a) a valid data point or an outlier?




The World's Greatest Science Protecting America

Applied Physics Division



Unreflected Spheres of Plutonium-Nitrate Solutions



⇒ Thermal cross sections for ^{239}Pu should be reviewed

REFLECTED HEU NITRATE SOLUTION WITH Cd

Source	k_{eff}	Δk
Benchmark	1.0012 ± 0.0019	—
ENDF/B-VI	0.9870 ± 0.0004	-0.0142 ± 0.0016
ENDF/B-VII.0	0.9918 ± 0.0004	-0.0094 ± 0.0016
ENDF/B-VII.0 + NNDC ^{113}Cd	1.0008 ± 0.0004	-0.0004 ± 0.0016

$$|\Delta k| > 5 \sigma$$

Benchmark is a reflected HEU nitrate solution in water that has a thermal spectrum even though cadmium is present in the solution

Reactivity deficiency was traced to problems with the ^{113}Cd cross sections, and a new evaluation for ^{113}Cd was generated by the National Nuclear Data Center at Brookhaven National Laboratory



The World's Greatest Science Protecting America

Applied Physics X Division



CONCLUSIONS

Improvements still are needed, particularly for:

- ^{235}U cross sections in the unresolved resonance region
- Fast copper cross section
- Neptunium cross sections
- Angular scattering distribution for ^2H (probably)
- Thermal plutonium (^{239}Pu) cross sections

One problem already has been resolved:

- Thermal ^{113}Cd cross sections

New cross sections for ^{113}Cd will be included in the next interim distribution, ENDF/B-VII.1

MCNPX 2.6.X Features (2006-2007)

by

Gregg McKinney, Joe Durkee, John Hendricks, Mike James,
Russ Johns, Denise Pelowitz, and Laurie Waters

M&C/SNA 2007, Monterey, CA, April 15-19, 2007

MCNPX 2.6.X Features (2006-2007)

- MCNPX Overview
- History of MCNPX
- User Base
- Features for 2006-2007
- Future of MCNPX



EST. 1943

Operated by the Los Alamos National Security, LLC for the DOE/NNSA


UNCLASSIFIED




MCNPX Overview

- Monte Carlo radiation transport code
 - Extends MCNP4C to virtually all particles and energies
 - 34 particles (n,p,e, 5 leptons, 11 baryons, 11 mesons, 4 LI)
 - Continuous energy (roughly 0-100 GeV)
 - Data libraries below ~ 150 MeV (n,p,e,h) & models otherwise
- General 3-D geometry
 - 1st & 2nd degree surfaces, tori, 10 macrobodies, lattices
- General sources and tallies
 - Interdependent source variables, 7 tally types, many modifiers
- Supported on virtually all computer platforms
 - Unix, Linux, Windows, OS X (parallel with MPI)

History of MCNPX



History of MCNPX



History of MCNPX

1992-1993	LAHET and Superhet Superconducting Super Collider	
1994-1995	Start of the APT program	
Version 1.0		April 22, 1997
Version 2.0		October 1, 1997
Version 2.1.3	HISTP writing, compatible with HTAPE, collisional energy loss model	April 17, 1998
Version 2.1.4	Mesh & radiography tallies, gridconv, bertin & phtlib binary support	July 24, 1998
Version 2.1.5	CEM, HTAPE3X, User's Manual, Beta test team	Nov 14, 1999
Version 2.1.6	Proton libraries (internal user only)	September 14, 1999
Version 2.3.0		April 27, 2002

History of MCNPX

Version 2.4.0

August 01, 2002

Update to MCNP4C3, F90, Windows PC, New user's manual

Version 2.5.C

April, 2003

MPI Multiprocessing, Mix & Match, CEM2K

Version 2.5.D

August, 2003

INCL4/ABLA physics models, Multiple particles on SDEF card, READ card

Version 2.5.E

February, 2004

MPI KCODE speedup, 64-bit integers, G5 support, 2-D color contour plots

Version 2.5.0

March, 2005

Mesh tally contour plots, Pulse-height tally with VR, PN improvements

Version 2.6.A

December, 2005

Transmutation, Long file names, STOP card

Version 2.6.B

June, 2006

CEM 03, new PHTLIB, predictor-corrector for transmutation

Version 2.6.C

December, 2006

Spherical weight windows, delayed particle production

Version 2.6.D

~May, 2007



EST. 1943
Operated by the Los Alamos National Security, LLC for the DOE/NNSA

UNCLASSIFIED



User Base

- ~2500 users world wide
 - Provide 6-8 workshops per year (4-6 US, ~2 international)
 - 150 workshop participants per year
 - Access to RSICC/NEA released versions only
 - <http://www-rsicc.ornl.gov/> (C00730) 2.5.0
 - <http://www.nea.fr/html/dbprog/> (CCC-0715) 2.4.0
 - Limited access to MCNPX web site
 - <http://mcnpx.lanl.gov> (some documentation)
- ~2000 registered Beta Testers
 - Full access to MCNPX web site
 - Access to intermediate versions
 - Increased user support

Application	# Groups	Percent
Medical (BNCT, proton therapy, etc.)	50	15
Spacecraft, Cosmic Rays, SEE, propulsion	42	12
Detectors, experiments, Threat Reduction	39	11
ATW, ADS, Energy Amplifiers	37	11
Fuel cycles, beginning to end, including storage	32	9
Accelerator Shielding and Health Physics	28	8
Theoretical Physics	23	7
Neutron Production for Scattering	21	6
Isotope Production	14	4
Radiography	12	4
MCNPX/MCNP code development	11	3
Homeland Security	10	3
Materials studies (IFMIF)	6	2
Radioactive Ion Beams	5	1
Irradiation Facilities	4	1
Neutrino Targets	4	1
Light Sources, electron machines	3	1

Features for 2006 – Version 2.6.A

- Transmutation using Cinder90 (BURN card)
 - Several keywords of options (MAT, POWER, etc.)
 - Automatic updating of material atom densities
- Long file names (40 vs. 8 characters)
- STOP card - terminate tallies at desired precision
- Corrections/enhancements/extensions
 - Proton step size control (HSTEP on M card)
 - New $S(\alpha,\beta)$ scattering law
 - Differential data tallies extended to table physics
 - Separate printout of induced fission multiplicity

BURN Card


Burn 7 fuel pins surrounded by H₂O in a hex lattice

```
1 1 -8.3 -1      u=1 imp:n=1    vol=192.287
2 8 -6.5 1 -2      u=1 imp:n=1
3 9 -0.7 2       u=1 imp:n=1
100 9 -1.8 -3    u=8 lat=2 imp:n=1 fill=-2:2 -2:2 0:0
                  8 8 8 8 8 8 8 1 1 8
                  8 1 1 1 8
                  8 1 1 8 8 8 8 8 8
101 0           -4      imp:n=1 fill=8
102 9 -1.8 4 -5      imp:n=1
103 0           5      imp:n=0

1 rcc 0 0 0 0 365 0.4095
2 rcc 0 0 -1 0 0 367 0.4750
3 rhp 0 0 -1 0 0 367 0.6565 0 0
4 rhp 0 0 -1 0 0 367 0 1.895 0
*5 rhp 0 0 -1.1 0 0 367.2 0 1.896 0
```

BURN TIME=50,500 MAT=1 POWER=0.066956 PFRAC=1.0,1.0
OMIT=1,6,6014,7016,8018,9018,90234,91232 BOPT=1.0,4

```
m1   8016.60c 2.0 92235.60c 0.01 92238.60c 0.99
m8   40000.60c 1.0
m9   1001.60c 2.0 8016.60c 1.0
mt9  lwtr.01t
kcode 100 1.0 10 50
ksrc  0 0 150 1.3 0 150 -1.3 0 150 0.66 1.14 150
      0.66 -1.14 150 -0.66 1.14 150 -0.66 -1.14 150
```



UNCLASSIFIED

1burnup summary table by material
210

print table

nuclides with atom fractions below 1.000E-10 for a material are zeroed and deleted from print tables after t=0

neutronics and burnup data

step	duration (days)	time (days)	power (MW)	keff	flux	ave. nu	ave. q	burnup (GWd/MTU)
0	0.000E+00	0.000E+00	6.696E-02	0.99763	3.641E+14	2.449	200.981	0.000E+00
1	5.000E+01	5.000E+01	6.696E-02	1.00012	3.701E+14	2.554	203.154	2.383E+00
2	5.000E+02	5.500E+02	6.696E-02	0.85037	4.638E+14	2.869	209.385	2.621E+01

actinide inventory for sum of materials at end of step 2, time 5.500E+02 (days), power 6.696E-02 (MW)

no.	zaid	mass (gm)	activity (Ci)	sp. act. (Ci/gm)	atom den. (a/b-cm)	atom fr.	mass fr.
1	92234	3.465E-04	2.154E-06	6.217E-03	4.636E-09	2.577E-07	2.533E-07
2	92235	3.935E-01	8.506E-07	2.161E-06	5.244E-06	2.914E-04	2.877E-04
3	92236	1.789E+00	1.157E-04	6.467E-05	2.374E-05	1.319E-03	1.308E-03
4	92237	7.849E-03	6.405E+02	8.160E+04	1.037E-07	5.763E-06	5.739E-06
5	92238	1.355E+03	4.553E-04	3.361E-07	1.782E-02	9.905E-01	9.904E-01
6	92239	1.539E-03	5.158E+04	3.351E+07	2.016E-08	1.121E-06	1.125E-06
14	94242	1.117E+00	4.418E-03	3.954E-03	1.446E-05	8.034E-04	8.169E-04
	totals	1.368E+03	1.041E+05	7.610E+01	1.799E-02	1.000E+00	1.000E+00

nonactinide inventory for sum of materials at end of step 2, time 5.500E+02 (days), power 6.696E-02 (MW)

no.	zaid	mass (gm)	activity (Ci)	sp. act. (Ci/gm)	atom den. (a/b-cm)	atom fr.	mass fr.
1	6012	2.336E-06	0.000E+00	0.000E+00	6.096E-10	1.638E-08	1.186E-08
2	6013	1.057E-02	0.000E+00	0.000E+00	2.545E-06	6.839E-05	5.366E-05
3	8016	1.891E+02	0.000E+00	0.000E+00	3.702E-02	9.946E-01	9.599E-01
4	8017	1.405E-02	0.000E+00	0.000E+00	2.588E-06	6.954E-05	7.132E-05
16	60145	3.469E-01	1.426E-14	4.112E-14	7.497E-06	2.014E-04	1.761E-03
	totals	1.970E+02	8.830E+01	4.483E-01	3.722E-02	1.000E+00	1.000E+00



UNCLASSIFIED

Operated by the Los Alamos National Security, LLC for the DOE/NNSA



Long File Names

```
E:\MCNPX\scratch>.\mcnpx inp=test_long_names.txt na=test_long_names.  
mcnpx ver=26bc1 ld=Sat Jul 01 08:00:00 MST 2006 11/12/06 20:08:04  
...  
dynamic storage =      0 words,      0 bytes.    cp0 = 0.00  
run terminated when      10 particle histories were done.  
dump 2 on file test_long_names.r  nps =      10  coll =      0  
          ctm =    0.00   nrn =      40  
mcrun is done
```

```
E:\MCNPX\scratch>dir
```

Directory of E:\MCNPX\scratch

11/12/2006 08:08 PM	<DIR>	.
11/12/2006 08:08 PM	<DIR>	..
09/11/2006 03:23 PM	6,574,080	mcnpx.exe
11/12/2006 08:08 PM	22,513	test_long_names.d
11/12/2006 08:08 PM	25,510	test_long_names.o
11/12/2006 08:08 PM	401,342	test_long_names.r
08/04/2006 01:21 PM	481	test_long_names.txt
	5 File(s)	7,023,926 bytes
	2 Dir(s)	31,759,495,168 bytes free



UNCLASSIFIED

Operated by the Los Alamos National Security, LLC for the DOE/NNSA



STOP Card

14 MeV neutrons in water - test STOP card

c cell cards

1 1 -1. -1 IMP:N=1
2 0 1 IMP:N=0

1 so 30.0

m1 1001.60c 2. 8016.60c 1.

mt1 hh2o.20t

xs1 hh2o.20t 0.998623 ct00 0 1 1 1237501 0 0 2.530E-08

sdef erg=14.1

e0 1.00000E-11 6251og 1.44544E+01

vol 1.

f44:n 1

f141:n 1

STOP F44 .01 NPS 10000 CTME 10.0

OUTPUT FILE

1 tally fluctuation charts

tally 44						tally 141					
nps	mean	error	vov	slope	fom	mean	error	vov	slope	fom	
1000	5.6661E+01	0.0144	0.0031	10.0	489705	3.7459E-01	0.0354	0.0003	0.0	81225	
2000	5.6824E+01	0.0103	0.0016	10.0	468574	3.7000E-01	0.0254	0.0002	10.0	77294	
3000	5.7028E+01	0.0083	0.0010	10.0	477186	3.6737E-01	0.0209	0.0001	10.0	75590	

dump no. 2 on file stop.r nps = 3000 coll = 599417 ctm = 0.03



UNCLASSIFIED


New S(α,β) Treatment

```
14 MeV neutrons in water - test S(alpha,beta)
c    cell cards
1    1   -1.   -1 IMP:N=1
2    0        1 IMP:N=0

1    so 30.0

m1    1001.60c 2. 8016.60c 1.
mt1   hh2o.20t $ Replace with lwtr.01t for old treatment
xs1   hh2o.20t 0.998623 ct00 0 1 1 1237501 0 0 2.530E-08
nps   1000000
sdef  erg=14.1
e0    1.00000E-11 625log 1.44544E+01
vol   1.
f44:n 1
f141:n 1
```

New S(α, β) Treatment



Features for 2006 – Version 2.6.B


- Transmutation improvements (BURN card)
 - Predictor/corrector
 - Automatic selection of FP dist. (thermal, fast, high)
- CEM INC model upgrade (from 2K to 03)
- FIELD card—planetary gravity effects for neutrons
- Corrections/enhancements/extensions
 - New photon emission data: PHTLIB
 - Geometry plot basis vectors
 - Extend ZAID identifiers

CEM Upgrade

```
1 GeV protons into Fe-56
1      1 -7.86 -1 IMP:N=1
2      0           1 IMP:N=0

1      so 1.0

mode  n h
m1   26056.24c 1.
nps  200000
sdef erg=1000 par=h
PHYS:N 1001.0
LCA  7j -2 1 $ Use CEM
e0   1 1991og 1000
c0   0 1
f1:n 1
f11:h 1
```



CEM Upgrade

Light-product yields (A<30)

Model	Proton energy (MeV)					
	300	500	750	1000	1500	2600
BERTINI	1035	26.1	50.5	13.8	4.93	3.35
ISABEL	---	256	49.1	17.0	5.99	4.02
INCL	233	215	51.5	38.1	26.1	12.1
CEM2K	---	12.6	21.1	7.83	4.87	4.02
CEM03	13.0	2.23	1.32	1.49	1.58	1.72

Heavy-product yields (A>30)

Model	Proton energy (MeV)						Ave. Dev.
	300	500	750	1000	1500	2600	
BERTINI	2.24	2.29	2.75	2.86	3.16	3.20	4.37
ISABEL	3.75	2.85	3.02	2.63	2.85	3.01	4.24
INCL	4.72	3.24	3.14	3.13	3.35	3.54	7.14
CEM2K	2.74	2.54	2.62	2.76	2.92	3.20	3.55
CEM03	1.84	1.89	1.89	1.92	2.04	3.17	2.26

Mean-squared deviation factors between model predictions and experimental data measured at ITEP.



UNCLASSIFIED

Operated by the Los Alamos National Security, LLC for the DOE/NNSA



FIELD Card

5 GeV protons into Mars, gravity reflection

```
1    1 -1.0      -1      imp:n=1
100   2 -1.35e-5  -101 +1      imp:n=1
101   2 -1.28e-5  -102 +101 imp:n=1
102   2 -1.22e-5  -103 +102 imp:n=1
103   2 -1.14e-5  -104 +103 imp:n=1
104   2 -1.08e-5  -105 +104 imp:n=1
105   2 -1.01e-5  -106 +105 imp:n=1
999    0          +106     imp:n=0
```

```
1    so 339000000.0
101   so 339060000.0
102   so 339110000.0
103   so 339180000.0
104   so 339240000.0
105   so 339310000.0
106   so 339380000.0
```

```
m1  8016.60c -0.6 14000.60c -0.3 26056.60c -0.1
m2  6000.60c -0.27 7014.60c -0.02 8016.60c -0.70
      18000.35c -0.01
```

FIELD GCUT=0.1320 GPAR=1 GRAD=3393.0 GSUR=106


```
mode h n p z / d t s a
lca 8j 1 $ Use CEM
sdef par=9 erg=5000 sur=106 nrm=-1
nps 10000
```

```
phys:n 5010 j j j 20
e11  0. 1024i 10. 5000.
full 0. 8016.00051 8016.00052 8016.00053
      8016.00102 8016.
      14028.14027 14028.14026 14028.13027
      14028.13026 14000.
      26056.00051 26056.00052 26056.00053
      26056.00102 26056.

f11:p 1
ft11 tag 1
e21  1e-10 99log 1e-7
f21:n 105
```

Undocumented Feature

FIELD Card



Features for 2006 – Version 2.6.C

- Transmutation improvements (BURN card)
 - Support for continue-runs & parallel execution
 - Printing of reaction rates sent to Cinder90
 - Reduced memory requirements
- Spherical weight windows
- Delayed neutrons & gammas
 - ~1000 nuclides treated with gamma line data
- Photon tally tagging
- Model treatment for library absorption reactions


Spherical Weight Windows

10 MeV photons into 1m H₂O surrounding HEU

```
1 1 -19.0 -1 imp:p=1
2 2 -1.0 +1 -2 imp:p=1
3 0 +2 -3 imp:p=1
4 0 -3 imp:p=0
```

```
1 sph 0 0 0 3
2 sph 0 0 0 100
3 sph 0 0 0 200
```

```
mode p
sdef erg=10 pos=-105 0 0 rad=d1 axs=1 0 0 ext=0
    vec=1 0 0 dir=d2
s1 0 10
sp1 -21 1
si2 0 1
sp2 0 1
m1 92235 .5 92238 .5
m2 1001 2 8016 1
nps 100000
f4:p 1
wwg 4 0
mesh geom rpt origin=0 0 0 ref=-99 1 1 axs 1 0 0 vec 0 1 0
    imesh 101. iints 20
    jmesh .25 .5 jint 4 8
    kmesh 1 kints 1
c wwp:p 4j -1 $ Add this card to use WW
```



Delayed Neutrons and Gammas

DN/DG from high-energy fission of U-235

```
1 1 -18.9 -1      imp:n=1
2 0          +1      imp:n=0
```

```
1 sph  0 0 0 0.1
```

```
mode n p
m1    92235.60c -1
phys:n 20 2j 101 20
phys:p 5j -102
cut:n j j 0 0
sdef erg=10.0 par=n
f01:p 1
e01   .100 499i 12.
t01   0.0001e8 100.e8
f11:p 1
t11   0.0000001e8 99log 100.e8
f21:n 1
e21   .001 199log 12.
t21   0.0001e8 100.e8
f31:n 1
t31   0.0000001e8 99log 100.e8
nps   2000000
```



Operated by the Los Alamos National Security, LLC for the DOE/NNSA

UNCLASSIFIED



Delayed Neutrons and Gammas


neutron creation	tracks	weight (per source particle)	energy (per source particle)	neutron loss	tracks	weight (per source particle)	energy (per source particle)
source	2000000	1.0000E+00	1.0000E+01	escape	2078492	1.0317E+00	9.9508E+00
nucl. interaction	0	0.	0.	energy cutoff	0	0.	0.
particle decay	0	0.	0.	time cutoff	0	0.	0.
weight window	0	0.	0.	weight window	0	0.	0.
cell importance	0	0.	0.	cell importance	0	0.	0.
weight cutoff	0	0.	0.	weight cutoff	0	0.	0.
energy importance	0	0.	0.	energy importance	0	0.	0.
dxtran	0	0.	0.	dxtran	0	0.	0.
forced collisions	0	0.	0.	forced collisions	0	0.	0.
exp. transform	0	0.	0.	exp. transform	0	0.	0.
upscattering	0	0.	0.	downscattering	0	0.	8.1324E-03
photonuclear	0	0.	0.	capture	98	3.8100E-05	1.6678E-04
(n,xn)	22458	1.1229E-02	1.2686E-02	loss to (n,xn)	7486	3.7430E-03	3.7421E-02
prompt fission	65553	3.2679E-02	6.7772E-02	loss to fission	17266	8.5948E-03	8.4020E-02
delayed fission	15331	1.2414E-04	5.7533E-05	nucl. interaction	0	0.	0.
tabular boundary	0	0.	0.	particle decay	0	0.	0.
tabular sampling	0	0.	0.	tabular boundary	0	0.	0.
total	2103342	1.0440E+00	1.0081E+01	total	2103342	1.0440E+00	1.0081E+01
photon creation	tracks	weight (per source particle)	energy (per source particle)	photon loss	tracks	weight (per source particle)	energy (per source particle)
source	0	0.	0.	escape	127772	1.2601E-01	1.3541E-01
nucl. interaction	0	0.	0.	energy cutoff	1	2.0234E-06	5.2097E-06
particle decay	113751	5.6552E-02	5.4380E-02	time cutoff	0	0.	0.
weight window	0	0.	0.	weight window	0	0.	0.

Total DG = 6.6 γ/f


Total DN = .014 n/f

UNCLASSIFIED

Delayed Neutrons and Gammas




Delayed Neutrons and Gammas



UNCLASSIFIED

Delayed Neutrons and Gammas



UNCLASSIFIED

Photon Tally Tagging

5 GeV protons into Mars with photon tagging

```
1    1 -1.0      -1      imp:n=1
100   2 -1.35e-5 -101 +1      imp:n=1
101   2 -1.28e-5 -102 +101  imp:n=1
102   2 -1.22e-5 -103 +102  imp:n=1
103   2 -1.14e-5 -104 +103  imp:n=1
104   2 -1.08e-5 -105 +104  imp:n=1
105   2 -1.01e-5 -106 +105  imp:n=1
999   0           +106     imp:n=0
```

```
1    so 339000000.0
101   so 339060000.0
102   so 339110000.0
103   so 339180000.0
104   so 339240000.0
105   so 339310000.0
106   so 339380000.0
```


```
m1    8016.60c -0.6 14000.60c -0.3 26056.60c -0.1
m2    6000.60c -0.27 7014.60c -0.02 8016.60c -0.70
                  18000.35c -0.01
FIELD GCUT=0.1320 GPAR=1 GRAD=3393.0 GSUR=106
mode  h n p z / d t s a
lca   8j 1 $ Use CEM
sdef  par=9 erg=5000 sur=106 nrm=-1
nps   10000
```

```
phys:n 5010 j j j 20
e11   0. 1024i 10. 5000.
full  0. 8016.00051 8016.00052 8016.00053
                  8016.00102 8016.
                  14028.14027 14028.14026 14028.13027
                  14028.13026 14000.
                  26056.00051 26056.00052 26056.00053
                  26056.00102 26056.

f11:p 1
ft11  tag 1
e21   1e-10 99log 1e-7
f21:n 105
```

Undocumented Feature

Photon Tally Tagging



Models for Library Interactions


2 MeV neutrons into He-3

```
1 1 -5.3540E-4 -1 imp:n=1
2 0 1 -2 imp:n=1
3 0 2 imp:n=0

1 so 4.0
2 so 100.0

mode n h d t s
sdef par=n erg=2 pos=-5 0 0 rad=d1
axs=1 0 0 ext=0 vec=1 0 0 dir=1
sil 0 3
sp1 -21 1
cut:n 2j 0 0
cut:h,d,t,s j .001
phys:n 6j 2
m1 2003.60c 1
nps 10000000
f6:h 1
f16:d 1
f26:t 1
f36:s 1
```

```
f8:n 1
e8 0. 99i 2.1
ft8 PHL 1 6 1 0
f18:n 1
e18 0. 99i 2.1
ft18 PHL 1 16 1 0
f28:n 1
e28 0. 99i 2.1
ft28 PHL 1 26 1 0
f38:n 1
e38 0. 99i 2.1
ft38 PHL 1 36 1 0
f58:n 1
e58 0. 99i 2.1
ft58 PHL 4 6 1 16 1 26 1 36 1 0
```



LA150N Library

Isotope	ZAID	Proton	Deuteron	Triton	Alpha
H-1	1001.24c	1.0E-11			
H-2	1002.24c	3.339		1.0E-11	
Be-9	4009.24c	14.266	16.301	11.709	0.667
C	6000.24c	20.0	20.0		20.0
N-14	7014.24c	20.0	20.00		20.0
O-16	8016.24c	20.0	20.0		20.0
Al-27	13027.24c	1.897	6.274	11.29	3.25
Si-28	14028.24c	4.0	20.0	20.0	2.746
Si-29	14029.24c	3.0	20.0	20.0	1.3
Si-30	14030.24c	8.012	20.0	20.0	4.345
P-31	15031.24c	20.0	20.0		20.0
Ca	20000.24c	20.0	20.0	20.0	20.0
Cr-50	24050.24c	1.0	20.0	20.0	2.25
Cr-52	24052.24c	3.256	20.0	20.0	1.233
Cr-53	24053.24c	2.69	20.0	20.0	1.0
Cr-54	24054.24c	6.33	20.0	20.0	1.581
Fe-54	26054.24c	0.7	20.0	20.0	3.0
Fe-56	26056.24c	2.966	20.0	20.0	0.862
Fe-57	26057.24c	1.943	20.0	20.0	0.8
Ni-58	28058.24c	0.5	20.0	20.0	0.5




UNCLASSIFIED

Operated by the Los Alamos National Security, LLC for the DOE/NNSA



Models for Library Interactions



Features for 2007 – Version 2.6.D

- Transmutation improvements (BURN card)
 - Time-dependent material changes (CONC keyword)
 - Repeated-structures power norm. (VOL keyword)
 - Fission-product tier improvements
- Coupled space-energy-time weight windows
- Activation neutrons and gammas
- Background radioactive sources
- Muon capture physics
- Heavy-ion transport (via LAQGSM)

Coupled Space-Energy-Time WW


7 MeV neutrons into H₂O surrounding 2kg HEU

```
1 1 -19.0 -1 imp:n=1
2 2 -1.0 +1 -2 imp:n=1
3 0          +2 imp:n=0
```

```
1 sph 0 0 0 3
2 sph 0 0 0 100
```

```
mode n p
phys:p 5j -101 $ turn on delayed gammas
sdef par=n erg=7 pos=-99 1 1 vec=1 0 0 dir=1
m1 92235 .5 92238 .5
m2 1001 2 8016 1
nps 100000
f1:p 2
t1 0.1e8 1e15 nt
f4:n 1
wwg 4 0
mesh geom=rpt origin=0 0 0 ref=-99 1 1
  axs=1 0 0 vec=0 1 0
  imesh 3.01 101. iints 3 5
  jmesh .5 jints 10
  kmesh 1 kints 1
c wwp:n 4j -1 $ Add this to use neutron WWS
```

7 MeV
Neutrons



Coupled Space-Energy-Time WW


7 MeV neutrons into H₂O surrounding 2kg HEU

```
1 1 -19.0 -1 imp:n,p=1
2 2 -1.0 +1 -2 imp:n,p=1
3 0 +2 imp:n,p=0
```


```
1 sph 0 0 0 3
2 sph 0 0 0 100
```

```
mode n p
totnu no
phys:p j 1 3j -101 $ turn on delayed gammas
sdef par=n erg=7 poss=-99 1 1 vec=1 0 0 dir=1
m1 92235 .5 92238 .5
m2 1001 2 8016 1
nps 300000
f1:p 2
t1 0.1e8 1e15 nt
e1 3 100 nt
wwg 1 0 5j 1
wwge:p 0.1e8 1e15
mesh geom=rpt origin=0 0 0 ref=-99 1 1 axs=1 0 0 vec=0 1 0
      imesh 3.01 101. iints 3 5
      jmesh .5 jints 10
      kmesh 1 kints 1
wwp:n 4j -1
c wwp:p 4j -1 $ Add this to use photon WWs
```

Neutron
WWs



Photon
WWs



Activation Neutrons & Gammas


14 MeV neutrons into H₂O surrounding 2kg HEU

```
1 1 -19.0 -1 imp:n,p=0  
2 2 -1.0 +1 -2 imp:n,p=1  
3 0 +2 imp:n,p=0
```


```
1 sph 0 0 0 3  
2 sph 0 0 0 100
```

```
mode n p  
phys:p 5j -102 $ turn on delayed gammas  
cut:n 2j 0 0 $ turn off implicit capture  
sdef par=n erg=14 pos=-99 1 1 vec=1 0 0 dir=1  
m1 92235 .5 92238 .5  
m2 1001 2 8016 1  
nps 1000000  
f1:p 2  
t1 0.1e8 1e15 nt  
e1 1 99i 10
```

14 MeV
Neutrons →



Activation Neutrons & Gammas




Background Radioactive Sources

Co-57, Co-60, and Cs-137 within soil


```
1 2 -1.6 -1      imp:p=1  
2 0           1      imp:p=0
```

```
1 so 100.0
```

```
sdef par=sp cel=1 pos=0 0 0 wgt=1 rad=d1  
sil 0 100  
sp1 -21 2  
mode p  
nps 100000  
m2   1001.66c -.002  8016.66c -.527  
     11023.66c -.021 13027.66c -.061  
     14028     -.345 19000      -.029  
     26056     -.016  
     27057 -0.00000001 27060 -.000001  
     55137 -.000323  
plib=.02p  
e0 0.0 2999i 3.0  
f4:p 1  
f2:p 1  
c2 0 1
```



Background Radioactive Sources



Muon Capture Physics


350 MeV muons into Pb surrounding HEU

```
1 1 -18.95 -1      imp:l,p=1
2 8 -11.35 1 -2    imp:l,p=1
3 3 -1.0 2 -3     imp:l,p=1
4 4 -7.8 3 -4     imp:l,p=1
5 5 -1.205e-3 4 -100 imp:l,p=1
100 0 100          imp:l,p=0
```


```
1 rcc -10.0 0.0 0.0 20.0 0.0 0.0 5.0
2 rcc -12.5 0.0 0.0 25.0 0.0 0.0 7.5
3 rpp -47.5 47.5 -47.5 47.5 -47.5 47.5
4 rpp -50.0 50.0 -50.0 50.0 -50.0 50.0
100 so 100.0
```

```
mode l p
phys:l 350.0
phys:p 350.0
sdef par=l erg=350.0 x=d1 y=d2 z=-60.0
  vec=0 0 1 dir=1
sil -12.5 12.5
sp1 0 1
si2 -7.5 7.5
sp2 0 1
m1 92238 -.20 92235 -.80
m3 1001 2       6012 1
m4 26054 5.9 26056 91.72 26057 2.1 26058 .28
```

m5	1000	-6e-4	8000	-0.2353	7000	-0.7513
	18000		18000	-0.0128		
m8	82204	1.4	82206	24.1	82207	22.1
	82208		82208	52.4		
e2	0.0	9999i	10.0			
f2:p	100					



Muon Capture Physics



Operated by the Los Alamos National Security, LLC for the DOE/NNSA



Heavy-ion Transport



```
969 MeV/n Fe into H2O
1      2   -1.0     -1      imp:n=1
2      1   -0.0012    1   -2 imp:n=1
3      0           2 imp:n=0

1      rpp -15 15 -15 15 0 40.0
2      rcc 0 0 -20 0 0 80 50
3      pz -1

m1    7014.24h 0.781 8016.24h 0.219
m2    1001.24h 0.667 8016.24h 0.333
mode  h a n #
phys:h 60000 j j j j j 0.1
phys:n 60000 j 0 -1 j j 1
lca 8j 3
sdef par=26056 erg=d2 sur=3 dir=1 vec=0 0 1 pos=0 0 -1 rad=d1
s1    0 0.5
sp1   -21
si2   L 54224.00 54251.00 54278.00
sp2   0.25 0.5 0.25
nps   20000
tmesh
rmesh3 total
cora3 -7.0 7.0
corb3 -7.0 7.0
corc3 0.0 400i 40.0
endmd
```

Undocumented Feature?

Heavy-ion Transport



Features for 2007 – Version 2.6.E

- Transmutation improvements
 - Accurate evolution of meta-stables
 - Treatment for minor actinides
- Plotting of spherical mesh tallies
- Enhanced tally tagging (other particles & tags)
- Time-dependent transformations
 - Allows moving universes (e.g., control-rod movement)
- Fission gamma multiplicity
- Nuclear resonance fluorescence physics

Future of MCNPX

- Possible public release of 2.6.0 (~Sept. 2007)
- MCNPX and MCNP merger
 - Hope to preserve all features of both codes
 - Preliminary version by Summer 2007
 - Public release perhaps in 2008
- Capabilities beyond 2.6.0 will be put in MCNP6/7
 - MCNPX will cease to be distributed
 - MCNP web site may allow for a small Beta group
 - MCNP & MCNPX workshops will be combined