Table 16. Tuberculosis Case Rates per 100,000 Population by Hispanic Ethnicity and Non-Hispanic Race, Sex, and Age Group: United States, 2004

Race/Ethnicity and Sex	Age Group						
	All Ages	Under 5	5–14	15–24	25–44	45 –64	≥65
Total Rate	4.9	2.8	1.0	3.8	5.9	5.9	7.8
Hispanic or Latino ¹	10.1	6.7	2.2	9.6	11.6	14.4	25.4
Male	12.5	5.9	2.2	11.5	14.4	20.2	36.2
Female	7.6	7.5	2.2	7.4	8.3	8.7	17.4
Non-Hispanic							
American Indian or Alaska Native	7.3	5.2	1.1	2.0	6.6	12.1	25.3
Male	10.5	4.6	1.0	3.5	10.5	19.5	34.9
Female	4.1	5.9	1.1	0.5	2.8	5.3	18.0
Asian	27.6	6.2	3.7	20.6	28.8	34.5	66.4
Male	31.0	5.1	3.7	19.3	30.2	43.4	92.0
Female	24.4	7.4	3.7	21.9	27.4	26.7	47.3
Black or African American	11.3	5.1	2.3	7.4	13.8	17.7	18.6
Male	14.6	5.6	2.1	7.2	17.8	27.6	26.9
Female	8.3	4.7	2.5	7.6	10.2	9.3	13.4
Native Hawaiian or Other Pac. Islander	16.3	18.4	7.7	21.2	12.8	24.4	15.8
Male	14.9	14.3	3.0	13.8	16.3	25.8	8.8
Female	17.8	22.8	12.7	29.2	9.2	23.0	21.5
White	1.3	0.4	0.1	0.4	1.0	1.8	3.3
Male	1.8	0.5	0.1	0.4	1.3	2.6	4.7
Female	0.9	0.3	0.1	0.4	0.7	0.9	2.2
Multiple Race ²	0.9	0.4	0.1	1.0	1.4	1.4	2.0
Male	0.9	0.4	0.2	0.0	1.2	2.6	4.7
Female	8.0	0.4	0.0	2.0	1.6	0.3	0.0

¹Persons of Hispanic or Latino origin may be of any race or multiple race.

Note: Denominators for computing case rates were obtained from the U.S.Census Bureau Monthly Postcensal Resident Population, by single year of age, sex, race, and Hispanic origin: July 1, 2004 (http://www.census.gov/popest/national/asrh/files/NC-EST2004-ALLDATA-R-File08.txt).

Case counts for race categories (American Indian or Alaska Native, Asian, Black or African American, Native Hawaiian or Other Pacific Islander, and White) are mutually exclusive and do not include persons of Hispanic ethnicity or multiple race. Multiple Race does not include persons of Hispanic ethnicity.

See Technical Notes (page 9).

See Surveillance Slide #10.

²Indicates two or more races reported for a person. Category first reported in 2003.