5. **REFERENCES**

Abou-Donia MB, Makkawy HM, Campbell GM. 1985. Pattern of neurotoxicity of *n*-hexane, methyl *n*-butyl ketone, 2,5-hexanediol, and 2,5-hexanedione alone and in combination with *o*-4-nitrophenyl phenylphosphonothioate in hens. J Toxicol Environ Health 16:85-100.

ACGIH. 1984. Threshold limit values—discussion and thirty-five year index with recommendations. American Conference of Governmental Industrial Hygienists. Cincinnati, OH, 365-368.

ACGIH. 2000. 2000 TLVs and BEIs. Threshold limit values for chemical substances and physical agents and biological exposure indices. American Conference of Governmental Industrial Hygienists. Cincinnati, OH, 80-82.

Ahlborg UG, Becking GC, Birnbaum LS, et al. 1994. Toxic equivalency factors for dioxin-like PCBs: Report on a WHO-ECEH and IPCS consultation, December 1993. Chemosphere 28:1049-1067.

Albert RE, Lewtas J, Nesnow S, et al. 1983. Comparative potency method for cancer risk assessment: Application to diesel particulate emissions. Risk Anal 3:101-107.

ATSDR. 1992. Public health assessment guidance manual. Atlanta, GA: U.S. Department of Health and Human Services, Public Health Service, Agency for Toxic Substances and Disease Registry.

ATSDR. 1994. Toxicological profile for chlorodibenzofurans. Atlanta, GA: U.S. Department of Health and Human Services, Public Health Service, Agency for Toxic Substances and Disease Registry.

ATSDR. 1995a. Toxicological profile for automotive gasoline. Atlanta, GA: U.S. Department of Health and Human Services, Public Health Service, Agency for Toxic Substances and Disease Registry. NTIS PB95-264206.

ATSDR. 1995b. Toxicological profile for polycyclic aromatic hydrocarbons (PAHs) (update). Atlanta, GA: U.S. Department of Health and Human Services, Public Health Service, Agency for Toxic Substances and Disease Registry.

ATSDR. 1995c. Toxicological profile for Stoddard solvent. Atlanta, GA: U.S. Department of Health and Human Services, Public Health Service, Agency for Toxic Substances and Disease Registry. NTIS PB95-264263.

ATSDR. 1996a. Minimal risk levels for priority substances and guidance for derivation; republication. Fed Reg 61(125):33511-33520.

ATSDR. 1996b. Toxicological profile for Toxaphene (update). Atlanta, GA: U.S. Department of Health and Human Services, Public Health Service, Agency for Toxic Substances and Disease Registry.

ATSDR. 1998a. Toxicological profile for jet fuels (JP-5 and JP-8) (February 1998 draft final). Atlanta, GA: U.S. Department of Health and Human Services, Public Health Service, Agency for Toxic Substances and Disease Registry.

ATSDR. 1998b. Toxicological profile for chlorinated dibenzo-*p*-dioxins (December 1998 final). Atlanta, GA: U.S. Department of Health and Human Services, Public Health Service, Agency for Toxic Substances and Disease Registry.

ATSDR. 1999. Toxicological profile for total petroleum hydrocarbons (TPHs). Atlanta, GA: U.S. Department of Health and Human Services, Public Health Service, Agency for Toxic Substances and Disease Registry.

ATSDR. 2000. Toxicological profile for polychlorinated biphenyls. Atlanta, GA: U.S. Department of Health and Human Services, Public Health Service, Agency for Toxic Substances and Disease Registry.

ATSDR. 2001. Guidance for the preparation of an interaction profile. Atlanta, GA: U.S. Department of Health and Human Services, Public Health Service, Agency for Toxic Substances and Disease Registry.

Berman E, House DE, Allis JW, et al. 1992. Hepatotoxic interactions of ethanol with allyl alcohol or carbon tetrachloride in rats. J Toxicol Environ Health 37(1):161-176.

Birnbaum LS, DeVito MJ. 1995. Use of toxic equivalency factors for risk assessment for dioxins and related compounds. Toxicology 105:391-401.

Bliss CI. 1939. The toxicity of poisons applied jointly. Ann Appl Biol 26:585-615.

Bond JA, Leavens TL, Seaton MJ, et al. 1998. Predicting the toxicity of chemical mixtures. Chemtech July 1998:16-23.

Calabrese EJ. 1991. Multiple chemical interactions. Chelsea, MI: Lewis Publishers, 619-622.

Caprino L, Borrelli F, Anonetti F, et al. 1983. Sex-related toxicity of somatostatin and its interaction with pentobarbital and strychnine. Toxicol Lett 17:145-149.

Carpenter DO, Arcaro K, Spink DC. 2002. Understanding the human health effects of chemical mixtures. Environ Health Perspect 110 (Suppl 1):25-42.

Cerklewski FL, Forbes RM. 1976. Influence of dietary zinc on lead toxicity in the rat. J Nutr 106:689-696.

Chen RW, Whanger PD, Weswig PH. 1975. Selenium-induced redistribution of cadmium binding to tissue proteins: A possible mechanism of protection against cadmium toxicity. Bioinorg Chem 4:125-133.

Chou T-C, Rideout DC, ed. 1991. Synergism and antagonism in chemotherapy. San Diego, CA: Academic Press, 46-47, 436.

Cogliano VJ. 1997. Plausible upper bounds: Are their sums plausible? Risk Anal 17:77-84.

Crump KS. 1984. A new method for determining allowable daily intakes. Fund Appl Toxicol 4:854-871.

Crump KS. 1995. Calculation of benchmark doses from continuous data. Risk Anal 15:79-89.

Dawson DA. 1994. Chemical mixture toxicity assessment using an alternative-species model: Applications, opportunities, and perspectives. In: Yang RSH, ed. Toxicology of chemical mixtures. New York, NY: Academic Press, 539-563.

De Rosa CT, Stevens Y-W, Johnson BL. 1993. Cancer policy framework for: Public health assessment of carcinogens in the environment. Toxicol Ind Health 9:559-575.

De Rosa CT, Johnson BL, Fay M, et al. 1996. Public health implications of hazardous waste sites: Findings, assessment and research. Food Chem Toxicol 34:1131-1138.

De Rosa CT, Brown D, Dhara R, et al. 1997a. Dioxin and dioxin-like compounds in soil, Part I: ATSDR interim policy guideline. Toxicol Ind Health 13:759-768.

De Rosa CT, Brown D, Dhara R, et al. 1997b. Dioxin and dioxin-like compounds in soil, Part II: Technical support document for ATSDR interim policy guideline. Toxicol Ind Health 13:769-804.

De Rosa CT, Brown D, Dhara R, et al. 1997c. Appendices for ATSDR interim policy guideline. J Clean Technol Environ Toxicol Occup Med 6:139-163.

De Rosa CT, Pohl HR, Williams M, et al. 1998. Public health implications of environmental exposures. Environ Health Perspect 106(Suppl 1):369-378.

Drott P, Meurling S, Gebre-Medhin M. 1993. Interactions of vitamins A and E and retinol-binding protein in healthy Swedish children—evidence of thresholds of essentiality and toxicity. Scand J Clin Lab Invest 53:275-280.

Durkin P. 1995. Development of mixtures assessment methods: Guidelines for application of the binary weight-of-evidence methodology. Submitted to The Kevric Company, Inc., Silver Spring, MD. SERA TR 95-018-001a.

El-Masri HA, Tessari JC, Yang RSH. 1996a. Exploration of an interaction threshold for the joint toxicity of trichloroethylene and 1,1-dichloroethylene: Utilization of a PBPK mode. Arch Toxicol 70:527-539.

El-Masri HA, Constan AA, Ramsdell HS, et al. 1996b. Physiologically based pharmacodynamic modeling of an interaction threshold between trichloroethylene and 1,1-dichloroethylene in Fischer 344 rats. Toxicol Appl Pharmacol 141:124-132.

El-Masri HA, Thomas RS, Sabados GR, et al. 1996c. Physiologically based pharmacodynamic modeling of the toxicologic interaction between carbon tetrachloride and Kepone. Arch Toxicol 70:704-713.

El-Masri HA, Reardon KF, Yang RSH. 1997. Integrated approaches for the analysis of toxicologic interactions of chemical mixtures. Crit Rev Toxicol 27:175-197.

EPA. 1986. U.S. Environmental Protection Agency. Guidelines for the health risk assessment of chemical mixtures. Fed Reg 51:34014-34025.

EPA. 1989a. Risk assessment guidance for superfund. Volume I. Human health evaluation manual (Part A). Washington, DC: U.S. Environmental Protection Agency, Office of Emergency and Remedial Response. EPA/540/1-89/001.

EPA. 1989b. Interim procedures for estimating risks associated with exposures to mixtures of chlorinated dibenzo-p-dioxins and -dibenzofurans (CDDs and CDFs) and 1989 update. Washington, DC: U.S. Environmental Protection Agency, Risk Assessment Forum. EPA/625/3-89/016.

EPA. 1990. Technical support document on health risk assessment of chemical mixtures. U.S. Environmental Protection Agency, Office of Research and Development. EPA/600/8-90/064.

EPA. 1993. Provisional guidance for quantitative risk assessment of polycyclic aromatic hydrocarbons. Cincinnati, OH: U.S. Environmental Protection Agency, Environmental Criteria and Assessment Office, Office of Health and Environmental Assessment. NTIS PB94-116571.

EPA. 1994. Estimating exposures and risks. In: Estimating exposure to dioxin-like compounds. Volume III: Site-specific assessment procedures. External Review Draft. Washington, DC: Office of Research and Development. EPA/600/6-88/005C, 2-1 to 2-6.

EPA. 1996. PCBs: Cancer dose-response assessment and application to environmental mixtures. Washington, DC: U.S. Environmental Protection Agency, Office of Research and Development, National Center for Environmental Assessment. EPA/600/P-96/001F, 61-63.

EPA. 2000. Supplementary guidance for conducting health risk assessment of chemical mixtures. Washington, DC: U.S. Environmental Protection Agency, Risk Assessment Forum. EPA/630R-00/002.

EPA. 2001. U.S. EPA's benchmark dose software home page. http://www.epa.gov/ncea/bmds.htm.

ERG and Durkin, P. 1995. A weight-of-evidence approach for mixtures risk assessment. Prepared by Syracuse Environmental Research Associates, Inc. and Eastern Research Group, Inc. under ERG Task No. 8495-61, as part of EPA Contract No. 68-C1-0030 for National Center for Environmental Assessment, U.S. EPA, Cincinnati, OH.

Fay RM, Feron VJ. 1996. Complex mixtures: hazard identification and risk assessment. Food Chem Toxicol 34:1175-1176.

Feron VJ, Groten JP, Van Zorge JA, et al. 1995. Toxicity studies in rats of simple mixtures of chemicals with the same or different target organs. Toxicol Lett 82-83:505-512.

Finney DJ. 1971. Probit analysis. 3rd ed. Cambridge University Press, 232-238, 246-247, 252-254, 262.

Gaughan LC, Engel J, Casida JE. 1980. Pesticide interactions: effects of organophosphorous pesticides on the metabolism, toxicity and persistence of selected pyrethroid insecticides. Pestic Biochem Physiol 14:81-85.

Gaylor D, Ryan L, Krewski D, et al. 1998. Procedures for calculating benchmark doses for health risk assessment. Reg Toxicol Pharmacol 28:150-164.

Goldstein A, Aronow L, Kalman SM. 1974. Principles of drug action: The basis of pharmacology. 2nd ed. New York, NY: John Wiley and Sons, Inc., 401, 404-405, 407.

Groten JP, Schoen ED, Van Bladeren PJ, et al. 1997. Subacute toxicity of a mixture of nine chemicals in rats: Detecting interactive effects with a fractionated two-level factorial design. Fund Appl Toxicol 36(1):15-29.

Haddad S, Krishnan K. 1998. Physiological modeling of toxicokinetic interactions: Implications for mixture risk assessment. Environ Health Perspect 106(Suppl 6):1377-1384.

Haddad S, Tardif R, Charest-Tardif G, et al. 1999a. Physiological modeling of the toxicokinetic interactions in a quaternary mixture of aromatic hydrocarbons. Toxicol Appl Pharmacol 161:249-257.

Haddad S, Tardif R, Viau C, et al. 1999b. A modeling approach to account for toxicokinetic interactions in the calculation of biological hazard index for chemical mixtures. Toxicol Let 108:303-308.

Hansen H, De Rosa CT, Pohl H, et al. 1998. Public health challenges posed by chemical mixtures. Environ Health Perspect 106(Suppl 6):1271-1280.

Harris LW, Lennox WJ, Talbot BG, et al. 1984. Toxicity of anticholinesterases: Interactions of pyridostigmine and physostigmine with soman. Drug Chem Toxicol 7:507-526.

Hermens J, Leeuwangh P, Musch A. 1985. Joint toxicity of mixtures of groups of organic aquatic pollutants to the guppy (poecilia reticulata). Ecotoxicol Environ Safety 9(3):321-326.

Hertzberg RC, Durkin PR. 1994. Influence of exposure timing and other factors on toxicologic interaction patterns. Conference on temporal aspects in risk assessment for noncancer endpoints, Dayton, OH, April 18-20, Registration and Abstract Booklet, 65.

Hertzberg RC, Rice G, Teuschler LK. 1999. Methods for health risk assessment of combustion mixtures. In: Roberts S, Teaf C, Bean J, eds. Hazardous waste incineration: Evaluating the human health and environmental risks. Boca Raton: CRC Press LLC, 105-148.

Johnson BL, De Rosa CT. 1995. Chemical mixtures released from hazardous waste sites: Implications for health risk assessment. Toxicology 105:145-156.

Jonker D, Woutersen RA, van Bladeren PJ, et al. 1990. 4-week oral toxicity study of a combination of eight chemicals in rats: Comparison with the toxicity of the individual compounds. Food Chem Toxicol 28:623-631.

Jonker D, Woutersen RA, van Bladeren PJ, et al. 1993. Subacute (4-wk) oral toxicity of a combination of four nephrotoxins in rats: Comparison with the toxicity of the individual compounds. Food Chem Toxicol 31(2):125-136.

Jonker D, Woutersen RA, Feron VJ. 1996. Toxicity of mixtures of nephrotoxicants with similar or dissimilar mode of action. Food Chem Toxicol 34:1075-1082.

Klaassen CD. 1996. Casarett and Doull's toxicology: The basic science of poisons. New York, NY: McGraw-Hill, 18, 96, 243.

Kligerman AD, Chapin RE, Erexson GL, et al. 1993. Analyses of cytogenetic damage in rodents following exposure to simulated groundwater contaminated with pesticides and a fertilizer. Mutat Res 300:125-134.

Kodell RL, Chen JJ. 1994. Reducing conservatism in risk estimation form mixtures of carcinogens. Risk Anal 14:327-332.

Konemann H. 1981. Fish toxicity tests with mixtures of more than two chemicals: A proposal for a quantitative approach and experimental results. Toxicology 19(3):229-238.

Krishnan K, Pelekis M. 1995. Hematotoxic interactions: occurrence, mechanisms and predictability. Toxicology 105(2-3):355-364.

Levander O, Argrett L. 1969. Effects of arsenic, mercury, thallium and lead on selenium metabolism in rats. Toxicol Appl Pharmacol 14:308-314.

Levin BC. 1996. New research avenues in toxicology: 7-gas N-Gas model., toxicant suppressant, and genetic toxicology. Toxicology 115:89-106.

Levin BC, Paabo M, Gurman JL, Harris SE. 1987a. Effects of exposure to single or multiple combinations of the predominant toxic gases and low oxygen atmospheres produced in fires. Fund Appl Toxicol 9:236-250.

Levin BC, Paabo M, Gurman JL, Harris SE, Braun E. 1987b. Toxicological interactions between carbon monoxide and carbon dioxide. Toxicology 47:135-164.

Levin BC, Paabo M, Schiller SB. 1991. A standard reference material for calibration of the cup furnace smoke toxicity methods for assessing the acute inhalation toxicity of combustion products. Journal of Research of the National Institute of Standards and Technology. 96:741-755.

Levin BC, Braun E, Navarro M, Paabo M. 1995. Further development of the N-Gas mathematical model: An approach for predicting the toxic potency of complex combustion mixtures. Chapter 20. In: Nelson G (ed.) Fire and Polymers II: Material and Tests for Hazard Prevention. American Chemical Society. ACS Symposium Series No. 599.

Levine RE. 1973. Pharmacology: Drug actions and reactions. Boston, MA: Little, Brown and Company, 279, 285, 353.

Lewtas J. 1985. Development of a comparative potency method for cancer risk assessment of complex mixtures using short-term *in vivo* and *in vitro* bioassays. Toxicol Ind Health 1:193-203.

Lewtas J. 1988. Genotoxicity of complex mixtures: Strategies for the identification and comparative assessment of airborne mutagens and carcinogens from combustion sources. Fund Appl Toxicol 10:571-589.

Mahaffey KR, Fowler BA. 1977. Effects pf concurrent administration of lead, cadmium, and arsenic in the rat. Environ Health Perspect 19:165-171.

Mahaffey KR, Capar SG, Gladen BC, et al. 1981. Concurrent exposure to lead, cadmium, and arsenic: Effects on toxicity and tissue metal concentrations in the rat. J Lab Clin Med 98:463-481.

Marlowe M, Cossairt A, Moon C, et al. 1985. Main and interaction effects of metallic toxins on classroom behavior. J Abnorm Child Psychol 13(2):185-198.

Marshall M, Arnott M, Jacobs M, et al. 1979. Selenium effects on the carcinogenicity and metabolism of 2-acetylaminofluorene. Cancer Lett 7:331-338.

Mehendale HM. 1994. Amplified interactive toxicity of chemicals at nontoxic levels: mechanistic considerations and implications to public health. Environ Health Perspect 102(Suppl. 9):139-149.

Moon C, Marlowe M, Stellern J, et al. 1985. Main and interaction effects of metallic pollutants on cognitive functioning. J Learn Disabil 18(4):217-221.

Mumtaz M, Colman J. 1992. The risk assessment of chemical mixtures: Fine tuning the hazard index. Presented at the Conference on Applications of Advances in Toxicology, May 19-21, Dayton, OH.

Mumtaz MM, Durkin PR. 1992. A weight-of-evidence approach for assessing interactions in chemical mixtures. Toxicol Ind Health 8:377-406.

Mumtaz MM, Hertzberg, RC. 1993. The status of interactions data in risk assessment of chemical mixtures. In: Saxena J, ed. Hazard assessment of chemicals. Vol 8. Washington, DC: Taylor and Francis, 47-79.

Mumtaz MM, De Rosa CT, Durkin PR. 1994a. Approaches and challenges in risk assessments of chemical mixtures. In: Yang RSH, ed. Toxicology of chemical mixtures: Case studies, mechanisms and novel approaches. New York, NY: Academic Press, 565-597.

Mumtaz MM, Durkin PR, Diamond GL, et al. 1994b. Exercises in the use of weight-of-evidence approach for chemical-mixture interactions. In: Andrews JS, Frumkin H, Johnson BL, et al., eds. Hazardous Waste and Public Health: International Congress on the Health Effects of Hazardous Waste, May 3-6, 1993, Atlanta, GA. Princeton, NJ: Princeton Scientific Publishing Co., Inc., 637-642.

Mumtaz MM, Cibulas W, De Rosa CT. 1995. An integrated framework to identify significant human exposures (SHELs). Chemosphere 31:2485-2489.

Mumtaz MM, Poirier KA, Colman JT. 1997. Risk assessment for chemical mixtures: Fine-tuning the hazard index approach. J Clean Technol Environ Toxicol Occup Med 6(2):189-204.

Mumtaz MM, De Rosa CT, Groten J, et al. 1998. Estimation of toxicity of chemical mixtures through modeling of chemical interactions. Environ Health Perspect 106(Suppl 6):1353-1360.

NAS. 1974. Water quality criteria, 1972. Section III—Freshwater aquatic life and wildlife: Mixtures of two or more toxicants. National Academy of Sciences, National Academy of Engineering. NTIS PB-236 199, ii-xix, 1-4, 106-108, 122-123.

NIOSH. 1976. Criteria for a recommended standard for occupational exposure to methylene chloride. Cincinnati, OH: National Institute for Occupational Safety and Health, 91–99.

NIOSH. 1992. NIOSH recommendations for occupational safety and health. Cincinnati, OH: National Institute for Occupational Safety and Health. DHHS (NIOSH) Publications No. 92-100. NTIS PB92-162536, 101.

NRC. 1988. National Research Council. Complex mixtures. Washington, DC: National Academy Press, 46-49.

NRC. 1989. Mixtures. In: Drinking water and health. Vol. 9. National Academy of Sciences, National Research Council, Safe Drinking Water Committee. Washington, DC: National Academy Press, 93–107, 121-132, 168-170.

OSHA. 1993. Occupational Safety and Health Administration. 29 CFR 1910.1000. Air contaminants; Rule. Federal Register. 58(124):35338–35351.

OSHA. 2001. Occupational Safety and Health Administration. OSHA regulations (Standards - 29 CFR): Air contaminants. - 1910.1000. http://www.osha-slc.gov/OshStd_data/1910_1000.html

Oskarsson A, Lind B. 1985. Increased lead levels in brain after long-term treatment with lead and dithiocarbamate or thiuram derivatives in rats. Acta Pharmacol Toxicol 56:309-315.

Oskarsson A, Ljungberg T, Stahle L, et al. 1986a. Behavioral and neurochemical effects after combined perinatal treatment of rats with disulfiram. Neurobehav Toxicol Teratol 8:591-599.

Oskarsson A, Olson L, Palmer MR, et al. 1986b. Increased lead concentration in brain and potentiation of lead-induced neuronal depression in rats after combined treatment with lead and disulfiram. Environ Res 41:623-632.

Pelekis M, Krishnan K. 1997. Assessing the relevance of rodent data on chemical interactions for health risk assessment purposes: A case study with dichloromethane–toluene mixture. Reg Toxicol Pharmacol 25:79-86.

Plackett RL, Hewlett PS. 1952. Quantal responses to mixtures of poisons. J Royal Stat Soc B 14:141-163.

Pohl H, Holler J. 1995. Halogenated aromatic hydrocarbons and toxicity equivalency factors (TEFs) from the public health assessment perspective. Chemosphere 31:2547-2559.

Pohl H, De Rosa C, Holler J. 1995. Public health assessment for dioxins exposure from soil. Chemosphere 31:2437-2454.

Pohl H, Hansen H, Chou C-HSC. 1997. Public health guidance values for chemical mixtures: Current practice and future directions. Reg Toxicol Pharmacol 26:322-329.

Safe SH. 1998. Hazard and risk assessment of chemical mixtures using the toxic equivalency factor approach. Environ Health Perspect 106(Suppl 4):1051-1058.

Sagripanti JL, Goering PL, Lamanna A. 1991. Interaction of copper with DNA and antagonism by other metals. Toxicol Appl Pharmacol 110:477-485.

Stacey NH. 1989. Toxicity of mixtures of trichloroethylene, tetrachloroethylene and 1,1,1-trichloroethylene: similarity of in vitro to in vivo responses. Toxicol Ind Health 5(3):441-450.

Smyth HF, Weil CS, West JS, et al. 1969. An exploration of joint toxic action: Twenty-seven industrial chemicals intubated in rats in all possible pairs. Toxicol Appl Pharmacol 14:340-347.

Smyth HF, Weil CS, West JS, et al. 1970. An exploration of joint toxic action. II. Equitoxic versus equivolume mixtures. Toxicol Appl Pharmacol 17:489-503.

Svendsgaard DJ, Hertzberg RC. 1994. Statistical methods for the toxicological evaluation of the additivity assumption as used in the environmental protection agency chemical mixture risk assessment guidelines. In: RSH Yang, ed. Toxicology of chemical mixtures. New York, NY: Academic Press, 100-155.

Tardif R, Lapare S, Charest-Tardif G, et al. 1995. Physiologically-based pharmacokinetic modeling of a mixture of toluene and xylene in humans. Risk Anal 15:335-342.

Tardif R, Charest-Tardif G, Brodeur J, et al. 1997. Physiologically-based pharmacokinetic modeling of a ternary mixture of alkyl benzenes in rats and humans. Toxicol Appl Pharmacol 144:120-134.

Thomas RS, Yang RSH, Morgan DG, et al. 1996. PBPK modeling/Monte Carlo simulation of methylene chloride kinetic changes in mice in relation to age and acute, subchronic, and chronic inhalation exposure. Environ Health Perspect 104(8):858-865.

Van den Berg M, Birnbaum L, Bosveld ATC et al. 1998. Toxic equivalency factors (TEFs) for PCBs, PCDDs, PCDFs for humans and wildlife. Environ Health Perspect 106(12):775-792.

Verhaar JMH, Morroni JR, Reardon KF et al. 1997. A proposed approach to study the toxicology of complex mixtures of petroleum products: The integrated use of QSAR, lumping analysis and PBPK/PD modeling. Environ Health Perspect 105(Suppl 1):179-195.

Withey RJ, Hall JW. 1975. The joint toxic action of perchloroethylene with benzene or toluene in rats. Toxicology 4(1):5-15.

Woo Y, Di Carlo FJ, Arcos JC, et al. 1994. Assessment of carcinogenic hazard of chemical mixtures through analysis of binary chemical interaction data. Environ Health Perspect 102(Suppl 9):113-118.

Yang RSH. 1994. Toxicology of chemical mixtures derived from hazardous waste sites or application of pesticides and fertilizers. In: RSH Yang, ed. Toxicology of chemical mixtures. New York, NY: Academic Press, 100-155.

Yang RSH. 2000. Health risks and preventive research strategy for deployed U.S. forces from toxicologic interactions among potentially harmful agents, in *Strategies to Protect The Health of Deployed U.S. Forces: Assessing Health Risks to Deployed U.S. Forces*, National Academy Press, Washington, DC, pp. 150-182.

Yang RSH, Thomas RS, Gustafson DL, et al. 1998. Approaches to developing alternative and predictive toxicology based on PBPK/PD and QSAR modeling. Environ Health Perspect 106(Suppl 6):1385-1393.