


Estimated Number and Percentage of AIDS Cases among Adult and Adolescent Men Who Have Sex with Men 1985–2006—United States and Dependent Areas



Note. The data have been adjusted for reporting delay and cases without risk factor information were proportionally redistributed.



Estimated AIDS Cases in Males, Cumulative through 2006—50 States and DC

Of 973,354 cases of AIDS in adults and adolescents

81% were in males

59% of cases in males were attributed to
male-to-male sexual contact

4% of cases in males were aged 13-24 years

64% of cases in males aged 13-24 were
attributed to male-to-male sexual contact


Of AIDS cases diagnosed during 2006, 43% of cases in
adults and adolescents were attributed to male-to-male
sexual contact



Note. The data have been adjusted for reporting delay and cases without risk factor information were proportionally redistributed.




Estimated Number of AIDS Cases and Deaths among Men Who Have Sex with Men, 1985–2006—United States and Dependent Areas



Note. The data have been adjusted for reporting delay and cases without risk factor information were proportionally redistributed.



Estimated Number of HIV/AIDS Cases among Adults and Adolescents, by Transmission Category, 1994–2006—25 States



Note. The data have been adjusted for reporting delay and cases without risk factor information were proportionally redistributed.

*Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

†Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.



Estimated Number and Proportion of HIV/AIDS Cases among Adults and Adolescents by Transmission Category, 2006—33 States

	No.	%
Male-to-male sexual contact	17,465	50
Injection drug use	4,728	13
Male-to-male sexual contact and injection drug use	1,180	3
High-risk heterosexual contact*	11,584	33
Other/not identified†	223	1




Note. The data have been adjusted for reporting delay and cases without risk factor information were proportionally redistributed.

*Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

†Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.




Estimated Number of AIDS Cases among Adult and Adolescent Men Who Have Sex with Men, by Race/Ethnicity, 1985–2006—United States and Dependent Areas



Note. The data have been adjusted for reporting delay and cases without risk factor information were proportionally redistributed.



Proportion of Estimated HIV/AIDS Cases among Adult and Adolescent Men Who Have Sex with Men, by Race/Ethnicity 1994–2006—25 States



Note. The data have been adjusted for reporting delay and cases without risk factor information were proportionally redistributed.



Estimated Number of Adult and Adolescent Men Who Have Sex with Men Living with HIV/AIDS by Race/Ethnicity, 2001–2006—33 States

Year	White, not Hispanic	Black, not Hispanic	Hispanic	Asian/ Pacific Islander	American Indian/ Alaska Native	Total*
2001	83,582	49,326	23,560	843	669	159,212
2002	88,395	53,341	26,050	934	718	170,776
2003	92,941	57,188	28,381	1,137	769	181,770
2004	97,799	61,110	30,813	1,292	815	193,245
2005	102,843	65,372	33,372	1,480	869	205,424
2006	108,259	70,242	36,043	1,669	938	218,677




Note. The data have been adjusted for reporting delay and cases without risk factor information were proportionally redistributed.

*Total includes persons of unknown race or multiple races.




Proportion of Estimated HIV/AIDS Cases among Adult and Adolescent Men Who Have Sex with Men by Race/Ethnicity, 2006—33 States



Note. The data have been adjusted for reporting delay and cases without risk factor information were proportionally redistributed.




Estimated Number of AIDS Cases among Adult and Adolescent Men Who Have Sex with Men, by Region and Race/Ethnicity, 2006—50 States and DC



Note. The data have been adjusted for reporting delay and cases without risk factor information were proportionally redistributed.




Estimated Number of HIV/AIDS Cases among Adult and Adolescent Men Who Have Sex with Men, by Age Group, 2001–2006—33 States



Note. The data have been adjusted for reporting delay and cases without risk factor information were proportionally redistributed.




Estimated Number of HIV/AIDS Cases among Men Who Have Sex with Men, Aged 13–24 years, by Race/Ethnicity 2001–2006—33 States



Note. The data have been adjusted for reporting delay and cases without risk factor information were proportionally redistributed.



Proportion of HIV/AIDS Diagnoses among Men Who Have Sex with Men, Aged 13–24 years by Race/Ethnicity, 2006—33 States



Note. The data have been adjusted for reporting delay and cases without risk factor information were proportionally redistributed.



Estimated Number of HIV/AIDS Cases among Adult and Adolescent Men Who Have Sex with Men by Race/Ethnicity and Population of Area of Residence, 2006—33 States

	≥50,000	50,000— 499,999	Non-metropolitan area
White, not Hispanic	5,604	1,025	513
Black, not Hispanic	5,190	852	537
Hispanic	2,972	245	110
Asian/Pacific Islander	197	18	4
American Indian/Alaska Native	56	16	19
Unknown or multiple races	59	11	6



Note. The data have been adjusted for reporting delay and cases without risk factor information were proportionally redistributed.

