

APPENDIX A

ATSDR MINIMAL RISK LEVELS AND WORKSHEETS

The Comprehensive Environmental Response, Compensation, and Liability Act (CERCLA) [42 U.S.C. 9601 et seq.], as amended by the Superfund Amendments and Reauthorization Act (SARA) [Pub. L. 99–499], requires that the Agency for Toxic Substances and Disease Registry (ATSDR) develop jointly with the U.S. Environmental Protection Agency (EPA), in order of priority, a list of hazardous substances most commonly found at facilities on the CERCLA National Priorities List (NPL); prepare toxicological profiles for each substance included on the priority list of hazardous substances; and assure the initiation of a research program to fill identified data needs associated with the substances.

The toxicological profiles include an examination, summary, and interpretation of available toxicological information and epidemiologic evaluations of a hazardous substance. During the development of toxicological profiles, Minimal Risk Levels (MRLs) are derived when reliable and sufficient data exist to identify the target organ(s) of effect or the most sensitive health effect(s) for a specific duration for a given route of exposure. An MRL is an estimate of the daily human exposure to a hazardous substance that is likely to be without appreciable risk of adverse noncancer health effects over a specified duration of exposure. MRLs are based on noncancer health effects only and are not based on a consideration of cancer effects. These substance-specific estimates, which are intended to serve as screening levels, are used by ATSDR health assessors to identify contaminants and potential health effects that may be of concern at hazardous waste sites. It is important to note that MRLs are not intended to define clean-up or action levels.

MRLs are derived for hazardous substances using the no-observed-adverse-effect level/uncertainty factor approach. They are below levels that might cause adverse health effects in the people most sensitive to such chemical-induced effects. MRLs are derived for acute (1–14 days), intermediate (15–364 days), and chronic (365 days and longer) durations and for the oral and inhalation routes of exposure. Currently, MRLs for the dermal route of exposure are not derived because ATSDR has not yet identified a method suitable for this route of exposure. MRLs are generally based on the most sensitive chemical-induced end point considered to be of relevance to humans. Serious health effects (such as irreparable damage to the liver or kidneys, or birth defects) are not used as a basis for establishing MRLs. Exposure to a level above the MRL does not mean that adverse health effects will occur.

APPENDIX A

MRLs are intended only to serve as a screening tool to help public health professionals decide where to look more closely. They may also be viewed as a mechanism to identify those hazardous waste sites that are not expected to cause adverse health effects. Most MRLs contain a degree of uncertainty because of the lack of precise toxicological information on the people who might be most sensitive (e.g., infants, elderly, nutritionally or immunologically compromised) to the effects of hazardous substances. ATSDR uses a conservative (i.e., protective) approach to address this uncertainty consistent with the public health principle of prevention. Although human data are preferred, MRLs often must be based on animal studies because relevant human studies are lacking. In the absence of evidence to the contrary, ATSDR assumes that humans are more sensitive to the effects of hazardous substance than animals and that certain persons may be particularly sensitive. Thus, the resulting MRL may be as much as a hundredfold below levels that have been shown to be nontoxic in laboratory animals.

Proposed MRLs undergo a rigorous review process: Health Effects/MRL Workgroup reviews within the Division of Toxicology, expert panel peer reviews, and agencywide MRL Workgroup reviews, with participation from other federal agencies and comments from the public. They are subject to change as new information becomes available concomitant with updating the toxicological profiles. Thus, MRLs in the most recent toxicological profiles supersede previously published levels. For additional information regarding MRLs, please contact the Division of Toxicology, Agency for Toxic Substances and Disease Registry, 1600 Clifton Road, Mailstop E-29, Atlanta, Georgia 30333.

APPENDIX A

MINIMAL RISK LEVEL WORKSHEET

Chemical Name: Endosulfan
CAS Number: 115-29-7
Date: April 6, 2000
Profile Status: Third Draft Post Public
Route: Inhalation Oral
Duration: Acute Intermediate Chronic
Graph Key: 66r
Species: Rat

Minimal Risk Level: 0.005 mg/kg/day ppm

Reference: Banerjee BD, Hussain QZ. 1986. Effect of sub-chronic endosulfan exposure on humoral and cell-mediated immune responses in albino rats. Arch Toxicol 59:279-284.

Experimental design: Groups of male Wistar (10–12/group) (85–90 g body weight) received technical grade endosulfan (α - and β -endosulfan in the ratio of 7:3) in their diets at dietary levels of 0, 5, 10, or 20 ppm (equivalent to 0, 0.45, 0.9, and 1.8 mg/kg/day, using the EPA [1988d] food factor for male Wistar rats, subchronic duration). Test diets were prepared by dissolving the endosulfan in groundnut oil and mixing this into standard laboratory diet. Samples analyzed from each batch of diet indicated that the actual levels of endosulfan in the diet were within 10% of the desired levels. Control animals received a diet with an equal amount of groundnut oil mixed in. Rats were randomly allocated to groups and were caged 4 to a stainless steel, mesh-bottom cage. Food and water were available to these rats on “as needed” basis for between 8 and 22 weeks. At weeks 8, 12, 18, and 22, between 10 and 12 rats were selected from each group and sacrificed. Twenty days before sacrifice, the rats were immunized by injecting 0.2 mL of tetanus toxin mixed with an equal volume of Freund’s adjuvant subcutaneously. An additional group of 10–12 rats per dose were sacrificed at the time periods indicated, but these rats were not immunized with the tetanus toxin and adjuvant. At the time of sacrifice, the liver, spleen, and thymus were removed and weighed, blood samples were collected by cardiac puncture, and peritoneal exudate was collected by washing the peritoneal cavity with between 10 and 15 mL of RMPI medium.

The antibody titer to tetanus toxin was estimated using an indirect hemagglutination technique. Briefly, a suspension of sheep red blood cells was treated with tannic acid (1:20,000 dilution) and used for antigen coating. Tetanus toxin was then mixed with the treated sheep red blood cell and antibody titers were determined using the first dilution where no visible agglutination was observed. Serum proteins were determined using zone electrophoresis. Quantitation of serum levels of IgG and IgM was performed using radial immunodiffusion in agarose containing either anti IgG or anti IgM. The leukocyte migration inhibition test was performed using leukocytes isolated from rat blood by sequential centrifugation and washing. Migration from micro capillary tubes was measured using a camera lucida and migration into control medium was compared with migration into medium containing tetanus toxin. The macrophage migration inhibition test was performed using macrophages isolated from the peritoneal exudate by sequential centrifugation and washing. Migration was measured as described above for the leukocytes.

APPENDIX A

Effects noted in study and corresponding doses: No difference between the controls and rats given diets containing 5 ppm endosulfan was observed in any of the parameters measured. Rats consuming diets containing 10 ppm endosulfan and treated with tetanus toxin had significantly decreased serum IgG levels at weeks 12, 18, and 22. These rats also had significantly decreased antibody titer to tetanus toxin at weeks 8, 12, 18, and 22. Leukocyte and macrophage migration was also significantly inhibited at weeks 8, 12, 18, and 22. The magnitude of the differences between the 10 ppm rats and the controls increased at each later time point. These rats also had a significantly increased albumin to globulin ratio at week 22. Rats consuming diets containing 20 ppm showed all of the same changes as the rats at 10 ppm but to a greater degree. In addition, at weeks 2, 18, and 22, these rats showed a significantly increased albumin to globulin ratio, and at 22 weeks, these rats showed a significant decrease in relative spleen weight. No effect on the relative thymus weight was observed at any dose at any of the times tested.

Dose and end point used for MRL derivation: 0.45 mg/kg/day; depressed immune response.

[X] NOAEL [] LOAEL

Uncertainty Factors used in MRL derivation:

- 10 for use of a LOAEL
- 10 for extrapolation from animals to humans
- 10 for human variability

Was a conversion factor used from ppm in food or water to a mg/body weight dose?

Yes, 0.45 mg/kg/day was calculated by multiplying the dietary level of 5 ppm (5 mg endosulfan/kg diet) by the food factor for male Wistar rats in a subchronic study of 0.09 kg diet/ kg body weight/day (EPA 1988d).

If an inhalation study in animals, list conversion factors used in determining human equivalent dose:

Not applicable

Was a conversion used from intermittent to continuous exposure?

No.

Other additional studies or pertinent information that lend support to this MRL: Choice of this immunological end point as the basis for derivation of the intermediate-duration oral MRL is supported by the observation of similar effects in rats following ingestion of higher doses of endosulfan for shorter periods of time (Banerjee and Hussain 1986, 1987). Additional support of the positive findings is provided in Khurana et al. (1998) in which decreased macrophage functionality was observed, in the absence of any other apparent toxicological effects, in 1-day-old broiler chicks fed 30 ppm endosulfan in the diet for 4 or 8 weeks. The absence of observed immunotoxicity in the study by Vos et al. (1982) in rats does not contradict these findings since not all of the same immunological parameters were evaluated in the by Vos et al. (1982) and a shorter period of exposure was used. The only other intermediate-duration study that reported effects at relatively low doses of endosulfan is that by Hoechst (1984a) in which a decrease in litter weight was observed in F1B pups during the lactation to weaning period at an endosulfan dose level of 0.8 mg/kg/day. However, EPA review the study and because there was no corroborative finding of a decrease in the number of pups per litter or in pup weight, the decrease in litter weight was not considered to be treatment-related (IRIS 2000).

Agency Contact (Chemical Manager): Lori L. Miller, M.P.H.

APPENDIX A

MINIMAL RISK LEVEL WORKSHEET

Chemical Name: Endosulfan
CAS Number: 115-29-7
Date: April 6, 2000
Profile Status: Final
Route: Inhalation Oral
Duration: Acute Intermediate Chronic
Graph Key: 103d
Species: Dog

Minimal Risk Level: 0.002 mg/kg/day ppm

Reference: Hoechst. 1989c. Endosulfan - substance technical (code HOE 02671 OI ZD96 0002): Testing for toxicity by repeated oral administration (1-year feeding study) to Beagle dogs. Conducted for Hoechst Aktiengesellschaft, Frankfurt, Germany. Project no. 87.0643.

Experimental design: Male and female beagle dogs (6/sex/dose) were fed diets containing endosulfan at dietary concentrations of 0, 3, 10, and 30 ppm for 1 year. An additional group was fed diets containing endosulfan at concentrations of 30 ppm for 54 days, 45 ppm for 52 days, and 60 ppm for up to 40 days, but this group is considered separately since all surviving animals were sacrificed by day 147. When multiplied by estimated food factors of 0.067 and 0.060 kg food/kg body weight/day for males and females, respectively, these dietary concentrations are equivalent to 0, 0.2, 0.67, and 2 mg/kg/day for males and 0, 0.18, 0.6, and 1.8 mg/kg/day for females. The food factors were calculated based upon the actual average body weight and food consumption data from the study. Test diets were prepared daily immediately before feeding from endosulfan-containing cornmeal premixes which were mixed into the daily feed ration of 1 kg (males) and 0.8 kg (females) wet diet. Control animals received the basal diet containing the same proportion of cornmeal as the 30/45/60 ppm group's diet. Dogs were randomly allocated to dose groups and were housed in separate kennels.

Animals were observed twice daily for deaths and behavioral abnormalities. General health and food consumption were checked only daily and body weights were checked once weekly. Tests for hepatic function (bromsulphthalein) and renal function (phenosulphthalein), tests for reflexes and righting reactions, ophthalmological exams, hearing tests, and dental inspections for all dogs were performed before the start of the study and before termination. In addition, the two highest dose groups were tested after 6 weeks and every 3 months. Hematology, clinical chemistry, and urinalysis were performed at the start and at termination of the study, after 6 weeks, and again every 3 months. At termination, all dogs were examined grossly, the weights of the major organs were determined, and a wide spectrum of tissues were examined microscopically.

Effects noted in study and corresponding doses: Because the highest dose group (30/45/60 ppm) was sacrificed before 6 months, this group is not discussed here. No effects were observed in the dogs receiving diets containing 3 ppm endosulfan. No deaths occurred during the study that could be directly attributed to consumption of diets containing endosulfan. One male in the 30 ppm dietary group was sacrificed in extremis and necropsy revealed severe inflammation of the mediastinum (origin unknown). Approximately 2.5–6 hours after consuming test diets containing 30 ppm endosulfan, three males and two females exhibited convulsive spasms of the abdominal muscles and jaws without vomiting. It is not clear whether this effect was neurological or gastrointestinal in origin. Routine histopathological examination of the central and peripheral nervous system and eyes revealed no adverse effects. No abnormalities were detected in animal's pupillary or blink reflexes; flexor patellar, anal, or cutaneous reflexes; or extensor postural thrust reaction, placing reaction, or righting abilities. Cholinesterase activities were also not

APPENDIX A

affected by treatment with endosulfan. Serum alkaline phosphatase was significantly elevated in males at 10 and 30 ppm at most test intervals and in females at early intervals at 10 ppm and at most intervals at 30 ppm, indicating some form of tissue damage. In dogs, alkaline phosphatase is primarily of hepatic origin, but gross and microscopic pathology was unremarkable. Lactate dehydrogenase was also elevated in males and females from the 10 ppm group at 9 months and at 30 ppm at 12 months. Routine and gross histopathological examination of the major organs and tissues including skin, reproductive organs, lymphoreticular organs, and endocrine and exocrine glands revealed no adverse effects. In addition, ophthalmology, hematology, urinalysis, and renal function tests revealed no adverse effects of endosulfan consumption.

Dose and end point used for MRL derivation: 0.18 mg/kg/day; for increase in serum alkaline phosphatase activity.

NOAEL LOAEL

Uncertainty Factors used in MRL derivation:

- 10 for use of a LOAEL
- 10 for extrapolation from animals to humans
- 10 for human variability

Was a conversion factor used from ppm in food or water to a mg/body weight dose?

Yes, 0.18 mg/kg/day was calculated by multiplying the dietary level of 3 ppm (3 mg endosulfan/kg diet) by food factors of 0.067 and 0.06 kg food/kg body weight/day for males and females, respectively, derived from food consumption and body weight data provided in the study. The female food factor (0.06 kg food/kg body weight/day) was used because this yielded the most conservative dose estimate.

If an inhalation study in animals, list conversion factors used in determining human equivalent dose: Not applicable.

Was a conversion used from intermittent to continuous exposure?

No.

Other additional studies or pertinent information that lend support to this MRL: The choice of this end point (hepatic toxicity as evidenced by increased serum alkaline phosphatase activity) is supported by the observation of hydropic hepatic cells in male rats that consumed a diet that provided a dose of endosulfan of approximately 5 mg/kg/day for 2 years (FMC 1959b). The hepatic NOAEL in that study was approximately 1 mg/kg/day. In another 2-year dietary study in rats it was observed that male and female rats that received doses of endosulfan of approximately 0.6–0.6 mg/kg/day gained 11–15% less weight (estimated as the difference between starting weight and final weight) than the controls; however, the biological significance of this observation is unclear (Hack et al. 1995).

Agency Contact (Chemical Manager): Lori L. Miller, M.P.H.

APPENDIX B

USER'S GUIDE

Chapter 1

Public Health Statement

This chapter of the profile is a health effects summary written in non-technical language. Its intended audience is the general public especially people living in the vicinity of a hazardous waste site or chemical release. If the Public Health Statement were removed from the rest of the document, it would still communicate to the lay public essential information about the chemical.

The major headings in the Public Health Statement are useful to find specific topics of concern. The topics are written in a question and answer format. The answer to each question includes a sentence that will direct the reader to chapters in the profile that will provide more information on the given topic.

Chapter 2

Tables and Figures for Levels of Significant Exposure (LSE)

Tables (2-1, 2-2, and 2-3) and figures (2-1 and 2-2) are used to summarize health effects and illustrate graphically levels of exposure associated with those effects. These levels cover health effects observed at increasing dose concentrations and durations, differences in response by species, minimal risk levels (MRLs) to humans for noncancer end points, and EPA's estimated range associated with an upper-bound individual lifetime cancer risk of 1 in 10,000 to 1 in 10,000,000. Use the LSE tables and figures for a quick review of the health effects and to locate data for a specific exposure scenario. The LSE tables and figures should always be used in conjunction with the text. All entries in these tables and figures represent studies that provide reliable, quantitative estimates of No-Observed-Adverse-Effect Levels (NOAELs), Lowest-Observed-Adverse-Effect Levels (LOAELs), or Cancer Effect Levels (CELs).

The legends presented below demonstrate the application of these tables and figures. Representative examples of LSE Table 2-1 and Figure 2-1 are shown. The numbers in the left column of the legends correspond to the numbers in the example table and figure.

LEGEND

See LSE Table 2-1

- (1) Route of Exposure One of the first considerations when reviewing the toxicity of a substance using these tables and figures should be the relevant and appropriate route of exposure. When sufficient data exists, three LSE tables and two LSE figures are presented in the document. The three LSE tables present data on the three principal routes of exposure, i.e., inhalation, oral, and dermal (LSE Table 2-1, 2-2, and 2-3, respectively). LSE figures are limited to the inhalation (LSE Figure 2-1) and oral (LSE Figure 2-2) routes. Not all substances will have data on each route of exposure and will not therefore have all five of the tables and figures.

APPENDIX B

- (2) Exposure Period Three exposure periods - acute (less than 15 days), intermediate (15–364 days), and chronic (365 days or more) are presented within each relevant route of exposure. In this example, an inhalation study of intermediate exposure duration is reported. For quick reference to health effects occurring from a known length of exposure, locate the applicable exposure period within the LSE table and figure.
- (3) Health Effect The major categories of health effects included in LSE tables and figures are death, systemic, immunological, neurological, developmental, reproductive, and cancer. NOAELs and LOAELs can be reported in the tables and figures for all effects but cancer. Systemic effects are further defined in the "System" column of the LSE table (see key number 18).
- (4) Key to Figure Each key number in the LSE table links study information to one or more data points using the same key number in the corresponding LSE figure. In this example, the study represented by key number 18 has been used to derive a NOAEL and a Less Serious LOAEL (also see the 2 "18r" data points in Figure 2-1).
- (5) Species The test species, whether animal or human, are identified in this column. Section 2.5, "Relevance to Public Health," covers the relevance of animal data to human toxicity and Section 2.3, "Toxicokinetics," contains any available information on comparative toxicokinetics. Although NOAELs and LOAELs are species specific, the levels are extrapolated to equivalent human doses to derive an MRL.
- (6) Exposure Frequency/Duration The duration of the study and the weekly and daily exposure regimen are provided in this column. This permits comparison of NOAELs and LOAELs from different studies. In this case (key number 18), rats were exposed to 1,1,2,2-tetrachloroethane via inhalation for 6 hours per day, 5 days per week, for 3 weeks. For a more complete review of the dosing regimen refer to the appropriate sections of the text or the original reference paper, i.e., Nitschke et al. 1981.
- (7) System This column further defines the systemic effects. These systems include: respiratory, cardiovascular, gastrointestinal, hematological, musculoskeletal, hepatic, renal, and dermal/ocular. "Other" refers to any systemic effect (e.g., a decrease in body weight) not covered in these systems. In the example of key number 18, 1 systemic effect (respiratory) was investigated.
- (8) NOAEL A No-Observed-Adverse-Effect Level (NOAEL) is the highest exposure level at which no harmful effects were seen in the organ system studied. Key number 18 reports a NOAEL of 3 ppm for the respiratory system which was used to derive an intermediate exposure, inhalation MRL of 0.005 ppm (see footnote "b").
- (9) LOAEL A Lowest-Observed-Adverse-Effect Level (LOAEL) is the lowest dose used in the study that caused a harmful health effect. LOAELs have been classified into "Less Serious" and "Serious" effects. These distinctions help readers identify the levels of exposure at which adverse health effects first appear and the gradation of effects with increasing dose. A brief description of the specific endpoint used to quantify the adverse effect accompanies the LOAEL. The respiratory effect reported in key number 18 (hyperplasia) is a Less serious LOAEL of 10 ppm. MRLs are not derived from Serious LOAELs.
- (10) Reference The complete reference citation is given in chapter 8 of the profile.
- (11) CEL A Cancer Effect Level (CEL) is the lowest exposure level associated with the onset of carcinogenesis in experimental or epidemiologic studies. CELs are always considered serious

APPENDIX B

effects. The LSE tables and figures do not contain NOAELs for cancer, but the text may report doses not causing measurable cancer increases.

- (12) Footnotes Explanations of abbreviations or reference notes for data in the LSE tables are found in the footnotes. Footnote "b" indicates the NOAEL of 3 ppm in key number 18 was used to derive an MRL of 0.005 ppm.

LEGEND**See Figure 2-1**

LSE figures graphically illustrate the data presented in the corresponding LSE tables. Figures help the reader quickly compare health effects according to exposure concentrations for particular exposure periods.

- (13) Exposure Period The same exposure periods appear as in the LSE table. In this example, health effects observed within the intermediate and chronic exposure periods are illustrated.
- (14) Health Effect These are the categories of health effects for which reliable quantitative data exists. The same health effects appear in the LSE table.
- (15) Levels of Exposure concentrations or doses for each health effect in the LSE tables are graphically displayed in the LSE figures. Exposure concentration or dose is measured on the log scale "y" axis. Inhalation exposure is reported in mg/m³ or ppm and oral exposure is reported in mg/kg/day.
- (16) NOAEL In this example, 18r NOAEL is the critical endpoint for which an intermediate inhalation exposure MRL is based. As you can see from the LSE figure key, the open-circle symbol indicates to a NOAEL for the test species-rat. The key number 18 corresponds to the entry in the LSE table. The dashed descending arrow indicates the extrapolation from the exposure level of 3 ppm (see entry 18 in the Table) to the MRL of 0.005 ppm (see footnote "b" in the LSE table).
- (17) CEL Key number 38r is 1 of 3 studies for which Cancer Effect Levels were derived. The diamond symbol refers to a Cancer Effect Level for the test species-mouse. The number 38 corresponds to the entry in the LSE table.
- (18) Estimated Upper-Bound Human Cancer Risk Levels This is the range associated with the upper-bound for lifetime cancer risk of 1 in 10,000 to 1 in 10,000,000. These risk levels are derived from the EPA's Human Health Assessment Group's upper-bound estimates of the slope of the cancer dose response curve at low dose levels (q_1^*).
- (19) Key to LSE Figure The Key explains the abbreviations and symbols used in the figure.

SAMPLE

1⁶

TABLE 2-1. Levels of Significant Exposure to [Chemical x] – Inhalation

Key to figure ^a	Species	Exposure frequency/ duration	System	NOAEL (ppm)	LOAEL (effect)		Reference
					Less serious (ppm)	Serious (ppm)	
INTERMEDIATE EXPOSURE							
2 ⁶	5	6	7	8	9		10
3 ⁶	Systemic	9	9	9	9		9
4 ⁶	18	Rat	13 wk 5d/wk 6hr/d	Resp	3 ^b	10 (hyperplasia)	Nitschke et al. 1981
<hr style="border-top: 1px dashed black;"/>							
CHRONIC EXPOSURE							
						11	
	Cancer					9	
38	Rat	18 mo 5d/wk 7hr/d				20	(CEL, multiple organs) Wong et al. 1982
39	Rat	89–104 wk 5d/wk 6hr/d				10	(CEL, lung tumors, nasal tumors) NTP 1982
40	Mouse	79–103 wk 5d/wk 6hr/d				10	(CEL, lung tumors, hemangiosarcomas) NTP 1982

^a The number corresponds to entries in Figure 2-1.

12⁶

^b Used to derive an intermediate inhalation Minimal Risk Level (MRL) of 5×10^{-3} ppm; dose adjusted for intermittent exposure and divided by an uncertainty factor of 100 (10 for extrapolation from animal to humans, 10 for human variability).

SAMPLE


13

Figure 2-1. Levels of Significant Exposure to [Chemical X] – Inhalation

14

15

16



17

18

19

Chapter 2 (Section 2.5)

Relevance to Public Health

The Relevance to Public Health section provides a health effects summary based on evaluations of existing toxicologic, epidemiologic, and toxicokinetic information. This summary is designed to present interpretive, weight-of-evidence discussions for human health end points by addressing the following questions.

1. What effects are known to occur in humans?
2. What effects observed in animals are likely to be of concern to humans?
3. What exposure conditions are likely to be of concern to humans, especially around hazardous waste sites?

The section covers end points in the same order they appear within the Discussion of Health Effects by Route of Exposure section, by route (inhalation, oral, dermal) and within route by effect. Human data are presented first, then animal data. Both are organized by duration (acute, intermediate, chronic). *In vitro* data and data from parenteral routes (intramuscular, intravenous, subcutaneous, etc.) are also considered in this section. If data are located in the scientific literature, a table of genotoxicity information is included.

The carcinogenic potential of the profiled substance is qualitatively evaluated, when appropriate, using existing toxicokinetic, genotoxic, and carcinogenic data. ATSDR does not currently assess cancer potency or perform cancer risk assessments. Minimal risk levels (MRLs) for noncancer end points (if derived) and the end points from which they were derived are indicated and discussed.

Limitations to existing scientific literature that prevent a satisfactory evaluation of the relevance to public health are identified in the Data Needs section.

Interpretation of Minimal Risk Levels

Where sufficient toxicologic information is available, we have derived minimal risk levels (MRLs) for inhalation and oral routes of entry at each duration of exposure (acute, intermediate, and chronic). These MRLs are not meant to support regulatory action; but to acquaint health professionals with exposure levels at which adverse health effects are not expected to occur in humans. They should help physicians and public health officials determine the safety of a community living near a chemical emission, given the concentration of a contaminant in air or the estimated daily dose in water. MRLs are based largely on toxicological studies in animals and on reports of human occupational exposure.

MRL users should be familiar with the toxicologic information on which the number is based. Chapter 2.5, "Relevance to Public Health," contains basic information known about the substance. Other sections such as 2.8, "Interactions with Other Substances," and 2.9, "Populations that are Unusually Susceptible" provide important supplemental information.

MRL users should also understand the MRL derivation methodology. MRLs are derived using a modified version of the risk assessment methodology the Environmental Protection Agency (EPA) provides (Barnes and Dourson 1988) to determine reference doses for lifetime exposure (RfDs).

APPENDIX B

To derive an MRL, ATSDR generally selects the most sensitive endpoint which, in its best judgement, represents the most sensitive human health effect for a given exposure route and duration. ATSDR cannot make this judgement or derive an MRL unless information (quantitative or qualitative) is available for all potential systemic, neurological, and developmental effects. If this information and reliable quantitative data on the chosen endpoint are available, ATSDR derives an MRL using the most sensitive species (when information from multiple species is available) with the highest NOAEL that does not exceed any adverse effect levels. When a NOAEL is not available, a lowest-observed-adverse-effect level (LOAEL) can be used to derive an MRL, and an uncertainty factor (UF) of 10 must be employed. Additional uncertainty factors of 10 must be used both for human variability to protect sensitive subpopulations (people who are most susceptible to the health effects caused by the substance) and for interspecies variability (extrapolation from animals to humans). In deriving an MRL, these individual uncertainty factors are multiplied together. The product is then divided into the inhalation concentration or oral dosage selected from the study. Uncertainty factors used in developing a substance-specific MRL are provided in the footnotes of the LSE Tables.

APPENDIX C

ACRONYMS, ABBREVIATIONS, AND SYMBOLS

ACGIH	American Conference of Governmental Industrial Hygienists
ADI	Acceptable Daily Intake
ADME	Absorption, Distribution, Metabolism, and Excretion
AFID	alkali flame ionization detector
AFOSH	Air Force Office of Safety and Health
AML	acute myeloid leukemia
AOAC	Association of Official Analytical Chemists
atm	atmosphere
ATSDR	Agency for Toxic Substances and Disease Registry
AWQC	Ambient Water Quality Criteria
BAT	Best Available Technology
BCF	bioconcentration factor
BEI	Biological Exposure Index
BSC	Board of Scientific Counselors
C	Centigrade
CAA	Clean Air Act
CAG	Cancer Assessment Group of the U.S. Environmental Protection Agency
CAS	Chemical Abstract Services
CDC	Centers for Disease Control and Prevention
CEL	Cancer Effect Level
CELDS	Computer-Environmental Legislative Data System
CERCLA	Comprehensive Environmental Response, Compensation, and Liability Act
CFR	Code of Federal Regulations
Ci	curie
CL	ceiling limit value
CLP	Contract Laboratory Program
cm	centimeter
CML	chronic myeloid leukemia
CNS	central nervous system
CPSC	Consumer Products Safety Commission
CWA	Clean Water Act
d	day
Derm	dermal
DHEW	Department of Health, Education, and Welfare
DHHS	Department of Health and Human Services
DNA	deoxyribonucleic acid
DOD	Department of Defense
DOE	Department of Energy
DOL	Department of Labor
DOT	Department of Transportation
DOT/UN/ NA/IMCO	Department of Transportation/United Nations/ North America/International Maritime Dangerous Goods Code
DWEL	Drinking Water Exposure Level
ECD	electron capture detection
ECG/EKG	electrocardiogram

APPENDIX C

EEG	electroencephalogram
EEGL	Emergency Exposure Guidance Level
EPA	Environmental Protection Agency
F	Fahrenheit
F ₁	first-filial generation
FAO	Food and Agricultural Organization of the United Nations
FDA	Food and Drug Administration
FEMA	Federal Emergency Management Agency
FIFRA	Federal Insecticide, Fungicide, and Rodenticide Act
FPD	flame photometric detection
fpm	feet per minute
ft	foot
FR	<i>Federal Register</i>
g	gram
GC	gas chromatography
Gd	gestational day
gen	generation
GLC	gas liquid chromatography
GPC	gel permeation chromatography
HPLC	high-performance liquid chromatography
hr	hour
HRGC	high resolution gas chromatography
HSDB	Hazardous Substance Data Bank
IDLH	Immediately Dangerous to Life and Health
IARC	International Agency for Research on Cancer
ILO	International Labor Organization
in	inch
IRIS	Integrated Risk Information System
K _d	adsorption ratio
kg	kilogram
kgg	metric ton
K _{oc}	organic carbon partition coefficient
K _{ow}	octanol-water partition coefficient
L	liter
LC	liquid chromatography
LC _{Lo}	lethal concentration, low
LC ₅₀	lethal concentration, 50% kill
LD _{Lo}	lethal dose, low
LD ₅₀	lethal dose, 50% kill
LT ₅₀	lethal time, 50% kill
LOAEL	lowest-observed-adverse-effect level
LSE	Levels of Significant Exposure
m	meter
MA	<i>trans,trans</i> -muconic acid
MAL	Maximum Allowable Level
mCi	millicurie
MCL	Maximum Contaminant Level
MCLG	Maximum Contaminant Level Goal
mg	milligram
min	minute
mL	milliliter

APPENDIX C

mm	millimeter
mm Hg	millimeters of mercury
mmol	millimole
mo	month
mppcf	millions of particles per cubic foot
MRL	Minimal Risk Level
MS	mass spectrometry
NAAQS	National Ambient Air Quality Standard
NAS	National Academy of Science
NATICH	National Air Toxics Information Clearinghouse
NATO	North Atlantic Treaty Organization
NCE	normochromatic erythrocytes
NCI	National Cancer Institute
NIEHS	National Institute of Environmental Health Sciences
NIOSH	National Institute for Occupational Safety and Health
NIOSHTIC	NIOSH's Computerized Information Retrieval System
NFPA	National Fire Protection Association
ng	nanogram
NLM	National Library of Medicine
nm	nanometer
NHANES	National Health and Nutrition Examination Survey
nmol	nanomole
NOAEL	no-observed-adverse-effect level
NOES	National Occupational Exposure Survey
NOHS	National Occupational Hazard Survey
NPD	nitrogen phosphorus detection
NPDES	National Pollutant Discharge Elimination System
NPL	National Priorities List
NR	not reported
NRC	National Research Council
NS	not specified
NSPS	New Source Performance Standards
NTIS	National Technical Information Service
NTP	National Toxicology Program
ODW	Office of Drinking Water, EPA
OERR	Office of Emergency and Remedial Response, EPA
OHM/TADS	Oil and Hazardous Materials/Technical Assistance Data System
OPP	Office of Pesticide Programs, EPA
OPPTS	Office of Prevention, Pesticides and Toxic Substances, EPA
OPPT	Office of Pollution Prevention and Toxics, EPA
OSHA	Occupational Safety and Health Administration
OSW	Office of Solid Waste, EPA
OTS	Office of Toxic Substances
OW	Office of Water
OWRS	Office of Water Regulations and Standards, EPA
PAH	Polycyclic Aromatic Hydrocarbon
PBPD	Physiologically Based Pharmacodynamic
PBPK	Physiologically Based Pharmacokinetic
PCE	polychromatic erythrocytes
PEL	permissible exposure limit
PID	photo ionization detector

APPENDIX C

pg	picogram
pmol	picomole
PHS	Public Health Service
PMR	proportionate mortality ratio
ppb	parts per billion
ppm	parts per million
ppt	parts per trillion
PSNS	Pretreatment Standards for New Sources
REL	recommended exposure level/limit
RfC	Reference Concentration
RfD	Reference Dose
RNA	ribonucleic acid
RTECS	Registry of Toxic Effects of Chemical Substances
RQ	Reportable Quantity
SARA	Superfund Amendments and Reauthorization Act
SCE	sister chromatid exchange
sec	second
SIC	Standard Industrial Classification
SIM	selected ion monitoring
SMCL	Secondary Maximum Contaminant Level
SMR	standard mortality ratio
SNARL	Suggested No Adverse Response Level
SPEGL	Short-Term Public Emergency Guidance Level
STEL	short term exposure limit
STORET	Storage and Retrieval
TD ₅₀	toxic dose, 50% specific toxic effect
TLV	threshold limit value
TOC	Total Organic Compound
TPQ	Threshold Planning Quantity
TRI	Toxics Release Inventory
TSCA	Toxic Substances Control Act
TRI	Toxics Release Inventory
TWA	time-weighted average
U.S.	United States
UF	uncertainty factor
VOC	Volatile Organic Compound
yr	year
WHO	World Health Organization
wk	week
>	greater than
≥	greater than or equal to
=	equal to
<	less than
≤	less than or equal to
%	percent
α	alpha
β	beta
γ	gamma
δ	delta
μm	micrometer

APPENDIX C

μg	microgram
q_1^*	cancer slope factor
-	negative
+	positive
(+)	weakly positive result
(-)	weakly negative result