

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All causes												
All races, both sexes												
2004.....	2,397,615	27,936	4,785	2,888	3,946	13,706	19,715	18,771	22,097	31,944	53,418	77,927
2003.....	2,448,288	28,025	4,965	2,898	4,056	13,595	19,973	18,250	23,050	34,055	55,406	78,173
2002.....	2,443,387	28,034	4,858	3,018	4,132	13,812	19,234	17,959	23,396	35,347	55,793	76,065
2001.....	2,416,425	27,568	5,107	3,093	4,002	13,555	18,697	18,211	23,472	36,895	54,779	73,684
2000.....	2,403,351	28,035	4,979	3,253	4,160	13,563	17,744	17,681	22,770	36,140	53,658	70,832
1999.....	2,391,399	27,937	5,249	3,474	4,121	13,778	16,878	17,703	23,363	36,804	52,452	68,120
All races, male												
2004.....	1,181,668	15,718	2,649	1,645	2,354	9,678	14,909	13,554	14,805	20,485	33,192	48,514
2003.....	1,201,964	15,902	2,826	1,640	2,510	9,706	14,964	13,043	15,559	21,741	34,694	49,139
2002.....	1,199,264	15,717	2,806	1,702	2,496	9,844	14,572	12,954	15,782	22,626	34,967	47,701
2001.....	1,183,421	15,477	2,899	1,727	2,441	9,766	14,197	13,006	15,751	23,727	34,437	46,567
2000.....	1,177,578	15,718	2,824	1,850	2,551	9,697	13,374	12,619	15,271	23,252	34,045	45,121
1999.....	1,175,460	15,646	2,975	1,964	2,528	9,780	12,634	12,542	15,734	23,681	33,437	43,303
All races, female												
2004.....	1,215,947	12,218	2,136	1,243	1,592	4,028	4,806	5,217	7,292	11,459	20,226	29,413
2003.....	1,246,324	12,123	2,139	1,258	1,546	3,889	5,009	5,207	7,491	12,314	20,712	29,034
2002.....	1,244,123	12,317	2,052	1,316	1,636	3,968	4,662	5,005	7,614	12,721	20,826	28,364
2001.....	1,233,004	12,091	2,208	1,366	1,561	3,789	4,500	5,205	7,721	13,168	20,342	27,117
2000.....	1,225,773	12,317	2,155	1,403	1,609	3,866	4,370	5,062	7,499	12,888	19,613	25,711
1999.....	1,215,939	12,291	2,274	1,510	1,593	3,998	4,244	5,161	7,629	13,123	19,015	24,817
White, both sexes												
2004.....	2,056,643	18,231	3,363	2,043	2,857	10,450	14,563	13,421	15,854	23,686	40,354	59,181
2003.....	2,103,714	18,440	3,509	2,111	2,920	10,392	14,584	12,991	16,698	25,012	41,736	59,135
2002.....	2,102,589	18,369	3,406	2,168	2,970	10,613	14,028	12,703	16,868	25,912	42,181	57,690
2001.....	2,079,691	17,955	3,670	2,250	2,920	10,235	13,513	12,988	17,006	27,238	40,899	55,634
2000.....	2,071,287	18,144	3,494	2,359	3,091	10,273	12,745	12,427	16,292	26,633	39,863	53,131
1999.....	2,061,348	18,067	3,690	2,450	3,043	10,355	11,997	12,569	16,831	27,024	38,583	50,895

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All causes												
White, male												
2004.....	1,007,266	10,265	1,878	1,178	1,727	7,267	11,039	9,795	10,789	15,483	25,732	37,461
2003.....	1,025,650	10,500	1,981	1,205	1,803	7,308	10,911	9,342	11,472	16,344	26,739	37,898
2002.....	1,025,196	10,433	1,958	1,219	1,786	7,474	10,608	9,219	11,579	16,840	27,115	36,894
2001.....	1,011,218	10,099	2,094	1,232	1,782	7,264	10,265	9,381	11,597	17,880	26,363	35,774
2000.....	1,007,191	10,177	2,004	1,348	1,907	7,242	9,626	8,943	11,197	17,529	25,849	34,599
1999.....	1,005,335	10,197	2,084	1,376	1,852	7,222	8,981	9,015	11,534	17,792	25,122	32,778
White, female												
2004.....	1,049,377	7,966	1,485	865	1,130	3,183	3,524	3,626	5,065	8,203	14,622	21,720
2003.....	1,078,064	7,940	1,528	906	1,117	3,084	3,673	3,649	5,226	8,668	14,997	21,237
2002.....	1,077,393	7,936	1,448	949	1,184	3,139	3,420	3,484	5,289	9,072	15,066	20,796
2001.....	1,068,473	7,856	1,576	1,018	1,138	2,971	3,248	3,607	5,409	9,358	14,536	19,860
2000.....	1,064,096	7,967	1,490	1,011	1,184	3,031	3,119	3,484	5,095	9,104	14,014	18,532
1999.....	1,056,013	7,870	1,606	1,074	1,191	3,133	3,016	3,554	5,297	9,232	13,461	18,117
Black, both sexes												
2004.....	287,315	8,494	1,164	685	899	2,680	4,369	4,616	5,309	7,098	11,376	16,423
2003.....	291,300	8,402	1,199	637	914	2,572	4,573	4,524	5,418	7,823	11,961	16,680
2002.....	290,051	8,524	1,196	701	965	2,602	4,443	4,550	5,621	8,227	12,008	16,184
2001.....	287,709	8,498	1,181	700	887	2,710	4,437	4,467	5,561	8,385	12,302	15,867
2000.....	285,826	8,771	1,248	756	887	2,717	4,332	4,541	5,698	8,352	12,205	15,735
1999.....	285,064	8,822	1,311	880	909	2,849	4,217	4,444	5,671	8,663	12,378	15,304
Black, male												
2004.....	145,970	4,769	641	372	516	2,019	3,317	3,264	3,420	4,262	6,438	9,640
2003.....	148,022	4,740	699	355	567	1,974	3,474	3,219	3,474	4,620	6,924	9,824
2002.....	146,835	4,652	702	402	596	1,963	3,401	3,229	3,619	4,996	6,853	9,501
2001.....	145,908	4,764	672	409	551	2,080	3,399	3,117	3,556	5,022	7,113	9,495
2000.....	145,184	4,901	692	431	543	2,045	3,273	3,163	3,587	5,015	7,236	9,350
1999.....	145,703	4,897	746	519	577	2,157	3,191	3,073	3,650	5,171	7,396	9,392

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All causes												
Black, female												
2004.....	141,345	3,725	523	313	383	661	1,052	1,352	1,889	2,836	4,938	6,783
2003.....	143,278	3,662	500	282	347	598	1,099	1,305	1,944	3,203	5,037	6,856
2002.....	143,216	3,872	494	299	369	639	1,042	1,321	2,002	3,231	5,155	6,683
2001.....	141,801	3,734	509	291	336	630	1,038	1,350	2,005	3,363	5,189	6,372
2000.....	140,642	3,870	556	325	344	672	1,059	1,378	2,111	3,337	4,969	6,385
1999.....	139,361	3,925	565	361	332	692	1,026	1,371	2,021	3,492	4,982	5,912
American Indian 1/, both sexes												
2004.....	13,124	389	97	53	74	260	333	314	381	516	678	833
2003.....	13,147	335	95	55	82	286	335	288	355	539	705	842
2002.....	12,415	343	90	48	78	267	316	290	362	512	657	751
2001.....	11,977	373	103	44	85	273	284	265	331	531	601	747
2000.....	11,363	323	91	50	63	251	268	272	324	472	609	611
1999.....	11,312	343	82	48	57	245	295	281	364	453	574	628
American Indian 1/, male												
2004.....	7,134	237	50	28	42	171	238	222	239	338	409	508
2003.....	7,106	195	55	30	56	195	255	193	251	351	433	516
2002.....	6,750	191	49	21	44	178	238	218	239	335	414	455
2001.....	6,466	211	52	29	46	186	214	194	223	355	367	437
2000.....	6,185	193	49	28	33	178	191	192	218	284	391	389
1999.....	6,092	180	48	21	34	179	217	180	238	293	352	415
American Indian 1/, female												
2004.....	5,990	152	47	25	32	89	95	92	142	178	269	325
2003.....	6,041	140	40	25	26	91	80	95	104	188	272	326
2002.....	5,665	152	41	27	34	89	78	72	123	177	243	296
2001.....	5,511	162	51	15	39	87	70	71	108	176	234	310
2000.....	5,178	130	42	22	30	73	77	80	106	188	218	222
1999.....	5,220	163	34	27	23	66	78	101	126	160	222	213

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

All causes

Asian or Pacific Islander 2/, both sexes

2004.....	40,533	822	161	107	116	316	450	420	553	644	1,010	1,490
2003.....	40,127	848	162	95	140	345	481	447	579	681	1,004	1,516
2002.....	38,332	798	166	101	119	330	447	416	545	696	947	1,440
2001.....	37,048	742	153	99	110	337	463	491	574	741	977	1,436
2000.....	34,875	797	146	88	119	322	399	441	456	683	981	1,355
1999.....	33,675	705	166	96	112	329	369	409	497	664	917	1,293

Asian or Pacific Islander 2/, male

2004.....	21,298	447	80	67	69	221	315	273	357	402	613	905
2003.....	21,186	467	91	50	84	229	324	289	362	426	598	901
2002.....	20,483	441	97	60	70	229	325	288	345	455	585	851
2001.....	19,829	403	81	57	62	236	319	314	375	470	594	861
2000.....	19,018	447	79	43	68	232	284	321	269	424	569	783
1999.....	18,330	372	97	48	65	222	245	274	312	425	567	718

Asian or Pacific Islander 2/, female

2004.....	19,235	375	81	40	47	95	135	147	196	242	397	585
2003.....	18,941	381	71	45	56	116	157	158	217	255	406	615
2002.....	17,849	357	69	41	49	101	122	128	200	241	362	589
2001.....	17,219	339	72	42	48	101	144	177	199	271	383	575
2000.....	15,857	350	67	45	51	90	115	120	187	259	412	572
1999.....	15,345	333	69	48	47	107	124	135	185	239	350	575

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
I. Certain infectious and parasitic diseases (A00-B99)												
All races, both sexes												
2004.....	64,618	480	187	77	81	142	332	653	1,394	2,637	4,187	5,034
2003.....	64,661	494	197	90	96	167	320	659	1,481	3,076	4,351	5,217
2002.....	63,981	582	170	94	135	167	300	746	1,710	3,343	4,498	5,254
2001.....	60,336	563	225	98	110	197	334	775	1,870	3,451	4,483	4,817
2000.....	59,007	510	222	99	98	160	303	850	2,127	3,632	4,369	4,719
1999.....	59,984	562	236	143	114	139	341	999	2,386	3,772	4,635	4,510
All races, male												
2004.....	34,303	288	94	41	40	73	196	356	869	1,774	2,856	3,510
2003.....	34,292	257	115	38	48	92	167	379	969	2,116	3,015	3,683
2002.....	34,352	306	99	52	69	78	165	422	1,112	2,369	3,135	3,739
2001.....	32,553	304	127	53	61	106	171	453	1,255	2,461	3,175	3,467
2000.....	31,799	275	120	55	48	96	171	483	1,423	2,611	3,090	3,447
1999.....	32,480	331	143	72	65	73	176	624	1,651	2,706	3,371	3,274
All races, female												
2004.....	30,315	192	93	36	41	69	136	297	525	863	1,331	1,524
2003.....	30,369	237	82	52	48	75	153	280	512	960	1,336	1,534
2002.....	29,629	276	71	42	66	89	135	324	598	974	1,363	1,515
2001.....	27,783	259	98	45	49	91	163	322	615	990	1,308	1,350
2000.....	27,208	235	102	44	50	64	132	367	704	1,021	1,279	1,272
1999.....	27,504	231	93	71	49	66	165	375	735	1,066	1,264	1,236
White, both sexes												
2004.....	47,439	315	135	58	47	84	152	270	597	1,384	2,261	2,982
2003.....	47,191	280	124	56	58	107	164	277	700	1,526	2,410	3,155
2002.....	46,293	389	117	69	87	111	149	342	752	1,696	2,402	3,142
2001.....	43,252	334	159	74	68	111	150	315	838	1,763	2,400	2,816
2000.....	41,965	315	144	67	68	101	144	360	948	1,904	2,358	2,715
1999.....	42,522	354	160	91	68	93	161	449	1,174	2,013	2,425	2,513

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

I. Certain infectious and parasitic diseases (A00-B99)

White, male

2004.....	24,631	189	68	31	27	43	100	158	425	992	1,653	2,146
2003.....	24,399	137	73	24	29	65	88	176	498	1,145	1,768	2,324
2002.....	24,210	204	72	41	51	51	89	226	537	1,298	1,784	2,323
2001.....	22,722	182	89	39	39	61	85	207	613	1,360	1,830	2,090
2000.....	21,933	171	75	38	31	62	94	225	710	1,478	1,749	2,049
1999.....	22,301	218	97	48	37	50	101	302	892	1,571	1,860	1,844

White, female

2004.....	22,808	126	67	27	20	41	52	112	172	392	608	836
2003.....	22,792	143	51	32	29	42	76	101	202	381	642	831
2002.....	22,083	185	45	28	36	60	60	116	215	398	618	819
2001.....	20,530	152	70	35	29	50	65	108	225	403	570	726
2000.....	20,032	144	69	29	37	39	50	135	238	426	609	666
1999.....	20,221	136	63	43	31	43	60	147	282	442	565	669

Black, both sexes

2004.....	15,593	144	38	14	29	52	168	364	755	1,186	1,835	1,928
2003.....	15,919	186	58	29	31	53	141	362	743	1,477	1,866	1,948
2002.....	16,204	166	42	16	41	48	139	390	918	1,592	1,998	1,998
2001.....	15,745	203	55	21	37	74	173	440	991	1,623	2,001	1,904
2000.....	15,792	184	61	29	23	49	151	476	1,147	1,672	1,931	1,908
1999.....	16,090	188	63	45	37	40	165	532	1,174	1,696	2,121	1,902

Black, male

2004.....	8,797	82	19	7	12	28	88	186	414	730	1,135	1,286
2003.....	9,088	107	33	14	15	22	71	191	444	918	1,194	1,292
2002.....	9,293	87	21	8	16	22	68	185	542	1,034	1,277	1,336
2001.....	9,074	111	33	11	19	39	80	235	609	1,054	1,298	1,307
2000.....	9,167	98	35	15	11	27	73	248	688	1,095	1,283	1,332
1999.....	9,396	102	40	23	23	21	68	307	735	1,086	1,450	1,361

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

I. Certain infectious and parasitic diseases (A00-B99)

Black, female

2004.....	6,796	62	19	7	17	24	80	178	341	456	700	642
2003.....	6,831	79	25	15	16	31	70	171	299	559	672	656
2002.....	6,911	79	21	8	25	26	71	205	376	558	721	662
2001.....	6,671	92	22	10	18	35	93	205	382	569	703	597
2000.....	6,625	86	26	14	12	22	78	228	459	577	648	576
1999.....	6,694	86	23	22	14	19	97	225	439	610	671	541

American Indian 1/, both sexes

2004.....	444	13	6	2	3	-	2	6	17	29	48	62
2003.....	461	10	8	3	4	3	6	5	16	36	39	49
2002.....	407	9	6	1	3	-	3	4	16	27	45	47
2001.....	355	9	7	2	3	2	4	5	15	35	36	47
2000.....	343	4	6	1	1	2	1	3	13	27	32	39
1999.....	352	7	2	3	2	1	5	5	17	30	32	32

American Indian 1/, male

2004.....	244	12	4	-	1	-	1	4	10	23	33	35
2003.....	215	5	4	-	3	3	3	2	14	22	24	19
2002.....	225	5	4	1	2	-	1	3	14	16	32	30
2001.....	200	4	4	2	2	2	2	2	12	22	18	31
2000.....	178	2	2	1	-	1	-	1	8	16	22	28
1999.....	191	4	-	1	-	-	3	4	12	22	19	23

American Indian 1/, female

2004.....	200	1	2	2	2	-	1	2	7	6	15	27
2003.....	246	5	4	3	1	-	3	3	2	14	15	30
2002.....	182	4	2	-	1	-	2	1	2	11	13	17
2001.....	155	5	3	-	1	-	2	3	3	13	18	16
2000.....	165	2	4	-	1	1	2	5	5	11	10	11
1999.....	161	3	2	2	2	1	2	1	5	8	13	9

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

I. Certain infectious and parasitic diseases (A00-B99)

Asian or Pacific Islander 2/, both sexes

2004.....	1,142	8	8	3	2	6	10	13	25	38	43	62
2003.....	1,090	18	7	2	3	4	9	15	22	37	36	65
2002.....	1,077	18	5	8	4	8	9	10	24	28	53	67
2001.....	984	17	4	1	2	10	7	15	26	30	46	50
2000.....	907	7	11	2	6	8	7	11	19	29	48	57
1999.....	1,020	13	11	4	7	5	10	13	21	33	57	63

Asian or Pacific Islander 2/, male

2004.....	631	5	3	3	-	2	7	8	20	29	35	43
2003.....	590	8	5	-	1	2	5	10	13	31	29	48
2002.....	624	10	2	2	-	5	7	8	19	21	42	50
2001.....	557	7	1	1	1	4	4	9	21	25	29	39
2000.....	521	4	8	1	6	6	4	9	17	22	36	38
1999.....	592	7	6	-	5	2	4	11	12	27	42	46

Asian or Pacific Islander 2/, female

2004.....	511	3	5	-	2	4	3	5	5	9	8	19
2003.....	500	10	2	2	2	2	4	5	9	6	7	17
2002.....	453	8	3	6	4	3	2	2	5	7	11	17
2001.....	427	10	3	-	1	6	3	6	5	5	17	11
2000.....	386	3	3	1	-	2	3	2	2	7	12	19
1999.....	428	6	5	4	2	3	6	2	9	6	15	17

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Intestinal infectious diseases (A00-A09)

All races, both sexes

2004.....	4,391	14	15	1	1	1	2	9	4	11	21	39
2003.....	3,149	14	4	5	3	1	3	1	8	6	15	23
2002.....	2,501	16	9	1	2	5	2	3	1	5	17	30
2001.....	1,591	14	19	2	3	-	-	1	5	5	8	13
2000.....	1,380	17	12	-	-	1	1	2	3	7	13	14
1999.....	1,169	21	9	3	1	1	6	3	3	2	8	11

All races, male

2004.....	1,647	9	5	-	1	-	1	4	1	8	12	20
2003.....	1,143	8	4	1	1	1	1	-	4	4	6	13
2002.....	967	4	5	1	-	4	-	1	-	2	8	15
2001.....	603	9	12	2	1	-	-	-	3	5	4	8
2000.....	554	13	6	-	-	1	-	2	3	4	7	10
1999.....	458	15	5	3	1	1	4	2	3	1	5	5

All races, female

2004.....	2,744	5	10	1	-	1	1	5	3	3	9	19
2003.....	2,006	6	-	4	2	-	2	1	4	2	9	10
2002.....	1,534	12	4	-	2	1	2	2	1	3	9	15
2001.....	988	5	7	-	2	-	-	1	2	-	4	5
2000.....	826	4	6	-	-	-	1	-	-	3	6	4
1999.....	711	6	4	-	-	-	2	1	-	1	3	6

White, both sexes

2004.....	4,099	11	12	1	1	1	2	7	3	9	13	27
2003.....	2,922	7	3	5	2	1	2	-	6	5	10	20
2002.....	2,319	9	6	1	2	5	2	2	-	4	15	26
2001.....	1,461	9	16	2	2	-	-	-	4	3	7	9
2000.....	1,270	11	9	-	-	-	1	1	3	3	8	11
1999.....	1,080	16	8	1	1	1	4	3	3	2	5	8

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Intestinal infectious diseases (A00-A09)

White, male

2004.....	1,519	7	3	-	1	-	1	3	1	6	7	13
2003.....	1,063	3	3	1	1	1	1	-	3	3	5	11
2002.....	891	2	4	1	-	4	-	1	-	1	7	14
2001.....	545	6	10	2	-	-	-	-	3	3	3	5
2000.....	504	8	5	-	-	-	-	1	3	2	5	8
1999.....	421	12	4	1	1	1	3	2	3	1	4	3

White, female

2004.....	2,580	4	9	1	-	1	1	4	2	3	6	14
2003.....	1,859	4	-	4	1	-	1	-	3	2	5	9
2002.....	1,428	7	2	-	2	1	2	1	-	3	8	12
2001.....	916	3	6	-	2	-	-	-	1	-	4	4
2000.....	766	3	4	-	-	-	1	-	-	1	3	3
1999.....	659	4	4	-	-	-	1	1	-	1	1	5

Black, both sexes

2004.....	241	1	2	-	-	-	-	2	1	1	6	10
2003.....	195	7	1	-	1	-	1	1	1	1	3	3
2002.....	159	6	1	-	-	-	-	1	1	1	2	3
2001.....	108	4	3	-	1	-	-	1	1	1	1	4
2000.....	88	5	2	-	-	1	-	1	-	3	4	3
1999.....	79	5	-	2	-	-	2	-	-	-	3	2

Black, male

2004.....	99	1	1	-	-	-	-	1	-	1	3	6
2003.....	70	5	1	-	-	-	-	-	1	1	-	2
2002.....	67	1	1	-	-	-	-	-	-	1	1	1
2001.....	45	2	2	-	1	-	-	-	-	1	1	3
2000.....	41	4	1	-	-	1	-	1	-	1	2	2
1999.....	29	3	-	2	-	-	1	-	-	-	1	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other intestinal infectious diseases (A01-A08)												
All races, both sexes												
2004.....	4,280	14	14	-	1	1	2	9	3	11	21	39
2003.....	3,051	13	3	5	3	1	3	1	8	5	15	23
2002.....	2,408	13	8	1	1	5	2	3	1	5	17	26
2001.....	1,535	12	18	2	3	-	-	1	5	5	7	13
2000.....	1,314	16	12	-	-	1	1	1	3	7	12	13
1999.....	1,072	21	8	3	1	1	5	3	3	2	5	10
All races, male												
2004.....	1,612	9	4	-	1	-	1	4	1	8	12	20
2003.....	1,106	7	3	1	1	1	1	-	4	3	6	13
2002.....	934	3	5	1	-	4	-	1	-	2	8	13
2001.....	582	8	12	2	1	-	-	-	3	5	3	8
2000.....	525	13	6	-	-	1	-	1	3	4	6	9
1999.....	428	15	5	3	1	1	4	2	3	1	4	5
All races, female												
2004.....	2,668	5	10	-	-	1	1	5	2	3	9	19
2003.....	1,945	6	-	4	2	-	2	1	4	2	9	10
2002.....	1,474	10	3	-	1	1	2	2	1	3	9	13
2001.....	953	4	6	-	2	-	-	1	2	-	4	5
2000.....	789	3	6	-	-	-	1	-	-	3	6	4
1999.....	644	6	3	-	-	-	1	1	-	1	1	5
White, both sexes												
2004.....	4,001	11	12	-	1	1	2	7	3	9	13	27
2003.....	2,832	7	2	5	2	1	2	-	6	4	10	20
2002.....	2,234	8	5	1	1	5	2	2	-	4	15	22
2001.....	1,409	9	15	2	2	-	-	-	4	3	6	9
2000.....	1,209	11	9	-	-	-	1	-	3	3	7	11
1999.....	993	16	7	1	1	1	4	3	3	2	4	7

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other intestinal infectious diseases (A01-A08)

White, male

2004.....	1,491	7	3	-	1	-	1	3	1	6	7	13
2003.....	1,029	3	2	1	1	1	1	-	3	2	5	11
2002.....	863	2	4	1	-	4	-	1	-	1	7	12
2001.....	525	6	10	2	-	-	-	-	3	3	2	5
2000.....	477	8	5	-	-	-	-	-	3	2	4	8
1999.....	393	12	4	1	1	1	3	2	3	1	3	3

White, female

2004.....	2,510	4	9	-	-	1	1	4	2	3	6	14
2003.....	1,803	4	-	4	1	-	1	-	3	2	5	9
2002.....	1,371	6	1	-	1	1	2	1	-	3	8	10
2001.....	884	3	5	-	2	-	-	-	1	-	4	4
2000.....	732	3	4	-	-	-	1	-	-	1	3	3
1999.....	600	4	3	-	-	-	1	1	-	1	1	4

Black, both sexes

2004.....	230	1	1	-	-	-	-	2	-	1	6	10
2003.....	187	6	1	-	1	-	1	1	1	1	3	3
2002.....	152	5	1	-	-	-	-	1	1	1	2	3
2001.....	104	2	3	-	1	-	-	1	1	1	1	4
2000.....	83	4	2	-	-	1	-	1	-	3	4	2
1999.....	69	5	-	2	-	-	1	-	-	-	1	2

Black, male

2004.....	93	1	-	-	-	-	-	1	-	1	3	6
2003.....	67	4	1	-	-	-	-	-	1	1	-	2
2002.....	63	1	1	-	-	-	-	-	-	1	1	1
2001.....	44	1	2	-	1	-	-	-	-	1	1	3
2000.....	39	4	1	-	-	1	-	1	-	1	2	1
1999.....	27	3	-	2	-	-	1	-	-	-	1	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Paratyphoid fevers and other salmonella infections (A01.1-A01.4,A02)												
All races, both sexes												
2004.....	30	1	-	-	-	-	-	-	1	1	-	2
2003.....	43	3	-	1	-	-	-	1	-	-	3	1
2002.....	21	3	-	-	-	-	-	-	-	-	2	1
2001.....	40	2	-	-	-	-	-	-	2	1	-	1
2000.....	28	3	2	-	-	1	-	-	1	-	-	1
1999.....	38	3	-	-	-	-	2	-	-	-	2	-
All races, male												
2004.....	22	1	-	-	-	-	-	-	-	1	-	2
2003.....	23	1	-	-	-	-	-	-	-	-	2	1
2002.....	10	1	-	-	-	-	-	-	-	-	-	-
2001.....	21	-	-	-	-	-	-	-	1	1	-	1
2000.....	15	2	-	-	-	1	-	-	1	-	-	1
1999.....	21	2	-	-	-	-	2	-	-	-	1	-
All races, female												
2004.....	8	-	-	-	-	-	-	-	1	-	-	-
2003.....	20	2	-	1	-	-	-	1	-	-	1	-
2002.....	11	2	-	-	-	-	-	-	-	-	2	1
2001.....	19	2	-	-	-	-	-	-	1	-	-	-
2000.....	13	1	2	-	-	-	-	-	-	-	-	-
1999.....	17	1	-	-	-	-	-	-	-	-	1	-
White, both sexes												
2004.....	20	1	-	-	-	-	-	-	1	-	-	1
2003.....	31	1	-	1	-	-	-	-	-	-	1	1
2002.....	18	2	-	-	-	-	-	-	-	-	2	1
2001.....	32	1	-	-	-	-	-	-	2	-	-	1
2000.....	25	2	2	-	-	-	-	-	1	-	-	1
1999.....	32	3	-	-	-	-	1	-	-	-	2	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Paratyphoid fevers and other salmonella infections (A01.1-A01.4,A02)												
White, male												
2004.....	14	1	-	-	-	-	-	-	-	-	-	1
2003.....	17	-	-	-	-	-	-	-	-	-	1	1
2002.....	9	1	-	-	-	-	-	-	-	-	-	-
2001.....	18	-	-	-	-	-	-	-	1	-	-	1
2000.....	13	1	-	-	-	-	-	-	1	-	-	1
1999.....	16	2	-	-	-	-	1	-	-	-	1	-
White, female												
2004.....	6	-	-	-	-	-	-	-	1	-	-	-
2003.....	14	1	-	1	-	-	-	-	-	-	-	-
2002.....	9	1	-	-	-	-	-	-	-	-	2	1
2001.....	14	1	-	-	-	-	-	-	1	-	-	-
2000.....	12	1	2	-	-	-	-	-	-	-	-	-
1999.....	16	1	-	-	-	-	-	-	-	-	1	-
Black, both sexes												
2004.....	4	-	-	-	-	-	-	-	-	-	-	1
2003.....	8	2	-	-	-	-	-	1	-	-	1	-
2002.....	3	1	-	-	-	-	-	-	-	-	-	-
2001.....	5	1	-	-	-	-	-	-	-	-	-	-
2000.....	2	1	-	-	-	1	-	-	-	-	-	-
1999.....	4	-	-	-	-	-	1	-	-	-	-	-
Black, male												
2004.....	3	-	-	-	-	-	-	-	-	-	-	1
2003.....	3	1	-	-	-	-	-	-	-	-	-	-
2002.....	1	-	-	-	-	-	-	-	-	-	-	-
2001.....	1	-	-	-	-	-	-	-	-	-	-	-
2000.....	2	1	-	-	-	1	-	-	-	-	-	-
1999.....	3	-	-	-	-	-	1	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Intestinal infections due to other specified organisms (A04,A07-A08)

All races, both sexes

2004.....	4,246	13	14	-	-	1	1	9	2	10	21	37
2003.....	2,992	9	3	4	3	1	3	-	8	3	10	21
2002.....	2,372	10	7	1	1	4	2	3	1	5	13	21
2001.....	1,481	8	17	1	3	-	-	1	3	4	7	9
2000.....	1,263	12	10	-	-	-	1	1	1	7	11	10
1999.....	1,012	12	7	2	1	1	3	3	3	2	2	10

All races, male

2004.....	1,587	8	4	-	-	-	1	4	1	7	12	18
2003.....	1,072	5	3	1	1	1	1	-	4	2	2	11
2002.....	914	2	5	1	-	3	-	1	-	2	6	10
2001.....	551	6	11	1	1	-	-	-	2	4	3	4
2000.....	496	10	6	-	-	-	-	1	1	4	5	6
1999.....	391	8	4	2	1	1	2	2	3	1	2	5

All races, female

2004.....	2,659	5	10	-	-	1	-	5	1	3	9	19
2003.....	1,920	4	-	3	2	-	2	-	4	1	8	10
2002.....	1,458	8	2	-	1	1	2	2	1	3	7	11
2001.....	930	2	6	-	2	-	-	1	1	-	4	5
2000.....	767	2	4	-	-	-	1	-	-	3	6	4
1999.....	621	4	3	-	-	-	1	1	-	1	-	5

White, both sexes

2004.....	3,977	10	12	-	-	1	1	7	2	9	13	26
2003.....	2,787	5	2	4	2	1	2	-	6	3	7	18
2002.....	2,202	6	5	1	1	4	2	2	-	4	11	17
2001.....	1,365	6	14	1	2	-	-	-	2	3	6	6
2000.....	1,164	8	7	-	-	-	1	-	1	3	7	8
1999.....	940	8	6	-	1	1	3	3	3	2	1	7

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Tuberculosis (A16-A19)

All races, both sexes

2004.....	657	1	1	-	2	1	6	6	10	24	29	33
2003.....	711	-	1	-	2	3	7	8	10	11	24	54
2002.....	784	1	-	-	-	-	9	7	11	18	34	35
2001.....	764	2	1	-	-	6	6	11	14	18	22	37
2000.....	776	1	6	-	1	2	9	7	13	17	25	38
1999.....	930	2	2	1	2	1	10	3	16	25	32	51

All races, male

2004.....	406	-	1	-	1	1	3	2	10	19	21	21
2003.....	427	-	1	-	2	1	2	6	7	8	17	39
2002.....	462	-	-	-	-	-	5	4	7	11	27	20
2001.....	453	-	-	-	-	5	3	7	7	12	15	27
2000.....	468	-	1	-	-	2	5	3	11	11	17	35
1999.....	584	-	2	-	2	1	3	1	10	20	23	42

All races, female

2004.....	251	1	-	-	1	-	3	4	-	5	8	12
2003.....	284	-	-	-	-	2	5	2	3	3	7	15
2002.....	322	1	-	-	-	-	4	3	4	7	7	15
2001.....	311	2	1	-	-	1	3	4	7	6	7	10
2000.....	308	1	5	-	1	-	4	4	2	6	8	3
1999.....	346	2	-	1	-	-	7	2	6	5	9	9

White, both sexes

2004.....	390	1	1	-	1	-	3	5	3	12	17	18
2003.....	437	-	1	-	2	2	4	4	4	4	7	22
2002.....	461	-	-	-	-	-	3	3	3	8	14	13
2001.....	469	1	1	-	-	4	4	7	3	7	15	20
2000.....	439	-	4	-	1	1	5	5	6	8	5	13
1999.....	564	1	-	-	1	1	4	2	3	12	13	25

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Tuberculosis (A16-A19)

White, male

2004.....	237	-	1	-	1	-	2	2	3	8	11	12
2003.....	255	-	1	-	2	1	2	3	2	4	5	18
2002.....	272	-	-	-	-	-	2	2	1	5	12	7
2001.....	271	-	-	-	-	4	2	6	2	5	11	14
2000.....	253	-	-	-	-	1	3	2	5	8	3	12
1999.....	348	-	-	-	1	1	3	1	3	9	10	21

White, female

2004.....	153	1	-	-	-	-	1	3	-	4	6	6
2003.....	182	-	-	-	-	1	2	1	2	-	2	4
2002.....	189	-	-	-	-	-	1	1	2	3	2	6
2001.....	198	1	1	-	-	-	2	1	1	2	4	6
2000.....	186	-	4	-	1	-	2	3	1	-	2	1
1999.....	216	1	-	-	-	-	1	1	-	3	3	4

Black, both sexes

2004.....	189	-	-	-	1	1	2	1	5	11	10	12
2003.....	172	-	-	-	-	1	3	1	5	5	11	24
2002.....	204	-	-	-	-	-	4	4	4	9	15	17
2001.....	203	-	-	-	-	-	1	3	7	9	7	14
2000.....	244	-	2	-	-	-	2	2	5	8	17	21
1999.....	245	1	2	1	-	-	4	1	8	10	15	20

Black, male

2004.....	118	-	-	-	-	1	1	-	5	10	8	6
2003.....	114	-	-	-	-	-	-	1	4	3	8	17
2002.....	127	-	-	-	-	-	2	2	3	5	12	11
2001.....	127	-	-	-	-	-	-	1	3	6	4	11
2000.....	161	-	1	-	-	-	1	1	4	2	12	19
1999.....	159	-	2	-	-	-	-	-	4	8	11	17

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Tuberculosis (A16-A19)												
Black, female												
2004.....	71	-	-	-	1	-	1	1	-	1	2	6
2003.....	58	-	-	-	-	1	3	-	1	2	3	7
2002.....	77	-	-	-	-	-	2	2	1	4	3	6
2001.....	76	-	-	-	-	-	1	2	4	3	3	3
2000.....	83	-	1	-	-	-	1	1	1	6	5	2
1999.....	86	1	-	1	-	-	4	1	4	2	4	3
American Indian 1/, both sexes												
2004.....	13	-	-	-	-	-	-	-	-	-	-	-
2003.....	15	-	-	-	-	-	-	-	-	1	1	4
2002.....	27	-	-	-	-	-	-	-	1	-	2	2
2001.....	13	-	-	-	-	-	-	1	-	-	-	2
2000.....	20	1	-	-	-	-	-	-	-	-	-	2
1999.....	20	-	-	-	-	-	-	-	1	1	-	3
American Indian 1/, male												
2004.....	9	-	-	-	-	-	-	-	-	-	-	-
2003.....	6	-	-	-	-	-	-	-	-	-	1	2
2002.....	14	-	-	-	-	-	-	-	1	-	-	1
2001.....	5	-	-	-	-	-	-	-	-	-	-	1
2000.....	12	-	-	-	-	-	-	-	-	-	-	2
1999.....	11	-	-	-	-	-	-	-	-	1	-	3
American Indian 1/, female												
2004.....	4	-	-	-	-	-	-	-	-	-	-	-
2003.....	9	-	-	-	-	-	-	-	-	1	-	2
2002.....	13	-	-	-	-	-	-	-	-	-	2	1
2001.....	8	-	-	-	-	-	-	1	-	-	-	1
2000.....	8	1	-	-	-	-	-	-	-	-	-	-
1999.....	9	-	-	-	-	-	-	-	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Tuberculosis (A16-A19)												
Asian or Pacific Islander 2/, both sexes												
2004.....	65	-	-	-	-	-	1	-	2	1	2	3
2003.....	87	-	-	-	-	-	-	3	1	1	5	4
2002.....	92	1	-	-	-	-	2	-	3	1	3	3
2001.....	79	1	-	-	-	2	1	-	4	2	-	1
2000.....	73	-	-	-	-	1	2	-	2	1	3	2
1999.....	101	-	-	-	1	-	2	-	4	2	4	3
Asian or Pacific Islander 2/, male												
2004.....	42	-	-	-	-	-	-	-	2	1	2	3
2003.....	52	-	-	-	-	-	-	2	1	1	3	2
2002.....	49	-	-	-	-	-	1	-	2	1	3	1
2001.....	50	-	-	-	-	1	1	-	2	1	-	1
2000.....	42	-	-	-	-	1	1	-	2	1	2	2
1999.....	66	-	-	-	1	-	-	-	3	2	2	1
Asian or Pacific Islander 2/, female												
2004.....	23	-	-	-	-	-	1	-	-	-	-	-
2003.....	35	-	-	-	-	-	-	1	-	-	2	2
2002.....	43	1	-	-	-	-	1	-	1	-	-	2
2001.....	29	1	-	-	-	1	-	-	2	1	-	-
2000.....	31	-	-	-	-	-	1	-	-	-	1	-
1999.....	35	-	-	-	-	-	2	-	1	-	2	2

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Respiratory tuberculosis (A16)

All races, both sexes

2004.....	504	1	1	-	1	1	3	4	7	16	20	24
2003.....	562	-	1	-	2	2	7	2	4	5	21	44
2002.....	626	-	-	-	-	-	6	6	8	13	27	28
2001.....	600	2	1	-	-	3	4	8	9	14	14	27
2000.....	615	-	3	-	-	2	5	5	6	15	20	30
1999.....	764	1	1	-	1	-	8	2	12	19	26	38

All races, male

2004.....	316	-	1	-	-	1	1	1	7	12	16	14
2003.....	344	-	1	-	2	1	2	2	4	3	14	32
2002.....	372	-	-	-	-	-	4	4	5	9	22	16
2001.....	371	-	-	-	-	3	2	5	5	9	9	20
2000.....	393	-	1	-	-	2	2	3	5	11	15	29
1999.....	486	-	1	-	1	-	2	-	8	15	18	32

All races, female

2004.....	188	1	-	-	1	-	2	3	-	4	4	10
2003.....	218	-	-	-	-	1	5	-	-	2	7	12
2002.....	254	-	-	-	-	-	2	2	3	4	5	12
2001.....	229	2	1	-	-	-	2	3	4	5	5	7
2000.....	222	-	2	-	-	-	3	2	1	4	5	1
1999.....	278	1	-	-	-	-	6	2	4	4	8	6

White, both sexes

2004.....	300	1	1	-	-	-	1	3	1	10	10	13
2003.....	347	-	1	-	2	2	4	2	2	3	6	18
2002.....	374	-	-	-	-	-	2	2	3	8	12	12
2001.....	368	1	1	-	-	2	2	5	2	4	10	14
2000.....	351	-	1	-	-	1	3	3	2	8	3	9
1999.....	472	-	-	-	-	-	3	1	2	8	12	20

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Respiratory tuberculosis (A16)

White, male

2004.....	185	-	1	-	-	-	-	1	1	7	8	7
2003.....	208	-	1	-	2	1	2	2	2	3	4	16
2002.....	223	-	-	-	-	-	2	2	1	5	11	6
2001.....	221	-	-	-	-	2	1	4	1	3	7	9
2000.....	216	-	-	-	-	1	2	2	2	8	3	9
1999.....	296	-	-	-	-	-	2	-	2	6	9	17

White, female

2004.....	115	1	-	-	-	-	1	2	-	3	2	6
2003.....	139	-	-	-	-	1	2	-	-	-	2	2
2002.....	151	-	-	-	-	-	-	-	2	3	1	6
2001.....	147	1	1	-	-	-	1	1	1	1	3	5
2000.....	135	-	1	-	-	-	1	1	-	-	-	-
1999.....	176	-	-	-	-	-	1	1	-	2	3	3

Black, both sexes

2004.....	141	-	-	-	1	1	1	1	4	6	9	8
2003.....	138	-	-	-	-	-	3	-	2	1	10	20
2002.....	164	-	-	-	-	-	2	4	4	4	11	13
2001.....	160	-	-	-	-	-	1	3	4	8	4	12
2000.....	192	-	2	-	-	-	1	2	4	6	14	17
1999.....	199	1	1	-	-	-	4	1	7	9	11	15

Black, male

2004.....	88	-	-	-	-	1	1	-	4	5	7	4
2003.....	93	-	-	-	-	-	-	-	2	-	7	13
2002.....	104	-	-	-	-	-	1	2	3	3	8	8
2001.....	106	-	-	-	-	-	-	1	2	5	2	10
2000.....	134	-	1	-	-	-	-	1	3	2	10	16
1999.....	126	-	1	-	-	-	-	-	3	7	7	13

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Respiratory tuberculosis (A16)												
Asian or Pacific Islander 2/, both sexes												
2004.....	54	-	-	-	-	-	1	-	2	-	1	3
2003.....	67	-	-	-	-	-	-	-	-	-	5	3
2002.....	70	-	-	-	-	-	2	-	1	1	3	2
2001.....	64	1	-	-	-	1	1	-	3	2	-	1
2000.....	60	-	-	-	-	1	1	-	-	1	3	2
1999.....	83	-	-	-	1	-	1	-	3	2	3	2
Asian or Pacific Islander 2/, male												
2004.....	37	-	-	-	-	-	-	-	2	-	1	3
2003.....	39	-	-	-	-	-	-	-	-	-	3	1
2002.....	36	-	-	-	-	-	1	-	1	1	3	1
2001.....	41	-	-	-	-	1	1	-	2	1	-	1
2000.....	35	-	-	-	-	1	-	-	-	1	2	2
1999.....	59	-	-	-	1	-	-	-	3	2	2	1
Asian or Pacific Islander 2/, female												
2004.....	17	-	-	-	-	-	1	-	-	-	-	-
2003.....	28	-	-	-	-	-	-	-	-	-	2	2
2002.....	34	-	-	-	-	-	1	-	-	-	-	1
2001.....	23	1	-	-	-	-	-	-	1	1	-	-
2000.....	25	-	-	-	-	-	1	-	-	-	1	-
1999.....	24	-	-	-	-	-	1	-	-	-	1	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other tuberculosis (A17-A19)												
All races, both sexes												
2004.....	153	-	-	-	1	-	3	2	3	8	9	9
2003.....	149	-	-	-	-	1	-	6	6	6	3	10
2002.....	158	1	-	-	-	-	3	1	3	5	7	7
2001.....	164	-	-	-	-	3	2	3	5	4	8	10
2000.....	161	1	3	-	1	-	4	2	7	2	5	8
1999.....	166	1	1	1	1	1	2	1	4	6	6	13
All races, male												
2004.....	90	-	-	-	1	-	2	1	3	7	5	7
2003.....	83	-	-	-	-	-	-	4	3	5	3	7
2002.....	90	-	-	-	-	-	1	-	2	2	5	4
2001.....	82	-	-	-	-	2	1	2	2	3	6	7
2000.....	75	-	-	-	-	-	3	-	6	-	2	6
1999.....	98	-	1	-	1	1	1	1	2	5	5	10
All races, female												
2004.....	63	-	-	-	-	-	1	1	-	1	4	2
2003.....	66	-	-	-	-	1	-	2	3	1	-	3
2002.....	68	1	-	-	-	-	2	1	1	3	2	3
2001.....	82	-	-	-	-	1	1	1	3	1	2	3
2000.....	86	1	3	-	1	-	1	2	1	2	3	2
1999.....	68	1	-	1	-	-	1	-	2	1	1	3
White, both sexes												
2004.....	90	-	-	-	1	-	2	2	2	2	7	5
2003.....	90	-	-	-	-	-	-	2	2	1	1	4
2002.....	87	-	-	-	-	-	1	1	-	-	2	1
2001.....	101	-	-	-	-	2	2	2	1	3	5	6
2000.....	88	-	3	-	1	-	2	2	4	-	2	4
1999.....	92	1	-	-	1	1	1	1	1	4	1	5

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other tuberculosis (A17-A19)												
White, male												
2004.....	52	-	-	-	1	-	2	1	2	1	3	5
2003.....	47	-	-	-	-	-	-	1	-	1	1	2
2002.....	49	-	-	-	-	-	-	-	-	-	1	1
2001.....	50	-	-	-	-	2	1	2	1	2	4	5
2000.....	37	-	-	-	-	-	1	-	3	-	-	3
1999.....	52	-	-	-	1	1	1	1	1	3	1	4
White, female												
2004.....	38	-	-	-	-	-	-	1	-	1	4	-
2003.....	43	-	-	-	-	-	-	1	2	-	-	2
2002.....	38	-	-	-	-	-	1	1	-	-	1	-
2001.....	51	-	-	-	-	-	1	-	-	1	1	1
2000.....	51	-	3	-	1	-	1	2	1	-	2	1
1999.....	40	1	-	-	-	-	-	-	-	1	-	1
Black, both sexes												
2004.....	48	-	-	-	-	-	1	-	1	5	1	4
2003.....	34	-	-	-	-	1	-	1	3	4	1	4
2002.....	40	-	-	-	-	-	2	-	-	5	4	4
2001.....	43	-	-	-	-	-	-	-	3	1	3	2
2000.....	52	-	-	-	-	-	1	-	1	2	3	4
1999.....	46	-	1	1	-	-	-	-	1	1	4	5
Black, male												
2004.....	30	-	-	-	-	-	-	-	1	5	1	2
2003.....	21	-	-	-	-	-	-	1	2	3	1	4
2002.....	23	-	-	-	-	-	1	-	-	2	4	3
2001.....	21	-	-	-	-	-	-	-	1	1	2	1
2000.....	27	-	-	-	-	-	1	-	1	-	2	3
1999.....	33	-	1	-	-	-	-	-	1	1	4	4

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other tuberculosis (A17-A19)												
Black, female												
2004.....	18	-	-	-	-	-	1	-	-	-	-	2
2003.....	13	-	-	-	-	1	-	-	1	1	-	-
2002.....	17	-	-	-	-	-	1	-	-	3	-	1
2001.....	22	-	-	-	-	-	-	-	2	-	1	1
2000.....	25	-	-	-	-	-	-	-	-	2	1	1
1999.....	13	-	-	1	-	-	-	-	-	-	-	1
American Indian 1/, both sexes												
2004.....	4	-	-	-	-	-	-	-	-	-	-	-
2003.....	5	-	-	-	-	-	-	-	-	-	1	1
2002.....	9	-	-	-	-	-	-	-	1	-	1	1
2001.....	5	-	-	-	-	-	-	1	-	-	-	2
2000.....	8	1	-	-	-	-	-	-	-	-	-	-
1999.....	10	-	-	-	-	-	-	-	1	1	-	2
American Indian 1/, male												
2004.....	3	-	-	-	-	-	-	-	-	-	-	-
2003.....	2	-	-	-	-	-	-	-	-	-	1	-
2002.....	5	-	-	-	-	-	-	-	1	-	-	-
2001.....	2	-	-	-	-	-	-	-	-	-	-	1
2000.....	4	-	-	-	-	-	-	-	-	-	-	-
1999.....	6	-	-	-	-	-	-	-	-	1	-	2
American Indian 1/, female												
2004.....	1	-	-	-	-	-	-	-	-	-	-	-
2003.....	3	-	-	-	-	-	-	-	-	-	-	1
2002.....	4	-	-	-	-	-	-	-	-	-	1	1
2001.....	3	-	-	-	-	-	-	1	-	-	-	1
2000.....	4	1	-	-	-	-	-	-	-	-	-	-
1999.....	4	-	-	-	-	-	-	-	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other tuberculosis (A17-A19)												
Asian or Pacific Islander 2/, both sexes												
2004.....	11	-	-	-	-	-	-	-	-	1	1	-
2003.....	20	-	-	-	-	-	-	3	1	1	-	1
2002.....	22	1	-	-	-	-	-	-	2	-	-	1
2001.....	15	-	-	-	-	1	-	-	1	-	-	-
2000.....	13	-	-	-	-	-	1	-	2	-	-	-
1999.....	18	-	-	-	-	-	1	-	1	-	1	1
Asian or Pacific Islander 2/, male												
2004.....	5	-	-	-	-	-	-	-	-	1	1	-
2003.....	13	-	-	-	-	-	-	2	1	1	-	1
2002.....	13	-	-	-	-	-	-	-	1	-	-	-
2001.....	9	-	-	-	-	-	-	-	-	-	-	-
2000.....	7	-	-	-	-	-	1	-	2	-	-	-
1999.....	7	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	6	-	-	-	-	-	-	-	-	-	-	-
2003.....	7	-	-	-	-	-	-	1	-	-	-	-
2002.....	9	1	-	-	-	-	-	-	1	-	-	1
2001.....	6	-	-	-	-	1	-	-	1	-	-	-
2000.....	6	-	-	-	-	-	-	-	-	-	-	-
1999.....	11	-	-	-	-	-	1	-	1	-	1	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Zoonotic and other bacterial diseases (A20-A49)												
All races, both sexes												
2004.....	35,152	307	107	48	38	65	109	132	233	332	579	964
2003.....	35,778	318	107	44	50	82	120	136	203	351	625	988
2002.....	35,548	345	96	56	73	79	90	140	206	352	582	967
2001.....	33,884	360	132	43	60	105	92	111	208	341	554	894
2000.....	32,745	329	126	52	33	80	89	120	194	358	584	838
1999.....	32,714	344	123	62	48	64	111	129	207	366	556	804
All races, male												
2004.....	15,895	186	55	24	18	36	59	63	126	175	292	524
2003.....	15,853	171	66	17	23	44	58	60	118	184	312	541
2002.....	15,770	178	52	34	38	37	47	76	109	176	313	530
2001.....	15,121	196	72	24	34	55	53	58	102	188	297	478
2000.....	14,314	174	70	29	18	50	51	68	104	183	304	470
1999.....	14,378	202	70	31	23	32	62	66	113	197	319	444
All races, female												
2004.....	19,257	121	52	24	20	29	50	69	107	157	287	440
2003.....	19,925	147	41	27	27	38	62	76	85	167	313	447
2002.....	19,778	167	44	22	35	42	43	64	97	176	269	437
2001.....	18,763	164	60	19	26	50	39	53	106	153	257	416
2000.....	18,431	155	56	23	15	30	38	52	90	175	280	368
1999.....	18,336	142	53	31	25	32	49	63	94	169	237	360
White, both sexes												
2004.....	28,139	195	78	39	27	46	64	88	142	238	363	630
2003.....	28,603	174	67	28	35	55	84	83	128	227	417	666
2002.....	28,494	230	70	41	47	62	64	89	137	229	390	675
2001.....	27,160	213	97	35	45	73	54	72	123	217	364	594
2000.....	26,053	211	88	34	28	57	58	74	115	222	397	538
1999.....	25,943	216	90	47	31	49	69	79	141	233	361	513

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Zoonotic and other bacterial diseases (A20-A49)												
White, male												
2004.....	12,702	119	40	21	14	26	35	42	76	131	186	343
2003.....	12,612	89	43	11	18	32	40	36	71	124	208	373
2002.....	12,601	117	42	29	29	28	37	55	69	114	211	381
2001.....	12,047	120	55	19	28	41	29	39	61	128	205	318
2000.....	11,341	110	48	21	15	37	35	39	64	119	213	311
1999.....	11,279	130	51	25	15	24	44	44	75	129	207	284
White, female												
2004.....	15,437	76	38	18	13	20	29	46	66	107	177	287
2003.....	15,991	85	24	17	17	23	44	47	57	103	209	293
2002.....	15,893	113	28	12	18	34	27	34	68	115	179	294
2001.....	15,113	93	42	16	17	32	25	33	62	89	159	276
2000.....	14,712	101	40	13	13	20	23	35	51	103	184	227
1999.....	14,664	86	39	22	16	25	25	35	66	104	154	229
Black, both sexes												
2004.....	6,269	100	24	6	8	17	37	40	75	78	197	297
2003.....	6,452	129	32	13	10	22	27	46	63	110	193	296
2002.....	6,386	100	21	10	22	13	20	47	63	110	177	257
2001.....	6,129	132	28	6	11	29	29	31	78	114	169	270
2000.....	6,163	113	31	16	4	17	26	42	75	123	171	272
1999.....	6,160	115	23	9	13	13	32	45	60	123	183	267
Black, male												
2004.....	2,825	57	12	2	3	9	19	19	41	34	99	159
2003.....	2,914	74	18	6	3	9	13	20	39	52	94	158
2002.....	2,823	54	7	3	8	7	7	17	34	56	92	129
2001.....	2,756	69	14	3	4	13	20	14	37	55	82	140
2000.....	2,716	62	17	7	2	10	14	27	37	58	80	144
1999.....	2,795	63	14	5	7	8	13	18	36	63	105	145

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Zoonotic and other bacterial diseases (A20-A49)

Black, female

2004.....	3,444	43	12	4	5	8	18	21	34	44	98	138
2003.....	3,538	55	14	7	7	13	14	26	24	58	99	138
2002.....	3,563	46	14	7	14	6	13	30	29	54	85	128
2001.....	3,373	63	14	3	7	16	9	17	41	59	87	130
2000.....	3,447	51	14	9	2	7	12	15	38	65	91	128
1999.....	3,365	52	9	4	6	5	19	27	24	60	78	122

American Indian 1/, both sexes

2004.....	203	6	2	2	3	-	2	3	8	8	11	18
2003.....	240	5	4	2	2	3	4	3	5	10	13	10
2002.....	213	5	4	1	2	-	2	1	3	8	7	17
2001.....	167	4	4	1	2	-	3	2	2	6	10	18
2000.....	174	2	3	1	-	2	1	1	2	7	7	12
1999.....	177	4	1	3	1	-	4	1	2	7	5	10

American Indian 1/, male

2004.....	95	6	2	-	1	-	1	1	3	6	4	9
2003.....	94	3	2	-	1	3	2	1	4	5	9	2
2002.....	101	2	3	1	1	-	-	1	3	3	4	9
2001.....	86	2	2	1	1	-	1	1	1	2	5	12
2000.....	81	2	2	1	-	1	-	-	2	3	5	7
1999.....	83	4	-	1	-	-	2	1	1	3	4	6

American Indian 1/, female

2004.....	108	-	-	2	2	-	1	2	5	2	7	9
2003.....	146	2	2	2	1	-	2	2	1	5	4	8
2002.....	112	3	1	-	1	-	2	-	-	5	3	8
2001.....	81	2	2	-	1	-	2	1	1	4	5	6
2000.....	93	-	1	-	-	1	1	1	-	4	2	5
1999.....	94	-	1	2	1	-	2	-	1	4	1	4

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Zoonotic and other bacterial diseases (A20-A49)

Asian or Pacific Islander 2/, both sexes

2004.....	541	6	3	1	-	2	6	1	8	8	8	19
2003.....	483	10	4	1	3	2	5	4	7	4	2	16
2002.....	455	10	1	4	2	4	4	3	3	5	8	18
2001.....	428	11	3	1	2	3	6	6	5	4	11	12
2000.....	355	3	4	1	1	4	4	3	2	6	9	16
1999.....	434	9	9	3	3	2	6	4	4	3	7	14

Asian or Pacific Islander 2/, male

2004.....	273	4	1	1	-	1	4	1	6	4	3	13
2003.....	233	5	3	-	1	-	3	3	4	3	1	8
2002.....	245	5	-	1	-	2	3	3	3	3	6	11
2001.....	232	5	1	1	1	1	3	4	3	3	5	8
2000.....	176	-	3	-	1	2	2	2	1	3	6	8
1999.....	221	5	5	-	1	-	3	3	1	2	3	9

Asian or Pacific Islander 2/, female

2004.....	268	2	2	-	-	1	2	-	2	4	5	6
2003.....	250	5	1	1	2	2	2	1	3	1	1	8
2002.....	210	5	1	3	2	2	1	-	-	2	2	7
2001.....	196	6	2	-	1	2	3	2	2	1	6	4
2000.....	179	3	1	1	-	2	2	1	1	3	3	8
1999.....	213	4	4	3	2	2	3	1	3	1	4	5

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Tetanus (A33-A35)												
All races, both sexes												
2004.....	4	-	-	-	-	-	-	-	-	-	-	-
2003.....	4	-	-	-	-	-	-	-	-	-	-	-
2002.....	5	-	-	-	-	-	-	-	-	-	-	-
2001.....	5	-	-	-	-	-	-	-	-	-	-	-
2000.....	5	-	-	-	-	-	-	-	-	-	-	-
1999.....	7	-	-	-	-	-	-	-	1	-	-	-
All races, male												
2004.....	1	-	-	-	-	-	-	-	-	-	-	-
2003.....	2	-	-	-	-	-	-	-	-	-	-	-
2002.....	3	-	-	-	-	-	-	-	-	-	-	-
2001.....	2	-	-	-	-	-	-	-	-	-	-	-
2000.....	2	-	-	-	-	-	-	-	-	-	-	-
1999.....	4	-	-	-	-	-	-	-	-	-	-	-
All races, female												
2004.....	3	-	-	-	-	-	-	-	-	-	-	-
2003.....	2	-	-	-	-	-	-	-	-	-	-	-
2002.....	2	-	-	-	-	-	-	-	-	-	-	-
2001.....	3	-	-	-	-	-	-	-	-	-	-	-
2000.....	3	-	-	-	-	-	-	-	-	-	-	-
1999.....	3	-	-	-	-	-	-	-	1	-	-	-
White, both sexes												
2004.....	4	-	-	-	-	-	-	-	-	-	-	-
2003.....	3	-	-	-	-	-	-	-	-	-	-	-
2002.....	5	-	-	-	-	-	-	-	-	-	-	-
2001.....	4	-	-	-	-	-	-	-	-	-	-	-
2000.....	4	-	-	-	-	-	-	-	-	-	-	-
1999.....	6	-	-	-	-	-	-	-	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Meningococcal infection (A39)

All races, both sexes

2004.....	138	8	18	8	5	16	15	9	5	6	9	7
2003.....	161	12	18	2	9	18	21	8	4	10	9	8
2002.....	161	17	12	6	14	21	12	12	4	4	5	10
2001.....	199	16	19	12	14	43	12	6	4	8	5	9
2000.....	211	22	19	12	6	34	22	9	6	7	6	13
1999.....	227	32	30	13	12	28	11	11	12	12	15	11

All races, male

2004.....	79	2	11	5	4	11	8	7	3	3	5	5
2003.....	75	9	10	1	2	8	10	3	4	4	5	5
2002.....	83	8	9	5	8	11	6	8	2	2	3	5
2001.....	94	6	9	6	9	20	7	4	2	3	3	6
2000.....	106	14	12	5	2	24	11	3	3	6	3	5
1999.....	101	14	19	7	6	13	6	8	4	3	6	1

All races, female

2004.....	59	6	7	3	1	5	7	2	2	3	4	2
2003.....	86	3	8	1	7	10	11	5	-	6	4	3
2002.....	78	9	3	1	6	10	6	4	2	2	2	5
2001.....	105	10	10	6	5	23	5	2	2	5	2	3
2000.....	105	8	7	7	4	10	11	6	3	1	3	8
1999.....	126	18	11	6	6	15	5	3	8	9	9	10

White, both sexes

2004.....	110	7	15	7	5	12	12	8	2	5	5	6
2003.....	121	12	11	1	6	14	16	6	2	9	5	6
2002.....	125	14	8	5	5	17	10	9	4	1	4	9
2001.....	158	14	14	8	12	31	9	5	2	6	5	8
2000.....	161	19	14	5	5	29	16	7	3	3	4	7
1999.....	184	30	26	10	8	20	9	6	8	8	11	10

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Meningococcal infection (A39)												
White, male												
2004.....	62	2	10	4	4	7	6	6	2	3	3	5
2003.....	56	9	6	-	2	6	8	3	2	4	3	4
2002.....	67	8	6	5	4	8	6	7	2	1	2	5
2001.....	72	5	7	4	8	15	5	3	1	1	3	5
2000.....	75	11	8	2	2	20	8	2	-	2	3	2
1999.....	79	14	17	5	4	8	5	4	2	2	4	1
White, female												
2004.....	48	5	5	3	1	5	6	2	-	2	2	1
2003.....	65	3	5	1	4	8	8	3	-	5	2	2
2002.....	58	6	2	-	1	9	4	2	2	-	2	4
2001.....	86	9	7	4	4	16	4	2	1	5	2	3
2000.....	86	8	6	3	3	9	8	5	3	1	1	5
1999.....	105	16	9	5	4	12	4	2	6	6	7	9
Black, both sexes												
2004.....	22	-	3	1	-	3	2	1	2	1	3	1
2003.....	36	-	7	1	3	3	4	2	1	1	4	2
2002.....	29	3	3	-	8	3	2	2	-	3	1	1
2001.....	36	2	4	4	2	10	3	-	2	1	-	1
2000.....	44	3	5	7	1	3	4	1	3	4	2	5
1999.....	39	2	4	2	4	7	2	4	4	4	4	1
Black, male												
2004.....	13	-	1	1	-	3	1	1	1	-	1	-
2003.....	17	-	4	1	-	1	2	-	1	-	2	1
2002.....	12	-	2	-	4	2	-	-	-	1	1	-
2001.....	18	1	1	2	1	4	2	-	1	1	-	1
2000.....	27	3	4	3	-	3	2	-	3	4	-	2
1999.....	20	-	2	1	2	5	1	3	2	1	2	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Meningococcal infection (A39)												
Asian or Pacific Islander 2/, both sexes												
2004.....	3	1	-	-	-	1	1	-	-	-	-	-
2003.....	1	-	-	-	-	-	-	-	1	-	-	-
2002.....	5	-	-	1	-	1	-	1	-	-	-	-
2001.....	4	-	-	-	-	2	-	1	-	1	-	-
2000.....	5	-	-	-	-	1	2	1	-	-	-	1
1999.....	2	-	-	-	-	1	-	1	-	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	2	-	-	-	-	1	1	-	-	-	-	-
2003.....	1	-	-	-	-	-	-	-	1	-	-	-
2002.....	3	-	-	-	-	1	-	1	-	-	-	-
2001.....	3	-	-	-	-	1	-	1	-	1	-	-
2000.....	4	-	-	-	-	1	1	1	-	-	-	1
1999.....	1	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	1	1	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	2	-	-	1	-	-	-	-	-	-	-	-
2001.....	1	-	-	-	-	1	-	-	-	-	-	-
2000.....	1	-	-	-	-	-	1	-	-	-	-	-
1999.....	1	-	-	-	-	1	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Septicemia (A40-A41)												
All races, both sexes												
2004.....	33,373	271	84	38	28	48	87	115	213	306	540	898
2003.....	34,069	278	85	39	38	59	95	123	186	315	595	916
2002.....	33,865	296	79	42	53	49	69	119	195	318	538	904
2001.....	32,238	312	108	29	40	57	76	96	185	309	520	837
2000.....	31,224	274	99	38	25	40	60	104	176	324	553	781
1999.....	30,680	283	87	47	30	32	89	106	169	314	498	731
All races, male												
2004.....	15,011	167	41	18	13	25	46	51	114	162	271	487
2003.....	14,987	151	54	15	19	35	47	56	105	167	288	498
2002.....	14,947	158	41	26	26	22	37	63	103	159	282	495
2001.....	14,307	174	60	17	22	31	44	50	89	170	281	439
2000.....	13,537	139	53	24	14	23	37	62	96	162	289	439
1999.....	13,395	175	48	22	14	17	50	53	94	170	290	406
All races, female												
2004.....	18,362	104	43	20	15	23	41	64	99	144	269	411
2003.....	19,082	127	31	24	19	24	48	67	81	148	307	418
2002.....	18,918	138	38	16	27	27	32	56	92	159	256	409
2001.....	17,931	138	48	12	18	26	32	46	96	139	239	398
2000.....	17,687	135	46	14	11	17	23	42	80	162	264	342
1999.....	17,285	108	39	25	16	15	39	53	75	144	208	325
White, both sexes												
2004.....	26,678	165	60	31	19	34	48	75	131	216	343	584
2003.....	27,180	142	53	26	28	37	66	74	116	202	396	613
2002.....	27,110	191	58	31	38	38	48	75	127	210	364	628
2001.....	25,806	171	80	25	29	37	45	61	108	191	340	547
2000.....	24,804	169	66	28	21	24	36	61	104	204	374	495
1999.....	24,287	164	60	35	20	26	53	66	116	194	316	457

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Septicemia (A40-A41)												
White, male												
2004.....	11,986	103	29	17	9	19	27	33	68	120	174	315
2003.....	11,909	74	35	11	15	25	31	32	63	112	191	343
2002.....	11,919	100	34	21	21	16	29	45	64	104	190	354
2001.....	11,371	100	46	14	18	22	24	33	52	115	194	286
2000.....	10,716	88	35	19	11	15	24	34	61	110	201	289
1999.....	10,489	106	32	18	10	14	35	37	65	109	185	256
White, female												
2004.....	14,692	62	31	14	10	15	21	42	63	96	169	269
2003.....	15,271	68	18	15	13	12	35	42	53	90	205	270
2002.....	15,191	91	24	10	17	22	19	30	63	106	174	274
2001.....	14,435	71	34	11	11	15	21	28	56	76	146	261
2000.....	14,088	81	31	9	10	9	12	27	43	94	173	206
1999.....	13,798	58	28	17	10	12	18	29	51	85	131	201
Black, both sexes												
2004.....	6,010	96	19	5	7	13	33	36	68	75	182	279
2003.....	6,206	123	24	10	5	18	23	42	60	101	184	278
2002.....	6,137	91	17	7	13	9	15	41	62	97	161	243
2001.....	5,880	126	22	2	8	19	22	28	70	109	160	261
2000.....	5,928	101	26	8	3	12	21	40	68	109	163	259
1999.....	5,826	107	17	7	7	5	28	37	48	110	172	251
Black, male												
2004.....	2,690	55	9	1	3	6	16	16	37	33	93	151
2003.....	2,775	71	14	4	2	8	11	20	36	48	87	145
2002.....	2,703	51	5	3	4	5	5	15	33	50	83	122
2001.....	2,635	67	12	1	2	9	16	13	33	51	77	133
2000.....	2,587	50	13	4	2	6	12	27	32	48	77	136
1999.....	2,623	61	11	4	3	3	11	14	27	56	99	136

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Septicemia (A40-A41)

Black, female

2004.....	3,320	41	10	4	4	7	17	20	31	42	89	128
2003.....	3,431	52	10	6	3	10	12	22	24	53	97	133
2002.....	3,434	40	12	4	9	4	10	26	29	47	78	121
2001.....	3,245	59	10	1	6	10	6	15	37	58	83	128
2000.....	3,341	51	13	4	1	6	9	13	36	61	86	123
1999.....	3,203	46	6	3	4	2	17	23	21	54	73	115

American Indian 1/, both sexes

2004.....	183	5	2	2	2	-	2	3	6	7	9	16
2003.....	227	4	4	2	2	2	2	3	5	9	13	9
2002.....	195	4	3	1	1	-	2	1	3	7	6	16
2001.....	155	4	3	1	1	-	3	2	2	6	9	18
2000.....	162	1	3	1	-	1	1	1	2	6	7	12
1999.....	169	3	1	2	1	-	4	1	2	7	5	10

American Indian 1/, male

2004.....	85	5	2	-	1	-	1	1	3	5	3	8
2003.....	89	2	2	-	1	2	2	1	4	5	9	2
2002.....	95	2	2	1	1	-	-	1	3	3	3	8
2001.....	83	2	1	1	1	-	1	1	1	2	5	12
2000.....	74	1	2	1	-	1	-	-	2	2	5	7
1999.....	81	3	-	-	-	-	2	1	1	3	4	6

American Indian 1/, female

2004.....	98	-	-	2	1	-	1	2	3	2	6	8
2003.....	138	2	2	2	1	-	-	2	1	4	4	7
2002.....	100	2	1	-	-	-	2	-	-	4	3	8
2001.....	72	2	2	-	-	-	2	1	1	4	4	6
2000.....	88	-	1	-	-	-	1	1	-	4	2	5
1999.....	88	-	1	2	1	-	2	-	1	4	1	4

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Septicemia (A40-A41)												
Asian or Pacific Islander 2/, both sexes												
2004.....	502	5	3	-	-	1	4	1	8	8	6	19
2003.....	456	9	4	1	3	2	4	4	5	3	2	16
2002.....	423	10	1	3	1	2	4	2	3	4	7	17
2001.....	397	11	3	1	2	1	6	5	5	3	11	11
2000.....	330	3	4	1	1	3	2	2	2	5	9	15
1999.....	398	9	9	3	2	1	4	2	3	3	5	13
Asian or Pacific Islander 2/, male												
2004.....	250	4	1	-	-	-	2	1	6	4	1	13
2003.....	214	4	3	-	1	-	3	3	2	2	1	8
2002.....	230	5	-	1	-	1	3	2	3	2	6	11
2001.....	218	5	1	1	1	-	3	3	3	2	5	8
2000.....	160	-	3	-	1	1	1	1	1	2	6	7
1999.....	202	5	5	-	1	-	2	1	1	2	2	8
Asian or Pacific Islander 2/, female												
2004.....	252	1	2	-	-	1	2	-	2	4	5	6
2003.....	242	5	1	1	2	2	1	1	3	1	1	8
2002.....	193	5	1	2	1	1	1	-	-	2	1	6
2001.....	179	6	2	-	1	1	3	2	2	1	6	3
2000.....	170	3	1	1	-	2	1	1	1	3	3	8
1999.....	196	4	4	3	1	1	2	1	2	1	3	5

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other zoonotic and bacterial diseases (A21-A22,A24-A32,A42-A44,A48-A49)

All races, both sexes

2004.....	1,617	13	5	2	4	1	7	8	15	20	30	59
2003.....	1,531	18	3	3	3	5	4	5	13	26	21	64
2002.....	1,496	14	5	8	6	9	9	9	7	30	39	53
2001.....	1,422	16	4	1	6	5	4	9	19	24	29	48
2000.....	1,290	20	8	2	2	6	7	7	12	27	25	43
1999.....	1,786	22	6	2	6	4	11	12	25	40	43	62

All races, male

2004.....	792	8	3	1	-	-	5	5	9	10	16	32
2003.....	787	10	2	1	2	1	1	1	9	13	19	38
2002.....	731	6	2	3	4	4	4	5	4	15	28	30
2001.....	709	10	2	-	3	4	2	4	11	15	13	33
2000.....	663	16	5	-	2	3	3	3	5	15	12	26
1999.....	876	12	3	2	3	2	6	5	15	24	23	37

All races, female

2004.....	825	5	2	1	4	1	2	3	6	10	14	27
2003.....	744	8	1	2	1	4	3	4	4	13	2	26
2002.....	765	8	3	5	2	5	5	4	3	15	11	23
2001.....	713	6	2	1	3	1	2	5	8	9	16	15
2000.....	627	4	3	2	-	3	4	4	7	12	13	17
1999.....	910	10	3	-	3	2	5	7	10	16	20	25

White, both sexes

2004.....	1,330	11	3	1	2	-	4	5	9	17	15	40
2003.....	1,287	11	2	1	1	4	2	3	10	16	16	47
2002.....	1,236	10	4	5	4	7	6	5	6	18	22	38
2001.....	1,173	13	2	1	4	5	-	6	13	20	19	39
2000.....	1,070	11	8	1	2	4	6	6	8	15	19	35
1999.....	1,454	16	4	2	3	3	7	7	16	31	34	46

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other zoonotic and bacterial diseases (A21-A22,A24-A32,A42-A44,A48-A49)

White, male

2004.....	643	7	1	-	-	-	2	3	6	8	9	23
2003.....	644	6	2	-	1	1	1	1	6	8	14	26
2002.....	607	4	2	3	4	4	2	3	3	9	19	22
2001.....	593	9	1	-	2	4	-	3	8	12	8	27
2000.....	543	7	5	-	2	2	3	3	3	7	9	20
1999.....	706	9	2	2	1	2	4	3	8	18	18	27

White, female

2004.....	687	4	2	1	2	-	2	2	3	9	6	17
2003.....	643	5	-	1	-	3	1	2	4	8	2	21
2002.....	629	6	2	2	-	3	4	2	3	9	3	16
2001.....	580	4	1	1	2	1	-	3	5	8	11	12
2000.....	527	4	3	1	-	2	3	3	5	8	10	15
1999.....	748	7	2	-	2	1	3	4	8	13	16	19

Black, both sexes

2004.....	235	2	2	-	1	1	2	3	5	2	12	17
2003.....	209	6	1	2	2	1	-	2	2	8	5	16
2002.....	218	4	1	3	1	1	3	4	1	10	15	13
2001.....	211	3	2	-	1	-	4	3	6	4	9	8
2000.....	189	8	-	1	-	2	1	1	4	10	6	8
1999.....	292	5	2	-	2	1	2	4	8	9	7	15

Black, male

2004.....	121	1	2	-	-	-	2	2	3	1	5	8
2003.....	122	3	-	1	1	-	-	-	2	4	5	12
2002.....	107	2	-	-	-	-	2	2	1	5	8	7
2001.....	103	1	1	-	1	-	2	1	3	3	5	6
2000.....	101	8	-	-	-	1	-	-	2	6	3	6
1999.....	151	2	1	-	2	-	1	1	7	6	4	9

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other zoonotic and bacterial diseases (A21-A22,A24-A32,A42-A44,A48-A49)												
Asian or Pacific Islander 2/, both sexes												
2004.....	36	-	-	1	-	-	1	-	-	-	2	-
2003.....	26	1	-	-	-	-	1	-	1	1	-	-
2002.....	27	-	-	-	1	1	-	-	-	1	1	1
2001.....	27	-	-	-	-	-	-	-	-	-	-	1
2000.....	20	-	-	-	-	-	-	-	-	1	-	-
1999.....	34	-	-	-	1	-	2	1	1	-	2	1
Asian or Pacific Islander 2/, male												
2004.....	21	-	-	1	-	-	1	-	-	-	2	-
2003.....	18	1	-	-	-	-	-	-	1	1	-	-
2002.....	12	-	-	-	-	-	-	-	-	1	-	-
2001.....	11	-	-	-	-	-	-	-	-	-	-	-
2000.....	12	-	-	-	-	-	-	-	-	1	-	-
1999.....	18	-	-	-	-	-	1	1	-	-	1	1
Asian or Pacific Islander 2/, female												
2004.....	15	-	-	-	-	-	-	-	-	-	-	-
2003.....	8	-	-	-	-	-	1	-	-	-	-	-
2002.....	15	-	-	-	1	1	-	-	-	-	1	1
2001.....	16	-	-	-	-	-	-	-	-	-	-	1
2000.....	8	-	-	-	-	-	-	-	-	-	-	-
1999.....	16	-	-	-	1	-	1	-	1	-	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Infections with a predominately sexual mode of transmission (A50-A64)												
All races, both sexes												
2004.....	52	-	-	-	-	-	-	-	3	1	2	2
2003.....	49	-	-	-	-	1	-	-	-	1	2	2
2002.....	49	2	1	-	-	1	1	1	-	1	1	1
2001.....	44	1	1	-	1	-	-	-	2	4	-	2
2000.....	60	1	-	-	-	1	2	1	1	3	-	2
1999.....	47	2	-	-	-	2	1	-	1	1	2	3
All races, male												
2004.....	26	-	-	-	-	-	-	-	2	1	1	1
2003.....	27	-	-	-	-	1	-	-	-	1	2	2
2002.....	24	-	1	-	-	-	-	1	-	1	1	-
2001.....	22	1	-	-	1	-	-	-	-	2	-	2
2000.....	35	1	-	-	-	1	1	-	1	2	-	1
1999.....	19	-	-	-	-	2	1	-	1	-	-	2
All races, female												
2004.....	26	-	-	-	-	-	-	-	1	-	1	1
2003.....	22	-	-	-	-	-	-	-	-	-	-	-
2002.....	25	2	-	-	-	1	1	-	-	-	-	1
2001.....	22	-	1	-	-	-	-	-	2	2	-	-
2000.....	25	-	-	-	-	-	1	1	-	1	-	1
1999.....	28	2	-	-	-	-	-	-	-	1	2	1
White, both sexes												
2004.....	26	-	-	-	-	-	-	-	-	1	2	2
2003.....	31	-	-	-	-	1	-	-	-	1	-	-
2002.....	24	-	-	-	-	-	1	1	-	-	1	-
2001.....	26	1	1	-	-	-	-	-	1	2	-	-
2000.....	36	-	-	-	-	-	1	1	1	3	-	1
1999.....	29	2	-	-	-	1	-	-	1	-	1	3

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Infections with a predominately sexual mode of transmission (A50-A64)												
White, male												
2004.....	12	-	-	-	-	-	-	-	-	1	1	1
2003.....	15	-	-	-	-	1	-	-	-	1	-	-
2002.....	12	-	-	-	-	-	-	1	-	-	1	-
2001.....	10	1	-	-	-	-	-	-	-	-	-	-
2000.....	15	-	-	-	-	-	-	-	1	2	-	-
1999.....	11	-	-	-	-	1	-	-	1	-	-	2
White, female												
2004.....	14	-	-	-	-	-	-	-	-	-	1	1
2003.....	16	-	-	-	-	-	-	-	-	-	-	-
2002.....	12	-	-	-	-	-	1	-	-	-	-	-
2001.....	16	-	1	-	-	-	-	-	1	2	-	-
2000.....	21	-	-	-	-	-	1	1	-	1	-	1
1999.....	18	2	-	-	-	-	-	-	-	-	1	1
Black, both sexes												
2004.....	23	-	-	-	-	-	-	-	3	-	-	-
2003.....	16	-	-	-	-	-	-	-	-	-	2	2
2002.....	25	2	1	-	-	1	-	-	-	1	-	1
2001.....	15	-	-	-	1	-	-	-	1	2	-	2
2000.....	24	1	-	-	-	1	1	-	-	-	-	1
1999.....	17	-	-	-	-	1	1	-	-	1	1	-
Black, male												
2004.....	13	-	-	-	-	-	-	-	2	-	-	-
2003.....	11	-	-	-	-	-	-	-	-	-	2	2
2002.....	12	-	1	-	-	-	-	-	-	1	-	-
2001.....	10	-	-	-	1	-	-	-	-	2	-	2
2000.....	20	1	-	-	-	1	1	-	-	-	-	1
1999.....	8	-	-	-	-	1	1	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Syphilis (A50-A53)												
All races, both sexes												
2004.....	43	-	-	-	-	-	-	-	3	1	2	-
2003.....	34	-	-	-	-	-	-	-	-	1	2	2
2002.....	41	1	1	-	-	-	-	-	-	1	1	1
2001.....	36	1	-	-	1	-	-	-	1	3	-	2
2000.....	41	1	-	-	-	-	-	-	1	-	-	2
1999.....	33	2	-	-	-	-	-	-	1	-	1	3
All races, male												
2004.....	21	-	-	-	-	-	-	-	2	1	1	-
2003.....	22	-	-	-	-	-	-	-	-	1	2	2
2002.....	21	-	1	-	-	-	-	-	-	1	1	-
2001.....	19	1	-	-	1	-	-	-	-	2	-	2
2000.....	24	1	-	-	-	-	-	-	1	-	-	1
1999.....	13	-	-	-	-	-	-	-	1	-	-	2
All races, female												
2004.....	22	-	-	-	-	-	-	-	1	-	1	-
2003.....	12	-	-	-	-	-	-	-	-	-	-	-
2002.....	20	1	-	-	-	-	-	-	-	-	-	1
2001.....	17	-	-	-	-	-	-	-	1	1	-	-
2000.....	17	-	-	-	-	-	-	-	-	-	-	1
1999.....	20	2	-	-	-	-	-	-	-	-	1	1
White, both sexes												
2004.....	17	-	-	-	-	-	-	-	-	1	2	-
2003.....	17	-	-	-	-	-	-	-	-	1	-	-
2002.....	18	-	-	-	-	-	-	-	-	-	1	-
2001.....	21	1	-	-	-	-	-	-	-	1	-	-
2000.....	21	-	-	-	-	-	-	-	1	-	-	1
1999.....	20	2	-	-	-	-	-	-	1	-	1	3

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Neurosyphilis (A52.1-A52.3)

All races, both sexes

2004.....	30	-	-	-	-	-	-	-	3	1	1	-
2003.....	18	-	-	-	-	-	-	-	-	1	2	2
2002.....	26	-	-	-	-	-	-	-	-	-	1	1
2001.....	19	-	-	-	-	-	-	-	-	1	-	1
2000.....	28	-	-	-	-	-	-	-	-	-	-	1
1999.....	16	-	-	-	-	-	-	-	1	-	1	-

All races, male

2004.....	16	-	-	-	-	-	-	-	2	1	1	-
2003.....	13	-	-	-	-	-	-	-	-	1	2	2
2002.....	11	-	-	-	-	-	-	-	-	-	1	-
2001.....	9	-	-	-	-	-	-	-	-	-	-	1
2000.....	17	-	-	-	-	-	-	-	-	-	-	1
1999.....	9	-	-	-	-	-	-	-	1	-	-	-

All races, female

2004.....	14	-	-	-	-	-	-	-	1	-	-	-
2003.....	5	-	-	-	-	-	-	-	-	-	-	-
2002.....	15	-	-	-	-	-	-	-	-	-	-	1
2001.....	10	-	-	-	-	-	-	-	-	1	-	-
2000.....	11	-	-	-	-	-	-	-	-	-	-	-
1999.....	7	-	-	-	-	-	-	-	-	-	1	-

White, both sexes

2004.....	11	-	-	-	-	-	-	-	-	1	1	-
2003.....	10	-	-	-	-	-	-	-	-	1	-	-
2002.....	12	-	-	-	-	-	-	-	-	-	1	-
2001.....	14	-	-	-	-	-	-	-	-	1	-	-
2000.....	13	-	-	-	-	-	-	-	-	-	-	-
1999.....	8	-	-	-	-	-	-	-	1	-	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gonococcal infection (A54)												
All races, both sexes												
2004.....	2	-	-	-	-	-	-	-	-	-	-	1
2003.....	6	-	-	-	-	1	-	-	-	-	-	-
2002.....	7	1	-	-	-	1	1	1	-	-	-	-
2001.....	7	-	1	-	-	-	-	-	1	1	-	-
2000.....	12	-	-	-	-	1	2	1	-	3	-	-
1999.....	9	-	-	-	-	2	1	-	-	1	1	-
All races, male												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	1	-	-	-	-	1	-	-	-	-	-	-
2002.....	3	-	-	-	-	-	-	1	-	-	-	-
2001.....	2	-	-	-	-	-	-	-	-	-	-	-
2000.....	6	-	-	-	-	1	1	-	-	2	-	-
1999.....	4	-	-	-	-	2	1	-	-	-	-	-
All races, female												
2004.....	2	-	-	-	-	-	-	-	-	-	-	1
2003.....	5	-	-	-	-	-	-	-	-	-	-	-
2002.....	4	1	-	-	-	1	1	-	-	-	-	-
2001.....	5	-	1	-	-	-	-	-	1	1	-	-
2000.....	6	-	-	-	-	-	1	1	-	1	-	-
1999.....	5	-	-	-	-	-	-	-	-	1	1	-
White, both sexes												
2004.....	2	-	-	-	-	-	-	-	-	-	-	1
2003.....	6	-	-	-	-	1	-	-	-	-	-	-
2002.....	5	-	-	-	-	-	1	1	-	-	-	-
2001.....	5	-	1	-	-	-	-	-	1	1	-	-
2000.....	9	-	-	-	-	-	1	1	-	3	-	-
1999.....	5	-	-	-	-	1	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gonococcal infection (A54)												
White, male												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	1	-	-	-	-	1	-	-	-	-	-	-
2002.....	3	-	-	-	-	-	-	1	-	-	-	-
2001.....	1	-	-	-	-	-	-	-	-	-	-	-
2000.....	4	-	-	-	-	-	-	-	-	2	-	-
1999.....	2	-	-	-	-	1	-	-	-	-	-	-
White, female												
2004.....	2	-	-	-	-	-	-	-	-	-	-	1
2003.....	5	-	-	-	-	-	-	-	-	-	-	-
2002.....	2	-	-	-	-	-	1	-	-	-	-	-
2001.....	4	-	1	-	-	-	-	-	1	1	-	-
2000.....	5	-	-	-	-	-	1	1	-	1	-	-
1999.....	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	2	1	-	-	-	1	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	3	-	-	-	-	1	1	-	-	-	-	-
1999.....	4	-	-	-	-	1	1	-	-	1	1	-
Black, male												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	2	-	-	-	-	1	1	-	-	-	-	-
1999.....	2	-	-	-	-	1	1	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other infections caused by spirochetes, chlamydia or rickettsia (A65-A79)												
White, male												
2004.....	9	-	-	-	-	-	-	-	-	-	1	3
2003.....	10	-	1	-	-	-	-	-	1	1	1	1
2002.....	7	-	-	-	-	-	-	-	-	1	-	1
2001.....	5	-	-	-	-	-	-	-	-	-	-	-
2000.....	6	-	1	-	-	-	1	-	-	-	1	1
1999.....	10	-	-	-	-	-	-	-	-	-	1	-
White, female												
2004.....	6	-	-	1	-	-	-	-	-	-	2	-
2003.....	3	-	-	-	-	1	-	-	-	-	-	-
2002.....	9	-	-	2	-	-	-	-	-	-	-	-
2001.....	4	-	-	-	-	-	-	-	-	-	-	1
2000.....	5	-	-	-	-	1	-	-	-	-	-	-
1999.....	7	-	1	-	-	-	-	-	-	1	-	-
Black, both sexes												
2004.....	2	-	-	-	-	-	-	-	-	-	-	-
2003.....	2	-	-	-	-	-	1	-	-	-	-	-
2002.....	1	-	-	-	-	-	-	-	-	-	-	-
2001.....	1	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	1	-	-	-	1	-	-	-	-	-	-	-
Black, male												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	2	-	-	-	-	-	1	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	1	-	-	-	1	-	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other infections caused by spirochetes, chlamydia or rickettsia (A65-A79)

Asian or Pacific Islander 2/, both sexes

2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	1	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	1	-	-	-	-	-	-	-	-	-	-	1
1999.....	2	-	-	-	-	-	-	-	1	-	-	-

Asian or Pacific Islander 2/, male

2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	1	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	1	-	-	-	-	-	-	-	-	-	-	1
1999.....	-	-	-	-	-	-	-	-	-	-	-	-

Asian or Pacific Islander 2/, female

2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	2	-	-	-	-	-	-	-	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

All other infections caused by spirochetes, chlamydia or rickettsia (A65-A68,A69.0-A69.1,A69.8-A69.9,A70-A74,A77-A79)

White, male

2004.....	5	-	-	-	-	-	-	-	-	-	1	1
2003.....	8	-	1	-	-	-	-	-	1	1	1	1
2002.....	2	-	-	-	-	-	-	-	-	1	-	-
2001.....	4	-	-	-	-	-	-	-	-	-	-	-
2000.....	5	-	1	-	-	-	1	-	-	-	1	1
1999.....	7	-	-	-	-	-	-	-	-	-	1	-

White, female

2004.....	4	-	-	1	-	-	-	-	-	-	2	-
2003.....	2	-	-	-	-	1	-	-	-	-	-	-
2002.....	8	-	-	2	-	-	-	-	-	-	-	-
2001.....	3	-	-	-	-	-	-	-	-	-	-	1
2000.....	1	-	-	-	-	-	-	-	-	-	-	-
1999.....	3	-	1	-	-	-	-	-	-	1	-	-

Black, both sexes

2004.....	1	-	-	-	-	-	-	-	-	-	-	-
2003.....	1	-	-	-	-	-	1	-	-	-	-	-
2002.....	1	-	-	-	-	-	-	-	-	-	-	-
2001.....	1	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	1	-	-	-	1	-	-	-	-	-	-	-

Black, male

2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	1	-	-	-	-	-	1	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	1	-	-	-	1	-	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

All other infections caused by spirochetes, chlamydia or rickettsia (A65-A68,A69.0-A69.1,A69.8-A69.9,A70-A74,A77-A79)

Asian or Pacific Islander 2/, both sexes

2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	1	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	2	-	-	-	-	-	-	-	1	-	-	-

Asian or Pacific Islander 2/, male

2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	1	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-

Asian or Pacific Islander 2/, female

2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	2	-	-	-	-	-	-	-	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Viral diseases (A80-B34)												
All races, both sexes												
2004.....	19,823	117	54	22	37	64	193	474	1,094	2,167	3,336	3,560
2003.....	20,652	116	68	34	35	63	176	484	1,206	2,599	3,468	3,734
2002.....	21,408	164	54	23	44	69	176	558	1,427	2,873	3,667	3,925
2001.....	20,938	121	66	44	35	73	214	616	1,588	2,980	3,756	3,639
2000.....	21,122	111	63	36	49	61	182	695	1,864	3,146	3,616	3,637
1999.....	21,080	134	78	60	45	55	189	800	2,060	3,225	3,808	3,359
All races, male												
2004.....	13,710	66	29	13	18	29	113	273	695	1,511	2,391	2,634
2003.....	14,424	51	34	16	19	33	98	293	807	1,849	2,536	2,802
2002.....	14,976	94	35	11	20	31	102	320	958	2,117	2,649	2,957
2001.....	14,684	65	40	23	19	39	101	367	1,113	2,191	2,760	2,789
2000.....	14,844	57	34	19	21	35	102	398	1,273	2,342	2,674	2,813
1999.....	14,908	77	51	31	27	28	94	519	1,470	2,394	2,882	2,613
All races, female												
2004.....	6,113	51	25	9	19	35	80	201	399	656	945	926
2003.....	6,228	65	34	18	16	30	78	191	399	750	932	932
2002.....	6,432	70	19	12	24	38	74	238	469	756	1,018	968
2001.....	6,254	56	26	21	16	34	113	249	475	789	996	850
2000.....	6,278	54	29	17	28	26	80	297	591	804	942	824
1999.....	6,172	57	27	29	18	27	95	281	590	831	926	746
White, both sexes												
2004.....	11,029	82	36	13	15	30	65	148	420	1,050	1,693	1,940
2003.....	11,669	72	45	19	14	37	66	174	527	1,213	1,801	2,113
2002.....	12,004	116	35	14	25	35	68	222	572	1,394	1,840	2,194
2001.....	11,637	72	41	30	14	27	77	211	678	1,465	1,913	2,018
2000.....	11,807	63	33	25	28	32	64	266	794	1,607	1,855	2,008
1999.....	11,818	83	50	30	23	29	68	327	976	1,673	1,910	1,782

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Viral diseases (A80-B34)												
White, male												
2004.....	7,979	45	21	7	9	12	46	101	326	802	1,339	1,514
2003.....	8,465	31	21	9	6	22	41	124	402	960	1,432	1,683
2002.....	8,696	66	23	5	13	15	45	156	444	1,138	1,451	1,746
2001.....	8,512	38	23	15	8	13	45	145	528	1,181	1,543	1,628
2000.....	8,552	33	16	12	9	20	46	176	619	1,305	1,463	1,626
1999.....	8,587	52	34	15	12	17	42	233	781	1,367	1,543	1,416
White, female												
2004.....	3,050	37	15	6	6	18	19	47	94	248	354	426
2003.....	3,204	41	24	10	8	15	25	50	125	253	369	430
2002.....	3,308	50	12	9	12	20	23	66	128	256	389	448
2001.....	3,125	34	18	15	6	14	32	66	150	284	370	390
2000.....	3,255	30	17	13	19	12	18	90	175	302	392	382
1999.....	3,231	31	16	15	11	12	26	94	195	306	367	366
Black, both sexes												
2004.....	8,283	28	11	8	20	30	126	312	656	1,073	1,579	1,556
2003.....	8,492	33	18	14	19	25	106	303	658	1,334	1,619	1,552
2002.....	8,880	39	15	6	17	31	106	326	827	1,441	1,759	1,664
2001.....	8,802	43	21	13	20	42	137	396	883	1,469	1,788	1,564
2000.....	8,802	45	23	10	17	28	117	423	1,045	1,503	1,702	1,567
1999.....	8,781	44	28	30	21	24	118	462	1,060	1,506	1,835	1,523
Black, male												
2004.....	5,409	15	5	5	9	16	65	163	356	673	998	1,076
2003.....	5,657	16	10	7	11	10	55	165	388	848	1,069	1,072
2002.....	5,939	22	9	5	7	14	55	157	492	952	1,143	1,157
2001.....	5,871	25	15	7	10	23	56	216	560	974	1,183	1,117
2000.....	5,970	22	14	6	9	14	55	216	635	1,011	1,167	1,141
1999.....	6,013	23	17	16	14	10	50	277	670	988	1,294	1,153

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Viral diseases (A80-B34)

Black, female

2004.....	2,874	13	6	3	11	14	61	149	300	400	581	480
2003.....	2,835	17	8	7	8	15	51	138	270	486	550	480
2002.....	2,941	17	6	1	10	17	51	169	335	489	616	507
2001.....	2,931	18	6	6	10	19	81	180	323	495	605	447
2000.....	2,832	23	9	4	8	14	62	207	410	492	535	426
1999.....	2,768	21	11	14	7	14	68	185	390	518	541	370

American Indian 1/, both sexes

2004.....	164	6	3	-	-	-	-	3	7	20	34	34
2003.....	152	3	2	-	2	-	2	1	11	22	24	29
2002.....	133	4	1	-	-	-	1	3	12	19	30	25
2001.....	140	4	3	1	1	1	-	2	13	26	23	24
2000.....	128	1	2	-	1	-	-	1	11	17	24	25
1999.....	117	3	-	-	-	1	1	4	14	21	24	17

American Indian 1/, male

2004.....	105	5	2	-	-	-	-	3	5	16	27	21
2003.....	90	1	1	-	2	-	1	-	10	15	14	12
2002.....	92	3	1	-	-	-	1	2	10	13	25	18
2001.....	93	2	2	1	1	1	-	1	11	18	12	17
2000.....	78	-	-	-	-	-	-	1	6	11	17	19
1999.....	78	-	-	-	-	-	1	3	11	17	14	14

American Indian 1/, female

2004.....	59	1	1	-	-	-	-	-	2	4	7	13
2003.....	62	2	1	-	-	-	1	1	1	7	10	17
2002.....	41	1	-	-	-	-	-	1	2	6	5	7
2001.....	47	2	1	-	-	-	-	1	2	8	11	7
2000.....	50	1	2	-	1	-	-	-	5	6	7	6
1999.....	39	3	-	-	-	1	-	1	3	4	10	3

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Viral diseases (A80-B34)												
Asian or Pacific Islander 2/, both sexes												
2004.....	347	1	4	1	2	4	2	11	11	24	30	30
2003.....	339	8	3	1	-	1	2	6	10	30	24	40
2002.....	391	5	3	3	2	3	1	7	16	19	38	42
2001.....	359	2	1	-	-	3	-	7	14	20	32	33
2000.....	385	2	5	1	3	1	1	5	14	19	35	37
1999.....	364	4	-	-	1	1	2	7	10	25	39	37
Asian or Pacific Islander 2/, male												
2004.....	217	1	1	1	-	1	2	6	8	20	27	23
2003.....	212	3	2	-	-	1	1	4	7	26	21	35
2002.....	249	3	2	1	-	2	1	5	12	14	30	36
2001.....	208	-	-	-	-	2	-	5	14	18	22	27
2000.....	244	2	4	1	3	1	1	5	13	15	27	27
1999.....	230	2	-	-	1	1	1	6	8	22	31	30
Asian or Pacific Islander 2/, female												
2004.....	130	-	3	-	2	3	-	5	3	4	3	7
2003.....	127	5	1	1	-	-	1	2	3	4	3	5
2002.....	142	2	1	2	2	1	-	2	4	5	8	6
2001.....	151	2	1	-	-	1	-	2	-	2	10	6
2000.....	141	-	1	-	-	-	-	-	1	4	8	10
1999.....	134	2	-	-	-	-	1	1	2	3	8	7

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified arthropod-borne viral and hemorrhagic fevers (A90-A94,A96-A99)												
All races, both sexes												
2004.....	81	-	-	-	-	-	-	-	-	1	2	-
2003.....	192	-	-	-	-	1	-	1	1	2	1	1
2002.....	167	-	-	-	-	1	-	1	2	2	2	3
2001.....	3	-	-	-	-	-	-	-	-	1	1	-
2000.....	3	-	-	-	-	-	-	-	-	-	1	-
1999.....	1	-	-	-	-	-	-	1	-	-	-	-
All races, male												
2004.....	51	-	-	-	-	-	-	-	-	1	2	-
2003.....	122	-	-	-	-	1	-	-	1	1	1	1
2002.....	103	-	-	-	-	1	-	1	2	1	-	2
2001.....	1	-	-	-	-	-	-	-	-	-	-	-
2000.....	2	-	-	-	-	-	-	-	-	-	1	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-
All races, female												
2004.....	30	-	-	-	-	-	-	-	-	-	-	-
2003.....	70	-	-	-	-	-	-	1	-	1	-	-
2002.....	64	-	-	-	-	-	-	-	-	1	2	1
2001.....	2	-	-	-	-	-	-	-	-	1	1	-
2000.....	1	-	-	-	-	-	-	-	-	-	-	-
1999.....	1	-	-	-	-	-	-	1	-	-	-	-
White, both sexes												
2004.....	72	-	-	-	-	-	-	-	-	-	-	-
2003.....	173	-	-	-	-	1	-	1	-	2	1	1
2002.....	140	-	-	-	-	1	-	-	2	-	1	3
2001.....	2	-	-	-	-	-	-	-	-	-	1	-
2000.....	2	-	-	-	-	-	-	-	-	-	-	-
1999.....	1	-	-	-	-	-	-	1	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Herpesviral (herpes simplex) infections (B00)

All races, both sexes

2004.....	194	19	3	1	1	2	3	2	1	3	4	7
2003.....	183	22	5	1	1	1	3	1	1	-	6	5
2002.....	177	24	2	1	1	1	3	2	1	3	6	3
2001.....	183	24	1	3	1	1	2	1	3	4	2	5
2000.....	144	24	3	3	-	-	2	1	3	-	1	4
1999.....	145	15	3	-	3	3	-	-	1	2	1	6

All races, male

2004.....	86	9	3	1	-	-	1	-	-	1	2	3
2003.....	76	12	2	-	1	-	2	1	1	-	3	3
2002.....	76	14	1	1	1	-	1	1	-	1	4	2
2001.....	89	12	1	2	1	-	-	-	1	2	-	3
2000.....	60	9	1	-	-	-	-	1	1	-	-	1
1999.....	63	12	2	-	2	1	-	-	1	2	-	2

All races, female

2004.....	108	10	-	-	1	2	2	2	1	2	2	4
2003.....	107	10	3	1	-	1	1	-	-	-	3	2
2002.....	101	10	1	-	-	1	2	1	1	2	2	1
2001.....	94	12	-	1	-	1	2	1	2	2	2	2
2000.....	84	15	2	3	-	-	2	-	2	-	1	3
1999.....	82	3	1	-	1	2	-	-	-	-	1	4

White, both sexes

2004.....	172	12	3	1	-	1	3	2	-	2	4	6
2003.....	161	15	4	1	1	1	3	1	1	-	4	4
2002.....	150	15	2	1	1	1	2	2	-	1	4	-
2001.....	158	16	1	2	1	1	1	1	2	4	-	4
2000.....	117	12	2	3	-	-	1	1	1	-	1	2
1999.....	121	9	3	-	1	3	-	-	1	2	-	4

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Herpesviral (herpes simplex) infections (B00)

White, male

2004.....	75	6	3	1	-	-	1	-	-	1	2	3
2003.....	66	7	2	-	1	-	2	1	1	-	2	2
2002.....	63	10	1	1	1	-	-	1	-	-	2	-
2001.....	75	6	1	1	1	-	-	-	1	2	-	3
2000.....	52	6	1	-	-	-	-	1	1	-	-	1
1999.....	52	9	2	-	-	1	-	-	1	2	-	1

White, female

2004.....	97	6	-	-	-	1	2	2	-	1	2	3
2003.....	95	8	2	1	-	1	1	-	-	-	2	2
2002.....	87	5	1	-	-	1	2	1	-	1	2	-
2001.....	83	10	-	1	-	1	1	1	1	2	-	1
2000.....	65	6	1	3	-	-	1	-	-	-	1	1
1999.....	69	-	1	-	1	2	-	-	-	-	-	3

Black, both sexes

2004.....	16	6	-	-	1	-	-	-	1	1	-	1
2003.....	17	5	1	-	-	-	-	-	-	-	2	-
2002.....	24	8	-	-	-	-	1	-	1	1	2	3
2001.....	22	8	-	1	-	-	1	-	1	-	2	1
2000.....	25	12	1	-	-	-	1	-	2	-	-	2
1999.....	21	5	-	-	2	-	-	-	-	-	1	2

Black, male

2004.....	6	2	-	-	-	-	-	-	-	-	-	-
2003.....	8	5	-	-	-	-	-	-	-	-	1	-
2002.....	12	4	-	-	-	-	1	-	-	-	2	2
2001.....	11	6	-	1	-	-	-	-	-	-	-	-
2000.....	7	3	-	-	-	-	-	-	-	-	-	-
1999.....	11	3	-	-	2	-	-	-	-	-	-	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Zoster (herpes zoster)(B02)												
All races, both sexes												
2004.....	151	-	1	-	-	-	-	-	-	1	-	2
2003.....	157	-	-	-	-	-	1	-	2	-	-	2
2002.....	130	-	1	2	-	-	-	1	-	1	1	-
2001.....	123	1	-	-	-	1	-	-	-	-	-	-
2000.....	131	-	-	-	-	-	1	-	-	2	-	-
1999.....	152	1	-	-	-	-	-	-	2	2	-	1
All races, male												
2004.....	59	-	1	-	-	-	-	-	-	1	-	1
2003.....	50	-	-	-	-	-	-	-	-	-	-	2
2002.....	53	-	1	2	-	-	-	1	-	1	-	-
2001.....	38	1	-	-	-	1	-	-	-	-	-	-
2000.....	43	-	-	-	-	-	1	-	-	1	-	-
1999.....	47	-	-	-	-	-	-	-	2	1	-	1
All races, female												
2004.....	92	-	-	-	-	-	-	-	-	-	-	1
2003.....	107	-	-	-	-	-	1	-	2	-	-	-
2002.....	77	-	-	-	-	-	-	-	-	-	1	-
2001.....	85	-	-	-	-	-	-	-	-	-	-	-
2000.....	88	-	-	-	-	-	-	-	-	1	-	-
1999.....	105	1	-	-	-	-	-	-	-	1	-	-
White, both sexes												
2004.....	140	-	1	-	-	-	-	-	-	1	-	1
2003.....	144	-	-	-	-	-	1	-	1	-	-	1
2002.....	120	-	1	2	-	-	-	1	-	1	1	-
2001.....	117	-	-	-	-	-	-	-	-	-	-	-
2000.....	119	-	-	-	-	-	1	-	-	2	-	-
1999.....	146	1	-	-	-	-	-	-	2	2	-	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Viral hepatitis (B15-B19)												
All races, both sexes												
2004.....	5,382	1	-	-	2	3	8	25	36	140	470	956
2003.....	5,431	1	-	-	-	5	9	21	43	174	478	1,053
2002.....	5,793	2	3	-	2	12	11	37	57	216	524	1,192
2001.....	5,585	1	-	3	-	9	11	19	55	202	595	1,101
2000.....	5,357	6	1	2	-	4	14	25	54	219	560	1,044
1999.....	4,853	2	3	1	2	3	11	19	69	204	518	871
All races, male												
2004.....	3,513	-	-	-	1	1	6	10	23	83	316	701
2003.....	3,546	-	-	-	-	2	5	12	30	126	335	763
2002.....	3,770	1	3	-	1	4	9	20	43	154	365	887
2001.....	3,539	1	-	2	-	6	5	7	35	133	420	833
2000.....	3,381	2	1	1	-	2	9	13	33	151	389	787
1999.....	3,043	2	2	-	2	3	7	9	44	139	381	651
All races, female												
2004.....	1,869	1	-	-	1	2	2	15	13	57	154	255
2003.....	1,885	1	-	-	-	3	4	9	13	48	143	290
2002.....	2,023	1	-	-	1	8	2	17	14	62	159	305
2001.....	2,046	-	-	1	-	3	6	12	20	69	175	268
2000.....	1,976	4	-	1	-	2	5	12	21	68	171	257
1999.....	1,810	-	1	1	-	-	4	10	25	65	137	220
White, both sexes												
2004.....	4,246	1	-	-	1	2	3	13	27	109	391	787
2003.....	4,295	-	-	-	-	5	4	17	32	146	385	878
2002.....	4,620	1	-	-	2	9	11	21	44	178	422	961
2001.....	4,451	1	-	3	-	6	7	12	41	166	480	898
2000.....	4,216	3	-	2	-	2	6	17	34	180	445	827
1999.....	3,835	1	3	1	1	3	6	11	57	169	402	683

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Viral hepatitis (B15-B19)												
White, male												
2004.....	2,799	-	-	-	1	1	2	6	17	67	271	579
2003.....	2,848	-	-	-	-	2	1	10	20	104	275	648
2002.....	3,049	1	-	-	1	2	9	12	34	133	296	727
2001.....	2,900	1	-	2	-	4	2	5	26	111	345	691
2000.....	2,686	1	-	1	-	2	4	9	21	123	316	636
1999.....	2,436	1	2	-	1	3	4	4	37	118	301	501
White, female												
2004.....	1,447	1	-	-	-	1	1	7	10	42	120	208
2003.....	1,447	-	-	-	-	3	3	7	12	42	110	230
2002.....	1,571	-	-	-	1	7	2	9	10	45	126	234
2001.....	1,551	-	-	1	-	2	5	7	15	55	135	207
2000.....	1,530	2	-	1	-	-	2	8	13	57	129	191
1999.....	1,399	-	1	1	-	-	2	7	20	51	101	182
Black, both sexes												
2004.....	863	-	-	-	1	-	4	8	6	23	59	140
2003.....	854	-	-	-	-	-	4	2	9	20	78	138
2002.....	865	1	1	-	-	2	-	15	7	31	81	185
2001.....	831	-	-	-	-	2	4	7	8	27	94	167
2000.....	793	3	-	-	-	2	8	7	14	30	88	172
1999.....	731	1	-	-	1	-	4	6	11	25	96	161
Black, male												
2004.....	564	-	-	-	-	-	3	2	4	9	31	104
2003.....	544	-	-	-	-	-	3	1	8	15	49	93
2002.....	541	-	1	-	-	1	-	7	4	16	52	125
2001.....	484	-	-	-	-	1	3	2	5	15	64	115
2000.....	501	1	-	-	-	-	5	3	8	22	56	121
1999.....	443	1	-	-	1	-	2	3	6	15	66	126

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Viral hepatitis (B15-B19)

Black, female

2004.....	299	-	-	-	1	-	1	6	2	14	28	36
2003.....	310	-	-	-	-	-	1	1	1	5	29	45
2002.....	324	1	-	-	-	1	-	8	3	15	29	60
2001.....	347	-	-	-	-	1	1	5	3	12	30	52
2000.....	292	2	-	-	-	2	3	4	6	8	32	51
1999.....	288	-	-	-	-	-	2	3	5	10	30	35

American Indian 1/, both sexes

2004.....	62	-	-	-	-	-	-	-	-	3	10	18
2003.....	62	-	-	-	-	-	1	1	2	2	7	17
2002.....	56	-	-	-	-	-	-	-	2	3	9	17
2001.....	55	-	-	-	-	1	-	-	4	4	9	10
2000.....	63	-	-	-	-	-	-	-	4	3	12	12
1999.....	37	-	-	-	-	-	1	-	1	2	6	8

American Indian 1/, male

2004.....	31	-	-	-	-	-	-	-	-	2	6	10
2003.....	32	-	-	-	-	-	1	-	2	1	5	4
2002.....	36	-	-	-	-	-	-	-	2	1	7	11
2001.....	30	-	-	-	-	1	-	-	2	3	3	6
2000.....	33	-	-	-	-	-	-	-	2	2	6	7
1999.....	22	-	-	-	-	-	1	-	1	-	3	8

American Indian 1/, female

2004.....	31	-	-	-	-	-	-	-	-	1	4	8
2003.....	30	-	-	-	-	-	-	1	-	1	2	13
2002.....	20	-	-	-	-	-	-	-	-	2	2	6
2001.....	25	-	-	-	-	-	-	-	2	1	6	4
2000.....	30	-	-	-	-	-	-	-	2	1	6	5
1999.....	15	-	-	-	-	-	-	-	-	2	3	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Viral hepatitis (B15-B19)

Asian or Pacific Islander 2/, both sexes

2004.....	211	-	-	-	-	1	1	4	3	5	10	11
2003.....	220	1	-	-	-	-	-	1	-	6	8	20
2002.....	252	-	2	-	-	1	-	1	4	4	12	29
2001.....	248	-	-	-	-	-	-	-	2	5	12	26
2000.....	285	-	1	-	-	-	-	1	2	6	15	33
1999.....	250	-	-	-	-	-	-	2	-	8	14	19

Asian or Pacific Islander 2/, male

2004.....	119	-	-	-	-	-	1	2	2	5	8	8
2003.....	122	-	-	-	-	-	-	1	-	6	6	18
2002.....	144	-	2	-	-	1	-	1	3	4	10	24
2001.....	125	-	-	-	-	-	-	-	2	4	8	21
2000.....	161	-	1	-	-	-	-	1	2	4	11	23
1999.....	142	-	-	-	-	-	-	2	-	6	11	16

Asian or Pacific Islander 2/, female

2004.....	92	-	-	-	-	1	-	2	1	-	2	3
2003.....	98	1	-	-	-	-	-	-	-	-	2	2
2002.....	108	-	-	-	-	-	-	-	1	-	2	5
2001.....	123	-	-	-	-	-	-	-	-	1	4	5
2000.....	124	-	-	-	-	-	-	-	-	2	4	10
1999.....	108	-	-	-	-	-	-	-	-	2	3	3

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Human immunodeficiency virus (HIV) disease (B20-B24)												
All races, both sexes												
2004.....	13,063	3	4	8	17	31	160	432	1,036	1,995	2,831	2,554
2003.....	13,658	5	2	15	21	32	146	444	1,144	2,401	2,939	2,636
2002.....	14,095	7	5	4	21	38	140	500	1,339	2,618	3,089	2,680
2001.....	14,175	10	13	17	21	42	183	584	1,517	2,741	3,126	2,511
2000.....	14,478	10	14	17	29	35	144	647	1,790	2,896	3,023	2,551
1999.....	14,802	14	24	38	28	31	167	763	1,966	2,981	3,251	2,440
All races, male												
2004.....	9,506	1	1	6	6	14	93	259	658	1,411	2,055	1,911
2003.....	10,119	2	2	7	12	15	80	272	766	1,710	2,174	2,013
2002.....	10,468	4	1	4	10	17	76	288	899	1,943	2,255	2,039
2001.....	10,590	7	8	11	12	21	84	354	1,067	2,042	2,318	1,942
2000.....	10,868	8	6	9	13	19	80	371	1,231	2,172	2,269	2,008
1999.....	11,234	9	16	20	16	15	80	504	1,410	2,224	2,482	1,936
All races, female												
2004.....	3,557	2	3	2	11	17	67	173	378	584	776	643
2003.....	3,539	3	-	8	9	17	66	172	378	691	765	623
2002.....	3,627	3	4	-	11	21	64	212	440	675	834	641
2001.....	3,585	3	5	6	9	21	99	230	450	699	808	569
2000.....	3,610	2	8	8	16	16	64	276	559	724	754	543
1999.....	3,568	5	8	18	12	16	87	259	556	757	769	504
White, both sexes												
2004.....	5,608	2	2	2	4	7	41	126	377	917	1,279	1,113
2003.....	6,019	1	1	2	3	12	45	139	480	1,049	1,383	1,199
2002.....	6,089	4	-	-	6	12	40	184	504	1,192	1,378	1,193
2001.....	6,171	3	3	7	4	6	54	191	623	1,275	1,406	1,097
2000.....	6,498	1	1	8	13	13	40	235	748	1,402	1,383	1,150
1999.....	6,752	4	7	12	13	10	53	304	898	1,472	1,479	1,065

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Human immunodeficiency virus (HIV) disease (B20-B24)												
White, male												
2004.....	4,593	1	1	1	1	1	32	94	298	722	1,055	918
2003.....	4,964	1	1	-	2	6	29	106	373	846	1,140	1,014
2002.....	5,028	3	-	-	4	5	26	135	397	990	1,133	997
2001.....	5,135	2	-	5	2	2	33	136	492	1,059	1,179	924
2000.....	5,360	1	-	4	5	8	30	158	591	1,165	1,135	977
1999.....	5,613	3	5	6	7	7	33	226	730	1,222	1,227	897
White, female												
2004.....	1,015	1	1	1	3	6	9	32	79	195	224	195
2003.....	1,055	-	-	2	1	6	16	33	107	203	243	185
2002.....	1,061	1	-	-	2	7	14	49	107	202	245	196
2001.....	1,036	1	3	2	2	4	21	55	131	216	227	173
2000.....	1,138	-	1	4	8	5	10	77	157	237	248	173
1999.....	1,139	1	2	6	6	3	20	78	168	250	252	168
Black, both sexes												
2004.....	7,271	1	2	5	13	24	118	299	645	1,042	1,512	1,407
2003.....	7,479	3	1	13	18	20	99	301	647	1,309	1,523	1,407
2002.....	7,835	3	5	4	15	25	99	309	814	1,397	1,666	1,466
2001.....	7,844	7	10	9	17	34	129	384	874	1,432	1,687	1,393
2000.....	7,848	9	12	9	14	22	103	409	1,023	1,470	1,610	1,385
1999.....	7,893	10	17	26	15	21	112	452	1,045	1,473	1,733	1,350
Black, male												
2004.....	4,769	-	-	4	5	13	60	159	349	660	964	967
2003.....	5,033	1	1	7	10	9	50	164	379	830	1,010	975
2002.....	5,301	1	1	4	6	11	49	148	486	932	1,085	1,023
2001.....	5,328	5	8	5	10	18	51	212	555	955	1,116	1,001
2000.....	5,400	7	6	5	7	11	49	210	625	988	1,108	1,016
1999.....	5,493	6	11	14	9	8	46	273	662	969	1,225	1,019

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Human immunodeficiency virus (HIV) disease (B20-B24)												
Black, female												
2004.....	2,502	1	2	1	8	11	58	140	296	382	548	440
2003.....	2,446	2	-	6	8	11	49	137	268	479	513	432
2002.....	2,534	2	4	-	9	14	50	161	328	465	581	443
2001.....	2,516	2	2	4	7	16	78	172	319	477	571	392
2000.....	2,448	2	6	4	7	11	54	199	398	482	502	369
1999.....	2,400	4	6	12	6	13	66	179	383	504	508	331
American Indian 1/, both sexes												
2004.....	84	-	-	-	-	-	-	3	6	17	22	16
2003.....	70	-	-	-	-	-	-	-	8	20	17	12
2002.....	64	-	-	-	-	-	1	3	10	16	20	8
2001.....	74	-	-	1	-	-	-	2	9	21	14	14
2000.....	57	-	1	-	1	-	-	1	7	12	12	12
1999.....	68	-	-	-	-	-	-	4	13	19	15	8
American Indian 1/, male												
2004.....	63	-	-	-	-	-	-	3	5	14	19	11
2003.....	49	-	-	-	-	-	-	-	7	14	9	8
2002.....	50	-	-	-	-	-	1	2	8	12	17	7
2001.....	57	-	-	1	-	-	-	1	9	15	9	11
2000.....	43	-	-	-	-	-	-	1	4	9	11	11
1999.....	54	-	-	-	-	-	-	3	10	17	10	6
American Indian 1/, female												
2004.....	21	-	-	-	-	-	-	-	1	3	3	5
2003.....	21	-	-	-	-	-	-	-	1	6	8	4
2002.....	14	-	-	-	-	-	-	1	2	4	3	1
2001.....	17	-	-	-	-	-	-	1	-	6	5	3
2000.....	14	-	1	-	1	-	-	-	3	3	1	1
1999.....	14	-	-	-	-	-	-	1	3	2	5	2

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Human immunodeficiency virus (HIV) disease (B20-B24)												
Asian or Pacific Islander 2/, both sexes												
2004.....	100	-	-	1	-	-	1	4	8	19	18	18
2003.....	90	1	-	-	-	-	2	4	9	23	16	18
2002.....	107	-	-	-	-	1	-	4	11	13	25	13
2001.....	86	-	-	-	-	2	-	7	11	13	19	7
2000.....	75	-	-	-	1	-	1	2	12	12	18	4
1999.....	89	-	-	-	-	-	2	3	10	17	24	17
Asian or Pacific Islander 2/, male												
2004.....	81	-	-	1	-	-	1	3	6	15	17	15
2003.....	73	-	-	-	-	-	1	2	7	20	15	16
2002.....	89	-	-	-	-	1	-	3	8	9	20	12
2001.....	70	-	-	-	-	1	-	5	11	13	14	6
2000.....	65	-	-	-	1	-	1	2	11	10	15	4
1999.....	74	-	-	-	-	-	1	2	8	16	20	14
Asian or Pacific Islander 2/, female												
2004.....	19	-	-	-	-	-	-	1	2	4	1	3
2003.....	17	1	-	-	-	-	1	2	2	3	1	2
2002.....	18	-	-	-	-	-	-	1	3	4	5	1
2001.....	16	-	-	-	-	1	-	2	-	-	5	1
2000.....	10	-	-	-	-	-	-	-	1	2	3	-
1999.....	15	-	-	-	-	-	1	1	2	1	4	3

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Human immunodeficiency virus (HIV) disease resulting in infectious and parasitic diseases (B20)

All races, both sexes

2004.....	5,138	1	1	1	5	10	69	182	434	778	1,154	1,027
2003.....	5,222	1	-	5	9	15	53	184	457	915	1,095	1,019
2002.....	5,603	2	2	2	5	15	66	206	533	1,075	1,215	1,080
2001.....	5,553	6	9	6	7	19	78	245	602	1,070	1,214	1,020
2000.....	5,722	4	7	8	6	15	70	274	700	1,179	1,208	1,038
1999.....	5,931	7	13	12	12	11	78	332	788	1,228	1,279	1,020

All races, male

2004.....	3,695	-	1	1	3	3	38	105	275	553	842	755
2003.....	3,874	-	-	3	6	9	30	112	317	634	815	801
2002.....	4,132	1	1	2	4	7	42	114	351	774	897	823
2001.....	4,106	4	5	4	2	10	44	150	410	788	908	777
2000.....	4,256	3	2	2	2	8	37	165	488	878	881	812
1999.....	4,456	5	11	7	7	5	35	216	546	912	980	796

All races, female

2004.....	1,443	1	-	-	2	7	31	77	159	225	312	272
2003.....	1,348	1	-	2	3	6	23	72	140	281	280	218
2002.....	1,471	1	1	-	1	8	24	92	182	301	318	257
2001.....	1,447	2	4	2	5	9	34	95	192	282	306	243
2000.....	1,466	1	5	6	4	7	33	109	212	301	327	226
1999.....	1,475	2	2	5	5	6	43	116	242	316	299	224

White, both sexes

2004.....	2,277	-	1	-	-	2	13	62	157	382	545	476
2003.....	2,427	-	-	-	1	7	19	64	199	426	562	503
2002.....	2,510	-	-	-	3	6	22	81	202	520	567	484
2001.....	2,525	2	3	3	1	4	25	93	260	524	572	467
2000.....	2,628	-	-	4	1	6	24	109	300	573	563	477
1999.....	2,729	1	3	1	7	3	21	142	358	612	587	434

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Human immunodeficiency virus (HIV) disease resulting in infectious and parasitic diseases (B20)

White, male

2004.....	1,841	-	1	-	-	-	11	47	121	298	449	384
2003.....	1,962	-	-	-	1	4	14	47	158	331	458	421
2002.....	2,011	-	-	-	2	4	17	59	145	411	454	401
2001.....	2,051	1	-	2	-	2	16	70	201	415	473	392
2000.....	2,135	-	-	1	-	4	17	73	239	462	453	402
1999.....	2,225	1	3	-	3	3	13	107	279	495	480	361

White, female

2004.....	436	-	-	-	-	2	2	15	36	84	96	92
2003.....	465	-	-	-	-	3	5	17	41	95	104	82
2002.....	499	-	-	-	1	2	5	22	57	109	113	83
2001.....	474	1	3	1	1	2	9	23	59	109	99	75
2000.....	493	-	-	3	1	2	7	36	61	111	110	75
1999.....	504	-	-	1	4	-	8	35	79	117	107	73

Black, both sexes

2004.....	2,779	1	-	1	5	8	56	114	271	377	594	535
2003.....	2,715	-	-	5	8	8	34	118	248	468	518	500
2002.....	3,012	2	2	2	2	8	43	119	319	540	629	587
2001.....	2,963	4	6	3	6	15	53	149	333	532	630	547
2000.....	3,034	4	7	4	5	9	46	164	392	595	629	555
1999.....	3,125	6	10	11	5	8	57	187	418	593	675	574

Black, male

2004.....	1,786	-	-	1	3	3	27	53	149	239	380	358
2003.....	1,848	-	-	3	5	5	16	63	151	288	345	366
2002.....	2,053	1	1	2	2	2	24	50	196	352	427	413
2001.....	2,001	3	5	2	2	8	28	77	200	361	425	381
2000.....	2,074	3	2	1	2	4	20	91	244	409	414	405
1999.....	2,169	4	8	7	4	2	22	107	258	395	487	427

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Human immunodeficiency virus (HIV) disease resulting in infectious and parasitic diseases (B20)

Black, female

2004.....	993	1	-	-	2	5	29	61	122	138	214	177
2003.....	867	-	-	2	3	3	18	55	97	180	173	134
2002.....	959	1	1	-	-	6	19	69	123	188	202	174
2001.....	962	1	1	1	4	7	25	72	133	171	205	166
2000.....	960	1	5	3	3	5	26	73	148	186	215	150
1999.....	956	2	2	4	1	6	35	80	160	198	188	147

American Indian 1/, both sexes

2004.....	40	-	-	-	-	-	-	3	3	9	10	7
2003.....	35	-	-	-	-	-	-	-	4	11	6	6
2002.....	29	-	-	-	-	-	1	3	4	9	6	3
2001.....	27	-	-	-	-	-	-	-	3	8	7	3
2000.....	30	-	-	-	-	-	-	-	4	6	8	5
1999.....	34	-	-	-	-	-	-	1	6	12	5	5

American Indian 1/, male

2004.....	34	-	-	-	-	-	-	3	3	8	9	5
2003.....	25	-	-	-	-	-	-	-	3	7	3	5
2002.....	24	-	-	-	-	-	1	2	3	7	6	3
2001.....	23	-	-	-	-	-	-	-	3	6	6	2
2000.....	23	-	-	-	-	-	-	-	2	4	8	4
1999.....	27	-	-	-	-	-	-	1	4	11	3	3

American Indian 1/, female

2004.....	6	-	-	-	-	-	-	-	-	1	1	2
2003.....	10	-	-	-	-	-	-	-	1	4	3	1
2002.....	5	-	-	-	-	-	-	1	1	2	-	-
2001.....	4	-	-	-	-	-	-	-	-	2	1	1
2000.....	7	-	-	-	-	-	-	-	2	2	-	1
1999.....	7	-	-	-	-	-	-	-	2	1	2	2

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Human immunodeficiency virus (HIV) disease resulting in infectious and parasitic diseases (B20)

Asian or Pacific Islander 2/, both sexes

2004.....	42	-	-	-	-	-	-	3	3	10	5	9
2003.....	45	1	-	-	-	-	-	2	6	10	9	10
2002.....	52	-	-	-	-	1	-	3	8	6	13	6
2001.....	38	-	-	-	-	-	-	3	6	6	5	3
2000.....	30	-	-	-	-	-	-	1	4	5	8	1
1999.....	43	-	-	-	-	-	-	2	6	11	12	7

Asian or Pacific Islander 2/, male

2004.....	34	-	-	-	-	-	-	2	2	8	4	8
2003.....	39	-	-	-	-	-	-	2	5	8	9	9
2002.....	44	-	-	-	-	1	-	3	7	4	10	6
2001.....	31	-	-	-	-	-	-	3	6	6	4	2
2000.....	24	-	-	-	-	-	-	1	3	3	6	1
1999.....	35	-	-	-	-	-	-	1	5	11	10	5

Asian or Pacific Islander 2/, female

2004.....	8	-	-	-	-	-	-	1	1	2	1	1
2003.....	6	1	-	-	-	-	-	-	1	2	-	1
2002.....	8	-	-	-	-	-	-	-	1	2	3	-
2001.....	7	-	-	-	-	-	-	-	-	-	1	1
2000.....	6	-	-	-	-	-	-	-	1	2	2	-
1999.....	8	-	-	-	-	-	-	1	1	-	2	2

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Human immunodeficiency virus (HIV) disease resulting in malignant neoplasms (B21)

All races, both sexes

2004.....	916	-	-	3	1	4	12	19	62	138	171	176
2003.....	895	-	-	-	-	-	9	22	59	142	200	168
2002.....	920	-	-	-	2	2	12	21	66	155	204	189
2001.....	962	-	-	-	-	2	5	33	94	164	226	159
2000.....	1,016	1	-	-	3	-	10	33	107	197	219	186
1999.....	1,117	-	-	1	-	1	8	43	164	228	235	169

All races, male

2004.....	761	-	-	3	1	3	6	15	47	116	139	149
2003.....	717	-	-	-	-	-	7	19	43	122	167	124
2002.....	765	-	-	-	-	1	7	14	55	132	169	161
2001.....	797	-	-	-	-	1	1	20	74	146	193	127
2000.....	863	1	-	-	1	-	9	21	86	170	187	158
1999.....	948	-	-	-	-	1	6	33	139	193	202	148

All races, female

2004.....	155	-	-	-	-	1	6	4	15	22	32	27
2003.....	178	-	-	-	-	-	2	3	16	20	33	44
2002.....	155	-	-	-	2	1	5	7	11	23	35	28
2001.....	165	-	-	-	-	1	4	13	20	18	33	32
2000.....	153	-	-	-	2	-	1	12	21	27	32	28
1999.....	169	-	-	1	-	-	2	10	25	35	33	21

White, both sexes

2004.....	508	-	-	1	-	1	4	11	31	84	93	108
2003.....	510	-	-	-	-	-	3	8	33	84	123	99
2002.....	507	-	-	-	-	-	3	10	38	94	106	109
2001.....	540	-	-	-	-	-	2	14	48	103	130	91
2000.....	586	-	-	-	-	-	4	12	65	120	121	113
1999.....	675	-	-	1	-	-	5	19	104	152	136	102

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Human immunodeficiency virus (HIV) disease resulting in malignant neoplasms (B21)

White, male

2004.....	456	-	-	1	-	-	1	9	27	74	84	102
2003.....	450	-	-	-	-	-	2	7	27	78	112	85
2002.....	466	-	-	-	-	-	1	10	36	86	99	103
2001.....	480	-	-	-	-	-	-	9	38	99	118	80
2000.....	530	-	-	-	-	-	4	8	56	109	112	101
1999.....	604	-	-	-	-	-	3	15	96	137	121	91

White, female

2004.....	52	-	-	-	-	1	3	2	4	10	9	6
2003.....	60	-	-	-	-	-	1	1	6	6	11	14
2002.....	41	-	-	-	-	-	2	-	2	8	7	6
2001.....	60	-	-	-	-	-	2	5	10	4	12	11
2000.....	56	-	-	-	-	-	-	4	9	11	9	12
1999.....	71	-	-	1	-	-	2	4	8	15	15	11

Black, both sexes

2004.....	393	-	-	1	1	3	8	8	28	53	76	65
2003.....	374	-	-	-	-	-	6	14	25	54	73	68
2002.....	402	-	-	-	2	2	9	11	28	61	91	80
2001.....	413	-	-	-	-	2	3	17	45	61	92	67
2000.....	422	1	-	-	2	-	5	20	41	76	98	72
1999.....	429	-	-	-	-	1	3	22	59	74	95	64

Black, male

2004.....	293	-	-	1	1	3	5	6	17	41	53	45
2003.....	258	-	-	-	-	-	5	12	15	40	52	38
2002.....	289	-	-	-	-	1	6	4	19	46	64	58
2001.....	309	-	-	-	-	1	1	10	35	47	71	46
2000.....	326	1	-	-	-	-	4	12	29	60	75	56
1999.....	333	-	-	-	-	1	3	17	42	54	78	54

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Human immunodeficiency virus (HIV) disease resulting in malignant neoplasms (B21)												
Black, female												
2004.....	100	-	-	-	-	-	3	2	11	12	23	20
2003.....	116	-	-	-	-	-	1	2	10	14	21	30
2002.....	113	-	-	-	2	1	3	7	9	15	27	22
2001.....	104	-	-	-	-	1	2	7	10	14	21	21
2000.....	96	-	-	-	2	-	1	8	12	16	23	16
1999.....	96	-	-	-	-	-	-	5	17	20	17	10
American Indian 1/, both sexes												
2004.....	5	-	-	-	-	-	-	-	-	1	1	1
2003.....	4	-	-	-	-	-	-	-	-	2	1	1
2002.....	4	-	-	-	-	-	-	-	-	-	4	-
2001.....	2	-	-	-	-	-	-	-	-	-	1	1
2000.....	1	-	-	-	-	-	-	-	-	-	-	1
1999.....	9	-	-	-	-	-	-	2	1	2	2	1
American Indian 1/, male												
2004.....	4	-	-	-	-	-	-	-	-	1	1	1
2003.....	4	-	-	-	-	-	-	-	-	2	1	1
2002.....	3	-	-	-	-	-	-	-	-	-	3	-
2001.....	2	-	-	-	-	-	-	-	-	-	1	1
2000.....	1	-	-	-	-	-	-	-	-	-	-	1
1999.....	7	-	-	-	-	-	-	1	1	2	1	1
American Indian 1/, female												
2004.....	1	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	1	-	-	-	-	-	-	-	-	-	1	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	2	-	-	-	-	-	-	1	-	-	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Human immunodeficiency virus (HIV) disease resulting in other specified diseases (B22)												
All races, both sexes												
2004.....	1,299	-	-	1	4	5	13	42	94	181	276	267
2003.....	1,319	1	1	1	-	3	13	40	96	249	293	257
2002.....	1,348	-	-	-	-	2	8	58	153	221	314	236
2001.....	1,418	-	1	5	4	3	14	58	164	298	274	230
2000.....	1,609	-	-	4	4	2	12	66	222	330	338	262
1999.....	1,608	2	2	4	5	2	14	76	217	354	360	242
All races, male												
2004.....	981	-	-	1	-	2	10	28	58	137	208	207
2003.....	1,052	1	1	1	-	2	9	29	65	188	237	207
2002.....	1,066	-	-	-	-	-	4	34	116	181	245	187
2001.....	1,111	-	1	3	1	1	8	42	121	242	213	189
2000.....	1,274	-	-	3	2	1	6	43	165	255	275	216
1999.....	1,289	1	1	4	3	1	7	55	168	281	290	202
All races, female												
2004.....	318	-	-	-	4	3	3	14	36	44	68	60
2003.....	267	-	-	-	-	1	4	11	31	61	56	50
2002.....	282	-	-	-	-	2	4	24	37	40	69	49
2001.....	307	-	-	2	3	2	6	16	43	56	61	41
2000.....	335	-	-	1	2	1	6	23	57	75	63	46
1999.....	319	1	1	-	2	1	7	21	49	73	70	40
White, both sexes												
2004.....	640	-	-	-	2	1	8	10	44	93	137	140
2003.....	664	-	-	-	-	2	5	13	48	140	144	133
2002.....	691	-	-	-	-	2	2	20	75	113	165	136
2001.....	684	-	-	2	-	1	2	22	78	154	126	118
2000.....	806	-	-	2	3	1	1	24	109	185	177	124
1999.....	837	-	1	1	2	2	6	35	110	200	177	144

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Human immunodeficiency virus (HIV) disease resulting in other specified diseases (B22)												
White, male												
2004.....	536	-	-	-	-	-	6	9	36	82	115	117
2003.....	578	-	-	-	-	1	4	13	35	118	123	119
2002.....	609	-	-	-	-	-	2	17	67	103	145	119
2001.....	602	-	-	1	-	-	2	19	66	138	109	106
2000.....	703	-	-	2	1	1	1	20	92	164	153	112
1999.....	730	-	1	1	1	1	4	27	94	176	155	127
White, female												
2004.....	104	-	-	-	2	1	2	1	8	11	22	23
2003.....	86	-	-	-	-	1	1	-	13	22	21	14
2002.....	82	-	-	-	-	2	-	3	8	10	20	17
2001.....	82	-	-	1	-	1	-	3	12	16	17	12
2000.....	103	-	-	-	2	-	-	4	17	21	24	12
1999.....	107	-	-	-	1	1	2	8	16	24	22	17
Black, both sexes												
2004.....	643	-	-	1	2	4	5	32	49	86	134	124
2003.....	642	1	1	1	-	1	8	27	46	104	146	124
2002.....	639	-	-	-	-	-	6	37	74	103	146	98
2001.....	715	-	1	3	4	2	12	35	85	139	145	108
2000.....	785	-	-	2	1	1	11	42	110	140	157	135
1999.....	757	2	1	3	3	-	8	40	106	151	178	97
Black, male												
2004.....	431	-	-	1	-	2	4	19	21	53	89	87
2003.....	462	1	1	1	-	1	5	16	28	65	112	88
2002.....	445	-	-	-	-	-	2	17	46	75	98	66
2001.....	494	-	1	2	1	1	6	22	54	100	102	80
2000.....	554	-	-	1	1	-	5	23	70	86	119	101
1999.....	546	1	-	3	2	-	3	27	74	102	130	74

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Human immunodeficiency virus (HIV) disease resulting in other specified diseases (B22)												
Asian or Pacific Islander 2/, both sexes												
2004.....	8	-	-	-	-	-	-	-	-	2	2	1
2003.....	5	-	-	-	-	-	-	-	-	3	-	-
2002.....	9	-	-	-	-	-	-	1	2	2	-	1
2001.....	10	-	-	-	-	-	-	1	1	2	2	1
2000.....	11	-	-	-	-	-	-	-	3	1	4	1
1999.....	12	-	-	-	-	-	-	-	1	3	4	1
Asian or Pacific Islander 2/, male												
2004.....	7	-	-	-	-	-	-	-	-	2	2	1
2003.....	5	-	-	-	-	-	-	-	-	3	-	-
2002.....	6	-	-	-	-	-	-	-	1	2	-	1
2001.....	9	-	-	-	-	-	-	1	1	2	1	1
2000.....	10	-	-	-	-	-	-	-	3	1	3	1
1999.....	11	-	-	-	-	-	-	-	-	3	4	1
Asian or Pacific Islander 2/, female												
2004.....	1	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	3	-	-	-	-	-	-	1	1	-	-	-
2001.....	1	-	-	-	-	-	-	-	-	-	1	-
2000.....	1	-	-	-	-	-	-	-	-	-	1	-
1999.....	1	-	-	-	-	-	-	-	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Human immunodeficiency virus (HIV) disease resulting in other conditions (B23)

All races, both sexes

2004.....	2,169	1	2	-	4	7	19	65	154	303	457	402
2003.....	3,265	1	1	6	9	10	36	96	246	519	700	662
2002.....	3,414	1	3	2	10	13	22	106	294	579	751	661
2001.....	3,471	2	1	5	8	7	53	143	339	655	756	621
2000.....	3,388	3	4	4	13	11	25	143	391	651	674	574
1999.....	3,433	2	6	13	9	14	39	164	407	636	734	562

All races, male

2004.....	1,562	1	-	-	1	4	11	36	105	205	318	304
2003.....	2,307	-	1	3	5	3	16	46	153	354	489	477
2002.....	2,449	-	-	2	4	7	10	61	196	430	502	479
2001.....	2,486	2	1	4	7	3	18	76	234	469	516	457
2000.....	2,423	3	3	4	7	7	9	73	246	467	485	434
1999.....	2,490	2	3	7	5	6	19	108	277	443	519	434

All races, female

2004.....	607	-	2	-	3	3	8	29	49	98	139	98
2003.....	958	1	-	3	4	7	20	50	93	165	211	185
2002.....	965	1	3	-	6	6	12	45	98	149	249	182
2001.....	985	-	-	1	1	4	35	67	105	186	240	164
2000.....	965	-	1	-	6	4	16	70	145	184	189	140
1999.....	943	-	3	6	4	8	20	56	130	193	215	128

White, both sexes

2004.....	773	1	1	-	1	1	4	15	46	105	174	132
2003.....	1,202	-	1	1	2	2	5	21	79	185	279	241
2002.....	1,233	1	-	-	1	2	6	28	91	234	286	229
2001.....	1,311	-	-	2	2	1	17	32	122	272	296	232
2000.....	1,335	1	1	1	7	4	6	39	137	285	276	222
1999.....	1,299	1	3	7	3	4	13	52	149	259	282	201

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Human immunodeficiency virus (HIV) disease resulting in other conditions (B23)

White, male

2004.....	626	1	-	-	1	-	3	11	37	79	138	105
2003.....	970	-	1	-	1	1	2	14	58	146	220	200
2002.....	993	-	-	-	1	-	3	19	72	193	223	178
2001.....	1,042	-	-	2	1	-	10	17	92	215	234	183
2000.....	1,046	1	-	1	3	3	4	21	98	232	215	179
1999.....	1,039	1	1	4	2	2	9	38	117	206	221	162

White, female

2004.....	147	-	1	-	-	1	1	4	9	26	36	27
2003.....	232	-	-	1	1	1	3	7	21	39	59	41
2002.....	240	1	-	-	-	2	3	9	19	41	63	51
2001.....	269	-	-	-	1	1	7	15	30	57	62	49
2000.....	289	-	1	-	4	1	2	18	39	53	61	43
1999.....	260	-	2	3	1	2	4	14	32	53	61	39

Black, both sexes

2004.....	1,365	-	1	-	3	6	14	50	108	192	276	264
2003.....	2,035	1	-	5	7	8	30	74	167	326	416	414
2002.....	2,152	-	3	2	9	11	16	78	199	340	458	428
2001.....	2,118	2	1	2	6	5	36	109	211	374	452	383
2000.....	2,023	2	3	3	5	7	19	104	250	362	392	348
1999.....	2,109	1	3	6	6	10	24	112	254	374	448	356

Black, male

2004.....	915	-	-	-	-	4	7	25	68	122	173	195
2003.....	1,319	-	-	3	4	2	14	32	95	203	265	272
2002.....	1,434	-	-	2	3	7	7	42	122	232	274	299
2001.....	1,414	2	1	1	6	3	8	59	136	247	278	269
2000.....	1,352	2	3	3	4	4	5	52	144	231	265	251
1999.....	1,433	1	2	3	3	4	9	70	156	236	295	268

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Human immunodeficiency virus (HIV) disease resulting in other conditions (B23)												
Black, female												
2004.....	450	-	1	-	3	2	7	25	40	70	103	69
2003.....	716	1	-	2	3	6	16	42	72	123	151	142
2002.....	718	-	3	-	6	4	9	36	77	108	184	129
2001.....	704	-	-	1	-	2	28	50	75	127	174	114
2000.....	671	-	-	-	1	3	14	52	106	131	127	97
1999.....	676	-	1	3	3	6	15	42	98	138	153	88
American Indian 1/, both sexes												
2004.....	16	-	-	-	-	-	-	-	-	5	3	3
2003.....	9	-	-	-	-	-	-	-	-	3	3	2
2002.....	11	-	-	-	-	-	-	-	3	3	4	1
2001.....	21	-	-	1	-	-	-	1	4	6	3	4
2000.....	10	-	-	-	1	-	-	-	1	-	4	2
1999.....	9	-	-	-	-	-	-	-	2	1	2	1
American Indian 1/, male												
2004.....	8	-	-	-	-	-	-	-	-	3	3	1
2003.....	4	-	-	-	-	-	-	-	-	1	2	-
2002.....	8	-	-	-	-	-	-	-	2	3	3	-
2001.....	14	-	-	1	-	-	-	-	4	4	-	3
2000.....	7	-	-	-	-	-	-	-	1	-	3	2
1999.....	6	-	-	-	-	-	-	-	2	-	1	1
American Indian 1/, female												
2004.....	8	-	-	-	-	-	-	-	-	2	-	2
2003.....	5	-	-	-	-	-	-	-	-	2	1	2
2002.....	3	-	-	-	-	-	-	-	1	-	1	1
2001.....	7	-	-	-	-	-	-	1	-	2	3	1
2000.....	3	-	-	-	1	-	-	-	-	-	1	-
1999.....	3	-	-	-	-	-	-	-	-	1	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Human immunodeficiency virus (HIV) disease resulting in other conditions (B23)

Asian or Pacific Islander 2/, both sexes

2004.....	15	-	-	-	-	-	1	-	-	1	4	3
2003.....	19	-	-	-	-	-	1	1	-	5	2	5
2002.....	18	-	-	-	-	-	-	-	1	2	3	3
2001.....	21	-	-	-	-	1	-	1	2	3	5	2
2000.....	20	-	-	-	-	-	-	-	3	4	2	2
1999.....	16	-	-	-	-	-	2	-	2	2	2	4

Asian or Pacific Islander 2/, male

2004.....	13	-	-	-	-	-	1	-	-	1	4	3
2003.....	14	-	-	-	-	-	-	-	-	4	2	5
2002.....	14	-	-	-	-	-	-	-	-	2	2	2
2001.....	16	-	-	-	-	-	-	-	2	3	4	2
2000.....	18	-	-	-	-	-	-	-	3	4	2	2
1999.....	12	-	-	-	-	-	1	-	2	1	2	3

Asian or Pacific Islander 2/, female

2004.....	2	-	-	-	-	-	-	-	-	-	-	-
2003.....	5	-	-	-	-	-	1	1	-	1	-	-
2002.....	4	-	-	-	-	-	-	-	1	-	1	1
2001.....	5	-	-	-	-	1	-	1	-	-	1	-
2000.....	2	-	-	-	-	-	-	-	-	-	-	-
1999.....	4	-	-	-	-	-	1	-	-	1	-	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Unspecified human immunodeficiency virus (HIV) disease (B24)												
All races, both sexes												
2004.....	3,541	1	1	3	3	5	47	124	292	595	773	682
2003.....	2,957	2	-	3	3	4	35	102	286	576	651	530
2002.....	2,810	4	-	-	4	6	32	109	293	588	605	514
2001.....	2,771	2	2	1	2	11	33	105	318	554	656	481
2000.....	2,743	2	3	1	3	7	27	131	370	539	584	491
1999.....	2,713	3	3	8	2	3	28	148	390	535	643	447
All races, male												
2004.....	2,507	-	-	1	1	2	28	75	173	400	548	496
2003.....	2,169	1	-	-	1	1	18	66	188	412	466	404
2002.....	2,056	3	-	-	2	2	13	65	181	426	442	389
2001.....	2,090	1	1	-	2	6	13	66	228	397	488	392
2000.....	2,052	1	1	-	1	3	19	69	246	402	441	388
1999.....	2,051	1	1	2	1	2	13	92	280	395	491	356
All races, female												
2004.....	1,034	1	1	2	2	3	19	49	119	195	225	186
2003.....	788	1	-	3	2	3	17	36	98	164	185	126
2002.....	754	1	-	-	2	4	19	44	112	162	163	125
2001.....	681	1	1	1	-	5	20	39	90	157	168	89
2000.....	691	1	2	1	2	4	8	62	124	137	143	103
1999.....	662	2	2	6	1	1	15	56	110	140	152	91
White, both sexes												
2004.....	1,410	1	-	1	1	2	12	28	99	253	330	257
2003.....	1,216	1	-	1	-	1	13	33	121	214	275	223
2002.....	1,148	3	-	-	2	2	7	45	98	231	254	235
2001.....	1,111	1	-	-	1	-	8	30	115	222	282	189
2000.....	1,143	-	-	1	2	2	5	51	137	239	246	214
1999.....	1,212	2	-	2	1	1	8	56	177	249	297	184

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	-------------	-----------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Unspecified human immunodeficiency virus (HIV) disease (B24)

White, male

2004.....	1,134	-	-	-	-	1	11	18	77	189	269	210
2003.....	1,004	1	-	-	-	-	7	25	95	173	227	189
2002.....	949	3	-	-	1	1	3	30	77	197	212	196
2001.....	960	1	-	-	1	-	5	21	95	192	245	163
2000.....	946	-	-	-	1	-	4	36	106	198	202	183
1999.....	1,015	1	-	1	1	1	4	39	144	208	250	156

White, female

2004.....	276	1	-	1	1	1	1	10	22	64	61	47
2003.....	212	-	-	1	-	1	6	8	26	41	48	34
2002.....	199	-	-	-	1	1	4	15	21	34	42	39
2001.....	151	-	-	-	-	-	3	9	20	30	37	26
2000.....	197	-	-	1	1	2	1	15	31	41	44	31
1999.....	197	1	-	1	-	-	4	17	33	41	47	28

Black, both sexes

2004.....	2,091	-	1	2	2	3	35	95	189	334	432	419
2003.....	1,713	1	-	2	3	3	21	68	161	357	370	301
2002.....	1,630	1	-	-	2	4	25	64	194	353	342	273
2001.....	1,635	1	2	1	1	10	25	74	200	326	368	288
2000.....	1,584	2	2	-	1	5	22	79	230	297	334	275
1999.....	1,473	1	3	6	1	2	20	91	208	281	337	259

Black, male

2004.....	1,344	-	-	1	1	1	17	56	94	205	269	282
2003.....	1,146	-	-	-	1	1	10	41	90	234	236	211
2002.....	1,080	-	-	-	1	1	10	35	103	227	222	187
2001.....	1,110	-	1	-	1	5	8	44	130	200	240	225
2000.....	1,094	1	1	-	-	3	15	32	138	202	235	203
1999.....	1,012	-	1	1	-	1	9	52	132	182	235	196

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Unspecified human immunodeficiency virus (HIV) disease (B24)												
Black, female												
2004.....	747	-	1	1	1	2	18	39	95	129	163	137
2003.....	567	1	-	2	2	2	11	27	71	123	134	90
2002.....	550	1	-	-	1	3	15	29	91	126	120	86
2001.....	525	1	1	1	-	5	17	30	70	126	128	63
2000.....	490	1	1	-	1	2	7	47	92	95	99	72
1999.....	461	1	2	5	1	1	11	39	76	99	102	63
American Indian 1/, both sexes												
2004.....	15	-	-	-	-	-	-	-	2	2	5	3
2003.....	14	-	-	-	-	-	-	-	2	2	4	3
2002.....	11	-	-	-	-	-	-	-	1	1	3	3
2001.....	15	-	-	-	-	-	-	1	2	4	2	3
2000.....	9	-	1	-	-	-	-	1	2	2	-	2
1999.....	14	-	-	-	-	-	-	-	4	4	5	1
American Indian 1/, male												
2004.....	10	-	-	-	-	-	-	-	1	2	4	2
2003.....	9	-	-	-	-	-	-	-	2	2	1	2
2002.....	9	-	-	-	-	-	-	-	1	1	3	3
2001.....	12	-	-	-	-	-	-	1	2	3	1	3
2000.....	5	-	-	-	-	-	-	1	1	1	-	2
1999.....	12	-	-	-	-	-	-	-	3	4	4	1
American Indian 1/, female												
2004.....	5	-	-	-	-	-	-	-	1	-	1	1
2003.....	5	-	-	-	-	-	-	-	-	-	3	1
2002.....	2	-	-	-	-	-	-	-	-	-	-	-
2001.....	3	-	-	-	-	-	-	-	-	1	1	-
2000.....	4	-	1	-	-	-	-	-	1	1	-	-
1999.....	2	-	-	-	-	-	-	-	1	-	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

All other and unspecified viral diseases (A81,A85.0-A85.1,A85.8,A86-A89,B01,B04,B07-B09,B25-B34)

All races, both sexes

2004.....	943	94	46	13	17	27	21	15	21	25	28	41
2003.....	1,014	87	60	18	12	24	17	16	15	22	44	37
2002.....	1,033	131	43	16	19	17	20	15	28	33	45	46
2001.....	857	84	52	21	13	20	17	11	13	31	32	22
2000.....	1,003	70	45	14	20	22	21	21	17	29	30	36
1999.....	1,121	101	48	21	12	18	11	17	22	36	38	40

All races, male

2004.....	486	56	24	6	11	13	12	4	14	12	15	18
2003.....	498	36	29	9	5	15	11	7	9	12	23	20
2002.....	498	75	29	4	7	9	15	8	14	17	25	26
2001.....	418	44	31	8	6	11	11	5	10	13	22	11
2000.....	485	38	26	9	8	14	12	12	8	18	14	15
1999.....	517	54	31	11	7	9	7	6	13	28	19	22

All races, female

2004.....	457	38	22	7	6	14	9	11	7	13	13	23
2003.....	516	51	31	9	7	9	6	9	6	10	21	17
2002.....	535	56	14	12	12	8	5	7	14	16	20	20
2001.....	439	40	21	13	7	9	6	6	3	18	10	11
2000.....	518	32	19	5	12	8	9	9	9	11	16	21
1999.....	604	47	17	10	5	9	4	11	9	8	19	18

White, both sexes

2004.....	785	67	30	10	10	19	17	7	16	20	19	33
2003.....	863	56	39	16	10	18	13	15	13	16	28	30
2002.....	876	96	32	11	15	12	14	12	22	22	34	36
2001.....	729	52	37	18	9	14	14	6	12	19	26	19
2000.....	849	46	30	12	15	17	16	12	11	23	25	27
1999.....	958	68	37	17	8	13	9	11	18	28	29	28

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

All other and unspecified viral diseases (A81,A85.0-A85.1,A85.8,A86-A89,B01,B04,B07-B09,B25-B34)

White, male

2004.....	410	38	16	5	7	9	10	1	11	10	11	14
2003.....	422	23	17	9	3	13	9	6	8	9	14	17
2002.....	420	52	21	2	6	7	10	6	11	14	20	19
2001.....	358	29	22	7	5	7	9	3	9	8	19	10
2000.....	408	25	15	7	4	10	11	7	6	16	11	10
1999.....	437	39	25	9	4	6	5	3	11	24	15	15

White, female

2004.....	375	29	14	5	3	10	7	6	5	10	8	19
2003.....	441	33	22	7	7	5	4	9	5	7	14	13
2002.....	456	44	11	9	9	5	4	6	11	8	14	17
2001.....	371	23	15	11	4	7	5	3	3	11	7	9
2000.....	441	21	15	5	11	7	5	5	5	7	14	17
1999.....	521	29	12	8	4	7	4	8	7	4	14	13

Black, both sexes

2004.....	117	21	9	3	5	6	4	5	4	5	6	7
2003.....	116	25	16	1	1	5	3	-	1	5	16	6
2002.....	120	27	9	2	2	4	5	1	5	10	9	10
2001.....	100	27	11	3	3	5	3	5	-	9	5	3
2000.....	125	21	10	1	3	4	5	7	6	3	4	8
1999.....	130	27	11	4	3	3	2	4	4	8	5	10

Black, male

2004.....	56	13	5	1	4	3	2	2	3	2	1	4
2003.....	58	10	9	-	1	1	2	-	1	3	9	3
2002.....	61	17	7	1	1	2	4	1	2	3	4	7
2001.....	47	14	7	1	-	3	2	2	-	4	3	1
2000.....	59	11	8	1	2	3	1	3	2	1	3	4
1999.....	64	13	6	2	2	2	2	1	2	4	3	7

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

All other and unspecified viral diseases (A81,A85.0-A85.1,A85.8,A86-A89,B01,B04,B07-B09,B25-B34)

Black, female

2004.....	61	8	4	2	1	3	2	3	1	3	5	3
2003.....	58	15	7	1	-	4	1	-	-	2	7	3
2002.....	59	10	2	1	1	2	1	-	3	7	5	3
2001.....	53	13	4	2	3	2	1	3	-	5	2	2
2000.....	66	10	2	-	1	1	4	4	4	2	1	4
1999.....	66	14	5	2	1	1	-	3	2	4	2	3

American Indian 1/, both sexes

2004.....	14	5	3	-	-	-	-	-	1	-	1	-
2003.....	10	1	2	-	1	-	1	-	-	-	-	-
2002.....	11	4	1	-	-	-	-	-	-	-	1	-
2001.....	10	3	3	-	1	-	-	-	-	1	-	-
2000.....	7	1	1	-	-	-	-	-	-	2	-	1
1999.....	10	2	-	-	-	1	-	-	-	-	3	1

American Indian 1/, male

2004.....	8	4	2	-	-	-	-	-	-	-	1	-
2003.....	3	-	1	-	1	-	-	-	-	-	-	-
2002.....	5	3	1	-	-	-	-	-	-	-	1	-
2001.....	5	1	2	-	1	-	-	-	-	-	-	-
2000.....	2	-	-	-	-	-	-	-	-	-	-	1
1999.....	2	-	-	-	-	-	-	-	-	-	1	-

American Indian 1/, female

2004.....	6	1	1	-	-	-	-	-	1	-	-	-
2003.....	7	1	1	-	-	-	1	-	-	-	-	-
2002.....	6	1	-	-	-	-	-	-	-	-	-	-
2001.....	5	2	1	-	-	-	-	-	-	1	-	-
2000.....	5	1	1	-	-	-	-	-	-	2	-	-
1999.....	8	2	-	-	-	1	-	-	-	-	2	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other and unspecified infectious and parasitic diseases and their sequelae (B35-B99)

All races, both sexes

2004.....	4,526	41	10	5	3	11	22	32	50	102	217	433
2003.....	4,304	46	15	6	6	16	13	30	53	107	216	415
2002.....	3,674	54	10	12	11	13	22	37	65	93	197	295
2001.....	3,104	65	6	9	11	13	22	36	53	102	143	231
2000.....	2,912	51	14	11	15	14	19	25	52	101	130	188
1999.....	4,024	59	23	17	17	16	24	64	98	152	228	282

All races, male

2004.....	2,610	27	4	4	2	7	20	14	35	60	138	307
2003.....	2,404	27	8	4	3	12	7	20	32	69	141	285
2002.....	2,146	30	6	6	11	6	11	20	38	61	137	216
2001.....	1,664	33	3	4	6	7	14	21	30	62	99	163
2000.....	1,577	30	8	7	9	7	11	12	31	69	87	116
1999.....	2,122	37	15	7	11	9	12	36	54	94	141	168

All races, female

2004.....	1,916	14	6	1	1	4	2	18	15	42	79	126
2003.....	1,900	19	7	2	3	4	6	10	21	38	75	130
2002.....	1,528	24	4	6	5	7	11	17	27	32	60	79
2001.....	1,440	32	3	5	5	6	8	15	23	40	44	68
2000.....	1,335	21	6	4	6	7	8	13	21	32	43	72
1999.....	1,902	22	8	10	6	7	12	28	44	58	87	114

White, both sexes

2004.....	3,741	26	8	4	3	7	18	22	29	74	170	362
2003.....	3,516	27	7	4	5	10	8	16	34	75	174	333
2002.....	2,975	34	6	11	13	9	11	25	40	60	142	233
2001.....	2,490	38	3	7	7	7	15	25	29	69	101	174
2000.....	2,349	30	9	8	11	10	14	13	29	61	92	143
1999.....	3,071	36	11	13	12	12	16	38	50	92	134	182

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other and unspecified infectious and parasitic diseases and their sequelae (B35-B99)

White, male

2004.....	2,173	18	3	3	2	5	16	10	19	44	108	260
2003.....	1,979	14	4	3	2	8	4	13	19	52	117	238
2002.....	1,731	19	3	6	9	4	5	11	23	39	102	174
2001.....	1,332	17	1	3	3	3	9	17	19	43	68	125
2000.....	1,262	20	5	5	7	4	9	7	18	42	64	91
1999.....	1,645	24	8	7	8	6	9	22	29	65	95	118

White, female

2004.....	1,568	8	5	1	1	2	2	12	10	30	62	102
2003.....	1,537	13	3	1	3	2	4	3	15	23	57	95
2002.....	1,244	15	3	5	4	5	6	14	17	21	40	59
2001.....	1,158	21	2	4	4	4	6	8	10	26	33	49
2000.....	1,087	10	4	3	4	6	5	6	11	19	28	52
1999.....	1,426	12	3	6	4	6	7	16	21	27	39	64

Black, both sexes

2004.....	586	15	1	-	-	4	3	9	15	23	43	53
2003.....	590	17	7	2	1	5	3	11	16	27	38	71
2002.....	549	19	4	-	2	3	9	12	23	30	45	56
2001.....	487	24	3	2	4	3	6	9	21	28	36	50
2000.....	471	20	3	3	2	2	5	8	22	35	37	44
1999.....	807	23	10	3	2	2	8	24	46	56	84	90

Black, male

2004.....	333	9	1	-	-	2	3	3	10	12	27	39
2003.....	320	12	4	1	1	3	2	5	12	14	21	41
2002.....	325	10	3	-	1	1	4	9	13	19	29	38
2001.....	265	15	2	1	3	3	4	4	9	16	28	34
2000.....	259	9	2	2	-	1	2	3	12	23	22	25
1999.....	391	13	7	-	1	2	3	12	25	27	39	45

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other and unspecified infectious and parasitic diseases and their sequelae (B35-B99)

Black, female

2004.....	253	6	-	-	-	2	-	6	5	11	16	14
2003.....	270	5	3	1	-	2	1	6	4	13	17	30
2002.....	224	9	1	-	1	2	5	3	10	11	16	18
2001.....	222	9	1	1	1	-	2	5	12	12	8	16
2000.....	212	11	1	1	2	1	3	5	10	12	15	19
1999.....	416	10	3	3	1	-	5	12	21	29	45	45

American Indian 1/, both sexes

2004.....	44	-	1	-	-	-	-	-	2	-	2	9
2003.....	43	2	1	-	-	-	-	1	-	3	1	6
2002.....	26	-	-	-	1	-	-	-	-	-	6	2
2001.....	28	1	-	-	-	1	1	-	-	2	3	3
2000.....	15	-	1	-	-	-	-	1	-	2	-	-
1999.....	35	-	1	-	1	-	-	-	-	1	3	2

American Indian 1/, male

2004.....	24	-	-	-	-	-	-	-	2	-	1	4
2003.....	22	1	-	-	-	-	-	1	-	2	-	3
2002.....	17	-	-	-	1	-	-	-	-	-	3	2
2001.....	12	-	-	-	-	1	1	-	-	1	1	1
2000.....	4	-	-	-	-	-	-	-	-	1	-	-
1999.....	17	-	-	-	-	-	-	-	-	1	1	-

American Indian 1/, female

2004.....	20	-	1	-	-	-	-	-	-	-	1	5
2003.....	21	1	1	-	-	-	-	-	-	1	1	3
2002.....	9	-	-	-	-	-	-	-	-	-	3	-
2001.....	16	1	-	-	-	-	-	-	-	1	2	2
2000.....	11	-	1	-	-	-	-	1	-	1	-	-
1999.....	18	-	1	-	1	-	-	-	-	-	2	2

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified infectious and parasitic diseases and their sequelae (B35-B99)												
Asian or Pacific Islander 2/, both sexes												
2004.....	155	-	-	1	-	-	1	1	4	5	2	9
2003.....	155	-	-	-	-	1	2	2	3	2	3	5
2002.....	124	1	-	1	-	1	2	-	2	3	4	4
2001.....	99	2	-	-	-	2	-	2	3	3	3	4
2000.....	77	1	1	-	2	2	-	3	1	3	1	1
1999.....	111	-	1	1	2	2	-	2	2	3	7	8
Asian or Pacific Islander 2/, male												
2004.....	80	-	-	1	-	-	1	1	4	4	2	4
2003.....	83	-	-	-	-	1	1	1	1	1	3	3
2002.....	73	1	-	-	-	1	2	-	2	3	3	2
2001.....	55	1	-	-	-	-	-	-	2	2	2	3
2000.....	52	1	1	-	2	2	-	2	1	3	1	-
1999.....	69	-	-	-	2	1	-	2	-	1	6	5
Asian or Pacific Islander 2/, female												
2004.....	75	-	-	-	-	-	-	-	-	1	-	5
2003.....	72	-	-	-	-	-	1	1	2	1	-	2
2002.....	51	-	-	1	-	-	-	-	-	-	1	2
2001.....	44	1	-	-	-	2	-	2	1	1	1	1
2000.....	25	-	-	-	-	-	-	1	-	-	-	1
1999.....	42	-	1	1	-	1	-	-	2	2	1	3

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Mycoses (B35-B49)

All races, both sexes

2004.....	1,064	35	8	3	2	7	10	13	21	30	39	56
2003.....	1,086	40	7	5	-	8	8	14	21	30	42	53
2002.....	1,236	47	6	6	5	6	12	19	32	33	51	60
2001.....	1,144	58	2	3	7	12	10	12	17	30	46	69
2000.....	1,245	48	10	7	8	8	8	15	22	42	50	61
1999.....	1,845	52	12	8	11	9	16	38	44	58	81	112

All races, male

2004.....	602	21	3	3	1	5	9	6	15	17	28	39
2003.....	589	23	4	3	-	7	5	10	14	19	29	30
2002.....	763	25	2	2	3	4	8	11	22	21	36	43
2001.....	614	29	-	-	3	7	7	5	10	19	29	47
2000.....	694	29	7	6	4	5	5	7	13	31	30	33
1999.....	1,030	32	7	3	8	4	7	22	26	43	50	62

All races, female

2004.....	462	14	5	-	1	2	1	7	6	13	11	17
2003.....	497	17	3	2	-	1	3	4	7	11	13	23
2002.....	473	22	4	4	2	2	4	8	10	12	15	17
2001.....	530	29	2	3	4	5	3	7	7	11	17	22
2000.....	551	19	3	1	4	3	3	8	9	11	20	28
1999.....	815	20	5	5	3	5	9	16	18	15	31	50

White, both sexes

2004.....	804	22	6	2	2	4	8	8	9	20	26	41
2003.....	833	23	2	3	-	6	5	7	9	22	28	34
2002.....	943	29	5	6	4	3	5	12	16	21	30	44
2001.....	860	32	1	3	4	7	7	10	10	17	32	50
2000.....	967	29	6	6	5	5	5	8	11	29	30	46
1999.....	1,406	31	5	5	10	7	11	22	25	40	47	79

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Mycoses (B35-B49)

White, male

2004.....	457	14	2	2	1	3	7	3	6	12	18	28
2003.....	455	12	2	2	-	5	3	6	5	14	20	20
2002.....	574	15	2	2	2	2	3	4	10	14	24	32
2001.....	456	13	-	-	1	3	5	4	6	9	19	36
2000.....	541	19	5	5	3	3	4	5	7	22	19	25
1999.....	805	21	5	3	7	3	5	12	15	32	30	44

White, female

2004.....	347	8	4	-	1	1	1	5	3	8	8	13
2003.....	378	11	-	1	-	1	2	1	4	8	8	14
2002.....	369	14	3	4	2	1	2	8	6	7	6	12
2001.....	404	19	1	3	3	4	2	6	4	8	13	14
2000.....	426	10	1	1	2	2	1	3	4	7	11	21
1999.....	601	10	-	2	3	4	6	10	10	8	17	35

Black, both sexes

2004.....	209	13	1	-	-	3	1	4	7	8	13	14
2003.....	212	15	4	2	-	2	2	6	10	7	14	18
2002.....	245	17	1	-	1	2	5	7	14	12	18	14
2001.....	237	23	1	-	3	3	2	2	6	11	12	17
2000.....	233	18	2	1	2	2	3	3	11	12	19	15
1999.....	365	21	5	3	-	1	5	14	18	15	31	30

Black, male

2004.....	115	7	1	-	-	2	1	2	4	4	10	10
2003.....	112	10	2	1	-	2	1	3	8	4	9	10
2002.....	154	9	-	-	1	1	3	7	10	7	11	10
2001.....	133	15	-	-	2	3	1	1	4	8	10	10
2000.....	122	9	1	1	-	1	1	-	6	8	10	8
1999.....	185	11	2	-	-	1	2	8	11	9	18	17

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Mycoses (B35-B49)												
Asian or Pacific Islander 2/, both sexes												
2004.....	39	-	-	1	-	-	1	1	3	2	-	-
2003.....	29	-	-	-	-	-	1	-	2	-	-	-
2002.....	38	1	-	-	-	1	2	-	2	-	1	1
2001.....	32	2	-	-	-	1	-	-	1	1	-	1
2000.....	37	1	1	-	1	1	-	3	-	1	1	-
1999.....	57	-	1	-	1	1	-	2	1	2	2	1
Asian or Pacific Islander 2/, male												
2004.....	25	-	-	1	-	-	1	1	3	1	-	-
2003.....	16	-	-	-	-	-	1	-	1	-	-	-
2002.....	27	1	-	-	-	1	2	-	2	-	-	-
2001.....	17	1	-	-	-	-	-	-	-	1	-	-
2000.....	28	1	1	-	1	1	-	2	-	1	1	-
1999.....	34	-	-	-	1	-	-	2	-	1	2	1
Asian or Pacific Islander 2/, female												
2004.....	14	-	-	-	-	-	-	-	-	1	-	-
2003.....	13	-	-	-	-	-	-	-	1	-	-	-
2002.....	11	-	-	-	-	-	-	-	-	-	1	1
2001.....	15	1	-	-	-	1	-	-	1	-	-	1
2000.....	9	-	-	-	-	-	-	1	-	-	-	-
1999.....	23	-	1	-	-	1	-	-	1	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Protozoal diseases (B50-B64)												
All races, both sexes												
2004.....	185	2	1	-	1	1	7	5	7	14	22	15
2003.....	178	3	1	-	1	2	2	7	11	18	15	16
2002.....	223	3	2	3	6	2	4	9	16	24	19	17
2001.....	277	2	-	3	1	-	7	12	16	28	20	26
2000.....	237	3	1	2	1	1	5	3	13	31	20	24
1999.....	289	6	4	1	1	2	2	14	17	33	39	23
All races, male												
2004.....	128	2	-	-	1	-	7	1	3	11	17	11
2003.....	104	1	-	-	-	1	2	3	7	10	9	12
2002.....	129	3	2	2	4	1	1	7	7	14	13	16
2001.....	157	1	-	2	1	-	4	7	8	14	16	16
2000.....	149	1	-	-	-	-	2	1	7	21	14	15
1999.....	158	4	1	-	1	2	2	8	8	20	21	12
All races, female												
2004.....	57	-	1	-	-	1	-	4	4	3	5	4
2003.....	74	2	1	-	1	1	-	4	4	8	6	4
2002.....	94	-	-	1	2	1	3	2	9	10	6	1
2001.....	120	1	-	1	-	-	3	5	8	14	4	10
2000.....	88	2	1	2	1	1	3	2	6	10	6	9
1999.....	131	2	3	1	-	-	-	6	9	13	18	11
White, both sexes												
2004.....	130	1	1	-	1	-	5	4	4	7	10	11
2003.....	122	3	1	-	1	1	1	2	7	6	8	8
2002.....	145	3	1	3	5	2	1	6	9	11	11	6
2001.....	198	2	-	2	-	-	3	6	9	19	11	13
2000.....	165	1	1	1	1	1	3	1	8	15	13	12
1999.....	203	4	3	-	-	2	2	5	10	20	25	13

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Protozoal diseases (B50-B64)												
White, male												
2004.....	94	1	-	-	1	-	5	1	2	6	9	8
2003.....	73	1	-	-	-	1	1	2	3	4	5	8
2002.....	88	3	1	2	4	1	-	5	5	6	9	6
2001.....	116	1	-	2	-	-	1	5	6	11	10	8
2000.....	107	1	-	-	-	-	1	-	4	10	10	9
1999.....	117	2	-	-	-	2	2	4	4	14	17	8
White, female												
2004.....	36	-	1	-	-	-	-	3	2	1	1	3
2003.....	49	2	1	-	1	-	-	-	4	2	3	-
2002.....	57	-	-	1	1	1	1	1	4	5	2	-
2001.....	82	1	-	-	-	-	2	1	3	8	1	5
2000.....	58	-	1	1	1	1	2	1	4	5	3	3
1999.....	86	2	3	-	-	-	-	1	6	6	8	5
Black, both sexes												
2004.....	48	1	-	-	-	1	2	1	3	7	12	4
2003.....	49	-	-	-	-	1	1	4	4	12	6	7
2002.....	69	-	1	-	1	-	3	3	7	11	8	11
2001.....	71	-	-	1	1	-	4	6	6	7	8	13
2000.....	70	2	-	1	-	-	2	2	5	16	7	12
1999.....	79	2	1	-	-	-	-	9	7	13	14	8
Black, male												
2004.....	29	1	-	-	-	-	2	-	1	5	8	3
2003.....	25	-	-	-	-	-	1	1	4	6	3	3
2002.....	37	-	1	-	-	-	1	2	2	6	4	10
2001.....	36	-	-	-	1	-	3	2	1	2	5	8
2000.....	40	-	-	-	-	-	1	1	3	11	4	6
1999.....	37	2	1	-	-	-	-	4	4	6	4	3

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Malaria (B50-B54)												
All races, both sexes												
2004.....	8	-	-	-	-	-	1	-	-	1	3	1
2003.....	4	-	-	-	-	-	-	-	-	-	1	-
2002.....	12	-	-	-	-	-	1	-	1	-	1	1
2001.....	9	-	-	-	1	-	-	-	-	2	-	2
2000.....	3	-	-	-	-	-	1	-	-	-	-	-
1999.....	7	1	-	-	-	-	-	2	-	-	1	-
All races, male												
2004.....	8	-	-	-	-	-	1	-	-	1	3	1
2003.....	4	-	-	-	-	-	-	-	-	-	1	-
2002.....	8	-	-	-	-	-	-	-	-	-	1	1
2001.....	5	-	-	-	1	-	-	-	-	-	-	1
2000.....	2	-	-	-	-	-	-	-	-	-	-	-
1999.....	4	1	-	-	-	-	-	1	-	-	-	-
All races, female												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	4	-	-	-	-	-	1	-	1	-	-	-
2001.....	4	-	-	-	-	-	-	-	-	2	-	1
2000.....	1	-	-	-	-	-	1	-	-	-	-	-
1999.....	3	-	-	-	-	-	-	1	-	-	1	-
White, both sexes												
2004.....	3	-	-	-	-	-	1	-	-	1	1	-
2003.....	4	-	-	-	-	-	-	-	-	-	1	-
2002.....	7	-	-	-	-	-	-	-	-	-	1	-
2001.....	5	-	-	-	-	-	-	-	-	2	-	1
2000.....	1	-	-	-	-	-	1	-	-	-	-	-
1999.....	4	-	-	-	-	-	-	1	-	-	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Trypanosomiasis (B56-B57)												
All races, both sexes												
2004.....	2	-	-	-	-	-	-	-	1	-	-	-
2003.....	1	-	-	-	-	-	-	-	-	-	-	-
2002.....	1	-	-	-	-	-	-	-	-	-	1	-
2001.....	3	-	-	-	-	-	-	-	-	2	-	-
2000.....	3	-	-	-	-	-	-	1	-	-	-	-
1999.....	3	-	-	-	-	-	-	-	-	2	-	1
All races, male												
2004.....	2	-	-	-	-	-	-	-	1	-	-	-
2003.....	1	-	-	-	-	-	-	-	-	-	-	-
2002.....	1	-	-	-	-	-	-	-	-	-	1	-
2001.....	2	-	-	-	-	-	-	-	-	1	-	-
2000.....	1	-	-	-	-	-	-	-	-	-	-	-
1999.....	1	-	-	-	-	-	-	-	-	1	-	-
All races, female												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	1	-	-	-	-	-	-	-	-	1	-	-
2000.....	2	-	-	-	-	-	-	1	-	-	-	-
1999.....	2	-	-	-	-	-	-	-	-	1	-	1
White, both sexes												
2004.....	2	-	-	-	-	-	-	-	1	-	-	-
2003.....	1	-	-	-	-	-	-	-	-	-	-	-
2002.....	1	-	-	-	-	-	-	-	-	-	1	-
2001.....	3	-	-	-	-	-	-	-	-	2	-	-
2000.....	3	-	-	-	-	-	-	1	-	-	-	-
1999.....	3	-	-	-	-	-	-	-	-	2	-	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumocystosis (B59)												
All races, both sexes												
2004.....	158	2	1	-	1	1	5	2	4	11	18	12
2003.....	156	3	1	-	1	1	2	7	10	12	13	16
2002.....	188	2	2	2	2	2	3	7	14	19	16	13
2001.....	245	2	-	1	-	-	6	12	16	23	15	24
2000.....	206	3	1	1	1	1	4	1	8	25	18	23
1999.....	244	4	3	1	1	1	2	9	14	26	32	19
All races, male												
2004.....	108	2	-	-	1	-	5	-	2	8	14	8
2003.....	89	1	-	-	-	-	2	3	6	7	7	12
2002.....	109	2	2	1	1	1	1	6	6	13	11	13
2001.....	139	1	-	1	-	-	4	7	8	12	13	15
2000.....	129	1	-	-	-	-	2	1	4	16	12	15
1999.....	132	2	-	-	1	1	2	4	7	15	18	10
All races, female												
2004.....	50	-	1	-	-	1	-	2	2	3	4	4
2003.....	67	2	1	-	1	1	-	4	4	5	6	4
2002.....	79	-	-	1	1	1	2	1	8	6	5	-
2001.....	106	1	-	-	-	-	2	5	8	11	2	9
2000.....	77	2	1	1	1	1	2	-	4	9	6	8
1999.....	112	2	3	1	-	-	-	5	7	11	14	9
White, both sexes												
2004.....	113	1	1	-	1	-	3	2	2	6	9	9
2003.....	111	3	1	-	1	-	1	2	7	6	7	8
2002.....	127	2	1	2	1	2	1	5	9	10	9	6
2001.....	181	2	-	1	-	-	3	6	9	14	10	12
2000.....	146	1	1	-	1	1	2	-	6	12	12	12
1999.....	177	3	3	-	-	1	2	3	8	16	20	11

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumocystosis (B59)												
White, male												
2004.....	81	1	-	-	1	-	3	-	1	5	8	6
2003.....	64	1	-	-	-	-	1	2	3	4	4	8
2002.....	73	2	1	1	1	1	-	4	5	5	7	6
2001.....	105	1	-	1	-	-	1	5	6	9	9	8
2000.....	95	1	-	-	-	-	1	-	2	8	9	9
1999.....	101	1	-	-	-	1	2	2	4	11	14	7
White, female												
2004.....	32	-	1	-	-	-	-	2	1	1	1	3
2003.....	47	2	1	-	1	-	-	-	4	2	3	-
2002.....	54	-	-	1	-	1	1	1	4	5	2	-
2001.....	76	1	-	-	-	-	2	1	3	5	1	4
2000.....	51	-	1	-	1	1	1	-	4	4	3	3
1999.....	76	2	3	-	-	-	-	1	4	5	6	4
Black, both sexes												
2004.....	40	1	-	-	-	1	2	-	2	5	9	3
2003.....	39	-	-	-	-	1	1	4	3	6	5	7
2002.....	54	-	1	-	1	-	2	2	5	7	7	7
2001.....	57	-	-	-	-	-	3	6	6	7	4	12
2000.....	59	2	-	1	-	-	2	1	2	13	6	11
1999.....	60	1	-	-	-	-	-	6	6	10	12	6
Black, male												
2004.....	24	1	-	-	-	-	2	-	1	3	6	2
2003.....	20	-	-	-	-	-	1	1	3	3	2	3
2002.....	33	-	1	-	-	-	1	2	1	6	4	7
2001.....	30	-	-	-	-	-	3	2	1	2	3	7
2000.....	33	-	-	-	-	-	1	1	2	8	3	6
1999.....	27	1	-	-	-	-	-	2	3	4	4	2

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified protozoal diseases (B58,B60-B64)												
All races, both sexes												
2004.....	17	-	-	-	-	-	1	3	2	2	1	2
2003.....	17	-	-	-	-	1	-	-	1	6	1	-
2002.....	21	1	-	1	4	-	-	2	1	5	1	3
2001.....	20	-	-	2	-	-	1	-	-	1	5	-
2000.....	25	-	-	1	-	-	-	1	5	6	2	1
1999.....	35	1	1	-	-	1	-	3	3	5	6	3
All races, male												
2004.....	10	-	-	-	-	-	1	1	-	2	-	2
2003.....	10	-	-	-	-	1	-	-	1	3	1	-
2002.....	11	1	-	1	3	-	-	1	1	1	-	2
2001.....	11	-	-	1	-	-	-	-	-	1	3	-
2000.....	17	-	-	-	-	-	-	-	3	5	2	-
1999.....	21	1	1	-	-	1	-	3	1	4	3	2
All races, female												
2004.....	7	-	-	-	-	-	-	2	2	-	1	-
2003.....	7	-	-	-	-	-	-	-	-	3	-	-
2002.....	10	-	-	-	1	-	-	1	-	4	1	1
2001.....	9	-	-	1	-	-	1	-	-	-	2	-
2000.....	8	-	-	1	-	-	-	1	2	1	-	1
1999.....	14	-	-	-	-	-	-	-	2	1	3	1
White, both sexes												
2004.....	12	-	-	-	-	-	1	2	1	-	-	2
2003.....	6	-	-	-	-	1	-	-	-	-	-	-
2002.....	9	1	-	1	4	-	-	1	-	1	-	-
2001.....	9	-	-	1	-	-	-	-	-	1	1	-
2000.....	15	-	-	1	-	-	-	-	2	3	1	-
1999.....	19	1	-	-	-	1	-	1	2	2	4	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other and unspecified protozoal diseases (B58,B60-B64)

White, male

2004.....	8	-	-	-	-	-	1	1	-	-	-	2
2003.....	4	-	-	-	-	1	-	-	-	-	-	-
2002.....	7	1	-	1	3	-	-	1	-	1	-	-
2001.....	7	-	-	1	-	-	-	-	-	1	1	-
2000.....	11	-	-	-	-	-	-	-	2	2	1	-
1999.....	12	1	-	-	-	1	-	1	-	2	3	1

White, female

2004.....	4	-	-	-	-	-	-	1	1	-	-	-
2003.....	2	-	-	-	-	-	-	-	-	-	-	-
2002.....	2	-	-	-	1	-	-	-	-	-	-	-
2001.....	2	-	-	-	-	-	-	-	-	-	-	-
2000.....	4	-	-	1	-	-	-	-	-	1	-	-
1999.....	7	-	-	-	-	-	-	-	2	-	1	-

Black, both sexes

2004.....	5	-	-	-	-	-	-	1	1	2	1	-
2003.....	10	-	-	-	-	-	-	-	1	6	1	-
2002.....	11	-	-	-	-	-	-	1	1	4	1	3
2001.....	11	-	-	1	-	-	1	-	-	-	4	-
2000.....	10	-	-	-	-	-	-	1	3	3	1	1
1999.....	16	-	1	-	-	-	-	2	1	3	2	2

Black, male

2004.....	2	-	-	-	-	-	-	-	-	2	-	-
2003.....	5	-	-	-	-	-	-	-	1	3	1	-
2002.....	3	-	-	-	-	-	-	-	1	-	-	2
2001.....	4	-	-	-	-	-	-	-	-	-	2	-
2000.....	6	-	-	-	-	-	-	-	1	3	1	-
1999.....	9	-	1	-	-	-	-	2	1	2	-	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Helminthiases (B65-B83)												
All races, both sexes												
2004.....	36	-	-	-	-	-	1	4	2	5	2	2
2003.....	20	-	-	-	-	-	1	1	1	1	2	2
2002.....	15	-	-	-	-	-	1	1	2	1	-	1
2001.....	25	-	-	-	-	-	1	3	2	2	-	2
2000.....	33	-	1	-	1	1	2	3	2	2	1	3
1999.....	14	-	-	1	-	-	2	1	-	1	1	1
All races, male												
2004.....	22	-	-	-	-	-	1	3	-	3	1	2
2003.....	14	-	-	-	-	-	-	1	1	-	1	2
2002.....	8	-	-	-	-	-	-	-	1	1	-	1
2001.....	19	-	-	-	-	-	1	3	2	1	-	1
2000.....	22	-	-	-	1	-	1	2	1	2	1	2
1999.....	8	-	-	-	-	-	1	1	-	-	1	1
All races, female												
2004.....	14	-	-	-	-	-	-	1	2	2	1	-
2003.....	6	-	-	-	-	-	1	-	-	1	1	-
2002.....	7	-	-	-	-	-	1	1	1	-	-	-
2001.....	6	-	-	-	-	-	-	-	-	1	-	1
2000.....	11	-	1	-	-	1	1	1	1	-	-	1
1999.....	6	-	-	1	-	-	1	-	-	1	-	-
White, both sexes												
2004.....	23	-	-	-	-	-	1	3	2	4	1	1
2003.....	18	-	-	-	-	-	1	1	1	-	2	2
2002.....	12	-	-	-	-	-	1	1	2	1	-	-
2001.....	23	-	-	-	-	-	1	3	1	2	-	2
2000.....	28	-	1	-	1	1	2	3	1	2	1	2
1999.....	12	-	-	1	-	-	2	1	-	1	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified helminthiasis (B66-B83)												
All races, both sexes												
2004.....	31	-	-	-	-	-	1	4	2	4	2	1
2003.....	20	-	-	-	-	-	1	1	1	1	2	2
2002.....	15	-	-	-	-	-	1	1	2	1	-	1
2001.....	25	-	-	-	-	-	1	3	2	2	-	2
2000.....	32	-	1	-	1	1	2	3	2	2	1	3
1999.....	13	-	-	1	-	-	2	1	-	1	1	1
All races, male												
2004.....	18	-	-	-	-	-	1	3	-	2	1	1
2003.....	14	-	-	-	-	-	-	1	1	-	1	2
2002.....	8	-	-	-	-	-	-	-	1	1	-	1
2001.....	19	-	-	-	-	-	1	3	2	1	-	1
2000.....	21	-	-	-	1	-	1	2	1	2	1	2
1999.....	8	-	-	-	-	-	1	1	-	-	1	1
All races, female												
2004.....	13	-	-	-	-	-	-	1	2	2	1	-
2003.....	6	-	-	-	-	-	1	-	-	1	1	-
2002.....	7	-	-	-	-	-	1	1	1	-	-	-
2001.....	6	-	-	-	-	-	-	-	-	1	-	1
2000.....	11	-	1	-	-	1	1	1	1	-	-	1
1999.....	5	-	-	1	-	-	1	-	-	1	-	-
White, both sexes												
2004.....	23	-	-	-	-	-	1	3	2	4	1	1
2003.....	18	-	-	-	-	-	1	1	1	-	2	2
2002.....	12	-	-	-	-	-	1	1	2	1	-	-
2001.....	23	-	-	-	-	-	1	3	1	2	-	2
2000.....	28	-	1	-	1	1	2	3	1	2	1	2
1999.....	11	-	-	1	-	-	2	1	-	1	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Sequelae of tuberculosis (B90)

All races, both sexes

2004.....	82	...	-	-	-	-	-	-	-	-	-	4
2003.....	80	...	-	-	-	-	-	-	1	-	1	3
2002.....	108	...	-	1	-	-	-	1	-	1	2	4
2001.....	111	...	-	-	-	-	-	-	2	1	1	3
2000.....	125	...	-	-	1	-	-	-	1	1	2	4
1999.....	189	...	-	1	-	-	-	3	1	3	4	10

All races, male

2004.....	41	...	-	-	-	-	-	-	-	-	-	3
2003.....	33	...	-	-	-	-	-	-	-	-	1	2
2002.....	46	...	-	-	-	-	-	-	-	1	1	3
2001.....	47	...	-	-	-	-	-	-	2	-	-	2
2000.....	66	...	-	-	1	-	-	-	1	1	1	3
1999.....	77	...	-	-	-	-	-	1	-	1	2	4

All races, female

2004.....	41	...	-	-	-	-	-	-	-	-	-	1
2003.....	47	...	-	-	-	-	-	-	1	-	-	1
2002.....	62	...	-	1	-	-	-	1	-	-	1	1
2001.....	64	...	-	-	-	-	-	-	-	1	1	1
2000.....	59	...	-	-	-	-	-	-	-	-	1	1
1999.....	112	...	-	1	-	-	-	2	1	2	2	6

White, both sexes

2004.....	51	...	-	-	-	-	-	-	-	-	-	1
2003.....	59	...	-	-	-	-	-	-	1	-	1	1
2002.....	80	...	-	-	-	-	-	1	-	-	1	1
2001.....	84	...	-	-	-	-	-	-	1	-	1	2
2000.....	89	...	-	-	1	-	-	-	-	-	-	1
1999.....	140	...	-	1	-	-	-	3	-	-	1	5

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Sequelae of tuberculosis (B90)												
White, male												
2004.....	24	...	-	-	-	-	-	-	-	-	-	1
2003.....	22	...	-	-	-	-	-	-	-	-	1	1
2002.....	28	...	-	-	-	-	-	-	-	-	-	1
2001.....	38	...	-	-	-	-	-	-	1	-	-	1
2000.....	42	...	-	-	1	-	-	-	-	-	-	1
1999.....	50	...	-	-	-	-	-	1	-	-	1	1
White, female												
2004.....	27	...	-	-	-	-	-	-	-	-	-	-
2003.....	37	...	-	-	-	-	-	-	1	-	-	-
2002.....	52	...	-	-	-	-	-	1	-	-	1	-
2001.....	46	...	-	-	-	-	-	-	-	-	1	1
2000.....	47	...	-	-	-	-	-	-	-	-	-	-
1999.....	90	...	-	1	-	-	-	2	-	-	-	4
Black, both sexes												
2004.....	15	...	-	-	-	-	-	-	-	-	-	2
2003.....	8	...	-	-	-	-	-	-	-	-	-	1
2002.....	11	...	-	-	-	-	-	-	-	-	-	2
2001.....	10	...	-	-	-	-	-	-	1	1	-	1
2000.....	19	...	-	-	-	-	-	-	-	-	2	2
1999.....	22	...	-	-	-	-	-	-	-	2	2	4
Black, male												
2004.....	9	...	-	-	-	-	-	-	-	-	-	2
2003.....	4	...	-	-	-	-	-	-	-	-	-	-
2002.....	7	...	-	-	-	-	-	-	-	-	-	1
2001.....	5	...	-	-	-	-	-	-	1	-	-	1
2000.....	14	...	-	-	-	-	-	-	-	-	1	2
1999.....	10	...	-	-	-	-	-	-	-	1	1	3

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Sequelae of tuberculosis (B90)												
Asian or Pacific Islander 2/, both sexes												
2004.....	14	...	-	-	-	-	-	-	-	-	-	-
2003.....	10	...	-	-	-	-	-	-	-	-	-	1
2002.....	16	...	-	1	-	-	-	-	-	1	1	1
2001.....	15	...	-	-	-	-	-	-	-	-	-	-
2000.....	14	...	-	-	-	-	-	-	1	1	-	1
1999.....	19	...	-	-	-	-	-	-	1	1	-	1
Asian or Pacific Islander 2/, male												
2004.....	8	...	-	-	-	-	-	-	-	-	-	-
2003.....	6	...	-	-	-	-	-	-	-	-	-	1
2002.....	10	...	-	-	-	-	-	-	-	1	1	1
2001.....	4	...	-	-	-	-	-	-	-	-	-	-
2000.....	10	...	-	-	-	-	-	-	1	1	-	-
1999.....	12	...	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	6	...	-	-	-	-	-	-	-	-	-	-
2003.....	4	...	-	-	-	-	-	-	-	-	-	-
2002.....	6	...	-	1	-	-	-	-	-	-	-	-
2001.....	11	...	-	-	-	-	-	-	-	-	-	-
2000.....	4	...	-	-	-	-	-	-	-	-	-	1
1999.....	7	...	-	-	-	-	-	-	1	1	-	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Sequelae of poliomyelitis (B91)												
All races, both sexes												
2004.....	313	...	-	-	-	-	-	-	-	-	-	5
2003.....	301	...	-	-	-	2	-	-	2	1	1	6
2002.....	323	...	-	-	-	-	-	-	-	1	-	4
2001.....	338	...	-	-	-	-	-	1	-	-	2	4
2000.....	299	...	-	-	-	1	-	-	1	-	2	6
1999.....	263	...	-	-	1	-	-	-	-	-	-	8
All races, male												
2004.....	152	...	-	-	-	-	-	-	-	-	-	1
2003.....	140	...	-	-	-	2	-	-	1	1	-	3
2002.....	146	...	-	-	-	-	-	-	-	1	-	1
2001.....	163	...	-	-	-	-	-	-	-	-	1	3
2000.....	133	...	-	-	-	-	-	-	1	-	-	3
1999.....	121	...	-	-	1	-	-	-	-	-	-	6
All races, female												
2004.....	161	...	-	-	-	-	-	-	-	-	-	4
2003.....	161	...	-	-	-	-	-	-	1	-	1	3
2002.....	177	...	-	-	-	-	-	-	-	-	-	3
2001.....	175	...	-	-	-	-	-	1	-	-	1	1
2000.....	166	...	-	-	-	1	-	-	-	-	2	3
1999.....	142	...	-	-	-	-	-	-	-	-	-	2
White, both sexes												
2004.....	304	...	-	-	-	-	-	-	-	-	-	3
2003.....	286	...	-	-	-	1	-	-	1	-	1	4
2002.....	314	...	-	-	-	-	-	-	-	1	-	4
2001.....	325	...	-	-	-	-	-	-	-	-	1	3
2000.....	291	...	-	-	-	1	-	-	-	-	2	5
1999.....	259	...	-	-	1	-	-	-	-	-	-	8

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

All other and unspecified infectious and parasitic diseases and their sequelae (B85-B89,B92-B99)

All races, both sexes

2004.....	2,846	4	1	2	-	3	4	10	20	53	154	351
2003.....	2,639	3	7	1	5	4	2	8	17	57	155	335
2002.....	1,769	4	2	2	5	5	5	7	15	33	125	209
2001.....	1,209	5	4	3	3	1	4	8	16	41	74	127
2000.....	973	-	2	2	4	3	4	4	13	25	55	90
1999.....	1,424	1	7	6	4	5	4	8	36	57	103	128

All races, male

2004.....	1,665	4	1	1	-	2	3	4	17	29	92	251
2003.....	1,524	3	4	1	3	2	-	6	9	39	101	236
2002.....	1,054	2	2	2	4	1	2	2	8	23	87	152
2001.....	664	3	3	2	2	-	2	6	8	28	53	94
2000.....	513	-	1	1	3	2	3	2	8	14	41	60
1999.....	728	1	7	4	1	3	2	4	20	30	67	83

All races, female

2004.....	1,181	-	-	1	-	1	1	6	3	24	62	100
2003.....	1,115	-	3	-	2	2	2	2	8	18	54	99
2002.....	715	2	-	-	1	4	3	5	7	10	38	57
2001.....	545	2	1	1	1	1	2	2	8	13	21	33
2000.....	460	-	1	1	1	1	1	2	5	11	14	30
1999.....	696	-	-	2	3	2	2	4	16	27	36	45

White, both sexes

2004.....	2,429	3	1	2	-	3	4	7	14	43	133	305
2003.....	2,198	1	4	1	4	2	1	6	15	47	134	284
2002.....	1,481	2	-	2	4	4	4	5	13	26	100	178
2001.....	1,000	4	2	2	3	-	4	6	8	31	56	104
2000.....	809	-	1	1	3	2	4	1	9	15	46	77
1999.....	1,051	1	3	6	1	3	1	7	15	31	60	77

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

All other and unspecified infectious and parasitic diseases and their sequelae (B85-B89,B92-B99)

White, male

2004.....	1,435	3	1	1	-	2	3	3	11	24	80	221
2003.....	1,284	1	2	1	2	1	-	4	9	34	90	205
2002.....	891	1	-	2	3	1	2	2	7	17	69	134
2001.....	552	3	1	1	2	-	2	5	5	22	39	77
2000.....	424	-	-	-	2	1	3	-	6	8	34	53
1999.....	548	1	3	4	-	1	1	4	10	19	46	59

White, female

2004.....	994	-	-	1	-	1	1	4	3	19	53	84
2003.....	914	-	2	-	2	1	1	2	6	13	44	79
2002.....	590	1	-	-	1	3	2	3	6	9	31	44
2001.....	448	1	1	1	1	-	2	1	3	9	17	27
2000.....	385	-	1	1	1	1	1	1	3	7	12	24
1999.....	503	-	-	2	1	2	-	3	5	12	14	18

Black, both sexes

2004.....	297	1	-	-	-	-	-	3	5	7	17	31
2003.....	311	2	3	-	1	2	-	1	2	7	18	43
2002.....	214	2	2	-	-	1	1	2	2	7	19	28
2001.....	161	1	2	1	-	-	-	1	7	9	16	18
2000.....	139	-	1	1	-	-	-	3	4	7	9	13
1999.....	337	-	4	-	2	1	3	1	21	26	37	47

Black, male

2004.....	172	1	-	-	-	-	-	1	5	2	9	23
2003.....	174	2	2	-	1	1	-	1	-	4	9	27
2002.....	124	1	2	-	-	-	-	-	1	6	14	16
2001.....	85	-	2	1	-	-	-	1	2	6	13	14
2000.....	78	-	1	1	-	-	-	2	2	4	7	7
1999.....	157	-	4	-	1	1	1	-	10	11	16	21

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

All other and unspecified infectious and parasitic diseases and their sequelae (B85-B89,B92-B99)

Black, female

2004.....	125	-	-	-	-	-	-	2	-	5	8	8
2003.....	137	-	1	-	-	1	-	-	2	3	9	16
2002.....	90	1	-	-	-	1	1	2	1	1	5	12
2001.....	76	1	-	-	-	-	-	-	5	3	3	4
2000.....	61	-	-	-	-	-	-	1	2	3	2	6
1999.....	180	-	-	-	1	-	2	1	11	15	21	26

American Indian 1/, both sexes

2004.....	29	-	-	-	-	-	-	-	-	-	2	7
2003.....	25	-	-	-	-	-	-	-	-	2	1	5
2002.....	14	-	-	-	1	-	-	-	-	-	4	1
2001.....	9	-	-	-	-	-	-	-	-	1	-	2
2000.....	4	-	-	-	-	-	-	-	-	2	-	-
1999.....	9	-	-	-	1	-	-	-	-	-	1	-

American Indian 1/, male

2004.....	18	-	-	-	-	-	-	-	-	-	1	3
2003.....	14	-	-	-	-	-	-	-	-	1	-	3
2002.....	8	-	-	-	1	-	-	-	-	-	2	1
2001.....	3	-	-	-	-	-	-	-	-	-	-	-
2000.....	1	-	-	-	-	-	-	-	-	1	-	-
1999.....	5	-	-	-	-	-	-	-	-	-	1	-

American Indian 1/, female

2004.....	11	-	-	-	-	-	-	-	-	-	1	4
2003.....	11	-	-	-	-	-	-	-	-	1	1	2
2002.....	6	-	-	-	-	-	-	-	-	-	2	-
2001.....	6	-	-	-	-	-	-	-	-	1	-	2
2000.....	3	-	-	-	-	-	-	-	-	1	-	-
1999.....	4	-	-	-	1	-	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

II. Neoplasms (C00-D48)

All races, both sexes

2004.....	567,468	145	452	567	536	781	1,038	1,349	2,457	4,650	10,391	19,498
2003.....	570,465	137	443	554	601	731	1,016	1,376	2,543	5,061	10,787	19,847
2002.....	570,570	143	462	581	580	764	1,053	1,320	2,715	5,287	11,164	19,622
2001.....	567,444	117	478	545	568	779	1,027	1,449	2,696	5,622	11,313	19,503
2000.....	566,637	152	473	551	562	785	1,023	1,427	2,654	5,720	11,168	19,096
1999.....	563,065	126	481	573	540	797	1,025	1,484	2,692	5,904	11,170	18,688

All races, male

2004.....	293,594	72	240	323	295	460	630	714	1,130	1,977	4,603	9,553
2003.....	294,677	66	258	314	320	462	571	704	1,198	2,149	4,871	9,875
2002.....	295,304	72	246	319	330	445	621	711	1,241	2,243	5,009	9,587
2001.....	293,810	57	259	279	306	467	610	727	1,210	2,410	5,080	9,550
2000.....	292,641	82	266	301	317	473	611	722	1,214	2,488	5,044	9,403
1999.....	292,352	66	260	309	295	474	603	787	1,268	2,546	5,122	9,293

All races, female

2004.....	273,874	73	212	244	241	321	408	635	1,327	2,673	5,788	9,945
2003.....	275,788	71	185	240	281	269	445	672	1,345	2,912	5,916	9,972
2002.....	275,266	71	216	262	250	319	432	609	1,474	3,044	6,155	10,035
2001.....	273,634	60	219	266	262	312	417	722	1,486	3,212	6,233	9,953
2000.....	273,996	70	207	250	245	312	412	705	1,440	3,232	6,124	9,693
1999.....	270,713	60	221	264	245	323	422	697	1,424	3,358	6,048	9,395

White, both sexes

2004.....	490,226	122	354	445	407	616	793	1,034	1,878	3,609	8,017	15,228
2003.....	493,737	112	356	453	455	594	808	1,069	1,943	3,919	8,409	15,457
2002.....	494,394	115	349	464	449	608	859	1,017	2,101	4,121	8,747	15,454
2001.....	491,912	91	401	425	432	620	813	1,127	2,093	4,402	8,847	15,384
2000.....	492,182	123	374	443	451	605	801	1,128	2,073	4,440	8,616	14,968
1999.....	489,087	94	401	449	415	605	792	1,176	2,095	4,602	8,727	14,727

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
II. Neoplasms (C00-D48)												
White, male												
2004.....	253,521	61	183	258	228	370	483	570	905	1,594	3,606	7,496
2003.....	255,089	54	204	261	241	377	459	574	935	1,741	3,839	7,732
2002.....	255,769	62	189	245	251	362	506	555	987	1,775	3,994	7,638
2001.....	254,235	44	213	214	221	372	489	578	979	1,925	4,017	7,570
2000.....	253,303	67	215	239	256	362	484	583	987	1,975	3,911	7,419
1999.....	253,287	52	216	248	226	352	470	627	1,001	2,047	4,051	7,287
White, female												
2004.....	236,705	61	171	187	179	246	310	464	973	2,015	4,411	7,732
2003.....	238,648	58	152	192	214	217	349	495	1,008	2,178	4,570	7,725
2002.....	238,625	53	160	219	198	246	353	462	1,114	2,346	4,753	7,816
2001.....	237,677	47	188	211	211	248	324	549	1,114	2,477	4,830	7,814
2000.....	238,879	56	159	204	195	243	317	545	1,086	2,465	4,705	7,549
1999.....	235,800	42	185	201	189	253	322	549	1,094	2,555	4,676	7,440
Black, both sexes												
2004.....	63,691	17	73	88	103	137	193	259	446	821	1,928	3,544
2003.....	63,783	18	66	83	108	102	161	243	453	917	1,967	3,705
2002.....	63,760	21	88	92	98	121	144	225	475	938	2,035	3,499
2001.....	63,327	17	53	93	109	125	158	248	467	985	2,057	3,430
2000.....	63,081	20	79	85	84	144	173	233	449	1,041	2,130	3,493
1999.....	63,124	26	62	97	103	152	192	242	459	1,088	2,058	3,358
Black, male												
2004.....	32,994	6	42	44	52	73	113	111	170	276	792	1,705
2003.....	32,967	9	39	46	59	61	89	99	188	306	852	1,812
2002.....	33,127	6	42	61	54	69	80	108	190	357	841	1,645
2001.....	33,204	8	33	49	66	75	91	110	171	383	864	1,658
2000.....	33,337	8	42	48	47	89	99	99	174	396	945	1,688
1999.....	33,371	11	36	47	58	95	104	122	195	400	897	1,721

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

II. Neoplasms (C00-D48)

Black, female

2004.....	30,697	11	31	44	51	64	80	148	276	545	1,136	1,839
2003.....	30,816	9	27	37	49	41	72	144	265	611	1,115	1,893
2002.....	30,633	15	46	31	44	52	64	117	285	581	1,194	1,854
2001.....	30,123	9	20	44	43	50	67	138	296	602	1,193	1,772
2000.....	29,744	12	37	37	37	55	74	134	275	645	1,185	1,805
1999.....	29,753	15	26	50	45	57	88	120	264	688	1,161	1,637

American Indian 1/, both sexes

2004.....	2,440	1	4	7	2	7	10	8	18	39	71	133
2003.....	2,193	-	7	4	5	9	10	10	16	37	66	115
2002.....	2,209	1	5	6	7	9	11	12	24	39	64	109
2001.....	2,213	4	4	1	5	12	14	16	19	47	73	107
2000.....	1,959	2	3	6	3	8	9	11	17	37	72	89
1999.....	1,878	1	4	2	4	5	9	13	14	28	57	87

American Indian 1/, male

2004.....	1,264	1	2	6	2	4	5	6	6	16	29	69
2003.....	1,112	-	7	1	2	6	7	4	8	15	29	62
2002.....	1,096	-	2	3	5	4	6	6	8	14	30	44
2001.....	1,132	4	3	1	3	8	9	12	7	17	38	49
2000.....	1,017	2	2	4	1	5	5	6	8	17	33	44
1999.....	971	-	1	1	4	2	6	7	8	13	22	52

American Indian 1/, female

2004.....	1,176	-	2	1	-	3	5	2	12	23	42	64
2003.....	1,081	-	-	3	3	3	3	6	8	22	37	53
2002.....	1,113	1	3	3	2	5	5	6	16	25	34	65
2001.....	1,081	-	1	-	2	4	5	4	12	30	35	58
2000.....	942	-	1	2	2	3	4	5	9	20	39	45
1999.....	907	1	3	1	-	3	3	6	6	15	35	35

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

II. Neoplasms (C00-D48)

Asian or Pacific Islander 2/, both sexes

2004.....	11,111	5	21	27	24	21	42	48	115	181	375	593
2003.....	10,752	7	14	14	33	26	37	54	131	188	345	570
2002.....	10,207	6	20	19	26	26	39	66	115	189	318	560
2001.....	9,992	5	20	26	22	22	42	58	117	188	336	582
2000.....	9,415	7	17	17	24	28	40	55	115	202	350	546
1999.....	8,976	5	14	25	18	35	32	53	124	186	328	516

Asian or Pacific Islander 2/, male

2004.....	5,815	4	13	15	13	13	29	27	49	91	176	283
2003.....	5,509	3	8	6	18	18	16	27	67	87	151	269
2002.....	5,312	4	13	10	20	10	29	42	56	97	144	260
2001.....	5,239	1	10	15	16	12	21	27	53	85	161	273
2000.....	4,984	5	7	10	13	17	23	34	45	100	155	252
1999.....	4,723	3	7	13	7	25	23	31	64	86	152	233

Asian or Pacific Islander 2/, female

2004.....	5,296	1	8	12	11	8	13	21	66	90	199	310
2003.....	5,243	4	6	8	15	8	21	27	64	101	194	301
2002.....	4,895	2	7	9	6	16	10	24	59	92	174	300
2001.....	4,753	4	10	11	6	10	21	31	64	103	175	309
2000.....	4,431	2	10	7	11	11	17	21	70	102	195	294
1999.....	4,253	2	7	12	11	10	9	22	60	100	176	283

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Malignant neoplasms (C00-C97)												
All races, both sexes												
2004.....	553,888	74	399	526	493	731	978	1,286	2,347	4,518	10,205	19,190
2003.....	556,902	75	392	516	560	690	961	1,307	2,434	4,919	10,590	19,552
2002.....	557,271	74	402	537	535	723	1,007	1,258	2,614	5,150	10,935	19,295
2001.....	553,768	66	420	493	515	732	972	1,389	2,605	5,466	11,103	19,199
2000.....	553,091	92	420	489	525	745	968	1,359	2,557	5,565	10,955	18,803
1999.....	549,838	68	418	509	503	745	979	1,429	2,576	5,764	10,968	18,412
All races, male												
2004.....	286,830	37	212	303	269	435	592	679	1,071	1,909	4,499	9,395
2003.....	287,990	34	227	293	298	438	541	664	1,140	2,070	4,755	9,727
2002.....	288,768	42	216	297	303	424	592	678	1,190	2,166	4,890	9,418
2001.....	287,075	31	229	253	280	447	577	694	1,166	2,331	4,966	9,377
2000.....	286,082	50	233	269	300	446	576	692	1,162	2,402	4,942	9,243
1999.....	285,832	36	227	279	276	448	575	755	1,204	2,467	5,004	9,151
All races, female												
2004.....	267,058	37	187	223	224	296	386	607	1,276	2,609	5,706	9,795
2003.....	268,912	41	165	223	262	252	420	643	1,294	2,849	5,835	9,825
2002.....	268,503	32	186	240	232	299	415	580	1,424	2,984	6,045	9,877
2001.....	266,693	35	191	240	235	285	395	695	1,439	3,135	6,137	9,822
2000.....	267,009	42	187	220	225	299	392	667	1,395	3,163	6,013	9,560
1999.....	264,006	32	191	230	227	297	404	674	1,372	3,297	5,964	9,261
White, both sexes												
2004.....	478,134	63	311	415	378	581	742	985	1,798	3,508	7,878	14,977
2003.....	481,556	64	317	427	426	563	761	1,015	1,853	3,818	8,248	15,226
2002.....	482,481	56	306	428	418	574	825	975	2,022	4,009	8,573	15,180
2001.....	479,651	54	353	387	393	587	772	1,074	2,020	4,278	8,681	15,127
2000.....	480,011	75	330	397	421	576	757	1,081	1,991	4,316	8,448	14,743
1999.....	477,238	53	353	397	383	568	759	1,137	2,010	4,495	8,565	14,513

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Malignant neoplasms (C00-C97)												
White, male												
2004.....	247,450	31	161	242	211	350	451	542	859	1,544	3,524	7,365
2003.....	249,053	29	182	248	224	358	432	543	889	1,679	3,743	7,612
2002.....	249,867	35	168	229	230	344	486	531	941	1,710	3,894	7,492
2001.....	248,146	24	189	195	203	360	464	549	944	1,861	3,922	7,421
2000.....	247,403	42	190	215	241	344	454	559	941	1,907	3,832	7,292
1999.....	247,396	31	194	224	210	335	449	606	955	1,984	3,950	7,171
White, female												
2004.....	230,684	32	150	173	167	231	291	443	939	1,964	4,354	7,612
2003.....	232,503	35	135	179	202	205	329	472	964	2,139	4,505	7,614
2002.....	232,614	21	138	199	188	230	339	444	1,081	2,299	4,679	7,688
2001.....	231,505	30	164	192	190	227	308	525	1,076	2,417	4,759	7,706
2000.....	232,608	33	140	182	180	232	303	522	1,050	2,409	4,616	7,451
1999.....	229,842	22	159	173	173	233	310	531	1,055	2,511	4,615	7,342
Black, both sexes												
2004.....	62,499	7	65	79	91	126	185	247	420	793	1,889	3,500
2003.....	62,660	7	57	73	100	94	153	229	438	880	1,937	3,646
2002.....	62,617	12	74	85	87	115	135	212	457	918	1,985	3,449
2001.....	62,170	6	48	80	96	113	149	242	455	959	2,022	3,395
2000.....	61,945	11	70	72	78	136	165	215	434	1,017	2,091	3,437
1999.....	61,951	11	49	85	98	139	179	233	438	1,058	2,022	3,308
Black, male												
2004.....	32,443	2	37	40	44	69	107	105	160	260	775	1,684
2003.....	32,442	4	31	40	56	57	86	90	179	291	836	1,786
2002.....	32,627	4	34	55	51	66	74	103	187	349	823	1,625
2001.....	32,679	3	30	43	59	68	85	107	163	370	847	1,642
2000.....	32,817	4	34	42	45	82	97	94	168	383	926	1,664
1999.....	32,850	3	26	41	55	87	97	114	183	385	882	1,702

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Malignant neoplasms (C00-C97)												
Black, female												
2004.....	30,056	5	28	39	47	57	78	142	260	533	1,114	1,816
2003.....	30,218	3	26	33	44	37	67	139	259	589	1,101	1,860
2002.....	29,990	8	40	30	36	49	61	109	270	569	1,162	1,824
2001.....	29,491	3	18	37	37	45	64	135	292	589	1,175	1,753
2000.....	29,128	7	36	30	33	54	68	121	266	634	1,165	1,773
1999.....	29,101	8	23	44	43	52	82	119	255	673	1,140	1,606
American Indian 1/, both sexes												
2004.....	2,392	1	3	7	1	6	10	7	18	37	69	132
2003.....	2,154	-	7	4	4	9	10	10	14	36	65	114
2002.....	2,175	1	3	6	6	9	11	11	24	39	61	109
2001.....	2,155	3	4	-	5	10	12	16	18	45	69	105
2000.....	1,914	2	3	5	3	6	7	9	17	34	71	88
1999.....	1,836	1	4	2	4	4	9	13	13	28	57	86
American Indian 1/, male												
2004.....	1,241	1	2	6	1	4	5	5	6	15	27	69
2003.....	1,092	-	7	1	1	6	7	4	7	15	29	61
2002.....	1,081	-	2	3	4	4	6	6	8	14	30	44
2001.....	1,103	3	3	-	3	7	8	12	7	16	37	47
2000.....	997	2	2	3	1	4	3	5	8	16	33	44
1999.....	949	-	1	1	4	1	6	7	7	13	22	51
American Indian 1/, female												
2004.....	1,151	-	1	1	-	2	5	2	12	22	42	63
2003.....	1,062	-	-	3	3	3	3	6	7	21	36	53
2002.....	1,094	1	1	3	2	5	5	5	16	25	31	65
2001.....	1,052	-	1	-	2	3	4	4	11	29	32	58
2000.....	917	-	1	2	2	2	4	4	9	18	38	44
1999.....	887	1	3	1	-	3	3	6	6	15	35	35

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Malignant neoplasms (C00-C97)

Asian or Pacific Islander 2/, both sexes

2004.....	10,863	3	20	25	23	18	41	47	111	180	369	581
2003.....	10,532	4	11	12	30	24	37	53	129	185	340	566
2002.....	9,998	5	19	18	24	25	36	60	111	184	316	557
2001.....	9,792	3	15	26	21	22	39	57	112	184	331	572
2000.....	9,221	4	17	15	23	27	39	54	115	198	345	535
1999.....	8,813	3	12	25	18	34	32	46	115	183	324	505

Asian or Pacific Islander 2/, male

2004.....	5,696	3	12	15	13	12	29	27	46	90	173	277
2003.....	5,403	1	7	4	17	17	16	27	65	85	147	268
2002.....	5,193	3	12	10	18	10	26	38	54	93	143	257
2001.....	5,147	1	7	15	15	12	20	26	52	84	160	267
2000.....	4,865	2	7	9	13	16	22	34	45	96	151	243
1999.....	4,637	2	6	13	7	25	23	28	59	85	150	227

Asian or Pacific Islander 2/, female

2004.....	5,167	-	8	10	10	6	12	20	65	90	196	304
2003.....	5,129	3	4	8	13	7	21	26	64	100	193	298
2002.....	4,805	2	7	8	6	15	10	22	57	91	173	300
2001.....	4,645	2	8	11	6	10	19	31	60	100	171	305
2000.....	4,356	2	10	6	10	11	17	20	70	102	194	292
1999.....	4,176	1	6	12	11	9	9	18	56	98	174	278

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)

All races, both sexes

2004.....	7,826	-	2	1	4	10	13	13	30	56	200	450
2003.....	7,778	-	4	4	4	11	19	19	40	64	210	445
2002.....	7,737	-	3	1	7	12	11	22	35	77	181	431
2001.....	7,702	-	-	2	2	10	19	19	37	93	190	427
2000.....	7,492	-	-	-	4	9	17	18	32	88	205	408
1999.....	7,486	-	2	1	5	9	18	14	29	96	195	449

All races, male

2004.....	5,312	-	-	-	2	8	10	11	24	33	143	346
2003.....	5,298	-	2	1	3	8	13	11	28	40	155	346
2002.....	5,184	-	2	1	2	7	4	14	25	47	132	336
2001.....	5,156	-	-	1	-	3	9	12	23	61	141	325
2000.....	4,952	-	-	-	4	6	7	12	16	60	155	309
1999.....	4,944	-	1	-	3	6	14	10	17	58	146	346

All races, female

2004.....	2,514	-	2	1	2	2	3	2	6	23	57	104
2003.....	2,480	-	2	3	1	3	6	8	12	24	55	99
2002.....	2,553	-	1	-	5	5	7	8	10	30	49	95
2001.....	2,546	-	-	1	2	7	10	7	14	32	49	102
2000.....	2,540	-	-	-	-	3	10	6	16	28	50	99
1999.....	2,542	-	1	1	2	3	4	4	12	38	49	103

White, both sexes

2004.....	6,443	-	2	1	2	7	7	9	20	34	128	306
2003.....	6,409	-	3	4	1	5	14	14	27	47	149	322
2002.....	6,432	-	2	1	4	8	7	14	25	52	128	304
2001.....	6,369	-	-	1	1	7	14	14	23	55	123	296
2000.....	6,135	-	-	-	2	6	12	12	23	60	138	291
1999.....	6,116	-	2	1	3	5	9	10	20	60	134	287

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)

White, male

2004.....	4,270	-	-	-	-	6	4	9	16	21	88	236
2003.....	4,282	-	1	1	1	4	9	8	17	32	121	243
2002.....	4,256	-	2	1	1	5	3	9	19	35	92	240
2001.....	4,183	-	-	1	-	1	6	10	13	37	93	227
2000.....	3,942	-	-	-	2	3	4	8	12	38	100	222
1999.....	3,955	-	1	-	2	3	6	6	11	34	100	218

White, female

2004.....	2,173	-	2	1	2	1	3	-	4	13	40	70
2003.....	2,127	-	2	3	-	1	5	6	10	15	28	79
2002.....	2,176	-	-	-	3	3	4	5	6	17	36	64
2001.....	2,186	-	-	-	1	6	8	4	10	18	30	69
2000.....	2,193	-	-	-	-	3	8	4	11	22	38	69
1999.....	2,161	-	1	1	1	2	3	4	9	26	34	69

Black, both sexes

2004.....	1,149	-	-	-	2	1	4	3	8	19	57	123
2003.....	1,107	-	1	-	3	5	4	4	7	14	49	104
2002.....	1,044	-	-	-	2	4	2	3	8	22	43	106
2001.....	1,065	-	-	1	-	1	3	2	4	24	48	106
2000.....	1,102	-	-	-	2	3	4	5	2	15	46	96
1999.....	1,145	-	-	-	2	4	4	2	6	25	42	135

Black, male

2004.....	874	-	-	-	2	1	4	2	7	11	40	96
2003.....	837	-	1	-	2	3	3	3	7	6	26	86
2002.....	766	-	-	-	1	2	1	2	5	10	33	80
2001.....	782	-	-	-	-	1	1	1	4	14	32	80
2000.....	833	-	-	-	2	3	2	3	1	11	36	69
1999.....	828	-	-	-	1	3	3	2	4	17	30	109

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)												
Asian or Pacific Islander 2/, both sexes												
2004.....	195	-	-	-	-	1	2	1	2	1	12	19
2003.....	229	-	-	-	-	1	1	1	6	3	12	18
2002.....	218	-	1	-	1	-	2	4	2	3	8	20
2001.....	228	-	-	-	1	2	2	3	8	12	16	24
2000.....	214	-	-	-	-	-	1	1	7	11	19	21
1999.....	192	-	-	-	-	-	5	1	3	11	18	24
Asian or Pacific Islander 2/, male												
2004.....	137	-	-	-	-	-	2	-	1	-	12	12
2003.....	162	-	-	-	-	1	1	-	4	2	8	16
2002.....	135	-	-	-	-	-	-	2	1	2	6	15
2001.....	158	-	-	-	-	1	2	1	5	8	14	17
2000.....	150	-	-	-	-	-	1	1	3	9	17	18
1999.....	139	-	-	-	-	-	5	1	2	7	15	16
Asian or Pacific Islander 2/, female												
2004.....	58	-	-	-	-	1	-	1	1	1	-	7
2003.....	67	-	-	-	-	-	-	1	2	1	4	2
2002.....	83	-	1	-	1	-	2	2	1	1	2	5
2001.....	70	-	-	-	1	1	-	2	3	4	2	7
2000.....	64	-	-	-	-	-	-	-	4	2	2	3
1999.....	53	-	-	-	-	-	-	-	1	4	3	8

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Of tongue (C01-C02)												
All races, both sexes												
2004.....	1,881	-	-	-	-	-	6	3	12	27	56	111
2003.....	1,875	-	-	-	-	1	4	10	12	18	58	119
2002.....	1,887	-	1	-	-	1	3	8	17	27	52	104
2001.....	1,818	-	-	-	1	2	3	11	17	28	51	110
2000.....	1,767	-	-	-	-	1	5	7	15	31	59	103
1999.....	1,738	-	-	-	-	-	4	9	9	27	66	114
All races, male												
2004.....	1,228	-	-	-	-	-	4	3	7	15	34	74
2003.....	1,241	-	-	-	-	-	3	6	7	10	41	91
2002.....	1,220	-	1	-	-	-	1	5	11	24	38	77
2001.....	1,191	-	-	-	-	-	2	6	10	19	34	78
2000.....	1,115	-	-	-	-	1	1	4	7	18	38	75
1999.....	1,138	-	-	-	-	-	3	5	4	12	46	87
All races, female												
2004.....	653	-	-	-	-	-	2	-	5	12	22	37
2003.....	634	-	-	-	-	1	1	4	5	8	17	28
2002.....	667	-	-	-	-	1	2	3	6	3	14	27
2001.....	627	-	-	-	1	2	1	5	7	9	17	32
2000.....	652	-	-	-	-	-	4	3	8	13	21	28
1999.....	600	-	-	-	-	-	1	4	5	15	20	27
White, both sexes												
2004.....	1,633	-	-	-	-	-	5	3	10	20	38	84
2003.....	1,608	-	-	-	-	1	4	9	10	16	43	94
2002.....	1,621	-	1	-	-	1	3	6	13	20	42	73
2001.....	1,569	-	-	-	-	2	2	10	15	18	37	85
2000.....	1,522	-	-	-	-	1	5	6	14	26	44	87
1999.....	1,478	-	-	-	-	-	3	8	9	19	55	80

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Of tongue (C01-C02)												
White, male												
2004.....	1,049	-	-	-	-	-	3	3	6	13	22	59
2003.....	1,061	-	-	-	-	-	3	6	5	9	36	69
2002.....	1,032	-	1	-	-	-	1	4	10	18	29	55
2001.....	1,014	-	-	-	-	-	1	6	9	11	25	57
2000.....	939	-	-	-	-	1	1	4	6	15	28	61
1999.....	957	-	-	-	-	-	2	4	4	7	39	59
White, female												
2004.....	584	-	-	-	-	-	2	-	4	7	16	25
2003.....	547	-	-	-	-	1	1	3	5	7	7	25
2002.....	589	-	-	-	-	1	2	2	3	2	13	18
2001.....	555	-	-	-	-	2	1	4	6	7	12	28
2000.....	583	-	-	-	-	-	4	2	8	11	16	26
1999.....	521	-	-	-	-	-	1	4	5	12	16	21
Black, both sexes												
2004.....	204	-	-	-	-	-	-	-	1	6	13	24
2003.....	215	-	-	-	-	-	-	1	1	2	14	23
2002.....	212	-	-	-	-	-	-	-	3	7	8	26
2001.....	199	-	-	-	-	-	-	-	-	7	10	24
2000.....	205	-	-	-	-	-	-	1	-	2	14	14
1999.....	224	-	-	-	-	-	-	-	-	5	8	30
Black, male												
2004.....	152	-	-	-	-	-	-	-	1	2	7	15
2003.....	153	-	-	-	-	-	-	-	1	1	5	20
2002.....	157	-	-	-	-	-	-	-	1	6	7	18
2001.....	150	-	-	-	-	-	-	-	-	5	6	20
2000.....	157	-	-	-	-	-	-	-	-	1	9	12
1999.....	157	-	-	-	-	-	-	-	-	3	4	24

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Of tongue (C01-C02)												
Asian or Pacific Islander 2/, both sexes												
2004.....	36	-	-	-	-	-	1	-	1	-	3	3
2003.....	44	-	-	-	-	-	-	-	1	-	1	2
2002.....	41	-	-	-	-	-	-	1	1	-	1	4
2001.....	43	-	-	-	1	-	1	1	2	1	4	1
2000.....	35	-	-	-	-	-	-	-	1	2	1	2
1999.....	27	-	-	-	-	-	1	-	-	3	3	4
Asian or Pacific Islander 2/, male												
2004.....	21	-	-	-	-	-	1	-	-	-	3	-
2003.....	23	-	-	-	-	-	-	-	1	-	-	2
2002.....	22	-	-	-	-	-	-	-	-	-	1	3
2001.....	21	-	-	-	-	-	1	-	1	1	3	1
2000.....	17	-	-	-	-	-	-	-	1	1	1	2
1999.....	20	-	-	-	-	-	1	-	-	2	3	4
Asian or Pacific Islander 2/, female												
2004.....	15	-	-	-	-	-	-	-	1	-	-	3
2003.....	21	-	-	-	-	-	-	-	-	-	1	-
2002.....	19	-	-	-	-	-	-	1	1	-	-	1
2001.....	22	-	-	-	1	-	-	1	1	-	1	-
2000.....	18	-	-	-	-	-	-	-	-	1	-	-
1999.....	7	-	-	-	-	-	-	-	-	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Of pharynx (C10-C13,C14.0)

All races, both sexes

2004.....	3,292	-	1	-	-	4	2	6	9	12	89	218
2003.....	3,203	-	1	1	3	6	9	6	18	26	77	177
2002.....	3,115	-	1	-	4	6	7	9	8	29	74	177
2001.....	3,194	-	-	-	-	3	8	8	11	40	83	191
2000.....	3,111	-	-	-	3	6	9	8	11	29	81	163
1999.....	3,102	-	1	-	4	3	11	3	11	38	81	191

All races, male

2004.....	2,400	-	-	-	-	4	2	4	9	7	66	176
2003.....	2,342	-	1	1	2	4	7	3	14	20	59	141
2002.....	2,219	-	-	-	1	4	2	7	7	14	55	142
2001.....	2,288	-	-	-	-	1	5	6	8	27	66	152
2000.....	2,223	-	-	-	3	3	5	5	6	24	67	128
1999.....	2,210	-	1	-	3	2	9	3	6	26	62	152

All races, female

2004.....	892	-	1	-	-	-	-	2	-	5	23	42
2003.....	861	-	-	-	1	2	2	3	4	6	18	36
2002.....	896	-	1	-	3	2	5	2	1	15	19	35
2001.....	906	-	-	-	-	2	3	2	3	13	17	39
2000.....	888	-	-	-	-	3	4	3	5	5	14	35
1999.....	892	-	-	-	1	1	2	-	5	12	19	39

White, both sexes

2004.....	2,517	-	1	-	-	2	1	2	6	3	52	139
2003.....	2,458	-	1	1	1	-	4	3	9	18	47	108
2002.....	2,412	-	-	-	1	4	3	4	5	15	41	112
2001.....	2,468	-	-	-	-	1	5	4	2	20	44	115
2000.....	2,351	-	-	-	1	3	5	3	5	13	39	97
1999.....	2,362	-	1	-	2	1	4	1	3	22	44	105

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Of pharynx (C10-C13,C14.0)												
White, male												
2004.....	1,804	-	-	-	-	2	1	2	6	2	38	113
2003.....	1,759	-	1	1	1	-	3	1	6	15	39	82
2002.....	1,710	-	-	-	-	3	1	3	5	9	32	90
2001.....	1,736	-	-	-	-	-	3	4	1	15	35	97
2000.....	1,637	-	-	-	1	-	2	1	3	9	31	77
1999.....	1,653	-	1	-	2	1	3	1	1	14	32	83
White, female												
2004.....	713	-	1	-	-	-	-	-	-	1	14	26
2003.....	699	-	-	-	-	-	1	2	3	3	8	26
2002.....	702	-	-	-	1	1	2	1	-	6	9	22
2001.....	732	-	-	-	-	1	2	-	1	5	9	18
2000.....	714	-	-	-	-	3	3	2	2	4	8	20
1999.....	709	-	-	-	-	-	1	-	2	8	12	22
Black, both sexes												
2004.....	643	-	-	-	-	1	-	3	2	8	29	62
2003.....	584	-	-	-	2	5	4	2	6	6	20	54
2002.....	534	-	-	-	2	2	2	3	3	11	25	50
2001.....	549	-	-	-	-	1	2	2	2	11	26	53
2000.....	596	-	-	-	2	3	3	4	1	9	23	50
1999.....	596	-	-	-	2	2	4	1	6	8	23	68
Black, male												
2004.....	493	-	-	-	-	1	-	2	2	4	20	50
2003.....	464	-	-	-	1	3	3	2	6	3	12	46
2002.....	398	-	-	-	1	1	1	2	2	3	18	40
2001.....	416	-	-	-	-	1	1	1	2	7	19	39
2000.....	461	-	-	-	2	3	2	3	1	9	19	38
1999.....	449	-	-	-	1	1	3	1	4	7	18	57

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Of pharynx (C10-C13,C14.0)												
Asian or Pacific Islander 2/, both sexes												
2004.....	113	-	-	-	-	-	1	1	1	-	8	15
2003.....	143	-	-	-	-	1	1	1	3	2	10	15
2002.....	149	-	1	-	1	-	2	2	-	3	7	15
2001.....	156	-	-	-	-	1	1	2	6	9	12	22
2000.....	142	-	-	-	-	-	1	1	5	7	17	16
1999.....	130	-	-	-	-	-	3	1	2	8	13	17
Asian or Pacific Islander 2/, male												
2004.....	86	-	-	-	-	-	1	-	1	-	8	11
2003.....	111	-	-	-	-	1	1	-	2	2	8	13
2002.....	99	-	-	-	-	-	-	2	-	2	5	12
2001.....	118	-	-	-	-	-	1	1	4	5	11	15
2000.....	109	-	-	-	-	-	1	1	2	6	15	13
1999.....	98	-	-	-	-	-	3	1	1	5	11	11
Asian or Pacific Islander 2/, female												
2004.....	27	-	-	-	-	-	-	1	-	-	-	4
2003.....	32	-	-	-	-	-	-	1	1	-	2	2
2002.....	50	-	1	-	1	-	2	-	-	1	2	3
2001.....	38	-	-	-	-	1	-	1	2	4	1	7
2000.....	33	-	-	-	-	-	-	-	3	1	2	3
1999.....	32	-	-	-	-	-	-	-	1	3	2	6

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Of other and unspecified sites within the lip, oral cavity and pharynx (C03-C09,C14.2-C14.8)

All races, both sexes

2004.....	2,579	-	1	1	4	6	5	3	9	16	55	117
2003.....	2,628	-	3	3	1	4	6	3	10	20	73	148
2002.....	2,661	-	-	1	3	5	1	4	9	20	52	146
2001.....	2,613	-	-	2	1	5	8	-	8	24	53	122
2000.....	2,547	-	-	-	1	2	3	3	6	27	63	140
1999.....	2,594	-	1	1	1	6	3	2	9	31	48	143

All races, male

2004.....	1,635	-	-	-	2	4	4	3	8	10	43	94
2003.....	1,663	-	1	-	1	4	3	2	7	10	53	113
2002.....	1,691	-	-	1	1	3	1	1	6	9	36	113
2001.....	1,617	-	-	1	-	2	2	-	4	14	39	91
2000.....	1,561	-	-	-	1	2	1	3	3	17	48	104
1999.....	1,560	-	-	-	-	4	2	2	7	20	38	106

All races, female

2004.....	944	-	1	1	2	2	1	-	1	6	12	23
2003.....	965	-	2	3	-	-	3	1	3	10	20	35
2002.....	970	-	-	-	2	2	-	3	3	11	16	33
2001.....	996	-	-	1	1	3	6	-	4	10	14	31
2000.....	986	-	-	-	-	-	2	-	3	10	15	36
1999.....	1,034	-	1	1	1	2	1	-	2	11	10	37

White, both sexes

2004.....	2,224	-	1	1	2	5	1	3	4	10	38	79
2003.....	2,273	-	2	3	-	4	6	2	8	13	57	119
2002.....	2,330	-	-	1	3	3	1	3	6	16	42	115
2001.....	2,257	-	-	1	1	4	7	-	5	16	39	92
2000.....	2,197	-	-	-	1	2	2	3	4	20	54	105
1999.....	2,225	-	1	1	1	4	2	1	8	19	35	101

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Of other and unspecified sites within the lip, oral cavity and pharynx (C03-C09,C14.2-C14.8)

White, male

2004.....	1,371	-	-	-	-	4	-	3	4	5	28	62
2003.....	1,411	-	-	-	-	4	3	1	6	8	44	91
2002.....	1,462	-	-	1	1	2	1	1	3	8	28	91
2001.....	1,374	-	-	1	-	1	2	-	2	10	31	69
2000.....	1,314	-	-	-	1	2	1	3	3	13	40	82
1999.....	1,309	-	-	-	-	2	1	1	6	13	29	75

White, female

2004.....	853	-	1	1	2	1	1	-	-	5	10	17
2003.....	862	-	2	3	-	-	3	1	2	5	13	28
2002.....	868	-	-	-	2	1	-	2	3	8	14	24
2001.....	883	-	-	-	1	3	5	-	3	6	8	23
2000.....	883	-	-	-	-	-	1	-	1	7	14	23
1999.....	916	-	1	1	1	2	1	-	2	6	6	26

Black, both sexes

2004.....	299	-	-	-	2	-	4	-	5	5	15	37
2003.....	307	-	1	-	1	-	-	1	-	6	15	27
2002.....	294	-	-	-	-	2	-	-	2	4	10	30
2001.....	315	-	-	1	-	-	1	-	2	6	12	29
2000.....	299	-	-	-	-	-	1	-	1	4	8	32
1999.....	325	-	-	-	-	2	-	1	-	12	11	37

Black, male

2004.....	227	-	-	-	2	-	4	-	4	5	13	31
2003.....	219	-	1	-	1	-	-	1	-	2	9	20
2002.....	210	-	-	-	-	1	-	-	2	1	8	22
2001.....	215	-	-	-	-	-	-	-	2	2	7	21
2000.....	214	-	-	-	-	-	-	-	-	1	7	19
1999.....	222	-	-	-	-	2	-	1	-	7	8	28

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Malignant neoplasms of digestive organs (C15-C26)												
All races, both sexes												
2004.....	131,331	1	22	7	11	33	98	175	431	906	2,210	4,533
2003.....	132,234	6	23	7	14	28	89	192	428	994	2,130	4,622
2002.....	131,725	5	30	7	14	28	91	190	421	999	2,313	4,486
2001.....	130,982	5	19	8	10	34	71	167	432	1,108	2,222	4,495
2000.....	130,702	5	24	9	17	33	62	176	424	1,019	2,238	4,354
1999.....	129,441	10	29	5	8	36	77	192	412	1,045	2,271	4,287
All races, male												
2004.....	71,958	1	12	3	7	20	58	106	245	529	1,347	2,958
2003.....	72,034	2	18	5	10	19	46	109	237	587	1,302	3,000
2002.....	71,799	3	21	4	9	17	51	112	243	579	1,448	2,889
2001.....	70,559	3	8	4	4	19	38	90	247	668	1,313	2,877
2000.....	70,066	3	15	6	5	20	38	103	256	627	1,370	2,805
1999.....	69,306	5	20	5	6	20	41	118	237	608	1,377	2,803
All races, female												
2004.....	59,373	-	10	4	4	13	40	69	186	377	863	1,575
2003.....	60,200	4	5	2	4	9	43	83	191	407	828	1,622
2002.....	59,926	2	9	3	5	11	40	78	178	420	865	1,597
2001.....	60,423	2	11	4	6	15	33	77	185	440	909	1,618
2000.....	60,636	2	9	3	12	13	24	73	168	392	868	1,549
1999.....	60,135	5	9	-	2	16	36	74	175	437	894	1,484
White, both sexes												
2004.....	110,236	1	19	3	9	22	62	121	305	678	1,628	3,382
2003.....	111,150	4	19	5	8	23	61	140	297	732	1,546	3,427
2002.....	111,031	3	24	7	9	20	71	136	308	751	1,720	3,383
2001.....	110,565	5	15	3	8	23	53	120	312	807	1,653	3,374
2000.....	110,555	3	18	8	10	26	44	120	304	743	1,633	3,271
1999.....	109,699	6	26	3	4	23	52	128	290	768	1,667	3,267

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Malignant neoplasms of digestive organs (C15-C26)												
White, male												
2004.....	60,546	1	11	2	5	13	34	71	180	415	1,006	2,245
2003.....	60,742	2	15	4	7	15	34	80	162	445	956	2,242
2002.....	60,665	3	16	4	4	11	36	77	174	426	1,094	2,224
2001.....	59,748	3	6	1	4	15	26	62	180	491	982	2,184
2000.....	59,305	2	12	5	4	15	25	72	187	463	1,005	2,131
1999.....	58,795	3	18	3	3	14	27	80	169	455	1,032	2,146
White, female												
2004.....	49,690	-	8	1	4	9	28	50	125	263	622	1,137
2003.....	50,408	2	4	1	1	8	27	60	135	287	590	1,185
2002.....	50,366	-	8	3	5	9	35	59	134	325	626	1,159
2001.....	50,817	2	9	2	4	8	27	58	132	316	671	1,190
2000.....	51,250	1	6	3	6	11	19	48	117	280	628	1,140
1999.....	50,904	3	8	-	1	9	25	48	121	313	635	1,121
Black, both sexes												
2004.....	16,494	-	2	4	1	10	27	39	86	151	436	910
2003.....	16,658	2	1	2	3	4	17	31	86	193	452	941
2002.....	16,470	2	4	-	3	7	15	38	76	172	471	885
2001.....	16,394	-	3	3	2	5	12	31	86	226	450	903
2000.....	16,298	2	2	-	3	5	12	35	90	204	471	892
1999.....	16,146	3	3	1	4	8	19	48	85	212	469	827
Black, male												
2004.....	8,778	-	-	1	1	7	19	22	45	75	251	557
2003.....	8,813	-	-	1	3	3	9	17	48	99	268	601
2002.....	8,724	-	3	-	3	5	10	24	47	104	274	529
2001.....	8,511	-	2	2	-	2	8	18	48	131	257	553
2000.....	8,561	1	1	-	1	3	9	15	54	118	284	550
1999.....	8,488	1	2	1	3	3	10	30	48	110	273	533

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Malignant neoplasms of digestive organs (C15-C26)												
Black, female												
2004.....	7,716	-	2	3	-	3	8	17	41	76	185	353
2003.....	7,845	2	1	1	-	1	8	14	38	94	184	340
2002.....	7,746	2	1	-	-	2	5	14	29	68	197	356
2001.....	7,883	-	1	1	2	3	4	13	38	95	193	350
2000.....	7,737	1	1	-	2	2	3	20	36	86	187	342
1999.....	7,658	2	1	-	1	5	9	18	37	102	196	294
American Indian 1/, both sexes												
2004.....	688	-	-	-	-	-	1	3	3	8	18	38
2003.....	589	-	2	-	1	-	1	2	4	10	20	48
2002.....	619	-	1	-	-	-	-	2	4	9	17	32
2001.....	560	-	-	-	-	1	1	1	4	9	16	24
2000.....	529	-	-	1	-	-	-	2	3	6	26	18
1999.....	473	-	-	-	-	1	2	1	4	7	13	24
American Indian 1/, male												
2004.....	388	-	-	-	-	-	1	3	3	4	8	26
2003.....	341	-	2	-	-	-	-	1	2	5	10	31
2002.....	335	-	1	-	-	-	-	1	2	7	11	19
2001.....	311	-	-	-	-	-	1	1	1	6	7	14
2000.....	305	-	-	1	-	-	-	1	1	2	20	15
1999.....	257	-	-	-	-	-	2	-	3	4	6	18
American Indian 1/, female												
2004.....	300	-	-	-	-	-	-	-	-	4	10	12
2003.....	248	-	-	-	1	-	1	1	2	5	10	17
2002.....	284	-	-	-	-	-	-	1	2	2	6	13
2001.....	249	-	-	-	-	1	-	-	3	3	9	10
2000.....	224	-	-	-	-	-	-	1	2	4	6	3
1999.....	216	-	-	-	-	1	-	1	1	3	7	6

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Malignant neoplasms of digestive organs (C15-C26)												
Asian or Pacific Islander 2/, both sexes												
2004.....	3,913	-	1	-	1	1	8	12	37	69	128	203
2003.....	3,837	-	1	-	2	1	10	19	41	59	112	206
2002.....	3,605	-	1	-	2	1	5	14	33	67	105	186
2001.....	3,463	-	1	2	-	5	5	15	30	66	103	194
2000.....	3,320	-	4	-	4	2	6	19	27	66	108	173
1999.....	3,123	1	-	1	-	4	4	15	33	58	122	169
Asian or Pacific Islander 2/, male												
2004.....	2,246	-	1	-	1	-	4	10	17	35	82	130
2003.....	2,138	-	1	-	-	1	3	11	25	38	68	126
2002.....	2,075	-	1	-	2	1	5	10	20	42	69	117
2001.....	1,989	-	-	1	-	2	3	9	18	40	67	126
2000.....	1,895	-	2	-	-	2	4	15	14	44	61	109
1999.....	1,766	1	-	1	-	3	2	8	17	39	66	106
Asian or Pacific Islander 2/, female												
2004.....	1,667	-	-	-	-	1	4	2	20	34	46	73
2003.....	1,699	-	-	-	2	-	7	8	16	21	44	80
2002.....	1,530	-	-	-	-	-	-	4	13	25	36	69
2001.....	1,474	-	1	1	-	3	2	6	12	26	36	68
2000.....	1,425	-	2	-	4	-	2	4	13	22	47	64
1999.....	1,357	-	-	-	-	1	2	7	16	19	56	63

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Of esophagus (C15)												
All races, both sexes												
2004.....	13,023	-	-	-	-	-	5	11	25	69	222	513
2003.....	12,860	-	-	-	-	2	3	11	18	76	212	526
2002.....	12,701	-	-	-	1	-	6	4	15	84	227	503
2001.....	12,530	-	-	-	-	2	4	9	20	68	241	492
2000.....	12,232	1	-	-	-	1	2	9	30	78	219	478
1999.....	11,918	-	-	-	-	-	-	10	27	75	218	469
All races, male												
2004.....	10,081	-	-	-	-	-	4	8	20	58	184	440
2003.....	9,921	-	-	-	-	2	2	8	15	66	174	445
2002.....	9,789	-	-	-	-	-	3	4	10	71	186	426
2001.....	9,607	-	-	-	-	1	4	8	18	52	203	417
2000.....	9,279	1	-	-	-	-	2	7	24	69	196	415
1999.....	9,094	-	-	-	-	-	-	7	22	65	190	410
All races, female												
2004.....	2,942	-	-	-	-	-	1	3	5	11	38	73
2003.....	2,939	-	-	-	-	-	1	3	3	10	38	81
2002.....	2,912	-	-	-	1	-	3	-	5	13	41	77
2001.....	2,923	-	-	-	-	1	-	1	2	16	38	75
2000.....	2,953	-	-	-	-	1	-	2	6	9	23	63
1999.....	2,824	-	-	-	-	-	-	3	5	10	28	59
White, both sexes												
2004.....	11,251	-	-	-	-	-	3	9	18	62	184	398
2003.....	11,058	-	-	-	-	2	3	9	16	65	174	406
2002.....	10,910	-	-	-	1	-	6	3	14	73	178	403
2001.....	10,624	-	-	-	-	2	4	7	17	59	191	389
2000.....	10,406	-	-	-	-	1	2	9	29	74	175	377
1999.....	10,079	-	-	-	-	-	-	9	23	67	162	362

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Of esophagus (C15)												
White, male												
2004.....	8,831	-	-	-	-	-	3	6	15	54	159	355
2003.....	8,617	-	-	-	-	2	2	7	14	58	146	349
2002.....	8,510	-	-	-	-	-	3	3	10	65	151	353
2001.....	8,205	-	-	-	-	1	4	6	15	45	164	338
2000.....	7,986	-	-	-	-	-	2	7	24	67	162	341
1999.....	7,740	-	-	-	-	-	-	6	18	60	143	323
White, female												
2004.....	2,420	-	-	-	-	-	-	3	3	8	25	43
2003.....	2,441	-	-	-	-	-	1	2	2	7	28	57
2002.....	2,400	-	-	-	1	-	3	-	4	8	27	50
2001.....	2,419	-	-	-	-	1	-	1	2	14	27	51
2000.....	2,420	-	-	-	-	1	-	2	5	7	13	36
1999.....	2,339	-	-	-	-	-	-	3	5	7	19	39
Black, both sexes												
2004.....	1,527	-	-	-	-	-	2	1	5	5	34	107
2003.....	1,591	-	-	-	-	-	-	1	2	10	37	109
2002.....	1,590	-	-	-	-	-	-	1	1	9	47	93
2001.....	1,688	-	-	-	-	-	-	2	1	9	46	92
2000.....	1,655	1	-	-	-	-	-	-	1	3	41	96
1999.....	1,648	-	-	-	-	-	-	-	3	6	48	99
Black, male												
2004.....	1,060	-	-	-	-	-	1	1	4	3	23	79
2003.....	1,133	-	-	-	-	-	-	1	1	8	28	87
2002.....	1,133	-	-	-	-	-	-	1	-	4	34	68
2001.....	1,231	-	-	-	-	-	-	2	1	7	35	69
2000.....	1,170	1	-	-	-	-	-	-	-	1	31	69
1999.....	1,203	-	-	-	-	-	-	-	3	4	41	79

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Of esophagus (C15)												
Asian or Pacific Islander 2/, both sexes												
2004.....	187	-	-	-	-	-	-	1	2	1	2	5
2003.....	162	-	-	-	-	-	-	1	-	1	1	5
2002.....	154	-	-	-	-	-	-	-	-	2	-	6
2001.....	166	-	-	-	-	-	-	-	2	-	1	9
2000.....	128	-	-	-	-	-	-	-	-	1	1	2
1999.....	153	-	-	-	-	-	-	1	1	2	7	5
Asian or Pacific Islander 2/, male												
2004.....	145	-	-	-	-	-	-	1	1	1	1	4
2003.....	133	-	-	-	-	-	-	-	-	-	-	4
2002.....	109	-	-	-	-	-	-	-	-	2	-	4
2001.....	133	-	-	-	-	-	-	-	2	-	1	8
2000.....	87	-	-	-	-	-	-	-	-	1	1	2
1999.....	120	-	-	-	-	-	-	1	1	1	5	5
Asian or Pacific Islander 2/, female												
2004.....	42	-	-	-	-	-	-	-	1	-	1	1
2003.....	29	-	-	-	-	-	-	1	-	1	1	1
2002.....	45	-	-	-	-	-	-	-	-	-	-	2
2001.....	33	-	-	-	-	-	-	-	-	-	-	1
2000.....	41	-	-	-	-	-	-	-	-	-	-	-
1999.....	33	-	-	-	-	-	-	-	-	1	2	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Of stomach (C16)

All races, both sexes

2004.....	11,859	-	-	-	-	2	10	29	93	145	333	457
2003.....	12,110	-	-	-	1	5	14	34	93	178	302	493
2002.....	12,198	-	1	-	-	7	22	48	90	160	276	443
2001.....	12,319	-	-	-	-	5	15	33	89	178	286	452
2000.....	12,645	-	-	-	1	3	11	36	88	167	317	468
1999.....	12,711	-	-	-	1	5	8	47	89	189	316	450

All races, male

2004.....	6,958	-	-	-	-	1	6	17	46	78	191	280
2003.....	7,055	-	-	-	1	3	4	16	47	91	171	305
2002.....	7,133	-	1	-	-	-	8	31	49	82	157	285
2001.....	7,191	-	-	-	-	4	5	20	51	98	155	288
2000.....	7,328	-	-	-	-	-	6	22	47	100	184	290
1999.....	7,459	-	-	-	1	2	4	22	54	101	182	303

All races, female

2004.....	4,901	-	-	-	-	1	4	12	47	67	142	177
2003.....	5,055	-	-	-	-	2	10	18	46	87	131	188
2002.....	5,065	-	-	-	-	7	14	17	41	78	119	158
2001.....	5,128	-	-	-	-	1	10	13	38	80	131	164
2000.....	5,317	-	-	-	1	3	5	14	41	67	133	178
1999.....	5,252	-	-	-	-	3	4	25	35	88	134	147

White, both sexes

2004.....	9,067	-	-	-	-	2	5	17	65	116	218	309
2003.....	9,282	-	-	-	1	5	10	24	64	123	200	330
2002.....	9,319	-	1	-	-	5	16	33	61	119	182	317
2001.....	9,501	-	-	-	-	2	8	25	59	112	198	321
2000.....	9,723	-	-	-	1	2	7	21	57	113	205	314
1999.....	9,779	-	-	-	-	2	6	29	59	128	193	320

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	-------------	-----------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Of stomach (C16)

White, male

2004.....	5,332	-	-	-	-	1	4	8	38	64	129	192
2003.....	5,428	-	-	-	1	3	4	12	34	71	112	206
2002.....	5,542	-	1	-	-	-	5	21	36	60	108	209
2001.....	5,568	-	-	-	-	2	2	14	34	67	109	207
2000.....	5,688	-	-	-	-	-	3	13	32	66	115	192
1999.....	5,786	-	-	-	-	1	3	15	40	70	112	222

White, female

2004.....	3,735	-	-	-	-	1	1	9	27	52	89	117
2003.....	3,854	-	-	-	-	2	6	12	30	52	88	124
2002.....	3,777	-	-	-	-	5	11	12	25	59	74	108
2001.....	3,933	-	-	-	-	-	6	11	25	45	89	114
2000.....	4,035	-	-	-	1	2	4	8	25	47	90	122
1999.....	3,993	-	-	-	-	1	3	14	19	58	81	98

Black, both sexes

2004.....	2,034	-	-	-	-	-	4	9	19	14	79	106
2003.....	2,041	-	-	-	-	-	1	6	16	34	69	114
2002.....	2,131	-	-	-	-	2	6	13	20	27	65	92
2001.....	2,067	-	-	-	-	1	4	4	21	49	72	90
2000.....	2,176	-	-	-	-	1	3	11	25	37	81	110
1999.....	2,203	-	-	-	1	1	2	14	22	40	87	89

Black, male

2004.....	1,191	-	-	-	-	-	2	6	6	7	43	67
2003.....	1,187	-	-	-	-	-	-	3	7	10	45	76
2002.....	1,194	-	-	-	-	-	3	10	10	16	33	60
2001.....	1,186	-	-	-	-	1	1	3	11	25	38	57
2000.....	1,218	-	-	-	-	-	2	5	15	26	52	72
1999.....	1,263	-	-	-	1	-	1	7	12	20	52	60

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Of stomach (C16)												
Black, female												
2004.....	843	-	-	-	-	-	2	3	13	7	36	39
2003.....	854	-	-	-	-	-	1	3	9	24	24	38
2002.....	937	-	-	-	-	2	3	3	10	11	32	32
2001.....	881	-	-	-	-	-	3	1	10	24	34	33
2000.....	958	-	-	-	-	1	1	6	10	11	29	38
1999.....	940	-	-	-	-	1	1	7	10	20	35	29
American Indian 1/, both sexes												
2004.....	78	-	-	-	-	-	-	-	-	3	3	8
2003.....	84	-	-	-	-	-	1	1	3	1	6	8
2002.....	91	-	-	-	-	-	-	-	1	1	5	6
2001.....	79	-	-	-	-	-	1	1	1	1	-	4
2000.....	73	-	-	-	-	-	-	-	-	1	3	-
1999.....	67	-	-	-	-	1	-	-	1	2	3	3
American Indian 1/, male												
2004.....	46	-	-	-	-	-	-	-	-	2	-	4
2003.....	50	-	-	-	-	-	-	1	1	-	2	4
2002.....	45	-	-	-	-	-	-	-	1	1	3	3
2001.....	51	-	-	-	-	-	1	1	1	1	-	2
2000.....	41	-	-	-	-	-	-	-	-	-	3	-
1999.....	44	-	-	-	-	-	-	-	1	1	2	2
American Indian 1/, female												
2004.....	32	-	-	-	-	-	-	-	-	1	3	4
2003.....	34	-	-	-	-	-	1	-	2	1	4	4
2002.....	46	-	-	-	-	-	-	-	-	-	2	3
2001.....	28	-	-	-	-	-	-	-	-	-	-	2
2000.....	32	-	-	-	-	-	-	-	-	1	-	-
1999.....	23	-	-	-	-	1	-	-	-	1	1	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Of stomach (C16)												
Asian or Pacific Islander 2/, both sexes												
2004.....	680	-	-	-	-	-	1	3	9	12	33	34
2003.....	703	-	-	-	-	-	2	3	10	20	27	41
2002.....	657	-	-	-	-	-	-	2	8	13	24	28
2001.....	672	-	-	-	-	2	2	3	8	16	16	37
2000.....	673	-	-	-	-	-	1	4	6	16	28	44
1999.....	662	-	-	-	-	1	-	4	7	19	33	38
Asian or Pacific Islander 2/, male												
2004.....	389	-	-	-	-	-	-	3	2	5	19	17
2003.....	390	-	-	-	-	-	-	-	5	10	12	19
2002.....	352	-	-	-	-	-	-	-	2	5	13	13
2001.....	386	-	-	-	-	1	1	2	5	5	8	22
2000.....	381	-	-	-	-	-	1	4	-	8	14	26
1999.....	366	-	-	-	-	1	-	-	1	10	16	19
Asian or Pacific Islander 2/, female												
2004.....	291	-	-	-	-	-	1	-	7	7	14	17
2003.....	313	-	-	-	-	-	2	3	5	10	15	22
2002.....	305	-	-	-	-	-	-	2	6	8	11	15
2001.....	286	-	-	-	-	1	1	1	3	11	8	15
2000.....	292	-	-	-	-	-	-	-	6	8	14	18
1999.....	296	-	-	-	-	-	-	4	6	9	17	19

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Of small intestine (C17)

All races, both sexes

2004.....	1,115	-	-	-	-	-	2	2	5	16	30	46
2003.....	1,070	-	-	-	-	-	-	3	9	11	25	47
2002.....	1,017	-	-	-	-	-	-	4	4	13	30	47
2001.....	1,082	-	-	-	-	-	1	2	9	24	25	50
2000.....	1,057	-	-	-	-	-	1	2	4	14	32	57
1999.....	1,037	-	-	-	-	-	-	1	4	16	31	70

All races, male

2004.....	592	-	-	-	-	-	-	1	3	10	14	22
2003.....	539	-	-	-	-	-	-	2	5	5	14	27
2002.....	530	-	-	-	-	-	-	4	2	6	20	28
2001.....	568	-	-	-	-	-	1	2	5	16	13	28
2000.....	546	-	-	-	-	-	1	-	3	10	16	32
1999.....	574	-	-	-	-	-	-	1	3	8	20	42

All races, female

2004.....	523	-	-	-	-	-	2	1	2	6	16	24
2003.....	531	-	-	-	-	-	-	1	4	6	11	20
2002.....	487	-	-	-	-	-	-	-	2	7	10	19
2001.....	514	-	-	-	-	-	-	-	4	8	12	22
2000.....	511	-	-	-	-	-	-	2	1	4	16	25
1999.....	463	-	-	-	-	-	-	-	1	8	11	28

White, both sexes

2004.....	908	-	-	-	-	-	2	-	3	6	18	40
2003.....	889	-	-	-	-	-	-	2	3	7	13	34
2002.....	842	-	-	-	-	-	-	3	2	12	21	32
2001.....	883	-	-	-	-	-	1	-	4	14	20	37
2000.....	895	-	-	-	-	-	1	1	3	7	24	41
1999.....	864	-	-	-	-	-	-	1	3	10	25	55

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Of small intestine (C17)												
White, male												
2004.....	487	-	-	-	-	-	-	-	2	4	11	20
2003.....	443	-	-	-	-	-	-	2	1	3	8	21
2002.....	441	-	-	-	-	-	-	3	2	5	14	21
2001.....	462	-	-	-	-	-	1	-	2	9	11	18
2000.....	462	-	-	-	-	-	1	-	2	5	12	25
1999.....	481	-	-	-	-	-	-	1	2	4	16	36
White, female												
2004.....	421	-	-	-	-	-	2	-	1	2	7	20
2003.....	446	-	-	-	-	-	-	-	2	4	5	13
2002.....	401	-	-	-	-	-	-	-	-	7	7	11
2001.....	421	-	-	-	-	-	-	-	2	5	9	19
2000.....	433	-	-	-	-	-	-	1	1	2	12	16
1999.....	383	-	-	-	-	-	-	-	1	6	9	19
Black, both sexes												
2004.....	175	-	-	-	-	-	-	2	1	10	8	4
2003.....	154	-	-	-	-	-	-	-	6	3	11	12
2002.....	150	-	-	-	-	-	-	1	2	1	8	14
2001.....	174	-	-	-	-	-	-	2	5	9	5	13
2000.....	133	-	-	-	-	-	-	1	1	6	8	11
1999.....	150	-	-	-	-	-	-	-	1	5	6	14
Black, male												
2004.....	85	-	-	-	-	-	-	1	-	6	1	2
2003.....	80	-	-	-	-	-	-	-	4	1	6	5
2002.....	72	-	-	-	-	-	-	1	-	1	5	6
2001.....	89	-	-	-	-	-	-	2	3	6	2	10
2000.....	70	-	-	-	-	-	-	-	1	4	4	6
1999.....	83	-	-	-	-	-	-	-	1	3	4	6

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Of colon, rectum and anus (C18-C21)												
All races, both sexes												
2004.....	53,772	-	-	-	1	14	40	87	192	428	898	1,637
2003.....	55,958	-	-	1	3	8	32	96	195	445	870	1,724
2002.....	56,741	1	-	-	1	9	41	88	202	439	986	1,755
2001.....	56,887	-	-	-	-	11	35	81	218	505	928	1,744
2000.....	57,477	-	-	-	3	13	27	79	193	480	889	1,656
1999.....	57,155	1	-	-	1	11	38	87	196	439	894	1,623
All races, male												
2004.....	26,930	-	-	-	1	9	25	50	102	218	477	911
2003.....	28,007	-	-	-	3	6	19	59	99	235	479	965
2002.....	28,501	1	-	-	1	6	28	48	113	233	564	958
2001.....	28,225	-	-	-	-	6	18	38	119	290	491	967
2000.....	28,462	-	-	-	2	8	16	42	111	271	488	893
1999.....	28,240	-	-	-	1	5	22	57	103	226	452	876
All races, female												
2004.....	26,842	-	-	-	-	5	15	37	90	210	421	726
2003.....	27,951	-	-	1	-	2	13	37	96	210	391	759
2002.....	28,240	-	-	-	-	3	13	40	89	206	422	797
2001.....	28,662	-	-	-	-	5	17	43	99	215	437	777
2000.....	29,015	-	-	-	1	5	11	37	82	209	401	763
1999.....	28,915	1	-	-	-	6	16	30	93	213	442	747
White, both sexes												
2004.....	45,824	-	-	-	1	10	25	67	142	320	677	1,242
2003.....	47,711	-	-	1	3	7	22	72	145	329	636	1,310
2002.....	48,562	1	-	-	1	7	31	70	164	337	734	1,313
2001.....	48,748	-	-	-	-	7	27	60	163	384	707	1,324
2000.....	49,488	-	-	-	3	12	19	57	144	349	666	1,250
1999.....	49,374	-	-	-	1	6	26	62	150	336	684	1,264

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	-------------	-----------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Of colon, rectum and anus (C18-C21)

White, male

2004.....	23,038	-	-	-	1	6	15	37	77	178	364	708
2003.....	24,013	-	-	-	3	5	15	45	73	177	362	746
2002.....	24,439	1	-	-	1	4	20	39	87	175	425	736
2001.....	24,335	-	-	-	-	5	11	28	91	226	373	746
2000.....	24,517	-	-	-	2	7	9	32	87	204	363	692
1999.....	24,539	-	-	-	1	2	14	40	78	180	360	692

White, female

2004.....	22,786	-	-	-	-	4	10	30	65	142	313	534
2003.....	23,698	-	-	1	-	2	7	27	72	152	274	564
2002.....	24,123	-	-	-	-	3	11	31	77	162	309	577
2001.....	24,413	-	-	-	-	2	16	32	72	158	334	578
2000.....	24,971	-	-	-	1	5	10	25	57	145	303	558
1999.....	24,835	-	-	-	-	4	12	22	72	156	324	572

Black, both sexes

2004.....	6,640	-	-	-	-	3	9	15	38	78	174	329
2003.....	6,943	-	-	-	-	1	7	17	33	96	192	348
2002.....	6,874	-	-	-	-	2	8	11	29	75	209	370
2001.....	6,905	-	-	-	-	2	6	13	43	99	187	356
2000.....	6,867	-	-	-	-	-	6	12	42	109	184	359
1999.....	6,777	1	-	-	-	5	11	21	35	92	181	305

Black, male

2004.....	3,205	-	-	-	-	3	7	9	18	31	91	166
2003.....	3,335	-	-	-	-	1	3	9	16	44	97	187
2002.....	3,351	-	-	-	-	2	6	5	20	43	114	186
2001.....	3,251	-	-	-	-	1	6	6	24	53	102	187
2000.....	3,326	-	-	-	-	-	5	4	22	56	105	180
1999.....	3,196	-	-	-	-	3	7	14	17	43	82	159

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Of colon, rectum and anus (C18-C21)

Black, female

2004.....	3,435	-	-	-	-	-	2	6	20	47	83	163
2003.....	3,608	-	-	-	-	-	4	8	17	52	95	161
2002.....	3,523	-	-	-	-	-	2	6	9	32	95	184
2001.....	3,654	-	-	-	-	1	-	7	19	46	85	169
2000.....	3,541	-	-	-	-	-	1	8	20	53	79	179
1999.....	3,581	1	-	-	-	2	4	7	18	49	99	146

American Indian 1/, both sexes

2004.....	226	-	-	-	-	-	1	1	2	3	6	10
2003.....	205	-	-	-	-	-	-	-	-	3	7	18
2002.....	246	-	-	-	-	-	-	2	1	7	4	11
2001.....	192	-	-	-	-	-	-	-	1	5	6	3
2000.....	194	-	-	-	-	-	-	2	1	2	11	7
1999.....	170	-	-	-	-	-	1	-	2	1	6	7

American Indian 1/, male

2004.....	131	-	-	-	-	-	1	1	2	2	2	6
2003.....	111	-	-	-	-	-	-	-	-	2	3	11
2002.....	131	-	-	-	-	-	-	1	-	5	1	6
2001.....	95	-	-	-	-	-	-	-	-	3	2	2
2000.....	110	-	-	-	-	-	-	1	-	1	7	4
1999.....	87	-	-	-	-	-	1	-	2	-	2	6

American Indian 1/, female

2004.....	95	-	-	-	-	-	-	-	-	1	4	4
2003.....	94	-	-	-	-	-	-	-	-	1	4	7
2002.....	115	-	-	-	-	-	-	1	1	2	3	5
2001.....	97	-	-	-	-	-	-	-	1	2	4	1
2000.....	84	-	-	-	-	-	-	1	1	1	4	3
1999.....	83	-	-	-	-	-	-	-	-	1	4	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Of colon, rectum and anus (C18-C21)

Asian or Pacific Islander 2/, both sexes

2004.....	1,082	-	-	-	-	1	5	4	10	27	41	56
2003.....	1,099	-	-	-	-	-	3	7	17	17	35	48
2002.....	1,059	-	-	-	-	-	2	5	8	20	39	61
2001.....	1,042	-	-	-	-	2	2	8	11	17	28	61
2000.....	928	-	-	-	-	1	2	8	6	20	28	40
1999.....	834	-	-	-	-	-	-	4	9	10	23	47

Asian or Pacific Islander 2/, male

2004.....	556	-	-	-	-	-	2	3	5	7	20	31
2003.....	548	-	-	-	-	-	1	5	10	12	17	21
2002.....	580	-	-	-	-	-	2	3	6	10	24	30
2001.....	544	-	-	-	-	-	1	4	4	8	14	32
2000.....	509	-	-	-	-	1	2	5	2	10	13	17
1999.....	418	-	-	-	-	-	-	3	6	3	8	19

Asian or Pacific Islander 2/, female

2004.....	526	-	-	-	-	1	3	1	5	20	21	25
2003.....	551	-	-	-	-	-	2	2	7	5	18	27
2002.....	479	-	-	-	-	-	-	2	2	10	15	31
2001.....	498	-	-	-	-	2	1	4	7	9	14	29
2000.....	419	-	-	-	-	-	-	3	4	10	15	23
1999.....	416	-	-	-	-	-	-	1	3	7	15	28

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Colon (C18)												
All races, both sexes												
2004.....	44,591	-	-	-	1	12	29	59	137	297	639	1,219
2003.....	46,868	-	-	1	2	8	28	82	141	359	639	1,328
2002.....	47,586	-	-	-	1	7	37	59	151	318	755	1,337
2001.....	47,860	-	-	-	-	11	28	57	175	380	709	1,331
2000.....	48,570	-	-	-	3	9	19	55	156	360	699	1,267
1999.....	48,433	1	-	-	1	9	29	62	154	342	688	1,265
All races, male												
2004.....	21,883	-	-	-	1	7	18	35	66	146	330	676
2003.....	23,081	-	-	-	2	6	17	51	69	180	346	729
2002.....	23,486	-	-	-	1	4	25	31	83	166	423	709
2001.....	23,302	-	-	-	-	6	13	25	94	210	363	730
2000.....	23,634	-	-	-	2	7	12	26	92	197	380	665
1999.....	23,596	-	-	-	1	4	17	36	78	171	347	675
All races, female												
2004.....	22,708	-	-	-	-	5	11	24	71	151	309	543
2003.....	23,787	-	-	1	-	2	11	31	72	179	293	599
2002.....	24,100	-	-	-	-	3	12	28	68	152	332	628
2001.....	24,558	-	-	-	-	5	15	32	81	170	346	601
2000.....	24,936	-	-	-	1	2	7	29	64	163	319	602
1999.....	24,837	1	-	-	-	5	12	26	76	171	341	590
White, both sexes												
2004.....	37,866	-	-	-	1	9	17	47	102	224	485	920
2003.....	39,867	-	-	1	2	7	18	64	104	268	454	991
2002.....	40,626	-	-	-	1	5	27	48	121	246	562	972
2001.....	40,860	-	-	-	-	7	22	40	131	287	527	991
2000.....	41,724	-	-	-	3	8	13	42	118	265	512	935
1999.....	41,735	-	-	-	1	5	19	41	115	259	516	966

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Colon (C18)

White, male

2004.....	18,657	-	-	-	1	5	9	27	51	121	264	529
2003.....	19,753	-	-	-	2	5	13	40	51	134	259	562
2002.....	20,097	-	-	-	1	2	17	28	64	127	321	539
2001.....	20,026	-	-	-	-	5	8	18	74	166	273	549
2000.....	20,332	-	-	-	2	6	7	22	73	155	276	509
1999.....	20,494	-	-	-	1	2	10	23	58	138	275	527

White, female

2004.....	19,209	-	-	-	-	4	8	20	51	103	221	391
2003.....	20,114	-	-	1	-	2	5	24	53	134	195	429
2002.....	20,529	-	-	-	-	3	10	20	57	119	241	433
2001.....	20,834	-	-	-	-	2	14	22	57	121	254	442
2000.....	21,392	-	-	-	1	2	6	20	45	110	236	426
1999.....	21,241	-	-	-	-	3	9	18	57	121	241	439

Black, both sexes

2004.....	5,668	-	-	-	-	2	8	7	27	53	123	248
2003.....	5,951	-	-	-	-	1	7	11	25	76	152	289
2002.....	5,916	-	-	-	-	2	8	7	24	55	165	308
2001.....	5,985	-	-	-	-	2	5	12	34	78	157	290
2000.....	5,948	-	-	-	-	-	4	7	32	81	152	296
1999.....	5,900	1	-	-	-	4	10	18	29	75	152	261

Black, male

2004.....	2,677	-	-	-	-	2	7	4	11	19	54	117
2003.....	2,804	-	-	-	-	1	3	6	11	34	72	145
2002.....	2,834	-	-	-	-	2	6	2	15	31	86	144
2001.....	2,761	-	-	-	-	1	5	5	16	37	80	154
2000.....	2,810	-	-	-	-	-	3	1	17	36	87	141
1999.....	2,711	-	-	-	-	2	7	11	12	32	66	130

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Colon (C18)

Black, female

2004.....	2,991	-	-	-	-	-	1	3	16	34	69	131
2003.....	3,147	-	-	-	-	-	4	5	14	42	80	144
2002.....	3,082	-	-	-	-	-	2	5	9	24	79	164
2001.....	3,224	-	-	-	-	1	-	7	18	41	77	136
2000.....	3,138	-	-	-	-	-	1	6	15	45	65	155
1999.....	3,189	1	-	-	-	2	3	7	17	43	86	131

American Indian 1/, both sexes

2004.....	187	-	-	-	-	-	1	1	2	2	3	9
2003.....	164	-	-	-	-	-	-	-	-	2	4	11
2002.....	192	-	-	-	-	-	-	1	1	3	3	9
2001.....	150	-	-	-	-	-	-	-	1	3	3	3
2000.....	153	-	-	-	-	-	-	-	1	2	11	6
1999.....	133	-	-	-	-	-	-	-	2	1	4	5

American Indian 1/, male

2004.....	106	-	-	-	-	-	1	1	2	1	1	5
2003.....	90	-	-	-	-	-	-	-	-	2	2	7
2002.....	95	-	-	-	-	-	-	-	-	2	1	5
2001.....	69	-	-	-	-	-	-	-	-	1	-	2
2000.....	88	-	-	-	-	-	-	-	-	1	7	3
1999.....	64	-	-	-	-	-	-	-	2	-	1	4

American Indian 1/, female

2004.....	81	-	-	-	-	-	-	-	-	1	2	4
2003.....	74	-	-	-	-	-	-	-	-	-	2	4
2002.....	97	-	-	-	-	-	-	1	1	1	2	4
2001.....	81	-	-	-	-	-	-	-	1	2	3	1
2000.....	65	-	-	-	-	-	-	-	1	1	4	3
1999.....	69	-	-	-	-	-	-	-	-	1	3	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Colon (C18)

Asian or Pacific Islander 2/, both sexes

2004.....	870	-	-	-	-	1	3	4	6	18	28	42
2003.....	886	-	-	-	-	-	3	7	12	13	29	37
2002.....	852	-	-	-	-	-	2	3	5	14	25	48
2001.....	865	-	-	-	-	2	1	5	9	12	22	47
2000.....	745	-	-	-	-	1	2	6	5	12	24	30
1999.....	665	-	-	-	-	-	-	3	8	7	16	33

Asian or Pacific Islander 2/, male

2004.....	443	-	-	-	-	-	1	3	2	5	11	25
2003.....	434	-	-	-	-	-	1	5	7	10	13	15
2002.....	460	-	-	-	-	-	2	1	4	6	15	21
2001.....	446	-	-	-	-	-	-	2	4	6	10	25
2000.....	404	-	-	-	-	1	2	3	2	5	10	12
1999.....	327	-	-	-	-	-	-	2	6	1	5	14

Asian or Pacific Islander 2/, female

2004.....	427	-	-	-	-	1	2	1	4	13	17	17
2003.....	452	-	-	-	-	-	2	2	5	3	16	22
2002.....	392	-	-	-	-	-	-	2	1	8	10	27
2001.....	419	-	-	-	-	2	1	3	5	6	12	22
2000.....	341	-	-	-	-	-	-	3	3	7	14	18
1999.....	338	-	-	-	-	-	-	1	2	6	11	19

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rectosigmoid junction and rectum (C19-C20)												
All races, both sexes												
2004.....	8,592	-	-	-	-	2	11	27	52	112	229	363
2003.....	8,535	-	-	-	1	-	4	13	45	75	205	352
2002.....	8,616	1	-	-	-	2	4	28	47	106	198	373
2001.....	8,516	-	-	-	-	-	7	24	39	107	193	381
2000.....	8,415	-	-	-	-	4	7	24	33	104	163	347
1999.....	8,260	-	-	-	-	2	9	24	37	79	186	320
All races, male												
2004.....	4,807	-	-	-	-	2	7	14	34	61	129	216
2003.....	4,740	-	-	-	1	-	2	8	27	46	119	217
2002.....	4,802	1	-	-	-	2	3	17	28	59	123	227
2001.....	4,738	-	-	-	-	-	5	13	23	71	116	223
2000.....	4,634	-	-	-	-	1	4	16	16	65	97	215
1999.....	4,476	-	-	-	-	1	5	20	20	46	95	181
All races, female												
2004.....	3,785	-	-	-	-	-	4	13	18	51	100	147
2003.....	3,795	-	-	-	-	-	2	5	18	29	86	135
2002.....	3,814	-	-	-	-	-	1	11	19	47	75	146
2001.....	3,778	-	-	-	-	-	2	11	16	36	77	158
2000.....	3,781	-	-	-	-	3	3	8	17	39	66	132
1999.....	3,784	-	-	-	-	1	4	4	17	33	91	139
White, both sexes												
2004.....	7,446	-	-	-	-	1	8	20	39	82	168	279
2003.....	7,364	-	-	-	1	-	4	8	35	53	160	280
2002.....	7,451	1	-	-	-	2	4	21	41	79	146	299
2001.....	7,426	-	-	-	-	-	5	20	29	81	157	304
2000.....	7,328	-	-	-	-	4	5	15	23	72	131	277
1999.....	7,234	-	-	-	-	1	7	21	32	65	152	264

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Rectosigmoid junction and rectum (C19-C20)

White, male

2004.....	4,179	-	-	-	-	1	6	10	25	50	87	165
2003.....	4,106	-	-	-	1	-	2	5	21	37	93	169
2002.....	4,158	1	-	-	-	2	3	11	23	43	91	176
2001.....	4,145	-	-	-	-	-	3	10	16	53	89	184
2000.....	4,019	-	-	-	-	1	2	10	11	42	79	171
1999.....	3,901	-	-	-	-	-	4	17	17	37	77	148

White, female

2004.....	3,267	-	-	-	-	-	2	10	14	32	81	114
2003.....	3,258	-	-	-	-	-	2	3	14	16	67	111
2002.....	3,293	-	-	-	-	-	1	10	18	36	55	123
2001.....	3,281	-	-	-	-	-	2	10	13	28	68	120
2000.....	3,309	-	-	-	-	3	3	5	12	30	52	106
1999.....	3,333	-	-	-	-	1	3	4	15	28	75	116

Black, both sexes

2004.....	911	-	-	-	-	1	1	7	9	20	46	72
2003.....	930	-	-	-	-	-	-	5	5	17	36	56
2002.....	907	-	-	-	-	-	-	4	3	18	37	59
2001.....	877	-	-	-	-	-	1	1	8	19	27	63
2000.....	869	-	-	-	-	-	2	5	9	24	28	60
1999.....	826	-	-	-	-	1	1	2	4	11	26	40

Black, male

2004.....	496	-	-	-	-	1	-	4	6	8	33	45
2003.....	502	-	-	-	-	-	-	3	3	7	21	39
2002.....	490	-	-	-	-	-	-	3	3	10	23	41
2001.....	472	-	-	-	-	-	1	1	7	14	21	32
2000.....	490	-	-	-	-	-	2	3	5	18	15	38
1999.....	462	-	-	-	-	1	-	2	3	7	14	26

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rectosigmoid junction and rectum (C19-C20)												
Asian or Pacific Islander 2/, both sexes												
2004.....	197	-	-	-	-	-	2	-	4	9	13	11
2003.....	203	-	-	-	-	-	-	-	5	4	6	11
2002.....	205	-	-	-	-	-	-	2	3	6	14	13
2001.....	174	-	-	-	-	-	1	3	2	5	6	14
2000.....	179	-	-	-	-	-	-	2	1	8	4	9
1999.....	164	-	-	-	-	-	-	1	1	3	6	14
Asian or Pacific Islander 2/, male												
2004.....	108	-	-	-	-	-	1	-	3	2	9	5
2003.....	112	-	-	-	-	-	-	-	3	2	4	6
2002.....	119	-	-	-	-	-	-	2	2	4	9	9
2001.....	96	-	-	-	-	-	1	2	-	2	4	7
2000.....	105	-	-	-	-	-	-	2	-	5	3	5
1999.....	90	-	-	-	-	-	-	1	-	2	3	5
Asian or Pacific Islander 2/, female												
2004.....	89	-	-	-	-	-	1	-	1	7	4	6
2003.....	91	-	-	-	-	-	-	-	2	2	2	5
2002.....	86	-	-	-	-	-	-	-	1	2	5	4
2001.....	78	-	-	-	-	-	-	1	2	3	2	7
2000.....	74	-	-	-	-	-	-	-	1	3	1	4
1999.....	74	-	-	-	-	-	-	-	1	1	3	9

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Anus and anus canal (C21)

All races, both sexes

2004.....	589	-	-	-	-	-	-	1	3	19	30	55
2003.....	555	-	-	-	-	-	-	1	9	11	26	44
2002.....	539	-	-	-	-	-	-	1	4	15	33	45
2001.....	511	-	-	-	-	-	-	-	4	18	26	32
2000.....	492	-	-	-	-	-	1	-	4	16	27	42
1999.....	462	-	-	-	-	-	-	1	5	18	20	38

All races, male

2004.....	240	-	-	-	-	-	-	1	2	11	18	19
2003.....	186	-	-	-	-	-	-	-	3	9	14	19
2002.....	213	-	-	-	-	-	-	-	2	8	18	22
2001.....	185	-	-	-	-	-	-	-	2	9	12	14
2000.....	194	-	-	-	-	-	-	-	3	9	11	13
1999.....	168	-	-	-	-	-	-	1	5	9	10	20

All races, female

2004.....	349	-	-	-	-	-	-	-	1	8	12	36
2003.....	369	-	-	-	-	-	-	1	6	2	12	25
2002.....	326	-	-	-	-	-	-	1	2	7	15	23
2001.....	326	-	-	-	-	-	-	-	2	9	14	18
2000.....	298	-	-	-	-	-	1	-	1	7	16	29
1999.....	294	-	-	-	-	-	-	-	-	9	10	18

White, both sexes

2004.....	512	-	-	-	-	-	-	-	1	14	24	43
2003.....	480	-	-	-	-	-	-	-	6	8	22	39
2002.....	485	-	-	-	-	-	-	1	2	12	26	42
2001.....	462	-	-	-	-	-	-	-	3	16	23	29
2000.....	436	-	-	-	-	-	1	-	3	12	23	38
1999.....	405	-	-	-	-	-	-	-	3	12	16	34

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Anus and anus canal (C21)												
White, male												
2004.....	202	-	-	-	-	-	-	-	1	7	13	14
2003.....	154	-	-	-	-	-	-	-	1	6	10	15
2002.....	184	-	-	-	-	-	-	-	-	5	13	21
2001.....	164	-	-	-	-	-	-	-	1	7	11	13
2000.....	166	-	-	-	-	-	-	-	3	7	8	12
1999.....	144	-	-	-	-	-	-	-	3	5	8	17
White, female												
2004.....	310	-	-	-	-	-	-	-	-	7	11	29
2003.....	326	-	-	-	-	-	-	-	5	2	12	24
2002.....	301	-	-	-	-	-	-	1	2	7	13	21
2001.....	298	-	-	-	-	-	-	-	2	9	12	16
2000.....	270	-	-	-	-	-	1	-	-	5	15	26
1999.....	261	-	-	-	-	-	-	-	-	7	8	17
Black, both sexes												
2004.....	61	-	-	-	-	-	-	1	2	5	5	9
2003.....	62	-	-	-	-	-	-	1	3	3	4	3
2002.....	51	-	-	-	-	-	-	-	2	2	7	3
2001.....	43	-	-	-	-	-	-	-	1	2	3	3
2000.....	50	-	-	-	-	-	-	-	1	4	4	3
1999.....	51	-	-	-	-	-	-	1	2	6	3	4
Black, male												
2004.....	32	-	-	-	-	-	-	1	1	4	4	4
2003.....	29	-	-	-	-	-	-	-	2	3	4	3
2002.....	27	-	-	-	-	-	-	-	2	2	5	1
2001.....	18	-	-	-	-	-	-	-	1	2	1	1
2000.....	26	-	-	-	-	-	-	-	-	2	3	1
1999.....	23	-	-	-	-	-	-	1	2	4	2	3

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	-------------	-----------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Of liver and intrahepatic bile ducts (C22)

All races, both sexes

2004.....	15,321	1	22	7	9	16	27	28	56	108	272	872
2003.....	14,706	6	23	6	8	10	26	25	64	113	282	815
2002.....	14,047	3	28	7	11	10	17	33	72	123	291	731
2001.....	13,351	5	18	7	8	14	15	29	46	137	264	795
2000.....	12,916	2	23	9	11	15	15	28	58	113	316	742
1999.....	12,382	8	29	5	6	19	24	33	45	127	317	666

All races, male

2004.....	9,832	1	12	3	5	10	15	20	35	82	207	673
2003.....	9,482	2	18	5	4	6	14	12	41	83	205	643
2002.....	9,011	2	19	4	8	9	9	20	42	85	223	596
2001.....	8,483	3	7	4	3	6	9	16	27	100	202	618
2000.....	8,176	1	14	6	2	11	9	18	41	74	223	595
1999.....	7,689	5	20	5	4	12	12	22	26	88	230	533

All races, female

2004.....	5,489	-	10	4	4	6	12	8	21	26	65	199
2003.....	5,224	4	5	1	4	4	12	13	23	30	77	172
2002.....	5,036	1	9	3	3	1	8	13	30	38	68	135
2001.....	4,868	2	11	3	5	8	6	13	19	37	62	177
2000.....	4,740	1	9	3	9	4	6	10	17	39	93	147
1999.....	4,693	3	9	-	2	7	12	11	19	39	87	133

White, both sexes

2004.....	12,111	1	19	3	7	9	18	17	30	63	185	592
2003.....	11,660	4	19	4	4	7	16	12	33	73	191	555
2002.....	11,250	2	22	7	6	6	13	18	39	74	201	521
2001.....	10,697	5	14	2	6	10	12	19	29	82	155	549
2000.....	10,304	1	17	8	5	10	10	12	32	65	196	510
1999.....	9,986	6	26	3	3	14	15	17	19	73	210	448

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Of liver and intrahepatic bile ducts (C22)

White, male

2004.....	7,694	1	11	2	3	6	8	13	18	46	137	455
2003.....	7,489	2	15	4	3	4	9	3	20	50	133	441
2002.....	7,114	2	14	4	3	5	5	8	19	48	153	426
2001.....	6,774	3	5	1	3	5	7	9	15	56	119	427
2000.....	6,447	1	11	5	2	7	6	8	19	37	142	401
1999.....	6,151	3	18	3	2	10	7	11	10	45	155	351

White, female

2004.....	4,417	-	8	1	4	3	10	4	12	17	48	137
2003.....	4,171	2	4	-	1	3	7	9	13	23	58	114
2002.....	4,136	-	8	3	3	1	8	10	20	26	48	95
2001.....	3,923	2	9	1	3	5	5	10	14	26	36	122
2000.....	3,857	-	6	3	3	3	4	4	13	28	54	109
1999.....	3,835	3	8	-	1	4	8	6	9	28	55	97

Black, both sexes

2004.....	2,014	-	2	4	1	7	7	7	15	27	52	191
2003.....	1,877	2	1	2	1	2	5	6	19	23	55	164
2002.....	1,734	1	4	-	3	3	1	9	18	23	61	147
2001.....	1,670	-	3	3	2	2	2	6	9	30	66	175
2000.....	1,598	1	2	-	2	4	2	10	11	25	77	166
1999.....	1,507	1	3	1	3	2	5	9	12	31	60	150

Black, male

2004.....	1,349	-	-	1	1	4	5	4	10	20	40	147
2003.....	1,234	-	-	1	1	1	3	3	12	18	41	126
2002.....	1,165	-	3	-	3	3	1	6	11	16	45	116
2001.....	1,048	-	2	2	-	-	1	4	6	23	48	133
2000.....	1,061	-	1	-	-	3	2	5	10	16	50	138
1999.....	947	1	2	1	2	-	2	7	7	21	44	125

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Of liver and intrahepatic bile ducts (C22)												
Black, female												
2004.....	665	-	2	3	-	3	2	3	5	7	12	44
2003.....	643	2	1	1	-	1	2	3	7	5	14	38
2002.....	569	1	1	-	-	-	-	3	7	7	16	31
2001.....	622	-	1	1	2	2	1	2	3	7	18	42
2000.....	537	1	1	-	2	1	-	5	1	9	27	28
1999.....	560	-	1	-	1	2	3	2	5	10	16	25
American Indian 1/, both sexes												
2004.....	143	-	-	-	-	-	-	2	-	-	2	9
2003.....	105	-	2	-	1	-	-	-	1	1	3	9
2002.....	94	-	1	-	-	-	-	-	1	-	2	6
2001.....	109	-	-	-	-	1	-	-	1	2	1	9
2000.....	98	-	-	1	-	-	-	-	2	1	3	7
1999.....	76	-	-	-	-	-	1	1	-	3	-	7
American Indian 1/, male												
2004.....	77	-	-	-	-	-	-	2	-	-	2	8
2003.....	71	-	2	-	-	-	-	-	1	1	3	7
2002.....	60	-	1	-	-	-	-	-	1	-	2	4
2001.....	71	-	-	-	-	-	-	-	-	1	-	7
2000.....	65	-	-	1	-	-	-	-	1	1	3	7
1999.....	42	-	-	-	-	-	1	-	-	2	-	5
American Indian 1/, female												
2004.....	66	-	-	-	-	-	-	-	-	-	-	1
2003.....	34	-	-	-	1	-	-	-	-	-	-	2
2002.....	34	-	-	-	-	-	-	-	-	-	-	2
2001.....	38	-	-	-	-	1	-	-	1	1	1	2
2000.....	33	-	-	-	-	-	-	-	1	-	-	-
1999.....	34	-	-	-	-	-	-	1	-	1	-	2

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Of liver and intrahepatic bile ducts (C22)												
Asian or Pacific Islander 2/, both sexes												
2004.....	1,053	-	1	-	1	-	2	2	11	18	33	80
2003.....	1,064	-	1	-	2	1	5	7	11	16	33	87
2002.....	969	-	1	-	2	1	3	6	14	26	27	57
2001.....	875	-	1	2	-	1	1	4	7	23	42	62
2000.....	916	-	4	-	4	1	3	6	13	22	40	59
1999.....	813	1	-	1	-	3	3	6	14	20	47	61
Asian or Pacific Islander 2/, male												
2004.....	712	-	1	-	1	-	2	1	7	16	28	63
2003.....	688	-	1	-	-	1	2	6	8	14	28	69
2002.....	672	-	1	-	2	1	3	6	11	21	23	50
2001.....	590	-	-	1	-	1	1	3	6	20	35	51
2000.....	603	-	2	-	-	1	1	5	11	20	28	49
1999.....	549	1	-	1	-	2	2	4	9	20	31	52
Asian or Pacific Islander 2/, female												
2004.....	341	-	-	-	-	-	-	1	4	2	5	17
2003.....	376	-	-	-	2	-	3	1	3	2	5	18
2002.....	297	-	-	-	-	-	-	-	3	5	4	7
2001.....	285	-	1	1	-	-	-	1	1	3	7	11
2000.....	313	-	2	-	4	-	2	1	2	2	12	10
1999.....	264	-	-	-	-	1	1	2	5	-	16	9

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Liver (C22.0,C22.2-C22.9)												
All races, both sexes												
2004.....	11,831	1	21	7	9	15	24	21	41	68	216	741
2003.....	11,423	6	23	6	8	10	23	21	42	85	227	687
2002.....	11,048	3	28	7	11	10	16	31	51	91	239	641
2001.....	10,489	5	18	7	8	12	14	20	40	109	215	707
2000.....	10,192	2	22	9	11	13	15	25	44	94	264	644
1999.....	9,830	8	29	5	6	19	24	27	35	93	266	584
All races, male												
2004.....	8,125	1	12	3	5	9	14	15	30	57	178	602
2003.....	7,823	2	18	5	4	6	11	11	30	69	179	582
2002.....	7,531	2	19	4	8	9	9	20	33	66	193	539
2001.....	7,075	3	7	4	3	4	9	11	23	84	173	577
2000.....	6,843	1	14	6	2	10	9	16	30	64	195	536
1999.....	6,481	5	20	5	4	12	12	18	20	71	204	490
All races, female												
2004.....	3,706	-	9	4	4	6	10	6	11	11	38	139
2003.....	3,600	4	5	1	4	4	12	10	12	16	48	105
2002.....	3,517	1	9	3	3	1	7	11	18	25	46	102
2001.....	3,414	2	11	3	5	8	5	9	17	25	42	130
2000.....	3,349	1	8	3	9	3	6	9	14	30	69	108
1999.....	3,349	3	9	-	2	7	12	9	15	22	62	94
White, both sexes												
2004.....	9,083	1	18	3	7	8	16	10	17	32	141	490
2003.....	8,824	4	19	4	4	7	13	8	16	47	142	461
2002.....	8,673	2	22	7	6	6	12	16	24	47	159	450
2001.....	8,224	5	14	2	6	8	11	12	25	58	117	482
2000.....	7,914	1	16	8	5	9	10	10	22	50	155	430
1999.....	7,730	6	26	3	3	14	15	12	11	49	168	376

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Liver (C22.0,C22.2-C22.9)												
White, male												
2004.....	6,194	1	11	2	3	5	8	8	14	24	113	397
2003.....	6,053	2	15	4	3	4	6	2	11	36	111	394
2002.....	5,839	2	14	4	3	5	5	8	14	32	128	381
2001.....	5,557	3	5	1	3	3	7	5	12	41	95	393
2000.....	5,273	1	11	5	2	7	6	7	10	30	121	353
1999.....	5,081	3	18	3	2	10	7	8	5	33	133	313
White, female												
2004.....	2,889	-	7	1	4	3	8	2	3	8	28	93
2003.....	2,771	2	4	-	1	3	7	6	5	11	31	67
2002.....	2,834	-	8	3	3	1	7	8	10	15	31	69
2001.....	2,667	2	9	1	3	5	4	7	13	17	22	89
2000.....	2,641	-	5	3	3	2	4	3	12	20	34	77
1999.....	2,649	3	8	-	1	4	8	4	6	16	35	63
Black, both sexes												
2004.....	1,737	-	2	4	1	7	6	7	13	20	42	172
2003.....	1,598	2	1	2	1	2	5	6	15	22	52	141
2002.....	1,494	1	4	-	3	3	1	9	12	19	56	135
2001.....	1,437	-	3	3	2	2	2	5	8	27	61	159
2000.....	1,389	1	2	-	2	3	2	10	9	22	69	154
1999.....	1,319	1	3	1	3	2	5	8	11	24	52	145
Black, male												
2004.....	1,226	-	-	1	1	4	4	4	9	18	35	138
2003.....	1,100	-	-	1	1	1	3	3	11	18	40	118
2002.....	1,049	-	3	-	3	3	1	6	7	13	43	109
2001.....	938	-	2	2	-	-	1	4	5	23	46	128
2000.....	966	-	1	-	-	2	2	5	9	14	45	131
1999.....	866	1	2	1	2	-	2	6	6	19	40	122

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Liver (C22.0,C22.2-C22.9)												
Black, female												
2004.....	511	-	2	3	-	3	2	3	4	2	7	34
2003.....	498	2	1	1	-	1	2	3	4	4	12	23
2002.....	445	1	1	-	-	-	-	3	5	6	13	26
2001.....	499	-	1	1	2	2	1	1	3	4	15	31
2000.....	423	1	1	-	2	1	-	5	-	8	24	23
1999.....	453	-	1	-	1	2	3	2	5	5	12	23
American Indian 1/, both sexes												
2004.....	116	-	-	-	-	-	-	2	-	-	2	9
2003.....	83	-	2	-	1	-	-	-	1	1	3	8
2002.....	67	-	1	-	-	-	-	-	1	-	2	5
2001.....	86	-	-	-	-	1	-	-	-	2	1	8
2000.....	85	-	-	1	-	-	-	-	1	1	3	6
1999.....	64	-	-	-	-	-	1	1	-	2	-	7
American Indian 1/, male												
2004.....	69	-	-	-	-	-	-	2	-	-	2	8
2003.....	60	-	2	-	-	-	-	-	1	1	3	7
2002.....	48	-	1	-	-	-	-	-	1	-	2	4
2001.....	60	-	-	-	-	-	-	-	-	1	-	7
2000.....	58	-	-	1	-	-	-	-	1	1	3	6
1999.....	35	-	-	-	-	-	1	-	-	1	-	5
American Indian 1/, female												
2004.....	47	-	-	-	-	-	-	-	-	-	-	1
2003.....	23	-	-	-	1	-	-	-	-	-	-	1
2002.....	19	-	-	-	-	-	-	-	-	-	-	1
2001.....	26	-	-	-	-	1	-	-	-	1	1	1
2000.....	27	-	-	-	-	-	-	-	-	-	-	-
1999.....	29	-	-	-	-	-	-	1	-	1	-	2

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Liver (C22.0,C22.2-C22.9)												
Asian or Pacific Islander 2/, both sexes												
2004.....	895	-	1	-	1	-	2	2	11	16	31	70
2003.....	918	-	1	-	2	1	5	7	10	15	30	77
2002.....	814	-	1	-	2	1	3	6	14	25	22	51
2001.....	742	-	1	2	-	1	1	3	7	22	36	58
2000.....	804	-	4	-	4	1	3	5	12	21	37	54
1999.....	717	1	-	1	-	3	3	6	13	18	46	56
Asian or Pacific Islander 2/, male												
2004.....	636	-	1	-	1	-	2	1	7	15	28	59
2003.....	610	-	1	-	-	1	2	6	7	14	25	63
2002.....	595	-	1	-	2	1	3	6	11	21	20	45
2001.....	520	-	-	1	-	1	1	2	6	19	32	49
2000.....	546	-	2	-	-	1	1	4	10	19	26	46
1999.....	499	1	-	1	-	2	2	4	9	18	31	50
Asian or Pacific Islander 2/, female												
2004.....	259	-	-	-	-	-	-	1	4	1	3	11
2003.....	308	-	-	-	2	-	3	1	3	1	5	14
2002.....	219	-	-	-	-	-	-	-	3	4	2	6
2001.....	222	-	1	1	-	-	-	1	1	3	4	9
2000.....	258	-	2	-	4	-	2	1	2	2	11	8
1999.....	218	-	-	-	-	1	1	2	4	-	15	6

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intrahepatic bile duct carcinoma (C22.1)												
All races, both sexes												
2004.....	3,490	-	1	-	-	1	3	7	15	40	56	131
2003.....	3,283	-	-	-	-	-	3	4	22	28	55	128
2002.....	2,999	-	-	-	-	-	1	2	21	32	52	90
2001.....	2,862	-	-	-	-	2	1	9	6	28	49	88
2000.....	2,724	-	1	-	-	2	-	3	14	19	52	98
1999.....	2,552	-	-	-	-	-	-	6	10	34	51	82
All races, male												
2004.....	1,707	-	-	-	-	1	1	5	5	25	29	71
2003.....	1,659	-	-	-	-	-	3	1	11	14	26	61
2002.....	1,480	-	-	-	-	-	-	-	9	19	30	57
2001.....	1,408	-	-	-	-	2	-	5	4	16	29	41
2000.....	1,333	-	-	-	-	1	-	2	11	10	28	59
1999.....	1,208	-	-	-	-	-	-	4	6	17	26	43
All races, female												
2004.....	1,783	-	1	-	-	-	2	2	10	15	27	60
2003.....	1,624	-	-	-	-	-	-	3	11	14	29	67
2002.....	1,519	-	-	-	-	-	1	2	12	13	22	33
2001.....	1,454	-	-	-	-	-	1	4	2	12	20	47
2000.....	1,391	-	1	-	-	1	-	1	3	9	24	39
1999.....	1,344	-	-	-	-	-	-	2	4	17	25	39
White, both sexes												
2004.....	3,028	-	1	-	-	1	2	7	13	31	44	102
2003.....	2,836	-	-	-	-	-	3	4	17	26	49	94
2002.....	2,577	-	-	-	-	-	1	2	15	27	42	71
2001.....	2,473	-	-	-	-	2	1	7	4	24	38	67
2000.....	2,390	-	1	-	-	1	-	2	10	15	41	80
1999.....	2,256	-	-	-	-	-	-	5	8	24	42	72

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intrahepatic bile duct carcinoma (C22.1)												
White, male												
2004.....	1,500	-	-	-	-	1	-	5	4	22	24	58
2003.....	1,436	-	-	-	-	-	3	1	9	14	22	47
2002.....	1,275	-	-	-	-	-	-	-	5	16	25	45
2001.....	1,217	-	-	-	-	2	-	4	3	15	24	34
2000.....	1,174	-	-	-	-	-	-	1	9	7	21	48
1999.....	1,070	-	-	-	-	-	-	3	5	12	22	38
White, female												
2004.....	1,528	-	1	-	-	-	2	2	9	9	20	44
2003.....	1,400	-	-	-	-	-	-	3	8	12	27	47
2002.....	1,302	-	-	-	-	-	1	2	10	11	17	26
2001.....	1,256	-	-	-	-	-	1	3	1	9	14	33
2000.....	1,216	-	1	-	-	1	-	1	1	8	20	32
1999.....	1,186	-	-	-	-	-	-	2	3	12	20	34
Black, both sexes												
2004.....	277	-	-	-	-	-	1	-	2	7	10	19
2003.....	279	-	-	-	-	-	-	-	4	1	3	23
2002.....	240	-	-	-	-	-	-	-	6	4	5	12
2001.....	233	-	-	-	-	-	-	1	1	3	5	16
2000.....	209	-	-	-	-	1	-	-	2	3	8	12
1999.....	188	-	-	-	-	-	-	1	1	7	8	5
Black, male												
2004.....	123	-	-	-	-	-	1	-	1	2	5	9
2003.....	134	-	-	-	-	-	-	-	1	-	1	8
2002.....	116	-	-	-	-	-	-	-	4	3	2	7
2001.....	110	-	-	-	-	-	-	-	1	-	2	5
2000.....	95	-	-	-	-	1	-	-	1	2	5	7
1999.....	81	-	-	-	-	-	-	1	1	2	4	3

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intrahepatic bile duct carcinoma (C22.1)												
Asian or Pacific Islander 2/, both sexes												
2004.....	158	-	-	-	-	-	-	-	-	2	2	10
2003.....	146	-	-	-	-	-	-	-	1	1	3	10
2002.....	155	-	-	-	-	-	-	-	-	1	5	6
2001.....	133	-	-	-	-	-	-	1	-	1	6	4
2000.....	112	-	-	-	-	-	-	1	1	1	3	5
1999.....	96	-	-	-	-	-	-	-	1	2	1	5
Asian or Pacific Islander 2/, male												
2004.....	76	-	-	-	-	-	-	-	-	1	-	4
2003.....	78	-	-	-	-	-	-	-	1	-	3	6
2002.....	77	-	-	-	-	-	-	-	-	-	3	5
2001.....	70	-	-	-	-	-	-	1	-	1	3	2
2000.....	57	-	-	-	-	-	-	1	1	1	2	3
1999.....	50	-	-	-	-	-	-	-	-	2	-	2
Asian or Pacific Islander 2/, female												
2004.....	82	-	-	-	-	-	-	-	-	1	2	6
2003.....	68	-	-	-	-	-	-	-	-	1	-	4
2002.....	78	-	-	-	-	-	-	-	-	1	2	1
2001.....	63	-	-	-	-	-	-	-	-	-	3	2
2000.....	55	-	-	-	-	-	-	-	-	-	1	2
1999.....	46	-	-	-	-	-	-	-	1	-	1	3

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Of gallbladder and extrahepatic bile ducts (C23-C24)

All races, both sexes

2004.....	3,397	-	-	-	1	-	1	4	8	10	39	51
2003.....	3,406	-	-	-	-	1	2	-	5	10	30	91
2002.....	3,408	-	-	-	1	-	-	5	2	18	45	89
2001.....	3,601	-	-	-	1	-	-	2	4	21	53	97
2000.....	3,666	-	-	-	-	-	1	2	5	17	40	88
1999.....	3,590	-	-	-	-	-	2	1	2	14	50	71

All races, male

2004.....	1,285	-	-	-	1	-	-	4	3	3	15	22
2003.....	1,306	-	-	-	-	1	1	-	3	5	15	40
2002.....	1,288	-	-	-	-	-	-	2	1	9	16	23
2001.....	1,332	-	-	-	1	-	-	1	-	9	17	34
2000.....	1,372	-	-	-	-	-	1	2	2	9	17	33
1999.....	1,300	-	-	-	-	-	1	1	-	6	20	31

All races, female

2004.....	2,112	-	-	-	-	-	1	-	5	7	24	29
2003.....	2,100	-	-	-	-	-	1	-	2	5	15	51
2002.....	2,120	-	-	-	1	-	-	3	1	9	29	66
2001.....	2,269	-	-	-	-	-	-	1	4	12	36	63
2000.....	2,294	-	-	-	-	-	-	-	3	8	23	55
1999.....	2,290	-	-	-	-	-	1	-	2	8	30	40

White, both sexes

2004.....	2,917	-	-	-	1	-	1	3	7	9	29	41
2003.....	2,969	-	-	-	-	-	2	-	4	9	19	70
2002.....	2,945	-	-	-	1	-	-	4	2	13	36	63
2001.....	3,140	-	-	-	1	-	-	2	3	16	41	80
2000.....	3,254	-	-	-	-	-	1	2	4	15	35	73
1999.....	3,156	-	-	-	-	-	1	-	1	11	39	54

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Of gallbladder and extrahepatic bile ducts (C23-C24)

White, male

2004.....	1,117	-	-	-	1	-	-	3	3	2	13	20
2003.....	1,149	-	-	-	-	-	1	-	3	5	10	33
2002.....	1,119	-	-	-	-	-	-	2	1	6	11	15
2001.....	1,173	-	-	-	1	-	-	1	-	8	15	30
2000.....	1,233	-	-	-	-	-	1	2	2	8	14	29
1999.....	1,156	-	-	-	-	-	1	-	-	5	18	24

White, female

2004.....	1,800	-	-	-	-	-	1	-	4	7	16	21
2003.....	1,820	-	-	-	-	-	1	-	1	4	9	37
2002.....	1,826	-	-	-	1	-	-	2	1	7	25	48
2001.....	1,967	-	-	-	-	-	-	1	3	8	26	50
2000.....	2,021	-	-	-	-	-	-	-	2	7	21	44
1999.....	2,000	-	-	-	-	-	-	-	1	6	21	30

Black, both sexes

2004.....	314	-	-	-	-	-	-	-	-	-	7	6
2003.....	285	-	-	-	-	1	-	-	-	1	7	18
2002.....	346	-	-	-	-	-	-	1	-	5	5	19
2001.....	327	-	-	-	-	-	-	-	1	3	8	13
2000.....	290	-	-	-	-	-	-	-	-	-	3	10
1999.....	306	-	-	-	-	-	1	1	1	2	5	13

Black, male

2004.....	96	-	-	-	-	-	-	-	-	-	1	1
2003.....	90	-	-	-	-	1	-	-	-	-	3	5
2002.....	111	-	-	-	-	-	-	-	-	3	3	5
2001.....	104	-	-	-	-	-	-	-	-	1	1	3
2000.....	87	-	-	-	-	-	-	-	-	-	2	3
1999.....	98	-	-	-	-	-	-	1	-	-	1	5

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Of pancreas (C25)												
All races, both sexes												
2004.....	31,772	-	-	-	-	1	10	11	51	125	396	926
2003.....	30,777	-	-	-	1	-	10	20	40	153	395	891
2002.....	30,264	1	-	-	-	2	5	7	32	156	440	889
2001.....	29,803	-	1	1	1	1	1	11	44	167	407	833
2000.....	29,332	2	1	-	1	1	4	17	38	140	408	834
1999.....	29,082	1	-	-	-	-	3	9	46	178	416	887
All races, male												
2004.....	15,776	-	-	-	-	-	6	4	35	76	247	596
2003.....	15,060	-	-	-	1	-	6	11	24	97	232	556
2002.....	14,877	-	-	-	-	2	3	3	23	89	275	555
2001.....	14,467	-	1	-	-	1	1	5	25	99	222	503
2000.....	14,238	1	1	-	-	1	2	11	22	87	237	531
1999.....	14,176	-	-	-	-	-	1	7	26	107	262	575
All races, female												
2004.....	15,996	-	-	-	-	1	4	7	16	49	149	330
2003.....	15,717	-	-	-	-	-	4	9	16	56	163	335
2002.....	15,387	1	-	-	-	-	2	4	9	67	165	334
2001.....	15,336	-	-	1	1	-	-	6	19	68	185	330
2000.....	15,094	1	-	-	1	-	2	6	16	53	171	303
1999.....	14,906	1	-	-	-	-	2	2	20	71	154	312
White, both sexes												
2004.....	27,229	-	-	-	-	1	5	6	39	99	303	735
2003.....	26,418	-	-	-	-	-	7	18	30	121	300	699
2002.....	26,031	-	-	-	-	2	5	4	24	119	354	715
2001.....	25,769	-	1	1	1	1	1	7	35	134	327	652
2000.....	25,301	2	1	-	1	1	3	15	30	110	319	679
1999.....	25,086	-	-	-	-	-	2	7	33	137	331	729

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Of pancreas (C25)

White, male

2004.....	13,610	-	-	-	-	-	2	3	26	64	186	482
2003.....	13,033	-	-	-	-	-	3	10	16	78	173	434
2002.....	12,921	-	-	-	-	2	3	1	18	65	226	451
2001.....	12,641	-	1	-	-	1	1	4	21	77	182	403
2000.....	12,392	1	1	-	-	1	2	9	18	69	189	437
1999.....	12,273	-	-	-	-	-	1	6	19	85	212	476

White, female

2004.....	13,619	-	-	-	-	1	3	3	13	35	117	253
2003.....	13,385	-	-	-	-	-	4	8	14	43	127	265
2002.....	13,110	-	-	-	-	-	2	3	6	54	128	264
2001.....	13,128	-	-	1	1	-	-	3	14	57	145	249
2000.....	12,909	1	-	-	1	-	1	6	12	41	130	242
1999.....	12,813	-	-	-	-	-	1	1	14	52	119	253

Black, both sexes

2004.....	3,674	-	-	-	-	-	5	4	8	16	76	162
2003.....	3,616	-	-	-	1	-	3	1	9	23	80	168
2002.....	3,494	1	-	-	-	-	-	2	4	30	72	144
2001.....	3,394	-	-	-	-	-	-	4	6	26	64	157
2000.....	3,419	-	-	-	-	-	1	1	8	24	73	136
1999.....	3,391	1	-	-	-	-	-	2	10	36	76	141

Black, male

2004.....	1,737	-	-	-	-	-	4	-	7	8	47	95
2003.....	1,675	-	-	-	1	-	3	1	7	16	48	110
2002.....	1,617	-	-	-	-	-	-	1	4	19	39	85
2001.....	1,523	-	-	-	-	-	-	1	3	16	31	90
2000.....	1,561	-	-	-	-	-	-	1	4	15	39	80
1999.....	1,611	-	-	-	-	-	-	1	7	19	44	88

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Of pancreas (C25)												
Black, female												
2004.....	1,937	-	-	-	-	-	1	4	1	8	29	67
2003.....	1,941	-	-	-	-	-	-	-	2	7	32	58
2002.....	1,877	1	-	-	-	-	-	1	-	11	33	59
2001.....	1,871	-	-	-	-	-	-	3	3	10	33	67
2000.....	1,858	-	-	-	-	-	1	-	4	9	34	56
1999.....	1,780	1	-	-	-	-	-	1	3	17	32	53
American Indian 1/, both sexes												
2004.....	139	-	-	-	-	-	-	-	-	1	3	6
2003.....	103	-	-	-	-	-	-	1	-	4	3	6
2002.....	112	-	-	-	-	-	-	-	1	1	3	6
2001.....	89	-	-	-	-	-	-	-	1	-	3	5
2000.....	91	-	-	-	-	-	-	-	-	2	6	1
1999.....	84	-	-	-	-	-	-	-	1	-	2	3
American Indian 1/, male												
2004.....	71	-	-	-	-	-	-	-	-	-	2	5
2003.....	51	-	-	-	-	-	-	-	-	1	2	3
2002.....	50	-	-	-	-	-	-	-	-	1	3	5
2001.....	38	-	-	-	-	-	-	-	-	-	1	1
2000.....	46	-	-	-	-	-	-	-	-	-	5	1
1999.....	40	-	-	-	-	-	-	-	-	-	1	2
American Indian 1/, female												
2004.....	68	-	-	-	-	-	-	-	-	1	1	1
2003.....	52	-	-	-	-	-	-	1	-	3	1	3
2002.....	62	-	-	-	-	-	-	-	1	-	-	1
2001.....	51	-	-	-	-	-	-	-	1	-	2	4
2000.....	45	-	-	-	-	-	-	-	-	2	1	-
1999.....	44	-	-	-	-	-	-	-	1	-	1	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Of pancreas (C25)												
Asian or Pacific Islander 2/, both sexes												
2004.....	730	-	-	-	-	-	-	1	4	9	14	23
2003.....	640	-	-	-	-	-	-	-	1	5	12	18
2002.....	627	-	-	-	-	-	-	1	3	6	11	24
2001.....	551	-	-	-	-	-	-	-	2	7	13	19
2000.....	521	-	-	-	-	-	-	1	-	4	10	18
1999.....	521	-	-	-	-	-	1	-	2	5	7	14
Asian or Pacific Islander 2/, male												
2004.....	358	-	-	-	-	-	-	1	2	4	12	14
2003.....	301	-	-	-	-	-	-	-	1	2	9	9
2002.....	289	-	-	-	-	-	-	1	1	4	7	14
2001.....	265	-	-	-	-	-	-	-	1	6	8	9
2000.....	239	-	-	-	-	-	-	1	-	3	4	13
1999.....	252	-	-	-	-	-	-	-	-	3	5	9
Asian or Pacific Islander 2/, female												
2004.....	372	-	-	-	-	-	-	-	2	5	2	9
2003.....	339	-	-	-	-	-	-	-	-	3	3	9
2002.....	338	-	-	-	-	-	-	-	2	2	4	10
2001.....	286	-	-	-	-	-	-	-	1	1	5	10
2000.....	282	-	-	-	-	-	-	-	-	1	6	5
1999.....	269	-	-	-	-	-	1	-	2	2	2	5

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Of other and ill-defined digestive organs (C26)

All races, both sexes

2004.....	1,072	-	-	-	-	-	3	3	1	5	20	31
2003.....	1,347	-	-	-	1	2	2	3	4	8	14	35
2002.....	1,349	-	1	-	-	-	-	1	4	6	18	29
2001.....	1,409	-	-	-	-	1	-	-	2	8	18	32
2000.....	1,377	-	-	-	1	-	1	3	8	10	17	31
1999.....	1,566	-	-	-	-	1	2	4	3	7	29	51

All races, male

2004.....	504	-	-	-	-	-	2	2	1	4	12	14
2003.....	664	-	-	-	1	1	-	1	3	5	12	19
2002.....	670	-	1	-	-	-	-	-	3	4	7	18
2001.....	686	-	-	-	-	1	-	-	2	4	10	22
2000.....	665	-	-	-	1	-	1	1	6	7	9	16
1999.....	774	-	-	-	-	1	1	1	3	7	21	33

All races, female

2004.....	568	-	-	-	-	-	1	1	-	1	8	17
2003.....	683	-	-	-	-	1	2	2	1	3	2	16
2002.....	679	-	-	-	-	-	-	1	1	2	11	11
2001.....	723	-	-	-	-	-	-	-	-	4	8	10
2000.....	712	-	-	-	-	-	-	2	2	3	8	15
1999.....	792	-	-	-	-	-	1	3	-	-	8	18

White, both sexes

2004.....	929	-	-	-	-	-	3	2	1	3	14	25
2003.....	1,163	-	-	-	-	2	1	3	2	5	13	23
2002.....	1,172	-	1	-	-	-	-	1	2	4	14	19
2001.....	1,203	-	-	-	-	1	-	-	2	6	14	22
2000.....	1,184	-	-	-	-	-	1	3	5	10	13	27
1999.....	1,375	-	-	-	-	1	2	3	2	6	23	35

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Of other and ill-defined digestive organs (C26)

White, male

2004.....	437	-	-	-	-	-	2	1	1	3	7	13
2003.....	570	-	-	-	-	1	-	1	1	3	12	12
2002.....	579	-	1	-	-	-	-	-	1	2	6	13
2001.....	590	-	-	-	-	1	-	-	2	3	9	15
2000.....	580	-	-	-	-	-	1	1	3	7	8	14
1999.....	669	-	-	-	-	1	1	1	2	6	16	22

White, female

2004.....	492	-	-	-	-	-	1	1	-	-	7	12
2003.....	593	-	-	-	-	1	1	2	1	2	1	11
2002.....	593	-	-	-	-	-	-	1	1	2	8	6
2001.....	613	-	-	-	-	-	-	-	-	3	5	7
2000.....	604	-	-	-	-	-	-	2	2	3	5	13
1999.....	706	-	-	-	-	-	1	2	-	-	7	13

Black, both sexes

2004.....	116	-	-	-	-	-	-	1	-	1	6	5
2003.....	151	-	-	-	1	-	1	-	1	3	1	8
2002.....	151	-	-	-	-	-	-	-	2	2	4	6
2001.....	169	-	-	-	-	-	-	-	-	1	2	7
2000.....	160	-	-	-	1	-	-	-	2	-	4	4
1999.....	164	-	-	-	-	-	-	1	1	-	6	16

Black, male

2004.....	55	-	-	-	-	-	-	1	-	-	5	-
2003.....	79	-	-	-	1	-	-	-	1	2	-	5
2002.....	81	-	-	-	-	-	-	-	2	2	1	3
2001.....	79	-	-	-	-	-	-	-	-	-	-	4
2000.....	68	-	-	-	1	-	-	-	2	-	1	2
1999.....	87	-	-	-	-	-	-	-	1	-	5	11

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Malignant neoplasms of respiratory and intrathoracic organs (C30-C39)

All races, both sexes

2004.....	162,664	2	4	2	3	13	20	53	121	501	2,040	4,783
2003.....	162,841	-	3	-	2	11	27	44	135	552	2,072	4,828
2002.....	162,388	2	1	3	3	8	34	47	156	633	2,201	4,570
2001.....	160,876	-	6	3	5	11	23	45	136	686	2,270	4,477
2000.....	160,288	2	2	1	3	9	28	46	176	708	2,194	4,421
1999.....	156,914	1	-	6	4	8	21	54	156	716	2,163	4,112

All races, male

2004.....	93,073	1	1	1	2	8	12	36	71	252	1,106	2,736
2003.....	93,544	-	2	-	1	6	18	20	77	279	1,165	2,823
2002.....	93,656	2	1	2	3	5	29	30	85	327	1,155	2,651
2001.....	94,004	-	4	3	2	5	16	21	68	343	1,269	2,653
2000.....	94,053	2	1	1	-	8	18	37	85	400	1,248	2,690
1999.....	92,965	1	-	4	3	6	12	33	83	361	1,228	2,506

All races, female

2004.....	69,591	1	3	1	1	5	8	17	50	249	934	2,047
2003.....	69,297	-	1	-	1	5	9	24	58	273	907	2,005
2002.....	68,732	-	-	1	-	3	5	17	71	306	1,046	1,919
2001.....	66,872	-	2	-	3	6	7	24	68	343	1,001	1,824
2000.....	66,235	-	1	-	3	1	10	9	91	308	946	1,731
1999.....	63,949	-	-	2	1	2	9	21	73	355	935	1,606

White, both sexes

2004.....	142,174	1	3	2	3	9	18	43	90	401	1,590	3,767
2003.....	142,626	-	3	-	1	9	21	34	111	438	1,627	3,725
2002.....	142,503	2	1	3	1	7	23	34	121	500	1,715	3,588
2001.....	141,193	-	5	3	4	8	20	36	116	552	1,777	3,484
2000.....	140,768	1	2	1	3	5	21	35	141	531	1,664	3,406
1999.....	137,674	1	-	4	2	4	15	42	130	546	1,687	3,159

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Malignant neoplasms of respiratory and intrathoracic organs (C30-C39)

White, male

2004.....	80,530	1	-	1	2	5	11	29	53	195	855	2,117
2003.....	81,101	-	2	-	-	4	13	16	61	222	899	2,167
2002.....	81,301	2	1	2	1	5	19	22	66	247	883	2,041
2001.....	81,555	-	3	3	1	4	14	19	58	275	972	2,024
2000.....	81,638	1	1	1	-	4	15	28	71	302	911	2,060
1999.....	80,716	1	-	2	2	3	8	25	72	276	939	1,891

White, female

2004.....	61,644	-	3	1	1	4	7	14	37	206	735	1,650
2003.....	61,525	-	1	-	1	5	8	18	50	216	728	1,558
2002.....	61,202	-	-	1	-	2	4	12	55	253	832	1,547
2001.....	59,638	-	2	-	3	4	6	17	58	277	805	1,460
2000.....	59,130	-	1	-	3	1	6	7	70	229	753	1,346
1999.....	56,958	-	-	2	-	1	7	17	58	270	748	1,268

Black, both sexes

2004.....	17,216	1	1	-	-	3	2	8	20	75	392	881
2003.....	17,169	-	-	-	1	1	5	6	19	83	385	986
2002.....	17,086	-	-	-	2	1	8	7	23	110	433	888
2001.....	16,812	-	1	-	1	2	2	5	13	116	442	863
2000.....	16,908	1	-	-	-	4	4	9	27	153	473	892
1999.....	16,729	-	-	1	2	3	5	6	21	146	433	859

Black, male

2004.....	10,581	-	1	-	-	3	1	6	9	38	213	545
2003.....	10,624	-	-	-	1	1	4	3	13	40	228	590
2002.....	10,677	-	-	-	2	-	7	5	15	68	247	552
2001.....	10,737	-	1	-	1	-	1	1	6	58	264	559
2000.....	10,799	1	-	-	-	4	1	7	11	84	305	560
1999.....	10,727	-	-	1	1	2	3	5	9	71	267	565

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Malignant neoplasms of respiratory and intrathoracic organs (C30-C39)

Black, female

2004.....	6,635	1	-	-	-	-	1	2	11	37	179	336
2003.....	6,545	-	-	-	-	-	1	3	6	43	157	396
2002.....	6,409	-	-	-	-	1	1	2	8	42	186	336
2001.....	6,075	-	-	-	-	2	1	4	7	58	178	304
2000.....	6,109	-	-	-	-	-	3	2	16	69	168	332
1999.....	6,002	-	-	-	1	1	2	1	12	75	166	294

American Indian 1/, both sexes

2004.....	698	-	-	-	-	-	-	-	1	4	11	32
2003.....	575	-	-	-	-	-	-	-	-	3	11	23
2002.....	572	-	-	-	-	-	-	-	1	-	6	15
2001.....	577	-	-	-	-	-	-	-	1	2	8	26
2000.....	497	-	-	-	-	-	-	-	1	3	8	17
1999.....	513	-	-	-	-	-	-	-	-	3	5	14

American Indian 1/, male

2004.....	387	-	-	-	-	-	-	-	1	2	7	15
2003.....	315	-	-	-	-	-	-	-	-	2	7	12
2002.....	317	-	-	-	-	-	-	-	-	-	3	8
2001.....	321	-	-	-	-	-	-	-	-	1	8	15
2000.....	281	-	-	-	-	-	-	-	1	1	4	9
1999.....	296	-	-	-	-	-	-	-	-	2	1	9

American Indian 1/, female

2004.....	311	-	-	-	-	-	-	-	-	2	4	17
2003.....	260	-	-	-	-	-	-	-	-	1	4	11
2002.....	255	-	-	-	-	-	-	-	1	-	3	7
2001.....	256	-	-	-	-	-	-	-	1	1	-	11
2000.....	216	-	-	-	-	-	-	-	-	2	4	8
1999.....	217	-	-	-	-	-	-	-	-	1	4	5

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Malignant neoplasms of respiratory and intrathoracic organs (C30-C39)												
Asian or Pacific Islander 2/, both sexes												
2004.....	2,576	-	-	-	-	1	-	2	10	21	47	103
2003.....	2,471	-	-	-	-	1	1	4	5	28	49	94
2002.....	2,227	-	-	-	-	-	3	6	11	23	47	79
2001.....	2,294	-	-	-	-	1	1	4	6	16	43	104
2000.....	2,115	-	-	-	-	-	3	2	7	21	49	106
1999.....	1,998	-	-	1	-	1	1	6	5	21	38	80
Asian or Pacific Islander 2/, male												
2004.....	1,575	-	-	-	-	-	-	1	8	17	31	59
2003.....	1,504	-	-	-	-	1	1	1	3	15	31	54
2002.....	1,361	-	-	-	-	-	3	3	4	12	22	50
2001.....	1,391	-	-	-	-	1	1	1	4	9	25	55
2000.....	1,335	-	-	-	-	-	2	2	2	13	28	61
1999.....	1,226	-	-	1	-	1	1	3	2	12	21	41
Asian or Pacific Islander 2/, female												
2004.....	1,001	-	-	-	-	1	-	1	2	4	16	44
2003.....	967	-	-	-	-	-	-	3	2	13	18	40
2002.....	866	-	-	-	-	-	-	3	7	11	25	29
2001.....	903	-	-	-	-	-	-	3	2	7	18	49
2000.....	780	-	-	-	-	-	1	-	5	8	21	45
1999.....	772	-	-	-	-	-	-	3	3	9	17	39

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Of nasal cavity, middle ear and accessory sinuses (C30-C31)

All races, both sexes

2004.....	458	-	-	-	-	-	-	2	5	13	15	25
2003.....	457	-	-	-	-	1	2	2	1	9	24	21
2002.....	444	-	-	-	-	-	3	1	4	12	11	33
2001.....	485	-	-	-	1	3	2	1	2	11	20	31
2000.....	419	-	-	-	-	2	2	2	2	13	20	18
1999.....	456	-	-	-	1	1	3	1	4	8	29	23

All races, male

2004.....	260	-	-	-	-	-	-	2	5	5	10	16
2003.....	282	-	-	-	-	1	2	2	-	5	12	15
2002.....	275	-	-	-	-	-	3	1	2	9	8	28
2001.....	262	-	-	-	-	1	-	-	1	6	12	22
2000.....	253	-	-	-	-	2	2	2	1	10	15	11
1999.....	257	-	-	-	-	-	1	1	1	5	17	17

All races, female

2004.....	198	-	-	-	-	-	-	-	-	8	5	9
2003.....	175	-	-	-	-	-	-	-	1	4	12	6
2002.....	169	-	-	-	-	-	-	-	2	3	3	5
2001.....	223	-	-	-	1	2	2	1	1	5	8	9
2000.....	166	-	-	-	-	-	-	-	1	3	5	7
1999.....	199	-	-	-	1	1	2	-	3	3	12	6

White, both sexes

2004.....	394	-	-	-	-	-	-	1	5	12	15	18
2003.....	399	-	-	-	-	-	1	2	1	8	21	18
2002.....	398	-	-	-	-	-	2	-	4	9	9	28
2001.....	420	-	-	-	1	2	2	-	2	8	14	26
2000.....	356	-	-	-	-	1	2	1	1	9	15	13
1999.....	408	-	-	-	-	1	3	1	2	6	25	17

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Of nasal cavity, middle ear and accessory sinuses (C30-C31)

White, male

2004.....	219	-	-	-	-	-	-	1	5	5	10	11
2003.....	240	-	-	-	-	-	1	2	-	5	10	13
2002.....	243	-	-	-	-	-	2	-	2	7	7	24
2001.....	224	-	-	-	-	1	-	-	1	5	9	19
2000.....	213	-	-	-	-	1	2	1	-	8	10	7
1999.....	227	-	-	-	-	-	1	1	1	4	14	11

White, female

2004.....	175	-	-	-	-	-	-	-	-	7	5	7
2003.....	159	-	-	-	-	-	-	-	1	3	11	5
2002.....	155	-	-	-	-	-	-	-	2	2	2	4
2001.....	196	-	-	-	1	1	2	-	1	3	5	7
2000.....	143	-	-	-	-	-	-	-	1	1	5	6
1999.....	181	-	-	-	-	1	2	-	1	2	11	6

Black, both sexes

2004.....	46	-	-	-	-	-	-	1	-	1	-	6
2003.....	47	-	-	-	-	1	1	-	-	1	2	3
2002.....	38	-	-	-	-	-	-	1	-	3	2	5
2001.....	50	-	-	-	-	1	-	1	-	2	5	3
2000.....	57	-	-	-	-	1	-	1	1	4	5	4
1999.....	42	-	-	-	1	-	-	-	2	2	3	5

Black, male

2004.....	31	-	-	-	-	-	-	1	-	-	-	4
2003.....	34	-	-	-	-	1	1	-	-	-	1	2
2002.....	27	-	-	-	-	-	-	1	-	2	1	4
2001.....	31	-	-	-	-	-	-	-	-	1	2	2
2000.....	36	-	-	-	-	1	-	1	1	2	5	3
1999.....	27	-	-	-	-	-	-	-	-	1	3	5

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Of larynx (C32)												
All races, both sexes												
2004.....	3,668	-	-	1	-	1	-	-	3	7	52	161
2003.....	3,792	-	-	-	-	-	-	-	1	10	69	157
2002.....	3,723	-	-	-	-	-	-	3	8	19	57	156
2001.....	3,797	-	-	-	-	-	-	1	2	18	68	149
2000.....	3,861	-	-	-	2	1	-	-	4	9	55	156
1999.....	3,815	-	-	-	-	-	-	-	6	20	58	151
All races, male												
2004.....	2,935	-	-	-	-	1	-	-	3	5	43	134
2003.....	3,020	-	-	-	-	-	-	-	1	7	53	129
2002.....	2,922	-	-	-	-	-	-	3	3	16	42	130
2001.....	3,018	-	-	-	-	-	-	1	1	12	54	123
2000.....	3,041	-	-	-	-	-	-	-	2	6	43	122
1999.....	2,992	-	-	-	-	-	-	-	5	18	46	130
All races, female												
2004.....	733	-	-	1	-	-	-	-	-	2	9	27
2003.....	772	-	-	-	-	-	-	-	-	3	16	28
2002.....	801	-	-	-	-	-	-	-	5	3	15	26
2001.....	779	-	-	-	-	-	-	-	1	6	14	26
2000.....	820	-	-	-	2	1	-	-	2	3	12	34
1999.....	823	-	-	-	-	-	-	-	1	2	12	21
White, both sexes												
2004.....	2,985	-	-	1	-	1	-	-	3	5	37	127
2003.....	3,052	-	-	-	-	-	-	-	1	9	53	110
2002.....	3,012	-	-	-	-	-	-	2	5	10	38	104
2001.....	3,049	-	-	-	-	-	-	1	2	11	50	98
2000.....	3,096	-	-	-	2	1	-	-	4	8	41	108
1999.....	3,106	-	-	-	-	-	-	-	4	14	31	101

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Of larynx (C32)												
White, male												
2004.....	2,370	-	-	-	-	1	-	-	3	4	32	107
2003.....	2,410	-	-	-	-	-	-	-	1	6	43	94
2002.....	2,351	-	-	-	-	-	-	2	1	8	29	88
2001.....	2,417	-	-	-	-	-	-	1	1	10	40	79
2000.....	2,426	-	-	-	-	-	-	-	2	5	30	86
1999.....	2,419	-	-	-	-	-	-	-	3	14	24	87
White, female												
2004.....	615	-	-	1	-	-	-	-	-	1	5	20
2003.....	642	-	-	-	-	-	-	-	-	3	10	16
2002.....	661	-	-	-	-	-	-	-	4	2	9	16
2001.....	632	-	-	-	-	-	-	-	1	1	10	19
2000.....	670	-	-	-	2	1	-	-	2	3	11	22
1999.....	687	-	-	-	-	-	-	-	1	-	7	14
Black, both sexes												
2004.....	638	-	-	-	-	-	-	-	-	2	15	33
2003.....	693	-	-	-	-	-	-	-	-	1	15	44
2002.....	675	-	-	-	-	-	-	1	3	8	19	48
2001.....	678	-	-	-	-	-	-	-	-	7	17	48
2000.....	719	-	-	-	-	-	-	-	-	1	14	47
1999.....	663	-	-	-	-	-	-	-	2	6	26	48
Black, male												
2004.....	530	-	-	-	-	-	-	-	-	1	11	27
2003.....	571	-	-	-	-	-	-	-	-	1	9	33
2002.....	542	-	-	-	-	-	-	1	2	7	13	38
2001.....	541	-	-	-	-	-	-	-	-	2	13	41
2000.....	578	-	-	-	-	-	-	-	-	1	13	35
1999.....	536	-	-	-	-	-	-	-	2	4	21	41

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Of trachea, bronchus and lung (C33-C34)

All races, both sexes

2004.....	158,091	2	4	1	3	6	17	44	103	470	1,956	4,564
2003.....	158,086	-	2	-	2	7	18	29	125	520	1,958	4,617
2002.....	157,713	-	1	2	2	4	17	35	130	583	2,115	4,341
2001.....	156,058	-	5	1	1	6	14	36	121	640	2,156	4,255
2000.....	155,521	2	2	-	1	3	19	37	160	664	2,092	4,220
1999.....	152,156	-	-	3	2	6	14	43	136	668	2,056	3,916

All races, male

2004.....	89,630	1	1	1	2	4	10	30	57	236	1,044	2,566
2003.....	89,964	-	1	-	1	4	10	9	69	258	1,086	2,659
2002.....	90,171	-	1	1	2	2	14	19	71	290	1,096	2,471
2001.....	90,426	-	4	1	-	2	10	17	60	315	1,189	2,487
2000.....	90,469	2	1	-	-	3	11	28	74	370	1,172	2,540
1999.....	89,453	-	-	2	2	5	8	24	73	323	1,151	2,346

All races, female

2004.....	68,461	1	3	-	1	2	7	14	46	234	912	1,998
2003.....	68,122	-	1	-	1	3	8	20	56	262	872	1,958
2002.....	67,542	-	-	1	-	2	3	16	59	293	1,019	1,870
2001.....	65,632	-	1	-	1	4	4	19	61	325	967	1,768
2000.....	65,052	-	1	-	1	-	8	9	86	294	920	1,680
1999.....	62,703	-	-	1	-	1	6	19	63	345	905	1,570

White, both sexes

2004.....	138,425	1	3	1	3	3	15	36	73	376	1,524	3,594
2003.....	138,743	-	2	-	1	7	15	24	103	409	1,541	3,571
2002.....	138,661	-	1	2	1	3	14	25	102	465	1,655	3,426
2001.....	137,274	-	4	1	1	5	11	29	101	522	1,692	3,328
2000.....	136,912	1	2	-	1	1	13	27	128	499	1,594	3,265
1999.....	133,752	-	-	2	1	3	8	33	116	511	1,617	3,022

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Of trachea, bronchus and lung (C33-C34)

White, male

2004.....	77,730	1	-	1	2	2	9	25	39	182	806	1,980
2003.....	78,215	-	1	-	-	4	8	8	54	203	838	2,042
2002.....	78,465	-	1	1	1	2	11	14	58	222	841	1,913
2001.....	78,665	-	3	1	-	2	8	15	50	253	912	1,907
2000.....	78,747	1	1	-	-	1	9	20	62	277	859	1,953
1999.....	77,846	-	-	1	1	3	4	17	64	247	891	1,782

White, female

2004.....	60,695	-	3	-	1	1	6	11	34	194	718	1,614
2003.....	60,528	-	1	-	1	3	7	16	49	206	703	1,529
2002.....	60,196	-	-	1	-	1	3	11	44	243	814	1,513
2001.....	58,609	-	1	-	1	3	3	14	51	269	780	1,421
2000.....	58,165	-	1	-	1	-	4	7	66	222	735	1,312
1999.....	55,906	-	-	1	-	-	4	16	52	264	726	1,240

Black, both sexes

2004.....	16,472	1	1	-	-	2	2	7	19	70	374	838
2003.....	16,372	-	-	-	1	-	3	2	17	81	361	932
2002.....	16,316	-	-	-	1	1	3	4	19	96	409	827
2001.....	16,019	-	1	-	-	1	2	3	13	103	416	804
2000.....	16,066	1	-	-	-	2	3	8	24	142	442	837
1999.....	15,963	-	-	1	1	3	5	5	15	134	399	804

Black, male

2004.....	9,995	-	1	-	-	2	1	5	9	36	200	513
2003.....	9,988	-	-	-	1	-	2	1	12	39	213	553
2002.....	10,075	-	-	-	1	-	3	2	12	57	232	505
2001.....	10,131	-	1	-	-	-	1	1	6	54	247	515
2000.....	10,153	1	-	-	-	2	-	6	9	80	282	520
1999.....	10,135	-	-	1	1	2	3	4	7	63	240	518

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Of trachea, bronchus and lung (C33-C34)

Black, female

2004.....	6,477	1	-	-	-	-	1	2	10	34	174	325
2003.....	6,384	-	-	-	-	-	1	1	5	42	148	379
2002.....	6,241	-	-	-	-	1	-	2	7	39	177	322
2001.....	5,888	-	-	-	-	1	1	2	7	49	169	289
2000.....	5,913	-	-	-	-	-	3	2	15	62	160	317
1999.....	5,828	-	-	-	-	1	2	1	8	71	159	286

American Indian 1/, both sexes

2004.....	677	-	-	-	-	-	-	-	1	4	11	32
2003.....	556	-	-	-	-	-	-	-	-	3	8	21
2002.....	556	-	-	-	-	-	-	-	1	-	5	14
2001.....	547	-	-	-	-	-	-	-	1	1	7	24
2000.....	480	-	-	-	-	-	-	-	1	3	8	17
1999.....	497	-	-	-	-	-	-	-	-	3	5	13

American Indian 1/, male

2004.....	373	-	-	-	-	-	-	-	1	2	7	15
2003.....	301	-	-	-	-	-	-	-	-	2	5	11
2002.....	304	-	-	-	-	-	-	-	-	-	2	7
2001.....	299	-	-	-	-	-	-	-	-	1	7	13
2000.....	270	-	-	-	-	-	-	-	1	1	4	9
1999.....	285	-	-	-	-	-	-	-	-	2	1	8

American Indian 1/, female

2004.....	304	-	-	-	-	-	-	-	-	2	4	17
2003.....	255	-	-	-	-	-	-	-	-	1	3	10
2002.....	252	-	-	-	-	-	-	-	1	-	3	7
2001.....	248	-	-	-	-	-	-	-	1	-	-	11
2000.....	210	-	-	-	-	-	-	-	-	2	4	8
1999.....	212	-	-	-	-	-	-	-	-	1	4	5

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Of trachea, bronchus and lung (C33-C34)												
Asian or Pacific Islander 2/, both sexes												
2004.....	2,517	-	-	-	-	1	-	1	10	20	47	100
2003.....	2,415	-	-	-	-	-	-	3	5	27	48	93
2002.....	2,180	-	-	-	-	-	-	6	8	22	46	74
2001.....	2,218	-	-	-	-	-	1	4	6	14	41	99
2000.....	2,063	-	-	-	-	-	3	2	7	20	48	101
1999.....	1,944	-	-	-	-	-	1	5	5	20	35	77
Asian or Pacific Islander 2/, male												
2004.....	1,532	-	-	-	-	-	-	-	8	16	31	58
2003.....	1,460	-	-	-	-	-	-	-	3	14	30	53
2002.....	1,327	-	-	-	-	-	-	3	1	11	21	46
2001.....	1,331	-	-	-	-	-	1	1	4	7	23	52
2000.....	1,299	-	-	-	-	-	2	2	2	12	27	58
1999.....	1,187	-	-	-	-	-	1	3	2	11	19	38
Asian or Pacific Islander 2/, female												
2004.....	985	-	-	-	-	1	-	1	2	4	16	42
2003.....	955	-	-	-	-	-	-	3	2	13	18	40
2002.....	853	-	-	-	-	-	-	3	7	11	25	28
2001.....	887	-	-	-	-	-	-	3	2	7	18	47
2000.....	764	-	-	-	-	-	1	-	5	8	21	43
1999.....	757	-	-	-	-	-	-	2	3	9	16	39

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Of thymus, heart, mediastinum and pleura (C37-C38)												
All races, both sexes												
2004.....	436	-	-	-	-	6	3	7	10	11	17	33
2003.....	452	-	1	-	-	3	7	12	8	12	21	29
2002.....	460	2	-	1	1	4	14	8	14	18	17	36
2001.....	484	-	1	2	3	2	7	7	11	17	26	39
2000.....	441	-	-	1	-	3	7	7	10	20	25	25
1999.....	445	1	-	2	1	1	4	10	9	20	19	21
All races, male												
2004.....	242	-	-	-	-	3	2	4	6	6	9	20
2003.....	245	-	1	-	-	1	6	9	7	8	14	17
2002.....	261	2	-	1	1	3	12	7	9	11	9	21
2001.....	264	-	-	2	2	2	6	3	6	10	14	21
2000.....	258	-	-	1	-	3	5	7	8	13	16	15
1999.....	235	1	-	2	1	1	3	8	3	15	13	12
All races, female												
2004.....	194	-	-	-	-	3	1	3	4	5	8	13
2003.....	207	-	-	-	-	2	1	3	1	4	7	12
2002.....	199	-	-	-	-	1	2	1	5	7	8	15
2001.....	220	-	1	-	1	-	1	4	5	7	12	18
2000.....	183	-	-	-	-	-	2	-	2	7	9	10
1999.....	210	-	-	-	-	-	1	2	6	5	6	9
White, both sexes												
2004.....	364	-	-	-	-	5	3	6	9	8	14	28
2003.....	384	-	1	-	-	2	5	7	6	11	12	23
2002.....	390	2	-	1	-	4	7	7	10	15	12	28
2001.....	409	-	1	2	2	1	7	6	11	11	21	31
2000.....	363	-	-	1	-	2	6	7	8	14	12	18
1999.....	371	1	-	1	1	-	4	8	7	15	14	18

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Of thymus, heart, mediastinum and pleura (C37-C38)

White, male

2004.....	208	-	-	-	-	2	2	3	6	4	7	19
2003.....	206	-	1	-	-	-	4	6	6	7	8	15
2002.....	218	2	-	1	-	3	6	6	5	9	6	15
2001.....	223	-	-	2	1	1	6	3	6	7	11	19
2000.....	222	-	-	1	-	2	4	7	7	11	10	12
1999.....	201	1	-	1	1	-	3	7	3	11	10	10

White, female

2004.....	156	-	-	-	-	3	1	3	3	4	7	9
2003.....	178	-	-	-	-	2	1	1	-	4	4	8
2002.....	172	-	-	-	-	1	1	1	5	6	6	13
2001.....	186	-	1	-	1	-	1	3	5	4	10	12
2000.....	141	-	-	-	-	-	2	-	1	3	2	6
1999.....	170	-	-	-	-	-	1	1	4	4	4	8

Black, both sexes

2004.....	55	-	-	-	-	1	-	-	1	2	3	4
2003.....	51	-	-	-	-	-	1	4	2	-	7	6
2002.....	51	-	-	-	1	-	5	1	1	3	3	6
2001.....	56	-	-	-	1	-	-	1	-	4	4	6
2000.....	61	-	-	-	-	1	1	-	2	5	12	4
1999.....	57	-	-	-	-	-	-	1	2	4	4	2

Black, male

2004.....	22	-	-	-	-	1	-	-	-	1	2	1
2003.....	28	-	-	-	-	-	1	2	1	-	5	2
2002.....	30	-	-	-	1	-	4	1	1	2	1	5
2001.....	27	-	-	-	1	-	-	-	-	1	2	1
2000.....	30	-	-	-	-	1	1	-	1	1	5	2
1999.....	25	-	-	-	-	-	-	1	-	3	2	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Of thymus, heart, mediastinum and pleura (C37-C38)												
Asian or Pacific Islander 2/, both sexes												
2004.....	13	-	-	-	-	-	-	1	-	1	-	1
2003.....	14	-	-	-	-	1	1	1	-	1	-	-
2002.....	17	-	-	-	-	-	2	-	3	-	1	2
2001.....	16	-	-	-	-	1	-	-	-	2	1	2
2000.....	17	-	-	-	-	-	-	-	-	1	1	3
1999.....	15	-	-	1	-	1	-	1	-	1	1	1
Asian or Pacific Islander 2/, male												
2004.....	10	-	-	-	-	-	-	1	-	1	-	-
2003.....	10	-	-	-	-	1	1	1	-	1	-	-
2002.....	11	-	-	-	-	-	2	-	3	-	1	1
2001.....	12	-	-	-	-	1	-	-	-	2	1	1
2000.....	6	-	-	-	-	-	-	-	-	1	1	1
1999.....	9	-	-	1	-	1	-	-	-	1	1	1
Asian or Pacific Islander 2/, female												
2004.....	3	-	-	-	-	-	-	-	-	-	-	1
2003.....	4	-	-	-	-	-	-	-	-	-	-	-
2002.....	6	-	-	-	-	-	-	-	-	-	-	1
2001.....	4	-	-	-	-	-	-	-	-	-	-	1
2000.....	11	-	-	-	-	-	-	-	-	-	-	2
1999.....	6	-	-	-	-	-	-	1	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Of other and ill-defined sites in respiratory system and intrathoracic organs (C39)

All races, both sexes

2004.....	11	-	-	-	-	-	-	-	-	-	-	-
2003.....	54	-	-	-	-	-	-	1	-	1	-	4
2002.....	48	-	-	-	-	-	-	-	-	1	1	4
2001.....	52	-	-	-	-	-	-	-	-	-	-	3
2000.....	46	-	-	-	-	-	-	-	-	2	2	2
1999.....	42	-	-	1	-	-	-	-	1	-	1	1

All races, male

2004.....	6	-	-	-	-	-	-	-	-	-	-	-
2003.....	33	-	-	-	-	-	-	-	-	1	-	3
2002.....	27	-	-	-	-	-	-	-	-	1	-	1
2001.....	34	-	-	-	-	-	-	-	-	-	-	-
2000.....	32	-	-	-	-	-	-	-	-	1	2	2
1999.....	28	-	-	-	-	-	-	-	1	-	1	1

All races, female

2004.....	5	-	-	-	-	-	-	-	-	-	-	-
2003.....	21	-	-	-	-	-	-	1	-	-	-	1
2002.....	21	-	-	-	-	-	-	-	-	-	1	3
2001.....	18	-	-	-	-	-	-	-	-	-	-	3
2000.....	14	-	-	-	-	-	-	-	-	1	-	-
1999.....	14	-	-	1	-	-	-	-	-	-	-	-

White, both sexes

2004.....	6	-	-	-	-	-	-	-	-	-	-	-
2003.....	48	-	-	-	-	-	-	1	-	1	-	3
2002.....	42	-	-	-	-	-	-	-	-	1	1	2
2001.....	41	-	-	-	-	-	-	-	-	-	-	1
2000.....	41	-	-	-	-	-	-	-	-	1	2	2
1999.....	37	-	-	1	-	-	-	-	1	-	-	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasms of bone and articular cartilage (C40-C41)

All races, both sexes

2004.....	1,301	3	3	12	57	114	81	53	41	42	45	54
2003.....	1,262	-	5	13	64	104	86	52	40	44	63	55
2002.....	1,194	1	2	21	61	94	81	55	44	40	44	57
2001.....	1,298	-	2	14	72	101	94	52	41	44	50	53
2000.....	1,212	-	2	19	62	98	77	39	40	39	39	55
1999.....	1,224	-	3	24	55	127	75	42	45	43	48	67

All races, male

2004.....	715	3	-	8	30	77	54	33	23	26	27	42
2003.....	723	-	4	9	36	73	52	28	25	36	39	36
2002.....	664	-	1	12	29	54	52	40	26	26	30	38
2001.....	717	-	2	8	39	65	69	30	20	24	33	37
2000.....	682	-	2	5	37	61	50	26	29	24	25	36
1999.....	667	-	-	12	29	76	47	30	24	24	29	38

All races, female

2004.....	586	-	3	4	27	37	27	20	18	16	18	12
2003.....	539	-	1	4	28	31	34	24	15	8	24	19
2002.....	530	1	1	9	32	40	29	15	18	14	14	19
2001.....	581	-	-	6	33	36	25	22	21	20	17	16
2000.....	530	-	-	14	25	37	27	13	11	15	14	19
1999.....	557	-	3	12	26	51	28	12	21	19	19	29

White, both sexes

2004.....	1,098	3	3	9	44	88	65	46	32	27	39	42
2003.....	1,087	-	3	12	52	89	66	41	35	39	51	45
2002.....	1,030	1	2	17	50	81	68	47	36	34	35	48
2001.....	1,118	-	2	12	57	88	80	47	31	33	43	43
2000.....	1,061	-	2	16	53	73	67	35	35	32	31	48
1999.....	1,066	-	2	18	43	105	63	35	40	37	39	54

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Malignant neoplasms of bone and articular cartilage (C40-C41)

White, male

2004.....	602	3	-	6	24	59	45	29	19	17	24	34
2003.....	624	-	2	8	29	64	37	22	23	33	30	27
2002.....	574	-	1	10	22	46	42	35	19	25	24	32
2001.....	622	-	2	6	30	57	59	26	15	21	30	30
2000.....	592	-	2	5	32	43	43	24	28	19	19	29
1999.....	574	-	-	10	25	59	39	24	22	20	22	31

White, female

2004.....	496	-	3	3	20	29	20	17	13	10	15	8
2003.....	463	-	1	4	23	25	29	19	12	6	21	18
2002.....	456	1	1	7	28	35	26	12	17	9	11	16
2001.....	496	-	-	6	27	31	21	21	16	12	13	13
2000.....	469	-	-	11	21	30	24	11	7	13	12	19
1999.....	492	-	2	8	18	46	24	11	18	17	17	23

Black, both sexes

2004.....	170	-	-	2	12	21	13	6	6	11	5	10
2003.....	137	-	1	1	10	9	15	8	2	3	12	6
2002.....	136	-	-	3	10	10	9	5	7	6	7	6
2001.....	147	-	-	2	11	10	13	4	8	9	5	8
2000.....	127	-	-	2	8	22	8	3	4	5	8	6
1999.....	123	-	1	4	8	16	6	3	3	5	8	12

Black, male

2004.....	93	-	-	1	5	16	8	4	3	5	3	7
2003.....	78	-	1	1	5	5	13	5	-	2	9	6
2002.....	75	-	-	1	6	7	7	3	6	1	5	5
2001.....	75	-	-	2	7	5	10	3	4	2	1	5
2000.....	76	-	-	-	5	16	6	1	-	4	6	6
1999.....	71	-	-	2	2	12	3	3	1	3	6	6

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasms of bone and articular cartilage (C40-C41)

Black, female

2004.....	77	-	-	1	7	5	5	2	3	6	2	3
2003.....	59	-	-	-	5	4	2	3	2	1	3	-
2002.....	61	-	-	2	4	3	2	2	1	5	2	1
2001.....	72	-	-	-	4	5	3	1	4	7	4	3
2000.....	51	-	-	2	3	6	2	2	4	1	2	-
1999.....	52	-	1	2	6	4	3	-	2	2	2	6

American Indian 1/, both sexes

2004.....	10	-	-	1	-	3	1	-	-	-	-	-
2003.....	9	-	1	-	-	1	-	1	2	1	-	1
2002.....	6	-	-	-	-	1	-	-	-	-	-	1
2001.....	7	-	-	-	1	2	1	-	-	-	-	-
2000.....	4	-	-	-	-	-	1	-	-	1	-	-
1999.....	8	-	-	-	1	-	3	2	-	-	-	-

American Indian 1/, male

2004.....	6	-	-	1	-	1	-	-	-	-	-	-
2003.....	5	-	1	-	-	1	-	-	1	-	-	1
2002.....	2	-	-	-	-	-	-	-	-	-	-	-
2001.....	4	-	-	-	1	2	-	-	-	-	-	-
2000.....	2	-	-	-	-	-	-	-	-	-	-	-
1999.....	5	-	-	-	1	-	2	1	-	-	-	-

American Indian 1/, female

2004.....	4	-	-	-	-	2	1	-	-	-	-	-
2003.....	4	-	-	-	-	-	-	1	1	1	-	-
2002.....	4	-	-	-	-	1	-	-	-	-	-	1
2001.....	3	-	-	-	-	-	1	-	-	-	-	-
2000.....	2	-	-	-	-	-	1	-	-	1	-	-
1999.....	3	-	-	-	-	-	1	1	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Malignant neoplasms of bone and articular cartilage (C40-C41)												
Asian or Pacific Islander 2/, both sexes												
2004.....	23	-	-	-	1	2	2	1	3	4	1	2
2003.....	29	-	-	-	2	5	5	2	1	1	-	3
2002.....	22	-	-	1	1	2	4	3	1	-	2	2
2001.....	26	-	-	-	3	1	-	1	2	2	2	2
2000.....	20	-	-	1	1	3	1	1	1	1	-	1
1999.....	27	-	-	2	3	6	3	2	2	1	1	1
Asian or Pacific Islander 2/, male												
2004.....	14	-	-	-	1	1	1	-	1	4	-	1
2003.....	16	-	-	-	2	3	2	1	1	1	-	2
2002.....	13	-	-	1	1	1	3	2	1	-	1	1
2001.....	16	-	-	-	1	1	-	1	1	1	2	2
2000.....	12	-	-	-	-	2	1	1	1	1	-	1
1999.....	17	-	-	-	1	5	3	2	1	1	1	1
Asian or Pacific Islander 2/, female												
2004.....	9	-	-	-	-	1	1	1	2	-	1	1
2003.....	13	-	-	-	-	2	3	1	-	-	-	1
2002.....	9	-	-	-	-	1	1	1	-	-	1	1
2001.....	10	-	-	-	2	-	-	-	1	1	-	-
2000.....	8	-	-	1	1	1	-	-	-	-	-	-
1999.....	10	-	-	2	2	1	-	-	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Melanoma and other malignant neoplasms of skin (C43-C44)												
All races, both sexes												
2004.....	10,301	-	2	-	-	11	30	71	128	207	360	577
2003.....	10,215	-	1	1	1	10	30	68	136	202	415	616
2002.....	9,905	1	3	-	1	7	34	72	130	236	421	559
2001.....	9,992	-	-	-	-	11	34	77	118	227	442	605
2000.....	9,672	-	-	1	1	7	35	63	128	248	423	597
1999.....	9,530	1	-	1	-	10	30	62	138	264	423	604
All races, male												
2004.....	6,770	-	2	-	-	3	15	42	77	120	225	370
2003.....	6,612	-	-	-	-	7	19	44	84	127	254	389
2002.....	6,371	-	2	-	1	3	17	39	85	132	269	395
2001.....	6,421	-	-	-	-	9	17	45	66	140	273	387
2000.....	6,146	-	-	1	1	4	18	36	77	127	263	368
1999.....	6,119	1	-	-	-	8	16	38	78	168	280	406
All races, female												
2004.....	3,531	-	-	-	-	8	15	29	51	87	135	207
2003.....	3,603	-	1	1	1	3	11	24	52	75	161	227
2002.....	3,534	1	1	-	-	4	17	33	45	104	152	164
2001.....	3,571	-	-	-	-	2	17	32	52	87	169	218
2000.....	3,526	-	-	-	-	3	17	27	51	121	160	229
1999.....	3,411	-	-	1	-	2	14	24	60	96	143	198
White, both sexes												
2004.....	9,989	-	1	-	-	9	28	69	126	202	347	561
2003.....	9,890	-	1	1	1	10	28	66	130	194	400	587
2002.....	9,570	1	3	-	1	6	34	70	129	223	410	540
2001.....	9,671	-	-	-	-	10	33	74	116	224	432	585
2000.....	9,316	-	-	1	1	6	35	59	125	239	401	575
1999.....	9,230	1	-	1	-	8	28	61	131	257	412	575

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Melanoma and other malignant neoplasms of skin (C43-C44)												
White, male												
2004.....	6,578	-	1	-	-	3	14	42	75	118	213	356
2003.....	6,429	-	-	-	-	7	19	44	80	121	245	372
2002.....	6,159	-	2	-	1	2	17	37	85	126	261	383
2001.....	6,222	-	-	-	-	8	17	42	65	138	265	371
2000.....	5,945	-	-	1	1	4	18	34	76	122	249	357
1999.....	5,921	1	-	-	-	6	14	37	73	164	272	385
White, female												
2004.....	3,411	-	-	-	-	6	14	27	51	84	134	205
2003.....	3,461	-	1	1	1	3	9	22	50	73	155	215
2002.....	3,411	1	1	-	-	4	17	33	44	97	149	157
2001.....	3,449	-	-	-	-	2	16	32	51	86	167	214
2000.....	3,371	-	-	-	-	2	17	25	49	117	152	218
1999.....	3,309	-	-	1	-	2	14	24	58	93	140	190
Black, both sexes												
2004.....	255	-	1	-	-	2	2	2	1	4	11	13
2003.....	248	-	-	-	-	-	1	1	2	7	14	24
2002.....	261	-	-	-	-	1	-	1	1	10	8	12
2001.....	257	-	-	-	-	1	-	1	1	3	10	15
2000.....	287	-	-	-	-	1	-	2	1	6	17	18
1999.....	243	-	-	-	-	2	-	1	5	4	9	24
Black, male												
2004.....	160	-	1	-	-	-	1	-	1	2	10	11
2003.....	138	-	-	-	-	-	-	-	2	6	8	15
2002.....	170	-	-	-	-	1	-	1	-	5	6	8
2001.....	154	-	-	-	-	1	-	1	1	2	8	14
2000.....	160	-	-	-	-	-	-	1	1	3	12	9
1999.....	164	-	-	-	-	2	-	1	3	2	6	19

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Melanoma and other malignant neoplasms of skin (C43-C44)												
Asian or Pacific Islander 2/, both sexes												
2004.....	39	-	-	-	-	-	-	-	1	1	-	3
2003.....	54	-	-	-	-	-	-	1	4	1	-	5
2002.....	52	-	-	-	-	-	-	1	-	3	3	6
2001.....	49	-	-	-	-	-	1	1	1	-	-	5
2000.....	57	-	-	-	-	-	-	2	2	3	5	3
1999.....	41	-	-	-	-	-	1	-	2	3	2	2
Asian or Pacific Islander 2/, male												
2004.....	19	-	-	-	-	-	-	-	1	-	-	3
2003.....	34	-	-	-	-	-	-	-	2	-	-	2
2002.....	29	-	-	-	-	-	-	1	-	1	2	3
2001.....	32	-	-	-	-	-	-	1	-	-	-	2
2000.....	32	-	-	-	-	-	-	1	-	2	2	2
1999.....	24	-	-	-	-	-	1	-	2	2	2	-
Asian or Pacific Islander 2/, female												
2004.....	20	-	-	-	-	-	-	-	-	1	-	-
2003.....	20	-	-	-	-	-	-	1	2	1	-	3
2002.....	23	-	-	-	-	-	-	-	-	2	1	3
2001.....	17	-	-	-	-	-	1	-	1	-	-	3
2000.....	25	-	-	-	-	-	-	1	2	1	3	1
1999.....	17	-	-	-	-	-	-	-	-	1	-	2

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Malignant melanoma of skin (C43)

All races, both sexes

2004.....	7,952	-	2	-	-	10	29	68	126	194	332	512
2003.....	7,818	-	1	1	1	9	29	63	128	189	372	541
2002.....	7,514	-	3	-	1	7	34	70	124	219	385	495
2001.....	7,542	-	-	-	-	11	30	72	114	213	409	536
2000.....	7,420	-	-	-	1	7	34	58	128	234	392	534
1999.....	7,215	1	-	1	-	9	30	62	132	239	386	518

All races, male

2004.....	5,125	-	2	-	-	3	14	41	76	109	203	321
2003.....	4,963	-	-	-	-	6	18	40	79	114	222	328
2002.....	4,725	-	2	-	1	3	17	38	81	120	243	343
2001.....	4,763	-	-	-	-	9	15	40	63	130	250	331
2000.....	4,592	-	-	-	1	4	17	33	77	118	237	321
1999.....	4,529	1	-	-	-	7	16	38	72	151	246	337

All races, female

2004.....	2,827	-	-	-	-	7	15	27	50	85	129	191
2003.....	2,855	-	1	1	1	3	11	23	49	75	150	213
2002.....	2,789	-	1	-	-	4	17	32	43	99	142	152
2001.....	2,779	-	-	-	-	2	15	32	51	83	159	205
2000.....	2,828	-	-	-	-	3	17	25	51	116	155	213
1999.....	2,686	-	-	1	-	2	14	24	60	88	140	181

White, both sexes

2004.....	7,801	-	1	-	-	9	27	67	124	191	325	507
2003.....	7,648	-	1	1	1	9	27	61	125	186	363	528
2002.....	7,353	-	3	-	1	6	34	68	124	210	380	485
2001.....	7,403	-	-	-	-	10	30	70	114	211	404	529
2000.....	7,245	-	-	-	1	6	34	55	125	229	377	525
1999.....	7,086	1	-	1	-	8	28	61	127	236	382	509

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Malignant melanoma of skin (C43)												
White, male												
2004.....	5,053	-	1	-	-	3	13	41	74	109	197	317
2003.....	4,887	-	-	-	-	6	18	40	77	113	219	322
2002.....	4,639	-	2	-	1	2	17	36	81	118	239	337
2001.....	4,702	-	-	-	-	8	15	38	63	129	247	328
2000.....	4,514	-	-	-	1	4	17	32	76	116	229	319
1999.....	4,459	1	-	-	-	6	14	37	69	149	245	331
White, female												
2004.....	2,748	-	-	-	-	6	14	26	50	82	128	190
2003.....	2,761	-	1	1	1	3	9	21	48	73	144	206
2002.....	2,714	-	1	-	-	4	17	32	43	92	141	148
2001.....	2,701	-	-	-	-	2	15	32	51	82	157	201
2000.....	2,731	-	-	-	-	2	17	23	49	113	148	206
1999.....	2,627	-	-	1	-	2	14	24	58	87	137	178
Black, both sexes												
2004.....	119	-	1	-	-	1	2	1	1	2	6	3
2003.....	121	-	-	-	-	-	1	1	2	2	9	8
2002.....	102	-	-	-	-	1	-	1	-	7	2	3
2001.....	104	-	-	-	-	1	-	1	-	2	5	3
2000.....	135	-	-	-	-	1	-	2	1	4	11	6
1999.....	96	-	-	-	-	1	-	1	3	1	3	6
Black, male												
2004.....	57	-	1	-	-	-	1	-	1	-	5	2
2003.....	50	-	-	-	-	-	-	-	2	1	3	4
2002.....	52	-	-	-	-	1	-	1	-	2	2	2
2001.....	37	-	-	-	-	1	-	1	-	1	3	2
2000.....	60	-	-	-	-	-	-	1	1	2	7	1
1999.....	48	-	-	-	-	1	-	1	1	1	-	4

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant melanoma of skin (C43)

Asian or Pacific Islander 2/, both sexes

2004.....	21	-	-	-	-	-	-	-	1	1	-	2
2003.....	35	-	-	-	-	-	-	1	1	1	-	5
2002.....	42	-	-	-	-	-	-	1	-	2	3	6
2001.....	25	-	-	-	-	-	-	-	-	-	-	4
2000.....	32	-	-	-	-	-	-	1	2	1	4	2
1999.....	23	-	-	-	-	-	1	-	2	2	1	1

Asian or Pacific Islander 2/, male

2004.....	8	-	-	-	-	-	-	-	1	-	-	2
2003.....	21	-	-	-	-	-	-	-	-	-	-	2
2002.....	25	-	-	-	-	-	-	1	-	-	2	3
2001.....	15	-	-	-	-	-	-	-	-	-	-	1
2000.....	13	-	-	-	-	-	-	-	-	-	1	1
1999.....	15	-	-	-	-	-	1	-	2	1	1	-

Asian or Pacific Islander 2/, female

2004.....	13	-	-	-	-	-	-	-	-	1	-	-
2003.....	14	-	-	-	-	-	-	1	1	1	-	3
2002.....	17	-	-	-	-	-	-	-	-	2	1	3
2001.....	10	-	-	-	-	-	-	-	-	-	-	3
2000.....	19	-	-	-	-	-	-	1	2	1	3	1
1999.....	8	-	-	-	-	-	-	-	-	1	-	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other malignant neoplasms of skin (C44)												
All races, both sexes												
2004.....	2,349	-	-	-	-	1	1	3	2	13	28	65
2003.....	2,397	-	-	-	-	1	1	5	8	13	43	75
2002.....	2,391	1	-	-	-	-	-	2	6	17	36	64
2001.....	2,450	-	-	-	-	-	4	5	4	14	33	69
2000.....	2,252	-	-	1	-	-	1	5	-	14	31	63
1999.....	2,315	-	-	-	-	1	-	-	6	25	37	86
All races, male												
2004.....	1,645	-	-	-	-	-	1	1	1	11	22	49
2003.....	1,649	-	-	-	-	1	1	4	5	13	32	61
2002.....	1,646	-	-	-	-	-	-	1	4	12	26	52
2001.....	1,658	-	-	-	-	-	2	5	3	10	23	56
2000.....	1,554	-	-	1	-	-	1	3	-	9	26	47
1999.....	1,590	-	-	-	-	1	-	-	6	17	34	69
All races, female												
2004.....	704	-	-	-	-	1	-	2	1	2	6	16
2003.....	748	-	-	-	-	-	-	1	3	-	11	14
2002.....	745	1	-	-	-	-	-	1	2	5	10	12
2001.....	792	-	-	-	-	-	2	-	1	4	10	13
2000.....	698	-	-	-	-	-	-	2	-	5	5	16
1999.....	725	-	-	-	-	-	-	-	-	8	3	17
White, both sexes												
2004.....	2,188	-	-	-	-	-	1	2	2	11	22	54
2003.....	2,242	-	-	-	-	1	1	5	5	8	37	59
2002.....	2,217	1	-	-	-	-	-	2	5	13	30	55
2001.....	2,268	-	-	-	-	-	3	4	2	13	28	56
2000.....	2,071	-	-	1	-	-	1	4	-	10	24	50
1999.....	2,144	-	-	-	-	-	-	-	4	21	30	66

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other malignant neoplasms of skin (C44)

White, male

2004.....	1,525	-	-	-	-	-	1	1	1	9	16	39
2003.....	1,542	-	-	-	-	1	1	4	3	8	26	50
2002.....	1,520	-	-	-	-	-	-	1	4	8	22	46
2001.....	1,520	-	-	-	-	-	2	4	2	9	18	43
2000.....	1,431	-	-	1	-	-	1	2	-	6	20	38
1999.....	1,462	-	-	-	-	-	-	-	4	15	27	54

White, female

2004.....	663	-	-	-	-	-	-	1	1	2	6	15
2003.....	700	-	-	-	-	-	-	1	2	-	11	9
2002.....	697	1	-	-	-	-	-	1	1	5	8	9
2001.....	748	-	-	-	-	-	1	-	-	4	10	13
2000.....	640	-	-	-	-	-	-	2	-	4	4	12
1999.....	682	-	-	-	-	-	-	-	-	6	3	12

Black, both sexes

2004.....	136	-	-	-	-	1	-	1	-	2	5	10
2003.....	127	-	-	-	-	-	-	-	-	5	5	16
2002.....	159	-	-	-	-	-	-	-	1	3	6	9
2001.....	153	-	-	-	-	-	-	-	1	1	5	12
2000.....	152	-	-	-	-	-	-	-	-	2	6	12
1999.....	147	-	-	-	-	1	-	-	2	3	6	18

Black, male

2004.....	103	-	-	-	-	-	-	-	-	2	5	9
2003.....	88	-	-	-	-	-	-	-	-	5	5	11
2002.....	118	-	-	-	-	-	-	-	-	3	4	6
2001.....	117	-	-	-	-	-	-	-	1	1	5	12
2000.....	100	-	-	-	-	-	-	-	-	1	5	8
1999.....	116	-	-	-	-	1	-	-	2	1	6	15

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Malignant neoplasms of mesothelial and soft tissue (C45-C49)												
All races, both sexes												
2004.....	7,104	5	24	29	43	64	76	71	106	137	215	285
2003.....	6,911	4	13	26	37	71	79	77	109	132	190	287
2002.....	6,808	3	23	28	42	60	73	84	91	145	217	256
2001.....	6,745	6	19	26	35	66	67	79	117	134	211	281
2000.....	6,817	7	21	17	27	73	72	88	122	150	205	286
1999.....	6,649	4	18	27	37	65	72	86	108	168	218	281
All races, male												
2004.....	4,087	3	10	16	22	33	43	48	62	87	101	143
2003.....	3,986	3	11	18	14	42	46	44	67	73	100	159
2002.....	3,955	-	13	16	21	32	40	50	57	89	122	139
2001.....	3,897	5	15	13	18	38	45	44	75	70	108	151
2000.....	3,947	5	11	10	11	37	46	52	64	87	100	151
1999.....	3,898	1	10	14	20	46	37	49	68	97	139	136
All races, female												
2004.....	3,017	2	14	13	21	31	33	23	44	50	114	142
2003.....	2,925	1	2	8	23	29	33	33	42	59	90	128
2002.....	2,853	3	10	12	21	28	33	34	34	56	95	117
2001.....	2,848	1	4	13	17	28	22	35	42	64	103	130
2000.....	2,870	2	10	7	16	36	26	36	58	63	105	135
1999.....	2,751	3	8	13	17	19	35	37	40	71	79	145
White, both sexes												
2004.....	6,374	2	17	25	32	45	52	59	80	106	168	234
2003.....	6,195	3	8	23	29	60	67	57	79	112	146	226
2002.....	6,107	3	17	21	33	44	63	63	68	119	171	210
2001.....	6,075	5	14	19	28	53	53	61	91	117	163	223
2000.....	6,123	6	16	16	23	56	51	73	99	116	161	235
1999.....	5,929	3	13	20	29	50	52	66	78	131	164	239

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Malignant neoplasms of mesothelial and soft tissue (C45-C49)

White, male

2004.....	3,736	1	8	13	19	22	31	43	48	68	82	118
2003.....	3,617	2	7	17	11	34	39	37	50	63	75	121
2002.....	3,606	-	9	12	15	22	37	38	39	75	96	112
2001.....	3,561	5	11	10	14	27	32	33	58	61	85	121
2000.....	3,602	4	9	9	8	32	31	44	53	66	79	129
1999.....	3,538	1	7	12	17	35	23	39	47	77	111	118

White, female

2004.....	2,638	1	9	12	13	23	21	16	32	38	86	116
2003.....	2,578	1	1	6	18	26	28	20	29	49	71	105
2002.....	2,501	3	8	9	18	22	26	25	29	44	75	98
2001.....	2,514	-	3	9	14	26	21	28	33	56	78	102
2000.....	2,521	2	7	7	15	24	20	29	46	50	82	106
1999.....	2,391	2	6	8	12	15	29	27	31	54	53	121

Black, both sexes

2004.....	592	2	6	4	9	16	18	10	21	27	36	40
2003.....	594	1	4	3	8	11	11	16	28	18	35	54
2002.....	575	-	5	6	6	15	9	18	16	23	39	43
2001.....	532	1	4	6	5	12	12	15	19	13	36	42
2000.....	582	1	3	1	2	14	17	13	17	31	39	44
1999.....	587	1	3	6	8	14	18	18	20	33	42	34

Black, male

2004.....	282	1	2	3	3	8	8	4	12	16	14	19
2003.....	305	1	3	1	3	8	6	5	16	9	20	35
2002.....	289	-	3	4	4	9	3	9	12	13	23	27
2001.....	265	-	4	2	2	10	11	9	12	9	20	18
2000.....	287	1	2	1	1	4	13	8	8	19	19	20
1999.....	296	-	2	2	3	10	13	9	13	19	25	14

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Malignant neoplasms of mesothelial and soft tissue (C45-C49)

Black, female

2004.....	310	1	4	1	6	8	10	6	9	11	22	21
2003.....	289	-	1	2	5	3	5	11	12	9	15	19
2002.....	286	-	2	2	2	6	6	9	4	10	16	16
2001.....	267	1	-	4	3	2	1	6	7	4	16	24
2000.....	295	-	1	-	1	10	4	5	9	12	20	24
1999.....	291	1	1	4	5	4	5	9	7	14	17	20

American Indian 1/, both sexes

2004.....	21	-	-	-	-	1	1	1	-	1	1	3
2003.....	28	-	-	-	-	-	-	1	-	-	3	2
2002.....	26	-	-	-	1	1	-	1	-	2	3	1
2001.....	25	-	1	-	-	-	1	2	2	-	2	3
2000.....	13	-	-	-	-	1	1	1	-	-	-	2
1999.....	27	-	-	-	-	-	-	1	2	-	4	2

American Indian 1/, male

2004.....	9	-	-	-	-	1	-	-	-	-	1	2
2003.....	12	-	-	-	-	-	-	1	-	-	1	1
2002.....	15	-	-	-	1	1	-	1	-	-	1	-
2001.....	12	-	-	-	-	-	1	1	1	-	-	2
2000.....	8	-	-	-	-	1	-	-	-	-	-	1
1999.....	10	-	-	-	-	-	-	-	2	-	1	1

American Indian 1/, female

2004.....	12	-	-	-	-	-	1	1	-	1	-	1
2003.....	16	-	-	-	-	-	-	-	-	-	2	1
2002.....	11	-	-	-	-	-	-	-	-	2	2	1
2001.....	13	-	1	-	-	-	-	1	1	-	2	1
2000.....	5	-	-	-	-	-	1	1	-	-	-	1
1999.....	17	-	-	-	-	-	-	1	-	-	3	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasms of mesothelial and soft tissue (C45-C49)

Asian or Pacific Islander 2/, both sexes

2004.....	117	1	1	-	2	2	5	1	5	3	10	8
2003.....	94	-	1	-	-	-	1	3	2	2	6	5
2002.....	100	-	1	1	2	-	1	2	7	1	4	2
2001.....	113	-	-	1	2	1	1	1	5	4	10	13
2000.....	99	-	2	-	2	2	3	1	6	3	5	5
1999.....	106	-	2	1	-	1	2	1	8	4	8	6

Asian or Pacific Islander 2/, male

2004.....	60	1	-	-	-	2	4	1	2	3	4	4
2003.....	52	-	1	-	-	-	1	1	1	1	4	2
2002.....	45	-	1	-	1	-	-	2	6	1	2	-
2001.....	59	-	-	1	2	1	1	1	4	-	3	10
2000.....	50	-	-	-	2	-	2	-	3	2	2	1
1999.....	54	-	1	-	-	1	1	1	6	1	2	3

Asian or Pacific Islander 2/, female

2004.....	57	-	1	-	2	-	1	-	3	-	6	4
2003.....	42	-	-	-	-	-	-	2	1	1	2	3
2002.....	55	-	-	1	1	-	1	-	1	-	2	2
2001.....	54	-	-	-	-	-	-	-	1	4	7	3
2000.....	49	-	2	-	-	2	1	1	3	1	3	4
1999.....	52	-	1	1	-	-	1	-	2	3	6	3

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Mesothelioma (C45)

All races, both sexes

2004.....	2,504	-	-	-	-	-	1	4	5	9	31	39
2003.....	2,476	-	-	-	-	3	-	-	7	13	23	53
2002.....	2,431	-	-	-	1	2	1	5	13	24	29	
2001.....	2,371	-	-	-	-	1	3	1	6	13	20	47
2000.....	2,384	-	-	-	-	-	1	1	4	12	21	40
1999.....	2,343	-	-	-	-	-	1	1	3	16	15	48

All races, male

2004.....	2,010	-	-	-	-	-	1	2	2	6	16	24
2003.....	2,007	-	-	-	-	2	-	-	4	8	13	40
2002.....	2,005	-	-	-	-	-	-	4	4	11	18	19
2001.....	1,907	-	-	-	-	-	2	1	5	7	9	34
2000.....	1,925	-	-	-	-	-	-	1	-	9	10	22
1999.....	1,881	-	-	-	-	-	-	-	3	10	10	35

All races, female

2004.....	494	-	-	-	-	-	-	2	3	3	15	15
2003.....	469	-	-	-	-	1	-	-	3	5	10	13
2002.....	426	-	-	-	1	2	1	1	1	2	6	10
2001.....	464	-	-	-	-	1	1	-	1	6	11	13
2000.....	459	-	-	-	-	-	1	-	4	3	11	18
1999.....	462	-	-	-	-	-	1	1	-	6	5	13

White, both sexes

2004.....	2,392	-	-	-	-	-	1	4	4	8	26	36
2003.....	2,345	-	-	-	-	2	-	-	5	11	21	44
2002.....	2,313	-	-	-	-	1	1	3	5	13	21	27
2001.....	2,281	-	-	-	-	1	2	1	4	11	19	40
2000.....	2,258	-	-	-	-	-	1	1	3	11	17	33
1999.....	2,222	-	-	-	-	-	1	1	3	15	13	43

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Kaposi's sarcoma (C46)												
All races, both sexes												
2004.....	48	-	-	-	1	-	-	-	3	3	3	5
2003.....	55	1	-	-	-	-	-	5	5	-	2	3
2002.....	54	-	-	-	-	-	-	-	3	5	5	2
2001.....	40	-	-	-	-	-	-	1	3	2	-	4
2000.....	62	-	-	-	-	1	-	-	11	1	2	3
1999.....	42	-	-	-	-	-	-	1	6	3	4	-
All races, male												
2004.....	37	-	-	-	1	-	-	-	3	3	3	5
2003.....	34	1	-	-	-	-	-	4	5	-	2	3
2002.....	31	-	-	-	-	-	-	-	3	5	3	2
2001.....	29	-	-	-	-	-	-	1	3	2	-	4
2000.....	43	-	-	-	-	-	-	-	11	1	2	3
1999.....	32	-	-	-	-	-	-	1	6	3	4	-
All races, female												
2004.....	11	-	-	-	-	-	-	-	-	-	-	-
2003.....	21	-	-	-	-	-	-	1	-	-	-	-
2002.....	23	-	-	-	-	-	-	-	-	-	2	-
2001.....	11	-	-	-	-	-	-	-	-	-	-	-
2000.....	19	-	-	-	-	1	-	-	-	-	-	-
1999.....	10	-	-	-	-	-	-	-	-	-	-	-
White, both sexes												
2004.....	38	-	-	-	1	-	-	-	1	1	1	3
2003.....	48	-	-	-	-	-	-	3	3	-	2	3
2002.....	43	-	-	-	-	-	-	-	-	1	4	2
2001.....	33	-	-	-	-	-	-	-	1	2	-	2
2000.....	42	-	-	-	-	1	-	-	4	-	-	1
1999.....	35	-	-	-	-	-	-	-	4	3	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Kaposi's sarcoma (C46)												
White, male												
2004.....	28	-	-	-	1	-	-	-	1	1	1	3
2003.....	28	-	-	-	-	-	-	3	3	-	2	3
2002.....	22	-	-	-	-	-	-	-	-	1	3	2
2001.....	22	-	-	-	-	-	-	-	1	2	-	2
2000.....	23	-	-	-	-	-	-	-	4	-	-	1
1999.....	25	-	-	-	-	-	-	-	4	3	1	-
White, female												
2004.....	10	-	-	-	-	-	-	-	-	-	-	-
2003.....	20	-	-	-	-	-	-	-	-	-	-	-
2002.....	21	-	-	-	-	-	-	-	-	-	1	-
2001.....	11	-	-	-	-	-	-	-	-	-	-	-
2000.....	19	-	-	-	-	1	-	-	-	-	-	-
1999.....	10	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes												
2004.....	10	-	-	-	-	-	-	-	2	2	2	2
2003.....	7	1	-	-	-	-	-	2	2	-	-	-
2002.....	9	-	-	-	-	-	-	-	3	4	1	-
2001.....	6	-	-	-	-	-	-	1	1	-	-	2
2000.....	19	-	-	-	-	-	-	-	6	1	2	2
1999.....	7	-	-	-	-	-	-	1	2	-	3	-
Black, male												
2004.....	9	-	-	-	-	-	-	-	2	2	2	2
2003.....	6	1	-	-	-	-	-	1	2	-	-	-
2002.....	8	-	-	-	-	-	-	-	3	4	-	-
2001.....	6	-	-	-	-	-	-	1	1	-	-	2
2000.....	19	-	-	-	-	-	-	-	6	1	2	2
1999.....	7	-	-	-	-	-	-	1	2	-	3	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Of other mesothelial and soft tissue (C47-C49)

All races, both sexes

2004.....	4,552	5	24	29	42	64	75	67	98	125	181	241
2003.....	4,380	3	13	26	37	68	79	72	97	119	165	231
2002.....	4,323	3	23	28	41	58	72	79	83	127	188	225
2001.....	4,334	6	19	26	35	65	64	77	108	119	191	230
2000.....	4,371	7	21	17	27	72	71	87	107	137	182	243
1999.....	4,264	4	18	27	37	65	71	84	99	149	199	233

All races, male

2004.....	2,040	3	10	16	21	33	42	46	57	78	82	114
2003.....	1,945	2	11	18	14	40	46	40	58	65	85	116
2002.....	1,919	-	13	16	21	32	40	46	50	73	101	118
2001.....	1,961	5	15	13	18	38	43	42	67	61	99	113
2000.....	1,979	5	11	10	11	37	46	51	53	77	88	126
1999.....	1,985	1	10	14	20	46	37	48	59	84	125	101

All races, female

2004.....	2,512	2	14	13	21	31	33	21	41	47	99	127
2003.....	2,435	1	2	8	23	28	33	32	39	54	80	115
2002.....	2,404	3	10	12	20	26	32	33	33	54	87	107
2001.....	2,373	1	4	13	17	27	21	35	41	58	92	117
2000.....	2,392	2	10	7	16	35	25	36	54	60	94	117
1999.....	2,279	3	8	13	17	19	34	36	40	65	74	132

White, both sexes

2004.....	3,944	2	17	25	31	45	51	55	75	97	141	195
2003.....	3,802	3	8	23	29	58	67	54	71	101	123	179
2002.....	3,751	3	17	21	33	43	62	60	63	105	146	181
2001.....	3,761	5	14	19	28	52	51	60	86	104	144	181
2000.....	3,823	6	16	16	23	55	50	72	92	105	144	201
1999.....	3,672	3	13	20	29	50	51	65	71	113	150	196

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Of other mesothelial and soft tissue (C47-C49)

White, male

2004.....	1,787	1	8	13	18	22	30	41	45	62	67	93
2003.....	1,693	2	7	17	11	33	39	34	44	57	61	87
2002.....	1,671	-	9	12	15	22	37	35	35	63	78	93
2001.....	1,704	5	11	10	14	27	31	32	54	53	77	90
2000.....	1,753	4	9	9	8	32	31	43	49	58	70	111
1999.....	1,718	1	7	12	17	35	23	39	40	64	102	85

White, female

2004.....	2,157	1	9	12	13	23	21	14	30	35	74	102
2003.....	2,109	1	1	6	18	25	28	20	27	44	62	92
2002.....	2,080	3	8	9	18	21	25	25	28	42	68	88
2001.....	2,057	-	3	9	14	25	20	28	32	51	67	91
2000.....	2,070	2	7	7	15	23	19	29	43	47	74	90
1999.....	1,954	2	6	8	12	15	28	26	31	49	48	111

Black, both sexes

2004.....	493	2	6	4	9	16	18	10	18	24	31	36
2003.....	485	-	4	3	8	10	11	14	25	17	34	46
2002.....	471	-	5	6	5	14	9	16	13	19	35	41
2001.....	461	1	4	6	5	12	12	14	17	12	35	35
2000.....	459	1	3	1	2	14	17	13	10	29	35	37
1999.....	482	1	3	6	8	14	18	17	18	32	37	30

Black, male

2004.....	200	1	2	3	3	8	8	4	10	13	11	16
2003.....	210	-	3	1	3	7	6	4	13	8	19	27
2002.....	205	-	3	4	4	9	3	8	9	9	20	25
2001.....	207	-	4	2	2	10	11	8	10	8	19	13
2000.....	186	1	2	1	1	4	13	8	2	17	16	15
1999.....	218	-	2	2	3	10	13	8	11	19	20	13

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Of other mesothelial and soft tissue (C47-C49)												
Black, female												
2004.....	293	1	4	1	6	8	10	6	8	11	20	20
2003.....	275	-	1	2	5	3	5	10	12	9	15	19
2002.....	266	-	2	2	1	5	6	8	4	10	15	16
2001.....	254	1	-	4	3	2	1	6	7	4	16	22
2000.....	273	-	1	-	1	10	4	5	8	12	19	22
1999.....	264	1	1	4	5	4	5	9	7	13	17	17
American Indian 1/, both sexes												
2004.....	16	-	-	-	-	1	1	1	-	1	-	3
2003.....	24	-	-	-	-	-	-	1	-	-	3	2
2002.....	18	-	-	-	1	1	-	1	-	2	3	1
2001.....	21	-	1	-	-	-	1	2	1	-	2	3
2000.....	8	-	-	-	-	1	1	1	-	-	-	1
1999.....	19	-	-	-	-	-	-	1	2	-	4	1
American Indian 1/, male												
2004.....	6	-	-	-	-	1	-	-	-	-	-	2
2003.....	10	-	-	-	-	-	-	1	-	-	1	1
2002.....	9	-	-	-	1	1	-	1	-	-	1	-
2001.....	8	-	-	-	-	-	1	1	-	-	-	2
2000.....	3	-	-	-	-	1	-	-	-	-	-	-
1999.....	6	-	-	-	-	-	-	-	2	-	1	-
American Indian 1/, female												
2004.....	10	-	-	-	-	-	1	1	-	1	-	1
2003.....	14	-	-	-	-	-	-	-	-	-	2	1
2002.....	9	-	-	-	-	-	-	-	-	2	2	1
2001.....	13	-	1	-	-	-	-	1	1	-	2	1
2000.....	5	-	-	-	-	-	1	1	-	-	-	1
1999.....	13	-	-	-	-	-	-	1	-	-	3	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Of other mesothelial and soft tissue (C47-C49)												
Asian or Pacific Islander 2/, both sexes												
2004.....	99	1	1	-	2	2	5	1	5	3	9	7
2003.....	69	-	1	-	-	-	1	3	1	1	5	4
2002.....	83	-	1	1	2	-	1	2	7	1	4	2
2001.....	91	-	-	1	2	1	-	1	4	3	10	11
2000.....	81	-	2	-	2	2	3	1	5	3	3	4
1999.....	91	-	2	1	-	1	2	1	8	4	8	6
Asian or Pacific Islander 2/, male												
2004.....	47	1	-	-	-	2	4	1	2	3	4	3
2003.....	32	-	1	-	-	-	1	1	1	-	4	1
2002.....	34	-	1	-	1	-	-	2	6	1	2	-
2001.....	42	-	-	1	2	1	-	1	3	-	3	8
2000.....	37	-	-	-	2	-	2	-	2	2	2	-
1999.....	43	-	1	-	-	1	1	1	6	1	2	3
Asian or Pacific Islander 2/, female												
2004.....	52	-	1	-	2	-	1	-	3	-	5	4
2003.....	37	-	-	-	-	-	-	2	-	1	1	3
2002.....	49	-	-	1	1	-	1	-	1	-	2	2
2001.....	49	-	-	-	-	-	-	-	1	3	7	3
2000.....	44	-	2	-	-	2	1	1	3	1	1	4
1999.....	48	-	1	1	-	-	1	-	2	3	6	3

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Malignant neoplasm of breast (C50)												
All races, both sexes												
2004.....	41,316	-	-	-	-	-	11	76	320	823	1,688	2,593
2003.....	42,000	-	-	-	-	1	15	55	352	912	1,804	2,734
2002.....	41,883	-	-	-	-	1	9	63	362	904	1,818	2,747
2001.....	41,809	-	-	-	-	1	8	88	375	960	1,863	2,790
2000.....	42,300	-	-	-	-	1	10	83	384	982	1,859	2,730
1999.....	41,528	-	1	-	1	3	10	86	353	986	1,755	2,619
All races, male												
2004.....	362	-	-	-	-	-	1	1	-	3	7	13
2003.....	380	-	-	-	-	-	-	1	1	2	7	14
2002.....	369	-	-	-	-	-	-	2	1	4	8	16
2001.....	415	-	-	-	-	-	-	-	2	5	9	13
2000.....	428	-	-	-	-	-	-	2	1	12	7	18
1999.....	384	-	-	-	-	-	-	-	3	6	9	11
All races, female												
2004.....	40,954	-	-	-	-	-	10	75	320	820	1,681	2,580
2003.....	41,620	-	-	-	-	1	15	54	351	910	1,797	2,720
2002.....	41,514	-	-	-	-	1	9	61	361	900	1,810	2,731
2001.....	41,394	-	-	-	-	1	8	88	373	955	1,854	2,777
2000.....	41,872	-	-	-	-	1	10	81	383	970	1,852	2,712
1999.....	41,144	-	1	-	1	3	10	86	350	980	1,746	2,608
White, both sexes												
2004.....	34,759	-	-	-	-	-	7	47	216	569	1,212	1,920
2003.....	35,358	-	-	-	-	1	10	27	242	652	1,329	2,052
2002.....	35,382	-	-	-	-	-	6	40	245	634	1,316	2,059
2001.....	35,422	-	-	-	-	-	4	58	248	697	1,372	2,135
2000.....	36,127	-	-	-	-	1	4	60	265	717	1,380	2,066
1999.....	35,385	-	-	-	1	3	5	59	245	700	1,301	1,977

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Malignant neoplasm of breast (C50)												
White, male												
2004.....	301	-	-	-	-	-	-	-	-	1	5	11
2003.....	326	-	-	-	-	-	-	-	1	1	3	9
2002.....	300	-	-	-	-	-	-	2	-	2	7	11
2001.....	344	-	-	-	-	-	-	-	1	4	4	10
2000.....	360	-	-	-	-	-	-	-	1	8	3	13
1999.....	312	-	-	-	-	-	-	-	2	-	8	8
White, female												
2004.....	34,458	-	-	-	-	-	7	47	216	568	1,207	1,909
2003.....	35,032	-	-	-	-	1	10	27	241	651	1,326	2,043
2002.....	35,082	-	-	-	-	-	6	38	245	632	1,309	2,048
2001.....	35,078	-	-	-	-	-	4	58	247	693	1,368	2,125
2000.....	35,767	-	-	-	-	1	4	60	264	709	1,377	2,053
1999.....	35,073	-	-	-	1	3	5	59	243	700	1,293	1,969
Black, both sexes												
2004.....	5,597	-	-	-	-	-	2	26	90	225	402	572
2003.....	5,746	-	-	-	-	-	5	24	95	228	412	595
2002.....	5,636	-	-	-	-	1	2	22	103	234	445	585
2001.....	5,571	-	-	-	-	1	4	30	112	221	424	556
2000.....	5,424	-	-	-	-	-	5	20	103	228	417	564
1999.....	5,425	-	1	-	-	-	5	25	101	261	404	553
Black, male												
2004.....	56	-	-	-	-	-	-	1	-	2	1	2
2003.....	51	-	-	-	-	-	-	1	-	1	4	4
2002.....	64	-	-	-	-	-	-	-	1	2	1	5
2001.....	65	-	-	-	-	-	-	-	1	-	5	3
2000.....	63	-	-	-	-	-	-	2	-	4	4	4
1999.....	67	-	-	-	-	-	-	-	1	6	1	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Malignant neoplasm of breast (C50)												
Black, female												
2004.....	5,541	-	-	-	-	-	2	25	90	223	401	570
2003.....	5,695	-	-	-	-	-	5	23	95	227	408	591
2002.....	5,572	-	-	-	-	1	2	22	102	232	444	580
2001.....	5,506	-	-	-	-	1	4	30	111	221	419	553
2000.....	5,361	-	-	-	-	-	5	18	103	224	413	560
1999.....	5,358	-	1	-	-	-	5	25	100	255	403	552
American Indian 1/, both sexes												
2004.....	169	-	-	-	-	-	-	-	3	5	10	17
2003.....	148	-	-	-	-	-	-	2	2	4	5	8
2002.....	148	-	-	-	-	-	-	1	4	10	7	14
2001.....	127	-	-	-	-	-	-	-	2	9	6	15
2000.....	131	-	-	-	-	-	-	1	1	3	13	11
1999.....	131	-	-	-	-	-	-	1	1	3	8	9
American Indian 1/, male												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	1	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	1	-	-	-	-	-	-	-	-	-	-	-
2000.....	1	-	-	-	-	-	-	-	-	-	-	-
1999.....	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, female												
2004.....	169	-	-	-	-	-	-	-	3	5	10	17
2003.....	147	-	-	-	-	-	-	2	2	4	5	8
2002.....	148	-	-	-	-	-	-	1	4	10	7	14
2001.....	126	-	-	-	-	-	-	-	2	9	6	15
2000.....	130	-	-	-	-	-	-	1	1	3	13	11
1999.....	130	-	-	-	-	-	-	1	1	3	8	9

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasm of breast (C50)

Asian or Pacific Islander 2/, both sexes

2004.....	791	-	-	-	-	-	2	3	11	24	64	84
2003.....	748	-	-	-	-	-	-	2	13	28	58	79
2002.....	717	-	-	-	-	-	1	-	10	26	50	89
2001.....	689	-	-	-	-	-	-	-	13	33	61	84
2000.....	618	-	-	-	-	-	1	2	15	34	49	89
1999.....	587	-	-	-	-	-	-	1	6	22	42	80

Asian or Pacific Islander 2/, male

2004.....	5	-	-	-	-	-	1	-	-	-	1	-
2003.....	2	-	-	-	-	-	-	-	-	-	-	1
2002.....	5	-	-	-	-	-	-	-	-	-	-	-
2001.....	5	-	-	-	-	-	-	-	-	1	-	-
2000.....	4	-	-	-	-	-	-	-	-	-	-	1
1999.....	4	-	-	-	-	-	-	-	-	-	-	2

Asian or Pacific Islander 2/, female

2004.....	786	-	-	-	-	-	1	3	11	24	63	84
2003.....	746	-	-	-	-	-	-	2	13	28	58	78
2002.....	712	-	-	-	-	-	1	-	10	26	50	89
2001.....	684	-	-	-	-	-	-	-	13	32	61	84
2000.....	614	-	-	-	-	-	1	2	15	34	49	88
1999.....	583	-	-	-	-	-	-	1	6	22	42	78

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasms of female genital organs (C51-C58)

All races, both sexes

2004.....	27,049	-	-	-	2	12	44	110	227	426	872	1,361
2003.....	27,011	-	-	-	-	12	39	105	220	435	847	1,290
2002.....	27,091	1	1	1	1	13	46	83	259	491	847	1,360
2001.....	26,839	-	-	-	3	5	30	104	257	501	865	1,326
2000.....	26,412	-	-	2	1	10	35	119	249	501	854	1,300
1999.....	25,882	-	1	-	2	11	32	102	224	508	871	1,302

All races, male

2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

All races, female

2004.....	27,049	-	-	-	2	12	44	110	227	426	872	1,361
2003.....	27,011	-	-	-	-	12	39	105	220	435	847	1,290
2002.....	27,091	1	1	1	1	13	46	83	259	491	847	1,360
2001.....	26,839	-	-	-	3	5	30	104	257	501	865	1,326
2000.....	26,412	-	-	2	1	10	35	119	249	501	854	1,300
1999.....	25,882	-	1	-	2	11	32	102	224	508	871	1,302

White, both sexes

2004.....	23,065	-	-	-	2	10	29	75	175	333	682	1,094
2003.....	23,087	-	-	-	-	10	29	73	160	323	669	1,040
2002.....	23,177	1	1	1	1	9	36	65	199	383	673	1,090
2001.....	23,050	-	-	-	1	3	22	80	196	390	693	1,093
2000.....	22,684	-	-	1	-	8	25	101	185	391	658	1,068
1999.....	22,326	-	1	-	2	8	26	82	176	400	701	1,091

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Malignant neoplasms of female genital organs (C51-C58)

Black, female

2004.....	3,301	-	-	-	-	2	12	31	39	79	143	214
2003.....	3,244	-	-	-	-	2	8	28	45	90	132	204
2002.....	3,248	-	-	-	-	4	8	17	48	96	142	203
2001.....	3,152	-	-	-	2	1	6	23	47	87	149	173
2000.....	3,151	-	-	1	1	-	7	16	53	91	157	173
1999.....	3,052	-	-	-	-	3	5	19	37	85	141	148

American Indian 1/, both sexes

2004.....	116	-	-	-	-	-	-	-	4	3	12	8
2003.....	105	-	-	-	-	-	-	2	1	4	7	7
2002.....	122	-	-	-	-	-	2	1	3	4	7	19
2001.....	133	-	-	-	-	-	-	-	1	10	3	14
2000.....	79	-	-	-	-	-	1	-	3	3	6	6
1999.....	65	-	-	-	-	-	-	1	1	4	2	7

American Indian 1/, male

2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

American Indian 1/, female

2004.....	116	-	-	-	-	-	-	-	4	3	12	8
2003.....	105	-	-	-	-	-	-	2	1	4	7	7
2002.....	122	-	-	-	-	-	2	1	3	4	7	19
2001.....	133	-	-	-	-	-	-	-	1	10	3	14
2000.....	79	-	-	-	-	-	1	-	3	3	6	6
1999.....	65	-	-	-	-	-	-	1	1	4	2	7

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasms of female genital organs (C51-C58)

Asian or Pacific Islander 2/, both sexes

2004.....	567	-	-	-	-	-	3	4	9	11	35	45
2003.....	575	-	-	-	-	-	2	2	14	18	39	39
2002.....	544	-	-	-	-	-	-	-	9	8	25	48
2001.....	504	-	-	-	-	1	2	1	13	14	20	46
2000.....	498	-	-	-	-	2	2	2	8	16	33	53
1999.....	439	-	-	-	-	-	1	-	10	19	27	56

Asian or Pacific Islander 2/, male

2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

Asian or Pacific Islander 2/, female

2004.....	567	-	-	-	-	-	3	4	9	11	35	45
2003.....	575	-	-	-	-	-	2	2	14	18	39	39
2002.....	544	-	-	-	-	-	-	-	9	8	25	48
2001.....	504	-	-	-	-	1	2	1	13	14	20	46
2000.....	498	-	-	-	-	2	2	2	8	16	33	53
1999.....	439	-	-	-	-	-	1	-	10	19	27	56

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Of vulva and vagina (C51-C52)

All races, both sexes

2004.....	1,222	-	-	-	-	-	1	1	5	10	31	41
2003.....	1,166	-	-	-	-	-	1	4	4	8	17	38
2002.....	1,172	-	-	-	-	-	3	-	7	14	23	29
2001.....	1,147	-	-	-	-	-	-	2	2	8	27	34
2000.....	1,157	-	-	-	-	-	1	3	2	16	23	28
1999.....	1,165	-	-	-	-	-	-	1	4	12	20	26

All races, male

2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

All races, female

2004.....	1,222	-	-	-	-	-	1	1	5	10	31	41
2003.....	1,166	-	-	-	-	-	1	4	4	8	17	38
2002.....	1,172	-	-	-	-	-	3	-	7	14	23	29
2001.....	1,147	-	-	-	-	-	-	2	2	8	27	34
2000.....	1,157	-	-	-	-	-	1	3	2	16	23	28
1999.....	1,165	-	-	-	-	-	-	1	4	12	20	26

White, both sexes

2004.....	1,090	-	-	-	-	-	1	-	3	6	26	35
2003.....	1,051	-	-	-	-	-	-	2	2	5	15	33
2002.....	1,068	-	-	-	-	-	2	-	4	13	20	24
2001.....	1,005	-	-	-	-	-	-	2	1	8	19	25
2000.....	1,047	-	-	-	-	-	1	3	2	12	15	24
1999.....	1,046	-	-	-	-	-	-	1	3	9	17	22

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Of cervix uteri (C53)

All races, both sexes

2004.....	3,850	-	-	-	-	2	17	70	143	232	390	455
2003.....	3,919	-	-	-	-	1	15	59	142	250	383	440
2002.....	3,952	-	-	-	-	1	16	47	161	275	391	439
2001.....	4,092	-	-	-	1	-	12	51	158	291	420	475
2000.....	4,200	-	-	-	1	3	14	77	153	294	400	448
1999.....	4,205	-	-	-	-	2	10	62	155	292	407	462

All races, male

2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

All races, female

2004.....	3,850	-	-	-	-	2	17	70	143	232	390	455
2003.....	3,919	-	-	-	-	1	15	59	142	250	383	440
2002.....	3,952	-	-	-	-	1	16	47	161	275	391	439
2001.....	4,092	-	-	-	1	-	12	51	158	291	420	475
2000.....	4,200	-	-	-	1	3	14	77	153	294	400	448
1999.....	4,205	-	-	-	-	2	10	62	155	292	407	462

White, both sexes

2004.....	2,899	-	-	-	-	2	12	48	110	179	288	344
2003.....	2,953	-	-	-	-	1	12	40	99	182	296	334
2002.....	2,967	-	-	-	-	-	12	34	126	211	292	324
2001.....	3,136	-	-	-	-	-	11	40	122	226	325	371
2000.....	3,158	-	-	-	-	3	10	63	111	223	288	334
1999.....	3,229	-	-	-	-	-	9	52	124	221	314	371

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Of cervix uteri (C53)												
Black, female												
2004.....	789	-	-	-	-	-	4	20	25	44	84	95
2003.....	794	-	-	-	-	-	2	17	33	57	70	92
2002.....	829	-	-	-	-	1	2	13	30	60	86	95
2001.....	787	-	-	-	1	-	1	11	31	54	85	80
2000.....	869	-	-	-	1	-	4	12	39	62	88	88
1999.....	838	-	-	-	-	2	1	9	27	61	83	68
American Indian 1/, both sexes												
2004.....	31	-	-	-	-	-	-	-	4	3	7	3
2003.....	31	-	-	-	-	-	-	1	1	3	4	2
2002.....	34	-	-	-	-	-	2	-	2	-	3	9
2001.....	32	-	-	-	-	-	-	-	1	5	2	9
2000.....	30	-	-	-	-	-	-	-	2	2	5	3
1999.....	19	-	-	-	-	-	-	1	1	3	-	4
American Indian 1/, male												
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....
American Indian 1/, female												
2004.....	31	-	-	-	-	-	-	-	4	3	7	3
2003.....	31	-	-	-	-	-	-	1	1	3	4	2
2002.....	34	-	-	-	-	-	2	-	2	-	3	9
2001.....	32	-	-	-	-	-	-	-	1	5	2	9
2000.....	30	-	-	-	-	-	-	-	2	2	5	3
1999.....	19	-	-	-	-	-	-	1	1	3	-	4

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Of cervix uteri (C53)

Asian or Pacific Islander 2/, both sexes

2004.....	131	-	-	-	-	-	1	2	4	6	11	13
2003.....	141	-	-	-	-	-	1	1	9	8	13	12
2002.....	122	-	-	-	-	-	-	-	3	4	10	11
2001.....	137	-	-	-	-	-	-	-	4	6	8	15
2000.....	143	-	-	-	-	-	-	2	1	7	19	23
1999.....	119	-	-	-	-	-	-	-	3	7	10	19

Asian or Pacific Islander 2/, male

2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

Asian or Pacific Islander 2/, female

2004.....	131	-	-	-	-	-	1	2	4	6	11	13
2003.....	141	-	-	-	-	-	1	1	9	8	13	12
2002.....	122	-	-	-	-	-	-	-	3	4	10	11
2001.....	137	-	-	-	-	-	-	-	4	6	8	15
2000.....	143	-	-	-	-	-	-	2	1	7	19	23
1999.....	119	-	-	-	-	-	-	-	3	7	10	19

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Of corpus uteri and uterus, part unspecified (C54-C55)

All races, both sexes

2004.....	6,990	-	-	-	-	2	3	6	14	51	117	206
2003.....	6,899	-	-	-	-	1	3	8	19	37	117	171
2002.....	6,853	-	-	-	-	-	-	2	21	46	100	189
2001.....	6,784	-	-	-	-	-	-	10	28	47	92	192
2000.....	6,586	-	-	-	-	-	3	2	25	41	97	190
1999.....	6,468	-	-	-	-	-	2	11	12	38	104	182

All races, male

2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

All races, female

2004.....	6,990	-	-	-	-	2	3	6	14	51	117	206
2003.....	6,899	-	-	-	-	1	3	8	19	37	117	171
2002.....	6,853	-	-	-	-	-	-	2	21	46	100	189
2001.....	6,784	-	-	-	-	-	-	10	28	47	92	192
2000.....	6,586	-	-	-	-	-	3	2	25	41	97	190
1999.....	6,468	-	-	-	-	-	2	11	12	38	104	182

White, both sexes

2004.....	5,713	-	-	-	-	2	1	5	11	42	89	168
2003.....	5,641	-	-	-	-	1	2	5	11	29	79	133
2002.....	5,567	-	-	-	-	-	-	1	16	34	82	147
2001.....	5,606	-	-	-	-	-	-	3	23	33	74	150
2000.....	5,397	-	-	-	-	-	1	2	18	33	73	158
1999.....	5,386	-	-	-	-	-	2	8	10	31	78	150

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Of corpus uteri and uterus, part unspecified (C54-C55)

Asian or Pacific Islander 2/, both sexes

2004.....	142	-	-	-	-	-	-	-	-	1	6	8
2003.....	140	-	-	-	-	-	-	-	2	2	12	8
2002.....	135	-	-	-	-	-	-	-	1	1	3	10
2001.....	106	-	-	-	-	-	-	1	2	4	4	8
2000.....	109	-	-	-	-	-	1	-	1	2	7	7
1999.....	82	-	-	-	-	-	-	-	-	1	2	12

Asian or Pacific Islander 2/, male

2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

Asian or Pacific Islander 2/, female

2004.....	142	-	-	-	-	-	-	-	-	1	6	8
2003.....	140	-	-	-	-	-	-	-	2	2	12	8
2002.....	135	-	-	-	-	-	-	-	1	1	3	10
2001.....	106	-	-	-	-	-	-	1	2	4	4	8
2000.....	109	-	-	-	-	-	1	-	1	2	7	7
1999.....	82	-	-	-	-	-	-	-	-	1	2	12

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Of ovary (C56)

All races, both sexes

2004.....	14,716	-	-	-	2	8	21	31	63	128	326	647
2003.....	14,657	-	-	-	-	10	18	31	53	134	318	632
2002.....	14,682	1	1	1	1	12	25	33	64	150	319	688
2001.....	14,414	-	-	-	2	5	16	37	65	152	320	610
2000.....	14,060	-	-	2	-	7	16	36	65	145	327	617
1999.....	13,627	-	1	-	2	9	18	26	51	162	334	616

All races, male

2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

All races, female

2004.....	14,716	-	-	-	2	8	21	31	63	128	326	647
2003.....	14,657	-	-	-	-	10	18	31	53	134	318	632
2002.....	14,682	1	1	1	1	12	25	33	64	150	319	688
2001.....	14,414	-	-	-	2	5	16	37	65	152	320	610
2000.....	14,060	-	-	2	-	7	16	36	65	145	327	617
1999.....	13,627	-	1	-	2	9	18	26	51	162	334	616

White, both sexes

2004.....	13,127	-	-	-	2	6	14	20	50	102	273	539
2003.....	13,123	-	-	-	-	8	13	23	46	103	270	536
2002.....	13,202	1	1	1	1	9	21	29	48	122	269	582
2001.....	12,950	-	-	-	1	3	11	32	49	120	270	536
2000.....	12,731	-	-	1	-	5	12	32	52	120	277	541
1999.....	12,302	-	1	-	2	8	14	20	37	136	288	535

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Of ovary (C56)												
Asian or Pacific Islander 2/, both sexes												
2004.....	276	-	-	-	-	-	2	2	5	4	17	23
2003.....	276	-	-	-	-	-	1	1	2	7	12	18
2002.....	267	-	-	-	-	-	-	-	5	3	12	27
2001.....	243	-	-	-	-	1	2	-	6	4	8	21
2000.....	228	-	-	-	-	2	1	-	5	7	7	23
1999.....	217	-	-	-	-	-	1	-	7	10	14	25
Asian or Pacific Islander 2/, male												
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....
Asian or Pacific Islander 2/, female												
2004.....	276	-	-	-	-	-	2	2	5	4	17	23
2003.....	276	-	-	-	-	-	1	1	2	7	12	18
2002.....	267	-	-	-	-	-	-	-	5	3	12	27
2001.....	243	-	-	-	-	1	2	-	6	4	8	21
2000.....	228	-	-	-	-	2	1	-	5	7	7	23
1999.....	217	-	-	-	-	-	1	-	7	10	14	25

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Of other and unspecified female genital organs (C57)

All races, both sexes

2004.....	251	-	-	-	-	-	-	-	-	2	7	10
2003.....	355	-	-	-	-	-	1	2	2	1	7	7
2002.....	418	-	-	-	-	-	-	1	4	5	11	13
2001.....	386	-	-	-	-	-	-	2	1	2	4	13
2000.....	393	-	-	-	-	-	-	-	3	1	5	14
1999.....	408	-	-	-	-	-	-	-	-	3	4	16

All races, male

2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

All races, female

2004.....	251	-	-	-	-	-	-	-	-	2	7	10
2003.....	355	-	-	-	-	-	1	2	2	1	7	7
2002.....	418	-	-	-	-	-	-	1	4	5	11	13
2001.....	386	-	-	-	-	-	-	2	1	2	4	13
2000.....	393	-	-	-	-	-	-	-	3	1	5	14
1999.....	408	-	-	-	-	-	-	-	-	3	4	16

White, both sexes

2004.....	222	-	-	-	-	-	-	-	-	2	5	7
2003.....	311	-	-	-	-	-	1	2	2	1	7	4
2002.....	363	-	-	-	-	-	-	1	3	3	8	12
2001.....	345	-	-	-	-	-	-	1	1	2	3	10
2000.....	342	-	-	-	-	-	-	-	2	1	4	9
1999.....	359	-	-	-	-	-	-	-	-	3	4	13

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Of placenta (C58)												
All races, both sexes												
2004.....	20	-	-	-	-	-	2	2	2	3	1	2
2003.....	15	-	-	-	-	-	1	1	-	5	5	2
2002.....	14	-	-	-	-	-	2	-	2	1	3	2
2001.....	16	-	-	-	-	-	2	2	3	1	2	2
2000.....	16	-	-	-	-	-	1	1	1	4	2	3
1999.....	9	-	-	-	-	-	2	2	2	1	2	-
All races, male												
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....
All races, female												
2004.....	20	-	-	-	-	-	2	2	2	3	1	2
2003.....	15	-	-	-	-	-	1	1	-	5	5	2
2002.....	14	-	-	-	-	-	2	-	2	1	3	2
2001.....	16	-	-	-	-	-	2	2	3	1	2	2
2000.....	16	-	-	-	-	-	1	1	1	4	2	3
1999.....	9	-	-	-	-	-	2	2	2	1	2	-
White, both sexes												
2004.....	14	-	-	-	-	-	1	2	1	2	1	1
2003.....	8	-	-	-	-	-	1	1	-	3	2	-
2002.....	10	-	-	-	-	-	1	-	2	-	2	1
2001.....	8	-	-	-	-	-	-	2	-	1	2	1
2000.....	9	-	-	-	-	-	1	1	-	2	1	2
1999.....	4	-	-	-	-	-	1	1	2	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Of placenta (C58)

Asian or Pacific Islander 2/, both sexes

2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	2	-	-	-	-	-	-	-	-	1	1	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	2	-	-	-	-	-	-	-	1	-	-	1
2000.....	2	-	-	-	-	-	-	-	1	-	-	-
1999.....	1	-	-	-	-	-	-	-	-	1	-	-

Asian or Pacific Islander 2/, male

2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

Asian or Pacific Islander 2/, female

2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	2	-	-	-	-	-	-	-	-	1	1	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	2	-	-	-	-	-	-	-	1	-	-	1
2000.....	2	-	-	-	-	-	-	-	1	-	-	-
1999.....	1	-	-	-	-	-	-	-	-	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasms of male genital organs (C60-C63)

All races, both sexes

2004.....	29,629	-	1	1	1	11	38	38	40	51	79	157
2003.....	30,176	-	-	2	-	10	25	40	47	62	73	146
2002.....	31,084	-	-	-	-	9	36	43	55	56	88	134
2001.....	31,300	-	-	-	-	8	28	45	39	50	84	155
2000.....	31,675	-	-	-	1	9	35	43	46	68	74	148
1999.....	32,350	1	-	-	2	5	28	49	54	66	96	148

All races, male

2004.....	29,629	-	1	1	1	11	38	38	40	51	79	157
2003.....	30,176	-	-	2	-	10	25	40	47	62	73	146
2002.....	31,084	-	-	-	-	9	36	43	55	56	88	134
2001.....	31,300	-	-	-	-	8	28	45	39	50	84	155
2000.....	31,675	-	-	-	1	9	35	43	46	68	74	148
1999.....	32,350	1	-	-	2	5	28	49	54	66	96	148

All races, female

2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

White, both sexes

2004.....	24,284	-	1	1	1	9	32	33	37	45	69	114
2003.....	24,781	-	-	2	-	10	21	35	42	56	63	111
2002.....	25,485	-	-	-	-	9	30	37	50	50	78	95
2001.....	25,577	-	-	-	-	6	26	41	34	46	73	116
2000.....	25,877	-	-	-	1	7	32	40	41	60	60	104
1999.....	26,561	1	-	-	2	4	24	45	49	60	82	114

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Malignant neoplasms of male genital organs (C60-C63)

White, male

2004.....	24,284	-	1	1	1	9	32	33	37	45	69	114
2003.....	24,781	-	-	2	-	10	21	35	42	56	63	111
2002.....	25,485	-	-	-	-	9	30	37	50	50	78	95
2001.....	25,577	-	-	-	-	6	26	41	34	46	73	116
2000.....	25,877	-	-	-	1	7	32	40	41	60	60	104
1999.....	26,561	1	-	-	2	4	24	45	49	60	82	114

White, female

2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

Black, both sexes

2004.....	4,868	-	-	-	-	-	5	4	2	4	7	40
2003.....	4,952	-	-	-	-	-	1	4	2	6	10	32
2002.....	5,200	-	-	-	-	-	4	4	4	6	9	38
2001.....	5,318	-	-	-	-	1	1	3	3	4	11	35
2000.....	5,395	-	-	-	-	1	3	2	3	8	13	40
1999.....	5,408	-	-	-	-	1	2	3	3	6	12	32

Black, male

2004.....	4,868	-	-	-	-	-	5	4	2	4	7	40
2003.....	4,952	-	-	-	-	-	1	4	2	6	10	32
2002.....	5,200	-	-	-	-	-	4	4	4	6	9	38
2001.....	5,318	-	-	-	-	1	1	3	3	4	11	35
2000.....	5,395	-	-	-	-	1	3	2	3	8	13	40
1999.....	5,408	-	-	-	-	1	2	3	3	6	12	32

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Of prostate (C61)

All races, both sexes

2004.....	29,004	-	1	1	-	-	1	1	1	1	22	102
2003.....	29,554	-	-	1	-	1	-	1	-	3	22	99
2002.....	30,446	-	-	-	-	-	1	-	1	7	25	80
2001.....	30,719	-	-	-	-	-	1	1	-	3	27	99
2000.....	31,078	-	-	-	-	-	-	1	-	6	19	108
1999.....	31,729	1	-	-	-	-	1	1	1	5	28	100

All races, male

2004.....	29,004	-	1	1	-	-	1	1	1	1	22	102
2003.....	29,554	-	-	1	-	1	-	1	-	3	22	99
2002.....	30,446	-	-	-	-	-	1	-	1	7	25	80
2001.....	30,719	-	-	-	-	-	1	1	-	3	27	99
2000.....	31,078	-	-	-	-	-	-	1	-	6	19	108
1999.....	31,729	1	-	-	-	-	1	1	1	5	28	100

All races, female

2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

White, both sexes

2004.....	23,727	-	1	1	-	-	1	1	1	-	17	65
2003.....	24,232	-	-	1	-	1	-	1	-	2	15	71
2002.....	24,918	-	-	-	-	-	-	-	1	6	19	48
2001.....	25,066	-	-	-	-	-	1	-	-	3	16	70
2000.....	25,340	-	-	-	-	-	-	1	-	4	12	70
1999.....	26,005	1	-	-	-	-	1	1	1	4	21	68

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Of prostate (C61)

White, male

2004.....	23,727	-	1	1	-	-	1	1	1	-	17	65
2003.....	24,232	-	-	1	-	1	-	1	-	2	15	71
2002.....	24,918	-	-	-	-	-	-	-	1	6	19	48
2001.....	25,066	-	-	-	-	-	1	-	-	3	16	70
2000.....	25,340	-	-	-	-	-	-	1	-	4	12	70
1999.....	26,005	1	-	-	-	-	1	1	1	4	21	68

White, female

2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

Black, both sexes

2004.....	4,816	-	-	-	-	-	-	-	-	-	4	35
2003.....	4,894	-	-	-	-	-	-	-	-	1	7	28
2002.....	5,145	-	-	-	-	-	1	-	-	1	6	31
2001.....	5,265	-	-	-	-	-	-	1	-	-	11	28
2000.....	5,346	-	-	-	-	-	-	-	-	2	7	35
1999.....	5,357	-	-	-	-	-	-	-	-	1	6	31

Black, male

2004.....	4,816	-	-	-	-	-	-	-	-	-	4	35
2003.....	4,894	-	-	-	-	-	-	-	-	1	7	28
2002.....	5,145	-	-	-	-	-	1	-	-	1	6	31
2001.....	5,265	-	-	-	-	-	-	1	-	-	11	28
2000.....	5,346	-	-	-	-	-	-	-	-	2	7	35
1999.....	5,357	-	-	-	-	-	-	-	-	1	6	31

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Of testis (C62)												
All races, both sexes												
2004.....	357	-	-	-	1	11	37	36	37	47	50	43
2003.....	344	-	-	1	-	9	25	39	45	54	39	34
2002.....	393	-	-	-	-	9	34	42	52	45	54	42
2001.....	335	-	-	-	-	8	27	42	38	39	51	42
2000.....	338	-	-	-	1	9	33	41	45	54	49	30
1999.....	378	-	-	-	2	5	27	48	52	55	59	43
All races, male												
2004.....	357	-	-	-	1	11	37	36	37	47	50	43
2003.....	344	-	-	1	-	9	25	39	45	54	39	34
2002.....	393	-	-	-	-	9	34	42	52	45	54	42
2001.....	335	-	-	-	-	8	27	42	38	39	51	42
2000.....	338	-	-	-	1	9	33	41	45	54	49	30
1999.....	378	-	-	-	2	5	27	48	52	55	59	43
All races, female												
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....
White, both sexes												
2004.....	321	-	-	-	1	9	31	32	34	43	45	38
2003.....	312	-	-	1	-	9	21	34	40	49	38	30
2002.....	357	-	-	-	-	9	29	37	47	41	51	38
2001.....	303	-	-	-	-	6	25	39	34	36	51	36
2000.....	306	-	-	-	1	7	30	38	40	51	44	25
1999.....	347	-	-	-	2	4	23	44	47	50	55	41

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Of testis (C62)

White, male

2004.....	321	-	-	-	1	9	31	32	34	43	45	38
2003.....	312	-	-	1	-	9	21	34	40	49	38	30
2002.....	357	-	-	-	-	9	29	37	47	41	51	38
2001.....	303	-	-	-	-	6	25	39	34	36	51	36
2000.....	306	-	-	-	1	7	30	38	40	51	44	25
1999.....	347	-	-	-	2	4	23	44	47	50	55	41

White, female

2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

Black, both sexes

2004.....	25	-	-	-	-	-	5	3	2	3	3	4
2003.....	24	-	-	-	-	-	1	4	2	5	1	3
2002.....	28	-	-	-	-	-	3	4	4	4	2	4
2001.....	22	-	-	-	-	1	1	2	2	3	-	3
2000.....	25	-	-	-	-	1	3	2	3	3	4	4
1999.....	23	-	-	-	-	1	2	3	3	5	3	1

Black, male

2004.....	25	-	-	-	-	-	5	3	2	3	3	4
2003.....	24	-	-	-	-	-	1	4	2	5	1	3
2002.....	28	-	-	-	-	-	3	4	4	4	2	4
2001.....	22	-	-	-	-	1	1	2	2	3	-	3
2000.....	25	-	-	-	-	1	3	2	3	3	4	4
1999.....	23	-	-	-	-	1	2	3	3	5	3	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Of penis and other and unspecified male genital organs (C60,C63)

All races, both sexes

2004.....	268	-	-	-	-	-	-	1	2	3	7	12
2003.....	278	-	-	-	-	-	-	-	2	5	12	13
2002.....	245	-	-	-	-	-	1	1	2	4	9	12
2001.....	246	-	-	-	-	-	-	2	1	8	6	14
2000.....	259	-	-	-	-	-	2	1	1	8	6	10
1999.....	243	-	-	-	-	-	-	-	1	6	9	5

All races, male

2004.....	268	-	-	-	-	-	-	1	2	3	7	12
2003.....	278	-	-	-	-	-	-	-	2	5	12	13
2002.....	245	-	-	-	-	-	1	1	2	4	9	12
2001.....	246	-	-	-	-	-	-	2	1	8	6	14
2000.....	259	-	-	-	-	-	2	1	1	8	6	10
1999.....	243	-	-	-	-	-	-	-	1	6	9	5

All races, female

2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

White, both sexes

2004.....	236	-	-	-	-	-	-	-	2	2	7	11
2003.....	237	-	-	-	-	-	-	-	2	5	10	10
2002.....	210	-	-	-	-	-	1	-	2	3	8	9
2001.....	208	-	-	-	-	-	-	2	-	7	6	10
2000.....	231	-	-	-	-	-	2	1	1	5	4	9
1999.....	209	-	-	-	-	-	-	-	1	6	6	5

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Malignant neoplasms of urinary tract (C64-C68)

All races, both sexes

2004.....	25,928	4	18	23	7	8	9	24	38	113	282	675
2003.....	25,422	1	14	28	11	11	15	20	40	105	318	639
2002.....	25,444	4	8	25	8	7	14	19	46	123	312	706
2001.....	24,910	1	22	31	5	9	14	25	42	110	383	670
2000.....	24,344	2	15	21	9	10	13	15	37	125	330	622
1999.....	23,667	2	19	23	11	8	10	32	39	140	312	577

All races, male

2004.....	16,955	2	9	9	3	4	6	16	25	69	187	465
2003.....	16,569	1	4	15	4	7	11	10	28	71	227	483
2002.....	16,448	1	-	15	5	2	5	12	31	84	212	492
2001.....	16,205	1	10	7	2	8	11	12	23	72	241	480
2000.....	15,742	1	8	12	3	6	7	7	21	76	220	424
1999.....	15,309	1	11	12	5	4	7	19	28	91	213	420

All races, female

2004.....	8,973	2	9	14	4	4	3	8	13	44	95	210
2003.....	8,853	-	10	13	7	4	4	10	12	34	91	156
2002.....	8,996	3	8	10	3	5	9	7	15	39	100	214
2001.....	8,705	-	12	24	3	1	3	13	19	38	142	190
2000.....	8,602	1	7	9	6	4	6	8	16	49	110	198
1999.....	8,358	1	8	11	6	4	3	13	11	49	99	157

White, both sexes

2004.....	23,470	4	13	18	7	7	4	14	23	87	217	563
2003.....	22,927	1	14	24	6	8	10	13	26	77	247	517
2002.....	22,948	4	8	20	7	5	11	14	28	96	263	599
2001.....	22,443	-	16	22	2	3	8	16	29	81	303	567
2000.....	22,070	-	10	12	8	7	8	9	31	97	278	528
1999.....	21,319	2	12	19	7	5	5	18	25	102	256	467

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Malignant neoplasms of urinary tract (C64-C68)

White, male

2004.....	15,546	2	6	7	3	4	3	8	14	52	147	395
2003.....	15,105	1	4	12	2	5	8	5	17	54	177	406
2002.....	15,040	1	-	12	4	1	4	8	19	64	181	427
2001.....	14,786	-	6	4	1	2	6	8	16	54	198	422
2000.....	14,402	-	5	6	3	4	5	2	17	64	190	367
1999.....	13,948	1	8	10	2	4	3	9	17	65	177	344

White, female

2004.....	7,924	2	7	11	4	3	1	6	9	35	70	168
2003.....	7,822	-	10	12	4	3	2	8	9	23	70	111
2002.....	7,908	3	8	8	3	4	7	6	9	32	82	172
2001.....	7,657	-	10	18	1	1	2	8	13	27	105	145
2000.....	7,668	-	5	6	5	3	3	7	14	33	88	161
1999.....	7,371	1	4	9	5	1	2	9	8	37	79	123

Black, both sexes

2004.....	2,052	-	4	4	-	1	5	9	14	19	52	97
2003.....	2,084	-	-	4	5	3	5	7	12	23	64	102
2002.....	2,066	-	-	5	1	2	1	4	17	22	37	90
2001.....	2,045	-	4	9	3	5	5	9	11	26	70	86
2000.....	1,939	1	5	7	1	3	4	6	4	24	42	80
1999.....	2,002	-	4	3	3	3	5	13	12	36	46	93

Black, male

2004.....	1,150	-	3	1	-	-	3	7	11	12	35	61
2003.....	1,196	-	-	3	2	2	3	5	9	12	46	66
2002.....	1,129	-	-	3	1	1	1	3	11	16	23	52
2001.....	1,146	-	2	3	1	5	5	4	6	16	37	50
2000.....	1,117	-	3	4	-	2	1	5	3	9	23	46
1999.....	1,133	-	2	1	2	-	4	9	9	26	30	62

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasms of urinary tract (C64-C68)

Asian or Pacific Islander 2/, both sexes

2004.....	295	-	1	-	-	-	-	1	1	6	11	5
2003.....	302	-	-	-	-	-	-	-	2	2	5	16
2002.....	298	-	-	-	-	-	1	1	-	4	6	10
2001.....	308	-	1	-	-	-	1	-	2	2	7	11
2000.....	250	-	-	1	-	-	-	-	1	2	6	8
1999.....	246	-	1	1	1	-	-	-	2	2	7	13

Asian or Pacific Islander 2/, male

2004.....	185	-	-	-	-	-	-	1	-	4	5	3
2003.....	204	-	-	-	-	-	-	-	2	2	4	9
2002.....	196	-	-	-	-	-	-	1	-	4	4	9
2001.....	202	-	1	-	-	-	-	-	1	1	5	4
2000.....	168	-	-	1	-	-	-	-	-	1	4	8
1999.....	161	-	1	1	1	-	-	-	2	-	5	11

Asian or Pacific Islander 2/, female

2004.....	110	-	1	-	-	-	-	-	1	2	6	2
2003.....	98	-	-	-	-	-	-	-	-	-	1	7
2002.....	102	-	-	-	-	-	1	-	-	-	2	1
2001.....	106	-	-	-	-	-	1	-	1	1	2	7
2000.....	82	-	-	-	-	-	-	-	1	1	2	-
1999.....	85	-	-	-	-	-	-	-	-	2	2	2

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Of kidney and renal pelvis (C64-C65)

All races, both sexes

2004.....	12,313	4	18	23	6	8	8	22	27	76	192	470
2003.....	12,286	1	14	28	11	11	13	19	37	81	240	438
2002.....	12,165	4	8	23	7	7	13	14	34	91	215	514
2001.....	12,078	1	22	30	5	8	12	19	33	74	284	498
2000.....	11,736	2	15	21	8	8	12	14	28	83	251	450
1999.....	11,116	2	19	23	10	8	9	25	29	105	237	427

All races, male

2004.....	7,640	2	9	9	2	4	5	15	19	50	133	322
2003.....	7,667	1	4	15	4	7	9	9	27	56	175	324
2002.....	7,587	1	-	14	5	2	5	8	23	63	154	355
2001.....	7,524	1	10	7	2	7	9	11	16	58	182	355
2000.....	7,249	1	8	12	3	4	6	7	19	53	175	309
1999.....	6,867	1	11	12	4	4	7	15	22	70	166	310

All races, female

2004.....	4,673	2	9	14	4	4	3	7	8	26	59	148
2003.....	4,619	-	10	13	7	4	4	10	10	25	65	114
2002.....	4,578	3	8	9	2	5	8	6	11	28	61	159
2001.....	4,554	-	12	23	3	1	3	8	17	16	102	143
2000.....	4,487	1	7	9	5	4	6	7	9	30	76	141
1999.....	4,249	1	8	11	6	4	2	10	7	35	71	117

White, both sexes

2004.....	10,988	4	13	18	6	7	4	12	16	61	152	398
2003.....	10,930	1	14	24	6	8	8	12	23	56	191	358
2002.....	10,856	4	8	18	6	5	11	10	20	66	182	428
2001.....	10,720	-	16	21	2	2	6	10	21	51	223	424
2000.....	10,480	-	10	12	7	6	7	8	23	59	211	381
1999.....	9,878	2	12	19	6	5	5	14	18	76	195	346

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Of kidney and renal pelvis (C64-C65)

White, male

2004.....	6,839	2	6	7	2	4	3	7	10	39	108	276
2003.....	6,840	1	4	12	2	5	6	4	16	41	139	274
2002.....	6,803	1	-	11	4	1	4	5	14	45	131	304
2001.....	6,695	-	6	4	1	1	4	7	9	41	151	312
2000.....	6,475	-	5	6	3	3	4	2	15	42	150	266
1999.....	6,112	1	8	10	1	4	3	7	13	51	138	255

White, female

2004.....	4,149	2	7	11	4	3	1	5	6	22	44	122
2003.....	4,090	-	10	12	4	3	2	8	7	15	52	84
2002.....	4,053	3	8	7	2	4	7	5	6	21	51	124
2001.....	4,025	-	10	17	1	1	2	3	12	10	72	112
2000.....	4,005	-	5	6	4	3	3	6	8	17	61	115
1999.....	3,766	1	4	9	5	1	2	7	5	25	57	91

Black, both sexes

2004.....	1,091	-	4	4	-	1	4	9	11	10	34	62
2003.....	1,116	-	-	4	5	3	5	7	12	20	44	65
2002.....	1,069	-	-	5	1	2	1	3	14	21	25	72
2001.....	1,113	-	4	9	3	5	5	9	10	21	51	64
2000.....	1,044	1	5	7	1	2	4	6	4	21	31	58
1999.....	1,035	-	4	3	3	3	4	11	10	27	34	67

Black, male

2004.....	655	-	3	1	-	-	2	7	9	7	22	40
2003.....	677	-	-	3	2	2	3	5	9	10	34	44
2002.....	630	-	-	3	1	1	1	2	9	15	17	41
2001.....	678	-	2	3	1	5	5	4	6	15	25	40
2000.....	632	-	3	4	-	1	1	5	3	9	18	35
1999.....	622	-	2	1	2	-	4	8	8	19	23	44

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Of kidney and renal pelvis (C64-C65)

Asian or Pacific Islander 2/, both sexes

2004.....	143	-	1	-	-	-	-	1	-	4	5	2
2003.....	161	-	-	-	-	-	-	-	2	2	3	11
2002.....	142	-	-	-	-	-	1	1	-	3	2	7
2001.....	158	-	1	-	-	-	1	-	2	1	7	6
2000.....	148	-	-	1	-	-	-	-	-	2	5	5
1999.....	125	-	1	1	1	-	-	-	1	2	5	10

Asian or Pacific Islander 2/, male

2004.....	87	-	-	-	-	-	-	1	-	3	3	1
2003.....	102	-	-	-	-	-	-	-	2	2	2	4
2002.....	91	-	-	-	-	-	-	1	-	3	2	6
2001.....	97	-	1	-	-	-	-	-	1	1	5	1
2000.....	99	-	-	1	-	-	-	-	-	1	4	5
1999.....	83	-	1	1	1	-	-	-	1	-	4	8

Asian or Pacific Islander 2/, female

2004.....	56	-	1	-	-	-	-	-	-	1	2	1
2003.....	59	-	-	-	-	-	-	-	-	-	1	7
2002.....	51	-	-	-	-	-	1	-	-	-	-	1
2001.....	61	-	-	-	-	-	1	-	1	-	2	5
2000.....	49	-	-	-	-	-	-	-	-	1	1	-
1999.....	42	-	-	-	-	-	-	-	-	2	1	2

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Of bladder (C67)												
All races, both sexes												
2004.....	13,030	-	-	-	1	-	1	2	10	35	85	199
2003.....	12,483	-	-	-	-	-	1	1	3	24	74	181
2002.....	12,628	-	-	2	1	-	1	3	11	31	92	182
2001.....	12,225	-	-	1	-	1	2	6	9	35	93	161
2000.....	12,002	-	-	-	1	2	1	1	9	40	71	160
1999.....	11,911	-	-	-	1	-	1	6	10	32	68	140
All races, male												
2004.....	8,995	-	-	-	1	-	1	1	6	17	52	141
2003.....	8,576	-	-	-	-	-	1	1	1	15	49	149
2002.....	8,504	-	-	1	-	-	-	2	7	21	57	129
2001.....	8,343	-	-	-	-	1	2	1	7	14	56	118
2000.....	8,163	-	-	-	-	2	1	-	2	21	41	110
1999.....	8,105	-	-	-	1	-	-	3	6	20	43	107
All races, female												
2004.....	4,035	-	-	-	-	-	-	1	4	18	33	58
2003.....	3,907	-	-	-	-	-	-	-	2	9	25	32
2002.....	4,124	-	-	1	1	-	1	1	4	10	35	53
2001.....	3,882	-	-	1	-	-	-	5	2	21	37	43
2000.....	3,839	-	-	-	1	-	-	1	7	19	30	50
1999.....	3,806	-	-	-	-	-	1	3	4	12	25	33
White, both sexes												
2004.....	11,957	-	-	-	1	-	-	2	7	24	61	162
2003.....	11,409	-	-	-	-	-	1	1	3	21	54	145
2002.....	11,512	-	-	2	1	-	-	2	8	29	76	163
2001.....	11,180	-	-	1	-	1	2	6	8	30	74	133
2000.....	11,036	-	-	-	1	1	1	1	8	36	63	137
1999.....	10,863	-	-	-	1	-	-	3	7	23	57	114

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Of bladder (C67)												
White, male												
2004.....	8,407	-	-	-	1	-	-	1	4	11	37	117
2003.....	7,965	-	-	-	-	-	1	1	1	13	36	123
2002.....	7,908	-	-	1	-	-	-	1	5	19	49	116
2001.....	7,780	-	-	-	-	1	2	1	7	13	44	103
2000.....	7,620	-	-	-	-	1	1	-	2	20	37	96
1999.....	7,520	-	-	-	1	-	-	1	4	13	36	87
White, female												
2004.....	3,550	-	-	-	-	-	-	1	3	13	24	45
2003.....	3,444	-	-	-	-	-	-	-	2	8	18	22
2002.....	3,604	-	-	1	1	-	-	1	3	10	27	47
2001.....	3,400	-	-	1	-	-	-	5	1	17	30	30
2000.....	3,416	-	-	-	1	-	-	1	6	16	26	41
1999.....	3,343	-	-	-	-	-	-	2	3	10	21	27
Black, both sexes												
2004.....	913	-	-	-	-	-	1	-	3	9	18	32
2003.....	914	-	-	-	-	-	-	-	-	3	18	31
2002.....	940	-	-	-	-	-	-	1	3	1	12	16
2001.....	877	-	-	-	-	-	-	-	1	4	19	21
2000.....	852	-	-	-	-	1	-	-	-	3	7	20
1999.....	918	-	-	-	-	-	1	2	2	9	9	23
Black, male												
2004.....	478	-	-	-	-	-	1	-	2	5	13	21
2003.....	502	-	-	-	-	-	-	-	-	2	11	21
2002.....	481	-	-	-	-	-	-	1	2	1	6	10
2001.....	447	-	-	-	-	-	-	-	-	1	12	10
2000.....	466	-	-	-	-	1	-	-	-	-	4	11
1999.....	496	-	-	-	-	-	-	1	1	7	6	17

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Of other and unspecified sites within the urinary tract (C66,C68)

All races, both sexes

2004.....	585	-	-	-	-	-	-	-	1	2	5	6
2003.....	653	-	-	-	-	-	1	-	-	-	4	20
2002.....	651	-	-	-	-	-	-	2	1	1	5	10
2001.....	607	-	-	-	-	-	-	-	-	1	6	11
2000.....	606	-	-	-	-	-	-	-	-	2	8	12
1999.....	640	-	-	-	-	-	-	1	-	3	7	10

All races, male

2004.....	320	-	-	-	-	-	-	-	-	2	2	2
2003.....	326	-	-	-	-	-	1	-	-	-	3	10
2002.....	357	-	-	-	-	-	-	2	1	-	1	8
2001.....	338	-	-	-	-	-	-	-	-	-	3	7
2000.....	330	-	-	-	-	-	-	-	-	2	4	5
1999.....	337	-	-	-	-	-	-	1	-	1	4	3

All races, female

2004.....	265	-	-	-	-	-	-	-	1	-	3	4
2003.....	327	-	-	-	-	-	-	-	-	-	1	10
2002.....	294	-	-	-	-	-	-	-	-	1	4	2
2001.....	269	-	-	-	-	-	-	-	-	1	3	4
2000.....	276	-	-	-	-	-	-	-	-	-	4	7
1999.....	303	-	-	-	-	-	-	-	-	2	3	7

White, both sexes

2004.....	525	-	-	-	-	-	-	-	-	2	4	3
2003.....	588	-	-	-	-	-	1	-	-	-	2	14
2002.....	580	-	-	-	-	-	-	2	-	1	5	8
2001.....	543	-	-	-	-	-	-	-	-	-	6	10
2000.....	554	-	-	-	-	-	-	-	-	2	4	10
1999.....	578	-	-	-	-	-	-	1	-	3	4	7

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Malignant neoplasm of eye and adnexa (C69)												
All races, both sexes												
2004.....	208	1	6	2	2	1	-	2	-	4	6	13
2003.....	231	-	5	-	2	1	1	-	2	4	7	9
2002.....	240	-	3	3	2	-	2	-	-	6	7	14
2001.....	226	-	8	5	-	3	1	3	3	1	6	10
2000.....	236	-	8	7	-	1	-	-	3	1	7	10
1999.....	227	-	6	2	-	1	1	-	-	9	7	12
All races, male												
2004.....	112	1	2	2	1	1	-	1	-	3	3	8
2003.....	122	-	4	-	1	1	1	-	2	1	3	5
2002.....	117	-	1	1	1	-	1	-	-	1	2	11
2001.....	116	-	6	2	-	1	1	1	-	1	3	7
2000.....	130	-	5	2	-	1	-	-	1	1	4	7
1999.....	106	-	3	2	-	1	1	-	-	1	4	3
All races, female												
2004.....	96	-	4	-	1	-	-	1	-	1	3	5
2003.....	109	-	1	-	1	-	-	-	-	3	4	4
2002.....	123	-	2	2	1	-	1	-	-	5	5	3
2001.....	110	-	2	3	-	2	-	2	3	-	3	3
2000.....	106	-	3	5	-	-	-	-	2	-	3	3
1999.....	121	-	3	-	-	-	-	-	-	8	3	9
White, both sexes												
2004.....	204	-	4	2	2	1	-	2	-	4	6	13
2003.....	220	-	1	-	1	1	1	-	2	4	7	9
2002.....	226	-	2	1	2	-	1	-	-	6	7	12
2001.....	212	-	5	1	-	2	1	3	1	1	6	10
2000.....	218	-	6	5	-	1	-	-	2	-	5	8
1999.....	218	-	6	1	-	1	-	-	-	7	6	11

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Malignant neoplasm of eye and adnexa (C69)												
White, male												
2004.....	109	-	1	2	1	1	-	1	-	3	3	8
2003.....	115	-	1	-	1	1	1	-	2	1	3	5
2002.....	111	-	1	-	1	-	1	-	-	1	2	9
2001.....	112	-	5	1	-	-	1	1	-	1	3	7
2000.....	121	-	4	2	-	1	-	-	1	-	2	7
1999.....	103	-	3	1	-	1	-	-	-	1	3	3
White, female												
2004.....	95	-	3	-	1	-	-	1	-	1	3	5
2003.....	105	-	-	-	-	-	-	-	-	3	4	4
2002.....	115	-	1	1	1	-	-	-	-	5	5	3
2001.....	100	-	-	-	-	2	-	2	1	-	3	3
2000.....	97	-	2	3	-	-	-	-	1	-	3	1
1999.....	115	-	3	-	-	-	-	-	-	6	3	8
Black, both sexes												
2004.....	3	1	1	-	-	-	-	-	-	-	-	-
2003.....	10	-	3	-	1	-	-	-	-	-	-	-
2002.....	10	-	1	2	-	-	1	-	-	-	-	1
2001.....	8	-	3	1	-	1	-	-	-	-	-	-
2000.....	16	-	2	2	-	-	-	-	1	1	2	2
1999.....	8	-	-	-	-	-	1	-	-	2	1	1
Black, male												
2004.....	2	1	-	-	-	-	-	-	-	-	-	-
2003.....	6	-	2	-	-	-	-	-	-	-	-	-
2002.....	4	-	-	1	-	-	-	-	-	-	-	1
2001.....	4	-	1	1	-	1	-	-	-	-	-	-
2000.....	8	-	1	-	-	-	-	-	-	1	2	-
1999.....	2	-	-	-	-	-	1	-	-	-	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)

All races, both sexes

2004.....	12,829	19	102	186	139	109	114	137	222	352	552	832
2003.....	12,901	18	104	175	161	119	117	165	240	411	668	857
2002.....	12,830	12	114	202	135	108	102	129	259	375	634	900
2001.....	12,609	16	100	170	130	104	120	134	282	399	629	830
2000.....	12,655	20	104	166	150	105	120	159	247	454	642	815
1999.....	12,765	10	113	184	124	115	103	177	267	443	622	842

All races, male

2004.....	7,074	10	56	102	71	62	71	80	138	224	336	516
2003.....	7,122	9	56	89	86	68	68	85	147	246	393	544
2002.....	7,125	8	63	112	79	65	71	67	156	222	396	560
2001.....	7,015	8	52	81	64	63	66	88	174	245	393	500
2000.....	7,011	10	58	89	87	63	69	94	170	293	411	481
1999.....	7,001	8	56	96	58	69	62	104	172	268	370	507

All races, female

2004.....	5,755	9	46	84	68	47	43	57	84	128	216	316
2003.....	5,779	9	48	86	75	51	49	80	93	165	275	313
2002.....	5,705	4	51	90	56	43	31	62	103	153	238	340
2001.....	5,594	8	48	89	66	41	54	46	108	154	236	330
2000.....	5,644	10	46	77	63	42	51	65	77	161	231	334
1999.....	5,764	2	57	88	66	46	41	73	95	175	252	335

White, both sexes

2004.....	11,793	18	82	145	104	91	93	115	191	319	484	741
2003.....	11,849	15	85	140	131	99	98	144	211	366	610	767
2002.....	11,794	7	80	159	106	88	86	109	229	341	581	826
2001.....	11,638	12	84	144	100	88	95	120	254	374	568	753
2000.....	11,672	16	82	135	120	84	99	140	220	405	564	745
1999.....	11,819	7	96	139	103	86	85	165	238	408	558	775

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)

White, male

2004.....	6,508	9	44	79	55	54	59	66	120	203	289	459
2003.....	6,543	8	48	70	69	55	57	75	131	217	355	495
2002.....	6,554	5	46	84	59	55	62	56	135	203	369	514
2001.....	6,498	6	46	66	47	54	53	81	157	232	357	461
2000.....	6,486	9	46	72	69	52	57	82	145	264	366	449
1999.....	6,492	6	49	75	50	50	51	97	153	249	328	463

White, female

2004.....	5,285	9	38	66	49	37	34	49	71	116	195	282
2003.....	5,306	7	37	70	62	44	41	69	80	149	255	272
2002.....	5,240	2	34	75	47	33	24	53	94	138	212	312
2001.....	5,140	6	38	78	53	34	42	39	97	142	211	292
2000.....	5,186	7	36	63	51	32	42	58	75	141	198	296
1999.....	5,327	1	47	64	53	36	34	68	85	159	230	312

Black, both sexes

2004.....	765	-	18	25	25	17	16	16	21	20	52	60
2003.....	811	2	17	27	21	17	17	19	17	35	48	70
2002.....	781	3	27	34	20	16	12	11	22	30	38	49
2001.....	741	1	10	19	25	12	17	8	20	18	45	64
2000.....	772	2	18	26	26	16	19	15	21	34	57	52
1999.....	764	3	12	37	18	23	14	10	26	26	51	59

Black, male

2004.....	409	-	10	12	12	7	7	8	12	11	39	37
2003.....	447	1	6	16	11	11	10	9	10	23	33	38
2002.....	409	2	12	23	12	9	5	5	17	16	19	25
2001.....	382	-	3	12	13	6	10	4	10	8	25	34
2000.....	414	1	10	15	16	9	12	9	19	19	35	24
1999.....	404	2	5	18	7	15	9	6	17	15	33	39

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)

Black, female

2004.....	356	-	8	13	13	10	9	8	9	9	13	23
2003.....	364	1	11	11	10	6	7	10	7	12	15	32
2002.....	372	1	15	11	8	7	7	6	5	14	19	24
2001.....	359	1	7	7	12	6	7	4	10	10	20	30
2000.....	358	1	8	11	10	7	7	6	2	15	22	28
1999.....	360	1	7	19	11	8	5	4	9	11	18	20

American Indian 1/, both sexes

2004.....	34	1	1	2	-	-	2	1	-	1	-	8
2003.....	45	-	1	2	2	1	1	-	1	2	1	5
2002.....	41	1	1	4	3	2	-	1	3	-	3	4
2001.....	43	2	1	-	-	1	-	1	1	5	2	2
2000.....	41	-	2	2	2	2	-	1	-	7	4	4
1999.....	34	-	1	1	-	1	1	-	-	2	7	-

American Indian 1/, male

2004.....	25	1	1	2	-	-	2	1	-	1	-	6
2003.....	26	-	1	1	1	1	-	-	-	1	1	4
2002.....	23	-	-	2	2	1	-	1	-	-	2	4
2001.....	25	2	1	-	-	-	-	1	1	4	2	-
2000.....	20	-	1	1	1	1	-	-	-	4	1	1
1999.....	19	-	-	-	-	1	-	-	-	2	5	-

American Indian 1/, female

2004.....	9	-	-	-	-	-	-	-	-	-	-	2
2003.....	19	-	-	1	1	-	1	-	1	1	-	1
2002.....	18	1	1	2	1	1	-	-	3	-	1	-
2001.....	18	-	-	-	-	1	-	-	-	1	-	2
2000.....	21	-	1	1	1	1	-	1	-	3	3	3
1999.....	15	-	1	1	-	-	1	-	-	-	2	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)

Asian or Pacific Islander 2/, both sexes

2004.....	237	-	1	14	10	1	3	5	10	12	16	23
2003.....	196	1	1	6	7	2	1	2	11	8	9	15
2002.....	214	1	6	5	6	2	4	8	5	4	12	21
2001.....	187	1	5	7	5	3	8	5	7	2	14	11
2000.....	170	2	2	3	2	3	2	3	6	8	17	14
1999.....	148	-	4	7	3	5	3	2	3	7	6	8

Asian or Pacific Islander 2/, male

2004.....	132	-	1	9	4	1	3	5	6	9	8	14
2003.....	106	-	1	2	5	1	1	1	6	5	4	7
2002.....	139	1	5	3	6	-	4	5	4	3	6	17
2001.....	110	-	2	3	4	3	3	2	6	1	9	5
2000.....	91	-	1	1	1	1	-	3	6	6	9	7
1999.....	86	-	2	3	1	3	2	1	2	2	4	5

Asian or Pacific Islander 2/, female

2004.....	105	-	-	5	6	-	-	-	4	3	8	9
2003.....	90	1	-	4	2	1	-	1	5	3	5	8
2002.....	75	-	1	2	-	2	-	3	1	1	6	4
2001.....	77	1	3	4	1	-	5	3	1	1	5	6
2000.....	79	2	1	2	1	2	2	-	-	2	8	7
1999.....	62	-	2	4	2	2	1	1	1	5	2	3

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Of brain (C71)

All races, both sexes

2004.....	12,606	19	97	184	130	106	109	134	218	343	546	824
2003.....	12,663	15	100	170	158	118	111	161	236	399	661	846
2002.....	12,556	11	109	199	132	101	100	126	251	369	623	885
2001.....	12,372	15	94	166	128	99	117	132	277	395	623	819
2000.....	12,412	20	100	164	147	102	115	153	238	450	626	797
1999.....	12,484	8	110	180	118	108	100	172	260	430	606	828

All races, male

2004.....	6,964	10	55	102	66	61	69	78	136	217	331	510
2003.....	7,015	7	54	85	85	68	65	84	147	242	390	537
2002.....	6,971	7	62	111	76	61	70	65	152	218	389	552
2001.....	6,889	8	50	80	63	60	63	87	170	243	388	494
2000.....	6,892	10	54	89	85	62	65	89	163	292	401	472
1999.....	6,852	6	54	92	58	67	59	99	168	256	360	499

All races, female

2004.....	5,642	9	42	82	64	45	40	56	82	126	215	314
2003.....	5,648	8	46	85	73	50	46	77	89	157	271	309
2002.....	5,585	4	47	88	56	40	30	61	99	151	234	333
2001.....	5,483	7	44	86	65	39	54	45	107	152	235	325
2000.....	5,520	10	46	75	62	40	50	64	75	158	225	325
1999.....	5,632	2	56	88	60	41	41	73	92	174	246	329

White, both sexes

2004.....	11,602	18	77	143	97	88	91	114	188	312	480	734
2003.....	11,638	12	82	136	129	98	94	140	207	354	603	756
2002.....	11,561	6	77	156	105	82	84	108	221	335	571	814
2001.....	11,422	11	78	140	98	84	92	118	251	371	564	743
2000.....	11,457	16	78	133	117	81	95	134	213	402	552	728
1999.....	11,576	5	94	135	98	80	83	161	234	396	544	762

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Of brain (C71)												
White, male												
2004.....	6,413	9	43	79	50	53	58	66	119	196	285	454
2003.....	6,448	6	46	67	69	55	55	74	131	213	352	488
2002.....	6,423	4	45	83	58	51	61	55	131	199	362	507
2001.....	6,385	6	44	65	46	52	50	80	155	230	353	456
2000.....	6,380	9	42	72	67	51	53	77	140	263	357	440
1999.....	6,365	4	48	71	50	48	49	93	152	238	319	456
White, female												
2004.....	5,189	9	34	64	47	35	33	48	69	116	195	280
2003.....	5,190	6	36	69	60	43	39	66	76	141	251	268
2002.....	5,138	2	32	73	47	31	23	53	90	136	209	307
2001.....	5,037	5	34	75	52	32	42	38	96	141	211	287
2000.....	5,077	7	36	61	50	30	42	57	73	139	195	288
1999.....	5,211	1	46	64	48	32	34	68	82	158	225	306
Black, both sexes												
2004.....	741	-	18	25	23	17	14	15	21	19	51	59
2003.....	788	2	16	26	20	17	16	19	17	35	48	70
2002.....	749	3	26	34	19	15	12	10	22	30	38	47
2001.....	726	1	10	19	25	11	17	8	19	17	43	63
2000.....	749	2	18	26	26	16	18	15	19	34	53	51
1999.....	733	3	12	37	17	22	13	10	23	25	49	59
Black, male												
2004.....	397	-	10	12	12	7	7	7	12	11	38	36
2003.....	437	1	6	15	10	11	9	9	10	23	33	38
2002.....	391	2	12	23	11	9	5	5	17	16	19	24
2001.....	373	-	3	12	13	5	10	4	9	8	24	33
2000.....	403	1	10	15	16	9	12	9	17	19	34	24
1999.....	387	2	5	18	7	15	8	6	14	14	32	39

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Of brain (C71)												
Black, female												
2004.....	344	-	8	13	11	10	7	8	9	8	13	23
2003.....	351	1	10	11	10	6	7	10	7	12	15	32
2002.....	358	1	14	11	8	6	7	5	5	14	19	23
2001.....	353	1	7	7	12	6	7	4	10	9	19	30
2000.....	346	1	8	11	10	7	6	6	2	15	19	27
1999.....	346	1	7	19	10	7	5	4	9	11	17	20
American Indian 1/, both sexes												
2004.....	34	1	1	2	-	-	2	1	-	1	-	8
2003.....	44	-	1	2	2	1	-	-	1	2	1	5
2002.....	40	1	-	4	3	2	-	1	3	-	3	4
2001.....	43	2	1	-	-	1	-	1	1	5	2	2
2000.....	41	-	2	2	2	2	-	1	-	7	4	4
1999.....	32	-	1	1	-	1	1	-	-	2	7	-
American Indian 1/, male												
2004.....	25	1	1	2	-	-	2	1	-	1	-	6
2003.....	26	-	1	1	1	1	-	-	-	1	1	4
2002.....	23	-	-	2	2	1	-	1	-	-	2	4
2001.....	25	2	1	-	-	-	-	1	1	4	2	-
2000.....	20	-	1	1	1	1	-	-	-	4	1	1
1999.....	18	-	-	-	-	1	-	-	-	2	5	-
American Indian 1/, female												
2004.....	9	-	-	-	-	-	-	-	-	-	-	2
2003.....	18	-	-	1	1	-	-	-	1	1	-	1
2002.....	17	1	-	2	1	1	-	-	3	-	1	-
2001.....	18	-	-	-	-	1	-	-	-	1	-	2
2000.....	21	-	1	1	1	1	-	1	-	3	3	3
1999.....	14	-	1	1	-	-	1	-	-	-	2	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Of brain (C71)												
Asian or Pacific Islander 2/, both sexes												
2004.....	229	-	1	14	10	1	2	4	9	11	15	23
2003.....	193	1	1	6	7	2	1	2	11	8	9	15
2002.....	206	1	6	5	5	2	4	7	5	4	11	20
2001.....	181	1	5	7	5	3	8	5	6	2	14	11
2000.....	165	2	2	3	2	3	2	3	6	7	17	14
1999.....	143	-	3	7	3	5	3	1	3	7	6	7
Asian or Pacific Islander 2/, male												
2004.....	129	-	1	9	4	1	2	4	5	9	8	14
2003.....	104	-	1	2	5	1	1	1	6	5	4	7
2002.....	134	1	5	3	5	-	4	4	4	3	6	17
2001.....	106	-	2	3	4	3	3	2	5	1	9	5
2000.....	89	-	1	1	1	1	-	3	6	6	9	7
1999.....	82	-	1	3	1	3	2	-	2	2	4	4
Asian or Pacific Islander 2/, female												
2004.....	100	-	-	5	6	-	-	-	4	2	7	9
2003.....	89	1	-	4	2	1	-	1	5	3	5	8
2002.....	72	-	1	2	-	2	-	3	1	1	5	3
2001.....	75	1	3	4	1	-	5	3	1	1	5	6
2000.....	76	2	1	2	1	2	2	-	-	1	8	7
1999.....	61	-	2	4	2	2	1	1	1	5	2	3

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Of other parts of central nervous system (C70,C72)												
All races, both sexes												
2004.....	223	-	5	2	9	3	5	3	4	9	6	8
2003.....	238	3	4	5	3	1	6	4	4	12	7	11
2002.....	274	1	5	3	3	7	2	3	8	6	11	15
2001.....	237	1	6	4	2	5	3	2	5	4	6	11
2000.....	243	-	4	2	3	3	5	6	9	4	16	18
1999.....	281	2	3	4	6	7	3	5	7	13	16	14
All races, male												
2004.....	110	-	1	-	5	1	2	2	2	7	5	6
2003.....	107	2	2	4	1	-	3	1	-	4	3	7
2002.....	154	1	1	1	3	4	1	2	4	4	7	8
2001.....	126	-	2	1	1	3	3	1	4	2	5	6
2000.....	119	-	4	-	2	1	4	5	7	1	10	9
1999.....	149	2	2	4	-	2	3	5	4	12	10	8
All races, female												
2004.....	113	-	4	2	4	2	3	1	2	2	1	2
2003.....	131	1	2	1	2	1	3	3	4	8	4	4
2002.....	120	-	4	2	-	3	1	1	4	2	4	7
2001.....	111	1	4	3	1	2	-	1	1	2	1	5
2000.....	124	-	-	2	1	2	1	1	2	3	6	9
1999.....	132	-	1	-	6	5	-	-	3	1	6	6
White, both sexes												
2004.....	191	-	5	2	7	3	2	1	3	7	4	7
2003.....	211	3	3	4	2	1	4	4	4	12	7	11
2002.....	233	1	3	3	1	6	2	1	8	6	10	12
2001.....	216	1	6	4	2	4	3	2	3	3	4	10
2000.....	215	-	4	2	3	3	4	6	7	3	12	17
1999.....	243	2	2	4	5	6	2	4	4	12	14	13

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Of other parts of central nervous system (C70,C72)

White, male

2004.....	95	-	1	-	5	1	1	-	1	7	4	5
2003.....	95	2	2	3	-	-	2	1	-	4	3	7
2002.....	131	1	1	1	1	4	1	1	4	4	7	7
2001.....	113	-	2	1	1	2	3	1	2	2	4	5
2000.....	106	-	4	-	2	1	4	5	5	1	9	9
1999.....	127	2	1	4	-	2	2	4	1	11	9	7

White, female

2004.....	96	-	4	2	2	2	1	1	2	-	-	2
2003.....	116	1	1	1	2	1	2	3	4	8	4	4
2002.....	102	-	2	2	-	2	1	-	4	2	3	5
2001.....	103	1	4	3	1	2	-	1	1	1	-	5
2000.....	109	-	-	2	1	2	-	1	2	2	3	8
1999.....	116	-	1	-	5	4	-	-	3	1	5	6

Black, both sexes

2004.....	24	-	-	-	2	-	2	1	-	1	1	1
2003.....	23	-	1	1	1	-	1	-	-	-	-	-
2002.....	32	-	1	-	1	1	-	1	-	-	-	2
2001.....	15	-	-	-	-	1	-	-	1	1	2	1
2000.....	23	-	-	-	-	-	1	-	2	-	4	1
1999.....	31	-	-	-	1	1	1	-	3	1	2	-

Black, male

2004.....	12	-	-	-	-	-	-	1	-	-	1	1
2003.....	10	-	-	1	1	-	1	-	-	-	-	-
2002.....	18	-	-	-	1	-	-	-	-	-	-	1
2001.....	9	-	-	-	-	1	-	-	1	-	1	1
2000.....	11	-	-	-	-	-	-	-	2	-	1	-
1999.....	17	-	-	-	-	-	1	-	3	1	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	-------------	-----------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Malignant neoplasms of thyroid and other endocrine glands (C73-C75)

All races, both sexes

2004.....	2,086	11	75	69	26	15	23	15	14	32	42	81
2003.....	1,970	5	66	80	23	4	7	18	22	44	40	85
2002.....	2,059	7	74	71	29	12	14	14	23	31	54	96
2001.....	2,104	7	72	54	27	15	14	23	17	34	54	100
2000.....	2,068	10	76	63	25	14	18	11	20	40	53	79
1999.....	2,007	5	74	77	25	13	13	10	21	50	52	85

All races, male

2004.....	923	5	38	43	14	9	14	6	10	15	26	30
2003.....	864	3	38	46	11	2	3	11	9	23	21	44
2002.....	907	6	35	41	13	9	11	5	7	19	26	50
2001.....	947	2	42	32	14	6	8	10	10	19	29	51
2000.....	920	7	42	35	13	8	8	4	11	18	24	46
1999.....	913	3	43	41	14	7	7	6	10	25	30	41

All races, female

2004.....	1,163	6	37	26	12	6	9	9	4	17	16	51
2003.....	1,106	2	28	34	12	2	4	7	13	21	19	41
2002.....	1,152	1	39	30	16	3	3	9	16	12	28	46
2001.....	1,157	5	30	22	13	9	6	13	7	15	25	49
2000.....	1,148	3	34	28	12	6	10	7	9	22	29	33
1999.....	1,094	2	31	36	11	6	6	4	11	25	22	44

White, both sexes

2004.....	1,807	11	61	53	17	13	15	15	11	30	32	68
2003.....	1,737	4	54	63	14	3	7	15	17	37	36	78
2002.....	1,780	7	53	54	23	8	7	11	17	28	43	80
2001.....	1,830	6	63	43	19	12	12	19	11	24	48	84
2000.....	1,779	8	57	56	18	12	16	10	18	31	41	66
1999.....	1,747	4	65	62	15	8	10	10	17	40	39	76

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Malignant neoplasms of thyroid and other endocrine glands (C73-C75)

White, male

2004.....	810	5	30	34	12	8	8	6	8	15	19	26
2003.....	777	3	30	36	6	2	3	10	6	20	20	41
2002.....	792	6	27	30	11	5	6	3	6	17	20	39
2001.....	841	1	34	25	9	5	7	6	5	14	26	48
2000.....	819	6	35	31	9	7	6	4	11	16	19	38
1999.....	802	2	39	34	7	3	6	6	8	20	23	37

White, female

2004.....	997	6	31	19	5	5	7	9	3	15	13	42
2003.....	960	1	24	27	8	1	4	5	11	17	16	37
2002.....	988	1	26	24	12	3	1	8	11	11	23	41
2001.....	989	5	29	18	10	7	5	13	6	10	22	36
2000.....	960	2	22	25	9	5	10	6	7	15	22	28
1999.....	945	2	26	28	8	5	4	4	9	20	16	39

Black, both sexes

2004.....	196	-	9	14	9	2	6	-	3	1	9	7
2003.....	169	-	11	16	8	-	-	3	5	3	3	5
2002.....	206	-	15	14	5	3	6	2	6	2	9	11
2001.....	205	-	8	9	5	3	1	4	5	9	4	13
2000.....	215	1	15	7	6	2	2	1	2	8	7	11
1999.....	201	-	8	12	8	5	3	-	3	9	11	8

Black, male

2004.....	74	-	5	7	2	1	4	-	2	-	6	1
2003.....	65	-	7	9	5	-	-	1	3	1	1	2
2002.....	85	-	5	9	2	3	4	2	1	1	5	9
2001.....	80	-	7	6	3	1	1	4	4	4	2	2
2000.....	75	-	6	4	4	1	2	-	-	1	1	7
1999.....	89	-	4	5	6	4	1	-	1	4	6	4

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasms of thyroid and other endocrine glands (C73-C75)

Asian or Pacific Islander 2/, both sexes

2004.....	72	-	5	2	-	-	2	-	-	-	1	5
2003.....	58	1	1	1	1	1	-	-	-	2	1	1
2002.....	65	-	6	1	1	1	1	1	-	1	2	5
2001.....	61	1	1	2	3	-	1	-	-	1	1	3
2000.....	65	1	4	-	1	-	-	-	-	1	5	2
1999.....	52	1	1	3	2	-	-	-	1	1	1	-

Asian or Pacific Islander 2/, male

2004.....	34	-	3	2	-	-	2	-	-	-	1	2
2003.....	19	-	1	1	-	-	-	-	-	1	-	-
2002.....	27	-	3	1	-	1	1	-	-	1	1	2
2001.....	24	1	1	1	2	-	-	-	-	1	1	1
2000.....	23	1	1	-	-	-	-	-	-	1	4	1
1999.....	20	1	-	2	1	-	-	-	1	1	1	-

Asian or Pacific Islander 2/, female

2004.....	38	-	2	-	-	-	-	-	-	-	-	3
2003.....	39	1	-	-	1	1	-	-	-	1	1	1
2002.....	38	-	3	-	1	-	-	1	-	-	1	3
2001.....	37	-	-	1	1	-	1	-	-	-	-	2
2000.....	42	-	3	-	1	-	-	-	-	-	1	1
1999.....	32	-	1	1	1	-	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other malignant neoplasms of other and unspecified sites (C76-C80,C97)

All races, both sexes

2004.....	39,671	3	18	15	18	46	60	92	163	299	669	1,319
2003.....	40,271	10	16	7	20	45	58	79	131	303	680	1,382
2002.....	40,658	10	21	9	19	54	67	69	185	339	742	1,380
2001.....	40,035	4	15	15	24	43	55	92	157	351	745	1,306
2000.....	40,700	7	22	15	21	43	51	87	143	366	755	1,303
1999.....	43,850	4	15	17	20	31	72	100	193	393	859	1,379

All races, male

2004.....	20,338	1	9	7	13	28	42	47	81	160	334	730
2003.....	20,484	6	8	6	9	29	30	41	73	143	353	791
2002.....	20,787	7	14	6	10	33	46	39	99	171	373	754
2001.....	20,263	-	8	7	15	30	35	50	91	169	401	739
2000.....	20,442	1	15	10	13	21	27	50	72	171	404	745
1999.....	22,149	3	8	9	14	20	46	58	114	197	436	800

All races, female

2004.....	19,333	2	9	8	5	18	18	45	82	139	335	589
2003.....	19,787	4	8	1	11	16	28	38	58	160	327	591
2002.....	19,871	3	7	3	9	21	21	30	86	168	369	626
2001.....	19,772	4	7	8	9	13	20	42	66	182	344	567
2000.....	20,258	6	7	5	8	22	24	37	71	195	351	558
1999.....	21,701	1	7	8	6	11	26	42	79	196	423	579

White, both sexes

2004.....	34,198	3	11	11	11	40	45	72	131	232	517	985
2003.....	34,820	8	11	7	14	34	46	64	96	232	511	1,062
2002.....	35,073	8	15	7	15	45	51	48	144	250	570	1,070
2001.....	34,499	4	13	14	17	31	42	67	123	265	569	1,000
2000.....	35,232	5	16	10	12	35	43	72	110	277	566	992
1999.....	37,960	4	12	15	15	26	53	84	152	302	657	1,059

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other malignant neoplasms of other and unspecified sites (C76-C80,C97)

White, male

2004.....	17,426	1	7	5	6	25	32	37	71	125	249	537
2003.....	17,668	5	5	6	6	21	26	36	53	114	260	598
2002.....	17,937	7	10	5	10	28	32	27	83	123	271	591
2001.....	17,330	-	7	6	12	22	30	38	73	127	302	567
2000.....	17,548	1	12	6	9	17	22	41	58	131	293	559
1999.....	19,121	3	6	7	10	17	35	52	87	155	335	597

White, female

2004.....	16,772	2	4	6	5	15	13	35	60	107	268	448
2003.....	17,152	3	6	1	8	13	20	28	43	118	251	464
2002.....	17,136	1	5	2	5	17	19	21	61	127	299	479
2001.....	17,169	4	6	8	5	9	12	29	50	138	267	433
2000.....	17,684	4	4	4	3	18	21	31	52	146	273	433
1999.....	18,839	1	6	8	5	9	18	32	65	147	322	462

Black, both sexes

2004.....	4,608	-	4	4	6	5	13	16	22	56	133	287
2003.....	4,628	1	5	-	3	8	10	13	27	60	137	273
2002.....	4,747	2	6	2	4	6	13	18	31	74	139	264
2001.....	4,696	-	1	1	6	10	10	22	28	70	138	274
2000.....	4,658	2	5	5	7	7	6	13	21	79	166	272
1999.....	5,030	-	2	-	3	3	12	13	28	73	174	284

Black, male

2004.....	2,466	-	1	2	6	2	8	8	8	27	74	166
2003.....	2,393	1	3	-	2	5	4	5	13	23	74	165
2002.....	2,413	-	4	1	-	4	12	10	10	37	80	139
2001.....	2,484	-	1	1	2	6	5	11	12	32	75	153
2000.....	2,471	-	2	4	4	3	4	8	10	36	99	166
1999.....	2,583	-	2	-	2	2	7	3	19	32	84	177

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other malignant neoplasms of other and unspecified sites (C76-C80,C97)

Black, female

2004.....	2,142	-	3	2	-	3	5	8	14	29	59	121
2003.....	2,235	-	2	-	1	3	6	8	14	37	63	108
2002.....	2,334	2	2	1	4	2	1	8	21	37	59	125
2001.....	2,212	-	-	-	4	4	5	11	16	38	63	121
2000.....	2,187	2	3	1	3	4	2	5	11	43	67	106
1999.....	2,447	-	-	-	1	1	5	10	9	41	90	107

American Indian 1/, both sexes

2004.....	176	-	1	-	-	-	-	1	2	3	5	5
2003.....	170	-	-	-	-	-	-	-	-	2	8	5
2002.....	173	-	-	-	-	1	1	-	4	3	6	7
2001.....	183	-	-	-	-	1	1	-	1	2	12	7
2000.....	176	-	-	-	-	-	1	1	1	3	2	11
1999.....	200	-	-	1	1	1	-	1	1	6	6	12

American Indian 1/, male

2004.....	89	-	-	-	-	-	-	-	-	2	3	4
2003.....	94	-	-	-	-	-	-	-	-	1	5	3
2002.....	83	-	-	-	-	-	-	-	3	1	4	4
2001.....	99	-	-	-	-	1	-	-	1	2	8	6
2000.....	88	-	-	-	-	-	1	1	1	2	1	6
1999.....	100	-	-	1	1	-	-	1	-	4	3	10

American Indian 1/, female

2004.....	87	-	1	-	-	-	-	1	2	1	2	1
2003.....	76	-	-	-	-	-	-	-	-	1	3	2
2002.....	90	-	-	-	-	1	1	-	1	2	2	3
2001.....	84	-	-	-	-	-	1	-	-	-	4	1
2000.....	88	-	-	-	-	-	-	-	-	1	1	5
1999.....	100	-	-	-	-	1	-	-	1	2	3	2

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other malignant neoplasms of other and unspecified sites (C76-C80,C97)

Asian or Pacific Islander 2/, both sexes

2004.....	689	-	2	-	1	1	2	3	8	8	14	42
2003.....	653	1	-	-	3	3	2	2	8	9	24	42
2002.....	665	-	-	-	-	2	3	6	12	27	39	
2001.....	657	-	1	-	1	2	3	5	14	26	25	
2000.....	634	-	1	-	2	1	1	11	7	21	28	
1999.....	660	-	1	1	1	1	7	2	12	12	22	24

Asian or Pacific Islander 2/, male

2004.....	357	-	1	-	1	1	2	2	2	6	8	23
2003.....	329	-	-	-	1	3	-	7	5	14	25	
2002.....	354	-	-	-	-	1	2	3	10	18	20	
2001.....	350	-	-	-	1	1	-	5	8	16	13	
2000.....	335	-	1	-	-	1	-	3	2	11	14	
1999.....	345	-	-	1	1	1	4	2	8	6	14	16

Asian or Pacific Islander 2/, female

2004.....	332	-	1	-	-	-	-	1	6	2	6	19
2003.....	324	1	-	-	2	-	2	2	1	4	10	17
2002.....	311	-	-	-	-	1	-	1	3	2	9	19
2001.....	307	-	1	-	-	-	2	2	-	6	10	12
2000.....	299	-	-	-	2	-	1	1	8	5	10	14
1999.....	315	-	1	-	-	-	3	-	4	6	8	8

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)

All races, both sexes

2004.....	54,645	25	122	179	180	284	361	356	466	569	945	1,477
2003.....	55,679	31	138	173	221	252	354	373	492	655	1,073	1,557
2002.....	56,225	28	119	166	213	310	393	368	548	695	1,056	1,599
2001.....	56,341	27	157	165	202	311	394	436	552	768	1,089	1,674
2000.....	56,518	39	146	168	204	323	395	412	506	776	1,077	1,675
1999.....	56,318	30	137	142	209	303	417	423	537	837	1,076	1,648

All races, male

2004.....	29,522	10	72	111	103	171	228	214	275	337	578	881
2003.....	30,076	10	80	102	123	166	209	220	315	380	663	947
2002.....	30,302	15	63	87	130	188	229	225	320	409	629	953
2001.....	30,060	12	82	95	122	192	234	246	328	464	669	1,002
2000.....	29,888	21	76	98	125	202	253	226	313	438	637	1,015
1999.....	29,721	12	75	84	122	180	257	241	316	497	647	986

All races, female

2004.....	25,123	15	50	68	77	113	133	142	191	232	367	596
2003.....	25,603	21	58	71	98	86	145	153	177	275	410	610
2002.....	25,923	13	56	79	83	122	164	143	228	286	427	646
2001.....	26,281	15	75	70	80	119	160	190	224	304	420	672
2000.....	26,630	18	70	70	79	121	142	186	193	338	440	660
1999.....	26,597	18	62	58	87	123	160	182	221	340	429	662

White, both sexes

2004.....	48,240	20	94	145	144	230	285	265	361	441	759	1,187
2003.....	49,420	29	115	146	168	201	282	292	378	509	857	1,258
2002.....	49,943	19	98	137	166	244	331	287	423	542	863	1,276
2001.....	49,989	22	136	125	156	253	309	318	435	612	858	1,364
2000.....	50,394	36	121	136	170	249	300	315	392	617	868	1,340
1999.....	50,189	24	118	114	157	232	332	332	419	677	862	1,362

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)

White, male

2004.....	26,204	8	52	92	83	141	178	168	218	266	475	709
2003.....	26,943	8	67	92	92	136	165	175	244	300	536	775
2002.....	27,087	11	53	69	101	155	197	180	246	316	516	774
2001.....	26,767	9	69	72	85	159	187	182	269	360	532	833
2000.....	26,766	19	64	77	103	155	196	180	240	354	536	827
1999.....	26,558	12	63	70	90	136	213	186	245	408	518	816

White, female

2004.....	22,036	12	42	53	61	89	107	97	143	175	284	478
2003.....	22,477	21	48	54	76	65	117	117	134	209	321	483
2002.....	22,856	8	45	68	65	89	134	107	177	226	347	502
2001.....	23,222	13	67	53	71	94	122	136	166	252	326	531
2000.....	23,628	17	57	59	67	94	104	135	152	263	332	513
1999.....	23,631	12	55	44	67	96	119	146	174	269	344	546

Black, both sexes

2004.....	5,233	3	19	22	27	46	60	77	87	102	154	246
2003.....	5,103	1	14	20	37	34	54	65	91	117	184	250
2002.....	5,151	5	16	19	34	45	45	62	95	111	165	268
2001.....	5,227	4	14	29	36	48	63	85	98	133	190	257
2000.....	5,071	1	20	21	22	58	74	75	85	130	176	295
1999.....	5,088	4	15	21	42	54	80	72	88	135	179	239

Black, male

2004.....	2,650	-	14	13	13	24	39	39	48	57	82	142
2003.....	2,537	1	8	9	22	19	33	32	56	63	109	146
2002.....	2,622	2	7	13	20	25	20	35	58	70	98	155
2001.....	2,676	3	9	14	30	29	32	48	52	90	110	136
2000.....	2,558	-	9	14	12	36	44	33	58	66	87	163
1999.....	2,590	-	9	11	28	33	41	43	55	74	108	141

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)

Black, female

2004.....	2,583	3	5	9	14	22	21	38	39	45	72	104
2003.....	2,566	-	6	11	15	15	21	33	35	54	75	104
2002.....	2,529	3	9	6	14	20	25	27	37	41	67	113
2001.....	2,551	1	5	15	6	19	31	37	46	43	80	121
2000.....	2,513	1	11	7	10	22	30	42	27	64	89	132
1999.....	2,498	4	6	10	14	21	39	29	33	61	71	98

American Indian 1/, both sexes

2004.....	199	-	1	3	1	1	4	1	5	8	4	7
2003.....	210	-	3	2	1	7	5	2	3	5	7	9
2002.....	171	-	1	-	2	4	7	3	4	10	4	7
2001.....	225	-	1	-	4	3	7	11	3	5	13	7
2000.....	199	1	1	1	1	3	2	2	7	4	5	9
1999.....	161	1	1	-	2	1	2	4	3	3	6	7

American Indian 1/, male

2004.....	112	-	1	2	1	1	1	1	2	4	2	6
2003.....	99	-	3	-	-	4	5	2	3	2	4	5
2002.....	89	-	1	-	1	2	6	1	2	6	4	3
2001.....	113	-	1	-	2	2	5	8	1	-	9	5
2000.....	101	1	1	-	-	2	1	2	4	3	1	6
1999.....	92	-	1	-	2	-	1	3	1	1	3	5

American Indian 1/, female

2004.....	87	-	-	1	-	-	3	-	3	4	2	1
2003.....	111	-	-	2	1	3	-	-	-	3	3	4
2002.....	82	-	-	-	1	2	1	2	2	4	-	4
2001.....	112	-	-	-	2	1	2	3	2	5	4	2
2000.....	98	-	-	1	1	1	1	-	3	1	4	3
1999.....	69	1	-	-	-	1	1	1	2	2	3	2

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)

Asian or Pacific Islander 2/, both sexes

2004.....	973	2	8	9	8	7	12	13	13	18	28	37
2003.....	946	1	6	5	15	10	13	14	20	24	25	40
2002.....	960	4	4	10	11	17	10	16	26	32	24	48
2001.....	900	1	6	11	6	7	15	22	16	18	28	46
2000.....	854	1	4	10	11	13	19	20	22	25	28	31
1999.....	880	1	3	7	8	16	3	15	27	22	29	40

Asian or Pacific Islander 2/, male

2004.....	556	2	5	4	6	5	10	6	7	10	19	24
2003.....	497	1	2	1	9	7	6	11	12	15	14	21
2002.....	504	2	2	5	8	6	6	9	14	17	11	21
2001.....	504	-	3	9	5	2	10	8	6	14	18	28
2000.....	463	1	2	7	10	9	12	11	11	15	13	19
1999.....	481	-	2	3	2	11	2	9	15	14	18	24

Asian or Pacific Islander 2/, female

2004.....	417	-	3	5	2	2	2	7	6	8	9	13
2003.....	449	-	4	4	6	3	7	3	8	9	11	19
2002.....	456	2	2	5	3	11	4	7	12	15	13	27
2001.....	396	1	3	2	1	5	5	14	10	4	10	18
2000.....	391	-	2	3	1	4	7	9	11	10	15	12
1999.....	399	1	1	4	6	5	1	6	12	8	11	16

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hodgkin's disease (C81)												
All races, both sexes												
2004.....	1,276	-	-	1	5	17	48	63	71	53	67	87
2003.....	1,347	-	-	-	5	16	48	71	96	76	78	71
2002.....	1,352	-	-	3	3	27	59	64	88	81	78	83
2001.....	1,323	-	-	1	5	25	53	76	91	81	80	91
2000.....	1,287	-	-	3	5	21	52	60	85	78	73	73
1999.....	1,403	-	1	-	9	31	63	77	71	90	75	82
All races, male												
2004.....	730	-	-	-	4	7	30	33	37	32	38	52
2003.....	725	-	-	-	4	7	23	35	53	43	52	44
2002.....	751	-	-	2	3	15	26	37	44	46	46	53
2001.....	725	-	-	1	2	15	28	39	51	50	51	54
2000.....	686	-	-	1	3	6	29	31	48	42	40	46
1999.....	785	-	1	-	3	20	35	42	44	56	47	48
All races, female												
2004.....	546	-	-	1	1	10	18	30	34	21	29	35
2003.....	622	-	-	-	1	9	25	36	43	33	26	27
2002.....	601	-	-	1	-	12	33	27	44	35	32	30
2001.....	598	-	-	-	3	10	25	37	40	31	29	37
2000.....	601	-	-	2	2	15	23	29	37	36	33	27
1999.....	618	-	-	-	6	11	28	35	27	34	28	34
White, both sexes												
2004.....	1,118	-	-	-	5	15	36	44	54	43	58	76
2003.....	1,192	-	-	-	5	15	36	56	78	65	67	64
2002.....	1,193	-	-	3	2	21	51	52	75	64	66	72
2001.....	1,170	-	-	1	4	21	41	56	69	69	69	77
2000.....	1,156	-	-	2	3	17	40	54	76	66	61	66
1999.....	1,263	-	1	-	7	26	49	60	62	80	66	72

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hodgkin's disease (C81)												
White, male												
2004.....	647	-	-	-	4	6	23	24	32	24	35	44
2003.....	638	-	-	-	4	7	17	30	44	39	45	40
2002.....	655	-	-	2	2	13	24	28	37	36	37	44
2001.....	644	-	-	1	2	13	25	27	42	41	44	46
2000.....	616	-	-	1	1	4	26	28	42	36	34	42
1999.....	708	-	1	-	3	17	28	32	40	49	43	41
White, female												
2004.....	471	-	-	-	1	9	13	20	22	19	23	32
2003.....	554	-	-	-	1	8	19	26	34	26	22	24
2002.....	538	-	-	1	-	8	27	24	38	28	29	28
2001.....	526	-	-	-	2	8	16	29	27	28	25	31
2000.....	540	-	-	1	2	13	14	26	34	30	27	24
1999.....	555	-	-	-	4	9	21	28	22	31	23	31
Black, both sexes												
2004.....	135	-	-	1	-	2	12	17	15	8	9	10
2003.....	120	-	-	-	-	-	10	15	16	9	9	6
2002.....	130	-	-	-	1	4	7	11	13	15	11	10
2001.....	137	-	-	-	1	4	11	19	21	11	10	12
2000.....	110	-	-	1	1	4	10	6	7	12	12	7
1999.....	128	-	-	-	1	5	14	17	9	10	8	10
Black, male												
2004.....	71	-	-	-	-	1	7	9	4	6	3	8
2003.....	68	-	-	-	-	-	5	5	7	4	6	3
2002.....	82	-	-	-	1	2	2	8	7	9	9	8
2001.....	73	-	-	-	-	2	2	12	9	8	7	7
2000.....	57	-	-	-	1	2	3	3	5	6	6	4
1999.....	75	-	-	-	-	3	7	10	4	7	4	7

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hodgkin's disease (C81)												
Asian or Pacific Islander 2/, both sexes												
2004.....	19	-	-	-	-	-	-	2	-	1	-	1
2003.....	26	-	-	-	-	1	2	-	2	2	1	1
2002.....	28	-	-	-	-	2	1	1	-	2	1	1
2001.....	13	-	-	-	-	-	1	1	-	1	-	2
2000.....	16	-	-	-	1	-	2	-	2	-	-	-
1999.....	7	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	9	-	-	-	-	-	-	-	-	1	-	-
2003.....	15	-	-	-	-	-	1	-	2	-	-	1
2002.....	14	-	-	-	-	-	-	1	-	1	-	1
2001.....	8	-	-	-	-	-	1	-	-	1	-	1
2000.....	9	-	-	-	1	-	-	-	1	-	-	-
1999.....	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	10	-	-	-	-	-	-	2	-	-	-	1
2003.....	11	-	-	-	-	1	1	-	-	2	1	-
2002.....	14	-	-	-	-	2	1	-	-	1	1	-
2001.....	5	-	-	-	-	-	-	1	-	-	-	1
2000.....	7	-	-	-	-	-	2	-	1	-	-	-
1999.....	6	-	-	-	1	-	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Of other lymphoid, hematopoietic and related tissue (C82-C85,C96)

All races, both sexes

2004.....	21,009	1	2	22	19	56	88	119	139	198	347	612
2003.....	21,530	1	5	17	29	53	87	96	157	233	398	653
2002.....	21,983	5	15	15	42	44	99	111	170	253	434	640
2001.....	22,395	2	2	11	27	61	87	129	170	302	458	712
2000.....	22,829	1	9	23	25	60	121	124	169	292	463	735
1999.....	22,885	3	17	12	25	51	93	109	181	326	467	699

All races, male

2004.....	11,091	-	2	15	13	37	56	85	81	130	239	406
2003.....	11,430	1	5	12	14	34	54	59	107	153	254	417
2002.....	11,582	4	6	10	30	31	63	71	107	161	278	406
2001.....	11,761	2	1	8	21	43	51	72	108	196	307	451
2000.....	11,872	-	7	16	13	46	81	64	108	193	292	470
1999.....	11,833	1	9	9	21	30	61	69	110	204	307	460

All races, female

2004.....	9,918	1	-	7	6	19	32	34	58	68	108	206
2003.....	10,100	-	-	5	15	19	33	37	50	80	144	236
2002.....	10,401	1	9	5	12	13	36	40	63	92	156	234
2001.....	10,634	-	1	3	6	18	36	57	62	106	151	261
2000.....	10,957	1	2	7	12	14	40	60	61	99	171	265
1999.....	11,052	2	8	3	4	21	32	40	71	122	160	239

White, both sexes

2004.....	19,097	1	2	18	12	46	64	85	102	159	282	489
2003.....	19,648	1	4	14	18	39	73	66	112	183	317	526
2002.....	20,057	2	14	15	30	28	79	84	119	194	346	534
2001.....	20,412	2	2	10	22	44	65	90	131	234	370	585
2000.....	20,981	-	6	18	23	46	85	89	129	227	386	606
1999.....	21,017	1	14	8	18	36	74	79	136	255	375	584

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Of other lymphoid, hematopoietic and related tissue (C82-C85,C96)

White, male

2004.....	10,077	-	2	14	8	30	39	65	59	108	200	324
2003.....	10,475	1	4	10	7	27	45	44	77	125	203	333
2002.....	10,573	1	6	10	22	23	53	58	73	123	220	351
2001.....	10,686	2	1	7	16	31	37	52	88	149	249	378
2000.....	10,889	-	6	12	13	38	58	50	81	150	253	390
1999.....	10,804	1	8	7	16	22	49	48	83	163	245	387

White, female

2004.....	9,020	1	-	4	4	16	25	20	43	51	82	165
2003.....	9,173	-	-	4	11	12	28	22	35	58	114	193
2002.....	9,484	1	8	5	8	5	26	26	46	71	126	183
2001.....	9,726	-	1	3	6	13	28	38	43	85	121	207
2000.....	10,092	-	-	6	10	8	27	39	48	77	133	216
1999.....	10,213	-	6	1	2	14	25	31	53	92	130	197

Black, both sexes

2004.....	1,422	-	-	4	4	9	21	30	29	34	51	104
2003.....	1,374	-	-	3	7	11	12	24	34	42	66	104
2002.....	1,480	1	1	-	10	11	14	22	41	40	78	90
2001.....	1,508	-	-	1	4	16	16	29	35	60	70	105
2000.....	1,378	1	2	4	1	8	28	30	26	56	65	109
1999.....	1,449	2	2	3	6	13	19	24	34	62	77	94

Black, male

2004.....	740	-	-	1	2	6	14	17	19	19	31	68
2003.....	688	-	-	2	6	4	8	10	24	22	41	71
2002.....	777	1	-	-	7	4	6	11	29	27	52	48
2001.....	811	-	-	1	4	11	11	15	19	43	46	58
2000.....	735	-	-	3	-	3	17	12	19	37	34	69
1999.....	789	-	1	2	4	6	12	17	22	36	51	59

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Of other lymphoid, hematopoietic and related tissue (C82-C85,C96)

Black, female

2004.....	682	-	-	3	2	3	7	13	10	15	20	36
2003.....	686	-	-	1	1	7	4	14	10	20	25	33
2002.....	703	-	1	-	3	7	8	11	12	13	26	42
2001.....	697	-	-	-	-	5	5	14	16	17	24	47
2000.....	643	1	2	1	1	5	11	18	7	19	31	40
1999.....	660	2	1	1	2	7	7	7	12	26	26	35

American Indian 1/, both sexes

2004.....	67	-	-	-	-	-	-	-	2	2	3	2
2003.....	69	-	1	-	-	-	-	1	1	1	2	7
2002.....	54	-	-	-	1	-	1	-	1	6	2	3
2001.....	94	-	-	-	1	1	-	3	1	3	7	3
2000.....	84	-	-	-	-	1	-	-	4	1	3	5
1999.....	57	-	-	-	1	-	-	1	2	1	5	2

American Indian 1/, male

2004.....	39	-	-	-	-	-	-	-	-	1	1	2
2003.....	34	-	1	-	-	-	-	1	1	-	2	3
2002.....	29	-	-	-	1	-	1	-	-	5	2	1
2001.....	48	-	-	-	1	1	-	2	-	-	5	3
2000.....	42	-	-	-	-	1	-	-	3	1	-	3
1999.....	31	-	-	-	1	-	-	1	-	1	3	1

American Indian 1/, female

2004.....	28	-	-	-	-	-	-	-	2	1	2	-
2003.....	35	-	-	-	-	-	-	-	-	1	-	4
2002.....	25	-	-	-	-	-	-	-	1	1	-	2
2001.....	46	-	-	-	-	-	-	1	1	3	2	-
2000.....	42	-	-	-	-	-	-	-	1	-	3	2
1999.....	26	-	-	-	-	-	-	-	2	-	2	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Of other lymphoid, hematopoietic and related tissue (C82-C85,C96)

Asian or Pacific Islander 2/, both sexes

2004.....	423	-	-	-	3	1	3	4	6	3	11	17
2003.....	439	-	-	-	4	3	2	5	10	7	13	16
2002.....	392	2	-	-	1	5	5	5	9	13	8	13
2001.....	381	-	-	-	-	-	6	7	3	5	11	19
2000.....	386	-	1	1	1	5	8	5	10	8	9	15
1999.....	362	-	1	1	-	2	-	5	9	8	10	19

Asian or Pacific Islander 2/, male

2004.....	235	-	-	-	3	1	3	3	3	2	7	12
2003.....	233	-	-	-	1	3	1	4	5	6	8	10
2002.....	203	2	-	-	-	4	3	2	5	6	4	6
2001.....	216	-	-	-	-	-	3	3	1	4	7	12
2000.....	206	-	1	1	-	4	6	2	5	5	5	8
1999.....	209	-	-	-	-	2	-	3	5	4	8	13

Asian or Pacific Islander 2/, female

2004.....	188	-	-	-	-	-	-	1	3	1	4	5
2003.....	206	-	-	-	3	-	1	1	5	1	5	6
2002.....	189	-	-	-	1	1	2	3	4	7	4	7
2001.....	165	-	-	-	-	-	3	4	2	1	4	7
2000.....	180	-	-	-	1	1	2	3	5	3	4	7
1999.....	153	-	1	1	-	-	-	2	4	4	2	6

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Non-Hodgkin's lymphoma (C82-C85)												
All races, both sexes												
2004.....	20,938	1	2	22	19	56	88	117	139	197	346	611
2003.....	21,475	-	5	17	28	53	87	96	156	233	397	649
2002.....	21,910	2	15	15	42	44	98	111	170	253	434	638
2001.....	22,305	1	1	11	27	61	87	129	169	301	455	711
2000.....	22,729	1	7	23	24	60	120	124	169	290	461	730
1999.....	22,802	2	16	12	25	51	93	109	180	324	465	696
All races, male												
2004.....	11,056	-	2	15	13	37	56	83	81	129	238	405
2003.....	11,399	-	5	12	14	34	54	59	106	153	253	414
2002.....	11,538	1	6	10	30	31	62	71	107	161	278	404
2001.....	11,710	1	1	8	21	43	51	72	107	195	305	450
2000.....	11,812	-	5	16	12	46	80	64	108	192	290	466
1999.....	11,794	-	9	9	21	30	61	69	109	204	306	459
All races, female												
2004.....	9,882	1	-	7	6	19	32	34	58	68	108	206
2003.....	10,076	-	-	5	14	19	33	37	50	80	144	235
2002.....	10,372	1	9	5	12	13	36	40	63	92	156	234
2001.....	10,595	-	-	3	6	18	36	57	62	106	150	261
2000.....	10,917	1	2	7	12	14	40	60	61	98	171	264
1999.....	11,008	2	7	3	4	21	32	40	71	120	159	237
White, both sexes												
2004.....	19,036	1	2	18	12	46	64	84	102	158	281	488
2003.....	19,598	-	4	14	17	39	73	66	111	183	316	524
2002.....	19,996	1	14	15	30	28	78	84	119	194	346	532
2001.....	20,335	1	1	10	22	44	65	90	131	234	367	584
2000.....	20,892	-	4	18	22	46	85	89	129	226	384	603
1999.....	20,941	-	13	8	18	36	74	79	135	253	374	582

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Non-Hodgkin's lymphoma (C82-C85)

White, male

2004.....	10,045	-	2	14	8	30	39	64	59	107	199	323
2003.....	10,447	-	4	10	7	27	45	44	76	125	202	332
2002.....	10,535	-	6	10	22	23	52	58	73	123	220	349
2001.....	10,642	1	1	7	16	31	37	52	88	149	247	377
2000.....	10,838	-	4	12	12	38	58	50	81	150	251	388
1999.....	10,768	-	8	7	16	22	49	48	82	163	244	386

White, female

2004.....	8,991	1	-	4	4	16	25	20	43	51	82	165
2003.....	9,151	-	-	4	10	12	28	22	35	58	114	192
2002.....	9,461	1	8	5	8	5	26	26	46	71	126	183
2001.....	9,693	-	-	3	6	13	28	38	43	85	120	207
2000.....	10,054	-	-	6	10	8	27	39	48	76	133	215
1999.....	10,173	-	5	1	2	14	25	31	53	90	130	196

Black, both sexes

2004.....	1,414	-	-	4	4	9	21	30	29	34	51	104
2003.....	1,371	-	-	3	7	11	12	24	34	42	66	103
2002.....	1,474	1	1	-	10	11	14	22	41	40	78	90
2001.....	1,496	-	-	1	4	16	16	29	34	59	70	105
2000.....	1,369	1	2	4	1	8	27	30	26	56	65	107
1999.....	1,443	2	2	3	6	13	19	24	34	62	76	93

Black, male

2004.....	739	-	-	1	2	6	14	17	19	19	31	68
2003.....	687	-	-	2	6	4	8	10	24	22	41	70
2002.....	775	1	-	-	7	4	6	11	29	27	52	48
2001.....	804	-	-	1	4	11	11	15	18	42	46	58
2000.....	728	-	-	3	-	3	16	12	19	37	34	67
1999.....	786	-	1	2	4	6	12	17	22	36	51	59

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Non-Hodgkin's lymphoma (C82-C85)												
Black, female												
2004.....	675	-	-	3	2	3	7	13	10	15	20	36
2003.....	684	-	-	1	1	7	4	14	10	20	25	33
2002.....	699	-	1	-	3	7	8	11	12	13	26	42
2001.....	692	-	-	-	-	5	5	14	16	17	24	47
2000.....	641	1	2	1	1	5	11	18	7	19	31	40
1999.....	657	2	1	1	2	7	7	7	12	26	25	34
American Indian 1/, both sexes												
2004.....	67	-	-	-	-	-	-	-	2	2	3	2
2003.....	69	-	1	-	-	-	-	1	1	1	2	7
2002.....	53	-	-	-	1	-	1	-	1	6	2	3
2001.....	93	-	-	-	1	1	-	3	1	3	7	3
2000.....	83	-	-	-	-	1	-	-	4	1	3	5
1999.....	57	-	-	-	1	-	-	1	2	1	5	2
American Indian 1/, male												
2004.....	39	-	-	-	-	-	-	-	-	1	1	2
2003.....	34	-	1	-	-	-	-	1	1	-	2	3
2002.....	29	-	-	-	1	-	1	-	-	5	2	1
2001.....	48	-	-	-	1	1	-	2	-	-	5	3
2000.....	41	-	-	-	-	1	-	-	3	1	-	3
1999.....	31	-	-	-	1	-	-	1	-	1	3	1
American Indian 1/, female												
2004.....	28	-	-	-	-	-	-	-	2	1	2	-
2003.....	35	-	-	-	-	-	-	-	-	1	-	4
2002.....	24	-	-	-	-	-	-	-	1	1	-	2
2001.....	45	-	-	-	-	-	-	1	1	3	2	-
2000.....	42	-	-	-	-	-	-	-	1	-	3	2
1999.....	26	-	-	-	-	-	-	-	2	-	2	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Non-Hodgkin's lymphoma (C82-C85)												
Asian or Pacific Islander 2/, both sexes												
2004.....	421	-	-	-	3	1	3	3	6	3	11	17
2003.....	437	-	-	-	4	3	2	5	10	7	13	15
2002.....	387	-	-	-	1	5	5	5	9	13	8	13
2001.....	381	-	-	-	-	-	6	7	3	5	11	19
2000.....	385	-	1	1	1	5	8	5	10	7	9	15
1999.....	361	-	1	1	-	2	-	5	9	8	10	19
Asian or Pacific Islander 2/, male												
2004.....	233	-	-	-	3	1	3	2	3	2	7	12
2003.....	231	-	-	-	1	3	1	4	5	6	8	9
2002.....	199	-	-	-	-	4	3	2	5	6	4	6
2001.....	216	-	-	-	-	-	3	3	1	4	7	12
2000.....	205	-	1	1	-	4	6	2	5	4	5	8
1999.....	209	-	-	-	-	2	-	3	5	4	8	13
Asian or Pacific Islander 2/, female												
2004.....	188	-	-	-	-	-	-	1	3	1	4	5
2003.....	206	-	-	-	3	-	1	1	5	1	5	6
2002.....	188	-	-	-	1	1	2	3	4	7	4	7
2001.....	165	-	-	-	-	-	3	4	2	1	4	7
2000.....	180	-	-	-	1	1	2	3	5	3	4	7
1999.....	152	-	1	1	-	-	-	2	4	4	2	6

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Of other and unspecified lymphoid, hematopoietic and related tissue (C96)

All races, both sexes

2004.....	71	-	-	-	-	-	-	2	-	1	1	1
2003.....	55	1	-	-	1	-	-	-	1	-	1	4
2002.....	73	3	-	-	-	-	1	-	-	-	-	2
2001.....	90	1	1	-	-	-	-	-	1	1	3	1
2000.....	100	-	2	-	1	-	1	-	-	2	2	5
1999.....	83	1	1	-	-	-	-	-	1	2	2	3

All races, male

2004.....	35	-	-	-	-	-	-	2	-	1	1	1
2003.....	31	1	-	-	-	-	-	-	1	-	1	3
2002.....	44	3	-	-	-	-	1	-	-	-	-	2
2001.....	51	1	-	-	-	-	-	-	1	1	2	1
2000.....	60	-	2	-	1	-	1	-	-	1	2	4
1999.....	39	1	-	-	-	-	-	-	1	-	1	1

All races, female

2004.....	36	-	-	-	-	-	-	-	-	-	-	-
2003.....	24	-	-	-	1	-	-	-	-	-	-	1
2002.....	29	-	-	-	-	-	-	-	-	-	-	-
2001.....	39	-	1	-	-	-	-	-	-	-	1	-
2000.....	40	-	-	-	-	-	-	-	-	1	-	1
1999.....	44	-	1	-	-	-	-	-	-	2	1	2

White, both sexes

2004.....	61	-	-	-	-	-	-	1	-	1	1	1
2003.....	50	1	-	-	1	-	-	-	1	-	1	2
2002.....	61	1	-	-	-	-	1	-	-	-	-	2
2001.....	77	1	1	-	-	-	-	-	-	-	3	1
2000.....	89	-	2	-	1	-	-	-	-	1	2	3
1999.....	76	1	1	-	-	-	-	-	1	2	1	2

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Malignant immunoproliferative diseases (C88)												
White, male												
2004.....	171	1	-	-	-	-	-	-	-	1	1	1
2003.....	221	-	-	-	-	-	-	-	-	-	-	2
2002.....	227	-	-	-	-	-	-	-	-	-	1	1
2001.....	189	-	-	-	-	-	-	-	-	-	1	1
2000.....	196	-	-	-	-	-	-	-	-	-	-	4
1999.....	228	-	-	-	-	-	-	-	-	-	-	3
White, female												
2004.....	131	-	-	-	-	-	-	-	-	1	-	2
2003.....	145	-	-	-	-	-	-	-	-	-	-	1
2002.....	148	-	-	-	-	-	-	-	-	-	-	2
2001.....	166	-	-	-	-	-	-	-	1	1	-	-
2000.....	150	-	-	-	-	-	-	-	-	1	-	-
1999.....	193	-	-	-	-	-	-	-	-	-	-	3
Black, both sexes												
2004.....	7	-	-	-	-	-	-	-	-	-	-	-
2003.....	18	-	-	-	-	-	-	-	-	-	-	1
2002.....	17	-	-	-	-	-	-	-	-	-	-	1
2001.....	17	-	-	-	-	-	-	-	-	-	-	1
2000.....	18	-	-	-	-	-	-	-	-	-	-	-
1999.....	23	-	-	-	-	-	-	-	1	1	-	-
Black, male												
2004.....	7	-	-	-	-	-	-	-	-	-	-	-
2003.....	8	-	-	-	-	-	-	-	-	-	-	1
2002.....	9	-	-	-	-	-	-	-	-	-	-	-
2001.....	7	-	-	-	-	-	-	-	-	-	-	-
2000.....	10	-	-	-	-	-	-	-	-	-	-	-
1999.....	13	-	-	-	-	-	-	-	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Multiple myeloma and malignant plasma cell neoplasms (C90)

All races, both sexes

2004.....	10,655	-	-	-	-	1	-	1	13	29	104	220
2003.....	10,882	-	-	-	1	-	-	1	11	25	122	234
2002.....	10,996	2	-	-	-	2	-	3	9	36	101	266
2001.....	10,795	-	-	-	-	-	-	1	7	38	116	239
2000.....	10,697	-	-	-	1	-	-	-	7	37	108	241
1999.....	10,565	-	-	-	-	-	-	3	18	46	109	235

All races, male

2004.....	5,516	-	-	-	-	1	-	1	10	15	61	119
2003.....	5,619	-	-	-	-	-	-	1	6	17	80	136
2002.....	5,713	-	-	-	-	-	-	-	6	24	53	158
2001.....	5,524	-	-	-	-	-	-	-	4	22	73	150
2000.....	5,347	-	-	-	-	-	-	-	5	18	67	139
1999.....	5,338	-	-	-	-	-	-	2	13	33	62	129

All races, female

2004.....	5,139	-	-	-	-	-	-	-	3	14	43	101
2003.....	5,263	-	-	-	1	-	-	-	5	8	42	98
2002.....	5,283	2	-	-	-	2	-	3	3	12	48	108
2001.....	5,271	-	-	-	-	-	-	1	3	16	43	89
2000.....	5,350	-	-	-	1	-	-	-	2	19	41	102
1999.....	5,227	-	-	-	-	-	-	1	5	13	47	106

White, both sexes

2004.....	8,646	-	-	-	-	1	-	1	10	19	73	159
2003.....	8,861	-	-	-	1	-	-	1	8	17	86	155
2002.....	9,045	1	-	-	-	2	-	2	8	22	68	186
2001.....	8,813	-	-	-	-	-	-	1	3	29	69	170
2000.....	8,720	-	-	-	1	-	-	-	4	23	73	164
1999.....	8,589	-	-	-	-	-	-	2	11	35	68	176

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Multiple myeloma and malignant plasma cell neoplasms (C90)												
White, male												
2004.....	4,556	-	-	-	-	1	-	1	8	11	45	88
2003.....	4,671	-	-	-	-	-	-	1	5	13	60	94
2002.....	4,797	-	-	-	-	-	-	-	5	15	38	115
2001.....	4,576	-	-	-	-	-	-	-	3	15	51	109
2000.....	4,448	-	-	-	-	-	-	-	3	14	50	99
1999.....	4,423	-	-	-	-	-	-	2	7	26	38	99
White, female												
2004.....	4,090	-	-	-	-	-	-	-	2	8	28	71
2003.....	4,190	-	-	-	1	-	-	-	3	4	26	61
2002.....	4,248	1	-	-	-	2	-	2	3	7	30	71
2001.....	4,237	-	-	-	-	-	-	1	-	14	18	61
2000.....	4,272	-	-	-	1	-	-	-	1	9	23	65
1999.....	4,166	-	-	-	-	-	-	-	4	9	30	77
Black, both sexes												
2004.....	1,809	-	-	-	-	-	-	-	2	10	30	55
2003.....	1,829	-	-	-	-	-	-	-	2	8	35	72
2002.....	1,754	-	-	-	-	-	-	1	1	13	28	74
2001.....	1,800	-	-	-	-	-	-	-	4	9	44	64
2000.....	1,823	-	-	-	-	-	-	-	3	13	31	72
1999.....	1,780	-	-	-	-	-	-	1	6	9	40	56
Black, male												
2004.....	852	-	-	-	-	-	-	-	1	4	16	27
2003.....	855	-	-	-	-	-	-	-	1	4	20	38
2002.....	831	-	-	-	-	-	-	-	1	8	14	42
2001.....	857	-	-	-	-	-	-	-	1	7	21	39
2000.....	827	-	-	-	-	-	-	-	2	3	14	36
1999.....	821	-	-	-	-	-	-	-	6	6	24	28

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Multiple myeloma and malignant plasma cell neoplasms (C90)												
Black, female												
2004.....	957	-	-	-	-	-	-	-	1	6	14	28
2003.....	974	-	-	-	-	-	-	-	1	4	15	34
2002.....	923	-	-	-	-	-	-	1	-	5	14	32
2001.....	943	-	-	-	-	-	-	-	3	2	23	25
2000.....	996	-	-	-	-	-	-	-	1	10	17	36
1999.....	959	-	-	-	-	-	-	1	-	3	16	28
American Indian 1/, both sexes												
2004.....	52	-	-	-	-	-	-	-	-	-	-	4
2003.....	54	-	-	-	-	-	-	-	-	-	1	-
2002.....	42	-	-	-	-	-	-	-	-	1	1	1
2001.....	49	-	-	-	-	-	-	-	-	-	2	2
2000.....	37	-	-	-	-	-	-	-	-	-	1	2
1999.....	43	-	-	-	-	-	-	-	-	1	-	2
American Indian 1/, male												
2004.....	28	-	-	-	-	-	-	-	-	-	-	3
2003.....	23	-	-	-	-	-	-	-	-	-	-	-
2002.....	19	-	-	-	-	-	-	-	-	1	1	-
2001.....	21	-	-	-	-	-	-	-	-	-	1	-
2000.....	17	-	-	-	-	-	-	-	-	-	1	2
1999.....	22	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, female												
2004.....	24	-	-	-	-	-	-	-	-	-	-	1
2003.....	31	-	-	-	-	-	-	-	-	-	1	-
2002.....	23	-	-	-	-	-	-	-	-	-	-	1
2001.....	28	-	-	-	-	-	-	-	-	-	1	2
2000.....	20	-	-	-	-	-	-	-	-	-	-	-
1999.....	21	-	-	-	-	-	-	-	-	1	-	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Multiple myeloma and malignant plasma cell neoplasms (C90)

Asian or Pacific Islander 2/, both sexes

2004.....	148	-	-	-	-	-	-	-	1	-	1	2
2003.....	138	-	-	-	-	-	-	-	1	-	-	7
2002.....	155	1	-	-	-	-	-	-	-	-	4	5
2001.....	133	-	-	-	-	-	-	-	-	-	1	3
2000.....	117	-	-	-	-	-	-	-	-	1	3	3
1999.....	153	-	-	-	-	-	-	-	1	1	1	1

Asian or Pacific Islander 2/, male

2004.....	80	-	-	-	-	-	-	-	1	-	-	1
2003.....	70	-	-	-	-	-	-	-	-	-	-	4
2002.....	66	-	-	-	-	-	-	-	-	-	-	1
2001.....	70	-	-	-	-	-	-	-	-	-	-	2
2000.....	55	-	-	-	-	-	-	-	-	1	2	2
1999.....	72	-	-	-	-	-	-	-	-	1	-	1

Asian or Pacific Islander 2/, female

2004.....	68	-	-	-	-	-	-	-	-	-	1	1
2003.....	68	-	-	-	-	-	-	-	1	-	-	3
2002.....	89	1	-	-	-	-	-	-	-	-	4	4
2001.....	63	-	-	-	-	-	-	-	-	-	1	1
2000.....	62	-	-	-	-	-	-	-	-	-	1	1
1999.....	81	-	-	-	-	-	-	-	1	-	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Leukemia (C91-C95)

All races, both sexes

2004.....	21,395	23	120	156	156	210	225	173	243	287	426	555
2003.....	21,535	30	133	156	186	183	219	205	228	321	475	595
2002.....	21,498	21	104	148	168	237	235	190	281	325	442	606
2001.....	21,451	25	155	153	170	225	254	230	283	346	433	630
2000.....	21,339	38	137	142	173	242	222	228	245	368	433	622
1999.....	21,014	27	119	130	175	221	261	234	266	374	425	626

All races, male

2004.....	12,007	9	70	96	86	126	142	95	147	159	239	303
2003.....	12,073	9	75	90	105	125	132	125	149	167	277	347
2002.....	12,016	11	57	75	97	142	140	117	163	178	251	335
2001.....	11,850	10	81	86	99	134	155	135	165	196	236	346
2000.....	11,777	21	69	81	109	150	143	131	152	185	238	356
1999.....	11,519	11	65	75	98	130	161	128	148	204	231	346

All races, female

2004.....	9,388	14	50	60	70	84	83	78	96	128	187	252
2003.....	9,462	21	58	66	81	58	87	80	79	154	198	248
2002.....	9,482	10	47	73	71	95	95	73	118	147	191	271
2001.....	9,601	15	74	67	71	91	99	95	118	150	197	284
2000.....	9,562	17	68	61	64	92	79	97	93	183	195	266
1999.....	9,495	16	54	55	77	91	100	106	118	170	194	280

White, both sexes

2004.....	19,077	18	92	127	127	168	185	135	195	218	345	460
2003.....	19,353	28	111	132	144	147	173	169	180	244	387	510
2002.....	19,273	16	84	119	134	193	201	149	221	262	382	481
2001.....	19,239	20	134	114	130	188	203	171	231	279	349	531
2000.....	19,191	36	115	116	143	186	175	172	183	300	348	500
1999.....	18,899	23	103	106	132	170	209	191	210	307	353	524

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Leukemia (C91-C95)												
White, male												
2004.....	10,753	7	50	78	71	104	116	78	119	122	194	252
2003.....	10,938	7	63	82	81	102	103	100	118	123	228	306
2002.....	10,835	10	47	57	77	119	120	94	131	142	220	263
2001.....	10,672	7	68	64	67	115	125	103	136	155	187	299
2000.....	10,617	19	58	64	89	113	112	102	114	154	199	292
1999.....	10,395	11	54	63	71	97	136	104	115	170	192	286
White, female												
2004.....	8,324	11	42	49	56	64	69	57	76	96	151	208
2003.....	8,415	21	48	50	63	45	70	69	62	121	159	204
2002.....	8,438	6	37	62	57	74	81	55	90	120	162	218
2001.....	8,567	13	66	50	63	73	78	68	95	124	162	232
2000.....	8,574	17	57	52	54	73	63	70	69	146	149	208
1999.....	8,504	12	49	43	61	73	73	87	95	137	161	238
Black, both sexes												
2004.....	1,860	3	19	17	23	35	27	30	41	50	64	77
2003.....	1,762	1	14	17	30	23	32	26	39	58	74	67
2002.....	1,770	4	15	19	23	30	24	28	40	43	48	93
2001.....	1,765	4	14	28	31	28	36	37	38	53	66	75
2000.....	1,742	-	18	16	20	46	36	39	49	49	68	107
1999.....	1,708	2	13	18	35	36	47	30	38	53	54	79
Black, male												
2004.....	980	-	14	12	11	17	18	13	24	28	32	39
2003.....	918	1	8	7	16	15	20	17	24	33	42	33
2002.....	923	1	7	13	12	19	12	16	21	26	23	57
2001.....	928	3	9	13	26	16	19	21	23	32	36	32
2000.....	929	-	9	11	11	31	24	18	32	20	33	54
1999.....	892	-	8	9	24	24	22	16	22	25	29	47

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Leukemia (C91-C95)												
Black, female												
2004.....	880	3	5	5	12	18	9	17	17	22	32	38
2003.....	844	-	6	10	14	8	12	9	15	25	32	34
2002.....	847	3	8	6	11	11	12	12	19	17	25	36
2001.....	837	1	5	15	5	12	17	16	15	21	30	43
2000.....	813	-	9	5	9	15	12	21	17	29	35	53
1999.....	816	2	5	9	11	12	25	14	16	28	25	32
American Indian 1/, both sexes												
2004.....	76	-	1	3	1	1	4	1	1	5	1	1
2003.....	77	-	2	2	1	7	5	1	2	4	3	2
2002.....	73	-	1	-	1	4	6	3	3	3	1	3
2001.....	77	-	1	-	3	2	7	8	1	2	3	2
2000.....	72	1	1	1	1	2	2	2	3	3	1	2
1999.....	56	1	1	-	1	1	2	3	1	1	-	3
American Indian 1/, male												
2004.....	42	-	1	2	1	1	1	1	1	2	1	1
2003.....	38	-	2	-	-	4	5	1	2	2	1	2
2002.....	40	-	1	-	-	2	5	1	2	-	1	2
2001.....	42	-	1	-	1	1	5	6	1	-	3	2
2000.....	38	1	1	-	-	1	1	2	1	2	-	1
1999.....	38	-	1	-	1	-	1	2	1	-	-	3
American Indian 1/, female												
2004.....	34	-	-	1	-	-	3	-	-	3	-	-
2003.....	39	-	-	2	1	3	-	-	-	2	2	-
2002.....	33	-	-	-	1	2	1	2	1	3	-	1
2001.....	35	-	-	-	2	1	2	2	-	2	-	-
2000.....	34	-	-	1	1	1	1	-	2	1	1	1
1999.....	18	1	-	-	-	1	1	1	-	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Leukemia (C91-C95)												
Asian or Pacific Islander 2/, both sexes												
2004.....	382	2	8	9	5	6	9	7	6	14	16	17
2003.....	343	1	6	5	11	6	9	9	7	15	11	16
2002.....	382	1	4	10	10	10	4	10	17	17	11	29
2001.....	370	1	6	11	6	7	8	14	13	12	15	22
2000.....	334	1	3	9	9	8	9	15	10	16	16	13
1999.....	351	1	2	6	7	14	3	10	17	13	18	20
Asian or Pacific Islander 2/, male												
2004.....	232	2	5	4	3	4	7	3	3	7	12	11
2003.....	179	1	2	1	8	4	4	7	5	9	6	6
2002.....	218	-	2	5	8	2	3	6	9	10	7	13
2001.....	208	-	3	9	5	2	6	5	5	9	10	13
2000.....	193	1	1	6	9	5	6	9	5	9	6	9
1999.....	194	-	2	3	2	9	2	6	10	9	10	10
Asian or Pacific Islander 2/, female												
2004.....	150	-	3	5	2	2	2	4	3	7	4	6
2003.....	164	-	4	4	3	2	5	2	2	6	5	10
2002.....	164	1	2	5	2	8	1	4	8	7	4	16
2001.....	162	1	3	2	1	5	2	9	8	3	5	9
2000.....	141	-	2	3	-	3	3	6	5	7	10	4
1999.....	157	1	-	3	5	5	1	4	7	4	8	10

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	-------------	-----------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Lymphoid leukemia (C91)

All races, both sexes

2004.....	6,131	5	44	89	73	96	90	60	66	70	90	121
2003.....	6,301	11	61	88	93	85	98	67	59	75	99	139
2002.....	6,271	11	34	83	93	108	108	55	81	88	105	140
2001.....	6,234	5	42	85	81	105	106	69	78	78	88	126
2000.....	6,167	16	49	80	95	103	84	58	50	66	79	128
1999.....	6,252	10	46	65	79	94	99	69	67	78	76	116

All races, male

2004.....	3,568	2	29	54	41	59	64	39	46	41	56	72
2003.....	3,713	2	34	55	64	59	69	41	45	43	64	86
2002.....	3,613	7	17	41	54	71	66	37	45	55	72	81
2001.....	3,554	3	21	45	53	74	72	37	48	50	51	75
2000.....	3,525	9	23	47	65	68	63	36	36	42	52	79
1999.....	3,597	4	26	41	45	65	64	44	43	46	47	71

All races, female

2004.....	2,563	3	15	35	32	37	26	21	20	29	34	49
2003.....	2,588	9	27	33	29	26	29	26	14	32	35	53
2002.....	2,658	4	17	42	39	37	42	18	36	33	33	59
2001.....	2,680	2	21	40	28	31	34	32	30	28	37	51
2000.....	2,642	7	26	33	30	35	21	22	14	24	27	49
1999.....	2,655	6	20	24	34	29	35	25	24	32	29	45

White, both sexes

2004.....	5,493	5	35	71	64	89	72	47	55	52	72	101
2003.....	5,714	11	53	82	72	73	77	56	48	60	84	116
2002.....	5,674	9	30	69	82	88	92	45	66	76	92	119
2001.....	5,608	5	36	67	62	88	91	53	66	62	74	101
2000.....	5,616	15	42	68	84	86	69	51	40	53	70	103
1999.....	5,676	7	43	58	59	76	83	59	58	62	66	98

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Lymphoid leukemia (C91)

White, male

2004.....	3,187	2	22	43	35	55	49	32	39	30	45	60
2003.....	3,380	2	30	53	51	52	52	34	36	36	52	74
2002.....	3,285	6	15	33	45	61	57	31	38	47	65	72
2001.....	3,207	3	18	36	38	62	64	30	40	40	43	63
2000.....	3,200	8	20	41	57	56	52	33	29	35	49	63
1999.....	3,266	4	24	37	32	50	57	38	37	39	38	60

White, female

2004.....	2,306	3	13	28	29	34	23	15	16	22	27	41
2003.....	2,334	9	23	29	21	21	25	22	12	24	32	42
2002.....	2,389	3	15	36	37	27	35	14	28	29	27	47
2001.....	2,401	2	18	31	24	26	27	23	26	22	31	38
2000.....	2,416	7	22	27	27	30	17	18	11	18	21	40
1999.....	2,410	3	19	21	27	26	26	21	21	23	28	38

Black, both sexes

2004.....	532	-	4	8	5	5	10	11	8	11	13	19
2003.....	497	-	5	4	13	6	15	8	9	10	15	16
2002.....	506	2	3	7	7	12	11	6	9	8	11	18
2001.....	540	-	4	12	16	13	11	10	7	10	7	21
2000.....	474	-	6	8	8	14	10	5	10	7	7	24
1999.....	507	1	2	6	15	15	13	8	7	14	8	14

Black, male

2004.....	323	-	4	6	3	3	9	6	5	7	7	11
2003.....	282	-	3	1	9	3	12	6	7	6	12	10
2002.....	280	1	2	5	6	7	5	4	5	4	6	9
2001.....	299	-	3	4	13	9	4	5	5	6	3	10
2000.....	276	-	2	4	5	10	7	1	7	2	3	15
1999.....	293	-	1	4	10	13	6	5	5	6	7	9

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Lymphoid leukemia (C91)												
Black, female												
2004.....	209	-	-	2	2	2	1	5	3	4	6	8
2003.....	215	-	2	3	4	3	3	2	2	4	3	6
2002.....	226	1	1	2	1	5	6	2	4	4	5	9
2001.....	241	-	1	8	3	4	7	5	2	4	4	11
2000.....	198	-	4	4	3	4	3	4	3	5	4	9
1999.....	214	1	1	2	5	2	7	3	2	8	1	5
American Indian 1/, both sexes												
2004.....	27	-	1	3	1	-	3	1	-	1	-	-
2003.....	25	-	1	1	1	4	5	-	-	-	-	-
2002.....	20	-	-	-	1	3	3	1	2	-	-	-
2001.....	23	-	-	-	2	1	4	3	-	2	2	1
2000.....	19	-	-	1	-	-	1	-	-	1	-	-
1999.....	18	1	1	-	1	1	1	1	-	1	-	2
American Indian 1/, male												
2004.....	14	-	1	2	1	-	1	1	-	1	-	-
2003.....	16	-	1	-	-	3	5	-	-	-	-	-
2002.....	10	-	-	-	-	2	3	-	1	-	-	-
2001.....	13	-	-	-	1	1	4	1	-	-	2	1
2000.....	11	-	-	-	-	-	1	-	-	1	-	-
1999.....	12	-	1	-	1	-	-	1	-	-	-	2
American Indian 1/, female												
2004.....	13	-	-	1	-	-	2	-	-	-	-	-
2003.....	9	-	-	1	1	1	-	-	-	-	-	-
2002.....	10	-	-	-	1	1	-	1	1	-	-	-
2001.....	10	-	-	-	1	-	-	2	-	2	-	-
2000.....	8	-	-	1	-	-	-	-	-	-	-	-
1999.....	6	1	-	-	-	1	1	-	-	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Lymphoid leukemia (C91)												
Asian or Pacific Islander 2/, both sexes												
2004.....	79	-	4	7	3	2	5	1	3	6	5	1
2003.....	65	-	2	1	7	2	1	3	2	5	-	7
2002.....	71	-	1	7	3	5	2	3	4	4	2	3
2001.....	63	-	2	6	1	3	-	3	5	4	5	3
2000.....	58	1	1	3	3	3	4	2	-	5	2	1
1999.....	51	1	-	1	4	2	2	1	2	1	2	2
Asian or Pacific Islander 2/, male												
2004.....	44	-	2	3	2	1	5	-	2	3	4	1
2003.....	35	-	-	1	4	1	-	1	2	1	-	2
2002.....	38	-	-	3	3	1	1	2	1	4	1	-
2001.....	35	-	-	5	1	2	-	1	3	4	3	1
2000.....	38	1	1	2	3	2	3	2	-	4	-	1
1999.....	26	-	-	-	2	2	1	-	1	1	2	-
Asian or Pacific Islander 2/, female												
2004.....	35	-	2	4	1	1	-	1	1	3	1	-
2003.....	30	-	2	-	3	1	1	2	-	4	-	5
2002.....	33	-	1	4	-	4	1	1	3	-	1	3
2001.....	28	-	2	1	-	1	-	2	2	-	2	2
2000.....	20	-	-	1	-	1	1	-	-	1	2	-
1999.....	25	1	-	1	2	-	1	1	1	-	-	2

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	-------------	-----------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Myeloid leukemia (C92)

All races, both sexes

2004.....	9,949	7	56	42	55	75	100	92	130	165	243	345
2003.....	9,893	9	42	40	66	75	84	103	126	181	285	336
2002.....	9,900	7	49	35	50	83	85	98	150	185	253	347
2001.....	9,938	8	77	37	50	79	104	114	156	212	275	387
2000.....	9,764	9	55	34	55	93	94	114	156	240	267	360
1999.....	9,284	8	54	32	63	76	112	124	142	220	267	388

All races, male

2004.....	5,496	2	32	27	27	44	53	43	68	88	132	172
2003.....	5,502	4	24	21	29	48	43	61	78	91	158	190
2002.....	5,533	3	32	15	27	42	45	58	88	97	132	188
2001.....	5,423	3	39	23	23	40	54	68	88	112	143	201
2000.....	5,377	8	30	18	31	57	56	60	91	106	135	199
1999.....	5,019	3	29	14	33	38	65	62	76	118	133	203

All races, female

2004.....	4,453	5	24	15	28	31	47	49	62	77	111	173
2003.....	4,391	5	18	19	37	27	41	42	48	90	127	146
2002.....	4,367	4	17	20	23	41	40	40	62	88	121	159
2001.....	4,515	5	38	14	27	39	50	46	68	100	132	186
2000.....	4,387	1	25	16	24	36	38	54	65	134	132	161
1999.....	4,265	5	25	18	30	38	47	62	66	102	134	185

White, both sexes

2004.....	8,873	3	42	35	40	53	81	72	102	123	205	283
2003.....	8,854	8	32	29	51	51	67	85	98	135	231	287
2002.....	8,856	6	35	27	38	65	70	72	114	142	221	280
2001.....	8,901	6	65	26	36	67	81	88	127	170	223	329
2000.....	8,717	9	44	25	41	65	70	81	110	196	208	292
1999.....	8,303	8	43	21	51	53	84	99	109	185	215	326

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Myeloid leukemia (C92)

White, male

2004.....	4,938	1	22	23	20	33	44	34	53	67	113	143
2003.....	4,971	3	19	18	21	32	33	47	61	61	131	170
2002.....	4,974	3	24	11	21	33	37	44	68	73	116	148
2001.....	4,881	2	31	16	13	36	40	50	72	87	115	173
2000.....	4,844	8	25	10	23	39	41	44	65	89	108	167
1999.....	4,500	3	21	9	28	27	50	49	56	100	108	168

White, female

2004.....	3,935	2	20	12	20	20	37	38	49	56	92	140
2003.....	3,883	5	13	11	30	19	34	38	37	74	100	117
2002.....	3,882	3	11	16	17	32	33	28	46	69	105	132
2001.....	4,020	4	34	10	23	31	41	38	55	83	108	156
2000.....	3,873	1	19	15	18	26	29	37	45	107	100	125
1999.....	3,803	5	22	12	23	26	34	50	53	85	107	158

Black, both sexes

2004.....	838	3	13	5	13	19	14	16	25	31	31	46
2003.....	815	-	6	8	14	17	13	13	22	33	43	41
2002.....	788	1	10	7	10	13	11	19	24	29	24	48
2001.....	785	1	8	9	10	10	16	16	24	35	45	40
2000.....	834	-	9	7	9	24	20	24	34	33	49	60
1999.....	755	-	9	10	10	16	27	18	22	27	39	47

Black, male

2004.....	412	-	9	3	6	9	7	7	13	17	15	19
2003.....	422	-	3	3	7	12	7	9	13	21	21	16
2002.....	412	-	5	4	4	8	5	10	13	19	10	30
2001.....	404	1	5	5	8	4	9	10	15	20	23	16
2000.....	419	-	4	6	4	16	13	13	21	13	24	27
1999.....	395	-	6	4	5	8	14	9	14	13	18	27

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Myeloid leukemia (C92)												
Black, female												
2004.....	426	3	4	2	7	10	7	9	12	14	16	27
2003.....	393	-	3	5	7	5	6	4	9	12	22	25
2002.....	376	1	5	3	6	5	6	9	11	10	14	18
2001.....	381	-	3	4	2	6	7	6	9	15	22	24
2000.....	415	-	5	1	5	8	7	11	13	20	25	33
1999.....	360	-	3	6	5	8	13	9	8	14	21	20
American Indian 1/, both sexes												
2004.....	31	-	-	-	-	1	1	-	1	4	-	1
2003.....	37	-	1	-	-	3	-	1	2	4	2	2
2002.....	38	-	1	-	-	1	3	2	1	2	-	2
2001.....	32	-	1	-	1	-	2	4	1	-	-	1
2000.....	40	-	1	-	1	1	1	-	2	2	-	1
1999.....	26	-	-	-	-	-	1	2	-	-	-	1
American Indian 1/, male												
2004.....	17	-	-	-	-	1	-	-	1	1	-	1
2003.....	18	-	1	-	-	1	-	1	2	2	1	2
2002.....	21	-	1	-	-	-	2	1	1	-	-	1
2001.....	16	-	1	-	-	-	-	4	1	-	-	1
2000.....	19	-	1	-	-	1	-	-	-	1	-	-
1999.....	17	-	-	-	-	-	1	1	-	-	-	1
American Indian 1/, female												
2004.....	14	-	-	-	-	-	1	-	-	3	-	-
2003.....	19	-	-	-	-	2	-	-	-	2	1	-
2002.....	17	-	-	-	-	1	1	1	-	2	-	1
2001.....	16	-	-	-	1	-	2	-	-	-	-	-
2000.....	21	-	-	-	1	-	1	-	2	1	-	1
1999.....	9	-	-	-	-	-	-	1	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Myeloid leukemia (C92)

Asian or Pacific Islander 2/, both sexes

2004.....	207	1	1	2	2	2	4	4	2	7	7	15
2003.....	187	1	3	3	1	4	4	4	4	9	9	6
2002.....	218	-	3	1	2	4	1	5	11	12	8	17
2001.....	220	1	3	2	3	2	5	6	4	7	7	17
2000.....	173	-	1	2	4	3	3	9	10	9	10	7
1999.....	200	-	2	1	2	7	-	5	11	8	13	14

Asian or Pacific Islander 2/, male

2004.....	129	1	1	1	1	1	2	2	1	3	4	9
2003.....	91	1	1	-	1	3	3	4	2	7	5	2
2002.....	126	-	2	-	2	1	1	3	6	5	6	9
2001.....	122	-	2	2	2	-	5	4	-	5	5	11
2000.....	95	-	-	2	4	1	2	3	5	3	3	5
1999.....	107	-	2	1	-	3	-	3	6	5	7	7

Asian or Pacific Islander 2/, female

2004.....	78	-	-	1	1	1	2	2	1	4	3	6
2003.....	96	-	2	3	-	1	1	-	2	2	4	4
2002.....	92	-	1	1	-	3	-	2	5	7	2	8
2001.....	98	1	1	-	1	2	-	2	4	2	2	6
2000.....	78	-	1	-	-	2	1	6	5	6	7	2
1999.....	93	-	-	-	2	4	-	2	5	3	6	7

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Monocytic leukemia (C93)												
All races, both sexes												
2004.....	121	-	-	-	-	-	2	-	-	1	-	2
2003.....	144	-	-	-	-	1	-	-	2	3	5	1
2002.....	146	-	4	-	-	1	-	1	-	2	3	4
2001.....	152	1	-	2	-	2	2	1	2	1	1	3
2000.....	149	-	3	-	1	1	1	-	2	2	5	4
1999.....	179	1	1	-	1	1	3	1	1	2	2	6
All races, male												
2004.....	79	-	-	-	-	-	2	-	-	-	-	2
2003.....	82	-	-	-	-	1	-	-	1	2	2	1
2002.....	87	-	1	-	-	1	-	1	-	1	-	3
2001.....	79	-	-	-	-	1	1	1	-	1	1	1
2000.....	93	-	2	-	1	1	1	-	1	2	5	1
1999.....	95	-	-	-	1	-	2	1	-	1	2	2
All races, female												
2004.....	42	-	-	-	-	-	-	-	-	1	-	-
2003.....	62	-	-	-	-	-	-	-	1	1	3	-
2002.....	59	-	3	-	-	-	-	-	-	1	3	1
2001.....	73	1	-	2	-	1	1	-	2	-	-	2
2000.....	56	-	1	-	-	-	-	-	1	-	-	3
1999.....	84	1	1	-	-	1	1	-	1	1	-	4
White, both sexes												
2004.....	112	-	-	-	-	-	2	-	-	1	-	2
2003.....	138	-	-	-	-	1	-	-	2	3	4	1
2002.....	134	-	4	-	-	1	-	-	-	2	2	2
2001.....	143	1	-	1	-	2	-	-	2	1	1	2
2000.....	135	-	2	-	1	1	1	-	-	2	5	2
1999.....	162	-	1	-	-	1	3	-	1	2	2	3

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Monocytic leukemia (C93)												
White, male												
2004.....	72	-	-	-	-	-	2	-	-	-	-	2
2003.....	80	-	-	-	-	1	-	-	1	2	2	1
2002.....	80	-	1	-	-	1	-	-	-	1	-	1
2001.....	76	-	-	-	-	1	-	-	-	1	1	1
2000.....	86	-	1	-	1	1	1	-	-	2	5	1
1999.....	86	-	-	-	-	-	2	-	-	1	2	-
White, female												
2004.....	40	-	-	-	-	-	-	-	-	1	-	-
2003.....	58	-	-	-	-	-	-	-	1	1	2	-
2002.....	54	-	3	-	-	-	-	-	-	1	2	1
2001.....	67	1	-	1	-	1	-	-	2	-	-	1
2000.....	49	-	1	-	-	-	-	-	-	-	-	1
1999.....	76	-	1	-	-	1	1	-	1	1	-	3
Black, both sexes												
2004.....	3	-	-	-	-	-	-	-	-	-	-	-
2003.....	6	-	-	-	-	-	-	-	-	-	1	-
2002.....	12	-	-	-	-	-	-	1	-	-	1	2
2001.....	9	-	-	1	-	-	2	1	-	-	-	1
2000.....	10	-	1	-	-	-	-	-	1	-	-	2
1999.....	14	1	-	-	1	-	-	1	-	-	-	2
Black, male												
2004.....	2	-	-	-	-	-	-	-	-	-	-	-
2003.....	2	-	-	-	-	-	-	-	-	-	-	-
2002.....	7	-	-	-	-	-	-	1	-	-	-	2
2001.....	3	-	-	-	-	-	1	1	-	-	-	-
2000.....	4	-	1	-	-	-	-	-	-	-	-	-
1999.....	6	-	-	-	1	-	-	1	-	-	-	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other and unspecified leukemia (C94-C95)

All races, both sexes

2004.....	5,194	11	20	25	28	39	33	21	47	51	93	87
2003.....	5,197	10	30	28	27	22	37	35	41	62	86	119
2002.....	5,181	3	17	30	25	45	42	36	50	50	81	115
2001.....	5,127	11	36	29	39	39	42	46	47	55	69	114
2000.....	5,259	13	30	28	22	45	43	56	37	60	82	130
1999.....	5,299	8	18	33	32	50	47	40	56	74	80	116

All races, male

2004.....	2,864	5	9	15	18	23	23	13	33	30	51	57
2003.....	2,776	3	17	14	12	17	20	23	25	31	53	70
2002.....	2,783	1	7	19	16	28	29	21	30	25	47	63
2001.....	2,794	4	21	18	23	19	28	29	29	33	41	69
2000.....	2,782	4	14	16	12	24	23	35	24	35	46	77
1999.....	2,808	4	10	20	19	27	30	21	29	39	49	70

All races, female

2004.....	2,330	6	11	10	10	16	10	8	14	21	42	30
2003.....	2,421	7	13	14	15	5	17	12	16	31	33	49
2002.....	2,398	2	10	11	9	17	13	15	20	25	34	52
2001.....	2,333	7	15	11	16	20	14	17	18	22	28	45
2000.....	2,477	9	16	12	10	21	20	21	13	25	36	53
1999.....	2,491	4	8	13	13	23	17	19	27	35	31	46

White, both sexes

2004.....	4,599	10	15	21	23	26	30	16	38	42	68	74
2003.....	4,647	9	26	21	21	22	29	28	32	46	68	106
2002.....	4,609	1	15	23	14	39	39	32	41	42	67	80
2001.....	4,587	8	33	20	32	31	31	30	36	46	51	99
2000.....	4,723	12	27	23	17	34	35	40	33	49	65	103
1999.....	4,758	8	16	27	22	40	39	33	42	58	70	97

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other and unspecified leukemia (C94-C95)

White, male

2004.....	2,556	4	6	12	16	16	21	12	27	25	36	47
2003.....	2,507	2	14	11	9	17	18	19	20	24	43	61
2002.....	2,496	1	7	13	11	24	26	19	25	21	39	42
2001.....	2,508	2	19	12	16	16	21	23	24	27	28	62
2000.....	2,487	3	12	13	8	17	18	25	20	28	37	61
1999.....	2,543	4	9	17	11	20	27	17	22	30	44	58

White, female

2004.....	2,043	6	9	9	7	10	9	4	11	17	32	27
2003.....	2,140	7	12	10	12	5	11	9	12	22	25	45
2002.....	2,113	-	8	10	3	15	13	13	16	21	28	38
2001.....	2,079	6	14	8	16	15	10	7	12	19	23	37
2000.....	2,236	9	15	10	9	17	17	15	13	21	28	42
1999.....	2,215	4	7	10	11	20	12	16	20	28	26	39

Black, both sexes

2004.....	487	-	2	4	5	11	3	3	8	8	20	12
2003.....	444	1	3	5	3	-	4	5	8	15	15	10
2002.....	464	1	2	5	6	5	2	2	7	6	12	25
2001.....	431	3	2	6	5	5	7	10	7	8	14	13
2000.....	424	-	2	1	3	8	6	10	4	9	12	21
1999.....	432	-	2	2	9	5	7	3	9	12	7	16

Black, male

2004.....	243	-	1	3	2	5	2	-	6	4	10	9
2003.....	212	1	2	3	-	-	1	2	4	6	9	7
2002.....	224	-	-	4	2	4	2	1	3	3	7	16
2001.....	222	2	1	4	5	3	5	5	3	6	10	6
2000.....	230	-	2	1	2	5	4	4	4	5	6	12
1999.....	198	-	1	1	8	3	2	1	3	6	4	10

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified leukemia (C94-C95)												
Asian or Pacific Islander 2/, both sexes												
2004.....	90	1	3	-	-	2	-	2	1	1	4	1
2003.....	91	-	1	1	3	-	4	2	1	1	2	3
2002.....	93	1	-	2	5	1	1	2	2	1	1	9
2001.....	87	-	1	3	2	2	3	5	4	1	3	2
2000.....	101	-	1	4	2	2	2	4	-	2	4	5
1999.....	97	-	-	4	1	5	1	4	4	4	3	3
Asian or Pacific Islander 2/, male												
2004.....	54	1	2	-	-	2	-	1	-	1	4	1
2003.....	53	-	1	-	3	-	1	2	1	1	1	2
2002.....	54	-	-	2	3	-	1	1	2	1	-	4
2001.....	51	-	1	2	2	-	1	-	2	-	2	1
2000.....	59	-	-	2	2	2	1	4	-	2	3	3
1999.....	58	-	-	2	-	4	1	3	3	3	1	2
Asian or Pacific Islander 2/, female												
2004.....	36	-	1	-	-	-	-	1	1	-	-	-
2003.....	38	-	-	1	-	-	3	-	-	-	1	1
2002.....	39	1	-	-	2	1	-	1	-	-	1	5
2001.....	36	-	-	1	-	2	2	5	2	1	1	1
2000.....	42	-	1	2	-	-	1	-	-	-	1	2
1999.....	39	-	-	2	1	1	-	1	1	1	2	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

In situ neoplasms (D00-D09)

All races, both sexes

2004.....	8	-	-	-	-	-	-	-	-	-	-	-
2003.....	13	-	-	-	-	-	-	-	-	-	1	-
2002.....	18	-	-	-	-	1	-	-	-	-	-	1
2001.....	31	-	-	-	-	-	-	-	1	-	-	-
2000.....	18	-	-	1	-	-	-	-	-	-	2	1
1999.....	19	-	-	-	-	-	-	-	2	1	1	1

All races, male

2004.....	3	-	-	-	-	-	-	-	-	-	-	-
2003.....	7	-	-	-	-	-	-	-	-	-	1	-
2002.....	7	-	-	-	-	1	-	-	-	-	-	-
2001.....	16	-	-	-	-	-	-	-	-	-	-	-
2000.....	11	-	-	1	-	-	-	-	-	-	-	1
1999.....	9	-	-	-	-	-	-	-	1	1	-	1

All races, female

2004.....	5	-	-	-	-	-	-	-	-	-	-	-
2003.....	6	-	-	-	-	-	-	-	-	-	-	-
2002.....	11	-	-	-	-	-	-	-	-	-	-	1
2001.....	15	-	-	-	-	-	-	-	1	-	-	-
2000.....	7	-	-	-	-	-	-	-	-	-	2	-
1999.....	10	-	-	-	-	-	-	-	1	-	1	-

White, both sexes

2004.....	6	-	-	-	-	-	-	-	-	-	-	-
2003.....	10	-	-	-	-	-	-	-	-	-	1	-
2002.....	17	-	-	-	-	1	-	-	-	-	-	1
2001.....	25	-	-	-	-	-	-	-	1	-	-	-
2000.....	13	-	-	1	-	-	-	-	-	-	2	-
1999.....	18	-	-	-	-	-	-	-	2	-	1	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
In situ neoplasms (D00-D09)												
White, male												
2004.....	2	-	-	-	-	-	-	-	-	-	-	-
2003.....	5	-	-	-	-	-	-	-	-	-	1	-
2002.....	7	-	-	-	-	1	-	-	-	-	-	-
2001.....	14	-	-	-	-	-	-	-	-	-	-	-
2000.....	7	-	-	1	-	-	-	-	-	-	-	-
1999.....	8	-	-	-	-	-	-	-	1	-	-	1
White, female												
2004.....	4	-	-	-	-	-	-	-	-	-	-	-
2003.....	5	-	-	-	-	-	-	-	-	-	-	-
2002.....	10	-	-	-	-	-	-	-	-	-	-	1
2001.....	11	-	-	-	-	-	-	-	1	-	-	-
2000.....	6	-	-	-	-	-	-	-	-	-	2	-
1999.....	10	-	-	-	-	-	-	-	1	-	1	-
Black, both sexes												
2004.....	2	-	-	-	-	-	-	-	-	-	-	-
2003.....	3	-	-	-	-	-	-	-	-	-	-	-
2002.....	1	-	-	-	-	-	-	-	-	-	-	-
2001.....	6	-	-	-	-	-	-	-	-	-	-	-
2000.....	5	-	-	-	-	-	-	-	-	-	-	1
1999.....	1	-	-	-	-	-	-	-	-	1	-	-
Black, male												
2004.....	1	-	-	-	-	-	-	-	-	-	-	-
2003.....	2	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	2	-	-	-	-	-	-	-	-	-	-	-
2000.....	4	-	-	-	-	-	-	-	-	-	-	1
1999.....	1	-	-	-	-	-	-	-	-	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Carcinoma in situ of other and unspecified sites (D00-D04,D09)

All races, both sexes

2004.....	8	-	-	-	-	-	-	-	-	-	-	-
2003.....	10	-	-	-	-	-	-	-	-	-	1	-
2002.....	13	-	-	-	-	1	-	-	-	-	-	-
2001.....	23	-	-	-	-	-	-	-	-	-	-	-
2000.....	14	-	-	1	-	-	-	-	-	-	1	-
1999.....	14	-	-	-	-	-	-	-	2	1	-	1

All races, male

2004.....	3	-	-	-	-	-	-	-	-	-	-	-
2003.....	6	-	-	-	-	-	-	-	-	-	1	-
2002.....	7	-	-	-	-	1	-	-	-	-	-	-
2001.....	15	-	-	-	-	-	-	-	-	-	-	-
2000.....	9	-	-	1	-	-	-	-	-	-	-	-
1999.....	8	-	-	-	-	-	-	-	1	1	-	1

All races, female

2004.....	5	-	-	-	-	-	-	-	-	-	-	-
2003.....	4	-	-	-	-	-	-	-	-	-	-	-
2002.....	6	-	-	-	-	-	-	-	-	-	-	-
2001.....	8	-	-	-	-	-	-	-	-	-	-	-
2000.....	5	-	-	-	-	-	-	-	-	-	1	-
1999.....	6	-	-	-	-	-	-	-	1	-	-	-

White, both sexes

2004.....	6	-	-	-	-	-	-	-	-	-	-	-
2003.....	7	-	-	-	-	-	-	-	-	-	1	-
2002.....	13	-	-	-	-	1	-	-	-	-	-	-
2001.....	18	-	-	-	-	-	-	-	-	-	-	-
2000.....	10	-	-	1	-	-	-	-	-	-	1	-
1999.....	13	-	-	-	-	-	-	-	2	-	-	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Carcinoma in situ of other and unspecified sites (D00-D04,D09)												
White, male												
2004.....	2	-	-	-	-	-	-	-	-	-	-	-
2003.....	4	-	-	-	-	-	-	-	-	-	1	-
2002.....	7	-	-	-	-	1	-	-	-	-	-	-
2001.....	13	-	-	-	-	-	-	-	-	-	-	-
2000.....	6	-	-	1	-	-	-	-	-	-	-	-
1999.....	7	-	-	-	-	-	-	-	1	-	-	1
White, female												
2004.....	4	-	-	-	-	-	-	-	-	-	-	-
2003.....	3	-	-	-	-	-	-	-	-	-	-	-
2002.....	6	-	-	-	-	-	-	-	-	-	-	-
2001.....	5	-	-	-	-	-	-	-	-	-	-	-
2000.....	4	-	-	-	-	-	-	-	-	-	1	-
1999.....	6	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes												
2004.....	2	-	-	-	-	-	-	-	-	-	-	-
2003.....	3	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	5	-	-	-	-	-	-	-	-	-	-	-
2000.....	4	-	-	-	-	-	-	-	-	-	-	-
1999.....	1	-	-	-	-	-	-	-	-	1	-	-
Black, male												
2004.....	1	-	-	-	-	-	-	-	-	-	-	-
2003.....	2	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	2	-	-	-	-	-	-	-	-	-	-	-
2000.....	3	-	-	-	-	-	-	-	-	-	-	-
1999.....	1	-	-	-	-	-	-	-	-	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Benign neoplasms (D10-D36)												
All races, both sexes												
2004.....	1,553	19	1	-	4	5	10	14	34	26	34	61
2003.....	1,541	12	9	4	7	3	12	14	19	31	50	66
2002.....	1,590	14	7	4	2	4	6	16	21	36	50	78
2001.....	1,554	15	10	2	3	12	10	14	23	25	37	49
2000.....	1,589	21	2	2	3	6	9	13	17	35	49	50
1999.....	1,537	14	7	5	2	5	6	8	24	31	30	41
All races, male												
2004.....	581	10	-	-	3	1	8	5	15	15	10	21
2003.....	624	6	5	2	4	-	6	7	6	10	22	26
2002.....	616	8	3	3	1	3	4	10	10	18	19	30
2001.....	639	9	4	1	1	4	7	5	9	9	17	19
2000.....	590	13	1	-	1	5	3	1	9	13	13	27
1999.....	608	6	2	2	-	2	2	3	10	15	20	18
All races, female												
2004.....	972	9	1	-	1	4	2	9	19	11	24	40
2003.....	917	6	4	2	3	3	6	7	13	21	28	40
2002.....	974	6	4	1	1	1	2	6	11	18	31	48
2001.....	915	6	6	1	2	8	3	9	14	16	20	30
2000.....	999	8	1	2	2	1	6	12	8	22	36	23
1999.....	929	8	5	3	2	3	4	5	14	16	10	23
White, both sexes												
2004.....	1,273	16	1	-	2	4	8	8	23	22	21	47
2003.....	1,292	9	8	2	4	1	11	12	16	20	31	48
2002.....	1,312	13	3	3	-	3	5	10	16	27	37	60
2001.....	1,263	9	8	1	2	10	8	10	19	16	29	34
2000.....	1,326	15	1	-	3	6	7	6	13	23	28	29
1999.....	1,259	9	6	4	2	4	4	5	13	19	21	25

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Benign neoplasms (D10-D36)												
White, male												
2004.....	482	9	-	-	2	1	7	3	10	13	6	19
2003.....	514	4	4	1	2	-	6	6	4	8	15	20
2002.....	509	7	1	2	-	2	3	7	8	13	17	24
2001.....	519	6	4	-	-	3	5	3	7	4	15	15
2000.....	494	9	-	-	1	5	3	1	8	8	8	21
1999.....	496	4	2	1	-	2	2	2	5	9	16	13
White, female												
2004.....	791	7	1	-	-	3	1	5	13	9	15	28
2003.....	778	5	4	1	2	1	5	6	12	12	16	28
2002.....	803	6	2	1	-	1	2	3	8	14	20	36
2001.....	744	3	4	1	2	7	3	7	12	12	14	19
2000.....	832	6	1	-	2	1	4	5	5	15	20	8
1999.....	763	5	4	3	2	2	2	3	8	10	5	12
Black, both sexes												
2004.....	236	3	-	-	1	-	1	6	11	4	13	12
2003.....	211	3	1	2	2	1	1	2	3	9	17	18
2002.....	229	-	4	1	2	1	1	5	2	8	11	17
2001.....	248	5	2	1	1	2	1	4	3	8	5	12
2000.....	226	4	1	2	-	-	2	6	4	11	19	19
1999.....	236	4	1	1	-	1	2	1	7	10	7	13
Black, male												
2004.....	89	1	-	-	-	-	1	2	5	2	4	2
2003.....	93	2	1	1	1	-	-	1	2	1	7	6
2002.....	86	-	2	1	1	1	1	2	-	4	2	5
2001.....	104	3	-	1	1	1	1	2	2	5	1	3
2000.....	76	2	1	-	-	-	-	-	1	4	4	5
1999.....	95	1	-	1	-	-	-	1	4	5	3	4

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Benign neoplasms (D10-D36)												
Asian or Pacific Islander 2/, both sexes												
2004.....	41	-	-	-	-	1	1	-	-	-	-	2
2003.....	32	-	-	-	-	1	-	-	-	2	1	-
2002.....	43	1	-	-	-	-	-	1	3	1	1	1
2001.....	40	1	-	-	-	-	1	-	1	-	3	3
2000.....	29	2	-	-	-	-	-	1	-	-	2	1
1999.....	38	1	-	-	-	-	-	2	4	2	2	3
Asian or Pacific Islander 2/, male												
2004.....	9	-	-	-	-	-	-	-	-	-	-	-
2003.....	15	-	-	-	-	-	-	-	-	1	-	-
2002.....	19	1	-	-	-	-	-	1	2	1	-	1
2001.....	16	-	-	-	-	-	1	-	-	-	1	1
2000.....	16	2	-	-	-	-	-	-	-	-	1	1
1999.....	16	1	-	-	-	-	-	-	1	1	1	1
Asian or Pacific Islander 2/, female												
2004.....	32	-	-	-	-	1	1	-	-	-	-	2
2003.....	17	-	-	-	-	1	-	-	-	1	1	-
2002.....	24	-	-	-	-	-	-	-	1	-	1	-
2001.....	24	1	-	-	-	-	-	-	1	-	2	2
2000.....	13	-	-	-	-	-	-	1	-	-	1	-
1999.....	22	-	-	-	-	-	-	2	3	1	1	2

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Of female genital organs (D25-D28)												
All races, both sexes												
2004.....	40	2	-	-	-	-	1	-	1	1	7	6
2003.....	41	-	-	-	-	1	-	-	-	4	5	11
2002.....	50	1	-	-	-	-	-	1	1	2	7	12
2001.....	51	1	-	-	1	1	-	2	3	6	2	8
2000.....	54	-	-	-	-	-	-	3	1	5	13	7
1999.....	33	1	-	-	1	1	1	-	1	2	6	1
All races, male												
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....
All races, female												
2004.....	40	2	-	-	-	-	1	-	1	1	7	6
2003.....	41	-	-	-	-	1	-	-	-	4	5	11
2002.....	50	1	-	-	-	-	-	1	1	2	7	12
2001.....	51	1	-	-	1	1	-	2	3	6	2	8
2000.....	54	-	-	-	-	-	-	3	1	5	13	7
1999.....	33	1	-	-	1	1	1	-	1	2	6	1
White, both sexes												
2004.....	21	1	-	-	-	-	-	-	-	1	3	2
2003.....	23	-	-	-	-	1	-	-	-	2	1	3
2002.....	27	1	-	-	-	-	-	-	-	1	1	4
2001.....	34	-	-	-	1	1	-	1	2	5	-	5
2000.....	25	-	-	-	-	-	-	-	-	1	3	2
1999.....	18	-	-	-	1	1	1	-	1	-	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Of eye, brain and other parts of central nervous system (D31-D33)												
All races, both sexes												
2004.....	968	-	1	-	1	1	3	5	12	9	13	32
2003.....	955	-	3	-	1	1	5	3	6	10	20	27
2002.....	953	1	2	-	1	2	1	6	7	13	22	33
2001.....	921	1	1	-	-	-	2	6	10	7	13	20
2000.....	981	1	-	-	1	3	4	4	6	11	19	22
1999.....	947	-	-	-	-	1	-	4	9	18	5	15
All races, male												
2004.....	334	-	-	-	1	-	2	1	7	7	6	15
2003.....	365	-	1	-	-	-	4	2	1	4	7	12
2002.....	330	1	2	-	1	1	1	4	5	7	8	17
2001.....	334	-	-	-	-	-	2	2	4	4	6	7
2000.....	346	1	-	-	-	3	1	-	4	3	8	13
1999.....	342	-	-	-	-	-	-	2	4	9	5	5
All races, female												
2004.....	634	-	1	-	-	1	1	4	5	2	7	17
2003.....	590	-	2	-	1	1	1	1	5	6	13	15
2002.....	623	-	-	-	-	1	-	2	2	6	14	16
2001.....	587	1	1	-	-	-	-	4	6	3	7	13
2000.....	635	-	-	-	1	-	3	4	2	8	11	9
1999.....	605	-	-	-	-	1	-	2	5	9	-	10
White, both sexes												
2004.....	801	-	1	-	1	1	2	4	10	7	7	27
2003.....	815	-	3	-	1	-	5	3	6	6	13	23
2002.....	809	1	1	-	-	2	1	5	5	10	19	30
2001.....	746	-	1	-	-	-	1	5	8	5	11	13
2000.....	844	1	-	-	1	3	2	2	5	9	13	15
1999.....	793	-	-	-	-	1	-	3	5	12	4	10

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Of eye, brain and other parts of central nervous system (D31-D33)

White, male

2004.....	276	-	-	-	1	-	1	1	6	6	4	14
2003.....	312	-	1	-	-	-	4	2	1	4	5	9
2002.....	276	1	1	-	-	1	1	3	3	6	7	15
2001.....	270	-	-	-	-	-	1	1	3	3	5	4
2000.....	295	1	-	-	-	3	1	-	3	2	5	11
1999.....	287	-	-	-	-	-	-	1	3	5	4	4

White, female

2004.....	525	-	1	-	-	1	1	3	4	1	3	13
2003.....	503	-	2	-	1	-	1	1	5	2	8	14
2002.....	533	-	-	-	-	1	-	2	2	4	12	15
2001.....	476	-	1	-	-	-	-	4	5	2	6	9
2000.....	549	-	-	-	1	-	1	2	2	7	8	4
1999.....	506	-	-	-	-	1	-	2	2	7	-	6

Black, both sexes

2004.....	139	-	-	-	-	-	1	1	2	2	6	3
2003.....	117	-	-	-	-	1	-	-	-	4	5	4
2002.....	116	-	1	-	1	-	-	1	-	3	3	3
2001.....	145	-	-	-	-	-	-	1	1	2	-	5
2000.....	114	-	-	-	-	-	2	1	1	2	6	5
1999.....	136	-	-	-	-	-	-	1	2	4	-	5

Black, male

2004.....	53	-	-	-	-	-	1	-	1	1	2	1
2003.....	45	-	-	-	-	-	-	-	-	-	2	3
2002.....	42	-	1	-	1	-	-	1	-	1	1	2
2001.....	53	-	-	-	-	-	-	1	1	1	-	2
2000.....	39	-	-	-	-	-	-	-	1	1	3	1
1999.....	48	-	-	-	-	-	-	1	1	3	-	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Of eye, brain and other parts of central nervous system (D31-D33)

Asian or Pacific Islander 2/, both sexes

2004.....	27	-	-	-	-	-	-	-	-	-	-	2
2003.....	18	-	-	-	-	-	-	-	-	-	1	-
2002.....	24	-	-	-	-	-	-	-	2	-	-	-
2001.....	28	1	-	-	-	-	1	-	1	-	2	2
2000.....	18	-	-	-	-	-	-	1	-	-	-	1
1999.....	16	-	-	-	-	-	-	-	2	2	1	-

Asian or Pacific Islander 2/, male

2004.....	5	-	-	-	-	-	-	-	-	-	-	-
2003.....	7	-	-	-	-	-	-	-	-	-	-	-
2002.....	10	-	-	-	-	-	-	-	2	-	-	-
2001.....	11	-	-	-	-	-	1	-	-	-	1	1
2000.....	10	-	-	-	-	-	-	-	-	-	-	1
1999.....	7	-	-	-	-	-	-	-	-	1	1	-

Asian or Pacific Islander 2/, female

2004.....	22	-	-	-	-	-	-	-	-	-	-	2
2003.....	11	-	-	-	-	-	-	-	-	-	1	-
2002.....	14	-	-	-	-	-	-	-	-	-	-	-
2001.....	17	1	-	-	-	-	-	-	1	-	1	1
2000.....	8	-	-	-	-	-	-	1	-	-	-	-
1999.....	9	-	-	-	-	-	-	-	2	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Of other and unspecified sites (D10-D24,D29-D30,D34-D36)

All races, both sexes

2004.....	545	17	-	-	3	4	6	9	21	16	14	23
2003.....	545	12	6	4	6	1	7	11	13	17	25	28
2002.....	587	12	5	4	1	2	5	9	13	21	21	33
2001.....	582	13	9	2	2	11	8	6	10	12	22	21
2000.....	554	20	2	2	2	3	5	6	10	19	17	21
1999.....	557	13	7	5	1	3	5	4	14	11	19	25

All races, male

2004.....	247	10	-	-	2	1	6	4	8	8	4	6
2003.....	259	6	4	2	4	-	2	5	5	6	15	14
2002.....	286	7	1	3	-	2	3	6	5	11	11	13
2001.....	305	9	4	1	1	4	5	3	5	5	11	12
2000.....	244	12	1	-	1	2	2	1	5	10	5	14
1999.....	266	6	2	2	-	2	2	1	6	6	15	13

All races, female

2004.....	298	7	-	-	1	3	-	5	13	8	10	17
2003.....	286	6	2	2	2	1	5	6	8	11	10	14
2002.....	301	5	4	1	1	-	2	3	8	10	10	20
2001.....	277	4	5	1	1	7	3	3	5	7	11	9
2000.....	310	8	1	2	1	1	3	5	5	9	12	7
1999.....	291	7	5	3	1	1	3	3	8	5	4	12

White, both sexes

2004.....	451	15	-	-	1	3	6	4	13	14	11	18
2003.....	454	9	5	2	3	-	6	9	10	12	17	22
2002.....	476	11	2	3	-	1	4	5	11	16	17	26
2001.....	483	9	7	1	1	9	7	4	9	6	18	16
2000.....	457	14	1	-	2	3	5	4	8	13	12	12
1999.....	448	9	6	4	1	2	3	2	7	7	16	15

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Of other and unspecified sites (D10-D24,D29-D30,D34-D36)

White, male

2004.....	206	9	-	-	1	1	6	2	4	7	2	5
2003.....	202	4	3	1	2	-	2	4	3	4	10	11
2002.....	233	6	-	2	-	1	2	4	5	7	10	9
2001.....	249	6	4	-	-	3	4	2	4	1	10	11
2000.....	199	8	-	-	1	2	2	1	5	6	3	10
1999.....	209	4	2	1	-	2	2	1	2	4	12	9

White, female

2004.....	245	6	-	-	-	2	-	2	9	7	9	13
2003.....	252	5	2	1	1	-	4	5	7	8	7	11
2002.....	243	5	2	1	-	-	2	1	6	9	7	17
2001.....	234	3	3	1	1	6	3	2	5	5	8	5
2000.....	258	6	1	-	1	1	3	3	3	7	9	2
1999.....	239	5	4	3	1	-	1	1	5	3	4	6

Black, both sexes

2004.....	80	2	-	-	1	-	-	5	8	2	3	5
2003.....	78	3	1	2	2	-	1	2	3	4	8	6
2002.....	92	-	3	1	1	1	1	3	1	4	3	6
2001.....	86	4	2	1	1	2	1	2	1	5	3	4
2000.....	85	4	1	2	-	-	-	2	2	5	3	9
1999.....	87	3	1	1	-	1	2	-	5	4	3	7

Black, male

2004.....	36	1	-	-	-	-	-	2	4	1	2	1
2003.....	48	2	1	1	1	-	-	1	2	1	5	3
2002.....	44	-	1	1	-	1	1	1	-	3	1	3
2001.....	51	3	-	1	1	1	1	1	1	4	1	1
2000.....	37	2	1	-	-	-	-	-	-	3	1	4
1999.....	47	1	-	1	-	-	-	-	3	2	3	3

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Of other and unspecified sites (D10-D24,D29-D30,D34-D36)												
Asian or Pacific Islander 2/, both sexes												
2004.....	12	-	-	-	-	1	-	-	-	-	-	-
2003.....	12	-	-	-	-	1	-	-	-	1	-	-
2002.....	17	1	-	-	-	-	-	1	1	1	-	1
2001.....	12	-	-	-	-	-	-	-	-	-	1	1
2000.....	9	2	-	-	-	-	-	-	-	-	2	-
1999.....	20	1	-	-	-	-	-	2	2	-	-	3
Asian or Pacific Islander 2/, male												
2004.....	4	-	-	-	-	-	-	-	-	-	-	-
2003.....	8	-	-	-	-	-	-	-	-	1	-	-
2002.....	9	1	-	-	-	-	-	1	-	1	-	1
2001.....	5	-	-	-	-	-	-	-	-	-	-	-
2000.....	6	2	-	-	-	-	-	-	-	-	1	-
1999.....	9	1	-	-	-	-	-	-	1	-	-	1
Asian or Pacific Islander 2/, female												
2004.....	8	-	-	-	-	1	-	-	-	-	-	-
2003.....	4	-	-	-	-	1	-	-	-	-	-	-
2002.....	8	-	-	-	-	-	-	-	1	-	-	-
2001.....	7	-	-	-	-	-	-	-	-	-	1	1
2000.....	3	-	-	-	-	-	-	-	-	-	1	-
1999.....	11	-	-	-	-	-	-	2	1	-	-	2

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Neoplasms of uncertain or unknown behavior of specified sites (D37-D47)

All races, both sexes

2004.....	11,268	32	45	38	33	34	39	46	66	89	140	213
2003.....	11,423	35	39	31	31	30	33	38	79	101	134	216
2002.....	11,037	36	46	35	38	30	36	39	67	85	152	221
2001.....	11,471	22	44	44	47	29	32	37	57	120	163	230
2000.....	11,334	25	48	54	32	29	39	47	69	109	149	218
1999.....	11,144	32	54	53	31	40	38	39	82	101	162	219

All races, male

2004.....	5,860	16	26	19	19	16	23	29	39	47	86	115
2003.....	5,814	17	23	16	17	19	16	24	45	65	85	116
2002.....	5,669	17	22	16	22	11	23	20	34	47	87	128
2001.....	5,851	12	23	22	22	12	18	21	31	64	91	139
2000.....	5,741	13	29	29	14	19	27	26	35	69	84	119
1999.....	5,702	19	30	25	17	19	24	24	47	58	94	116

All races, female

2004.....	5,408	16	19	19	14	18	16	17	27	42	54	98
2003.....	5,609	18	16	15	14	11	17	14	34	36	49	100
2002.....	5,368	19	24	19	16	19	13	19	33	38	65	93
2001.....	5,620	10	21	22	25	17	14	16	26	56	72	91
2000.....	5,593	12	19	25	18	10	12	21	34	40	65	99
1999.....	5,442	13	24	28	14	21	14	15	35	43	68	103

White, both sexes

2004.....	10,162	24	35	29	23	24	33	39	47	67	110	176
2003.....	10,360	26	29	22	24	24	28	28	64	73	119	173
2002.....	10,015	32	36	28	29	25	25	28	52	73	118	191
2001.....	10,424	16	36	32	34	20	22	35	45	99	128	200
2000.....	10,295	20	40	42	25	18	32	35	58	91	127	176
1999.....	10,130	21	41	42	28	29	28	30	64	83	133	178

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Neoplasms of uncertain or unknown behavior of specified sites (D37-D47)

White, male

2004.....	5,309	12	20	15	12	13	19	25	31	34	71	94
2003.....	5,303	12	16	10	15	15	14	19	36	50	73	95
2002.....	5,176	16	17	11	19	10	15	16	31	43	73	114
2001.....	5,355	10	17	16	15	6	14	20	25	55	75	120
2000.....	5,211	11	22	22	12	10	23	20	30	56	68	96
1999.....	5,217	13	20	20	15	12	18	16	36	50	81	97

White, female

2004.....	4,853	12	15	14	11	11	14	14	16	33	39	82
2003.....	5,057	14	13	12	9	9	14	9	28	23	46	78
2002.....	4,839	16	19	17	10	15	10	12	21	30	45	77
2001.....	5,069	6	19	16	19	14	8	15	20	44	53	80
2000.....	5,084	9	18	20	13	8	9	15	28	35	59	80
1999.....	4,913	8	21	22	13	17	10	14	28	33	52	81

Black, both sexes

2004.....	869	6	8	8	9	8	6	5	15	19	23	27
2003.....	852	7	7	7	4	5	5	9	11	26	11	39
2002.....	849	4	8	6	8	4	8	8	15	10	32	28
2001.....	849	5	3	11	12	7	7	1	9	17	29	21
2000.....	843	4	8	9	6	8	5	10	11	12	18	33
1999.....	862	10	11	11	3	9	10	5	13	17	27	32

Black, male

2004.....	425	3	5	4	7	2	4	3	5	11	10	16
2003.....	407	3	6	4	1	3	2	5	6	14	8	19
2002.....	391	1	5	5	2	1	5	3	3	2	13	12
2001.....	397	1	3	5	6	5	4	-	6	8	15	12
2000.....	413	1	7	5	2	7	2	5	5	9	13	16
1999.....	395	6	9	5	2	6	6	5	7	8	12	13

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Neoplasms of uncertain or unknown behavior of specified sites (D37-D47)												
Asian or Pacific Islander 2/, both sexes												
2004.....	193	2	1	1	1	1	-	1	4	1	5	9
2003.....	180	2	3	2	3	1	-	1	2	1	4	3
2002.....	147	-	-	1	-	1	3	2	-	2	1	2
2001.....	148	-	5	-	1	-	2	1	2	3	2	7
2000.....	162	1	-	2	1	1	1	-	-	4	3	9
1999.....	117	1	2	-	-	1	-	4	5	1	2	8
Asian or Pacific Islander 2/, male												
2004.....	104	1	1	-	-	1	-	-	3	1	3	5
2003.....	87	2	1	2	1	1	-	-	2	1	4	1
2002.....	89	-	-	-	-	-	3	1	-	2	1	2
2001.....	73	-	3	-	1	-	-	1	-	-	-	5
2000.....	102	1	-	1	-	1	1	-	-	4	3	7
1999.....	70	-	1	-	-	-	-	3	4	-	1	5
Asian or Pacific Islander 2/, female												
2004.....	89	1	-	1	1	-	-	1	1	-	2	4
2003.....	93	-	2	-	2	-	-	1	-	-	-	2
2002.....	58	-	-	1	-	1	-	1	-	-	-	-
2001.....	75	-	2	-	-	-	2	-	2	3	2	2
2000.....	60	-	-	1	1	-	-	-	-	-	-	2
1999.....	47	1	1	-	-	1	-	1	1	1	1	3

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Neoplasms of uncertain or unknown behavior of unspecified sites (D48)

All races, both sexes

2004.....	751	20	7	3	6	11	11	3	10	17	12	34
2003.....	586	15	3	3	3	8	10	17	11	10	12	13
2002.....	654	19	7	5	5	6	4	7	13	16	27	27
2001.....	620	14	4	6	3	6	13	9	10	11	10	25
2000.....	605	14	3	5	2	5	7	8	11	11	13	24
1999.....	527	12	2	6	4	7	2	8	8	7	9	15

All races, male

2004.....	320	9	2	1	4	8	7	1	5	6	8	22
2003.....	242	9	3	3	1	5	8	9	7	4	8	6
2002.....	244	5	5	3	4	6	2	3	7	12	13	11
2001.....	229	5	3	3	3	4	8	7	4	6	6	15
2000.....	217	6	3	2	2	3	5	3	8	4	5	13
1999.....	201	5	1	3	2	5	2	5	6	5	4	7

All races, female

2004.....	431	11	5	2	2	3	4	2	5	11	4	12
2003.....	344	6	-	-	2	3	2	8	4	6	4	7
2002.....	410	14	2	2	1	-	2	4	6	4	14	16
2001.....	391	9	1	3	-	2	5	2	6	5	4	10
2000.....	388	8	-	3	-	2	2	5	3	7	8	11
1999.....	326	7	1	3	2	2	-	3	2	2	5	8

White, both sexes

2004.....	651	19	7	1	4	7	10	2	10	12	8	28
2003.....	519	13	2	2	1	6	8	14	10	8	10	10
2002.....	569	14	4	5	2	5	4	4	11	12	19	22
2001.....	549	12	4	5	3	3	11	8	8	9	9	23
2000.....	537	13	3	3	2	5	5	6	11	10	11	20
1999.....	442	11	1	6	2	4	1	4	6	5	7	10

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Neoplasms of uncertain or unknown behavior of unspecified sites (D48)

White, male

2004.....	278	9	2	1	3	6	6	-	5	3	5	18
2003.....	214	9	2	2	-	4	7	6	6	4	7	5
2002.....	210	4	3	3	2	5	2	1	7	9	10	8
2001.....	201	4	3	3	3	3	6	6	3	5	5	14
2000.....	188	5	3	1	2	3	4	3	8	4	3	10
1999.....	170	4	-	3	1	3	1	3	4	4	4	5

White, female

2004.....	373	10	5	-	1	1	4	2	5	9	3	10
2003.....	305	4	-	-	1	2	1	8	4	4	3	5
2002.....	359	10	1	2	-	-	2	3	4	3	9	14
2001.....	348	8	1	2	-	-	5	2	5	4	4	9
2000.....	349	8	-	2	-	2	1	3	3	6	8	10
1999.....	272	7	1	3	1	1	-	1	2	1	3	5

Black, both sexes

2004.....	85	1	-	1	2	3	1	1	-	5	3	5
2003.....	57	1	1	1	2	2	2	3	1	2	2	2
2002.....	64	5	2	-	1	1	-	-	1	2	7	5
2001.....	54	1	-	1	-	3	1	1	-	1	1	2
2000.....	62	1	-	2	-	-	1	2	-	1	2	3
1999.....	74	1	1	-	2	3	1	3	1	2	2	5

Black, male

2004.....	36	-	-	-	1	2	1	1	-	3	3	3
2003.....	23	-	1	1	1	1	1	3	1	-	1	1
2002.....	23	1	1	-	-	1	-	-	-	2	3	3
2001.....	22	1	-	-	-	1	1	1	-	-	1	1
2000.....	27	1	-	1	-	-	-	-	-	-	2	2
1999.....	30	1	1	-	1	2	1	2	1	1	-	2

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Neoplasms of uncertain or unknown behavior of unspecified sites (D48)

Asian or Pacific Islander 2/, both sexes

2004.....	14	-	-	1	-	1	-	-	-	-	1	1
2003.....	8	1	-	-	-	-	-	-	-	-	-	1
2002.....	19	-	1	-	2	-	-	3	1	2	-	-
2001.....	12	1	-	-	-	-	-	-	2	1	-	-
2000.....	3	-	-	-	-	-	-	-	-	-	-	1
1999.....	8	-	-	-	-	-	-	1	-	-	-	-

Asian or Pacific Islander 2/, male

2004.....	6	-	-	-	-	-	-	-	-	-	-	1
2003.....	4	-	-	-	-	-	-	-	-	-	-	-
2002.....	11	-	1	-	2	-	-	2	-	1	-	-
2001.....	3	-	-	-	-	-	-	-	1	1	-	-
2000.....	1	-	-	-	-	-	-	-	-	-	-	1
1999.....	-	-	-	-	-	-	-	-	-	-	-	-

Asian or Pacific Islander 2/, female

2004.....	8	-	-	1	-	1	-	-	-	-	1	-
2003.....	4	1	-	-	-	-	-	-	-	-	-	1
2002.....	8	-	-	-	-	-	-	1	1	1	-	-
2001.....	9	1	-	-	-	-	-	-	1	-	-	-
2000.....	2	-	-	-	-	-	-	-	-	-	-	-
1999.....	8	-	-	-	-	-	-	1	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

III. Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)

All races, both sexes

2004.....	9,424	94	68	50	44	82	106	145	168	234	376	438
2003.....	9,585	105	73	32	60	60	114	120	192	256	376	445
2002.....	9,661	82	77	41	40	72	111	136	193	270	390	474
2001.....	9,535	106	73	40	47	72	96	112	218	260	417	447
2000.....	9,315	70	82	43	46	61	94	119	194	286	355	417
1999.....	9,067	94	82	49	50	80	116	122	210	281	360	407

All races, male

2004.....	4,067	59	38	25	24	46	57	72	93	128	193	228
2003.....	4,085	60	39	16	34	32	59	54	99	131	201	240
2002.....	4,065	52	40	17	24	44	63	84	107	136	181	222
2001.....	3,938	59	47	20	29	37	52	66	120	142	186	214
2000.....	3,852	40	40	31	23	29	48	60	95	140	170	219
1999.....	3,826	58	50	21	31	43	64	70	109	159	173	211

All races, female

2004.....	5,357	35	30	25	20	36	49	73	75	106	183	210
2003.....	5,500	45	34	16	26	28	55	66	93	125	175	205
2002.....	5,596	30	37	24	16	28	48	52	86	134	209	252
2001.....	5,597	47	26	20	18	35	44	46	98	118	231	233
2000.....	5,463	30	42	12	23	32	46	59	99	146	185	198
1999.....	5,241	36	32	28	19	37	52	52	101	122	187	196

White, both sexes

2004.....	7,043	62	40	25	31	42	37	62	50	86	166	219
2003.....	7,108	71	44	16	37	33	41	43	72	103	148	210
2002.....	7,225	58	51	25	19	39	43	49	81	105	159	235
2001.....	7,156	72	52	25	28	32	36	44	84	93	166	188
2000.....	7,036	50	50	21	26	36	32	33	61	114	151	191
1999.....	6,800	53	46	28	27	51	36	38	69	103	151	202

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

III. Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)

White, male

2004.....	3,012	35	22	17	16	22	25	31	28	51	95	127
2003.....	2,986	45	20	10	21	16	24	18	40	60	86	124
2002.....	2,996	40	28	9	11	24	23	29	47	62	80	122
2001.....	2,925	39	34	14	21	16	20	27	45	56	82	99
2000.....	2,884	26	27	16	14	17	19	19	35	59	86	111
1999.....	2,821	33	29	12	16	26	18	25	38	67	83	117

White, female

2004.....	4,031	27	18	8	15	20	12	31	22	35	71	92
2003.....	4,122	26	24	6	16	17	17	25	32	43	62	86
2002.....	4,229	18	23	16	8	15	20	20	34	43	79	113
2001.....	4,231	33	18	11	7	16	16	17	39	37	84	89
2000.....	4,152	24	23	5	12	19	13	14	26	55	65	80
1999.....	3,979	20	17	16	11	25	18	13	31	36	68	85

Black, both sexes

2004.....	2,190	23	23	19	13	37	69	81	113	143	201	210
2003.....	2,271	20	23	14	20	24	72	75	111	149	222	225
2002.....	2,248	17	21	14	18	30	64	84	106	159	220	230
2001.....	2,180	23	16	13	17	36	59	64	129	160	243	252
2000.....	2,110	16	29	20	20	21	60	82	130	168	196	217
1999.....	2,079	33	34	20	20	27	76	79	140	172	204	190

Black, male

2004.....	977	17	15	7	8	23	32	40	62	74	95	97
2003.....	1,010	11	15	6	11	15	34	35	54	71	111	114
2002.....	985	7	9	8	11	19	40	52	57	71	96	96
2001.....	927	13	10	5	8	20	32	37	74	82	102	110
2000.....	880	12	12	13	9	10	28	39	58	80	81	103
1999.....	911	20	19	9	13	15	44	43	71	89	88	87

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

III. Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)

Black, female

2004.....	1,213	6	8	12	5	14	37	41	51	69	106	113
2003.....	1,261	9	8	8	9	9	38	40	57	78	111	111
2002.....	1,263	10	12	6	7	11	24	32	49	88	124	134
2001.....	1,253	10	6	8	9	16	27	27	55	78	141	142
2000.....	1,230	4	17	7	11	11	32	43	72	88	115	114
1999.....	1,168	13	15	11	7	12	32	36	69	83	116	103

American Indian 1/, both sexes

2004.....	37	2	-	2	-	1	-	-	1	2	4	3
2003.....	42	-	1	-	1	-	-	1	2	2	2	4
2002.....	40	-	2	-	2	2	2	1	1	1	4	2
2001.....	57	3	2	-	2	3	-	1	-	1	5	2
2000.....	48	2	-	-	-	2	1	1	2	-	5	3
1999.....	47	2	1	1	3	1	2	1	1	2	1	4

American Indian 1/, male

2004.....	17	2	-	-	-	-	-	-	1	2	2	1
2003.....	17	-	1	-	1	-	-	-	1	-	1	1
2002.....	16	-	1	-	1	1	-	1	-	1	2	1
2001.....	20	2	1	-	-	1	-	-	-	1	1	1
2000.....	23	1	-	-	-	1	-	1	2	-	3	1
1999.....	23	2	1	-	2	1	2	-	-	2	-	3

American Indian 1/, female

2004.....	20	-	-	2	-	1	-	-	-	-	2	2
2003.....	25	-	-	-	-	-	-	1	1	2	1	3
2002.....	24	-	1	-	1	1	2	-	1	-	2	1
2001.....	37	1	1	-	2	2	-	1	-	-	4	1
2000.....	25	1	-	-	-	1	1	-	-	-	2	2
1999.....	24	-	-	1	1	-	-	1	1	-	1	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

III. Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)

Asian or Pacific Islander 2/, both sexes

2004.....	154	7	5	4	-	2	-	2	4	3	5	6
2003.....	164	14	5	2	2	3	1	1	7	2	4	6
2002.....	148	7	3	2	1	1	2	2	5	5	7	7
2001.....	142	8	3	2	-	1	1	3	5	6	3	5
2000.....	121	2	3	2	-	2	1	3	1	4	3	6
1999.....	141	6	1	-	-	1	2	4	-	4	4	11

Asian or Pacific Islander 2/, male

2004.....	61	5	1	1	-	1	-	1	2	1	1	3
2003.....	72	4	3	-	1	1	1	1	4	-	3	1
2002.....	68	5	2	-	1	-	-	2	3	2	3	3
2001.....	66	5	2	1	-	-	-	2	1	3	1	4
2000.....	65	1	1	2	-	1	1	1	-	1	-	4
1999.....	71	3	1	-	-	1	-	2	-	1	2	4

Asian or Pacific Islander 2/, female

2004.....	93	2	4	3	-	1	-	1	2	2	4	3
2003.....	92	10	2	2	1	2	-	-	3	2	1	5
2002.....	80	2	1	2	-	1	2	-	2	3	4	4
2001.....	76	3	1	1	-	1	1	1	4	3	2	1
2000.....	56	1	2	-	-	1	-	2	1	3	3	2
1999.....	70	3	-	-	-	-	2	2	-	3	2	7

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Anemias (D50-D64)												
All races, both sexes												
2004.....	4,575	15	33	25	21	50	78	84	80	84	117	120
2003.....	4,594	22	24	16	31	37	68	69	88	81	102	137
2002.....	4,614	18	21	29	24	42	69	83	97	99	97	131
2001.....	4,627	16	22	25	30	48	61	65	95	83	132	112
2000.....	4,494	10	35	24	28	27	58	62	91	85	105	110
1999.....	4,503	12	30	24	29	43	78	64	96	90	104	134
All races, male												
2004.....	1,833	9	18	10	11	31	38	39	51	50	57	56
2003.....	1,832	11	14	8	15	19	33	37	46	37	54	77
2002.....	1,781	12	9	13	13	23	44	53	48	53	43	57
2001.....	1,774	11	12	12	15	26	33	40	55	42	49	57
2000.....	1,669	6	16	17	12	9	30	29	41	34	44	52
1999.....	1,746	8	17	8	15	24	43	33	48	47	54	68
All races, female												
2004.....	2,742	6	15	15	10	19	40	45	29	34	60	64
2003.....	2,762	11	10	8	16	18	35	32	42	44	48	60
2002.....	2,833	6	12	16	11	19	25	30	49	46	54	74
2001.....	2,853	5	10	13	15	22	28	25	40	41	83	55
2000.....	2,825	4	19	7	16	18	28	33	50	51	61	58
1999.....	2,757	4	13	16	14	19	35	31	48	43	50	66
White, both sexes												
2004.....	3,456	8	13	10	10	20	15	20	13	14	41	49
2003.....	3,476	9	10	4	17	15	8	16	23	17	28	46
2002.....	3,460	8	12	15	8	13	17	15	28	22	23	49
2001.....	3,543	6	11	11	16	17	16	15	24	16	37	31
2000.....	3,450	4	15	9	12	10	11	15	14	17	26	38
1999.....	3,393	6	9	7	13	18	16	6	11	22	26	55

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Anemias (D50-D64)												
White, male												
2004.....	1,322	4	6	5	3	11	8	7	11	9	22	27
2003.....	1,319	4	5	3	8	6	5	8	14	8	14	29
2002.....	1,267	4	5	6	5	7	10	10	14	14	12	25
2001.....	1,303	5	5	6	10	8	10	9	11	9	12	12
2000.....	1,235	2	8	7	5	2	6	9	6	7	14	16
1999.....	1,244	4	6	1	6	9	5	1	5	10	14	31
White, female												
2004.....	2,134	4	7	5	7	9	7	13	2	5	19	22
2003.....	2,157	5	5	1	9	9	3	8	9	9	14	17
2002.....	2,193	4	7	9	3	6	7	5	14	8	11	24
2001.....	2,240	1	6	5	6	9	6	6	13	7	25	19
2000.....	2,215	2	7	2	7	8	5	6	8	10	12	22
1999.....	2,149	2	3	6	7	9	11	5	6	12	12	24
Black, both sexes												
2004.....	1,029	4	19	12	11	30	63	62	67	68	72	67
2003.....	1,037	6	12	12	13	19	60	52	62	62	72	87
2002.....	1,064	6	9	12	14	27	52	65	65	74	71	80
2001.....	1,001	6	11	13	13	28	45	50	69	62	92	79
2000.....	967	5	19	15	16	16	46	47	77	65	76	69
1999.....	1,019	3	21	17	16	23	61	56	84	66	76	74
Black, male												
2004.....	472	3	12	4	8	20	30	31	40	40	34	27
2003.....	487	4	7	5	7	12	28	28	30	29	40	48
2002.....	479	5	4	7	7	16	34	40	33	37	31	31
2001.....	446	4	7	5	5	18	23	31	43	30	36	45
2000.....	395	4	8	10	7	7	23	20	35	26	29	35
1999.....	457	2	11	7	9	13	37	30	43	35	38	35

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Anemias (D50-D64)												
Black, female												
2004.....	557	1	7	8	3	10	33	31	27	28	38	40
2003.....	550	2	5	7	6	7	32	24	32	33	32	39
2002.....	585	1	5	5	7	11	18	25	32	37	40	49
2001.....	555	2	4	8	8	10	22	19	26	32	56	34
2000.....	572	1	11	5	9	9	23	27	42	39	47	34
1999.....	562	1	10	10	7	10	24	26	41	31	38	39
American Indian 1/, both sexes												
2004.....	13	-	-	1	-	-	-	-	-	1	-	-
2003.....	15	-	-	-	-	-	-	-	1	1	1	1
2002.....	15	-	-	-	1	1	-	1	1	-	1	-
2001.....	27	-	-	-	1	2	-	-	-	1	3	1
2000.....	20	-	-	-	-	-	-	-	-	-	2	1
1999.....	21	-	-	-	-	1	1	-	1	2	-	2
American Indian 1/, male												
2004.....	6	-	-	-	-	-	-	-	-	1	-	-
2003.....	4	-	-	-	-	-	-	-	1	-	-	-
2002.....	4	-	-	-	-	-	-	1	-	-	-	-
2001.....	7	-	-	-	-	-	-	-	-	1	1	-
2000.....	9	-	-	-	-	-	-	-	-	-	1	-
1999.....	9	-	-	-	-	1	1	-	-	2	-	1
American Indian 1/, female												
2004.....	7	-	-	1	-	-	-	-	-	-	-	-
2003.....	11	-	-	-	-	-	-	-	-	1	1	1
2002.....	11	-	-	-	1	1	-	-	1	-	1	-
2001.....	20	-	-	-	1	2	-	-	-	-	2	1
2000.....	11	-	-	-	-	-	-	-	-	-	1	1
1999.....	12	-	-	-	-	-	-	-	1	-	-	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hemolytic anemias (D55-D59)												
All races, both sexes												
2004.....	849	9	19	19	8	30	63	67	68	60	74	61
2003.....	822	16	17	12	13	19	58	59	64	61	71	76
2002.....	894	13	15	12	15	26	54	68	83	73	70	79
2001.....	853	8	16	12	12	27	44	50	76	61	87	71
2000.....	838	8	27	13	18	15	44	52	69	61	74	59
1999.....	866	8	22	18	11	24	57	59	82	63	76	72
All races, male												
2004.....	417	5	12	7	6	20	31	32	42	34	38	26
2003.....	377	8	11	5	5	10	25	29	31	29	38	41
2002.....	399	9	5	6	8	14	33	43	42	40	30	32
2001.....	400	5	10	4	4	18	25	29	48	28	34	38
2000.....	354	5	13	8	8	7	22	25	28	27	31	27
1999.....	408	6	11	7	6	14	37	31	42	33	39	35
All races, female												
2004.....	432	4	7	12	2	10	32	35	26	26	36	35
2003.....	445	8	6	7	8	9	33	30	33	32	33	35
2002.....	495	4	10	6	7	12	21	25	41	33	40	47
2001.....	453	3	6	8	8	9	19	21	28	33	53	33
2000.....	484	3	14	5	10	8	22	27	41	34	43	32
1999.....	458	2	11	11	5	10	20	28	40	30	37	37
White, both sexes												
2004.....	278	3	5	7	-	5	3	6	3	1	11	9
2003.....	300	6	5	1	3	4	4	8	8	6	5	9
2002.....	343	7	6	1	3	2	7	8	18	6	6	12
2001.....	331	2	7	2	2	4	4	7	12	5	7	5
2000.....	331	4	10	2	3	1	2	9	4	3	9	5
1999.....	316	2	2	2	1	4	5	4	3	1	12	13

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hemolytic anemias (D55-D59)												
White, male												
2004.....	128	1	2	4	-	3	2	1	3	1	7	4
2003.....	116	2	3	-	-	1	2	2	4	4	2	7
2002.....	136	4	1	-	2	1	3	6	9	5	2	6
2001.....	140	2	4	-	1	3	3	4	6	2	2	3
2000.....	131	2	5	1	1	-	1	5	1	1	5	1
1999.....	117	2	-	-	-	1	2	1	1	-	5	9
White, female												
2004.....	150	2	3	3	-	2	1	5	-	-	4	5
2003.....	184	4	2	1	3	3	2	6	4	2	3	2
2002.....	207	3	5	1	1	1	4	2	9	1	4	6
2001.....	191	-	3	2	1	1	1	3	6	3	5	2
2000.....	200	2	5	1	2	1	1	4	3	2	4	4
1999.....	199	-	2	2	1	3	3	3	2	1	7	4
Black, both sexes												
2004.....	562	3	14	11	8	25	60	60	65	59	63	52
2003.....	512	4	10	11	9	15	54	51	56	55	65	67
2002.....	538	2	9	10	12	23	47	59	64	66	64	67
2001.....	506	2	9	9	10	22	40	43	63	54	79	66
2000.....	497	3	17	11	15	14	41	43	65	58	65	52
1999.....	538	3	20	16	10	20	52	53	78	61	64	57
Black, male												
2004.....	284	2	10	3	6	17	29	30	39	33	31	22
2003.....	256	3	6	5	5	9	23	27	27	25	36	34
2002.....	257	2	4	6	6	13	30	36	32	34	28	26
2001.....	256	1	6	3	3	15	22	25	41	26	32	35
2000.....	219	3	8	7	7	7	20	20	27	26	26	25
1999.....	284	2	11	7	6	13	35	28	41	32	34	25

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hemolytic anemias (D55-D59)												
Black, female												
2004.....	278	1	4	8	2	8	31	30	26	26	32	30
2003.....	256	1	4	6	4	6	31	24	29	30	29	33
2002.....	281	-	5	4	6	10	17	23	32	32	36	41
2001.....	250	1	3	6	7	7	18	18	22	28	47	31
2000.....	278	-	9	4	8	7	21	23	38	32	39	27
1999.....	254	1	9	9	4	7	17	25	37	29	30	32
American Indian 1/, both sexes												
2004.....	1	-	-	1	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	2	-	-	-	-	-	-	1	-	-	-	-
2001.....	3	-	-	-	-	1	-	-	-	-	1	-
2000.....	1	-	-	-	-	-	-	-	-	-	-	-
1999.....	3	-	-	-	-	-	-	-	1	1	-	1
American Indian 1/, male												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	1	-	-	-	-	-	-	1	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	1	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, female												
2004.....	1	-	-	1	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	1	-	-	-	-	-	-	-	-	-	-	-
2001.....	3	-	-	-	-	1	-	-	-	-	1	-
2000.....	1	-	-	-	-	-	-	-	-	-	-	-
1999.....	2	-	-	-	-	-	-	-	1	-	-	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Aplastic anemias (D60-D61)

All races, both sexes

2004.....	967	5	9	6	13	17	13	11	6	13	25	29
2003.....	952	4	4	3	15	17	7	9	17	14	16	30
2002.....	944	1	5	16	8	14	11	10	11	19	17	27
2001.....	1,096	5	6	12	16	19	13	10	15	10	23	26
2000.....	1,066	1	7	10	9	9	11	6	16	21	19	29
1999.....	1,069	4	8	6	15	18	14	4	10	22	13	31

All races, male

2004.....	437	4	5	3	5	9	5	5	4	9	11	14
2003.....	457	2	2	2	8	8	5	8	12	5	8	17
2002.....	434	-	3	6	5	9	8	6	5	9	9	13
2001.....	462	5	2	7	10	7	7	9	6	7	8	9
2000.....	439	-	2	8	4	1	5	3	9	6	8	13
1999.....	479	2	6	1	7	9	6	1	5	11	7	16

All races, female

2004.....	530	1	4	3	8	8	8	6	2	4	14	15
2003.....	495	2	2	1	7	9	2	1	5	9	8	13
2002.....	510	1	2	10	3	5	3	4	6	10	8	14
2001.....	634	-	4	5	6	12	6	1	9	3	15	17
2000.....	627	1	5	2	5	8	6	3	7	15	11	16
1999.....	590	2	2	5	8	9	8	3	5	11	6	15

White, both sexes

2004.....	834	4	5	3	10	12	11	8	5	8	19	25
2003.....	810	1	3	2	11	11	2	7	13	9	12	23
2002.....	797	-	5	13	4	9	7	6	8	11	13	20
2001.....	958	2	4	8	14	12	10	4	10	6	19	22
2000.....	918	-	4	7	8	6	8	4	9	13	11	20
1999.....	922	4	7	5	11	13	8	2	7	18	10	24

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Aplastic anemias (D60-D61)												
White, male												
2004.....	378	3	4	1	3	6	5	4	4	6	9	13
2003.....	395	1	1	2	6	5	1	6	9	3	5	15
2002.....	367	-	3	5	3	6	5	3	4	5	8	9
2001.....	400	2	1	5	9	4	6	4	4	4	6	7
2000.....	373	-	2	6	4	1	4	3	5	5	5	10
1999.....	412	2	6	1	5	7	3	-	4	8	7	12
White, female												
2004.....	456	1	1	2	7	6	6	4	1	2	10	12
2003.....	415	-	2	-	5	6	1	1	4	6	7	8
2002.....	430	-	2	8	1	3	2	3	4	6	5	11
2001.....	558	-	3	3	5	8	4	-	6	2	13	15
2000.....	545	-	2	1	4	5	4	1	4	8	6	10
1999.....	510	2	1	4	6	6	5	2	3	10	3	12
Black, both sexes												
2004.....	98	1	3	1	3	5	2	2	1	3	5	1
2003.....	113	2	1	1	4	3	5	1	1	3	4	5
2002.....	114	1	-	2	2	4	4	3	1	6	2	5
2001.....	106	3	2	4	2	5	3	6	4	2	3	2
2000.....	113	1	2	3	1	2	3	2	7	5	5	9
1999.....	117	-	1	1	4	3	5	2	3	4	3	5
Black, male												
2004.....	45	1	1	1	2	3	-	1	-	2	2	-
2003.....	50	1	1	-	2	2	4	1	1	2	3	2
2002.....	57	-	-	1	1	3	3	2	1	3	1	3
2001.....	54	3	1	2	1	3	1	5	2	1	2	2
2000.....	48	-	-	2	-	-	1	-	4	-	2	3
1999.....	51	-	-	-	2	-	2	1	1	3	-	4

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute posthemorrhagic and other anemias (D62,D64)												
All races, both sexes												
2004.....	2,520	1	5	-	-	3	2	5	5	10	17	29
2003.....	2,528	2	3	1	3	1	2	1	6	5	11	27
2002.....	2,508	4	-	1	1	-	4	5	2	7	8	23
2001.....	2,396	2	-	1	2	2	4	5	4	12	18	15
2000.....	2,301	1	1	1	1	3	3	2	6	3	12	18
1999.....	2,281	-	-	-	3	1	6	1	4	5	15	29
All races, male												
2004.....	902	-	1	-	-	2	2	1	5	6	8	15
2003.....	891	1	1	1	2	1	2	-	2	2	5	18
2002.....	868	3	-	1	-	-	3	4	-	4	4	11
2001.....	813	-	-	1	1	1	1	2	1	7	7	10
2000.....	781	1	1	1	-	1	3	1	4	1	5	9
1999.....	760	-	-	-	2	1	-	1	1	3	8	16
All races, female												
2004.....	1,618	1	4	-	-	1	-	4	-	4	9	14
2003.....	1,637	1	2	-	1	-	-	1	4	3	6	9
2002.....	1,640	1	-	-	1	-	1	1	2	3	4	12
2001.....	1,583	2	-	-	1	1	3	3	3	5	11	5
2000.....	1,520	-	-	-	1	2	-	1	2	2	7	9
1999.....	1,521	-	-	-	1	-	6	-	3	2	7	13
White, both sexes												
2004.....	2,129	1	3	-	-	3	1	5	4	5	11	14
2003.....	2,118	2	2	1	3	-	2	1	2	1	7	13
2002.....	2,083	1	-	1	1	-	3	1	1	5	4	15
2001.....	2,008	1	-	1	-	1	2	4	2	5	10	4
2000.....	1,954	-	1	-	1	3	1	2	1	1	6	13
1999.....	1,910	-	-	-	1	1	3	-	1	3	4	16

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Acute posthemorrhagic and other anemias (D62,D64)

White, male

2004.....	745	-	-	-	-	2	1	1	4	2	6	9
2003.....	720	1	1	1	2	-	2	-	1	-	4	7
2002.....	695	-	-	1	-	2	1	1	-	4	2	9
2001.....	678	-	-	1	-	1	1	1	1	3	4	2
2000.....	652	-	1	-	-	1	1	1	-	1	4	5
1999.....	631	-	-	-	1	1	-	-	-	2	2	9

White, female

2004.....	1,384	1	3	-	-	1	-	4	-	3	5	5
2003.....	1,398	1	1	-	1	-	1	1	1	1	3	6
2002.....	1,388	1	-	-	1	-	1	-	1	1	2	6
2001.....	1,330	1	-	-	-	-	1	3	1	2	6	2
2000.....	1,302	-	-	-	1	2	-	1	1	-	2	8
1999.....	1,279	-	-	-	-	-	3	-	1	1	2	7

Black, both sexes

2004.....	349	-	2	-	-	-	1	-	1	5	3	14
2003.....	375	-	1	-	-	1	-	-	4	4	3	12
2002.....	387	3	-	-	-	-	1	3	-	2	4	8
2001.....	358	1	-	-	1	1	2	1	2	6	7	11
2000.....	321	1	-	1	-	-	2	-	5	2	6	5
1999.....	327	-	-	-	2	-	3	1	3	1	9	12

Black, male

2004.....	138	-	1	-	-	-	1	-	1	4	1	5
2003.....	164	-	-	-	-	1	-	-	1	2	1	11
2002.....	156	3	-	-	-	-	1	2	-	-	2	2
2001.....	124	-	-	-	1	-	-	1	-	3	2	8
2000.....	114	1	-	1	-	-	2	-	4	-	1	4
1999.....	109	-	-	-	1	-	-	1	1	-	4	6

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Coagulation defects, purpura and other hemorrhagic conditions (D65-D69)

All races, both sexes

2004.....	2,639	37	5	11	5	10	11	25	36	55	94	127
2003.....	2,657	41	9	2	6	9	24	24	36	47	89	129
2002.....	2,683	24	16	6	4	15	26	25	45	53	93	143
2001.....	2,632	34	12	7	6	12	17	25	51	56	104	129
2000.....	2,639	23	15	8	6	14	20	21	30	63	81	107
1999.....	2,601	43	11	13	8	16	21	20	39	66	98	112

All races, male

2004.....	1,216	24	2	7	1	4	6	15	15	27	42	68
2003.....	1,204	23	4	1	2	4	14	8	19	20	50	68
2002.....	1,188	13	7	1	3	8	11	14	27	22	47	59
2001.....	1,153	19	9	3	4	6	8	13	15	29	48	63
2000.....	1,157	13	7	6	4	7	6	11	10	26	43	65
1999.....	1,147	27	4	8	8	7	12	11	15	34	47	58

All races, female

2004.....	1,423	13	3	4	4	6	5	10	21	28	52	59
2003.....	1,453	18	5	1	4	5	10	16	17	27	39	61
2002.....	1,495	11	9	5	1	7	15	11	18	31	46	84
2001.....	1,479	15	3	4	2	6	9	12	36	27	56	66
2000.....	1,482	10	8	2	2	7	14	10	20	37	38	42
1999.....	1,454	16	7	5	-	9	9	9	24	32	51	54

White, both sexes

2004.....	2,175	26	3	7	3	5	7	22	22	38	59	94
2003.....	2,159	28	5	1	3	8	20	15	22	35	63	85
2002.....	2,239	19	9	5	2	15	17	20	32	43	71	103
2001.....	2,157	23	10	7	4	7	8	18	29	34	73	79
2000.....	2,179	18	10	4	4	12	10	8	17	47	55	80
1999.....	2,112	26	8	13	4	16	10	12	26	39	66	80

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Coagulation defects, purpura and other hemorrhagic conditions (D65-D69)

White, male

2004.....	1,014	15	2	6	1	2	4	13	9	21	29	50
2003.....	984	19	1	-	1	3	11	5	12	20	34	50
2002.....	1,000	12	4	1	1	8	9	11	19	19	35	46
2001.....	946	12	7	3	3	4	4	11	9	19	36	37
2000.....	965	10	5	4	3	6	4	4	7	19	32	50
1999.....	931	17	3	8	4	7	7	8	11	24	36	43

White, female

2004.....	1,161	11	1	1	2	3	3	9	13	17	30	44
2003.....	1,175	9	4	1	2	5	9	10	10	15	29	35
2002.....	1,239	7	5	4	1	7	8	9	13	24	36	57
2001.....	1,211	11	3	4	1	3	4	7	20	15	37	42
2000.....	1,214	8	5	-	1	6	6	4	10	28	23	30
1999.....	1,181	9	5	5	-	9	3	4	15	15	30	37

Black, both sexes

2004.....	410	10	1	2	2	5	4	3	11	17	32	30
2003.....	412	7	3	1	3	1	3	8	11	10	23	40
2002.....	382	4	5	1	2	-	6	5	12	7	16	36
2001.....	410	9	1	-	2	4	8	6	19	21	30	46
2000.....	404	5	4	3	2	2	10	10	12	15	22	25
1999.....	424	15	3	-	3	-	9	5	13	25	30	26

Black, male

2004.....	182	8	-	1	-	2	2	2	5	6	11	17
2003.....	178	3	2	1	1	1	2	3	6	-	13	16
2002.....	157	-	2	-	2	-	2	3	7	2	9	12
2001.....	168	5	1	-	1	1	4	2	6	10	12	22
2000.....	165	3	2	1	1	1	2	6	3	7	9	13
1999.....	184	9	1	-	3	-	5	3	4	10	11	12

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Coagulation defects, purpura and other hemorrhagic conditions (D65-D69)												
Black, female												
2004.....	228	2	1	1	2	3	2	1	6	11	21	13
2003.....	234	4	1	-	2	-	1	5	5	10	10	24
2002.....	225	4	3	1	-	-	4	2	5	5	7	24
2001.....	242	4	-	-	1	3	4	4	13	11	18	24
2000.....	239	2	2	2	1	1	8	4	9	8	13	12
1999.....	240	6	2	-	-	-	4	2	9	15	19	14
American Indian 1/, both sexes												
2004.....	13	-	-	1	-	-	-	-	-	-	3	3
2003.....	17	-	-	-	-	-	-	1	1	1	1	2
2002.....	15	-	-	-	-	-	2	-	-	1	3	1
2001.....	15	1	-	-	-	1	-	1	-	-	1	-
2000.....	12	-	-	-	-	-	-	1	-	-	2	-
1999.....	16	-	-	-	1	-	-	1	-	-	-	1
American Indian 1/, male												
2004.....	6	-	-	-	-	-	-	-	-	-	2	1
2003.....	7	-	-	-	-	-	-	-	-	-	1	1
2002.....	7	-	-	-	-	-	-	-	-	1	2	-
2001.....	8	1	-	-	-	1	-	-	-	-	-	-
2000.....	6	-	-	-	-	-	-	1	-	-	2	-
1999.....	8	-	-	-	1	-	-	-	-	-	-	1
American Indian 1/, female												
2004.....	7	-	-	1	-	-	-	-	-	-	1	2
2003.....	10	-	-	-	-	-	-	1	1	1	-	1
2002.....	8	-	-	-	-	-	2	-	-	-	1	1
2001.....	7	-	-	-	-	-	-	1	-	-	1	-
2000.....	6	-	-	-	-	-	-	-	-	-	-	-
1999.....	8	-	-	-	-	-	-	1	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Coagulation defects, purpura and other hemorrhagic conditions (D65-D69)												
Asian or Pacific Islander 2/, both sexes												
2004.....	41	1	1	1	-	-	-	-	3	-	-	-
2003.....	69	6	1	-	-	-	1	-	2	1	2	2
2002.....	47	1	2	-	-	-	1	-	1	2	3	3
2001.....	50	1	1	-	-	-	1	-	3	1	-	4
2000.....	44	-	1	1	-	-	-	2	1	1	2	2
1999.....	49	2	-	-	-	-	2	2	-	2	2	5
Asian or Pacific Islander 2/, male												
2004.....	14	1	-	-	-	-	-	-	1	-	-	-
2003.....	35	1	1	-	-	-	1	-	1	-	2	1
2002.....	24	1	1	-	-	-	-	-	1	-	1	1
2001.....	31	1	1	-	-	-	-	-	-	-	-	4
2000.....	21	-	-	1	-	-	-	-	-	-	-	2
1999.....	24	1	-	-	-	-	-	-	-	-	-	2
Asian or Pacific Islander 2/, female												
2004.....	27	-	1	1	-	-	-	-	2	-	-	-
2003.....	34	5	-	-	-	-	-	-	1	1	-	1
2002.....	23	-	1	-	-	-	1	-	-	2	2	2
2001.....	19	-	-	-	-	-	1	-	3	1	-	-
2000.....	23	-	1	-	-	-	-	2	1	1	2	-
1999.....	25	1	-	-	-	-	2	2	-	2	2	3

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other diseases of blood and blood-forming organs (D70-D76)

All races, both sexes

2004.....	811	24	23	6	11	12	6	10	9	15	22	32
2003.....	860	22	22	6	16	7	5	7	14	19	20	26
2002.....	892	19	27	4	4	6	6	11	7	12	30	30
2001.....	846	25	22	3	3	6	7	8	7	11	29	23
2000.....	836	18	16	3	4	11	9	10	16	18	25	33
1999.....	806	21	16	4	6	10	10	14	9	21	27	22

All races, male

2004.....	394	16	13	4	6	4	5	5	3	8	14	17
2003.....	402	14	11	3	10	6	4	3	6	11	11	16
2002.....	435	13	20	3	4	4	2	7	5	6	15	16
2001.....	385	11	12	2	3	2	4	4	4	5	14	17
2000.....	398	9	7	1	2	6	6	5	11	13	12	20
1999.....	394	11	11	-	4	7	7	11	6	16	12	12

All races, female

2004.....	417	8	10	2	5	8	1	5	6	7	8	15
2003.....	458	8	11	3	6	1	1	4	8	8	9	10
2002.....	457	6	7	1	-	2	4	4	2	6	15	14
2001.....	461	14	10	1	-	4	3	4	3	6	15	6
2000.....	438	9	9	2	2	5	3	5	5	5	13	13
1999.....	412	10	5	4	2	3	3	3	3	5	15	10

White, both sexes

2004.....	708	15	18	5	11	10	6	10	5	10	17	24
2003.....	760	19	16	5	13	6	4	5	11	16	16	21
2002.....	778	13	23	3	3	6	5	8	5	7	25	25
2001.....	733	19	15	3	3	4	7	5	6	9	22	20
2000.....	719	15	15	2	3	8	8	5	10	13	19	25
1999.....	696	10	13	4	6	10	7	11	7	14	23	16

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other diseases of blood and blood-forming organs (D70-D76)

White, male

2004.....	345	10	10	4	6	3	5	5	2	4	13	12
2003.....	356	12	8	3	8	5	3	2	5	10	9	13
2002.....	376	11	16	2	3	4	2	6	4	4	12	13
2001.....	336	8	8	2	3	2	4	2	4	4	12	16
2000.....	337	7	7	-	2	4	6	3	6	10	10	16
1999.....	338	6	9	-	4	7	5	8	4	11	9	9

White, female

2004.....	363	5	8	1	5	7	1	5	3	6	4	12
2003.....	404	7	8	2	5	1	1	3	6	6	7	8
2002.....	402	2	7	1	-	2	3	2	1	3	13	12
2001.....	397	11	7	1	-	2	3	3	2	5	10	4
2000.....	382	8	8	2	1	4	2	2	4	3	9	9
1999.....	358	4	4	4	2	3	2	3	3	3	14	7

Black, both sexes

2004.....	76	7	2	-	-	1	-	-	3	3	4	7
2003.....	80	3	3	-	2	1	1	2	2	3	4	5
2002.....	94	4	2	1	1	-	1	3	2	5	4	4
2001.....	81	2	4	-	-	2	-	-	1	1	5	3
2000.....	95	2	1	-	1	2	-	5	4	5	6	7
1999.....	96	9	3	-	-	-	2	3	2	6	4	5

Black, male

2004.....	37	4	2	-	-	1	-	-	-	3	1	4
2003.....	36	2	2	-	1	1	1	1	1	1	2	3
2002.....	48	1	2	1	1	-	-	1	1	2	2	2
2001.....	34	1	2	-	-	-	-	-	-	-	2	1
2000.....	48	2	-	-	-	1	-	2	3	3	2	4
1999.....	48	4	2	-	-	-	1	3	2	5	3	3

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of blood and blood-forming organs (D70-D76)												
Asian or Pacific Islander 2/, both sexes												
2004.....	23	-	3	1	-	1	-	-	1	2	1	1
2003.....	15	-	2	1	1	-	-	-	1	-	-	-
2002.....	17	2	1	-	-	-	-	-	-	-	1	1
2001.....	26	3	2	-	-	-	-	3	-	1	1	-
2000.....	13	-	-	1	-	-	-	-	-	-	-	1
1999.....	11	1	-	-	-	-	-	-	-	1	-	1
Asian or Pacific Islander 2/, male												
2004.....	10	-	1	-	-	-	-	-	1	1	-	1
2003.....	8	-	-	-	1	-	-	-	-	-	-	-
2002.....	9	1	1	-	-	-	-	-	-	-	1	1
2001.....	14	2	1	-	-	-	-	2	-	1	-	-
2000.....	7	-	-	1	-	-	-	-	-	-	-	-
1999.....	6	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	13	-	2	1	-	1	-	-	-	1	1	-
2003.....	7	-	2	1	-	-	-	-	1	-	-	-
2002.....	8	1	-	-	-	-	-	-	-	-	-	-
2001.....	12	1	1	-	-	-	-	1	-	-	1	-
2000.....	6	-	-	-	-	-	-	-	-	-	-	1
1999.....	5	1	-	-	-	-	-	-	-	1	-	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Certain disorders involving the immune mechanism (D80-D89)

All races, both sexes

2004.....	1,399	18	7	8	7	10	11	26	43	80	143	159
2003.....	1,474	20	18	8	7	7	17	20	54	109	165	153
2002.....	1,472	21	13	2	8	9	10	17	44	106	170	170
2001.....	1,430	31	17	5	8	6	11	14	65	110	152	183
2000.....	1,346	19	16	8	8	9	7	26	57	120	144	167
1999.....	1,157	18	25	8	7	11	7	24	66	104	131	139

All races, male

2004.....	624	10	5	4	6	7	8	13	24	43	80	87
2003.....	647	12	10	4	7	3	8	6	28	63	86	79
2002.....	661	14	4	-	4	9	6	10	27	55	76	90
2001.....	626	18	14	3	7	3	7	9	46	66	75	77
2000.....	628	12	10	7	5	7	6	15	33	67	71	82
1999.....	539	12	18	5	4	5	2	15	40	62	60	73

All races, female

2004.....	775	8	2	4	1	3	3	13	19	37	63	72
2003.....	827	8	8	4	-	4	9	14	26	46	79	74
2002.....	811	7	9	2	4	-	4	7	17	51	94	80
2001.....	804	13	3	2	1	3	4	5	19	44	77	106
2000.....	718	7	6	1	3	2	1	11	24	53	73	85
1999.....	618	6	7	3	3	6	5	9	26	42	71	66

White, both sexes

2004.....	704	13	6	3	7	7	9	10	10	24	49	52
2003.....	713	15	13	6	4	4	9	7	16	35	41	58
2002.....	748	18	7	2	6	5	4	6	16	33	40	58
2001.....	723	24	16	4	5	4	5	6	25	34	34	58
2000.....	688	13	10	6	7	6	3	5	20	37	51	48
1999.....	599	11	16	4	4	7	3	9	25	28	36	51

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Certain disorders involving the immune mechanism (D80-D89)

White, male

2004.....	331	6	4	2	6	6	8	6	6	17	31	38
2003.....	327	10	6	4	4	2	5	3	9	22	29	32
2002.....	353	13	3	-	2	5	2	2	10	25	21	38
2001.....	340	14	14	3	5	2	2	5	21	24	22	34
2000.....	347	7	7	5	4	5	3	3	16	23	30	29
1999.....	308	6	11	3	2	3	1	8	18	22	24	34

White, female

2004.....	373	7	2	1	1	1	1	4	4	7	18	14
2003.....	386	5	7	2	-	2	4	4	7	13	12	26
2002.....	395	5	4	2	4	-	2	4	6	8	19	20
2001.....	383	10	2	1	-	2	3	1	4	10	12	24
2000.....	341	6	3	1	3	1	-	2	4	14	21	19
1999.....	291	5	5	1	2	4	2	1	7	6	12	17

Black, both sexes

2004.....	675	2	1	5	-	1	2	16	32	55	93	106
2003.....	742	4	5	1	2	3	8	13	36	74	123	93
2002.....	708	3	5	-	1	3	5	11	27	73	129	110
2001.....	688	6	-	-	2	2	6	8	40	76	116	124
2000.....	644	4	5	2	1	1	4	20	37	83	92	116
1999.....	540	6	7	3	1	4	4	15	41	75	94	85

Black, male

2004.....	286	2	1	2	-	-	-	7	17	25	49	49
2003.....	309	2	4	-	2	1	3	3	17	41	56	47
2002.....	301	1	1	-	1	3	4	8	16	30	54	51
2001.....	279	3	-	-	2	1	5	4	25	42	52	42
2000.....	272	3	2	2	1	1	3	11	17	44	41	51
1999.....	222	5	5	2	1	2	1	7	22	39	36	37

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Certain disorders involving the immune mechanism (D80-D89)												
Black, female												
2004.....	389	-	-	3	-	1	2	9	15	30	44	57
2003.....	433	2	1	1	-	2	5	10	19	33	67	46
2002.....	407	2	4	-	-	-	1	3	11	43	75	59
2001.....	409	3	-	-	-	1	1	4	15	34	64	82
2000.....	372	1	3	-	-	-	1	9	20	39	51	65
1999.....	318	1	2	1	-	2	3	8	19	36	58	48
American Indian 1/, both sexes												
2004.....	7	-	-	-	-	1	-	-	1	1	1	-
2003.....	5	-	-	-	1	-	-	-	-	-	-	1
2002.....	7	-	1	-	1	1	-	-	-	-	-	1
2001.....	9	1	1	-	1	-	-	-	-	-	-	1
2000.....	7	1	-	-	-	1	-	-	-	-	1	2
1999.....	7	1	1	1	2	-	-	-	-	-	1	1
American Indian 1/, male												
2004.....	3	-	-	-	-	-	-	-	1	1	-	-
2003.....	4	-	-	-	1	-	-	-	-	-	-	-
2002.....	3	-	-	-	1	1	-	-	-	-	-	1
2001.....	4	1	-	-	-	-	-	-	-	-	-	1
2000.....	2	1	-	-	-	-	-	-	-	-	-	1
1999.....	4	1	1	-	1	-	-	-	-	-	-	1
American Indian 1/, female												
2004.....	4	-	-	-	-	1	-	-	-	-	1	-
2003.....	1	-	-	-	-	-	-	-	-	-	-	1
2002.....	4	-	1	-	-	-	-	-	-	-	-	-
2001.....	5	-	1	-	1	-	-	-	-	-	-	-
2000.....	5	-	-	-	-	1	-	-	-	-	1	1
1999.....	3	-	-	1	1	-	-	-	-	-	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
IV. Endocrine, nutritional and metabolic diseases (E00-E88)												
All races, both sexes												
2004.....	99,042	233	144	95	125	177	343	438	745	1,155	1,975	3,228
2003.....	100,443	248	190	91	119	207	332	503	745	1,169	2,029	3,219
2002.....	99,458	266	145	91	129	188	349	477	767	1,217	2,119	2,976
2001.....	96,820	292	183	94	124	195	303	430	661	1,184	1,819	2,903
2000.....	94,345	260	160	77	134	204	303	447	698	1,120	1,881	2,675
1999.....	92,872	261	153	97	120	223	301	422	642	1,126	1,795	2,610
All races, male												
2004.....	46,499	149	78	47	69	95	181	244	408	679	1,175	1,921
2003.....	46,520	140	109	59	68	113	179	274	450	646	1,231	1,887
2002.....	45,132	152	66	58	74	98	193	264	465	695	1,303	1,709
2001.....	43,188	166	92	45	72	101	153	229	377	679	1,076	1,716
2000.....	41,720	144	87	42	78	107	169	257	424	629	1,091	1,580
1999.....	40,789	141	86	48	64	114	142	215	369	657	1,038	1,556
All races, female												
2004.....	52,543	84	66	48	56	82	162	194	337	476	800	1,307
2003.....	53,923	108	81	32	51	94	153	229	295	523	798	1,332
2002.....	54,326	114	79	33	55	90	156	213	302	522	816	1,267
2001.....	53,632	126	91	49	52	94	150	201	284	505	743	1,187
2000.....	52,625	116	73	35	56	97	134	190	274	491	790	1,095
1999.....	52,083	120	67	49	56	109	159	207	273	469	757	1,054
White, both sexes												
2004.....	80,256	184	118	77	95	136	264	311	510	788	1,432	2,397
2003.....	81,573	169	151	80	95	159	257	335	526	824	1,491	2,348
2002.....	80,833	194	117	74	103	152	251	342	540	878	1,564	2,173
2001.....	79,016	213	149	77	97	151	223	297	468	843	1,344	2,102
2000.....	77,123	186	133	64	113	167	229	298	493	819	1,417	1,943
1999.....	75,648	179	127	72	97	182	232	316	452	797	1,312	1,876

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

IV. Endocrine, nutritional and metabolic diseases (E00-E88)

White, male

2004.....	38,277	117	63	36	52	69	144	173	282	474	878	1,460
2003.....	38,456	101	86	52	58	89	142	176	325	475	931	1,427
2002.....	37,344	122	52	52	62	75	134	190	328	512	989	1,283
2001.....	35,740	116	77	38	56	79	113	154	267	483	836	1,270
2000.....	34,725	103	75	35	69	86	129	181	309	467	850	1,195
1999.....	33,704	96	72	37	49	95	108	166	257	477	783	1,157

White, female

2004.....	41,979	67	55	41	43	67	120	138	228	314	554	937
2003.....	43,117	68	65	28	37	70	115	159	201	349	560	921
2002.....	43,489	72	65	22	41	77	117	152	212	366	575	890
2001.....	43,276	97	72	39	41	72	110	143	201	360	508	832
2000.....	42,398	83	58	29	44	81	100	117	184	352	567	748
1999.....	41,944	83	55	35	48	87	124	150	195	320	529	719

Black, both sexes

2004.....	16,118	39	17	9	23	36	74	116	214	339	487	735
2003.....	16,145	65	25	8	19	38	67	148	199	312	480	769
2002.....	16,075	56	21	14	25	32	89	130	209	311	504	721
2001.....	15,479	67	20	13	22	34	73	120	178	309	435	707
2000.....	15,146	55	20	11	18	35	69	143	186	283	426	667
1999.....	14,991	69	21	20	21	33	60	99	176	300	448	658

Black, male

2004.....	6,936	26	10	7	13	23	35	64	116	187	264	396
2003.....	6,755	30	15	5	8	18	32	87	112	152	263	404
2002.....	6,576	24	9	5	11	20	53	70	123	164	281	379
2001.....	6,358	39	9	5	12	17	35	66	102	176	218	387
2000.....	6,018	31	10	6	7	19	37	71	104	149	218	346
1999.....	6,048	39	10	8	14	16	30	44	102	162	232	355

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

IV. Endocrine, nutritional and metabolic diseases (E00-E88)

Black, female

2004.....	9,182	13	7	2	10	13	39	52	98	152	223	339
2003.....	9,390	35	10	3	11	20	35	61	87	160	217	365
2002.....	9,499	32	12	9	14	12	36	60	86	147	223	342
2001.....	9,121	28	11	8	10	17	38	54	76	133	217	320
2000.....	9,128	24	10	5	11	16	32	72	82	134	208	321
1999.....	8,943	30	11	12	7	17	30	55	74	138	216	303

American Indian 1/, both sexes

2004.....	874	1	3	2	2	-	2	3	13	13	35	50
2003.....	918	4	4	1	1	2	2	11	10	21	29	56
2002.....	882	5	2	1	-	1	5	2	6	17	27	49
2001.....	779	-	2	1	2	3	4	5	3	12	19	49
2000.....	734	7	5	-	1	2	1	2	10	9	20	30
1999.....	829	6	-	1	-	5	3	5	9	15	18	40

American Indian 1/, male

2004.....	421	-	1	-	1	-	-	2	6	10	21	34
2003.....	418	2	3	1	-	1	2	8	7	11	18	29
2002.....	399	1	2	-	-	1	3	2	5	11	17	29
2001.....	340	-	1	-	2	2	2	4	1	8	7	27
2000.....	323	2	2	-	1	2	1	2	8	6	12	19
1999.....	370	2	-	1	-	1	3	3	6	10	13	25

American Indian 1/, female

2004.....	453	1	2	2	1	-	2	1	7	3	14	16
2003.....	500	2	1	-	1	1	-	3	3	10	11	27
2002.....	483	4	-	1	-	-	2	-	1	6	10	20
2001.....	439	-	1	1	-	1	2	1	2	4	12	22
2000.....	411	5	3	-	-	-	-	-	2	3	8	11
1999.....	459	4	-	-	-	4	-	2	3	5	5	15

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

IV. Endocrine, nutritional and metabolic diseases (E00-E88)

Asian or Pacific Islander 2/, both sexes

2004.....	1,794	9	6	7	5	5	3	8	8	15	21	46
2003.....	1,807	10	10	2	4	8	6	9	10	12	29	46
2002.....	1,668	11	5	2	1	3	4	3	12	11	24	33
2001.....	1,546	12	12	3	3	7	3	8	12	20	21	45
2000.....	1,342	12	2	2	2	-	4	4	9	9	18	35
1999.....	1,404	7	5	4	2	3	6	2	5	14	17	36

Asian or Pacific Islander 2/, male

2004.....	865	6	4	4	3	3	2	5	4	8	12	31
2003.....	891	7	5	1	2	5	3	3	6	8	19	27
2002.....	813	5	3	1	1	2	3	2	9	8	16	18
2001.....	750	11	5	2	2	3	3	5	7	12	15	32
2000.....	654	8	-	1	1	-	2	3	3	7	11	20
1999.....	667	4	4	2	1	2	1	2	4	8	10	19

Asian or Pacific Islander 2/, female

2004.....	929	3	2	3	2	2	1	3	4	7	9	15
2003.....	916	3	5	1	2	3	3	6	4	4	10	19
2002.....	855	6	2	1	-	1	1	1	3	3	8	15
2001.....	796	1	7	1	1	4	-	3	5	8	6	13
2000.....	688	4	2	1	1	-	2	1	6	2	7	15
1999.....	737	3	1	2	1	1	5	-	1	6	7	17

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Endocrine diseases (E00-E34)												
All races, both sexes												
2004.....	75,885	26	20	15	38	59	139	215	441	745	1,372	2,292
2003.....	77,135	30	20	15	26	55	137	255	462	745	1,393	2,321
2002.....	76,159	40	12	8	31	64	139	248	446	807	1,439	2,171
2001.....	74,280	44	18	10	38	55	115	230	404	776	1,284	2,119
2000.....	71,987	45	22	14	22	58	121	235	444	732	1,301	1,973
1999.....	71,109	39	21	13	26	62	122	217	414	771	1,260	2,016
All races, male												
2004.....	36,136	19	11	9	20	34	69	127	245	461	837	1,391
2003.....	36,234	14	15	10	11	33	80	132	274	430	871	1,387
2002.....	35,095	23	3	5	20	35	78	139	274	462	899	1,271
2001.....	33,698	27	8	4	25	30	66	119	239	457	771	1,260
2000.....	32,325	25	12	6	10	36	67	128	274	414	774	1,148
1999.....	31,928	15	11	5	12	30	62	107	234	454	723	1,211
All races, female												
2004.....	39,749	7	9	6	18	25	70	88	196	284	535	901
2003.....	40,901	16	5	5	15	22	57	123	188	315	522	934
2002.....	41,064	17	9	3	11	29	61	109	172	345	540	900
2001.....	40,582	17	10	6	13	25	49	111	165	319	513	859
2000.....	39,662	20	10	8	12	22	54	107	170	318	527	825
1999.....	39,181	24	10	8	14	32	60	110	180	317	537	805
White, both sexes												
2004.....	60,399	21	15	10	19	36	88	143	289	504	978	1,669
2003.....	61,610	18	16	12	17	37	95	151	316	510	998	1,636
2002.....	60,986	33	11	7	18	43	89	163	310	570	1,035	1,536
2001.....	59,698	34	13	6	24	32	67	135	276	540	928	1,513
2000.....	57,849	29	17	9	14	41	78	142	309	520	966	1,389
1999.....	56,894	29	16	9	16	46	82	155	281	540	902	1,412

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Endocrine diseases (E00-E34)

White, male

2004.....	29,324	14	8	5	11	19	48	83	164	317	613	1,023
2003.....	29,592	7	12	8	8	23	56	71	193	300	645	1,014
2002.....	28,777	21	3	5	15	22	50	94	190	334	663	922
2001.....	27,632	18	6	3	18	21	40	66	158	314	592	916
2000.....	26,603	17	9	4	7	27	46	81	201	295	596	829
1999.....	26,165	9	10	4	6	25	44	82	159	328	526	877

White, female

2004.....	31,075	7	7	5	8	17	40	60	125	187	365	646
2003.....	32,018	11	4	4	9	14	39	80	123	210	353	622
2002.....	32,209	12	8	2	3	21	39	69	120	236	372	614
2001.....	32,066	16	7	3	6	11	27	69	118	226	336	597
2000.....	31,246	12	8	5	7	14	32	61	108	225	370	560
1999.....	30,729	20	6	5	10	21	38	73	122	212	376	535

Black, both sexes

2004.....	13,215	4	3	5	15	22	48	64	137	222	350	542
2003.....	13,247	10	4	3	7	17	40	94	132	213	354	605
2002.....	13,021	5	1	1	13	18	48	81	129	221	366	562
2001.....	12,645	9	3	4	11	20	42	90	121	213	328	532
2000.....	12,386	11	4	5	8	16	42	90	124	197	304	528
1999.....	12,302	9	5	3	9	16	34	58	123	211	326	542

Black, male

2004.....	5,712	4	1	4	7	15	20	38	76	131	197	315
2003.....	5,552	6	3	2	3	10	22	57	72	116	201	325
2002.....	5,312	1	-	-	5	10	27	42	79	119	210	307
2001.....	5,177	8	2	1	5	8	22	51	76	128	166	300
2000.....	4,884	4	3	2	3	8	20	45	65	106	159	283
1999.....	4,898	5	1	1	6	5	16	23	68	114	175	300

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Endocrine diseases (E00-E34)

Black, female

2004.....	7,503	-	2	1	8	7	28	26	61	91	153	227
2003.....	7,695	4	1	1	4	7	18	37	60	97	153	280
2002.....	7,709	4	1	1	8	8	21	39	50	102	156	255
2001.....	7,468	1	1	3	6	12	20	39	45	85	162	232
2000.....	7,502	7	1	3	5	8	22	45	59	91	145	245
1999.....	7,404	4	4	2	3	11	18	35	55	97	151	242

American Indian 1/, both sexes

2004.....	762	-	1	-	-	-	1	1	11	9	29	41
2003.....	792	-	-	-	-	-	-	7	7	18	24	46
2002.....	757	-	-	-	-	1	2	2	1	10	23	44
2001.....	659	-	1	-	1	2	4	2	2	8	11	40
2000.....	624	1	-	-	-	1	1	-	7	7	16	25
1999.....	735	-	-	-	-	-	2	2	7	13	16	35

American Indian 1/, male

2004.....	371	-	1	-	-	-	-	1	4	8	18	27
2003.....	369	-	-	-	-	-	-	4	5	10	16	25
2002.....	343	-	-	-	-	1	1	2	1	6	15	27
2001.....	281	-	-	-	1	1	2	1	1	6	2	21
2000.....	278	-	-	-	-	1	1	-	6	6	10	18
1999.....	325	-	-	-	-	-	2	-	5	8	12	21

American Indian 1/, female

2004.....	391	-	-	-	-	-	1	-	7	1	11	14
2003.....	423	-	-	-	-	-	-	3	2	8	8	21
2002.....	414	-	-	-	-	-	1	-	-	4	8	17
2001.....	378	-	1	-	-	1	2	1	1	2	9	19
2000.....	346	1	-	-	-	-	-	-	1	1	6	7
1999.....	410	-	-	-	-	-	-	2	2	5	4	14

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Endocrine diseases (E00-E34)												
Asian or Pacific Islander 2/, both sexes												
2004.....	1,509	1	1	-	4	1	2	7	4	10	15	40
2003.....	1,486	2	-	-	2	1	2	3	7	4	17	34
2002.....	1,395	2	-	-	-	2	-	2	6	6	15	29
2001.....	1,278	1	1	-	2	1	2	3	5	15	17	34
2000.....	1,128	4	1	-	-	-	-	3	4	8	15	31
1999.....	1,178	1	-	1	1	-	4	2	3	7	16	27
Asian or Pacific Islander 2/, male												
2004.....	729	1	1	-	2	-	1	5	1	5	9	26
2003.....	721	1	-	-	-	-	2	-	4	4	9	23
2002.....	663	1	-	-	-	2	-	1	4	3	11	15
2001.....	608	1	-	-	1	-	2	1	4	9	11	23
2000.....	560	4	-	-	-	-	-	2	2	7	9	18
1999.....	540	1	-	-	-	-	-	2	2	4	10	13
Asian or Pacific Islander 2/, female												
2004.....	780	-	-	-	2	1	1	2	3	5	6	14
2003.....	765	1	-	-	2	1	-	3	3	-	8	11
2002.....	732	1	-	-	-	-	-	1	2	3	4	14
2001.....	670	-	1	-	1	1	-	2	1	6	6	11
2000.....	568	-	1	-	-	-	-	1	2	1	6	13
1999.....	638	-	-	1	1	-	4	-	1	3	6	14

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Disorders of thyroid gland (E00-E07)

All races, both sexes

2004.....	1,777	3	2	1	2	2	10	16	13	19	21	37
2003.....	1,957	4	1	-	1	-	13	14	15	11	24	34
2002.....	1,942	1	-	1	1	1	8	9	16	16	20	29
2001.....	1,918	1	-	1	-	4	1	5	14	20	20	25
2000.....	1,780	3	-	-	1	2	2	20	12	27	27	19
1999.....	1,811	2	1	-	-	3	4	13	8	15	21	39

All races, male

2004.....	441	-	-	1	1	-	6	12	6	7	8	15
2003.....	390	3	1	-	-	-	6	6	5	3	6	10
2002.....	408	1	-	-	1	-	4	5	4	3	7	8
2001.....	427	1	-	-	-	1	1	1	7	8	5	10
2000.....	376	1	-	-	1	-	-	8	9	12	6	5
1999.....	403	1	-	-	-	2	-	5	3	5	5	12

All races, female

2004.....	1,336	3	2	-	1	2	4	4	7	12	13	22
2003.....	1,567	1	-	-	1	-	7	8	10	8	18	24
2002.....	1,534	-	-	1	-	1	4	4	12	13	13	21
2001.....	1,491	-	-	1	-	3	-	4	7	12	15	15
2000.....	1,404	2	-	-	-	2	2	12	3	15	21	14
1999.....	1,408	1	1	-	-	1	4	8	5	10	16	27

White, both sexes

2004.....	1,542	3	1	1	1	1	5	10	5	13	14	22
2003.....	1,740	2	1	-	1	-	5	8	7	8	18	23
2002.....	1,733	1	-	1	1	1	4	3	10	8	11	19
2001.....	1,689	-	-	1	-	2	1	4	8	14	8	18
2000.....	1,554	2	-	-	1	1	1	7	7	12	15	13
1999.....	1,581	2	1	-	-	2	2	7	1	8	13	20

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Disorders of thyroid gland (E00-E07)

White, male

2004.....	371	-	-	1	1	-	4	8	2	5	4	8
2003.....	337	2	1	-	-	-	2	4	2	1	6	5
2002.....	357	1	-	-	1	-	4	2	3	1	5	6
2001.....	370	-	-	-	-	1	1	1	5	4	2	7
2000.....	314	1	-	-	1	-	-	3	5	5	4	5
1999.....	336	1	-	-	-	1	-	3	1	3	2	7

White, female

2004.....	1,171	3	1	-	-	1	1	2	3	8	10	14
2003.....	1,403	-	-	-	1	-	3	4	5	7	12	18
2002.....	1,376	-	-	1	-	1	-	1	7	7	6	13
2001.....	1,319	-	-	1	-	1	-	3	3	10	6	11
2000.....	1,240	1	-	-	-	1	1	4	2	7	11	8
1999.....	1,245	1	1	-	-	1	2	4	-	5	11	13

Black, both sexes

2004.....	207	-	1	-	1	1	5	5	8	6	6	13
2003.....	192	1	-	-	-	-	8	6	7	1	6	10
2002.....	183	-	-	-	-	-	4	5	5	7	8	10
2001.....	197	1	-	-	-	2	-	1	5	5	11	6
2000.....	205	-	-	-	-	1	1	13	5	14	12	6
1999.....	205	-	-	-	-	1	2	5	6	7	7	17

Black, male

2004.....	58	-	-	-	-	-	2	3	4	2	3	5
2003.....	47	-	-	-	-	-	4	2	3	-	-	5
2002.....	42	-	-	-	-	-	-	2	1	2	2	2
2001.....	53	1	-	-	-	-	-	-	2	4	3	3
2000.....	56	-	-	-	-	-	-	5	4	6	2	-
1999.....	54	-	-	-	-	1	-	1	1	2	2	4

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diabetes mellitus (E10-E14)												
All races, both sexes												
2004.....	73,138	1	8	6	31	48	113	180	419	709	1,317	2,217
2003.....	74,219	3	8	10	21	46	114	227	430	715	1,334	2,245
2002.....	73,249	1	3	5	29	55	116	229	413	776	1,388	2,106
2001.....	71,372	2	6	7	32	48	103	215	380	735	1,223	2,063
2000.....	69,301	4	6	8	18	48	114	206	417	684	1,242	1,917
1999.....	68,399	2	8	9	23	51	104	189	393	730	1,212	1,942
All races, male												
2004.....	35,267	1	5	1	15	30	53	103	237	444	814	1,359
2003.....	35,438	-	6	6	10	28	68	119	260	420	844	1,354
2002.....	34,301	-	1	4	18	30	68	128	257	452	878	1,248
2001.....	32,841	1	2	2	21	27	58	114	227	436	745	1,234
2000.....	31,602	4	2	5	8	31	63	118	258	396	755	1,124
1999.....	31,150	-	4	4	12	25	55	94	225	437	702	1,189
All races, female												
2004.....	37,871	-	3	5	16	18	60	77	182	265	503	858
2003.....	38,781	3	2	4	11	18	46	108	170	295	490	891
2002.....	38,948	1	2	1	11	25	48	101	156	324	510	858
2001.....	38,531	1	4	5	11	21	45	101	153	299	478	829
2000.....	37,699	-	4	3	10	17	51	88	159	288	487	793
1999.....	37,249	2	4	5	11	26	49	95	168	293	510	753
White, both sexes												
2004.....	58,087	-	6	4	15	28	71	118	278	484	937	1,621
2003.....	59,099	2	6	9	13	29	82	135	296	487	952	1,583
2002.....	58,459	1	3	4	16	36	73	152	287	549	1,005	1,489
2001.....	57,180	1	6	3	18	27	55	124	259	511	888	1,474
2000.....	55,561	3	6	5	12	33	72	128	292	493	926	1,344
1999.....	54,599	2	6	8	14	38	69	137	272	515	869	1,367

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diabetes mellitus (E10-E14)												
White, male												
2004.....	28,629	-	4	-	7	15	37	66	161	308	597	1,005
2003.....	28,939	-	5	5	8	18	49	63	183	295	622	994
2002.....	28,110	-	1	4	13	18	42	88	178	327	651	906
2001.....	26,917	-	2	1	14	18	32	63	149	302	574	898
2000.....	26,009	3	2	3	6	22	42	76	191	285	581	808
1999.....	25,545	-	3	4	6	21	38	72	154	318	511	864
White, female												
2004.....	29,458	-	2	4	8	13	34	52	117	176	340	616
2003.....	30,160	2	1	4	5	11	33	72	113	192	330	589
2002.....	30,349	1	2	-	3	18	31	64	109	222	354	583
2001.....	30,263	1	4	2	4	9	23	61	110	209	314	576
2000.....	29,552	-	4	2	6	11	30	52	101	208	345	536
1999.....	29,054	2	3	4	8	17	31	65	118	197	358	503
Black, both sexes												
2004.....	12,834	1	1	2	14	20	41	56	127	208	339	519
2003.....	12,892	1	2	1	6	16	30	85	123	208	341	583
2002.....	12,687	-	-	1	13	17	42	74	121	212	348	545
2001.....	12,305	1	-	4	11	18	42	86	116	203	308	517
2000.....	12,021	1	-	3	6	14	41	75	114	177	285	517
1999.....	11,927	-	2	1	9	13	31	49	113	195	312	515
Black, male												
2004.....	5,565	1	-	1	7	15	16	33	71	125	192	304
2003.....	5,425	-	1	1	2	10	17	53	69	113	197	312
2002.....	5,207	-	-	-	5	10	26	38	75	116	203	301
2001.....	5,049	1	-	1	5	8	22	49	74	120	158	292
2000.....	4,771	1	-	2	2	8	20	40	59	99	155	280
1999.....	4,759	-	1	-	6	4	15	21	65	107	170	292

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diabetes mellitus (E10-E14)												
Black, female												
2004.....	7,269	-	1	1	7	5	25	23	56	83	147	215
2003.....	7,467	1	1	-	4	6	13	32	54	95	144	271
2002.....	7,480	-	-	1	8	7	16	36	46	96	145	244
2001.....	7,256	-	-	3	6	10	20	37	42	83	150	225
2000.....	7,250	-	-	1	4	6	21	35	55	78	130	237
1999.....	7,168	-	1	1	3	9	16	28	48	88	142	223
American Indian 1/, both sexes												
2004.....	746	-	-	-	-	-	1	1	10	8	28	40
2003.....	783	-	-	-	-	-	-	5	7	18	24	46
2002.....	744	-	-	-	-	1	1	2	1	10	23	43
2001.....	644	-	-	-	1	2	4	2	2	7	11	40
2000.....	616	-	-	-	-	1	1	-	7	7	16	25
1999.....	725	-	-	-	-	-	2	2	6	13	16	35
American Indian 1/, male												
2004.....	365	-	-	-	-	-	-	1	4	7	18	26
2003.....	367	-	-	-	-	-	-	3	5	10	16	25
2002.....	336	-	-	-	-	1	-	2	1	6	15	26
2001.....	276	-	-	-	1	1	2	1	1	5	2	21
2000.....	275	-	-	-	-	1	1	-	6	6	10	18
1999.....	323	-	-	-	-	-	2	-	5	8	12	21
American Indian 1/, female												
2004.....	381	-	-	-	-	-	1	-	6	1	10	14
2003.....	416	-	-	-	-	-	-	2	2	8	8	21
2002.....	408	-	-	-	-	-	1	-	-	4	8	17
2001.....	368	-	-	-	-	1	2	1	1	2	9	19
2000.....	341	-	-	-	-	-	-	-	1	1	6	7
1999.....	402	-	-	-	-	-	-	2	1	5	4	14

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Diabetes mellitus (E10-E14)

Asian or Pacific Islander 2/, both sexes

2004.....	1,471	-	1	-	2	-	-	5	4	9	13	37
2003.....	1,445	-	-	-	2	1	2	2	4	2	17	33
2002.....	1,359	-	-	-	-	1	-	1	4	5	12	29
2001.....	1,243	-	-	-	2	1	2	3	3	14	16	32
2000.....	1,103	-	-	-	-	-	-	3	4	7	15	31
1999.....	1,148	-	-	-	-	-	2	1	2	7	15	25

Asian or Pacific Islander 2/, male

2004.....	708	-	1	-	1	-	-	3	1	4	7	24
2003.....	707	-	-	-	-	-	2	-	3	2	9	23
2002.....	648	-	-	-	-	1	-	-	3	3	9	15
2001.....	599	-	-	-	1	-	2	1	3	9	11	23
2000.....	547	-	-	-	-	-	-	2	2	6	9	18
1999.....	523	-	-	-	-	-	-	1	1	4	9	12

Asian or Pacific Islander 2/, female

2004.....	763	-	-	-	1	-	-	2	3	5	6	13
2003.....	738	-	-	-	2	1	-	2	1	-	8	10
2002.....	711	-	-	-	-	-	-	1	1	2	3	14
2001.....	644	-	-	-	1	1	-	2	-	5	5	9
2000.....	556	-	-	-	-	-	-	1	2	1	6	13
1999.....	625	-	-	-	-	-	2	-	1	3	6	13

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Disorders of pituitary gland (E22-E23)

All races, both sexes

2004.....	232	5	4	6	2	5	6	6	1	4	5	8
2003.....	205	2	6	-	1	4	5	6	4	4	6	4
2002.....	209	3	4	-	-	3	5	5	2	5	10	1
2001.....	220	6	7	-	4	-	2	1	2	4	12	11
2000.....	224	5	8	4	1	2	1	2	4	4	5	6
1999.....	223	4	8	2	2	2	2	5	2	6	5	7

All races, male

2004.....	106	4	2	5	1	2	4	3	-	2	2	3
2003.....	94	1	4	-	-	2	2	4	2	-	3	3
2002.....	90	1	2	-	-	2	3	3	1	2	7	-
2001.....	99	5	3	-	2	-	2	1	1	3	5	5
2000.....	91	3	4	1	-	1	-	1	2	-	4	3
1999.....	88	1	4	-	-	-	-	3	1	2	1	3

All races, female

2004.....	126	1	2	1	1	3	2	3	1	2	3	5
2003.....	111	1	2	-	1	2	3	2	2	4	3	1
2002.....	119	2	2	-	-	1	2	2	1	3	3	1
2001.....	121	1	4	-	2	-	-	-	1	1	7	6
2000.....	133	2	4	3	1	1	1	1	2	4	1	3
1999.....	135	3	4	2	2	2	2	2	1	4	4	4

White, both sexes

2004.....	193	5	3	5	1	4	3	4	1	1	5	7
2003.....	168	2	4	-	1	4	4	4	2	3	4	1
2002.....	177	2	4	-	-	1	3	4	2	5	4	1
2001.....	193	4	4	-	4	-	2	-	2	4	9	9
2000.....	191	2	6	2	-	2	1	1	3	4	4	5
1999.....	167	4	6	-	1	1	1	3	1	2	2	3

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of pituitary gland (E22-E23)												
White, male												
2004.....	86	4	1	4	1	2	2	2	-	-	2	2
2003.....	76	1	2	-	-	2	1	2	1	-	3	1
2002.....	74	1	2	-	-	1	1	2	1	2	3	-
2001.....	83	3	2	-	2	-	2	-	1	3	3	4
2000.....	73	2	3	1	-	1	-	1	1	-	3	2
1999.....	64	1	4	-	-	-	-	2	1	-	-	1
White, female												
2004.....	107	1	2	1	-	2	1	2	1	1	3	5
2003.....	92	1	2	-	1	2	3	2	1	3	1	-
2002.....	103	1	2	-	-	-	2	2	1	3	1	1
2001.....	110	1	2	-	2	-	-	-	1	1	6	5
2000.....	118	-	3	1	-	1	1	-	2	4	1	3
1999.....	103	3	2	-	1	1	1	1	-	2	2	2
Black, both sexes												
2004.....	31	-	-	1	-	1	1	1	-	2	-	1
2003.....	28	-	2	-	-	-	1	1	-	1	2	3
2002.....	24	1	-	-	-	1	1	1	-	-	4	-
2001.....	22	1	2	-	-	-	-	1	-	-	3	1
2000.....	31	3	1	2	1	-	-	1	1	-	1	1
1999.....	46	-	2	1	-	1	-	2	-	4	3	4
Black, male												
2004.....	16	-	-	1	-	-	1	-	-	1	-	1
2003.....	13	-	2	-	-	-	1	1	-	-	-	2
2002.....	10	-	-	-	-	-	1	1	-	-	2	-
2001.....	14	1	1	-	-	-	-	1	-	-	2	1
2000.....	17	1	1	-	-	-	-	-	1	-	1	1
1999.....	21	-	-	-	-	-	-	1	-	2	1	2

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of pituitary gland (E22-E23)												
Black, female												
2004.....	15	-	-	-	-	1	-	1	-	1	-	-
2003.....	15	-	-	-	-	-	-	-	-	1	2	1
2002.....	14	1	-	-	-	1	-	-	-	-	2	-
2001.....	8	-	1	-	-	-	-	-	-	-	1	-
2000.....	14	2	-	2	1	-	-	1	-	-	-	-
1999.....	25	-	2	1	-	1	-	1	-	2	2	2
American Indian 1/, both sexes												
2004.....	2	-	1	-	-	-	-	-	-	-	-	-
2003.....	2	-	-	-	-	-	-	1	-	-	-	-
2002.....	1	-	-	-	-	-	1	-	-	-	-	-
2001.....	2	-	1	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	2	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, male												
2004.....	1	-	1	-	-	-	-	-	-	-	-	-
2003.....	2	-	-	-	-	-	-	1	-	-	-	-
2002.....	1	-	-	-	-	-	1	-	-	-	-	-
2001.....	1	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, female												
2004.....	1	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	1	-	1	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	2	-	-	-	-	-	-	-	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Disorders of pituitary gland (E22-E23)

Asian or Pacific Islander 2/, both sexes

2004.....	6	-	-	-	1	-	2	1	-	1	-	-
2003.....	7	-	-	-	-	-	-	-	2	-	-	-
2002.....	7	-	-	-	-	1	-	-	-	-	2	-
2001.....	3	1	-	-	-	-	-	-	-	-	-	1
2000.....	2	-	1	-	-	-	-	-	-	-	-	-
1999.....	8	-	-	1	1	-	1	-	-	-	-	-

Asian or Pacific Islander 2/, male

2004.....	3	-	-	-	-	-	1	1	-	1	-	-
2003.....	3	-	-	-	-	-	-	-	1	-	-	-
2002.....	5	-	-	-	-	1	-	-	-	-	2	-
2001.....	1	1	-	-	-	-	-	-	-	-	-	-
2000.....	1	-	-	-	-	-	-	-	-	-	-	-
1999.....	3	-	-	-	-	-	-	-	-	-	-	-

Asian or Pacific Islander 2/, female

2004.....	3	-	-	-	1	-	1	-	-	-	-	-
2003.....	4	-	-	-	-	-	-	-	1	-	-	-
2002.....	2	-	-	-	-	-	-	-	-	-	-	-
2001.....	2	-	-	-	-	-	-	-	-	-	-	1
2000.....	1	-	1	-	-	-	-	-	-	-	-	-
1999.....	5	-	-	1	1	-	1	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of adrenal glands (E24-E27)												
All races, both sexes												
2004.....	273	7	2	1	2	-	5	6	4	4	16	10
2003.....	308	4	1	3	2	2	3	4	5	8	11	19
2002.....	284	5	2	1	1	3	7	2	11	6	9	12
2001.....	338	6	5	2	2	2	3	5	2	7	14	9
2000.....	281	6	4	1	2	3	4	2	6	9	10	11
1999.....	253	4	4	2	1	2	9	7	4	8	9	9
All races, male												
2004.....	107	5	1	1	2	-	3	6	1	2	7	3
2003.....	116	3	-	2	-	1	2	2	3	5	5	7
2002.....	108	3	-	1	2	2	3	1	9	2	5	5
2001.....	156	4	3	2	2	2	2	2	1	5	7	4
2000.....	102	3	3	-	1	3	4	-	2	3	2	3
1999.....	101	2	3	1	-	2	4	4	2	4	7	-
All races, female												
2004.....	166	2	1	-	-	-	2	-	3	2	9	7
2003.....	192	1	1	1	2	1	1	2	2	3	6	12
2002.....	176	2	2	-	-	1	4	1	2	4	4	7
2001.....	182	2	2	-	-	-	1	3	1	2	7	5
2000.....	179	3	1	1	1	-	-	2	4	6	8	8
1999.....	152	2	1	1	1	-	5	3	2	4	2	9
White, both sexes												
2004.....	227	5	1	-	2	-	5	4	3	3	10	6
2003.....	266	1	1	2	2	2	2	2	5	5	11	16
2002.....	242	4	2	1	1	3	6	2	8	5	6	12
2001.....	299	5	3	2	2	2	3	4	2	4	11	7
2000.....	241	4	2	1	1	3	4	1	4	5	8	10
1999.....	222	2	3	1	1	2	7	6	4	7	8	7

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Disorders of adrenal glands (E24-E27)

White, male

2004.....	84	3	-	-	2	-	3	4	1	2	4	2
2003.....	96	-	-	2	-	1	2	2	3	2	5	5
2002.....	87	2	-	1	1	2	3	1	6	2	2	5
2001.....	137	3	2	2	2	2	2	2	1	2	6	4
2000.....	91	2	2	-	-	3	4	-	2	3	2	3
1999.....	87	-	3	-	-	2	3	4	2	3	6	-

White, female

2004.....	143	2	1	-	-	-	2	-	2	1	6	4
2003.....	170	1	1	-	2	1	-	-	2	3	6	11
2002.....	155	2	2	-	-	1	3	1	2	3	4	7
2001.....	162	2	1	-	-	-	1	2	1	2	5	3
2000.....	150	2	-	1	1	-	-	1	2	2	6	7
1999.....	135	2	-	1	1	-	4	2	2	4	2	7

Black, both sexes

2004.....	38	1	1	1	-	-	-	2	-	1	4	3
2003.....	36	3	-	1	-	-	1	1	-	3	-	3
2002.....	33	-	-	-	-	-	1	-	2	1	3	-
2001.....	35	1	1	-	-	-	-	1	-	3	3	2
2000.....	38	1	2	-	1	-	-	1	2	4	2	1
1999.....	29	1	1	1	-	-	1	1	-	1	1	2

Black, male

2004.....	18	1	1	1	-	-	-	2	-	-	2	1
2003.....	19	3	-	-	-	-	-	-	-	3	-	2
2002.....	16	-	-	-	-	-	-	-	2	-	3	-
2001.....	17	1	1	-	-	-	-	-	-	3	1	-
2000.....	9	-	1	-	1	-	-	-	-	-	-	-
1999.....	13	1	-	1	-	-	1	-	-	1	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of adrenal glands (E24-E27)												
Asian or Pacific Islander 2/, both sexes												
2004.....	6	1	-	-	-	-	-	-	-	-	1	1
2003.....	2	-	-	-	-	-	-	-	-	-	-	-
2002.....	6	1	-	-	-	-	-	-	1	-	-	-
2001.....	2	-	1	-	-	-	-	-	-	-	-	-
2000.....	1	1	-	-	-	-	-	-	-	-	-	-
1999.....	2	1	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	5	1	-	-	-	-	-	-	-	-	1	-
2003.....	1	-	-	-	-	-	-	-	-	-	-	-
2002.....	3	1	-	-	-	-	-	-	1	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	1	1	-	-	-	-	-	-	-	-	-	-
1999.....	1	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	1	-	-	-	-	-	-	-	-	-	-	1
2003.....	1	-	-	-	-	-	-	-	-	-	-	-
2002.....	3	-	-	-	-	-	-	-	-	-	-	-
2001.....	2	-	1	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	1	-	-	-	-	-	1	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other endocrine diseases (E15-E21,E28-E31,E34)

All races, both sexes

2004.....	463	8	4	1	1	4	5	7	4	9	13	20
2003.....	443	15	4	2	1	3	2	4	8	7	18	19
2002.....	472	28	3	1	-	2	3	3	4	4	12	23
2001.....	429	28	-	-	-	1	6	4	6	10	15	11
2000.....	397	26	3	1	-	3	-	5	4	8	17	20
1999.....	422	27	-	-	-	4	3	3	7	12	13	19

All races, male

2004.....	214	8	3	1	1	2	3	3	1	6	6	11
2003.....	195	7	4	2	1	2	2	1	4	2	13	13
2002.....	187	17	-	-	-	1	-	2	3	3	2	10
2001.....	174	16	-	-	-	-	3	1	3	5	9	7
2000.....	152	14	2	-	-	1	-	1	3	3	7	13
1999.....	186	11	-	-	-	1	3	1	3	6	8	7

All races, female

2004.....	249	-	1	-	-	2	2	4	3	3	7	9
2003.....	248	8	-	-	-	1	-	3	4	5	5	6
2002.....	285	11	3	1	-	1	3	1	1	1	10	13
2001.....	255	12	-	-	-	1	3	3	3	5	6	4
2000.....	245	12	1	1	-	2	-	4	1	5	10	7
1999.....	236	16	-	-	-	3	-	2	4	6	5	12

White, both sexes

2004.....	349	7	4	-	-	3	4	7	2	3	12	13
2003.....	334	9	4	1	-	2	2	2	6	7	13	13
2002.....	374	25	2	1	-	2	3	2	3	3	9	15
2001.....	334	23	-	-	-	1	6	3	5	7	12	5
2000.....	300	17	3	1	-	2	-	5	3	6	13	17
1999.....	324	19	-	-	-	3	3	2	3	8	10	15

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other endocrine diseases (E15-E21,E28-E31,E34)												
White, male												
2004.....	154	7	3	-	-	2	2	3	-	2	6	6
2003.....	143	4	4	1	-	2	2	-	4	2	9	9
2002.....	149	17	-	-	-	1	-	1	2	2	2	5
2001.....	124	12	-	-	-	-	3	-	2	3	7	3
2000.....	115	9	2	-	-	1	-	1	2	2	6	11
1999.....	133	7	-	-	-	1	3	1	1	4	7	5
White, female												
2004.....	195	-	1	-	-	1	2	4	2	1	6	7
2003.....	191	5	-	-	-	-	-	2	2	5	4	4
2002.....	225	8	2	1	-	1	3	1	1	1	7	10
2001.....	210	11	-	-	-	1	3	3	3	4	5	2
2000.....	185	8	1	1	-	1	-	4	1	4	7	6
1999.....	191	12	-	-	-	2	-	1	2	4	3	10
Black, both sexes												
2004.....	104	1	-	1	-	-	1	-	2	5	1	6
2003.....	99	5	-	1	1	1	-	1	2	-	5	6
2002.....	92	2	1	-	-	-	-	1	1	1	3	7
2001.....	86	5	-	-	-	-	-	1	-	2	3	6
2000.....	89	6	-	-	-	1	-	-	1	2	4	3
1999.....	95	8	-	-	-	1	-	1	4	4	3	4
Black, male												
2004.....	54	1	-	1	-	-	1	-	1	3	-	4
2003.....	48	3	-	1	1	-	-	1	-	-	4	4
2002.....	36	-	-	-	-	-	-	1	1	1	-	4
2001.....	44	4	-	-	-	-	-	1	-	1	2	4
2000.....	30	2	-	-	-	-	-	-	1	1	1	2
1999.....	51	4	-	-	-	-	-	-	2	2	1	2

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Nutritional diseases (E40-E68)

All races, both sexes

2004.....	7,223	7	4	3	5	16	42	120	183	261	391	555
2003.....	7,359	10	3	2	3	25	53	121	174	269	399	551
2002.....	7,470	4	1	2	5	14	64	106	194	270	442	475
2001.....	6,843	4	5	1	2	12	59	84	152	263	314	437
2000.....	7,233	15	4	1	9	19	36	89	137	240	365	397
1999.....	6,889	9	3	2	-	21	40	78	123	210	337	330

All races, male

2004.....	3,043	5	2	1	3	11	28	64	92	139	209	269
2003.....	3,048	7	3	2	3	11	29	77	112	131	204	277
2002.....	3,047	3	1	2	2	9	38	56	116	152	254	232
2001.....	2,622	4	1	1	2	6	27	53	80	134	170	223
2000.....	2,829	11	3	-	6	11	22	54	89	139	188	213
1999.....	2,704	4	2	1	-	13	17	40	82	121	195	170

All races, female

2004.....	4,180	2	2	2	2	5	14	56	91	122	182	286
2003.....	4,311	3	-	-	-	14	24	44	62	138	195	274
2002.....	4,423	1	-	-	3	5	26	50	78	118	188	243
2001.....	4,221	-	4	-	-	6	32	31	72	129	144	214
2000.....	4,404	4	1	1	3	8	14	35	48	101	177	184
1999.....	4,185	5	1	1	-	8	23	38	41	89	142	160

White, both sexes

2004.....	5,826	5	2	2	4	9	25	75	117	165	289	418
2003.....	6,080	6	2	2	2	15	35	76	116	191	303	432
2002.....	6,108	3	1	1	4	7	37	68	120	193	336	377
2001.....	5,590	3	4	1	1	5	34	60	99	177	227	322
2000.....	5,988	8	3	-	6	12	22	51	87	174	277	301
1999.....	5,684	5	2	1	-	12	23	44	82	144	262	252

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Nutritional diseases (E40-E68)

White, male

2004.....	2,506	3	-	-	2	7	17	41	58	95	166	219
2003.....	2,551	5	2	2	2	8	22	50	80	104	160	228
2002.....	2,525	2	1	1	2	5	22	36	72	112	200	199
2001.....	2,130	3	1	1	1	2	18	38	56	93	128	171
2000.....	2,349	6	2	-	6	8	15	35	59	108	149	180
1999.....	2,207	2	1	-	-	9	8	24	55	85	163	139

White, female

2004.....	3,320	2	2	2	2	2	8	34	59	70	123	199
2003.....	3,529	1	-	-	-	7	13	26	36	87	143	204
2002.....	3,583	1	-	-	2	2	15	32	48	81	136	178
2001.....	3,460	-	3	-	-	3	16	22	43	84	99	151
2000.....	3,639	2	1	-	-	4	7	16	28	66	128	121
1999.....	3,477	3	1	1	-	3	15	20	27	59	99	113

Black, both sexes

2004.....	1,269	1	2	1	-	5	17	43	62	92	93	127
2003.....	1,161	3	1	-	1	8	15	41	54	72	87	109
2002.....	1,247	-	-	1	1	7	22	38	68	67	99	94
2001.....	1,132	1	1	-	1	7	24	20	47	82	82	105
2000.....	1,135	6	1	1	3	7	13	36	45	63	83	90
1999.....	1,102	4	1	1	-	6	17	33	37	62	73	73

Black, male

2004.....	480	1	2	1	-	3	11	22	31	42	39	42
2003.....	444	1	1	-	1	2	5	23	30	25	38	45
2002.....	465	-	-	1	-	4	12	20	39	32	50	31
2001.....	437	1	-	-	1	4	8	11	22	39	39	45
2000.....	443	4	1	-	-	3	6	17	27	31	37	31
1999.....	444	2	1	1	-	3	9	15	24	33	31	28

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Nutritional diseases (E40-E68)

Black, female

2004.....	789	-	-	-	-	2	6	21	31	50	54	85
2003.....	717	2	-	-	-	6	10	18	24	47	49	64
2002.....	782	-	-	-	1	3	10	18	29	35	49	63
2001.....	695	-	1	-	-	3	16	9	25	43	43	60
2000.....	692	2	-	1	3	4	7	19	18	32	46	59
1999.....	658	2	-	-	-	3	8	18	13	29	42	45

American Indian 1/, both sexes

2004.....	54	-	-	-	1	-	-	2	2	4	4	6
2003.....	52	-	-	-	-	-	1	3	2	3	4	6
2002.....	54	-	-	-	-	-	2	-	4	7	2	3
2001.....	55	-	-	-	-	-	-	1	1	2	5	4
2000.....	50	1	-	-	-	-	-	1	2	2	3	3
1999.....	47	-	-	-	-	2	-	1	2	1	2	2

American Indian 1/, male

2004.....	27	-	-	-	1	-	-	1	2	2	2	5
2003.....	21	-	-	-	-	-	1	3	1	1	2	2
2002.....	23	-	-	-	-	-	2	-	3	5	1	1
2001.....	23	-	-	-	-	-	-	1	-	1	3	2
2000.....	15	1	-	-	-	-	-	1	2	-	1	1
1999.....	21	-	-	-	-	1	-	1	1	1	1	1

American Indian 1/, female

2004.....	27	-	-	-	-	-	-	1	-	2	2	1
2003.....	31	-	-	-	-	-	-	-	1	2	2	4
2002.....	31	-	-	-	-	-	-	-	1	2	1	2
2001.....	32	-	-	-	-	-	-	-	1	1	2	2
2000.....	35	-	-	-	-	-	-	-	-	2	2	2
1999.....	26	-	-	-	-	1	-	-	1	-	1	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nutritional diseases (E40-E68)												
Asian or Pacific Islander 2/, both sexes												
2004.....	74	1	-	-	-	2	-	-	2	-	5	4
2003.....	66	1	-	-	-	2	2	1	2	3	5	4
2002.....	61	1	-	-	-	-	3	-	2	3	5	1
2001.....	66	-	-	-	-	-	1	3	5	2	-	6
2000.....	60	-	-	-	-	-	1	1	3	1	2	3
1999.....	56	-	-	-	-	1	-	-	2	3	-	3
Asian or Pacific Islander 2/, male												
2004.....	30	1	-	-	-	1	-	-	1	-	2	3
2003.....	32	1	-	-	-	1	1	1	1	1	4	2
2002.....	34	1	-	-	-	-	2	-	2	3	3	1
2001.....	32	-	-	-	-	-	1	3	2	1	-	5
2000.....	22	-	-	-	-	-	1	1	1	-	1	1
1999.....	32	-	-	-	-	-	-	-	2	2	-	2
Asian or Pacific Islander 2/, female												
2004.....	44	-	-	-	-	1	-	-	1	-	3	1
2003.....	34	-	-	-	-	1	1	-	1	2	1	2
2002.....	27	-	-	-	-	-	1	-	-	-	2	-
2001.....	34	-	-	-	-	-	-	-	3	1	-	1
2000.....	38	-	-	-	-	-	-	-	2	1	1	2
1999.....	24	-	-	-	-	1	-	-	-	1	-	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Malnutrition (E40-E46)

All races, both sexes

2004.....	2,847	6	1	1	1	2	3	6	5	14	21	36
2003.....	3,153	5	1	1	-	2	4	4	4	12	21	36
2002.....	3,510	2	1	-	1	1	5	4	9	25	30	36
2001.....	3,454	2	-	-	2	1	8	4	9	16	23	42
2000.....	3,926	11	1	1	2	3	5	9	7	29	24	44
1999.....	3,985	7	2	1	-	2	5	4	7	20	41	42

All races, male

2004.....	1,026	5	1	1	1	2	3	3	3	7	6	16
2003.....	1,098	4	1	1	-	-	3	3	3	7	12	19
2002.....	1,233	1	1	-	-	-	4	2	3	11	17	18
2001.....	1,148	2	-	-	2	-	6	2	2	8	11	22
2000.....	1,302	8	1	-	1	1	1	5	3	9	6	23
1999.....	1,371	4	2	1	-	2	2	1	3	8	19	21

All races, female

2004.....	1,821	1	-	-	-	-	-	3	2	7	15	20
2003.....	2,055	1	-	-	-	2	1	1	1	5	9	17
2002.....	2,277	1	-	-	1	1	1	2	6	14	13	18
2001.....	2,306	-	-	-	-	1	2	2	7	8	12	20
2000.....	2,624	3	-	1	1	2	4	4	4	20	18	21
1999.....	2,614	3	-	-	-	-	3	3	4	12	22	21

White, both sexes

2004.....	2,373	4	-	-	1	1	2	4	4	7	16	25
2003.....	2,657	2	1	1	-	1	4	3	3	10	18	25
2002.....	2,979	2	1	-	-	1	4	4	6	15	23	29
2001.....	2,939	1	-	-	1	1	6	4	5	11	14	30
2000.....	3,352	7	-	-	1	3	4	5	3	20	17	35
1999.....	3,368	4	1	-	-	2	3	3	5	15	32	27

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Malnutrition (E40-E46)

White, male

2004.....	829	3	-	-	1	1	2	2	3	3	3	8
2003.....	899	2	1	1	-	-	3	3	2	6	10	12
2002.....	1,029	1	1	-	-	-	3	2	2	7	13	12
2001.....	932	1	-	-	1	-	6	2	1	5	5	11
2000.....	1,079	5	-	-	1	1	1	3	1	7	3	18
1999.....	1,100	2	1	-	-	2	1	-	3	5	14	15

White, female

2004.....	1,544	1	-	-	-	-	-	2	1	4	13	17
2003.....	1,758	-	-	-	-	1	1	-	1	4	8	13
2002.....	1,950	1	-	-	-	1	1	2	4	8	10	17
2001.....	2,007	-	-	-	-	1	-	2	4	6	9	19
2000.....	2,273	2	-	-	-	2	3	2	2	13	14	17
1999.....	2,268	2	-	-	-	-	2	3	2	10	18	12

Black, both sexes

2004.....	402	1	1	1	-	1	1	2	1	6	4	10
2003.....	442	2	-	-	-	1	-	1	1	1	3	11
2002.....	472	-	-	-	1	-	1	-	3	7	7	7
2001.....	441	1	-	-	1	-	2	-	4	5	9	10
2000.....	508	3	1	1	1	-	1	4	3	9	7	8
1999.....	547	3	1	1	-	-	2	1	2	5	9	13

Black, male

2004.....	167	1	1	1	-	1	1	1	-	4	2	7
2003.....	179	1	-	-	-	-	-	-	1	-	2	7
2002.....	175	-	-	-	-	-	1	-	1	2	4	6
2001.....	188	1	-	-	1	-	-	-	1	3	6	9
2000.....	203	2	1	-	-	-	-	2	2	2	3	5
1999.....	234	2	1	1	-	-	1	1	-	3	5	5

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Nutritional marasmus (E41)

All races, both sexes

2004.....	120	1	1	-	-	-	-	-	-	1	-	1
2003.....	150	-	1	-	-	-	-	-	-	-	1	2
2002.....	189	-	-	-	-	-	-	-	2	-	2	-
2001.....	195	-	-	-	-	-	1	1	-	2	-	1
2000.....	206	1	1	-	-	-	-	1	-	1	1	1
1999.....	200	-	-	-	-	-	-	-	2	-	2	3

All races, male

2004.....	40	1	1	-	-	-	-	-	-	-	-	-
2003.....	51	-	1	-	-	-	-	-	-	-	-	2
2002.....	52	-	-	-	-	-	-	-	1	-	2	-
2001.....	68	-	-	-	-	-	1	-	-	1	-	1
2000.....	74	1	1	-	-	-	-	1	-	-	-	-
1999.....	82	-	-	-	-	-	-	-	1	-	1	1

All races, female

2004.....	80	-	-	-	-	-	-	-	-	1	-	1
2003.....	99	-	-	-	-	-	-	-	-	-	1	-
2002.....	137	-	-	-	-	-	-	-	1	-	-	-
2001.....	127	-	-	-	-	-	-	1	-	1	-	-
2000.....	132	-	-	-	-	-	-	-	-	1	1	1
1999.....	118	-	-	-	-	-	-	-	1	-	1	2

White, both sexes

2004.....	106	1	-	-	-	-	-	-	-	1	-	1
2003.....	126	-	1	-	-	-	-	-	-	-	-	1
2002.....	161	-	-	-	-	-	-	-	1	-	1	-
2001.....	167	-	-	-	-	-	1	1	-	-	-	1
2000.....	182	1	-	-	-	-	-	-	-	1	1	1
1999.....	167	-	-	-	-	-	-	-	1	-	1	2

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified malnutrition (E40,E42-E46)												
All races, both sexes												
2004.....	2,727	5	-	1	1	2	3	6	5	13	21	35
2003.....	3,003	5	-	1	-	2	4	4	4	12	20	34
2002.....	3,321	2	1	-	1	1	5	4	7	25	28	36
2001.....	3,259	2	-	-	2	1	7	3	9	14	23	41
2000.....	3,720	10	-	1	2	3	5	8	7	28	23	43
1999.....	3,785	7	2	1	-	2	5	4	5	20	39	39
All races, male												
2004.....	986	4	-	1	1	2	3	3	3	7	6	16
2003.....	1,047	4	-	1	-	-	3	3	3	7	12	17
2002.....	1,181	1	1	-	-	-	4	2	2	11	15	18
2001.....	1,080	2	-	-	2	-	5	2	2	7	11	21
2000.....	1,228	7	-	-	1	1	1	4	3	9	6	23
1999.....	1,289	4	2	1	-	2	2	1	2	8	18	20
All races, female												
2004.....	1,741	1	-	-	-	-	-	3	2	6	15	19
2003.....	1,956	1	-	-	-	2	1	1	1	5	8	17
2002.....	2,140	1	-	-	1	1	1	2	5	14	13	18
2001.....	2,179	-	-	-	-	1	2	1	7	7	12	20
2000.....	2,492	3	-	1	1	2	4	4	4	19	17	20
1999.....	2,496	3	-	-	-	-	3	3	3	12	21	19
White, both sexes												
2004.....	2,267	3	-	-	1	1	2	4	4	6	16	24
2003.....	2,531	2	-	1	-	1	4	3	3	10	18	24
2002.....	2,818	2	1	-	-	1	4	4	5	15	22	29
2001.....	2,772	1	-	-	1	1	5	3	5	11	14	29
2000.....	3,170	6	-	-	1	3	4	5	3	19	16	34
1999.....	3,201	4	1	-	-	2	3	3	4	15	31	25

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	-------------	-----------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Other and unspecified malnutrition (E40,E42-E46)

White, male

2004.....	796	2	-	-	1	1	2	2	3	3	3	8
2003.....	857	2	-	1	-	-	3	3	2	6	10	11
2002.....	987	1	1	-	-	-	3	2	1	7	12	12
2001.....	876	1	-	-	1	-	5	2	1	5	5	10
2000.....	1,019	4	-	-	1	1	1	3	1	7	3	18
1999.....	1,032	2	1	-	-	2	1	-	2	5	14	14

White, female

2004.....	1,471	1	-	-	-	-	-	2	1	3	13	16
2003.....	1,674	-	-	-	-	1	1	-	1	4	8	13
2002.....	1,831	1	-	-	-	1	1	2	4	8	10	17
2001.....	1,896	-	-	-	-	1	-	1	4	6	9	19
2000.....	2,151	2	-	-	-	2	3	2	2	12	13	16
1999.....	2,169	2	-	-	-	-	2	3	2	10	17	11

Black, both sexes

2004.....	393	1	-	1	-	1	1	2	1	6	4	10
2003.....	423	2	-	-	-	1	-	1	1	1	2	10
2002.....	450	-	-	-	1	-	1	-	2	7	6	7
2001.....	420	1	-	-	1	-	2	-	4	3	9	10
2000.....	485	3	-	1	1	-	1	3	3	9	7	8
1999.....	517	3	1	1	-	-	2	1	1	5	8	12

Black, male

2004.....	163	1	-	1	-	1	1	1	-	4	2	7
2003.....	170	1	-	-	-	-	-	-	1	-	2	6
2002.....	166	-	-	-	-	-	1	-	1	2	3	6
2001.....	180	1	-	-	1	-	-	-	1	2	6	9
2000.....	189	2	-	-	-	-	-	1	2	2	3	5
1999.....	223	2	1	1	-	-	1	1	-	3	4	5

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other nutritional deficiencies (E50-E64)												
All races, both sexes												
2004.....	172	1	1	1	-	1	-	-	2	2	2	4
2003.....	185	5	1	1	-	1	-	1	1	2	-	3
2002.....	269	2	-	-	-	-	1	1	2	2	3	3
2001.....	250	2	4	-	-	-	1	1	-	-	5	6
2000.....	316	3	1	-	2	1	1	-	2	1	2	2
1999.....	304	2	-	-	-	-	1	-	1	-	3	3
All races, male												
2004.....	61	-	-	-	-	-	-	-	-	1	1	-
2003.....	78	3	1	1	-	1	-	1	-	-	-	-
2002.....	88	2	-	-	-	-	-	1	1	1	2	2
2001.....	85	2	1	-	-	-	1	-	-	-	2	5
2000.....	110	2	1	-	1	-	-	-	1	-	1	2
1999.....	115	-	-	-	-	-	1	-	-	-	1	1
All races, female												
2004.....	111	1	1	1	-	1	-	-	2	1	1	4
2003.....	107	2	-	-	-	-	-	-	1	2	-	3
2002.....	181	-	-	-	-	-	1	-	1	1	1	1
2001.....	165	-	3	-	-	-	-	1	-	-	3	1
2000.....	206	1	-	-	1	1	1	-	1	1	1	-
1999.....	189	2	-	-	-	-	-	-	1	-	2	2
White, both sexes												
2004.....	156	1	1	1	-	-	-	-	2	1	1	2
2003.....	169	4	1	1	-	-	-	1	1	1	-	2
2002.....	234	1	-	-	-	-	1	1	2	-	3	3
2001.....	213	2	4	-	-	-	1	1	-	-	4	4
2000.....	275	1	1	-	1	1	-	-	1	1	2	2
1999.....	260	1	-	-	-	-	-	-	1	-	2	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other nutritional deficiencies (E50-E64)

White, male

2004.....	55	-	-	-	-	-	-	-	-	-	1	-
2003.....	70	3	1	1	-	-	-	1	-	-	-	-
2002.....	73	1	-	-	-	-	-	1	1	-	2	2
2001.....	68	2	1	-	-	-	1	-	-	-	2	3
2000.....	96	1	1	-	1	-	-	-	-	-	1	2
1999.....	99	-	-	-	-	-	-	-	-	-	1	1

White, female

2004.....	101	1	1	1	-	-	-	-	2	1	-	2
2003.....	99	1	-	-	-	-	-	-	1	1	-	2
2002.....	161	-	-	-	-	-	1	-	1	-	1	1
2001.....	145	-	3	-	-	-	-	1	-	-	2	1
2000.....	179	-	-	-	-	1	-	-	1	1	1	-
1999.....	161	1	-	-	-	-	-	-	1	-	1	-

Black, both sexes

2004.....	14	-	-	-	-	-	-	-	-	1	1	2
2003.....	13	1	-	-	-	-	-	-	-	-	-	1
2002.....	30	-	-	-	-	-	-	-	-	2	-	-
2001.....	34	-	-	-	-	-	-	-	-	-	1	2
2000.....	32	2	-	-	1	-	1	-	1	-	-	-
1999.....	42	1	-	-	-	-	1	-	-	-	1	2

Black, male

2004.....	6	-	-	-	-	-	-	-	-	1	-	-
2003.....	6	-	-	-	-	-	-	-	-	-	-	-
2002.....	13	-	-	-	-	-	-	-	-	1	-	-
2001.....	16	-	-	-	-	-	-	-	-	-	-	2
2000.....	12	1	-	-	-	-	-	-	1	-	-	-
1999.....	16	-	-	-	-	-	1	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	-------------	-----------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Obesity and other hyperalimentation (E65-E68)

All races, both sexes

2004.....	4,204	-	2	1	4	13	39	114	176	245	368	515
2003.....	4,021	-	1	-	3	22	49	116	169	255	378	512
2002.....	3,691	-	-	2	4	13	58	101	183	243	409	436
2001.....	3,139	-	1	1	-	11	50	79	143	247	286	389
2000.....	2,991	1	2	-	5	15	30	80	128	210	339	351
1999.....	2,600	-	1	1	-	19	34	74	115	190	293	285

All races, male

2004.....	1,956	-	1	-	2	9	25	61	89	131	202	253
2003.....	1,872	-	1	-	3	10	26	73	109	124	192	258
2002.....	1,726	-	-	2	2	9	34	53	112	140	235	212
2001.....	1,389	-	-	1	-	6	20	51	78	126	157	196
2000.....	1,417	1	1	-	4	10	21	49	85	130	181	188
1999.....	1,218	-	-	-	-	11	14	39	79	113	175	148

All races, female

2004.....	2,248	-	1	1	2	4	14	53	87	114	166	262
2003.....	2,149	-	-	-	-	12	23	43	60	131	186	254
2002.....	1,965	-	-	-	2	4	24	48	71	103	174	224
2001.....	1,750	-	1	-	-	5	30	28	65	121	129	193
2000.....	1,574	-	1	-	1	5	9	31	43	80	158	163
1999.....	1,382	-	1	1	-	8	20	35	36	77	118	137

White, both sexes

2004.....	3,297	-	1	1	3	8	23	71	111	157	272	391
2003.....	3,254	-	-	-	2	14	31	72	112	180	285	405
2002.....	2,895	-	-	1	4	6	32	63	112	178	310	345
2001.....	2,438	-	-	1	-	4	27	55	94	166	209	288
2000.....	2,361	-	2	-	4	8	18	46	83	153	258	264
1999.....	2,056	-	1	1	-	10	20	41	76	129	228	224

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Obesity and other hyperalimentation (E65-E68)

White, male

2004.....	1,622	-	-	-	1	6	15	39	55	92	162	211
2003.....	1,582	-	-	-	2	8	19	46	78	98	150	216
2002.....	1,423	-	-	1	2	5	19	33	69	105	185	185
2001.....	1,130	-	-	1	-	2	11	36	55	88	121	157
2000.....	1,174	-	1	-	4	7	14	32	58	101	145	160
1999.....	1,008	-	-	-	-	7	7	24	52	80	148	123

White, female

2004.....	1,675	-	1	1	2	2	8	32	56	65	110	180
2003.....	1,672	-	-	-	2	6	12	26	34	82	135	189
2002.....	1,472	-	-	-	2	1	13	30	43	73	125	160
2001.....	1,308	-	-	-	-	2	16	19	39	78	88	131
2000.....	1,187	-	1	-	-	1	4	14	25	52	113	104
1999.....	1,048	-	1	1	-	3	13	17	24	49	80	101

Black, both sexes

2004.....	853	-	1	-	-	4	16	41	61	85	88	115
2003.....	706	-	1	-	1	7	15	40	53	71	84	97
2002.....	745	-	-	1	-	7	21	38	65	58	92	87
2001.....	657	-	1	-	-	7	22	20	43	77	72	93
2000.....	595	1	-	-	1	7	11	32	41	54	76	82
1999.....	513	-	-	-	-	6	14	32	35	57	63	58

Black, male

2004.....	307	-	1	-	-	2	10	21	31	37	37	35
2003.....	259	-	1	-	1	2	5	23	29	25	36	38
2002.....	277	-	-	1	-	4	11	20	38	29	46	25
2001.....	233	-	-	-	-	4	8	11	21	36	33	34
2000.....	228	1	-	-	-	3	6	15	24	29	34	26
1999.....	194	-	-	-	-	3	7	14	24	30	26	23

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Obesity and other hyperalimentation (E65-E68)												
Black, female												
2004.....	546	-	-	-	-	2	6	20	30	48	51	80
2003.....	447	-	-	-	-	5	10	17	24	46	48	59
2002.....	468	-	-	-	-	3	10	18	27	29	46	62
2001.....	424	-	1	-	-	3	14	9	22	41	39	59
2000.....	367	-	-	-	1	4	5	17	17	25	42	56
1999.....	319	-	-	-	-	3	7	18	11	27	37	35
American Indian 1/, both sexes												
2004.....	29	-	-	-	1	-	-	2	2	3	3	6
2003.....	31	-	-	-	-	-	1	3	2	2	4	6
2002.....	29	-	-	-	-	-	2	-	4	6	2	3
2001.....	27	-	-	-	-	-	-	1	1	2	5	4
2000.....	19	-	-	-	-	-	-	1	2	2	3	3
1999.....	19	-	-	-	-	2	-	1	2	1	2	-
American Indian 1/, male												
2004.....	17	-	-	-	1	-	-	1	2	2	1	5
2003.....	17	-	-	-	-	-	1	3	1	1	2	2
2002.....	15	-	-	-	-	-	2	-	3	5	1	1
2001.....	15	-	-	-	-	-	-	1	-	1	3	2
2000.....	7	-	-	-	-	-	-	1	2	-	1	1
1999.....	9	-	-	-	-	1	-	1	1	1	1	-
American Indian 1/, female												
2004.....	12	-	-	-	-	-	-	1	-	1	2	1
2003.....	14	-	-	-	-	-	-	-	1	1	2	4
2002.....	14	-	-	-	-	-	-	-	1	1	1	2
2001.....	12	-	-	-	-	-	-	-	1	1	2	2
2000.....	12	-	-	-	-	-	-	-	-	2	2	2
1999.....	10	-	-	-	-	1	-	-	1	-	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Obesity and other hyperalimentation (E65-E68)												
Asian or Pacific Islander 2/, both sexes												
2004.....	25	-	-	-	-	1	-	-	2	-	5	3
2003.....	30	-	-	-	-	1	2	1	2	2	5	4
2002.....	22	-	-	-	-	-	3	-	2	1	5	1
2001.....	17	-	-	-	-	-	1	3	5	2	-	4
2000.....	16	-	-	-	-	-	1	1	2	1	2	2
1999.....	12	-	-	-	-	1	-	-	2	3	-	3
Asian or Pacific Islander 2/, male												
2004.....	10	-	-	-	-	1	-	-	1	-	2	2
2003.....	14	-	-	-	-	-	1	1	1	-	4	2
2002.....	11	-	-	-	-	-	2	-	2	1	3	1
2001.....	11	-	-	-	-	-	1	3	2	1	-	3
2000.....	8	-	-	-	-	-	1	1	1	-	1	1
1999.....	7	-	-	-	-	-	-	-	2	2	-	2
Asian or Pacific Islander 2/, female												
2004.....	15	-	-	-	-	-	-	-	1	-	3	1
2003.....	16	-	-	-	-	1	1	-	1	2	1	2
2002.....	11	-	-	-	-	-	1	-	-	-	2	-
2001.....	6	-	-	-	-	-	-	-	3	1	-	1
2000.....	8	-	-	-	-	-	-	-	1	1	1	1
1999.....	5	-	-	-	-	1	-	-	-	1	-	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Metabolic disorders (E70-E88)												
All races, both sexes												
2004.....	15,934	200	120	77	82	102	162	103	121	149	212	381
2003.....	15,949	208	167	74	90	127	142	127	109	155	237	347
2002.....	15,829	222	132	81	93	110	146	123	127	140	238	330
2001.....	15,697	244	160	83	84	128	129	116	105	145	221	347
2000.....	15,125	200	134	62	103	127	146	123	117	148	215	305
1999.....	14,874	213	129	82	94	140	139	127	105	145	198	264
All races, male												
2004.....	7,320	125	65	37	46	50	84	53	71	79	129	261
2003.....	7,238	119	91	47	54	69	70	65	64	85	156	223
2002.....	6,990	126	62	51	52	54	77	69	75	81	150	206
2001.....	6,868	135	83	40	45	65	60	57	58	88	135	233
2000.....	6,566	108	72	36	62	60	80	75	61	76	129	219
1999.....	6,157	122	73	42	52	71	63	68	53	82	120	175
All races, female												
2004.....	8,614	75	55	40	36	52	78	50	50	70	83	120
2003.....	8,711	89	76	27	36	58	72	62	45	70	81	124
2002.....	8,839	96	70	30	41	56	69	54	52	59	88	124
2001.....	8,829	109	77	43	39	63	69	59	47	57	86	114
2000.....	8,559	92	62	26	41	67	66	48	56	72	86	86
1999.....	8,717	91	56	40	42	69	76	59	52	63	78	89
White, both sexes												
2004.....	14,031	158	101	65	72	91	151	93	104	119	165	310
2003.....	13,883	145	133	66	76	107	127	108	94	123	190	280
2002.....	13,739	158	105	66	81	102	125	111	110	115	193	260
2001.....	13,728	176	132	70	72	114	122	102	93	126	189	267
2000.....	13,286	149	113	55	93	114	129	105	97	125	174	253
1999.....	13,070	145	109	62	81	124	127	117	89	113	148	212

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Metabolic disorders (E70-E88)

White, male

2004.....	6,447	100	55	31	39	43	79	49	60	62	99	218
2003.....	6,313	89	72	42	48	58	64	55	52	71	126	185
2002.....	6,042	99	48	46	45	48	62	60	66	66	126	162
2001.....	5,978	95	70	34	37	56	55	50	53	76	116	183
2000.....	5,773	80	64	31	56	51	68	65	49	64	105	186
1999.....	5,332	85	61	33	43	61	56	60	43	64	94	141

White, female

2004.....	7,584	58	46	34	33	48	72	44	44	57	66	92
2003.....	7,570	56	61	24	28	49	63	53	42	52	64	95
2002.....	7,697	59	57	20	36	54	63	51	44	49	67	98
2001.....	7,750	81	62	36	35	58	67	52	40	50	73	84
2000.....	7,513	69	49	24	37	63	61	40	48	61	69	67
1999.....	7,738	60	48	29	38	63	71	57	46	49	54	71

Black, both sexes

2004.....	1,634	34	12	3	8	9	9	9	15	25	44	66
2003.....	1,737	52	20	5	11	13	12	13	13	27	39	55
2002.....	1,807	51	20	12	11	7	19	11	12	23	39	65
2001.....	1,702	57	16	9	10	7	7	10	10	14	25	70
2000.....	1,625	38	15	5	7	12	14	17	17	23	39	49
1999.....	1,587	56	15	16	12	11	9	8	16	27	49	43

Black, male

2004.....	744	21	7	2	6	5	4	4	9	14	28	39
2003.....	759	23	11	3	4	6	5	7	10	11	24	34
2002.....	799	23	9	4	6	6	14	8	5	13	21	41
2001.....	744	30	7	4	6	5	5	4	4	9	13	42
2000.....	691	23	6	4	4	8	11	9	12	12	22	32
1999.....	706	32	8	6	8	8	5	6	10	15	26	27

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Metabolic disorders (E70-E88)

Black, female

2004.....	890	13	5	1	2	4	5	5	6	11	16	27
2003.....	978	29	9	2	7	7	7	6	3	16	15	21
2002.....	1,008	28	11	8	5	1	5	3	7	10	18	24
2001.....	958	27	9	5	4	2	2	6	6	5	12	28
2000.....	934	15	9	1	3	4	3	8	5	11	17	17
1999.....	881	24	7	10	4	3	4	2	6	12	23	16

American Indian 1/, both sexes

2004.....	58	1	2	2	1	-	1	-	-	-	2	3
2003.....	74	4	4	1	1	2	1	1	1	-	1	4
2002.....	71	5	2	1	-	-	-	-	1	-	2	2
2001.....	65	-	1	1	1	1	-	2	-	2	3	5
2000.....	60	5	5	-	1	1	-	1	1	-	1	2
1999.....	47	6	-	1	-	3	1	2	-	1	-	3

American Indian 1/, male

2004.....	23	-	-	-	-	-	-	-	-	-	1	2
2003.....	28	2	3	1	-	1	1	1	1	-	-	2
2002.....	33	1	2	-	-	-	-	-	1	-	1	1
2001.....	36	-	1	-	1	1	-	2	-	1	2	4
2000.....	30	1	2	-	1	1	-	1	-	-	1	-
1999.....	24	2	-	1	-	-	1	2	-	1	-	3

American Indian 1/, female

2004.....	35	1	2	2	1	-	1	-	-	-	1	1
2003.....	46	2	1	-	1	1	-	-	-	-	1	2
2002.....	38	4	-	1	-	-	1	-	-	-	1	1
2001.....	29	-	-	1	-	-	-	-	-	1	1	1
2000.....	30	4	3	-	-	-	-	-	1	-	-	2
1999.....	23	4	-	-	-	3	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Metabolic disorders (E70-E88)

Asian or Pacific Islander 2/, both sexes

2004.....	211	7	5	7	1	2	1	1	2	5	1	2
2003.....	255	7	10	2	2	5	2	5	1	5	7	8
2002.....	212	8	5	2	1	1	1	1	4	2	4	3
2001.....	202	11	11	3	1	6	-	2	2	3	4	5
2000.....	154	8	1	2	2	-	3	-	2	-	1	1
1999.....	170	6	5	3	1	2	2	-	-	4	1	6

Asian or Pacific Islander 2/, male

2004.....	106	4	3	4	1	2	1	-	2	3	1	2
2003.....	138	5	5	1	2	4	-	2	1	3	6	2
2002.....	116	3	3	1	1	-	1	1	3	2	2	2
2001.....	110	10	5	2	1	3	-	1	1	2	4	4
2000.....	72	4	-	1	1	-	1	-	-	-	1	1
1999.....	95	3	4	2	1	2	1	-	-	2	-	4

Asian or Pacific Islander 2/, female

2004.....	105	3	2	3	-	-	-	1	-	2	-	-
2003.....	117	2	5	1	-	1	2	3	-	2	1	6
2002.....	96	5	2	1	-	1	-	-	1	-	2	1
2001.....	92	1	6	1	-	3	-	1	1	1	-	1
2000.....	82	4	1	1	1	-	2	-	2	-	-	-
1999.....	75	3	1	1	-	-	1	-	-	2	1	2

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Cystic fibrosis (E84)

All races, both sexes

2004.....	460	7	3	8	28	60	108	61	60	48	19	18
2003.....	451	5	5	5	22	64	92	83	48	40	39	15
2002.....	484	10	8	18	29	60	93	72	66	45	35	13
2001.....	482	6	7	8	31	76	94	75	51	43	30	29
2000.....	494	4	5	9	37	73	92	71	68	61	36	18
1999.....	485	12	1	14	36	81	102	83	53	38	25	9

All races, male

2004.....	224	6	-	3	14	28	52	29	33	22	6	12
2003.....	228	3	2	3	9	30	39	41	31	22	22	11
2002.....	241	7	2	8	10	22	43	41	42	23	17	6
2001.....	237	4	2	2	15	32	40	36	30	27	21	17
2000.....	241	1	2	5	16	30	49	41	28	28	16	14
1999.....	234	6	-	5	17	34	48	43	29	18	16	3

All races, female

2004.....	236	1	3	5	14	32	56	32	27	26	13	6
2003.....	223	2	3	2	13	34	53	42	17	18	17	4
2002.....	243	3	6	10	19	38	50	31	24	22	18	7
2001.....	245	2	5	6	16	44	54	39	21	16	9	12
2000.....	253	3	3	4	21	43	43	30	40	33	20	4
1999.....	251	6	1	9	19	47	54	40	24	20	9	6

White, both sexes

2004.....	442	7	3	8	26	55	107	59	56	47	18	17
2003.....	420	5	4	5	17	59	87	76	45	39	38	15
2002.....	458	9	6	16	26	59	86	70	65	43	35	13
2001.....	464	6	7	7	28	73	93	70	50	42	29	29
2000.....	473	4	4	9	36	70	90	68	65	58	36	17
1999.....	470	12	1	13	34	77	98	83	51	38	25	9

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cystic fibrosis (E84)												
White, male												
2004.....	217	6	-	3	14	26	52	29	31	21	6	11
2003.....	215	3	2	3	8	28	37	37	29	21	22	11
2002.....	232	7	1	8	10	21	39	40	42	23	17	6
2001.....	226	4	2	1	14	29	39	33	29	27	20	17
2000.....	231	1	1	5	16	28	48	40	27	27	16	13
1999.....	226	6	-	4	17	32	45	43	28	18	16	3
White, female												
2004.....	225	1	3	5	12	29	55	30	25	26	12	6
2003.....	205	2	2	2	9	31	50	39	16	18	16	4
2002.....	226	2	5	8	16	38	47	30	23	20	18	7
2001.....	238	2	5	6	14	44	54	37	21	15	9	12
2000.....	242	3	3	4	20	42	42	28	38	31	20	4
1999.....	244	6	1	9	17	45	53	40	23	20	9	6
Black, both sexes												
2004.....	16	-	-	-	1	5	1	2	4	1	1	1
2003.....	25	-	1	-	3	4	4	5	3	1	1	-
2002.....	25	1	2	2	3	1	7	2	1	2	-	-
2001.....	9	-	-	-	3	-	1	3	1	1	-	-
2000.....	17	-	1	-	1	2	1	3	2	3	-	1
1999.....	12	-	-	-	2	3	3	-	2	-	-	-
Black, male												
2004.....	6	-	-	-	-	2	-	-	2	1	-	1
2003.....	11	-	-	-	-	2	2	3	2	1	-	-
2002.....	9	-	1	-	-	1	4	1	-	-	-	-
2001.....	4	-	-	-	1	-	1	1	1	-	-	-
2000.....	9	-	1	-	-	1	1	1	1	1	-	1
1999.....	6	-	-	-	-	2	2	-	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Cystic fibrosis (E84)

Black, female

2004.....	10	-	-	-	1	3	1	2	2	-	1	-
2003.....	14	-	1	-	3	2	2	2	1	-	1	-
2002.....	16	1	1	2	3	-	3	1	1	2	-	-
2001.....	5	-	-	-	2	-	-	2	-	1	-	-
2000.....	8	-	-	-	1	1	-	2	1	2	-	-
1999.....	6	-	-	-	2	1	1	-	1	-	-	-

American Indian 1/, both sexes

2004.....	1	-	-	-	1	-	-	-	-	-	-	-
2003.....	3	-	-	-	1	1	-	1	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	4	-	-	-	-	1	-	2	-	-	1	-
2000.....	2	-	-	-	-	1	-	-	1	-	-	-
1999.....	1	-	-	-	-	1	-	-	-	-	-	-

American Indian 1/, male

2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	1	-	-	-	-	-	-	1	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	4	-	-	-	-	1	-	2	-	-	1	-
2000.....	1	-	-	-	-	1	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-

American Indian 1/, female

2004.....	1	-	-	-	1	-	-	-	-	-	-	-
2003.....	2	-	-	-	1	1	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	1	-	-	-	-	-	-	-	1	-	-	-
1999.....	1	-	-	-	-	1	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Volume depletion, disorders of fluid, electrolyte and acid-base balance (E86-E87)

All races, both sexes

2004.....	6,550	66	12	5	6	5	15	14	22	33	54	107
2003.....	7,025	78	28	7	5	9	17	15	25	44	53	91
2002.....	7,345	74	18	8	3	9	14	16	23	33	60	99
2001.....	7,793	90	30	9	6	5	11	17	23	39	67	94
2000.....	7,780	73	13	4	7	12	13	17	17	33	80	87
1999.....	8,163	73	22	7	7	13	13	17	20	48	65	82

All races, male

2004.....	2,314	40	7	-	2	4	7	8	15	15	33	63
2003.....	2,431	45	12	4	-	6	9	8	16	16	28	51
2002.....	2,454	44	8	5	3	5	5	6	11	15	34	57
2001.....	2,586	46	14	5	2	3	7	7	11	20	32	54
2000.....	2,701	36	7	2	5	7	6	9	7	18	44	52
1999.....	2,697	45	11	5	6	7	6	11	8	25	40	50

All races, female

2004.....	4,236	26	5	5	4	1	8	6	7	18	21	44
2003.....	4,594	33	16	3	5	3	8	7	9	28	25	40
2002.....	4,891	30	10	3	-	4	9	10	12	18	26	42
2001.....	5,207	44	16	4	4	2	4	10	12	19	35	40
2000.....	5,079	37	6	2	2	5	7	8	10	15	36	35
1999.....	5,466	28	11	2	1	6	7	6	12	23	25	32

White, both sexes

2004.....	5,653	43	8	4	5	4	9	10	15	19	33	78
2003.....	5,984	43	19	5	4	4	11	10	19	27	35	64
2002.....	6,262	45	13	6	2	6	9	11	17	20	37	68
2001.....	6,667	54	25	5	3	3	6	12	19	30	49	57
2000.....	6,712	44	10	1	7	7	7	10	11	20	55	56
1999.....	7,084	40	15	5	3	11	8	13	14	29	36	58

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Volume depletion, disorders of fluid, electrolyte and acid-base balance (E86-E87)

White, male

2004.....	1,944	25	5	-	1	3	5	6	10	9	19	48
2003.....	2,030	26	8	3	-	2	6	5	11	11	18	37
2002.....	2,014	28	6	4	2	3	2	3	6	7	21	36
2001.....	2,144	26	12	3	-	2	3	4	10	15	21	33
2000.....	2,282	21	6	1	5	4	3	6	4	10	29	33
1999.....	2,209	25	6	4	2	5	3	7	5	13	20	37

White, female

2004.....	3,709	18	3	4	4	1	4	4	5	10	14	30
2003.....	3,954	17	11	2	4	2	5	5	8	16	17	27
2002.....	4,248	17	7	2	-	3	7	8	11	13	16	32
2001.....	4,523	28	13	2	3	1	3	8	9	15	28	24
2000.....	4,430	23	4	-	2	3	4	4	7	10	26	23
1999.....	4,875	15	9	1	1	6	5	6	9	16	16	21

Black, both sexes

2004.....	799	19	1	-	1	1	5	4	6	11	21	28
2003.....	931	30	7	1	1	4	5	4	5	13	18	22
2002.....	964	24	4	1	1	3	4	5	3	11	22	31
2001.....	1,027	33	3	4	3	2	5	4	3	8	15	34
2000.....	976	20	2	2	-	5	5	7	6	13	23	30
1999.....	977	29	6	2	4	1	4	3	6	17	29	23

Black, male

2004.....	338	13	-	-	1	1	2	2	4	5	14	14
2003.....	363	17	2	1	-	3	2	2	4	3	10	14
2002.....	383	13	1	1	1	2	3	3	3	6	13	21
2001.....	397	17	-	2	2	1	4	2	1	5	9	18
2000.....	384	12	-	1	-	3	3	3	3	8	13	19
1999.....	435	17	4	1	4	1	2	3	3	11	20	12

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Volume depletion, disorders of fluid, electrolyte and acid-base balance (E86-E87)

Asian or Pacific Islander 2/, both sexes

2004.....	72	4	3	1	-	-	-	-	1	3	-	-
2003.....	74	3	1	1	-	1	-	1	-	4	-	3
2002.....	87	4	1	-	-	-	-	-	2	2	1	-
2001.....	67	3	1	-	-	-	-	1	1	1	1	2
2000.....	61	4	-	1	-	-	1	-	-	-	1	-
1999.....	79	1	1	-	-	1	-	-	-	2	-	-

Asian or Pacific Islander 2/, male

2004.....	26	2	2	-	-	-	-	-	1	1	-	-
2003.....	30	1	1	-	-	1	-	1	-	2	-	-
2002.....	46	3	1	-	-	-	-	-	1	2	-	-
2001.....	31	3	1	-	-	-	-	1	-	-	1	2
2000.....	24	2	-	-	-	-	-	-	-	-	1	-
1999.....	42	1	1	-	-	1	-	-	-	1	-	-

Asian or Pacific Islander 2/, female

2004.....	46	2	1	1	-	-	-	-	-	2	-	-
2003.....	44	2	-	1	-	-	-	-	-	2	-	3
2002.....	41	1	-	-	-	-	-	-	1	-	1	-
2001.....	36	-	-	-	-	-	-	-	1	1	-	-
2000.....	37	2	-	1	-	-	1	-	-	-	-	-
1999.....	37	-	-	-	-	-	-	-	-	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other metabolic disorders (E70-E83,E85,E88)

All races, both sexes

2004.....	8,924	127	105	64	48	37	39	28	39	68	139	256
2003.....	8,473	125	134	62	63	54	33	29	36	71	145	241
2002.....	8,000	138	106	55	61	41	39	35	38	62	143	218
2001.....	7,422	148	123	66	47	47	24	24	31	63	124	224
2000.....	6,851	123	116	49	59	42	41	35	32	54	99	200
1999.....	6,226	128	106	61	51	46	24	27	32	59	108	173

All races, male

2004.....	4,782	79	58	34	30	18	25	16	23	42	90	186
2003.....	4,579	71	77	40	45	33	22	16	17	47	106	161
2002.....	4,295	75	52	38	39	27	29	22	22	43	99	143
2001.....	4,045	85	67	33	28	30	13	14	17	41	82	162
2000.....	3,624	71	63	29	41	23	25	25	26	30	69	153
1999.....	3,226	71	62	32	29	30	9	14	16	39	64	122

All races, female

2004.....	4,142	48	47	30	18	19	14	12	16	26	49	70
2003.....	3,894	54	57	22	18	21	11	13	19	24	39	80
2002.....	3,705	63	54	17	22	14	10	13	16	19	44	75
2001.....	3,377	63	56	33	19	17	11	10	14	22	42	62
2000.....	3,227	52	53	20	18	19	16	10	6	24	30	47
1999.....	3,000	57	44	29	22	16	15	13	16	20	44	51

White, both sexes

2004.....	7,936	108	90	53	41	32	35	24	33	53	114	215
2003.....	7,479	97	110	56	55	44	29	22	30	57	117	201
2002.....	7,019	104	86	44	53	37	30	30	28	52	121	179
2001.....	6,597	116	100	58	41	38	23	20	24	54	111	181
2000.....	6,101	101	99	45	50	37	32	27	21	47	83	180
1999.....	5,516	93	93	44	44	36	21	21	24	46	87	145

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	-------------	-----------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Other metabolic disorders (E70-E83,E85,E88)

White, male

2004.....	4,286	69	50	28	24	14	22	14	19	32	74	159
2003.....	4,068	60	62	36	40	28	21	13	12	39	86	137
2002.....	3,796	64	41	34	33	24	21	17	18	36	88	120
2001.....	3,608	65	56	30	23	25	13	13	14	34	75	133
2000.....	3,260	58	57	25	35	19	17	19	18	27	60	140
1999.....	2,897	54	55	25	24	24	8	10	10	33	58	101

White, female

2004.....	3,650	39	40	25	17	18	13	10	14	21	40	56
2003.....	3,411	37	48	20	15	16	8	9	18	18	31	64
2002.....	3,223	40	45	10	20	13	9	13	10	16	33	59
2001.....	2,989	51	44	28	18	13	10	7	10	20	36	48
2000.....	2,841	43	42	20	15	18	15	8	3	20	23	40
1999.....	2,619	39	38	19	20	12	13	11	14	13	29	44

Black, both sexes

2004.....	819	15	11	3	6	3	3	3	5	13	22	37
2003.....	781	22	12	4	7	5	3	4	5	13	20	33
2002.....	818	26	14	9	7	3	8	4	8	10	17	34
2001.....	666	24	13	5	4	5	1	3	6	5	10	36
2000.....	632	18	12	3	6	5	8	7	9	7	16	18
1999.....	598	27	9	14	6	7	2	5	8	10	20	20

Black, male

2004.....	400	8	7	2	5	2	2	2	3	8	14	24
2003.....	385	6	9	2	4	1	1	2	4	7	14	20
2002.....	407	10	7	3	5	3	7	4	2	7	8	20
2001.....	343	13	7	2	3	4	-	1	2	4	4	24
2000.....	298	11	5	3	4	4	7	5	8	3	9	12
1999.....	265	15	4	5	4	5	1	3	6	4	6	15

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other metabolic disorders (E70-E83,E85,E88)

Black, female

2004.....	419	7	4	1	1	1	1	1	2	5	8	13
2003.....	396	16	3	2	3	4	2	2	1	6	6	13
2002.....	411	16	7	6	2	-	1	-	6	3	9	14
2001.....	323	11	6	3	1	1	1	2	4	1	6	12
2000.....	334	7	7	-	2	1	1	2	1	4	7	6
1999.....	333	12	5	9	2	2	1	2	2	6	14	5

American Indian 1/, both sexes

2004.....	31	1	2	2	-	-	-	-	-	-	2	2
2003.....	35	2	3	1	-	1	-	-	-	-	1	2
2002.....	39	4	2	-	-	-	-	-	-	-	2	2
2001.....	29	-	-	1	1	-	-	-	-	2	-	4
2000.....	27	-	4	-	1	-	-	1	-	-	-	1
1999.....	23	3	-	1	-	2	-	1	-	1	-	2

American Indian 1/, male

2004.....	17	-	-	-	-	-	-	-	-	-	1	1
2003.....	19	1	2	1	-	1	-	-	-	-	-	2
2002.....	22	1	2	-	-	-	-	-	-	-	1	1
2001.....	18	-	-	-	1	-	-	-	-	1	-	3
2000.....	18	-	1	-	1	-	-	1	-	-	-	-
1999.....	13	-	-	1	-	-	-	1	-	1	-	2

American Indian 1/, female

2004.....	14	1	2	2	-	-	-	-	-	-	1	1
2003.....	16	1	1	-	-	-	-	-	-	-	1	-
2002.....	17	3	-	-	-	-	-	-	-	-	1	1
2001.....	11	-	-	1	-	-	-	-	-	1	-	1
2000.....	9	-	3	-	-	-	-	-	-	-	-	1
1999.....	10	3	-	-	-	2	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other metabolic disorders (E70-E83,E85,E88)												
Asian or Pacific Islander 2/, both sexes												
2004.....	138	3	2	6	1	2	1	1	1	2	1	2
2003.....	178	4	9	1	1	4	1	3	1	1	7	5
2002.....	124	4	4	2	1	1	1	1	2	-	3	3
2001.....	130	8	10	2	1	4	-	1	1	2	3	3
2000.....	91	4	1	1	2	-	1	-	2	-	-	1
1999.....	89	5	4	2	1	1	1	-	-	2	1	6
Asian or Pacific Islander 2/, male												
2004.....	79	2	1	4	1	2	1	-	1	2	1	2
2003.....	107	4	4	1	1	3	-	1	1	1	6	2
2002.....	70	-	2	1	1	-	1	1	2	-	2	2
2001.....	76	7	4	1	1	1	-	-	1	2	3	2
2000.....	48	2	-	1	1	-	1	-	-	-	-	1
1999.....	51	2	3	1	1	1	-	-	-	1	-	4
Asian or Pacific Islander 2/, female												
2004.....	59	1	1	2	-	-	-	1	-	-	-	-
2003.....	71	-	5	-	-	1	1	2	-	-	1	3
2002.....	54	4	2	1	-	1	-	-	-	-	1	1
2001.....	54	1	6	1	-	3	-	1	-	-	-	1
2000.....	43	2	1	-	1	-	-	-	2	-	-	-
1999.....	38	3	1	1	-	-	1	-	-	1	1	2

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
V. Mental and behavioral disorders (F01-F99)												
All races, both sexes												
2004.....	61,524	2	9	5	24	90	215	316	438	796	1,390	1,913
2003.....	60,058	1	10	11	17	82	199	292	431	854	1,446	1,828
2002.....	55,788	8	10	13	22	70	168	239	420	873	1,488	1,689
2001.....	50,811	6	10	10	16	79	153	268	454	823	1,368	1,651
2000.....	46,015	13	11	11	18	79	175	256	471	880	1,311	1,507
1999.....	42,128	12	11	12	24	71	147	260	478	935	1,352	1,457
All races, male												
2004.....	23,382	2	4	4	14	54	148	240	311	577	1,014	1,367
2003.....	22,423	-	6	3	7	48	150	223	303	606	1,045	1,360
2002.....	21,000	3	5	5	12	46	119	176	281	611	1,066	1,256
2001.....	19,343	4	5	5	5	52	121	198	332	582	987	1,248
2000.....	17,672	7	5	7	8	53	129	188	339	676	946	1,146
1999.....	16,763	9	3	7	14	56	106	199	337	688	1,038	1,113
All races, female												
2004.....	38,142	-	5	1	10	36	67	76	127	219	376	546
2003.....	37,635	1	4	8	10	34	49	69	128	248	401	468
2002.....	34,788	5	5	8	10	24	49	63	139	262	422	433
2001.....	31,468	2	5	5	11	27	32	70	122	241	381	403
2000.....	28,343	6	6	4	10	26	46	68	132	204	365	361
1999.....	25,365	3	8	5	10	15	41	61	141	247	314	344
White, both sexes												
2004.....	54,979	2	5	3	17	74	187	255	339	645	1,125	1,526
2003.....	53,776	1	7	6	15	67	164	239	340	672	1,138	1,452
2002.....	49,967	5	9	12	16	61	138	193	344	688	1,173	1,323
2001.....	45,444	2	7	9	13	63	126	213	360	648	1,026	1,226
2000.....	41,082	9	4	9	15	67	141	211	362	687	1,000	1,116
1999.....	37,693	7	8	8	17	62	117	213	395	700	1,006	1,066

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
V. Mental and behavioral disorders (F01-F99)												
White, male												
2004.....	20,309	2	2	2	9	45	133	192	237	474	836	1,116
2003.....	19,507	-	4	1	7	39	125	189	244	486	845	1,088
2002.....	18,205	2	5	5	8	41	101	144	236	489	865	993
2001.....	16,672	2	3	5	4	43	97	156	266	462	753	935
2000.....	15,174	5	1	6	8	44	101	154	263	535	733	857
1999.....	14,390	6	3	5	11	47	85	164	288	529	792	833
White, female												
2004.....	34,670	-	3	1	8	29	54	63	102	171	289	410
2003.....	34,269	1	3	5	8	28	39	50	96	186	293	364
2002.....	31,762	3	4	7	8	20	37	49	108	199	308	330
2001.....	28,772	-	4	4	9	20	29	57	94	186	273	291
2000.....	25,908	4	3	3	7	23	40	57	99	152	267	259
1999.....	23,303	1	5	3	6	15	32	49	107	171	214	233
Black, both sexes												
2004.....	5,565	-	4	2	6	8	19	47	68	115	198	339
2003.....	5,343	-	2	3	1	9	22	42	62	149	246	320
2002.....	4,999	3	-	-	4	5	22	33	65	141	277	322
2001.....	4,674	3	2	1	2	13	23	38	65	145	296	378
2000.....	4,326	4	7	2	2	12	30	38	92	155	263	359
1999.....	3,873	4	2	3	5	7	24	36	66	201	290	345
Black, male												
2004.....	2,581	-	2	2	4	5	9	40	54	76	136	220
2003.....	2,441	-	2	1	-	7	15	29	36	99	157	232
2002.....	2,360	1	-	-	3	2	15	23	39	94	174	231
2001.....	2,307	2	1	-	1	7	20	32	45	98	201	280
2000.....	2,180	2	4	1	-	9	24	29	60	111	179	268
1999.....	2,047	2	-	2	3	7	16	26	40	136	199	245

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

V. Mental and behavioral disorders (F01-F99)

Black, female

2004.....	2,984	-	2	-	2	3	10	7	14	39	62	119
2003.....	2,902	-	-	2	1	2	7	13	26	50	89	88
2002.....	2,639	2	-	-	1	3	7	10	26	47	103	91
2001.....	2,367	1	1	1	1	6	3	6	20	47	95	98
2000.....	2,146	2	3	1	2	3	6	9	32	44	84	91
1999.....	1,826	2	2	1	2	-	8	10	26	65	91	100

American Indian 1/, both sexes

2004.....	379	-	-	-	1	6	8	11	24	20	53	35
2003.....	377	-	1	1	-	6	10	3	16	25	48	45
2002.....	317	-	-	-	1	3	6	10	9	30	28	29
2001.....	318	1	-	-	1	3	2	14	23	27	37	39
2000.....	258	-	-	-	-	-	2	5	14	33	41	23
1999.....	291	-	1	-	-	2	2	9	13	29	46	37

American Indian 1/, male

2004.....	226	-	-	-	1	3	5	5	14	13	32	23
2003.....	225	-	-	-	-	2	8	2	12	14	30	31
2002.....	190	-	-	-	1	2	2	8	4	19	18	19
2001.....	192	-	-	-	-	2	2	9	16	19	25	27
2000.....	166	-	-	-	-	-	2	4	14	25	29	14
1999.....	192	-	-	-	-	2	2	7	8	19	37	27

American Indian 1/, female

2004.....	153	-	-	-	-	3	3	6	10	7	21	12
2003.....	152	-	1	1	-	4	2	1	4	11	18	14
2002.....	127	-	-	-	-	1	4	2	5	11	10	10
2001.....	126	1	-	-	1	1	-	5	7	8	12	12
2000.....	92	-	-	-	-	-	-	1	-	8	12	9
1999.....	99	-	1	-	-	-	-	2	5	10	9	10

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
V. Mental and behavioral disorders (F01-F99)												
Asian or Pacific Islander 2/, both sexes												
2004.....	601	-	-	-	-	2	1	3	7	16	14	13
2003.....	562	-	-	1	1	-	3	8	13	8	14	11
2002.....	505	-	1	1	1	1	2	3	2	14	10	15
2001.....	375	-	1	-	-	-	2	3	6	3	9	8
2000.....	349	-	-	-	1	-	2	2	3	5	7	9
1999.....	271	1	-	1	2	-	4	2	4	5	10	9
Asian or Pacific Islander 2/, male												
2004.....	266	-	-	-	-	1	1	3	6	14	10	8
2003.....	250	-	-	1	-	-	2	3	11	7	13	9
2002.....	245	-	-	-	-	1	1	1	2	9	9	13
2001.....	172	-	1	-	-	-	2	1	5	3	8	6
2000.....	152	-	-	-	-	-	2	1	2	5	5	7
1999.....	134	1	-	-	-	-	3	2	1	4	10	8
Asian or Pacific Islander 2/, female												
2004.....	335	-	-	-	-	1	-	-	1	2	4	5
2003.....	312	-	-	-	1	-	1	5	2	1	1	2
2002.....	260	-	1	1	1	-	1	2	-	5	1	2
2001.....	203	-	-	-	-	-	-	2	1	-	1	2
2000.....	197	-	-	-	1	-	-	1	1	-	2	2
1999.....	137	-	-	1	2	-	1	-	3	1	-	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Organic dementia (F01,F03)												
All races, both sexes												
2004.....	46,662	-	-	-	-	-	-	1	1	-	8	21
2003.....	45,550	-	-	-	-	-	1	1	-	4	4	12
2002.....	41,839	-	-	-	-	1	-	-	3	-	10	21
2001.....	37,327	-	-	-	-	-	-	-	-	1	9	17
2000.....	32,822	-	-	-	-	-	-	-	1	3	5	22
1999.....	28,709	-	-	-	-	-	-	-	-	2	5	9
All races, male												
2004.....	13,750	-	-	-	-	-	-	-	1	-	3	11
2003.....	13,242	-	-	-	-	-	1	1	-	3	4	7
2002.....	12,283	-	-	-	-	-	-	-	1	-	5	10
2001.....	10,926	-	-	-	-	-	-	-	-	1	6	13
2000.....	9,505	-	-	-	-	-	-	-	-	2	1	9
1999.....	8,477	-	-	-	-	-	-	-	-	1	5	3
All races, female												
2004.....	32,912	-	-	-	-	-	-	1	-	-	5	10
2003.....	32,308	-	-	-	-	-	-	-	-	1	-	5
2002.....	29,556	-	-	-	-	1	-	-	2	-	5	11
2001.....	26,401	-	-	-	-	-	-	-	-	-	3	4
2000.....	23,317	-	-	-	-	-	-	-	1	1	4	13
1999.....	20,232	-	-	-	-	-	-	-	-	1	-	6
White, both sexes												
2004.....	42,531	-	-	-	-	-	-	1	1	-	7	19
2003.....	41,656	-	-	-	-	-	1	1	-	4	4	6
2002.....	38,391	-	-	-	-	1	-	-	2	-	5	16
2001.....	34,369	-	-	-	-	-	-	-	-	1	8	13
2000.....	30,154	-	-	-	-	-	-	-	-	2	5	18
1999.....	26,651	-	-	-	-	-	-	-	-	1	5	6

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Organic dementia (F01,F03)												
White, male												
2004.....	12,392	-	-	-	-	-	-	-	1	-	3	11
2003.....	11,979	-	-	-	-	-	1	1	-	3	4	3
2002.....	11,124	-	-	-	-	-	-	-	-	-	1	5
2001.....	9,962	-	-	-	-	-	-	-	-	1	5	9
2000.....	8,601	-	-	-	-	-	-	-	-	1	1	6
1999.....	7,765	-	-	-	-	-	-	-	-	-	5	2
White, female												
2004.....	30,139	-	-	-	-	-	-	1	-	-	4	8
2003.....	29,677	-	-	-	-	-	-	-	-	1	-	3
2002.....	27,267	-	-	-	-	1	-	-	2	-	4	11
2001.....	24,407	-	-	-	-	-	-	-	-	-	3	4
2000.....	21,553	-	-	-	-	-	-	-	-	1	4	12
1999.....	18,886	-	-	-	-	-	-	-	-	1	-	4
Black, both sexes												
2004.....	3,562	-	-	-	-	-	-	-	-	-	1	1
2003.....	3,354	-	-	-	-	-	-	-	-	-	-	6
2002.....	2,970	-	-	-	-	-	-	-	1	-	5	4
2001.....	2,589	-	-	-	-	-	-	-	-	-	1	3
2000.....	2,326	-	-	-	-	-	-	-	1	1	-	4
1999.....	1,799	-	-	-	-	-	-	-	-	1	-	3
Black, male												
2004.....	1,154	-	-	-	-	-	-	-	-	-	-	-
2003.....	1,062	-	-	-	-	-	-	-	-	-	-	4
2002.....	957	-	-	-	-	-	-	-	1	-	4	4
2001.....	822	-	-	-	-	-	-	-	-	-	1	3
2000.....	776	-	-	-	-	-	-	-	-	1	-	3
1999.....	608	-	-	-	-	-	-	-	-	1	-	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other and unspecified organic mental disorders (F04-F09)

All races, both sexes

2004.....	1,768	1	1	-	-	-	2	8	2	4	5	9
2003.....	1,875	-	-	-	-	-	3	7	4	6	3	9
2002.....	1,986	1	-	-	-	1	3	1	1	4	7	18
2001.....	2,153	-	-	-	-	1	2	5	9	7	12	7
2000.....	2,287	2	-	1	-	1	2	2	5	5	10	8
1999.....	2,455	2	-	1	-	-	2	5	2	5	4	8

All races, male

2004.....	624	1	1	-	-	-	1	7	1	2	4	6
2003.....	654	-	-	-	-	-	3	7	2	4	2	6
2002.....	624	-	-	-	-	1	3	-	-	4	5	11
2001.....	697	-	-	-	-	-	2	5	8	6	11	2
2000.....	705	2	-	1	-	-	2	1	4	3	7	4
1999.....	758	1	-	1	-	-	2	5	2	5	4	6

All races, female

2004.....	1,144	-	-	-	-	-	1	1	1	2	1	3
2003.....	1,221	-	-	-	-	-	-	-	2	2	1	3
2002.....	1,362	1	-	-	-	-	-	1	1	-	2	7
2001.....	1,456	-	-	-	-	1	-	-	1	1	1	5
2000.....	1,582	-	-	-	-	1	-	1	1	2	3	4
1999.....	1,697	1	-	-	-	-	-	-	-	-	-	2

White, both sexes

2004.....	1,631	1	1	-	-	-	2	4	-	1	4	5
2003.....	1,704	-	-	-	-	-	2	4	2	5	2	6
2002.....	1,822	1	-	-	-	1	3	-	1	4	5	15
2001.....	1,968	-	-	-	-	1	2	4	6	6	8	4
2000.....	2,103	2	-	1	-	1	2	1	4	3	6	4
1999.....	2,248	1	-	1	-	-	1	4	1	3	3	5

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other and unspecified organic mental disorders (F04-F09)

White, male

2004.....	570	1	1	-	-	-	1	4	-	-	4	4
2003.....	580	-	-	-	-	-	2	4	1	3	1	3
2002.....	569	-	-	-	-	1	3	-	-	4	3	9
2001.....	624	-	-	-	-	-	2	4	5	5	7	1
2000.....	631	2	-	1	-	-	2	-	3	2	3	2
1999.....	671	1	-	1	-	-	1	4	1	3	3	4

White, female

2004.....	1,061	-	-	-	-	-	1	-	-	1	-	1
2003.....	1,124	-	-	-	-	-	-	-	1	2	1	3
2002.....	1,253	1	-	-	-	-	-	-	1	-	2	6
2001.....	1,344	-	-	-	-	1	-	-	1	1	1	3
2000.....	1,472	-	-	-	-	1	-	1	1	1	3	2
1999.....	1,577	-	-	-	-	-	-	-	-	-	-	1

Black, both sexes

2004.....	125	-	-	-	-	-	-	4	2	3	1	4
2003.....	148	-	-	-	-	-	1	2	2	1	1	2
2002.....	142	-	-	-	-	-	-	-	-	-	2	3
2001.....	170	-	-	-	-	-	-	1	2	1	4	3
2000.....	166	-	-	-	-	-	-	1	1	2	4	3
1999.....	189	1	-	-	-	-	1	1	1	2	1	3

Black, male

2004.....	47	-	-	-	-	-	-	3	1	2	-	2
2003.....	65	-	-	-	-	-	1	2	1	1	1	2
2002.....	44	-	-	-	-	-	-	-	-	-	2	2
2001.....	66	-	-	-	-	-	-	1	2	1	4	1
2000.....	62	-	-	-	-	-	-	1	1	1	4	2
1999.....	79	-	-	-	-	-	1	1	1	2	1	2

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Mental and behavioral disorders due to psychoactive substance use (F10-F19)

All races, both sexes

2004.....	10,683	-	-	-	4	65	181	274	403	737	1,286	1,760
2003.....	10,104	-	-	-	5	49	158	248	393	785	1,364	1,698
2002.....	9,528	3	-	-	4	49	135	209	374	814	1,378	1,549
2001.....	9,087	2	-	-	3	59	127	227	404	760	1,264	1,547
2000.....	8,721	2	2	-	4	49	140	221	424	826	1,229	1,402
1999.....	8,619	1	-	-	2	53	120	209	427	860	1,268	1,365

All races, male

2004.....	8,041	-	-	-	3	46	136	216	294	546	959	1,282
2003.....	7,578	-	-	-	3	32	131	199	285	565	1,005	1,285
2002.....	7,192	-	-	-	2	36	101	162	268	578	1,008	1,180
2001.....	6,842	2	-	-	-	46	106	174	298	545	926	1,185
2000.....	6,629	1	1	-	3	34	113	167	311	642	904	1,094
1999.....	6,604	-	-	-	1	44	92	171	307	642	990	1,059

All races, female

2004.....	2,642	-	-	-	1	19	45	58	109	191	327	478
2003.....	2,526	-	-	-	2	17	27	49	108	220	359	413
2002.....	2,336	3	-	-	2	13	34	47	106	236	370	369
2001.....	2,245	-	-	-	3	13	21	53	106	215	338	362
2000.....	2,092	1	1	-	1	15	27	54	113	184	325	308
1999.....	2,015	1	-	-	1	9	28	38	120	218	278	306

White, both sexes

2004.....	8,659	-	-	-	4	56	160	228	313	596	1,047	1,397
2003.....	8,133	-	-	-	4	41	133	204	310	617	1,074	1,351
2002.....	7,570	2	-	-	3	43	113	171	307	638	1,091	1,206
2001.....	7,084	1	-	-	2	50	107	182	321	596	943	1,147
2000.....	6,827	2	-	-	4	44	113	184	322	647	929	1,028
1999.....	6,649	-	-	-	2	45	99	169	350	643	931	991

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Mental and behavioral disorders due to psychoactive substance use (F10-F19)

White, male

2004.....	6,512	-	-	-	3	40	123	181	227	447	795	1,041
2003.....	6,104	-	-	-	3	28	111	171	230	455	813	1,027
2002.....	5,731	-	-	-	1	33	88	134	227	463	825	931
2001.....	5,322	1	-	-	-	39	86	141	241	431	705	889
2000.....	5,192	1	-	-	3	30	89	138	239	512	702	814
1999.....	5,128	-	-	-	1	36	76	139	262	494	749	788

White, female

2004.....	2,147	-	-	-	1	16	37	47	86	149	252	356
2003.....	2,029	-	-	-	1	13	22	33	80	162	261	324
2002.....	1,839	2	-	-	2	10	25	37	80	175	266	275
2001.....	1,762	-	-	-	2	11	21	41	80	165	238	258
2000.....	1,635	1	-	-	1	14	24	46	83	135	227	214
1999.....	1,521	-	-	-	1	9	23	30	88	149	182	203

Black, both sexes

2004.....	1,663	-	-	-	-	2	12	33	60	106	177	319
2003.....	1,633	-	-	-	1	2	12	37	55	137	230	294
2002.....	1,678	1	-	-	-	3	14	28	57	136	251	303
2001.....	1,727	1	-	-	-	7	16	31	58	134	276	357
2000.....	1,666	-	2	-	-	5	24	31	85	141	252	345
1999.....	1,710	1	-	-	-	6	16	29	61	184	281	328

Black, male

2004.....	1,269	-	-	-	-	2	7	28	48	72	125	210
2003.....	1,224	-	-	-	-	2	10	25	33	91	149	219
2002.....	1,260	-	-	-	-	1	10	20	36	90	157	219
2001.....	1,318	1	-	-	-	5	16	25	39	92	189	264
2000.....	1,264	-	1	-	-	4	21	25	56	100	168	259
1999.....	1,273	-	-	-	-	6	11	23	36	126	194	236

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Mental and behavioral disorders due to psychoactive substance use (F10-F19)

Black, female

2004.....	394	-	-	-	-	-	5	5	12	34	52	109
2003.....	409	-	-	-	1	-	2	12	22	46	81	75
2002.....	418	1	-	-	-	2	4	8	21	46	94	84
2001.....	409	-	-	-	-	2	-	6	19	42	87	93
2000.....	402	-	1	-	-	1	3	6	29	41	84	86
1999.....	437	1	-	-	-	-	5	6	25	58	87	92

American Indian 1/, both sexes

2004.....	264	-	-	-	-	6	8	11	24	20	50	33
2003.....	262	-	-	-	-	6	10	3	15	25	47	44
2002.....	197	-	-	-	1	3	6	9	9	29	26	26
2001.....	225	-	-	-	1	2	2	12	23	27	37	37
2000.....	182	-	-	-	-	-	2	5	14	33	41	22
1999.....	209	-	-	-	-	2	2	9	13	28	46	37

American Indian 1/, male

2004.....	178	-	-	-	-	3	5	5	14	13	30	23
2003.....	184	-	-	-	-	2	8	2	11	14	30	31
2002.....	135	-	-	-	1	2	2	7	4	18	17	18
2001.....	161	-	-	-	-	2	2	7	16	19	25	27
2000.....	136	-	-	-	-	-	2	4	14	25	29	14
1999.....	159	-	-	-	-	2	2	7	8	18	37	27

American Indian 1/, female

2004.....	86	-	-	-	-	3	3	6	10	7	20	10
2003.....	78	-	-	-	-	4	2	1	4	11	17	13
2002.....	62	-	-	-	-	1	4	2	5	11	9	8
2001.....	64	-	-	-	1	-	-	5	7	8	12	10
2000.....	46	-	-	-	-	-	-	1	-	8	12	8
1999.....	50	-	-	-	-	-	-	2	5	10	9	10

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Mental and behavioral disorders due to psychoactive substance use (F10-F19)												
Asian or Pacific Islander 2/, both sexes												
2004.....	97	-	-	-	-	1	1	2	6	15	12	11
2003.....	76	-	-	-	-	-	3	4	13	6	13	9
2002.....	83	-	-	-	-	-	2	1	1	11	10	14
2001.....	51	-	-	-	-	-	2	2	2	3	8	6
2000.....	46	-	-	-	-	-	1	1	3	5	7	7
1999.....	51	-	-	-	-	-	3	2	3	5	10	9
Asian or Pacific Islander 2/, male												
2004.....	82	-	-	-	-	1	1	2	5	14	9	8
2003.....	66	-	-	-	-	-	2	1	11	5	13	8
2002.....	66	-	-	-	-	-	1	1	1	7	9	12
2001.....	41	-	-	-	-	-	2	1	2	3	7	5
2000.....	37	-	-	-	-	-	1	-	2	5	5	7
1999.....	44	-	-	-	-	-	3	2	1	4	10	8
Asian or Pacific Islander 2/, female												
2004.....	15	-	-	-	-	-	-	-	1	1	3	3
2003.....	10	-	-	-	-	-	1	3	2	1	-	1
2002.....	17	-	-	-	-	-	1	-	-	4	1	2
2001.....	10	-	-	-	-	-	-	1	-	-	1	1
2000.....	9	-	-	-	-	-	-	1	1	-	2	-
1999.....	7	-	-	-	-	-	-	-	2	1	-	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Mental and behavioral disorders due to use of alcohol (F10)

All races, both sexes

2004.....	7,242	-	-	-	4	23	61	97	199	412	813	1,204
2003.....	7,023	-	-	-	5	26	53	85	200	437	905	1,210
2002.....	6,842	1	-	-	2	22	42	87	172	488	908	1,122
2001.....	6,627	2	-	-	2	29	45	91	208	479	843	1,120
2000.....	6,262	2	1	-	2	19	38	85	203	500	814	1,026
1999.....	6,157	-	-	-	1	17	46	83	209	517	810	996

All races, male

2004.....	5,704	-	-	-	3	14	49	78	147	312	638	923
2003.....	5,465	-	-	-	3	18	49	73	157	325	693	929
2002.....	5,402	-	-	-	1	17	33	73	133	366	691	872
2001.....	5,171	2	-	-	-	21	43	70	155	363	639	878
2000.....	4,977	1	-	-	2	16	31	72	158	402	632	820
1999.....	4,858	-	-	-	-	13	37	72	148	391	646	787

All races, female

2004.....	1,538	-	-	-	1	9	12	19	52	100	175	281
2003.....	1,558	-	-	-	2	8	4	12	43	112	212	281
2002.....	1,440	1	-	-	1	5	9	14	39	122	217	250
2001.....	1,456	-	-	-	2	8	2	21	53	116	204	242
2000.....	1,285	1	1	-	-	3	7	13	45	98	182	206
1999.....	1,299	-	-	-	1	4	9	11	61	126	164	209

White, both sexes

2004.....	5,991	-	-	-	4	18	52	75	150	341	687	1,017
2003.....	5,773	-	-	-	4	20	39	71	158	360	745	1,005
2002.....	5,562	-	-	-	1	20	34	70	139	391	757	931
2001.....	5,260	1	-	-	2	22	34	71	161	390	654	891
2000.....	4,986	2	-	-	2	17	30	69	155	413	642	791
1999.....	4,846	-	-	-	1	14	40	69	173	395	632	774

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Mental and behavioral disorders due to use of alcohol (F10)

White, male

2004.....	4,707	-	-	-	3	11	44	63	110	259	545	780
2003.....	4,472	-	-	-	3	16	37	61	123	269	580	771
2002.....	4,390	-	-	-	-	16	28	58	108	299	587	722
2001.....	4,084	1	-	-	-	16	32	56	121	292	498	700
2000.....	3,959	1	-	-	2	14	24	58	118	339	506	640
1999.....	3,837	-	-	-	-	10	32	61	125	307	513	618

White, female

2004.....	1,284	-	-	-	1	7	8	12	40	82	142	237
2003.....	1,301	-	-	-	1	4	2	10	35	91	165	234
2002.....	1,172	-	-	-	1	4	6	12	31	92	170	209
2001.....	1,176	-	-	-	2	6	2	15	40	98	156	191
2000.....	1,027	1	-	-	-	3	6	11	37	74	136	151
1999.....	1,009	-	-	-	1	4	8	8	48	88	119	156

Black, both sexes

2004.....	967	-	-	-	-	1	5	12	26	44	78	152
2003.....	968	-	-	-	1	1	5	11	20	54	106	158
2002.....	1,037	1	-	-	-	-	3	8	25	65	118	157
2001.....	1,129	1	-	-	-	5	9	8	24	61	150	194
2000.....	1,071	-	1	-	-	2	6	13	34	53	131	207
1999.....	1,089	-	-	-	-	1	2	7	23	95	134	182

Black, male

2004.....	794	-	-	-	-	1	3	11	21	34	61	118
2003.....	780	-	-	-	-	1	4	10	16	39	76	124
2002.....	831	-	-	-	-	-	3	8	20	47	81	123
2001.....	913	1	-	-	-	3	9	8	18	50	113	153
2000.....	858	-	-	-	-	2	5	12	26	36	96	159
1999.....	847	-	-	-	-	1	1	5	15	66	97	137

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Mental and behavioral disorders due to use of alcohol (F10)

Black, female

2004.....	173	-	-	-	-	-	2	1	5	10	17	34
2003.....	188	-	-	-	1	-	1	1	4	15	30	34
2002.....	206	1	-	-	-	-	-	-	5	18	37	34
2001.....	216	-	-	-	-	2	-	-	6	11	37	41
2000.....	213	-	1	-	-	-	1	1	8	17	35	48
1999.....	242	-	-	-	-	-	1	2	8	29	37	45

American Indian 1/, both sexes

2004.....	219	-	-	-	-	4	4	9	20	17	39	28
2003.....	229	-	-	-	-	5	7	2	12	19	45	40
2002.....	179	-	-	-	1	2	4	9	7	25	25	24
2001.....	199	-	-	-	-	2	1	11	22	25	32	32
2000.....	164	-	-	-	-	-	1	3	12	29	36	21
1999.....	182	-	-	-	-	2	2	5	10	24	37	32

American Indian 1/, male

2004.....	148	-	-	-	-	2	2	3	13	10	24	20
2003.....	165	-	-	-	-	1	7	2	9	13	28	28
2002.....	126	-	-	-	1	1	1	7	4	15	16	17
2001.....	144	-	-	-	-	2	1	6	15	18	22	23
2000.....	126	-	-	-	-	-	1	2	12	22	27	14
1999.....	140	-	-	-	-	2	2	4	7	15	29	25

American Indian 1/, female

2004.....	71	-	-	-	-	2	2	6	7	7	15	8
2003.....	64	-	-	-	-	4	-	-	3	6	17	12
2002.....	53	-	-	-	-	1	3	2	3	10	9	7
2001.....	55	-	-	-	-	-	-	5	7	7	10	9
2000.....	38	-	-	-	-	-	-	1	-	7	9	7
1999.....	42	-	-	-	-	-	-	1	3	9	8	7

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Mental and behavioral disorders due to use of alcohol (F10)												
Asian or Pacific Islander 2/, both sexes												
2004.....	65	-	-	-	-	-	-	1	3	10	9	7
2003.....	53	-	-	-	-	-	2	1	10	4	9	7
2002.....	64	-	-	-	-	-	1	-	1	7	8	10
2001.....	39	-	-	-	-	-	1	1	1	3	7	3
2000.....	41	-	-	-	-	-	1	-	2	5	5	7
1999.....	40	-	-	-	-	-	2	2	3	3	7	8
Asian or Pacific Islander 2/, male												
2004.....	55	-	-	-	-	-	-	1	3	9	8	5
2003.....	48	-	-	-	-	-	1	-	9	4	9	6
2002.....	55	-	-	-	-	-	1	-	1	5	7	10
2001.....	30	-	-	-	-	-	1	-	1	3	6	2
2000.....	34	-	-	-	-	-	1	-	2	5	3	7
1999.....	34	-	-	-	-	-	2	2	1	3	7	7
Asian or Pacific Islander 2/, female												
2004.....	10	-	-	-	-	-	-	-	-	1	1	2
2003.....	5	-	-	-	-	-	1	1	1	-	-	1
2002.....	9	-	-	-	-	-	-	-	-	2	1	-
2001.....	9	-	-	-	-	-	-	1	-	-	1	1
2000.....	7	-	-	-	-	-	-	-	-	-	2	-
1999.....	6	-	-	-	-	-	-	-	2	-	-	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Mental and behavioral disorders due to other psychoactive substance use (F11-F19)

All races, both sexes

2004.....	3,441	-	-	-	-	42	120	177	204	325	473	556
2003.....	3,081	-	-	-	-	23	105	163	193	348	459	488
2002.....	2,686	2	-	-	2	27	93	122	202	326	470	427
2001.....	2,460	-	-	-	1	30	82	136	196	281	421	427
2000.....	2,459	-	1	-	2	30	102	136	221	326	415	376
1999.....	2,462	1	-	-	1	36	74	126	218	343	458	369

All races, male

2004.....	2,337	-	-	-	-	32	87	138	147	234	321	359
2003.....	2,113	-	-	-	-	14	82	126	128	240	312	356
2002.....	1,790	-	-	-	1	19	68	89	135	212	317	308
2001.....	1,671	-	-	-	-	25	63	104	143	182	287	307
2000.....	1,652	-	1	-	1	18	82	95	153	240	272	274
1999.....	1,746	-	-	-	1	31	55	99	159	251	344	272

All races, female

2004.....	1,104	-	-	-	-	10	33	39	57	91	152	197
2003.....	968	-	-	-	-	9	23	37	65	108	147	132
2002.....	896	2	-	-	1	8	25	33	67	114	153	119
2001.....	789	-	-	-	1	5	19	32	53	99	134	120
2000.....	807	-	-	-	1	12	20	41	68	86	143	102
1999.....	716	1	-	-	-	5	19	27	59	92	114	97

White, both sexes

2004.....	2,668	-	-	-	-	38	108	153	163	255	360	380
2003.....	2,360	-	-	-	-	21	94	133	152	257	329	346
2002.....	2,008	2	-	-	2	23	79	101	168	247	334	275
2001.....	1,824	-	-	-	-	28	73	111	160	206	289	256
2000.....	1,841	-	-	-	2	27	83	115	167	234	287	237
1999.....	1,803	-	-	-	1	31	59	100	177	248	299	217

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Mental and behavioral disorders due to other psychoactive substance use (F11-F19)

White, male

2004.....	1,805	-	-	-	-	29	79	118	117	188	250	261
2003.....	1,632	-	-	-	-	12	74	110	107	186	233	256
2002.....	1,341	-	-	-	1	17	60	76	119	164	238	209
2001.....	1,238	-	-	-	-	23	54	85	120	139	207	189
2000.....	1,233	-	-	-	1	16	65	80	121	173	196	174
1999.....	1,291	-	-	-	1	26	44	78	137	187	236	170

White, female

2004.....	863	-	-	-	-	9	29	35	46	67	110	119
2003.....	728	-	-	-	-	9	20	23	45	71	96	90
2002.....	667	2	-	-	1	6	19	25	49	83	96	66
2001.....	586	-	-	-	-	5	19	26	40	67	82	67
2000.....	608	-	-	-	1	11	18	35	46	61	91	63
1999.....	512	-	-	-	-	5	15	22	40	61	63	47

Black, both sexes

2004.....	696	-	-	-	-	1	7	21	34	62	99	167
2003.....	665	-	-	-	-	1	7	26	35	83	124	136
2002.....	641	-	-	-	-	3	11	20	32	71	133	146
2001.....	598	-	-	-	-	2	7	23	34	73	126	163
2000.....	595	-	1	-	-	3	18	18	51	88	121	138
1999.....	621	1	-	-	-	5	14	22	38	89	147	146

Black, male

2004.....	475	-	-	-	-	1	4	17	27	38	64	92
2003.....	444	-	-	-	-	1	6	15	17	52	73	95
2002.....	429	-	-	-	-	1	7	12	16	43	76	96
2001.....	405	-	-	-	-	2	7	17	21	42	76	111
2000.....	406	-	1	-	-	2	16	13	30	64	72	100
1999.....	426	-	-	-	-	5	10	18	21	60	97	99

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Mental and behavioral disorders due to other psychoactive substance use (F11-F19)

Black, female

2004.....	221	-	-	-	-	-	3	4	7	24	35	75
2003.....	221	-	-	-	-	-	1	11	18	31	51	41
2002.....	212	-	-	-	-	2	4	8	16	28	57	50
2001.....	193	-	-	-	-	-	-	6	13	31	50	52
2000.....	189	-	-	-	-	1	2	5	21	24	49	38
1999.....	195	1	-	-	-	-	4	4	17	29	50	47

American Indian 1/, both sexes

2004.....	45	-	-	-	-	2	4	2	4	3	11	5
2003.....	33	-	-	-	-	1	3	1	3	6	2	4
2002.....	18	-	-	-	-	1	2	-	2	4	1	2
2001.....	26	-	-	-	1	-	1	1	1	2	5	5
2000.....	18	-	-	-	-	-	1	2	2	4	5	1
1999.....	27	-	-	-	-	-	-	4	3	4	9	5

American Indian 1/, male

2004.....	30	-	-	-	-	1	3	2	1	3	6	3
2003.....	19	-	-	-	-	1	1	-	2	1	2	3
2002.....	9	-	-	-	-	1	1	-	-	3	1	1
2001.....	17	-	-	-	-	-	1	1	1	1	3	4
2000.....	10	-	-	-	-	-	1	2	2	3	2	-
1999.....	19	-	-	-	-	-	-	3	1	3	8	2

American Indian 1/, female

2004.....	15	-	-	-	-	1	1	-	3	-	5	2
2003.....	14	-	-	-	-	-	2	1	1	5	-	1
2002.....	9	-	-	-	-	-	1	-	2	1	-	1
2001.....	9	-	-	-	1	-	-	-	-	1	2	1
2000.....	8	-	-	-	-	-	-	-	-	1	3	1
1999.....	8	-	-	-	-	-	-	1	2	1	1	3

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Mental and behavioral disorders due to other psychoactive substance use (F11-F19)												
Asian or Pacific Islander 2/, both sexes												
2004.....	32	-	-	-	-	1	1	1	3	5	3	4
2003.....	23	-	-	-	-	-	1	3	3	2	4	2
2002.....	19	-	-	-	-	-	1	1	-	4	2	4
2001.....	12	-	-	-	-	-	1	1	1	-	1	3
2000.....	5	-	-	-	-	-	-	1	1	-	2	-
1999.....	11	-	-	-	-	-	1	-	-	2	3	1
Asian or Pacific Islander 2/, male												
2004.....	27	-	-	-	-	1	1	1	2	5	1	3
2003.....	18	-	-	-	-	-	1	1	2	1	4	2
2002.....	11	-	-	-	-	-	-	1	-	2	2	2
2001.....	11	-	-	-	-	-	1	1	1	-	1	3
2000.....	3	-	-	-	-	-	-	-	-	-	2	-
1999.....	10	-	-	-	-	-	1	-	-	1	3	1
Asian or Pacific Islander 2/, female												
2004.....	5	-	-	-	-	-	-	-	1	-	2	1
2003.....	5	-	-	-	-	-	-	2	1	1	-	-
2002.....	8	-	-	-	-	-	1	-	-	2	-	2
2001.....	1	-	-	-	-	-	-	-	-	-	-	-
2000.....	2	-	-	-	-	-	-	1	1	-	-	-
1999.....	1	-	-	-	-	-	-	-	-	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Schizophrenia, schizotypal and delusional disorders (F20-F29)

All races, both sexes

2004.....	513	-	-	-	-	-	1	2	5	10	21	26
2003.....	497	-	-	-	-	1	4	1	5	11	16	27
2002.....	508	-	1	-	-	-	1	5	6	8	20	21
2001.....	456	-	-	-	-	-	1	4	6	13	18	24
2000.....	498	-	-	-	-	2	1	4	6	6	16	17
1999.....	557	-	-	-	-	-	-	2	7	14	6	17

All races, male

2004.....	227	-	-	-	-	-	1	2	4	7	10	21
2003.....	181	-	-	-	-	1	2	-	4	9	7	17
2002.....	190	-	1	-	-	-	-	4	2	5	12	15
2001.....	187	-	-	-	-	-	1	3	6	9	13	12
2000.....	196	-	-	-	-	1	1	4	5	5	11	11
1999.....	210	-	-	-	-	-	-	1	7	12	4	8

All races, female

2004.....	286	-	-	-	-	-	-	-	1	3	11	5
2003.....	316	-	-	-	-	-	2	1	1	2	9	10
2002.....	318	-	-	-	-	-	1	1	4	3	8	6
2001.....	269	-	-	-	-	-	-	1	-	4	5	12
2000.....	302	-	-	-	-	1	-	-	1	1	5	6
1999.....	347	-	-	-	-	-	-	1	-	2	2	9

White, both sexes

2004.....	438	-	-	-	-	-	-	1	2	7	16	20
2003.....	428	-	-	-	-	-	3	1	2	5	9	17
2002.....	432	-	1	-	-	-	-	3	3	7	11	15
2001.....	397	-	-	-	-	-	1	2	5	9	15	19
2000.....	445	-	-	-	-	-	1	2	4	2	13	15
1999.....	498	-	-	-	-	-	-	2	6	10	5	16

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Schizophrenia, schizotypal and delusional disorders (F20-F29)												
White, male												
2004.....	183	-	-	-	-	-	-	1	1	6	8	17
2003.....	150	-	-	-	-	-	2	-	2	4	4	12
2002.....	154	-	1	-	-	-	-	3	1	4	7	12
2001.....	158	-	-	-	-	-	1	1	5	6	11	10
2000.....	173	-	-	-	-	-	1	2	4	2	8	10
1999.....	181	-	-	-	-	-	-	1	6	9	4	8
White, female												
2004.....	255	-	-	-	-	-	-	-	1	1	8	3
2003.....	278	-	-	-	-	-	1	1	-	1	5	5
2002.....	278	-	-	-	-	-	-	-	2	3	4	3
2001.....	239	-	-	-	-	-	-	1	-	3	4	9
2000.....	272	-	-	-	-	-	-	-	-	-	5	5
1999.....	317	-	-	-	-	-	-	1	-	1	1	8
Black, both sexes												
2004.....	71	-	-	-	-	-	1	1	2	3	4	6
2003.....	60	-	-	-	-	1	1	-	2	4	6	9
2002.....	68	-	-	-	-	-	1	1	3	1	9	5
2001.....	54	-	-	-	-	-	-	2	1	4	2	4
2000.....	50	-	-	-	-	2	-	1	2	4	3	2
1999.....	52	-	-	-	-	-	-	-	1	3	1	1
Black, male												
2004.....	40	-	-	-	-	-	1	1	2	1	1	4
2003.....	28	-	-	-	-	1	-	-	1	3	3	5
2002.....	32	-	-	-	-	-	-	-	1	1	5	3
2001.....	27	-	-	-	-	-	-	2	1	3	1	2
2000.....	21	-	-	-	-	1	-	1	1	3	3	1
1999.....	22	-	-	-	-	-	-	-	1	2	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Mood (affective) disorders (F30-F39)												
All races, both sexes												
2004.....	822	-	-	-	-	-	2	1	2	4	9	19
2003.....	926	-	-	-	-	2	2	1	4	7	10	12
2002.....	874	3	-	1	-	2	3	-	2	7	8	16
2001.....	879	3	-	-	-	2	2	2	3	8	21	14
2000.....	819	8	-	-	-	2	4	3	5	6	9	13
1999.....	850	6	-	-	-	3	1	5	7	7	12	8
All races, male												
2004.....	260	-	-	-	-	-	1	1	1	1	4	7
2003.....	280	-	-	-	-	1	1	1	2	7	1	4
2002.....	248	2	-	1	-	2	2	-	2	2	4	7
2001.....	294	2	-	-	-	1	2	2	1	6	10	12
2000.....	265	3	-	-	-	2	3	2	4	5	5	6
1999.....	300	5	-	-	-	2	1	5	6	6	8	7
All races, female												
2004.....	562	-	-	-	-	-	1	-	1	3	5	12
2003.....	646	-	-	-	-	1	1	-	2	-	9	8
2002.....	626	1	-	-	-	-	1	-	-	5	4	9
2001.....	585	1	-	-	-	1	-	-	2	2	11	2
2000.....	554	5	-	-	-	-	1	1	1	1	4	7
1999.....	550	1	-	-	-	1	-	-	1	1	4	1
White, both sexes												
2004.....	779	-	-	-	-	-	2	-	1	4	6	15
2003.....	879	-	-	-	-	1	2	1	3	6	8	10
2002.....	828	2	-	1	-	2	2	-	2	7	6	15
2001.....	828	1	-	-	-	1	2	-	3	8	19	9
2000.....	781	5	-	-	-	2	4	2	5	6	6	12
1999.....	819	4	-	-	-	3	1	5	7	7	11	8

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Neurotic, stress-related and somatoform disorders (F40-F48)												
All races, both sexes												
2004.....	120	1	-	-	-	-	-	1	-	1	3	5
2003.....	111	-	-	-	-	-	-	-	1	1	2	2
2002.....	117	1	-	-	-	-	3	-	1	-	5	3
2001.....	80	-	-	-	-	-	-	2	1	-	2	1
2000.....	73	-	-	-	-	3	-	1	1	-	3	-
1999.....	97	-	-	-	-	-	-	2	2	1	3	2
All races, male												
2004.....	52	1	-	-	-	-	-	1	-	1	1	1
2003.....	44	-	-	-	-	-	-	-	1	1	1	1
2002.....	52	1	-	-	-	-	2	-	-	-	2	2
2001.....	38	-	-	-	-	-	-	1	1	-	1	1
2000.....	30	-	-	-	-	1	-	1	1	-	-	-
1999.....	42	-	-	-	-	-	-	2	2	1	2	1
All races, female												
2004.....	68	-	-	-	-	-	-	-	-	-	2	4
2003.....	67	-	-	-	-	-	-	-	-	-	1	1
2002.....	65	-	-	-	-	-	1	-	1	-	3	1
2001.....	42	-	-	-	-	-	-	1	-	-	1	-
2000.....	43	-	-	-	-	2	-	-	-	-	3	-
1999.....	55	-	-	-	-	-	-	-	-	-	1	1
White, both sexes												
2004.....	107	1	-	-	-	-	-	-	-	-	1	3
2003.....	104	-	-	-	-	-	-	-	1	-	2	2
2002.....	99	-	-	-	-	-	3	-	1	-	4	3
2001.....	71	-	-	-	-	-	-	1	1	-	1	1
2000.....	68	-	-	-	-	2	-	1	1	-	3	-
1999.....	91	-	-	-	-	-	-	2	2	1	3	2

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Mental retardation (F70-F79)

All races, both sexes

2004.....	492	-	4	4	11	13	13	20	15	24	40	55
2003.....	487	1	4	5	6	11	16	22	15	27	34	47
2002.....	464	-	7	6	9	8	17	15	14	26	42	48
2001.....	423	-	6	7	6	10	11	13	25	25	32	36
2000.....	413	1	5	7	9	16	16	19	17	29	28	37
1999.....	447	2	8	8	18	13	17	26	27	35	37	34

All races, male

2004.....	283	-	2	4	8	6	8	12	8	15	25	33
2003.....	288	-	2	2	4	7	10	15	8	15	24	35
2002.....	263	-	3	3	7	5	10	10	7	17	23	28
2001.....	224	-	3	3	3	3	6	7	14	14	18	21
2000.....	231	1	1	5	4	12	9	12	9	18	16	22
1999.....	261	2	3	4	12	10	9	14	13	17	20	22

All races, female

2004.....	209	-	2	-	3	7	5	8	7	9	15	22
2003.....	199	1	2	3	2	4	6	7	7	12	10	12
2002.....	201	-	4	3	2	3	7	5	7	9	19	20
2001.....	199	-	3	4	3	7	5	6	11	11	14	15
2000.....	182	-	4	2	5	4	7	7	8	11	12	15
1999.....	186	-	5	4	6	3	8	12	14	18	17	12

White, both sexes

2004.....	424	-	2	2	7	9	8	13	12	22	29	53
2003.....	424	1	4	2	6	9	12	18	13	24	29	41
2002.....	402	-	6	6	4	6	11	11	9	19	36	44
2001.....	360	-	4	6	5	6	7	12	20	20	24	30
2000.....	362	-	2	5	7	13	10	16	15	23	27	33
1999.....	377	2	5	5	11	12	12	20	24	26	34	28

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Mental retardation (F70-F79)												
White, male												
2004.....	243	-	1	2	5	5	7	6	6	15	16	31
2003.....	251	-	2	-	4	6	7	12	7	14	21	33
2002.....	225	-	3	3	4	3	6	7	5	12	21	26
2001.....	187	-	2	3	2	2	4	6	11	12	14	16
2000.....	203	-	-	4	4	9	6	11	7	13	15	20
1999.....	226	2	3	3	9	9	7	12	11	13	18	19
White, female												
2004.....	181	-	1	-	2	4	1	7	6	7	13	22
2003.....	173	1	2	2	2	3	5	6	6	10	8	8
2002.....	177	-	3	3	-	3	5	4	4	7	15	18
2001.....	173	-	2	3	3	4	3	6	9	8	10	14
2000.....	159	-	2	1	3	4	4	5	8	10	12	13
1999.....	151	-	2	2	2	3	5	8	13	13	16	9
Black, both sexes												
2004.....	65	-	2	2	4	3	5	7	3	2	9	2
2003.....	59	-	-	2	-	2	4	3	2	3	5	6
2002.....	49	-	-	-	4	1	6	4	4	4	5	2
2001.....	57	-	2	1	1	4	4	-	3	5	8	5
2000.....	47	1	3	2	1	3	5	3	2	6	1	3
1999.....	65	-	2	2	5	1	5	6	3	9	3	6
Black, male												
2004.....	38	-	1	2	3	1	1	6	2	-	7	2
2003.....	33	-	-	1	-	1	3	2	1	1	3	2
2002.....	30	-	-	-	3	1	4	3	1	3	2	1
2001.....	34	-	1	-	1	1	2	-	2	2	4	5
2000.....	26	1	1	1	-	3	2	1	2	5	1	2
1999.....	35	-	-	1	3	1	2	2	2	4	2	3

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Mental retardation (F70-F79)												
Asian or Pacific Islander 2/, both sexes												
2004.....	1	-	-	-	-	1	-	-	-	-	-	-
2003.....	2	-	-	1	-	-	-	1	-	-	-	-
2002.....	9	-	1	-	1	1	-	-	1	2	-	1
2001.....	3	-	-	-	-	-	-	-	2	-	-	-
2000.....	3	-	-	-	1	-	1	-	-	-	-	-
1999.....	4	-	-	1	2	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	2	-	-	1	-	-	-	1	-	-	-	-
2002.....	6	-	-	-	-	1	-	-	1	1	-	1
2001.....	2	-	-	-	-	-	-	-	1	-	-	-
2000.....	2	-	-	-	-	-	1	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	1	-	-	-	-	1	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	3	-	1	-	1	-	-	-	-	1	-	-
2001.....	1	-	-	-	-	-	-	-	1	-	-	-
2000.....	1	-	-	-	1	-	-	-	-	-	-	-
1999.....	4	-	-	1	2	-	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other and unspecified mental and behavioral disorders (F50-F69,F80-F99)

All races, both sexes

2004.....	464	-	4	1	9	12	16	9	10	16	18	18
2003.....	508	-	6	6	6	19	15	12	9	13	13	21
2002.....	472	-	2	6	9	9	6	9	19	14	18	13
2001.....	406	1	4	3	7	7	10	15	6	9	10	5
2000.....	382	-	4	3	5	6	12	6	12	5	11	8
1999.....	394	1	3	3	4	2	7	11	6	11	17	14

All races, male

2004.....	145	-	1	-	3	2	1	1	2	5	8	6
2003.....	156	-	4	1	-	7	2	-	1	2	1	5
2002.....	148	-	1	1	3	2	1	-	1	5	7	3
2001.....	135	-	2	2	2	2	4	6	4	1	2	2
2000.....	111	-	3	1	1	3	1	1	5	1	2	-
1999.....	111	1	-	2	1	-	2	1	-	4	5	7

All races, female

2004.....	319	-	3	1	6	10	15	8	8	11	10	12
2003.....	352	-	2	5	6	12	13	12	8	11	12	16
2002.....	324	-	1	5	6	7	5	9	18	9	11	10
2001.....	271	1	2	1	5	5	6	9	2	8	8	3
2000.....	271	-	1	2	4	3	11	5	7	4	9	8
1999.....	283	-	3	1	3	2	5	10	6	7	12	7

White, both sexes

2004.....	410	-	2	1	6	9	15	8	10	15	15	14
2003.....	448	-	3	4	5	16	11	10	9	11	10	19
2002.....	423	-	2	5	9	8	6	8	19	13	15	9
2001.....	367	-	3	3	6	5	7	12	4	8	8	3
2000.....	342	-	2	3	4	5	11	5	11	4	11	6
1999.....	360	-	3	2	4	2	4	11	5	9	14	10

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other and unspecified mental and behavioral disorders (F50-F69,F80-F99)

White, male

2004.....	119	-	-	-	1	-	1	-	2	5	6	5
2003.....	135	-	2	1	-	4	1	-	1	1	-	5
2002.....	128	-	1	1	3	2	1	-	1	4	4	1
2001.....	116	-	1	2	2	2	2	4	2	1	2	1
2000.....	93	-	1	1	1	2	-	-	5	-	2	-
1999.....	94	-	-	1	1	-	-	1	-	3	3	4

White, female

2004.....	291	-	2	1	5	9	14	8	8	10	9	9
2003.....	313	-	1	3	5	12	10	10	8	10	10	14
2002.....	295	-	1	4	6	6	5	8	18	9	11	8
2001.....	251	-	2	1	4	3	5	8	2	7	6	2
2000.....	249	-	1	2	3	3	11	5	6	4	9	6
1999.....	266	-	3	1	3	2	4	10	5	6	11	6

Black, both sexes

2004.....	37	-	2	-	2	3	1	-	-	-	2	3
2003.....	46	-	2	1	-	3	4	-	-	2	2	1
2002.....	38	-	-	-	-	1	-	-	-	-	2	4
2001.....	31	-	-	-	1	1	3	2	1	1	2	2
2000.....	34	-	2	-	1	1	1	1	1	1	-	1
1999.....	27	-	-	1	-	-	2	-	-	2	3	4

Black, male

2004.....	17	-	1	-	1	2	-	-	-	-	2	1
2003.....	19	-	2	-	-	3	1	-	-	1	1	-
2002.....	14	-	-	-	-	-	-	-	-	-	2	2
2001.....	16	-	-	-	-	-	2	2	1	-	-	1
2000.....	17	-	2	-	-	1	1	1	-	1	-	-
1999.....	14	-	-	1	-	-	2	-	-	1	2	3

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified mental and behavioral disorders (F50-F69,F80-F99)												
Asian or Pacific Islander 2/, both sexes												
2004.....	14	-	-	-	-	-	-	1	-	1	1	1
2003.....	8	-	-	-	1	-	-	2	-	-	-	-
2002.....	8	-	-	1	-	-	-	1	-	1	-	-
2001.....	6	-	1	-	-	-	-	1	1	-	-	-
2000.....	5	-	-	-	-	-	-	-	-	-	-	1
1999.....	7	1	-	-	-	-	1	-	1	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	7	-	-	-	-	-	-	1	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	4	-	-	-	-	-	-	-	-	1	-	-
2001.....	3	-	1	-	-	-	-	-	1	-	-	-
2000.....	1	-	-	-	-	-	-	-	-	-	-	-
1999.....	3	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	7	-	-	-	-	-	-	-	-	1	1	1
2003.....	8	-	-	-	1	-	-	2	-	-	-	-
2002.....	4	-	-	1	-	-	-	1	-	-	-	-
2001.....	3	-	-	-	-	-	-	1	-	-	-	-
2000.....	4	-	-	-	-	-	-	-	-	-	-	1
1999.....	4	-	-	-	-	-	1	-	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

VI. Diseases of the nervous system (G00-G98)

All races, both sexes

2004.....	111,751	377	322	221	305	391	445	427	495	683	1,127	1,672
2003.....	109,640	400	280	251	296	399	496	437	493	703	1,149	1,680
2002.....	103,064	382	322	222	277	362	447	405	488	715	1,163	1,654
2001.....	96,524	360	315	229	263	404	434	388	476	710	1,137	1,514
2000.....	91,014	389	316	227	291	362	373	370	481	721	1,133	1,474
1999.....	84,077	441	361	263	253	375	415	381	447	738	1,026	1,369

All races, male

2004.....	43,313	218	164	122	183	275	278	263	278	378	643	913
2003.....	42,736	224	144	142	179	270	325	260	279	385	680	924
2002.....	40,332	228	192	126	161	244	292	250	297	436	649	894
2001.....	38,310	207	174	111	153	282	291	240	242	427	619	788
2000.....	36,238	225	169	112	169	230	249	228	267	397	633	817
1999.....	34,208	238	201	147	143	262	285	228	260	425	566	695

All races, female

2004.....	68,438	159	158	99	122	116	167	164	217	305	484	759
2003.....	66,904	176	136	109	117	129	171	177	214	318	469	756
2002.....	62,732	154	130	96	116	118	155	155	191	279	514	760
2001.....	58,214	153	141	118	110	122	143	148	234	283	518	726
2000.....	54,776	164	147	115	122	132	124	142	214	324	500	657
1999.....	49,869	203	160	116	110	113	130	153	187	313	460	674

White, both sexes

2004.....	101,812	290	219	159	225	284	328	321	348	526	908	1,340
2003.....	100,267	286	197	184	202	311	385	337	363	515	912	1,337
2002.....	94,455	290	218	167	193	275	319	301	361	542	920	1,330
2001.....	88,364	249	220	171	196	298	329	272	346	539	905	1,212
2000.....	83,440	278	215	169	207	272	293	276	365	547	880	1,166
1999.....	77,098	311	259	187	182	284	307	298	335	574	806	1,080

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

VI. Diseases of the nervous system (G00-G98)

White, male

2004.....	39,438	173	114	95	137	199	210	216	205	303	531	745
2003.....	38,903	177	94	112	118	216	263	204	202	287	554	747
2002.....	36,751	176	132	96	110	195	207	189	225	337	520	721
2001.....	34,930	140	129	85	119	210	228	169	176	332	496	640
2000.....	33,067	157	121	88	120	187	202	178	213	305	486	670
1999.....	31,244	181	152	102	97	208	213	181	198	340	435	560

White, female

2004.....	62,374	117	105	64	88	85	118	105	143	223	377	595
2003.....	61,364	109	103	72	84	95	122	133	161	228	358	590
2002.....	57,704	114	86	71	83	80	112	112	136	205	400	609
2001.....	53,434	109	91	86	77	88	101	103	170	207	409	572
2000.....	50,373	121	94	81	87	85	91	98	152	242	394	496
1999.....	45,854	130	107	85	85	76	94	117	137	234	371	520

Black, both sexes

2004.....	8,449	69	84	54	67	92	100	91	134	140	202	301
2003.....	7,952	96	71	55	79	74	92	96	117	174	213	314
2002.....	7,373	78	85	42	70	72	107	94	114	160	224	299
2001.....	7,053	94	78	50	61	93	88	103	117	158	212	264
2000.....	6,552	88	84	50	74	78	71	85	105	158	223	285
1999.....	6,030	108	82	67	59	77	95	75	96	155	197	268

Black, male

2004.....	3,191	35	42	20	38	65	55	37	65	66	103	153
2003.....	3,178	42	42	22	51	46	51	52	69	89	110	160
2002.....	2,994	41	51	21	43	40	71	54	63	91	116	160
2001.....	2,858	57	37	20	30	63	51	63	59	89	115	132
2000.....	2,682	56	40	20	45	36	41	45	50	81	127	134
1999.....	2,511	49	40	41	39	46	66	42	51	77	115	130

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

VI. Diseases of the nervous system (G00-G98)

Black, female

2004.....	5,258	34	42	34	29	27	45	54	69	74	99	148
2003.....	4,774	54	29	33	28	28	41	44	48	85	103	154
2002.....	4,379	37	34	21	27	32	36	40	51	69	108	139
2001.....	4,195	37	41	30	31	30	37	40	58	69	97	132
2000.....	3,870	32	44	30	29	42	30	40	55	77	96	151
1999.....	3,519	59	42	26	20	31	29	33	45	78	82	138

American Indian 1/, both sexes

2004.....	313	6	9	2	5	5	4	6	2	11	9	16
2003.....	318	4	4	3	4	6	5	-	7	5	9	8
2002.....	298	3	5	5	2	8	7	1	4	6	6	8
2001.....	264	6	6	1	-	6	6	7	2	7	7	15
2000.....	262	5	9	2	2	3	1	6	4	7	10	9
1999.....	227	3	8	4	2	4	6	2	5	2	7	7

American Indian 1/, male

2004.....	144	4	3	1	3	4	2	4	1	5	4	9
2003.....	132	-	4	2	3	5	3	-	2	3	6	7
2002.....	152	2	3	3	1	6	5	1	3	4	5	6
2001.....	116	4	2	1	-	4	5	4	1	3	5	5
2000.....	108	3	3	2	1	1	1	3	2	3	9	6
1999.....	99	2	4	1	-	1	2	1	5	2	4	2

American Indian 1/, female

2004.....	169	2	6	1	2	1	2	2	1	6	5	7
2003.....	186	4	-	1	1	1	2	-	5	2	3	1
2002.....	146	1	2	2	1	2	2	-	1	2	1	2
2001.....	148	2	4	-	-	2	1	3	1	4	2	10
2000.....	154	2	6	-	1	2	-	3	2	4	1	3
1999.....	128	1	4	3	2	3	4	1	-	-	3	5

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
-------------	-----------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

VI. Diseases of the nervous system (G00-G98)

Asian or Pacific Islander 2/, both sexes

2004.....	1,177	12	10	6	8	10	13	9	11	6	8	15
2003.....	1,103	14	8	9	11	8	14	4	6	9	15	21
2002.....	938	11	14	8	12	7	14	9	9	7	13	17
2001.....	843	11	11	7	6	7	11	6	11	6	13	23
2000.....	760	18	8	6	8	9	8	3	7	9	20	14
1999.....	722	19	12	5	10	10	7	6	11	7	16	14

Asian or Pacific Islander 2/, male

2004.....	540	6	5	6	5	7	11	6	7	4	5	6
2003.....	523	5	4	6	7	3	8	4	6	6	10	10
2002.....	435	9	6	6	7	3	9	6	6	4	8	7
2001.....	406	6	6	5	4	5	7	4	6	3	3	11
2000.....	381	9	5	2	3	6	5	2	2	8	11	7
1999.....	354	6	5	3	7	7	4	4	6	6	12	3

Asian or Pacific Islander 2/, female

2004.....	637	6	5	-	3	3	2	3	4	2	3	9
2003.....	580	9	4	3	4	5	6	-	-	3	5	11
2002.....	503	2	8	2	5	4	5	3	3	3	5	10
2001.....	437	5	5	2	2	2	4	2	5	3	10	12
2000.....	379	9	3	4	5	3	3	1	5	1	9	7
1999.....	368	13	7	2	3	3	3	2	5	1	4	11

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Meningitis (G00,G03)												
All races, both sexes												
2004.....	674	66	25	11	14	8	18	17	20	21	46	57
2003.....	730	77	24	10	7	14	23	18	26	31	36	59
2002.....	700	74	19	10	21	12	14	16	20	23	38	63
2001.....	727	78	26	15	9	19	21	13	28	31	38	55
2000.....	758	91	36	12	18	7	8	16	27	35	37	45
1999.....	850	116	45	14	13	19	22	21	27	33	43	62
All races, male												
2004.....	349	32	15	6	7	3	7	9	13	15	31	38
2003.....	387	38	12	7	3	9	11	10	18	21	19	35
2002.....	389	41	13	5	11	8	7	9	10	17	26	41
2001.....	394	39	16	9	6	12	11	11	14	23	19	30
2000.....	405	48	20	6	10	2	2	7	13	22	24	25
1999.....	442	61	26	9	9	10	16	8	16	18	27	39
All races, female												
2004.....	325	34	10	5	7	5	11	8	7	6	15	19
2003.....	343	39	12	3	4	5	12	8	8	10	17	24
2002.....	311	33	6	5	10	4	7	7	10	6	12	22
2001.....	333	39	10	6	3	7	10	2	14	8	19	25
2000.....	353	43	16	6	8	5	6	9	14	13	13	20
1999.....	408	55	19	5	4	9	6	13	11	15	16	23
White, both sexes												
2004.....	489	46	19	10	9	6	13	9	10	16	33	36
2003.....	524	52	18	8	5	10	15	10	16	17	26	41
2002.....	498	52	13	9	11	10	8	9	11	18	21	39
2001.....	526	47	19	10	8	14	12	8	21	15	25	37
2000.....	536	54	23	11	10	4	6	8	17	24	20	35
1999.....	601	68	33	9	10	10	11	16	18	22	27	39

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Meningitis (G00,G03)												
White, male												
2004.....	259	25	12	6	5	3	4	6	7	11	23	26
2003.....	293	32	9	6	3	6	10	6	11	12	14	27
2002.....	270	29	8	5	6	7	3	5	4	14	12	24
2001.....	275	22	13	5	5	7	6	7	11	10	14	20
2000.....	294	31	14	6	4	1	2	3	7	15	12	21
1999.....	308	46	20	5	7	5	7	4	9	12	15	22
White, female												
2004.....	230	21	7	4	4	3	9	3	3	5	10	10
2003.....	231	20	9	2	2	4	5	4	5	5	12	14
2002.....	228	23	5	4	5	3	5	4	7	4	9	15
2001.....	251	25	6	5	3	7	6	1	10	5	11	17
2000.....	242	23	9	5	6	3	4	5	10	9	8	14
1999.....	293	22	13	4	3	5	4	12	9	10	12	17
Black, both sexes												
2004.....	159	17	5	1	5	2	3	7	8	5	13	20
2003.....	187	23	5	2	1	4	8	8	10	13	9	16
2002.....	178	20	6	1	9	1	5	7	8	5	15	21
2001.....	175	29	5	4	-	4	7	5	6	13	13	18
2000.....	204	35	13	1	8	3	2	8	10	10	17	9
1999.....	222	42	11	5	3	9	8	5	6	10	12	22
Black, male												
2004.....	76	7	2	-	2	-	2	2	5	4	8	11
2003.....	84	6	2	1	-	3	1	4	7	8	4	7
2002.....	107	11	5	-	5	1	4	4	5	3	12	15
2001.....	105	16	2	3	-	4	4	4	2	12	5	10
2000.....	102	16	6	-	6	1	-	4	6	6	12	3
1999.....	121	13	6	4	2	5	6	4	5	6	9	17

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Meningitis (G00,G03)												
Asian or Pacific Islander 2/, both sexes												
2004.....	17	2	1	-	-	-	2	1	1	-	-	-
2003.....	11	1	-	-	1	-	-	-	-	-	1	1
2002.....	16	1	-	-	-	1	-	-	1	-	1	3
2001.....	16	1	-	1	1	-	2	-	1	3	-	-
2000.....	13	1	-	-	-	-	-	-	-	-	-	1
1999.....	17	4	1	-	-	-	1	-	1	1	3	-
Asian or Pacific Islander 2/, male												
2004.....	8	-	1	-	-	-	1	1	-	-	-	-
2003.....	5	-	-	-	-	-	-	-	-	-	1	-
2002.....	7	-	-	-	-	-	-	-	1	-	1	2
2001.....	9	1	-	1	1	-	1	-	1	1	-	-
2000.....	6	-	-	-	-	-	-	-	-	-	-	1
1999.....	6	1	-	-	-	-	1	-	-	-	3	-
Asian or Pacific Islander 2/, female												
2004.....	9	2	-	-	-	-	1	-	1	-	-	-
2003.....	6	1	-	-	1	-	-	-	-	-	-	1
2002.....	9	1	-	-	-	1	-	-	-	-	-	1
2001.....	7	-	-	-	-	-	1	-	-	2	-	-
2000.....	7	1	-	-	-	-	-	-	-	-	-	-
1999.....	11	3	1	-	-	-	-	-	1	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other inflammatory diseases of central nervous system (G04-G09)

All races, both sexes

2004.....	655	11	12	10	16	13	13	17	23	17	35	37
2003.....	706	7	15	15	10	12	17	16	20	29	45	43
2002.....	722	8	14	15	10	17	20	20	29	30	37	52
2001.....	618	9	14	6	9	15	11	22	23	17	32	46
2000.....	606	8	9	8	22	10	12	16	12	24	39	28
1999.....	568	4	19	23	10	13	16	14	17	25	33	32

All races, male

2004.....	363	7	5	5	13	8	5	8	9	7	24	23
2003.....	379	4	7	7	5	6	10	4	10	15	27	29
2002.....	391	3	6	10	4	11	12	10	15	16	25	28
2001.....	309	6	9	2	6	8	5	6	7	8	18	25
2000.....	326	7	7	6	9	3	9	7	1	12	27	13
1999.....	303	1	12	15	6	5	9	7	11	15	22	16

All races, female

2004.....	292	4	7	5	3	5	8	9	14	10	11	14
2003.....	327	3	8	8	5	6	7	12	10	14	18	14
2002.....	331	5	8	5	6	6	8	10	14	14	12	24
2001.....	309	3	5	4	3	7	6	16	16	9	14	21
2000.....	280	1	2	2	13	7	3	9	11	12	12	15
1999.....	265	3	7	8	4	8	7	7	6	10	11	16

White, both sexes

2004.....	532	8	11	5	12	6	11	10	14	10	25	30
2003.....	575	6	11	10	8	7	12	10	13	14	36	30
2002.....	578	6	10	12	7	15	15	14	21	18	26	34
2001.....	486	7	9	4	6	9	10	12	14	12	24	33
2000.....	489	4	7	4	16	8	9	12	7	20	29	19
1999.....	452	4	11	17	8	11	9	10	10	17	22	26

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other inflammatory diseases of central nervous system (G04-G09)

White, male

2004.....	303	4	4	3	10	4	5	7	6	3	18	20
2003.....	311	4	5	4	3	4	8	3	5	8	22	22
2002.....	314	2	5	7	3	9	9	8	11	10	17	20
2001.....	250	5	7	1	4	6	4	3	4	7	13	19
2000.....	273	3	5	3	7	3	7	6	1	9	20	7
1999.....	240	1	7	11	5	4	5	6	6	11	13	12

White, female

2004.....	229	4	7	2	2	2	6	3	8	7	7	10
2003.....	264	2	6	6	5	3	4	7	8	6	14	8
2002.....	264	4	5	5	4	6	6	6	10	8	9	14
2001.....	236	2	2	3	2	3	6	9	10	5	11	14
2000.....	216	1	2	1	9	5	2	6	6	11	9	12
1999.....	212	3	4	6	3	7	4	4	4	6	9	14

Black, both sexes

2004.....	111	2	1	5	3	7	2	5	8	7	9	7
2003.....	110	1	4	4	2	3	5	6	7	15	6	11
2002.....	121	2	4	1	2	2	4	5	6	12	10	16
2001.....	116	2	4	2	3	5	1	9	8	5	8	9
2000.....	101	4	2	4	5	2	3	3	5	4	10	9
1999.....	104	-	6	5	2	1	7	4	6	8	11	6

Black, male

2004.....	51	2	1	2	2	4	-	-	2	4	5	3
2003.....	56	-	2	2	2	1	2	1	5	7	3	5
2002.....	66	1	1	1	1	2	3	2	2	6	7	8
2001.....	53	1	2	1	2	2	1	2	3	1	5	5
2000.....	48	4	2	3	2	-	2	-	-	3	7	6
1999.....	56	-	3	4	1	-	4	1	4	4	9	4

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Parkinson's disease (G20-G21)												
All races, both sexes												
2004.....	17,989	-	1	-	-	1	-	2	-	2	6	16
2003.....	17,997	1	-	1	-	-	-	-	2	3	8	18
2002.....	16,959	-	-	-	-	-	-	-	-	1	5	19
2001.....	16,544	-	-	-	-	-	-	1	1	3	12	13
2000.....	15,682	-	1	-	-	-	-	2	2	4	3	16
1999.....	14,593	1	-	-	-	-	2	1	1	1	4	8
All races, male												
2004.....	10,366	-	-	-	-	1	-	2	-	2	4	12
2003.....	10,180	1	-	-	-	-	-	-	-	2	4	10
2002.....	9,593	-	-	-	-	-	-	-	-	-	3	14
2001.....	9,405	-	-	-	-	-	-	1	-	1	10	8
2000.....	8,804	-	-	-	-	-	-	1	2	3	2	10
1999.....	8,262	-	-	-	-	-	2	1	1	1	2	5
All races, female												
2004.....	7,623	-	1	-	-	-	-	-	-	-	2	4
2003.....	7,817	-	-	1	-	-	-	-	2	1	4	8
2002.....	7,366	-	-	-	-	-	-	-	-	1	2	5
2001.....	7,139	-	-	-	-	-	-	-	1	2	2	5
2000.....	6,878	-	1	-	-	-	-	1	-	1	1	6
1999.....	6,331	1	-	-	-	-	-	-	-	-	2	3
White, both sexes												
2004.....	17,041	-	1	-	-	-	-	2	-	1	6	15
2003.....	17,045	1	-	-	-	-	-	-	2	2	8	18
2002.....	16,135	-	-	-	-	-	-	-	-	-	3	15
2001.....	15,757	-	-	-	-	-	-	-	1	2	9	10
2000.....	14,942	-	-	-	-	-	-	1	2	3	1	12
1999.....	13,886	1	-	-	-	-	2	1	-	1	4	8

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Alzheimer's disease (G30)												
All races, both sexes												
2004.....	65,965	-	-	-	-	-	1	-	1	1	6	36
2003.....	63,457	-	-	-	-	-	-	-	-	2	9	21
2002.....	58,866	-	-	-	1	-	-	-	2	3	9	12
2001.....	53,852	-	-	-	-	-	-	-	2	1	2	17
2000.....	49,558	-	-	-	-	-	-	1	1	2	4	17
1999.....	44,536	-	-	-	-	-	1	-	-	1	6	20
All races, male												
2004.....	18,974	-	-	-	-	-	-	-	1	1	3	8
2003.....	18,335	-	-	-	-	-	-	-	-	1	3	9
2002.....	16,989	-	-	-	-	-	-	-	2	3	5	5
2001.....	15,762	-	-	-	-	-	-	-	2	1	2	11
2000.....	14,438	-	-	-	-	-	-	1	-	1	3	10
1999.....	13,391	-	-	-	-	-	1	-	-	-	3	10
All races, female												
2004.....	46,991	-	-	-	-	-	1	-	-	-	3	28
2003.....	45,122	-	-	-	-	-	-	-	-	1	6	12
2002.....	41,877	-	-	-	1	-	-	-	-	-	4	7
2001.....	38,090	-	-	-	-	-	-	-	-	-	-	6
2000.....	35,120	-	-	-	-	-	-	-	1	1	1	7
1999.....	31,145	-	-	-	-	-	-	-	-	1	3	10
White, both sexes												
2004.....	61,087	-	-	-	-	-	1	-	1	1	5	36
2003.....	59,184	-	-	-	-	-	-	-	-	2	9	19
2002.....	55,058	-	-	-	1	-	-	-	1	1	6	12
2001.....	50,348	-	-	-	-	-	-	-	2	1	2	13
2000.....	46,460	-	-	-	-	-	-	1	1	2	4	14
1999.....	41,877	-	-	-	-	-	1	-	-	1	4	18

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Multiple sclerosis (G35)

All races, both sexes

2004.....	3,324	1	-	1	-	2	6	9	34	85	164	315
2003.....	3,215	-	-	1	-	3	3	13	39	89	183	335
2002.....	3,124	-	-	-	-	2	5	9	30	92	185	359
2001.....	3,053	-	1	-	1	1	5	14	38	96	207	331
2000.....	2,844	-	-	-	-	2	2	20	38	88	210	289
1999.....	2,651	-	-	-	1	5	9	9	37	97	207	284

All races, male

2004.....	1,105	1	-	-	-	1	1	2	12	35	54	112
2003.....	1,126	-	-	-	-	2	3	5	19	28	74	122
2002.....	1,064	-	-	-	-	-	-	3	12	39	54	113
2001.....	1,043	-	-	-	-	-	2	6	13	36	65	118
2000.....	994	-	-	-	-	1	1	6	10	33	83	107
1999.....	886	-	-	-	-	1	3	1	12	35	74	80

All races, female

2004.....	2,219	-	-	1	-	1	5	7	22	50	110	203
2003.....	2,089	-	-	1	-	1	-	8	20	61	109	213
2002.....	2,060	-	-	-	-	2	5	6	18	53	131	246
2001.....	2,010	-	1	-	1	1	3	8	25	60	142	213
2000.....	1,850	-	-	-	-	1	1	14	28	55	127	182
1999.....	1,765	-	-	-	1	4	6	8	25	62	133	204

White, both sexes

2004.....	2,972	1	-	1	-	2	1	7	19	59	134	265
2003.....	2,849	-	-	1	-	1	2	5	22	62	150	272
2002.....	2,770	-	-	-	-	-	4	6	17	65	148	301
2001.....	2,739	-	1	-	-	-	3	12	29	65	171	279
2000.....	2,519	-	-	-	-	-	2	14	25	59	166	236
1999.....	2,391	-	-	-	1	3	4	5	26	75	171	249

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Multiple sclerosis (G35)												
White, male												
2004.....	999	1	-	-	-	1	-	2	6	28	44	96
2003.....	1,007	-	-	-	-	1	2	2	8	23	64	103
2002.....	971	-	-	-	-	-	-	2	7	28	49	96
2001.....	955	-	-	-	-	-	-	6	11	27	56	101
2000.....	895	-	-	-	-	-	1	4	8	21	66	96
1999.....	809	-	-	-	-	1	1	1	9	31	62	70
White, female												
2004.....	1,973	-	-	1	-	1	1	5	13	31	90	169
2003.....	1,842	-	-	1	-	-	-	3	14	39	86	169
2002.....	1,799	-	-	-	-	-	4	4	10	37	99	205
2001.....	1,784	-	1	-	-	-	3	6	18	38	115	178
2000.....	1,624	-	-	-	-	-	1	10	17	38	100	140
1999.....	1,582	-	-	-	1	2	3	4	17	44	109	179
Black, both sexes												
2004.....	335	-	-	-	-	-	4	2	14	24	28	47
2003.....	352	-	-	-	-	2	1	8	16	26	31	59
2002.....	340	-	-	-	-	2	1	3	13	25	37	56
2001.....	300	-	-	-	1	1	2	1	9	30	34	49
2000.....	307	-	-	-	-	2	-	5	11	28	41	50
1999.....	247	-	-	-	-	2	5	4	10	22	35	32
Black, male												
2004.....	101	-	-	-	-	-	-	-	6	6	10	15
2003.....	116	-	-	-	-	1	1	3	10	5	9	18
2002.....	88	-	-	-	-	-	-	1	5	10	5	16
2001.....	84	-	-	-	-	-	2	-	2	9	9	16
2000.....	95	-	-	-	-	1	-	1	2	12	15	11
1999.....	76	-	-	-	-	-	2	-	2	4	12	10

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Multiple sclerosis (G35)

Asian or Pacific Islander 2/, both sexes

2004.....	10	-	-	-	-	-	1	-	1	1	-	2
2003.....	10	-	-	-	-	-	-	-	1	-	1	3
2002.....	9	-	-	-	-	-	-	-	2	-	-	2
2001.....	9	-	-	-	-	-	-	-	1	2	2	2
2000.....	7	-	-	-	-	-	-	1	-	1	1	1
1999.....	9	-	-	-	-	-	-	-	-	1	2	2

Asian or Pacific Islander 2/, male

2004.....	4	-	-	-	-	-	1	-	-	1	-	1
2003.....	1	-	-	-	-	-	-	-	1	-	-	-
2002.....	3	-	-	-	-	-	-	-	-	1	-	1
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-

Asian or Pacific Islander 2/, female

2004.....	6	-	-	-	-	-	-	1	-	-	-	1
2003.....	9	-	-	-	-	-	-	-	-	-	1	3
2002.....	6	-	-	-	-	-	-	-	1	-	-	1
2001.....	9	-	-	-	-	-	-	-	1	2	2	2
2000.....	7	-	-	-	-	-	-	1	-	1	1	1
1999.....	9	-	-	-	-	-	-	-	-	1	2	2

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Epilepsy (G40-G41)

All races, both sexes

2004.....	1,347	13	28	12	25	36	65	86	104	95	128	124
2003.....	1,416	13	25	15	33	36	76	79	87	111	125	145
2002.....	1,379	11	23	21	20	32	71	73	89	109	141	136
2001.....	1,292	8	18	24	16	42	53	76	84	115	143	104
2000.....	1,323	11	25	14	22	38	54	62	86	106	136	116
1999.....	1,338	14	31	21	24	43	59	86	86	120	125	108

All races, male

2004.....	748	10	18	5	14	21	37	52	66	58	86	79
2003.....	756	6	11	10	21	24	44	49	50	63	76	86
2002.....	746	5	13	14	9	14	39	46	52	66	91	86
2001.....	700	3	8	12	11	24	40	47	46	77	86	63
2000.....	701	4	16	8	13	14	36	41	50	56	86	76
1999.....	730	4	21	9	12	31	34	59	51	81	80	69

All races, female

2004.....	599	3	10	7	11	15	28	34	38	37	42	45
2003.....	660	7	14	5	12	12	32	30	37	48	49	59
2002.....	633	6	10	7	11	18	32	27	37	43	50	50
2001.....	592	5	10	12	5	18	13	29	38	38	57	41
2000.....	622	7	9	6	9	24	18	21	36	50	50	40
1999.....	608	10	10	12	12	12	25	27	35	39	45	39

White, both sexes

2004.....	1,005	10	23	8	21	27	44	64	71	71	93	94
2003.....	1,052	11	21	13	23	29	59	62	62	82	94	103
2002.....	1,034	11	15	18	16	25	49	55	68	83	108	97
2001.....	968	6	15	20	14	29	38	52	64	80	105	84
2000.....	990	7	21	11	17	23	39	52	63	79	107	83
1999.....	979	8	27	16	17	29	45	59	64	88	88	81

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Epilepsy (G40-G41)

White, male

2004.....	557	8	15	4	11	14	21	41	48	47	63	61
2003.....	547	6	9	9	14	19	34	40	35	47	57	61
2002.....	547	5	8	11	7	10	26	32	40	51	68	60
2001.....	518	2	7	10	10	15	29	32	33	52	63	51
2000.....	524	3	13	7	11	10	26	36	37	40	65	59
1999.....	529	3	20	7	9	20	25	40	41	57	54	51

White, female

2004.....	448	2	8	4	10	13	23	23	23	24	30	33
2003.....	505	5	12	4	9	10	25	22	27	35	37	42
2002.....	487	6	7	7	9	15	23	23	28	32	40	37
2001.....	450	4	8	10	4	14	9	20	31	28	42	33
2000.....	466	4	8	4	6	13	13	16	26	39	42	24
1999.....	450	5	7	9	8	9	20	19	23	31	34	30

Black, both sexes

2004.....	299	2	4	4	3	8	19	18	28	19	32	25
2003.....	327	1	-	2	10	6	15	16	21	25	29	39
2002.....	315	-	7	2	4	7	20	18	17	21	30	36
2001.....	290	2	2	3	2	10	14	20	19	33	34	17
2000.....	302	2	2	3	5	12	15	8	22	20	25	31
1999.....	332	4	3	4	6	11	14	26	18	30	32	26

Black, male

2004.....	164	1	2	1	2	6	14	9	14	8	21	16
2003.....	189	-	-	1	7	4	10	8	12	13	18	24
2002.....	180	-	4	2	2	4	11	14	11	12	20	24
2001.....	164	1	-	1	1	6	10	13	12	24	22	11
2000.....	159	1	2	1	2	4	10	4	13	10	17	16
1999.....	188	1	1	2	3	9	9	19	8	22	22	18

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Epilepsy (G40-G41)												
Black, female												
2004.....	135	1	2	3	1	2	5	9	14	11	11	9
2003.....	138	1	-	1	3	2	5	8	9	12	11	15
2002.....	135	-	3	-	2	3	9	4	6	9	10	12
2001.....	126	1	2	2	1	4	4	7	7	9	12	6
2000.....	143	1	-	2	3	8	5	4	9	10	8	15
1999.....	144	3	2	2	3	2	5	7	10	8	10	8
American Indian 1/, both sexes												
2004.....	18	-	1	-	1	-	-	-	-	4	2	3
2003.....	14	1	1	-	-	1	1	-	1	1	-	1
2002.....	10	-	-	-	-	-	1	-	2	2	-	2
2001.....	15	-	1	-	-	-	-	2	-	2	2	1
2000.....	11	1	1	-	-	1	-	-	-	3	1	1
1999.....	10	-	1	-	1	1	-	-	1	1	1	-
American Indian 1/, male												
2004.....	12	-	1	-	1	-	-	-	-	2	1	2
2003.....	8	-	1	-	-	1	-	-	-	1	-	-
2002.....	7	-	-	-	-	-	1	-	1	2	-	2
2001.....	7	-	1	-	-	-	-	1	-	1	1	-
2000.....	5	-	-	-	-	-	-	-	-	2	1	1
1999.....	5	-	-	-	-	-	-	-	1	1	1	-
American Indian 1/, female												
2004.....	6	-	-	-	-	-	-	-	-	2	1	1
2003.....	6	1	-	-	-	-	1	-	1	-	-	1
2002.....	3	-	-	-	-	-	-	-	1	-	-	-
2001.....	8	-	-	-	-	-	-	1	-	1	1	1
2000.....	6	1	1	-	-	1	-	-	-	1	-	-
1999.....	5	-	1	-	1	1	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Epilepsy (G40-G41)												
Asian or Pacific Islander 2/, both sexes												
2004.....	25	1	-	-	-	1	2	4	5	1	1	2
2003.....	23	-	3	-	-	-	1	1	3	3	2	2
2002.....	20	-	1	1	-	-	1	-	2	3	3	1
2001.....	19	-	-	1	-	3	1	2	1	-	2	2
2000.....	20	1	1	-	-	2	-	2	1	4	3	1
1999.....	17	2	-	1	-	2	-	1	3	1	4	1
Asian or Pacific Islander 2/, male												
2004.....	15	1	-	-	-	1	2	2	4	1	1	-
2003.....	12	-	1	-	-	-	-	1	3	2	1	1
2002.....	12	-	1	1	-	-	1	-	-	1	3	-
2001.....	11	-	-	1	-	3	1	1	1	-	-	1
2000.....	13	-	1	-	-	-	-	1	-	4	3	-
1999.....	8	-	-	-	-	2	-	-	1	1	3	-
Asian or Pacific Islander 2/, female												
2004.....	10	-	-	-	-	-	-	2	1	-	-	2
2003.....	11	-	2	-	-	-	1	-	-	1	1	1
2002.....	8	-	-	-	-	-	-	-	2	2	-	1
2001.....	8	-	-	-	-	-	-	1	-	-	2	1
2000.....	7	1	-	-	-	2	-	1	1	-	-	1
1999.....	9	2	-	1	-	-	-	1	2	-	1	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Transient cerebral ischemic attacks and related syndromes (G45)												
All races, both sexes												
2004.....	175	-	-	-	-	-	-	-	-	-	1	1
2003.....	267	-	-	-	-	-	-	1	-	-	1	4
2002.....	239	-	-	-	-	-	-	-	1	-	1	1
2001.....	271	-	-	-	-	-	-	1	-	-	1	1
2000.....	289	-	-	-	-	-	-	-	-	-	2	4
1999.....	317	-	-	-	-	-	-	1	-	2	1	2
All races, male												
2004.....	50	-	-	-	-	-	-	-	-	-	-	1
2003.....	75	-	-	-	-	-	-	-	-	-	1	2
2002.....	78	-	-	-	-	-	-	-	1	-	-	-
2001.....	106	-	-	-	-	-	-	1	-	-	-	1
2000.....	89	-	-	-	-	-	-	-	-	-	-	3
1999.....	107	-	-	-	-	-	-	-	-	-	-	1
All races, female												
2004.....	125	-	-	-	-	-	-	-	-	-	1	-
2003.....	192	-	-	-	-	-	-	1	-	-	-	2
2002.....	161	-	-	-	-	-	-	-	-	-	1	1
2001.....	165	-	-	-	-	-	-	-	-	-	1	-
2000.....	200	-	-	-	-	-	-	-	-	-	2	1
1999.....	210	-	-	-	-	-	-	1	-	2	1	1
White, both sexes												
2004.....	161	-	-	-	-	-	-	-	-	-	1	1
2003.....	250	-	-	-	-	-	-	1	-	-	1	3
2002.....	221	-	-	-	-	-	-	-	1	-	1	1
2001.....	250	-	-	-	-	-	-	1	-	-	1	-
2000.....	269	-	-	-	-	-	-	-	-	-	2	3
1999.....	300	-	-	-	-	-	-	1	-	-	1	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Infantile cerebral palsy (G80)												
All races, both sexes												
2004.....	1,417	12	74	93	138	163	153	113	70	66	60	71
2003.....	1,334	11	68	113	127	143	138	108	74	62	48	75
2002.....	1,200	9	93	109	104	132	133	90	50	53	45	54
2001.....	1,155	19	71	87	110	141	121	70	68	50	52	50
2000.....	1,091	10	59	105	122	125	107	67	64	54	49	50
1999.....	1,008	13	83	92	93	107	84	79	59	60	47	49
All races, male												
2004.....	801	5	33	52	80	109	91	70	36	31	28	35
2003.....	761	6	41	67	74	87	79	57	40	31	31	40
2002.....	675	6	57	58	56	82	86	50	28	33	25	19
2001.....	657	12	43	41	59	85	75	38	41	23	35	25
2000.....	603	8	26	48	77	77	65	36	38	27	25	28
1999.....	572	5	50	55	50	65	54	44	32	28	24	26
All races, female												
2004.....	616	7	41	41	58	54	62	43	34	35	32	36
2003.....	573	5	27	46	53	56	59	51	34	31	17	35
2002.....	525	3	36	51	48	50	47	40	22	20	20	35
2001.....	498	7	28	46	51	56	46	32	27	27	17	25
2000.....	488	2	33	57	45	48	42	31	26	27	24	22
1999.....	436	8	33	37	43	42	30	35	27	32	23	23
White, both sexes												
2004.....	1,097	8	41	62	95	111	109	91	52	53	47	64
2003.....	1,035	5	42	81	84	102	101	86	55	52	37	58
2002.....	911	5	64	75	67	95	85	62	31	37	38	51
2001.....	910	7	50	61	77	99	92	47	52	45	45	42
2000.....	854	9	30	73	85	94	80	50	49	40	45	44
1999.....	799	6	54	62	68	75	60	63	50	55	37	44

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Infantile cerebral palsy (G80)												
White, male												
2004.....	625	5	19	39	56	72	66	60	26	24	21	30
2003.....	591	3	25	53	48	65	58	42	30	25	23	31
2002.....	513	4	40	42	33	64	53	34	17	23	23	19
2001.....	521	5	33	28	43	59	59	27	29	21	30	23
2000.....	482	7	15	37	54	59	50	28	29	20	23	24
1999.....	446	2	35	34	34	47	38	33	28	27	15	25
White, female												
2004.....	472	3	22	23	39	39	43	31	26	29	26	34
2003.....	444	2	17	28	36	37	43	44	25	27	14	27
2002.....	398	1	24	33	34	31	32	28	14	14	15	32
2001.....	389	2	17	33	34	40	33	20	23	24	15	19
2000.....	372	2	15	36	31	35	30	22	20	20	22	20
1999.....	353	4	19	28	34	28	22	30	22	28	22	19
Black, both sexes												
2004.....	288	3	29	29	36	45	42	21	17	11	10	7
2003.....	268	6	23	28	35	37	33	20	17	10	10	15
2002.....	253	3	25	28	31	33	39	24	19	16	7	3
2001.....	216	9	18	22	30	39	23	23	15	5	5	5
2000.....	217	1	27	29	35	30	22	14	13	14	2	6
1999.....	190	7	28	25	21	29	22	12	9	5	10	5
Black, male												
2004.....	159	-	12	12	21	33	23	9	9	6	6	5
2003.....	150	3	14	11	21	19	18	13	10	6	7	8
2002.....	140	2	15	12	20	16	27	12	11	10	2	-
2001.....	120	5	9	10	15	25	12	11	11	2	4	1
2000.....	110	1	10	10	22	17	12	7	7	7	1	4
1999.....	115	3	15	18	13	17	15	8	4	1	9	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Infantile cerebral palsy (G80)												
Black, female												
2004.....	129	3	17	17	15	12	19	12	8	5	4	2
2003.....	118	3	9	17	14	18	15	7	7	4	3	7
2002.....	113	1	10	16	11	17	12	12	8	6	5	3
2001.....	96	4	9	12	15	14	11	12	4	3	1	4
2000.....	107	-	17	19	13	13	10	7	6	7	1	2
1999.....	75	4	13	7	8	12	7	4	5	4	1	4
American Indian 1/, both sexes												
2004.....	13	1	2	1	3	1	1	1	-	1	-	-
2003.....	11	-	1	2	3	2	1	-	2	-	-	-
2002.....	12	-	-	3	-	3	4	1	-	-	-	-
2001.....	11	-	1	1	-	2	3	-	1	-	1	1
2000.....	7	-	2	-	-	1	1	3	-	-	-	-
1999.....	5	-	-	2	-	1	1	1	-	-	-	-
American Indian 1/, male												
2004.....	5	-	1	-	1	-	1	1	-	-	-	-
2003.....	7	-	1	2	2	2	-	-	-	-	-	-
2002.....	9	-	-	2	-	2	4	1	-	-	-	-
2001.....	9	-	-	1	-	1	3	-	1	-	1	1
2000.....	4	-	1	-	-	1	1	1	-	-	-	-
1999.....	1	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, female												
2004.....	8	1	1	1	2	1	-	-	-	1	-	-
2003.....	4	-	-	-	1	-	1	-	2	-	-	-
2002.....	3	-	-	1	-	1	-	-	-	-	-	-
2001.....	2	-	1	-	-	1	-	-	-	-	-	-
2000.....	3	-	1	-	-	-	-	2	-	-	-	-
1999.....	4	-	-	1	-	1	1	1	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Infantile cerebral palsy (G80)												
Asian or Pacific Islander 2/, both sexes												
2004.....	19	-	2	1	4	6	1	-	1	1	3	-
2003.....	20	-	2	2	5	2	3	2	-	-	1	2
2002.....	24	1	4	3	6	1	5	3	-	-	-	-
2001.....	18	3	2	3	3	1	3	-	-	-	1	2
2000.....	13	-	-	3	2	-	4	-	2	-	2	-
1999.....	14	-	1	3	4	2	1	3	-	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	12	-	1	1	2	4	1	-	1	1	1	-
2003.....	13	-	1	1	3	1	3	2	-	-	1	1
2002.....	13	-	2	2	3	-	2	3	-	-	-	-
2001.....	7	2	1	2	1	-	1	-	-	-	-	-
2000.....	7	-	-	1	1	-	2	-	2	-	1	-
1999.....	10	-	-	2	3	1	1	3	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	7	-	1	-	2	2	-	-	-	-	2	-
2003.....	7	-	1	1	2	1	-	-	-	-	-	1
2002.....	11	1	2	1	3	1	3	-	-	-	-	-
2001.....	11	1	1	1	2	1	2	-	-	-	1	2
2000.....	6	-	-	2	1	-	2	-	-	-	1	-
1999.....	4	-	1	1	1	1	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

All other diseases of nervous system (G10-G12,G23-G25,G31,G36-G37,G43-G44,G47-G72,G81-G98)

All races, both sexes

2004.....	20,205	274	182	94	112	168	189	183	243	396	681	1,015
2003.....	20,518	291	148	96	119	191	239	202	245	376	694	980
2002.....	19,875	280	173	67	121	167	204	197	267	404	702	958
2001.....	19,012	246	185	97	118	186	223	191	232	397	650	897
2000.....	18,863	269	186	88	107	180	190	186	251	408	653	909
1999.....	18,216	293	183	113	112	188	222	170	220	399	560	804

All races, male

2004.....	10,557	163	93	54	69	132	137	120	141	229	413	605
2003.....	10,737	169	73	51	76	142	178	135	142	224	445	591
2002.....	10,407	173	103	39	81	129	148	132	177	262	420	588
2001.....	9,934	147	98	47	71	153	158	130	119	258	384	507
2000.....	9,878	158	100	44	60	133	136	129	153	243	383	545
1999.....	9,515	167	92	59	66	150	166	108	137	247	334	449

All races, female

2004.....	9,648	111	89	40	43	36	52	63	102	167	268	410
2003.....	9,781	122	75	45	43	49	61	67	103	152	249	389
2002.....	9,468	107	70	28	40	38	56	65	90	142	282	370
2001.....	9,078	99	87	50	47	33	65	61	113	139	266	390
2000.....	8,985	111	86	44	47	47	54	57	98	165	270	364
1999.....	8,701	126	91	54	46	38	56	62	83	152	226	355

White, both sexes

2004.....	17,428	217	124	73	88	132	149	138	181	315	564	799
2003.....	17,753	211	105	71	82	162	196	163	193	284	551	793
2002.....	17,250	216	116	53	91	130	158	155	211	320	569	780
2001.....	16,380	182	126	76	91	147	174	140	163	319	523	714
2000.....	16,381	204	134	70	79	143	157	138	201	320	506	720
1999.....	15,813	224	134	83	78	156	175	143	167	315	452	614

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

All other diseases of nervous system (G10-G12,G23-G25,G31,G36-G37,G43-G44,G47-G72,G81-G98)

White, male

2004.....	9,213	130	64	43	55	105	114	98	111	188	356	491
2003.....	9,366	131	46	40	50	121	151	111	113	169	366	483
2002.....	9,062	136	71	31	61	105	116	108	144	210	345	487
2001.....	8,617	106	69	41	57	123	130	93	86	214	310	410
2000.....	8,627	113	74	35	44	114	116	99	129	197	296	445
1999.....	8,348	129	70	45	42	131	134	96	105	201	272	364

White, female

2004.....	8,215	87	60	30	33	27	35	40	70	127	208	308
2003.....	8,387	80	59	31	32	41	45	52	80	115	185	310
2002.....	8,188	80	45	22	30	25	42	47	67	110	224	293
2001.....	7,763	76	57	35	34	24	44	47	77	105	213	304
2000.....	7,754	91	60	35	35	29	41	39	72	123	210	275
1999.....	7,465	95	64	38	36	25	41	47	62	114	180	250

Black, both sexes

2004.....	2,388	45	45	15	20	30	30	38	59	73	109	195
2003.....	2,347	65	39	18	31	22	30	38	46	84	128	171
2002.....	2,252	53	43	10	24	27	38	37	50	78	123	164
2001.....	2,270	52	49	19	25	34	41	44	60	71	115	159
2000.....	2,141	46	40	13	21	29	29	46	44	81	126	176
1999.....	2,070	55	34	28	27	25	39	24	47	78	96	174

Black, male

2004.....	1,146	25	25	5	11	22	16	17	29	37	52	103
2003.....	1,142	33	24	7	21	18	19	23	25	50	69	97
2002.....	1,134	27	26	6	15	17	26	21	28	48	69	94
2001.....	1,120	34	24	5	12	26	22	32	29	40	68	86
2000.....	1,078	34	20	6	13	13	17	29	22	42	74	91
1999.....	986	32	15	13	20	15	30	10	28	40	54	80

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

All other diseases of nervous system (G10-G12,G23-G25,G31,G36-G37,G43-G44,G47-G72,G81-G98)

Black, female

2004.....	1,242	20	20	10	9	8	14	21	30	36	57	92
2003.....	1,205	32	15	11	10	4	11	15	21	34	59	74
2002.....	1,118	26	17	4	9	10	12	16	22	30	54	70
2001.....	1,150	18	25	14	13	8	19	12	31	31	47	73
2000.....	1,063	12	20	7	8	16	12	17	22	39	52	85
1999.....	1,084	23	19	15	7	10	9	14	19	38	42	94

American Indian 1/, both sexes

2004.....	103	3	6	1	-	3	3	4	1	5	5	10
2003.....	92	2	1	1	1	3	3	-	4	2	7	5
2002.....	90	2	5	1	1	5	1	-	1	4	3	6
2001.....	89	5	1	-	-	2	3	3	1	5	4	9
2000.....	83	3	5	2	2	1	-	2	3	2	7	6
1999.....	82	1	6	1	1	2	3	1	1	1	5	5

American Indian 1/, male

2004.....	56	3	1	1	-	3	1	2	-	3	3	6
2003.....	53	-	1	-	1	2	3	-	2	1	4	5
2002.....	52	1	3	-	1	4	-	-	1	2	2	4
2001.....	43	4	-	-	-	2	2	2	-	2	3	2
2000.....	39	2	2	2	1	-	-	1	2	-	6	5
1999.....	44	1	3	-	-	1	-	1	1	1	3	2

American Indian 1/, female

2004.....	47	-	5	-	-	-	2	2	1	2	2	4
2003.....	39	2	-	1	-	1	-	-	2	1	3	-
2002.....	38	1	2	1	-	1	1	-	-	2	1	2
2001.....	46	1	1	-	-	-	1	1	1	3	1	7
2000.....	44	1	3	-	1	1	-	1	1	2	1	1
1999.....	38	-	3	1	1	1	3	-	-	-	2	3

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

All other diseases of nervous system (G10-G12,G23-G25,G31,G36-G37,G43-G44,G47-G72,G81-G98)

Asian or Pacific Islander 2/, both sexes

2004.....	286	9	7	5	4	3	7	3	2	3	3	11
2003.....	326	13	3	6	5	4	10	1	2	6	8	11
2002.....	283	9	9	3	5	5	7	5	5	2	7	8
2001.....	273	7	9	2	2	3	5	4	8	2	8	15
2000.....	258	16	7	3	5	7	4	-	3	5	14	7
1999.....	251	13	9	1	6	5	5	2	5	5	7	11

Asian or Pacific Islander 2/, male

2004.....	142	5	3	5	3	2	6	3	1	1	2	5
2003.....	176	5	2	4	4	1	5	1	2	4	6	6
2002.....	159	9	3	2	4	3	6	3	4	2	4	3
2001.....	154	3	5	1	2	2	4	3	4	2	3	9
2000.....	134	9	4	1	2	6	3	-	-	4	7	4
1999.....	137	5	4	1	4	3	2	1	3	5	5	3

Asian or Pacific Islander 2/, female

2004.....	144	4	4	-	1	1	1	-	1	2	1	6
2003.....	150	8	1	2	1	3	5	-	-	2	2	5
2002.....	124	-	6	1	1	2	1	2	1	-	3	5
2001.....	119	4	4	1	-	1	1	1	4	-	5	6
2000.....	124	7	3	2	3	1	1	-	3	1	7	3
1999.....	114	8	5	-	2	2	3	1	2	-	2	8

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

VII. Diseases of the eye and adnexa (H00-H57)

All races, both sexes

2004.....	43	2	3	1	-	-	-	-	-	-	1	1
2003.....	41	1	-	1	-	1	-	1	2	-	-	2
2002.....	40	2	1	1	1	1	-	-	-	-	-	3
2001.....	37	2	-	-	1	1	1	-	-	1	-	1
2000.....	59	-	1	1	-	2	3	1	-	-	-	2
1999.....	44	-	-	-	-	-	-	-	-	1	3	2

All races, male

2004.....	18	2	2	1	-	-	-	-	-	-	1	1
2003.....	18	1	-	1	-	1	-	1	-	-	-	2
2002.....	15	1	-	1	1	-	-	-	-	-	-	1
2001.....	11	2	-	-	-	-	-	-	-	-	-	1
2000.....	29	-	1	-	-	1	2	-	-	-	-	1
1999.....	17	-	-	-	-	-	-	-	-	-	2	-

All races, female

2004.....	25	-	1	-	-	-	-	-	-	-	-	-
2003.....	23	-	-	-	-	-	-	-	2	-	-	-
2002.....	25	1	1	-	-	1	-	-	-	-	-	2
2001.....	26	-	-	-	1	1	1	-	-	1	-	-
2000.....	30	-	-	1	-	1	1	1	-	-	-	1
1999.....	27	-	-	-	-	-	-	-	-	1	1	2

White, both sexes

2004.....	40	2	3	1	-	-	-	-	-	-	1	1
2003.....	40	1	-	1	-	1	-	1	2	-	-	2
2002.....	37	2	1	1	1	1	-	-	-	-	-	3
2001.....	32	2	-	-	1	1	1	-	-	-	-	1
2000.....	44	-	1	1	-	-	2	-	-	-	-	2
1999.....	33	-	-	-	-	-	-	-	-	1	2	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

VIII. Diseases of the ear and mastoid process (H60-H93)

All races, both sexes

2004.....	61	4	5	-	-	-	-	-	-	1	6	3
2003.....	72	8	1	-	1	-	1	-	2	-	5	9
2002.....	69	6	5	-	4	-	-	2	1	3	2	7
2001.....	75	3	5	1	1	-	-	1	2	6	4	2
2000.....	67	6	5	1	-	1	1	-	2	6	2	7
1999.....	74	4	2	1	1	3	2	2	8	4	4	3

All races, male

2004.....	28	2	2	-	-	-	-	-	-	1	4	2
2003.....	32	6	1	-	1	-	1	-	2	-	2	4
2002.....	41	6	1	-	2	-	-	1	1	1	1	6
2001.....	33	2	2	-	-	-	-	-	1	4	4	2
2000.....	34	2	2	-	-	1	1	-	2	3	2	3
1999.....	40	4	1	-	-	2	1	2	2	3	3	2

All races, female

2004.....	33	2	3	-	-	-	-	-	-	-	2	1
2003.....	40	2	-	-	-	-	-	-	-	-	3	5
2002.....	28	-	4	-	2	-	-	1	-	2	1	1
2001.....	42	1	3	1	1	-	-	1	1	2	-	-
2000.....	33	4	3	1	-	-	-	-	-	3	-	4
1999.....	34	-	1	1	1	1	1	-	6	1	1	1

White, both sexes

2004.....	45	1	4	-	-	-	-	-	-	-	5	1
2003.....	55	6	-	-	1	-	1	-	2	-	3	5
2002.....	57	2	4	-	4	-	-	2	1	2	2	6
2001.....	65	2	5	1	1	-	-	1	1	4	3	2
2000.....	55	3	5	1	-	1	1	-	1	5	1	6
1999.....	55	3	2	1	1	1	-	2	7	3	2	2

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VIII. Diseases of the ear and mastoid process (H60-H93)												
White, male												
2004.....	20	1	2	-	-	-	-	-	-	-	3	1
2003.....	25	5	-	-	1	-	1	-	2	-	2	1
2002.....	35	2	1	-	2	-	-	1	1	1	1	5
2001.....	28	1	2	-	-	-	-	-	1	3	3	2
2000.....	27	1	2	-	-	1	1	-	1	3	1	3
1999.....	34	3	1	-	-	1	-	2	2	3	2	1
White, female												
2004.....	25	-	2	-	-	-	-	-	-	-	2	-
2003.....	30	1	-	-	-	-	-	-	-	-	1	4
2002.....	22	-	3	-	2	-	-	1	-	1	1	1
2001.....	37	1	3	1	1	-	-	1	-	1	-	-
2000.....	28	2	3	1	-	-	-	-	-	2	-	3
1999.....	21	-	1	1	1	-	-	-	5	-	-	1
Black, both sexes												
2004.....	14	2	1	-	-	-	-	-	-	1	1	2
2003.....	17	2	1	-	-	-	-	-	-	-	2	4
2002.....	10	3	1	-	-	-	-	-	-	1	-	1
2001.....	8	1	-	-	-	-	-	-	1	2	-	-
2000.....	11	3	-	-	-	-	-	-	1	1	1	1
1999.....	14	-	-	-	-	1	2	-	1	1	2	-
Black, male												
2004.....	7	1	-	-	-	-	-	-	-	1	1	1
2003.....	7	1	1	-	-	-	-	-	-	-	-	3
2002.....	5	3	-	-	-	-	-	-	-	-	-	1
2001.....	3	1	-	-	-	-	-	-	-	1	-	-
2000.....	7	1	-	-	-	-	-	-	1	-	1	-
1999.....	3	-	-	-	-	-	1	-	-	-	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
IX. Diseases of the circulatory system (I00-I99)												
All races, both sexes												
2004.....	865,863	593	236	121	220	461	907	1,474	2,669	5,225	11,226	19,297
2003.....	907,180	591	238	135	216	533	964	1,405	2,823	5,501	11,771	19,364
2002.....	923,339	667	227	128	232	505	813	1,386	2,739	5,669	11,598	18,967
2001.....	926,999	622	290	139	226	453	865	1,396	2,766	5,668	11,303	17,990
2000.....	941,526	663	238	141	233	503	843	1,353	2,572	5,616	11,327	18,034
1999.....	954,339	667	235	145	211	559	786	1,300	2,666	6,016	11,265	17,556
All races, male												
2004.....	408,541	325	120	54	119	301	585	945	1,729	3,449	7,519	13,283
2003.....	425,776	331	119	76	130	328	585	853	1,816	3,642	7,827	13,362
2002.....	431,626	358	123	57	137	337	529	881	1,756	3,669	7,841	13,135
2001.....	430,046	326	154	65	119	283	541	858	1,750	3,723	7,606	12,545
2000.....	437,893	357	130	81	131	301	498	818	1,617	3,608	7,692	12,689
1999.....	443,541	360	128	85	115	351	461	783	1,703	3,953	7,677	12,252
All races, female												
2004.....	457,322	268	116	67	101	160	322	529	940	1,776	3,707	6,014
2003.....	481,404	260	119	59	86	205	379	552	1,007	1,859	3,944	6,002
2002.....	491,713	309	104	71	95	168	284	505	983	2,000	3,757	5,832
2001.....	496,953	296	136	74	107	170	324	538	1,016	1,945	3,697	5,445
2000.....	503,633	306	108	60	102	202	345	535	955	2,008	3,635	5,345
1999.....	510,798	307	107	60	96	208	325	517	963	2,063	3,588	5,304
White, both sexes												
2004.....	746,570	399	151	83	148	306	611	920	1,736	3,624	7,992	13,845
2003.....	783,863	371	152	89	148	364	617	897	1,853	3,754	8,289	13,871
2002.....	800,601	441	162	92	153	349	535	881	1,768	3,895	8,337	13,686
2001.....	804,895	400	188	100	158	312	547	906	1,852	3,919	7,898	12,952
2000.....	820,006	454	164	99	152	335	545	878	1,709	3,899	8,010	12,983
1999.....	832,809	452	161	109	143	388	513	855	1,807	4,181	7,886	12,705

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
IX. Diseases of the circulatory system (I00-I99)												
White, male												
2004.....	351,489	222	82	38	81	199	399	618	1,160	2,451	5,642	9,922
2003.....	366,959	200	70	49	92	223	381	555	1,235	2,602	5,785	9,982
2002.....	373,633	243	92	42	89	230	359	583	1,166	2,591	5,910	9,919
2001.....	372,521	220	98	49	87	192	353	574	1,208	2,654	5,602	9,448
2000.....	380,713	246	97	59	92	204	333	549	1,130	2,620	5,710	9,633
1999.....	385,971	243	90	65	76	240	299	527	1,200	2,844	5,645	9,287
White, female												
2004.....	395,081	177	69	45	67	107	212	302	576	1,173	2,350	3,923
2003.....	416,904	171	82	40	56	141	236	342	618	1,152	2,504	3,889
2002.....	426,968	198	70	50	64	119	176	298	602	1,304	2,427	3,767
2001.....	432,374	180	90	51	71	120	194	332	644	1,265	2,296	3,504
2000.....	439,293	208	67	40	60	131	212	329	579	1,279	2,300	3,350
1999.....	446,838	209	71	44	67	148	214	328	607	1,337	2,241	3,418
Black, both sexes												
2004.....	101,276	152	75	36	65	138	251	485	821	1,425	2,943	4,910
2003.....	105,005	179	73	39	56	146	311	450	850	1,542	3,105	4,914
2002.....	105,012	193	52	27	69	130	246	454	867	1,576	2,933	4,758
2001.....	105,058	191	89	34	54	127	286	449	817	1,545	3,087	4,534
2000.....	105,041	176	64	35	64	150	263	430	790	1,546	2,997	4,589
1999.....	105,309	181	62	35	58	155	253	401	761	1,659	3,082	4,372
Black, male												
2004.....	47,719	83	37	16	33	94	161	282	480	881	1,667	2,975
2003.....	49,221	110	43	23	30	94	186	264	491	892	1,797	2,983
2002.....	48,630	99	26	10	41	86	147	266	510	931	1,700	2,848
2001.....	48,576	97	48	13	27	82	167	263	473	922	1,780	2,733
2000.....	48,315	93	28	18	31	83	143	239	440	876	1,762	2,744
1999.....	48,860	97	33	19	34	101	147	232	442	981	1,819	2,643

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

IX. Diseases of the circulatory system (I00-I99)

Black, female

2004.....	53,557	69	38	20	32	44	90	203	341	544	1,276	1,935
2003.....	55,784	69	30	16	26	52	125	186	359	650	1,308	1,931
2002.....	56,382	94	26	17	28	44	99	188	357	645	1,233	1,910
2001.....	56,482	94	41	21	27	45	119	186	344	623	1,307	1,801
2000.....	56,726	83	36	17	33	67	120	191	350	670	1,235	1,845
1999.....	56,449	84	29	16	24	54	106	169	319	678	1,263	1,729

American Indian 1/, both sexes

2004.....	3,428	12	2	1	2	8	7	28	32	62	92	152
2003.....	3,503	9	5	1	2	5	9	13	39	64	114	169
2002.....	3,282	6	6	2	5	10	10	15	26	48	101	132
2001.....	3,187	11	3	3	1	3	8	12	25	65	99	155
2000.....	3,216	7	2	2	5	2	11	13	18	53	97	131
1999.....	3,134	8	5	-	6	2	8	11	33	50	90	119

American Indian 1/, male

2004.....	1,798	6	-	-	2	2	3	19	24	38	60	99
2003.....	1,889	5	2	-	-	3	7	6	32	48	68	114
2002.....	1,765	4	2	1	3	7	9	8	21	36	67	95
2001.....	1,677	3	2	2	-	1	7	4	17	40	64	106
2000.....	1,678	3	1	1	2	2	6	6	11	30	66	90
1999.....	1,617	5	2	-	2	1	6	6	13	34	59	81

American Indian 1/, female

2004.....	1,630	6	2	1	-	6	4	9	8	24	32	53
2003.....	1,614	4	3	1	2	2	2	7	7	16	46	55
2002.....	1,517	2	4	1	2	3	1	7	5	12	34	37
2001.....	1,510	8	1	1	1	2	1	8	8	25	35	49
2000.....	1,538	4	1	1	3	-	5	7	7	23	31	41
1999.....	1,517	3	3	-	4	1	2	5	20	16	31	38

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
IX. Diseases of the circulatory system (I00-I99)												
Asian or Pacific Islander 2/, both sexes												
2004.....	14,589	30	8	1	5	9	38	41	80	114	199	390
2003.....	14,809	32	8	6	10	18	27	45	81	141	263	410
2002.....	14,444	27	7	7	5	16	22	36	78	150	227	391
2001.....	13,859	20	10	2	13	11	24	29	72	139	219	349
2000.....	13,263	26	8	5	12	16	24	32	55	118	223	331
1999.....	13,087	26	7	1	4	14	12	33	65	126	207	360
Asian or Pacific Islander 2/, male												
2004.....	7,535	14	1	-	3	6	22	26	65	79	150	287
2003.....	7,707	16	4	4	8	8	11	28	58	100	177	283
2002.....	7,598	12	3	4	4	14	14	24	59	111	164	273
2001.....	7,272	6	6	1	5	8	14	17	52	107	160	258
2000.....	7,187	15	4	3	6	12	16	24	36	82	154	222
1999.....	7,093	15	3	1	3	9	9	18	48	94	154	241
Asian or Pacific Islander 2/, female												
2004.....	7,054	16	7	1	2	3	16	15	15	35	49	103
2003.....	7,102	16	4	2	2	10	16	17	23	41	86	127
2002.....	6,846	15	4	3	1	2	8	12	19	39	63	118
2001.....	6,587	14	4	1	8	3	10	12	20	32	59	91
2000.....	6,076	11	4	2	6	4	8	8	19	36	69	109
1999.....	5,994	11	4	-	1	5	3	15	17	32	53	119

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
All races, both sexes												
2004.....	3,254	2	3	1	3	3	4	14	13	32	64	85
2003.....	3,624	1	1	2	4	4	7	11	14	41	58	107
2002.....	3,579	4	1	4	2	8	10	10	26	38	57	97
2001.....	3,489	3	1	1	4	3	8	11	23	45	85	107
2000.....	3,582	6	2	2	2	9	8	10	23	49	92	119
1999.....	3,676	6	1	2	3	5	8	12	30	47	77	101
All races, male												
2004.....	1,009	2	3	1	1	2	2	4	5	14	24	39
2003.....	1,153	1	-	2	2	3	4	8	6	19	25	39
2002.....	1,078	3	1	1	1	3	5	6	12	19	26	40
2001.....	1,031	3	1	-	2	2	5	4	11	18	36	46
2000.....	1,072	5	2	1	2	5	6	2	10	25	36	49
1999.....	1,042	2	1	1	1	4	6	5	12	19	30	40
All races, female												
2004.....	2,245	-	-	-	2	1	2	10	8	18	40	46
2003.....	2,471	-	1	-	2	1	3	3	8	22	33	68
2002.....	2,501	1	-	3	1	5	5	4	14	19	31	57
2001.....	2,458	-	-	1	2	1	3	7	12	27	49	61
2000.....	2,510	1	-	1	-	4	2	8	13	24	56	70
1999.....	2,634	4	-	1	2	1	2	7	18	28	47	61
White, both sexes												
2004.....	2,911	2	2	1	1	2	2	9	7	19	38	57
2003.....	3,226	1	1	2	2	-	5	9	9	25	36	71
2002.....	3,204	4	1	1	2	3	7	6	18	32	36	70
2001.....	3,086	1	1	1	2	3	7	5	10	28	57	70
2000.....	3,229	5	2	2	2	6	4	7	18	34	62	90
1999.....	3,302	4	1	1	2	2	6	8	17	33	51	79

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)

White, male

2004.....	898	2	2	1	-	2	1	1	1	8	15	29
2003.....	1,023	1	-	2	1	-	4	6	3	12	14	29
2002.....	950	3	1	1	1	-	4	3	8	15	15	30
2001.....	910	1	1	-	1	2	5	2	7	12	28	32
2000.....	965	4	2	1	2	3	3	1	9	19	27	39
1999.....	933	1	1	-	-	1	4	4	8	14	21	32

White, female

2004.....	2,013	-	-	-	1	-	1	8	6	11	23	28
2003.....	2,203	-	1	-	1	-	1	3	6	13	22	42
2002.....	2,254	1	-	-	1	3	3	3	10	17	21	40
2001.....	2,176	-	-	1	1	1	2	3	3	16	29	38
2000.....	2,264	1	-	1	-	3	1	6	9	15	35	51
1999.....	2,369	3	-	1	2	1	2	4	9	19	30	47

Black, both sexes

2004.....	247	-	1	-	2	1	2	3	3	11	22	27
2003.....	308	-	-	-	2	2	2	2	4	13	16	32
2002.....	252	-	-	2	-	3	2	3	6	4	16	21
2001.....	302	2	-	-	1	-	1	3	10	15	26	27
2000.....	265	1	-	-	-	3	4	3	5	11	24	23
1999.....	275	1	-	1	1	2	2	4	9	12	20	20

Black, male

2004.....	84	-	1	-	1	-	1	2	2	6	8	9
2003.....	110	-	-	-	1	2	-	2	2	6	9	9
2002.....	93	-	-	-	-	2	1	2	2	2	10	7
2001.....	86	2	-	-	-	-	-	-	3	6	7	11
2000.....	79	1	-	-	-	2	3	1	1	5	7	8
1999.....	88	1	-	1	1	2	2	1	2	4	8	8

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Black, female												
2004.....	163	-	-	-	1	1	1	1	1	5	14	18
2003.....	198	-	-	-	1	-	2	-	2	7	7	23
2002.....	159	-	-	2	-	1	1	1	4	2	6	14
2001.....	216	-	-	-	1	-	1	3	7	9	19	16
2000.....	186	-	-	-	-	1	1	2	4	6	17	15
1999.....	187	-	-	-	-	-	-	3	7	8	12	12
American Indian 1/, both sexes												
2004.....	15	-	-	-	-	-	-	-	1	-	1	-
2003.....	11	-	-	-	-	-	-	-	-	1	-	-
2002.....	23	-	-	-	-	1	-	1	-	-	1	-
2001.....	21	-	-	-	-	-	-	-	-	-	1	3
2000.....	18	-	-	-	-	-	-	-	-	1	1	1
1999.....	19	-	-	-	-	-	-	-	2	1	-	-
American Indian 1/, male												
2004.....	6	-	-	-	-	-	-	-	-	-	1	-
2003.....	3	-	-	-	-	-	-	-	-	-	-	-
2002.....	9	-	-	-	-	-	-	1	-	-	-	-
2001.....	11	-	-	-	-	-	-	-	-	-	1	2
2000.....	3	-	-	-	-	-	-	-	-	-	-	-
1999.....	2	-	-	-	-	-	-	-	1	1	-	-
American Indian 1/, female												
2004.....	9	-	-	-	-	-	-	-	1	-	-	-
2003.....	8	-	-	-	-	-	-	-	-	1	-	-
2002.....	14	-	-	-	-	1	-	-	-	-	1	-
2001.....	10	-	-	-	-	-	-	-	-	-	-	1
2000.....	15	-	-	-	-	-	-	-	-	1	1	1
1999.....	17	-	-	-	-	-	-	-	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Asian or Pacific Islander 2/, both sexes												
2004.....	81	-	-	-	-	-	-	2	2	2	3	1
2003.....	79	-	-	-	-	2	-	-	1	2	6	4
2002.....	100	-	-	1	-	1	1	-	2	2	4	6
2001.....	80	-	-	-	1	-	-	3	3	2	1	7
2000.....	70	-	-	-	-	-	-	-	-	3	5	5
1999.....	80	1	-	-	-	1	-	-	2	1	6	2
Asian or Pacific Islander 2/, male												
2004.....	21	-	-	-	-	-	-	1	2	-	-	1
2003.....	17	-	-	-	-	1	-	-	1	1	2	1
2002.....	26	-	-	-	-	1	-	-	2	2	1	3
2001.....	24	-	-	-	1	-	-	2	1	-	-	1
2000.....	25	-	-	-	-	-	-	-	-	1	2	2
1999.....	19	-	-	-	-	1	-	-	1	-	1	-
Asian or Pacific Islander 2/, female												
2004.....	60	-	-	-	-	-	-	1	-	2	3	-
2003.....	62	-	-	-	-	1	-	-	-	1	4	3
2002.....	74	-	-	1	-	-	1	-	-	-	3	3
2001.....	56	-	-	-	-	-	-	1	2	2	1	6
2000.....	45	-	-	-	-	-	-	-	-	2	3	3
1999.....	61	1	-	-	-	-	-	-	1	1	5	2

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever (I00-I02)												
All races, both sexes												
2004.....	33	-	-	-	-	-	-	-	-	-	1	1
2003.....	41	-	-	-	-	-	2	1	1	-	1	2
2002.....	53	-	-	3	-	-	-	-	-	3	1	3
2001.....	65	-	-	-	2	-	-	-	3	-	2	3
2000.....	37	-	-	-	1	1	-	-	1	1	1	4
1999.....	30	-	-	-	2	-	-	-	1	-	1	1
All races, male												
2004.....	13	-	-	-	-	-	-	-	-	-	1	1
2003.....	19	-	-	-	-	-	2	1	-	-	1	1
2002.....	18	-	-	-	-	-	-	-	-	2	1	3
2001.....	23	-	-	-	1	-	-	-	1	-	1	2
2000.....	16	-	-	-	1	-	-	-	-	1	-	1
1999.....	14	-	-	-	1	-	-	-	-	-	-	1
All races, female												
2004.....	20	-	-	-	-	-	-	-	-	-	-	-
2003.....	22	-	-	-	-	-	-	-	1	-	-	1
2002.....	35	-	-	3	-	-	-	-	-	1	-	-
2001.....	42	-	-	-	1	-	-	-	2	-	1	1
2000.....	21	-	-	-	-	1	-	-	1	-	1	3
1999.....	16	-	-	-	1	-	-	-	1	-	1	-
White, both sexes												
2004.....	28	-	-	-	-	-	-	-	-	-	1	1
2003.....	31	-	-	-	-	-	2	1	1	-	-	2
2002.....	45	-	-	-	-	-	-	-	-	3	-	3
2001.....	53	-	-	-	1	-	-	-	1	-	1	3
2000.....	32	-	-	-	1	1	-	-	-	1	1	4
1999.....	23	-	-	-	1	-	-	-	1	-	-	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever (I00-I02)												
White, male												
2004.....	11	-	-	-	-	-	-	-	-	-	1	1
2003.....	15	-	-	-	-	-	2	1	-	-	-	1
2002.....	17	-	-	-	-	-	-	-	-	2	-	3
2001.....	20	-	-	-	1	-	-	-	1	-	-	2
2000.....	15	-	-	-	1	-	-	-	-	1	-	1
1999.....	11	-	-	-	-	-	-	-	-	-	-	1
White, female												
2004.....	17	-	-	-	-	-	-	-	-	-	-	-
2003.....	16	-	-	-	-	-	-	-	1	-	-	1
2002.....	28	-	-	-	-	-	-	-	-	1	-	-
2001.....	33	-	-	-	-	-	-	-	-	-	1	1
2000.....	17	-	-	-	-	1	-	-	-	-	1	3
1999.....	12	-	-	-	1	-	-	-	1	-	-	-
Black, both sexes												
2004.....	5	-	-	-	-	-	-	-	-	-	-	-
2003.....	9	-	-	-	-	-	-	-	-	-	1	-
2002.....	6	-	-	2	-	-	-	-	-	-	1	-
2001.....	10	-	-	-	1	-	-	-	1	-	1	-
2000.....	5	-	-	-	-	-	-	-	1	-	-	-
1999.....	7	-	-	-	1	-	-	-	-	-	1	-
Black, male												
2004.....	2	-	-	-	-	-	-	-	-	-	-	-
2003.....	3	-	-	-	-	-	-	-	-	-	1	-
2002.....	1	-	-	-	-	-	-	-	-	-	1	-
2001.....	3	-	-	-	-	-	-	-	-	-	1	-
2000.....	1	-	-	-	-	-	-	-	-	-	-	-
1999.....	3	-	-	-	1	-	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Chronic rheumatic heart diseases (I05-I09)

All races, both sexes

2004.....	3,221	2	3	1	3	3	4	14	13	32	63	84
2003.....	3,583	1	1	2	4	4	5	10	13	41	57	105
2002.....	3,526	4	1	1	2	8	10	10	26	35	56	94
2001.....	3,424	3	1	1	2	3	8	11	20	45	83	104
2000.....	3,545	6	2	2	1	8	8	10	22	48	91	115
1999.....	3,646	6	1	2	1	5	8	12	29	47	76	100

All races, male

2004.....	996	2	3	1	1	2	2	4	5	14	23	38
2003.....	1,134	1	-	2	2	3	2	7	6	19	24	38
2002.....	1,060	3	1	1	1	3	5	6	12	17	25	37
2001.....	1,008	3	1	-	1	2	5	4	10	18	35	44
2000.....	1,056	5	2	1	1	5	6	2	10	24	36	48
1999.....	1,028	2	1	1	-	4	6	5	12	19	30	39

All races, female

2004.....	2,225	-	-	-	2	1	2	10	8	18	40	46
2003.....	2,449	-	1	-	2	1	3	3	7	22	33	67
2002.....	2,466	1	-	-	1	5	5	4	14	18	31	57
2001.....	2,416	-	-	1	1	1	3	7	10	27	48	60
2000.....	2,489	1	-	1	-	3	2	8	12	24	55	67
1999.....	2,618	4	-	1	1	1	2	7	17	28	46	61

White, both sexes

2004.....	2,883	2	2	1	1	2	2	9	7	19	37	56
2003.....	3,195	1	1	2	2	-	3	8	8	25	36	69
2002.....	3,159	4	1	1	2	3	7	6	18	29	36	67
2001.....	3,033	1	1	1	1	3	7	5	9	28	56	67
2000.....	3,197	5	2	2	1	5	4	7	18	33	61	86
1999.....	3,279	4	1	1	1	2	6	8	16	33	51	78

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Chronic rheumatic heart diseases (I05-I09)

White, male

2004.....	887	2	2	1	-	2	1	1	1	8	14	28
2003.....	1,008	1	-	2	1	-	2	5	3	12	14	28
2002.....	933	3	1	1	1	-	4	3	8	13	15	27
2001.....	890	1	1	-	-	2	5	2	6	12	28	30
2000.....	950	4	2	1	1	3	3	1	9	18	27	38
1999.....	922	1	1	-	-	1	4	4	8	14	21	31

White, female

2004.....	1,996	-	-	-	1	-	1	8	6	11	23	28
2003.....	2,187	-	1	-	1	-	1	3	5	13	22	41
2002.....	2,226	1	-	-	1	3	3	3	10	16	21	40
2001.....	2,143	-	-	1	1	1	2	3	3	16	28	37
2000.....	2,247	1	-	1	-	2	1	6	9	15	34	48
1999.....	2,357	3	-	1	1	1	2	4	8	19	30	47

Black, both sexes

2004.....	242	-	1	-	2	1	2	3	3	11	22	27
2003.....	299	-	-	-	2	2	2	2	4	13	15	32
2002.....	246	-	-	-	-	3	2	3	6	4	15	21
2001.....	292	2	-	-	-	-	1	3	9	15	25	27
2000.....	260	1	-	-	-	3	4	3	4	11	24	23
1999.....	268	1	-	1	-	2	2	4	9	12	19	20

Black, male

2004.....	82	-	1	-	1	-	1	2	2	6	8	9
2003.....	107	-	-	-	1	2	-	2	2	6	8	9
2002.....	92	-	-	-	-	2	1	2	2	2	9	7
2001.....	83	2	-	-	-	-	-	-	3	6	6	11
2000.....	78	1	-	-	-	2	3	1	1	5	7	8
1999.....	85	1	-	1	-	2	2	1	2	4	8	8

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic rheumatic heart diseases (I05-I09)												
Black, female												
2004.....	160	-	-	-	1	1	1	1	1	5	14	18
2003.....	192	-	-	-	1	-	2	-	2	7	7	23
2002.....	154	-	-	-	-	1	1	1	4	2	6	14
2001.....	209	-	-	-	-	-	1	3	6	9	19	16
2000.....	182	-	-	-	-	1	1	2	3	6	17	15
1999.....	183	-	-	-	-	-	-	3	7	8	11	12
American Indian 1/, both sexes												
2004.....	15	-	-	-	-	-	-	-	1	-	1	-
2003.....	11	-	-	-	-	-	-	-	-	1	-	-
2002.....	23	-	-	-	-	1	-	1	-	-	1	-
2001.....	20	-	-	-	-	-	-	-	-	-	1	3
2000.....	18	-	-	-	-	-	-	-	-	1	1	1
1999.....	19	-	-	-	-	-	-	-	2	1	-	-
American Indian 1/, male												
2004.....	6	-	-	-	-	-	-	-	-	-	1	-
2003.....	3	-	-	-	-	-	-	-	-	-	-	-
2002.....	9	-	-	-	-	-	-	1	-	-	-	-
2001.....	11	-	-	-	-	-	-	-	-	-	1	2
2000.....	3	-	-	-	-	-	-	-	-	-	-	-
1999.....	2	-	-	-	-	-	-	-	1	1	-	-
American Indian 1/, female												
2004.....	9	-	-	-	-	-	-	-	1	-	-	-
2003.....	8	-	-	-	-	-	-	-	-	1	-	-
2002.....	14	-	-	-	-	1	-	-	-	-	1	-
2001.....	9	-	-	-	-	-	-	-	-	-	-	1
2000.....	15	-	-	-	-	-	-	-	-	1	1	1
1999.....	17	-	-	-	-	-	-	-	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic rheumatic heart diseases (I05-I09)												
Asian or Pacific Islander 2/, both sexes												
2004.....	81	-	-	-	-	-	-	2	2	2	3	1
2003.....	78	-	-	-	-	2	-	-	1	2	6	4
2002.....	98	-	-	-	-	1	1	-	2	2	4	6
2001.....	79	-	-	-	1	-	-	3	2	2	1	7
2000.....	70	-	-	-	-	-	-	-	-	3	5	5
1999.....	80	1	-	-	-	1	-	-	2	1	6	2
Asian or Pacific Islander 2/, male												
2004.....	21	-	-	-	-	-	-	1	2	-	-	1
2003.....	16	-	-	-	-	1	-	-	1	1	2	1
2002.....	26	-	-	-	-	1	-	-	2	2	1	3
2001.....	24	-	-	-	1	-	-	2	1	-	-	1
2000.....	25	-	-	-	-	-	-	-	-	1	2	2
1999.....	19	-	-	-	-	1	-	-	1	-	1	-
Asian or Pacific Islander 2/, female												
2004.....	60	-	-	-	-	-	-	1	-	2	3	-
2003.....	62	-	-	-	-	1	-	-	-	1	4	3
2002.....	72	-	-	-	-	-	1	-	-	-	3	3
2001.....	55	-	-	-	-	-	-	1	1	2	1	6
2000.....	45	-	-	-	-	-	-	-	-	2	3	3
1999.....	61	1	-	-	-	-	-	-	1	1	5	2

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Rheumatic mitral valve diseases (I05)

All races, both sexes

2004.....	1,475	2	2	-	1	1	1	4	5	26	31	49
2003.....	1,719	1	-	1	-	3	2	6	5	18	32	67
2002.....	1,829	3	1	1	1	6	5	7	16	18	35	53
2001.....	1,902	3	-	-	2	2	5	9	12	31	48	68
2000.....	2,038	4	1	-	-	4	3	8	16	29	53	75
1999.....	2,050	6	-	-	1	3	4	7	12	24	37	67

All races, male

2004.....	444	2	2	-	1	1	1	-	1	10	10	23
2003.....	552	1	-	1	-	2	-	5	1	9	10	24
2002.....	529	2	1	1	-	2	1	5	7	6	18	18
2001.....	535	3	-	-	1	2	2	3	5	10	20	27
2000.....	609	3	1	-	-	2	2	1	6	16	20	33
1999.....	562	2	-	-	-	2	3	3	4	10	14	27

All races, female

2004.....	1,031	-	-	-	-	-	-	4	4	16	21	26
2003.....	1,167	-	-	-	-	1	2	1	4	9	22	43
2002.....	1,300	1	-	-	1	4	4	2	9	12	17	35
2001.....	1,367	-	-	-	1	-	3	6	7	21	28	41
2000.....	1,429	1	-	-	-	2	1	7	10	13	33	42
1999.....	1,488	4	-	-	1	1	1	4	8	14	23	40

White, both sexes

2004.....	1,299	2	1	-	-	1	1	4	4	14	15	34
2003.....	1,512	1	-	1	-	-	-	5	3	14	24	43
2002.....	1,631	3	1	1	1	3	3	3	10	16	25	36
2001.....	1,686	1	-	-	1	2	4	5	5	23	35	47
2000.....	1,853	3	1	-	-	3	1	5	12	18	40	59
1999.....	1,838	4	-	-	1	2	3	5	6	19	26	54

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic mitral valve diseases (I05)												
White, male												
2004.....	390	2	1	-	-	1	1	-	-	5	4	16
2003.....	482	1	-	1	-	-	-	4	-	7	8	17
2002.....	462	2	1	1	-	-	-	2	5	6	12	11
2001.....	477	1	-	-	-	2	2	2	3	10	18	20
2000.....	552	2	1	-	-	1	1	-	5	12	16	26
1999.....	503	1	-	-	-	1	2	3	3	8	11	23
White, female												
2004.....	909	-	-	-	-	-	-	4	4	9	11	18
2003.....	1,030	-	-	-	-	-	-	1	3	7	16	26
2002.....	1,169	1	-	-	1	3	3	1	5	10	13	25
2001.....	1,209	-	-	-	1	-	2	3	2	13	17	27
2000.....	1,301	1	-	-	-	2	-	5	7	6	24	33
1999.....	1,335	3	-	-	1	1	1	2	3	11	15	31
Black, both sexes												
2004.....	125	-	1	-	1	-	-	-	-	10	12	14
2003.....	158	-	-	-	-	1	2	1	2	4	5	21
2002.....	140	-	-	-	-	2	1	3	6	2	9	14
2001.....	167	2	-	-	-	-	1	2	6	8	12	17
2000.....	131	1	-	-	-	1	2	3	4	7	12	11
1999.....	166	1	-	-	-	1	1	2	4	5	8	11
Black, male												
2004.....	41	-	1	-	1	-	-	-	-	5	5	6
2003.....	58	-	-	-	-	1	-	1	1	2	1	6
2002.....	52	-	-	-	-	2	1	2	2	-	6	6
2001.....	44	2	-	-	-	-	-	-	2	-	2	7
2000.....	41	1	-	-	-	1	1	1	1	3	4	5
1999.....	53	1	-	-	-	1	1	-	-	2	3	4

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic mitral valve diseases (I05)												
Asian or Pacific Islander 2/, both sexes												
2004.....	43	-	-	-	-	-	-	-	1	2	3	1
2003.....	42	-	-	-	-	2	-	-	-	-	3	3
2002.....	45	-	-	-	-	-	1	-	-	-	-	3
2001.....	40	-	-	-	1	-	-	2	1	-	1	3
2000.....	45	-	-	-	-	-	-	-	-	3	1	4
1999.....	37	1	-	-	-	-	-	-	1	-	3	2
Asian or Pacific Islander 2/, male												
2004.....	9	-	-	-	-	-	-	-	1	-	-	1
2003.....	9	-	-	-	-	1	-	-	-	-	1	1
2002.....	10	-	-	-	-	-	-	-	-	-	-	1
2001.....	11	-	-	-	1	-	-	1	-	-	-	-
2000.....	15	-	-	-	-	-	-	-	-	1	-	2
1999.....	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	34	-	-	-	-	-	-	-	-	2	3	-
2003.....	33	-	-	-	-	1	-	-	-	-	2	2
2002.....	35	-	-	-	-	-	1	-	-	-	-	2
2001.....	29	-	-	-	-	-	-	1	1	-	1	3
2000.....	30	-	-	-	-	-	-	-	-	2	1	2
1999.....	32	1	-	-	-	-	-	-	1	-	3	2

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic aortic valve diseases (I06)												
All races, both sexes												
2004.....	103	-	-	-	-	-	-	-	-	-	1	1
2003.....	103	-	-	-	-	-	1	-	-	1	3	1
2002.....	128	-	-	-	-	-	-	-	-	-	2	1
2001.....	66	-	-	-	-	-	-	-	1	-	2	3
2000.....	35	-	-	-	-	1	-	-	-	1	1	2
1999.....	53	-	-	-	-	-	1	-	-	1	1	2
All races, male												
2004.....	40	-	-	-	-	-	-	-	-	-	1	1
2003.....	36	-	-	-	-	-	1	-	-	-	2	-
2002.....	40	-	-	-	-	-	-	-	-	-	-	-
2001.....	26	-	-	-	-	-	-	-	1	-	1	2
2000.....	18	-	-	-	-	1	-	-	-	1	1	1
1999.....	18	-	-	-	-	-	1	-	-	1	1	-
All races, female												
2004.....	63	-	-	-	-	-	-	-	-	-	-	-
2003.....	67	-	-	-	-	-	-	-	-	1	1	1
2002.....	88	-	-	-	-	-	-	-	-	-	2	1
2001.....	40	-	-	-	-	-	-	-	-	-	1	1
2000.....	17	-	-	-	-	-	-	-	-	-	-	1
1999.....	35	-	-	-	-	-	-	-	-	-	-	2
White, both sexes												
2004.....	97	-	-	-	-	-	-	-	-	-	1	1
2003.....	96	-	-	-	-	-	1	-	-	-	1	-
2002.....	114	-	-	-	-	-	-	-	-	-	1	1
2001.....	56	-	-	-	-	-	-	-	-	-	-	1
2000.....	30	-	-	-	-	1	-	-	-	1	-	2
1999.....	50	-	-	-	-	-	1	-	-	1	1	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of both mitral and aortic valves (I08.0)												
All races, both sexes												
2004.....	716	-	-	-	-	-	-	4	1	3	11	10
2003.....	775	-	-	-	1	-	-	-	3	10	5	8
2002.....	507	-	-	-	-	-	1	1	-	5	4	11
2001.....	383	-	-	-	-	1	2	1	4	3	4	8
2000.....	317	1	-	-	-	-	-	-	1	2	5	5
1999.....	325	-	1	-	-	-	1	1	3	2	12	8
All races, male												
2004.....	261	-	-	-	-	-	-	2	-	3	5	5
2003.....	278	-	-	-	-	-	-	-	2	6	3	4
2002.....	178	-	-	-	-	-	1	1	-	4	3	7
2001.....	145	-	-	-	-	-	2	1	1	2	2	4
2000.....	112	1	-	-	-	-	-	-	-	1	2	2
1999.....	119	-	1	-	-	-	-	1	2	1	6	4
All races, female												
2004.....	455	-	-	-	-	-	-	2	1	-	6	5
2003.....	497	-	-	-	1	-	-	-	1	4	2	4
2002.....	329	-	-	-	-	-	-	-	-	1	1	4
2001.....	238	-	-	-	-	1	-	-	3	1	2	4
2000.....	205	-	-	-	-	-	-	-	1	1	3	3
1999.....	206	-	-	-	-	-	1	-	1	1	6	4
White, both sexes												
2004.....	658	-	-	-	-	-	-	2	-	2	8	7
2003.....	710	-	-	-	-	-	-	-	-	6	3	6
2002.....	467	-	-	-	-	-	1	1	-	3	3	11
2001.....	349	-	-	-	-	1	2	-	2	3	4	4
2000.....	289	1	-	-	-	-	-	-	1	2	5	4
1999.....	291	-	1	-	-	-	1	-	2	1	7	6

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of both mitral and aortic valves (I08.0)												
White, male												
2004.....	237	-	-	-	-	-	-	-	-	2	4	4
2003.....	253	-	-	-	-	-	-	-	-	3	1	3
2002.....	160	-	-	-	-	-	1	1	-	2	2	7
2001.....	131	-	-	-	-	-	2	-	1	2	2	2
2000.....	100	1	-	-	-	-	-	-	-	1	2	2
1999.....	108	-	1	-	-	-	-	-	1	1	4	2
White, female												
2004.....	421	-	-	-	-	-	-	2	-	-	4	3
2003.....	457	-	-	-	-	-	-	-	-	3	2	3
2002.....	307	-	-	-	-	-	-	-	-	1	1	4
2001.....	218	-	-	-	-	1	-	-	1	1	2	2
2000.....	189	-	-	-	-	-	-	-	1	1	3	2
1999.....	183	-	-	-	-	-	1	-	1	-	3	4
Black, both sexes												
2004.....	43	-	-	-	-	-	-	1	-	1	3	3
2003.....	48	-	-	-	1	-	-	-	2	4	1	2
2002.....	30	-	-	-	-	-	-	-	-	1	1	-
2001.....	26	-	-	-	-	-	-	-	2	-	-	2
2000.....	20	-	-	-	-	-	-	-	-	-	-	-
1999.....	23	-	-	-	-	-	-	1	1	1	5	2
Black, male												
2004.....	20	-	-	-	-	-	-	1	-	1	1	1
2003.....	21	-	-	-	-	-	-	-	1	3	1	1
2002.....	14	-	-	-	-	-	-	-	-	1	1	-
2001.....	9	-	-	-	-	-	-	-	-	-	-	1
2000.....	7	-	-	-	-	-	-	-	-	-	-	-
1999.....	10	-	-	-	-	-	-	1	1	-	2	2

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)

All races, both sexes

2004.....	927	-	1	1	2	2	3	6	7	3	20	24
2003.....	986	-	1	1	3	1	2	4	5	12	17	29
2002.....	1,062	1	-	-	1	2	4	2	10	12	15	29
2001.....	1,073	-	1	1	-	-	1	1	3	11	29	25
2000.....	1,155	1	1	2	1	3	5	2	5	16	32	33
1999.....	1,218	-	-	2	-	2	2	4	14	20	26	23

All races, male

2004.....	251	-	1	1	-	1	1	2	4	1	7	9
2003.....	268	-	-	1	2	1	1	2	3	4	9	10
2002.....	313	1	-	-	1	1	3	-	5	7	4	12
2001.....	302	-	1	-	-	-	1	-	3	6	12	11
2000.....	317	1	1	1	1	2	4	1	4	6	13	12
1999.....	329	-	-	1	-	2	2	1	6	7	9	8

All races, female

2004.....	676	-	-	-	2	1	2	4	3	2	13	15
2003.....	718	-	1	-	1	-	1	2	2	8	8	19
2002.....	749	-	-	-	-	1	1	2	5	5	11	17
2001.....	771	-	-	1	-	-	-	1	-	5	17	14
2000.....	838	-	-	1	-	1	1	1	1	10	19	21
1999.....	889	-	-	1	-	-	-	3	8	13	17	15

White, both sexes

2004.....	829	-	1	1	1	1	1	3	3	3	13	14
2003.....	877	-	1	1	2	-	2	3	5	5	8	20
2002.....	947	1	-	-	1	-	3	2	8	10	7	19
2001.....	942	-	1	1	-	-	1	-	2	2	17	15
2000.....	1,025	1	1	2	1	1	3	2	5	12	16	21
1999.....	1,100	-	-	1	-	-	1	3	8	12	17	17

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)

White, male

2004.....	223	-	1	1	-	1	-	1	1	1	5	7
2003.....	239	-	-	1	1	-	1	1	3	2	4	8
2002.....	272	1	-	-	1	-	3	-	3	5	1	9
2001.....	259	-	1	-	-	-	1	-	2	-	8	7
2000.....	285	1	1	1	1	1	2	1	4	4	9	9
1999.....	294	-	-	-	-	-	1	1	4	4	5	6

White, female

2004.....	606	-	-	-	1	-	1	2	2	2	8	7
2003.....	638	-	1	-	1	-	1	2	2	3	4	12
2002.....	675	-	-	-	-	-	-	2	5	5	6	10
2001.....	683	-	-	1	-	-	-	-	-	2	9	8
2000.....	740	-	-	1	-	-	1	1	1	8	7	12
1999.....	806	-	-	1	-	-	-	2	4	8	12	11

Black, both sexes

2004.....	70	-	-	-	1	1	2	2	3	-	7	10
2003.....	88	-	-	-	1	1	-	1	-	4	8	8
2002.....	69	-	-	-	-	1	1	-	-	1	4	7
2001.....	92	-	-	-	-	-	-	1	1	7	12	6
2000.....	106	-	-	-	-	2	2	-	-	4	11	12
1999.....	76	-	-	1	-	1	1	1	4	6	6	6

Black, male

2004.....	19	-	-	-	-	-	1	1	2	-	2	2
2003.....	26	-	-	-	1	1	-	1	-	1	5	2
2002.....	25	-	-	-	-	-	-	-	-	1	2	1
2001.....	29	-	-	-	-	-	-	-	1	6	4	2
2000.....	27	-	-	-	-	1	2	-	-	2	2	3
1999.....	21	-	-	1	-	1	1	-	1	2	3	2

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)

Black, female

2004.....	51	-	-	-	1	1	1	1	1	-	5	8
2003.....	62	-	-	-	-	-	-	-	-	3	3	6
2002.....	44	-	-	-	-	1	1	-	-	-	2	6
2001.....	63	-	-	-	-	-	-	1	-	1	8	4
2000.....	79	-	-	-	-	1	-	-	-	2	9	9
1999.....	55	-	-	-	-	-	-	1	3	4	3	4

American Indian 1/, both sexes

2004.....	4	-	-	-	-	-	-	-	-	-	-	-
2003.....	3	-	-	-	-	-	-	-	-	1	-	-
2002.....	5	-	-	-	-	-	-	-	-	-	-	-
2001.....	9	-	-	-	-	-	-	-	-	-	-	2
2000.....	8	-	-	-	-	-	-	-	-	-	1	-
1999.....	7	-	-	-	-	-	-	-	1	1	-	-

American Indian 1/, male

2004.....	2	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	2	-	-	-	-	-	-	-	-	-	-	-
2001.....	6	-	-	-	-	-	-	-	-	-	-	2
2000.....	1	-	-	-	-	-	-	-	-	-	-	-
1999.....	1	-	-	-	-	-	-	-	-	1	-	-

American Indian 1/, female

2004.....	2	-	-	-	-	-	-	-	-	-	-	-
2003.....	3	-	-	-	-	-	-	-	-	1	-	-
2002.....	3	-	-	-	-	-	-	-	-	-	-	-
2001.....	3	-	-	-	-	-	-	-	-	-	-	-
2000.....	7	-	-	-	-	-	-	-	-	-	1	-
1999.....	6	-	-	-	-	-	-	-	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)

Asian or Pacific Islander 2/, both sexes

2004.....	24	-	-	-	-	-	-	1	1	-	-	-
2003.....	18	-	-	-	-	-	-	-	-	2	1	1
2002.....	41	-	-	-	-	1	-	-	2	1	4	3
2001.....	30	-	-	-	-	-	-	-	-	2	-	2
2000.....	16	-	-	-	-	-	-	-	-	-	4	-
1999.....	35	-	-	-	-	1	-	-	1	1	3	-

Asian or Pacific Islander 2/, male

2004.....	7	-	-	-	-	-	-	-	1	-	-	-
2003.....	3	-	-	-	-	-	-	-	-	1	-	-
2002.....	14	-	-	-	-	1	-	-	2	1	1	2
2001.....	8	-	-	-	-	-	-	-	-	-	-	-
2000.....	4	-	-	-	-	-	-	-	-	-	2	-
1999.....	13	-	-	-	-	1	-	-	1	-	1	-

Asian or Pacific Islander 2/, female

2004.....	17	-	-	-	-	-	-	1	-	-	-	-
2003.....	15	-	-	-	-	-	-	-	-	1	1	1
2002.....	27	-	-	-	-	-	-	-	-	-	3	1
2001.....	22	-	-	-	-	-	-	-	-	2	-	2
2000.....	12	-	-	-	-	-	-	-	-	-	2	-
1999.....	22	-	-	-	-	-	-	-	-	1	2	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive diseases (I10-I13)												
All races, both sexes												
2004.....	54,707	2	1	1	3	24	48	145	283	567	1,194	2,089
2003.....	53,393	1	4	-	2	16	64	125	306	570	1,201	1,933
2002.....	49,707	5	1	1	2	9	47	128	282	556	1,088	1,723
2001.....	46,765	1	1	1	1	13	41	107	271	493	1,028	1,499
2000.....	44,619	1	2	2	5	12	43	101	210	450	962	1,372
1999.....	42,997	4	3	2	1	11	41	79	230	495	896	1,298
All races, male												
2004.....	23,099	1	1	-	-	18	31	100	196	412	775	1,355
2003.....	22,075	1	4	-	1	11	45	77	202	383	771	1,290
2002.....	20,512	2	1	1	1	4	36	94	187	383	729	1,131
2001.....	18,874	1	-	-	-	6	31	64	181	313	696	1,000
2000.....	17,934	1	-	1	3	4	28	62	134	283	625	892
1999.....	17,194	2	1	1	1	9	27	57	156	325	597	832
All races, female												
2004.....	31,608	1	-	1	3	6	17	45	87	155	419	734
2003.....	31,318	-	-	-	1	5	19	48	104	187	430	643
2002.....	29,195	3	-	-	1	5	11	34	95	173	359	592
2001.....	27,891	-	1	1	1	7	10	43	90	180	332	499
2000.....	26,685	-	2	1	2	8	15	39	76	167	337	480
1999.....	25,803	2	2	1	-	2	14	22	74	170	299	466
White, both sexes												
2004.....	40,920	-	1	1	1	11	27	57	139	279	666	1,141
2003.....	39,934	-	2	-	1	9	32	65	142	297	644	1,018
2002.....	37,042	4	1	1	1	5	24	60	127	271	586	957
2001.....	34,906	-	1	-	1	8	19	51	126	249	502	772
2000.....	33,104	1	1	1	4	3	20	49	86	224	513	711
1999.....	31,711	3	1	1	-	5	14	37	115	221	429	662

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive diseases (I10-I13)												
White, male												
2004.....	16,704	-	1	-	-	10	20	49	107	214	471	804
2003.....	15,921	-	2	-	-	5	24	47	97	226	462	735
2002.....	14,713	2	1	1	1	1	20	46	89	193	433	675
2001.....	13,508	-	-	-	-	5	14	33	94	179	372	542
2000.....	12,822	1	-	1	3	1	15	34	63	161	355	512
1999.....	12,149	1	-	1	-	4	8	30	94	161	318	460
White, female												
2004.....	24,216	-	-	1	1	1	7	8	32	65	195	337
2003.....	24,013	-	-	-	1	4	8	18	45	71	182	283
2002.....	22,329	2	-	-	-	4	4	14	38	78	153	282
2001.....	21,398	-	1	-	1	3	5	18	32	70	130	230
2000.....	20,282	-	1	-	1	2	5	15	23	63	158	199
1999.....	19,562	2	1	-	-	1	6	7	21	60	111	202
Black, both sexes												
2004.....	12,426	2	-	-	2	12	21	84	134	272	503	895
2003.....	12,245	1	2	-	-	7	31	59	151	260	531	875
2002.....	11,579	1	-	-	1	4	20	63	149	270	475	738
2001.....	10,840	1	-	1	-	5	22	54	137	231	495	696
2000.....	10,582	-	1	1	1	9	23	48	120	220	429	636
1999.....	10,351	1	1	1	1	5	27	40	109	263	449	611
Black, male												
2004.....	5,762	1	-	-	-	8	11	50	79	186	285	513
2003.....	5,606	1	2	-	-	6	21	29	95	147	295	527
2002.....	5,268	-	-	-	-	3	13	45	93	178	277	434
2001.....	4,904	1	-	-	-	1	17	30	81	126	301	436
2000.....	4,670	-	-	-	-	3	13	26	68	118	257	364
1999.....	4,605	1	1	-	1	4	19	25	57	155	262	359

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Hypertensive diseases (I10-I13)

Black, female

2004.....	6,664	1	-	-	2	4	10	34	55	86	218	382
2003.....	6,639	-	-	-	-	1	10	30	56	113	236	348
2002.....	6,311	1	-	-	1	1	7	18	56	92	198	304
2001.....	5,936	-	-	1	-	4	5	24	56	105	194	260
2000.....	5,912	-	1	1	1	6	10	22	52	102	172	272
1999.....	5,746	-	-	1	-	1	8	15	52	108	187	252

American Indian 1/, both sexes

2004.....	229	-	-	-	-	1	-	2	3	7	7	14
2003.....	208	-	-	-	-	-	-	1	6	4	7	11
2002.....	149	-	-	-	-	-	2	1	-	7	8	6
2001.....	135	-	-	-	-	-	-	1	3	5	8	3
2000.....	141	-	-	-	-	-	-	1	-	1	2	4
1999.....	125	-	-	-	-	-	-	1	1	4	5	2

American Indian 1/, male

2004.....	114	-	-	-	-	-	-	-	3	5	5	10
2003.....	99	-	-	-	-	-	-	1	4	3	2	6
2002.....	64	-	-	-	-	-	2	-	-	5	5	5
2001.....	59	-	-	-	-	-	-	-	2	4	4	2
2000.....	59	-	-	-	-	-	-	-	-	-	2	2
1999.....	59	-	-	-	-	-	-	1	-	3	5	2

American Indian 1/, female

2004.....	115	-	-	-	-	1	-	2	-	2	2	4
2003.....	109	-	-	-	-	-	-	-	2	1	5	5
2002.....	85	-	-	-	-	-	-	1	-	2	3	1
2001.....	76	-	-	-	-	-	-	1	1	1	4	1
2000.....	82	-	-	-	-	-	-	1	-	1	-	2
1999.....	66	-	-	-	-	-	-	-	1	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive diseases (I10-I13)												
Asian or Pacific Islander 2/, both sexes												
2004.....	1,132	-	-	-	-	-	-	2	7	9	18	39
2003.....	1,006	-	-	-	1	-	1	-	7	9	19	29
2002.....	937	-	-	-	-	-	1	4	6	8	19	22
2001.....	884	-	-	-	-	-	-	1	5	8	23	28
2000.....	792	-	-	-	-	-	-	3	4	5	18	21
1999.....	810	-	1	-	-	1	-	1	5	7	13	23
Asian or Pacific Islander 2/, male												
2004.....	519	-	-	-	-	-	-	1	7	7	14	28
2003.....	449	-	-	-	1	-	-	-	6	7	12	22
2002.....	467	-	-	-	-	-	1	3	5	7	14	17
2001.....	403	-	-	-	-	-	-	1	4	4	19	20
2000.....	383	-	-	-	-	-	-	2	3	4	11	14
1999.....	381	-	-	-	-	1	-	1	5	6	12	11
Asian or Pacific Islander 2/, female												
2004.....	613	-	-	-	-	-	-	1	-	2	4	11
2003.....	557	-	-	-	-	-	1	-	1	2	7	7
2002.....	470	-	-	-	-	-	-	1	1	1	5	5
2001.....	481	-	-	-	-	-	-	-	1	4	4	8
2000.....	409	-	-	-	-	-	-	1	1	1	7	7
1999.....	429	-	1	-	-	-	-	-	-	1	1	12

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Essential (primary) hypertension (I10)												
All races, both sexes												
2004.....	13,456	2	1	-	-	2	5	6	20	41	109	268
2003.....	12,913	-	2	-	-	1	3	8	23	55	122	207
2002.....	11,658	4	1	1	-	1	8	8	15	46	120	219
2001.....	11,077	-	-	-	-	1	4	11	23	47	102	202
2000.....	10,296	1	1	1	2	5	3	10	17	37	120	211
1999.....	9,917	3	1	-	-	3	3	6	23	42	89	177
All races, male												
2004.....	4,691	1	1	-	-	1	3	3	11	32	69	161
2003.....	4,336	-	2	-	-	1	2	3	14	30	67	137
2002.....	3,965	2	1	1	-	-	6	5	9	28	77	141
2001.....	3,846	-	-	-	-	1	2	8	15	31	61	126
2000.....	3,631	1	-	1	2	-	1	6	13	23	72	147
1999.....	3,457	2	-	-	-	2	2	3	14	26	56	109
All races, female												
2004.....	8,765	1	-	-	-	1	2	3	9	9	40	107
2003.....	8,577	-	-	-	-	-	1	5	9	25	55	70
2002.....	7,693	2	-	-	-	1	2	3	6	18	43	78
2001.....	7,231	-	-	-	-	-	2	3	8	16	41	76
2000.....	6,665	-	1	-	-	5	2	4	4	14	48	64
1999.....	6,460	1	1	-	-	1	1	3	9	16	33	68
White, both sexes												
2004.....	11,020	-	1	-	-	-	4	2	8	23	68	156
2003.....	10,533	-	1	-	-	1	1	5	14	34	78	132
2002.....	9,541	3	1	1	-	-	6	4	6	21	74	126
2001.....	9,016	-	-	-	-	1	2	7	13	28	52	115
2000.....	8,375	1	1	1	2	2	1	7	7	20	76	129
1999.....	8,047	2	1	-	-	3	1	2	10	19	48	117

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Essential (primary) hypertension (I10)												
White, male												
2004.....	3,753	-	1	-	-	-	3	-	5	18	46	106
2003.....	3,378	-	1	-	-	1	1	2	7	22	45	91
2002.....	3,139	2	1	1	-	-	5	3	4	13	50	90
2001.....	2,986	-	-	-	-	1	1	4	11	19	34	80
2000.....	2,836	1	-	1	2	-	1	4	5	16	50	93
1999.....	2,706	1	-	-	-	2	-	2	9	15	33	77
White, female												
2004.....	7,267	-	-	-	-	-	1	2	3	5	22	50
2003.....	7,155	-	-	-	-	-	-	3	7	12	33	41
2002.....	6,402	1	-	-	-	-	1	1	2	8	24	36
2001.....	6,030	-	-	-	-	-	1	3	2	9	18	35
2000.....	5,539	-	1	-	-	2	-	3	2	4	26	36
1999.....	5,341	1	1	-	-	1	1	-	1	4	15	40
Black, both sexes												
2004.....	2,108	2	-	-	-	1	1	4	12	16	39	102
2003.....	2,114	-	1	-	-	-	2	3	9	21	42	70
2002.....	1,874	1	-	-	-	1	2	4	9	24	44	88
2001.....	1,835	-	-	-	-	-	2	4	9	17	46	83
2000.....	1,716	-	-	-	-	3	2	1	10	16	39	76
1999.....	1,667	1	-	-	-	-	2	4	12	23	40	54
Black, male												
2004.....	807	1	-	-	-	1	-	3	6	13	21	48
2003.....	853	-	1	-	-	-	1	1	7	8	21	45
2002.....	723	-	-	-	-	-	1	2	5	14	25	48
2001.....	768	-	-	-	-	-	1	4	3	11	23	44
2000.....	700	-	-	-	-	-	-	1	8	7	18	49
1999.....	661	1	-	-	-	-	2	1	5	11	22	28

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Essential (primary) hypertension (I10)												
Black, female												
2004.....	1,301	1	-	-	-	-	1	1	6	3	18	54
2003.....	1,261	-	-	-	-	-	1	2	2	13	21	25
2002.....	1,151	1	-	-	-	1	1	2	4	10	19	40
2001.....	1,067	-	-	-	-	-	1	-	6	6	23	39
2000.....	1,016	-	-	-	-	3	2	-	2	9	21	27
1999.....	1,006	-	-	-	-	-	-	3	7	12	18	26
American Indian 1/, both sexes												
2004.....	54	-	-	-	-	1	-	-	-	1	1	4
2003.....	47	-	-	-	-	-	-	-	-	-	1	2
2002.....	40	-	-	-	-	-	-	-	-	-	1	2
2001.....	29	-	-	-	-	-	-	-	1	1	1	-
2000.....	41	-	-	-	-	-	-	1	-	1	1	2
1999.....	25	-	-	-	-	-	-	-	1	-	-	1
American Indian 1/, male												
2004.....	24	-	-	-	-	-	-	-	-	1	1	3
2003.....	22	-	-	-	-	-	-	-	-	-	-	-
2002.....	11	-	-	-	-	-	-	-	-	-	1	1
2001.....	12	-	-	-	-	-	-	-	1	-	1	-
2000.....	17	-	-	-	-	-	-	-	-	-	1	1
1999.....	14	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, female												
2004.....	30	-	-	-	-	1	-	-	-	-	-	1
2003.....	25	-	-	-	-	-	-	-	-	-	1	2
2002.....	29	-	-	-	-	-	-	-	-	-	-	1
2001.....	17	-	-	-	-	-	-	-	-	1	-	-
2000.....	24	-	-	-	-	-	-	1	-	1	-	1
1999.....	11	-	-	-	-	-	-	-	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Hypertensive heart disease (I11)

All races, both sexes

2004.....	28,585	-	-	-	2	16	28	100	202	401	916	1,505
2003.....	28,345	-	1	-	-	9	44	90	215	409	897	1,446
2002.....	26,551	-	-	-	2	4	26	81	216	402	811	1,266
2001.....	24,689	-	-	1	-	6	18	65	181	352	739	1,050
2000.....	23,761	-	1	-	2	5	27	69	144	323	658	952
1999.....	22,702	1	-	-	-	6	27	50	160	347	617	889

All races, male

2004.....	12,929	-	-	-	-	14	18	71	152	296	600	1,007
2003.....	12,536	-	1	-	-	6	34	64	144	285	592	962
2002.....	11,660	-	-	-	1	4	22	64	151	289	560	844
2001.....	10,451	-	-	-	-	4	13	40	129	234	506	715
2000.....	9,876	-	-	-	1	3	21	43	96	210	433	625
1999.....	9,316	-	-	-	-	6	19	41	108	232	428	586

All races, female

2004.....	15,656	-	-	-	2	2	10	29	50	105	316	498
2003.....	15,809	-	-	-	-	3	10	26	71	124	305	484
2002.....	14,891	-	-	-	1	-	4	17	65	113	251	422
2001.....	14,238	-	-	1	-	2	5	25	52	118	233	335
2000.....	13,885	-	1	-	1	2	6	26	48	113	225	327
1999.....	13,386	1	-	-	-	-	8	9	52	115	189	303

White, both sexes

2004.....	21,156	-	-	-	1	8	15	42	102	223	534	878
2003.....	20,904	-	-	-	-	4	23	49	108	231	503	785
2002.....	19,530	-	-	-	1	1	12	39	102	222	458	753
2001.....	18,232	-	-	-	-	4	6	36	89	197	389	572
2000.....	17,425	-	-	-	2	1	14	31	64	174	368	521
1999.....	16,646	1	-	-	-	2	10	24	85	170	321	469

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	-------------	-----------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Hypertensive heart disease (I11)

White, male

2004.....	9,218	-	-	-	-	7	11	38	82	172	385	635
2003.....	8,948	-	-	-	-	2	19	41	80	179	377	572
2002.....	8,234	-	-	-	1	1	11	32	76	164	347	539
2001.....	7,395	-	-	-	-	3	4	24	72	149	292	412
2000.....	6,927	-	-	-	1	1	11	22	47	126	259	383
1999.....	6,499	-	-	-	-	2	7	22	69	125	245	340

White, female

2004.....	11,938	-	-	-	1	1	4	4	20	51	149	243
2003.....	11,956	-	-	-	-	2	4	8	28	52	126	213
2002.....	11,296	-	-	-	-	-	1	7	26	58	111	214
2001.....	10,837	-	-	-	-	1	2	12	17	48	97	160
2000.....	10,498	-	-	-	1	-	3	9	17	48	109	138
1999.....	10,147	1	-	-	-	-	3	2	16	45	76	129

Black, both sexes

2004.....	6,796	-	-	-	1	8	13	58	90	169	361	595
2003.....	6,843	-	1	-	-	5	20	40	97	170	374	629
2002.....	6,480	-	-	-	1	3	12	40	110	168	332	493
2001.....	5,970	-	-	1	-	2	12	28	87	149	330	459
2000.....	5,886	-	1	-	-	4	13	37	77	145	279	420
1999.....	5,619	-	-	-	-	3	17	26	72	169	281	409

Black, male

2004.....	3,391	-	-	-	-	7	7	33	60	116	200	349
2003.....	3,298	-	1	-	-	4	15	22	57	98	204	365
2002.....	3,146	-	-	-	-	3	9	30	71	116	199	289
2001.....	2,813	-	-	-	-	1	9	15	53	81	199	287
2000.....	2,722	-	-	-	-	2	10	20	46	81	167	235
1999.....	2,606	-	-	-	-	3	12	19	36	101	169	239

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Hypertensive heart disease (I11)

Black, female

2004.....	3,405	-	-	-	1	1	6	25	30	53	161	246
2003.....	3,545	-	-	-	-	1	5	18	40	72	170	264
2002.....	3,334	-	-	-	1	-	3	10	39	52	133	204
2001.....	3,157	-	-	1	-	1	3	13	34	68	131	172
2000.....	3,164	-	1	-	-	2	3	17	31	64	112	185
1999.....	3,013	-	-	-	-	-	5	7	36	68	112	170

American Indian 1/, both sexes

2004.....	109	-	-	-	-	-	-	-	3	3	5	6
2003.....	112	-	-	-	-	-	-	1	5	3	5	9
2002.....	68	-	-	-	-	-	2	-	-	6	5	4
2001.....	64	-	-	-	-	-	-	-	2	1	5	2
2000.....	59	-	-	-	-	-	-	-	-	-	-	1
1999.....	56	-	-	-	-	-	-	-	-	3	3	-

American Indian 1/, male

2004.....	59	-	-	-	-	-	-	-	3	2	3	4
2003.....	54	-	-	-	-	-	-	1	3	3	2	6
2002.....	32	-	-	-	-	-	2	-	-	4	2	4
2001.....	29	-	-	-	-	-	-	-	1	1	2	2
2000.....	32	-	-	-	-	-	-	-	-	-	-	1
1999.....	26	-	-	-	-	-	-	-	-	2	3	-

American Indian 1/, female

2004.....	50	-	-	-	-	-	-	-	-	1	2	2
2003.....	58	-	-	-	-	-	-	-	2	-	3	3
2002.....	36	-	-	-	-	-	-	-	-	2	3	-
2001.....	35	-	-	-	-	-	-	-	1	-	3	-
2000.....	27	-	-	-	-	-	-	-	-	-	-	-
1999.....	30	-	-	-	-	-	-	-	-	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart disease (I11)												
Asian or Pacific Islander 2/, both sexes												
2004.....	524	-	-	-	-	-	-	-	7	6	16	26
2003.....	486	-	-	-	-	-	1	-	5	5	15	23
2002.....	473	-	-	-	-	-	-	2	4	6	16	16
2001.....	423	-	-	-	-	-	-	1	3	5	15	17
2000.....	391	-	-	-	-	-	-	1	3	4	11	10
1999.....	381	-	-	-	-	1	-	-	3	5	12	11
Asian or Pacific Islander 2/, male												
2004.....	261	-	-	-	-	-	-	-	7	6	12	19
2003.....	236	-	-	-	-	-	-	-	4	5	9	19
2002.....	248	-	-	-	-	-	-	2	4	5	12	12
2001.....	214	-	-	-	-	-	-	1	3	3	13	14
2000.....	195	-	-	-	-	-	-	1	3	3	7	6
1999.....	185	-	-	-	-	1	-	-	3	4	11	7
Asian or Pacific Islander 2/, female												
2004.....	263	-	-	-	-	-	-	-	-	-	4	7
2003.....	250	-	-	-	-	-	1	-	1	-	6	4
2002.....	225	-	-	-	-	-	-	-	-	1	4	4
2001.....	209	-	-	-	-	-	-	-	-	2	2	3
2000.....	196	-	-	-	-	-	-	-	-	1	4	4
1999.....	196	-	-	-	-	-	-	-	-	1	1	4

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Hypertensive renal disease (I12)

All races, both sexes

2004.....	9,620	-	-	-	1	4	10	31	46	87	111	212
2003.....	9,027	1	-	-	2	6	12	19	48	71	124	193
2002.....	8,603	1	-	-	-	3	9	28	34	80	108	163
2001.....	8,173	-	1	-	1	4	14	21	49	54	128	183
2000.....	7,777	-	-	1	-	1	8	11	32	68	123	141
1999.....	7,051	-	2	-	1	-	7	18	31	63	119	152

All races, male

2004.....	4,178	-	-	-	-	3	6	19	26	55	66	127
2003.....	3,877	1	-	-	1	4	5	6	35	42	76	128
2002.....	3,682	-	-	-	-	-	5	16	17	47	67	97
2001.....	3,408	-	-	-	-	1	12	12	24	28	86	113
2000.....	3,264	-	-	-	-	-	3	6	14	38	83	77
1999.....	3,031	-	1	-	1	-	4	10	21	43	70	87

All races, female

2004.....	5,442	-	-	-	1	1	4	12	20	32	45	85
2003.....	5,150	-	-	-	1	2	7	13	13	29	48	65
2002.....	4,921	1	-	-	-	3	4	12	17	33	41	66
2001.....	4,765	-	1	-	1	3	2	9	25	26	42	70
2000.....	4,513	-	-	1	-	1	5	5	18	30	40	64
1999.....	4,020	-	1	-	-	-	3	8	10	20	49	65

White, both sexes

2004.....	6,673	-	-	-	-	3	6	10	24	26	44	81
2003.....	6,394	-	-	-	1	4	6	7	17	24	50	76
2002.....	6,046	1	-	-	-	3	5	11	14	23	38	59
2001.....	5,736	-	1	-	1	3	8	6	19	17	43	68
2000.....	5,400	-	-	-	-	-	4	5	13	21	41	40
1999.....	4,783	-	-	-	-	-	3	10	14	18	46	55

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Hypertensive renal disease (I12)

White, male

2004.....	2,887	-	-	-	-	3	4	8	16	19	28	46
2003.....	2,730	-	-	-	-	2	3	2	10	17	34	53
2002.....	2,580	-	-	-	-	-	3	7	7	11	26	33
2001.....	2,365	-	-	-	-	1	6	4	9	8	32	39
2000.....	2,284	-	-	-	-	-	2	4	9	14	29	24
1999.....	2,038	-	-	-	-	-	1	5	11	14	30	32

White, female

2004.....	3,786	-	-	-	-	-	2	2	8	7	16	35
2003.....	3,664	-	-	-	1	2	3	5	7	7	16	23
2002.....	3,466	1	-	-	-	3	2	4	7	12	12	26
2001.....	3,371	-	1	-	1	2	2	2	10	9	11	29
2000.....	3,116	-	-	-	-	-	2	1	4	7	12	16
1999.....	2,745	-	-	-	-	-	2	5	3	4	16	23

Black, both sexes

2004.....	2,639	-	-	-	1	1	4	17	22	57	66	123
2003.....	2,375	1	-	-	-	2	6	12	29	42	71	116
2002.....	2,322	-	-	-	-	-	4	14	18	56	67	101
2001.....	2,203	-	-	-	-	1	6	14	28	36	80	108
2000.....	2,171	-	-	1	-	1	4	6	18	46	78	95
1999.....	2,065	-	1	-	1	-	4	7	15	43	71	91

Black, male

2004.....	1,157	-	-	-	-	-	2	10	10	33	37	75
2003.....	1,042	1	-	-	-	2	2	4	23	23	41	75
2002.....	994	-	-	-	-	-	2	8	9	35	39	61
2001.....	947	-	-	-	-	-	6	8	14	19	51	71
2000.....	892	-	-	-	-	-	1	2	5	23	52	50
1999.....	904	-	1	-	1	-	3	4	8	27	38	55

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart and renal disease (I13)												
All races, both sexes												
2004.....	3,046	-	-	1	-	2	5	8	15	38	58	104
2003.....	3,108	-	1	-	-	-	5	8	20	35	58	87
2002.....	2,895	-	-	-	-	1	4	11	17	28	49	75
2001.....	2,826	1	-	-	-	2	5	10	18	40	59	64
2000.....	2,785	-	-	-	1	1	5	11	17	22	61	68
1999.....	3,327	-	-	2	-	2	4	5	16	43	71	80
All races, male												
2004.....	1,301	-	-	-	-	-	4	7	7	29	40	60
2003.....	1,326	-	1	-	-	-	4	4	9	26	36	63
2002.....	1,205	-	-	-	-	-	3	9	10	19	25	49
2001.....	1,169	1	-	-	-	-	4	4	13	20	43	46
2000.....	1,163	-	-	-	-	1	3	7	11	12	37	43
1999.....	1,390	-	-	1	-	1	2	3	13	24	43	50
All races, female												
2004.....	1,745	-	-	1	-	2	1	1	8	9	18	44
2003.....	1,782	-	-	-	-	-	1	4	11	9	22	24
2002.....	1,690	-	-	-	-	1	1	2	7	9	24	26
2001.....	1,657	-	-	-	-	2	1	6	5	20	16	18
2000.....	1,622	-	-	-	1	-	2	4	6	10	24	25
1999.....	1,937	-	-	1	-	1	2	2	3	19	28	30
White, both sexes												
2004.....	2,071	-	-	1	-	-	2	3	5	7	20	26
2003.....	2,103	-	1	-	-	-	2	4	3	8	13	25
2002.....	1,925	-	-	-	-	1	1	6	5	5	16	19
2001.....	1,922	-	-	-	-	-	3	2	5	7	18	17
2000.....	1,904	-	-	-	-	-	1	6	2	9	28	21
1999.....	2,235	-	-	1	-	-	-	1	6	14	14	21

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Hypertensive heart and renal disease (I13)

White, male

2004.....	846	-	-	-	-	-	2	3	4	5	12	17
2003.....	865	-	1	-	-	-	1	2	-	8	6	19
2002.....	760	-	-	-	-	-	1	4	2	5	10	13
2001.....	762	-	-	-	-	-	3	1	2	3	14	11
2000.....	775	-	-	-	-	-	1	4	2	5	17	12
1999.....	906	-	-	1	-	-	-	1	5	7	10	11

White, female

2004.....	1,225	-	-	1	-	-	-	-	1	2	8	9
2003.....	1,238	-	-	-	-	-	1	2	3	-	7	6
2002.....	1,165	-	-	-	-	1	-	2	3	-	6	6
2001.....	1,160	-	-	-	-	-	-	1	3	4	4	6
2000.....	1,129	-	-	-	-	-	-	2	-	4	11	9
1999.....	1,329	-	-	-	-	-	-	-	1	7	4	10

Black, both sexes

2004.....	883	-	-	-	-	2	3	5	10	30	37	75
2003.....	913	-	-	-	-	-	3	4	16	27	44	60
2002.....	903	-	-	-	-	-	2	5	12	22	32	56
2001.....	832	1	-	-	-	2	2	8	13	29	39	46
2000.....	809	-	-	-	1	1	4	4	15	13	33	45
1999.....	1,000	-	-	1	-	2	4	3	10	28	57	57

Black, male

2004.....	407	-	-	-	-	-	2	4	3	24	27	41
2003.....	413	-	-	-	-	-	3	2	8	18	29	42
2002.....	405	-	-	-	-	-	1	5	8	13	14	36
2001.....	376	1	-	-	-	-	1	3	11	15	28	34
2000.....	356	-	-	-	-	1	2	3	9	7	20	30
1999.....	434	-	-	-	-	1	2	1	8	16	33	37

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Ischemic heart diseases (I20-I25)												
All races, both sexes												
2004.....	451,326	13	10	8	9	24	101	292	800	1,977	5,155	9,731
2003.....	480,028	22	12	12	12	32	128	274	808	2,152	5,395	9,989
2002.....	494,382	23	7	4	16	30	103	251	745	2,211	5,472	9,892
2001.....	502,189	26	9	4	16	31	132	279	837	2,212	5,322	9,461
2000.....	515,204	30	10	8	8	33	90	236	699	2,233	5,286	9,638
1999.....	529,659	27	5	13	9	52	85	245	769	2,387	5,364	9,524
All races, male												
2004.....	233,538	8	1	4	5	20	76	206	616	1,467	3,899	7,442
2003.....	246,342	13	6	4	6	21	86	197	616	1,630	4,049	7,661
2002.....	252,760	15	4	1	12	18	74	191	550	1,670	4,170	7,582
2001.....	254,005	17	3	2	11	24	95	211	619	1,667	4,045	7,323
2000.....	260,574	17	7	5	4	21	60	181	523	1,665	4,122	7,601
1999.....	267,268	13	3	10	7	36	57	169	570	1,801	4,169	7,447
All races, female												
2004.....	217,788	5	9	4	4	4	25	86	184	510	1,256	2,289
2003.....	233,686	9	6	8	6	11	42	77	192	522	1,346	2,328
2002.....	241,622	8	3	3	4	12	29	60	195	541	1,302	2,310
2001.....	248,184	9	6	2	5	7	37	68	218	545	1,277	2,138
2000.....	254,630	13	3	3	4	12	30	55	176	568	1,164	2,037
1999.....	262,391	14	2	3	2	16	28	76	199	586	1,195	2,077
White, both sexes												
2004.....	395,705	9	7	5	6	14	58	199	585	1,555	4,055	7,657
2003.....	421,482	16	6	7	8	25	87	192	604	1,680	4,206	7,811
2002.....	435,170	14	5	3	12	24	70	185	555	1,700	4,310	7,782
2001.....	442,536	19	6	3	10	22	82	197	623	1,714	4,128	7,426
2000.....	455,400	23	10	6	7	28	64	160	516	1,745	4,106	7,570
1999.....	469,149	18	4	9	5	36	49	180	573	1,855	4,133	7,541

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Ischemic heart diseases (I20-I25)												
White, male												
2004.....	205,475	6	1	3	4	12	45	148	459	1,167	3,178	6,022
2003.....	216,751	9	3	2	5	17	59	143	465	1,304	3,238	6,156
2002.....	223,262	10	2	1	9	14	52	149	424	1,306	3,390	6,172
2001.....	224,464	14	2	2	7	16	62	156	468	1,316	3,244	5,958
2000.....	230,951	14	7	3	4	18	43	124	405	1,336	3,326	6,194
1999.....	237,193	8	3	7	5	25	38	124	431	1,425	3,317	6,089
White, female												
2004.....	190,230	3	6	2	2	2	13	51	126	388	877	1,635
2003.....	204,731	7	3	5	3	8	28	49	139	376	968	1,655
2002.....	211,908	4	3	2	3	10	18	36	131	394	920	1,610
2001.....	218,072	5	4	1	3	6	20	41	155	398	884	1,468
2000.....	224,449	9	3	3	3	10	21	36	111	409	780	1,376
1999.....	231,956	10	1	2	-	11	11	56	142	430	816	1,452
Black, both sexes												
2004.....	46,695	3	1	3	3	8	38	79	180	354	978	1,821
2003.....	49,274	4	5	4	4	7	36	64	170	399	1,026	1,884
2002.....	50,174	8	2	1	4	4	27	55	158	438	1,022	1,858
2001.....	50,933	7	3	1	5	9	44	73	177	437	1,049	1,787
2000.....	51,265	4	-	2	-	4	20	66	166	432	1,041	1,845
1999.....	51,995	8	1	4	4	14	33	53	160	474	1,096	1,761
Black, male												
2004.....	23,060	2	-	1	1	7	29	45	126	242	619	1,215
2003.....	24,273	3	3	2	1	4	24	40	122	262	686	1,266
2002.....	24,322	4	2	-	3	2	16	34	99	300	659	1,204
2001.....	24,593	3	1	-	3	8	28	49	122	298	685	1,165
2000.....	24,625	2	-	2	-	2	12	48	104	281	682	1,223
1999.....	25,104	4	-	3	2	9	17	38	110	323	734	1,171

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Ischemic heart diseases (I20-I25)

Black, female

2004.....	23,635	1	1	2	2	1	9	34	54	112	359	606
2003.....	25,001	1	2	2	3	3	12	24	48	137	340	618
2002.....	25,852	4	-	1	1	2	11	21	59	138	363	654
2001.....	26,340	4	2	1	2	1	16	24	55	139	364	622
2000.....	26,640	2	-	-	-	2	8	18	62	151	359	622
1999.....	26,891	4	1	1	2	5	16	15	50	151	362	590

American Indian 1/, both sexes

2004.....	1,843	1	1	-	-	-	1	9	12	21	47	81
2003.....	1,905	-	-	-	-	-	2	2	14	25	57	95
2002.....	1,777	1	-	-	-	1	3	6	10	15	44	61
2001.....	1,710	-	-	-	-	-	3	1	10	18	46	77
2000.....	1,740	1	-	-	-	-	4	4	7	18	55	75
1999.....	1,720	1	-	-	-	-	1	5	11	17	39	65

American Indian 1/, male

2004.....	1,058	-	-	-	-	-	1	8	10	16	35	57
2003.....	1,133	-	-	-	-	-	1	1	10	22	41	70
2002.....	1,041	1	-	-	-	1	3	4	9	13	35	49
2001.....	995	-	-	-	-	-	3	-	8	12	34	58
2000.....	1,008	-	-	-	-	-	3	3	6	14	42	61
1999.....	977	1	-	-	-	-	1	2	6	13	32	52

American Indian 1/, female

2004.....	785	1	1	-	-	-	-	1	2	5	12	24
2003.....	772	-	-	-	-	-	1	1	4	3	16	25
2002.....	736	-	-	-	-	-	-	2	1	2	9	12
2001.....	715	-	-	-	-	-	-	1	2	6	12	19
2000.....	732	1	-	-	-	-	1	1	1	4	13	14
1999.....	743	-	-	-	-	-	-	3	5	4	7	13

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Ischemic heart diseases (I20-I25)												
Asian or Pacific Islander 2/, both sexes												
2004.....	7,083	-	1	-	-	2	4	5	23	47	75	172
2003.....	7,367	2	1	1	-	-	3	16	20	48	106	199
2002.....	7,261	-	-	-	-	1	3	5	22	58	96	191
2001.....	7,010	-	-	-	1	-	3	8	27	43	99	171
2000.....	6,799	2	-	-	1	1	2	6	10	38	84	148
1999.....	6,795	-	-	-	-	2	2	7	25	41	96	157
Asian or Pacific Islander 2/, male												
2004.....	3,945	-	-	-	-	1	1	5	21	42	67	148
2003.....	4,185	1	-	-	-	-	2	13	19	42	84	169
2002.....	4,135	-	-	-	-	1	3	4	18	51	86	157
2001.....	3,953	-	-	-	1	-	2	6	21	41	82	142
2000.....	3,990	1	-	-	-	1	2	6	8	34	72	123
1999.....	3,994	-	-	-	-	2	1	5	23	40	86	135
Asian or Pacific Islander 2/, female												
2004.....	3,138	-	1	-	-	1	3	-	2	5	8	24
2003.....	3,182	1	1	1	-	-	1	3	1	6	22	30
2002.....	3,126	-	-	-	-	-	-	1	4	7	10	34
2001.....	3,057	-	-	-	-	-	1	2	6	2	17	29
2000.....	2,809	1	-	-	1	-	-	-	2	4	12	25
1999.....	2,801	-	-	-	-	-	1	2	2	1	10	22

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute myocardial infarction (I21-I22)												
All races, both sexes												
2004.....	156,816	9	2	3	2	7	44	120	327	796	2,042	3,917
2003.....	170,564	9	6	6	6	17	56	105	308	881	2,213	4,092
2002.....	179,514	11	5	2	5	15	47	91	309	882	2,279	4,055
2001.....	184,757	15	2	1	6	15	49	105	370	924	2,252	4,097
2000.....	192,898	20	1	5	1	14	38	108	267	904	2,228	4,230
1999.....	199,454	13	2	5	6	25	35	101	333	935	2,314	4,225
All races, male												
2004.....	82,909	7	-	2	2	5	34	80	242	598	1,515	2,989
2003.....	89,515	4	2	2	4	11	35	75	225	664	1,637	3,130
2002.....	93,830	7	2	-	2	10	31	67	225	656	1,691	3,066
2001.....	95,875	10	2	-	4	14	34	69	264	691	1,693	3,146
2000.....	100,306	10	1	3	1	8	22	75	194	657	1,730	3,319
1999.....	104,169	6	1	3	4	16	24	68	238	707	1,763	3,261
All races, female												
2004.....	73,907	2	2	1	-	2	10	40	85	198	527	928
2003.....	81,049	5	4	4	2	6	21	30	83	217	576	962
2002.....	85,684	4	3	2	3	5	16	24	84	226	588	989
2001.....	88,882	5	-	1	2	1	15	36	106	233	559	951
2000.....	92,592	10	-	2	-	6	16	33	73	247	498	911
1999.....	95,285	7	1	2	2	9	11	33	95	228	551	964
White, both sexes												
2004.....	137,643	6	2	1	2	3	24	82	255	636	1,633	3,161
2003.....	149,915	7	3	5	4	14	39	75	235	705	1,747	3,263
2002.....	157,922	9	5	1	4	13	30	63	238	698	1,826	3,247
2001.....	162,943	12	2	-	2	12	30	70	285	730	1,783	3,321
2000.....	170,584	16	1	3	1	11	28	71	201	744	1,752	3,442
1999.....	177,075	9	2	3	3	19	20	76	249	757	1,839	3,453

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Acute myocardial infarction (I21-I22)

White, male

2004.....	73,448	5	-	1	2	2	19	56	199	479	1,267	2,472
2003.....	79,157	3	1	1	3	10	25	58	171	548	1,328	2,573
2002.....	83,288	6	2	-	2	8	22	50	179	531	1,390	2,543
2001.....	85,280	9	2	-	1	11	23	46	208	556	1,395	2,654
2000.....	89,383	9	1	1	1	6	18	50	155	556	1,413	2,797
1999.....	93,189	3	1	1	3	13	16	51	178	585	1,447	2,757

White, female

2004.....	64,195	1	2	-	-	1	5	26	56	157	366	689
2003.....	70,758	4	2	4	1	4	14	17	64	157	419	690
2002.....	74,634	3	3	1	2	5	8	13	59	167	436	704
2001.....	77,663	3	-	-	1	1	7	24	77	174	388	667
2000.....	81,201	7	-	2	-	5	10	21	46	188	339	645
1999.....	83,886	6	1	2	-	6	4	25	71	172	392	696

Black, both sexes

2004.....	16,206	2	-	2	-	3	18	32	62	138	364	676
2003.....	17,441	1	2	1	2	3	16	24	64	160	420	726
2002.....	18,322	2	-	1	1	-	15	25	64	158	403	707
2001.....	18,581	3	-	1	3	3	18	29	70	169	410	684
2000.....	19,112	2	-	2	-	2	9	32	62	146	423	699
1999.....	19,276	4	-	2	3	6	13	22	66	163	424	688

Black, male

2004.....	7,811	2	-	1	-	3	14	18	36	102	212	456
2003.....	8,472	-	1	1	1	1	10	13	46	101	273	473
2002.....	8,680	1	-	-	-	-	7	15	40	103	259	440
2001.....	8,717	1	-	-	2	3	10	19	45	110	250	416
2000.....	9,045	1	-	2	-	1	4	21	35	90	272	449
1999.....	9,186	3	-	2	1	3	7	16	45	107	272	436

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Acute myocardial infarction (I21-I22)

Black, female

2004.....	8,395	-	-	1	-	-	4	14	26	36	152	220
2003.....	8,969	1	1	-	1	2	6	11	18	59	147	253
2002.....	9,642	1	-	1	1	-	8	10	24	55	144	267
2001.....	9,864	2	-	1	1	-	8	10	25	59	160	268
2000.....	10,067	1	-	-	-	1	5	11	27	56	151	250
1999.....	10,090	1	-	-	2	3	6	6	21	56	152	252

American Indian 1/, both sexes

2004.....	634	1	-	-	-	-	-	3	4	8	18	33
2003.....	706	-	-	-	-	-	-	2	3	7	20	40
2002.....	657	-	-	-	-	1	1	-	3	6	15	32
2001.....	660	-	-	-	-	-	-	1	6	9	21	30
2000.....	656	1	-	-	-	-	1	2	3	7	23	30
1999.....	674	-	-	-	-	-	-	2	6	3	19	26

American Indian 1/, male

2004.....	359	-	-	-	-	-	-	3	3	5	13	23
2003.....	430	-	-	-	-	-	-	1	2	6	16	32
2002.....	384	-	-	-	-	1	1	-	3	5	12	26
2001.....	404	-	-	-	-	-	-	-	5	9	17	24
2000.....	384	-	-	-	-	-	-	1	3	5	18	25
1999.....	382	-	-	-	-	-	-	-	4	3	17	20

American Indian 1/, female

2004.....	275	1	-	-	-	-	-	-	1	3	5	10
2003.....	276	-	-	-	-	-	-	1	1	1	4	8
2002.....	273	-	-	-	-	-	-	-	-	1	3	6
2001.....	256	-	-	-	-	-	-	1	1	-	4	6
2000.....	272	1	-	-	-	-	1	1	-	2	5	5
1999.....	292	-	-	-	-	-	-	2	2	-	2	6

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute myocardial infarction (I21-I22)												
Asian or Pacific Islander 2/, both sexes												
2004.....	2,333	-	-	-	-	1	2	3	6	14	27	47
2003.....	2,502	1	1	-	-	-	1	4	6	9	26	63
2002.....	2,613	-	-	-	-	1	1	3	4	20	35	69
2001.....	2,573	-	-	-	1	-	1	5	9	16	38	62
2000.....	2,546	1	-	-	-	1	-	3	1	7	30	59
1999.....	2,429	-	-	-	-	-	2	1	12	12	32	58
Asian or Pacific Islander 2/, male												
2004.....	1,291	-	-	-	-	-	1	3	4	12	23	38
2003.....	1,456	1	-	-	-	-	-	3	6	9	20	52
2002.....	1,478	-	-	-	-	1	1	2	3	17	30	57
2001.....	1,474	-	-	-	1	-	1	4	6	16	31	52
2000.....	1,494	-	-	-	-	1	-	3	1	6	27	48
1999.....	1,412	-	-	-	-	-	1	1	11	12	27	48
Asian or Pacific Islander 2/, female												
2004.....	1,042	-	-	-	-	1	1	-	2	2	4	9
2003.....	1,046	-	1	-	-	-	1	1	-	-	6	11
2002.....	1,135	-	-	-	-	-	-	1	1	3	5	12
2001.....	1,099	-	-	-	-	-	-	1	3	-	7	10
2000.....	1,052	1	-	-	-	-	-	-	-	1	3	11
1999.....	1,017	-	-	-	-	-	1	-	1	-	5	10

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other acute ischemic heart diseases (I24)

All races, both sexes

2004.....	2,462	1	2	1	-	1	4	2	8	15	65	81
2003.....	3,158	4	1	1	-	-	5	3	8	27	71	106
2002.....	3,407	2	-	1	-	2	2	4	14	17	60	119
2001.....	3,351	3	-	-	2	-	7	2	12	32	62	120
2000.....	3,363	1	1	-	-	4	4	4	12	30	62	93
1999.....	3,232	3	1	1	-	-	3	2	4	25	56	92

All races, male

2004.....	1,301	-	-	-	-	1	3	1	5	13	46	55
2003.....	1,621	2	1	-	-	-	5	2	5	20	53	75
2002.....	1,767	1	-	1	-	-	1	2	10	9	41	88
2001.....	1,720	2	-	-	1	-	6	2	8	18	47	89
2000.....	1,749	1	1	-	-	3	3	3	10	24	47	76
1999.....	1,665	1	1	1	-	-	3	1	3	19	44	69

All races, female

2004.....	1,161	1	2	1	-	-	1	1	3	2	19	26
2003.....	1,537	2	-	1	-	-	-	1	3	7	18	31
2002.....	1,640	1	-	-	-	2	1	2	4	8	19	31
2001.....	1,631	1	-	-	1	-	1	-	4	14	15	31
2000.....	1,614	-	-	-	-	1	1	1	2	6	15	17
1999.....	1,567	2	-	-	-	-	-	1	1	6	12	23

White, both sexes

2004.....	2,025	-	2	1	-	1	2	-	3	12	48	52
2003.....	2,663	2	-	1	-	-	4	2	4	20	49	73
2002.....	2,907	1	-	1	-	2	1	3	9	15	40	90
2001.....	2,821	3	-	-	2	-	4	2	9	23	40	89
2000.....	2,839	1	1	-	-	4	2	4	8	23	45	69
1999.....	2,752	1	1	1	-	-	3	2	3	13	40	56

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute ischemic heart diseases (I24)												
White, male												
2004.....	1,042	-	-	-	-	1	1	-	3	11	35	37
2003.....	1,338	1	-	-	-	-	4	2	3	16	40	51
2002.....	1,494	1	-	1	-	-	1	2	6	8	34	64
2001.....	1,431	2	-	-	1	-	3	2	7	14	32	69
2000.....	1,453	1	1	-	-	3	1	3	6	19	35	61
1999.....	1,389	1	1	1	-	-	3	1	3	10	31	42
White, female												
2004.....	983	-	2	1	-	-	1	-	-	1	13	15
2003.....	1,325	1	-	1	-	-	-	-	1	4	9	22
2002.....	1,413	-	-	-	-	2	-	1	3	7	6	26
2001.....	1,390	1	-	-	1	-	1	-	2	9	8	20
2000.....	1,386	-	-	-	-	1	1	1	2	4	10	8
1999.....	1,363	-	-	-	-	-	-	1	-	3	9	14
Black, both sexes												
2004.....	395	1	-	-	-	-	1	2	5	2	16	27
2003.....	436	1	1	-	-	-	1	1	2	7	20	30
2002.....	446	1	-	-	-	-	1	1	5	2	18	27
2001.....	454	-	-	-	-	-	3	-	2	8	21	27
2000.....	456	-	-	-	-	-	1	-	4	7	15	23
1999.....	431	2	-	-	-	-	-	-	1	12	14	34
Black, male												
2004.....	235	-	-	-	-	-	1	1	2	2	10	16
2003.....	256	1	1	-	-	-	1	-	1	4	12	21
2002.....	243	-	-	-	-	-	-	-	4	1	5	23
2001.....	249	-	-	-	-	-	3	-	-	4	15	17
2000.....	258	-	-	-	-	-	1	-	4	5	11	15
1999.....	248	-	-	-	-	-	-	-	-	9	11	26

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other forms of chronic ischemic heart disease (I20,I25)

All races, both sexes

2004.....	292,048	3	6	4	7	16	53	170	465	1,166	3,048	5,733
2003.....	306,306	9	5	5	6	15	67	166	492	1,244	3,111	5,791
2002.....	311,461	10	2	1	11	13	54	156	422	1,312	3,133	5,718
2001.....	314,081	8	7	3	8	16	76	172	455	1,256	3,008	5,244
2000.....	318,943	9	8	3	7	15	48	124	420	1,299	2,996	5,315
1999.....	326,973	11	2	7	3	27	47	142	432	1,427	2,994	5,207

All races, male

2004.....	149,328	1	1	2	3	14	39	125	369	856	2,338	4,398
2003.....	155,206	7	3	2	2	10	46	120	386	946	2,359	4,456
2002.....	157,163	7	2	-	10	8	42	122	315	1,005	2,438	4,428
2001.....	156,410	5	1	2	6	10	55	140	347	958	2,305	4,088
2000.....	158,519	6	5	2	3	10	35	103	319	984	2,345	4,206
1999.....	161,434	6	1	6	3	20	30	100	329	1,075	2,362	4,117

All races, female

2004.....	142,720	2	5	2	4	2	14	45	96	310	710	1,335
2003.....	151,100	2	2	3	4	5	21	46	106	298	752	1,335
2002.....	154,298	3	-	1	1	5	12	34	107	307	695	1,290
2001.....	157,671	3	6	1	2	6	21	32	108	298	703	1,156
2000.....	160,424	3	3	1	4	5	13	21	101	315	651	1,109
1999.....	165,539	5	1	1	-	7	17	42	103	352	632	1,090

White, both sexes

2004.....	256,037	3	3	3	4	10	32	117	327	907	2,374	4,444
2003.....	268,904	7	3	1	4	11	44	115	365	955	2,410	4,475
2002.....	274,341	4	-	1	8	9	39	119	308	987	2,444	4,445
2001.....	276,772	4	4	3	6	10	48	125	329	961	2,305	4,016
2000.....	281,977	6	8	3	6	13	34	85	307	978	2,309	4,059
1999.....	289,322	8	1	5	2	17	26	102	321	1,085	2,254	4,032

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other forms of chronic ischemic heart disease (I20,I25)

White, male

2004.....	130,985	1	1	2	2	9	25	92	257	677	1,876	3,513
2003.....	136,256	5	2	1	2	7	30	83	291	740	1,870	3,532
2002.....	138,480	3	-	-	7	6	29	97	239	767	1,966	3,565
2001.....	137,753	3	-	2	5	5	36	108	253	746	1,817	3,235
2000.....	140,115	4	5	2	3	9	24	71	244	761	1,878	3,336
1999.....	142,615	4	1	5	2	12	19	72	250	830	1,839	3,290

White, female

2004.....	125,052	2	2	1	2	1	7	25	70	230	498	931
2003.....	132,648	2	1	-	2	4	14	32	74	215	540	943
2002.....	135,861	1	-	1	1	3	10	22	69	220	478	880
2001.....	139,019	1	4	1	1	5	12	17	76	215	488	781
2000.....	141,862	2	3	1	3	4	10	14	63	217	431	723
1999.....	146,707	4	-	-	-	5	7	30	71	255	415	742

Black, both sexes

2004.....	30,094	-	1	1	3	5	19	45	113	214	598	1,118
2003.....	31,397	2	2	3	2	4	19	39	104	232	586	1,128
2002.....	31,406	5	2	-	3	4	11	29	89	278	601	1,124
2001.....	31,898	4	3	-	2	6	23	44	105	260	618	1,076
2000.....	31,697	2	-	-	-	2	10	34	100	279	603	1,123
1999.....	32,288	2	1	2	1	8	20	31	93	299	658	1,039

Black, male

2004.....	15,014	-	-	-	1	4	14	26	88	138	397	743
2003.....	15,545	2	1	1	-	3	13	27	75	157	401	772
2002.....	15,399	3	2	-	3	2	9	19	55	196	395	741
2001.....	15,627	2	1	-	1	5	15	30	77	184	420	732
2000.....	15,322	1	-	-	-	1	7	27	65	186	399	759
1999.....	15,670	1	-	1	1	6	10	22	65	207	451	709

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other forms of chronic ischemic heart disease (I20,I25)

Black, female

2004.....	15,080	-	1	1	2	1	5	19	25	76	201	375
2003.....	15,852	-	1	2	2	1	6	12	29	75	185	356
2002.....	16,007	2	-	-	-	2	2	10	34	82	206	383
2001.....	16,271	2	2	-	1	1	8	14	28	76	198	344
2000.....	16,375	1	-	-	-	1	3	7	35	93	204	364
1999.....	16,618	1	1	1	-	2	10	9	28	92	207	330

American Indian 1/, both sexes

2004.....	1,186	-	1	-	-	-	-	6	8	12	28	47
2003.....	1,162	-	-	-	-	-	2	-	10	18	35	53
2002.....	1,084	1	-	-	-	-	2	6	7	9	27	27
2001.....	1,003	-	-	-	-	-	3	-	3	8	25	44
2000.....	1,041	-	-	-	-	-	2	2	4	11	30	44
1999.....	1,009	1	-	-	-	-	1	3	5	14	19	37

American Indian 1/, male

2004.....	688	-	-	-	-	-	-	5	7	11	21	33
2003.....	688	-	-	-	-	-	1	-	8	16	24	36
2002.....	636	1	-	-	-	-	2	4	6	8	21	22
2001.....	563	-	-	-	-	-	3	-	2	3	17	32
2000.....	598	-	-	-	-	-	2	2	3	9	23	36
1999.....	575	1	-	-	-	-	1	2	2	10	14	31

American Indian 1/, female

2004.....	498	-	1	-	-	-	-	1	1	1	7	14
2003.....	474	-	-	-	-	-	1	-	2	2	11	17
2002.....	448	-	-	-	-	-	-	2	1	1	6	5
2001.....	440	-	-	-	-	-	-	-	1	5	8	12
2000.....	443	-	-	-	-	-	-	-	1	2	7	8
1999.....	434	-	-	-	-	-	-	1	3	4	5	6

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other forms of chronic ischemic heart disease (I20,I25)

Asian or Pacific Islander 2/, both sexes

2004.....	4,731	-	1	-	-	1	2	2	17	33	48	124
2003.....	4,843	-	-	1	-	-	2	12	13	39	80	135
2002.....	4,630	-	-	-	-	-	2	2	18	38	61	122
2001.....	4,408	-	-	-	-	-	2	3	18	27	60	108
2000.....	4,228	1	-	-	1	-	2	3	9	31	54	89
1999.....	4,354	-	-	-	-	2	-	6	13	29	63	99

Asian or Pacific Islander 2/, male

2004.....	2,641	-	-	-	-	1	-	2	17	30	44	109
2003.....	2,717	-	-	-	-	-	2	10	12	33	64	116
2002.....	2,648	-	-	-	-	-	2	2	15	34	56	100
2001.....	2,467	-	-	-	-	-	1	2	15	25	51	89
2000.....	2,484	1	-	-	-	-	2	3	7	28	45	75
1999.....	2,574	-	-	-	-	2	-	4	12	28	58	87

Asian or Pacific Islander 2/, female

2004.....	2,090	-	1	-	-	-	2	-	-	3	4	15
2003.....	2,126	-	-	1	-	-	-	2	1	6	16	19
2002.....	1,982	-	-	-	-	-	-	-	3	4	5	22
2001.....	1,941	-	-	-	-	-	1	1	3	2	9	19
2000.....	1,744	-	-	-	1	-	-	-	2	3	9	14
1999.....	1,780	-	-	-	-	-	-	2	1	1	5	12

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerotic cardiovascular disease, so described (I25.0)												
All races, both sexes												
2004.....	63,661	-	1	-	-	1	8	37	156	351	1,112	2,110
2003.....	67,232	2	-	-	-	2	14	50	161	414	1,074	2,151
2002.....	68,129	2	-	-	1	4	13	39	142	410	1,105	2,081
2001.....	68,103	-	-	-	-	3	17	39	140	402	1,055	1,838
2000.....	69,554	1	-	-	1	-	15	33	111	391	985	1,894
1999.....	71,926	-	1	-	1	-	13	32	128	453	1,035	1,792
All races, male												
2004.....	34,574	-	-	-	-	1	7	30	127	266	877	1,674
2003.....	36,042	2	-	-	-	1	11	40	141	315	842	1,715
2002.....	36,047	1	-	-	-	3	10	31	113	325	891	1,634
2001.....	35,440	-	-	-	-	2	14	30	103	315	815	1,488
2000.....	35,896	-	-	-	1	-	12	26	89	312	787	1,533
1999.....	36,633	-	-	-	1	-	10	21	98	355	837	1,438
All races, female												
2004.....	29,087	-	1	-	-	-	1	7	29	85	235	436
2003.....	31,190	-	-	-	-	1	3	10	20	99	232	436
2002.....	32,082	1	-	-	1	1	3	8	29	85	214	447
2001.....	32,663	-	-	-	-	1	3	9	37	87	240	350
2000.....	33,658	1	-	-	-	-	3	7	22	79	198	361
1999.....	35,293	-	1	-	-	-	3	11	30	98	198	354
White, both sexes												
2004.....	52,585	-	-	-	-	1	6	27	105	278	860	1,579
2003.....	55,760	1	-	-	-	2	10	35	121	324	820	1,609
2002.....	56,569	1	-	-	1	4	11	32	100	296	834	1,602
2001.....	56,414	-	-	-	-	2	10	28	105	304	785	1,370
2000.....	57,794	-	-	-	1	-	13	19	83	291	743	1,393
1999.....	59,711	-	-	-	1	-	5	20	90	336	761	1,305

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Atherosclerotic cardiovascular disease, so described (I25.0)

White, male

2004.....	28,370	-	-	-	-	1	5	23	84	210	703	1,305
2003.....	29,560	1	-	-	-	1	8	27	106	246	653	1,323
2002.....	29,604	1	-	-	-	3	8	26	83	239	696	1,293
2001.....	29,039	-	-	-	-	1	7	23	80	244	620	1,145
2000.....	29,467	-	-	-	1	-	10	14	69	237	615	1,164
1999.....	29,977	-	-	-	1	-	5	14	71	269	640	1,076

White, female

2004.....	24,215	-	-	-	-	-	1	4	21	68	157	274
2003.....	26,200	-	-	-	-	1	2	8	15	78	167	286
2002.....	26,965	-	-	-	1	1	3	6	17	57	138	309
2001.....	27,375	-	-	-	-	1	3	5	25	60	165	225
2000.....	28,327	-	-	-	-	-	3	5	14	54	128	229
1999.....	29,734	-	-	-	-	-	-	6	19	67	121	229

Black, both sexes

2004.....	9,666	-	1	-	-	-	2	8	41	55	222	473
2003.....	9,923	1	-	-	-	-	3	12	35	73	212	459
2002.....	10,195	1	-	-	-	-	2	5	36	100	235	434
2001.....	10,321	-	-	-	-	1	6	11	27	89	236	413
2000.....	10,418	1	-	-	-	-	1	12	24	87	208	459
1999.....	10,823	-	1	-	-	-	8	8	33	110	251	437

Black, male

2004.....	5,331	-	-	-	-	-	2	5	34	39	148	316
2003.....	5,543	1	-	-	-	-	2	10	30	52	155	324
2002.....	5,598	-	-	-	-	-	2	3	25	73	161	302
2001.....	5,578	-	-	-	-	1	6	7	17	64	167	294
2000.....	5,615	-	-	-	-	-	1	10	16	65	145	332
1999.....	5,815	-	-	-	-	-	5	5	22	80	177	317

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerotic cardiovascular disease, so described (I25.0)												
Black, female												
2004.....	4,335	-	1	-	-	-	-	3	7	16	74	157
2003.....	4,380	-	-	-	-	-	1	2	5	21	57	135
2002.....	4,597	1	-	-	-	-	-	2	11	27	74	132
2001.....	4,743	-	-	-	-	-	-	4	10	25	69	119
2000.....	4,803	1	-	-	-	-	-	2	8	22	63	127
1999.....	5,008	-	1	-	-	-	3	3	11	30	74	120
American Indian 1/, both sexes												
2004.....	331	-	-	-	-	-	-	2	3	7	7	16
2003.....	358	-	-	-	-	-	-	-	2	6	12	22
2002.....	318	-	-	-	-	-	-	2	1	2	16	8
2001.....	335	-	-	-	-	-	1	-	2	3	11	15
2000.....	266	-	-	-	-	-	1	1	3	3	9	13
1999.....	295	-	-	-	-	-	-	1	2	3	3	13
American Indian 1/, male												
2004.....	207	-	-	-	-	-	-	2	2	6	5	11
2003.....	227	-	-	-	-	-	-	-	2	6	9	16
2002.....	196	-	-	-	-	-	-	2	-	2	14	7
2001.....	196	-	-	-	-	-	1	-	1	1	8	10
2000.....	159	-	-	-	-	-	1	1	3	2	6	11
1999.....	179	-	-	-	-	-	-	-	2	2	2	12
American Indian 1/, female												
2004.....	124	-	-	-	-	-	-	-	1	1	2	5
2003.....	131	-	-	-	-	-	-	-	-	-	3	6
2002.....	122	-	-	-	-	-	-	-	1	-	2	1
2001.....	139	-	-	-	-	-	-	-	1	2	3	5
2000.....	107	-	-	-	-	-	-	-	-	1	3	2
1999.....	116	-	-	-	-	-	-	1	-	1	1	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerotic cardiovascular disease, so described (I25.0)												
Asian or Pacific Islander 2/, both sexes												
2004.....	1,079	-	-	-	-	-	-	-	7	11	23	42
2003.....	1,191	-	-	-	-	-	1	3	3	11	30	61
2002.....	1,047	-	-	-	-	-	-	-	5	12	20	37
2001.....	1,033	-	-	-	-	-	-	-	6	6	23	40
2000.....	1,076	-	-	-	-	-	-	1	1	10	25	29
1999.....	1,097	-	-	-	-	-	-	3	3	4	20	37
Asian or Pacific Islander 2/, male												
2004.....	666	-	-	-	-	-	-	-	7	11	21	42
2003.....	712	-	-	-	-	-	1	3	3	11	25	52
2002.....	649	-	-	-	-	-	-	-	5	11	20	32
2001.....	627	-	-	-	-	-	-	-	5	6	20	39
2000.....	655	-	-	-	-	-	-	1	1	8	21	26
1999.....	662	-	-	-	-	-	-	2	3	4	18	33
Asian or Pacific Islander 2/, female												
2004.....	413	-	-	-	-	-	-	-	-	-	2	-
2003.....	479	-	-	-	-	-	-	-	-	-	5	9
2002.....	398	-	-	-	-	-	-	-	-	1	-	5
2001.....	406	-	-	-	-	-	-	-	1	-	3	1
2000.....	421	-	-	-	-	-	-	-	-	2	4	3
1999.....	435	-	-	-	-	-	-	1	-	-	2	4

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)

All races, both sexes

2004.....	228,387	3	5	4	7	15	45	133	309	815	1,936	3,623
2003.....	239,074	7	5	5	6	13	53	116	331	830	2,037	3,640
2002.....	243,332	8	2	1	10	9	41	117	280	902	2,028	3,637
2001.....	245,978	8	7	3	8	13	59	133	315	854	1,953	3,406
2000.....	249,389	8	8	3	6	15	33	91	309	908	2,011	3,421
1999.....	255,047	11	1	7	2	27	34	110	304	974	1,959	3,415

All races, male

2004.....	114,754	1	1	2	3	13	32	95	242	590	1,461	2,724
2003.....	119,164	5	3	2	2	9	35	80	245	631	1,517	2,741
2002.....	121,116	6	2	-	10	5	32	91	202	680	1,547	2,794
2001.....	120,970	5	1	2	6	8	41	110	244	643	1,490	2,600
2000.....	122,623	6	5	2	2	10	23	77	230	672	1,558	2,673
1999.....	124,801	6	1	6	2	20	20	79	231	720	1,525	2,679

All races, female

2004.....	113,633	2	4	2	4	2	13	38	67	225	475	899
2003.....	119,910	2	2	3	4	4	18	36	86	199	520	899
2002.....	122,216	2	-	1	-	4	9	26	78	222	481	843
2001.....	125,008	3	6	1	2	5	18	23	71	211	463	806
2000.....	126,766	2	3	1	4	5	10	14	79	236	453	748
1999.....	130,246	5	-	1	-	7	14	31	73	254	434	736

White, both sexes

2004.....	203,452	3	3	3	4	9	26	90	222	629	1,514	2,865
2003.....	213,144	6	3	1	4	9	34	80	244	631	1,590	2,866
2002.....	217,772	3	-	1	7	5	28	87	208	691	1,610	2,843
2001.....	220,358	4	4	3	6	8	38	97	224	657	1,520	2,646
2000.....	224,183	6	8	3	5	13	21	66	224	687	1,566	2,666
1999.....	229,611	8	1	5	1	17	21	82	231	749	1,493	2,727

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	-------------	-----------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)

White, male

2004.....	102,615	1	1	2	2	8	20	69	173	467	1,173	2,208
2003.....	106,696	4	2	1	2	6	22	56	185	494	1,217	2,209
2002.....	108,876	2	-	-	7	3	21	71	156	528	1,270	2,272
2001.....	108,714	3	-	2	5	4	29	85	173	502	1,197	2,090
2000.....	110,648	4	5	2	2	9	14	57	175	524	1,263	2,172
1999.....	112,638	4	1	5	1	12	14	58	179	561	1,199	2,214

White, female

2004.....	100,837	2	2	1	2	1	6	21	49	162	341	657
2003.....	106,448	2	1	-	2	3	12	24	59	137	373	657
2002.....	108,896	1	-	1	-	2	7	16	52	163	340	571
2001.....	111,644	1	4	1	1	4	9	12	51	155	323	556
2000.....	113,535	2	3	1	3	4	7	9	49	163	303	494
1999.....	116,973	4	-	-	-	5	7	24	52	188	294	513

Black, both sexes

2004.....	20,428	-	-	1	3	5	17	37	72	159	376	645
2003.....	21,474	1	2	3	2	4	16	27	69	159	374	669
2002.....	21,211	4	2	-	3	4	9	24	53	178	366	690
2001.....	21,577	4	3	-	2	5	17	33	78	171	382	663
2000.....	21,279	1	-	-	-	2	9	22	76	192	395	664
1999.....	21,465	2	-	2	1	8	12	23	60	189	407	602

Black, male

2004.....	9,683	-	-	-	1	4	12	21	54	99	249	427
2003.....	10,002	1	1	1	-	3	11	17	45	105	246	448
2002.....	9,801	3	2	-	3	2	7	16	30	123	234	439
2001.....	10,049	2	1	-	1	4	9	23	60	120	253	438
2000.....	9,707	1	-	-	-	1	6	17	49	121	254	427
1999.....	9,855	1	-	1	1	6	5	17	43	127	274	392

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)

Black, female

2004.....	10,745	-	-	1	2	1	5	16	18	60	127	218
2003.....	11,472	-	1	2	2	1	5	10	24	54	128	221
2002.....	11,410	1	-	-	-	2	2	8	23	55	132	251
2001.....	11,528	2	2	-	1	1	8	10	18	51	129	225
2000.....	11,572	-	-	-	-	1	3	5	27	71	141	237
1999.....	11,610	1	-	1	-	2	7	6	17	62	133	210

American Indian 1/, both sexes

2004.....	855	-	1	-	-	-	-	4	5	5	21	31
2003.....	804	-	-	-	-	-	2	4	8	12	23	31
2002.....	766	1	-	-	-	-	2	4	6	7	11	19
2001.....	668	-	-	-	-	-	2	-	1	5	14	29
2000.....	775	-	-	-	-	-	1	1	1	8	21	31
1999.....	714	1	-	-	-	-	1	2	3	11	16	24

American Indian 1/, male

2004.....	481	-	-	-	-	-	-	3	5	5	16	22
2003.....	461	-	-	-	-	-	1	-	6	10	15	20
2002.....	440	1	-	-	-	-	2	2	6	6	7	15
2001.....	367	-	-	-	-	-	2	-	1	2	9	22
2000.....	439	-	-	-	-	-	1	1	-	7	17	25
1999.....	396	1	-	-	-	-	1	2	-	8	12	19

American Indian 1/, female

2004.....	374	-	1	-	-	-	-	1	-	-	5	9
2003.....	343	-	-	-	-	-	1	-	2	2	8	11
2002.....	326	-	-	-	-	-	-	2	-	1	4	4
2001.....	301	-	-	-	-	-	-	-	-	3	5	7
2000.....	336	-	-	-	-	-	-	-	1	1	4	6
1999.....	318	-	-	-	-	-	-	-	3	3	4	5

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Asian or Pacific Islander 2/, both sexes												
2004.....	3,652	-	1	-	-	1	2	2	10	22	25	82
2003.....	3,652	-	-	1	-	-	1	9	10	28	50	74
2002.....	3,583	-	-	-	-	-	2	2	13	26	41	85
2001.....	3,375	-	-	-	-	-	2	3	12	21	37	68
2000.....	3,152	1	-	-	1	-	2	2	8	21	29	60
1999.....	3,257	-	-	-	-	2	-	3	10	25	43	62
Asian or Pacific Islander 2/, male												
2004.....	1,975	-	-	-	-	1	-	2	10	19	23	67
2003.....	2,005	-	-	-	-	-	1	7	9	22	39	64
2002.....	1,999	-	-	-	-	-	2	2	10	23	36	68
2001.....	1,840	-	-	-	-	-	1	2	10	19	31	50
2000.....	1,829	1	-	-	-	-	2	2	6	20	24	49
1999.....	1,912	-	-	-	-	2	-	2	9	24	40	54
Asian or Pacific Islander 2/, female												
2004.....	1,677	-	1	-	-	-	2	-	-	3	2	15
2003.....	1,647	-	-	1	-	-	-	2	1	6	11	10
2002.....	1,584	-	-	-	-	-	-	-	3	3	5	17
2001.....	1,535	-	-	-	-	-	1	1	2	2	6	18
2000.....	1,323	-	-	-	1	-	-	-	2	1	5	11
1999.....	1,345	-	-	-	-	-	-	1	1	1	3	8

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other heart diseases (I26-I51)												
All races, both sexes												
2004.....	166,275	406	174	73	148	321	534	696	1,023	1,637	2,647	3,965
2003.....	169,984	416	171	90	144	348	556	692	1,118	1,673	2,882	3,891
2002.....	169,540	473	157	84	143	362	474	723	1,085	1,769	2,851	3,849
2001.....	166,949	449	215	92	154	305	489	687	1,049	1,770	2,702	3,708
2000.....	165,428	457	168	96	152	355	498	687	1,062	1,706	2,751	3,536
1999.....	165,828	490	177	99	149	398	482	680	1,099	1,879	2,768	3,502
All races, male												
2004.....	73,196	218	90	28	83	204	351	447	628	994	1,648	2,530
2003.....	74,738	233	81	50	91	219	341	404	677	1,017	1,784	2,439
2002.....	74,230	249	82	38	87	250	307	446	692	1,051	1,809	2,468
2001.....	72,439	222	117	41	82	193	307	413	624	1,099	1,678	2,361
2000.....	72,122	238	98	58	92	218	300	407	642	1,033	1,713	2,278
1999.....	72,601	255	100	54	83	250	283	400	667	1,176	1,718	2,223
All races, female												
2004.....	93,079	188	84	45	65	117	183	249	395	643	999	1,435
2003.....	95,246	183	90	40	53	129	215	288	441	656	1,098	1,452
2002.....	95,310	224	75	46	56	112	167	277	393	718	1,042	1,381
2001.....	94,510	227	98	51	72	112	182	274	425	671	1,024	1,347
2000.....	93,306	219	70	38	60	137	198	280	420	673	1,038	1,258
1999.....	93,227	235	77	45	66	148	199	280	432	703	1,050	1,279
White, both sexes												
2004.....	143,860	276	107	47	97	209	368	437	651	1,088	1,812	2,740
2003.....	147,127	256	109	57	98	235	355	425	717	1,106	1,972	2,736
2002.....	147,047	314	109	61	96	245	303	446	678	1,172	2,016	2,647
2001.....	144,862	292	134	64	107	203	309	438	681	1,166	1,806	2,561
2000.....	143,761	315	114	63	95	229	309	433	700	1,121	1,885	2,443
1999.....	143,786	329	119	72	103	281	314	425	729	1,252	1,875	2,390

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other heart diseases (I26-I51)

White, male

2004.....	62,477	148	58	18	56	132	247	285	410	683	1,171	1,783
2003.....	63,973	140	48	31	64	147	215	252	457	686	1,269	1,763
2002.....	63,698	168	62	28	58	170	204	280	442	709	1,327	1,743
2001.....	62,025	155	71	29	59	124	195	268	408	736	1,187	1,687
2000.....	61,933	164	71	38	62	139	194	267	435	699	1,206	1,639
1999.....	62,054	174	70	39	56	170	182	259	453	805	1,216	1,580

White, female

2004.....	81,383	128	49	29	41	77	121	152	241	405	641	957
2003.....	83,154	116	61	26	34	88	140	173	260	420	703	973
2002.....	83,349	146	47	33	38	75	99	166	236	463	689	904
2001.....	82,837	137	63	35	48	79	114	170	273	430	619	874
2000.....	81,828	151	43	25	33	90	115	166	265	422	679	804
1999.....	81,732	155	49	33	47	111	132	166	276	447	659	810

Black, both sexes

2004.....	19,604	98	60	24	47	99	142	222	330	500	774	1,125
2003.....	20,034	129	51	29	37	97	182	238	356	516	843	1,075
2002.....	19,812	136	38	16	41	99	154	252	363	549	773	1,113
2001.....	19,637	131	71	26	39	92	162	230	341	538	839	1,061
2000.....	19,298	117	48	28	45	113	167	236	328	526	801	1,023
1999.....	19,685	135	49	26	40	109	154	233	344	576	833	1,028

Black, male

2004.....	9,280	53	31	10	24	66	88	139	183	283	436	683
2003.....	9,372	78	28	16	22	64	116	136	189	301	474	626
2002.....	9,128	72	16	6	24	65	93	150	218	311	436	672
2001.....	9,148	59	40	11	21	61	101	135	195	319	454	612
2000.....	8,958	59	24	16	24	68	91	129	186	300	459	595
1999.....	9,344	67	27	14	24	76	91	132	199	337	468	595

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other heart diseases (I26-I51)												
Black, female												
2004.....	10,324	45	29	14	23	33	54	83	147	217	338	442
2003.....	10,662	51	23	13	15	33	66	102	167	215	369	449
2002.....	10,684	64	22	10	17	34	61	102	145	238	337	441
2001.....	10,489	72	31	15	18	31	61	95	146	219	385	449
2000.....	10,340	58	24	12	21	45	76	107	142	226	342	428
1999.....	10,341	68	22	12	16	33	63	101	145	239	365	433
American Indian 1/, both sexes												
2004.....	618	10	1	1	1	7	5	13	10	19	22	33
2003.....	676	9	4	-	2	4	5	7	14	19	25	31
2002.....	594	4	6	2	3	6	4	5	12	15	21	30
2001.....	599	9	3	2	1	2	5	9	7	28	24	38
2000.....	591	6	1	1	4	2	5	4	9	25	22	25
1999.....	595	5	5	-	3	2	6	5	7	19	24	30
American Indian 1/, male												
2004.....	307	6	-	-	1	2	2	8	8	13	13	19
2003.....	333	5	1	-	-	2	4	3	13	12	15	20
2002.....	328	2	2	1	3	5	3	2	9	12	16	21
2001.....	318	3	2	1	-	1	4	4	5	18	15	30
2000.....	295	3	-	1	2	2	3	2	5	12	15	15
1999.....	289	2	2	-	1	1	4	3	2	13	11	17
American Indian 1/, female												
2004.....	311	4	1	1	-	5	3	5	2	6	9	14
2003.....	343	4	3	-	2	2	1	4	1	7	10	11
2002.....	266	2	4	1	-	1	1	3	3	3	5	9
2001.....	281	6	1	1	1	1	1	5	2	10	9	8
2000.....	296	3	1	-	2	-	2	2	4	13	7	10
1999.....	306	3	3	-	2	1	2	2	5	6	13	13

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other heart diseases (I26-I51)

Asian or Pacific Islander 2/, both sexes

2004.....	2,193	22	6	1	3	6	19	24	32	30	39	67
2003.....	2,147	22	7	4	7	12	14	22	31	32	42	49
2002.....	2,087	19	4	5	3	12	13	20	32	33	41	59
2001.....	1,851	17	7	-	7	8	13	10	20	38	33	48
2000.....	1,778	19	5	4	8	11	17	14	25	34	43	45
1999.....	1,762	21	4	1	3	6	8	17	19	32	36	54

Asian or Pacific Islander 2/, male

2004.....	1,132	11	1	-	2	4	14	15	27	15	28	45
2003.....	1,060	10	4	3	5	6	6	13	18	18	26	30
2002.....	1,076	7	2	3	2	10	7	14	23	19	30	32
2001.....	948	5	4	-	2	7	7	6	16	26	22	32
2000.....	936	12	3	3	4	9	12	9	16	22	33	29
1999.....	914	12	1	1	2	3	6	6	13	21	23	31

Asian or Pacific Islander 2/, female

2004.....	1,061	11	5	1	1	2	5	9	5	15	11	22
2003.....	1,087	12	3	1	2	6	8	9	13	14	16	19
2002.....	1,011	12	2	2	1	2	6	6	9	14	11	27
2001.....	903	12	3	-	5	1	6	4	4	12	11	16
2000.....	842	7	2	1	4	2	5	5	9	12	10	16
1999.....	848	9	3	-	1	3	2	11	6	11	13	23

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	-------------	-----------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)

All races, both sexes

2004.....	13,122	155	24	6	19	45	82	128	182	305	427	640
2003.....	13,989	132	24	15	15	51	92	125	220	329	512	600
2002.....	13,722	186	28	11	22	41	70	136	189	317	438	628
2001.....	13,320	170	21	11	19	34	88	146	207	307	470	599
2000.....	13,047	185	32	17	12	49	83	125	218	319	471	572
1999.....	13,244	218	19	21	14	60	97	132	211	339	459	557

All races, male

2004.....	5,145	85	10	3	6	18	33	54	70	118	196	297
2003.....	5,628	75	13	8	7	21	31	40	85	139	211	287
2002.....	5,372	98	13	5	9	17	24	63	74	139	182	293
2001.....	5,299	80	12	9	10	17	34	59	71	122	213	267
2000.....	5,205	99	18	11	7	15	28	50	92	137	187	267
1999.....	5,369	114	11	12	5	26	33	46	84	147	197	246

All races, female

2004.....	7,977	70	14	3	13	27	49	74	112	187	231	343
2003.....	8,361	57	11	7	8	30	61	85	135	190	301	313
2002.....	8,350	88	15	6	13	24	46	73	115	178	256	335
2001.....	8,021	90	9	2	9	17	54	87	136	185	257	332
2000.....	7,842	86	14	6	5	34	55	75	126	182	284	305
1999.....	7,875	104	8	9	9	34	64	86	127	192	262	311

White, both sexes

2004.....	10,749	103	17	6	14	33	45	79	119	201	299	469
2003.....	11,456	78	15	9	11	37	53	82	143	208	346	445
2002.....	11,324	121	16	7	17	27	43	73	115	222	321	455
2001.....	10,988	114	16	9	14	21	60	90	140	206	333	433
2000.....	10,781	122	21	11	11	33	53	81	143	216	324	425
1999.....	10,832	154	13	18	13	49	65	78	134	234	313	399

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)

White, male

2004.....	4,201	56	8	3	4	13	17	32	49	72	150	231
2003.....	4,616	47	7	3	4	15	15	22	56	90	143	211
2002.....	4,493	65	8	4	8	11	15	35	47	97	141	215
2001.....	4,380	54	10	7	8	7	24	38	48	83	157	199
2000.....	4,350	66	12	7	7	11	21	37	60	101	129	210
1999.....	4,428	83	8	10	4	20	19	30	51	110	143	188

White, female

2004.....	6,548	47	9	3	10	20	28	47	70	129	149	238
2003.....	6,840	31	8	6	7	22	38	60	87	118	203	234
2002.....	6,831	56	8	3	9	16	28	38	68	125	180	240
2001.....	6,608	60	6	2	6	14	36	52	92	123	176	234
2000.....	6,431	56	9	4	4	22	32	44	83	115	195	215
1999.....	6,404	71	5	8	9	29	46	48	83	124	170	211

Black, both sexes

2004.....	2,168	39	6	-	4	10	34	43	62	95	125	160
2003.....	2,321	43	8	5	2	10	36	38	69	109	152	147
2002.....	2,237	55	10	2	4	14	24	59	69	91	114	164
2001.....	2,163	44	5	2	5	12	23	51	65	93	130	156
2000.....	2,100	49	9	6	1	15	29	42	70	92	137	138
1999.....	2,246	57	4	2	1	10	28	49	73	97	134	143

Black, male

2004.....	850	22	2	-	2	3	15	21	20	44	44	61
2003.....	939	24	5	4	1	6	16	15	27	46	64	72
2002.....	812	31	3	-	1	6	9	28	24	40	39	73
2001.....	845	23	2	2	2	9	9	18	22	38	51	63
2000.....	799	24	5	4	-	3	7	11	32	35	52	53
1999.....	877	27	2	1	1	5	12	16	33	33	51	53

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)

Black, female

2004.....	1,318	17	4	-	2	7	19	22	42	51	81	99
2003.....	1,382	19	3	1	1	4	20	23	42	63	88	75
2002.....	1,425	24	7	2	3	8	15	31	45	51	75	91
2001.....	1,318	21	3	-	3	3	14	33	43	55	79	93
2000.....	1,301	25	4	2	1	12	22	31	38	57	85	85
1999.....	1,369	30	2	1	-	5	16	33	40	64	83	90

American Indian 1/, both sexes

2004.....	58	5	1	-	-	1	-	2	-	3	3	5
2003.....	74	3	1	-	-	1	1	4	1	4	5	5
2002.....	51	1	1	-	-	-	1	2	-	1	3	3
2001.....	44	-	-	-	-	-	2	2	-	3	3	6
2000.....	43	3	-	-	-	-	-	-	1	4	5	4
1999.....	56	1	1	-	-	-	2	1	1	3	5	7

American Indian 1/, male

2004.....	32	4	-	-	-	1	-	-	-	2	2	2
2003.....	28	-	1	-	-	-	-	2	1	1	1	4
2002.....	19	1	1	-	-	-	-	-	-	1	2	1
2001.....	21	-	-	-	-	-	1	1	-	-	3	3
2000.....	14	2	-	-	-	-	-	-	-	1	2	2
1999.....	21	-	1	-	-	-	1	-	-	2	-	2

American Indian 1/, female

2004.....	26	1	1	-	-	-	-	2	-	1	1	3
2003.....	46	3	-	-	-	1	1	2	-	3	4	1
2002.....	32	-	-	-	-	-	1	2	-	-	1	2
2001.....	23	-	-	-	-	-	1	1	-	3	-	3
2000.....	29	1	-	-	-	-	-	-	1	3	3	2
1999.....	35	1	-	-	-	-	1	1	1	1	5	5

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Asian or Pacific Islander 2/, both sexes												
2004.....	147	8	-	-	1	1	3	4	1	6	-	6
2003.....	138	8	-	1	2	3	2	1	7	8	9	3
2002.....	110	9	1	2	1	-	2	2	5	3	-	6
2001.....	125	12	-	-	-	1	3	3	2	5	4	4
2000.....	123	11	2	-	-	1	1	2	4	7	5	5
1999.....	110	6	1	1	-	1	2	4	3	5	7	8
Asian or Pacific Islander 2/, male												
2004.....	62	3	-	-	-	1	1	1	1	-	-	3
2003.....	45	4	-	1	2	-	-	1	1	2	3	-
2002.....	48	1	1	1	-	-	-	-	3	1	-	4
2001.....	53	3	-	-	-	1	-	2	1	1	2	2
2000.....	42	7	1	-	-	1	-	2	-	-	4	2
1999.....	43	4	-	1	-	1	1	-	-	2	3	3
Asian or Pacific Islander 2/, female												
2004.....	85	5	-	-	1	-	2	3	-	6	-	3
2003.....	93	4	-	-	-	3	2	-	6	6	6	3
2002.....	62	8	-	1	1	-	2	2	2	2	-	2
2001.....	72	9	-	-	-	-	3	1	1	4	2	2
2000.....	81	4	1	-	-	-	1	-	4	7	1	3
1999.....	67	2	1	-	-	-	1	4	3	3	4	5

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary embolism (I26)												
All races, both sexes												
2004.....	8,113	3	1	1	5	22	54	83	130	212	285	450
2003.....	8,702	3	2	2	4	21	68	80	146	227	335	407
2002.....	8,620	2	-	1	6	22	52	94	121	220	297	420
2001.....	8,627	2	1	-	2	15	56	94	143	205	316	398
2000.....	8,437	5	1	5	3	25	59	80	138	216	308	387
1999.....	9,008	2	-	1	2	31	65	86	145	217	310	389
All races, male												
2004.....	3,385	2	1	1	1	8	24	39	54	77	136	209
2003.....	3,736	1	2	1	2	11	24	30	61	100	142	212
2002.....	3,514	1	-	-	4	8	19	45	50	94	131	215
2001.....	3,572	-	-	-	2	7	22	40	51	88	149	179
2000.....	3,539	3	1	4	2	8	19	34	64	96	126	193
1999.....	3,724	-	-	-	1	12	24	31	65	100	136	183
All races, female												
2004.....	4,728	1	-	-	4	14	30	44	76	135	149	241
2003.....	4,966	2	-	1	2	10	44	50	85	127	193	195
2002.....	5,106	1	-	1	2	14	33	49	71	126	166	205
2001.....	5,055	2	1	-	-	8	34	54	92	117	167	219
2000.....	4,898	2	-	1	1	17	40	46	74	120	182	194
1999.....	5,284	2	-	1	1	19	41	55	80	117	174	206
White, both sexes												
2004.....	6,636	2	1	1	3	15	35	55	91	142	195	326
2003.....	7,092	1	2	2	3	13	36	49	91	142	232	309
2002.....	7,119	2	-	1	4	15	33	51	79	157	226	313
2001.....	7,130	2	-	-	1	10	41	53	93	132	224	288
2000.....	6,942	3	1	4	3	16	34	54	94	144	211	282
1999.....	7,377	1	-	1	2	21	40	49	88	150	218	281

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary embolism (I26)												
White, male												
2004.....	2,743	2	1	1	1	5	15	22	39	44	101	161
2003.....	3,033	-	2	1	1	7	10	14	40	63	100	154
2002.....	2,933	1	-	-	3	6	12	25	31	66	107	158
2001.....	2,947	-	-	-	1	3	16	23	31	59	106	133
2000.....	2,932	2	1	3	2	6	13	28	44	67	84	148
1999.....	3,044	-	-	-	1	6	13	18	39	76	103	139
White, female												
2004.....	3,893	-	-	-	2	10	20	33	52	98	94	165
2003.....	4,059	1	-	1	2	6	26	35	51	79	132	155
2002.....	4,186	1	-	1	1	9	21	26	48	91	119	155
2001.....	4,183	2	-	-	-	7	25	30	62	73	118	155
2000.....	4,010	1	-	1	1	10	21	26	50	77	127	134
1999.....	4,333	1	-	1	1	15	27	31	49	74	115	142
Black, both sexes												
2004.....	1,380	1	-	-	2	6	18	26	39	67	87	118
2003.....	1,521	2	-	-	-	7	31	27	54	83	97	95
2002.....	1,424	-	-	-	2	7	18	41	41	60	71	105
2001.....	1,423	-	1	-	1	5	13	38	48	71	88	109
2000.....	1,414	2	-	1	-	9	24	24	43	68	92	102
1999.....	1,553	1	-	-	-	9	21	34	56	60	88	101
Black, male												
2004.....	589	-	-	-	-	2	9	17	15	32	33	45
2003.....	663	1	-	-	-	4	14	13	21	36	41	55
2002.....	547	-	-	-	1	2	7	20	18	27	24	55
2001.....	588	-	-	-	1	4	5	15	19	29	39	46
2000.....	579	1	-	1	-	2	6	4	20	28	39	43
1999.....	651	-	-	-	-	5	9	13	26	20	33	40

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary embolism (I26)												
Black, female												
2004.....	791	1	-	-	2	4	9	9	24	35	54	73
2003.....	858	1	-	-	-	3	17	14	33	47	56	40
2002.....	877	-	-	-	1	5	11	21	23	33	47	50
2001.....	835	-	1	-	-	1	8	23	29	42	49	63
2000.....	835	1	-	-	-	7	18	20	23	40	53	59
1999.....	902	1	-	-	-	4	12	21	30	40	55	61
American Indian 1/, both sexes												
2004.....	30	-	-	-	-	1	-	1	-	1	3	2
2003.....	44	-	-	-	-	-	1	3	-	2	4	3
2002.....	32	-	-	-	-	-	-	2	-	1	-	-
2001.....	26	-	-	-	-	-	2	2	-	1	2	1
2000.....	23	-	-	-	-	-	-	-	1	2	2	2
1999.....	31	-	-	-	-	-	2	1	1	2	1	4
American Indian 1/, male												
2004.....	19	-	-	-	-	1	-	-	-	1	2	1
2003.....	21	-	-	-	-	-	-	2	-	1	1	3
2002.....	10	-	-	-	-	-	-	-	-	1	-	-
2001.....	12	-	-	-	-	-	1	1	-	-	2	-
2000.....	7	-	-	-	-	-	-	-	-	1	1	1
1999.....	13	-	-	-	-	-	1	-	-	2	-	2
American Indian 1/, female												
2004.....	11	-	-	-	-	-	-	1	-	-	1	1
2003.....	23	-	-	-	-	-	1	1	-	1	3	-
2002.....	22	-	-	-	-	-	-	2	-	-	-	-
2001.....	14	-	-	-	-	-	1	1	-	1	-	1
2000.....	16	-	-	-	-	-	-	-	1	1	1	1
1999.....	18	-	-	-	-	-	1	1	1	-	1	2

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary embolism (I26)												
Asian or Pacific Islander 2/, both sexes												
2004.....	67	-	-	-	-	-	1	1	-	2	-	4
2003.....	45	-	-	-	1	1	-	1	1	-	2	-
2002.....	45	-	-	-	-	-	1	-	1	2	-	2
2001.....	48	-	-	-	-	-	-	1	2	1	2	-
2000.....	58	-	-	-	-	-	1	2	-	2	3	1
1999.....	47	-	-	-	-	1	2	2	-	5	3	3
Asian or Pacific Islander 2/, male												
2004.....	34	-	-	-	-	-	-	-	-	-	-	2
2003.....	19	-	-	-	1	-	-	1	-	-	-	-
2002.....	24	-	-	-	-	-	-	-	1	-	-	2
2001.....	25	-	-	-	-	-	-	1	1	-	2	-
2000.....	21	-	-	-	-	-	-	2	-	-	2	1
1999.....	16	-	-	-	-	1	1	-	-	2	-	2
Asian or Pacific Islander 2/, female												
2004.....	33	-	-	-	-	-	1	1	-	2	-	2
2003.....	26	-	-	-	-	1	-	-	1	-	2	-
2002.....	21	-	-	-	-	-	1	-	-	2	-	-
2001.....	23	-	-	-	-	-	-	-	1	1	-	-
2000.....	37	-	-	-	-	-	1	-	-	2	1	-
1999.....	31	-	-	-	-	-	1	2	-	3	3	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)

All races, both sexes

2004.....	5,009	152	23	5	14	23	28	45	52	93	142	190
2003.....	5,287	129	22	13	11	30	24	45	74	102	177	193
2002.....	5,102	184	28	10	16	19	18	42	68	97	141	208
2001.....	4,693	168	20	11	17	19	32	52	64	102	154	201
2000.....	4,610	180	31	12	9	24	24	45	80	103	163	185
1999.....	4,236	216	19	20	12	29	32	46	66	122	149	168

All races, male

2004.....	1,760	83	9	2	5	10	9	15	16	41	60	88
2003.....	1,892	74	11	7	5	10	7	10	24	39	69	75
2002.....	1,858	97	13	5	5	9	5	18	24	45	51	78
2001.....	1,727	80	12	9	8	10	12	19	20	34	64	88
2000.....	1,666	96	17	7	5	7	9	16	28	41	61	74
1999.....	1,645	114	11	12	4	14	9	15	19	47	61	63

All races, female

2004.....	3,249	69	14	3	9	13	19	30	36	52	82	102
2003.....	3,395	55	11	6	6	20	17	35	50	63	108	118
2002.....	3,244	87	15	5	11	10	13	24	44	52	90	130
2001.....	2,966	88	8	2	9	9	20	33	44	68	90	113
2000.....	2,944	84	14	5	4	17	15	29	52	62	102	111
1999.....	2,591	102	8	8	8	15	23	31	47	75	88	105

White, both sexes

2004.....	4,113	101	16	5	11	18	10	24	28	59	104	143
2003.....	4,364	77	13	7	8	24	17	33	52	66	114	136
2002.....	4,205	119	16	6	13	12	10	22	36	65	95	142
2001.....	3,858	112	16	9	13	11	19	37	47	74	109	145
2000.....	3,839	119	20	7	8	17	19	27	49	72	113	143
1999.....	3,455	153	13	17	11	28	25	29	46	84	95	118

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)

White, male

2004.....	1,458	54	7	2	3	8	2	10	10	28	49	70
2003.....	1,583	47	5	2	3	8	5	8	16	27	43	57
2002.....	1,560	64	8	4	5	5	3	10	16	31	34	57
2001.....	1,433	54	10	7	7	4	8	15	17	24	51	66
2000.....	1,418	64	11	4	5	5	8	9	16	34	45	62
1999.....	1,384	83	8	10	3	14	6	12	12	34	40	49

White, female

2004.....	2,655	47	9	3	8	10	8	14	18	31	55	73
2003.....	2,781	30	8	5	5	16	12	25	36	39	71	79
2002.....	2,645	55	8	2	8	7	7	12	20	34	61	85
2001.....	2,425	58	6	2	6	7	11	22	30	50	58	79
2000.....	2,421	55	9	3	3	12	11	18	33	38	68	81
1999.....	2,071	70	5	7	8	14	19	17	34	50	55	69

Black, both sexes

2004.....	788	38	6	-	2	4	16	17	23	28	38	42
2003.....	800	41	8	5	2	3	5	11	15	26	55	52
2002.....	813	55	10	2	2	7	6	18	28	31	43	59
2001.....	740	44	4	2	4	7	10	13	17	22	42	47
2000.....	686	47	9	5	1	6	5	18	27	24	45	36
1999.....	693	56	4	2	1	1	7	15	17	37	46	42

Black, male

2004.....	261	22	2	-	2	1	6	4	5	12	11	16
2003.....	276	23	5	4	1	2	2	2	6	10	23	17
2002.....	265	31	3	-	-	4	2	8	6	13	15	18
2001.....	257	23	2	2	1	5	4	3	3	9	12	17
2000.....	220	23	5	3	-	1	1	7	12	7	13	10
1999.....	226	27	2	1	1	-	3	3	7	13	18	13

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)

Black, female

2004.....	527	16	4	-	-	3	10	13	18	16	27	26
2003.....	524	18	3	1	1	1	3	9	9	16	32	35
2002.....	548	24	7	2	2	3	4	10	22	18	28	41
2001.....	483	21	2	-	3	2	6	10	14	13	30	30
2000.....	466	24	4	2	1	5	4	11	15	17	32	26
1999.....	467	29	2	1	-	1	4	12	10	24	28	29

American Indian 1/, both sexes

2004.....	28	5	1	-	-	-	-	1	-	2	-	3
2003.....	30	3	1	-	-	1	-	1	1	2	1	2
2002.....	19	1	1	-	-	-	1	-	-	-	3	3
2001.....	18	-	-	-	-	-	-	-	-	2	1	5
2000.....	20	3	-	-	-	-	-	-	-	2	3	2
1999.....	25	1	1	-	-	-	-	-	-	1	4	3

American Indian 1/, male

2004.....	13	4	-	-	-	-	-	-	-	1	-	1
2003.....	7	-	1	-	-	-	-	-	1	-	-	1
2002.....	9	1	1	-	-	-	-	-	-	-	2	1
2001.....	9	-	-	-	-	-	-	-	-	-	1	3
2000.....	7	2	-	-	-	-	-	-	-	-	1	1
1999.....	8	-	1	-	-	-	-	-	-	-	-	-

American Indian 1/, female

2004.....	15	1	1	-	-	-	-	1	-	1	-	2
2003.....	23	3	-	-	-	1	-	1	-	2	1	1
2002.....	10	-	-	-	-	-	1	-	-	-	1	2
2001.....	9	-	-	-	-	-	-	-	-	2	-	2
2000.....	13	1	-	-	-	-	-	-	-	2	2	1
1999.....	17	1	-	-	-	-	-	-	-	1	4	3

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)

Asian or Pacific Islander 2/, both sexes

2004.....	80	8	-	-	1	1	2	3	1	4	-	2
2003.....	93	8	-	1	1	2	2	-	6	8	7	3
2002.....	65	9	1	2	1	-	1	2	4	1	-	4
2001.....	77	12	-	-	-	1	3	2	-	4	2	4
2000.....	65	11	2	-	-	1	-	-	4	5	2	4
1999.....	63	6	1	1	-	-	-	2	3	-	4	5

Asian or Pacific Islander 2/, male

2004.....	28	3	-	-	-	1	1	1	1	-	-	1
2003.....	26	4	-	1	1	-	-	-	1	2	3	-
2002.....	24	1	1	1	-	-	-	-	2	1	-	2
2001.....	28	3	-	-	-	1	-	1	-	1	-	2
2000.....	21	7	1	-	-	1	-	-	-	-	2	1
1999.....	27	4	-	1	-	-	-	-	-	-	3	1

Asian or Pacific Islander 2/, female

2004.....	52	5	-	-	1	-	1	2	-	4	-	1
2003.....	67	4	-	-	-	2	2	-	5	6	4	3
2002.....	41	8	-	1	1	-	1	2	2	-	-	2
2001.....	49	9	-	-	-	-	3	1	-	3	2	2
2000.....	44	4	1	-	-	-	-	-	4	5	-	3
1999.....	36	2	1	-	-	-	-	2	3	-	1	4

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other non-pulmonary forms of heart disease (I30-I51)

All races, both sexes

2004.....	153,153	251	150	67	129	276	452	568	841	1,332	2,220	3,325
2003.....	155,995	284	147	75	129	297	464	567	898	1,344	2,370	3,291
2002.....	155,818	287	129	73	121	321	404	587	896	1,452	2,413	3,221
2001.....	153,629	279	194	81	135	271	401	541	842	1,463	2,232	3,109
2000.....	152,381	272	136	79	140	306	415	562	844	1,387	2,280	2,964
1999.....	152,584	272	158	78	135	338	385	548	888	1,540	2,309	2,945

All races, male

2004.....	68,051	133	80	25	77	186	318	393	558	876	1,452	2,233
2003.....	69,110	158	68	42	84	198	310	364	592	878	1,573	2,152
2002.....	68,858	151	69	33	78	233	283	383	618	912	1,627	2,175
2001.....	67,140	142	105	32	72	176	273	354	553	977	1,465	2,094
2000.....	66,917	139	80	47	85	203	272	357	550	896	1,526	2,011
1999.....	67,232	141	89	42	78	224	250	354	583	1,029	1,521	1,977

All races, female

2004.....	85,102	118	70	42	52	90	134	175	283	456	768	1,092
2003.....	86,885	126	79	33	45	99	154	203	306	466	797	1,139
2002.....	86,960	136	60	40	43	88	121	204	278	540	786	1,046
2001.....	86,489	137	89	49	63	95	128	187	289	486	767	1,015
2000.....	85,464	133	56	32	55	103	143	205	294	491	754	953
1999.....	85,352	131	69	36	57	114	135	194	305	511	788	968

White, both sexes

2004.....	133,111	173	90	41	83	176	323	358	532	887	1,513	2,271
2003.....	135,671	178	94	48	87	198	302	343	574	898	1,626	2,291
2002.....	135,723	193	93	54	79	218	260	373	563	950	1,695	2,192
2001.....	133,874	178	118	55	93	182	249	348	541	960	1,473	2,128
2000.....	132,980	193	93	52	84	196	256	352	557	905	1,561	2,018
1999.....	132,954	175	106	54	90	232	249	347	595	1,018	1,562	1,991

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other non-pulmonary forms of heart disease (I30-I51)												
White, male												
2004.....	58,276	92	50	15	52	119	230	253	361	611	1,021	1,552
2003.....	59,357	93	41	28	60	132	200	230	401	596	1,126	1,552
2002.....	59,205	103	54	24	50	159	189	245	395	612	1,186	1,528
2001.....	57,645	101	61	22	51	117	171	230	360	653	1,030	1,488
2000.....	57,583	98	59	31	55	128	173	230	375	598	1,077	1,429
1999.....	57,626	91	62	29	52	150	163	229	402	695	1,073	1,392
White, female												
2004.....	74,835	81	40	26	31	57	93	105	171	276	492	719
2003.....	76,314	85	53	20	27	66	102	113	173	302	500	739
2002.....	76,518	90	39	30	29	59	71	128	168	338	509	664
2001.....	76,229	77	57	33	42	65	78	118	181	307	443	640
2000.....	75,397	95	34	21	29	68	83	122	182	307	484	589
1999.....	75,328	84	44	25	38	82	86	118	193	323	489	599
Black, both sexes												
2004.....	17,436	59	54	24	43	89	108	179	268	405	649	965
2003.....	17,713	86	43	24	35	87	146	200	287	407	691	928
2002.....	17,575	81	28	14	37	85	130	193	294	458	659	949
2001.....	17,474	87	66	24	34	80	139	179	276	445	709	905
2000.....	17,198	68	39	22	44	98	138	194	258	434	664	885
1999.....	17,439	78	45	24	39	99	126	184	271	479	699	885
Black, male												
2004.....	8,430	31	29	10	22	63	73	118	163	239	392	622
2003.....	8,433	54	23	12	21	58	100	121	162	255	410	554
2002.....	8,316	41	13	6	23	59	84	122	194	271	397	599
2001.....	8,303	36	38	9	19	52	92	117	173	281	403	549
2000.....	8,159	35	19	12	24	65	84	118	154	265	407	542
1999.....	8,467	40	25	13	23	71	79	116	166	304	417	542

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other non-pulmonary forms of heart disease (I30-I51)

Black, female

2004.....	9,006	28	25	14	21	26	35	61	105	166	257	343
2003.....	9,280	32	20	12	14	29	46	79	125	152	281	374
2002.....	9,259	40	15	8	14	26	46	71	100	187	262	350
2001.....	9,171	51	28	15	15	28	47	62	103	164	306	356
2000.....	9,039	33	20	10	20	33	54	76	104	169	257	343
1999.....	8,972	38	20	11	16	28	47	68	105	175	282	343

American Indian 1/, both sexes

2004.....	560	5	-	1	1	6	5	11	10	16	19	28
2003.....	602	6	3	-	2	3	4	3	13	15	20	26
2002.....	543	3	5	2	3	6	3	3	12	14	18	27
2001.....	555	9	3	2	1	2	3	7	7	25	21	32
2000.....	548	3	1	1	4	2	5	4	8	21	17	21
1999.....	539	4	4	-	3	2	4	4	6	16	19	23

American Indian 1/, male

2004.....	275	2	-	-	1	1	2	8	8	11	11	17
2003.....	305	5	-	-	-	2	4	1	12	11	14	16
2002.....	309	1	1	1	3	5	3	2	9	11	14	20
2001.....	297	3	2	1	-	1	3	3	5	18	12	27
2000.....	281	1	-	1	2	2	3	2	5	11	13	13
1999.....	268	2	1	-	1	1	3	3	2	11	11	15

American Indian 1/, female

2004.....	285	3	-	1	-	5	3	3	2	5	8	11
2003.....	297	1	3	-	2	1	-	2	1	4	6	10
2002.....	234	2	4	1	-	1	-	1	3	3	4	7
2001.....	258	6	1	1	1	1	-	4	2	7	9	5
2000.....	267	2	1	-	2	-	2	2	3	10	4	8
1999.....	271	2	3	-	2	1	1	1	4	5	8	8

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other non-pulmonary forms of heart disease (I30-I51)

Asian or Pacific Islander 2/, both sexes

2004.....	2,046	14	6	1	2	5	16	20	31	24	39	61
2003.....	2,009	14	7	3	5	9	12	21	24	24	33	46
2002.....	1,977	10	3	3	2	12	11	18	27	30	41	53
2001.....	1,726	5	7	-	7	7	10	7	18	33	29	44
2000.....	1,655	8	3	4	8	10	16	12	21	27	38	40
1999.....	1,652	15	3	-	3	5	6	13	16	27	29	46

Asian or Pacific Islander 2/, male

2004.....	1,070	8	1	-	2	3	13	14	26	15	28	42
2003.....	1,015	6	4	2	3	6	6	12	17	16	23	30
2002.....	1,028	6	1	2	2	10	7	14	20	18	30	28
2001.....	895	2	4	-	2	6	7	4	15	25	20	30
2000.....	894	5	2	3	4	8	12	7	16	22	29	27
1999.....	871	8	1	-	2	2	5	6	13	19	20	28

Asian or Pacific Islander 2/, female

2004.....	976	6	5	1	-	2	3	6	5	9	11	19
2003.....	994	8	3	1	2	3	6	9	7	8	10	16
2002.....	949	4	2	1	-	2	4	4	7	12	11	25
2001.....	831	3	3	-	5	1	3	3	3	8	9	14
2000.....	761	3	1	1	4	2	4	5	5	5	9	13
1999.....	781	7	2	-	1	3	1	7	3	8	9	18

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Acute and subacute endocarditis (I33)

All races, both sexes

2004.....	1,196	1	1	-	-	1	11	8	27	37	47	90
2003.....	1,245	1	-	-	1	3	13	18	15	43	69	89
2002.....	1,154	2	2	1	1	2	5	16	28	26	64	84
2001.....	1,178	1	-	1	2	3	5	9	20	46	71	93
2000.....	1,172	3	1	2	1	5	8	13	27	41	75	89
1999.....	1,116	2	1	-	-	2	5	12	18	54	60	68

All races, male

2004.....	659	-	-	-	-	1	7	5	17	24	31	58
2003.....	682	1	-	-	1	-	6	11	8	22	41	60
2002.....	643	1	1	-	1	2	4	9	22	19	41	55
2001.....	668	1	-	-	1	2	3	5	12	25	37	68
2000.....	665	2	-	1	-	1	6	7	17	24	49	59
1999.....	639	2	1	-	-	-	3	9	15	34	41	49

All races, female

2004.....	537	1	1	-	-	-	4	3	10	13	16	32
2003.....	563	-	-	-	-	3	7	7	7	21	28	29
2002.....	511	1	1	1	-	-	1	7	6	7	23	29
2001.....	510	-	-	1	1	1	2	4	8	21	34	25
2000.....	507	1	1	1	1	4	2	6	10	17	26	30
1999.....	477	-	-	-	-	2	2	3	3	20	19	19

White, both sexes

2004.....	957	1	-	-	-	-	10	7	20	31	35	55
2003.....	990	-	-	-	1	2	12	14	11	27	51	57
2002.....	930	1	2	-	-	2	4	11	23	20	46	49
2001.....	917	1	-	1	2	1	5	8	13	30	44	64
2000.....	927	1	1	1	1	1	5	10	17	27	53	57
1999.....	879	1	1	-	-	2	4	10	14	32	37	41

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute and subacute endocarditis (I33)												
White, male												
2004.....	536	-	-	-	-	-	6	5	11	23	24	36
2003.....	541	-	-	-	1	-	6	9	5	14	32	36
2002.....	524	-	1	-	-	2	3	7	19	14	31	34
2001.....	517	1	-	-	1	1	3	4	8	16	25	45
2000.....	536	1	-	-	-	-	3	6	13	14	36	39
1999.....	509	1	1	-	-	-	2	7	11	21	24	32
White, female												
2004.....	421	1	-	-	-	-	4	2	9	8	11	19
2003.....	449	-	-	-	-	2	6	5	6	13	19	21
2002.....	406	1	1	-	-	-	1	4	4	6	15	15
2001.....	400	-	-	1	1	-	2	4	5	14	19	19
2000.....	391	-	1	1	1	1	2	4	4	13	17	18
1999.....	370	-	-	-	-	2	2	3	3	11	13	9
Black, both sexes												
2004.....	204	-	1	-	-	1	-	1	5	5	9	30
2003.....	225	1	-	-	-	1	1	2	3	15	17	29
2002.....	199	-	-	1	-	-	1	5	5	6	17	29
2001.....	235	-	-	-	-	1	-	1	7	15	26	27
2000.....	214	2	-	-	-	3	3	3	8	11	20	30
1999.....	210	-	-	-	-	-	1	2	4	20	21	26
Black, male												
2004.....	104	-	-	-	-	1	-	-	4	1	6	19
2003.....	122	1	-	-	-	-	-	2	2	7	8	21
2002.....	106	-	-	-	-	-	1	2	3	5	10	18
2001.....	135	-	-	-	-	-	-	1	4	8	12	21
2000.....	111	1	-	-	-	1	3	1	2	9	12	20
1999.....	117	-	-	-	-	-	1	2	4	12	15	16

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute and subacute endocarditis (I33)												
Black, female												
2004.....	100	-	1	-	-	-	-	1	1	4	3	11
2003.....	103	-	-	-	-	1	1	-	1	8	9	8
2002.....	93	-	-	1	-	-	-	3	2	1	7	11
2001.....	100	-	-	-	-	1	-	-	3	7	14	6
2000.....	103	1	-	-	-	2	-	2	6	2	8	10
1999.....	93	-	-	-	-	-	-	-	-	8	6	10
American Indian 1/, both sexes												
2004.....	13	-	-	-	-	-	-	-	1	1	1	4
2003.....	12	-	-	-	-	-	-	1	1	1	1	2
2002.....	6	-	-	-	-	-	-	-	-	-	-	3
2001.....	6	-	-	-	-	1	-	-	-	-	-	2
2000.....	13	-	-	-	-	-	-	-	-	1	2	2
1999.....	7	-	-	-	-	-	-	-	-	2	1	-
American Indian 1/, male												
2004.....	5	-	-	-	-	-	-	-	1	-	-	2
2003.....	7	-	-	-	-	-	-	-	1	1	1	2
2002.....	2	-	-	-	-	-	-	-	-	-	-	1
2001.....	5	-	-	-	-	1	-	-	-	-	-	2
2000.....	7	-	-	-	-	-	-	-	-	1	1	-
1999.....	4	-	-	-	-	-	-	-	-	1	1	-
American Indian 1/, female												
2004.....	8	-	-	-	-	-	-	-	-	1	1	2
2003.....	5	-	-	-	-	-	-	1	-	-	-	-
2002.....	4	-	-	-	-	-	-	-	-	-	-	2
2001.....	1	-	-	-	-	-	-	-	-	-	-	-
2000.....	6	-	-	-	-	-	-	-	-	-	1	2
1999.....	3	-	-	-	-	-	-	-	-	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
All races, both sexes												
2004.....	779	21	25	7	10	17	14	16	23	29	38	49
2003.....	868	33	22	14	14	21	34	23	25	32	54	63
2002.....	848	26	22	12	10	17	19	16	30	35	40	56
2001.....	839	32	26	12	9	14	26	20	30	43	50	57
2000.....	767	20	24	9	11	17	21	29	24	45	41	45
1999.....	890	27	25	11	10	12	20	19	31	54	60	59
All races, male												
2004.....	404	11	15	5	5	9	9	8	10	20	20	27
2003.....	449	15	8	4	8	14	20	13	16	21	31	41
2002.....	434	9	11	4	3	12	12	9	23	17	32	40
2001.....	401	18	15	2	5	7	15	16	17	24	17	30
2000.....	404	10	13	4	5	10	11	17	12	30	28	28
1999.....	442	13	14	5	2	8	12	14	20	32	32	31
All races, female												
2004.....	375	10	10	2	5	8	5	8	13	9	18	22
2003.....	419	18	14	10	6	7	14	10	9	11	23	22
2002.....	414	17	11	8	7	5	7	7	7	18	8	16
2001.....	438	14	11	10	4	7	11	4	13	19	33	27
2000.....	363	10	11	5	6	7	10	12	12	15	13	17
1999.....	448	14	11	6	8	4	8	5	11	22	28	28
White, both sexes												
2004.....	595	11	12	2	7	10	8	9	14	17	23	33
2003.....	677	15	13	9	8	17	21	10	14	17	38	46
2002.....	666	13	13	8	4	10	16	14	19	22	33	43
2001.....	658	14	14	9	7	10	17	15	23	28	32	43
2000.....	590	11	15	8	2	15	10	21	13	27	31	26
1999.....	703	12	12	7	6	9	15	13	22	41	38	47

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Diseases of pericardium and acute myocarditis (I30-I31,I40)

White, male

2004.....	305	7	7	2	3	8	6	4	6	15	11	15
2003.....	359	9	4	2	5	13	12	5	8	13	23	31
2002.....	344	6	7	2	1	7	10	8	16	9	27	29
2001.....	310	10	8	2	4	3	9	12	13	15	14	21
2000.....	309	6	11	4	1	8	5	11	7	18	21	17
1999.....	351	7	5	4	2	7	8	10	14	24	23	24

White, female

2004.....	290	4	5	-	4	2	2	5	8	2	12	18
2003.....	318	6	9	7	3	4	9	5	6	4	15	15
2002.....	322	7	6	6	3	3	6	6	3	13	6	14
2001.....	348	4	6	7	3	7	8	3	10	13	18	22
2000.....	281	5	4	4	1	7	5	10	6	9	10	9
1999.....	352	5	7	3	4	2	7	3	8	17	15	23

Black, both sexes

2004.....	164	8	12	5	3	4	6	4	8	12	14	16
2003.....	168	14	9	4	5	3	12	12	10	14	16	16
2002.....	154	12	7	2	6	5	3	1	9	10	7	12
2001.....	158	14	12	3	1	4	9	5	5	10	15	14
2000.....	142	7	8	1	7	1	7	6	10	14	10	14
1999.....	160	10	12	4	4	3	3	6	8	12	19	9

Black, male

2004.....	89	3	7	3	2	1	3	3	3	5	9	12
2003.....	81	4	4	2	3	1	8	8	7	8	8	9
2002.....	75	3	3	1	2	3	2	-	6	5	5	10
2001.....	79	5	7	-	1	4	6	4	3	5	3	9
2000.....	72	4	2	-	3	1	2	5	5	9	7	7
1999.....	77	4	9	1	-	1	2	4	6	7	7	5

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Black, female												
2004.....	75	5	5	2	1	3	3	1	5	7	5	4
2003.....	87	10	5	2	2	2	4	4	3	6	8	7
2002.....	79	9	4	1	4	2	1	1	3	5	2	2
2001.....	79	9	5	3	-	-	3	1	2	5	12	5
2000.....	70	3	6	1	4	-	5	1	5	5	3	7
1999.....	83	6	3	3	4	2	1	2	2	5	12	4
American Indian 1/, both sexes												
2004.....	5	1	-	-	-	1	-	1	-	-	1	-
2003.....	6	3	-	-	-	-	-	-	-	-	-	-
2002.....	8	-	1	1	-	1	-	-	-	1	-	1
2001.....	8	3	-	-	1	-	-	-	-	2	1	-
2000.....	6	1	-	-	-	-	1	-	-	-	-	1
1999.....	9	1	1	-	-	-	1	-	1	-	1	1
American Indian 1/, male												
2004.....	1	-	-	-	-	-	-	-	-	-	-	-
2003.....	5	2	-	-	-	-	-	-	-	-	-	-
2002.....	5	-	-	1	-	1	-	-	-	1	-	1
2001.....	5	2	-	-	-	-	-	-	-	2	-	-
2000.....	4	-	-	-	-	-	1	-	-	-	-	1
1999.....	2	-	-	-	-	-	1	-	-	-	-	1
American Indian 1/, female												
2004.....	4	1	-	-	-	1	-	1	-	-	1	-
2003.....	1	1	-	-	-	-	-	-	-	-	-	-
2002.....	3	-	1	-	-	-	-	-	-	-	-	-
2001.....	3	1	-	-	1	-	-	-	-	-	1	-
2000.....	2	1	-	-	-	-	-	-	-	-	-	-
1999.....	7	1	1	-	-	-	-	-	1	-	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other diseases of endocardium (I34-I38)

All races, both sexes

2004.....	20,260	2	8	-	5	23	27	34	78	109	194	325
2003.....	20,056	2	7	2	5	11	29	46	77	112	203	264
2002.....	19,989	6	6	4	3	18	22	53	67	127	219	286
2001.....	19,553	6	7	1	8	11	28	53	64	130	172	259
2000.....	19,737	8	7	4	10	16	30	48	71	141	199	286
1999.....	19,612	5	2	5	7	30	36	43	98	138	241	263

All races, male

2004.....	7,890	-	6	-	4	11	14	18	44	60	120	217
2003.....	7,767	1	4	-	4	8	20	17	41	67	122	154
2002.....	7,658	4	4	3	2	14	11	20	46	74	130	180
2001.....	7,594	5	3	-	4	2	13	32	32	75	105	166
2000.....	7,544	3	3	3	6	8	18	26	31	76	116	188
1999.....	7,586	4	-	3	1	17	20	18	51	86	135	144

All races, female

2004.....	12,370	2	2	-	1	12	13	16	34	49	74	108
2003.....	12,289	1	3	2	1	3	9	29	36	45	81	110
2002.....	12,331	2	2	1	1	4	11	33	21	53	89	106
2001.....	11,959	1	4	1	4	9	15	21	32	55	67	93
2000.....	12,193	5	4	1	4	8	12	22	40	65	83	98
1999.....	12,026	1	2	2	6	13	16	25	47	52	106	119

White, both sexes

2004.....	18,660	1	6	-	4	14	21	22	61	81	140	231
2003.....	18,469	2	7	1	5	10	18	33	53	90	134	189
2002.....	18,494	5	3	4	2	15	14	40	50	95	161	190
2001.....	18,090	4	2	1	7	8	19	36	47	97	126	183
2000.....	18,230	8	6	4	6	13	19	34	53	101	144	211
1999.....	18,067	5	2	4	5	23	27	29	79	108	174	173

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of endocardium (I34-I38)												
White, male												
2004.....	7,201	-	4	-	3	6	12	10	38	46	87	157
2003.....	7,109	1	4	-	4	8	12	14	30	53	84	111
2002.....	7,063	3	3	3	1	11	8	15	33	57	103	124
2001.....	6,954	4	1	-	3	2	9	21	23	57	77	120
2000.....	6,923	3	3	3	4	7	10	19	22	54	90	141
1999.....	6,940	4	-	2	1	13	15	11	41	68	101	93
White, female												
2004.....	11,459	1	2	-	1	8	9	12	23	35	53	74
2003.....	11,360	1	3	1	1	2	6	19	23	37	50	78
2002.....	11,431	2	-	1	1	4	6	25	17	38	58	66
2001.....	11,136	-	1	1	4	6	10	15	24	40	49	63
2000.....	11,307	5	3	1	2	6	9	15	31	47	54	70
1999.....	11,127	1	2	2	4	10	12	18	38	40	73	80
Black, both sexes												
2004.....	1,293	1	2	-	1	8	4	12	15	24	48	85
2003.....	1,278	-	-	1	-	1	8	12	20	20	62	71
2002.....	1,218	1	3	-	-	2	7	12	13	29	52	91
2001.....	1,215	2	4	-	1	3	7	16	16	29	42	71
2000.....	1,273	-	1	-	3	3	8	13	18	33	53	70
1999.....	1,270	-	-	1	1	6	8	14	18	27	62	73
Black, male												
2004.....	541	-	2	-	1	4	1	8	5	13	28	54
2003.....	528	-	-	-	-	-	6	3	8	12	35	41
2002.....	473	1	1	-	-	2	3	4	10	15	21	53
2001.....	516	1	1	-	1	-	2	11	8	16	25	43
2000.....	514	-	-	-	2	1	6	6	9	18	24	45
1999.....	515	-	-	1	-	4	4	7	9	17	30	40

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other diseases of endocardium (I34-I38)

Black, female

2004.....	752	1	-	-	-	4	3	4	10	11	20	31
2003.....	750	-	-	1	-	1	2	9	12	8	27	30
2002.....	745	-	2	-	-	-	4	8	3	14	31	38
2001.....	699	1	3	-	-	3	5	5	8	13	17	28
2000.....	759	-	1	-	1	2	2	7	9	15	29	25
1999.....	755	-	-	-	1	2	4	7	9	10	32	33

American Indian 1/, both sexes

2004.....	60	-	-	-	-	-	1	-	-	1	-	3
2003.....	60	-	-	-	-	-	1	-	2	-	3	2
2002.....	58	-	-	-	1	1	-	-	3	1	1	2
2001.....	41	-	-	-	-	-	-	-	1	1	1	2
2000.....	49	-	-	-	-	-	-	-	-	3	-	3
1999.....	56	-	-	-	1	-	1	-	-	1	2	4

American Indian 1/, male

2004.....	31	-	-	-	-	-	1	-	-	-	-	1
2003.....	33	-	-	-	-	-	1	-	2	-	2	1
2002.....	27	-	-	-	1	1	-	-	2	1	1	1
2001.....	22	-	-	-	-	-	-	-	1	1	1	2
2000.....	14	-	-	-	-	-	-	-	-	-	-	1
1999.....	26	-	-	-	-	-	1	-	-	-	1	2

American Indian 1/, female

2004.....	29	-	-	-	-	-	-	-	-	1	-	2
2003.....	27	-	-	-	-	-	-	-	-	-	1	1
2002.....	31	-	-	-	-	-	-	-	1	-	-	1
2001.....	19	-	-	-	-	-	-	-	-	-	-	-
2000.....	35	-	-	-	-	-	-	-	-	3	-	2
1999.....	30	-	-	-	1	-	-	-	-	1	1	2

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of endocardium (I34-I38)												
Asian or Pacific Islander 2/, both sexes												
2004.....	247	-	-	-	-	1	1	-	2	3	6	6
2003.....	249	-	-	-	-	-	2	1	2	2	4	2
2002.....	219	-	-	-	-	-	1	1	1	2	5	3
2001.....	207	-	1	-	-	-	2	1	-	3	3	3
2000.....	185	-	-	-	1	-	3	1	-	4	2	2
1999.....	219	-	-	-	-	1	-	-	1	2	3	13
Asian or Pacific Islander 2/, male												
2004.....	117	-	-	-	-	1	-	-	1	1	5	5
2003.....	97	-	-	-	-	-	1	-	1	2	1	1
2002.....	95	-	-	-	-	-	-	1	1	1	5	2
2001.....	102	-	1	-	-	-	2	-	-	1	2	1
2000.....	93	-	-	-	-	-	2	1	-	4	2	1
1999.....	105	-	-	-	-	-	-	-	1	1	3	9
Asian or Pacific Islander 2/, female												
2004.....	130	-	-	-	-	-	1	-	1	2	1	1
2003.....	152	-	-	-	-	-	1	1	1	-	3	1
2002.....	124	-	-	-	-	-	1	-	-	1	-	1
2001.....	105	-	-	-	-	-	-	1	-	2	1	2
2000.....	92	-	-	-	1	-	1	-	-	-	-	1
1999.....	114	-	-	-	-	1	-	-	-	1	-	4

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic mitral valve disorders (I34)												
All races, both sexes												
2004.....	2,554	1	1	-	3	9	7	7	29	32	57	63
2003.....	2,542	-	3	-	3	5	10	19	23	30	48	48
2002.....	2,759	-	1	1	-	13	9	19	25	38	48	57
2001.....	2,785	1	-	-	2	9	16	26	28	40	53	66
2000.....	2,865	1	3	2	5	6	14	14	28	44	59	64
1999.....	2,895	-	-	-	4	9	15	19	35	40	71	66
All races, male												
2004.....	880	-	-	-	3	3	5	2	11	12	26	34
2003.....	893	-	1	-	2	3	7	4	11	13	21	20
2002.....	938	-	1	1	-	10	3	4	13	18	25	23
2001.....	939	-	-	-	1	1	6	12	10	17	22	35
2000.....	1,001	-	-	1	2	2	6	6	8	16	22	42
1999.....	1,010	-	-	-	1	1	4	5	14	25	30	29
All races, female												
2004.....	1,674	1	1	-	-	6	2	5	18	20	31	29
2003.....	1,649	-	2	-	1	2	3	15	12	17	27	28
2002.....	1,821	-	-	-	-	3	6	15	12	20	23	34
2001.....	1,846	1	-	-	1	8	10	14	18	23	31	31
2000.....	1,864	1	3	1	3	4	8	8	20	28	37	22
1999.....	1,885	-	-	-	3	8	11	14	21	15	41	37
White, both sexes												
2004.....	2,309	-	1	-	2	5	6	4	22	28	46	46
2003.....	2,316	-	3	-	3	5	8	15	14	23	35	41
2002.....	2,539	-	1	1	-	10	5	16	18	32	41	43
2001.....	2,564	-	-	-	2	6	11	21	25	30	43	53
2000.....	2,636	1	3	2	3	5	11	10	24	37	47	50
1999.....	2,643	-	-	-	3	7	11	13	30	37	57	48

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic mitral valve disorders (I34)												
White, male												
2004.....	791	-	-	-	2	1	4	1	10	10	20	25
2003.....	810	-	1	-	2	3	5	4	7	9	15	17
2002.....	856	-	1	1	-	7	1	4	8	12	23	20
2001.....	870	-	-	-	1	1	5	10	9	11	17	28
2000.....	928	-	-	1	1	2	6	5	6	13	17	33
1999.....	930	-	-	-	1	-	3	3	13	23	24	20
White, female												
2004.....	1,518	-	1	-	-	4	2	3	12	18	26	21
2003.....	1,506	-	2	-	1	2	3	11	7	14	20	24
2002.....	1,683	-	-	-	-	3	4	12	10	20	18	23
2001.....	1,694	-	-	-	1	5	6	11	16	19	26	25
2000.....	1,708	1	3	1	2	3	5	5	18	24	30	17
1999.....	1,713	-	-	-	2	7	8	10	17	14	33	28
Black, both sexes												
2004.....	191	1	-	-	1	4	1	3	6	3	10	13
2003.....	178	-	-	-	-	-	2	4	8	6	12	7
2002.....	173	-	-	-	-	2	4	3	6	6	6	12
2001.....	185	1	-	-	-	3	5	4	3	9	10	13
2000.....	192	-	-	-	1	1	2	4	4	7	11	14
1999.....	194	-	-	-	1	1	4	6	5	3	13	12
Black, male												
2004.....	66	-	-	-	1	2	1	1	1	2	5	5
2003.....	65	-	-	-	-	-	2	-	4	3	5	3
2002.....	61	-	-	-	-	2	2	-	4	6	1	2
2001.....	56	-	-	-	-	-	1	2	1	5	5	7
2000.....	59	-	-	-	1	-	-	1	2	3	4	9
1999.....	57	-	-	-	-	1	1	2	1	2	5	6

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic mitral valve disorders (I34)												
Asian or Pacific Islander 2/, both sexes												
2004.....	49	-	-	-	-	-	-	-	1	1	1	4
2003.....	39	-	-	-	-	-	-	-	1	1	-	-
2002.....	41	-	-	-	-	-	-	-	1	-	1	1
2001.....	31	-	-	-	-	-	-	1	-	1	-	-
2000.....	32	-	-	-	1	-	1	-	-	-	1	-
1999.....	48	-	-	-	-	1	-	-	-	-	1	4
Asian or Pacific Islander 2/, male												
2004.....	22	-	-	-	-	-	-	-	-	-	1	4
2003.....	15	-	-	-	-	-	-	-	-	1	-	-
2002.....	18	-	-	-	-	-	-	-	1	-	1	1
2001.....	13	-	-	-	-	-	-	-	-	1	-	-
2000.....	13	-	-	-	-	-	-	-	-	-	1	-
1999.....	18	-	-	-	-	-	-	-	-	-	1	3
Asian or Pacific Islander 2/, female												
2004.....	27	-	-	-	-	-	-	-	1	1	-	-
2003.....	24	-	-	-	-	-	-	-	1	-	-	-
2002.....	23	-	-	-	-	-	-	-	-	-	-	-
2001.....	18	-	-	-	-	-	-	1	-	-	-	-
2000.....	19	-	-	-	1	-	1	-	-	-	-	-
1999.....	30	-	-	-	-	1	-	-	-	-	-	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic aortic valve disorders (I35)												
All races, both sexes												
2004.....	12,665	1	5	-	2	6	8	14	20	25	64	100
2003.....	12,548	1	2	-	1	4	11	7	23	44	60	88
2002.....	12,471	4	2	2	3	2	7	12	17	39	70	98
2001.....	12,189	3	1	-	6	2	4	11	18	38	50	104
2000.....	12,380	4	2	-	3	7	6	21	16	38	67	99
1999.....	12,212	3	2	4	1	9	8	12	20	47	75	101
All races, male												
2004.....	4,941	-	4	-	1	5	6	9	14	18	49	75
2003.....	4,951	1	2	-	1	4	9	4	17	31	43	62
2002.....	4,909	3	2	1	2	2	5	9	15	26	46	75
2001.....	4,847	3	1	-	3	1	4	8	14	31	39	77
2000.....	4,836	2	2	-	2	5	5	13	11	24	49	75
1999.....	4,851	3	-	3	-	7	7	8	16	34	48	66
All races, female												
2004.....	7,724	1	1	-	1	1	2	5	6	7	15	25
2003.....	7,597	-	-	-	-	-	2	3	6	13	17	26
2002.....	7,562	1	-	1	1	-	2	3	2	13	24	23
2001.....	7,342	-	-	-	3	1	-	3	4	7	11	27
2000.....	7,544	2	-	-	1	2	1	8	5	14	18	24
1999.....	7,361	-	2	1	1	2	1	4	4	13	27	35
White, both sexes												
2004.....	11,904	1	3	-	2	3	6	8	19	18	49	85
2003.....	11,802	1	2	-	1	4	6	3	19	37	44	69
2002.....	11,725	3	2	2	2	2	5	9	11	25	59	74
2001.....	11,469	3	-	-	5	2	3	8	14	33	35	85
2000.....	11,633	4	2	-	2	7	3	17	12	24	51	77
1999.....	11,432	3	2	3	1	8	3	7	16	37	57	72

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Nonrheumatic aortic valve disorders (I35)

White, male

2004.....	4,608	-	2	-	1	2	5	4	13	15	40	64
2003.....	4,648	1	2	-	1	4	5	2	13	26	32	49
2002.....	4,620	2	2	1	1	2	4	7	9	21	40	57
2001.....	4,507	3	-	-	2	1	3	6	10	28	28	62
2000.....	4,520	2	2	-	2	5	2	11	8	16	39	59
1999.....	4,503	3	-	2	-	6	3	5	12	27	37	47

White, female

2004.....	7,296	1	1	-	1	1	1	4	6	3	9	21
2003.....	7,154	-	-	-	-	-	1	1	6	11	12	20
2002.....	7,105	1	-	1	1	-	1	2	2	4	19	17
2001.....	6,962	-	-	-	3	1	-	2	4	5	7	23
2000.....	7,113	2	-	-	-	2	1	6	4	8	12	18
1999.....	6,929	-	2	1	1	2	-	2	4	10	20	25

Black, both sexes

2004.....	590	-	2	-	-	2	-	6	1	7	12	14
2003.....	574	-	-	-	-	-	2	4	2	6	16	19
2002.....	585	1	-	-	-	-	1	2	5	13	9	22
2001.....	576	-	1	-	1	-	-	3	3	4	14	18
2000.....	614	-	-	-	1	-	2	3	4	9	15	20
1999.....	625	-	-	1	-	1	4	5	4	9	14	24

Black, male

2004.....	244	-	2	-	-	2	-	5	1	3	7	10
2003.....	225	-	-	-	-	-	2	2	2	4	11	13
2002.....	221	1	-	-	-	-	1	1	5	4	4	16
2001.....	257	-	1	-	1	-	-	2	3	3	11	14
2000.....	250	-	-	-	-	-	2	1	3	5	9	15
1999.....	271	-	-	1	-	1	3	3	4	7	8	15

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic aortic valve disorders (I35)												
Asian or Pacific Islander 2/, both sexes												
2004.....	140	-	-	-	-	1	1	-	-	-	3	1
2003.....	136	-	-	-	-	-	2	-	-	1	-	-
2002.....	125	-	-	-	-	-	1	1	-	-	2	1
2001.....	120	-	-	-	-	-	1	-	-	1	1	-
2000.....	107	-	-	-	-	-	1	1	-	3	1	1
1999.....	126	-	-	-	-	-	-	-	-	1	2	4
Asian or Pacific Islander 2/, male												
2004.....	71	-	-	-	-	1	-	-	-	-	2	1
2003.....	55	-	-	-	-	-	1	-	-	1	-	-
2002.....	51	-	-	-	-	-	-	1	-	-	2	1
2001.....	67	-	-	-	-	-	1	-	-	-	-	-
2000.....	59	-	-	-	-	-	1	1	-	3	1	1
1999.....	66	-	-	-	-	-	-	-	-	-	2	3
Asian or Pacific Islander 2/, female												
2004.....	69	-	-	-	-	-	1	-	-	-	1	-
2003.....	81	-	-	-	-	-	1	-	-	-	-	-
2002.....	74	-	-	-	-	-	1	-	-	-	-	-
2001.....	53	-	-	-	-	-	-	-	-	1	1	-
2000.....	48	-	-	-	-	-	-	-	-	-	-	-
1999.....	60	-	-	-	-	-	-	-	-	1	-	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

All other diseases of endocardium (I36-I38)

All races, both sexes

2004.....	5,041	-	2	-	-	8	12	13	29	52	73	162
2003.....	4,966	1	2	2	1	2	8	20	31	38	95	128
2002.....	4,759	2	3	1	-	3	6	22	25	50	101	131
2001.....	4,579	2	6	1	-	-	8	16	18	52	69	89
2000.....	4,492	3	2	2	2	3	10	13	27	59	73	123
1999.....	4,505	2	-	1	2	12	13	12	43	51	95	96

All races, male

2004.....	2,069	-	2	-	-	3	3	7	19	30	45	108
2003.....	1,923	-	1	-	1	1	4	9	13	23	58	72
2002.....	1,811	1	1	1	-	2	3	7	18	30	59	82
2001.....	1,808	2	2	-	-	-	3	12	8	27	44	54
2000.....	1,707	1	1	2	2	1	7	7	12	36	45	71
1999.....	1,725	1	-	-	-	9	9	5	21	27	57	49

All races, female

2004.....	2,972	-	-	-	-	5	9	6	10	22	28	54
2003.....	3,043	1	1	2	-	1	4	11	18	15	37	56
2002.....	2,948	1	2	-	-	1	3	15	7	20	42	49
2001.....	2,771	-	4	1	-	-	5	4	10	25	25	35
2000.....	2,785	2	1	-	-	2	3	6	15	23	28	52
1999.....	2,780	1	-	1	2	3	4	7	22	24	38	47

White, both sexes

2004.....	4,447	-	2	-	-	6	9	10	20	35	45	100
2003.....	4,351	1	2	1	1	1	4	15	20	30	55	79
2002.....	4,230	2	-	1	-	3	4	15	21	38	61	73
2001.....	4,057	1	2	1	-	-	5	7	8	34	48	45
2000.....	3,961	3	1	2	1	1	5	7	17	40	46	84
1999.....	3,992	2	-	1	1	8	13	9	33	34	60	53

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of endocardium (I36-I38)												
White, male												
2004.....	1,802	-	2	-	-	3	3	5	15	21	27	68
2003.....	1,651	-	1	-	1	1	2	8	10	18	37	45
2002.....	1,587	1	-	1	-	2	3	4	16	24	40	47
2001.....	1,577	1	1	-	-	-	1	5	4	18	32	30
2000.....	1,475	1	1	2	1	-	2	3	8	25	34	49
1999.....	1,507	1	-	-	-	7	9	3	16	18	40	26
White, female												
2004.....	2,645	-	-	-	-	3	6	5	5	14	18	32
2003.....	2,700	1	1	1	-	-	2	7	10	12	18	34
2002.....	2,643	1	-	-	-	1	1	11	5	14	21	26
2001.....	2,480	-	1	1	-	-	4	2	4	16	16	15
2000.....	2,486	2	-	-	-	1	3	4	9	15	12	35
1999.....	2,485	1	-	1	1	1	4	6	17	16	20	27
Black, both sexes												
2004.....	512	-	-	-	-	2	3	3	8	14	26	58
2003.....	526	-	-	1	-	1	4	4	10	8	34	45
2002.....	460	-	3	-	-	-	2	7	2	10	37	57
2001.....	454	1	3	-	-	-	2	9	10	16	18	40
2000.....	467	-	1	-	1	2	4	6	10	17	27	36
1999.....	451	-	-	-	-	4	-	3	9	15	35	37
Black, male												
2004.....	231	-	-	-	-	-	-	2	3	8	16	39
2003.....	238	-	-	-	-	-	2	1	2	5	19	25
2002.....	191	-	1	-	-	-	-	3	1	5	16	35
2001.....	203	1	-	-	-	-	1	7	4	8	9	22
2000.....	205	-	-	-	1	1	4	4	4	10	11	21
1999.....	187	-	-	-	-	2	-	2	4	8	17	19

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of endocardium (I36-I38)												
Black, female												
2004.....	281	-	-	-	-	2	3	1	5	6	10	19
2003.....	288	-	-	1	-	1	2	3	8	3	15	20
2002.....	269	-	2	-	-	-	2	4	1	5	21	22
2001.....	251	-	3	-	-	-	1	2	6	8	9	18
2000.....	262	-	1	-	-	1	-	2	6	7	16	15
1999.....	264	-	-	-	-	2	-	1	5	7	18	18
American Indian 1/, both sexes												
2004.....	24	-	-	-	-	-	-	-	-	1	-	3
2003.....	15	-	-	-	-	-	-	-	-	-	2	2
2002.....	16	-	-	-	-	-	-	-	2	-	1	-
2001.....	12	-	-	-	-	-	-	-	-	1	1	1
2000.....	18	-	-	-	-	-	-	-	-	1	-	2
1999.....	17	-	-	-	1	-	-	-	-	1	-	1
American Indian 1/, male												
2004.....	12	-	-	-	-	-	-	-	-	-	-	1
2003.....	7	-	-	-	-	-	-	-	-	-	1	1
2002.....	7	-	-	-	-	-	-	-	1	-	1	-
2001.....	6	-	-	-	-	-	-	-	-	1	1	1
2000.....	6	-	-	-	-	-	-	-	-	-	-	1
1999.....	10	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, female												
2004.....	12	-	-	-	-	-	-	-	-	1	-	2
2003.....	8	-	-	-	-	-	-	-	-	-	1	1
2002.....	9	-	-	-	-	-	-	-	1	-	-	-
2001.....	6	-	-	-	-	-	-	-	-	-	-	-
2000.....	12	-	-	-	-	-	-	-	-	1	-	1
1999.....	7	-	-	-	1	-	-	-	-	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiomyopathy (I42)												
All races, both sexes												
2004.....	25,580	116	38	24	47	103	166	228	336	576	813	1,166
2003.....	26,858	126	56	20	34	104	172	250	361	553	916	1,225
2002.....	27,728	118	34	23	49	115	161	262	387	623	946	1,247
2001.....	26,863	137	61	23	41	112	154	213	346	588	917	1,140
2000.....	27,213	118	39	25	52	102	168	232	376	591	912	1,128
1999.....	27,260	132	36	25	49	125	129	236	366	686	922	1,124
All races, male												
2004.....	14,863	56	21	6	26	74	129	169	249	402	576	845
2003.....	15,436	70	28	11	24	65	118	180	251	392	646	863
2002.....	16,062	56	18	10	34	84	118	191	278	403	673	899
2001.....	15,334	60	36	10	23	73	112	146	253	420	653	805
2000.....	15,696	54	22	15	38	74	115	153	267	410	660	781
1999.....	15,688	61	18	12	33	93	86	161	248	488	666	789
All races, female												
2004.....	10,717	60	17	18	21	29	37	59	87	174	237	321
2003.....	11,422	56	28	9	10	39	54	70	110	161	270	362
2002.....	11,666	62	16	13	15	31	43	71	109	220	273	348
2001.....	11,529	77	25	13	18	39	42	67	93	168	264	335
2000.....	11,517	64	17	10	14	28	53	79	109	181	252	347
1999.....	11,572	71	18	13	16	32	43	75	118	198	256	335
White, both sexes												
2004.....	20,697	86	26	19	31	64	112	135	190	356	522	755
2003.....	21,761	95	35	13	22	62	107	135	216	335	611	822
2002.....	22,516	88	29	18	33	71	91	146	222	373	620	825
2001.....	21,721	88	34	15	27	68	88	122	207	358	561	724
2000.....	21,945	90	28	16	33	55	97	130	225	354	585	710
1999.....	21,910	91	28	19	27	69	75	137	234	424	567	715

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiomyopathy (I42)												
White, male												
2004.....	12,004	42	18	5	18	44	91	107	145	258	379	560
2003.....	12,465	47	16	9	16	43	74	103	158	245	450	599
2002.....	13,014	39	16	7	22	51	74	109	163	250	454	614
2001.....	12,383	44	22	7	12	43	67	89	154	263	422	535
2000.....	12,596	41	13	9	26	38	72	90	162	253	425	521
1999.....	12,514	40	16	8	18	49	51	96	165	303	433	518
White, female												
2004.....	8,693	44	8	14	13	20	21	28	45	98	143	195
2003.....	9,296	48	19	4	6	19	33	32	58	90	161	223
2002.....	9,502	49	13	11	11	20	17	37	59	123	166	211
2001.....	9,338	44	12	8	15	25	21	33	53	95	139	189
2000.....	9,349	49	15	7	7	17	25	40	63	101	160	189
1999.....	9,396	51	12	11	9	20	24	41	69	121	134	197
Black, both sexes												
2004.....	4,327	18	10	5	14	37	47	77	128	197	269	372
2003.....	4,535	23	15	6	8	35	59	102	130	199	285	372
2002.....	4,616	23	2	3	14	37	63	106	147	230	303	389
2001.....	4,623	44	22	7	14	40	61	84	128	207	339	381
2000.....	4,710	22	9	8	16	42	65	98	133	214	301	393
1999.....	4,770	31	5	6	20	54	51	86	121	237	333	380
Black, male												
2004.....	2,526	9	3	1	6	30	34	51	89	128	181	259
2003.....	2,637	18	9	1	6	18	40	68	83	132	184	243
2002.....	2,679	12	2	2	10	27	39	75	101	141	202	265
2001.....	2,644	16	10	3	11	27	42	52	88	137	220	245
2000.....	2,742	9	7	5	10	32	40	60	91	138	214	241
1999.....	2,823	17	1	4	14	42	33	58	76	165	217	249

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiomyopathy (I42)												
Black, female												
2004.....	1,801	9	7	4	8	7	13	26	39	69	88	113
2003.....	1,898	5	6	5	2	17	19	34	47	67	101	129
2002.....	1,937	11	-	1	4	10	24	31	46	89	101	124
2001.....	1,979	28	12	4	3	13	19	32	40	70	119	136
2000.....	1,968	13	2	3	6	10	25	38	42	76	87	152
1999.....	1,947	14	4	2	6	12	18	28	45	72	116	131
American Indian 1/, both sexes												
2004.....	132	3	-	-	-	2	2	6	5	10	8	10
2003.....	148	1	1	-	1	2	-	2	5	8	8	12
2002.....	132	1	2	1	2	1	2	2	6	9	6	12
2001.....	124	3	1	1	-	-	2	5	2	8	8	12
2000.....	128	1	-	1	-	1	2	3	2	11	6	9
1999.....	134	2	3	-	1	-	-	4	4	6	7	9
American Indian 1/, male												
2004.....	73	1	-	-	-	-	-	5	4	8	4	6
2003.....	89	1	-	-	-	1	-	1	5	6	5	9
2002.....	90	1	-	-	2	1	2	1	4	7	5	10
2001.....	75	-	1	-	-	-	2	3	2	7	5	9
2000.....	85	-	-	1	-	1	-	2	1	8	6	7
1999.....	78	1	1	-	-	-	-	3	1	5	5	8
American Indian 1/, female												
2004.....	59	2	-	-	-	2	2	1	1	2	4	4
2003.....	59	-	1	-	1	1	-	1	-	2	3	3
2002.....	42	-	2	1	-	-	-	1	2	2	1	2
2001.....	49	3	-	1	-	-	-	2	-	1	3	3
2000.....	43	1	-	-	-	-	2	1	1	3	-	2
1999.....	56	1	2	-	1	-	-	1	3	1	2	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiomyopathy (I42)												
Asian or Pacific Islander 2/, both sexes												
2004.....	424	9	2	-	2	-	5	10	13	13	14	29
2003.....	414	7	5	1	3	5	6	11	10	11	12	19
2002.....	464	6	1	1	-	6	5	8	12	11	17	21
2001.....	395	2	4	-	-	4	3	2	9	15	9	23
2000.....	430	5	2	-	3	4	4	1	16	12	20	16
1999.....	446	8	-	-	1	2	3	9	7	19	15	20
Asian or Pacific Islander 2/, male												
2004.....	260	4	-	-	2	-	4	6	11	8	12	20
2003.....	245	4	3	1	2	3	4	8	5	9	7	12
2002.....	279	4	-	1	-	5	3	6	10	5	12	10
2001.....	232	-	3	-	-	3	1	2	9	13	6	16
2000.....	273	4	2	-	2	3	3	1	13	11	15	12
1999.....	273	3	-	-	1	2	2	4	6	15	11	14
Asian or Pacific Islander 2/, female												
2004.....	164	5	2	-	-	-	1	4	2	5	2	9
2003.....	169	3	2	-	1	2	2	3	5	2	5	7
2002.....	185	2	1	-	-	1	2	2	2	6	5	11
2001.....	163	2	1	-	-	1	2	-	-	2	3	7
2000.....	157	1	-	-	1	1	1	-	3	1	5	4
1999.....	173	5	-	-	-	-	1	5	1	4	4	6

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Conduction disorders and cardiac dysrhythmias (I44-I49)												
All races, both sexes												
2004.....	37,606	32	21	10	32	68	116	128	194	273	561	861
2003.....	39,104	32	22	18	32	78	99	112	197	304	561	877
2002.....	39,335	56	21	20	27	88	97	126	185	294	598	767
2001.....	38,554	38	30	16	42	60	96	109	174	287	495	806
2000.....	38,330	39	28	9	36	82	97	115	163	297	556	761
1999.....	39,976	39	40	17	37	84	98	130	187	291	550	805
All races, male												
2004.....	16,613	18	12	4	18	44	72	77	113	164	351	550
2003.....	17,178	17	13	15	20	53	61	62	117	178	354	546
2002.....	17,245	32	13	10	19	64	71	75	114	173	393	508
2001.....	16,714	19	17	7	21	44	69	68	103	173	322	521
2000.....	16,748	21	18	7	21	53	56	74	93	186	336	509
1999.....	17,683	24	24	11	23	49	61	79	116	173	344	551
All races, female												
2004.....	20,993	14	9	6	14	24	44	51	81	109	210	311
2003.....	21,926	15	9	3	12	25	38	50	80	126	207	331
2002.....	22,090	24	8	10	8	24	26	51	71	121	205	259
2001.....	21,840	19	13	9	21	16	27	41	71	114	173	285
2000.....	21,582	18	10	2	15	29	41	41	70	111	220	252
1999.....	22,293	15	16	6	14	35	37	51	71	118	206	254
White, both sexes												
2004.....	32,095	22	13	7	20	50	86	81	130	188	394	628
2003.....	33,525	15	13	12	20	55	71	72	133	225	412	642
2002.....	33,817	36	14	15	22	72	67	85	120	200	459	563
2001.....	33,151	28	23	11	28	44	63	78	117	183	362	596
2000.....	33,231	29	21	5	24	60	72	81	113	216	406	553
1999.....	34,726	25	31	10	30	67	66	91	126	197	425	592

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Conduction disorders and cardiac dysrhythmias (I44-I49)

White, male

2004.....	13,991	11	8	3	11	34	52	50	78	115	267	409
2003.....	14,556	8	8	10	13	35	44	40	88	135	272	418
2002.....	14,673	26	8	8	14	55	49	54	75	120	317	386
2001.....	14,173	15	11	5	18	33	46	47	73	115	249	402
2000.....	14,399	15	15	4	15	41	44	55	69	136	259	385
1999.....	15,172	16	21	7	18	40	41	57	83	126	278	429

White, female

2004.....	18,104	11	5	4	9	16	34	31	52	73	127	219
2003.....	18,969	7	5	2	7	20	27	32	45	90	140	224
2002.....	19,144	10	6	7	8	17	18	31	45	80	142	177
2001.....	18,978	13	12	6	10	11	17	31	44	68	113	194
2000.....	18,832	14	6	1	9	19	28	26	44	80	147	168
1999.....	19,554	9	10	3	12	27	25	34	43	71	147	163

Black, both sexes

2004.....	4,664	9	7	3	11	16	24	39	52	82	152	210
2003.....	4,730	14	7	5	12	20	24	37	55	68	139	217
2002.....	4,732	16	6	4	4	11	27	34	55	84	128	184
2001.....	4,682	7	6	5	11	15	30	27	52	95	116	196
2000.....	4,486	10	7	3	8	19	23	29	46	74	135	197
1999.....	4,708	13	8	7	6	16	30	38	55	92	115	203

Black, male

2004.....	2,185	6	4	1	6	10	15	20	25	47	74	125
2003.....	2,175	8	5	4	7	15	15	20	22	35	74	120
2002.....	2,147	5	4	1	4	6	19	15	33	47	69	110
2001.....	2,165	4	5	2	3	10	21	19	28	53	59	108
2000.....	2,033	6	3	2	4	9	10	18	22	46	67	118
1999.....	2,215	7	3	4	5	9	19	21	29	46	61	118

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Conduction disorders and cardiac dysrhythmias (I44-I49)

Black, female

2004.....	2,479	3	3	2	5	6	9	19	27	35	78	85
2003.....	2,555	6	2	1	5	5	9	17	33	33	65	97
2002.....	2,585	11	2	3	-	5	8	19	22	37	59	74
2001.....	2,517	3	1	3	8	5	9	8	24	42	57	88
2000.....	2,453	4	4	1	4	10	13	11	24	28	68	79
1999.....	2,493	6	5	3	1	7	11	17	26	46	54	85

American Indian 1/, both sexes

2004.....	122	1	-	-	1	2	-	2	2	2	4	7
2003.....	144	1	1	-	-	1	2	-	2	3	2	6
2002.....	113	2	1	-	-	2	-	-	1	-	4	4
2001.....	142	2	1	-	-	-	-	-	3	4	5	7
2000.....	135	-	-	-	2	-	1	1	3	3	6	1
1999.....	127	1	-	-	-	-	1	-	1	1	6	6

American Indian 1/, male

2004.....	56	1	-	-	1	-	-	2	1	1	3	5
2003.....	73	1	-	-	-	1	2	-	1	3	1	1
2002.....	62	-	1	-	-	1	-	-	1	-	2	3
2001.....	76	-	1	-	-	-	-	-	1	2	4	5
2000.....	67	-	-	-	1	-	1	-	2	1	3	1
1999.....	75	1	-	-	-	-	-	-	1	1	4	3

American Indian 1/, female

2004.....	66	-	-	-	-	2	-	-	1	1	1	2
2003.....	71	-	1	-	-	-	-	-	1	-	1	5
2002.....	51	2	-	-	-	1	-	-	-	-	2	1
2001.....	66	2	-	-	-	-	-	-	2	2	1	2
2000.....	68	-	-	-	1	-	-	1	1	2	3	-
1999.....	52	-	-	-	-	-	1	-	-	-	2	3

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Asian or Pacific Islander 2/, both sexes												
2004.....	725	-	1	-	-	-	6	6	10	1	11	16
2003.....	705	2	1	1	-	2	2	3	7	8	8	12
2002.....	673	2	-	1	1	3	3	7	9	10	7	16
2001.....	579	1	-	-	3	1	3	4	2	5	12	7
2000.....	478	-	-	1	2	3	1	4	1	4	9	10
1999.....	415	-	1	-	1	1	1	1	5	1	4	4
Asian or Pacific Islander 2/, male												
2004.....	381	-	-	-	-	-	5	5	9	1	7	11
2003.....	374	-	-	1	-	2	-	2	6	5	7	7
2002.....	363	1	-	1	1	2	3	6	5	6	5	9
2001.....	300	-	-	-	-	1	2	2	1	3	10	6
2000.....	249	-	-	1	1	3	1	1	-	3	7	5
1999.....	221	-	-	-	-	-	1	1	3	-	1	1
Asian or Pacific Islander 2/, female												
2004.....	344	-	1	-	-	-	1	1	1	-	4	5
2003.....	331	2	1	-	-	-	2	1	1	3	1	5
2002.....	310	1	-	-	-	1	-	1	4	4	2	7
2001.....	279	1	-	-	3	-	1	2	1	2	2	1
2000.....	229	-	-	-	1	-	-	3	1	1	2	5
1999.....	194	-	1	-	1	1	-	-	2	1	3	3

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Heart failure (I50)												
All races, both sexes												
2004.....	57,120	30	11	3	7	12	21	38	52	110	203	386
2003.....	57,448	38	5	5	7	18	20	28	69	91	224	358
2002.....	56,494	23	11	3	7	10	17	33	68	102	180	355
2001.....	56,934	24	17	8	7	19	15	41	51	115	192	353
2000.....	55,704	31	9	10	11	21	13	27	36	77	185	283
1999.....	54,913	11	10	5	7	14	17	27	38	88	184	298
All races, male												
2004.....	22,292	16	4	1	6	6	13	28	31	69	117	237
2003.....	22,427	28	3	4	3	13	11	16	44	50	143	223
2002.....	21,698	14	6	1	4	6	9	22	38	56	117	210
2001.....	21,632	16	5	5	3	11	5	25	31	76	106	227
2000.....	21,175	22	7	4	5	11	9	15	21	40	119	191
1999.....	20,755	5	7	3	3	12	11	13	19	52	112	186
All races, female												
2004.....	34,828	14	7	2	1	6	8	10	21	41	86	149
2003.....	35,021	10	2	1	4	5	9	12	25	41	81	135
2002.....	34,796	9	5	2	3	4	8	11	30	46	63	145
2001.....	35,302	8	12	3	4	8	10	16	20	39	86	126
2000.....	34,529	9	2	6	6	10	4	12	15	37	66	92
1999.....	34,158	6	3	2	4	2	6	14	19	36	72	112
White, both sexes												
2004.....	51,205	20	10	-	5	7	14	22	28	62	131	246
2003.....	51,454	20	4	3	4	9	11	19	39	56	122	229
2002.....	50,672	15	8	3	4	5	11	16	37	61	113	211
2001.....	51,252	19	15	5	6	13	5	25	32	76	112	237
2000.....	50,142	19	5	7	9	12	5	15	23	44	120	180
1999.....	49,317	8	8	3	6	9	9	13	21	58	116	183

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Heart failure (I50)												
White, male												
2004.....	19,806	10	4	-	4	4	9	17	13	44	76	147
2003.....	19,987	13	2	2	2	5	7	13	25	30	86	151
2002.....	19,335	10	5	1	3	2	5	11	19	34	78	130
2001.....	19,332	12	5	3	2	9	1	12	19	49	69	166
2000.....	18,949	14	5	3	4	5	5	9	13	25	84	127
1999.....	18,469	3	5	2	3	8	6	6	11	34	77	116
White, female												
2004.....	31,399	10	6	-	1	3	5	5	15	18	55	99
2003.....	31,467	7	2	1	2	4	4	6	14	26	36	78
2002.....	31,337	5	3	2	1	3	6	5	18	27	35	81
2001.....	31,920	7	10	2	4	4	4	13	13	27	43	71
2000.....	31,193	5	-	4	5	7	-	6	10	19	36	53
1999.....	30,848	5	3	1	3	1	3	7	10	24	39	67
Black, both sexes												
2004.....	5,230	8	1	2	2	5	6	15	21	46	65	135
2003.....	5,330	15	-	2	3	9	9	8	29	34	97	124
2002.....	5,186	8	3	-	3	3	6	16	29	40	64	140
2001.....	5,122	5	2	3	-	6	10	16	18	35	77	111
2000.....	4,965	12	4	2	2	8	5	12	12	32	62	98
1999.....	4,998	3	2	2	1	4	8	13	17	28	66	111
Black, male												
2004.....	2,174	5	-	1	2	2	4	10	15	23	36	86
2003.....	2,150	13	-	2	1	8	4	2	18	20	53	68
2002.....	2,069	4	1	-	1	2	4	10	17	22	37	78
2001.....	2,052	4	-	2	-	2	4	13	11	26	35	57
2000.....	1,951	8	2	1	1	5	3	6	7	14	33	60
1999.....	2,021	2	2	1	-	3	5	7	8	16	35	68

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Heart failure (I50)												
Asian or Pacific Islander 2/, both sexes												
2004.....	492	2	-	-	-	-	-	1	1	1	3	4
2003.....	484	2	1	-	-	-	-	1	-	1	3	2
2002.....	452	-	-	-	-	1	-	-	1	-	1	2
2001.....	396	-	-	-	1	-	-	-	1	2	2	2
2000.....	422	-	-	1	-	1	2	-	1	-	3	3
1999.....	428	-	-	-	-	-	-	1	-	-	-	2
Asian or Pacific Islander 2/, male												
2004.....	225	1	-	-	-	-	-	1	1	1	2	3
2003.....	215	1	1	-	-	-	-	1	-	-	3	2
2002.....	202	-	-	-	-	1	-	-	1	-	-	-
2001.....	177	-	-	-	1	-	-	-	1	1	2	1
2000.....	196	-	-	-	-	1	-	-	1	-	2	3
1999.....	200	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, female												
2004.....	267	1	-	-	-	-	-	-	-	-	1	1
2003.....	269	1	-	-	-	-	-	-	-	1	-	-
2002.....	250	-	-	-	-	-	-	-	-	-	1	2
2001.....	219	-	-	-	-	-	-	-	-	1	-	1
2000.....	226	-	-	1	-	-	2	-	-	-	1	-
1999.....	228	-	-	-	-	-	-	1	-	-	-	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Congestive heart failure (I50.0)												
All races, both sexes												
2004.....	53,191	17	7	1	3	6	14	29	45	92	159	324
2003.....	53,595	24	3	3	4	12	13	22	56	75	179	298
2002.....	52,434	14	7	1	4	5	10	27	56	86	147	279
2001.....	52,828	13	9	5	4	15	11	30	42	90	152	299
2000.....	51,546	10	6	4	7	13	8	20	28	57	150	231
1999.....	50,824	4	7	5	4	10	12	20	26	67	146	235
All races, male												
2004.....	20,522	10	3	-	2	4	8	22	28	57	91	195
2003.....	20,672	15	2	3	3	9	10	13	32	38	109	184
2002.....	19,900	9	2	-	2	3	7	18	34	46	93	168
2001.....	19,805	10	3	4	2	8	3	17	26	60	85	192
2000.....	19,384	7	4	2	4	6	5	10	16	26	91	155
1999.....	18,987	1	4	3	2	9	7	9	15	41	89	148
All races, female												
2004.....	32,669	7	4	1	1	2	6	7	17	35	68	129
2003.....	32,923	9	1	-	1	3	3	9	24	37	70	114
2002.....	32,534	5	5	1	2	2	3	9	22	40	54	111
2001.....	33,023	3	6	1	2	7	8	13	16	30	67	107
2000.....	32,162	3	2	2	3	7	3	10	12	31	59	76
1999.....	31,837	3	3	2	2	1	5	11	11	26	57	87
White, both sexes												
2004.....	47,906	13	7	-	2	4	10	17	23	52	104	206
2003.....	48,250	14	2	2	3	5	8	13	31	46	92	184
2002.....	47,228	9	5	1	2	3	7	12	29	50	90	160
2001.....	47,724	11	8	3	3	9	4	18	28	57	85	197
2000.....	46,583	6	3	2	5	8	4	11	19	32	97	142
1999.....	45,798	3	6	3	4	8	6	7	14	43	94	135

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Congestive heart failure (I50.0)												
White, male												
2004.....	18,336	8	3	-	1	3	5	13	12	37	61	123
2003.....	18,533	8	1	2	2	3	7	10	17	22	62	121
2002.....	17,816	6	2	-	1	2	4	9	16	27	61	101
2001.....	17,782	8	3	2	1	6	-	8	18	38	52	138
2000.....	17,440	4	3	1	3	4	4	6	10	16	62	99
1999.....	16,954	1	3	2	2	7	4	2	9	24	62	87
White, female												
2004.....	29,570	5	4	-	1	1	5	4	11	15	43	83
2003.....	29,717	6	1	-	1	2	1	3	14	24	30	63
2002.....	29,412	3	3	1	1	1	3	3	13	23	29	59
2001.....	29,942	3	5	1	2	3	4	10	10	19	33	59
2000.....	29,143	2	-	1	2	4	-	5	9	16	35	43
1999.....	28,844	2	3	1	2	1	2	5	5	19	32	48
Black, both sexes												
2004.....	4,658	3	-	-	1	2	4	11	19	38	49	114
2003.....	4,748	8	-	1	1	7	5	8	24	28	84	109
2002.....	4,647	5	2	-	2	1	3	15	26	36	55	117
2001.....	4,599	2	1	2	-	6	7	12	13	30	64	99
2000.....	4,427	4	3	1	2	5	1	9	9	24	50	84
1999.....	4,488	1	1	2	-	2	6	13	12	22	51	96
Black, male												
2004.....	1,904	1	-	-	1	1	3	8	13	18	26	69
2003.....	1,883	6	-	1	1	6	3	2	14	16	45	59
2002.....	1,831	3	-	-	1	-	3	9	17	19	30	66
2001.....	1,802	2	-	2	-	2	3	9	7	21	31	52
2000.....	1,701	3	1	1	1	2	-	4	6	9	27	52
1999.....	1,797	-	1	1	-	2	3	7	6	15	27	59

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other and unspecified heart failure (I50.1-I50.9)

All races, both sexes

2004.....	3,929	13	4	2	4	6	7	9	7	18	44	62
2003.....	3,853	14	2	2	3	6	7	6	13	16	45	60
2002.....	4,060	9	4	2	3	5	7	6	12	16	33	76
2001.....	4,106	11	8	3	3	4	4	11	9	25	40	54
2000.....	4,158	21	3	6	4	8	5	7	8	20	35	52
1999.....	4,089	7	3	-	3	4	5	7	12	21	38	63

All races, male

2004.....	1,770	6	1	1	4	2	5	6	3	12	26	42
2003.....	1,755	13	1	1	-	4	1	3	12	12	34	39
2002.....	1,798	5	4	1	2	3	2	4	4	10	24	42
2001.....	1,827	6	2	1	1	3	2	8	5	16	21	35
2000.....	1,791	15	3	2	1	5	4	5	5	14	28	36
1999.....	1,768	4	3	-	1	3	4	4	4	11	23	38

All races, female

2004.....	2,159	7	3	1	-	4	2	3	4	6	18	20
2003.....	2,098	1	1	1	3	2	6	3	1	4	11	21
2002.....	2,262	4	-	1	1	2	5	2	8	6	9	34
2001.....	2,279	5	6	2	2	1	2	3	4	9	19	19
2000.....	2,367	6	-	4	3	3	1	2	3	6	7	16
1999.....	2,321	3	-	-	2	1	1	3	8	10	15	25

White, both sexes

2004.....	3,299	7	3	-	3	3	4	5	5	10	27	40
2003.....	3,204	6	2	1	1	4	3	6	8	10	30	45
2002.....	3,444	6	3	2	2	2	4	4	8	11	23	51
2001.....	3,528	8	7	2	3	4	1	7	4	19	27	40
2000.....	3,559	13	2	5	4	4	1	4	4	12	23	38
1999.....	3,519	5	2	-	2	1	3	6	7	15	22	48

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified heart failure (I50.1-I50.9)												
White, male												
2004.....	1,470	2	1	-	3	1	4	4	1	7	15	24
2003.....	1,454	5	1	-	-	2	-	3	8	8	24	30
2002.....	1,519	4	3	1	2	-	1	2	3	7	17	29
2001.....	1,550	4	2	1	1	3	1	4	1	11	17	28
2000.....	1,509	10	2	2	1	1	1	3	3	9	22	28
1999.....	1,515	2	2	-	1	1	2	4	2	10	15	29
White, female												
2004.....	1,829	5	2	-	-	2	-	1	4	3	12	16
2003.....	1,750	1	1	1	1	2	3	3	-	2	6	15
2002.....	1,925	2	-	1	-	2	3	2	5	4	6	22
2001.....	1,978	4	5	1	2	1	-	3	3	8	10	12
2000.....	2,050	3	-	3	3	3	-	1	1	3	1	10
1999.....	2,004	3	-	-	1	-	1	2	5	5	7	19
Black, both sexes												
2004.....	572	5	1	2	1	3	2	4	2	8	16	21
2003.....	582	7	-	1	2	2	4	-	5	6	13	15
2002.....	539	3	1	-	1	2	3	1	3	4	9	23
2001.....	523	3	1	1	-	-	3	4	5	5	13	12
2000.....	538	8	1	1	-	3	4	3	3	8	12	14
1999.....	510	2	1	-	1	2	2	-	5	6	15	15
Black, male												
2004.....	270	4	-	1	1	1	1	2	2	5	10	17
2003.....	267	7	-	1	-	2	1	-	4	4	8	9
2002.....	238	1	1	-	-	2	1	1	-	3	7	12
2001.....	250	2	-	-	-	-	1	4	4	5	4	5
2000.....	250	5	1	-	-	3	3	2	1	5	6	8
1999.....	224	2	1	-	-	1	2	-	2	1	8	9

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified heart failure (I50.1-I50.9)												
Asian or Pacific Islander 2/, both sexes												
2004.....	40	1	-	-	-	-	-	-	-	-	1	-
2003.....	43	1	-	-	-	-	-	-	-	-	1	-
2002.....	50	-	-	-	-	-	-	-	-	-	1	1
2001.....	29	-	-	-	-	-	-	-	-	-	-	-
2000.....	40	-	-	-	-	1	-	-	1	-	-	-
1999.....	38	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	22	-	-	-	-	-	-	-	-	-	1	-
2003.....	20	1	-	-	-	-	-	-	-	-	1	-
2002.....	26	-	-	-	-	-	-	-	-	-	-	-
2001.....	12	-	-	-	-	-	-	-	-	-	-	-
2000.....	21	-	-	-	-	1	-	-	1	-	-	-
1999.....	18	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	18	1	-	-	-	-	-	-	-	-	-	-
2003.....	23	-	-	-	-	-	-	-	-	-	-	-
2002.....	24	-	-	-	-	-	-	-	-	-	1	1
2001.....	17	-	-	-	-	-	-	-	-	-	-	-
2000.....	19	-	-	-	-	-	-	-	-	-	-	-
1999.....	20	-	-	-	-	-	-	1	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
-------------	-----------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)

All races, both sexes

2004.....	653	25	30	16	12	22	31	27	27	42	58	67
2003.....	647	30	25	10	22	29	28	33	34	50	44	53
2002.....	881	32	21	5	11	35	27	25	36	60	75	63
2001.....	733	23	39	13	12	24	23	36	41	70	63	54
2000.....	687	22	22	10	14	28	24	33	44	55	59	54
1999.....	725	26	34	11	13	36	37	32	47	59	62	51

All races, male

2004.....	324	13	15	5	9	18	21	16	14	27	32	34
2003.....	322	18	8	4	16	20	19	17	23	27	14	22
2002.....	430	19	10	3	6	16	14	17	21	37	40	34
2001.....	377	13	22	5	5	13	13	16	18	45	32	27
2000.....	337	13	11	6	8	16	11	19	25	30	33	23
1999.....	376	17	18	6	9	19	24	20	32	39	30	25

All races, female

2004.....	329	12	15	11	3	4	10	11	13	15	26	33
2003.....	325	12	17	6	6	9	9	16	11	23	30	31
2002.....	451	13	11	2	5	19	13	8	15	23	35	29
2001.....	356	10	17	8	7	11	10	20	23	25	31	27
2000.....	350	9	11	4	6	12	13	14	19	25	26	31
1999.....	349	9	16	5	4	17	13	12	15	20	32	26

White, both sexes

2004.....	477	13	13	7	9	13	25	20	18	33	41	45
2003.....	505	17	13	6	18	20	18	25	22	34	37	39
2002.....	694	17	16	3	6	22	18	20	30	38	50	50
2001.....	546	9	22	7	8	16	18	22	23	58	44	38
2000.....	542	15	13	7	6	21	19	22	36	41	42	37
1999.....	555	12	17	8	9	28	27	21	29	38	49	40

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)

White, male

2004.....	236	8	7	2	8	10	16	13	12	21	22	21
2003.....	244	9	4	3	12	13	9	14	14	19	12	17
2002.....	330	8	9	2	3	11	8	13	17	23	29	24
2001.....	277	5	10	3	4	7	9	10	12	37	24	17
2000.....	265	9	8	5	4	12	9	13	22	23	24	16
1999.....	279	8	8	5	7	14	19	14	21	24	25	19

White, female

2004.....	241	5	6	5	1	3	9	7	6	12	19	24
2003.....	261	8	9	3	6	7	9	11	8	15	25	22
2002.....	364	9	7	1	3	11	10	7	13	15	21	26
2001.....	269	4	12	4	4	9	9	12	11	21	20	21
2000.....	277	6	5	2	2	9	10	9	14	18	18	21
1999.....	276	4	9	3	2	14	8	7	8	14	24	21

Black, both sexes

2004.....	158	11	15	8	3	8	6	6	9	7	16	20
2003.....	126	12	11	4	3	8	9	6	8	15	6	14
2002.....	168	15	3	2	5	12	7	5	6	19	22	13
2001.....	161	12	14	5	3	5	4	13	17	9	18	14
2000.....	129	7	8	3	6	7	5	10	7	14	15	15
1999.....	147	13	15	3	3	7	10	9	16	20	12	11

Black, male

2004.....	80	4	8	3	1	7	5	3	2	4	10	12
2003.....	68	8	4	1	3	6	9	3	5	8	2	5
2002.....	91	11	1	1	3	4	4	4	4	12	10	10
2001.....	84	6	12	1	1	4	3	6	6	5	8	8
2000.....	62	4	3	1	3	4	2	5	2	7	7	6
1999.....	84	8	9	1	1	5	5	5	9	14	5	6

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Black, female												
2004.....	78	7	7	5	2	1	1	3	7	3	6	8
2003.....	58	4	7	3	-	2	-	3	3	7	4	9
2002.....	77	4	2	1	2	8	3	1	2	7	12	3
2001.....	77	6	2	4	2	1	1	7	11	4	10	6
2000.....	67	3	5	2	3	3	3	5	5	7	8	9
1999.....	63	5	6	2	2	2	5	4	7	6	7	5
American Indian 1/, both sexes												
2004.....	6	-	-	-	-	1	-	1	-	-	-	2
2003.....	2	-	1	-	-	-	1	-	-	-	-	-
2002.....	3	-	1	-	-	-	1	-	-	-	1	-
2001.....	9	1	1	1	-	1	-	1	-	-	1	1
2000.....	6	-	1	-	2	-	-	-	1	-	-	1
1999.....	5	-	-	-	1	1	-	-	-	-	-	-
American Indian 1/, male												
2004.....	2	-	-	-	-	1	-	-	-	-	-	1
2003.....	1	-	-	-	-	-	1	-	-	-	-	-
2002.....	1	-	-	-	-	-	1	-	-	-	-	-
2001.....	4	1	-	1	-	-	-	-	-	-	-	1
2000.....	2	-	-	-	1	-	-	-	1	-	-	-
1999.....	3	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, female												
2004.....	4	-	-	-	-	-	-	1	-	-	-	1
2003.....	1	-	1	-	-	-	-	-	-	-	-	-
2002.....	2	-	1	-	-	-	-	-	-	-	1	-
2001.....	5	-	1	-	-	1	-	1	-	-	1	-
2000.....	4	-	1	-	1	-	-	-	-	-	-	1
1999.....	2	-	-	-	-	1	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Asian or Pacific Islander 2/, both sexes												
2004.....	12	1	2	1	-	-	-	-	-	2	1	-
2003.....	14	1	-	-	1	1	-	2	4	1	1	-
2002.....	16	-	1	-	-	1	1	-	-	3	2	-
2001.....	17	1	2	-	1	2	1	-	1	3	-	1
2000.....	10	-	-	-	-	-	-	1	-	-	2	1
1999.....	18	1	2	-	-	-	-	2	2	1	1	-
Asian or Pacific Islander 2/, male												
2004.....	6	1	-	-	-	-	-	-	-	2	-	-
2003.....	9	1	-	-	1	1	-	-	4	-	-	-
2002.....	8	-	-	-	-	1	1	-	-	2	1	-
2001.....	12	1	-	-	-	2	1	-	-	3	-	1
2000.....	8	-	-	-	-	-	-	1	-	-	2	1
1999.....	10	1	1	-	-	-	-	1	2	1	-	-
Asian or Pacific Islander 2/, female												
2004.....	6	-	2	1	-	-	-	-	-	-	1	-
2003.....	5	-	-	-	-	-	-	2	-	1	1	-
2002.....	8	-	1	-	-	-	-	-	-	1	1	-
2001.....	5	-	2	-	1	-	-	-	1	-	-	-
2000.....	2	-	-	-	-	-	-	-	-	-	-	-
1999.....	8	-	1	-	-	-	-	1	-	-	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

All other and ill-defined forms of heart disease (I50.0-I51.3,I51.6-I51.9)

All races, both sexes

2004.....	9,959	24	16	7	16	30	66	89	104	156	306	381
2003.....	9,769	22	10	6	14	33	69	57	120	159	299	362
2002.....	9,389	24	12	5	13	36	56	56	95	185	291	363
2001.....	8,975	18	14	7	14	28	54	60	116	184	272	347
2000.....	8,771	31	6	10	5	35	54	65	103	140	253	318
1999.....	8,092	30	10	4	12	35	43	49	103	170	230	277

All races, male

2004.....	5,006	19	7	4	9	23	53	72	80	110	205	265
2003.....	4,849	8	4	4	8	25	55	48	92	121	222	243
2002.....	4,688	16	6	2	9	35	44	40	76	133	201	249
2001.....	4,420	10	7	3	10	24	43	46	87	139	193	250
2000.....	4,348	14	6	7	2	30	46	46	84	100	185	232
1999.....	4,063	15	7	2	7	26	33	40	82	125	161	202

All races, female

2004.....	4,953	5	9	3	7	7	13	17	24	46	101	116
2003.....	4,920	14	6	2	6	8	14	9	28	38	77	119
2002.....	4,701	8	6	3	4	1	12	16	19	52	90	114
2001.....	4,555	8	7	4	4	4	11	14	29	45	79	97
2000.....	4,423	17	-	3	3	5	8	19	19	40	68	86
1999.....	4,029	15	3	2	5	9	10	9	21	45	69	75

White, both sexes

2004.....	8,425	19	10	6	7	18	47	62	71	119	227	278
2003.....	8,290	14	9	4	9	23	44	35	86	114	221	267
2002.....	7,934	18	8	3	8	21	39	41	62	141	213	261
2001.....	7,539	15	8	6	8	22	34	42	79	130	192	243
2000.....	7,373	20	4	4	3	19	29	39	77	95	180	244
1999.....	6,797	21	7	3	7	25	26	33	70	120	156	200

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

All other and ill-defined forms of heart disease (I50.0-I51.3,I51.6-I51.9)

White, male

2004.....	4,197	14	2	3	5	13	38	47	58	89	155	207
2003.....	4,096	6	3	2	7	15	36	32	73	87	167	189
2002.....	3,922	11	5	1	6	20	32	28	53	105	147	187
2001.....	3,699	10	4	2	7	19	27	35	58	101	150	182
2000.....	3,606	9	4	3	1	17	25	27	67	75	138	183
1999.....	3,392	12	6	1	3	19	21	28	56	95	112	161

White, female

2004.....	4,228	5	8	3	2	5	9	15	13	30	72	71
2003.....	4,194	8	6	2	2	8	8	3	13	27	54	78
2002.....	4,012	7	3	2	2	1	7	13	9	36	66	74
2001.....	3,840	5	4	4	1	3	7	7	21	29	42	61
2000.....	3,767	11	-	1	2	2	4	12	10	20	42	61
1999.....	3,405	9	1	2	4	6	5	5	14	25	44	39

Black, both sexes

2004.....	1,396	4	6	1	9	10	15	25	30	32	76	97
2003.....	1,321	7	1	2	4	10	24	21	32	42	69	85
2002.....	1,302	6	4	2	5	15	16	14	30	40	66	91
2001.....	1,278	3	6	1	4	6	18	17	33	45	76	91
2000.....	1,279	8	2	5	2	15	22	23	24	42	68	68
1999.....	1,176	8	3	1	4	9	15	16	32	43	71	72

Black, male

2004.....	731	4	5	1	4	8	11	23	20	18	48	55
2003.....	672	2	1	2	1	10	18	15	17	33	46	47
2002.....	676	5	1	1	3	15	12	12	20	24	43	55
2001.....	628	-	3	1	2	5	14	11	25	31	41	58
2000.....	674	3	2	3	1	12	18	17	16	24	43	45
1999.....	615	2	1	1	3	7	10	12	25	27	47	40

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

All other and ill-defined forms of heart disease (I50.0-I51.3,I51.6-I51.9)

Asian or Pacific Islander 2/, both sexes

2004.....	109	1	-	-	-	2	3	1	3	4	2	5
2003.....	108	1	-	-	-	-	1	1	-	-	5	9
2002.....	114	-	-	-	-	-	1	1	2	2	8	8
2001.....	97	-	-	-	2	-	1	-	3	1	-	8
2000.....	83	2	-	1	-	-	3	3	-	1	2	4
1999.....	88	1	-	-	1	1	1	-	1	3	3	4

Asian or Pacific Islander 2/, male

2004.....	58	1	-	-	-	2	3	1	2	2	1	2
2003.....	59	-	-	-	-	-	1	1	-	-	5	6
2002.....	60	-	-	-	-	-	-	-	2	2	7	5
2001.....	54	-	-	-	1	-	1	-	3	1	-	5
2000.....	45	1	-	1	-	-	3	2	-	1	1	2
1999.....	41	1	-	-	1	-	1	-	1	1	2	1

Asian or Pacific Islander 2/, female

2004.....	51	-	-	-	-	-	-	-	1	2	1	3
2003.....	49	1	-	-	-	-	-	-	-	-	-	3
2002.....	54	-	-	-	-	-	1	1	-	-	1	3
2001.....	43	-	-	-	1	-	-	-	-	-	-	3
2000.....	38	1	-	-	-	-	-	1	-	-	1	2
1999.....	47	-	-	-	-	1	-	-	-	2	1	3

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebrovascular diseases (I60-I69)												
All races, both sexes												
2004.....	150,074	127	41	34	43	69	142	208	359	719	1,642	2,622
2003.....	157,689	101	46	29	40	88	133	188	395	761	1,699	2,577
2002.....	162,672	117	53	33	58	53	118	183	384	797	1,628	2,615
2001.....	163,538	108	54	38	42	68	128	199	402	840	1,651	2,507
2000.....	167,661	124	47	25	51	67	132	209	393	880	1,719	2,625
1999.....	167,366	103	46	26	39	67	115	202	378	916	1,658	2,455
All races, male												
2004.....	58,800	71	23	18	22	41	75	112	172	389	851	1,394
2003.....	61,426	57	25	19	22	51	65	101	213	411	872	1,389
2002.....	62,622	67	30	16	31	34	69	100	184	389	789	1,402
2001.....	63,177	63	25	21	21	36	67	105	206	435	832	1,354
2000.....	64,769	74	19	12	21	38	68	111	197	430	867	1,401
1999.....	64,485	65	22	17	17	36	59	102	197	449	859	1,282
All races, female												
2004.....	91,274	56	18	16	21	28	67	96	187	330	791	1,228
2003.....	96,263	44	21	10	18	37	68	87	182	350	827	1,188
2002.....	100,050	50	23	17	27	19	49	83	200	408	839	1,213
2001.....	100,361	45	29	17	21	32	61	94	196	405	819	1,153
2000.....	102,892	50	28	13	30	29	64	98	196	450	852	1,224
1999.....	102,881	38	24	9	22	31	56	100	181	467	799	1,173
White, both sexes												
2004.....	127,868	79	28	26	32	50	102	140	233	492	1,065	1,656
2003.....	134,705	63	30	21	30	63	89	126	262	457	1,051	1,611
2002.....	139,719	71	40	25	38	38	92	124	252	509	1,038	1,671
2001.....	140,465	60	39	29	30	53	85	133	283	553	1,052	1,597
2000.....	144,580	75	30	21	34	51	97	160	260	558	1,087	1,627
1999.....	144,827	70	34	23	26	46	89	145	257	602	1,044	1,535

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebrovascular diseases (I60-I69)												
White, male												
2004.....	49,296	45	18	14	15	27	55	85	113	265	577	883
2003.....	51,646	32	14	13	18	36	51	66	144	245	557	903
2002.....	52,959	41	22	11	18	23	56	75	120	254	519	930
2001.....	53,428	35	18	17	17	28	53	68	152	281	553	876
2000.....	54,938	46	14	12	14	33	51	89	136	282	560	894
1999.....	54,867	43	16	16	11	28	44	73	140	300	556	802
White, female												
2004.....	78,572	34	10	12	17	23	47	55	120	227	488	773
2003.....	83,059	31	16	8	12	27	38	60	118	212	494	708
2002.....	86,760	30	18	14	20	15	36	49	132	255	519	741
2001.....	87,037	25	21	12	13	25	32	65	131	272	499	721
2000.....	89,642	29	16	9	20	18	46	71	124	276	527	733
1999.....	89,960	27	18	7	15	18	45	72	117	302	488	733
Black, both sexes												
2004.....	18,118	40	12	8	9	18	31	60	113	196	507	851
2003.....	18,806	32	15	6	10	21	38	54	114	250	543	828
2002.....	18,856	37	11	7	17	13	25	51	117	238	506	818
2001.....	19,002	43	12	6	8	13	38	59	99	236	529	802
2000.....	19,221	46	14	2	13	13	30	41	119	283	553	878
1999.....	18,884	28	10	3	12	18	23	50	100	268	548	787
Black, male												
2004.....	7,644	23	5	4	6	13	18	23	54	112	233	445
2003.....	7,866	20	10	5	4	13	13	33	54	130	257	419
2002.....	7,828	20	7	4	12	10	12	21	53	102	230	407
2001.....	7,907	27	5	2	3	8	12	35	44	121	240	418
2000.....	8,026	27	3	-	5	4	15	18	53	127	270	449
1999.....	7,894	18	4	1	6	7	12	24	48	124	269	412

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebrovascular diseases (I60-I69)												
Black, female												
2004.....	10,474	17	7	4	3	5	13	37	59	84	274	406
2003.....	10,940	12	5	1	6	8	25	21	60	120	286	409
2002.....	11,028	17	4	3	5	3	13	30	64	136	276	411
2001.....	11,095	16	7	4	5	5	26	24	55	115	289	384
2000.....	11,195	19	11	2	8	9	15	23	66	156	283	429
1999.....	10,990	10	6	2	6	11	11	26	52	144	279	375
American Indian 1/, both sexes												
2004.....	581	1	-	-	1	-	1	2	3	10	14	19
2003.....	552	-	1	1	-	1	-	2	3	13	21	26
2002.....	567	1	-	-	2	2	-	2	3	8	25	27
2001.....	574	2	-	1	-	-	-	1	4	11	19	27
2000.....	572	-	1	1	1	-	1	3	2	7	15	22
1999.....	546	2	-	-	1	-	1	-	11	9	21	21
American Indian 1/, male												
2004.....	246	-	-	-	1	-	-	1	1	1	5	9
2003.....	242	-	1	-	-	1	-	1	3	10	8	15
2002.....	236	1	-	-	-	1	-	1	3	3	11	14
2001.....	217	-	-	1	-	-	-	-	1	4	9	9
2000.....	250	-	1	-	-	-	-	-	-	4	7	10
1999.....	236	2	-	-	-	-	1	-	4	4	11	10
American Indian 1/, female												
2004.....	335	1	-	-	-	-	1	1	2	9	9	10
2003.....	310	-	-	1	-	-	-	1	-	3	13	11
2002.....	331	-	-	-	2	1	-	1	-	5	14	13
2001.....	357	2	-	-	-	-	-	1	3	7	10	18
2000.....	322	-	-	1	1	-	1	3	2	3	8	12
1999.....	310	-	-	-	1	-	-	-	7	5	10	11

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebrovascular diseases (I60-I69)												
Asian or Pacific Islander 2/, both sexes												
2004.....	3,507	7	1	-	1	1	8	6	10	21	56	96
2003.....	3,626	6	-	1	-	3	6	6	16	41	84	112
2002.....	3,530	8	2	1	1	-	1	6	12	42	59	99
2001.....	3,497	3	3	2	4	2	5	6	16	40	51	81
2000.....	3,288	3	2	1	3	3	4	5	12	32	64	98
1999.....	3,109	3	2	-	-	3	2	7	10	37	45	112
Asian or Pacific Islander 2/, male												
2004.....	1,614	3	-	-	-	1	2	3	4	11	36	57
2003.....	1,672	5	-	1	-	1	1	1	12	26	50	52
2002.....	1,599	5	1	1	1	-	1	3	8	30	29	51
2001.....	1,625	1	2	1	1	-	2	2	9	29	30	51
2000.....	1,555	1	1	-	2	1	2	4	8	17	30	48
1999.....	1,488	2	2	-	-	1	2	5	5	21	23	58
Asian or Pacific Islander 2/, female												
2004.....	1,893	4	1	-	1	-	6	3	6	10	20	39
2003.....	1,954	1	-	-	-	2	5	5	4	15	34	60
2002.....	1,931	3	1	-	-	-	-	3	4	12	30	48
2001.....	1,872	2	1	1	3	2	3	4	7	11	21	30
2000.....	1,733	2	1	1	1	2	2	1	4	15	34	50
1999.....	1,621	1	-	-	-	2	-	2	5	16	22	54

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Subarachnoid hemorrhage (I60)												
All races, both sexes												
2004.....	6,088	1	1	5	15	16	49	67	126	259	470	625
2003.....	6,033	3	-	7	8	18	41	74	132	262	474	573
2002.....	6,192	-	4	3	17	9	47	59	138	281	518	647
2001.....	6,142	2	3	5	8	21	46	67	146	283	507	617
2000.....	6,538	2	-	8	14	18	60	83	153	302	551	667
1999.....	6,489	4	4	3	8	24	39	80	153	360	582	657
All races, male												
2004.....	2,143	1	1	3	5	12	22	34	62	108	171	259
2003.....	2,135	3	-	3	4	9	24	35	68	119	185	220
2002.....	2,123	-	4	-	9	6	27	26	60	116	204	257
2001.....	2,151	-	2	3	2	9	25	39	71	111	211	242
2000.....	2,230	1	-	5	5	10	31	44	68	115	219	257
1999.....	2,239	2	2	2	2	14	17	44	74	152	241	265
All races, female												
2004.....	3,945	-	-	2	10	4	27	33	64	151	299	366
2003.....	3,898	-	-	4	4	9	17	39	64	143	289	353
2002.....	4,069	-	-	3	8	3	20	33	78	165	314	390
2001.....	3,991	2	1	2	6	12	21	28	75	172	296	375
2000.....	4,308	1	-	3	9	8	29	39	85	187	332	410
1999.....	4,250	2	2	1	6	10	22	36	79	208	341	392
White, both sexes												
2004.....	4,907	1	-	4	10	11	33	43	89	190	336	468
2003.....	4,813	2	-	4	6	14	30	50	102	169	333	419
2002.....	4,981	-	4	2	11	7	36	43	90	201	380	486
2001.....	4,897	-	3	3	5	15	32	45	107	196	363	460
2000.....	5,272	2	-	7	9	14	43	71	113	219	391	485
1999.....	5,264	4	4	3	4	18	34	60	115	263	421	479

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Subarachnoid hemorrhage (I60)												
White, male												
2004.....	1,738	1	-	2	3	9	16	26	46	77	131	195
2003.....	1,720	2	-	2	3	6	21	24	57	78	134	172
2002.....	1,739	-	4	-	6	5	22	22	38	87	159	212
2001.....	1,731	-	2	3	1	7	21	28	53	80	156	182
2000.....	1,845	1	-	5	3	9	23	40	57	90	164	198
1999.....	1,857	2	2	2	1	11	15	34	61	115	190	205
White, female												
2004.....	3,169	-	-	2	7	2	17	17	43	113	205	273
2003.....	3,093	-	-	2	3	8	9	26	45	91	199	247
2002.....	3,242	-	-	2	5	2	14	21	52	114	221	274
2001.....	3,166	-	1	-	4	8	11	17	54	116	207	278
2000.....	3,427	1	-	2	6	5	20	31	56	129	227	287
1999.....	3,407	2	2	1	3	7	19	26	54	148	231	274
Black, both sexes												
2004.....	866	-	1	1	4	5	11	19	31	54	114	125
2003.....	880	1	-	1	2	3	10	22	26	73	112	115
2002.....	895	-	-	1	5	2	10	15	39	65	118	136
2001.....	924	2	-	2	1	5	11	19	27	69	126	126
2000.....	971	-	-	-	4	3	15	11	36	73	132	144
1999.....	937	-	-	-	3	6	5	18	32	79	139	138
Black, male												
2004.....	300	-	1	1	2	3	4	6	13	27	34	47
2003.....	300	1	-	-	1	2	3	11	8	29	38	34
2002.....	284	-	-	-	3	1	4	4	16	20	37	37
2001.....	307	-	-	-	1	2	3	10	11	23	46	45
2000.....	292	-	-	-	2	-	8	3	9	22	48	46
1999.....	307	-	-	-	1	3	2	8	12	31	45	48

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Subarachnoid hemorrhage (I60)												
Black, female												
2004.....	566	-	-	-	2	2	7	13	18	27	80	78
2003.....	580	-	-	1	1	1	7	11	18	44	74	81
2002.....	611	-	-	1	2	1	6	11	23	45	81	99
2001.....	617	2	-	2	-	3	8	9	16	46	80	81
2000.....	679	-	-	-	2	3	7	8	27	51	84	98
1999.....	630	-	-	-	2	3	3	10	20	48	94	90
American Indian 1/, both sexes												
2004.....	56	-	-	-	-	-	1	1	2	7	5	8
2003.....	50	-	-	1	-	-	-	1	-	5	5	6
2002.....	38	-	-	-	1	-	-	-	2	3	4	3
2001.....	46	-	-	-	-	-	-	1	3	4	3	8
2000.....	41	-	-	1	1	-	1	-	1	2	5	5
1999.....	38	-	-	-	1	-	-	-	2	5	6	6
American Indian 1/, male												
2004.....	14	-	-	-	-	-	-	-	-	1	1	3
2003.....	14	-	-	-	-	-	-	-	-	4	1	2
2002.....	15	-	-	-	-	-	-	-	2	1	1	1
2001.....	16	-	-	-	-	-	-	-	1	1	1	3
2000.....	14	-	-	-	-	-	-	-	-	-	1	3
1999.....	7	-	-	-	-	-	-	-	-	2	2	1
American Indian 1/, female												
2004.....	42	-	-	-	-	-	1	1	2	6	4	5
2003.....	36	-	-	1	-	-	-	1	-	1	4	4
2002.....	23	-	-	-	1	-	-	-	-	2	3	2
2001.....	30	-	-	-	-	-	-	1	2	3	2	5
2000.....	27	-	-	1	1	-	1	-	1	2	4	2
1999.....	31	-	-	-	1	-	-	-	2	3	4	5

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Subarachnoid hemorrhage (I60)												
Asian or Pacific Islander 2/, both sexes												
2004.....	259	-	-	-	1	-	4	4	4	8	15	24
2003.....	290	-	-	1	-	1	1	1	4	15	24	33
2002.....	278	-	-	-	-	-	1	1	7	12	16	22
2001.....	275	-	-	-	2	1	3	2	9	14	15	23
2000.....	254	-	-	-	-	1	1	1	3	8	23	33
1999.....	250	-	-	-	-	-	-	2	4	13	16	34
Asian or Pacific Islander 2/, male												
2004.....	91	-	-	-	-	-	2	2	3	3	5	14
2003.....	101	-	-	1	-	1	-	-	3	8	12	12
2002.....	85	-	-	-	-	-	1	-	4	8	7	7
2001.....	97	-	-	-	-	-	1	1	6	7	8	12
2000.....	79	-	-	-	-	1	-	1	2	3	6	10
1999.....	68	-	-	-	-	-	-	2	1	4	4	11
Asian or Pacific Islander 2/, female												
2004.....	168	-	-	-	1	-	2	2	1	5	10	10
2003.....	189	-	-	-	-	-	1	1	1	7	12	21
2002.....	193	-	-	-	-	-	-	1	3	4	9	15
2001.....	178	-	-	-	2	1	2	1	3	7	7	11
2000.....	175	-	-	-	-	-	1	-	1	5	17	23
1999.....	182	-	-	-	-	-	-	-	3	9	12	23

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Intracerebral and other intracranial hemorrhage (I61-I62)

All races, both sexes

2004.....	27,555	54	12	9	9	26	60	84	123	253	665	1,126
2003.....	27,212	31	12	10	17	35	55	68	132	279	704	1,138
2002.....	26,976	42	9	13	18	25	40	64	133	286	642	1,056
2001.....	26,636	40	15	14	18	28	60	81	151	324	696	1,074
2000.....	27,402	41	6	9	15	25	49	63	141	342	715	1,122
1999.....	25,461	41	9	10	11	24	39	68	124	352	632	957

All races, male

2004.....	12,901	29	5	4	6	15	35	49	65	173	400	655
2003.....	12,662	17	7	7	9	21	27	41	68	168	391	655
2002.....	12,376	22	4	6	8	17	28	39	72	147	365	629
2001.....	12,266	25	7	8	9	20	35	47	81	200	385	646
2000.....	12,533	26	4	4	5	14	25	40	80	196	403	677
1999.....	11,564	25	5	7	3	15	23	33	77	199	378	537

All races, female

2004.....	14,654	25	7	5	3	11	25	35	58	80	265	471
2003.....	14,550	14	5	3	8	14	28	27	64	111	313	483
2002.....	14,600	20	5	7	10	8	12	25	61	139	277	427
2001.....	14,370	15	8	6	9	8	25	34	70	124	311	428
2000.....	14,869	15	2	5	10	11	24	23	61	146	312	445
1999.....	13,897	16	4	3	8	9	16	35	47	153	254	420

White, both sexes

2004.....	22,510	32	10	5	7	19	41	52	74	160	401	639
2003.....	22,030	18	7	9	13	24	36	40	80	161	397	638
2002.....	22,209	28	7	12	12	16	31	42	83	170	343	599
2001.....	21,822	22	10	12	11	23	40	49	100	206	407	610
2000.....	22,615	22	5	7	9	17	38	49	83	194	399	619
1999.....	21,037	25	5	8	10	15	27	47	73	199	344	534

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intracerebral and other intracranial hemorrhage (I61-I62)												
White, male												
2004.....	10,397	17	5	2	4	10	24	35	41	107	259	384
2003.....	10,106	9	4	6	8	15	21	23	39	95	240	390
2002.....	10,038	13	3	5	3	11	22	29	43	88	203	365
2001.....	9,910	13	4	8	6	16	25	28	58	123	234	383
2000.....	10,131	13	3	4	3	11	20	30	47	114	230	389
1999.....	9,380	15	2	7	3	12	15	21	50	114	209	302
White, female												
2004.....	12,113	15	5	3	3	9	17	17	33	53	142	255
2003.....	11,924	9	3	3	5	9	15	17	41	66	157	248
2002.....	12,171	15	4	7	9	5	9	13	40	82	140	234
2001.....	11,912	9	6	4	5	7	15	21	42	83	173	227
2000.....	12,484	9	2	3	6	6	18	19	36	80	169	230
1999.....	11,657	10	3	1	7	3	12	26	23	85	135	232
Black, both sexes												
2004.....	3,939	17	1	4	2	6	16	29	44	84	232	430
2003.....	4,020	12	5	1	4	9	16	23	42	91	259	432
2002.....	3,674	13	1	1	6	8	9	17	47	91	257	400
2001.....	3,733	17	5	1	5	4	18	30	44	96	249	409
2000.....	3,777	17	-	1	4	7	8	11	54	131	280	449
1999.....	3,443	16	3	2	1	9	10	18	40	131	259	368
Black, male												
2004.....	1,966	10	-	2	2	4	11	12	23	60	119	239
2003.....	2,002	7	3	1	1	5	5	16	21	54	125	230
2002.....	1,818	8	1	1	5	6	6	8	26	42	142	229
2001.....	1,825	12	3	-	2	4	9	19	20	58	126	232
2000.....	1,898	12	-	-	1	3	3	7	29	72	153	257
1999.....	1,689	10	2	-	-	3	6	11	20	69	150	204

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Intracerebral and other intracranial hemorrhage (I61-I62)

Black, female

2004.....	1,973	7	1	2	-	2	5	17	21	24	113	191
2003.....	2,018	5	2	-	3	4	11	7	21	37	134	202
2002.....	1,856	5	-	-	1	2	3	9	21	49	115	171
2001.....	1,908	5	2	1	3	-	9	11	24	38	123	177
2000.....	1,879	5	-	1	3	4	5	4	25	59	127	192
1999.....	1,754	6	1	2	1	6	4	7	20	62	109	164

American Indian 1/, both sexes

2004.....	119	1	-	-	-	-	-	1	-	1	7	8
2003.....	133	-	-	-	-	1	-	1	2	5	13	9
2002.....	122	-	-	-	-	1	-	1	1	4	9	10
2001.....	131	1	-	-	-	-	-	-	1	5	13	14
2000.....	116	-	1	-	-	-	-	-	-	1	6	10
1999.....	114	-	-	-	-	-	1	-	5	2	11	9

American Indian 1/, male

2004.....	58	-	-	-	-	-	-	1	-	-	3	5
2003.....	61	-	-	-	-	1	-	1	2	4	5	5
2002.....	56	-	-	-	-	-	-	-	1	2	4	8
2001.....	54	-	-	-	-	-	-	-	-	3	7	3
2000.....	52	-	1	-	-	-	-	-	-	1	3	4
1999.....	50	-	-	-	-	-	1	-	3	2	7	5

American Indian 1/, female

2004.....	61	1	-	-	-	-	-	-	-	1	4	3
2003.....	72	-	-	-	-	-	-	-	-	1	8	4
2002.....	66	-	-	-	-	1	-	1	-	2	5	2
2001.....	77	1	-	-	-	-	-	-	1	2	6	11
2000.....	64	-	-	-	-	-	-	-	-	-	3	6
1999.....	64	-	-	-	-	-	-	-	2	-	4	4

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Intracerebral and other intracranial hemorrhage (I61-I62)

Asian or Pacific Islander 2/, both sexes

2004.....	987	4	1	-	-	1	3	2	5	8	25	49
2003.....	1,029	1	-	-	-	1	3	4	8	22	35	59
2002.....	971	1	1	-	-	-	-	4	2	21	33	47
2001.....	950	-	-	1	2	1	2	2	6	17	27	41
2000.....	894	2	-	1	2	1	3	3	4	16	30	44
1999.....	867	-	1	-	-	-	1	3	6	20	18	46

Asian or Pacific Islander 2/, male

2004.....	480	2	-	-	-	1	-	1	1	6	19	27
2003.....	493	1	-	-	-	-	1	1	6	15	21	30
2002.....	464	1	-	-	-	-	-	2	2	15	16	27
2001.....	477	-	-	-	1	-	1	-	3	16	18	28
2000.....	452	1	-	-	1	-	2	3	4	9	17	27
1999.....	445	-	1	-	-	-	1	1	4	14	12	26

Asian or Pacific Islander 2/, female

2004.....	507	2	1	-	-	-	3	1	4	2	6	22
2003.....	536	-	-	-	-	1	2	3	2	7	14	29
2002.....	507	-	1	-	-	-	-	2	-	6	17	20
2001.....	473	-	-	1	1	1	1	2	3	1	9	13
2000.....	442	1	-	1	1	1	1	-	-	7	13	17
1999.....	422	-	-	-	-	-	-	2	2	6	6	20

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Cerebral infarction (I63)

All races, both sexes

2004.....	10,995	10	3	2	1	5	10	18	23	41	94	142
2003.....	12,094	14	8	3	1	8	5	7	28	45	92	164
2002.....	12,335	15	2	3	2	5	10	18	27	34	77	150
2001.....	12,654	16	2	5	5	2	6	12	27	51	85	163
2000.....	13,747	17	5	3	8	5	7	12	26	52	101	139
1999.....	14,804	13	9	6	3	2	9	12	23	52	100	140

All races, male

2004.....	4,398	7	3	-	1	3	4	9	11	19	51	76
2003.....	4,753	8	3	1	-	5	2	2	16	25	58	112
2002.....	4,841	7	-	1	1	2	3	11	14	17	38	93
2001.....	4,943	8	-	3	4	1	1	4	21	29	45	93
2000.....	5,386	11	-	-	2	1	4	4	8	28	55	86
1999.....	5,777	9	4	3	3	1	4	8	12	30	53	87

All races, female

2004.....	6,597	3	-	2	-	2	6	9	12	22	43	66
2003.....	7,341	6	5	2	1	3	3	5	12	20	34	52
2002.....	7,494	8	2	2	1	3	7	7	13	17	39	57
2001.....	7,711	8	2	2	1	1	5	8	6	22	40	70
2000.....	8,361	6	5	3	6	4	3	8	18	24	46	53
1999.....	9,027	4	5	3	-	1	5	4	11	22	47	53

White, both sexes

2004.....	9,455	8	2	2	-	4	8	16	16	30	63	99
2003.....	10,504	9	6	3	1	5	3	5	22	25	62	112
2002.....	10,747	13	2	2	2	4	7	9	18	22	59	112
2001.....	11,057	10	2	4	5	2	4	11	21	31	60	109
2000.....	11,962	12	3	2	5	5	4	8	12	37	60	101
1999.....	12,962	8	6	6	2	2	5	12	16	42	68	95

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Cerebral infarction (I63)

White, male

2004.....	3,717	6	2	-	-	2	3	7	7	17	34	50
2003.....	4,082	4	1	1	-	3	1	1	16	13	42	81
2002.....	4,166	6	-	1	1	1	3	7	10	11	30	75
2001.....	4,255	4	-	2	4	1	1	4	17	17	37	62
2000.....	4,604	7	-	-	2	1	2	4	2	20	36	64
1999.....	4,967	4	2	3	2	1	2	8	7	26	38	54

White, female

2004.....	5,738	2	-	2	-	2	5	9	9	13	29	49
2003.....	6,422	5	5	2	1	2	2	4	6	12	20	31
2002.....	6,581	7	2	1	1	3	4	2	8	11	29	37
2001.....	6,802	6	2	2	1	1	3	7	4	14	23	47
2000.....	7,358	5	3	2	3	4	2	4	10	17	24	37
1999.....	7,995	4	4	3	-	1	3	4	9	16	30	41

Black, both sexes

2004.....	1,282	2	1	-	1	1	2	2	7	9	30	41
2003.....	1,331	3	2	-	-	2	2	1	5	20	25	42
2002.....	1,317	1	-	1	-	-	3	7	9	11	17	34
2001.....	1,360	4	-	-	-	-	2	-	6	16	25	50
2000.....	1,506	5	2	1	3	-	3	2	12	12	35	31
1999.....	1,588	3	2	-	1	-	3	-	6	9	29	41

Black, male

2004.....	565	1	1	-	1	1	1	2	4	2	16	25
2003.....	560	2	2	-	-	2	1	1	-	12	12	26
2002.....	552	1	-	-	-	-	-	2	4	6	7	17
2001.....	585	3	-	-	-	-	-	-	4	9	8	28
2000.....	641	4	-	-	-	-	2	-	6	6	15	18
1999.....	689	3	1	-	1	-	1	-	4	3	12	29

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Stroke, not specified as hemorrhage or infarction (I64)												
All races, both sexes												
2004.....	76,481	4	1	5	5	7	12	22	59	118	299	566
2003.....	82,472	5	4	1	1	14	15	24	77	124	336	533
2002.....	86,879	3	6	4	3	3	12	25	56	133	284	595
2001.....	88,670	1	4	1	3	6	10	21	44	118	278	510
2000.....	90,379	2	6	1	4	7	7	33	51	123	262	530
1999.....	91,051	5	2	-	3	6	13	24	51	97	249	535
All races, male												
2004.....	28,433	3	-	3	4	5	8	11	22	65	167	323
2003.....	30,749	5	2	-	1	8	6	14	46	72	199	309
2002.....	32,029	3	2	1	1	2	6	14	19	78	142	339
2001.....	32,798	-	1	1	3	3	5	7	16	60	151	304
2000.....	33,425	-	3	-	2	4	3	16	32	64	141	303
1999.....	33,756	2	1	-	1	3	5	11	23	42	146	312
All races, female												
2004.....	48,048	1	1	2	1	2	4	11	37	53	132	243
2003.....	51,723	-	2	1	-	6	9	10	31	52	137	224
2002.....	54,850	-	4	3	2	1	6	11	37	55	142	256
2001.....	55,872	1	3	-	-	3	5	14	28	58	127	206
2000.....	56,954	2	3	1	2	3	4	17	19	59	121	227
1999.....	57,295	3	1	-	2	3	8	13	28	55	103	223
White, both sexes												
2004.....	65,877	4	1	4	4	5	11	15	32	72	181	349
2003.....	71,331	4	2	1	1	9	8	16	39	69	194	332
2002.....	75,302	3	6	3	1	2	9	18	41	72	180	363
2001.....	76,977	1	4	1	3	3	7	14	29	72	167	320
2000.....	78,787	2	3	1	4	4	4	19	35	68	177	310
1999.....	79,569	5	2	-	2	4	9	16	33	61	145	317

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Stroke, not specified as hemorrhage or infarction (I64)

White, male

2004.....	24,123	3	-	3	3	3	7	8	10	45	103	206
2003.....	26,179	4	1	-	1	5	3	9	21	41	115	199
2002.....	27,270	3	2	1	-	1	4	11	15	42	100	218
2001.....	27,968	-	1	1	3	2	5	2	11	36	99	201
2000.....	28,632	-	2	-	2	3	2	8	23	40	99	184
1999.....	29,042	2	1	-	-	2	3	7	12	28	93	190

White, female

2004.....	41,754	1	1	1	1	2	4	7	22	27	78	143
2003.....	45,152	-	1	1	-	4	5	7	18	28	79	133
2002.....	48,032	-	4	2	1	1	5	7	26	30	80	145
2001.....	49,009	1	3	-	-	1	2	12	18	36	68	119
2000.....	50,155	2	1	1	2	1	2	11	12	28	78	126
1999.....	50,527	3	1	-	2	2	6	9	21	33	52	127

Black, both sexes

2004.....	8,898	-	-	1	-	2	1	7	26	42	109	200
2003.....	9,390	1	1	-	-	5	5	8	34	51	126	187
2002.....	9,809	-	-	1	1	1	3	7	14	55	92	203
2001.....	9,891	-	-	-	-	3	3	6	15	41	104	174
2000.....	9,970	-	3	-	-	3	3	14	12	50	81	203
1999.....	9,931	-	-	-	1	-	4	7	16	32	95	195

Black, male

2004.....	3,521	-	-	-	-	2	1	3	11	18	57	107
2003.....	3,742	1	-	-	-	3	3	5	21	29	72	99
2002.....	3,958	-	-	-	1	1	2	3	4	30	34	105
2001.....	4,016	-	-	-	-	1	-	4	5	22	48	94
2000.....	4,041	-	1	-	-	1	1	8	7	21	39	110
1999.....	3,995	-	-	-	1	-	2	3	11	13	49	107

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Stroke, not specified as hemorrhage or infarction (I64)												
Black, female												
2004.....	5,377	-	-	1	-	-	-	4	15	24	52	93
2003.....	5,648	-	1	-	-	2	2	3	13	22	54	88
2002.....	5,851	-	-	1	-	-	1	4	10	25	58	98
2001.....	5,875	-	-	-	-	2	3	2	10	19	56	80
2000.....	5,929	-	2	-	-	2	2	6	5	29	42	93
1999.....	5,936	-	-	-	-	-	2	4	5	19	46	88
American Indian 1/, both sexes												
2004.....	285	-	-	-	1	-	-	-	1	1	2	3
2003.....	250	-	1	-	-	-	-	-	1	2	2	5
2002.....	274	-	-	-	1	-	-	-	-	-	7	11
2001.....	285	-	-	-	-	-	-	-	-	2	3	5
2000.....	272	-	-	-	-	-	-	-	-	1	1	3
1999.....	273	-	-	-	-	-	-	-	2	2	2	4
American Indian 1/, male												
2004.....	128	-	-	-	1	-	-	-	1	-	1	1
2003.....	115	-	1	-	-	-	-	-	1	1	1	4
2002.....	113	-	-	-	-	-	-	-	-	-	4	4
2001.....	105	-	-	-	-	-	-	-	-	-	1	3
2000.....	119	-	-	-	-	-	-	-	-	1	1	2
1999.....	123	-	-	-	-	-	-	-	-	-	1	4
American Indian 1/, female												
2004.....	157	-	-	-	-	-	-	-	-	1	1	2
2003.....	135	-	-	-	-	-	-	-	-	1	1	1
2002.....	161	-	-	-	1	-	-	-	-	-	3	7
2001.....	180	-	-	-	-	-	-	-	-	2	2	2
2000.....	153	-	-	-	-	-	-	-	-	-	-	1
1999.....	150	-	-	-	-	-	-	-	2	2	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Stroke, not specified as hemorrhage or infarction (I64)												
Asian or Pacific Islander 2/, both sexes												
2004.....	1,421	-	-	-	-	-	-	-	-	3	7	14
2003.....	1,501	-	-	-	-	-	2	-	3	2	14	9
2002.....	1,494	-	-	-	-	-	-	-	1	6	5	18
2001.....	1,517	-	-	-	-	-	-	1	-	3	4	11
2000.....	1,350	-	-	-	-	-	-	-	4	4	3	14
1999.....	1,278	-	-	-	-	2	-	1	-	2	7	19
Asian or Pacific Islander 2/, male												
2004.....	661	-	-	-	-	-	-	-	-	2	6	9
2003.....	713	-	-	-	-	-	-	-	3	1	11	7
2002.....	688	-	-	-	-	-	-	-	-	6	4	12
2001.....	709	-	-	-	-	-	-	1	-	2	3	6
2000.....	633	-	-	-	-	-	-	-	2	2	2	7
1999.....	596	-	-	-	-	1	-	1	-	1	3	11
Asian or Pacific Islander 2/, female												
2004.....	760	-	-	-	-	-	-	-	-	1	1	5
2003.....	788	-	-	-	-	-	2	-	-	1	3	2
2002.....	806	-	-	-	-	-	-	-	1	-	1	6
2001.....	808	-	-	-	-	-	-	-	-	1	1	5
2000.....	717	-	-	-	-	-	-	-	2	2	1	7
1999.....	682	-	-	-	-	1	-	-	-	1	4	8

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other cerebrovascular diseases and their sequelae (I67,I69)												
All races, both sexes												
2004.....	28,955	58	24	13	13	15	11	17	28	48	114	163
2003.....	29,878	48	22	8	13	13	17	15	26	51	93	169
2002.....	30,290	57	32	10	18	11	9	17	30	63	107	167
2001.....	29,436	49	30	13	8	11	6	18	34	64	85	143
2000.....	29,595	62	30	4	10	12	9	18	22	61	90	167
1999.....	29,561	40	22	7	14	11	15	18	27	55	95	166
All races, male												
2004.....	10,925	31	14	8	6	6	6	9	12	24	62	81
2003.....	11,127	24	13	8	8	8	6	9	15	27	39	93
2002.....	11,253	35	20	8	12	7	5	10	19	31	40	84
2001.....	11,019	30	15	6	3	3	1	8	17	35	40	69
2000.....	11,195	36	12	3	7	9	5	7	9	27	49	78
1999.....	11,149	27	10	5	8	3	10	6	11	26	41	81
All races, female												
2004.....	18,030	27	10	5	7	9	5	8	16	24	52	82
2003.....	18,751	24	9	-	5	5	11	6	11	24	54	76
2002.....	19,037	22	12	2	6	4	4	7	11	32	67	83
2001.....	18,417	19	15	7	5	8	5	10	17	29	45	74
2000.....	18,400	26	18	1	3	3	4	11	13	34	41	89
1999.....	18,412	13	12	2	6	8	5	12	16	29	54	85
White, both sexes												
2004.....	25,119	34	15	11	11	11	9	14	22	40	84	101
2003.....	26,027	30	15	4	9	11	12	15	19	33	65	110
2002.....	26,480	27	21	6	12	9	9	12	20	44	76	111
2001.....	25,712	27	20	9	6	10	2	14	26	48	55	98
2000.....	25,944	37	19	4	7	11	8	13	17	40	60	112
1999.....	25,995	28	17	6	8	7	14	10	20	37	66	110

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other cerebrovascular diseases and their sequelae (I67,I69)												
White, male												
2004.....	9,321	18	11	7	5	3	5	9	9	19	50	48
2003.....	9,559	13	8	4	6	7	5	9	11	18	26	61
2002.....	9,746	19	13	4	8	5	5	6	14	26	27	60
2001.....	9,564	18	11	3	3	2	1	6	13	25	27	48
2000.....	9,726	25	9	3	4	9	4	7	7	18	31	59
1999.....	9,621	20	9	4	5	2	9	3	10	17	26	51
White, female												
2004.....	15,798	16	4	4	6	8	4	5	13	21	34	53
2003.....	16,468	17	7	-	3	4	7	6	8	15	39	49
2002.....	16,734	8	8	2	4	4	4	6	6	18	49	51
2001.....	16,148	9	9	6	3	8	1	8	13	23	28	50
2000.....	16,218	12	10	1	3	2	4	6	10	22	29	53
1999.....	16,374	8	8	2	3	5	5	7	10	20	40	59
Black, both sexes												
2004.....	3,133	21	9	2	2	4	1	3	5	7	22	55
2003.....	3,185	15	7	4	4	2	5	-	7	15	21	52
2002.....	3,161	23	10	3	5	2	-	5	8	16	22	45
2001.....	3,094	20	7	3	2	1	4	4	7	14	25	43
2000.....	2,997	24	9	-	2	-	1	3	5	17	25	51
1999.....	2,985	9	5	1	6	3	1	7	6	17	26	45
Black, male												
2004.....	1,292	12	3	1	1	3	1	-	3	5	7	27
2003.....	1,262	9	5	4	2	1	1	-	4	6	10	30
2002.....	1,216	11	6	3	3	2	-	4	3	4	10	19
2001.....	1,174	12	2	2	-	1	-	2	4	9	12	19
2000.....	1,154	11	2	-	2	-	1	-	2	6	15	18
1999.....	1,214	5	1	1	3	1	1	2	1	8	13	24

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other cerebrovascular diseases and their sequelae (I67,I69)

Black, female

2004.....	1,841	9	6	1	1	1	-	3	2	2	15	28
2003.....	1,923	6	2	-	2	1	4	-	3	9	11	22
2002.....	1,945	12	4	-	2	-	-	1	5	12	12	26
2001.....	1,920	8	5	1	2	-	4	2	3	5	13	24
2000.....	1,843	13	7	-	-	-	-	3	3	11	10	33
1999.....	1,771	4	4	-	3	2	-	5	5	9	13	21

American Indian 1/, both sexes

2004.....	86	-	-	-	-	-	-	-	-	-	-	-
2003.....	82	-	-	-	-	-	-	-	-	1	-	2
2002.....	80	1	-	-	-	-	-	-	-	1	4	2
2001.....	81	-	-	1	-	-	-	-	-	-	-	-
2000.....	93	-	-	-	-	-	-	2	-	1	1	1
1999.....	84	2	-	-	-	-	-	-	1	-	2	2

American Indian 1/, male

2004.....	29	-	-	-	-	-	-	-	-	-	-	-
2003.....	36	-	-	-	-	-	-	-	-	1	-	1
2002.....	27	1	-	-	-	-	-	-	-	-	1	1
2001.....	31	-	-	1	-	-	-	-	-	-	-	-
2000.....	44	-	-	-	-	-	-	-	-	1	-	-
1999.....	41	2	-	-	-	-	-	-	-	-	1	-

American Indian 1/, female

2004.....	57	-	-	-	-	-	-	-	-	-	-	-
2003.....	46	-	-	-	-	-	-	-	-	-	-	1
2002.....	53	-	-	-	-	-	-	-	-	1	3	1
2001.....	50	-	-	-	-	-	-	-	-	-	-	-
2000.....	49	-	-	-	-	-	-	2	-	-	1	1
1999.....	43	-	-	-	-	-	-	-	1	-	1	2

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other cerebrovascular diseases and their sequelae (I67,I69)												
Asian or Pacific Islander 2/, both sexes												
2004.....	617	3	-	-	-	-	1	-	1	1	8	7
2003.....	584	3	-	-	-	-	-	-	-	2	7	5
2002.....	569	6	1	1	1	-	-	-	2	2	5	9
2001.....	549	2	3	-	-	-	-	-	1	2	5	2
2000.....	561	1	2	-	1	1	-	-	-	3	4	3
1999.....	497	1	-	-	-	1	-	1	-	1	1	9
Asian or Pacific Islander 2/, male												
2004.....	283	1	-	-	-	-	-	-	-	-	5	6
2003.....	270	2	-	-	-	-	-	-	-	2	3	1
2002.....	264	4	1	1	1	-	-	-	2	1	2	4
2001.....	250	-	2	-	-	-	-	-	-	1	1	2
2000.....	271	-	1	-	1	-	-	-	-	2	3	1
1999.....	273	-	-	-	-	-	-	1	-	1	1	6
Asian or Pacific Islander 2/, female												
2004.....	334	2	-	-	-	-	1	-	1	1	3	1
2003.....	314	1	-	-	-	-	-	-	-	-	4	4
2002.....	305	2	-	-	-	-	-	-	-	1	3	5
2001.....	299	2	1	-	-	-	-	-	1	1	4	-
2000.....	290	1	1	-	-	1	-	-	-	1	1	2
1999.....	224	1	-	-	-	1	-	-	-	-	-	3

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of arteries, arterioles and capillaries (I70-I78)												
All races, both sexes												
2004.....	35,554	14	5	3	10	14	49	65	108	186	318	523
2003.....	37,725	14	-	1	11	31	40	62	96	178	320	594
2002.....	38,748	18	4	1	10	31	38	53	116	170	325	540
2001.....	39,404	13	5	3	9	23	39	68	105	208	330	486
2000.....	40,429	18	5	3	8	17	45	52	110	173	323	495
1999.....	40,788	14	2	2	9	15	36	52	97	194	329	474
All races, male												
2004.....	16,816	7	2	2	6	13	34	47	69	122	211	366
2003.....	18,076	5	-	-	7	18	29	43	65	125	216	397
2002.....	18,480	7	4	-	5	19	26	32	74	90	231	375
2001.....	18,539	9	3	1	3	17	24	43	74	140	221	334
2000.....	19,441	6	2	1	6	9	26	37	72	102	209	323
1999.....	19,237	10	-	1	6	10	22	36	71	128	200	326
All races, female												
2004.....	18,738	7	3	1	4	1	15	18	39	64	107	157
2003.....	19,649	9	-	1	4	13	11	19	31	53	104	197
2002.....	20,268	11	-	1	5	12	12	21	42	80	94	165
2001.....	20,865	4	2	2	6	6	15	25	31	68	109	152
2000.....	20,988	12	3	2	2	8	19	15	38	71	114	172
1999.....	21,551	4	2	1	3	5	14	16	26	66	129	148
White, both sexes												
2004.....	31,511	11	4	2	9	14	37	47	70	123	226	387
2003.....	33,546	7	-	1	7	23	28	44	68	111	234	422
2002.....	34,566	14	4	1	3	26	26	37	77	129	230	374
2001.....	35,197	11	4	3	8	18	29	52	79	148	225	363
2000.....	36,145	15	4	2	5	14	33	36	80	130	214	353
1999.....	36,689	10	2	2	6	14	30	39	73	152	237	352

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of arteries, arterioles and capillaries (I70-I78)												
White, male												
2004.....	14,956	6	2	1	5	13	24	34	45	82	154	276
2003.....	16,076	2	-	-	4	15	20	30	44	84	161	293
2002.....	16,506	6	4	-	2	16	17	24	48	70	168	272
2001.....	16,583	7	3	1	3	14	17	36	56	98	151	251
2000.....	17,486	4	2	1	5	7	21	25	57	76	142	244
1999.....	17,376	7	-	1	4	9	18	27	54	102	148	251
White, female												
2004.....	16,555	5	2	1	4	1	13	13	25	41	72	111
2003.....	17,470	5	-	1	3	8	8	14	24	27	73	129
2002.....	18,060	8	-	1	1	10	9	13	29	59	62	102
2001.....	18,614	4	1	2	5	4	12	16	23	50	74	112
2000.....	18,659	11	2	1	-	7	12	11	23	54	72	109
1999.....	19,313	3	2	1	2	5	12	12	19	50	89	101
Black, both sexes												
2004.....	3,387	2	1	1	1	-	7	16	31	55	85	126
2003.....	3,514	7	-	-	3	7	8	16	23	59	79	151
2002.....	3,542	4	-	-	6	3	9	15	35	34	89	150
2001.....	3,602	2	1	-	1	5	8	15	25	52	96	109
2000.....	3,649	3	-	1	3	2	11	13	26	38	98	126
1999.....	3,503	4	-	-	-	1	6	12	20	37	84	110
Black, male												
2004.....	1,530	1	-	1	1	-	7	12	19	33	52	83
2003.....	1,626	3	-	-	2	3	5	12	18	35	51	94
2002.....	1,625	1	-	-	2	1	6	8	24	17	60	88
2001.....	1,599	2	-	-	-	3	5	7	17	35	63	71
2000.....	1,622	2	-	-	1	1	5	9	14	23	61	71
1999.....	1,543	3	-	-	-	1	4	8	16	22	45	69

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of arteries, arterioles and capillaries (I70-I78)												
Black, female												
2004.....	1,857	1	1	-	-	-	-	4	12	22	33	43
2003.....	1,888	4	-	-	1	4	3	4	5	24	28	57
2002.....	1,917	3	-	-	4	2	3	7	11	17	29	62
2001.....	2,003	-	1	-	1	2	3	8	8	17	33	38
2000.....	2,027	1	-	1	2	1	6	4	12	15	37	55
1999.....	1,960	1	-	-	-	-	2	4	4	15	39	41
American Indian 1/, both sexes												
2004.....	122	-	-	-	-	-	-	-	3	4	-	3
2003.....	136	-	-	-	-	-	2	1	2	2	2	5
2002.....	149	-	-	-	-	-	1	-	1	2	1	4
2001.....	123	-	-	-	-	-	-	-	-	1	1	3
2000.....	136	-	-	-	-	-	-	-	-	1	2	3
1999.....	116	-	-	-	2	-	-	-	-	-	-	1
American Indian 1/, male												
2004.....	55	-	-	-	-	-	-	-	2	3	-	2
2003.....	72	-	-	-	-	-	2	-	2	1	1	2
2002.....	74	-	-	-	-	-	1	-	-	2	-	4
2001.....	63	-	-	-	-	-	-	-	-	1	1	2
2000.....	55	-	-	-	-	-	-	-	-	-	-	2
1999.....	50	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, female												
2004.....	67	-	-	-	-	-	-	-	1	1	-	1
2003.....	64	-	-	-	-	-	-	1	-	1	1	3
2002.....	75	-	-	-	-	-	-	-	1	-	1	-
2001.....	60	-	-	-	-	-	-	-	-	-	-	1
2000.....	81	-	-	-	-	-	-	-	-	1	2	1
1999.....	66	-	-	-	1	-	-	-	-	-	-	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of arteries, arterioles and capillaries (I70-I78)												
Asian or Pacific Islander 2/, both sexes												
2004.....	534	1	-	-	-	-	5	2	4	4	7	7
2003.....	529	-	-	-	1	1	2	1	3	6	5	16
2002.....	491	-	-	-	1	2	2	1	3	5	5	12
2001.....	482	-	-	-	-	-	2	1	1	7	8	11
2000.....	499	-	1	-	-	1	1	3	4	4	9	13
1999.....	480	-	-	-	1	-	-	1	4	5	8	11
Asian or Pacific Islander 2/, male												
2004.....	275	-	-	-	-	-	3	1	3	4	5	5
2003.....	302	-	-	-	1	-	2	1	1	5	3	8
2002.....	275	-	-	-	1	2	2	-	2	1	3	11
2001.....	294	-	-	-	-	-	2	-	1	6	6	10
2000.....	278	-	-	-	-	1	-	3	1	3	6	6
1999.....	268	-	-	-	1	-	-	1	1	4	7	6
Asian or Pacific Islander 2/, female												
2004.....	259	1	-	-	-	-	2	1	1	-	2	2
2003.....	227	-	-	-	-	1	-	-	2	1	2	8
2002.....	216	-	-	-	-	-	-	1	1	4	2	1
2001.....	188	-	-	-	-	-	-	1	-	1	2	1
2000.....	221	-	1	-	-	-	1	-	3	1	3	7
1999.....	212	-	-	-	-	-	-	-	3	1	1	5

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerosis (I70)												
All races, both sexes												
2004.....	11,861	-	1	-	-	-	2	-	6	12	31	60
2003.....	13,053	3	-	-	-	1	2	-	7	11	36	70
2002.....	13,821	4	-	-	-	-	3	2	7	16	34	61
2001.....	14,086	1	1	1	-	-	1	1	2	14	26	65
2000.....	14,393	2	-	-	-	-	3	3	5	16	32	71
1999.....	14,979	2	-	-	-	1	1	1	7	16	29	56
All races, male												
2004.....	4,510	-	-	-	-	-	2	-	3	7	18	42
2003.....	4,859	2	-	-	-	1	1	-	5	8	27	54
2002.....	5,171	1	-	-	-	-	2	2	3	9	21	44
2001.....	5,143	1	1	1	-	-	1	-	2	8	18	43
2000.....	5,357	-	-	-	-	-	2	1	5	10	25	48
1999.....	5,443	1	-	-	-	-	-	1	4	11	21	37
All races, female												
2004.....	7,351	-	1	-	-	-	-	-	3	5	13	18
2003.....	8,194	1	-	-	-	-	1	-	2	3	9	16
2002.....	8,650	3	-	-	-	-	1	-	4	7	13	17
2001.....	8,943	-	-	-	-	-	-	1	-	6	8	22
2000.....	9,036	2	-	-	-	-	1	2	-	6	7	23
1999.....	9,536	1	-	-	-	1	1	-	3	5	8	19
White, both sexes												
2004.....	10,877	-	1	-	-	-	2	-	5	7	23	52
2003.....	11,885	1	-	-	-	-	2	-	5	7	32	58
2002.....	12,631	3	-	-	-	-	3	2	4	14	26	44
2001.....	12,961	1	1	1	-	-	-	-	2	11	22	54
2000.....	13,182	2	-	-	-	-	3	1	4	12	24	50
1999.....	13,737	1	-	-	-	1	1	1	5	14	19	48

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerosis (I70)												
White, male												
2004.....	4,146	-	-	-	-	-	2	-	3	4	15	38
2003.....	4,368	-	-	-	-	-	1	-	4	5	23	43
2002.....	4,688	1	-	-	-	-	2	2	2	7	17	34
2001.....	4,696	1	1	1	-	-	-	-	2	5	16	35
2000.....	4,875	-	-	-	-	-	2	1	4	9	19	36
1999.....	4,932	1	-	-	-	-	-	1	3	10	14	32
White, female												
2004.....	6,731	-	1	-	-	-	-	-	2	3	8	14
2003.....	7,517	1	-	-	-	-	1	-	1	2	9	15
2002.....	7,943	2	-	-	-	-	1	-	2	7	9	10
2001.....	8,265	-	-	-	-	-	-	-	-	6	6	19
2000.....	8,307	2	-	-	-	-	1	-	-	3	5	14
1999.....	8,805	-	-	-	-	1	1	-	2	4	5	16
Black, both sexes												
2004.....	837	-	-	-	-	-	-	-	1	4	8	8
2003.....	995	2	-	-	-	1	-	-	2	2	3	11
2002.....	1,017	1	-	-	-	-	-	-	3	2	7	15
2001.....	970	-	-	-	-	-	1	1	-	2	4	11
2000.....	1,054	-	-	-	-	-	-	2	1	4	8	19
1999.....	1,069	1	-	-	-	-	-	-	2	2	7	6
Black, male												
2004.....	305	-	-	-	-	-	-	-	-	2	3	4
2003.....	410	2	-	-	-	1	-	-	1	1	3	10
2002.....	414	-	-	-	-	-	-	-	1	2	4	8
2001.....	370	-	-	-	-	-	1	-	-	2	2	8
2000.....	415	-	-	-	-	-	-	-	1	1	6	11
1999.....	432	-	-	-	-	-	-	-	1	1	5	4

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Aortic aneurysm and dissection (I71)

All races, both sexes

2004.....	13,753	1	1	1	4	9	36	49	77	131	187	306
2003.....	14,810	1	-	-	3	16	25	43	69	124	192	368
2002.....	14,818	2	-	1	4	20	19	30	80	104	201	327
2001.....	15,234	-	2	-	4	15	28	46	71	134	198	287
2000.....	15,810	-	-	-	5	10	26	38	69	103	200	268
1999.....	15,807	1	1	-	3	8	23	33	60	134	195	283

All races, male

2004.....	8,140	1	-	-	3	8	28	39	54	100	146	233
2003.....	8,976	-	-	-	2	10	21	32	51	96	150	269
2002.....	9,013	-	-	-	3	17	15	25	57	67	162	249
2001.....	9,114	-	2	-	3	12	19	34	56	105	159	219
2000.....	9,661	-	-	-	3	7	16	30	48	72	147	191
1999.....	9,615	1	-	-	2	6	14	25	53	98	134	216

All races, female

2004.....	5,613	-	1	1	1	1	8	10	23	31	41	73
2003.....	5,834	1	-	-	1	6	4	11	18	28	42	99
2002.....	5,805	2	-	1	1	3	4	5	23	37	39	78
2001.....	6,120	-	-	-	1	3	9	12	15	29	39	68
2000.....	6,149	-	-	-	2	3	10	8	21	31	53	77
1999.....	6,192	-	1	-	1	2	9	8	7	36	61	67

White, both sexes

2004.....	12,189	1	-	1	4	9	26	36	49	85	129	213
2003.....	13,246	-	-	-	1	14	19	28	49	77	137	252
2002.....	13,285	2	-	1	1	18	13	21	52	82	137	220
2001.....	13,652	-	2	-	4	11	21	37	55	96	131	208
2000.....	14,210	-	-	-	2	9	17	27	50	75	129	193
1999.....	14,336	1	1	-	2	8	19	25	45	105	135	194

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	-------------	-----------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Aortic aneurysm and dissection (I71)

White, male

2004.....	7,226	1	-	-	3	8	20	29	33	67	102	165
2003.....	8,074	-	-	-	-	10	15	20	34	64	110	189
2002.....	8,132	-	-	-	1	15	9	18	37	52	112	172
2001.....	8,224	-	2	-	3	9	14	29	44	77	106	163
2000.....	8,797	-	-	-	2	6	12	19	37	52	97	147
1999.....	8,817	1	-	-	1	6	11	19	39	77	95	154

White, female

2004.....	4,963	-	-	1	1	1	6	7	16	18	27	48
2003.....	5,172	-	-	-	1	4	4	8	15	13	27	63
2002.....	5,153	2	-	1	-	3	4	3	15	30	25	48
2001.....	5,428	-	-	-	1	2	7	8	11	19	25	45
2000.....	5,413	-	-	-	-	3	5	8	13	23	32	46
1999.....	5,519	-	1	-	1	2	8	6	6	28	40	40

Black, both sexes

2004.....	1,228	-	1	-	-	-	6	11	22	41	52	87
2003.....	1,230	1	-	-	1	1	3	13	16	43	52	101
2002.....	1,218	-	-	-	3	1	4	9	25	19	61	96
2001.....	1,251	-	-	-	-	4	6	8	15	33	59	67
2000.....	1,277	-	-	-	3	1	8	8	17	25	65	68
1999.....	1,181	-	-	-	-	-	4	8	13	24	56	81

Black, male

2004.....	697	-	-	-	-	-	6	9	17	29	39	64
2003.....	686	-	-	-	1	-	3	11	15	28	38	71
2002.....	677	-	-	-	2	1	4	7	18	13	48	67
2001.....	671	-	-	-	-	3	4	5	11	24	46	45
2000.....	671	-	-	-	1	1	4	8	11	18	46	39
1999.....	619	-	-	-	-	-	3	6	13	17	35	57

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Aortic aneurysm and dissection (I71)												
Asian or Pacific Islander 2/, both sexes												
2004.....	298	-	-	-	-	-	4	2	4	3	6	5
2003.....	288	-	-	-	1	1	2	1	3	4	2	12
2002.....	266	-	-	-	-	1	2	-	2	2	3	9
2001.....	284	-	-	-	-	-	1	1	1	4	7	10
2000.....	282	-	-	-	-	-	1	3	2	3	6	6
1999.....	254	-	-	-	-	-	-	-	2	5	4	8
Asian or Pacific Islander 2/, male												
2004.....	194	-	-	-	-	-	2	1	3	3	5	3
2003.....	190	-	-	-	1	-	2	1	1	4	2	7
2002.....	174	-	-	-	-	1	2	-	2	1	2	8
2001.....	189	-	-	-	-	-	1	-	1	3	6	9
2000.....	172	-	-	-	-	-	-	3	-	2	4	4
1999.....	160	-	-	-	-	-	-	-	1	4	4	5
Asian or Pacific Islander 2/, female												
2004.....	104	-	-	-	-	-	2	1	1	-	1	2
2003.....	98	-	-	-	-	1	-	-	2	-	-	5
2002.....	92	-	-	-	-	-	-	-	-	1	1	1
2001.....	95	-	-	-	-	-	-	1	-	1	1	1
2000.....	110	-	-	-	-	-	1	-	2	1	2	2
1999.....	94	-	-	-	-	-	-	-	1	1	-	3

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other diseases of arteries, arterioles and capillaries (I72-I78)

All races, both sexes

2004.....	9,940	13	3	2	6	5	11	16	25	43	100	157
2003.....	9,862	10	-	1	8	14	13	19	20	43	92	156
2002.....	10,109	12	4	-	6	11	16	21	29	50	90	152
2001.....	10,084	12	2	2	5	8	10	21	32	60	106	134
2000.....	10,226	16	5	3	3	7	16	11	36	54	91	156
1999.....	10,002	11	1	2	6	6	12	18	30	44	105	135

All races, male

2004.....	4,166	6	2	2	3	5	4	8	12	15	47	91
2003.....	4,241	3	-	-	5	7	7	11	9	21	39	74
2002.....	4,296	6	4	-	2	2	9	5	14	14	48	82
2001.....	4,282	8	-	-	-	5	4	9	16	27	44	72
2000.....	4,423	6	2	1	3	2	8	6	19	20	37	84
1999.....	4,179	8	-	1	4	4	8	10	14	19	45	73

All races, female

2004.....	5,774	7	1	-	3	-	7	8	13	28	53	66
2003.....	5,621	7	-	1	3	7	6	8	11	22	53	82
2002.....	5,813	6	-	-	4	9	7	16	15	36	42	70
2001.....	5,802	4	2	2	5	3	6	12	16	33	62	62
2000.....	5,803	10	3	2	-	5	8	5	17	34	54	72
1999.....	5,823	3	1	1	2	2	4	8	16	25	60	62

White, both sexes

2004.....	8,445	10	3	1	5	5	9	11	16	31	74	122
2003.....	8,415	6	-	1	6	9	7	16	14	27	65	112
2002.....	8,650	9	4	-	2	8	10	14	21	33	67	110
2001.....	8,584	10	1	2	4	7	8	15	22	41	72	101
2000.....	8,753	13	4	2	3	5	13	8	26	43	61	110
1999.....	8,616	8	1	2	4	5	10	13	23	33	83	110

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other diseases of arteries, arterioles and capillaries (I72-I78)

White, male

2004.....	3,584	5	2	1	2	5	2	5	9	11	37	73
2003.....	3,634	2	-	-	4	5	4	10	6	15	28	61
2002.....	3,686	5	4	-	1	1	6	4	9	11	39	66
2001.....	3,663	6	-	-	-	5	3	7	10	16	29	53
2000.....	3,814	4	2	1	3	1	7	5	16	15	26	61
1999.....	3,627	5	-	1	3	3	7	7	12	15	39	65

White, female

2004.....	4,861	5	1	-	3	-	7	6	7	20	37	49
2003.....	4,781	4	-	1	2	4	3	6	8	12	37	51
2002.....	4,964	4	-	-	1	7	4	10	12	22	28	44
2001.....	4,921	4	1	2	4	2	5	8	12	25	43	48
2000.....	4,939	9	2	1	-	4	6	3	10	28	35	49
1999.....	4,989	3	1	1	1	2	3	6	11	18	44	45

Black, both sexes

2004.....	1,322	2	-	1	1	-	1	5	8	10	25	31
2003.....	1,289	4	-	-	2	5	5	3	5	14	24	39
2002.....	1,307	3	-	-	3	2	5	6	7	13	21	39
2001.....	1,381	2	1	-	1	1	1	6	10	17	33	31
2000.....	1,318	3	-	1	-	1	3	3	8	9	25	39
1999.....	1,253	3	-	-	-	1	2	4	5	11	21	23

Black, male

2004.....	528	1	-	1	1	-	1	3	2	2	10	15
2003.....	530	1	-	-	1	2	2	1	2	6	10	13
2002.....	534	1	-	-	-	-	2	1	5	2	8	13
2001.....	558	2	-	-	-	-	-	2	6	9	15	18
2000.....	536	2	-	-	-	-	1	1	2	4	9	21
1999.....	492	3	-	-	-	1	1	2	2	4	5	8

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Black, female												
2004.....	794	1	-	-	-	-	-	2	6	8	15	16
2003.....	759	3	-	-	1	3	3	2	3	8	14	26
2002.....	773	2	-	-	3	2	3	5	2	11	13	26
2001.....	823	-	1	-	1	1	1	4	4	8	18	13
2000.....	782	1	-	1	-	1	2	2	6	5	16	18
1999.....	761	-	-	-	-	-	1	2	3	7	16	15
American Indian 1/, both sexes												
2004.....	48	-	-	-	-	-	-	-	1	1	-	2
2003.....	56	-	-	-	-	-	1	-	1	1	-	2
2002.....	46	-	-	-	-	-	1	-	-	1	-	1
2001.....	32	-	-	-	-	-	-	-	-	-	-	1
2000.....	46	-	-	-	-	-	-	-	-	1	2	2
1999.....	36	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, male												
2004.....	15	-	-	-	-	-	-	-	1	1	-	1
2003.....	29	-	-	-	-	-	1	-	1	-	-	-
2002.....	23	-	-	-	-	-	1	-	-	1	-	1
2001.....	12	-	-	-	-	-	-	-	-	-	-	-
2000.....	16	-	-	-	-	-	-	-	-	-	-	1
1999.....	13	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, female												
2004.....	33	-	-	-	-	-	-	-	-	-	-	1
2003.....	27	-	-	-	-	-	-	-	-	1	-	2
2002.....	23	-	-	-	-	-	-	-	-	-	-	-
2001.....	20	-	-	-	-	-	-	-	-	-	-	1
2000.....	30	-	-	-	-	-	-	-	-	1	2	1
1999.....	23	-	-	-	1	-	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Asian or Pacific Islander 2/, both sexes												
2004.....	125	1	-	-	-	-	1	-	-	1	1	2
2003.....	102	-	-	-	-	-	-	-	-	1	3	3
2002.....	106	-	-	-	1	1	-	1	1	3	2	2
2001.....	87	-	-	-	-	-	1	-	-	2	1	1
2000.....	109	-	1	-	-	1	-	-	2	1	3	5
1999.....	97	-	-	-	1	-	-	1	2	-	1	2
Asian or Pacific Islander 2/, male												
2004.....	39	-	-	-	-	-	1	-	-	1	-	2
2003.....	48	-	-	-	-	-	-	-	-	-	1	-
2002.....	53	-	-	-	1	1	-	-	-	-	1	2
2001.....	49	-	-	-	-	-	1	-	-	2	-	1
2000.....	57	-	-	-	-	1	-	-	1	1	2	1
1999.....	47	-	-	-	1	-	-	1	-	-	1	-
Asian or Pacific Islander 2/, female												
2004.....	86	1	-	-	-	-	-	-	-	-	1	-
2003.....	54	-	-	-	-	-	-	-	-	1	2	3
2002.....	53	-	-	-	-	-	-	1	1	3	1	-
2001.....	38	-	-	-	-	-	-	-	-	-	1	-
2000.....	52	-	1	-	-	-	-	-	1	-	1	4
1999.....	50	-	-	-	-	-	-	-	2	-	-	2

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of circulatory system (I80-I99)												
All races, both sexes												
2004.....	4,673	29	2	1	4	6	29	54	83	107	206	282
2003.....	4,737	36	4	1	3	14	36	53	86	126	216	273
2002.....	4,711	27	4	1	1	12	23	38	101	128	177	251
2001.....	4,665	22	5	-	-	10	28	45	79	100	185	222
2000.....	4,603	27	4	5	7	10	27	58	75	125	194	249
1999.....	4,025	23	1	1	1	11	19	30	63	98	173	202
All races, male												
2004.....	2,083	18	-	1	2	3	16	29	43	51	111	157
2003.....	1,966	21	3	1	1	5	15	23	37	57	110	147
2002.....	1,944	15	1	-	-	9	12	12	57	67	87	137
2001.....	1,981	11	5	-	-	5	12	18	35	51	98	127
2000.....	1,981	16	2	3	3	6	10	18	39	70	120	145
1999.....	1,714	13	1	1	-	6	7	14	30	55	104	102
All races, female												
2004.....	2,590	11	2	-	2	3	13	25	40	56	95	125
2003.....	2,771	15	1	-	2	9	21	30	49	69	106	126
2002.....	2,767	12	3	1	1	3	11	26	44	61	90	114
2001.....	2,684	11	-	-	-	5	16	27	44	49	87	95
2000.....	2,622	11	2	2	4	4	17	40	36	55	74	104
1999.....	2,311	10	-	-	1	5	12	16	33	43	69	100
White, both sexes												
2004.....	3,795	22	2	1	2	6	17	31	51	68	130	207
2003.....	3,843	28	4	1	2	9	21	36	51	78	146	202
2002.....	3,853	20	2	-	1	8	13	23	61	82	121	185
2001.....	3,843	17	3	-	-	5	16	30	50	61	128	163
2000.....	3,787	20	3	4	5	4	18	33	49	87	143	189
1999.....	3,345	18	-	1	1	4	11	21	43	66	117	146

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
-------------	-----------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Other disorders of circulatory system (I80-I99)

White, male

2004.....	1,683	15	-	1	1	3	7	16	25	32	76	125
2003.....	1,569	16	3	1	-	3	8	11	25	45	84	103
2002.....	1,545	13	-	-	-	6	6	35	44	58	97	
2001.....	1,603	8	3	-	-	3	7	11	23	32	67	102
2000.....	1,618	13	1	3	2	3	6	9	25	47	94	111
1999.....	1,399	9	-	1	-	3	5	10	20	37	69	73

White, female

2004.....	2,112	7	2	-	1	3	10	15	26	36	54	82
2003.....	2,274	12	1	-	2	6	13	25	26	33	62	99
2002.....	2,308	7	2	-	1	2	7	17	26	38	63	88
2001.....	2,240	9	-	-	-	2	9	19	27	29	61	61
2000.....	2,169	7	2	1	3	1	12	24	24	40	49	78
1999.....	1,946	9	-	-	1	1	6	11	23	29	48	73

Black, both sexes

2004.....	799	7	-	-	1	-	10	21	30	37	74	65
2003.....	824	6	-	-	-	5	14	17	32	45	67	69
2002.....	797	7	1	1	-	4	9	15	39	43	52	60
2001.....	742	5	2	-	-	3	11	15	28	36	53	52
2000.....	761	5	1	1	2	6	8	23	26	36	51	58
1999.....	616	4	1	-	-	6	8	9	19	29	52	55

Black, male

2004.....	359	3	-	-	-	-	7	11	17	19	34	27
2003.....	368	5	-	-	-	2	7	12	11	11	25	42
2002.....	366	2	1	-	-	3	6	6	21	21	28	36
2001.....	339	3	2	-	-	1	4	7	11	17	30	20
2000.....	335	2	1	-	1	3	4	8	14	22	26	34
1999.....	282	3	1	-	-	2	2	4	10	16	33	29

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of circulatory system (I80-I99)												
Black, female												
2004.....	440	4	-	-	1	-	3	10	13	18	40	38
2003.....	456	1	-	-	-	3	7	5	21	34	42	27
2002.....	431	5	-	1	-	1	3	9	18	22	24	24
2001.....	403	2	-	-	-	2	7	8	17	19	23	32
2000.....	426	3	-	1	1	3	4	15	12	14	25	24
1999.....	334	1	-	-	-	4	6	5	9	13	19	26
American Indian 1/, both sexes												
2004.....	20	-	-	-	-	-	-	2	-	1	1	2
2003.....	15	-	-	-	-	-	-	-	-	-	2	1
2002.....	23	-	-	-	-	-	-	-	-	1	1	4
2001.....	25	-	-	-	-	1	-	-	1	2	-	4
2000.....	18	-	-	-	-	-	1	1	-	-	-	1
1999.....	13	-	-	-	-	-	-	-	1	-	1	-
American Indian 1/, male												
2004.....	12	-	-	-	-	-	-	2	-	-	1	2
2003.....	7	-	-	-	-	-	-	-	-	-	1	1
2002.....	13	-	-	-	-	-	-	-	-	1	-	2
2001.....	14	-	-	-	-	-	-	-	1	1	-	3
2000.....	8	-	-	-	-	-	-	1	-	-	-	-
1999.....	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, female												
2004.....	8	-	-	-	-	-	-	-	-	1	-	-
2003.....	8	-	-	-	-	-	-	-	-	-	1	-
2002.....	10	-	-	-	-	-	-	-	-	-	1	2
2001.....	11	-	-	-	-	1	-	-	-	1	-	1
2000.....	10	-	-	-	-	-	1	-	-	-	-	1
1999.....	9	-	-	-	-	-	-	-	1	-	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of circulatory system (I80-I99)												
Asian or Pacific Islander 2/, both sexes												
2004.....	59	-	-	-	1	-	2	-	2	1	1	8
2003.....	55	2	-	-	1	-	1	-	3	3	1	1
2002.....	38	-	1	-	-	-	1	-	1	2	3	2
2001.....	55	-	-	-	-	1	1	-	-	1	4	3
2000.....	37	2	-	-	-	-	-	1	-	2	-	1
1999.....	51	1	-	-	-	1	-	-	-	3	3	1
Asian or Pacific Islander 2/, male												
2004.....	29	-	-	-	1	-	2	-	1	-	-	3
2003.....	22	-	-	-	1	-	-	-	1	1	-	1
2002.....	20	-	-	-	-	-	-	-	1	1	1	2
2001.....	25	-	-	-	-	1	1	-	-	1	1	2
2000.....	20	1	-	-	-	-	-	-	-	1	-	-
1999.....	29	1	-	-	-	1	-	-	-	2	2	-
Asian or Pacific Islander 2/, female												
2004.....	30	-	-	-	-	-	-	-	1	1	1	5
2003.....	33	2	-	-	-	-	1	-	2	2	1	-
2002.....	18	-	1	-	-	-	1	-	-	1	2	-
2001.....	30	-	-	-	-	-	-	-	-	-	3	1
2000.....	17	1	-	-	-	-	-	1	-	1	-	1
1999.....	22	-	-	-	-	-	-	-	-	1	1	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)

All races, both sexes

2004.....	3,489	6	-	-	4	5	23	50	76	94	175	206
2003.....	3,623	7	-	1	3	14	33	52	77	107	180	210
2002.....	3,599	6	-	-	-	10	23	35	92	108	150	194
2001.....	3,531	3	2	-	-	7	26	44	72	86	146	165
2000.....	3,410	7	2	3	5	10	27	51	70	107	142	181
1999.....	2,625	4	1	1	1	9	13	28	55	79	130	135

All races, male

2004.....	1,471	4	-	-	2	3	12	27	39	45	88	98
2003.....	1,426	2	-	1	1	5	13	23	32	51	85	104
2002.....	1,394	2	-	-	-	8	12	11	50	54	66	93
2001.....	1,398	2	2	-	-	3	11	18	31	42	69	81
2000.....	1,368	6	1	2	2	6	10	17	35	58	82	93
1999.....	1,034	3	1	1	-	4	6	13	26	43	67	62

All races, female

2004.....	2,018	2	-	-	2	2	11	23	37	49	87	108
2003.....	2,197	5	-	-	2	9	20	29	45	56	95	106
2002.....	2,205	4	-	-	-	2	11	24	42	54	84	101
2001.....	2,133	1	-	-	-	4	15	26	41	44	77	84
2000.....	2,042	1	1	1	3	4	17	34	35	49	60	88
1999.....	1,591	1	-	-	1	5	7	15	29	36	63	73

White, both sexes

2004.....	2,795	4	-	-	2	5	15	29	45	58	109	141
2003.....	2,914	5	-	1	2	9	18	35	45	70	118	151
2002.....	2,923	4	-	-	-	7	13	22	53	68	100	144
2001.....	2,884	1	1	-	-	4	14	29	44	51	99	116
2000.....	2,786	6	2	2	3	4	18	27	46	71	105	134
1999.....	2,163	4	-	1	1	3	6	19	38	52	87	97

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)

White, male

2004.....	1,170	3	-	-	1	3	6	14	22	28	60	72
2003.....	1,124	1	-	1	-	3	6	11	22	40	63	69
2002.....	1,099	2	-	-	-	6	6	6	29	35	42	66
2001.....	1,122	1	1	-	-	2	6	11	19	25	45	63
2000.....	1,112	5	1	2	1	3	6	9	23	37	63	69
1999.....	843	3	-	1	-	2	4	9	16	28	45	47

White, female

2004.....	1,625	1	-	-	1	2	9	15	23	30	49	69
2003.....	1,790	4	-	-	2	6	12	24	23	30	55	82
2002.....	1,824	2	-	-	-	1	7	16	24	33	58	78
2001.....	1,762	-	-	-	-	2	8	18	25	26	54	53
2000.....	1,674	1	1	-	2	1	12	18	23	34	42	65
1999.....	1,320	1	-	-	1	1	2	10	22	24	42	50

Black, both sexes

2004.....	644	2	-	-	1	-	7	19	30	34	65	57
2003.....	668	2	-	-	-	5	14	17	30	34	61	58
2002.....	639	2	-	-	-	3	9	13	38	38	47	47
2001.....	602	2	1	-	-	3	11	15	27	33	43	45
2000.....	595	1	-	1	2	6	8	23	24	35	37	45
1999.....	436	-	1	-	-	6	7	9	17	25	41	37

Black, male

2004.....	277	1	-	-	-	-	5	11	17	17	28	22
2003.....	286	1	-	-	-	2	7	12	10	10	22	34
2002.....	275	-	-	-	-	2	6	5	20	18	23	25
2001.....	260	1	1	-	-	1	4	7	11	16	23	16
2000.....	243	1	-	-	1	3	4	8	12	21	19	24
1999.....	183	-	1	-	-	2	2	4	10	13	22	15

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

All other and unspecified disorders of circulatory system (I83-I99)

All races, both sexes

2004.....	1,184	23	2	1	-	1	6	4	7	13	31	76
2003.....	1,114	29	4	-	-	-	3	1	9	19	36	63
2002.....	1,112	21	4	1	1	2	-	3	9	20	27	57
2001.....	1,134	19	3	-	-	3	2	1	7	14	39	57
2000.....	1,193	20	2	2	2	-	-	7	5	18	52	68
1999.....	1,400	19	-	-	-	2	6	2	8	19	43	67

All races, male

2004.....	612	14	-	1	-	-	4	2	4	6	23	59
2003.....	540	19	3	-	-	-	2	-	5	6	25	43
2002.....	550	13	1	-	-	1	-	1	7	13	21	44
2001.....	583	9	3	-	-	2	1	-	4	9	29	46
2000.....	613	10	1	1	1	-	-	1	4	12	38	52
1999.....	680	10	-	-	-	2	1	1	4	12	37	40

All races, female

2004.....	572	9	2	-	-	1	2	2	3	7	8	17
2003.....	574	10	1	-	-	-	1	1	4	13	11	20
2002.....	562	8	3	1	1	1	-	2	2	7	6	13
2001.....	551	10	-	-	-	1	1	1	3	5	10	11
2000.....	580	10	1	1	1	-	-	6	1	6	14	16
1999.....	720	9	-	-	-	-	5	1	4	7	6	27

White, both sexes

2004.....	1,000	18	2	1	-	1	2	2	6	10	21	66
2003.....	929	23	4	-	-	-	3	1	6	8	28	51
2002.....	930	16	2	-	1	1	-	1	8	14	21	41
2001.....	959	16	2	-	-	1	2	1	6	10	29	47
2000.....	1,001	14	1	2	2	-	-	6	3	16	38	55
1999.....	1,182	14	-	-	-	1	5	2	5	14	30	49

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

All other and unspecified disorders of circulatory system (I83-I99)

White, male

2004.....	513	12	-	1	-	-	1	2	3	4	16	53
2003.....	445	15	3	-	-	-	2	-	3	5	21	34
2002.....	446	11	-	-	-	-	-	-	6	9	16	31
2001.....	481	7	2	-	-	1	1	-	4	7	22	39
2000.....	506	8	-	1	1	-	-	-	2	10	31	42
1999.....	556	6	-	-	-	1	1	1	4	9	24	26

White, female

2004.....	487	6	2	-	-	1	1	-	3	6	5	13
2003.....	484	8	1	-	-	-	1	1	3	3	7	17
2002.....	484	5	2	-	1	1	-	1	2	5	5	10
2001.....	478	9	-	-	-	-	1	1	2	3	7	8
2000.....	495	6	1	1	1	-	-	6	1	6	7	13
1999.....	626	8	-	-	-	-	4	1	1	5	6	23

Black, both sexes

2004.....	155	5	-	-	-	-	3	2	-	3	9	8
2003.....	156	4	-	-	-	-	-	-	2	11	6	11
2002.....	158	5	1	1	-	1	-	2	1	5	5	13
2001.....	140	3	1	-	-	-	-	-	1	3	10	7
2000.....	166	4	1	-	-	-	-	-	2	1	14	13
1999.....	180	4	-	-	-	-	1	-	2	4	11	18

Black, male

2004.....	82	2	-	-	-	-	2	-	-	2	6	5
2003.....	82	4	-	-	-	-	-	-	1	1	3	8
2002.....	91	2	1	-	-	1	-	1	1	3	5	11
2001.....	79	2	1	-	-	-	-	-	-	1	7	4
2000.....	92	1	1	-	-	-	-	-	2	1	7	10
1999.....	99	3	-	-	-	-	-	-	-	3	11	14

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

All other and unspecified disorders of circulatory system (I83-I99)

Black, female

2004.....	73	3	-	-	-	-	1	2	-	1	3	3
2003.....	74	-	-	-	-	-	-	-	1	10	3	3
2002.....	67	3	-	1	-	-	-	1	-	2	-	2
2001.....	61	1	-	-	-	-	-	-	1	2	3	3
2000.....	74	3	-	-	-	-	-	-	-	-	7	3
1999.....	81	1	-	-	-	-	1	-	2	1	-	4

American Indian 1/, both sexes

2004.....	10	-	-	-	-	-	-	-	-	-	1	1
2003.....	6	-	-	-	-	-	-	-	-	-	2	-
2002.....	9	-	-	-	-	-	-	-	-	-	1	2
2001.....	11	-	-	-	-	1	-	-	-	1	-	1
2000.....	6	-	-	-	-	-	-	1	-	-	-	-
1999.....	10	-	-	-	-	-	-	-	1	-	-	-

American Indian 1/, male

2004.....	6	-	-	-	-	-	-	-	-	-	1	1
2003.....	3	-	-	-	-	-	-	-	-	-	1	-
2002.....	5	-	-	-	-	-	-	-	-	-	-	1
2001.....	6	-	-	-	-	-	-	-	-	1	-	1
2000.....	3	-	-	-	-	-	-	1	-	-	-	-
1999.....	3	-	-	-	-	-	-	-	-	-	-	-

American Indian 1/, female

2004.....	4	-	-	-	-	-	-	-	-	-	-	-
2003.....	3	-	-	-	-	-	-	-	-	-	1	-
2002.....	4	-	-	-	-	-	-	-	-	-	1	1
2001.....	5	-	-	-	-	1	-	-	-	-	-	-
2000.....	3	-	-	-	-	-	-	-	-	-	-	-
1999.....	7	-	-	-	-	-	-	-	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

All other and unspecified disorders of circulatory system (I83-I99)

Asian or Pacific Islander 2/, both sexes

2004.....	19	-	-	-	-	-	1	-	1	-	-	1
2003.....	23	2	-	-	-	-	-	-	1	-	-	1
2002.....	15	-	1	-	-	-	-	-	-	1	-	1
2001.....	24	-	-	-	-	1	-	-	-	-	-	2
2000.....	20	2	-	-	-	-	-	-	-	1	-	-
1999.....	28	1	-	-	-	1	-	-	-	1	2	-

Asian or Pacific Islander 2/, male

2004.....	11	-	-	-	-	-	1	-	1	-	-	-
2003.....	10	-	-	-	-	-	-	-	1	-	-	1
2002.....	8	-	-	-	-	-	-	-	-	1	-	1
2001.....	17	-	-	-	-	1	-	-	-	-	-	2
2000.....	12	1	-	-	-	-	-	-	-	1	-	-
1999.....	22	1	-	-	-	1	-	-	-	-	2	-

Asian or Pacific Islander 2/, female

2004.....	8	-	-	-	-	-	-	-	-	-	-	1
2003.....	13	2	-	-	-	-	-	-	-	-	-	-
2002.....	7	-	1	-	-	-	-	-	-	-	-	-
2001.....	7	-	-	-	-	-	-	-	-	-	-	-
2000.....	8	1	-	-	-	-	-	-	-	-	-	-
1999.....	6	-	-	-	-	-	-	-	-	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
X. Diseases of the respiratory system (J00-J98)												
All races, both sexes												
2004.....	227,093	646	297	115	161	223	308	323	480	839	1,648	2,790
2003.....	235,935	699	342	144	203	237	367	373	564	937	1,684	2,966
2002.....	234,629	636	290	134	201	236	296	355	555	972	1,740	2,789
2001.....	230,009	696	281	126	166	205	323	355	572	1,037	1,681	2,628
2000.....	231,079	651	263	142	178	227	325	394	592	1,016	1,797	2,654
1999.....	229,863	687	315	138	191	252	289	412	596	1,060	1,653	2,541
All races, male												
2004.....	108,664	366	150	69	95	131	182	191	286	446	855	1,481
2003.....	111,820	421	200	82	112	137	211	222	322	491	850	1,626
2002.....	111,775	388	175	70	97	148	164	187	299	499	951	1,494
2001.....	109,322	407	148	70	90	114	190	179	303	544	900	1,467
2000.....	110,716	369	136	81	97	135	187	219	321	535	995	1,468
1999.....	111,273	380	172	76	117	140	166	213	313	558	868	1,407
All races, female												
2004.....	118,429	280	147	46	66	92	126	132	194	393	793	1,309
2003.....	124,115	278	142	62	91	100	156	151	242	446	834	1,340
2002.....	122,854	248	115	64	104	88	132	168	256	473	789	1,295
2001.....	120,687	289	133	56	76	91	133	176	269	493	781	1,161
2000.....	120,363	282	127	61	81	92	138	175	271	481	802	1,186
1999.....	118,590	307	143	62	74	112	123	199	283	502	785	1,134
White, both sexes												
2004.....	205,826	388	181	58	103	145	213	216	335	578	1,212	2,090
2003.....	214,096	450	217	91	133	144	251	244	386	666	1,192	2,242
2002.....	213,080	375	174	88	115	149	200	249	383	673	1,257	2,134
2001.....	208,523	386	185	80	108	132	205	223	368	678	1,212	1,960
2000.....	209,758	394	177	93	108	141	218	258	405	697	1,291	1,942
1999.....	208,671	396	191	95	119	179	183	247	389	732	1,119	1,839

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	-------------	-----------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

X. Diseases of the respiratory system (J00-J98)

White, male

2004.....	97,611	219	94	30	60	85	134	121	199	302	627	1,119
2003.....	100,310	270	124	50	69	76	146	146	215	357	607	1,253
2002.....	100,565	234	105	49	50	89	113	126	210	334	692	1,139
2001.....	98,223	230	91	46	57	68	130	109	192	370	661	1,126
2000.....	99,472	230	81	55	55	81	131	134	210	366	724	1,075
1999.....	100,060	224	91	49	74	100	103	135	210	389	578	1,023

White, female

2004.....	108,215	169	87	28	43	60	79	95	136	276	585	971
2003.....	113,786	180	93	41	64	68	105	98	171	309	585	989
2002.....	112,515	141	69	39	65	60	87	123	173	339	565	995
2001.....	110,300	156	94	34	51	64	75	114	176	308	551	834
2000.....	110,286	164	96	38	53	60	87	124	195	331	567	867
1999.....	108,611	172	100	46	45	79	80	112	179	343	541	816

Black, both sexes

2004.....	17,077	221	95	49	53	73	81	97	133	237	402	624
2003.....	17,509	216	89	46	59	77	105	117	159	241	441	651
2002.....	17,621	230	104	41	81	78	89	93	156	273	451	590
2001.....	17,541	258	79	39	53	62	105	114	189	326	429	613
2000.....	17,580	232	77	41	62	78	96	118	167	282	448	640
1999.....	17,783	252	100	38	67	63	97	154	190	300	485	642

Black, male

2004.....	8,818	125	46	35	32	43	41	65	80	130	210	327
2003.....	9,097	130	59	30	36	50	60	71	98	118	210	336
2002.....	9,028	135	60	19	43	55	48	50	83	146	244	315
2001.....	8,943	146	48	21	30	41	51	58	103	156	218	312
2000.....	9,128	123	50	25	37	49	50	75	100	148	244	354
1999.....	9,286	137	64	25	41	36	57	72	95	151	256	353

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

X. Diseases of the respiratory system (J00-J98)

Black, female

2004.....	8,259	96	49	14	21	30	40	32	53	107	192	297
2003.....	8,412	86	30	16	23	27	45	46	61	123	231	315
2002.....	8,593	95	44	22	38	23	41	43	73	127	207	275
2001.....	8,598	112	31	18	23	21	54	56	86	170	211	301
2000.....	8,452	109	27	16	25	29	46	43	67	134	204	286
1999.....	8,497	115	36	13	26	27	40	82	95	149	229	289

American Indian 1/, both sexes

2004.....	1,026	21	7	4	-	1	7	5	5	11	16	36
2003.....	1,144	16	8	2	3	6	2	3	7	16	26	41
2002.....	972	12	3	1	-	4	4	3	9	13	16	29
2001.....	992	27	9	2	3	5	4	9	7	18	22	22
2000.....	927	9	4	2	2	3	3	11	6	19	27	26
1999.....	927	14	6	2	2	2	5	5	7	7	29	30

American Indian 1/, male

2004.....	502	12	2	2	-	1	3	2	4	5	9	16
2003.....	564	10	2	1	2	4	1	-	2	7	18	21
2002.....	476	7	3	-	-	1	2	3	3	10	6	18
2001.....	480	18	5	-	1	1	4	8	4	11	12	10
2000.....	479	3	2	-	-	3	2	5	4	10	17	13
1999.....	453	8	6	1	1	2	3	2	3	2	18	17

American Indian 1/, female

2004.....	524	9	5	2	-	-	4	3	1	6	7	20
2003.....	580	6	6	1	1	2	1	3	5	9	8	20
2002.....	496	5	-	1	-	3	2	-	6	3	10	11
2001.....	512	9	4	2	2	4	-	1	3	7	10	12
2000.....	448	6	2	2	2	-	1	6	2	9	10	13
1999.....	474	6	-	1	1	-	2	3	4	5	11	13

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

X. Diseases of the respiratory system (J00-J98)

Asian or Pacific Islander 2/, both sexes

2004.....	3,164	16	14	4	5	4	7	5	7	13	18	40
2003.....	3,186	17	28	5	8	10	9	9	12	14	25	32
2002.....	2,956	19	9	4	5	5	3	10	7	13	16	36
2001.....	2,953	25	8	5	2	6	9	9	8	15	18	33
2000.....	2,814	16	5	6	6	5	8	7	14	18	31	46
1999.....	2,482	25	18	3	3	8	4	6	10	21	20	30

Asian or Pacific Islander 2/, male

2004.....	1,733	10	8	2	3	2	4	3	3	9	9	19
2003.....	1,849	11	15	1	5	7	4	5	7	9	15	16
2002.....	1,706	12	7	2	4	3	1	8	3	9	9	22
2001.....	1,676	13	4	3	2	4	5	4	4	7	9	19
2000.....	1,637	13	3	1	5	2	4	5	7	11	10	26
1999.....	1,474	11	11	1	1	2	3	4	5	16	16	14

Asian or Pacific Islander 2/, female

2004.....	1,431	6	6	2	2	2	3	2	4	4	9	21
2003.....	1,337	6	13	4	3	3	5	4	5	5	10	16
2002.....	1,250	7	2	2	1	2	2	2	4	4	7	14
2001.....	1,277	12	4	2	-	2	4	5	4	8	9	14
2000.....	1,177	3	2	5	1	3	4	2	7	7	21	20
1999.....	1,008	14	7	2	2	6	1	2	5	5	4	16

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute upper respiratory infections (J00-J06)												
All races, both sexes												
2004.....	149	5	13	5	1	6	2	2	2	4	8	5
2003.....	192	14	15	1	2	3	4	2	6	7	6	4
2002.....	164	9	13	1	2	2	5	1	1	2	8	7
2001.....	169	9	16	4	4	5	2	4	2	1	6	7
2000.....	205	6	7	3	1	3	6	6	6	3	8	6
1999.....	212	10	15	3	2	2	-	4	7	8	6	7
All races, male												
2004.....	67	4	6	1	1	3	1	1	2	3	6	5
2003.....	95	9	8	-	2	1	1	2	4	6	4	1
2002.....	66	5	7	-	-	2	2	-	1	2	5	3
2001.....	61	7	8	3	2	2	2	3	1	1	4	5
2000.....	79	5	2	1	-	-	3	5	5	2	6	2
1999.....	98	6	11	3	1	2	-	3	6	6	6	5
All races, female												
2004.....	82	1	7	4	-	3	1	1	-	1	2	-
2003.....	97	5	7	1	-	2	3	-	2	1	2	3
2002.....	98	4	6	1	2	-	3	1	-	-	3	4
2001.....	108	2	8	1	2	3	-	1	1	-	2	2
2000.....	126	1	5	2	1	3	3	1	1	1	2	4
1999.....	114	4	4	-	1	-	-	1	1	2	-	2
White, both sexes												
2004.....	128	5	11	5	1	5	2	1	2	2	4	5
2003.....	156	11	12	-	1	3	3	2	5	4	3	2
2002.....	138	4	11	1	2	2	2	1	1	2	6	5
2001.....	142	5	12	4	2	4	2	3	1	-	4	6
2000.....	180	5	6	1	-	3	3	4	6	2	8	4
1999.....	176	5	12	1	2	1	-	4	3	5	6	5

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Acute upper respiratory infections (J00-J06)

White, male

2004.....	54	4	5	1	1	2	1	1	2	2	3	5
2003.....	71	8	6	-	1	1	-	2	3	4	2	-
2002.....	52	1	7	-	-	2	1	-	1	2	3	3
2001.....	46	3	5	3	1	1	2	3	-	-	3	4
2000.....	66	5	1	-	-	-	1	4	5	1	6	2
1999.....	74	3	8	1	1	1	-	3	3	5	6	3

White, female

2004.....	74	1	6	4	-	3	1	-	-	-	1	-
2003.....	85	3	6	-	-	2	3	-	2	-	1	2
2002.....	86	3	4	1	2	-	1	1	-	-	3	2
2001.....	96	2	7	1	1	3	-	-	1	-	1	2
2000.....	114	-	5	1	-	3	2	-	1	1	2	2
1999.....	102	2	4	-	1	-	-	1	-	-	-	2

Black, both sexes

2004.....	15	-	-	-	-	1	-	-	-	2	4	-
2003.....	30	3	3	1	-	-	1	-	-	3	3	2
2002.....	19	4	2	-	-	-	2	-	-	-	2	2
2001.....	24	3	4	-	2	1	-	-	1	1	2	1
2000.....	17	1	1	1	1	-	3	2	-	-	-	2
1999.....	30	4	3	2	-	1	-	-	4	2	-	1

Black, male

2004.....	10	-	-	-	-	1	-	-	-	1	3	-
2003.....	19	1	2	-	-	-	1	-	-	2	2	1
2002.....	10	3	-	-	-	-	1	-	-	-	2	-
2001.....	14	3	3	-	1	1	-	-	1	1	1	1
2000.....	10	-	1	1	-	-	2	1	-	-	-	-
1999.....	19	2	3	2	-	1	-	-	3	-	-	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute pharyngitis and tonsillitis (J02-J03)												
All races, both sexes												
2004.....	22	1	-	4	-	2	1	-	1	1	1	1
2003.....	20	1	1	-	-	1	3	1	1	1	1	2
2002.....	20	-	1	-	-	-	-	-	-	-	5	2
2001.....	30	1	3	-	2	2	1	2	1	1	1	1
2000.....	25	-	2	-	-	-	2	1	1	1	4	-
1999.....	17	1	1	-	-	1	-	2	1	2	-	1
All races, male												
2004.....	10	1	-	1	-	-	-	-	1	1	1	1
2003.....	7	-	-	-	-	-	1	1	1	1	-	-
2002.....	10	-	1	-	-	-	-	-	-	-	3	-
2001.....	13	1	1	-	-	1	1	2	-	1	-	1
2000.....	12	-	2	-	-	-	-	1	1	1	2	-
1999.....	9	-	-	-	-	1	-	2	1	2	-	-
All races, female												
2004.....	12	-	-	3	-	2	1	-	-	-	-	-
2003.....	13	1	1	-	-	1	2	-	-	-	1	2
2002.....	10	-	-	-	-	-	-	-	-	-	2	2
2001.....	17	-	2	-	2	1	-	-	1	-	1	-
2000.....	13	-	-	-	-	-	2	-	-	-	2	-
1999.....	8	1	1	-	-	-	-	-	-	-	-	1
White, both sexes												
2004.....	21	1	-	4	-	2	1	-	1	1	1	1
2003.....	13	-	1	-	-	1	2	1	-	1	-	2
2002.....	13	-	1	-	-	-	-	-	-	-	3	-
2001.....	26	1	2	-	1	2	1	2	1	-	-	1
2000.....	22	-	1	-	-	-	1	1	1	1	4	-
1999.....	13	-	1	-	-	-	-	2	1	2	-	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute pharyngitis and tonsillitis (J02-J03)												
White, male												
2004.....	9	1	-	1	-	-	-	-	1	1	1	1
2003.....	3	-	-	-	-	-	-	1	-	1	-	-
2002.....	6	-	1	-	-	-	-	-	-	-	1	-
2001.....	11	1	-	-	-	1	1	2	-	-	-	1
2000.....	11	-	1	-	-	-	-	1	1	1	2	-
1999.....	6	-	-	-	-	-	-	2	1	2	-	-
White, female												
2004.....	12	-	-	3	-	2	1	-	-	-	-	-
2003.....	10	-	1	-	-	1	2	-	-	-	-	2
2002.....	7	-	-	-	-	-	-	-	-	-	2	-
2001.....	15	-	2	-	1	1	-	-	1	-	-	-
2000.....	11	-	-	-	-	-	1	-	-	-	2	-
1999.....	7	-	1	-	-	-	-	-	-	-	-	1
Black, both sexes												
2004.....	1	-	-	-	-	-	-	-	-	-	-	-
2003.....	4	1	-	-	-	-	1	-	-	-	1	-
2002.....	7	-	-	-	-	-	-	-	-	-	2	2
2001.....	4	-	1	-	1	-	-	-	-	1	1	-
2000.....	3	-	1	-	-	-	1	-	-	-	-	-
1999.....	4	1	-	-	-	1	-	-	-	-	-	-
Black, male												
2004.....	1	-	-	-	-	-	-	-	-	-	-	-
2003.....	2	-	-	-	-	-	1	-	-	-	-	-
2002.....	4	-	-	-	-	-	-	-	-	-	2	-
2001.....	2	-	1	-	-	-	-	-	-	1	-	-
2000.....	1	-	1	-	-	-	-	-	-	-	-	-
1999.....	3	-	-	-	-	1	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Acute laryngitis and tracheitis (J04)

All races, both sexes

2004.....	25	2	6	1	1	2	1	-	-	-	-	1
2003.....	26	6	7	1	1	-	-	-	-	1	-	-
2002.....	23	8	5	-	-	-	3	-	-	-	1	1
2001.....	21	3	6	1	1	1	-	-	-	-	1	1
2000.....	16	3	2	1	-	1	1	1	1	-	-	1
1999.....	28	4	8	2	1	1	-	-	1	1	1	2

All races, male

2004.....	15	1	2	-	1	2	1	-	-	-	-	1
2003.....	13	2	3	-	1	-	-	-	-	1	-	-
2002.....	9	4	3	-	-	-	-	-	-	-	-	-
2001.....	13	3	3	1	1	-	-	-	-	-	1	1
2000.....	8	2	-	-	-	-	1	1	1	-	-	-
1999.....	19	3	7	2	1	1	-	-	1	1	1	1

All races, female

2004.....	10	1	4	1	-	-	-	-	-	-	-	-
2003.....	13	4	4	1	-	-	-	-	-	-	-	-
2002.....	14	4	2	-	-	-	3	-	-	-	1	1
2001.....	8	-	3	-	-	1	-	-	-	-	-	-
2000.....	8	1	2	1	-	1	-	-	-	-	-	1
1999.....	9	1	1	-	-	-	-	-	-	-	-	1

White, both sexes

2004.....	22	2	6	1	1	1	1	-	-	-	-	1
2003.....	19	4	7	-	1	-	-	-	-	-	-	-
2002.....	17	4	5	-	-	-	1	-	-	-	1	1
2001.....	17	1	4	1	1	1	-	-	-	-	1	1
2000.....	9	2	2	-	-	1	-	-	1	-	-	1
1999.....	21	1	7	-	1	1	-	-	-	1	1	2

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute laryngitis and tracheitis (J04)												
White, male												
2004.....	12	1	2	-	1	1	1	-	-	-	-	1
2003.....	9	1	3	-	1	-	-	-	-	-	-	-
2002.....	6	1	3	-	-	-	-	-	-	-	-	-
2001.....	10	1	2	1	1	-	-	-	-	-	1	1
2000.....	4	2	-	-	-	-	-	-	1	-	-	-
1999.....	13	1	6	-	1	1	-	-	-	1	1	1
White, female												
2004.....	10	1	4	1	-	-	-	-	-	-	-	-
2003.....	10	3	4	-	-	-	-	-	-	-	-	-
2002.....	11	3	2	-	-	-	1	-	-	-	1	1
2001.....	7	-	2	-	-	1	-	-	-	-	-	-
2000.....	5	-	2	-	-	1	-	-	-	-	-	1
1999.....	8	-	1	-	-	-	-	-	-	-	-	1
Black, both sexes												
2004.....	3	-	-	-	-	1	-	-	-	-	-	-
2003.....	7	2	-	1	-	-	-	-	-	1	-	-
2002.....	4	3	-	-	-	-	1	-	-	-	-	-
2001.....	4	2	2	-	-	-	-	-	-	-	-	-
2000.....	4	1	-	-	-	-	1	1	-	-	-	-
1999.....	6	2	1	2	-	-	-	-	1	-	-	-
Black, male												
2004.....	3	-	-	-	-	1	-	-	-	-	-	-
2003.....	4	1	-	-	-	-	-	-	-	1	-	-
2002.....	2	2	-	-	-	-	-	-	-	-	-	-
2001.....	3	2	1	-	-	-	-	-	-	-	-	-
2000.....	3	-	-	-	-	-	1	1	-	-	-	-
1999.....	5	1	1	2	-	-	-	-	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other and unspecified acute upper respiratory infections (J00-J01,J05-J06)

All races, both sexes

2004.....	102	2	7	-	-	2	-	2	1	3	7	3
2003.....	146	7	7	-	1	2	1	1	5	5	5	2
2002.....	121	1	7	1	2	2	1	1	2	2	2	4
2001.....	118	5	7	3	1	2	1	2	1	-	4	5
2000.....	164	3	3	2	1	2	3	4	4	2	4	5
1999.....	167	5	6	1	1	-	-	2	5	5	5	4

All races, male

2004.....	42	2	4	-	-	1	-	1	1	2	5	3
2003.....	75	7	5	-	1	1	-	1	3	4	4	1
2002.....	47	1	3	-	-	2	2	-	1	2	2	3
2001.....	35	3	4	2	1	1	1	1	1	-	3	3
2000.....	59	3	-	1	-	-	2	3	3	1	4	2
1999.....	70	3	4	1	-	-	-	1	4	3	5	4

All races, female

2004.....	60	-	3	-	-	1	-	1	-	1	2	-
2003.....	71	-	2	-	-	1	1	-	2	1	1	1
2002.....	74	-	4	1	2	-	-	1	-	-	-	1
2001.....	83	2	3	1	-	1	-	1	-	-	1	2
2000.....	105	-	3	1	1	2	1	1	1	1	-	3
1999.....	97	2	2	-	1	-	-	1	1	2	-	-

White, both sexes

2004.....	85	2	5	-	-	2	-	1	1	1	3	3
2003.....	124	7	4	-	-	2	1	1	5	3	3	-
2002.....	108	-	5	1	2	2	1	1	1	2	2	4
2001.....	99	3	6	3	-	1	1	1	-	-	3	4
2000.....	149	3	3	1	-	2	2	3	4	1	4	3
1999.....	142	4	4	1	1	-	-	2	2	2	5	2

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other and unspecified acute upper respiratory infections (J00-J01,J05-J06)

White, male

2004.....	33	2	3	-	-	1	-	1	1	1	2	3
2003.....	59	7	3	-	-	1	-	1	3	3	2	-
2002.....	40	-	3	-	-	2	1	-	1	2	2	3
2001.....	25	1	3	2	-	-	1	1	-	-	2	2
2000.....	51	3	-	-	-	-	1	3	3	-	4	2
1999.....	55	2	2	1	-	-	-	1	2	2	5	2

White, female

2004.....	52	-	2	-	-	1	-	-	-	-	1	-
2003.....	65	-	1	-	-	1	1	-	2	-	1	-
2002.....	68	-	2	1	2	-	-	1	-	-	-	1
2001.....	74	2	3	1	-	1	-	-	-	-	1	2
2000.....	98	-	3	1	-	2	1	-	1	1	-	1
1999.....	87	2	2	-	1	-	-	1	-	-	-	-

Black, both sexes

2004.....	11	-	-	-	-	-	-	-	-	2	4	-
2003.....	19	-	3	-	-	-	-	-	-	2	2	2
2002.....	8	1	2	-	-	-	1	-	-	-	-	-
2001.....	16	1	1	-	1	1	-	-	1	-	1	1
2000.....	10	-	-	1	1	-	1	1	-	-	-	2
1999.....	20	1	2	-	-	-	-	-	3	2	-	1

Black, male

2004.....	6	-	-	-	-	-	-	-	-	1	3	-
2003.....	13	-	2	-	-	-	-	-	-	1	2	1
2002.....	4	1	-	-	-	-	1	-	-	-	-	-
2001.....	9	1	1	-	1	1	-	-	1	-	1	1
2000.....	6	-	-	1	-	-	1	-	-	-	-	-
1999.....	11	1	2	-	-	-	-	-	2	-	-	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other diseases of the respiratory system (J10-J98)

All races, both sexes

2004.....	226,944	641	284	110	160	217	306	321	478	835	1,640	2,785
2003.....	235,743	685	327	143	201	234	363	371	558	930	1,678	2,962
2002.....	234,465	627	277	133	199	234	291	354	554	970	1,732	2,782
2001.....	229,840	687	265	122	162	200	321	351	570	1,036	1,675	2,621
2000.....	230,874	645	256	139	177	224	319	388	586	1,013	1,789	2,648
1999.....	229,651	677	300	135	189	250	289	408	589	1,052	1,647	2,534

All races, male

2004.....	108,597	362	144	68	94	128	181	190	284	443	849	1,476
2003.....	111,725	412	192	82	110	136	210	220	318	485	846	1,625
2002.....	111,709	383	168	70	97	146	162	187	298	497	946	1,491
2001.....	109,261	400	140	67	88	112	188	176	302	543	896	1,462
2000.....	110,637	364	134	80	97	135	184	214	316	533	989	1,466
1999.....	111,175	374	161	73	116	138	166	210	307	552	862	1,402

All races, female

2004.....	118,347	279	140	42	66	89	125	131	194	392	791	1,309
2003.....	124,018	273	135	61	91	98	153	151	240	445	832	1,337
2002.....	122,756	244	109	63	102	88	129	167	256	473	786	1,291
2001.....	120,579	287	125	55	74	88	133	175	268	493	779	1,159
2000.....	120,237	281	122	59	80	89	135	174	270	480	800	1,182
1999.....	118,476	303	139	62	73	112	123	198	282	500	785	1,132

White, both sexes

2004.....	205,698	383	170	53	102	140	211	215	333	576	1,208	2,085
2003.....	213,940	439	205	91	132	141	248	242	381	662	1,189	2,240
2002.....	212,942	371	163	87	113	147	198	248	382	671	1,251	2,129
2001.....	208,381	381	173	76	106	128	203	220	367	678	1,208	1,954
2000.....	209,578	389	171	92	108	138	215	254	399	695	1,283	1,938
1999.....	208,495	391	179	94	117	178	183	243	386	727	1,113	1,834

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other diseases of the respiratory system (J10-J98)

White, male

2004.....	97,557	215	89	29	59	83	133	120	197	300	624	1,114
2003.....	100,239	262	118	50	68	75	146	144	212	353	605	1,253
2002.....	100,513	233	98	49	50	87	112	126	209	332	689	1,136
2001.....	98,177	227	86	43	56	67	128	106	192	370	658	1,122
2000.....	99,406	225	80	55	55	81	130	130	205	365	718	1,073
1999.....	99,986	221	83	48	73	99	103	132	207	384	572	1,020

White, female

2004.....	108,141	168	81	24	43	57	78	95	136	276	584	971
2003.....	113,701	177	87	41	64	66	102	98	169	309	584	987
2002.....	112,429	138	65	38	63	60	86	122	173	339	562	993
2001.....	110,204	154	87	33	50	61	75	114	175	308	550	832
2000.....	110,172	164	91	37	53	57	85	124	194	330	565	865
1999.....	108,509	170	96	46	44	79	80	111	179	343	541	814

Black, both sexes

2004.....	17,062	221	95	49	53	72	81	97	133	235	398	624
2003.....	17,479	213	86	45	59	77	104	117	159	238	438	649
2002.....	17,602	226	102	41	81	78	87	93	156	273	449	588
2001.....	17,517	255	75	39	51	61	105	114	188	325	427	612
2000.....	17,563	231	76	40	61	78	93	116	167	282	448	638
1999.....	17,753	248	97	36	67	62	97	154	186	298	485	641

Black, male

2004.....	8,808	125	46	35	32	42	41	65	80	129	207	327
2003.....	9,078	129	57	30	36	50	59	71	98	116	208	335
2002.....	9,018	132	60	19	43	55	47	50	83	146	242	315
2001.....	8,929	143	45	21	29	40	51	58	102	155	217	311
2000.....	9,118	123	49	24	37	49	48	74	100	148	244	354
1999.....	9,267	135	61	23	41	35	57	72	92	151	256	352

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other diseases of the respiratory system (J10-J98)

Black, female

2004.....	8,254	96	49	14	21	30	40	32	53	106	191	297
2003.....	8,401	84	29	15	23	27	45	46	61	122	230	314
2002.....	8,584	94	42	22	38	23	40	43	73	127	207	273
2001.....	8,588	112	30	18	22	21	54	56	86	170	210	301
2000.....	8,445	108	27	16	24	29	45	42	67	134	204	284
1999.....	8,486	113	36	13	26	27	40	82	94	147	229	289

American Indian 1/, both sexes

2004.....	1,025	21	7	4	-	1	7	5	5	11	16	36
2003.....	1,143	16	8	2	3	6	2	3	7	16	26	41
2002.....	970	11	3	1	-	4	3	3	9	13	16	29
2001.....	992	27	9	2	3	5	4	9	7	18	22	22
2000.....	923	9	4	2	2	3	3	11	6	18	27	26
1999.....	925	14	6	2	2	2	5	5	7	7	29	29

American Indian 1/, male

2004.....	501	12	2	2	-	1	3	2	4	5	9	16
2003.....	564	10	2	1	2	4	1	-	2	7	18	21
2002.....	475	6	3	-	-	1	2	3	3	10	6	18
2001.....	480	18	5	-	1	1	4	8	4	11	12	10
2000.....	476	3	2	-	-	3	2	5	4	9	17	13
1999.....	452	8	6	1	1	2	3	2	3	2	18	16

American Indian 1/, female

2004.....	524	9	5	2	-	-	4	3	1	6	7	20
2003.....	579	6	6	1	1	2	1	3	5	9	8	20
2002.....	495	5	-	1	-	3	1	-	6	3	10	11
2001.....	512	9	4	2	2	4	-	1	3	7	10	12
2000.....	447	6	2	2	2	-	1	6	2	9	10	13
1999.....	473	6	-	1	1	-	2	3	4	5	11	13

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other diseases of the respiratory system (J10-J98)

Asian or Pacific Islander 2/, both sexes

2004.....	3,159	16	12	4	5	4	7	4	7	13	18	40
2003.....	3,181	17	28	5	7	10	9	9	11	14	25	32
2002.....	2,951	19	9	4	5	5	3	10	7	13	16	36
2001.....	2,950	24	8	5	2	6	9	8	8	15	18	33
2000.....	2,810	16	5	5	6	5	8	7	14	18	31	46
1999.....	2,478	24	18	3	3	8	4	6	10	20	20	30

Asian or Pacific Islander 2/, male

2004.....	1,731	10	7	2	3	2	4	3	3	9	9	19
2003.....	1,844	11	15	1	4	7	4	5	6	9	15	16
2002.....	1,703	12	7	2	4	3	1	8	3	9	9	22
2001.....	1,675	12	4	3	2	4	5	4	4	7	9	19
2000.....	1,637	13	3	1	5	2	4	5	7	11	10	26
1999.....	1,470	10	11	1	1	2	3	4	5	15	16	14

Asian or Pacific Islander 2/, female

2004.....	1,428	6	5	2	2	2	3	1	4	4	9	21
2003.....	1,337	6	13	4	3	3	5	4	5	5	10	16
2002.....	1,248	7	2	2	1	2	2	2	4	4	7	14
2001.....	1,275	12	4	2	-	2	4	4	4	8	9	14
2000.....	1,173	3	2	4	1	3	4	2	7	7	21	20
1999.....	1,008	14	7	2	2	6	1	2	5	5	4	16

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Influenza (J10-J11)												
All races, both sexes												
2004.....	1,100	18	18	3	10	4	3	-	4	8	8	14
2003.....	1,792	31	59	28	18	18	12	5	8	10	14	29
2002.....	727	7	5	5	7	5	2	3	1	3	5	8
2001.....	257	7	6	9	3	4	3	4	6	1	5	8
2000.....	1,765	9	10	3	8	8	5	8	5	8	14	24
1999.....	1,665	13	12	6	5	7	4	2	7	8	8	13
All races, male												
2004.....	421	10	8	2	6	1	1	-	2	1	6	3
2003.....	699	18	33	13	6	9	3	4	5	5	5	15
2002.....	289	5	3	1	1	2	2	2	-	2	2	7
2001.....	101	6	2	3	1	3	3	3	3	1	1	5
2000.....	664	5	4	1	4	3	3	4	1	5	9	11
1999.....	602	6	9	4	1	1	1	-	5	5	4	8
All races, female												
2004.....	679	8	10	1	4	3	2	-	2	7	2	11
2003.....	1,093	13	26	15	12	9	9	1	3	5	9	14
2002.....	438	2	2	4	6	3	-	1	1	1	3	1
2001.....	156	1	4	6	2	1	-	1	3	-	4	3
2000.....	1,101	4	6	2	4	5	2	4	4	3	5	13
1999.....	1,063	7	3	2	4	6	3	2	2	3	4	5
White, both sexes												
2004.....	1,033	12	14	2	9	3	3	-	3	7	7	11
2003.....	1,653	23	42	20	10	11	9	3	7	8	12	22
2002.....	686	5	2	4	6	3	2	3	1	3	5	8
2001.....	229	7	5	5	2	3	2	2	5	1	3	8
2000.....	1,662	7	9	2	7	8	4	7	4	7	11	18
1999.....	1,574	7	8	6	4	6	4	1	7	4	6	12

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Influenza (J10-J11)												
White, male												
2004.....	388	5	5	2	5	1	1	-	1	-	6	1
2003.....	640	14	25	9	3	5	2	3	5	3	3	11
2002.....	270	3	1	1	1	-	2	2	-	2	2	7
2001.....	88	6	2	2	1	2	2	1	2	1	1	5
2000.....	617	5	4	1	3	3	2	3	-	4	6	7
1999.....	569	1	6	4	1	1	1	-	5	3	3	7
White, female												
2004.....	645	7	9	-	4	2	2	-	2	7	1	10
2003.....	1,013	9	17	11	7	6	7	-	2	5	9	11
2002.....	416	2	1	3	5	3	-	1	1	1	3	1
2001.....	141	1	3	3	1	1	-	1	3	-	2	3
2000.....	1,045	2	5	1	4	5	2	4	4	3	5	11
1999.....	1,005	6	2	2	3	5	3	1	2	1	3	5
Black, both sexes												
2004.....	50	4	3	1	-	1	-	-	1	-	1	2
2003.....	97	7	6	6	5	4	3	2	1	-	1	7
2002.....	30	1	3	1	1	2	-	-	-	-	-	-
2001.....	20	-	-	2	-	1	-	1	1	-	1	-
2000.....	81	2	1	-	1	-	1	1	-	-	3	5
1999.....	66	4	1	-	-	1	-	1	-	3	2	1
Black, male												
2004.....	25	4	2	-	-	-	-	-	1	-	-	2
2003.....	37	3	3	3	1	2	1	1	-	-	1	4
2002.....	15	1	2	-	-	2	-	-	-	-	-	-
2001.....	10	-	-	-	-	1	-	1	1	-	-	-
2000.....	35	-	-	-	1	-	1	1	-	-	3	3
1999.....	24	3	1	-	-	-	-	-	-	1	1	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Influenza (J10-J11)												
Black, female												
2004.....	25	-	1	1	-	1	-	-	-	-	1	-
2003.....	60	4	3	3	4	2	2	1	1	-	-	3
2002.....	15	-	1	1	1	-	-	-	-	-	-	-
2001.....	10	-	-	2	-	-	-	-	-	-	1	-
2000.....	46	2	1	-	-	-	-	-	-	-	-	2
1999.....	42	1	-	-	-	1	-	1	-	2	1	-
American Indian 1/, both sexes												
2004.....	3	-	-	-	-	-	-	-	-	-	-	1
2003.....	17	1	3	-	1	2	-	-	-	1	-	-
2002.....	5	-	-	-	-	-	-	-	-	-	-	-
2001.....	3	-	1	1	1	-	-	-	-	-	-	-
2000.....	4	-	-	1	-	-	-	-	-	-	-	-
1999.....	5	1	-	-	1	-	-	-	-	-	-	-
American Indian 1/, male												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	7	1	1	-	-	1	-	-	-	1	-	-
2002.....	3	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	1	-	-	-	-	-	-	-	-	-	-	-
1999.....	2	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, female												
2004.....	3	-	-	-	-	-	-	-	-	-	-	1
2003.....	10	-	2	-	1	1	-	-	-	-	-	-
2002.....	2	-	-	-	-	-	-	-	-	-	-	-
2001.....	3	-	1	1	1	-	-	-	-	-	-	-
2000.....	3	-	-	1	-	-	-	-	-	-	-	-
1999.....	3	-	-	-	1	-	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonia (J12-J18)												
All races, both sexes												
2004.....	58,564	255	101	30	39	63	115	107	192	302	573	789
2003.....	63,371	291	104	47	54	62	132	144	216	366	602	889
2002.....	64,954	256	105	33	46	70	90	132	209	364	599	879
2001.....	61,777	292	106	37	43	62	112	120	209	392	585	802
2000.....	63,548	280	93	44	32	57	119	134	217	397	649	759
1999.....	62,065	307	118	40	42	66	102	133	197	454	593	785
All races, male												
2004.....	26,440	156	39	17	22	41	75	71	121	178	309	463
2003.....	28,079	174	58	24	27	41	71	86	134	203	347	513
2002.....	28,629	157	60	14	22	39	54	66	122	195	347	512
2001.....	27,241	161	55	20	23	33	67	61	124	218	336	492
2000.....	27,994	158	50	26	17	32	64	78	121	206	410	453
1999.....	27,116	165	57	23	25	33	55	76	116	246	337	492
All races, female												
2004.....	32,124	99	62	13	17	22	40	36	71	124	264	326
2003.....	35,292	117	46	23	27	21	61	58	82	163	255	376
2002.....	36,325	99	45	19	24	31	36	66	87	169	252	367
2001.....	34,536	131	51	17	20	29	45	59	85	174	249	310
2000.....	35,554	122	43	18	15	25	55	56	96	191	239	306
1999.....	34,949	142	61	17	17	33	47	57	81	208	256	293
White, both sexes												
2004.....	51,397	149	63	17	29	50	86	80	145	208	413	579
2003.....	55,992	186	73	30	46	46	107	97	151	264	425	645
2002.....	57,660	157	61	25	34	49	69	99	145	255	434	665
2001.....	54,545	161	73	28	32	43	78	81	144	274	400	563
2000.....	56,252	165	60	34	27	40	86	91	159	298	451	543
1999.....	55,120	177	67	30	32	48	63	84	130	323	381	545

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonia (J12-J18)												
White, male												
2004.....	22,960	95	27	9	16	33	63	51	91	123	219	344
2003.....	24,369	113	42	14	24	30	58	64	93	150	240	374
2002.....	25,111	102	34	9	15	31	41	49	86	134	255	380
2001.....	23,656	94	39	15	16	22	53	39	85	150	230	352
2000.....	24,385	93	26	21	14	24	50	48	87	157	288	318
1999.....	23,712	102	27	17	20	25	35	50	79	178	206	339
White, female												
2004.....	28,437	54	36	8	13	17	23	29	54	85	194	235
2003.....	31,623	73	31	16	22	16	49	33	58	114	185	271
2002.....	32,549	55	27	16	19	18	28	50	59	121	179	285
2001.....	30,889	67	34	13	16	21	25	42	59	124	170	211
2000.....	31,867	72	34	13	13	16	36	43	72	141	163	225
1999.....	31,408	75	40	13	12	23	28	34	51	145	175	206
Black, both sexes												
2004.....	5,650	89	32	11	8	13	26	24	41	83	145	187
2003.....	5,775	86	22	14	5	13	22	42	57	93	158	218
2002.....	5,841	86	38	6	11	16	20	27	56	101	151	192
2001.....	5,751	106	26	7	8	15	29	31	56	109	166	221
2000.....	5,909	101	29	7	3	14	29	37	55	85	169	192
1999.....	5,810	111	42	8	7	13	35	46	60	118	189	222
Black, male												
2004.....	2,750	51	10	8	4	8	11	18	25	49	81	105
2003.....	2,869	51	11	9	2	8	13	21	37	47	95	123
2002.....	2,753	47	20	4	6	6	12	13	32	54	84	117
2001.....	2,803	53	14	4	5	10	12	16	35	62	94	130
2000.....	2,880	56	21	4	2	6	14	26	32	42	107	121
1999.....	2,801	55	24	5	3	6	17	25	33	61	115	141

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
-------------	-----------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Pneumonia (J12-J18)

Black, female

2004.....	2,900	38	22	3	4	5	15	6	16	34	64	82
2003.....	2,906	35	11	5	3	5	9	21	20	46	63	95
2002.....	3,088	39	18	2	5	10	8	14	24	47	67	75
2001.....	2,948	53	12	3	3	5	17	15	21	47	72	91
2000.....	3,029	45	8	3	1	8	15	11	23	43	62	71
1999.....	3,009	56	18	3	4	7	18	21	27	57	74	81

American Indian 1/, both sexes

2004.....	288	12	4	1	-	-	-	2	5	4	9	11
2003.....	373	11	2	1	1	1	1	3	3	6	14	18
2002.....	288	6	1	-	-	2	1	2	5	5	8	10
2001.....	315	17	2	-	2	2	1	5	4	6	13	6
2000.....	285	6	-	1	1	1	1	4	1	11	17	8
1999.....	310	8	3	1	1	1	4	1	3	4	16	14

American Indian 1/, male

2004.....	142	7	1	-	-	-	-	1	4	1	6	7
2003.....	177	6	1	1	1	1	-	-	1	3	10	12
2002.....	130	3	1	-	-	-	1	2	2	5	4	7
2001.....	160	11	-	-	1	-	1	5	1	5	8	3
2000.....	148	2	-	-	-	1	-	2	1	6	12	5
1999.....	145	3	3	1	1	1	3	-	1	-	12	8

American Indian 1/, female

2004.....	146	5	3	1	-	-	-	1	1	3	3	4
2003.....	196	5	1	-	-	-	1	3	2	3	4	6
2002.....	158	3	-	-	-	2	-	-	3	-	4	3
2001.....	155	6	2	-	1	2	-	-	3	1	5	3
2000.....	137	4	-	1	1	-	1	2	-	5	5	3
1999.....	165	5	-	-	-	-	1	1	2	4	4	6

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonia (J12-J18)												
Asian or Pacific Islander 2/, both sexes												
2004.....	1,229	5	2	1	2	-	3	1	1	7	6	12
2003.....	1,231	8	7	2	2	2	2	2	5	3	5	8
2002.....	1,165	7	5	2	1	3	-	4	3	3	6	12
2001.....	1,166	8	5	2	1	2	4	3	5	3	6	12
2000.....	1,102	8	4	2	1	2	3	2	2	3	12	16
1999.....	825	11	6	1	2	4	-	2	4	9	7	4
Asian or Pacific Islander 2/, male												
2004.....	588	3	1	-	2	-	1	1	1	5	3	7
2003.....	664	4	4	-	-	2	-	1	3	3	2	4
2002.....	635	5	5	1	1	2	-	2	2	2	4	8
2001.....	622	3	2	1	1	1	1	1	3	1	4	7
2000.....	581	7	3	1	1	1	-	2	1	1	3	9
1999.....	458	5	3	-	1	1	-	1	3	7	4	4
Asian or Pacific Islander 2/, female												
2004.....	641	2	1	1	-	-	2	-	-	2	3	5
2003.....	567	4	3	2	2	-	2	1	2	-	3	4
2002.....	530	2	-	1	-	1	-	2	1	1	2	4
2001.....	544	5	3	1	-	1	3	2	2	2	2	5
2000.....	521	1	1	1	-	1	3	-	1	2	9	7
1999.....	367	6	3	1	1	3	-	1	1	2	3	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Viral pneumonia, not elsewhere classified (J12)

All races, both sexes

2004.....	253	55	20	2	4	-	-	4	4	4	7	5
2003.....	313	63	11	4	3	2	5	2	2	4	3	5
2002.....	293	57	13	1	-	5	-	3	4	2	4	4
2001.....	282	76	12	-	1	-	1	3	3	3	6	9
2000.....	392	55	18	1	2	4	2	2	3	4	1	4
1999.....	407	79	22	5	5	1	4	4	2	10	8	3

All races, male

2004.....	121	36	13	2	4	-	-	3	2	1	4	4
2003.....	143	37	7	1	1	2	2	-	1	2	3	1
2002.....	143	38	9	1	-	3	-	2	2	-	1	1
2001.....	151	42	7	-	-	-	-	1	2	1	1	6
2000.....	161	33	11	1	-	2	1	-	1	2	-	1
1999.....	183	43	9	1	4	-	2	2	2	5	4	1

All races, female

2004.....	132	19	7	-	-	-	-	1	2	3	3	1
2003.....	170	26	4	3	2	-	3	2	1	2	-	4
2002.....	150	19	4	-	-	2	-	1	2	2	3	3
2001.....	131	34	5	-	1	-	1	2	1	2	5	3
2000.....	231	22	7	-	2	2	1	2	2	2	1	3
1999.....	224	36	13	4	1	1	2	2	-	5	4	2

White, both sexes

2004.....	211	33	16	-	4	-	-	2	2	3	5	5
2003.....	275	43	9	3	3	2	5	1	2	4	3	4
2002.....	256	37	8	1	-	5	-	3	3	2	4	4
2001.....	231	42	11	-	1	-	-	2	2	3	6	7
2000.....	341	34	12	1	2	2	1	2	3	3	-	4
1999.....	334	50	11	4	3	-	1	1	2	9	5	2

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Viral pneumonia, not elsewhere classified (J12)

White, male

2004.....	99	26	10	-	4	-	-	1	1	1	3	4
2003.....	123	26	7	-	1	2	2	-	1	2	3	1
2002.....	118	23	5	1	-	3	-	2	2	-	1	1
2001.....	123	23	6	-	-	-	-	1	2	1	1	5
2000.....	133	18	5	1	-	2	1	-	1	1	-	1
1999.....	147	29	4	-	3	-	-	1	2	5	1	-

White, female

2004.....	112	7	6	-	-	-	-	1	1	2	2	1
2003.....	152	17	2	3	2	-	3	1	1	2	-	3
2002.....	138	14	3	-	-	2	-	1	1	2	3	3
2001.....	108	19	5	-	1	-	-	1	-	2	5	2
2000.....	208	16	7	-	2	-	-	2	2	2	-	3
1999.....	187	21	7	4	-	-	1	-	-	4	4	2

Black, both sexes

2004.....	30	15	3	2	-	-	-	1	2	1	2	-
2003.....	31	18	2	1	-	-	-	1	-	-	-	1
2002.....	31	19	5	-	-	-	-	-	1	-	-	-
2001.....	32	23	-	-	-	-	-	1	1	-	-	2
2000.....	38	18	6	-	-	2	1	-	-	1	1	-
1999.....	58	25	10	1	1	1	2	3	-	1	2	1

Black, male

2004.....	16	7	2	2	-	-	-	1	1	-	1	-
2003.....	17	11	-	1	-	-	-	-	-	-	-	-
2002.....	22	14	4	-	-	-	-	-	-	-	-	-
2001.....	18	14	-	-	-	-	-	-	-	-	-	1
2000.....	23	12	6	-	-	-	-	-	-	1	-	-
1999.....	29	11	5	1	1	-	1	1	-	-	2	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Viral pneumonia, not elsewhere classified (J12)												
Asian or Pacific Islander 2/, both sexes												
2004.....	5	2	-	-	-	-	-	-	-	-	-	-
2003.....	5	-	-	-	-	-	-	-	-	-	-	-
2002.....	4	1	-	-	-	-	-	-	-	-	-	-
2001.....	11	3	1	-	-	-	1	-	-	-	-	-
2000.....	8	2	-	-	-	-	-	-	-	-	-	-
1999.....	8	1	1	-	1	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, male												
2004.....	2	1	-	-	-	-	-	-	-	-	-	-
2003.....	3	-	-	-	-	-	-	-	-	-	-	-
2002.....	2	1	-	-	-	-	-	-	-	-	-	-
2001.....	6	1	1	-	-	-	-	-	-	-	-	-
2000.....	4	2	-	-	-	-	-	-	-	-	-	-
1999.....	4	1	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, female												
2004.....	3	1	-	-	-	-	-	-	-	-	-	-
2003.....	2	-	-	-	-	-	-	-	-	-	-	-
2002.....	2	-	-	-	-	-	-	-	-	-	-	-
2001.....	5	2	-	-	-	-	1	-	-	-	-	-
2000.....	4	-	-	-	-	-	-	-	-	-	-	-
1999.....	4	-	1	-	1	-	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Bacterial pneumonia (J13-J15)

All races, both sexes

2004.....	4,376	31	17	7	6	6	13	17	24	45	78	115
2003.....	4,770	42	26	15	14	11	10	17	25	52	97	117
2002.....	4,650	43	17	4	9	5	10	23	29	48	98	125
2001.....	4,421	32	15	4	2	6	7	17	27	52	79	108
2000.....	4,486	31	12	10	2	3	14	19	28	68	106	115
1999.....	3,872	51	20	6	6	7	11	16	19	78	102	102

All races, male

2004.....	2,206	17	4	2	2	2	8	11	18	19	37	66
2003.....	2,416	25	19	7	6	6	5	12	17	30	58	62
2002.....	2,282	25	8	2	4	3	6	12	14	25	50	67
2001.....	2,234	18	7	2	1	5	3	9	17	27	51	59
2000.....	2,326	22	7	7	1	3	6	7	16	36	64	63
1999.....	1,873	24	11	4	2	3	6	11	8	35	51	59

All races, female

2004.....	2,170	14	13	5	4	4	5	6	6	26	41	49
2003.....	2,354	17	7	8	8	5	5	5	8	22	39	55
2002.....	2,368	18	9	2	5	2	4	11	15	23	48	58
2001.....	2,187	14	8	2	1	1	4	8	10	25	28	49
2000.....	2,160	9	5	3	1	-	8	12	12	32	42	52
1999.....	1,999	27	9	2	4	4	5	5	11	43	51	43

White, both sexes

2004.....	3,725	23	13	6	6	5	11	14	18	35	57	88
2003.....	4,100	26	18	10	12	9	10	10	20	39	67	85
2002.....	4,034	29	8	3	5	2	7	20	18	33	67	92
2001.....	3,830	24	8	3	2	4	6	12	19	39	52	85
2000.....	3,857	15	9	8	2	2	12	14	24	53	81	83
1999.....	3,326	28	11	6	5	5	7	12	11	55	65	74

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bacterial pneumonia (J13-J15)												
White, male												
2004.....	1,871	14	3	2	2	2	7	9	13	16	27	53
2003.....	2,050	16	12	4	5	6	5	7	14	23	35	44
2002.....	1,959	19	4	1	2	2	3	10	10	15	37	48
2001.....	1,900	14	5	1	1	3	3	7	12	18	33	45
2000.....	1,983	10	4	6	1	2	6	3	13	28	55	46
1999.....	1,586	15	4	4	2	3	4	8	6	22	33	41
White, female												
2004.....	1,854	9	10	4	4	3	4	5	5	19	30	35
2003.....	2,050	10	6	6	7	3	5	3	6	16	32	41
2002.....	2,075	10	4	2	3	-	4	10	8	18	30	44
2001.....	1,930	10	3	2	1	1	3	5	7	21	19	40
2000.....	1,874	5	5	2	1	-	6	11	11	25	26	37
1999.....	1,740	13	7	2	3	2	3	4	5	33	32	33
Black, both sexes												
2004.....	519	6	3	-	-	1	2	3	6	9	17	24
2003.....	505	10	5	3	2	2	-	7	3	10	24	27
2002.....	457	12	8	-	4	3	2	2	9	13	24	28
2001.....	461	6	6	1	-	2	1	2	7	12	20	18
2000.....	481	14	2	2	-	1	2	4	3	12	17	24
1999.....	424	17	5	-	1	1	2	4	7	17	29	24
Black, male												
2004.....	265	2	1	-	-	-	1	2	5	3	7	11
2003.....	269	6	4	3	1	-	-	5	3	5	19	15
2002.....	227	5	3	-	2	1	2	1	3	9	9	15
2001.....	260	2	2	1	-	2	-	-	4	9	14	13
2000.....	260	11	2	1	-	1	-	3	2	7	8	11
1999.....	221	7	4	-	-	-	1	3	2	9	14	15

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bacterial pneumonia (J13-J15)												
Black, female												
2004.....	254	4	2	-	-	1	1	1	1	6	10	13
2003.....	236	4	1	-	1	2	-	2	-	5	5	12
2002.....	230	7	5	-	2	2	-	1	6	4	15	13
2001.....	201	4	4	-	-	-	1	2	3	3	6	5
2000.....	221	3	-	1	-	-	2	1	1	5	9	13
1999.....	203	10	1	-	1	1	1	1	5	8	15	9
American Indian 1/, both sexes												
2004.....	40	2	1	1	-	-	-	-	-	1	3	2
2003.....	49	3	-	-	-	-	-	-	1	3	4	2
2002.....	37	2	-	-	-	-	1	-	2	1	4	3
2001.....	25	2	-	-	-	-	-	2	-	-	5	1
2000.....	37	1	-	-	-	-	-	1	-	2	2	4
1999.....	42	2	1	-	-	-	2	-	-	2	6	4
American Indian 1/, male												
2004.....	19	1	-	-	-	-	-	-	-	-	3	1
2003.....	28	1	-	-	-	-	-	-	-	2	2	1
2002.....	20	1	-	-	-	-	1	-	1	1	2	2
2001.....	15	2	-	-	-	-	-	2	-	-	2	-
2000.....	21	-	-	-	-	-	-	1	-	1	-	4
1999.....	18	-	1	-	-	-	1	-	-	-	3	3
American Indian 1/, female												
2004.....	21	1	1	1	-	-	-	-	-	1	-	1
2003.....	21	2	-	-	-	-	-	-	1	1	2	1
2002.....	17	1	-	-	-	-	-	-	1	-	2	1
2001.....	10	-	-	-	-	-	-	-	-	-	3	1
2000.....	16	1	-	-	-	-	-	-	-	1	2	-
1999.....	24	2	-	-	-	-	1	-	-	2	3	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bacterial pneumonia (J13-J15)												
Asian or Pacific Islander 2/, both sexes												
2004.....	92	-	-	-	-	-	-	-	-	-	1	1
2003.....	116	3	3	2	-	-	-	-	1	-	2	3
2002.....	122	-	1	1	-	-	-	1	-	1	3	2
2001.....	105	-	1	-	-	-	-	1	1	1	2	4
2000.....	111	1	1	-	-	-	-	-	1	1	6	4
1999.....	80	4	3	-	-	1	-	-	1	4	2	-
Asian or Pacific Islander 2/, male												
2004.....	51	-	-	-	-	-	-	-	-	-	-	1
2003.....	69	2	3	-	-	-	-	-	-	-	2	2
2002.....	76	-	1	1	-	-	-	1	-	-	2	2
2001.....	59	-	-	-	-	-	-	-	1	-	2	1
2000.....	62	1	1	-	-	-	-	-	1	-	1	2
1999.....	48	2	2	-	-	-	-	-	-	4	1	-
Asian or Pacific Islander 2/, female												
2004.....	41	-	-	-	-	-	-	-	-	-	1	-
2003.....	47	1	-	2	-	-	-	-	1	-	-	1
2002.....	46	-	-	-	-	-	-	-	-	1	1	-
2001.....	46	-	1	-	-	-	-	1	-	1	-	3
2000.....	49	-	-	-	-	-	-	-	-	1	5	2
1999.....	32	2	1	-	-	1	-	-	1	-	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonia due to other or unspecified organisms (J16,J18)												
All races, both sexes												
2004.....	53,935	169	64	21	29	57	102	86	164	253	488	669
2003.....	58,288	186	67	28	37	49	117	125	189	310	502	767
2002.....	60,011	156	75	28	37	60	80	106	176	314	497	750
2001.....	57,074	184	79	33	40	56	104	100	179	337	500	685
2000.....	58,670	194	63	33	28	50	103	113	186	325	542	640
1999.....	57,786	177	76	29	31	58	87	113	176	366	483	680
All races, male												
2004.....	24,113	103	22	13	16	39	67	57	101	158	268	393
2003.....	25,520	112	32	16	20	33	64	74	116	171	286	450
2002.....	26,204	94	43	11	18	33	48	52	106	170	296	444
2001.....	24,856	101	41	18	22	28	64	51	105	190	284	427
2000.....	25,507	103	32	18	16	27	57	71	104	168	346	389
1999.....	25,060	98	37	18	19	30	47	63	106	206	282	432
All races, female												
2004.....	29,822	66	42	8	13	18	35	29	63	95	220	276
2003.....	32,768	74	35	12	17	16	53	51	73	139	216	317
2002.....	33,807	62	32	17	19	27	32	54	70	144	201	306
2001.....	32,218	83	38	15	18	28	40	49	74	147	216	258
2000.....	33,163	91	31	15	12	23	46	42	82	157	196	251
1999.....	32,726	79	39	11	12	28	40	50	70	160	201	248
White, both sexes												
2004.....	47,461	93	34	11	19	45	75	64	125	170	351	486
2003.....	51,617	117	46	17	31	35	92	86	129	221	355	556
2002.....	53,370	91	45	21	29	42	62	76	124	220	363	569
2001.....	50,484	95	54	25	29	39	72	67	123	232	342	471
2000.....	52,054	116	39	25	23	36	73	75	132	242	370	456
1999.....	51,460	99	45	20	24	43	55	71	117	259	311	469

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonia due to other or unspecified organisms (J16,J18)												
White, male												
2004.....	20,990	55	14	7	10	31	56	41	77	106	189	287
2003.....	22,196	71	23	10	18	22	51	57	78	125	202	329
2002.....	23,034	60	25	7	13	26	38	37	74	119	217	331
2001.....	21,633	57	28	14	15	19	50	31	71	131	196	302
2000.....	22,269	65	17	14	13	20	43	45	73	128	233	271
1999.....	21,979	58	19	13	15	22	31	41	71	151	172	298
White, female												
2004.....	26,471	38	20	4	9	14	19	23	48	64	162	199
2003.....	29,421	46	23	7	13	13	41	29	51	96	153	227
2002.....	30,336	31	20	14	16	16	24	39	50	101	146	238
2001.....	28,851	38	26	11	14	20	22	36	52	101	146	169
2000.....	29,785	51	22	11	10	16	30	30	59	114	137	185
1999.....	29,481	41	26	7	9	21	24	30	46	108	139	171
Black, both sexes												
2004.....	5,101	68	26	9	8	12	24	20	33	73	126	163
2003.....	5,239	58	15	10	3	11	22	34	54	83	134	190
2002.....	5,353	55	25	6	7	13	18	25	46	88	127	164
2001.....	5,258	77	20	6	8	13	28	28	48	97	146	201
2000.....	5,390	69	21	5	3	11	26	33	52	72	151	168
1999.....	5,328	69	27	7	5	11	31	39	53	100	158	197
Black, male												
2004.....	2,469	42	7	6	4	8	10	15	19	46	73	94
2003.....	2,583	34	7	5	1	8	13	16	34	42	76	108
2002.....	2,504	28	13	4	4	5	10	12	29	45	75	102
2001.....	2,525	37	12	3	5	8	12	16	31	53	80	116
2000.....	2,597	33	13	3	2	5	14	23	30	34	99	110
1999.....	2,551	37	15	4	2	6	15	21	31	52	99	125

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Pneumonia due to other or unspecified organisms (J16,J18)

Black, female

2004.....	2,632	26	19	3	4	4	14	5	14	27	53	69
2003.....	2,656	24	8	5	2	3	9	18	20	41	58	82
2002.....	2,849	27	12	2	3	8	8	13	17	43	52	62
2001.....	2,733	40	8	3	3	5	16	12	17	44	66	85
2000.....	2,793	36	8	2	1	6	12	10	22	38	52	58
1999.....	2,777	32	12	3	3	5	16	18	22	48	59	72

American Indian 1/, both sexes

2004.....	241	5	2	-	-	-	-	1	5	3	6	9
2003.....	322	6	2	1	1	1	1	3	2	3	10	16
2002.....	249	4	1	-	-	2	-	2	3	4	4	7
2001.....	282	7	2	-	2	2	1	3	4	6	8	5
2000.....	243	4	-	1	1	1	1	3	1	9	15	4
1999.....	261	3	2	1	1	1	1	1	3	2	10	10

American Indian 1/, male

2004.....	119	4	-	-	-	-	-	-	4	1	3	6
2003.....	149	5	1	1	1	1	-	-	1	1	8	11
2002.....	109	2	1	-	-	-	-	2	1	4	2	5
2001.....	141	5	-	-	1	-	1	3	1	5	6	3
2000.....	126	1	-	-	-	1	-	1	1	5	12	1
1999.....	124	1	2	1	1	1	1	-	1	-	9	5

American Indian 1/, female

2004.....	122	1	2	-	-	-	-	1	1	2	3	3
2003.....	173	1	1	-	-	-	1	3	1	2	2	5
2002.....	140	2	-	-	-	2	-	-	2	-	2	2
2001.....	141	2	2	-	1	2	-	-	3	1	2	2
2000.....	117	3	-	1	-	1	1	2	-	4	3	3
1999.....	137	2	-	-	-	-	-	1	2	2	1	5

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonia due to other or unspecified organisms (J16,J18)												
Asian or Pacific Islander 2/, both sexes												
2004.....	1,132	3	2	1	2	-	3	1	1	7	5	11
2003.....	1,110	5	4	-	2	2	2	2	4	3	3	5
2002.....	1,039	6	4	1	1	3	-	3	3	2	3	10
2001.....	1,050	5	3	2	1	2	3	2	4	2	4	8
2000.....	983	5	3	2	1	2	3	2	1	2	6	12
1999.....	737	6	2	1	1	3	-	2	3	5	4	4
Asian or Pacific Islander 2/, male												
2004.....	535	2	1	-	2	-	1	1	1	5	3	6
2003.....	592	2	1	-	-	2	-	1	3	3	-	2
2002.....	557	4	4	-	1	2	-	1	2	2	2	6
2001.....	557	2	1	1	1	1	1	1	2	1	2	6
2000.....	515	4	2	1	1	1	-	2	-	1	2	7
1999.....	406	2	1	-	1	1	-	1	3	3	2	4
Asian or Pacific Islander 2/, female												
2004.....	597	1	1	1	-	-	2	-	-	2	2	5
2003.....	518	3	3	-	2	-	2	1	1	-	3	3
2002.....	482	2	-	1	-	1	-	2	1	-	1	4
2001.....	493	3	2	1	-	1	2	1	2	1	2	2
2000.....	468	1	1	1	-	1	3	-	1	1	4	5
1999.....	331	4	1	1	-	2	-	1	-	2	2	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other acute lower respiratory infections (J20-J22)

All races, both sexes

2004.....	319	60	16	1	-	3	1	-	3	4	5	5
2003.....	421	54	28	3	3	1	5	-	2	4	7	7
2002.....	386	52	23	7	1	1	2	3	2	2	6	6
2001.....	347	50	18	3	-	-	1	1	1	7	5	4
2000.....	425	43	11	3	4	4	1	5	3	3	9	8
1999.....	540	63	25	2	3	3	2	2	2	6	5	7

All races, male

2004.....	137	34	9	1	-	2	1	-	2	3	3	3
2003.....	170	35	17	2	-	-	2	-	1	3	3	2
2002.....	154	39	9	7	-	-	-	1	-	1	2	1
2001.....	117	26	8	2	-	-	1	-	-	4	3	1
2000.....	160	25	4	2	1	4	-	5	1	1	6	2
1999.....	196	36	15	-	2	1	-	2	1	3	2	5

All races, female

2004.....	182	26	7	-	-	1	-	-	1	1	2	2
2003.....	251	19	11	1	3	1	3	-	1	1	4	5
2002.....	232	13	14	-	1	1	2	2	2	1	4	5
2001.....	230	24	10	1	-	-	-	1	1	3	2	3
2000.....	265	18	7	1	3	-	1	-	2	2	3	6
1999.....	344	27	10	2	1	2	2	-	1	3	3	2

White, both sexes

2004.....	279	43	9	1	-	2	1	-	3	4	5	5
2003.....	375	42	17	3	2	1	2	-	1	4	7	6
2002.....	334	34	20	7	1	1	1	2	2	2	5	4
2001.....	299	31	13	3	-	-	1	-	1	7	5	4
2000.....	372	25	8	2	1	4	1	4	1	2	6	5
1999.....	462	36	15	1	3	2	2	1	2	6	3	6

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other acute lower respiratory infections (J20-J22)

White, male

2004.....	115	24	4	1	-	1	1	-	2	3	3	3
2003.....	147	27	10	2	-	-	-	-	1	3	3	2
2002.....	130	26	7	7	-	-	-	-	-	1	2	1
2001.....	94	16	5	2	-	-	1	-	-	4	3	1
2000.....	134	12	4	2	-	4	-	4	-	1	5	2
1999.....	156	19	8	-	2	1	-	1	1	3	1	5

White, female

2004.....	164	19	5	-	-	1	-	-	1	1	2	2
2003.....	228	15	7	1	2	1	2	-	-	1	4	4
2002.....	204	8	13	-	1	1	1	2	2	1	3	3
2001.....	205	15	8	1	-	-	-	-	1	3	2	3
2000.....	238	13	4	-	1	-	1	-	1	1	1	3
1999.....	306	17	7	1	1	1	2	-	1	3	2	1

Black, both sexes

2004.....	31	13	6	-	-	1	-	-	-	-	-	-
2003.....	40	12	7	-	1	-	3	-	1	-	-	1
2002.....	45	16	2	-	-	-	1	1	-	-	1	2
2001.....	34	15	2	-	-	-	-	1	-	-	-	-
2000.....	46	14	3	-	3	-	-	1	1	1	3	3
1999.....	64	24	6	-	-	1	-	1	-	-	2	1

Black, male

2004.....	15	6	4	-	-	1	-	-	-	-	-	-
2003.....	21	8	6	-	-	-	2	-	-	-	-	-
2002.....	23	13	1	-	-	-	-	1	-	-	-	-
2001.....	16	9	1	-	-	-	-	-	-	-	-	-
2000.....	22	10	-	-	1	-	-	1	1	-	1	-
1999.....	30	15	3	-	-	-	-	1	-	-	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Acute bronchitis and bronchiolitis (J20-J21)

All races, both sexes

2004.....	226	58	16	1	-	3	1	-	3	3	5	3
2003.....	304	53	27	3	3	1	5	-	2	4	5	7
2002.....	279	49	23	7	1	-	2	3	2	2	4	6
2001.....	263	49	18	3	-	-	1	1	1	6	5	4
2000.....	290	42	11	2	4	3	1	3	3	2	8	8
1999.....	388	62	24	2	2	3	2	1	2	6	4	7

All races, male

2004.....	108	33	9	1	-	2	1	-	2	2	3	1
2003.....	126	34	16	2	-	-	2	-	1	3	2	2
2002.....	122	37	9	7	-	-	-	1	-	1	2	1
2001.....	89	25	8	2	-	-	1	-	-	3	3	1
2000.....	119	24	4	1	1	3	-	3	1	1	5	2
1999.....	154	36	14	-	2	1	-	1	1	3	1	5

All races, female

2004.....	118	25	7	-	-	1	-	-	1	1	2	2
2003.....	178	19	11	1	3	1	3	-	1	1	3	5
2002.....	157	12	14	-	1	-	2	2	2	1	2	5
2001.....	174	24	10	1	-	-	-	1	1	3	2	3
2000.....	171	18	7	1	3	-	1	-	2	1	3	6
1999.....	234	26	10	2	-	2	2	-	1	3	3	2

White, both sexes

2004.....	190	42	9	1	-	2	1	-	3	3	5	3
2003.....	263	41	16	3	2	1	2	-	1	4	5	6
2002.....	237	31	20	7	1	-	1	2	2	2	4	4
2001.....	218	30	13	3	-	-	1	-	1	6	5	4
2000.....	244	25	8	1	244	3	1	2	1	1	5	5
1999.....	320	35	15	1	2	2	2	1	2	6	2	6

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Unspecified acute lower respiratory infection (J22)												
All races, both sexes												
2004.....	93	2	-	-	-	-	-	-	-	1	-	2
2003.....	117	1	1	-	-	-	-	-	-	-	2	-
2002.....	107	3	-	-	-	1	-	-	-	-	2	-
2001.....	84	1	-	-	-	-	-	-	-	1	-	-
2000.....	135	1	-	1	-	1	-	2	-	1	1	-
1999.....	152	1	1	-	1	-	-	1	-	-	1	-
All races, male												
2004.....	29	1	-	-	-	-	-	-	-	1	-	2
2003.....	44	1	1	-	-	-	-	-	-	-	1	-
2002.....	32	2	-	-	-	-	-	-	-	-	-	-
2001.....	28	1	-	-	-	-	-	-	-	1	-	-
2000.....	41	1	-	1	-	1	-	2	-	-	1	-
1999.....	42	-	1	-	-	-	-	1	-	-	1	-
All races, female												
2004.....	64	1	-	-	-	-	-	-	-	-	-	-
2003.....	73	-	-	-	-	-	-	-	-	-	1	-
2002.....	75	1	-	-	-	1	-	-	-	-	2	-
2001.....	56	-	-	-	-	-	-	-	-	-	-	-
2000.....	94	-	-	-	-	-	-	-	-	1	-	-
1999.....	110	1	-	-	1	-	-	-	-	-	-	-
White, both sexes												
2004.....	89	1	-	-	-	-	-	-	-	1	-	2
2003.....	112	1	1	-	-	-	-	-	-	-	2	-
2002.....	97	3	-	-	-	1	-	-	-	-	1	-
2001.....	81	1	-	-	-	-	-	-	-	1	-	-
2000.....	128	-	-	1	-	1	-	2	-	1	1	-
1999.....	142	1	-	-	1	-	-	-	-	-	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Unspecified acute lower respiratory infection (J22)

White, male

2004.....	28	1	-	-	-	-	-	-	-	1	-	2
2003.....	43	1	1	-	-	-	-	-	-	-	1	-
2002.....	30	2	-	-	-	-	-	-	-	-	-	-
2001.....	27	1	-	-	-	-	-	-	-	1	-	-
2000.....	36	-	-	1	-	1	-	2	-	-	1	-
1999.....	37	-	-	-	-	-	-	-	-	-	1	-

White, female

2004.....	61	-	-	-	-	-	-	-	-	-	-	-
2003.....	69	-	-	-	-	-	-	-	-	-	1	-
2002.....	67	1	-	-	-	1	-	-	-	-	1	-
2001.....	54	-	-	-	-	-	-	-	-	-	-	-
2000.....	92	-	-	-	-	-	-	-	-	1	-	-
1999.....	105	1	-	-	1	-	-	-	-	-	-	-

Black, both sexes

2004.....	3	1	-	-	-	-	-	-	-	-	-	-
2003.....	5	-	-	-	-	-	-	-	-	-	-	-
2002.....	9	-	-	-	-	-	-	-	-	-	1	-
2001.....	2	-	-	-	-	-	-	-	-	-	-	-
2000.....	6	1	-	-	-	-	-	-	-	-	-	-
1999.....	9	-	1	-	-	-	-	1	-	-	-	-

Black, male

2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	1	-	-	-	-	-	-	-	-	-	-	-
2002.....	2	-	-	-	-	-	-	-	-	-	-	-
2001.....	1	-	-	-	-	-	-	-	-	-	-	-
2000.....	4	1	-	-	-	-	-	-	-	-	-	-
1999.....	4	-	1	-	-	-	-	1	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	-------------	-----------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Other diseases of upper respiratory tract (J30-J39)

All races, both sexes

2004.....	426	11	16	2	1	7	5	6	10	10	10	25
2003.....	438	9	5	4	4	8	9	9	5	6	11	19
2002.....	401	15	8	7	9	4	5	3	7	6	5	24
2001.....	454	15	11	4	5	9	5	7	8	12	14	22
2000.....	446	13	13	1	3	4	1	11	6	10	15	22
1999.....	449	14	9	1	6	5	6	6	3	12	9	18

All races, male

2004.....	224	6	6	1	-	5	5	5	7	4	4	15
2003.....	234	3	5	3	3	4	4	7	2	3	7	12
2002.....	187	5	5	6	3	4	4	1	5	4	2	13
2001.....	225	9	6	1	2	7	3	5	5	8	9	12
2000.....	222	6	6	-	1	4	1	6	3	7	9	14
1999.....	234	7	6	-	2	4	4	3	2	10	6	10

All races, female

2004.....	202	5	10	1	1	2	-	1	3	6	6	10
2003.....	204	6	-	1	1	4	5	2	3	3	4	7
2002.....	214	10	3	1	6	-	1	2	2	2	3	11
2001.....	229	6	5	3	3	2	2	2	3	4	5	10
2000.....	224	7	7	1	2	-	-	5	3	3	6	8
1999.....	215	7	3	1	4	1	2	3	1	2	3	8

White, both sexes

2004.....	348	9	8	1	1	6	3	4	7	6	7	20
2003.....	368	8	2	4	1	6	4	8	3	6	8	12
2002.....	335	9	5	6	5	2	3	3	7	6	2	18
2001.....	354	9	6	1	2	7	3	3	6	6	10	10
2000.....	371	9	10	1	1	1	-	7	6	7	12	17
1999.....	368	12	6	1	5	4	2	4	2	7	3	13

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of upper respiratory tract (J30-J39)												
White, male												
2004.....	181	5	2	1	-	4	3	3	4	2	3	11
2003.....	201	3	2	3	1	2	2	6	2	3	6	7
2002.....	152	4	3	6	-	2	2	1	5	4	1	7
2001.....	170	5	3	1	-	7	2	2	3	5	6	6
2000.....	183	4	3	-	1	1	-	3	3	5	7	11
1999.....	193	6	3	-	2	3	1	3	1	6	3	9
White, female												
2004.....	167	4	6	-	1	2	-	1	3	4	4	9
2003.....	167	5	-	1	-	4	2	2	1	3	2	5
2002.....	183	5	2	-	5	-	1	2	2	2	1	11
2001.....	184	4	3	-	2	-	1	1	3	1	4	4
2000.....	188	5	7	1	-	-	-	4	3	2	5	6
1999.....	175	6	3	1	3	1	1	1	1	1	-	4
Black, both sexes												
2004.....	70	2	7	1	-	1	2	2	3	4	3	3
2003.....	60	1	3	-	3	1	4	1	2	-	2	7
2002.....	56	5	3	1	3	2	2	-	-	-	3	4
2001.....	87	4	5	2	3	2	2	3	2	4	4	11
2000.....	63	4	3	-	1	2	1	4	-	3	2	4
1999.....	70	1	2	-	1	1	4	2	1	5	5	4
Black, male												
2004.....	37	1	4	-	-	1	2	2	3	2	1	2
2003.....	25	-	3	-	2	1	1	1	-	-	-	5
2002.....	32	1	2	-	2	2	2	-	-	-	1	4
2001.....	44	2	3	-	2	-	1	2	2	1	3	5
2000.....	32	2	3	-	-	2	1	3	-	2	2	3
1999.....	34	-	2	-	-	1	3	-	1	4	2	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic lower respiratory diseases (J40-J47)												
All races, both sexes												
2004.....	121,987	35	48	46	74	85	94	112	143	277	610	1,232
2003.....	126,382	31	55	37	81	84	107	118	164	301	649	1,276
2002.....	124,816	39	65	41	95	93	99	116	185	340	668	1,226
2001.....	123,013	41	43	42	62	74	97	102	188	355	614	1,127
2000.....	122,009	36	51	48	91	86	104	125	171	310	620	1,174
1999.....	124,181	36	54	49	90	107	102	145	185	302	588	1,079
All races, male												
2004.....	58,646	24	30	30	45	55	55	63	88	132	299	613
2003.....	60,714	23	38	23	52	50	64	67	89	142	287	681
2002.....	60,713	23	44	24	50	68	52	70	95	159	338	617
2001.....	59,697	22	29	24	37	43	56	55	91	171	284	591
2000.....	60,004	23	28	26	54	58	62	72	93	145	292	617
1999.....	62,415	19	30	29	64	60	63	69	83	142	268	547
All races, female												
2004.....	63,341	11	18	16	29	30	39	49	55	145	311	619
2003.....	65,668	8	17	14	29	34	43	51	75	159	362	595
2002.....	64,103	16	21	17	45	25	47	46	90	181	330	609
2001.....	63,316	19	14	18	25	31	41	47	97	184	330	536
2000.....	62,005	13	23	22	37	28	42	53	78	165	328	557
1999.....	61,766	17	24	20	26	47	39	76	102	160	320	532
White, both sexes												
2004.....	112,914	23	22	17	36	39	52	62	88	188	456	946
2003.....	116,917	23	24	14	40	38	63	70	99	211	464	1,012
2002.....	115,395	25	30	17	38	47	62	68	121	226	470	970
2001.....	113,819	22	20	18	30	40	45	53	103	212	453	882
2000.....	112,840	23	28	23	40	40	63	67	99	194	430	882
1999.....	114,735	16	31	29	42	68	57	73	103	200	408	801

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	-------------	-----------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Chronic lower respiratory diseases (J40-J47)

White, male

2004.....	53,622	13	15	8	22	25	30	29	53	87	225	477
2003.....	55,397	20	14	7	24	20	39	34	45	100	205	549
2002.....	55,409	16	21	12	19	31	29	37	64	96	240	489
2001.....	54,561	13	12	10	18	19	26	29	45	109	207	480
2000.....	54,816	16	13	12	21	26	39	36	48	85	207	472
1999.....	57,000	10	14	15	30	36	32	32	46	90	181	410

White, female

2004.....	59,292	10	7	9	14	14	22	33	35	101	231	469
2003.....	61,520	3	10	7	16	18	24	36	54	111	259	463
2002.....	59,986	9	9	5	19	16	33	31	57	130	230	481
2001.....	59,258	9	8	8	12	21	19	24	58	103	246	402
2000.....	58,024	7	15	11	19	14	24	31	51	109	223	410
1999.....	57,735	6	17	14	12	32	25	41	57	110	227	391

Black, both sexes

2004.....	7,400	11	22	27	37	42	37	47	54	83	142	258
2003.....	7,709	8	26	23	40	41	41	45	59	81	166	241
2002.....	7,831	14	30	21	55	43	36	43	61	110	188	233
2001.....	7,589	17	23	23	31	30	47	47	82	132	147	224
2000.....	7,607	11	21	24	46	44	37	51	65	104	171	259
1999.....	7,915	18	18	20	48	36	42	68	76	95	169	257

Black, male

2004.....	4,046	10	14	21	23	27	21	32	34	41	67	124
2003.....	4,247	3	21	16	27	26	23	31	41	38	70	120
2002.....	4,341	7	20	11	30	36	23	28	30	61	94	117
2001.....	4,187	7	17	14	18	22	26	26	44	59	69	102
2000.....	4,238	6	14	14	29	31	20	31	41	52	78	125
1999.....	4,500	9	13	14	34	23	29	34	35	46	80	126

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	-------------	-----------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Chronic lower respiratory diseases (J40-J47)

Black, female

2004.....	3,354	1	8	6	14	15	16	15	20	42	75	134
2003.....	3,462	5	5	7	13	15	18	14	18	43	96	121
2002.....	3,490	7	10	10	25	7	13	15	31	49	94	116
2001.....	3,402	10	6	9	13	8	21	21	38	73	78	122
2000.....	3,369	5	7	10	17	13	17	20	24	52	93	134
1999.....	3,415	9	5	6	14	13	13	34	41	49	89	131

American Indian 1/, both sexes

2004.....	486	1	1	1	-	1	2	2	-	3	5	11
2003.....	512	-	1	-	-	1	1	-	3	5	9	12
2002.....	452	-	1	1	-	1	1	1	-	2	4	10
2001.....	427	1	-	1	-	2	1	1	1	5	5	8
2000.....	429	1	2	-	-	1	2	4	-	4	6	10
1999.....	406	1	-	-	-	-	1	-	2	2	7	8

American Indian 1/, male

2004.....	235	1	-	-	-	1	1	1	-	2	3	4
2003.....	269	-	-	-	-	1	1	-	1	2	6	6
2002.....	220	-	1	-	-	-	-	1	-	2	-	5
2001.....	200	1	-	-	-	1	1	-	1	-	3	3
2000.....	228	1	1	-	-	1	2	3	-	1	3	7
1999.....	197	-	-	-	-	-	-	-	1	2	3	5

American Indian 1/, female

2004.....	251	-	1	1	-	-	1	1	-	1	2	7
2003.....	243	-	1	-	-	-	-	-	2	3	3	6
2002.....	232	-	-	1	-	1	1	-	-	-	4	5
2001.....	227	-	-	1	-	1	-	1	-	5	2	5
2000.....	201	-	1	-	-	-	-	1	-	3	3	3
1999.....	209	1	-	-	-	-	1	-	1	-	4	3

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Chronic lower respiratory diseases (J40-J47)

Asian or Pacific Islander 2/, both sexes

2004.....	1,187	-	3	1	1	3	3	1	1	3	7	17
2003.....	1,244	-	4	-	1	4	2	3	3	4	10	11
2002.....	1,138	-	4	2	2	2	-	4	3	2	6	13
2001.....	1,178	1	-	-	1	2	4	1	2	6	9	13
2000.....	1,133	1	-	1	5	1	2	3	7	8	13	23
1999.....	1,125	1	5	-	-	3	2	4	4	5	4	13

Asian or Pacific Islander 2/, male

2004.....	743	-	1	1	-	2	3	1	1	2	4	8
2003.....	801	-	3	-	1	3	1	2	2	2	6	6
2002.....	743	-	2	1	1	1	-	4	1	-	4	6
2001.....	749	1	-	-	1	1	3	-	1	3	5	6
2000.....	722	-	-	-	4	-	1	2	4	7	4	13
1999.....	718	-	3	-	-	1	2	3	1	4	4	6

Asian or Pacific Islander 2/, female

2004.....	444	-	2	-	1	1	-	-	-	1	3	9
2003.....	443	-	1	-	-	1	1	1	1	2	4	5
2002.....	395	-	2	1	1	1	-	-	2	2	2	7
2001.....	429	-	-	-	-	1	1	1	1	3	4	7
2000.....	411	1	-	1	1	1	1	1	3	1	9	10
1999.....	407	1	2	-	-	2	-	1	3	1	-	7

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)												
All races, both sexes												
2004.....	121,029	35	48	46	72	84	93	112	140	271	604	1,224
2003.....	125,448	31	55	37	81	81	107	116	160	297	638	1,263
2002.....	123,838	39	65	41	95	93	97	115	180	336	662	1,217
2001.....	122,042	40	43	42	62	72	96	99	184	349	610	1,119
2000.....	121,058	36	51	46	91	84	103	122	169	306	610	1,165
1999.....	123,211	36	53	49	89	107	100	137	183	301	582	1,066
All races, male												
2004.....	58,375	24	30	30	45	54	54	63	86	130	296	608
2003.....	60,451	23	38	23	52	47	64	66	87	140	281	677
2002.....	60,440	23	44	24	50	68	51	69	93	157	336	614
2001.....	59,432	22	29	24	37	42	55	53	91	169	284	586
2000.....	59,738	23	28	24	54	57	62	70	93	144	288	609
1999.....	62,089	19	30	29	63	60	61	66	83	141	265	542
All races, female												
2004.....	62,654	11	18	16	27	30	39	49	54	141	308	616
2003.....	64,997	8	17	14	29	34	43	50	73	157	357	586
2002.....	63,398	16	21	17	45	25	46	46	87	179	326	603
2001.....	62,610	18	14	18	25	30	41	46	93	180	326	533
2000.....	61,320	13	23	22	37	27	41	52	76	162	322	556
1999.....	61,122	17	23	20	26	47	39	71	100	160	317	524
White, both sexes												
2004.....	112,015	23	22	17	36	38	51	62	85	183	451	941
2003.....	116,063	23	24	14	40	36	63	70	95	208	460	1,002
2002.....	114,485	25	30	17	38	47	60	67	117	222	465	966
2001.....	112,927	21	20	18	30	39	45	51	101	208	450	877
2000.....	111,964	23	28	21	40	39	62	64	98	191	423	875
1999.....	113,845	16	31	29	41	68	55	68	103	199	403	792

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	-------------	-----------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)

White, male

2004.....	53,368	13	15	8	22	24	29	29	51	86	222	474
2003.....	55,164	20	14	7	24	18	39	34	43	99	204	545
2002.....	55,164	16	21	12	19	31	28	36	63	94	238	487
2001.....	54,337	13	12	10	18	18	26	27	45	107	207	476
2000.....	54,575	16	13	10	21	26	39	34	48	84	203	465
1999.....	56,706	10	14	15	29	36	30	30	46	89	179	406

White, female

2004.....	58,647	10	7	9	14	14	22	33	34	97	229	467
2003.....	60,899	3	10	7	16	18	24	36	52	109	256	457
2002.....	59,321	9	9	5	19	16	32	31	54	128	227	479
2001.....	58,590	8	8	8	12	21	19	24	56	101	243	401
2000.....	57,389	7	15	11	19	13	23	30	50	107	220	410
1999.....	57,139	6	17	14	12	32	25	38	57	110	224	386

Black, both sexes

2004.....	7,371	11	22	27	35	42	37	47	54	82	142	256
2003.....	7,679	8	26	23	40	41	41	43	59	80	163	239
2002.....	7,799	14	30	21	55	43	36	43	61	110	188	230
2001.....	7,560	17	23	23	31	30	47	46	81	131	147	222
2000.....	7,572	11	21	24	46	43	37	51	64	103	170	258
1999.....	7,875	18	17	20	48	36	42	66	76	95	168	255

Black, male

2004.....	4,038	10	14	21	23	27	21	32	34	40	67	123
2003.....	4,234	3	21	16	27	26	23	30	41	37	69	120
2002.....	4,326	7	20	11	30	36	23	28	30	61	94	116
2001.....	4,171	7	17	14	18	22	26	26	44	59	69	101
2000.....	4,222	6	14	14	29	30	20	31	41	52	78	124
1999.....	4,485	9	13	14	34	23	29	34	35	46	79	125

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)

Black, female

2004.....	3,333	1	8	6	12	15	16	15	20	42	75	133
2003.....	3,445	5	5	7	13	15	18	13	18	43	94	119
2002.....	3,473	7	10	10	25	7	13	15	31	49	94	114
2001.....	3,389	10	6	9	13	8	21	20	37	72	78	121
2000.....	3,350	5	7	10	17	13	17	20	23	51	92	134
1999.....	3,390	9	4	6	14	13	13	32	41	49	89	130

American Indian 1/, both sexes

2004.....	483	1	1	1	-	1	2	2	-	3	5	10
2003.....	503	-	1	-	-	1	1	-	3	5	7	11
2002.....	445	-	1	1	-	1	1	1	-	2	4	10
2001.....	417	1	-	1	-	1	1	1	1	5	5	8
2000.....	426	1	2	-	-	1	2	4	-	4	6	10
1999.....	399	1	-	-	-	-	1	-	2	2	7	7

American Indian 1/, male

2004.....	233	1	-	-	-	1	1	1	-	2	3	3
2003.....	266	-	-	-	-	1	1	-	1	2	4	6
2002.....	216	-	1	-	-	-	-	1	-	2	-	5
2001.....	194	1	-	-	-	1	1	-	1	-	3	3
2000.....	227	1	1	-	-	1	2	3	-	1	3	7
1999.....	195	-	-	-	-	-	-	-	1	2	3	5

American Indian 1/, female

2004.....	250	-	1	1	-	-	1	1	-	1	2	7
2003.....	237	-	1	-	-	-	-	-	2	3	3	5
2002.....	229	-	-	1	-	1	1	-	-	-	4	5
2001.....	223	-	-	1	-	-	-	1	-	5	2	5
2000.....	199	-	1	-	-	-	-	1	-	3	3	3
1999.....	204	1	-	-	-	-	1	-	1	-	4	2

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)												
Asian or Pacific Islander 2/, both sexes												
2004.....	1,160	-	3	1	1	3	3	1	1	3	6	17
2003.....	1,203	-	4	-	1	3	2	3	3	4	8	11
2002.....	1,109	-	4	2	2	2	-	4	2	2	5	11
2001.....	1,138	1	-	-	1	2	3	1	1	5	8	12
2000.....	1,096	1	-	1	5	1	2	3	7	8	11	22
1999.....	1,092	1	5	-	-	3	2	3	2	5	4	12
Asian or Pacific Islander 2/, male												
2004.....	736	-	1	1	-	2	3	1	1	2	4	8
2003.....	787	-	3	-	1	2	1	2	2	2	4	6
2002.....	734	-	2	1	1	1	-	4	-	-	4	6
2001.....	730	1	-	-	1	1	2	-	1	3	5	6
2000.....	714	-	-	-	4	-	1	2	4	7	4	13
1999.....	703	-	3	-	-	1	2	2	1	4	4	6
Asian or Pacific Islander 2/, female												
2004.....	424	-	2	-	1	1	-	-	-	1	2	9
2003.....	416	-	1	-	-	1	1	1	1	2	4	5
2002.....	375	-	2	1	1	1	-	-	2	2	1	5
2001.....	408	-	-	-	-	1	1	1	-	2	3	6
2000.....	382	1	-	1	1	1	1	1	3	1	7	9
1999.....	389	1	2	-	-	2	-	1	1	1	-	6

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bronchitis, not specified as acute or chronic (J40)												
All races, both sexes												
2004.....	411	13	8	2	1	1	4	1	2	1	8	12
2003.....	453	11	10	1	4	-	3	2	2	5	6	10
2002.....	504	16	5	3	-	3	1	1	-	7	2	6
2001.....	530	14	7	-	-	2	2	4	2	6	7	15
2000.....	645	11	9	2	3	3	2	1	3	4	9	11
1999.....	640	11	13	3	-	2	2	2	5	8	6	8
All races, male												
2004.....	182	10	6	2	1	1	1	1	1	1	3	7
2003.....	174	8	5	-	3	-	2	1	1	3	3	7
2002.....	185	10	4	2	-	2	-	-	-	5	-	4
2001.....	207	8	4	-	-	2	1	2	-	1	3	10
2000.....	268	6	6	2	1	2	2	1	2	3	7	4
1999.....	247	5	7	2	-	1	2	-	2	2	1	5
All races, female												
2004.....	229	3	2	-	-	-	3	-	1	-	5	5
2003.....	279	3	5	1	1	-	1	1	1	2	3	3
2002.....	319	6	1	1	-	1	1	1	-	2	2	2
2001.....	323	6	3	-	-	-	1	2	2	5	4	5
2000.....	377	5	3	-	2	1	-	-	1	1	2	7
1999.....	393	6	6	1	-	1	-	2	3	6	5	3
White, both sexes												
2004.....	378	10	6	2	1	-	3	1	2	-	6	10
2003.....	402	8	7	1	2	-	3	-	2	3	5	7
2002.....	467	11	2	2	-	1	1	-	-	6	2	5
2001.....	481	7	5	-	-	1	2	3	2	6	6	10
2000.....	589	5	5	2	2	2	2	1	2	4	4	9
1999.....	570	7	11	3	-	2	1	1	5	8	4	6

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bronchitis, not specified as acute or chronic (J40)												
White, male												
2004.....	164	7	5	2	1	-	1	1	1	-	2	5
2003.....	154	6	4	-	1	-	2	-	1	2	2	5
2002.....	170	7	1	2	-	1	-	-	-	4	-	3
2001.....	178	5	4	-	-	1	1	1	-	1	3	7
2000.....	243	4	3	2	-	1	2	1	1	3	2	4
1999.....	216	3	6	2	-	1	1	-	2	2	1	4
White, female												
2004.....	214	3	1	-	-	-	2	-	1	-	4	5
2003.....	248	2	3	1	1	-	1	-	1	1	3	2
2002.....	297	4	1	-	-	-	1	-	-	2	2	2
2001.....	303	2	1	-	-	-	1	2	2	5	3	3
2000.....	346	1	2	-	2	1	-	-	1	1	2	5
1999.....	354	4	5	1	-	1	-	1	3	6	3	2
Black, both sexes												
2004.....	26	3	2	-	-	1	1	-	-	1	2	2
2003.....	39	3	1	-	2	-	-	1	-	2	1	3
2002.....	35	5	3	1	-	2	-	1	-	1	-	1
2001.....	42	7	2	-	-	1	-	1	-	-	1	4
2000.....	47	4	3	-	1	1	-	-	-	-	5	2
1999.....	56	4	1	-	-	-	1	1	-	-	2	2
Black, male												
2004.....	15	3	1	-	-	1	-	-	-	1	1	2
2003.....	14	2	-	-	2	-	-	1	-	1	1	2
2002.....	14	3	3	-	-	1	-	-	-	1	-	1
2001.....	26	3	-	-	-	1	-	1	-	-	-	3
2000.....	20	1	2	-	1	1	-	-	-	-	5	-
1999.....	25	2	1	-	-	-	1	-	-	-	-	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Chronic bronchitis (J41-J42)

All races, both sexes

2004.....	362	4	4	1	-	1	1	1	1	1	2	3
2003.....	397	9	3	-	1	2	2	3	3	3	3	7
2002.....	451	8	8	-	2	-	2	1	4	5	4	10
2001.....	429	7	1	-	-	1	-	-	-	3	5	6
2000.....	522	7	5	-	-	-	-	2	2	3	4	8
1999.....	532	9	6	-	1	1	2	1	3	2	2	6

All races, male

2004.....	156	2	3	1	-	1	1	1	1	1	-	3
2003.....	177	7	1	-	-	2	2	-	1	2	1	6
2002.....	235	5	4	-	-	-	1	-	2	4	3	9
2001.....	189	3	1	-	-	-	-	-	-	2	2	2
2000.....	235	3	1	-	-	-	-	2	2	1	3	4
1999.....	251	4	4	-	1	1	-	-	1	1	1	4

All races, female

2004.....	206	2	1	-	-	-	-	-	-	-	2	-
2003.....	220	2	2	-	1	-	-	3	2	1	2	1
2002.....	216	3	4	-	2	-	1	1	2	1	1	1
2001.....	240	4	-	-	-	1	-	-	-	1	3	4
2000.....	287	4	4	-	-	-	-	-	-	2	1	4
1999.....	281	5	2	-	-	-	2	1	2	1	1	2

White, both sexes

2004.....	313	2	-	-	-	1	1	1	1	-	2	3
2003.....	349	7	1	-	-	1	2	1	3	3	2	4
2002.....	409	5	5	-	1	-	2	1	4	5	3	7
2001.....	397	4	1	-	-	1	-	-	-	3	3	6
2000.....	484	4	2	-	-	-	-	2	1	3	3	8
1999.....	486	4	4	-	1	1	2	1	1	2	2	3

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Chronic bronchitis (J41-J42)

White, male

2004.....	131	-	-	-	-	1	1	1	1	-	-	3
2003.....	155	6	1	-	-	1	2	-	1	2	1	4
2002.....	206	4	2	-	-	-	1	-	2	4	2	6
2001.....	176	2	1	-	-	-	-	-	-	2	1	2
2000.....	215	1	-	-	-	-	-	2	1	1	2	4
1999.....	224	3	2	-	1	1	-	-	-	1	1	1

White, female

2004.....	182	2	-	-	-	-	-	-	-	-	2	-
2003.....	194	1	-	-	-	-	-	1	2	1	1	-
2002.....	203	1	3	-	1	-	1	1	2	1	1	1
2001.....	221	2	-	-	-	1	-	-	-	1	2	4
2000.....	269	3	2	-	-	-	-	-	-	2	1	4
1999.....	262	1	2	-	-	-	2	1	1	1	1	2

Black, both sexes

2004.....	39	2	3	1	-	-	-	-	-	1	-	-
2003.....	36	2	2	-	1	1	-	2	-	-	1	3
2002.....	31	3	3	-	1	-	-	-	-	-	1	3
2001.....	21	3	-	-	-	-	-	-	-	-	2	-
2000.....	33	3	3	-	-	-	-	-	1	-	1	-
1999.....	33	3	-	-	-	-	-	-	2	-	-	2

Black, male

2004.....	20	2	3	1	-	-	-	-	-	1	-	-
2003.....	16	1	-	-	-	1	-	-	-	-	-	2
2002.....	21	1	2	-	-	-	-	-	-	-	1	3
2001.....	6	1	-	-	-	-	-	-	-	-	1	-
2000.....	15	2	1	-	-	-	-	-	1	-	1	-
1999.....	19	1	-	-	-	-	-	-	1	-	-	2

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Emphysema (J43)												
All races, both sexes												
2004.....	13,639	3	2	-	1	-	3	1	9	23	57	130
2003.....	14,861	-	1	-	-	1	3	2	3	21	66	157
2002.....	15,489	1	3	-	-	1	1	2	8	21	73	156
2001.....	16,242	3	-	-	1	1	1	1	6	24	51	141
2000.....	16,700	3	2	-	-	1	4	6	5	23	59	162
1999.....	17,787	1	-	1	-	-	1	1	8	19	78	136
All races, male												
2004.....	7,075	2	-	-	1	-	3	-	5	14	35	87
2003.....	7,549	-	1	-	-	-	2	2	1	14	35	105
2002.....	8,084	-	3	-	-	-	-	1	7	14	52	95
2001.....	8,415	1	-	-	1	1	-	-	4	17	35	89
2000.....	8,671	3	-	-	-	1	4	1	4	15	31	101
1999.....	9,480	-	-	-	-	-	1	-	6	9	40	85
All races, female												
2004.....	6,564	1	2	-	-	-	-	1	4	9	22	43
2003.....	7,312	-	-	-	-	1	1	-	2	7	31	52
2002.....	7,405	1	-	-	-	1	1	1	1	7	21	61
2001.....	7,827	2	-	-	-	-	1	1	2	7	16	52
2000.....	8,029	-	2	-	-	-	-	5	1	8	28	61
1999.....	8,307	1	-	1	-	-	-	1	2	10	38	51
White, both sexes												
2004.....	12,757	1	1	-	1	-	2	-	7	20	41	110
2003.....	13,856	-	1	-	-	1	3	1	2	17	53	129
2002.....	14,476	1	1	-	-	1	-	2	6	17	50	129
2001.....	15,243	2	-	-	-	1	-	1	3	22	42	127
2000.....	15,607	3	-	-	-	-	2	6	4	18	45	121
1999.....	16,674	-	-	1	-	-	-	1	6	15	55	104

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Emphysema (J43)												
White, male												
2004.....	6,511	-	-	-	1	-	2	-	3	11	24	71
2003.....	6,915	-	1	-	-	-	2	1	-	11	28	86
2002.....	7,426	-	1	-	-	-	-	1	5	11	37	78
2001.....	7,798	-	-	-	-	1	-	-	2	16	28	82
2000.....	7,985	3	-	-	-	-	2	1	3	11	23	72
1999.....	8,754	-	-	-	-	-	-	-	4	5	30	64
White, female												
2004.....	6,246	1	1	-	-	-	-	-	4	9	17	39
2003.....	6,941	-	-	-	-	1	1	-	2	6	25	43
2002.....	7,050	1	-	-	-	1	-	1	1	6	13	51
2001.....	7,445	2	-	-	-	-	-	1	1	6	14	45
2000.....	7,622	-	-	-	-	-	-	5	1	7	22	49
1999.....	7,920	-	-	1	-	-	-	1	2	10	25	40
Black, both sexes												
2004.....	719	2	-	-	-	-	1	1	1	3	12	17
2003.....	810	-	-	-	-	-	-	1	1	3	12	27
2002.....	850	-	1	-	-	-	1	-	2	4	22	26
2001.....	811	1	-	-	1	-	1	-	3	1	9	13
2000.....	901	-	1	-	-	1	2	-	1	4	13	38
1999.....	930	1	-	-	-	-	-	-	-	3	22	32
Black, male												
2004.....	460	2	-	-	-	-	1	-	1	3	8	14
2003.....	498	-	-	-	-	-	-	1	1	2	6	18
2002.....	541	-	1	-	-	-	-	-	2	3	15	16
2001.....	487	1	-	-	1	-	-	-	2	1	7	6
2000.....	559	-	-	-	-	1	2	-	1	3	8	26
1999.....	598	-	-	-	-	-	-	-	-	3	10	21

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic obstructive pulmonary disease (J44)												
All races, both sexes												
2004.....	102,801	8	5	4	4	6	2	12	28	89	318	786
2003.....	105,638	4	4	1	1	8	11	11	23	116	304	772
2002.....	103,133	10	6	3	5	5	8	9	35	96	318	742
2001.....	100,572	6	4	1	3	8	13	6	31	109	299	678
2000.....	98,704	7	3	1	2	2	8	8	15	93	263	666
1999.....	99,595	10	7	5	2	9	8	12	26	66	255	604
All races, male												
2004.....	49,576	3	2	4	2	2	-	9	18	50	164	389
2003.....	51,058	4	2	-	1	5	7	5	15	61	142	432
2002.....	50,356	5	3	2	1	3	3	5	18	42	175	393
2001.....	49,142	4	2	-	3	4	9	2	20	56	142	382
2000.....	48,932	5	1	-	1	2	6	2	7	44	132	372
1999.....	50,491	7	3	2	2	5	3	2	10	35	133	352
All races, female												
2004.....	53,225	5	3	-	2	4	2	3	10	39	154	397
2003.....	54,580	-	2	1	-	3	4	6	8	55	162	340
2002.....	52,777	5	3	1	4	2	5	4	17	54	143	349
2001.....	51,430	2	2	1	-	4	4	4	11	53	157	296
2000.....	49,772	2	2	1	1	-	2	6	8	49	131	294
1999.....	49,104	3	4	3	-	4	5	10	16	31	122	252
White, both sexes												
2004.....	95,909	5	2	1	3	5	1	10	22	67	272	666
2003.....	98,568	4	2	1	1	3	8	9	16	95	248	665
2002.....	96,119	6	4	-	5	4	8	7	19	73	265	638
2001.....	93,816	3	3	1	2	5	7	4	25	78	251	570
2000.....	92,140	6	3	1	2	1	5	6	8	71	210	559
1999.....	92,787	4	5	5	1	6	5	10	19	51	199	488

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic obstructive pulmonary disease (J44)												
White, male												
2004.....	45,711	1	1	1	1	1	-	7	15	38	141	337
2003.....	46,979	4	-	-	1	1	5	4	10	52	114	372
2002.....	46,345	4	2	-	1	2	3	4	13	32	145	332
2001.....	45,248	2	1	-	2	1	5	1	17	40	119	327
2000.....	45,075	5	1	-	1	1	4	1	2	32	108	306
1999.....	46,466	3	1	2	1	2	1	2	10	28	101	282
White, female												
2004.....	50,198	4	1	-	2	4	1	3	7	29	131	329
2003.....	51,589	-	2	1	-	2	3	5	6	43	134	293
2002.....	49,774	2	2	-	4	2	5	3	6	41	120	306
2001.....	48,568	1	2	1	-	4	2	3	8	38	132	243
2000.....	47,065	1	2	1	1	-	1	5	6	39	102	253
1999.....	46,321	1	4	3	-	4	4	8	9	23	98	206
Black, both sexes												
2004.....	5,579	2	3	3	1	1	1	2	6	20	43	108
2003.....	5,764	-	1	-	-	4	2	2	5	18	49	97
2002.....	5,787	4	1	3	-	1	-	1	15	22	49	96
2001.....	5,578	2	1	-	1	2	6	1	6	30	45	98
2000.....	5,433	1	-	-	-	1	1	2	7	20	51	90
1999.....	5,711	6	2	-	1	3	3	2	7	15	53	108
Black, male												
2004.....	3,088	1	1	3	1	1	-	2	3	11	21	48
2003.....	3,261	-	1	-	-	3	2	1	4	8	25	55
2002.....	3,253	1	-	2	-	1	-	-	5	10	27	56
2001.....	3,193	1	1	-	1	2	4	1	3	16	22	49
2000.....	3,147	-	-	-	-	1	1	1	5	11	23	54
1999.....	3,362	4	2	-	1	3	2	-	-	7	31	63

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Asthma (J45-J46)												
All races, both sexes												
2004.....	3,816	7	29	39	66	76	83	97	100	157	219	293
2003.....	4,099	7	37	35	75	70	88	98	129	152	259	317
2002.....	4,261	4	43	35	88	84	85	102	133	207	265	303
2001.....	4,269	10	31	41	58	60	80	88	145	207	248	279
2000.....	4,487	8	32	43	86	78	89	105	144	183	275	318
1999.....	4,657	5	27	40	86	95	87	121	141	206	241	312
All races, male												
2004.....	1,386	7	19	23	41	50	49	52	61	64	94	122
2003.....	1,493	4	29	23	48	40	51	58	69	60	100	127
2002.....	1,580	3	30	20	49	63	47	63	66	92	106	113
2001.....	1,479	6	22	24	33	35	45	49	67	93	102	103
2000.....	1,632	6	20	22	52	52	50	64	78	81	115	128
1999.....	1,620	3	16	25	60	53	55	64	64	94	90	96
All races, female												
2004.....	2,430	-	10	16	25	26	34	45	39	93	125	171
2003.....	2,606	3	8	12	27	30	37	40	60	92	159	190
2002.....	2,681	1	13	15	39	21	38	39	67	115	159	190
2001.....	2,790	4	9	17	25	25	35	39	78	114	146	176
2000.....	2,855	2	12	21	34	26	39	41	66	102	160	190
1999.....	3,037	2	11	15	26	42	32	57	77	112	151	216
White, both sexes												
2004.....	2,658	5	13	14	31	32	44	50	53	96	130	152
2003.....	2,888	4	13	12	37	31	47	59	72	90	152	197
2002.....	3,014	2	18	15	32	41	49	57	88	121	145	187
2001.....	2,990	5	11	17	28	31	36	43	71	99	148	164
2000.....	3,144	5	18	18	36	36	53	49	83	95	161	178
1999.....	3,328	1	11	20	39	59	47	55	72	123	143	191

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Asthma (J45-J46)												
White, male												
2004.....	851	5	9	5	19	22	25	20	31	37	55	58
2003.....	961	4	8	7	22	16	28	29	31	32	59	78
2002.....	1,017	1	15	10	18	28	24	31	43	43	54	68
2001.....	937	4	6	10	16	15	20	25	26	48	56	58
2000.....	1,057	3	9	8	20	24	31	29	41	37	68	79
1999.....	1,046	1	5	11	27	32	28	28	30	53	46	55
White, female												
2004.....	1,807	-	4	9	12	10	19	30	22	59	75	94
2003.....	1,927	-	5	5	15	15	19	30	41	58	93	119
2002.....	1,997	1	3	5	14	13	25	26	45	78	91	119
2001.....	2,053	1	5	7	12	16	16	18	45	51	92	106
2000.....	2,087	2	9	10	16	12	22	20	42	58	93	99
1999.....	2,282	-	6	9	12	27	19	27	42	70	97	136
Black, both sexes												
2004.....	1,008	2	14	23	34	40	34	44	47	57	85	129
2003.....	1,030	3	22	23	37	36	39	37	53	57	100	109
2002.....	1,096	2	22	17	54	40	35	41	44	83	116	104
2001.....	1,108	4	20	23	29	27	40	44	72	100	90	107
2000.....	1,158	3	14	24	45	40	34	49	55	79	100	128
1999.....	1,145	4	14	20	47	33	38	63	67	77	91	111
Black, male												
2004.....	455	2	9	17	22	25	20	30	30	24	37	59
2003.....	445	-	20	16	25	22	21	27	36	26	37	43
2002.....	497	2	14	9	30	34	23	28	23	47	51	40
2001.....	459	1	16	14	16	19	22	24	39	42	39	43
2000.....	481	3	11	14	28	27	17	30	34	38	41	44
1999.....	481	2	10	14	33	20	26	34	34	36	38	38

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Asthma (J45-J46)												
Black, female												
2004.....	553	-	5	6	12	15	14	14	17	33	48	70
2003.....	585	3	2	7	12	14	18	10	17	31	63	66
2002.....	599	-	8	8	24	6	12	13	21	36	65	64
2001.....	649	3	4	9	13	8	18	20	33	58	51	64
2000.....	677	-	3	10	17	13	17	19	21	41	59	84
1999.....	664	2	4	6	14	13	12	29	33	41	53	73
American Indian 1/, both sexes												
2004.....	19	-	1	1	-	1	2	2	-	1	1	2
2003.....	38	-	-	-	-	-	1	-	3	3	2	5
2002.....	25	-	-	1	-	1	1	-	-	2	2	3
2001.....	33	1	-	1	-	-	1	1	1	4	3	2
2000.....	31	-	-	-	-	1	1	4	-	2	5	1
1999.....	29	-	-	-	-	-	1	-	1	1	3	2
American Indian 1/, male												
2004.....	8	-	-	-	-	1	1	1	-	1	-	2
2003.....	18	-	-	-	-	-	1	-	1	2	2	2
2002.....	9	-	-	-	-	-	-	-	-	2	-	1
2001.....	11	1	-	-	-	-	1	-	1	-	2	-
2000.....	14	-	-	-	-	1	1	3	-	-	2	-
1999.....	12	-	-	-	-	-	-	-	-	1	2	-
American Indian 1/, female												
2004.....	11	-	1	1	-	-	1	1	-	-	1	-
2003.....	20	-	-	-	-	-	-	-	2	1	-	3
2002.....	16	-	-	1	-	1	1	-	-	-	2	2
2001.....	22	-	-	1	-	-	-	1	-	4	1	2
2000.....	17	-	-	-	-	-	-	1	-	2	3	1
1999.....	17	-	-	-	-	-	1	-	1	-	1	2

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Asthma (J45-J46)												
Asian or Pacific Islander 2/, both sexes												
2004.....	131	-	1	1	1	3	3	1	-	3	3	10
2003.....	143	-	2	-	1	3	1	2	1	2	5	6
2002.....	126	-	3	2	2	2	-	4	1	1	2	9
2001.....	138	-	-	-	1	2	3	-	1	4	7	6
2000.....	154	-	-	1	5	1	1	3	6	7	9	11
1999.....	155	-	2	-	-	3	1	3	1	5	4	8
Asian or Pacific Islander 2/, male												
2004.....	72	-	1	1	-	2	3	1	-	2	2	3
2003.....	69	-	1	-	1	2	1	2	1	-	2	4
2002.....	57	-	1	1	1	1	-	4	-	-	1	4
2001.....	72	-	-	-	1	1	2	-	1	3	5	2
2000.....	80	-	-	-	4	-	1	2	3	6	4	5
1999.....	81	-	1	-	-	1	1	2	-	4	4	3
Asian or Pacific Islander 2/, female												
2004.....	59	-	-	-	1	1	-	-	-	1	1	7
2003.....	74	-	1	-	-	1	-	-	-	2	3	2
2002.....	69	-	2	1	1	1	-	-	1	1	1	5
2001.....	66	-	-	-	-	1	1	-	-	1	2	4
2000.....	74	-	-	1	1	1	-	1	3	1	5	6
1999.....	74	-	1	-	-	2	-	1	1	1	-	5

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bronchiectasis (J47)												
All races, both sexes												
2004.....	958	-	-	-	2	1	1	-	3	6	6	8
2003.....	934	-	-	-	-	3	-	2	4	4	11	13
2002.....	978	-	-	-	-	-	2	1	5	4	6	9
2001.....	971	1	-	-	-	2	1	3	4	6	4	8
2000.....	951	-	-	2	-	2	1	3	2	4	10	9
1999.....	970	-	1	-	1	-	2	8	2	1	6	13
All races, male												
2004.....	271	-	-	-	-	1	1	-	2	2	3	5
2003.....	263	-	-	-	-	3	-	1	2	2	6	4
2002.....	273	-	-	-	-	-	1	1	2	2	2	3
2001.....	265	-	-	-	-	1	1	2	-	2	-	5
2000.....	266	-	-	2	-	1	-	2	-	1	4	8
1999.....	326	-	-	-	1	-	2	3	-	1	3	5
All races, female												
2004.....	687	-	-	-	2	-	-	-	1	4	3	3
2003.....	671	-	-	-	-	-	-	1	2	2	5	9
2002.....	705	-	-	-	-	-	1	-	3	2	4	6
2001.....	706	1	-	-	-	1	-	1	4	4	4	3
2000.....	685	-	-	-	-	1	1	1	2	3	6	1
1999.....	644	-	1	-	-	-	-	5	2	-	3	8
White, both sexes												
2004.....	899	-	-	-	-	1	1	-	3	5	5	5
2003.....	854	-	-	-	-	2	-	-	4	3	4	10
2002.....	910	-	-	-	-	-	2	1	4	4	5	4
2001.....	892	1	-	-	-	1	-	2	2	4	3	5
2000.....	876	-	-	2	-	1	1	3	1	3	7	7
1999.....	890	-	-	-	1	-	2	5	-	1	5	9

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bronchiectasis (J47)												
Asian or Pacific Islander 2/, both sexes												
2004.....	27	-	-	-	-	-	-	-	-	-	1	-
2003.....	41	-	-	-	-	1	-	-	-	-	2	-
2002.....	29	-	-	-	-	-	-	-	1	-	1	2
2001.....	40	-	-	-	-	-	1	-	1	1	1	1
2000.....	37	-	-	-	-	-	-	-	-	-	2	1
1999.....	33	-	-	-	-	-	-	1	2	-	-	1
Asian or Pacific Islander 2/, male												
2004.....	7	-	-	-	-	-	-	-	-	-	-	-
2003.....	14	-	-	-	-	1	-	-	-	-	2	-
2002.....	9	-	-	-	-	-	-	-	1	-	-	-
2001.....	19	-	-	-	-	-	1	-	-	-	-	-
2000.....	8	-	-	-	-	-	-	-	-	-	-	-
1999.....	15	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	20	-	-	-	-	-	-	-	-	-	1	-
2003.....	27	-	-	-	-	-	-	-	-	-	-	-
2002.....	20	-	-	-	-	-	-	-	-	-	1	2
2001.....	21	-	-	-	-	-	-	-	1	1	1	1
2000.....	29	-	-	-	-	-	-	-	-	-	2	1
1999.....	18	-	-	-	-	-	-	-	2	-	-	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
-------------	-----------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Lung diseases due to external agents (J60-J70)

All races, both sexes

2004.....	17,913	12	9	3	9	13	27	35	34	58	107	176
2003.....	18,542	14	10	4	13	13	36	26	44	69	106	196
2002.....	18,791	4	7	7	9	16	29	31	44	72	128	148
2001.....	18,524	12	15	5	10	10	24	32	51	65	123	145
2000.....	17,863	10	11	8	5	12	25	24	42	67	122	166
1999.....	16,471	16	15	4	12	10	23	35	49	63	111	136

All races, male

2004.....	9,643	4	5	3	3	7	14	23	18	35	67	110
2003.....	9,819	9	5	4	6	6	27	16	27	42	54	116
2002.....	9,974	-	3	3	6	10	18	17	25	41	82	94
2001.....	9,725	6	7	3	4	5	17	15	27	45	80	83
2000.....	9,543	4	5	5	4	8	15	12	26	46	76	105
1999.....	8,878	9	6	-	6	8	15	21	28	38	65	86

All races, female

2004.....	8,270	8	4	-	6	6	13	12	16	23	40	66
2003.....	8,723	5	5	-	7	7	9	10	17	27	52	80
2002.....	8,817	4	4	4	3	6	11	14	19	31	46	54
2001.....	8,799	6	8	2	6	5	7	17	24	20	43	62
2000.....	8,320	6	6	3	1	4	10	12	16	21	46	61
1999.....	7,593	7	9	4	6	2	8	14	21	25	46	50

White, both sexes

2004.....	16,040	5	7	2	8	10	19	27	26	48	80	144
2003.....	16,536	8	8	3	11	8	24	19	36	52	79	158
2002.....	16,903	3	5	6	8	13	22	27	33	55	99	121
2001.....	16,582	9	13	4	7	8	17	27	39	49	101	115
2000.....	16,027	8	9	8	4	10	17	18	32	44	103	124
1999.....	14,702	13	8	4	8	8	14	24	38	45	81	105

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Lung diseases due to external agents (J60-J70)												
White, male												
2004.....	8,638	2	4	2	3	6	10	18	15	32	50	91
2003.....	8,791	5	5	3	5	4	19	11	22	33	44	94
2002.....	8,992	-	3	3	5	8	13	16	19	31	62	76
2001.....	8,792	6	5	3	2	4	14	14	23	34	67	72
2000.....	8,576	4	4	5	3	7	11	8	17	32	65	78
1999.....	7,900	7	2	-	5	6	8	13	22	28	47	69
White, female												
2004.....	7,402	3	3	-	5	4	9	9	11	16	30	53
2003.....	7,745	3	3	-	6	4	5	8	14	19	35	64
2002.....	7,911	3	2	3	7	5	9	11	14	24	37	45
2001.....	7,790	3	8	1	5	4	3	13	16	15	34	43
2000.....	7,451	4	5	3	1	3	6	10	15	12	38	46
1999.....	6,802	6	6	4	3	2	6	11	16	17	34	36
Black, both sexes												
2004.....	1,552	6	2	-	1	3	8	7	7	9	26	28
2003.....	1,661	4	1	1	2	4	10	6	7	15	27	36
2002.....	1,597	1	2	1	1	2	6	4	10	11	28	23
2001.....	1,659	3	2	1	3	1	6	5	11	13	22	27
2000.....	1,583	1	2	-	1	2	7	6	9	22	19	39
1999.....	1,579	2	7	-	3	2	9	10	9	16	28	28
Black, male												
2004.....	820	1	1	-	-	1	4	5	3	3	17	17
2003.....	828	3	-	1	1	2	6	5	5	7	10	22
2002.....	813	-	-	-	1	1	4	1	5	5	19	14
2001.....	797	-	2	-	2	-	2	1	3	8	13	9
2000.....	823	-	1	-	1	1	3	4	8	13	11	26
1999.....	842	1	4	-	1	2	7	7	5	9	16	15

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Lung diseases due to external agents (J60-J70)												
Black, female												
2004.....	732	5	1	-	1	2	4	2	4	6	9	11
2003.....	833	1	1	-	1	2	4	1	2	8	17	14
2002.....	784	1	2	1	-	1	2	3	5	6	9	9
2001.....	862	3	-	1	1	1	4	4	8	5	9	18
2000.....	760	1	1	-	-	1	4	2	1	9	8	13
1999.....	737	1	3	-	2	-	2	3	4	7	12	13
American Indian 1/, both sexes												
2004.....	66	1	-	1	-	-	-	-	-	1	-	2
2003.....	75	2	1	-	-	-	-	-	1	-	-	1
2002.....	78	-	-	-	-	1	-	-	1	3	1	2
2001.....	63	-	-	-	-	-	1	-	1	2	-	2
2000.....	50	-	-	-	-	-	-	-	1	-	-	2
1999.....	53	-	-	-	-	-	-	1	2	1	1	1
American Indian 1/, male												
2004.....	31	1	-	1	-	-	-	-	-	-	-	-
2003.....	38	1	-	-	-	-	-	-	-	-	-	-
2002.....	49	-	-	-	-	1	-	-	1	2	1	2
2001.....	27	-	-	-	-	-	1	-	1	2	-	1
2000.....	28	-	-	-	-	-	-	-	1	-	-	-
1999.....	35	-	-	-	-	-	-	1	1	-	1	1
American Indian 1/, female												
2004.....	35	-	-	-	-	-	-	-	-	1	-	2
2003.....	37	1	1	-	-	-	-	-	1	-	-	1
2002.....	29	-	-	-	-	-	-	-	-	1	-	-
2001.....	36	-	-	-	-	-	-	-	-	-	-	1
2000.....	22	-	-	-	-	-	-	-	-	-	-	2
1999.....	18	-	-	-	-	-	-	-	1	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Lung diseases due to external agents (J60-J70)												
Asian or Pacific Islander 2/, both sexes												
2004.....	255	-	-	-	-	-	-	1	1	-	1	2
2003.....	270	-	-	-	-	1	2	1	-	2	-	1
2002.....	213	-	-	-	-	-	1	-	-	3	-	2
2001.....	220	-	-	-	-	1	-	-	-	1	-	1
2000.....	203	1	-	-	-	-	1	-	-	1	-	1
1999.....	137	1	-	-	1	-	-	-	-	1	1	2
Asian or Pacific Islander 2/, male												
2004.....	154	-	-	-	-	-	-	-	-	-	-	2
2003.....	162	-	-	-	-	-	2	-	-	2	-	-
2002.....	120	-	-	-	-	-	1	-	-	3	-	2
2001.....	109	-	-	-	-	1	-	-	-	1	-	1
2000.....	116	-	-	-	-	-	1	-	-	1	-	1
1999.....	101	1	-	-	-	-	-	-	-	1	1	1
Asian or Pacific Islander 2/, female												
2004.....	101	-	-	-	-	-	-	1	1	-	1	-
2003.....	108	-	-	-	-	1	-	1	-	-	-	1
2002.....	93	-	-	-	-	-	-	-	-	-	-	-
2001.....	111	-	-	-	-	-	-	-	-	-	-	-
2000.....	87	1	-	-	-	-	-	-	-	-	-	-
1999.....	36	-	-	-	1	-	-	-	-	-	-	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumoconioses and chemical effects (J60-J66,J68)												
All races, both sexes												
2004.....	1,033	-	-	-	-	1	-	-	-	2	2	4
2003.....	1,114	-	-	-	-	-	1	-	1	2	5	4
2002.....	1,114	2	-	-	-	-	-	-	-	-	5	5
2001.....	1,136	-	1	-	-	-	-	-	1	3	2	3
2000.....	1,161	-	-	-	-	-	-	1	2	2	5	5
1999.....	1,113	-	3	-	-	1	-	-	-	3	4	3
All races, male												
2004.....	993	-	-	-	-	1	-	-	-	2	2	2
2003.....	1,065	-	-	-	-	-	1	-	1	2	4	4
2002.....	1,057	-	-	-	-	-	-	-	-	-	4	4
2001.....	1,095	-	1	-	-	-	-	-	1	3	2	2
2000.....	1,113	-	-	-	-	-	-	1	2	1	5	5
1999.....	1,079	-	3	-	-	1	-	-	-	2	3	3
All races, female												
2004.....	40	-	-	-	-	-	-	-	-	-	-	2
2003.....	49	-	-	-	-	-	-	-	-	-	1	-
2002.....	57	2	-	-	-	-	-	-	-	-	1	1
2001.....	41	-	-	-	-	-	-	-	-	-	-	1
2000.....	48	-	-	-	-	-	-	-	-	1	-	-
1999.....	34	-	-	-	-	-	-	-	-	1	1	-
White, both sexes												
2004.....	977	-	-	-	-	1	-	-	-	2	2	4
2003.....	1,058	-	-	-	-	-	1	-	-	1	4	3
2002.....	1,048	1	-	-	-	-	-	-	-	-	4	3
2001.....	1,073	-	1	-	-	-	-	-	1	3	2	1
2000.....	1,099	-	-	-	-	-	-	1	2	1	3	5
1999.....	1,037	-	-	-	-	-	-	-	-	1	3	3

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumoconioses and chemical effects (J60-J66,J68)												
White, male												
2004.....	938	-	-	-	-	1	-	-	-	2	2	2
2003.....	1,012	-	-	-	-	-	1	-	-	1	4	3
2002.....	998	-	-	-	-	-	-	-	-	-	3	2
2001.....	1,034	-	1	-	-	-	-	-	1	3	2	1
2000.....	1,053	-	-	-	-	-	-	1	2	-	3	5
1999.....	1,006	-	-	-	-	-	-	-	-	1	2	3
White, female												
2004.....	39	-	-	-	-	-	-	-	-	-	-	2
2003.....	46	-	-	-	-	-	-	-	-	-	-	-
2002.....	50	1	-	-	-	-	-	-	-	-	1	1
2001.....	39	-	-	-	-	-	-	-	-	-	-	-
2000.....	46	-	-	-	-	-	-	-	-	1	-	-
1999.....	31	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes												
2004.....	54	-	-	-	-	-	-	-	-	-	-	-
2003.....	48	-	-	-	-	-	-	-	1	-	1	1
2002.....	58	1	-	-	-	-	-	-	-	-	1	2
2001.....	57	-	-	-	-	-	-	-	-	-	-	2
2000.....	55	-	-	-	-	-	-	-	-	1	2	-
1999.....	67	-	3	-	-	1	-	-	-	2	1	-
Black, male												
2004.....	54	-	-	-	-	-	-	-	-	-	-	-
2003.....	45	-	-	-	-	-	-	-	1	-	-	1
2002.....	52	-	-	-	-	-	-	-	-	-	1	2
2001.....	55	-	-	-	-	-	-	-	-	-	-	1
2000.....	53	-	-	-	-	-	-	-	-	1	2	-
1999.....	64	-	3	-	-	1	-	-	-	1	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Pneumonitis due to solids and liquids (J69)

All races, both sexes

2004.....	16,780	12	9	3	8	12	27	34	34	55	104	168
2003.....	17,335	13	10	4	12	13	33	26	43	66	100	188
2002.....	17,593	2	7	7	9	16	28	31	43	72	120	143
2001.....	17,301	12	14	5	10	10	24	31	48	61	118	142
2000.....	16,636	10	11	8	5	12	25	22	40	65	116	157
1999.....	15,268	16	12	4	12	9	23	33	45	59	107	129

All races, male

2004.....	8,593	4	5	3	3	6	14	22	18	33	64	107
2003.....	8,702	8	5	4	5	6	25	16	26	39	50	110
2002.....	8,876	-	3	3	6	10	17	17	25	41	78	90
2001.....	8,576	6	6	3	4	5	17	15	25	42	75	81
2000.....	8,390	4	5	5	4	8	15	11	24	45	70	99
1999.....	7,755	9	3	-	6	7	15	21	26	35	62	83

All races, female

2004.....	8,187	8	4	-	5	6	13	12	16	22	40	61
2003.....	8,633	5	5	-	7	7	8	10	17	27	50	78
2002.....	8,717	2	4	4	3	6	11	14	18	31	42	53
2001.....	8,725	6	8	2	6	5	7	16	23	19	43	61
2000.....	8,246	6	6	3	1	4	10	11	16	20	46	58
1999.....	7,513	7	9	4	6	2	8	12	19	24	45	46

White, both sexes

2004.....	14,968	5	7	2	7	9	19	26	26	45	77	136
2003.....	15,395	7	8	3	10	8	21	19	36	50	74	152
2002.....	15,773	2	5	6	8	13	21	27	32	55	92	118
2001.....	15,425	9	12	4	7	8	17	26	36	45	97	114
2000.....	14,869	8	9	8	4	10	17	16	30	43	99	115
1999.....	13,582	13	8	4	8	8	14	23	34	43	78	98

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	-------------	-----------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Pneumonitis due to solids and liquids (J69)

White, male

2004.....	7,644	2	4	2	3	5	10	17	15	30	47	88
2003.....	7,731	4	5	3	4	4	17	11	22	31	40	89
2002.....	7,954	-	3	3	5	8	12	16	19	31	59	74
2001.....	7,705	6	4	3	2	4	14	14	21	31	63	71
2000.....	7,488	4	4	5	3	7	11	7	15	32	61	72
1999.....	6,853	7	2	-	5	6	8	13	20	26	45	66

White, female

2004.....	7,324	3	3	-	4	4	9	9	11	15	30	48
2003.....	7,664	3	3	-	6	4	4	8	14	19	34	63
2002.....	7,819	2	2	3	3	5	9	11	13	24	33	44
2001.....	7,720	3	8	1	5	4	3	12	15	14	34	43
2000.....	7,381	4	5	3	1	3	6	9	15	11	38	43
1999.....	6,729	6	6	4	3	2	6	10	14	17	33	32

Black, both sexes

2004.....	1,495	6	2	-	1	3	8	7	7	9	26	28
2003.....	1,607	4	1	1	2	4	10	6	6	15	26	34
2002.....	1,537	-	2	1	1	2	6	4	10	11	27	21
2001.....	1,599	3	2	1	3	1	6	5	11	13	21	25
2000.....	1,521	1	2	-	1	2	7	6	9	21	17	39
1999.....	1,505	2	4	-	3	1	9	9	9	14	27	28

Black, male

2004.....	766	1	1	-	-	1	4	5	3	3	17	17
2003.....	781	3	-	1	1	2	6	5	4	7	10	21
2002.....	760	-	-	-	1	1	4	1	5	5	18	12
2001.....	741	-	2	-	2	-	2	1	3	8	12	8
2000.....	765	-	1	-	1	1	3	4	8	12	9	26
1999.....	775	1	1	-	1	1	7	7	5	8	15	15

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonitis due to solids and liquids (J69)												
Black, female												
2004.....	729	5	1	-	1	2	4	2	4	6	9	11
2003.....	826	1	1	-	1	2	4	1	2	8	16	13
2002.....	777	-	2	1	-	1	2	3	5	6	9	9
2001.....	858	3	-	1	1	1	4	4	8	5	9	17
2000.....	756	1	1	-	-	1	4	2	1	9	8	13
1999.....	730	1	3	-	2	-	2	2	4	6	12	13
American Indian 1/, both sexes												
2004.....	64	1	-	1	-	-	-	-	-	1	-	2
2003.....	70	2	1	-	-	-	-	-	1	-	-	1
2002.....	73	-	-	-	-	1	-	-	1	3	1	2
2001.....	59	-	-	-	-	-	1	-	1	2	-	2
2000.....	49	-	-	-	-	-	-	-	1	-	-	2
1999.....	46	-	-	-	-	-	-	1	2	1	1	1
American Indian 1/, male												
2004.....	31	1	-	1	-	-	-	-	-	-	-	-
2003.....	33	1	-	-	-	-	-	-	-	-	-	-
2002.....	44	-	-	-	-	1	-	-	1	2	1	2
2001.....	23	-	-	-	-	-	1	-	1	2	-	1
2000.....	27	-	-	-	-	-	-	-	1	-	-	-
1999.....	28	-	-	-	-	-	-	1	1	-	1	1
American Indian 1/, female												
2004.....	33	-	-	-	-	-	-	-	-	1	-	2
2003.....	37	1	1	-	-	-	-	-	1	-	-	1
2002.....	29	-	-	-	-	-	-	-	-	1	-	-
2001.....	36	-	-	-	-	-	-	-	-	-	-	1
2000.....	22	-	-	-	-	-	-	-	-	-	-	2
1999.....	18	-	-	-	-	-	-	-	1	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonitis due to solids and liquids (J69)												
Asian or Pacific Islander 2/, both sexes												
2004.....	253	-	-	-	-	-	-	1	1	-	1	2
2003.....	263	-	-	-	-	1	2	1	-	1	-	1
2002.....	210	-	-	-	-	-	1	-	-	3	-	2
2001.....	218	-	-	-	-	1	-	-	-	1	-	1
2000.....	197	1	-	-	-	-	1	-	-	1	-	1
1999.....	135	1	-	-	1	-	-	-	-	1	1	2
Asian or Pacific Islander 2/, male												
2004.....	152	-	-	-	-	-	-	-	-	-	-	2
2003.....	157	-	-	-	-	-	2	-	-	1	-	-
2002.....	118	-	-	-	-	-	1	-	-	3	-	2
2001.....	107	-	-	-	-	1	-	-	-	1	-	1
2000.....	110	-	-	-	-	-	1	-	-	1	-	1
1999.....	99	1	-	-	-	-	-	-	-	1	1	1
Asian or Pacific Islander 2/, female												
2004.....	101	-	-	-	-	-	-	1	1	-	1	-
2003.....	106	-	-	-	-	1	-	1	-	-	-	1
2002.....	92	-	-	-	-	-	-	-	-	-	-	-
2001.....	111	-	-	-	-	-	-	-	-	-	-	-
2000.....	87	1	-	-	-	-	-	-	-	-	-	-
1999.....	36	-	-	-	1	-	-	-	-	-	-	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	-------------	-----------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Pneumonitis due to food and vomit (J69.0)

All races, both sexes

2004.....	16,777	12	9	3	8	12	27	34	34	55	104	168
2003.....	17,329	13	10	4	12	12	33	26	43	66	100	188
2002.....	17,586	2	6	7	9	16	28	31	43	72	120	143
2001.....	17,294	11	14	5	10	10	24	31	48	61	118	142
2000.....	16,632	10	11	8	5	12	25	22	40	65	116	157
1999.....	15,257	16	12	4	12	9	23	33	45	59	106	129

All races, male

2004.....	8,591	4	5	3	3	6	14	22	18	33	64	107
2003.....	8,700	8	5	4	5	5	25	16	26	39	50	110
2002.....	8,875	-	3	3	6	10	17	17	25	41	78	90
2001.....	8,571	5	6	3	4	5	17	15	25	42	75	81
2000.....	8,388	4	5	5	4	8	15	11	24	45	70	99
1999.....	7,752	9	3	-	6	7	15	21	26	35	62	83

All races, female

2004.....	8,186	8	4	-	5	6	13	12	16	22	40	61
2003.....	8,629	5	5	-	7	7	8	10	17	27	50	78
2002.....	8,711	2	3	4	3	6	11	14	18	31	42	53
2001.....	8,723	6	8	2	6	5	7	16	23	19	43	61
2000.....	8,244	6	6	3	1	4	10	11	16	20	46	58
1999.....	7,505	7	9	4	6	2	8	12	19	24	44	46

White, both sexes

2004.....	14,966	5	7	2	7	9	19	26	26	45	77	136
2003.....	15,391	7	8	3	10	8	21	19	36	50	74	152
2002.....	15,768	2	5	6	8	13	21	27	32	55	92	118
2001.....	15,418	8	12	4	7	8	17	26	36	45	97	114
2000.....	14,865	8	9	8	4	10	17	16	30	43	99	115
1999.....	13,573	13	8	4	8	8	14	23	34	43	77	98

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Pneumonitis due to food and vomit (J69.0)

White, male

2004.....	7,643	2	4	2	3	5	10	17	15	30	47	88
2003.....	7,730	4	5	3	4	4	17	11	22	31	40	89
2002.....	7,953	-	3	3	5	8	12	16	19	31	59	74
2001.....	7,700	5	4	3	2	4	14	14	21	31	63	71
2000.....	7,486	4	4	5	3	7	11	7	15	32	61	72
1999.....	6,851	7	2	-	5	6	8	13	20	26	45	66

White, female

2004.....	7,323	3	3	-	4	4	9	9	11	15	30	48
2003.....	7,661	3	3	-	6	4	4	8	14	19	34	63
2002.....	7,815	2	2	3	3	5	9	11	13	24	33	44
2001.....	7,718	3	8	1	5	4	3	12	15	14	34	43
2000.....	7,379	4	5	3	1	3	6	9	15	11	38	43
1999.....	6,722	6	6	4	3	2	6	10	14	17	32	32

Black, both sexes

2004.....	1,494	6	2	-	1	3	8	7	7	9	26	28
2003.....	1,605	4	1	1	2	3	10	6	6	15	26	34
2002.....	1,535	-	1	1	1	2	6	4	10	11	27	21
2001.....	1,599	3	2	1	3	1	6	5	11	13	21	25
2000.....	1,521	1	2	-	1	2	7	6	9	21	17	39
1999.....	1,503	2	4	-	3	1	9	9	9	14	27	28

Black, male

2004.....	765	1	1	-	-	1	4	5	3	3	17	17
2003.....	780	3	-	1	1	1	6	5	4	7	10	21
2002.....	760	-	-	-	1	1	4	1	5	5	18	12
2001.....	741	-	2	-	2	-	2	1	3	8	12	8
2000.....	765	-	1	-	1	1	3	4	8	12	9	26
1999.....	774	1	1	-	1	1	7	7	5	8	15	15

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonitis due to food and vomit (J69.0)												
Black, female												
2004.....	729	5	1	-	1	2	4	2	4	6	9	11
2003.....	825	1	1	-	1	2	4	1	2	8	16	13
2002.....	775	-	1	1	-	1	2	3	5	6	9	9
2001.....	858	3	-	1	1	1	4	4	8	5	9	17
2000.....	756	1	1	-	-	1	4	2	1	9	8	13
1999.....	729	1	3	-	2	-	2	2	4	6	12	13
American Indian 1/, both sexes												
2004.....	64	1	-	1	-	-	-	-	-	1	-	2
2003.....	70	2	1	-	-	-	-	-	1	-	-	1
2002.....	73	-	-	-	-	1	-	-	1	3	1	2
2001.....	59	-	-	-	-	-	1	-	1	2	-	2
2000.....	49	-	-	-	-	-	-	-	1	-	-	2
1999.....	46	-	-	-	-	-	-	1	2	1	1	1
American Indian 1/, male												
2004.....	31	1	-	1	-	-	-	-	-	-	-	-
2003.....	33	1	-	-	-	-	-	-	-	-	-	-
2002.....	44	-	-	-	-	1	-	-	1	2	1	2
2001.....	23	-	-	-	-	-	1	-	1	2	-	1
2000.....	27	-	-	-	-	-	-	-	1	-	-	-
1999.....	28	-	-	-	-	-	-	1	1	-	1	1
American Indian 1/, female												
2004.....	33	-	-	-	-	-	-	-	-	1	-	2
2003.....	37	1	1	-	-	-	-	-	1	-	-	1
2002.....	29	-	-	-	-	-	-	-	-	1	-	-
2001.....	36	-	-	-	-	-	-	-	-	-	-	1
2000.....	22	-	-	-	-	-	-	-	-	-	-	2
1999.....	18	-	-	-	-	-	-	-	1	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonitis due to food and vomit (J69.0)												
Asian or Pacific Islander 2/, both sexes												
2004.....	253	-	-	-	-	-	-	1	1	-	1	2
2003.....	263	-	-	-	-	1	2	1	-	1	-	1
2002.....	210	-	-	-	-	-	1	-	-	3	-	2
2001.....	218	-	-	-	-	1	-	-	-	1	-	1
2000.....	197	1	-	-	-	-	1	-	-	1	-	1
1999.....	135	1	-	-	1	-	-	-	-	1	1	2
Asian or Pacific Islander 2/, male												
2004.....	152	-	-	-	-	-	-	-	-	-	-	2
2003.....	157	-	-	-	-	-	2	-	-	1	-	-
2002.....	118	-	-	-	-	-	1	-	-	3	-	2
2001.....	107	-	-	-	-	1	-	-	-	1	-	1
2000.....	110	-	-	-	-	-	1	-	-	1	-	1
1999.....	99	1	-	-	-	-	-	-	-	1	1	1
Asian or Pacific Islander 2/, female												
2004.....	101	-	-	-	-	-	-	1	1	-	1	-
2003.....	106	-	-	-	-	1	-	1	-	-	-	1
2002.....	92	-	-	-	-	-	-	-	-	-	-	-
2001.....	111	-	-	-	-	-	-	-	-	-	-	-
2000.....	87	1	-	-	-	-	-	-	-	-	-	-
1999.....	36	-	-	-	1	-	-	-	-	-	-	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other lung diseases due to external agents (J67,J70)												
All races, both sexes												
2004.....	100	-	-	-	1	-	-	1	-	1	1	4
2003.....	93	1	-	-	1	-	2	-	-	1	1	4
2002.....	84	-	-	-	-	-	1	-	1	-	3	-
2001.....	87	-	-	-	-	-	-	1	2	1	3	-
2000.....	66	-	-	-	-	-	-	1	-	-	1	4
1999.....	90	-	-	-	-	-	-	2	4	1	-	4
All races, male												
2004.....	57	-	-	-	-	-	-	1	-	-	1	1
2003.....	52	1	-	-	1	-	1	-	-	1	-	2
2002.....	41	-	-	-	-	-	1	-	-	-	-	-
2001.....	54	-	-	-	-	-	-	-	1	-	3	-
2000.....	40	-	-	-	-	-	-	-	-	-	1	1
1999.....	44	-	-	-	-	-	-	-	2	1	-	-
All races, female												
2004.....	43	-	-	-	1	-	-	-	-	1	-	3
2003.....	41	-	-	-	-	-	1	-	-	-	1	2
2002.....	43	-	-	-	-	-	-	-	1	-	3	-
2001.....	33	-	-	-	-	-	-	1	1	1	-	-
2000.....	26	-	-	-	-	-	-	1	-	-	-	3
1999.....	46	-	-	-	-	-	-	2	2	-	-	4
White, both sexes												
2004.....	95	-	-	-	1	-	-	1	-	1	1	4
2003.....	83	1	-	-	1	-	2	-	-	1	1	3
2002.....	82	-	-	-	-	-	1	-	1	-	3	-
2001.....	84	-	-	-	-	-	-	1	2	1	2	-
2000.....	59	-	-	-	-	-	-	1	-	-	1	4
1999.....	83	-	-	-	-	-	-	1	4	1	-	4

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Suppurative and necrotic conditions of lower respiratory tract (J85-J86)												
All races, both sexes												
2004.....	669	8	1	2	1	3	5	2	7	13	33	50
2003.....	725	6	3	2	5	3	8	7	3	13	34	58
2002.....	719	6	3	4	2	4	4	1	10	14	31	48
2001.....	785	4	1	-	2	2	8	5	9	23	32	41
2000.....	727	8	5	1	1	-	5	2	13	12	37	36
1999.....	799	5	6	2	1	6	5	11	13	24	38	47
All races, male												
2004.....	404	3	-	-	1	1	3	1	6	6	21	25
2003.....	422	4	1	1	3	-	4	5	3	8	21	37
2002.....	455	3	3	1	1	2	3	1	7	11	21	37
2001.....	458	2	1	-	1	1	3	4	6	10	24	25
2000.....	447	4	2	1	1	-	2	2	7	9	23	26
1999.....	491	2	2	1	1	2	2	5	9	15	29	33
All races, female												
2004.....	265	5	1	2	-	2	2	1	1	7	12	25
2003.....	303	2	2	1	2	3	4	2	-	5	13	21
2002.....	264	3	-	3	1	2	1	-	3	3	10	11
2001.....	327	2	-	-	1	1	5	1	3	13	8	16
2000.....	280	4	3	-	-	-	3	-	6	3	14	10
1999.....	308	3	4	1	-	4	3	6	4	9	9	14
White, both sexes												
2004.....	558	6	-	-	1	-	4	-	6	11	27	45
2003.....	596	4	1	2	4	1	7	4	2	11	27	43
2002.....	569	3	1	3	2	3	3	-	7	9	21	29
2001.....	659	1	-	-	2	1	6	3	7	17	22	36
2000.....	592	3	2	1	-	-	2	1	8	10	25	25
1999.....	642	5	3	1	1	4	2	6	11	15	29	34

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Suppurative and necrotic conditions of lower respiratory tract (J85-J86)

White, male

2004.....	341	2	-	-	1	-	2	-	5	4	17	22
2003.....	343	2	-	1	2	-	4	2	2	6	17	28
2002.....	361	1	1	1	1	2	3	-	5	7	14	23
2001.....	372	-	-	-	1	1	2	2	5	7	17	22
2000.....	363	2	1	1	-	-	1	1	5	8	14	17
1999.....	392	2	-	1	1	2	2	3	7	8	22	24

White, female

2004.....	217	4	-	-	-	-	2	-	1	7	10	23
2003.....	253	2	1	1	2	1	3	2	-	5	10	15
2002.....	208	2	-	2	1	1	-	-	2	2	7	6
2001.....	287	1	-	-	1	-	4	1	2	10	5	14
2000.....	229	1	1	-	-	-	1	-	3	2	11	8
1999.....	250	3	3	-	-	2	-	3	4	7	7	10

Black, both sexes

2004.....	92	2	1	2	-	3	1	2	1	2	5	5
2003.....	108	2	2	-	1	1	-	2	1	2	6	12
2002.....	126	3	2	1	-	1	1	1	2	4	9	17
2001.....	101	2	-	-	-	1	2	1	2	5	9	5
2000.....	120	5	2	-	1	-	3	1	4	2	11	9
1999.....	131	-	3	1	-	2	2	4	2	9	5	12

Black, male

2004.....	52	1	-	-	-	1	1	1	1	2	4	3
2003.....	69	2	1	-	1	-	-	2	1	2	3	8
2002.....	79	2	2	-	-	-	-	1	2	3	6	12
2001.....	66	1	-	-	-	-	1	1	1	3	6	3
2000.....	76	2	1	-	1	-	1	1	2	1	9	8
1999.....	84	-	2	-	-	-	-	1	2	7	4	9

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Suppurative and necrotic conditions of lower respiratory tract (J85-J86)

Asian or Pacific Islander 2/, both sexes

2004.....	15	-	-	-	-	-	-	-	-	-	-	-
2003.....	18	-	-	-	-	1	1	1	-	-	1	1
2002.....	13	-	-	-	-	-	-	-	-	1	1	2
2001.....	15	-	-	-	-	-	-	1	-	1	-	-
2000.....	7	-	-	-	-	-	-	-	-	-	-	-
1999.....	21	-	-	-	-	-	1	-	-	-	3	1

Asian or Pacific Islander 2/, male

2004.....	8	-	-	-	-	-	-	-	-	-	-	-
2003.....	10	-	-	-	-	-	-	1	-	-	1	1
2002.....	10	-	-	-	-	-	-	-	-	1	1	2
2001.....	11	-	-	-	-	-	-	1	-	-	-	-
2000.....	5	-	-	-	-	-	-	-	-	-	-	-
1999.....	13	-	-	-	-	-	-	-	-	-	3	-

Asian or Pacific Islander 2/, female

2004.....	7	-	-	-	-	-	-	-	-	-	-	-
2003.....	8	-	-	-	-	1	1	-	-	-	-	-
2002.....	3	-	-	-	-	-	-	-	-	-	-	-
2001.....	4	-	-	-	-	-	-	-	-	1	-	-
2000.....	2	-	-	-	-	-	-	-	-	-	-	-
1999.....	8	-	-	-	-	-	1	-	-	-	-	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pleural effusion and plaque (J90,J92)												
All races, both sexes												
2004.....	969	3	2	3	-	1	1	1	-	2	2	9
2003.....	930	-	-	-	-	1	-	1	5	3	3	7
2002.....	970	2	-	1	-	-	-	1	-	6	4	10
2001.....	962	2	-	-	-	-	1	1	-	1	5	10
2000.....	870	2	2	2	-	1	-	1	4	6	11	10
1999.....	863	3	1	-	1	-	1	-	-	3	4	11
All races, male												
2004.....	441	1	-	2	-	-	1	-	-	2	-	4
2003.....	407	-	-	-	-	-	-	1	3	1	2	5
2002.....	402	1	-	1	-	-	-	-	-	4	3	4
2001.....	405	1	-	-	-	-	-	-	-	1	1	5
2000.....	388	1	1	2	-	1	-	-	3	5	7	8
1999.....	390	1	1	-	1	-	-	-	-	1	2	5
All races, female												
2004.....	528	2	2	1	-	1	-	1	-	-	2	5
2003.....	523	-	-	-	-	1	-	-	2	2	1	2
2002.....	568	1	-	-	-	-	-	1	-	2	1	6
2001.....	557	1	-	-	-	-	1	1	-	-	4	5
2000.....	482	1	1	-	-	-	-	1	1	1	4	2
1999.....	473	2	-	-	-	-	1	-	-	2	2	6
White, both sexes												
2004.....	872	2	2	2	-	1	1	1	-	2	2	7
2003.....	829	-	-	-	-	1	-	1	3	1	2	3
2002.....	875	2	-	1	-	-	-	1	-	2	4	9
2001.....	845	1	-	-	-	-	1	1	-	-	2	8
2000.....	773	2	1	-	-	1	-	1	2	4	9	5
1999.....	774	3	1	-	1	-	1	-	-	3	3	8

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of respiratory system (J80-J84,J93-J98)												
All races, both sexes												
2004.....	24,997	239	73	20	26	38	55	58	85	161	292	485
2003.....	23,142	249	63	18	23	44	54	61	111	158	252	481
2002.....	22,701	246	61	28	30	41	60	64	96	163	286	433
2001.....	23,721	264	65	22	37	39	70	79	98	180	292	462
2000.....	23,221	244	60	29	33	52	59	78	125	200	312	449
1999.....	22,618	220	60	31	29	46	44	74	133	180	291	438
All races, male												
2004.....	12,241	124	47	12	17	16	26	27	40	82	140	240
2003.....	11,181	146	35	12	13	26	35	34	54	78	120	244
2002.....	10,906	150	41	13	14	21	29	29	44	80	149	206
2001.....	11,292	167	32	14	20	20	38	33	46	85	158	248
2000.....	11,215	138	34	17	15	25	37	35	61	109	157	230
1999.....	10,853	129	35	16	14	29	26	34	63	92	149	216
All races, female												
2004.....	12,756	115	26	8	9	22	29	31	45	79	152	245
2003.....	11,961	103	28	6	10	18	19	27	57	80	132	237
2002.....	11,795	96	20	15	16	20	31	35	52	83	137	227
2001.....	12,429	97	33	8	17	19	32	46	52	95	134	214
2000.....	12,006	106	26	12	18	27	22	43	64	91	155	219
1999.....	11,765	91	25	15	15	17	18	40	70	88	142	222
White, both sexes												
2004.....	22,257	134	45	11	18	29	42	41	55	102	211	328
2003.....	20,674	145	38	15	18	29	32	40	79	105	165	339
2002.....	20,185	133	39	18	19	29	36	45	66	113	211	305
2001.....	21,049	140	43	17	31	26	50	50	62	112	212	328
2000.....	20,689	147	44	21	28	34	42	58	88	129	236	319
1999.....	20,118	122	40	22	21	38	38	50	93	124	199	310

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

All other diseases of respiratory system (J80-J84,J93-J98)

White, male

2004.....	10,911	69	32	5	12	13	22	19	26	47	101	163
2003.....	9,993	78	20	11	9	14	22	23	40	55	85	186
2002.....	9,725	80	28	9	9	13	22	21	30	56	110	150
2001.....	10,083	86	20	10	18	12	28	19	29	60	127	180
2000.....	9,994	88	25	13	13	15	27	27	44	70	119	163
1999.....	9,715	73	22	11	11	25	24	30	46	67	108	154

White, female

2004.....	11,346	65	13	6	6	16	20	22	29	55	110	165
2003.....	10,681	67	18	4	9	15	10	17	39	50	80	153
2002.....	10,460	53	11	9	10	16	14	24	36	57	101	155
2001.....	10,966	54	23	7	13	14	22	31	33	52	85	148
2000.....	10,695	59	19	8	15	19	15	31	44	59	117	156
1999.....	10,403	49	18	11	10	13	14	20	47	57	91	156

Black, both sexes

2004.....	2,139	93	22	7	7	8	7	15	26	54	76	139
2003.....	1,945	93	19	1	2	13	21	19	29	45	77	125
2002.....	1,991	100	22	10	10	12	21	17	27	44	69	116
2001.....	2,174	108	17	4	6	11	19	25	34	61	76	122
2000.....	2,069	93	14	7	5	16	15	15	31	63	68	122
1999.....	2,043	88	18	7	8	6	5	22	38	52	84	114

Black, male

2004.....	1,028	50	11	6	5	3	2	7	13	32	37	72
2003.....	942	59	12	1	2	11	13	10	13	21	29	51
2002.....	926	61	13	4	4	8	6	6	14	20	38	50
2001.....	965	71	8	3	2	7	9	11	16	21	31	61
2000.....	969	47	8	4	2	9	8	7	14	36	33	65
1999.....	918	52	12	4	3	3	1	4	16	23	36	57

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

All other diseases of respiratory system (J80-J84,J93-J98)

Black, female

2004.....	1,111	43	11	1	2	5	5	8	13	22	39	67
2003.....	1,003	34	7	-	-	2	8	9	16	24	48	74
2002.....	1,065	39	9	6	6	4	15	11	13	24	31	66
2001.....	1,209	37	9	1	4	4	10	14	18	40	45	61
2000.....	1,100	46	6	3	3	7	7	8	17	27	35	57
1999.....	1,125	36	6	3	5	3	4	18	22	29	48	57

American Indian 1/, both sexes

2004.....	168	6	1	1	-	-	5	1	-	3	1	9
2003.....	154	2	-	1	1	2	-	-	-	4	3	8
2002.....	127	4	-	-	-	-	1	-	2	2	3	6
2001.....	161	5	2	-	-	1	1	2	1	4	3	6
2000.....	136	1	1	-	-	-	-	3	3	3	3	3
1999.....	135	2	1	-	-	1	-	2	-	-	3	4

American Indian 1/, male

2004.....	84	2	1	1	-	-	2	-	-	2	-	3
2003.....	71	2	-	-	1	1	-	-	-	1	2	3
2002.....	66	3	-	-	-	-	1	-	-	1	1	3
2001.....	75	3	2	-	-	-	1	2	1	3	-	3
2000.....	64	-	1	-	-	-	-	-	2	2	2	-
1999.....	65	2	1	-	-	1	-	-	-	-	1	2

American Indian 1/, female

2004.....	84	4	-	-	-	-	3	1	-	1	1	6
2003.....	83	-	-	1	-	1	-	-	-	3	1	5
2002.....	61	1	-	-	-	-	-	-	2	1	2	3
2001.....	86	2	-	-	-	1	-	-	-	1	3	3
2000.....	72	1	-	-	-	-	-	3	1	1	1	3
1999.....	70	-	-	-	-	-	-	2	-	-	2	2

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

All other diseases of respiratory system (J80-J84,J93-J98)

Asian or Pacific Islander 2/, both sexes

2004.....	433	6	5	1	1	1	1	1	4	2	4	9
2003.....	369	9	6	1	2	-	1	2	3	4	7	9
2002.....	398	9	-	-	1	-	2	2	1	4	3	6
2001.....	337	11	3	1	-	1	-	2	1	3	1	6
2000.....	327	3	1	1	-	2	2	2	3	5	5	5
1999.....	322	8	1	2	-	1	1	-	2	4	5	10

Asian or Pacific Islander 2/, male

2004.....	218	3	3	-	-	-	-	1	1	1	2	2
2003.....	175	7	3	-	1	-	-	1	1	1	4	4
2002.....	189	6	-	-	1	-	-	2	-	3	-	3
2001.....	169	7	2	1	-	1	-	1	-	1	-	4
2000.....	188	3	-	-	-	1	2	1	1	1	3	2
1999.....	155	2	-	1	-	-	1	-	1	2	4	3

Asian or Pacific Islander 2/, female

2004.....	215	3	2	1	1	1	1	-	3	1	2	7
2003.....	194	2	3	1	1	-	1	1	2	3	3	5
2002.....	209	3	-	-	-	-	2	-	1	1	3	3
2001.....	168	4	1	-	-	-	-	1	1	2	1	2
2000.....	139	-	1	1	-	1	-	1	2	4	2	3
1999.....	167	6	1	1	-	1	-	-	1	2	1	7

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
XI. Diseases of the digestive system (K00-K92)												
All races, both sexes												
2004.....	86,337	580	121	45	56	93	182	254	601	1,475	3,319	5,449
2003.....	87,764	551	130	58	64	106	156	299	654	1,562	3,528	5,551
2002.....	86,710	509	129	48	68	86	175	262	674	1,605	3,664	5,611
2001.....	85,833	478	118	56	72	74	142	271	666	1,812	3,469	5,567
2000.....	84,015	499	107	58	65	72	183	275	680	1,815	3,564	5,192
1999.....	82,876	500	115	55	79	94	138	256	718	1,814	3,506	4,978
All races, male												
2004.....	43,347	325	74	26	22	43	98	159	402	948	2,153	3,649
2003.....	44,264	339	73	34	36	64	90	175	421	963	2,267	3,799
2002.....	43,440	312	74	33	31	42	96	142	437	995	2,362	3,884
2001.....	42,739	276	72	31	40	42	78	158	416	1,141	2,270	3,836
2000.....	41,867	288	70	32	29	42	108	179	424	1,157	2,388	3,676
1999.....	41,541	321	56	25	37	54	69	150	443	1,135	2,416	3,581
All races, female												
2004.....	42,990	255	47	19	34	50	84	95	199	527	1,166	1,800
2003.....	43,500	212	57	24	28	42	66	124	233	599	1,261	1,752
2002.....	43,270	197	55	15	37	44	79	120	237	610	1,302	1,727
2001.....	43,094	202	46	25	32	32	64	113	250	671	1,199	1,731
2000.....	42,148	211	37	26	36	30	75	96	256	658	1,176	1,516
1999.....	41,335	179	59	30	42	40	69	106	275	679	1,090	1,397
White, both sexes												
2004.....	74,952	340	76	34	36	65	113	169	459	1,143	2,690	4,435
2003.....	76,317	343	92	39	48	75	102	197	512	1,228	2,872	4,533
2002.....	75,673	315	86	33	49	63	122	177	497	1,277	3,002	4,525
2001.....	74,518	274	80	43	47	52	86	183	504	1,423	2,754	4,489
2000.....	72,862	297	56	37	50	52	128	180	503	1,401	2,834	4,159
1999.....	71,944	300	81	39	59	71	86	188	548	1,427	2,747	3,897

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
XI. Diseases of the digestive system (K00-K92)												
White, male												
2004.....	37,292	186	45	20	17	31	64	109	310	734	1,799	3,001
2003.....	38,141	221	57	24	24	48	61	117	336	776	1,885	3,172
2002.....	37,601	194	46	22	25	31	75	94	342	807	1,983	3,190
2001.....	36,633	150	52	22	23	31	51	106	320	897	1,822	3,146
2000.....	35,987	170	35	22	24	30	74	130	328	914	1,933	3,033
1999.....	35,738	191	33	16	28	42	45	114	342	917	1,938	2,850
White, female												
2004.....	37,660	154	31	14	19	34	49	60	149	409	891	1,434
2003.....	38,176	122	35	15	24	27	41	80	176	452	987	1,361
2002.....	38,072	121	40	11	24	32	47	83	155	470	1,019	1,335
2001.....	37,885	124	28	21	24	21	35	77	184	526	932	1,343
2000.....	36,875	127	21	15	26	22	54	50	175	487	901	1,126
1999.....	36,206	109	48	23	31	29	41	74	206	510	809	1,047
Black, both sexes												
2004.....	9,137	209	36	8	18	23	53	62	99	241	482	832
2003.....	9,232	191	32	10	14	24	45	84	101	253	505	814
2002.....	8,929	175	36	14	18	20	45	65	121	231	511	895
2001.....	9,326	186	31	11	20	20	43	70	118	295	576	902
2000.....	9,235	182	38	16	12	17	46	79	130	320	599	877
1999.....	9,080	179	27	14	16	22	48	57	127	298	624	918
Black, male												
2004.....	4,871	122	23	4	5	10	26	34	66	150	268	536
2003.....	4,928	109	13	4	10	11	24	46	59	134	291	500
2002.....	4,672	108	22	10	6	9	17	36	65	124	285	565
2001.....	5,026	112	15	7	13	10	18	42	69	182	358	587
2000.....	4,829	106	26	6	4	10	28	39	67	188	369	544
1999.....	4,775	112	16	7	7	11	21	29	75	169	388	628

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

XI. Diseases of the digestive system (K00-K92)

Black, female

2004.....	4,266	87	13	4	13	13	27	28	33	91	214	296
2003.....	4,304	82	19	6	4	13	21	38	42	119	214	314
2002.....	4,257	67	14	4	12	11	28	29	56	107	226	330
2001.....	4,300	74	16	4	7	10	25	28	49	113	218	315
2000.....	4,406	76	12	10	8	7	18	40	63	132	230	333
1999.....	4,305	67	11	7	9	11	27	28	52	129	236	290

American Indian 1/, both sexes

2004.....	1,021	5	3	1	1	2	8	10	26	65	108	130
2003.....	1,020	3	3	2	1	5	5	10	29	64	119	137
2002.....	945	3	2	-	1	2	2	12	43	73	109	127
2001.....	917	6	4	-	2	2	7	12	31	71	99	121
2000.....	871	6	5	-	2	2	4	12	33	71	91	98
1999.....	845	7	2	-	1	1	3	6	25	72	103	108

American Indian 1/, male

2004.....	514	5	3	1	-	1	4	6	12	41	56	72
2003.....	531	1	1	2	1	3	2	7	17	37	67	78
2002.....	525	1	2	-	-	1	1	8	22	47	61	80
2001.....	506	5	3	-	2	1	6	7	18	44	61	66
2000.....	460	4	2	-	1	2	2	7	21	36	57	58
1999.....	445	5	2	-	1	1	2	4	13	36	63	65

American Indian 1/, female

2004.....	507	-	-	-	1	1	4	4	14	24	52	58
2003.....	489	2	2	-	-	2	3	3	12	27	52	59
2002.....	420	2	-	-	1	1	1	4	21	26	48	47
2001.....	411	1	1	-	-	1	1	5	13	27	38	55
2000.....	411	2	3	-	1	-	2	5	12	35	34	40
1999.....	400	2	-	-	-	-	1	2	12	36	40	43

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

XI. Diseases of the digestive system (K00-K92)

Asian or Pacific Islander 2/, both sexes

2004.....	1,227	26	6	2	1	3	8	13	17	26	39	52
2003.....	1,195	14	3	7	1	2	4	8	12	17	32	67
2002.....	1,163	16	5	1	-	1	6	8	13	24	42	64
2001.....	1,072	12	3	2	3	-	6	6	13	23	40	55
2000.....	1,047	14	8	5	1	1	5	4	14	23	40	58
1999.....	1,007	14	5	2	3	-	1	5	18	17	32	55

Asian or Pacific Islander 2/, male

2004.....	670	12	3	1	-	1	4	10	14	23	30	40
2003.....	664	8	2	4	1	2	3	5	9	16	24	49
2002.....	642	9	4	1	-	1	3	4	8	17	33	49
2001.....	574	9	2	2	2	-	3	3	9	18	29	37
2000.....	591	8	7	4	-	-	4	3	8	19	29	41
1999.....	583	13	5	2	1	-	1	3	13	13	27	38

Asian or Pacific Islander 2/, female

2004.....	557	14	3	1	1	2	4	3	3	3	9	12
2003.....	531	6	1	3	-	-	1	3	3	1	8	18
2002.....	521	7	1	-	-	-	3	4	5	7	9	15
2001.....	498	3	1	-	1	-	3	3	4	5	11	18
2000.....	456	6	1	1	1	1	1	1	6	4	11	17
1999.....	424	1	-	-	2	-	-	2	5	4	5	17

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of oral cavity, salivary glands and jaws (K00-K14)												
All races, both sexes												
2004.....	230	6	-	-	-	-	1	6	2	3	1	7
2003.....	141	2	1	-	-	1	3	3	-	2	7	3
2002.....	163	3	2	1	-	-	3	-	3	4	2	6
2001.....	134	2	1	1	-	-	1	4	1	-	3	4
2000.....	112	3	-	-	-	-	-	1	4	1	4	3
1999.....	121	2	-	-	-	-	2	2	3	3	6	7
All races, male												
2004.....	86	2	-	-	-	-	-	2	2	2	1	4
2003.....	53	1	-	-	-	1	2	1	-	1	4	2
2002.....	74	2	2	1	-	-	1	-	2	3	2	4
2001.....	60	1	1	1	-	-	1	3	1	-	2	4
2000.....	43	-	-	-	-	-	-	1	2	-	3	3
1999.....	51	2	-	-	-	-	1	1	2	1	5	2
All races, female												
2004.....	144	4	-	-	-	-	1	4	-	1	-	3
2003.....	88	1	1	-	-	-	1	2	-	1	3	1
2002.....	89	1	-	-	-	-	2	-	1	1	-	2
2001.....	74	1	-	-	-	-	-	1	-	-	1	-
2000.....	69	3	-	-	-	-	-	-	2	1	1	-
1999.....	70	-	-	-	-	-	1	1	1	2	1	5
White, both sexes												
2004.....	205	6	-	-	-	-	-	2	1	1	1	5
2003.....	129	1	1	-	-	1	2	1	-	2	6	3
2002.....	145	3	2	-	-	-	1	-	1	3	2	4
2001.....	120	2	1	-	-	-	1	3	1	-	3	4
2000.....	100	2	-	-	-	-	-	1	2	-	4	3
1999.....	106	2	-	-	-	-	2	2	3	2	4	6

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of other parts of digestive system (K20-K92)												
All races, both sexes												
2004.....	86,107	574	121	45	56	93	181	248	599	1,472	3,318	5,442
2003.....	87,623	549	129	58	64	105	153	296	654	1,560	3,521	5,548
2002.....	86,547	506	127	47	68	86	172	262	671	1,601	3,662	5,605
2001.....	85,699	476	117	55	72	74	141	267	665	1,812	3,466	5,563
2000.....	83,903	496	107	58	65	72	183	274	676	1,814	3,560	5,189
1999.....	82,755	498	115	55	79	94	136	254	715	1,811	3,500	4,971
All races, male												
2004.....	43,261	323	74	26	22	43	98	157	400	946	2,152	3,645
2003.....	44,211	338	73	34	36	63	88	174	421	962	2,263	3,797
2002.....	43,366	310	72	32	31	42	95	142	435	992	2,360	3,880
2001.....	42,679	275	71	30	40	42	77	155	415	1,141	2,268	3,832
2000.....	41,824	288	70	32	29	42	108	178	422	1,157	2,385	3,673
1999.....	41,490	319	56	25	37	54	68	149	441	1,134	2,411	3,579
All races, female												
2004.....	42,846	251	47	19	34	50	83	91	199	526	1,166	1,797
2003.....	43,412	211	56	24	28	42	65	122	233	598	1,258	1,751
2002.....	43,181	196	55	15	37	44	77	120	236	609	1,302	1,725
2001.....	43,020	201	46	25	32	32	64	112	250	671	1,198	1,731
2000.....	42,079	208	37	26	36	30	75	96	254	657	1,175	1,516
1999.....	41,265	179	59	30	42	40	68	105	274	677	1,089	1,392
White, both sexes												
2004.....	74,747	334	76	34	36	65	113	167	458	1,142	2,689	4,430
2003.....	76,188	342	91	39	48	74	100	196	512	1,226	2,866	4,530
2002.....	75,528	312	84	33	49	63	121	177	496	1,274	3,000	4,521
2001.....	74,398	272	79	43	47	52	85	180	503	1,423	2,751	4,485
2000.....	72,762	295	56	37	50	52	128	179	501	1,401	2,830	4,156
1999.....	71,838	298	81	39	59	71	84	186	545	1,425	2,743	3,891

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	-------------	-----------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Diseases of other parts of digestive system (K20-K92)

White, male

2004.....	37,217	184	45	20	17	31	64	108	309	733	1,798	2,999
2003.....	38,094	220	57	24	24	47	60	117	336	775	1,882	3,170
2002.....	37,537	192	44	22	25	31	74	94	341	805	1,981	3,188
2001.....	36,581	149	51	22	23	31	50	104	319	897	1,820	3,142
2000.....	35,946	170	35	22	24	30	74	129	327	914	1,930	3,030
1999.....	35,693	189	33	16	28	42	44	113	340	916	1,935	2,848

White, female

2004.....	37,530	150	31	14	19	34	49	59	149	409	891	1,431
2003.....	38,094	122	34	15	24	27	40	79	176	451	984	1,360
2002.....	37,991	120	40	11	24	32	47	83	155	469	1,019	1,333
2001.....	37,817	123	28	21	24	21	35	76	184	526	931	1,343
2000.....	36,816	125	21	15	26	22	54	50	174	487	900	1,126
1999.....	36,145	109	48	23	31	29	40	73	205	509	808	1,043

Black, both sexes

2004.....	9,116	209	36	8	18	23	52	59	98	239	482	830
2003.....	9,222	190	32	10	14	24	44	82	101	253	504	814
2002.....	8,914	175	36	14	18	20	43	65	119	230	511	893
2001.....	9,314	186	31	11	20	20	43	69	118	295	576	902
2000.....	9,225	181	38	16	12	17	46	79	128	319	599	877
1999.....	9,068	179	27	14	16	22	48	57	127	297	622	917

Black, male

2004.....	4,861	122	23	4	5	10	26	33	65	149	268	534
2003.....	4,923	109	13	4	10	11	23	45	59	134	290	500
2002.....	4,665	108	22	10	6	9	17	36	64	123	285	563
2001.....	5,020	112	15	7	13	10	18	41	69	182	358	587
2000.....	4,827	106	26	6	4	10	28	39	66	188	369	544
1999.....	4,771	112	16	7	7	11	21	29	75	169	386	628

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Diseases of other parts of digestive system (K20-K92)

Black, female

2004.....	4,255	87	13	4	13	13	26	26	33	90	214	296
2003.....	4,299	81	19	6	4	13	21	37	42	119	214	314
2002.....	4,249	67	14	4	12	11	26	29	55	107	226	330
2001.....	4,294	74	16	4	7	10	25	28	49	113	218	315
2000.....	4,398	75	12	10	8	7	18	40	62	131	230	333
1999.....	4,297	67	11	7	9	11	27	28	52	128	236	289

American Indian 1/, both sexes

2004.....	1,020	5	3	1	1	2	8	10	26	65	108	130
2003.....	1,020	3	3	2	1	5	5	10	29	64	119	137
2002.....	944	3	2	-	1	2	2	12	43	73	109	127
2001.....	917	6	4	-	2	2	7	12	31	71	99	121
2000.....	871	6	5	-	2	2	4	12	33	71	91	98
1999.....	845	7	2	-	1	1	3	6	25	72	103	108

American Indian 1/, male

2004.....	513	5	3	1	-	1	4	6	12	41	56	72
2003.....	531	1	1	2	1	3	2	7	17	37	67	78
2002.....	524	1	2	-	-	1	1	8	22	47	61	80
2001.....	506	5	3	-	2	1	6	7	18	44	61	66
2000.....	460	4	2	-	1	2	2	7	21	36	57	58
1999.....	445	5	2	-	1	1	2	4	13	36	63	65

American Indian 1/, female

2004.....	507	-	-	-	1	1	4	4	14	24	52	58
2003.....	489	2	2	-	-	2	3	3	12	27	52	59
2002.....	420	2	-	-	1	1	1	4	21	26	48	47
2001.....	411	1	1	-	-	1	1	5	13	27	38	55
2000.....	411	2	3	-	1	-	2	5	12	35	34	40
1999.....	400	2	-	-	-	-	1	2	12	36	40	43

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Diseases of other parts of digestive system (K20-K92)

Asian or Pacific Islander 2/, both sexes

2004.....	1,224	26	6	2	1	3	8	12	17	26	39	52
2003.....	1,193	14	3	7	1	2	4	8	12	17	32	67
2002.....	1,161	16	5	-	-	1	6	8	13	24	42	64
2001.....	1,070	12	3	1	3	-	6	6	13	23	40	55
2000.....	1,045	14	8	5	1	1	5	4	14	23	40	58
1999.....	1,004	14	5	2	3	-	1	5	18	17	32	55

Asian or Pacific Islander 2/, male

2004.....	670	12	3	1	-	1	4	10	14	23	30	40
2003.....	663	8	2	4	1	2	3	5	9	16	24	49
2002.....	640	9	4	-	-	1	3	4	8	17	33	49
2001.....	572	9	2	1	2	-	3	3	9	18	29	37
2000.....	591	8	7	4	-	-	4	3	8	19	29	41
1999.....	581	13	5	2	1	-	1	3	13	13	27	38

Asian or Pacific Islander 2/, female

2004.....	554	14	3	1	1	2	4	2	3	3	9	12
2003.....	530	6	1	3	-	-	1	3	3	1	8	18
2002.....	521	7	1	-	-	-	3	4	5	7	9	15
2001.....	498	3	1	-	1	-	3	3	4	5	11	18
2000.....	454	6	1	1	1	1	1	1	6	4	11	17
1999.....	423	1	-	-	2	-	-	2	5	4	5	17

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Diseases of esophagus (K20-K22)

All races, both sexes

2004.....	1,912	13	3	2	1	3	4	6	2	14	30	37
2003.....	2,022	16	10	5	6	4	4	7	10	23	25	31
2002.....	2,122	13	11	1	5	-	6	5	18	12	35	56
2001.....	2,112	18	6	3	11	5	-	9	4	19	19	28
2000.....	2,210	20	7	4	1	4	6	10	8	13	32	38
1999.....	2,160	13	8	2	1	6	2	8	13	15	22	38

All races, male

2004.....	856	8	3	1	-	2	3	3	1	11	20	25
2003.....	944	8	6	2	6	2	1	7	6	15	19	23
2002.....	978	5	5	1	1	-	5	2	14	7	27	39
2001.....	924	8	3	-	3	4	-	6	2	13	13	19
2000.....	1,034	13	5	4	-	3	1	8	6	12	20	27
1999.....	1,001	9	4	-	1	3	1	4	8	13	19	26

All races, female

2004.....	1,056	5	-	1	1	1	1	3	1	3	10	12
2003.....	1,078	8	4	3	-	2	3	-	4	8	6	8
2002.....	1,144	8	6	-	4	-	1	3	4	5	8	17
2001.....	1,188	10	3	3	8	1	-	3	2	6	6	9
2000.....	1,176	7	2	-	1	1	5	2	2	1	12	11
1999.....	1,159	4	4	2	-	3	1	4	5	2	3	12

White, both sexes

2004.....	1,730	11	1	2	-	-	4	4	1	12	25	26
2003.....	1,848	12	5	4	5	3	3	4	10	17	23	27
2002.....	1,923	8	8	1	5	-	4	5	14	8	30	45
2001.....	1,928	9	2	2	8	4	-	8	3	16	16	25
2000.....	2,022	18	5	2	1	4	5	7	7	10	23	30
1999.....	1,968	10	4	2	1	5	2	6	9	12	14	28

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Diseases of esophagus (K20-K22)

White, male

2004.....	760	7	1	1	-	-	3	2	1	9	16	17
2003.....	856	6	2	2	5	2	-	4	6	12	17	20
2002.....	876	3	4	1	1	-	3	2	13	5	24	31
2001.....	827	5	2	-	3	3	-	5	1	11	11	17
2000.....	923	12	4	2	-	3	1	6	5	9	13	21
1999.....	890	7	-	-	1	2	1	2	5	10	12	18

White, female

2004.....	970	4	-	1	-	-	1	2	-	3	9	9
2003.....	992	6	3	2	-	1	3	-	4	5	6	7
2002.....	1,047	5	4	-	4	-	1	3	1	3	6	14
2001.....	1,101	4	-	2	5	1	-	3	2	5	5	8
2000.....	1,099	6	1	-	1	1	4	1	2	1	10	9
1999.....	1,078	3	4	2	-	3	1	4	4	2	2	10

Black, both sexes

2004.....	154	1	2	-	1	2	-	2	1	2	3	10
2003.....	143	3	3	1	1	1	1	3	-	6	1	3
2002.....	164	4	3	-	-	-	1	-	4	3	4	10
2001.....	156	8	4	1	3	1	-	1	1	3	2	3
2000.....	168	2	2	-	-	-	-	2	1	3	9	7
1999.....	166	3	2	-	-	1	-	2	4	3	8	8

Black, male

2004.....	82	-	2	-	-	1	-	1	-	2	2	7
2003.....	70	1	2	-	1	-	1	3	-	3	1	3
2002.....	82	1	1	-	-	-	1	-	1	2	2	7
2001.....	80	2	1	-	-	1	-	1	1	2	1	2
2000.....	97	1	1	-	-	-	-	1	1	3	7	5
1999.....	94	2	2	-	-	1	-	2	3	3	7	6

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Peptic ulcer (K25-K28)												
All races, both sexes												
2004.....	3,682	2	3	1	1	4	6	7	10	28	62	119
2003.....	3,913	2	1	-	3	3	4	8	17	33	68	122
2002.....	4,079	1	1	-	1	-	-	5	17	40	73	118
2001.....	4,491	-	1	3	2	3	3	13	19	36	82	137
2000.....	4,558	-	-	-	-	4	5	3	11	42	71	130
1999.....	4,604	2	2	-	-	1	1	7	14	38	85	113
All races, male												
2004.....	1,690	1	2	1	-	1	5	6	8	19	38	79
2003.....	1,880	-	1	-	1	3	3	4	9	20	40	82
2002.....	1,951	-	1	-	1	-	-	2	12	28	44	71
2001.....	2,074	-	1	2	-	2	1	10	11	25	62	91
2000.....	2,201	-	-	-	-	3	4	1	5	31	48	86
1999.....	2,189	-	-	-	-	1	1	5	7	27	49	67
All races, female												
2004.....	1,992	1	1	-	1	3	1	1	2	9	24	40
2003.....	2,033	2	-	-	2	-	1	4	8	13	28	40
2002.....	2,128	1	-	-	-	-	-	3	5	12	29	47
2001.....	2,417	-	-	1	2	1	2	3	8	11	20	46
2000.....	2,357	-	-	-	-	1	1	2	6	11	23	44
1999.....	2,415	2	2	-	-	-	-	2	7	11	36	46
White, both sexes												
2004.....	3,215	1	1	1	-	4	5	4	6	17	45	80
2003.....	3,408	1	1	-	2	2	2	6	14	22	50	90
2002.....	3,587	1	1	-	-	-	-	2	10	30	47	94
2001.....	3,942	-	1	3	1	2	1	5	11	19	59	102
2000.....	4,021	-	-	-	-	4	5	3	6	31	50	95
1999.....	4,040	1	2	-	-	1	1	6	9	21	56	80

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Peptic ulcer (K25-K28)

White, male

2004.....	1,450	-	1	1	-	1	4	4	4	10	30	52
2003.....	1,592	-	1	-	1	2	1	4	6	13	29	62
2002.....	1,685	-	1	-	-	-	-	2	8	20	30	56
2001.....	1,751	-	1	2	-	1	1	4	7	12	44	65
2000.....	1,900	-	-	-	-	3	4	1	3	22	34	68
1999.....	1,887	-	-	-	-	1	1	5	3	16	28	45

White, female

2004.....	1,765	1	-	-	-	3	1	-	2	7	15	28
2003.....	1,816	1	-	-	1	-	1	2	8	9	21	28
2002.....	1,902	1	-	-	-	-	-	-	2	10	17	38
2001.....	2,191	-	-	1	1	1	-	1	4	7	15	37
2000.....	2,121	-	-	-	-	1	1	2	3	9	16	27
1999.....	2,153	1	2	-	-	-	-	1	6	5	28	35

Black, both sexes

2004.....	366	-	1	-	1	-	1	3	3	11	17	34
2003.....	383	1	-	-	1	1	-	2	3	7	16	32
2002.....	380	-	-	-	1	-	-	3	7	9	26	23
2001.....	409	-	-	-	1	1	1	8	7	16	19	27
2000.....	425	-	-	-	-	-	-	-	4	10	19	32
1999.....	448	1	-	-	-	-	-	-	4	16	28	28

Black, male

2004.....	191	-	1	-	-	-	1	2	3	9	8	26
2003.....	223	-	-	-	-	1	-	-	3	3	10	20
2002.....	212	-	-	-	1	-	-	-	4	7	14	14
2001.....	243	-	-	-	-	1	-	6	4	13	14	21
2000.....	229	-	-	-	-	-	-	-	2	8	13	17
1999.....	243	-	-	-	-	-	-	-	3	11	20	20

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Peptic ulcer (K25-K28)												
Black, female												
2004.....	175	-	-	-	1	-	-	1	-	2	9	8
2003.....	160	1	-	-	1	-	-	2	-	4	6	12
2002.....	168	-	-	-	-	-	-	3	3	2	12	9
2001.....	166	-	-	-	1	-	1	2	3	3	5	6
2000.....	196	-	-	-	-	-	-	-	2	2	6	15
1999.....	205	1	-	-	-	-	-	-	1	5	8	8
American Indian 1/, both sexes												
2004.....	13	-	-	-	-	-	-	-	-	-	-	3
2003.....	16	-	-	-	-	-	1	-	-	1	2	-
2002.....	17	-	-	-	-	-	-	-	-	-	-	-
2001.....	28	-	-	-	-	-	-	-	1	1	1	4
2000.....	19	-	-	-	-	-	-	-	-	1	-	2
1999.....	10	-	-	-	-	-	-	-	-	1	1	1
American Indian 1/, male												
2004.....	2	-	-	-	-	-	-	-	-	-	-	-
2003.....	7	-	-	-	-	-	1	-	-	1	1	-
2002.....	10	-	-	-	-	-	-	-	-	-	-	-
2001.....	17	-	-	-	-	-	-	-	-	-	1	3
2000.....	11	-	-	-	-	-	-	-	-	1	-	-
1999.....	5	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, female												
2004.....	11	-	-	-	-	-	-	-	-	-	-	3
2003.....	9	-	-	-	-	-	-	-	-	-	1	-
2002.....	7	-	-	-	-	-	-	-	-	-	-	-
2001.....	11	-	-	-	-	-	-	-	1	1	-	1
2000.....	8	-	-	-	-	-	-	-	-	-	-	2
1999.....	5	-	-	-	-	-	-	-	-	1	-	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Peptic ulcer (K25-K28)												
Asian or Pacific Islander 2/, both sexes												
2004.....	88	1	1	-	-	-	-	-	1	-	-	2
2003.....	106	-	-	-	-	-	1	-	-	3	-	-
2002.....	95	-	-	-	-	-	-	-	-	1	-	1
2001.....	112	-	-	-	-	-	1	-	-	-	3	4
2000.....	93	-	-	-	-	-	-	-	1	-	2	1
1999.....	106	-	-	-	-	-	-	1	1	-	-	4
Asian or Pacific Islander 2/, male												
2004.....	47	1	-	-	-	-	-	-	1	-	-	1
2003.....	58	-	-	-	-	-	1	-	-	3	-	-
2002.....	44	-	-	-	-	-	-	-	-	1	-	1
2001.....	63	-	-	-	-	-	-	-	-	-	3	2
2000.....	61	-	-	-	-	-	-	-	-	-	1	1
1999.....	54	-	-	-	-	-	-	-	1	-	-	2
Asian or Pacific Islander 2/, female												
2004.....	41	-	1	-	-	-	-	-	-	-	-	1
2003.....	48	-	-	-	-	-	-	-	-	-	-	-
2002.....	51	-	-	-	-	-	-	-	-	-	-	-
2001.....	49	-	-	-	-	-	1	-	-	-	-	2
2000.....	32	-	-	-	-	-	-	-	1	-	1	-
1999.....	52	-	-	-	-	-	-	1	-	-	-	2

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Gastric ulcer (K25)

All races, both sexes

2004.....	1,129	1	1	1	1	-	2	4	5	15	21	51
2003.....	1,154	1	1	-	3	-	1	4	5	9	24	42
2002.....	1,200	1	-	-	-	-	-	3	3	13	30	43
2001.....	1,277	-	1	3	-	-	1	7	8	11	24	52
2000.....	1,322	-	-	-	-	4	4	1	3	11	28	52
1999.....	1,333	-	1	-	-	-	-	1	4	16	37	45

All races, male

2004.....	493	-	-	1	-	-	2	3	4	9	12	32
2003.....	498	-	1	-	1	-	1	2	2	5	13	27
2002.....	547	-	-	-	-	-	-	1	2	9	19	26
2001.....	545	-	1	2	-	-	-	5	4	8	20	34
2000.....	598	-	-	-	-	3	3	-	1	9	16	35
1999.....	589	-	-	-	-	-	-	1	2	9	21	28

All races, female

2004.....	636	1	1	-	1	-	-	1	1	6	9	19
2003.....	656	1	-	-	2	-	-	2	3	4	11	15
2002.....	653	1	-	-	-	-	-	2	1	4	11	17
2001.....	732	-	-	1	-	-	1	2	4	3	4	18
2000.....	724	-	-	-	-	1	1	1	2	2	12	17
1999.....	744	-	1	-	-	-	-	-	2	7	16	17

White, both sexes

2004.....	952	1	-	1	-	-	1	1	3	10	14	30
2003.....	978	-	1	-	2	-	1	3	5	7	18	29
2002.....	1,022	1	-	-	-	-	-	1	3	10	17	31
2001.....	1,073	-	1	3	-	-	-	2	4	6	14	35
2000.....	1,123	-	-	-	-	4	4	1	1	7	19	33
1999.....	1,127	-	1	-	-	-	-	1	3	9	22	33

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gastric ulcer (K25)												
White, male												
2004.....	407	-	-	1	-	-	1	1	2	6	8	18
2003.....	400	-	1	-	1	-	1	2	2	3	8	20
2002.....	449	-	-	-	-	-	-	1	2	7	12	17
2001.....	431	-	1	2	-	-	-	2	2	3	11	22
2000.....	479	-	-	-	-	3	3	-	-	5	11	25
1999.....	472	-	-	-	-	-	-	1	1	6	10	20
White, female												
2004.....	545	1	-	-	-	-	-	-	1	4	6	12
2003.....	578	-	-	-	1	-	-	1	3	4	10	9
2002.....	573	1	-	-	-	-	-	-	1	3	5	14
2001.....	642	-	-	1	-	-	-	-	2	3	3	13
2000.....	644	-	-	-	-	1	1	1	1	2	8	8
1999.....	655	-	1	-	-	-	-	-	2	3	12	13
Black, both sexes												
2004.....	139	-	-	-	1	-	1	3	1	5	7	17
2003.....	145	1	-	-	1	-	-	1	-	2	5	13
2002.....	135	-	-	-	-	-	-	2	-	3	13	11
2001.....	150	-	-	-	-	-	1	5	3	5	9	13
2000.....	165	-	-	-	-	-	-	-	1	4	8	18
1999.....	173	-	-	-	-	-	-	-	1	6	14	10
Black, male												
2004.....	71	-	-	-	-	-	1	2	1	3	4	13
2003.....	79	-	-	-	-	-	-	-	-	2	4	7
2002.....	73	-	-	-	-	-	-	-	-	2	7	8
2001.....	87	-	-	-	-	-	-	3	2	5	8	10
2000.....	101	-	-	-	-	-	-	-	1	4	5	9
1999.....	101	-	-	-	-	-	-	-	1	3	10	8

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Gastric ulcer (K25)

Black, female

2004.....	68	-	-	-	1	-	-	1	-	2	3	4
2003.....	66	1	-	-	1	-	-	1	-	-	1	6
2002.....	62	-	-	-	-	-	-	2	-	1	6	3
2001.....	63	-	-	-	-	-	1	2	1	-	1	3
2000.....	64	-	-	-	-	-	-	-	-	-	3	9
1999.....	72	-	-	-	-	-	-	-	-	3	4	2

American Indian 1/, both sexes

2004.....	6	-	-	-	-	-	-	-	-	-	-	2
2003.....	7	-	-	-	-	-	-	-	-	-	1	-
2002.....	10	-	-	-	-	-	-	-	-	-	-	-
2001.....	12	-	-	-	-	-	-	-	1	-	-	2
2000.....	6	-	-	-	-	-	-	-	-	-	-	-
1999.....	5	-	-	-	-	-	-	-	-	1	1	1

American Indian 1/, male

2004.....	1	-	-	-	-	-	-	-	-	-	-	-
2003.....	4	-	-	-	-	-	-	-	-	-	1	-
2002.....	6	-	-	-	-	-	-	-	-	-	-	-
2001.....	6	-	-	-	-	-	-	-	-	-	-	1
2000.....	3	-	-	-	-	-	-	-	-	-	-	-
1999.....	2	-	-	-	-	-	-	-	-	-	1	-

American Indian 1/, female

2004.....	5	-	-	-	-	-	-	-	-	-	-	2
2003.....	3	-	-	-	-	-	-	-	-	-	-	-
2002.....	4	-	-	-	-	-	-	-	-	-	-	-
2001.....	6	-	-	-	-	-	-	-	1	-	-	1
2000.....	3	-	-	-	-	-	-	-	-	-	-	-
1999.....	3	-	-	-	-	-	-	-	-	1	-	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Duodenal ulcer (K26)												
All races, both sexes												
2004.....	1,264	1	2	-	-	2	3	3	3	7	26	42
2003.....	1,324	1	-	-	-	-	2	4	8	11	26	55
2002.....	1,426	-	-	-	-	-	-	2	10	19	30	48
2001.....	1,544	-	-	-	1	3	-	2	7	14	40	55
2000.....	1,518	-	-	-	-	-	1	2	3	21	27	41
1999.....	1,476	2	1	-	-	1	1	5	7	13	33	43
All races, male												
2004.....	642	1	2	-	-	1	2	3	2	6	18	33
2003.....	698	-	-	-	-	-	2	2	5	7	17	41
2002.....	718	-	-	-	-	-	-	1	8	14	18	27
2001.....	782	-	-	-	-	2	-	1	5	9	31	40
2000.....	775	-	-	-	-	-	1	1	2	16	20	29
1999.....	793	-	-	-	-	1	1	4	4	11	19	26
All races, female												
2004.....	622	-	-	-	-	1	1	-	1	1	8	9
2003.....	626	1	-	-	-	-	-	2	3	4	9	14
2002.....	708	-	-	-	-	-	-	1	2	5	12	21
2001.....	762	-	-	-	1	1	-	1	2	5	9	15
2000.....	743	-	-	-	-	-	-	1	1	5	7	12
1999.....	683	2	1	-	-	-	-	1	3	2	14	17
White, both sexes												
2004.....	1,133	-	1	-	-	2	3	3	2	3	19	31
2003.....	1,161	1	-	-	-	-	-	3	6	7	17	44
2002.....	1,282	-	-	-	-	-	-	1	7	13	22	40
2001.....	1,387	-	-	-	-	2	-	1	5	8	30	40
2000.....	1,358	-	-	-	-	-	1	2	2	14	18	34
1999.....	1,313	1	1	-	-	1	1	4	4	7	25	28

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Duodenal ulcer (K26)												
White, male												
2004.....	563	-	1	-	-	1	2	3	1	2	14	24
2003.....	601	-	-	-	-	-	-	2	3	4	12	31
2002.....	644	-	-	-	-	-	-	1	6	9	14	25
2001.....	681	-	-	-	-	1	-	-	4	5	24	28
2000.....	684	-	-	-	-	-	1	1	2	11	13	25
1999.....	703	-	-	-	-	1	1	4	2	6	13	15
White, female												
2004.....	570	-	-	-	-	1	1	-	1	1	5	7
2003.....	560	1	-	-	-	-	-	1	3	3	5	13
2002.....	638	-	-	-	-	-	-	-	1	4	8	15
2001.....	706	-	-	-	-	1	-	1	1	3	6	12
2000.....	674	-	-	-	-	-	-	1	-	3	5	9
1999.....	610	1	1	-	-	-	-	-	2	1	12	13
Black, both sexes												
2004.....	101	-	1	-	-	-	-	-	1	4	7	10
2003.....	118	-	-	-	-	-	-	1	2	1	9	11
2002.....	115	-	-	-	-	-	-	1	3	5	8	8
2001.....	119	-	-	-	1	1	-	1	2	6	8	12
2000.....	122	-	-	-	-	-	-	-	1	6	8	7
1999.....	123	1	-	-	-	-	-	-	2	6	8	12
Black, male												
2004.....	62	-	1	-	-	-	-	-	1	4	4	9
2003.....	72	-	-	-	-	-	-	-	2	-	5	10
2002.....	65	-	-	-	-	-	-	-	2	4	4	2
2001.....	74	-	-	-	-	1	-	1	1	4	5	10
2000.....	67	-	-	-	-	-	-	-	-	4	6	4
1999.....	68	-	-	-	-	-	-	-	1	5	6	9

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Duodenal ulcer (K26)												
Asian or Pacific Islander 2/, both sexes												
2004.....	26	1	-	-	-	-	-	-	-	-	-	-
2003.....	41	-	-	-	-	-	1	-	-	3	-	-
2002.....	27	-	-	-	-	-	-	-	-	1	-	-
2001.....	32	-	-	-	-	-	-	-	-	-	2	2
2000.....	31	-	-	-	-	-	-	-	-	-	1	-
1999.....	40	-	-	-	-	-	-	1	1	-	-	3
Asian or Pacific Islander 2/, male												
2004.....	16	1	-	-	-	-	-	-	-	-	-	-
2003.....	23	-	-	-	-	-	1	-	-	3	-	-
2002.....	8	-	-	-	-	-	-	-	-	1	-	-
2001.....	22	-	-	-	-	-	-	-	-	-	2	1
2000.....	20	-	-	-	-	-	-	-	-	-	1	-
1999.....	22	-	-	-	-	-	-	-	1	-	-	2
Asian or Pacific Islander 2/, female												
2004.....	10	-	-	-	-	-	-	-	-	-	-	-
2003.....	18	-	-	-	-	-	-	-	-	-	-	-
2002.....	19	-	-	-	-	-	-	-	-	-	-	-
2001.....	10	-	-	-	-	-	-	-	-	-	-	1
2000.....	11	-	-	-	-	-	-	-	-	-	-	-
1999.....	18	-	-	-	-	-	-	1	-	-	-	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Peptic ulcer, site unspecified (K27)												
All races, both sexes												
2004.....	1,242	-	-	-	-	2	-	-	2	6	15	26
2003.....	1,379	-	-	-	-	3	1	-	4	12	15	23
2002.....	1,404	-	1	-	1	-	-	-	4	6	12	27
2001.....	1,631	-	-	-	1	-	2	4	4	11	17	29
2000.....	1,664	-	-	-	-	-	-	-	5	8	16	36
1999.....	1,758	-	-	-	-	-	-	1	3	9	14	23
All races, male												
2004.....	534	-	-	-	-	-	-	-	2	4	8	14
2003.....	653	-	-	-	-	3	-	-	2	7	9	13
2002.....	659	-	1	-	1	-	-	-	2	3	7	18
2001.....	729	-	-	-	-	-	1	4	2	8	10	17
2000.....	803	-	-	-	-	-	-	-	2	5	12	21
1999.....	789	-	-	-	-	-	-	-	1	7	9	13
All races, female												
2004.....	708	-	-	-	-	2	-	-	-	2	7	12
2003.....	726	-	-	-	-	-	1	-	2	5	6	10
2002.....	745	-	-	-	-	-	-	-	2	3	5	9
2001.....	902	-	-	-	1	-	1	-	2	3	7	12
2000.....	861	-	-	-	-	-	-	-	3	3	4	15
1999.....	969	-	-	-	-	-	-	1	2	2	5	10
White, both sexes												
2004.....	1,090	-	-	-	-	2	-	-	1	4	12	19
2003.....	1,218	-	-	-	-	2	1	-	3	7	12	15
2002.....	1,240	-	1	-	-	-	-	-	-	5	7	23
2001.....	1,445	-	-	-	1	-	1	2	2	5	14	26
2000.....	1,496	-	-	-	-	-	-	-	3	8	13	27
1999.....	1,570	-	-	-	-	-	-	1	2	5	8	17

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Peptic ulcer, site unspecified (K27)												
White, male												
2004.....	461	-	-	-	-	-	-	-	1	2	8	10
2003.....	564	-	-	-	-	2	-	-	1	5	8	10
2002.....	567	-	1	-	-	-	-	-	-	2	4	14
2001.....	622	-	-	-	-	-	1	2	1	4	8	15
2000.....	717	-	-	-	-	-	-	-	1	5	10	17
1999.....	698	-	-	-	-	-	-	-	-	4	5	10
White, female												
2004.....	629	-	-	-	-	2	-	-	-	2	4	9
2003.....	654	-	-	-	-	-	1	-	2	2	4	5
2002.....	673	-	-	-	-	-	-	-	-	3	3	9
2001.....	823	-	-	-	1	-	-	-	1	1	6	11
2000.....	779	-	-	-	-	-	-	-	2	3	3	10
1999.....	872	-	-	-	-	-	-	1	2	1	3	7
Black, both sexes												
2004.....	119	-	-	-	-	-	-	-	1	2	3	7
2003.....	117	-	-	-	-	1	-	-	1	4	2	8
2002.....	126	-	-	-	1	-	-	-	4	1	5	4
2001.....	139	-	-	-	-	-	-	2	2	5	2	2
2000.....	130	-	-	-	-	-	-	-	2	-	3	7
1999.....	150	-	-	-	-	-	-	-	1	4	6	6
Black, male												
2004.....	56	-	-	-	-	-	-	-	1	2	-	4
2003.....	70	-	-	-	-	1	-	-	1	1	1	3
2002.....	72	-	-	-	1	-	-	-	2	1	3	4
2001.....	82	-	-	-	-	-	-	2	1	4	1	1
2000.....	58	-	-	-	-	-	-	-	1	-	2	4
1999.....	74	-	-	-	-	-	-	-	1	3	4	3

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Gastritis and duodenitis (K29)

All races, both sexes

2004.....	368	-	-	1	-	3	-	1	1	9	9	14
2003.....	387	-	-	-	1	1	1	1	6	4	15	18
2002.....	365	-	1	1	-	-	1	3	5	3	13	11
2001.....	375	-	-	-	1	-	2	3	3	4	10	7
2000.....	394	-	-	-	-	-	2	-	3	7	11	12
1999.....	404	-	-	1	-	-	1	2	1	5	15	15

All races, male

2004.....	183	-	-	-	-	1	-	1	-	8	6	9
2003.....	191	-	-	-	1	1	1	1	5	1	11	16
2002.....	166	-	1	-	-	-	1	-	3	3	8	8
2001.....	161	-	-	-	1	-	1	-	2	2	5	3
2000.....	173	-	-	-	-	-	2	-	2	3	10	8
1999.....	179	-	-	1	-	-	-	2	-	3	10	12

All races, female

2004.....	185	-	-	1	-	2	-	-	1	1	3	5
2003.....	196	-	-	-	-	-	-	-	1	3	4	2
2002.....	199	-	-	1	-	-	-	3	2	-	5	3
2001.....	214	-	-	-	-	-	1	3	1	2	5	4
2000.....	221	-	-	-	-	-	-	-	1	4	1	4
1999.....	225	-	-	-	-	-	1	-	1	2	5	3

White, both sexes

2004.....	305	-	-	-	-	3	-	-	-	6	5	10
2003.....	326	-	-	-	-	1	1	-	5	1	13	12
2002.....	320	-	1	1	-	-	1	3	4	3	11	9
2001.....	334	-	-	-	-	-	1	2	3	3	10	5
2000.....	349	-	-	-	-	-	2	-	2	6	10	8
1999.....	342	-	-	-	-	-	1	1	1	3	12	11

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gastritis and duodenitis (K29)												
White, male												
2004.....	152	-	-	-	-	1	-	-	-	5	3	7
2003.....	158	-	-	-	-	1	1	-	4	-	10	12
2002.....	150	-	1	-	-	-	1	-	3	3	7	6
2001.....	141	-	-	-	-	-	1	-	2	2	5	1
2000.....	149	-	-	-	-	-	2	-	1	2	9	5
1999.....	145	-	-	-	-	-	-	1	-	2	8	9
White, female												
2004.....	153	-	-	-	-	2	-	-	-	1	2	3
2003.....	168	-	-	-	-	-	-	-	1	1	3	-
2002.....	170	-	-	1	-	-	-	3	1	-	4	3
2001.....	193	-	-	-	-	-	-	2	1	1	5	4
2000.....	200	-	-	-	-	-	-	-	1	4	1	3
1999.....	197	-	-	-	-	-	1	-	1	1	4	2
Black, both sexes												
2004.....	47	-	-	1	-	-	-	1	1	3	3	3
2003.....	46	-	-	-	-	-	-	1	1	3	2	6
2002.....	36	-	-	-	-	-	-	-	1	-	2	2
2001.....	34	-	-	-	1	-	1	1	-	1	-	2
2000.....	38	-	-	-	-	-	-	-	-	1	1	3
1999.....	53	-	-	1	-	-	-	1	-	2	2	4
Black, male												
2004.....	22	-	-	-	-	-	-	1	-	3	2	2
2003.....	24	-	-	-	-	-	-	1	1	1	1	4
2002.....	14	-	-	-	-	-	-	-	-	-	1	2
2001.....	18	-	-	-	1	-	-	-	-	-	-	2
2000.....	18	-	-	-	-	-	-	-	-	1	1	2
1999.....	28	-	-	1	-	-	-	1	-	1	1	3

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Dyspepsia and other diseases of stomach and duodenum (K30-K31)

All races, both sexes

2004.....	612	6	2	1	1	4	3	3	8	24	12	23
2003.....	607	9	2	2	1	4	3	3	7	8	21	17
2002.....	594	12	2	1	1	5	3	2	6	16	20	17
2001.....	542	2	1	-	3	2	5	2	7	8	17	17
2000.....	570	6	1	1	3	-	1	2	8	5	9	14
1999.....	557	4	-	-	2	1	2	5	4	9	16	21

All races, male

2004.....	265	3	1	-	-	2	2	1	3	10	6	7
2003.....	232	6	-	1	1	3	3	1	2	3	8	6
2002.....	232	9	2	-	-	3	-	-	2	6	7	8
2001.....	218	2	-	-	2	-	4	1	2	3	9	8
2000.....	238	3	1	-	1	-	-	2	4	1	2	5
1999.....	228	3	-	-	-	1	-	2	2	2	11	11

All races, female

2004.....	347	3	1	1	1	2	1	2	5	14	6	16
2003.....	375	3	2	1	-	1	-	2	5	5	13	11
2002.....	362	3	-	1	1	2	3	2	4	10	13	9
2001.....	324	-	1	-	1	2	1	1	5	5	8	9
2000.....	332	3	-	1	2	-	1	-	4	4	7	9
1999.....	329	1	-	-	2	-	2	3	2	7	5	10

White, both sexes

2004.....	525	5	2	-	1	4	3	1	7	21	10	18
2003.....	519	7	1	1	1	3	1	2	5	6	14	13
2002.....	509	5	2	-	1	5	2	1	5	15	13	14
2001.....	477	1	1	-	2	2	4	1	6	6	13	14
2000.....	506	3	-	1	2	-	1	2	8	4	7	12
1999.....	495	3	-	-	1	1	1	4	2	7	15	18

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Dyspepsia and other diseases of stomach and duodenum (K30-K31)

White, male

2004.....	217	3	1	-	-	2	2	-	3	8	4	4
2003.....	199	5	-	-	1	2	1	1	1	3	5	5
2002.....	200	4	2	-	-	3	-	-	2	6	6	7
2001.....	188	1	-	-	1	-	3	-	2	3	7	7
2000.....	206	1	-	-	1	-	-	2	4	1	1	5
1999.....	202	2	-	-	-	1	-	1	1	2	10	10

White, female

2004.....	308	2	1	-	1	2	1	1	4	13	6	14
2003.....	320	2	1	1	-	1	-	1	4	3	9	8
2002.....	309	1	-	-	1	2	2	1	3	9	7	7
2001.....	289	-	1	-	1	2	1	1	4	3	6	7
2000.....	300	2	-	1	1	-	1	-	4	3	6	7
1999.....	293	1	-	-	1	-	1	3	1	5	5	8

Black, both sexes

2004.....	84	-	-	1	-	-	-	2	1	3	2	5
2003.....	68	2	-	-	-	-	1	1	2	2	7	3
2002.....	73	7	-	1	-	-	1	1	1	1	7	3
2001.....	59	1	-	-	1	-	-	1	1	2	3	2
2000.....	58	3	1	-	1	-	-	-	-	1	2	2
1999.....	54	1	-	-	1	-	1	1	2	2	1	3

Black, male

2004.....	47	-	-	-	-	-	-	1	-	2	2	3
2003.....	22	1	-	-	-	-	1	-	1	-	3	-
2002.....	26	5	-	-	-	-	-	-	-	-	1	1
2001.....	27	1	-	-	1	-	-	1	-	-	1	1
2000.....	28	2	1	-	-	-	-	-	-	-	1	-
1999.....	20	1	-	-	-	-	-	1	1	-	1	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Diseases of appendix (K35-K38)

All races, both sexes

2004.....	452	3	4	7	6	3	5	7	2	9	5	14
2003.....	439	4	5	6	5	6	5	5	11	10	15	25
2002.....	480	3	-	3	4	7	5	4	6	8	19	24
2001.....	428	1	8	3	3	5	10	7	8	9	17	25
2000.....	435	3	5	4	11	5	6	3	8	11	16	18
1999.....	390	3	2	12	14	2	3	3	3	13	11	25

All races, male

2004.....	253	3	4	6	5	2	2	6	1	9	4	10
2003.....	258	2	2	2	3	4	5	2	7	5	11	14
2002.....	275	1	-	1	2	5	2	4	5	7	10	19
2001.....	252	-	7	3	2	3	5	5	6	7	10	17
2000.....	237	2	2	1	3	2	5	3	7	5	11	12
1999.....	236	1	2	6	9	2	3	3	2	10	7	18

All races, female

2004.....	199	-	-	1	1	1	3	1	1	-	1	4
2003.....	181	2	3	4	2	2	-	3	4	5	4	11
2002.....	205	2	-	2	2	2	3	-	1	1	9	5
2001.....	176	1	1	-	1	2	5	2	2	2	7	8
2000.....	198	1	3	3	8	3	1	-	1	6	5	6
1999.....	154	2	-	6	5	-	-	-	1	3	4	7

White, both sexes

2004.....	377	3	3	4	4	2	3	4	1	7	3	11
2003.....	355	3	3	4	2	4	3	3	7	6	8	19
2002.....	404	2	-	2	2	4	3	3	5	5	13	17
2001.....	342	1	5	2	2	5	5	3	5	7	10	14
2000.....	348	3	4	3	10	5	3	1	3	7	11	10
1999.....	314	3	-	8	10	1	2	3	1	10	9	16

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of appendix (K35-K38)												
White, male												
2004.....	210	3	3	4	4	1	2	3	-	7	2	8
2003.....	213	2	2	1	-	3	3	1	4	5	7	11
2002.....	225	-	-	1	1	2	1	3	4	4	9	13
2001.....	198	-	4	2	1	3	2	2	3	6	6	10
2000.....	183	2	2	1	3	2	3	1	3	3	6	7
1999.....	184	1	-	4	6	1	2	3	-	8	5	11
White, female												
2004.....	167	-	-	-	-	1	1	1	1	-	1	3
2003.....	142	1	1	3	2	1	-	2	3	1	1	8
2002.....	179	2	-	1	1	2	2	-	1	1	4	4
2001.....	144	1	1	-	1	2	3	1	2	1	4	4
2000.....	165	1	2	2	7	3	-	-	-	4	5	3
1999.....	130	2	-	4	4	-	-	-	1	2	4	5
Black, both sexes												
2004.....	59	-	1	1	2	1	1	3	1	1	2	2
2003.....	70	-	2	2	3	2	2	2	3	4	7	5
2002.....	63	1	-	1	2	3	2	1	1	-	5	7
2001.....	73	-	2	1	1	-	5	3	3	2	6	9
2000.....	77	-	-	1	1	-	2	2	5	4	5	8
1999.....	68	-	2	4	3	-	1	-	2	3	2	8
Black, male												
2004.....	39	-	1	1	1	1	-	3	1	1	2	2
2003.....	37	-	-	1	3	1	2	1	2	-	4	3
2002.....	42	1	-	-	1	3	1	1	1	-	1	6
2001.....	47	-	2	1	1	-	3	3	3	1	4	6
2000.....	48	-	-	-	-	-	1	2	4	2	5	5
1999.....	48	-	2	2	2	-	1	-	2	2	2	7

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Hernia (K40-K46)

All races, both sexes

2004.....	1,602	40	4	-	4	7	2	5	5	11	25	43
2003.....	1,613	37	6	4	2	-	3	1	4	7	21	45
2002.....	1,595	29	3	4	3	2	3	4	7	14	21	33
2001.....	1,522	43	2	1	2	2	1	6	9	13	21	24
2000.....	1,522	46	4	1	2	2	3	2	3	13	18	26
1999.....	1,464	44	3	1	1	1	2	6	10	15	24	16

All races, male

2004.....	652	20	1	-	2	4	1	3	4	6	16	20
2003.....	668	20	4	2	2	-	1	1	3	6	10	24
2002.....	599	18	3	3	2	1	3	2	5	8	9	16
2001.....	587	28	1	-	-	1	-	4	6	5	11	13
2000.....	586	28	3	1	2	1	2	-	1	8	5	13
1999.....	597	26	2	1	-	1	2	5	3	10	13	12

All races, female

2004.....	950	20	3	-	2	3	1	2	1	5	9	23
2003.....	945	17	2	2	-	-	2	-	1	1	11	21
2002.....	996	11	-	1	1	1	-	2	2	6	12	17
2001.....	935	15	1	1	2	1	1	2	3	8	10	11
2000.....	936	18	1	-	-	1	1	2	2	5	13	13
1999.....	867	18	1	-	1	-	-	1	7	5	11	4

White, both sexes

2004.....	1,423	32	2	-	4	5	2	3	5	6	19	30
2003.....	1,421	29	3	2	1	-	2	1	2	4	16	37
2002.....	1,433	24	2	4	3	2	3	2	4	11	15	24
2001.....	1,343	31	2	-	1	1	1	6	7	10	15	20
2000.....	1,354	36	2	1	2	2	1	2	2	9	13	20
1999.....	1,298	34	3	1	1	1	1	4	6	14	16	11

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hernia (K40-K46)												
White, male												
2004.....	572	15	-	-	2	3	1	1	4	2	13	16
2003.....	585	16	3	1	1	-	1	1	1	3	8	19
2002.....	533	14	2	3	2	1	3	-	3	6	8	13
2001.....	502	18	1	-	-	1	-	4	4	5	8	9
2000.....	520	23	2	1	2	1	-	-	1	6	4	9
1999.....	535	20	2	1	-	1	1	3	2	9	11	9
White, female												
2004.....	851	17	2	-	2	2	1	2	1	4	6	14
2003.....	836	13	-	1	-	-	1	-	1	1	8	18
2002.....	900	10	-	1	1	1	-	2	1	5	7	11
2001.....	841	13	1	-	1	-	1	2	3	5	7	11
2000.....	834	13	-	-	-	1	1	2	1	3	9	11
1999.....	763	14	1	-	1	-	-	1	4	5	5	2
Black, both sexes												
2004.....	150	6	2	-	-	2	-	2	-	3	6	11
2003.....	164	8	3	2	1	-	-	-	2	3	5	7
2002.....	137	4	1	-	-	-	-	2	1	3	5	9
2001.....	157	11	-	1	-	1	-	-	2	3	6	4
2000.....	145	7	2	-	-	-	2	-	1	3	5	5
1999.....	140	9	-	-	-	-	1	-	4	1	8	5
Black, male												
2004.....	67	4	1	-	-	1	-	2	-	2	3	4
2003.....	72	4	1	1	1	-	-	-	2	3	2	4
2002.....	56	3	1	-	-	-	-	2	-	2	1	3
2001.....	76	9	-	-	-	-	-	-	2	-	3	4
2000.....	54	3	1	-	-	-	2	-	-	1	1	4
1999.....	53	6	-	-	-	-	1	-	1	1	2	3

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Crohn's disease and ulcerative colitis (K50-K51)

All races, both sexes

2004.....	930	-	-	1	-	5	7	9	20	15	30	46
2003.....	865	1	-	-	3	5	8	16	15	9	31	51
2002.....	825	1	-	-	1	1	3	6	12	28	41	48
2001.....	831	-	2	-	-	4	9	10	11	21	19	32
2000.....	763	-	-	-	1	4	7	10	14	22	18	30
1999.....	799	1	-	-	1	6	5	7	17	19	32	47

All races, male

2004.....	391	-	-	1	-	2	4	8	11	7	15	22
2003.....	377	-	-	-	2	4	5	6	9	6	17	20
2002.....	333	1	-	-	-	1	-	1	8	10	21	24
2001.....	357	-	2	-	-	3	5	6	7	11	8	14
2000.....	290	-	-	-	-	3	4	6	8	8	7	19
1999.....	315	1	-	-	1	4	2	6	11	4	15	15

All races, female

2004.....	539	-	-	-	-	3	3	1	9	8	15	24
2003.....	488	1	-	-	1	1	3	10	6	3	14	31
2002.....	492	-	-	-	1	-	3	5	4	18	20	24
2001.....	474	-	-	-	-	1	4	4	4	10	11	18
2000.....	473	-	-	-	1	1	3	4	6	14	11	11
1999.....	484	-	-	-	-	2	3	1	6	15	17	32

White, both sexes

2004.....	862	-	-	1	-	2	3	7	19	14	26	40
2003.....	801	-	-	-	3	4	7	13	12	8	27	43
2002.....	765	1	-	-	-	1	2	5	11	19	33	43
2001.....	767	-	2	-	-	2	8	9	8	20	14	27
2000.....	708	-	-	-	1	3	4	7	12	14	14	26
1999.....	738	-	-	-	1	4	3	6	12	14	30	39

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Crohn's disease and ulcerative colitis (K50-K51)

White, male

2004.....	351	-	-	1	-	1	-	6	10	6	15	20
2003.....	348	-	-	-	2	3	5	5	6	5	14	18
2002.....	303	1	-	-	-	1	-	-	8	8	16	21
2001.....	327	-	2	-	-	2	4	5	6	10	6	13
2000.....	269	-	-	-	-	2	2	4	8	5	6	16
1999.....	290	-	-	-	1	2	1	5	8	4	14	12

White, female

2004.....	511	-	-	-	-	1	3	1	9	8	11	20
2003.....	453	-	-	-	1	1	2	8	6	3	13	25
2002.....	462	-	-	-	-	-	2	5	3	11	17	22
2001.....	440	-	-	-	-	-	4	4	2	10	8	14
2000.....	439	-	-	-	1	1	2	3	4	9	8	10
1999.....	448	-	-	-	-	2	2	1	4	10	16	27

Black, both sexes

2004.....	61	-	-	-	-	3	3	1	1	-	3	6
2003.....	56	1	-	-	-	1	1	2	3	1	4	8
2002.....	52	-	-	-	1	-	1	1	1	8	8	5
2001.....	56	-	-	-	-	2	1	1	2	1	5	5
2000.....	47	-	-	-	-	1	3	3	1	7	4	3
1999.....	56	1	-	-	-	2	2	1	5	5	2	8

Black, male

2004.....	36	-	-	-	-	1	3	1	1	-	-	2
2003.....	25	-	-	-	-	1	-	1	3	1	3	2
2002.....	27	-	-	-	-	-	-	1	-	2	5	3
2001.....	25	-	-	-	-	1	1	1	1	1	2	1
2000.....	17	-	-	-	-	1	2	2	-	2	1	2
1999.....	24	1	-	-	-	2	1	1	3	-	1	3

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Vascular disorders and obstruction of intestine without hernia (K55-K56)

All races, both sexes

2004.....	14,500	280	32	7	13	15	18	17	37	57	135	183
2003.....	14,851	271	23	16	12	13	26	25	25	49	118	194
2002.....	14,534	233	23	11	15	16	23	24	26	55	118	187
2001.....	14,357	234	23	14	12	10	8	25	24	47	99	176
2000.....	14,281	230	20	15	16	15	22	20	30	59	107	175
1999.....	13,966	230	22	12	10	16	14	21	38	54	121	150

All races, male

2004.....	5,109	161	20	6	5	9	10	6	17	31	72	87
2003.....	5,265	168	17	12	8	10	18	14	16	24	55	87
2002.....	5,104	134	17	8	3	5	16	8	20	29	69	85
2001.....	5,050	134	14	9	10	5	4	11	14	23	52	80
2000.....	5,002	122	14	12	11	11	14	11	15	32	47	88
1999.....	4,820	148	13	5	7	10	9	6	17	23	68	79

All races, female

2004.....	9,391	119	12	1	8	6	8	11	20	26	63	96
2003.....	9,586	103	6	4	4	3	8	11	9	25	63	107
2002.....	9,430	99	6	3	12	11	7	16	6	26	49	102
2001.....	9,307	100	9	5	2	5	4	14	10	24	47	96
2000.....	9,279	108	6	3	5	4	8	9	15	27	60	87
1999.....	9,146	82	9	7	3	6	5	15	21	31	53	71

White, both sexes

2004.....	12,655	145	21	6	9	9	8	11	20	39	98	136
2003.....	13,013	150	18	9	10	12	19	19	17	28	83	152
2002.....	12,779	131	12	9	10	8	18	17	20	43	89	136
2001.....	12,609	128	18	10	8	8	5	14	19	34	74	130
2000.....	12,474	114	9	10	15	9	20	15	17	39	77	111
1999.....	12,268	126	12	10	8	14	9	15	30	39	81	105

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Vascular disorders and obstruction of intestine without hernia (K55-K56)

White, male

2004.....	4,314	83	14	5	4	6	4	5	7	21	58	64
2003.....	4,442	98	15	8	6	9	13	10	11	13	40	62
2002.....	4,368	73	8	6	3	4	13	4	15	24	50	61
2001.....	4,320	70	11	6	6	4	3	6	12	14	37	58
2000.....	4,215	58	4	8	10	6	12	10	10	21	31	51
1999.....	4,123	84	6	4	6	10	4	5	12	18	45	54

White, female

2004.....	8,341	62	7	1	5	3	4	6	13	18	40	72
2003.....	8,571	52	3	1	4	3	6	9	6	15	43	90
2002.....	8,411	58	4	3	7	4	5	13	5	19	39	75
2001.....	8,289	58	7	4	2	4	2	8	7	20	37	72
2000.....	8,259	56	5	2	5	3	8	5	7	18	46	60
1999.....	8,145	42	6	6	2	4	5	10	18	21	36	51

Black, both sexes

2004.....	1,587	124	9	1	4	6	10	6	17	15	32	43
2003.....	1,571	119	5	3	2	1	7	5	7	18	30	39
2002.....	1,515	95	9	2	5	6	4	7	6	11	28	46
2001.....	1,541	103	4	4	4	2	3	8	4	12	22	45
2000.....	1,588	103	7	4	1	5	2	5	10	19	26	56
1999.....	1,517	94	8	2	2	2	5	5	8	13	40	41

Black, male

2004.....	677	72	4	1	1	3	6	1	10	7	11	22
2003.....	699	69	2	1	2	1	5	3	4	8	11	23
2002.....	626	57	7	2	-	-	2	4	5	5	18	22
2001.....	660	62	2	3	4	1	1	4	1	8	13	21
2000.....	688	58	6	3	1	5	2	1	4	10	15	33
1999.....	607	55	5	1	1	-	5	1	5	5	23	24

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Vascular disorders and obstruction of intestine without hernia (K55-K56)

Black, female

2004.....	910	52	5	-	3	3	4	5	7	8	21	21
2003.....	872	50	3	2	-	-	2	2	3	10	19	16
2002.....	889	38	2	-	5	6	2	3	1	6	10	24
2001.....	881	41	2	1	-	1	2	4	3	4	9	24
2000.....	900	45	1	1	-	-	-	4	6	9	11	23
1999.....	910	39	3	1	1	2	-	4	3	8	17	17

American Indian 1/, both sexes

2004.....	79	1	-	-	-	-	-	-	-	2	4	3
2003.....	85	1	-	1	-	-	-	-	1	1	3	1
2002.....	70	1	-	-	-	1	-	-	-	-	-	2
2001.....	78	-	1	-	-	-	-	-	1	-	2	-
2000.....	66	2	2	-	-	-	-	-	1	1	1	4
1999.....	59	3	1	-	-	-	-	1	-	2	-	3

American Indian 1/, male

2004.....	21	1	-	-	-	-	-	-	-	2	2	-
2003.....	32	-	-	1	-	-	-	-	1	1	2	-
2002.....	26	-	-	-	-	-	-	-	-	-	-	1
2001.....	24	-	1	-	-	-	-	-	1	-	1	-
2000.....	32	1	2	-	-	-	-	-	1	1	1	3
1999.....	18	2	1	-	-	-	-	-	-	-	-	1

American Indian 1/, female

2004.....	58	-	-	-	-	-	-	-	-	-	2	3
2003.....	53	1	-	-	-	-	-	-	-	-	1	1
2002.....	44	1	-	-	-	1	-	-	-	-	-	1
2001.....	54	-	-	-	-	-	-	-	-	-	1	-
2000.....	34	1	-	-	-	-	-	-	-	-	-	1
1999.....	41	1	-	-	-	-	-	1	-	2	-	2

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Diverticular disease of intestine (K57)

All races, both sexes

2004.....	3,370	-	-	-	-	-	1	2	5	8	23	18
2003.....	3,480	-	-	-	-	-	-	3	-	8	21	30
2002.....	3,324	-	-	-	-	-	-	2	3	7	18	43
2001.....	3,438	-	-	-	-	-	2	1	7	8	14	43
2000.....	3,290	-	-	-	-	-	1	3	5	8	21	34
1999.....	3,427	-	-	-	-	-	1	-	6	12	16	36

All races, male

2004.....	1,071	-	-	-	-	-	-	1	4	6	13	10
2003.....	1,080	-	-	-	-	-	-	3	-	5	14	16
2002.....	1,063	-	-	-	-	-	-	1	3	6	15	27
2001.....	1,078	-	-	-	-	-	-	1	2	6	6	26
2000.....	1,012	-	-	-	-	-	1	2	4	4	12	20
1999.....	1,047	-	-	-	-	-	1	-	3	7	10	19

All races, female

2004.....	2,299	-	-	-	-	-	1	1	1	2	10	8
2003.....	2,400	-	-	-	-	-	-	-	-	3	7	14
2002.....	2,261	-	-	-	-	-	-	1	-	1	3	16
2001.....	2,360	-	-	-	-	-	2	-	5	2	8	17
2000.....	2,278	-	-	-	-	-	-	1	1	4	9	14
1999.....	2,380	-	-	-	-	-	-	-	3	5	6	17

White, both sexes

2004.....	3,082	-	-	-	-	-	1	1	4	6	17	12
2003.....	3,193	-	-	-	-	-	-	2	-	7	15	26
2002.....	3,033	-	-	-	-	-	-	1	2	7	16	39
2001.....	3,152	-	-	-	-	-	2	-	6	7	13	38
2000.....	3,018	-	-	-	-	-	1	3	5	5	16	27
1999.....	3,151	-	-	-	-	-	1	-	5	9	14	31

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diverticular disease of intestine (K57)												
White, male												
2004.....	969	-	-	-	-	-	-	1	3	5	12	7
2003.....	994	-	-	-	-	-	-	2	-	5	10	13
2002.....	956	-	-	-	-	-	-	1	2	6	13	23
2001.....	987	-	-	-	-	-	-	-	2	6	6	24
2000.....	927	-	-	-	-	-	1	2	4	2	8	16
1999.....	963	-	-	-	-	-	1	-	3	5	9	17
White, female												
2004.....	2,113	-	-	-	-	-	1	-	1	1	5	5
2003.....	2,199	-	-	-	-	-	-	-	-	2	5	13
2002.....	2,077	-	-	-	-	-	-	-	-	1	3	16
2001.....	2,165	-	-	-	-	-	2	-	4	1	7	14
2000.....	2,091	-	-	-	-	-	-	1	1	3	8	11
1999.....	2,188	-	-	-	-	-	-	-	2	4	5	14
Black, both sexes												
2004.....	243	-	-	-	-	-	-	-	-	2	5	6
2003.....	242	-	-	-	-	-	-	1	-	1	5	3
2002.....	247	-	-	-	-	-	-	-	1	-	1	4
2001.....	250	-	-	-	-	-	-	1	1	1	-	4
2000.....	244	-	-	-	-	-	-	-	-	3	5	7
1999.....	246	-	-	-	-	-	-	-	1	3	1	5
Black, male												
2004.....	83	-	-	-	-	-	-	-	-	1	1	3
2003.....	70	-	-	-	-	-	-	1	-	-	3	2
2002.....	86	-	-	-	-	-	-	-	1	-	1	4
2001.....	73	-	-	-	-	-	-	1	-	-	-	2
2000.....	74	-	-	-	-	-	-	-	-	2	4	4
1999.....	70	-	-	-	-	-	-	-	-	2	-	2

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other diseases of intestines and peritoneum (K52,K58-K63)

All races, both sexes

2004.....	4,964	82	24	7	7	8	15	9	18	37	49	83
2003.....	4,910	95	23	7	12	10	16	12	22	39	51	85
2002.....	4,841	80	35	12	8	6	14	8	19	37	69	92
2001.....	4,757	55	26	8	7	7	10	5	16	37	58	91
2000.....	4,525	66	19	7	6	7	10	17	20	28	57	95
1999.....	4,434	56	26	8	10	5	10	11	17	40	61	85

All races, male

2004.....	1,811	48	15	3	5	6	6	6	8	15	16	48
2003.....	1,868	60	12	4	5	8	10	5	15	21	18	47
2002.....	1,845	50	19	9	5	4	6	4	9	17	34	53
2001.....	1,742	33	13	4	5	6	6	2	12	17	26	40
2000.....	1,648	36	12	4	2	3	6	11	8	12	32	42
1999.....	1,687	39	11	2	6	5	5	5	9	21	32	55

All races, female

2004.....	3,153	34	9	4	2	2	9	3	10	22	33	35
2003.....	3,042	35	11	3	7	2	6	7	7	18	33	38
2002.....	2,996	30	16	3	3	2	8	4	10	20	35	39
2001.....	3,015	22	13	4	2	1	4	3	4	20	32	51
2000.....	2,877	30	7	3	4	4	4	6	12	16	25	53
1999.....	2,747	17	15	6	4	-	5	6	8	19	29	30

White, both sexes

2004.....	4,411	50	13	6	4	7	10	7	11	27	38	62
2003.....	4,338	65	19	3	9	7	12	9	16	25	44	65
2002.....	4,326	49	24	7	7	6	10	4	15	30	53	75
2001.....	4,200	26	16	7	5	5	8	5	13	27	47	69
2000.....	4,004	42	8	5	1	4	7	13	18	21	43	78
1999.....	3,963	36	20	5	7	4	7	8	15	31	48	65

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other diseases of intestines and peritoneum (K52,K58-K63)

White, male

2004.....	1,561	29	7	3	3	5	4	5	4	7	15	36
2003.....	1,623	43	11	3	3	6	8	5	11	16	15	36
2002.....	1,613	35	13	5	4	4	6	2	8	14	26	42
2001.....	1,503	12	8	3	4	4	5	2	10	12	21	35
2000.....	1,421	21	6	2	-	1	4	9	7	10	25	36
1999.....	1,471	24	6	1	5	4	4	3	8	16	27	43

White, female

2004.....	2,850	21	6	3	1	2	6	2	7	20	23	26
2003.....	2,715	22	8	-	6	1	4	4	5	9	29	29
2002.....	2,713	14	11	2	3	2	4	2	7	16	27	33
2001.....	2,697	14	8	4	1	1	3	3	3	15	26	34
2000.....	2,583	21	2	3	1	3	3	4	11	11	18	42
1999.....	2,492	12	14	4	2	-	3	5	7	15	21	22

Black, both sexes

2004.....	456	24	8	1	3	1	5	2	5	9	9	19
2003.....	497	25	4	1	3	2	4	3	5	14	7	18
2002.....	437	26	8	5	1	-	3	3	3	7	14	11
2001.....	488	25	8	1	2	2	1	-	3	10	11	20
2000.....	432	23	8	1	3	2	2	4	2	7	14	13
1999.....	408	18	6	2	3	1	3	2	1	8	12	20

Black, male

2004.....	209	15	6	-	2	1	2	1	4	7	1	11
2003.....	210	14	1	-	2	2	2	-	3	5	3	10
2002.....	197	12	4	4	1	-	-	2	1	3	7	7
2001.....	210	17	4	1	1	2	-	-	2	5	5	4
2000.....	179	14	5	1	1	1	1	2	1	2	7	5
1999.....	182	13	5	-	1	1	1	1	-	4	5	12

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Diseases of peritoneum (K65-K66)

All races, both sexes

2004.....	1,781	6	3	2	5	5	7	6	14	33	48	78
2003.....	1,862	15	2	2	3	9	5	17	15	23	51	78
2002.....	1,939	8	1	-	4	6	8	7	19	34	47	78
2001.....	1,874	12	3	2	6	1	6	5	11	32	50	93
2000.....	1,948	16	2	3	2	6	4	5	23	36	66	72
1999.....	1,894	20	5	4	6	3	7	9	18	35	57	75

All races, male

2004.....	778	4	2	1	2	2	-	2	7	18	24	43
2003.....	813	11	1	2	-	4	3	8	5	11	32	36
2002.....	755	4	-	-	2	1	3	3	10	14	21	41
2001.....	819	7	1	2	3	1	2	2	6	20	30	54
2000.....	832	9	-	-	1	5	1	3	12	17	35	43
1999.....	868	12	3	2	4	-	3	4	13	18	34	52

All races, female

2004.....	1,003	2	1	1	3	3	7	4	7	15	24	35
2003.....	1,049	4	1	-	3	5	2	9	10	12	19	42
2002.....	1,184	4	1	-	2	5	5	4	9	20	26	37
2001.....	1,055	5	2	-	3	-	4	3	5	12	20	39
2000.....	1,116	7	2	3	1	1	3	2	11	19	31	29
1999.....	1,026	8	2	2	2	3	4	5	5	17	23	23

White, both sexes

2004.....	1,507	5	1	2	2	3	3	3	12	19	34	57
2003.....	1,557	13	1	1	2	6	4	10	14	19	37	51
2002.....	1,640	2	1	-	3	5	5	3	10	22	39	48
2001.....	1,542	7	2	2	4	1	3	4	8	19	29	65
2000.....	1,610	10	1	1	-	5	2	1	14	21	46	48
1999.....	1,577	11	5	2	3	1	4	4	16	30	45	61

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of peritoneum (K65-K66)												
White, male												
2004.....	657	3	1	1	1	1	-	2	7	12	16	35
2003.....	686	10	-	1	-	3	2	5	5	9	25	30
2002.....	631	1	-	-	2	1	3	1	7	9	16	25
2001.....	668	4	1	2	2	1	1	1	5	11	19	41
2000.....	676	7	-	-	-	4	-	1	9	10	23	29
1999.....	726	7	3	2	2	-	2	2	12	17	29	42
White, female												
2004.....	850	2	-	1	1	2	3	1	5	7	18	22
2003.....	871	3	1	-	2	3	2	5	9	10	12	21
2002.....	1,009	1	1	-	1	4	2	2	3	13	23	23
2001.....	874	3	1	-	2	-	2	3	3	8	10	24
2000.....	934	3	1	1	-	1	2	-	5	11	23	19
1999.....	851	4	2	-	1	1	2	2	4	13	16	19
Black, both sexes												
2004.....	235	1	2	-	3	1	3	3	1	13	12	16
2003.....	263	1	1	-	1	2	1	7	1	3	12	24
2002.....	261	5	-	-	1	1	3	3	6	12	7	26
2001.....	286	4	1	-	1	-	3	1	2	11	19	27
2000.....	299	6	1	2	2	1	2	3	8	12	18	23
1999.....	279	8	-	2	1	2	3	4	2	4	9	13
Black, male												
2004.....	97	1	1	-	1	-	-	-	-	5	6	6
2003.....	106	1	1	-	-	-	1	3	-	1	6	6
2002.....	105	3	-	-	-	-	-	1	2	5	5	13
2001.....	127	2	-	-	-	-	1	1	1	8	9	13
2000.....	137	2	-	-	1	1	1	1	2	5	10	14
1999.....	124	4	-	-	1	-	1	1	1	1	3	9

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of peritoneum (K65-K66)												
Asian or Pacific Islander 2/, both sexes												
2004.....	25	-	-	-	-	1	-	-	1	-	1	3
2003.....	30	1	-	1	-	1	-	-	-	1	1	2
2002.....	28	1	-	-	-	-	-	-	1	-	1	4
2001.....	37	-	-	-	1	-	-	-	1	1	1	-
2000.....	22	-	-	-	-	-	-	-	-	1	1	-
1999.....	32	1	-	-	1	-	-	1	-	1	2	1
Asian or Pacific Islander 2/, male												
2004.....	16	-	-	-	-	1	-	-	-	-	1	1
2003.....	16	-	-	1	-	1	-	-	-	1	-	-
2002.....	15	-	-	-	-	-	-	-	1	-	-	3
2001.....	18	-	-	-	1	-	-	-	-	-	1	-
2000.....	13	-	-	-	-	-	-	-	-	1	1	-
1999.....	16	1	-	-	-	-	-	1	-	-	2	1
Asian or Pacific Islander 2/, female												
2004.....	9	-	-	-	-	-	-	-	1	-	-	2
2003.....	14	1	-	-	-	-	-	-	-	-	1	2
2002.....	13	1	-	-	-	-	-	-	-	-	1	1
2001.....	19	-	-	-	-	-	-	-	1	1	-	-
2000.....	9	-	-	-	-	-	-	-	-	-	-	-
1999.....	16	-	-	-	1	-	-	-	-	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of liver (K70-K76)												
All races, both sexes												
2004.....	36,022	101	31	8	6	20	63	111	374	1,028	2,564	4,280
2003.....	36,523	61	36	6	5	32	51	142	407	1,145	2,735	4,392
2002.....	35,671	89	27	10	19	20	55	134	405	1,171	2,835	4,401
2001.....	34,840	78	28	15	15	23	55	112	416	1,354	2,715	4,376
2000.....	33,853	76	27	11	14	14	68	125	433	1,359	2,778	4,101
1999.....	33,619	86	23	11	16	31	47	112	456	1,367	2,695	3,916
All races, male												
2004.....	22,659	53	15	4	1	7	35	69	255	662	1,715	2,946
2003.....	23,064	44	20	3	2	13	22	86	265	716	1,800	3,133
2002.....	22,403	63	11	6	11	9	26	74	256	740	1,858	3,128
2001.....	21,919	47	15	9	9	10	28	66	260	864	1,806	3,129
2000.....	21,351	52	15	6	6	7	39	82	273	886	1,911	3,016
1999.....	21,350	57	11	6	3	16	18	64	291	871	1,919	2,921
All races, female												
2004.....	13,363	48	16	4	5	13	28	42	119	366	849	1,334
2003.....	13,459	17	16	3	3	19	29	56	142	429	935	1,259
2002.....	13,268	26	16	4	8	11	29	60	149	431	977	1,273
2001.....	12,921	31	13	6	6	13	27	46	156	490	909	1,247
2000.....	12,502	24	12	5	8	7	29	43	160	473	867	1,085
1999.....	12,269	29	12	5	13	15	29	48	165	496	776	995
White, both sexes												
2004.....	30,987	59	24	6	4	13	37	78	302	818	2,119	3,561
2003.....	31,505	37	26	6	4	17	28	92	325	937	2,295	3,659
2002.....	30,792	60	19	6	13	15	37	92	311	953	2,377	3,618
2001.....	29,879	51	21	11	9	12	30	74	330	1,097	2,205	3,601
2000.....	28,983	48	16	6	11	7	43	79	330	1,078	2,263	3,380
1999.....	28,673	50	18	7	13	23	28	82	353	1,099	2,147	3,117

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of liver (K70-K76)												
White, male												
2004.....	19,570	30	11	2	1	6	22	48	211	531	1,446	2,470
2003.....	19,961	28	16	3	1	8	12	57	221	599	1,554	2,659
2002.....	19,453	43	7	3	9	5	21	53	208	616	1,595	2,633
2001.....	18,810	31	14	6	4	6	18	46	207	699	1,480	2,621
2000.....	18,410	30	8	4	5	5	24	58	216	720	1,590	2,555
1999.....	18,323	30	10	2	2	14	11	52	234	718	1,564	2,371
White, female												
2004.....	11,417	29	13	4	3	7	15	30	91	287	673	1,091
2003.....	11,544	9	10	3	3	9	16	35	104	338	741	1,000
2002.....	11,339	17	12	3	4	10	16	39	103	337	782	985
2001.....	11,069	20	7	5	5	6	12	28	123	398	725	980
2000.....	10,573	18	8	2	6	2	19	21	114	358	673	825
1999.....	10,350	20	8	5	11	9	17	30	119	381	583	746
Black, both sexes												
2004.....	3,776	36	5	2	2	5	20	20	38	134	316	568
2003.....	3,828	20	7	-	1	12	20	38	50	142	312	550
2002.....	3,732	25	6	4	5	5	17	27	54	130	327	622
2001.....	3,877	22	5	3	4	9	18	25	53	170	391	629
2000.....	3,817	26	9	4	3	6	20	33	64	203	399	594
1999.....	3,919	30	4	3	3	8	17	25	68	187	430	663
Black, male												
2004.....	2,373	19	2	2	-	1	10	11	24	81	193	376
2003.....	2,419	13	3	-	1	3	9	21	25	78	169	357
2002.....	2,271	19	2	3	2	4	4	12	29	66	182	388
2001.....	2,476	13	-	2	3	3	5	12	31	108	251	422
2000.....	2,348	20	5	2	1	1	12	17	31	123	244	379
1999.....	2,425	22	-	3	1	2	6	9	35	106	274	461

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Diseases of liver (K70-K76)

Black, female

2004.....	1,403	17	3	-	2	4	10	9	14	53	123	192
2003.....	1,409	7	4	-	-	9	11	17	25	64	143	193
2002.....	1,461	6	4	1	3	1	13	15	25	64	145	234
2001.....	1,401	9	5	1	1	6	13	13	22	62	140	207
2000.....	1,469	6	4	2	2	5	8	16	33	80	155	215
1999.....	1,494	8	4	-	2	6	11	16	33	81	156	202

American Indian 1/, both sexes

2004.....	736	1	1	-	-	-	4	7	24	58	97	113
2003.....	725	-	1	-	-	3	3	8	25	57	104	126
2002.....	681	1	1	-	1	-	1	10	33	70	98	114
2001.....	650	1	-	-	2	2	5	11	24	67	89	107
2000.....	623	-	-	-	-	1	2	10	30	64	85	84
1999.....	616	1	-	-	-	-	2	3	24	66	92	98

American Indian 1/, male

2004.....	398	1	1	-	-	-	2	4	11	35	50	68
2003.....	399	-	-	-	-	2	1	6	14	31	58	74
2002.....	400	-	1	-	-	-	1	6	14	45	54	69
2001.....	376	1	-	-	2	1	4	7	15	41	54	57
2000.....	341	-	-	-	-	1	1	5	19	31	53	50
1999.....	353	1	-	-	-	-	1	2	13	35	59	59

American Indian 1/, female

2004.....	338	-	-	-	-	-	2	3	13	23	47	45
2003.....	326	-	1	-	-	1	2	2	11	26	46	52
2002.....	281	1	-	-	1	-	-	4	19	25	44	45
2001.....	274	-	-	-	-	1	1	4	9	26	35	50
2000.....	282	-	-	-	-	-	1	5	11	33	32	34
1999.....	263	-	-	-	-	-	1	1	11	31	33	39

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Diseases of liver (K70-K76)

Asian or Pacific Islander 2/, both sexes

2004.....	523	5	1	-	-	2	2	6	10	18	32	38
2003.....	465	4	2	-	-	-	-	4	7	9	24	57
2002.....	466	3	1	-	-	-	-	5	7	18	33	47
2001.....	434	4	2	1	-	-	2	2	9	20	30	39
2000.....	430	2	2	1	-	-	3	3	9	14	31	43
1999.....	411	5	1	1	-	-	-	2	11	15	26	38

Asian or Pacific Islander 2/, male

2004.....	318	3	1	-	-	-	1	6	9	15	26	32
2003.....	285	3	1	-	-	-	-	2	5	8	19	43
2002.....	279	1	1	-	-	-	-	3	5	13	27	38
2001.....	257	2	1	1	-	-	1	1	7	16	21	29
2000.....	252	2	2	-	-	-	2	2	7	12	24	32
1999.....	249	4	1	1	-	-	-	1	9	12	22	30

Asian or Pacific Islander 2/, female

2004.....	205	2	-	-	-	2	1	-	1	3	6	6
2003.....	180	1	1	-	-	-	-	2	2	1	5	14
2002.....	187	2	-	-	-	-	-	2	2	5	6	9
2001.....	177	2	1	-	-	-	1	1	2	4	9	10
2000.....	178	-	-	1	-	-	1	1	2	2	7	11
1999.....	162	1	-	-	-	-	-	1	2	3	4	8

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Alcoholic liver disease (K70)

All races, both sexes

2004.....	12,548	-	-	-	-	-	11	42	186	533	1,338	2,023
2003.....	12,360	-	-	-	-	1	5	49	194	612	1,359	2,050
2002.....	12,121	-	1	-	-	-	8	43	199	607	1,364	1,998
2001.....	12,207	-	-	-	-	1	7	45	219	745	1,400	2,121
2000.....	12,109	-	-	-	-	-	8	56	223	741	1,485	2,000
1999.....	11,958	-	-	-	-	-	8	37	222	763	1,423	1,965

All races, male

2004.....	9,155	-	-	-	-	-	8	28	139	346	921	1,404
2003.....	9,104	-	-	-	-	-	3	39	143	402	915	1,524
2002.....	8,814	-	1	-	-	-	6	32	130	392	913	1,450
2001.....	8,932	-	-	-	-	-	5	31	148	492	954	1,563
2000.....	8,985	-	-	-	-	-	5	45	157	486	1,047	1,510
1999.....	8,922	-	-	-	-	-	6	24	141	491	1,029	1,501

All races, female

2004.....	3,393	-	-	-	-	-	3	14	47	187	417	619
2003.....	3,256	-	-	-	-	1	2	10	51	210	444	526
2002.....	3,307	-	-	-	-	-	2	11	69	215	451	548
2001.....	3,275	-	-	-	-	1	2	14	71	253	446	558
2000.....	3,124	-	-	-	-	-	3	11	66	255	438	490
1999.....	3,036	-	-	-	-	-	2	13	81	272	394	464

White, both sexes

2004.....	10,859	-	-	-	-	-	7	33	156	439	1,143	1,731
2003.....	10,625	-	-	-	-	-	4	37	161	499	1,159	1,743
2002.....	10,423	-	1	-	-	-	4	31	156	501	1,159	1,700
2001.....	10,387	-	-	-	-	-	2	32	179	613	1,158	1,812
2000.....	10,272	-	-	-	-	-	6	42	176	596	1,214	1,670
1999.....	10,041	-	-	-	-	-	5	31	174	623	1,141	1,605

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Alcoholic liver disease (K70)

White, male

2004.....	7,992	-	-	-	-	-	6	20	120	283	799	1,210
2003.....	7,905	-	-	-	-	-	2	31	118	339	792	1,306
2002.....	7,688	-	1	-	-	-	4	25	110	326	786	1,257
2001.....	7,689	-	-	-	-	-	2	26	125	407	795	1,363
2000.....	7,770	-	-	-	-	-	4	36	132	397	867	1,301
1999.....	7,602	-	-	-	-	-	4	21	115	409	840	1,252

White, female

2004.....	2,867	-	-	-	-	-	1	13	36	156	344	521
2003.....	2,720	-	-	-	-	-	2	6	43	160	367	437
2002.....	2,735	-	-	-	-	-	-	6	46	175	373	443
2001.....	2,698	-	-	-	-	-	-	6	54	206	363	449
2000.....	2,502	-	-	-	-	-	2	6	44	199	347	369
1999.....	2,439	-	-	-	-	-	1	10	59	214	301	353

Black, both sexes

2004.....	1,169	-	-	-	-	-	2	1	8	43	121	202
2003.....	1,222	-	-	-	-	-	-	6	13	67	121	199
2002.....	1,191	-	-	-	-	-	3	6	20	47	123	209
2001.....	1,348	-	-	-	-	-	1	7	20	75	170	221
2000.....	1,380	-	-	-	-	-	-	7	27	94	193	259
1999.....	1,466	-	-	-	-	-	2	4	25	84	199	292

Black, male

2004.....	812	-	-	-	-	-	-	1	6	29	77	135
2003.....	865	-	-	-	-	-	-	4	10	37	74	147
2002.....	797	-	-	-	-	-	1	4	10	27	76	136
2001.....	935	-	-	-	-	-	-	2	10	51	110	148
2000.....	918	-	-	-	-	-	-	5	12	61	126	164
1999.....	1,018	-	-	-	-	-	2	2	13	49	130	204

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Alcoholic liver disease (K70)

Black, female

2004.....	357	-	-	-	-	-	2	-	2	14	44	67
2003.....	357	-	-	-	-	-	-	2	3	30	47	52
2002.....	394	-	-	-	-	-	2	2	10	20	47	73
2001.....	413	-	-	-	-	-	1	5	10	24	60	73
2000.....	462	-	-	-	-	-	-	2	15	33	67	95
1999.....	448	-	-	-	-	-	-	2	12	35	69	88

American Indian 1/, both sexes

2004.....	389	-	-	-	-	-	2	4	16	43	61	69
2003.....	403	-	-	-	-	1	1	5	16	42	66	81
2002.....	381	-	-	-	-	-	1	6	22	49	63	71
2001.....	359	-	-	-	-	1	4	5	15	47	63	68
2000.....	339	-	-	-	-	-	1	7	18	42	57	52
1999.....	356	-	-	-	-	-	1	2	17	48	70	54

American Indian 1/, male

2004.....	240	-	-	-	-	-	2	3	8	28	34	40
2003.....	241	-	-	-	-	-	1	3	11	23	38	47
2002.....	230	-	-	-	-	-	1	3	10	31	35	41
2001.....	218	-	-	-	-	-	3	3	10	27	41	34
2000.....	205	-	-	-	-	-	-	4	11	20	36	30
1999.....	224	-	-	-	-	-	-	1	8	26	47	33

American Indian 1/, female

2004.....	149	-	-	-	-	-	-	1	8	15	27	29
2003.....	162	-	-	-	-	1	-	2	5	19	28	34
2002.....	151	-	-	-	-	-	-	3	12	18	28	30
2001.....	141	-	-	-	-	1	1	2	5	20	22	34
2000.....	134	-	-	-	-	-	1	3	7	22	21	22
1999.....	132	-	-	-	-	-	1	1	9	22	23	21

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Alcoholic liver disease (K70)												
Asian or Pacific Islander 2/, both sexes												
2004.....	131	-	-	-	-	-	-	4	6	8	13	21
2003.....	110	-	-	-	-	-	-	1	4	4	13	27
2002.....	126	-	-	-	-	-	-	-	1	10	19	18
2001.....	113	-	-	-	-	-	-	1	5	10	9	20
2000.....	118	-	-	-	-	-	1	-	2	9	21	19
1999.....	95	-	-	-	-	-	-	-	6	8	13	14
Asian or Pacific Islander 2/, male												
2004.....	111	-	-	-	-	-	-	4	5	6	11	19
2003.....	93	-	-	-	-	-	-	1	4	3	11	24
2002.....	99	-	-	-	-	-	-	-	-	8	16	16
2001.....	90	-	-	-	-	-	-	-	3	7	8	18
2000.....	92	-	-	-	-	-	1	-	2	8	18	15
1999.....	78	-	-	-	-	-	-	-	5	7	12	12
Asian or Pacific Islander 2/, female												
2004.....	20	-	-	-	-	-	-	-	1	2	2	2
2003.....	17	-	-	-	-	-	-	-	-	1	2	3
2002.....	27	-	-	-	-	-	-	-	1	2	3	2
2001.....	23	-	-	-	-	-	-	1	2	3	1	2
2000.....	26	-	-	-	-	-	-	-	-	1	3	4
1999.....	17	-	-	-	-	-	-	-	1	1	1	2

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Chronic hepatitis, not elsewhere classified (K73)

All races, both sexes

2004.....	119	-	-	-	1	-	-	-	2	1	6	21
2003.....	144	-	-	-	-	-	-	-	2	5	10	13
2002.....	534	1	2	1	3	6	7	7	11	13	26	40
2001.....	467	-	-	-	1	2	9	7	9	13	16	37
2000.....	440	-	2	-	3	1	3	8	6	9	23	22
1999.....	439	-	-	1	1	4	4	8	7	10	19	33

All races, male

2004.....	64	-	-	-	-	-	-	-	1	-	5	17
2003.....	76	-	-	-	-	-	-	-	1	2	10	9
2002.....	145	1	-	1	1	3	1	1	3	3	13	19
2001.....	127	-	-	-	1	-	3	4	2	6	8	11
2000.....	105	-	1	-	2	-	2	4	-	4	10	11
1999.....	128	-	-	1	-	-	-	1	3	-	12	16

All races, female

2004.....	55	-	-	-	1	-	-	-	1	1	1	4
2003.....	68	-	-	-	-	-	-	-	1	3	-	4
2002.....	389	-	2	-	2	3	6	6	8	10	13	21
2001.....	340	-	-	-	-	2	6	3	7	7	8	26
2000.....	335	-	1	-	1	1	1	4	6	5	13	11
1999.....	311	-	-	-	1	4	4	7	4	10	7	17

White, both sexes

2004.....	106	-	-	-	1	-	-	-	-	1	5	19
2003.....	127	-	-	-	-	-	-	-	1	5	9	12
2002.....	460	1	2	1	3	6	6	6	5	10	17	28
2001.....	405	-	-	-	1	2	5	5	7	10	10	28
2000.....	374	-	2	-	2	-	1	3	5	7	17	15
1999.....	355	-	-	-	1	2	2	4	3	4	16	25

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Chronic hepatitis, not elsewhere classified (K73)

White, male

2004.....	55	-	-	-	-	-	-	-	-	-	4	15
2003.....	70	-	-	-	-	-	-	-	1	2	9	8
2002.....	128	1	-	1	1	3	1	1	1	3	11	13
2001.....	115	-	-	-	1	-	3	4	2	4	6	9
2000.....	88	-	1	-	1	-	1	3	-	3	7	9
1999.....	106	-	-	-	-	-	-	-	1	-	11	12

White, female

2004.....	51	-	-	-	1	-	-	-	-	1	1	4
2003.....	57	-	-	-	-	-	-	-	-	3	-	4
2002.....	332	-	2	-	2	3	5	5	4	7	6	15
2001.....	290	-	-	-	-	2	2	1	5	6	4	19
2000.....	286	-	1	-	1	-	-	-	5	4	10	6
1999.....	249	-	-	-	1	2	2	4	2	4	5	13

Black, both sexes

2004.....	7	-	-	-	-	-	-	-	-	-	1	2
2003.....	12	-	-	-	-	-	-	-	-	-	1	1
2002.....	59	-	-	-	-	-	1	-	5	3	9	11
2001.....	48	-	-	-	-	-	4	2	2	2	6	7
2000.....	48	-	-	-	1	1	2	5	1	1	5	5
1999.....	61	-	-	1	-	2	2	4	2	5	3	6

Black, male

2004.....	6	-	-	-	-	-	-	-	-	-	1	2
2003.....	5	-	-	-	-	-	-	-	-	-	1	1
2002.....	13	-	-	-	-	-	-	-	1	-	2	5
2001.....	9	-	-	-	-	-	-	-	-	1	2	2
2000.....	13	-	-	-	1	-	1	1	-	1	3	1
1999.....	13	-	-	1	-	-	-	1	1	-	1	3

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic hepatitis, not elsewhere classified (K73)												
Black, female												
2004.....	1	-	-	-	-	-	-	-	-	-	-	-
2003.....	7	-	-	-	-	-	-	-	-	-	-	-
2002.....	46	-	-	-	-	-	1	-	4	3	7	6
2001.....	39	-	-	-	-	-	4	2	2	1	4	5
2000.....	35	-	-	-	-	1	1	4	1	-	2	4
1999.....	48	-	-	-	-	2	2	3	1	5	2	3
American Indian 1/, both sexes												
2004.....	3	-	-	-	-	-	-	-	1	-	-	-
2003.....	1	-	-	-	-	-	-	-	-	-	-	-
2002.....	3	-	-	-	-	-	-	-	-	-	-	-
2001.....	2	-	-	-	-	-	-	-	-	-	-	-
2000.....	7	-	-	-	-	-	-	-	-	-	1	2
1999.....	6	-	-	-	-	-	-	-	1	-	-	2
American Indian 1/, male												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	1	-	-	-	-	-	-	-	-	-	-	-
2002.....	1	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	1	-	-	-	-	-	-	-	-	-	-	1
1999.....	1	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, female												
2004.....	3	-	-	-	-	-	-	-	1	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	2	-	-	-	-	-	-	-	-	-	-	-
2001.....	2	-	-	-	-	-	-	-	-	-	-	-
2000.....	6	-	-	-	-	-	-	-	-	-	1	1
1999.....	5	-	-	-	-	-	-	-	1	-	-	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic hepatitis, not elsewhere classified (K73)												
Asian or Pacific Islander 2/, both sexes												
2004.....	3	-	-	-	-	-	-	-	1	-	-	-
2003.....	4	-	-	-	-	-	-	-	1	-	-	-
2002.....	12	-	-	-	-	-	-	1	1	-	-	1
2001.....	12	-	-	-	-	-	-	-	-	1	-	2
2000.....	11	-	-	-	-	-	-	-	-	1	-	-
1999.....	17	-	-	-	-	-	-	-	1	1	-	-
Asian or Pacific Islander 2/, male												
2004.....	3	-	-	-	-	-	-	-	1	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	3	-	-	-	-	-	-	-	1	-	-	1
2001.....	3	-	-	-	-	-	-	-	-	1	-	-
2000.....	3	-	-	-	-	-	-	-	-	-	-	-
1999.....	8	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	4	-	-	-	-	-	-	-	1	-	-	-
2002.....	9	-	-	-	-	-	-	1	-	-	-	-
2001.....	9	-	-	-	-	-	-	-	-	-	-	2
2000.....	8	-	-	-	-	-	-	-	-	1	-	-
1999.....	9	-	-	-	-	-	-	-	-	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	-------------	-----------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Fibrosis and cirrhosis of liver (K74)

All races, both sexes

2004.....	14,346	9	1	-	-	-	8	10	69	248	673	1,307
2003.....	14,999	2	-	-	-	2	8	29	84	265	769	1,394
2002.....	14,602	5	-	-	2	1	9	27	87	309	835	1,366
2001.....	14,361	3	-	1	-	3	4	25	82	350	812	1,399
2000.....	14,003	3	1	-	-	6	18	24	98	333	780	1,352
1999.....	13,862	5	2	-	-	5	5	25	108	330	757	1,230

All races, male

2004.....	8,326	6	-	-	-	-	1	6	47	176	468	923
2003.....	8,732	2	-	-	-	1	4	15	54	170	534	977
2002.....	8,442	5	-	-	1	-	4	13	57	210	548	973
2001.....	8,334	1	-	1	-	2	2	18	56	215	531	1,007
2000.....	8,124	3	-	-	-	3	9	14	60	226	541	994
1999.....	8,065	4	-	-	-	3	2	16	72	225	557	929

All races, female

2004.....	6,020	3	1	-	-	-	7	4	22	72	205	384
2003.....	6,267	-	-	-	-	1	4	14	30	95	235	417
2002.....	6,160	-	-	-	1	1	5	14	30	99	287	393
2001.....	6,027	2	-	-	-	1	2	7	26	135	281	392
2000.....	5,879	-	1	-	-	3	9	10	38	107	239	358
1999.....	5,797	1	2	-	-	2	3	9	36	105	200	301

White, both sexes

2004.....	12,597	7	1	-	-	-	5	7	55	198	541	1,088
2003.....	13,253	1	-	-	-	2	3	16	71	225	643	1,155
2002.....	12,926	4	-	-	1	1	8	17	70	255	714	1,114
2001.....	12,616	3	-	-	-	-	1	14	70	296	658	1,118
2000.....	12,336	-	1	-	-	2	12	12	74	269	642	1,132
1999.....	12,224	4	-	-	-	5	1	17	94	278	614	966

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Fibrosis and cirrhosis of liver (K74)

White, male

2004.....	7,287	4	-	-	-	-	1	4	38	143	385	770
2003.....	7,687	1	-	-	-	1	2	7	48	145	461	837
2002.....	7,462	4	-	-	1	-	4	8	48	175	482	819
2001.....	7,244	1	-	-	-	-	1	8	46	179	435	812
2000.....	7,144	-	-	-	-	1	5	7	46	188	459	850
1999.....	7,082	4	-	-	-	3	1	14	62	191	455	738

White, female

2004.....	5,310	3	1	-	-	-	4	3	17	55	156	318
2003.....	5,566	-	-	-	-	1	1	9	23	80	182	318
2002.....	5,464	-	-	-	-	1	4	9	22	80	232	295
2001.....	5,372	2	-	-	-	-	-	6	24	117	223	306
2000.....	5,192	-	1	-	-	1	7	5	28	81	183	282
1999.....	5,142	-	-	-	-	2	-	3	32	87	159	228

Black, both sexes

2004.....	1,358	2	-	-	-	-	2	2	11	40	105	186
2003.....	1,387	1	-	-	-	-	3	12	10	34	103	194
2002.....	1,338	1	-	-	1	-	1	8	12	37	99	224
2001.....	1,379	-	-	1	-	2	3	9	7	41	128	244
2000.....	1,309	2	-	-	-	3	6	9	8	49	109	189
1999.....	1,306	-	2	-	-	-	4	8	12	40	126	218

Black, male

2004.....	852	2	-	-	-	-	-	1	6	25	67	131
2003.....	857	1	-	-	-	-	2	7	4	21	58	115
2002.....	812	1	-	-	-	-	-	4	8	23	55	136
2001.....	899	-	-	1	-	1	1	8	6	26	84	173
2000.....	799	2	-	-	-	1	4	6	3	30	63	124
1999.....	811	-	-	-	-	-	1	2	9	26	90	159

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Fibrosis and cirrhosis of liver (K74)												
Black, female												
2004.....	506	-	-	-	-	-	2	1	5	15	38	55
2003.....	530	-	-	-	-	-	1	5	6	13	45	79
2002.....	526	-	-	-	1	-	1	4	4	14	44	88
2001.....	480	-	-	-	-	1	2	1	1	15	44	71
2000.....	510	-	-	-	-	2	2	3	5	19	46	65
1999.....	495	-	2	-	-	-	3	6	3	14	36	59
American Indian 1/, both sexes												
2004.....	185	-	-	-	-	-	1	-	-	7	17	24
2003.....	166	-	-	-	-	-	2	1	3	3	16	27
2002.....	163	-	-	-	-	-	-	2	5	13	17	18
2001.....	172	-	-	-	-	1	-	2	3	10	15	27
2000.....	188	-	-	-	-	1	-	3	11	14	23	18
1999.....	151	-	-	-	-	-	-	-	1	9	8	32
American Indian 1/, male												
2004.....	83	-	-	-	-	-	-	-	-	5	8	14
2003.....	75	-	-	-	-	-	-	1	2	1	10	14
2002.....	88	-	-	-	-	-	-	1	1	8	8	13
2001.....	91	-	-	-	-	1	-	2	2	7	7	14
2000.....	92	-	-	-	-	1	-	1	7	7	15	11
1999.....	74	-	-	-	-	-	-	-	1	6	5	20
American Indian 1/, female												
2004.....	102	-	-	-	-	-	1	-	-	2	9	10
2003.....	91	-	-	-	-	-	2	-	1	2	6	13
2002.....	75	-	-	-	-	-	-	1	4	5	9	5
2001.....	81	-	-	-	-	-	-	-	1	3	8	13
2000.....	96	-	-	-	-	-	-	2	4	7	8	7
1999.....	77	-	-	-	-	-	-	-	-	3	3	12

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Fibrosis and cirrhosis of liver (K74)

Asian or Pacific Islander 2/, both sexes

2004.....	206	-	-	-	-	-	1	3	3	10	9
2003.....	193	-	-	-	-	-	-	-	3	7	18
2002.....	175	-	-	-	-	-	-	-	4	5	10
2001.....	194	-	-	-	-	-	-	2	3	11	10
2000.....	170	1	-	-	-	-	-	5	1	6	13
1999.....	181	1	-	-	-	-	-	1	3	9	14

Asian or Pacific Islander 2/, male

2004.....	104	-	-	-	-	-	1	3	3	8	8
2003.....	113	-	-	-	-	-	-	-	3	5	11
2002.....	80	-	-	-	-	-	-	-	4	3	5
2001.....	100	-	-	-	-	-	-	2	3	5	8
2000.....	89	1	-	-	-	-	-	4	1	4	9
1999.....	98	-	-	-	-	-	-	-	2	7	12

Asian or Pacific Islander 2/, female

2004.....	102	-	-	-	-	-	-	-	-	2	1
2003.....	80	-	-	-	-	-	-	-	-	2	7
2002.....	95	-	-	-	-	-	-	-	-	2	5
2001.....	94	-	-	-	-	-	-	-	-	6	2
2000.....	81	-	-	-	-	-	-	1	-	2	4
1999.....	83	1	-	-	-	-	-	1	1	2	2

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other diseases of liver (K71-K72,K75-K76)

All races, both sexes

2004.....	9,009	92	30	8	5	20	44	59	117	246	547	929
2003.....	9,020	59	36	6	5	29	38	64	127	263	597	935
2002.....	8,414	83	24	9	14	13	31	57	108	242	610	997
2001.....	7,805	75	28	14	14	17	35	35	106	246	487	819
2000.....	7,301	73	24	11	11	7	39	37	106	276	490	727
1999.....	7,360	81	21	10	15	22	30	42	119	264	496	688

All races, male

2004.....	5,114	47	15	4	1	7	26	35	68	140	321	602
2003.....	5,152	42	20	3	2	12	15	32	67	142	341	623
2002.....	5,002	57	10	5	9	6	15	28	66	135	384	686
2001.....	4,526	46	15	8	8	8	18	13	54	151	313	548
2000.....	4,137	49	14	6	4	4	23	19	56	170	313	501
1999.....	4,235	53	11	5	3	13	10	23	75	155	321	475

All races, female

2004.....	3,895	45	15	4	4	13	18	24	49	106	226	327
2003.....	3,868	17	16	3	3	17	23	32	60	121	256	312
2002.....	3,412	26	14	4	5	7	16	29	42	107	226	311
2001.....	3,279	29	13	6	6	9	17	22	52	95	174	271
2000.....	3,164	24	10	5	7	3	16	18	50	106	177	226
1999.....	3,125	28	10	5	12	9	20	19	44	109	175	213

White, both sexes

2004.....	7,425	52	23	6	3	13	25	38	91	180	430	723
2003.....	7,500	36	26	6	4	15	21	39	92	208	484	749
2002.....	6,983	55	16	5	9	8	19	38	80	187	487	776
2001.....	6,471	48	21	11	8	10	22	23	74	178	379	643
2000.....	6,001	48	13	6	9	5	24	22	75	206	390	563
1999.....	6,053	46	18	7	12	16	20	30	82	194	376	521

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other diseases of liver (K71-K72,K75-K76)

White, male

2004.....	4,236	26	11	2	1	6	15	24	53	105	258	475
2003.....	4,299	27	16	3	1	7	8	19	54	113	292	508
2002.....	4,175	38	6	2	7	2	12	19	49	112	316	544
2001.....	3,762	30	14	6	3	6	12	8	34	109	244	437
2000.....	3,408	30	7	4	4	4	14	12	38	132	257	395
1999.....	3,533	26	10	2	2	11	6	17	56	118	258	369

White, female

2004.....	3,189	26	12	4	2	7	10	14	38	75	172	248
2003.....	3,201	9	10	3	3	8	13	20	38	95	192	241
2002.....	2,808	17	10	3	2	6	7	19	31	75	171	232
2001.....	2,709	18	7	5	5	4	10	15	40	69	135	206
2000.....	2,593	18	6	2	5	1	10	10	37	74	133	168
1999.....	2,520	20	8	5	10	5	14	13	26	76	118	152

Black, both sexes

2004.....	1,242	34	5	2	2	5	16	17	19	51	89	178
2003.....	1,207	19	7	-	1	12	17	20	27	41	87	156
2002.....	1,144	24	6	4	4	5	12	13	17	43	96	178
2001.....	1,102	22	5	2	4	7	10	7	24	52	87	157
2000.....	1,080	24	9	4	2	2	12	12	28	59	92	141
1999.....	1,086	30	2	2	3	6	9	9	29	58	102	147

Black, male

2004.....	703	17	2	2	-	1	10	9	12	27	48	108
2003.....	692	12	3	-	1	3	7	10	11	20	36	94
2002.....	649	18	2	3	2	4	3	4	10	16	49	111
2001.....	633	13	-	1	3	2	4	2	15	30	55	99
2000.....	618	18	5	2	-	-	7	5	16	31	52	90
1999.....	583	22	-	2	1	2	3	4	12	31	53	95

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other diseases of liver (K71-K72,K75-K76)

Black, female

2004.....	539	17	3	-	2	4	6	8	7	24	41	70
2003.....	515	7	4	-	-	9	10	10	16	21	51	62
2002.....	495	6	4	1	2	1	9	9	7	27	47	67
2001.....	469	9	5	1	1	5	6	5	9	22	32	58
2000.....	462	6	4	2	2	2	5	7	12	28	40	51
1999.....	503	8	2	-	2	4	6	5	17	27	49	52

American Indian 1/, both sexes

2004.....	159	1	1	-	-	-	1	3	7	8	19	20
2003.....	155	-	1	-	-	2	-	2	6	12	22	18
2002.....	134	1	1	-	1	-	-	2	6	8	18	25
2001.....	117	1	-	-	2	-	1	4	6	10	11	12
2000.....	89	-	-	-	-	-	1	-	1	8	4	12
1999.....	103	1	-	-	-	-	1	1	5	9	14	10

American Indian 1/, male

2004.....	75	1	1	-	-	-	-	1	3	2	8	14
2003.....	82	-	-	-	-	2	-	2	1	7	10	13
2002.....	81	-	1	-	-	-	-	2	3	6	11	15
2001.....	67	1	-	-	2	-	1	2	3	7	6	9
2000.....	43	-	-	-	-	-	1	-	1	4	2	8
1999.....	54	1	-	-	-	-	1	1	4	3	7	5

American Indian 1/, female

2004.....	84	-	-	-	-	-	1	2	4	6	11	6
2003.....	73	-	1	-	-	-	-	-	5	5	12	5
2002.....	53	1	-	-	1	-	-	-	3	2	7	10
2001.....	50	-	-	-	-	-	-	2	3	3	5	3
2000.....	46	-	-	-	-	-	-	-	-	4	2	4
1999.....	49	-	-	-	-	-	-	-	1	6	7	5

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of liver (K71-K72,K75-K76)												
Asian or Pacific Islander 2/, both sexes												
2004.....	183	5	1	-	-	2	2	1	-	7	9	8
2003.....	158	4	2	-	-	-	-	3	2	2	4	12
2002.....	153	3	1	-	-	-	-	4	5	4	9	18
2001.....	115	4	2	1	-	-	2	1	2	6	10	7
2000.....	131	1	2	1	-	-	2	3	2	3	4	11
1999.....	118	4	1	1	-	-	-	2	3	3	4	10
Asian or Pacific Islander 2/, male												
2004.....	100	3	1	-	-	-	1	1	-	6	7	5
2003.....	79	3	1	-	-	-	-	1	1	2	3	8
2002.....	97	1	1	-	-	-	-	3	4	1	8	16
2001.....	64	2	1	1	-	-	1	1	2	5	8	3
2000.....	68	1	2	-	-	-	1	2	1	3	2	8
1999.....	65	4	1	1	-	-	-	1	3	3	3	6
Asian or Pacific Islander 2/, female												
2004.....	83	2	-	-	-	2	1	-	-	1	2	3
2003.....	79	1	1	-	-	-	-	2	1	-	1	4
2002.....	56	2	-	-	-	-	-	1	1	3	1	2
2001.....	51	2	1	-	-	-	1	-	-	1	2	4
2000.....	63	-	-	1	-	-	1	1	1	-	2	3
1999.....	53	-	-	-	-	-	-	1	-	-	1	4

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	-------------	-----------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Cholelithiasis and other disorders of gallbladder (K80-K82)

All races, both sexes

2004.....	3,086	-	-	-	-	1	7	8	9	17	38	61
2003.....	2,948	-	-	-	1	1	3	9	16	14	26	46
2002.....	2,979	-	-	-	-	2	7	9	12	15	39	52
2001.....	2,965	1	-	-	1	2	5	11	8	21	23	69
2000.....	2,810	-	-	1	1	2	7	6	13	22	31	43
1999.....	2,830	2	-	-	1	4	6	5	11	25	34	55

All races, male

2004.....	1,370	-	-	-	-	-	3	4	6	11	19	32
2003.....	1,318	-	-	-	-	1	3	3	8	7	14	20
2002.....	1,341	-	-	-	-	-	3	4	7	7	22	35
2001.....	1,341	1	-	-	-	1	2	5	4	9	16	39
2000.....	1,244	-	-	-	1	1	3	2	5	12	15	23
1999.....	1,248	1	-	-	1	2	3	-	4	14	16	28

All races, female

2004.....	1,716	-	-	-	-	1	4	4	3	6	19	29
2003.....	1,630	-	-	-	1	-	-	6	8	7	12	26
2002.....	1,638	-	-	-	-	2	4	5	5	8	17	17
2001.....	1,624	-	-	-	1	1	3	6	4	12	7	30
2000.....	1,566	-	-	1	-	1	4	4	8	10	16	20
1999.....	1,582	1	-	-	-	2	3	5	7	11	18	27

White, both sexes

2004.....	2,672	-	-	-	-	1	5	6	4	13	32	43
2003.....	2,573	-	-	-	1	1	2	8	10	11	17	33
2002.....	2,614	-	-	-	-	2	5	6	10	9	34	38
2001.....	2,567	1	-	-	1	2	3	9	6	14	19	49
2000.....	2,457	-	-	-	1	1	6	3	7	17	21	26
1999.....	2,487	-	-	-	1	3	3	4	9	17	20	43

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years years

Cholelithiasis and other disorders of gallbladder (K80-K82)

White, male

2004.....	1,205	-	-	-	-	-	3	4	3	8	16	22
2003.....	1,150	-	-	-	-	1	2	2	4	6	7	16
2002.....	1,179	-	-	-	-	-	3	2	6	5	19	24
2001.....	1,157	1	-	-	-	1	1	3	3	7	13	26
2000.....	1,111	-	-	-	1	1	2	2	3	10	12	16
1999.....	1,101	-	-	-	1	1	3	-	2	10	10	21

White, female

2004.....	1,467	-	-	-	-	1	2	2	1	5	16	21
2003.....	1,423	-	-	-	1	-	-	6	6	5	10	17
2002.....	1,435	-	-	-	-	2	2	4	4	4	15	14
2001.....	1,410	-	-	-	1	1	2	6	3	7	6	23
2000.....	1,346	-	-	-	-	-	4	1	4	7	9	10
1999.....	1,386	-	-	-	-	2	-	4	7	7	10	22

Black, both sexes

2004.....	317	-	-	-	-	-	-	-	4	3	6	15
2003.....	286	-	-	-	-	-	1	1	6	3	8	13
2002.....	273	-	-	-	-	-	1	3	2	6	3	10
2001.....	310	-	-	-	-	-	2	2	2	7	4	14
2000.....	280	-	-	1	-	1	1	3	5	4	10	14
1999.....	257	2	-	-	-	1	3	1	2	7	13	8

Black, male

2004.....	122	-	-	-	-	-	-	-	2	2	3	9
2003.....	127	-	-	-	-	-	1	1	4	1	7	4
2002.....	110	-	-	-	-	-	-	2	1	2	2	8
2001.....	130	-	-	-	-	-	1	2	1	2	3	9
2000.....	96	-	-	-	-	-	1	-	2	1	3	5
1999.....	102	1	-	-	-	1	-	-	2	3	6	5

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	-------------	-----------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Disorders of biliary tract and pancreas (K83-K86)

All races, both sexes

2004.....	4,883	13	4	2	9	9	31	50	78	139	212	290
2003.....	4,938	8	4	5	6	11	18	41	83	135	222	273
2002.....	4,912	12	2	1	3	16	33	41	93	135	222	280
2001.....	4,958	7	2	2	6	10	21	46	98	165	228	301
2000.....	4,653	9	5	8	6	4	35	55	82	149	234	268
1999.....	4,447	11	4	1	12	16	29	50	92	131	229	251

All races, male

2004.....	2,567	8	3	-	2	2	19	38	66	108	141	194
2003.....	2,562	4	1	2	2	9	9	29	60	90	141	181
2002.....	2,647	9	-	1	1	10	22	31	67	92	146	202
2001.....	2,511	4	2	1	4	6	15	29	63	108	148	196
2000.....	2,428	5	4	2	1	1	22	38	59	103	163	174
1999.....	2,302	7	1	1	4	9	14	36	62	87	153	172

All races, female

2004.....	2,316	5	1	2	7	7	12	12	12	31	71	96
2003.....	2,376	4	3	3	4	2	9	12	23	45	81	92
2002.....	2,265	3	2	-	2	6	11	10	26	43	76	78
2001.....	2,447	3	-	1	2	4	6	17	35	57	80	105
2000.....	2,225	4	1	6	5	3	13	17	23	46	71	94
1999.....	2,145	4	3	-	8	7	15	14	30	44	76	79

White, both sexes

2004.....	4,055	8	3	2	6	6	19	35	55	99	157	219
2003.....	4,097	4	4	4	5	8	13	23	63	92	147	193
2002.....	4,147	10	2	1	1	11	23	30	61	101	166	201
2001.....	4,137	4	2	2	3	8	11	34	59	115	164	217
2000.....	3,874	3	4	5	5	4	23	34	60	113	173	190
1999.....	3,703	7	3	1	9	11	17	37	65	91	176	168

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of biliary tract and pancreas (K83-K86)												
White, male												
2004.....	2,109	4	3	-	2	1	12	26	45	78	114	147
2003.....	2,087	2	1	1	2	6	8	18	47	61	89	136
2002.....	2,202	7	-	1	-	8	15	22	45	66	115	145
2001.....	2,069	2	2	1	1	5	8	21	40	78	111	139
2000.....	1,988	1	3	2	1	1	16	25	44	78	120	126
1999.....	1,880	4	1	1	3	5	9	25	43	61	118	115
White, female												
2004.....	1,946	4	-	2	4	5	7	9	10	21	43	72
2003.....	2,010	2	3	3	3	2	5	5	16	31	58	57
2002.....	1,945	3	2	-	1	3	8	8	16	35	51	56
2001.....	2,068	2	-	1	2	3	3	13	19	37	53	78
2000.....	1,886	2	1	3	4	3	7	9	16	35	53	64
1999.....	1,823	3	2	-	6	6	8	12	22	30	58	53
Black, both sexes												
2004.....	704	5	1	-	2	2	7	10	20	35	52	66
2003.....	710	3	-	1	1	2	3	14	16	36	67	75
2002.....	648	2	-	-	2	5	8	10	24	32	51	74
2001.....	707	1	-	-	2	2	8	11	32	48	63	80
2000.....	670	5	1	3	-	-	11	20	22	28	57	73
1999.....	641	4	1	-	2	5	11	13	22	38	47	75
Black, male												
2004.....	393	4	-	-	-	1	3	7	18	26	25	44
2003.....	403	2	-	1	-	2	-	8	10	23	49	43
2002.....	377	2	-	-	1	2	7	8	15	24	28	53
2001.....	384	-	-	-	2	1	6	7	19	28	36	54
2000.....	379	3	1	-	-	-	5	12	15	20	40	43
1999.....	370	3	-	-	1	4	4	11	18	26	32	53

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of biliary tract and pancreas (K83-K86)												
Black, female												
2004.....	311	1	1	-	2	1	4	3	2	9	27	22
2003.....	307	1	-	-	1	-	3	6	6	13	18	32
2002.....	271	-	-	-	1	3	1	2	9	8	23	21
2001.....	323	1	-	-	-	1	2	4	13	20	27	26
2000.....	291	2	-	3	-	-	6	8	7	8	17	30
1999.....	271	1	1	-	1	1	7	2	4	12	15	22
American Indian 1/, both sexes												
2004.....	29	-	-	-	1	1	1	2	-	2	1	1
2003.....	32	-	-	-	-	-	-	2	1	5	5	2
2002.....	37	-	-	-	-	-	-	-	5	1	3	3
2001.....	37	-	-	-	-	-	-	-	4	1	1	3
2000.....	27	1	-	-	1	-	-	-	-	2	2	2
1999.....	31	-	-	-	-	-	1	-	1	2	3	3
American Indian 1/, male												
2004.....	15	-	-	-	-	-	1	2	-	1	1	-
2003.....	18	-	-	-	-	-	-	1	1	4	2	-
2002.....	17	-	-	-	-	-	-	-	5	1	1	2
2001.....	18	-	-	-	-	-	-	-	2	1	1	2
2000.....	14	1	-	-	-	-	-	-	-	1	1	2
1999.....	14	-	-	-	-	-	1	-	-	-	1	2
American Indian 1/, female												
2004.....	14	-	-	-	1	1	-	-	-	1	-	1
2003.....	14	-	-	-	-	-	-	1	-	1	3	2
2002.....	20	-	-	-	-	-	-	-	-	-	2	1
2001.....	19	-	-	-	-	-	-	-	2	-	-	1
2000.....	13	-	-	-	1	-	-	-	-	1	1	-
1999.....	17	-	-	-	-	-	-	-	1	2	2	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of biliary tract and pancreas (K83-K86)												
Asian or Pacific Islander 2/, both sexes												
2004.....	95	-	-	-	-	-	4	3	3	3	2	4
2003.....	99	1	-	-	-	1	2	2	3	2	3	3
2002.....	80	-	-	-	-	-	2	1	3	1	2	2
2001.....	77	2	-	-	1	-	2	1	3	1	-	1
2000.....	82	-	-	-	-	-	1	1	-	6	2	3
1999.....	72	-	-	-	1	-	-	-	4	-	3	5
Asian or Pacific Islander 2/, male												
2004.....	50	-	-	-	-	-	3	3	3	3	1	3
2003.....	54	-	-	-	-	1	1	2	2	2	1	2
2002.....	51	-	-	-	-	-	-	1	2	1	2	2
2001.....	40	2	-	-	1	-	1	1	2	1	-	1
2000.....	47	-	-	-	-	-	1	1	-	4	2	3
1999.....	38	-	-	-	-	-	-	-	1	-	2	2
Asian or Pacific Islander 2/, female												
2004.....	45	-	-	-	-	-	1	-	-	-	1	1
2003.....	45	1	-	-	-	-	1	-	1	-	2	1
2002.....	29	-	-	-	-	-	2	-	1	-	-	-
2001.....	37	-	-	-	-	-	1	-	1	-	-	-
2000.....	35	-	-	-	-	-	-	-	-	2	-	-
1999.....	34	-	-	-	1	-	-	-	3	-	1	3

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pancreatitis (K85,K86.0-K86.1)												
All races, both sexes												
2004.....	3,475	3	2	1	9	8	26	48	70	123	191	264
2003.....	3,490	4	1	4	6	8	16	34	72	119	199	238
2002.....	3,532	2	2	1	2	14	31	36	80	125	197	251
2001.....	3,648	1	1	2	6	9	20	39	85	144	209	257
2000.....	3,377	3	1	8	6	4	35	47	79	136	211	225
1999.....	3,289	1	1	1	10	12	26	47	88	121	206	223
All races, male												
2004.....	1,928	2	1	-	2	2	17	36	59	99	132	183
2003.....	1,904	2	1	2	2	7	8	24	55	81	131	163
2002.....	1,985	1	-	1	-	9	20	27	61	86	131	185
2001.....	1,926	1	1	1	4	5	14	27	56	96	136	176
2000.....	1,830	1	1	2	1	1	22	31	58	100	153	141
1999.....	1,769	1	-	1	3	6	12	34	59	82	142	152
All races, female												
2004.....	1,547	1	1	1	7	6	9	12	11	24	59	81
2003.....	1,586	2	-	2	4	1	8	10	17	38	68	75
2002.....	1,547	1	2	-	2	5	11	9	19	39	66	66
2001.....	1,722	-	-	1	2	4	6	12	29	48	73	81
2000.....	1,547	2	-	6	5	3	13	16	21	36	58	84
1999.....	1,520	-	1	-	7	6	14	13	29	39	64	71
White, both sexes												
2004.....	2,833	3	1	1	6	5	17	34	52	89	142	197
2003.....	2,843	2	1	3	5	6	11	20	55	83	129	168
2002.....	2,968	2	2	1	1	9	21	28	54	93	151	181
2001.....	3,014	1	1	2	3	8	10	29	52	104	149	184
2000.....	2,778	1	1	5	5	4	23	30	57	103	154	157
1999.....	2,702	1	1	1	8	8	15	35	63	84	158	147

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Pancreatitis (K85,K86.0-K86.1)

White, male

2004.....	1,550	2	1	-	2	1	11	25	43	72	106	138
2003.....	1,511	1	1	1	2	5	7	16	42	56	83	120
2002.....	1,633	1	-	1	-	7	13	20	40	62	103	135
2001.....	1,569	1	1	1	1	5	7	19	36	71	100	125
2000.....	1,472	-	1	2	1	1	16	21	43	76	110	101
1999.....	1,423	1	-	1	3	3	7	24	42	57	108	100

White, female

2004.....	1,283	1	-	1	4	4	6	9	9	17	36	59
2003.....	1,332	1	-	2	3	1	4	4	13	27	46	48
2002.....	1,335	1	2	-	1	2	8	8	14	31	48	46
2001.....	1,445	-	-	1	2	3	3	10	16	33	49	59
2000.....	1,306	1	-	3	4	3	7	9	14	27	44	56
1999.....	1,279	-	1	-	5	5	8	11	21	27	50	47

Black, both sexes

2004.....	557	-	1	-	2	2	4	9	16	29	46	62
2003.....	552	2	-	1	1	2	3	10	14	30	63	66
2002.....	487	-	-	-	1	5	8	7	19	31	42	65
2001.....	556	-	-	-	2	1	8	9	26	38	59	70
2000.....	520	1	-	3	-	-	11	16	22	26	54	63
1999.....	513	-	-	-	1	4	10	12	20	35	43	68

Black, male

2004.....	327	-	-	-	-	1	2	6	14	23	24	42
2003.....	337	1	-	1	-	2	-	5	10	20	45	41
2002.....	302	-	-	-	-	2	7	6	14	23	25	46
2001.....	315	-	-	-	2	-	6	7	16	23	35	48
2000.....	310	-	-	-	-	-	5	9	15	20	40	35
1999.....	309	-	-	-	-	3	4	10	16	25	31	48

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pancreatitis (K85,K86.0-K86.1)												
Black, female												
2004.....	230	-	1	-	2	1	2	3	2	6	22	20
2003.....	215	1	-	-	1	-	3	5	4	10	18	25
2002.....	185	-	-	-	1	3	1	1	5	8	17	19
2001.....	241	-	-	-	-	1	2	2	10	15	24	22
2000.....	210	1	-	3	-	-	6	7	7	6	14	28
1999.....	204	-	-	-	1	1	6	2	4	10	12	20
American Indian 1/, both sexes												
2004.....	24	-	-	-	1	1	1	2	-	2	1	1
2003.....	27	-	-	-	-	-	-	2	1	4	5	1
2002.....	27	-	-	-	-	-	-	-	5	-	2	3
2001.....	26	-	-	-	-	-	-	-	4	1	1	2
2000.....	20	1	-	-	1	-	-	-	-	2	1	2
1999.....	26	-	-	-	-	-	1	-	1	2	3	3
American Indian 1/, male												
2004.....	14	-	-	-	-	-	1	2	-	1	1	-
2003.....	17	-	-	-	-	-	-	1	1	3	2	-
2002.....	13	-	-	-	-	-	-	-	5	-	1	2
2001.....	15	-	-	-	-	-	-	-	2	1	1	2
2000.....	11	1	-	-	-	-	-	-	-	1	1	2
1999.....	12	-	-	-	-	-	1	-	-	-	1	2
American Indian 1/, female												
2004.....	10	-	-	-	1	1	-	-	-	1	-	1
2003.....	10	-	-	-	-	-	-	1	-	1	3	1
2002.....	14	-	-	-	-	-	-	-	-	-	1	1
2001.....	11	-	-	-	-	-	-	-	2	-	-	-
2000.....	9	-	-	-	1	-	-	-	-	1	-	-
1999.....	14	-	-	-	-	-	-	-	1	2	2	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pancreatitis (K85,K86.0-K86.1)												
Asian or Pacific Islander 2/, both sexes												
2004.....	61	-	-	-	-	-	4	3	2	3	2	4
2003.....	68	-	-	-	-	-	2	2	2	2	2	3
2002.....	50	-	-	-	-	-	2	1	2	1	2	2
2001.....	52	-	-	-	1	-	2	1	3	1	-	1
2000.....	59	-	-	-	-	-	1	1	-	5	2	3
1999.....	48	-	-	-	1	-	-	-	4	-	2	5
Asian or Pacific Islander 2/, male												
2004.....	37	-	-	-	-	-	3	3	2	3	1	3
2003.....	39	-	-	-	-	-	1	2	2	2	1	2
2002.....	37	-	-	-	-	-	-	1	2	1	2	2
2001.....	27	-	-	-	1	-	1	1	2	1	-	1
2000.....	37	-	-	-	-	-	1	1	-	3	2	3
1999.....	25	-	-	-	-	-	-	-	1	-	2	2
Asian or Pacific Islander 2/, female												
2004.....	24	-	-	-	-	-	1	-	-	-	1	1
2003.....	29	-	-	-	-	-	1	-	-	-	1	1
2002.....	13	-	-	-	-	-	2	-	-	-	-	-
2001.....	25	-	-	-	-	-	1	-	1	-	-	-
2000.....	22	-	-	-	-	-	-	-	-	2	-	-
1999.....	23	-	-	-	1	-	-	-	3	-	-	3

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other disorders of biliary tract and pancreas (K83,K86.2-K86.9)

All races, both sexes

2004.....	1,408	10	2	1	-	1	5	2	8	16	21	26
2003.....	1,448	4	3	1	-	3	2	7	11	16	23	35
2002.....	1,380	10	-	-	1	2	2	5	13	10	25	29
2001.....	1,310	6	1	-	-	1	1	7	13	21	19	44
2000.....	1,276	6	4	-	-	-	-	8	3	13	23	43
1999.....	1,158	10	3	-	2	4	3	3	4	10	23	28

All races, male

2004.....	639	6	2	-	-	-	2	2	7	9	9	11
2003.....	658	2	-	-	-	2	1	5	5	9	10	18
2002.....	662	8	-	-	1	1	2	4	6	6	15	17
2001.....	585	3	1	-	-	1	1	2	7	12	12	20
2000.....	598	4	3	-	-	-	-	7	1	3	10	33
1999.....	533	6	1	-	1	3	2	2	3	5	11	20

All races, female

2004.....	769	4	-	1	-	1	3	-	1	7	12	15
2003.....	790	2	3	1	-	1	1	2	6	7	13	17
2002.....	718	2	-	-	-	1	-	1	7	4	10	12
2001.....	725	3	-	-	-	-	-	5	6	9	7	24
2000.....	678	2	1	-	-	-	-	1	2	10	13	10
1999.....	625	4	2	-	1	1	1	1	1	5	12	8

White, both sexes

2004.....	1,222	5	2	1	-	1	2	1	3	10	15	22
2003.....	1,254	2	3	1	-	2	2	3	8	9	18	25
2002.....	1,179	8	-	-	-	2	2	2	7	8	15	20
2001.....	1,123	3	1	-	-	-	1	5	7	11	15	33
2000.....	1,096	2	3	-	-	-	-	4	3	10	19	33
1999.....	1,001	6	2	-	1	3	2	2	2	7	18	21

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other disorders of biliary tract and pancreas (K83,K86.2-K86.9)

White, male

2004.....	559	2	2	-	-	-	1	1	2	6	8	9
2003.....	576	1	-	-	-	1	1	2	5	5	6	16
2002.....	569	6	-	-	-	1	2	2	5	4	12	10
2001.....	500	1	1	-	-	-	1	2	4	7	11	14
2000.....	516	1	2	-	-	-	-	4	1	2	10	25
1999.....	457	3	1	-	-	2	2	1	1	4	10	15

White, female

2004.....	663	3	-	1	-	1	1	-	1	4	7	13
2003.....	678	1	3	1	-	1	1	1	3	4	12	9
2002.....	610	2	-	-	-	1	-	-	2	4	3	10
2001.....	623	2	-	-	-	-	-	3	3	4	4	19
2000.....	580	1	1	-	-	-	-	-	2	8	9	8
1999.....	544	3	1	-	1	1	-	1	1	3	8	6

Black, both sexes

2004.....	147	5	-	-	-	-	3	1	4	6	6	4
2003.....	158	1	-	-	-	-	-	4	2	6	4	9
2002.....	161	2	-	-	1	-	-	3	5	1	9	9
2001.....	151	1	-	-	-	1	-	2	6	10	4	10
2000.....	150	4	1	-	-	-	-	4	-	2	3	10
1999.....	128	4	1	-	1	1	1	1	2	3	4	7

Black, male

2004.....	66	4	-	-	-	-	1	1	4	3	1	2
2003.....	66	1	-	-	-	-	-	3	-	3	4	2
2002.....	75	2	-	-	1	-	-	2	1	1	3	7
2001.....	69	-	-	-	-	1	-	-	3	5	1	6
2000.....	69	3	1	-	-	-	-	3	-	-	-	8
1999.....	61	3	-	-	1	1	-	1	2	1	1	5

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

All other diseases of digestive system (K90,K92)

All races, both sexes

2004.....	7,943	28	11	6	3	6	12	7	16	43	76	153
2003.....	8,265	30	17	5	4	6	6	6	16	53	101	141
2002.....	8,287	25	21	3	4	5	11	8	23	26	92	165
2001.....	8,209	25	15	4	3	-	4	12	24	38	94	144
2000.....	8,091	24	17	3	2	5	6	13	15	40	91	133
1999.....	7,760	26	20	3	5	2	6	8	15	33	82	128

All races, male

2004.....	3,606	14	8	3	-	3	8	3	9	25	47	113
2003.....	3,691	15	9	4	3	1	4	4	11	32	73	92
2002.....	3,674	16	13	3	3	3	8	6	14	18	69	124
2001.....	3,646	11	12	-	1	-	4	7	18	28	66	103
2000.....	3,548	18	14	2	1	2	4	9	13	23	67	97
1999.....	3,423	15	9	1	1	-	6	7	9	24	55	92

All races, female

2004.....	4,337	14	3	3	3	3	4	4	7	18	29	40
2003.....	4,574	15	8	1	1	5	2	2	5	21	28	49
2002.....	4,613	9	8	-	1	2	3	2	9	8	23	41
2001.....	4,563	14	3	4	2	-	-	5	6	10	28	41
2000.....	4,543	6	3	1	1	3	2	4	2	17	24	36
1999.....	4,337	11	11	2	4	2	-	1	6	9	27	36

White, both sexes

2004.....	6,941	15	5	4	2	6	10	3	11	38	61	125
2003.....	7,234	21	10	5	3	6	3	4	12	43	77	110
2002.....	7,256	19	12	2	4	4	8	3	14	18	64	120
2001.....	7,179	13	7	4	3	-	3	6	19	29	63	109
2000.....	7,034	18	7	3	1	4	5	9	10	26	63	95
1999.....	6,821	17	14	3	4	2	4	6	12	28	60	98

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

All other diseases of digestive system (K90,K92)

White, male

2004.....	3,120	7	3	2	-	3	7	1	7	24	38	94
2003.....	3,200	10	6	4	2	1	3	2	9	25	52	71
2002.....	3,163	11	6	2	3	2	5	2	9	13	47	88
2001.....	3,133	5	5	-	1	-	3	5	15	21	46	76
2000.....	3,048	15	6	2	1	1	3	8	9	15	48	70
1999.....	2,973	10	5	1	1	-	4	6	7	20	45	71

White, female

2004.....	3,821	8	2	2	2	3	3	2	4	14	23	31
2003.....	4,034	11	4	1	1	5	-	2	3	18	25	39
2002.....	4,093	8	6	-	1	2	3	1	5	5	17	32
2001.....	4,046	8	2	4	2	-	-	1	4	8	17	33
2000.....	3,986	3	1	1	-	3	2	1	1	11	15	25
1999.....	3,848	7	9	2	3	2	-	-	5	8	15	27

Black, both sexes

2004.....	877	12	5	1	-	-	2	4	5	5	14	26
2003.....	895	7	7	-	-	-	3	2	2	10	21	28
2002.....	896	6	9	1	-	-	2	4	7	8	23	41
2001.....	911	11	7	-	-	-	-	6	5	8	25	31
2000.....	937	6	7	-	1	1	1	4	5	14	25	37
1999.....	816	8	4	-	1	-	1	2	2	5	19	28

Black, male

2004.....	423	7	5	-	-	-	1	2	2	1	9	17
2003.....	416	4	3	-	-	-	1	2	1	7	18	19
2002.....	434	5	7	1	-	-	2	3	4	5	17	32
2001.....	444	6	6	-	-	-	-	2	3	6	16	25
2000.....	435	3	6	-	-	1	1	1	4	8	17	26
1999.....	381	4	2	-	-	-	1	1	1	4	9	19

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

XII. Diseases of the skin and subcutaneous tissue (L00-L98)

All races, both sexes

2004.....	4,180	5	3	-	2	13	19	28	26	55	81	129
2003.....	4,132	2	6	5	2	6	13	25	49	57	83	116
2002.....	3,937	1	4	2	3	8	13	18	32	43	85	100
2001.....	3,749	2	2	3	4	9	6	22	27	49	65	108
2000.....	3,753	3	4	-	3	3	15	23	21	40	67	85
1999.....	3,697	2	2	4	3	5	12	11	30	41	66	98

All races, male

2004.....	1,540	5	1	-	1	6	7	11	9	28	37	70
2003.....	1,501	2	1	3	-	-	1	5	13	25	40	56
2002.....	1,439	1	4	2	2	2	1	4	12	19	36	54
2001.....	1,318	1	2	1	1	3	1	9	7	21	27	58
2000.....	1,301	3	2	-	1	1	4	5	9	16	33	38
1999.....	1,296	2	1	1	-	3	5	2	16	21	22	50

All races, female

2004.....	2,640	-	2	-	1	7	12	17	17	27	44	59
2003.....	2,631	-	5	2	2	6	12	20	36	32	43	60
2002.....	2,498	-	-	-	1	6	12	14	20	24	49	46
2001.....	2,431	1	-	2	3	6	5	13	20	28	38	50
2000.....	2,452	-	2	-	2	2	11	18	12	24	34	47
1999.....	2,401	-	1	3	3	2	7	9	14	20	44	48

White, both sexes

2004.....	3,186	4	3	-	2	6	8	9	9	31	57	92
2003.....	3,129	1	4	4	1	3	6	9	21	31	58	74
2002.....	2,902	1	3	2	-	2	5	4	19	17	53	70
2001.....	2,822	2	-	2	2	5	3	9	15	22	36	72
2000.....	2,777	1	3	-	-	2	8	12	12	22	41	49
1999.....	2,679	2	2	2	3	3	5	8	19	23	36	58

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years years

XII. Diseases of the skin and subcutaneous tissue (L00-L98)

White, male

2004.....	1,195	4	1	-	1	3	4	5	4	17	28	49
2003.....	1,130	1	-	2	-	-	1	1	6	13	31	37
2002.....	1,107	1	3	2	-	-	1	1	8	10	26	44
2001.....	998	1	-	1	-	2	1	5	3	12	16	40
2000.....	961	1	1	-	-	1	3	1	6	10	19	29
1999.....	944	2	1	-	-	2	3	2	10	13	12	27

White, female

2004.....	1,991	-	2	-	1	3	4	4	5	14	29	43
2003.....	1,999	-	4	2	1	3	5	8	15	18	27	37
2002.....	1,795	-	-	-	-	2	4	3	11	7	27	26
2001.....	1,824	1	-	1	2	3	2	4	12	10	20	32
2000.....	1,816	-	2	-	-	1	5	11	6	12	22	20
1999.....	1,735	-	1	2	3	1	2	6	9	10	24	31

Black, both sexes

2004.....	920	1	-	-	-	7	10	18	15	22	21	35
2003.....	929	1	2	-	-	3	7	15	26	26	22	37
2002.....	962	-	-	-	3	5	7	14	13	24	31	25
2001.....	866	-	2	1	1	4	2	13	10	24	24	31
2000.....	896	1	1	-	2	-	6	11	9	17	22	33
1999.....	963	-	-	1	-	2	5	3	10	17	28	39

Black, male

2004.....	308	1	-	-	-	3	2	6	4	9	8	19
2003.....	341	1	1	-	-	-	-	4	7	12	8	16
2002.....	305	-	-	-	2	2	-	3	4	9	10	7
2001.....	298	-	2	-	1	1	-	4	2	8	9	15
2000.....	310	1	1	-	1	-	1	4	3	6	13	8
1999.....	330	-	-	1	-	1	2	-	5	8	9	23

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years years

XII. Diseases of the skin and subcutaneous tissue (L00-L98)

Black, female

2004.....	612	-	-	-	-	4	8	12	11	13	13	16
2003.....	588	-	1	-	-	3	7	11	19	14	14	21
2002.....	657	-	-	-	1	3	7	11	9	15	21	18
2001.....	568	-	-	1	-	3	2	9	8	16	15	16
2000.....	586	-	-	-	1	-	5	7	6	11	9	25
1999.....	633	-	-	-	-	1	3	3	5	9	19	16

American Indian 1/, both sexes

2004.....	21	-	-	-	-	-	-	1	2	1	3	1
2003.....	27	-	-	-	-	-	-	-	1	-	2	3
2002.....	22	-	-	-	-	-	-	-	-	2	-	2
2001.....	21	-	-	-	1	-	-	-	-	2	3	3
2000.....	29	-	-	-	-	-	-	-	-	-	2	2
1999.....	22	-	-	-	-	-	-	-	1	1	2	1

American Indian 1/, male

2004.....	8	-	-	-	-	-	-	-	1	1	1	1
2003.....	7	-	-	-	-	-	-	-	-	-	-	1
2002.....	6	-	-	-	-	-	-	-	-	-	-	1
2001.....	4	-	-	-	-	-	-	-	-	-	1	1
2000.....	11	-	-	-	-	-	-	-	-	-	-	1
1999.....	5	-	-	-	-	-	-	-	1	-	1	-

American Indian 1/, female

2004.....	13	-	-	-	-	-	-	1	1	-	2	-
2003.....	20	-	-	-	-	-	-	-	1	-	2	2
2002.....	16	-	-	-	-	-	-	-	-	2	-	1
2001.....	17	-	-	-	1	-	-	-	-	2	2	2
2000.....	18	-	-	-	-	-	-	-	-	-	2	1
1999.....	17	-	-	-	-	-	-	-	-	1	1	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
XII. Diseases of the skin and subcutaneous tissue (L00-L98)												
Asian or Pacific Islander 2/, both sexes												
2004.....	53	-	-	-	-	-	1	-	-	1	-	1
2003.....	47	-	-	1	1	-	-	1	1	-	1	2
2002.....	51	-	1	-	-	1	1	-	-	-	1	3
2001.....	40	-	-	-	-	-	1	-	2	1	2	2
2000.....	51	1	-	-	1	1	1	-	-	1	2	1
1999.....	33	-	-	1	-	-	2	-	-	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	29	-	-	-	-	-	1	-	-	1	-	1
2003.....	23	-	-	1	-	-	-	-	-	-	1	2
2002.....	21	-	1	-	-	-	-	-	-	-	-	2
2001.....	18	-	-	-	-	-	-	-	2	1	1	2
2000.....	19	1	-	-	-	-	-	-	-	-	1	-
1999.....	17	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	24	-	-	-	-	-	-	-	-	-	-	-
2003.....	24	-	-	-	1	-	-	1	1	-	-	-
2002.....	30	-	-	-	-	1	1	-	-	-	1	1
2001.....	22	-	-	-	-	-	1	-	-	-	1	-
2000.....	32	-	-	-	1	1	1	-	-	1	1	1
1999.....	16	-	-	1	-	-	2	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Infections of skin and subcutaneous tissue (L00-L08)

All races, both sexes

2004.....	1,696	2	1	-	1	-	4	9	10	22	40	71
2003.....	1,587	-	-	1	-	1	3	4	13	27	29	52
2002.....	1,283	1	2	1	-	-	4	2	10	10	31	44
2001.....	1,249	-	-	1	2	3	3	5	8	21	26	56
2000.....	1,141	1	-	-	-	-	1	5	7	15	24	29
1999.....	983	-	1	-	-	1	2	2	12	12	17	46

All races, male

2004.....	720	2	-	-	1	-	3	6	5	15	24	46
2003.....	670	-	-	1	-	-	1	1	7	14	15	32
2002.....	562	1	2	1	-	-	-	1	7	7	22	32
2001.....	511	-	-	-	-	1	-	5	4	13	15	38
2000.....	481	1	-	-	-	-	-	1	4	6	13	19
1999.....	408	-	-	-	-	1	1	2	7	9	9	30

All races, female

2004.....	976	-	1	-	-	-	1	3	5	7	16	25
2003.....	917	-	-	-	-	1	2	3	6	13	14	20
2002.....	721	-	-	-	-	-	4	1	3	3	9	12
2001.....	738	-	-	1	2	2	3	-	4	8	11	18
2000.....	660	-	-	-	-	-	1	4	3	9	11	10
1999.....	575	-	1	-	-	-	1	-	5	3	8	16

White, both sexes

2004.....	1,460	2	1	-	1	-	4	3	6	15	32	56
2003.....	1,376	-	-	1	-	1	1	4	8	20	25	36
2002.....	1,100	1	1	1	-	-	-	1	8	4	21	35
2001.....	1,072	-	-	-	1	1	1	2	4	13	16	40
2000.....	989	-	-	-	-	-	-	3	6	11	17	23
1999.....	842	-	1	-	-	1	2	2	8	9	10	27

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Infections of skin and subcutaneous tissue (L00-L08)

White, male

2004.....	620	2	-	-	1	-	3	3	3	11	19	35
2003.....	571	-	-	1	-	-	1	1	4	8	14	21
2002.....	485	1	1	1	-	-	-	-	6	4	15	27
2001.....	441	-	-	-	-	-	-	2	2	9	9	28
2000.....	416	-	-	-	-	-	-	-	4	6	9	16
1999.....	343	-	-	-	-	1	1	2	5	7	7	17

White, female

2004.....	840	-	1	-	-	-	1	-	3	4	13	21
2003.....	805	-	-	-	-	1	-	3	4	12	11	15
2002.....	615	-	-	-	-	-	-	1	2	-	6	8
2001.....	631	-	-	-	1	1	1	-	2	4	7	12
2000.....	573	-	-	-	-	-	-	3	2	5	8	7
1999.....	499	-	1	-	-	-	1	-	3	2	3	10

Black, both sexes

2004.....	202	-	-	-	-	-	-	5	3	6	5	13
2003.....	180	-	-	-	-	-	2	-	5	7	3	13
2002.....	155	-	-	-	-	-	3	1	2	4	10	6
2001.....	152	-	-	1	-	2	2	3	3	6	8	12
2000.....	128	1	-	-	-	-	1	2	1	4	5	6
1999.....	124	-	-	-	-	-	-	-	3	2	6	18

Black, male

2004.....	83	-	-	-	-	-	-	3	1	3	4	9
2003.....	83	-	-	-	-	-	-	-	3	6	1	9
2002.....	65	-	-	-	-	-	-	1	1	3	7	2
2001.....	60	-	-	-	-	1	-	3	1	3	5	8
2000.....	52	1	-	-	-	-	-	1	-	-	4	3
1999.....	57	-	-	-	-	-	-	-	1	2	1	13

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Infections of skin and subcutaneous tissue (L00-L08)												
Black, female												
2004.....	119	-	-	-	-	-	-	2	2	3	1	4
2003.....	97	-	-	-	-	-	2	-	2	1	2	4
2002.....	90	-	-	-	-	-	3	-	1	1	3	4
2001.....	92	-	-	1	-	1	2	-	2	3	3	4
2000.....	76	-	-	-	-	-	1	1	1	4	1	3
1999.....	67	-	-	-	-	-	-	-	2	-	5	5
American Indian 1/, both sexes												
2004.....	14	-	-	-	-	-	-	1	1	1	3	1
2003.....	11	-	-	-	-	-	-	-	-	-	1	2
2002.....	10	-	-	-	-	-	-	-	-	2	-	1
2001.....	12	-	-	-	1	-	-	-	-	1	2	2
2000.....	11	-	-	-	-	-	-	-	-	-	1	-
1999.....	11	-	-	-	-	-	-	-	1	1	1	1
American Indian 1/, male												
2004.....	6	-	-	-	-	-	-	-	1	1	1	1
2003.....	4	-	-	-	-	-	-	-	-	-	-	1
2002.....	2	-	-	-	-	-	-	-	-	-	-	1
2001.....	3	-	-	-	-	-	-	-	-	-	1	-
2000.....	6	-	-	-	-	-	-	-	-	-	-	-
1999.....	4	-	-	-	-	-	-	-	1	-	1	-
American Indian 1/, female												
2004.....	8	-	-	-	-	-	-	1	-	-	2	-
2003.....	7	-	-	-	-	-	-	-	-	-	1	1
2002.....	8	-	-	-	-	-	-	-	-	2	-	-
2001.....	9	-	-	-	1	-	-	-	-	1	1	2
2000.....	5	-	-	-	-	-	-	-	-	-	1	-
1999.....	7	-	-	-	-	-	-	-	-	1	-	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified diseases of skin and subcutaneous tissue (L10-L98)												
All races, both sexes												
2004.....	2,484	3	2	-	1	13	15	19	16	33	41	58
2003.....	2,545	2	6	4	2	5	10	21	36	30	54	64
2002.....	2,654	-	2	1	3	8	9	16	22	33	54	56
2001.....	2,500	2	2	2	2	6	3	17	19	28	39	52
2000.....	2,612	2	4	-	3	3	14	18	14	25	43	56
1999.....	2,714	2	1	4	3	4	10	9	18	29	49	52
All races, male												
2004.....	820	3	1	-	-	6	4	5	4	13	13	24
2003.....	831	2	1	2	-	-	-	4	6	11	25	24
2002.....	877	-	2	1	2	2	1	3	5	12	14	22
2001.....	807	1	2	1	1	2	1	4	3	8	12	20
2000.....	820	2	2	-	1	1	4	4	5	10	20	19
1999.....	888	2	1	1	-	2	4	-	9	12	13	20
All races, female												
2004.....	1,664	-	1	-	1	7	11	14	12	20	28	34
2003.....	1,714	-	5	2	2	5	10	17	30	19	29	40
2002.....	1,777	-	-	-	1	6	8	13	17	21	40	34
2001.....	1,693	1	-	1	1	4	2	13	16	20	27	32
2000.....	1,792	-	2	-	2	2	10	14	9	15	23	37
1999.....	1,826	-	-	3	3	2	6	9	9	17	36	32
White, both sexes												
2004.....	1,726	2	2	-	1	6	4	6	3	16	25	36
2003.....	1,753	1	4	3	1	2	5	5	13	11	33	38
2002.....	1,802	-	2	1	-	2	5	3	11	13	32	35
2001.....	1,750	2	-	2	1	4	2	7	11	9	20	32
2000.....	1,788	1	3	-	-	2	8	9	6	11	24	26
1999.....	1,837	2	1	2	3	2	3	6	11	14	26	31

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other and unspecified diseases of skin and subcutaneous tissue (L10-L98)

White, male

2004.....	575	2	1	-	-	3	1	2	1	6	9	14
2003.....	559	1	-	1	-	-	-	-	2	5	17	16
2002.....	622	-	2	1	-	-	1	1	2	6	11	17
2001.....	557	1	-	1	-	2	1	3	1	3	7	12
2000.....	545	1	1	-	-	1	3	1	2	4	10	13
1999.....	601	2	1	-	-	1	2	-	5	6	5	10

White, female

2004.....	1,151	-	1	-	1	3	3	4	2	10	16	22
2003.....	1,194	-	4	2	1	2	5	5	11	6	16	22
2002.....	1,180	-	-	-	-	2	4	2	9	7	21	18
2001.....	1,193	1	-	1	1	2	1	4	10	6	13	20
2000.....	1,243	-	2	-	-	1	5	8	4	7	14	13
1999.....	1,236	-	-	2	3	1	1	6	6	8	21	21

Black, both sexes

2004.....	718	1	-	-	-	7	10	13	12	16	16	22
2003.....	749	1	2	-	-	3	5	15	21	19	19	24
2002.....	807	-	-	-	3	5	4	13	11	20	21	19
2001.....	714	-	2	-	1	2	-	10	7	18	16	19
2000.....	768	-	1	-	2	-	5	9	8	13	17	27
1999.....	839	-	-	1	-	2	5	3	7	15	22	21

Black, male

2004.....	225	1	-	-	-	3	2	3	3	6	4	10
2003.....	258	1	1	-	-	-	-	4	4	6	7	7
2002.....	240	-	-	-	2	2	-	2	3	6	3	5
2001.....	238	-	2	-	1	-	-	1	1	5	4	7
2000.....	258	-	1	-	1	-	1	3	3	6	9	5
1999.....	273	-	-	1	-	1	2	-	4	6	8	10

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified diseases of skin and subcutaneous tissue (L10-L98)												
Asian or Pacific Islander 2/, both sexes												
2004.....	33	-	-	-	-	-	1	-	-	1	-	-
2003.....	27	-	-	1	1	-	-	1	1	-	1	1
2002.....	33	-	-	-	-	1	-	-	-	-	1	1
2001.....	27	-	-	-	-	-	1	-	1	-	2	-
2000.....	38	1	-	-	1	1	1	-	-	1	1	1
1999.....	27	-	-	1	-	-	2	-	-	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	18	-	-	-	-	-	1	-	-	1	-	-
2003.....	11	-	-	1	-	-	-	-	-	-	1	1
2002.....	11	-	-	-	-	-	-	-	-	-	-	-
2001.....	11	-	-	-	-	-	-	-	1	-	1	-
2000.....	12	1	-	-	-	-	-	-	-	-	1	-
1999.....	13	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	15	-	-	-	-	-	-	-	-	-	-	-
2003.....	16	-	-	-	1	-	-	1	1	-	-	-
2002.....	22	-	-	-	-	1	-	-	-	-	1	1
2001.....	16	-	-	-	-	-	1	-	-	-	1	-
2000.....	26	-	-	-	1	1	1	-	-	1	-	1
1999.....	14	-	-	1	-	-	2	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

XIII. Diseases of the musculoskeletal system and connective tissue (M00-M99)

All races, both sexes

2004.....	14,284	16	9	11	32	53	101	132	144	242	350	497
2003.....	14,489	13	10	6	25	71	106	137	187	239	376	484
2002.....	14,400	13	14	16	25	50	104	126	156	258	353	495
2001.....	14,057	9	4	11	24	68	114	123	171	296	378	443
2000.....	13,764	14	10	7	27	45	107	145	172	258	387	470
1999.....	13,347	9	8	10	28	59	89	124	167	278	358	471

All races, male

2004.....	4,239	10	6	2	12	22	32	28	41	75	102	167
2003.....	4,292	8	7	3	13	23	25	36	44	87	133	157
2002.....	4,194	11	7	5	7	13	21	37	48	81	113	174
2001.....	4,071	7	3	3	13	17	25	32	49	109	122	150
2000.....	4,035	9	6	4	8	17	36	43	52	72	136	161
1999.....	3,813	6	4	6	7	16	23	29	48	86	119	143

All races, female

2004.....	10,045	6	3	9	20	31	69	104	103	167	248	330
2003.....	10,197	5	3	3	12	48	81	101	143	152	243	327
2002.....	10,206	2	7	11	18	37	83	89	108	177	240	321
2001.....	9,986	2	1	8	11	51	89	91	122	187	256	293
2000.....	9,729	5	4	3	19	28	71	102	120	186	251	309
1999.....	9,534	3	4	4	21	43	66	95	119	192	239	328

White, both sexes

2004.....	12,014	11	7	5	20	31	44	53	64	130	210	300
2003.....	12,259	11	6	4	18	41	57	64	94	128	266	297
2002.....	12,260	8	11	13	17	27	41	60	74	133	209	325
2001.....	11,964	7	2	9	18	41	55	51	92	181	222	294
2000.....	11,679	9	5	3	14	19	48	79	100	136	222	281
1999.....	11,378	5	7	4	19	31	40	63	100	140	200	300

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
XIII. Diseases of the musculoskeletal system and connective tissue (M00-M99)												
White, male												
2004.....	3,532	8	6	2	7	17	14	10	17	46	69	102
2003.....	3,576	6	5	3	9	17	20	23	24	48	102	108
2002.....	3,529	6	6	4	4	6	11	22	23	52	74	122
2001.....	3,435	6	1	3	8	14	16	17	29	78	73	100
2000.....	3,359	4	4	3	6	4	17	26	32	37	87	109
1999.....	3,229	5	3	3	4	12	10	18	36	47	73	97
White, female												
2004.....	8,482	3	1	3	13	14	30	43	47	84	141	198
2003.....	8,683	5	1	1	9	24	37	41	70	80	164	189
2002.....	8,731	2	5	9	13	21	30	38	51	81	135	203
2001.....	8,529	1	1	6	10	27	39	34	63	103	149	194
2000.....	8,320	5	1	-	8	15	31	53	68	99	135	172
1999.....	8,149	-	4	1	15	19	30	45	64	93	127	203
Black, both sexes												
2004.....	1,915	5	1	6	9	18	52	71	69	98	119	177
2003.....	1,869	2	3	2	5	28	46	61	76	97	100	175
2002.....	1,811	4	2	3	6	17	56	60	74	113	131	152
2001.....	1,779	2	1	2	4	21	48	62	70	104	142	135
2000.....	1,787	5	4	2	13	23	52	62	69	110	153	170
1999.....	1,689	4	1	4	8	23	46	55	60	121	150	156
Black, male												
2004.....	604	2	-	-	3	5	16	18	22	23	29	57
2003.....	601	2	1	-	3	6	5	12	15	35	27	43
2002.....	563	4	1	1	3	5	10	13	24	27	34	46
2001.....	554	1	1	-	4	3	8	12	18	29	46	48
2000.....	588	5	1	1	2	12	16	16	19	33	47	46
1999.....	506	1	1	3	3	2	13	11	11	33	44	42

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

XIII. Diseases of the musculoskeletal system and connective tissue (M00-M99)

Black, female

2004.....	1,311	3	1	6	6	13	36	53	47	75	90	120
2003.....	1,268	-	2	2	2	22	41	49	61	62	73	132
2002.....	1,248	-	1	2	3	12	46	47	50	86	97	106
2001.....	1,225	1	-	2	-	18	40	50	52	75	96	87
2000.....	1,199	-	3	1	11	11	36	46	50	77	106	124
1999.....	1,183	3	-	1	5	21	33	44	49	88	106	114

American Indian 1/, both sexes

2004.....	101	-	-	-	-	-	-	3	5	4	4	6
2003.....	106	-	-	-	2	-	-	2	4	2	3	3
2002.....	92	-	-	-	1	-	2	2	3	1	3	9
2001.....	93	-	-	-	-	1	1	1	2	1	5	6
2000.....	81	-	-	-	-	-	2	2	1	2	2	3
1999.....	90	-	-	-	-	1	2	1	2	6	4	5

American Indian 1/, male

2004.....	34	-	-	-	-	-	-	-	-	2	1	2
2003.....	34	-	-	-	1	-	-	-	1	2	-	3
2002.....	22	-	-	-	-	-	-	1	-	-	1	2
2001.....	24	-	-	-	-	-	-	1	-	1	-	-
2000.....	24	-	-	-	-	-	2	1	-	1	-	-
1999.....	23	-	-	-	-	1	-	-	-	1	1	1

American Indian 1/, female

2004.....	67	-	-	-	-	-	-	3	5	2	3	4
2003.....	72	-	-	-	1	-	-	2	3	-	3	-
2002.....	70	-	-	-	1	-	2	1	3	1	2	7
2001.....	69	-	-	-	-	1	1	-	2	-	5	6
2000.....	57	-	-	-	-	-	-	1	1	1	2	3
1999.....	67	-	-	-	-	-	2	1	2	5	3	4

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
XIII. Diseases of the musculoskeletal system and connective tissue (M00-M99)												
Asian or Pacific Islander 2/, both sexes												
2004.....	254	-	1	-	3	4	5	5	6	10	17	14
2003.....	255	-	1	-	-	2	3	10	13	12	7	9
2002.....	237	1	1	-	1	6	5	4	5	11	10	9
2001.....	221	-	1	-	2	5	10	9	7	10	9	8
2000.....	217	-	1	2	-	3	5	2	2	10	10	16
1999.....	190	-	-	2	1	4	1	5	5	11	4	10
Asian or Pacific Islander 2/, male												
2004.....	69	-	-	-	2	-	2	-	2	4	3	6
2003.....	81	-	1	-	-	-	-	1	4	2	4	3
2002.....	80	1	-	-	-	2	-	1	1	2	4	4
2001.....	58	-	1	-	1	-	1	2	2	1	3	2
2000.....	64	-	1	-	-	1	1	-	1	1	2	6
1999.....	55	-	-	-	-	1	-	-	1	5	1	3
Asian or Pacific Islander 2/, female												
2004.....	185	-	1	-	1	4	3	5	4	6	14	8
2003.....	174	-	-	-	-	2	3	9	9	10	3	6
2002.....	157	-	1	-	1	4	5	3	4	9	6	5
2001.....	163	-	-	-	1	5	9	7	5	9	6	6
2000.....	153	-	-	2	-	2	4	2	1	9	8	10
1999.....	135	-	-	2	1	3	1	5	4	6	3	7

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Rheumatoid arthritis and related inflammatory polyarthropathies (M05-M08)

All races, both sexes

2004.....	2,571	1	-	1	1	1	2	3	3	11	15	38
2003.....	2,677	-	1	1	1	2	1	6	4	7	22	33
2002.....	2,809	-	-	4	2	2	-	2	2	5	17	43
2001.....	2,657	-	-	-	-	-	2	1	2	16	17	29
2000.....	2,689	-	1	-	1	2	1	3	2	7	16	35
1999.....	2,763	-	-	1	2	-	-	2	9	7	13	37

All races, male

2004.....	600	1	-	-	-	-	1	-	-	3	4	10
2003.....	640	-	1	1	1	1	-	-	-	4	8	10
2002.....	685	-	-	3	-	-	-	1	-	3	2	16
2001.....	616	-	-	-	-	-	-	1	1	4	3	9
2000.....	635	-	-	-	-	-	-	-	-	1	6	8
1999.....	657	-	-	-	1	-	-	-	3	1	6	10

All races, female

2004.....	1,971	-	-	1	1	1	1	3	3	8	11	28
2003.....	2,037	-	-	-	-	1	1	6	4	3	14	23
2002.....	2,124	-	-	1	2	2	-	1	2	2	15	27
2001.....	2,041	-	-	-	-	-	2	-	1	12	14	20
2000.....	2,054	-	1	-	1	2	1	3	2	6	10	27
1999.....	2,106	-	-	1	1	-	-	2	6	6	7	27

White, both sexes

2004.....	2,318	1	-	-	-	1	1	3	2	5	12	28
2003.....	2,449	-	1	1	1	-	1	4	4	6	19	27
2002.....	2,566	-	-	2	2	1	-	-	1	4	13	38
2001.....	2,439	-	-	-	-	-	1	1	2	11	14	24
2000.....	2,489	-	-	-	1	2	-	2	-	7	14	27
1999.....	2,585	-	-	-	1	-	-	1	8	5	9	31

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Rheumatoid arthritis and related inflammatory polyarthropathies (M05-M08)

White, male

2004.....	551	1	-	-	-	-	1	-	-	3	4	7
2003.....	582	-	1	1	1	-	-	-	-	3	6	8
2002.....	626	-	-	2	-	-	-	-	-	3	1	15
2001.....	564	-	-	-	-	-	-	1	1	3	2	7
2000.....	592	-	-	-	-	-	-	-	-	1	5	7
1999.....	613	-	-	-	-	-	-	-	3	1	5	9

White, female

2004.....	1,767	-	-	-	-	1	-	3	2	2	8	21
2003.....	1,867	-	-	-	-	-	1	4	4	3	13	19
2002.....	1,940	-	-	-	2	1	-	-	1	1	12	23
2001.....	1,875	-	-	-	-	-	1	-	1	8	12	17
2000.....	1,897	-	-	-	1	2	-	2	-	6	9	20
1999.....	1,972	-	-	-	1	-	-	1	5	4	4	22

Black, both sexes

2004.....	184	-	-	1	1	-	1	-	1	4	2	5
2003.....	169	-	-	-	-	2	-	2	-	1	3	4
2002.....	174	-	-	2	-	1	-	1	1	1	4	5
2001.....	164	-	-	-	-	-	1	-	-	3	2	3
2000.....	148	-	1	-	-	-	1	1	2	-	2	7
1999.....	141	-	-	-	1	-	-	1	1	2	4	5

Black, male

2004.....	38	-	-	-	-	-	-	-	-	-	-	1
2003.....	44	-	-	-	-	1	-	-	-	1	2	1
2002.....	43	-	-	1	-	-	-	1	-	-	1	1
2001.....	40	-	-	-	-	-	-	-	-	1	1	1
2000.....	34	-	-	-	-	-	-	-	-	-	1	1
1999.....	38	-	-	-	1	-	-	-	-	-	1	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Systemic lupus erythematosus (M32)

All races, both sexes

2004.....	1,303	-	-	4	3	19	59	73	73	96	112	105
2003.....	1,324	-	-	1	5	31	58	64	80	91	115	120
2002.....	1,292	-	-	1	8	23	60	56	65	105	111	138
2001.....	1,359	-	-	2	6	35	72	73	75	105	111	126
2000.....	1,339	1	-	1	9	22	59	69	61	116	138	129
1999.....	1,322	-	-	-	6	25	57	62	74	115	126	133

All races, male

2004.....	210	-	-	-	1	2	11	8	9	23	18	10
2003.....	193	-	-	-	1	5	4	10	8	12	23	18
2002.....	168	-	-	-	1	3	7	7	10	7	13	16
2001.....	196	-	-	-	3	2	10	11	9	15	14	13
2000.....	201	1	-	-	-	7	10	7	4	18	18	15
1999.....	188	-	-	-	1	4	10	4	12	14	15	16

All races, female

2004.....	1,093	-	-	4	2	17	48	65	64	73	94	95
2003.....	1,131	-	-	1	4	26	54	54	72	79	92	102
2002.....	1,124	-	-	1	7	20	53	49	55	98	98	122
2001.....	1,163	-	-	2	3	33	62	62	66	90	97	113
2000.....	1,138	-	-	1	9	15	49	62	57	98	120	114
1999.....	1,134	-	-	-	5	21	47	58	62	101	111	117

White, both sexes

2004.....	764	-	-	1	2	7	22	25	27	43	53	48
2003.....	849	-	-	-	3	11	26	27	37	43	70	57
2002.....	781	-	-	1	3	8	18	17	25	36	54	76
2001.....	851	-	-	2	4	17	30	22	31	45	46	73
2000.....	838	-	-	-	2	5	24	29	27	53	69	58
1999.....	833	-	-	-	2	8	22	29	32	47	58	77

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Systemic lupus erythematosus (M32)

White, male

2004.....	125	-	-	-	1	2	3	1	3	10	7	3
2003.....	140	-	-	-	1	2	3	6	3	7	16	9
2002.....	121	-	-	-	-	-	2	3	3	4	7	13
2001.....	136	-	-	-	2	1	7	3	3	8	7	8
2000.....	134	-	-	-	-	1	4	4	1	6	10	7
1999.....	130	-	-	-	1	2	3	2	6	6	6	10

White, female

2004.....	639	-	-	1	1	5	19	24	24	33	46	45
2003.....	709	-	-	-	2	9	23	21	34	36	54	48
2002.....	660	-	-	1	3	8	16	14	22	32	47	63
2001.....	715	-	-	2	2	16	23	19	28	37	39	65
2000.....	704	-	-	-	2	4	20	25	26	47	59	51
1999.....	703	-	-	-	1	6	19	27	26	41	52	67

Black, both sexes

2004.....	476	-	-	3	1	9	34	41	40	44	51	54
2003.....	421	-	-	1	2	20	30	29	36	40	41	63
2002.....	457	-	-	-	4	11	37	35	37	61	52	56
2001.....	435	-	-	-	-	14	32	42	38	55	59	49
2000.....	447	1	-	1	7	14	32	38	32	56	62	62
1999.....	432	-	-	-	3	14	33	29	36	59	65	54

Black, male

2004.....	75	-	-	-	-	-	7	7	5	9	10	7
2003.....	42	-	-	-	-	3	1	3	2	4	5	9
2002.....	40	-	-	-	1	2	5	3	7	2	4	2
2001.....	53	-	-	-	-	1	2	5	6	6	7	5
2000.....	62	1	-	-	-	5	6	3	3	12	8	6
1999.....	48	-	-	-	-	1	7	2	5	6	8	6

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Systemic lupus erythematosus (M32)												
Black, female												
2004.....	401	-	-	3	1	9	27	34	35	35	41	47
2003.....	379	-	-	1	2	17	29	26	34	36	36	54
2002.....	417	-	-	-	3	9	32	32	30	59	48	54
2001.....	382	-	-	-	-	13	30	37	32	49	52	44
2000.....	385	-	-	1	7	9	26	35	29	44	54	56
1999.....	384	-	-	-	3	13	26	27	31	53	57	48
American Indian 1/, both sexes												
2004.....	15	-	-	-	-	-	-	3	1	1	-	-
2003.....	13	-	-	-	-	-	-	2	1	-	1	-
2002.....	13	-	-	-	-	-	-	1	-	-	2	3
2001.....	14	-	-	-	-	-	1	1	1	-	1	1
2000.....	8	-	-	-	-	-	-	-	1	1	1	-
1999.....	12	-	-	-	-	-	2	-	1	2	-	-
American Indian 1/, male												
2004.....	2	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	1	-	-	-	-	-	-	1	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, female												
2004.....	13	-	-	-	-	-	-	3	1	1	-	-
2003.....	13	-	-	-	-	-	-	2	1	-	1	-
2002.....	13	-	-	-	-	-	-	1	-	-	2	3
2001.....	13	-	-	-	-	-	1	-	1	-	1	1
2000.....	8	-	-	-	-	-	-	-	1	1	1	-
1999.....	12	-	-	-	-	-	2	-	1	2	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Systemic lupus erythematosus (M32)												
Asian or Pacific Islander 2/, both sexes												
2004.....	48	-	-	-	-	3	3	4	5	8	8	3
2003.....	41	-	-	-	-	-	2	6	6	8	3	-
2002.....	41	-	-	-	1	4	5	3	3	8	3	3
2001.....	59	-	-	-	2	4	9	8	5	5	5	3
2000.....	46	-	-	-	-	3	3	2	1	6	6	9
1999.....	45	-	-	-	1	3	-	4	5	7	3	2
Asian or Pacific Islander 2/, male												
2004.....	8	-	-	-	-	-	1	-	1	4	1	-
2003.....	11	-	-	-	-	-	-	1	3	1	2	-
2002.....	7	-	-	-	-	1	-	1	-	1	2	1
2001.....	6	-	-	-	1	-	1	2	-	1	-	-
2000.....	5	-	-	-	-	1	-	-	-	-	-	2
1999.....	10	-	-	-	-	1	-	-	1	2	1	-
Asian or Pacific Islander 2/, female												
2004.....	40	-	-	-	-	3	2	4	4	4	7	3
2003.....	30	-	-	-	-	-	2	5	3	7	1	-
2002.....	34	-	-	-	1	3	5	2	3	7	1	2
2001.....	53	-	-	-	1	4	8	6	5	4	5	3
2000.....	41	-	-	-	-	2	3	2	1	6	6	7
1999.....	35	-	-	-	1	2	-	4	4	5	2	2

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other arthropathies and related disorders (M00,M02,M10-M31,M33-M35)

All races, both sexes

2004.....	4,713	6	5	3	8	18	25	42	40	94	147	195
2003.....	4,770	2	4	2	8	20	25	46	61	92	164	198
2002.....	4,815	3	5	3	3	12	24	44	60	99	157	201
2001.....	4,805	-	2	4	5	18	28	40	56	114	170	194
2000.....	4,790	2	3	2	6	12	29	47	78	90	161	193
1999.....	4,652	5	2	4	5	21	19	38	57	111	152	207

All races, male

2004.....	1,380	3	3	-	4	8	9	14	18	26	41	61
2003.....	1,465	2	4	1	4	7	7	13	18	40	59	63
2002.....	1,421	2	4	1	-	3	4	15	20	46	51	70
2001.....	1,448	-	2	2	2	5	9	14	18	45	55	77
2000.....	1,458	2	1	2	2	4	13	20	32	23	62	69
1999.....	1,420	3	2	2	2	6	5	14	17	46	58	63

All races, female

2004.....	3,333	3	2	3	4	10	16	28	22	68	106	134
2003.....	3,305	-	-	1	4	13	18	33	43	52	105	135
2002.....	3,394	1	1	2	3	9	20	29	40	53	106	131
2001.....	3,357	-	-	2	3	13	19	26	38	69	115	117
2000.....	3,332	-	2	-	4	8	16	27	46	67	99	124
1999.....	3,232	2	-	2	3	15	14	24	40	65	94	144

White, both sexes

2004.....	3,937	5	4	1	4	11	15	17	22	52	86	105
2003.....	3,939	2	3	1	6	12	15	19	22	46	114	115
2002.....	4,069	2	4	3	3	8	11	24	30	56	91	127
2001.....	4,073	-	1	3	4	13	16	21	32	71	101	122
2000.....	4,020	1	1	2	4	9	15	30	47	41	91	115
1999.....	3,877	2	2	2	2	13	11	18	33	61	83	121

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other arthropathies and related disorders (M00,M02,M10-M31,M33-M35)

White, male

2004.....	1,144	3	3	-	2	5	5	6	7	17	28	29
2003.....	1,200	2	3	1	3	5	7	6	9	19	44	44
2002.....	1,174	1	3	1	-	2	2	8	10	29	32	43
2001.....	1,207	-	1	2	1	4	5	9	11	28	30	47
2000.....	1,188	1	1	2	2	2	9	13	19	7	37	49
1999.....	1,178	2	2	1	-	4	3	8	11	25	32	37

White, female

2004.....	2,793	2	1	1	2	6	10	11	15	35	58	76
2003.....	2,739	-	-	-	3	7	8	13	13	27	70	71
2002.....	2,895	1	1	2	3	6	9	16	20	27	59	84
2001.....	2,866	-	-	1	3	9	11	12	21	43	71	75
2000.....	2,832	-	-	-	2	7	6	17	28	34	54	66
1999.....	2,699	-	-	1	2	9	8	10	22	36	51	84

Black, both sexes

2004.....	652	1	-	2	3	6	8	24	18	40	51	83
2003.....	689	-	-	1	1	6	9	23	31	42	45	75
2002.....	641	-	1	-	-	2	11	19	26	41	61	65
2001.....	636	-	1	1	1	3	11	18	24	40	63	67
2000.....	656	1	2	-	2	3	12	15	31	46	66	71
1999.....	670	3	-	2	3	6	7	18	23	47	65	76

Black, male

2004.....	200	-	-	-	1	3	3	8	11	8	12	29
2003.....	225	-	-	-	1	2	-	7	9	20	14	16
2002.....	214	-	1	-	-	-	2	6	10	17	18	24
2001.....	210	-	1	-	1	1	4	5	7	16	22	30
2000.....	227	1	-	-	-	2	3	6	13	15	24	18
1999.....	217	1	-	1	2	1	2	6	6	21	26	24

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other arthropathies and related disorders (M00,M02,M10-M31,M33-M35)												
Black, female												
2004.....	452	1	-	2	2	3	5	16	7	32	39	54
2003.....	464	-	-	1	-	4	9	16	22	22	31	59
2002.....	427	-	-	-	-	2	9	13	16	24	43	41
2001.....	426	-	-	1	-	2	7	13	17	24	41	37
2000.....	429	-	2	-	2	1	9	9	18	31	42	53
1999.....	453	2	-	1	1	5	5	12	17	26	39	52
American Indian 1/, both sexes												
2004.....	30	-	-	-	-	-	-	-	-	1	4	1
2003.....	30	-	-	-	1	-	-	-	2	-	2	1
2002.....	27	-	-	-	-	-	2	1	3	-	-	4
2001.....	30	-	-	-	-	1	-	-	-	1	3	2
2000.....	33	-	-	-	-	-	-	2	-	-	1	2
1999.....	34	-	-	-	-	1	-	1	1	2	3	3
American Indian 1/, male												
2004.....	11	-	-	-	-	-	-	-	-	1	1	-
2003.....	5	-	-	-	-	-	-	-	-	-	-	1
2002.....	6	-	-	-	-	-	-	1	-	-	-	1
2001.....	8	-	-	-	-	-	-	-	-	1	-	-
2000.....	10	-	-	-	-	-	-	1	-	-	-	-
1999.....	6	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, female												
2004.....	19	-	-	-	-	-	-	-	-	-	3	1
2003.....	25	-	-	-	1	-	-	-	2	-	2	-
2002.....	21	-	-	-	-	-	2	-	3	-	-	3
2001.....	22	-	-	-	-	1	-	-	-	-	3	2
2000.....	23	-	-	-	-	-	-	1	-	-	1	2
1999.....	28	-	-	-	-	-	-	1	1	2	3	3

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other arthropathies and related disorders (M00,M02,M10-M31,M33-M35)												
Asian or Pacific Islander 2/, both sexes												
2004.....	94	-	1	-	1	1	2	1	-	1	6	6
2003.....	112	-	1	-	-	2	1	4	6	4	3	7
2002.....	78	1	-	-	-	2	-	-	1	2	5	5
2001.....	66	-	-	-	-	1	1	1	-	2	3	3
2000.....	81	-	-	-	-	-	2	-	-	3	3	5
1999.....	71	-	-	-	-	1	1	1	-	1	1	7
Asian or Pacific Islander 2/, male												
2004.....	25	-	-	-	1	-	1	-	-	-	-	3
2003.....	35	-	1	-	-	-	-	-	-	1	1	2
2002.....	27	1	-	-	-	1	-	-	-	-	1	2
2001.....	23	-	-	-	-	-	-	-	-	-	3	-
2000.....	33	-	-	-	-	-	1	-	-	1	1	2
1999.....	19	-	-	-	-	-	-	-	-	-	-	2
Asian or Pacific Islander 2/, female												
2004.....	69	-	1	-	-	1	1	1	-	1	6	3
2003.....	77	-	-	-	-	2	1	4	6	3	2	5
2002.....	51	-	-	-	-	1	-	-	1	2	4	3
2001.....	43	-	-	-	-	1	1	1	-	2	-	3
2000.....	48	-	-	-	-	-	1	-	-	2	2	3
1999.....	52	-	-	-	-	1	1	1	-	1	1	5

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Dorsopathies (M40-M54)												
All races, both sexes												
2004.....	1,075	2	3	1	15	9	5	6	5	10	13	25
2003.....	1,015	2	1	2	5	13	6	3	16	11	12	24
2002.....	947	3	-	4	10	10	7	9	7	12	13	25
2001.....	959	3	-	3	10	10	6	2	9	16	17	14
2000.....	869	4	-	2	8	6	5	7	8	8	13	23
1999.....	787	-	3	2	11	8	7	6	13	8	13	21
All races, male												
2004.....	443	1	2	-	6	6	3	2	3	4	6	14
2003.....	426	2	-	1	2	10	2	2	3	6	7	11
2002.....	395	3	-	1	5	5	3	6	3	5	6	16
2001.....	379	3	-	-	7	7	3	-	5	10	10	6
2000.....	375	3	-	1	3	4	3	3	3	6	9	12
1999.....	339	-	1	2	3	2	4	3	8	5	7	12
All races, female												
2004.....	632	1	1	1	9	3	2	4	2	6	7	11
2003.....	589	-	1	1	3	3	4	1	13	5	5	13
2002.....	552	-	-	3	5	5	4	3	4	7	7	9
2001.....	580	-	-	3	3	3	3	2	4	6	7	8
2000.....	494	1	-	1	5	2	2	4	5	2	4	11
1999.....	448	-	2	-	8	6	3	3	5	3	6	9
White, both sexes												
2004.....	956	2	2	1	11	6	3	4	3	8	10	18
2003.....	887	2	1	2	4	13	4	3	13	6	11	23
2002.....	841	-	-	4	8	8	5	7	7	8	9	22
2001.....	852	2	-	2	7	8	3	2	9	14	15	13
2000.....	775	2	-	-	6	2	3	5	7	8	11	20
1999.....	702	-	3	1	10	5	5	4	13	6	12	18

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Dorsopathies (M40-M54)												
White, male												
2004.....	376	1	2	-	3	4	3	1	2	3	5	9
2003.....	359	2	-	1	1	10	1	2	2	4	6	11
2002.....	333	-	-	1	4	3	2	4	3	2	4	15
2001.....	331	2	-	-	4	7	2	-	5	8	8	6
2000.....	324	1	-	-	3	1	1	1	2	6	8	9
1999.....	298	-	1	1	3	2	3	2	8	3	6	11
White, female												
2004.....	580	1	-	1	8	2	-	3	1	5	5	9
2003.....	528	-	1	1	3	3	3	1	11	2	5	12
2002.....	508	-	-	3	4	5	3	3	4	6	5	7
2001.....	521	-	-	2	3	1	1	2	4	6	7	7
2000.....	451	1	-	-	3	1	2	4	5	2	3	11
1999.....	404	-	2	-	7	3	2	2	5	3	6	7
Black, both sexes												
2004.....	97	-	1	-	3	3	2	2	2	2	3	6
2003.....	106	-	-	-	-	-	2	-	3	5	1	1
2002.....	85	3	-	-	1	2	2	2	-	3	4	3
2001.....	91	1	-	1	3	2	3	-	-	2	2	-
2000.....	82	2	-	1	2	4	2	2	-	-	2	3
1999.....	70	-	-	1	1	3	2	2	-	-	1	2
Black, male												
2004.....	54	-	-	-	2	2	-	1	1	1	1	4
2003.....	57	-	-	-	-	-	1	-	1	2	1	-
2002.....	51	3	-	-	1	2	1	2	-	3	2	1
2001.....	41	1	-	-	3	-	1	-	-	2	2	-
2000.....	46	2	-	1	-	3	2	2	-	-	1	3
1999.....	32	-	-	1	-	-	1	1	-	-	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Soft tissue disorders (M60-M79)

All races, both sexes

2004.....	1,278	2	-	1	5	4	9	6	21	20	41	85
2003.....	1,286	7	3	-	4	2	13	12	21	31	41	63
2002.....	1,161	7	8	1	-	3	11	10	18	25	44	60
2001.....	1,068	4	2	2	2	4	5	5	18	32	47	47
2000.....	1,015	3	6	1	1	1	9	13	13	23	44	71
1999.....	956	3	2	1	4	4	5	13	11	31	43	47

All races, male

2004.....	662	1	-	1	1	4	7	3	10	13	23	44
2003.....	637	4	2	-	3	-	11	8	14	18	22	31
2002.....	610	6	3	-	-	2	6	6	13	15	32	41
2001.....	578	2	1	1	-	2	2	4	11	25	30	28
2000.....	539	1	5	1	1	1	7	9	9	13	27	44
1999.....	517	2	-	1	-	3	3	7	6	17	28	26

All races, female

2004.....	616	1	-	-	4	-	2	3	11	7	18	41
2003.....	649	3	1	-	1	2	2	4	7	13	19	32
2002.....	551	1	5	1	-	1	5	4	5	10	12	19
2001.....	490	2	1	1	2	2	3	1	7	7	17	19
2000.....	476	2	1	-	-	-	2	4	4	10	17	27
1999.....	439	1	2	-	4	1	2	6	5	14	15	21

White, both sexes

2004.....	1,020	1	-	1	3	4	3	3	9	14	29	67
2003.....	1,057	5	1	-	3	2	8	9	14	25	32	43
2002.....	942	6	7	1	-	2	6	9	8	18	32	40
2001.....	851	3	1	2	2	2	4	5	10	27	34	38
2000.....	785	2	4	1	-	-	2	9	11	16	25	49
1999.....	757	2	2	-	4	4	2	9	11	18	33	30

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Soft tissue disorders (M60-M79)

White, male

2004.....	528	1	-	1	1	4	2	2	5	9	17	35
2003.....	527	2	1	-	2	-	8	8	9	13	18	20
2002.....	492	5	3	-	-	1	4	6	6	10	22	25
2001.....	465	2	-	1	-	1	1	4	6	21	19	21
2000.....	424	-	3	1	-	-	-	6	7	9	16	29
1999.....	411	2	-	-	-	3	1	5	6	10	22	17

White, female

2004.....	492	-	-	-	2	-	1	1	4	5	12	32
2003.....	530	3	-	-	1	2	-	1	5	12	14	23
2002.....	450	1	4	1	-	1	2	3	2	8	10	15
2001.....	386	1	1	1	2	1	3	1	4	6	15	17
2000.....	361	2	1	-	-	-	2	3	4	7	9	20
1999.....	346	-	2	-	4	1	1	4	5	8	11	13

Black, both sexes

2004.....	222	1	-	-	1	-	6	3	7	5	10	16
2003.....	200	2	2	-	1	-	5	3	5	5	8	19
2002.....	190	1	1	-	-	1	5	1	9	6	9	20
2001.....	189	1	-	-	-	2	1	-	5	4	12	8
2000.....	205	1	1	-	1	1	5	4	2	5	18	20
1999.....	177	1	-	1	-	-	3	4	-	11	10	16

Black, male

2004.....	114	-	-	-	-	-	5	1	4	3	4	8
2003.....	93	2	1	-	1	-	3	-	3	4	3	10
2002.....	102	1	-	-	-	1	2	-	6	4	8	16
2001.....	101	-	-	-	-	1	1	-	3	4	11	6
2000.....	102	1	1	-	1	1	5	3	2	3	10	13
1999.....	94	-	-	1	-	-	2	2	-	6	6	8

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Soft tissue disorders (M60-M79)												
Black, female												
2004.....	108	1	-	-	1	-	1	2	3	2	6	8
2003.....	107	-	1	-	-	-	2	3	2	1	5	9
2002.....	88	-	1	-	-	-	3	1	3	2	1	4
2001.....	88	1	-	-	-	1	-	-	2	-	1	2
2000.....	103	-	-	-	-	-	-	1	-	2	8	7
1999.....	83	1	-	-	-	-	1	2	-	5	4	8
American Indian 1/, both sexes												
2004.....	14	-	-	-	-	-	-	-	4	1	-	-
2003.....	12	-	-	-	-	-	-	-	1	1	-	1
2002.....	9	-	-	-	-	-	-	-	-	-	1	-
2001.....	10	-	-	-	-	-	-	-	1	-	-	-
2000.....	10	-	-	-	-	-	2	-	-	1	-	-
1999.....	7	-	-	-	-	-	-	-	-	1	-	1
American Indian 1/, male												
2004.....	6	-	-	-	-	-	-	-	-	1	-	-
2003.....	9	-	-	-	-	-	-	-	1	1	-	1
2002.....	2	-	-	-	-	-	-	-	-	-	1	-
2001.....	2	-	-	-	-	-	-	-	-	-	-	-
2000.....	3	-	-	-	-	-	2	-	-	1	-	-
1999.....	2	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, female												
2004.....	8	-	-	-	-	-	-	-	4	-	-	-
2003.....	3	-	-	-	-	-	-	-	-	-	-	-
2002.....	7	-	-	-	-	-	-	-	-	-	-	-
2001.....	8	-	-	-	-	-	-	-	1	-	-	-
2000.....	7	-	-	-	-	-	-	-	-	-	-	-
1999.....	5	-	-	-	-	-	-	-	-	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Osteopathies, chondropathies and other disorders of musculoskeletal system and connective tissue (M80-M99)

All races, both sexes

2004.....	3,344	5	1	1	-	2	1	2	2	11	22	49
2003.....	3,417	2	1	-	2	3	3	6	5	7	22	46
2002.....	3,376	-	1	3	2	-	2	5	4	12	11	28
2001.....	3,209	2	-	-	1	1	1	2	11	13	16	33
2000.....	3,062	4	-	1	2	2	4	6	10	14	15	19
1999.....	2,867	1	1	2	-	1	1	3	3	6	11	26

All races, male

2004.....	944	4	1	1	-	2	1	1	1	6	10	28
2003.....	931	-	-	-	2	-	1	3	1	7	14	24
2002.....	915	-	-	-	1	-	1	2	2	5	9	15
2001.....	854	2	-	-	1	1	1	2	5	10	10	17
2000.....	827	2	-	-	2	1	3	4	4	11	14	13
1999.....	692	1	1	1	-	1	1	1	2	3	5	16

All races, female

2004.....	2,400	1	-	-	-	-	-	1	1	5	12	21
2003.....	2,486	2	1	-	-	3	2	3	4	-	8	22
2002.....	2,461	-	1	3	1	-	1	3	2	7	2	13
2001.....	2,355	-	-	-	-	-	-	-	6	3	6	16
2000.....	2,235	2	-	1	-	1	1	2	6	3	1	6
1999.....	2,175	-	-	1	-	-	-	2	1	3	6	10

White, both sexes

2004.....	3,019	2	1	1	-	2	-	1	1	8	20	34
2003.....	3,078	2	-	-	1	3	3	2	4	2	20	32
2002.....	3,061	-	-	2	1	-	1	3	3	11	10	22
2001.....	2,898	2	-	-	1	1	1	-	8	13	12	24
2000.....	2,772	4	-	-	1	1	4	4	8	11	12	12
1999.....	2,624	1	-	1	-	1	-	2	3	3	5	23

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Osteopathies, chondropathies and other disorders of musculoskeletal system and connective tissue (M80-M99)

White, male

2004.....	808	2	1	1	-	2	-	-	-	4	8	19
2003.....	768	-	-	-	1	-	1	1	1	2	12	16
2002.....	783	-	-	-	-	-	1	1	1	4	8	11
2001.....	732	2	-	-	1	1	1	-	3	10	7	11
2000.....	697	2	-	-	1	-	3	2	3	8	11	8
1999.....	599	1	-	1	-	1	-	1	2	2	2	13

White, female

2004.....	2,211	-	-	-	-	-	-	1	1	4	12	15
2003.....	2,310	2	-	-	-	3	2	1	3	-	8	16
2002.....	2,278	-	-	2	1	-	-	2	2	7	2	11
2001.....	2,166	-	-	-	-	-	-	-	5	3	5	13
2000.....	2,075	2	-	-	-	1	1	2	5	3	1	4
1999.....	2,025	-	-	-	-	-	-	1	1	1	3	10

Black, both sexes

2004.....	284	3	-	-	-	-	1	1	1	3	2	13
2003.....	284	-	1	-	1	-	-	4	1	4	2	13
2002.....	264	-	-	1	1	-	1	2	1	1	1	3
2001.....	264	-	-	-	-	-	-	2	3	-	4	8
2000.....	249	-	-	-	1	1	-	2	2	3	3	7
1999.....	199	-	1	-	-	-	1	1	-	2	5	3

Black, male

2004.....	123	2	-	-	-	-	1	1	1	2	2	8
2003.....	140	-	-	-	1	-	-	2	-	4	2	7
2002.....	113	-	-	-	1	-	-	1	1	1	1	2
2001.....	109	-	-	-	-	-	-	2	2	-	3	6
2000.....	117	-	-	-	1	1	-	2	1	3	3	5
1999.....	77	-	1	-	-	-	1	-	-	-	2	3

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Osteoporosis (M80-M81)												
All races, both sexes												
2004.....	1,697	1	-	-	-	2	-	-	-	-	2	4
2003.....	1,784	1	-	-	-	-	-	1	1	-	1	3
2002.....	1,862	-	-	1	-	-	-	-	2	1	-	1
2001.....	1,684	1	-	-	1	-	-	-	-	-	2	-
2000.....	1,716	2	-	-	-	-	-	-	1	1	1	1
1999.....	1,642	1	-	1	-	-	-	-	1	-	1	2
All races, male												
2004.....	203	1	-	-	-	2	-	-	-	-	1	3
2003.....	190	-	-	-	-	-	-	-	-	-	1	1
2002.....	213	-	-	-	-	-	-	-	1	-	-	-
2001.....	196	1	-	-	1	-	-	-	-	-	2	-
2000.....	201	2	-	-	-	-	-	-	-	1	-	-
1999.....	160	1	-	1	-	-	-	-	-	-	-	-
All races, female												
2004.....	1,494	-	-	-	-	-	-	-	-	-	1	1
2003.....	1,594	1	-	-	-	-	-	1	1	-	-	2
2002.....	1,649	-	-	1	-	-	-	-	1	1	-	1
2001.....	1,488	-	-	-	-	-	-	-	-	-	-	-
2000.....	1,515	-	-	-	-	-	-	-	1	-	1	1
1999.....	1,482	-	-	-	-	-	-	-	1	-	1	2
White, both sexes												
2004.....	1,649	1	-	-	-	2	-	-	-	-	2	3
2003.....	1,735	1	-	-	-	-	-	1	1	-	1	1
2002.....	1,806	-	-	1	-	-	-	-	1	1	-	1
2001.....	1,634	1	-	-	1	-	-	-	-	-	2	-
2000.....	1,669	2	-	-	-	-	-	-	1	1	1	1
1999.....	1,601	1	-	1	-	-	-	-	1	-	1	2

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other osteopathies, chondropathies and disorders of musculoskeletal system and connective tissue (M83-M99)

All races, both sexes

2004.....	1,647	4	1	1	-	-	1	2	2	11	20	45
2003.....	1,633	1	1	-	2	3	3	5	4	7	21	43
2002.....	1,514	-	1	2	2	-	2	5	2	11	11	27
2001.....	1,525	1	-	-	-	1	1	2	11	13	14	33
2000.....	1,346	2	-	1	2	2	4	6	9	13	14	18
1999.....	1,225	-	1	1	-	1	1	3	2	6	10	24

All races, male

2004.....	741	3	1	1	-	-	1	1	1	6	9	25
2003.....	741	-	-	-	2	-	1	3	1	7	13	23
2002.....	702	-	-	-	1	-	1	2	1	5	9	15
2001.....	658	1	-	-	-	1	1	2	5	10	8	17
2000.....	626	-	-	-	2	1	3	4	4	10	14	13
1999.....	532	-	1	-	-	1	1	1	2	3	5	16

All races, female

2004.....	906	1	-	-	-	-	-	1	1	5	11	20
2003.....	892	1	1	-	-	3	2	2	3	-	8	20
2002.....	812	-	1	2	1	-	1	3	1	6	2	12
2001.....	867	-	-	-	-	-	-	-	6	3	6	16
2000.....	720	2	-	1	-	1	1	2	5	3	-	5
1999.....	693	-	-	1	-	-	-	2	-	3	5	8

White, both sexes

2004.....	1,370	1	1	1	-	-	-	1	1	8	18	31
2003.....	1,343	1	-	-	1	3	3	1	3	2	19	31
2002.....	1,255	-	-	1	1	-	1	3	2	10	10	21
2001.....	1,264	1	-	-	-	1	1	-	8	13	10	24
2000.....	1,103	2	-	-	1	1	4	4	7	10	11	11
1999.....	1,023	-	-	-	-	1	-	2	2	3	4	21

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other osteopathies, chondropathies and disorders of musculoskeletal system and connective tissue (M83-M99)

White, male

2004.....	612	1	1	1	-	-	-	-	-	4	7	16
2003.....	590	-	-	-	1	-	1	1	1	2	11	16
2002.....	579	-	-	-	-	-	1	1	1	4	8	11
2001.....	540	1	-	-	-	1	1	-	3	10	5	11
2000.....	507	-	-	-	1	-	3	2	3	7	11	8
1999.....	446	-	-	-	-	1	-	1	2	2	2	13

White, female

2004.....	758	-	-	-	-	-	-	1	1	4	11	15
2003.....	753	1	-	-	-	3	2	-	2	-	8	15
2002.....	676	-	-	1	1	-	-	2	1	6	2	10
2001.....	724	-	-	-	-	-	-	-	5	3	5	13
2000.....	596	2	-	-	-	1	1	2	4	3	-	3
1999.....	577	-	-	-	-	-	-	1	-	1	2	8

Black, both sexes

2004.....	260	3	-	-	-	-	1	1	1	3	2	13
2003.....	265	-	1	-	1	-	-	4	1	4	2	11
2002.....	236	-	-	1	1	-	1	2	-	1	1	3
2001.....	239	-	-	-	-	-	-	2	3	-	4	8
2000.....	228	-	-	-	1	1	-	2	2	3	3	7
1999.....	178	-	1	-	-	-	1	1	-	2	5	3

Black, male

2004.....	120	2	-	-	-	-	1	1	1	2	2	8
2003.....	133	-	-	-	1	-	-	2	-	4	2	6
2002.....	110	-	-	-	1	-	-	1	-	1	1	2
2001.....	106	-	-	-	-	-	-	2	2	-	3	6
2000.....	110	-	-	-	1	1	-	2	1	3	3	5
1999.....	73	-	1	-	-	-	1	-	-	-	2	3

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

XIV. Diseases of the genitourinary system (N00-N98)

All races, both sexes

2004.....	60,326	200	26	12	17	35	81	123	195	360	631	1,027
2003.....	60,081	206	21	9	22	43	92	130	231	338	653	1,003
2002.....	58,334	187	19	13	17	32	80	135	205	348	587	964
2001.....	57,026	157	25	9	10	26	88	112	191	377	582	930
2000.....	54,560	191	22	13	19	42	73	105	203	342	574	855
1999.....	53,127	192	24	21	27	35	59	132	189	339	544	764

All races, male

2004.....	26,792	121	16	5	10	16	46	57	81	186	349	574
2003.....	26,624	122	10	3	10	23	46	61	119	173	356	542
2002.....	25,673	116	10	5	9	14	38	54	103	170	309	532
2001.....	24,771	87	10	3	4	17	40	57	89	196	306	516
2000.....	23,749	118	9	7	10	23	30	53	101	174	325	505
1999.....	23,062	97	14	12	14	15	26	56	96	180	292	436

All races, female

2004.....	33,534	79	10	7	7	19	35	66	114	174	282	453
2003.....	33,457	84	11	6	12	20	46	69	112	165	297	461
2002.....	32,661	71	9	8	8	18	42	81	102	178	278	432
2001.....	32,255	70	15	6	6	9	48	55	102	181	276	414
2000.....	30,811	73	13	6	9	19	43	52	102	168	249	350
1999.....	30,065	95	10	9	13	20	33	76	93	159	252	328

White, both sexes

2004.....	49,145	113	18	9	15	25	51	72	106	215	386	603
2003.....	49,003	129	14	8	15	30	53	81	131	185	370	597
2002.....	47,680	124	15	10	11	16	43	70	126	196	352	580
2001.....	46,518	94	19	5	6	18	59	67	115	223	354	534
2000.....	44,591	108	15	9	13	27	58	69	116	188	342	505
1999.....	43,289	118	19	18	17	23	32	75	111	194	312	435

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
XIV. Diseases of the genitourinary system (N00-N98)												
White, male												
2004.....	21,854	69	11	3	8	11	28	34	37	104	229	339
2003.....	21,735	77	7	2	8	17	26	40	66	90	216	329
2002.....	20,982	75	7	4	6	8	23	29	68	101	186	320
2001.....	20,315	54	6	3	3	13	25	29	47	136	192	289
2000.....	19,460	70	6	6	8	15	27	35	57	94	196	290
1999.....	18,732	69	9	10	10	12	12	33	58	104	163	244
White, female												
2004.....	27,291	44	7	6	7	14	23	38	69	111	157	264
2003.....	27,268	52	7	6	7	13	27	41	65	95	154	268
2002.....	26,698	49	8	6	5	8	20	41	58	95	166	260
2001.....	26,203	40	13	2	3	5	34	38	68	87	162	245
2000.....	25,131	38	9	3	5	12	31	34	59	94	146	215
1999.....	24,557	49	10	8	7	11	20	42	53	90	149	191
Black, both sexes												
2004.....	9,904	77	7	3	1	9	28	44	79	136	228	393
2003.....	9,907	73	6	1	7	13	31	43	95	145	261	369
2002.....	9,500	56	4	3	3	14	32	61	73	140	221	347
2001.....	9,354	60	4	4	3	5	24	43	71	137	217	358
2000.....	8,928	78	5	3	5	14	14	33	80	144	211	331
1999.....	8,808	67	4	3	9	11	26	50	68	136	218	309
Black, male												
2004.....	4,318	44	5	2	1	5	16	18	41	77	111	214
2003.....	4,372	42	2	1	2	6	16	18	52	80	126	190
2002.....	4,169	36	3	1	1	5	12	24	32	63	113	195
2001.....	3,933	31	3	-	-	2	12	27	41	52	105	209
2000.....	3,817	44	2	1	1	8	3	15	40	77	117	206
1999.....	3,847	25	4	2	4	3	13	19	31	73	119	183

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years years

XIV. Diseases of the genitourinary system (N00-N98)

Black, female

2004.....	5,586	33	2	1	-	4	12	26	38	59	117	179
2003.....	5,535	31	4	-	5	7	15	25	43	65	135	179
2002.....	5,331	20	1	2	2	9	20	37	41	77	108	152
2001.....	5,421	29	1	4	3	3	12	16	30	85	112	149
2000.....	5,111	34	3	2	4	6	11	18	40	67	94	125
1999.....	4,961	42	-	1	5	8	13	31	37	63	99	126

American Indian 1/, both sexes

2004.....	322	1	-	-	1	1	1	2	8	6	7	12
2003.....	341	1	-	-	-	-	5	2	2	4	13	19
2002.....	304	2	-	-	2	-	2	2	4	4	7	20
2001.....	324	2	1	-	1	-	4	-	2	7	6	18
2000.....	284	2	-	-	-	-	-	2	3	3	13	8
1999.....	248	-	-	-	-	-	1	2	5	2	9	11

American Indian 1/, male

2004.....	142	1	-	-	1	-	1	2	1	2	3	6
2003.....	148	1	-	-	-	-	1	1	1	-	9	10
2002.....	126	2	-	-	2	-	2	1	2	3	6	7
2001.....	136	2	-	-	1	-	3	-	-	3	5	9
2000.....	123	1	-	-	-	-	-	2	1	-	8	2
1999.....	100	-	-	-	-	-	1	2	4	1	7	5

American Indian 1/, female

2004.....	180	-	-	-	-	1	-	-	7	4	4	6
2003.....	193	-	-	-	-	-	4	1	1	4	4	9
2002.....	178	-	-	-	-	-	-	1	2	1	1	13
2001.....	188	-	1	-	-	-	1	-	2	4	1	9
2000.....	161	1	-	-	-	-	-	-	2	3	5	6
1999.....	148	-	-	-	-	-	-	-	1	1	2	6

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
XIV. Diseases of the genitourinary system (N00-N98)												
Asian or Pacific Islander 2/, both sexes												
2004.....	955	9	1	-	-	-	1	5	2	3	10	19
2003.....	830	3	1	-	-	-	3	4	3	4	9	18
2002.....	850	5	-	-	1	2	3	2	2	8	7	17
2001.....	830	1	1	-	-	3	1	2	3	10	5	20
2000.....	757	3	2	1	1	1	1	1	4	7	8	11
1999.....	782	7	1	-	1	1	-	5	5	7	5	9
Asian or Pacific Islander 2/, male												
2004.....	478	7	-	-	-	-	1	3	2	3	6	15
2003.....	369	2	1	-	-	-	3	2	-	3	5	13
2002.....	396	3	-	-	-	1	1	-	1	3	4	10
2001.....	387	-	1	-	-	2	-	1	1	5	4	9
2000.....	349	3	1	-	1	-	-	1	3	3	4	7
1999.....	383	3	1	-	-	-	-	2	3	2	3	4
Asian or Pacific Islander 2/, female												
2004.....	477	2	1	-	-	-	-	2	-	-	4	4
2003.....	461	1	-	-	-	-	-	2	3	1	4	5
2002.....	454	2	-	-	1	1	2	2	1	5	3	7
2001.....	443	1	-	-	-	1	1	1	2	5	1	11
2000.....	408	-	1	1	-	1	1	-	1	4	4	4
1999.....	399	4	-	-	1	1	-	3	2	5	2	5

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of urinary system (N00-N39)												
All races, both sexes												
2004.....	59,195	196	26	12	17	33	75	119	190	351	612	997
2003.....	59,046	202	21	9	21	41	83	126	226	321	629	983
2002.....	57,394	184	19	13	17	30	78	134	196	325	572	940
2001.....	56,054	150	25	9	10	26	86	109	187	368	565	902
2000.....	53,615	184	22	13	18	42	69	101	193	325	556	837
1999.....	52,189	177	23	21	25	34	56	128	182	330	528	748
All races, male												
2004.....	26,062	121	16	5	10	16	43	55	80	185	344	562
2003.....	26,012	121	10	3	10	23	46	61	119	169	353	539
2002.....	25,096	116	10	5	9	14	38	54	103	169	309	529
2001.....	24,178	86	10	3	4	17	40	57	89	193	303	507
2000.....	23,187	117	9	7	10	23	30	53	99	172	324	504
1999.....	22,470	97	14	12	14	15	26	56	96	179	290	434
All races, female												
2004.....	33,133	75	10	7	7	17	32	64	110	166	268	435
2003.....	33,034	81	11	6	11	18	37	65	107	152	276	444
2002.....	32,298	68	9	8	8	16	40	80	93	156	263	411
2001.....	31,876	64	15	6	6	9	46	52	98	175	262	395
2000.....	30,428	67	13	6	8	19	39	48	94	153	232	333
1999.....	29,719	80	9	9	11	19	30	72	86	151	238	314
White, both sexes												
2004.....	48,182	112	18	9	15	24	47	68	103	209	373	585
2003.....	48,101	128	14	8	14	28	49	79	130	172	353	583
2002.....	46,889	121	15	10	11	15	42	70	119	185	341	566
2001.....	45,686	89	19	5	6	18	57	64	111	221	345	515
2000.....	43,786	104	15	9	13	27	56	66	110	179	333	491
1999.....	42,505	110	18	18	15	22	31	74	107	190	301	427

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Diseases of urinary system (N00-N39)

White, male

2004.....	21,220	69	11	3	8	11	26	32	37	103	225	329
2003.....	21,189	77	7	2	8	17	26	40	66	87	213	327
2002.....	20,485	75	7	4	6	8	23	29	68	101	186	318
2001.....	19,791	53	6	3	3	13	25	29	47	135	192	282
2000.....	18,974	69	6	6	8	15	27	35	56	93	196	289
1999.....	18,215	69	9	10	10	12	12	33	58	103	161	243

White, female

2004.....	26,962	43	7	6	7	13	21	36	66	106	148	256
2003.....	26,912	51	7	6	6	11	23	39	64	85	140	256
2002.....	26,404	46	8	6	5	7	19	41	51	84	155	248
2001.....	25,895	36	13	2	3	5	32	35	64	86	153	233
2000.....	24,812	35	9	3	5	12	29	31	54	86	137	202
1999.....	24,290	41	9	8	5	10	19	41	49	87	140	184

Black, both sexes

2004.....	9,764	74	7	3	1	8	26	44	77	133	225	382
2003.....	9,791	70	6	1	7	13	27	41	91	141	254	363
2002.....	9,368	56	4	3	3	13	31	60	71	129	217	340
2001.....	9,223	58	4	4	3	5	24	43	71	130	209	349
2000.....	8,807	75	5	3	4	14	12	32	77	139	202	327
1999.....	8,661	60	4	3	9	11	24	47	66	132	213	301

Black, male

2004.....	4,243	44	5	2	1	5	15	18	40	77	110	212
2003.....	4,316	41	2	1	2	6	16	18	52	79	126	189
2002.....	4,101	36	3	1	1	5	12	24	32	62	113	194
2001.....	3,869	31	3	-	-	2	12	27	41	50	102	207
2000.....	3,752	44	2	1	1	8	3	15	39	76	116	206
1999.....	3,776	25	4	2	4	3	13	19	31	73	119	182

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Diseases of urinary system (N00-N39)

Black, female

2004.....	5,521	30	2	1	-	3	11	26	37	56	115	170
2003.....	5,475	29	4	-	5	7	11	23	39	62	128	174
2002.....	5,267	20	1	2	2	8	19	36	39	67	104	146
2001.....	5,354	27	1	4	3	3	12	16	30	80	107	142
2000.....	5,055	31	3	2	3	6	9	17	38	63	86	121
1999.....	4,885	35	-	1	5	8	11	28	35	59	94	119

American Indian 1/, both sexes

2004.....	316	1	-	-	1	1	1	2	8	6	6	12
2003.....	336	1	-	-	-	-	4	2	2	4	13	19
2002.....	300	2	-	-	2	-	2	2	4	4	7	18
2001.....	321	2	1	-	1	-	4	-	2	7	6	18
2000.....	278	2	-	-	-	-	-	2	2	2	13	8
1999.....	245	-	-	-	-	-	1	2	5	2	9	11

American Indian 1/, male

2004.....	138	1	-	-	1	-	1	2	1	2	3	6
2003.....	145	1	-	-	-	-	1	1	1	-	9	10
2002.....	124	2	-	-	2	-	2	1	2	3	6	7
2001.....	135	2	-	-	1	-	3	-	-	3	5	9
2000.....	120	1	-	-	-	-	-	2	1	-	8	2
1999.....	98	-	-	-	-	-	1	2	4	1	7	5

American Indian 1/, female

2004.....	178	-	-	-	-	1	-	-	7	4	3	6
2003.....	191	-	-	-	-	-	3	1	1	4	4	9
2002.....	176	-	-	-	-	-	-	1	2	1	1	11
2001.....	186	-	1	-	-	-	1	-	2	4	1	9
2000.....	158	1	-	-	-	-	-	-	1	2	5	6
1999.....	147	-	-	-	-	-	-	-	1	1	2	6

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of urinary system (N00-N39)												
Asian or Pacific Islander 2/, both sexes												
2004.....	933	9	1	-	-	-	1	5	2	3	8	18
2003.....	818	3	1	-	-	-	3	4	3	4	9	18
2002.....	837	5	-	-	1	2	3	2	2	7	7	16
2001.....	824	1	1	-	-	3	1	2	3	10	5	20
2000.....	744	3	2	1	1	1	1	1	4	5	8	11
1999.....	778	7	1	-	1	1	-	5	4	6	5	9
Asian or Pacific Islander 2/, male												
2004.....	461	7	-	-	-	-	1	3	2	3	6	15
2003.....	362	2	1	-	-	-	3	2	-	3	5	13
2002.....	386	3	-	-	-	1	1	-	1	3	4	10
2001.....	383	-	1	-	-	2	-	1	1	5	4	9
2000.....	341	3	1	-	1	-	-	1	3	3	4	7
1999.....	381	3	1	-	-	-	-	2	3	2	3	4
Asian or Pacific Islander 2/, female												
2004.....	472	2	1	-	-	-	-	2	-	-	2	3
2003.....	456	1	-	-	-	-	-	2	3	1	4	5
2002.....	451	2	-	-	1	1	2	2	1	4	3	6
2001.....	441	1	-	-	-	1	1	1	2	5	1	11
2000.....	403	-	1	1	-	1	1	-	1	2	4	4
1999.....	397	4	-	-	1	1	-	3	1	4	2	5

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Glomerular and renal tubulo-interstitial diseases (N00-N15)

All races, both sexes

2004.....	1,914	16	10	2	1	3	11	13	21	28	43	51
2003.....	1,946	19	5	1	1	7	16	17	18	18	42	56
2002.....	1,930	13	5	1	6	6	8	15	19	27	34	52
2001.....	1,851	17	11	1	1	4	10	12	18	44	33	38
2000.....	1,934	20	6	1	6	2	9	11	19	26	34	50
1999.....	2,076	7	7	5	2	5	8	23	10	28	37	49

All races, male

2004.....	846	7	6	-	-	-	4	5	5	10	18	22
2003.....	847	16	1	-	1	2	9	7	4	9	19	29
2002.....	866	9	2	1	3	2	5	7	11	10	18	19
2001.....	809	10	7	-	-	3	5	4	7	20	15	15
2000.....	818	12	2	1	2	1	2	5	10	12	19	24
1999.....	924	5	5	2	1	3	6	13	5	8	20	22

All races, female

2004.....	1,068	9	4	2	1	3	7	8	16	18	25	29
2003.....	1,099	3	4	1	-	5	7	10	14	9	23	27
2002.....	1,064	4	3	-	3	4	3	8	8	17	16	33
2001.....	1,042	7	4	1	1	1	5	8	11	24	18	23
2000.....	1,116	8	4	-	4	1	7	6	9	14	15	26
1999.....	1,152	2	2	3	1	2	2	10	5	20	17	27

White, both sexes

2004.....	1,626	9	7	2	1	3	8	7	14	21	32	36
2003.....	1,638	11	2	1	1	5	11	9	14	14	33	43
2002.....	1,663	10	4	1	3	3	4	9	13	20	29	32
2001.....	1,576	13	8	-	-	2	8	7	15	28	28	25
2000.....	1,603	16	4	1	4	-	9	7	14	10	21	37
1999.....	1,762	5	5	5	1	2	4	17	7	18	26	42

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Glomerular and renal tubulo-interstitial diseases (N00-N15)												
White, male												
2004.....	715	4	4	-	-	-	3	2	4	8	12	17
2003.....	716	8	-	-	1	1	6	4	4	8	17	25
2002.....	750	7	1	1	1	2	4	6	9	9	13	14
2001.....	680	7	5	-	-	2	4	1	6	17	12	8
2000.....	674	10	1	1	1	-	2	3	7	3	12	20
1999.....	775	5	3	2	1	2	2	10	3	6	14	18
White, female												
2004.....	911	5	3	2	1	3	5	5	10	13	20	19
2003.....	922	3	2	1	-	4	5	5	10	6	16	18
2002.....	913	3	3	-	2	1	-	3	4	11	16	18
2001.....	896	6	3	-	-	-	4	6	9	11	16	17
2000.....	929	6	3	-	3	-	7	4	7	7	9	17
1999.....	987	-	2	3	-	-	2	7	4	12	12	24
Black, both sexes												
2004.....	236	5	3	-	-	-	3	5	7	6	10	11
2003.....	237	7	2	-	-	2	3	4	2	3	9	11
2002.....	211	2	1	-	-	2	4	5	5	6	5	17
2001.....	216	3	1	1	1	-	1	5	2	11	5	10
2000.....	266	4	2	-	1	2	-	3	4	14	8	11
1999.....	261	1	2	-	1	3	3	4	2	8	10	6
Black, male												
2004.....	104	2	2	-	-	-	1	2	1	2	5	4
2003.....	104	7	-	-	-	1	1	1	-	1	2	4
2002.....	91	1	1	-	-	-	1	1	2	-	5	4
2001.....	103	2	1	-	-	-	1	3	1	3	3	6
2000.....	114	2	1	-	-	1	-	1	2	7	3	3
1999.....	126	-	2	-	-	1	3	2	1	2	5	4

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Glomerular and renal tubulo-interstitial diseases (N00-N15)												
Black, female												
2004.....	132	3	1	-	-	-	2	3	6	4	5	7
2003.....	133	-	2	-	-	1	2	3	2	2	7	7
2002.....	120	1	-	-	-	2	3	4	3	6	-	13
2001.....	113	1	-	1	1	-	-	2	1	8	2	4
2000.....	152	2	1	-	1	1	-	2	2	7	5	8
1999.....	135	1	-	-	1	2	-	2	1	6	5	2
American Indian 1/, both sexes												
2004.....	15	1	-	-	-	-	-	1	-	1	-	1
2003.....	16	-	-	-	-	-	-	2	1	1	-	1
2002.....	16	-	-	-	2	-	-	-	1	1	-	2
2001.....	17	1	1	-	-	-	1	-	-	1	-	2
2000.....	15	-	-	-	-	-	-	1	-	-	2	1
1999.....	12	-	-	-	-	-	1	-	1	1	1	-
American Indian 1/, male												
2004.....	6	1	-	-	-	-	-	1	-	-	-	-
2003.....	6	-	-	-	-	-	-	1	-	-	-	-
2002.....	8	-	-	-	2	-	-	-	-	1	-	-
2001.....	10	1	-	-	-	-	-	-	-	-	-	1
2000.....	8	-	-	-	-	-	-	1	-	-	2	-
1999.....	6	-	-	-	-	-	1	-	1	-	1	-
American Indian 1/, female												
2004.....	9	-	-	-	-	-	-	-	-	1	-	1
2003.....	10	-	-	-	-	-	-	1	1	1	-	1
2002.....	8	-	-	-	-	-	-	-	1	-	-	2
2001.....	7	-	1	-	-	-	1	-	-	1	-	1
2000.....	7	-	-	-	-	-	-	-	-	-	-	1
1999.....	6	-	-	-	-	-	-	-	-	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Glomerular and renal tubulo-interstitial diseases (N00-N15)												
Asian or Pacific Islander 2/, both sexes												
2004.....	37	1	-	-	-	-	-	-	-	-	1	3
2003.....	55	1	1	-	-	-	2	2	1	-	-	1
2002.....	40	1	-	-	1	1	-	1	-	-	-	1
2001.....	42	-	1	-	-	2	-	-	1	4	-	1
2000.....	50	-	-	-	1	-	-	-	1	2	3	1
1999.....	41	1	-	-	-	-	-	2	-	1	-	1
Asian or Pacific Islander 2/, male												
2004.....	21	-	-	-	-	-	-	-	-	-	1	1
2003.....	21	1	1	-	-	-	2	1	-	-	-	-
2002.....	17	1	-	-	-	-	-	-	-	-	-	1
2001.....	16	-	1	-	-	1	-	-	-	-	-	-
2000.....	22	-	-	-	1	-	-	-	1	2	2	1
1999.....	17	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	16	1	-	-	-	-	-	-	-	-	-	2
2003.....	34	-	-	-	-	-	-	1	1	-	-	1
2002.....	23	-	-	-	1	1	-	1	-	-	-	-
2001.....	26	-	-	-	-	1	-	-	1	4	-	1
2000.....	28	-	-	-	-	-	-	-	-	-	1	-
1999.....	24	1	-	-	-	-	-	1	-	1	-	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Glomerular diseases (N00-N07)												
All races, both sexes												
2004.....	662	8	5	2	1	1	5	7	9	15	17	18
2003.....	628	9	2	1	-	4	7	7	6	8	14	16
2002.....	660	7	5	1	3	4	3	9	11	15	15	28
2001.....	633	8	6	1	1	3	4	5	8	20	14	14
2000.....	683	8	3	1	4	1	7	6	13	13	21	21
1999.....	716	3	3	4	2	2	6	13	6	13	16	21
All races, male												
2004.....	319	3	4	-	-	-	2	3	2	7	11	10
2003.....	307	8	1	-	-	2	6	4	1	6	10	12
2002.....	316	3	2	1	2	2	2	4	8	7	7	11
2001.....	324	4	4	-	-	3	2	2	1	11	8	8
2000.....	347	3	2	1	2	-	2	3	9	6	13	13
1999.....	359	3	2	2	1	2	5	9	5	5	11	12
All races, female												
2004.....	343	5	1	2	1	1	3	4	7	8	6	8
2003.....	321	1	1	1	-	2	1	3	5	2	4	4
2002.....	344	4	3	-	1	2	1	5	3	8	8	17
2001.....	309	4	2	1	1	-	2	3	7	9	6	6
2000.....	336	5	1	-	2	1	5	3	4	7	8	8
1999.....	357	-	1	2	1	-	1	4	1	8	5	9
White, both sexes												
2004.....	534	4	4	2	1	1	3	4	5	11	9	9
2003.....	501	7	1	1	-	2	4	4	3	8	9	14
2002.....	531	5	4	1	2	3	1	5	7	12	13	18
2001.....	518	7	5	-	-	2	4	2	7	12	11	8
2000.....	538	6	1	1	3	-	7	2	10	5	13	15
1999.....	569	3	3	4	1	1	3	9	3	9	8	17

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Glomerular diseases (N00-N07)												
White, male												
2004.....	257	1	4	-	-	-	1	2	1	5	7	6
2003.....	240	6	-	-	-	1	3	2	1	6	8	10
2002.....	261	2	1	1	1	2	1	3	6	6	5	9
2001.....	266	3	3	-	-	2	2	-	1	9	5	3
2000.....	274	3	1	1	1	-	2	1	6	3	8	11
1999.....	286	3	2	2	1	1	2	6	3	3	7	10
White, female												
2004.....	277	3	-	2	1	1	2	2	4	6	2	3
2003.....	261	1	1	1	-	1	1	2	2	2	1	4
2002.....	270	3	3	-	1	1	-	2	1	6	8	9
2001.....	252	4	2	-	-	-	2	2	6	3	6	5
2000.....	264	3	-	-	2	-	5	1	4	2	5	4
1999.....	283	-	1	2	-	-	1	3	-	6	1	7
Black, both sexes												
2004.....	107	2	1	-	-	-	2	2	4	4	8	8
2003.....	86	1	-	-	-	2	1	2	2	-	5	2
2002.....	102	2	1	-	-	1	2	4	3	2	2	10
2001.....	87	-	1	1	1	-	-	3	-	4	3	5
2000.....	113	2	2	-	-	1	-	3	2	8	4	6
1999.....	113	-	-	-	1	1	3	2	2	4	7	3
Black, male												
2004.....	47	1	-	-	-	-	1	-	1	2	4	4
2003.....	45	1	-	-	-	1	1	1	-	-	2	2
2002.....	44	1	1	-	-	-	1	1	2	-	2	2
2001.....	45	-	1	-	-	-	-	2	-	2	3	4
2000.....	53	-	1	-	-	-	-	1	2	3	2	2
1999.....	56	-	-	-	-	1	3	2	1	2	3	2

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Glomerular diseases (N00-N07)												
Asian or Pacific Islander 2/, both sexes												
2004.....	14	1	-	-	-	-	-	-	-	-	-	1
2003.....	34	1	1	-	-	-	2	1	-	-	-	-
2002.....	19	-	-	-	-	-	-	-	-	-	-	-
2001.....	20	-	-	-	-	1	-	-	1	3	-	-
2000.....	23	-	-	-	1	-	-	-	1	-	2	-
1999.....	27	-	-	-	-	-	-	2	-	-	-	1
Asian or Pacific Islander 2/, male												
2004.....	10	-	-	-	-	-	-	-	-	-	-	-
2003.....	17	1	1	-	-	-	2	1	-	-	-	-
2002.....	6	-	-	-	-	-	-	-	-	-	-	-
2001.....	8	-	-	-	-	1	-	-	-	-	-	-
2000.....	13	-	-	-	1	-	-	-	1	-	1	-
1999.....	12	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	4	1	-	-	-	-	-	-	-	-	-	1
2003.....	17	-	-	-	-	-	-	-	-	-	-	-
2002.....	13	-	-	-	-	-	-	-	-	-	-	-
2001.....	12	-	-	-	-	-	-	-	1	3	-	-
2000.....	10	-	-	-	-	-	-	-	-	-	1	-
1999.....	15	-	-	-	-	-	-	1	-	-	-	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute and rapidly progressive nephritic syndrome (N00-N01)												
All races, both sexes												
2004.....	30	-	-	-	-	-	2	-	-	-	-	2
2003.....	18	-	-	-	-	-	-	-	-	1	-	-
2002.....	24	-	-	-	-	-	-	-	-	-	1	-
2001.....	25	-	-	-	-	-	-	-	1	2	2	-
2000.....	24	-	-	-	1	-	-	-	-	-	-	-
1999.....	28	-	-	1	-	-	-	-	1	1	-	-
All races, male												
2004.....	14	-	-	-	-	-	2	-	-	-	-	-
2003.....	7	-	-	-	-	-	-	-	-	-	-	-
2002.....	9	-	-	-	-	-	-	-	-	-	-	-
2001.....	11	-	-	-	-	-	-	-	-	2	1	-
2000.....	15	-	-	-	1	-	-	-	-	-	-	-
1999.....	15	-	-	1	-	-	-	-	1	-	-	-
All races, female												
2004.....	16	-	-	-	-	-	-	-	-	-	-	2
2003.....	11	-	-	-	-	-	-	-	-	1	-	-
2002.....	15	-	-	-	-	-	-	-	-	-	1	-
2001.....	14	-	-	-	-	-	-	-	1	-	1	-
2000.....	9	-	-	-	-	-	-	-	-	-	-	-
1999.....	13	-	-	-	-	-	-	-	-	1	-	-
White, both sexes												
2004.....	24	-	-	-	-	-	1	-	-	-	-	1
2003.....	16	-	-	-	-	-	-	-	-	1	-	-
2002.....	20	-	-	-	-	-	-	-	-	-	1	-
2001.....	22	-	-	-	-	-	-	-	1	2	2	-
2000.....	20	-	-	-	-	-	-	-	-	-	-	-
1999.....	23	-	-	1	-	-	-	-	-	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nephrotic syndrome (N04)												
All races, both sexes												
2004.....	127	7	5	1	-	-	-	1	-	1	1	4
2003.....	130	8	2	1	-	1	3	1	1	-	3	3
2002.....	142	6	5	1	1	2	1	-	1	2	1	5
2001.....	131	8	4	1	-	-	-	2	-	2	2	1
2000.....	153	8	3	1	-	-	1	-	4	1	2	5
1999.....	154	3	2	2	2	-	2	3	1	1	3	6
All races, male												
2004.....	59	2	4	-	-	-	-	1	-	1	-	4
2003.....	58	7	1	-	-	1	3	1	-	-	2	2
2002.....	61	3	2	1	1	1	-	-	1	-	-	4
2001.....	62	4	3	-	-	-	-	2	-	2	1	1
2000.....	76	3	2	1	-	-	1	-	3	-	1	2
1999.....	75	3	2	-	1	-	2	2	1	-	2	3
All races, female												
2004.....	68	5	1	1	-	-	-	-	-	-	1	-
2003.....	72	1	1	1	-	-	-	-	1	-	1	1
2002.....	81	3	3	-	-	1	1	-	-	2	1	1
2001.....	69	4	1	1	-	-	-	-	-	-	1	-
2000.....	77	5	1	-	-	-	-	-	1	1	1	3
1999.....	79	-	-	2	1	-	-	1	-	1	1	3
White, both sexes												
2004.....	103	3	4	1	-	-	-	-	-	-	-	2
2003.....	104	6	1	1	-	1	2	-	1	-	2	3
2002.....	110	5	4	1	1	1	-	-	-	1	1	3
2001.....	108	7	3	-	-	-	-	-	-	2	2	1
2000.....	121	6	1	1	-	-	1	-	3	-	1	2
1999.....	121	3	2	2	1	-	1	1	1	1	1	4

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nephrotic syndrome (N04)												
White, male												
2004.....	48	-	4	-	-	-	-	-	-	-	-	2
2003.....	41	5	-	-	-	1	2	-	-	-	2	2
2002.....	49	2	1	1	1	1	-	-	-	-	-	3
2001.....	50	3	2	-	-	-	-	-	-	2	1	1
2000.....	59	3	1	1	-	-	1	-	2	-	1	2
1999.....	57	3	2	-	1	-	1	-	1	-	1	2
White, female												
2004.....	55	3	-	1	-	-	-	-	-	-	-	-
2003.....	63	1	1	1	-	-	-	-	1	-	-	1
2002.....	61	3	3	-	-	-	-	-	-	1	1	-
2001.....	58	4	1	-	-	-	-	-	-	-	1	-
2000.....	62	3	-	-	-	-	-	-	1	-	-	-
1999.....	64	-	-	2	-	-	-	1	-	1	-	2
Black, both sexes												
2004.....	18	2	1	-	-	-	-	-	-	1	1	2
2003.....	17	1	-	-	-	-	-	1	-	-	1	-
2002.....	25	1	1	-	-	1	1	-	1	1	-	2
2001.....	19	-	1	1	-	-	-	2	-	-	-	-
2000.....	27	2	2	-	-	-	-	-	1	1	1	3
1999.....	26	-	-	-	1	-	1	2	-	-	2	2
Black, male												
2004.....	7	1	-	-	-	-	-	-	-	1	-	2
2003.....	11	1	-	-	-	-	-	1	-	-	-	-
2002.....	9	1	1	-	-	-	-	-	1	-	-	1
2001.....	9	-	1	-	-	-	-	2	-	-	-	-
2000.....	12	-	1	-	-	-	-	-	1	-	-	-
1999.....	14	-	-	-	-	-	1	2	-	-	1	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other and unspecified glomerular diseases (N02-N03,N05-N07)

All races, both sexes

2004.....	505	1	-	1	1	1	3	6	9	14	16	12
2003.....	480	1	-	-	-	3	4	6	5	7	11	13
2002.....	494	1	-	-	2	2	2	9	10	13	13	23
2001.....	477	-	2	-	1	3	4	3	7	16	10	13
2000.....	506	-	-	-	3	1	6	6	9	12	19	16
1999.....	534	-	1	1	-	2	4	10	4	11	13	15

All races, male

2004.....	246	1	-	-	-	-	-	2	2	6	11	6
2003.....	242	1	-	-	-	1	3	3	1	6	8	10
2002.....	246	-	-	-	1	1	2	4	7	7	7	7
2001.....	251	-	1	-	-	3	2	-	1	7	6	7
2000.....	256	-	-	-	1	-	1	3	6	6	12	11
1999.....	269	-	-	1	-	2	3	7	3	5	9	9

All races, female

2004.....	259	-	-	1	1	1	3	4	7	8	5	6
2003.....	238	-	-	-	-	2	1	3	4	1	3	3
2002.....	248	1	-	-	1	1	-	5	3	6	6	16
2001.....	226	-	1	-	1	-	2	3	6	9	4	6
2000.....	250	-	-	-	2	1	5	3	3	6	7	5
1999.....	265	-	1	-	-	-	1	3	1	6	4	6

White, both sexes

2004.....	407	1	-	1	1	1	2	4	5	11	9	6
2003.....	381	1	-	-	-	1	2	4	2	7	7	11
2002.....	401	-	-	-	1	2	1	5	7	11	11	15
2001.....	388	-	2	-	-	2	4	2	6	8	7	7
2000.....	397	-	-	-	3	-	6	2	7	5	12	13
1999.....	425	-	1	1	-	1	2	8	2	7	7	13

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other and unspecified glomerular diseases (N02-N03,N05-N07)

White, male

2004.....	198	1	-	-	-	-	-	2	1	5	7	4
2003.....	193	1	-	-	-	-	1	2	1	6	6	8
2002.....	205	-	-	-	-	1	1	3	6	6	5	6
2001.....	206	-	1	-	-	2	2	-	1	5	3	2
2000.....	202	-	-	-	1	-	1	1	4	3	7	9
1999.....	219	-	-	1	-	1	1	6	2	3	6	8

White, female

2004.....	209	-	-	1	1	1	2	2	4	6	2	2
2003.....	188	-	-	-	-	1	1	2	1	1	1	3
2002.....	196	-	-	-	1	1	-	2	1	5	6	9
2001.....	182	-	1	-	-	-	2	2	5	3	4	5
2000.....	195	-	-	-	2	-	5	1	3	2	5	4
1999.....	206	-	1	-	-	-	1	2	-	4	1	5

Black, both sexes

2004.....	83	-	-	-	-	-	1	2	4	3	7	5
2003.....	68	-	-	-	-	2	1	1	2	-	4	2
2002.....	73	1	-	-	-	-	1	4	2	1	2	8
2001.....	66	-	-	-	1	-	-	1	-	4	3	5
2000.....	83	-	-	-	-	1	-	3	1	7	3	3
1999.....	85	-	-	-	-	1	2	-	2	4	5	1

Black, male

2004.....	37	-	-	-	-	-	-	-	1	1	4	2
2003.....	33	-	-	-	-	1	1	-	-	-	2	2
2002.....	33	-	-	-	-	-	1	1	1	-	2	1
2001.....	36	-	-	-	-	-	-	-	-	2	3	4
2000.....	40	-	-	-	-	-	-	1	1	3	2	2
1999.....	40	-	-	-	-	1	2	-	1	2	2	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other and unspecified glomerular diseases (N02-N03,N05-N07)

Asian or Pacific Islander 2/, both sexes

2004.....	10	-	-	-	-	-	-	-	-	-	-	1
2003.....	25	-	-	-	-	-	1	1	-	-	-	-
2002.....	14	-	-	-	-	-	-	-	-	-	-	-
2001.....	17	-	-	-	-	1	-	-	1	3	-	-
2000.....	18	-	-	-	-	-	-	-	1	-	2	-
1999.....	20	-	-	-	-	-	-	2	-	-	-	1

Asian or Pacific Islander 2/, male

2004.....	8	-	-	-	-	-	-	-	-	-	-	-
2003.....	11	-	-	-	-	-	1	1	-	-	-	-
2002.....	4	-	-	-	-	-	-	-	-	-	-	-
2001.....	6	-	-	-	-	1	-	-	-	-	-	-
2000.....	8	-	-	-	-	-	-	-	1	-	1	-
1999.....	7	-	-	-	-	-	-	1	-	-	-	-

Asian or Pacific Islander 2/, female

2004.....	2	-	-	-	-	-	-	-	-	-	-	1
2003.....	14	-	-	-	-	-	-	-	-	-	-	-
2002.....	10	-	-	-	-	-	-	-	-	-	-	-
2001.....	11	-	-	-	-	-	-	-	1	3	-	-
2000.....	10	-	-	-	-	-	-	-	-	-	1	-
1999.....	13	-	-	-	-	-	-	1	-	-	-	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Renal tubulo-interstitial diseases (N10-N15)												
All races, both sexes												
2004.....	1,252	8	5	-	-	2	6	6	12	13	26	33
2003.....	1,318	10	3	-	1	3	9	10	12	10	28	40
2002.....	1,270	6	-	-	3	2	5	6	8	12	19	24
2001.....	1,218	9	5	-	-	1	6	7	10	24	19	24
2000.....	1,251	12	3	-	2	1	2	5	6	13	13	29
1999.....	1,360	4	4	1	-	3	2	10	4	15	21	28
All races, male												
2004.....	527	4	2	-	-	-	2	2	3	3	7	12
2003.....	540	8	-	-	1	-	3	3	3	3	9	17
2002.....	550	6	-	-	1	-	3	3	3	3	11	8
2001.....	485	6	3	-	-	-	3	2	6	9	7	7
2000.....	471	9	-	-	-	1	-	2	1	6	6	11
1999.....	565	2	3	-	-	1	1	4	-	3	9	10
All races, female												
2004.....	725	4	3	-	-	2	4	4	9	10	19	21
2003.....	778	2	3	-	-	3	6	7	9	7	19	23
2002.....	720	-	-	-	2	2	2	3	5	9	8	16
2001.....	733	3	2	-	-	1	3	5	4	15	12	17
2000.....	780	3	3	-	2	-	2	3	5	7	7	18
1999.....	795	2	1	1	-	2	1	6	4	12	12	18
White, both sexes												
2004.....	1,092	5	3	-	-	2	5	3	9	10	23	27
2003.....	1,137	4	1	-	1	3	7	5	11	6	24	29
2002.....	1,132	5	-	-	1	-	3	4	6	8	16	14
2001.....	1,058	6	3	-	-	-	4	5	8	16	17	17
2000.....	1,065	10	3	-	1	-	2	5	4	5	8	22
1999.....	1,193	2	2	1	-	1	1	8	4	9	18	25

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Renal tubulo-interstitial diseases (N10-N15)												
Black, female												
2004.....	72	2	-	-	-	-	1	1	3	2	1	3
2003.....	92	-	2	-	-	-	2	2	-	2	4	7
2002.....	62	-	-	-	-	1	2	1	2	4	-	5
2001.....	71	1	-	-	-	-	-	1	1	6	2	3
2000.....	92	-	-	-	1	-	-	-	2	2	3	4
1999.....	78	1	-	-	-	2	-	2	-	4	1	1
American Indian 1/, both sexes												
2004.....	8	-	-	-	-	-	-	-	-	1	-	1
2003.....	9	-	-	-	-	-	-	2	-	1	-	1
2002.....	8	-	-	-	1	-	-	-	-	-	-	2
2001.....	9	-	1	-	-	-	1	-	-	-	-	1
2000.....	6	-	-	-	-	-	-	-	-	-	-	1
1999.....	5	-	-	-	-	-	1	-	-	1	-	-
American Indian 1/, male												
2004.....	1	-	-	-	-	-	-	-	-	-	-	-
2003.....	1	-	-	-	-	-	-	1	-	-	-	-
2002.....	3	-	-	-	1	-	-	-	-	-	-	-
2001.....	5	-	-	-	-	-	-	-	-	-	-	-
2000.....	1	-	-	-	-	-	-	-	-	-	-	-
1999.....	1	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, female												
2004.....	7	-	-	-	-	-	-	-	-	1	-	1
2003.....	8	-	-	-	-	-	-	1	-	1	-	1
2002.....	5	-	-	-	-	-	-	-	-	-	-	2
2001.....	4	-	1	-	-	-	1	-	-	-	-	1
2000.....	5	-	-	-	-	-	-	-	-	-	-	1
1999.....	4	-	-	-	-	-	-	-	-	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Renal tubulo-interstitial diseases (N10-N15)												
Asian or Pacific Islander 2/, both sexes												
2004.....	23	-	-	-	-	-	-	-	-	-	1	2
2003.....	21	-	-	-	-	-	-	1	1	-	-	1
2002.....	21	1	-	-	1	1	-	1	-	-	-	1
2001.....	22	-	1	-	-	1	-	-	-	1	-	1
2000.....	27	-	-	-	-	-	-	-	-	2	1	1
1999.....	14	1	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, male												
2004.....	11	-	-	-	-	-	-	-	-	-	1	1
2003.....	4	-	-	-	-	-	-	-	-	-	-	-
2002.....	11	1	-	-	-	-	-	-	-	-	-	1
2001.....	8	-	1	-	-	-	-	-	-	-	-	-
2000.....	9	-	-	-	-	-	-	-	-	2	1	1
1999.....	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	12	-	-	-	-	-	-	-	-	-	-	1
2003.....	17	-	-	-	-	-	-	1	1	-	-	1
2002.....	10	-	-	-	1	1	-	1	-	-	-	-
2001.....	14	-	-	-	-	1	-	-	-	1	-	1
2000.....	18	-	-	-	-	-	-	-	-	-	-	-
1999.....	9	1	-	-	-	-	-	-	-	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Renal failure (N17-N19)												
All races, both sexes												
2004.....	41,732	165	9	7	11	26	53	83	140	277	490	801
2003.....	41,737	170	11	6	15	25	57	95	171	261	510	813
2002.....	40,222	165	9	6	9	17	60	95	148	258	460	757
2001.....	38,784	125	9	8	8	18	64	76	133	278	454	729
2000.....	36,471	152	11	9	10	30	48	72	149	262	439	658
1999.....	34,719	163	12	9	16	21	42	83	147	250	423	586
All races, male												
2004.....	20,006	106	6	3	8	13	33	42	63	149	284	465
2003.....	20,135	98	5	3	6	17	32	49	92	142	301	465
2002.....	19,341	105	5	3	5	8	29	39	75	141	252	440
2001.....	18,509	72	3	3	3	11	29	43	66	146	248	434
2000.....	17,416	100	5	6	8	18	22	39	76	144	263	415
1999.....	16,611	87	7	6	9	8	17	34	76	143	239	352
All races, female												
2004.....	21,726	59	3	4	3	13	20	41	77	128	206	336
2003.....	21,602	72	6	3	9	8	25	46	79	119	209	348
2002.....	20,881	60	4	3	4	9	31	56	73	117	208	317
2001.....	20,275	53	6	5	5	7	35	33	67	132	206	295
2000.....	19,055	52	6	3	2	12	26	33	73	118	176	243
1999.....	18,108	76	5	3	7	13	25	49	71	107	184	234
White, both sexes												
2004.....	33,084	92	6	4	10	18	30	47	67	159	285	450
2003.....	33,133	108	8	5	9	16	34	58	90	129	268	454
2002.....	32,008	107	6	5	6	7	31	48	87	136	255	441
2001.....	30,777	70	7	5	5	13	40	42	72	153	255	394
2000.....	28,982	80	7	7	9	19	36	44	77	137	253	356
1999.....	27,407	99	10	7	10	14	23	42	80	135	227	302

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Renal failure (N17-N19)												
White, male												
2004.....	16,055	59	3	1	7	9	18	25	24	78	182	260
2003.....	16,139	65	4	2	5	13	19	31	46	67	172	267
2002.....	15,558	68	3	3	4	3	16	19	46	76	142	253
2001.....	14,961	42	1	3	2	9	17	22	32	93	148	233
2000.....	14,070	55	4	5	7	11	20	25	39	75	154	223
1999.....	13,312	59	5	5	7	7	8	18	42	79	128	181
White, female												
2004.....	17,029	33	3	3	3	9	12	22	43	81	103	190
2003.....	16,994	43	4	3	4	3	15	27	44	62	96	187
2002.....	16,450	39	3	2	2	4	15	29	41	60	113	188
2001.....	15,816	28	6	2	3	4	23	20	40	60	107	161
2000.....	14,912	25	3	2	2	8	16	19	38	62	99	133
1999.....	14,095	40	5	2	3	7	15	24	38	56	99	121
Black, both sexes												
2004.....	7,715	65	3	3	-	7	21	32	63	111	193	330
2003.....	7,755	59	3	1	6	9	20	35	79	125	226	328
2002.....	7,373	52	3	1	3	9	24	45	56	113	192	287
2001.....	7,178	53	2	3	2	5	21	32	59	113	188	303
2000.....	6,782	68	3	2	1	11	11	26	68	121	172	291
1999.....	6,577	58	1	2	5	7	19	38	61	112	185	267
Black, male												
2004.....	3,483	40	3	2	-	4	13	14	36	67	94	188
2003.....	3,587	31	1	1	1	4	12	17	45	72	118	177
2002.....	3,377	33	2	-	1	4	10	19	26	60	101	171
2001.....	3,138	29	2	-	-	2	10	20	34	45	91	184
2000.....	3,015	42	1	1	1	7	2	12	35	68	102	187
1999.....	2,943	25	1	1	2	1	9	15	29	63	103	162

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Renal failure (N17-N19)												
Asian or Pacific Islander 2/, both sexes												
2004.....	693	8	-	-	-	-	1	3	2	3	7	13
2003.....	601	2	-	-	-	-	-	2	1	4	8	13
2002.....	627	4	-	-	-	1	3	-	2	6	7	15
2001.....	603	1	-	-	-	-	1	2	1	6	5	17
2000.....	502	3	1	-	-	-	1	1	3	3	5	6
1999.....	548	6	1	-	1	-	-	3	3	3	5	6
Asian or Pacific Islander 2/, male												
2004.....	361	7	-	-	-	-	1	2	2	3	5	12
2003.....	287	1	-	-	-	-	-	1	-	3	5	11
2002.....	313	2	-	-	-	1	1	-	1	3	4	9
2001.....	311	-	-	-	-	-	-	1	-	5	4	9
2000.....	241	3	-	-	-	-	-	1	2	1	2	3
1999.....	282	3	1	-	-	-	-	1	2	1	3	4
Asian or Pacific Islander 2/, female												
2004.....	332	1	-	-	-	-	-	1	-	-	2	1
2003.....	314	1	-	-	-	-	-	1	1	1	3	2
2002.....	314	2	-	-	-	-	2	-	1	3	3	6
2001.....	292	1	-	-	-	-	1	1	1	1	1	8
2000.....	261	-	1	-	-	-	1	-	1	2	3	3
1999.....	266	3	-	-	1	-	-	2	1	2	2	2

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Urolithiasis (N20-N23)												
All races, both sexes												
2004.....	301	-	-	-	-	-	-	-	2	1	7	10
2003.....	302	-	-	-	-	-	1	2	3	2	5	8
2002.....	265	-	-	-	-	-	-	3	3	1	6	6
2001.....	271	-	-	-	-	-	2	4	5	1	5	7
2000.....	252	-	-	-	-	1	-	-	-	2	6	6
1999.....	235	-	-	1	-	-	-	-	-	4	3	3
All races, male												
2004.....	121	-	-	-	-	-	-	-	2	1	5	5
2003.....	125	-	-	-	-	-	1	1	2	-	1	3
2002.....	105	-	-	-	-	-	-	-	2	1	2	4
2001.....	117	-	-	-	-	-	1	3	2	1	4	3
2000.....	93	-	-	-	-	1	-	-	-	1	1	3
1999.....	95	-	-	1	-	-	-	-	-	2	1	1
All races, female												
2004.....	180	-	-	-	-	-	-	-	-	-	2	5
2003.....	177	-	-	-	-	-	-	1	1	2	4	5
2002.....	160	-	-	-	-	-	-	3	1	-	4	2
2001.....	154	-	-	-	-	-	1	1	3	-	1	4
2000.....	159	-	-	-	-	-	-	-	-	1	5	3
1999.....	140	-	-	-	-	-	-	-	-	2	2	2
White, both sexes												
2004.....	267	-	-	-	-	-	-	-	2	1	6	6
2003.....	271	-	-	-	-	-	1	2	1	2	4	6
2002.....	240	-	-	-	-	-	-	3	2	1	4	6
2001.....	243	-	-	-	-	-	2	3	4	1	5	6
2000.....	228	-	-	-	-	1	-	-	-	2	6	5
1999.....	211	-	-	1	-	-	-	-	-	4	3	3

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified disorders of kidney (N25-N27)												
All races, both sexes												
2004.....	86	1	-	-	-	-	2	4	3	-	2	3
2003.....	88	2	1	-	-	-	2	1	2	1	2	6
2002.....	92	1	-	2	-	-	-	3	3	1	-	5
2001.....	63	-	1	-	-	-	3	1	2	1	1	6
2000.....	97	2	-	1	-	1	1	2	2	3	2	8
1999.....	90	-	-	-	-	1	-	1	2	5	-	1
All races, male												
2004.....	45	1	-	-	-	-	2	2	1	-	-	3
2003.....	39	-	1	-	-	-	2	-	1	-	-	5
2002.....	38	1	-	1	-	-	-	1	2	-	-	3
2001.....	19	-	-	-	-	-	-	-	-	1	-	3
2000.....	48	-	-	-	-	-	1	2	-	2	1	5
1999.....	46	-	-	-	-	1	-	-	1	4	-	1
All races, female												
2004.....	41	-	-	-	-	-	-	2	2	-	2	-
2003.....	49	2	-	-	-	-	-	1	1	1	2	1
2002.....	54	-	-	1	-	-	-	2	1	1	-	2
2001.....	44	-	1	-	-	-	3	1	2	-	1	3
2000.....	49	2	-	1	-	1	-	-	2	1	1	3
1999.....	44	-	-	-	-	-	-	1	1	1	-	-
White, both sexes												
2004.....	73	1	-	-	-	-	1	2	3	-	1	1
2003.....	73	2	1	-	-	-	-	1	1	1	2	4
2002.....	76	-	-	1	-	-	-	3	2	1	-	4
2001.....	50	-	1	-	-	-	2	1	1	1	-	4
2000.....	78	1	-	-	-	1	-	2	1	2	2	5
1999.....	65	-	-	-	-	1	-	1	2	2	-	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified disorders of kidney (N25-N27)												
White, male												
2004.....	37	1	-	-	-	-	1	2	1	-	-	1
2003.....	29	-	1	-	-	-	-	-	-	-	-	3
2002.....	31	-	-	-	-	-	-	1	1	-	-	2
2001.....	14	-	-	-	-	-	-	-	-	1	-	2
2000.....	41	-	-	-	-	-	-	2	-	2	1	3
1999.....	34	-	-	-	-	1	-	-	1	1	-	1
White, female												
2004.....	36	-	-	-	-	-	-	-	2	-	1	-
2003.....	44	2	-	-	-	-	-	1	1	1	2	1
2002.....	45	-	-	1	-	-	-	2	1	1	-	2
2001.....	36	-	1	-	-	-	2	1	1	-	-	2
2000.....	37	1	-	-	-	1	-	-	1	-	1	2
1999.....	31	-	-	-	-	-	-	1	1	1	-	-
Black, both sexes												
2004.....	12	-	-	-	-	-	1	2	-	-	1	2
2003.....	14	-	-	-	-	-	2	-	1	-	-	2
2002.....	13	1	-	1	-	-	-	-	1	-	-	1
2001.....	9	-	-	-	-	-	1	-	1	-	1	1
2000.....	16	1	-	1	-	-	1	-	1	1	-	3
1999.....	21	-	-	-	-	-	-	-	-	2	-	-
Black, male												
2004.....	7	-	-	-	-	-	1	-	-	-	-	2
2003.....	10	-	-	-	-	-	2	-	1	-	-	2
2002.....	6	1	-	1	-	-	-	-	1	-	-	1
2001.....	3	-	-	-	-	-	-	-	-	-	-	1
2000.....	6	-	-	-	-	-	1	-	-	-	-	2
1999.....	9	-	-	-	-	-	-	-	-	2	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Urinary tract infection, site not specified (N39.0)

All races, both sexes

2004.....	13,913	5	3	2	2	2	5	13	16	28	53	94
2003.....	13,836	2	2	2	1	5	4	6	19	20	49	74
2002.....	13,754	4	3	3	2	4	8	6	15	25	49	83
2001.....	13,981	4	3	-	-	4	5	12	17	23	47	89
2000.....	13,744	5	3	2	1	3	7	13	17	22	35	82
1999.....	14,021	2	2	2	3	5	5	14	19	24	49	83

All races, male

2004.....	4,413	2	1	2	1	2	-	2	5	15	25	45
2003.....	4,302	1	1	-	1	3	2	2	11	9	22	28
2002.....	4,195	1	1	-	1	1	3	1	6	10	24	37
2001.....	4,197	3	-	-	-	3	3	5	7	14	19	34
2000.....	4,257	1	1	-	-	1	2	4	11	8	17	38
1999.....	4,260	1	1	1	1	3	3	6	10	12	20	37

All races, female

2004.....	9,500	3	2	-	1	-	5	11	11	13	28	49
2003.....	9,534	1	1	2	-	2	2	4	8	11	27	46
2002.....	9,559	3	2	3	1	3	5	5	9	15	25	46
2001.....	9,784	1	3	-	-	1	2	7	10	9	28	55
2000.....	9,487	4	2	2	1	2	5	9	6	14	18	44
1999.....	9,761	1	1	1	2	2	2	8	9	12	29	46

White, both sexes

2004.....	12,109	2	2	2	1	1	4	10	13	19	38	64
2003.....	12,042	1	1	2	1	4	1	6	17	16	31	54
2002.....	11,976	3	3	2	2	3	7	2	11	17	35	64
2001.....	12,126	3	2	-	-	3	4	8	13	20	40	63
2000.....	11,974	4	2	1	-	3	7	12	12	20	22	65
1999.....	12,204	1	2	2	2	3	3	9	14	17	34	61

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Urinary tract infection, site not specified (N39.0)

White, male

2004.....	3,776	1	1	2	-	1	-	2	4	11	17	32
2003.....	3,734	-	-	-	1	2	-	2	10	7	17	23
2002.....	3,600	-	1	-	1	1	3	-	6	9	18	31
2001.....	3,595	3	-	-	-	2	2	3	6	12	17	23
2000.....	3,640	1	-	-	-	1	2	4	8	8	10	28
1999.....	3,583	1	1	1	-	2	2	4	8	9	12	28

White, female

2004.....	8,333	1	1	-	1	-	4	8	9	8	21	32
2003.....	8,308	1	1	2	-	2	1	4	7	9	14	31
2002.....	8,376	3	2	2	1	2	4	2	5	8	17	33
2001.....	8,531	-	2	-	-	1	2	5	7	8	23	40
2000.....	8,334	3	2	1	-	2	5	8	4	12	12	37
1999.....	8,621	-	1	1	2	1	1	5	6	8	22	33

Black, both sexes

2004.....	1,568	3	-	-	1	1	1	3	3	8	14	27
2003.....	1,586	1	1	-	-	1	1	-	2	4	14	18
2002.....	1,565	1	-	1	-	1	1	4	4	7	13	17
2001.....	1,627	1	1	-	-	-	1	4	4	3	7	24
2000.....	1,558	1	-	-	1	-	-	1	4	1	11	13
1999.....	1,615	1	-	-	1	1	2	4	3	6	14	21

Black, male

2004.....	547	1	-	-	1	1	-	-	1	3	8	12
2003.....	510	1	1	-	-	1	1	-	1	2	3	4
2002.....	527	1	-	-	-	-	-	1	-	1	5	6
2001.....	532	-	-	-	-	-	1	2	1	2	2	11
2000.....	534	-	-	-	-	-	-	-	2	-	6	9
1999.....	596	-	-	-	1	1	1	1	1	3	8	9

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Urinary tract infection, site not specified (N39.0)												
Black, female												
2004.....	1,021	2	-	-	-	-	1	3	2	5	6	15
2003.....	1,076	-	-	-	-	-	-	-	1	2	11	14
2002.....	1,038	-	-	1	-	1	1	3	4	6	8	11
2001.....	1,095	1	1	-	-	-	-	2	3	1	5	13
2000.....	1,024	1	-	-	1	-	-	1	2	1	5	4
1999.....	1,019	1	-	-	-	-	1	3	2	3	6	12
American Indian 1/, both sexes												
2004.....	57	-	-	-	-	-	-	-	-	1	1	3
2003.....	65	-	-	-	-	-	1	-	-	-	3	-
2002.....	62	-	-	-	-	-	-	-	-	-	1	2
2001.....	69	-	-	-	-	-	-	-	-	-	-	-
2000.....	51	-	-	-	-	-	-	-	1	1	2	2
1999.....	38	-	-	-	-	-	-	1	1	-	1	-
American Indian 1/, male												
2004.....	21	-	-	-	-	-	-	-	-	1	-	1
2003.....	14	-	-	-	-	-	-	-	-	-	2	-
2002.....	18	-	-	-	-	-	-	-	-	-	1	-
2001.....	21	-	-	-	-	-	-	-	-	-	-	-
2000.....	18	-	-	-	-	-	-	-	1	-	1	-
1999.....	11	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, female												
2004.....	36	-	-	-	-	-	-	-	-	-	1	2
2003.....	51	-	-	-	-	-	1	-	-	-	1	-
2002.....	44	-	-	-	-	-	-	-	-	-	-	2
2001.....	48	-	-	-	-	-	-	-	-	-	-	-
2000.....	33	-	-	-	-	-	-	-	-	1	1	2
1999.....	27	-	-	-	-	-	-	-	1	-	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Urinary tract infection, site not specified (N39.0)												
Asian or Pacific Islander 2/, both sexes												
2004.....	179	-	1	-	-	-	-	-	-	-	-	-
2003.....	143	-	-	-	-	-	1	-	-	-	1	2
2002.....	151	-	-	-	-	-	-	-	-	1	-	-
2001.....	159	-	-	-	-	1	-	-	-	-	-	2
2000.....	161	-	1	1	-	-	-	-	-	-	-	2
1999.....	164	-	-	-	-	1	-	-	1	1	-	1
Asian or Pacific Islander 2/, male												
2004.....	69	-	-	-	-	-	-	-	-	-	-	-
2003.....	44	-	-	-	-	-	1	-	-	-	-	1
2002.....	50	-	-	-	-	-	-	-	-	-	-	-
2001.....	49	-	-	-	-	1	-	-	-	-	-	-
2000.....	65	-	1	-	-	-	-	-	-	-	-	1
1999.....	70	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	110	-	1	-	-	-	-	-	-	-	-	-
2003.....	99	-	-	-	-	-	-	-	-	-	1	1
2002.....	101	-	-	-	-	-	-	-	-	1	-	-
2001.....	110	-	-	-	-	-	-	-	-	-	-	2
2000.....	96	-	-	1	-	-	-	-	-	-	-	1
1999.....	94	-	-	-	-	1	-	-	-	1	-	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other diseases of urinary system (N28-N36,N39.1-N39.9)

All races, both sexes

2004.....	1,249	9	4	1	3	2	4	6	8	17	17	38
2003.....	1,137	9	2	-	4	4	3	5	13	19	21	26
2002.....	1,131	1	2	1	-	3	2	12	8	13	23	37
2001.....	1,104	4	1	-	1	-	2	4	12	21	25	33
2000.....	1,117	5	2	-	1	5	4	3	6	10	40	33
1999.....	1,048	5	2	4	4	2	1	7	4	19	16	26

All races, male

2004.....	631	5	3	-	1	1	4	4	4	10	12	22
2003.....	564	6	2	-	2	1	-	2	9	9	10	9
2002.....	551	-	2	-	-	3	1	6	7	7	13	26
2001.....	527	1	-	-	1	-	2	2	7	11	17	18
2000.....	555	4	1	-	-	2	3	3	2	5	23	19
1999.....	534	4	1	2	3	-	-	3	4	10	10	21

All races, female

2004.....	618	4	1	1	2	1	-	2	4	7	5	16
2003.....	573	3	-	-	2	3	3	3	4	10	11	17
2002.....	580	1	-	1	-	-	1	6	1	6	10	11
2001.....	577	3	1	-	-	-	-	2	5	10	8	15
2000.....	562	1	1	-	1	3	1	-	4	5	17	14
1999.....	514	1	1	2	1	2	1	4	-	9	6	5

White, both sexes

2004.....	1,023	8	3	1	3	2	4	2	4	9	11	28
2003.....	944	6	2	-	3	3	2	3	7	10	15	22
2002.....	926	1	2	1	-	2	-	5	4	10	18	19
2001.....	914	3	1	-	1	-	1	3	6	18	17	23
2000.....	921	3	2	-	-	3	4	1	6	8	29	23
1999.....	856	5	1	3	2	2	1	5	4	14	11	18

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other diseases of urinary system (N28-N36,N39.1-N39.9)

White, male

2004.....	529	4	3	-	1	1	4	1	2	5	9	17
2003.....	464	4	2	-	1	1	-	2	5	5	7	7
2002.....	451	-	2	-	-	2	-	3	4	6	11	14
2001.....	436	1	-	-	1	-	1	1	2	11	11	13
2000.....	462	3	1	-	-	2	3	1	2	4	18	13
1999.....	428	4	-	1	2	-	-	1	4	6	6	14

White, female

2004.....	494	4	-	1	2	1	-	1	2	4	2	11
2003.....	480	2	-	-	2	2	2	1	2	5	8	15
2002.....	475	1	-	1	-	-	-	2	-	4	7	5
2001.....	478	2	1	-	-	-	-	2	4	7	6	10
2000.....	459	-	1	-	-	1	1	-	4	4	11	10
1999.....	428	1	1	2	-	2	1	4	-	8	5	4

Black, both sexes

2004.....	207	1	1	-	-	-	-	2	4	8	6	9
2003.....	177	3	-	-	1	1	1	2	6	9	5	4
2002.....	184	-	-	-	-	1	2	6	4	3	5	18
2001.....	168	1	-	-	-	-	-	1	4	3	8	10
2000.....	168	1	-	-	1	1	-	2	-	2	11	8
1999.....	168	-	1	1	2	-	-	1	-	4	4	7

Black, male

2004.....	93	1	-	-	-	-	-	2	2	5	3	4
2003.....	92	2	-	-	1	-	-	-	4	4	3	2
2002.....	90	-	-	-	-	1	1	3	3	1	2	12
2001.....	81	-	-	-	-	-	-	1	4	-	6	5
2000.....	78	-	-	-	-	-	-	2	-	1	5	4
1999.....	93	-	1	1	1	-	-	1	-	3	3	7

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Diseases of male genital organs (N40-N50)

All races, both sexes

2004.....	727	-	-	-	-	-	3	2	1	1	5	12
2003.....	610	1	-	-	-	-	-	-	-	4	3	3
2002.....	574	-	-	-	-	-	-	-	-	1	-	3
2001.....	592	1	-	-	-	-	-	-	-	3	3	9
2000.....	561	1	-	-	-	-	-	-	2	2	1	1
1999.....	590	-	-	-	-	-	-	-	-	1	2	2

All races, male

2004.....	727	-	-	-	-	-	3	2	1	1	5	12
2003.....	610	1	-	-	-	-	-	-	-	4	3	3
2002.....	574	-	-	-	-	-	-	-	-	1	-	3
2001.....	592	1	-	-	-	-	-	-	-	3	3	9
2000.....	561	1	-	-	-	-	-	-	2	2	1	1
1999.....	590	-	-	-	-	-	-	-	-	1	2	2

All races, female

2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

White, both sexes

2004.....	632	-	-	-	-	-	2	2	-	1	4	10
2003.....	544	-	-	-	-	-	-	-	-	3	3	2
2002.....	495	-	-	-	-	-	-	-	-	-	-	2
2001.....	523	1	-	-	-	-	-	-	-	1	-	7
2000.....	485	1	-	-	-	-	-	-	1	1	-	1
1999.....	515	-	-	-	-	-	-	-	-	1	2	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Disorders of female genital organs (N70-N98)

All races, both sexes

2004.....	342	4	-	-	-	2	3	2	4	7	11	18
2003.....	374	3	-	-	1	2	8	4	5	13	19	16
2002.....	315	3	-	-	-	2	2	1	9	21	11	20
2001.....	315	6	-	-	-	-	2	3	4	6	14	15
2000.....	341	6	-	-	1	-	4	3	8	14	16	16
1999.....	307	15	1	-	2	1	2	3	7	8	13	11

All races, male

2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

All races, female

2004.....	342	4	-	-	-	2	3	2	4	7	11	18
2003.....	374	3	-	-	1	2	8	4	5	13	19	16
2002.....	315	3	-	-	-	2	2	1	9	21	11	20
2001.....	315	6	-	-	-	-	2	3	4	6	14	15
2000.....	341	6	-	-	1	-	4	3	8	14	16	16
1999.....	307	15	1	-	2	1	2	3	7	8	13	11

White, both sexes

2004.....	278	1	-	-	-	1	2	2	3	5	7	8
2003.....	320	1	-	-	1	2	4	2	1	10	12	12
2002.....	255	3	-	-	-	1	1	-	7	10	9	11
2001.....	253	4	-	-	-	-	2	3	4	1	9	10
2000.....	285	3	-	-	-	-	2	2	5	8	9	12
1999.....	239	8	1	-	2	1	1	1	4	3	8	5

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of female genital organs (N70-N98)												
Asian or Pacific Islander 2/, both sexes												
2004.....	4	-	-	-	-	-	-	-	-	-	1	1
2003.....	3	-	-	-	-	-	-	-	-	-	-	-
2002.....	3	-	-	-	-	-	-	-	-	1	-	1
2001.....	2	-	-	-	-	-	-	-	-	-	-	-
2000.....	4	-	-	-	-	-	-	-	-	1	-	-
1999.....	2	-	-	-	-	-	-	-	1	1	-	-
Asian or Pacific Islander 2/, male												
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....
Asian or Pacific Islander 2/, female												
2004.....	4	-	-	-	-	-	-	-	-	-	1	1
2003.....	3	-	-	-	-	-	-	-	-	-	-	-
2002.....	3	-	-	-	-	-	-	-	-	1	-	1
2001.....	2	-	-	-	-	-	-	-	-	-	-	-
2000.....	4	-	-	-	-	-	-	-	-	1	-	-
1999.....	2	-	-	-	-	-	-	-	1	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Inflammatory diseases of female pelvic organs (N70-N76)												
All races, both sexes												
2004.....	112	1	-	-	-	1	1	2	2	4	4	8
2003.....	125	1	-	-	1	1	5	1	2	7	7	7
2002.....	114	1	-	-	-	1	1	-	4	12	5	4
2001.....	127	1	-	-	-	-	2	2	1	2	8	9
2000.....	120	2	-	-	1	-	2	2	4	9	6	6
1999.....	108	2	1	-	-	1	2	3	3	4	8	4
All races, male												
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....
All races, female												
2004.....	112	1	-	-	-	1	1	2	2	4	4	8
2003.....	125	1	-	-	1	1	5	1	2	7	7	7
2002.....	114	1	-	-	-	1	1	-	4	12	5	4
2001.....	127	1	-	-	-	-	2	2	1	2	8	9
2000.....	120	2	-	-	1	-	2	2	4	9	6	6
1999.....	108	2	1	-	-	1	2	3	3	4	8	4
White, both sexes												
2004.....	92	1	-	-	-	1	1	2	1	4	2	4
2003.....	101	-	-	-	1	1	2	1	1	6	4	5
2002.....	92	1	-	-	-	-	-	-	3	4	5	2
2001.....	103	1	-	-	-	-	2	2	1	-	6	6
2000.....	94	1	-	-	-	-	2	2	2	5	2	4
1999.....	80	1	1	-	-	1	1	1	2	1	3	2

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Inflammatory diseases of female pelvic organs (N70-N76)

Asian or Pacific Islander 2/, both sexes

2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	2	-	-	-	-	-	-	-	-	-	-	-
2002.....	1	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	1	-	-	-	-	-	-	-	-	1	-	-
1999.....	2	-	-	-	-	-	-	-	1	1	-	-

Asian or Pacific Islander 2/, male

2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

Asian or Pacific Islander 2/, female

2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	2	-	-	-	-	-	-	-	-	-	-	-
2002.....	1	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	1	-	-	-	-	-	-	-	-	1	-	-
1999.....	2	-	-	-	-	-	-	-	1	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Noninflammatory disorders of female genital tract (N80-N98)

All races, both sexes

2004.....	230	3	-	-	-	1	2	-	2	3	7	10
2003.....	249	2	-	-	-	1	3	3	3	6	12	9
2002.....	201	2	-	-	-	1	1	1	5	9	6	16
2001.....	188	5	-	-	-	-	-	1	3	4	6	6
2000.....	221	4	-	-	-	-	2	1	4	5	10	10
1999.....	199	13	-	-	2	-	-	-	4	4	5	7

All races, male

2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

All races, female

2004.....	230	3	-	-	-	1	2	-	2	3	7	10
2003.....	249	2	-	-	-	1	3	3	3	6	12	9
2002.....	201	2	-	-	-	1	1	1	5	9	6	16
2001.....	188	5	-	-	-	-	-	1	3	4	6	6
2000.....	221	4	-	-	-	-	2	1	4	5	10	10
1999.....	199	13	-	-	2	-	-	-	4	4	5	7

White, both sexes

2004.....	186	-	-	-	-	-	1	-	2	1	5	4
2003.....	219	1	-	-	-	1	2	1	-	4	8	7
2002.....	163	2	-	-	-	1	1	-	4	6	4	9
2001.....	150	3	-	-	-	-	-	1	3	1	3	4
2000.....	191	2	-	-	-	-	-	-	3	3	7	8
1999.....	159	7	-	-	2	-	-	-	2	2	5	3

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

XV. Pregnancy, childbirth and the puerperium (000-099)

All races, both sexes

2004.....	697	2	37	131	142	145	96	61	39
2003.....	545	1	27	89	110	119	106	60	19
2002.....	379	1	28	61	82	93	79	26	8
2001.....	416	3	37	73	93	97	73	33	5
2000.....	404	-	18	77	87	95	86	35	5
1999.....	406	-	33	64	89	92	88	33	5

All races, male

2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

All races, female

2004.....	697	2	37	131	142	145	96	61	39
2003.....	545	1	27	89	110	119	106	60	19
2002.....	379	1	28	61	82	93	79	26	8
2001.....	416	3	37	73	93	97	73	33	5
2000.....	404	-	18	77	87	95	86	35	5
1999.....	406	-	33	64	89	92	88	33	5

White, both sexes

2004.....	411	-	24	64	76	92	60	39	21
2003.....	317	-	15	51	73	59	56	38	14
2002.....	203	-	12	28	41	56	41	19	5
2001.....	235	2	22	42	50	64	33	18	2
2000.....	243	-	11	44	53	55	56	21	2
1999.....	220	-	14	31	47	53	55	16	4

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
XV. Pregnancy, childbirth and the puerperium (O00-O99)												
Black, female												
2004.....	251	-	12	62	62	47	30	15	16
2003.....	193	1	12	32	32	51	40	18	5
2002.....	157	1	16	30	38	28	35	7	2
2001.....	158	1	14	31	36	30	30	13	3
2000.....	142	-	4	32	31	34	26	12	3
1999.....	162	-	18	29	38	33	28	14	-
American Indian 1/, both sexes												
2004.....	6	-	-	1	1	-	1	2	1
2003.....	8	-	-	1	1	1	4	1	-
2002.....	-	-	-	-	-	-	-	-	-
2001.....	5	-	-	-	3	2	-	-	-
2000.....	6	-	1	1	1	3	-	-	-
1999.....	5	-	1	2	-	1	1	-	-
American Indian 1/, male												
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....
American Indian 1/, female												
2004.....	6	-	-	1	1	-	1	2	1
2003.....	8	-	-	1	1	1	4	1	-
2002.....	-	-	-	-	-	-	-	-	-
2001.....	5	-	-	-	3	2	-	-	-
2000.....	6	-	1	1	1	3	-	-	-
1999.....	5	-	1	2	-	1	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
XV. Pregnancy, childbirth and the puerperium (000-099)												
Asian or Pacific Islander 2/, both sexes												
2004.....	29	2	1	4	3	6	5	5	1
2003.....	27	-	-	5	4	8	6	3	-
2002.....	19	-	-	3	3	9	3	-	1
2001.....	18	-	1	-	4	1	10	2	-
2000.....	13	-	2	-	2	3	4	2	-
1999.....	19	-	-	2	4	5	4	3	1
Asian or Pacific Islander 2/, male												
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....
Asian or Pacific Islander 2/, female												
2004.....	29	2	1	4	3	6	5	5	1
2003.....	27	-	-	5	4	8	6	3	-
2002.....	19	-	-	3	3	9	3	-	1
2001.....	18	-	1	-	4	1	10	2	-
2000.....	13	-	2	-	2	3	4	2	-
1999.....	19	-	-	2	4	5	4	3	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Pregnancy with abortive outcome (000-007)

All races, both sexes

2004.....	32	-	1	13	6	5	4	2	-
2003.....	40	-	6	5	8	11	6	3	1
2002.....	22	1	-	4	4	5	6	1	1
2001.....	38	2	6	7	6	8	8	1	-
2000.....	37	-	-	6	12	5	10	4	-
1999.....	31	-	3	5	5	10	6	2	-

All races, male

2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

All races, female

2004.....	32	-	1	13	6	5	4	2	-
2003.....	40	-	6	5	8	11	6	3	1
2002.....	22	1	-	4	4	5	6	1	1
2001.....	38	2	6	7	6	8	8	1	-
2000.....	37	-	-	6	12	5	10	4	-
1999.....	31	-	3	5	5	10	6	2	-

White, both sexes

2004.....	11	-	-	5	3	2	-	-	-
2003.....	18	-	1	2	7	4	3	1	-
2002.....	7	-	-	1	2	3	-	1	-
2001.....	20	1	3	4	4	4	3	1	-
2000.....	18	-	-	1	7	1	6	3	-
1999.....	15	-	2	2	1	5	3	2	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Pregnancy with abortive outcome (000-007)

Black, female

2004.....	19	-	1	7	3	3	4	1	-
2003.....	18	-	5	3	-	5	2	2	1
2002.....	12	1	-	2	2	2	5	-	-
2001.....	16	1	3	3	2	3	4	-	-
2000.....	19	-	-	5	5	4	4	1	-
1999.....	15	-	1	3	4	4	3	-	-

American Indian 1/, both sexes

2004.....	-	-	-	-	-	-	-	-	-
2003.....	1	-	-	-	-	-	1	-	-
2002.....	-	-	-	-	-	-	-	-	-
2001.....	1	-	-	-	-	1	-	-	-
2000.....	-	-	-	-	-	-	-	-	-
1999.....	1	-	-	-	-	1	-	-	-

American Indian 1/, male

2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

American Indian 1/, female

2004.....	-	-	-	-	-	-	-	-	-
2003.....	1	-	-	-	-	-	1	-	-
2002.....	-	-	-	-	-	-	-	-	-
2001.....	1	-	-	-	-	1	-	-	-
2000.....	-	-	-	-	-	-	-	-	-
1999.....	1	-	-	-	-	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Ectopic pregnancy (000)												
All races, both sexes												
2004.....	14	-	-	5	3	3	2	1	-
2003.....	23	-	4	3	5	6	3	2	-
2002.....	12	-	-	1	3	3	4	1	-
2001.....	23	2	4	4	3	3	7	-	-
2000.....	25	-	-	4	8	4	6	3	-
1999.....	19	-	1	4	2	7	3	2	-
All races, male												
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....
All races, female												
2004.....	14	-	-	5	3	3	2	1	-
2003.....	23	-	4	3	5	6	3	2	-
2002.....	12	-	-	1	3	3	4	1	-
2001.....	23	2	4	4	3	3	7	-	-
2000.....	25	-	-	4	8	4	6	3	-
1999.....	19	-	1	4	2	7	3	2	-
White, both sexes												
2004.....	5	-	-	2	1	2	-	-	-
2003.....	12	-	-	2	5	2	2	1	-
2002.....	3	-	-	-	1	1	-	1	-
2001.....	11	1	3	2	2	1	2	-	-
2000.....	13	-	-	1	5	1	4	2	-
1999.....	8	-	1	1	-	3	1	2	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Ectopic pregnancy (000)												
Black, female												
2004.....	9	-	-	3	2	1	2	1	-
2003.....	11	-	4	1	-	4	1	1	-
2002.....	8	-	-	-	2	2	4	-	-
2001.....	10	1	1	2	1	1	4	-	-
2000.....	12	-	-	3	3	3	2	1	-
1999.....	10	-	-	3	2	3	2	-	-
American Indian 1/, both sexes												
2004.....	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-
2001.....	1	-	-	-	-	1	-	-	-
2000.....	-	-	-	-	-	-	-	-	-
1999.....	1	-	-	-	-	1	-	-	-
American Indian 1/, male												
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....
American Indian 1/, female												
2004.....	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-
2001.....	1	-	-	-	-	1	-	-	-
2000.....	-	-	-	-	-	-	-	-	-
1999.....	1	-	-	-	-	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Spontaneous abortion (003)												
All races, both sexes												
2004.....	5	-	-	2	2	-	-	-	-
2003.....	6	-	-	-	1	3	2	-	-
2002.....	2	1	-	1	-	-	-	-	-
2001.....	7	-	-	-	2	3	1	1	-
2000.....	4	-	-	-	1	-	3	-	-
1999.....	4	-	-	-	1	2	1	-	-
All races, male												
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....
All races, female												
2004.....	5	-	-	2	2	-	-	-	-
2003.....	6	-	-	-	1	3	2	-	-
2002.....	2	1	-	1	-	-	-	-	-
2001.....	7	-	-	-	2	3	1	1	-
2000.....	4	-	-	-	1	-	3	-	-
1999.....	4	-	-	-	1	2	1	-	-
White, both sexes												
2004.....	4	-	-	1	2	-	-	-	-
2003.....	2	-	-	-	-	2	-	-	-
2002.....	1	-	-	1	-	-	-	-	-
2001.....	5	-	-	-	1	2	1	1	-
2000.....	3	-	-	-	1	-	2	-	-
1999.....	2	-	-	-	-	1	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Medical abortion (004)

All races, both sexes

2004.....	2	-	-	-	-	1	1	-	-
2003.....	3	-	2	-	-	-	1	-	-
2002.....	1	-	-	1	-	-	-	-	-
2001.....	1	-	-	1	-	-	-	-	-
2000.....	3	-	-	1	1	-	-	1	-
1999.....	1	-	-	-	1	-	-	-	-

All races, male

2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

All races, female

2004.....	2	-	-	-	-	1	1	-	-
2003.....	3	-	2	-	-	-	1	-	-
2002.....	1	-	-	1	-	-	-	-	-
2001.....	1	-	-	1	-	-	-	-	-
2000.....	3	-	-	1	1	-	-	1	-
1999.....	1	-	-	-	1	-	-	-	-

White, both sexes

2004.....	-	-	-	-	-	-	-	-	-
2003.....	2	-	1	-	-	-	1	-	-
2002.....	-	-	-	-	-	-	-	-	-
2001.....	1	-	-	1	-	-	-	-	-
2000.....	2	-	-	-	1	-	-	1	-
1999.....	-	-	-	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other abortion (005)

All races, both sexes

2004.....	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-

All races, male

2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

All races, female

2004.....	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-

White, both sexes

2004.....	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified pregnancy with abortive outcome (001-002,006-007)												
All races, both sexes												
2004.....	11	-	1	6	1	1	1	1	-
2003.....	8	-	-	2	2	2	-	1	1
2002.....	7	-	-	1	1	2	2	-	1
2001.....	7	-	2	2	1	2	-	-	-
2000.....	5	-	-	1	2	1	1	-	-
1999.....	7	-	2	1	1	1	2	-	-
All races, male												
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....
All races, female												
2004.....	11	-	1	6	1	1	1	1	-
2003.....	8	-	-	2	2	2	-	1	1
2002.....	7	-	-	1	1	2	2	-	1
2001.....	7	-	2	2	1	2	-	-	-
2000.....	5	-	-	1	2	1	1	-	-
1999.....	7	-	2	1	1	1	2	-	-
White, both sexes												
2004.....	2	-	-	2	-	-	-	-	-
2003.....	2	-	-	-	2	-	-	-	-
2002.....	3	-	-	-	1	2	-	-	-
2001.....	3	-	-	1	1	1	-	-	-
2000.....	-	-	-	-	-	-	-	-	-
1999.....	5	-	1	1	1	1	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other direct obstetric deaths (O10-O92)												
All races, both sexes												
2004.....	343	-	21	62	80	81	52	27	10
2003.....	338	1	15	55	64	72	74	41	11
2002.....	285	-	23	43	61	73	62	19	4
2001.....	309	1	27	52	74	75	51	25	4
2000.....	300	-	14	57	60	77	63	24	4
1999.....	308	-	28	44	68	70	65	29	4
All races, male												
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....
All races, female												
2004.....	343	-	21	62	80	81	52	27	10
2003.....	338	1	15	55	64	72	74	41	11
2002.....	285	-	23	43	61	73	62	19	4
2001.....	309	1	27	52	74	75	51	25	4
2000.....	300	-	14	57	60	77	63	24	4
1999.....	308	-	28	44	68	70	65	29	4
White, both sexes												
2004.....	188	-	11	26	37	49	35	17	4
2003.....	180	-	11	29	38	31	37	24	8
2002.....	154	-	10	18	32	42	34	16	2
2001.....	179	1	17	30	41	51	25	12	2
2000.....	185	-	8	33	36	47	43	15	2
1999.....	168	-	12	23	36	41	39	14	3

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other direct obstetric deaths (O10-O92)

Black, female

2004.....	138	-	9	32	41	29	13	7	6
2003.....	135	1	4	20	24	36	32	13	3
2002.....	117	-	13	23	26	24	26	3	2
2001.....	114	-	9	22	28	23	19	11	2
2000.....	99	-	4	23	21	24	17	8	2
1999.....	120	-	15	18	29	25	21	12	-

American Indian 1/, both sexes

2004.....	3	-	-	1	-	-	1	1	-
2003.....	5	-	-	1	-	1	2	1	-
2002.....	-	-	-	-	-	-	-	-	-
2001.....	4	-	-	-	3	1	-	-	-
2000.....	5	-	-	1	1	3	-	-	-
1999.....	4	-	1	2	-	-	1	-	-

American Indian 1/, male

2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

American Indian 1/, female

2004.....	3	-	-	1	-	-	1	1	-
2003.....	5	-	-	1	-	1	2	1	-
2002.....	-	-	-	-	-	-	-	-	-
2001.....	4	-	-	-	3	1	-	-	-
2000.....	5	-	-	1	1	3	-	-	-
1999.....	4	-	1	2	-	-	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other direct obstetric deaths (O10-O92)												
Asian or Pacific Islander 2/, both sexes												
2004.....	14	-	1	3	2	3	3	2	-
2003.....	18	-	-	5	2	4	3	3	-
2002.....	14	-	-	2	3	7	2	-	-
2001.....	12	-	1	-	2	-	7	2	-
2000.....	11	-	2	-	2	3	3	1	-
1999.....	16	-	-	1	3	4	4	3	1
Asian or Pacific Islander 2/, male												
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....
Asian or Pacific Islander 2/, female												
2004.....	14	-	1	3	2	3	3	2	-
2003.....	18	-	-	5	2	4	3	3	-
2002.....	14	-	-	2	3	7	2	-	-
2001.....	12	-	1	-	2	-	7	2	-
2000.....	11	-	2	-	2	3	3	1	-
1999.....	16	-	-	1	3	4	4	3	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Eclampsia and pre-eclampsia (O11,O13-O16)												
All races, both sexes												
2004.....	54	-	8	11	12	13	4	5	1
2003.....	68	-	2	13	16	13	18	5	1
2002.....	56	-	6	6	9	19	9	5	2
2001.....	72	1	6	16	17	17	8	7	-
2000.....	72	-	3	11	18	21	14	3	1
1999.....	90	-	9	14	22	17	19	8	1
All races, male												
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....
All races, female												
2004.....	54	-	8	11	12	13	4	5	1
2003.....	68	-	2	13	16	13	18	5	1
2002.....	56	-	6	6	9	19	9	5	2
2001.....	72	1	6	16	17	17	8	7	-
2000.....	72	-	3	11	18	21	14	3	1
1999.....	90	-	9	14	22	17	19	8	1
White, both sexes												
2004.....	27	-	4	5	3	8	2	4	1
2003.....	33	-	2	5	11	5	7	2	1
2002.....	32	-	2	5	4	9	6	5	1
2001.....	43	1	3	10	11	12	3	3	-
2000.....	46	-	2	6	12	13	10	2	-
1999.....	50	-	5	7	11	10	12	4	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Eclampsia and pre-eclampsia (O11,O13-O16)

Black, female

2004.....	25	-	3	6	9	5	2	-	-
2003.....	29	-	-	6	4	8	10	1	-
2002.....	23	-	4	1	4	10	3	-	1
2001.....	26	-	2	6	6	5	4	3	-
2000.....	25	-	1	4	6	8	4	1	1
1999.....	35	-	3	6	10	7	5	4	-

American Indian 1/, both sexes

2004.....	-	-	-	-	-	-	-	-	-
2003.....	2	-	-	1	-	-	-	1	-
2002.....	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-
2000.....	1	-	-	1	-	-	-	-	-
1999.....	2	-	1	1	-	-	-	-	-

American Indian 1/, male

2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

American Indian 1/, female

2004.....	-	-	-	-	-	-	-	-	-
2003.....	2	-	-	1	-	-	-	1	-
2002.....	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-
2000.....	1	-	-	1	-	-	-	-	-
1999.....	2	-	1	1	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Eclampsia and pre-eclampsia (O11,O13-O16)												
Asian or Pacific Islander 2/, both sexes												
2004.....	2	-	1	-	-	-	-	1	-
2003.....	4	-	-	1	1	-	1	1	-
2002.....	1	-	-	-	1	-	-	-	-
2001.....	3	-	1	-	-	-	1	1	-
2000.....	-	-	-	-	-	-	-	-	-
1999.....	3	-	-	-	1	-	2	-	-
Asian or Pacific Islander 2/, male												
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....
Asian or Pacific Islander 2/, female												
2004.....	2	-	1	-	-	-	-	1	-
2003.....	4	-	-	1	1	-	1	1	-
2002.....	1	-	-	-	1	-	-	-	-
2001.....	3	-	1	-	-	-	1	1	-
2000.....	-	-	-	-	-	-	-	-	-
1999.....	3	-	-	-	1	-	2	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hemorrhage of pregnancy and childbirth and placenta previa (O20,O44-O46,O67,072)												
All races, both sexes												
2004.....	38	-	2	5	11	9	7	3	1
2003.....	47	-	1	7	12	9	10	8	-
2002.....	26	-	1	5	7	7	5	-	1
2001.....	44	-	2	5	4	14	11	7	1
2000.....	30	-	-	3	7	7	9	4	-
1999.....	45	-	4	5	8	11	13	4	-
All races, male												
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....
All races, female												
2004.....	38	-	2	5	11	9	7	3	1
2003.....	47	-	1	7	12	9	10	8	-
2002.....	26	-	1	5	7	7	5	-	1
2001.....	44	-	2	5	4	14	11	7	1
2000.....	30	-	-	3	7	7	9	4	-
1999.....	45	-	4	5	8	11	13	4	-
White, both sexes												
2004.....	28	-	1	3	8	7	5	3	1
2003.....	28	-	1	5	8	4	6	4	-
2002.....	16	-	1	3	5	4	2	-	1
2001.....	26	-	1	4	2	10	6	3	-
2000.....	21	-	-	2	4	5	8	2	-
1999.....	27	-	-	3	5	9	7	3	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hemorrhage of pregnancy and childbirth and placenta previa (O20,O44-O46,O67,O72)												
Black, female												
2004.....	10	-	1	2	3	2	2	-	-
2003.....	16	-	-	1	3	5	3	4	-
2002.....	9	-	-	2	2	2	3	-	-
2001.....	15	-	1	1	2	3	4	3	1
2000.....	6	-	-	1	2	1	1	1	-
1999.....	13	-	4	1	2	-	5	1	-
American Indian 1/, both sexes												
2004.....	-	-	-	-	-	-	-	-	-
2003.....	1	-	-	-	-	-	1	-	-
2002.....	-	-	-	-	-	-	-	-	-
2001.....	1	-	-	-	-	1	-	-	-
2000.....	2	-	-	-	1	1	-	-	-
1999.....	1	-	-	1	-	-	-	-	-
American Indian 1/, male												
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....
American Indian 1/, female												
2004.....	-	-	-	-	-	-	-	-	-
2003.....	1	-	-	-	-	-	1	-	-
2002.....	-	-	-	-	-	-	-	-	-
2001.....	1	-	-	-	-	1	-	-	-
2000.....	2	-	-	-	1	1	-	-	-
1999.....	1	-	-	1	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Hemorrhage of pregnancy and childbirth and placenta previa (O20,044-046,067,072)

Asian or Pacific Islander 2/, both sexes

2004.....	-	-	-	-	-	-	-	-	-
2003.....	2	-	-	1	1	-	-	-	-
2002.....	1	-	-	-	-	1	-	-	-
2001.....	2	-	-	-	-	-	1	1	-
2000.....	1	-	-	-	-	-	-	1	-
1999.....	4	-	-	-	1	2	1	-	-

Asian or Pacific Islander 2/, male

2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

Asian or Pacific Islander 2/, female

2004.....	-	-	-	-	-	-	-	-	-
2003.....	2	-	-	1	1	-	-	-	-
2002.....	1	-	-	-	-	1	-	-	-
2001.....	2	-	-	-	-	-	1	1	-
2000.....	1	-	-	-	-	-	-	1	-
1999.....	4	-	-	-	1	2	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Complications predominately related to the puerperium (085-092)

All races, both sexes

2004.....	102	-	5	18	23	24	20	8	3
2003.....	88	-	7	12	20	17	20	8	4
2002.....	82	-	6	12	17	20	18	8	1
2001.....	101	-	13	14	28	20	19	6	1
2000.....	93	-	6	21	16	26	15	7	2
1999.....	84	-	6	12	18	23	16	6	3

All races, male

2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

All races, female

2004.....	102	-	5	18	23	24	20	8	3
2003.....	88	-	7	12	20	17	20	8	4
2002.....	82	-	6	12	17	20	18	8	1
2001.....	101	-	13	14	28	20	19	6	1
2000.....	93	-	6	21	16	26	15	7	2
1999.....	84	-	6	12	18	23	16	6	3

White, both sexes

2004.....	53	-	2	7	11	15	14	3	-
2003.....	47	-	4	7	10	8	10	5	3
2002.....	39	-	3	3	7	12	7	7	-
2001.....	55	-	7	7	14	12	11	3	1
2000.....	55	-	4	12	7	16	9	5	2
1999.....	47	-	4	8	9	12	9	3	2

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Complications predominately related to the puerperium (085-092)												
Black, female												
2004.....	45	-	3	10	11	9	5	4	3
2003.....	34	-	3	4	10	5	8	3	1
2002.....	38	-	3	9	9	5	10	1	1
2001.....	39	-	6	7	10	8	5	3	-
2000.....	31	-	1	9	7	7	5	2	-
1999.....	32	-	2	4	8	10	5	3	-
American Indian 1/, both sexes												
2004.....	-	-	-	-	-	-	-	-	-
2003.....	2	-	-	-	-	1	1	-	-
2002.....	-	-	-	-	-	-	-	-	-
2001.....	2	-	-	-	2	-	-	-	-
2000.....	2	-	-	-	-	2	-	-	-
1999.....	1	-	-	-	-	-	1	-	-
American Indian 1/, male												
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....
American Indian 1/, female												
2004.....	-	-	-	-	-	-	-	-	-
2003.....	2	-	-	-	-	1	1	-	-
2002.....	-	-	-	-	-	-	-	-	-
2001.....	2	-	-	-	2	-	-	-	-
2000.....	2	-	-	-	-	2	-	-	-
1999.....	1	-	-	-	-	-	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Complications predominately related to the puerperium (O85-O92)

Asian or Pacific Islander 2/, both sexes

2004.....	4	-	-	1	1	-	1	1	-
2003.....	5	-	-	1	-	3	1	-	-
2002.....	5	-	-	-	1	3	1	-	-
2001.....	5	-	-	-	2	-	3	-	-
2000.....	5	-	1	-	2	1	1	-	-
1999.....	4	-	-	-	1	1	1	-	1

Asian or Pacific Islander 2/, male

2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

Asian or Pacific Islander 2/, female

2004.....	4	-	-	1	1	-	1	1	-
2003.....	5	-	-	1	-	3	1	-	-
2002.....	5	-	-	-	1	3	1	-	-
2001.....	5	-	-	-	2	-	3	-	-
2000.....	5	-	1	-	2	1	1	-	-
1999.....	4	-	-	-	1	1	1	-	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Obstetric embolism (O88)

All races, both sexes

2004.....	48	-	1	5	12	14	13	2	1
2003.....	49	-	4	6	10	12	11	6	-
2002.....	47	-	4	7	10	11	11	4	-
2001.....	69	-	10	9	17	11	18	4	-
2000.....	58	-	2	13	8	20	10	4	1
1999.....	53	-	2	9	10	15	13	2	2

All races, male

2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

All races, female

2004.....	48	-	1	5	12	14	13	2	1
2003.....	49	-	4	6	10	12	11	6	-
2002.....	47	-	4	7	10	11	11	4	-
2001.....	69	-	10	9	17	11	18	4	-
2000.....	58	-	2	13	8	20	10	4	1
1999.....	53	-	2	9	10	15	13	2	2

White, both sexes

2004.....	27	-	1	2	6	9	9	-	-
2003.....	26	-	2	4	5	6	6	3	-
2002.....	26	-	3	2	6	7	4	4	-
2001.....	39	-	6	3	8	9	11	2	-
2000.....	38	-	1	8	5	13	6	4	1
1999.....	32	-	1	7	6	10	6	1	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Obstetric embolism (O88)

Black, female

2004.....	19	-	-	3	5	5	4	1	1
2003.....	18	-	2	2	5	3	3	3	-
2002.....	16	-	1	5	3	1	6	-	-
2001.....	24	-	4	6	6	2	4	2	-
2000.....	15	-	-	5	2	5	3	-	-
1999.....	16	-	1	2	3	4	5	1	-

American Indian 1/, both sexes

2004.....	-	-	-	-	-	-	-	-	-
2003.....	1	-	-	-	-	-	1	-	-
2002.....	-	-	-	-	-	-	-	-	-
2001.....	1	-	-	-	1	-	-	-	-
2000.....	1	-	-	-	-	1	-	-	-
1999.....	1	-	-	-	-	-	1	-	-

American Indian 1/, male

2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

American Indian 1/, female

2004.....	-	-	-	-	-	-	-	-	-
2003.....	1	-	-	-	-	-	1	-	-
2002.....	-	-	-	-	-	-	-	-	-
2001.....	1	-	-	-	1	-	-	-	-
2000.....	1	-	-	-	-	1	-	-	-
1999.....	1	-	-	-	-	-	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Obstetric embolism (O88)												
Asian or Pacific Islander 2/, both sexes												
2004.....	2	-	-	-	1	-	-	1	-
2003.....	4	-	-	-	-	3	1	-	-
2002.....	5	-	-	-	1	3	1	-	-
2001.....	5	-	-	-	2	-	3	-	-
2000.....	4	-	1	-	1	1	1	-	-
1999.....	4	-	-	-	1	1	1	-	1
Asian or Pacific Islander 2/, male												
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....
Asian or Pacific Islander 2/, female												
2004.....	2	-	-	-	1	-	-	1	-
2003.....	4	-	-	-	-	3	1	-	-
2002.....	5	-	-	-	1	3	1	-	-
2001.....	5	-	-	-	2	-	3	-	-
2000.....	4	-	1	-	1	1	1	-	-
1999.....	4	-	-	-	1	1	1	-	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other complications predominately related to the puerperium (O85-O87,O89-O92)												
All races, both sexes												
2004.....	54	-	4	13	11	10	7	6	2
2003.....	39	-	3	6	10	5	9	2	4
2002.....	35	-	2	5	7	9	7	4	1
2001.....	32	-	3	5	11	9	1	2	1
2000.....	35	-	4	8	8	6	5	3	1
1999.....	31	-	4	3	8	8	3	4	1
All races, male												
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....
All races, female												
2004.....	54	-	4	13	11	10	7	6	2
2003.....	39	-	3	6	10	5	9	2	4
2002.....	35	-	2	5	7	9	7	4	1
2001.....	32	-	3	5	11	9	1	2	1
2000.....	35	-	4	8	8	6	5	3	1
1999.....	31	-	4	3	8	8	3	4	1
White, both sexes												
2004.....	26	-	1	5	5	6	5	3	-
2003.....	21	-	2	3	5	2	4	2	3
2002.....	13	-	-	1	1	5	3	3	-
2001.....	16	-	1	4	6	3	-	1	1
2000.....	17	-	3	4	2	3	3	1	1
1999.....	15	-	3	1	3	2	3	2	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other direct obstetric causes (O10,O12,O21-O43,O47-O66,O68-O71,O73-O75)												
All races, both sexes												
2004.....	149	-	6	28	34	35	21	11	5
2003.....	135	1	5	23	16	33	26	20	6
2002.....	121	-	10	20	28	27	30	6	-
2001.....	92	-	6	17	25	24	13	5	2
2000.....	105	-	5	22	19	23	25	10	1
1999.....	89	-	9	13	20	19	17	11	-
All races, male												
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....
All races, female												
2004.....	149	-	6	28	34	35	21	11	5
2003.....	135	1	5	23	16	33	26	20	6
2002.....	121	-	10	20	28	27	30	6	-
2001.....	92	-	6	17	25	24	13	5	2
2000.....	105	-	5	22	19	23	25	10	1
1999.....	89	-	9	13	20	19	17	11	-
White, both sexes												
2004.....	80	-	4	11	15	19	14	7	2
2003.....	72	-	4	12	9	14	14	13	4
2002.....	67	-	4	7	16	17	19	4	-
2001.....	55	-	6	9	14	17	5	3	1
2000.....	63	-	2	13	13	13	16	6	-
1999.....	44	-	3	5	11	10	11	4	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

All other direct obstetric causes (O10,O12,O21-O43,O47-O66,O68-O71,O73-O75)

Black, female

2004.....	58	-	2	14	18	13	4	3	3
2003.....	56	1	1	9	7	18	11	5	2
2002.....	47	-	6	11	11	7	10	2	-
2001.....	34	-	-	8	10	7	6	2	1
2000.....	37	-	2	9	6	8	7	4	1
1999.....	40	-	6	7	9	8	6	4	-

American Indian 1/, both sexes

2004.....	3	-	-	1	-	-	1	1	-
2003.....	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-
2001.....	1	-	-	-	1	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-

American Indian 1/, male

2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

American Indian 1/, female

2004.....	3	-	-	1	-	-	1	1	-
2003.....	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-
2001.....	1	-	-	-	1	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other direct obstetric causes (O10,O12,O21-O43,O47-O66,O68-O71,O73-O75)												
Asian or Pacific Islander 2/, both sexes												
2004.....	8	-	-	2	1	3	2	-	-
2003.....	7	-	-	2	-	1	1	2	-
2002.....	7	-	-	2	1	3	1	-	-
2001.....	2	-	-	-	-	-	2	-	-
2000.....	5	-	1	-	-	2	2	-	-
1999.....	5	-	-	1	-	1	-	3	-
Asian or Pacific Islander 2/, male												
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....
Asian or Pacific Islander 2/, female												
2004.....	8	-	-	2	1	3	2	-	-
2003.....	7	-	-	2	-	1	1	2	-
2002.....	7	-	-	2	1	3	1	-	-
2001.....	2	-	-	-	-	-	2	-	-
2000.....	5	-	1	-	-	2	2	-	-
1999.....	5	-	-	1	-	1	-	3	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Obstetric death of unspecified cause (O95)

All races, both sexes

2004.....	32	-	-	9	10	6	4	2	1
2003.....	22	-	-	3	4	5	3	4	2
2002.....	4	-	-	1	1	2	-	-	-
2001.....	1	-	-	1	-	-	-	-	-
2000.....	4	-	-	1	1	-	1	1	-
1999.....	5	-	-	-	2	-	3	-	-

All races, male

2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

All races, female

2004.....	32	-	-	9	10	6	4	2	1
2003.....	22	-	-	3	4	5	3	4	2
2002.....	4	-	-	1	1	2	-	-	-
2001.....	1	-	-	1	-	-	-	-	-
2000.....	4	-	-	1	1	-	1	1	-
1999.....	5	-	-	-	2	-	3	-	-

White, both sexes

2004.....	19	-	-	6	3	5	2	2	1
2003.....	17	-	-	2	3	5	1	3	2
2002.....	1	-	-	-	-	1	-	-	-
2001.....	1	-	-	1	-	-	-	-	-
2000.....	3	-	-	1	-	-	1	1	-
1999.....	3	-	-	-	2	-	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other deaths related to pregnancy, childbirth and the puerperium (096-097)

All races, both sexes

2004.....	157	-	11	19	21	31	14	19	19
2003.....	50	-	2	10	16	13	4	2	1
2002.....	22	-	-	2	3	5	6	3	2
2001.....	17	-	-	3	3	2	6	1	-
2000.....	8	-	-	2	1	2	-	2	1
1999.....	15	-	1	3	1	2	5	1	-

All races, male

2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

All races, female

2004.....	157	-	11	19	21	31	14	19	19
2003.....	50	-	2	10	16	13	4	2	1
2002.....	22	-	-	2	3	5	6	3	2
2001.....	17	-	-	3	3	2	6	1	-
2000.....	8	-	-	2	1	2	-	2	1
1999.....	15	-	1	3	1	2	5	1	-

White, both sexes

2004.....	111	-	10	13	15	22	11	13	11
2003.....	37	-	2	9	13	5	3	2	1
2002.....	13	-	-	1	2	4	3	-	2
2001.....	7	-	-	1	1	1	1	1	-
2000.....	3	-	-	1	1	1	-	-	-
1999.....	6	-	-	1	-	1	4	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other deaths related to pregnancy, childbirth and the puerperium (096-097)

Black, female

2004.....	37	-	1	6	5	7	3	4	6
2003.....	10	-	-	1	3	6	-	-	-
2002.....	9	-	-	1	1	1	3	3	-
2001.....	8	-	-	2	2	1	3	-	-
2000.....	5	-	-	1	-	1	-	2	1
1999.....	8	-	1	2	1	-	1	1	-

American Indian 1/, both sexes

2004.....	2	-	-	-	1	-	-	-	1
2003.....	1	-	-	-	-	-	1	-	-
2002.....	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-

American Indian 1/, male

2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

American Indian 1/, female

2004.....	2	-	-	-	1	-	-	-	1
2003.....	1	-	-	-	-	-	1	-	-
2002.....	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other deaths related to pregnancy, childbirth and the puerperium (096-097)

Asian or Pacific Islander 2/, both sexes

2004.....	7	-	-	-	-	2	-	2	1
2003.....	2	-	-	-	-	2	-	-	-
2002.....	-	-	-	-	-	-	-	-	-
2001.....	2	-	-	-	-	-	2	-	-
2000.....	-	-	-	-	-	-	-	-	-
1999.....	1	-	-	-	-	1	-	-	-

Asian or Pacific Islander 2/, male

2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

Asian or Pacific Islander 2/, female

2004.....	7	-	-	-	-	2	-	2	1
2003.....	2	-	-	-	-	2	-	-	-
2002.....	-	-	-	-	-	-	-	-	-
2001.....	2	-	-	-	-	-	2	-	-
2000.....	-	-	-	-	-	-	-	-	-
1999.....	1	-	-	-	-	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Indirect obstetric deaths (O98-O99)												
All races, both sexes												
2004.....	133	2	4	28	25	22	22	11	9
2003.....	95	-	4	16	18	18	19	10	4
2002.....	46	-	5	11	13	8	5	3	1
2001.....	51	-	4	10	10	12	8	6	1
2000.....	55	-	4	11	13	11	12	4	-
1999.....	47	-	1	12	13	10	9	1	1
All races, male												
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....
All races, female												
2004.....	133	2	4	28	25	22	22	11	9
2003.....	95	-	4	16	18	18	19	10	4
2002.....	46	-	5	11	13	8	5	3	1
2001.....	51	-	4	10	10	12	8	6	1
2000.....	55	-	4	11	13	11	12	4	-
1999.....	47	-	1	12	13	10	9	1	1
White, both sexes												
2004.....	82	-	3	14	18	14	12	7	5
2003.....	65	-	1	9	12	14	12	8	3
2002.....	28	-	2	8	5	6	4	2	1
2001.....	28	-	2	6	4	8	4	4	-
2000.....	34	-	3	8	9	6	6	2	-
1999.....	28	-	-	5	8	6	8	-	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Indirect obstetric deaths (O98-O99)

Black, female

2004.....	45	-	1	14	7	7	8	3	4
2003.....	26	-	3	7	4	4	5	2	1
2002.....	16	-	3	3	8	-	1	1	-
2001.....	20	-	2	4	4	3	4	2	1
2000.....	18	-	-	3	4	5	5	1	-
1999.....	17	-	1	6	4	4	1	1	-

American Indian 1/, both sexes

2004.....	1	-	-	-	-	-	-	1	-
2003.....	1	-	-	-	1	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-
2000.....	1	-	1	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-

American Indian 1/, male

2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

American Indian 1/, female

2004.....	1	-	-	-	-	-	-	1	-
2003.....	1	-	-	-	1	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-
2000.....	1	-	1	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Indirect obstetric deaths (098-099)												
Asian or Pacific Islander 2/, both sexes												
2004.....	5	2	-	-	-	1	2	-	-
2003.....	3	-	-	-	1	-	2	-	-
2002.....	2	-	-	-	-	2	-	-	-
2001.....	3	-	-	-	2	1	-	-	-
2000.....	2	-	-	-	-	-	1	1	-
1999.....	2	-	-	1	1	-	-	-	-
Asian or Pacific Islander 2/, male												
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....
Asian or Pacific Islander 2/, female												
2004.....	5	2	-	-	-	1	2	-	-
2003.....	3	-	-	-	1	-	2	-	-
2002.....	2	-	-	-	-	2	-	-	-
2001.....	3	-	-	-	2	1	-	-	-
2000.....	2	-	-	-	-	-	1	1	-
1999.....	2	-	-	1	1	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

XVI. Certain conditions originating in the perinatal period (P00-P96)

All races, both sexes

2004.....	14,213	14,079	61	25	12	4	7	6	4	3	3	2
2003.....	14,378	14,254	79	10	9	12	4	2	1	1	2	1
2002.....	14,254	14,106	65	15	16	14	7	6	4	7	3	2
2001.....	13,887	13,734	72	22	14	8	9	5	1	6	3	3
2000.....	14,069	13,918	79	25	16	10	5	5	2	2	2	1
1999.....	14,259	14,084	92	25	18	11	5	2	5	2	1	2

All races, male

2004.....	8,093	8,013	38	15	8	3	2	1	2	3	2	1
2003.....	8,197	8,129	43	5	6	6	2	1	1	1	-	-
2002.....	8,047	7,960	40	10	9	9	3	3	1	4	1	2
2001.....	7,868	7,775	44	16	8	3	4	2	-	4	2	3
2000.....	7,968	7,872	52	13	10	9	2	4	2	-	2	1
1999.....	8,044	7,941	56	14	9	6	4	-	2	2	-	2

All races, female

2004.....	6,120	6,066	23	10	4	1	5	5	2	-	1	1
2003.....	6,181	6,125	36	5	3	6	2	1	-	-	2	1
2002.....	6,207	6,146	25	5	7	5	4	3	3	3	2	-
2001.....	6,019	5,959	28	6	6	5	5	3	1	2	1	-
2000.....	6,101	6,046	27	12	6	1	3	1	-	2	-	-
1999.....	6,215	6,143	36	11	9	5	1	2	3	-	1	-

White, both sexes

2004.....	8,732	8,643	33	18	8	4	6	4	4	3	1	2
2003.....	8,975	8,891	45	9	7	12	3	2	1	1	2	1
2002.....	8,777	8,679	36	9	12	13	4	5	2	5	3	2
2001.....	8,551	8,444	47	14	7	5	8	5	1	6	2	3
2000.....	8,532	8,431	46	18	12	8	5	2	1	2	2	1
1999.....	8,648	8,522	56	19	14	11	4	2	5	2	1	2

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

XVI. Certain conditions originating in the perinatal period (P00-P96)

White, male

2004.....	5,025	4,971	22	11	5	3	2	1	2	3	-	1
2003.....	5,134	5,087	25	5	5	6	2	1	1	1	-	-
2002.....	5,011	4,959	19	7	5	8	1	3	1	2	1	2
2001.....	4,846	4,780	28	11	5	2	4	2	-	4	1	3
2000.....	4,828	4,762	30	10	10	8	2	1	1	-	2	1
1999.....	4,884	4,815	30	10	7	6	4	-	2	2	-	2

White, female

2004.....	3,707	3,672	11	7	3	1	4	3	2	-	1	1
2003.....	3,841	3,804	20	4	2	6	1	1	-	-	2	1
2002.....	3,766	3,720	17	2	7	5	3	2	1	3	2	-
2001.....	3,705	3,664	19	3	2	3	4	3	1	2	1	-
2000.....	3,704	3,669	16	8	2	-	3	1	-	2	-	-
1999.....	3,764	3,707	26	9	7	5	-	2	3	-	1	-

Black, both sexes

2004.....	4,920	4,883	23	6	3	-	1	2	-	-	2	-
2003.....	4,879	4,844	30	-	2	-	1	-	-	-	-	-
2002.....	4,931	4,889	26	4	4	1	2	1	1	1	-	-
2001.....	4,847	4,808	23	6	5	3	1	-	-	-	1	-
2000.....	5,021	4,979	27	6	4	1	-	3	1	-	-	-
1999.....	5,150	5,113	29	2	3	-	1	-	-	-	-	-

Black, male

2004.....	2,753	2,733	13	3	2	-	-	-	-	-	2	-
2003.....	2,775	2,756	16	-	1	-	-	-	-	-	-	-
2002.....	2,725	2,696	18	2	4	1	1	-	-	1	-	-
2001.....	2,746	2,725	14	3	2	1	-	-	-	-	1	-
2000.....	2,844	2,818	18	3	-	1	-	3	1	-	-	-
1999.....	2,907	2,883	19	2	1	-	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Newborn affected by maternal factors and by complications of pregnancy, labor and delivery (P00-P04)

All races, both sexes

2004.....	3,117	3,108	3	4	1	-	-	-	-	-	1	-
2003.....	3,183	3,166	10	3	2	2	-	-	-	-	-	-
2002.....	3,073	3,063	5	-	3	-	-	-	-	-	-	-
2001.....	2,845	2,832	7	1	2	-	2	-	-	-	-	1
2000.....	2,781	2,768	9	2	-	-	1	-	-	-	-	-
1999.....	2,750	2,738	4	3	2	1	1	-	-	-	-	-

All races, male

2004.....	1,714	1,708	2	2	1	-	-	-	-	-	1	-
2003.....	1,761	1,752	5	2	1	1	-	-	-	-	-	-
2002.....	1,726	1,718	4	-	2	-	-	-	-	-	-	-
2001.....	1,613	1,606	4	1	1	-	-	-	-	-	-	1
2000.....	1,542	1,533	7	1	-	-	1	-	-	-	-	-
1999.....	1,498	1,490	3	3	1	-	-	-	-	-	-	-

All races, female

2004.....	1,403	1,400	1	2	-	-	-	-	-	-	-	-
2003.....	1,422	1,414	5	1	1	1	-	-	-	-	-	-
2002.....	1,347	1,345	1	-	1	-	-	-	-	-	-	-
2001.....	1,232	1,226	3	-	1	-	2	-	-	-	-	-
2000.....	1,239	1,235	2	1	-	-	-	-	-	-	-	-
1999.....	1,252	1,248	1	-	1	1	1	-	-	-	-	-

White, both sexes

2004.....	1,911	1,906	2	3	-	-	-	-	-	-	-	-
2003.....	2,055	2,044	6	2	1	2	-	-	-	-	-	-
2002.....	1,917	1,912	2	-	2	-	-	-	-	-	-	-
2001.....	1,815	1,806	5	-	1	-	2	-	-	-	-	1
2000.....	1,773	1,766	4	1	-	-	1	-	-	-	-	-
1999.....	1,693	1,684	3	2	2	1	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Birth trauma (P10-P15)												
All races, both sexes												
2004.....	28	22	-	1	-	1	-	-	1	-	-	1
2003.....	32	26	2	-	-	2	-	-	1	-	-	-
2002.....	352	345	2	2	2	-	-	1	-	-	-	-
2001.....	158	145	2	3	1	-	1	-	-	2	1	1
2000.....	44	31	3	1	4	2	1	-	1	-	-	-
1999.....	30	20	4	-	1	1	-	-	-	1	-	-
All races, male												
2004.....	17	12	-	1	-	1	-	-	-	-	-	1
2003.....	17	12	1	-	-	2	-	-	1	-	-	-
2002.....	223	218	2	1	1	-	-	1	-	-	-	-
2001.....	98	86	2	2	1	-	1	-	-	2	1	1
2000.....	20	13	1	1	2	2	-	-	1	-	-	-
1999.....	19	14	2	-	-	1	-	-	-	1	-	-
All races, female												
2004.....	11	10	-	-	-	-	-	-	1	-	-	-
2003.....	15	14	1	-	-	-	-	-	-	-	-	-
2002.....	129	127	-	1	1	-	-	-	-	-	-	-
2001.....	60	59	-	1	-	-	-	-	-	-	-	-
2000.....	24	18	2	-	2	-	1	-	-	-	-	-
1999.....	11	6	2	-	1	-	-	-	-	-	-	-
White, both sexes												
2004.....	23	17	-	1	-	1	-	-	1	-	-	1
2003.....	23	19	1	-	-	2	-	-	1	-	-	-
2002.....	251	245	2	1	2	-	-	1	-	-	-	-
2001.....	107	95	2	3	1	-	1	-	-	2	-	1
2000.....	28	19	2	1	2	2	1	-	-	-	-	-
1999.....	19	11	2	-	1	1	-	-	-	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intrauterine hypoxia and birth asphyxia (P20-P21)												
All races, both sexes												
2004.....	569	530	13	9	5	1	3	2	1	1	2	-
2003.....	581	558	14	4	-	1	-	1	-	-	-	1
2002.....	619	583	14	2	4	4	4	2	3	1	1	1
2001.....	572	534	18	8	3	2	2	2	-	1	-	-
2000.....	660	630	11	11	4	1	2	-	-	-	1	-
1999.....	654	613	20	8	3	4	4	-	-	-	-	-
All races, male												
2004.....	328	305	7	7	3	1	1	-	1	1	1	-
2003.....	316	300	10	2	-	1	-	1	-	-	-	-
2002.....	316	299	6	2	3	1	2	1	1	-	-	1
2001.....	320	295	12	5	1	1	1	2	-	1	-	-
2000.....	369	352	6	6	2	1	1	-	-	-	1	-
1999.....	360	335	11	4	2	2	4	-	-	-	-	-
All races, female												
2004.....	241	225	6	2	2	-	2	2	-	-	1	-
2003.....	265	258	4	2	-	-	-	-	-	-	-	1
2002.....	303	284	8	-	1	3	2	1	2	1	1	-
2001.....	252	239	6	3	2	1	1	-	-	-	-	-
2000.....	291	278	5	5	2	-	1	-	-	-	-	-
1999.....	294	278	9	4	1	2	-	-	-	-	-	-
White, both sexes												
2004.....	405	376	7	8	4	1	3	1	1	1	1	-
2003.....	409	391	10	4	-	1	-	1	-	-	-	1
2002.....	438	413	7	2	3	4	2	2	2	1	1	1
2001.....	410	385	12	6	-	1	2	2	-	1	-	-
2000.....	459	435	7	10	3	1	2	-	-	-	1	-
1999.....	462	429	14	6	3	4	4	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other respiratory conditions originating in the perinatal period (P23-P28)

All races, both sexes

2004.....	1,257	1,217	29	4	1	1	2	1	1	-	-	-
2003.....	1,250	1,217	29	1	1	1	1	-	-	-	-	-
2002.....	1,278	1,245	22	5	-	2	-	1	-	1	-	-
2001.....	1,536	1,503	26	3	-	-	2	1	-	1	-	-
2000.....	1,573	1,529	39	2	1	2	-	-	-	-	-	-
1999.....	1,768	1,718	39	5	5	1	-	-	-	-	-	-

All races, male

2004.....	732	708	17	2	1	1	-	1	1	-	-	-
2003.....	759	739	17	-	1	1	1	-	-	-	-	-
2002.....	752	732	13	3	-	2	-	1	-	-	-	-
2001.....	882	869	11	1	-	-	1	-	-	-	-	-
2000.....	910	881	26	-	1	2	-	-	-	-	-	-
1999.....	1,035	1,000	27	4	4	-	-	-	-	-	-	-

All races, female

2004.....	525	509	12	2	-	-	2	-	-	-	-	-
2003.....	491	478	12	1	-	-	-	-	-	-	-	-
2002.....	526	513	9	2	-	-	-	-	-	1	-	-
2001.....	654	634	15	2	-	-	1	1	-	1	-	-
2000.....	663	648	13	2	-	-	-	-	-	-	-	-
1999.....	733	718	12	1	1	1	-	-	-	-	-	-

White, both sexes

2004.....	795	772	17	1	1	1	1	1	1	-	-	-
2003.....	819	802	13	1	1	1	1	-	-	-	-	-
2002.....	812	798	9	1	-	1	-	1	-	1	-	-
2001.....	960	941	15	1	-	-	1	1	-	1	-	-
2000.....	984	961	21	-	1	1	-	-	-	-	-	-
1999.....	1,176	1,151	18	3	3	1	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Infections specific to the perinatal period (P35-P39)

All races, both sexes

2004.....	1,055	1,038	3	3	2	1	1	2	1	2	-	1
2003.....	982	967	4	-	2	4	1	1	-	1	2	-
2002.....	975	948	7	1	5	4	3	2	-	3	1	-
2001.....	887	869	2	6	3	3	1	-	1	-	-	1
2000.....	944	927	2	4	3	4	1	2	-	1	-	-
1999.....	900	871	8	6	5	2	-	2	3	-	1	1

All races, male

2004.....	611	603	1	2	1	-	1	-	-	2	-	-
2003.....	579	576	1	-	-	-	1	-	-	1	-	-
2002.....	532	518	5	1	2	2	1	-	-	2	-	-
2001.....	518	505	2	6	1	2	1	-	-	-	-	1
2000.....	548	538	1	3	2	3	-	1	-	-	-	-
1999.....	505	495	4	1	1	2	-	-	1	-	-	1

All races, female

2004.....	444	435	2	1	1	1	-	2	1	-	-	1
2003.....	403	391	3	-	2	4	-	1	-	-	2	-
2002.....	443	430	2	-	3	2	2	2	-	1	1	-
2001.....	369	364	-	-	2	1	-	-	1	-	-	-
2000.....	396	389	1	1	1	1	1	1	-	1	-	-
1999.....	395	376	4	5	4	-	-	2	2	-	1	-

White, both sexes

2004.....	664	650	2	1	2	1	1	2	1	2	-	1
2003.....	629	615	3	-	2	4	1	1	-	1	2	-
2002.....	643	621	7	1	3	4	2	1	-	2	1	-
2001.....	564	552	2	3	1	2	1	-	1	-	-	1
2000.....	588	576	1	3	2	3	1	1	-	1	-	-
1999.....	559	537	4	5	3	2	-	2	3	-	1	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Infections specific to the perinatal period (P35-P39)												
White, male												
2004.....	378	372	-	1	1	-	1	-	-	2	-	-
2003.....	371	368	1	-	-	-	1	-	-	1	-	-
2002.....	361	351	5	1	-	2	-	-	-	1	-	-
2001.....	344	336	2	3	-	1	1	-	-	-	-	1
2000.....	348	340	-	3	2	3	-	-	-	-	-	-
1999.....	302	296	1	1	-	2	-	-	1	-	-	1
White, female												
2004.....	286	278	2	-	1	1	-	2	1	-	-	1
2003.....	258	247	2	-	2	4	-	1	-	-	2	-
2002.....	282	270	2	-	3	2	2	1	-	1	1	-
2001.....	220	216	-	-	1	1	-	-	1	-	-	-
2000.....	240	236	1	-	-	-	1	1	-	1	-	-
1999.....	257	241	3	4	3	-	-	2	2	-	1	-
Black, both sexes												
2004.....	350	347	1	2	-	-	-	-	-	-	-	-
2003.....	326	325	1	-	-	-	-	-	-	-	-	-
2002.....	308	304	-	-	2	-	1	1	-	-	-	-
2001.....	283	280	-	1	1	1	-	-	-	-	-	-
2000.....	321	318	1	-	1	-	-	1	-	-	-	-
1999.....	313	308	3	-	2	-	-	-	-	-	-	-
Black, male												
2004.....	204	202	1	1	-	-	-	-	-	-	-	-
2003.....	190	190	-	-	-	-	-	-	-	-	-	-
2002.....	159	156	-	-	2	-	1	-	-	-	-	-
2001.....	154	152	-	1	-	1	-	-	-	-	-	-
2000.....	185	183	1	-	-	-	-	1	-	-	-	-
1999.....	189	186	2	-	1	-	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other and unspecified conditions originating in the perinatal period (P05,P08,P29,P50-P96)

All races, both sexes

2004.....	2,657	2,647	4	2	2	-	1	1	-	-	-	-
2003.....	2,661	2,640	12	1	4	2	2	-	-	-	-	-
2002.....	2,370	2,342	9	4	2	4	-	-	1	2	1	1
2001.....	2,455	2,430	6	1	4	3	1	2	-	2	2	-
2000.....	2,662	2,637	8	4	4	1	-	3	1	1	-	1
1999.....	2,645	2,622	11	2	2	2	-	-	2	1	-	1

All races, male

2004.....	1,541	1,537	3	-	1	-	-	-	-	-	-	-
2003.....	1,532	1,521	6	-	4	1	-	-	-	-	-	-
2002.....	1,362	1,344	5	3	1	4	-	-	-	2	1	1
2001.....	1,429	1,414	6	1	4	-	-	-	-	1	1	-
2000.....	1,538	1,521	6	1	3	1	-	3	1	-	-	1
1999.....	1,513	1,497	7	2	1	1	-	-	1	1	-	1

All races, female

2004.....	1,116	1,110	1	2	1	-	1	1	-	-	-	-
2003.....	1,129	1,119	6	1	-	1	2	-	-	-	-	-
2002.....	1,008	998	4	1	1	-	-	-	1	-	-	-
2001.....	1,026	1,016	-	-	-	3	1	2	-	1	1	-
2000.....	1,124	1,116	2	3	1	-	-	-	-	1	-	-
1999.....	1,132	1,125	4	-	1	1	-	-	1	-	-	-

White, both sexes

2004.....	1,714	1,708	2	2	1	-	1	-	-	-	-	-
2003.....	1,726	1,709	10	1	3	2	1	-	-	-	-	-
2002.....	1,558	1,535	7	3	2	4	-	-	-	1	1	1
2001.....	1,606	1,584	5	1	3	2	1	2	-	2	2	-
2000.....	1,689	1,667	8	3	4	1	-	1	1	1	-	1
1999.....	1,662	1,640	10	2	2	2	-	-	2	1	-	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

XVII. Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)

All races, both sexes

2004.....	10,498	5,622	569	205	184	257	226	210	210	208	285	334
2003.....	10,518	5,621	541	180	206	223	228	198	228	242	322	324
2002.....	10,687	5,623	530	199	218	248	244	213	262	264	308	346
2001.....	10,442	5,513	557	182	194	255	250	225	233	259	327	303
2000.....	10,578	5,743	495	198	201	225	216	208	269	277	292	287
1999.....	10,393	5,473	549	207	221	222	212	212	253	279	273	342

All races, male

2004.....	5,423	2,898	318	111	100	145	137	122	113	128	151	189
2003.....	5,460	3,020	275	94	104	144	128	119	114	129	175	177
2002.....	5,515	2,942	282	99	116	146	136	132	152	132	164	195
2001.....	5,448	2,954	282	101	109	161	145	121	128	137	180	156
2000.....	5,517	3,028	259	105	102	137	131	120	159	154	162	157
1999.....	5,346	2,875	310	102	113	139	123	110	132	152	144	185

All races, female

2004.....	5,075	2,724	251	94	84	112	89	88	97	80	134	145
2003.....	5,058	2,601	266	86	102	79	100	79	114	113	147	147
2002.....	5,172	2,681	248	100	102	102	108	81	110	132	144	151
2001.....	4,994	2,559	275	81	85	94	105	104	105	122	147	147
2000.....	5,061	2,715	236	93	99	88	85	88	110	123	130	130
1999.....	5,047	2,598	239	105	108	83	89	102	121	127	129	157

White, both sexes

2004.....	8,331	4,271	400	147	140	185	167	174	171	165	244	283
2003.....	8,378	4,292	392	131	157	171	171	159	184	199	278	264
2002.....	8,556	4,330	395	155	166	189	194	169	215	213	250	291
2001.....	8,338	4,247	387	137	145	189	185	178	197	206	273	249
2000.....	8,396	4,371	347	147	145	159	173	163	228	223	236	245
1999.....	8,297	4,211	385	144	157	170	162	178	208	232	229	292

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

XVII. Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)

White, male

2004.....	4,260	2,182	222	85	77	104	92	102	87	97	130	166
2003.....	4,288	2,282	198	67	79	110	96	95	94	108	152	138
2002.....	4,416	2,297	211	75	84	111	107	104	127	107	135	162
2001.....	4,376	2,301	195	79	90	117	105	104	108	113	148	131
2000.....	4,368	2,316	180	72	75	93	108	93	136	127	132	132
1999.....	4,269	2,240	218	69	79	109	95	93	109	128	117	156

White, female

2004.....	4,071	2,089	178	62	63	81	75	72	84	68	114	117
2003.....	4,090	2,010	194	64	78	61	75	64	90	91	126	126
2002.....	4,140	2,033	184	80	82	78	87	65	88	106	115	129
2001.....	3,962	1,946	192	58	55	72	80	74	89	93	125	118
2000.....	4,028	2,055	167	75	70	66	65	70	92	96	104	113
1999.....	4,028	1,971	167	75	78	61	67	85	99	104	112	136

Black, both sexes

2004.....	1,745	1,077	132	48	37	59	52	30	30	34	35	50
2003.....	1,700	1,039	120	36	38	40	52	33	43	34	36	51
2002.....	1,758	1,047	114	40	43	54	46	35	40	40	53	47
2001.....	1,720	1,032	131	34	41	61	53	36	32	45	46	44
2000.....	1,790	1,102	126	44	47	56	35	38	39	48	45	39
1999.....	1,720	1,025	138	50	56	43	40	28	33	42	35	43

Black, male

2004.....	942	573	77	20	19	33	40	16	21	23	19	23
2003.....	923	574	62	20	19	27	28	19	19	15	18	32
2002.....	909	522	59	23	25	34	26	22	21	18	28	27
2001.....	882	537	69	17	18	41	31	12	19	19	28	18
2000.....	944	567	70	29	23	39	19	24	22	25	26	22
1999.....	899	531	79	25	29	27	24	14	18	19	23	27

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

VII. Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)

Black, female

2004.....	803	504	55	28	18	26	12	14	9	11	16	27
2003.....	777	465	58	16	19	13	24	14	24	19	18	19
2002.....	849	525	55	17	18	20	20	13	19	22	25	20
2001.....	838	495	62	17	23	20	22	24	13	26	18	26
2000.....	846	535	56	15	24	17	16	14	17	23	19	17
1999.....	821	494	59	25	27	16	16	14	15	23	12	16

American Indian 1/, both sexes

2004.....	125	90	8	5	2	3	2	3	4	2	2	-
2003.....	118	72	10	5	6	2	4	1	-	3	3	2
2002.....	93	61	5	3	3	1	-	4	2	1	2	2
2001.....	98	60	15	1	2	-	5	5	-	-	2	3
2000.....	95	68	4	3	-	1	2	-	-	2	2	-
1999.....	106	67	5	3	3	3	4	2	5	-	1	3

American Indian 1/, male

2004.....	70	50	3	4	1	3	-	2	3	2	1	-
2003.....	58	35	4	2	3	1	3	1	-	2	2	2
2002.....	42	29	2	1	2	-	-	2	1	-	1	1
2001.....	46	30	5	-	1	-	5	1	-	-	1	2
2000.....	56	42	1	2	-	-	2	-	-	-	1	-
1999.....	50	28	3	2	2	1	2	-	3	-	-	2

American Indian 1/, female

2004.....	55	40	5	1	1	-	2	1	1	-	1	-
2003.....	60	37	6	3	3	1	1	-	-	1	1	-
2002.....	51	32	3	2	1	1	-	2	1	1	1	1
2001.....	52	30	10	1	1	-	-	4	-	-	1	1
2000.....	39	26	3	1	-	1	-	-	-	2	1	-
1999.....	56	39	2	1	1	2	2	2	2	-	1	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

XVII. Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)

Asian or Pacific Islander 2/, both sexes

2004.....	297	184	29	5	5	10	5	3	5	7	4	1
2003.....	322	218	19	8	5	10	1	5	1	6	5	7
2002.....	280	185	16	1	6	4	4	5	5	10	3	6
2001.....	286	174	24	10	6	5	7	6	4	8	6	7
2000.....	297	202	18	4	9	9	6	7	2	4	9	3
1999.....	270	170	21	10	5	6	6	4	7	5	8	4

Asian or Pacific Islander 2/, male

2004.....	151	93	16	2	3	5	5	2	2	6	1	-
2003.....	191	129	11	5	3	6	1	4	1	4	3	5
2002.....	148	94	10	-	5	1	3	4	3	7	-	5
2001.....	144	86	13	5	-	3	4	4	1	5	3	5
2000.....	149	103	8	2	4	5	2	3	1	2	3	3
1999.....	128	76	10	6	3	2	2	3	2	5	4	-

Asian or Pacific Islander 2/, female

2004.....	146	91	13	3	2	5	-	1	3	1	3	1
2003.....	131	89	8	3	2	4	-	1	-	2	2	2
2002.....	132	91	6	1	1	3	1	1	2	3	3	1
2001.....	142	88	11	5	6	2	3	2	3	3	3	2
2000.....	148	99	10	2	5	4	4	4	1	2	6	-
1999.....	142	94	11	4	2	4	4	1	5	-	4	4

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Anencephaly and similar malformations (Q00)												
All races, both sexes												
2004.....	296	293	1	-	-	-	1	-	1	-	-	-
2003.....	341	334	3	-	1	-	1	-	-	-	-	-
2002.....	301	297	2	-	1	-	1	-	-	-	-	-
2001.....	275	263	5	2	-	-	-	1	1	-	-	-
2000.....	328	324	2	-	-	1	-	-	-	1	-	-
1999.....	317	311	2	3	-	1	-	-	-	-	-	-
All races, male												
2004.....	128	127	-	-	-	-	-	-	1	-	-	-
2003.....	154	153	-	-	-	-	-	-	-	-	-	-
2002.....	139	137	1	-	1	-	-	-	-	-	-	-
2001.....	116	110	2	1	-	-	-	1	-	-	-	-
2000.....	158	156	2	-	-	-	-	-	-	-	-	-
1999.....	126	123	1	1	-	1	-	-	-	-	-	-
All races, female												
2004.....	168	166	1	-	-	-	1	-	-	-	-	-
2003.....	187	181	3	-	1	-	1	-	-	-	-	-
2002.....	162	160	1	-	-	-	1	-	-	-	-	-
2001.....	159	153	3	1	-	-	-	-	1	-	-	-
2000.....	170	168	-	-	-	1	-	-	-	1	-	-
1999.....	191	188	1	2	-	-	-	-	-	-	-	-
White, both sexes												
2004.....	251	248	1	-	-	-	1	-	1	-	-	-
2003.....	286	282	-	-	1	-	1	-	-	-	-	-
2002.....	242	239	1	-	1	-	1	-	-	-	-	-
2001.....	225	218	2	1	-	-	-	1	1	-	-	-
2000.....	266	263	2	-	-	-	-	-	-	1	-	-
1999.....	258	253	1	3	-	1	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Spina bifida (Q05)												
All races, both sexes												
2004.....	141	18	4	3	9	11	11	13	21	12	16	4
2003.....	127	11	3	6	10	12	16	14	14	9	7	5
2002.....	154	20	7	7	11	13	13	13	15	16	10	4
2001.....	138	19	7	12	6	13	12	14	16	10	7	7
2000.....	134	26	5	5	9	11	17	11	11	12	6	7
1999.....	112	22	6	4	14	9	12	10	10	4	5	3
All races, male												
2004.....	62	9	3	-	4	5	7	7	5	7	7	2
2003.....	48	4	1	3	2	5	5	4	4	5	3	4
2002.....	77	12	5	2	5	6	5	6	9	8	6	1
2001.....	57	8	2	6	1	5	5	7	5	6	4	2
2000.....	73	15	1	1	7	5	13	6	4	6	2	4
1999.....	49	12	3	1	7	5	4	2	3	2	2	2
All races, female												
2004.....	79	9	1	3	5	6	4	6	16	5	9	2
2003.....	79	7	2	3	8	7	11	10	10	4	4	1
2002.....	77	8	2	5	6	7	8	7	6	8	4	3
2001.....	81	11	5	6	5	8	7	7	11	4	3	5
2000.....	61	11	4	4	2	6	4	5	7	6	4	3
1999.....	63	10	3	3	7	4	8	8	7	2	3	1
White, both sexes												
2004.....	127	15	1	3	9	10	9	12	20	11	16	3
2003.....	116	10	2	4	9	12	13	12	13	9	7	5
2002.....	134	15	7	7	9	10	11	11	13	15	9	4
2001.....	122	15	4	12	6	11	11	12	14	10	6	6
2000.....	116	23	4	4	7	9	15	10	9	12	4	5
1999.....	102	18	5	4	12	7	12	9	10	4	5	3

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Spina bifida (Q05)												
White, male												
2004.....	54	7	1	-	4	4	5	7	5	6	7	2
2003.....	45	4	-	3	2	5	4	3	4	5	3	4
2002.....	64	10	5	2	3	4	5	4	7	8	5	1
2001.....	50	5	2	6	1	4	5	6	4	6	3	2
2000.....	65	14	1	1	6	3	12	5	4	6	1	3
1999.....	44	9	3	1	6	4	4	2	3	2	2	2
White, female												
2004.....	73	8	-	3	5	6	4	5	15	5	9	1
2003.....	71	6	2	1	7	7	9	9	9	4	4	1
2002.....	70	5	2	5	6	6	6	7	6	7	4	3
2001.....	72	10	2	6	5	7	6	6	10	4	3	4
2000.....	51	9	3	3	1	6	3	5	5	6	3	2
1999.....	58	9	2	3	6	3	8	7	7	2	3	1
Black, both sexes												
2004.....	13	2	3	-	-	1	2	1	1	1	-	1
2003.....	9	-	1	1	1	-	3	2	1	-	-	-
2002.....	17	4	-	-	2	3	1	2	1	1	1	-
2001.....	13	3	1	-	-	2	1	2	2	-	1	1
2000.....	13	2	-	-	2	1	1	1	2	-	2	2
1999.....	9	3	1	-	2	2	-	1	-	-	-	-
Black, male												
2004.....	8	2	2	-	-	1	2	-	-	1	-	-
2003.....	3	-	1	-	-	-	1	1	-	-	-	-
2002.....	12	2	-	-	2	2	-	2	1	-	1	-
2001.....	6	2	-	-	-	1	-	1	1	-	1	-
2000.....	5	-	-	-	1	1	-	1	-	-	1	1
1999.....	4	2	-	-	1	1	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	-------------	-----------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

All other congenital malformations of nervous system (Q01-Q04,Q06-Q07)

All races, both sexes

2004.....	859	439	133	54	43	40	18	19	5	18	14	21
2003.....	816	420	102	50	33	38	29	19	20	12	19	16
2002.....	803	376	127	60	43	31	26	16	14	20	20	17
2001.....	793	391	130	40	36	33	20	22	18	12	18	12
2000.....	762	408	97	61	43	21	21	16	13	12	14	13
1999.....	747	357	105	60	44	26	16	20	20	11	16	17

All races, male

2004.....	404	192	72	25	22	20	12	12	2	9	7	9
2003.....	421	220	58	24	18	23	16	10	7	4	10	9
2002.....	381	186	55	30	15	15	9	11	7	7	10	8
2001.....	403	194	66	21	20	17	10	13	8	9	11	5
2000.....	394	206	44	30	20	13	14	11	6	5	8	11
1999.....	361	168	53	27	21	13	7	11	5	6	11	10

All races, female

2004.....	455	247	61	29	21	20	6	7	3	9	7	12
2003.....	395	200	44	26	15	15	13	9	13	8	9	7
2002.....	422	190	72	30	28	16	17	5	7	13	10	9
2001.....	390	197	64	19	16	16	10	9	10	3	7	7
2000.....	368	202	53	31	23	8	7	5	7	7	6	2
1999.....	386	189	52	33	23	13	9	9	15	5	5	7

White, both sexes

2004.....	642	339	94	39	30	31	10	14	3	12	12	14
2003.....	633	317	79	37	27	35	24	13	14	10	15	10
2002.....	632	295	89	47	31	25	25	15	11	15	17	13
2001.....	613	303	89	31	28	29	15	17	15	10	14	7
2000.....	578	311	74	41	31	15	14	13	10	8	13	10
1999.....	557	261	76	39	34	19	13	18	15	8	14	14

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

All other congenital malformations of nervous system (Q01-Q04,Q06-Q07)

White, male

2004.....	298	148	51	20	15	12	6	8	1	5	6	6
2003.....	320	163	41	17	17	21	15	7	4	4	6	5
2002.....	302	145	39	25	12	12	9	10	5	3	9	7
2001.....	326	154	45	18	20	15	9	12	7	7	9	3
2000.....	302	158	36	19	15	9	9	10	5	3	7	9
1999.....	271	124	39	15	15	10	6	11	4	4	10	7

White, female

2004.....	344	191	43	19	15	19	4	6	2	7	6	8
2003.....	313	154	38	20	10	14	9	6	10	6	9	5
2002.....	330	150	50	22	19	13	16	5	6	12	8	6
2001.....	287	149	44	13	8	14	6	5	8	3	5	4
2000.....	276	153	38	22	16	6	5	3	5	5	6	1
1999.....	286	137	37	24	19	9	7	7	11	4	4	7

Black, both sexes

2004.....	171	80	32	10	11	7	7	3	1	4	2	7
2003.....	143	79	16	11	3	3	4	6	6	2	2	5
2002.....	150	66	33	13	12	5	1	1	3	5	3	4
2001.....	151	73	34	7	7	4	4	4	3	2	4	4
2000.....	153	82	18	16	8	5	7	2	3	4	1	3
1999.....	157	78	24	16	8	7	3	1	5	3	2	3

Black, male

2004.....	81	32	19	3	6	6	6	2	-	2	1	3
2003.....	80	45	12	7	-	2	1	3	3	-	2	3
2002.....	71	35	14	5	3	3	-	1	2	4	1	1
2001.....	67	36	17	2	-	2	-	1	1	2	2	2
2000.....	78	41	6	9	3	3	5	1	1	2	1	2
1999.....	74	35	13	8	6	3	1	-	1	2	1	3

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

All other congenital malformations of nervous system (Q01-Q04,Q06-Q07)

Black, female

2004.....	90	48	13	7	5	1	1	1	1	2	1	4
2003.....	63	34	4	4	3	1	3	3	3	2	-	2
2002.....	79	31	19	8	9	2	1	-	1	1	2	3
2001.....	84	37	17	5	7	2	4	3	2	-	2	2
2000.....	75	41	12	7	5	2	2	1	2	2	-	1
1999.....	83	43	11	8	2	4	2	1	4	1	1	-

American Indian 1/, both sexes

2004.....	16	7	2	2	-	1	1	2	-	1	-	-
2003.....	12	9	1	-	-	-	1	-	-	-	1	-
2002.....	10	8	2	-	-	-	-	-	-	-	-	-
2001.....	13	7	3	-	-	-	1	1	-	-	-	-
2000.....	7	2	3	1	-	-	-	-	-	-	-	-
1999.....	8	4	1	-	1	-	-	1	-	-	-	-

American Indian 1/, male

2004.....	10	5	-	1	-	1	-	2	-	1	-	-
2003.....	4	3	-	-	-	-	-	-	-	-	1	-
2002.....	3	3	-	-	-	-	-	-	-	-	-	-
2001.....	5	3	1	-	-	-	1	-	-	-	-	-
2000.....	4	2	1	1	-	-	-	-	-	-	-	-
1999.....	5	3	1	-	-	-	-	-	-	-	-	-

American Indian 1/, female

2004.....	6	2	2	1	-	-	1	-	-	-	-	-
2003.....	8	6	1	-	-	-	1	-	-	-	-	-
2002.....	7	5	2	-	-	-	-	-	-	-	-	-
2001.....	8	4	2	-	-	-	-	1	-	-	-	-
2000.....	3	-	2	-	-	-	-	-	-	-	-	-
1999.....	3	1	-	-	1	-	-	1	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

All other congenital malformations of nervous system (Q01-Q04,Q06-Q07)

Asian or Pacific Islander 2/, both sexes

2004.....	30	13	5	3	2	1	-	-	1	1	-	-
2003.....	28	15	6	2	3	-	-	-	-	-	1	1
2002.....	11	7	3	-	-	1	-	-	-	-	-	-
2001.....	16	8	4	2	1	-	-	-	-	-	-	1
2000.....	24	13	2	3	4	1	-	1	-	-	-	-
1999.....	25	14	4	5	1	-	-	-	-	-	-	-

Asian or Pacific Islander 2/, male

2004.....	15	7	2	1	1	1	-	-	1	1	-	-
2003.....	17	9	5	-	1	-	-	-	-	-	1	1
2002.....	5	3	2	-	-	-	-	-	-	-	-	-
2001.....	5	1	3	1	-	-	-	-	-	-	-	-
2000.....	10	5	1	1	2	1	-	-	-	-	-	-
1999.....	11	6	-	4	-	-	-	-	-	-	-	-

Asian or Pacific Islander 2/, female

2004.....	15	6	3	2	1	-	-	-	-	-	-	-
2003.....	11	6	1	2	2	-	-	-	-	-	-	-
2002.....	6	4	1	-	-	1	-	-	-	-	-	-
2001.....	11	7	1	1	1	-	-	-	-	-	-	1
2000.....	14	8	1	2	2	-	-	1	-	-	-	-
1999.....	14	8	4	1	1	-	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Congenital malformations of heart (Q20-Q24)

All races, both sexes

2004.....	2,968	1,391	191	77	65	116	98	100	98	82	110	88
2003.....	3,131	1,445	223	57	87	88	84	93	107	117	137	106
2002.....	3,277	1,510	194	63	84	119	93	99	136	119	123	119
2001.....	3,313	1,556	197	47	74	123	124	102	111	114	132	114
2000.....	3,422	1,634	191	58	79	118	100	99	141	120	126	102
1999.....	3,551	1,598	203	63	96	120	102	113	132	138	135	127

All races, male

2004.....	1,621	766	114	44	37	70	60	64	59	53	59	49
2003.....	1,669	821	117	32	47	64	55	58	59	65	73	56
2002.....	1,751	833	119	32	53	78	63	65	80	67	62	59
2001.....	1,783	863	109	27	43	92	73	56	70	62	68	60
2000.....	1,818	867	103	32	45	77	60	59	89	71	69	54
1999.....	1,883	889	118	35	57	83	63	60	71	76	65	67

All races, female

2004.....	1,347	625	77	33	28	46	38	36	39	29	51	39
2003.....	1,462	624	106	25	40	24	29	35	48	52	64	50
2002.....	1,526	677	75	31	31	41	30	34	56	52	61	60
2001.....	1,530	693	88	20	31	31	51	46	41	52	64	54
2000.....	1,604	767	88	26	34	41	40	40	52	49	57	48
1999.....	1,668	709	85	28	39	37	39	53	61	62	70	60

White, both sexes

2004.....	2,349	1,056	135	52	51	80	69	84	79	69	90	77
2003.....	2,462	1,091	157	47	62	57	55	75	85	93	122	84
2002.....	2,610	1,140	153	49	61	92	71	76	114	98	101	105
2001.....	2,606	1,175	140	34	54	84	89	79	94	94	108	97
2000.....	2,706	1,234	134	47	54	78	76	79	123	96	103	88
1999.....	2,859	1,252	137	44	67	93	75	93	114	113	112	112

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Congenital malformations of heart (Q20-Q24)

White, male

2004.....	1,281	583	82	31	30	49	38	55	46	44	49	43
2003.....	1,303	626	82	25	33	42	37	47	47	54	69	43
2002.....	1,392	641	95	23	36	61	48	48	68	58	50	50
2001.....	1,417	665	81	21	34	61	51	48	61	53	56	51
2000.....	1,436	665	71	22	32	49	47	48	77	56	60	47
1999.....	1,510	706	80	24	44	65	46	50	62	63	52	59

White, female

2004.....	1,068	473	53	21	21	31	31	29	33	25	41	34
2003.....	1,159	465	75	22	29	15	18	28	38	39	53	41
2002.....	1,218	499	58	26	25	31	23	28	46	40	51	55
2001.....	1,189	510	59	13	20	23	38	31	33	41	52	46
2000.....	1,270	569	63	25	22	29	29	31	46	40	43	41
1999.....	1,349	546	57	20	23	28	29	43	52	50	60	53

Black, both sexes

2004.....	512	279	43	22	11	31	23	15	17	11	15	11
2003.....	532	275	55	6	22	23	25	14	22	18	13	21
2002.....	539	299	33	12	18	24	22	17	20	16	20	10
2001.....	584	311	46	10	15	38	30	19	13	18	20	13
2000.....	582	317	46	9	23	33	20	17	17	21	19	14
1999.....	579	290	55	15	26	23	24	17	11	23	18	12

Black, male

2004.....	281	151	26	11	5	18	17	8	12	7	8	6
2003.....	285	151	29	3	11	17	14	8	12	6	4	13
2002.....	288	148	19	8	14	16	15	12	12	6	12	7
2001.....	303	163	22	5	8	30	18	6	8	8	11	5
2000.....	313	156	27	8	11	25	11	10	12	14	9	7
1999.....	312	154	31	9	11	16	15	9	7	11	10	8

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Congenital malformations of heart (Q20-Q24)												
Black, female												
2004.....	231	128	17	11	6	13	6	7	5	4	7	5
2003.....	247	124	26	3	11	6	11	6	10	12	9	8
2002.....	251	151	14	4	4	8	7	5	8	10	8	3
2001.....	281	148	24	5	7	8	12	13	5	10	9	8
2000.....	269	161	19	1	12	8	9	7	5	7	10	7
1999.....	267	136	24	6	15	7	9	8	4	12	8	4
American Indian 1/, both sexes												
2004.....	25	15	-	2	1	1	1	-	1	1	2	-
2003.....	38	23	5	1	1	1	3	-	-	2	1	-
2002.....	29	16	1	2	2	-	-	2	-	-	1	2
2001.....	33	18	5	1	2	-	2	2	-	-	1	1
2000.....	36	25	-	1	-	1	1	-	-	1	1	-
1999.....	30	16	2	1	1	1	1	1	3	-	-	-
American Indian 1/, male												
2004.....	17	10	-	2	1	1	-	-	1	1	1	-
2003.....	21	11	3	1	1	-	3	-	-	2	-	-
2002.....	15	11	-	1	1	-	-	1	-	-	-	1
2001.....	15	8	2	-	1	-	2	1	-	-	-	1
2000.....	25	19	-	1	-	-	1	-	-	-	-	-
1999.....	16	7	1	1	1	1	1	-	1	-	-	-
American Indian 1/, female												
2004.....	8	5	-	-	-	-	1	-	-	-	1	-
2003.....	17	12	2	-	-	1	-	-	-	-	1	-
2002.....	14	5	1	1	1	-	-	1	-	-	1	1
2001.....	18	10	3	1	1	-	-	1	-	-	1	-
2000.....	11	6	-	-	-	1	-	-	-	1	1	-
1999.....	14	9	1	-	-	-	-	1	2	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Congenital malformations of heart (Q20-Q24)												
Asian or Pacific Islander 2/, both sexes												
2004.....	82	41	13	1	2	4	5	1	1	1	3	-
2003.....	99	56	6	3	2	7	1	4	-	4	1	1
2002.....	99	55	7	-	3	3	-	4	2	5	1	2
2001.....	90	52	6	2	3	1	3	2	4	2	3	3
2000.....	98	58	11	1	2	6	3	3	1	2	3	-
1999.....	83	40	9	3	2	3	2	2	4	2	5	3
Asian or Pacific Islander 2/, male												
2004.....	42	22	6	-	1	2	5	1	-	1	1	-
2003.....	60	33	3	3	2	5	1	3	-	3	-	-
2002.....	56	33	5	-	2	1	-	4	-	3	-	1
2001.....	48	27	4	1	-	1	2	1	1	1	1	3
2000.....	44	27	5	1	2	3	1	1	-	1	-	-
1999.....	45	22	6	1	1	1	1	1	1	2	3	-
Asian or Pacific Islander 2/, female												
2004.....	40	19	7	1	1	2	-	-	1	-	2	-
2003.....	39	23	3	-	-	2	-	1	-	1	1	1
2002.....	43	22	2	-	1	2	-	-	2	2	1	1
2001.....	42	25	2	1	3	-	1	1	3	1	2	-
2000.....	54	31	6	-	-	3	2	2	1	1	3	-
1999.....	38	18	3	2	1	2	1	1	3	-	2	3

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other congenital malformations of circulatory system (Q25-Q28)

All races, both sexes

2004.....	893	255	27	21	23	23	22	13	12	21	38	55
2003.....	852	237	30	14	22	15	18	12	20	21	44	36
2002.....	901	223	24	9	21	14	19	26	30	29	32	53
2001.....	796	192	15	28	17	23	18	15	12	18	36	44
2000.....	888	208	25	19	22	16	21	17	24	38	44	49
1999.....	885	220	25	15	12	19	19	19	23	46	27	55

All races, male

2004.....	466	140	14	13	10	9	16	9	5	12	23	34
2003.....	446	132	11	3	10	11	8	11	14	14	23	19
2002.....	450	120	13	5	9	3	11	16	21	14	15	30
2001.....	416	101	5	15	12	12	12	9	6	12	21	27
2000.....	464	108	16	9	10	9	14	10	16	22	25	28
1999.....	447	114	15	8	4	13	9	10	13	27	15	30

All races, female

2004.....	427	115	13	8	13	14	6	4	7	9	15	21
2003.....	406	105	19	11	12	4	10	1	6	7	21	17
2002.....	451	103	11	4	12	11	8	10	9	15	17	23
2001.....	380	91	10	13	5	11	6	6	6	6	15	17
2000.....	424	100	9	10	12	7	7	7	8	16	19	21
1999.....	438	106	10	7	8	6	10	9	10	19	12	25

White, both sexes

2004.....	713	177	21	16	19	15	18	10	11	13	31	46
2003.....	693	183	20	10	20	13	13	10	18	15	36	27
2002.....	697	158	19	6	18	9	13	18	26	18	19	38
2001.....	646	144	9	20	12	17	14	12	10	13	28	31
2000.....	707	151	14	15	19	11	19	10	21	28	30	35
1999.....	704	154	19	13	6	13	13	16	19	39	23	45

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other congenital malformations of circulatory system (Q25-Q28)

White, male

2004.....	359	95	10	11	8	5	12	7	4	6	18	30
2003.....	363	108	8	1	10	9	5	9	13	11	20	11
2002.....	332	78	10	3	7	3	7	11	18	9	9	22
2001.....	342	80	4	11	9	8	8	8	5	8	17	20
2000.....	367	79	10	7	8	6	13	5	14	16	15	21
1999.....	360	84	12	7	2	10	7	7	11	24	12	24

White, female

2004.....	354	82	11	5	11	10	6	3	7	7	13	16
2003.....	330	75	12	9	10	4	8	1	5	4	16	16
2002.....	365	80	9	3	11	6	6	7	8	9	10	16
2001.....	304	64	5	9	3	9	6	4	5	5	11	11
2000.....	340	72	4	8	11	5	6	5	7	12	15	14
1999.....	344	70	7	6	4	3	6	9	8	15	11	21

Black, both sexes

2004.....	145	66	3	4	4	4	4	2	1	6	6	8
2003.....	124	39	9	4	-	1	5	1	1	6	7	6
2002.....	167	50	4	3	2	5	4	7	3	8	12	13
2001.....	118	39	4	4	5	5	2	3	2	4	6	10
2000.....	148	49	10	4	3	5	-	5	2	10	7	13
1999.....	147	57	4	2	6	4	4	2	1	5	3	8

Black, male

2004.....	83	35	2	1	2	2	4	1	1	4	5	4
2003.....	66	20	3	2	-	1	3	1	-	3	2	5
2002.....	91	31	2	2	1	-	2	5	2	4	5	6
2001.....	60	18	1	2	3	4	2	1	1	3	3	4
2000.....	80	26	6	2	2	3	-	4	1	6	6	6
1999.....	71	26	2	1	2	3	2	2	1	1	3	5

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other congenital malformations of circulatory system (Q25-Q28)												
Black, female												
2004.....	62	31	1	3	2	2	-	1	-	2	1	4
2003.....	58	19	6	2	-	-	2	-	1	3	5	1
2002.....	76	19	2	1	1	5	2	2	1	4	7	7
2001.....	58	21	3	2	2	1	-	2	1	1	3	6
2000.....	68	23	4	2	1	2	-	1	1	4	1	7
1999.....	76	31	2	1	4	1	2	-	-	4	-	3
American Indian 1/, both sexes												
2004.....	5	3	-	-	-	-	-	-	-	-	-	-
2003.....	8	2	-	-	2	-	-	1	-	-	1	-
2002.....	7	3	-	-	-	-	-	-	-	1	1	-
2001.....	7	3	-	-	-	-	1	-	-	-	-	1
2000.....	6	2	-	-	-	-	-	-	-	-	1	-
1999.....	12	2	2	-	-	1	1	-	1	-	1	1
American Indian 1/, male												
2004.....	4	3	-	-	-	-	-	-	-	-	-	-
2003.....	4	-	-	-	-	-	-	1	-	-	1	-
2002.....	6	3	-	-	-	-	-	-	-	-	1	-
2001.....	4	1	-	-	-	-	1	-	-	-	-	1
2000.....	3	-	-	-	-	-	-	-	-	-	1	-
1999.....	6	1	1	-	-	-	-	-	1	-	-	1
American Indian 1/, female												
2004.....	1	-	-	-	-	-	-	-	-	-	-	-
2003.....	4	2	-	-	2	-	-	-	-	-	-	-
2002.....	1	-	-	-	-	-	-	-	-	1	-	-
2001.....	3	2	-	-	-	-	-	-	-	-	-	-
2000.....	3	2	-	-	-	-	-	-	-	-	-	-
1999.....	6	1	1	-	-	1	1	-	-	-	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other congenital malformations of circulatory system (Q25-Q28)												
Asian or Pacific Islander 2/, both sexes												
2004.....	30	9	3	1	-	4	-	1	-	2	1	1
2003.....	27	13	1	-	-	1	-	-	1	-	-	3
2002.....	30	12	1	-	1	-	2	1	1	2	-	2
2001.....	25	6	2	4	-	1	1	-	-	1	2	2
2000.....	27	6	1	-	-	-	2	2	1	-	6	1
1999.....	22	7	-	-	-	1	1	1	2	2	-	1
Asian or Pacific Islander 2/, male												
2004.....	20	7	2	1	-	2	-	1	-	2	-	-
2003.....	13	4	-	-	-	1	-	-	1	-	-	3
2002.....	21	8	1	-	1	-	2	-	1	1	-	2
2001.....	10	2	-	2	-	-	1	-	-	1	1	2
2000.....	14	3	-	-	-	-	1	1	1	-	3	1
1999.....	10	3	-	-	-	-	-	1	-	2	-	-
Asian or Pacific Islander 2/, female												
2004.....	10	2	1	-	-	2	-	-	-	-	1	1
2003.....	14	9	1	-	-	-	-	-	-	-	-	-
2002.....	9	4	-	-	-	-	-	1	-	1	-	-
2001.....	15	4	2	2	-	1	-	-	-	-	1	-
2000.....	13	3	1	-	-	-	1	1	-	-	3	-
1999.....	12	4	-	-	-	1	1	-	2	-	-	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Congenital malformations of respiratory system (Q30-Q34)												
All races, both sexes												
2004.....	683	633	12	6	2	-	-	3	1	-	1	2
2003.....	668	619	13	2	1	3	6	2	1	1	3	1
2002.....	674	638	10	1	1	1	2	1	1	2	3	2
2001.....	649	594	18	3	4	-	2	2	2	1	-	2
2000.....	728	685	11	-	-	1	2	2	1	1	3	2
1999.....	607	564	13	5	-	1	1	2	1	3	2	-
All races, male												
2004.....	399	375	6	5	-	-	-	-	1	-	-	1
2003.....	378	351	7	1	-	2	3	1	-	1	2	-
2002.....	371	354	6	1	1	1	1	-	-	-	3	1
2001.....	373	344	12	1	3	-	1	1	1	1	-	1
2000.....	408	385	6	-	-	-	2	1	-	1	2	2
1999.....	359	332	11	1	-	-	1	1	1	1	2	-
All races, female												
2004.....	284	258	6	1	2	-	-	3	-	-	1	1
2003.....	290	268	6	1	1	1	3	1	1	-	1	1
2002.....	303	284	4	-	-	-	1	1	1	2	-	1
2001.....	276	250	6	2	1	-	1	1	1	-	-	1
2000.....	320	300	5	-	-	1	-	1	1	-	1	-
1999.....	248	232	2	4	-	1	-	1	-	2	-	-
White, both sexes												
2004.....	499	461	9	3	2	-	-	2	-	-	1	2
2003.....	509	466	10	2	1	3	5	2	1	1	3	1
2002.....	488	459	6	1	1	1	2	-	1	1	2	2
2001.....	490	446	12	2	3	-	1	1	2	1	-	2
2000.....	535	500	9	-	-	-	1	1	1	1	2	2
1999.....	453	420	9	3	-	1	-	1	1	2	2	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Congenital malformations of respiratory system (Q30-Q34)

White, male

2004.....	300	282	5	3	-	-	-	-	-	-	-	1
2003.....	271	247	6	1	-	2	2	1	-	1	2	-
2002.....	278	266	2	1	1	1	1	-	-	-	2	1
2001.....	276	256	7	-	2	-	1	-	1	1	-	1
2000.....	300	282	5	-	-	-	1	-	-	1	1	2
1999.....	262	242	8	-	-	-	-	-	1	1	2	-

White, female

2004.....	199	179	4	-	2	-	-	2	-	-	1	1
2003.....	238	219	4	1	1	1	3	1	1	-	1	1
2002.....	210	193	4	-	-	-	1	-	1	1	-	1
2001.....	214	190	5	2	1	-	-	1	1	-	-	1
2000.....	235	218	4	-	-	-	-	1	1	-	1	-
1999.....	191	178	1	3	-	1	-	1	-	1	-	-

Black, both sexes

2004.....	154	144	2	3	-	-	-	1	1	-	-	-
2003.....	123	118	2	-	-	-	1	-	-	-	-	-
2002.....	158	152	3	-	-	-	-	1	-	1	1	-
2001.....	135	127	6	1	1	-	-	-	-	-	-	-
2000.....	162	154	2	-	-	1	1	1	-	-	1	-
1999.....	130	121	4	1	-	-	1	1	-	1	-	-

Black, male

2004.....	87	83	-	2	-	-	-	-	1	-	-	-
2003.....	83	81	-	-	-	-	1	-	-	-	-	-
2002.....	78	74	3	-	-	-	-	-	-	-	1	-
2001.....	87	80	5	1	1	-	-	-	-	-	-	-
2000.....	88	83	1	-	-	-	1	1	-	-	1	-
1999.....	86	79	3	1	-	-	1	1	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other congenital malformations and deformations (Q10-Q18,Q35-Q89)

All races, both sexes

2004.....	2,687	1,484	129	36	26	51	52	39	46	45	54	69
2003.....	2,687	1,500	110	33	36	47	51	41	47	50	50	71
2002.....	2,665	1,462	110	38	37	51	57	37	37	41	66	80
2001.....	2,619	1,465	116	32	35	44	48	42	33	65	69	58
2000.....	2,593	1,443	103	40	37	36	42	40	42	58	62	57
1999.....	2,584	1,452	142	38	29	30	38	32	37	53	54	68

All races, male

2004.....	1,416	807	77	18	19	30	31	20	22	27	30	42
2003.....	1,466	876	57	22	17	28	28	26	24	25	28	42
2002.....	1,441	809	50	21	21	31	32	22	19	24	41	54
2001.....	1,430	849	59	17	19	26	34	20	17	31	44	28
2000.....	1,401	820	57	24	17	23	22	24	27	31	37	35
1999.....	1,406	826	85	22	13	18	25	16	23	26	34	36

All races, female

2004.....	1,271	677	52	18	7	21	21	19	24	18	24	27
2003.....	1,221	624	53	11	19	19	23	15	23	25	22	29
2002.....	1,224	653	60	17	16	20	25	15	18	17	25	26
2001.....	1,189	616	57	15	16	18	14	22	16	34	25	30
2000.....	1,192	623	46	16	20	13	20	16	15	27	25	22
1999.....	1,178	626	57	16	16	12	13	16	14	27	20	32

White, both sexes

2004.....	2,148	1,132	94	28	20	36	40	34	40	37	48	56
2003.....	2,125	1,135	82	20	24	34	40	35	37	43	44	54
2002.....	2,174	1,157	79	29	29	36	44	33	28	35	55	67
2001.....	2,094	1,131	82	25	24	33	36	34	27	49	59	46
2000.....	2,088	1,111	74	27	27	29	38	33	30	48	51	52
1999.....	2,071	1,097	104	29	21	26	32	28	30	45	44	55

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other congenital malformations and deformations (Q10-Q18,Q35-Q89)

White, male

2004.....	1,117	610	52	16	14	25	23	18	18	21	27	36
2003.....	1,138	653	43	12	10	21	22	21	21	21	25	31
2002.....	1,179	647	36	16	16	20	24	21	17	19	36	45
2001.....	1,133	654	39	13	15	22	24	16	12	25	35	23
2000.....	1,117	630	41	15	12	17	20	19	20	28	30	30
1999.....	1,131	644	62	16	6	16	20	14	19	23	26	28

White, female

2004.....	1,031	522	42	12	6	11	17	16	22	16	21	20
2003.....	987	482	39	8	14	13	18	14	16	22	19	23
2002.....	995	510	43	13	13	16	20	12	11	16	19	22
2001.....	961	477	43	12	9	11	12	18	15	24	24	23
2000.....	971	481	33	12	15	12	18	14	10	20	21	22
1999.....	940	453	42	13	15	10	12	14	11	22	18	27

Black, both sexes

2004.....	425	266	29	7	5	13	12	4	4	7	6	13
2003.....	451	293	23	10	9	11	11	5	10	6	5	14
2002.....	412	250	26	8	6	14	12	4	8	5	9	13
2001.....	429	273	23	7	10	9	9	4	6	15	9	11
2000.....	417	264	25	13	8	6	3	6	12	7	11	5
1999.....	417	280	34	7	5	3	4	4	6	8	9	12

Black, male

2004.....	249	160	20	1	4	4	8	2	4	5	3	6
2003.....	261	175	12	8	5	6	6	4	3	4	3	8
2002.....	227	138	12	5	3	11	7	1	1	4	5	9
2001.....	242	159	14	4	4	3	8	2	5	5	8	5
2000.....	235	150	14	9	5	6	2	4	7	2	7	5
1999.....	231	153	20	5	4	2	4	2	3	3	7	7

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other congenital malformations and deformations (Q10-Q18,Q35-Q89)												
Asian or Pacific Islander 2/, both sexes												
2004.....	71	50	4	-	1	1	-	-	1	1	-	-
2003.....	75	53	1	2	-	1	-	1	-	1	1	1
2002.....	57	40	4	-	1	-	1	-	1	1	2	-
2001.....	71	45	7	-	1	2	2	2	-	1	1	-
2000.....	63	46	4	-	2	1	1	1	-	2	-	-
1999.....	61	44	4	1	2	1	2	-	1	-	1	-
Asian or Pacific Islander 2/, male												
2004.....	29	20	3	-	1	-	-	-	-	1	-	-
2003.....	48	36	1	2	-	-	-	1	-	-	-	1
2002.....	27	19	1	-	1	-	1	-	1	1	-	-
2001.....	43	26	5	-	-	1	1	2	-	1	1	-
2000.....	36	28	2	-	-	-	-	1	-	1	-	-
1999.....	28	17	3	-	2	-	1	-	1	-	1	-
Asian or Pacific Islander 2/, female												
2004.....	42	30	1	-	-	1	-	-	1	-	-	-
2003.....	27	17	-	-	-	1	-	-	-	1	1	-
2002.....	30	21	3	-	-	-	-	-	-	-	2	-
2001.....	28	19	2	-	1	1	1	-	-	-	-	-
2000.....	27	18	2	-	2	1	1	-	-	1	-	-
1999.....	33	27	1	1	-	1	1	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Down's syndrome (Q90)

All races, both sexes

2004.....	866	113	21	2	5	10	15	17	23	28	48	95
2003.....	837	116	11	3	7	11	13	11	16	30	58	88
2002.....	833	118	21	8	10	13	17	18	25	36	52	68
2001.....	828	119	20	6	10	8	21	24	39	37	60	63
2000.....	713	93	19	7	4	14	8	23	32	32	34	56
1999.....	619	94	13	4	9	7	14	12	27	21	34	70

All races, male

2004.....	458	60	14	2	4	8	8	6	16	18	25	52
2003.....	415	54	6	2	6	5	7	5	6	15	34	46
2002.....	435	64	16	5	8	8	9	11	13	12	27	40
2001.....	411	71	12	3	6	6	8	13	20	15	30	32
2000.....	350	54	12	6	1	9	3	9	16	17	17	23
1999.....	308	47	8	2	5	4	8	6	15	14	15	40

All races, female

2004.....	408	53	7	-	1	2	7	11	7	10	23	43
2003.....	422	62	5	1	1	6	6	6	10	15	24	42
2002.....	398	54	5	3	2	5	8	7	12	24	25	28
2001.....	417	48	8	3	4	2	13	11	19	22	30	31
2000.....	363	39	7	1	3	5	5	14	16	15	17	33
1999.....	311	47	5	2	4	3	6	6	12	7	19	30

White, both sexes

2004.....	765	83	15	2	3	7	12	13	15	22	42	85
2003.....	739	91	7	2	5	8	11	9	13	26	47	82
2002.....	729	98	13	7	7	11	13	13	19	30	46	60
2001.....	730	96	14	4	7	7	14	20	33	27	53	57
2000.....	631	72	13	6	2	11	6	17	29	26	30	52
1999.....	536	79	7	3	6	5	8	9	17	18	29	62

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Down's syndrome (Q90)

White, male

2004.....	400	44	10	2	2	6	5	3	10	14	23	48
2003.....	360	39	4	2	4	4	6	5	5	12	25	43
2002.....	375	53	9	5	5	7	7	9	10	10	24	34
2001.....	367	59	8	2	5	5	5	12	17	12	26	30
2000.....	316	46	8	5	1	8	3	6	15	16	16	20
1999.....	267	39	4	2	2	2	6	5	9	11	13	36

White, female

2004.....	365	39	5	-	1	1	7	10	5	8	19	37
2003.....	379	52	3	-	1	4	5	4	8	14	22	39
2002.....	354	45	4	2	2	4	6	4	9	20	22	26
2001.....	363	37	6	2	2	2	9	8	16	15	27	27
2000.....	315	26	5	1	1	3	3	11	14	10	14	32
1999.....	269	40	3	1	4	3	2	4	8	7	16	26

Black, both sexes

2004.....	83	21	6	-	2	3	3	4	4	4	6	10
2003.....	80	17	3	-	2	2	2	2	3	2	9	5
2002.....	87	16	8	-	2	2	4	3	4	4	6	6
2001.....	78	15	5	2	2	1	7	3	6	6	6	5
2000.....	74	19	6	1	1	3	2	6	3	6	4	2
1999.....	67	11	6	1	3	1	4	2	9	2	3	8

Black, male

2004.....	48	12	4	-	2	2	3	3	3	3	2	4
2003.....	45	10	1	-	2	1	1	-	1	2	7	3
2002.....	50	9	7	-	2	1	2	1	2	-	3	4
2001.....	34	7	3	1	1	1	3	1	3	1	3	2
2000.....	28	7	4	1	-	1	-	3	1	1	1	1
1999.....	33	6	4	-	3	1	1	-	5	2	2	4

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Down's syndrome (Q90)

Asian or Pacific Islander 2/, both sexes

2004.....	12	5	-	-	-	-	-	-	2	2	-	-
2003.....	14	7	1	-	-	1	-	-	-	1	2	1
2002.....	13	3	-	1	1	-	-	-	1	2	-	2
2001.....	15	5	-	-	1	-	-	1	-	4	-	1
2000.....	7	2	-	-	1	-	-	-	-	-	-	2
1999.....	12	3	-	-	-	1	1	1	-	1	2	-

Asian or Pacific Islander 2/, male

2004.....	6	2	-	-	-	-	-	-	1	1	-	-
2003.....	10	5	1	-	-	-	-	-	-	1	2	-
2002.....	8	1	-	-	1	-	-	-	1	2	-	2
2001.....	5	2	-	-	-	-	-	-	-	2	-	-
2000.....	5	1	-	-	-	-	-	-	-	-	-	2
1999.....	6	2	-	-	-	1	-	1	-	1	-	-

Asian or Pacific Islander 2/, female

2004.....	6	3	-	-	-	-	-	-	1	1	-	-
2003.....	4	2	-	-	-	1	-	-	-	-	-	1
2002.....	5	2	-	1	-	-	-	-	-	-	-	-
2001.....	10	3	-	-	1	-	-	1	-	2	-	1
2000.....	2	1	-	-	1	-	-	-	-	-	-	-
1999.....	6	1	-	-	-	-	1	-	-	-	2	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

All other chromosomal abnormalities, not elsewhere classified (Q91-Q99)

All races, both sexes

2004.....	1,105	996	51	6	11	6	9	6	3	2	4	-
2003.....	1,059	939	46	15	9	9	10	6	3	2	4	1
2002.....	1,079	979	35	13	10	6	16	3	4	1	2	3
2001.....	1,031	914	49	12	12	11	5	3	1	2	5	3
2000.....	1,010	922	42	8	7	7	5	-	5	3	3	1
1999.....	971	855	40	15	17	9	10	4	3	3	-	2

All races, male

2004.....	469	422	18	4	4	3	3	4	2	2	-	-
2003.....	463	409	18	7	4	6	6	4	-	-	2	1
2002.....	470	427	17	3	3	4	6	1	3	-	-	2
2001.....	459	414	15	10	5	3	2	1	1	1	2	1
2000.....	451	417	18	3	2	1	3	-	1	1	2	-
1999.....	407	364	16	5	6	2	6	4	1	-	-	-

All races, female

2004.....	636	574	33	2	7	3	6	2	1	-	4	-
2003.....	596	530	28	8	5	3	4	2	3	2	2	-
2002.....	609	552	18	10	7	2	10	2	1	1	2	1
2001.....	572	500	34	2	7	8	3	2	-	1	3	2
2000.....	559	505	24	5	5	6	2	-	4	2	1	1
1999.....	564	491	24	10	11	7	4	-	2	3	-	2

White, both sexes

2004.....	837	760	30	4	6	6	8	5	2	1	4	-
2003.....	815	717	35	9	8	9	9	3	3	2	4	1
2002.....	850	769	28	9	9	5	14	3	3	1	1	2
2001.....	812	719	35	8	11	8	5	2	1	2	5	3
2000.....	769	706	23	7	5	6	4	-	5	3	3	1
1999.....	757	677	27	6	11	5	9	4	2	3	-	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

All other chromosomal abnormalities, not elsewhere classified (Q91-Q99)

White, male

2004.....	346	309	11	2	4	3	3	4	2	1	-	-
2003.....	358	313	14	6	3	6	5	2	-	-	2	1
2002.....	379	344	14	-	3	3	6	1	2	-	-	2
2001.....	366	334	8	7	4	2	2	1	1	1	2	1
2000.....	340	319	6	3	1	1	3	-	1	1	2	-
1999.....	323	293	10	3	4	1	6	4	-	-	-	-

White, female

2004.....	491	451	19	2	2	3	5	1	-	-	4	-
2003.....	457	404	21	3	5	3	4	1	3	2	2	-
2002.....	471	425	14	9	6	2	8	2	1	1	1	-
2001.....	446	385	27	1	7	6	3	1	-	1	3	2
2000.....	429	387	17	4	4	5	1	-	4	2	1	1
1999.....	434	384	17	3	7	4	3	-	2	3	-	1

Black, both sexes

2004.....	208	185	14	2	4	-	1	-	1	1	-	-
2003.....	197	180	8	4	1	-	1	3	-	-	-	-
2002.....	186	169	6	4	1	1	2	-	1	-	1	1
2001.....	174	157	9	3	1	2	-	1	-	-	-	-
2000.....	197	172	19	1	2	1	1	-	-	-	-	-
1999.....	174	146	9	8	6	3	-	-	1	-	-	-

Black, male

2004.....	89	82	4	2	-	-	-	-	-	1	-	-
2003.....	85	77	4	-	1	-	1	2	-	-	-	-
2002.....	77	70	2	3	-	1	-	-	1	-	-	-
2001.....	72	62	6	2	1	-	-	-	-	-	-	-
2000.....	96	83	12	-	1	-	-	-	-	-	-	-
1999.....	72	61	5	1	2	1	-	-	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

All other chromosomal abnormalities, not elsewhere classified (Q91-Q99)

Black, female

2004.....	119	103	10	-	4	-	1	-	1	-	-	-
2003.....	112	103	4	4	-	-	-	1	-	-	-	-
2002.....	109	99	4	1	1	-	2	-	-	-	1	1
2001.....	102	95	3	1	-	2	-	1	-	-	-	-
2000.....	101	89	7	1	1	1	1	-	-	-	-	-
1999.....	102	85	4	7	4	2	-	-	-	-	-	-

American Indian 1/, both sexes

2004.....	17	12	4	-	1	-	-	-	-	-	-	-
2003.....	8	7	-	1	-	-	-	-	-	-	-	-
2002.....	7	6	-	-	-	-	-	-	-	-	-	-
2001.....	6	5	1	-	-	-	-	-	-	-	-	-
2000.....	12	12	-	-	-	-	-	-	-	-	-	-
1999.....	6	3	-	-	-	1	1	-	-	-	-	1

American Indian 1/, male

2004.....	7	6	1	-	-	-	-	-	-	-	-	-
2003.....	4	3	-	1	-	-	-	-	-	-	-	-
2002.....	3	3	-	-	-	-	-	-	-	-	-	-
2001.....	1	1	-	-	-	-	-	-	-	-	-	-
2000.....	6	6	-	-	-	-	-	-	-	-	-	-
1999.....	1	1	-	-	-	-	-	-	-	-	-	-

American Indian 1/, female

2004.....	10	6	3	-	1	-	-	-	-	-	-	-
2003.....	4	4	-	-	-	-	-	-	-	-	-	-
2002.....	4	3	-	-	-	-	-	-	-	-	-	-
2001.....	5	4	1	-	-	-	-	-	-	-	-	-
2000.....	6	6	-	-	-	-	-	-	-	-	-	-
1999.....	5	2	-	-	-	1	1	-	-	-	-	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

XVIII. Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

All races, both sexes

2004.....	30,126	3,361	189	55	65	208	450	523	569	768	1,132	1,270
2003.....	31,444	3,318	256	75	103	300	537	617	710	1,037	1,345	1,578
2002.....	29,975	3,456	248	61	78	256	484	562	757	1,013	1,389	1,368
2001.....	32,326	3,491	272	79	102	361	588	663	867	1,379	1,747	1,882
2000.....	31,876	3,639	248	85	92	317	551	663	891	1,284	1,703	1,727
1999.....	26,672	3,538	236	65	72	228	381	446	649	940	1,122	1,168

All races, male

2004.....	13,712	2,021	100	33	26	133	319	361	371	490	745	813
2003.....	14,583	1,974	143	30	62	197	366	434	473	685	841	1,011
2002.....	14,269	2,032	145	37	45	172	329	391	514	670	870	895
2001.....	15,779	2,041	162	44	55	249	405	461	579	888	1,119	1,220
2000.....	15,863	2,156	137	48	51	196	385	456	604	857	1,137	1,157
1999.....	12,877	2,072	136	35	42	153	269	287	434	599	740	796

All races, female

2004.....	16,414	1,340	89	22	39	75	131	162	198	278	387	457
2003.....	16,861	1,344	113	45	41	103	171	183	237	352	504	567
2002.....	15,706	1,424	103	24	33	84	155	171	243	343	519	473
2001.....	16,547	1,450	110	35	47	112	183	202	288	491	628	662
2000.....	16,013	1,483	111	37	41	121	166	207	287	427	566	570
1999.....	13,795	1,466	100	30	30	75	112	159	215	341	382	372

White, both sexes

2004.....	25,030	2,091	127	37	40	160	355	400	435	604	880	1,014
2003.....	26,147	2,125	166	57	61	244	415	464	550	806	1,043	1,188
2002.....	24,749	2,224	161	35	49	192	369	422	580	775	1,079	1,024
2001.....	26,877	2,231	185	53	71	288	459	519	683	1,089	1,423	1,517
2000.....	26,285	2,294	182	57	63	240	416	495	690	1,035	1,360	1,364
1999.....	21,591	2,227	159	49	47	163	270	312	486	699	808	849

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

XVIII. Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

White, male

2004.....	11,082	1,250	68	27	15	103	255	287	289	395	590	651
2003.....	11,802	1,313	91	26	35	163	290	328	378	534	655	772
2002.....	11,421	1,323	89	22	30	130	261	303	397	517	681	669
2001.....	12,868	1,320	106	25	41	198	318	379	461	702	933	994
2000.....	12,827	1,363	106	35	33	157	301	349	481	714	935	929
1999.....	10,094	1,321	96	26	28	110	199	211	318	463	542	572

White, female

2004.....	13,948	841	59	10	25	57	100	113	146	209	290	363
2003.....	14,345	812	75	31	26	81	125	136	172	272	388	416
2002.....	13,328	901	72	13	19	62	108	119	183	258	398	355
2001.....	14,009	911	79	28	30	90	141	140	222	387	490	523
2000.....	13,458	931	76	22	30	83	115	146	209	321	425	435
1999.....	11,497	906	63	23	19	53	71	101	168	236	266	277

Black, both sexes

2004.....	4,490	1,135	50	14	22	43	73	107	116	146	225	233
2003.....	4,708	1,051	80	14	37	47	107	132	139	203	284	354
2002.....	4,638	1,095	67	24	26	58	96	121	158	218	277	308
2001.....	4,691	1,104	69	20	22	55	111	123	147	241	282	326
2000.....	5,004	1,212	57	21	25	67	119	142	174	214	314	325
1999.....	4,591	1,167	68	15	18	54	100	116	150	231	283	299

Black, male

2004.....	2,312	689	28	5	11	27	49	64	73	83	133	148
2003.....	2,459	574	44	3	24	29	65	94	82	129	171	215
2002.....	2,536	630	45	14	14	39	54	73	104	140	169	203
2001.....	2,465	626	44	15	11	38	75	69	85	145	158	198
2000.....	2,693	710	29	11	17	32	76	91	107	115	178	200
1999.....	2,517	669	35	9	11	38	63	67	109	131	176	209

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

XVIII. Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

Black, female

2004.....	2,178	446	22	9	11	16	24	43	43	63	92	85
2003.....	2,249	477	36	11	13	18	42	38	57	74	113	139
2002.....	2,102	465	22	10	12	19	42	48	54	78	108	105
2001.....	2,226	478	25	5	11	17	36	54	62	96	124	128
2000.....	2,311	502	28	10	8	35	43	51	67	99	136	125
1999.....	2,074	498	33	6	7	16	37	49	41	100	107	90

American Indian 1/, both sexes

2004.....	214	64	5	2	1	3	13	6	10	9	13	10
2003.....	231	76	4	1	3	5	5	6	6	19	9	17
2002.....	248	72	7	1	-	3	6	8	12	9	22	17
2001.....	263	81	6	1	3	4	5	6	14	14	14	10
2000.....	209	61	3	1	-	4	5	5	12	12	9	13
1999.....	215	72	3	-	2	3	3	8	4	5	10	9

American Indian 1/, male

2004.....	122	41	1	-	-	2	11	4	5	7	10	8
2003.....	135	44	3	-	3	3	5	4	5	15	6	14
2002.....	130	41	3	-	-	1	5	6	7	4	14	9
2001.....	160	52	5	1	-	3	4	3	13	12	10	7
2000.....	127	42	1	-	-	3	3	2	5	7	7	12
1999.....	117	39	1	-	1	2	3	4	1	3	6	7

American Indian 1/, female

2004.....	92	23	4	2	1	1	2	2	5	2	3	2
2003.....	96	32	1	1	-	2	-	2	1	4	3	3
2002.....	118	31	4	1	-	2	1	2	5	5	8	8
2001.....	103	29	1	-	3	1	1	3	1	2	4	3
2000.....	82	19	2	1	-	1	2	3	7	5	2	1
1999.....	98	33	2	-	1	1	-	4	3	2	4	2

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

XVIII. Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

Asian or Pacific Islander 2/, both sexes

2004.....	392	71	7	2	2	2	9	10	8	9	14	13
2003.....	358	66	6	3	2	4	10	15	15	9	9	19
2002.....	340	65	13	1	3	3	13	11	7	11	11	19
2001.....	495	75	12	5	6	14	13	15	23	35	28	29
2000.....	378	72	6	6	4	6	11	21	15	23	20	25
1999.....	275	72	6	1	5	8	8	10	9	5	21	11

Asian or Pacific Islander 2/, male

2004.....	196	41	3	1	-	1	4	6	4	5	12	6
2003.....	187	43	5	1	-	2	6	8	8	7	9	10
2002.....	182	38	8	1	1	2	9	9	6	9	6	14
2001.....	286	43	7	3	3	10	8	10	20	29	18	21
2000.....	216	41	1	2	1	4	5	14	11	21	17	16
1999.....	149	43	4	-	2	3	4	5	6	2	16	8

Asian or Pacific Islander 2/, female

2004.....	196	30	4	1	2	1	5	4	4	4	2	7
2003.....	171	23	1	2	2	2	4	7	7	2	-	9
2002.....	158	27	5	-	2	1	4	2	1	2	5	5
2001.....	209	32	5	2	3	4	5	5	3	6	10	8
2000.....	162	31	5	4	3	2	6	7	4	2	3	9
1999.....	126	29	2	1	3	5	4	5	3	3	5	3

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Senility (R54)												
All races, both sexes												
2004.....	4,806	-	-	-	-
2003.....	4,676	-	-	-	1
2002.....	4,269	-	-	1	2
2001.....	4,189	-	1	-	-
2000.....	3,785	-	-	-	-
1999.....	3,150	-	-	1	-
All races, male												
2004.....	1,336	-	-	-	-
2003.....	1,238	-	-	-	1
2002.....	1,179	-	-	-	1
2001.....	1,154	-	1	-	-
2000.....	1,018	-	-	-	-
1999.....	890	-	-	-	-
All races, female												
2004.....	3,470	-	-	-	-
2003.....	3,438	-	-	-	-
2002.....	3,090	-	-	1	1
2001.....	3,035	-	-	-	-
2000.....	2,767	-	-	-	-
1999.....	2,260	-	-	1	-
White, both sexes												
2004.....	4,444	-	-	-	-
2003.....	4,360	-	-	-	1
2002.....	3,959	-	-	1	-
2001.....	3,872	-	1	-	-
2000.....	3,470	-	-	-	-
1999.....	2,923	-	-	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Senility (R54)												
White, male												
2004.....	1,236	-	-	-	-
2003.....	1,150	-	-	-	1
2002.....	1,093	-	-	-	-
2001.....	1,061	-	1	-	-
2000.....	941	-	-	-	-
1999.....	812	-	-	-	-
White, female												
2004.....	3,208	-	-	-	-
2003.....	3,210	-	-	-	-
2002.....	2,866	-	-	1	-
2001.....	2,811	-	-	-	-
2000.....	2,529	-	-	-	-
1999.....	2,111	-	-	1	-
Black, both sexes												
2004.....	303	-	-	-	-
2003.....	257	-	-	-	-
2002.....	257	-	-	-	2
2001.....	275	-	-	-	-
2000.....	283	-	-	-	-
1999.....	213	-	-	-	-
Black, male												
2004.....	84	-	-	-	-
2003.....	67	-	-	-	-
2002.....	73	-	-	-	1
2001.....	83	-	-	-	-
2000.....	68	-	-	-	-
1999.....	72	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Senility (R54)

Black, female

2004.....	219	-	-	-	-
2003.....	190	-	-	-	-
2002.....	184	-	-	-	1
2001.....	192	-	-	-	-
2000.....	215	-	-	-	-
1999.....	141	-	-	-	-

American Indian 1/, both sexes

2004.....	11	-	-	-	-
2003.....	21	-	-	-	-
2002.....	9	-	-	-	-
2001.....	17	-	-	-	-
2000.....	14	-	-	-	-
1999.....	11	-	-	-	-

American Indian 1/, male

2004.....	2	-	-	-	-
2003.....	7	-	-	-	-
2002.....	1	-	-	-	-
2001.....	4	-	-	-	-
2000.....	4	-	-	-	-
1999.....	5	-	-	-	-

American Indian 1/, female

2004.....	9	-	-	-	-
2003.....	14	-	-	-	-
2002.....	8	-	-	-	-
2001.....	13	-	-	-	-
2000.....	10	-	-	-	-
1999.....	6	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Senility (R54)												
Asian or Pacific Islander 2/, both sexes												
2004.....	48	-	-	-	-
2003.....	38	-	-	-	-
2002.....	44	-	-	-	-
2001.....	25	-	-	-	-
2000.....	18	-	-	-	-
1999.....	3	-	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	14	-	-	-	-
2003.....	14	-	-	-	-
2002.....	12	-	-	-	-
2001.....	6	-	-	-	-
2000.....	5	-	-	-	-
1999.....	1	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	34	-	-	-	-
2003.....	24	-	-	-	-
2002.....	32	-	-	-	-
2001.....	19	-	-	-	-
2000.....	13	-	-	-	-
1999.....	2	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Ill-defined and unknown causes of mortality (R95-R99)

All races, both sexes

2004.....	15,545	3,290	157	40	34	165	350	410	447	630	934	1,026
2003.....	17,417	3,252	203	55	75	250	414	476	566	856	1,132	1,293
2002.....	17,200	3,379	201	45	55	198	392	440	587	821	1,160	1,122
2001.....	20,310	3,419	229	54	71	300	483	559	742	1,194	1,514	1,652
2000.....	20,645	3,572	190	60	69	259	449	530	753	1,101	1,464	1,496
1999.....	16,553	3,466	192	48	55	175	312	335	531	754	935	972

All races, male

2004.....	8,691	1,979	86	24	9	102	252	293	297	401	632	677
2003.....	9,765	1,929	112	20	44	165	290	334	389	567	723	837
2002.....	9,669	1,992	116	27	31	134	273	310	404	554	746	753
2001.....	11,478	1,993	139	31	35	204	342	398	507	771	970	1,073
2000.....	11,709	2,123	111	36	39	165	316	367	528	732	980	1,020
1999.....	9,110	2,027	110	21	32	119	231	220	366	480	622	666

All races, female

2004.....	6,854	1,311	71	16	25	63	98	117	150	229	302	349
2003.....	7,652	1,323	91	35	31	85	124	142	177	289	409	456
2002.....	7,531	1,387	85	18	24	64	119	130	183	267	414	369
2001.....	8,832	1,426	90	23	36	96	141	161	235	423	544	579
2000.....	8,936	1,449	79	24	30	94	133	163	225	369	484	476
1999.....	7,443	1,439	82	27	23	56	81	115	165	274	313	306

White, both sexes

2004.....	12,346	2,047	105	26	22	131	289	328	362	509	747	827
2003.....	13,905	2,070	131	41	47	204	327	369	448	684	886	1,003
2002.....	13,633	2,177	130	26	32	158	313	333	471	641	922	854
2001.....	16,431	2,179	157	34	47	246	392	451	589	964	1,262	1,345
2000.....	16,583	2,247	138	40	43	198	344	404	605	900	1,183	1,204
1999.....	12,872	2,180	126	34	37	127	229	242	406	574	681	720

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Ill-defined and unknown causes of mortality (R95-R99)

White, male

2004.....	6,857	1,230	60	19	5	81	210	246	244	336	519	547
2003.....	7,732	1,278	68	17	27	138	235	258	317	456	570	658
2002.....	7,559	1,300	71	17	20	110	224	245	326	433	594	572
2001.....	9,259	1,285	90	17	25	167	280	332	409	624	831	886
2000.....	9,337	1,339	85	27	22	132	252	286	432	618	813	835
1999.....	6,952	1,292	77	14	20	89	176	164	278	375	464	488

White, female

2004.....	5,489	817	45	7	17	50	79	82	118	173	228	280
2003.....	6,173	792	63	24	20	66	92	111	131	228	316	345
2002.....	6,074	877	59	9	12	48	89	88	145	208	328	282
2001.....	7,172	894	67	17	22	79	112	119	180	340	431	459
2000.....	7,246	908	53	13	21	66	92	118	173	282	370	369
1999.....	5,920	888	49	20	17	38	53	78	128	199	217	232

Black, both sexes

2004.....	2,824	1,114	43	11	9	30	43	69	71	106	165	180
2003.....	3,137	1,043	64	11	24	39	79	91	105	154	228	264
2002.....	3,160	1,066	58	17	20	35	62	90	100	162	214	242
2001.....	3,280	1,090	57	14	16	39	73	89	121	185	213	269
2000.....	3,614	1,193	44	13	22	52	92	102	127	172	258	260
1999.....	3,291	1,145	58	13	13	40	74	77	112	172	227	233

Black, male

2004.....	1,612	671	23	4	4	18	29	39	46	55	94	118
2003.....	1,804	567	37	2	15	23	49	67	62	97	138	161
2002.....	1,873	613	37	9	10	22	37	51	67	109	135	164
2001.....	1,839	617	37	10	8	27	50	53	70	109	113	160
2000.....	2,095	701	24	7	16	27	57	66	82	90	148	162
1999.....	1,931	654	29	7	11	27	48	48	81	101	139	163

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Ill-defined and unknown causes of mortality (R95-R99)

Black, female

2004.....	1,212	443	20	7	5	12	14	30	25	51	71	62
2003.....	1,333	476	27	9	9	16	30	24	43	57	90	103
2002.....	1,287	453	21	8	10	13	25	39	33	53	79	78
2001.....	1,441	473	20	4	8	12	23	36	51	76	100	109
2000.....	1,519	492	20	6	6	25	35	36	45	82	110	98
1999.....	1,360	491	29	6	2	13	26	29	31	71	88	70

American Indian 1/, both sexes

2004.....	156	62	4	2	1	2	10	4	7	6	9	8
2003.....	166	74	2	1	2	3	3	5	3	11	9	11
2002.....	194	71	5	1	-	3	6	6	11	8	15	13
2001.....	205	78	6	1	3	3	5	5	11	12	13	9
2000.....	153	61	3	1	-	3	4	4	7	7	6	9
1999.....	172	71	3	-	1	1	3	7	4	3	7	9

American Indian 1/, male

2004.....	96	40	1	-	-	2	9	3	4	5	8	6
2003.....	103	42	2	-	2	2	3	3	3	9	6	9
2002.....	109	41	3	-	-	1	5	5	6	4	11	7
2001.....	136	50	5	1	-	2	4	3	10	10	9	6
2000.....	99	42	1	-	-	2	3	2	3	4	5	8
1999.....	99	39	1	-	-	1	3	3	1	2	4	7

American Indian 1/, female

2004.....	60	22	3	2	1	-	1	1	3	1	1	2
2003.....	63	32	-	1	-	1	-	2	-	2	3	2
2002.....	85	30	2	1	-	2	1	1	5	4	4	6
2001.....	69	28	1	-	3	1	1	2	1	2	4	3
2000.....	54	19	2	1	-	1	1	2	4	3	1	1
1999.....	73	32	2	-	1	-	-	4	3	1	3	2

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Ill-defined and unknown causes of mortality (R95-R99)

Asian or Pacific Islander 2/, both sexes

2004.....	219	67	5	1	2	2	8	9	7	9	13	11
2003.....	209	65	6	2	2	4	5	11	10	7	9	15
2002.....	213	65	8	1	3	2	11	11	5	10	9	13
2001.....	394	72	9	5	5	12	13	14	21	33	26	29
2000.....	295	71	5	6	4	6	9	20	14	22	17	23
1999.....	218	70	5	1	4	7	6	9	9	5	20	10

Asian or Pacific Islander 2/, male

2004.....	126	38	2	1	-	1	4	5	3	5	11	6
2003.....	126	42	5	1	-	2	3	6	7	5	9	9
2002.....	128	38	5	1	1	1	7	9	5	8	6	10
2001.....	244	41	7	3	2	8	8	10	18	28	17	21
2000.....	178	41	1	2	1	4	4	13	11	20	14	15
1999.....	128	42	3	-	1	2	4	5	6	2	15	8

Asian or Pacific Islander 2/, female

2004.....	93	29	3	-	2	1	4	4	4	4	2	5
2003.....	83	23	1	1	2	2	2	5	3	2	-	6
2002.....	85	27	3	-	2	1	4	2	-	2	3	3
2001.....	150	31	2	2	3	4	5	4	3	5	9	8
2000.....	117	30	4	4	3	2	5	7	3	2	3	8
1999.....	90	28	2	1	3	5	2	4	3	3	5	2

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other ill-defined and unknown causes of mortality (R96-R99)												
All races, both sexes												
2004.....	13,299	1,044	157	40	34	165	350	410	447	630	934	1,026
2003.....	15,255	1,090	203	55	75	250	414	476	566	856	1,132	1,293
2002.....	14,905	1,084	201	45	55	198	392	440	587	821	1,160	1,122
2001.....	18,076	1,185	229	54	71	300	483	559	742	1,194	1,514	1,652
2000.....	18,122	1,049	190	60	69	259	449	530	753	1,101	1,464	1,496
1999.....	13,905	818	192	48	55	175	312	335	531	754	935	972
All races, male												
2004.....	7,363	651	86	24	9	102	252	293	297	401	632	677
2003.....	8,485	649	112	20	44	165	290	334	389	567	723	837
2002.....	8,318	641	116	27	31	134	273	310	404	554	746	753
2001.....	10,161	676	139	31	35	204	342	398	507	771	970	1,073
2000.....	10,200	614	111	36	39	165	316	367	528	732	980	1,020
1999.....	7,557	474	110	21	32	119	231	220	366	480	622	666
All races, female												
2004.....	5,936	393	71	16	25	63	98	117	150	229	302	349
2003.....	6,770	441	91	35	31	85	124	142	177	289	409	456
2002.....	6,587	443	85	18	24	64	119	130	183	267	414	369
2001.....	7,915	509	90	23	36	96	141	161	235	423	544	579
2000.....	7,922	435	79	24	30	94	133	163	225	369	484	476
1999.....	6,348	344	82	27	23	56	81	115	165	274	313	306
White, both sexes												
2004.....	10,905	606	105	26	22	131	289	328	362	509	747	827
2003.....	12,538	703	131	41	47	204	327	369	448	684	886	1,003
2002.....	12,139	683	130	26	32	158	313	333	471	641	922	854
2001.....	15,025	773	157	34	47	246	392	451	589	964	1,262	1,345
2000.....	14,975	639	138	40	43	198	344	404	605	900	1,183	1,204
1999.....	11,161	469	126	34	37	127	229	242	406	574	681	720

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other ill-defined and unknown causes of mortality (R96-R99)

White, male

2004.....	6,009	382	60	19	5	81	210	246	244	336	519	547
2003.....	6,889	435	68	17	27	138	235	258	317	456	570	658
2002.....	6,669	410	71	17	20	110	224	245	326	433	594	572
2001.....	8,418	444	90	17	25	167	280	332	409	624	831	886
2000.....	8,370	372	85	27	22	132	252	286	432	618	813	835
1999.....	5,939	279	77	14	20	89	176	164	278	375	464	488

White, female

2004.....	4,896	224	45	7	17	50	79	82	118	173	228	280
2003.....	5,649	268	63	24	20	66	92	111	131	228	316	345
2002.....	5,470	273	59	9	12	48	89	88	145	208	328	282
2001.....	6,607	329	67	17	22	79	112	119	180	340	431	459
2000.....	6,605	267	53	13	21	66	92	118	173	282	370	369
1999.....	5,222	190	49	20	17	38	53	78	128	199	217	232

Black, both sexes

2004.....	2,098	388	43	11	9	30	43	69	71	106	165	180
2003.....	2,446	352	64	11	24	39	79	91	105	154	228	264
2002.....	2,457	363	58	17	20	35	62	90	100	162	214	242
2001.....	2,535	345	57	14	16	39	73	89	121	185	213	269
2000.....	2,788	367	44	13	22	52	92	102	127	172	258	260
1999.....	2,452	306	58	13	13	40	74	77	112	172	227	233

Black, male

2004.....	1,176	235	23	4	4	18	29	39	46	55	94	118
2003.....	1,437	200	37	2	15	23	49	67	62	97	138	161
2002.....	1,469	209	37	9	10	22	37	51	67	109	135	164
2001.....	1,417	195	37	10	8	27	50	53	70	109	113	160
2000.....	1,613	219	24	7	16	27	57	66	82	90	148	162
1999.....	1,454	177	29	7	11	27	48	48	81	101	139	163

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other ill-defined and unknown causes of mortality (R96-R99)

Black, female

2004.....	922	153	20	7	5	12	14	30	25	51	71	62
2003.....	1,009	152	27	9	9	16	30	24	43	57	90	103
2002.....	988	154	21	8	10	13	25	39	33	53	79	78
2001.....	1,118	150	20	4	8	12	23	36	51	76	100	109
2000.....	1,175	148	20	6	6	25	35	36	45	82	110	98
1999.....	998	129	29	6	2	13	26	29	31	71	88	70

American Indian 1/, both sexes

2004.....	115	21	4	2	1	2	10	4	7	6	9	8
2003.....	112	20	2	1	2	3	3	5	3	11	9	11
2002.....	141	18	5	1	-	3	6	6	11	8	15	13
2001.....	153	26	6	1	3	3	5	5	11	12	13	9
2000.....	106	14	3	1	-	3	4	4	7	7	6	9
1999.....	124	23	3	-	1	1	3	7	4	3	7	9

American Indian 1/, male

2004.....	73	17	1	-	-	2	9	3	4	5	8	6
2003.....	69	8	2	-	2	2	3	3	3	9	6	9
2002.....	75	7	3	-	-	1	5	5	6	4	11	7
2001.....	101	15	5	1	-	2	4	3	10	10	9	6
2000.....	65	8	1	-	-	2	3	2	3	4	5	8
1999.....	70	10	1	-	-	1	3	3	1	2	4	7

American Indian 1/, female

2004.....	42	4	3	2	1	-	1	1	3	1	1	2
2003.....	43	12	-	1	-	1	-	2	-	2	3	2
2002.....	66	11	2	1	-	2	1	1	5	4	4	6
2001.....	52	11	1	-	3	1	1	2	1	2	4	3
2000.....	41	6	2	1	-	1	1	2	4	3	1	1
1999.....	54	13	2	-	1	-	-	4	3	1	3	2

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other ill-defined and unknown causes of mortality (R96-R99)												
Asian or Pacific Islander 2/, both sexes												
2004.....	181	29	5	1	2	2	8	9	7	9	13	11
2003.....	159	15	6	2	2	4	5	11	10	7	9	15
2002.....	168	20	8	1	3	2	11	11	5	10	9	13
2001.....	363	41	9	5	5	12	13	14	21	33	26	29
2000.....	253	29	5	6	4	6	9	20	14	22	17	23
1999.....	168	20	5	1	4	7	6	9	9	5	20	10
Asian or Pacific Islander 2/, male												
2004.....	105	17	2	1	-	1	4	5	3	5	11	6
2003.....	90	6	5	1	-	2	3	6	7	5	9	9
2002.....	105	15	5	1	1	1	7	9	5	8	6	10
2001.....	225	22	7	3	2	8	8	10	18	28	17	21
2000.....	152	15	1	2	1	4	4	13	11	20	14	15
1999.....	94	8	3	-	1	2	4	5	6	2	15	8
Asian or Pacific Islander 2/, female												
2004.....	76	12	3	-	2	1	4	4	4	4	2	5
2003.....	69	9	1	1	2	2	2	5	3	2	-	6
2002.....	63	5	3	-	2	1	4	2	-	2	3	3
2001.....	138	19	2	2	3	4	5	4	3	5	9	8
2000.....	101	14	4	4	3	2	5	7	3	2	3	8
1999.....	74	12	2	1	3	5	2	4	3	3	5	2

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

All other symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R53,R55-R94)

All races, both sexes

2004.....	9,775	71	32	15	31	43	100	113	122	138	198	244
2003.....	9,351	66	53	20	28	50	123	141	144	181	213	284
2002.....	8,506	77	47	16	23	58	92	122	170	192	228	244
2001.....	7,827	72	43	25	31	61	105	104	125	184	233	230
2000.....	7,446	67	58	25	23	58	102	133	138	183	239	231
1999.....	6,969	72	44	17	17	53	69	111	118	186	186	196

All races, male

2004.....	3,685	42	14	9	17	31	67	68	74	89	113	136
2003.....	3,580	45	31	10	18	32	76	100	84	118	118	173
2002.....	3,421	40	29	10	14	38	56	81	110	116	124	141
2001.....	3,147	48	23	13	20	45	63	63	72	116	149	147
2000.....	3,136	33	26	12	12	31	69	89	76	125	157	137
1999.....	2,877	45	26	14	10	34	38	67	68	119	118	130

All races, female

2004.....	6,090	29	18	6	14	12	33	45	48	49	85	108
2003.....	5,771	21	22	10	10	18	47	41	60	63	95	111
2002.....	5,085	37	18	6	9	20	36	41	60	76	104	103
2001.....	4,680	24	20	12	11	16	42	41	53	68	84	83
2000.....	4,310	34	32	13	11	27	33	44	62	58	82	94
1999.....	4,092	27	18	3	7	19	31	44	50	67	68	66

White, both sexes

2004.....	8,240	44	22	11	18	29	66	72	73	95	133	187
2003.....	7,882	55	35	16	14	40	88	95	102	122	157	184
2002.....	7,157	47	31	9	17	34	56	89	109	134	156	170
2001.....	6,574	52	28	19	24	42	67	68	94	124	161	172
2000.....	6,232	47	44	17	20	42	72	91	85	135	177	160
1999.....	5,796	47	33	15	10	36	41	70	80	125	126	129

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

All other symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R53,R55-R94)

White, male

2004.....	2,989	20	8	8	10	22	45	41	45	59	71	104
2003.....	2,920	35	23	9	8	25	55	70	61	78	85	113
2002.....	2,769	23	18	5	10	20	37	58	71	84	87	97
2001.....	2,548	35	16	8	16	31	38	47	52	77	102	108
2000.....	2,549	24	21	8	11	25	49	63	49	96	122	94
1999.....	2,330	29	19	12	8	21	23	47	40	88	78	84

White, female

2004.....	5,251	24	14	3	8	7	21	31	28	36	62	83
2003.....	4,962	20	12	7	6	15	33	25	41	44	72	71
2002.....	4,388	24	13	4	7	14	19	31	38	50	69	73
2001.....	4,026	17	12	11	8	11	29	21	42	47	59	64
2000.....	3,683	23	23	9	9	17	23	28	36	39	55	66
1999.....	3,466	18	14	3	2	15	18	23	40	37	48	45

Black, both sexes

2004.....	1,363	21	7	3	13	13	30	38	45	40	60	53
2003.....	1,314	8	16	3	13	8	28	41	34	49	56	90
2002.....	1,221	29	9	7	6	23	34	31	58	56	63	64
2001.....	1,136	14	12	6	6	16	38	34	26	56	69	57
2000.....	1,107	19	13	8	3	15	27	40	47	42	56	65
1999.....	1,087	22	10	2	5	14	26	39	38	59	56	66

Black, male

2004.....	616	18	5	1	7	9	20	25	27	28	39	30
2003.....	588	7	7	1	9	6	16	27	20	32	33	54
2002.....	590	17	8	5	4	17	17	22	37	31	34	38
2001.....	543	9	7	5	3	11	25	16	15	36	45	38
2000.....	530	9	5	4	1	5	19	25	25	25	30	38
1999.....	514	15	6	2	-	11	15	19	28	30	37	46

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

All other symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R53,R55-R94)

Black, female

2004.....	747	3	2	2	6	4	10	13	18	12	21	23
2003.....	726	1	9	2	4	2	12	14	14	17	23	36
2002.....	631	12	1	2	2	6	17	9	21	25	29	26
2001.....	593	5	5	1	3	5	13	18	11	20	24	19
2000.....	577	10	8	4	2	10	8	15	22	17	26	27
1999.....	573	7	4	-	5	3	11	20	10	29	19	20

American Indian 1/, both sexes

2004.....	47	2	1	-	-	1	3	2	3	3	4	2
2003.....	44	2	2	-	1	2	2	1	3	8	-	6
2002.....	45	1	2	-	-	-	-	2	1	1	7	4
2001.....	41	3	-	-	-	1	-	1	3	2	1	1
2000.....	42	-	-	-	-	1	1	1	5	5	3	4
1999.....	32	1	-	-	1	2	-	1	-	2	3	-

American Indian 1/, male

2004.....	24	1	-	-	-	-	2	1	1	2	2	2
2003.....	25	2	1	-	1	1	2	1	2	6	-	5
2002.....	20	-	-	-	-	-	-	1	1	-	3	2
2001.....	20	2	-	-	-	1	-	-	3	2	1	1
2000.....	24	-	-	-	-	1	-	-	2	3	2	4
1999.....	13	-	-	-	1	1	-	1	-	1	2	-

American Indian 1/, female

2004.....	23	1	1	-	-	1	1	1	2	1	2	-
2003.....	19	-	1	-	-	1	-	-	1	2	-	1
2002.....	25	1	2	-	-	-	-	1	-	1	4	2
2001.....	21	1	-	-	-	-	-	1	-	-	-	-
2000.....	18	-	-	-	-	-	1	1	3	2	1	-
1999.....	19	1	-	-	-	1	-	-	-	1	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

All other symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R53,R55-R94)

Asian or Pacific Islander 2/, both sexes

2004.....	125	4	2	1	-	-	1	1	1	-	1	2
2003.....	111	1	-	1	-	-	5	4	5	2	-	4
2002.....	83	-	5	-	-	1	2	-	2	1	2	6
2001.....	76	3	3	-	1	2	-	1	2	2	2	-
2000.....	65	1	1	-	-	-	2	1	1	1	3	2
1999.....	54	2	1	-	1	1	2	1	-	-	1	1

Asian or Pacific Islander 2/, male

2004.....	56	3	1	-	-	-	-	1	1	-	1	-
2003.....	47	1	-	-	-	-	3	2	1	2	-	1
2002.....	42	-	3	-	-	1	2	-	1	1	-	4
2001.....	36	2	-	-	1	2	-	-	2	1	1	-
2000.....	33	-	-	-	-	-	1	1	-	1	3	1
1999.....	20	1	1	-	1	1	-	-	-	-	1	-

Asian or Pacific Islander 2/, female

2004.....	69	1	1	1	-	-	1	-	-	-	-	2
2003.....	64	-	-	1	-	-	2	2	4	-	-	3
2002.....	41	-	2	-	-	-	-	-	1	-	2	2
2001.....	40	1	3	-	-	-	-	1	-	1	1	-
2000.....	32	1	1	-	-	-	1	-	1	-	-	1
1999.....	34	1	-	-	-	-	2	1	-	-	-	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
XX. External causes of mortality (*U01-*U03,V01-Y89)												
All races, both sexes												
2004.....	170,067	1,497	2,084	1,283	2,080	10,659	14,824	12,228	11,357	12,517	15,229	15,306
2003.....	166,857	1,376	2,148	1,246	2,015	10,390	14,939	11,566	11,595	12,916	15,439	14,520
2002.....	164,112	1,365	2,140	1,359	2,085	10,725	14,529	11,489	11,625	13,381	15,216	13,736
2001 3/.....	160,099	1,417	2,197	1,449	2,057	10,332	13,891	11,523	11,504	13,882	14,650	12,987
2000.....	151,268	1,314	2,243	1,574	2,177	10,447	13,074	10,953	10,646	13,039	13,691	11,625
1999.....	151,109	1,285	2,347	1,666	2,169	10,592	12,496	11,049	11,135	13,186	13,286	11,149
All races, male												
2004.....	116,113	842	1,204	767	1,336	7,875	12,011	9,790	8,682	9,218	10,790	10,792
2003.....	114,664	802	1,283	737	1,380	7,766	12,058	9,242	8,936	9,512	11,160	10,434
2002.....	113,045	777	1,297	806	1,370	8,006	11,802	9,215	8,956	9,896	10,976	9,922
2001 3/.....	110,873	802	1,316	880	1,376	7,832	11,370	9,216	8,893	10,259	10,778	9,630
2000.....	104,684	743	1,333	931	1,469	7,846	10,613	8,784	8,218	9,735	10,199	8,653
1999.....	104,192	745	1,354	1,004	1,462	7,879	10,111	8,787	8,551	9,811	9,846	8,307
All races, female												
2004.....	53,954	655	880	516	744	2,784	2,813	2,438	2,675	3,299	4,439	4,514
2003.....	52,193	574	865	509	635	2,624	2,881	2,324	2,659	3,404	4,279	4,086
2002.....	51,067	588	843	553	715	2,719	2,727	2,274	2,669	3,485	4,240	3,814
2001 3/.....	49,226	615	881	569	681	2,500	2,521	2,307	2,611	3,623	3,872	3,357
2000.....	46,584	571	910	643	708	2,601	2,461	2,169	2,428	3,304	3,492	2,972
1999.....	46,917	540	993	662	707	2,713	2,385	2,262	2,584	3,375	3,440	2,842
White, both sexes												
2004.....	140,606	993	1,489	884	1,523	8,263	11,170	9,075	8,721	10,095	12,728	12,802
2003.....	137,483	901	1,542	883	1,469	8,021	11,038	8,500	8,959	10,399	12,817	12,088
2002.....	134,847	817	1,497	919	1,526	8,354	10,728	8,379	8,968	10,655	12,653	11,382
2001 3/.....	131,209	905	1,584	1,025	1,520	7,895	10,186	8,528	8,925	11,166	12,016	10,631
2000.....	123,231	821	1,573	1,121	1,654	8,030	9,459	7,932	8,170	10,458	11,081	9,493
1999.....	122,886	833	1,626	1,135	1,658	8,024	9,026	8,102	8,578	10,546	10,798	9,047

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

XX. External causes of mortality (*U01-*U03,V01-Y89)

White, male

2004.....	94,701	574	871	522	987	5,963	8,952	7,168	6,602	7,446	9,015	9,019
2003.....	93,193	523	923	516	1,007	5,845	8,786	6,698	6,871	7,621	9,281	8,662
2002.....	91,608	492	901	543	997	6,113	8,597	6,620	6,876	7,845	9,194	8,241
2001 3/.....	89,740	511	970	598	1,008	5,846	8,230	6,765	6,882	8,293	8,898	7,890
2000.....	84,081	485	947	664	1,106	5,890	7,599	6,285	6,298	7,825	8,295	7,063
1999.....	83,621	498	943	676	1,110	5,810	7,216	6,415	6,573	7,851	8,047	6,721

White, female

2004.....	45,905	419	618	362	536	2,300	2,218	1,907	2,119	2,649	3,713	3,783
2003.....	44,290	378	619	367	462	2,176	2,252	1,802	2,088	2,778	3,536	3,426
2002.....	43,239	325	596	376	529	2,241	2,131	1,759	2,092	2,810	3,459	3,141
2001 3/.....	41,469	394	614	427	512	2,049	1,956	1,763	2,043	2,873	3,118	2,741
2000.....	39,150	336	626	457	548	2,140	1,860	1,647	1,872	2,633	2,786	2,430
1999.....	39,265	335	683	459	548	2,214	1,810	1,687	2,005	2,695	2,751	2,326

Black, both sexes

2004.....	24,057	440	505	329	450	1,936	3,083	2,680	2,170	1,984	2,052	2,094
2003.....	23,938	419	518	297	437	1,882	3,281	2,591	2,193	2,064	2,193	2,025
2002.....	24,061	491	533	367	455	1,901	3,229	2,652	2,203	2,274	2,135	1,990
2001 3/.....	23,898	449	528	358	435	1,963	3,159	2,508	2,129	2,255	2,241	1,986
2000.....	23,373	434	569	391	432	1,967	3,114	2,536	2,095	2,167	2,234	1,798
1999.....	23,598	406	618	466	429	2,121	2,958	2,479	2,127	2,218	2,138	1,804

Black, male

2004.....	17,841	230	282	200	283	1,582	2,634	2,283	1,752	1,476	1,465	1,483
2003.....	17,858	242	311	180	298	1,582	2,798	2,198	1,748	1,570	1,589	1,492
2002.....	17,957	253	336	219	319	1,555	2,759	2,250	1,762	1,726	1,485	1,447
2001 3/.....	17,754	258	305	243	309	1,640	2,728	2,087	1,686	1,626	1,612	1,501
2000.....	17,439	226	324	234	308	1,620	2,634	2,126	1,653	1,635	1,646	1,355
1999.....	17,484	219	350	296	297	1,738	2,522	2,045	1,670	1,656	1,579	1,385

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

XX. External causes of mortality (*U01-*U03,V01-Y89)

Black, female

2004.....	6,216	210	223	129	167	354	449	397	418	508	587	611
2003.....	6,080	177	207	117	139	300	483	393	445	494	604	533
2002.....	6,104	238	197	148	136	346	470	402	441	548	650	543
2001 3/.....	6,144	191	223	115	126	323	431	421	443	629	629	485
2000.....	5,934	208	245	157	124	347	480	410	442	532	588	443
1999.....	6,114	187	268	170	132	383	436	434	457	562	559	419

American Indian 1/, both sexes

2004.....	2,236	37	49	25	54	223	268	221	214	241	211	186
2003.....	2,232	33	40	31	50	237	271	220	199	237	222	174
2002.....	2,162	28	47	28	51	224	255	214	203	241	223	169
2001 3/.....	1,964	39	43	31	59	229	220	169	186	224	173	150
2000.....	1,917	29	48	33	47	220	225	198	188	197	186	137
1999.....	1,955	19	45	30	32	214	240	211	222	203	165	135

American Indian 1/, male

2004.....	1,549	24	31	14	30	151	203	166	151	171	147	133
2003.....	1,552	23	24	21	37	164	213	158	149	175	155	124
2002.....	1,493	13	25	12	27	154	201	168	149	170	154	113
2001 3/.....	1,360	17	20	21	34	161	165	139	134	174	118	96
2000.....	1,340	17	32	18	27	158	165	152	134	133	127	101
1999.....	1,362	12	28	13	21	164	182	140	161	148	102	105

American Indian 1/, female

2004.....	687	13	18	11	24	72	65	55	63	70	64	53
2003.....	680	10	16	10	13	73	58	62	50	62	67	50
2002.....	669	15	22	16	24	70	54	46	54	71	69	56
2001 3/.....	604	22	23	10	25	68	55	30	52	50	55	54
2000.....	577	12	16	15	20	62	60	46	54	64	59	36
1999.....	593	7	17	17	11	50	58	71	61	55	63	30

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

XX. External causes of mortality (*U01-*U03,V01-Y89)

Asian or Pacific Islander 2/, both sexes

2004.....	3,168	27	41	45	53	237	303	252	252	197	238	224
2003.....	3,204	23	48	35	59	250	349	255	244	216	207	233
2002.....	3,042	29	63	45	53	246	317	244	251	211	205	195
2001 3/.....	3,028	24	42	35	43	245	326	318	264	237	220	220
2000.....	2,747	30	53	29	44	230	276	287	193	217	190	197
1999.....	2,670	27	58	35	50	233	272	257	208	219	185	163

Asian or Pacific Islander 2/, male

2004.....	2,022	14	20	31	36	179	222	173	177	125	163	157
2003.....	2,061	14	25	20	38	175	261	188	168	146	135	156
2002.....	1,987	19	35	32	27	184	245	177	169	155	143	121
2001 3/.....	2,019	16	21	18	25	185	247	225	191	166	150	143
2000.....	1,824	15	30	15	28	178	215	221	133	142	131	134
1999.....	1,725	16	33	19	34	167	191	187	147	156	118	96

Asian or Pacific Islander 2/, female

2004.....	1,146	13	21	14	17	58	81	79	75	72	75	67
2003.....	1,143	9	23	15	21	75	88	67	76	70	72	77
2002.....	1,055	10	28	13	26	62	72	67	82	56	62	74
2001 3/.....	1,009	8	21	17	18	60	79	93	73	71	70	77
2000.....	923	15	23	14	16	52	61	66	60	75	59	63
1999.....	945	11	25	16	16	66	81	70	61	63	67	67

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Accidents (unintentional injuries) (V01-X59,Y85-Y86)												
All races, both sexes												
2004.....	112,012	1,052	1,641	1,126	1,540	6,825	8,624	6,771	6,261	7,319	9,152	9,426
2003.....	109,277	945	1,717	1,096	1,522	6,755	8,517	6,284	6,257	7,495	9,271	8,865
2002.....	106,742	946	1,641	1,176	1,542	7,137	8,275	6,122	6,447	7,642	9,068	8,241
2001.....	101,537	976	1,714	1,283	1,553	6,646	7,765	5,935	5,904	7,589	8,356	7,469
2000.....	97,900	881	1,826	1,391	1,588	6,755	7,358	5,898	5,871	7,471	7,942	6,870
1999.....	97,860	845	1,898	1,459	1,632	6,688	6,968	5,881	6,009	7,366	7,865	6,696
All races, male												
2004.....	72,050	604	964	682	979	4,744	6,679	5,222	4,649	5,252	6,419	6,602
2003.....	70,532	539	1,048	650	1,012	4,706	6,585	4,826	4,708	5,390	6,613	6,325
2002.....	69,257	553	1,024	720	993	4,981	6,457	4,768	4,867	5,535	6,477	5,940
2001.....	66,060	541	1,045	786	1,017	4,722	6,110	4,605	4,487	5,502	6,110	5,519
2000.....	63,817	501	1,098	824	1,037	4,757	5,703	4,606	4,453	5,537	5,907	5,103
1999.....	63,535	494	1,127	897	1,077	4,624	5,416	4,555	4,542	5,411	5,837	4,996
All races, female												
2004.....	39,962	448	677	444	561	2,081	1,945	1,549	1,612	2,067	2,733	2,824
2003.....	38,745	406	669	446	510	2,049	1,932	1,458	1,549	2,105	2,658	2,540
2002.....	37,485	393	617	456	549	2,156	1,818	1,354	1,580	2,107	2,591	2,301
2001.....	35,477	435	669	497	536	1,924	1,655	1,330	1,417	2,087	2,246	1,950
2000.....	34,083	380	728	567	551	1,998	1,655	1,292	1,418	1,934	2,035	1,767
1999.....	34,325	351	771	562	555	2,064	1,552	1,326	1,467	1,955	2,028	1,700
White, both sexes												
2004.....	95,890	706	1,242	792	1,161	5,850	7,324	5,617	5,170	6,090	7,751	7,872
2003.....	93,381	624	1,304	805	1,149	5,771	7,219	5,217	5,183	6,226	7,790	7,353
2002.....	90,866	583	1,199	810	1,155	6,072	6,976	5,004	5,374	6,270	7,631	6,784
2001.....	85,964	647	1,309	906	1,154	5,563	6,475	4,877	4,850	6,269	6,874	6,011
2000.....	82,592	569	1,335	996	1,223	5,731	6,092	4,736	4,783	6,171	6,492	5,554
1999.....	82,245	562	1,374	1,011	1,259	5,647	5,717	4,790	4,871	6,051	6,449	5,369

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Accidents (unintentional injuries) (V01-X59,Y85-Y86)

White, male

2004.....	61,267	414	734	474	749	4,045	5,725	4,343	3,837	4,403	5,462	5,542
2003.....	59,912	359	807	472	761	4,005	5,591	4,041	3,923	4,480	5,608	5,247
2002.....	58,467	358	736	497	743	4,214	5,465	3,907	4,079	4,546	5,494	4,949
2001.....	55,493	356	815	536	749	3,930	5,137	3,803	3,715	4,571	5,076	4,443
2000.....	53,329	331	808	587	791	4,007	4,763	3,704	3,665	4,608	4,854	4,143
1999.....	52,898	333	811	611	823	3,862	4,465	3,741	3,701	4,477	4,811	4,002

White, female

2004.....	34,623	292	508	318	412	1,805	1,599	1,274	1,333	1,687	2,289	2,330
2003.....	33,469	265	497	333	388	1,766	1,628	1,176	1,260	1,746	2,182	2,106
2002.....	32,399	225	463	313	412	1,858	1,511	1,097	1,295	1,724	2,137	1,835
2001.....	30,471	291	494	370	405	1,633	1,338	1,074	1,135	1,698	1,798	1,568
2000.....	29,263	238	527	409	432	1,724	1,329	1,032	1,118	1,563	1,638	1,411
1999.....	29,347	229	563	400	436	1,785	1,252	1,049	1,170	1,574	1,638	1,367

Black, both sexes

2004.....	12,670	303	324	269	301	712	949	890	850	973	1,131	1,324
2003.....	12,351	287	343	235	288	667	951	800	835	1,010	1,214	1,263
2002.....	12,513	330	359	306	315	758	968	867	844	1,105	1,163	1,232
2001.....	12,462	281	339	319	325	775	967	806	832	1,049	1,272	1,233
2000.....	12,277	274	405	338	293	737	958	892	878	1,058	1,229	1,109
1999.....	12,728	252	441	396	311	774	945	855	910	1,081	1,204	1,153

Black, male

2004.....	8,603	163	188	168	183	520	708	689	644	679	778	908
2003.....	8,385	160	204	142	192	490	737	601	620	735	830	907
2002.....	8,612	176	239	185	215	558	740	677	632	803	789	853
2001.....	8,537	162	194	216	226	576	733	609	611	735	898	934
2000.....	8,531	149	239	209	205	542	719	693	647	772	906	817
1999.....	8,773	141	266	258	214	574	727	658	677	769	891	877

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Accidents (unintentional injuries) (V01-X59,Y85-Y86)

Black, female

2004.....	4,067	140	136	101	118	192	241	201	206	294	353	416
2003.....	3,966	127	139	93	96	177	214	199	215	275	384	356
2002.....	3,901	154	120	121	100	200	228	190	212	302	374	379
2001.....	3,925	119	145	103	99	199	234	197	221	314	374	299
2000.....	3,746	125	166	129	88	195	239	199	231	286	323	292
1999.....	3,955	111	175	138	97	200	218	197	233	312	313	276

American Indian 1/, both sexes

2004.....	1,520	23	42	22	39	122	173	127	135	149	141	130
2003.....	1,573	20	33	28	40	153	162	137	127	157	156	127
2002.....	1,488	15	33	23	37	146	154	132	122	166	167	117
2001.....	1,361	32	30	29	44	145	148	102	114	152	108	113
2000.....	1,353	20	39	31	39	149	150	123	116	129	127	108
1999.....	1,327	13	37	25	26	130	159	119	131	129	123	96

American Indian 1/, male

2004.....	1,010	15	26	11	22	78	125	94	90	99	92	87
2003.....	1,043	14	19	19	29	102	126	88	89	111	104	89
2002.....	1,003	8	19	11	20	91	121	101	85	114	114	78
2001.....	908	14	15	20	26	95	108	80	83	117	71	70
2000.....	900	11	25	16	21	102	100	94	78	81	85	80
1999.....	890	8	24	12	16	95	119	72	91	90	77	74

American Indian 1/, female

2004.....	510	8	16	11	17	44	48	33	45	50	49	43
2003.....	530	6	14	9	11	51	36	49	38	46	52	38
2002.....	485	7	14	12	17	55	33	31	37	52	53	39
2001.....	453	18	15	9	18	50	40	22	31	35	37	43
2000.....	453	9	14	15	18	47	50	29	38	48	42	28
1999.....	437	5	13	13	10	35	40	47	40	39	46	22

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Accidents (unintentional injuries) (V01-X59,Y85-Y86)

Asian or Pacific Islander 2/, both sexes

2004.....	1,932	20	33	43	39	141	178	137	106	107	129	100
2003.....	1,972	14	37	28	45	164	185	130	112	102	111	122
2002.....	1,875	18	50	37	35	161	177	119	107	101	107	108
2001.....	1,750	16	36	29	30	163	175	150	108	119	102	112
2000.....	1,678	18	47	26	33	138	158	147	94	113	94	99
1999.....	1,560	18	46	27	36	137	147	117	97	105	89	78

Asian or Pacific Islander 2/, male

2004.....	1,170	12	16	29	25	101	121	96	78	71	87	65
2003.....	1,192	6	18	17	30	109	131	96	76	64	71	82
2002.....	1,175	11	30	27	15	118	131	83	71	72	80	60
2001.....	1,122	9	21	14	16	121	132	113	78	79	65	72
2000.....	1,057	10	26	12	20	106	121	115	63	76	62	63
1999.....	974	12	26	16	24	93	105	84	73	75	58	43

Asian or Pacific Islander 2/, female

2004.....	762	8	17	14	14	40	57	41	28	36	42	35
2003.....	780	8	19	11	15	55	54	34	36	38	40	40
2002.....	700	7	20	10	20	43	46	36	36	29	27	48
2001.....	628	7	15	15	14	42	43	37	30	40	37	40
2000.....	621	8	21	14	13	32	37	32	31	37	32	36
1999.....	586	6	20	11	12	44	42	33	24	30	31	35

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Transport accidents (V01-V99,Y85)

All races, both sexes

2004.....	48,053	145	659	681	1,074	5,394	5,983	4,094	3,427	3,460	3,787	3,775
2003.....	48,071	148	649	694	1,071	5,471	5,961	3,855	3,425	3,579	4,015	3,656
2002.....	48,366	127	640	705	1,033	5,782	6,034	3,891	3,593	3,750	3,938	3,593
2001.....	47,288	151	678	769	1,054	5,393	5,746	3,953	3,501	3,881	3,946	3,379
2000.....	46,749	174	691	842	1,080	5,475	5,553	3,957	3,443	3,889	3,772	3,248
1999.....	46,423	189	712	897	1,078	5,452	5,237	3,922	3,512	3,869	3,684	3,121

All races, male

2004.....	33,311	81	359	392	647	3,577	4,536	3,117	2,524	2,539	2,758	2,717
2003.....	33,291	81	366	401	683	3,653	4,518	2,915	2,596	2,585	2,912	2,667
2002.....	33,533	72	381	405	634	3,875	4,596	2,990	2,686	2,703	2,796	2,638
2001.....	32,794	68	392	447	668	3,694	4,417	3,015	2,646	2,744	2,869	2,452
2000.....	32,166	94	391	464	674	3,693	4,197	3,016	2,554	2,802	2,758	2,338
1999.....	31,642	101	400	532	674	3,593	3,965	2,952	2,559	2,741	2,635	2,234

All races, female

2004.....	14,742	64	300	289	427	1,817	1,447	977	903	921	1,029	1,058
2003.....	14,780	67	283	293	388	1,818	1,443	940	829	994	1,103	989
2002.....	14,833	55	259	300	399	1,907	1,438	901	907	1,047	1,142	955
2001.....	14,494	83	286	322	386	1,699	1,329	938	855	1,137	1,077	927
2000.....	14,583	80	300	378	406	1,782	1,356	941	889	1,087	1,014	910
1999.....	14,781	88	312	365	404	1,859	1,272	970	953	1,128	1,049	887

White, both sexes

2004.....	40,279	104	499	520	857	4,647	5,001	3,284	2,738	2,818	3,147	3,139
2003.....	40,282	111	491	538	859	4,686	4,957	3,125	2,760	2,925	3,328	3,043
2002.....	40,614	84	478	525	822	4,968	5,023	3,101	2,936	3,032	3,290	2,982
2001.....	39,544	115	511	558	820	4,563	4,722	3,227	2,826	3,179	3,272	2,743
2000.....	39,088	121	513	646	873	4,713	4,566	3,121	2,769	3,208	3,110	2,679
1999.....	38,767	137	497	664	878	4,661	4,292	3,171	2,855	3,140	3,059	2,563

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Transport accidents (V01-V99,Y85)												
White, male												
2004.....	27,918	54	269	301	528	3,067	3,834	2,505	2,005	2,083	2,305	2,273
2003.....	27,925	63	285	305	546	3,114	3,753	2,383	2,106	2,110	2,430	2,225
2002.....	28,118	50	278	305	509	3,316	3,845	2,388	2,202	2,189	2,351	2,220
2001.....	27,407	52	299	320	515	3,107	3,667	2,473	2,147	2,249	2,383	1,992
2000.....	26,815	68	290	358	538	3,158	3,474	2,377	2,064	2,312	2,277	1,945
1999.....	26,385	80	273	390	546	3,047	3,269	2,401	2,076	2,232	2,196	1,837
White, female												
2004.....	12,361	50	230	219	329	1,580	1,167	779	733	735	842	866
2003.....	12,357	48	206	233	313	1,572	1,204	742	654	815	898	818
2002.....	12,496	34	200	220	313	1,652	1,178	713	734	843	939	762
2001.....	12,137	63	212	238	305	1,456	1,055	754	679	930	889	751
2000.....	12,273	53	223	288	335	1,555	1,092	744	705	896	833	734
1999.....	12,382	57	224	274	332	1,614	1,023	770	779	908	863	726
Black, both sexes												
2004.....	5,868	32	132	126	169	546	714	618	536	492	490	516
2003.....	5,771	28	129	117	161	521	721	546	510	508	546	476
2002.....	5,796	36	126	151	161	569	745	608	504	557	497	482
2001.....	5,906	19	130	179	184	590	757	556	524	531	556	504
2000.....	5,839	40	141	157	160	539	744	634	535	527	542	457
1999.....	5,901	44	177	202	162	580	704	578	511	587	498	449
Black, male												
2004.....	4,204	20	71	70	95	383	524	481	415	367	355	372
2003.....	4,119	12	66	73	102	371	562	410	389	381	387	357
2002.....	4,156	18	83	83	100	395	554	473	383	406	343	345
2001.....	4,214	7	72	113	127	423	563	419	394	375	413	387
2000.....	4,199	21	79	87	110	380	548	489	401	396	408	321
1999.....	4,162	18	101	127	105	405	521	443	385	412	366	331

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Transport accidents (V01-V99,Y85)												
Black, female												
2004.....	1,664	12	61	56	74	163	190	137	121	125	135	144
2003.....	1,652	16	63	44	59	150	159	136	121	127	159	119
2002.....	1,640	18	43	68	61	174	191	135	121	151	154	137
2001.....	1,692	12	58	66	57	167	194	137	130	156	143	117
2000.....	1,640	19	62	70	50	159	196	145	134	131	134	136
1999.....	1,739	26	76	75	57	175	183	135	126	175	132	118
American Indian 1/, both sexes												
2004.....	851	5	20	13	22	97	129	94	80	77	67	62
2003.....	915	7	18	20	26	126	137	98	80	82	72	67
2002.....	903	4	16	12	27	123	127	89	82	94	89	56
2001.....	832	11	17	16	32	121	126	73	74	91	55	61
2000.....	829	7	16	19	27	120	121	94	73	80	66	53
1999.....	816	4	15	16	19	107	126	87	76	76	72	52
American Indian 1/, male												
2004.....	569	4	12	6	10	60	88	67	57	45	48	40
2003.....	595	5	8	11	20	82	103	64	57	55	52	41
2002.....	603	1	9	6	14	74	95	66	56	63	60	36
2001.....	558	5	8	8	18	78	86	56	55	73	39	33
2000.....	537	2	10	9	14	80	82	69	48	50	42	38
1999.....	547	2	12	8	13	77	91	50	50	51	45	39
American Indian 1/, female												
2004.....	282	1	8	7	12	37	41	27	23	32	19	22
2003.....	320	2	10	9	6	44	34	34	23	27	20	26
2002.....	300	3	7	6	13	49	32	23	26	31	29	20
2001.....	274	6	9	8	14	43	40	17	19	18	16	28
2000.....	292	5	6	10	13	40	39	25	25	30	24	15
1999.....	269	2	3	8	6	30	35	37	26	25	27	13

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Transport accidents (V01-V99,Y85)												
Asian or Pacific Islander 2/, both sexes												
2004.....	1,055	4	8	22	26	104	139	98	73	73	83	58
2003.....	1,103	2	11	19	25	138	146	86	75	64	69	70
2002.....	1,053	3	20	17	23	122	139	93	71	67	62	73
2001.....	1,006	6	20	16	18	119	141	97	77	80	63	71
2000.....	993	6	21	20	20	103	122	108	66	74	54	59
1999.....	939	4	23	15	19	104	115	86	70	66	55	57
Asian or Pacific Islander 2/, male												
2004.....	620	3	7	15	14	67	90	64	47	44	50	32
2003.....	652	1	7	12	15	86	100	58	44	39	43	44
2002.....	656	3	11	11	11	90	102	63	45	45	42	37
2001.....	615	4	13	6	8	86	101	67	50	47	34	40
2000.....	615	3	12	10	12	75	93	81	41	44	31	34
1999.....	548	1	14	7	10	64	84	58	48	46	28	27
Asian or Pacific Islander 2/, female												
2004.....	435	1	1	7	12	37	49	34	26	29	33	26
2003.....	451	1	4	7	10	52	46	28	31	25	26	26
2002.....	397	-	9	6	12	32	37	30	26	22	20	36
2001.....	391	2	7	10	10	33	40	30	27	33	29	31
2000.....	378	3	9	10	8	28	29	27	25	30	23	25
1999.....	391	3	9	8	9	40	31	28	22	20	27	30

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Railway accidents (V05,V15,V80.6,V81.2-V81.9)												
All races, both sexes												
2004.....	457	-	-	2	11	35	32	47	47	43	52	65
2003.....	426	-	2	2	14	30	38	34	42	58	49	59
2002.....	473	-	2	4	13	39	51	37	46	51	71	52
2001.....	491	-	2	1	18	38	42	46	45	60	61	52
2000.....	482	1	5	2	14	47	46	38	42	71	67	47
1999.....	493	1	2	8	17	35	57	46	50	74	66	44
All races, male												
2004.....	387	-	-	1	8	29	31	40	41	35	46	55
2003.....	364	-	1	2	11	27	34	29	39	50	42	51
2002.....	409	-	1	2	10	29	47	31	42	41	65	49
2001.....	416	-	2	1	13	31	34	42	42	53	53	46
2000.....	417	1	4	2	10	42	39	35	34	61	62	41
1999.....	423	1	1	7	15	29	54	39	46	65	58	35
All races, female												
2004.....	70	-	-	1	3	6	1	7	6	8	6	10
2003.....	62	-	1	-	3	3	4	5	3	8	7	8
2002.....	64	-	1	2	3	10	4	6	4	10	6	3
2001.....	75	-	-	-	5	7	8	4	3	7	8	6
2000.....	65	-	1	-	4	5	7	3	8	10	5	6
1999.....	70	-	1	1	2	6	3	7	4	9	8	9
White, both sexes												
2004.....	346	-	-	2	10	22	26	37	35	35	46	44
2003.....	348	-	2	1	12	23	32	29	36	50	38	42
2002.....	362	-	1	3	9	32	45	30	38	42	50	34
2001.....	388	-	1	1	12	28	38	41	33	44	49	37
2000.....	375	1	2	2	9	39	36	34	30	57	50	35
1999.....	389	-	2	5	14	27	43	37	40	58	56	33

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Railway accidents (V05,V15,V80.6,V81.2-V81.9)												
White, male												
2004.....	297	-	-	1	7	18	25	32	31	29	41	38
2003.....	298	-	1	1	9	20	29	24	34	43	35	36
2002.....	313	-	-	2	7	24	43	24	34	35	46	31
2001.....	327	-	1	1	8	23	31	39	32	38	42	32
2000.....	326	1	1	2	6	35	30	31	25	50	46	31
1999.....	334	-	1	5	12	23	40	30	38	51	50	27
White, female												
2004.....	49	-	-	1	3	4	1	5	4	6	5	6
2003.....	50	-	1	-	3	3	3	5	2	7	3	6
2002.....	49	-	1	1	2	8	2	6	4	7	4	3
2001.....	61	-	-	-	4	5	7	2	1	6	7	5
2000.....	49	-	1	-	3	4	6	3	5	7	4	4
1999.....	55	-	1	-	2	4	3	7	2	7	6	6
Black, both sexes												
2004.....	88	-	-	-	1	12	6	8	7	6	5	16
2003.....	68	-	-	1	2	6	5	4	4	6	10	16
2002.....	94	-	1	1	3	7	5	7	6	7	16	18
2001.....	82	-	1	-	6	8	2	3	8	11	12	12
2000.....	92	-	2	-	3	7	10	4	11	12	14	11
1999.....	76	1	-	2	2	4	6	7	7	13	8	10
Black, male												
2004.....	72	-	-	-	1	10	6	7	5	5	4	13
2003.....	59	-	-	1	2	6	4	4	4	6	7	14
2002.....	83	-	1	-	3	5	3	7	6	5	14	18
2001.....	71	-	1	-	5	6	1	2	7	11	11	11
2000.....	80	-	2	-	3	6	9	4	8	10	14	10
1999.....	64	1	-	2	2	3	6	7	6	11	6	7

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Railway accidents (V05,V15,V80.6,V81.2-V81.9)												
Black, female												
2004.....	16	-	-	-	-	2	-	1	2	1	1	3
2003.....	9	-	-	-	-	-	1	-	-	-	3	2
2002.....	11	-	-	1	-	2	2	-	-	2	2	-
2001.....	11	-	-	-	1	2	1	1	1	-	1	1
2000.....	12	-	-	-	-	1	1	-	3	2	-	1
1999.....	12	-	-	-	-	1	-	-	1	2	2	3
American Indian 1/, both sexes												
2004.....	19	-	-	-	-	1	-	1	5	1	1	4
2003.....	9	-	-	-	-	1	1	1	2	2	1	-
2002.....	9	-	-	-	-	-	1	-	2	2	3	-
2001.....	14	-	-	-	-	1	1	2	3	3	-	2
2000.....	11	-	-	-	1	1	-	-	1	2	2	1
1999.....	20	-	-	1	1	3	5	1	3	3	2	1
American Indian 1/, male												
2004.....	15	-	-	-	-	1	-	1	5	-	1	3
2003.....	6	-	-	-	-	1	1	1	1	1	-	-
2002.....	8	-	-	-	-	-	1	-	2	1	3	-
2001.....	13	-	-	-	-	1	1	1	3	3	-	2
2000.....	7	-	-	-	-	1	-	-	1	1	1	-
1999.....	17	-	-	-	1	2	5	1	2	3	2	1
American Indian 1/, female												
2004.....	4	-	-	-	-	-	-	-	-	1	-	1
2003.....	3	-	-	-	-	-	-	-	1	1	1	-
2002.....	1	-	-	-	-	-	-	-	-	1	-	-
2001.....	1	-	-	-	-	-	-	1	-	-	-	-
2000.....	4	-	-	-	1	-	-	-	-	1	1	1
1999.....	3	-	-	1	-	1	-	-	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)												
All races, both sexes												
2004.....	44,933	143	635	640	1,013	5,224	5,763	3,849	3,187	3,190	3,473	3,422
2003.....	44,757	145	621	646	996	5,269	5,703	3,644	3,189	3,289	3,672	3,286
2002.....	45,380	123	610	664	950	5,634	5,825	3,731	3,361	3,462	3,615	3,240
2001.....	43,788	144	626	718	968	5,208	5,517	3,710	3,227	3,545	3,538	3,011
2000.....	43,354	168	651	780	992	5,251	5,309	3,715	3,169	3,545	3,382	2,936
1999.....	42,401	184	650	802	969	5,198	4,930	3,604	3,174	3,460	3,278	2,747
All races, male												
2004.....	30,837	80	344	369	606	3,453	4,361	2,924	2,321	2,318	2,508	2,431
2003.....	30,655	80	353	373	636	3,500	4,306	2,753	2,403	2,350	2,634	2,366
2002.....	31,064	68	361	385	578	3,765	4,413	2,855	2,494	2,459	2,525	2,328
2001.....	30,064	66	356	418	611	3,546	4,232	2,826	2,410	2,473	2,543	2,164
2000.....	29,451	90	368	429	614	3,527	3,989	2,804	2,331	2,507	2,433	2,091
1999.....	28,552	97	366	469	593	3,412	3,715	2,690	2,303	2,424	2,311	1,936
All races, female												
2004.....	14,096	63	291	271	407	1,771	1,402	925	866	872	965	991
2003.....	14,102	65	268	273	360	1,769	1,397	891	786	939	1,038	920
2002.....	14,316	55	249	279	372	1,869	1,412	876	867	1,003	1,090	912
2001.....	13,724	78	270	300	357	1,662	1,285	884	817	1,072	995	847
2000.....	13,903	78	283	351	378	1,724	1,320	911	838	1,038	949	845
1999.....	13,849	87	284	333	376	1,786	1,215	914	871	1,036	967	811
White, both sexes												
2004.....	37,618	103	480	487	805	4,508	4,817	3,083	2,534	2,585	2,881	2,843
2003.....	37,424	108	467	501	793	4,517	4,740	2,942	2,565	2,675	3,030	2,735
2002.....	38,054	81	452	494	760	4,840	4,840	2,963	2,734	2,787	3,020	2,683
2001.....	36,576	109	466	522	753	4,409	4,520	3,014	2,599	2,904	2,924	2,447
2000.....	36,160	115	484	593	796	4,530	4,351	2,908	2,540	2,918	2,772	2,421
1999.....	35,343	133	455	591	786	4,450	4,035	2,897	2,572	2,791	2,706	2,244

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)												
White, male												
2004.....	25,816	54	258	283	492	2,969	3,693	2,349	1,835	1,894	2,095	2,029
2003.....	25,660	62	273	286	508	2,986	3,576	2,242	1,947	1,909	2,187	1,975
2002.....	26,000	47	260	288	467	3,221	3,683	2,273	2,033	1,979	2,125	1,955
2001.....	25,084	50	268	298	472	2,982	3,506	2,306	1,949	2,026	2,107	1,756
2000.....	24,470	64	275	330	485	3,026	3,291	2,191	1,876	2,061	1,996	1,739
1999.....	23,763	77	253	344	479	2,899	3,060	2,175	1,863	1,961	1,918	1,583
White, female												
2004.....	11,802	49	222	204	313	1,539	1,124	734	699	691	786	814
2003.....	11,764	46	194	215	285	1,531	1,164	700	618	766	843	760
2002.....	12,054	34	192	206	293	1,619	1,157	690	701	808	895	728
2001.....	11,492	59	198	224	281	1,427	1,014	708	650	878	817	691
2000.....	11,690	51	209	263	311	1,504	1,060	717	664	857	776	682
1999.....	11,580	56	202	247	307	1,551	975	722	709	830	788	661
Black, both sexes												
2004.....	5,513	31	128	121	161	520	685	585	510	464	454	468
2003.....	5,431	28	125	110	153	497	692	524	483	476	511	428
2002.....	5,462	35	123	141	146	553	726	591	481	519	461	432
2001.....	5,506	19	123	165	166	568	740	533	492	491	506	447
2000.....	5,499	40	134	149	153	513	721	614	497	485	502	415
1999.....	5,470	43	160	184	150	553	677	547	471	538	457	408
Black, male												
2004.....	3,910	19	68	67	91	361	497	451	392	343	324	336
2003.....	3,834	12	65	67	94	352	535	393	366	353	361	317
2002.....	3,874	17	81	80	89	382	539	457	364	376	313	303
2001.....	3,900	7	67	106	113	406	547	402	369	343	370	342
2000.....	3,922	21	74	80	105	358	528	470	372	361	374	288
1999.....	3,821	17	89	111	94	384	501	418	353	374	330	299

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)

Black, female

2004.....	1,603	12	60	54	70	159	188	134	118	121	130	132
2003.....	1,597	16	60	43	59	145	157	131	117	123	150	111
2002.....	1,588	18	42	61	57	171	187	134	117	143	148	129
2001.....	1,606	12	56	59	53	162	193	131	123	148	136	105
2000.....	1,577	19	60	69	48	155	193	144	125	124	128	127
1999.....	1,649	26	71	73	56	169	176	129	118	164	127	109

American Indian 1/, both sexes

2004.....	802	5	19	12	22	94	125	89	74	75	63	58
2003.....	865	7	18	19	25	121	133	95	69	77	68	63
2002.....	866	4	16	12	26	122	121	87	78	91	77	54
2001.....	764	10	17	16	31	116	121	69	65	79	50	53
2000.....	779	7	14	19	25	113	118	92	71	74	61	45
1999.....	730	4	14	14	16	98	112	82	68	70	63	42

American Indian 1/, male

2004.....	529	4	11	5	10	58	84	63	51	44	44	37
2003.....	557	5	8	11	19	78	99	61	48	52	49	38
2002.....	572	1	9	6	13	74	90	65	53	61	49	34
2001.....	506	5	8	8	18	74	82	53	47	63	35	30
2000.....	500	2	9	9	14	75	79	67	47	45	38	32
1999.....	481	2	11	7	12	70	79	45	45	46	38	30

American Indian 1/, female

2004.....	273	1	8	7	12	36	41	26	23	31	19	21
2003.....	308	2	10	8	6	43	34	34	21	25	19	25
2002.....	294	3	7	6	13	48	31	22	25	30	28	20
2001.....	258	5	9	8	13	42	39	16	18	16	15	23
2000.....	279	5	5	10	11	38	39	25	24	29	23	13
1999.....	249	2	3	7	4	28	33	37	23	24	25	12

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)

Asian or Pacific Islander 2/, both sexes

2004.....	1,000	4	8	20	25	102	136	92	69	66	75	53
2003.....	1,037	2	11	16	25	134	138	83	72	61	63	60
2002.....	998	3	19	17	18	119	138	90	68	65	57	71
2001.....	942	6	20	15	18	115	136	94	71	71	58	64
2000.....	916	6	19	19	18	95	119	101	61	68	47	55
1999.....	858	4	21	13	17	97	106	78	63	61	52	53

Asian or Pacific Islander 2/, male

2004.....	582	3	7	14	13	65	87	61	43	37	45	29
2003.....	604	1	7	9	15	84	96	57	42	36	37	36
2002.....	618	3	11	11	9	88	101	60	44	43	38	36
2001.....	574	4	13	6	8	84	97	65	45	41	31	36
2000.....	559	3	10	10	10	68	91	76	36	40	25	32
1999.....	487	1	13	7	8	59	75	52	42	43	25	24

Asian or Pacific Islander 2/, female

2004.....	418	1	1	6	12	37	49	31	26	29	30	24
2003.....	433	1	4	7	10	50	42	26	30	25	26	24
2002.....	380	-	8	6	9	31	37	30	24	22	19	35
2001.....	368	2	7	9	10	31	39	29	26	30	27	28
2000.....	357	3	9	9	8	27	28	25	25	28	22	23
1999.....	371	3	8	6	9	38	31	26	21	18	27	29

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pedestrian involved in collision with motor vehicle (V02-V04)												
All races, both sexes												
2004.....	3,364	5	158	103	99	157	210	186	188	236	283	302
2003.....	3,561	10	179	106	112	203	232	203	193	248	320	324
2002.....	3,624	7	155	119	109	201	226	197	220	283	326	304
2001.....	3,557	8	170	121	148	176	222	184	204	301	331	296
2000.....	3,434	9	179	146	114	162	186	187	200	272	312	276
1999.....	3,389	9	170	160	111	167	197	195	205	289	284	257
All races, male												
2004.....	2,314	4	99	62	66	118	159	142	141	164	214	215
2003.....	2,457	4	116	69	72	142	175	155	147	181	237	228
2002.....	2,472	4	106	70	67	144	180	152	167	200	227	231
2001.....	2,469	5	112	82	110	125	178	153	152	209	249	208
2000.....	2,340	4	111	88	73	107	146	148	154	194	227	216
1999.....	2,337	5	96	93	71	122	153	148	159	220	209	193
All races, female												
2004.....	1,050	1	59	41	33	39	51	44	47	72	69	87
2003.....	1,104	6	63	37	40	61	57	48	46	67	83	96
2002.....	1,152	3	49	49	42	57	46	45	53	83	99	73
2001.....	1,088	3	58	39	38	51	44	31	52	92	82	88
2000.....	1,094	5	68	58	41	55	40	39	46	78	85	60
1999.....	1,052	4	74	67	40	45	44	47	46	69	75	64
White, both sexes												
2004.....	2,638	4	125	70	74	128	161	150	137	187	216	226
2003.....	2,779	9	137	69	81	159	185	158	146	191	233	249
2002.....	2,788	5	112	81	83	148	182	146	170	193	244	231
2001.....	2,744	7	125	75	112	134	179	150	158	219	233	223
2000.....	2,644	7	123	99	90	129	157	134	154	197	217	203
1999.....	2,536	6	121	98	86	134	152	135	150	200	187	189

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	-------------	-----------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Pedestrian involved in collision with motor vehicle (V02-V04)

White, male

2004.....	1,820	3	77	43	51	96	125	118	106	131	169	166
2003.....	1,927	4	88	46	54	112	142	130	108	139	175	175
2002.....	1,903	3	76	52	50	107	145	117	131	138	171	175
2001.....	1,919	4	85	49	84	97	144	127	121	161	177	159
2000.....	1,773	3	76	60	56	81	121	105	121	139	160	160
1999.....	1,728	4	65	57	55	99	118	105	115	152	136	137

White, female

2004.....	818	1	48	27	23	32	36	32	31	56	47	60
2003.....	852	5	49	23	27	47	43	28	38	52	58	74
2002.....	885	2	36	29	33	41	37	29	39	55	73	56
2001.....	825	3	40	26	28	37	35	23	37	58	56	64
2000.....	871	4	47	39	34	48	36	29	33	58	57	43
1999.....	808	2	56	41	31	35	34	30	35	48	51	52

Black, both sexes

2004.....	537	1	29	30	19	18	36	27	34	37	51	66
2003.....	577	1	39	31	27	31	32	29	37	41	73	62
2002.....	610	2	32	32	21	42	29	37	38	70	58	66
2001.....	618	1	37	42	32	33	30	26	33	65	78	62
2000.....	604	2	47	42	23	24	22	36	30	62	78	63
1999.....	654	2	41	55	23	25	31	45	42	71	85	54

Black, male

2004.....	385	1	18	18	13	12	26	17	28	25	34	44
2003.....	412	-	25	21	15	20	21	14	31	34	51	46
2002.....	434	1	22	15	16	28	22	26	29	49	40	52
2001.....	429	1	21	31	25	21	23	20	22	38	59	42
2000.....	441	1	29	26	17	19	19	27	21	47	53	48
1999.....	479	-	25	32	15	16	24	33	35	54	66	45

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Pedestrian involved in collision with motor vehicle (V02-V04)

Black, female

2004.....	152	-	11	12	6	6	10	10	6	12	17	22
2003.....	165	1	14	10	12	11	11	15	6	7	22	16
2002.....	176	1	10	17	5	14	7	11	9	21	18	14
2001.....	189	-	16	11	7	12	7	6	11	27	19	20
2000.....	163	1	18	16	6	5	3	9	9	15	25	15
1999.....	175	2	16	23	8	9	7	12	7	17	19	9

American Indian 1/, both sexes

2004.....	68	-	3	1	1	6	5	4	8	10	10	5
2003.....	72	-	-	1	4	5	5	8	5	13	8	8
2002.....	87	-	5	3	-	6	7	10	7	13	11	4
2001.....	70	-	3	-	2	7	8	2	7	13	10	5
2000.....	73	-	4	2	-	6	3	10	10	7	7	8
1999.....	88	-	1	2	1	4	10	10	7	11	7	8

American Indian 1/, male

2004.....	52	-	3	-	-	5	4	4	5	7	7	4
2003.....	53	-	-	-	3	4	5	7	4	8	7	5
2002.....	66	-	4	2	-	5	7	7	5	8	8	4
2001.....	57	-	3	-	-	7	7	1	6	9	8	4
2000.....	61	-	3	-	-	6	3	9	9	6	6	6
1999.....	67	-	1	1	1	4	8	8	5	9	4	8

American Indian 1/, female

2004.....	16	-	-	1	1	1	1	-	3	3	3	1
2003.....	19	-	-	1	1	1	-	1	1	5	1	3
2002.....	21	-	1	1	-	1	-	3	2	5	3	-
2001.....	13	-	-	-	2	-	1	1	1	4	2	1
2000.....	12	-	1	2	-	-	-	1	1	1	1	2
1999.....	21	-	-	1	-	-	2	2	2	2	3	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pedestrian involved in collision with motor vehicle (V02-V04)												
Asian or Pacific Islander 2/, both sexes												
2004.....	121	-	1	2	5	5	8	5	9	2	6	5
2003.....	133	-	3	5	-	8	10	8	5	3	6	5
2002.....	139	-	6	3	5	5	8	4	5	7	13	3
2001.....	125	-	5	4	2	2	5	6	6	4	10	6
2000.....	113	-	5	3	1	3	4	7	6	6	10	2
1999.....	111	1	7	5	1	4	4	5	6	7	5	6
Asian or Pacific Islander 2/, male												
2004.....	57	-	1	1	2	5	4	3	2	1	4	1
2003.....	65	-	3	2	-	6	7	4	4	-	4	2
2002.....	69	-	4	1	1	4	6	2	2	5	8	-
2001.....	64	-	3	2	1	-	4	5	3	1	5	3
2000.....	65	-	3	2	-	1	3	7	3	2	8	2
1999.....	63	1	5	3	-	3	3	2	4	5	3	3
Asian or Pacific Islander 2/, female												
2004.....	64	-	-	1	3	-	4	2	7	1	2	4
2003.....	68	-	-	3	-	2	3	4	1	3	2	3
2002.....	70	-	2	2	4	1	2	2	3	2	5	3
2001.....	61	-	2	2	1	2	1	1	3	3	5	3
2000.....	48	-	2	1	1	2	1	-	3	4	2	-
1999.....	48	-	2	2	1	1	1	3	2	2	2	3

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Pedalcyclist involved in collision with motor vehicle (V12-V14,V19.0-V19.2,V19.4-V19.6)

All races, both sexes

2004.....	656	-	11	37	69	52	34	29	27	53	75	67
2003.....	587	-	2	34	80	54	30	26	21	48	70	55
2002.....	570	-	4	34	71	46	29	25	45	41	66	76
2001.....	621	1	3	47	64	59	31	39	33	47	71	63
2000.....	589	-	4	47	93	47	28	23	33	43	61	61
1999.....	652	-	6	64	83	70	41	32	49	55	79	51

All races, male

2004.....	569	-	4	29	56	41	32	26	23	48	70	57
2003.....	514	-	2	25	69	44	28	21	19	42	65	48
2002.....	510	-	4	26	60	43	23	22	41	38	62	72
2001.....	569	1	2	34	56	57	26	34	31	44	68	59
2000.....	531	-	4	37	84	45	25	21	26	38	59	57
1999.....	573	-	5	53	61	66	35	28	43	51	70	49

All races, female

2004.....	87	-	7	8	13	11	2	3	4	5	5	10
2003.....	73	-	-	9	11	10	2	5	2	6	5	7
2002.....	60	-	-	8	11	3	6	3	4	3	4	4
2001.....	52	-	1	13	8	2	5	5	2	3	3	4
2000.....	58	-	-	10	9	2	3	2	7	5	2	4
1999.....	79	-	1	11	22	4	6	4	6	4	9	2

White, both sexes

2004.....	537	-	8	28	56	40	31	26	21	45	58	47
2003.....	470	-	2	26	58	42	25	25	19	37	52	43
2002.....	469	-	4	23	55	38	29	24	39	29	54	59
2001.....	494	1	2	36	43	44	26	34	29	40	53	55
2000.....	454	-	3	33	67	40	24	15	26	30	42	52
1999.....	521	-	5	47	63	56	33	27	42	38	67	40

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Pedalcyclist involved in collision with motor vehicle (V12-V14,V19.0-V19.2,V19.4-V19.6)

White, male

2004.....	466	-	2	22	47	31	29	23	18	40	54	41
2003.....	408	-	2	17	50	32	25	20	18	33	47	37
2002.....	418	-	4	19	47	35	23	21	36	26	50	56
2001.....	450	1	2	26	36	42	21	30	28	37	51	51
2000.....	406	-	3	28	60	38	21	13	21	25	40	48
1999.....	454	-	4	41	45	52	27	23	37	35	59	38

White, female

2004.....	71	-	6	6	9	9	2	3	3	5	4	6
2003.....	62	-	-	9	8	10	-	5	1	4	5	6
2002.....	51	-	-	4	8	3	6	3	3	3	4	3
2001.....	44	-	-	10	7	2	5	4	1	3	2	4
2000.....	48	-	-	5	7	2	3	2	5	5	2	4
1999.....	67	-	1	6	18	4	6	4	5	3	8	2

Black, both sexes

2004.....	93	-	3	8	10	8	3	3	4	8	15	15
2003.....	90	-	-	7	15	7	4	1	1	10	16	12
2002.....	85	-	-	10	16	7	-	1	5	11	10	13
2001.....	109	-	-	10	19	15	4	-	2	7	18	8
2000.....	118	-	1	11	24	6	3	7	6	11	19	8
1999.....	103	-	1	13	19	10	7	3	6	11	11	11

Black, male

2004.....	82	-	2	6	7	7	3	3	4	8	14	13
2003.....	81	-	-	7	13	7	2	1	-	8	16	11
2002.....	79	-	-	7	13	7	-	1	5	11	10	13
2001.....	104	-	-	7	18	15	4	-	2	7	17	8
2000.....	110	-	1	7	22	6	3	7	4	11	19	8
1999.....	95	-	1	10	15	10	7	3	5	11	11	11

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Pedalcyclist involved in collision with motor vehicle (V12-V14,V19.0-V19.2,V19.4-V19.6)

Black, female

2004.....	11	-	1	2	3	1	-	-	-	-	1	2
2003.....	9	-	-	-	2	-	2	-	1	2	-	1
2002.....	6	-	-	3	3	-	-	-	-	-	-	-
2001.....	5	-	-	3	1	-	-	-	-	-	1	-
2000.....	8	-	-	4	2	-	-	-	2	-	-	-
1999.....	8	-	-	3	4	-	-	-	1	-	-	-

American Indian 1/, both sexes

2004.....	9	-	-	-	-	3	-	-	1	-	-	4
2003.....	9	-	-	1	2	3	-	-	1	-	1	-
2002.....	5	-	-	1	-	-	-	-	-	-	2	2
2001.....	3	-	-	1	2	-	-	-	-	-	-	-
2000.....	8	-	-	2	2	1	-	-	1	1	-	1
1999.....	14	-	-	4	1	2	-	1	1	2	-	-

American Indian 1/, male

2004.....	6	-	-	-	-	2	-	-	1	-	-	2
2003.....	9	-	-	1	2	3	-	-	1	-	1	-
2002.....	3	-	-	-	-	-	-	-	-	-	2	1
2001.....	3	-	-	1	2	-	-	-	-	-	-	-
2000.....	7	-	-	1	2	1	-	-	1	1	-	1
1999.....	12	-	-	2	1	2	-	1	1	2	-	-

American Indian 1/, female

2004.....	3	-	-	-	-	1	-	-	-	-	-	2
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	2	-	-	1	-	-	-	-	-	-	-	1
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	1	-	-	1	-	-	-	-	-	-	-	-
1999.....	2	-	-	2	-	-	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pedalcyclist involved in collision with motor vehicle (V12-V14,V19.0-V19.2,V19.4-V19.6)												
Asian or Pacific Islander 2/, both sexes												
2004.....	17	-	-	1	3	1	-	-	1	-	2	1
2003.....	18	-	-	-	5	2	1	-	-	1	1	-
2002.....	11	-	-	-	-	1	-	-	1	1	-	2
2001.....	15	-	1	-	-	-	1	5	2	-	-	-
2000.....	9	-	-	1	-	-	1	1	-	1	-	-
1999.....	14	-	-	-	-	2	1	1	-	4	1	-
Asian or Pacific Islander 2/, male												
2004.....	15	-	-	1	2	1	-	-	-	-	2	1
2003.....	16	-	-	-	4	2	1	-	-	1	1	-
2002.....	10	-	-	-	-	1	-	-	-	1	-	2
2001.....	12	-	-	-	-	-	1	4	1	-	-	-
2000.....	8	-	-	1	-	-	1	1	-	1	-	-
1999.....	12	-	-	-	-	2	1	1	-	3	-	-
Asian or Pacific Islander 2/, female												
2004.....	2	-	-	-	1	-	-	-	1	-	-	-
2003.....	2	-	-	-	1	-	-	-	-	-	-	-
2002.....	1	-	-	-	-	-	-	-	1	-	-	-
2001.....	3	-	1	-	-	-	-	1	1	-	-	-
2000.....	1	-	-	-	-	-	-	-	-	-	-	-
1999.....	2	-	-	-	-	-	-	-	-	1	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Motorcyclist involved in any accident except collision with railway train (V20-V24,V26-V34,V36-V39)

All races, both sexes

2004.....	4,026	-	-	12	42	232	574	450	433	416	494	468
2003.....	3,680	-	-	7	36	202	514	407	413	404	478	425
2002.....	3,229	-	-	4	21	164	455	385	380	377	406	361
2001.....	3,051	-	1	10	29	172	451	419	388	371	382	312
2000.....	2,785	-	-	5	25	170	391	398	326	349	367	288
1999.....	2,343	-	-	6	25	126	363	345	270	292	304	225

All races, male

2004.....	3,621	-	-	8	39	209	536	419	400	369	422	408
2003.....	3,295	-	-	6	32	182	474	380	368	363	409	366
2002.....	2,922	-	-	2	19	153	429	359	346	325	360	322
2001.....	2,765	-	-	8	27	159	424	393	352	318	340	278
2000.....	2,528	-	-	5	21	153	372	378	301	308	323	251
1999.....	2,127	-	-	3	21	113	343	324	253	270	257	195

All races, female

2004.....	405	-	-	4	3	23	38	31	33	47	72	60
2003.....	385	-	-	1	4	20	40	27	45	41	69	59
2002.....	307	-	-	2	2	11	26	26	34	52	46	39
2001.....	286	-	1	2	2	13	27	26	36	53	42	34
2000.....	257	-	-	-	4	17	19	20	25	41	44	37
1999.....	216	-	-	3	4	13	20	21	17	22	47	30

White, both sexes

2004.....	3,521	-	-	10	39	199	496	353	333	340	450	440
2003.....	3,203	-	-	5	31	183	442	311	308	320	431	408
2002.....	2,832	-	-	4	17	149	372	297	300	330	371	342
2001.....	2,661	-	1	9	26	156	369	332	303	326	358	292
2000.....	2,401	-	-	5	21	150	319	302	243	302	343	278
1999.....	2,018	-	-	6	24	105	293	253	216	255	281	218

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Motorcyclist involved in any accident except collision with railway train (V20-V24,V26-V34,V36-V39)

White, male

2004.....	3,146	-	-	6	36	180	462	326	306	299	379	381
2003.....	2,847	-	-	4	28	165	405	286	273	282	363	352
2002.....	2,544	-	-	2	15	140	349	277	267	280	326	305
2001.....	2,396	-	-	7	24	144	349	310	270	275	317	260
2000.....	2,159	-	-	5	17	135	305	282	221	262	300	241
1999.....	1,818	-	-	3	20	95	278	233	200	233	238	188

White, female

2004.....	375	-	-	4	3	19	34	27	27	41	71	59
2003.....	356	-	-	1	3	18	37	25	35	38	68	56
2002.....	288	-	-	2	2	9	23	20	33	50	45	37
2001.....	265	-	1	2	2	12	20	22	33	51	41	32
2000.....	242	-	-	-	4	15	14	20	22	40	43	37
1999.....	200	-	-	3	4	10	15	20	16	22	43	30

Black, both sexes

2004.....	413	-	-	1	3	27	58	83	88	68	38	21
2003.....	412	-	-	1	4	14	52	91	95	78	39	16
2002.....	326	-	-	-	3	9	66	73	68	44	27	15
2001.....	320	-	-	1	2	13	61	76	74	38	20	18
2000.....	336	-	-	-	4	18	61	85	79	44	19	7
1999.....	282	-	-	-	1	17	64	83	46	34	18	5

Black, male

2004.....	396	-	-	1	3	24	55	80	84	65	37	21
2003.....	394	-	-	1	3	12	49	90	89	77	39	14
2002.....	316	-	-	-	3	8	64	69	67	44	27	13
2001.....	308	-	-	1	2	13	57	74	71	37	20	17
2000.....	324	-	-	-	4	16	57	85	76	43	19	7
1999.....	273	-	-	-	1	14	60	83	46	34	16	5

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Motorcyclist involved in any accident except collision with railway train (V20-V24,V26-V34,V36-V39)

Black, female

2004.....	17	-	-	-	-	3	3	3	4	3	1	-
2003.....	18	-	-	-	1	2	3	1	6	1	-	2
2002.....	10	-	-	-	-	1	2	4	1	-	-	2
2001.....	12	-	-	-	-	-	4	2	3	1	-	1
2000.....	12	-	-	-	-	2	4	-	3	1	-	-
1999.....	9	-	-	-	-	3	4	-	-	-	2	-

American Indian 1/, both sexes

2004.....	33	-	-	-	-	2	8	2	4	3	2	5
2003.....	18	-	-	-	-	1	3	2	3	3	1	1
2002.....	22	-	-	-	1	2	2	3	5	2	4	1
2001.....	20	-	-	-	1	2	3	1	3	2	1	1
2000.....	14	-	-	-	-	-	2	1	-	1	3	2
1999.....	18	-	-	-	-	-	3	2	3	2	3	2

American Indian 1/, male

2004.....	30	-	-	-	-	2	7	2	3	2	2	5
2003.....	11	-	-	-	-	1	3	1	1	1	1	-
2002.....	17	-	-	-	1	1	1	3	5	1	3	1
2001.....	17	-	-	-	1	1	3	1	3	1	1	-
2000.....	12	-	-	-	-	-	1	1	-	1	3	2
1999.....	16	-	-	-	-	-	2	2	3	2	2	2

American Indian 1/, female

2004.....	3	-	-	-	-	-	1	-	1	1	-	-
2003.....	7	-	-	-	-	-	-	1	2	2	-	1
2002.....	5	-	-	-	-	1	1	-	-	1	1	-
2001.....	3	-	-	-	-	1	-	-	-	1	-	1
2000.....	2	-	-	-	-	-	1	-	-	-	-	-
1999.....	2	-	-	-	-	-	1	-	-	-	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Motorcyclist involved in any accident except collision with railway train (V20-V24,V26-V34,V36-V39)

Asian or Pacific Islander 2/, both sexes

2004.....	59	-	-	1	-	4	12	12	8	5	4	2
2003.....	47	-	-	1	1	4	17	3	7	3	7	-
2002.....	49	-	-	-	-	4	15	12	7	1	4	3
2001.....	50	-	-	-	-	1	18	10	8	5	3	1
2000.....	34	-	-	-	-	2	9	10	4	2	2	1
1999.....	25	-	-	-	-	4	3	7	5	1	2	-

Asian or Pacific Islander 2/, male

2004.....	49	-	-	1	-	3	12	11	7	3	4	1
2003.....	43	-	-	1	1	4	17	3	5	3	6	-
2002.....	45	-	-	-	-	4	15	10	7	-	4	3
2001.....	44	-	-	-	-	1	15	8	8	5	2	1
2000.....	33	-	-	-	-	2	9	10	4	2	1	1
1999.....	20	-	-	-	-	4	3	6	4	1	1	-

Asian or Pacific Islander 2/, female

2004.....	10	-	-	-	-	1	-	1	1	2	-	1
2003.....	4	-	-	-	-	-	-	-	2	-	1	-
2002.....	4	-	-	-	-	-	-	2	-	1	-	-
2001.....	6	-	-	-	-	-	3	2	-	-	1	-
2000.....	1	-	-	-	-	-	-	-	-	-	1	-
1999.....	5	-	-	-	-	-	-	1	1	-	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Motor vehicle accident involving collision with railway train (V25,V35,V45,V55,V65,V75,V81.0-V81.1,V87.6,V88.6)

All races, both sexes

2004.....	213	1	-	-	7	29	24	16	26	12	9	18
2003.....	219	-	2	2	5	26	21	12	16	18	13	20
2002.....	247	-	3	10	7	23	29	15	15	20	22	16
2001.....	255	1	7	6	8	17	32	27	13	27	32	16
2000.....	282	1	7	8	5	39	25	24	29	14	24	18
1999.....	269	-	6	6	6	33	32	18	18	14	24	24

All races, male

2004.....	157	1	-	-	4	20	19	13	22	4	4	13
2003.....	159	-	-	1	4	17	19	10	11	14	8	13
2002.....	160	-	2	4	2	15	21	12	11	14	17	12
2001.....	172	1	7	5	3	11	21	21	9	16	24	10
2000.....	200	-	4	4	2	25	20	16	21	10	19	13
1999.....	176	-	5	3	3	20	20	11	13	10	17	20

All races, female

2004.....	56	-	-	-	3	9	5	3	4	8	5	5
2003.....	60	-	2	1	1	9	2	2	5	4	5	7
2002.....	87	-	1	6	5	8	8	3	4	6	5	4
2001.....	83	-	-	1	5	6	11	6	4	11	8	6
2000.....	82	1	3	4	3	14	5	8	8	4	5	5
1999.....	93	-	1	3	3	13	12	7	5	4	7	4

White, both sexes

2004.....	179	-	-	-	6	29	19	9	24	9	8	16
2003.....	185	-	-	2	3	22	18	8	15	18	9	16
2002.....	215	-	3	10	5	22	26	13	12	14	16	14
2001.....	229	1	4	5	7	17	27	25	12	26	28	13
2000.....	253	1	6	8	4	36	22	22	25	13	23	16
1999.....	228	-	4	4	6	28	25	15	16	12	19	19

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Motor vehicle accident involving collision with railway train (V25,V35,V45,V55,V65,V75,V81.0-V81.1,V87.6,V88.6)

White, male

2004.....	128	-	-	-	4	20	15	6	20	2	4	11
2003.....	139	-	-	1	3	16	16	7	10	14	5	10
2002.....	136	-	2	4	1	14	18	11	8	9	13	11
2001.....	152	1	4	4	2	11	17	19	9	15	20	8
2000.....	178	-	3	4	2	23	17	15	17	9	19	11
1999.....	150	-	3	3	3	15	16	10	11	9	14	16

White, female

2004.....	51	-	-	-	2	9	4	3	4	7	4	5
2003.....	46	-	-	1	-	6	2	1	5	4	4	6
2002.....	79	-	1	6	4	8	8	2	4	5	3	3
2001.....	77	-	-	1	5	6	10	6	3	11	8	5
2000.....	75	1	3	4	2	13	5	7	8	4	4	5
1999.....	78	-	1	1	3	13	9	5	5	3	5	3

Black, both sexes

2004.....	33	-	-	-	1	-	5	7	2	3	1	2
2003.....	21	-	2	-	1	3	2	1	-	-	3	2
2002.....	25	-	-	-	2	-	2	2	2	4	6	1
2001.....	24	-	3	1	1	-	5	2	1	1	3	3
2000.....	26	-	-	-	1	3	2	2	3	1	1	2
1999.....	34	-	2	2	-	1	7	3	1	1	5	4

Black, male

2004.....	28	-	-	-	-	-	4	7	2	2	-	2
2003.....	13	-	-	-	-	1	2	1	-	-	2	2
2002.....	19	-	-	-	1	-	2	1	2	4	4	1
2001.....	18	-	3	1	1	-	4	2	-	1	3	2
2000.....	19	-	-	-	-	2	2	1	3	1	-	2
1999.....	21	-	2	-	-	1	4	1	1	1	3	4

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Motor vehicle accident involving collision with railway train (V25,V35,V45,V55,V65,V75,V81.0-V81.1,V87.6,V88.6)												
Black, female												
2004.....	5	-	-	-	1	-	1	-	-	1	1	-
2003.....	8	-	2	-	1	2	-	-	-	-	1	-
2002.....	6	-	-	-	1	-	-	1	-	-	2	-
2001.....	6	-	-	-	-	-	1	-	1	-	-	1
2000.....	7	-	-	-	1	1	-	1	-	-	1	-
1999.....	13	-	-	2	-	-	3	2	-	-	2	-
American Indian 1/, both sexes												
2004.....	1	1	-	-	-	-	-	-	-	-	-	-
2003.....	7	-	-	-	1	-	1	1	-	-	1	1
2002.....	3	-	-	-	-	1	1	-	-	1	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	2	-	-	-	-	-	1	-	1	-	-	-
1999.....	3	-	-	-	-	2	-	-	-	1	-	-
American Indian 1/, male												
2004.....	1	1	-	-	-	-	-	-	-	-	-	-
2003.....	5	-	-	-	1	-	1	1	-	-	1	1
2002.....	3	-	-	-	-	1	1	-	-	1	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	2	-	-	-	-	-	1	-	1	-	-	-
1999.....	2	-	-	-	-	2	-	-	-	-	-	-
American Indian 1/, female												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	2	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	1	-	-	-	-	-	-	-	-	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Motorcyclist involved in collision with railway train (V25,V35)

All races, both sexes

2004.....	5	-	-	-	-	1	1	2	1	-	-	-
2003.....	9	-	-	-	-	1	2	1	-	1	1	-
2002.....	7	-	-	-	2	1	-	1	2	1	-	-
2001.....	2	-	-	-	-	-	-	-	-	1	1	-
2000.....	3	-	-	-	-	-	1	1	-	-	-	-
1999.....	6	-	-	-	1	-	1	1	1	1	-	1

All races, male

2004.....	4	-	-	-	-	1	1	1	1	-	-	-
2003.....	8	-	-	-	-	-	2	1	-	1	1	-
2002.....	5	-	-	-	1	-	-	1	2	1	-	-
2001.....	1	-	-	-	-	-	-	-	-	-	1	-
2000.....	3	-	-	-	-	-	1	1	-	-	-	-
1999.....	6	-	-	-	1	-	1	1	1	1	-	1

All races, female

2004.....	1	-	-	-	-	-	-	1	-	-	-	-
2003.....	1	-	-	-	-	1	-	-	-	-	-	-
2002.....	2	-	-	-	1	1	-	-	-	-	-	-
2001.....	1	-	-	-	-	-	-	-	-	1	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-

White, both sexes

2004.....	5	-	-	-	-	1	1	2	1	-	-	-
2003.....	7	-	-	-	-	-	1	1	-	1	1	-
2002.....	6	-	-	-	2	1	-	1	1	1	-	-
2001.....	2	-	-	-	-	-	-	-	-	1	1	-
2000.....	3	-	-	-	-	-	1	1	-	-	-	-
1999.....	5	-	-	-	1	-	1	-	1	1	-	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Motorcyclist involved in collision with railway train (V25,V35)												
White, male												
2004.....	4	-	-	-	-	1	1	1	1	-	-	-
2003.....	7	-	-	-	-	-	1	1	-	1	1	-
2002.....	4	-	-	-	1	-	-	1	1	1	-	-
2001.....	1	-	-	-	-	-	-	-	-	-	1	-
2000.....	3	-	-	-	-	-	1	1	-	-	-	-
1999.....	5	-	-	-	1	-	1	-	1	1	-	1
White, female												
2004.....	1	-	-	-	-	-	-	1	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	2	-	-	-	1	1	-	-	-	-	-	-
2001.....	1	-	-	-	-	-	-	-	-	1	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	1	-	-	-	-	-	1	-	-	-	-	-
2002.....	1	-	-	-	-	-	-	-	1	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	1	-	-	-	-	-	-	1	-	-	-	-
Black, male												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	1	-	-	-	-	-	1	-	-	-	-	-
2002.....	1	-	-	-	-	-	-	-	1	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	1	-	-	-	-	-	-	1	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other motor vehicle accident involving collision with railway train (V45,V55,V65,V75,V81.0-V81.1,V87.6,V88.6)

All races, both sexes

2004.....	208	1	-	-	7	28	23	14	25	12	9	18
2003.....	210	-	2	2	5	25	19	11	16	17	12	20
2002.....	240	-	3	10	5	22	29	14	13	19	22	16
2001.....	253	1	7	6	8	17	32	27	13	26	31	16
2000.....	279	1	7	8	5	39	24	23	29	14	24	18
1999.....	263	-	6	6	5	33	31	17	17	13	24	23

All races, male

2004.....	153	1	-	-	4	19	18	12	21	4	4	13
2003.....	151	-	-	1	4	17	17	9	11	13	7	13
2002.....	155	-	2	4	1	15	21	11	9	13	17	12
2001.....	171	1	7	5	3	11	21	21	9	16	23	10
2000.....	197	-	4	4	2	25	19	15	21	10	19	13
1999.....	170	-	5	3	2	20	19	10	12	9	17	19

All races, female

2004.....	55	-	-	-	3	9	5	2	4	8	5	5
2003.....	59	-	2	1	1	8	2	2	5	4	5	7
2002.....	85	-	1	6	4	7	8	3	4	6	5	4
2001.....	82	-	-	1	5	6	11	6	4	10	8	6
2000.....	82	1	3	4	3	14	5	8	8	4	5	5
1999.....	93	-	1	3	3	13	12	7	5	4	7	4

White, both sexes

2004.....	174	-	-	-	6	28	18	7	23	9	8	16
2003.....	178	-	-	2	3	22	17	7	15	17	8	16
2002.....	209	-	3	10	3	21	26	12	11	13	16	14
2001.....	227	1	4	5	7	17	27	25	12	25	27	13
2000.....	250	1	6	8	4	36	21	21	25	13	23	16
1999.....	223	-	4	4	5	28	24	15	15	11	19	18

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other motor vehicle accident involving collision with railway train (V45,V55,V65,V75,V81.0-V81.1,V87.6,V88.6)

White, male

2004.....	124	-	-	-	4	19	14	5	19	2	4	11
2003.....	132	-	-	1	3	16	15	6	10	13	4	10
2002.....	132	-	2	4	-	14	18	10	7	8	13	11
2001.....	151	1	4	4	2	11	17	19	9	15	19	8
2000.....	175	-	3	4	2	23	16	14	17	9	19	11
1999.....	145	-	3	3	2	15	15	10	10	8	14	15

White, female

2004.....	50	-	-	-	2	9	4	2	4	7	4	5
2003.....	46	-	-	1	-	6	2	1	5	4	4	6
2002.....	77	-	1	6	3	7	8	2	4	5	3	3
2001.....	76	-	-	1	5	6	10	6	3	10	8	5
2000.....	75	1	3	4	2	13	5	7	8	4	4	5
1999.....	78	-	1	1	3	13	9	5	5	3	5	3

Black, both sexes

2004.....	33	-	-	-	1	-	5	7	2	3	1	2
2003.....	20	-	2	-	1	3	1	1	-	-	3	2
2002.....	24	-	-	-	2	-	2	2	1	4	6	1
2001.....	24	-	3	1	1	-	5	2	1	1	3	3
2000.....	26	-	-	-	1	3	2	2	3	1	1	2
1999.....	33	-	2	2	-	1	7	2	1	1	5	4

Black, male

2004.....	28	-	-	-	-	-	4	7	2	2	-	2
2003.....	12	-	-	-	-	1	1	1	-	-	2	2
2002.....	18	-	-	-	1	-	2	1	1	4	4	1
2001.....	18	-	3	1	1	-	4	2	-	1	3	2
2000.....	19	-	-	-	-	2	2	1	3	1	-	2
1999.....	20	-	2	-	-	1	4	-	1	1	3	4

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other motor vehicle accident involving collision with railway train (V45,V55,V65,V75,V81.0-V81.1,V87.6,V88.6)												
Black, female												
2004.....	5	-	-	-	1	-	1	-	-	1	1	-
2003.....	8	-	2	-	1	2	-	-	-	-	1	-
2002.....	6	-	-	-	1	-	-	1	-	-	2	-
2001.....	6	-	-	-	-	-	1	-	1	-	-	1
2000.....	7	-	-	-	1	1	-	1	-	-	1	-
1999.....	13	-	-	2	-	-	3	2	-	-	2	-
American Indian 1/, both sexes												
2004.....	1	1	-	-	-	-	-	-	-	-	-	-
2003.....	7	-	-	-	1	-	1	1	-	-	1	1
2002.....	3	-	-	-	-	1	1	-	-	1	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	2	-	-	-	-	-	1	-	1	-	-	-
1999.....	3	-	-	-	-	2	-	-	-	1	-	-
American Indian 1/, male												
2004.....	1	1	-	-	-	-	-	-	-	-	-	-
2003.....	5	-	-	-	1	-	1	1	-	-	1	1
2002.....	3	-	-	-	-	1	1	-	-	1	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	2	-	-	-	-	-	1	-	1	-	-	-
1999.....	2	-	-	-	-	2	-	-	-	-	-	-
American Indian 1/, female												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	2	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	1	-	-	-	-	-	-	-	-	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Occupant of motor vehicle involved in collision with other (non-motorized) road vehicle, streetcar, animal or pedestrian (V40-V41,V46-V47,V50-V51,V56-V57, V60-V61,V66-V67,V70-V71,V76-V77)

All races, both sexes

2004.....	4,670	8	26	34	64	800	914	490	341	345	316	289
2003.....	4,972	7	35	30	90	827	912	541	394	346	365	326
2002.....	5,060	13	33	39	75	916	920	523	388	394	376	314
2001.....	4,257	3	23	31	52	745	778	418	364	326	317	274
2000.....	3,909	9	26	32	63	684	672	384	306	366	276	220
1999.....	3,805	12	20	26	64	672	620	389	333	316	268	218

All races, male

2004.....	3,403	2	10	22	31	565	716	380	258	270	230	217
2003.....	3,653	2	14	22	62	604	729	418	302	257	270	236
2002.....	3,695	5	17	21	41	658	729	422	303	291	270	217
2001.....	3,142	1	10	19	30	538	618	328	281	235	247	208
2000.....	2,816	4	10	13	32	501	549	309	228	257	197	152
1999.....	2,786	8	9	17	42	457	505	314	260	237	199	153

All races, female

2004.....	1,267	6	16	12	33	235	198	110	83	75	86	72
2003.....	1,319	5	21	8	28	223	183	123	92	89	95	90
2002.....	1,365	8	16	18	34	258	191	101	85	103	106	97
2001.....	1,115	2	13	12	22	207	160	90	83	91	70	66
2000.....	1,093	5	16	19	31	183	123	75	78	109	79	68
1999.....	1,019	4	11	9	22	215	115	75	73	79	69	65

White, both sexes

2004.....	3,879	5	18	24	49	705	755	391	262	285	259	242
2003.....	4,189	5	25	28	73	707	777	428	331	296	316	268
2002.....	4,196	8	25	25	54	778	759	422	318	311	335	258
2001.....	3,466	3	14	21	47	611	604	335	288	269	254	215
2000.....	3,179	5	18	24	52	573	528	291	255	307	223	175
1999.....	3,080	5	12	21	52	565	484	306	266	252	218	175

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Occupant of motor vehicle involved in collision with other (non-motorized) road vehicle, streetcar, animal or pedestrian (V40-V41,V46-V47,V50-V51,V56-V57, V60-V61,V66-V67,V70-V71,V76-V77)

White, male

2004.....	2,839	1	6	16	24	502	604	305	192	230	185	185
2003.....	3,066	1	11	20	48	516	617	324	255	221	234	193
2002.....	3,043	3	13	14	29	550	604	336	252	225	247	175
2001.....	2,557	1	6	14	28	437	488	263	220	195	200	165
2000.....	2,263	3	5	11	25	412	434	232	185	212	160	123
1999.....	2,265	4	8	13	34	382	400	247	206	192	161	121

White, female

2004.....	1,040	4	12	8	25	203	151	86	70	55	74	57
2003.....	1,123	4	14	8	25	191	160	104	76	75	82	75
2002.....	1,153	5	12	11	25	228	155	86	66	86	88	83
2001.....	909	2	8	7	19	174	116	72	68	74	54	50
2000.....	916	2	13	13	27	161	94	59	70	95	63	52
1999.....	815	1	4	8	18	183	84	59	60	60	57	54

Black, both sexes

2004.....	641	2	8	8	14	71	120	82	65	51	47	37
2003.....	619	2	9	2	14	90	105	86	47	38	37	50
2002.....	715	5	8	14	18	106	129	83	58	69	38	50
2001.....	652	-	8	8	4	112	138	69	63	50	59	45
2000.....	574	3	6	8	11	80	104	72	38	50	47	38
1999.....	596	6	8	5	12	87	111	67	57	54	43	32

Black, male

2004.....	465	1	4	5	7	51	83	65	54	34	38	24
2003.....	475	1	3	2	11	65	92	73	36	31	28	39
2002.....	543	2	4	7	10	83	100	73	44	54	20	37
2001.....	493	-	3	4	2	85	107	53	49	36	46	35
2000.....	433	1	3	2	7	61	80	61	34	38	34	24
1999.....	440	3	1	4	8	63	85	56	47	37	34	26

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Occupant of motor vehicle involved in collision with other (non-motorized) road vehicle, streetcar, animal or pedestrian (V40-V41,V46-V47,V50-V51,V56-V57, V60-V61,V66-V67,V70-V71,V76-V77)

Black, female

2004.....	176	1	4	3	7	20	37	17	11	17	9	13
2003.....	144	1	6	-	3	25	13	13	11	7	9	11
2002.....	172	3	4	7	8	23	29	10	14	15	18	13
2001.....	159	-	5	4	2	27	31	16	14	14	13	10
2000.....	141	2	3	6	4	19	24	11	4	12	13	14
1999.....	156	3	7	1	4	24	26	11	10	17	9	6

American Indian 1/, both sexes

2004.....	61	1	-	1	-	9	10	12	6	3	5	4
2003.....	64	-	1	-	-	10	14	6	11	8	3	4
2002.....	74	-	-	-	1	16	13	12	7	7	3	3
2001.....	56	-	-	2	-	6	17	7	4	4	1	4
2000.....	57	-	1	-	-	12	11	10	7	7	3	3
1999.....	56	1	-	-	-	8	7	8	7	4	4	4

American Indian 1/, male

2004.....	40	-	-	-	-	3	7	7	5	2	4	4
2003.....	45	-	-	-	-	6	11	4	8	5	2	2
2002.....	52	-	-	-	1	10	9	9	6	5	3	3
2001.....	37	-	-	1	-	4	10	6	4	3	-	3
2000.....	42	-	1	-	-	9	8	7	4	6	2	3
1999.....	32	1	-	-	-	5	5	5	5	2	1	1

American Indian 1/, female

2004.....	21	1	-	1	-	6	3	5	1	1	1	-
2003.....	19	-	1	-	-	4	3	2	3	3	1	2
2002.....	22	-	-	-	-	6	4	3	1	2	-	-
2001.....	19	-	-	1	-	2	7	1	-	1	1	1
2000.....	15	-	-	-	-	3	3	3	3	1	1	-
1999.....	24	-	-	-	-	3	2	3	2	2	3	3

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Occupant of motor vehicle involved in collision with other (non-motorized) road vehicle, streetcar, animal or pedestrian (V40-V41,V46-V47,V50-V51,V56-V57, V60-V61,V66-V67,V70-V71,V76-V77)

Asian or Pacific Islander 2/, both sexes

2004.....	89	-	-	1	1	15	29	5	8	6	5	6
2003.....	100	-	-	-	3	20	16	21	5	4	9	4
2002.....	75	-	-	-	2	16	19	6	5	7	-	3
2001.....	83	-	1	-	1	16	19	7	9	3	3	10
2000.....	99	1	1	-	-	19	29	11	6	2	3	4
1999.....	73	-	-	-	-	12	18	8	3	6	3	7

Asian or Pacific Islander 2/, male

2004.....	59	-	-	1	-	9	22	3	7	4	3	4
2003.....	67	-	-	-	3	17	9	17	3	-	6	2
2002.....	57	-	-	-	1	15	16	4	1	7	-	2
2001.....	55	-	1	-	-	12	13	6	8	1	1	5
2000.....	78	-	1	-	-	19	27	9	5	1	1	2
1999.....	49	-	-	-	-	7	15	6	2	6	3	5

Asian or Pacific Islander 2/, female

2004.....	30	-	-	-	1	6	7	2	1	2	2	2
2003.....	33	-	-	-	-	3	7	4	2	4	3	2
2002.....	18	-	-	-	1	1	3	2	4	-	-	1
2001.....	28	-	-	-	1	4	6	1	1	2	2	5
2000.....	21	1	-	-	-	-	2	2	1	1	2	2
1999.....	24	-	-	-	-	5	3	2	1	-	-	2

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Occupant of car, pickup truck or van involved in collision with other motor vehicle (V42-V44,V49,V52-V54,V59)

All races, both sexes

2004.....	10,565	61	133	159	218	1,281	1,202	786	645	635	667	688
2003.....	10,735	59	141	151	169	1,262	1,248	786	706	678	751	674
2002.....	11,267	46	150	166	192	1,402	1,373	805	711	735	778	676
2001.....	10,528	64	149	167	210	1,282	1,163	803	648	732	756	652
2000.....	10,390	57	141	181	177	1,246	1,129	803	666	740	672	651
1999.....	10,238	59	132	178	207	1,201	1,053	781	720	678	724	626

All races, male

2004.....	6,307	39	62	83	116	775	807	536	427	418	410	408
2003.....	6,393	29	71	67	96	735	839	528	472	436	479	422
2002.....	6,845	30	82	90	98	874	938	578	458	457	479	415
2001.....	6,350	32	70	78	112	796	812	549	444	466	465	403
2000.....	6,186	29	73	95	94	739	773	518	444	464	417	413
1999.....	6,072	33	77	90	103	730	711	515	478	398	453	384

All races, female

2004.....	4,258	22	71	76	102	506	395	250	218	217	257	280
2003.....	4,342	30	70	84	73	527	409	258	234	242	272	252
2002.....	4,422	16	68	76	94	528	435	227	253	278	299	261
2001.....	4,178	32	79	89	98	486	351	254	204	266	291	249
2000.....	4,204	28	68	86	83	507	356	285	222	276	255	238
1999.....	4,166	26	55	88	104	471	342	266	242	280	271	242

White, both sexes

2004.....	9,003	49	105	129	179	1,111	1,024	630	509	515	556	571
2003.....	9,119	45	108	117	137	1,112	1,035	649	586	551	612	569
2002.....	9,578	32	113	129	160	1,212	1,155	622	566	605	651	551
2001.....	8,924	48	117	126	171	1,078	972	663	529	590	627	517
2000.....	8,813	44	108	145	143	1,082	924	637	535	604	546	534
1999.....	8,678	44	97	140	169	1,041	868	634	587	533	593	523

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Occupant of car, pickup truck or van involved in collision with other motor vehicle (V42-V44,V49,V52-V54,V59)

White, male

2004.....	5,352	31	54	68	99	659	698	435	324	344	350	331
2003.....	5,431	24	56	52	82	642	693	441	395	364	388	362
2002.....	5,805	23	60	75	83	740	800	454	367	378	401	344
2001.....	5,351	25	55	59	93	666	682	455	361	376	378	309
2000.....	5,203	23	55	78	71	641	625	418	351	384	337	339
1999.....	5,101	28	56	70	79	625	587	426	392	305	368	322

White, female

2004.....	3,651	18	51	61	80	452	326	195	185	171	206	240
2003.....	3,688	21	52	65	55	470	342	208	191	187	224	207
2002.....	3,773	9	53	54	77	472	355	168	199	227	250	207
2001.....	3,573	23	62	67	78	412	290	208	168	214	249	208
2000.....	3,610	21	53	67	72	441	299	219	184	220	209	195
1999.....	3,577	16	41	70	90	416	281	208	195	228	225	201

Black, both sexes

2004.....	1,181	10	23	25	29	121	132	124	107	86	82	98
2003.....	1,227	11	24	26	25	111	161	103	95	95	111	79
2002.....	1,262	13	31	27	25	130	167	145	114	99	109	90
2001.....	1,242	11	27	31	27	139	145	113	98	109	106	101
2000.....	1,191	8	25	24	25	114	155	126	102	96	102	94
1999.....	1,215	13	27	32	33	112	139	114	112	124	102	81

Black, male

2004.....	730	6	8	10	11	83	81	84	82	55	46	64
2003.....	744	3	11	10	11	75	111	67	63	49	74	46
2002.....	791	7	21	10	11	92	104	98	76	61	66	55
2001.....	786	5	14	15	15	90	99	74	68	66	75	76
2000.....	765	4	14	12	18	70	109	77	75	56	71	62
1999.....	773	4	15	18	19	78	92	73	72	79	71	51

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Occupant of car, pickup truck or van involved in collision with other motor vehicle (V42-V44,V49,V52-V54,V59)

Black, female

2004.....	451	4	15	15	18	38	51	40	25	31	36	34
2003.....	483	8	13	16	14	36	50	36	32	46	37	33
2002.....	471	6	10	17	14	38	63	47	38	38	43	35
2001.....	456	6	13	16	12	49	46	39	30	43	31	25
2000.....	426	4	11	12	7	44	46	49	27	40	31	32
1999.....	442	9	12	14	14	34	47	41	40	45	31	30

American Indian 1/, both sexes

2004.....	151	1	5	1	7	23	21	11	12	12	8	9
2003.....	155	2	7	6	3	10	24	19	8	13	13	9
2002.....	159	1	2	6	4	23	20	12	11	16	8	10
2001.....	152	4	4	6	6	29	18	10	9	14	10	15
2000.....	158	3	4	6	4	24	22	12	10	18	12	6
1999.....	144	1	4	3	1	22	22	20	8	11	14	6

American Indian 1/, male

2004.....	95	1	-	1	3	18	13	5	10	7	5	5
2003.....	91	1	2	5	2	6	17	10	6	10	11	3
2002.....	91	-	-	3	2	14	13	7	5	9	6	5
2001.....	85	2	-	3	1	14	13	8	7	12	6	7
2000.....	89	1	3	2	2	10	16	9	7	13	4	3
1999.....	84	1	4	1	1	14	15	7	4	6	8	4

American Indian 1/, female

2004.....	56	-	5	-	4	5	8	6	2	5	3	4
2003.....	64	1	5	1	1	4	7	9	2	3	2	6
2002.....	68	1	2	3	2	9	7	5	6	7	2	5
2001.....	67	2	4	3	5	15	5	2	2	2	4	8
2000.....	69	2	1	4	2	14	6	3	3	5	8	3
1999.....	60	-	-	2	-	8	7	13	4	5	6	2

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Occupant of car, pickup truck or van involved in collision with other motor vehicle (V42-V44,V49,V52-V54,V59)

Asian or Pacific Islander 2/, both sexes

2004.....	230	1	-	4	3	26	25	21	17	22	21	10
2003.....	234	1	2	2	4	29	28	15	17	19	15	17
2002.....	268	-	4	4	3	37	31	26	20	15	10	25
2001.....	210	1	1	4	6	36	28	17	12	19	13	19
2000.....	228	2	4	6	5	26	28	28	19	22	12	17
1999.....	201	1	4	3	4	26	24	13	13	10	15	16

Asian or Pacific Islander 2/, male

2004.....	130	1	-	4	3	15	15	12	11	12	9	8
2003.....	127	1	2	-	1	12	18	10	8	13	6	11
2002.....	158	-	1	2	2	28	21	19	10	9	6	11
2001.....	128	-	1	1	3	26	18	12	8	12	6	11
2000.....	129	1	1	3	3	18	23	14	11	11	5	9
1999.....	114	-	2	1	4	13	17	9	10	8	6	7

Asian or Pacific Islander 2/, female

2004.....	100	-	-	-	-	11	10	9	6	10	12	2
2003.....	107	-	-	2	3	17	10	5	9	6	9	6
2002.....	110	-	3	2	1	9	10	7	10	6	4	14
2001.....	82	1	-	3	3	10	10	5	4	7	7	8
2000.....	99	1	3	3	2	8	5	14	8	11	7	8
1999.....	87	1	2	2	-	13	7	4	3	2	9	9

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Occupant of heavy transport vehicle or bus involved in collision with other motor vehicle (V62-V64,V69,V72-V74,V79)

All races, both sexes

2004.....	213	-	2	-	6	8	9	11	17	20	18	27
2003.....	214	-	-	2	1	2	8	12	17	21	24	27
2002.....	218	-	2	2	1	4	11	14	23	22	29	30
2001.....	166	-	1	2	4	3	15	13	16	15	17	22
2000.....	166	-	-	1	6	7	5	7	18	21	25	23
1999.....	207	-	-	3	3	6	6	11	15	25	28	23

All races, male

2004.....	187	-	-	-	5	6	9	10	15	20	17	24
2003.....	175	-	-	1	-	1	8	10	15	17	22	25
2002.....	191	-	-	2	-	4	11	12	19	21	27	27
2001.....	137	-	1	1	3	3	14	11	13	12	10	20
2000.....	143	-	-	1	3	6	4	5	17	19	21	21
1999.....	163	-	-	2	2	4	4	11	12	21	27	22

All races, female

2004.....	26	-	2	-	1	2	-	1	2	-	1	3
2003.....	39	-	-	1	1	1	-	2	2	4	2	2
2002.....	27	-	2	-	1	-	-	2	4	1	2	3
2001.....	29	-	-	1	1	-	1	2	3	3	7	2
2000.....	23	-	-	-	3	1	1	2	1	2	4	2
1999.....	44	-	-	1	1	2	2	-	3	4	1	1

White, both sexes

2004.....	177	-	2	-	4	7	8	8	14	17	14	24
2003.....	186	-	-	2	1	2	7	10	15	17	21	22
2002.....	180	-	2	2	1	4	9	13	20	16	23	26
2001.....	147	-	1	2	2	3	15	12	10	14	15	19
2000.....	137	-	-	1	5	6	5	6	14	16	18	20
1999.....	181	-	-	2	3	6	6	9	12	22	26	20

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Occupant of heavy transport vehicle or bus involved in collision with other motor vehicle (V62-V64,V69,V72-V74,V79)

White, male

2004.....	158	-	-	-	3	5	8	7	13	17	13	22
2003.....	150	-	-	1	-	1	7	8	13	14	20	20
2002.....	158	-	-	2	-	4	9	11	18	15	21	23
2001.....	122	-	1	1	2	3	14	10	9	11	9	17
2000.....	119	-	-	1	3	6	4	4	13	15	15	19
1999.....	143	-	-	1	2	4	4	9	10	18	25	19

White, female

2004.....	19	-	2	-	1	2	-	1	1	-	1	2
2003.....	36	-	-	1	1	1	-	2	2	3	1	2
2002.....	22	-	2	-	1	-	-	2	2	1	2	3
2001.....	25	-	-	1	-	-	1	2	1	3	6	2
2000.....	18	-	-	-	2	-	1	2	1	1	3	1
1999.....	38	-	-	1	1	2	2	-	2	4	1	1

Black, both sexes

2004.....	34	-	-	-	2	-	1	3	3	3	4	3
2003.....	23	-	-	-	-	-	1	2	2	3	3	4
2002.....	35	-	-	-	-	-	2	1	3	5	5	3
2001.....	19	-	-	-	2	-	-	1	6	1	2	3
2000.....	25	-	-	-	-	-	-	1	4	4	7	2
1999.....	22	-	-	1	-	-	-	-	3	3	1	3

Black, male

2004.....	27	-	-	-	2	-	1	3	2	3	4	2
2003.....	22	-	-	-	-	-	1	2	2	3	2	4
2002.....	30	-	-	-	-	-	2	1	1	5	5	3
2001.....	15	-	-	-	1	-	-	1	4	1	1	3
2000.....	24	-	-	-	-	-	-	1	4	4	6	2
1999.....	17	-	-	1	-	-	-	-	2	3	1	3

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Occupant of motor vehicle involved in noncollision accident (V48,V58,V68,V78)												
All races, both sexes												
2004.....	4,030	17	46	70	85	611	667	375	313	336	344	283
2003.....	4,324	18	49	59	102	704	673	422	339	343	357	307
2002.....	4,363	12	47	56	102	746	679	437	373	383	341	304
2001.....	3,908	15	35	47	100	563	651	361	355	357	317	267
2000.....	3,743	22	31	51	98	593	603	424	308	344	299	251
1999.....	3,688	16	48	57	58	611	579	384	308	350	310	237
All races, male												
2004.....	2,815	9	26	42	43	382	508	289	219	247	241	195
2003.....	2,961	12	21	26	61	454	508	315	252	231	249	218
2002.....	3,058	6	32	30	57	483	516	331	273	260	248	230
2001.....	2,742	3	18	26	50	355	502	271	272	253	225	205
2000.....	2,677	15	13	24	60	407	455	335	229	241	217	188
1999.....	2,580	10	24	30	33	400	438	296	221	243	225	169
All races, female												
2004.....	1,215	8	20	28	42	229	159	86	94	89	103	88
2003.....	1,363	6	28	33	41	250	165	107	87	112	108	89
2002.....	1,305	6	15	26	45	263	163	106	100	123	93	74
2001.....	1,166	12	17	21	50	208	149	90	83	104	92	62
2000.....	1,066	7	18	27	38	186	148	89	79	103	82	63
1999.....	1,108	6	24	27	25	211	141	88	87	107	85	68
White, both sexes												
2004.....	3,365	10	31	55	71	541	547	312	250	277	279	229
2003.....	3,627	12	35	47	75	616	565	347	272	282	312	263
2002.....	3,682	11	32	47	84	640	570	367	321	313	275	256
2001.....	3,282	14	26	34	80	490	551	281	285	302	274	213
2000.....	3,133	14	23	38	71	517	506	350	258	285	248	218
1999.....	3,066	11	31	44	49	530	479	317	254	280	255	191

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Occupant of motor vehicle involved in noncollision accident (V48,V58,V68,V78)

White, male

2004.....	2,371	6	19	32	37	336	428	242	184	209	195	156
2003.....	2,493	8	15	18	41	399	428	269	212	181	218	185
2002.....	2,578	6	20	23	49	420	438	273	232	215	198	198
2001.....	2,309	3	14	17	41	305	440	215	217	215	193	163
2000.....	2,266	11	12	17	45	357	388	274	198	202	183	162
1999.....	2,173	7	19	23	28	346	370	244	186	200	191	140

White, female

2004.....	994	4	12	23	34	205	119	70	66	68	84	73
2003.....	1,134	4	20	29	34	217	137	78	60	101	94	78
2002.....	1,104	5	12	24	35	220	132	94	89	98	77	58
2001.....	973	11	12	17	39	185	111	66	68	87	81	50
2000.....	867	3	11	21	26	160	118	76	60	83	65	56
1999.....	893	4	12	21	21	184	109	73	68	80	64	51

Black, both sexes

2004.....	436	7	8	7	5	47	68	41	47	35	44	36
2003.....	445	3	10	7	18	40	65	44	50	43	35	27
2002.....	443	1	14	8	17	57	67	44	34	41	48	34
2001.....	403	-	6	12	13	42	58	47	48	29	33	37
2000.....	406	5	7	9	20	42	69	44	30	38	38	19
1999.....	418	4	14	12	5	52	60	44	38	48	41	33

Black, male

2004.....	305	3	3	5	1	34	47	32	26	25	31	29
2003.....	309	2	4	6	13	28	48	26	31	35	23	22
2002.....	322	-	11	6	7	35	48	38	29	24	36	27
2001.....	290	-	2	9	6	29	34	32	42	21	25	30
2000.....	287	2	1	6	11	28	47	38	22	32	25	14
1999.....	285	3	4	7	2	35	39	36	26	33	27	22

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Occupant of motor vehicle involved in noncollision accident (V48,V58,V68,V78)

Black, female

2004.....	131	4	5	2	4	13	21	9	21	10	13	7
2003.....	136	1	6	1	5	12	17	18	19	8	12	5
2002.....	121	1	3	2	10	22	19	6	5	17	12	7
2001.....	113	-	4	3	7	13	24	15	6	8	8	7
2000.....	119	3	6	3	9	14	22	6	8	6	13	5
1999.....	133	1	10	5	3	17	21	8	12	15	14	11

American Indian 1/, both sexes

2004.....	152	-	6	3	8	19	32	15	11	18	13	11
2003.....	161	3	3	3	7	31	30	22	10	14	8	7
2002.....	158	-	-	-	1	33	28	18	15	20	13	7
2001.....	136	1	1	1	5	19	32	21	10	17	3	8
2000.....	128	2	1	3	5	27	18	17	11	12	10	8
1999.....	124	1	1	1	3	17	27	17	13	13	10	5

American Indian 1/, male

2004.....	91	-	3	2	4	9	20	10	6	10	7	7
2003.....	102	2	1	-	5	17	22	16	6	11	7	4
2002.....	105	-	-	-	1	18	18	14	9	16	9	2
2001.....	90	-	-	-	2	12	21	18	6	11	3	6
2000.....	79	1	-	1	2	19	14	11	5	2	7	7
1999.....	81	-	-	-	2	11	19	12	9	5	5	4

American Indian 1/, female

2004.....	61	-	3	1	4	10	12	5	5	8	6	4
2003.....	59	1	2	3	2	14	8	6	4	3	1	3
2002.....	53	-	-	-	-	15	10	4	6	4	4	5
2001.....	46	1	1	1	3	7	11	3	4	6	-	2
2000.....	49	1	1	2	3	8	4	6	6	10	3	1
1999.....	43	1	1	1	1	6	8	5	4	8	5	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Occupant of motor vehicle involved in noncollision accident (V48,V58,V68,V78)

Asian or Pacific Islander 2/, both sexes

2004.....	77	-	1	5	1	4	20	7	5	6	8	7
2003.....	91	-	1	2	2	17	13	9	7	4	2	10
2002.....	80	-	1	1	-	16	14	8	3	9	5	7
2001.....	87	-	2	-	2	12	10	12	12	9	7	9
2000.....	76	1	-	1	2	7	10	13	9	9	3	6
1999.....	80	-	2	-	1	12	13	6	3	9	4	8

Asian or Pacific Islander 2/, male

2004.....	48	-	1	3	1	3	13	5	3	3	8	3
2003.....	57	-	1	2	2	10	10	4	3	4	1	7
2002.....	53	-	1	1	-	10	12	6	3	5	5	3
2001.....	53	-	2	-	1	9	7	6	7	6	4	6
2000.....	45	1	-	-	2	3	6	12	4	5	2	5
1999.....	41	-	1	-	1	8	10	4	-	5	2	3

Asian or Pacific Islander 2/, female

2004.....	29	-	-	2	-	1	7	2	2	3	-	4
2003.....	34	-	-	-	-	7	3	5	4	-	1	3
2002.....	27	-	-	-	-	6	2	2	-	4	-	4
2001.....	34	-	-	-	1	3	3	6	5	3	3	3
2000.....	31	-	-	1	-	4	4	1	5	4	1	1
1999.....	39	-	1	-	-	4	3	2	3	4	2	5

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Occupant of special-use motor vehicle involved in any accident (V83-V86)

All races, both sexes

2004.....	1,095	1	11	37	119	135	120	97	76	74	86	84
2003.....	1,113	-	13	46	109	113	115	85	81	73	90	86
2002.....	972	-	7	36	100	113	95	78	63	85	60	74
2001.....	1,018	-	13	42	88	130	92	76	80	75	79	44
2000.....	1,042	-	14	43	90	138	101	73	68	73	70	59
1999.....	1,007	-	15	34	76	89	84	71	78	67	69	59

All races, male

2004.....	951	-	9	27	93	107	108	87	64	69	76	74
2003.....	971	-	11	36	77	100	104	72	78	63	83	75
2002.....	850	-	5	27	78	96	87	69	58	77	52	69
2001.....	913	-	10	28	70	117	82	66	75	70	73	40
2000.....	920	-	11	27	65	119	91	70	61	65	62	53
1999.....	924	-	11	28	67	77	79	63	72	63	64	55

All races, female

2004.....	144	1	2	10	26	28	12	10	12	5	10	10
2003.....	142	-	2	10	32	13	11	13	3	10	7	11
2002.....	122	-	2	9	22	17	8	9	5	8	8	5
2001.....	105	-	3	14	18	13	10	10	5	5	6	4
2000.....	122	-	3	16	25	19	10	3	7	8	8	6
1999.....	83	-	4	6	9	12	5	8	6	4	5	4

White, both sexes

2004.....	1,013	1	9	34	112	129	107	91	73	69	82	76
2003.....	1,042	-	12	42	101	101	107	80	76	69	89	81
2002.....	930	-	6	32	94	109	92	75	62	80	56	71
2001.....	948	-	11	40	76	121	85	72	76	72	73	37
2000.....	965	-	14	40	82	119	91	65	68	70	66	55
1999.....	948	-	13	29	71	80	82	69	77	62	67	52

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Occupant of special-use motor vehicle involved in any accident (V83-V86)

White, male

2004.....	875	-	7	25	87	101	95	81	62	65	72	67
2003.....	908	-	10	33	72	88	96	68	73	60	82	70
2002.....	814	-	5	23	74	93	84	66	57	72	48	67
2001.....	853	-	9	27	61	109	76	62	71	67	67	35
2000.....	852	-	11	25	61	103	82	62	61	62	58	49
1999.....	868	-	9	24	63	68	77	61	71	58	62	49

White, female

2004.....	138	1	2	9	25	28	12	10	11	4	10	9
2003.....	134	-	2	9	29	13	11	12	3	9	7	11
2002.....	116	-	1	9	20	16	8	9	5	8	8	4
2001.....	95	-	2	13	15	12	9	10	5	5	6	2
2000.....	113	-	3	15	21	16	9	3	7	8	8	6
1999.....	80	-	4	5	8	12	5	8	6	4	5	3

Black, both sexes

2004.....	57	-	2	2	6	6	8	4	1	3	3	7
2003.....	42	-	1	3	4	8	2	3	3	2	-	5
2002.....	26	-	-	4	3	2	2	2	1	4	3	1
2001.....	46	-	1	-	7	5	5	4	4	2	3	6
2000.....	46	-	-	3	3	9	3	6	-	2	3	4
1999.....	38	-	2	4	4	3	2	1	1	2	2	5

Black, male

2004.....	56	-	2	2	5	6	8	4	1	3	3	7
2003.....	39	-	1	2	4	8	2	2	3	1	-	5
2002.....	25	-	-	4	3	2	2	2	1	4	3	1
2001.....	42	-	1	-	5	5	5	4	4	2	3	5
2000.....	41	-	-	2	-	8	3	6	-	2	3	4
1999.....	36	-	2	3	3	3	2	1	1	2	2	5

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other and unspecified motor vehicle accidents (V09.0,V09.2,V80.3-V80.5,V82.0-V82.1,V87.0-V87.5,V87.7-V87.8,V88.0-V88.5,V88.7-V88.8,V89.0,V89.2)

All races, both sexes

2004.....	16,101	50	248	188	304	1,919	2,009	1,409	1,121	1,063	1,181	1,196
2003.....	15,352	51	200	209	292	1,876	1,950	1,150	1,009	1,110	1,204	1,042
2002.....	15,830	45	209	198	272	2,019	2,008	1,252	1,143	1,122	1,211	1,085
2001.....	16,427	52	224	245	265	2,061	2,082	1,370	1,126	1,294	1,236	1,065
2000.....	17,014	70	249	266	321	2,165	2,169	1,392	1,215	1,323	1,276	1,089
1999.....	16,803	88	253	268	336	2,223	1,955	1,378	1,178	1,374	1,188	1,027

All races, male

2004.....	10,513	25	134	96	153	1,230	1,467	1,022	752	709	824	820
2003.....	10,077	33	118	120	163	1,221	1,422	844	739	746	812	735
2002.....	10,361	23	113	113	156	1,295	1,479	898	818	776	783	733
2001.....	10,805	23	126	137	150	1,385	1,555	1,000	781	850	842	733
2000.....	11,110	38	142	135	180	1,425	1,554	1,004	850	911	891	727
1999.....	10,814	41	139	150	190	1,423	1,427	980	792	911	790	696

All races, female

2004.....	5,588	25	114	92	151	689	542	387	369	354	357	376
2003.....	5,275	18	82	89	129	655	528	306	270	364	392	307
2002.....	5,469	22	96	85	116	724	529	354	325	346	428	352
2001.....	5,622	29	98	108	115	676	527	370	345	444	394	332
2000.....	5,904	32	107	131	141	740	615	388	365	412	385	362
1999.....	5,989	47	114	118	146	800	528	398	386	463	398	331

White, both sexes

2004.....	13,306	34	182	137	215	1,619	1,669	1,113	911	841	959	972
2003.....	12,624	37	148	163	233	1,573	1,579	926	797	894	955	816
2002.....	13,184	25	155	141	207	1,740	1,646	984	926	896	995	875
2001.....	13,681	35	165	174	189	1,755	1,692	1,110	909	1,046	1,009	863
2000.....	14,181	44	189	200	261	1,878	1,775	1,086	962	1,094	1,046	870
1999.....	14,087	67	172	200	263	1,905	1,613	1,132	952	1,137	993	817

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	-------------	-----------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Other and unspecified motor vehicle accidents (V09.0,V09.2,V80.3-V80.5,V82.0-V82.1,V87.0-V87.5,V87.7-V87.8,V88.0-V88.5,V88.7-V88.8,V89.0,V89.2)

White, male

2004.....	8,661	13	93	71	104	1,039	1,229	806	610	557	674	669
2003.....	8,291	25	91	94	130	1,015	1,147	689	590	601	655	571
2002.....	8,601	12	80	74	119	1,118	1,213	707	665	621	650	601
2001.....	8,975	15	92	94	101	1,168	1,275	815	643	674	695	589
2000.....	9,251	24	110	101	145	1,230	1,294	786	688	751	724	587
1999.....	9,063	34	89	109	150	1,213	1,183	817	635	759	664	553

White, female

2004.....	4,645	21	89	66	111	580	440	307	301	284	285	303
2003.....	4,333	12	57	69	103	558	432	237	207	293	300	245
2002.....	4,583	13	75	67	88	622	433	277	261	275	345	274
2001.....	4,706	20	73	80	88	587	417	295	266	372	314	274
2000.....	4,930	20	79	99	116	648	481	300	274	343	322	283
1999.....	5,024	33	83	91	113	692	430	315	317	378	329	264

Black, both sexes

2004.....	2,088	11	55	40	72	222	254	211	159	170	169	183
2003.....	1,975	11	40	33	45	193	268	164	153	166	194	171
2002.....	1,935	14	38	46	41	200	262	203	158	172	157	159
2001.....	2,073	7	41	60	59	209	294	195	163	189	184	164
2000.....	2,173	22	48	52	42	217	302	235	205	177	188	178
1999.....	2,108	18	65	60	53	246	256	187	165	190	149	180

Black, male

2004.....	1,436	8	31	20	42	144	189	156	109	123	117	130
2003.....	1,345	6	21	18	24	136	207	117	111	115	126	128
2002.....	1,315	7	23	31	25	127	195	148	110	120	102	101
2001.....	1,415	1	23	38	38	148	214	142	107	134	121	124
2000.....	1,478	13	26	25	26	148	208	167	133	127	144	117
1999.....	1,402	7	39	36	31	164	188	132	118	120	99	127

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other and unspecified motor vehicle accidents (V09.0,V09.2,V80.3-V80.5,V82.0-V82.1,V87.0-V87.5,V87.7-V87.8,V88.0-V88.5,V88.7-V88.8,V89.0,V89.2)

Black, female

2004.....	652	3	24	20	30	78	65	55	50	47	52	53
2003.....	630	5	19	15	21	57	61	47	42	51	68	43
2002.....	620	7	15	15	16	73	67	55	48	52	55	58
2001.....	658	6	18	22	21	61	80	53	56	55	63	40
2000.....	695	9	22	27	16	69	94	68	72	50	44	61
1999.....	706	11	26	24	22	82	68	55	47	70	50	53

American Indian 1/, both sexes

2004.....	307	2	5	5	5	32	45	43	30	27	24	19
2003.....	352	2	7	7	5	58	50	35	29	23	32	32
2002.....	341	3	8	2	16	39	49	31	33	30	34	25
2001.....	307	5	8	4	10	50	43	28	32	28	22	19
2000.....	311	2	4	6	8	35	55	40	31	28	25	17
1999.....	263	1	8	3	9	37	43	23	29	23	24	15

American Indian 1/, male

2004.....	199	2	5	2	2	19	29	33	20	15	18	10
2003.....	219	2	5	4	5	38	34	20	20	15	18	22
2002.....	223	1	5	1	7	24	40	24	23	19	16	17
2001.....	201	3	5	2	8	33	28	19	21	26	14	10
2000.....	184	-	2	5	4	23	31	28	20	16	15	10
1999.....	168	-	6	2	6	26	30	9	18	17	17	10

American Indian 1/, female

2004.....	108	-	-	3	3	13	16	10	10	12	6	9
2003.....	133	-	2	3	-	20	16	15	9	8	14	10
2002.....	118	2	3	1	9	15	9	7	10	11	18	8
2001.....	106	2	3	2	2	17	15	9	11	2	8	9
2000.....	127	2	2	1	4	12	24	12	11	12	10	7
1999.....	95	1	2	1	3	11	13	14	11	6	7	5

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other and unspecified motor vehicle accidents (V09.0,V09.2,V80.3-V80.5,V82.0-V82.1,V87.0-V87.5,V87.7-V87.8,V88.0-V88.5,V88.7-V88.8,V89.0,V89.2)

Asian or Pacific Islander 2/, both sexes

2004.....	400	3	6	6	12	46	41	42	21	25	29	22
2003.....	401	1	5	6	9	52	53	25	30	27	23	23
2002.....	370	3	8	9	8	40	51	34	26	24	25	26
2001.....	366	5	10	7	7	47	53	37	22	31	21	19
2000.....	349	2	8	8	10	35	37	31	17	24	17	24
1999.....	345	2	8	5	11	35	43	36	32	24	22	15

Asian or Pacific Islander 2/, male

2004.....	217	2	5	3	5	28	20	27	13	14	15	11
2003.....	222	-	1	4	4	32	34	18	18	15	13	14
2002.....	222	3	5	7	5	26	31	19	20	16	15	14
2001.....	214	4	6	3	3	36	38	24	10	16	12	10
2000.....	197	1	4	4	5	24	21	23	9	17	8	13
1999.....	181	-	5	3	3	20	26	22	21	15	10	6

Asian or Pacific Islander 2/, female

2004.....	183	1	1	3	7	18	21	15	8	11	14	11
2003.....	179	1	4	2	5	20	19	7	12	12	10	9
2002.....	148	-	3	2	3	14	20	15	6	8	10	12
2001.....	152	1	4	4	4	11	15	13	12	15	9	9
2000.....	152	1	4	4	5	11	16	8	8	7	9	11
1999.....	164	2	3	2	8	15	17	14	11	9	12	9

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other and unspecified land transport accidents (V01,V06,V09.1,V09.3,V09.9,V10-V11,V16-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.7-V80.9,V87.9,V88.9,V89.1,V89.3,V89.9)

All races, both sexes

2004.....	741	1	19	23	21	61	61	48	42	47	69	61
2003.....	929	2	15	21	23	99	97	65	62	49	83	77
2002.....	610	2	15	17	36	24	45	22	33	50	51	69
2001.....	911	3	26	24	37	57	61	38	47	81	83	90
2000.....	1,009	1	26	37	45	81	69	56	76	78	101	78
1999.....	1,624	3	46	54	49	137	133	106	115	131	137	135

All races, male

2004.....	538	1	12	12	17	45	46	36	33	37	50	46
2003.....	679	-	8	10	15	68	76	51	47	35	59	62
2002.....	473	2	9	9	26	20	38	16	19	44	37	58
2001.....	650	-	18	12	23	41	52	29	39	64	65	59
2000.....	707	1	13	19	30	57	56	46	56	60	75	51
1999.....	1,136	3	25	36	33	90	106	87	72	92	104	100

All races, female

2004.....	203	-	7	11	4	16	15	12	9	10	19	15
2003.....	250	2	7	11	8	31	21	14	15	14	24	15
2002.....	137	-	6	8	10	4	7	6	14	6	14	11
2001.....	261	3	8	12	14	16	9	9	8	17	18	31
2000.....	302	-	13	18	15	24	13	10	20	18	26	27
1999.....	488	-	21	18	16	47	27	19	43	39	33	35

White, both sexes

2004.....	598	-	15	17	17	49	43	33	33	42	55	48
2003.....	772	2	12	15	21	87	85	55	44	38	68	61
2002.....	505	1	12	14	27	22	37	19	28	38	40	57
2001.....	714	2	22	15	28	44	51	32	34	55	60	69
2000.....	845	1	19	32	40	70	58	43	64	61	82	59
1999.....	1,365	3	30	39	41	114	111	89	93	111	115	113

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other and unspecified land transport accidents (V01,V06,V09.1,V09.3,V09.9,V10-V11,V16-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.7-V80.9,V87.9,V88.9,V89.1,V89.3,V89.9)

White, male

2004.....	425	-	9	9	14	36	29	23	25	33	37	37
2003.....	558	-	7	6	13	61	67	43	33	26	50	46
2002.....	395	1	8	9	20	19	32	14	17	34	28	48
2001.....	522	-	16	8	17	34	44	24	29	46	47	48
2000.....	597	1	9	16	28	48	48	36	47	48	62	39
1999.....	947	3	14	24	27	73	87	74	57	79	86	81

White, female

2004.....	173	-	6	8	3	13	14	10	8	9	18	11
2003.....	214	2	5	9	8	26	18	12	11	12	18	15
2002.....	110	-	4	5	7	3	5	5	11	4	12	9
2001.....	192	2	6	7	11	10	7	8	5	9	13	21
2000.....	248	-	10	16	12	22	10	7	17	13	20	20
1999.....	418	-	16	15	14	41	24	15	36	32	29	32

Black, both sexes

2004.....	111	1	3	4	3	10	14	11	8	5	12	12
2003.....	121	-	3	4	1	8	8	9	13	10	13	12
2002.....	78	1	2	3	7	2	4	3	3	11	6	11
2001.....	144	-	4	9	8	8	5	5	9	17	20	17
2000.....	114	-	5	4	4	7	8	7	10	14	14	14
1999.....	177	-	13	13	4	14	10	11	17	16	15	14

Black, male

2004.....	91	1	2	2	2	8	13	11	7	4	12	8
2003.....	95	-	1	3	1	4	7	7	10	8	7	12
2002.....	62	1	1	-	4	1	3	2	2	9	6	10
2001.....	97	-	2	4	6	5	5	4	6	11	16	11
2000.....	78	-	3	3	2	5	6	6	7	11	8	10
1999.....	132	-	9	12	4	11	8	9	12	11	13	12

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other and unspecified land transport accidents (V01,V06,V09.1,V09.3,V09.9,V10-V11,V16-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.7-V80.9,V87.9,V88.9,V89.1,V89.3,V89.9)

Black, female

2004.....	20	-	1	2	1	2	1	-	1	1	-	4
2003.....	26	-	2	1	-	4	1	2	3	2	6	-
2002.....	16	-	1	3	3	1	1	1	1	2	-	1
2001.....	47	-	2	5	2	3	-	1	3	6	4	6
2000.....	36	-	2	1	2	2	2	1	3	3	6	4
1999.....	45	-	4	1	-	3	2	2	5	5	2	2

American Indian 1/, both sexes

2004.....	13	-	1	1	-	2	3	1	-	-	1	-
2003.....	19	-	-	1	1	4	1	1	5	1	-	2
2002.....	12	-	-	-	1	-	4	-	1	1	3	-
2001.....	25	1	-	-	1	3	3	1	3	5	1	2
2000.....	17	-	1	-	1	3	1	2	-	1	3	2
1999.....	40	-	1	-	2	5	7	1	3	2	5	8

American Indian 1/, male

2004.....	12	-	1	1	-	1	3	1	-	-	1	-
2003.....	15	-	-	-	1	3	1	1	4	1	-	2
2002.....	9	-	-	-	1	-	3	-	-	1	2	-
2001.....	17	-	-	-	-	2	2	1	3	4	1	-
2000.....	12	-	-	-	-	3	1	2	-	1	3	1
1999.....	28	-	1	-	-	4	6	1	1	1	3	7

American Indian 1/, female

2004.....	1	-	-	-	-	1	-	-	-	-	-	-
2003.....	4	-	-	1	-	1	-	-	1	-	-	-
2002.....	3	-	-	-	-	-	1	-	1	-	1	-
2001.....	8	1	-	-	1	1	1	-	-	1	-	2
2000.....	5	-	1	-	1	-	-	-	-	-	-	1
1999.....	12	-	-	-	2	1	1	-	2	1	2	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other and unspecified land transport accidents (V01,V06,V09.1,V09.3,V09.9,V10-V11,V16-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.7-V80.9,V87.9,V88.9,V89.1,V89.3,V89.9)

Asian or Pacific Islander 2/, both sexes

2004.....	19	-	-	1	1	-	1	3	1	-	1	1
2003.....	17	-	-	1	-	-	3	-	-	-	2	2
2002.....	15	-	1	-	1	-	-	-	1	-	2	1
2001.....	28	-	-	-	-	2	2	-	1	4	2	2
2000.....	33	-	1	1	-	1	2	4	2	2	2	3
1999.....	42	-	2	2	2	4	5	5	2	2	2	-

Asian or Pacific Islander 2/, male

2004.....	10	-	-	-	1	-	1	1	1	-	-	1
2003.....	11	-	-	1	-	-	1	-	-	-	2	2
2002.....	7	-	-	-	1	-	-	-	-	-	1	-
2001.....	14	-	-	-	-	-	1	-	1	3	1	-
2000.....	20	-	1	-	-	1	1	2	2	-	2	1
1999.....	29	-	1	-	2	2	5	3	2	1	2	-

Asian or Pacific Islander 2/, female

2004.....	9	-	-	1	-	-	-	2	-	-	1	-
2003.....	6	-	-	-	-	-	2	-	-	-	-	-
2002.....	8	-	1	-	-	-	-	-	1	-	1	1
2001.....	14	-	-	-	-	2	1	-	-	1	1	2
2000.....	13	-	-	1	-	-	1	2	-	2	-	2
1999.....	13	-	1	2	-	2	-	2	-	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Water transport accidents (V90-V94)

All races, both sexes

2004.....	574	-	1	10	13	38	52	54	52	50	52	58
2003.....	573	-	6	8	22	47	57	39	41	63	61	52
2002.....	617	2	5	8	16	49	51	42	64	72	62	47
2001.....	591	1	5	14	15	56	54	54	45	50	60	53
2000.....	630	-	4	14	16	61	55	57	52	69	78	47
1999.....	679	1	12	18	29	48	63	64	63	67	70	69

All races, male

2004.....	486	-	-	6	8	30	46	42	43	44	42	51
2003.....	502	-	2	5	15	38	48	34	37	56	54	47
2002.....	555	2	3	2	11	35	46	40	59	68	56	45
2001.....	526	1	4	10	12	48	49	47	40	43	52	49
2000.....	566	-	3	11	12	47	50	55	46	64	71	46
1999.....	583	-	7	12	22	37	53	53	54	58	59	65

All races, female

2004.....	88	-	1	4	5	8	6	12	9	6	10	7
2003.....	71	-	4	3	7	9	9	5	4	7	7	5
2002.....	62	-	2	6	5	14	5	2	5	4	6	2
2001.....	65	-	1	4	3	8	5	7	5	7	8	4
2000.....	64	-	1	3	4	14	5	2	6	5	7	1
1999.....	96	1	5	6	7	11	10	11	9	9	11	4

White, both sexes

2004.....	495	-	1	9	11	32	45	47	48	38	40	53
2003.....	488	-	6	8	19	40	43	33	35	54	57	43
2002.....	533	2	5	5	15	40	47	36	60	64	53	42
2001.....	513	1	5	11	15	52	47	45	40	45	53	44
2000.....	543	-	4	10	15	49	52	52	42	57	66	39
1999.....	567	1	8	17	23	42	58	54	53	54	58	58

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Water transport accidents (V90-V94)

White, male

2004.....	419	-	-	5	7	24	39	36	39	33	33	48
2003.....	424	-	2	5	12	32	36	29	31	49	50	38
2002.....	477	2	3	1	10	28	42	34	55	61	47	41
2001.....	450	1	4	8	12	44	42	39	35	38	45	40
2000.....	482	-	3	7	11	35	47	50	37	52	59	38
1999.....	487	-	4	11	17	32	49	47	45	49	47	54

White, female

2004.....	76	-	1	4	4	8	6	11	9	5	7	5
2003.....	64	-	4	3	7	8	7	4	4	5	7	5
2002.....	56	-	2	4	5	12	5	2	5	3	6	1
2001.....	63	-	1	3	3	8	5	6	5	7	8	4
2000.....	61	-	1	3	4	14	5	2	5	5	7	1
1999.....	80	1	4	6	6	10	9	7	8	5	11	4

Black, both sexes

2004.....	60	-	-	1	2	4	5	5	3	9	8	4
2003.....	60	-	-	-	3	5	10	4	3	7	2	6
2002.....	65	-	-	3	1	6	3	3	3	8	6	4
2001.....	59	-	-	2	-	2	5	7	2	3	6	8
2000.....	63	-	-	4	-	8	2	4	9	7	9	5
1999.....	88	-	4	-	6	4	4	7	9	12	11	8

Black, male

2004.....	52	-	-	1	1	4	5	4	3	8	6	3
2003.....	59	-	-	-	3	5	10	4	3	6	2	6
2002.....	61	-	-	1	1	6	3	3	3	7	6	3
2001.....	58	-	-	2	-	2	5	6	2	3	6	8
2000.....	61	-	-	4	-	8	2	4	8	7	9	5
1999.....	76	-	3	-	5	3	3	3	8	9	11	8

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Water transport accidents (V90-V94)												
Asian or Pacific Islander 2/, both sexes												
2004.....	13	-	-	-	-	2	1	1	1	3	2	1
2003.....	15	-	-	-	-	2	3	1	1	1	1	3
2002.....	9	-	-	-	-	2	-	2	-	-	-	-
2001.....	9	-	-	1	-	1	1	1	2	-	1	-
2000.....	16	-	-	-	1	3	-	1	1	3	3	1
1999.....	12	-	-	-	-	1	-	1	1	1	1	3
Asian or Pacific Islander 2/, male												
2004.....	10	-	-	-	-	2	1	1	1	3	1	-
2003.....	11	-	-	-	-	1	1	-	1	1	1	3
2002.....	8	-	-	-	-	1	-	2	-	-	-	-
2001.....	8	-	-	-	-	1	1	1	2	-	1	-
2000.....	15	-	-	-	1	3	-	1	1	3	3	1
1999.....	9	-	-	-	-	1	-	1	1	-	1	3
Asian or Pacific Islander 2/, female												
2004.....	3	-	-	-	-	-	-	-	-	-	1	1
2003.....	4	-	-	-	-	1	2	1	-	-	-	-
2002.....	1	-	-	-	-	1	-	-	-	-	-	-
2001.....	1	-	-	1	-	-	-	-	-	-	-	-
2000.....	1	-	-	-	-	-	-	-	-	-	-	-
1999.....	3	-	-	-	-	-	-	-	-	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Air and space transport accidents (V95-V97)

All races, both sexes

2004.....	679	1	2	3	8	16	36	48	55	68	60	76
2003.....	742	1	3	13	10	13	32	35	46	63	74	101
2002.....	653	-	2	3	11	19	32	27	43	56	69	89
2001.....	918	3	14	8	11	13	49	69	90	94	117	107
2000.....	777	4	4	4	10	20	49	61	71	69	75	90
1999.....	715	-	1	9	11	12	31	64	65	74	78	81

All races, male

2004.....	566	-	1	2	3	7	25	38	49	58	51	60
2003.....	624	1	1	8	3	11	28	26	36	49	67	85
2002.....	564	-	2	3	5	14	29	25	36	47	62	81
2001.....	696	1	10	5	5	10	30	48	77	73	90	76
2000.....	653	2	2	1	6	11	45	51	58	64	69	74
1999.....	587	-	1	6	8	9	25	53	54	58	60	67

All races, female

2004.....	113	1	1	1	5	9	11	10	6	10	9	16
2003.....	118	-	2	5	7	2	4	9	10	14	7	16
2002.....	89	-	-	-	6	5	3	2	7	9	7	8
2001.....	222	2	4	3	6	3	19	21	13	21	27	31
2000.....	124	2	2	3	4	9	4	10	13	5	6	16
1999.....	128	-	-	3	3	3	6	11	11	16	18	14

White, both sexes

2004.....	658	1	2	3	8	16	34	47	53	64	55	74
2003.....	717	1	3	11	10	10	30	34	43	61	72	98
2002.....	633	-	2	2	8	18	31	26	40	54	69	85
2001.....	874	3	13	8	9	13	47	63	82	89	111	101
2000.....	744	4	4	4	10	16	47	57	67	67	75	86
1999.....	691	-	1	8	11	12	29	61	60	71	78	78

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Air and space transport accidents (V95-V97)												
White, male												
2004.....	545	-	1	2	3	7	23	37	47	54	46	58
2003.....	602	1	1	6	3	9	26	25	34	47	65	82
2002.....	550	-	2	2	4	13	28	24	34	46	62	78
2001.....	668	1	9	5	4	10	28	44	70	71	86	75
2000.....	628	2	2	1	6	9	43	47	55	63	69	70
1999.....	565	-	1	5	8	9	24	50	49	55	60	64
White, female												
2004.....	113	1	1	1	5	9	11	10	6	10	9	16
2003.....	115	-	2	5	7	1	4	9	9	14	7	16
2002.....	83	-	-	-	4	5	3	2	6	8	7	7
2001.....	206	2	4	3	5	3	19	19	12	18	25	26
2000.....	116	2	2	3	4	7	4	10	12	4	6	16
1999.....	126	-	-	3	3	3	5	11	11	16	18	14
Black, both sexes												
2004.....	8	-	-	-	-	-	1	-	-	2	2	1
2003.....	6	-	-	-	-	1	1	-	-	1	1	-
2002.....	8	-	-	1	-	-	-	-	2	1	-	3
2001.....	25	-	1	-	2	-	1	5	5	3	3	2
2000.....	12	-	-	-	-	-	1	3	2	1	-	2
1999.....	11	-	-	1	-	-	-	1	3	-	-	2
Black, male												
2004.....	8	-	-	-	-	-	1	-	-	2	2	1
2003.....	6	-	-	-	-	1	1	-	-	1	1	-
2002.....	6	-	-	1	-	-	-	-	2	-	-	2
2001.....	15	-	1	-	1	-	1	3	4	1	2	-
2000.....	10	-	-	-	-	-	1	3	1	-	-	2
1999.....	11	-	-	1	-	-	-	1	3	-	-	2

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Air and space transport accidents (V95-V97)												
Black, female												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	2	-	-	-	-	-	-	-	-	1	-	1
2001.....	10	-	-	-	1	-	-	2	1	2	1	2
2000.....	2	-	-	-	-	-	-	-	1	1	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes												
2004.....	3	-	-	-	-	-	-	-	-	-	-	-
2003.....	3	-	-	-	-	-	-	-	1	-	-	1
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	13	-	-	-	-	-	-	-	1	2	3	3
2000.....	8	-	-	-	-	2	-	-	-	1	-	2
1999.....	7	-	-	-	-	-	1	1	1	1	-	1
American Indian 1/, male												
2004.....	3	-	-	-	-	-	-	-	-	-	-	-
2003.....	3	-	-	-	-	-	-	-	1	-	-	1
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	7	-	-	-	-	-	-	-	1	1	2	-
2000.....	5	-	-	-	-	-	-	-	-	1	-	2
1999.....	5	-	-	-	-	-	-	1	1	1	-	1
American Indian 1/, female												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	6	-	-	-	-	-	-	-	-	1	1	3
2000.....	3	-	-	-	-	2	-	-	-	-	-	-
1999.....	2	-	-	-	-	-	1	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other and unspecified transport accidents and their sequelae (V98-V99,Y85)

All races, both sexes

2004.....	668	-	2	3	8	19	39	48	44	62	81	93
2003.....	642	-	2	4	6	13	34	37	45	57	76	80
2002.....	630	-	6	9	7	17	30	32	46	59	70	96
2001.....	586	-	5	4	5	21	23	35	47	51	87	66
2000.....	496	-	1	5	3	15	25	30	33	57	69	50
1999.....	510	-	1	6	3	22	23	38	45	63	55	45

All races, male

2004.....	496	-	2	2	5	12	27	37	37	47	61	74
2003.....	466	-	1	3	3	9	26	22	34	45	56	55
2002.....	468	-	5	4	4	12	23	23	36	44	51	77
2001.....	441	-	2	1	4	18	20	23	38	38	66	58
2000.....	371	-	1	2	2	9	18	25	29	46	48	35
1999.....	361	-	-	2	3	16	12	30	30	44	43	31

All races, female

2004.....	172	-	-	1	3	7	12	11	7	15	20	19
2003.....	176	-	1	1	3	4	8	15	11	12	20	25
2002.....	162	-	1	5	3	5	7	9	10	15	19	19
2001.....	145	-	3	3	1	3	3	12	9	13	21	8
2000.....	125	-	-	3	1	6	7	5	4	11	21	15
1999.....	149	-	1	4	-	6	11	8	15	19	12	14

White, both sexes

2004.....	563	-	1	2	6	19	36	37	35	54	70	77
2003.....	532	-	1	2	4	9	27	31	37	47	63	64
2002.....	524	-	6	7	3	16	23	27	36	47	58	81
2001.....	476	-	4	1	3	17	19	31	38	42	75	45
2000.....	420	-	-	5	3	9	22	27	26	48	65	39
1999.....	411	-	1	4	3	16	16	33	37	55	46	37

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other and unspecified transport accidents and their sequelae (V98-V99,Y85)

White, male

2004.....	415	-	1	1	5	12	25	28	28	40	53	63
2003.....	383	-	1	1	1	6	19	20	27	36	43	48
2002.....	383	-	5	3	1	11	17	19	29	34	43	67
2001.....	355	-	1	-	2	14	16	21	32	30	56	41
2000.....	311	-	-	2	2	5	15	22	24	38	45	28
1999.....	289	-	-	1	3	11	9	25	24	37	35	28

White, female

2004.....	148	-	-	1	1	7	11	9	7	14	17	14
2003.....	149	-	-	1	3	3	8	11	10	11	20	16
2002.....	141	-	1	4	2	5	6	8	7	13	15	14
2001.....	121	-	3	1	1	3	3	10	6	12	19	4
2000.....	109	-	-	3	1	4	7	5	2	10	20	11
1999.....	122	-	1	3	-	5	7	8	13	18	11	9

Black, both sexes

2004.....	88	-	1	-	2	-	3	9	8	6	9	15
2003.....	84	-	1	2	2	4	5	5	7	8	9	13
2002.....	89	-	-	2	4	1	7	4	9	11	8	14
2001.....	90	-	1	3	2	4	4	3	8	6	9	18
2000.....	59	-	-	-	-	4	2	2	6	8	3	10
1999.....	79	-	-	2	-	5	7	5	4	8	7	7

Black, male

2004.....	71	-	1	-	-	-	2	8	8	5	7	11
2003.....	65	-	-	2	2	3	5	2	6	7	9	7
2002.....	70	-	-	1	3	1	6	4	6	9	4	9
2001.....	73	-	1	1	2	4	4	2	6	6	8	15
2000.....	48	-	-	-	-	3	2	2	5	7	3	6
1999.....	58	-	-	1	-	4	3	5	3	7	6	3

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other and unspecified transport accidents and their sequelae (V98-V99,Y85)

Asian or Pacific Islander 2/, both sexes

2004.....	9	-	-	1	-	-	-	-	-	1	2	1
2003.....	17	-	-	-	-	-	1	1	-	1	2	2
2002.....	11	-	-	-	-	-	-	-	1	1	1	-
2001.....	14	-	-	-	-	-	-	1	-	3	2	3
2000.....	11	-	-	-	-	2	-	1	-	1	1	-
1999.....	13	-	-	-	-	1	-	-	3	-	-	1

Asian or Pacific Islander 2/, male

2004.....	5	-	-	1	-	-	-	-	-	1	1	-
2003.....	12	-	-	-	-	-	1	-	-	1	2	-
2002.....	10	-	-	-	-	-	-	-	1	1	1	-
2001.....	8	-	-	-	-	-	-	-	-	2	1	2
2000.....	7	-	-	-	-	1	-	1	-	1	-	-
1999.....	9	-	-	-	-	1	-	-	2	-	-	-

Asian or Pacific Islander 2/, female

2004.....	4	-	-	-	-	-	-	-	-	-	1	1
2003.....	5	-	-	-	-	-	-	1	-	-	-	2
2002.....	1	-	-	-	-	-	-	-	-	-	-	-
2001.....	6	-	-	-	-	-	-	1	-	1	1	1
2000.....	4	-	-	-	-	1	-	-	-	-	1	-
1999.....	4	-	-	-	-	-	-	-	1	-	-	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Falls (W00-W19)

All races, both sexes

2004.....	18,807	23	47	11	26	87	154	151	169	241	418	567
2003.....	17,229	14	54	14	31	90	140	139	146	255	381	508
2002.....	16,257	16	37	18	24	83	164	130	177	256	408	487
2001.....	15,019	23	32	33	33	88	168	160	180	272	375	520
2000.....	13,322	8	36	16	21	99	138	138	165	238	370	409
1999.....	13,162	12	55	25	28	112	130	156	187	276	369	447

All races, male

2004.....	9,856	13	31	5	17	70	136	130	144	191	335	447
2003.....	8,910	9	39	9	21	71	115	123	114	218	307	391
2002.....	8,463	9	28	9	18	65	146	118	151	191	331	378
2001.....	8,089	15	25	22	24	79	153	142	153	233	311	426
2000.....	7,122	6	23	14	16	88	115	122	150	199	297	332
1999.....	7,109	8	37	19	21	95	114	137	157	225	303	366

All races, female

2004.....	8,951	10	16	6	9	17	18	21	25	50	83	120
2003.....	8,319	5	15	5	10	19	25	16	32	37	74	117
2002.....	7,794	7	9	9	6	18	18	12	26	65	77	109
2001.....	6,930	8	7	11	9	9	15	18	27	39	64	94
2000.....	6,200	2	13	2	5	11	23	16	15	39	73	77
1999.....	6,053	4	18	6	7	17	16	19	30	51	66	81

White, both sexes

2004.....	17,383	18	33	9	21	79	138	125	146	201	355	477
2003.....	15,855	10	38	13	26	79	126	122	127	211	320	413
2002.....	15,027	12	23	14	19	72	147	119	152	217	343	410
2001.....	13,827	15	28	27	29	80	157	140	157	227	310	429
2000.....	12,252	5	23	12	18	90	127	125	140	194	310	352
1999.....	11,992	8	43	18	24	100	112	135	149	239	308	362

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992),
race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Falls (W00-W19)

White, male

2004.....	8,988	11	23	4	13	64	123	108	126	167	291	378
2003.....	8,099	6	29	8	17	63	106	108	101	183	261	322
2002.....	7,699	6	17	7	15	56	130	108	131	160	280	321
2001.....	7,331	10	21	20	20	71	146	126	134	195	261	348
2000.....	6,449	3	13	10	15	81	106	112	127	163	253	290
1999.....	6,352	7	29	14	18	86	100	120	126	193	249	298

White, female

2004.....	8,395	7	10	5	8	15	15	17	20	34	64	99
2003.....	7,756	4	9	5	9	16	20	14	26	28	59	91
2002.....	7,328	6	6	7	4	16	17	11	21	57	63	89
2001.....	6,496	5	7	7	9	9	11	14	23	32	49	81
2000.....	5,803	2	10	2	3	9	21	13	13	31	57	62
1999.....	5,640	1	14	4	6	14	12	15	23	46	59	64

Black, both sexes

2004.....	999	3	9	2	3	5	9	16	20	32	49	69
2003.....	920	2	13	1	2	7	10	13	14	31	45	79
2002.....	838	3	8	3	3	8	9	8	16	31	46	65
2001.....	854	6	3	5	2	4	9	10	18	33	58	64
2000.....	760	2	13	4	3	6	7	5	18	29	45	47
1999.....	878	2	11	6	4	8	13	13	30	27	48	73

Black, male

2004.....	628	-	7	1	2	3	6	14	16	20	36	51
2003.....	537	1	9	1	2	5	5	11	9	23	36	59
2002.....	532	2	7	2	3	6	9	7	14	25	35	49
2001.....	547	4	3	2	2	4	5	7	14	27	45	56
2000.....	485	2	10	4	1	4	7	3	16	23	34	37
1999.....	564	1	8	4	3	6	10	11	24	24	43	59

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Falls (W00-W19)												
Black, female												
2004.....	371	3	2	1	1	2	3	2	4	12	13	18
2003.....	383	1	4	-	-	2	5	2	5	8	9	20
2002.....	306	1	1	1	-	2	-	1	2	6	11	16
2001.....	307	2	-	3	-	-	4	3	4	6	13	8
2000.....	275	-	3	-	2	2	-	2	2	6	11	10
1999.....	314	1	3	2	1	2	3	2	6	3	5	14
American Indian 1/, both sexes												
2004.....	110	2	1	-	-	1	4	5	3	3	7	12
2003.....	120	-	-	-	-	2	2	2	3	8	8	6
2002.....	97	1	-	1	1	1	3	2	4	3	10	9
2001.....	86	-	-	-	1	2	1	4	-	5	4	14
2000.....	81	1	-	-	-	1	-	6	3	7	6	5
1999.....	75	-	1	-	-	2	4	4	2	3	4	5
American Indian 1/, male												
2004.....	69	2	-	-	-	1	4	5	2	-	3	9
2003.....	83	-	-	-	-	2	2	2	3	8	5	3
2002.....	54	1	-	-	-	1	3	2	3	2	8	6
2001.....	51	-	-	-	1	2	1	3	-	5	2	11
2000.....	57	1	-	-	-	1	-	5	3	5	5	1
1999.....	42	-	-	-	-	2	3	3	1	3	4	4
American Indian 1/, female												
2004.....	41	-	1	-	-	-	-	-	1	3	4	3
2003.....	37	-	-	-	-	-	-	-	-	-	3	3
2002.....	43	-	-	1	1	-	-	-	1	1	2	3
2001.....	35	-	-	-	-	-	-	1	-	-	2	3
2000.....	24	-	-	-	-	-	-	1	-	2	1	4
1999.....	33	-	1	-	-	-	1	1	1	-	-	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Falls (W00-W19)												
Asian or Pacific Islander 2/, both sexes												
2004.....	315	-	4	-	2	2	3	5	-	5	7	9
2003.....	334	2	3	-	3	2	2	2	2	5	8	10
2002.....	295	-	6	-	1	2	5	1	5	5	9	3
2001.....	252	2	1	1	1	2	1	6	5	7	3	13
2000.....	229	-	-	-	-	2	4	2	4	8	9	5
1999.....	217	2	-	1	-	2	1	4	6	7	9	7
Asian or Pacific Islander 2/, male												
2004.....	171	-	1	-	2	2	3	3	-	4	5	9
2003.....	191	2	1	-	2	1	2	2	1	4	5	7
2002.....	178	-	4	-	-	2	4	1	3	4	8	2
2001.....	160	1	1	-	1	2	1	6	5	6	3	11
2000.....	131	-	-	-	-	2	2	2	4	8	5	4
1999.....	151	-	-	1	-	1	1	3	6	5	7	5
Asian or Pacific Islander 2/, female												
2004.....	144	-	3	-	-	-	-	2	-	1	2	-
2003.....	143	-	2	-	1	1	-	-	1	1	3	3
2002.....	117	-	2	-	1	-	1	-	2	1	1	1
2001.....	92	1	-	1	-	-	-	-	-	1	-	2
2000.....	98	-	-	-	-	-	2	-	-	-	4	1
1999.....	66	2	-	-	-	1	-	1	-	2	2	2

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Fall on same level (W00-W09,W18)

All races, both sexes

2004.....	6,188	9	12	4	11	21	25	35	35	47	73	129
2003.....	5,620	8	13	5	12	16	15	20	25	53	81	94
2002.....	5,279	12	8	8	9	19	29	14	34	42	74	105
2001.....	4,042	11	8	11	12	16	26	19	30	46	52	112
2000.....	3,264	6	6	5	6	11	17	15	15	30	55	53
1999.....	2,241	6	9	3	6	19	11	13	14	27	31	65

All races, male

2004.....	2,992	5	6	2	8	12	22	28	25	28	52	98
2003.....	2,679	5	8	3	9	10	7	16	17	41	54	62
2002.....	2,500	7	7	2	4	14	26	13	25	26	58	82
2001.....	1,964	5	5	8	8	12	22	15	26	38	38	85
2000.....	1,518	4	2	5	5	11	10	12	11	24	42	43
1999.....	1,077	4	8	1	4	17	8	10	9	18	21	46

All races, female

2004.....	3,196	4	6	2	3	9	3	7	10	19	21	31
2003.....	2,941	3	5	2	3	6	8	4	8	12	27	32
2002.....	2,779	5	1	6	5	5	3	1	9	16	16	23
2001.....	2,078	6	3	3	4	4	4	4	4	8	14	27
2000.....	1,746	2	4	-	1	-	7	3	4	6	13	10
1999.....	1,164	2	1	2	2	2	3	3	5	9	10	19

White, both sexes

2004.....	5,752	6	9	3	10	20	23	27	29	33	65	107
2003.....	5,235	6	10	5	10	15	12	18	20	40	67	79
2002.....	4,937	8	7	7	7	15	25	11	30	38	63	87
2001.....	3,746	7	8	9	9	14	23	16	27	41	38	95
2000.....	3,040	3	3	5	5	10	16	13	12	23	48	49
1999.....	2,038	4	6	2	6	17	9	12	11	22	28	52

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Fall on same level (W00-W09,W18)

White, male

2004.....	2,750	3	4	1	7	12	21	23	21	21	49	81
2003.....	2,461	4	6	3	7	10	7	14	14	30	46	52
2002.....	2,298	4	6	2	4	12	22	10	22	24	50	69
2001.....	1,804	3	5	7	5	10	21	13	24	33	29	72
2000.....	1,395	1	1	5	5	10	9	11	9	18	38	39
1999.....	965	3	5	1	4	15	7	10	6	13	18	34

White, female

2004.....	3,002	3	5	2	3	8	2	4	8	12	16	26
2003.....	2,774	2	4	2	3	5	5	4	6	10	21	27
2002.....	2,639	4	1	5	3	3	3	1	8	14	13	18
2001.....	1,942	4	3	2	4	4	2	3	3	8	9	23
2000.....	1,645	2	2	-	-	-	7	2	3	5	10	10
1999.....	1,073	1	1	1	2	2	2	2	5	9	10	18

Black, both sexes

2004.....	292	1	3	1	1	1	1	2	5	11	4	16
2003.....	196	1	3	-	2	-	3	2	3	8	11	12
2002.....	204	3	1	-	-	4	2	3	2	2	10	14
2001.....	187	3	-	2	1	1	2	2	2	3	12	9
2000.....	133	2	3	-	1	-	-	-	2	3	3	4
1999.....	149	2	3	1	-	1	2	-	1	5	2	11

Black, male

2004.....	163	-	2	1	1	-	-	1	4	6	3	12
2003.....	112	-	2	-	2	-	-	2	2	6	7	9
2002.....	124	2	1	-	-	2	2	3	2	1	7	10
2001.....	102	2	-	1	1	1	-	1	1	3	8	7
2000.....	69	2	1	-	-	-	-	-	1	3	1	4
1999.....	82	1	3	-	-	1	1	-	1	5	2	10

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Fall on same level (W00-W09,W18)												
Black, female												
2004.....	129	1	1	-	-	1	1	1	1	5	1	4
2003.....	84	1	1	-	-	-	3	-	1	2	4	3
2002.....	80	1	-	-	-	2	-	-	-	1	3	4
2001.....	85	1	-	1	-	-	2	1	1	-	4	2
2000.....	64	-	2	-	1	-	-	-	1	-	2	-
1999.....	67	1	-	1	-	-	1	-	-	-	-	1
American Indian 1/, both sexes												
2004.....	38	2	-	-	-	-	1	2	1	1	3	4
2003.....	55	-	-	-	-	-	-	-	1	5	1	3
2002.....	37	1	-	1	1	-	1	-	-	2	1	4
2001.....	28	-	-	-	1	-	-	1	-	1	1	4
2000.....	26	1	-	-	-	-	-	2	1	3	2	-
1999.....	16	-	-	-	-	-	-	1	-	-	1	1
American Indian 1/, male												
2004.....	20	2	-	-	-	-	1	2	-	-	-	3
2003.....	37	-	-	-	-	-	-	-	1	5	-	1
2002.....	19	1	-	-	-	-	1	-	-	1	1	3
2001.....	16	-	-	-	1	-	-	1	-	1	-	3
2000.....	20	1	-	-	-	-	-	1	1	2	2	-
1999.....	5	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, female												
2004.....	18	-	-	-	-	-	-	-	1	1	3	1
2003.....	18	-	-	-	-	-	-	-	-	-	1	2
2002.....	18	-	-	1	1	-	-	-	-	1	-	1
2001.....	12	-	-	-	-	-	-	-	-	-	1	1
2000.....	6	-	-	-	-	-	-	1	-	1	-	-
1999.....	11	-	-	-	-	-	-	1	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Fall from one level to another (W10-W17)

All races, both sexes

2004.....	3,341	5	20	5	10	45	91	78	90	124	191	221
2003.....	3,303	5	27	6	14	64	109	100	83	132	160	233
2002.....	3,324	4	23	7	11	54	118	97	113	147	194	203
2001.....	3,317	4	20	16	13	61	120	118	109	149	180	229
2000.....	2,912	1	24	7	12	71	103	93	123	132	196	200
1999.....	3,114	5	36	11	15	73	82	115	116	145	191	187

All races, male

2004.....	2,394	4	13	3	5	40	82	71	83	107	169	189
2003.....	2,341	3	21	4	7	53	92	91	72	118	146	196
2002.....	2,386	2	16	5	10	44	105	90	99	122	167	177
2001.....	2,432	4	16	9	11	56	112	109	93	135	161	203
2000.....	2,181	1	16	5	9	62	89	84	115	116	168	168
1999.....	2,266	3	24	9	11	62	73	103	108	126	169	166

All races, female

2004.....	947	1	7	2	5	5	9	7	7	17	22	32
2003.....	962	2	6	2	7	11	17	9	11	14	14	37
2002.....	938	2	7	2	1	10	13	7	14	25	27	26
2001.....	885	-	4	7	2	5	8	9	16	14	19	26
2000.....	731	-	8	2	3	9	14	9	8	16	28	32
1999.....	848	2	12	2	4	11	9	12	8	19	22	21

White, both sexes

2004.....	3,023	5	12	4	6	39	80	69	81	107	162	190
2003.....	2,949	3	20	5	11	55	100	89	73	112	137	187
2002.....	3,020	4	13	5	8	48	107	93	103	126	166	178
2001.....	2,989	2	16	12	12	57	115	105	95	127	155	187
2000.....	2,636	1	15	5	12	68	96	86	108	112	172	174
1999.....	2,809	3	29	7	13	66	73	98	96	127	162	162

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Fall from one level to another (W10-W17)

White, male

2004.....	2,160	4	8	3	2	35	72	62	75	94	146	163
2003.....	2,086	1	16	3	5	46	85	82	64	100	126	164
2002.....	2,176	2	9	4	7	38	95	86	92	104	147	156
2001.....	2,168	2	12	8	10	52	109	97	81	116	138	165
2000.....	1,960	1	8	3	9	59	84	78	101	98	149	147
1999.....	2,022	3	20	5	10	57	66	89	90	111	142	144

White, female

2004.....	863	1	4	1	4	4	8	7	6	13	16	27
2003.....	863	2	4	2	6	9	15	7	9	12	11	23
2002.....	844	2	4	1	1	10	12	7	11	22	19	22
2001.....	821	-	4	4	2	5	6	8	14	11	17	22
2000.....	676	-	7	2	3	9	12	8	7	14	23	27
1999.....	787	-	9	2	3	9	7	9	6	16	20	18

Black, both sexes

2004.....	228	-	4	1	2	4	6	8	8	12	26	25
2003.....	250	1	6	1	-	6	5	7	7	15	16	35
2002.....	212	-	5	2	3	3	5	3	6	17	19	22
2001.....	245	2	3	3	1	1	5	5	11	16	22	30
2000.....	204	-	9	2	-	2	4	3	10	11	17	22
1999.....	213	-	6	3	2	4	5	10	15	12	23	18

Black, male

2004.....	176	-	4	-	1	3	5	8	7	10	20	20
2003.....	178	1	5	1	-	4	3	5	5	13	15	25
2002.....	148	-	4	1	3	3	5	3	4	15	13	18
2001.....	195	2	3	1	1	1	3	4	9	14	20	27
2000.....	165	-	8	2	-	2	4	2	9	10	14	19
1999.....	169	-	4	3	1	3	4	8	13	10	22	15

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Fall from one level to another (W10-W17)												
Black, female												
2004.....	52	-	-	1	1	1	1	-	1	2	6	5
2003.....	72	-	1	-	-	2	2	2	2	2	1	10
2002.....	64	-	1	1	-	-	-	-	2	2	6	4
2001.....	50	-	-	2	-	-	2	1	2	2	2	3
2000.....	39	-	1	-	-	-	-	1	1	1	3	3
1999.....	44	-	2	-	1	1	1	2	2	2	1	3
American Indian 1/, both sexes												
2004.....	25	-	1	-	-	-	2	1	1	2	1	4
2003.....	32	-	-	-	-	2	2	2	2	2	4	3
2002.....	20	-	-	-	-	1	2	-	2	-	6	2
2001.....	21	-	-	-	-	2	-	2	-	1	1	6
2000.....	17	-	-	-	-	-	-	3	1	4	2	1
1999.....	25	-	1	-	-	2	3	3	1	-	1	3
American Indian 1/, male												
2004.....	21	-	-	-	-	-	2	1	1	-	1	4
2003.....	28	-	-	-	-	2	2	2	2	2	3	2
2002.....	15	-	-	-	-	1	2	-	2	-	4	2
2001.....	16	-	-	-	-	2	-	2	-	1	1	5
2000.....	14	-	-	-	-	-	-	3	1	3	1	-
1999.....	21	-	-	-	-	2	2	3	1	-	1	3
American Indian 1/, female												
2004.....	4	-	1	-	-	-	-	-	-	2	-	-
2003.....	4	-	-	-	-	-	-	-	-	-	1	1
2002.....	5	-	-	-	-	-	-	-	-	-	2	-
2001.....	5	-	-	-	-	-	-	-	-	-	-	1
2000.....	3	-	-	-	-	-	-	-	-	1	1	1
1999.....	4	-	1	-	-	-	1	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Fall from one level to another (W10-W17)

Asian or Pacific Islander 2/, both sexes

2004.....	65	-	3	-	2	2	3	-	-	3	2	2
2003.....	72	1	1	-	3	1	2	2	1	3	3	8
2002.....	72	-	5	-	-	2	4	1	2	4	3	1
2001.....	62	-	1	1	-	1	-	6	3	5	2	6
2000.....	55	-	-	-	-	1	3	1	4	5	5	3
1999.....	67	2	-	1	-	1	1	4	4	6	5	4

Asian or Pacific Islander 2/, male

2004.....	37	-	1	-	2	2	3	-	-	3	2	2
2003.....	49	1	-	-	2	1	2	2	1	3	2	5
2002.....	47	-	3	-	-	2	3	1	1	3	3	1
2001.....	53	-	1	-	-	1	-	6	3	4	2	6
2000.....	42	-	-	-	-	1	1	1	4	5	4	2
1999.....	54	-	-	1	-	-	1	3	4	5	4	4

Asian or Pacific Islander 2/, female

2004.....	28	-	2	-	-	-	-	-	-	-	-	-
2003.....	23	-	1	-	1	-	-	-	-	-	1	3
2002.....	25	-	2	-	-	-	1	-	1	1	-	-
2001.....	9	-	-	1	-	-	-	-	-	1	-	-
2000.....	13	-	-	-	-	-	2	-	-	-	1	1
1999.....	13	2	-	-	-	1	-	1	-	1	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Unspecified fall (W19)												
All races, both sexes												
2004.....	9,278	9	15	2	5	21	38	38	44	70	154	217
2003.....	8,306	1	14	3	5	10	16	19	38	70	140	181
2002.....	7,654	-	6	3	4	10	17	19	30	67	140	179
2001.....	7,660	8	4	6	8	11	22	23	41	77	143	179
2000.....	7,146	1	6	4	3	17	18	30	27	76	119	156
1999.....	7,807	1	10	11	7	20	37	28	57	104	147	195
All races, male												
2004.....	4,470	4	12	-	4	18	32	31	36	56	114	160
2003.....	3,890	1	10	2	5	8	16	16	25	59	107	133
2002.....	3,577	-	5	2	4	7	15	15	27	43	106	119
2001.....	3,693	6	4	5	5	11	19	18	34	60	112	138
2000.....	3,423	1	5	4	2	15	16	26	24	59	87	121
1999.....	3,766	1	5	9	6	16	33	24	40	81	113	154
All races, female												
2004.....	4,808	5	3	2	1	3	6	7	8	14	40	57
2003.....	4,416	-	4	1	-	2	-	3	13	11	33	48
2002.....	4,077	-	1	1	-	3	2	4	3	24	34	60
2001.....	3,967	2	-	1	3	-	3	5	7	17	31	41
2000.....	3,723	-	1	-	1	2	2	4	3	17	32	35
1999.....	4,041	-	5	2	1	4	4	4	17	23	34	41
White, both sexes												
2004.....	8,608	7	12	2	5	20	35	29	36	61	128	180
2003.....	7,671	1	8	3	5	9	14	15	34	59	116	147
2002.....	7,070	-	3	2	4	9	15	15	19	53	114	145
2001.....	7,092	6	4	6	8	9	19	19	35	59	117	147
2000.....	6,576	1	5	2	1	12	15	26	20	59	90	129
1999.....	7,145	1	8	9	5	17	30	25	42	90	118	148

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Unspecified fall (W19)

White, male

2004.....	4,078	4	11	-	4	17	30	23	30	52	96	134
2003.....	3,552	1	7	2	5	7	14	12	23	53	89	106
2002.....	3,225	-	2	1	4	6	13	12	17	32	83	96
2001.....	3,359	5	4	5	5	9	16	16	29	46	94	111
2000.....	3,094	1	4	2	1	12	13	23	17	47	66	104
1999.....	3,365	1	4	8	4	14	27	21	30	69	89	120

White, female

2004.....	4,530	3	1	2	1	3	5	6	6	9	32	46
2003.....	4,119	-	1	1	-	2	-	3	11	6	27	41
2002.....	3,845	-	1	1	-	3	2	3	2	21	31	49
2001.....	3,733	1	-	1	3	-	3	3	6	13	23	36
2000.....	3,482	-	1	-	-	-	2	3	3	12	24	25
1999.....	3,780	-	4	1	1	3	3	4	12	21	29	28

Black, both sexes

2004.....	479	2	2	-	-	-	2	6	7	9	19	28
2003.....	474	-	4	-	-	1	2	4	4	8	18	32
2002.....	422	-	2	1	-	1	2	2	8	12	17	29
2001.....	422	1	-	-	-	2	2	3	5	14	24	25
2000.....	423	-	1	2	2	4	3	2	6	15	25	21
1999.....	516	-	2	2	2	3	6	3	14	10	23	44

Black, male

2004.....	289	-	1	-	-	-	1	5	5	4	13	19
2003.....	247	-	2	-	-	1	2	4	2	4	14	25
2002.....	260	-	2	1	-	1	2	1	8	9	15	21
2001.....	250	-	-	-	-	2	2	2	4	10	17	22
2000.....	251	-	1	2	1	2	3	1	6	10	19	14
1999.....	313	-	1	1	2	2	5	3	10	9	19	34

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Unspecified fall (W19)												
Black, female												
2004.....	190	2	1	-	-	-	1	1	2	5	6	9
2003.....	227	-	2	-	-	-	-	-	2	4	4	7
2002.....	162	-	-	-	-	-	-	1	-	3	2	8
2001.....	172	1	-	-	-	-	-	1	1	4	7	3
2000.....	172	-	-	-	1	2	-	1	-	5	6	7
1999.....	203	-	1	1	-	1	1	-	4	1	4	10
American Indian 1/, both sexes												
2004.....	47	-	-	-	-	1	1	2	1	-	3	4
2003.....	33	-	-	-	-	-	-	-	-	1	3	-
2002.....	40	-	-	-	-	-	-	2	2	1	3	3
2001.....	37	-	-	-	-	-	1	1	-	3	2	4
2000.....	38	-	-	-	-	1	-	1	1	-	2	4
1999.....	34	-	-	-	-	-	1	-	1	3	2	1
American Indian 1/, male												
2004.....	28	-	-	-	-	1	1	2	1	-	2	2
2003.....	18	-	-	-	-	-	-	-	-	1	2	-
2002.....	20	-	-	-	-	-	-	2	1	1	3	1
2001.....	19	-	-	-	-	-	1	-	-	3	1	3
2000.....	23	-	-	-	-	1	-	1	1	-	2	1
1999.....	16	-	-	-	-	-	1	-	-	3	2	-
American Indian 1/, female												
2004.....	19	-	-	-	-	-	-	-	-	-	1	2
2003.....	15	-	-	-	-	-	-	-	-	-	1	-
2002.....	20	-	-	-	-	-	-	-	1	-	-	2
2001.....	18	-	-	-	-	-	-	1	-	-	1	1
2000.....	15	-	-	-	-	-	-	-	-	-	-	3
1999.....	18	-	-	-	-	-	-	-	1	-	-	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Unspecified fall (W19)												
Asian or Pacific Islander 2/, both sexes												
2004.....	144	-	1	-	-	-	-	1	-	-	4	5
2003.....	128	-	2	-	-	-	-	-	-	2	3	2
2002.....	122	-	1	-	-	-	-	-	1	1	6	2
2001.....	109	1	-	-	-	-	-	-	1	1	-	3
2000.....	109	-	-	-	-	-	-	1	-	2	2	2
1999.....	112	-	-	-	-	-	-	-	-	1	4	2
Asian or Pacific Islander 2/, male												
2004.....	75	-	-	-	-	-	-	1	-	-	3	5
2003.....	73	-	1	-	-	-	-	-	-	1	2	2
2002.....	72	-	1	-	-	-	-	-	1	1	5	1
2001.....	65	1	-	-	-	-	-	-	1	1	-	2
2000.....	55	-	-	-	-	-	-	1	-	2	-	2
1999.....	72	-	-	-	-	-	-	-	-	-	3	-
Asian or Pacific Islander 2/, female												
2004.....	69	-	1	-	-	-	-	-	-	-	1	-
2003.....	55	-	1	-	-	-	-	-	-	1	1	-
2002.....	50	-	-	-	-	-	-	-	-	-	1	1
2001.....	44	-	-	-	-	-	-	-	-	-	-	1
2000.....	54	-	-	-	-	-	-	-	-	-	2	-
1999.....	40	-	-	-	-	-	-	-	-	1	1	2

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Accidental discharge of firearms (W32-W34)												
All races, both sexes												
2004.....	649	1	14	13	35	80	92	58	42	44	53	41
2003.....	730	-	7	13	36	95	105	74	46	57	61	48
2002.....	762	1	11	14	34	107	103	75	68	52	71	52
2001.....	802	-	15	18	39	110	96	71	51	70	76	61
2000.....	776	1	18	18	49	107	95	74	57	75	78	38
1999.....	824	-	12	19	57	126	125	76	67	67	69	51
All races, male												
2004.....	573	-	11	9	31	74	86	53	39	40	47	34
2003.....	656	-	7	11	35	88	94	69	41	50	55	43
2002.....	667	1	7	10	29	101	94	70	59	42	56	45
2001.....	690	-	12	14	35	102	91	63	40	57	64	51
2000.....	671	-	11	16	46	97	86	65	44	60	65	31
1999.....	707	-	7	13	47	115	114	62	60	56	56	46
All races, female												
2004.....	76	1	3	4	4	6	6	5	3	4	6	7
2003.....	74	-	-	2	1	7	11	5	5	7	6	5
2002.....	95	-	4	4	5	6	9	5	9	10	15	7
2001.....	112	-	3	4	4	8	5	8	11	13	12	10
2000.....	105	1	7	2	3	10	9	9	13	15	13	7
1999.....	117	-	5	6	10	11	11	14	7	11	13	5
White, both sexes												
2004.....	504	1	6	6	31	57	60	39	26	35	46	38
2003.....	586	-	4	10	28	72	75	47	36	49	55	39
2002.....	591	1	9	10	26	69	73	51	49	39	63	44
2001.....	675	-	10	14	31	79	73	55	48	64	63	57
2000.....	611	1	11	11	41	76	65	52	42	60	68	30
1999.....	649	-	5	9	48	88	86	62	53	56	60	45

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Accidental discharge of firearms (W32-W34)												
White, male												
2004.....	443	-	5	4	27	53	56	35	24	31	40	32
2003.....	525	-	4	8	27	66	66	43	32	44	51	34
2002.....	510	1	5	6	22	64	67	46	42	32	48	38
2001.....	580	-	8	11	30	72	69	48	39	52	55	47
2000.....	530	-	6	10	38	70	59	44	31	49	58	26
1999.....	554	-	3	4	42	80	78	49	47	46	48	41
White, female												
2004.....	61	1	1	2	4	4	4	4	2	4	6	6
2003.....	61	-	-	2	1	6	9	4	4	5	4	5
2002.....	81	-	4	4	4	5	6	5	7	7	15	6
2001.....	95	-	2	3	1	7	4	7	9	12	8	10
2000.....	81	1	5	1	3	6	6	8	11	11	10	4
1999.....	95	-	2	5	6	8	8	13	6	10	12	4
Black, both sexes												
2004.....	129	-	7	7	2	19	28	19	14	7	6	3
2003.....	128	-	3	3	6	22	28	26	8	6	6	4
2002.....	155	-	2	3	6	36	25	23	17	13	7	8
2001.....	112	-	5	2	6	26	23	14	3	4	12	4
2000.....	137	-	5	5	7	28	24	18	14	13	6	7
1999.....	156	-	7	9	8	36	31	12	11	9	9	6
Black, male												
2004.....	117	-	6	5	2	19	26	18	13	7	6	2
2003.....	116	-	3	3	6	21	26	25	8	4	4	4
2002.....	142	-	2	3	5	35	23	23	15	10	7	7
2001.....	99	-	4	1	5	25	22	13	1	4	9	4
2000.....	117	-	3	5	7	25	21	17	12	9	5	4
1999.....	135	-	4	8	4	33	28	12	10	8	8	5

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Accidental discharge of firearms (W32-W34)												
Black, female												
2004.....	12	-	1	2	-	-	2	1	1	-	-	1
2003.....	12	-	-	-	-	1	2	1	-	2	2	-
2002.....	13	-	-	-	1	1	2	-	2	3	-	1
2001.....	13	-	1	1	1	1	1	1	2	-	3	-
2000.....	20	-	2	-	-	3	3	1	2	4	1	3
1999.....	21	-	3	1	4	3	3	-	1	1	1	1
American Indian 1/, both sexes												
2004.....	13	-	1	-	2	4	3	-	1	1	1	-
2003.....	13	-	-	-	2	1	2	1	1	2	-	3
2002.....	9	-	-	1	1	2	1	1	1	-	-	-
2001.....	9	-	-	2	1	3	-	1	-	1	1	-
2000.....	14	-	2	1	1	3	3	1	-	-	1	1
1999.....	14	-	-	-	1	1	8	1	2	1	-	-
American Indian 1/, male												
2004.....	10	-	-	-	2	2	3	-	1	1	1	-
2003.....	12	-	-	-	2	1	2	1	-	2	-	3
2002.....	9	-	-	1	1	2	1	1	1	-	-	-
2001.....	6	-	-	2	-	3	-	1	-	-	-	-
2000.....	12	-	2	1	1	2	3	1	-	-	-	1
1999.....	13	-	-	-	1	1	8	-	2	1	-	-
American Indian 1/, female												
2004.....	3	-	1	-	-	2	-	-	-	-	-	-
2003.....	1	-	-	-	-	-	-	-	1	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	3	-	-	-	1	-	-	-	-	1	1	-
2000.....	2	-	-	-	-	1	-	-	-	-	1	-
1999.....	1	-	-	-	-	-	-	1	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Accidental explosion (W39-W40)

All races, both sexes

2004.....	150	-	2	1	1	4	9	8	6	18	24	17
2003.....	158	1	1	1	2	5	16	11	10	15	19	13
2002.....	142	-	2	-	4	15	7	8	17	19	12	12
2001.....	144	-	-	-	-	5	12	7	14	15	14	6
2000.....	172	1	2	2	2	5	10	16	15	11	24	24
1999.....	173	-	-	1	5	5	10	13	14	15	25	19

All races, male

2004.....	125	-	1	1	1	4	9	7	5	14	22	16
2003.....	138	1	1	-	1	4	14	11	9	15	17	13
2002.....	126	-	2	-	3	3	15	6	7	17	19	11
2001.....	117	-	-	-	-	4	10	5	14	14	11	6
2000.....	143	-	2	-	2	4	10	14	15	8	23	20
1999.....	146	-	-	-	5	5	9	11	12	13	23	16

All races, female

2004.....	25	-	1	-	-	-	-	1	1	4	2	1
2003.....	20	-	-	1	1	1	2	-	1	-	2	-
2002.....	16	-	-	-	1	1	-	1	1	-	-	1
2001.....	27	-	-	-	-	1	2	2	-	1	3	-
2000.....	29	1	-	2	-	1	-	2	-	3	1	4
1999.....	27	-	-	1	-	-	1	2	2	2	2	3

White, both sexes

2004.....	132	-	1	1	1	4	9	6	6	17	17	14
2003.....	146	1	1	1	2	5	14	11	9	15	19	13
2002.....	128	-	2	-	3	4	13	5	8	14	18	12
2001.....	127	-	-	-	-	5	11	7	13	14	11	5
2000.....	152	1	1	2	2	5	9	14	15	10	24	18
1999.....	156	-	-	1	5	5	9	13	11	15	21	17

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Accidental explosion (W39-W40)												
White, male												
2004.....	109	-	1	1	1	4	9	5	5	13	15	13
2003.....	130	1	1	-	1	4	12	11	9	15	17	13
2002.....	112	-	2	-	2	3	13	4	7	14	18	11
2001.....	102	-	-	-	-	4	9	5	13	13	9	5
2000.....	126	-	1	-	2	4	9	12	15	7	23	15
1999.....	132	-	-	-	5	5	8	11	9	13	19	14
White, female												
2004.....	23	-	-	-	-	-	-	1	1	4	2	1
2003.....	16	-	-	1	1	1	2	-	-	-	2	-
2002.....	16	-	-	-	1	1	-	1	1	-	-	1
2001.....	25	-	-	-	-	1	2	2	-	1	2	-
2000.....	26	1	-	2	-	1	-	2	-	3	1	3
1999.....	24	-	-	1	-	-	1	2	2	2	2	3
Black, both sexes												
2004.....	15	-	-	-	-	-	-	2	-	-	7	2
2003.....	10	-	-	-	-	-	2	-	1	-	-	-
2002.....	11	-	-	-	1	-	1	2	-	2	1	-
2001.....	16	-	-	-	-	-	1	-	1	1	3	1
2000.....	17	-	1	-	-	-	-	-	-	1	-	6
1999.....	12	-	-	-	-	-	1	-	3	-	2	2
Black, male												
2004.....	14	-	-	-	-	-	-	2	-	-	7	2
2003.....	8	-	-	-	-	-	2	-	-	-	-	-
2002.....	11	-	-	-	1	-	1	2	-	2	1	-
2001.....	14	-	-	-	-	-	1	-	1	1	2	1
2000.....	14	-	1	-	-	-	-	-	-	1	-	5
1999.....	11	-	-	-	-	-	1	-	3	-	2	2

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Accidental drowning and submersion (W65-W74)												
All races, both sexes												
2004.....	3,308	62	430	131	138	304	270	220	165	198	237	240
2003.....	3,306	58	456	126	142	280	287	191	165	217	245	211
2002.....	3,447	63	454	159	162	320	309	227	206	263	263	205
2001.....	3,281	68	458	168	165	322	274	212	162	221	241	214
2000.....	3,482	75	493	201	174	371	275	213	206	250	230	182
1999.....	3,529	68	490	192	177	359	288	212	234	250	225	198
All races, male												
2004.....	2,594	38	265	88	106	277	245	191	141	172	195	200
2003.....	2,632	34	301	90	109	262	263	177	146	186	206	173
2002.....	2,761	40	312	126	118	288	290	201	179	223	216	164
2001.....	2,560	30	291	120	117	295	249	186	142	181	195	182
2000.....	2,735	43	316	137	130	339	244	184	180	212	194	156
1999.....	2,794	39	310	139	136	330	262	193	204	215	187	166
All races, female												
2004.....	714	24	165	43	32	27	25	29	24	26	42	40
2003.....	674	24	155	36	33	18	24	14	19	31	39	38
2002.....	686	23	142	33	44	32	19	26	27	40	47	41
2001.....	721	38	167	48	48	27	25	26	20	40	46	32
2000.....	747	32	177	64	44	32	31	29	26	38	36	26
1999.....	735	29	180	53	41	29	26	19	30	35	38	32
White, both sexes												
2004.....	2,565	51	368	77	65	196	211	164	131	156	194	203
2003.....	2,622	44	386	85	75	201	218	144	125	177	195	173
2002.....	2,624	49	368	88	76	220	235	167	163	201	217	158
2001.....	2,515	52	397	97	78	207	217	150	122	173	188	170
2000.....	2,683	47	415	125	93	253	207	162	160	205	180	147
1999.....	2,715	46	416	106	80	247	215	151	179	205	188	175

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	-------------	-----------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Accidental drowning and submersion (W65-W74)

White, male

2004.....	2,004	29	225	53	49	179	190	143	110	132	163	167
2003.....	2,067	24	252	61	56	188	201	134	108	148	166	138
2002.....	2,093	30	251	68	52	198	219	147	144	167	178	125
2001.....	1,936	25	257	61	54	191	193	128	104	138	154	145
2000.....	2,105	28	268	84	67	231	182	141	143	172	151	125
1999.....	2,120	26	262	71	61	226	193	135	158	172	155	143

White, female

2004.....	561	22	143	24	16	17	21	21	21	24	31	36
2003.....	555	20	134	24	19	13	17	10	17	29	29	35
2002.....	531	19	117	20	24	22	16	20	19	34	39	33
2001.....	579	27	140	36	24	16	24	22	18	35	34	25
2000.....	578	19	147	41	26	22	25	21	17	33	29	22
1999.....	595	20	154	35	19	21	22	16	21	33	33	32

Black, both sexes

2004.....	533	9	49	40	64	79	39	37	24	31	31	30
2003.....	480	13	54	33	56	58	43	30	31	24	27	23
2002.....	613	11	63	58	78	72	57	42	33	43	31	33
2001.....	579	12	48	56	78	90	44	33	30	35	37	34
2000.....	593	26	56	67	69	86	48	34	24	36	43	23
1999.....	634	21	60	76	83	85	53	48	40	32	28	17

Black, male

2004.....	426	7	37	26	49	71	37	30	21	29	24	27
2003.....	407	9	39	23	44	55	40	27	29	23	25	22
2002.....	509	7	44	46	62	70	55	39	26	38	28	28
2001.....	478	3	28	47	56	83	44	33	29	31	29	27
2000.....	478	14	35	48	54	81	45	29	23	32	37	21
1999.....	530	12	40	60	65	78	51	45	35	30	25	17

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Accidental drowning and submersion (W65-W74)												
Black, female												
2004.....	107	2	12	14	15	8	2	7	3	2	7	3
2003.....	73	4	15	10	12	3	3	3	2	1	2	1
2002.....	104	4	19	12	16	2	2	3	7	5	3	5
2001.....	101	9	20	9	22	7	-	-	1	4	8	7
2000.....	115	12	21	19	15	5	3	5	1	4	6	2
1999.....	104	9	20	16	18	7	2	3	5	2	3	-
American Indian 1/, both sexes												
2004.....	57	1	4	3	6	4	8	4	2	6	4	5
2003.....	62	1	6	4	4	4	6	1	1	10	11	8
2002.....	66	-	8	2	1	8	4	10	2	12	8	5
2001.....	52	2	6	6	3	3	2	8	5	4	5	3
2000.....	79	2	10	7	4	12	6	2	12	4	3	5
1999.....	68	-	7	2	3	6	5	5	10	7	6	4
American Indian 1/, male												
2004.....	46	1	1	2	5	3	7	4	2	6	3	4
2003.....	51	1	4	4	3	3	6	1	1	9	7	8
2002.....	53	-	6	1	1	7	4	9	2	11	5	4
2001.....	46	1	3	6	3	3	2	8	5	3	4	3
2000.....	61	1	6	4	4	11	4	2	7	4	3	4
1999.....	55	-	4	2	1	6	5	5	6	7	5	4
American Indian 1/, female												
2004.....	11	-	3	1	1	1	1	-	-	-	1	1
2003.....	11	-	2	-	1	1	-	-	-	1	4	-
2002.....	13	-	2	1	-	1	-	1	-	1	3	1
2001.....	6	1	3	-	-	-	-	-	-	1	1	-
2000.....	18	1	4	3	-	1	2	-	5	-	-	1
1999.....	13	-	3	-	2	-	-	-	4	-	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Accidental drowning and submersion (W65-W74)												
Asian or Pacific Islander 2/, both sexes												
2004.....	153	1	9	11	3	25	12	15	8	5	8	2
2003.....	142	-	10	4	7	17	20	16	8	6	12	7
2002.....	144	3	15	11	7	20	13	8	8	7	7	9
2001.....	135	2	7	9	6	22	11	21	5	9	11	7
2000.....	127	-	12	2	8	20	14	15	10	5	4	7
1999.....	112	1	7	8	11	21	15	8	5	6	3	2
Asian or Pacific Islander 2/, male												
2004.....	118	1	2	7	3	24	11	14	8	5	5	2
2003.....	107	-	6	2	6	16	16	15	8	6	8	5
2002.....	106	3	11	11	3	13	12	6	7	7	5	7
2001.....	100	1	3	6	4	18	10	17	4	9	8	7
2000.....	91	-	7	1	5	16	13	12	7	4	3	6
1999.....	89	1	4	6	9	20	13	8	5	6	2	2
Asian or Pacific Islander 2/, female												
2004.....	35	-	7	4	-	1	1	1	-	-	3	-
2003.....	35	-	4	2	1	1	4	1	-	-	4	2
2002.....	38	-	4	-	4	7	1	2	1	-	2	2
2001.....	35	1	4	3	2	4	1	4	1	-	3	-
2000.....	36	-	5	1	3	4	1	3	3	1	1	1
1999.....	23	-	3	2	2	1	2	-	-	-	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Accidental inhalation and ingestion of food or other objects causing obstruction of respiratory tract (W78-W80)

All races, both sexes

2004.....	4,470	58	70	19	12	30	38	37	51	59	113	152
2003.....	4,272	61	88	15	8	21	35	41	48	74	114	155
2002.....	4,128	61	72	17	18	19	33	40	42	67	100	117
2001.....	4,185	62	74	16	17	19	36	23	34	73	103	163
2000.....	4,313	77	92	13	11	20	25	37	57	76	100	140
1999.....	3,885	63	88	24	22	17	24	27	52	59	100	107

All races, male

2004.....	2,264	47	45	10	7	21	21	26	29	32	64	94
2003.....	2,222	32	57	7	5	14	21	27	35	50	63	91
2002.....	2,114	41	49	8	13	13	21	27	28	39	66	73
2001.....	2,113	42	45	7	9	11	23	14	22	40	71	102
2000.....	2,152	44	56	7	7	13	18	24	37	55	58	95
1999.....	1,917	37	60	14	11	9	18	17	35	38	67	55

All races, female

2004.....	2,206	11	25	9	5	9	17	11	22	27	49	58
2003.....	2,050	29	31	8	3	7	14	14	13	24	51	64
2002.....	2,014	20	23	9	5	6	12	13	14	28	34	44
2001.....	2,072	20	29	9	8	8	13	9	12	33	32	61
2000.....	2,161	33	36	6	4	7	7	13	20	21	42	45
1999.....	1,968	26	28	10	11	8	6	10	17	21	33	52

White, both sexes

2004.....	3,867	45	41	16	7	25	32	30	37	49	100	134
2003.....	3,685	42	65	10	7	17	27	25	40	57	91	133
2002.....	3,521	43	47	9	13	15	22	34	34	56	83	96
2001.....	3,550	41	54	13	13	17	26	18	26	59	83	128
2000.....	3,695	51	62	10	9	12	18	27	44	65	85	115
1999.....	3,293	43	55	16	16	14	20	18	37	47	86	84

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Accidental inhalation and ingestion of food or other objects causing obstruction of respiratory tract (W78-W80)

White, male

2004.....	1,955	38	27	8	5	17	18	22	18	26	55	83
2003.....	1,908	20	44	5	4	10	16	19	28	38	50	75
2002.....	1,796	30	34	5	9	10	16	23	22	32	55	64
2001.....	1,762	29	33	4	6	9	17	10	17	34	56	80
2000.....	1,820	29	44	6	6	8	12	19	28	48	51	77
1999.....	1,607	26	39	9	6	6	15	12	25	29	56	46

White, female

2004.....	1,912	7	14	8	2	8	14	8	19	23	45	51
2003.....	1,777	22	21	5	3	7	11	6	12	19	41	58
2002.....	1,725	13	13	4	4	5	6	11	12	24	28	32
2001.....	1,788	12	21	9	7	8	9	8	9	25	27	48
2000.....	1,875	22	18	4	3	4	6	8	16	17	34	38
1999.....	1,686	17	16	7	10	8	5	6	12	18	30	38

Black, both sexes

2004.....	517	10	22	3	3	5	6	7	11	10	12	17
2003.....	486	18	18	4	1	4	8	15	7	12	22	17
2002.....	546	16	21	8	5	3	10	5	8	9	15	19
2001.....	559	20	19	2	3	2	10	5	8	14	17	31
2000.....	543	23	24	3	2	8	6	10	11	10	13	20
1999.....	540	19	28	7	6	2	4	7	14	10	14	23

Black, male

2004.....	262	6	11	2	2	4	3	4	8	6	8	10
2003.....	259	12	11	1	1	4	5	7	7	9	12	14
2002.....	286	11	13	3	4	2	5	3	6	5	9	8
2001.....	308	13	11	2	3	2	6	4	5	6	13	19
2000.....	283	12	9	1	1	5	5	5	7	6	6	13
1999.....	285	10	18	5	5	2	3	5	9	7	11	9

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Accidental inhalation and ingestion of food or other objects causing obstruction of respiratory tract (W78-W80)

Black, female

2004.....	255	4	11	1	1	1	3	3	3	4	4	7
2003.....	227	6	7	3	-	-	3	8	-	3	10	3
2002.....	260	5	8	5	1	1	5	2	2	4	6	11
2001.....	251	7	8	-	-	-	4	1	3	8	4	12
2000.....	260	11	15	2	1	3	1	5	4	4	7	7
1999.....	255	9	10	2	1	-	1	2	5	3	3	14

American Indian 1/, both sexes

2004.....	28	-	5	-	1	-	-	-	2	-	-	1
2003.....	29	-	1	1	-	-	-	-	1	4	1	1
2002.....	16	-	-	-	-	-	-	1	-	1	-	2
2001.....	26	-	-	1	-	-	-	-	-	-	2	2
2000.....	31	3	1	-	-	-	-	-	2	1	1	5
1999.....	18	-	-	1	-	1	-	2	-	2	-	-

American Indian 1/, male

2004.....	19	-	5	-	-	-	-	-	2	-	-	1
2003.....	17	-	1	1	-	-	-	-	-	3	1	-
2002.....	7	-	-	-	-	-	-	1	-	1	-	1
2001.....	14	-	-	1	-	-	-	-	-	-	1	2
2000.....	25	3	1	-	-	-	-	-	2	1	-	5
1999.....	8	-	-	-	-	1	-	-	-	2	-	-

American Indian 1/, female

2004.....	9	-	-	-	1	-	-	-	-	-	-	-
2003.....	12	-	-	-	-	-	-	-	1	1	-	1
2002.....	9	-	-	-	-	-	-	-	-	-	-	1
2001.....	12	-	-	-	-	-	-	-	-	-	1	-
2000.....	6	-	-	-	-	-	-	-	-	-	1	-
1999.....	10	-	-	1	-	-	-	2	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other accidental and unspecified threats to breathing (W75-W77,W81-W84)

All races, both sexes

2004.....	1,421	667	55	26	56	38	47	37	55	67	76	71
2003.....	1,307	558	71	22	36	30	48	58	48	79	73	55
2002.....	1,389	575	67	23	52	49	54	47	61	71	67	65
2001.....	1,370	552	64	28	51	46	44	47	52	80	88	73
2000.....	1,335	449	59	32	61	50	52	48	62	88	78	47
1999.....	1,618	409	74	30	56	49	49	52	47	64	79	74

All races, male

2004.....	956	370	36	21	50	37	43	26	44	54	68	57
2003.....	909	334	44	17	33	29	47	48	40	65	58	47
2002.....	934	323	39	14	43	45	47	38	53	60	53	55
2001.....	929	316	38	20	43	37	40	39	48	69	73	63
2000.....	931	258	36	19	54	44	48	42	53	81	63	36
1999.....	1,040	244	36	23	53	43	43	49	41	55	66	62

All races, female

2004.....	465	297	19	5	6	1	4	11	11	13	8	14
2003.....	398	224	27	5	3	1	1	10	8	14	15	8
2002.....	455	252	28	9	9	4	7	9	8	11	14	10
2001.....	441	236	26	8	8	9	4	8	4	11	15	10
2000.....	404	191	23	13	7	6	4	6	9	7	15	11
1999.....	578	165	38	7	3	6	6	3	6	9	13	12

White, both sexes

2004.....	1,101	420	46	20	46	36	46	36	48	62	67	64
2003.....	1,010	346	57	16	29	28	45	50	44	72	69	46
2002.....	1,055	334	52	17	47	45	51	41	57	63	59	57
2001.....	1,059	341	45	26	44	39	43	41	45	74	78	65
2000.....	1,053	278	50	24	49	41	45	41	54	77	69	45
1999.....	1,318	258	59	24	47	45	43	44	43	58	71	65

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other accidental and unspecified threats to breathing (W75-W77,W81-W84)

White, male

2004.....	770	243	32	16	40	36	42	25	38	50	60	50
2003.....	720	210	35	12	26	27	44	44	36	59	54	41
2002.....	739	197	29	11	39	42	44	33	50	52	48	49
2001.....	739	192	26	18	38	32	39	35	41	64	67	56
2000.....	752	164	30	13	45	37	41	35	46	72	55	35
1999.....	847	149	28	20	44	39	37	43	37	53	61	54

White, female

2004.....	331	177	14	4	6	-	4	11	10	12	7	14
2003.....	290	136	22	4	3	1	1	6	8	13	15	5
2002.....	316	137	23	6	8	3	7	8	7	11	11	8
2001.....	320	149	19	8	6	7	4	6	4	10	11	9
2000.....	301	114	20	11	4	4	4	6	8	5	14	10
1999.....	471	109	31	4	3	6	6	1	6	5	10	11

Black, both sexes

2004.....	275	221	7	4	6	2	1	1	5	4	6	6
2003.....	260	198	9	6	6	1	1	5	3	7	4	4
2002.....	292	224	11	3	5	4	2	4	3	5	2	6
2001.....	266	191	16	2	5	5	-	5	5	4	7	6
2000.....	244	153	7	8	10	9	6	6	5	7	7	2
1999.....	260	135	11	4	6	2	5	7	4	4	8	7

Black, male

2004.....	158	115	3	3	6	1	1	1	4	3	5	6
2003.....	167	115	7	5	6	1	1	4	3	6	4	3
2002.....	166	116	7	1	4	3	2	3	3	5	-	6
2001.....	163	115	10	2	3	3	-	3	5	3	4	5
2000.....	156	83	6	6	9	7	6	6	5	6	6	1
1999.....	163	83	6	3	6	2	5	5	4	1	5	7

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other accidental and unspecified threats to breathing (W75-W77,W81-W84)

Asian or Pacific Islander 2/, both sexes

2004.....	20	11	2	1	2	-	-	-	1	-	-	1
2003.....	16	4	3	-	1	-	1	-	1	-	-	2
2002.....	18	9	-	1	-	-	-	-	-	1	2	2
2001.....	15	4	-	-	1	2	1	1	2	1	-	1
2000.....	21	11	1	-	1	-	1	-	1	3	-	-
1999.....	25	9	3	1	3	2	1	-	-	2	-	-

Asian or Pacific Islander 2/, male

2004.....	11	4	1	1	2	-	-	-	1	-	-	1
2003.....	8	2	1	-	1	-	1	-	1	-	-	-
2002.....	10	5	-	-	-	-	-	-	-	1	1	-
2001.....	12	2	-	-	1	2	1	1	2	1	-	1
2000.....	14	7	-	-	-	-	1	-	1	3	-	-
1999.....	20	8	1	-	3	2	1	-	-	1	-	-

Asian or Pacific Islander 2/, female

2004.....	9	7	1	-	-	-	-	-	-	-	-	-
2003.....	8	2	2	-	-	-	-	-	-	-	-	2
2002.....	8	4	-	1	-	-	-	-	-	-	1	2
2001.....	3	2	-	-	-	-	-	-	-	-	-	-
2000.....	7	4	1	-	1	-	-	-	-	-	-	-
1999.....	5	1	2	1	-	-	-	-	-	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Accidental exposure to electric current (W85-W87)

All races, both sexes

2004.....	382	1	5	5	7	16	50	39	50	37	40	50
2003.....	376	5	3	4	8	20	44	51	37	47	43	39
2002.....	431	1	5	7	8	32	50	49	46	54	61	49
2001.....	409	1	4	6	10	18	43	42	48	57	62	39
2000.....	395	-	2	1	14	26	47	47	48	54	44	47
1999.....	437	1	6	6	7	32	45	46	41	59	60	45

All races, male

2004.....	360	1	4	3	5	15	48	39	48	33	39	48
2003.....	361	4	3	3	6	19	44	50	36	43	43	39
2002.....	403	-	2	3	6	32	46	48	43	52	60	47
2001.....	391	-	4	4	9	17	43	41	48	55	59	38
2000.....	375	-	2	1	10	24	45	44	45	54	43	46
1999.....	414	1	3	5	6	31	43	44	39	58	57	42

All races, female

2004.....	22	-	1	2	2	1	2	-	2	4	1	2
2003.....	15	1	-	1	2	1	-	1	1	4	-	-
2002.....	28	1	3	4	2	-	4	1	3	2	1	2
2001.....	18	1	-	2	1	1	-	1	-	2	3	1
2000.....	20	-	-	-	4	2	2	3	3	-	1	1
1999.....	23	-	3	1	1	1	2	2	2	1	3	3

White, both sexes

2004.....	343	1	3	4	6	16	46	37	47	32	36	40
2003.....	339	2	3	3	7	20	42	44	34	42	37	37
2002.....	390	1	5	6	6	29	45	47	42	50	57	44
2001.....	359	-	2	5	7	17	36	37	42	55	57	32
2000.....	358	-	1	1	11	24	45	40	44	53	42	41
1999.....	394	-	6	6	6	30	42	43	36	56	53	37

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Accidental exposure to electric current (W85-W87)												
White, male												
2004.....	325	1	2	2	5	15	44	37	45	31	35	38
2003.....	326	1	3	2	6	19	42	43	33	38	37	37
2002.....	365	-	2	3	4	29	41	46	39	48	56	42
2001.....	344	-	2	3	7	16	36	36	42	53	54	31
2000.....	341	-	1	1	8	22	43	38	42	53	41	40
1999.....	374	-	3	5	5	29	40	41	35	55	50	34
White, female												
2004.....	18	-	1	2	1	1	2	-	2	1	1	2
2003.....	13	1	-	1	1	1	-	1	1	4	-	-
2002.....	25	1	3	3	2	-	4	1	3	2	1	2
2001.....	15	-	-	2	-	1	-	1	-	2	3	1
2000.....	17	-	-	-	3	2	2	2	2	-	1	1
1999.....	20	-	3	1	1	1	2	2	1	1	3	3
Black, both sexes												
2004.....	32	-	2	1	-	-	4	2	3	3	4	7
2003.....	31	3	-	1	1	-	2	6	1	4	5	2
2002.....	34	-	-	1	2	1	5	2	2	3	4	5
2001.....	35	1	2	-	3	1	3	3	2	-	4	7
2000.....	31	-	1	-	3	-	2	6	4	1	2	6
1999.....	40	1	-	-	1	2	2	3	4	3	7	8
Black, male												
2004.....	30	-	2	1	-	-	4	2	3	1	4	7
2003.....	29	3	-	1	-	-	2	6	1	4	5	2
2002.....	32	-	-	-	2	1	5	2	2	3	4	5
2001.....	32	-	2	-	2	1	3	3	2	-	4	7
2000.....	28	-	1	-	2	-	2	5	3	1	2	6
1999.....	37	1	-	-	1	2	2	3	3	3	7	8

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	-------------	-----------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Accidental exposure to smoke, fire and flames (X00-X09)

All races, both sexes

2004.....	3,229	26	223	168	87	66	120	131	109	142	171	252
2003.....	3,369	30	224	137	78	75	124	118	134	167	228	270
2002.....	3,159	36	221	152	101	86	107	112	154	147	214	213
2001.....	3,309	50	225	164	88	76	133	114	133	190	251	213
2000.....	3,377	37	290	182	84	79	113	120	121	188	214	214
1999.....	3,348	41	302	170	92	88	113	142	140	200	221	198

All races, male

2004.....	1,902	14	120	107	45	42	65	91	74	95	114	160
2003.....	1,948	8	131	72	40	43	78	70	84	115	142	181
2002.....	1,935	21	126	89	62	61	71	75	95	101	150	146
2001.....	1,998	30	142	104	48	45	92	73	88	132	176	151
2000.....	2,025	20	174	109	36	44	77	78	79	133	135	145
1999.....	2,025	21	173	95	52	58	75	93	104	138	162	144

All races, female

2004.....	1,327	12	103	61	42	24	55	40	35	47	57	92
2003.....	1,421	22	93	65	38	32	46	48	50	52	86	89
2002.....	1,224	15	95	63	39	25	36	37	59	46	64	67
2001.....	1,311	20	83	60	40	31	41	41	45	58	75	62
2000.....	1,352	17	116	73	48	35	36	42	42	55	79	69
1999.....	1,323	20	129	75	40	30	38	49	36	62	59	54

White, both sexes

2004.....	2,334	15	139	85	51	50	90	92	87	112	122	175
2003.....	2,526	22	144	74	41	53	103	88	107	137	180	207
2002.....	2,287	19	127	78	61	61	77	72	118	110	175	167
2001.....	2,387	34	142	102	57	55	102	71	103	139	187	154
2000.....	2,376	23	160	104	54	60	81	83	86	134	157	155
1999.....	2,380	26	178	96	63	70	83	107	108	141	169	145

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Accidental exposure to smoke, fire and flames (X00-X09)

White, male

2004.....	1,375	9	76	51	29	30	52	65	59	72	81	108
2003.....	1,480	7	85	39	19	33	67	52	68	96	118	138
2002.....	1,419	14	70	48	42	44	53	52	74	75	119	117
2001.....	1,436	19	95	62	30	33	73	52	68	94	134	107
2000.....	1,435	11	98	61	24	32	58	54	59	97	98	106
1999.....	1,466	14	103	52	36	47	56	74	82	97	126	106

White, female

2004.....	959	6	63	34	22	20	38	27	28	40	41	67
2003.....	1,046	15	59	35	22	20	36	36	39	41	62	69
2002.....	868	5	57	30	19	17	24	20	44	35	56	50
2001.....	951	15	47	40	27	22	29	19	35	45	53	47
2000.....	941	12	62	43	30	28	23	29	27	37	59	49
1999.....	914	12	75	44	27	23	27	33	26	44	43	39

Black, both sexes

2004.....	799	11	73	72	33	14	25	32	21	25	41	70
2003.....	775	7	76	60	34	20	19	25	22	28	40	61
2002.....	774	15	88	64	32	21	27	31	30	29	31	40
2001.....	846	15	82	60	28	14	28	38	28	45	59	56
2000.....	930	14	121	74	27	16	28	31	34	44	55	52
1999.....	892	15	111	74	28	16	24	29	24	55	47	46

Black, male

2004.....	475	5	38	50	14	10	12	21	14	18	27	49
2003.....	433	1	42	31	19	9	10	16	12	18	21	41
2002.....	461	7	53	37	17	14	17	17	18	21	24	25
2001.....	511	10	46	40	16	8	16	17	18	34	39	42
2000.....	551	9	71	46	11	10	17	19	20	30	35	35
1999.....	510	7	63	43	15	9	15	15	15	39	32	32

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Accidental exposure to smoke, fire and flames (X00-X09)

Black, female

2004.....	324	6	35	22	19	4	13	11	7	7	14	21
2003.....	342	6	34	29	15	11	9	9	10	10	19	20
2002.....	313	8	35	27	15	7	10	14	12	8	7	15
2001.....	335	5	36	20	12	6	12	21	10	11	20	14
2000.....	379	5	50	28	16	6	11	12	14	14	20	17
1999.....	382	8	48	31	13	7	9	14	9	16	15	14

American Indian 1/, both sexes

2004.....	47	-	7	4	1	1	2	3	-	2	5	4
2003.....	33	-	4	1	2	1	1	3	2	1	6	-
2002.....	56	1	3	3	5	1	2	8	3	5	6	5
2001.....	45	1	1	1	3	4	1	3	1	6	3	1
2000.....	37	-	5	2	2	3	1	6	1	5	2	1
1999.....	40	-	5	-	-	1	3	1	6	3	1	4

American Indian 1/, male

2004.....	28	-	5	2	1	1	1	2	-	2	3	2
2003.....	16	-	4	1	1	-	1	1	1	1	3	-
2002.....	35	-	1	-	3	-	1	5	3	3	5	3
2001.....	30	1	1	1	2	2	1	2	1	4	1	1
2000.....	23	-	3	1	1	2	1	5	-	2	2	1
1999.....	26	-	3	-	-	1	3	-	5	2	-	3

American Indian 1/, female

2004.....	19	-	2	2	-	-	1	1	-	-	2	2
2003.....	17	-	-	-	1	1	-	2	1	-	3	-
2002.....	21	1	2	3	2	1	1	3	-	2	1	2
2001.....	15	-	-	-	1	2	-	1	-	2	2	-
2000.....	14	-	2	1	1	1	-	1	1	3	-	-
1999.....	14	-	2	-	-	-	-	1	1	1	1	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Accidental exposure to smoke, fire and flames (X00-X09)												
Asian or Pacific Islander 2/, both sexes												
2004.....	49	-	4	7	2	1	3	4	1	3	3	3
2003.....	35	1	-	2	1	1	1	2	3	1	2	2
2002.....	42	1	3	7	3	3	1	1	3	3	2	1
2001.....	31	-	-	1	-	3	2	2	1	-	2	2
2000.....	34	-	4	2	1	-	3	-	-	5	-	6
1999.....	36	-	8	-	1	1	3	5	2	1	4	3
Asian or Pacific Islander 2/, male												
2004.....	24	-	1	4	1	1	-	3	1	3	3	1
2003.....	19	-	-	1	1	1	-	1	3	-	-	2
2002.....	20	-	2	4	-	3	-	1	-	2	2	1
2001.....	21	-	-	1	-	2	2	2	1	-	2	1
2000.....	16	-	2	1	-	-	1	-	-	4	-	3
1999.....	23	-	4	-	1	1	1	4	2	-	4	3
Asian or Pacific Islander 2/, female												
2004.....	25	-	3	3	1	-	3	1	-	-	-	2
2003.....	16	1	-	1	-	-	1	1	-	1	2	-
2002.....	22	1	1	3	3	-	1	-	3	1	-	-
2001.....	10	-	-	-	-	1	-	-	-	-	-	1
2000.....	18	-	2	1	1	-	2	-	-	1	-	3
1999.....	13	-	4	-	-	-	2	1	-	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Accidental contact with heat and hot substances (X10-X19)												
All races, both sexes												
2004.....	93	2	5	1	-	-	-	2	3	3	4	5
2003.....	88	2	5	-	-	-	1	3	1	3	4	5
2002.....	102	4	5	1	-	-	3	1	2	1	6	8
2001.....	114	-	5	-	-	-	5	-	3	2	5	5
2000.....	110	2	7	1	-	-	1	4	1	6	1	5
1999.....	123	3	6	1	-	-	2	3	3	2	7	6
All races, male												
2004.....	52	1	2	1	-	-	-	2	2	2	2	2
2003.....	45	1	4	-	-	-	-	2	1	1	3	2
2002.....	52	-	4	1	-	-	2	1	1	1	4	6
2001.....	58	-	1	-	-	-	3	-	2	2	5	5
2000.....	57	1	4	1	-	-	-	3	1	3	-	4
1999.....	61	1	4	1	-	-	2	1	3	2	5	3
All races, female												
2004.....	41	1	3	-	-	-	-	-	1	1	2	3
2003.....	43	1	1	-	-	-	1	1	-	2	1	3
2002.....	50	4	1	-	-	-	1	-	1	-	2	2
2001.....	56	-	4	-	-	-	2	-	1	-	-	-
2000.....	53	1	3	-	-	-	1	1	-	3	1	1
1999.....	62	2	2	-	-	-	-	2	-	-	2	3
White, both sexes												
2004.....	68	1	3	-	-	-	-	2	2	2	4	3
2003.....	66	2	4	-	-	-	1	3	1	3	4	3
2002.....	67	2	2	-	-	-	3	1	1	1	4	4
2001.....	87	-	2	-	-	-	4	-	2	1	3	2
2000.....	79	1	4	1	-	-	1	4	1	5	-	3
1999.....	87	1	3	1	-	-	2	1	2	2	6	4

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Accidental contact with heat and hot substances (X10-X19)												
White, male												
2004.....	39	1	2	-	-	-	-	2	1	1	2	2
2003.....	32	1	3	-	-	-	-	2	1	1	3	1
2002.....	33	-	1	-	-	-	2	1	1	1	3	4
2001.....	41	-	1	-	-	-	2	-	1	1	3	2
2000.....	42	1	2	1	-	-	-	3	1	2	-	3
1999.....	40	-	2	1	-	-	2	-	2	2	5	1
White, female												
2004.....	29	-	1	-	-	-	-	-	1	1	2	1
2003.....	34	1	1	-	-	-	1	1	-	2	1	2
2002.....	34	2	1	-	-	-	1	-	-	-	1	-
2001.....	46	-	1	-	-	-	2	-	1	-	-	-
2000.....	37	-	2	-	-	-	1	1	-	3	-	-
1999.....	47	1	1	-	-	-	-	1	-	-	1	3
Black, both sexes												
2004.....	23	1	2	1	-	-	-	-	1	1	-	2
2003.....	19	-	1	-	-	-	-	-	-	-	-	2
2002.....	27	2	3	-	-	-	-	-	1	-	2	3
2001.....	25	-	3	-	-	-	-	-	1	1	2	3
2000.....	25	1	2	-	-	-	-	-	-	1	-	1
1999.....	29	1	3	-	-	-	-	2	1	-	1	1
Black, male												
2004.....	11	-	-	1	-	-	-	-	1	1	-	-
2003.....	11	-	1	-	-	-	-	-	-	-	-	1
2002.....	16	-	3	-	-	-	-	-	-	-	1	2
2001.....	15	-	-	-	-	-	-	-	1	1	2	3
2000.....	12	-	1	-	-	-	-	-	-	1	-	-
1999.....	18	-	2	-	-	-	-	1	1	-	-	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Lightning (X33)												
All races, both sexes												
2004.....	46	-	-	-	2	6	4	4	6	4	6	8
2003.....	47	-	1	-	-	4	6	6	8	4	3	3
2002.....	66	-	1	1	1	3	2	9	6	5	8	9
2001.....	44	-	-	1	2	5	5	4	5	5	4	5
2000.....	50	-	-	2	5	4	4	10	3	1	6	4
1999.....	64	-	1	1	4	4	6	4	10	10	7	5
All races, male												
2004.....	36	-	-	-	1	5	4	2	6	4	4	6
2003.....	38	-	1	-	-	3	5	4	7	3	3	1
2002.....	52	-	1	-	1	2	1	6	4	5	7	7
2001.....	38	-	-	1	2	5	4	3	5	4	3	4
2000.....	41	-	-	1	4	3	3	10	3	1	2	3
1999.....	53	-	-	1	1	4	6	4	7	9	5	5
All races, female												
2004.....	10	-	-	-	1	1	-	2	-	-	2	2
2003.....	9	-	-	-	-	1	1	2	1	1	-	2
2002.....	14	-	-	1	-	1	1	3	2	-	1	2
2001.....	6	-	-	-	-	-	1	1	-	1	1	1
2000.....	9	-	-	1	1	1	1	-	-	-	4	1
1999.....	11	-	1	-	3	-	-	-	3	1	2	-
White, both sexes												
2004.....	42	-	-	-	1	5	3	4	6	4	6	8
2003.....	43	-	1	-	-	3	6	5	7	4	2	3
2002.....	57	-	1	-	1	3	2	7	6	4	7	8
2001.....	40	-	-	1	2	3	5	4	5	4	4	4
2000.....	44	-	-	2	5	3	3	10	3	1	5	4
1999.....	57	-	-	1	4	4	5	4	8	9	7	5

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Lightning (X33)												
White, male												
2004.....	33	-	-	-	1	4	3	2	6	4	4	6
2003.....	35	-	1	-	-	3	5	3	6	3	2	1
2002.....	47	-	1	-	1	2	1	6	4	4	7	6
2001.....	34	-	-	1	2	3	4	3	5	3	3	3
2000.....	35	-	-	1	4	2	2	10	3	1	1	3
1999.....	48	-	-	1	1	4	5	4	6	8	5	5
White, female												
2004.....	9	-	-	-	-	1	-	2	-	-	2	2
2003.....	8	-	-	-	-	-	1	2	1	1	-	2
2002.....	10	-	-	-	-	1	1	1	2	-	-	2
2001.....	6	-	-	-	-	-	1	1	-	1	1	1
2000.....	9	-	-	1	1	1	1	-	-	-	4	1
1999.....	9	-	-	-	3	-	-	-	2	1	2	-
Black, both sexes												
2004.....	2	-	-	-	1	-	-	-	-	-	-	-
2003.....	1	-	-	-	-	-	-	-	-	-	1	-
2002.....	6	-	-	1	-	-	-	1	-	-	-	1
2001.....	2	-	-	-	-	-	-	-	-	1	-	1
2000.....	4	-	-	-	-	-	1	-	-	-	1	-
1999.....	3	-	-	-	-	-	1	-	1	-	-	-
Black, male												
2004.....	1	-	-	-	-	-	-	-	-	-	-	-
2003.....	1	-	-	-	-	-	-	-	-	-	1	-
2002.....	4	-	-	-	-	-	-	-	-	-	-	1
2001.....	2	-	-	-	-	-	-	-	-	1	-	1
2000.....	4	-	-	-	-	-	1	-	-	-	1	-
1999.....	3	-	-	-	-	-	1	-	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Lightning (X33)												
Black, female												
2004.....	1	-	-	-	1	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	2	-	-	1	-	-	-	1	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes												
2004.....	1	-	-	-	-	1	-	-	-	-	-	-
2003.....	2	-	-	-	-	1	-	-	1	-	-	-
2002.....	2	-	-	-	-	-	-	-	-	1	1	-
2001.....	2	-	-	-	-	2	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	4	-	1	-	-	-	-	-	1	1	-	-
American Indian 1/, male												
2004.....	1	-	-	-	-	1	-	-	-	-	-	-
2003.....	1	-	-	-	-	-	-	-	1	-	-	-
2002.....	1	-	-	-	-	-	-	-	-	1	-	-
2001.....	2	-	-	-	-	2	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	2	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, female												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	1	-	-	-	-	1	-	-	-	-	-	-
2002.....	1	-	-	-	-	-	-	-	-	-	1	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	2	-	1	-	-	-	-	-	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Earthquake, volcanic eruption, avalanche, landslide and other earth movements (X34-X36)												
All races, both sexes												
2004.....	30	-	1	-	-	1	7	4	5	1	3	3
2003.....	32	-	-	-	-	1	7	1	5	2	6	3
2002.....	31	-	-	-	1	5	-	6	7	7	2	1
2001.....	28	-	-	-	-	1	4	2	8	7	3	-
2000.....	35	-	-	1	1	1	1	10	3	2	7	3
1999.....	46	-	-	1	-	4	10	8	6	7	3	3
All races, male												
2004.....	28	-	1	-	-	1	7	4	5	1	3	2
2003.....	27	-	-	-	-	1	5	1	5	1	6	3
2002.....	28	-	-	-	-	5	-	6	6	6	2	1
2001.....	23	-	-	-	-	1	4	2	7	7	1	-
2000.....	28	-	-	-	1	1	1	8	3	2	4	3
1999.....	39	-	-	-	-	4	8	8	5	5	3	3
All races, female												
2004.....	2	-	-	-	-	-	-	-	-	-	-	1
2003.....	5	-	-	-	-	-	2	-	-	1	-	-
2002.....	3	-	-	-	1	-	-	-	1	1	-	-
2001.....	5	-	-	-	-	-	-	-	1	-	2	-
2000.....	7	-	-	1	-	-	-	2	-	-	3	-
1999.....	7	-	-	1	-	-	2	-	1	2	-	-
White, both sexes												
2004.....	29	-	1	-	-	1	6	4	5	1	3	3
2003.....	31	-	-	-	-	1	7	1	5	2	6	3
2002.....	31	-	-	-	1	5	-	6	7	7	2	1
2001.....	28	-	-	-	-	1	4	2	8	7	3	-
2000.....	33	-	-	1	1	1	1	9	3	2	7	3
1999.....	44	-	-	-	-	4	10	8	6	6	3	3

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Earthquake, volcanic eruption, avalanche, landslide and other earth movements (X34-X36)

Asian or Pacific Islander 2/, both sexes

2004.....	1	-	-	-	-	-	1	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	1	-	-	-	-	-	-	-	-	1	-	-

Asian or Pacific Islander 2/, male

2004.....	1	-	-	-	-	-	1	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-

Asian or Pacific Islander 2/, female

2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	1	-	-	-	-	-	-	-	-	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Cataclysmic storm and flood (X37-X38)

All races, both sexes

2004.....	85	1	2	1	1	3	5	6	2	3	4	8
2003.....	101	3	3	11	3	1	3	6	9	6	11	8
2002.....	72	1	1	-	2	5	1	2	3	8	6	9
2001.....	89	-	6	2	7	1	6	5	6	6	9	2
2000.....	54	2	3	2	-	3	2	5	4	3	2	1
1999.....	144	3	2	3	3	2	6	10	4	14	18	10

All races, male

2004.....	52	-	-	-	1	3	4	5	2	-	3	7
2003.....	57	3	2	7	3	-	1	4	5	1	4	3
2002.....	42	-	-	-	2	2	1	1	1	3	5	7
2001.....	52	-	4	2	4	1	4	1	2	5	4	2
2000.....	27	-	1	2	-	1	2	1	3	1	1	1
1999.....	79	3	-	1	1	2	5	3	2	7	11	4

All races, female

2004.....	33	1	2	1	-	-	1	1	-	3	1	1
2003.....	44	-	1	4	-	1	2	2	4	5	7	5
2002.....	30	1	1	-	-	3	-	1	2	5	1	2
2001.....	37	-	2	-	3	-	2	4	4	1	5	-
2000.....	27	2	2	-	-	2	-	4	1	2	1	-
1999.....	65	-	2	2	2	-	1	7	2	7	7	6

White, both sexes

2004.....	78	1	2	1	1	3	4	5	2	2	3	8
2003.....	84	2	1	7	3	1	2	5	9	5	9	7
2002.....	69	1	1	-	2	4	1	2	3	8	6	9
2001.....	66	-	5	2	4	1	6	3	3	4	7	2
2000.....	40	1	3	2	-	1	2	3	1	3	2	1
1999.....	117	3	1	2	3	2	5	8	3	12	14	7

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cataclysmic storm and flood (X37-X38)												
Black, female												
2004.....	1	-	-	-	-	-	-	-	-	-	-	-
2003.....	3	-	-	-	-	-	1	-	-	-	1	1
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	8	-	-	-	1	-	-	1	1	-	1	-
2000.....	10	1	-	-	-	2	-	2	1	-	-	-
1999.....	7	-	-	1	-	-	1	1	1	-	-	1
American Indian 1/, both sexes												
2004.....	2	-	-	-	-	-	-	-	-	1	-	-
2003.....	2	-	-	2	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	4	-	1	-	-	-	-	-	1	-	1	-
2000.....	1	-	-	-	-	-	-	-	-	-	-	-
1999.....	5	-	1	-	-	-	-	-	-	2	-	-
American Indian 1/, male												
2004.....	1	-	-	-	-	-	-	-	-	-	-	-
2003.....	2	-	-	2	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	3	-	1	-	-	-	-	-	1	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	3	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, female												
2004.....	1	-	-	-	-	-	-	-	-	1	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	1	-	-	-	-	-	-	-	-	-	1	-
2000.....	1	-	-	-	-	-	-	-	-	-	-	-
1999.....	2	-	1	-	-	-	-	-	-	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cataclysmic storm and flood (X37-X38)												
Asian or Pacific Islander 2/, both sexes												
2004.....	1	-	-	-	-	-	-	1	-	-	-	-
2003.....	1	-	1	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	2	-	-	-	1	-	-	1	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	1	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	2	-	-	-	1	-	-	1	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	1	-	-	-	-	-	-	1	-	-	-	-
2003.....	1	-	1	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	1	-	-	-	-	-	-	1	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Accidental poisoning and exposure to noxious substances (X40-X49)												
All races, both sexes												
2004.....	20,950	13	18	8	47	643	1,616	1,747	1,894	2,694	3,750	3,696
2003.....	19,457	20	49	18	43	520	1,479	1,517	1,918	2,674	3,556	3,351
2002.....	17,550	26	31	15	28	486	1,193	1,310	1,806	2,597	3,410	2,946
2001.....	14,078	15	31	18	32	406	956	1,078	1,429	2,307	2,729	2,266
2000.....	12,757	14	32	17	28	351	809	980	1,400	2,159	2,504	1,992
1999.....	12,186	12	34	12	28	260	704	952	1,403	2,082	2,467	1,926
All races, male												
2004.....	13,934	5	13	3	27	494	1,271	1,328	1,345	1,781	2,395	2,360
2003.....	13,176	9	27	4	27	398	1,164	1,145	1,369	1,804	2,362	2,230
2002.....	12,059	21	14	10	18	366	940	1,003	1,316	1,807	2,315	1,960
2001.....	9,885	8	21	7	16	311	769	832	1,030	1,626	1,896	1,610
2000.....	9,138	11	21	12	15	273	647	779	1,041	1,554	1,830	1,455
1999.....	8,887	11	17	6	19	195	559	757	1,071	1,523	1,826	1,456
All races, female												
2004.....	7,016	8	5	5	20	149	345	419	549	913	1,355	1,336
2003.....	6,281	11	22	14	16	122	315	372	549	870	1,194	1,121
2002.....	5,491	5	17	5	10	120	253	307	490	790	1,095	986
2001.....	4,193	7	10	11	16	95	187	246	399	681	833	656
2000.....	3,619	3	11	5	13	78	162	201	359	605	674	537
1999.....	3,299	1	17	6	9	65	145	195	332	559	641	470
White, both sexes												
2004.....	18,042	10	12	5	35	601	1,494	1,592	1,667	2,326	3,255	3,111
2003.....	16,690	12	31	12	23	490	1,390	1,367	1,671	2,278	3,058	2,814
2002.....	14,868	13	19	12	25	452	1,103	1,165	1,580	2,199	2,901	2,407
2001.....	11,662	9	21	14	26	362	866	949	1,227	1,939	2,258	1,801
2000.....	10,508	10	19	10	20	318	732	847	1,184	1,811	2,031	1,568
1999.....	9,883	10	20	4	23	231	612	816	1,149	1,734	1,979	1,493

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	-------------	-----------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Accidental poisoning and exposure to noxious substances (X40-X49)

White, male

2004.....	12,022	3	7	3	22	465	1,180	1,215	1,198	1,558	2,090	2,001
2003.....	11,333	8	16	3	17	379	1,093	1,042	1,206	1,544	2,050	1,870
2002.....	10,213	12	6	8	17	342	870	891	1,161	1,542	1,985	1,609
2001.....	8,232	5	15	6	13	286	694	731	907	1,384	1,600	1,274
2000.....	7,558	8	11	7	11	251	601	674	894	1,330	1,498	1,143
1999.....	7,240	9	9	1	16	173	489	656	895	1,289	1,488	1,125

White, female

2004.....	6,020	7	5	2	13	136	314	377	469	768	1,165	1,110
2003.....	5,357	4	15	9	6	111	297	325	465	734	1,008	944
2002.....	4,655	1	13	4	8	110	233	274	419	657	916	798
2001.....	3,430	4	6	8	13	76	172	218	320	555	658	527
2000.....	2,950	2	8	3	9	67	131	173	290	481	533	425
1999.....	2,643	1	11	3	7	58	123	160	254	445	491	368

Black, both sexes

2004.....	2,498	3	6	3	8	28	91	123	173	307	429	529
2003.....	2,396	7	16	4	14	17	70	107	205	335	442	495
2002.....	2,365	12	11	3	3	19	69	116	190	345	456	494
2001.....	2,169	6	8	4	2	27	68	107	175	326	437	437
2000.....	2,028	4	12	6	3	21	58	115	188	317	429	389
1999.....	2,062	1	12	6	2	18	76	115	221	305	445	411

Black, male

2004.....	1,638	2	6	-	3	18	69	87	115	178	263	324
2003.....	1,609	1	9	-	7	10	54	73	135	224	280	327
2002.....	1,631	8	8	2	1	14	53	89	131	232	294	323
2001.....	1,490	3	5	1	1	15	54	85	108	213	274	320
2000.....	1,433	3	9	5	2	13	37	91	125	206	301	288
1999.....	1,494	1	7	4	2	15	62	87	150	207	312	315

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Accidental poisoning and exposure to noxious substances (X40-X49)

Black, female

2004.....	860	1	-	3	5	10	22	36	58	129	166	205
2003.....	787	6	7	4	7	7	16	34	70	111	162	168
2002.....	734	4	3	1	2	5	16	27	59	113	162	171
2001.....	679	3	3	3	1	12	14	22	67	113	163	117
2000.....	595	1	3	1	1	8	21	24	63	111	128	101
1999.....	568	-	5	2	-	3	14	28	71	98	133	96

American Indian 1/, both sexes

2004.....	262	-	-	-	4	9	19	19	40	45	45	36
2003.....	231	-	-	-	3	10	10	23	26	42	43	26
2002.....	188	1	-	-	-	7	9	16	22	38	33	31
2001.....	149	-	1	-	3	9	9	11	20	28	21	21
2000.....	137	-	-	1	3	6	12	11	19	21	28	23
1999.....	149	1	2	2	2	7	8	13	24	28	28	17

American Indian 1/, male

2004.....	162	-	-	-	2	6	14	15	20	33	25	21
2003.....	141	-	-	-	2	7	9	14	15	26	22	20
2002.....	124	1	-	-	-	4	9	12	14	22	19	20
2001.....	90	-	-	-	1	4	9	8	8	18	13	12
2000.....	84	-	-	-	-	3	4	9	15	10	17	16
1999.....	91	1	1	1	-	4	6	8	18	17	12	12

American Indian 1/, female

2004.....	100	-	-	-	2	3	5	4	20	12	20	15
2003.....	90	-	-	-	1	3	1	9	11	16	21	6
2002.....	64	-	-	-	-	3	-	4	8	16	14	11
2001.....	59	-	1	-	2	5	-	3	12	10	8	9
2000.....	53	-	-	1	3	3	8	2	4	11	11	7
1999.....	58	-	1	1	2	3	2	5	6	11	16	5

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Accidental poisoning and exposure to noxious substances (X40-X49)												
Asian or Pacific Islander 2/, both sexes												
2004.....	148	-	-	-	-	5	12	13	14	16	21	20
2003.....	140	1	2	2	3	3	9	20	16	19	13	16
2002.....	129	-	1	-	-	8	12	13	14	15	20	14
2001.....	98	-	1	-	1	8	13	11	7	14	13	7
2000.....	84	-	1	-	2	6	7	7	9	10	16	12
1999.....	92	-	-	-	1	4	8	8	9	15	15	5
Asian or Pacific Islander 2/, male												
2004.....	112	-	-	-	-	5	8	11	12	12	17	14
2003.....	93	-	2	1	1	2	8	16	13	10	10	13
2002.....	91	-	-	-	-	6	8	11	10	11	17	8
2001.....	73	-	1	-	1	6	12	8	7	11	9	4
2000.....	63	-	1	-	2	6	5	5	7	8	14	8
1999.....	62	-	-	-	1	3	2	6	8	10	14	4
Asian or Pacific Islander 2/, female												
2004.....	36	-	-	-	-	-	4	2	2	4	4	6
2003.....	47	1	-	1	2	1	1	4	3	9	3	3
2002.....	38	-	1	-	-	2	4	2	4	4	3	6
2001.....	25	-	-	-	-	2	1	3	-	3	4	3
2000.....	21	-	-	-	-	-	2	2	2	2	2	4
1999.....	30	-	-	-	-	1	6	2	1	5	1	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Accidental poisoning by and exposure to drugs and other biological substances (X40-X44)												
All races, both sexes												
2004.....	19,838	10	11	3	26	591	1,563	1,688	1,823	2,583	3,625	3,548
2003.....	18,294	16	26	8	19	471	1,411	1,453	1,850	2,558	3,404	3,200
2002.....	16,394	18	18	2	15	439	1,134	1,250	1,721	2,488	3,258	2,816
2001.....	13,024	9	18	5	18	352	885	1,021	1,367	2,180	2,593	2,156
2000.....	11,712	9	10	3	14	291	754	920	1,323	2,056	2,385	1,874
1999.....	11,155	8	16	4	12	208	641	908	1,324	1,971	2,322	1,814
All races, male												
2004.....	13,114	3	8	2	15	456	1,225	1,276	1,290	1,695	2,294	2,245
2003.....	12,297	8	15	1	12	355	1,110	1,094	1,311	1,717	2,237	2,103
2002.....	11,191	15	7	2	11	330	890	950	1,247	1,718	2,192	1,857
2001.....	9,120	4	15	2	9	270	710	790	987	1,530	1,790	1,522
2000.....	8,368	6	8	2	7	223	601	731	972	1,466	1,733	1,372
1999.....	8,111	7	8	-	9	155	508	716	1,013	1,430	1,708	1,368
All races, female												
2004.....	6,724	7	3	1	11	135	338	412	533	888	1,331	1,303
2003.....	5,997	8	11	7	7	116	301	359	539	841	1,167	1,097
2002.....	5,203	3	11	-	4	109	244	300	474	770	1,066	959
2001.....	3,904	5	3	3	9	82	175	231	380	650	803	634
2000.....	3,344	3	2	1	7	68	153	189	351	590	652	502
1999.....	3,044	1	8	4	3	53	133	192	311	541	614	446
White, both sexes												
2004.....	17,112	8	8	2	20	558	1,449	1,543	1,607	2,230	3,151	2,982
2003.....	15,724	9	18	6	12	450	1,328	1,311	1,614	2,183	2,926	2,689
2002.....	13,927	8	10	1	13	410	1,051	1,113	1,511	2,117	2,778	2,307
2001.....	10,820	6	13	4	16	320	809	903	1,174	1,840	2,147	1,707
2000.....	9,649	6	5	2	13	269	688	799	1,118	1,727	1,932	1,470
1999.....	9,062	7	9	2	11	189	561	775	1,093	1,647	1,860	1,406

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Accidental poisoning by and exposure to drugs and other biological substances (X40-X44)

White, male

2004.....	11,325	2	5	2	13	432	1,140	1,170	1,151	1,482	2,002	1,900
2003.....	10,606	7	11	1	9	344	1,044	998	1,156	1,471	1,942	1,769
2002.....	9,506	7	2	1	10	309	825	846	1,105	1,476	1,883	1,528
2001.....	7,606	2	11	2	7	252	647	698	868	1,307	1,512	1,197
2000.....	6,919	4	3	2	6	209	562	633	834	1,259	1,416	1,075
1999.....	6,608	6	5	-	8	139	447	617	852	1,215	1,387	1,055

White, female

2004.....	5,787	6	3	-	7	126	309	373	456	748	1,149	1,082
2003.....	5,118	2	7	5	3	106	284	313	458	712	984	920
2002.....	4,421	1	8	-	3	101	226	267	406	641	895	779
2001.....	3,214	4	2	2	9	68	162	205	306	533	635	510
2000.....	2,730	2	2	-	7	60	126	166	284	468	516	395
1999.....	2,454	1	4	2	3	50	114	158	241	432	473	351

Black, both sexes

2004.....	2,370	2	3	1	4	23	85	119	166	298	418	514
2003.....	2,261	7	8	2	6	13	65	100	197	328	426	478
2002.....	2,204	9	7	1	2	16	66	111	180	324	436	471
2001.....	2,012	3	4	1	1	19	62	100	169	307	418	423
2000.....	1,896	3	4	1	1	19	56	106	182	305	411	374
1999.....	1,920	1	7	1	1	14	69	114	209	288	432	391

Black, male

2004.....	1,551	1	3	-	2	16	64	85	110	172	254	312
2003.....	1,499	1	4	-	3	6	50	66	128	218	267	310
2002.....	1,514	7	5	1	1	12	51	84	123	215	280	306
2001.....	1,390	2	3	-	1	12	50	80	105	200	259	311
2000.....	1,337	2	4	-	1	11	36	85	121	194	287	276
1999.....	1,391	1	3	-	1	12	56	86	142	193	303	300

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Accidental poisoning by and exposure to drugs and other biological substances (X40-X44)

Black, female

2004.....	819	1	-	1	2	7	21	34	56	126	164	202
2003.....	762	6	4	2	3	7	15	34	69	110	159	168
2002.....	690	2	2	-	1	4	15	27	57	109	156	165
2001.....	622	1	1	1	-	7	12	20	64	107	159	112
2000.....	559	1	-	1	-	8	20	21	61	111	124	98
1999.....	529	-	4	1	-	2	13	28	67	95	129	91

American Indian 1/, both sexes

2004.....	228	-	-	-	2	5	17	16	38	40	37	33
2003.....	196	-	-	-	1	6	9	23	24	33	41	20
2002.....	154	1	-	-	-	7	8	13	18	34	27	26
2001.....	110	-	-	-	-	6	5	8	18	20	17	20
2000.....	98	-	-	-	-	-	5	8	15	15	26	20
1999.....	96	-	-	1	-	3	3	11	14	22	19	12

American Indian 1/, male

2004.....	140	-	-	-	-	3	13	13	19	29	22	19
2003.....	114	-	-	-	-	4	8	14	14	20	20	14
2002.....	95	1	-	-	-	4	8	9	11	18	15	16
2001.....	61	-	-	-	-	1	5	5	8	12	10	11
2000.....	61	-	-	-	-	-	-	8	11	6	16	13
1999.....	60	-	-	-	-	2	3	7	12	13	8	9

American Indian 1/, female

2004.....	88	-	-	-	2	2	4	3	19	11	15	14
2003.....	82	-	-	-	1	2	1	9	10	13	21	6
2002.....	59	-	-	-	-	3	-	4	7	16	12	10
2001.....	49	-	-	-	-	5	-	3	10	8	7	9
2000.....	37	-	-	-	-	-	5	-	4	9	10	7
1999.....	36	-	-	1	-	1	-	4	2	9	11	3

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Accidental poisoning by and exposure to drugs and other biological substances (X40-X44)

Asian or Pacific Islander 2/, both sexes

2004.....	128	-	-	-	-	5	12	10	12	15	19	19
2003.....	113	-	-	-	-	2	9	19	15	14	11	13
2002.....	109	-	1	-	-	6	9	13	12	13	17	12
2001.....	82	-	1	-	1	7	9	10	6	13	11	6
2000.....	69	-	1	-	-	3	5	7	8	9	16	10
1999.....	77	-	-	-	-	2	8	8	8	14	11	5

Asian or Pacific Islander 2/, male

2004.....	98	-	-	-	-	5	8	8	10	12	16	14
2003.....	78	-	-	-	-	1	8	16	13	8	8	10
2002.....	76	-	-	-	-	5	6	11	8	9	14	7
2001.....	63	-	1	-	1	5	8	7	6	11	9	3
2000.....	51	-	1	-	-	3	3	5	6	7	14	8
1999.....	52	-	-	-	-	2	2	6	7	9	10	4

Asian or Pacific Islander 2/, female

2004.....	30	-	-	-	-	-	4	2	2	3	3	5
2003.....	35	-	-	-	-	1	1	3	2	6	3	3
2002.....	33	-	1	-	-	1	3	2	4	4	3	5
2001.....	19	-	-	-	-	2	1	3	-	2	2	3
2000.....	18	-	-	-	-	-	2	2	2	2	2	2
1999.....	25	-	-	-	-	-	6	2	1	5	1	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Accidental poisoning by and exposure to other and unspecified solid or liquids substances (X45-X46,X48-X49)

All races, both sexes

2004.....	550	1	-	-	5	27	31	29	31	71	75	86
2003.....	533	-	3	-	4	31	22	26	30	65	96	94
2002.....	516	3	3	-	2	20	18	29	49	54	86	73
2001.....	461	1	3	1	4	19	22	23	30	74	80	57
2000.....	452	2	7	1	4	23	23	23	32	55	67	65
1999.....	497	1	4	2	10	23	27	25	36	59	85	63

All races, male

2004.....	412	-	-	-	4	20	30	25	25	56	60	65
2003.....	416	-	1	-	2	28	18	21	27	49	77	77
2002.....	401	2	2	-	2	19	16	26	40	43	66	57
2001.....	340	1	2	-	3	17	16	17	20	57	62	43
2000.....	335	2	5	-	3	19	18	18	29	47	53	47
1999.....	366	1	3	1	6	19	22	24	23	48	66	49

All races, female

2004.....	138	1	-	-	1	7	1	4	6	15	15	21
2003.....	117	-	2	-	2	3	4	5	3	16	19	17
2002.....	115	1	1	-	-	1	2	3	9	11	20	16
2001.....	121	-	1	1	1	2	6	6	10	17	18	14
2000.....	117	-	2	1	1	4	5	5	3	8	14	18
1999.....	131	-	1	1	4	4	5	1	13	11	19	14

White, both sexes

2004.....	462	1	-	-	1	20	30	24	28	63	58	75
2003.....	447	-	3	-	2	25	21	24	25	53	82	76
2002.....	402	2	1	-	2	20	16	23	40	40	70	55
2001.....	374	1	1	1	2	17	17	20	24	56	67	50
2000.....	364	2	4	1	4	19	16	15	24	43	58	53
1999.....	389	-	3	1	8	17	19	22	22	49	69	49

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Accidental poisoning by and exposure to other and unspecified solid or liquids substances (X45-X46,X48-X49)

White, male

2004.....	350	-	-	-	1	15	29	21	23	50	50	57
2003.....	341	-	1	-	1	23	17	19	23	40	64	59
2002.....	313	2	1	-	2	19	14	20	32	30	56	42
2001.....	284	1	1	-	2	15	13	14	19	44	53	39
2000.....	270	2	3	-	3	17	15	14	22	37	45	36
1999.....	294	-	2	-	6	15	16	22	13	41	57	40

White, female

2004.....	112	1	-	-	-	5	1	3	5	13	8	18
2003.....	106	-	2	-	1	2	4	5	2	13	18	17
2002.....	89	-	-	-	-	1	2	3	8	10	14	13
2001.....	90	-	-	1	-	2	4	6	5	12	14	11
2000.....	94	-	1	1	1	2	1	1	2	6	13	17
1999.....	95	-	1	1	2	2	3	-	9	8	12	9

Black, both sexes

2004.....	51	-	-	-	2	3	1	-	1	3	8	8
2003.....	53	-	-	-	2	2	-	2	3	4	10	11
2002.....	78	1	2	-	-	-	1	3	4	9	10	13
2001.....	54	-	2	-	-	-	4	1	4	9	8	6
2000.....	57	-	3	-	-	-	1	5	4	5	8	9
1999.....	64	-	1	1	-	3	3	1	7	7	4	9

Black, male

2004.....	38	-	-	-	1	2	1	-	1	3	6	6
2003.....	49	-	-	-	1	2	-	2	3	4	9	11
2002.....	57	-	1	-	-	-	1	3	4	8	6	11
2001.....	34	-	1	-	-	-	2	1	1	7	6	3
2000.....	44	-	2	-	-	-	-	3	3	5	7	8
1999.....	46	-	1	1	-	3	3	1	5	6	2	6

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Accidental poisoning by and exposure to other and unspecified solid or liquids substances (X45-X46,X48-X49)

Black, female

2004.....	13	-	-	-	1	1	-	-	-	-	2	2
2003.....	4	-	-	-	1	-	-	-	-	-	1	-
2002.....	21	1	1	-	-	-	-	-	-	1	4	2
2001.....	20	-	1	-	-	-	2	-	3	2	2	3
2000.....	13	-	1	-	-	-	1	2	1	-	1	1
1999.....	18	-	-	-	-	-	-	-	2	1	2	3

American Indian 1/, both sexes

2004.....	30	-	-	-	2	4	-	3	2	4	8	3
2003.....	25	-	-	-	-	3	1	-	2	7	2	5
2002.....	30	-	-	-	-	-	1	3	4	4	5	4
2001.....	30	-	-	-	2	2	1	2	2	8	4	1
2000.....	27	-	-	-	-	3	4	3	4	6	1	3
1999.....	41	1	-	-	2	3	5	2	7	3	9	5

American Indian 1/, male

2004.....	19	-	-	-	2	3	-	2	1	3	3	2
2003.....	19	-	-	-	-	2	1	-	1	4	2	5
2002.....	25	-	-	-	-	-	1	3	3	4	3	3
2001.....	22	-	-	-	1	2	1	2	-	6	3	1
2000.....	17	-	-	-	-	1	1	1	4	4	1	3
1999.....	23	1	-	-	-	1	3	1	5	1	4	3

American Indian 1/, female

2004.....	11	-	-	-	-	1	-	1	1	1	5	1
2003.....	6	-	-	-	-	1	-	-	1	3	-	-
2002.....	5	-	-	-	-	-	-	-	1	-	2	1
2001.....	8	-	-	-	1	-	-	-	2	2	1	-
2000.....	10	-	-	-	-	2	3	2	-	2	-	-
1999.....	18	-	-	-	2	2	2	1	2	2	5	2

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Accidental poisoning by and exposure to other gases and vapors (X47)

All races, both sexes

2004.....	562	2	7	5	16	25	22	30	40	40	50	62
2003.....	630	4	20	10	20	18	46	38	38	51	56	57
2002.....	640	5	10	13	11	27	41	31	36	55	66	57
2001.....	593	5	10	12	10	35	49	34	32	53	56	53
2000.....	593	3	15	13	10	37	32	37	45	48	52	53
1999.....	534	3	14	6	6	29	36	19	43	52	60	49

All races, male

2004.....	408	2	5	1	8	18	16	27	30	30	41	50
2003.....	463	1	11	3	13	15	36	30	31	38	48	50
2002.....	467	4	5	8	5	17	34	27	29	46	57	46
2001.....	425	3	4	5	4	24	43	25	23	39	44	45
2000.....	435	3	8	10	5	31	28	30	40	41	44	36
1999.....	410	3	6	5	4	21	29	17	35	45	52	39

All races, female

2004.....	154	-	2	4	8	7	6	3	10	10	9	12
2003.....	167	3	9	7	7	3	10	8	7	13	8	7
2002.....	173	1	5	5	6	10	7	4	7	9	9	11
2001.....	168	2	6	7	6	11	6	9	9	14	12	8
2000.....	158	-	7	3	5	6	4	7	5	7	8	17
1999.....	124	-	8	1	2	8	7	2	8	7	8	10

White, both sexes

2004.....	468	1	4	3	14	23	15	25	32	33	46	54
2003.....	519	3	10	6	9	15	41	32	32	42	50	49
2002.....	539	3	8	11	10	22	36	29	29	42	53	45
2001.....	468	2	7	9	8	25	40	26	29	43	44	44
2000.....	495	2	10	7	3	30	28	33	42	41	41	45
1999.....	432	3	8	1	4	25	32	19	34	38	50	38

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Accidental poisoning by and exposure to other gases and vapors (X47)

White, male

2004.....	347	1	2	1	8	18	11	24	24	26	38	44
2003.....	386	1	4	2	7	12	32	25	27	33	44	42
2002.....	394	3	3	7	5	14	31	25	24	36	46	39
2001.....	342	2	3	4	4	19	34	19	20	33	35	38
2000.....	369	2	5	5	2	25	24	27	38	34	37	32
1999.....	338	3	2	1	2	19	26	17	30	33	44	30

White, female

2004.....	121	-	2	2	6	5	4	1	8	7	8	10
2003.....	133	2	6	4	2	3	9	7	5	9	6	7
2002.....	145	-	5	4	5	8	5	4	5	6	7	6
2001.....	126	-	4	5	4	6	6	7	9	10	9	6
2000.....	126	-	5	2	1	5	4	6	4	7	4	13
1999.....	94	-	6	-	2	6	6	2	4	5	6	8

Black, both sexes

2004.....	77	1	3	2	2	2	5	4	6	6	3	7
2003.....	82	-	8	2	6	2	5	5	5	3	6	6
2002.....	83	2	2	2	1	3	2	2	6	12	10	10
2001.....	103	3	2	3	1	8	2	6	2	10	11	8
2000.....	75	1	5	5	2	2	1	4	2	7	10	6
1999.....	78	-	4	4	1	1	4	-	5	10	9	11

Black, male

2004.....	49	1	3	-	-	-	4	2	4	3	3	6
2003.....	61	-	5	-	3	2	4	5	4	2	4	6
2002.....	60	1	2	1	-	2	1	2	4	9	8	6
2001.....	66	1	1	1	-	3	2	4	2	6	9	6
2000.....	52	1	3	5	1	2	1	3	1	7	7	4
1999.....	57	-	3	3	1	-	3	-	3	8	7	9

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Accidental poisoning by and exposure to other gases and vapors (X47)												
Black, female												
2004.....	28	-	-	2	2	2	1	2	2	3	-	1
2003.....	21	-	3	2	3	-	1	-	1	1	2	-
2002.....	23	1	-	1	1	1	1	-	2	3	2	4
2001.....	37	2	1	2	1	5	-	2	-	4	2	2
2000.....	23	-	2	-	1	-	-	1	1	-	3	2
1999.....	21	-	1	1	-	1	1	-	2	2	2	2
American Indian 1/, both sexes												
2004.....	4	-	-	-	-	-	2	-	-	1	-	-
2003.....	10	-	-	-	2	1	-	-	-	2	-	1
2002.....	4	-	-	-	-	-	-	-	-	-	1	1
2001.....	9	-	1	-	1	1	3	1	-	-	-	-
2000.....	12	-	-	1	3	3	3	-	-	-	1	-
1999.....	12	-	2	1	-	1	-	-	3	3	-	-
American Indian 1/, male												
2004.....	3	-	-	-	-	-	1	-	-	1	-	-
2003.....	8	-	-	-	2	1	-	-	-	2	-	1
2002.....	4	-	-	-	-	-	-	-	-	-	1	1
2001.....	7	-	-	-	-	1	3	1	-	-	-	-
2000.....	6	-	-	-	-	2	3	-	-	-	-	-
1999.....	8	-	1	1	-	1	-	-	1	3	-	-
American Indian 1/, female												
2004.....	1	-	-	-	-	-	1	-	-	-	-	-
2003.....	2	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	2	-	1	-	1	-	-	-	-	-	-	-
2000.....	6	-	-	1	3	1	-	-	-	-	1	-
1999.....	4	-	1	-	-	-	-	-	2	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Accidental poisoning by and exposure to other gases and vapors (X47)												
Asian or Pacific Islander 2/, both sexes												
2004.....	13	-	-	-	-	-	-	1	2	-	1	1
2003.....	19	1	2	2	3	-	-	1	1	4	-	1
2002.....	14	-	-	-	-	2	3	-	1	1	2	1
2001.....	13	-	-	-	-	1	4	1	1	-	1	1
2000.....	11	-	-	-	2	2	-	-	1	-	-	2
1999.....	12	-	-	-	1	2	-	-	1	1	1	-
Asian or Pacific Islander 2/, male												
2004.....	9	-	-	-	-	-	-	1	2	-	-	-
2003.....	8	-	2	1	1	-	-	-	-	1	-	1
2002.....	9	-	-	-	-	1	2	-	1	1	2	-
2001.....	10	-	-	-	-	1	4	1	1	-	-	1
2000.....	8	-	-	-	2	2	-	-	1	-	-	-
1999.....	7	-	-	-	1	1	-	-	1	1	1	-
Asian or Pacific Islander 2/, female												
2004.....	4	-	-	-	-	-	-	-	-	-	1	1
2003.....	11	1	-	1	2	-	-	1	1	3	-	-
2002.....	5	-	-	-	-	1	1	-	-	-	-	1
2001.....	3	-	-	-	-	-	-	-	-	-	1	-
2000.....	3	-	-	-	-	-	-	-	-	-	-	2
1999.....	5	-	-	-	-	1	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

All other and unspecified accidents and their sequelae (W20-W31,W35-W38,W41-W64,W92-W99,X20-X32,X39,X50-X59,Y86)

All races, both sexes

2004.....	10,339	53	110	61	54	153	229	233	277	348	466	541
2003.....	10,734	45	106	41	64	142	261	213	257	316	512	540
2002.....	10,840	35	94	64	74	156	207	216	268	347	495	475
2001.....	11,377	54	122	60	55	156	237	217	278	403	450	523
2000.....	10,973	41	101	61	58	164	233	239	286	431	512	516
1999.....	11,898	44	116	77	75	178	219	258	289	392	531	486

All races, male

2004.....	6,007	34	76	42	41	124	204	201	241	294	370	452
2003.....	6,122	23	65	29	49	121	216	180	220	253	432	441
2002.....	6,088	25	59	45	46	123	187	178	238	285	397	402
2001.....	6,323	32	70	38	42	120	208	189	240	333	372	427
2000.....	6,206	24	61	41	42	133	210	216	245	372	434	438
1999.....	6,622	28	80	48	51	140	193	224	243	326	431	394

All races, female

2004.....	4,332	19	34	19	13	29	25	32	36	54	96	89
2003.....	4,612	22	41	12	15	21	45	33	37	63	80	99
2002.....	4,752	10	35	19	28	33	20	38	30	62	98	73
2001.....	5,054	22	52	22	13	36	29	28	38	70	78	96
2000.....	4,767	17	40	20	16	31	23	23	41	59	78	78
1999.....	5,276	16	36	29	24	38	26	34	46	66	100	92

White, both sexes

2004.....	9,123	39	88	48	39	130	184	197	222	273	396	455
2003.....	9,416	30	78	36	49	115	206	180	208	249	417	419
2002.....	9,537	24	65	51	53	125	181	186	218	269	406	385
2001.....	10,038	40	92	47	43	134	203	173	223	330	350	419
2000.....	9,620	30	73	45	47	134	190	198	237	343	402	393
1999.....	10,393	30	91	63	62	146	181	209	232	331	425	364

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

All other and unspecified accidents and their sequelae (W20-W31,W35-W38,W41-W64,W92-W99,X20-X32,X39,X50-X59,Y86)

White, male

2004.....	5,211	25	64	31	28	107	165	170	195	234	316	382
2003.....	5,262	16	48	26	39	98	180	153	179	200	360	346
2002.....	5,256	18	40	36	29	102	163	155	195	221	339	335
2001.....	5,488	24	55	28	32	104	180	154	190	281	292	351
2000.....	5,271	19	43	33	32	109	173	177	208	299	343	331
1999.....	5,632	19	60	42	42	114	160	184	196	277	343	293

White, female

2004.....	3,912	14	24	17	11	23	19	27	27	39	80	73
2003.....	4,154	14	30	10	10	17	26	27	29	49	57	73
2002.....	4,281	6	25	15	24	23	18	31	23	48	67	50
2001.....	4,550	16	37	19	11	30	23	19	33	49	58	68
2000.....	4,349	11	30	12	15	25	17	21	29	44	59	62
1999.....	4,761	11	31	21	20	32	21	25	36	54	82	71

Black, both sexes

2004.....	976	13	15	10	12	14	31	33	42	61	55	73
2003.....	1,059	10	23	4	7	17	46	26	33	54	74	99
2002.....	1,053	11	26	11	19	24	18	25	40	68	71	76
2001.....	1,076	11	23	9	12	16	24	34	35	52	79	85
2000.....	1,111	10	22	14	9	22	34	30	42	72	86	99
1999.....	1,299	13	21	10	11	25	30	40	45	49	93	107

Black, male

2004.....	636	8	7	9	10	11	25	29	34	49	42	58
2003.....	677	5	16	2	5	14	30	21	27	42	54	77
2002.....	663	7	19	8	16	17	16	19	34	56	43	54
2001.....	655	7	13	8	10	12	19	25	32	37	64	62
2000.....	766	5	14	7	8	17	30	28	33	61	71	86
1999.....	847	8	17	4	8	22	28	31	37	38	76	89

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

All other and unspecified accidents and their sequelae (W20-W31,W35-W38,W41-W64,W92-W99,X20-X32,X39,X50-X59,Y86)

Black, female

2004.....	340	5	8	1	2	3	6	4	8	12	13	15
2003.....	382	5	7	2	2	3	16	5	6	12	20	22
2002.....	390	4	7	3	3	7	2	6	6	12	28	22
2001.....	421	4	10	1	2	4	5	9	3	15	15	23
2000.....	345	5	8	7	1	5	4	2	9	11	15	13
1999.....	452	5	4	6	3	3	2	9	8	11	17	18

American Indian 1/, both sexes

2004.....	120	-	4	1	-	5	8	2	6	11	9	9
2003.....	139	2	2	-	3	7	3	5	11	8	15	13
2002.....	122	-	2	2	2	4	7	3	6	8	16	9
2001.....	117	2	1	2	-	1	6	2	11	15	13	10
2000.....	124	-	3	1	1	4	7	2	4	10	17	15
1999.....	117	-	4	3	1	5	4	5	10	6	11	12

American Indian 1/, male

2004.....	85	-	3	-	-	4	8	1	5	9	6	9
2003.....	108	1	1	-	1	6	2	4	10	7	14	11
2002.....	94	-	-	1	1	3	7	3	5	7	13	8
2001.....	84	-	-	1	-	1	6	2	11	12	9	7
2000.....	90	-	2	1	1	3	6	2	2	9	14	14
1999.....	89	-	3	1	1	3	2	5	9	5	10	11

American Indian 1/, female

2004.....	35	-	1	1	-	1	-	1	1	2	3	-
2003.....	31	1	1	-	2	1	1	1	1	1	1	2
2002.....	28	-	2	1	1	1	-	-	1	1	3	1
2001.....	33	2	1	1	-	-	-	-	-	3	4	3
2000.....	34	-	1	-	-	1	1	-	2	1	3	1
1999.....	28	-	1	2	-	2	2	-	1	1	1	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

All other and unspecified accidents and their sequelae (W20-W31,W35-W38,W41-W64,W92-W99,X20-X32,X39,X50-X59,Y86)

Asian or Pacific Islander 2/, both sexes

2004.....	120	1	3	2	3	4	6	1	7	3	6	4
2003.....	120	3	3	1	5	3	6	2	5	5	6	9
2002.....	128	-	1	-	-	3	1	2	4	2	2	5
2001.....	146	1	6	2	-	5	4	8	9	6	8	9
2000.....	118	1	3	1	1	4	2	9	3	6	7	9
1999.....	89	1	-	1	1	2	4	4	2	6	2	3

Asian or Pacific Islander 2/, male

2004.....	75	1	2	2	3	2	6	1	7	2	6	3
2003.....	75	1	-	1	4	3	4	2	4	4	4	7
2002.....	75	-	-	-	-	1	1	1	4	1	2	5
2001.....	96	1	2	1	-	3	3	8	7	3	7	7
2000.....	79	-	2	-	1	4	1	9	2	3	6	7
1999.....	54	1	-	1	-	1	3	4	1	6	2	1

Asian or Pacific Islander 2/, female

2004.....	45	-	1	-	-	2	-	-	-	1	-	1
2003.....	45	2	3	-	1	-	2	-	1	1	2	2
2002.....	53	-	1	-	-	2	-	1	-	1	-	-
2001.....	50	-	4	1	-	2	1	-	2	3	1	2
2000.....	39	1	1	1	-	-	1	-	1	3	1	2
1999.....	35	-	-	-	1	1	1	-	1	-	-	2

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)												
All races, both sexes												
2004.....	32,439	2	283	1,700	2,616	2,412	2,662	2,948	3,690	3,743
2003.....	31,484	6	244	1,487	2,501	2,276	2,789	2,958	3,644	3,485
2002.....	31,655	4	260	1,513	2,497	2,423	2,623	3,141	3,710	3,473
2001 3/.....	30,622	7	272	1,611	2,360	2,389	2,681	3,176	3,459	3,260
2000.....	29,350	7	300	1,621	2,373	2,341	2,451	3,112	3,450	2,987
1999.....	29,199	2	242	1,615	2,286	2,398	2,708	3,286	3,180	2,817
All races, male												
2004.....	25,566	2	185	1,345	2,251	2,015	2,127	2,292	2,775	2,749
2003.....	25,203	6	188	1,222	2,159	1,901	2,255	2,347	2,791	2,616
2002.....	25,409	3	196	1,280	2,152	2,003	2,132	2,484	2,821	2,608
2001 3/.....	24,672	7	207	1,345	2,064	2,015	2,184	2,518	2,662	2,484
2000.....	23,618	6	238	1,351	2,073	1,956	1,982	2,457	2,657	2,307
1999.....	23,458	2	192	1,347	1,979	2,005	2,189	2,590	2,429	2,152
All races, female												
2004.....	6,873	-	98	355	365	397	535	656	915	994
2003.....	6,281	-	56	265	342	375	534	611	853	869
2002.....	6,246	1	64	233	345	420	491	657	889	865
2001 3/.....	5,950	-	65	266	296	374	497	658	797	776
2000.....	5,732	1	62	270	300	385	469	655	793	680
1999.....	5,741	-	50	268	307	393	519	696	751	665
White, both sexes												
2004.....	29,251	1	217	1,425	2,185	2,001	2,287	2,636	3,395	3,453
2003.....	28,485	5	188	1,274	2,044	1,895	2,410	2,629	3,354	3,254
2002.....	28,731	3	211	1,308	2,098	2,021	2,277	2,799	3,420	3,245
2001 3/.....	27,710	5	213	1,379	1,955	2,012	2,344	2,849	3,168	3,030
2000.....	26,475	6	248	1,354	1,961	1,941	2,143	2,796	3,146	2,762
1999.....	26,300	2	204	1,344	1,875	1,996	2,350	2,933	2,927	2,599

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)												
White, male												
2004.....	23,081	1	142	1,132	1,884	1,684	1,842	2,057	2,549	2,525
2003.....	22,830	5	147	1,047	1,781	1,580	1,950	2,080	2,571	2,447
2002.....	23,049	2	157	1,109	1,810	1,651	1,844	2,205	2,608	2,423
2001 3/.....	22,328	5	163	1,156	1,704	1,699	1,905	2,248	2,443	2,325
2000.....	21,293	5	195	1,130	1,723	1,631	1,732	2,197	2,423	2,128
1999.....	21,107	2	162	1,122	1,629	1,667	1,900	2,291	2,244	1,987
White, female												
2004.....	6,170	-	75	293	301	317	445	579	846	928
2003.....	5,655	-	41	227	263	315	460	549	783	807
2002.....	5,682	1	54	199	288	370	433	594	812	822
2001 3/.....	5,382	-	50	223	251	313	439	601	725	705
2000.....	5,182	1	53	224	238	310	411	599	723	634
1999.....	5,193	-	42	222	246	329	450	642	683	612
Black, both sexes												
2004.....	2,019	1	48	155	310	280	241	194	187	174
2003.....	1,955	1	43	121	326	255	254	212	204	141
2002.....	1,939	1	34	128	275	273	231	235	207	150
2001 3/.....	1,957	2	39	136	297	254	238	230	195	159
2000.....	1,962	-	40	172	308	266	210	214	224	156
1999.....	1,950	-	29	179	291	261	242	248	187	146
Black, male												
2004.....	1,655	1	29	124	272	240	189	154	150	133
2003.....	1,597	1	34	107	278	224	208	173	156	109
2002.....	1,633	1	28	111	240	246	201	197	157	130
2001 3/.....	1,627	2	30	116	279	223	202	191	145	115
2000.....	1,636	-	35	150	265	224	178	180	176	129
1999.....	1,651	-	22	155	259	231	205	217	139	116

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)												
Black, female												
2004.....	364	-	19	31	38	40	52	40	37	41
2003.....	358	-	9	14	48	31	46	39	48	32
2002.....	306	-	6	17	35	27	30	38	50	20
2001 3/.....	330	-	9	20	18	31	36	39	50	44
2000.....	326	-	5	22	43	42	32	34	48	27
1999.....	299	-	7	24	32	30	37	31	48	30
American Indian 1/, both sexes												
2004.....	404	-	10	69	53	53	49	55	35	36
2003.....	322	-	7	50	53	44	42	42	35	15
2002.....	324	-	6	42	58	50	34	44	24	25
2001 3/.....	321	-	12	52	34	40	36	44	36	19
2000.....	297	1	6	44	45	43	35	37	30	16
1999.....	291	-	2	52	43	53	38	35	14	20
American Indian 1/, male												
2004.....	306	-	6	49	43	39	37	42	29	29
2003.....	265	-	6	37	43	38	38	36	30	11
2002.....	258	-	4	34	46	45	29	35	17	15
2001 3/.....	259	-	7	41	28	35	28	37	28	14
2000.....	237	1	5	34	37	33	26	31	22	11
1999.....	233	-	2	43	36	42	30	29	9	17
American Indian 1/, female												
2004.....	98	-	4	20	10	14	12	13	6	7
2003.....	57	-	1	13	10	6	4	6	5	4
2002.....	66	-	2	8	12	5	5	9	7	10
2001 3/.....	62	-	5	11	6	5	8	7	8	5
2000.....	60	-	1	10	8	10	9	6	8	5
1999.....	58	-	-	9	7	11	8	6	5	3

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)

Asian or Pacific Islander 2/, both sexes

2004.....	765	-	8	51	68	78	85	63	73	80
2003.....	722	-	6	42	78	82	83	75	51	75
2002.....	661	-	9	35	66	79	81	63	59	53
2001 3/.....	634	-	8	44	74	83	63	53	60	52
2000.....	616	-	6	51	59	91	63	65	50	53
1999.....	658	-	7	40	77	88	78	70	52	52

Asian or Pacific Islander 2/, male

2004.....	524	-	8	40	52	52	59	39	47	62
2003.....	511	-	1	31	57	59	59	58	34	49
2002.....	469	-	7	26	56	61	58	47	39	40
2001 3/.....	458	-	7	32	53	58	49	42	46	30
2000.....	452	-	3	37	48	68	46	49	36	39
1999.....	467	-	6	27	55	65	54	53	37	32

Asian or Pacific Islander 2/, female

2004.....	241	-	-	11	16	26	26	24	26	18
2003.....	211	-	5	11	21	23	24	17	17	26
2002.....	192	-	2	9	10	18	23	16	20	13
2001 3/.....	176	-	1	12	21	25	14	11	14	22
2000.....	164	-	3	14	11	23	17	16	14	14
1999.....	191	-	1	13	22	23	24	17	15	20

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Intentional self-poisoning (suicide) by and exposure to drugs and other biological substances (X60-X64)

All races, both sexes

2004.....	4,208	-	15	95	177	243	333	482	653	686
2003.....	3,890	-	9	79	146	207	339	458	637	652
2002.....	3,884	-	9	81	151	217	325	473	673	660
2001.....	3,559	-	10	84	145	188	309	473	625	610
2000.....	3,243	-	14	74	124	198	272	420	590	507
1999.....	3,181	-	6	69	121	205	265	437	560	478

All races, male

2004.....	1,966	-	-	43	106	133	173	253	284	305
2003.....	1,837	-	3	43	75	110	164	235	327	293
2002.....	1,851	-	-	46	70	122	184	245	316	286
2001.....	1,713	-	2	39	72	102	163	267	321	286
2000.....	1,533	-	8	38	66	92	151	215	276	247
1999.....	1,511	-	3	27	63	103	126	226	279	239

All races, female

2004.....	2,242	-	15	52	71	110	160	229	369	381
2003.....	2,053	-	6	36	71	97	175	223	310	359
2002.....	2,033	-	9	35	81	95	141	228	357	374
2001.....	1,846	-	8	45	73	86	146	206	304	324
2000.....	1,710	-	6	36	58	106	121	205	314	260
1999.....	1,670	-	3	42	58	102	139	211	281	239

White, both sexes

2004.....	3,928	-	11	84	159	214	300	437	621	652
2003.....	3,618	-	9	71	126	182	307	421	588	602
2002.....	3,618	-	8	71	138	190	283	433	632	632
2001.....	3,319	-	9	75	128	168	282	436	582	566
2000.....	3,017	-	12	66	107	170	255	383	548	472
1999.....	2,969	-	6	62	107	178	233	408	521	454

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Intentional self-poisoning (suicide) by and exposure to drugs and other biological substances (X60-X64)

White, male

2004.....	1,846	-	-	38	97	121	160	233	274	291
2003.....	1,732	-	3	40	71	101	150	220	302	276
2002.....	1,722	-	-	39	66	105	160	223	300	274
2001.....	1,606	-	2	38	62	93	152	250	299	267
2000.....	1,418	-	7	36	54	79	141	197	258	226
1999.....	1,410	-	3	24	59	87	111	211	265	227

White, female

2004.....	2,082	-	11	46	62	93	140	204	347	361
2003.....	1,886	-	6	31	55	81	157	201	286	326
2002.....	1,896	-	8	32	72	85	123	210	332	358
2001.....	1,713	-	7	37	66	75	130	186	283	299
2000.....	1,599	-	5	30	53	91	114	186	290	246
1999.....	1,559	-	3	38	48	91	122	197	256	227

Black, both sexes

2004.....	164	-	3	3	12	18	22	25	24	21
2003.....	167	-	-	3	8	12	19	27	36	35
2002.....	164	-	1	6	7	15	27	22	32	17
2001.....	157	-	1	5	10	9	17	28	29	31
2000.....	155	-	1	4	8	17	15	23	32	24
1999.....	141	-	-	3	7	16	22	21	32	20

Black, male

2004.....	67	-	-	1	7	9	7	12	6	8
2003.....	60	-	-	-	2	4	7	12	17	10
2002.....	80	-	-	5	2	9	17	12	14	8
2001.....	71	-	-	-	8	4	9	12	15	13
2000.....	80	-	-	-	6	9	9	12	13	15
1999.....	74	-	-	2	2	11	12	12	13	10

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Intentional self-poisoning (suicide) by and exposure to drugs and other biological substances (X60-X64)

Black, female

2004.....	97	-	3	2	5	9	15	13	18	13
2003.....	107	-	-	3	6	8	12	15	19	25
2002.....	84	-	1	1	5	6	10	10	18	9
2001.....	86	-	1	5	2	5	8	16	14	18
2000.....	75	-	1	4	2	8	6	11	19	9
1999.....	67	-	-	1	5	5	10	9	19	10

American Indian 1/, both sexes

2004.....	51	-	1	6	2	2	6	11	4	6
2003.....	39	-	-	2	4	3	5	7	8	4
2002.....	40	-	-	2	1	5	4	8	6	6
2001.....	34	-	-	3	2	5	5	3	6	5
2000.....	23	-	1	1	4	5	1	2	4	2
1999.....	25	-	-	1	2	6	3	4	3	2

American Indian 1/, male

2004.....	26	-	-	4	1	1	3	4	2	2
2003.....	18	-	-	1	1	2	4	2	6	1
2002.....	22	-	-	2	-	4	3	4	1	2
2001.....	12	-	-	-	2	2	-	2	1	2
2000.....	10	-	1	-	3	2	-	1	1	1
1999.....	7	-	-	-	1	3	-	1	-	1

American Indian 1/, female

2004.....	25	-	1	2	1	1	3	7	2	4
2003.....	21	-	-	1	3	1	1	5	2	3
2002.....	18	-	-	-	1	1	1	4	5	4
2001.....	22	-	-	3	-	3	5	1	5	3
2000.....	13	-	-	1	1	3	1	1	3	1
1999.....	18	-	-	1	1	3	3	3	3	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Intentional self-poisoning (suicide) by and exposure to drugs and other biological substances (X60-X64)

Asian or Pacific Islander 2/, both sexes

2004.....	65	-	-	2	4	9	5	9	4	7
2003.....	66	-	-	3	8	10	8	3	5	11
2002.....	62	-	-	2	5	7	11	10	3	5
2001.....	49	-	-	1	5	6	5	6	8	8
2000.....	48	-	-	3	5	6	1	12	6	9
1999.....	46	-	-	3	5	5	7	4	4	2

Asian or Pacific Islander 2/, male

2004.....	27	-	-	-	1	2	3	4	2	4
2003.....	27	-	-	2	1	3	3	1	2	6
2002.....	27	-	-	-	2	4	4	6	1	2
2001.....	24	-	-	1	-	3	2	3	6	4
2000.....	25	-	-	2	3	2	1	5	4	5
1999.....	20	-	-	1	1	2	3	2	1	1

Asian or Pacific Islander 2/, female

2004.....	38	-	-	2	3	7	2	5	2	3
2003.....	39	-	-	1	7	7	5	2	3	5
2002.....	35	-	-	2	3	3	7	4	2	3
2001.....	25	-	-	-	5	3	3	3	2	4
2000.....	23	-	-	1	2	4	-	7	2	4
1999.....	26	-	-	2	4	3	4	2	3	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-poisoning (suicide) by and exposure to other and unspecified solid or liquid substances and their vapors (X65-X66,X68-X69)

All races, both sexes

2004.....	264	-	-	4	15	21	28	20	40	31
2003.....	216	-	-	6	13	17	22	29	22	25
2002.....	183	-	1	5	8	17	16	15	27	30
2001.....	190	-	-	8	8	23	13	22	37	19
2000.....	178	-	-	3	9	14	15	27	27	23
1999.....	177	-	-	1	5	19	16	21	27	22

All races, male

2004.....	187	-	-	3	13	18	20	17	30	19
2003.....	145	-	-	3	9	13	16	25	18	17
2002.....	132	-	-	2	8	14	10	11	23	24
2001.....	144	-	-	6	7	14	10	18	31	14
2000.....	136	-	-	3	9	11	13	22	21	17
1999.....	125	-	-	1	4	16	11	16	20	14

All races, female

2004.....	77	-	-	1	2	3	8	3	10	12
2003.....	71	-	-	3	4	4	6	4	4	8
2002.....	51	-	1	3	-	3	6	4	4	6
2001.....	46	-	-	2	1	9	3	4	6	5
2000.....	42	-	-	-	-	3	2	5	6	6
1999.....	52	-	-	-	1	3	5	5	7	8

White, both sexes

2004.....	234	-	-	4	15	16	23	17	35	29
2003.....	188	-	-	6	9	15	18	24	19	25
2002.....	162	-	-	4	6	12	13	12	26	28
2001.....	151	-	-	7	5	19	10	17	32	18
2000.....	152	-	-	3	7	12	14	25	25	18
1999.....	155	-	-	1	3	15	13	18	24	19

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Intentional self-poisoning (suicide) by and exposure to other and unspecified solid or liquid substances and their vapors (X65-X66,X68-X69)

White, male

2004.....	164	-	-	3	13	13	17	15	26	17
2003.....	126	-	-	3	6	13	14	20	15	17
2002.....	117	-	-	2	6	10	8	8	22	22
2001.....	112	-	-	5	4	12	7	14	26	13
2000.....	119	-	-	3	7	10	12	20	21	13
1999.....	111	-	-	1	3	13	11	14	18	12

White, female

2004.....	70	-	-	1	2	3	6	2	9	12
2003.....	62	-	-	3	3	2	4	4	4	8
2002.....	45	-	-	2	-	2	5	4	4	6
2001.....	39	-	-	2	1	7	3	3	6	5
2000.....	33	-	-	-	-	2	2	5	4	5
1999.....	44	-	-	-	-	2	2	4	6	7

Black, both sexes

2004.....	19	-	-	-	-	2	5	3	2	2
2003.....	10	-	-	-	-	1	2	2	1	-
2002.....	12	-	-	-	2	1	2	2	1	1
2001.....	19	-	-	-	2	2	2	2	4	1
2000.....	18	-	-	-	1	-	1	2	2	3
1999.....	10	-	-	-	-	1	2	3	2	-

Black, male

2004.....	15	-	-	-	-	2	3	2	2	2
2003.....	6	-	-	-	-	-	1	2	1	-
2002.....	9	-	-	-	2	-	2	2	1	1
2001.....	16	-	-	-	2	1	2	2	4	1
2000.....	12	-	-	-	1	-	1	2	-	3
1999.....	6	-	-	-	-	1	-	2	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Intentional self-poisoning (suicide) by and exposure to other and unspecified solid or liquid substances and their vapors (X65-X66,X68-X69)

Asian or Pacific Islander 2/, both sexes

2004.....	7	-	-	-	-	1	-	-	2	-
2003.....	16	-	-	-	3	1	2	3	2	-
2002.....	7	-	1	1	-	3	1	-	-	1
2001.....	18	-	-	1	1	2	1	3	1	-
2000.....	8	-	-	-	1	2	-	-	-	2
1999.....	11	-	-	-	2	3	1	-	1	2

Asian or Pacific Islander 2/, male

2004.....	4	-	-	-	-	1	-	-	1	-
2003.....	11	-	-	-	2	-	1	3	2	-
2002.....	4	-	-	-	-	3	-	-	-	1
2001.....	14	-	-	1	1	1	1	2	1	-
2000.....	5	-	-	-	1	1	-	-	-	1
1999.....	7	-	-	-	1	2	-	-	1	1

Asian or Pacific Islander 2/, female

2004.....	3	-	-	-	-	-	-	-	1	-
2003.....	5	-	-	-	1	1	1	-	-	-
2002.....	3	-	1	1	-	-	1	-	-	-
2001.....	4	-	-	-	-	1	-	1	-	-
2000.....	3	-	-	-	-	1	-	-	-	1
1999.....	4	-	-	-	1	1	1	-	-	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-poisoning (suicide) by and exposure to other gases and vapors (X67)

All races, both sexes

2004.....	1,328	-	-	26	46	84	108	151	200	202
2003.....	1,356	-	-	18	48	65	147	183	203	182
2002.....	1,419	-	1	32	48	80	139	167	214	221
2001.....	1,442	-	1	25	67	81	139	179	205	210
2000.....	1,438	-	1	34	60	107	140	197	204	211
1999.....	1,535	-	-	33	99	101	179	199	212	198

All races, male

2004.....	1,047	-	-	24	42	68	93	116	161	158
2003.....	1,066	-	-	14	40	56	124	147	161	144
2002.....	1,114	-	-	27	38	66	108	134	171	174
2001.....	1,115	-	-	21	61	63	118	143	156	165
2000.....	1,123	-	-	29	53	86	112	157	160	168
1999.....	1,191	-	-	28	86	85	139	159	170	149

All races, female

2004.....	281	-	-	2	4	16	15	35	39	44
2003.....	290	-	-	4	8	9	23	36	42	38
2002.....	305	-	1	5	10	14	31	33	43	47
2001.....	327	-	-	4	6	18	21	36	49	45
2000.....	315	-	1	5	7	21	28	40	44	43
1999.....	344	-	-	5	13	16	40	40	42	49

White, both sexes

2004.....	1,263	-	-	25	42	76	96	144	189	195
2003.....	1,301	-	-	18	43	60	138	175	195	178
2002.....	1,369	-	1	31	44	76	133	162	204	214
2001.....	1,381	-	-	24	65	76	129	171	197	198
2000.....	1,384	-	1	33	55	96	133	191	196	204
1999.....	1,470	-	-	32	94	86	172	192	199	187

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-poisoning (suicide) by and exposure to other gases and vapors (X67)

White, male

2004.....	999	-	-	23	38	63	85	111	154	152
2003.....	1,026	-	-	14	36	52	119	142	154	142
2002.....	1,072	-	-	27	36	63	103	129	162	167
2001.....	1,066	-	-	20	60	59	112	136	149	156
2000.....	1,080	-	-	28	51	77	105	153	153	162
1999.....	1,143	-	-	27	82	74	134	154	161	140

White, female

2004.....	264	-	-	2	4	13	11	33	35	43
2003.....	275	-	-	4	7	8	19	33	41	36
2002.....	297	-	1	4	8	13	30	33	42	47
2001.....	315	-	-	4	5	17	17	35	48	42
2000.....	304	-	1	5	4	19	28	38	43	42
1999.....	327	-	-	5	12	12	38	38	38	47

Black, both sexes

2004.....	41	-	-	1	3	3	6	5	8	6
2003.....	38	-	-	-	4	4	6	7	4	2
2002.....	34	-	-	1	1	4	5	4	6	7
2001.....	38	-	-	-	1	3	8	5	6	8
2000.....	36	-	-	-	4	6	4	4	4	6
1999.....	41	-	-	1	1	9	4	4	12	7

Black, male

2004.....	36	-	-	1	3	3	5	5	6	5
2003.....	30	-	-	-	3	4	4	5	4	1
2002.....	30	-	-	-	1	3	4	4	6	7
2001.....	32	-	-	-	-	3	5	5	5	7
2000.....	29	-	-	-	2	6	4	2	3	5
1999.....	33	-	-	1	1	7	3	4	8	7

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Intentional self-poisoning (suicide) by and exposure to other gases and vapors (X67)

Black, female

2004.....	5	-	-	-	-	-	1	-	2	1
2003.....	8	-	-	-	1	-	2	2	-	1
2002.....	4	-	-	1	-	1	1	-	-	-
2001.....	6	-	-	-	1	-	3	-	1	1
2000.....	7	-	-	-	2	-	-	2	1	1
1999.....	8	-	-	-	-	2	1	-	4	-

American Indian 1/, both sexes

2004.....	6	-	-	-	-	1	1	1	2	1
2003.....	1	-	-	-	-	-	-	-	-	-
2002.....	4	-	-	-	1	-	-	-	1	-
2001.....	6	-	-	-	-	1	-	2	-	1
2000.....	8	-	-	-	-	4	-	-	3	1
1999.....	9	-	-	-	1	3	2	1	-	1

American Indian 1/, male

2004.....	3	-	-	-	-	-	1	-	1	1
2003.....	1	-	-	-	-	-	-	-	-	-
2002.....	3	-	-	-	-	-	-	-	1	-
2001.....	4	-	-	-	-	1	-	2	-	-
2000.....	6	-	-	-	-	2	-	-	3	1
1999.....	7	-	-	-	1	2	2	1	-	1

American Indian 1/, female

2004.....	3	-	-	-	-	1	-	1	1	-
2003.....	-	-	-	-	-	-	-	-	-	-
2002.....	1	-	-	-	1	-	-	-	-	-
2001.....	2	-	-	-	-	-	-	-	-	1
2000.....	2	-	-	-	-	2	-	-	-	-
1999.....	2	-	-	-	-	1	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-poisoning (suicide) by and exposure to other gases and vapors (X67)

Asian or Pacific Islander 2/, both sexes

2004.....	18	-	-	-	1	4	5	1	1	-
2003.....	16	-	-	-	1	1	3	1	4	2
2002.....	12	-	-	-	2	-	1	1	3	-
2001.....	17	-	-	1	1	-	2	1	2	3
2000.....	10	-	-	1	1	1	3	2	1	-
1999.....	15	-	-	-	3	3	1	2	1	3

Asian or Pacific Islander 2/, male

2004.....	9	-	-	-	1	2	2	-	-	-
2003.....	9	-	-	-	1	-	1	-	3	1
2002.....	9	-	-	-	1	-	1	1	2	-
2001.....	13	-	-	1	1	-	1	-	2	2
2000.....	8	-	-	1	-	1	3	2	1	-
1999.....	8	-	-	-	2	2	-	-	1	1

Asian or Pacific Islander 2/, female

2004.....	9	-	-	-	-	2	3	1	1	-
2003.....	7	-	-	-	-	1	2	1	1	1
2002.....	3	-	-	-	1	-	-	-	1	-
2001.....	4	-	-	-	-	1	1	1	-	1
2000.....	2	-	-	-	1	-	-	-	-	-
1999.....	7	-	-	-	1	1	1	2	-	2

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intentional self-harm (suicide) by hanging, strangulation and suffocation (X70)												
All races, both sexes												
2004.....	7,336	2	204	675	841	761	831	812	855	734
2003.....	6,635	4	152	568	780	688	791	742	789	626
2002.....	6,462	4	154	550	731	688	723	754	771	632
2001.....	6,198	6	163	551	684	656	717	810	724	564
2000.....	5,688	7	168	514	620	637	615	683	698	507
1999.....	5,427	2	125	434	532	615	708	738	614	418
All races, male												
2004.....	5,980	2	134	501	716	656	705	691	731	590
2003.....	5,496	4	120	444	669	582	673	640	661	522
2002.....	5,355	3	120	453	637	579	638	641	639	522
2001.....	5,210	6	130	453	600	572	616	688	617	484
2000.....	4,733	6	135	414	548	552	533	591	600	427
1999.....	4,490	2	103	347	456	526	613	644	520	349
All races, female												
2004.....	1,356	-	70	174	125	105	126	121	124	144
2003.....	1,139	-	32	124	111	106	118	102	128	104
2002.....	1,107	1	34	97	94	109	85	113	132	110
2001.....	988	-	33	98	84	84	101	122	107	80
2000.....	955	1	33	100	72	85	82	92	98	80
1999.....	937	-	22	87	76	89	95	94	94	69
White, both sexes												
2004.....	6,314	1	153	542	708	631	702	724	774	651
2003.....	5,707	4	107	481	644	576	688	633	703	564
2002.....	5,642	3	123	484	631	574	610	655	689	566
2001.....	5,387	4	125	470	592	544	600	715	653	519
2000.....	4,885	6	126	433	514	527	518	594	612	454
1999.....	4,672	2	99	351	442	510	599	641	557	375

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intentional self-harm (suicide) by hanging, strangulation and suffocation (X70)												
White, male												
2004.....	5,180	1	100	404	603	555	603	618	661	518
2003.....	4,749	4	87	378	560	488	586	549	588	471
2002.....	4,696	2	93	402	555	487	535	557	572	463
2001.....	4,551	4	98	392	525	479	515	604	558	450
2000.....	4,087	5	98	351	460	469	452	515	528	383
1999.....	3,886	2	83	284	384	441	523	559	472	311
White, female												
2004.....	1,134	-	53	138	105	76	99	106	113	133
2003.....	958	-	20	103	84	88	102	84	115	93
2002.....	946	1	30	82	76	87	75	98	117	103
2001.....	836	-	27	78	67	65	85	111	95	69
2000.....	798	1	28	82	54	58	66	79	84	71
1999.....	786	-	16	67	58	69	76	82	85	64
Black, both sexes												
2004.....	535	1	36	71	81	81	68	42	47	39
2003.....	489	-	34	42	84	63	57	57	51	34
2002.....	427	1	23	31	60	62	64	54	54	30
2001.....	414	2	22	35	52	66	72	52	38	26
2000.....	431	-	34	41	64	63	52	49	57	28
1999.....	377	-	22	39	48	55	62	57	35	20
Black, male												
2004.....	445	1	22	56	69	67	58	37	44	32
2003.....	415	-	28	36	65	54	48	49	47	32
2002.....	368	1	20	26	47	51	57	49	48	29
2001.....	361	2	19	30	45	59	63	46	30	24
2000.....	371	-	30	33	56	50	48	42	53	25
1999.....	334	-	16	30	44	49	55	55	31	20

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Intentional self-harm (suicide) by hanging, strangulation and suffocation (X70)

Black, female

2004.....	90	-	14	15	12	14	10	5	3	7
2003.....	74	-	6	6	19	9	9	8	4	2
2002.....	59	-	3	5	13	11	7	5	6	1
2001.....	53	-	3	5	7	7	9	6	8	2
2000.....	60	-	4	8	8	13	4	7	4	3
1999.....	43	-	6	9	4	6	7	2	4	-

American Indian 1/, both sexes

2004.....	165	-	8	39	27	20	25	20	13	10
2003.....	128	-	6	24	25	15	20	19	13	1
2002.....	108	-	5	23	16	19	15	16	4	6
2001.....	131	-	10	26	12	21	13	21	11	2
2000.....	114	1	4	20	18	16	18	21	7	3
1999.....	106	-	1	26	13	18	20	13	4	5

American Indian 1/, male

2004.....	124	-	5	23	22	13	18	18	12	10
2003.....	109	-	5	17	20	15	18	18	11	1
2002.....	89	-	4	16	13	16	15	12	4	6
2001.....	114	-	7	20	9	20	12	19	10	2
2000.....	98	1	4	15	14	14	15	20	6	3
1999.....	87	-	1	21	10	14	17	11	3	5

American Indian 1/, female

2004.....	41	-	3	16	5	7	7	2	1	-
2003.....	19	-	1	7	5	-	2	1	2	-
2002.....	19	-	1	7	3	3	-	4	-	-
2001.....	17	-	3	6	3	1	1	2	1	-
2000.....	16	-	-	5	4	2	3	1	1	-
1999.....	19	-	-	5	3	4	3	2	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Intentional self-harm (suicide) by hanging, strangulation and suffocation (X70)

Asian or Pacific Islander 2/, both sexes

2004.....	322	-	7	23	25	29	36	26	21	34
2003.....	311	-	5	21	27	34	26	33	22	27
2002.....	285	-	3	12	24	33	34	29	24	30
2001.....	266	-	6	20	28	25	32	22	22	17
2000.....	258	-	4	20	24	31	27	19	22	22
1999.....	272	-	3	18	29	32	27	27	18	18

Asian or Pacific Islander 2/, male

2004.....	231	-	7	18	22	21	26	18	14	30
2003.....	223	-	-	13	24	25	21	24	15	18
2002.....	202	-	3	9	22	25	31	23	15	24
2001.....	184	-	6	11	21	14	26	19	19	8
2000.....	177	-	3	15	18	19	18	14	13	16
1999.....	183	-	3	12	18	22	18	19	14	13

Asian or Pacific Islander 2/, female

2004.....	91	-	-	5	3	8	10	8	7	4
2003.....	88	-	5	8	3	9	5	9	7	9
2002.....	83	-	-	3	2	8	3	6	9	6
2001.....	82	-	-	9	7	11	6	3	3	9
2000.....	81	-	1	5	6	12	9	5	9	6
1999.....	89	-	-	6	11	10	9	8	4	5

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intentional self-harm (suicide) by discharge of firearms (X72-X74)												
All races, both sexes												
2004.....	16,750	-	59	787	1,317	1,115	1,168	1,233	1,635	1,784
2003.....	16,907	1	73	736	1,339	1,114	1,267	1,280	1,647	1,705
2002.....	17,108	-	86	742	1,346	1,217	1,182	1,450	1,707	1,652
2001.....	16,869	-	90	838	1,292	1,263	1,301	1,429	1,601	1,552
2000.....	16,586	-	110	897	1,370	1,223	1,230	1,521	1,670	1,491
1999.....	16,599	-	103	975	1,340	1,292	1,329	1,617	1,480	1,462
All races, male												
2004.....	14,523	-	50	688	1,201	987	991	1,039	1,337	1,465
2003.....	14,827	1	62	659	1,220	990	1,109	1,092	1,378	1,419
2002.....	15,045	-	68	668	1,229	1,068	1,010	1,234	1,438	1,394
2001.....	14,758	-	69	743	1,192	1,122	1,124	1,200	1,351	1,309
2000.....	14,454	-	90	793	1,250	1,084	1,039	1,255	1,402	1,263
1999.....	14,479	-	80	867	1,216	1,137	1,133	1,349	1,237	1,221
All races, female												
2004.....	2,227	-	9	99	116	128	177	194	298	319
2003.....	2,080	-	11	77	119	124	158	188	269	286
2002.....	2,063	-	18	74	117	149	172	216	269	258
2001.....	2,111	-	21	95	100	141	177	229	250	243
2000.....	2,132	-	20	104	120	139	191	266	268	228
1999.....	2,120	-	23	108	124	155	196	268	243	241
White, both sexes												
2004.....	15,407	-	49	676	1,094	923	1,014	1,116	1,520	1,679
2003.....	15,587	1	64	638	1,090	927	1,091	1,166	1,550	1,627
2002.....	15,733	-	73	632	1,115	1,011	1,040	1,303	1,597	1,560
2001.....	15,467	-	73	714	1,039	1,068	1,152	1,296	1,474	1,467
2000.....	15,174	-	103	741	1,137	1,012	1,086	1,384	1,549	1,390
1999.....	15,135	-	94	817	1,082	1,081	1,172	1,452	1,381	1,360

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Intentional self-harm (suicide) by discharge of firearms (X72-X74)

White, male

2004.....	13,341	-	41	592	998	815	856	940	1,238	1,376
2003.....	13,649	1	55	569	997	815	950	985	1,299	1,351
2002.....	13,809	-	58	570	1,016	869	888	1,104	1,347	1,313
2001.....	13,521	-	58	628	953	946	990	1,082	1,249	1,243
2000.....	13,214	-	86	653	1,040	893	922	1,133	1,298	1,178
1999.....	13,167	-	72	724	983	945	993	1,200	1,149	1,139

White, female

2004.....	2,066	-	8	84	96	108	158	176	282	303
2003.....	1,938	-	9	69	93	112	141	181	251	276
2002.....	1,924	-	15	62	99	142	152	199	250	247
2001.....	1,946	-	15	86	86	122	162	214	225	224
2000.....	1,960	-	17	88	97	119	164	251	251	212
1999.....	1,968	-	22	93	99	136	179	252	232	221

Black, both sexes

2004.....	1,009	-	8	74	186	148	117	82	74	68
2003.....	1,004	-	7	67	202	148	133	80	71	50
2002.....	1,059	-	9	78	177	162	112	115	86	73
2001.....	1,091	-	13	88	208	153	114	109	91	64
2000.....	1,090	-	4	117	191	161	111	101	93	77
1999.....	1,125	-	5	120	207	156	116	125	74	77

Black, male

2004.....	904	-	7	62	169	136	103	72	67	59
2003.....	902	-	5	64	184	142	120	74	57	46
2002.....	962	-	8	69	165	157	102	101	71	67
2001.....	971	-	10	80	200	137	105	98	73	50
2000.....	965	-	4	109	172	145	95	89	80	66
1999.....	1,018	-	4	109	189	144	106	113	66	64

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Intentional self-harm (suicide) by discharge of firearms (X72-X74)

Black, female

2004.....	105	-	1	12	17	12	14	10	7	9
2003.....	102	-	2	3	18	6	13	6	14	4
2002.....	97	-	1	9	12	5	10	14	15	6
2001.....	120	-	3	8	8	16	9	11	18	14
2000.....	125	-	-	8	19	16	16	12	13	11
1999.....	107	-	1	11	18	12	10	12	8	13

American Indian 1/, both sexes

2004.....	152	-	1	23	19	23	14	19	13	16
2003.....	127	-	1	21	18	19	16	13	13	8
2002.....	146	-	1	17	36	20	13	17	10	9
2001.....	132	-	2	22	18	12	17	14	18	9
2000.....	133	-	1	20	23	14	13	12	13	8
1999.....	136	-	1	25	24	24	13	16	6	9

American Indian 1/, male

2004.....	130	-	1	21	18	19	13	17	11	14
2003.....	117	-	1	18	17	16	15	13	12	8
2002.....	128	-	-	16	30	20	10	16	9	6
2001.....	115	-	-	21	16	11	15	10	16	9
2000.....	109	-	-	17	20	12	8	10	10	4
1999.....	120	-	1	22	21	21	11	15	5	8

American Indian 1/, female

2004.....	22	-	-	2	1	4	1	2	2	2
2003.....	10	-	-	3	1	3	1	-	1	-
2002.....	18	-	1	1	6	-	3	1	1	3
2001.....	17	-	2	1	2	1	2	4	2	-
2000.....	24	-	1	3	3	2	5	2	3	4
1999.....	16	-	-	3	3	3	2	1	1	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Intentional self-harm (suicide) by discharge of firearms (X72-X74)

Asian or Pacific Islander 2/, both sexes

2004.....	182	-	1	14	18	21	23	16	28	21
2003.....	189	-	1	10	29	20	27	21	13	20
2002.....	170	-	3	15	18	24	17	15	14	10
2001.....	179	-	2	14	27	30	18	10	18	12
2000.....	189	-	2	19	19	36	20	24	15	16
1999.....	203	-	3	13	27	31	28	24	19	16

Asian or Pacific Islander 2/, male

2004.....	148	-	1	13	16	17	19	10	21	16
2003.....	159	-	1	8	22	17	24	20	10	14
2002.....	146	-	2	13	18	22	10	13	11	8
2001.....	151	-	1	14	23	28	14	10	13	7
2000.....	166	-	-	14	18	34	14	23	14	15
1999.....	174	-	3	12	23	27	23	21	17	10

Asian or Pacific Islander 2/, female

2004.....	34	-	-	1	2	4	4	6	7	5
2003.....	30	-	-	2	7	3	3	1	3	6
2002.....	24	-	1	2	-	2	7	2	3	2
2001.....	28	-	1	-	4	2	4	-	5	5
2000.....	23	-	2	5	1	2	6	1	1	1
1999.....	29	-	-	1	4	4	5	3	2	6

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Intentional self-harm (suicide) by jumping from a high place (X80)

All races, both sexes

2004.....	676	-	-	39	71	62	61	59	80	55
2003.....	723	-	4	34	67	53	70	80	89	74
2002.....	740	-	3	32	83	68	88	75	82	65
2001.....	651	-	4	37	47	49	60	64	75	68
2000.....	607	-	2	37	71	54	52	69	54	65
1999.....	693	-	2	38	74	57	58	91	82	69

All races, male

2004.....	488	-	-	28	55	49	48	39	61	38
2003.....	509	-	1	26	58	39	53	58	59	53
2002.....	514	-	3	25	67	54	59	54	57	40
2001.....	443	-	3	28	35	35	45	47	48	45
2000.....	432	-	1	29	51	42	36	55	38	50
1999.....	504	-	2	26	61	46	46	65	62	53

All races, female

2004.....	188	-	-	11	16	13	13	20	19	17
2003.....	214	-	3	8	9	14	17	22	30	21
2002.....	226	-	-	7	16	14	29	21	25	25
2001.....	208	-	1	9	12	14	15	17	27	23
2000.....	175	-	1	8	20	12	16	14	16	15
1999.....	189	-	-	12	13	11	12	26	20	16

White, both sexes

2004.....	542	-	-	32	53	41	45	47	67	43
2003.....	592	-	3	24	50	39	55	61	77	60
2002.....	615	-	2	22	65	54	70	63	67	59
2001.....	542	-	2	31	40	37	47	49	61	57
2000.....	503	-	2	29	55	35	41	60	43	61
1999.....	565	-	2	29	55	43	45	72	69	57

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Intentional self-harm (suicide) by jumping from a high place (X80)

White, male

2004.....	400	-	-	24	44	35	37	31	52	31
2003.....	418	-	1	18	45	29	42	45	52	43
2002.....	428	-	2	16	52	42	44	47	50	37
2001.....	366	-	2	24	30	25	36	35	38	39
2000.....	357	-	1	24	40	28	27	46	31	46
1999.....	412	-	2	20	45	35	34	50	56	44

White, female

2004.....	142	-	-	8	9	6	8	16	15	12
2003.....	174	-	2	6	5	10	13	16	25	17
2002.....	187	-	-	6	13	12	26	16	17	22
2001.....	176	-	-	7	10	12	11	14	23	18
2000.....	146	-	1	5	15	7	14	14	12	15
1999.....	153	-	-	9	10	8	11	22	13	13

Black, both sexes

2004.....	67	-	-	2	6	11	10	9	6	9
2003.....	70	-	1	3	9	6	10	13	10	8
2002.....	62	-	-	7	8	9	7	7	9	3
2001.....	69	-	2	3	5	4	9	8	10	9
2000.....	62	-	-	2	13	8	8	7	8	3
1999.....	79	-	-	5	13	9	10	13	8	7

Black, male

2004.....	50	-	-	2	6	8	7	6	4	6
2003.....	51	-	-	3	8	4	8	9	5	8
2002.....	45	-	-	6	7	9	6	4	4	2
2001.....	52	-	1	2	5	4	5	7	7	6
2000.....	46	-	-	1	9	7	6	7	6	3
1999.....	60	-	-	5	10	7	9	10	6	5

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intentional self-harm (suicide) by jumping from a high place (X80)												
Asian or Pacific Islander 2/, both sexes												
2004.....	61	-	-	5	10	8	5	2	7	3
2003.....	52	-	-	5	6	6	5	6	1	5
2002.....	59	-	1	3	8	5	11	4	6	3
2001.....	38	-	-	3	2	8	4	6	4	1
2000.....	36	-	-	5	3	9	3	2	2	-
1999.....	49	-	-	4	6	5	3	6	5	5
Asian or Pacific Islander 2/, male												
2004.....	33	-	-	2	4	4	3	1	5	1
2003.....	36	-	-	5	4	5	3	4	1	2
2002.....	37	-	1	3	6	3	9	2	3	1
2001.....	24	-	-	2	-	6	4	4	3	-
2000.....	25	-	-	4	2	6	3	2	-	-
1999.....	32	-	-	1	6	4	3	5	-	4
Asian or Pacific Islander 2/, female												
2004.....	28	-	-	3	6	4	2	1	2	2
2003.....	16	-	-	-	2	1	2	2	-	3
2002.....	22	-	-	-	2	2	2	2	3	2
2001.....	14	-	-	1	2	2	-	2	1	1
2000.....	11	-	-	1	1	3	-	-	2	-
1999.....	17	-	-	3	-	1	-	1	5	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-harm (suicide) by all other and unspecified means and their sequelae (*U03,X71,X75-X79,X81-X84,X87.0)

All races, both sexes

2004.....	1,877	-	5	74	149	126	133	191	227	251
2003.....	1,757	1	6	46	108	132	153	186	257	221
2002.....	1,859	-	6	71	130	136	150	207	236	213
2001 3/.....	1,713	1	5	68	117	129	142	199	192	237
2000.....	1,610	-	5	62	119	108	127	195	207	183
1999.....	1,587	-	6	65	115	109	153	183	205	170

All races, male

2004.....	1,375	-	1	58	118	104	97	137	171	174
2003.....	1,323	1	2	33	88	111	116	150	187	168
2002.....	1,398	-	5	59	103	100	123	165	177	168
2001 3/.....	1,289	1	3	55	97	107	108	155	138	181
2000.....	1,207	-	4	45	96	89	98	162	160	135
1999.....	1,158	-	4	51	93	92	121	131	141	127

All races, female

2004.....	502	-	4	16	31	22	36	54	56	77
2003.....	434	-	4	13	20	21	37	36	70	53
2002.....	461	-	1	12	27	36	27	42	59	45
2001 3/.....	424	-	2	13	20	22	34	44	54	56
2000.....	403	-	1	17	23	19	29	33	47	48
1999.....	429	-	2	14	22	17	32	52	64	43

White, both sexes

2004.....	1,563	-	4	62	114	100	107	151	189	204
2003.....	1,492	-	5	36	82	96	113	149	222	198
2002.....	1,592	-	4	64	99	104	128	171	205	186
2001 3/.....	1,463	1	4	58	86	100	124	165	169	205
2000.....	1,360	-	4	49	86	89	96	159	173	163
1999.....	1,334	-	3	52	92	83	116	150	176	147

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Intentional self-harm (suicide) by all other and unspecified means and their sequelae (*U03,X71,X75-X79,X81-X84,X87.0)

White, male

2004.....	1,151	-	1	48	91	82	84	109	144	140
2003.....	1,130	-	1	25	66	82	89	119	161	147
2002.....	1,205	-	4	53	79	75	106	137	155	147
2001 3/.....	1,106	1	3	49	70	85	93	127	124	157
2000.....	1,018	-	3	35	71	75	73	133	134	120
1999.....	978	-	2	42	73	72	94	103	123	114

White, female

2004.....	412	-	3	14	23	18	23	42	45	64
2003.....	362	-	4	11	16	14	24	30	61	51
2002.....	387	-	-	11	20	29	22	34	50	39
2001 3/.....	357	-	1	9	16	15	31	38	45	48
2000.....	342	-	1	14	15	14	23	26	39	43
1999.....	356	-	1	10	19	11	22	47	53	33

Black, both sexes

2004.....	184	-	1	4	22	17	13	28	26	29
2003.....	177	1	1	6	19	21	27	26	31	12
2002.....	181	-	1	5	20	20	14	31	19	19
2001 3/.....	169	-	1	5	19	17	16	26	17	20
2000.....	170	-	1	8	27	11	19	28	28	15
1999.....	177	-	2	11	15	15	26	25	24	15

Black, male

2004.....	138	-	-	2	18	15	6	20	21	21
2003.....	133	1	1	4	16	16	20	22	25	12
2002.....	139	-	-	5	16	17	13	25	13	16
2001 3/.....	124	-	-	4	19	15	13	21	11	14
2000.....	133	-	1	7	19	7	15	26	21	12
1999.....	126	-	2	8	13	12	20	21	14	10

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-harm (suicide) by all other and unspecified means and their sequelae (*U03,X71,X75-X79,X81-X84,X87.0)

Asian or Pacific Islander 2/, both sexes

2004.....	110	-	-	7	10	6	11	9	10	15
2003.....	72	-	-	3	4	10	12	8	4	10
2002.....	66	-	1	2	9	7	6	4	9	4
2001 3/.....	67	-	-	4	10	11	1	5	5	11
2000.....	67	-	-	3	6	6	9	6	4	4
1999.....	62	-	1	2	5	9	11	7	4	6

Asian or Pacific Islander 2/, male

2004.....	72	-	-	7	8	5	6	6	4	11
2003.....	46	-	-	3	3	9	6	6	1	8
2002.....	44	-	1	1	7	4	3	2	7	4
2001 3/.....	48	-	-	2	7	6	1	4	2	9
2000.....	46	-	-	1	6	5	7	3	4	2
1999.....	43	-	-	1	4	6	7	6	3	2

Asian or Pacific Islander 2/, female

2004.....	38	-	-	-	2	1	5	3	6	4
2003.....	26	-	-	-	1	1	6	2	3	2
2002.....	22	-	-	1	2	3	3	2	2	-
2001 3/.....	19	-	-	2	3	5	-	1	3	2
2000.....	21	-	-	2	-	1	2	3	-	2
1999.....	19	-	1	1	1	3	4	1	1	4

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Assault (homicide) (*U01-*U02,X85-Y09,X87.1)												
All races, both sexes												
2004.....	17,357	325	377	122	207	1,932	3,153	2,598	1,897	1,530	1,454	1,182
2003.....	17,732	341	376	122	202	1,938	3,430	2,566	1,950	1,636	1,474	1,219
2002.....	17,638	303	423	140	216	1,892	3,327	2,514	1,975	1,734	1,505	1,155
2001 3/.....	20,308	332	415	137	189	1,899	3,398	2,817	2,387	2,306	1,962	1,463
2000.....	16,765	349	356	140	231	1,914	3,025	2,312	1,852	1,732	1,487	1,059
1999.....	16,889	331	376	186	246	2,093	2,905	2,362	1,869	1,751	1,455	998
All races, male												
2004.....	13,578	167	204	62	138	1,631	2,749	2,227	1,537	1,187	1,037	851
2003.....	13,882	205	202	70	145	1,677	2,947	2,191	1,550	1,224	1,084	893
2002.....	13,640	164	234	64	133	1,604	2,870	2,121	1,548	1,288	1,098	841
2001 3/.....	15,555	197	238	75	118	1,632	2,909	2,308	1,849	1,698	1,424	1,085
2000.....	12,820	203	198	77	150	1,613	2,590	1,918	1,443	1,235	1,084	773
1999.....	12,785	187	194	95	155	1,748	2,443	1,922	1,424	1,257	1,050	746
All races, female												
2004.....	3,779	158	173	60	69	301	404	371	360	343	417	331
2003.....	3,850	136	174	52	57	261	483	375	400	412	390	326
2002.....	3,998	139	189	76	83	288	457	393	427	446	407	314
2001 3/.....	4,753	135	177	62	71	267	489	509	538	608	538	378
2000.....	3,945	146	158	63	81	301	435	394	409	497	403	286
1999.....	4,104	144	182	91	91	345	462	440	445	494	405	252
White, both sexes												
2004.....	8,643	208	201	74	110	822	1,304	1,078	842	779	813	683
2003.....	8,709	216	204	61	98	798	1,373	1,042	869	857	796	702
2002.....	8,685	170	239	83	112	830	1,307	1,015	850	891	848	664
2001 3/.....	11,328	196	228	102	116	809	1,455	1,329	1,294	1,406	1,266	958
2000.....	8,339	197	199	98	142	823	1,158	945	869	909	816	616
1999.....	8,628	200	208	106	155	884	1,180	1,006	920	946	820	587

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Assault (homicide) (*U01-*U02,X85-Y09,X87.1)												
White, male												
2004.....	6,302	112	113	36	73	660	1,067	865	636	584	546	470
2003.....	6,337	129	96	34	72	660	1,121	830	657	601	546	487
2002.....	6,282	100	133	35	63	681	1,065	812	623	624	616	446
2001 3/.....	8,254	118	132	52	67	652	1,159	1,032	956	1,058	914	700
2000.....	5,925	125	114	59	89	659	924	721	635	614	588	423
1999.....	6,162	121	108	55	93	704	918	784	661	659	581	429
White, female												
2004.....	2,341	96	88	38	37	162	237	213	206	195	267	213
2003.....	2,372	87	108	27	26	138	252	212	212	256	250	215
2002.....	2,403	70	106	48	49	149	242	203	227	267	232	218
2001 3/.....	3,074	78	96	50	49	157	296	297	338	348	352	258
2000.....	2,414	72	85	39	53	164	234	224	234	295	228	193
1999.....	2,466	79	100	51	62	180	262	222	259	287	239	158
Black, both sexes												
2004.....	8,135	103	163	44	87	1,042	1,761	1,458	987	703	594	453
2003.....	8,392	106	156	54	95	1,068	1,938	1,462	1,013	721	624	466
2002.....	8,287	117	161	45	90	983	1,917	1,432	1,037	779	598	449
2001 3/.....	8,226	123	169	29	65	1,027	1,843	1,392	979	827	623	445
2000.....	7,867	140	145	40	84	1,028	1,791	1,301	923	764	612	397
1999.....	7,648	118	152	67	80	1,131	1,650	1,278	879	741	584	371
Black, male												
2004.....	6,839	48	83	22	60	915	1,607	1,312	851	568	454	349
2003.....	7,083	60	96	33	66	962	1,727	1,310	846	585	502	368
2002.....	6,896	56	92	24	62	859	1,723	1,262	867	616	442	365
2001 3/.....	6,780	70	101	20	48	928	1,672	1,211	817	590	461	343
2000.....	6,482	73	75	16	56	899	1,602	1,146	765	590	450	315
1999.....	6,214	60	75	36	55	978	1,475	1,084	716	556	440	289

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	-------------	-----------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Assault (homicide) (*U01-*U02,X85-Y09,X87.1)

Black, female

2004.....	1,296	55	80	22	27	127	154	146	136	135	140	104
2003.....	1,309	46	60	21	29	106	211	152	167	136	122	98
2002.....	1,391	61	69	21	28	124	194	170	170	163	156	84
2001 3/.....	1,446	53	68	9	17	99	171	181	162	237	162	102
2000.....	1,385	67	70	24	28	129	189	155	158	174	162	82
1999.....	1,434	58	77	31	25	153	175	194	163	185	144	82

American Indian 1/, both sexes

2004.....	228	9	7	2	4	29	38	33	21	24	22	15
2003.....	237	10	7	2	2	31	47	26	25	25	19	23
2002.....	267	10	12	4	7	32	38	29	34	27	25	15
2001 3/.....	211	6	12	1	3	27	32	25	29	17	19	13
2000.....	203	5	8	1	-	25	26	28	31	24	22	9
1999.....	253	5	6	5	4	27	33	34	39	25	19	15

American Indian 1/, male

2004.....	181	5	5	2	2	22	31	28	19	22	16	12
2003.....	175	8	5	1	1	23	35	23	18	17	12	18
2002.....	185	4	5	-	3	25	29	21	25	20	20	13
2001 3/.....	146	3	5	-	1	21	24	22	17	11	13	9
2000.....	159	3	6	1	-	21	25	22	24	16	16	7
1999.....	179	3	4	1	3	22	23	22	28	18	12	11

American Indian 1/, female

2004.....	47	4	2	-	2	7	7	5	2	2	6	3
2003.....	62	2	2	1	1	8	12	3	7	8	7	5
2002.....	82	6	7	4	4	7	9	8	9	7	5	2
2001 3/.....	65	3	7	1	2	6	8	3	12	6	6	4
2000.....	44	2	2	-	-	4	1	6	7	8	6	2
1999.....	74	2	2	4	1	5	10	12	11	7	7	4

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Assault (homicide) (*U01-*U02,X85-Y09,X87.1)

Asian or Pacific Islander 2/, both sexes

2004.....	351	5	6	2	6	39	50	29	47	24	25	31
2003.....	394	9	9	5	7	41	72	36	43	33	35	28
2002.....	399	6	11	8	7	47	65	38	54	37	34	27
2001 3/.....	543	7	6	5	5	36	68	71	85	56	54	47
2000.....	356	7	4	1	5	38	50	38	29	35	37	37
1999.....	360	8	10	8	7	51	42	44	31	39	32	25

Asian or Pacific Islander 2/, male

2004.....	256	2	3	2	3	34	44	22	31	13	21	20
2003.....	287	8	5	2	6	32	64	28	29	21	24	20
2002.....	277	4	4	5	5	39	53	26	33	28	20	17
2001 3/.....	375	6	-	3	2	31	54	43	59	39	36	33
2000.....	254	2	3	1	5	34	39	29	19	15	30	28
1999.....	230	3	7	3	4	44	27	32	19	24	17	17

Asian or Pacific Islander 2/, female

2004.....	95	3	3	-	3	5	6	7	16	11	4	11
2003.....	107	1	4	3	1	9	8	8	14	12	11	8
2002.....	122	2	7	3	2	8	12	12	21	9	14	10
2001 3/.....	168	1	6	2	3	5	14	28	26	17	18	14
2000.....	102	5	1	-	-	4	11	9	10	20	7	9
1999.....	130	5	3	5	3	7	15	12	12	15	15	8

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Assault (homicide) by drugs, biological substances and other and unspecified noxious substances (*U01.6-*U01.7,X85-X90)

All races, both sexes

2004.....	103	14	11	8	5	6	12	6	5	5	8	8
2003.....	81	7	8	6	6	8	7	6	4	6	6	2
2002.....	63	4	11	4	1	4	5	3	5	1	4	2
2001.....	64	10	3	4	3	2	2	2	2	6	8	6
2000.....	56	6	2	2	2	2	1	2	6	8	7	3
1999.....	67	6	5	6	1	2	7	6	5	3	8	4

All races, male

2004.....	61	5	10	3	3	5	5	5	2	4	4	6
2003.....	49	5	6	2	4	3	5	4	2	5	3	1
2002.....	38	2	5	3	-	1	3	2	4	1	4	2
2001.....	37	8	2	2	1	1	-	2	2	3	4	3
2000.....	25	4	-	2	2	1	1	1	3	3	2	2
1999.....	34	4	2	2	1	-	2	2	4	3	5	4

All races, female

2004.....	42	9	1	5	2	1	7	1	3	1	4	2
2003.....	32	2	2	4	2	5	2	2	2	1	3	1
2002.....	25	2	6	1	1	3	2	1	1	-	-	-
2001.....	27	2	1	2	2	1	2	-	-	3	4	3
2000.....	31	2	2	-	-	1	-	1	3	5	5	1
1999.....	33	2	3	4	-	2	5	4	1	-	3	-

White, both sexes

2004.....	77	6	8	6	5	6	11	4	4	4	5	7
2003.....	53	6	4	4	2	7	6	3	3	4	4	2
2002.....	51	2	6	4	1	3	4	3	5	-	4	2
2001.....	43	1	1	3	2	1	2	2	1	5	5	5
2000.....	44	2	2	2	2	2	1	2	5	5	7	2
1999.....	60	5	4	6	1	2	7	5	4	3	6	4

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Assault (homicide) by drugs, biological substances and other and unspecified noxious substances (*U01.6-*U01.7,X85-X90)

White, male

2004.....	47	3	7	3	3	5	4	3	1	4	2	5
2003.....	31	4	3	2	2	2	4	3	2	3	1	1
2002.....	34	2	3	3	-	1	3	2	4	-	4	2
2001.....	23	-	1	1	1	1	-	2	1	3	2	2
2000.....	20	2	-	2	2	1	1	1	3	2	2	1
1999.....	30	4	2	2	1	-	2	2	3	3	3	4

White, female

2004.....	30	3	1	3	2	1	7	1	3	-	3	2
2003.....	22	2	1	2	-	5	2	-	1	1	3	1
2002.....	17	-	3	1	1	2	1	1	1	-	-	-
2001.....	20	1	-	2	1	-	2	-	-	2	3	3
2000.....	24	-	2	-	-	1	-	1	2	3	5	1
1999.....	30	1	2	4	-	2	5	3	1	-	3	-

Black, both sexes

2004.....	23	7	3	2	-	-	1	2	1	1	3	1
2003.....	26	1	3	2	3	1	1	3	1	2	2	-
2002.....	7	1	4	-	-	-	1	-	-	-	-	-
2001.....	18	8	2	1	1	1	-	-	1	1	2	-
2000.....	10	4	-	-	-	-	-	-	1	2	-	1
1999.....	7	1	1	-	-	-	-	1	1	-	2	-

Black, male

2004.....	13	1	3	-	-	-	1	2	1	-	2	1
2003.....	18	1	3	-	2	1	1	1	-	2	2	-
2002.....	3	-	2	-	-	-	-	-	-	-	-	-
2001.....	11	7	1	1	-	-	-	-	1	-	1	-
2000.....	4	2	-	-	-	-	-	-	-	-	-	1
1999.....	4	-	-	-	-	-	-	-	1	-	2	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Assault (homicide) by hanging, strangulation and suffocation (X91)

All races, both sexes

2004.....	664	42	29	12	15	29	77	68	57	62	85	56
2003.....	670	39	17	14	10	51	68	62	65	76	67	48
2002.....	679	32	30	14	12	46	70	65	60	88	63	72
2001.....	690	40	18	10	6	39	76	63	69	73	72	57
2000.....	658	26	26	13	19	41	78	64	72	69	68	50
1999.....	708	34	21	14	19	45	69	81	90	88	65	37

All races, male

2004.....	214	16	14	4	5	6	23	14	20	17	23	18
2003.....	220	21	6	6	5	11	19	12	17	28	25	12
2002.....	238	17	13	3	2	16	21	22	15	24	15	35
2001.....	244	24	10	3	-	9	23	19	16	23	20	25
2000.....	224	15	10	6	7	8	18	12	29	19	29	19
1999.....	238	22	11	6	7	9	15	20	27	28	17	14

All races, female

2004.....	450	26	15	8	10	23	54	54	37	45	62	38
2003.....	450	18	11	8	5	40	49	50	48	48	42	36
2002.....	441	15	17	11	10	30	49	43	45	64	48	37
2001.....	446	16	8	7	6	30	53	44	53	50	52	32
2000.....	434	11	16	7	12	33	60	52	43	50	39	31
1999.....	470	12	10	8	12	36	54	61	63	60	48	23

White, both sexes

2004.....	425	29	17	6	9	17	51	44	34	32	48	38
2003.....	437	27	13	7	6	28	41	42	43	48	51	25
2002.....	427	20	15	10	10	23	49	38	33	52	36	44
2001.....	454	23	12	9	4	29	45	42	46	47	39	37
2000.....	431	19	22	11	14	25	48	40	40	44	36	37
1999.....	450	22	14	12	16	25	43	46	53	51	37	24

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Assault (homicide) by hanging, strangulation and suffocation (X91)

White, male

2004.....	141	8	10	3	3	1	17	10	14	8	13	12
2003.....	149	15	5	3	3	7	10	7	15	20	19	6
2002.....	153	12	4	3	1	12	12	14	8	14	12	18
2001.....	165	15	6	2	-	6	12	13	11	19	7	18
2000.....	163	11	10	5	5	7	9	9	19	15	18	14
1999.....	161	14	7	5	7	5	8	12	17	17	11	9

White, female

2004.....	284	21	7	3	6	16	34	34	20	24	35	26
2003.....	288	12	8	4	3	21	31	35	28	28	32	19
2002.....	274	8	11	7	9	11	37	24	25	38	24	26
2001.....	289	8	6	7	4	23	33	29	35	28	32	19
2000.....	268	8	12	6	9	18	39	31	21	29	18	23
1999.....	289	8	7	7	9	20	35	34	36	34	26	15

Black, both sexes

2004.....	212	11	11	6	6	11	21	22	16	27	37	16
2003.....	202	11	4	7	4	21	23	18	18	25	14	19
2002.....	216	7	10	4	2	17	20	24	25	33	23	24
2001.....	212	16	6	1	-	10	28	19	20	26	29	17
2000.....	204	6	4	2	5	15	27	23	28	23	30	9
1999.....	222	11	6	2	2	18	21	31	31	33	22	9

Black, male

2004.....	63	7	4	1	2	5	3	3	4	8	10	5
2003.....	61	5	1	3	2	4	7	4	2	7	6	4
2002.....	69	3	6	-	1	2	9	6	6	9	1	14
2001.....	70	8	4	1	-	3	9	5	5	4	10	6
2000.....	49	3	-	1	2	1	7	3	7	4	10	2
1999.....	67	7	3	1	-	3	7	8	8	10	4	3

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Assault (homicide) by hanging, strangulation and suffocation (X91)												
Black, female												
2004.....	149	4	7	5	4	6	18	19	12	19	27	11
2003.....	141	6	3	4	2	17	16	14	16	18	8	15
2002.....	147	4	4	4	1	15	11	18	19	24	22	10
2001.....	142	8	2	-	-	7	19	14	15	22	19	11
2000.....	155	3	4	1	3	14	20	20	21	19	20	7
1999.....	155	4	3	1	2	15	14	23	23	23	18	6
American Indian 1/, both sexes												
2004.....	5	-	-	-	-	1	2	-	-	-	-	1
2003.....	9	-	-	-	-	1	1	-	2	2	-	1
2002.....	4	1	1	-	-	-	-	1	-	-	-	-
2001.....	7	1	-	-	-	-	-	-	1	-	2	1
2000.....	9	1	-	-	-	1	1	-	3	2	-	1
1999.....	11	-	1	-	-	-	1	3	2	2	1	1
American Indian 1/, male												
2004.....	2	-	-	-	-	-	1	-	-	-	-	1
2003.....	3	-	-	-	-	-	-	-	-	1	-	1
2002.....	2	-	1	-	-	-	-	1	-	-	-	-
2001.....	3	1	-	-	-	-	-	-	-	-	2	-
2000.....	5	1	-	-	-	-	1	-	2	-	-	1
1999.....	4	-	1	-	-	-	-	-	1	-	1	1
American Indian 1/, female												
2004.....	3	-	-	-	-	1	1	-	-	-	-	-
2003.....	6	-	-	-	-	1	1	-	2	1	-	-
2002.....	2	1	-	-	-	-	-	-	-	-	-	-
2001.....	4	-	-	-	-	-	-	-	1	-	-	1
2000.....	4	-	-	-	-	1	-	-	1	2	-	-
1999.....	7	-	-	-	-	-	1	3	1	2	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Assault (homicide) by hanging, strangulation and suffocation (X91)

Asian or Pacific Islander 2/, both sexes

2004.....	22	2	1	-	-	-	3	2	7	3	-	1
2003.....	22	1	-	-	-	1	3	2	2	1	2	3
2002.....	32	4	4	-	-	6	1	2	2	3	4	4
2001.....	17	-	-	-	2	-	3	2	2	-	2	2
2000.....	14	-	-	-	-	-	2	1	1	-	2	3
1999.....	25	1	-	-	1	2	4	1	4	2	5	3

Asian or Pacific Islander 2/, male

2004.....	8	1	-	-	-	-	2	1	2	1	-	-
2003.....	7	1	-	-	-	-	2	1	-	-	-	1
2002.....	14	2	2	-	-	2	-	1	1	1	2	3
2001.....	6	-	-	-	-	-	2	1	-	-	1	1
2000.....	7	-	-	-	-	-	1	-	1	-	1	2
1999.....	6	1	-	-	-	1	-	-	1	1	1	1

Asian or Pacific Islander 2/, female

2004.....	14	1	1	-	-	-	1	1	5	2	-	1
2003.....	15	-	-	-	-	1	1	1	2	1	2	2
2002.....	18	2	2	-	-	4	1	1	1	2	2	1
2001.....	11	-	-	-	2	-	1	1	2	-	1	1
2000.....	7	-	-	-	-	-	1	1	-	-	1	1
1999.....	19	-	-	-	1	1	4	1	3	1	4	2

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)												
All races, both sexes												
2004.....	11,624	6	36	45	139	1,578	2,549	2,063	1,440	1,025	870	638
2003.....	11,920	8	40	48	139	1,587	2,823	2,070	1,470	1,066	875	683
2002.....	11,829	9	49	55	150	1,567	2,750	2,018	1,447	1,135	907	604
2001.....	11,348	11	55	59	121	1,525	2,675	1,940	1,368	1,142	836	552
2000.....	10,801	12	28	50	137	1,549	2,414	1,789	1,285	1,064	842	582
1999.....	10,828	8	50	61	163	1,708	2,330	1,795	1,253	1,054	814	551
All races, male												
2004.....	9,921	2	18	24	110	1,396	2,322	1,876	1,235	877	679	485
2003.....	10,126	7	20	27	107	1,436	2,541	1,867	1,258	860	675	534
2002.....	9,899	3	30	26	101	1,384	2,467	1,790	1,227	918	705	445
2001.....	9,532	7	27	35	90	1,377	2,419	1,708	1,156	892	636	434
2000.....	9,006	9	17	28	110	1,381	2,167	1,574	1,076	842	645	445
1999.....	8,944	5	26	37	118	1,498	2,071	1,568	1,041	807	607	426
All races, female												
2004.....	1,703	4	18	21	29	182	227	187	205	148	191	153
2003.....	1,794	1	20	21	32	151	282	203	212	206	200	149
2002.....	1,930	6	19	29	49	183	283	228	220	217	202	159
2001.....	1,816	4	28	24	31	148	256	232	212	250	200	118
2000.....	1,795	3	11	22	27	168	247	215	209	222	197	137
1999.....	1,884	3	24	24	45	210	259	227	212	247	207	125
White, both sexes												
2004.....	5,119	3	11	26	66	617	913	748	594	485	469	359
2003.....	5,173	2	20	22	59	604	1,030	742	589	520	446	388
2002.....	5,185	3	33	35	71	649	951	738	563	533	494	343
2001.....	5,188	8	33	47	72	593	985	741	586	583	482	324
2000.....	4,806	5	17	39	75	617	817	645	547	525	460	338
1999.....	4,918	4	26	32	95	662	822	675	562	555	456	313

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Assault (homicide) by discharge of firearms (*U01.4,X93-X95)

White, male

2004.....	4,078	1	5	11	53	524	795	659	482	401	333	259
2003.....	4,108	1	8	10	48	528	898	641	476	384	321	281
2002.....	4,050	1	20	18	46	556	824	621	449	399	374	228
2001.....	4,042	5	17	28	50	508	843	627	456	443	348	242
2000.....	3,686	2	9	22	59	531	693	528	427	384	340	246
1999.....	3,820	4	14	20	66	567	688	568	440	408	328	230

White, female

2004.....	1,041	2	6	15	13	93	118	89	112	84	136	100
2003.....	1,065	1	12	12	11	76	132	101	113	136	125	107
2002.....	1,135	2	13	17	25	93	127	117	114	134	120	115
2001.....	1,146	3	16	19	22	85	142	114	130	140	134	82
2000.....	1,120	3	8	17	16	86	124	117	120	141	120	92
1999.....	1,098	-	12	12	29	95	134	107	122	147	128	83

Black, both sexes

2004.....	6,201	3	22	16	65	914	1,586	1,277	812	515	378	253
2003.....	6,397	6	19	23	72	932	1,715	1,286	839	515	400	269
2002.....	6,285	6	14	13	68	865	1,724	1,240	842	564	381	244
2001.....	5,885	2	18	12	45	902	1,635	1,155	746	540	333	207
2000.....	5,699	7	8	11	58	888	1,546	1,097	715	509	354	223
1999.....	5,577	2	18	23	59	991	1,459	1,068	656	468	336	222

Black, male

2004.....	5,589	1	11	10	52	832	1,483	1,186	726	456	323	207
2003.....	5,740	6	11	15	52	869	1,574	1,191	746	454	332	234
2002.....	5,575	2	9	6	48	781	1,583	1,139	746	487	310	206
2001.....	5,279	1	9	7	37	842	1,531	1,050	673	434	274	175
2000.....	5,084	7	5	6	47	812	1,431	1,009	628	439	284	183
1999.....	4,883	1	7	15	46	883	1,344	961	576	378	265	183

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Assault (homicide) by discharge of firearms (*U01.4,X93-X95)

Black, female

2004.....	612	2	11	6	13	82	103	91	86	59	55	46
2003.....	657	-	8	8	20	63	141	95	93	61	68	35
2002.....	710	4	5	7	20	84	141	101	96	77	71	38
2001.....	606	1	9	5	8	60	104	105	73	106	59	32
2000.....	615	-	3	5	11	76	115	88	87	70	70	40
1999.....	694	1	11	8	13	108	115	107	80	90	71	39

American Indian 1/, both sexes

2004.....	104	-	2	2	4	17	21	21	8	12	7	6
2003.....	109	-	-	2	2	20	29	13	12	7	10	9
2002.....	117	-	1	2	5	20	24	15	14	12	10	5
2001.....	87	1	4	-	2	12	17	13	14	4	8	4
2000.....	86	-	2	-	-	14	18	17	10	9	8	2
1999.....	104	2	2	2	4	16	17	16	14	7	4	4

American Indian 1/, male

2004.....	87	-	2	2	2	15	16	17	7	11	7	4
2003.....	81	-	-	1	1	14	24	12	10	4	6	7
2002.....	85	-	-	-	2	15	17	11	11	10	8	3
2001.....	64	1	1	-	1	10	14	12	8	2	6	2
2000.....	70	-	2	-	-	12	17	13	9	7	4	2
1999.....	82	-	2	1	3	15	14	12	12	5	2	3

American Indian 1/, female

2004.....	17	-	-	-	2	2	5	4	1	1	-	2
2003.....	28	-	-	1	1	6	5	1	2	3	4	2
2002.....	32	-	1	2	3	5	7	4	3	2	2	2
2001.....	23	-	3	-	1	2	3	1	6	2	2	2
2000.....	16	-	-	-	-	2	1	4	1	2	4	-
1999.....	22	2	-	1	1	1	3	4	2	2	2	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Assault (homicide) by discharge of firearms (*U01.4,X93-X95)

Asian or Pacific Islander 2/, both sexes

2004.....	200	-	1	1	4	30	29	17	26	13	16	20
2003.....	241	-	1	1	6	31	49	29	30	24	19	17
2002.....	242	-	1	5	6	33	51	25	28	26	22	12
2001.....	188	-	-	-	2	18	38	31	22	15	13	17
2000.....	210	-	1	-	4	30	33	30	13	21	20	19
1999.....	229	-	4	4	5	39	32	36	21	24	18	12

Asian or Pacific Islander 2/, male

2004.....	167	-	-	1	3	25	28	14	20	9	16	15
2003.....	197	-	1	1	6	25	45	23	26	18	16	12
2002.....	189	-	1	2	5	32	43	19	21	22	13	8
2001.....	147	-	-	-	2	17	31	19	19	13	8	15
2000.....	166	-	1	-	4	26	26	24	12	12	17	14
1999.....	159	-	3	1	3	33	25	27	13	16	12	10

Asian or Pacific Islander 2/, female

2004.....	33	-	1	-	1	5	1	3	6	4	-	5
2003.....	44	-	-	-	-	6	4	6	4	6	3	5
2002.....	53	-	-	3	1	1	8	6	7	4	9	4
2001.....	41	-	-	-	-	1	7	12	3	2	5	2
2000.....	44	-	-	-	-	4	7	6	1	9	3	5
1999.....	70	-	1	3	2	6	7	9	8	8	6	2

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	-------------	-----------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Assault (homicide) by sharp object (X99)

All races, both sexes

2004.....	2,079	3	16	12	22	171	313	264	215	219	231	202
2003.....	2,049	4	12	10	14	169	302	242	224	238	241	198
2002.....	2,074	6	8	21	16	144	281	255	244	248	259	184
2001.....	1,971	2	10	8	16	165	316	261	211	247	211	160
2000.....	1,805	7	12	13	20	156	253	205	215	242	224	134
1999.....	1,879	3	18	13	20	159	252	218	249	249	221	148

All races, male

2004.....	1,474	2	7	7	10	134	248	190	154	146	161	134
2003.....	1,453	2	6	4	10	136	228	180	156	169	170	141
2002.....	1,470	5	5	12	11	113	228	185	154	171	174	140
2001.....	1,375	1	5	2	9	132	243	195	147	153	146	122
2000.....	1,283	3	8	7	8	116	208	155	153	157	151	96
1999.....	1,350	1	4	10	7	124	187	161	178	178	174	109

All races, female

2004.....	605	1	9	5	12	37	65	74	61	73	70	68
2003.....	596	2	6	6	4	33	74	62	68	69	71	57
2002.....	604	1	3	9	5	31	53	70	90	77	85	44
2001.....	596	1	5	6	7	33	73	66	64	94	65	38
2000.....	522	4	4	6	12	40	45	50	62	85	73	38
1999.....	529	2	14	3	13	35	65	57	71	71	47	39

White, both sexes

2004.....	1,161	3	10	9	17	90	191	158	107	125	115	103
2003.....	1,108	2	8	7	9	79	150	141	125	124	110	108
2002.....	1,139	4	4	11	9	70	164	133	121	145	136	92
2001.....	1,104	1	5	8	9	81	190	143	110	131	105	79
2000.....	1,024	5	6	11	14	95	138	118	125	134	107	64
1999.....	1,056	1	9	11	12	87	142	135	135	121	118	77

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Assault (homicide) by sharp object (X99)

White, male

2004.....	803	2	4	5	8	75	150	107	69	85	80	63
2003.....	765	1	4	3	7	66	111	103	91	90	71	73
2002.....	809	3	2	6	6	57	133	99	74	110	92	68
2001.....	767	-	3	2	4	69	150	107	79	83	71	56
2000.....	731	2	5	6	5	73	115	91	87	89	76	43
1999.....	750	-	2	8	5	67	109	105	95	81	89	60

White, female

2004.....	358	1	6	4	9	15	41	51	38	40	35	40
2003.....	343	1	4	4	2	13	39	38	34	34	39	35
2002.....	330	1	2	5	3	13	31	34	47	35	44	24
2001.....	337	1	2	6	5	12	40	36	31	48	34	23
2000.....	293	3	1	5	9	22	23	27	38	45	31	21
1999.....	306	1	7	3	7	20	33	30	40	40	29	17

Black, both sexes

2004.....	818	-	5	2	5	70	103	94	92	85	107	92
2003.....	837	1	2	-	5	78	131	95	87	104	121	82
2002.....	813	1	4	7	6	64	100	110	95	92	112	80
2001.....	759	-	3	-	7	70	108	108	79	104	101	73
2000.....	693	2	6	2	6	49	99	78	73	100	107	67
1999.....	730	2	6	2	8	63	102	71	99	114	93	63

Black, male

2004.....	586	-	2	1	2	49	81	72	71	53	75	65
2003.....	619	-	1	-	3	60	100	72	59	72	93	62
2002.....	581	1	3	4	4	49	80	79	65	56	73	63
2001.....	539	-	2	-	5	53	78	79	55	65	71	62
2000.....	488	1	3	1	3	32	78	58	57	64	67	51
1999.....	534	1	1	2	2	49	71	47	72	89	77	44

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Assault (homicide) by sharp object (X99)

Black, female

2004.....	232	-	3	1	3	21	22	22	21	32	32	27
2003.....	218	1	1	-	2	18	31	23	28	32	28	20
2002.....	232	-	1	3	2	15	20	31	30	36	39	17
2001.....	220	-	1	-	2	17	30	29	24	39	30	11
2000.....	205	1	3	1	3	17	21	20	16	36	40	16
1999.....	196	1	5	-	6	14	31	24	27	25	16	19

American Indian 1/, both sexes

2004.....	47	-	-	-	-	7	7	8	5	8	5	4
2003.....	40	-	-	-	-	7	8	3	5	8	4	2
2002.....	60	1	-	1	1	8	7	5	11	8	7	5
2001.....	41	-	-	-	-	4	7	5	8	6	3	4
2000.....	38	-	-	-	-	6	6	5	10	2	3	-
1999.....	57	-	1	-	-	5	7	8	13	6	5	7

American Indian 1/, male

2004.....	44	-	-	-	-	6	7	7	5	8	5	3
2003.....	30	-	-	-	-	6	6	3	4	6	2	1
2002.....	43	1	-	-	1	7	6	3	7	4	7	5
2001.....	29	-	-	-	-	3	5	5	5	3	2	3
2000.....	30	-	-	-	-	5	6	4	7	2	3	-
1999.....	42	-	-	-	-	4	7	6	9	5	4	4

American Indian 1/, female

2004.....	3	-	-	-	-	1	-	1	-	-	-	1
2003.....	10	-	-	-	-	1	2	-	1	2	2	1
2002.....	17	-	-	1	-	1	1	2	4	4	-	-
2001.....	12	-	-	-	-	1	2	-	3	3	1	1
2000.....	8	-	-	-	-	1	-	1	3	-	-	-
1999.....	15	-	1	-	-	1	-	2	4	1	1	3

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Assault (homicide) by sharp object (X99)												
Asian or Pacific Islander 2/, both sexes												
2004.....	53	-	1	1	-	4	12	4	11	1	4	3
2003.....	64	1	2	3	-	5	13	3	7	2	6	6
2002.....	62	-	-	2	-	2	10	7	17	3	4	7
2001.....	67	1	2	-	-	10	11	5	14	6	2	4
2000.....	50	-	-	-	-	6	10	4	7	6	7	3
1999.....	36	-	2	-	-	4	1	4	2	8	5	1
Asian or Pacific Islander 2/, male												
2004.....	41	-	1	1	-	4	10	4	9	-	1	3
2003.....	39	1	1	1	-	4	11	2	2	1	4	5
2002.....	37	-	-	2	-	-	9	4	8	1	2	4
2001.....	40	1	-	-	-	7	10	4	8	2	2	1
2000.....	34	-	-	-	-	6	9	2	2	2	5	2
1999.....	24	-	1	-	-	4	-	3	2	3	4	1
Asian or Pacific Islander 2/, female												
2004.....	12	-	-	-	-	-	2	-	2	1	3	-
2003.....	25	-	1	2	-	1	2	1	5	1	2	1
2002.....	25	-	-	-	-	2	1	3	9	2	2	3
2001.....	27	-	2	-	-	3	1	1	6	4	-	3
2000.....	16	-	-	-	-	-	1	2	5	4	2	1
1999.....	12	-	1	-	-	-	1	1	-	5	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Assault (homicide) by blunt object (Y00)

All races, both sexes

2004.....	102	3	5	-	-	8	7	5	8	9	9	9
2003.....	129	6	4	2	2	4	8	9	10	5	15	15
2002.....	184	5	10	1	1	10	16	11	18	29	15	18
2001.....	199	7	7	-	-	7	10	18	9	17	31	30
2000.....	210	5	9	1	2	11	13	11	19	17	33	26
1999.....	242	7	13	4	4	7	25	27	20	21	24	15

All races, male

2004.....	66	1	3	-	-	6	6	2	6	5	5	8
2003.....	94	5	2	2	2	4	5	5	6	4	12	12
2002.....	129	3	5	-	-	8	13	9	11	21	9	14
2001.....	131	6	3	-	-	4	4	11	5	8	21	23
2000.....	142	2	4	1	1	9	11	6	12	10	24	20
1999.....	158	3	4	-	2	6	15	13	14	17	19	12

All races, female

2004.....	36	2	2	-	-	2	1	3	2	4	4	1
2003.....	35	1	2	-	-	-	3	4	4	1	3	3
2002.....	55	2	5	1	1	2	3	2	7	8	6	4
2001.....	68	1	4	-	-	3	6	7	4	9	10	7
2000.....	68	3	5	-	1	2	2	5	7	7	9	6
1999.....	84	4	9	4	2	1	10	14	6	4	5	3

White, both sexes

2004.....	66	3	2	-	-	6	5	3	3	7	7	4
2003.....	82	3	2	2	1	1	3	5	6	3	10	9
2002.....	131	4	8	1	1	8	9	7	12	20	9	13
2001.....	127	5	2	-	-	5	9	9	3	14	19	20
2000.....	128	2	6	1	2	7	11	4	9	11	20	17
1999.....	167	4	6	2	3	4	17	16	12	15	17	9

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Assault (homicide) by blunt object (Y00)												
White, male												
2004.....	46	1	2	-	-	4	5	1	3	4	4	4
2003.....	59	2	1	2	1	1	2	5	4	2	8	7
2002.....	88	3	4	-	-	6	8	6	7	13	5	10
2001.....	80	4	-	-	-	3	4	7	2	7	13	14
2000.....	87	2	2	1	1	6	9	2	7	5	13	13
1999.....	113	2	2	-	1	3	11	10	9	13	14	6
White, female												
2004.....	20	2	-	-	-	2	-	2	-	3	3	-
2003.....	23	1	1	-	-	-	1	-	2	1	2	2
2002.....	43	1	4	1	1	2	1	1	5	7	4	3
2001.....	47	1	2	-	-	2	5	2	1	7	6	6
2000.....	41	-	4	-	1	1	2	2	2	6	7	4
1999.....	54	2	4	2	2	1	6	6	3	2	3	3
Black, both sexes												
2004.....	29	-	2	-	-	2	2	1	3	2	1	4
2003.....	36	3	2	-	1	2	3	3	4	1	4	4
2002.....	51	1	2	-	-	2	7	4	6	8	5	5
2001.....	61	2	5	-	-	1	1	8	5	3	10	10
2000.....	70	2	3	-	-	4	2	6	10	6	10	6
1999.....	65	2	7	2	1	2	7	9	6	6	7	5
Black, male												
2004.....	17	-	1	-	-	2	1	-	2	1	-	4
2003.....	27	3	1	-	1	2	2	-	2	1	4	3
2002.....	40	-	1	-	-	2	5	3	4	7	4	4
2001.....	45	2	3	-	-	1	-	4	2	1	8	9
2000.....	47	-	2	-	-	3	2	3	5	5	8	4
1999.....	40	1	2	-	1	2	3	3	4	4	5	5

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Assault (homicide) by blunt object (Y00)

Black, female

2004.....	12	-	1	-	-	-	1	1	1	1	1	-
2003.....	9	-	1	-	-	-	1	3	2	-	-	1
2002.....	11	1	1	-	-	-	2	1	2	1	1	1
2001.....	16	-	2	-	-	-	1	4	3	2	2	1
2000.....	23	2	1	-	-	1	-	3	5	1	2	2
1999.....	25	1	5	2	-	-	4	6	2	2	2	-

American Indian 1/, both sexes

2004.....	4	-	-	-	-	-	-	1	1	-	1	-
2003.....	6	-	-	-	-	-	1	1	-	1	-	2
2002.....	1	-	-	-	-	-	-	-	-	1	-	-
2001.....	2	-	-	-	-	-	-	-	1	-	-	-
2000.....	5	1	-	-	-	-	-	1	-	-	1	1
1999.....	5	-	-	-	-	-	1	1	2	-	-	1

American Indian 1/, male

2004.....	3	-	-	-	-	-	-	1	1	-	1	-
2003.....	4	-	-	-	-	-	-	-	-	1	-	2
2002.....	1	-	-	-	-	-	-	-	-	1	-	-
2001.....	2	-	-	-	-	-	-	-	1	-	-	-
2000.....	3	-	-	-	-	-	-	1	-	-	1	1
1999.....	3	-	-	-	-	-	1	-	1	-	-	1

American Indian 1/, female

2004.....	1	-	-	-	-	-	-	-	-	-	-	-
2003.....	2	-	-	-	-	-	1	1	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	2	1	-	-	-	-	-	-	-	-	-	-
1999.....	2	-	-	-	-	-	-	1	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Assault (homicide) by blunt object (Y00)												
Asian or Pacific Islander 2/, both sexes												
2004.....	3	-	1	-	-	-	-	-	1	-	-	1
2003.....	5	-	-	-	-	1	1	-	-	-	1	-
2002.....	1	-	-	-	-	-	-	-	-	-	1	-
2001.....	9	-	-	-	-	1	-	1	-	-	2	-
2000.....	7	-	-	-	-	-	-	-	-	-	2	2
1999.....	5	1	-	-	-	1	-	1	-	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	4	-	-	-	-	1	1	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	4	-	-	-	-	-	-	-	-	-	-	-
2000.....	5	-	-	-	-	-	-	-	-	-	2	2
1999.....	2	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	3	-	1	-	-	-	-	-	1	-	-	1
2003.....	1	-	-	-	-	-	-	-	-	-	1	-
2002.....	1	-	-	-	-	-	-	-	-	-	1	-
2001.....	5	-	-	-	-	1	-	1	-	-	2	-
2000.....	2	-	-	-	-	-	-	-	-	-	-	-
1999.....	3	1	-	-	-	-	-	1	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Assault (homicide) by bodily force (Y04)

All races, both sexes

2004.....	61	-	1	-	1	7	6	4	3	4	12	7
2003.....	93	2	1	-	-	7	6	7	9	8	8	17
2002.....	103	1	3	-	1	7	13	9	11	7	11	17
2001.....	142	3	4	-	-	9	14	9	13	13	24	18
2000.....	139	1	3	1	-	8	13	16	18	16	22	15
1999.....	166	4	2	1	-	11	17	20	14	26	23	15

All races, male

2004.....	49	-	1	-	-	7	6	4	-	4	9	5
2003.....	82	1	1	-	-	7	6	7	9	7	7	14
2002.....	88	-	-	-	1	7	10	8	10	7	10	15
2001.....	106	3	3	-	-	9	12	6	9	8	17	12
2000.....	124	1	2	1	-	7	12	16	17	14	21	15
1999.....	139	3	1	-	-	10	16	17	13	21	18	13

All races, female

2004.....	12	-	-	-	1	-	-	-	3	-	3	2
2003.....	11	1	-	-	-	-	-	-	-	1	1	3
2002.....	15	1	3	-	-	-	3	1	1	-	1	2
2001.....	36	-	1	-	-	-	2	3	4	5	7	6
2000.....	15	-	1	-	-	1	1	-	1	2	1	-
1999.....	27	1	1	1	-	1	1	3	1	5	5	2

White, both sexes

2004.....	36	-	-	-	1	3	3	2	1	2	8	5
2003.....	63	2	-	-	-	3	5	6	5	7	7	10
2002.....	63	-	3	-	-	6	10	4	5	3	6	10
2001.....	102	3	3	-	-	7	10	7	8	7	17	12
2000.....	85	-	2	1	-	3	6	9	12	8	14	11
1999.....	97	3	1	-	-	6	9	6	10	15	13	11

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Assault (homicide) by bodily force (Y04)

White, male

2004.....	29	-	-	-	-	3	3	2	-	2	6	4
2003.....	56	1	-	-	-	3	5	6	5	6	6	9
2002.....	52	-	-	-	-	6	7	4	4	3	6	9
2001.....	76	3	2	-	-	7	8	5	6	4	11	9
2000.....	75	-	1	1	-	3	6	9	11	8	13	11
1999.....	82	3	1	-	-	5	9	6	9	12	10	9

White, female

2004.....	7	-	-	-	1	-	-	-	1	-	2	1
2003.....	7	1	-	-	-	-	-	-	-	1	1	1
2002.....	11	-	3	-	-	-	3	-	1	-	-	1
2001.....	26	-	1	-	-	-	2	2	2	3	6	3
2000.....	10	-	1	-	-	-	-	-	1	-	1	-
1999.....	15	-	-	-	-	1	-	-	1	3	3	2

Black, both sexes

2004.....	22	-	1	-	-	3	2	2	2	1	4	2
2003.....	27	-	1	-	-	4	1	-	4	1	1	7
2002.....	33	1	-	-	1	1	2	3	4	4	4	7
2001.....	33	-	1	-	-	2	4	2	4	3	6	5
2000.....	46	1	1	-	-	4	6	7	4	7	8	3
1999.....	60	1	1	1	-	2	7	13	4	10	9	4

Black, male

2004.....	17	-	1	-	-	3	2	2	-	1	3	1
2003.....	23	-	1	-	-	4	1	-	4	1	1	5
2002.....	30	-	-	-	1	1	2	3	4	4	3	6
2001.....	25	-	1	-	-	2	4	1	3	2	5	2
2000.....	41	1	1	-	-	3	5	7	4	5	8	3
1999.....	49	-	-	-	-	2	6	11	4	8	7	4

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Assault (homicide) by bodily force (Y04)												
Black, female												
2004.....	5	-	-	-	-	-	-	-	2	-	1	1
2003.....	4	-	-	-	-	-	-	-	-	-	-	2
2002.....	3	1	-	-	-	-	-	-	-	-	1	1
2001.....	8	-	-	-	-	-	-	1	1	1	1	3
2000.....	5	-	-	-	-	1	1	-	-	2	-	-
1999.....	11	1	1	1	-	-	1	2	-	2	2	-
American Indian 1/, both sexes												
2004.....	2	-	-	-	-	1	1	-	-	-	-	-
2003.....	2	-	-	-	-	-	-	1	-	-	-	-
2002.....	3	-	-	-	-	-	1	1	1	-	-	-
2001.....	4	-	-	-	-	-	-	-	-	2	1	1
2000.....	7	-	-	-	-	1	1	-	2	1	-	-
1999.....	5	-	-	-	-	2	-	1	-	1	1	-
American Indian 1/, male												
2004.....	2	-	-	-	-	1	1	-	-	-	-	-
2003.....	2	-	-	-	-	-	-	1	-	-	-	-
2002.....	3	-	-	-	-	-	1	1	1	-	-	-
2001.....	3	-	-	-	-	-	-	-	-	1	1	1
2000.....	7	-	-	-	-	1	1	-	2	1	-	-
1999.....	4	-	-	-	-	2	-	-	-	1	1	-
American Indian 1/, female												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	1	-	-	-	-	-	-	-	-	1	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	1	-	-	-	-	-	-	1	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Neglect, abandonment and other maltreatment syndromes (Y06-Y07)												
All races, both sexes												
2004.....	202	101	69	6	3	1	-	2	-	3	3	-
2003.....	218	100	84	6	3	-	-	1	-	1	1	-
2002.....	238	98	89	13	3	1	2	-	-	2	1	1
2001.....	242	116	79	8	3	4	2	1	2	1	1	2
2000.....	248	129	72	6	3	2	3	3	1	4	1	3
1999.....	227	125	64	11	-	1	2	1	5	4	2	-
All races, male												
2004.....	117	61	41	2	2	1	-	1	-	1	2	-
2003.....	127	67	41	5	2	-	-	1	-	1	1	-
2002.....	116	51	47	7	2	-	1	-	-	1	1	-
2001.....	133	66	51	4	1	4	2	-	1	-	-	1
2000.....	143	78	40	5	2	2	1	1	-	1	1	2
1999.....	122	71	33	5	-	-	1	-	5	3	1	-
All races, female												
2004.....	85	40	28	4	1	-	-	1	-	2	1	-
2003.....	91	33	43	1	1	-	-	-	-	-	-	-
2002.....	122	47	42	6	1	1	1	-	-	1	-	1
2001.....	109	50	28	4	2	-	-	1	1	1	1	1
2000.....	105	51	32	1	1	-	2	2	1	3	-	1
1999.....	105	54	31	6	-	1	1	1	-	1	1	-
White, both sexes												
2004.....	135	67	43	4	2	1	-	1	-	2	3	-
2003.....	137	66	48	4	2	-	-	1	-	-	1	-
2002.....	149	58	56	4	2	-	2	-	-	-	-	1
2001.....	146	69	45	3	2	3	1	1	2	-	-	1
2000.....	151	76	44	3	2	1	1	3	1	4	-	1
1999.....	127	76	29	6	-	-	2	-	4	1	2	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Neglect, abandonment and other maltreatment syndromes (Y06-Y07)												
White, male												
2004.....	77	40	23	2	1	1	-	1	-	1	2	-
2003.....	79	44	20	4	2	-	-	1	-	-	1	-
2002.....	75	33	31	2	2	-	1	-	-	-	-	-
2001.....	82	41	30	2	1	3	1	-	1	-	-	-
2000.....	93	53	25	3	2	1	-	1	-	1	-	-
1999.....	71	46	15	2	-	-	1	-	4	1	1	-
White, female												
2004.....	58	27	20	2	1	-	-	-	-	1	1	-
2003.....	58	22	28	-	-	-	-	-	-	-	-	-
2002.....	74	25	25	2	-	-	1	-	-	-	-	1
2001.....	64	28	15	1	1	-	-	1	1	-	-	1
2000.....	58	23	19	-	-	-	1	2	1	3	-	1
1999.....	56	30	14	4	-	-	1	-	-	-	1	-
Black, both sexes												
2004.....	58	31	23	2	1	-	-	-	-	1	-	-
2003.....	69	26	34	2	1	-	-	-	-	1	-	-
2002.....	81	38	30	8	1	1	-	-	-	2	1	-
2001.....	88	43	33	5	1	1	1	-	-	-	-	-
2000.....	85	49	27	1	1	1	1	-	-	-	1	-
1999.....	90	45	33	4	-	1	-	1	-	3	-	-
Black, male												
2004.....	36	19	16	-	1	-	-	-	-	-	-	-
2003.....	40	16	20	1	-	-	-	-	-	1	-	-
2002.....	40	18	15	5	-	-	-	-	-	1	1	-
2001.....	48	24	20	2	-	1	1	-	-	-	-	-
2000.....	45	24	15	-	-	1	1	-	-	-	1	-
1999.....	44	22	17	2	-	-	-	-	-	2	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Assault (homicide) by all other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5,*U01.8-*U01.9,*U02,X92,X96-X98,Y01-Y03,Y08-Y09,Y87.1)

All races, both sexes

2004.....	2,521	156	210	39	22	132	189	186	169	203	236	262
2003.....	2,570	175	210	35	28	112	215	169	168	236	261	256
2002.....	2,461	148	223	30	31	113	189	153	190	222	245	257
2001 3/.....	5,645	143	239	45	40	146	302	523	713	807	779	637
2000.....	2,845	163	203	54	48	145	250	221	236	312	290	246
1999.....	2,768	144	203	76	38	160	203	214	233	306	298	226

All races, male

2004.....	1,676	80	110	22	8	76	139	135	120	133	154	195
2003.....	1,731	97	120	24	15	80	143	115	102	150	191	179
2002.....	1,660	83	129	12	16	75	127	105	127	144	180	190
2001 3/.....	3,996	82	137	29	17	95	206	367	513	611	580	465
2000.....	1,873	91	117	27	20	89	172	153	153	189	211	174
1999.....	1,799	78	113	35	20	101	136	141	142	200	209	167

All races, female

2004.....	845	76	100	17	14	56	50	51	49	70	82	67
2003.....	839	78	90	11	13	32	72	54	66	86	70	77
2002.....	801	65	94	18	15	38	62	48	63	78	65	67
2001 3/.....	1,649	61	102	16	23	51	96	156	200	196	199	172
2000.....	972	72	86	27	28	56	78	68	83	123	79	72
1999.....	969	66	90	41	18	59	67	73	91	106	89	59

White, both sexes

2004.....	1,623	97	110	23	10	82	130	118	99	122	158	167
2003.....	1,654	108	109	14	19	76	137	102	98	151	167	160
2002.....	1,535	79	114	16	17	71	118	92	111	137	163	159
2001 3/.....	4,160	86	127	30	27	89	212	384	538	619	599	480
2000.....	1,669	88	100	30	33	73	136	123	130	178	172	146
1999.....	1,751	85	119	37	28	98	138	123	140	185	171	147

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Assault (homicide) by all other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5,*U01.8-*U01.9,*U02,X92,X96-X98,Y01-Y03,Y08-Y09,Y87.1)

White, male

2004.....	1,081	57	62	12	5	47	93	82	67	79	106	123
2003.....	1,090	61	55	10	9	53	91	64	64	96	119	110
2002.....	1,020	46	69	2	8	43	77	66	77	85	123	111
2001 3/.....	3,019	50	73	17	11	55	141	271	400	499	462	359
2000.....	1,070	53	62	19	15	37	91	80	81	110	126	95
1999.....	1,134	48	65	18	13	57	90	81	84	124	125	110

White, female

2004.....	542	40	48	11	5	35	37	36	32	43	52	44
2003.....	564	47	54	4	10	23	46	38	34	55	48	50
2002.....	515	33	45	14	9	28	41	26	34	52	40	48
2001 3/.....	1,141	36	54	13	16	34	71	113	138	120	137	121
2000.....	599	35	38	11	18	36	45	43	49	68	46	51
1999.....	617	37	54	19	15	41	48	42	56	61	46	37

Black, both sexes

2004.....	772	51	96	16	10	42	46	60	61	71	64	85
2003.....	798	58	91	20	9	30	64	57	60	72	82	85
2002.....	799	62	97	13	12	33	62	51	65	75	72	89
2001 3/.....	1,170	52	101	10	11	40	66	100	124	150	142	133
2000.....	1,058	69	95	24	14	67	110	90	92	117	102	88
1999.....	895	54	80	33	9	54	54	84	82	107	115	68

Black, male

2004.....	518	20	45	10	3	24	36	47	47	49	41	66
2003.....	555	29	58	14	6	22	42	42	33	47	64	60
2002.....	557	32	56	9	8	24	44	32	42	51	50	72
2001 3/.....	763	28	61	9	6	26	49	72	78	84	92	89
2000.....	724	35	49	8	4	47	78	66	64	73	72	71
1999.....	593	28	45	16	6	39	44	54	51	65	80	50

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Assault (homicide) by all other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5,*U01.8-*U01.9,*U02,X92,X96-X98,Y01-Y03,Y08-Y09,Y87.1)

Black, female

2004.....	254	31	51	6	7	18	10	13	14	22	23	19
2003.....	243	29	33	6	3	8	22	15	27	25	18	25
2002.....	242	30	41	4	4	9	18	19	23	24	22	17
2001 3/.....	407	24	40	1	5	14	17	28	46	66	50	44
2000.....	334	34	46	16	10	20	32	24	28	44	30	17
1999.....	302	26	35	17	3	15	10	30	31	42	35	18

American Indian 1/, both sexes

2004.....	60	7	2	-	-	3	7	3	7	4	9	4
2003.....	64	5	5	-	-	3	8	8	6	7	5	9
2002.....	74	5	8	-	1	3	6	7	8	6	8	5
2001 3/.....	64	1	7	1	1	10	8	7	5	5	4	3
2000.....	53	2	5	-	-	3	-	5	6	9	10	4
1999.....	64	1	-	3	-	4	7	5	7	9	8	2

American Indian 1/, male

2004.....	39	3	1	-	-	-	6	3	6	3	3	4
2003.....	49	3	4	-	-	3	5	7	4	5	4	7
2002.....	51	3	4	-	-	3	5	5	6	5	5	5
2001 3/.....	43	1	3	-	-	7	5	5	3	5	2	3
2000.....	41	2	4	-	-	3	-	4	4	5	8	2
1999.....	39	1	-	-	-	1	1	4	4	7	4	2

American Indian 1/, female

2004.....	21	4	1	-	-	3	1	-	1	1	6	-
2003.....	15	2	1	-	-	-	3	1	2	2	1	2
2002.....	23	2	4	-	1	-	1	2	2	1	3	-
2001 3/.....	21	-	4	1	1	3	3	2	2	-	2	-
2000.....	12	-	1	-	-	-	-	1	2	4	2	2
1999.....	25	-	-	3	-	3	6	1	3	2	4	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Assault (homicide) by all other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5,*U01.8-*U01.9,*U02,X92,X96-X98,Y01-Y03,Y08-Y09,Y87.1)

Asian or Pacific Islander 2/, both sexes

2004.....	66	1	2	-	2	5	6	5	2	6	5	6
2003.....	54	4	5	1	-	3	6	2	4	6	7	2
2002.....	53	2	4	1	1	6	3	3	6	4	2	4
2001 3/.....	251	4	4	4	1	7	16	32	46	33	34	21
2000.....	65	4	3	-	1	2	4	3	8	8	6	8
1999.....	58	4	4	3	1	4	4	2	4	5	4	9

Asian or Pacific Islander 2/, male

2004.....	38	-	2	-	-	5	4	3	-	2	4	2
2003.....	37	4	3	-	-	2	5	2	1	2	4	2
2002.....	32	2	-	1	-	5	1	2	2	3	2	2
2001 3/.....	171	3	-	3	-	7	11	19	32	23	24	14
2000.....	38	1	2	-	1	2	3	3	4	1	5	6
1999.....	33	1	3	1	1	4	1	2	3	4	-	5

Asian or Pacific Islander 2/, female

2004.....	28	1	-	-	2	-	2	2	2	4	1	4
2003.....	17	-	2	1	-	1	1	-	3	4	3	-
2002.....	21	-	4	-	1	1	2	1	4	1	-	2
2001 3/.....	80	1	4	1	1	-	5	13	14	10	10	7
2000.....	27	3	1	-	-	-	1	-	4	7	1	2
1999.....	25	3	1	2	-	-	3	-	1	1	4	4

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Event of undetermined intent (Y10-Y34,Y87.2,Y89.9)												
All races, both sexes												
2004.....	4,976	104	47	25	39	153	348	366	440	604	785	790
2003.....	5,072	75	31	16	39	169	394	351	484	686	895	777
2002.....	4,830	101	57	26	42	143	344	352	476	727	789	709
2001.....	4,198	83	44	17	34	132	280	311	417	674	722	619
2000.....	3,819	65	39	18	42	120	250	306	373	599	687	543
1999.....	3,917	55	43	12	37	159	271	313	430	644	651	528
All races, male												
2004.....	3,164	65	26	16	27	117	263	261	298	401	471	495
2003.....	3,295	50	17	9	32	124	288	249	331	450	563	493
2002.....	3,114	52	28	11	27	108	250	256	329	494	487	432
2001.....	2,833	49	19	10	28	102	214	231	284	438	493	433
2000.....	2,635	27	26	13	37	93	192	223	266	415	470	374
1999.....	2,692	33	21	6	32	132	213	230	293	446	441	357
All races, female												
2004.....	1,812	39	21	9	12	36	85	105	142	203	314	295
2003.....	1,777	25	14	7	7	45	106	102	153	236	332	284
2002.....	1,716	49	29	15	15	35	94	96	147	233	302	277
2001.....	1,365	34	25	7	6	30	66	80	133	236	229	186
2000.....	1,184	38	13	5	5	27	58	83	107	184	217	169
1999.....	1,225	22	22	6	5	27	58	83	137	198	210	171
White, both sexes												
2004.....	4,142	68	30	10	26	125	307	325	352	505	658	668
2003.....	4,255	52	21	7	30	149	339	300	419	578	755	654
2002.....	3,962	56	45	16	33	120	298	291	388	599	653	581
2001.....	3,453	45	31	9	30	116	235	259	353	548	598	508
2000.....	3,053	41	27	9	29	103	199	252	301	492	539	432
1999.....	3,122	36	23	10	32	125	217	255	351	516	507	409

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Event of undetermined intent (Y10-Y34,Y87.2,Y89.9)												
White, male												
2004.....	2,607	44	16	6	18	95	234	234	237	334	390	407
2003.....	2,709	30	11	3	26	108	243	209	280	380	468	405
2002.....	2,524	29	23	5	21	89	217	210	267	399	406	353
2001.....	2,283	28	13	4	25	89	174	190	241	347	398	345
2000.....	2,078	18	17	6	26	79	149	181	211	342	368	292
1999.....	2,105	22	13	5	28	106	172	180	236	349	349	262
White, female												
2004.....	1,535	24	14	4	8	30	73	91	115	171	268	261
2003.....	1,546	22	10	4	4	41	96	91	139	198	287	249
2002.....	1,438	27	22	11	12	31	81	81	121	200	247	228
2001.....	1,170	17	18	5	5	27	61	69	112	201	200	163
2000.....	975	23	10	3	3	24	50	71	90	150	171	140
1999.....	1,017	14	10	5	4	19	45	75	115	167	158	147
Black, both sexes												
2004.....	702	32	15	14	12	22	34	27	73	87	104	110
2003.....	687	21	9	7	7	16	35	38	58	92	124	115
2002.....	759	37	8	9	6	17	40	54	71	118	126	116
2001.....	646	36	13	7	4	12	32	39	54	112	115	101
2000.....	668	15	11	9	12	13	39	45	60	98	136	103
1999.....	696	18	16	2	5	25	46	51	70	114	126	109
Black, male												
2004.....	479	18	9	9	9	19	24	20	53	60	64	78
2003.....	498	19	5	4	4	13	30	32	45	59	85	83
2002.....	521	18	3	5	6	14	29	41	49	89	75	74
2001.....	480	20	6	5	3	10	28	31	36	81	89	80
2000.....	488	3	8	7	10	13	34	35	45	67	95	77
1999.....	523	10	8	1	4	20	36	44	49	87	84	88

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Event of undetermined intent (Y10-Y34,Y87.2,Y89.9)												
Black, female												
2004.....	223	14	6	5	3	3	10	7	20	27	40	32
2003.....	189	2	4	3	3	3	5	6	13	33	39	32
2002.....	238	19	5	4	-	3	11	13	22	29	51	42
2001.....	166	16	7	2	1	2	4	8	18	31	26	21
2000.....	180	12	3	2	2	-	5	10	15	31	41	26
1999.....	173	8	8	1	1	5	10	7	21	27	42	21
American Indian 1/, both sexes												
2004.....	68	3	-	1	1	3	3	8	8	11	12	3
2003.....	64	2	-	1	1	1	7	8	3	11	10	5
2002.....	48	3	2	1	1	4	1	2	10	3	5	7
2001.....	43	1	-	-	-	3	4	2	5	8	6	5
2000.....	46	4	1	-	1	1	4	3	6	6	6	4
1999.....	54	1	2	-	-	4	5	2	9	9	9	4
American Indian 1/, male												
2004.....	41	3	-	1	-	2	3	5	4	6	9	3
2003.....	47	1	-	1	1	-	7	4	3	9	7	3
2002.....	26	1	1	1	-	4	1	-	7	1	2	3
2001.....	32	-	-	-	-	3	4	2	5	6	3	3
2000.....	36	3	1	-	1	-	3	2	6	5	4	3
1999.....	37	1	-	-	-	3	4	2	8	6	4	3
American Indian 1/, female												
2004.....	27	-	-	-	1	1	-	3	4	5	3	-
2003.....	17	1	-	-	-	1	-	4	-	2	3	2
2002.....	22	2	1	-	1	-	-	2	3	2	3	4
2001.....	11	1	-	-	-	-	-	-	-	2	3	2
2000.....	10	1	-	-	-	1	1	1	-	1	2	1
1999.....	17	-	2	-	-	1	1	-	1	3	5	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Event of undetermined intent (Y10-Y34,Y87.2,Y89.9)												
Asian or Pacific Islander 2/, both sexes												
2004.....	64	1	2	-	-	3	4	6	7	1	11	9
2003.....	66	-	1	1	1	3	13	5	4	5	6	3
2002.....	61	5	2	-	2	2	5	5	7	7	5	5
2001.....	56	1	-	1	-	1	9	11	5	6	3	5
2000.....	52	5	-	-	-	3	8	6	6	3	6	4
1999.....	45	-	2	-	-	5	3	5	-	5	9	6
Asian or Pacific Islander 2/, male												
2004.....	37	-	1	-	-	1	2	2	4	1	8	7
2003.....	41	-	1	1	1	3	8	4	3	2	3	2
2002.....	43	4	1	-	-	1	3	5	6	5	4	2
2001.....	38	1	-	1	-	-	8	8	2	4	3	5
2000.....	33	3	-	-	-	1	6	5	4	1	3	2
1999.....	27	-	-	-	-	3	1	4	-	4	4	4
Asian or Pacific Islander 2/, female												
2004.....	27	1	1	-	-	2	2	4	3	-	3	2
2003.....	25	-	-	-	-	-	5	1	1	3	3	1
2002.....	18	1	1	-	2	1	2	-	1	2	1	3
2001.....	18	-	-	-	-	1	1	3	3	2	-	-
2000.....	19	2	-	-	-	2	2	1	2	2	3	2
1999.....	18	-	2	-	-	2	2	1	-	1	5	2

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the
International Classification of
Diseases, Tenth Revision, 1992),
race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	-------------	-----------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Poisoning by and exposure to drugs and biological substances, undetermined intent (Y10-Y14)

All races, both sexes

2004.....	3,306	8	7	3	8	81	230	264	319	479	609	615
2003.....	3,553	3	5	1	3	89	282	259	373	551	740	609
2002.....	3,197	5	6	3	9	74	209	259	348	550	618	550
2001.....	2,769	4	6	1	8	70	162	206	315	513	575	465
2000.....	2,429	5	4	5	5	55	134	195	255	466	544	417
1999.....	2,473	8	5	1	3	52	142	208	317	487	511	389

All races, male

2004.....	1,997	4	3	-	4	64	166	182	208	307	351	369
2003.....	2,232	3	2	1	2	63	207	174	248	355	448	380
2002.....	1,959	3	-	2	6	51	145	183	225	359	366	314
2001.....	1,798	3	4	-	8	51	119	144	203	322	376	324
2000.....	1,643	2	4	3	3	38	98	138	179	321	366	278
1999.....	1,617	5	3	1	1	38	102	147	212	328	336	248

All races, female

2004.....	1,309	4	4	3	4	17	64	82	111	172	258	246
2003.....	1,321	-	3	-	1	26	75	85	125	196	292	229
2002.....	1,238	2	6	1	3	23	64	76	123	191	252	236
2001.....	971	1	2	1	-	19	43	62	112	191	199	141
2000.....	786	3	-	2	2	17	36	57	76	145	178	139
1999.....	856	3	2	-	2	14	40	61	105	159	175	141

White, both sexes

2004.....	2,859	4	3	2	7	73	220	246	271	404	521	525
2003.....	3,074	2	1	1	2	82	266	239	332	474	631	512
2002.....	2,707	2	5	1	9	65	193	227	292	462	517	457
2001.....	2,340	3	3	-	7	66	151	181	266	426	476	387
2000.....	2,013	3	2	3	4	51	123	168	213	391	431	336
1999.....	2,017	3	2	-	2	49	128	172	274	394	403	305

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Poisoning by and exposure to drugs and biological substances, undetermined intent (Y10-Y14)

White, male

2004.....	1,713	2	-	-	4	57	160	173	177	258	296	308
2003.....	1,904	2	-	1	2	59	194	160	218	308	378	311
2002.....	1,649	2	-	-	6	45	135	161	190	297	311	263
2001.....	1,483	2	3	-	7	49	109	125	173	263	299	261
2000.....	1,353	1	2	2	2	35	89	118	148	272	290	223
1999.....	1,288	1	2	-	1	37	93	117	182	260	268	183

White, female

2004.....	1,146	2	3	2	3	16	60	73	94	146	225	217
2003.....	1,170	-	1	-	-	23	72	79	114	166	253	201
2002.....	1,058	-	5	1	3	20	58	66	102	165	206	194
2001.....	857	1	-	-	-	17	42	56	93	163	177	126
2000.....	660	2	-	1	2	16	34	50	65	119	141	113
1999.....	729	2	-	-	1	12	35	55	92	134	135	122

Black, both sexes

2004.....	391	4	4	1	1	6	6	12	44	69	77	81
2003.....	434	1	4	-	1	6	11	16	39	69	100	93
2002.....	461	3	1	1	-	7	14	29	47	85	99	88
2001.....	396	1	3	1	1	2	8	21	43	81	95	74
2000.....	383	1	1	2	-	3	7	25	35	72	107	77
1999.....	422	5	3	1	1	1	11	33	41	87	98	82

Black, male

2004.....	255	2	3	-	-	6	3	7	29	48	46	53
2003.....	302	1	2	-	-	3	10	13	28	42	65	67
2002.....	292	1	-	1	-	4	9	20	29	60	54	50
2001.....	296	1	1	-	1	1	8	16	26	55	76	60
2000.....	267	-	1	1	-	3	5	18	26	46	74	54
1999.....	308	4	1	1	-	-	7	27	29	65	62	64

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Poisoning by and exposure to drugs and biological substances, undetermined intent (Y10-Y14)

Black, female

2004.....	136	2	1	1	1	-	3	5	15	21	31	28
2003.....	132	-	2	-	1	3	1	3	11	27	35	26
2002.....	169	2	1	-	-	3	5	9	18	25	45	38
2001.....	100	-	2	1	-	1	-	5	17	26	19	14
2000.....	116	1	-	1	-	-	2	7	9	26	33	23
1999.....	114	1	2	-	1	1	4	6	12	22	36	18

American Indian 1/, both sexes

2004.....	31	-	-	-	-	1	2	2	3	6	7	3
2003.....	28	-	-	-	-	-	2	4	1	7	5	3
2002.....	17	-	-	1	-	2	-	1	6	-	2	4
2001.....	16	-	-	-	-	1	-	1	2	3	3	4
2000.....	19	1	1	-	1	-	2	1	3	2	3	2
1999.....	18	-	-	-	-	1	2	1	2	3	5	-

American Indian 1/, male

2004.....	17	-	-	-	-	1	2	1	2	1	6	3
2003.....	17	-	-	-	-	-	2	1	1	5	3	1
2002.....	10	-	-	1	-	2	-	-	4	-	1	1
2001.....	10	-	-	-	-	1	-	1	2	2	-	3
2000.....	16	1	1	-	1	-	2	1	3	2	1	1
1999.....	11	-	-	-	-	-	2	1	1	1	3	-

American Indian 1/, female

2004.....	14	-	-	-	-	-	-	1	1	5	1	-
2003.....	11	-	-	-	-	-	-	3	-	2	2	2
2002.....	7	-	-	-	-	-	-	1	2	-	1	3
2001.....	6	-	-	-	-	-	-	-	-	1	3	1
2000.....	3	-	-	-	-	-	-	-	-	-	2	1
1999.....	7	-	-	-	-	1	-	-	1	2	2	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Poisoning by and exposure to drugs and biological substances, undetermined intent (Y10-Y14)

Asian or Pacific Islander 2/, both sexes

2004.....	25	-	-	-	-	1	2	4	1	-	4	6
2003.....	17	-	-	-	-	1	3	-	1	1	4	1
2002.....	12	-	-	-	-	-	2	2	3	3	-	1
2001.....	17	-	-	-	-	1	3	3	4	3	1	-
2000.....	14	-	-	-	-	1	2	1	4	1	3	2
1999.....	16	-	-	-	-	1	1	2	-	3	5	2

Asian or Pacific Islander 2/, male

2004.....	12	-	-	-	-	-	1	1	-	-	3	5
2003.....	9	-	-	-	-	1	1	-	1	-	2	1
2002.....	8	-	-	-	-	-	1	2	2	2	-	-
2001.....	9	-	-	-	-	-	2	2	2	2	1	-
2000.....	7	-	-	-	-	-	2	1	2	1	1	-
1999.....	10	-	-	-	-	1	-	2	-	2	3	1

Asian or Pacific Islander 2/, female

2004.....	13	-	-	-	-	1	1	3	1	-	1	1
2003.....	8	-	-	-	-	-	2	-	-	1	2	-
2002.....	4	-	-	-	-	-	1	-	1	1	-	1
2001.....	8	-	-	-	-	1	1	1	2	1	-	-
2000.....	7	-	-	-	-	1	-	-	2	-	2	2
1999.....	6	-	-	-	-	-	1	-	-	1	2	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Poisoning by and exposure to other and unspecified solid or liquid substance, undetermined intent (Y15-Y16,Y18-Y19)

All races, both sexes

2004.....	63	-	-	-	-	1	5	2	5	7	11	10
2003.....	58	1	1	1	-	-	2	4	5	5	8	10
2002.....	68	-	-	-	1	1	5	7	6	9	5	15
2001.....	61	-	-	-	-	2	1	4	7	7	9	9
2000.....	53	-	-	-	1	2	2	2	6	10	8	6
1999.....	42	-	-	-	-	3	1	3	5	5	7	5

All races, male

2004.....	45	-	-	-	-	-	4	1	4	5	6	7
2003.....	38	1	1	-	-	-	-	2	4	4	7	5
2002.....	50	-	-	-	1	-	5	6	5	6	4	11
2001.....	47	-	-	-	-	1	-	4	6	6	7	8
2000.....	41	-	-	-	1	2	2	2	5	9	8	5
1999.....	24	-	-	-	-	2	1	2	2	4	5	3

All races, female

2004.....	18	-	-	-	-	1	1	1	1	2	5	3
2003.....	20	-	-	1	-	-	2	2	1	1	1	5
2002.....	18	-	-	-	-	1	-	1	1	3	1	4
2001.....	14	-	-	-	-	1	1	-	1	1	2	1
2000.....	12	-	-	-	-	-	-	-	1	1	-	1
1999.....	18	-	-	-	-	1	-	1	3	1	2	2

White, both sexes

2004.....	57	-	-	-	-	1	5	2	5	7	7	9
2003.....	51	1	1	1	-	-	1	4	5	5	6	9
2002.....	61	-	-	-	1	1	5	5	5	9	4	15
2001.....	52	-	-	-	-	2	1	2	6	5	9	7
2000.....	38	-	-	-	-	2	1	2	4	7	5	6
1999.....	36	-	-	-	-	2	-	3	5	5	6	4

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Poisoning by and exposure to other and unspecified solid or liquid substance, undetermined intent (Y15-Y16,Y18-Y19)

White, male

2004.....	39	-	-	-	-	-	4	1	4	5	2	6
2003.....	32	1	1	-	-	-	-	2	4	4	5	4
2002.....	45	-	-	-	1	-	5	4	4	6	4	11
2001.....	39	-	-	-	-	1	-	2	5	4	7	6
2000.....	29	-	-	-	-	2	1	2	4	6	5	5
1999.....	21	-	-	-	-	2	-	2	2	4	5	2

White, female

2004.....	18	-	-	-	-	1	1	1	1	2	5	3
2003.....	19	-	-	1	-	-	1	2	1	1	1	5
2002.....	16	-	-	-	-	1	-	1	1	3	-	4
2001.....	13	-	-	-	-	1	1	-	1	1	2	1
2000.....	9	-	-	-	-	-	-	-	-	1	-	1
1999.....	15	-	-	-	-	-	-	1	3	1	1	2

Black, both sexes

2004.....	4	-	-	-	-	-	-	-	-	-	2	1
2003.....	6	-	-	-	-	-	1	-	-	-	2	1
2002.....	4	-	-	-	-	-	-	2	1	-	-	-
2001.....	9	-	-	-	-	-	-	2	1	2	-	2
2000.....	9	-	-	-	1	-	1	-	1	2	2	-
1999.....	4	-	-	-	-	-	1	-	-	-	1	1

Black, male

2004.....	4	-	-	-	-	-	-	-	-	-	2	1
2003.....	5	-	-	-	-	-	-	-	-	-	2	1
2002.....	4	-	-	-	-	-	-	2	1	-	-	-
2001.....	8	-	-	-	-	-	-	2	1	2	-	2
2000.....	7	-	-	-	1	-	1	-	-	2	2	-
1999.....	3	-	-	-	-	-	1	-	-	-	-	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Poisoning by and exposure to other and unspecified solid or liquid substance, undetermined intent (Y15-Y16,Y18-Y19)

Asian or Pacific Islander 2/, both sexes

2004.....	2	-	-	-	-	-	-	-	-	-	2	-
2003.....	1	-	-	-	-	-	-	-	-	-	-	-
2002.....	2	-	-	-	-	-	-	-	-	-	1	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	2	-	-	-	-	-	-	-	-	-	-	-
1999.....	1	-	-	-	-	1	-	-	-	-	-	-

Asian or Pacific Islander 2/, male

2004.....	2	-	-	-	-	-	-	-	-	-	2	-
2003.....	1	-	-	-	-	-	-	-	-	-	-	-
2002.....	1	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	2	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-

Asian or Pacific Islander 2/, female

2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	1	-	-	-	-	-	-	-	-	-	1	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	1	-	-	-	-	1	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Poisoning by and exposure to other gases and vapors, undetermined intent (Y17)

All races, both sexes

2004.....	86	1	1	3	4	7	5	2	10	8	2	12
2003.....	89	3	-	-	-	5	8	6	9	8	8	10
2002.....	71	1	-	-	-	2	4	4	7	5	16	8
2001.....	79	3	1	-	1	1	5	5	12	10	7	12
2000.....	75	-	-	2	1	3	4	9	8	7	13	6
1999.....	80	1	2	-	-	1	5	4	6	9	12	6

All races, male

2004.....	68	1	1	3	2	5	4	2	8	7	1	9
2003.....	61	1	-	-	-	2	4	5	7	7	6	8
2002.....	54	-	-	-	-	2	3	3	5	4	14	8
2001.....	60	2	1	-	-	1	4	3	11	6	7	9
2000.....	55	-	-	2	1	2	3	9	5	5	9	3
1999.....	56	-	-	-	-	1	5	4	2	6	11	3

All races, female

2004.....	18	-	-	-	2	2	1	-	2	1	1	3
2003.....	28	2	-	-	-	3	4	1	2	1	2	2
2002.....	17	1	-	-	-	-	1	1	2	1	2	-
2001.....	19	1	-	-	1	-	1	2	1	4	-	3
2000.....	20	-	-	-	-	1	1	-	3	2	4	3
1999.....	24	1	2	-	-	-	-	-	4	3	1	3

White, both sexes

2004.....	69	-	-	1	1	3	4	2	9	8	2	10
2003.....	76	2	-	-	-	5	8	6	5	8	7	9
2002.....	63	1	-	-	-	2	3	3	5	5	15	7
2001.....	72	2	1	-	1	1	5	5	12	8	7	10
2000.....	64	-	-	1	1	2	2	9	7	7	11	4
1999.....	69	1	-	-	-	1	4	3	5	8	10	6

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Poisoning by and exposure to other gases and vapors, undetermined intent (Y17)												
White, male												
2004.....	55	-	-	1	-	3	3	2	7	7	1	8
2003.....	50	-	-	-	-	2	4	5	4	7	5	7
2002.....	51	-	-	-	-	2	3	2	5	4	13	7
2001.....	55	1	1	-	-	1	4	3	11	4	7	8
2000.....	48	-	-	1	1	2	1	9	5	5	8	2
1999.....	48	-	-	-	-	1	4	3	1	5	10	3
White, female												
2004.....	14	-	-	-	1	-	1	-	2	1	1	2
2003.....	26	2	-	-	-	3	4	1	1	1	2	2
2002.....	12	1	-	-	-	-	-	1	-	1	2	-
2001.....	17	1	-	-	1	-	1	2	1	4	-	2
2000.....	16	-	-	-	-	-	1	-	2	2	3	2
1999.....	21	1	-	-	-	-	-	-	4	3	-	3
Black, both sexes												
2004.....	17	1	1	2	3	4	1	-	1	-	-	2
2003.....	8	1	-	-	-	-	-	-	2	-	-	1
2002.....	8	-	-	-	-	-	1	1	2	-	1	1
2001.....	5	1	-	-	-	-	-	-	-	1	-	2
2000.....	8	-	-	1	-	-	1	-	1	-	2	2
1999.....	10	-	1	-	-	-	1	1	1	1	2	-
Black, male												
2004.....	13	1	1	2	2	2	1	-	1	-	-	1
2003.....	6	1	-	-	-	-	-	-	1	-	-	1
2002.....	3	-	-	-	-	-	-	1	-	-	1	1
2001.....	3	1	-	-	-	-	-	-	-	1	-	1
2000.....	5	-	-	1	-	-	1	-	-	-	1	1
1999.....	8	-	-	-	-	-	1	1	1	1	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Discharge of firearms, undetermined intent (Y22-Y24)

All races, both sexes

2004.....	235	-	1	3	6	22	36	22	23	16	28	12
2003.....	232	-	1	1	11	31	38	21	17	21	22	19
2002.....	243	-	1	2	4	35	44	27	19	31	19	19
2001.....	231	-	-	2	4	24	40	32	16	33	17	16
2000.....	230	-	-	2	8	26	45	32	20	18	21	9
1999.....	324	-	3	-	12	68	59	32	19	28	29	18

All races, male

2004.....	179	-	-	2	5	18	32	19	14	11	19	9
2003.....	186	-	1	1	11	29	31	18	15	12	14	11
2002.....	199	-	1	2	3	34	37	23	16	22	17	14
2001.....	190	-	-	2	4	23	36	27	15	20	13	12
2000.....	190	-	-	2	7	25	37	26	17	12	18	6
1999.....	277	-	3	-	12	61	50	28	16	24	21	15

All races, female

2004.....	56	-	1	1	1	4	4	3	9	5	9	3
2003.....	46	-	-	-	-	2	7	3	2	9	8	8
2002.....	44	-	-	-	1	1	7	4	3	9	2	5
2001.....	41	-	-	-	-	1	4	5	1	13	4	4
2000.....	40	-	-	-	1	1	8	6	3	6	3	3
1999.....	47	-	-	-	-	7	9	4	3	4	8	3

White, both sexes

2004.....	187	-	-	1	3	15	27	16	15	15	24	11
2003.....	183	-	-	-	7	23	24	14	13	18	20	18
2002.....	192	-	-	-	4	24	35	20	13	25	16	18
2001.....	188	-	-	-	4	17	27	26	14	31	14	14
2000.....	167	-	-	-	5	20	28	23	11	13	18	8
1999.....	247	-	1	-	11	42	43	22	14	25	25	13

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Discharge of firearms, undetermined intent (Y22-Y24)

White, male

2004.....	142	-	-	1	3	12	25	13	9	10	16	8
2003.....	144	-	-	-	7	21	19	12	11	10	12	11
2002.....	153	-	-	-	3	23	28	16	11	18	14	13
2001.....	150	-	-	-	4	16	24	22	13	18	11	10
2000.....	137	-	-	-	5	19	23	18	10	8	17	5
1999.....	212	-	1	-	11	39	37	19	12	21	20	10

White, female

2004.....	45	-	-	-	-	3	2	3	6	5	8	3
2003.....	39	-	-	-	-	2	5	2	2	8	8	7
2002.....	39	-	-	-	1	1	7	4	2	7	2	5
2001.....	38	-	-	-	-	1	3	4	1	13	3	4
2000.....	30	-	-	-	-	1	5	5	1	5	1	3
1999.....	35	-	-	-	-	3	6	3	2	4	5	3

Black, both sexes

2004.....	39	-	1	2	2	7	8	4	5	1	3	-
2003.....	35	-	1	1	3	7	10	4	3	2	1	1
2002.....	41	-	1	2	-	9	8	7	4	4	2	-
2001.....	32	-	-	2	-	7	8	4	1	-	2	2
2000.....	54	-	-	2	3	6	14	8	7	4	3	1
1999.....	66	-	2	-	1	20	15	10	4	3	3	4

Black, male

2004.....	31	-	-	1	2	6	6	4	4	1	2	-
2003.....	32	-	1	1	3	7	9	4	3	1	1	-
2002.....	39	-	1	2	-	9	8	7	4	3	2	-
2001.....	29	-	-	2	-	7	7	3	1	-	1	2
2000.....	47	-	-	2	2	6	13	7	5	4	1	1
1999.....	56	-	2	-	1	17	12	9	3	3	1	4

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Discharge of firearms, undetermined intent (Y22-Y24)												
Asian or Pacific Islander 2/, both sexes												
2004.....	2	-	-	-	-	-	-	-	1	-	-	1
2003.....	5	-	-	-	-	1	2	-	-	-	-	-
2002.....	3	-	-	-	-	-	-	-	-	2	-	1
2001.....	7	-	-	-	-	-	3	2	-	1	1	-
2000.....	5	-	-	-	-	-	1	1	1	1	-	-
1999.....	5	-	-	-	-	3	-	-	-	-	-	1
Asian or Pacific Islander 2/, male												
2004.....	2	-	-	-	-	-	-	-	1	-	-	1
2003.....	2	-	-	-	-	1	1	-	-	-	-	-
2002.....	2	-	-	-	-	-	-	-	-	1	-	1
2001.....	7	-	-	-	-	-	3	2	-	1	1	-
2000.....	3	-	-	-	-	-	-	1	1	-	-	-
1999.....	4	-	-	-	-	2	-	-	-	-	-	1
Asian or Pacific Islander 2/, female												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	3	-	-	-	-	-	1	-	-	-	-	-
2002.....	1	-	-	-	-	-	-	-	-	1	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	2	-	-	-	-	-	1	-	-	1	-	-
1999.....	1	-	-	-	-	1	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Falling, jumping or pushed from a high place, undetermined intent (Y30)												
All races, both sexes												
2004.....	71	1	2	-	-	4	8	7	5	6	9	9
2003.....	77	-	2	-	1	2	11	3	9	12	5	8
2002.....	103	12	1	-	1	4	7	6	13	12	11	8
2001.....	77	6	1	-	-	5	6	7	8	9	10	6
2000.....	55	1	2	-	-	3	2	7	1	9	6	4
1999.....	59	-	2	-	1	4	4	5	8	11	5	10
All races, male												
2004.....	50	-	1	-	-	2	6	7	3	6	7	6
2003.....	54	-	2	-	1	1	8	2	6	10	5	6
2002.....	73	8	1	-	-	3	6	5	11	9	8	4
2001.....	54	3	-	-	-	3	6	7	4	7	8	4
2000.....	42	1	2	-	-	2	2	6	1	7	4	4
1999.....	46	-	1	-	1	4	3	4	7	9	5	7
All races, female												
2004.....	21	1	1	-	-	2	2	-	2	-	2	3
2003.....	23	-	-	-	-	1	3	1	3	2	-	2
2002.....	30	4	-	-	1	1	1	1	2	3	3	4
2001.....	23	3	1	-	-	2	-	-	4	2	2	2
2000.....	13	-	-	-	-	1	-	1	-	2	2	-
1999.....	13	-	1	-	-	-	1	1	1	2	-	3
White, both sexes												
2004.....	58	-	1	-	-	2	6	5	4	5	8	8
2003.....	61	-	2	-	1	2	9	-	9	10	2	7
2002.....	77	3	1	-	-	4	6	4	10	10	9	6
2001.....	57	4	-	-	-	4	6	4	6	7	8	3
2000.....	43	1	2	-	-	2	1	6	1	5	5	4
1999.....	47	-	-	-	1	3	4	4	4	9	5	9

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Falling, jumping or pushed from a high place, undetermined intent (Y30)												
White, male												
2004.....	41	-	-	-	-	1	5	5	2	5	6	6
2003.....	44	-	2	-	1	1	7	-	6	9	2	5
2002.....	55	2	1	-	-	3	6	3	8	8	6	3
2001.....	41	3	-	-	-	3	6	4	3	6	6	2
2000.....	31	1	2	-	-	1	1	5	1	4	3	4
1999.....	37	-	-	-	1	3	3	3	4	7	5	7
White, female												
2004.....	17	-	1	-	-	1	1	-	2	-	2	2
2003.....	17	-	-	-	-	1	2	-	3	1	-	2
2002.....	22	1	-	-	-	1	-	1	2	2	3	3
2001.....	16	1	-	-	-	1	-	-	3	1	2	1
2000.....	12	-	-	-	-	1	-	1	-	1	2	-
1999.....	10	-	-	-	-	-	1	1	-	2	-	2
Black, both sexes												
2004.....	6	1	-	-	-	1	2	1	-	-	1	-
2003.....	9	-	-	-	-	-	1	3	-	1	1	-
2002.....	16	7	-	-	-	-	1	2	1	1	1	1
2001.....	10	2	1	-	-	1	-	1	1	1	1	-
2000.....	10	-	-	-	-	1	-	-	-	4	1	-
1999.....	10	-	2	-	-	1	-	1	3	1	-	1
Black, male												
2004.....	4	-	-	-	-	1	1	1	-	-	1	-
2003.....	6	-	-	-	-	-	1	2	-	-	1	-
2002.....	12	5	-	-	-	-	-	2	1	1	1	1
2001.....	6	-	-	-	-	-	-	1	1	1	1	-
2000.....	9	-	-	-	-	1	-	-	-	3	1	-
1999.....	7	-	1	-	-	1	-	1	2	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

All other and unspecified events of undetermined intent and their sequelae (Y20-Y21,Y25-Y29,Y31-Y34,Y87.2,Y89.9)

All races, both sexes

2004.....	1,215	94	36	16	21	38	64	69	78	88	126	132
2003.....	1,063	68	22	13	24	42	53	58	71	89	112	121
2002.....	1,148	83	49	21	27	27	75	49	83	120	120	109
2001.....	981	70	36	14	21	30	66	57	59	102	104	111
2000.....	977	59	33	9	27	31	63	61	83	89	95	101
1999.....	939	46	31	11	21	31	60	61	75	104	87	100

All races, male

2004.....	825	60	21	11	16	28	51	50	61	65	87	95
2003.....	724	45	11	7	18	29	38	48	51	62	83	83
2002.....	779	41	26	7	17	18	54	36	67	94	78	81
2001.....	684	41	14	8	16	23	49	46	45	77	82	76
2000.....	664	24	20	6	25	24	50	42	59	61	65	78
1999.....	672	28	14	5	18	26	52	45	54	75	63	81

All races, female

2004.....	390	34	15	5	5	10	13	19	17	23	39	37
2003.....	339	23	11	6	6	13	15	10	20	27	29	38
2002.....	369	42	23	14	10	9	21	13	16	26	42	28
2001.....	297	29	22	6	5	7	17	11	14	25	22	35
2000.....	313	35	13	3	2	7	13	19	24	28	30	23
1999.....	267	18	17	6	3	5	8	16	21	29	24	19

White, both sexes

2004.....	912	64	26	6	15	31	45	54	48	66	96	105
2003.....	810	47	17	5	20	37	31	37	55	63	89	99
2002.....	862	50	39	15	19	24	56	32	63	88	92	78
2001.....	744	36	27	9	18	26	45	41	49	71	84	87
2000.....	728	37	23	5	19	26	44	44	65	69	69	74
1999.....	706	32	20	10	18	28	38	51	49	75	58	72

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

All other and unspecified events of undetermined intent and their sequelae (Y20-Y21,Y25-Y29,Y31-Y34,Y87.2,Y89.9)

White, male

2004.....	617	42	16	4	11	22	37	40	38	49	69	71
2003.....	535	27	8	2	16	25	19	30	37	42	66	67
2002.....	571	25	22	5	11	16	40	24	49	66	58	56
2001.....	515	22	9	4	14	19	31	34	36	52	68	58
2000.....	480	16	13	3	18	20	34	29	43	47	45	53
1999.....	499	21	10	5	15	24	35	36	35	52	41	57

White, female

2004.....	295	22	10	2	4	9	8	14	10	17	27	34
2003.....	275	20	9	3	4	12	12	7	18	21	23	32
2002.....	291	25	17	10	8	8	16	8	14	22	34	22
2001.....	229	14	18	5	4	7	14	7	13	19	16	29
2000.....	248	21	10	2	1	6	10	15	22	22	24	21
1999.....	207	11	10	5	3	4	3	15	14	23	17	15

Black, both sexes

2004.....	245	26	9	9	6	4	17	10	23	17	21	26
2003.....	195	19	4	6	3	3	12	15	14	20	20	19
2002.....	229	27	6	6	6	1	16	13	16	28	23	26
2001.....	194	32	9	4	3	2	16	11	8	27	17	21
2000.....	204	14	10	4	8	3	16	12	16	16	21	23
1999.....	184	13	8	1	3	3	18	6	21	22	22	21

Black, male

2004.....	172	15	5	6	5	4	13	8	19	11	13	23
2003.....	147	17	2	3	1	3	10	13	13	16	16	14
2002.....	171	12	2	2	6	1	12	9	14	25	17	22
2001.....	138	18	5	3	2	2	13	9	7	22	11	15
2000.....	153	3	7	3	7	3	14	10	14	12	16	21
1999.....	141	6	4	-	3	2	15	6	14	17	20	19

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

All other and unspecified events of undetermined intent and their sequelae (Y20-Y21,Y25-Y29,Y31-Y34,Y87.2,Y89.9)

Black, female

2004.....	73	11	4	3	1	-	4	2	4	6	8	3
2003.....	48	2	2	3	2	-	2	2	1	4	4	5
2002.....	58	15	4	4	-	-	4	4	2	3	6	4
2001.....	56	14	4	1	1	-	3	2	1	5	6	6
2000.....	51	11	3	1	1	-	2	2	2	4	5	2
1999.....	43	7	4	1	-	1	3	-	7	5	2	2

American Indian 1/, both sexes

2004.....	27	3	-	1	-	2	-	3	3	4	4	-
2003.....	22	2	-	1	-	1	3	1	-	3	2	2
2002.....	22	3	2	-	1	-	-	1	2	2	2	3
2001.....	18	1	-	-	-	2	2	1	2	3	2	-
2000.....	19	3	-	-	-	1	-	2	1	3	2	2
1999.....	26	1	1	-	-	-	2	1	5	5	3	4

American Indian 1/, male

2004.....	17	3	-	1	-	1	-	1	2	4	2	-
2003.....	17	1	-	1	-	-	3	1	-	3	1	2
2002.....	11	1	1	-	-	-	-	-	2	1	-	2
2001.....	15	-	-	-	-	2	2	1	2	3	2	-
2000.....	14	2	-	-	-	-	-	1	1	2	2	2
1999.....	19	1	-	-	-	-	1	1	5	4	1	3

American Indian 1/, female

2004.....	10	-	-	-	-	1	-	2	1	-	2	-
2003.....	5	1	-	-	-	1	-	-	-	-	1	-
2002.....	11	2	1	-	1	-	-	1	-	1	2	1
2001.....	3	1	-	-	-	-	-	-	-	-	-	-
2000.....	5	1	-	-	-	1	-	1	-	1	-	-
1999.....	7	-	1	-	-	-	1	-	-	1	2	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and unspecified events of undetermined intent and their sequelae (Y20-Y21,Y25-Y29,Y31-Y34,Y87.2,Y89.9)												
Asian or Pacific Islander 2/, both sexes												
2004.....	31	1	1	-	-	1	2	2	4	1	5	1
2003.....	36	-	1	1	1	1	7	5	2	3	1	1
2002.....	35	3	2	-	1	2	3	3	2	2	3	2
2001.....	25	1	-	1	-	-	3	4	-	1	1	3
2000.....	26	5	-	-	-	1	3	3	1	1	3	2
1999.....	23	-	2	-	-	-	2	3	-	2	4	3
Asian or Pacific Islander 2/, male												
2004.....	19	-	-	-	-	1	1	1	2	1	3	1
2003.....	25	-	1	1	1	1	6	4	1	1	-	-
2002.....	26	3	1	-	-	1	2	3	2	2	3	1
2001.....	16	1	-	1	-	-	3	2	-	-	1	3
2000.....	17	3	-	-	-	1	2	2	1	-	2	2
1999.....	13	-	-	-	-	-	1	2	-	2	1	2
Asian or Pacific Islander 2/, female												
2004.....	12	1	1	-	-	-	1	1	2	-	2	-
2003.....	11	-	-	-	-	-	1	1	1	2	1	1
2002.....	9	-	1	-	1	1	1	-	-	-	-	1
2001.....	9	-	-	-	-	-	-	2	-	1	-	-
2000.....	9	2	-	-	-	-	1	1	-	1	1	-
1999.....	10	-	2	-	-	-	1	1	-	-	3	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Legal intervention (Y35,Y89.0)												
All races, both sexes												
2004.....	372	-	-	-	-	27	53	50	56	57	47	31
2003.....	423	-	-	-	2	20	73	63	72	66	51	29
2002.....	384	-	-	-	3	24	67	51	51	64	45	43
2001.....	396	-	-	-	-	26	62	47	60	65	50	33
2000.....	359	-	-	-	3	27	46	59	51	60	41	38
1999.....	398	-	-	-	1	20	47	66	70	72	49	19
All races, male												
2004.....	363	-	-	-	-	26	53	48	56	57	46	28
2003.....	405	-	-	-	1	19	70	61	70	65	49	27
2002.....	369	-	-	-	3	23	64	51	48	60	45	40
2001.....	380	-	-	-	-	25	59	46	58	63	45	33
2000.....	348	-	-	-	3	27	44	58	50	58	38	37
1999.....	392	-	-	-	1	20	47	66	69	69	48	19
All races, female												
2004.....	9	-	-	-	-	1	-	2	-	-	1	3
2003.....	18	-	-	-	1	1	3	2	2	1	2	2
2002.....	15	-	-	-	-	1	3	-	3	4	-	3
2001.....	16	-	-	-	-	1	3	1	2	2	5	-
2000.....	11	-	-	-	-	-	2	1	1	2	3	1
1999.....	6	-	-	-	-	-	-	-	1	3	1	-
White, both sexes												
2004.....	262	-	-	-	-	22	33	27	39	44	37	24
2003.....	284	-	-	-	1	11	43	30	46	52	38	22
2002.....	257	-	-	-	1	14	35	28	41	45	33	28
2001.....	292	-	-	-	-	15	45	35	43	46	40	23
2000.....	237	-	-	-	1	11	33	36	36	38	25	32
1999.....	255	-	-	-	-	9	22	39	47	47	33	15

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Legal intervention (Y35,Y89.0)

White, male

2004.....	256	-	-	-	-	21	33	26	39	44	36	22
2003.....	271	-	-	-	-	10	42	30	44	51	36	20
2002.....	246	-	-	-	1	13	33	28	39	43	33	25
2001.....	283	-	-	-	-	14	44	34	43	44	37	23
2000.....	231	-	-	-	1	11	31	35	35	37	25	32
1999.....	250	-	-	-	-	9	22	39	47	44	32	15

White, female

2004.....	6	-	-	-	-	1	-	1	-	-	1	2
2003.....	13	-	-	-	1	1	1	-	2	1	2	2
2002.....	11	-	-	-	-	1	2	-	2	2	-	3
2001.....	9	-	-	-	-	1	1	1	-	2	3	-
2000.....	6	-	-	-	-	-	2	1	1	1	-	-
1999.....	5	-	-	-	-	-	-	-	-	3	1	-

Black, both sexes

2004.....	92	-	-	-	-	2	18	21	13	10	9	5
2003.....	120	-	-	-	1	8	27	29	24	12	10	6
2002.....	110	-	-	-	2	10	27	20	9	17	11	11
2001.....	84	-	-	-	-	10	16	9	12	15	8	9
2000.....	111	-	-	-	2	15	13	19	14	20	15	6
1999.....	126	-	-	-	1	10	22	24	18	21	15	4

Black, male

2004.....	90	-	-	-	-	2	18	21	13	10	9	4
2003.....	117	-	-	-	1	8	25	28	24	12	10	6
2002.....	106	-	-	-	2	10	26	20	8	15	11	11
2001.....	79	-	-	-	-	10	14	9	11	15	6	9
2000.....	108	-	-	-	2	15	13	19	14	20	13	5
1999.....	125	-	-	-	1	10	22	24	17	21	15	4

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Legal execution (Y35.5)												
All races, both sexes												
2004.....	52	-	-	-	-	-	-	4	12	10	8	7
2003.....	60	-	-	-	-	-	-	3	16	10	11	9
2002.....	67	-	-	-	-	-	1	6	7	16	13	9
2001.....	63	-	-	-	-	-	-	3	8	14	14	9
2000.....	80	-	-	-	-	-	1	9	18	18	11	11
1999.....	88	-	-	-	-	-	1	11	8	25	18	5
All races, male												
2004.....	52	-	-	-	-	-	-	4	12	10	8	7
2003.....	60	-	-	-	-	-	-	3	16	10	11	9
2002.....	65	-	-	-	-	-	1	6	7	16	13	8
2001.....	60	-	-	-	-	-	-	3	8	14	12	9
2000.....	78	-	-	-	-	-	1	8	18	18	11	11
1999.....	88	-	-	-	-	-	1	11	8	25	18	5
All races, female												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	2	-	-	-	-	-	-	-	-	-	-	1
2001.....	3	-	-	-	-	-	-	-	-	-	2	-
2000.....	2	-	-	-	-	-	-	1	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes												
2004.....	33	-	-	-	-	-	-	1	7	7	6	6
2003.....	41	-	-	-	-	-	-	1	9	7	9	6
2002.....	49	-	-	-	-	-	-	2	6	12	9	6
2001.....	43	-	-	-	-	-	-	3	5	9	10	3
2000.....	47	-	-	-	-	-	1	4	9	10	5	9
1999.....	54	-	-	-	-	-	1	4	7	14	10	4

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Legal execution (Y35.5)												
White, male												
2004.....	33	-	-	-	-	-	-	1	7	7	6	6
2003.....	41	-	-	-	-	-	-	1	9	7	9	6
2002.....	47	-	-	-	-	-	-	2	6	12	9	5
2001.....	42	-	-	-	-	-	-	3	5	9	9	3
2000.....	45	-	-	-	-	-	1	3	9	10	5	9
1999.....	54	-	-	-	-	-	1	4	7	14	10	4
White, female												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	2	-	-	-	-	-	-	-	-	-	-	1
2001.....	1	-	-	-	-	-	-	-	-	-	1	-
2000.....	2	-	-	-	-	-	-	1	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes												
2004.....	18	-	-	-	-	-	-	3	5	2	2	1
2003.....	18	-	-	-	-	-	-	2	7	3	2	3
2002.....	18	-	-	-	-	-	1	4	1	4	4	3
2001.....	17	-	-	-	-	-	-	-	3	4	3	6
2000.....	32	-	-	-	-	-	-	5	9	8	6	2
1999.....	30	-	-	-	-	-	-	6	1	9	7	1
Black, male												
2004.....	18	-	-	-	-	-	-	3	5	2	2	1
2003.....	18	-	-	-	-	-	-	2	7	3	2	3
2002.....	18	-	-	-	-	-	1	4	1	4	4	3
2001.....	16	-	-	-	-	-	-	-	3	4	2	6
2000.....	32	-	-	-	-	-	-	5	9	8	6	2
1999.....	30	-	-	-	-	-	-	6	1	9	7	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other legal intervention and their sequelae (Y35.0-Y35.4,Y35.6-Y35.7,Y89.0)

All races, both sexes

2004.....	320	-	-	-	-	27	53	46	44	47	39	24
2003.....	363	-	-	-	2	20	73	60	56	56	40	20
2002.....	317	-	-	-	3	24	66	45	44	48	32	34
2001.....	333	-	-	-	-	26	62	44	52	51	36	24
2000.....	279	-	-	-	3	27	45	50	33	42	30	27
1999.....	310	-	-	-	1	20	46	55	62	47	31	14

All races, male

2004.....	311	-	-	-	-	26	53	44	44	47	38	21
2003.....	345	-	-	-	1	19	70	58	54	55	38	18
2002.....	304	-	-	-	3	23	63	45	41	44	32	32
2001.....	320	-	-	-	-	25	59	43	50	49	33	24
2000.....	270	-	-	-	3	27	43	50	32	40	27	26
1999.....	304	-	-	-	1	20	46	55	61	44	30	14

All races, female

2004.....	9	-	-	-	-	1	-	2	-	-	1	3
2003.....	18	-	-	-	1	1	3	2	2	1	2	2
2002.....	13	-	-	-	-	1	3	-	3	4	-	2
2001.....	13	-	-	-	-	1	3	1	2	2	3	-
2000.....	9	-	-	-	-	-	2	-	1	2	3	1
1999.....	6	-	-	-	-	-	-	-	1	3	1	-

White, both sexes

2004.....	229	-	-	-	-	22	33	26	32	37	31	18
2003.....	243	-	-	-	1	11	43	29	37	45	29	16
2002.....	208	-	-	-	1	14	35	26	35	33	24	22
2001.....	249	-	-	-	-	15	45	32	38	37	30	20
2000.....	190	-	-	-	1	11	32	32	27	28	20	23
1999.....	201	-	-	-	-	9	21	35	40	33	23	11

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other legal intervention and their sequelae (Y35.0-Y35.4,Y35.6-Y35.7,Y89.0)												
White, male												
2004.....	223	-	-	-	-	21	33	25	32	37	30	16
2003.....	230	-	-	-	-	10	42	29	35	44	27	14
2002.....	199	-	-	-	1	13	33	26	33	31	24	20
2001.....	241	-	-	-	-	14	44	31	38	35	28	20
2000.....	186	-	-	-	1	11	30	32	26	27	20	23
1999.....	196	-	-	-	-	9	21	35	40	30	22	11
White, female												
2004.....	6	-	-	-	-	1	-	1	-	-	1	2
2003.....	13	-	-	-	1	1	1	-	2	1	2	2
2002.....	9	-	-	-	-	1	2	-	2	2	-	2
2001.....	8	-	-	-	-	1	1	1	-	2	2	-
2000.....	4	-	-	-	-	-	2	-	1	1	-	-
1999.....	5	-	-	-	-	-	-	-	-	3	1	-
Black, both sexes												
2004.....	74	-	-	-	-	2	18	18	8	8	7	4
2003.....	102	-	-	-	1	8	27	27	17	9	8	3
2002.....	92	-	-	-	2	10	26	16	8	13	7	8
2001.....	67	-	-	-	-	10	16	9	9	11	5	3
2000.....	79	-	-	-	2	15	13	14	5	12	9	4
1999.....	96	-	-	-	1	10	22	18	17	12	8	3
Black, male												
2004.....	72	-	-	-	-	2	18	18	8	8	7	3
2003.....	99	-	-	-	1	8	25	26	17	9	8	3
2002.....	88	-	-	-	2	10	25	16	7	11	7	8
2001.....	63	-	-	-	-	10	14	9	8	11	4	3
2000.....	76	-	-	-	2	15	13	14	5	12	7	3
1999.....	95	-	-	-	1	10	22	18	16	12	8	3

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Operations of war and their sequelae (Y36,Y89.1)												
All races, both sexes												
2004.....	28	-	-	-	-	-	4	1	-	3	-	-
2003.....	14	-	-	-	-	-	1	-	-	-	-	-
2002.....	20	-	-	-	-	-	-	-	1	-	1	1
2001.....	17	-	-	-	-	-	1	-	-	1	-	1
2000.....	16	-	-	-	-	-	-	-	-	-	-	-
1999.....	23	-	-	-	-	-	-	-	-	1	-	-
All races, male												
2004.....	28	-	-	-	-	-	4	1	-	3	-	-
2003.....	14	-	-	-	-	-	1	-	-	-	-	-
2002.....	20	-	-	-	-	-	-	-	1	-	1	1
2001.....	16	-	-	-	-	-	1	-	-	1	-	1
2000.....	16	-	-	-	-	-	-	-	-	-	-	-
1999.....	22	-	-	-	-	-	-	-	-	1	-	-
All races, female												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	1	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes												
2004.....	26	-	-	-	-	-	3	1	-	2	-	-
2003.....	13	-	-	-	-	-	1	-	-	-	-	-
2002.....	15	-	-	-	-	-	-	-	1	-	-	-
2001.....	16	-	-	-	-	-	1	-	-	1	-	1
2000.....	15	-	-	-	-	-	-	-	-	-	-	-
1999.....	22	-	-	-	-	-	-	-	-	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Complications of medical and surgical care (Y40-Y84,Y88)

All races, both sexes

2004.....	2,883	16	19	8	11	22	26	30	41	56	101	134
2003.....	2,855	15	24	6	6	21	23	26	43	75	104	145
2002.....	2,843	15	19	13	22	16	19	27	52	73	98	114
2001.....	3,021	26	24	5	9	18	25	24	55	71	101	142
2000.....	3,059	19	22	18	13	10	22	37	48	65	84	128
1999.....	2,823	54	30	7	11	17	19	29	49	66	86	91

All races, male

2004.....	1,364	6	10	5	7	12	12	16	15	26	42	67
2003.....	1,333	8	16	2	2	18	8	14	22	36	60	80
2002.....	1,236	8	11	8	18	10	9	16	31	35	47	60
2001.....	1,357	15	14	2	6	6	13	11	31	39	44	75
2000.....	1,430	12	11	11	4	5	11	23	24	33	43	59
1999.....	1,308	31	12	4	5	8	13	9	34	37	41	37

All races, female

2004.....	1,519	10	9	3	4	10	14	14	26	30	59	67
2003.....	1,522	7	8	4	4	3	15	12	21	39	44	65
2002.....	1,607	7	8	5	4	6	10	11	21	38	51	54
2001.....	1,664	11	10	3	3	12	12	13	24	32	57	67
2000.....	1,629	7	11	7	9	5	11	14	24	32	41	69
1999.....	1,515	23	18	3	6	9	6	20	15	29	45	54

White, both sexes

2004.....	2,392	11	16	7	9	19	14	26	31	39	74	102
2003.....	2,356	9	13	5	3	18	19	16	32	57	84	103
2002.....	2,331	8	14	7	14	10	14	20	37	51	68	80
2001.....	2,446	17	16	3	7	13	20	16	41	47	70	100
2000.....	2,520	14	12	12	11	8	16	22	38	52	63	97
1999.....	2,314	35	21	6	8	15	15	16	39	52	62	68

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Complications of medical and surgical care (Y40-Y84,Y88)

White, male

2004.....	1,162	4	8	5	5	10	6	15	11	22	32	53
2003.....	1,121	5	9	2	1	15	7	8	17	29	52	56
2002.....	1,025	5	9	4	12	7	7	12	23	28	37	45
2001.....	1,084	9	10	1	4	5	11	7	22	24	30	53
2000.....	1,210	11	8	7	4	4	9	13	20	27	37	45
1999.....	1,078	22	11	3	4	7	10	4	28	30	30	26

White, female

2004.....	1,230	7	8	2	4	9	8	11	20	17	42	49
2003.....	1,235	4	4	3	2	3	12	8	15	28	32	47
2002.....	1,306	3	5	3	2	3	7	8	14	23	31	35
2001.....	1,362	8	6	2	3	8	9	9	19	23	40	47
2000.....	1,310	3	4	5	7	4	7	9	18	25	26	52
1999.....	1,236	13	10	3	4	8	5	12	11	22	32	42

Black, both sexes

2004.....	437	2	3	1	2	3	10	4	6	16	27	28
2003.....	432	5	10	-	3	2	4	7	9	17	17	34
2002.....	448	7	5	6	8	5	2	6	11	20	29	31
2001.....	522	9	7	1	2	3	4	8	14	22	28	39
2000.....	487	5	8	4	1	2	5	13	10	13	18	27
1999.....	449	18	9	1	3	2	4	10	8	13	22	21

Black, male

2004.....	173	1	2	-	2	2	4	1	2	4	10	11
2003.....	177	3	6	-	1	2	1	3	5	6	6	19
2002.....	184	3	2	4	6	3	1	4	5	6	10	13
2001.....	250	6	4	-	2	-	2	4	9	14	13	20
2000.....	193	1	2	2	-	1	1	9	4	6	6	12
1999.....	197	8	1	1	1	1	3	4	6	6	10	11

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Complications of medical and surgical care (Y40-Y84,Y88)												
Black, female												
2004.....	264	1	1	1	-	1	6	3	4	12	17	17
2003.....	255	2	4	-	2	-	3	4	4	11	11	15
2002.....	264	4	3	2	2	2	1	2	6	14	19	18
2001.....	272	3	3	1	-	3	2	4	5	8	15	19
2000.....	294	4	6	2	1	1	4	4	6	7	12	15
1999.....	252	10	8	-	2	1	1	6	2	7	12	10
American Indian 1/, both sexes												
2004.....	11	2	-	-	-	-	-	-	-	-	-	2
2003.....	24	1	-	-	-	1	-	2	1	1	-	4
2002.....	26	-	-	-	-	-	1	-	2	1	1	2
2001.....	17	-	1	1	-	1	1	-	-	-	2	-
2000.....	14	-	-	-	1	-	-	-	-	-	1	-
1999.....	20	-	-	-	-	-	-	2	1	1	-	-
American Indian 1/, male												
2004.....	6	1	-	-	-	-	-	-	-	-	-	2
2003.....	10	-	-	-	-	1	-	2	-	1	-	3
2002.....	12	-	-	-	-	-	1	-	2	-	-	1
2001.....	6	-	-	1	-	-	-	-	-	-	1	-
2000.....	5	-	-	-	-	-	-	-	-	-	-	-
1999.....	13	-	-	-	-	-	-	1	-	1	-	-
American Indian 1/, female												
2004.....	5	1	-	-	-	-	-	-	-	-	-	-
2003.....	14	1	-	-	-	-	-	-	1	-	-	1
2002.....	14	-	-	-	-	-	-	-	-	1	1	1
2001.....	11	-	1	-	-	1	1	-	-	-	1	-
2000.....	9	-	-	-	1	-	-	-	-	-	1	-
1999.....	7	-	-	-	-	-	-	1	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Complications of medical and surgical care (Y40-Y84,Y88)												
Asian or Pacific Islander 2/, both sexes												
2004.....	43	1	-	-	-	-	2	-	4	1	-	2
2003.....	43	-	1	1	-	-	-	1	1	-	3	4
2002.....	38	-	-	-	-	1	2	1	2	1	-	1
2001.....	36	-	-	-	-	1	-	-	-	2	1	3
2000.....	38	-	2	2	-	-	1	2	-	-	2	4
1999.....	40	1	-	-	-	-	-	1	1	-	2	2
Asian or Pacific Islander 2/, male												
2004.....	23	-	-	-	-	-	2	-	2	-	-	1
2003.....	25	-	1	-	-	-	-	1	-	-	2	2
2002.....	15	-	-	-	-	-	-	-	1	1	-	1
2001.....	17	-	-	-	-	1	-	-	-	1	-	2
2000.....	22	-	1	2	-	-	1	1	-	-	-	2
1999.....	20	1	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, female												
2004.....	20	1	-	-	-	-	-	-	2	1	-	1
2003.....	18	-	-	1	-	-	-	-	1	-	1	2
2002.....	23	-	-	-	-	1	2	1	1	-	-	-
2001.....	19	-	-	-	-	-	-	-	-	1	1	1
2000.....	16	-	1	-	-	-	-	1	-	-	2	2
1999.....	20	-	-	-	-	-	-	1	1	-	1	2

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Drugs, medicaments and biological substances causing adverse effects in therapeutic use, and their sequelae (Y40-Y59,Y88.0)

All races, both sexes

2004.....	370	3	3	1	2	3	14	11	8	8	29	35
2003.....	320	2	4	-	-	5	10	9	15	19	23	26
2002.....	247	1	1	2	7	3	7	10	12	13	14	16
2001.....	277	2	1	-	-	4	7	9	17	19	21	24
2000.....	255	-	-	1	1	1	6	8	11	16	13	18
1999.....	232	3	6	-	3	5	4	9	13	20	14	16

All races, male

2004.....	168	-	1	1	1	-	6	7	3	5	11	18
2003.....	167	1	1	-	-	5	5	7	10	11	16	17
2002.....	118	1	1	2	4	3	3	5	10	10	8	10
2001.....	135	2	1	-	-	2	5	7	13	15	14	10
2000.....	129	-	-	1	-	-	4	6	8	11	6	8
1999.....	117	1	2	-	1	3	3	2	12	13	7	10

All races, female

2004.....	202	3	2	-	1	3	8	4	5	3	18	17
2003.....	153	1	3	-	-	-	5	2	5	8	7	9
2002.....	129	-	-	-	3	-	4	5	2	3	6	6
2001.....	142	-	-	-	-	2	2	2	4	4	7	14
2000.....	126	-	-	-	1	1	2	2	3	5	7	10
1999.....	115	2	4	-	2	2	1	7	1	7	7	6

White, both sexes

2004.....	302	2	3	1	1	3	7	10	6	7	22	25
2003.....	276	2	2	-	-	5	9	8	14	15	19	22
2002.....	202	-	1	2	5	2	5	7	8	10	12	11
2001.....	220	1	-	-	-	3	6	5	14	16	18	10
2000.....	211	-	-	1	1	1	5	4	8	14	12	14
1999.....	189	3	5	-	3	4	4	4	10	17	10	11

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Drugs, medicaments and biological substances causing adverse effects in therapeutic use, and their sequelae (Y40-Y59,Y88.0)

White, male

2004.....	140	-	1	1	-	-	3	7	3	5	9	11
2003.....	143	1	-	-	-	5	5	6	9	10	14	15
2002.....	98	-	1	2	3	2	3	2	6	9	6	8
2001.....	109	1	-	-	-	2	4	4	11	13	11	3
2000.....	112	-	-	1	-	-	3	3	7	10	6	7
1999.....	93	1	2	-	1	2	3	-	9	11	5	6

White, female

2004.....	162	2	2	-	1	3	4	3	3	2	13	14
2003.....	133	1	2	-	-	-	4	2	5	5	5	7
2002.....	104	-	-	-	2	-	2	5	2	1	6	3
2001.....	111	-	-	-	-	1	2	1	3	3	7	7
2000.....	99	-	-	-	1	1	2	1	1	4	6	7
1999.....	96	2	3	-	2	2	1	4	1	6	5	5

Black, both sexes

2004.....	60	-	-	-	1	-	6	1	2	1	7	8
2003.....	37	-	2	-	-	-	1	1	1	4	3	3
2002.....	38	1	-	-	2	1	1	3	2	3	2	4
2001.....	52	1	1	-	-	1	1	4	3	3	3	12
2000.....	40	-	-	-	-	-	1	3	3	2	1	4
1999.....	37	-	1	-	-	1	-	4	3	3	4	4

Black, male

2004.....	23	-	-	-	1	-	2	-	-	-	2	5
2003.....	20	-	1	-	-	-	-	1	1	1	1	1
2002.....	16	1	-	-	1	1	-	3	2	1	2	1
2001.....	25	1	1	-	-	-	1	3	2	2	3	6
2000.....	14	-	-	-	-	-	1	2	1	1	-	1
1999.....	21	-	-	-	-	1	-	2	3	2	2	4

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Drugs, medicaments and biological substances causing adverse effects in therapeutic use, and their sequelae (Y40-Y59,Y88.0)

Asian or Pacific Islander 2/, both sexes

2004.....	7	1	-	-	-	-	1	-	-	-	-	1
2003.....	6	-	-	-	-	-	-	-	-	-	1	1
2002.....	4	-	-	-	-	-	1	-	1	-	-	-
2001.....	5	-	-	-	-	-	-	-	-	-	-	2
2000.....	3	-	-	-	-	-	-	1	-	-	-	-
1999.....	5	-	-	-	-	-	-	1	-	-	-	1

Asian or Pacific Islander 2/, male

2004.....	4	-	-	-	-	-	1	-	-	-	-	1
2003.....	4	-	-	-	-	-	-	-	-	-	1	1
2002.....	2	-	-	-	-	-	-	-	1	-	-	-
2001.....	1	-	-	-	-	-	-	-	-	-	-	1
2000.....	3	-	-	-	-	-	-	1	-	-	-	-
1999.....	2	-	-	-	-	-	-	-	-	-	-	-

Asian or Pacific Islander 2/, female

2004.....	3	1	-	-	-	-	-	-	-	-	-	-
2003.....	2	-	-	-	-	-	-	-	-	-	-	-
2002.....	2	-	-	-	-	-	1	-	-	-	-	-
2001.....	4	-	-	-	-	-	-	-	-	-	-	1
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	3	-	-	-	-	-	-	1	-	-	-	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Misadventures to patients during medical and surgical care and their sequelae (Y60-Y69,Y88.1)

All races, both sexes

2004.....	171	1	2	-	-	3	-	2	6	4	6	11
2003.....	166	4	-	-	-	4	1	4	4	6	7	12
2002.....	169	1	6	2	1	2	-	1	5	5	7	7
2001.....	177	6	5	-	2	-	-	1	4	5	3	9
2000.....	183	1	4	1	-	1	-	4	4	5	6	10
1999.....	179	6	3	-	-	1	1	1	3	8	7	8

All races, male

2004.....	65	-	-	-	-	2	-	1	3	1	1	4
2003.....	65	3	-	-	-	3	-	2	-	4	3	1
2002.....	69	-	5	1	1	1	-	-	2	1	4	3
2001.....	78	4	4	-	1	-	-	-	1	1	2	4
2000.....	82	1	2	-	-	-	-	1	3	4	2	3
1999.....	74	3	-	-	-	1	1	-	2	3	4	1

All races, female

2004.....	106	1	2	-	-	1	-	1	3	3	5	7
2003.....	101	1	-	-	-	1	1	2	4	2	4	11
2002.....	100	1	1	1	-	1	-	1	3	4	3	4
2001.....	99	2	1	-	1	-	-	1	3	4	1	5
2000.....	101	-	2	1	-	1	-	3	1	1	4	7
1999.....	105	3	3	-	-	-	-	1	1	5	3	7

White, both sexes

2004.....	142	1	2	-	-	3	-	2	5	2	4	9
2003.....	124	2	-	-	-	3	1	1	-	5	6	7
2002.....	128	1	5	1	1	-	-	1	2	3	5	3
2001.....	139	5	5	-	2	-	-	-	2	3	1	6
2000.....	139	1	1	1	-	1	-	2	2	5	4	6
1999.....	140	2	2	-	-	1	-	1	2	6	6	5

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Misadventures to patients during medical and surgical care and their sequelae (Y60-Y69,Y88.1)

White, male

2004.....	47	-	-	-	-	2	-	1	2	-	-	4
2003.....	51	2	-	-	-	2	-	1	-	4	2	-
2002.....	54	-	4	-	1	-	-	-	1	1	3	2
2001.....	62	3	4	-	1	-	-	-	1	-	1	2
2000.....	66	1	1	-	-	-	-	-	2	4	2	3
1999.....	60	1	-	-	-	1	-	-	2	2	4	1

White, female

2004.....	95	1	2	-	-	1	-	1	3	2	4	5
2003.....	73	-	-	-	-	1	1	-	-	1	4	7
2002.....	74	1	1	1	-	-	-	1	1	2	2	1
2001.....	77	2	1	-	1	-	-	-	1	3	-	4
2000.....	73	-	-	1	-	1	-	2	-	1	2	3
1999.....	80	1	2	-	-	-	-	1	-	4	2	4

Black, both sexes

2004.....	24	-	-	-	-	-	-	-	1	2	2	2
2003.....	37	2	-	-	-	-	-	3	3	1	1	4
2002.....	38	-	1	1	-	2	-	-	2	2	2	4
2001.....	34	1	-	-	-	-	-	1	2	2	1	3
2000.....	40	-	3	-	-	-	-	2	2	-	2	3
1999.....	35	4	1	-	-	-	1	-	-	2	1	3

Black, male

2004.....	13	-	-	-	-	-	-	-	1	1	1	-
2003.....	12	1	-	-	-	-	-	1	-	-	1	1
2002.....	14	-	1	1	-	1	-	-	1	-	1	1
2001.....	15	1	-	-	-	-	-	-	-	1	1	2
2000.....	14	-	1	-	-	-	-	1	1	-	-	-
1999.....	11	2	-	-	-	-	1	-	-	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Misadventures to patients during medical and surgical care and their sequelae (Y60-Y69,Y88.1)												
Black, female												
2004.....	11	-	-	-	-	-	-	-	-	1	1	2
2003.....	25	1	-	-	-	-	-	2	3	1	-	3
2002.....	24	-	-	-	-	1	-	-	1	2	1	3
2001.....	19	-	-	-	-	-	-	1	2	1	-	1
2000.....	26	-	2	-	-	-	-	1	1	-	2	3
1999.....	24	2	1	-	-	-	-	-	-	1	1	3
American Indian 1/, both sexes												
2004.....	1	-	-	-	-	-	-	-	-	-	-	-
2003.....	1	-	-	-	-	1	-	-	-	-	-	-
2002.....	1	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	1	-	-	-	-	-	-	-	-	-	-	-
1999.....	2	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, male												
2004.....	1	-	-	-	-	-	-	-	-	-	-	-
2003.....	1	-	-	-	-	1	-	-	-	-	-	-
2002.....	1	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	1	-	-	-	-	-	-	-	-	-	-	-
1999.....	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, female												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	1	-	-	-	-	-	-	-	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other complications of medical and surgical care and their sequelae (Y70-Y84,Y88.2-Y88.3)

All races, both sexes

2004.....	2,342	12	14	7	9	16	12	17	27	44	66	88
2003.....	2,369	9	20	6	6	12	12	13	24	50	74	107
2002.....	2,427	13	12	9	14	11	12	16	35	55	77	91
2001.....	2,567	18	18	5	7	14	18	14	34	47	77	109
2000.....	2,621	18	18	16	12	8	16	25	33	44	65	100
1999.....	2,412	45	21	7	8	11	14	19	33	38	65	67

All races, male

2004.....	1,131	6	9	4	6	10	6	8	9	20	30	45
2003.....	1,101	4	15	2	2	10	3	5	12	21	41	62
2002.....	1,049	7	5	5	13	6	6	11	19	24	35	47
2001.....	1,144	9	9	2	5	4	8	4	17	23	28	61
2000.....	1,219	11	9	10	4	5	7	16	13	18	35	48
1999.....	1,117	27	10	4	4	4	9	7	20	21	30	26

All races, female

2004.....	1,211	6	5	3	3	6	6	9	18	24	36	43
2003.....	1,268	5	5	4	4	2	9	8	12	29	33	45
2002.....	1,378	6	7	4	1	5	6	5	16	31	42	44
2001.....	1,423	9	9	3	2	10	10	10	17	24	49	48
2000.....	1,402	7	9	6	8	3	9	9	20	26	30	52
1999.....	1,295	18	11	3	4	7	5	12	13	17	35	41

White, both sexes

2004.....	1,948	8	11	6	8	13	7	14	20	30	48	68
2003.....	1,956	5	11	5	3	10	9	7	18	37	59	74
2002.....	2,001	7	8	4	8	8	9	12	27	38	51	66
2001.....	2,087	11	11	3	5	10	14	11	25	28	51	84
2000.....	2,170	13	11	10	10	6	11	16	28	33	47	77
1999.....	1,985	30	14	6	5	10	11	11	27	29	46	52

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other complications of medical and surgical care and their sequelae (Y70-Y84,Y88.2-Y88.3)

White, male

2004.....	975	4	7	4	5	8	3	7	6	17	23	38
2003.....	927	2	9	2	1	8	2	1	8	15	36	41
2002.....	873	5	4	2	8	5	4	10	16	18	28	35
2001.....	913	5	6	1	3	3	7	3	10	11	18	48
2000.....	1,032	10	7	6	4	4	6	10	11	13	29	35
1999.....	925	20	9	3	3	4	7	4	17	17	21	19

White, female

2004.....	973	4	4	2	3	5	4	7	14	13	25	30
2003.....	1,029	3	2	3	2	2	7	6	10	22	23	33
2002.....	1,128	2	4	2	-	3	5	2	11	20	23	31
2001.....	1,174	6	5	2	2	7	7	8	15	17	33	36
2000.....	1,138	3	4	4	6	2	5	6	17	20	18	42
1999.....	1,060	10	5	3	2	6	4	7	10	12	25	33

Black, both sexes

2004.....	353	2	3	1	1	3	4	3	3	13	18	18
2003.....	358	3	8	-	3	2	3	3	5	12	13	27
2002.....	372	6	4	5	6	2	1	3	7	15	25	23
2001.....	436	7	6	1	2	2	3	3	9	17	24	24
2000.....	407	5	5	4	1	2	4	8	5	11	15	20
1999.....	377	14	7	1	3	1	3	6	5	8	17	14

Black, male

2004.....	137	1	2	-	1	2	2	1	1	3	7	6
2003.....	145	2	5	-	1	2	1	1	4	5	4	17
2002.....	154	2	1	3	5	1	1	1	2	5	7	11
2001.....	210	4	3	-	2	-	1	1	7	11	9	12
2000.....	165	1	1	2	-	1	-	6	2	5	6	11
1999.....	165	6	1	1	1	-	2	2	3	3	8	7

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other complications of medical and surgical care and their sequelae (Y70-Y84,Y88.2-Y88.3)

Black, female

2004.....	216	1	1	1	-	1	2	2	2	10	11	12
2003.....	213	1	3	-	2	-	2	2	1	7	9	10
2002.....	218	4	3	2	1	1	-	2	5	10	18	12
2001.....	226	3	3	1	-	2	2	2	2	6	15	12
2000.....	242	4	4	2	1	1	4	2	3	6	9	9
1999.....	212	8	6	-	2	1	1	4	2	5	9	7

American Indian 1/, both sexes

2004.....	9	2	-	-	-	-	-	-	-	-	-	1
2003.....	22	1	-	-	-	-	-	2	1	1	-	4
2002.....	22	-	-	-	-	-	1	-	1	1	1	1
2001.....	17	-	1	1	-	1	1	-	-	-	2	-
2000.....	12	-	-	-	1	-	-	-	-	-	1	-
1999.....	17	-	-	-	-	-	-	2	-	1	-	-

American Indian 1/, male

2004.....	4	1	-	-	-	-	-	-	-	-	-	1
2003.....	9	-	-	-	-	-	-	2	-	1	-	3
2002.....	9	-	-	-	-	-	1	-	1	-	-	-
2001.....	6	-	-	1	-	-	-	-	-	-	1	-
2000.....	4	-	-	-	-	-	-	-	-	-	-	-
1999.....	11	-	-	-	-	-	-	1	-	1	-	-

American Indian 1/, female

2004.....	5	1	-	-	-	-	-	-	-	-	-	-
2003.....	13	1	-	-	-	-	-	-	1	-	-	1
2002.....	13	-	-	-	-	-	-	-	-	1	1	1
2001.....	11	-	1	-	-	1	1	-	-	-	1	-
2000.....	8	-	-	-	1	-	-	-	-	-	1	-
1999.....	6	-	-	-	-	-	-	1	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other complications of medical and surgical care and their sequelae (Y70-Y84,Y88.2-Y88.3)												
Asian or Pacific Islander 2/, both sexes												
2004.....	32	-	-	-	-	-	1	-	4	1	-	1
2003.....	33	-	1	1	-	-	-	1	-	-	2	2
2002.....	32	-	-	-	-	1	1	1	-	1	-	1
2001.....	27	-	-	-	-	1	-	-	-	2	-	1
2000.....	32	-	2	2	-	-	1	1	-	-	2	3
1999.....	33	1	-	-	-	-	-	-	1	-	2	1
Asian or Pacific Islander 2/, male												
2004.....	15	-	-	-	-	-	1	-	2	-	-	-
2003.....	20	-	1	-	-	-	-	1	-	-	1	1
2002.....	13	-	-	-	-	-	-	-	-	1	-	1
2001.....	15	-	-	-	-	1	-	-	-	1	-	1
2000.....	18	-	1	2	-	-	1	-	-	-	-	2
1999.....	16	1	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, female												
2004.....	17	-	-	-	-	-	-	-	2	1	-	1
2003.....	13	-	-	1	-	-	-	-	-	-	1	1
2002.....	19	-	-	-	-	1	1	1	-	-	-	-
2001.....	12	-	-	-	-	-	-	-	-	1	-	-
2000.....	14	-	1	-	-	-	-	1	-	-	2	1
1999.....	17	-	-	-	-	-	-	-	1	-	1	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All causes												
All races, both sexes												
2004.....	99,770	121,163	143,534	171,984	227,682	310,746	373,484	338,140	227,661	87,446	18,526	346
2003.....	98,608	119,339	143,180	174,817	238,680	325,185	377,839	346,847	231,403	90,271	19,331	342
2002.....	96,320	115,441	137,901	175,591	247,399	330,140	377,514	345,385	228,587	88,282	18,822	357
2001.....	94,381	109,523	134,616	178,232	252,728	333,677	368,252	339,775	221,314	85,954	18,488	422
2000.....	89,509	106,751	134,095	181,739	259,470	337,700	362,745	335,813	219,283	84,641	18,434	356
1999.....	84,854	104,884	134,095	186,695	265,905	338,349	360,241	330,903	213,991	83,334	17,913	356
All races, male												
2004.....	62,649	73,400	84,632	98,455	124,436	160,308	173,361	133,667	70,168	19,820	2,995	274
2003.....	61,543	72,188	84,273	100,315	131,106	167,354	174,978	135,177	70,162	19,872	3,001	271
2002.....	60,021	69,830	81,533	100,590	136,431	169,625	173,879	134,637	68,817	19,522	2,930	282
2001.....	58,281	65,801	79,157	102,110	139,471	171,311	169,431	130,405	65,527	18,732	2,869	331
2000.....	55,277	64,425	78,896	103,935	143,473	173,327	166,892	128,877	64,439	18,552	2,874	289
1999.....	52,356	63,441	79,283	107,529	147,391	174,853	166,117	126,531	62,511	18,111	2,836	277
All races, female												
2004.....	37,121	47,763	58,902	73,529	103,246	150,438	200,123	204,473	157,493	67,626	15,531	72
2003.....	37,065	47,151	58,907	74,502	107,574	157,831	202,861	211,670	161,241	70,399	16,330	71
2002.....	36,299	45,611	56,368	75,001	110,968	160,515	203,635	210,748	159,770	68,760	15,892	75
2001.....	36,100	43,722	55,459	76,122	113,257	162,366	198,821	209,370	155,787	67,222	15,619	91
2000.....	34,232	42,326	55,199	77,804	115,997	164,373	195,853	206,936	154,844	66,089	15,560	67
1999.....	32,498	41,443	54,812	79,166	118,514	163,496	194,124	204,372	151,480	65,223	15,077	79
White, both sexes												
2004.....	76,538	96,434	117,737	143,232	195,030	273,291	336,369	309,335	208,415	79,849	16,141	269
2003.....	75,377	95,720	117,101	145,645	204,904	286,819	340,633	317,904	212,499	82,525	16,767	292
2002.....	74,126	92,712	112,702	146,491	213,556	291,430	340,923	316,444	210,030	80,543	16,438	286
2001.....	73,065	87,807	110,046	148,608	218,520	295,048	332,704	311,165	203,570	78,462	16,068	320
2000.....	69,543	85,840	109,701	152,597	224,389	299,257	328,472	307,495	201,942	77,167	16,157	275
1999.....	66,116	83,876	109,706	157,260	230,177	299,701	326,788	302,550	197,437	76,142	15,820	271

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All causes												
White, male												
2004.....	48,805	58,771	69,804	82,606	107,517	142,168	157,174	122,938	64,196	17,919	2,536	218
2003.....	47,689	58,348	69,404	84,313	113,540	148,708	158,342	124,593	64,481	17,945	2,556	228
2002.....	46,757	56,535	67,016	84,741	118,994	150,862	157,792	124,020	63,038	17,634	2,452	230
2001.....	45,629	53,120	65,233	86,088	121,705	152,517	153,761	119,811	60,027	16,916	2,425	255
2000.....	43,267	52,048	65,066	88,182	125,197	154,451	151,919	118,321	59,009	16,624	2,455	231
1999.....	41,130	51,133	65,404	91,520	128,922	155,859	151,591	115,599	57,211	16,329	2,471	213
White, female												
2004.....	27,733	37,663	47,933	60,626	87,513	131,123	179,195	186,397	144,219	61,930	13,605	51
2003.....	27,688	37,372	47,697	61,332	91,364	138,111	182,291	193,311	148,018	64,580	14,211	64
2002.....	27,369	36,177	45,686	61,750	94,562	140,568	183,131	192,424	146,992	62,909	13,986	56
2001.....	27,436	34,687	44,813	62,520	96,815	142,531	178,943	191,354	143,543	61,546	13,643	65
2000.....	26,276	33,792	44,635	64,415	99,192	144,806	176,553	189,174	142,933	60,543	13,702	44
1999.....	24,986	32,743	44,302	65,740	101,255	143,842	175,197	186,951	140,226	59,813	13,349	58
Black, both sexes												
2004.....	20,287	21,336	22,154	24,310	27,372	31,111	30,268	23,184	15,748	6,320	2,048	64
2003.....	20,326	20,458	22,365	24,747	28,381	31,859	30,513	23,582	15,606	6,517	2,203	40
2002.....	19,439	19,693	21,643	24,783	28,531	32,431	30,431	23,920	15,458	6,578	2,066	57
2001.....	18,651	18,852	21,124	25,367	29,012	32,590	29,643	23,803	14,999	6,466	2,119	88
2000.....	17,554	18,161	21,120	25,064	30,131	32,541	28,903	23,919	14,680	6,407	2,032	72
1999.....	16,455	18,359	21,117	25,461	30,661	32,927	28,274	24,046	14,101	6,281	1,857	77
Black, male												
2004.....	12,116	12,646	12,706	13,384	14,171	14,830	12,865	8,253	4,520	1,412	363	46
2003.....	12,162	12,040	12,730	13,544	14,747	15,288	13,312	8,197	4,311	1,445	341	35
2002.....	11,621	11,550	12,450	13,465	14,733	15,434	13,044	8,292	4,455	1,454	383	40
2001.....	11,099	10,984	11,921	13,763	15,012	15,616	12,779	8,389	4,292	1,428	380	67
2000.....	10,563	10,787	11,891	13,505	15,566	15,682	12,344	8,428	4,249	1,532	351	50
1999.....	9,906	10,751	12,018	13,862	15,732	15,873	11,966	8,828	4,167	1,462	311	58

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All causes												
Black, female												
2004.....	8,171	8,690	9,448	10,926	13,201	16,281	17,403	14,931	11,228	4,908	1,685	18
2003.....	8,164	8,418	9,635	11,203	13,634	16,571	17,201	15,385	11,295	5,072	1,862	5
2002.....	7,818	8,143	9,193	11,318	13,798	16,997	17,387	15,628	11,003	5,124	1,683	17
2001.....	7,552	7,868	9,203	11,604	14,000	16,974	16,864	15,414	10,707	5,038	1,739	21
2000.....	6,991	7,374	9,229	11,559	14,565	16,859	16,559	15,491	10,431	4,875	1,681	22
1999.....	6,549	7,608	9,099	11,599	14,929	17,054	16,308	15,218	9,934	4,819	1,546	19
American Indian 1/, both sexes												
2004.....	906	1,006	1,067	1,177	1,152	1,204	1,171	809	453	187	54	10
2003.....	970	951	1,091	1,111	1,145	1,233	1,130	837	503	175	79	5
2002.....	851	894	992	1,095	1,196	1,160	1,055	746	470	182	53	7
2001.....	812	821	947	1,060	1,146	1,148	998	782	407	167	46	6
2000.....	743	813	885	1,049	1,092	1,130	932	728	442	169	43	3
1999.....	700	783	891	1,037	1,127	1,151	913	733	390	157	59	1
American Indian 1/, male												
2004.....	537	597	615	647	578	632	514	313	148	49	14	8
2003.....	574	543	593	583	608	595	497	326	177	52	23	5
2002.....	523	514	560	590	611	586	462	317	135	47	17	6
2001.....	503	475	529	546	619	547	436	308	131	46	7	5
2000.....	469	459	505	556	557	567	422	286	155	50	10	3
1999.....	430	452	468	558	590	587	400	261	128	46	14	1
American Indian 1/, female												
2004.....	369	409	452	530	574	572	657	496	305	138	40	2
2003.....	396	408	498	528	537	638	633	511	326	123	56	-
2002.....	328	380	432	505	585	574	593	429	335	135	36	1
2001.....	309	346	418	514	527	601	562	474	276	121	39	1
2000.....	274	354	380	493	535	563	510	442	287	119	33	-
1999.....	270	331	423	479	537	564	513	472	262	111	45	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All causes												
Asian or Pacific Islander 2/, both sexes												
2004.....	2,039	2,387	2,576	3,265	4,128	5,140	5,676	4,812	3,045	1,090	283	3
2003.....	1,935	2,210	2,623	3,314	4,250	5,274	5,563	4,524	2,795	1,054	282	5
2002.....	1,904	2,142	2,564	3,222	4,116	5,119	5,105	4,275	2,629	979	265	7
2001.....	1,853	2,043	2,499	3,197	4,050	4,891	4,907	4,025	2,338	859	255	8
2000.....	1,669	1,937	2,389	3,029	3,858	4,772	4,438	3,671	2,219	898	202	6
1999.....	1,583	1,866	2,381	2,937	3,940	4,570	4,266	3,574	2,063	754	177	7
Asian or Pacific Islander 2/, male												
2004.....	1,191	1,386	1,507	1,818	2,170	2,678	2,808	2,163	1,304	440	82	2
2003.....	1,118	1,257	1,546	1,875	2,211	2,763	2,827	2,061	1,193	430	81	3
2002.....	1,120	1,231	1,507	1,794	2,093	2,743	2,581	2,008	1,189	387	78	6
2001.....	1,050	1,222	1,474	1,713	2,135	2,631	2,455	1,897	1,077	342	57	4
2000.....	978	1,131	1,434	1,692	2,153	2,627	2,207	1,842	1,026	346	58	5
1999.....	890	1,105	1,393	1,589	2,147	2,534	2,160	1,843	1,005	274	40	5
Asian or Pacific Islander 2/, female												
2004.....	848	1,001	1,069	1,447	1,958	2,462	2,868	2,649	1,741	650	201	1
2003.....	817	953	1,077	1,439	2,039	2,511	2,736	2,463	1,602	624	201	2
2002.....	784	911	1,057	1,428	2,023	2,376	2,524	2,267	1,440	592	187	1
2001.....	803	821	1,025	1,484	1,915	2,260	2,452	2,128	1,261	517	198	4
2000.....	691	806	955	1,337	1,705	2,145	2,231	1,829	1,193	552	144	1
1999.....	693	761	988	1,348	1,793	2,036	2,106	1,731	1,058	480	137	2

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
I. Certain infectious and parasitic diseases (A00-B99)												
All races, both sexes												
2004.....	5,298	4,364	3,770	4,002	5,131	6,787	7,932	6,608	3,923	1,366	226	7
2003.....	5,021	4,098	3,596	3,861	5,174	7,056	7,671	6,600	3,848	1,356	226	6
2002.....	4,714	3,594	3,345	3,979	5,286	6,977	7,467	6,338	3,678	1,357	240	7
2001.....	4,154	3,244	3,190	3,760	5,022	6,499	6,694	5,856	3,557	1,213	220	4
2000.....	3,909	2,946	2,969	3,575	4,845	6,310	6,505	5,890	3,460	1,249	256	4
1999.....	3,577	2,909	2,831	3,845	5,005	6,379	6,471	5,976	3,560	1,356	236	2
All races, male												
2004.....	3,622	2,863	2,149	2,157	2,610	3,213	3,469	2,576	1,167	336	39	5
2003.....	3,497	2,653	2,036	2,095	2,599	3,292	3,348	2,429	1,126	299	34	5
2002.....	3,357	2,267	1,952	2,126	2,686	3,350	3,296	2,347	1,063	311	44	7
2001.....	2,909	2,086	1,796	1,975	2,547	3,132	2,967	2,149	1,053	270	32	4
2000.....	2,782	1,877	1,715	1,928	2,446	3,037	2,841	2,084	985	240	41	4
1999.....	2,502	1,837	1,625	2,084	2,547	3,091	2,844	2,147	1,000	291	24	2
All races, female												
2004.....	1,676	1,501	1,621	1,845	2,521	3,574	4,463	4,032	2,756	1,030	187	2
2003.....	1,524	1,445	1,560	1,766	2,575	3,764	4,323	4,171	2,722	1,057	192	1
2002.....	1,357	1,327	1,393	1,853	2,600	3,627	4,171	3,991	2,615	1,046	196	-
2001.....	1,245	1,158	1,394	1,785	2,475	3,367	3,727	3,707	2,504	943	188	-
2000.....	1,127	1,069	1,254	1,647	2,399	3,273	3,664	3,806	2,475	1,009	215	-
1999.....	1,075	1,072	1,206	1,761	2,458	3,288	3,627	3,829	2,560	1,065	212	-
White, both sexes												
2004.....	3,371	2,960	2,662	2,975	4,088	5,678	6,809	5,790	3,451	1,188	177	5
2003.....	3,135	2,754	2,534	2,841	4,139	5,867	6,632	5,748	3,353	1,156	171	4
2002.....	2,890	2,395	2,336	2,949	4,208	5,772	6,421	5,524	3,198	1,153	186	5
2001.....	2,528	2,168	2,209	2,802	3,985	5,404	5,747	5,080	3,096	1,034	170	1
2000.....	2,359	1,946	2,039	2,643	3,837	5,183	5,510	5,095	3,009	1,018	199	3
1999.....	2,195	1,867	1,989	2,877	3,899	5,206	5,501	5,102	3,049	1,158	178	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
I. Certain infectious and parasitic diseases (A00-B99)												
White, male												
2004.....	2,362	1,921	1,498	1,607	2,082	2,722	2,974	2,290	1,015	294	30	4
2003.....	2,228	1,759	1,423	1,533	2,078	2,750	2,908	2,121	985	257	27	3
2002.....	2,066	1,510	1,361	1,569	2,161	2,759	2,830	2,065	914	262	32	5
2001.....	1,804	1,384	1,242	1,474	2,019	2,622	2,562	1,865	901	233	20	1
2000.....	1,685	1,232	1,170	1,422	1,966	2,485	2,422	1,798	845	193	30	3
1999.....	1,551	1,162	1,122	1,555	1,994	2,564	2,414	1,816	843	241	19	-
White, female												
2004.....	1,009	1,039	1,164	1,368	2,006	2,956	3,835	3,500	2,436	894	147	1
2003.....	907	995	1,111	1,308	2,061	3,117	3,724	3,627	2,368	899	144	1
2002.....	824	885	975	1,380	2,047	3,013	3,591	3,459	2,284	891	154	-
2001.....	724	784	967	1,328	1,966	2,782	3,185	3,215	2,195	801	150	-
2000.....	674	714	869	1,221	1,871	2,698	3,088	3,297	2,164	825	169	-
1999.....	644	705	867	1,322	1,905	2,642	3,087	3,286	2,206	917	159	-
Black, both sexes												
2004.....	1,789	1,284	979	885	885	955	970	708	418	159	46	2
2003.....	1,770	1,210	947	879	876	1,007	906	753	440	184	51	2
2002.....	1,715	1,096	876	892	915	1,059	910	721	435	183	52	2
2001.....	1,518	989	884	830	887	957	837	697	409	165	47	3
2000.....	1,450	894	816	814	894	999	897	711	417	212	56	1
1999.....	1,273	945	740	842	973	1,018	863	777	452	184	58	2
Black, male												
2004.....	1,176	870	581	475	456	423	417	241	126	35	9	1
2003.....	1,202	823	548	498	440	450	386	268	125	38	7	2
2002.....	1,221	699	513	495	453	507	389	237	129	41	11	2
2001.....	1,031	651	504	439	449	439	347	245	127	32	11	3
2000.....	1,037	587	484	446	431	485	369	245	124	42	11	1
1999.....	883	610	451	462	490	450	373	282	127	45	5	2

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
I. Certain infectious and parasitic diseases (A00-B99)												
Black, female												
2004.....	613	414	398	410	429	532	553	467	292	124	37	1
2003.....	568	387	399	381	436	557	520	485	315	146	44	-
2002.....	494	397	363	397	462	552	521	484	306	142	41	-
2001.....	487	338	380	391	438	518	490	452	282	133	36	-
2000.....	413	307	332	368	463	514	528	466	293	170	45	-
1999.....	390	335	289	380	483	568	490	495	325	139	53	-
American Indian 1/, both sexes												
2004.....	51	31	40	31	21	30	21	21	7	3	-	-
2003.....	46	49	33	34	31	29	29	16	7	5	3	-
2002.....	30	32	29	34	35	28	29	14	10	4	1	-
2001.....	33	17	21	30	29	19	20	14	5	2	-	-
2000.....	32	29	31	34	16	27	19	14	6	6	-	-
1999.....	33	18	29	29	28	24	23	15	12	5	-	-
American Indian 1/, male												
2004.....	24	15	20	17	8	16	10	9	2	-	-	-
2003.....	24	27	16	15	9	9	8	6	1	1	-	-
2002.....	18	16	13	18	13	15	16	4	1	2	1	-
2001.....	24	9	10	13	12	12	10	6	2	1	-	-
2000.....	21	12	15	16	7	10	10	3	2	1	-	-
1999.....	20	10	11	18	13	7	12	3	6	3	-	-
American Indian 1/, female												
2004.....	27	16	20	14	13	14	11	12	5	3	-	-
2003.....	22	22	17	19	22	20	21	10	6	4	3	-
2002.....	12	16	16	16	22	13	13	10	9	2	-	-
2001.....	9	8	11	17	17	7	10	8	3	1	-	-
2000.....	11	17	16	18	9	17	9	11	4	5	-	-
1999.....	13	8	18	11	15	17	11	12	6	2	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
I. Certain infectious and parasitic diseases (A00-B99)												
Asian or Pacific Islander 2/, both sexes												
2004.....	87	89	89	111	137	124	132	89	47	16	3	-
2003.....	70	85	82	107	128	153	104	83	48	11	1	-
2002.....	79	71	104	104	128	118	107	79	35	17	1	-
2001.....	75	70	76	98	121	119	90	65	47	12	3	-
2000.....	68	77	83	84	98	101	79	70	28	13	1	-
1999.....	76	79	73	97	105	131	84	82	47	9	-	-
Asian or Pacific Islander 2/, male												
2004.....	60	57	50	58	64	52	68	36	24	7	-	-
2003.....	43	44	49	49	72	83	46	34	15	3	-	-
2002.....	52	42	65	44	59	69	61	41	19	6	-	-
2001.....	50	42	40	49	67	59	48	33	23	4	1	-
2000.....	39	46	46	44	42	57	40	38	14	4	-	-
1999.....	48	55	41	49	50	70	45	46	24	2	-	-
Asian or Pacific Islander 2/, female												
2004.....	27	32	39	53	73	72	64	53	23	9	3	-
2003.....	27	41	33	58	56	70	58	49	33	8	1	-
2002.....	27	29	39	60	69	49	46	38	16	11	1	-
2001.....	25	28	36	49	54	60	42	32	24	8	2	-
2000.....	29	31	37	40	56	44	39	32	14	9	1	-
1999.....	28	24	32	48	55	61	39	36	23	7	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intestinal infectious diseases (A00-A09)												
All races, both sexes												
2004.....	58	96	160	221	396	694	975	877	590	181	25	-
2003.....	45	75	110	163	291	533	671	653	387	121	17	-
2002.....	34	64	79	138	231	377	521	510	319	114	23	-
2001.....	37	30	54	77	165	244	339	299	191	75	10	-
2000.....	24	36	38	87	144	209	268	269	170	53	12	-
1999.....	16	16	37	72	116	178	226	216	149	61	14	-
All races, male												
2004.....	31	48	81	104	165	291	380	297	153	35	1	-
2003.....	23	38	49	70	120	218	266	202	88	24	2	-
2002.....	19	29	42	70	120	157	206	161	91	27	5	-
2001.....	15	16	21	35	79	110	125	94	50	14	-	-
2000.....	12	23	20	45	66	92	109	81	49	9	2	-
1999.....	10	10	19	33	51	82	92	55	46	13	2	-
All races, female												
2004.....	27	48	79	117	231	403	595	580	437	146	24	-
2003.....	22	37	61	93	171	315	405	451	299	97	15	-
2002.....	15	35	37	68	111	220	315	349	228	87	18	-
2001.....	22	14	33	42	86	134	214	205	141	61	10	-
2000.....	12	13	18	42	78	117	159	188	121	44	10	-
1999.....	6	6	18	39	65	96	134	161	103	48	12	-
White, both sexes												
2004.....	51	82	143	194	365	648	921	837	570	177	24	-
2003.....	34	65	93	149	271	484	645	624	367	113	16	-
2002.....	31	48	64	129	206	353	500	483	304	109	20	-
2001.....	35	26	43	62	153	229	313	290	180	68	10	-
2000.....	19	33	33	77	132	197	259	249	164	49	11	-
1999.....	14	14	30	66	106	164	212	209	142	58	13	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intestinal infectious diseases (A00-A09)												
White, male												
2004.....	26	36	74	90	152	273	361	285	145	34	1	-
2003.....	19	30	42	66	112	203	256	192	87	22	2	-
2002.....	18	22	34	64	104	146	199	153	85	27	5	-
2001.....	14	14	19	29	71	102	114	89	49	12	-	-
2000.....	9	20	18	41	61	87	103	76	47	8	2	-
1999.....	8	9	15	31	49	77	87	54	43	12	1	-
White, female												
2004.....	25	46	69	104	213	375	560	552	425	143	23	-
2003.....	15	35	51	83	159	281	389	432	280	91	14	-
2002.....	13	26	30	65	102	207	301	330	219	82	15	-
2001.....	21	12	24	33	82	127	199	201	131	56	10	-
2000.....	10	13	15	36	71	110	156	173	117	41	9	-
1999.....	6	5	15	35	57	87	125	155	99	46	12	-
Black, both sexes												
2004.....	7	14	13	22	25	35	46	35	16	4	1	-
2003.....	8	8	15	13	13	43	24	26	18	7	1	-
2002.....	2	15	14	9	23	20	17	25	12	4	3	-
2001.....	2	4	9	12	12	11	21	7	9	5	-	-
2000.....	5	3	3	8	7	10	7	17	5	4	-	-
1999.....	1	2	4	6	9	14	13	5	7	3	1	-
Black, male												
2004.....	5	12	5	12	10	11	15	10	5	1	-	-
2003.....	3	6	6	4	5	15	10	8	1	2	-	-
2002.....	1	7	8	6	15	8	5	7	5	-	-	-
2001.....	1	2	2	4	8	4	8	3	1	2	-	-
2000.....	3	3	1	3	3	4	5	4	2	1	-	-
1999.....	1	1	1	2	1	5	5	-	3	1	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intestinal infectious diseases (A00-A09)												
Black, female												
2004.....	2	2	8	10	15	24	31	25	11	3	1	-
2003.....	5	2	9	9	8	28	14	18	17	5	1	-
2002.....	1	8	6	3	8	12	12	18	7	4	3	-
2001.....	1	2	7	8	4	7	13	4	8	3	-	-
2000.....	2	-	2	5	4	6	2	13	3	3	-	-
1999.....	-	1	3	4	8	9	8	5	4	2	-	-
American Indian 1/, both sexes												
2004.....	-	-	2	3	3	4	2	-	-	-	-	-
2003.....	1	-	2	1	3	2	-	-	-	-	-	-
2002.....	-	-	1	-	1	1	1	-	2	-	-	-
2001.....	-	-	1	-	-	2	2	-	-	1	-	-
2000.....	-	-	2	1	-	-	-	1	-	-	-	-
1999.....	1	-	1	-	-	-	1	-	-	-	-	-
American Indian 1/, male												
2004.....	-	-	-	1	1	3	2	-	-	-	-	-
2003.....	-	-	1	-	1	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	1	-	-	-	-	-	-
2001.....	-	-	-	-	-	2	1	-	-	-	-	-
2000.....	-	-	1	1	-	-	-	-	-	-	-	-
1999.....	1	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, female												
2004.....	-	-	2	2	2	1	-	-	-	-	-	-
2003.....	1	-	1	1	2	2	-	-	-	-	-	-
2002.....	-	-	1	-	1	-	1	-	2	-	-	-
2001.....	-	-	1	-	-	-	1	-	-	1	-	-
2000.....	-	-	1	-	-	-	-	1	-	-	-	-
1999.....	-	-	-	-	-	-	1	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intestinal infectious diseases (A00-A09)												
Asian or Pacific Islander 2/, both sexes												
2004.....	-	-	2	2	3	7	6	5	4	-	-	-
2003.....	2	2	-	-	4	4	2	3	2	1	-	-
2002.....	1	1	-	-	1	3	3	2	1	1	-	-
2001.....	-	-	1	3	-	2	3	2	2	1	-	-
2000.....	-	-	-	1	5	2	2	2	1	-	1	-
1999.....	-	-	2	-	1	-	-	2	-	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	-	-	2	1	2	4	2	2	3	-	-	-
2003.....	1	2	-	-	2	-	-	2	-	-	-	-
2002.....	-	-	-	-	1	2	2	1	1	-	-	-
2001.....	-	-	-	2	-	2	2	2	-	-	-	-
2000.....	-	-	-	-	2	1	1	1	-	-	-	-
1999.....	-	-	2	-	1	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	-	-	-	1	1	3	4	3	1	-	-	-
2003.....	1	-	-	-	2	4	2	1	2	1	-	-
2002.....	1	1	-	-	-	1	1	1	-	1	-	-
2001.....	-	-	1	1	-	-	1	-	2	1	-	-
2000.....	-	-	-	1	3	1	1	1	1	-	1	-
1999.....	-	-	-	-	-	-	-	1	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other intestinal infectious diseases (A01-A08)												
All races, both sexes												
2004.....	54	93	157	215	386	675	951	858	576	175	25	-
2003.....	44	72	106	160	281	515	654	635	370	118	16	-
2002.....	34	62	76	133	222	370	501	492	306	108	22	-
2001.....	34	27	54	75	159	234	329	288	185	74	10	-
2000.....	23	36	35	80	136	201	260	255	161	50	11	-
1999.....	14	16	32	63	102	160	211	198	142	59	13	-
All races, male												
2004.....	29	45	80	101	163	285	371	291	151	35	1	-
2003.....	23	35	48	69	118	209	259	196	84	23	2	-
2002.....	19	29	40	67	115	154	199	158	85	26	5	-
2001.....	15	14	21	34	76	103	123	91	49	14	-	-
2000.....	12	23	19	42	62	88	106	73	47	8	2	-
1999.....	8	10	17	29	44	77	88	53	43	13	2	-
All races, female												
2004.....	25	48	77	114	223	390	580	567	425	140	24	-
2003.....	21	37	58	91	163	306	395	439	286	95	14	-
2002.....	15	33	36	66	107	216	302	334	221	82	17	-
2001.....	19	13	33	41	83	131	206	197	136	60	10	-
2000.....	11	13	16	38	74	113	154	182	114	42	9	-
1999.....	6	6	15	34	58	83	123	145	99	46	11	-
White, both sexes												
2004.....	48	81	140	188	356	632	898	819	557	172	24	-
2003.....	34	63	91	146	262	467	628	607	350	110	15	-
2002.....	31	46	62	124	199	347	482	465	291	103	19	-
2001.....	33	23	43	60	148	219	303	279	174	67	10	-
2000.....	19	33	30	70	124	189	251	237	155	46	10	-
1999.....	13	14	26	57	93	148	198	192	135	56	12	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other intestinal infectious diseases (A01-A08)												
White, male												
2004.....	25	35	73	87	150	268	352	280	144	34	1	-
2003.....	19	28	41	65	110	194	249	187	83	21	2	-
2002.....	18	22	33	61	101	143	193	150	79	26	5	-
2001.....	14	12	19	28	68	95	112	86	48	12	-	-
2000.....	9	20	17	38	57	83	100	69	45	7	2	-
1999.....	7	9	13	27	42	72	84	52	40	12	1	-
White, female												
2004.....	23	46	67	101	206	364	546	539	413	138	23	-
2003.....	15	35	50	81	152	273	379	420	267	89	13	-
2002.....	13	24	29	63	98	204	289	315	212	77	14	-
2001.....	19	11	24	32	80	124	191	193	126	55	10	-
2000.....	10	13	13	32	67	106	151	168	110	39	8	-
1999.....	6	5	13	30	51	76	114	140	95	44	11	-
Black, both sexes												
2004.....	6	12	13	22	25	33	45	34	15	3	1	-
2003.....	7	7	13	13	12	42	24	25	18	7	1	-
2002.....	2	15	13	9	21	19	15	25	12	4	3	-
2001.....	1	4	9	12	11	11	21	7	9	5	-	-
2000.....	4	3	3	8	7	10	7	15	5	4	-	-
1999.....	-	2	3	6	8	12	12	4	7	3	1	-
Black, male												
2004.....	4	10	5	12	10	11	15	9	4	1	-	-
2003.....	3	5	6	4	5	15	10	7	1	2	-	-
2002.....	1	7	7	6	13	8	4	7	5	-	-	-
2001.....	1	2	2	4	8	4	8	3	1	2	-	-
2000.....	3	3	1	3	3	4	5	3	2	1	-	-
1999.....	-	1	1	2	1	5	4	-	3	1	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other intestinal infectious diseases (A01-A08)												
Black, female												
2004.....	2	2	8	10	15	22	30	25	11	2	1	-
2003.....	4	2	7	9	7	27	14	18	17	5	1	-
2002.....	1	8	6	3	8	11	11	18	7	4	3	-
2001.....	-	2	7	8	3	7	13	4	8	3	-	-
2000.....	1	-	2	5	4	6	2	12	3	3	-	-
1999.....	-	1	2	4	7	7	8	4	4	2	-	-
American Indian 1/, both sexes												
2004.....	-	-	2	3	2	3	2	-	-	-	-	-
2003.....	1	-	2	1	3	2	-	-	-	-	-	-
2002.....	-	-	1	-	1	1	1	-	2	-	-	-
2001.....	-	-	1	-	-	2	2	-	-	1	-	-
2000.....	-	-	2	1	-	-	-	1	-	-	-	-
1999.....	1	-	1	-	-	-	1	-	-	-	-	-
American Indian 1/, male												
2004.....	-	-	-	1	1	2	2	-	-	-	-	-
2003.....	-	-	1	-	1	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	1	-	-	-	-	-	-
2001.....	-	-	-	-	-	2	1	-	-	-	-	-
2000.....	-	-	1	1	-	-	-	-	-	-	-	-
1999.....	1	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, female												
2004.....	-	-	2	2	1	1	-	-	-	-	-	-
2003.....	1	-	1	1	2	2	-	-	-	-	-	-
2002.....	-	-	1	-	1	-	1	-	2	-	-	-
2001.....	-	-	1	-	-	-	1	-	-	1	-	-
2000.....	-	-	1	-	-	-	-	1	-	-	-	-
1999.....	-	-	-	-	-	-	1	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other intestinal infectious diseases (A01-A08)												
Asian or Pacific Islander 2/, both sexes												
2004.....	-	-	2	2	3	7	6	5	4	-	-	-
2003.....	2	2	-	-	4	4	2	3	2	1	-	-
2002.....	1	1	-	-	1	3	3	2	1	1	-	-
2001.....	-	-	1	3	-	2	3	2	2	1	-	-
2000.....	-	-	-	1	5	2	2	2	1	-	1	-
1999.....	-	-	2	-	1	-	-	2	-	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	-	-	2	1	2	4	2	2	3	-	-	-
2003.....	1	2	-	-	2	-	-	2	-	-	-	-
2002.....	-	-	-	-	1	2	2	1	1	-	-	-
2001.....	-	-	-	2	-	2	2	2	-	-	-	-
2000.....	-	-	-	-	2	1	1	1	-	-	-	-
1999.....	-	-	2	-	1	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	-	-	-	1	1	3	4	3	1	-	-	-
2003.....	1	-	-	-	2	4	2	1	2	1	-	-
2002.....	1	1	-	-	-	1	1	1	-	1	-	-
2001.....	-	-	1	1	-	-	1	-	2	1	-	-
2000.....	-	-	-	1	3	1	1	1	1	-	1	-
1999.....	-	-	-	-	-	-	-	1	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Paratyphoid fevers and other salmonella infections (A01.1-A01.4,A02)												
All races, both sexes												
2004.....	1	2	2	4	3	4	3	6	-	-	-	-
2003.....	1	4	2	2	6	8	2	5	3	1	-	-
2002.....	2	-	-	3	1	3	5	1	-	-	-	-
2001.....	2	2	1	4	3	5	9	6	2	-	-	-
2000.....	3	2	-	1	6	3	2	1	2	-	-	-
1999.....	2	1	2	2	5	8	4	4	2	1	-	-
All races, male												
2004.....	1	2	2	2	1	4	2	4	-	-	-	-
2003.....	1	3	1	2	2	4	-	4	1	1	-	-
2002.....	1	-	-	1	1	1	4	1	-	-	-	-
2001.....	2	2	-	3	3	3	4	1	-	-	-	-
2000.....	1	2	-	1	4	1	-	-	1	-	-	-
1999.....	2	-	2	2	3	2	3	1	1	-	-	-
All races, female												
2004.....	-	-	-	2	2	-	1	2	-	-	-	-
2003.....	-	1	1	-	4	4	2	1	2	-	-	-
2002.....	1	-	-	2	-	2	1	-	-	-	-	-
2001.....	-	-	1	1	-	2	5	5	2	-	-	-
2000.....	2	-	-	-	2	2	2	1	1	-	-	-
1999.....	-	1	-	-	2	6	1	3	1	1	-	-
White, both sexes												
2004.....	1	1	1	1	2	3	3	5	-	-	-	-
2003.....	1	3	2	2	4	5	2	4	3	1	-	-
2002.....	2	-	-	3	-	2	5	1	-	-	-	-
2001.....	2	2	1	2	2	5	7	6	1	-	-	-
2000.....	3	2	-	1	5	3	2	1	2	-	-	-
1999.....	2	1	1	1	4	7	3	4	2	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Paratyphoid fevers and other salmonella infections (A01.1-A01.4,A02)												
White, male												
2004.....	1	1	1	1	-	3	2	3	-	-	-	-
2003.....	1	2	1	2	2	2	-	3	1	1	-	-
2002.....	1	-	-	1	-	1	4	1	-	-	-	-
2001.....	2	2	-	2	2	3	4	1	-	-	-	-
2000.....	1	2	-	1	4	1	-	-	1	-	-	-
1999.....	2	-	1	1	2	2	2	1	1	-	-	-
White, female												
2004.....	-	-	-	-	2	-	1	2	-	-	-	-
2003.....	-	1	1	-	2	3	2	1	2	-	-	-
2002.....	1	-	-	2	-	1	1	-	-	-	-	-
2001.....	-	-	1	-	-	2	3	5	1	-	-	-
2000.....	2	-	-	-	1	2	2	1	1	-	-	-
1999.....	-	1	-	-	2	5	1	3	1	1	-	-
Black, both sexes												
2004.....	-	1	-	2	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	1	3	-	-	-	-	-	-
2002.....	-	-	-	-	1	1	-	-	-	-	-	-
2001.....	-	-	-	-	1	-	2	-	1	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	1	-	1	1	-	-	-	-	-
Black, male												
2004.....	-	1	-	1	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	2	-	-	-	-	-	-
2002.....	-	-	-	-	1	-	-	-	-	-	-	-
2001.....	-	-	-	-	1	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	1	-	-	1	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Shigellosis (A03)												
All races, both sexes												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	1	-	-	-	1	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	1	-	-	-
2001.....	1	-	-	-	-	-	-	-	-	-	-	-
2000.....	1	-	1	-	-	2	3	-	-	-	1	-
1999.....	-	-	-	-	1	-	-	1	1	2	-	-
All races, male												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	1	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	1	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	1	-	-	1	2	-	-	-	-	-
1999.....	-	-	-	-	1	-	-	-	1	-	-	-
All races, female												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	1	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	1	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	1	-	-	-	-	1	1	-	-	-	1	-
1999.....	-	-	-	-	-	-	-	1	-	2	-	-
White, both sexes												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	1	-	-	-	1	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	1	-	-	-
2001.....	1	-	-	-	-	-	-	-	-	-	-	-
2000.....	1	-	1	-	-	2	3	-	-	-	1	-
1999.....	-	-	-	-	1	-	-	1	1	2	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other bacterial food-borne intoxications (A05)												
All races, both sexes												
2004.....	-	-	-	1	-	-	-	-	-	-	-	-
2003.....	1	-	1	-	1	-	1	1	1	-	-	-
2002.....	1	-	-	2	1	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	1	1	-	1	-	-	-
2000.....	2	1	-	-	1	1	1	1	-	-	-	-
1999.....	2	-	-	2	1	1	-	-	-	-	-	-
All races, male												
2004.....	-	-	-	1	-	-	-	-	-	-	-	-
2003.....	1	-	-	-	-	-	1	-	1	-	-	-
2002.....	1	-	-	2	1	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	1	-	-	-	-	-
2000.....	2	1	-	-	1	1	-	-	-	-	-	-
1999.....	2	-	-	2	1	-	-	-	-	-	-	-
All races, female												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	1	-	1	-	-	1	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	1	-	-	1	-	-	-
2000.....	-	-	-	-	-	-	1	1	-	-	-	-
1999.....	-	-	-	-	-	1	-	-	-	-	-	-
White, both sexes												
2004.....	-	-	-	1	-	-	-	-	-	-	-	-
2003.....	1	-	1	-	-	-	1	1	1	-	-	-
2002.....	1	-	-	2	1	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	1	1	-	-	-	-	-
2000.....	1	1	-	-	1	1	1	1	-	-	-	-
1999.....	2	-	-	2	1	1	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Amebiasis (A06)												
All races, both sexes												
2004.....	-	-	-	-	1	-	-	-	-	-	-	-
2003.....	-	-	1	-	1	-	-	-	-	-	-	-
2002.....	-	1	-	-	-	-	1	-	-	-	-	-
2001.....	1	-	-	-	-	1	-	-	-	-	-	-
2000.....	-	-	-	1	-	-	2	-	-	-	-	-
1999.....	-	-	1	-	-	1	-	-	-	-	-	-
All races, male												
2004.....	-	-	-	-	1	-	-	-	-	-	-	-
2003.....	-	-	1	-	1	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	1	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	1	-	-	-	-	-	-	-	-	-
All races, female												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	1	-	-	-	-	1	-	-	-	-	-
2001.....	-	-	-	-	-	1	-	-	-	-	-	-
2000.....	-	-	-	1	-	-	2	-	-	-	-	-
1999.....	-	-	-	-	-	1	-	-	-	-	-	-
White, both sexes												
2004.....	-	-	-	-	1	-	-	-	-	-	-	-
2003.....	-	-	1	-	1	-	-	-	-	-	-	-
2002.....	-	1	-	-	-	-	1	-	-	-	-	-
2001.....	1	-	-	-	-	1	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	2	-	-	-	-	-
1999.....	-	-	1	-	-	1	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intestinal infections due to other specified organisms (A04,A07-A08)												
All races, both sexes												
2004.....	53	91	155	210	382	671	948	852	576	175	25	-
2003.....	42	67	102	158	273	506	651	629	366	117	16	-
2002.....	31	61	76	128	220	367	495	491	305	108	22	-
2001.....	30	25	53	71	156	227	319	282	181	74	10	-
2000.....	17	33	34	78	129	195	252	253	159	50	10	-
1999.....	10	15	29	59	95	150	207	193	139	56	13	-
All races, male												
2004.....	28	43	78	98	161	281	369	287	151	35	1	-
2003.....	21	32	46	67	115	204	258	192	82	22	2	-
2002.....	17	29	40	64	113	153	195	157	85	26	5	-
2001.....	11	12	21	31	73	100	118	90	49	14	-	-
2000.....	9	20	18	41	57	85	104	73	46	8	2	-
1999.....	4	10	14	25	39	75	85	52	41	13	2	-
All races, female												
2004.....	25	48	77	112	221	390	579	565	425	140	24	-
2003.....	21	35	56	91	158	302	393	437	284	95	14	-
2002.....	14	32	36	64	107	214	300	334	220	82	17	-
2001.....	19	13	32	40	83	127	201	192	132	60	10	-
2000.....	8	13	16	37	72	110	148	180	113	42	8	-
1999.....	6	5	15	34	56	75	122	141	98	43	11	-
White, both sexes												
2004.....	47	80	139	186	353	629	895	814	557	172	24	-
2003.....	32	59	87	144	257	461	625	602	346	109	15	-
2002.....	28	45	62	119	198	345	476	464	290	103	19	-
2001.....	29	21	42	58	146	212	295	273	172	67	10	-
2000.....	14	30	29	69	118	183	243	235	153	46	9	-
1999.....	9	13	24	54	87	139	195	187	132	53	12	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Intestinal infections due to other specified organisms (A04,A07-A08)

White, male

2004.....	24	34	72	85	149	265	350	277	144	34	1	-
2003.....	17	26	39	63	107	191	248	184	81	20	2	-
2002.....	16	22	33	58	100	142	189	149	79	26	5	-
2001.....	10	10	19	26	66	92	107	85	48	12	-	-
2000.....	7	17	16	37	52	80	98	69	44	7	2	-
1999.....	3	9	11	24	38	70	82	51	38	12	1	-

White, female

2004.....	23	46	67	101	204	364	545	537	413	138	23	-
2003.....	15	33	48	81	150	270	377	418	265	89	13	-
2002.....	12	23	29	61	98	203	287	315	211	77	14	-
2001.....	19	11	23	32	80	120	188	188	124	55	10	-
2000.....	7	13	13	32	66	103	145	166	109	39	7	-
1999.....	6	4	13	30	49	69	113	136	94	41	11	-

Black, both sexes

2004.....	6	11	13	20	25	33	45	34	15	3	1	-
2003.....	7	7	13	13	11	39	24	25	18	7	1	-
2002.....	2	15	13	9	20	18	15	25	12	4	3	-
2001.....	1	4	9	12	10	11	19	7	7	5	-	-
2000.....	3	3	3	8	7	10	7	15	5	4	-	-
1999.....	-	2	3	5	8	11	11	4	7	3	1	-

Black, male

2004.....	4	9	5	11	10	11	15	9	4	1	-	-
2003.....	3	5	6	4	5	13	10	7	1	2	-	-
2002.....	1	7	7	6	12	8	4	7	5	-	-	-
2001.....	1	2	2	4	7	4	8	3	1	2	-	-
2000.....	2	3	1	3	3	4	5	3	2	1	-	-
1999.....	-	1	1	1	1	5	3	-	3	1	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intestinal infections due to other specified organisms (A04,A07-A08)												
Black, female												
2004.....	2	2	8	9	15	22	30	25	11	2	1	-
2003.....	4	2	7	9	6	26	14	18	17	5	1	-
2002.....	1	8	6	3	8	10	11	18	7	4	3	-
2001.....	-	2	7	8	3	7	11	4	6	3	-	-
2000.....	1	-	2	5	4	6	2	12	3	3	-	-
1999.....	-	1	2	4	7	6	8	4	4	2	-	-
American Indian 1/, both sexes												
2004.....	-	-	2	3	2	3	2	-	-	-	-	-
2003.....	1	-	2	1	2	2	-	-	-	-	-	-
2002.....	-	-	1	-	1	1	1	-	2	-	-	-
2001.....	-	-	1	-	-	2	2	-	-	1	-	-
2000.....	-	-	2	1	-	-	-	1	-	-	-	-
1999.....	1	-	1	-	-	-	1	-	-	-	-	-
American Indian 1/, male												
2004.....	-	-	-	1	1	2	2	-	-	-	-	-
2003.....	-	-	1	-	1	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	1	-	-	-	-	-	-
2001.....	-	-	-	-	-	2	1	-	-	-	-	-
2000.....	-	-	1	1	-	-	-	-	-	-	-	-
1999.....	1	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, female												
2004.....	-	-	2	2	1	1	-	-	-	-	-	-
2003.....	1	-	1	1	1	2	-	-	-	-	-	-
2002.....	-	-	1	-	1	-	1	-	2	-	-	-
2001.....	-	-	1	-	-	-	1	-	-	1	-	-
2000.....	-	-	1	-	-	-	-	1	-	-	-	-
1999.....	-	-	-	-	-	-	1	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intestinal infections due to other specified organisms (A04,A07-A08)												
Asian or Pacific Islander 2/, both sexes												
2004.....	-	-	1	1	2	6	6	4	4	-	-	-
2003.....	2	1	-	-	3	4	2	2	2	1	-	-
2002.....	1	1	-	-	1	3	3	2	1	1	-	-
2001.....	-	-	1	1	-	2	3	2	2	1	-	-
2000.....	-	-	-	-	4	2	2	2	1	-	1	-
1999.....	-	-	1	-	-	-	-	2	-	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	-	-	1	1	1	3	2	1	3	-	-	-
2003.....	1	1	-	-	2	-	-	1	-	-	-	-
2002.....	-	-	-	-	1	2	2	1	1	-	-	-
2001.....	-	-	-	1	-	2	2	2	-	-	-	-
2000.....	-	-	-	-	2	1	1	1	-	-	-	-
1999.....	-	-	1	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	-	-	-	-	1	3	4	3	1	-	-	-
2003.....	1	-	-	-	1	4	2	1	2	1	-	-
2002.....	1	1	-	-	-	1	1	1	-	1	-	-
2001.....	-	-	1	-	-	-	1	-	2	1	-	-
2000.....	-	-	-	-	2	1	1	1	1	-	1	-
1999.....	-	-	-	-	-	-	-	1	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Diarrhea and gastroenteritis of infectious origin (A09)

All races, both sexes

2004.....	4	3	3	6	10	19	24	19	14	6	-	-
2003.....	1	3	4	3	10	18	17	18	17	3	1	-
2002.....	-	2	3	5	9	7	20	18	13	6	1	-
2001.....	3	3	-	2	6	10	10	11	6	1	-	-
2000.....	1	-	3	7	8	7	8	14	9	3	1	-
1999.....	2	-	5	8	14	18	15	18	7	2	1	-

All races, male

2004.....	2	3	1	3	2	6	9	6	2	-	-	-
2003.....	-	3	1	1	2	9	7	6	4	1	-	-
2002.....	-	-	2	3	5	3	7	3	6	1	-	-
2001.....	-	2	-	1	3	7	2	3	1	-	-	-
2000.....	-	-	1	3	4	3	3	8	2	1	-	-
1999.....	2	-	2	3	7	5	4	2	3	-	-	-

All races, female

2004.....	2	-	2	3	8	13	15	13	12	6	-	-
2003.....	1	-	3	2	8	9	10	12	13	2	1	-
2002.....	-	2	1	2	4	4	13	15	7	5	1	-
2001.....	3	1	-	1	3	3	8	8	5	1	-	-
2000.....	1	-	2	4	4	4	5	6	7	2	1	-
1999.....	-	-	3	5	7	13	11	16	4	2	1	-

White, both sexes

2004.....	3	1	3	6	9	16	23	18	13	5	-	-
2003.....	-	2	2	3	9	17	17	17	17	3	1	-
2002.....	-	2	2	5	7	6	18	18	13	6	1	-
2001.....	2	3	-	2	5	10	10	11	6	1	-	-
2000.....	-	-	3	7	8	7	8	12	9	3	1	-
1999.....	1	-	4	8	13	16	14	17	7	2	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diarrhea and gastroenteritis of infectious origin (A09)												
White, male												
2004.....	1	1	1	3	2	5	9	5	1	-	-	-
2003.....	-	2	1	1	2	9	7	5	4	1	-	-
2002.....	-	-	1	3	3	3	6	3	6	1	-	-
2001.....	-	2	-	1	3	7	2	3	1	-	-	-
2000.....	-	-	1	3	4	3	3	7	2	1	-	-
1999.....	1	-	2	3	7	5	3	2	3	-	-	-
White, female												
2004.....	2	-	2	3	7	11	14	13	12	5	-	-
2003.....	-	-	1	2	7	8	10	12	13	2	1	-
2002.....	-	2	1	2	4	3	12	15	7	5	1	-
2001.....	2	1	-	1	2	3	8	8	5	1	-	-
2000.....	-	-	2	4	4	4	5	5	7	2	1	-
1999.....	-	-	2	5	6	11	11	15	4	2	1	-
Black, both sexes												
2004.....	1	2	-	-	-	2	1	1	1	1	-	-
2003.....	1	1	2	-	1	1	-	1	-	-	-	-
2002.....	-	-	1	-	2	1	2	-	-	-	-	-
2001.....	1	-	-	-	1	-	-	-	-	-	-	-
2000.....	1	-	-	-	-	-	-	2	-	-	-	-
1999.....	1	-	1	-	1	2	1	1	-	-	-	-
Black, male												
2004.....	1	2	-	-	-	-	-	1	1	-	-	-
2003.....	-	1	-	-	-	-	-	1	-	-	-	-
2002.....	-	-	1	-	2	-	1	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	1	-	-	-	-
1999.....	1	-	-	-	-	-	1	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Tuberculosis (A16-A19)												
All races, both sexes												
2004.....	47	35	44	55	82	81	84	68	37	11	-	-
2003.....	51	63	39	48	66	106	111	70	33	4	-	-
2002.....	46	44	63	82	79	114	120	68	42	9	2	-
2001.....	47	63	53	60	92	102	104	85	30	10	-	1
2000.....	53	31	50	70	85	132	120	74	31	10	-	1
1999.....	55	67	63	70	103	134	119	115	41	16	1	1
All races, male												
2004.....	32	26	34	38	55	37	47	38	17	3	-	-
2003.....	40	46	25	28	44	61	55	32	11	2	-	-
2002.....	35	36	39	54	48	56	65	36	16	2	1	-
2001.....	36	47	38	40	58	53	47	34	18	5	-	1
2000.....	41	23	36	46	41	69	70	38	13	5	-	1
1999.....	45	48	48	49	67	71	66	58	16	11	-	1
All races, female												
2004.....	15	9	10	17	27	44	37	30	20	8	-	-
2003.....	11	17	14	20	22	45	56	38	22	2	-	-
2002.....	11	8	24	28	31	58	55	32	26	7	1	-
2001.....	11	16	15	20	34	49	57	51	12	5	-	-
2000.....	12	8	14	24	44	63	50	36	18	5	-	-
1999.....	10	19	15	21	36	63	53	57	25	5	1	-
White, both sexes												
2004.....	20	19	23	29	49	55	58	47	24	5	-	-
2003.....	23	39	21	23	53	69	81	50	25	3	-	-
2002.....	23	20	37	41	47	78	85	48	30	7	1	-
2001.....	22	35	32	38	52	73	67	59	21	7	-	1
2000.....	31	8	25	38	49	79	76	53	26	6	-	-
1999.....	27	37	36	48	60	85	87	87	24	11	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Tuberculosis (A16-A19)												
White, male												
2004.....	14	15	19	19	34	28	30	27	11	-	-	-
2003.....	19	29	14	12	34	36	41	23	7	2	-	-
2002.....	19	15	22	31	33	37	46	27	11	1	1	-
2001.....	16	29	22	24	29	38	31	22	11	4	-	1
2000.....	22	6	18	26	27	40	38	27	12	3	-	-
1999.....	25	27	25	33	40	42	49	43	9	6	-	-
White, female												
2004.....	6	4	4	10	15	27	28	20	13	5	-	-
2003.....	4	10	7	11	19	33	40	27	18	1	-	-
2002.....	4	5	15	10	14	41	39	21	19	6	-	-
2001.....	6	6	10	14	23	35	36	37	10	3	-	-
2000.....	9	2	7	12	22	39	38	26	14	3	-	-
1999.....	2	10	11	15	20	43	38	44	15	5	-	-
Black, both sexes												
2004.....	20	11	15	15	26	14	18	16	7	4	-	-
2003.....	23	21	10	14	6	16	17	11	4	-	-	-
2002.....	18	18	16	25	18	17	20	10	9	-	-	-
2001.....	23	23	15	19	25	16	19	11	8	3	-	-
2000.....	21	18	16	25	27	29	31	13	2	4	-	1
1999.....	22	22	18	15	28	31	19	14	8	4	1	1
Black, male												
2004.....	12	9	11	10	16	4	13	8	3	1	-	-
2003.....	18	16	5	10	5	12	8	5	2	-	-	-
2002.....	13	15	10	16	9	11	11	3	4	-	-	-
2001.....	19	16	11	13	17	7	7	5	6	1	-	-
2000.....	18	13	13	17	11	17	23	6	-	2	-	1
1999.....	17	15	17	11	18	17	8	6	3	4	-	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Tuberculosis (A16-A19)												
Black, female												
2004.....	8	2	4	5	10	10	5	8	4	3	-	-
2003.....	5	5	5	4	1	4	9	6	2	-	-	-
2002.....	5	3	6	9	9	6	9	7	5	-	-	-
2001.....	4	7	4	6	8	9	12	6	2	2	-	-
2000.....	3	5	3	8	16	12	8	7	2	2	-	-
1999.....	5	7	1	4	10	14	11	8	5	-	1	-
American Indian 1/, both sexes												
2004.....	-	1	1	5	-	3	2	-	1	-	-	-
2003.....	2	-	-	-	1	2	2	1	-	1	-	-
2002.....	1	2	3	4	3	4	3	2	-	-	-	-
2001.....	-	1	1	-	2	1	3	2	-	-	-	-
2000.....	-	3	3	4	1	4	1	-	1	-	-	-
1999.....	1	1	1	1	5	1	-	1	3	1	-	-
American Indian 1/, male												
2004.....	-	1	1	4	-	2	1	-	-	-	-	-
2003.....	1	-	-	-	-	-	1	1	-	-	-	-
2002.....	-	2	1	2	1	2	2	2	-	-	-	-
2001.....	-	-	1	-	1	1	1	-	-	-	-	-
2000.....	-	2	2	2	1	2	1	-	-	-	-	-
1999.....	-	1	-	1	2	-	-	-	2	1	-	-
American Indian 1/, female												
2004.....	-	-	-	1	-	1	1	-	1	-	-	-
2003.....	1	-	-	-	1	2	1	-	-	1	-	-
2002.....	1	-	2	2	2	2	1	-	-	-	-	-
2001.....	-	1	-	-	1	-	2	2	-	-	-	-
2000.....	-	1	1	2	-	2	-	-	1	-	-	-
1999.....	1	-	1	-	3	1	-	1	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Tuberculosis (A16-A19)												
Asian or Pacific Islander 2/, both sexes												
2004.....	7	4	5	6	7	9	6	5	5	2	-	-
2003.....	3	3	8	11	6	19	11	8	4	-	-	-
2002.....	4	4	7	12	11	15	12	8	3	2	1	-
2001.....	2	4	5	3	13	12	15	13	1	-	-	-
2000.....	1	2	6	3	8	20	12	8	2	-	-	-
1999.....	5	7	8	6	10	17	13	13	6	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	6	1	3	5	5	3	3	3	3	2	-	-
2003.....	2	1	6	6	5	13	5	3	2	-	-	-
2002.....	3	4	6	5	5	6	6	4	1	1	-	-
2001.....	1	2	4	3	11	7	8	7	1	-	-	-
2000.....	1	2	3	1	2	10	8	5	1	-	-	-
1999.....	3	5	6	4	7	12	9	9	2	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	1	3	2	1	2	6	3	2	2	-	-	-
2003.....	1	2	2	5	1	6	6	5	2	-	-	-
2002.....	1	-	1	7	6	9	6	4	2	1	1	-
2001.....	1	2	1	-	2	5	7	6	-	-	-	-
2000.....	-	-	3	2	6	10	4	3	1	-	-	-
1999.....	2	2	2	2	3	5	4	4	4	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Respiratory tuberculosis (A16)												
All races, both sexes												
2004.....	34	26	32	41	67	57	63	62	35	9	-	-
2003.....	42	52	29	36	52	89	88	55	28	3	-	-
2002.....	36	37	47	61	59	87	105	60	37	7	2	-
2001.....	34	51	45	47	68	82	84	71	28	7	-	1
2000.....	42	24	36	52	67	107	102	64	26	8	-	1
1999.....	38	53	56	58	81	112	104	100	36	16	1	1
All races, male												
2004.....	26	18	25	30	44	28	36	36	17	3	-	-
2003.....	32	38	19	23	38	52	45	26	9	1	-	-
2002.....	27	30	30	44	36	40	56	32	15	1	1	-
2001.....	29	41	37	31	45	46	38	29	17	4	-	1
2000.....	35	21	29	38	35	58	58	33	13	4	-	1
1999.....	32	39	44	41	56	60	60	52	13	11	-	1
All races, female												
2004.....	8	8	7	11	23	29	27	26	18	6	-	-
2003.....	10	14	10	13	14	37	43	29	19	2	-	-
2002.....	9	7	17	17	23	47	49	28	22	6	1	-
2001.....	5	10	8	16	23	36	46	42	11	3	-	-
2000.....	7	3	7	14	32	49	44	31	13	4	-	-
1999.....	6	14	12	17	25	52	44	48	23	5	1	-
White, both sexes												
2004.....	14	14	18	24	39	37	45	42	23	4	-	-
2003.....	17	31	17	16	41	59	66	38	20	2	-	-
2002.....	17	17	26	34	36	59	70	42	27	6	1	-
2001.....	16	30	27	29	40	59	55	46	19	5	-	1
2000.....	26	7	19	27	39	65	66	45	22	5	-	-
1999.....	20	27	34	38	49	70	75	79	23	11	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Respiratory tuberculosis (A16)												
White, male												
2004.....	10	11	15	16	27	22	23	25	11	-	-	-
2003.....	13	23	12	10	29	30	34	18	5	1	-	-
2002.....	14	12	17	28	26	25	37	24	11	1	1	-
2001.....	14	26	22	18	24	34	25	17	10	3	-	1
2000.....	21	5	15	20	23	36	31	24	12	2	-	-
1999.....	19	21	24	27	34	36	43	41	9	6	-	-
White, female												
2004.....	4	3	3	8	12	15	22	17	12	4	-	-
2003.....	4	8	5	6	12	29	32	20	15	1	-	-
2002.....	3	5	9	6	10	34	33	18	16	5	-	-
2001.....	2	4	5	11	16	25	30	29	9	2	-	-
2000.....	5	2	4	7	16	29	35	21	10	3	-	-
1999.....	1	6	10	11	15	34	32	38	14	5	-	-
Black, both sexes												
2004.....	14	7	10	10	21	10	13	16	6	3	-	-
2003.....	21	18	7	11	6	12	14	9	4	-	-	-
2002.....	16	15	14	19	13	14	20	8	7	-	-	-
2001.....	17	18	13	15	17	13	15	10	8	2	-	-
2000.....	15	12	10	21	22	24	24	12	2	3	-	1
1999.....	14	20	16	13	20	26	17	11	7	4	1	1
Black, male												
2004.....	11	5	8	7	12	2	9	8	3	1	-	-
2003.....	16	14	4	8	5	10	7	5	2	-	-	-
2002.....	12	13	9	12	7	10	11	2	3	-	-	-
2001.....	15	14	10	10	13	6	6	5	6	1	-	-
2000.....	13	12	10	16	10	13	19	5	-	2	-	1
1999.....	11	13	16	9	13	14	8	4	2	4	-	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Respiratory tuberculosis (A16)												
Black, female												
2004.....	3	2	2	3	9	8	4	8	3	2	-	-
2003.....	5	4	3	3	1	2	7	4	2	-	-	-
2002.....	4	2	5	7	6	4	9	6	4	-	-	-
2001.....	2	4	3	5	4	7	9	5	2	1	-	-
2000.....	2	-	-	5	12	11	5	7	2	1	-	-
1999.....	3	7	-	4	7	12	9	7	5	-	1	-
American Indian 1/, both sexes												
2004.....	-	1	-	2	-	3	2	-	1	-	-	-
2003.....	2	-	-	-	-	1	2	-	-	1	-	-
2002.....	1	2	2	1	2	3	3	2	-	-	-	-
2001.....	-	-	1	-	1	1	3	2	-	-	-	-
2000.....	-	3	2	2	-	3	-	-	-	-	-	-
1999.....	-	-	1	1	4	1	-	-	1	1	-	-
American Indian 1/, male												
2004.....	-	1	-	2	-	2	1	-	-	-	-	-
2003.....	1	-	-	-	-	-	1	-	-	-	-	-
2002.....	-	2	-	1	-	1	2	2	-	-	-	-
2001.....	-	-	1	-	-	1	1	-	-	-	-	-
2000.....	-	2	2	1	-	1	-	-	-	-	-	-
1999.....	-	-	-	1	2	-	-	-	-	1	-	-
American Indian 1/, female												
2004.....	-	-	-	-	-	1	1	-	1	-	-	-
2003.....	1	-	-	-	-	1	1	-	-	1	-	-
2002.....	1	-	2	-	2	2	1	-	-	-	-	-
2001.....	-	-	-	-	1	-	2	2	-	-	-	-
2000.....	-	1	-	1	-	2	-	-	-	-	-	-
1999.....	-	-	1	-	2	1	-	-	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Respiratory tuberculosis (A16)												
Asian or Pacific Islander 2/, both sexes												
2004.....	6	4	4	5	7	7	3	4	5	2	-	-
2003.....	2	3	5	9	5	17	6	8	4	-	-	-
2002.....	2	3	5	7	8	11	12	8	3	1	1	-
2001.....	1	3	4	3	10	9	11	13	1	-	-	-
2000.....	1	2	5	2	6	15	12	7	2	-	-	-
1999.....	4	6	5	6	8	15	12	10	5	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	5	1	2	5	5	2	3	3	3	2	-	-
2003.....	2	1	3	5	4	12	3	3	2	-	-	-
2002.....	1	3	4	3	3	4	6	4	1	-	-	-
2001.....	-	1	4	3	8	5	6	7	1	-	-	-
2000.....	1	2	2	1	2	8	8	4	1	-	-	-
1999.....	2	5	4	4	7	10	9	7	2	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	1	3	2	-	2	5	-	1	2	-	-	-
2003.....	-	2	2	4	1	5	3	5	2	-	-	-
2002.....	1	-	1	4	5	7	6	4	2	1	1	-
2001.....	1	2	-	-	2	4	5	6	-	-	-	-
2000.....	-	-	3	1	4	7	4	3	1	-	-	-
1999.....	2	1	1	2	1	5	3	3	3	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other tuberculosis (A17-A19)												
All races, both sexes												
2004.....	13	9	12	14	15	24	21	6	2	2	-	-
2003.....	9	11	10	12	14	17	23	15	5	1	-	-
2002.....	10	7	16	21	20	27	15	8	5	2	-	-
2001.....	13	12	8	13	24	20	20	14	2	3	-	-
2000.....	11	7	14	18	18	25	18	10	5	2	-	-
1999.....	17	14	7	12	22	22	15	15	5	-	-	-
All races, male												
2004.....	6	8	9	8	11	9	11	2	-	-	-	-
2003.....	8	8	6	5	6	9	10	6	2	1	-	-
2002.....	8	6	9	10	12	16	9	4	1	1	-	-
2001.....	7	6	1	9	13	7	9	5	1	1	-	-
2000.....	6	2	7	8	6	11	12	5	-	1	-	-
1999.....	13	9	4	8	11	11	6	6	3	-	-	-
All races, female												
2004.....	7	1	3	6	4	15	10	4	2	2	-	-
2003.....	1	3	4	7	8	8	13	9	3	-	-	-
2002.....	2	1	7	11	8	11	6	4	4	1	-	-
2001.....	6	6	7	4	11	13	11	9	1	2	-	-
2000.....	5	5	7	10	12	14	6	5	5	1	-	-
1999.....	4	5	3	4	11	11	9	9	2	-	-	-
White, both sexes												
2004.....	6	5	5	5	10	18	13	5	1	1	-	-
2003.....	6	8	4	7	12	10	15	12	5	1	-	-
2002.....	6	3	11	7	11	19	15	6	3	1	-	-
2001.....	6	5	5	9	12	14	12	13	2	2	-	-
2000.....	5	1	6	11	10	14	10	8	4	1	-	-
1999.....	7	10	2	10	11	15	12	8	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other tuberculosis (A17-A19)												
White, male												
2004.....	4	4	4	3	7	6	7	2	-	-	-	-
2003.....	6	6	2	2	5	6	7	5	2	1	-	-
2002.....	5	3	5	3	7	12	9	3	-	-	-	-
2001.....	2	3	-	6	5	4	6	5	1	1	-	-
2000.....	1	1	3	6	4	4	7	3	-	1	-	-
1999.....	6	6	1	6	6	6	6	2	-	-	-	-
White, female												
2004.....	2	1	1	2	3	12	6	3	1	1	-	-
2003.....	-	2	2	5	7	4	8	7	3	-	-	-
2002.....	1	-	6	4	4	7	6	3	3	1	-	-
2001.....	4	2	5	3	7	10	6	8	1	1	-	-
2000.....	4	-	3	5	6	10	3	5	4	-	-	-
1999.....	1	4	1	4	5	9	6	6	1	-	-	-
Black, both sexes												
2004.....	6	4	5	5	5	4	5	-	1	1	-	-
2003.....	2	3	3	3	-	4	3	2	-	-	-	-
2002.....	2	3	2	6	5	3	-	2	2	-	-	-
2001.....	6	5	2	4	8	3	4	1	-	1	-	-
2000.....	6	6	6	4	5	5	7	1	-	1	-	-
1999.....	8	2	2	2	8	5	2	3	1	-	-	-
Black, male												
2004.....	1	4	3	3	4	2	4	-	-	-	-	-
2003.....	2	2	1	2	-	2	1	-	-	-	-	-
2002.....	1	2	1	4	2	1	-	1	1	-	-	-
2001.....	4	2	1	3	4	1	1	-	-	-	-	-
2000.....	5	1	3	1	1	4	4	1	-	-	-	-
1999.....	6	2	1	2	5	3	-	2	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other tuberculosis (A17-A19)												
Black, female												
2004.....	5	-	2	2	1	2	1	-	1	1	-	-
2003.....	-	1	2	1	-	2	2	2	-	-	-	-
2002.....	1	1	1	2	3	2	-	1	1	-	-	-
2001.....	2	3	1	1	4	2	3	1	-	1	-	-
2000.....	1	5	3	3	4	1	3	-	-	1	-	-
1999.....	2	-	1	-	3	2	2	1	-	-	-	-
American Indian 1/, both sexes												
2004.....	-	-	1	3	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	1	1	-	1	-	-	-	-
2002.....	-	-	1	3	1	1	-	-	-	-	-	-
2001.....	-	1	-	-	1	-	-	-	-	-	-	-
2000.....	-	-	1	2	1	1	1	-	1	-	-	-
1999.....	1	1	-	-	1	-	-	1	2	-	-	-
American Indian 1/, male												
2004.....	-	-	1	2	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	1	-	-	-	-
2002.....	-	-	1	1	1	1	-	-	-	-	-	-
2001.....	-	-	-	-	1	-	-	-	-	-	-	-
2000.....	-	-	-	1	1	1	1	-	-	-	-	-
1999.....	-	1	-	-	-	-	-	-	2	-	-	-
American Indian 1/, female												
2004.....	-	-	-	1	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	1	1	-	-	-	-	-	-
2002.....	-	-	-	2	-	-	-	-	-	-	-	-
2001.....	-	1	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	1	1	-	-	-	-	1	-	-	-
1999.....	1	-	-	-	1	-	-	1	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other tuberculosis (A17-A19)												
Asian or Pacific Islander 2/, both sexes												
2004.....	1	-	1	1	-	2	3	1	-	-	-	-
2003.....	1	-	3	2	1	2	5	-	-	-	-	-
2002.....	2	1	2	5	3	4	-	-	-	1	-	-
2001.....	1	1	1	-	3	3	4	-	-	-	-	-
2000.....	-	-	1	1	2	5	-	1	-	-	-	-
1999.....	1	1	3	-	2	2	1	3	1	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	1	-	1	-	-	1	-	-	-	-	-	-
2003.....	-	-	3	1	1	1	2	-	-	-	-	-
2002.....	2	1	2	2	2	2	-	-	-	1	-	-
2001.....	1	1	-	-	3	2	2	-	-	-	-	-
2000.....	-	-	1	-	-	2	-	1	-	-	-	-
1999.....	1	-	2	-	-	2	-	2	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	-	-	-	1	-	1	3	1	-	-	-	-
2003.....	1	-	-	1	-	1	3	-	-	-	-	-
2002.....	-	-	-	3	1	2	-	-	-	-	-	-
2001.....	-	-	1	-	-	1	2	-	-	-	-	-
2000.....	-	-	-	1	2	3	-	-	-	-	-	-
1999.....	-	1	1	-	2	-	1	1	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Zoonotic and other bacterial diseases (A20-A49)												
All races, both sexes												
2004.....	1,426	1,813	2,163	2,593	3,668	5,062	6,045	5,140	3,058	1,090	177	3
2003.....	1,310	1,750	2,131	2,578	3,694	5,322	6,090	5,373	3,176	1,136	191	3
2002.....	1,249	1,590	1,948	2,693	3,921	5,438	6,052	5,229	3,086	1,148	205	3
2001.....	1,219	1,470	1,816	2,533	3,740	5,222	5,574	5,033	3,115	1,066	196	-
2000.....	1,147	1,329	1,721	2,406	3,543	4,959	5,402	5,061	3,041	1,110	222	1
1999.....	1,030	1,259	1,629	2,531	3,592	4,958	5,295	5,097	3,114	1,193	202	-
All races, male												
2004.....	761	975	1,081	1,320	1,861	2,414	2,687	2,009	913	280	35	1
2003.....	695	938	1,083	1,367	1,824	2,491	2,634	2,001	943	253	28	2
2002.....	677	818	1,025	1,341	1,956	2,608	2,659	1,934	863	259	37	3
2001.....	651	777	933	1,296	1,892	2,515	2,466	1,852	914	236	32	-
2000.....	655	698	896	1,243	1,756	2,353	2,331	1,762	853	209	36	1
1999.....	553	653	858	1,318	1,802	2,389	2,302	1,816	859	247	22	-
All races, female												
2004.....	665	838	1,082	1,273	1,807	2,648	3,358	3,131	2,145	810	142	2
2003.....	615	812	1,048	1,211	1,870	2,831	3,456	3,372	2,233	883	163	1
2002.....	572	772	923	1,352	1,965	2,830	3,393	3,295	2,223	889	168	-
2001.....	568	693	883	1,237	1,848	2,707	3,108	3,181	2,201	830	164	-
2000.....	492	631	825	1,163	1,787	2,606	3,071	3,299	2,188	901	186	-
1999.....	477	606	771	1,213	1,790	2,569	2,993	3,281	2,255	946	180	-
White, both sexes												
2004.....	981	1,328	1,584	1,959	2,915	4,183	5,124	4,443	2,644	935	131	2
2003.....	870	1,269	1,575	1,914	2,929	4,415	5,226	4,625	2,725	951	138	2
2002.....	841	1,143	1,403	2,037	3,130	4,469	5,156	4,510	2,656	957	156	2
2001.....	821	1,036	1,331	1,927	2,974	4,312	4,784	4,341	2,696	902	149	-
2000.....	786	951	1,236	1,800	2,818	4,045	4,551	4,353	2,623	899	168	1
1999.....	706	876	1,192	1,915	2,803	4,015	4,468	4,324	2,657	1,010	148	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Zoonotic and other bacterial diseases (A20-A49)												
White, male												
2004.....	540	716	792	1,011	1,486	2,023	2,277	1,767	786	244	26	1
2003.....	470	669	809	1,019	1,462	2,082	2,276	1,735	810	213	21	1
2002.....	456	601	761	1,014	1,588	2,139	2,264	1,688	736	215	25	2
2001.....	445	539	700	990	1,505	2,096	2,130	1,598	778	203	20	-
2000.....	442	507	657	941	1,431	1,923	1,994	1,513	725	169	26	1
1999.....	380	449	614	1,000	1,414	1,969	1,941	1,536	724	206	18	-
White, female												
2004.....	441	612	792	948	1,429	2,160	2,847	2,676	1,858	691	105	1
2003.....	400	600	766	895	1,467	2,333	2,950	2,890	1,915	738	117	1
2002.....	385	542	642	1,023	1,542	2,330	2,892	2,822	1,920	742	131	-
2001.....	376	497	631	937	1,469	2,216	2,654	2,743	1,918	699	129	-
2000.....	344	444	579	859	1,387	2,122	2,557	2,840	1,898	730	142	-
1999.....	326	427	578	915	1,389	2,046	2,527	2,788	1,933	804	130	-
Black, both sexes												
2004.....	405	447	509	576	674	793	819	608	375	139	44	1
2003.....	401	425	510	599	685	810	779	677	403	172	49	1
2002.....	374	418	490	595	706	882	813	653	392	174	48	1
2001.....	360	406	452	554	686	828	731	643	376	152	44	-
2000.....	335	342	452	555	672	846	785	646	393	193	54	-
1999.....	293	350	404	555	726	846	760	706	413	170	54	-
Black, male												
2004.....	204	241	250	280	334	353	354	207	108	31	9	-
2003.....	210	243	250	317	320	361	327	237	119	36	7	1
2002.....	205	203	238	293	331	416	347	216	110	38	11	1
2001.....	186	225	215	275	339	378	303	231	114	28	11	-
2000.....	198	176	223	274	302	398	304	222	115	36	10	-
1999.....	158	184	228	291	354	372	328	248	114	37	4	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Zoonotic and other bacterial diseases (A20-A49)												
Black, female												
2004.....	201	206	259	296	340	440	465	401	267	108	35	1
2003.....	191	182	260	282	365	449	452	440	284	136	42	-
2002.....	169	215	252	302	375	466	466	437	282	136	37	-
2001.....	174	181	237	279	347	450	428	412	262	124	33	-
2000.....	137	166	229	281	370	448	481	424	278	157	44	-
1999.....	135	166	176	264	372	474	432	458	299	133	50	-
American Indian 1/, both sexes												
2004.....	17	13	22	14	14	20	14	18	5	3	-	-
2003.....	15	29	18	24	24	18	24	14	7	3	3	-
2002.....	15	13	18	23	26	21	22	12	8	4	1	-
2001.....	11	13	10	20	22	12	12	9	5	1	-	-
2000.....	11	19	13	22	12	20	16	13	4	6	-	-
1999.....	17	13	17	19	16	19	15	11	8	4	-	-
American Indian 1/, male												
2004.....	8	3	12	7	6	10	6	8	2	-	-	-
2003.....	5	13	9	10	7	4	7	5	1	1	-	-
2002.....	9	5	8	13	10	11	12	2	1	2	1	-
2001.....	5	7	4	11	10	8	6	4	2	1	-	-
2000.....	7	9	7	10	5	6	8	3	2	1	-	-
1999.....	9	8	5	10	8	7	7	2	3	2	-	-
American Indian 1/, female												
2004.....	9	10	10	7	8	10	8	10	3	3	-	-
2003.....	10	16	9	14	17	14	17	9	6	2	3	-
2002.....	6	8	10	10	16	10	10	10	7	2	-	-
2001.....	6	6	6	9	12	4	6	5	3	-	-	-
2000.....	4	10	6	12	7	14	8	10	2	5	-	-
1999.....	8	5	12	9	8	12	8	9	5	2	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Zoonotic and other bacterial diseases (A20-A49)												
Asian or Pacific Islander 2/, both sexes												
2004.....	23	25	48	44	65	66	88	71	34	13	2	-
2003.....	24	27	28	41	56	79	61	57	41	10	1	-
2002.....	19	16	37	38	59	66	61	54	30	13	-	-
2001.....	27	15	23	32	58	70	47	40	38	11	3	-
2000.....	15	17	20	29	41	48	50	49	21	12	-	-
1999.....	14	20	16	42	47	78	52	56	36	9	-	-
Asian or Pacific Islander 2/, male												
2004.....	9	15	27	22	35	28	50	27	17	5	-	-
2003.....	10	13	15	21	35	44	24	24	13	3	-	-
2002.....	7	9	18	21	27	42	36	28	16	4	-	-
2001.....	15	6	14	20	38	33	27	19	20	4	1	-
2000.....	8	6	9	18	18	26	25	24	11	3	-	-
1999.....	6	12	11	17	26	41	26	30	18	2	-	-
Asian or Pacific Islander 2/, female												
2004.....	14	10	21	22	30	38	38	44	17	8	2	-
2003.....	14	14	13	20	21	35	37	33	28	7	1	-
2002.....	12	7	19	17	32	24	25	26	14	9	-	-
2001.....	12	9	9	12	20	37	20	21	18	7	2	-
2000.....	7	11	11	11	23	22	25	25	10	9	-	-
1999.....	8	8	5	25	21	37	26	26	18	7	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Plague (A20)

All races, both sexes

2004.....	-	-	-	1	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	1	-	-	-	-	-	-

All races, male

2004.....	-	-	-	1	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-

All races, female

2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	1	-	-	-	-	-	-

White, both sexes

2004.....	-	-	-	1	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	1	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Tetanus (A33-A35)												
All races, both sexes												
2004.....	-	-	-	-	1	1	-	1	1	-	-	-
2003.....	-	2	-	1	-	-	-	1	-	-	-	-
2002.....	-	1	-	1	-	2	1	-	-	-	-	-
2001.....	-	1	1	-	-	-	1	2	-	-	-	-
2000.....	-	-	-	1	2	-	1	-	1	-	-	-
1999.....	-	1	1	-	-	3	-	1	-	-	-	-
All races, male												
2004.....	-	-	-	-	1	-	-	-	-	-	-	-
2003.....	-	1	-	1	-	-	-	-	-	-	-	-
2002.....	-	1	-	-	-	1	1	-	-	-	-	-
2001.....	-	1	-	-	-	-	-	1	-	-	-	-
2000.....	-	-	-	1	-	-	-	-	1	-	-	-
1999.....	-	1	1	-	-	2	-	-	-	-	-	-
All races, female												
2004.....	-	-	-	-	-	1	-	1	1	-	-	-
2003.....	-	1	-	-	-	-	-	1	-	-	-	-
2002.....	-	-	-	1	-	1	-	-	-	-	-	-
2001.....	-	-	1	-	-	-	1	1	-	-	-	-
2000.....	-	-	-	-	2	-	1	-	-	-	-	-
1999.....	-	-	-	-	-	1	-	1	-	-	-	-
White, both sexes												
2004.....	-	-	-	-	1	1	-	1	1	-	-	-
2003.....	-	1	-	1	-	-	-	1	-	-	-	-
2002.....	-	1	-	1	-	2	1	-	-	-	-	-
2001.....	-	1	-	-	-	-	1	2	-	-	-	-
2000.....	-	-	-	1	1	-	1	-	1	-	-	-
1999.....	-	1	1	-	-	2	-	1	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Tetanus (A33-A35)												
White, male												
2004.....	-	-	-	-	1	-	-	-	-	-	-	-
2003.....	-	1	-	1	-	-	-	-	-	-	-	-
2002.....	-	1	-	-	-	1	1	-	-	-	-	-
2001.....	-	1	-	-	-	-	-	1	-	-	-	-
2000.....	-	-	-	1	-	-	-	-	1	-	-	-
1999.....	-	1	1	-	-	1	-	-	-	-	-	-
White, female												
2004.....	-	-	-	-	-	1	-	1	1	-	-	-
2003.....	-	-	-	-	-	-	-	1	-	-	-	-
2002.....	-	-	-	1	-	1	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	1	1	-	-	-	-
2000.....	-	-	-	-	1	-	1	-	-	-	-	-
1999.....	-	-	-	-	-	1	-	1	-	-	-	-
Black, both sexes												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	1	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	1	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	1	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	1	-	-	-	-	-	-
Black, male												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	1	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Scarlet fever and erysipelas (A38,A46)												
All races, both sexes												
2004.....	-	-	-	1	-	-	-	1	-	1	-	-
2003.....	-	-	-	-	-	-	-	-	-	1	-	-
2002.....	-	-	-	-	-	-	1	1	-	-	-	-
2001.....	-	-	-	1	-	-	-	-	1	-	-	-
2000.....	-	-	-	-	-	1	-	-	-	-	-	-
1999.....	-	-	-	1	1	1	-	-	1	-	1	-
All races, male												
2004.....	-	-	-	1	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	1	-	-	-
2000.....	-	-	-	-	-	1	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-
All races, female												
2004.....	-	-	-	-	-	-	-	1	-	1	-	-
2003.....	-	-	-	-	-	-	-	-	-	1	-	-
2002.....	-	-	-	-	-	-	1	1	-	-	-	-
2001.....	-	-	-	1	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	1	-	-	-
1999.....	-	-	-	1	1	1	-	-	1	-	1	-
White, both sexes												
2004.....	-	-	-	1	-	-	-	1	-	1	-	-
2003.....	-	-	-	-	-	-	-	-	-	1	-	-
2002.....	-	-	-	-	-	-	1	1	-	-	-	-
2001.....	-	-	-	1	-	-	-	-	1	-	-	-
2000.....	-	-	-	-	-	1	-	-	-	-	-	-
1999.....	-	-	-	1	-	1	-	-	1	-	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Meningococcal infection (A39)												
All races, both sexes												
2004.....	8	4	2	3	4	1	2	3	2	2	1	-
2003.....	10	9	2	3	6	4	5	3	-	-	-	-
2002.....	9	9	4	3	4	3	7	2	-	3	-	-
2001.....	4	11	4	7	8	7	1	6	1	2	-	-
2000.....	16	6	3	3	5	7	8	2	4	1	-	-
1999.....	6	3	1	9	-	7	6	3	4	1	-	-
All races, male												
2004.....	7	2	2	2	1	-	1	-	-	-	-	-
2003.....	5	4	1	1	1	-	1	1	-	-	-	-
2002.....	5	2	2	1	1	1	2	1	-	1	-	-
2001.....	2	5	2	2	3	2	1	1	-	1	-	-
2000.....	6	1	1	-	2	4	3	-	-	1	-	-
1999.....	3	-	-	5	-	2	2	-	1	1	-	-
All races, female												
2004.....	1	2	-	1	3	1	1	3	2	2	1	-
2003.....	5	5	1	2	5	4	4	2	-	-	-	-
2002.....	4	7	2	2	3	2	5	1	-	2	-	-
2001.....	2	6	2	5	5	5	-	5	1	1	-	-
2000.....	10	5	2	3	3	3	5	2	4	-	-	-
1999.....	3	3	1	4	-	5	4	3	3	-	-	-
White, both sexes												
2004.....	5	3	2	3	3	1	2	2	2	2	1	-
2003.....	5	6	1	3	6	4	5	3	-	-	-	-
2002.....	7	7	4	2	4	3	7	2	-	3	-	-
2001.....	4	10	1	6	8	6	1	5	1	2	-	-
2000.....	12	5	3	2	5	7	8	2	4	1	-	-
1999.....	5	3	1	9	-	7	5	3	4	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Meningococcal infection (A39)												
Black, female												
2004.....	-	-	-	-	-	-	-	1	-	-	-	-
2003.....	2	1	-	-	-	-	-	-	-	-	-	-
2002.....	-	1	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	2	1	-	-	-	-	-	-	-	-
2000.....	1	1	-	1	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes												
2004.....	1	-	-	-	-	-	-	-	-	-	-	-
2003.....	1	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, male												
2004.....	1	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, female												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	1	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	1	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Septicemia (A40-A41)												
All races, both sexes												
2004.....	1,353	1,700	2,045	2,481	3,502	4,828	5,758	4,914	2,943	1,045	173	3
2003.....	1,241	1,631	2,020	2,459	3,511	5,096	5,836	5,178	3,064	1,112	189	3
2002.....	1,170	1,499	1,861	2,589	3,747	5,195	5,815	5,026	2,988	1,111	199	3
2001.....	1,140	1,392	1,719	2,409	3,589	5,000	5,351	4,856	3,001	1,020	192	-
2000.....	1,064	1,260	1,639	2,303	3,401	4,745	5,193	4,900	2,949	1,079	216	1
1999.....	954	1,178	1,536	2,352	3,398	4,686	5,024	4,854	2,984	1,133	195	-
All races, male												
2004.....	720	910	1,024	1,247	1,770	2,308	2,562	1,904	869	267	34	1
2003.....	657	864	1,023	1,299	1,724	2,377	2,517	1,920	894	247	28	2
2002.....	629	767	981	1,292	1,862	2,476	2,552	1,857	834	247	35	3
2001.....	604	728	878	1,227	1,817	2,402	2,360	1,788	869	225	32	-
2000.....	600	657	856	1,182	1,685	2,249	2,233	1,690	812	199	35	1
1999.....	513	613	799	1,218	1,701	2,252	2,174	1,712	817	236	21	-
All races, female												
2004.....	633	790	1,021	1,234	1,732	2,520	3,196	3,010	2,074	778	139	2
2003.....	584	767	997	1,160	1,787	2,719	3,319	3,258	2,170	865	161	1
2002.....	541	732	880	1,297	1,885	2,719	3,263	3,169	2,154	864	164	-
2001.....	536	664	841	1,182	1,772	2,598	2,991	3,068	2,132	795	160	-
2000.....	464	603	783	1,121	1,716	2,496	2,960	3,210	2,137	880	181	-
1999.....	441	565	737	1,134	1,697	2,434	2,850	3,142	2,167	897	174	-
White, both sexes												
2004.....	931	1,237	1,490	1,868	2,781	3,990	4,869	4,240	2,540	897	127	2
2003.....	826	1,180	1,486	1,817	2,770	4,218	4,992	4,448	2,625	927	136	2
2002.....	782	1,070	1,340	1,948	2,990	4,259	4,943	4,323	2,569	925	151	2
2001.....	765	980	1,260	1,830	2,847	4,123	4,585	4,179	2,596	861	146	-
2000.....	724	897	1,169	1,716	2,702	3,859	4,364	4,206	2,546	875	163	1
1999.....	650	813	1,115	1,769	2,642	3,799	4,236	4,110	2,546	957	143	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Septicemia (A40-A41)												
White, male												
2004.....	510	664	747	951	1,419	1,936	2,169	1,672	746	232	25	1
2003.....	448	616	762	965	1,379	1,981	2,169	1,661	767	207	21	1
2002.....	422	560	730	972	1,510	2,025	2,168	1,614	709	205	24	2
2001.....	409	502	658	937	1,441	1,995	2,034	1,541	737	193	20	-
2000.....	404	474	623	892	1,372	1,833	1,904	1,448	692	161	25	1
1999.....	351	418	565	917	1,330	1,858	1,837	1,443	690	195	18	-
White, female												
2004.....	421	573	743	917	1,362	2,054	2,700	2,568	1,794	665	102	1
2003.....	378	564	724	852	1,391	2,237	2,823	2,787	1,858	720	115	1
2002.....	360	510	610	976	1,480	2,234	2,775	2,709	1,860	720	127	-
2001.....	356	478	602	893	1,406	2,128	2,551	2,638	1,859	668	126	-
2000.....	320	423	546	824	1,330	2,026	2,460	2,758	1,854	714	138	-
1999.....	299	395	550	852	1,312	1,941	2,399	2,667	1,856	762	125	-
Black, both sexes												
2004.....	385	428	488	561	650	756	792	593	366	133	44	1
2003.....	378	400	491	582	665	784	762	662	392	172	49	1
2002.....	357	405	471	584	676	853	798	638	382	169	47	1
2001.....	339	385	429	532	666	801	713	630	366	149	43	-
2000.....	316	328	440	540	649	824	766	635	380	187	53	-
1999.....	274	333	389	524	699	800	725	682	396	163	52	-
Black, male												
2004.....	194	230	238	269	316	338	341	200	105	30	9	-
2003.....	195	227	239	304	305	351	319	232	113	36	7	1
2002.....	193	195	228	287	316	400	340	213	108	36	10	1
2001.....	176	213	204	261	330	368	295	225	111	27	11	-
2000.....	183	169	218	265	290	387	297	218	109	34	10	-
1999.....	147	175	219	275	339	350	306	240	107	37	3	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Septicemia (A40-A41)												
Black, female												
2004.....	191	198	250	292	334	418	451	393	261	103	35	1
2003.....	183	173	252	278	360	433	443	430	279	136	42	-
2002.....	164	210	243	297	360	453	458	425	274	133	37	-
2001.....	163	172	225	271	336	433	418	405	255	122	32	-
2000.....	133	159	222	275	359	437	469	417	271	153	43	-
1999.....	127	158	170	249	360	450	419	442	289	126	49	-
American Indian 1/, both sexes												
2004.....	15	11	22	12	13	19	12	17	5	3	-	-
2003.....	14	28	17	23	23	18	24	13	6	3	3	-
2002.....	13	11	16	20	26	20	20	12	8	4	1	-
2001.....	11	12	9	18	21	10	11	9	4	1	-	-
2000.....	9	18	11	22	11	19	15	12	4	6	-	-
1999.....	16	12	17	18	15	18	14	11	8	4	-	-
American Indian 1/, male												
2004.....	7	2	12	7	5	9	4	8	2	-	-	-
2003.....	5	12	8	10	7	4	7	4	1	1	-	-
2002.....	8	4	7	13	10	11	12	2	1	2	1	-
2001.....	5	7	4	10	10	7	6	4	2	1	-	-
2000.....	5	8	6	10	5	6	8	2	2	1	-	-
1999.....	9	8	5	10	8	7	7	2	3	2	-	-
American Indian 1/, female												
2004.....	8	9	10	5	8	10	8	9	3	3	-	-
2003.....	9	16	9	13	16	14	17	9	5	2	3	-
2002.....	5	7	9	7	16	9	8	10	7	2	-	-
2001.....	6	5	5	8	11	3	5	5	2	-	-	-
2000.....	4	10	5	12	6	13	7	10	2	5	-	-
1999.....	7	4	12	8	7	11	7	9	5	2	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Septicemia (A40-A41)												
Asian or Pacific Islander 2/, both sexes												
2004.....	22	24	45	40	58	63	85	64	32	12	2	-
2003.....	23	23	26	37	53	76	58	55	41	10	1	-
2002.....	18	13	34	37	55	63	54	53	29	13	-	-
2001.....	25	15	21	29	55	66	42	38	35	9	3	-
2000.....	15	17	19	25	39	43	48	47	19	11	-	-
1999.....	14	20	15	41	42	69	49	51	34	9	-	-
Asian or Pacific Islander 2/, male												
2004.....	9	14	27	20	30	25	48	24	16	5	-	-
2003.....	9	9	14	20	33	41	22	23	13	3	-	-
2002.....	6	8	16	20	26	40	32	28	16	4	-	-
2001.....	14	6	12	19	36	32	25	18	19	4	1	-
2000.....	8	6	9	15	18	23	24	22	9	3	-	-
1999.....	6	12	10	16	24	37	24	27	17	2	-	-
Asian or Pacific Islander 2/, female												
2004.....	13	10	18	20	28	38	37	40	16	7	2	-
2003.....	14	14	12	17	20	35	36	32	28	7	1	-
2002.....	12	5	18	17	29	23	22	25	13	9	-	-
2001.....	11	9	9	10	19	34	17	20	16	5	2	-
2000.....	7	11	10	10	21	20	24	25	10	8	-	-
1999.....	8	8	5	25	18	32	25	24	17	7	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other zoonotic and bacterial diseases (A21-A22,A24-A32,A42-A44,A48-A49)												
All races, both sexes												
2004.....	65	109	116	107	161	232	285	221	112	42	3	-
2003.....	59	108	108	115	177	222	249	191	112	23	2	-
2002.....	70	81	83	99	170	238	228	200	98	34	6	-
2001.....	75	66	92	116	143	215	221	169	112	44	4	-
2000.....	67	63	79	99	135	206	200	159	87	30	6	-
1999.....	70	77	90	169	193	260	265	239	125	59	6	-
All races, male												
2004.....	34	63	55	69	89	106	124	105	44	13	1	-
2003.....	33	69	58	66	99	114	116	80	49	6	-	-
2002.....	43	48	42	48	93	130	104	76	29	11	2	-
2001.....	45	43	53	67	72	111	105	62	44	10	-	-
2000.....	49	40	39	60	69	99	95	72	40	9	1	-
1999.....	37	39	57	95	101	133	126	104	41	10	1	-
All races, female												
2004.....	31	46	61	38	72	126	161	116	68	29	2	-
2003.....	26	39	50	49	78	108	133	111	63	17	2	-
2002.....	27	33	41	51	77	108	124	124	69	23	4	-
2001.....	30	23	39	49	71	104	116	107	68	34	4	-
2000.....	18	23	40	39	66	107	105	87	47	21	5	-
1999.....	33	38	33	74	92	127	139	135	84	49	5	-
White, both sexes												
2004.....	45	88	92	86	130	191	253	199	101	35	3	-
2003.....	39	82	87	93	153	193	229	173	100	23	2	-
2002.....	52	65	59	85	136	205	204	184	87	29	5	-
2001.....	52	45	70	90	119	183	197	155	98	39	3	-
2000.....	50	49	64	81	110	178	178	145	72	23	5	-
1999.....	51	59	74	136	161	205	227	210	106	52	4	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other zoonotic and bacterial diseases (A21-A22,A24-A32,A42-A44,A48-A49)												
White, male												
2004.....	26	51	43	56	66	87	107	95	40	12	1	-
2003.....	19	50	46	52	82	101	106	73	43	6	-	-
2002.....	31	38	29	42	77	112	93	73	27	9	1	-
2001.....	34	32	41	51	61	100	95	56	40	9	-	-
2000.....	35	32	33	48	57	85	87	65	32	7	1	-
1999.....	27	30	47	78	84	108	102	93	33	10	-	-
White, female												
2004.....	19	37	49	30	64	104	146	104	61	23	2	-
2003.....	20	32	41	41	71	92	123	100	57	17	2	-
2002.....	21	27	30	43	59	93	111	111	60	20	4	-
2001.....	18	13	29	39	58	83	102	99	58	30	3	-
2000.....	15	17	31	33	53	93	91	80	40	16	4	-
1999.....	24	29	27	58	77	97	125	117	73	42	4	-
Black, both sexes												
2004.....	18	18	21	15	23	37	27	14	9	6	-	-
2003.....	19	21	18	17	20	26	17	15	11	-	-	-
2002.....	15	12	19	11	30	29	15	15	10	5	1	-
2001.....	21	20	19	21	20	26	18	12	10	3	1	-
2000.....	15	13	12	14	22	22	19	11	13	6	1	-
1999.....	18	17	15	31	26	45	35	24	17	7	2	-
Black, male												
2004.....	8	10	12	11	17	15	13	7	3	1	-	-
2003.....	13	14	10	13	15	10	8	5	6	-	-	-
2002.....	10	8	10	6	15	16	7	3	2	2	1	-
2001.....	10	11	10	14	9	9	8	5	3	1	-	-
2000.....	12	7	5	9	12	11	7	4	6	2	-	-
1999.....	10	9	9	16	15	21	22	8	7	-	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other zoonotic and bacterial diseases (A21-A22,A24-A32,A42-A44,A48-A49)												
Black, female												
2004.....	10	8	9	4	6	22	14	7	6	5	-	-
2003.....	6	7	8	4	5	16	9	10	5	-	-	-
2002.....	5	4	9	5	15	13	8	12	8	3	-	-
2001.....	11	9	9	7	11	17	10	7	7	2	1	-
2000.....	3	6	7	5	10	11	12	7	7	4	1	-
1999.....	8	8	6	15	11	24	13	16	10	7	1	-
American Indian 1/, both sexes												
2004.....	1	2	-	2	1	1	2	1	-	-	-	-
2003.....	-	1	1	1	1	-	-	1	1	-	-	-
2002.....	2	2	2	3	-	1	2	-	-	-	-	-
2001.....	-	1	1	2	1	2	1	-	1	-	-	-
2000.....	2	1	2	-	1	1	1	1	-	-	-	-
1999.....	1	1	-	1	1	1	-	-	-	-	-	-
American Indian 1/, male												
2004.....	-	1	-	-	1	1	2	-	-	-	-	-
2003.....	-	1	1	-	-	-	-	1	-	-	-	-
2002.....	1	1	1	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	1	-	1	-	-	-	-	-	-
2000.....	2	1	1	-	-	-	-	1	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, female												
2004.....	1	1	-	2	-	-	-	1	-	-	-	-
2003.....	-	-	-	1	1	-	-	-	1	-	-	-
2002.....	1	1	1	3	-	1	2	-	-	-	-	-
2001.....	-	1	1	1	1	1	1	-	1	-	-	-
2000.....	-	-	1	-	1	1	1	-	-	1	-	-
1999.....	1	1	-	1	1	1	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other zoonotic and bacterial diseases (A21-A22,A24-A32,A42-A44,A48-A49)												
Asian or Pacific Islander 2/, both sexes												
2004.....	1	1	3	4	7	3	3	7	2	1	-	-
2003.....	1	4	2	4	3	3	3	2	-	-	-	-
2002.....	1	2	3	-	4	3	7	1	1	-	-	-
2001.....	2	-	2	3	3	4	5	2	3	2	-	-
2000.....	-	-	1	4	2	5	2	2	2	1	-	-
1999.....	-	-	1	1	5	9	3	5	2	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	-	1	-	2	5	3	2	3	1	-	-	-
2003.....	1	4	1	1	2	3	2	1	-	-	-	-
2002.....	1	1	2	-	1	2	4	-	-	-	-	-
2001.....	1	-	2	1	2	1	2	1	1	-	-	-
2000.....	-	-	-	3	-	3	1	2	2	-	-	-
1999.....	-	-	1	1	2	4	2	3	1	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	1	-	3	2	2	-	1	4	1	1	-	-
2003.....	-	-	1	3	1	-	1	1	-	-	-	-
2002.....	-	1	1	-	3	1	3	1	1	-	-	-
2001.....	1	-	-	2	1	3	3	1	2	2	-	-
2000.....	-	-	1	1	2	2	1	-	-	1	-	-
1999.....	-	-	-	-	3	5	1	2	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Infections with a predominately sexual mode of transmission (A50-A64)

All races, both sexes

2004.....	3	3	3	3	4	8	4	10	5	1	-	-
2003.....	4	2	1	4	3	7	6	8	4	4	-	-
2002.....	4	2	5	5	2	5	8	6	3	-	-	-
2001.....	2	3	1	4	2	5	9	3	2	2	-	-
2000.....	6	2	1	2	8	7	11	8	3	1	-	-
1999.....	2	2	3	2	4	8	6	2	2	4	-	-

All races, male

2004.....	1	2	2	3	2	4	2	2	3	-	-	-
2003.....	3	1	1	2	3	2	4	3	2	-	-	-
2002.....	4	1	3	4	1	3	1	2	1	-	-	-
2001.....	1	2	1	2	2	2	3	2	-	1	-	-
2000.....	5	2	1	2	2	3	7	4	2	-	-	-
1999.....	1	2	2	2	3	1	-	1	-	1	-	-

All races, female

2004.....	2	1	1	-	2	4	2	8	2	1	-	-
2003.....	1	1	-	2	-	5	2	5	2	4	-	-
2002.....	-	1	2	1	1	2	7	4	2	-	-	-
2001.....	1	1	-	2	-	3	6	1	2	1	-	-
2000.....	1	-	-	-	6	4	4	4	1	1	-	-
1999.....	1	-	1	-	1	7	6	1	2	3	-	-

White, both sexes

2004.....	1	1	2	-	4	5	1	4	3	-	-	-
2003.....	2	2	-	3	-	5	4	5	4	4	-	-
2002.....	2	-	3	3	1	1	5	4	2	-	-	-
2001.....	2	2	1	1	1	4	6	3	-	1	-	-
2000.....	1	-	-	1	5	3	9	6	3	1	-	-
1999.....	2	1	3	2	1	4	4	-	1	3	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Infections with a predominately sexual mode of transmission (A50-A64)												
White, male												
2004.....	-	1	2	-	2	2	-	-	2	-	-	-
2003.....	1	1	-	2	-	2	3	2	2	-	-	-
2002.....	2	-	2	2	1	1	-	2	-	-	-	-
2001.....	1	1	1	-	1	2	1	2	-	-	-	-
2000.....	1	-	-	1	1	-	5	2	2	-	-	-
1999.....	1	1	2	2	-	1	-	-	-	-	-	-
White, female												
2004.....	1	-	-	-	2	3	1	4	1	-	-	-
2003.....	1	1	-	1	-	3	1	3	2	4	-	-
2002.....	-	-	1	1	-	-	5	2	2	-	-	-
2001.....	1	1	-	1	-	2	5	1	-	1	-	-
2000.....	-	-	-	-	4	3	4	4	1	1	-	-
1999.....	1	-	1	-	1	3	4	-	1	3	-	-
Black, both sexes												
2004.....	1	2	1	3	-	3	2	5	2	1	-	-
2003.....	2	-	1	1	3	2	1	2	-	-	-	-
2002.....	2	2	2	2	1	4	3	2	1	-	-	-
2001.....	-	-	-	2	1	-	3	-	2	1	-	-
2000.....	5	2	1	1	3	4	2	2	-	-	-	-
1999.....	-	1	-	-	3	3	2	2	1	1	-	-
Black, male												
2004.....	1	1	-	3	-	2	1	2	1	-	-	-
2003.....	2	-	1	-	3	-	1	-	-	-	-	-
2002.....	2	1	1	2	-	2	1	-	1	-	-	-
2001.....	-	-	-	1	1	-	2	-	-	1	-	-
2000.....	4	2	1	1	1	3	2	2	-	-	-	-
1999.....	-	1	-	-	3	-	-	1	-	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Infections with a predominately sexual mode of transmission (A50-A64)												
Asian or Pacific Islander 2/, both sexes												
2004.....	-	-	-	-	-	-	1	-	-	-	-	-
2003.....	-	-	-	-	-	-	1	1	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	1	-	1	-	1	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	-	-	-	-	-	-	1	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	1	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	1	-	1	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	1	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	1	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	1	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Syphilis (A50-A53)												
All races, both sexes												
2004.....	3	2	2	3	3	8	4	8	3	1	-	-
2003.....	3	2	1	3	3	6	2	5	3	1	-	-
2002.....	4	2	3	4	2	5	7	6	3	-	-	-
2001.....	1	3	1	3	2	4	9	2	2	1	-	-
2000.....	4	2	1	1	6	7	6	6	3	1	-	-
1999.....	2	1	3	2	4	6	4	2	1	1	-	-
All races, male												
2004.....	1	1	1	3	1	4	2	2	2	-	-	-
2003.....	3	1	1	2	3	2	2	2	1	-	-	-
2002.....	4	1	1	4	1	3	1	2	1	-	-	-
2001.....	1	2	1	1	2	2	3	1	-	-	-	-
2000.....	4	2	1	1	1	3	4	3	2	-	-	-
1999.....	1	1	2	2	3	-	-	1	-	-	-	-
All races, female												
2004.....	2	1	1	-	2	4	2	6	1	1	-	-
2003.....	-	1	-	1	-	4	-	3	2	1	-	-
2002.....	-	1	2	-	1	2	6	4	2	-	-	-
2001.....	-	1	-	2	-	2	6	1	2	1	-	-
2000.....	-	-	-	-	5	4	2	3	1	1	-	-
1999.....	1	-	1	-	1	6	4	1	1	1	-	-
White, both sexes												
2004.....	1	-	1	-	3	5	1	2	1	-	-	-
2003.....	1	2	-	2	-	4	1	2	3	1	-	-
2002.....	2	-	1	2	1	1	4	4	2	-	-	-
2001.....	1	2	1	1	1	4	6	2	-	1	-	-
2000.....	1	-	-	-	3	3	4	4	3	1	-	-
1999.....	2	-	3	2	1	2	2	-	-	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Syphilis (A50-A53)												
White, male												
2004.....	-	-	1	-	1	2	-	-	1	-	-	-
2003.....	1	1	-	2	-	2	1	1	1	-	-	-
2002.....	2	-	-	2	1	1	-	2	-	-	-	-
2001.....	1	1	1	-	1	2	1	1	-	-	-	-
2000.....	1	-	-	-	-	-	2	1	2	-	-	-
1999.....	1	-	2	2	-	-	-	-	-	-	-	-
White, female												
2004.....	1	-	-	-	2	3	1	2	-	-	-	-
2003.....	-	1	-	-	-	2	-	1	2	1	-	-
2002.....	-	-	1	-	-	-	4	2	2	-	-	-
2001.....	-	1	-	1	-	2	5	1	-	1	-	-
2000.....	-	-	-	-	3	3	2	3	1	1	-	-
1999.....	1	-	1	-	1	2	2	-	-	1	-	-
Black, both sexes												
2004.....	1	2	1	3	-	3	2	5	2	1	-	-
2003.....	2	-	1	1	3	2	1	2	-	-	-	-
2002.....	2	2	2	2	1	4	3	2	1	-	-	-
2001.....	-	-	-	2	1	-	3	-	2	-	-	-
2000.....	3	2	1	1	3	4	2	2	-	-	-	-
1999.....	-	1	-	-	3	3	2	2	1	-	-	-
Black, male												
2004.....	1	1	-	3	-	2	1	2	1	-	-	-
2003.....	2	-	1	-	3	-	1	-	-	-	-	-
2002.....	2	1	1	2	-	2	1	-	1	-	-	-
2001.....	-	-	-	1	1	-	2	-	-	-	-	-
2000.....	3	2	1	1	3	3	2	2	-	-	-	-
1999.....	-	1	-	-	3	-	-	1	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Syphilis (A50-A53)												
Asian or Pacific Islander 2/, both sexes												
2004.....	-	-	-	-	-	-	1	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	1	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	1	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	-	-	-	-	-	-	1	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	1	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	1	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	1	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiovascular syphilis (A52.0)												
All races, both sexes												
2004.....	-	-	1	-	-	2	1	-	-	-	-	-
2003.....	1	-	-	1	1	1	-	-	1	-	-	-
2002.....	1	-	2	1	-	-	1	-	-	-	-	-
2001.....	-	-	-	-	1	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	1	-	-	-	-	-
1999.....	-	-	1	-	1	-	-	-	-	-	-	-
All races, male												
2004.....	-	-	1	-	-	1	-	-	-	-	-	-
2003.....	1	-	-	1	1	1	-	-	-	-	-	-
2002.....	1	-	1	1	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	1	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	1	-	-	-	-	-	-	-	-	-
All races, female												
2004.....	-	-	-	-	-	1	1	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	1	-	-	-
2002.....	-	-	1	-	-	-	1	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	1	-	-	-	-	-
1999.....	-	-	-	-	1	-	-	-	-	-	-	-
White, both sexes												
2004.....	-	-	1	-	-	1	1	-	-	-	-	-
2003.....	-	-	-	1	-	1	-	-	1	-	-	-
2002.....	1	-	-	1	-	-	1	-	-	-	-	-
2001.....	-	-	-	-	1	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	1	-	-	-	-	-
1999.....	-	-	1	-	1	-	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Neurosyphilis (A52.1-A52.3)												
All races, both sexes												
2004.....	3	2	1	3	3	5	1	4	2	1	-	-
2003.....	2	1	-	2	1	3	1	1	1	1	-	-
2002.....	1	2	1	3	1	4	6	4	2	-	-	-
2001.....	1	1	1	2	1	3	5	2	-	1	-	-
2000.....	4	1	1	-	5	6	5	4	1	-	-	-
1999.....	1	-	1	2	3	3	2	2	-	-	-	-
All races, male												
2004.....	1	1	-	3	1	3	-	1	2	-	-	-
2003.....	2	-	-	1	1	1	1	1	1	-	-	-
2002.....	1	1	-	3	1	2	1	1	-	-	-	-
2001.....	1	1	1	-	1	2	1	1	-	-	-	-
2000.....	4	1	1	-	1	3	4	1	1	-	-	-
1999.....	1	-	1	2	3	-	-	1	-	-	-	-
All races, female												
2004.....	2	1	1	-	2	2	1	3	-	1	-	-
2003.....	-	1	-	1	-	2	-	-	-	1	-	-
2002.....	-	1	1	-	-	2	5	3	2	-	-	-
2001.....	-	-	-	2	-	1	4	1	-	1	-	-
2000.....	-	-	-	-	4	3	1	3	-	-	-	-
1999.....	-	-	-	-	-	3	2	1	-	-	-	-
White, both sexes												
2004.....	1	-	-	-	3	3	-	1	1	-	-	-
2003.....	1	1	-	1	-	3	-	1	1	1	-	-
2002.....	-	-	1	1	1	1	3	2	2	-	-	-
2001.....	1	1	1	1	-	3	3	2	-	1	-	-
2000.....	1	-	-	-	2	3	3	3	1	-	-	-
1999.....	1	-	1	2	-	1	1	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Neurosyphilis (A52.1-A52.3)												
White, male												
2004.....	-	-	-	-	1	1	-	-	1	-	-	-
2003.....	1	-	-	1	-	1	-	1	1	-	-	-
2002.....	-	-	-	1	1	1	-	1	-	-	-	-
2001.....	1	1	1	-	-	2	-	1	-	-	-	-
2000.....	1	-	-	-	-	-	2	-	1	-	-	-
1999.....	1	-	1	2	-	-	-	-	-	-	-	-
White, female												
2004.....	1	-	-	-	2	2	-	1	-	-	-	-
2003.....	-	1	-	-	-	2	-	-	-	1	-	-
2002.....	-	-	1	-	-	-	3	1	2	-	-	-
2001.....	-	-	-	1	-	1	3	1	-	1	-	-
2000.....	-	-	-	-	2	3	1	3	-	-	-	-
1999.....	-	-	-	-	-	1	1	-	-	-	-	-
Black, both sexes												
2004.....	1	2	1	3	-	2	1	3	1	1	-	-
2003.....	1	-	-	1	1	-	1	-	-	-	-	-
2002.....	1	2	-	2	-	3	3	2	-	-	-	-
2001.....	-	-	-	1	1	-	2	-	-	-	-	-
2000.....	3	1	1	-	3	3	2	1	-	-	-	-
1999.....	-	-	-	-	3	1	1	2	-	-	-	-
Black, male												
2004.....	1	1	-	3	-	2	-	1	1	-	-	-
2003.....	1	-	-	-	1	-	1	-	-	-	-	-
2002.....	1	1	-	2	-	1	1	-	-	-	-	-
2001.....	-	-	-	-	1	-	1	-	-	-	-	-
2000.....	3	1	1	-	1	3	2	1	-	-	-	-
1999.....	-	-	-	-	3	-	-	1	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Neurosyphilis (A52.1-A52.3)

Asian or Pacific Islander 2/, both sexes

2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	1	-	-	-	-	-	-

Asian or Pacific Islander 2/, male

2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-

Asian or Pacific Islander 2/, female

2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	1	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified syphilis (A50-A51,A52.7-A52.9,A53)												
All races, both sexes												
2004.....	-	-	-	-	-	1	2	4	1	-	-	-
2003.....	-	1	1	-	1	2	1	4	1	-	-	-
2002.....	2	-	-	-	1	1	-	2	1	-	-	-
2001.....	-	2	-	1	-	1	4	-	2	-	-	-
2000.....	-	1	-	1	1	1	-	2	2	1	-	-
1999.....	1	1	1	-	-	3	2	-	1	1	-	-
All races, male												
2004.....	-	-	-	-	-	-	2	1	-	-	-	-
2003.....	-	1	1	-	1	-	1	1	-	-	-	-
2002.....	2	-	-	-	-	1	-	1	1	-	-	-
2001.....	-	1	-	1	-	-	2	-	-	-	-	-
2000.....	-	1	-	1	-	-	-	2	1	-	-	-
1999.....	-	1	-	-	-	-	-	-	-	-	-	-
All races, female												
2004.....	-	-	-	-	-	1	-	3	1	-	-	-
2003.....	-	-	-	-	-	2	-	3	1	-	-	-
2002.....	-	-	-	-	1	-	-	1	-	-	-	-
2001.....	-	1	-	-	-	1	2	-	2	-	-	-
2000.....	-	-	-	-	1	1	-	-	1	1	-	-
1999.....	1	-	1	-	-	3	2	-	1	1	-	-
White, both sexes												
2004.....	-	-	-	-	-	1	-	1	-	-	-	-
2003.....	-	1	-	-	-	-	1	1	1	-	-	-
2002.....	1	-	-	-	-	-	-	2	-	-	-	-
2001.....	-	1	-	-	-	1	3	-	-	-	-	-
2000.....	-	-	-	-	1	-	-	1	2	1	-	-
1999.....	1	-	1	-	-	1	1	-	-	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gonococcal infection (A54)												
All races, both sexes												
2004.....	-	-	-	-	-	-	-	-	1	-	-	-
2003.....	1	-	-	1	-	1	1	-	-	1	-	-
2002.....	-	-	2	1	-	-	-	-	-	-	-	-
2001.....	1	-	-	1	-	1	-	1	-	-	-	-
2000.....	1	-	-	1	1	-	2	-	-	-	-	-
1999.....	-	-	-	-	-	1	1	-	1	1	-	-
All races, male												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	2	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	1	-	-	-	1	-	-	-	-
2000.....	-	-	-	1	-	-	1	-	-	-	-	-
1999.....	-	-	-	-	-	1	-	-	-	-	-	-
All races, female												
2004.....	-	-	-	-	-	-	-	-	1	-	-	-
2003.....	1	-	-	1	-	1	1	-	-	1	-	-
2002.....	-	-	-	1	-	-	-	-	-	-	-	-
2001.....	1	-	-	-	-	1	-	-	-	-	-	-
2000.....	1	-	-	-	1	-	1	-	-	-	-	-
1999.....	-	-	-	-	-	-	1	-	1	1	-	-
White, both sexes												
2004.....	-	-	-	-	-	-	-	-	1	-	-	-
2003.....	1	-	-	1	-	1	1	-	-	1	-	-
2002.....	-	-	2	1	-	-	-	-	-	-	-	-
2001.....	1	-	-	-	-	-	-	1	-	-	-	-
2000.....	-	-	-	1	1	-	2	-	-	-	-	-
1999.....	-	-	-	-	-	1	1	-	1	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other infections with a predominately sexual mode of transmission (A55-A64)												
All races, both sexes												
2004.....	-	1	1	-	1	-	-	2	1	-	-	-
2003.....	-	-	-	-	-	-	3	3	1	2	-	-
2002.....	-	-	-	-	-	-	1	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	1	-	-
2000.....	1	-	-	-	1	-	3	2	-	-	-	-
1999.....	-	1	-	-	-	1	1	-	-	2	-	-
All races, male												
2004.....	-	1	1	-	1	-	-	-	1	-	-	-
2003.....	-	-	-	-	-	-	2	1	1	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	1	-	-
2000.....	1	-	-	-	1	-	2	1	-	-	-	-
1999.....	-	1	-	-	-	-	-	-	-	1	-	-
All races, female												
2004.....	-	-	-	-	-	-	-	2	-	-	-	-
2003.....	-	-	-	-	-	-	1	2	-	2	-	-
2002.....	-	-	-	-	-	-	1	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	1	1	-	-	-	-
1999.....	-	-	-	-	-	1	1	-	-	1	-	-
White, both sexes												
2004.....	-	1	1	-	1	-	-	2	1	-	-	-
2003.....	-	-	-	-	-	-	2	3	1	2	-	-
2002.....	-	-	-	-	-	-	1	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	1	-	3	2	-	-	-	-
1999.....	-	1	-	-	-	1	1	-	-	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other infections with a predominately sexual mode of transmission (A55-A64)

White, male

2004.....	-	1	1	-	1	-	-	-	1	-	-	-
2003.....	-	-	-	-	-	-	2	1	1	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	1	-	2	1	-	-	-	-
1999.....	-	1	-	-	-	-	-	-	-	-	-	-

White, female

2004.....	-	-	-	-	-	-	-	2	-	-	-	-
2003.....	-	-	-	-	-	-	-	2	-	2	-	-
2002.....	-	-	-	-	-	-	1	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	1	1	-	-	-	-
1999.....	-	-	-	-	-	1	1	-	-	1	-	-

Black, both sexes

2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	1	-	-
2000.....	1	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	1	-	-

Black, male

2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	1	-	-
2000.....	1	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other infections caused by spirochetes, chlamydia or rickettsia (A65-A79)												
All races, both sexes												
2004.....	1	3	-	-	1	2	1	2	-	-	-	-
2003.....	2	1	-	3	-	-	2	1	-	-	-	-
2002.....	2	1	2	-	2	-	2	3	-	1	-	-
2001.....	-	-	2	1	3	-	2	1	-	-	-	-
2000.....	-	1	-	-	2	2	-	-	-	1	-	-
1999.....	1	1	2	1	2	3	3	1	1	-	-	-
All races, male												
2004.....	1	2	-	-	-	1	-	1	-	-	-	-
2003.....	1	1	-	3	-	-	2	-	-	-	-	-
2002.....	-	1	-	-	1	-	2	1	-	-	-	-
2001.....	-	-	2	1	1	-	1	-	-	-	-	-
2000.....	-	1	-	-	1	-	-	-	-	-	-	-
1999.....	1	1	1	-	2	2	2	-	-	-	-	-
All races, female												
2004.....	-	1	-	-	1	1	1	1	-	-	-	-
2003.....	1	-	-	-	-	-	-	1	-	-	-	-
2002.....	2	-	2	-	1	-	-	2	-	1	-	-
2001.....	-	-	-	-	2	-	1	1	-	-	-	-
2000.....	-	-	-	-	1	2	-	-	-	1	-	-
1999.....	-	-	1	1	-	1	1	1	1	-	-	-
White, both sexes												
2004.....	1	3	-	-	-	2	1	1	-	-	-	-
2003.....	1	1	-	2	-	-	2	1	-	-	-	-
2002.....	2	1	2	-	2	-	2	3	-	-	-	-
2001.....	-	-	2	1	2	-	2	1	-	-	-	-
2000.....	-	1	-	-	2	2	-	-	-	1	-	-
1999.....	1	1	2	1	2	2	3	1	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other infections caused by spirochetes, chlamydia or rickettsia (A65-A79)

Asian or Pacific Islander 2/, both sexes

2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	1	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	1	-	-	-	-	-	-

Asian or Pacific Islander 2/, male

2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	1	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-

Asian or Pacific Islander 2/, female

2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	1	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Lyme disease (A69.2)												
All races, both sexes												
2004.....	1	1	-	-	-	1	1	-	-	-	-	-
2003.....	-	1	-	2	-	-	-	1	-	-	-	-
2002.....	-	1	1	-	1	-	1	1	-	-	-	-
2001.....	-	-	1	-	-	-	-	1	-	-	-	-
2000.....	-	-	-	-	1	2	-	-	-	1	-	-
1999.....	1	-	1	1	-	1	1	1	1	-	-	-
All races, male												
2004.....	1	1	-	-	-	-	-	-	-	-	-	-
2003.....	-	1	-	2	-	-	-	-	-	-	-	-
2002.....	-	1	-	-	1	-	1	1	-	-	-	-
2001.....	-	-	1	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	1	-	-	-	-	-	-	-
1999.....	1	-	1	-	-	1	-	-	-	-	-	-
All races, female												
2004.....	-	-	-	-	-	1	1	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	1	-	-	-	-
2002.....	-	-	1	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	1	-	-	-	-
2000.....	-	-	-	-	-	2	-	-	-	1	-	-
1999.....	-	-	-	1	-	-	1	1	1	-	-	-
White, both sexes												
2004.....	1	1	-	-	-	1	1	-	-	-	-	-
2003.....	-	1	-	1	-	-	-	1	-	-	-	-
2002.....	-	1	1	-	1	-	1	1	-	-	-	-
2001.....	-	-	1	-	-	-	-	1	-	-	-	-
2000.....	-	-	-	-	1	2	-	-	-	1	-	-
1999.....	1	-	1	1	-	1	1	1	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

All other infections caused by spirochetes, chlamydia or rickettsia (A65-A68,A69.0-A69.1,A69.8-A69.9,A70-A74,A77-A79)

All races, both sexes

2004.....	-	2	-	-	-	1	-	2	-	-	-	-
2003.....	2	-	-	1	-	-	2	-	-	-	-	-
2002.....	2	-	1	-	1	-	1	2	-	1	-	-
2001.....	-	-	1	1	3	-	2	-	-	-	-	-
2000.....	-	1	-	-	1	-	-	-	-	-	-	-
1999.....	-	1	1	-	2	2	2	-	-	-	-	-

All races, male

2004.....	-	1	-	-	-	1	-	1	-	-	-	-
2003.....	1	-	-	1	-	-	2	-	-	-	-	-
2002.....	-	-	-	-	-	-	1	-	-	-	-	-
2001.....	-	-	1	1	1	-	1	-	-	-	-	-
2000.....	-	1	-	-	-	-	-	-	-	-	-	-
1999.....	-	1	-	-	2	1	2	-	-	-	-	-

All races, female

2004.....	-	1	-	-	-	-	-	1	-	-	-	-
2003.....	1	-	-	-	-	-	-	-	-	-	-	-
2002.....	2	-	1	-	1	-	-	2	-	1	-	-
2001.....	-	-	-	-	2	-	1	-	-	-	-	-
2000.....	-	-	-	-	1	-	-	-	-	-	-	-
1999.....	-	-	1	-	-	1	-	-	-	-	-	-

White, both sexes

2004.....	-	2	-	-	-	1	-	1	-	-	-	-
2003.....	1	-	-	1	-	-	2	-	-	-	-	-
2002.....	2	-	1	-	1	-	1	2	-	-	-	-
2001.....	-	-	1	1	2	-	2	-	-	-	-	-
2000.....	-	1	-	-	1	-	-	-	-	-	-	-
1999.....	-	1	1	-	2	1	2	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

All other infections caused by spirochetes, chlamydia or rickettsia (A65-A68,A69.0-A69.1,A69.8-A69.9,A70-A74,A77-A79)

Asian or Pacific Islander 2/, both sexes

2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	1	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	1	-	-	-	-	-	-

Asian or Pacific Islander 2/, male

2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	1	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-

Asian or Pacific Islander 2/, female

2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	1	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Viral diseases (A80-B34)												
All races, both sexes												
2004.....	3,192	1,931	1,058	746	552	478	366	217	97	47	17	4
2003.....	3,079	1,803	976	721	688	563	406	254	116	45	15	3
2002.....	2,991	1,589	961	753	678	581	409	286	118	51	7	4
2001.....	2,622	1,456	1,011	770	671	545	362	201	117	40	8	3
2000.....	2,495	1,356	935	754	710	603	401	228	115	46	17	2
1999.....	2,178	1,300	797	780	718	587	424	276	146	44	16	1
All races, male												
2004.....	2,392	1,485	747	486	306	240	142	97	29	8	2	4
2003.....	2,370	1,361	680	441	403	284	197	97	36	10	4	3
2002.....	2,344	1,186	666	474	355	308	182	109	41	13	-	4
2001.....	2,064	1,103	663	443	355	254	171	79	31	11	-	3
2000.....	1,953	1,025	636	448	370	327	182	90	32	9	2	2
1999.....	1,720	981	533	471	382	279	196	108	44	7	-	1
All races, female												
2004.....	800	446	311	260	246	238	224	120	68	39	15	-
2003.....	709	442	296	280	285	279	209	157	80	35	11	-
2002.....	647	403	295	279	323	273	227	177	77	38	7	-
2001.....	558	353	348	327	316	291	191	122	86	29	8	-
2000.....	542	331	299	306	340	276	219	138	83	37	15	-
1999.....	458	319	264	309	336	308	228	168	102	37	16	-
White, both sexes												
2004.....	1,833	1,142	636	487	396	389	310	190	90	45	16	3
2003.....	1,774	1,059	582	460	515	460	339	230	111	42	14	2
2002.....	1,698	963	596	483	500	464	355	263	110	47	7	3
2001.....	1,473	900	605	516	506	448	307	182	106	40	8	-
2000.....	1,385	800	577	517	530	505	348	204	106	42	16	2
1999.....	1,244	740	510	534	533	490	380	244	137	41	14	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Viral diseases (A80-B34)												
White, male												
2004.....	1,440	895	447	316	217	200	119	82	28	8	2	3
2003.....	1,417	813	404	275	297	226	165	87	34	10	4	2
2002.....	1,354	729	400	302	256	246	157	100	38	9	-	3
2001.....	1,220	692	390	299	274	211	145	75	28	11	-	-
2000.....	1,124	610	382	295	264	273	156	80	31	8	2	2
1999.....	1,015	558	348	319	283	235	180	92	39	6	-	-
White, female												
2004.....	393	247	189	171	179	189	191	108	62	37	14	-
2003.....	357	246	178	185	218	234	174	143	77	32	10	-
2002.....	344	234	196	181	244	218	198	163	72	38	7	-
2001.....	253	208	215	217	232	237	162	107	78	29	8	-
2000.....	261	190	195	222	266	232	192	124	75	34	14	-
1999.....	229	182	162	215	250	255	200	152	98	35	14	-
Black, both sexes												
2004.....	1,293	740	387	218	116	63	40	20	5	1	-	1
2003.....	1,254	691	360	218	133	77	50	19	4	3	1	1
2002.....	1,239	572	316	228	137	95	36	14	8	3	-	1
2001.....	1,092	515	365	202	118	74	37	13	7	-	-	3
2000.....	1,043	495	304	190	145	77	40	18	6	3	1	-
1999.....	879	516	247	204	145	71	31	25	6	3	2	1
Black, male												
2004.....	908	556	281	152	74	28	17	11	-	-	-	1
2003.....	919	520	256	148	87	41	22	10	2	-	-	1
2002.....	948	424	231	159	83	53	17	4	3	3	-	1
2001.....	798	385	253	124	65	37	17	1	2	-	-	3
2000.....	788	376	222	130	92	42	22	7	1	-	-	-
1999.....	666	391	169	128	85	33	11	13	3	1	-	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Viral diseases (A80-B34)												
Black, female												
2004.....	385	184	106	66	42	35	23	9	5	1	-	-
2003.....	335	171	104	70	46	36	28	9	2	3	1	-
2002.....	291	148	85	69	54	42	19	10	5	-	-	-
2001.....	294	130	112	78	53	37	20	12	5	-	-	-
2000.....	255	119	82	60	53	35	18	11	5	3	1	-
1999.....	213	125	78	76	60	38	20	12	3	2	2	-
American Indian 1/, both sexes												
2004.....	23	12	11	6	1	2	1	1	-	-	-	-
2003.....	20	14	9	7	3	3	-	-	-	-	-	-
2002.....	11	14	4	4	3	1	1	-	-	-	-	-
2001.....	21	2	5	7	4	2	1	-	-	-	-	-
2000.....	20	5	7	6	3	2	2	-	1	-	-	-
1999.....	11	3	5	4	3	2	3	1	-	-	-	-
American Indian 1/, male												
2004.....	10	8	5	2	-	-	1	-	-	-	-	-
2003.....	13	10	4	3	1	3	-	-	-	-	-	-
2002.....	7	7	2	2	1	-	-	-	-	-	-	-
2001.....	18	1	4	2	1	-	1	-	-	-	-	-
2000.....	13	1	4	3	1	1	1	-	-	-	-	-
1999.....	9	1	2	3	1	-	2	-	-	-	-	-
American Indian 1/, female												
2004.....	13	4	6	4	1	2	-	1	-	-	-	-
2003.....	7	4	5	4	2	-	-	-	-	-	-	-
2002.....	4	7	2	2	2	1	1	-	-	-	-	-
2001.....	3	1	1	5	3	2	-	-	-	-	-	-
2000.....	7	4	3	3	2	1	1	-	1	-	-	-
1999.....	2	2	3	1	2	2	1	1	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Viral diseases (A80-B34)												
Asian or Pacific Islander 2/, both sexes												
2004.....	43	37	24	35	39	24	15	6	2	1	1	-
2003.....	31	39	25	36	37	23	17	5	1	-	-	-
2002.....	43	40	45	38	38	21	17	9	-	1	-	-
2001.....	36	39	36	45	43	21	17	6	4	-	-	-
2000.....	47	56	47	41	32	19	11	6	2	1	-	-
1999.....	44	41	35	38	37	24	10	6	3	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	34	26	14	16	15	12	5	4	1	-	-	-
2003.....	21	18	16	15	18	14	10	-	-	-	-	-
2002.....	35	26	33	11	15	9	8	5	-	1	-	-
2001.....	28	25	16	18	15	6	8	3	1	-	-	-
2000.....	28	38	28	20	13	11	3	3	-	1	-	-
1999.....	30	31	14	21	13	11	3	3	2	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	9	11	10	19	24	12	10	2	1	1	1	-
2003.....	10	21	9	21	19	9	7	5	1	-	-	-
2002.....	8	14	12	27	23	12	9	4	-	-	-	-
2001.....	8	14	20	27	28	15	9	3	3	-	-	-
2000.....	19	18	19	21	19	8	8	3	2	-	-	-
1999.....	14	10	21	17	24	13	7	3	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Arthropod-borne viral encephalitis (A83-A84,A85.2)												
All races, both sexes												
2004.....	-	-	1	2	1	-	-	-	-	-	-	-
2003.....	1	-	3	-	2	4	1	-	-	-	-	-
2002.....	2	-	1	-	1	1	1	1	-	-	-	-
2001.....	1	-	-	-	5	1	1	-	-	-	-	-
2000.....	-	-	-	-	-	1	-	-	-	-	-	-
1999.....	-	-	-	1	1	-	-	1	-	-	-	-
All races, male												
2004.....	-	-	1	2	1	-	-	-	-	-	-	-
2003.....	1	-	2	-	2	2	-	-	-	-	-	-
2002.....	1	-	1	-	1	1	-	-	-	-	-	-
2001.....	-	-	-	-	4	1	1	-	-	-	-	-
2000.....	-	-	-	-	-	1	-	-	-	-	-	-
1999.....	-	-	-	1	1	-	-	1	-	-	-	-
All races, female												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	1	-	-	2	1	-	-	-	-	-
2002.....	1	-	-	-	-	-	1	1	-	-	-	-
2001.....	1	-	-	-	1	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes												
2004.....	-	-	1	2	-	-	-	-	-	-	-	-
2003.....	-	-	3	-	2	4	1	-	-	-	-	-
2002.....	1	-	-	-	-	1	1	1	-	-	-	-
2001.....	1	-	-	-	4	1	-	-	-	-	-	-
2000.....	-	-	-	-	-	1	-	-	-	-	-	-
1999.....	-	-	-	1	1	-	-	1	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Herpesviral (herpes simplex) infections (B00)												
All races, both sexes												
2004.....	3	14	7	16	18	27	32	21	8	2	-	-
2003.....	9	10	10	11	21	18	24	29	3	2	-	-
2002.....	4	7	4	8	35	19	28	22	3	-	-	-
2001.....	5	5	17	16	24	25	23	11	9	1	-	-
2000.....	4	7	6	5	16	24	23	11	4	2	1	-
1999.....	7	7	8	13	21	22	13	9	8	2	1	-
All races, male												
2004.....	1	10	4	8	10	11	9	11	2	-	-	-
2003.....	6	4	4	5	8	9	9	6	-	-	-	-
2002.....	1	4	3	3	13	7	10	7	2	-	-	-
2001.....	3	3	8	9	15	9	11	6	3	-	-	-
2000.....	3	3	1	2	6	9	15	6	2	-	-	-
1999.....	3	2	3	5	8	9	4	4	2	1	-	-
All races, female												
2004.....	2	4	3	8	8	16	23	10	6	2	-	-
2003.....	3	6	6	6	13	9	15	23	3	2	-	-
2002.....	3	3	1	5	22	12	18	15	1	-	-	-
2001.....	2	2	9	7	9	16	12	5	6	1	-	-
2000.....	1	4	5	3	10	15	8	5	2	2	1	-
1999.....	4	5	5	8	13	13	9	5	6	1	1	-
White, both sexes												
2004.....	3	12	7	14	15	27	31	19	8	2	-	-
2003.....	6	9	9	11	21	16	22	27	3	2	-	-
2002.....	4	5	4	6	34	18	25	22	3	-	-	-
2001.....	4	5	16	16	20	23	20	11	9	1	-	-
2000.....	4	6	4	5	14	24	22	8	4	2	1	-
1999.....	5	3	8	13	20	21	11	8	7	2	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Herpesviral (herpes simplex) infections (B00)

White, male

2004.....	1	8	4	7	7	11	9	9	2	-	-	-
2003.....	4	4	4	5	8	8	9	6	-	-	-	-
2002.....	1	2	3	3	13	7	9	7	2	-	-	-
2001.....	3	3	8	9	13	7	8	6	3	-	-	-
2000.....	3	2	1	2	5	9	14	4	2	-	-	-
1999.....	3	-	3	5	8	8	2	4	2	1	-	-

White, female

2004.....	2	4	3	7	8	16	22	10	6	2	-	-
2003.....	2	5	5	6	13	8	13	21	3	2	-	-
2002.....	3	3	1	3	21	11	16	15	1	-	-	-
2001.....	1	2	8	7	7	16	12	5	6	1	-	-
2000.....	1	4	3	3	9	15	8	4	2	2	1	-
1999.....	2	3	5	8	12	13	9	4	5	1	-	-

Black, both sexes

2004.....	-	1	-	2	-	-	1	2	-	-	-	-
2003.....	2	1	-	-	-	2	2	2	-	-	-	-
2002.....	-	2	-	2	1	-	3	-	-	-	-	-
2001.....	1	-	1	-	4	-	2	-	-	-	-	-
2000.....	-	-	1	-	2	-	1	3	-	-	-	-
1999.....	1	4	-	-	1	1	2	-	1	-	1	-

Black, male

2004.....	-	1	-	1	-	-	-	2	-	-	-	-
2003.....	1	-	-	-	-	1	-	-	-	-	-	-
2002.....	-	2	-	-	-	-	1	-	-	-	-	-
2001.....	-	-	-	-	2	-	2	-	-	-	-	-
2000.....	-	-	-	-	1	-	1	2	-	-	-	-
1999.....	-	2	-	-	-	1	2	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Herpesviral (herpes simplex) infections (B00)												
Asian or Pacific Islander 2/, both sexes												
2004.....	-	1	-	-	3	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	1	-	-	-	-	-	-
2001.....	-	-	-	-	-	2	1	-	-	-	-	-
2000.....	-	1	1	-	-	-	-	-	-	-	-	-
1999.....	1	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	-	1	-	-	3	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	2	1	-	-	-	-	-
2000.....	-	1	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	1	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	1	-	-	-	-	-	-	-	-	-
1999.....	1	-	-	-	-	-	-	1	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Zoster (herpes zoster)(B02)												
All races, both sexes												
2004.....	-	2	2	4	14	11	22	36	27	20	9	-
2003.....	1	3	6	5	2	11	21	28	48	18	9	-
2002.....	1	2	2	7	8	10	17	32	29	14	2	-
2001.....	5	-	2	3	10	11	18	23	28	17	4	-
2000.....	-	3	6	2	11	19	16	30	21	12	8	-
1999.....	-	4	1	2	9	16	29	34	26	19	6	-
All races, male												
2004.....	-	1	1	3	10	5	5	18	8	4	1	-
2003.....	-	1	2	2	1	4	12	9	12	2	3	-
2002.....	1	1	1	5	5	5	9	8	9	4	-	-
2001.....	3	-	2	2	5	3	5	6	5	5	-	-
2000.....	-	1	2	1	3	13	6	8	4	1	2	-
1999.....	-	2	1	2	5	5	13	9	3	3	-	-
All races, female												
2004.....	-	1	1	1	4	6	17	18	19	16	8	-
2003.....	1	2	4	3	1	7	9	19	36	16	6	-
2002.....	-	1	1	2	3	5	8	24	20	10	2	-
2001.....	2	-	-	1	5	8	13	17	23	12	4	-
2000.....	-	2	4	1	8	6	10	22	17	11	6	-
1999.....	-	2	-	-	4	11	16	25	23	16	6	-
White, both sexes												
2004.....	-	2	2	4	11	9	21	33	26	20	9	-
2003.....	-	2	5	4	2	9	19	28	47	17	8	-
2002.....	1	2	1	6	6	8	15	30	29	14	2	-
2001.....	4	-	2	3	9	10	18	23	27	17	4	-
2000.....	-	1	5	2	7	18	15	28	20	12	8	-
1999.....	-	3	-	2	8	16	28	33	26	18	6	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Zoster (herpes zoster)(B02)												
Black, female												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	1	-	-	1	-	1	1	-	-	1	1	-
2002.....	-	-	-	-	-	2	-	1	-	-	-	-
2001.....	1	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	2	-	-	3	-	-	2	1	-	-	-
1999.....	-	-	-	-	-	-	1	1	-	1	-	-
American Indian 1/, both sexes												
2004.....	-	-	-	-	-	1	1	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	1	1	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	1	-	-	-	-	-	-	-
1999.....	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, male												
2004.....	-	-	-	-	-	-	1	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	1	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, female												
2004.....	-	-	-	-	-	1	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	1	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	1	-	-	-	-	-	-	-
1999.....	-	-	-	-	1	-	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Viral hepatitis (B15-B19)												
All races, both sexes												
2004.....	1,270	800	444	328	306	265	199	92	33	4	-	-
2003.....	1,206	709	376	330	368	323	220	87	24	4	-	-
2002.....	1,139	644	428	369	393	352	235	136	34	7	-	-
2001.....	959	586	428	407	429	375	238	120	41	5	-	1
2000.....	850	492	412	410	435	401	259	123	35	10	-	1
1999.....	635	438	348	427	454	389	262	145	45	5	2	-
All races, male												
2004.....	944	588	272	192	142	119	70	35	10	-	-	-
2003.....	903	506	223	169	183	147	102	30	8	2	-	-
2002.....	868	448	254	193	180	174	96	56	11	3	-	-
2001.....	737	406	237	207	189	156	107	42	13	2	-	1
2000.....	638	338	239	202	194	198	113	55	11	4	-	1
1999.....	487	298	193	217	209	185	132	61	20	1	-	-
All races, female												
2004.....	326	212	172	136	164	146	129	57	23	4	-	-
2003.....	303	203	153	161	185	176	118	57	16	2	-	-
2002.....	271	196	174	176	213	178	139	80	23	4	-	-
2001.....	222	180	191	200	240	219	131	78	28	3	-	-
2000.....	212	154	173	208	241	203	146	68	24	6	-	-
1999.....	148	140	155	210	245	204	130	84	25	4	2	-
White, both sexes												
2004.....	1,002	594	340	257	234	218	157	76	30	4	-	-
2003.....	955	517	278	244	289	262	185	74	21	3	-	-
2002.....	920	487	317	278	311	298	205	121	30	4	-	-
2001.....	734	447	331	314	341	322	205	104	34	5	-	-
2000.....	635	350	315	320	358	350	222	110	30	9	-	1
1999.....	481	323	269	330	364	326	235	124	40	4	2	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Viral hepatitis (B15-B19)												
White, male												
2004.....	757	438	206	149	113	101	54	28	9	-	-	-
2003.....	723	378	171	130	145	116	89	27	7	2	-	-
2002.....	705	342	185	154	148	152	87	51	10	-	-	-
2001.....	580	319	189	170	161	143	98	41	10	2	-	-
2000.....	487	246	186	156	161	175	98	49	11	3	-	1
1999.....	371	224	160	177	177	159	126	52	17	1	-	-
White, female												
2004.....	245	156	134	108	121	117	103	48	21	4	-	-
2003.....	232	139	107	114	144	146	96	47	14	1	-	-
2002.....	215	145	132	124	163	146	118	70	20	4	-	-
2001.....	154	128	142	144	180	179	107	63	24	3	-	-
2000.....	148	104	129	164	197	175	124	61	19	6	-	-
1999.....	110	99	109	153	187	167	109	72	23	3	2	-
Black, both sexes												
2004.....	229	171	80	41	39	23	27	10	2	-	-	-
2003.....	219	148	74	49	43	38	21	8	2	1	-	-
2002.....	189	115	72	57	46	35	15	7	4	2	-	-
2001.....	188	103	67	47	46	35	19	12	4	-	-	1
2000.....	164	93	50	47	47	32	25	9	2	-	-	-
1999.....	118	75	45	60	53	38	17	17	2	1	-	-
Black, male												
2004.....	163	126	53	28	20	6	11	4	-	-	-	-
2003.....	161	106	39	22	21	17	5	3	1	-	-	-
2002.....	141	83	41	32	18	14	3	-	1	2	-	-
2001.....	128	65	36	18	16	9	4	-	2	-	-	1
2000.....	120	63	25	26	22	13	12	4	-	-	-	-
1999.....	91	45	21	21	19	15	2	8	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Viral hepatitis (B15-B19)												
Black, female												
2004.....	66	45	27	13	19	17	16	6	2	-	-	-
2003.....	58	42	35	27	22	21	16	5	1	1	-	-
2002.....	48	32	31	25	28	21	12	7	3	-	-	-
2001.....	60	38	31	29	30	26	15	12	2	-	-	-
2000.....	44	30	25	21	25	19	13	5	2	-	-	-
1999.....	27	30	24	39	34	23	15	9	1	1	-	-
American Indian 1/, both sexes												
2004.....	13	5	6	4	1	1	-	1	-	-	-	-
2003.....	10	9	7	3	2	1	-	-	-	-	-	-
2002.....	8	10	3	1	3	-	-	-	-	-	-	-
2001.....	12	1	4	5	3	2	-	-	-	-	-	-
2000.....	13	4	6	4	1	1	2	-	1	-	-	-
1999.....	6	2	2	2	1	2	3	1	-	-	-	-
American Indian 1/, male												
2004.....	4	4	3	2	-	-	-	-	-	-	-	-
2003.....	5	7	3	2	1	1	-	-	-	-	-	-
2002.....	6	5	2	1	1	-	-	-	-	-	-	-
2001.....	10	-	3	2	-	-	-	-	-	-	-	-
2000.....	9	1	4	1	-	-	1	-	-	-	-	-
1999.....	4	1	1	1	-	-	2	-	-	-	-	-
American Indian 1/, female												
2004.....	9	1	3	2	1	1	-	1	-	-	-	-
2003.....	5	2	4	1	1	-	-	-	-	-	-	-
2002.....	2	5	1	-	2	-	-	-	-	-	-	-
2001.....	2	1	1	3	3	2	-	-	-	-	-	-
2000.....	4	3	2	3	1	1	1	-	1	-	-	-
1999.....	2	1	1	1	1	2	1	1	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Viral hepatitis (B15-B19)												
Asian or Pacific Islander 2/, both sexes												
2004.....	26	30	18	26	32	23	15	5	1	-	-	-
2003.....	22	35	17	34	34	22	14	5	1	-	-	-
2002.....	22	32	36	33	33	19	15	8	-	1	-	-
2001.....	25	35	26	41	39	16	14	4	3	-	-	-
2000.....	38	45	41	39	29	18	10	4	2	1	-	-
1999.....	30	38	32	35	36	23	7	3	3	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	20	20	10	13	9	12	5	3	1	-	-	-
2003.....	14	15	10	15	16	13	8	-	-	-	-	-
2002.....	16	18	26	6	13	8	6	5	-	1	-	-
2001.....	19	22	9	17	12	4	5	1	1	-	-	-
2000.....	22	28	24	19	11	10	2	2	-	1	-	-
1999.....	21	28	11	18	13	11	2	1	2	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	6	10	8	13	23	11	10	2	-	-	-	-
2003.....	8	20	7	19	18	9	6	5	1	-	-	-
2002.....	6	14	10	27	20	11	9	3	-	-	-	-
2001.....	6	13	17	24	27	12	9	3	2	-	-	-
2000.....	16	17	17	20	18	8	8	2	2	-	-	-
1999.....	9	10	21	17	23	12	5	2	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Human immunodeficiency virus (HIV) disease (B20-B24)												
All races, both sexes												
2004.....	1,868	1,038	524	308	128	79	27	11	2	2	1	4
2003.....	1,806	1,017	500	277	161	71	24	11	3	-	-	3
2002.....	1,794	885	462	275	126	71	25	8	3	1	-	4
2001.....	1,609	821	496	258	126	68	20	5	3	2	-	2
2000.....	1,591	797	442	250	147	69	16	4	3	2	-	1
1999.....	1,467	777	372	261	137	52	18	7	5	1	1	1
All races, male												
2004.....	1,416	840	422	235	98	53	18	4	-	1	-	4
2003.....	1,425	816	399	209	127	55	20	10	2	-	-	3
2002.....	1,446	710	373	223	100	53	18	4	-	1	-	4
2001.....	1,299	673	378	187	105	55	18	4	1	2	-	2
2000.....	1,288	655	358	199	114	54	11	1	1	-	-	1
1999.....	1,198	646	302	211	106	39	11	6	1	1	-	1
All races, female												
2004.....	452	198	102	73	30	26	9	7	2	1	1	-
2003.....	381	201	101	68	34	16	4	1	1	-	-	-
2002.....	348	175	89	52	26	18	7	4	3	-	-	-
2001.....	310	148	118	71	21	13	2	1	2	-	-	-
2000.....	303	142	84	51	33	15	5	3	2	2	-	-
1999.....	269	131	70	50	31	13	7	1	4	-	1	-
White, both sexes												
2004.....	787	467	218	131	60	42	19	7	1	2	1	3
2003.....	771	474	213	111	78	41	7	5	3	-	-	2
2002.....	728	425	218	106	45	30	12	5	3	1	-	3
2001.....	695	406	194	102	61	31	6	3	2	2	-	-
2000.....	706	391	197	110	56	33	7	2	1	-	-	1
1999.....	700	346	173	119	52	25	12	3	4	-	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Human immunodeficiency virus (HIV) disease (B20-B24)												
White, male												
2004.....	660	407	193	110	49	32	12	2	-	1	-	3
2003.....	663	399	181	87	66	35	7	4	2	-	-	2
2002.....	622	365	181	93	41	21	8	3	-	1	-	3
2001.....	616	350	157	82	58	27	6	2	1	2	-	-
2000.....	613	336	163	96	44	28	5	-	-	-	-	1
1999.....	613	305	156	103	42	22	7	2	-	-	-	-
White, female												
2004.....	127	60	25	21	11	10	7	5	1	1	1	-
2003.....	108	75	32	24	12	6	-	1	1	-	-	-
2002.....	106	60	37	13	4	9	4	2	3	-	-	-
2001.....	79	56	37	20	3	4	-	1	1	-	-	-
2000.....	93	55	34	14	12	5	2	2	1	-	-	-
1999.....	87	41	17	16	10	3	5	1	4	-	1	-
Black, both sexes												
2004.....	1,057	560	300	170	65	37	8	4	1	-	-	1
2003.....	1,021	535	282	164	83	29	17	6	-	-	-	1
2002.....	1,043	450	238	165	78	41	13	3	-	-	-	1
2001.....	897	410	293	151	63	37	13	1	1	-	-	2
2000.....	872	395	241	137	87	35	9	2	2	2	-	-
1999.....	753	429	196	139	83	27	6	3	1	1	-	1
Black, male												
2004.....	739	425	225	122	47	21	6	2	-	-	-	1
2003.....	750	411	214	122	61	19	13	6	-	-	-	1
2002.....	804	336	187	126	58	32	10	1	-	-	-	1
2001.....	667	319	214	104	46	28	11	1	-	-	-	2
2000.....	666	310	191	101	67	25	6	1	1	-	-	-
1999.....	573	339	144	105	63	17	4	3	1	1	-	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Human immunodeficiency virus (HIV) disease (B20-B24)												
Asian or Pacific Islander 2/, both sexes												
2004.....	14	5	3	6	3	-	-	-	-	-	-	-
2003.....	8	4	4	1	-	-	-	-	-	-	-	-
2002.....	20	7	6	4	3	-	-	-	-	-	-	-
2001.....	9	4	8	3	2	-	-	1	-	-	-	-
2000.....	7	10	4	1	3	-	-	-	-	-	-	-
1999.....	9	2	1	2	1	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	11	5	2	3	2	-	-	-	-	-	-	-
2003.....	6	3	3	-	-	-	-	-	-	-	-	-
2002.....	19	7	5	4	1	-	-	-	-	-	-	-
2001.....	8	3	6	1	1	-	-	1	-	-	-	-
2000.....	6	9	4	-	2	-	-	-	-	-	-	-
1999.....	7	2	1	2	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	3	-	1	3	1	-	-	-	-	-	-	-
2003.....	2	1	1	1	-	-	-	-	-	-	-	-
2002.....	1	-	1	-	2	-	-	-	-	-	-	-
2001.....	1	1	2	2	1	-	-	-	-	-	-	-
2000.....	1	1	-	1	1	-	-	-	-	-	-	-
1999.....	2	-	-	-	1	-	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Human immunodeficiency virus (HIV) disease resulting in infectious and parasitic diseases (B20)												
All races, both sexes												
2004.....	712	392	162	123	46	28	8	2	1	-	-	2
2003.....	733	380	183	89	56	14	8	3	2	-	-	1
2002.....	731	338	163	98	39	26	4	2	1	-	-	-
2001.....	629	314	170	82	52	21	6	1	1	-	-	1
2000.....	607	273	155	90	51	26	7	1	2	-	-	1
1999.....	603	255	122	83	51	26	6	3	-	-	1	1
All races, male												
2004.....	520	313	128	93	34	22	6	1	-	-	-	2
2003.....	575	294	143	69	40	12	8	3	2	-	-	1
2002.....	594	268	119	76	33	20	4	2	-	-	-	-
2001.....	503	253	124	55	42	19	6	1	-	-	-	1
2000.....	501	212	124	75	42	18	5	-	-	-	-	1
1999.....	490	213	96	69	37	22	6	2	-	-	-	1
All races, female												
2004.....	192	79	34	30	12	6	2	1	1	-	-	-
2003.....	158	86	40	20	16	2	-	-	-	-	-	-
2002.....	137	70	44	22	6	6	-	-	1	-	-	-
2001.....	126	61	46	27	10	2	-	-	1	-	-	-
2000.....	106	61	31	15	9	8	2	1	2	-	-	-
1999.....	113	42	26	14	14	4	-	1	-	-	1	-
White, both sexes												
2004.....	310	171	62	55	18	16	4	1	-	-	-	2
2003.....	309	178	80	41	25	7	2	1	2	-	-	1
2002.....	297	170	87	39	17	10	2	2	1	-	-	-
2001.....	267	160	71	32	24	13	3	1	-	-	-	-
2000.....	289	135	72	43	17	11	2	-	1	-	-	1
1999.....	298	123	60	40	18	16	3	1	-	-	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Human immunodeficiency virus (HIV) disease resulting in infectious and parasitic diseases (B20)												
White, male												
2004.....	244	152	55	46	15	12	4	-	-	-	-	2
2003.....	259	140	65	32	20	6	2	1	2	-	-	1
2002.....	249	144	63	36	16	6	2	2	-	-	-	-
2001.....	231	135	53	23	22	11	3	1	-	-	-	-
2000.....	253	109	58	38	14	9	2	-	-	-	-	1
1999.....	255	105	54	34	14	15	3	-	-	-	-	-
White, female												
2004.....	66	19	7	9	3	4	-	1	-	-	-	-
2003.....	50	38	15	9	5	1	-	-	-	-	-	-
2002.....	48	26	24	3	1	4	-	-	1	-	-	-
2001.....	36	25	18	9	2	2	-	-	-	-	-	-
2000.....	36	26	14	5	3	2	-	-	1	-	-	-
1999.....	43	18	6	6	4	1	-	1	-	-	1	-
Black, both sexes												
2004.....	391	216	99	66	27	12	4	1	1	-	-	-
2003.....	416	197	101	48	31	7	6	2	-	-	-	-
2002.....	426	162	76	57	20	16	2	-	-	-	-	-
2001.....	351	152	95	48	27	8	2	-	1	-	-	1
2000.....	309	133	81	46	33	15	5	1	1	-	-	-
1999.....	298	132	62	42	32	10	3	1	-	-	-	1
Black, male												
2004.....	266	158	72	46	18	10	2	1	-	-	-	-
2003.....	309	150	76	37	20	6	6	2	-	-	-	-
2002.....	338	119	56	38	16	14	2	-	-	-	-	-
2001.....	261	117	68	32	20	8	2	-	-	-	-	1
2000.....	241	99	64	36	27	9	3	-	-	-	-	-
1999.....	229	108	42	34	23	7	3	1	-	-	-	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Human immunodeficiency virus (HIV) disease resulting in infectious and parasitic diseases (B20)												
Asian or Pacific Islander 2/, both sexes												
2004.....	6	2	1	2	1	-	-	-	-	-	-	-
2003.....	3	2	2	-	-	-	-	-	-	-	-	-
2002.....	8	3	-	2	2	-	-	-	-	-	-	-
2001.....	6	2	4	2	1	-	-	-	-	-	-	-
2000.....	4	5	2	-	-	-	-	-	-	-	-	-
1999.....	3	-	-	-	1	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	5	2	1	1	1	-	-	-	-	-	-	-
2003.....	2	2	2	-	-	-	-	-	-	-	-	-
2002.....	7	3	-	2	1	-	-	-	-	-	-	-
2001.....	6	1	3	-	-	-	-	-	-	-	-	-
2000.....	4	4	2	-	-	-	-	-	-	-	-	-
1999.....	2	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	1	-	-	1	-	-	-	-	-	-	-	-
2003.....	1	-	-	-	-	-	-	-	-	-	-	-
2002.....	1	-	-	-	1	-	-	-	-	-	-	-
2001.....	-	1	1	2	1	-	-	-	-	-	-	-
2000.....	-	1	-	-	-	-	-	-	-	-	-	-
1999.....	1	-	-	-	1	-	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Human immunodeficiency virus (HIV) disease resulting in malignant neoplasms (B21)												
All races, both sexes												
2004.....	147	93	46	24	15	3	1	1	-	-	-	-
2003.....	121	88	38	27	12	7	1	1	-	-	-	-
2002.....	129	61	38	19	16	3	2	1	-	-	-	-
2001.....	113	80	49	21	11	5	-	-	-	-	-	-
2000.....	120	66	38	20	12	3	1	-	-	-	-	-
1999.....	122	84	28	19	12	3	-	-	-	-	-	-
All races, male												
2004.....	123	84	40	19	13	2	1	-	-	-	-	-
2003.....	96	71	30	20	12	5	1	-	-	-	-	-
2002.....	106	52	31	19	14	2	2	-	-	-	-	-
2001.....	98	65	43	16	8	5	-	-	-	-	-	-
2000.....	103	62	33	19	10	3	-	-	-	-	-	-
1999.....	101	72	25	16	9	3	-	-	-	-	-	-
All races, female												
2004.....	24	9	6	5	2	1	-	1	-	-	-	-
2003.....	25	17	8	7	-	2	-	1	-	-	-	-
2002.....	23	9	7	-	2	1	-	1	-	-	-	-
2001.....	15	15	6	5	3	-	-	-	-	-	-	-
2000.....	17	4	5	1	2	-	1	-	-	-	-	-
1999.....	21	12	3	3	3	-	-	-	-	-	-	-
White, both sexes												
2004.....	71	53	27	11	8	3	1	1	-	-	-	-
2003.....	64	47	21	14	7	5	1	1	-	-	-	-
2002.....	62	40	21	11	9	2	1	1	-	-	-	-
2001.....	69	40	25	10	6	2	-	-	-	-	-	-
2000.....	69	40	23	10	6	2	1	-	-	-	-	-
1999.....	73	53	14	10	5	1	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Human immunodeficiency virus (HIV) disease resulting in malignant neoplasms (B21)												
White, male												
2004.....	65	49	24	10	7	2	1	-	-	-	-	-
2003.....	56	41	18	11	7	5	1	-	-	-	-	-
2002.....	55	36	18	11	8	2	1	-	-	-	-	-
2001.....	63	35	22	8	6	2	-	-	-	-	-	-
2000.....	64	39	20	10	5	2	-	-	-	-	-	-
1999.....	64	51	11	10	4	1	-	-	-	-	-	-
White, female												
2004.....	6	4	3	1	1	1	-	1	-	-	-	-
2003.....	8	6	3	3	-	-	-	1	-	-	-	-
2002.....	7	4	3	-	1	-	-	1	-	-	-	-
2001.....	6	5	3	2	-	-	-	-	-	-	-	-
2000.....	5	1	3	-	1	-	1	-	-	-	-	-
1999.....	9	2	3	-	1	-	-	-	-	-	-	-
Black, both sexes												
2004.....	73	39	18	13	7	-	-	-	-	-	-	-
2003.....	56	41	17	13	5	2	-	-	-	-	-	-
2002.....	64	20	17	8	7	1	1	-	-	-	-	-
2001.....	44	39	24	11	5	3	-	-	-	-	-	-
2000.....	50	26	14	10	6	1	-	-	-	-	-	-
1999.....	48	31	14	9	7	2	-	-	-	-	-	-
Black, male												
2004.....	57	34	15	9	6	-	-	-	-	-	-	-
2003.....	40	30	12	9	5	-	-	-	-	-	-	-
2002.....	48	15	13	8	6	-	1	-	-	-	-	-
2001.....	35	29	21	8	2	3	-	-	-	-	-	-
2000.....	39	23	12	9	5	1	-	-	-	-	-	-
1999.....	36	21	14	6	5	2	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Human immunodeficiency virus (HIV) disease resulting in other specified diseases (B22)

All races, both sexes

2004.....	191	106	58	27	14	11	2	3	1	-	1	2
2003.....	159	95	55	27	18	6	3	2	-	-	-	-
2002.....	151	95	52	30	13	9	3	1	-	-	-	2
2001.....	148	92	61	39	12	9	5	-	1	-	-	-
2000.....	174	99	43	26	21	3	3	-	-	-	-	-
1999.....	148	86	40	29	17	7	1	1	1	-	-	-

All races, male

2004.....	154	90	45	20	8	8	2	1	-	-	-	2
2003.....	136	81	48	23	15	4	3	2	-	-	-	-
2002.....	128	81	47	25	10	5	1	-	10	-	-	2
2001.....	122	78	43	26	9	8	3	-	1	-	-	-
2000.....	147	85	39	18	14	3	2	-	-	-	-	-
1999.....	129	74	34	22	12	4	-	1	-	-	-	-

All races, female

2004.....	37	16	13	7	6	3	-	2	1	-	1	-
2003.....	23	14	7	4	3	2	-	-	-	-	-	-
2002.....	23	14	5	5	3	4	2	1	-	-	-	-
2001.....	26	14	18	13	3	1	2	-	-	-	-	-
2000.....	27	14	4	8	7	-	1	-	-	-	-	-
1999.....	19	12	6	7	5	3	1	-	1	-	-	-

White, both sexes

2004.....	97	49	28	13	5	8	-	2	1	-	1	1
2003.....	81	45	19	13	14	4	2	1	-	-	-	-
2002.....	77	48	26	13	5	6	1	1	-	-	-	1
2001.....	76	52	26	17	3	6	-	-	1	-	-	-
2000.....	86	52	18	14	8	1	1	-	-	1	-	-
1999.....	80	34	19	11	10	3	1	-	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Human immunodeficiency virus (HIV) disease resulting in other specified diseases (B22)

White, male

2004.....	85	43	23	9	3	7	-	-	-	-	-	1
2003.....	77	39	19	12	12	3	2	1	-	-	-	-
2002.....	69	43	25	11	4	3	-	-	-	-	-	1
2001.....	70	48	20	13	3	6	-	-	1	-	-	-
2000.....	75	46	18	12	5	1	-	-	-	-	-	-
1999.....	77	32	16	9	7	2	-	-	-	-	-	-

White, female

2004.....	12	6	5	4	2	1	-	2	1	-	1	-
2003.....	4	6	-	1	2	1	-	-	-	-	-	-
2002.....	8	5	1	2	1	3	1	1	-	-	-	-
2001.....	6	4	6	4	-	-	-	-	-	-	-	-
2000.....	11	6	-	2	3	-	1	-	-	-	-	-
1999.....	3	2	3	2	3	1	1	-	1	-	-	-

Black, both sexes

2004.....	93	55	30	14	7	3	2	1	-	-	-	1
2003.....	76	49	36	14	4	2	1	1	-	-	-	-
2002.....	73	46	25	17	8	3	2	-	-	-	-	1
2001.....	71	40	33	20	9	3	5	-	-	-	-	-
2000.....	87	47	25	11	12	2	2	-	-	-	-	-
1999.....	66	52	21	17	7	4	-	1	-	-	-	-

Black, male

2004.....	68	45	22	11	4	1	2	1	-	-	-	1
2003.....	57	41	29	11	3	1	1	1	-	-	-	-
2002.....	58	37	22	14	6	2	1	-	-	-	-	1
2001.....	51	30	21	12	6	2	3	-	-	-	-	-
2000.....	71	39	21	5	8	2	2	-	-	-	-	-
1999.....	50	42	18	12	5	2	-	1	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Human immunodeficiency virus (HIV) disease resulting in other conditions (B23)												
All races, both sexes												
2004.....	332	176	125	66	24	19	8	4	-	1	-	-
2003.....	440	251	130	77	47	22	7	3	1	-	-	1
2002.....	451	240	130	80	32	21	13	2	2	-	-	1
2001.....	410	203	141	62	31	21	8	3	1	-	-	1
2000.....	405	227	126	67	39	25	4	1	-	1	-	-
1999.....	372	216	116	87	33	9	7	2	4	1	-	-
All races, male												
2004.....	252	140	98	52	19	10	3	2	-	1	-	-
2003.....	338	204	98	55	38	18	5	3	-	-	-	1
2002.....	350	189	108	66	21	14	8	1	-	-	-	1
2001.....	323	169	107	50	26	13	8	2	-	-	-	1
2000.....	311	183	91	49	27	20	4	-	-	-	-	-
1999.....	293	172	94	69	28	5	2	2	1	1	-	-
All races, female												
2004.....	80	36	27	14	5	9	5	2	-	-	-	-
2003.....	102	47	32	22	9	4	2	-	1	-	-	-
2002.....	101	51	22	14	11	7	5	1	2	-	-	-
2001.....	87	34	34	12	5	8	-	1	1	-	-	-
2000.....	94	44	35	18	12	5	-	1	-	1	-	-
1999.....	79	44	22	18	5	4	5	-	3	-	-	-
White, both sexes												
2004.....	119	64	49	29	13	7	8	3	-	1	-	-
2003.....	167	108	48	24	20	15	1	2	1	-	-	-
2002.....	161	95	48	24	7	9	8	-	2	-	-	1
2001.....	147	90	50	24	12	6	3	2	1	-	-	-
2000.....	154	99	46	24	16	13	3	1	-	-	-	-
1999.....	137	79	53	36	8	2	5	2	3	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Human immunodeficiency virus (HIV) disease resulting in other conditions (B23)												
White, male												
2004.....	102	59	44	25	10	5	3	2	-	1	-	-
2003.....	140	93	39	20	18	14	1	2	-	-	-	-
2002.....	139	83	43	20	6	7	5	-	-	-	-	1
2001.....	127	76	45	21	11	4	3	1	11	-	-	-
2000.....	126	86	34	18	11	11	3	-	-	-	-	-
1999.....	119	66	50	29	7	2	1	2	-	-	-	-
White, female												
2004.....	17	5	5	4	3	2	5	1	-	-	-	-
2003.....	27	15	9	4	2	1	-	-	1	-	-	-
2002.....	22	12	5	4	1	2	3	-	2	-	-	-
2001.....	20	14	5	3	1	2	-	1	1	-	-	-
2000.....	28	13	12	6	5	2	-	1	-	-	-	-
1999.....	18	13	3	7	1	-	4	-	3	-	-	-
Black, both sexes												
2004.....	210	110	74	33	11	12	-	1	-	-	-	-
2003.....	271	142	80	52	27	7	6	1	-	-	-	1
2002.....	287	143	80	55	24	12	5	2	-	-	-	-
2001.....	259	111	89	38	18	15	5	1	-	-	-	1
2000.....	250	122	79	42	22	11	1	-	-	1	-	-
1999.....	231	137	61	51	24	7	2	-	1	1	-	-
Black, male												
2004.....	149	80	53	25	9	5	-	-	-	-	-	-
2003.....	196	110	58	35	20	4	4	1	-	-	-	1
2002.....	208	104	63	45	15	7	3	1	-	-	-	-
2001.....	193	91	61	29	14	9	5	1	-	-	-	1
2000.....	184	92	56	31	16	8	1	-	-	-	-	-
1999.....	171	106	43	40	20	3	1	-	1	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Unspecified human immunodeficiency virus (HIV) disease (B24)												
All races, both sexes												
2004.....	486	271	133	68	29	18	8	1	-	1	-	-
2003.....	353	203	94	57	28	22	5	2	-	-	-	1
2002.....	332	151	79	48	26	12	3	2	-	1	-	1
2001.....	309	132	75	54	20	12	1	1	-	2	-	-
2000.....	285	132	80	47	24	12	1	2	1	1	-	-
1999.....	222	136	66	43	24	7	4	1	-	-	-	-
All races, male												
2004.....	367	213	111	51	24	11	6	-	-	-	-	-
2003.....	280	166	80	42	22	16	3	2	-	-	-	1
2002.....	268	120	68	37	22	12	3	1	-	1	-	1
2001.....	253	108	61	40	20	10	1	1	-	2	-	-
2000.....	226	113	71	38	21	10	-	1	1	-	-	-
1999.....	185	115	53	35	20	5	3	1	-	-	-	-
All races, female												
2004.....	119	58	22	17	5	7	2	1	-	1	-	-
2003.....	73	37	14	15	6	6	2	-	-	-	-	-
2002.....	64	31	11	11	4	-	-	1	-	-	-	-
2001.....	56	24	14	14	-	2	-	-	-	-	-	-
2000.....	59	19	9	9	3	2	1	1	-	1	-	-
1999.....	37	21	13	8	4	2	1	-	-	-	-	-
White, both sexes												
2004.....	190	130	52	23	16	8	6	-	-	1	-	-
2003.....	150	96	45	19	12	10	1	-	-	-	-	1
2002.....	131	72	36	19	7	3	-	1	-	1	-	1
2001.....	136	64	22	19	16	4	-	-	-	2	-	-
2000.....	108	65	38	19	9	6	-	1	-	-	-	-
1999.....	112	57	27	22	11	3	3	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Unspecified human immunodeficiency virus (HIV) disease (B24)												
White, male												
2004.....	164	104	47	20	14	6	4	-	-	-	-	-
2003.....	131	86	40	12	9	7	1	-	-	-	-	1
2002.....	110	59	32	15	7	3	-	1	-	1	-	1
2001.....	125	56	17	17	16	4	-	-	-	2	-	-
2000.....	95	56	33	18	9	5	-	-	-	-	-	-
1999.....	98	51	25	21	10	2	3	-	-	-	-	-
White, female												
2004.....	26	26	5	3	2	2	2	-	-	1	-	-
2003.....	19	10	5	7	3	3	-	-	-	-	-	-
2002.....	21	13	4	4	-	-	-	-	-	-	-	-
2001.....	11	8	5	2	-	-	-	-	-	-	-	-
2000.....	13	9	5	1	-	1	-	1	-	-	-	-
1999.....	14	6	2	1	1	1	-	-	-	-	-	-
Black, both sexes												
2004.....	290	140	79	44	13	10	2	1	-	-	-	-
2003.....	202	106	48	37	16	11	4	2	-	-	-	-
2002.....	193	79	40	28	19	9	3	1	-	-	-	-
2001.....	172	68	52	34	4	8	1	-	-	-	-	-
2000.....	176	67	42	28	14	6	1	1	1	1	-	-
1999.....	110	77	38	20	13	4	1	1	-	-	-	-
Black, male												
2004.....	199	108	63	31	10	5	2	-	-	-	-	-
2003.....	148	80	39	30	13	8	2	2	-	-	-	-
2002.....	152	61	33	21	15	9	3	-	-	-	-	-
2001.....	127	52	43	23	4	6	1	-	-	-	-	-
2000.....	131	57	38	20	11	5	-	1	1	-	-	-
1999.....	87	62	27	13	10	3	-	1	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

All other and unspecified viral diseases (A81,A85.0-A85.1,A85.8,A86-A89,B01,B04,B07-B09,B25-B34)

All races, both sexes

2004.....	45	73	75	76	71	82	78	45	25	18	7	-
2003.....	47	63	74	76	100	93	78	70	34	21	6	-
2002.....	47	47	57	81	86	90	79	58	42	28	5	-
2001.....	43	44	68	86	77	65	62	41	36	15	4	-
2000.....	50	57	69	87	101	89	86	59	52	20	8	-
1999.....	69	73	68	76	96	108	102	80	62	17	6	-

All races, male

2004.....	27	42	44	38	36	46	34	22	9	2	1	-
2003.....	30	33	46	39	55	41	35	25	11	6	1	-
2002.....	25	21	28	40	37	43	36	21	14	4	-	-
2001.....	22	21	38	38	37	30	29	20	9	2	-	-
2000.....	24	28	36	44	53	52	36	20	14	4	-	-
1999.....	32	33	34	35	53	41	36	27	18	1	-	-

All races, female

2004.....	18	31	31	38	35	36	44	23	16	16	6	-
2003.....	17	30	28	37	45	52	43	45	23	15	5	-
2002.....	22	26	29	41	49	47	43	37	28	24	5	-
2001.....	21	23	30	48	40	35	33	21	27	13	4	-
2000.....	26	29	33	43	48	37	50	39	38	16	8	-
1999.....	37	40	34	41	43	67	66	53	44	16	6	-

White, both sexes

2004.....	38	63	63	68	63	80	74	43	23	16	6	-
2003.....	36	57	69	69	93	89	70	67	33	20	6	-
2002.....	40	41	51	74	81	80	75	57	39	27	5	-
2001.....	35	42	62	81	71	61	58	40	34	15	4	-
2000.....	40	52	56	80	95	79	81	55	51	19	7	-
1999.....	58	64	60	69	88	102	94	75	60	17	5	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

All other and unspecified viral diseases (A81,A85.0-A85.1,A85.8,A86-A89,B01,B04,B07-B09,B25-B34)

White, male

2004.....	20	37	39	37	33	45	34	21	9	2	1	-
2003.....	23	32	41	35	51	39	31	24	11	6	1	-
2002.....	22	18	26	38	37	40	34	21	13	3	-	-
2001.....	18	20	34	36	34	30	28	19	9	2	-	-
2000.....	21	25	31	40	51	48	33	19	14	4	-	-
1999.....	28	28	29	31	50	41	32	24	17	1	-	-

White, female

2004.....	18	26	24	31	30	35	40	22	14	14	5	-
2003.....	13	25	28	34	42	50	39	43	22	14	5	-
2002.....	18	23	25	36	44	40	41	36	26	24	5	-
2001.....	17	22	28	45	37	31	30	21	25	13	4	-
2000.....	19	27	25	40	44	31	48	36	37	15	7	-
1999.....	30	36	31	38	38	61	62	51	43	16	5	-

Black, both sexes

2004.....	4	8	7	5	7	2	4	2	2	1	-	-
2003.....	8	5	2	4	4	4	5	3	1	1	-	-
2002.....	6	4	3	4	4	8	3	-	3	1	-	-
2001.....	5	2	4	4	4	2	3	-	2	-	-	-
2000.....	7	5	11	6	6	9	4	2	1	1	1	-
1999.....	7	7	5	5	8	5	5	4	2	-	1	-

Black, male

2004.....	4	4	3	1	2	1	-	1	-	-	-	-
2003.....	5	1	2	4	2	2	2	1	-	-	-	-
2002.....	3	2	-	1	-	2	2	-	1	1	-	-
2001.....	3	1	3	2	1	-	-	-	-	-	-	-
2000.....	2	3	5	3	2	3	2	-	-	-	-	-
1999.....	2	4	3	2	3	-	3	2	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

All other and unspecified viral diseases (A81,A85.0-A85.1,A85.8,A86-A89,B01,B04,B07-B09,B25-B34)

Asian or Pacific Islander 2/, both sexes

2004.....	3	1	3	2	1	-	-	-	-	1	1	-
2003.....	1	-	3	1	3	-	3	-	-	-	-	-
2002.....	1	1	3	1	1	1	-	1	-	-	-	-
2001.....	2	-	2	1	1	2	1	1	-	-	-	-
2000.....	2	-	1	1	-	1	1	2	-	-	-	-
1999.....	4	1	2	1	-	1	3	1	-	-	-	-

Asian or Pacific Islander 2/, male

2004.....	3	-	2	-	1	-	-	-	-	-	-	-
2003.....	1	-	3	-	2	-	2	-	-	-	-	-
2002.....	-	1	2	1	-	1	-	-	-	-	-	-
2001.....	1	-	1	-	1	-	1	1	-	-	-	-
2000.....	-	-	-	1	-	1	1	1	-	-	-	-
1999.....	2	1	2	1	-	-	1	1	-	-	-	-

Asian or Pacific Islander 2/, female

2004.....	-	1	1	2	-	-	-	-	-	1	1	-
2003.....	-	-	-	1	1	-	1	-	-	-	-	-
2002.....	1	-	1	-	1	-	-	1	-	-	-	-
2001.....	1	-	1	1	-	2	-	-	-	-	-	-
2000.....	2	-	1	-	-	-	-	1	-	-	-	-
1999.....	2	-	-	-	-	1	2	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified infectious and parasitic diseases and their sequelae (B35-B99)												
All races, both sexes												
2004.....	571	483	342	384	428	462	457	294	136	36	7	-
2003.....	530	404	339	344	432	525	385	241	132	46	3	-
2002.....	388	304	287	308	373	462	355	236	110	34	3	-
2001.....	227	222	253	315	349	381	304	234	102	20	6	-
2000.....	184	191	224	256	353	398	303	250	100	28	5	-
1999.....	295	264	300	389	470	511	398	269	107	38	3	-
All races, male												
2004.....	404	325	204	206	221	226	211	132	52	10	1	-
2003.....	365	268	198	184	205	236	190	94	46	10	-	-
2002.....	278	196	177	183	205	218	181	104	51	10	1	-
2001.....	142	141	138	158	160	198	154	88	40	3	-	-
2000.....	116	105	126	144	210	193	142	109	36	8	1	-
1999.....	172	142	164	211	240	267	186	109	35	12	-	-
All races, female												
2004.....	167	158	138	178	207	236	246	162	84	26	6	-
2003.....	165	136	141	160	227	289	195	147	86	36	3	-
2002.....	110	108	110	125	168	244	174	132	59	24	2	-
2001.....	85	81	115	157	189	183	150	146	62	17	6	-
2000.....	68	86	98	112	143	205	161	141	64	20	4	-
1999.....	123	122	136	178	230	244	212	160	72	26	3	-
White, both sexes												
2004.....	484	385	274	306	359	396	394	268	120	26	6	-
2003.....	431	319	263	290	371	434	335	213	121	43	3	-
2002.....	293	220	231	256	322	407	318	213	96	33	2	-
2001.....	175	169	195	257	297	338	268	204	93	16	3	-
2000.....	137	153	168	210	301	352	267	230	87	20	4	-
1999.....	201	198	216	311	394	446	347	237	87	35	3	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified infectious and parasitic diseases and their sequelae (B35-B99)												
White, male												
2004.....	341	256	164	171	191	195	187	128	43	8	1	-
2003.....	302	216	154	157	173	201	165	82	45	10	-	-
2002.....	217	142	142	156	178	190	162	94	44	10	1	-
2001.....	108	109	108	131	138	173	140	79	35	3	-	-
2000.....	87	88	95	118	181	162	126	100	28	5	-	-
1999.....	121	117	117	170	206	238	155	91	28	11	-	-
White, female												
2004.....	143	129	110	135	168	201	207	140	77	18	5	-
2003.....	129	103	109	133	198	233	170	131	76	33	3	-
2002.....	76	78	89	100	144	217	156	119	52	23	1	-
2001.....	67	60	87	126	159	165	128	125	58	13	3	-
2000.....	50	65	73	92	120	190	141	130	59	15	4	-
1999.....	80	81	99	141	188	208	192	146	59	24	3	-
Black, both sexes												
2004.....	63	70	54	51	43	47	45	23	13	10	1	-
2003.....	82	65	51	33	36	59	35	18	11	2	-	-
2002.....	80	71	38	33	30	41	21	17	13	1	1	-
2001.....	41	41	43	41	44	28	26	23	7	4	3	-
2000.....	41	34	40	35	40	33	32	15	11	8	1	-
1999.....	78	54	67	62	62	53	38	25	17	3	-	-
Black, male												
2004.....	46	51	34	18	22	25	17	3	9	2	-	-
2003.....	50	38	30	18	20	21	18	8	1	-	-	-
2002.....	52	49	25	19	15	17	8	7	6	-	-	-
2001.....	27	23	23	22	19	13	10	5	4	-	-	-
2000.....	26	17	24	21	22	21	13	4	6	3	1	-
1999.....	41	18	36	30	29	23	21	14	4	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified infectious and parasitic diseases and their sequelae (B35-B99)												
Black, female												
2004.....	17	19	20	33	21	22	28	20	4	8	1	-
2003.....	32	27	21	15	16	38	17	10	10	2	-	-
2002.....	28	22	13	14	15	24	13	10	7	1	1	-
2001.....	14	18	20	19	25	15	16	18	3	4	3	-
2000.....	15	17	16	14	18	12	19	11	5	5	-	-
1999.....	37	36	31	32	33	30	17	11	13	2	-	-
American Indian 1/, both sexes												
2004.....	10	5	4	3	3	1	2	1	1	-	-	-
2003.....	8	6	4	2	-	4	3	1	-	1	-	-
2002.....	3	3	3	3	2	1	2	-	-	-	-	-
2001.....	1	1	4	3	1	2	2	3	-	-	-	-
2000.....	1	2	6	1	-	1	-	-	-	-	-	-
1999.....	3	1	5	5	4	2	4	2	1	-	-	-
American Indian 1/, male												
2004.....	6	3	2	3	1	1	-	1	-	-	-	-
2003.....	5	4	2	2	-	2	-	-	-	-	-	-
2002.....	2	2	2	1	1	1	2	-	-	-	-	-
2001.....	1	1	1	-	-	1	1	2	-	-	-	-
2000.....	1	-	1	-	-	1	-	-	-	-	-	-
1999.....	1	-	3	4	2	-	3	1	1	-	-	-
American Indian 1/, female												
2004.....	4	2	2	-	2	-	2	-	1	-	-	-
2003.....	3	2	2	-	-	2	3	1	-	1	-	-
2002.....	1	1	1	2	1	-	-	-	-	-	-	-
2001.....	-	-	3	3	1	1	1	1	-	-	-	-
2000.....	-	2	5	1	-	-	-	-	-	-	-	-
1999.....	2	1	2	1	2	2	1	1	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other and unspecified infectious and parasitic diseases and their sequelae (B35-B99)

Asian or Pacific Islander 2/, both sexes

2004.....	14	23	10	24	23	18	16	2	2	-	-	-
2003.....	9	14	21	19	25	28	12	9	-	-	-	-
2002.....	12	10	15	16	19	13	14	6	1	-	-	-
2001.....	10	11	11	14	7	13	8	4	2	-	-	-
2000.....	5	2	10	10	12	12	4	5	2	-	-	-
1999.....	13	11	12	11	10	10	9	5	2	-	-	-

Asian or Pacific Islander 2/, male

2004.....	11	15	4	14	7	5	7	-	-	-	-	-
2003.....	8	10	12	7	12	12	7	4	-	-	-	-
2002.....	7	3	8	7	11	10	9	3	1	-	-	-
2001.....	6	8	6	5	3	11	3	2	1	-	-	-
2000.....	2	-	6	5	7	9	3	5	2	-	-	-
1999.....	9	7	8	7	3	6	7	3	2	-	-	-

Asian or Pacific Islander 2/, female

2004.....	3	8	6	10	16	13	9	2	2	-	-	-
2003.....	1	4	9	12	13	16	5	5	-	-	-	-
2002.....	5	7	7	9	8	3	5	3	-	-	-	-
2001.....	4	3	5	9	4	2	5	2	1	-	-	-
2000.....	3	2	4	5	5	3	1	-	-	-	-	-
1999.....	4	4	4	4	7	4	2	2	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Mycoses (B35-B49)												
All races, both sexes												
2004.....	63	93	97	95	113	124	134	75	34	9	3	-
2003.....	70	81	90	106	127	158	114	65	37	10	-	-
2002.....	87	96	94	109	146	167	145	71	36	8	-	-
2001.....	64	77	94	114	124	172	103	89	33	7	1	-
2000.....	81	90	109	119	154	170	106	86	37	12	2	-
1999.....	130	132	150	201	229	238	160	106	42	15	1	-
All races, male												
2004.....	38	62	50	51	57	67	72	39	16	2	1	-
2003.....	37	46	54	48	64	83	65	33	14	1	-	-
2002.....	62	63	59	74	90	96	83	36	20	3	-	-
2001.....	37	47	52	62	58	97	55	37	12	1	-	-
2000.....	49	47	62	70	93	91	52	39	17	3	1	-
1999.....	70	74	84	104	133	134	88	56	18	5	-	-
All races, female												
2004.....	25	31	47	44	56	57	62	36	18	7	2	-
2003.....	33	35	36	58	63	75	49	32	23	9	-	-
2002.....	25	33	35	35	56	71	62	35	16	5	-	-
2001.....	27	30	42	52	66	75	48	52	21	6	1	-
2000.....	32	43	47	49	61	79	54	47	20	9	1	-
1999.....	60	58	66	97	96	104	72	50	24	10	1	-
White, both sexes												
2004.....	39	67	74	74	91	99	108	65	30	6	3	-
2003.....	54	54	64	88	109	131	98	57	30	9	-	-
2002.....	49	67	74	84	123	140	130	62	31	8	-	-
2001.....	48	57	67	85	96	146	85	70	29	4	-	-
2000.....	60	71	81	98	124	145	94	78	27	8	1	-
1999.....	96	101	109	160	189	206	132	86	31	13	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Mycoses (B35-B49)												
White, male												
2004.....	22	44	38	41	47	53	62	39	13	1	1	-
2003.....	29	29	39	43	52	72	57	30	14	1	-	-
2002.....	33	42	48	59	75	81	74	31	18	3	-	-
2001.....	27	35	38	45	44	82	49	30	9	1	-	-
2000.....	36	39	48	57	77	74	44	36	11	2	-	-
1999.....	50	64	60	84	115	121	73	42	15	4	-	-
White, female												
2004.....	17	23	36	33	44	46	46	26	17	5	2	-
2003.....	25	25	25	45	57	59	41	27	16	8	-	-
2002.....	16	25	26	25	48	59	56	31	13	5	-	-
2001.....	21	22	29	40	52	64	36	40	20	3	-	-
2000.....	24	32	33	41	47	71	50	42	16	6	1	-
1999.....	46	37	49	76	74	85	59	44	16	9	1	-
Black, both sexes												
2004.....	18	20	21	13	17	20	21	9	3	3	-	-
2003.....	15	22	20	11	16	22	12	6	7	1	-	-
2002.....	34	26	17	20	17	19	9	8	4	-	-	-
2001.....	15	17	23	23	23	19	14	15	4	3	1	-
2000.....	18	17	19	18	24	18	10	8	8	4	1	-
1999.....	27	24	29	35	33	27	19	16	10	2	-	-
Black, male												
2004.....	11	16	12	6	7	12	6	-	3	1	-	-
2003.....	7	13	12	3	10	9	6	2	-	-	-	-
2002.....	26	19	9	13	11	7	5	4	1	-	-	-
2001.....	9	9	13	16	11	10	5	3	3	-	-	-
2000.....	12	8	8	11	11	12	6	3	4	1	1	-
1999.....	16	7	17	17	16	11	8	11	2	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Mycoses (B35-B49)												
Black, female												
2004.....	7	4	9	7	10	8	15	9	-	2	-	-
2003.....	8	9	8	8	6	13	6	4	7	1	-	-
2002.....	8	7	8	7	6	12	4	4	3	-	-	-
2001.....	6	8	10	7	12	9	9	12	1	3	1	-
2000.....	6	9	11	7	13	6	4	5	4	3	-	-
1999.....	11	17	12	18	17	16	11	5	8	1	-	-
American Indian 1/, both sexes												
2004.....	1	1	1	2	1	-	1	-	1	-	-	-
2003.....	1	1	1	1	-	1	1	-	-	-	-	-
2002.....	1	1	1	1	1	1	1	-	-	-	-	-
2001.....	-	-	2	2	-	2	-	2	-	-	-	-
2000.....	1	1	3	-	-	1	-	-	-	-	-	-
1999.....	3	-	3	1	1	1	2	1	-	-	-	-
American Indian 1/, male												
2004.....	-	-	-	2	-	-	-	-	-	-	-	-
2003.....	1	1	-	1	-	-	-	-	-	-	-	-
2002.....	1	1	1	-	1	1	1	-	-	-	-	-
2001.....	-	-	1	-	-	1	-	2	-	-	-	-
2000.....	1	-	1	-	-	1	-	-	-	-	-	-
1999.....	1	-	1	-	1	-	1	1	-	-	-	-
American Indian 1/, female												
2004.....	1	1	1	-	1	-	1	-	1	-	-	-
2003.....	-	-	1	-	-	1	1	-	-	-	-	-
2002.....	-	-	-	1	-	-	-	-	-	-	-	-
2001.....	-	-	1	2	-	1	-	-	-	-	-	-
2000.....	-	1	2	-	-	-	-	-	-	-	-	-
1999.....	2	-	2	1	-	1	1	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Mycoses (B35-B49)												
Asian or Pacific Islander 2/, both sexes												
2004.....	5	5	1	6	4	5	4	1	-	-	-	-
2003.....	-	4	5	6	2	4	3	2	-	-	-	-
2002.....	3	2	2	4	5	7	5	1	1	-	-	-
2001.....	1	3	2	4	5	5	4	2	-	-	-	-
2000.....	2	1	6	3	6	6	2	-	2	-	-	-
1999.....	4	7	9	5	6	4	7	3	1	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	5	2	-	2	3	2	4	-	-	-	-	-
2003.....	-	3	3	1	2	2	2	1	-	-	-	-
2002.....	2	1	1	2	3	7	3	1	1	-	-	-
2001.....	1	3	-	1	3	4	1	2	-	-	-	-
2000.....	-	-	5	2	5	4	2	-	2	-	-	-
1999.....	3	3	6	3	1	2	6	2	1	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	-	3	1	4	1	3	-	1	-	-	-	-
2003.....	-	1	2	5	-	2	1	1	-	-	-	-
2002.....	1	1	1	2	2	-	2	-	-	-	-	-
2001.....	-	-	2	3	2	1	3	-	-	-	-	-
2000.....	2	1	1	1	1	2	-	-	-	-	-	-
1999.....	1	4	3	2	5	2	1	1	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Protozoal diseases (B50-B64)												
All races, both sexes												
2004.....	21	17	11	16	12	16	10	4	3	-	-	-
2003.....	18	17	15	15	12	13	8	2	1	1	-	-
2002.....	24	21	18	14	17	17	5	2	-	-	-	-
2001.....	20	25	29	33	28	13	10	4	-	-	-	-
2000.....	13	20	23	16	29	15	12	4	-	1	-	-
1999.....	27	20	16	20	27	23	10	3	1	-	-	-
All races, male												
2004.....	14	8	11	15	7	10	6	2	2	-	-	-
2003.....	10	9	12	9	5	7	4	1	1	1	-	-
2002.....	12	13	11	7	9	5	1	1	-	-	-	-
2001.....	12	13	18	18	14	5	6	2	-	-	-	-
2000.....	10	15	15	7	24	6	8	2	-	1	-	-
1999.....	20	12	9	9	14	9	4	2	-	-	-	-
All races, female												
2004.....	7	9	-	1	5	6	4	2	1	-	-	-
2003.....	8	8	3	6	7	6	4	1	-	-	-	-
2002.....	12	8	7	7	8	12	4	1	-	-	-	-
2001.....	8	12	11	15	14	8	4	2	-	-	-	-
2000.....	3	5	8	9	5	9	4	2	-	-	-	-
1999.....	7	8	7	11	13	14	6	1	1	-	-	-
White, both sexes												
2004.....	15	11	6	13	11	15	8	4	3	-	-	-
2003.....	12	16	13	9	12	12	7	2	-	1	-	-
2002.....	14	12	13	12	14	16	4	2	-	-	-	-
2001.....	13	19	18	29	27	13	10	4	-	-	-	-
2000.....	5	19	19	12	25	11	12	4	-	1	-	-
1999.....	17	16	10	17	25	20	10	3	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Protozoal diseases (B50-B64)

White, male

2004.....	12	6	6	13	7	9	4	2	2	-	-	-
2003.....	7	9	10	5	5	7	3	1	-	1	-	-
2002.....	9	8	8	6	8	5	1	1	-	-	-	-
2001.....	7	10	13	16	13	5	6	2	-	-	-	-
2000.....	4	15	11	7	21	3	8	2	-	1	-	-
1999.....	12	10	6	9	13	8	4	2	-	-	-	-

White, female

2004.....	3	5	-	-	4	6	4	2	1	-	-	-
2003.....	5	7	3	4	7	5	4	1	-	-	-	-
2002.....	5	4	5	6	6	11	3	1	-	-	-	-
2001.....	6	9	5	13	14	8	4	2	-	-	-	-
2000.....	1	4	8	5	4	8	4	2	-	-	-	-
1999.....	5	6	4	8	12	12	6	1	1	-	-	-

Black, both sexes

2004.....	5	4	3	2	1	1	1	-	-	-	-	-
2003.....	5	1	1	4	-	1	1	-	1	-	-	-
2002.....	9	8	2	2	1	1	1	-	-	-	-	-
2001.....	7	5	9	3	1	-	-	-	-	-	-	-
2000.....	8	1	3	4	4	3	-	-	-	-	-	-
1999.....	10	3	6	2	2	2	-	-	-	-	-	-

Black, male

2004.....	1	2	3	1	-	1	1	-	-	-	-	-
2003.....	2	-	1	2	-	-	1	-	1	-	-	-
2002.....	3	5	2	1	-	-	-	-	-	-	-	-
2001.....	5	3	3	2	1	-	-	-	-	-	-	-
2000.....	6	-	3	-	3	2	-	-	-	-	-	-
1999.....	8	1	3	-	1	-	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malaria (B50-B54)												
All races, both sexes												
2004.....	1	-	-	1	-	-	-	-	-	-	-	-
2003.....	-	2	-	1	-	-	-	-	-	-	-	-
2002.....	4	2	-	1	1	-	-	-	-	-	-	-
2001.....	-	-	2	1	-	1	-	-	-	-	-	-
2000.....	1	-	-	-	-	1	-	-	-	-	-	-
1999.....	-	1	-	1	1	-	-	-	-	-	-	-
All races, male												
2004.....	1	-	-	1	-	-	-	-	-	-	-	-
2003.....	-	2	-	1	-	-	-	-	-	-	-	-
2002.....	3	1	-	1	1	-	-	-	-	-	-	-
2001.....	-	-	1	1	-	1	-	-	-	-	-	-
2000.....	1	-	-	-	-	1	-	-	-	-	-	-
1999.....	-	1	-	-	1	-	-	-	-	-	-	-
All races, female												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	1	1	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	1	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	1	-	-	-	-	-	-	-	-
White, both sexes												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	2	-	1	-	-	-	-	-	-	-	-
2002.....	3	1	-	1	1	-	-	-	-	-	-	-
2001.....	-	-	-	1	-	1	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	1	-	-	1	-	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumocystosis (B59)												
All races, both sexes												
2004.....	17	17	11	14	10	16	9	4	3	-	-	-
2003.....	16	14	14	11	11	12	8	2	1	1	-	-
2002.....	20	19	17	13	16	14	5	2	-	-	-	-
2001.....	18	23	24	30	25	12	10	4	-	-	-	-
2000.....	12	18	22	15	25	13	11	3	-	1	-	-
1999.....	27	15	13	19	26	22	8	1	1	-	-	-
All races, male												
2004.....	10	8	11	13	6	10	6	2	2	-	-	-
2003.....	8	7	11	7	4	7	4	1	1	1	-	-
2002.....	9	12	10	6	8	5	1	1	-	-	-	-
2001.....	11	13	15	16	11	4	6	2	-	-	-	-
2000.....	9	13	14	7	20	5	7	2	-	1	-	-
1999.....	20	10	8	9	13	8	4	-	-	-	-	-
All races, female												
2004.....	7	9	-	1	4	6	3	2	1	-	-	-
2003.....	8	7	3	4	7	5	4	1	-	-	-	-
2002.....	11	7	7	7	8	9	4	1	-	-	-	-
2001.....	7	10	9	14	14	8	4	2	-	-	-	-
2000.....	3	5	8	8	5	8	4	1	-	-	-	-
1999.....	7	5	5	10	13	14	4	1	1	-	-	-
White, both sexes												
2004.....	12	11	6	12	9	15	7	4	3	-	-	-
2003.....	10	13	12	7	11	12	7	2	-	1	-	-
2002.....	11	11	13	11	13	14	4	2	-	-	-	-
2001.....	13	18	16	26	25	12	10	4	-	-	-	-
2000.....	5	17	18	11	22	10	11	3	-	1	-	-
1999.....	17	14	9	17	24	19	8	1	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumocystosis (B59)												
White, male												
2004.....	9	6	6	12	6	9	4	2	2	-	-	-
2003.....	5	7	9	4	4	7	3	1	-	1	-	-
2002.....	6	7	8	5	7	5	1	1	-	-	-	-
2001.....	7	10	11	14	11	4	6	2	-	-	-	-
2000.....	4	13	10	7	18	3	7	2	-	1	-	-
1999.....	12	9	6	9	12	7	4	-	-	-	-	-
White, female												
2004.....	3	5	-	-	3	6	3	2	1	-	-	-
2003.....	5	6	3	3	7	5	4	1	-	-	-	-
2002.....	5	4	5	6	6	9	3	1	-	-	-	-
2001.....	6	8	5	12	14	8	4	2	-	-	-	-
2000.....	1	4	8	4	4	7	4	1	-	-	-	-
1999.....	5	5	3	8	12	12	4	1	1	-	-	-
Black, both sexes												
2004.....	5	4	3	2	1	1	1	-	-	-	-	-
2003.....	5	1	1	3	-	-	1	-	1	-	-	-
2002.....	8	8	2	2	1	-	1	-	-	-	-	-
2001.....	5	4	7	3	-	-	-	-	-	-	-	-
2000.....	7	1	3	4	3	3	-	-	-	-	-	-
1999.....	10	-	4	1	2	2	-	-	-	-	-	-
Black, male												
2004.....	1	2	3	1	-	1	1	-	-	-	-	-
2003.....	2	-	1	2	-	-	1	-	1	-	-	-
2002.....	3	5	2	1	-	-	-	-	-	-	-	-
2001.....	4	3	3	2	-	-	-	-	-	-	-	-
2000.....	5	-	3	-	2	2	-	-	-	-	-	-
1999.....	8	-	2	-	1	-	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified protozoal diseases (B58,B60-B64)												
All races, both sexes												
2004.....	2	-	-	1	2	-	1	-	-	-	-	-
2003.....	1	1	1	3	1	1	-	-	-	-	-	-
2002.....	-	-	1	-	-	2	-	-	-	-	-	-
2001.....	2	2	3	2	2	-	-	-	-	-	-	-
2000.....	-	2	1	1	3	-	1	1	-	-	-	-
1999.....	-	4	3	-	-	1	2	2	-	-	-	-
All races, male												
2004.....	2	-	-	1	1	-	-	-	-	-	-	-
2003.....	1	-	1	1	1	-	-	-	-	-	-	-
2002.....	-	-	1	-	-	-	-	-	-	-	-	-
2001.....	1	-	2	1	2	-	-	-	-	-	-	-
2000.....	-	2	1	-	3	-	1	-	-	-	-	-
1999.....	-	1	1	-	-	1	-	2	-	-	-	-
All races, female												
2004.....	-	-	-	-	1	-	1	-	-	-	-	-
2003.....	-	1	-	2	-	1	-	-	-	-	-	-
2002.....	-	-	-	-	-	2	-	-	-	-	-	-
2001.....	1	2	1	1	-	-	-	-	-	-	-	-
2000.....	-	-	-	1	-	-	-	1	-	-	-	-
1999.....	-	3	2	-	-	-	2	-	-	-	-	-
White, both sexes												
2004.....	2	-	-	1	2	-	1	-	-	-	-	-
2003.....	1	1	1	1	1	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	1	-	-	-	-	-	-
2001.....	-	1	2	2	1	-	-	-	-	-	-	-
2000.....	-	2	1	1	2	-	1	1	-	-	-	-
1999.....	-	1	1	-	-	1	2	2	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Helminthiases (B65-B83)												
All races, both sexes												
2004.....	5	1	3	6	3	2	-	-	-	-	-	-
2003.....	-	3	3	2	2	1	-	1	-	-	-	-
2002.....	1	1	1	1	2	-	-	2	-	1	-	-
2001.....	-	1	5	-	2	3	1	2	1	-	-	-
2000.....	1	1	1	6	1	2	2	3	-	-	-	-
1999.....	-	-	3	1	1	1	1	-	-	-	-	-
All races, male												
2004.....	3	-	2	5	1	1	-	-	-	-	-	-
2003.....	-	3	2	1	2	-	-	1	-	-	-	-
2002.....	-	-	-	1	2	-	-	2	-	-	-	-
2001.....	-	1	4	-	1	3	-	2	-	-	-	-
2000.....	1	1	1	4	-	2	1	2	-	-	-	-
1999.....	-	-	2	1	-	1	-	-	-	-	-	-
All races, female												
2004.....	2	1	1	1	2	1	-	-	-	-	-	-
2003.....	-	-	1	1	-	1	-	-	-	-	-	-
2002.....	1	1	1	-	-	-	-	-	-	1	-	-
2001.....	-	-	1	-	1	-	1	-	1	-	-	-
2000.....	-	-	-	2	1	-	1	1	-	-	-	-
1999.....	-	-	1	-	1	-	1	-	-	-	-	-
White, both sexes												
2004.....	2	1	1	4	2	1	-	-	-	-	-	-
2003.....	-	2	3	2	2	1	-	1	-	-	-	-
2002.....	-	1	1	1	2	-	-	2	-	-	-	-
2001.....	-	1	5	-	2	2	1	2	1	-	-	-
2000.....	1	1	1	4	1	1	2	3	-	-	-	-
1999.....	-	-	2	1	1	1	1	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Schistosomiasis (bilharziasis) (B65)												
All races, both sexes												
2004.....	1	-	1	1	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	1	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	1	-	-	-	-	-	-	-
All races, male												
2004.....	1	-	-	1	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	1	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-
All races, female												
2004.....	-	-	1	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	1	-	-	-	-	-	-	-
White, both sexes												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	1	-	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified helminthiases (B66-B83)												
All races, both sexes												
2004.....	4	1	2	5	3	2	-	-	-	-	-	-
2003.....	-	3	3	2	2	1	-	1	-	-	-	-
2002.....	1	1	1	1	2	-	-	2	-	1	-	-
2001.....	-	1	5	-	2	3	1	2	1	-	-	-
2000.....	1	1	1	5	1	2	2	3	-	-	-	-
1999.....	-	-	3	1	-	1	1	-	-	-	-	-
All races, male												
2004.....	2	-	2	4	1	1	-	-	-	-	-	-
2003.....	-	3	2	1	2	-	-	1	-	-	-	-
2002.....	-	-	-	1	2	-	-	2	-	-	-	-
2001.....	-	1	4	-	1	3	-	2	-	-	-	-
2000.....	1	1	1	3	-	2	1	2	-	-	-	-
1999.....	-	-	2	1	-	1	-	-	-	-	-	-
All races, female												
2004.....	2	1	-	1	2	1	-	-	-	-	-	-
2003.....	-	-	1	1	-	1	-	-	-	-	-	-
2002.....	1	1	1	-	-	-	-	-	-	1	-	-
2001.....	-	-	1	-	1	-	1	-	1	-	-	-
2000.....	-	-	-	2	1	-	1	1	-	-	-	-
1999.....	-	-	1	-	-	-	1	-	-	-	-	-
White, both sexes												
2004.....	2	1	1	4	2	1	-	-	-	-	-	-
2003.....	-	2	3	2	2	1	-	1	-	-	-	-
2002.....	-	1	1	1	2	-	-	2	-	-	-	-
2001.....	-	1	5	-	2	2	1	2	1	-	-	-
2000.....	1	1	1	4	1	1	2	3	-	-	-	-
1999.....	-	-	2	1	-	1	1	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Sequelae of tuberculosis (B90)												
All races, both sexes												
2004.....	-	8	-	8	11	11	18	12	8	1	1	-
2003.....	1	3	3	5	8	12	21	14	6	2	-	-
2002.....	4	3	5	5	8	17	18	26	8	5	-	-
2001.....	2	6	4	6	16	18	24	23	5	-	-	-
2000.....	2	3	5	8	17	23	28	23	6	1	-	-
1999.....	7	7	7	8	27	30	47	23	9	2	-	-
All races, male												
2004.....	-	6	-	4	5	5	9	5	4	-	-	-
2003.....	1	1	3	3	6	3	6	4	3	-	-	-
2002.....	2	1	2	4	4	8	9	10	1	-	-	-
2001.....	-	3	2	1	6	9	12	8	2	-	-	-
2000.....	2	3	2	3	11	8	12	16	1	1	-	-
1999.....	4	2	5	5	12	14	16	7	3	1	-	-
All races, female												
2004.....	-	2	-	4	6	6	9	7	4	1	1	-
2003.....	-	2	-	2	2	9	15	10	3	2	-	-
2002.....	2	2	3	1	4	9	9	16	7	5	-	-
2001.....	2	3	2	5	10	9	12	15	3	-	-	-
2000.....	-	-	3	5	6	15	16	7	5	-	-	-
1999.....	3	5	2	3	15	16	31	16	6	1	-	-
White, both sexes												
2004.....	-	3	-	3	4	6	15	11	7	-	1	-
2003.....	-	3	-	2	8	5	18	12	6	2	-	-
2002.....	4	1	3	3	4	13	13	23	8	5	-	-
2001.....	1	4	2	4	13	13	21	19	3	-	-	-
2000.....	1	1	-	6	10	19	23	22	5	-	-	-
1999.....	1	6	6	2	19	24	41	21	8	2	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Sequelae of tuberculosis (B90)

White, male

2004.....	-	2	-	1	2	3	7	5	3	-	-	-
2003.....	-	1	-	1	6	1	5	3	3	-	-	-
2002.....	2	-	1	2	2	5	6	8	1	-	-	-
2001.....	-	2	1	1	6	6	11	8	1	-	-	-
2000.....	1	1	-	3	7	4	9	15	-	-	-	-
1999.....	-	2	4	1	8	10	13	6	2	1	-	-

White, female

2004.....	-	1	-	2	2	3	8	6	4	-	1	-
2003.....	-	2	-	1	2	4	13	9	3	2	-	-
2002.....	2	1	2	1	2	8	7	15	7	5	-	-
2001.....	1	2	1	3	7	7	10	11	2	-	-	-
2000.....	-	-	-	3	3	15	14	7	5	-	-	-
1999.....	1	4	2	1	11	14	28	15	6	1	-	-

Black, both sexes

2004.....	-	3	-	1	2	2	2	1	1	1	-	-
2003.....	1	-	1	2	-	3	-	-	-	-	-	-
2002.....	-	1	2	1	-	3	1	1	-	-	-	-
2001.....	-	1	1	-	1	3	-	1	-	-	-	-
2000.....	-	2	3	-	3	1	4	-	1	1	-	-
1999.....	3	1	-	3	2	2	2	1	-	-	-	-

Black, male

2004.....	-	2	-	-	1	1	2	-	1	-	-	-
2003.....	1	-	1	2	-	-	-	-	-	-	-	-
2002.....	-	1	1	1	-	3	-	-	-	-	-	-
2001.....	-	1	1	-	-	1	-	-	-	-	-	-
2000.....	-	2	2	-	2	1	2	-	1	1	-	-
1999.....	1	-	-	1	1	2	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Sequelae of tuberculosis (B90)												
Black, female												
2004.....	-	1	-	1	1	1	-	1	-	1	-	-
2003.....	-	-	-	-	-	3	-	-	-	-	-	-
2002.....	-	-	1	-	-	-	1	1	-	-	-	-
2001.....	-	-	-	-	1	2	-	1	-	-	-	-
2000.....	-	-	1	-	1	-	2	-	-	-	-	-
1999.....	2	1	-	2	1	-	2	1	-	-	-	-
American Indian 1/, both sexes												
2004.....	-	-	-	-	1	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	1	1	1	-	-	-	-
2002.....	-	-	-	-	-	-	1	-	-	-	-	-
2001.....	-	-	-	-	-	-	1	1	-	-	-	-
2000.....	-	-	2	1	-	-	-	-	-	-	-	-
1999.....	-	-	1	1	2	1	2	-	-	-	-	-
American Indian 1/, male												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	1	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	1	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	1	1	1	-	2	-	-	-	-	-
American Indian 1/, female												
2004.....	-	-	-	-	1	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	1	1	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	1	1	-	-	-	-
2000.....	-	-	2	1	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	1	1	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Sequelae of tuberculosis (B90)

Asian or Pacific Islander 2/, both sexes

2004.....	-	2	-	4	4	3	1	-	-	-	-	-
2003.....	-	-	2	1	-	3	2	1	-	-	-	-
2002.....	-	1	-	1	4	1	3	2	-	-	-	-
2001.....	1	1	1	2	2	2	2	2	2	-	-	-
2000.....	1	-	-	1	4	3	1	1	-	-	-	-
1999.....	3	-	-	2	4	3	2	1	1	-	-	-

Asian or Pacific Islander 2/, male

2004.....	-	2	-	3	2	1	-	-	-	-	-	-
2003.....	-	-	2	-	-	1	1	1	-	-	-	-
2002.....	-	-	-	1	2	-	2	2	-	-	-	-
2001.....	-	-	-	-	-	2	1	-	1	-	-	-
2000.....	1	-	-	-	2	3	1	1	-	-	-	-
1999.....	3	-	-	2	2	2	1	1	1	-	-	-

Asian or Pacific Islander 2/, female

2004.....	-	-	-	1	2	2	1	-	-	-	-	-
2003.....	-	-	-	1	-	2	1	-	-	-	-	-
2002.....	-	1	-	-	2	1	1	-	-	-	-	-
2001.....	1	1	1	2	2	-	1	2	1	-	-	-
2000.....	-	-	-	1	2	-	-	-	-	-	-	-
1999.....	-	-	-	-	2	1	1	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Sequelae of poliomyelitis (B91)												
All races, both sexes												
2004.....	9	10	26	36	48	55	57	41	20	6	-	-
2003.....	11	22	30	29	47	50	44	33	18	4	1	-
2002.....	13	21	22	41	48	65	48	40	15	5	-	-
2001.....	17	19	35	40	56	63	47	36	17	1	-	-
2000.....	14	23	22	32	47	59	40	37	12	2	1	-
1999.....	13	13	22	29	44	44	34	36	13	5	1	-
All races, male												
2004.....	4	6	13	19	24	26	30	21	6	2	-	-
2003.....	5	11	16	15	22	16	27	13	6	2	-	-
2002.....	5	9	15	21	21	23	23	17	7	3	-	-
2001.....	6	10	17	18	29	37	19	14	9	-	-	-
2000.....	9	9	12	18	24	24	15	17	1	-	-	-
1999.....	5	7	11	19	19	23	11	13	6	-	-	-
All races, female												
2004.....	5	4	13	17	24	29	27	20	14	4	-	-
2003.....	6	11	14	14	25	34	17	20	12	2	1	-
2002.....	8	12	7	20	27	42	25	23	8	2	-	-
2001.....	11	9	18	22	27	26	28	22	8	1	-	-
2000.....	5	14	10	14	23	35	25	20	11	2	1	-
1999.....	8	6	11	10	25	21	23	23	7	5	1	-
White, both sexes												
2004.....	9	7	25	35	48	54	56	41	20	6	-	-
2003.....	9	21	28	28	46	50	42	32	18	4	1	-
2002.....	12	20	20	40	45	65	48	39	15	5	-	-
2001.....	16	16	34	38	55	62	46	36	17	1	-	-
2000.....	13	22	22	31	46	58	39	37	12	2	1	-
1999.....	12	13	21	29	43	44	34	36	12	5	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Sequelae of poliomyelitis (B91)												
White, male												
2004.....	4	4	12	19	24	26	30	21	6	2	-	-
2003.....	4	11	14	15	21	16	27	12	6	2	-	-
2002.....	5	8	14	21	21	23	23	16	7	3	-	-
2001.....	5	8	16	16	28	36	19	14	9	-	-	-
2000.....	8	8	12	18	24	24	15	17	1	-	-	-
1999.....	5	7	11	19	18	23	11	13	5	-	-	-
White, female												
2004.....	5	3	13	16	24	28	26	20	14	4	-	-
2003.....	5	10	14	13	25	34	15	20	12	2	1	-
2002.....	7	12	6	19	24	42	25	23	8	2	-	-
2001.....	11	8	18	22	27	26	27	22	8	1	-	-
2000.....	5	14	10	13	22	34	24	20	11	2	1	-
1999.....	7	6	10	10	25	21	23	23	7	5	1	-
Black, both sexes												
2004.....	-	3	1	1	-	1	1	-	-	-	-	-
2003.....	1	1	2	1	1	-	1	-	-	-	-	-
2002.....	1	1	1	1	2	-	-	1	-	-	-	-
2001.....	-	2	1	-	1	1	1	-	-	-	-	-
2000.....	1	1	-	1	1	1	1	-	-	-	-	-
1999.....	1	-	1	-	1	-	-	-	-	-	-	-
Black, male												
2004.....	-	2	1	-	-	-	-	-	-	-	-	-
2003.....	1	-	2	-	1	-	-	-	-	-	-	-
2002.....	-	1	-	-	-	-	-	1	-	-	-	-
2001.....	-	1	1	-	1	1	-	-	-	-	-	-
2000.....	1	1	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	1	-	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

All other and unspecified infectious and parasitic diseases and their sequelae (B85-B89,B92-B99)

All races, both sexes

2004.....	473	354	205	223	241	254	238	162	71	20	3	-
2003.....	430	278	198	187	236	291	198	126	70	29	2	-
2002.....	259	162	147	138	152	196	139	95	51	15	3	-
2001.....	124	94	86	122	123	112	119	80	46	12	5	-
2000.....	73	54	64	75	105	129	115	97	45	12	2	-
1999.....	118	92	102	130	142	175	146	101	42	16	1	-

All races, male

2004.....	345	243	128	112	127	117	94	65	24	6	-	-
2003.....	312	198	111	108	106	127	88	42	22	6	-	-
2002.....	197	110	90	76	79	86	65	38	23	4	1	-
2001.....	87	67	45	59	52	47	62	25	17	2	-	-
2000.....	45	30	34	42	58	62	54	33	17	3	-	-
1999.....	73	47	53	73	62	86	67	31	8	6	-	-

All races, female

2004.....	128	111	77	111	114	137	144	97	47	14	3	-
2003.....	118	80	87	79	130	164	110	84	48	23	2	-
2002.....	62	52	57	62	73	110	74	57	28	11	2	-
2001.....	37	27	41	63	71	65	57	55	29	10	5	-
2000.....	28	24	30	33	47	67	61	64	28	9	2	-
1999.....	45	45	49	57	80	89	79	70	34	10	1	-

White, both sexes

2004.....	419	296	168	177	203	221	207	147	60	14	2	-
2003.....	356	223	155	161	194	235	170	109	67	27	2	-
2002.....	214	119	120	116	134	173	123	85	42	15	2	-
2001.....	97	72	69	101	104	102	105	73	43	11	3	-
2000.....	57	39	45	59	95	118	97	86	43	9	2	-
1999.....	75	62	68	102	117	151	129	91	35	15	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

All other and unspecified infectious and parasitic diseases and their sequelae (B85-B89,B92-B99)

White, male

2004.....	302	200	107	94	110	104	84	61	19	5	-	-
2003.....	262	164	89	92	87	105	73	35	22	6	-	-
2002.....	168	84	71	67	70	76	58	36	18	4	1	-
2001.....	69	53	36	53	46	42	55	23	16	2	-	-
2000.....	37	24	23	30	52	56	49	28	16	2	-	-
1999.....	54	34	35	56	52	75	54	28	6	6	-	-

White, female

2004.....	117	96	61	83	93	117	123	86	41	9	2	-
2003.....	94	59	66	69	107	130	97	74	45	21	2	-
2002.....	46	35	49	49	64	97	65	49	24	11	1	-
2001.....	28	19	33	48	58	60	50	50	27	9	3	-
2000.....	20	15	22	29	43	62	48	58	27	7	2	-
1999.....	21	28	33	46	65	76	75	63	29	9	1	-

Black, both sexes

2004.....	39	40	27	33	22	23	20	13	9	6	1	-
2003.....	60	41	27	15	19	33	21	12	3	1	-	-
2002.....	35	35	16	9	10	18	10	7	9	-	1	-
2001.....	19	16	9	15	18	5	11	7	3	1	2	-
2000.....	14	13	15	12	8	10	17	7	2	3	-	-
1999.....	37	26	31	22	24	22	17	8	7	1	-	-

Black, male

2004.....	33	29	17	10	14	11	8	3	5	1	-	-
2003.....	39	25	14	11	9	12	11	6	-	-	-	-
2002.....	23	23	13	4	4	7	3	2	5	-	-	-
2001.....	13	9	5	4	6	1	5	2	1	-	-	-
2000.....	7	6	11	10	6	6	5	1	1	1	-	-
1999.....	16	10	16	12	10	10	13	3	2	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

All other and unspecified infectious and parasitic diseases and their sequelae (B85-B89,B92-B99)

Black, female

2004.....	6	11	10	23	8	12	12	10	4	5	1	-
2003.....	21	16	13	4	10	21	10	6	3	1	-	-
2002.....	12	12	3	5	6	11	7	5	4	-	1	-
2001.....	6	7	4	11	12	4	6	5	2	1	2	-
2000.....	7	7	4	2	2	4	12	6	1	2	-	-
1999.....	21	16	15	10	14	12	4	5	5	1	-	-

American Indian 1/, both sexes

2004.....	8	4	3	1	1	1	1	1	-	-	-	-
2003.....	6	5	2	1	-	2	-	-	-	1	-	-
2002.....	2	2	2	2	-	-	-	-	-	-	-	-
2001.....	1	1	2	-	1	-	1	-	-	-	-	-
2000.....	-	1	1	-	-	-	-	-	-	-	-	-
1999.....	-	1	1	3	1	-	-	1	-	-	-	-

American Indian 1/, male

2004.....	5	3	2	1	1	1	-	1	-	-	-	-
2003.....	4	3	1	1	-	1	-	-	-	-	-	-
2002.....	1	1	1	1	-	-	-	-	-	-	-	-
2001.....	1	1	-	-	-	-	1	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	1	3	-	-	-	-	-	-	-	-

American Indian 1/, female

2004.....	3	1	1	-	-	-	1	-	-	-	-	-
2003.....	2	2	1	-	-	1	-	-	-	1	-	-
2002.....	1	1	1	1	-	-	-	-	-	-	-	-
2001.....	-	-	2	-	1	-	-	-	-	-	-	-
2000.....	-	1	1	-	-	-	-	-	-	-	-	-
1999.....	-	1	-	-	1	-	-	1	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

All other and unspecified infectious and parasitic diseases and their sequelae (B85-B89,B92-B99)

Asian or Pacific Islander 2/, both sexes

2004.....	7	14	7	12	15	9	10	1	2	-	-	-
2003.....	8	9	14	10	23	21	7	5	-	-	-	-
2002.....	8	6	9	11	8	5	6	3	-	-	-	-
2001.....	7	5	6	6	-	5	2	-	-	-	-	-
2000.....	2	1	3	4	2	1	1	4	-	-	-	-
1999.....	6	3	2	3	-	2	-	1	-	-	-	-

Asian or Pacific Islander 2/, male

2004.....	5	11	2	7	2	1	2	-	-	-	-	-
2003.....	7	6	7	4	10	9	4	1	-	-	-	-
2002.....	5	2	5	4	5	3	4	-	-	-	-	-
2001.....	4	4	4	2	-	4	1	-	-	-	-	-
2000.....	1	-	-	2	-	-	-	4	-	-	-	-
1999.....	3	3	1	2	-	1	-	-	-	-	-	-

Asian or Pacific Islander 2/, female

2004.....	2	3	5	5	13	8	8	1	2	-	-	-
2003.....	1	3	7	6	13	12	3	4	-	-	-	-
2002.....	3	4	4	7	3	2	2	3	-	-	-	-
2001.....	3	1	2	4	-	1	1	-	-	-	-	-
2000.....	1	1	3	2	2	1	1	-	-	-	-	-
1999.....	3	-	1	1	-	1	-	1	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
II. Neoplasms (C00-D48)												
All races, both sexes												
2004.....	30,720	43,705	54,561	64,424	77,336	88,886	81,843	53,023	24,020	6,237	839	10
2003.....	30,678	42,821	54,109	64,096	79,506	91,209	81,097	52,992	23,875	6,164	810	12
2002.....	30,729	42,048	52,536	64,526	82,579	91,702	79,957	52,350	23,485	6,138	822	7
2001.....	30,778	40,380	51,096	65,172	84,420	92,159	77,940	51,782	22,801	6,035	776	8
2000.....	29,628	39,443	50,786	66,850	85,827	92,797	76,947	51,258	22,707	5,967	804	12
1999.....	28,655	39,098	51,102	67,843	87,179	92,025	75,217	50,039	21,761	5,829	828	9
All races, male												
2004.....	16,261	23,519	29,969	35,609	42,200	47,660	41,452	24,777	9,751	2,161	230	8
2003.....	16,033	22,950	29,662	35,572	43,555	48,630	41,131	24,486	9,535	2,100	227	8
2002.....	15,970	22,737	29,058	35,917	45,483	48,879	40,405	24,245	9,504	2,051	226	5
2001.....	15,912	21,669	28,248	36,463	46,675	49,150	39,379	24,022	9,032	2,053	245	7
2000.....	15,197	21,182	27,902	37,317	47,289	49,404	38,822	23,595	8,868	1,936	197	11
1999.....	14,703	21,146	28,343	37,905	48,161	49,303	37,975	22,986	8,654	1,908	239	6
All races, female												
2004.....	14,459	20,186	24,592	28,815	35,136	41,226	40,391	28,246	14,269	4,076	609	2
2003.....	14,645	19,871	24,447	28,524	35,951	42,579	39,966	28,506	14,340	4,064	583	4
2002.....	14,759	19,311	23,478	28,609	37,096	42,823	39,552	28,105	13,981	4,087	596	2
2001.....	14,866	18,711	22,848	28,709	37,745	43,009	38,561	27,760	13,769	3,982	531	1
2000.....	14,431	18,261	22,884	29,533	38,538	43,393	38,125	27,663	13,839	4,031	607	1
1999.....	13,952	17,952	22,759	29,938	39,018	42,722	37,242	27,053	13,107	3,921	589	3
White, both sexes												
2004.....	24,357	36,024	46,142	54,961	67,271	78,944	73,797	48,222	21,698	5,592	707	8
2003.....	24,351	35,602	45,757	54,820	69,484	81,131	73,048	48,174	21,651	5,467	666	11
2002.....	24,645	34,895	44,268	55,138	72,411	81,639	72,108	47,636	21,238	5,438	690	4
2001.....	24,912	33,527	43,151	55,657	74,035	82,137	70,160	47,077	20,622	5,354	641	4
2000.....	24,068	32,887	42,930	57,561	75,248	82,767	69,661	46,526	20,554	5,286	665	7
1999.....	23,296	32,472	43,057	58,386	76,702	81,982	68,147	45,388	19,701	5,172	694	7

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
II. Neoplasms (C00-D48)												
White, male												
2004.....	12,924	19,366	25,285	30,431	36,701	42,349	37,327	22,567	8,714	1,907	190	6
2003.....	12,728	19,066	25,043	30,445	38,185	43,263	37,036	22,292	8,598	1,821	187	8
2002.....	12,827	18,858	24,356	30,792	39,985	43,463	36,341	22,051	8,538	1,808	183	3
2001.....	12,841	17,922	23,816	31,190	40,989	43,696	35,313	21,735	8,099	1,804	204	4
2000.....	12,253	17,515	23,506	32,151	41,380	43,881	35,001	21,321	7,952	1,682	156	7
1999.....	11,854	17,505	23,832	32,646	42,405	43,782	34,346	20,700	7,780	1,655	201	4
White, female												
2004.....	11,433	16,658	20,857	24,530	30,570	36,595	36,470	25,655	12,984	3,685	517	2
2003.....	11,623	16,536	20,714	24,375	31,299	37,868	36,012	25,882	13,053	3,646	479	3
2002.....	11,818	16,037	19,912	24,346	32,426	38,176	35,767	25,585	12,700	3,630	507	1
2001.....	12,071	15,605	19,335	24,467	33,046	38,441	34,847	25,342	12,523	3,550	437	-
2000.....	11,815	15,372	19,424	25,410	33,868	38,886	34,660	25,205	12,602	3,604	509	-
1999.....	11,442	14,967	19,225	25,740	34,297	38,200	33,801	24,688	11,921	3,517	493	3
Black, both sexes												
2004.....	5,321	6,423	7,053	7,877	8,216	8,163	6,534	3,930	1,901	542	120	2
2003.....	5,316	6,109	7,050	7,759	8,280	8,275	6,621	3,961	1,877	588	124	-
2002.....	5,065	6,031	6,938	7,945	8,512	8,442	6,524	3,964	1,900	586	114	3
2001.....	4,905	5,805	6,667	8,052	8,734	8,392	6,496	3,922	1,879	604	125	4
2000.....	4,768	5,556	6,698	7,893	8,991	8,488	6,157	4,015	1,847	611	122	4
1999.....	4,578	5,700	6,882	8,079	8,943	8,579	6,032	4,038	1,769	561	125	1
Black, male												
2004.....	2,841	3,505	3,954	4,325	4,516	4,310	3,328	1,777	817	200	35	2
2003.....	2,833	3,353	3,920	4,298	4,497	4,386	3,346	1,760	763	223	28	-
2002.....	2,642	3,304	3,962	4,380	4,650	4,524	3,360	1,816	798	198	38	2
2001.....	2,615	3,193	3,714	4,519	4,822	4,554	3,369	1,890	767	211	39	3
2000.....	2,559	3,141	3,764	4,431	5,021	4,657	3,229	1,877	760	222	38	3
1999.....	2,469	3,184	3,884	4,532	4,911	4,720	3,079	1,943	725	204	33	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
II. Neoplasms (C00-D48)												
Black, female												
2004.....	2,480	2,918	3,099	3,552	3,700	3,853	3,206	2,153	1,084	342	85	-
2003.....	2,483	2,756	3,130	3,461	3,783	3,889	3,275	2,201	1,114	365	96	-
2002.....	2,423	2,727	2,976	3,565	3,862	3,918	3,164	2,148	1,102	388	76	1
2001.....	2,290	2,612	2,953	3,533	3,912	3,838	3,127	2,032	1,112	393	86	1
2000.....	2,209	2,415	2,934	3,462	3,970	3,831	2,928	2,138	1,087	389	84	1
1999.....	2,109	2,516	2,998	3,547	4,032	3,859	2,953	2,095	1,044	357	92	-
American Indian 1/, both sexes												
2004.....	184	263	308	343	336	291	229	114	57	15	-	-
2003.....	189	224	249	263	303	287	216	112	49	17	5	-
2002.....	186	220	269	276	322	256	212	105	54	18	4	-
2001.....	178	211	262	302	310	273	194	116	52	13	-	-
2000.....	133	192	231	255	294	253	166	109	47	17	5	-
1999.....	133	191	222	278	270	232	182	95	38	11	2	-
American Indian 1/, male												
2004.....	98	144	163	186	177	168	102	48	25	7	-	-
2003.....	92	112	132	147	157	138	102	57	22	10	2	-
2002.....	94	123	145	143	168	133	108	42	13	4	1	-
2001.....	89	109	137	162	173	131	100	50	24	6	-	-
2000.....	72	106	123	137	160	134	82	46	20	9	1	-
1999.....	63	92	117	148	149	123	91	46	17	7	2	-
American Indian 1/, female												
2004.....	86	119	145	157	159	123	127	66	32	8	-	-
2003.....	97	112	117	116	146	149	114	55	27	7	3	-
2002.....	92	97	124	133	154	123	104	63	41	14	3	-
2001.....	89	102	125	140	137	142	94	66	28	7	-	-
2000.....	61	86	108	118	134	119	84	63	27	8	4	-
1999.....	70	99	105	130	121	109	91	49	21	4	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
II. Neoplasms (C00-D48)												
Asian or Pacific Islander 2/, both sexes												
2004.....	858	995	1,058	1,243	1,513	1,488	1,283	757	364	88	12	-
2003.....	822	886	1,053	1,254	1,439	1,516	1,212	745	298	92	15	1
2002.....	833	902	1,061	1,167	1,334	1,365	1,113	645	293	96	14	-
2001.....	783	837	1,016	1,161	1,341	1,357	1,090	667	248	64	10	-
2000.....	659	808	927	1,141	1,294	1,289	963	608	259	53	12	1
1999.....	648	735	941	1,100	1,264	1,232	856	518	253	85	7	1
Asian or Pacific Islander 2/, male												
2004.....	398	504	567	667	806	833	695	385	195	47	5	-
2003.....	380	419	567	682	716	843	647	377	152	46	10	-
2002.....	407	452	595	602	680	759	596	336	155	41	4	-
2001.....	367	445	581	592	691	769	597	347	142	32	2	-
2000.....	313	420	509	598	728	732	510	351	136	23	2	1
1999.....	317	365	510	579	696	678	459	297	132	42	3	1
Asian or Pacific Islander 2/, female												
2004.....	460	491	491	576	707	655	588	372	169	41	7	-
2003.....	442	467	486	572	723	673	565	368	146	46	5	1
2002.....	426	450	466	565	654	606	517	309	138	55	10	-
2001.....	416	392	435	569	650	588	493	320	106	32	8	-
2000.....	346	388	418	543	566	557	453	257	123	30	10	-
1999.....	331	370	431	521	568	554	397	221	121	43	4	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant neoplasms (C00-C97)												
All races, both sexes												
2004.....	30,330	43,127	53,829	63,503	75,914	86,865	79,220	50,851	22,844	5,855	795	8
2003.....	30,291	42,315	53,377	63,141	78,107	89,086	78,531	50,789	22,660	5,827	770	12
2002.....	30,342	41,487	51,904	63,598	81,159	89,565	77,497	50,298	22,323	5,772	789	7
2001.....	30,363	39,827	50,396	64,172	82,846	89,967	75,478	49,680	21,660	5,684	727	8
2000.....	29,231	38,930	50,075	65,843	84,288	90,655	74,444	49,155	21,564	5,653	764	11
1999.....	28,269	38,637	50,430	66,814	85,524	89,874	72,896	48,041	20,702	5,492	779	9
All races, male												
2004.....	16,037	23,196	29,542	35,065	41,422	46,582	40,152	23,831	9,325	2,050	221	6
2003.....	15,815	22,671	29,256	35,048	42,745	47,516	39,866	23,531	9,124	2,006	217	8
2002.....	15,774	22,407	28,708	35,374	44,711	47,734	39,218	23,345	9,106	1,952	218	5
2001.....	15,694	21,390	27,854	35,900	45,799	47,951	38,175	23,140	8,628	1,951	235	7
2000.....	14,972	20,912	27,515	36,730	46,461	48,284	37,623	22,714	8,506	1,850	190	10
1999.....	14,483	20,891	27,942	37,315	47,229	48,184	36,876	22,147	8,287	1,826	224	6
All races, female												
2004.....	14,293	19,931	24,287	28,438	34,492	40,283	39,068	27,020	13,519	3,805	574	2
2003.....	14,476	19,644	24,121	28,093	35,362	41,570	38,665	27,258	13,536	3,821	553	4
2002.....	14,568	19,080	23,196	28,224	36,448	41,831	38,279	26,953	13,217	3,820	571	2
2001.....	14,669	18,437	22,542	28,272	37,047	42,016	37,303	26,540	13,032	3,733	492	1
2000.....	14,259	18,018	22,560	29,113	37,827	42,371	36,821	26,441	13,058	3,803	574	1
1999.....	13,786	17,746	22,488	29,499	38,295	41,690	36,020	25,894	12,415	3,666	555	3
White, both sexes												
2004.....	24,051	35,551	45,522	54,161	66,006	77,122	71,375	46,199	20,593	5,244	668	6
2003.....	24,033	35,179	45,119	54,000	68,223	79,174	70,674	46,109	20,524	5,156	636	11
2002.....	24,336	34,425	43,715	54,326	71,132	79,678	69,847	45,714	20,162	5,114	662	4
2001.....	24,559	33,051	42,552	54,773	72,631	80,120	67,885	45,136	19,572	5,040	602	4
2000.....	23,735	32,440	42,319	56,673	73,897	80,805	67,321	44,566	19,488	4,995	630	7
1999.....	22,976	32,083	42,485	57,500	75,203	80,023	65,986	43,518	18,705	4,869	650	7

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant neoplasms (C00-C97)												
White, male												
2004.....	12,743	19,099	24,921	29,948	35,998	41,367	36,117	21,678	8,311	1,803	181	4
2003.....	12,546	18,836	24,690	29,981	37,449	42,224	35,860	21,387	8,219	1,735	179	8
2002.....	12,665	18,583	24,049	30,313	39,275	42,403	35,246	21,201	8,171	1,723	175	3
2001.....	12,648	17,679	23,476	30,684	40,191	42,585	34,197	20,922	7,722	1,710	196	4
2000.....	12,061	17,275	23,182	31,634	40,650	42,849	33,865	20,503	7,611	1,600	149	7
1999.....	11,668	17,289	23,479	32,130	41,549	42,755	33,311	19,908	7,428	1,579	187	4
White, female												
2004.....	11,308	16,452	20,601	24,213	30,008	35,755	35,258	24,521	12,282	3,441	487	2
2003.....	11,487	16,343	20,429	24,019	30,774	36,950	34,814	24,722	12,305	3,421	457	3
2002.....	11,671	15,842	19,666	24,013	31,857	37,275	34,601	24,513	11,991	3,391	487	1
2001.....	11,911	15,372	19,076	24,089	32,440	37,535	33,688	24,214	11,850	3,330	406	-
2000.....	11,674	15,165	19,137	25,039	33,247	37,956	33,456	24,063	11,877	3,395	481	-
1999.....	11,308	14,794	19,006	25,370	33,654	37,268	32,675	23,610	11,277	3,290	463	3
Black, both sexes												
2004.....	5,253	6,330	6,966	7,778	8,093	8,009	6,375	3,819	1,845	512	115	2
2003.....	5,257	6,043	6,972	7,652	8,164	8,142	6,464	3,863	1,807	566	116	-
2002.....	5,001	5,954	6,871	7,845	8,400	8,300	6,364	3,856	1,832	552	110	3
2001.....	4,850	5,741	6,582	7,951	8,596	8,251	6,339	3,803	1,801	571	116	4
2000.....	4,710	5,499	6,620	7,797	8,839	8,338	6,023	3,897	1,784	591	118	3
1999.....	4,522	5,635	6,798	7,955	8,809	8,414	5,898	3,930	1,717	531	121	1
Black, male												
2004.....	2,805	3,455	3,904	4,274	4,456	4,234	3,263	1,737	800	195	35	2
2003.....	2,804	3,314	3,876	4,248	4,432	4,328	3,274	1,726	740	217	27	-
2002.....	2,615	3,254	3,929	4,329	4,602	4,459	3,290	1,775	776	187	38	2
2001.....	2,593	3,164	3,669	4,468	4,754	4,486	3,296	1,843	745	204	37	3
2000.....	2,528	3,114	3,715	4,371	4,950	4,584	3,183	1,832	742	219	38	2
1999.....	2,442	3,148	3,845	4,467	4,847	4,644	3,030	1,905	715	198	33	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant neoplasms (C00-C97)												
Black, female												
2004.....	2,448	2,875	3,062	3,504	3,637	3,775	3,112	2,082	1,045	317	80	-
2003.....	2,453	2,729	3,096	3,404	3,732	3,814	3,190	2,137	1,067	349	89	-
2002.....	2,386	2,700	2,942	3,516	3,798	3,841	3,074	2,081	1,056	365	72	1
2001.....	2,257	2,577	2,913	3,483	3,842	3,765	3,043	1,960	1,056	367	79	1
2000.....	2,182	2,385	2,905	3,426	3,889	3,754	2,840	2,065	1,042	372	80	1
1999.....	2,080	2,487	2,953	3,488	3,962	3,770	2,868	2,025	1,002	333	88	-
American Indian 1/, both sexes												
2004.....	181	262	302	341	329	284	227	107	54	14	-	-
2003.....	189	220	245	257	300	283	211	108	48	16	4	-
2002.....	184	219	266	274	315	252	208	104	53	16	4	-
2001.....	177	209	261	298	298	263	190	109	50	13	-	-
2000.....	132	190	227	251	288	248	164	103	46	16	4	-
1999.....	131	187	219	271	264	228	177	91	36	10	1	-
American Indian 1/, male												
2004.....	97	144	161	185	173	163	102	46	23	6	-	-
2003.....	92	109	130	146	155	135	99	55	21	10	2	-
2002.....	94	123	142	141	166	129	107	41	13	3	1	-
2001.....	88	108	136	159	169	127	98	46	23	6	-	-
2000.....	72	106	120	136	155	133	80	44	20	9	1	-
1999.....	62	90	115	146	146	120	88	44	17	7	1	-
American Indian 1/, female												
2004.....	84	118	141	156	156	121	125	61	31	8	-	-
2003.....	97	111	115	111	145	148	112	53	27	6	2	-
2002.....	90	96	124	133	149	123	101	63	40	13	3	-
2001.....	89	101	125	139	129	136	92	63	27	7	-	-
2000.....	60	84	107	115	133	115	84	59	26	7	3	-
1999.....	69	97	104	125	118	108	89	47	19	3	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant neoplasms (C00-C97)												
Asian or Pacific Islander 2/, both sexes												
2004.....	845	984	1,039	1,223	1,486	1,450	1,243	726	352	85	12	-
2003.....	812	873	1,041	1,232	1,420	1,487	1,182	709	281	89	14	1
2002.....	821	889	1,052	1,153	1,312	1,335	1,078	624	276	90	13	-
2001.....	777	826	1,001	1,150	1,321	1,333	1,064	632	237	60	9	-
2000.....	654	801	909	1,122	1,264	1,264	936	589	246	51	12	1
1999.....	640	732	928	1,088	1,248	1,209	835	502	244	82	7	1
Asian or Pacific Islander 2/, male												
2004.....	392	498	556	658	795	818	670	370	191	46	5	-
2003.....	373	412	560	673	709	829	633	363	144	44	9	-
2002.....	400	447	588	591	668	743	575	328	146	39	4	-
2001.....	365	439	573	589	685	753	584	329	138	31	2	-
2000.....	311	417	498	589	706	718	495	335	133	22	2	1
1999.....	311	364	503	572	687	665	447	290	127	42	3	1
Asian or Pacific Islander 2/, female												
2004.....	453	486	483	565	691	632	573	356	161	39	7	-
2003.....	439	461	481	559	711	658	549	346	137	45	5	1
2002.....	421	442	464	562	644	592	503	296	130	51	9	-
2001.....	412	387	428	561	636	580	480	303	99	29	7	-
2000.....	343	384	411	533	558	546	441	254	113	29	10	-
1999.....	329	368	425	516	561	544	388	212	117	40	4	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)												
All races, both sexes												
2004.....	743	834	947	937	937	943	831	545	240	77	13	-
2003.....	686	867	959	950	924	955	737	543	242	78	16	1
2002.....	670	875	936	846	1,052	973	748	505	262	77	13	-
2001.....	649	861	879	893	1,019	1,004	783	502	240	67	5	1
2000.....	660	770	872	942	1,017	933	693	488	249	74	13	-
1999.....	612	753	863	889	1,007	993	751	501	220	64	15	-
All races, male												
2004.....	591	687	713	689	623	597	460	249	97	27	2	-
2003.....	575	694	729	678	626	589	410	255	108	20	6	1
2002.....	534	681	711	619	698	590	407	245	101	24	4	-
2001.....	521	687	685	631	696	598	427	232	83	18	2	1
2000.....	516	618	652	667	659	569	370	217	96	19	-	-
1999.....	474	568	663	629	679	605	402	229	79	14	1	-
All races, female												
2004.....	152	147	234	248	314	346	371	296	143	50	11	-
2003.....	111	173	230	272	298	366	327	288	134	58	10	-
2002.....	136	194	225	227	354	383	341	260	161	53	9	-
2001.....	128	174	194	262	323	406	356	270	157	49	3	-
2000.....	144	152	220	275	358	364	323	271	153	55	13	-
1999.....	138	185	200	260	328	388	349	272	141	50	14	-
White, both sexes												
2004.....	547	645	756	771	814	834	747	510	219	72	12	-
2003.....	502	706	761	781	780	832	651	504	216	73	16	1
2002.....	491	683	745	689	938	877	681	458	247	67	11	-
2001.....	497	664	697	720	881	900	720	463	222	66	5	-
2000.....	475	581	674	776	868	832	629	449	229	67	11	-
1999.....	430	560	688	737	866	873	687	457	213	60	14	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)

White, male

2004.....	431	535	564	561	527	522	412	229	82	25	2	-
2003.....	432	562	564	553	515	506	356	234	97	19	6	1
2002.....	390	537	560	499	619	536	368	221	97	19	3	-
2001.....	400	525	537	509	594	529	391	213	77	18	2	-
2000.....	358	467	499	541	555	495	332	202	86	18	-	-
1999.....	337	418	530	521	587	524	362	205	76	13	1	-

White, female

2004.....	116	110	192	210	287	312	335	281	137	47	10	-
2003.....	70	144	197	228	265	326	295	270	119	54	10	-
2002.....	101	146	185	190	319	341	313	237	150	48	8	-
2001.....	97	139	160	211	287	371	329	250	145	48	3	-
2000.....	117	114	175	235	313	337	297	247	143	49	11	-
1999.....	93	142	158	216	279	349	325	252	137	47	13	-

Black, both sexes

2004.....	174	160	163	134	105	88	64	24	15	4	1	-
2003.....	140	139	170	148	116	87	63	30	20	3	-	-
2002.....	149	150	169	125	93	66	48	33	12	7	2	-
2001.....	124	173	146	142	117	82	49	25	16	1	-	1
2000.....	156	164	166	135	121	82	56	26	15	6	2	-
1999.....	155	165	157	130	120	99	54	37	5	2	1	-

Black, male

2004.....	141	129	127	108	84	57	38	15	10	2	-	-
2003.....	113	116	144	112	92	58	41	15	9	-	-	-
2002.....	124	122	136	97	64	36	28	18	3	3	1	-
2001.....	100	142	122	100	87	54	26	12	5	-	-	1
2000.....	137	133	132	106	87	61	34	8	8	-	-	-
1999.....	118	127	119	93	78	70	33	19	1	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)

Black, female

2004.....	33	31	36	26	21	31	26	9	5	2	1	-
2003.....	27	23	26	36	24	29	22	15	11	3	-	-
2002.....	25	28	33	28	29	30	20	15	9	4	1	-
2001.....	24	31	24	42	30	28	23	13	11	1	-	-
2000.....	19	31	34	29	34	21	22	18	7	6	2	-
1999.....	37	38	38	37	42	29	21	18	4	1	1	-

American Indian 1/, both sexes

2004.....	1	6	7	6	6	3	-	-	2	-	-	-
2003.....	5	4	5	1	5	7	3	1	1	-	-	-
2002.....	4	9	6	8	6	3	3	-	-	-	-	-
2001.....	4	6	7	4	4	4	2	1	-	-	-	-
2000.....	4	6	4	7	10	3	-	3	-	-	-	-
1999.....	7	3	4	6	5	2	-	1	-	-	-	-

American Indian 1/, male

2004.....	1	5	5	5	4	2	-	-	2	-	-	-
2003.....	4	2	2	1	3	2	1	-	1	-	-	-
2002.....	3	5	4	3	4	3	2	-	-	-	-	-
2001.....	3	6	6	4	4	1	2	1	-	-	-	-
2000.....	3	4	1	5	7	2	-	1	-	-	-	-
1999.....	4	3	2	4	3	-	-	1	-	-	-	-

American Indian 1/, female

2004.....	-	1	2	1	2	1	-	-	-	-	-	-
2003.....	1	2	3	-	2	5	2	1	-	-	-	-
2002.....	1	4	2	5	2	-	1	-	-	-	-	-
2001.....	1	-	1	-	-	3	-	-	-	-	-	-
2000.....	1	2	3	2	3	1	-	2	-	-	-	-
1999.....	3	-	2	2	2	2	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)

Asian or Pacific Islander 2/, both sexes

2004.....	21	23	21	26	12	18	20	11	4	1	-	-
2003.....	39	18	23	20	23	29	20	8	5	2	-	-
2002.....	26	33	16	24	15	27	16	14	3	3	-	-
2001.....	24	18	29	27	17	18	12	13	2	-	-	-
2000.....	25	19	28	24	18	16	8	10	5	1	-	-
1999.....	20	25	14	16	16	19	10	6	2	2	-	-

Asian or Pacific Islander 2/, male

2004.....	18	18	17	15	8	16	10	5	3	-	-	-
2003.....	26	14	19	12	16	23	12	6	1	1	-	-
2002.....	17	17	11	20	11	15	9	6	1	2	-	-
2001.....	18	14	20	18	11	14	8	6	1	-	-	-
2000.....	18	14	20	15	10	11	4	6	2	1	-	-
1999.....	15	20	12	11	11	11	7	4	2	-	-	-

Asian or Pacific Islander 2/, female

2004.....	3	5	4	11	4	2	10	6	1	1	-	-
2003.....	13	4	4	8	7	6	8	2	4	1	-	-
2002.....	9	16	5	4	4	12	7	8	2	1	-	-
2001.....	6	4	9	9	6	4	4	7	1	-	-	-
2000.....	7	5	8	9	8	5	4	4	3	-	-	-
1999.....	5	5	2	5	5	8	3	2	-	2	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Of lip (C00)												
All races, both sexes												
2004.....	1	3	5	6	6	10	17	7	8	4	1	-
2003.....	2	6	4	10	9	17	8	4	6	3	-	-
2002.....	2	4	2	3	7	14	8	12	9	2	-	-
2001.....	3	7	5	6	8	10	8	12	5	3	1	-
2000.....	3	8	4	2	7	8	17	10	3	-	-	-
1999.....	2	2	4	7	10	8	7	7	3	-	1	-
All races, male												
2004.....	1	2	5	6	4	9	10	5	-	2	1	-
2003.....	1	5	3	8	7	14	5	1	4	1	-	-
2002.....	2	4	-	2	5	13	6	7	3	2	-	-
2001.....	3	7	5	4	6	8	5	8	5	1	-	-
2000.....	3	8	4	2	7	6	9	7	2	-	-	-
1999.....	2	2	4	5	9	4	4	4	1	-	-	-
All races, female												
2004.....	-	1	-	-	2	1	7	2	8	2	-	-
2003.....	1	1	1	2	2	3	3	3	2	2	-	-
2002.....	-	-	2	1	2	1	2	5	6	-	-	-
2001.....	-	-	-	2	2	2	3	4	-	2	1	-
2000.....	-	-	-	-	-	2	8	3	1	-	-	-
1999.....	-	-	-	2	1	4	3	3	2	-	1	-
White, both sexes												
2004.....	1	3	4	6	6	10	14	6	8	4	1	-
2003.....	2	6	3	9	9	17	8	4	6	3	-	-
2002.....	2	3	1	2	6	14	8	12	8	2	-	-
2001.....	3	7	4	6	8	10	7	12	5	3	1	-
2000.....	3	8	4	2	7	8	16	10	3	-	-	-
1999.....	2	2	4	7	10	7	7	7	3	-	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Of lip (C00)												
Asian or Pacific Islander 2/, both sexes												
2004.....	-	-	1	-	-	-	1	-	-	-	-	-
2003.....	-	-	-	1	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	-	-	1	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	-	-	-	-	-	-	1	-	-	-	-	-
2003.....	-	-	-	1	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	1	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Of tongue (C01-C02)												
All races, both sexes												
2004.....	198	219	217	244	210	188	191	141	41	15	2	-
2003.....	184	230	217	230	225	231	161	109	44	15	7	-
2002.....	169	211	246	202	244	237	180	109	60	15	1	-
2001.....	173	222	204	194	225	231	180	116	43	6	1	-
2000.....	167	183	231	216	214	208	157	103	49	14	4	-
1999.....	152	170	197	209	228	221	164	102	47	15	4	-
All races, male												
2004.....	155	181	166	176	130	114	96	53	14	5	1	-
2003.....	148	190	150	163	141	131	88	51	16	1	4	-
2002.....	128	165	174	154	147	137	86	44	23	4	1	-
2001.....	139	177	161	137	157	124	94	40	12	1	-	-
2000.....	127	143	176	155	130	109	69	46	14	2	-	-
1999.....	115	123	150	152	158	142	77	48	14	2	-	-
All races, female												
2004.....	43	38	51	68	80	74	95	88	27	10	1	-
2003.....	36	40	67	67	84	100	73	58	28	14	3	-
2002.....	41	46	72	48	97	100	94	65	37	11	-	-
2001.....	34	45	43	57	68	107	86	76	31	5	1	-
2000.....	40	40	55	61	84	99	88	57	35	12	4	-
1999.....	37	47	47	57	70	79	87	54	33	13	4	-
White, both sexes												
2004.....	159	187	195	209	188	176	174	134	35	14	2	-
2003.....	142	193	192	191	198	206	146	102	39	15	7	-
2002.....	137	177	203	162	224	220	170	95	59	14	1	-
2001.....	145	190	167	160	203	214	164	112	38	6	1	-
2000.....	126	152	188	190	187	191	146	98	46	12	3	-
1999.....	115	128	168	179	202	201	157	94	43	13	4	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Of tongue (C01-C02)												
White, male												
2004.....	123	154	145	150	112	108	85	49	11	5	1	-
2003.....	121	162	131	135	124	115	79	46	15	1	4	-
2002.....	106	139	141	122	134	126	81	39	22	3	1	-
2001.....	117	149	131	117	141	115	84	39	11	1	-	-
2000.....	92	121	145	134	114	97	63	42	13	2	-	-
1999.....	89	89	130	134	139	128	74	44	13	2	-	-
White, female												
2004.....	36	33	50	59	76	68	89	85	24	9	1	-
2003.....	21	31	61	56	74	91	67	56	24	14	3	-
2002.....	31	38	62	40	90	94	89	56	37	11	-	-
2001.....	28	41	36	43	62	99	80	73	27	5	1	-
2000.....	34	31	43	56	73	94	83	56	33	10	3	-
1999.....	26	39	38	45	63	73	83	50	30	11	4	-
Black, both sexes												
2004.....	34	27	19	29	18	11	16	3	2	1	-	-
2003.....	28	34	24	35	19	14	11	6	3	-	-	-
2002.....	26	29	39	36	15	11	5	6	-	1	-	-
2001.....	25	28	32	24	18	14	10	3	4	-	-	-
2000.....	37	28	34	23	23	14	7	3	2	2	1	-
1999.....	35	37	27	25	22	18	6	6	4	1	-	-
Black, male												
2004.....	28	23	18	25	15	5	11	2	-	-	-	-
2003.....	19	27	18	26	13	11	7	4	1	-	-	-
2002.....	19	21	31	29	10	8	3	3	-	1	-	-
2001.....	21	26	27	15	14	8	6	1	1	-	-	-
2000.....	34	22	26	20	14	11	5	2	1	-	-	-
1999.....	26	30	19	15	17	13	3	2	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Of tongue (C01-C02)												
Black, female												
2004.....	6	4	1	4	3	6	5	1	2	1	-	-
2003.....	9	7	6	9	6	3	4	2	2	-	-	-
2002.....	7	8	8	7	5	3	2	3	-	-	-	-
2001.....	4	2	5	9	4	6	4	2	3	-	-	-
2000.....	3	6	8	3	9	3	2	1	1	2	1	-
1999.....	9	7	8	10	5	5	3	4	3	1	-	-
American Indian 1/, both sexes												
2004.....	-	2	-	1	1	-	-	-	1	-	-	-
2003.....	2	-	-	-	3	3	-	-	-	-	-	-
2002.....	3	-	3	2	1	1	-	-	-	-	-	-
2001.....	-	-	3	1	1	-	-	-	-	-	-	-
2000.....	1	1	1	-	1	-	-	-	-	-	-	-
1999.....	1	1	1	2	2	1	-	-	-	-	-	-
American Indian 1/, male												
2004.....	-	2	-	-	1	-	-	-	1	-	-	-
2003.....	1	-	-	-	2	1	-	-	-	-	-	-
2002.....	2	-	2	1	-	1	-	-	-	-	-	-
2001.....	-	-	2	1	1	-	-	-	-	-	-	-
2000.....	-	-	-	-	1	-	-	-	-	-	-	-
1999.....	-	1	-	1	1	-	-	-	-	-	-	-
American Indian 1/, female												
2004.....	-	-	-	1	-	-	-	-	-	-	-	-
2003.....	1	-	-	-	1	2	-	-	-	-	-	-
2002.....	1	-	1	1	1	-	-	-	-	-	-	-
2001.....	-	-	1	-	-	-	-	-	-	-	-	-
2000.....	1	1	1	-	-	-	-	-	-	-	-	-
1999.....	1	-	1	1	1	1	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Of tongue (C01-C02)												
Asian or Pacific Islander 2/, both sexes												
2004.....	5	3	3	5	3	1	1	4	3	-	-	-
2003.....	12	3	1	4	5	8	4	1	2	-	-	-
2002.....	3	5	1	2	4	5	5	8	1	-	-	-
2001.....	3	4	2	9	3	3	6	1	1	-	-	-
2000.....	3	2	8	3	3	3	4	2	1	-	-	-
1999.....	1	4	1	3	2	1	1	2	-	1	-	-
Asian or Pacific Islander 2/, male												
2004.....	4	2	3	1	2	1	-	2	2	-	-	-
2003.....	7	1	1	2	2	4	2	1	-	-	-	-
2002.....	1	5	-	2	3	2	2	2	1	-	-	-
2001.....	1	2	1	4	1	1	4	-	-	-	-	-
2000.....	1	-	5	1	1	1	1	2	-	-	-	-
1999.....	-	3	1	2	1	1	-	2	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	1	1	-	4	1	-	1	2	1	-	-	-
2003.....	5	2	-	2	3	4	2	-	2	-	-	-
2002.....	2	-	1	-	1	3	3	6	-	-	-	-
2001.....	2	2	1	5	2	2	2	1	1	-	-	-
2000.....	2	2	3	2	2	2	3	-	1	-	-	-
1999.....	1	1	-	1	1	-	1	-	-	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Of pharynx (C10-C13,C14.0)												
All races, both sexes												
2004.....	317	367	455	421	414	417	311	172	60	14	3	-
2003.....	292	365	435	442	410	378	273	192	75	16	-	1
2002.....	297	381	412	362	456	385	259	166	58	21	3	-
2001.....	284	371	415	394	428	405	291	177	67	17	-	1
2000.....	280	367	397	414	456	389	261	151	69	14	3	-
1999.....	275	358	386	391	452	407	272	156	49	11	2	-
All races, male												
2004.....	256	303	349	322	298	284	189	93	30	8	-	-
2003.....	247	295	338	328	291	255	179	108	42	6	-	1
2002.....	237	287	325	268	318	251	165	95	32	8	1	-
2001.....	230	302	321	288	308	269	174	96	28	6	-	1
2000.....	222	297	305	300	319	261	157	83	34	4	-	-
1999.....	217	284	298	284	310	262	173	90	24	4	-	-
All races, female												
2004.....	61	64	106	99	116	133	122	79	30	6	3	-
2003.....	45	70	97	114	119	123	94	84	33	10	-	-
2002.....	60	94	87	94	138	134	94	71	26	13	2	-
2001.....	54	69	94	106	120	136	117	81	39	11	-	-
2000.....	58	70	92	114	137	128	104	68	35	10	3	-
1999.....	58	74	88	107	142	145	99	66	25	7	2	-
White, both sexes												
2004.....	207	259	329	323	346	350	272	157	53	12	3	-
2003.....	195	283	317	348	329	319	225	172	64	13	-	1
2002.....	195	278	302	287	396	333	220	147	53	15	1	-
2001.....	189	265	316	307	358	347	257	158	63	17	-	-
2000.....	188	255	285	319	371	330	232	131	59	12	3	-
1999.....	173	254	293	312	373	339	234	142	48	10	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Of pharynx (C10-C13,C14.0)												
White, male												
2004.....	165	219	250	245	246	235	168	81	24	7	-	-
2003.....	169	226	238	253	226	213	145	96	37	6	-	1
2002.....	153	214	236	207	279	220	139	83	31	5	-	-
2001.....	154	214	243	215	256	228	155	85	25	6	-	-
2000.....	144	207	214	234	254	215	140	76	26	3	-	-
1999.....	134	203	226	226	258	214	147	81	23	3	-	-
White, female												
2004.....	42	40	79	78	100	115	104	76	29	5	3	-
2003.....	26	57	79	95	103	106	80	76	27	7	-	-
2002.....	42	64	66	80	117	113	81	64	22	10	1	-
2001.....	35	51	73	92	102	119	102	73	38	11	-	-
2000.....	44	48	71	85	117	115	92	55	33	9	3	-
1999.....	39	51	67	86	115	125	87	61	25	7	1	-
Black, both sexes												
2004.....	95	90	113	76	63	54	27	13	5	2	-	-
2003.....	71	67	99	82	64	45	32	14	9	2	-	-
2002.....	80	76	93	54	48	31	29	14	5	4	2	-
2001.....	74	92	72	69	58	42	28	13	3	-	-	1
2000.....	71	96	95	74	68	47	26	15	8	1	-	-
1999.....	81	85	80	68	66	57	31	12	-	1	1	-
Black, male												
2004.....	78	70	90	61	50	37	13	10	4	1	-	-
2003.....	59	57	85	67	52	32	24	7	5	-	-	-
2002.....	69	62	77	43	31	17	19	8	1	2	1	-
2001.....	60	75	57	57	43	29	15	7	2	-	-	1
2000.....	61	77	77	54	53	36	15	5	6	-	-	-
1999.....	66	66	61	50	43	42	20	8	-	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Of pharynx (C10-C13,C14.0)												
Black, female												
2004.....	17	20	23	15	13	17	14	3	1	1	-	-
2003.....	12	10	14	15	12	13	8	7	4	2	-	-
2002.....	11	14	16	11	17	14	10	6	4	2	1	-
2001.....	14	17	15	12	15	13	13	6	1	-	-	-
2000.....	10	19	18	20	15	11	11	10	2	1	-	-
1999.....	15	19	19	18	23	15	11	4	-	-	1	-
American Indian 1/, both sexes												
2004.....	-	3	2	4	3	2	-	-	1	-	-	-
2003.....	2	3	3	-	2	4	3	1	-	-	-	-
2002.....	1	5	3	2	3	2	3	-	-	-	-	-
2001.....	3	2	4	2	2	3	1	1	-	-	-	-
2000.....	2	1	2	5	7	2	-	1	-	-	-	-
1999.....	4	1	3	-	3	1	-	-	-	-	-	-
American Indian 1/, male												
2004.....	-	3	2	4	1	2	-	-	1	-	-	-
2003.....	2	2	1	-	1	1	1	-	-	-	-	-
2002.....	1	2	2	1	2	2	2	-	-	-	-	-
2001.....	2	2	4	2	2	1	1	1	-	-	-	-
2000.....	2	1	1	3	5	2	-	-	-	-	-	-
1999.....	3	1	2	-	2	-	-	-	-	-	-	-
American Indian 1/, female												
2004.....	-	-	-	-	2	-	-	-	-	-	-	-
2003.....	-	1	2	-	1	3	2	1	-	-	-	-
2002.....	-	3	1	1	1	-	1	-	-	-	-	-
2001.....	1	-	-	-	-	2	-	-	-	-	-	-
2000.....	-	-	1	2	2	-	-	1	-	-	-	-
1999.....	1	-	1	-	1	1	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Of pharynx (C10-C13,C14.0)												
Asian or Pacific Islander 2/, both sexes												
2004.....	15	15	11	18	2	11	12	2	1	-	-	-
2003.....	24	12	16	12	15	10	13	5	2	1	-	-
2002.....	21	22	14	19	9	19	7	5	-	2	-	-
2001.....	18	12	23	16	10	13	5	5	1	-	-	-
2000.....	19	15	15	16	10	10	3	4	2	1	-	-
1999.....	17	18	10	11	10	10	7	2	1	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	13	11	7	12	1	10	8	2	1	-	-	-
2003.....	17	10	14	8	12	9	9	5	-	-	-	-
2002.....	14	9	10	17	6	12	5	4	-	1	-	-
2001.....	14	11	17	14	7	11	3	3	1	-	-	-
2000.....	15	12	13	9	7	8	2	2	2	1	-	-
1999.....	14	14	9	8	7	6	6	1	1	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	2	4	4	6	1	1	4	-	-	-	-	-
2003.....	7	2	2	4	3	1	4	-	2	1	-	-
2002.....	7	13	4	2	3	7	2	1	-	1	-	-
2001.....	4	1	6	2	3	2	2	2	-	-	-	-
2000.....	4	3	2	7	3	2	1	2	-	-	-	-
1999.....	3	4	1	3	3	4	1	1	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Of other and unspecified sites within the lip, oral cavity and pharynx (C03-C09,C14.2-C14.8)

All races, both sexes

2004.....	227	245	270	266	307	328	312	225	131	44	7	-
2003.....	208	266	303	268	280	329	295	238	117	44	9	-
2002.....	202	279	276	279	345	337	301	218	135	39	9	-
2001.....	189	261	255	299	358	358	304	197	125	41	3	-
2000.....	210	212	240	310	340	328	258	224	128	46	6	-
1999.....	183	223	276	282	317	357	308	236	121	38	8	-

All races, male

2004.....	179	201	193	185	191	190	165	98	53	12	-	-
2003.....	179	204	238	179	187	189	138	95	46	12	2	-
2002.....	167	225	212	195	228	189	150	99	43	10	2	-
2001.....	149	201	198	202	225	197	154	88	38	10	2	-
2000.....	164	170	167	210	203	193	135	81	46	13	-	-
1999.....	140	159	211	188	202	197	148	87	40	8	1	-

All races, female

2004.....	48	44	77	81	116	138	147	127	78	32	7	-
2003.....	29	62	65	89	93	140	157	143	71	32	7	-
2002.....	35	54	64	84	117	148	151	119	92	29	7	-
2001.....	40	60	57	97	133	161	150	109	87	31	1	-
2000.....	46	42	73	100	137	135	123	143	82	33	6	-
1999.....	43	64	65	94	115	160	160	149	81	30	7	-

White, both sexes

2004.....	180	196	228	233	274	298	287	213	123	42	6	-
2003.....	163	224	249	233	244	290	272	226	107	42	9	-
2002.....	157	225	239	238	312	310	283	204	127	36	9	-
2001.....	160	202	210	247	312	329	292	181	116	40	3	-
2000.....	158	166	197	265	303	303	235	210	121	43	5	-
1999.....	140	176	223	239	281	326	289	214	119	37	8	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Of other and unspecified sites within the lip, oral cavity and pharynx (C03-C09,C14.2-C14.8)												
White, male												
2004.....	142	160	165	160	165	170	151	94	47	11	-	-
2003.....	141	169	193	157	158	164	127	91	41	11	2	-
2002.....	129	181	183	168	202	177	142	92	41	9	2	-
2001.....	126	155	159	173	191	178	147	81	36	10	2	-
2000.....	119	131	136	171	180	177	120	77	45	13	-	-
1999.....	112	124	170	156	181	178	137	76	39	8	1	-
White, female												
2004.....	38	36	63	73	109	128	136	119	76	31	6	-
2003.....	22	55	56	76	86	126	145	135	66	31	7	-
2002.....	28	44	56	70	110	133	141	112	86	27	7	-
2001.....	34	47	51	74	121	151	145	100	80	30	1	-
2000.....	39	35	61	94	123	126	115	133	76	30	5	-
1999.....	28	52	53	83	100	148	152	138	80	29	7	-
Black, both sexes												
2004.....	45	43	31	29	24	23	19	7	8	1	1	-
2003.....	41	38	46	31	33	28	20	10	8	1	-	-
2002.....	43	45	36	34	29	24	14	13	6	2	-	-
2001.....	25	53	41	49	41	26	10	9	9	1	-	-
2000.....	48	40	37	38	30	21	22	8	5	3	1	-
1999.....	39	43	50	37	32	24	17	19	1	-	-	-
Black, male												
2004.....	35	36	19	22	19	15	12	3	6	1	-	-
2003.....	35	32	40	19	27	15	10	4	3	-	-	-
2002.....	36	39	28	25	22	11	6	7	2	-	-	-
2001.....	19	41	37	28	30	17	5	4	2	-	-	-
2000.....	42	34	29	32	20	14	14	1	1	-	-	-
1999.....	26	31	39	28	18	15	10	9	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Of other and unspecified sites within the lip, oral cavity and pharynx (C03-C09,C14.2-C14.8)

Asian or Pacific Islander 2/, both sexes

2004.....	1	5	6	3	7	6	6	5	-	1	-	-
2003.....	3	3	6	3	3	11	3	2	1	1	-	-
2002.....	2	6	1	3	2	3	4	1	2	1	-	-
2001.....	3	2	4	2	4	2	1	7	-	-	-	-
2000.....	3	2	5	5	5	3	1	4	2	-	-	-
1999.....	2	3	3	2	4	7	2	2	1	1	-	-

Asian or Pacific Islander 2/, male

2004.....	1	5	6	2	5	5	2	1	-	-	-	-
2003.....	2	3	4	2	2	10	1	-	1	1	-	-
2002.....	2	3	1	1	2	1	2	-	-	1	-	-
2001.....	3	1	2	-	3	2	1	3	-	-	-	-
2000.....	2	2	2	5	2	2	1	2	-	-	-	-
1999.....	1	3	2	1	3	4	1	1	1	-	-	-

Asian or Pacific Islander 2/, female

2004.....	-	-	-	1	2	1	4	4	-	1	-	-
2003.....	1	-	2	1	1	1	2	2	-	-	-	-
2002.....	-	3	-	2	-	2	2	1	2	-	-	-
2001.....	-	1	2	2	1	-	-	4	-	-	-	-
2000.....	1	-	3	-	3	1	-	2	2	-	-	-
1999.....	1	-	1	1	1	3	1	1	-	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant neoplasms of digestive organs (C15-C26)												
All races, both sexes												
2004.....	7,663	10,347	12,492	14,212	17,258	20,067	19,224	13,187	6,459	1,755	239	1
2003.....	7,627	9,903	12,155	14,430	17,974	20,643	19,230	13,313	6,468	1,743	214	1
2002.....	7,422	9,719	11,731	14,603	18,472	20,915	18,810	13,161	6,357	1,722	226	3
2001.....	7,310	8,996	11,357	14,731	18,922	20,955	18,652	13,151	6,399	1,716	220	2
2000.....	6,817	8,750	11,412	14,654	19,265	21,513	18,523	13,097	6,306	1,769	234	1
1999.....	6,441	8,567	11,400	15,051	19,385	21,215	17,930	12,982	6,165	1,697	233	3
All races, male												
2004.....	5,118	6,774	7,890	8,747	10,035	10,823	9,267	5,445	2,083	445	45	-
2003.....	5,003	6,453	7,670	8,877	10,392	11,042	9,361	5,356	2,070	426	48	1
2002.....	4,915	6,312	7,504	8,972	10,747	11,193	9,024	5,265	2,018	425	46	2
2001.....	4,751	5,791	7,199	8,977	10,808	11,200	8,854	5,233	1,999	422	52	2
2000.....	4,425	5,609	7,154	8,926	10,982	11,477	8,709	5,168	1,879	441	47	1
1999.....	4,172	5,490	7,048	9,052	11,099	11,283	8,513	5,017	1,943	408	40	1
All races, female												
2004.....	2,545	3,573	4,602	5,465	7,223	9,244	9,957	7,742	4,376	1,310	194	1
2003.....	2,624	3,450	4,485	5,553	7,582	9,601	9,869	7,957	4,398	1,317	166	-
2002.....	2,507	3,407	4,227	5,631	7,725	9,722	9,786	7,896	4,339	1,297	180	1
2001.....	2,559	3,205	4,158	5,754	8,114	9,755	9,798	7,918	4,400	1,294	168	-
2000.....	2,392	3,141	4,258	5,728	8,283	10,036	9,814	7,929	4,427	1,328	187	-
1999.....	2,269	3,077	4,352	5,999	8,286	9,932	9,417	7,965	4,222	1,289	193	2
White, both sexes												
2004.....	5,884	8,190	10,138	11,673	14,454	17,386	17,007	11,775	5,760	1,543	195	1
2003.....	5,726	7,907	9,854	11,977	15,187	17,938	16,981	11,835	5,801	1,507	174	1
2002.....	5,715	7,751	9,532	12,018	15,701	18,162	16,603	11,788	5,643	1,499	186	1
2001.....	5,666	7,138	9,265	12,123	16,117	18,210	16,477	11,816	5,696	1,504	178	2
2000.....	5,297	6,962	9,276	12,165	16,458	18,754	16,388	11,709	5,634	1,542	189	1
1999.....	5,000	6,758	9,271	12,588	16,613	18,502	15,911	11,598	5,548	1,495	179	2

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant neoplasms of digestive organs (C15-C26)												
White, male												
2004.....	3,995	5,407	6,477	7,264	8,476	9,481	8,269	4,923	1,848	384	39	-
2003.....	3,795	5,237	6,273	7,454	8,915	9,691	8,365	4,792	1,855	360	42	1
2002.....	3,818	5,090	6,161	7,510	9,250	9,844	8,000	4,735	1,787	370	30	1
2001.....	3,721	4,646	5,939	7,481	9,343	9,858	7,887	4,723	1,779	373	42	2
2000.....	3,465	4,459	5,874	7,529	9,474	10,117	7,755	4,642	1,666	367	35	1
1999.....	3,250	4,352	5,775	7,661	9,659	9,887	7,630	4,481	1,754	361	35	-
White, female												
2004.....	1,889	2,783	3,661	4,409	5,978	7,905	8,738	6,852	3,912	1,159	156	1
2003.....	1,931	2,670	3,581	4,523	6,272	8,247	8,616	7,043	3,946	1,147	132	-
2002.....	1,897	2,661	3,371	4,508	6,451	8,318	8,603	7,053	3,856	1,129	156	-
2001.....	1,945	2,492	3,326	4,642	6,774	8,352	8,590	7,093	3,917	1,131	136	-
2000.....	1,832	2,503	3,402	4,636	6,984	8,637	8,633	7,067	3,968	1,175	154	-
1999.....	1,750	2,406	3,496	4,927	6,954	8,615	8,281	7,117	3,794	1,134	144	2
Black, both sexes												
2004.....	1,411	1,724	1,855	2,035	2,171	2,081	1,692	1,093	555	175	36	-
2003.....	1,552	1,611	1,846	1,920	2,169	2,111	1,773	1,160	555	195	34	-
2002.....	1,357	1,597	1,747	2,047	2,207	2,212	1,720	1,118	582	177	31	2
2001.....	1,322	1,524	1,683	2,081	2,213	2,189	1,754	1,091	595	182	39	-
2000.....	1,274	1,426	1,733	2,002	2,266	2,228	1,713	1,143	559	198	40	-
1999.....	1,172	1,481	1,721	2,009	2,266	2,217	1,677	1,175	529	172	47	1
Black, male												
2004.....	890	1,085	1,103	1,181	1,206	1,008	748	365	168	42	4	-
2003.....	1,001	989	1,112	1,092	1,176	1,024	742	410	162	53	3	-
2002.....	872	991	1,045	1,156	1,183	1,065	786	402	172	40	12	1
2001.....	822	935	983	1,204	1,145	1,054	751	387	162	38	9	-
2000.....	802	923	1,025	1,113	1,207	1,076	747	405	152	64	11	-
1999.....	751	938	1,022	1,128	1,172	1,145	706	424	153	32	2	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant neoplasms of digestive organs (C15-C26)												
Black, female												
2004.....	521	639	752	854	965	1,073	944	728	387	133	32	-
2003.....	551	622	734	828	993	1,087	1,031	750	393	142	31	-
2002.....	485	606	702	891	1,024	1,147	934	716	410	137	19	1
2001.....	500	589	700	877	1,068	1,135	1,003	704	433	144	30	-
2000.....	472	503	708	889	1,059	1,152	966	738	407	134	29	-
1999.....	421	543	699	881	1,094	1,072	971	751	376	140	45	-
American Indian 1/, both sexes												
2004.....	57	76	99	86	95	78	73	33	15	5	-	-
2003.....	55	50	70	76	80	74	47	26	18	4	1	-
2002.....	60	62	74	78	91	65	73	30	15	5	1	-
2001.....	43	56	60	73	97	73	54	33	14	1	-	-
2000.....	41	64	60	65	76	60	53	32	15	6	1	-
1999.....	36	45	51	65	54	68	54	31	15	2	-	-
American Indian 1/, male												
2004.....	41	51	53	48	51	43	32	16	6	2	-	-
2003.....	35	26	40	54	45	43	22	16	8	1	-	-
2002.....	37	41	46	44	47	29	35	11	3	-	1	-
2001.....	32	34	34	45	58	29	28	12	9	-	-	-
2000.....	27	43	38	40	43	31	22	12	6	3	-	-
1999.....	18	25	31	41	30	34	27	12	5	1	-	-
American Indian 1/, female												
2004.....	16	25	46	38	44	35	41	17	9	3	-	-
2003.....	20	24	30	22	35	31	25	10	10	3	1	-
2002.....	23	21	28	34	44	36	38	19	12	5	-	-
2001.....	11	22	26	28	39	44	26	21	5	1	-	-
2000.....	14	21	22	25	33	29	31	20	9	3	1	-
1999.....	18	20	20	24	24	34	27	19	10	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant neoplasms of digestive organs (C15-C26)												
Asian or Pacific Islander 2/, both sexes												
2004.....	311	357	400	418	538	522	452	286	129	32	8	-
2003.....	294	335	385	457	538	520	429	292	94	37	5	-
2002.....	290	309	378	460	473	476	414	225	117	41	8	-
2001.....	279	278	349	454	495	483	367	211	94	29	3	-
2000.....	205	298	343	422	465	471	369	213	98	23	4	-
1999.....	233	283	357	389	452	428	288	178	73	28	7	-
Asian or Pacific Islander 2/, male												
2004.....	192	231	257	254	302	291	218	141	61	17	2	-
2003.....	172	201	245	277	256	284	232	138	45	12	3	-
2002.....	188	190	252	262	267	255	203	117	56	15	3	-
2001.....	176	176	243	247	262	259	188	111	49	11	1	-
2000.....	131	184	217	244	258	253	185	109	55	7	1	-
1999.....	153	175	220	222	238	217	150	100	31	14	3	-
Asian or Pacific Islander 2/, female												
2004.....	119	126	143	164	236	231	234	145	68	15	6	-
2003.....	122	134	140	180	282	236	197	154	49	25	2	-
2002.....	102	119	126	198	206	221	211	108	61	26	5	-
2001.....	103	102	106	207	233	224	179	100	45	18	2	-
2000.....	74	114	126	178	207	218	184	104	43	16	3	-
1999.....	80	108	137	167	214	211	138	78	42	14	4	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Of esophagus (C15)												
All races, both sexes												
2004.....	910	1,334	1,657	1,739	1,815	1,928	1,539	879	301	67	9	-
2003.....	881	1,389	1,551	1,714	1,905	1,895	1,497	806	306	63	5	-
2002.....	887	1,286	1,477	1,736	1,955	1,909	1,480	766	295	65	4	1
2001.....	916	1,114	1,440	1,765	2,014	1,941	1,366	757	317	56	7	1
2000.....	870	1,180	1,430	1,711	1,960	1,817	1,346	764	274	55	7	-
1999.....	793	1,139	1,392	1,739	1,955	1,784	1,202	755	279	72	8	1
All races, male												
2004.....	771	1,130	1,372	1,412	1,405	1,450	1,096	555	154	21	1	-
2003.....	757	1,175	1,308	1,385	1,466	1,415	1,049	476	156	22	-	-
2002.....	759	1,099	1,222	1,394	1,558	1,437	1,017	452	127	23	1	-
2001.....	787	940	1,197	1,436	1,553	1,423	942	452	153	18	2	1
2000.....	746	988	1,181	1,349	1,482	1,337	896	437	122	23	4	-
1999.....	684	969	1,112	1,396	1,529	1,328	808	421	127	22	3	1
All races, female												
2004.....	139	204	285	327	410	478	443	324	147	46	8	-
2003.....	124	214	243	329	439	480	448	330	150	41	5	-
2002.....	128	187	255	342	397	472	463	314	168	42	3	1
2001.....	129	174	243	329	461	518	424	305	164	38	5	-
2000.....	124	192	249	362	478	480	450	327	152	32	3	-
1999.....	109	170	280	343	426	456	394	334	152	50	5	-
White, both sexes												
2004.....	739	1,106	1,382	1,491	1,572	1,740	1,398	807	276	59	7	-
2003.....	682	1,145	1,279	1,426	1,689	1,710	1,355	751	286	55	5	-
2002.....	725	1,044	1,218	1,442	1,708	1,698	1,353	704	276	61	3	-
2001.....	718	860	1,168	1,455	1,714	1,741	1,251	700	287	53	7	1
2000.....	674	932	1,150	1,441	1,705	1,599	1,218	709	253	51	7	-
1999.....	617	895	1,114	1,448	1,691	1,564	1,102	694	259	67	5	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Of esophagus (C15)												
White, male												
2004.....	646	955	1,171	1,225	1,241	1,320	1,011	510	141	18	1	-
2003.....	601	990	1,098	1,159	1,317	1,292	972	446	147	17	-	-
2002.....	633	906	1,026	1,185	1,378	1,301	935	418	121	22	-	-
2001.....	624	741	985	1,200	1,350	1,290	867	420	135	17	2	1
2000.....	588	791	975	1,158	1,308	1,186	830	412	112	19	4	-
1999.....	542	773	905	1,191	1,339	1,171	751	383	115	18	2	-
White, female												
2004.....	93	151	211	266	331	420	387	297	135	41	6	-
2003.....	81	155	181	267	372	418	383	305	139	38	5	-
2002.....	92	138	192	257	330	397	418	286	155	39	3	-
2001.....	94	119	183	255	364	451	384	280	152	36	5	-
2000.....	86	141	175	283	397	413	388	297	141	32	3	-
1999.....	75	122	209	257	352	393	351	311	144	49	3	-
Black, both sexes												
2004.....	147	202	249	223	205	162	109	52	18	4	2	-
2003.....	181	222	245	249	188	158	116	50	16	7	-	-
2002.....	145	222	235	259	222	185	100	53	13	3	1	1
2001.....	172	225	245	274	266	177	102	51	23	3	-	-
2000.....	184	228	255	244	222	203	106	48	19	4	-	-
1999.....	166	225	252	260	231	200	82	51	17	5	2	1
Black, male												
2004.....	110	151	179	165	133	109	63	31	7	1	-	-
2003.....	139	166	188	189	126	104	59	26	7	4	-	-
2002.....	111	180	174	182	163	118	63	30	3	1	1	-
2001.....	141	176	191	204	178	119	67	27	13	1	-	-
2000.....	150	185	186	172	149	139	54	21	8	4	-	-
1999.....	133	179	190	179	165	143	42	30	10	4	-	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Of esophagus (C15)												
Black, female												
2004.....	37	51	70	58	72	53	46	21	11	3	2	-
2003.....	42	56	57	60	62	54	57	24	9	3	-	-
2002.....	34	42	61	77	59	67	37	23	10	2	-	1
2001.....	31	49	54	70	88	58	35	24	10	2	-	-
2000.....	34	43	69	72	73	64	52	27	11	-	-	-
1999.....	33	46	62	81	66	57	40	21	7	1	2	-
American Indian 1/, both sexes												
2004.....	2	9	8	11	13	1	2	4	2	-	-	-
2003.....	5	6	6	9	6	5	6	-	-	-	-	-
2002.....	4	2	7	5	7	7	9	-	2	1	-	-
2001.....	4	7	7	8	8	7	3	2	1	-	-	-
2000.....	3	3	8	6	7	3	6	1	1	-	-	-
1999.....	1	7	7	5	6	5	3	-	-	-	-	-
American Indian 1/, male												
2004.....	2	8	7	9	11	-	-	3	2	-	-	-
2003.....	5	5	2	8	5	5	3	-	-	-	-	-
2002.....	4	1	7	4	5	4	8	-	2	-	-	-
2001.....	3	6	4	6	7	4	1	1	1	-	-	-
2000.....	3	1	8	5	7	2	4	-	1	-	-	-
1999.....	1	6	4	5	6	4	1	-	-	-	-	-
American Indian 1/, female												
2004.....	-	1	1	2	2	1	2	1	-	-	-	-
2003.....	-	1	4	1	1	-	3	-	-	-	-	-
2002.....	-	1	-	1	2	3	1	-	-	1	-	-
2001.....	1	1	3	2	1	3	2	1	-	-	-	-
2000.....	-	2	-	1	-	1	2	1	-	-	-	-
1999.....	-	1	3	-	-	1	2	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Of esophagus (C15)												
Asian or Pacific Islander 2/, both sexes												
2004.....	22	17	18	14	25	25	30	16	5	4	-	-
2003.....	13	16	21	30	22	22	20	5	4	1	-	-
2002.....	13	18	17	30	18	19	18	9	4	-	-	-
2001.....	22	22	20	28	26	16	10	4	6	-	-	-
2000.....	9	17	17	20	26	12	16	6	1	-	-	-
1999.....	9	12	19	26	27	15	15	10	3	-	1	-
Asian or Pacific Islander 2/, male												
2004.....	13	16	15	13	20	21	22	11	4	2	-	-
2003.....	12	14	20	29	18	14	15	4	2	1	-	-
2002.....	11	12	15	23	12	14	11	4	1	-	-	-
2001.....	19	17	17	26	18	10	7	4	4	-	-	-
2000.....	5	11	12	14	18	10	8	4	1	-	-	-
1999.....	8	11	13	21	19	10	14	8	2	-	1	-
Asian or Pacific Islander 2/, female												
2004.....	9	1	3	1	5	4	8	5	1	2	-	-
2003.....	1	2	1	1	4	8	5	1	2	-	-	-
2002.....	2	6	2	7	6	5	7	5	3	-	-	-
2001.....	3	5	3	2	8	6	3	-	2	-	-	-
2000.....	4	6	5	6	8	2	8	2	-	-	-	-
1999.....	1	1	6	5	8	5	1	2	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Of stomach (C16)												
All races, both sexes												
2004.....	685	882	1,005	1,202	1,550	1,733	1,682	1,255	622	151	23	-
2003.....	717	830	1,044	1,211	1,582	1,805	1,744	1,263	619	161	14	-
2002.....	663	841	990	1,179	1,637	1,903	1,753	1,344	664	153	24	-
2001.....	675	751	1,020	1,308	1,713	2,004	1,789	1,216	620	149	16	-
2000.....	629	753	1,051	1,317	1,737	2,009	1,824	1,379	641	188	26	-
1999.....	651	808	1,052	1,359	1,793	2,064	1,796	1,277	607	170	29	-
All races, male												
2004.....	442	601	672	788	946	1,041	900	659	240	43	7	-
2003.....	462	574	708	810	1,016	1,061	919	574	238	51	4	-
2002.....	440	556	659	804	1,063	1,118	946	628	258	38	10	-
2001.....	443	514	724	858	1,070	1,194	954	550	222	36	5	-
2000.....	413	506	722	860	1,106	1,201	943	617	247	56	8	-
1999.....	443	558	694	893	1,112	1,250	989	568	223	52	8	-
All races, female												
2004.....	243	281	333	414	604	692	782	596	382	108	16	-
2003.....	255	256	336	401	566	744	825	689	381	110	10	-
2002.....	223	285	331	375	574	785	807	716	406	115	14	-
2001.....	232	237	296	450	643	810	835	666	398	113	11	-
2000.....	216	247	329	457	631	808	881	762	394	132	18	-
1999.....	208	250	358	466	681	814	807	709	384	118	21	-
White, both sexes												
2004.....	475	634	742	872	1,178	1,371	1,365	1,041	525	118	14	-
2003.....	497	622	788	922	1,227	1,420	1,403	1,026	498	114	8	-
2002.....	453	624	730	864	1,260	1,496	1,406	1,086	537	115	14	-
2001.....	473	565	750	964	1,350	1,595	1,441	1,008	503	116	11	-
2000.....	463	561	791	988	1,348	1,571	1,465	1,129	525	149	13	-
1999.....	451	583	765	1,066	1,393	1,636	1,462	1,039	497	131	19	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Of stomach (C16)												
White, male												
2004.....	314	443	495	579	712	831	727	553	201	35	6	-
2003.....	323	436	531	610	794	843	749	459	200	36	4	-
2002.....	307	427	491	605	834	904	773	521	207	29	4	-
2001.....	312	385	536	627	850	973	771	463	183	30	3	-
2000.....	311	379	557	660	874	952	770	510	205	45	4	-
1999.....	312	415	508	713	880	1,004	807	452	184	42	6	-
White, female												
2004.....	161	191	247	293	466	540	638	488	324	83	8	-
2003.....	174	186	257	312	433	577	654	567	298	78	4	-
2002.....	146	197	239	259	426	592	633	565	330	86	10	-
2001.....	161	180	214	337	500	622	670	545	320	86	8	-
2000.....	152	182	234	328	474	619	695	619	320	104	9	-
1999.....	139	168	257	353	513	632	655	587	313	89	13	-
Black, both sexes												
2004.....	154	178	193	264	260	259	242	153	68	25	7	-
2003.....	161	152	198	213	250	287	246	157	96	35	6	-
2002.....	158	155	185	232	290	307	241	200	98	31	9	-
2001.....	150	130	198	251	258	294	258	161	94	27	5	-
2000.....	130	149	189	243	286	320	266	197	86	30	12	-
1999.....	151	173	205	214	298	328	257	194	86	32	9	-
Black, male												
2004.....	97	116	135	168	173	140	130	68	28	5	-	-
2003.....	108	106	135	148	165	164	118	71	22	9	-	-
2002.....	106	102	121	148	179	160	118	75	41	7	5	-
2001.....	106	88	135	172	161	163	133	59	25	6	2	-
2000.....	83	100	125	148	171	176	125	80	27	8	3	-
1999.....	99	112	133	135	178	188	139	89	28	7	2	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Of stomach (C16)												
Black, female												
2004.....	57	62	58	96	87	119	112	85	40	20	7	-
2003.....	53	46	63	65	85	123	128	86	74	26	6	-
2002.....	52	53	64	84	111	147	123	125	57	24	4	-
2001.....	44	42	63	79	97	131	125	102	69	21	3	-
2000.....	47	49	64	95	115	144	141	117	59	22	9	-
1999.....	52	61	72	79	120	140	118	105	58	25	7	-
American Indian 1/, both sexes												
2004.....	4	5	11	4	16	11	4	6	2	1	-	-
2003.....	8	3	7	11	11	12	4	5	2	1	-	-
2002.....	10	5	11	11	10	13	12	4	2	-	-	-
2001.....	3	7	8	6	11	18	10	4	4	-	-	-
2000.....	5	7	9	15	10	9	6	6	2	-	-	-
1999.....	2	6	5	12	11	9	7	1	4	-	-	-
American Indian 1/, male												
2004.....	3	3	5	3	10	10	3	3	-	-	-	-
2003.....	6	2	4	9	8	5	3	3	2	-	-	-
2002.....	5	1	4	7	5	5	7	3	-	-	-	-
2001.....	2	7	6	4	6	7	7	2	4	-	-	-
2000.....	4	5	4	8	5	6	2	3	1	-	-	-
1999.....	1	3	4	8	7	5	6	1	3	-	-	-
American Indian 1/, female												
2004.....	1	2	6	1	6	1	1	3	2	1	-	-
2003.....	2	1	3	2	3	7	1	2	-	1	-	-
2002.....	5	4	7	4	5	8	5	1	2	-	-	-
2001.....	1	-	2	2	5	11	3	2	-	-	-	-
2000.....	1	2	5	7	5	3	4	3	1	-	-	-
1999.....	1	3	1	4	4	4	1	-	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Of stomach (C16)												
Asian or Pacific Islander 2/, both sexes												
2004.....	52	65	59	62	96	92	71	55	27	7	2	-
2003.....	51	53	51	65	94	86	91	75	23	11	-	-
2002.....	42	57	64	72	77	87	94	54	27	7	1	-
2001.....	49	49	64	87	94	97	80	43	19	6	-	-
2000.....	31	36	62	71	93	109	87	47	28	9	1	-
1999.....	47	46	77	67	91	91	70	43	20	7	1	-
Asian or Pacific Islander 2/, male												
2004.....	28	39	37	38	51	60	40	35	11	3	1	-
2003.....	25	30	38	43	49	49	49	41	14	6	-	-
2002.....	22	26	43	44	45	49	48	29	10	2	1	-
2001.....	23	34	47	55	53	51	43	26	10	-	-	-
2000.....	15	22	36	44	56	67	46	24	14	3	1	-
1999.....	31	28	49	37	47	53	37	26	8	3	-	-
Asian or Pacific Islander 2/, female												
2004.....	24	26	22	24	45	32	31	20	16	4	1	-
2003.....	26	23	13	22	45	37	42	34	9	5	-	-
2002.....	20	31	21	28	32	38	46	25	17	5	-	-
2001.....	26	15	17	32	41	46	37	17	9	6	-	-
2000.....	16	14	26	27	37	42	41	23	14	6	-	-
1999.....	16	18	28	30	44	38	33	17	12	4	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Of small intestine (C17)												
All races, both sexes												
2004.....	76	89	118	141	134	155	152	104	35	8	2	-
2003.....	63	102	107	105	139	155	164	95	42	3	-	-
2002.....	67	87	86	119	138	155	139	81	41	6	-	-
2001.....	65	75	116	99	165	178	132	94	43	4	-	-
2000.....	64	79	106	118	152	161	140	81	32	14	-	-
1999.....	79	56	95	113	150	164	126	92	32	6	1	1
All races, male												
2004.....	36	57	67	82	78	75	82	43	18	3	1	-
2003.....	28	66	60	66	75	69	71	39	10	2	-	-
2002.....	38	47	57	61	75	82	58	36	14	2	-	-
2001.....	42	38	67	59	96	84	64	38	13	2	-	-
2000.....	38	42	61	63	84	77	67	39	12	1	-	-
1999.....	45	35	70	66	79	89	64	39	12	1	-	-
All races, female												
2004.....	40	32	51	59	56	80	70	61	17	5	1	-
2003.....	35	36	47	39	64	86	93	56	32	1	-	-
2002.....	29	40	29	58	63	73	81	45	27	4	-	-
2001.....	23	37	49	40	69	94	68	56	30	2	-	-
2000.....	26	37	45	55	68	84	73	42	20	13	-	-
1999.....	34	21	25	47	71	75	62	53	20	5	1	1
White, both sexes												
2004.....	56	70	90	118	109	133	137	89	30	5	2	-
2003.....	50	80	88	89	123	134	144	83	36	3	-	-
2002.....	56	68	64	97	123	132	122	69	36	5	-	-
2001.....	46	55	91	80	141	157	116	80	37	4	-	-
2000.....	49	65	90	96	135	143	122	77	28	13	-	-
1999.....	61	44	78	100	128	139	105	79	28	6	1	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Of small intestine (C17)

White, male

2004.....	29	46	46	70	65	65	73	38	16	1	1	-
2003.....	23	55	49	54	62	59	63	34	7	2	-	-
2002.....	33	36	42	51	66	72	51	32	11	2	-	-
2001.....	33	28	53	47	85	73	58	31	11	2	-	-
2000.....	29	32	53	55	74	66	60	38	9	1	-	-
1999.....	34	27	58	59	66	76	54	37	10	1	-	-

White, female

2004.....	27	24	44	48	44	68	64	51	14	4	1	-
2003.....	27	25	39	35	61	75	81	49	29	1	-	-
2002.....	23	32	22	46	57	60	71	37	25	3	-	-
2001.....	13	27	38	33	56	84	58	49	26	2	-	-
2000.....	20	33	37	41	61	77	62	39	19	12	-	-
1999.....	27	17	20	41	62	63	51	42	18	5	1	1

Black, both sexes

2004.....	18	17	23	19	23	20	11	14	3	2	-	-
2003.....	12	20	16	14	12	18	15	10	5	-	-	-
2002.....	11	17	17	17	13	19	14	12	4	-	-	-
2001.....	16	16	23	17	20	17	13	13	5	-	-	-
2000.....	15	12	15	19	14	12	13	4	2	-	-	-
1999.....	15	9	15	11	17	22	18	13	4	-	-	-

Black, male

2004.....	6	10	17	9	13	8	7	4	-	1	-	-
2003.....	5	11	8	10	10	8	5	4	3	-	-	-
2002.....	5	9	10	7	7	8	7	4	2	-	-	-
2001.....	7	9	12	10	7	9	5	6	1	-	-	-
2000.....	9	8	8	7	9	6	6	1	1	-	-	-
1999.....	9	6	12	7	11	12	8	2	2	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Of small intestine (C17)												
Asian or Pacific Islander 2/, both sexes												
2004.....	2	2	5	3	2	1	2	1	2	1	-	-
2003.....	-	2	3	-	3	2	4	1	-	-	-	-
2002.....	-	2	3	5	1	3	2	-	1	1	-	-
2001.....	3	3	1	1	3	4	2	1	1	-	-	-
2000.....	-	2	1	3	3	5	4	-	2	-	-	-
1999.....	2	3	2	2	5	3	3	-	-	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	1	1	4	3	-	1	1	1	2	1	-	-
2003.....	-	-	3	-	2	1	2	1	-	-	-	-
2002.....	-	2	3	3	1	1	-	-	1	-	-	-
2001.....	2	1	1	1	3	2	-	1	1	-	-	-
2000.....	-	2	-	1	1	4	1	-	2	-	-	-
1999.....	1	2	-	-	2	1	2	-	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	1	1	1	-	2	-	1	-	-	-	-	-
2003.....	-	2	-	-	1	1	2	-	-	-	-	-
2002.....	-	-	-	2	-	2	2	-	-	1	-	-
2001.....	1	2	-	-	-	2	2	-	-	-	-	-
2000.....	-	-	1	2	2	1	3	-	-	-	-	-
1999.....	1	1	2	2	3	2	1	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Of colon, rectum and anus (C18-C21)												
All races, both sexes												
2004.....	2,636	3,691	4,517	5,297	6,657	8,245	8,469	6,337	3,474	1,023	129	-
2003.....	2,718	3,631	4,673	5,775	7,159	8,720	8,611	6,590	3,547	1,024	135	1
2002.....	2,818	3,694	4,727	5,901	7,491	8,826	8,672	6,453	3,433	1,075	128	1
2001.....	2,804	3,610	4,374	5,997	7,794	8,988	8,539	6,618	3,482	1,028	131	-
2000.....	2,581	3,397	4,543	5,913	8,188	9,410	8,643	6,735	3,518	1,061	148	-
1999.....	2,434	3,309	4,658	6,095	8,154	9,473	8,456	6,672	3,503	973	138	-
All races, male												
2004.....	1,531	2,161	2,624	3,074	3,718	4,307	3,882	2,464	1,092	264	20	-
2003.....	1,544	2,117	2,714	3,393	3,953	4,433	4,038	2,565	1,112	245	27	1
2002.....	1,606	2,185	2,850	3,455	4,125	4,536	3,968	2,508	1,024	267	24	1
2001.....	1,545	2,116	2,536	3,479	4,294	4,568	3,883	2,520	1,068	251	36	-
2000.....	1,466	1,989	2,619	3,407	4,517	4,852	3,908	2,571	1,012	260	30	-
1999.....	1,374	1,932	2,689	3,431	4,452	4,857	3,897	2,525	1,081	236	24	-
All races, female												
2004.....	1,105	1,530	1,893	2,223	2,939	3,938	4,587	3,873	2,382	759	109	-
2003.....	1,174	1,514	1,959	2,382	3,206	4,287	4,573	4,025	2,435	779	108	-
2002.....	1,212	1,509	1,877	2,446	3,366	4,290	4,704	3,945	2,409	808	104	-
2001.....	1,259	1,494	1,838	2,518	3,500	4,420	4,656	4,098	2,414	777	95	-
2000.....	1,115	1,408	1,924	2,506	3,671	4,558	4,735	4,164	2,506	801	118	-
1999.....	1,060	1,377	1,969	2,664	3,702	4,616	4,559	4,147	2,422	737	114	-
White, both sexes												
2004.....	2,052	2,985	3,693	4,376	5,643	7,195	7,536	5,710	3,128	913	109	-
2003.....	2,060	2,898	3,836	4,853	6,057	7,625	7,705	5,911	3,212	914	114	1
2002.....	2,240	2,985	3,873	4,933	6,460	7,731	7,686	5,857	3,073	955	110	1
2001.....	2,211	2,906	3,609	5,004	6,722	7,831	7,618	6,015	3,146	908	106	-
2000.....	2,050	2,737	3,752	4,991	7,094	8,322	7,710	6,090	3,180	933	129	-
1999.....	1,935	2,676	3,861	5,145	7,083	8,370	7,574	6,030	3,198	864	109	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Of colon, rectum and anus (C18-C21)												
White, male												
2004.....	1,211	1,786	2,174	2,572	3,184	3,771	3,477	2,246	982	231	18	-
2003.....	1,195	1,730	2,245	2,891	3,407	3,911	3,641	2,326	1,002	215	23	1
2002.....	1,287	1,796	2,360	2,932	3,615	3,998	3,521	2,270	916	237	18	1
2001.....	1,254	1,735	2,132	2,949	3,766	4,036	3,486	2,288	962	217	30	-
2000.....	1,182	1,607	2,181	2,906	3,955	4,332	3,485	2,327	905	218	23	-
1999.....	1,119	1,574	2,254	2,943	3,948	4,303	3,529	2,279	990	211	22	-
White, female												
2004.....	841	1,199	1,519	1,804	2,459	3,424	4,059	3,464	2,146	682	91	-
2003.....	865	1,168	1,591	1,962	2,650	3,714	4,064	3,585	2,210	699	91	-
2002.....	953	1,189	1,513	2,001	2,845	3,733	4,165	3,587	2,157	718	92	-
2001.....	957	1,171	1,477	2,055	2,956	3,795	4,132	3,727	2,184	691	76	-
2000.....	868	1,130	1,571	2,085	3,139	3,990	4,225	3,763	2,275	715	106	-
1999.....	816	1,102	1,607	2,202	3,135	4,067	4,045	3,751	2,208	653	87	-
Black, both sexes												
2004.....	489	606	693	774	865	878	764	517	294	96	18	-
2003.....	551	622	707	774	942	909	779	563	291	93	18	-
2002.....	493	596	712	817	857	934	836	509	309	94	13	-
2001.....	495	611	651	842	894	994	779	509	296	105	23	-
2000.....	464	538	687	787	945	942	801	561	295	117	18	-
1999.....	414	551	705	819	927	965	785	576	268	92	24	-
Black, male												
2004.....	269	318	373	415	454	437	334	170	85	24	1	-
2003.....	300	329	396	417	477	427	328	184	90	28	2	-
2002.....	269	319	401	438	414	455	373	197	86	19	4	-
2001.....	244	328	333	449	436	448	328	188	86	27	5	-
2000.....	251	312	369	420	479	442	354	199	82	39	7	-
1999.....	218	319	379	426	435	480	315	207	77	15	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Of colon, rectum and anus (C18-C21)

Black, female

2004.....	220	288	320	359	411	441	430	347	209	72	17	-
2003.....	251	293	311	357	465	482	451	379	201	65	16	-
2002.....	224	277	311	379	443	479	463	312	223	75	9	-
2001.....	251	283	318	393	458	546	451	321	210	78	18	-
2000.....	213	226	318	367	466	500	447	362	213	78	11	-
1999.....	196	232	326	393	492	485	470	369	191	77	24	-

American Indian 1/, both sexes

2004.....	14	26	28	35	26	31	27	9	5	2	-	-
2003.....	15	13	25	24	25	33	18	13	9	2	-	-
2002.....	19	31	33	34	34	25	21	14	7	2	1	-
2001.....	16	17	25	22	39	20	22	12	4	-	-	-
2000.....	11	22	20	22	29	27	16	13	8	3	-	-
1999.....	18	14	16	18	19	24	22	14	7	1	-	-

American Indian 1/, male

2004.....	9	15	20	19	14	20	12	5	2	1	-	-
2003.....	6	7	13	17	13	17	7	10	5	-	-	-
2002.....	12	23	24	17	16	11	10	3	1	-	1	-
2001.....	11	10	10	14	19	6	10	6	2	-	-	-
2000.....	5	17	13	14	16	15	7	6	3	1	-	-
1999.....	6	7	10	13	7	13	11	6	2	1	-	-

American Indian 1/, female

2004.....	5	11	8	16	12	11	15	4	3	1	-	-
2003.....	9	6	12	7	12	16	11	3	4	2	-	-
2002.....	7	8	9	17	18	14	11	11	6	2	-	-
2001.....	5	7	15	8	20	14	12	6	2	-	-	-
2000.....	6	5	7	8	13	12	9	7	5	2	-	-
1999.....	12	7	6	5	12	11	11	8	5	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Of colon, rectum and anus (C18-C21)												
Asian or Pacific Islander 2/, both sexes												
2004.....	81	74	103	112	123	141	142	101	47	12	2	-
2003.....	92	98	105	124	135	153	109	103	35	15	3	-
2002.....	66	82	109	117	140	136	129	73	44	24	4	-
2001.....	82	76	89	129	139	143	120	82	36	15	2	-
2000.....	56	100	84	113	120	119	116	71	35	8	1	-
1999.....	67	68	76	113	125	114	75	52	30	16	5	-
Asian or Pacific Islander 2/, male												
2004.....	42	42	57	68	66	79	59	43	23	8	1	-
2003.....	43	51	60	68	56	78	62	45	15	2	2	-
2002.....	38	47	65	68	80	72	64	38	21	11	1	-
2001.....	36	43	61	67	73	78	59	38	18	7	1	-
2000.....	28	53	56	67	67	63	62	39	22	2	-	-
1999.....	31	32	46	49	62	61	42	33	12	9	2	-
Asian or Pacific Islander 2/, female												
2004.....	39	32	46	44	57	62	83	58	24	4	1	-
2003.....	49	47	45	56	79	75	47	58	20	13	1	-
2002.....	28	35	44	49	60	64	65	35	23	13	3	-
2001.....	46	33	28	62	66	65	61	44	18	8	1	-
2000.....	28	47	28	46	53	56	54	32	13	6	1	-
1999.....	36	36	30	64	63	53	33	19	18	7	3	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Colon (C18)												
All races, both sexes												
2004.....	2,046	2,900	3,580	4,348	5,586	6,968	7,280	5,485	3,006	890	109	-
2003.....	2,131	2,830	3,768	4,817	6,019	7,438	7,455	5,711	3,104	893	113	1
2002.....	2,229	2,968	3,769	4,835	6,321	7,531	7,563	5,647	2,995	946	116	1
2001.....	2,168	2,901	3,602	4,994	6,640	7,717	7,388	5,756	3,025	869	109	-
2000.....	2,080	2,714	3,718	4,902	6,927	8,033	7,599	5,913	3,066	922	128	-
1999.....	1,969	2,663	3,843	5,085	6,900	8,128	7,402	5,857	3,080	831	124	-
All races, male												
2004.....	1,184	1,639	2,049	2,490	3,051	3,586	3,289	2,135	952	213	16	-
2003.....	1,196	1,626	2,137	2,816	3,271	3,758	3,465	2,209	969	211	22	1
2002.....	1,242	1,726	2,221	2,805	3,466	3,831	3,413	2,187	886	242	24	1
2001.....	1,166	1,680	2,035	2,843	3,602	3,862	3,338	2,183	914	207	31	-
2000.....	1,169	1,559	2,100	2,760	3,769	4,117	3,395	2,244	886	230	24	-
1999.....	1,092	1,527	2,197	2,831	3,739	4,139	3,374	2,192	964	191	21	-
All races, female												
2004.....	862	1,261	1,531	1,858	2,535	3,382	3,991	3,350	2,054	677	93	-
2003.....	935	1,204	1,631	2,001	2,748	3,680	3,990	3,502	2,135	682	91	-
2002.....	987	1,242	1,548	2,030	2,855	3,700	4,150	3,460	2,109	704	92	-
2001.....	1,002	1,221	1,567	2,151	3,038	3,855	4,050	3,573	2,111	662	78	-
2000.....	911	1,155	1,618	2,142	3,158	3,916	4,204	3,669	2,180	692	104	-
1999.....	877	1,136	1,646	2,254	3,161	3,989	4,028	3,665	2,116	640	103	-
White, both sexes												
2004.....	1,567	2,312	2,899	3,559	4,689	6,044	6,457	4,934	2,718	789	93	-
2003.....	1,596	2,236	3,068	4,027	5,071	6,496	6,656	5,108	2,804	799	96	1
2002.....	1,749	2,380	3,060	4,008	5,425	6,569	6,713	5,121	2,675	843	100	1
2001.....	1,686	2,314	2,938	4,124	5,707	6,681	6,589	5,227	2,736	765	88	-
2000.....	1,636	2,170	3,051	4,132	5,983	7,081	6,755	5,336	2,771	803	110	-
1999.....	1,559	2,130	3,168	4,277	5,970	7,161	6,618	5,294	2,805	734	97	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Colon (C18)												
White, male												
2004.....	918	1,333	1,683	2,071	2,586	3,117	2,938	1,947	859	184	14	-
2003.....	917	1,317	1,749	2,386	2,809	3,314	3,108	2,006	875	185	20	1
2002.....	981	1,412	1,820	2,360	3,026	3,356	3,041	1,976	793	214	18	1
2001.....	931	1,375	1,696	2,393	3,156	3,387	2,990	1,976	825	178	26	-
2000.....	936	1,256	1,734	2,360	3,297	3,664	3,014	2,026	787	190	18	-
1999.....	893	1,237	1,833	2,424	3,310	3,655	3,055	1,980	883	171	19	-
White, female												
2004.....	649	979	1,216	1,488	2,103	2,927	3,519	2,987	1,859	605	79	-
2003.....	679	919	1,319	1,641	2,262	3,182	3,548	3,102	1,929	614	76	-
2002.....	768	968	1,240	1,648	2,399	3,213	3,672	3,145	1,882	629	82	-
2001.....	755	939	1,242	1,731	2,551	3,294	3,599	3,251	1,911	587	62	-
2000.....	700	914	1,317	1,772	2,686	3,417	3,741	3,310	1,984	613	92	-
1999.....	666	893	1,335	1,853	2,660	3,506	3,563	3,314	1,922	563	78	-
Black, both sexes												
2004.....	402	513	583	667	770	779	680	458	247	87	14	-
2003.....	449	508	601	668	823	794	692	500	260	79	16	-
2002.....	414	501	600	704	748	831	727	455	273	83	11	-
2001.....	411	514	573	742	787	892	674	445	260	90	19	-
2000.....	392	452	587	661	822	836	732	508	259	110	17	-
1999.....	347	468	605	707	809	849	706	510	241	86	22	-
Black, male												
2004.....	226	264	310	349	396	382	291	149	75	20	1	-
2003.....	241	262	334	362	407	368	299	157	76	24	2	-
2002.....	222	264	335	378	357	407	312	177	73	19	4	-
2001.....	203	266	281	385	371	398	288	170	75	22	4	-
2000.....	209	252	311	334	407	390	322	180	76	38	6	-
1999.....	175	262	319	365	373	423	274	178	68	14	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Colon (C18)												
Black, female												
2004.....	176	249	273	318	374	397	389	309	172	67	13	-
2003.....	208	246	267	306	416	426	393	343	184	55	14	-
2002.....	192	237	265	326	391	424	415	278	200	64	7	-
2001.....	208	248	292	357	416	494	386	275	185	68	15	-
2000.....	183	200	276	327	415	446	410	328	183	72	11	-
1999.....	172	206	286	342	436	426	432	332	173	72	22	-
American Indian 1/, both sexes												
2004.....	12	22	21	29	26	26	21	7	3	2	-	-
2003.....	13	12	20	19	19	24	16	13	9	2	-	-
2002.....	16	20	25	27	29	22	17	11	6	1	1	-
2001.....	10	12	17	17	31	20	19	11	3	-	-	-
2000.....	8	17	16	18	24	18	13	11	6	2	-	-
1999.....	12	12	12	14	14	22	17	11	7	-	-	-
American Indian 1/, male												
2004.....	8	11	16	14	14	17	8	5	1	1	-	-
2003.....	6	7	10	13	10	12	6	10	5	-	-	-
2002.....	9	13	18	12	14	10	7	2	1	-	1	-
2001.....	5	7	8	11	15	6	8	5	1	-	-	-
2000.....	3	13	11	12	12	12	5	6	3	-	-	-
1999.....	4	5	7	9	2	13	9	6	2	-	-	-
American Indian 1/, female												
2004.....	4	11	5	15	12	9	13	2	2	1	-	-
2003.....	7	5	10	6	9	12	10	3	4	2	-	-
2002.....	7	7	7	15	15	12	10	9	5	1	-	-
2001.....	5	5	9	6	16	14	11	6	2	-	-	-
2000.....	5	4	5	6	12	6	8	5	3	2	-	-
1999.....	8	7	5	5	12	9	8	5	5	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Colon (C18)												
Asian or Pacific Islander 2/, both sexes												
2004.....	65	53	77	93	101	119	122	86	38	12	2	-
2003.....	73	74	79	103	106	124	91	90	31	13	1	-
2002.....	50	67	84	96	119	109	106	60	41	19	4	-
2001.....	61	61	74	111	115	124	106	73	26	14	2	-
2000.....	44	75	64	91	98	98	99	58	30	7	1	-
1999.....	51	53	58	87	107	96	61	42	27	11	5	-
Asian or Pacific Islander 2/, male												
2004.....	32	31	40	56	55	70	52	34	17	8	1	-
2003.....	32	40	44	55	45	64	52	36	13	2	-	-
2002.....	30	37	48	55	69	58	53	32	19	9	1	-
2001.....	27	32	50	54	60	71	52	32	13	7	1	-
2000.....	21	38	44	54	53	51	54	32	20	2	-	-
1999.....	20	23	38	33	54	48	36	28	11	6	2	-
Asian or Pacific Islander 2/, female												
2004.....	33	22	37	37	46	49	70	52	21	4	1	-
2003.....	41	34	35	48	61	60	39	54	18	11	1	-
2002.....	20	30	36	41	50	51	53	28	22	10	3	-
2001.....	34	29	24	57	55	53	54	41	13	7	1	-
2000.....	23	37	20	37	45	47	45	26	10	5	1	-
1999.....	31	30	20	54	53	48	25	14	16	5	3	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rectosigmoid junction and rectum (C19-C20)												
All races, both sexes												
2004.....	528	727	878	892	1,013	1,211	1,131	820	451	126	19	-
2003.....	536	735	850	908	1,083	1,214	1,104	837	427	125	21	-
2002.....	527	675	894	1,015	1,120	1,238	1,057	773	420	126	12	-
2001.....	583	660	730	952	1,096	1,207	1,100	824	439	154	20	-
2000.....	452	639	774	963	1,203	1,313	1,006	794	435	134	20	-
1999.....	427	603	777	966	1,196	1,291	1,008	777	405	139	14	-
All races, male												
2004.....	322	487	548	558	645	697	582	320	133	49	3	-
2003.....	333	465	558	552	664	657	565	344	143	34	5	-
2002.....	335	442	599	629	642	688	540	310	132	25	-	-
2001.....	359	415	485	615	667	683	534	329	152	43	5	-
2000.....	274	407	499	625	728	714	496	322	120	29	6	-
1999.....	266	390	478	583	695	702	508	325	113	45	3	-
All races, female												
2004.....	206	240	330	334	368	514	549	500	318	77	16	-
2003.....	203	270	292	356	419	557	539	493	284	91	16	-
2002.....	192	233	295	386	478	550	517	463	288	101	12	-
2001.....	224	245	245	337	429	524	566	495	287	111	15	-
2000.....	178	232	275	338	475	599	510	472	315	105	14	-
1999.....	161	213	299	383	501	589	500	452	292	94	11	-
White, both sexes												
2004.....	432	614	740	767	905	1,092	1,023	748	394	119	15	-
2003.....	423	604	721	783	938	1,071	1,001	763	393	109	17	-
2002.....	435	559	758	878	991	1,108	924	706	380	109	10	-
2001.....	483	549	631	835	959	1,089	984	752	393	139	16	-
2000.....	373	528	655	814	1,060	1,185	918	728	396	125	19	-
1999.....	348	506	657	830	1,064	1,157	916	700	375	127	12	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rectosigmoid junction and rectum (C19-C20)												
White, male												
2004.....	272	420	467	479	580	631	528	292	117	46	3	-
2003.....	268	390	477	485	581	582	525	310	127	30	3	-
2002.....	282	369	514	553	573	627	466	284	117	23	-	-
2001.....	307	342	421	538	586	627	487	304	135	39	4	-
2000.....	229	331	431	526	640	651	454	296	113	27	5	-
1999.....	215	323	408	503	622	632	460	292	103	40	3	-
White, female												
2004.....	160	194	273	288	325	461	495	456	277	73	12	-
2003.....	155	214	244	298	357	489	476	453	266	79	14	-
2002.....	153	190	244	325	418	481	458	422	263	86	10	-
2001.....	176	207	210	297	373	462	497	448	258	100	12	-
2000.....	144	197	224	288	420	534	464	432	283	98	14	-
1999.....	133	183	249	327	442	525	456	408	272	87	9	-
Black, both sexes												
2004.....	79	89	105	101	91	93	84	56	46	7	4	-
2003.....	93	107	101	100	111	107	84	62	30	14	2	-
2002.....	73	90	104	109	104	100	106	51	36	11	2	-
2001.....	75	92	77	95	105	99	99	62	36	14	4	-
2000.....	64	82	96	123	117	99	68	51	33	7	1	-
1999.....	58	80	98	108	109	114	74	65	27	6	2	-
Black, male												
2004.....	39	53	60	62	56	54	43	20	9	3	-	-
2003.....	54	65	62	50	69	56	29	26	14	4	-	-
2002.....	42	53	63	58	56	46	60	19	13	-	-	-
2001.....	39	59	51	62	64	49	38	18	11	4	1	-
2000.....	36	58	55	84	70	48	32	19	5	1	1	-
1999.....	38	56	59	60	60	57	40	29	9	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rectosigmoid junction and rectum (C19-C20)												
Black, female												
2004.....	40	36	45	39	35	39	41	36	37	4	4	-
2003.....	39	42	39	50	42	51	55	36	16	10	2	-
2002.....	31	37	41	51	48	54	46	32	23	11	2	-
2001.....	36	33	26	33	41	50	61	44	25	10	3	-
2000.....	28	24	41	39	47	51	36	32	28	6	-	-
1999.....	20	24	39	48	49	57	34	36	18	5	2	-
American Indian 1/, both sexes												
2004.....	2	4	7	6	-	5	6	2	2	-	-	-
2003.....	2	1	4	5	6	9	2	-	-	-	-	-
2002.....	3	11	8	7	5	3	4	3	1	1	-	-
2001.....	5	4	7	5	8	-	3	1	1	-	-	-
2000.....	3	4	3	4	5	9	3	2	2	1	-	-
1999.....	5	2	4	4	5	2	5	3	-	1	-	-
American Indian 1/, male												
2004.....	1	4	4	5	-	3	4	-	1	-	-	-
2003.....	-	-	3	4	3	5	1	-	-	-	-	-
2002.....	3	10	6	5	2	1	3	1	-	-	-	-
2001.....	5	3	2	3	4	-	2	1	1	-	-	-
2000.....	2	3	1	2	4	3	2	-	-	1	-	-
1999.....	2	2	3	4	5	-	2	-	-	1	-	-
American Indian 1/, female												
2004.....	1	-	3	1	-	2	2	2	1	-	-	-
2003.....	2	1	1	1	3	4	1	-	-	-	-	-
2002.....	-	1	2	2	3	2	1	2	1	1	-	-
2001.....	-	1	5	2	4	-	1	-	-	-	-	-
2000.....	1	1	2	2	1	6	1	2	2	-	-	-
1999.....	3	-	1	-	-	2	3	3	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rectosigmoid junction and rectum (C19-C20)												
Asian or Pacific Islander 2/, both sexes												
2004.....	15	20	26	18	17	21	18	14	9	-	-	-
2003.....	18	23	24	20	28	27	17	12	4	2	2	-
2002.....	16	15	24	21	20	27	23	13	3	5	-	-
2001.....	20	15	15	17	24	19	14	9	9	1	-	-
2000.....	12	25	20	22	21	20	17	13	4	1	-	-
1999.....	16	15	18	24	18	18	13	9	3	5	-	-
Asian or Pacific Islander 2/, male												
2004.....	10	10	17	12	9	9	7	8	6	-	-	-
2003.....	11	10	16	13	11	14	10	8	2	-	2	-
2002.....	8	10	16	13	11	14	11	6	2	2	-	-
2001.....	8	11	11	12	13	7	7	6	5	-	-	-
2000.....	7	15	12	13	14	12	8	7	2	-	-	-
1999.....	11	9	8	16	8	13	6	4	1	3	-	-
Asian or Pacific Islander 2/, female												
2004.....	5	10	9	6	8	12	11	6	3	-	-	-
2003.....	7	13	8	7	17	13	7	4	2	2	-	-
2002.....	8	5	8	8	9	13	12	7	1	3	-	-
2001.....	12	4	4	5	11	12	7	3	4	1	-	-
2000.....	5	10	8	9	7	8	9	6	2	1	-	-
1999.....	5	6	10	8	10	5	7	5	2	2	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Anus and anus canal (C21)

All races, both sexes

2004.....	62	64	59	57	58	66	58	32	17	7	1	-
2003.....	51	66	55	50	57	68	52	42	16	6	1	-
2002.....	62	51	64	51	50	57	52	33	18	3	-	-
2001.....	53	49	42	51	58	64	51	38	18	5	2	-
2000.....	49	44	51	48	58	64	38	28	17	5	-	-
1999.....	38	43	38	44	58	54	46	38	18	3	-	-

All races, male

2004.....	25	35	27	26	22	24	11	9	7	2	1	-
2003.....	15	26	19	25	18	18	8	12	-	-	-	-
2002.....	29	17	30	21	17	17	15	11	6	-	-	-
2001.....	20	21	16	21	25	23	11	8	2	1	-	-
2000.....	23	23	20	22	20	21	17	5	6	1	-	-
1999.....	16	15	14	17	18	16	15	8	4	-	-	-

All races, female

2004.....	37	29	32	31	36	42	47	23	10	5	-	-
2003.....	36	40	36	25	39	50	44	30	16	6	1	-
2002.....	33	34	34	30	33	40	37	22	12	3	-	-
2001.....	33	28	26	30	33	41	40	30	16	4	2	-
2000.....	26	21	31	26	38	43	21	23	11	4	-	-
1999.....	22	28	24	27	40	38	31	30	14	3	-	-

White, both sexes

2004.....	53	59	54	50	49	59	56	28	16	5	1	-
2003.....	41	58	47	43	48	58	48	40	15	6	1	-
2002.....	56	46	55	47	44	54	49	30	18	3	-	-
2001.....	42	43	40	45	56	61	45	36	17	4	2	-
2000.....	41	39	46	45	51	56	37	26	13	5	-	-
1999.....	28	40	36	38	49	52	40	36	18	3	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Anus and anus canal (C21)												
White, male												
2004.....	21	33	24	22	18	23	11	7	6	1	1	-
2003.....	10	23	19	20	17	15	8	10	-	-	-	-
2002.....	24	15	26	19	16	15	14	10	6	-	-	-
2001.....	16	18	15	18	24	22	9	8	2	-	-	-
2000.....	17	20	16	20	18	17	17	5	5	1	-	-
1999.....	11	14	13	16	16	16	14	7	4	-	-	-
White, female												
2004.....	32	26	30	28	31	36	45	21	10	4	-	-
2003.....	31	35	28	23	31	43	40	30	15	6	1	-
2002.....	32	31	29	28	28	39	35	20	12	3	-	-
2001.....	26	25	25	27	32	39	36	28	15	4	2	-
2000.....	24	19	30	25	33	39	20	21	8	4	-	-
1999.....	17	26	23	22	33	36	26	29	14	3	-	-
Black, both sexes												
2004.....	8	4	5	6	4	6	-	3	1	2	-	-
2003.....	9	7	5	6	8	8	3	1	1	-	-	-
2002.....	6	5	8	4	5	3	3	3	-	-	-	-
2001.....	9	5	1	5	2	3	6	2	-	1	-	-
2000.....	8	4	4	3	6	7	1	2	3	-	-	-
1999.....	9	3	2	4	9	2	5	1	-	-	-	-
Black, male												
2004.....	4	1	3	4	2	1	-	1	1	1	-	-
2003.....	5	2	-	5	1	3	-	1	-	-	-	-
2002.....	5	2	3	2	1	2	1	1	-	-	-	-
2001.....	2	3	1	2	1	1	2	-	-	1	-	-
2000.....	6	2	3	2	2	4	-	-	1	-	-	-
1999.....	5	1	1	1	2	-	1	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Anus and anus canal (C21)												
Asian or Pacific Islander 2/, both sexes												
2004.....	1	1	-	1	5	1	2	1	-	-	-	-
2003.....	1	1	2	1	1	2	1	1	-	-	-	-
2002.....	-	-	1	-	1	-	-	-	-	-	-	-
2001.....	1	-	-	1	-	-	-	-	1	-	-	-
2000.....	-	-	-	-	1	1	-	-	1	-	-	-
1999.....	-	-	-	2	-	-	1	1	-	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	-	1	-	-	2	-	-	1	-	-	-	-
2003.....	-	1	-	-	-	-	-	1	-	-	-	-
2002.....	-	-	1	-	-	-	-	-	-	-	-	-
2001.....	1	-	-	1	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	1	-	-	1	3	1	2	-	-	-	-	-
2003.....	1	-	2	1	1	2	1	-	-	-	-	-
2002.....	-	-	-	-	1	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	1	-	-	-
2000.....	-	-	-	-	1	1	-	-	1	-	-	-
1999.....	-	-	-	2	-	-	1	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Of liver and intrahepatic bile ducts (C22)												
All races, both sexes												
2004.....	1,476	1,554	1,569	1,654	2,016	2,136	1,923	1,030	429	104	12	-
2003.....	1,408	1,348	1,417	1,612	2,018	2,208	1,741	1,056	396	113	11	-
2002.....	1,245	1,208	1,350	1,618	1,963	2,049	1,679	1,094	419	80	16	-
2001.....	1,073	1,063	1,288	1,556	1,884	2,003	1,646	1,009	386	97	8	-
2000.....	944	1,040	1,216	1,526	1,913	2,011	1,547	943	359	81	4	-
1999.....	826	928	1,163	1,558	1,933	1,914	1,511	863	325	75	7	-
All races, male												
2004.....	1,216	1,205	1,113	1,092	1,290	1,202	1,025	460	134	29	3	-
2003.....	1,136	999	996	1,054	1,276	1,354	947	495	154	36	2	-
2002.....	1,015	899	959	1,100	1,267	1,212	900	463	155	22	2	-
2001.....	872	780	911	1,036	1,188	1,200	847	481	140	31	2	-
2000.....	729	761	846	1,039	1,207	1,196	856	407	114	26	1	-
1999.....	628	671	794	1,041	1,243	1,103	774	352	109	17	-	-
All races, female												
2004.....	260	349	456	562	726	934	898	570	295	75	9	-
2003.....	272	349	421	558	742	854	794	561	242	77	9	-
2002.....	230	309	391	518	696	837	779	631	264	58	14	-
2001.....	201	283	377	520	696	803	799	528	246	66	6	-
2000.....	215	279	370	487	706	815	691	536	245	55	3	-
1999.....	198	257	369	517	690	811	737	511	216	58	7	-
White, both sexes												
2004.....	1,077	1,121	1,213	1,295	1,623	1,790	1,682	899	370	87	10	-
2003.....	1,001	1,011	1,069	1,270	1,633	1,878	1,512	923	349	89	7	-
2002.....	862	875	1,053	1,298	1,611	1,766	1,471	973	354	67	11	-
2001.....	764	772	1,000	1,239	1,566	1,729	1,415	896	344	82	7	-
2000.....	654	756	924	1,204	1,580	1,717	1,376	830	325	70	2	-
1999.....	601	662	892	1,258	1,614	1,660	1,338	766	291	63	7	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Of liver and intrahepatic bile ducts (C22)

White, male

2004.....	893	855	858	863	1,040	1,039	903	408	109	23	3	-
2003.....	801	752	761	840	1,060	1,162	833	435	132	28	1	-
2002.....	703	651	746	889	1,041	1,049	781	417	131	18	1	-
2001.....	617	561	719	851	1,009	1,037	746	426	129	27	2	-
2000.....	500	542	641	827	1,006	1,048	766	352	105	21	-	-
1999.....	448	481	613	846	1,052	960	698	321	103	14	-	-

White, female

2004.....	184	266	355	432	583	751	779	491	261	64	7	-
2003.....	200	259	308	430	573	716	679	488	217	61	6	-
2002.....	159	224	307	409	570	717	690	556	223	49	10	-
2001.....	147	211	281	388	557	692	669	470	215	55	5	-
2000.....	154	214	283	377	574	669	610	478	220	49	2	-
1999.....	153	181	279	412	562	700	640	445	188	49	7	-

Black, both sexes

2004.....	284	301	203	225	218	199	143	75	39	14	-	-
2003.....	306	232	218	180	211	178	129	91	30	19	3	-
2002.....	258	231	161	184	204	176	122	77	43	6	2	-
2001.....	214	215	160	185	178	155	151	71	32	10	1	-
2000.....	199	171	173	189	202	162	101	71	21	7	2	-
1999.....	143	157	158	191	207	149	117	69	29	10	-	-

Black, male

2004.....	234	242	145	152	139	87	71	27	16	4	-	-
2003.....	253	171	148	109	129	96	61	39	14	7	1	-
2002.....	209	170	115	129	135	97	67	24	12	3	-	-
2001.....	172	162	105	110	97	85	61	31	6	-	-	-
2000.....	152	134	122	131	124	77	54	34	4	3	1	-
1999.....	115	110	102	124	124	85	49	18	6	2	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Of liver and intrahepatic bile ducts (C22)

Black, female

2004.....	50	59	58	73	79	112	72	48	23	10	-	-
2003.....	53	61	70	71	82	82	68	52	16	12	2	-
2002.....	49	61	46	55	69	79	55	53	31	3	2	-
2001.....	42	53	55	75	81	70	90	40	26	10	1	-
2000.....	47	37	51	58	78	85	47	37	17	4	1	-
1999.....	28	47	56	67	83	64	68	51	23	8	-	-

American Indian 1/, both sexes

2004.....	19	17	22	11	10	21	20	7	3	-	-	-
2003.....	19	15	12	10	12	12	4	3	-	-	1	-
2002.....	11	9	11	13	14	6	11	5	2	2	-	-
2001.....	13	9	11	15	21	9	10	6	1	-	-	-
2000.....	13	14	9	11	10	6	11	8	1	1	-	-
1999.....	4	9	9	12	8	12	5	4	1	-	-	-

American Indian 1/, male

2004.....	14	14	9	5	5	6	9	1	2	-	-	-
2003.....	15	7	8	7	7	10	2	1	-	-	-	-
2002.....	10	8	7	7	10	2	6	2	-	-	-	-
2001.....	12	6	7	10	17	5	5	1	-	-	-	-
2000.....	10	10	5	8	8	3	5	2	-	1	-	-
1999.....	3	6	7	5	3	7	2	1	-	-	-	-

American Indian 1/, female

2004.....	5	3	13	6	5	15	11	6	1	-	-	-
2003.....	4	8	4	3	5	2	2	2	-	-	1	-
2002.....	1	1	4	6	4	4	5	3	2	2	-	-
2001.....	1	3	4	5	4	4	5	5	1	-	-	-
2000.....	3	4	4	3	2	3	6	6	1	-	-	-
1999.....	1	3	2	7	5	5	3	3	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Of liver and intrahepatic bile ducts (C22)												
Asian or Pacific Islander 2/, both sexes												
2004.....	96	115	131	123	165	126	78	49	17	3	2	-
2003.....	82	90	118	152	162	140	96	39	17	5	-	-
2002.....	114	93	125	123	134	101	75	39	20	5	3	-
2001.....	82	67	117	117	119	110	70	36	9	5	-	-
2000.....	78	99	110	122	121	126	59	34	12	3	-	-
1999.....	78	100	104	97	104	93	51	24	4	2	-	-
Asian or Pacific Islander 2/, male												
2004.....	75	94	101	72	106	70	42	24	7	2	-	-
2003.....	67	69	79	98	80	86	51	20	8	1	-	-
2002.....	93	70	91	75	81	64	46	20	12	1	1	-
2001.....	71	51	80	65	65	73	35	23	5	4	-	-
2000.....	67	75	78	73	69	68	31	19	5	1	-	-
1999.....	62	74	72	66	64	51	25	12	-	1	-	-
Asian or Pacific Islander 2/, female												
2004.....	21	21	30	51	59	56	36	25	10	1	2	-
2003.....	15	21	39	54	82	54	45	19	9	4	-	-
2002.....	21	23	34	48	53	37	29	19	8	4	2	-
2001.....	11	16	37	52	54	37	35	13	4	1	-	-
2000.....	11	24	32	49	52	58	28	15	7	2	-	-
1999.....	16	26	32	31	40	42	26	12	4	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Liver (C22.0,C22.2-C22.9)												
All races, both sexes												
2004.....	1,289	1,285	1,180	1,263	1,547	1,579	1,397	746	297	76	8	-
2003.....	1,200	1,094	1,078	1,248	1,557	1,681	1,280	758	289	93	7	-
2002.....	1,078	964	1,057	1,289	1,573	1,572	1,253	768	297	58	11	-
2001.....	926	874	1,028	1,224	1,463	1,530	1,223	722	276	60	8	-
2000.....	788	857	967	1,208	1,511	1,536	1,191	680	252	55	4	-
1999.....	702	746	929	1,240	1,552	1,495	1,137	632	238	57	6	-
All races, male												
2004.....	1,109	1,055	893	902	1,049	948	781	351	86	23	2	-
2003.....	1,034	850	811	859	1,019	1,088	737	362	115	30	1	-
2002.....	922	769	803	926	1,064	965	699	348	116	15	2	-
2001.....	786	689	768	860	967	958	649	375	104	19	2	-
2000.....	651	667	719	867	994	956	690	314	86	15	1	-
1999.....	570	565	679	876	1,041	901	616	271	87	14	-	-
All races, female												
2004.....	180	230	287	361	498	631	616	395	211	53	6	-
2003.....	166	244	267	389	538	593	543	396	174	63	6	-
2002.....	156	195	254	363	509	607	554	420	181	43	9	-
2001.....	140	185	260	364	496	572	574	347	172	41	6	-
2000.....	137	190	248	341	517	580	501	366	166	40	3	-
1999.....	132	181	250	364	511	594	521	361	151	43	6	-
White, both sexes												
2004.....	925	895	869	970	1,215	1,298	1,209	636	256	61	6	-
2003.....	835	793	790	972	1,234	1,407	1,090	654	247	73	4	-
2002.....	730	678	806	1,018	1,268	1,346	1,094	676	250	49	7	-
2001.....	644	619	782	954	1,199	1,303	1,046	638	241	51	7	-
2000.....	524	607	717	925	1,220	1,295	1,047	591	223	47	2	-
1999.....	494	515	694	981	1,267	1,282	998	553	209	48	6	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Liver (C22.0,C22.2-C22.9)												
White, male												
2004.....	806	730	664	705	828	809	674	304	68	18	2	-
2003.....	722	623	607	681	838	921	640	315	94	23	1	-
2002.....	626	544	611	745	860	832	604	308	99	13	1	-
2001.....	545	489	603	698	818	823	568	330	95	18	2	-
2000.....	434	470	532	675	814	831	611	263	77	13	-	-
1999.....	397	393	511	702	868	780	556	245	82	12	-	-
White, female												
2004.....	119	165	205	265	387	489	535	332	188	43	4	-
2003.....	113	170	183	291	396	486	450	339	153	50	3	-
2002.....	104	134	195	273	408	514	490	368	151	36	6	-
2001.....	99	130	179	256	381	480	478	308	146	33	5	-
2000.....	90	137	185	250	406	464	436	328	146	34	2	-
1999.....	97	122	183	279	399	502	442	308	127	36	6	-
Black, both sexes												
2004.....	262	279	176	184	183	160	116	63	28	12	-	-
2003.....	277	210	183	141	177	141	107	67	28	16	2	-
2002.....	242	205	138	155	178	141	95	59	33	4	1	-
2001.....	198	190	135	155	152	131	116	54	26	7	1	-
2000.....	183	153	143	164	173	134	86	53	19	5	2	-
1999.....	134	137	140	162	182	127	93	57	25	7	-	-
Black, male												
2004.....	223	232	129	130	124	72	65	24	10	3	-	-
2003.....	237	157	132	88	109	83	49	29	14	6	-	-
2002.....	202	160	103	110	120	79	54	23	8	2	-	-
2001.....	164	148	88	94	83	69	51	25	5	-	-	-
2000.....	146	123	111	118	109	65	48	31	4	1	1	-
1999.....	112	103	96	110	110	73	40	15	5	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Liver (C22.0,C22.2-C22.9)												
Black, female												
2004.....	39	47	47	54	59	88	51	39	18	9	-	-
2003.....	40	53	51	53	68	58	58	38	14	10	2	-
2002.....	40	45	35	45	58	62	41	36	25	2	1	-
2001.....	34	42	47	61	69	62	65	29	21	7	1	-
2000.....	37	30	32	46	64	69	38	22	15	4	1	-
1999.....	22	34	44	52	72	54	53	42	20	6	-	-
American Indian 1/, both sexes												
2004.....	17	15	20	9	7	15	12	5	3	-	-	-
2003.....	16	13	7	7	7	11	3	2	-	-	1	-
2002.....	9	5	8	9	10	5	8	2	-	2	-	-
2001.....	13	9	9	13	15	5	7	2	1	-	-	-
2000.....	12	11	8	10	9	5	9	8	-	1	-	-
1999.....	4	7	8	11	4	10	4	4	1	-	-	-
American Indian 1/, male												
2004.....	12	13	8	5	4	6	6	1	2	-	-	-
2003.....	14	6	5	5	4	9	2	1	-	-	-	-
2002.....	8	5	6	5	7	2	6	1	-	-	-	-
2001.....	12	6	5	9	14	3	3	-	-	-	-	-
2000.....	10	7	5	8	7	2	4	2	-	1	-	-
1999.....	3	4	7	5	1	6	1	1	-	-	-	-
American Indian 1/, female												
2004.....	5	2	12	4	3	9	6	4	1	-	-	-
2003.....	2	7	2	2	3	2	1	1	-	-	1	-
2002.....	1	-	2	4	3	3	2	1	-	2	-	-
2001.....	1	3	4	4	1	2	4	2	1	-	-	-
2000.....	2	4	3	2	2	3	5	6	-	-	-	-
1999.....	1	3	1	6	3	4	3	3	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Liver (C22.0,C22.2-C22.9)												
Asian or Pacific Islander 2/, both sexes												
2004.....	85	96	115	100	142	106	60	42	10	3	2	-
2003.....	72	78	98	128	139	122	80	35	14	4	-	-
2002.....	97	76	105	107	117	80	56	31	14	3	3	-
2001.....	71	56	102	102	97	91	54	28	8	2	-	-
2000.....	69	86	99	109	109	102	49	28	10	2	-	-
1999.....	70	87	87	86	99	76	42	18	3	2	-	-
Asian or Pacific Islander 2/, male												
2004.....	68	80	92	62	93	61	36	22	6	2	-	-
2003.....	61	64	67	85	68	75	46	17	7	1	-	-
2002.....	86	60	83	66	77	52	35	16	9	-	1	-
2001.....	65	46	72	59	52	63	27	20	4	1	-	-
2000.....	61	67	71	66	64	58	27	18	5	-	-	-
1999.....	58	65	65	59	62	42	19	10	-	1	-	-
Asian or Pacific Islander 2/, female												
2004.....	17	16	23	38	49	45	24	20	4	1	2	-
2003.....	11	14	31	43	71	47	34	18	7	3	-	-
2002.....	11	16	22	41	40	28	21	15	5	3	2	-
2001.....	6	10	30	43	45	28	27	8	4	1	-	-
2000.....	8	19	28	43	45	44	22	10	5	2	-	-
1999.....	12	22	22	27	37	34	23	8	3	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intrahepatic bile duct carcinoma (C22.1)												
All races, both sexes												
2004.....	187	269	389	391	469	557	526	284	132	28	4	-
2003.....	208	254	339	364	461	527	461	298	107	20	4	-
2002.....	167	244	293	329	390	477	426	326	122	22	5	-
2001.....	147	189	260	332	421	473	423	287	110	37	-	-
2000.....	156	183	249	318	402	475	356	263	107	26	-	-
1999.....	124	182	234	318	381	419	374	231	87	18	1	-
All races, male												
2004.....	107	150	220	190	241	254	244	109	48	6	1	-
2003.....	102	149	185	195	257	266	210	133	39	6	1	-
2002.....	93	130	156	174	203	247	201	115	39	7	-	-
2001.....	86	91	143	176	221	242	198	106	36	12	-	-
2000.....	78	94	127	172	213	240	166	93	28	11	-	-
1999.....	58	106	115	165	202	202	158	81	22	3	-	-
All races, female												
2004.....	80	119	169	201	228	303	282	175	84	22	3	-
2003.....	106	105	154	169	204	261	251	165	68	14	3	-
2002.....	74	114	137	155	187	230	225	211	83	15	5	-
2001.....	61	98	117	156	200	231	225	181	74	25	-	-
2000.....	78	89	122	146	189	235	190	170	79	15	-	-
1999.....	66	76	119	153	179	217	216	150	65	15	1	-
White, both sexes												
2004.....	152	226	344	325	408	492	473	263	114	26	4	-
2003.....	166	218	279	298	399	471	422	269	102	16	3	-
2002.....	132	197	247	280	343	420	377	297	104	18	4	-
2001.....	120	153	218	285	367	426	369	258	103	31	-	-
2000.....	130	149	207	279	360	422	329	239	102	23	-	-
1999.....	107	147	198	277	347	378	340	213	82	15	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intrahepatic bile duct carcinoma (C22.1)												
White, male												
2004.....	87	125	194	158	212	230	229	104	41	5	1	-
2003.....	79	129	154	159	222	241	193	120	38	5	-	-
2002.....	77	107	135	144	181	217	177	109	32	5	-	-
2001.....	72	72	116	153	191	214	178	96	34	9	-	-
2000.....	66	72	109	152	192	217	155	89	28	8	-	-
1999.....	51	88	102	144	184	180	142	76	21	2	-	-
White, female												
2004.....	65	101	150	167	196	262	244	159	73	21	3	-
2003.....	87	89	125	139	177	230	229	149	64	11	3	-
2002.....	55	90	112	136	162	203	200	188	72	13	4	-
2001.....	48	81	102	132	176	212	191	162	69	22	-	-
2000.....	64	77	98	127	168	205	174	150	74	15	-	-
1999.....	56	59	96	133	163	198	198	137	61	13	1	-
Black, both sexes												
2004.....	22	22	27	41	35	39	27	12	11	2	-	-
2003.....	29	22	35	39	34	37	22	24	2	3	1	-
2002.....	16	26	23	29	26	35	27	18	10	2	1	-
2001.....	16	25	25	30	26	24	35	17	6	3	-	-
2000.....	16	18	30	25	29	28	15	18	2	2	-	-
1999.....	9	20	18	29	25	22	24	12	4	3	-	-
Black, male												
2004.....	11	10	16	22	15	15	6	3	6	1	-	-
2003.....	16	14	16	21	20	13	12	10	-	1	1	-
2002.....	7	10	12	19	15	18	13	1	4	1	-	-
2001.....	8	14	17	16	14	16	10	6	1	-	-	-
2000.....	6	11	11	13	15	12	6	3	-	2	-	-
1999.....	3	7	6	14	14	12	9	3	1	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Intrahepatic bile duct carcinoma (C22.1)

Black, female

2004.....	11	12	11	19	20	24	21	9	5	1	-	-
2003.....	13	8	19	18	14	24	10	14	2	2	-	-
2002.....	9	16	11	10	11	17	14	17	6	1	1	-
2001.....	8	11	8	14	12	8	25	11	5	3	-	-
2000.....	10	7	19	12	14	16	9	15	2	-	-	-
1999.....	6	13	12	15	11	10	15	9	3	2	-	-

American Indian 1/, both sexes

2004.....	2	2	2	2	3	6	8	2	-	-	-	-
2003.....	3	2	5	3	5	1	1	1	-	-	-	-
2002.....	2	4	3	4	4	1	3	3	2	-	-	-
2001.....	-	-	2	2	6	4	3	4	-	-	-	-
2000.....	1	3	1	1	1	1	2	-	1	-	-	-
1999.....	-	2	1	1	4	2	1	-	-	-	-	-

American Indian 1/, male

2004.....	2	1	1	-	1	-	3	-	-	-	-	-
2003.....	1	1	3	2	3	1	-	-	-	-	-	-
2002.....	2	3	1	2	3	-	-	1	-	-	-	-
2001.....	-	-	2	1	3	2	2	1	-	-	-	-
2000.....	-	3	-	-	1	1	1	-	-	-	-	-
1999.....	-	2	-	-	2	1	1	-	-	-	-	-

American Indian 1/, female

2004.....	-	1	1	2	2	6	5	2	-	-	-	-
2003.....	2	1	2	1	2	-	1	1	-	-	-	-
2002.....	-	1	2	2	1	1	3	2	2	-	-	-
2001.....	-	-	-	1	3	2	1	3	-	-	-	-
2000.....	1	-	1	1	-	-	1	-	1	-	-	-
1999.....	-	-	1	1	2	1	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intrahepatic bile duct carcinoma (C22.1)												
Asian or Pacific Islander 2/, both sexes												
2004.....	11	19	16	23	23	20	18	7	7	-	-	-
2003.....	10	12	20	24	23	18	16	4	3	1	-	-
2002.....	17	17	20	16	17	21	19	8	6	2	-	-
2001.....	11	11	15	15	22	19	16	8	1	3	-	-
2000.....	9	13	11	13	12	24	10	6	2	1	-	-
1999.....	8	13	17	11	5	17	9	6	1	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	7	14	9	10	13	9	6	2	1	-	-	-
2003.....	6	5	12	13	12	11	5	3	1	-	-	-
2002.....	7	10	8	9	4	12	11	4	3	1	-	-
2001.....	6	5	8	6	13	10	8	3	1	3	-	-
2000.....	6	8	7	7	5	10	4	1	-	1	-	-
1999.....	4	9	7	7	2	9	6	2	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	4	5	7	13	10	11	12	5	6	-	-	-
2003.....	4	7	8	11	11	7	11	1	2	1	-	-
2002.....	10	7	12	7	13	9	8	4	3	1	-	-
2001.....	5	6	7	9	9	9	8	5	-	-	-	-
2000.....	3	5	4	6	7	14	6	5	2	-	-	-
1999.....	4	4	10	4	3	8	3	4	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Of gallbladder and extrahepatic bile ducts (C23-C24)

All races, both sexes

2004.....	126	222	291	361	502	592	542	388	196	57	6	-
2003.....	145	192	288	319	458	603	588	435	182	50	7	-
2002.....	153	211	232	357	492	613	524	422	182	52	10	-
2001.....	149	197	257	356	525	618	625	418	212	57	9	-
2000.....	151	197	280	401	568	675	564	442	179	51	5	-
1999.....	136	213	268	400	535	645	579	442	170	54	8	-

All races, male

2004.....	49	97	101	148	197	252	199	130	49	15	-	-
2003.....	54	79	117	155	196	235	201	151	42	9	2	-
2002.....	56	84	97	146	212	251	183	139	59	10	-	-
2001.....	63	84	92	125	207	247	247	135	56	14	-	-
2000.....	50	72	115	172	242	262	201	143	42	8	1	-
1999.....	52	75	108	165	212	250	198	127	43	9	2	-

All races, female

2004.....	77	125	190	213	305	340	343	258	147	42	6	-
2003.....	91	113	171	164	262	368	387	284	140	41	5	-
2002.....	97	127	135	211	280	362	341	283	123	42	10	-
2001.....	86	113	165	231	318	371	378	283	156	43	9	-
2000.....	101	125	165	229	326	413	363	299	137	43	4	-
1999.....	84	138	160	235	323	395	381	315	127	45	6	-

White, both sexes

2004.....	100	181	240	291	427	517	481	354	175	54	6	-
2003.....	114	160	252	278	395	533	519	398	167	42	7	-
2002.....	120	169	200	301	428	536	468	384	165	45	10	-
2001.....	123	167	213	288	458	549	570	377	193	51	8	-
2000.....	123	171	229	347	496	614	518	410	167	44	5	-
1999.....	114	181	233	348	470	576	512	404	158	48	6	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Of gallbladder and extrahepatic bile ducts (C23-C24)

White, male

2004.....	42	76	88	117	172	223	177	122	43	15	-	-
2003.....	43	63	103	141	170	208	180	138	40	9	2	-
2002.....	44	66	84	126	185	223	164	128	55	9	-	-
2001.....	52	73	78	107	183	219	224	122	46	14	-	-
2000.....	44	64	96	157	210	238	188	134	39	6	1	-
1999.....	41	68	94	139	187	230	178	119	41	9	2	-

White, female

2004.....	58	105	152	174	255	294	304	232	132	39	6	-
2003.....	71	97	149	137	225	325	339	260	127	33	5	-
2002.....	76	103	116	175	243	313	304	256	110	36	10	-
2001.....	71	94	135	181	275	330	346	255	147	37	8	-
2000.....	79	107	133	190	286	376	330	276	128	38	4	-
1999.....	73	113	139	209	283	346	334	285	117	39	4	-

Black, both sexes

2004.....	14	28	37	48	47	48	39	23	14	3	-	-
2003.....	21	20	21	25	40	45	44	25	10	7	-	-
2002.....	24	33	29	44	44	58	36	29	13	6	-	-
2001.....	19	20	33	48	48	44	39	35	11	5	-	-
2000.....	25	14	33	46	50	43	29	21	9	7	-	-
1999.....	17	24	28	38	48	44	46	21	10	5	2	-

Black, male

2004.....	4	15	6	16	16	21	12	3	1	-	-	-
2003.....	8	10	6	8	15	16	11	6	1	-	-	-
2002.....	9	12	10	14	17	19	11	6	2	-	-	-
2001.....	5	10	7	14	19	17	14	9	4	-	-	-
2000.....	3	3	11	10	23	16	7	5	2	2	-	-
1999.....	7	3	12	17	19	15	14	2	2	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Of gallbladder and extrahepatic bile ducts (C23-C24)												
Black, female												
2004.....	10	13	31	32	31	27	27	20	13	3	-	-
2003.....	13	10	15	17	25	29	33	19	9	7	-	-
2002.....	15	21	19	30	27	39	25	23	11	6	-	-
2001.....	14	10	26	34	29	27	25	26	7	5	-	-
2000.....	22	11	22	36	27	27	22	16	7	5	-	-
1999.....	10	21	16	21	29	29	32	19	8	5	2	-
American Indian 1/, both sexes												
2004.....	1	2	5	3	7	3	4	4	1	-	-	-
2003.....	2	2	2	5	6	2	4	2	3	-	-	-
2002.....	2	2	-	4	2	1	3	2	-	-	-	-
2001.....	1	3	2	3	3	6	2	2	3	-	-	-
2000.....	-	2	5	1	5	5	3	1	1	-	-	-
1999.....	2	2	2	4	1	8	3	3	2	-	-	-
American Indian 1/, male												
2004.....	1	1	3	2	2	1	1	1	-	-	-	-
2003.....	-	1	-	2	3	2	1	1	1	-	-	-
2002.....	-	2	-	2	1	-	1	-	-	-	-	-
2001.....	1	-	1	-	1	3	1	-	2	-	-	-
2000.....	-	-	2	-	-	1	-	1	-	-	-	-
1999.....	2	-	-	2	-	1	-	1	-	-	-	-
American Indian 1/, female												
2004.....	-	1	2	1	5	2	3	3	1	-	-	-
2003.....	2	1	2	3	3	-	3	1	2	-	-	-
2002.....	2	-	-	2	1	1	2	2	-	-	-	-
2001.....	-	3	1	3	2	3	1	2	1	-	-	-
2000.....	-	2	3	1	5	4	3	-	1	-	-	-
1999.....	-	2	2	2	1	7	3	2	2	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Of gallbladder and extrahepatic bile ducts (C23-C24)												
Asian or Pacific Islander 2/, both sexes												
2004.....	11	11	9	19	21	24	18	7	6	-	-	-
2003.....	8	10	13	11	17	23	21	10	2	1	-	-
2002.....	7	7	3	8	18	18	17	7	4	1	-	-
2001.....	6	7	9	17	16	19	14	4	5	1	1	-
2000.....	3	10	13	7	17	13	14	10	2	-	-	-
1999.....	3	6	5	10	16	17	18	14	-	1	-	-
Asian or Pacific Islander 2/, male												
2004.....	2	5	4	13	7	7	9	4	5	-	-	-
2003.....	3	5	8	4	8	9	9	6	-	-	-	-
2002.....	3	4	3	4	9	9	7	5	2	1	-	-
2001.....	5	1	6	4	4	8	8	4	4	-	-	-
2000.....	3	5	6	5	9	7	6	3	1	-	-	-
1999.....	2	4	2	7	6	4	6	5	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	9	6	5	6	14	17	9	3	1	-	-	-
2003.....	5	5	5	7	9	14	12	4	2	1	-	-
2002.....	4	3	-	4	9	9	10	2	2	-	-	-
2001.....	1	6	3	13	12	11	6	-	1	1	1	-
2000.....	-	5	7	2	8	6	8	7	1	-	-	-
1999.....	1	2	3	3	10	13	12	9	-	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Of pancreas (C25)												
All races, both sexes												
2004.....	1,715	2,509	3,247	3,717	4,461	5,106	4,747	3,056	1,323	317	53	1
2003.....	1,649	2,327	2,993	3,550	4,554	5,038	4,670	2,897	1,264	293	32	-
2002.....	1,544	2,321	2,784	3,564	4,623	5,210	4,345	2,824	1,217	263	36	1
2001.....	1,577	2,111	2,757	3,525	4,633	4,987	4,335	2,861	1,214	293	42	1
2000.....	1,528	2,030	2,678	3,546	4,557	5,223	4,236	2,571	1,196	286	34	1
1999.....	1,466	2,039	2,668	3,632	4,661	4,931	4,002	2,673	1,120	309	40	1
All races, male												
2004.....	1,051	1,490	1,887	2,093	2,337	2,407	2,009	1,090	371	66	11	-
2003.....	992	1,391	1,715	1,924	2,327	2,358	2,044	992	327	55	8	-
2002.....	971	1,394	1,618	1,924	2,346	2,424	1,846	985	351	60	7	1
2001.....	964	1,269	1,605	1,921	2,294	2,360	1,831	984	310	64	7	1
2000.....	955	1,206	1,543	1,965	2,224	2,447	1,742	891	310	59	2	1
1999.....	907	1,204	1,517	1,976	2,371	2,281	1,659	910	311	59	3	-
All races, female												
2004.....	664	1,019	1,360	1,624	2,124	2,699	2,738	1,966	952	251	42	1
2003.....	657	936	1,278	1,626	2,227	2,680	2,626	1,905	937	238	24	-
2002.....	573	927	1,166	1,640	2,277	2,786	2,499	1,839	866	203	29	-
2001.....	613	842	1,152	1,604	2,339	2,627	2,504	1,877	904	229	35	-
2000.....	573	824	1,135	1,581	2,333	2,776	2,494	1,680	886	227	32	-
1999.....	559	835	1,151	1,656	2,290	2,650	2,343	1,763	809	250	37	1
White, both sexes												
2004.....	1,356	2,042	2,706	3,149	3,796	4,488	4,250	2,748	1,183	280	42	1
2003.....	1,287	1,924	2,475	3,013	3,923	4,449	4,152	2,586	1,150	259	25	-
2002.....	1,223	1,927	2,321	2,975	3,969	4,574	3,904	2,552	1,104	229	30	-
2001.....	1,293	1,753	2,354	2,989	3,998	4,398	3,876	2,581	1,072	261	33	1
2000.....	1,248	1,675	2,250	2,999	3,945	4,603	3,786	2,297	1,059	250	27	1
1999.....	1,184	1,660	2,237	3,087	4,053	4,339	3,574	2,397	1,003	282	30	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Of pancreas (C25)												
White, male												
2004.....	844	1,219	1,601	1,788	2,005	2,156	1,833	1,003	333	57	8	-
2003.....	784	1,168	1,443	1,678	2,033	2,115	1,844	898	301	47	8	-
2002.....	785	1,166	1,378	1,645	2,048	2,177	1,679	900	320	52	5	-
2001.....	801	1,081	1,385	1,647	2,008	2,118	1,657	907	281	60	5	1
2000.....	787	1,003	1,314	1,709	1,949	2,197	1,570	812	272	49	2	1
1999.....	728	978	1,287	1,699	2,096	2,031	1,496	820	280	56	3	-
White, female												
2004.....	512	823	1,105	1,361	1,791	2,332	2,417	1,745	850	223	34	1
2003.....	503	756	1,032	1,335	1,890	2,334	2,308	1,688	849	212	17	-
2002.....	438	761	943	1,330	1,921	2,397	2,225	1,652	784	177	25	-
2001.....	492	672	969	1,342	1,990	2,280	2,219	1,674	791	201	28	-
2000.....	461	672	936	1,290	1,996	2,406	2,216	1,485	787	201	25	-
1999.....	456	682	950	1,388	1,957	2,308	2,078	1,577	723	226	27	1
Black, both sexes												
2004.....	297	380	444	465	539	498	377	251	113	30	9	-
2003.....	313	327	427	449	514	491	423	252	100	29	6	-
2002.....	259	334	396	475	550	516	351	226	95	32	6	1
2001.....	244	294	353	449	526	489	387	233	124	29	9	-
2000.....	246	307	365	455	523	528	369	228	118	32	5	-
1999.....	249	328	345	461	518	490	362	236	101	25	10	-
Black, male												
2004.....	164	227	240	250	272	195	127	62	29	7	3	-
2003.....	184	187	222	203	246	195	152	73	22	5	-	-
2002.....	159	195	206	227	252	197	138	62	22	8	2	1
2001.....	140	155	188	236	234	205	136	60	22	4	2	-
2000.....	151	178	195	214	236	215	138	60	27	8	-	-
1999.....	158	201	186	231	231	212	135	73	22	3	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Of pancreas (C25)												
Black, female												
2004.....	133	153	204	215	267	303	250	189	84	23	6	-
2003.....	129	140	205	246	268	296	271	179	78	24	6	-
2002.....	100	139	190	248	298	319	213	164	73	24	4	-
2001.....	104	139	165	213	292	284	251	173	102	25	7	-
2000.....	95	129	170	241	287	313	231	168	91	24	5	-
1999.....	91	127	159	230	287	278	227	163	79	22	10	-
American Indian 1/, both sexes												
2004.....	17	17	24	20	22	9	13	3	2	2	-	-
2003.....	5	11	18	14	16	9	10	2	3	1	-	-
2002.....	14	12	10	11	23	12	13	5	1	-	-	-
2001.....	6	11	4	18	14	12	6	7	1	1	-	-
2000.....	8	16	9	10	14	8	10	3	2	1	1	-
1999.....	6	7	12	14	7	10	12	8	1	1	-	-
American Indian 1/, male												
2004.....	12	10	9	10	9	4	6	3	-	1	-	-
2003.....	3	4	13	8	7	3	5	1	-	1	-	-
2002.....	6	6	2	7	9	6	2	3	-	-	-	-
2001.....	3	5	3	10	7	3	3	2	-	-	-	-
2000.....	4	10	6	5	7	2	4	-	1	1	-	-
1999.....	3	3	6	8	6	4	5	2	-	-	-	-
American Indian 1/, female												
2004.....	5	7	15	10	13	5	7	-	2	1	-	-
2003.....	2	7	5	6	9	6	5	1	3	-	-	-
2002.....	8	6	8	4	14	6	11	2	1	-	-	-
2001.....	3	6	1	8	7	9	3	5	1	1	-	-
2000.....	4	6	3	5	7	6	6	3	1	-	1	-
1999.....	3	4	6	6	1	6	7	6	1	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Of pancreas (C25)												
Asian or Pacific Islander 2/, both sexes												
2004.....	45	70	73	83	104	111	107	54	25	5	2	-
2003.....	44	65	73	74	101	89	85	57	11	4	1	-
2002.....	48	48	57	103	81	108	77	41	17	2	-	-
2001.....	34	53	46	69	95	88	66	40	17	2	-	-
2000.....	26	32	54	82	75	84	71	43	17	3	1	-
1999.....	27	44	74	70	83	92	54	32	15	1	-	-
Asian or Pacific Islander 2/, male												
2004.....	31	34	37	45	51	52	43	22	9	1	-	-
2003.....	21	32	37	35	41	45	43	20	4	2	-	-
2002.....	21	27	32	45	37	44	27	20	9	-	-	-
2001.....	20	28	29	28	45	34	35	15	7	-	-	-
2000.....	13	15	28	37	32	33	30	19	10	1	-	-
1999.....	18	22	38	38	38	34	23	15	9	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	14	36	36	38	53	59	64	32	16	4	2	-
2003.....	23	33	36	39	60	44	42	37	7	2	1	-
2002.....	27	21	25	58	44	64	50	21	8	2	-	-
2001.....	14	25	17	41	50	54	31	25	10	2	-	-
2000.....	13	17	26	45	43	51	41	24	7	2	1	-
1999.....	9	22	36	32	45	58	31	17	6	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Of other and ill-defined digestive organs (C26)

All races, both sexes

2004.....	39	66	88	101	123	172	170	138	79	28	5	-
2003.....	46	84	82	144	159	219	215	171	112	36	10	-
2002.....	45	71	85	129	173	250	218	177	106	28	8	-
2001.....	51	75	105	125	194	236	220	178	125	32	7	-
2000.....	50	74	108	122	190	207	223	182	107	33	10	-
1999.....	56	75	104	155	204	240	258	208	129	38	2	-

All races, male

2004.....	22	33	54	58	64	89	74	44	25	4	2	-
2003.....	30	52	52	90	83	117	92	64	31	6	5	-
2002.....	30	48	42	88	101	133	106	54	30	3	2	-
2001.....	35	50	67	63	106	124	86	73	37	6	-	-
2000.....	28	45	67	71	120	105	96	63	20	8	1	-
1999.....	39	46	64	84	101	125	124	75	37	12	-	-

All races, female

2004.....	17	33	34	43	59	83	96	94	54	24	3	-
2003.....	16	32	30	54	76	102	123	107	81	30	5	-
2002.....	15	23	43	41	72	117	112	123	76	25	6	-
2001.....	16	25	38	62	88	112	134	105	88	26	7	-
2000.....	22	29	41	51	70	102	127	119	87	25	9	-
1999.....	17	29	40	71	103	115	134	133	92	26	2	-

White, both sexes

2004.....	29	51	72	81	106	152	158	127	73	27	5	-
2003.....	35	67	67	126	140	189	191	157	103	31	8	-
2002.....	36	59	73	108	142	229	193	163	98	22	8	-
2001.....	38	60	80	104	168	210	190	159	114	29	6	-
2000.....	36	65	90	99	155	185	193	167	97	32	6	-
1999.....	37	57	91	136	181	218	244	189	114	34	2	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Of other and ill-defined digestive organs (C26)												
White, male												
2004.....	16	27	44	50	57	76	68	43	23	4	2	-
2003.....	25	43	43	81	72	101	83	56	26	6	4	-
2002.....	26	42	34	77	83	120	96	49	26	1	2	-
2001.....	28	42	51	53	92	112	78	66	32	6	-	-
2000.....	24	41	57	57	98	98	86	57	19	8	1	-
1999.....	26	36	56	71	91	112	117	70	31	10	-	-
White, female												
2004.....	13	24	28	31	49	76	90	84	50	23	3	-
2003.....	10	24	24	45	68	88	108	101	77	25	4	-
2002.....	10	17	39	31	59	109	97	114	72	21	6	-
2001.....	10	18	29	51	76	98	112	93	82	23	6	-
2000.....	12	24	33	42	57	87	107	110	78	24	5	-
1999.....	11	21	35	65	90	106	127	119	83	24	2	-
Black, both sexes												
2004.....	8	12	13	17	14	17	7	8	6	1	-	-
2003.....	7	16	14	16	12	25	21	12	7	5	1	-
2002.....	9	9	12	19	27	17	20	12	7	5	-	-
2001.....	12	13	20	15	23	19	25	18	10	3	1	-
2000.....	11	7	16	19	24	18	28	13	9	1	3	-
1999.....	17	14	13	15	20	19	10	15	14	3	-	-
Black, male												
2004.....	6	6	8	6	6	11	4	-	2	-	-	-
2003.....	4	9	9	8	8	14	8	7	3	-	-	-
2002.....	4	4	8	11	16	11	9	4	4	2	-	-
2001.....	7	7	12	9	13	8	7	7	5	-	-	-
2000.....	3	3	9	11	16	5	9	5	1	-	-	-
1999.....	12	8	8	9	9	10	4	3	6	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Of other and ill-defined digestive organs (C26)												
Black, female												
2004.....	2	6	5	11	8	6	3	8	4	1	-	-
2003.....	3	7	5	8	4	11	13	5	4	5	1	-
2002.....	5	5	4	8	11	6	11	8	3	3	-	-
2001.....	5	6	8	6	10	11	18	11	5	3	1	-
2000.....	8	4	7	8	8	13	19	8	8	1	3	-
1999.....	5	6	5	6	11	9	6	12	8	2	-	-
American Indian 1/, both sexes												
2004.....	-	-	1	1	1	1	1	-	-	-	-	-
2003.....	-	-	-	1	3	-	-	-	-	-	-	-
2002.....	-	1	-	-	-	-	3	-	1	-	-	-
2001.....	-	1	2	-	-	1	-	-	-	-	-	-
2000.....	1	-	-	-	1	1	-	-	-	-	-	-
1999.....	2	-	-	-	2	-	2	1	-	-	-	-
American Indian 1/, male												
2004.....	-	-	-	-	-	1	-	-	-	-	-	-
2003.....	-	-	-	1	1	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	1	-	-	-	-	-
2001.....	-	-	2	-	-	1	-	-	-	-	-	-
2000.....	1	-	-	-	-	1	-	-	-	-	-	-
1999.....	1	-	-	-	1	-	2	1	-	-	-	-
American Indian 1/, female												
2004.....	-	-	1	1	1	-	1	-	-	-	-	-
2003.....	-	-	-	-	2	-	-	-	-	-	-	-
2002.....	-	1	-	-	-	-	2	-	1	-	-	-
2001.....	-	1	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	1	-	-	-	-	-	-	-
1999.....	1	-	-	-	1	-	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Of other and ill-defined digestive organs (C26)

Asian or Pacific Islander 2/, both sexes

2004.....	2	3	2	2	2	2	4	3	-	-	-	-
2003.....	4	1	1	1	4	5	3	2	2	-	1	-
2002.....	-	2	-	2	4	4	2	2	-	1	-	-
2001.....	1	1	3	6	3	6	5	1	1	-	-	-
2000.....	2	2	2	4	10	3	2	2	1	-	1	-
1999.....	-	4	-	4	1	3	2	3	1	1	-	-

Asian or Pacific Islander 2/, male

2004.....	-	-	2	2	1	1	2	1	-	-	-	-
2003.....	1	-	-	-	2	2	1	1	2	-	1	-
2002.....	-	2	-	-	2	2	-	1	-	-	-	-
2001.....	-	1	2	1	1	3	1	-	-	-	-	-
2000.....	-	1	1	3	6	1	1	1	-	-	-	-
1999.....	-	2	-	4	-	3	1	1	-	1	-	-

Asian or Pacific Islander 2/, female

2004.....	2	3	-	-	1	1	2	2	-	-	-	-
2003.....	3	1	1	1	2	3	2	1	-	-	-	-
2002.....	-	-	-	2	2	2	2	1	-	1	-	-
2001.....	1	-	1	5	2	3	4	1	1	-	-	-
2000.....	2	1	1	1	4	2	1	1	1	-	1	-
1999.....	-	2	-	-	1	-	1	2	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant neoplasms of respiratory and intrathoracic organs (C30-C39)												
All races, both sexes												
2004.....	8,228	13,350	18,832	23,172	27,106	28,243	21,304	10,607	3,529	669	79	3
2003.....	8,081	13,401	18,685	23,076	27,652	28,662	21,095	10,364	3,427	642	79	3
2002.....	8,144	13,412	18,327	22,964	28,596	28,679	20,593	10,023	3,323	587	81	1
2001.....	8,132	13,104	17,756	23,488	29,125	28,406	19,452	9,902	3,178	607	60	4
2000.....	8,026	12,912	17,838	24,115	29,566	28,016	19,043	9,549	3,008	559	61	5
1999.....	7,847	12,656	17,747	23,922	29,383	27,210	18,170	9,210	2,873	570	83	2
All races, male												
2004.....	5,013	8,063	11,232	13,734	15,692	16,227	11,559	5,439	1,607	255	23	3
2003.....	4,922	8,127	11,104	13,740	16,221	16,406	11,503	5,292	1,542	267	27	2
2002.....	4,982	8,155	11,105	13,665	16,838	16,356	11,331	5,177	1,513	218	25	1
2001.....	4,975	7,933	10,790	14,279	17,468	16,465	10,768	5,225	1,440	249	24	4
2000.....	4,840	7,912	10,850	14,694	17,759	16,245	10,611	5,022	1,396	213	16	5
1999.....	4,776	7,857	10,990	14,697	17,647	15,929	10,260	4,925	1,370	244	31	2
All races, female												
2004.....	3,215	5,287	7,600	9,438	11,414	12,016	9,745	5,168	1,922	414	56	-
2003.....	3,159	5,274	7,581	9,336	11,431	12,256	9,592	5,072	1,885	375	52	1
2002.....	3,162	5,257	7,222	9,299	11,758	12,323	9,262	4,846	1,810	369	56	-
2001.....	3,157	5,171	6,966	9,209	11,657	11,941	8,684	4,677	1,738	358	36	-
2000.....	3,186	5,000	6,988	9,421	11,807	11,771	8,432	4,527	1,612	346	45	-
1999.....	3,071	4,799	6,757	9,225	11,736	11,281	7,910	4,285	1,503	326	52	-
White, both sexes												
2004.....	6,534	11,045	16,183	20,092	24,071	25,395	19,314	9,737	3,196	606	72	2
2003.....	6,451	11,247	16,085	20,090	24,567	25,792	19,173	9,493	3,117	570	69	3
2002.....	6,535	11,249	15,637	19,975	25,569	25,868	18,844	9,197	3,041	529	64	-
2001.....	6,598	11,056	15,293	20,436	25,961	25,604	17,694	9,060	2,904	532	48	2
2000.....	6,510	10,899	15,352	21,090	26,285	25,277	17,514	8,731	2,752	495	50	3
1999.....	6,371	10,702	15,204	20,845	26,222	24,525	16,612	8,402	2,613	518	69	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant neoplasms of respiratory and intrathoracic organs (C30-C39)												
White, male												
2004.....	3,902	6,581	9,479	11,845	13,828	14,526	10,421	4,990	1,439	227	21	2
2003.....	3,875	6,703	9,420	11,847	14,341	14,657	10,388	4,832	1,393	233	26	2
2002.....	3,936	6,738	9,307	11,790	14,985	14,635	10,288	4,732	1,377	201	23	-
2001.....	3,948	6,570	9,185	12,304	15,459	14,705	9,711	4,739	1,312	226	21	2
2000.....	3,824	6,539	9,248	12,726	15,684	14,505	9,692	4,563	1,264	185	11	3
1999.....	3,788	6,551	9,294	12,705	15,631	14,265	9,305	4,469	1,241	224	23	1
White, female												
2004.....	2,632	4,464	6,704	8,247	10,243	10,869	8,893	4,747	1,757	379	51	-
2003.....	2,576	4,544	6,665	8,243	10,226	11,135	8,785	4,661	1,724	337	43	1
2002.....	2,599	4,511	6,330	8,185	10,584	11,233	8,556	4,465	1,664	328	41	-
2001.....	2,650	4,486	6,108	8,132	10,502	10,899	7,983	4,321	1,592	306	27	-
2000.....	2,686	4,360	6,104	8,364	10,601	10,772	7,822	4,168	1,488	310	39	-
1999.....	2,583	4,151	5,910	8,140	10,591	10,260	7,307	3,933	1,372	294	46	-
Black, both sexes												
2004.....	1,491	2,014	2,303	2,586	2,463	2,336	1,619	714	249	52	5	1
2003.....	1,423	1,916	2,287	2,559	2,593	2,359	1,541	689	247	60	9	-
2002.....	1,435	1,897	2,343	2,618	2,564	2,374	1,446	650	226	45	15	1
2001.....	1,361	1,808	2,143	2,638	2,699	2,361	1,410	665	207	62	11	2
2000.....	1,383	1,774	2,198	2,641	2,817	2,319	1,266	673	210	54	9	1
1999.....	1,333	1,748	2,251	2,667	2,694	2,319	1,299	679	211	39	13	-
Black, male												
2004.....	986	1,310	1,540	1,603	1,520	1,378	915	370	119	22	1	1
2003.....	928	1,281	1,484	1,629	1,592	1,424	897	362	120	26	1	-
2002.....	944	1,239	1,575	1,674	1,591	1,451	841	346	106	11	2	1
2001.....	924	1,208	1,405	1,732	1,729	1,488	854	393	93	15	3	2
2000.....	935	1,220	1,421	1,744	1,776	1,456	766	371	107	24	5	1
1999.....	905	1,188	1,517	1,738	1,714	1,432	797	391	100	14	7	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant neoplasms of respiratory and intrathoracic organs (C30-C39)												
Black, female												
2004.....	505	704	763	983	943	958	704	344	130	30	4	-
2003.....	495	635	803	930	1,001	935	644	327	127	34	8	-
2002.....	491	658	768	944	973	923	605	304	120	34	13	-
2001.....	437	600	738	906	970	873	556	272	114	47	8	-
2000.....	448	554	777	897	1,041	863	500	302	103	30	4	-
1999.....	428	560	734	929	980	887	502	288	111	25	6	-
American Indian 1/, both sexes												
2004.....	35	77	95	144	118	94	53	22	10	2	-	-
2003.....	44	67	69	95	99	77	58	20	8	1	-	-
2002.....	37	64	91	96	99	80	48	26	8	-	1	-
2001.....	41	52	84	96	95	87	55	19	8	3	-	-
2000.....	29	57	75	85	89	69	38	21	3	2	-	-
1999.....	29	62	86	91	96	57	46	15	6	2	1	-
American Indian 1/, male												
2004.....	24	45	57	72	66	55	29	6	6	2	-	-
2003.....	24	42	45	48	52	42	29	9	2	1	-	-
2002.....	22	44	54	47	51	45	28	13	2	-	-	-
2001.....	24	29	48	52	55	48	30	9	-	2	-	-
2000.....	16	32	48	45	43	45	24	9	3	1	-	-
1999.....	19	31	50	51	61	30	25	10	4	2	1	-
American Indian 1/, female												
2004.....	11	32	38	72	52	39	24	16	4	-	-	-
2003.....	20	25	24	47	47	35	29	11	6	-	-	-
2002.....	15	20	37	49	48	35	20	13	6	-	1	-
2001.....	17	23	36	44	40	39	25	10	8	1	-	-
2000.....	13	25	27	40	46	24	14	12	-	1	-	-
1999.....	10	31	36	40	35	27	21	5	2	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant neoplasms of respiratory and intrathoracic organs (C30-C39)												
Asian or Pacific Islander 2/, both sexes												
2004.....	168	214	251	350	454	418	318	134	74	9	2	-
2003.....	163	171	244	332	393	434	323	162	55	11	1	-
2002.....	137	202	256	275	364	357	255	150	48	13	1	-
2001.....	132	188	236	318	370	354	293	158	59	10	1	-
2000.....	104	182	213	299	375	351	225	124	43	8	2	1
1999.....	114	144	206	319	371	309	213	114	43	11	-	1
Asian or Pacific Islander 2/, male												
2004.....	101	127	156	214	278	268	194	73	43	4	1	-
2003.....	95	101	155	216	236	283	189	89	27	7	-	-
2002.....	80	134	169	154	211	225	174	86	28	6	-	-
2001.....	79	126	152	191	225	224	173	84	35	6	-	-
2000.....	65	121	133	179	256	239	129	79	22	3	-	1
1999.....	64	87	129	203	241	202	133	55	25	4	-	1
Asian or Pacific Islander 2/, female												
2004.....	67	87	95	136	176	150	124	61	31	5	1	-
2003.....	68	70	89	116	157	151	134	73	28	4	1	-
2002.....	57	68	87	121	153	132	81	64	20	7	1	-
2001.....	53	62	84	127	145	130	120	74	24	4	1	-
2000.....	39	61	80	120	119	112	96	45	21	5	2	-
1999.....	50	57	77	116	130	107	80	59	18	7	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Of nasal cavity, middle ear and accessory sinuses (C30-C31)

All races, both sexes

2004.....	28	43	39	46	56	59	61	39	20	7	-	-
2003.....	38	33	51	47	53	73	47	24	22	7	2	-
2002.....	35	36	42	43	62	55	51	33	20	3	-	-
2001.....	31	43	43	41	50	63	70	43	25	4	1	-
2000.....	31	39	44	46	55	48	46	29	20	2	-	-
1999.....	43	37	34	50	53	60	47	36	22	3	1	-

All races, male

2004.....	15	30	26	25	37	31	31	17	9	1	-	-
2003.....	30	21	38	32	36	44	26	11	4	2	1	-
2002.....	24	25	31	30	38	30	23	11	11	1	-	-
2001.....	23	30	28	23	30	27	36	12	8	3	-	-
2000.....	21	27	30	29	35	29	21	12	6	-	-	-
1999.....	32	30	21	31	24	36	20	16	5	-	-	-

All races, female

2004.....	13	13	13	21	19	28	30	22	11	6	-	-
2003.....	8	12	13	15	17	29	21	13	18	5	1	-
2002.....	11	11	11	13	24	25	28	22	9	2	-	-
2001.....	8	13	15	18	20	36	34	31	17	1	1	-
2000.....	10	12	14	17	20	19	25	17	14	2	-	-
1999.....	11	7	13	19	29	24	27	20	17	3	1	-

White, both sexes

2004.....	20	37	30	38	51	50	57	33	20	7	-	-
2003.....	34	28	47	35	49	62	42	21	21	7	2	-
2002.....	30	34	38	36	55	52	49	29	20	3	-	-
2001.....	26	37	36	34	42	55	66	40	24	4	1	-
2000.....	25	29	37	39	48	45	42	27	20	2	-	-
1999.....	37	32	27	44	52	60	43	34	20	3	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Of nasal cavity, middle ear and accessory sinuses (C30-C31)												
White, male												
2004.....	10	25	21	19	34	26	28	14	9	1	-	-
2003.....	27	16	36	23	32	35	23	10	4	2	1	-
2002.....	22	23	28	25	32	28	21	10	11	1	-	-
2001.....	19	24	23	17	26	24	35	11	7	3	-	-
2000.....	17	20	25	24	32	28	21	11	6	-	-	-
1999.....	27	25	17	28	23	36	18	16	5	-	-	-
White, female												
2004.....	10	12	9	19	17	24	29	19	11	6	-	-
2003.....	7	12	11	12	17	27	19	11	17	5	1	-
2002.....	8	11	10	11	23	24	28	19	9	2	-	-
2001.....	7	13	13	17	16	31	31	29	17	1	1	-
2000.....	8	9	12	15	16	17	21	16	14	2	-	-
1999.....	10	7	10	16	29	24	25	18	15	3	1	-
Black, both sexes												
2004.....	7	3	5	5	5	5	4	4	-	-	-	-
2003.....	3	4	4	9	3	9	3	3	1	-	-	-
2002.....	5	1	4	5	6	2	1	3	-	-	-	-
2001.....	4	5	7	7	6	4	3	1	1	-	-	-
2000.....	5	9	7	7	5	2	4	2	-	-	-	-
1999.....	6	4	6	5	1	-	4	1	2	-	-	-
Black, male												
2004.....	4	3	3	4	3	4	3	2	-	-	-	-
2003.....	2	4	2	8	3	7	2	1	-	-	-	-
2002.....	2	1	3	4	6	2	1	-	-	-	-	-
2001.....	3	5	5	6	4	1	1	-	1	-	-	-
2000.....	3	6	5	5	3	-	-	1	-	-	-	-
1999.....	5	4	4	2	1	-	2	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Of nasal cavity, middle ear and accessory sinuses (C30-C31)												
Asian or Pacific Islander 2/, both sexes												
2004.....	1	3	4	2	-	4	-	1	-	-	-	-
2003.....	1	1	-	3	1	1	2	-	-	-	-	-
2002.....	-	1	-	2	1	1	1	1	-	-	-	-
2001.....	-	1	-	-	2	3	1	2	-	-	-	-
2000.....	1	1	-	-	1	1	-	-	-	-	-	-
1999.....	-	1	-	1	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	1	2	2	1	-	1	-	-	-	-	-	-
2003.....	1	1	-	1	1	1	1	-	-	-	-	-
2002.....	-	1	-	1	-	-	1	1	-	-	-	-
2001.....	-	1	-	-	-	2	-	1	-	-	-	-
2000.....	1	1	-	-	-	1	-	-	-	-	-	-
1999.....	-	1	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	-	1	2	1	-	3	-	1	-	-	-	-
2003.....	-	-	-	2	-	-	1	-	-	-	-	-
2002.....	-	-	-	1	1	1	-	-	-	-	-	-
2001.....	-	-	-	-	2	1	1	1	-	-	-	-
2000.....	-	-	-	-	1	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	1	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Of larynx (C32)												
All races, both sexes												
2004.....	282	426	510	526	525	523	335	215	77	22	2	-
2003.....	263	404	552	568	568	490	401	225	70	13	-	1
2002.....	269	412	492	508	576	588	381	172	76	5	-	1
2001.....	270	414	499	531	639	553	357	208	74	13	1	-
2000.....	270	384	489	574	645	595	401	191	73	11	1	-
1999.....	254	410	494	584	655	529	363	200	72	16	3	-
All races, male												
2004.....	228	351	405	435	419	413	263	160	55	18	2	-
2003.....	225	324	453	464	435	391	317	164	47	10	-	-
2002.....	219	339	399	395	437	455	300	126	56	1	-	1
2001.....	223	339	398	417	502	442	268	171	58	8	1	-
2000.....	218	319	407	441	504	451	314	155	51	8	-	-
1999.....	216	339	396	458	490	397	284	154	46	10	3	-
All races, female												
2004.....	54	75	105	91	106	110	72	55	22	4	-	-
2003.....	38	80	99	104	133	99	84	61	23	3	-	1
2002.....	50	73	93	113	139	133	81	46	20	4	-	-
2001.....	47	75	101	114	137	111	89	37	16	5	-	-
2000.....	52	65	82	133	141	144	87	36	22	3	1	-
1999.....	38	71	98	126	165	132	79	46	26	6	-	-
White, both sexes												
2004.....	197	328	407	427	434	453	287	194	62	20	2	-
2003.....	182	312	432	459	464	420	344	199	57	9	-	1
2002.....	192	317	383	416	485	494	333	158	71	4	-	-
2001.....	190	308	392	432	522	465	315	183	68	11	1	-
2000.....	191	279	364	466	521	501	358	171	70	10	1	-
1999.....	183	308	383	475	560	469	325	176	59	15	3	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Of larynx (C32)												
White, male												
2004.....	155	265	320	352	347	356	224	141	45	16	2	-
2003.....	154	247	358	365	353	331	268	144	39	7	-	-
2002.....	159	258	303	322	367	381	266	115	52	-	-	-
2001.....	155	251	316	339	412	366	232	152	55	7	1	-
2000.....	151	228	302	356	407	383	281	140	48	7	-	-
1999.....	158	252	310	372	417	352	249	133	36	9	3	-
White, female												
2004.....	42	63	87	75	87	97	63	53	17	4	-	-
2003.....	28	65	74	94	111	89	76	55	18	2	-	1
2002.....	33	59	80	94	118	113	67	43	19	4	-	-
2001.....	35	57	76	93	110	99	83	31	13	4	-	-
2000.....	40	51	62	110	114	118	77	31	22	3	1	-
1999.....	25	56	73	103	143	117	76	43	23	6	-	-
Black, both sexes												
2004.....	79	92	98	90	89	66	45	19	9	1	-	-
2003.....	77	91	115	101	96	65	50	24	11	3	-	-
2002.....	75	91	107	86	88	89	43	12	3	1	-	1
2001.....	77	101	99	88	103	77	34	20	6	1	-	-
2000.....	76	101	118	96	115	88	41	19	2	1	-	-
1999.....	69	98	107	103	87	57	34	15	10	1	-	-
Black, male												
2004.....	68	83	80	77	70	53	36	17	6	1	-	-
2003.....	67	76	91	92	77	56	43	18	6	2	-	-
2002.....	58	77	95	68	67	72	30	10	2	1	-	1
2001.....	65	83	78	69	78	66	28	15	3	-	-	-
2000.....	65	87	100	76	90	62	32	14	2	1	-	-
1999.....	56	85	84	81	67	42	32	13	7	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Of larynx (C32)												
Black, female												
2004.....	11	9	18	13	19	13	9	2	3	-	-	-
2003.....	10	15	24	9	19	9	7	6	5	1	-	-
2002.....	17	14	12	18	21	17	13	2	1	-	-	-
2001.....	12	18	21	19	25	11	6	5	3	1	-	-
2000.....	11	14	18	20	25	26	9	5	-	-	-	-
1999.....	13	13	23	22	20	15	2	2	3	-	-	-
American Indian 1/, both sexes												
2004.....	2	3	1	6	1	1	1	-	-	-	-	-
2003.....	3	-	2	1	2	2	1	1	-	-	-	-
2002.....	2	3	-	3	1	3	1	-	-	-	-	-
2001.....	1	1	2	6	4	2	3	1	-	1	-	-
2000.....	1	1	5	7	1	1	-	-	-	-	-	-
1999.....	2	1	2	2	2	-	-	3	-	-	-	-
American Indian 1/, male												
2004.....	1	1	1	4	1	1	1	-	-	-	-	-
2003.....	3	-	2	1	-	2	1	1	-	-	-	-
2002.....	2	3	-	3	1	-	1	-	-	-	-	-
2001.....	1	1	1	4	3	1	3	1	-	1	-	-
2000.....	1	1	3	5	-	1	-	-	-	-	-	-
1999.....	2	1	1	2	1	-	-	3	-	-	-	-
American Indian 1/, female												
2004.....	1	2	-	2	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	2	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	3	-	-	-	-	-	-
2001.....	-	-	1	2	1	1	-	-	-	-	-	-
2000.....	-	-	2	2	1	-	-	-	-	-	-	-
1999.....	-	-	1	-	1	-	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Of larynx (C32)												
Asian or Pacific Islander 2/, both sexes												
2004.....	4	3	4	3	1	3	2	2	6	1	-	-
2003.....	1	1	3	7	6	3	6	1	2	1	-	-
2002.....	-	1	2	3	2	2	4	2	2	-	-	-
2001.....	2	4	6	5	10	9	5	4	-	-	-	-
2000.....	2	3	2	5	8	5	2	1	1	-	-	-
1999.....	-	3	2	4	6	3	4	6	3	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	4	2	4	2	1	3	2	2	4	1	-	-
2003.....	1	1	2	6	5	2	5	1	2	1	-	-
2002.....	-	1	1	2	2	2	3	1	2	-	-	-
2001.....	2	4	3	5	9	9	5	3	-	-	-	-
2000.....	1	3	2	4	7	5	1	1	1	-	-	-
1999.....	-	1	1	3	5	3	3	5	3	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	-	1	-	1	-	-	-	-	2	-	-	-
2003.....	-	-	1	1	1	1	1	-	-	-	-	-
2002.....	-	-	1	1	-	-	1	1	-	-	-	-
2001.....	-	-	3	-	1	-	-	1	-	-	-	-
2000.....	1	-	-	1	1	-	1	-	-	-	-	-
1999.....	-	2	1	1	1	-	1	1	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Of trachea, bronchus and lung (C33-C34)												
All races, both sexes												
2004.....	7,884	12,846	18,240	22,552	26,477	27,608	20,856	10,325	3,418	637	75	3
2003.....	7,757	12,920	18,036	22,414	26,972	28,031	20,588	10,076	3,319	617	76	2
2002.....	7,809	12,930	17,754	22,358	27,898	27,969	20,113	9,781	3,214	576	81	-
2001.....	7,786	12,595	17,182	22,850	28,375	27,723	18,977	9,623	3,064	587	57	4
2000.....	7,695	12,453	17,268	23,438	28,820	27,305	18,531	9,302	2,902	542	60	5
1999.....	7,518	12,165	17,179	23,239	28,609	26,556	17,705	8,945	2,766	549	79	2
All races, male												
2004.....	4,751	7,663	10,777	13,243	15,210	15,754	11,238	5,246	1,537	235	21	3
2003.....	4,654	7,758	10,589	13,218	15,717	15,934	11,130	5,099	1,485	255	26	2
2002.....	4,723	7,769	10,646	13,209	16,334	15,833	10,988	5,022	1,439	216	25	-
2001.....	4,703	7,523	10,350	13,798	16,901	15,968	10,438	5,026	1,370	237	23	4
2000.....	4,583	7,546	10,391	14,189	17,193	15,726	10,240	4,840	1,335	204	16	5
1999.....	4,505	7,465	10,548	14,186	17,101	15,461	9,936	4,739	1,314	234	28	2
All races, female												
2004.....	3,133	5,183	7,463	9,309	11,267	11,854	9,618	5,079	1,881	402	54	-
2003.....	3,103	5,162	7,447	9,196	11,255	12,097	9,458	4,977	1,834	362	50	-
2002.....	3,086	5,161	7,108	9,149	11,564	12,136	9,125	4,759	1,775	360	56	-
2001.....	3,083	5,072	6,832	9,052	11,474	11,755	8,539	4,597	1,694	350	34	-
2000.....	3,112	4,907	6,877	9,249	11,627	11,579	8,291	4,462	1,567	338	44	-
1999.....	3,013	4,700	6,631	9,053	11,508	11,095	7,769	4,206	1,452	315	51	-
White, both sexes												
2004.....	6,294	10,653	15,711	19,590	23,546	24,848	18,922	9,483	3,103	576	68	2
2003.....	6,215	10,870	15,565	19,553	24,005	25,257	18,731	9,234	3,023	549	66	2
2002.....	6,288	10,873	15,182	19,480	24,973	25,262	18,416	8,973	2,937	519	64	-
2001.....	6,346	10,669	14,837	19,915	25,344	25,025	17,272	8,812	2,799	514	45	2
2000.....	6,272	10,562	14,925	20,532	25,676	24,672	17,054	8,508	2,649	479	49	3
1999.....	6,125	10,324	14,766	20,284	25,556	23,939	16,192	8,166	2,523	498	65	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Of trachea, bronchus and lung (C33-C34)												
White, male												
2004.....	3,726	6,276	9,118	11,449	13,424	14,118	10,143	4,819	1,380	209	19	2
2003.....	3,682	6,420	9,006	11,435	13,929	14,262	10,068	4,660	1,344	224	25	2
2002.....	3,743	6,441	8,950	11,416	14,558	14,193	9,981	4,589	1,307	200	23	-
2001.....	3,753	6,261	8,834	11,917	14,992	14,290	9,421	4,562	1,247	215	20	2
2000.....	3,641	6,274	8,905	12,313	15,221	14,059	9,356	4,398	1,206	177	11	3
1999.....	3,583	6,252	8,949	12,286	15,164	13,846	9,018	4,306	1,196	215	20	1
White, female												
2004.....	2,568	4,377	6,593	8,141	10,122	10,730	8,779	4,664	1,723	367	49	-
2003.....	2,533	4,450	6,559	8,118	10,076	10,995	8,663	4,574	1,679	325	41	-
2002.....	2,545	4,432	6,232	8,064	10,415	11,069	8,435	4,384	1,630	319	41	-
2001.....	2,593	4,408	6,003	7,998	10,352	10,735	7,851	4,250	1,552	299	25	-
2000.....	2,631	4,288	6,020	8,219	10,455	10,613	7,698	4,110	1,443	302	38	-
1999.....	2,542	4,072	5,817	7,998	10,392	10,093	7,174	3,860	1,327	283	45	-
Black, both sexes												
2004.....	1,398	1,913	2,193	2,484	2,361	2,258	1,566	690	238	51	5	1
2003.....	1,340	1,814	2,165	2,446	2,486	2,275	1,486	662	235	57	9	-
2002.....	1,353	1,797	2,229	2,516	2,467	2,277	1,400	635	223	44	15	-
2001.....	1,275	1,692	2,034	2,534	2,583	2,275	1,367	643	199	61	11	2
2000.....	1,295	1,660	2,066	2,534	2,692	2,222	1,216	651	208	53	9	1
1999.....	1,252	1,643	2,127	2,553	2,595	2,257	1,258	662	198	38	13	-
Black, male												
2004.....	910	1,221	1,454	1,518	1,444	1,320	875	351	112	21	1	1
2003.....	858	1,197	1,389	1,528	1,507	1,355	851	343	114	24	1	-
2002.....	882	1,156	1,475	1,598	1,517	1,373	810	336	104	10	2	-
2001.....	854	1,113	1,321	1,648	1,642	1,420	822	377	89	15	3	2
2000.....	864	1,125	1,312	1,661	1,680	1,390	732	355	105	23	5	1
1999.....	841	1,099	1,422	1,653	1,641	1,387	763	377	93	13	7	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Of trachea, bronchus and lung (C33-C34)												
Black, female												
2004.....	488	692	739	966	917	938	691	339	126	30	4	-
2003.....	482	617	776	918	979	920	635	319	121	33	8	-
2002.....	471	641	754	918	950	904	590	299	119	34	13	-
2001.....	421	579	713	886	941	855	545	266	110	46	8	-
2000.....	431	535	754	873	1,012	832	484	296	103	30	4	-
1999.....	411	544	705	900	954	870	495	285	105	25	6	-
American Indian 1/, both sexes												
2004.....	33	73	94	136	117	91	52	21	10	2	-	-
2003.....	41	67	67	94	97	74	56	19	8	1	-	-
2002.....	35	61	90	93	98	77	47	26	8	-	1	-
2001.....	39	51	82	90	91	82	52	17	8	2	-	-
2000.....	28	56	70	78	87	68	38	21	3	2	-	-
1999.....	27	59	83	89	94	57	46	12	6	2	1	-
American Indian 1/, male												
2004.....	23	44	56	67	65	52	28	5	6	2	-	-
2003.....	21	42	43	47	52	39	28	8	2	1	-	-
2002.....	20	41	53	44	50	45	27	13	2	-	-	-
2001.....	22	28	47	48	52	46	27	7	-	1	-	-
2000.....	15	31	45	40	43	44	24	9	3	1	-	-
1999.....	17	30	49	49	60	30	25	7	4	2	1	-
American Indian 1/, female												
2004.....	10	29	38	69	52	39	24	16	4	-	-	-
2003.....	20	25	24	47	45	35	28	11	6	-	-	-
2002.....	15	20	37	49	48	32	20	13	6	-	1	-
2001.....	17	23	35	42	39	36	25	10	8	1	-	-
2000.....	13	25	25	38	44	24	14	12	-	1	-	-
1999.....	10	29	34	40	34	27	21	5	2	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Of trachea, bronchus and lung (C33-C34)												
Asian or Pacific Islander 2/, both sexes												
2004.....	159	207	242	342	453	411	316	131	67	8	2	-
2003.....	161	169	239	321	384	425	315	161	53	10	1	-
2002.....	133	199	253	269	360	353	250	147	46	13	1	-
2001.....	126	183	229	311	357	341	286	151	58	10	1	-
2000.....	100	175	207	294	365	343	223	122	42	8	2	1
1999.....	114	139	203	313	364	303	209	105	39	11	-	1
Asian or Pacific Islander 2/, male												
2004.....	92	122	149	209	277	264	192	71	39	3	1	-
2003.....	93	99	151	208	229	278	183	88	25	6	-	-
2002.....	78	131	168	151	209	222	170	84	26	6	-	-
2001.....	74	121	148	185	215	212	168	80	34	6	-	-
2000.....	63	116	129	175	249	233	128	78	21	3	-	1
1999.....	64	84	128	198	236	198	130	49	21	4	-	1
Asian or Pacific Islander 2/, female												
2004.....	67	85	93	133	176	147	124	60	28	5	1	-
2003.....	68	70	88	113	155	147	132	73	28	4	1	-
2002.....	55	68	85	118	151	131	80	63	20	7	1	-
2001.....	52	62	81	126	142	129	118	71	24	4	1	-
2000.....	37	59	78	119	116	110	95	44	21	5	2	-
1999.....	50	55	75	115	128	105	79	56	18	7	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Of thymus, heart, mediastinum and pleura (C37-C38)

All races, both sexes

2004.....	34	34	43	46	47	50	49	28	13	3	2	-
2003.....	19	39	41	38	52	60	50	39	16	4	1	-
2002.....	28	27	34	51	55	58	40	36	13	3	-	-
2001.....	41	44	29	54	55	56	45	26	15	3	1	-
2000.....	26	32	36	48	41	61	62	20	13	4	-	-
1999.....	30	41	35	41	59	59	51	27	12	2	-	-

All races, male

2004.....	19	18	24	30	25	27	27	16	5	1	-	-
2003.....	10	22	22	21	27	31	25	18	6	-	-	-
2002.....	16	19	26	28	24	31	17	17	7	-	-	-
2001.....	22	34	12	32	32	23	23	15	4	1	-	-
2000.....	15	16	22	28	24	35	33	12	4	1	-	-
1999.....	21	22	21	18	28	29	19	14	4	-	-	-

All races, female

2004.....	15	16	19	16	22	23	22	12	8	2	2	-
2003.....	9	17	19	17	25	29	25	21	10	4	1	-
2002.....	12	8	8	23	31	27	23	19	6	3	-	-
2001.....	19	10	17	22	23	33	22	11	11	2	1	-
2000.....	11	16	14	20	17	26	29	8	9	3	-	-
1999.....	9	19	14	23	31	30	32	13	8	2	-	-

White, both sexes

2004.....	23	26	35	37	40	41	46	27	11	3	2	-
2003.....	16	33	36	35	44	45	48	39	16	4	1	-
2002.....	22	19	29	40	51	53	38	36	13	3	-	-
2001.....	34	34	26	45	47	49	40	24	13	3	1	-
2000.....	19	25	25	45	36	53	57	18	13	4	-	-
1999.....	25	35	23	36	47	52	48	24	10	2	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Of thymus, heart, mediastinum and pleura (C37-C38)

White, male

2004.....	11	14	20	25	23	24	26	16	5	1	-	-
2003.....	9	18	18	19	23	23	25	18	6	-	-	-
2002.....	12	13	23	25	23	28	17	17	7	-	-	-
2001.....	19	27	11	24	26	21	22	13	3	1	-	-
2000.....	12	13	16	27	22	31	31	11	4	1	-	-
1999.....	19	21	14	17	23	26	19	12	3	-	-	-

White, female

2004.....	12	12	15	12	17	17	20	11	6	2	2	-
2003.....	7	15	18	16	21	22	23	21	10	4	1	-
2002.....	10	6	6	15	28	25	21	19	6	3	-	-
2001.....	15	7	15	21	21	28	18	11	10	2	1	-
2000.....	7	12	9	18	14	22	26	7	9	3	-	-
1999.....	6	14	9	19	24	26	29	12	7	2	-	-

Black, both sexes

2004.....	7	6	7	5	7	7	3	1	1	-	-	-
2003.....	3	6	3	2	6	10	1	-	-	-	-	-
2002.....	2	7	3	10	3	4	2	-	-	-	-	-
2001.....	3	10	2	7	7	5	4	1	1	-	-	-
2000.....	6	4	7	3	4	6	5	1	-	-	-	-
1999.....	5	3	11	5	11	4	3	1	1	-	-	-

Black, male

2004.....	4	3	3	3	2	1	1	-	-	-	-	-
2003.....	1	4	2	1	3	6	-	-	-	-	-	-
2002.....	2	5	2	3	1	2	-	-	-	-	-	-
2001.....	-	7	-	7	5	1	1	1	-	-	-	-
2000.....	3	2	4	1	2	4	2	1	-	-	-	-
1999.....	2	-	7	1	5	2	-	1	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Of thymus, heart, mediastinum and pleura (C37-C38)												
Asian or Pacific Islander 2/, both sexes												
2004.....	4	1	1	3	-	-	-	-	1	-	-	-
2003.....	-	-	2	1	2	5	-	-	-	-	-	-
2002.....	4	1	1	1	1	1	-	-	-	-	-	-
2001.....	4	-	1	2	1	-	1	-	1	-	-	-
2000.....	1	3	4	-	1	2	-	1	-	-	-	-
1999.....	-	1	1	-	1	3	-	2	1	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	4	1	1	2	-	-	-	-	-	-	-	-
2003.....	-	-	2	1	1	2	-	-	-	-	-	-
2002.....	2	1	-	-	-	1	-	-	-	-	-	-
2001.....	3	-	1	1	1	-	-	-	1	-	-	-
2000.....	-	1	2	-	-	-	-	-	-	-	-	-
1999.....	-	1	-	-	-	1	-	1	1	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	-	-	-	1	-	-	-	-	1	-	-	-
2003.....	-	-	-	-	1	3	-	-	-	-	-	-
2002.....	2	-	1	1	1	-	-	-	-	-	-	-
2001.....	1	-	-	1	-	-	1	-	-	-	-	-
2000.....	1	2	2	-	1	2	-	1	-	-	-	-
1999.....	-	-	1	-	1	2	-	1	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Of other and ill-defined sites in respiratory system and intrathoracic organs (C39)												
All races, both sexes												
2004.....	-	1	-	2	1	3	3	-	1	-	-	-
2003.....	4	5	5	9	7	8	9	-	-	1	-	-
2002.....	3	7	5	4	5	9	8	1	-	-	-	-
2001.....	4	8	3	12	6	11	3	2	-	-	-	-
2000.....	4	4	1	9	5	7	3	7	-	-	-	-
1999.....	2	3	5	8	7	6	4	2	1	-	-	-
All races, male												
2004.....	-	1	-	1	1	2	-	-	1	-	-	-
2003.....	3	2	2	5	6	6	5	-	-	-	-	-
2002.....	-	3	3	3	5	7	3	1	-	-	-	-
2001.....	4	7	2	9	3	5	3	1	-	-	-	-
2000.....	3	4	-	7	3	4	3	3	-	-	-	-
1999.....	2	1	4	4	4	6	1	2	1	-	-	-
All races, female												
2004.....	-	-	-	1	-	1	3	-	-	-	-	-
2003.....	1	3	3	4	1	2	4	-	-	1	-	-
2002.....	3	4	2	1	-	2	5	-	-	-	-	-
2001.....	-	1	1	3	3	6	-	1	-	-	-	-
2000.....	1	-	1	2	2	3	-	4	-	-	-	-
1999.....	-	2	1	4	3	-	3	-	-	-	-	-
White, both sexes												
2004.....	-	1	-	-	-	3	2	-	-	-	-	-
2003.....	4	4	5	8	5	8	8	-	-	1	-	-
2002.....	3	6	5	3	5	7	8	1	-	-	-	-
2001.....	2	8	2	10	6	10	1	1	-	-	-	-
2000.....	3	4	1	8	4	6	3	7	-	-	-	-
1999.....	1	3	5	6	7	5	4	2	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Of other and ill-defined sites in respiratory system and intrathoracic organs (C39)

White, male

2004.....	-	1	-	-	-	2	-	-	-	-	-	-
2003.....	3	2	2	5	4	6	4	-	-	-	-	-
2002.....	-	3	3	2	5	5	3	1	-	-	-	-
2001.....	2	7	1	7	3	4	1	1	-	-	-	-
2000.....	3	4	-	6	2	4	3	3	-	-	-	-
1999.....	1	1	4	2	4	5	1	2	1	-	-	-

White, female

2004.....	-	-	-	-	-	1	2	-	-	-	-	-
2003.....	1	2	3	3	1	2	4	-	-	1	-	-
2002.....	3	3	2	1	-	2	5	-	-	-	-	-
2001.....	-	1	1	3	3	6	-	-	-	-	-	-
2000.....	-	-	1	2	2	2	-	4	-	-	-	-
1999.....	-	2	1	4	3	-	3	-	-	-	-	-

Black, both sexes

2004.....	-	-	-	2	1	-	1	-	1	-	-	-
2003.....	-	1	-	1	2	-	1	-	-	-	-	-
2002.....	-	1	-	1	-	2	-	-	-	-	-	-
2001.....	2	-	1	2	-	-	2	-	-	-	-	-
2000.....	1	-	-	1	1	1	-	-	-	-	-	-
1999.....	1	-	-	1	-	1	-	-	-	-	-	-

Black, male

2004.....	-	-	-	1	1	-	-	-	1	-	-	-
2003.....	-	-	-	-	2	-	1	-	-	-	-	-
2002.....	-	-	-	1	-	2	-	-	-	-	-	-
2001.....	2	-	1	2	-	-	2	-	-	-	-	-
2000.....	-	-	-	1	1	-	-	-	-	-	-	-
1999.....	1	-	-	1	-	1	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant neoplasms of bone and articular cartilage (C40-C41)												
All races, both sexes												
2004.....	67	80	94	94	96	113	120	90	31	10	1	-
2003.....	68	66	65	63	98	109	133	79	40	13	2	-
2002.....	51	77	76	62	89	100	107	85	37	7	3	-
2001.....	54	70	65	83	112	125	113	97	40	14	2	-
2000.....	51	73	62	89	86	131	106	86	45	11	2	-
1999.....	63	44	63	81	101	118	101	79	36	8	1	-
All races, male												
2004.....	43	50	50	52	42	50	57	34	13	1	-	-
2003.....	38	35	48	36	52	52	73	29	17	4	1	-
2002.....	39	45	41	33	42	56	48	34	13	4	1	-
2001.....	32	34	38	49	65	69	56	36	8	3	-	-
2000.....	39	45	36	53	45	67	50	35	15	2	-	-
1999.....	39	28	40	41	61	55	40	38	14	2	-	-
All races, female												
2004.....	24	30	44	42	54	63	63	56	18	9	1	-
2003.....	30	31	17	27	46	57	60	50	23	9	1	-
2002.....	12	32	35	29	47	44	59	51	24	3	2	-
2001.....	22	36	27	34	47	56	57	61	32	11	2	-
2000.....	12	28	26	36	41	64	56	51	30	9	2	-
1999.....	24	16	23	40	40	63	61	41	22	6	1	-
White, both sexes												
2004.....	55	66	77	83	89	97	109	86	27	10	1	-
2003.....	60	52	56	54	91	96	121	74	36	12	2	-
2002.....	41	66	63	52	77	89	99	80	34	7	3	-
2001.....	46	58	54	64	97	115	106	90	38	12	2	-
2000.....	47	57	54	82	79	117	98	82	42	9	2	-
1999.....	57	34	50	74	91	111	96	74	34	8	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasms of bone and articular cartilage (C40-C41)

White, male

2004.....	36	38	40	46	40	46	52	32	11	1	-	-
2003.....	34	31	40	31	49	50	68	29	13	3	1	-
2002.....	34	40	33	29	37	49	46	33	12	4	1	-
2001.....	26	28	34	40	60	62	54	32	8	2	-	-
2000.....	36	36	31	50	40	61	44	34	14	2	-	-
1999.....	34	22	32	38	55	51	38	36	14	2	-	-

White, female

2004.....	19	28	37	37	49	51	57	54	16	9	1	-
2003.....	26	21	16	23	42	46	53	45	23	9	1	-
2002.....	7	26	30	23	40	40	53	47	22	3	2	-
2001.....	20	30	20	24	37	53	52	58	30	10	2	-
2000.....	11	21	23	32	39	56	54	48	28	7	2	-
1999.....	23	12	18	36	36	60	58	38	20	6	1	-

Black, both sexes

2004.....	11	12	13	10	6	15	10	3	4	-	-	-
2003.....	8	12	8	7	6	12	10	4	2	1	-	-
2002.....	10	10	11	10	10	8	8	4	2	-	-	-
2001.....	7	10	11	13	13	7	6	7	2	1	-	-
2000.....	2	12	7	6	7	12	6	4	3	2	-	-
1999.....	6	7	12	7	8	6	4	5	2	-	-	-

Black, male

2004.....	7	10	6	6	1	4	4	1	2	-	-	-
2003.....	4	4	7	4	3	2	4	-	2	1	-	-
2002.....	5	4	8	4	5	4	2	1	1	-	-	-
2001.....	5	4	4	7	5	5	1	4	-	1	-	-
2000.....	1	6	5	3	5	5	5	1	1	-	-	-
1999.....	5	4	8	3	6	4	1	2	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant neoplasms of bone and articular cartilage (C40-C41)												
Black, female												
2004.....	4	2	7	4	5	11	6	2	2	-	-	-
2003.....	4	8	1	3	3	10	6	4	-	-	-	-
2002.....	5	6	3	6	5	4	6	3	1	-	-	-
2001.....	2	6	7	6	8	2	5	3	2	-	-	-
2000.....	1	6	2	3	2	7	1	3	2	2	-	-
1999.....	1	3	4	4	2	2	3	3	2	-	-	-
American Indian 1/, both sexes												
2004.....	-	-	3	1	1	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	1	-	1	-	-	-
2002.....	-	-	-	-	1	2	-	-	1	-	-	-
2001.....	-	-	-	2	-	1	-	-	-	-	-	-
2000.....	1	-	-	-	-	1	-	-	-	-	-	-
1999.....	-	1	-	-	-	1	-	-	-	-	-	-
American Indian 1/, male												
2004.....	-	-	3	-	1	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	1	-	-	-
2002.....	-	-	-	-	-	2	-	-	-	-	-	-
2001.....	-	-	-	-	-	1	-	-	-	-	-	-
2000.....	1	-	-	-	-	1	-	-	-	-	-	-
1999.....	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, female												
2004.....	-	-	-	1	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	1	-	-	-	-	-
2002.....	-	-	-	-	1	-	-	-	1	-	-	-
2001.....	-	-	-	2	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	1	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant neoplasms of bone and articular cartilage (C40-C41)												
Asian or Pacific Islander 2/, both sexes												
2004.....	1	2	1	-	-	1	1	1	-	-	-	-
2003.....	-	2	1	2	1	1	1	1	1	-	-	-
2002.....	-	1	2	-	1	1	-	1	-	-	-	-
2001.....	1	2	-	4	2	2	1	-	-	1	-	-
2000.....	1	4	1	1	-	1	2	-	-	-	-	-
1999.....	-	2	1	-	2	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	-	2	1	-	-	-	1	1	-	-	-	-
2003.....	-	-	1	1	-	-	1	-	1	-	-	-
2002.....	-	1	-	-	-	1	-	-	-	-	-	-
2001.....	1	2	-	2	-	1	1	-	-	-	-	-
2000.....	1	3	-	-	-	-	1	-	-	-	-	-
1999.....	-	1	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	1	-	-	-	-	1	-	-	-	-	-	-
2003.....	-	2	-	1	1	1	-	1	-	-	-	-
2002.....	-	-	2	-	1	-	-	1	-	-	-	-
2001.....	-	-	-	2	2	1	-	-	-	1	-	-
2000.....	-	1	1	1	-	1	1	-	-	-	-	-
1999.....	-	1	1	-	2	-	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Melanoma and other malignant neoplasms of skin (C43-C44)												
All races, both sexes												
2004.....	776	949	988	1,005	1,168	1,323	1,240	877	426	134	29	-
2003.....	765	943	962	922	1,157	1,364	1,171	825	439	155	31	1
2002.....	776	809	895	956	1,149	1,324	1,161	801	397	137	35	1
2001.....	806	832	893	973	1,246	1,273	1,081	804	390	152	28	-
2000.....	786	779	843	1,000	1,180	1,229	1,002	783	420	115	32	-
1999.....	718	748	853	1,007	1,160	1,260	1,038	701	359	127	26	-
All races, male												
2004.....	523	665	711	706	806	897	795	541	213	49	10	-
2003.....	524	661	666	634	780	922	728	499	205	64	5	-
2002.....	515	559	633	663	781	869	710	443	194	47	14	-
2001.....	536	571	629	689	863	830	665	438	191	66	6	-
2000.....	541	543	567	682	795	807	636	457	183	35	5	-
1999.....	472	530	591	684	800	809	634	380	185	30	9	-
All races, female												
2004.....	253	284	277	299	362	426	445	336	213	85	19	-
2003.....	241	282	296	288	377	442	443	326	234	91	26	1
2002.....	261	250	262	293	368	455	451	358	203	90	21	1
2001.....	270	261	264	284	383	443	416	366	199	86	22	-
2000.....	245	236	276	318	385	422	366	326	237	80	27	-
1999.....	246	218	262	323	360	451	404	321	174	97	17	-
White, both sexes												
2004.....	744	917	946	969	1,139	1,296	1,201	861	415	131	27	-
2003.....	729	913	925	890	1,129	1,320	1,149	807	428	151	30	1
2002.....	737	767	857	914	1,105	1,295	1,140	785	387	132	33	1
2001.....	776	807	847	929	1,201	1,242	1,053	789	379	148	26	-
2000.....	755	735	805	958	1,135	1,194	975	766	408	113	30	-
1999.....	691	722	816	977	1,127	1,219	1,017	685	351	125	26	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Melanoma and other malignant neoplasms of skin (C43-C44)												
White, male												
2004.....	502	641	678	688	790	880	775	535	211	47	9	-
2003.....	501	642	642	611	767	900	718	493	200	62	5	-
2002.....	485	526	610	632	752	853	702	435	189	47	14	-
2001.....	513	551	599	657	836	814	656	431	187	66	6	-
2000.....	518	514	544	655	771	787	622	455	178	35	4	-
1999.....	450	511	564	666	779	782	623	375	181	29	9	-
White, female												
2004.....	242	276	268	281	349	416	426	326	204	84	18	-
2003.....	228	271	283	279	362	420	431	314	228	89	25	1
2002.....	252	241	247	282	353	442	438	350	198	85	19	1
2001.....	263	256	248	272	365	428	397	358	192	82	20	-
2000.....	237	221	261	303	364	407	353	311	230	78	26	-
1999.....	241	211	252	311	348	437	394	310	170	96	17	-
Black, both sexes												
2004.....	29	28	33	29	23	23	29	11	9	3	2	-
2003.....	24	23	30	30	22	33	14	13	8	1	1	-
2002.....	36	37	28	34	29	22	16	14	9	2	1	-
2001.....	25	22	34	37	38	22	22	11	9	4	2	-
2000.....	25	38	32	33	37	27	24	15	8	2	1	-
1999.....	26	22	31	25	27	30	16	13	7	1	-	-
Black, male												
2004.....	20	21	26	16	12	15	15	4	2	2	1	-
2003.....	16	15	20	22	11	14	5	1	3	-	-	-
2002.....	27	30	18	25	19	13	6	6	5	-	-	-
2001.....	18	18	20	27	23	9	6	4	2	-	-	-
2000.....	19	25	21	20	18	16	12	1	2	-	-	-
1999.....	21	15	24	14	19	22	8	4	3	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Melanoma and other malignant neoplasms of skin (C43-C44)												
Black, female												
2004.....	9	7	7	13	11	8	14	7	7	1	1	-
2003.....	8	8	10	8	11	19	9	12	5	1	1	-
2002.....	9	7	10	9	10	9	10	8	4	2	1	-
2001.....	7	4	14	10	15	13	16	7	7	4	2	-
2000.....	6	13	11	13	19	11	12	14	6	2	1	-
1999.....	5	7	7	11	8	8	8	9	4	-	-	-
American Indian 1/, both sexes												
2004.....	3	2	3	3	2	-	3	-	-	-	-	-
2003.....	5	2	2	1	4	2	3	-	1	1	-	-
2002.....	1	3	2	2	6	2	1	1	-	2	1	-
2001.....	1	-	5	1	-	3	1	3	-	-	-	-
2000.....	2	2	1	3	1	1	1	-	-	-	-	-
1999.....	1	2	3	4	-	-	1	-	-	1	-	-
American Indian 1/, male												
2004.....	1	2	3	2	2	-	1	-	-	-	-	-
2003.....	2	1	1	-	2	-	2	-	1	1	-	-
2002.....	1	3	1	2	4	-	-	1	-	-	-	-
2001.....	1	-	4	1	-	2	1	3	-	-	-	-
2000.....	2	1	1	3	1	-	1	-	-	-	-	-
1999.....	1	2	1	3	-	-	-	-	-	-	-	-
American Indian 1/, female												
2004.....	2	-	-	1	-	-	2	-	-	-	-	-
2003.....	3	1	1	1	2	2	1	-	-	-	-	-
2002.....	-	-	1	-	2	2	1	-	-	2	1	-
2001.....	-	-	1	-	-	1	-	-	-	-	-	-
2000.....	-	1	-	-	-	1	-	-	-	-	-	-
1999.....	-	-	2	1	-	-	1	-	-	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Melanoma and other malignant neoplasms of skin (C43-C44)												
Asian or Pacific Islander 2/, both sexes												
2004.....	-	2	6	4	4	4	7	5	2	-	-	-
2003.....	7	5	5	1	2	9	5	5	2	2	-	-
2002.....	2	2	8	6	9	5	4	1	1	1	-	-
2001.....	4	3	7	6	7	6	5	1	2	-	-	-
2000.....	4	4	5	6	7	7	2	2	4	-	1	-
1999.....	-	2	3	1	6	11	4	3	1	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	-	1	4	-	2	2	4	2	-	-	-	-
2003.....	5	3	3	1	-	8	3	5	1	1	-	-
2002.....	2	-	4	4	6	3	2	1	-	-	-	-
2001.....	4	2	6	4	4	5	2	-	2	-	-	-
2000.....	2	3	1	4	5	4	1	1	3	-	1	-
1999.....	-	2	2	1	2	5	3	1	1	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	-	1	2	4	2	2	3	3	2	-	-	-
2003.....	2	2	2	-	2	1	2	-	1	1	-	-
2002.....	-	2	4	2	3	2	2	-	1	1	-	-
2001.....	-	1	1	2	3	1	3	1	-	-	-	-
2000.....	2	1	4	2	2	3	1	1	1	-	-	-
1999.....	-	-	1	-	4	6	1	2	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant melanoma of skin (C43)												
All races, both sexes												
2004.....	650	776	771	806	905	1,008	904	534	244	72	9	-
2003.....	638	778	769	710	883	1,033	842	500	243	78	10	-
2002.....	640	659	692	748	868	954	794	512	233	58	17	1
2001.....	672	653	680	746	943	919	759	489	213	72	11	-
2000.....	665	627	676	753	884	887	737	495	252	52	4	-
1999.....	602	615	648	791	861	945	694	435	181	53	12	-
All races, male												
2004.....	428	525	533	546	612	674	565	326	116	26	5	-
2003.....	425	528	514	471	579	692	511	291	110	32	3	-
2002.....	403	445	470	500	575	609	469	264	115	21	6	-
2001.....	425	430	463	524	632	600	463	256	99	31	2	-
2000.....	448	420	446	495	572	559	454	277	100	13	-	-
1999.....	380	425	426	526	570	596	410	223	90	11	4	-
All races, female												
2004.....	222	251	238	260	293	334	339	208	128	46	4	-
2003.....	213	250	255	239	304	341	331	209	133	46	7	-
2002.....	237	214	222	248	293	345	325	248	118	37	11	1
2001.....	247	223	217	222	311	319	296	233	114	41	9	-
2000.....	217	207	230	258	312	328	283	218	152	39	4	-
1999.....	222	190	222	265	291	349	284	212	91	42	8	-
White, both sexes												
2004.....	638	766	759	785	891	992	879	525	238	70	7	-
2003.....	623	760	753	694	870	1,005	829	490	236	77	9	-
2002.....	628	642	677	731	843	938	783	501	227	55	16	1
2001.....	663	647	665	724	920	903	742	481	209	71	10	-
2000.....	654	611	662	736	862	871	717	483	243	50	4	-
1999.....	598	608	636	777	846	924	680	424	176	52	12	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant melanoma of skin (C43)												
White, male												
2004.....	423	518	527	540	606	665	553	323	114	25	4	-
2003.....	417	519	505	461	577	679	504	288	107	32	3	-
2002.....	395	433	462	490	561	600	466	259	113	21	6	-
2001.....	421	427	456	513	622	592	458	253	99	31	2	-
2000.....	441	411	440	484	563	552	444	276	96	13	-	-
1999.....	377	420	418	519	564	584	404	219	88	10	4	-
White, female												
2004.....	215	248	232	245	285	327	326	202	124	45	3	-
2003.....	206	241	248	233	293	326	325	202	129	45	6	-
2002.....	233	209	215	241	282	338	317	242	114	34	10	1
2001.....	242	220	209	211	298	311	284	228	110	40	8	-
2000.....	213	200	222	252	299	319	273	207	147	37	4	-
1999.....	221	188	218	258	282	340	276	205	88	42	8	-
Black, both sexes												
2004.....	9	8	8	14	13	14	20	7	5	2	2	-
2003.....	8	12	11	16	10	20	8	7	5	-	1	-
2002.....	9	13	8	11	14	10	8	10	5	-	-	-
2001.....	6	5	8	18	19	9	16	5	4	1	1	-
2000.....	7	11	9	11	19	14	18	12	7	2	-	-
1999.....	3	5	9	11	14	14	11	8	5	1	-	-
Black, male												
2004.....	4	6	4	4	5	8	10	2	2	1	1	-
2003.....	4	6	6	10	2	7	3	-	2	-	-	-
2002.....	5	10	3	5	7	6	2	4	2	-	-	-
2001.....	1	3	1	9	8	2	4	1	-	-	-	-
2000.....	5	6	5	7	7	6	9	1	2	-	-	-
1999.....	2	3	7	4	6	8	4	3	2	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant melanoma of skin (C43)												
Black, female												
2004.....	5	2	4	10	8	6	10	5	3	1	1	-
2003.....	4	6	5	6	8	13	5	7	3	-	1	-
2002.....	4	3	5	6	7	4	6	6	3	-	-	-
2001.....	5	2	7	9	11	7	12	4	4	1	1	-
2000.....	2	5	4	4	12	8	9	11	5	2	-	-
1999.....	1	2	2	7	8	6	7	5	3	-	-	-
American Indian 1/, both sexes												
2004.....	3	1	1	3	1	-	1	-	-	-	-	-
2003.....	2	2	2	-	2	2	3	-	-	-	-	-
2002.....	1	2	2	1	5	2	-	-	-	2	1	-
2001.....	1	-	3	-	-	3	-	2	-	-	-	-
2000.....	1	2	1	1	-	1	1	-	-	-	-	-
1999.....	1	-	3	2	-	-	1	-	-	-	-	-
American Indian 1/, male												
2004.....	1	1	1	2	1	-	-	-	-	-	-	-
2003.....	1	1	1	-	-	-	2	-	-	-	-	-
2002.....	1	2	1	1	3	-	-	-	-	-	-	-
2001.....	1	-	3	-	-	2	-	2	-	-	-	-
2000.....	1	1	1	1	-	-	1	-	-	-	-	-
1999.....	1	-	1	2	-	-	-	-	-	-	-	-
American Indian 1/, female												
2004.....	2	-	-	1	-	-	1	-	-	-	-	-
2003.....	1	1	1	-	2	2	1	-	-	-	-	-
2002.....	-	-	1	-	2	2	-	-	-	2	1	-
2001.....	-	-	-	-	-	1	-	-	-	-	-	-
2000.....	-	1	-	-	-	1	-	-	-	1	-	-
1999.....	-	-	2	-	-	-	1	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant melanoma of skin (C43)												
Asian or Pacific Islander 2/, both sexes												
2004.....	-	1	3	4	-	2	4	2	1	-	-	-
2003.....	5	4	3	-	1	6	2	3	2	1	-	-
2002.....	2	2	5	5	6	4	3	1	1	1	-	-
2001.....	2	1	4	4	4	4	1	1	-	-	-	-
2000.....	3	3	4	5	3	1	1	-	2	-	-	-
1999.....	-	2	-	1	1	7	2	3	-	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	-	-	1	-	-	1	2	1	-	-	-	-
2003.....	3	2	2	-	-	6	2	3	1	-	-	-
2002.....	2	-	4	4	4	3	1	1	-	-	-	-
2001.....	2	-	3	2	2	4	1	-	-	-	-	-
2000.....	1	2	-	3	2	1	-	-	2	-	-	-
1999.....	-	2	-	1	-	4	2	1	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	-	1	2	4	-	1	2	1	1	-	-	-
2003.....	2	2	1	-	1	-	-	-	1	1	-	-
2002.....	-	2	1	1	2	1	2	-	1	1	-	-
2001.....	-	1	1	2	2	-	-	1	-	-	-	-
2000.....	2	1	4	2	1	-	1	-	-	-	-	-
1999.....	-	-	-	-	1	3	-	2	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other malignant neoplasms of skin (C44)												
All races, both sexes												
2004.....	126	173	217	199	263	315	336	343	182	62	20	-
2003.....	127	165	193	212	274	331	329	325	196	77	21	1
2002.....	136	150	203	208	281	370	367	289	164	79	18	-
2001.....	134	179	213	227	303	354	322	315	177	80	17	-
2000.....	121	152	167	247	296	342	265	288	168	63	28	-
1999.....	116	133	205	216	299	315	344	266	178	74	14	-
All races, male												
2004.....	95	140	178	160	194	223	230	215	97	23	5	-
2003.....	99	133	152	163	201	230	217	208	95	32	2	-
2002.....	112	114	163	163	206	260	241	179	79	26	8	-
2001.....	111	141	166	165	231	230	202	182	92	35	4	-
2000.....	93	123	121	187	223	248	182	180	83	22	5	-
1999.....	92	105	165	158	230	213	224	157	95	19	5	-
All races, female												
2004.....	31	33	39	39	69	92	106	128	85	39	15	-
2003.....	28	32	41	49	73	101	112	117	101	45	19	1
2002.....	24	36	40	45	75	110	126	110	85	53	10	-
2001.....	23	38	47	62	72	124	120	133	85	45	13	-
2000.....	28	29	46	60	73	94	83	108	85	41	23	-
1999.....	24	28	40	58	69	102	120	109	83	55	9	-
White, both sexes												
2004.....	106	151	187	184	248	304	322	336	177	61	20	-
2003.....	106	153	172	196	259	315	320	317	192	74	21	1
2002.....	109	125	180	183	262	357	357	284	160	77	17	-
2001.....	113	160	182	205	281	339	311	308	170	77	16	-
2000.....	101	124	143	222	273	323	258	283	165	63	26	-
1999.....	93	114	180	200	281	295	337	261	175	73	14	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other malignant neoplasms of skin (C44)												
White, male												
2004.....	79	123	151	148	184	215	222	212	97	22	5	-
2003.....	84	123	137	150	190	221	214	205	93	30	2	-
2002.....	90	93	148	142	191	253	236	176	76	26	8	-
2001.....	92	124	143	144	214	222	198	178	88	35	4	-
2000.....	77	103	104	171	208	235	178	179	82	22	4	-
1999.....	73	91	146	147	215	198	219	156	93	19	5	-
White, female												
2004.....	27	28	36	36	64	89	100	124	80	39	15	-
2003.....	22	30	35	46	69	94	106	112	99	44	19	1
2002.....	19	32	32	41	71	104	121	108	84	51	9	-
2001.....	21	36	39	61	67	117	113	130	82	42	12	-
2000.....	24	21	39	51	65	88	80	104	83	41	22	-
1999.....	20	23	34	53	66	97	118	105	82	54	9	-
Black, both sexes												
2004.....	20	20	25	15	10	9	9	4	4	1	-	-
2003.....	16	11	19	14	12	13	6	6	3	1	-	-
2002.....	27	24	20	23	15	12	8	4	4	2	1	-
2001.....	19	17	26	19	19	13	6	6	5	3	1	-
2000.....	18	27	23	22	18	13	6	3	1	-	1	-
1999.....	23	17	22	14	13	16	5	5	2	-	-	-
Black, male												
2004.....	16	15	22	12	7	7	5	2	-	1	-	-
2003.....	12	9	14	12	9	7	2	1	1	-	-	-
2002.....	22	20	15	20	12	7	4	2	3	-	-	-
2001.....	17	15	19	18	15	7	2	3	2	-	-	-
2000.....	14	19	16	13	11	10	3	-	-	-	-	-
1999.....	19	12	17	10	13	14	4	1	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other malignant neoplasms of skin (C44)												
Black, female												
2004.....	4	5	3	3	3	2	4	2	4	-	-	-
2003.....	4	2	5	2	3	6	4	5	2	1	-	-
2002.....	5	4	5	3	3	5	4	2	1	2	1	-
2001.....	2	2	7	1	4	6	4	3	3	3	1	-
2000.....	4	8	7	9	7	3	3	3	1	-	1	-
1999.....	4	5	5	4	-	2	1	4	1	-	-	-
American Indian 1/, both sexes												
2004.....	-	1	2	-	1	-	2	-	-	-	-	-
2003.....	3	-	-	1	2	-	-	-	1	1	-	-
2002.....	-	1	-	1	1	-	1	1	-	-	-	-
2001.....	-	-	2	1	-	-	1	1	-	-	-	-
2000.....	1	-	-	2	1	-	-	-	-	-	-	-
1999.....	-	2	-	2	-	-	-	-	-	1	-	-
American Indian 1/, male												
2004.....	-	1	2	-	1	-	1	-	-	-	-	-
2003.....	1	-	-	-	2	-	-	-	1	1	-	-
2002.....	-	1	-	1	1	-	-	1	-	-	-	-
2001.....	-	-	1	1	-	-	1	1	-	-	-	-
2000.....	1	-	-	2	1	-	-	-	-	-	-	-
1999.....	-	2	-	1	-	-	-	-	-	-	-	-
American Indian 1/, female												
2004.....	-	-	-	-	-	-	1	-	-	-	-	-
2003.....	2	-	-	1	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	1	-	-	-	-	-
2001.....	-	-	1	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	1	-	-	-	-	-	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other malignant neoplasms of skin (C44)												
Asian or Pacific Islander 2/, both sexes												
2004.....	-	1	3	-	4	2	3	3	1	-	-	-
2003.....	2	1	2	1	1	3	3	2	-	1	-	-
2002.....	-	-	3	1	3	1	1	-	-	-	-	-
2001.....	2	2	3	2	3	2	4	-	2	-	-	-
2000.....	1	1	1	1	4	6	1	2	2	-	1	-
1999.....	-	-	3	-	5	4	2	-	1	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	-	1	3	-	2	1	2	1	-	-	-	-
2003.....	2	1	1	1	-	2	1	2	-	1	-	-
2002.....	-	-	-	-	2	-	1	-	-	-	-	-
2001.....	2	2	3	2	2	1	1	-	2	-	-	-
2000.....	1	1	1	1	3	3	1	1	1	-	1	-
1999.....	-	-	2	-	2	1	1	-	1	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	-	-	-	-	2	1	1	2	1	-	-	-
2003.....	-	-	1	-	1	1	2	-	-	-	-	-
2002.....	-	-	3	1	1	1	-	-	-	-	-	-
2001.....	-	-	-	-	1	1	3	-	-	-	-	-
2000.....	-	-	-	-	1	3	-	1	1	-	-	-
1999.....	-	-	1	-	3	3	1	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant neoplasms of mesothelial and soft tissue (C45-C49)												
All races, both sexes												
2004.....	378	550	660	759	848	1,108	983	539	182	31	11	-
2003.....	406	496	661	740	867	1,102	913	489	179	28	5	-
2002.....	369	510	635	707	947	1,026	868	504	175	36	8	1
2001.....	423	476	586	734	928	1,090	809	464	162	27	5	-
2000.....	386	499	572	773	988	1,066	801	453	167	37	7	-
1999.....	387	455	597	733	975	1,029	775	433	147	31	3	-
All races, male												
2004.....	185	272	401	446	506	700	589	305	95	17	3	-
2003.....	209	252	364	437	530	697	557	277	74	10	2	-
2002.....	184	269	369	416	602	639	505	296	80	13	2	1
2001.....	232	245	333	442	594	651	481	252	74	9	2	-
2000.....	191	276	341	476	593	668	480	246	88	13	1	-
1999.....	210	272	350	459	606	626	464	223	59	12	-	-
All races, female												
2004.....	193	278	259	313	342	408	394	234	87	14	8	-
2003.....	197	244	297	303	337	405	356	212	105	18	3	-
2002.....	185	241	266	291	345	387	363	208	95	23	6	-
2001.....	191	231	253	292	334	439	328	212	88	18	3	-
2000.....	195	223	231	297	395	398	321	207	79	24	6	-
1999.....	177	183	247	274	369	403	311	210	88	19	3	-
White, both sexes												
2004.....	308	470	609	691	787	1,041	926	515	170	30	7	-
2003.....	339	445	575	669	804	1,021	870	460	171	27	4	-
2002.....	318	441	562	637	872	964	816	475	165	36	8	1
2001.....	356	411	535	681	862	1,020	769	431	154	25	4	-
2000.....	323	441	509	689	915	1,007	767	425	155	34	6	-
1999.....	333	405	524	669	901	953	724	405	139	28	3	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant neoplasms of mesothelial and soft tissue (C45-C49)												
White, male												
2004.....	154	237	380	413	475	664	559	293	89	16	3	-
2003.....	175	230	320	399	499	655	542	259	71	10	1	-
2002.....	163	238	336	385	564	606	483	284	76	13	2	1
2001.....	204	214	310	417	565	617	459	237	72	8	1	-
2000.....	172	251	306	430	556	635	461	233	82	11	1	-
1999.....	186	249	312	425	565	588	443	214	58	11	-	-
White, female												
2004.....	154	233	229	278	312	377	367	222	81	14	4	-
2003.....	164	215	255	270	305	366	328	201	100	17	3	-
2002.....	155	203	226	252	308	358	333	191	89	23	6	-
2001.....	152	197	225	264	297	403	310	194	82	17	3	-
2000.....	151	190	203	259	359	372	306	192	73	23	5	-
1999.....	147	156	212	244	336	365	281	191	81	17	3	-
Black, both sexes												
2004.....	62	65	35	58	46	53	51	17	11	1	4	-
2003.....	55	40	69	56	52	66	33	25	8	1	-	-
2002.....	43	56	54	62	56	53	45	20	6	-	-	-
2001.....	59	56	38	42	56	50	30	27	7	1	1	-
2000.....	49	51	53	70	66	48	28	21	11	2	1	-
1999.....	42	38	65	46	62	64	40	25	6	2	-	-
Black, male												
2004.....	28	29	16	27	24	28	26	8	5	1	-	-
2003.....	28	21	36	26	23	34	11	16	3	-	-	-
2002.....	19	27	24	27	29	26	19	8	3	-	-	-
2001.....	27	28	16	20	24	23	15	12	2	-	1	-
2000.....	12	24	30	36	33	26	15	8	6	1	-	-
1999.....	20	21	34	23	31	31	16	8	1	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasms of mesothelial and soft tissue (C45-C49)

Black, female

2004.....	34	36	19	31	22	25	25	9	6	-	4	-
2003.....	27	19	33	30	29	32	22	9	5	1	-	-
2002.....	24	29	30	35	27	27	26	12	3	-	-	-
2001.....	32	28	22	22	32	27	15	15	5	1	-	-
2000.....	37	27	23	34	33	22	13	13	5	1	1	-
1999.....	22	17	31	23	31	33	24	17	5	1	-	-

American Indian 1/, both sexes

2004.....	2	3	1	1	2	1	2	1	-	-	-	-
2003.....	2	3	6	3	2	3	1	2	-	-	-	-
2002.....	-	2	6	-	3	1	2	1	2	-	-	-
2001.....	2	1	2	3	-	4	1	-	1	-	-	-
2000.....	-	-	1	3	1	3	-	-	-	-	-	-
1999.....	3	4	-	2	-	4	4	1	-	-	-	-

American Indian 1/, male

2004.....	-	1	-	-	2	-	1	1	-	-	-	-
2003.....	1	1	1	3	1	1	-	1	-	-	-	-
2002.....	-	1	4	-	3	1	2	-	-	-	-	-
2001.....	-	1	2	2	-	1	1	-	-	-	-	-
2000.....	-	-	-	2	1	3	-	-	-	-	-	-
1999.....	1	1	-	-	-	2	2	-	-	-	-	-

American Indian 1/, female

2004.....	2	2	1	1	-	1	1	-	-	-	-	-
2003.....	1	2	5	-	1	2	1	1	-	-	-	-
2002.....	-	1	2	-	-	-	-	1	2	-	-	-
2001.....	2	-	-	1	-	3	-	-	1	-	-	-
2000.....	-	-	1	1	-	-	-	-	-	-	-	-
1999.....	2	3	-	2	-	2	2	1	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant neoplasms of mesothelial and soft tissue (C45-C49)												
Asian or Pacific Islander 2/, both sexes												
2004.....	6	12	15	9	13	13	4	6	1	-	-	-
2003.....	10	8	11	12	9	12	9	2	-	-	1	-
2002.....	8	11	13	8	16	8	5	8	2	-	-	-
2001.....	6	8	11	8	10	16	9	6	-	1	-	-
2000.....	14	7	9	11	6	8	6	7	1	1	-	-
1999.....	9	8	8	16	12	8	7	2	2	1	-	-
Asian or Pacific Islander 2/, male												
2004.....	3	5	5	6	5	8	3	3	1	-	-	-
2003.....	5	-	7	9	7	7	4	1	-	-	1	-
2002.....	2	3	5	4	6	6	1	4	1	-	-	-
2001.....	1	2	5	3	5	10	6	3	-	1	-	-
2000.....	7	1	5	8	3	4	4	5	-	1	-	-
1999.....	3	1	4	11	10	5	3	1	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	3	7	10	3	8	5	1	3	-	-	-	-
2003.....	5	8	4	3	2	5	5	1	-	-	-	-
2002.....	6	8	8	4	10	2	4	4	1	-	-	-
2001.....	5	6	6	5	5	6	3	3	-	-	-	-
2000.....	7	6	4	3	3	4	2	2	1	-	-	-
1999.....	6	7	4	5	2	3	4	1	2	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Mesothelioma (C45)												
All races, both sexes												
2004.....	80	132	253	288	357	561	467	219	50	7	1	-
2003.....	89	134	234	311	376	529	437	209	51	4	3	-
2002.....	72	148	219	286	442	520	391	205	58	9	-	1
2001.....	93	128	212	294	421	536	353	187	51	3	2	-
2000.....	83	150	206	317	458	491	357	186	50	7	-	-
1999.....	88	159	210	320	452	483	349	160	34	4	-	-
All races, male												
2004.....	60	99	211	236	288	480	368	170	40	6	1	-
2003.....	69	101	196	255	318	445	351	171	29	4	1	-
2002.....	54	114	187	249	378	431	314	170	43	8	-	1
2001.....	69	102	170	249	354	432	284	147	39	2	1	-
2000.....	62	123	173	260	384	407	284	144	41	5	-	-
1999.....	66	122	168	267	374	389	285	122	27	3	-	-
All races, female												
2004.....	20	33	42	52	69	81	99	49	10	1	-	-
2003.....	20	33	38	56	58	84	86	38	22	-	2	-
2002.....	18	34	32	37	64	89	77	35	15	1	-	-
2001.....	24	26	42	45	67	104	69	40	12	1	1	-
2000.....	21	27	33	57	74	84	73	42	9	2	-	-
1999.....	22	37	42	53	78	94	64	38	7	1	-	-
White, both sexes												
2004.....	72	122	243	276	342	543	447	212	48	7	1	-
2003.....	77	130	215	293	361	506	424	200	50	4	2	-
2002.....	64	138	206	273	422	494	379	201	55	9	-	1
2001.....	84	119	207	284	408	524	340	182	50	3	2	-
2000.....	76	137	193	291	436	475	350	179	48	7	-	-
1999.....	79	150	194	307	435	459	332	153	33	4	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Mesothelioma (C45)												
White, male												
2004.....	53	93	202	227	276	463	349	165	40	6	1	-
2003.....	58	97	180	240	305	424	344	163	28	4	-	-
2002.....	49	106	179	237	361	409	304	168	41	8	-	1
2001.....	64	94	165	242	344	421	273	143	38	2	1	-
2000.....	55	113	162	240	365	395	277	140	39	5	-	-
1999.....	59	116	154	259	360	371	275	117	27	3	-	-
White, female												
2004.....	19	29	41	49	66	80	98	47	8	1	-	-
2003.....	19	33	35	53	56	82	80	37	22	-	2	-
2002.....	15	32	27	36	61	85	75	33	14	1	-	-
2001.....	20	25	42	42	64	103	67	39	12	1	1	-
2000.....	21	24	31	51	71	80	73	39	9	2	-	-
1999.....	20	34	40	48	75	88	57	36	6	1	-	-
Black, both sexes												
2004.....	8	7	9	9	12	12	17	6	2	-	-	-
2003.....	10	4	16	12	10	18	10	9	1	-	-	-
2002.....	6	9	7	12	15	20	11	4	2	-	-	-
2001.....	8	6	5	7	11	8	8	3	1	-	-	-
2000.....	5	11	11	22	20	13	7	4	2	-	-	-
1999.....	8	7	14	10	13	19	14	5	1	-	-	-
Black, male												
2004.....	7	5	9	6	10	12	17	4	-	-	-	-
2003.....	9	4	13	10	10	17	5	8	1	-	-	-
2002.....	5	8	5	11	12	16	9	2	2	-	-	-
2001.....	5	5	5	5	9	7	6	3	1	-	-	-
2000.....	5	9	9	17	17	9	7	2	2	-	-	-
1999.....	7	6	13	6	10	14	8	4	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Mesothelioma (C45)												
Black, female												
2004.....	1	2	-	3	2	-	-	2	2	-	-	-
2003.....	1	-	3	2	-	1	5	1	-	-	-	-
2002.....	1	1	2	1	3	4	2	2	-	-	-	-
2001.....	3	1	-	2	2	1	2	-	-	-	-	-
2000.....	-	2	2	5	3	4	-	2	-	-	-	-
1999.....	1	1	1	4	3	5	6	1	1	-	-	-
American Indian 1/, both sexes												
2004.....	-	-	-	-	1	1	2	-	-	-	-	-
2003.....	-	-	-	2	1	1	-	-	-	-	-	-
2002.....	-	1	2	-	2	1	-	-	1	-	-	-
2001.....	-	1	-	1	-	-	1	-	-	-	-	-
2000.....	-	-	-	2	1	1	-	-	-	-	-	-
1999.....	1	1	-	-	-	3	1	1	-	-	-	-
American Indian 1/, male												
2004.....	-	-	-	-	1	-	1	-	-	-	-	-
2003.....	-	-	-	2	-	-	-	-	-	-	-	-
2002.....	-	-	2	-	2	1	-	-	-	-	-	-
2001.....	-	1	-	1	-	-	1	-	-	-	-	-
2000.....	-	-	-	2	1	1	-	-	-	-	-	-
1999.....	-	-	-	-	-	2	1	-	-	-	-	-
American Indian 1/, female												
2004.....	-	-	-	-	-	1	1	-	-	-	-	-
2003.....	-	-	-	-	1	1	-	-	-	-	-	-
2002.....	-	1	-	-	-	-	-	-	1	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	1	1	-	-	-	1	-	1	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Mesothelioma (C45)												
Asian or Pacific Islander 2/, both sexes												
2004.....	-	3	1	3	2	5	1	1	-	-	-	-
2003.....	2	-	3	4	4	4	3	-	-	-	1	-
2002.....	2	-	4	1	3	5	1	-	-	-	-	-
2001.....	1	2	-	2	2	4	4	2	-	-	-	-
2000.....	2	2	2	2	1	2	-	3	-	-	-	-
1999.....	-	1	2	3	4	2	2	1	-	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	-	1	-	3	1	5	1	1	-	-	-	-
2003.....	2	-	3	3	3	4	2	-	-	-	1	-
2002.....	-	-	1	1	3	5	1	-	-	-	-	-
2001.....	-	2	-	1	1	4	4	1	-	-	-	-
2000.....	2	1	2	1	1	2	-	2	-	-	-	-
1999.....	-	-	1	2	4	2	1	1	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	-	2	1	-	1	-	-	-	-	-	-	-
2003.....	-	-	-	1	1	-	1	-	-	-	-	-
2002.....	2	-	3	-	-	-	-	-	-	-	-	-
2001.....	1	-	-	1	1	-	-	1	-	-	-	-
2000.....	-	1	-	1	-	-	-	1	-	-	-	-
1999.....	-	1	1	1	-	-	1	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Kaposi's sarcoma (C46)												
All races, both sexes												
2004.....	-	2	-	1	1	3	6	10	6	3	1	-
2003.....	1	-	3	1	2	9	4	8	7	4	-	-
2002.....	2	1	1	1	2	2	8	11	10	1	-	-
2001.....	3	1	-	-	2	2	6	9	7	-	-	-
2000.....	1	2	1	2	6	7	10	5	7	2	1	-
1999.....	1	-	-	-	1	3	6	7	9	1	-	-
All races, male												
2004.....	-	2	-	-	1	3	4	4	6	2	-	-
2003.....	-	-	2	1	2	7	3	2	1	1	-	-
2002.....	2	1	-	-	2	1	5	4	3	-	-	-
2001.....	3	1	-	-	1	1	4	7	2	-	-	-
2000.....	1	2	1	2	5	4	6	1	3	-	1	-
1999.....	1	-	-	-	1	3	6	4	3	-	-	-
All races, female												
2004.....	-	-	-	1	-	-	2	6	-	1	1	-
2003.....	1	-	1	-	-	2	1	6	6	3	-	-
2002.....	-	-	1	1	-	1	3	7	7	1	-	-
2001.....	-	-	-	-	1	1	2	2	5	-	-	-
2000.....	-	-	-	-	1	3	4	4	4	2	-	-
1999.....	-	-	-	-	-	-	-	3	6	1	-	-
White, both sexes												
2004.....	-	2	-	-	1	2	6	10	6	3	1	-
2003.....	1	-	2	1	2	9	3	8	7	4	-	-
2002.....	1	1	1	1	2	2	7	11	9	1	-	-
2001.....	2	-	-	-	2	2	6	9	7	-	-	-
2000.....	1	1	-	1	4	6	8	5	7	2	1	-
1999.....	1	-	-	-	1	3	6	6	9	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Kaposi's sarcoma (C46)												
White, male												
2004.....	-	2	-	-	1	2	4	4	6	2	-	-
2003.....	-	-	1	1	2	7	2	2	1	1	-	-
2002.....	1	1	-	-	2	1	4	4	3	-	-	-
2001.....	2	-	-	-	1	1	4	7	2	-	-	-
2000.....	1	1	-	1	3	3	4	1	3	-	1	-
1999.....	1	-	-	-	1	3	6	3	3	-	-	-
White, female												
2004.....	-	-	-	-	-	-	2	6	-	1	1	-
2003.....	1	-	1	-	-	2	1	6	6	3	-	-
2002.....	-	-	1	1	-	1	3	7	6	1	-	-
2001.....	-	-	-	-	1	1	2	2	5	-	-	-
2000.....	-	-	-	-	1	3	4	4	4	2	-	-
1999.....	-	-	-	-	-	-	-	3	6	1	-	-
Black, both sexes												
2004.....	-	-	-	1	-	1	-	-	-	-	-	-
2003.....	-	-	1	-	-	-	1	-	-	-	-	-
2002.....	1	-	-	-	-	-	-	-	-	-	-	-
2001.....	1	1	-	-	-	-	-	-	-	-	-	-
2000.....	-	1	1	1	2	1	2	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	1	-	-	-	-
Black, male												
2004.....	-	-	-	-	-	1	-	-	-	-	-	-
2003.....	-	-	1	-	-	-	1	-	-	-	-	-
2002.....	1	-	-	-	-	-	-	-	-	-	-	-
2001.....	1	1	-	-	-	-	-	-	-	-	-	-
2000.....	-	1	1	1	2	1	2	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	1	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Of other mesothelial and soft tissue (C47-C49)												
All races, both sexes												
2004.....	298	416	407	470	490	544	510	310	126	21	9	-
2003.....	316	362	424	428	489	564	472	272	121	20	2	-
2002.....	295	361	415	420	503	504	469	288	107	26	8	-
2001.....	327	347	374	440	505	552	450	268	104	24	3	-
2000.....	302	347	365	454	524	568	434	262	110	28	6	-
1999.....	298	296	387	413	522	543	420	266	104	26	3	-
All races, male												
2004.....	125	171	190	210	217	217	217	131	49	9	2	-
2003.....	140	151	166	181	210	245	203	104	44	5	1	-
2002.....	128	154	182	167	222	207	186	122	34	5	2	-
2001.....	160	142	163	193	239	218	193	98	33	7	1	-
2000.....	128	151	167	214	204	257	190	101	44	8	-	-
1999.....	143	150	182	192	231	234	173	97	29	9	-	-
All races, female												
2004.....	173	245	217	260	273	327	293	179	77	12	7	-
2003.....	176	211	258	247	279	319	269	168	77	15	1	-
2002.....	167	207	233	253	281	297	283	166	73	21	6	-
2001.....	167	205	211	247	266	334	257	170	71	17	2	-
2000.....	174	196	198	240	320	311	244	161	66	20	6	-
1999.....	155	146	205	221	291	309	247	169	75	17	3	-
White, both sexes												
2004.....	236	346	366	415	444	496	473	293	116	20	5	-
2003.....	261	315	358	375	441	506	443	252	114	19	2	-
2002.....	253	302	355	363	448	468	430	263	101	26	8	-
2001.....	270	292	328	397	452	494	423	240	97	22	2	-
2000.....	246	303	316	397	475	526	409	241	100	25	5	-
1999.....	253	255	330	362	465	491	386	246	97	23	3	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Of other mesothelial and soft tissue (C47-C49)												
White, male												
2004.....	101	142	178	186	198	199	206	124	43	8	2	-
2003.....	117	133	139	158	192	224	196	94	42	5	1	-
2002.....	113	131	157	148	201	196	175	112	32	5	2	-
2001.....	138	120	145	175	220	195	182	87	32	6	-	-
2000.....	116	137	144	189	188	237	180	92	40	6	-	-
1999.....	126	133	158	166	204	214	162	94	28	8	-	-
White, female												
2004.....	135	204	188	229	246	297	267	169	73	12	3	-
2003.....	144	182	219	217	249	282	247	158	72	14	1	-
2002.....	140	171	198	215	247	272	255	151	69	21	6	-
2001.....	132	172	183	222	232	299	241	153	65	16	2	-
2000.....	130	166	172	208	287	289	229	149	60	19	5	-
1999.....	127	122	172	196	261	277	224	152	69	15	3	-
Black, both sexes												
2004.....	54	58	26	48	34	40	34	11	9	1	4	-
2003.....	45	36	52	44	42	48	22	16	7	1	-	-
2002.....	36	47	47	50	41	33	34	16	4	-	-	-
2001.....	50	49	33	35	45	42	22	24	6	1	1	-
2000.....	44	39	41	47	44	34	19	17	9	2	1	-
1999.....	34	31	51	36	49	45	26	19	5	2	-	-
Black, male												
2004.....	21	24	7	21	14	15	9	4	5	1	-	-
2003.....	19	17	22	16	13	17	5	8	2	-	-	-
2002.....	13	19	19	16	17	10	10	6	1	-	-	-
2001.....	21	22	11	15	15	16	9	9	1	-	1	-
2000.....	7	14	20	18	14	16	6	6	4	1	-	-
1999.....	13	15	21	17	21	17	8	3	1	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Of other mesothelial and soft tissue (C47-C49)												
Black, female												
2004.....	33	34	19	27	20	25	25	7	4	-	4	-
2003.....	26	19	30	28	29	31	17	8	5	1	-	-
2002.....	23	28	28	34	24	23	24	10	3	-	-	-
2001.....	29	27	22	20	30	26	13	15	5	1	-	-
2000.....	37	25	21	29	30	18	13	11	5	1	1	-
1999.....	21	16	30	19	28	28	18	16	4	1	-	-
American Indian 1/, both sexes												
2004.....	2	3	1	1	1	-	-	1	-	-	-	-
2003.....	2	3	6	1	1	2	1	2	-	-	-	-
2002.....	-	1	4	-	1	-	1	1	1	-	-	-
2001.....	2	-	2	2	-	4	-	-	1	-	-	-
2000.....	-	-	1	1	-	2	-	-	-	-	-	-
1999.....	2	3	-	2	-	1	3	-	-	-	-	-
American Indian 1/, male												
2004.....	-	1	-	-	1	-	-	1	-	-	-	-
2003.....	1	1	1	1	1	1	-	1	-	-	-	-
2002.....	-	1	2	-	1	-	1	-	-	-	-	-
2001.....	-	-	2	1	-	1	-	-	-	-	-	-
2000.....	-	-	-	-	-	2	-	-	-	-	-	-
1999.....	1	1	-	-	-	-	1	-	-	-	-	-
American Indian 1/, female												
2004.....	2	2	1	1	-	-	-	-	-	-	-	-
2003.....	1	2	5	-	-	1	1	1	-	-	-	-
2002.....	-	-	2	-	-	-	-	1	1	-	-	-
2001.....	2	-	-	1	-	3	-	-	1	-	-	-
2000.....	-	-	1	1	-	-	-	-	-	-	-	-
1999.....	1	2	-	2	-	1	2	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Of other mesothelial and soft tissue (C47-C49)												
Asian or Pacific Islander 2/, both sexes												
2004.....	6	9	14	6	11	8	3	5	1	-	-	-
2003.....	8	8	8	8	5	8	6	2	-	-	-	-
2002.....	6	11	9	7	13	3	4	8	1	-	-	-
2001.....	5	6	11	6	8	12	5	4	-	1	-	-
2000.....	12	5	7	9	5	6	6	4	1	1	-	-
1999.....	9	7	6	13	8	6	5	1	2	1	-	-
Asian or Pacific Islander 2/, male												
2004.....	3	4	5	3	4	3	2	2	1	-	-	-
2003.....	3	-	4	6	4	3	2	1	-	-	-	-
2002.....	2	3	4	3	3	1	-	4	1	-	-	-
2001.....	1	-	5	2	4	6	2	2	-	1	-	-
2000.....	5	-	3	7	2	2	4	3	-	1	-	-
1999.....	3	1	3	9	6	3	2	-	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	3	5	9	3	7	5	1	3	-	-	-	-
2003.....	5	8	4	2	1	5	4	1	-	-	-	-
2002.....	4	8	5	4	10	2	4	4	-	-	-	-
2001.....	4	6	6	4	4	6	3	2	-	-	-	-
2000.....	7	5	4	2	3	4	2	1	1	-	-	-
1999.....	6	6	3	4	2	3	3	1	2	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant neoplasm of breast (C50)												
All races, both sexes												
2004.....	3,639	4,312	4,162	4,065	4,220	4,792	4,579	3,473	1,823	618	122	-
2003.....	3,631	4,206	4,061	3,887	4,451	4,916	4,728	3,530	1,965	630	120	2
2002.....	3,707	3,924	3,914	3,954	4,551	4,978	4,770	3,528	1,869	658	126	-
2001.....	3,803	3,843	3,789	3,972	4,743	5,130	4,533	3,330	1,848	644	89	-
2000.....	3,655	3,790	3,783	4,135	4,889	5,256	4,494	3,596	1,880	658	115	-
1999.....	3,555	3,682	3,717	4,147	4,962	5,140	4,511	3,428	1,837	611	124	-
All races, male												
2004.....	20	30	37	43	48	65	44	33	10	6	1	-
2003.....	34	30	41	42	47	46	58	35	17	4	-	1
2002.....	28	29	38	35	49	57	55	31	13	3	-	-
2001.....	33	32	40	54	66	69	46	31	9	6	-	-
2000.....	28	44	41	47	68	65	44	36	13	1	1	-
1999.....	30	36	34	47	56	56	42	33	13	7	1	-
All races, female												
2004.....	3,619	4,282	4,125	4,022	4,172	4,727	4,535	3,440	1,813	612	121	-
2003.....	3,597	4,176	4,020	3,845	4,404	4,870	4,670	3,495	1,948	626	120	1
2002.....	3,679	3,895	3,876	3,919	4,502	4,921	4,715	3,497	1,856	655	126	-
2001.....	3,770	3,811	3,749	3,918	4,677	5,061	4,487	3,299	1,839	638	89	-
2000.....	3,627	3,746	3,742	4,088	4,821	5,191	4,450	3,560	1,867	657	114	-
1999.....	3,525	3,646	3,683	4,100	4,906	5,084	4,469	3,395	1,824	604	123	-
White, both sexes												
2004.....	2,780	3,514	3,452	3,474	3,666	4,271	4,165	3,123	1,669	566	108	-
2003.....	2,823	3,406	3,380	3,279	3,879	4,374	4,264	3,196	1,779	568	96	1
2002.....	2,904	3,206	3,288	3,365	3,968	4,465	4,268	3,218	1,702	590	108	-
2001.....	3,008	3,137	3,170	3,362	4,150	4,604	4,103	3,021	1,699	585	69	-
2000.....	2,934	3,100	3,182	3,584	4,341	4,735	4,107	3,267	1,696	586	102	-
1999.....	2,849	3,006	3,105	3,550	4,372	4,598	4,116	3,154	1,681	555	108	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant neoplasm of breast (C50)												
White, male												
2004.....	14	24	29	37	39	61	39	26	9	5	1	-
2003.....	30	24	35	36	40	42	53	31	16	4	-	1
2002.....	21	23	28	24	44	47	49	29	10	3	-	-
2001.....	24	27	34	46	55	59	41	26	9	4	-	-
2000.....	24	34	33	43	60	60	38	29	12	1	1	-
1999.....	22	29	25	40	45	49	37	30	11	5	1	-
White, female												
2004.....	2,766	3,490	3,423	3,437	3,627	4,210	4,126	3,097	1,660	561	107	-
2003.....	2,793	3,382	3,345	3,243	3,839	4,332	4,211	3,165	1,763	564	96	-
2002.....	2,883	3,183	3,260	3,341	3,924	4,418	4,219	3,189	1,692	587	108	-
2001.....	2,984	3,110	3,136	3,316	4,095	4,545	4,062	2,995	1,690	581	69	-
2000.....	2,910	3,066	3,149	3,541	4,281	4,675	4,069	3,238	1,684	585	101	-
1999.....	2,827	2,977	3,080	3,510	4,327	4,549	4,079	3,124	1,670	550	107	-
Black, both sexes												
2004.....	708	671	597	496	480	455	364	308	141	46	14	-
2003.....	675	685	569	521	492	472	419	303	169	59	23	-
2002.....	673	604	533	513	501	457	449	286	148	64	16	-
2001.....	665	602	529	537	523	481	391	275	142	59	19	-
2000.....	614	592	518	477	492	474	356	307	173	71	13	-
1999.....	614	582	521	523	517	493	363	252	140	54	16	-
Black, male												
2004.....	6	4	8	6	8	4	5	7	1	1	-	-
2003.....	4	6	6	6	7	3	4	4	1	-	-	-
2002.....	6	6	9	11	5	8	5	2	3	-	-	-
2001.....	9	4	6	7	10	9	5	4	-	2	-	-
2000.....	4	8	8	3	8	5	5	7	1	-	-	-
1999.....	8	7	9	6	10	6	5	3	2	2	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant neoplasm of breast (C50)												
Black, female												
2004.....	702	667	589	490	472	451	359	301	140	45	14	-
2003.....	671	679	563	515	485	469	415	299	168	59	23	-
2002.....	667	598	524	502	496	449	444	284	145	64	16	-
2001.....	656	598	523	530	513	472	386	271	142	57	19	-
2000.....	610	584	510	474	484	469	351	300	172	71	13	-
1999.....	606	575	512	517	507	487	358	249	138	52	16	-
American Indian 1/, both sexes												
2004.....	25	17	22	17	11	8	21	8	5	-	-	-
2003.....	20	17	19	9	15	18	15	10	4	-	-	-
2002.....	22	17	14	18	11	9	9	4	5	2	1	-
2001.....	20	14	18	12	13	3	5	9	1	-	-	-
2000.....	14	20	19	14	9	11	9	2	4	-	-	-
1999.....	17	14	14	18	18	11	6	9	2	-	-	-
American Indian 1/, male												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	1	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	1	-	-	-	-	-	-	-
2000.....	-	1	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, female												
2004.....	25	17	22	17	11	8	21	8	5	-	-	-
2003.....	20	17	19	9	15	18	14	10	4	-	-	-
2002.....	22	17	14	18	11	9	9	4	5	2	1	-
2001.....	20	14	18	12	12	3	5	9	1	-	-	-
2000.....	14	19	19	14	9	11	9	2	4	-	-	-
1999.....	17	14	14	17	18	11	6	9	2	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant neoplasm of breast (C50)												
Asian or Pacific Islander 2/, both sexes												
2004.....	126	110	91	78	63	58	29	34	8	6	-	-
2003.....	113	98	93	78	65	52	30	21	13	3	1	1
2002.....	108	97	79	58	71	47	44	20	14	2	1	-
2001.....	110	90	72	61	57	42	34	25	6	-	1	-
2000.....	93	78	64	60	47	36	22	20	7	1	-	-
1999.....	75	80	77	56	55	38	26	13	14	2	-	-
Asian or Pacific Islander 2/, male												
2004.....	-	2	-	-	1	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	1	-	-	-	-	-	-
2002.....	1	-	1	-	-	2	1	-	-	-	-	-
2001.....	-	1	-	1	-	1	-	1	-	-	-	-
2000.....	-	1	-	1	-	-	1	-	-	-	-	-
1999.....	-	-	-	-	1	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	126	108	91	78	62	58	29	34	8	6	-	-
2003.....	113	98	93	78	65	51	30	21	13	3	1	1
2002.....	107	97	78	58	71	45	43	20	14	2	1	-
2001.....	110	89	72	60	57	41	34	24	6	-	1	-
2000.....	93	77	64	59	47	36	21	20	7	1	-	-
1999.....	75	80	77	56	54	37	26	13	14	2	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant neoplasms of female genital organs (C51-C58)												
All races, both sexes												
2004.....	1,824	2,399	2,701	3,078	3,351	3,614	3,495	2,209	1,058	239	27	-
2003.....	1,884	2,401	2,771	3,008	3,281	3,716	3,456	2,233	1,029	243	41	-
2002.....	1,941	2,261	2,613	2,902	3,467	3,861	3,382	2,231	1,002	283	45	-
2001.....	1,948	2,083	2,533	2,870	3,476	3,871	3,446	2,237	978	274	31	1
2000.....	1,824	2,093	2,402	2,953	3,448	3,897	3,233	2,187	1,002	265	36	1
1999.....	1,751	1,979	2,423	2,939	3,502	3,795	3,155	2,081	917	254	32	1
All races, male												
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....
All races, female												
2004.....	1,824	2,399	2,701	3,078	3,351	3,614	3,495	2,209	1,058	239	27	-
2003.....	1,884	2,401	2,771	3,008	3,281	3,716	3,456	2,233	1,029	243	41	-
2002.....	1,941	2,261	2,613	2,902	3,467	3,861	3,382	2,231	1,002	283	45	-
2001.....	1,948	2,083	2,533	2,870	3,476	3,871	3,446	2,237	978	274	31	1
2000.....	1,824	2,093	2,402	2,953	3,448	3,897	3,233	2,187	1,002	265	36	1
1999.....	1,751	1,979	2,423	2,939	3,502	3,795	3,155	2,081	917	254	32	1
White, both sexes												
2004.....	1,467	1,997	2,273	2,542	2,824	3,191	3,172	1,997	966	215	21	-
2003.....	1,542	2,049	2,310	2,515	2,806	3,258	3,100	2,018	930	222	33	-
2002.....	1,615	1,904	2,195	2,410	2,973	3,395	3,034	2,006	905	244	37	-
2001.....	1,614	1,762	2,088	2,384	3,014	3,424	3,111	2,027	883	244	21	-
2000.....	1,515	1,780	2,011	2,465	2,951	3,462	2,912	1,999	893	229	30	-
1999.....	1,478	1,667	2,044	2,460	3,017	3,343	2,860	1,895	822	224	28	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant neoplasms of female genital organs (C51-C58)												
Black, female												
2004.....	277	308	357	454	451	369	278	182	79	20	6	-
2003.....	265	283	380	417	400	383	301	186	91	21	8	-
2002.....	254	281	338	414	427	389	296	202	85	37	7	-
2001.....	249	254	355	413	409	386	288	184	88	27	10	1
2000.....	232	250	331	425	434	388	282	169	98	36	6	1
1999.....	210	263	336	416	450	396	256	167	90	26	4	-
American Indian 1/, both sexes												
2004.....	13	15	13	8	15	6	7	5	5	2	-	-
2003.....	12	10	14	8	11	13	9	5	2	-	-	-
2002.....	13	9	10	8	15	9	14	5	3	-	-	-
2001.....	14	13	15	17	11	13	10	7	2	3	-	-
2000.....	7	5	10	8	8	12	5	3	2	-	-	-
1999.....	4	8	3	12	5	8	6	3	-	1	-	-
American Indian 1/, male												
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....
American Indian 1/, female												
2004.....	13	15	13	8	15	6	7	5	5	2	-	-
2003.....	12	10	14	8	11	13	9	5	2	-	-	-
2002.....	13	9	10	8	15	9	14	5	3	-	-	-
2001.....	14	13	15	17	11	13	10	7	2	3	-	-
2000.....	7	5	10	8	8	12	5	3	2	-	-	-
1999.....	4	8	3	12	5	8	6	3	-	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant neoplasms of female genital organs (C51-C58)												
Asian or Pacific Islander 2/, both sexes												
2004.....	67	79	58	74	61	48	38	25	8	2	-	-
2003.....	65	59	67	68	64	62	46	24	6	-	-	-
2002.....	59	67	70	70	52	68	38	18	9	2	1	-
2001.....	71	54	75	56	42	48	37	19	5	-	-	-
2000.....	70	58	50	55	55	35	34	16	9	-	-	-
1999.....	59	41	40	51	30	48	33	16	5	3	-	-
Asian or Pacific Islander 2/, male												
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....
Asian or Pacific Islander 2/, female												
2004.....	67	79	58	74	61	48	38	25	8	2	-	-
2003.....	65	59	67	68	64	62	46	24	6	-	-	-
2002.....	59	67	70	70	52	68	38	18	9	2	1	-
2001.....	71	54	75	56	42	48	37	19	5	-	-	-
2000.....	70	58	50	55	55	35	34	16	9	-	-	-
1999.....	59	41	40	51	30	48	33	16	5	3	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Of vulva and vagina (C51-C52)												
All races, both sexes												
2004.....	48	64	86	88	136	158	216	177	123	32	5	-
2003.....	47	55	72	93	119	176	193	161	131	39	8	-
2002.....	50	51	75	67	129	185	187	180	128	31	13	-
2001.....	53	46	58	84	126	185	202	173	105	38	4	-
2000.....	50	44	62	99	137	187	178	176	103	41	7	-
1999.....	48	45	66	82	132	184	189	214	105	35	2	-
All races, male												
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....
All races, female												
2004.....	48	64	86	88	136	158	216	177	123	32	5	-
2003.....	47	55	72	93	119	176	193	161	131	39	8	-
2002.....	50	51	75	67	129	185	187	180	128	31	13	-
2001.....	53	46	58	84	126	185	202	173	105	38	4	-
2000.....	50	44	62	99	137	187	178	176	103	41	7	-
1999.....	48	45	66	82	132	184	189	214	105	35	2	-
White, both sexes												
2004.....	40	48	74	78	120	148	198	165	113	31	4	-
2003.....	33	46	62	84	108	164	179	156	120	34	8	-
2002.....	44	45	66	61	117	170	171	169	121	30	11	-
2001.....	43	36	42	73	110	164	189	163	93	35	2	-
2000.....	43	36	56	90	126	169	167	164	95	37	7	-
1999.....	40	33	55	72	117	163	175	203	101	33	2	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Of vulva and vagina (C51-C52)												
Black, female												
2004.....	7	14	10	9	14	7	16	9	8	1	1	-
2003.....	12	9	8	9	10	8	13	5	11	5	-	-
2002.....	6	6	7	5	10	12	11	10	7	1	1	-
2001.....	9	7	15	10	15	20	12	8	9	2	2	-
2000.....	5	7	6	9	9	14	10	11	8	4	-	-
1999.....	7	8	9	9	13	15	13	11	3	2	-	-
American Indian 1/, both sexes												
2004.....	-	-	1	1	-	-	1	1	2	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	1	-	1	-	2	1	-	-	-	-
2001.....	1	-	-	-	-	1	1	1	1	1	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	1	-	1	-	2	-	-	-	-	-	-
American Indian 1/, male												
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....
American Indian 1/, female												
2004.....	-	-	1	1	-	-	1	1	2	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	1	-	1	-	2	1	-	-	-	-
2001.....	1	-	-	-	-	1	1	1	1	1	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	1	-	1	-	2	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Of vulva and vagina (C51-C52)												
Asian or Pacific Islander 2/, both sexes												
2004.....	1	2	1	-	2	3	1	2	-	-	-	-
2003.....	2	-	2	-	1	4	1	-	-	-	-	-
2002.....	-	-	1	1	1	3	3	-	-	-	1	-
2001.....	-	3	1	1	1	-	-	1	2	-	-	-
2000.....	2	1	-	-	2	4	1	1	-	-	-	-
1999.....	1	3	2	-	2	4	1	-	1	-	-	-
Asian or Pacific Islander 2/, male												
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....
Asian or Pacific Islander 2/, female												
2004.....	1	2	1	-	2	3	1	2	-	-	-	-
2003.....	2	-	2	-	1	4	1	-	-	-	-	-
2002.....	-	-	1	1	1	3	3	-	-	-	1	-
2001.....	-	3	1	1	1	-	-	1	2	-	-	-
2000.....	2	1	-	-	2	4	1	1	-	-	-	-
1999.....	1	3	2	-	2	4	1	-	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Of cervix uteri (C53)												
All races, both sexes												
2004.....	395	444	371	318	283	243	241	151	69	23	3	-
2003.....	450	445	348	308	256	274	278	166	78	22	4	-
2002.....	455	401	324	315	315	305	223	155	99	24	6	-
2001.....	478	371	352	279	306	298	285	193	90	30	2	-
2000.....	467	416	332	342	344	320	279	180	92	35	3	-
1999.....	452	393	376	350	346	345	223	199	98	26	6	1
All races, male												
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....
All races, female												
2004.....	395	444	371	318	283	243	241	151	69	23	3	-
2003.....	450	445	348	308	256	274	278	166	78	22	4	-
2002.....	455	401	324	315	315	305	223	155	99	24	6	-
2001.....	478	371	352	279	306	298	285	193	90	30	2	-
2000.....	467	416	332	342	344	320	279	180	92	35	3	-
1999.....	452	393	376	350	346	345	223	199	98	26	6	1
White, both sexes												
2004.....	291	349	284	232	206	188	188	113	49	15	1	-
2003.....	341	348	265	231	183	188	223	132	54	20	4	-
2002.....	343	319	234	226	229	237	170	116	74	15	5	-
2001.....	363	283	272	195	235	227	219	150	69	26	2	-
2000.....	352	314	243	243	255	253	225	147	67	25	2	-
1999.....	360	298	275	261	242	265	181	161	76	14	4	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Of cervix uteri (C53)												
Black, female												
2004.....	87	79	75	69	62	47	41	31	17	7	2	-
2003.....	83	81	72	68	58	66	45	29	19	2	-	-
2002.....	96	62	69	77	75	56	48	30	19	9	1	-
2001.....	89	65	66	73	57	56	56	39	19	4	-	-
2000.....	92	83	70	85	73	61	48	30	22	10	1	-
1999.....	72	80	89	79	97	69	33	35	22	9	2	-
American Indian 1/, both sexes												
2004.....	4	2	2	-	3	-	1	2	-	-	-	-
2003.....	3	6	2	1	2	2	2	-	2	-	-	-
2002.....	3	2	2	2	3	1	2	1	2	-	-	-
2001.....	3	5	1	1	1	1	1	1	1	-	-	-
2000.....	6	2	3	1	2	2	-	1	1	-	-	-
1999.....	2	3	2	1	-	-	2	-	-	-	-	-
American Indian 1/, male												
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....
American Indian 1/, female												
2004.....	4	2	2	-	3	-	1	2	-	-	-	-
2003.....	3	6	2	1	2	2	2	-	2	-	-	-
2002.....	3	2	2	2	3	1	2	1	2	-	-	-
2001.....	3	5	1	1	1	1	1	1	1	-	-	-
2000.....	6	2	3	1	2	2	-	1	1	-	-	-
1999.....	2	3	2	1	-	-	2	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Of cervix uteri (C53)												
Asian or Pacific Islander 2/, both sexes												
2004.....	13	14	10	17	12	8	11	5	3	1	-	-
2003.....	23	10	9	8	13	18	8	5	3	-	-	-
2002.....	13	18	19	10	8	11	3	8	4	-	-	-
2001.....	23	18	13	10	13	14	9	3	1	-	-	-
2000.....	17	17	16	13	14	4	6	2	2	-	-	-
1999.....	18	12	10	9	7	11	7	3	-	3	-	-
Asian or Pacific Islander 2/, male												
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....
Asian or Pacific Islander 2/, female												
2004.....	13	14	10	17	12	8	11	5	3	1	-	-
2003.....	23	10	9	8	13	18	8	5	3	-	-	-
2002.....	13	18	19	10	8	11	3	8	4	-	-	-
2001.....	23	18	13	10	13	14	9	3	1	-	-	-
2000.....	17	17	16	13	14	4	6	2	2	-	-	-
1999.....	18	12	10	9	7	11	7	3	-	3	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Of corpus uteri and uterus, part unspecified (C54-C55)

All races, both sexes

2004.....	351	566	724	876	980	1,025	986	651	345	79	8	-
2003.....	363	557	744	878	973	1,025	926	668	316	76	17	-
2002.....	376	531	668	809	1,026	1,051	906	694	314	109	11	-
2001.....	374	457	648	785	999	1,075	988	662	312	104	10	1
2000.....	317	478	644	800	981	1,039	919	617	333	84	15	1
1999.....	286	427	592	835	972	1,061	947	593	290	100	16	-

All races, male

2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

All races, female

2004.....	351	566	724	876	980	1,025	986	651	345	79	8	-
2003.....	363	557	744	878	973	1,025	926	668	316	76	17	-
2002.....	376	531	668	809	1,026	1,051	906	694	314	109	11	-
2001.....	374	457	648	785	999	1,075	988	662	312	104	10	1
2000.....	317	478	644	800	981	1,039	919	617	333	84	15	1
1999.....	286	427	592	835	972	1,061	947	593	290	100	16	-

White, both sexes

2004.....	265	453	558	655	756	856	869	585	314	76	8	-
2003.....	295	455	561	663	789	860	790	597	287	72	12	-
2002.....	296	412	522	595	824	865	783	609	281	91	9	-
2001.....	296	380	492	595	809	911	865	601	275	92	7	-
2000.....	262	394	496	602	771	866	785	557	294	74	11	-
1999.....	236	342	474	630	805	904	835	523	253	91	14	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Of corpus uteri and uterus, part unspecified (C54-C55)												
Black, female												
2004.....	67	87	145	196	202	157	103	62	28	3	-	-
2003.....	55	83	165	190	169	150	123	66	28	4	5	-
2002.....	61	99	124	188	188	162	111	81	31	17	2	-
2001.....	59	65	130	174	180	153	115	55	36	11	3	1
2000.....	44	71	131	177	198	156	124	56	36	10	4	1
1999.....	39	76	113	189	161	144	103	65	35	9	2	-
American Indian 1/, both sexes												
2004.....	3	2	2	3	4	-	4	-	1	-	-	-
2003.....	1	-	1	1	3	-	2	-	-	-	-	-
2002.....	3	3	3	3	1	7	2	2	-	-	-	-
2001.....	3	-	6	2	4	3	2	2	-	1	-	-
2000.....	-	-	2	5	3	4	2	-	1	-	-	-
1999.....	-	2	-	2	2	1	-	2	-	-	-	-
American Indian 1/, male												
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....
American Indian 1/, female												
2004.....	3	2	2	3	4	-	4	-	1	-	-	-
2003.....	1	-	1	1	3	-	2	-	-	-	-	-
2002.....	3	3	3	3	1	7	2	2	-	-	-	-
2001.....	3	-	6	2	4	3	2	2	-	1	-	-
2000.....	-	-	2	5	3	4	2	-	1	-	-	-
1999.....	-	2	-	2	2	1	-	2	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Of corpus uteri and uterus, part unspecified (C54-C55)												
Asian or Pacific Islander 2/, both sexes												
2004.....	16	24	19	22	18	12	10	4	2	-	-	-
2003.....	12	19	17	24	12	15	11	5	1	-	-	-
2002.....	16	17	19	23	13	17	10	2	2	1	-	-
2001.....	16	12	20	14	6	8	6	4	1	-	-	-
2000.....	11	13	15	16	9	13	8	4	2	-	-	-
1999.....	11	7	5	14	4	12	9	3	2	-	-	-
Asian or Pacific Islander 2/, male												
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....
Asian or Pacific Islander 2/, female												
2004.....	16	24	19	22	18	12	10	4	2	-	-	-
2003.....	12	19	17	24	12	15	11	5	1	-	-	-
2002.....	16	17	19	23	13	17	10	2	2	1	-	-
2001.....	16	12	20	14	6	8	6	4	1	-	-	-
2000.....	11	13	15	16	9	13	8	4	2	-	-	-
1999.....	11	7	5	14	4	12	9	3	2	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Of ovary (C56)												
All races, both sexes												
2004.....	1,007	1,301	1,493	1,775	1,918	2,142	2,023	1,208	510	103	10	-
2003.....	997	1,315	1,576	1,687	1,882	2,193	2,001	1,207	490	102	11	-
2002.....	1,029	1,249	1,499	1,662	1,939	2,249	2,010	1,174	447	115	14	-
2001.....	1,016	1,171	1,441	1,682	1,984	2,246	1,924	1,181	451	96	15	-
2000.....	966	1,123	1,321	1,669	1,932	2,295	1,803	1,176	453	98	9	-
1999.....	933	1,081	1,345	1,626	1,986	2,143	1,732	1,052	412	91	7	-
All races, male												
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....
All races, female												
2004.....	1,007	1,301	1,493	1,775	1,918	2,142	2,023	1,208	510	103	10	-
2003.....	997	1,315	1,576	1,687	1,882	2,193	2,001	1,207	490	102	11	-
2002.....	1,029	1,249	1,499	1,662	1,939	2,249	2,010	1,174	447	115	14	-
2001.....	1,016	1,171	1,441	1,682	1,984	2,246	1,924	1,181	451	96	15	-
2000.....	966	1,123	1,321	1,669	1,932	2,295	1,803	1,176	453	98	9	-
1999.....	933	1,081	1,345	1,626	1,986	2,143	1,732	1,052	412	91	7	-
White, both sexes												
2004.....	854	1,126	1,334	1,560	1,710	1,957	1,889	1,114	479	91	7	-
2003.....	851	1,173	1,398	1,499	1,682	2,004	1,854	1,105	457	93	8	-
2002.....	906	1,102	1,339	1,484	1,751	2,058	1,859	1,087	416	105	11	-
2001.....	890	1,030	1,254	1,486	1,803	2,062	1,794	1,088	426	85	10	-
2000.....	838	1,008	1,179	1,489	1,752	2,130	1,684	1,099	418	86	8	-
1999.....	814	965	1,199	1,455	1,799	1,956	1,614	987	381	84	7	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Of ovary (C56)												
Black, female												
2004.....	110	126	124	177	172	155	117	78	26	9	3	-
2003.....	110	108	130	146	156	154	116	83	31	9	3	-
2002.....	87	114	126	140	149	154	124	78	27	9	3	-
2001.....	88	112	141	151	154	151	103	80	24	10	5	-
2000.....	89	86	119	152	147	147	97	66	31	12	1	-
1999.....	89	95	122	136	168	162	98	54	30	6	-	-
American Indian 1/, both sexes												
2004.....	6	11	7	4	8	6	1	2	2	2	-	-
2003.....	8	4	11	6	6	11	5	5	-	-	-	-
2002.....	7	4	4	3	9	1	8	1	1	-	-	-
2001.....	6	8	8	14	6	8	6	2	-	1	-	-
2000.....	1	3	4	2	3	6	3	2	-	-	-	-
1999.....	2	2	1	8	3	5	4	1	-	1	-	-
American Indian 1/, male												
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....
American Indian 1/, female												
2004.....	6	11	7	4	8	6	1	2	2	2	-	-
2003.....	8	4	11	6	6	11	5	5	-	-	-	-
2002.....	7	4	4	3	9	1	8	1	1	-	-	-
2001.....	6	8	8	14	6	8	6	2	-	1	-	-
2000.....	1	3	4	2	3	6	3	2	-	-	-	-
1999.....	2	2	1	8	3	5	4	1	-	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Of ovary (C56)												
Asian or Pacific Islander 2/, both sexes												
2004.....	37	38	28	34	28	24	16	14	3	1	-	-
2003.....	28	30	37	36	38	24	26	14	2	-	-	-
2002.....	29	29	30	35	30	36	19	8	3	1	-	-
2001.....	32	21	38	31	21	25	21	11	1	-	-	-
2000.....	38	26	19	26	30	12	19	9	4	-	-	-
1999.....	28	19	23	27	16	20	16	10	1	-	-	-
Asian or Pacific Islander 2/, male												
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....
Asian or Pacific Islander 2/, female												
2004.....	37	38	28	34	28	24	16	14	3	1	-	-
2003.....	28	30	37	36	38	24	26	14	2	-	-	-
2002.....	29	29	30	35	30	36	19	8	3	1	-	-
2001.....	32	21	38	31	21	25	21	11	1	-	-	-
2000.....	38	26	19	26	30	12	19	9	4	-	-	-
1999.....	28	19	23	27	16	20	16	10	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Of other and unspecified female genital organs (C57)												
All races, both sexes												
2004.....	22	22	25	20	34	45	28	22	11	2	1	-
2003.....	26	29	31	42	51	48	58	31	14	4	1	-
2002.....	30	29	46	49	56	71	56	28	14	4	1	-
2001.....	25	38	34	39	61	67	46	28	20	6	-	-
2000.....	22	32	43	43	54	55	54	37	21	7	2	-
1999.....	32	33	44	46	66	62	64	23	12	2	1	-
All races, male												
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....
All races, female												
2004.....	22	22	25	20	34	45	28	22	11	2	1	-
2003.....	26	29	31	42	51	48	58	31	14	4	1	-
2002.....	30	29	46	49	56	71	56	28	14	4	1	-
2001.....	25	38	34	39	61	67	46	28	20	6	-	-
2000.....	22	32	43	43	54	55	54	37	21	7	2	-
1999.....	32	33	44	46	66	62	64	23	12	2	1	-
White, both sexes												
2004.....	17	20	21	16	32	41	27	20	11	2	1	-
2003.....	21	27	24	38	44	42	54	28	12	3	1	-
2002.....	25	26	33	44	50	65	51	25	13	3	1	-
2001.....	21	33	28	34	57	60	44	25	20	6	-	-
2000.....	19	28	37	41	47	43	51	32	19	7	2	-
1999.....	28	29	41	42	54	55	55	21	11	2	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Of other and unspecified female genital organs (C57)

Asian or Pacific Islander 2/, both sexes

2004.....	-	1	-	1	1	1	-	-	-	-	-	-
2003.....	-	-	2	-	-	1	-	-	-	-	-	-
2002.....	1	3	1	1	-	1	3	-	-	-	-	-
2001.....	-	-	3	-	1	1	1	-	-	-	-	-
2000.....	1	1	-	-	-	2	-	-	1	-	-	-
1999.....	1	-	-	1	1	1	-	-	1	-	-	-

Asian or Pacific Islander 2/, male

2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

Asian or Pacific Islander 2/, female

2004.....	-	1	-	1	1	1	-	-	-	-	-	-
2003.....	-	-	2	-	-	1	-	-	-	-	-	-
2002.....	1	3	1	1	-	1	3	-	-	-	-	-
2001.....	-	-	3	-	1	1	1	-	-	-	-	-
2000.....	1	1	-	-	-	2	-	-	1	-	-	-
1999.....	1	-	-	1	1	1	-	-	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant neoplasms of male genital organs (C60-C63)												
All races, both sexes												
2004.....	381	727	1,392	2,233	3,566	5,490	6,589	5,351	2,641	753	87	2
2003.....	367	699	1,452	2,329	3,782	5,723	6,655	5,382	2,642	657	83	-
2002.....	342	734	1,367	2,358	4,032	5,957	6,877	5,496	2,724	704	72	-
2001.....	355	680	1,291	2,416	4,208	6,161	6,891	5,525	2,586	693	85	-
2000.....	299	651	1,345	2,505	4,376	6,354	6,812	5,555	2,649	630	75	-
1999.....	304	668	1,343	2,753	4,644	6,570	6,949	5,449	2,482	647	91	1
All races, male												
2004.....	381	727	1,392	2,233	3,566	5,490	6,589	5,351	2,641	753	87	2
2003.....	367	699	1,452	2,329	3,782	5,723	6,655	5,382	2,642	657	83	-
2002.....	342	734	1,367	2,358	4,032	5,957	6,877	5,496	2,724	704	72	-
2001.....	355	680	1,291	2,416	4,208	6,161	6,891	5,525	2,586	693	85	-
2000.....	299	651	1,345	2,505	4,376	6,354	6,812	5,555	2,649	630	75	-
1999.....	304	668	1,343	2,753	4,644	6,570	6,949	5,449	2,482	647	91	1
All races, female												
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....
White, both sexes												
2004.....	269	540	1,046	1,704	2,795	4,447	5,557	4,638	2,250	633	62	1
2003.....	263	497	1,072	1,753	3,009	4,660	5,595	4,696	2,291	546	59	-
2002.....	240	537	994	1,784	3,153	4,839	5,801	4,802	2,344	589	53	-
2001.....	249	502	947	1,849	3,306	5,030	5,766	4,754	2,204	561	67	-
2000.....	225	472	996	1,886	3,413	5,199	5,687	4,807	2,266	525	56	-
1999.....	222	501	958	2,103	3,684	5,381	5,936	4,662	2,135	525	72	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant neoplasms of male genital organs (C60-C63)												
White, male												
2004.....	269	540	1,046	1,704	2,795	4,447	5,557	4,638	2,250	633	62	1
2003.....	263	497	1,072	1,753	3,009	4,660	5,595	4,696	2,291	546	59	-
2002.....	240	537	994	1,784	3,153	4,839	5,801	4,802	2,344	589	53	-
2001.....	249	502	947	1,849	3,306	5,030	5,766	4,754	2,204	561	67	-
2000.....	225	472	996	1,886	3,413	5,199	5,687	4,807	2,266	525	56	-
1999.....	222	501	958	2,103	3,684	5,381	5,936	4,662	2,135	525	72	1
White, female												
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....
Black, both sexes												
2004.....	107	178	325	499	713	958	923	630	345	104	23	1
2003.....	98	192	359	539	729	979	959	613	308	101	20	-
2002.....	93	178	343	548	830	1,032	1,004	641	345	103	18	-
2001.....	102	171	324	538	857	1,046	1,039	704	344	118	17	-
2000.....	70	174	325	589	905	1,076	1,042	679	350	97	18	-
1999.....	78	159	366	624	917	1,105	947	719	312	103	19	-
Black, male												
2004.....	107	178	325	499	713	958	923	630	345	104	23	1
2003.....	98	192	359	539	729	979	959	613	308	101	20	-
2002.....	93	178	343	548	830	1,032	1,004	641	345	103	18	-
2001.....	102	171	324	538	857	1,046	1,039	704	344	118	17	-
2000.....	70	174	325	589	905	1,076	1,042	679	350	97	18	-
1999.....	78	159	366	624	917	1,105	947	719	312	103	19	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Of prostate (C61)												
All races, both sexes												
2004.....	332	688	1,356	2,196	3,515	5,444	6,549	5,328	2,630	747	87	2
2003.....	319	669	1,405	2,288	3,745	5,671	6,613	5,358	2,624	653	82	-
2002.....	302	696	1,324	2,316	3,984	5,915	6,838	5,474	2,708	703	72	-
2001.....	313	647	1,260	2,379	4,164	6,119	6,857	5,500	2,575	689	85	-
2000.....	263	623	1,306	2,466	4,342	6,314	6,757	5,532	2,638	628	75	-
1999.....	274	636	1,310	2,711	4,605	6,532	6,903	5,418	2,467	644	91	1
All races, male												
2004.....	332	688	1,356	2,196	3,515	5,444	6,549	5,328	2,630	747	87	2
2003.....	319	669	1,405	2,288	3,745	5,671	6,613	5,358	2,624	653	82	-
2002.....	302	696	1,324	2,316	3,984	5,915	6,838	5,474	2,708	703	72	-
2001.....	313	647	1,260	2,379	4,164	6,119	6,857	5,500	2,575	689	85	-
2000.....	263	623	1,306	2,466	4,342	6,314	6,757	5,532	2,638	628	75	-
1999.....	274	636	1,310	2,711	4,605	6,532	6,903	5,418	2,467	644	91	1
All races, female												
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....
White, both sexes												
2004.....	223	503	1,015	1,676	2,750	4,406	5,519	4,617	2,239	629	62	1
2003.....	218	472	1,031	1,716	2,974	4,616	5,559	4,676	2,279	542	58	-
2002.....	207	505	958	1,746	3,112	4,801	5,765	4,780	2,329	588	53	-
2001.....	215	474	922	1,820	3,267	4,994	5,736	4,729	2,195	557	67	-
2000.....	192	444	960	1,851	3,380	5,167	5,637	4,785	2,257	524	56	-
1999.....	198	471	933	2,065	3,650	5,348	5,892	4,635	2,120	523	72	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Of prostate (C61)												
White, male												
2004.....	223	503	1,015	1,676	2,750	4,406	5,519	4,617	2,239	629	62	1
2003.....	218	472	1,031	1,716	2,974	4,616	5,559	4,676	2,279	542	58	-
2002.....	207	505	958	1,746	3,112	4,801	5,765	4,780	2,329	588	53	-
2001.....	215	474	922	1,820	3,267	4,994	5,736	4,729	2,195	557	67	-
2000.....	192	444	960	1,851	3,380	5,167	5,637	4,785	2,257	524	56	-
1999.....	198	471	933	2,065	3,650	5,348	5,892	4,635	2,120	523	72	1
White, female												
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....
Black, both sexes												
2004.....	105	176	322	490	709	953	921	630	345	102	23	1
2003.....	97	187	354	535	727	972	954	609	302	101	20	-
2002.....	88	174	338	545	825	1,029	1,001	641	344	103	18	-
2001.....	96	167	320	532	852	1,041	1,035	704	343	118	17	-
2000.....	67	174	323	586	905	1,070	1,037	678	348	96	18	-
1999.....	75	157	358	622	912	1,101	945	716	312	102	19	-
Black, male												
2004.....	105	176	322	490	709	953	921	630	345	102	23	1
2003.....	97	187	354	535	727	972	954	609	302	101	20	-
2002.....	88	174	338	545	825	1,029	1,001	641	344	103	18	-
2001.....	96	167	320	532	852	1,041	1,035	704	343	118	17	-
2000.....	67	174	323	586	905	1,070	1,037	678	348	96	18	-
1999.....	75	157	358	622	912	1,101	945	716	312	102	19	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Of testis (C62)												
All races, both sexes												
2004.....	29	14	13	7	7	5	11	8	1	-	-	-
2003.....	27	12	10	10	9	13	7	6	2	1	1	-
2002.....	25	18	18	14	12	5	12	6	5	-	-	-
2001.....	23	14	8	10	13	11	5	2	-	2	-	-
2000.....	20	9	12	7	8	7	6	6	1	-	-	-
1999.....	19	16	11	11	10	2	13	5	-	-	-	-
All races, male												
2004.....	29	14	13	7	7	5	11	8	1	-	-	-
2003.....	27	12	10	10	9	13	7	6	2	1	1	-
2002.....	25	18	18	14	12	5	12	6	5	-	-	-
2001.....	23	14	8	10	13	11	5	2	-	2	-	-
2000.....	20	9	12	7	8	7	6	6	1	-	-	-
1999.....	19	16	11	11	10	2	13	5	-	-	-	-
All races, female												
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....
White, both sexes												
2004.....	27	13	13	7	5	4	10	8	1	-	-	-
2003.....	26	11	10	10	9	9	6	6	1	1	1	-
2002.....	21	16	17	12	12	5	11	6	5	-	-	-
2001.....	21	12	6	8	11	9	5	2	-	2	-	-
2000.....	18	9	12	5	8	5	6	6	1	-	-	-
1999.....	16	16	9	11	9	2	13	5	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Of testis (C62)												
White, male												
2004.....	27	13	13	7	5	4	10	8	1	-	-	-
2003.....	26	11	10	10	9	9	6	6	1	1	1	-
2002.....	21	16	17	12	12	5	11	6	5	-	-	-
2001.....	21	12	6	8	11	9	5	2	-	2	-	-
2000.....	18	9	12	5	8	5	6	6	1	-	-	-
1999.....	16	16	9	11	9	2	13	5	-	-	-	-
White, female												
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....
Black, both sexes												
2004.....	2	1	-	-	-	1	1	-	-	-	-	-
2003.....	1	1	-	-	-	4	1	-	1	-	-	-
2002.....	3	1	-	2	-	-	1	-	-	-	-	-
2001.....	1	2	1	2	2	2	-	-	-	-	-	-
2000.....	2	-	-	1	-	2	-	-	-	-	-	-
1999.....	2	-	2	-	1	-	-	-	-	-	-	-
Black, male												
2004.....	2	1	-	-	-	1	1	-	-	-	-	-
2003.....	1	1	-	-	-	4	1	-	1	-	-	-
2002.....	3	1	-	2	-	-	1	-	-	-	-	-
2001.....	1	2	1	2	2	2	-	-	-	-	-	-
2000.....	2	-	-	1	-	2	-	-	-	-	-	-
1999.....	2	-	2	-	1	-	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Of penis and other and unspecified male genital organs (C60,C63)												
All races, both sexes												
2004.....	20	25	23	30	44	41	29	15	10	6	-	-
2003.....	21	18	37	31	28	39	35	18	16	3	-	-
2002.....	15	20	25	28	36	37	27	16	11	1	-	-
2001.....	19	19	23	27	31	31	29	23	11	2	-	-
2000.....	16	19	27	32	26	33	49	17	10	2	-	-
1999.....	11	16	22	31	29	36	33	26	15	3	-	-
All races, male												
2004.....	20	25	23	30	44	41	29	15	10	6	-	-
2003.....	21	18	37	31	28	39	35	18	16	3	-	-
2002.....	15	20	25	28	36	37	27	16	11	1	-	-
2001.....	19	19	23	27	31	31	29	23	11	2	-	-
2000.....	16	19	27	32	26	33	49	17	10	2	-	-
1999.....	11	16	22	31	29	36	33	26	15	3	-	-
All races, female												
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....
White, both sexes												
2004.....	19	24	18	21	40	37	28	13	10	4	-	-
2003.....	19	14	31	27	26	35	30	14	11	3	-	-
2002.....	12	16	19	26	29	33	25	16	10	1	-	-
2001.....	13	16	19	21	28	27	25	23	9	2	-	-
2000.....	15	19	24	30	25	27	44	16	8	1	-	-
1999.....	8	14	16	27	25	31	31	22	15	2	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Of penis and other and unspecified male genital organs (C60,C63)												
White, male												
2004.....	19	24	18	21	40	37	28	13	10	4	-	-
2003.....	19	14	31	27	26	35	30	14	11	3	-	-
2002.....	12	16	19	26	29	33	25	16	10	1	-	-
2001.....	13	16	19	21	28	27	25	23	9	2	-	-
2000.....	15	19	24	30	25	27	44	16	8	1	-	-
1999.....	8	14	16	27	25	31	31	22	15	2	-	-
White, female												
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....
Black, both sexes												
2004.....	-	1	3	9	4	4	1	-	-	2	-	-
2003.....	-	4	5	4	2	3	4	4	5	-	-	-
2002.....	2	3	5	1	5	3	2	-	1	-	-	-
2001.....	5	2	3	4	3	3	4	-	1	-	-	-
2000.....	1	-	2	2	-	4	5	1	2	1	-	-
1999.....	1	2	6	2	4	4	2	3	-	1	-	-
Black, male												
2004.....	-	1	3	9	4	4	1	-	-	2	-	-
2003.....	-	4	5	4	2	3	4	4	5	-	-	-
2002.....	2	3	5	1	5	3	2	-	1	-	-	-
2001.....	5	2	3	4	3	3	4	-	1	-	-	-
2000.....	1	-	2	2	-	4	5	1	2	1	-	-
1999.....	1	2	6	2	4	4	2	3	-	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant neoplasms of urinary tract (C64-C68)												
All races, both sexes												
2004.....	1,157	1,743	2,310	2,651	3,321	4,247	4,307	3,037	1,495	409	50	-
2003.....	1,123	1,700	2,248	2,604	3,364	4,212	4,189	2,991	1,347	396	46	-
2002.....	1,154	1,682	2,049	2,747	3,567	4,317	3,969	2,922	1,371	353	40	1
2001.....	1,152	1,514	2,142	2,605	3,587	4,101	3,920	2,881	1,292	357	47	-
2000.....	1,086	1,541	1,938	2,617	3,553	4,090	3,827	2,807	1,301	347	38	-
1999.....	1,063	1,482	1,944	2,565	3,601	3,991	3,567	2,665	1,230	349	36	1
All races, male												
2004.....	849	1,257	1,671	1,857	2,273	2,776	2,720	1,758	797	183	19	-
2003.....	799	1,229	1,638	1,807	2,239	2,789	2,623	1,695	698	177	14	-
2002.....	828	1,217	1,479	1,897	2,368	2,844	2,403	1,668	712	156	16	1
2001.....	832	1,090	1,531	1,827	2,446	2,674	2,439	1,668	665	149	17	-
2000.....	788	1,110	1,370	1,828	2,381	2,661	2,379	1,625	653	147	15	-
1999.....	758	1,066	1,381	1,786	2,408	2,657	2,193	1,494	593	146	15	1
All races, female												
2004.....	308	486	639	794	1,048	1,471	1,587	1,279	698	226	31	-
2003.....	324	471	610	797	1,125	1,423	1,566	1,296	649	219	32	-
2002.....	326	465	570	850	1,199	1,473	1,566	1,254	659	197	24	-
2001.....	320	424	611	778	1,141	1,427	1,481	1,213	627	208	30	-
2000.....	298	431	568	789	1,172	1,429	1,448	1,182	648	200	23	-
1999.....	305	416	563	779	1,193	1,334	1,374	1,171	637	203	21	-
White, both sexes												
2004.....	992	1,540	2,083	2,389	2,981	3,894	4,013	2,834	1,368	375	44	-
2003.....	965	1,482	1,998	2,314	3,072	3,859	3,863	2,779	1,250	362	40	-
2002.....	984	1,476	1,819	2,455	3,236	3,943	3,665	2,692	1,267	318	37	1
2001.....	982	1,323	1,875	2,353	3,266	3,754	3,604	2,681	1,181	334	43	-
2000.....	954	1,364	1,730	2,350	3,220	3,740	3,533	2,614	1,219	325	33	-
1999.....	913	1,294	1,726	2,288	3,263	3,662	3,313	2,472	1,127	307	35	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant neoplasms of urinary tract (C64-C68)												
White, male												
2004.....	740	1,128	1,516	1,697	2,074	2,578	2,574	1,659	747	174	18	-
2003.....	694	1,092	1,473	1,619	2,064	2,578	2,456	1,605	659	163	11	-
2002.....	725	1,091	1,330	1,720	2,183	2,635	2,236	1,561	674	147	16	1
2001.....	711	971	1,349	1,674	2,258	2,471	2,277	1,572	625	146	15	-
2000.....	694	984	1,242	1,654	2,189	2,469	2,214	1,517	622	140	14	-
1999.....	646	943	1,245	1,616	2,219	2,468	2,066	1,410	551	128	15	1
White, female												
2004.....	252	412	567	692	907	1,316	1,439	1,175	621	201	26	-
2003.....	271	390	525	695	1,008	1,281	1,407	1,174	591	199	29	-
2002.....	259	385	489	735	1,053	1,308	1,429	1,131	593	171	21	-
2001.....	271	352	526	679	1,008	1,283	1,327	1,109	556	188	28	-
2000.....	260	380	488	696	1,031	1,271	1,319	1,097	597	185	19	-
1999.....	267	351	481	672	1,044	1,194	1,247	1,062	576	179	20	-
Black, both sexes												
2004.....	142	169	195	211	279	296	240	176	106	27	6	-
2003.....	138	180	195	244	230	293	274	188	83	29	5	-
2002.....	134	175	190	242	269	315	248	193	90	28	3	-
2001.....	139	158	221	208	256	290	263	163	94	22	3	-
2000.....	113	147	174	227	281	301	262	162	71	20	4	-
1999.....	131	168	182	235	290	275	211	170	85	36	1	-
Black, male												
2004.....	95	107	130	118	168	162	116	78	38	4	1	-
2003.....	92	114	122	156	133	172	138	78	32	9	2	-
2002.....	80	111	124	138	142	168	129	88	33	5	-	-
2001.....	99	95	149	122	149	164	127	76	31	3	2	-
2000.....	78	99	109	146	160	162	145	90	25	6	1	-
1999.....	97	110	113	139	154	154	103	72	31	15	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasms of urinary tract (C64-C68)

Black, female

2004.....	47	62	65	93	111	134	124	98	68	23	5	-
2003.....	46	66	73	88	97	121	136	110	51	20	3	-
2002.....	54	64	66	104	127	147	119	105	57	23	3	-
2001.....	40	63	72	86	107	126	136	87	63	19	1	-
2000.....	35	48	65	81	121	139	117	72	46	14	3	-
1999.....	34	58	69	96	136	121	108	98	54	21	1	-

American Indian 1/, both sexes

2004.....	9	14	11	14	20	14	8	4	2	1	-	-
2003.....	10	15	15	11	16	12	16	3	2	-	-	-
2002.....	13	9	14	23	20	11	14	9	1	2	-	-
2001.....	13	11	19	16	14	10	10	4	4	-	-	-
2000.....	7	7	6	9	14	15	4	4	3	-	-	-
1999.....	8	10	11	15	16	15	7	5	2	1	-	-

American Indian 1/, male

2004.....	8	9	8	11	11	10	6	2	1	-	-	-
2003.....	4	8	14	7	12	5	7	1	1	-	-	-
2002.....	9	6	6	18	15	7	9	3	-	1	-	-
2001.....	8	6	11	11	10	6	6	2	2	-	-	-
2000.....	5	6	4	7	7	9	2	2	1	-	-	-
1999.....	6	8	5	12	12	10	4	3	1	1	-	-

American Indian 1/, female

2004.....	1	5	3	3	9	4	2	2	1	1	-	-
2003.....	6	7	1	4	4	7	9	2	1	-	-	-
2002.....	4	3	8	5	5	4	5	6	1	1	-	-
2001.....	5	5	8	5	4	4	4	2	2	-	-	-
2000.....	2	1	2	2	7	6	2	2	2	-	-	-
1999.....	2	2	6	3	4	5	3	2	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant neoplasms of urinary tract (C64-C68)												
Asian or Pacific Islander 2/, both sexes												
2004.....	14	20	21	37	41	43	46	23	19	6	-	-
2003.....	10	23	40	35	46	48	36	21	12	5	1	-
2002.....	23	22	26	27	42	48	42	28	13	5	-	-
2001.....	18	22	27	28	51	47	43	33	13	1	1	-
2000.....	12	23	28	31	38	34	28	27	8	2	1	-
1999.....	11	10	25	27	32	39	36	18	16	5	-	-
Asian or Pacific Islander 2/, male												
2004.....	6	13	17	31	20	26	24	19	11	5	-	-
2003.....	9	15	29	25	30	34	22	11	6	5	1	-
2002.....	14	9	19	21	28	34	29	16	5	3	-	-
2001.....	14	18	22	20	29	33	29	18	7	-	-	-
2000.....	11	21	15	21	25	21	18	16	5	1	-	-
1999.....	9	5	18	19	23	25	20	9	10	2	-	-
Asian or Pacific Islander 2/, female												
2004.....	8	7	4	6	21	17	22	4	8	1	-	-
2003.....	1	8	11	10	16	14	14	10	6	-	-	-
2002.....	9	13	7	6	14	14	13	12	8	2	-	-
2001.....	4	4	5	8	22	14	14	15	6	1	1	-
2000.....	1	2	13	10	13	13	10	11	3	1	1	-
1999.....	2	5	7	8	9	14	16	9	6	3	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Of kidney and renal pelvis (C64-C65)												
All races, both sexes												
2004.....	786	1,114	1,420	1,451	1,624	1,819	1,655	1,003	443	133	11	-
2003.....	763	1,121	1,377	1,409	1,681	1,849	1,626	1,033	420	107	7	-
2002.....	808	1,075	1,265	1,533	1,706	1,829	1,526	967	429	90	7	-
2001.....	813	1,006	1,322	1,395	1,826	1,810	1,496	947	390	82	5	-
2000.....	789	1,026	1,169	1,432	1,714	1,819	1,463	945	397	83	7	-
1999.....	720	959	1,152	1,374	1,702	1,696	1,314	867	341	91	6	-
All races, male												
2004.....	567	788	998	961	1,054	1,060	933	460	197	49	3	-
2003.....	535	798	989	916	1,059	1,114	922	486	175	40	2	-
2002.....	592	771	887	1,020	1,067	1,123	801	457	201	34	4	-
2001.....	586	722	912	928	1,165	1,052	814	500	164	23	-	-
2000.....	571	727	803	942	1,068	1,054	811	477	165	31	3	-
1999.....	506	671	803	894	1,064	1,016	701	421	140	29	-	-
All races, female												
2004.....	219	326	422	490	570	759	722	543	246	84	8	-
2003.....	228	323	388	493	622	735	704	547	245	67	5	-
2002.....	216	304	378	513	639	706	725	510	228	56	3	-
2001.....	227	284	410	467	661	758	682	447	226	59	5	-
2000.....	218	299	366	490	646	765	652	468	232	52	4	-
1999.....	214	288	349	480	638	680	613	446	201	62	6	-
White, both sexes												
2004.....	679	985	1,267	1,291	1,439	1,635	1,526	935	409	122	9	-
2003.....	660	967	1,208	1,252	1,520	1,684	1,499	956	380	98	5	-
2002.....	688	951	1,114	1,367	1,543	1,654	1,402	888	403	81	7	-
2001.....	687	878	1,152	1,259	1,649	1,639	1,371	873	354	77	5	-
2000.....	685	903	1,031	1,271	1,542	1,657	1,344	880	361	76	6	-
1999.....	616	841	1,027	1,226	1,531	1,541	1,211	802	303	77	5	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Of kidney and renal pelvis (C64-C65)												
White, male												
2004.....	494	703	893	854	943	958	869	431	181	47	2	-
2003.....	462	703	871	819	954	1,015	858	456	161	36	1	-
2002.....	511	682	785	917	969	1,033	733	423	195	31	4	-
2001.....	495	640	786	841	1,060	947	750	467	151	22	-	-
2000.....	495	637	716	829	966	965	744	444	152	29	2	-
1999.....	427	586	724	798	961	927	660	394	124	20	-	-
White, female												
2004.....	185	282	374	437	496	677	657	504	228	75	7	-
2003.....	198	264	337	433	566	669	641	500	219	62	4	-
2002.....	177	269	329	450	574	621	669	465	208	50	3	-
2001.....	192	238	366	418	589	692	621	406	203	55	5	-
2000.....	190	266	315	442	576	692	600	436	209	47	4	-
1999.....	189	255	303	428	570	614	551	408	179	57	5	-
Black, both sexes												
2004.....	89	104	128	127	147	155	102	59	31	8	2	-
2003.....	89	130	132	128	122	131	103	71	36	7	2	-
2002.....	92	104	122	135	125	150	102	67	21	7	-	-
2001.....	101	105	136	108	143	138	104	63	29	5	-	-
2000.....	87	100	114	134	139	132	103	57	30	7	1	-
1999.....	87	102	104	122	143	129	84	55	31	11	1	-
Black, male												
2004.....	62	70	86	81	92	84	50	23	15	-	1	-
2003.....	64	82	88	79	79	78	53	26	13	2	1	-
2002.....	62	79	85	80	71	75	51	29	6	2	-	-
2001.....	72	67	101	67	87	84	53	30	10	1	-	-
2000.....	61	69	74	96	84	68	58	30	10	2	1	-
1999.....	65	74	66	77	83	72	32	21	13	8	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Of kidney and renal pelvis (C64-C65)

Black, female

2004.....	27	34	42	46	55	71	52	36	16	8	1	-
2003.....	25	48	44	49	43	53	50	45	23	5	1	-
2002.....	30	25	37	55	54	75	51	38	15	5	-	-
2001.....	29	38	35	41	56	54	51	33	19	4	-	-
2000.....	26	31	40	38	55	64	45	27	20	5	-	-
1999.....	22	28	38	45	60	57	52	34	18	3	1	-

American Indian 1/, both sexes

2004.....	7	12	10	12	18	12	6	1	2	-	-	-
2003.....	8	10	10	8	15	7	10	1	1	-	-	-
2002.....	11	7	12	16	18	7	8	4	-	1	-	-
2001.....	11	9	16	12	10	9	4	1	4	-	-	-
2000.....	6	5	5	7	12	9	1	3	1	-	-	-
1999.....	8	9	7	10	11	14	6	2	1	1	-	-

American Indian 1/, male

2004.....	6	7	7	9	10	8	4	-	1	-	-	-
2003.....	4	6	10	5	11	3	4	-	-	-	-	-
2002.....	7	5	6	12	14	4	6	1	-	-	-	-
2001.....	8	5	9	9	8	5	1	-	2	-	-	-
2000.....	5	5	3	5	6	7	-	1	-	-	-	-
1999.....	6	7	3	7	8	9	4	1	-	1	-	-

American Indian 1/, female

2004.....	1	5	3	3	8	4	2	1	1	-	-	-
2003.....	4	4	-	3	4	4	6	1	1	-	-	-
2002.....	4	2	6	4	4	3	2	3	-	1	-	-
2001.....	3	4	7	3	2	4	3	1	2	-	-	-
2000.....	1	-	2	2	6	2	1	2	1	-	-	-
1999.....	2	2	4	3	3	5	2	1	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Of kidney and renal pelvis (C64-C65)												
Asian or Pacific Islander 2/, both sexes												
2004.....	11	13	15	21	20	17	21	8	1	3	-	-
2003.....	6	14	27	21	24	27	14	5	3	2	-	-
2002.....	17	13	17	15	20	18	14	8	5	1	-	-
2001.....	14	14	18	16	24	24	17	10	3	-	-	-
2000.....	11	18	19	20	21	21	15	5	5	-	-	-
1999.....	9	7	14	16	17	12	13	8	6	2	-	-
Asian or Pacific Islander 2/, male												
2004.....	5	8	12	17	9	10	10	6	-	2	-	-
2003.....	5	7	20	13	15	18	7	4	1	2	-	-
2002.....	12	5	11	11	13	11	11	4	-	1	-	-
2001.....	11	10	16	11	10	16	10	3	1	-	-	-
2000.....	10	16	10	12	12	14	9	2	3	-	-	-
1999.....	8	4	10	12	12	8	5	5	3	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	6	5	3	4	11	7	11	2	1	1	-	-
2003.....	1	7	7	8	9	9	7	1	2	-	-	-
2002.....	5	8	6	4	7	7	3	4	5	-	-	-
2001.....	3	4	2	5	14	8	7	7	2	-	-	-
2000.....	1	2	9	8	9	7	6	3	2	-	-	-
1999.....	1	3	4	4	5	4	8	3	3	2	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Of bladder (C67)												
All races, both sexes												
2004.....	346	600	852	1,143	1,610	2,307	2,540	1,976	1,012	273	38	-
2003.....	341	554	837	1,148	1,588	2,239	2,440	1,850	888	278	36	-
2002.....	321	572	738	1,151	1,756	2,366	2,341	1,864	908	255	32	1
2001.....	314	484	781	1,147	1,667	2,179	2,309	1,865	864	265	42	-
2000.....	280	486	737	1,119	1,736	2,155	2,252	1,790	875	256	31	-
1999.....	323	492	752	1,126	1,797	2,178	2,139	1,714	852	249	30	1
All races, male												
2004.....	270	455	648	859	1,162	1,646	1,730	1,275	583	133	15	-
2003.....	252	419	630	865	1,133	1,603	1,651	1,157	508	131	11	-
2002.....	223	425	559	845	1,244	1,659	1,540	1,163	498	118	12	1
2001.....	233	359	597	859	1,229	1,554	1,562	1,129	482	123	17	-
2000.....	212	369	547	838	1,256	1,542	1,512	1,110	478	110	12	-
1999.....	243	374	559	849	1,283	1,575	1,433	1,039	438	116	15	1
All races, female												
2004.....	76	145	204	284	448	661	810	701	429	140	23	-
2003.....	89	135	207	283	455	636	789	693	380	147	25	-
2002.....	98	147	179	306	512	707	801	701	410	137	20	-
2001.....	81	125	184	288	438	625	747	736	382	142	25	-
2000.....	68	117	190	281	480	613	740	680	397	146	19	-
1999.....	80	118	193	277	514	603	706	675	414	133	15	-
White, both sexes												
2004.....	297	535	784	1,049	1,465	2,144	2,379	1,843	920	250	34	-
2003.....	288	492	761	1,024	1,472	2,058	2,252	1,717	835	253	32	-
2002.....	280	502	668	1,031	1,594	2,177	2,167	1,722	831	229	29	1
2001.....	274	426	694	1,039	1,534	2,012	2,126	1,743	792	247	38	-
2000.....	254	439	670	1,020	1,585	1,977	2,079	1,665	830	242	27	-
1999.....	283	429	664	1,005	1,638	2,011	1,997	1,592	787	221	30	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Of bladder (C67)												
White, male												
2004.....	236	415	600	810	1,079	1,551	1,649	1,205	550	126	15	-
2003.....	221	378	584	779	1,069	1,496	1,551	1,098	484	121	9	-
2002.....	203	390	516	774	1,157	1,544	1,446	1,095	467	112	12	1
2001.....	205	323	546	798	1,148	1,462	1,467	1,069	455	121	15	-
2000.....	195	335	508	779	1,170	1,447	1,414	1,037	460	106	12	-
1999.....	210	340	503	778	1,199	1,477	1,351	985	412	107	15	1
White, female												
2004.....	61	120	184	239	386	593	730	638	370	124	19	-
2003.....	67	114	177	245	403	562	701	619	351	132	23	-
2002.....	77	112	152	257	437	633	721	627	364	117	17	-
2001.....	69	103	148	241	386	550	659	674	337	126	23	-
2000.....	59	104	162	241	415	530	665	628	370	136	15	-
1999.....	73	89	161	227	439	534	646	607	375	114	15	-
Black, both sexes												
2004.....	45	56	62	77	126	136	135	115	75	19	4	-
2003.....	48	50	59	109	96	156	161	115	43	22	3	-
2002.....	33	60	60	103	139	157	142	121	68	21	3	-
2001.....	35	48	77	93	105	143	152	97	62	17	3	-
2000.....	24	41	59	86	134	160	158	103	41	12	3	-
1999.....	39	59	74	107	140	142	122	110	54	25	-	-
Black, male												
2004.....	31	33	43	34	71	77	65	55	23	4	-	-
2003.....	28	32	34	74	50	90	83	51	18	7	1	-
2002.....	16	30	36	57	71	91	75	56	26	3	-	-
2001.....	26	27	45	50	61	74	72	44	21	2	2	-
2000.....	16	28	35	48	72	87	87	59	15	3	-	-
1999.....	32	32	46	60	70	82	68	49	18	7	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Of bladder (C67)												
Black, female												
2004.....	14	23	19	43	55	59	70	60	52	15	4	-
2003.....	20	18	25	35	46	66	78	64	25	15	2	-
2002.....	17	30	24	46	68	66	67	65	42	18	3	-
2001.....	9	21	32	43	44	69	80	53	41	15	1	-
2000.....	8	13	24	38	62	73	71	44	26	9	3	-
1999.....	7	27	28	47	70	60	54	61	36	18	-	-
American Indian 1/, both sexes												
2004.....	2	2	1	2	2	2	2	3	-	1	-	-
2003.....	2	5	5	3	1	5	6	2	1	-	-	-
2002.....	2	2	2	7	2	3	5	5	1	1	-	-
2001.....	2	2	2	4	4	1	6	3	-	-	-	-
2000.....	1	1	1	2	1	5	2	1	1	-	-	-
1999.....	-	1	4	5	5	1	1	2	1	-	-	-
American Indian 1/, male												
2004.....	2	2	1	2	1	2	2	2	-	-	-	-
2003.....	-	2	4	2	1	2	3	1	1	-	-	-
2002.....	2	1	-	6	1	2	2	2	-	1	-	-
2001.....	-	1	1	2	2	1	5	2	-	-	-	-
2000.....	-	1	1	2	1	1	2	1	1	-	-	-
1999.....	-	1	2	5	4	1	-	1	1	-	-	-
American Indian 1/, female												
2004.....	-	-	-	-	1	-	-	1	-	1	-	-
2003.....	2	3	1	1	-	3	3	1	-	-	-	-
2002.....	-	1	2	1	1	1	3	3	1	-	-	-
2001.....	2	1	1	2	2	-	1	1	-	-	-	-
2000.....	1	-	-	-	-	4	-	-	-	-	-	-
1999.....	-	-	2	-	1	-	1	1	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Of bladder (C67)												
Asian or Pacific Islander 2/, both sexes												
2004.....	2	7	5	15	17	25	24	15	17	3	-	-
2003.....	3	7	12	12	19	20	21	16	9	3	1	-
2002.....	6	8	8	10	21	29	27	16	8	4	-	-
2001.....	3	8	8	11	24	23	25	22	10	1	1	-
2000.....	1	5	7	11	16	13	13	21	3	2	1	-
1999.....	1	3	10	9	14	24	19	10	10	3	-	-
Asian or Pacific Islander 2/, male												
2004.....	1	5	4	13	11	16	14	13	10	3	-	-
2003.....	3	7	8	10	13	15	14	7	5	3	1	-
2002.....	2	4	7	8	15	22	17	10	5	2	-	-
2001.....	2	8	5	9	18	17	18	14	6	-	-	-
2000.....	1	5	3	9	13	7	9	13	2	1	-	-
1999.....	1	1	8	6	10	15	14	4	7	2	-	-
Asian or Pacific Islander 2/, female												
2004.....	1	2	1	2	6	9	10	2	7	-	-	-
2003.....	-	-	4	2	6	5	7	9	4	-	-	-
2002.....	4	4	1	2	6	7	10	6	3	2	-	-
2001.....	1	-	3	2	6	6	7	8	4	1	1	-
2000.....	-	-	4	2	3	6	4	8	1	1	1	-
1999.....	-	2	2	3	4	9	5	6	3	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Of other and unspecified sites within the urinary tract (C66,C68)

All races, both sexes

2004.....	25	29	38	57	87	121	112	58	40	3	1	-
2003.....	19	25	34	47	95	124	123	108	39	11	3	-
2002.....	25	35	46	63	105	122	102	91	34	8	1	-
2001.....	25	24	39	63	94	112	115	69	38	10	-	-
2000.....	17	29	32	66	103	116	112	72	29	8	-	-
1999.....	20	31	40	65	102	117	114	84	37	9	-	-

All races, male

2004.....	12	14	25	37	57	70	57	23	17	1	1	-
2003.....	12	12	19	26	47	72	50	52	15	6	1	-
2002.....	13	21	33	32	57	62	62	48	13	4	-	-
2001.....	13	9	22	40	52	68	63	39	19	3	-	-
2000.....	5	14	20	48	57	65	56	38	10	6	-	-
1999.....	9	21	19	43	61	66	59	34	15	1	-	-

All races, female

2004.....	13	15	13	20	30	51	55	35	23	2	-	-
2003.....	7	13	15	21	48	52	73	56	24	5	2	-
2002.....	12	14	13	31	48	60	40	43	21	4	1	-
2001.....	12	15	17	23	42	44	52	30	19	7	-	-
2000.....	12	15	12	18	46	51	56	34	19	2	-	-
1999.....	11	10	21	22	41	51	55	50	22	8	-	-

White, both sexes

2004.....	16	20	32	49	77	115	108	56	39	3	1	-
2003.....	17	23	29	38	80	117	112	106	35	11	3	-
2002.....	16	23	37	57	99	112	96	82	33	8	1	-
2001.....	21	19	29	55	83	103	107	65	35	10	-	-
2000.....	15	22	29	59	93	106	110	69	28	7	-	-
1999.....	14	24	35	57	94	110	105	78	37	9	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Of other and unspecified sites within the urinary tract (C66,C68)

White, male

2004.....	10	10	23	33	52	69	56	23	16	1	1	-
2003.....	11	11	18	21	41	67	47	51	14	6	1	-
2002.....	11	19	29	29	57	58	57	43	12	4	-	-
2001.....	11	8	17	35	50	62	60	36	19	3	-	-
2000.....	4	12	18	46	53	57	56	36	10	5	-	-
1999.....	9	17	18	40	59	64	55	31	15	1	-	-

White, female

2004.....	6	10	9	16	25	46	52	33	23	2	-	-
2003.....	6	12	11	17	39	50	65	55	21	5	2	-
2002.....	5	4	8	28	42	54	39	39	21	4	1	-
2001.....	10	11	12	20	33	41	47	29	16	7	-	-
2000.....	11	10	11	13	40	49	54	33	18	2	-	-
1999.....	5	7	17	17	35	46	50	47	22	8	-	-

Black, both sexes

2004.....	8	9	5	7	6	5	3	2	-	-	-	-
2003.....	1	-	4	7	12	6	10	2	4	-	-	-
2002.....	9	11	8	4	5	8	4	5	1	-	-	-
2001.....	3	5	8	7	8	9	7	3	3	-	-	-
2000.....	2	6	1	7	8	9	1	2	-	1	-	-
1999.....	5	7	4	6	7	4	5	5	-	-	-	-

Black, male

2004.....	2	4	1	3	5	1	1	-	-	-	-	-
2003.....	-	-	-	3	4	4	2	1	1	-	-	-
2002.....	2	2	3	1	-	2	3	3	1	-	-	-
2001.....	1	1	3	5	1	6	2	2	-	-	-	-
2000.....	1	2	-	2	4	7	-	1	-	1	-	-
1999.....	-	4	1	2	1	-	3	2	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Of other and unspecified sites within the urinary tract (C66,C68)

Asian or Pacific Islander 2/, both sexes

2004.....	1	-	1	1	4	1	1	-	1	-	-	-
2003.....	1	2	1	2	3	1	1	-	-	-	-	-
2002.....	-	1	1	2	1	1	1	4	-	-	-	-
2001.....	1	-	1	1	3	-	1	1	-	-	-	-
2000.....	-	-	2	-	1	-	-	1	-	-	-	-
1999.....	1	-	1	2	1	3	4	-	-	-	-	-

Asian or Pacific Islander 2/, male

2004.....	-	-	1	1	-	-	-	-	1	-	-	-
2003.....	1	1	1	2	2	1	1	-	-	-	-	-
2002.....	-	-	1	2	-	1	1	2	-	-	-	-
2001.....	1	-	1	-	1	-	1	1	-	-	-	-
2000.....	-	-	2	-	-	-	-	1	-	-	-	-
1999.....	-	-	-	1	1	2	1	-	-	-	-	-

Asian or Pacific Islander 2/, female

2004.....	1	-	-	-	4	1	1	-	-	-	-	-
2003.....	-	1	-	-	1	-	-	-	-	-	-	-
2002.....	-	1	-	-	1	-	-	2	-	-	-	-
2001.....	-	-	-	1	2	-	-	-	-	-	-	-
2000.....	-	-	-	-	1	-	-	-	-	-	-	-
1999.....	1	-	1	1	-	1	3	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant neoplasm of eye and adnexa (C69)												
All races, both sexes												
2004.....	13	16	23	25	26	30	15	12	6	5	-	-
2003.....	17	27	23	18	26	29	28	15	12	2	3	-
2002.....	15	29	23	20	22	33	28	14	10	7	2	-
2001.....	15	12	18	24	26	32	32	12	12	2	1	-
2000.....	16	21	22	25	28	30	23	20	10	3	1	-
1999.....	14	14	18	20	31	26	31	17	14	4	-	-
All races, male												
2004.....	8	8	17	15	12	16	10	2	2	-	-	-
2003.....	7	13	15	11	12	16	16	9	2	1	2	-
2002.....	7	18	13	12	10	18	10	7	3	-	1	-
2001.....	11	7	10	14	14	15	16	4	3	-	-	-
2000.....	11	12	13	17	15	18	10	7	5	-	1	-
1999.....	9	6	11	13	20	14	7	7	3	1	-	-
All races, female												
2004.....	5	8	6	10	14	14	5	10	4	5	-	-
2003.....	10	14	8	7	14	13	12	6	10	1	1	-
2002.....	8	11	10	8	12	15	18	7	7	7	1	-
2001.....	4	5	8	10	12	17	16	8	9	2	1	-
2000.....	5	9	9	8	13	12	13	13	5	3	-	-
1999.....	5	8	7	7	11	12	24	10	11	3	-	-
White, both sexes												
2004.....	12	16	23	25	26	30	15	12	6	5	-	-
2003.....	17	24	23	17	26	28	28	14	12	2	3	-
2002.....	14	27	22	18	22	32	28	14	10	6	2	-
2001.....	15	12	17	24	26	29	32	12	12	2	1	-
2000.....	16	19	21	24	28	30	21	19	10	2	1	-
1999.....	14	12	18	19	31	26	31	17	14	4	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)												
All races, both sexes												
2004.....	1,070	1,378	1,372	1,392	1,503	1,407	1,136	564	204	34	5	-
2003.....	1,105	1,337	1,361	1,349	1,487	1,449	1,061	513	173	28	3	-
2002.....	1,158	1,276	1,284	1,407	1,547	1,454	981	536	174	39	4	-
2001.....	1,121	1,174	1,290	1,354	1,469	1,506	1,060	528	157	31	5	-
2000.....	1,109	1,110	1,227	1,409	1,623	1,437	1,041	521	166	28	2	-
1999.....	1,038	1,206	1,247	1,498	1,621	1,543	949	496	144	22	1	-
All races, male												
2004.....	675	832	821	741	810	717	520	219	60	11	2	-
2003.....	689	741	789	741	815	745	522	214	65	8	2	-
2002.....	665	773	721	819	859	750	460	213	54	10	2	-
2001.....	673	734	736	763	826	777	511	202	48	9	2	-
2000.....	667	660	719	789	889	705	482	212	53	9	1	-
1999.....	613	670	747	858	870	785	441	192	47	7	1	-
All races, female												
2004.....	395	546	551	651	693	690	616	345	144	23	3	-
2003.....	416	596	572	608	672	704	539	299	108	20	1	-
2002.....	493	503	563	588	688	704	521	323	120	29	2	-
2001.....	448	440	554	591	643	729	549	326	109	22	3	-
2000.....	442	450	508	620	734	732	559	309	113	19	1	-
1999.....	425	536	500	640	751	758	508	304	97	15	-	-
White, both sexes												
2004.....	969	1,284	1,280	1,298	1,413	1,324	1,083	536	191	29	3	-
2003.....	1,008	1,225	1,271	1,247	1,390	1,368	1,007	478	164	22	3	-
2002.....	1,056	1,165	1,194	1,310	1,454	1,366	926	507	164	36	4	-
2001.....	1,030	1,073	1,197	1,250	1,378	1,433	1,007	500	148	26	4	-
2000.....	1,026	1,019	1,142	1,321	1,543	1,362	978	486	157	27	1	-
1999.....	950	1,123	1,153	1,400	1,545	1,463	900	473	131	20	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)

White, male

2004.....	609	773	769	701	773	672	493	213	57	9	2	-
2003.....	625	682	741	684	764	702	494	201	61	7	2	-
2002.....	618	700	669	767	810	705	431	204	51	9	2	-
2001.....	620	676	679	717	783	734	488	190	43	7	1	-
2000.....	609	611	673	747	846	669	459	201	51	9	-	-
1999.....	560	625	691	807	834	745	424	183	45	6	1	-

White, female

2004.....	360	511	511	597	640	652	590	323	134	20	1	-
2003.....	383	543	530	563	626	666	513	277	103	15	1	-
2002.....	438	465	525	543	644	661	495	303	113	27	2	-
2001.....	410	397	518	533	595	699	519	310	105	19	3	-
2000.....	417	408	469	574	697	693	519	285	106	18	1	-
1999.....	390	498	462	593	711	718	476	290	86	14	-	-

Black, both sexes

2004.....	78	69	73	69	70	59	40	21	11	3	2	-
2003.....	77	87	66	79	73	62	41	23	7	6	-	-
2002.....	78	85	64	75	71	71	43	20	9	3	-	-
2001.....	69	76	72	83	70	54	40	24	8	5	1	-
2000.....	70	73	66	68	66	58	51	26	6	1	1	-
1999.....	70	70	72	77	58	67	40	21	9	1	-	-

Black, male

2004.....	51	45	42	32	25	31	20	4	3	1	-	-
2003.....	51	47	35	48	37	35	19	4	2	1	-	-
2002.....	32	55	40	38	35	34	23	4	2	1	-	-
2001.....	41	44	45	36	34	27	15	8	4	2	1	-
2000.....	49	40	37	30	37	25	18	6	2	-	1	-
1999.....	43	35	41	42	26	30	11	8	2	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)

Black, female

2004.....	27	24	31	37	45	28	20	17	8	2	2	-
2003.....	26	40	31	31	36	27	22	19	5	5	-	-
2002.....	46	30	24	37	36	37	20	16	7	2	-	-
2001.....	28	32	27	47	36	27	25	16	4	3	-	-
2000.....	21	33	29	38	29	33	33	20	4	1	-	-
1999.....	27	35	31	35	32	37	29	13	7	1	-	-

American Indian 1/, both sexes

2004.....	5	2	3	1	2	3	2	-	-	-	-	-
2003.....	6	4	5	5	3	2	1	3	-	-	-	-
2002.....	4	4	1	2	4	3	-	1	-	-	-	-
2001.....	7	5	3	5	3	3	2	-	-	-	-	-
2000.....	1	2	1	3	1	6	2	1	-	-	-	-
1999.....	4	3	5	4	2	2	-	-	-	1	-	-

American Indian 1/, male

2004.....	4	1	3	-	1	1	1	-	-	-	-	-
2003.....	4	2	2	2	1	2	1	2	-	-	-	-
2002.....	2	4	-	1	3	1	-	-	-	-	-	-
2001.....	1	3	2	3	1	3	1	-	-	-	-	-
2000.....	1	1	-	2	-	4	1	1	-	-	-	-
1999.....	1	3	3	1	1	1	-	-	-	1	-	-

American Indian 1/, female

2004.....	1	1	-	1	1	2	1	-	-	-	-	-
2003.....	2	2	3	3	2	-	-	1	-	-	-	-
2002.....	2	-	1	1	1	2	-	1	-	-	-	-
2001.....	6	2	1	2	2	-	1	-	-	-	-	-
2000.....	-	1	1	1	1	2	1	-	-	-	-	-
1999.....	3	-	2	3	1	1	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)												
Asian or Pacific Islander 2/, both sexes												
2004.....	18	23	16	24	18	21	11	7	2	2	-	-
2003.....	14	21	19	18	21	17	12	9	2	-	-	-
2002.....	20	22	25	20	18	14	12	8	1	-	-	-
2001.....	15	20	18	16	18	16	11	4	1	-	-	-
2000.....	12	16	18	17	13	11	10	8	3	-	-	-
1999.....	14	10	17	17	16	11	9	2	4	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	11	13	7	8	11	13	6	2	-	1	-	-
2003.....	9	10	11	7	13	6	8	7	2	-	-	-
2002.....	13	14	12	13	11	10	6	5	1	-	-	-
2001.....	11	11	10	7	8	13	7	4	1	-	-	-
2000.....	8	8	9	10	6	7	4	4	-	-	-	-
1999.....	9	7	12	8	9	9	6	1	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	7	10	9	16	7	8	5	5	2	1	-	-
2003.....	5	11	8	11	8	11	4	2	-	-	-	-
2002.....	7	8	13	7	7	4	6	3	-	-	-	-
2001.....	4	9	8	9	10	3	4	-	-	-	-	-
2000.....	4	8	9	7	7	4	6	4	3	-	-	-
1999.....	5	3	5	9	7	2	3	1	4	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Of brain (C71)												
All races, both sexes												
2004.....	1,051	1,359	1,344	1,379	1,481	1,386	1,115	547	198	31	5	-
2003.....	1,087	1,314	1,349	1,325	1,467	1,418	1,040	496	161	28	3	-
2002.....	1,135	1,253	1,262	1,377	1,507	1,428	954	525	166	39	4	-
2001.....	1,103	1,157	1,267	1,331	1,437	1,472	1,037	515	153	31	4	-
2000.....	1,088	1,091	1,209	1,386	1,598	1,415	1,020	504	161	26	2	-
1999.....	1,011	1,187	1,223	1,478	1,587	1,508	926	481	141	21	1	-
All races, male												
2004.....	666	819	804	736	799	706	511	215	60	11	2	-
2003.....	679	725	781	731	805	735	518	205	62	8	2	-
2002.....	655	758	711	800	834	735	441	211	51	10	2	-
2001.....	663	721	722	749	808	762	502	198	47	9	2	-
2000.....	655	651	712	777	879	695	473	208	51	8	1	-
1999.....	601	661	733	846	851	769	433	185	47	7	1	-
All races, female												
2004.....	385	540	540	643	682	680	604	332	138	20	3	-
2003.....	408	589	568	594	662	683	522	291	99	20	1	-
2002.....	480	495	551	577	673	693	513	314	115	29	2	-
2001.....	440	436	545	582	629	710	535	317	106	22	2	-
2000.....	433	440	497	609	719	720	547	296	110	18	1	-
1999.....	410	526	490	632	736	739	493	296	94	14	-	-
White, both sexes												
2004.....	955	1,268	1,256	1,287	1,393	1,305	1,063	519	185	26	3	-
2003.....	991	1,208	1,260	1,225	1,375	1,341	988	462	152	22	3	-
2002.....	1,037	1,148	1,177	1,284	1,419	1,342	902	496	157	36	4	-
2001.....	1,016	1,058	1,178	1,228	1,346	1,400	986	487	144	26	3	-
2000.....	1,007	1,003	1,126	1,301	1,518	1,345	958	471	152	26	1	-
1999.....	928	1,108	1,134	1,380	1,514	1,431	880	460	129	19	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Of brain (C71)												
White, male												
2004.....	601	763	753	696	764	662	485	209	57	9	2	-
2003.....	616	669	733	675	756	693	491	192	58	7	2	-
2002.....	611	688	662	750	789	691	414	202	49	9	2	-
2001.....	612	665	668	703	765	719	480	186	42	7	1	-
2000.....	598	604	667	736	836	662	451	198	49	8	-	-
1999.....	551	617	680	795	818	730	417	177	45	6	1	-
White, female												
2004.....	354	505	503	591	629	643	578	310	128	17	1	-
2003.....	375	539	527	550	619	648	497	270	94	15	1	-
2002.....	426	460	515	534	630	651	488	294	108	27	2	-
2001.....	404	393	510	525	581	681	506	301	102	19	2	-
2000.....	409	399	459	565	682	683	507	273	103	18	1	-
1999.....	377	491	454	585	696	701	463	283	84	13	-	-
Black, both sexes												
2004.....	73	66	70	68	68	58	39	21	11	3	2	-
2003.....	76	82	65	77	68	59	40	22	7	6	-	-
2002.....	75	79	61	71	67	69	40	20	8	3	-	-
2001.....	67	75	69	82	70	53	39	24	8	5	1	-
2000.....	68	71	64	65	66	55	50	25	6	-	1	-
1999.....	66	66	69	77	55	64	37	19	9	1	-	-
Black, male												
2004.....	50	42	41	32	23	30	19	4	3	1	-	-
2003.....	50	44	35	47	35	35	19	4	2	1	-	-
2002.....	30	52	38	36	32	33	21	4	1	1	-	-
2001.....	39	43	43	36	34	27	15	8	4	2	1	-
2000.....	48	39	36	29	37	23	17	5	2	-	1	-
1999.....	40	34	40	42	23	29	10	7	2	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Of brain (C71)												
Black, female												
2004.....	23	24	29	36	45	28	20	17	8	2	2	-
2003.....	26	38	30	30	33	24	21	18	5	5	-	-
2002.....	45	27	23	35	35	36	19	16	7	2	-	-
2001.....	28	32	26	46	36	26	24	16	4	3	-	-
2000.....	20	32	28	36	29	32	33	20	4	-	-	-
1999.....	26	32	29	35	32	35	27	12	7	1	-	-
American Indian 1/, both sexes												
2004.....	5	2	3	1	2	3	2	-	-	-	-	-
2003.....	6	4	5	5	3	2	1	3	-	-	-	-
2002.....	4	4	1	2	4	3	-	1	-	-	-	-
2001.....	7	5	3	5	3	3	2	-	-	-	-	-
2000.....	1	2	1	3	1	6	2	1	-	-	-	-
1999.....	3	3	4	4	2	2	-	-	-	1	-	-
American Indian 1/, male												
2004.....	4	1	3	-	1	1	1	-	-	-	-	-
2003.....	4	2	2	2	1	2	1	2	-	-	-	-
2002.....	2	4	-	1	3	1	-	-	-	-	-	-
2001.....	1	3	2	3	1	3	1	-	-	-	-	-
2000.....	1	1	-	2	-	4	1	1	-	-	-	-
1999.....	1	3	2	1	1	1	-	-	-	1	-	-
American Indian 1/, female												
2004.....	1	1	-	1	1	2	1	-	-	-	-	-
2003.....	2	2	3	3	2	-	-	1	-	-	-	-
2002.....	2	-	1	1	1	2	-	1	-	-	-	-
2001.....	6	2	1	2	2	-	1	-	-	-	-	-
2000.....	-	1	1	1	1	2	1	-	-	-	-	-
1999.....	2	-	2	3	1	1	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Of brain (C71)												
Asian or Pacific Islander 2/, both sexes												
2004.....	18	23	15	23	18	20	11	7	2	2	-	-
2003.....	14	20	19	18	21	16	11	9	2	-	-	-
2002.....	19	22	23	20	17	14	12	8	1	-	-	-
2001.....	13	19	17	16	18	16	10	4	1	-	-	-
2000.....	12	15	18	17	13	9	10	7	3	-	-	-
1999.....	14	10	16	17	16	11	9	2	3	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	11	13	7	8	11	13	6	2	-	1	-	-
2003.....	9	10	11	7	13	5	7	7	2	-	-	-
2002.....	12	14	11	13	10	10	6	5	1	-	-	-
2001.....	11	10	9	7	8	13	6	4	1	-	-	-
2000.....	8	7	9	10	6	6	4	4	-	-	-	-
1999.....	9	7	11	8	9	9	6	1	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	7	10	8	15	7	7	5	5	2	1	-	-
2003.....	5	10	8	11	8	11	4	2	-	-	-	-
2002.....	7	8	12	7	7	4	6	3	-	-	-	-
2001.....	2	9	8	9	10	3	4	-	-	-	-	-
2000.....	4	8	9	7	7	3	6	3	3	-	-	-
1999.....	5	3	5	9	7	2	3	1	3	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Of other parts of central nervous system (C70,C72)												
All races, both sexes												
2004.....	19	19	28	13	22	21	21	17	6	3	-	-
2003.....	18	23	12	24	20	31	21	17	12	-	-	-
2002.....	23	23	22	30	40	26	27	11	8	-	-	-
2001.....	18	17	23	23	32	34	23	13	4	-	1	-
2000.....	21	19	18	23	25	22	21	17	5	2	-	-
1999.....	27	19	24	20	34	35	23	15	3	1	-	-
All races, male												
2004.....	9	13	17	5	11	11	9	4	-	-	-	-
2003.....	10	16	8	10	10	10	4	9	3	-	-	-
2002.....	10	15	10	19	25	15	19	2	3	-	-	-
2001.....	10	13	14	14	18	15	9	4	1	-	-	-
2000.....	12	9	7	12	10	10	9	4	2	1	-	-
1999.....	12	9	14	12	19	16	8	7	-	-	-	-
All races, female												
2004.....	10	6	11	8	11	10	12	13	6	3	-	-
2003.....	8	7	4	14	10	21	17	8	9	-	-	-
2002.....	13	8	12	11	15	11	8	9	5	-	-	-
2001.....	8	4	9	9	14	19	14	9	3	-	1	-
2000.....	9	10	11	11	15	12	12	13	3	1	-	-
1999.....	15	10	10	8	15	19	15	8	3	1	-	-
White, both sexes												
2004.....	14	16	24	11	20	19	20	17	6	3	-	-
2003.....	17	17	11	22	15	27	19	16	12	-	-	-
2002.....	19	17	17	26	35	24	24	11	7	-	-	-
2001.....	14	15	19	22	32	33	21	13	4	-	1	-
2000.....	19	16	16	20	25	17	20	15	5	1	-	-
1999.....	22	15	19	20	31	32	20	13	2	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Of other parts of central nervous system (C70,C72)												
White, male												
2004.....	8	10	16	5	9	10	8	4	-	-	-	-
2003.....	9	13	8	9	8	9	3	9	3	-	-	-
2002.....	7	12	7	17	21	14	17	2	2	-	-	-
2001.....	8	11	11	14	18	15	8	4	1	-	-	-
2000.....	11	7	6	11	10	7	8	3	2	1	-	-
1999.....	9	8	11	12	16	15	7	6	-	-	-	-
White, female												
2004.....	6	6	8	6	11	9	12	13	6	3	-	-
2003.....	8	4	3	13	7	18	16	7	9	-	-	-
2002.....	12	5	10	9	14	10	7	9	5	-	-	-
2001.....	6	4	8	8	14	18	13	9	3	-	1	-
2000.....	8	9	10	9	15	10	12	12	3	-	-	-
1999.....	13	7	8	8	15	17	13	7	2	1	-	-
Black, both sexes												
2004.....	5	3	3	1	2	1	1	-	-	-	-	-
2003.....	1	5	1	2	5	3	1	1	-	-	-	-
2002.....	3	6	3	4	4	2	3	-	1	-	-	-
2001.....	2	1	3	1	-	1	1	-	-	-	-	-
2000.....	2	2	2	3	-	3	1	1	-	1	-	-
1999.....	4	4	3	-	3	3	3	2	-	-	-	-
Black, male												
2004.....	1	3	1	-	2	1	1	-	-	-	-	-
2003.....	1	3	-	1	2	-	-	-	-	-	-	-
2002.....	2	3	2	2	3	1	2	-	1	-	-	-
2001.....	2	1	2	-	-	-	-	-	-	-	-	-
2000.....	1	1	1	1	-	2	1	1	-	-	-	-
1999.....	3	1	1	-	3	1	1	1	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Of other parts of central nervous system (C70,C72)												
Asian or Pacific Islander 2/, both sexes												
2004.....	-	-	1	1	-	1	-	-	-	-	-	-
2003.....	-	1	-	-	-	1	1	-	-	-	-	-
2002.....	1	-	2	-	1	-	-	-	-	-	-	-
2001.....	2	1	1	-	-	-	1	-	-	-	-	-
2000.....	-	1	-	-	-	2	-	1	-	-	-	-
1999.....	-	-	1	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	1	1	-	-	-	-	-
2002.....	1	-	1	-	1	-	-	-	-	-	-	-
2001.....	-	1	1	-	-	-	1	-	-	-	-	-
2000.....	-	1	-	-	-	1	-	-	-	-	-	-
1999.....	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	-	-	1	1	-	1	-	-	-	-	-	-
2003.....	-	1	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	1	-	-	-	-	-	-	-	-	-
2001.....	2	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	1	-	1	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant neoplasms of thyroid and other endocrine glands (C73-C75)												
All races, both sexes												
2004.....	92	154	179	214	239	251	260	185	87	21	1	-
2003.....	123	151	172	172	227	261	228	171	59	11	1	-
2002.....	118	144	166	184	236	292	250	146	76	21	1	-
2001.....	123	139	188	210	273	279	240	166	54	13	2	-
2000.....	95	141	174	219	255	286	225	166	74	19	3	2
1999.....	101	144	173	193	230	284	216	152	70	16	3	-
All races, male												
2004.....	46	77	85	104	127	101	90	55	24	4	-	-
2003.....	67	69	84	76	109	107	76	48	16	1	-	-
2002.....	61	76	86	85	102	125	92	42	10	6	-	-
2001.....	63	71	96	112	128	105	90	43	12	2	2	-
2000.....	43	67	90	109	118	118	92	46	17	1	1	2
1999.....	59	67	85	93	119	120	82	40	16	4	1	-
All races, female												
2004.....	46	77	94	110	112	150	170	130	63	17	1	-
2003.....	56	82	88	96	118	154	152	123	43	10	1	-
2002.....	57	68	80	99	134	167	158	104	66	15	1	-
2001.....	60	68	92	98	145	174	150	123	42	11	-	-
2000.....	52	74	84	110	137	168	133	120	57	18	2	-
1999.....	42	77	88	100	111	164	134	112	54	12	2	-
White, both sexes												
2004.....	76	133	155	180	213	231	234	159	79	20	1	-
2003.....	108	132	155	155	203	232	201	158	53	11	1	-
2002.....	99	125	149	164	201	266	223	133	71	17	1	-
2001.....	102	116	161	187	254	247	209	150	50	12	1	-
2000.....	69	124	144	185	220	259	204	147	70	19	3	2
1999.....	90	122	147	169	202	253	199	142	61	14	2	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasms of thyroid and other endocrine glands (C73-C75)

White, male

2004.....	40	66	75	93	114	94	85	49	20	3	-	-
2003.....	57	63	79	73	99	97	72	44	15	1	-	-
2002.....	55	65	79	77	89	117	88	38	9	5	-	-
2001.....	56	61	82	104	125	98	82	40	11	1	1	-
2000.....	34	63	72	96	109	112	87	43	17	1	1	2
1999.....	53	54	75	84	107	114	77	35	14	4	-	-

White, female

2004.....	36	67	80	87	99	137	149	110	59	17	1	-
2003.....	51	69	76	82	104	135	129	114	38	10	1	-
2002.....	44	60	70	87	112	149	135	95	62	12	1	-
2001.....	46	55	79	83	129	149	127	110	39	11	-	-
2000.....	35	61	72	89	111	147	117	104	53	18	2	-
1999.....	37	68	72	85	95	139	122	107	47	10	2	-

Black, both sexes

2004.....	10	15	18	27	12	11	18	20	5	-	-	-
2003.....	9	14	10	11	17	21	21	11	1	-	-	-
2002.....	13	14	15	15	16	20	24	9	3	4	-	-
2001.....	19	14	21	18	12	22	25	9	2	1	1	-
2000.....	22	11	22	22	23	21	13	15	4	-	-	-
1999.....	8	20	19	17	23	19	11	8	7	1	1	-

Black, male

2004.....	4	7	8	9	2	4	4	5	3	-	-	-
2003.....	5	5	4	2	5	7	4	3	1	-	-	-
2002.....	3	10	6	7	6	5	3	3	-	1	-	-
2001.....	6	7	9	6	3	4	7	2	-	1	1	-
2000.....	9	3	13	8	8	4	3	1	-	-	-	-
1999.....	4	12	7	8	10	4	4	4	-	-	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant neoplasms of thyroid and other endocrine glands (C73-C75)												
Black, female												
2004.....	6	8	10	18	10	7	14	15	2	-	-	-
2003.....	4	9	6	9	12	14	17	8	-	-	-	-
2002.....	10	4	9	8	10	15	21	6	3	3	-	-
2001.....	13	7	12	12	9	18	18	7	2	-	-	-
2000.....	13	8	9	14	15	17	10	14	4	-	-	-
1999.....	4	8	12	9	13	15	7	4	7	1	-	-
American Indian 1/, both sexes												
2004.....	2	1	1	-	1	-	1	2	1	-	-	-
2003.....	-	-	-	1	1	-	1	-	-	-	-	-
2002.....	-	1	-	-	3	-	1	-	1	-	-	-
2001.....	-	1	-	2	2	1	-	-	-	-	-	-
2000.....	-	-	3	2	3	-	1	-	-	-	-	-
1999.....	-	-	1	1	1	1	1	-	-	-	-	-
American Indian 1/, male												
2004.....	1	1	1	-	-	-	-	-	1	-	-	-
2003.....	-	-	-	-	1	-	-	-	-	-	-	-
2002.....	-	-	-	-	1	-	1	-	-	-	-	-
2001.....	-	-	-	1	-	-	-	-	-	-	-	-
2000.....	-	-	2	1	-	-	-	-	-	-	-	-
1999.....	-	-	1	-	1	-	-	-	-	-	-	-
American Indian 1/, female												
2004.....	1	-	-	-	1	-	1	2	-	-	-	-
2003.....	-	-	-	1	-	-	1	-	-	-	-	-
2002.....	-	1	-	-	2	-	-	-	1	-	-	-
2001.....	-	1	-	1	2	1	-	-	-	-	-	-
2000.....	-	-	1	1	3	-	1	-	-	-	-	-
1999.....	-	-	-	1	-	1	1	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasms of thyroid and other endocrine glands (C73-C75)

Asian or Pacific Islander 2/, both sexes

2004.....	4	5	5	7	13	9	7	4	2	1	-	-
2003.....	6	5	7	5	6	8	5	2	5	-	-	-
2002.....	6	4	2	5	16	6	2	4	1	-	-	-
2001.....	2	8	6	3	5	9	6	7	2	-	-	-
2000.....	4	6	5	10	9	6	7	4	-	-	-	-
1999.....	3	2	6	6	4	11	5	2	2	1	-	-

Asian or Pacific Islander 2/, male

2004.....	1	3	1	2	11	3	1	1	-	1	-	-
2003.....	5	1	1	1	4	3	-	1	-	-	-	-
2002.....	3	1	1	1	6	3	-	1	1	-	-	-
2001.....	1	3	5	1	-	3	1	1	1	-	-	-
2000.....	-	1	3	4	1	2	2	2	-	-	-	-
1999.....	2	1	2	1	1	2	1	1	2	-	-	-

Asian or Pacific Islander 2/, female

2004.....	3	2	4	5	2	6	6	3	2	-	-	-
2003.....	1	4	6	4	2	5	5	1	5	-	-	-
2002.....	3	3	1	4	10	3	2	3	-	-	-	-
2001.....	1	5	1	2	5	6	5	6	1	-	-	-
2000.....	4	5	2	6	8	4	5	2	-	-	-	-
1999.....	1	1	4	5	3	9	4	1	-	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other malignant neoplasms of other and unspecified sites (C76-C80,C97)												
All races, both sexes												
2004.....	2,035	2,965	3,509	4,243	5,110	6,147	6,095	4,250	2,030	523	60	2
2003.....	2,180	2,947	3,502	4,209	5,316	6,448	5,962	4,254	2,097	557	64	4
2002.....	2,114	2,818	3,605	4,216	5,596	6,337	5,983	4,343	2,113	567	71	-
2001.....	2,082	2,819	3,378	4,217	5,655	6,454	5,789	4,233	1,980	551	70	-
2000.....	2,071	2,644	3,457	4,475	5,905	6,674	5,916	4,131	1,958	579	75	2
1999.....	2,073	2,893	3,834	5,020	6,586	7,104	6,260	4,379	1,975	566	76	1
All races, male												
2004.....	1,213	1,761	2,018	2,454	2,758	3,208	2,905	1,739	688	128	13	1
2003.....	1,234	1,739	2,028	2,385	2,845	3,314	2,828	1,718	725	165	11	3
2002.....	1,286	1,592	2,118	2,409	3,071	3,233	2,790	1,804	765	156	11	-
2001.....	1,240	1,615	1,964	2,331	3,074	3,272	2,669	1,714	664	157	18	-
2000.....	1,200	1,515	1,982	2,492	3,188	3,362	2,717	1,613	676	154	12	2
1999.....	1,185	1,676	2,190	2,767	3,589	3,573	2,851	1,783	667	149	13	1
All races, female												
2004.....	822	1,204	1,491	1,789	2,352	2,939	3,190	2,511	1,342	395	47	1
2003.....	946	1,208	1,474	1,824	2,471	3,134	3,134	2,536	1,372	392	53	1
2002.....	828	1,226	1,487	1,807	2,525	3,104	3,193	2,539	1,348	411	60	-
2001.....	842	1,204	1,414	1,886	2,581	3,182	3,120	2,519	1,316	394	52	-
2000.....	871	1,129	1,475	1,983	2,717	3,312	3,199	2,518	1,282	425	63	-
1999.....	888	1,217	1,644	2,253	2,997	3,531	3,409	2,596	1,308	417	63	-
White, both sexes												
2004.....	1,590	2,405	2,922	3,565	4,444	5,451	5,484	3,897	1,854	473	53	2
2003.....	1,703	2,384	2,935	3,558	4,619	5,763	5,412	3,900	1,908	496	53	4
2002.....	1,678	2,336	2,984	3,565	4,856	5,605	5,395	3,940	1,920	510	61	-
2001.....	1,640	2,288	2,792	3,555	4,866	5,757	5,194	3,891	1,815	495	61	-
2000.....	1,632	2,194	2,861	3,847	5,134	5,893	5,379	3,773	1,812	511	57	1
1999.....	1,659	2,341	3,177	4,324	5,740	6,336	5,675	3,989	1,771	503	65	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other malignant neoplasms of other and unspecified sites (C76-C80,C97)												
White, male												
2004.....	929	1,419	1,673	2,056	2,392	2,876	2,641	1,587	627	119	11	1
2003.....	964	1,405	1,706	2,020	2,483	2,981	2,570	1,591	665	141	9	3
2002.....	1,025	1,321	1,746	2,078	2,692	2,858	2,535	1,643	702	140	10	-
2001.....	956	1,294	1,608	1,962	2,663	2,922	2,400	1,580	601	144	16	-
2000.....	941	1,243	1,604	2,158	2,792	2,938	2,490	1,461	623	137	11	1
1999.....	943	1,355	1,823	2,383	3,145	3,204	2,602	1,625	595	128	13	1
White, female												
2004.....	661	986	1,249	1,509	2,052	2,575	2,843	2,310	1,227	354	42	1
2003.....	739	979	1,229	1,538	2,136	2,782	2,842	2,309	1,243	355	44	1
2002.....	653	1,015	1,238	1,487	2,164	2,747	2,860	2,297	1,218	370	51	-
2001.....	684	994	1,184	1,593	2,203	2,835	2,794	2,311	1,214	351	45	-
2000.....	691	951	1,257	1,689	2,342	2,955	2,889	2,312	1,189	374	46	-
1999.....	716	986	1,354	1,941	2,595	3,132	3,073	2,364	1,176	375	52	-
Black, both sexes												
2004.....	377	476	500	582	561	581	503	286	147	42	7	-
2003.....	419	486	497	554	581	573	467	300	160	46	8	-
2002.....	362	425	530	554	632	629	489	340	169	48	10	-
2001.....	377	444	503	573	659	587	495	288	152	50	8	-
2000.....	372	392	512	539	646	661	456	301	117	63	15	1
1999.....	355	475	573	593	709	657	498	340	172	55	11	-
Black, male												
2004.....	250	291	299	337	314	279	214	122	49	7	2	-
2003.....	241	294	282	313	304	273	224	103	48	15	1	-
2002.....	219	239	320	287	324	317	212	133	51	13	1	-
2001.....	246	268	306	323	349	298	212	113	57	12	2	-
2000.....	224	235	326	292	331	359	190	123	39	14	1	1
1999.....	209	289	325	338	372	306	210	128	59	19	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other malignant neoplasms of other and unspecified sites (C76-C80,C97)

Black, female

2004.....	127	185	201	245	247	302	289	164	98	35	5	-
2003.....	178	192	215	241	277	300	243	197	112	31	7	-
2002.....	143	186	210	267	308	312	277	207	118	35	9	-
2001.....	131	176	197	250	310	289	283	175	95	38	6	-
2000.....	148	157	186	247	315	302	266	178	78	49	14	-
1999.....	146	186	248	255	337	351	288	212	113	36	11	-

American Indian 1/, both sexes

2004.....	16	22	21	20	22	27	17	10	3	1	-	-
2003.....	18	20	15	18	22	28	14	10	2	7	1	-
2002.....	15	16	23	14	24	25	12	13	8	1	-	-
2001.....	12	25	17	21	27	22	18	12	4	1	-	-
2000.....	13	11	23	17	33	24	13	13	6	2	2	-
1999.....	13	23	24	26	30	21	23	8	3	-	-	-

American Indian 1/, male

2004.....	7	13	11	15	12	12	5	5	-	-	-	-
2003.....	10	13	9	9	12	15	6	5	1	5	-	-
2002.....	9	5	12	10	15	11	4	3	2	-	-	-
2001.....	8	14	8	11	18	9	8	4	1	-	-	-
2000.....	7	8	13	8	18	10	4	4	3	1	-	-
1999.....	6	9	11	13	15	13	9	3	1	-	-	-

American Indian 1/, female

2004.....	9	9	10	5	10	15	12	5	3	1	-	-
2003.....	8	7	6	9	10	13	8	5	1	2	1	-
2002.....	6	11	11	4	9	14	8	10	6	1	-	-
2001.....	4	11	9	10	9	13	10	8	3	1	-	-
2000.....	6	3	10	9	15	14	9	9	3	1	2	-
1999.....	7	14	13	13	15	8	14	5	2	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other malignant neoplasms of other and unspecified sites (C76-C80,C97)												
Asian or Pacific Islander 2/, both sexes												
2004.....	52	62	66	76	83	88	91	57	26	7	-	-
2003.....	40	57	55	79	94	84	69	44	27	8	2	-
2002.....	59	41	68	83	84	78	87	50	16	8	-	-
2001.....	53	62	66	68	103	88	82	42	9	5	1	-
2000.....	54	47	61	72	92	96	68	44	23	3	1	-
1999.....	46	54	60	77	107	90	64	42	29	8	-	-
Asian or Pacific Islander 2/, male												
2004.....	27	38	35	46	40	41	45	25	12	2	-	-
2003.....	19	27	31	43	46	45	28	19	11	4	1	-
2002.....	33	27	40	34	40	47	39	25	10	3	-	-
2001.....	30	39	42	35	44	43	49	17	5	1	-	-
2000.....	28	29	39	34	47	55	33	25	11	2	-	-
1999.....	27	23	31	33	57	50	30	27	12	2	-	-
Asian or Pacific Islander 2/, female												
2004.....	25	24	31	30	43	47	46	32	14	5	-	-
2003.....	21	30	24	36	48	39	41	25	16	4	1	-
2002.....	26	14	28	49	44	31	48	25	6	5	-	-
2001.....	23	23	24	33	59	45	33	25	4	4	1	-
2000.....	26	18	22	38	45	41	35	19	12	1	1	-
1999.....	19	31	29	44	50	40	34	15	17	6	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)												
All races, both sexes												
2004.....	2,264	3,323	4,168	5,423	7,165	9,090	9,042	5,925	2,633	577	71	-
2003.....	2,228	3,171	4,300	5,384	7,501	9,497	8,945	6,087	2,541	644	62	-
2002.....	2,361	3,217	4,283	5,672	7,836	9,319	8,970	6,003	2,433	574	62	-
2001.....	2,390	3,224	4,231	5,602	8,057	9,580	8,677	5,848	2,344	536	77	-
2000.....	2,350	3,156	4,128	5,932	8,109	9,743	8,705	5,716	2,329	559	70	-
1999.....	2,302	3,346	4,208	5,996	8,336	9,596	8,493	5,468	2,233	526	55	-
All races, male												
2004.....	1,372	1,993	2,504	3,244	4,124	4,915	4,547	2,661	995	171	16	-
2003.....	1,347	1,929	2,628	3,255	4,295	5,068	4,456	2,722	943	202	16	-
2002.....	1,388	1,947	2,523	3,391	4,512	5,047	4,506	2,624	906	186	24	-
2001.....	1,440	1,900	2,512	3,316	4,543	5,065	4,262	2,537	846	168	25	-
2000.....	1,384	1,850	2,355	3,445	4,593	5,168	4,231	2,475	783	185	15	-
1999.....	1,382	1,957	2,469	3,436	4,631	5,102	3,998	2,337	816	155	21	-
All races, female												
2004.....	892	1,330	1,664	2,179	3,041	4,175	4,495	3,264	1,638	406	55	-
2003.....	881	1,242	1,672	2,129	3,206	4,429	4,489	3,365	1,598	442	46	-
2002.....	973	1,270	1,760	2,281	3,324	4,272	4,464	3,379	1,527	388	38	-
2001.....	950	1,324	1,719	2,286	3,514	4,515	4,415	3,311	1,498	368	52	-
2000.....	966	1,306	1,773	2,487	3,516	4,575	4,474	3,241	1,546	374	55	-
1999.....	920	1,389	1,739	2,560	3,705	4,494	4,495	3,131	1,417	371	34	-
White, both sexes												
2004.....	1,824	2,789	3,579	4,705	6,290	8,234	8,348	5,519	2,423	536	62	-
2003.....	1,797	2,710	3,719	4,701	6,661	8,633	8,259	5,697	2,368	587	53	-
2002.....	1,909	2,692	3,674	4,970	7,007	8,512	8,324	5,619	2,262	534	54	-
2001.....	1,980	2,704	3,614	4,856	7,252	8,751	8,040	5,451	2,187	494	72	-
2000.....	1,957	2,693	3,562	5,251	7,307	8,944	8,129	5,292	2,145	511	59	-
1999.....	1,919	2,836	3,604	5,297	7,529	8,778	7,909	5,093	2,065	483	47	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)												
White, male												
2004.....	1,115	1,702	2,178	2,828	3,663	4,504	4,230	2,502	919	160	13	-
2003.....	1,094	1,657	2,310	2,891	3,892	4,690	4,167	2,571	881	185	15	-
2002.....	1,149	1,660	2,183	3,007	4,087	4,661	4,209	2,477	840	176	20	-
2001.....	1,209	1,607	2,163	2,910	4,130	4,672	3,969	2,381	791	154	24	-
2000.....	1,150	1,591	2,048	3,103	4,146	4,784	3,976	2,309	725	169	14	-
1999.....	1,168	1,673	2,144	3,068	4,219	4,683	3,761	2,176	750	142	17	-
White, female												
2004.....	709	1,087	1,401	1,877	2,627	3,730	4,118	3,017	1,504	376	49	-
2003.....	703	1,053	1,409	1,810	2,769	3,943	4,092	3,126	1,487	402	38	-
2002.....	760	1,032	1,491	1,963	2,920	3,851	4,115	3,142	1,422	358	34	-
2001.....	771	1,097	1,451	1,946	3,122	4,079	4,071	3,070	1,396	340	48	-
2000.....	807	1,102	1,514	2,148	3,161	4,160	4,153	2,983	1,420	342	45	-
1999.....	751	1,163	1,460	2,229	3,310	4,095	4,148	2,917	1,315	341	30	-
Black, both sexes												
2004.....	375	441	499	588	713	684	544	334	168	35	9	-
2003.....	374	372	486	566	684	690	548	317	148	43	8	-
2002.....	363	444	505	587	695	651	528	326	146	34	7	-
2001.....	332	429	501	628	674	672	527	330	135	38	4	-
2000.....	328	393	482	563	678	643	466	356	159	38	8	-
1999.....	322	435	492	585	668	667	482	319	142	39	8	-
Black, male												
2004.....	219	239	274	332	379	306	235	128	55	9	3	-
2003.....	223	228	265	298	320	302	226	117	49	11	-	-
2002.....	190	242	281	316	369	300	232	123	52	10	4	-
2001.....	194	240	280	346	339	304	238	124	45	12	1	-
2000.....	188	223	262	281	375	313	199	132	49	13	1	-
1999.....	183	243	260	311	338	335	189	123	51	10	4	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)												
Black, female												
2004.....	156	202	225	256	334	378	309	206	113	26	6	-
2003.....	151	144	221	268	364	388	322	200	99	32	8	-
2002.....	173	202	224	271	326	351	296	203	94	24	3	-
2001.....	138	189	221	282	335	368	289	206	90	26	3	-
2000.....	140	170	220	282	303	330	267	224	110	25	7	-
1999.....	139	192	232	274	330	332	293	196	91	29	4	-
American Indian 1/, both sexes												
2004.....	11	24	15	29	19	26	26	9	4	1	-	-
2003.....	10	23	18	19	29	29	21	11	4	1	1	-
2002.....	12	16	19	17	17	23	13	6	4	2	-	-
2001.....	17	21	25	34	21	19	17	10	6	1	-	-
2000.....	11	15	19	22	24	25	23	13	7	4	-	-
1999.....	8	9	14	22	28	26	11	8	5	-	-	-
American Indian 1/, male												
2004.....	8	13	9	21	8	16	13	3	-	-	-	-
2003.....	6	9	9	12	13	7	9	4	1	-	1	-
2002.....	8	7	9	9	8	11	8	2	1	-	-	-
2001.....	8	11	15	17	11	7	6	4	1	-	-	-
2000.....	8	9	8	10	16	10	11	5	1	2	-	-
1999.....	5	4	8	15	14	18	3	5	3	-	-	-
American Indian 1/, female												
2004.....	3	11	6	8	11	10	13	6	4	1	-	-
2003.....	4	14	9	7	16	22	12	7	3	1	-	-
2002.....	4	9	10	8	9	12	5	4	3	2	-	-
2001.....	9	10	10	17	10	12	11	6	5	1	-	-
2000.....	3	6	11	12	8	15	12	8	6	2	-	-
1999.....	3	5	6	7	14	8	8	3	2	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)												
Asian or Pacific Islander 2/, both sexes												
2004.....	54	69	75	101	143	146	124	63	38	5	-	-
2003.....	47	66	77	98	127	145	117	62	21	13	-	-
2002.....	77	65	85	98	117	133	105	52	21	4	1	-
2001.....	61	70	91	84	110	138	93	57	16	3	1	-
2000.....	54	55	65	96	100	131	87	55	18	6	3	-
1999.....	53	66	98	92	111	125	91	48	21	4	-	-
Asian or Pacific Islander 2/, male												
2004.....	30	39	43	63	74	89	69	28	21	2	-	-
2003.....	24	35	44	54	70	69	54	30	12	6	-	-
2002.....	41	38	50	59	48	75	57	22	13	-	-	-
2001.....	29	42	54	43	63	82	49	28	9	2	-	-
2000.....	38	27	37	51	56	61	45	29	8	1	-	-
1999.....	26	37	57	42	60	66	45	33	12	3	-	-
Asian or Pacific Islander 2/, female												
2004.....	24	30	32	38	69	57	55	35	17	3	-	-
2003.....	23	31	33	44	57	76	63	32	9	7	-	-
2002.....	36	27	35	39	69	58	48	30	8	4	1	-
2001.....	32	28	37	41	47	56	44	29	7	1	1	-
2000.....	16	28	28	45	44	70	42	26	10	5	3	-
1999.....	27	29	41	50	51	59	46	15	9	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hodgkin's disease (C81)												
All races, both sexes												
2004.....	81	110	95	104	118	131	118	81	24	2	-	-
2003.....	83	98	100	83	133	153	124	75	34	3	-	-
2002.....	94	74	94	94	138	126	121	83	35	7	-	-
2001.....	91	61	82	103	108	151	127	66	24	7	-	-
2000.....	88	64	94	92	140	134	121	70	24	10	-	-
1999.....	83	87	91	120	163	149	113	70	22	5	1	-
All races, male												
2004.....	51	71	61	64	72	62	67	41	8	-	-	-
2003.....	47	57	62	52	76	73	59	31	7	-	-	-
2002.....	52	53	53	62	84	60	65	37	11	2	-	-
2001.....	56	40	45	65	64	75	55	26	7	1	-	-
2000.....	53	30	57	58	76	71	50	30	10	5	-	-
1999.....	50	54	61	72	88	69	50	28	16	1	-	-
All races, female												
2004.....	30	39	34	40	46	69	51	40	16	2	-	-
2003.....	36	41	38	31	57	80	65	44	27	3	-	-
2002.....	42	21	41	32	54	66	56	46	24	5	-	-
2001.....	35	21	37	38	44	76	72	40	17	6	-	-
2000.....	35	34	37	34	64	63	71	40	14	5	-	-
1999.....	33	33	30	48	75	80	63	42	6	4	1	-
White, both sexes												
2004.....	70	97	83	91	110	124	112	76	22	2	-	-
2003.....	75	79	91	71	125	143	115	71	33	3	-	-
2002.....	79	61	80	85	128	119	115	81	33	6	-	-
2001.....	84	52	72	99	101	138	122	64	24	7	-	-
2000.....	75	56	82	86	125	134	114	68	22	9	-	-
1999.....	74	78	85	106	154	144	110	62	21	5	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hodgkin's disease (C81)												
White, male												
2004.....	46	63	53	54	68	58	65	40	8	-	-	-
2003.....	41	45	56	43	69	67	56	28	7	-	-	-
2002.....	41	43	43	57	79	57	63	37	10	2	-	-
2001.....	51	35	39	63	61	70	52	24	7	1	-	-
2000.....	45	26	48	55	67	71	47	29	9	5	-	-
1999.....	42	49	58	64	84	67	49	25	15	1	-	-
White, female												
2004.....	24	34	30	37	42	66	47	36	14	2	-	-
2003.....	34	34	35	28	56	76	59	43	26	3	-	-
2002.....	38	18	37	28	49	62	52	44	23	4	-	-
2001.....	33	17	33	36	40	68	70	40	17	6	-	-
2000.....	30	30	34	31	58	63	67	39	13	4	-	-
1999.....	32	29	27	42	70	77	61	37	6	4	1	-
Black, both sexes												
2004.....	10	10	9	10	8	6	5	2	1	-	-	-
2003.....	8	15	6	11	6	6	3	-	-	-	-	-
2002.....	10	10	11	6	9	5	4	1	1	1	-	-
2001.....	5	9	9	4	7	9	3	2	-	-	-	-
2000.....	10	8	9	4	11	-	5	1	2	-	-	-
1999.....	9	9	5	11	7	5	1	6	1	-	-	-
Black, male												
2004.....	5	6	6	7	4	3	2	-	-	-	-	-
2003.....	6	10	5	8	5	3	1	-	-	-	-	-
2002.....	7	8	8	3	5	3	1	-	1	-	-	-
2001.....	4	5	5	2	3	3	2	2	-	-	-	-
2000.....	5	4	7	2	5	-	3	-	1	-	-	-
1999.....	8	5	3	8	4	2	1	1	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hodgkin's disease (C81)												
Black, female												
2004.....	5	4	3	3	4	3	3	2	1	-	-	-
2003.....	2	5	1	3	1	3	2	-	-	-	-	-
2002.....	3	2	3	3	4	2	3	1	-	1	-	-
2001.....	1	4	4	2	4	6	1	-	-	-	-	-
2000.....	5	4	2	2	6	-	2	1	1	-	-	-
1999.....	1	4	2	3	3	3	-	5	-	-	-	-
American Indian 1/, both sexes												
2004.....	-	-	-	1	-	-	-	-	-	-	-	-
2003.....	-	2	-	1	-	1	3	-	1	-	-	-
2002.....	-	-	1	-	-	-	-	-	-	-	-	-
2001.....	1	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	1	1	3	-	-	-	-	-	-	-
1999.....	-	-	-	2	1	-	-	1	-	-	-	-
American Indian 1/, male												
2004.....	-	-	-	1	-	-	-	-	-	-	-	-
2003.....	-	1	-	1	-	1	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	1	3	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, female												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	1	-	-	-	-	3	-	1	-	-	-
2002.....	-	-	1	-	-	-	-	-	-	-	-	-
2001.....	1	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	1	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	2	1	-	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hodgkin's disease (C81)												
Asian or Pacific Islander 2/, both sexes												
2004.....	1	3	3	2	-	1	1	3	1	-	-	-
2003.....	-	2	3	-	2	3	3	4	-	-	-	-
2002.....	5	3	2	3	1	2	2	1	1	-	-	-
2001.....	1	-	1	-	-	4	2	-	-	-	-	-
2000.....	3	-	2	1	1	-	2	1	-	1	-	-
1999.....	-	-	1	1	1	-	2	1	-	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	-	2	2	2	-	1	-	1	-	-	-	-
2003.....	-	1	1	-	2	2	2	3	-	-	-	-
2002.....	4	2	2	2	-	-	1	-	-	-	-	-
2001.....	1	-	1	-	-	2	1	-	-	-	-	-
2000.....	3	-	2	-	1	-	-	1	-	-	-	-
1999.....	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	1	1	1	-	-	-	1	2	1	-	-	-
2003.....	-	1	2	-	-	1	1	1	-	-	-	-
2002.....	1	1	-	1	1	2	1	1	1	-	-	-
2001.....	-	-	-	-	-	2	1	-	-	-	-	-
2000.....	-	-	-	1	-	-	2	-	-	1	-	-
1999.....	-	-	1	1	1	-	2	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Of other lymphoid, hematopoietic and related tissue (C82-C85,C96)												
All races, both sexes												
2004.....	931	1,285	1,593	2,105	2,727	3,486	3,620	2,365	1,041	229	24	-
2003.....	902	1,273	1,736	2,110	2,824	3,685	3,546	2,530	952	213	30	-
2002.....	948	1,321	1,731	2,232	3,050	3,734	3,574	2,455	886	202	22	-
2001.....	1,024	1,369	1,737	2,204	3,177	3,787	3,610	2,403	912	187	24	-
2000.....	1,018	1,440	1,742	2,393	3,227	3,997	3,567	2,343	863	194	23	-
1999.....	1,051	1,420	1,746	2,451	3,414	4,023	3,556	2,207	838	178	18	-
All races, male												
2004.....	581	787	943	1,235	1,507	1,785	1,737	1,016	380	50	6	-
2003.....	559	774	1,076	1,263	1,575	1,889	1,682	1,069	346	78	9	-
2002.....	576	824	1,016	1,339	1,662	1,902	1,704	1,018	316	52	6	-
2001.....	665	818	1,012	1,294	1,749	1,878	1,694	1,024	307	55	5	-
2000.....	648	870	988	1,349	1,768	2,028	1,623	965	277	62	4	-
1999.....	655	856	998	1,360	1,824	2,061	1,549	895	292	53	9	-
All races, female												
2004.....	350	498	650	870	1,220	1,701	1,883	1,349	661	179	18	-
2003.....	343	499	660	847	1,249	1,796	1,864	1,461	606	135	21	-
2002.....	372	497	715	893	1,388	1,832	1,870	1,437	570	150	16	-
2001.....	359	551	725	910	1,428	1,909	1,916	1,379	605	132	19	-
2000.....	370	570	754	1,044	1,459	1,969	1,944	1,378	586	132	19	-
1999.....	396	564	748	1,091	1,590	1,962	2,007	1,312	546	125	9	-
White, both sexes												
2004.....	770	1,119	1,416	1,900	2,488	3,242	3,440	2,247	979	216	20	-
2003.....	757	1,139	1,544	1,912	2,580	3,443	3,363	2,430	907	191	29	-
2002.....	781	1,152	1,550	2,021	2,795	3,506	3,387	2,356	849	194	21	-
2001.....	880	1,188	1,549	1,977	2,947	3,563	3,412	2,277	863	179	22	-
2000.....	882	1,295	1,570	2,186	2,996	3,772	3,411	2,224	823	186	21	-
1999.....	927	1,256	1,573	2,259	3,173	3,771	3,409	2,097	791	165	16	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Of other lymphoid, hematopoietic and related tissue (C82-C85,C96)

White, male

2004.....	489	686	843	1,113	1,384	1,672	1,657	970	361	48	5	-
2003.....	474	699	967	1,166	1,463	1,784	1,607	1,034	327	69	9	-
2002.....	483	727	912	1,222	1,537	1,798	1,620	981	299	48	6	-
2001.....	572	710	903	1,173	1,625	1,770	1,607	968	291	52	5	-
2000.....	558	775	891	1,251	1,635	1,926	1,555	922	262	59	4	-
1999.....	578	755	894	1,253	1,708	1,919	1,499	842	273	47	7	-

White, female

2004.....	281	433	573	787	1,104	1,570	1,783	1,277	618	168	15	-
2003.....	283	440	577	746	1,117	1,659	1,756	1,396	580	122	20	-
2002.....	298	425	638	799	1,258	1,708	1,767	1,375	550	146	15	-
2001.....	308	478	646	804	1,322	1,793	1,805	1,309	572	127	17	-
2000.....	324	520	679	935	1,361	1,846	1,856	1,302	561	127	17	-
1999.....	349	501	679	1,006	1,465	1,852	1,910	1,255	518	118	9	-

Black, both sexes

2004.....	132	130	134	154	166	172	109	83	44	8	4	-
2003.....	127	96	150	140	172	156	122	65	29	13	1	-
2002.....	128	139	147	167	192	158	136	74	26	5	-	-
2001.....	119	141	138	178	175	149	143	84	36	7	2	-
2000.....	103	120	138	150	170	148	101	81	31	6	-	-
1999.....	100	139	126	154	177	180	109	80	35	11	2	-

Black, male

2004.....	80	76	77	81	88	70	48	30	11	1	1	-
2003.....	74	55	83	70	75	63	47	18	11	4	-	-
2002.....	71	77	84	93	103	69	55	27	9	4	-	-
2001.....	80	79	76	98	94	67	60	36	11	2	-	-
2000.....	68	83	76	74	96	66	38	24	13	3	-	-
1999.....	65	82	74	90	84	100	33	33	11	5	2	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Of other lymphoid, hematopoietic and related tissue (C82-C85,C96)

Black, female

2004.....	52	54	57	73	78	102	61	53	33	7	3	-
2003.....	53	41	67	70	97	93	75	47	18	9	1	-
2002.....	57	62	63	74	89	89	81	47	17	1	-	-
2001.....	39	62	62	80	81	82	83	48	25	5	2	-
2000.....	35	37	62	76	74	82	63	57	18	3	-	-
1999.....	35	57	52	64	93	80	76	47	24	6	-	-

American Indian 1/, both sexes

2004.....	2	6	8	14	7	10	6	4	-	1	-	-
2003.....	1	11	6	9	11	10	5	1	1	1	-	-
2002.....	4	4	7	4	7	7	3	1	2	1	-	-
2001.....	6	6	11	18	6	12	8	4	4	-	-	-
2000.....	7	4	6	11	11	11	15	4	1	-	-	-
1999.....	1	3	6	7	11	8	4	3	2	-	-	-

American Indian 1/, male

2004.....	1	5	4	12	4	6	1	2	-	-	-	-
2003.....	-	6	4	4	6	3	3	-	-	-	-	-
2002.....	2	2	5	3	3	4	-	-	-	-	-	-
2001.....	2	4	7	9	4	5	2	2	1	-	-	-
2000.....	5	2	3	3	6	6	8	1	-	-	-	-
1999.....	1	2	4	4	3	7	1	1	1	-	-	-

American Indian 1/, female

2004.....	1	1	4	2	3	4	5	2	-	1	-	-
2003.....	1	5	2	5	5	7	2	1	1	1	-	-
2002.....	2	2	2	1	4	3	3	1	2	1	-	-
2001.....	4	2	4	9	2	7	6	2	3	-	-	-
2000.....	2	2	3	8	5	5	7	3	1	-	-	-
1999.....	-	1	2	3	8	1	3	2	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Of other lymphoid, hematopoietic and related tissue (C82-C85,C96)												
Asian or Pacific Islander 2/, both sexes												
2004.....	27	30	35	37	66	62	65	31	18	4	-	-
2003.....	17	27	36	49	61	76	56	34	15	8	-	-
2002.....	35	26	27	40	56	63	48	24	9	2	1	-
2001.....	19	34	39	31	49	63	47	38	9	1	-	-
2000.....	26	21	28	46	50	66	40	34	8	2	2	-
1999.....	23	22	41	31	53	64	34	27	10	2	-	-
Asian or Pacific Islander 2/, male												
2004.....	11	20	19	29	31	37	31	14	8	1	-	-
2003.....	11	14	22	23	31	39	25	17	8	5	-	-
2002.....	20	18	15	21	19	31	29	10	8	-	-	-
2001.....	11	25	26	14	26	36	25	18	4	1	-	-
2000.....	17	10	18	21	31	30	22	18	2	-	-	-
1999.....	11	17	26	13	29	35	16	19	7	1	-	-
Asian or Pacific Islander 2/, female												
2004.....	16	10	16	8	35	25	34	17	10	3	-	-
2003.....	6	13	14	26	30	37	31	17	7	3	-	-
2002.....	15	8	12	19	37	32	19	14	1	2	1	-
2001.....	8	9	13	17	23	27	22	20	5	-	-	-
2000.....	9	11	10	25	19	36	18	16	6	2	2	-
1999.....	12	5	15	18	24	29	18	8	3	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Non-Hodgkin's lymphoma (C82-C85)												
All races, both sexes												
2004.....	927	1,277	1,589	2,100	2,720	3,477	3,603	2,359	1,036	228	24	-
2003.....	899	1,272	1,735	2,104	2,815	3,674	3,543	2,520	949	213	30	-
2002.....	943	1,319	1,730	2,225	3,039	3,722	3,561	2,446	881	200	22	-
2001.....	1,021	1,367	1,731	2,193	3,165	3,776	3,593	2,392	904	186	24	-
2000.....	1,015	1,434	1,736	2,386	3,212	3,984	3,551	2,330	856	193	23	-
1999.....	1,045	1,414	1,740	2,441	3,407	4,009	3,545	2,198	834	178	18	-
All races, male												
2004.....	578	784	941	1,235	1,503	1,782	1,729	1,012	378	49	6	-
2003.....	558	773	1,075	1,260	1,569	1,882	1,681	1,065	345	78	9	-
2002.....	574	823	1,015	1,334	1,652	1,898	1,696	1,013	315	51	6	-
2001.....	663	817	1,008	1,286	1,739	1,873	1,685	1,020	305	55	5	-
2000.....	646	865	984	1,342	1,763	2,018	1,616	959	274	62	4	-
1999.....	652	851	996	1,355	1,821	2,052	1,544	892	292	53	9	-
All races, female												
2004.....	349	493	648	865	1,217	1,695	1,874	1,347	658	179	18	-
2003.....	341	499	660	844	1,246	1,792	1,862	1,455	604	135	21	-
2002.....	369	496	715	891	1,387	1,824	1,865	1,433	566	149	16	-
2001.....	358	550	723	907	1,426	1,903	1,908	1,372	599	131	19	-
2000.....	369	569	752	1,044	1,449	1,966	1,935	1,371	582	131	19	-
1999.....	393	563	744	1,086	1,586	1,957	2,001	1,306	542	125	9	-
White, both sexes												
2004.....	767	1,113	1,412	1,895	2,481	3,234	3,426	2,241	975	216	20	-
2003.....	754	1,138	1,543	1,907	2,571	3,432	3,361	2,421	904	191	29	-
2002.....	777	1,150	1,549	2,015	2,784	3,497	3,376	2,347	847	192	21	-
2001.....	877	1,188	1,544	1,967	2,936	3,554	3,398	2,267	855	178	22	-
2000.....	879	1,290	1,564	2,179	2,982	3,761	3,396	2,212	817	185	21	-
1999.....	922	1,251	1,567	2,249	3,166	3,758	3,399	2,089	787	165	16	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Non-Hodgkin's lymphoma (C82-C85)												
White, male												
2004.....	486	683	841	1,113	1,380	1,669	1,650	966	359	48	5	-
2003.....	473	698	966	1,163	1,457	1,777	1,607	1,030	326	69	9	-
2002.....	481	726	911	1,218	1,527	1,794	1,614	976	299	47	6	-
2001.....	570	710	899	1,166	1,615	1,766	1,599	965	289	52	5	-
2000.....	556	771	887	1,244	1,631	1,917	1,548	917	260	59	4	-
1999.....	576	751	892	1,248	1,705	1,911	1,494	839	273	47	7	-
White, female												
2004.....	281	430	571	782	1,101	1,565	1,776	1,275	616	168	15	-
2003.....	281	440	577	744	1,114	1,655	1,754	1,391	578	122	20	-
2002.....	296	424	638	797	1,257	1,703	1,762	1,371	548	145	15	-
2001.....	307	478	645	801	1,321	1,788	1,799	1,302	566	126	17	-
2000.....	323	519	677	935	1,351	1,844	1,848	1,295	557	126	17	-
1999.....	346	500	675	1,001	1,461	1,847	1,905	1,250	514	118	9	-
Black, both sexes												
2004.....	131	128	134	154	166	171	107	83	43	7	4	-
2003.....	127	96	150	139	172	156	122	64	29	13	1	-
2002.....	128	139	147	166	192	156	135	74	24	5	-	-
2001.....	119	139	137	177	174	148	140	83	36	7	2	-
2000.....	103	119	138	150	170	146	100	80	30	6	-	-
1999.....	99	138	126	154	177	179	109	79	35	11	2	-
Black, male												
2004.....	80	76	77	81	88	70	48	30	11	-	1	-
2003.....	74	55	83	70	75	63	47	18	11	4	-	-
2002.....	71	77	84	92	103	69	54	27	9	4	-	-
2001.....	80	78	76	97	94	66	59	35	11	2	-	-
2000.....	68	82	76	74	96	65	38	23	12	3	-	-
1999.....	64	81	74	90	84	99	33	33	11	5	2	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Non-Hodgkin's lymphoma (C82-C85)												
Black, female												
2004.....	51	52	57	73	78	101	59	53	32	7	3	-
2003.....	53	41	67	69	97	93	75	46	18	9	1	-
2002.....	57	62	63	74	89	87	81	47	15	1	-	-
2001.....	39	61	61	80	80	82	81	48	25	5	2	-
2000.....	35	37	62	76	74	81	62	57	18	3	-	-
1999.....	35	57	52	64	93	80	76	46	24	6	-	-
American Indian 1/, both sexes												
2004.....	2	6	8	14	7	10	6	4	-	1	-	-
2003.....	1	11	6	9	11	10	5	1	1	1	-	-
2002.....	4	4	7	4	7	6	3	1	2	1	-	-
2001.....	6	6	11	18	6	11	8	4	4	-	-	-
2000.....	7	4	6	11	10	11	15	4	1	-	-	-
1999.....	1	3	6	7	11	8	4	3	2	-	-	-
American Indian 1/, male												
2004.....	1	5	4	12	4	6	1	2	-	-	-	-
2003.....	-	6	4	4	6	3	3	-	-	-	-	-
2002.....	2	2	5	3	3	4	-	-	-	-	-	-
2001.....	2	4	7	9	4	5	2	2	1	-	-	-
2000.....	5	2	3	3	5	6	8	1	-	-	-	-
1999.....	1	2	4	4	3	7	1	1	1	-	-	-
American Indian 1/, female												
2004.....	1	1	4	2	3	4	5	2	-	1	-	-
2003.....	1	5	2	5	5	7	2	1	1	1	-	-
2002.....	2	2	2	1	4	2	3	1	2	1	-	-
2001.....	4	2	4	9	2	6	6	2	3	-	-	-
2000.....	2	2	3	8	5	5	7	3	1	-	-	-
1999.....	-	1	2	3	8	1	3	2	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Non-Hodgkin's lymphoma (C82-C85)												
Asian or Pacific Islander 2/, both sexes												
2004.....	27	30	35	37	66	62	64	31	18	4	-	-
2003.....	17	27	36	49	61	76	55	34	15	8	-	-
2002.....	34	26	27	40	56	63	47	24	8	2	1	-
2001.....	19	34	39	31	49	63	47	38	9	1	-	-
2000.....	26	21	28	46	50	66	40	34	8	2	2	-
1999.....	23	22	41	31	53	64	33	27	10	2	-	-
Asian or Pacific Islander 2/, male												
2004.....	11	20	19	29	31	37	30	14	8	1	-	-
2003.....	11	14	22	23	31	39	24	17	8	5	-	-
2002.....	20	18	15	21	19	31	28	10	7	-	-	-
2001.....	11	25	26	14	26	36	25	18	4	1	-	-
2000.....	17	10	18	21	31	30	22	18	2	-	-	-
1999.....	11	17	26	13	29	35	16	19	7	1	-	-
Asian or Pacific Islander 2/, female												
2004.....	16	10	16	8	35	25	34	17	10	3	-	-
2003.....	6	13	14	26	30	37	31	17	7	3	-	-
2002.....	14	8	12	19	37	32	19	14	1	2	1	-
2001.....	8	9	13	17	23	27	22	20	5	-	-	-
2000.....	9	11	10	25	19	36	18	16	6	2	2	-
1999.....	12	5	15	18	24	29	17	8	3	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Of other and unspecified lymphoid, hematopoietic and related tissue (C96)												
All races, both sexes												
2004.....	4	8	4	5	7	9	17	6	5	1	-	-
2003.....	3	1	1	6	9	11	3	10	3	-	-	-
2002.....	5	2	1	7	11	12	13	9	5	2	-	-
2001.....	3	2	6	11	12	11	17	11	8	1	-	-
2000.....	3	6	6	7	15	13	16	13	7	1	-	-
1999.....	6	6	6	10	7	14	11	9	4	-	-	-
All races, male												
2004.....	3	3	2	-	4	3	8	4	2	1	-	-
2003.....	1	1	1	3	6	7	1	4	1	-	-	-
2002.....	2	1	1	5	10	4	8	5	1	1	-	-
2001.....	2	1	4	8	10	5	9	4	2	-	-	-
2000.....	2	5	4	7	5	10	7	6	3	-	-	-
1999.....	3	5	2	5	3	9	5	3	-	-	-	-
All races, female												
2004.....	1	5	2	5	3	6	9	2	3	-	-	-
2003.....	2	-	-	3	3	4	2	6	2	-	-	-
2002.....	3	1	-	2	1	8	5	4	4	1	-	-
2001.....	1	1	2	3	2	6	8	7	6	1	-	-
2000.....	1	1	2	-	10	3	9	7	4	1	-	-
1999.....	3	1	4	5	4	5	6	6	4	-	-	-
White, both sexes												
2004.....	3	6	4	5	7	8	14	6	4	-	-	-
2003.....	3	1	1	5	9	11	2	9	3	-	-	-
2002.....	4	2	1	6	11	9	11	9	2	2	-	-
2001.....	3	-	5	10	11	9	14	10	8	1	-	-
2000.....	3	5	6	7	14	11	15	12	6	1	-	-
1999.....	5	5	6	10	7	13	10	8	4	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Of other and unspecified lymphoid, hematopoietic and related tissue (C96)

White, male

2004.....	3	3	2	-	4	3	7	4	2	-	-	-
2003.....	1	1	1	3	6	7	-	4	1	-	-	-
2002.....	2	1	1	4	10	4	6	5	-	1	-	-
2001.....	2	-	4	7	10	4	8	3	2	-	-	-
2000.....	2	4	4	7	4	9	7	5	2	-	-	-
1999.....	2	4	2	5	3	8	5	3	-	-	-	-

White, female

2004.....	-	3	2	5	3	5	7	2	2	-	-	-
2003.....	2	-	-	2	3	4	2	5	2	-	-	-
2002.....	2	1	-	2	1	5	5	4	2	1	-	-
2001.....	1	-	1	3	1	5	6	7	6	1	-	-
2000.....	1	1	2	-	10	2	8	7	4	1	-	-
1999.....	3	1	4	5	4	5	5	5	4	-	-	-

Black, both sexes

2004.....	1	2	-	-	-	1	2	-	1	1	-	-
2003.....	-	-	-	1	-	-	-	1	-	-	-	-
2002.....	-	-	-	1	-	2	1	-	2	-	-	-
2001.....	-	2	1	1	1	1	3	1	-	-	-	-
2000.....	-	1	-	-	-	2	1	1	1	-	-	-
1999.....	1	1	-	-	-	1	-	1	-	-	-	-

Black, male

2004.....	-	-	-	-	-	-	-	-	-	1	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	1	-	-	1	-	-	-	-	-
2001.....	-	1	-	1	-	1	1	1	-	-	-	-
2000.....	-	1	-	-	-	1	-	1	1	-	-	-
1999.....	1	1	-	-	-	1	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Of other and unspecified lymphoid, hematopoietic and related tissue (C96)

Asian or Pacific Islander 2/, both sexes

2004.....	-	-	-	-	-	-	1	-	-	-	-	-
2003.....	-	-	-	-	-	-	1	-	-	-	-	-
2002.....	1	-	-	-	-	-	1	-	1	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	1	-	-	-	-	-

Asian or Pacific Islander 2/, male

2004.....	-	-	-	-	-	-	1	-	-	-	-	-
2003.....	-	-	-	-	-	-	1	-	-	-	-	-
2002.....	-	-	-	-	-	-	1	-	1	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-

Asian or Pacific Islander 2/, female

2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	1	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	1	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant immunoproliferative diseases (C88)												
All races, both sexes												
2004.....	4	7	16	26	48	67	60	48	22	4	1	-
2003.....	9	12	20	23	65	88	60	67	33	4	-	-
2002.....	10	11	29	26	48	86	90	53	32	6	-	-
2001.....	6	17	23	40	51	63	81	57	27	5	1	-
2000.....	6	17	19	27	51	65	86	64	21	5	-	-
1999.....	10	15	40	33	64	90	79	64	39	9	-	-
All races, male												
2004.....	3	5	10	16	32	37	33	26	10	2	-	-
2003.....	7	6	14	14	45	53	35	32	19	1	-	-
2002.....	5	7	23	20	33	53	58	25	12	2	-	-
2001.....	4	10	14	27	26	35	43	25	11	1	1	-
2000.....	4	9	12	19	31	41	44	32	9	1	-	-
1999.....	6	9	26	23	41	50	41	29	16	1	-	-
All races, female												
2004.....	1	2	6	10	16	30	27	22	12	2	1	-
2003.....	2	6	6	9	20	35	25	35	14	3	-	-
2002.....	5	4	6	6	15	33	32	28	20	4	-	-
2001.....	2	7	9	13	25	28	38	32	16	4	-	-
2000.....	2	8	7	8	20	24	42	32	12	4	-	-
1999.....	4	6	14	10	23	40	38	35	23	8	-	-
White, both sexes												
2004.....	4	6	16	25	46	65	58	48	22	4	1	-
2003.....	5	11	18	20	65	84	57	66	33	4	-	-
2002.....	10	9	27	23	48	83	85	51	29	6	-	-
2001.....	6	16	18	36	48	61	80	56	25	5	-	-
2000.....	3	16	19	26	49	61	82	59	21	5	-	-
1999.....	6	12	35	32	62	80	78	62	39	9	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant immunoproliferative diseases (C88)												
White, male												
2004.....	3	4	10	15	31	35	31	26	10	2	-	-
2003.....	5	5	13	13	45	51	35	32	19	1	-	-
2002.....	5	6	21	17	33	51	55	24	11	2	-	-
2001.....	4	9	12	25	24	35	42	25	10	1	-	-
2000.....	2	8	12	19	31	39	42	29	9	1	-	-
1999.....	4	7	22	22	39	45	40	29	16	1	-	-
White, female												
2004.....	1	2	6	10	15	30	27	22	12	2	1	-
2003.....	-	6	5	7	20	33	22	34	14	3	-	-
2002.....	5	3	6	6	15	32	30	27	18	4	-	-
2001.....	2	7	6	11	24	26	38	31	15	4	-	-
2000.....	1	8	7	7	18	22	40	30	12	4	-	-
1999.....	2	5	13	10	23	35	38	33	23	8	-	-
Black, both sexes												
2004.....	-	1	-	1	1	2	2	-	-	-	-	-
2003.....	4	1	2	3	-	4	3	-	-	-	-	-
2002.....	-	2	1	3	-	3	3	1	3	-	-	-
2001.....	-	-	5	3	1	2	1	1	2	-	1	-
2000.....	3	1	-	1	2	3	3	5	-	-	-	-
1999.....	4	2	5	1	1	5	1	2	-	-	-	-
Black, male												
2004.....	-	1	-	1	1	2	2	-	-	-	-	-
2003.....	2	1	1	1	-	2	-	-	-	-	-	-
2002.....	-	1	1	3	-	2	1	-	1	-	-	-
2001.....	-	-	2	2	-	-	1	-	1	-	1	-
2000.....	2	1	-	-	-	2	2	3	-	-	-	-
1999.....	2	1	4	1	1	2	1	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant immunoproliferative diseases (C88)												
Asian or Pacific Islander 2/, both sexes												
2004.....	-	-	-	-	1	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	1	-	-	-	1	1	-	-	-	-
2001.....	-	-	-	1	1	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	1	-	-	-	-	-	-
1999.....	-	1	-	-	1	5	-	-	-	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	1	-	-	-	1	1	-	-	-	-
2001.....	-	-	-	-	1	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	1	-	-	1	3	-	-	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	-	-	-	-	1	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	1	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	1	-	-	-	-	-	-
1999.....	-	-	-	-	-	2	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Multiple myeloma and malignant plasma cell neoplasms (C90)												
All races, both sexes												
2004.....	436	745	952	1,203	1,563	2,029	1,824	1,082	381	66	6	-
2003.....	464	686	941	1,257	1,684	2,039	1,810	1,098	431	74	4	-
2002.....	456	700	929	1,322	1,780	1,981	1,814	1,118	394	78	5	-
2001.....	424	704	927	1,292	1,798	2,038	1,704	1,066	373	66	2	-
2000.....	449	594	900	1,347	1,754	2,032	1,713	1,045	380	79	10	-
1999.....	414	712	927	1,314	1,761	1,933	1,686	976	350	76	5	-
All races, male												
2004.....	258	410	547	692	847	1,046	874	466	144	24	1	-
2003.....	280	409	566	711	884	999	854	488	155	32	1	-
2002.....	269	396	540	728	965	1,037	870	482	147	35	3	-
2001.....	248	400	521	714	958	1,054	794	421	141	22	2	-
2000.....	244	312	475	732	915	1,046	801	430	128	32	3	-
1999.....	240	402	531	706	899	976	779	421	117	25	3	-
All races, female												
2004.....	178	335	405	511	716	983	950	616	237	42	5	-
2003.....	184	277	375	546	800	1,040	956	610	276	42	3	-
2002.....	187	304	389	594	815	944	944	636	247	43	2	-
2001.....	176	304	406	578	840	984	910	645	232	44	-	-
2000.....	205	282	425	615	839	986	912	615	252	47	7	-
1999.....	174	310	396	608	862	957	907	555	233	51	2	-
White, both sexes												
2004.....	306	548	750	939	1,242	1,708	1,563	947	321	55	4	-
2003.....	325	519	743	996	1,369	1,698	1,556	950	376	60	1	-
2002.....	324	526	715	1,048	1,468	1,694	1,585	993	332	68	3	-
2001.....	314	520	693	1,025	1,489	1,734	1,458	922	329	55	2	-
2000.....	328	426	703	1,083	1,447	1,705	1,496	888	312	61	6	-
1999.....	293	515	690	1,061	1,448	1,617	1,471	835	299	66	2	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Multiple myeloma and malignant plasma cell neoplasms (C90)												
White, male												
2004.....	184	314	440	554	696	902	758	411	121	22	-	-
2003.....	198	315	455	579	746	863	744	430	138	30	-	-
2002.....	205	304	428	593	820	908	771	435	124	34	2	-
2001.....	190	298	397	579	809	918	689	372	125	19	2	-
2000.....	179	232	380	602	762	893	716	380	107	28	3	-
1999.....	184	299	419	588	745	837	689	369	96	24	1	-
White, female												
2004.....	122	234	310	385	546	806	805	536	200	33	4	-
2003.....	127	204	288	417	623	835	812	520	238	30	1	-
2002.....	119	222	287	455	648	786	814	558	208	34	1	-
2001.....	124	222	296	446	680	816	769	550	204	36	-	-
2000.....	149	194	323	481	685	812	780	508	205	33	3	-
1999.....	109	216	271	473	703	780	782	466	203	42	1	-
Black, both sexes												
2004.....	115	171	187	235	290	285	238	124	54	11	2	-
2003.....	124	147	177	241	282	305	231	137	52	14	2	-
2002.....	117	157	187	248	281	257	207	117	55	9	2	-
2001.....	96	167	212	236	283	274	225	134	42	10	-	-
2000.....	115	157	181	246	282	297	202	145	62	14	3	-
1999.....	113	179	210	218	277	281	194	135	48	10	3	-
Black, male												
2004.....	62	85	97	126	136	125	101	49	20	2	1	-
2003.....	74	86	101	118	120	124	99	53	15	2	-	-
2002.....	57	82	101	123	134	112	91	44	20	1	1	-
2001.....	49	93	113	117	134	123	99	43	15	3	-	-
2000.....	60	74	89	119	144	140	80	44	19	3	-	-
1999.....	52	94	98	100	137	123	84	47	19	1	2	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Multiple myeloma and malignant plasma cell neoplasms (C90)

Black, female

2004.....	53	86	90	109	154	160	137	75	34	9	1	-
2003.....	50	61	76	123	162	181	132	84	37	12	2	-
2002.....	60	75	86	125	147	145	116	73	35	8	1	-
2001.....	47	74	99	119	149	151	126	91	27	7	-	-
2000.....	55	83	92	127	138	157	122	101	43	11	3	-
1999.....	61	85	112	118	140	158	110	88	29	9	1	-

American Indian 1/, both sexes

2004.....	5	11	3	6	4	12	5	1	1	-	-	-
2003.....	3	6	6	5	8	12	7	3	2	-	1	-
2002.....	7	5	5	6	5	8	2	1	-	-	-	-
2001.....	4	4	6	8	7	6	5	4	1	-	-	-
2000.....	2	3	5	1	3	9	2	3	3	3	-	-
1999.....	1	2	4	6	10	12	3	1	1	-	-	-

American Indian 1/, male

2004.....	5	3	1	3	2	8	3	-	-	-	-	-
2003.....	2	1	3	4	3	2	3	3	1	-	1	-
2002.....	5	2	-	2	2	3	2	1	-	-	-	-
2001.....	3	1	3	5	2	2	2	2	-	-	-	-
2000.....	1	3	2	1	1	3	-	2	-	1	-	-
1999.....	-	1	1	4	7	5	1	1	1	-	-	-

American Indian 1/, female

2004.....	-	8	2	3	2	4	2	1	1	-	-	-
2003.....	1	5	3	1	5	10	4	-	1	-	-	-
2002.....	2	3	5	4	3	5	-	-	-	-	-	-
2001.....	1	3	3	3	5	4	3	2	1	-	-	-
2000.....	1	-	3	-	2	6	2	1	3	2	-	-
1999.....	1	1	3	2	3	7	2	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Multiple myeloma and malignant plasma cell neoplasms (C90)												
Asian or Pacific Islander 2/, both sexes												
2004.....	10	15	12	23	27	24	18	10	5	-	-	-
2003.....	12	14	15	15	25	24	16	8	1	-	-	-
2002.....	8	12	22	20	26	22	20	7	7	1	-	-
2001.....	10	13	16	23	19	24	16	6	1	1	-	-
2000.....	4	8	11	17	22	21	13	9	3	1	1	-
1999.....	7	16	23	29	26	23	18	5	2	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	7	8	9	9	13	11	12	6	3	-	-	-
2003.....	6	7	7	10	15	10	8	2	1	-	-	-
2002.....	2	8	11	10	9	14	6	2	3	-	-	-
2001.....	6	8	8	13	13	11	4	4	1	-	-	-
2000.....	4	3	4	10	8	10	5	4	2	-	-	-
1999.....	4	8	13	14	10	11	5	4	1	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	3	7	3	14	14	13	6	4	2	-	-	-
2003.....	6	7	8	5	10	14	8	6	-	-	-	-
2002.....	6	4	11	10	17	8	14	5	4	1	-	-
2001.....	4	5	8	10	6	13	12	2	-	1	-	-
2000.....	-	5	7	7	14	11	8	5	1	1	1	-
1999.....	3	8	10	15	16	12	13	1	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Leukemia (C91-C95)												
All races, both sexes												
2004.....	812	1,176	1,512	1,985	2,709	3,377	3,420	2,349	1,165	276	40	-
2003.....	770	1,102	1,503	1,911	2,795	3,532	3,405	2,317	1,091	350	28	-
2002.....	853	1,111	1,500	1,998	2,820	3,392	3,371	2,294	1,086	281	35	-
2001.....	845	1,073	1,462	1,963	2,923	3,541	3,155	2,256	1,008	271	50	-
2000.....	789	1,041	1,373	2,073	2,937	3,515	3,218	2,194	1,041	271	37	-
1999.....	744	1,112	1,404	2,078	2,934	3,401	3,059	2,151	984	258	31	-
All races, male												
2004.....	479	720	943	1,237	1,666	1,985	1,836	1,112	453	95	9	-
2003.....	454	683	910	1,215	1,715	2,054	1,826	1,102	416	91	6	-
2002.....	486	667	891	1,242	1,768	1,995	1,809	1,062	420	95	15	-
2001.....	467	632	920	1,216	1,746	2,023	1,676	1,041	380	89	17	-
2000.....	435	629	823	1,287	1,803	1,982	1,713	1,018	359	85	8	-
1999.....	431	636	853	1,275	1,779	1,946	1,579	964	375	75	9	-
All races, female												
2004.....	333	456	569	748	1,043	1,392	1,584	1,237	712	181	31	-
2003.....	316	419	593	696	1,080	1,478	1,579	1,215	675	259	22	-
2002.....	367	444	609	756	1,052	1,397	1,562	1,232	666	186	20	-
2001.....	378	441	542	747	1,177	1,518	1,479	1,215	628	182	33	-
2000.....	354	412	550	786	1,134	1,533	1,505	1,176	682	186	29	-
1999.....	313	476	551	803	1,155	1,455	1,480	1,187	609	183	22	-
White, both sexes												
2004.....	674	1,019	1,314	1,750	2,404	3,095	3,175	2,201	1,079	259	37	-
2003.....	635	962	1,323	1,702	2,522	3,265	3,168	2,180	1,019	329	23	-
2002.....	715	944	1,302	1,793	2,568	3,110	3,152	2,138	1,019	260	30	-
2001.....	696	928	1,282	1,719	2,667	3,255	2,968	2,132	946	248	48	-
2000.....	669	900	1,188	1,870	2,690	3,272	3,026	2,053	967	250	32	-
1999.....	619	975	1,221	1,839	2,692	3,166	2,841	2,037	915	238	28	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Leukemia (C91-C95)												
White, male												
2004.....	393	635	832	1,092	1,484	1,837	1,719	1,055	419	88	8	-
2003.....	376	593	819	1,090	1,569	1,925	1,725	1,047	390	85	6	-
2002.....	415	580	779	1,118	1,618	1,847	1,700	1,000	396	90	12	-
2001.....	392	555	812	1,070	1,611	1,879	1,579	992	358	81	17	-
2000.....	366	550	717	1,176	1,651	1,855	1,616	949	338	76	7	-
1999.....	360	563	751	1,141	1,643	1,815	1,484	911	350	69	9	-
White, female												
2004.....	281	384	482	658	920	1,258	1,456	1,146	660	171	29	-
2003.....	259	369	504	612	953	1,340	1,443	1,133	629	244	17	-
2002.....	300	364	523	675	950	1,263	1,452	1,138	623	170	18	-
2001.....	304	373	470	649	1,056	1,376	1,389	1,140	588	167	31	-
2000.....	303	350	471	694	1,039	1,417	1,410	1,104	629	174	25	-
1999.....	259	412	470	698	1,049	1,351	1,357	1,126	565	169	19	-
Black, both sexes												
2004.....	118	129	169	188	248	219	190	125	69	16	3	-
2003.....	111	113	151	171	224	219	189	115	67	16	5	-
2002.....	108	136	159	163	213	228	178	133	61	19	5	-
2001.....	112	112	137	207	208	238	155	109	55	21	1	-
2000.....	97	107	154	162	213	195	155	124	64	18	5	-
1999.....	96	106	146	201	206	196	177	96	58	18	3	-
Black, male												
2004.....	72	71	94	117	150	106	82	49	24	6	1	-
2003.....	67	76	75	101	120	110	79	46	23	5	-	-
2002.....	55	74	87	94	127	114	84	52	21	5	3	-
2001.....	61	63	84	127	108	111	76	43	18	7	-	-
2000.....	53	61	90	86	130	105	76	61	16	7	1	-
1999.....	56	61	81	112	112	108	70	42	20	4	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Leukemia (C91-C95)												
Black, female												
2004.....	46	58	75	71	98	113	108	76	45	10	2	-
2003.....	44	37	76	70	104	109	110	69	44	11	5	-
2002.....	53	62	72	69	86	114	94	81	40	14	2	-
2001.....	51	49	53	80	100	127	79	66	37	14	1	-
2000.....	44	46	64	76	83	90	79	63	48	11	4	-
1999.....	40	45	65	89	94	88	107	54	38	14	3	-
American Indian 1/, both sexes												
2004.....	4	7	4	8	8	4	15	4	3	-	-	-
2003.....	6	4	6	4	10	6	6	6	-	-	-	-
2002.....	1	7	6	7	5	8	7	4	2	1	-	-
2001.....	6	10	8	8	7	1	4	2	1	1	-	-
2000.....	2	8	7	9	7	5	5	6	3	1	-	-
1999.....	6	4	4	7	6	6	4	3	2	-	-	-
American Indian 1/, male												
2004.....	2	5	4	5	2	2	9	1	-	-	-	-
2003.....	4	1	2	3	4	1	3	1	-	-	-	-
2002.....	1	3	4	4	3	4	5	1	1	-	-	-
2001.....	3	5	5	3	4	-	2	-	-	-	-	-
2000.....	2	4	3	5	6	1	3	2	1	1	-	-
1999.....	4	1	3	7	4	6	1	2	1	-	-	-
American Indian 1/, female												
2004.....	2	2	-	3	6	2	6	3	3	-	-	-
2003.....	2	3	4	1	6	5	3	5	-	-	-	-
2002.....	-	4	2	3	2	4	2	3	1	1	-	-
2001.....	3	5	3	5	3	1	2	2	1	1	-	-
2000.....	-	4	4	4	1	4	2	4	2	-	-	-
1999.....	2	3	1	-	2	-	3	1	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Leukemia (C91-C95)												
Asian or Pacific Islander 2/, both sexes												
2004.....	16	21	25	39	49	59	40	19	14	1	-	-
2003.....	18	23	23	34	39	42	42	16	5	5	-	-
2002.....	29	24	33	35	34	46	34	19	4	1	-	-
2001.....	31	23	35	29	41	47	28	13	6	1	1	-
2000.....	21	26	24	32	27	43	32	11	7	2	-	-
1999.....	23	27	33	31	30	33	37	15	9	2	-	-
Asian or Pacific Islander 2/, male												
2004.....	12	9	13	23	30	40	26	7	10	1	-	-
2003.....	7	13	14	21	22	18	19	8	3	1	-	-
2002.....	15	10	21	26	20	30	20	9	2	-	-	-
2001.....	11	9	19	16	23	33	19	6	4	1	-	-
2000.....	14	14	13	20	16	21	18	6	4	1	-	-
1999.....	11	11	18	15	20	17	24	9	4	2	-	-
Asian or Pacific Islander 2/, female												
2004.....	4	12	12	16	19	19	14	12	4	-	-	-
2003.....	11	10	9	13	17	24	23	8	2	4	-	-
2002.....	14	14	12	9	14	16	14	10	2	1	-	-
2001.....	20	14	16	13	18	14	9	7	2	-	1	-
2000.....	7	12	11	12	11	22	14	5	3	1	-	-
1999.....	12	16	15	16	10	16	13	6	5	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Lymphoid leukemia (C91)												
All races, both sexes												
2004.....	198	310	381	493	711	886	945	769	487	128	19	-
2003.....	166	285	363	509	732	960	1,016	772	448	160	15	-
2002.....	210	272	383	510	714	905	988	804	437	127	15	-
2001.....	195	263	370	514	761	956	938	790	429	125	30	-
2000.....	168	242	343	524	788	973	963	760	455	125	18	-
1999.....	179	282	378	572	820	983	961	740	399	120	19	-
All races, male												
2004.....	123	217	276	315	456	526	523	379	201	44	5	-
2003.....	113	205	242	370	478	572	569	366	188	44	4	-
2002.....	140	173	245	347	481	543	540	383	166	43	6	-
2001.....	128	179	258	356	476	564	495	356	154	46	13	-
2000.....	102	174	237	361	499	574	521	350	148	36	3	-
1999.....	128	187	259	380	510	576	517	352	147	40	5	-
All races, female												
2004.....	75	93	105	178	255	360	422	390	286	84	14	-
2003.....	53	80	121	139	254	388	447	406	260	116	11	-
2002.....	70	99	138	163	233	362	448	421	271	84	9	-
2001.....	67	84	112	158	285	392	443	434	275	79	17	-
2000.....	66	68	106	163	289	399	442	410	307	89	15	-
1999.....	51	95	119	192	310	407	444	388	252	80	14	-
White, both sexes												
2004.....	167	274	320	436	626	806	877	726	457	123	18	-
2003.....	137	250	324	452	654	886	960	734	421	152	12	-
2002.....	184	229	328	456	648	814	933	761	416	123	14	-
2001.....	156	226	320	437	695	873	887	754	408	117	30	-
2000.....	144	217	299	473	716	906	908	713	427	117	15	-
1999.....	155	241	326	506	756	917	900	704	373	112	17	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Lymphoid leukemia (C91)												
White, male												
2004.....	103	197	233	275	397	484	491	363	187	40	5	-
2003.....	90	180	220	333	438	536	542	346	179	40	4	-
2002.....	125	149	213	316	439	491	515	362	157	43	5	-
2001.....	107	158	223	308	441	523	467	340	147	43	13	-
2000.....	87	157	204	326	456	535	493	327	138	32	2	-
1999.....	111	162	227	336	470	539	490	335	138	37	5	-
White, female												
2004.....	64	77	87	161	229	322	386	363	270	83	13	-
2003.....	47	70	104	119	216	350	418	388	242	112	8	-
2002.....	59	80	115	140	209	323	418	399	259	80	9	-
2001.....	49	68	97	129	254	350	420	414	261	74	17	-
2000.....	57	60	95	147	260	371	415	386	289	85	13	-
1999.....	44	79	99	170	286	378	410	369	235	75	12	-
Black, both sexes												
2004.....	27	34	51	50	77	71	58	39	26	4	1	-
2003.....	24	31	34	53	72	66	48	33	25	7	3	-
2002.....	21	39	49	49	59	83	50	36	21	4	1	-
2001.....	35	34	45	69	59	75	50	33	21	8	-	-
2000.....	21	20	38	41	68	59	48	45	26	6	3	-
1999.....	21	37	43	63	59	56	56	34	25	8	2	-
Black, male												
2004.....	18	20	36	37	53	39	28	16	12	3	-	-
2003.....	20	23	19	34	36	33	23	15	7	3	-	-
2002.....	12	22	29	28	35	47	24	19	9	-	1	-
2001.....	20	20	32	44	32	36	28	15	7	3	-	-
2000.....	13	14	29	28	41	35	25	23	9	2	1	-
1999.....	15	23	26	42	36	32	25	16	9	3	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Lymphoid leukemia (C91)												
Black, female												
2004.....	9	14	15	13	24	32	30	23	14	1	1	-
2003.....	4	8	15	19	36	33	25	18	18	4	3	-
2002.....	9	17	20	21	24	36	26	17	12	4	-	-
2001.....	15	14	13	25	27	39	22	18	14	5	-	-
2000.....	8	6	9	13	27	24	23	22	17	4	2	-
1999.....	6	14	17	21	23	24	31	18	16	5	2	-
American Indian 1/, both sexes												
2004.....	1	-	2	2	3	1	6	1	1	-	-	-
2003.....	1	1	3	1	4	1	1	1	-	-	-	-
2002.....	-	2	1	1	1	3	-	2	-	-	-	-
2001.....	1	-	1	2	3	1	-	-	-	-	-	-
2000.....	1	2	2	3	1	1	2	2	1	1	-	-
1999.....	1	-	-	2	1	3	1	1	-	-	-	-
American Indian 1/, male												
2004.....	1	-	2	1	1	-	2	-	-	-	-	-
2003.....	1	-	1	1	3	-	-	1	-	-	-	-
2002.....	-	1	1	-	1	1	-	-	-	-	-	-
2001.....	-	-	1	1	1	-	-	-	-	-	-	-
2000.....	1	1	1	2	1	-	2	-	-	1	-	-
1999.....	1	-	-	2	1	3	-	-	-	-	-	-
American Indian 1/, female												
2004.....	-	-	-	1	2	1	4	1	1	-	-	-
2003.....	-	1	2	-	1	1	1	-	-	-	-	-
2002.....	-	1	-	1	-	2	-	2	-	-	-	-
2001.....	1	-	-	1	2	1	-	-	-	-	-	-
2000.....	-	1	1	1	-	1	-	2	1	-	-	-
1999.....	-	-	-	-	-	-	1	1	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Lymphoid leukemia (C91)												
Asian or Pacific Islander 2/, both sexes												
2004.....	3	2	8	5	5	8	4	3	3	1	-	-
2003.....	4	3	2	3	2	7	7	4	2	1	-	-
2002.....	5	2	5	4	6	5	5	5	-	-	-	-
2001.....	3	3	4	6	4	7	1	3	-	-	-	-
2000.....	2	3	4	7	3	7	5	-	1	1	-	-
1999.....	2	4	9	1	4	7	4	1	1	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	1	-	5	2	5	3	2	-	2	1	-	-
2003.....	2	2	2	2	1	3	4	4	2	1	-	-
2002.....	3	1	2	3	6	4	1	2	-	-	-	-
2001.....	1	1	2	3	2	5	-	1	-	-	-	-
2000.....	1	2	3	5	1	4	1	-	1	1	-	-
1999.....	1	2	6	-	3	2	2	1	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	2	2	3	3	-	5	2	3	1	-	-	-
2003.....	2	1	-	1	1	4	3	-	-	-	-	-
2002.....	2	1	3	1	-	1	4	3	-	-	-	-
2001.....	2	2	2	3	2	2	1	2	-	-	-	-
2000.....	1	1	1	2	2	3	4	-	-	-	-	-
1999.....	1	2	3	1	1	5	2	-	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Myeloid leukemia (C92)												
All races, both sexes												
2004.....	465	614	826	1,069	1,317	1,610	1,495	849	340	52	2	-
2003.....	460	577	838	983	1,403	1,671	1,383	854	298	76	3	-
2002.....	469	605	825	1,045	1,401	1,602	1,432	798	307	66	8	-
2001.....	502	603	772	1,022	1,471	1,651	1,312	779	264	58	5	-
2000.....	479	578	737	1,057	1,416	1,628	1,320	754	263	48	7	-
1999.....	423	607	699	1,030	1,348	1,430	1,197	742	251	64	7	-
All races, male												
2004.....	266	354	476	673	786	927	781	390	136	19	-	-
2003.....	251	342	487	599	831	962	735	413	121	12	2	-
2002.....	253	363	476	625	855	936	775	368	126	24	5	-
2001.....	254	337	475	591	862	913	714	366	102	15	-	-
2000.....	269	323	419	620	854	903	718	357	102	18	3	-
1999.....	231	330	405	605	807	793	624	337	95	15	3	-
All races, female												
2004.....	199	260	350	396	531	683	714	459	204	33	2	-
2003.....	209	235	351	384	572	709	648	441	177	64	1	-
2002.....	216	242	349	420	546	666	657	430	181	42	3	-
2001.....	248	266	297	431	609	738	598	413	162	43	5	-
2000.....	210	255	318	437	562	725	602	397	161	30	4	-
1999.....	192	277	294	425	541	637	573	405	156	49	4	-
White, both sexes												
2004.....	384	536	729	951	1,194	1,484	1,385	800	319	51	1	-
2003.....	384	512	735	887	1,274	1,554	1,288	804	273	67	2	-
2002.....	397	512	725	937	1,288	1,488	1,338	746	287	61	7	-
2001.....	419	520	694	908	1,350	1,523	1,234	733	244	53	5	-
2000.....	406	497	631	962	1,313	1,533	1,240	702	245	41	6	-
1999.....	353	541	608	919	1,234	1,331	1,116	708	232	60	7	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Myeloid leukemia (C92)												
White, male												
2004.....	213	313	429	603	720	858	727	372	131	19	-	-
2003.....	212	299	439	539	765	906	698	394	109	12	2	-
2002.....	219	315	419	561	792	876	727	344	117	22	4	-
2001.....	215	293	430	527	793	849	679	352	93	15	-	-
2000.....	226	281	365	573	788	862	676	337	99	15	3	-
1999.....	191	297	352	542	745	739	589	322	87	14	3	-
White, female												
2004.....	171	223	300	348	474	626	658	428	188	32	1	-
2003.....	172	213	296	348	509	648	590	410	164	55	-	-
2002.....	178	197	306	376	496	612	611	402	170	39	3	-
2001.....	204	227	264	381	557	674	555	381	151	38	5	-
2000.....	180	216	266	389	525	671	564	365	146	26	3	-
1999.....	162	244	256	377	489	592	527	386	145	46	4	-
Black, both sexes												
2004.....	69	62	83	91	85	87	82	43	18	1	1	-
2003.....	62	48	83	74	103	91	72	42	24	5	1	-
2002.....	51	71	76	79	93	88	69	42	18	4	1	-
2001.....	56	60	51	96	95	99	58	36	16	4	-	-
2000.....	59	60	87	78	84	75	57	43	14	7	1	-
1999.....	49	44	72	87	92	79	62	27	15	3	-	-
Black, male												
2004.....	43	32	40	53	45	39	35	16	4	-	-	-
2003.....	32	34	37	47	54	49	27	18	12	-	-	-
2002.....	24	39	41	44	53	43	30	19	8	2	1	-
2001.....	29	34	31	55	53	45	24	10	7	-	-	-
2000.....	33	31	45	37	51	33	27	16	2	3	-	-
1999.....	30	26	41	48	49	42	24	11	6	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Myeloid leukemia (C92)												
Black, female												
2004.....	26	30	43	38	40	48	47	27	14	1	1	-
2003.....	30	14	46	27	49	42	45	24	12	5	1	-
2002.....	27	32	35	35	40	45	39	23	10	2	-	-
2001.....	27	26	20	41	42	54	34	26	9	4	-	-
2000.....	26	29	42	41	33	42	30	27	12	4	1	-
1999.....	19	18	31	39	43	37	38	16	9	3	-	-
American Indian 1/, both sexes												
2004.....	1	4	2	4	3	3	5	1	-	-	-	-
2003.....	3	1	3	3	4	2	4	2	-	-	-	-
2002.....	1	5	4	4	3	1	7	1	-	-	-	-
2001.....	3	4	3	3	3	-	3	2	-	1	-	-
2000.....	1	6	5	4	5	3	1	4	2	-	-	-
1999.....	5	4	2	4	4	1	1	-	1	-	-	-
American Indian 1/, male												
2004.....	-	3	2	2	1	2	3	-	-	-	-	-
2003.....	2	-	1	2	1	-	2	-	-	-	-	-
2002.....	1	2	2	2	2	1	5	-	-	-	-	-
2001.....	2	2	1	-	3	-	1	-	-	-	-	-
2000.....	1	3	2	1	4	1	1	2	1	-	-	-
1999.....	3	1	2	4	2	1	-	-	1	-	-	-
American Indian 1/, female												
2004.....	1	1	-	2	2	1	2	1	-	-	-	-
2003.....	1	1	2	1	3	2	2	2	-	-	-	-
2002.....	-	3	2	2	1	-	2	1	-	-	-	-
2001.....	1	2	2	3	-	-	2	2	-	1	-	-
2000.....	-	3	3	3	1	2	-	2	1	-	-	-
1999.....	2	3	-	-	2	-	1	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Myeloid leukemia (C92)												
Asian or Pacific Islander 2/, both sexes												
2004.....	11	12	12	23	35	36	23	5	3	-	-	-
2003.....	11	16	17	19	22	24	19	6	1	4	-	-
2002.....	20	17	20	25	17	25	18	9	2	1	-	-
2001.....	24	19	24	15	23	29	17	8	4	-	-	-
2000.....	13	15	14	13	14	17	22	5	2	-	-	-
1999.....	16	18	17	20	18	19	18	7	3	1	-	-
Asian or Pacific Islander 2/, male												
2004.....	10	6	5	15	20	28	16	2	1	-	-	-
2003.....	5	9	10	11	11	7	8	1	-	-	-	-
2002.....	9	7	14	18	8	16	13	5	1	-	-	-
2001.....	8	8	13	9	13	19	10	4	2	-	-	-
2000.....	9	8	7	9	11	7	14	2	-	-	-	-
1999.....	7	6	10	11	11	11	11	4	1	1	-	-
Asian or Pacific Islander 2/, female												
2004.....	1	6	7	8	15	8	7	3	2	-	-	-
2003.....	6	7	7	8	11	17	11	5	1	4	-	-
2002.....	11	10	6	7	9	9	5	4	1	1	-	-
2001.....	16	11	11	6	10	10	7	4	2	-	-	-
2000.....	4	7	7	4	3	10	8	3	2	-	-	-
1999.....	9	12	7	9	7	8	7	3	2	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Monocytic leukemia (C93)												
All races, both sexes												
2004.....	2	2	4	13	12	29	27	19	6	2	-	-
2003.....	4	11	8	12	15	19	34	16	8	3	2	-
2002.....	7	7	8	15	18	27	20	22	6	1	-	-
2001.....	3	4	11	12	18	29	28	22	9	1	-	-
2000.....	1	7	13	14	21	25	20	20	7	2	-	-
1999.....	4	7	2	21	25	34	30	23	11	3	-	-
All races, male												
2004.....	2	1	4	7	8	21	15	14	3	-	-	-
2003.....	2	6	7	5	10	13	17	9	5	1	-	-
2002.....	4	4	6	9	12	22	5	14	4	-	-	-
2001.....	2	2	7	6	11	16	12	13	3	1	-	-
2000.....	1	5	6	8	15	16	13	12	2	1	-	-
1999.....	2	3	1	11	14	22	17	11	5	-	-	-
All races, female												
2004.....	-	1	-	6	4	8	12	5	3	2	-	-
2003.....	2	5	1	7	5	6	17	7	3	2	2	-
2002.....	3	3	2	6	6	5	15	8	2	1	-	-
2001.....	1	2	4	6	7	13	16	9	6	-	-	-
2000.....	-	2	7	6	6	9	7	8	5	1	-	-
1999.....	2	4	1	10	11	12	13	12	6	3	-	-
White, both sexes												
2004.....	2	1	4	10	11	26	27	18	6	2	-	-
2003.....	4	10	8	11	13	18	34	16	8	3	2	-
2002.....	5	7	8	15	17	25	20	20	5	1	-	-
2001.....	3	4	10	12	18	28	27	22	8	1	-	-
2000.....	1	7	11	13	19	24	19	19	6	2	-	-
1999.....	4	6	2	19	23	31	27	23	11	3	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Monocytic leukemia (C93)												
White, male												
2004.....	2	-	4	5	7	19	15	13	3	-	-	-
2003.....	2	6	7	4	9	13	17	9	5	1	-	-
2002.....	2	4	6	9	12	21	5	13	4	-	-	-
2001.....	2	2	7	6	11	16	11	13	3	1	-	-
2000.....	1	5	5	8	14	15	12	11	2	1	-	-
1999.....	2	3	1	11	13	19	16	11	5	-	-	-
White, female												
2004.....	-	1	-	5	4	7	12	5	3	2	-	-
2003.....	2	4	1	7	4	5	17	7	3	2	2	-
2002.....	3	3	2	6	5	4	15	7	1	1	-	-
2001.....	1	2	3	6	7	12	16	9	5	-	-	-
2000.....	-	2	6	5	5	9	7	8	4	1	-	-
1999.....	2	3	1	8	10	12	11	12	6	3	-	-
Black, both sexes												
2004.....	-	-	-	1	-	1	-	1	-	-	-	-
2003.....	-	1	-	1	2	1	-	-	-	-	-	-
2002.....	2	-	-	-	1	2	-	2	1	-	-	-
2001.....	-	-	1	-	-	1	1	-	1	-	-	-
2000.....	-	-	1	1	1	-	1	1	1	-	-	-
1999.....	-	1	-	2	2	2	2	-	-	-	-	-
Black, male												
2004.....	-	-	-	-	-	1	-	1	-	-	-	-
2003.....	-	-	-	1	1	-	-	-	-	-	-	-
2002.....	2	-	-	-	-	1	-	1	-	-	-	-
2001.....	-	-	-	-	-	-	1	-	-	-	-	-
2000.....	-	-	1	-	-	-	1	1	-	-	-	-
1999.....	-	-	-	-	1	2	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified leukemia (C94-C95)												
All races, both sexes												
2004.....	147	250	301	410	669	852	953	712	332	94	19	-
2003.....	140	229	294	407	645	882	972	675	337	111	8	-
2002.....	167	227	284	428	687	858	931	670	336	87	12	-
2001.....	145	203	309	415	673	905	877	665	306	87	15	-
2000.....	141	214	280	478	712	889	915	660	316	96	12	-
1999.....	138	216	325	455	741	954	871	646	323	71	5	-
All races, male												
2004.....	88	148	187	242	416	511	517	329	113	32	4	-
2003.....	88	130	174	241	396	507	505	314	102	34	-	-
2002.....	89	127	164	261	420	494	489	297	124	28	4	-
2001.....	83	114	180	263	397	530	455	306	121	27	4	-
2000.....	63	127	161	298	435	489	461	299	107	30	2	-
1999.....	70	116	188	279	448	555	421	264	128	20	1	-
All races, female												
2004.....	59	102	114	168	253	341	436	383	219	62	15	-
2003.....	52	99	120	166	249	375	467	361	235	77	8	-
2002.....	78	100	120	167	267	364	442	373	212	59	8	-
2001.....	62	89	129	152	276	375	422	359	185	60	11	-
2000.....	78	87	119	180	277	400	454	361	209	66	10	-
1999.....	68	100	137	176	293	399	450	382	195	51	4	-
White, both sexes												
2004.....	121	208	261	353	573	779	886	657	297	83	18	-
2003.....	110	190	256	352	581	807	886	626	317	107	7	-
2002.....	129	196	241	385	615	783	861	611	311	75	9	-
2001.....	118	178	258	362	604	831	820	623	286	77	13	-
2000.....	118	179	247	422	642	809	859	619	289	90	11	-
1999.....	107	187	285	395	679	887	798	602	299	63	4	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified leukemia (C94-C95)												
White, male												
2004.....	75	125	166	209	360	476	486	307	98	29	3	-
2003.....	72	108	153	214	357	470	468	298	97	32	-	-
2002.....	69	112	141	232	375	459	453	281	118	25	3	-
2001.....	68	102	152	229	366	491	422	287	115	22	4	-
2000.....	52	107	143	269	393	443	435	274	99	28	2	-
1999.....	56	101	171	252	415	518	389	243	120	18	1	-
White, female												
2004.....	46	83	95	144	213	303	400	350	199	54	15	-
2003.....	38	82	103	138	224	337	418	328	220	75	7	-
2002.....	60	84	100	153	240	324	408	330	193	50	6	-
2001.....	50	76	106	133	238	340	398	336	171	55	9	-
2000.....	66	72	104	153	249	366	424	345	190	62	9	-
1999.....	51	86	114	143	264	369	409	359	179	45	3	-
Black, both sexes												
2004.....	22	33	35	46	86	60	50	42	25	11	1	-
2003.....	25	33	34	43	47	61	69	40	18	4	1	-
2002.....	34	26	34	35	60	55	59	53	21	11	3	-
2001.....	21	18	40	42	54	63	46	40	17	9	1	-
2000.....	17	27	28	42	60	61	49	35	23	5	1	-
1999.....	26	24	31	49	53	59	57	35	18	7	1	-
Black, male												
2004.....	11	19	18	27	52	27	19	16	8	3	1	-
2003.....	15	19	19	19	29	28	29	13	4	2	-	-
2002.....	17	13	17	22	39	23	30	13	4	3	1	-
2001.....	12	9	21	28	23	30	23	18	4	4	-	-
2000.....	7	16	15	21	38	37	23	21	5	2	-	-
1999.....	11	12	14	22	26	32	21	15	5	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified leukemia (C94-C95)												
Black, female												
2004.....	11	14	17	19	34	33	31	26	17	8	-	-
2003.....	10	14	15	24	18	33	40	27	14	2	1	-
2002.....	17	13	17	13	21	32	29	40	17	8	2	-
2001.....	9	9	19	14	31	33	23	22	13	5	1	-
2000.....	10	11	13	21	22	24	26	14	18	3	1	-
1999.....	15	12	17	27	27	27	36	20	13	6	1	-
American Indian 1/, both sexes												
2004.....	2	3	-	2	2	-	4	2	2	-	-	-
2003.....	2	2	-	-	2	3	1	3	-	-	-	-
2002.....	-	-	1	2	1	4	-	1	2	1	-	-
2001.....	2	6	4	3	1	-	1	-	1	-	-	-
2000.....	-	-	-	2	-	1	2	-	-	-	-	-
1999.....	-	-	2	1	1	2	2	2	1	-	-	-
American Indian 1/, male												
2004.....	1	2	-	2	-	-	4	1	-	-	-	-
2003.....	1	1	-	-	-	1	1	-	-	-	-	-
2002.....	-	-	1	2	-	2	-	1	1	-	-	-
2001.....	1	3	3	2	-	-	1	-	-	-	-	-
2000.....	-	-	-	2	-	-	-	-	-	-	-	-
1999.....	-	-	1	1	1	2	1	2	-	-	-	-
American Indian 1/, female												
2004.....	1	1	-	-	2	-	-	1	2	-	-	-
2003.....	1	1	-	-	2	2	-	3	-	-	-	-
2002.....	-	-	-	-	1	2	-	-	1	1	-	-
2001.....	1	3	1	1	1	-	-	-	1	-	-	-
2000.....	-	-	-	-	-	1	2	-	-	-	-	-
1999.....	-	-	1	-	-	-	1	-	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified leukemia (C94-C95)												
Asian or Pacific Islander 2/, both sexes												
2004.....	2	6	5	9	8	13	13	11	8	-	-	-
2003.....	3	4	4	12	15	11	16	6	2	-	-	-
2002.....	4	5	8	6	11	16	11	5	2	-	-	-
2001.....	4	1	7	8	14	11	10	2	2	1	1	-
2000.....	6	8	5	12	10	18	5	6	4	1	-	-
1999.....	5	5	7	10	8	6	14	7	5	1	-	-
Asian or Pacific Islander 2/, male												
2004.....	1	2	3	4	4	8	8	5	7	-	-	-
2003.....	-	2	2	8	10	8	7	3	1	-	-	-
2002.....	3	2	5	5	6	10	6	2	1	-	-	-
2001.....	2	-	4	4	8	9	9	1	2	1	-	-
2000.....	4	4	3	6	4	9	3	4	3	-	-	-
1999.....	3	3	2	4	6	3	10	4	3	1	-	-
Asian or Pacific Islander 2/, female												
2004.....	1	4	2	5	4	5	5	6	1	-	-	-
2003.....	3	2	2	4	5	3	9	3	1	-	-	-
2002.....	1	3	3	1	5	6	5	3	1	-	-	-
2001.....	2	1	3	4	6	2	1	1	-	-	1	-
2000.....	2	4	2	6	6	9	2	2	1	1	-	-
1999.....	2	2	5	6	2	3	4	3	2	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
In situ neoplasms (D00-D09)												
All races, both sexes												
2004.....	1	-	-	1	-	2	1	1	1	1	-	-
2003.....	-	-	2	1	2	2	2	-	3	-	-	-
2002.....	1	-	1	2	3	3	1	2	3	-	-	-
2001.....	1	1	1	2	5	7	5	5	3	-	-	-
2000.....	-	-	1	3	4	-	1	3	1	1	-	-
1999.....	-	1	-	1	2	2	3	4	-	1	-	-
All races, male												
2004.....	1	-	-	-	-	1	1	-	-	-	-	-
2003.....	-	-	-	1	2	1	-	-	2	-	-	-
2002.....	1	-	-	1	1	2	1	-	-	-	-	-
2001.....	-	1	1	1	3	4	3	2	1	-	-	-
2000.....	-	-	-	3	4	-	-	1	-	1	-	-
1999.....	-	1	-	1	-	1	2	1	-	-	-	-
All races, female												
2004.....	-	-	-	1	-	1	-	1	1	1	-	-
2003.....	-	-	2	-	-	1	2	-	1	-	-	-
2002.....	-	-	1	1	2	1	-	2	3	-	-	-
2001.....	1	-	-	1	2	3	2	3	2	-	-	-
2000.....	-	-	1	-	-	-	1	2	1	-	-	-
1999.....	-	-	-	-	2	1	1	3	-	1	-	-
White, both sexes												
2004.....	1	-	-	1	-	1	1	1	1	-	-	-
2003.....	-	-	1	-	2	2	2	-	2	-	-	-
2002.....	1	-	1	2	3	2	1	2	3	-	-	-
2001.....	1	1	1	1	3	7	5	3	2	-	-	-
2000.....	-	-	1	1	3	-	-	3	1	1	-	-
1999.....	-	1	-	1	2	2	3	4	-	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Carcinoma in situ of breast and genitourinary system (D05-D07)												
All races, both sexes												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	1	-	2	-	-	-
2002.....	-	-	1	-	-	1	-	1	1	-	-	-
2001.....	-	-	-	1	2	3	-	1	-	-	-	-
2000.....	-	-	-	-	-	-	-	1	1	-	-	-
1999.....	-	-	-	1	1	1	-	1	-	-	-	-
All races, male												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	1	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	1	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	1	-	-	-	-
1999.....	-	-	-	1	-	-	-	-	-	-	-	-
All races, female												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	1	-	1	-	-	-
2002.....	-	-	1	-	-	1	-	1	1	-	-	-
2001.....	-	-	-	1	2	2	-	1	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	1	-	-	-
1999.....	-	-	-	-	1	1	-	1	-	-	-	-
White, both sexes												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	1	-	2	-	-	-
2002.....	-	-	1	-	-	-	-	1	1	-	-	-
2001.....	-	-	-	1	1	3	-	1	-	-	-	-
2000.....	-	-	-	-	-	-	-	1	1	-	-	-
1999.....	-	-	-	1	1	1	-	1	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Carcinoma in situ of other and unspecified sites (D00-D04,D09)												
All races, both sexes												
2004.....	1	-	-	1	-	2	1	1	1	1	-	-
2003.....	-	-	2	1	2	2	1	-	1	-	-	-
2002.....	1	-	-	2	3	2	1	1	2	-	-	-
2001.....	1	1	1	1	3	4	5	4	3	-	-	-
2000.....	-	-	1	3	4	-	1	2	-	1	-	-
1999.....	-	1	-	-	1	1	3	3	-	1	-	-
All races, male												
2004.....	1	-	-	-	-	1	1	-	-	-	-	-
2003.....	-	-	-	1	2	1	-	-	1	-	-	-
2002.....	1	-	-	1	1	2	1	-	-	-	-	-
2001.....	-	1	1	1	3	3	3	2	1	-	-	-
2000.....	-	-	-	3	4	-	-	-	-	1	-	-
1999.....	-	1	-	-	-	1	2	1	-	-	-	-
All races, female												
2004.....	-	-	-	1	-	1	-	1	1	1	-	-
2003.....	-	-	2	-	-	1	1	-	-	-	-	-
2002.....	-	-	-	1	2	-	-	1	2	-	-	-
2001.....	1	-	-	-	-	1	2	2	2	-	-	-
2000.....	-	-	1	-	-	-	1	2	-	-	-	-
1999.....	-	-	-	-	1	-	1	2	-	1	-	-
White, both sexes												
2004.....	1	-	-	1	-	1	1	1	1	-	-	-
2003.....	-	-	1	-	2	2	1	-	-	-	-	-
2002.....	1	-	-	2	3	2	1	1	2	-	-	-
2001.....	1	1	1	-	2	4	5	2	2	-	-	-
2000.....	-	-	1	1	3	-	-	2	-	1	-	-
1999.....	-	1	-	-	1	1	3	3	-	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Benign neoplasms (D10-D36)												
All races, both sexes												
2004.....	66	89	98	93	143	221	251	223	120	37	3	1
2003.....	68	79	92	110	142	209	265	219	103	21	6	-
2002.....	75	76	88	137	159	233	228	209	115	29	3	-
2001.....	70	86	99	121	146	213	237	227	113	36	6	-
2000.....	72	57	78	114	166	241	274	222	130	22	6	-
1999.....	58	82	96	128	163	253	235	197	121	29	2	-
All races, male												
2004.....	27	38	38	39	58	91	89	75	28	9	-	1
2003.....	35	36	48	50	70	85	100	66	34	5	1	-
2002.....	32	37	45	62	72	95	77	62	22	2	1	-
2001.....	32	43	51	57	61	95	99	77	30	9	-	-
2000.....	36	24	26	58	66	97	90	67	33	6	1	-
1999.....	28	41	44	49	81	92	83	69	31	9	1	-
All races, female												
2004.....	39	51	60	54	85	130	162	148	92	28	3	-
2003.....	33	43	44	60	72	124	165	153	69	16	5	-
2002.....	43	39	43	75	87	138	151	147	93	27	2	-
2001.....	38	43	48	64	85	118	138	150	83	27	6	-
2000.....	36	33	52	56	100	144	184	155	97	16	5	-
1999.....	30	41	52	79	82	161	152	128	90	20	1	-
White, both sexes												
2004.....	48	64	84	69	112	183	219	201	106	31	3	1
2003.....	56	62	79	82	114	188	236	201	88	20	4	-
2002.....	56	61	70	109	135	203	195	180	104	20	2	-
2001.....	55	62	69	91	126	187	202	197	97	27	4	-
2000.....	62	47	67	87	134	211	246	201	114	20	6	-
1999.....	41	58	74	100	136	214	207	175	114	26	2	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Benign neoplasms (D10-D36)												
White, male												
2004.....	22	26	33	31	50	74	75	67	27	6	-	1
2003.....	27	27	39	38	56	74	88	61	29	5	-	-
2002.....	28	30	33	49	65	82	66	52	19	-	1	-
2001.....	27	29	35	46	51	88	82	66	26	7	-	-
2000.....	31	20	18	42	52	88	83	59	30	6	1	-
1999.....	19	27	34	41	71	75	73	62	29	8	1	-
White, female												
2004.....	26	38	51	38	62	109	144	134	79	25	3	-
2003.....	29	35	40	44	58	114	148	140	59	15	4	-
2002.....	28	31	37	60	70	121	129	128	85	20	1	-
2001.....	28	33	34	45	75	99	120	131	71	20	4	-
2000.....	31	27	49	45	82	123	163	142	84	14	5	-
1999.....	22	31	40	59	65	139	134	113	85	18	1	-
Black, both sexes												
2004.....	14	22	9	20	26	34	24	21	9	6	-	-
2003.....	11	14	11	24	25	15	25	13	12	1	1	-
2002.....	15	12	15	25	19	26	27	23	8	6	1	-
2001.....	14	19	26	28	15	22	30	24	15	9	2	-
2000.....	9	10	10	20	25	27	24	17	14	2	-	-
1999.....	14	24	18	24	22	35	25	18	6	3	-	-
Black, male												
2004.....	5	11	4	7	7	17	10	8	-	3	-	-
2003.....	7	7	8	12	13	8	10	1	5	-	-	-
2002.....	3	6	10	11	6	11	9	7	2	2	-	-
2001.....	5	10	14	10	9	6	15	9	4	2	-	-
2000.....	5	4	7	12	8	9	5	6	3	-	-	-
1999.....	7	14	8	7	8	16	8	6	1	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Benign neoplasms (D10-D36)												
Black, female												
2004.....	9	11	5	13	19	17	14	13	9	3	-	-
2003.....	4	7	3	12	12	7	15	12	7	1	1	-
2002.....	12	6	5	14	13	15	18	16	6	4	1	-
2001.....	9	9	12	18	6	16	15	15	11	7	2	-
2000.....	4	6	3	8	17	18	19	11	11	2	-	-
1999.....	7	10	10	17	14	19	17	12	5	2	-	-
American Indian 1/, both sexes												
2004.....	-	-	1	-	-	-	-	-	1	-	-	-
2003.....	-	-	1	1	1	-	-	1	-	-	-	-
2002.....	1	-	1	-	1	1	1	-	-	-	-	-
2001.....	-	1	-	-	-	-	-	1	-	-	-	-
2000.....	-	-	1	2	1	-	-	2	-	-	-	-
1999.....	1	-	-	1	1	-	1	-	-	-	-	-
American Indian 1/, male												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	1	-	-	-	-
2002.....	-	-	1	-	-	1	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	1	1	1	-	-	-	-	-	-	-
1999.....	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, female												
2004.....	-	-	1	-	-	-	-	-	1	-	-	-
2003.....	-	-	1	1	1	-	-	-	-	-	-	-
2002.....	1	-	-	-	1	-	1	-	-	-	-	-
2001.....	-	1	-	-	-	-	-	1	-	-	-	-
2000.....	-	-	-	1	-	-	-	2	-	-	-	-
1999.....	1	-	-	1	-	-	1	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Benign neoplasms (D10-D36)												
Asian or Pacific Islander 2/, both sexes												
2004.....	4	3	4	4	5	4	8	1	4	-	-	-
2003.....	1	3	1	3	2	6	4	4	3	-	1	-
2002.....	3	3	2	3	4	3	5	6	3	3	-	-
2001.....	1	4	4	2	5	4	5	5	1	-	-	-
2000.....	1	-	-	5	6	3	4	2	2	-	-	-
1999.....	2	-	4	3	4	4	2	4	1	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	-	1	1	1	1	-	4	-	1	-	-	-
2003.....	1	2	1	-	1	3	2	3	-	-	1	-
2002.....	1	1	1	2	1	1	2	3	1	-	-	-
2001.....	-	4	2	1	1	1	2	2	-	-	-	-
2000.....	-	-	-	3	5	-	2	2	-	-	-	-
1999.....	2	-	2	1	1	1	2	1	1	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	4	2	3	3	4	4	4	1	3	-	-	-
2003.....	-	1	-	3	1	3	2	1	3	-	-	-
2002.....	2	2	1	1	3	2	3	3	2	3	-	-
2001.....	1	-	2	1	4	3	3	3	1	-	-	-
2000.....	1	-	-	2	1	3	2	-	2	-	-	-
1999.....	-	-	2	2	3	3	-	3	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Of female genital organs (D25-D28)												
All races, both sexes												
2004.....	3	3	2	2	2	4	2	3	1	-	-	-
2003.....	1	2	1	1	2	4	5	1	3	-	-	-
2002.....	7	2	2	2	6	1	1	2	3	-	-	-
2001.....	4	1	2	3	1	3	3	2	4	2	2	-
2000.....	8	-	3	1	5	1	3	-	3	1	-	-
1999.....	5	2	-	2	1	2	3	1	2	1	-	-
All races, male												
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....
All races, female												
2004.....	3	3	2	2	2	4	2	3	1	-	-	-
2003.....	1	2	1	1	2	4	5	1	3	-	-	-
2002.....	7	2	2	2	6	1	1	2	3	-	-	-
2001.....	4	1	2	3	1	3	3	2	4	2	2	-
2000.....	8	-	3	1	5	1	3	-	3	1	-	-
1999.....	5	2	-	2	1	2	3	1	2	1	-	-
White, both sexes												
2004.....	-	1	1	1	2	4	1	3	1	-	-	-
2003.....	-	1	1	1	2	3	4	1	3	-	-	-
2002.....	3	2	-	2	6	1	1	2	3	-	-	-
2001.....	3	1	1	2	1	2	3	2	3	1	-	-
2000.....	6	-	3	-	4	1	2	-	2	1	-	-
1999.....	4	1	-	1	1	2	2	-	2	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Of female genital organs (D25-D28)

Asian or Pacific Islander 2/, both sexes

2004.....	-	-	1	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	1	-	-	-	-	-	-
2002.....	-	-	1	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	1	-	-	-	1	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	1	-	-	-	-

Asian or Pacific Islander 2/, male

2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

Asian or Pacific Islander 2/, female

2004.....	-	-	1	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	1	-	-	-	-	-	-
2002.....	-	-	1	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	1	-	-	-	1	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	1	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Of eye, brain and other parts of central nervous system (D31-D33)

All races, both sexes

2004.....	40	47	57	57	100	126	192	157	83	30	2	-
2003.....	36	43	55	74	96	137	182	162	73	15	6	-
2002.....	43	42	50	85	102	155	156	140	73	19	-	-
2001.....	33	46	52	74	105	137	153	157	81	19	4	-
2000.....	33	33	51	71	105	168	185	157	90	14	3	-
1999.....	30	51	59	90	102	164	170	125	84	19	1	-

All races, male

2004.....	18	15	15	17	39	46	67	52	20	6	-	-
2003.....	19	21	25	33	42	49	70	49	20	5	1	-
2002.....	18	16	23	35	45	53	49	34	10	-	-	-
2001.....	12	22	22	35	40	56	52	48	18	4	-	-
2000.....	16	10	16	36	41	62	60	45	23	3	1	-
1999.....	16	22	20	33	46	57	57	39	20	6	1	-

All races, female

2004.....	22	32	42	40	61	80	125	105	63	24	2	-
2003.....	17	22	30	41	54	88	112	113	53	10	5	-
2002.....	25	26	27	50	57	102	107	106	63	19	-	-
2001.....	21	24	30	39	65	81	101	109	63	15	4	-
2000.....	17	23	35	35	64	106	125	112	67	11	2	-
1999.....	14	29	39	57	56	107	113	86	64	13	-	-

White, both sexes

2004.....	31	34	45	37	77	104	165	142	77	27	2	-
2003.....	31	32	49	51	75	124	164	148	63	14	4	-
2002.....	33	34	42	70	88	131	132	122	67	16	-	-
2001.....	26	29	34	54	89	118	132	135	68	13	4	-
2000.....	28	26	42	56	88	147	165	144	81	13	3	-
1999.....	21	36	46	72	85	139	149	113	79	17	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Of eye, brain and other parts of central nervous system (D31-D33)												
White, male												
2004.....	15	10	10	11	33	38	55	46	20	5	-	-
2003.....	16	15	21	24	33	44	64	46	18	5	-	-
2002.....	15	13	17	29	42	44	41	28	9	-	-	-
2001.....	11	12	14	29	34	51	44	41	15	2	-	-
2000.....	14	7	10	27	33	58	54	40	22	3	1	-
1999.....	10	15	17	31	42	46	50	35	18	5	1	-
White, female												
2004.....	16	24	35	26	44	66	110	96	57	22	2	-
2003.....	15	17	28	27	42	80	100	102	45	9	4	-
2002.....	18	21	25	41	46	87	91	94	58	16	-	-
2001.....	15	17	20	25	55	67	88	94	53	11	4	-
2000.....	14	19	32	29	55	89	111	104	59	10	2	-
1999.....	11	21	29	41	43	93	99	78	61	12	-	-
Black, both sexes												
2004.....	7	12	9	16	20	19	20	15	3	3	-	-
2003.....	5	9	4	19	19	9	17	11	8	1	1	-
2002.....	8	6	6	14	12	20	18	14	4	2	-	-
2001.....	6	14	16	19	11	16	18	18	12	6	-	-
2000.....	5	7	9	10	12	19	17	10	7	1	-	-
1999.....	8	15	12	16	16	22	18	11	4	2	-	-
Black, male												
2004.....	3	5	4	5	6	8	9	6	-	1	-	-
2003.....	3	5	3	9	9	3	6	-	2	-	-	-
2002.....	2	3	4	6	3	7	6	4	-	-	-	-
2001.....	1	8	7	6	5	4	7	5	3	2	-	-
2000.....	2	3	6	6	3	4	4	4	1	-	-	-
1999.....	5	7	3	2	4	11	5	3	1	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Of eye, brain and other parts of central nervous system (D31-D33)												
Black, female												
2004.....	4	7	5	11	14	11	11	9	3	2	-	-
2003.....	2	4	1	10	10	6	11	11	6	1	1	-
2002.....	6	3	2	8	9	13	12	10	4	2	-	-
2001.....	5	6	9	13	6	12	11	13	9	4	-	-
2000.....	3	4	3	4	9	15	13	6	6	1	-	-
1999.....	3	8	9	14	12	11	13	8	3	1	-	-
American Indian 1/, both sexes												
2004.....	-	-	-	-	-	-	-	-	1	-	-	-
2003.....	-	-	1	1	1	-	-	1	-	-	-	-
2002.....	-	-	1	-	1	1	1	-	-	-	-	-
2001.....	-	1	-	-	-	-	-	1	-	-	-	-
2000.....	-	-	-	1	1	-	-	2	-	-	-	-
1999.....	-	-	-	1	-	-	1	-	-	-	-	-
American Indian 1/, male												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	1	-	-	-	-
2002.....	-	-	1	-	-	1	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	1	1	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, female												
2004.....	-	-	-	-	-	-	-	-	1	-	-	-
2003.....	-	-	1	1	1	-	-	-	-	-	-	-
2002.....	-	-	-	-	1	-	1	-	-	-	-	-
2001.....	-	1	-	-	-	-	-	1	-	-	-	-
2000.....	-	-	-	-	-	-	-	2	-	-	-	-
1999.....	-	-	-	1	-	-	1	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Of eye, brain and other parts of central nervous system (D31-D33)												
Asian or Pacific Islander 2/, both sexes												
2004.....	2	1	3	4	3	3	7	-	2	-	-	-
2003.....	-	2	1	3	1	4	1	2	2	-	1	-
2002.....	2	2	1	1	1	3	5	4	2	1	-	-
2001.....	1	2	2	1	5	3	3	3	1	-	-	-
2000.....	-	-	-	4	4	2	3	1	2	-	-	-
1999.....	1	-	1	1	1	3	2	1	1	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	-	-	1	1	-	-	3	-	-	-	-	-
2003.....	-	1	1	-	-	2	-	2	-	-	1	-
2002.....	1	-	1	-	-	1	2	2	1	-	-	-
2001.....	-	2	1	-	1	1	1	2	-	-	-	-
2000.....	-	-	-	2	4	-	2	1	-	-	-	-
1999.....	1	-	-	-	-	-	2	1	1	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	2	1	2	3	3	3	4	-	2	-	-	-
2003.....	-	1	-	3	1	2	1	-	2	-	-	-
2002.....	1	2	-	1	1	2	3	2	1	1	-	-
2001.....	1	-	1	1	4	2	2	1	1	-	-	-
2000.....	-	-	-	2	-	2	1	-	2	-	-	-
1999.....	-	-	1	1	1	3	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Of other and unspecified sites (D10-D24,D29-D30,D34-D36)

All races, both sexes

2004.....	23	39	39	34	41	91	57	63	36	7	1	1
2003.....	31	34	36	35	44	68	78	56	27	6	-	-
2002.....	25	32	36	50	51	77	71	67	39	10	3	-
2001.....	33	39	45	44	40	73	81	68	28	15	-	-
2000.....	31	24	24	42	56	72	86	65	37	7	3	-
1999.....	23	29	37	36	60	87	62	71	35	9	1	-

All races, male

2004.....	9	23	23	22	19	45	22	23	8	3	-	1
2003.....	16	15	23	17	28	36	30	17	14	-	-	-
2002.....	14	21	22	27	27	42	28	28	12	2	1	-
2001.....	20	21	29	22	21	39	47	29	12	5	-	-
2000.....	20	14	10	22	25	35	30	22	10	3	-	-
1999.....	12	19	24	16	35	35	26	30	11	3	-	-

All races, female

2004.....	14	16	16	12	22	46	35	40	28	4	1	-
2003.....	15	19	13	18	16	32	48	39	13	6	-	-
2002.....	11	11	14	23	24	35	43	39	27	8	2	-
2001.....	13	18	16	22	19	34	34	39	16	10	-	-
2000.....	11	10	14	20	31	37	56	43	27	4	3	-
1999.....	11	10	13	20	25	52	36	41	24	6	1	-

White, both sexes

2004.....	17	29	38	31	33	75	53	56	28	4	1	1
2003.....	25	29	29	30	37	61	68	52	22	6	-	-
2002.....	20	25	28	37	41	71	62	56	34	4	2	-
2001.....	26	32	34	35	36	67	67	60	26	13	-	-
2000.....	28	21	22	31	42	63	79	57	31	6	3	-
1999.....	16	21	28	27	50	73	56	62	33	9	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Of other and unspecified sites (D10-D24,D29-D30,D34-D36)												
White, male												
2004.....	7	16	23	20	17	36	20	21	7	1	-	1
2003.....	11	12	18	14	23	30	24	15	11	-	-	-
2002.....	13	17	16	20	23	38	25	24	10	-	1	-
2001.....	16	17	21	17	17	37	38	25	11	5	-	-
2000.....	17	13	8	15	19	30	29	19	8	3	-	-
1999.....	9	12	17	10	29	29	23	27	11	3	-	-
White, female												
2004.....	10	13	15	11	16	39	33	35	21	3	1	-
2003.....	14	17	11	16	14	31	44	37	11	6	-	-
2002.....	7	8	12	17	18	33	37	32	24	4	1	-
2001.....	10	15	13	18	19	30	29	35	15	8	-	-
2000.....	11	8	14	16	23	33	50	38	23	3	3	-
1999.....	7	9	11	17	21	44	33	35	22	6	1	-
Black, both sexes												
2004.....	4	8	-	3	6	15	3	6	6	3	-	-
2003.....	5	4	7	5	6	6	7	2	4	-	-	-
2002.....	3	6	8	11	7	6	9	9	4	4	1	-
2001.....	7	5	9	8	4	5	12	6	2	2	-	-
2000.....	3	3	1	9	13	8	6	7	6	1	-	-
1999.....	5	8	6	7	6	13	6	7	2	-	-	-
Black, male												
2004.....	2	6	-	2	1	9	1	2	-	2	-	-
2003.....	4	2	5	3	4	5	4	1	3	-	-	-
2002.....	1	3	6	5	3	4	3	3	2	2	-	-
2001.....	4	2	7	4	4	2	8	4	1	-	-	-
2000.....	3	1	1	6	5	5	1	2	2	-	-	-
1999.....	2	7	5	5	4	5	3	3	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Of other and unspecified sites (D10-D24,D29-D30,D34-D36)

Asian or Pacific Islander 2/, both sexes

2004.....	2	2	-	-	2	1	1	1	2	-	-	-
2003.....	1	1	-	-	1	1	3	2	1	-	-	-
2002.....	1	1	-	2	3	-	-	2	1	2	-	-
2001.....	-	2	2	1	-	1	2	2	-	-	-	-
2000.....	-	-	-	1	1	1	1	1	-	-	-	-
1999.....	1	-	3	2	3	1	-	2	-	-	-	-

Asian or Pacific Islander 2/, male

2004.....	-	1	-	-	1	-	1	-	1	-	-	-
2003.....	1	1	-	-	1	1	2	1	-	-	-	-
2002.....	-	1	-	2	1	-	-	1	-	-	-	-
2001.....	-	2	1	1	-	-	1	-	-	-	-	-
2000.....	-	-	-	1	1	-	-	1	-	-	-	-
1999.....	1	-	2	1	1	1	-	-	-	-	-	-

Asian or Pacific Islander 2/, female

2004.....	2	1	-	-	1	1	-	1	1	-	-	-
2003.....	-	-	-	-	-	-	1	1	1	-	-	-
2002.....	1	-	-	-	2	-	-	1	1	2	-	-
2001.....	-	-	1	-	-	1	1	2	-	-	-	-
2000.....	-	-	-	-	-	1	1	-	-	-	-	-
1999.....	-	-	1	1	2	-	-	2	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Neoplasms of uncertain or unknown behavior of specified sites (D37-D47)												
All races, both sexes												
2004.....	285	441	591	784	1,215	1,715	2,274	1,850	975	324	38	1
2003.....	292	401	607	805	1,209	1,845	2,216	1,901	1,049	299	32	-
2002.....	284	455	507	754	1,214	1,832	2,128	1,759	987	305	27	-
2001.....	312	432	572	832	1,368	1,904	2,139	1,793	970	286	38	-
2000.....	310	426	592	860	1,319	1,821	2,135	1,799	956	264	32	1
1999.....	302	360	542	870	1,437	1,831	2,009	1,728	886	287	41	-
All races, male												
2004.....	173	258	368	485	687	954	1,173	844	376	98	8	1
2003.....	171	228	343	456	718	997	1,134	864	364	88	8	-
2002.....	155	280	291	464	676	1,020	1,079	813	364	94	6	-
2001.....	170	223	331	481	789	1,072	1,075	787	368	90	10	-
2000.....	183	233	343	511	739	991	1,079	798	318	75	6	1
1999.....	180	205	338	520	827	1,000	992	755	328	71	13	-
All races, female												
2004.....	112	183	223	299	528	761	1,101	1,006	599	226	30	-
2003.....	121	173	264	349	491	848	1,082	1,037	685	211	24	-
2002.....	129	175	216	290	538	812	1,049	946	623	211	21	-
2001.....	142	209	241	351	579	832	1,064	1,006	602	196	28	-
2000.....	127	193	249	349	580	830	1,056	1,001	638	189	26	-
1999.....	122	155	204	350	610	831	1,017	973	558	216	28	-
White, both sexes												
2004.....	227	370	501	695	1,095	1,561	2,117	1,732	920	303	33	1
2003.....	239	338	532	708	1,100	1,706	2,057	1,788	982	276	24	-
2002.....	234	381	450	672	1,103	1,693	1,969	1,663	915	275	23	-
2001.....	270	385	504	756	1,233	1,761	1,993	1,667	899	259	30	-
2000.....	258	376	510	772	1,173	1,683	2,009	1,682	897	244	27	-
1999.....	258	315	471	760	1,315	1,688	1,883	1,630	838	258	37	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Neoplasms of uncertain or unknown behavior of specified sites (D37-D47)												
White, male												
2004.....	139	216	314	436	622	878	1,103	797	354	95	8	1
2003.....	145	190	301	411	658	936	1,058	822	340	80	7	-
2002.....	127	234	263	415	624	948	1,001	775	336	82	6	-
2001.....	153	203	293	440	728	994	1,004	730	345	84	8	-
2000.....	156	209	292	461	660	917	1,025	744	300	71	6	-
1999.....	156	181	303	456	765	930	939	716	315	66	12	-
White, female												
2004.....	88	154	187	259	473	683	1,014	935	566	208	25	-
2003.....	94	148	231	297	442	770	999	966	642	196	17	-
2002.....	107	147	187	257	479	745	968	888	579	193	17	-
2001.....	117	182	211	316	505	767	989	937	554	175	22	-
2000.....	102	167	218	311	513	766	984	938	597	173	21	-
1999.....	102	134	168	304	550	758	944	914	523	192	25	-
Black, both sexes												
2004.....	46	63	72	72	92	116	124	83	45	17	5	-
2003.....	45	50	62	74	90	112	128	81	53	19	7	-
2002.....	41	63	49	69	89	112	127	81	57	27	3	-
2001.....	38	40	57	64	110	114	123	91	60	23	7	-
2000.....	47	41	62	72	117	113	101	96	47	18	4	1
1999.....	37	38	60	95	105	121	105	84	41	25	3	-
Black, male												
2004.....	27	38	43	40	51	57	50	31	17	1	-	-
2003.....	21	31	34	36	52	47	61	31	16	6	1	-
2002.....	23	42	21	38	40	54	59	33	20	9	-	-
2001.....	15	17	31	36	53	59	58	38	18	5	2	-
2000.....	25	21	38	44	58	59	39	38	15	3	-	1
1999.....	19	21	28	56	52	55	40	31	9	5	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Neoplasms of uncertain or unknown behavior of specified sites (D37-D47)

Black, female

2004.....	19	25	29	32	41	59	74	52	28	16	5	-
2003.....	24	19	28	38	38	65	67	50	37	13	6	-
2002.....	18	21	28	31	49	58	68	48	37	18	3	-
2001.....	23	23	26	28	57	55	65	53	42	18	5	-
2000.....	22	20	24	28	59	54	62	58	32	15	4	-
1999.....	18	17	32	39	53	66	65	53	32	20	3	-

American Indian 1/, both sexes

2004.....	3	1	5	2	7	7	2	6	2	1	-	-
2003.....	-	3	2	5	2	4	5	3	1	1	1	-
2002.....	1	1	2	2	6	3	3	1	1	1	-	-
2001.....	-	1	1	4	11	9	4	6	1	-	-	-
2000.....	1	2	3	2	5	3	2	4	1	1	1	-
1999.....	1	4	3	6	5	3	4	4	1	1	1	-

American Indian 1/, male

2004.....	1	-	2	1	4	5	-	2	2	1	-	-
2003.....	-	2	2	1	2	3	3	1	1	-	-	-
2002.....	-	-	2	2	2	3	1	1	-	1	-	-
2001.....	-	1	1	3	3	4	2	4	1	-	-	-
2000.....	-	-	2	-	4	1	2	2	-	-	-	-
1999.....	1	2	2	2	2	3	3	2	-	-	1	-

American Indian 1/, female

2004.....	2	1	3	1	3	2	2	4	-	-	-	-
2003.....	-	1	-	4	-	1	2	2	-	1	1	-
2002.....	1	1	-	-	4	-	2	-	1	-	-	-
2001.....	-	-	-	1	8	5	2	2	-	-	-	-
2000.....	1	2	1	2	1	2	-	2	1	1	1	-
1999.....	-	2	1	4	3	-	1	2	1	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Neoplasms of uncertain or unknown behavior of specified sites (D37-D47)												
Asian or Pacific Islander 2/, both sexes												
2004.....	9	7	13	15	21	31	31	29	8	3	-	-
2003.....	8	10	11	18	17	23	26	29	13	3	-	-
2002.....	8	10	6	11	16	24	29	14	14	2	1	-
2001.....	4	6	10	8	14	20	19	29	10	4	1	-
2000.....	4	7	17	14	24	22	23	17	11	1	-	-
1999.....	6	3	8	9	12	19	17	10	6	3	-	-
Asian or Pacific Islander 2/, male												
2004.....	6	4	9	8	10	14	20	14	3	1	-	-
2003.....	5	5	6	8	6	11	12	10	7	2	-	-
2002.....	5	4	5	9	10	15	18	4	8	2	-	-
2001.....	2	2	6	2	5	15	11	15	4	1	-	-
2000.....	2	3	11	6	17	14	13	14	3	1	-	-
1999.....	4	1	5	6	8	12	10	6	4	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	3	3	4	7	11	17	11	15	5	2	-	-
2003.....	3	5	5	10	11	12	14	19	6	1	-	-
2002.....	3	6	1	2	6	9	11	10	6	-	1	-
2001.....	2	4	4	6	9	5	8	14	6	3	1	-
2000.....	2	4	6	8	7	8	10	3	8	-	-	-
1999.....	2	2	3	3	4	7	7	4	2	3	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Neoplasms of uncertain or unknown behavior of unspecified sites (D48)

All races, both sexes

2004.....	38	48	43	43	64	83	97	98	80	20	3	-
2003.....	27	26	31	39	46	67	83	83	60	17	2	-
2002.....	27	30	36	35	44	69	103	82	57	32	3	-
2001.....	32	34	28	45	55	68	81	77	55	29	5	-
2000.....	15	30	40	30	50	80	93	79	56	27	2	-
1999.....	26	18	34	30	53	65	74	69	52	20	6	-

All races, male

2004.....	23	27	21	20	33	32	37	27	22	4	1	-
2003.....	12	15	15	17	20	31	31	25	11	1	1	-
2002.....	8	13	14	16	23	28	30	25	12	3	1	-
2001.....	16	12	11	24	23	28	27	16	5	3	-	-
2000.....	6	13	18	15	19	32	30	15	11	4	-	-
1999.....	12	8	19	20	24	26	22	14	8	2	1	-

All races, female

2004.....	15	21	22	23	31	51	60	71	58	16	2	-
2003.....	15	11	16	22	26	36	52	58	49	16	1	-
2002.....	19	17	22	19	21	41	73	57	45	29	2	-
2001.....	16	22	17	21	32	40	54	61	50	26	5	-
2000.....	9	17	22	15	31	48	63	64	45	23	2	-
1999.....	14	10	15	10	29	39	52	55	44	18	5	-

White, both sexes

2004.....	30	39	35	35	58	77	85	89	78	14	3	-
2003.....	23	23	26	30	45	61	79	76	55	15	2	-
2002.....	18	28	32	29	38	63	96	77	54	29	3	-
2001.....	27	28	25	36	42	62	75	74	52	28	5	-
2000.....	13	24	33	28	41	68	85	74	54	26	2	-
1999.....	21	15	27	25	46	55	68	61	44	18	5	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Neoplasms of uncertain or unknown behavior of unspecified sites (D48)												
White, male												
2004.....	19	25	17	16	31	30	31	25	22	3	1	-
2003.....	10	13	13	15	20	28	30	22	9	1	1	-
2002.....	6	11	11	14	20	28	27	23	12	3	1	-
2001.....	13	10	11	20	17	25	27	15	5	3	-	-
2000.....	5	11	14	13	15	27	28	14	11	4	-	-
1999.....	11	7	16	18	20	21	21	13	8	2	1	-
White, female												
2004.....	11	14	18	19	27	47	54	64	56	11	2	-
2003.....	13	10	13	15	25	33	49	54	46	14	1	-
2002.....	12	17	21	15	18	35	69	54	42	26	2	-
2001.....	14	18	14	16	25	37	48	59	47	25	5	-
2000.....	8	13	19	15	26	41	57	60	43	22	2	-
1999.....	10	8	11	7	26	34	47	48	36	16	4	-
Black, both sexes												
2004.....	8	8	6	7	5	3	11	7	2	6	-	-
2003.....	3	2	4	8	1	6	4	4	4	2	-	-
2002.....	8	2	3	6	4	3	6	4	3	1	-	-
2001.....	3	5	2	8	11	5	4	2	2	1	-	-
2000.....	2	6	6	2	9	10	8	5	2	-	-	-
1999.....	5	3	6	5	7	9	4	6	5	2	1	-
Black, male												
2004.....	4	1	3	4	2	1	5	1	-	1	-	-
2003.....	1	1	2	1	-	3	1	2	1	-	-	-
2002.....	1	2	2	2	2	-	2	1	-	-	-	-
2001.....	2	2	-	4	5	3	-	-	-	-	-	-
2000.....	1	2	4	2	4	5	2	1	-	-	-	-
1999.....	1	1	3	2	4	5	1	1	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Neoplasms of uncertain or unknown behavior of unspecified sites (D48)												
Black, female												
2004.....	4	7	3	3	3	2	6	6	2	5	-	-
2003.....	2	1	2	7	1	3	3	2	3	2	-	-
2002.....	7	-	1	4	2	3	4	3	3	1	-	-
2001.....	1	3	2	4	6	2	4	2	2	1	-	-
2000.....	1	4	2	-	5	5	6	4	2	-	-	-
1999.....	4	2	3	3	3	4	3	5	5	2	1	-
American Indian 1/, both sexes												
2004.....	-	-	-	-	-	-	-	1	-	-	-	-
2003.....	-	1	1	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	1	-	-
2001.....	1	-	-	-	1	1	-	-	1	-	-	-
2000.....	-	-	-	-	-	2	-	-	-	-	-	-
1999.....	-	-	-	-	-	1	-	-	1	-	-	-
American Indian 1/, male												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	1	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	1	-	-	-	1	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, female												
2004.....	-	-	-	-	-	-	-	1	-	-	-	-
2003.....	-	-	1	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	1	-	-
2001.....	-	-	-	-	-	1	-	-	1	-	-	-
2000.....	-	-	-	-	-	2	-	-	-	-	-	-
1999.....	-	-	-	-	-	1	-	-	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Neoplasms of uncertain or unknown behavior of unspecified sites (D48)												
Asian or Pacific Islander 2/, both sexes												
2004.....	-	1	2	1	1	3	1	1	-	-	-	-
2003.....	1	-	-	1	-	-	-	3	1	-	-	-
2002.....	1	-	1	-	2	3	1	1	-	1	-	-
2001.....	1	1	1	1	1	-	2	1	-	-	-	-
2000.....	-	-	1	-	-	-	-	-	-	1	-	-
1999.....	-	-	1	-	-	-	2	2	2	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	-	1	1	-	-	1	1	1	-	-	-	-
2003.....	1	-	-	1	-	-	-	1	1	-	-	-
2002.....	1	-	1	-	1	-	1	1	-	-	-	-
2001.....	-	-	-	-	-	-	-	1	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	-	-	1	1	1	2	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	2	-	-	-	-
2002.....	-	-	-	-	1	3	-	-	-	1	-	-
2001.....	1	1	1	1	1	-	2	-	-	-	-	-
2000.....	-	-	1	-	-	-	-	-	-	1	-	-
1999.....	-	-	1	-	-	-	2	2	2	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
III. Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)												
All races, both sexes												
2004.....	473	493	533	569	732	981	1,266	1,197	896	385	93	1
2003.....	488	477	499	567	784	1,015	1,248	1,224	948	419	83	-
2002.....	484	456	499	571	815	1,032	1,310	1,232	890	395	91	-
2001.....	447	432	473	577	766	1,114	1,262	1,234	870	367	105	-
2000.....	440	424	467	611	781	1,048	1,280	1,208	831	367	91	-
1999.....	360	376	415	517	822	1,062	1,158	1,170	855	389	92	-
All races, male												
2004.....	232	263	261	278	339	422	522	442	246	82	16	1
2003.....	233	218	243	274	385	459	512	430	263	94	9	-
2002.....	234	222	249	244	392	455	504	426	251	102	16	-
2001.....	216	186	219	259	336	468	527	435	236	68	16	-
2000.....	207	191	215	288	381	433	501	405	243	75	18	-
1999.....	173	171	185	239	384	474	457	410	238	91	15	-
All races, female												
2004.....	241	230	272	291	393	559	744	755	650	303	77	-
2003.....	255	259	256	293	399	556	736	794	685	325	74	-
2002.....	250	234	250	327	423	577	806	806	639	293	75	-
2001.....	231	246	254	318	430	646	735	799	634	299	89	-
2000.....	233	233	252	323	400	615	779	803	588	292	73	-
1999.....	187	205	230	278	438	588	701	760	617	298	77	-
White, both sexes												
2004.....	248	289	370	434	593	851	1,134	1,068	809	348	78	1
2003.....	241	273	340	418	648	863	1,104	1,111	838	380	74	-
2002.....	247	290	345	438	673	867	1,167	1,115	789	348	82	-
2001.....	240	277	347	449	628	963	1,109	1,116	796	328	83	-
2000.....	237	273	328	474	652	911	1,140	1,102	746	329	79	-
1999.....	195	219	298	393	681	924	1,042	1,052	772	344	76	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

III. Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)

White, male

2004.....	138	157	195	215	285	370	467	397	230	73	15	1
2003.....	118	125	168	213	320	393	465	392	236	83	9	-
2002.....	122	149	182	193	327	393	447	386	222	85	15	-
2001.....	124	124	166	209	274	409	473	404	215	62	12	-
2000.....	115	135	154	239	322	380	448	370	210	66	16	-
1999.....	101	106	131	189	321	415	408	367	222	83	14	-

White, female

2004.....	110	132	175	219	308	481	667	671	579	275	63	-
2003.....	123	148	172	205	328	470	639	719	602	297	65	-
2002.....	125	141	163	245	346	474	720	729	567	263	67	-
2001.....	116	153	181	240	354	554	636	712	581	266	71	-
2000.....	122	138	174	235	330	531	692	732	536	263	63	-
1999.....	94	113	167	204	360	509	634	685	550	261	62	-

Black, both sexes

2004.....	217	191	154	126	117	117	103	112	77	31	13	-
2003.....	234	189	142	132	125	129	127	99	100	33	6	-
2002.....	225	156	139	126	126	149	122	105	89	41	7	-
2001.....	200	141	113	114	125	132	132	97	62	33	19	-
2000.....	197	139	129	118	110	118	127	93	71	37	12	-
1999.....	154	148	108	110	115	125	101	99	74	36	14	-

Black, male

2004.....	92	100	63	60	47	46	37	41	13	7	1	-
2003.....	108	86	65	50	61	58	41	32	23	9	-	-
2002.....	107	70	61	49	58	54	47	35	26	11	1	-
2001.....	90	55	47	44	54	53	45	22	15	6	3	-
2000.....	91	49	57	42	47	42	43	29	25	8	2	-
1999.....	67	60	49	45	49	49	40	35	13	5	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

III. Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)

Black, female

2004.....	125	91	91	66	70	71	66	71	64	24	12	-
2003.....	126	103	77	82	64	71	86	67	77	24	6	-
2002.....	118	86	78	77	68	95	75	70	63	30	6	-
2001.....	110	86	66	70	71	79	87	75	47	27	16	-
2000.....	106	90	72	76	63	76	84	64	46	29	10	-
1999.....	87	88	59	65	66	76	61	64	61	31	13	-

American Indian 1/, both sexes

2004.....	3	2	1	4	3	2	2	2	3	-	-	-
2003.....	7	4	6	6	1	1	2	1	1	-	-	-
2002.....	6	5	2	1	2	2	1	1	1	1	1	-
2001.....	2	3	2	3	3	8	5	7	-	4	1	-
2000.....	1	6	1	5	3	9	2	2	3	-	-	-
1999.....	3	1	5	3	6	1	3	2	1	1	2	-

American Indian 1/, male

2004.....	1	2	1	1	1	-	2	-	1	-	-	-
2003.....	5	1	3	2	1	-	-	-	-	-	-	-
2002.....	3	2	1	-	-	-	1	-	-	1	-	-
2001.....	-	2	1	2	2	2	1	3	-	-	-	-
2000.....	1	2	-	1	1	8	1	-	-	-	-	-
1999.....	1	1	3	-	1	1	1	1	-	1	-	-

American Indian 1/, female

2004.....	2	-	-	3	2	2	-	2	2	-	-	-
2003.....	2	3	3	4	-	1	2	1	1	-	-	-
2002.....	3	3	1	1	2	2	-	1	1	-	1	-
2001.....	2	1	1	1	1	6	4	4	-	4	1	-
2000.....	-	4	1	4	2	1	1	2	3	-	-	-
1999.....	2	-	2	3	5	-	2	1	1	-	2	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
III. Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)												
Asian or Pacific Islander 2/, both sexes												
2004.....	5	11	8	5	19	11	27	15	7	6	2	-
2003.....	6	11	11	11	10	22	15	13	9	6	3	-
2002.....	6	5	13	6	14	14	20	11	11	5	1	-
2001.....	5	11	11	11	10	11	16	14	12	2	2	-
2000.....	5	6	9	14	16	10	11	11	11	1	-	-
1999.....	8	8	4	11	20	12	12	17	8	8	-	-
Asian or Pacific Islander 2/, male												
2004.....	1	4	2	2	6	6	16	4	2	2	-	-
2003.....	2	6	7	9	3	8	6	6	4	2	-	-
2002.....	2	1	5	2	7	8	9	5	3	5	-	-
2001.....	2	5	5	4	6	4	8	6	6	-	1	-
2000.....	-	5	4	6	11	3	9	6	8	1	-	-
1999.....	4	4	2	5	13	9	8	7	3	2	-	-
Asian or Pacific Islander 2/, female												
2004.....	4	7	6	3	13	5	11	11	5	4	2	-
2003.....	4	5	4	2	7	14	9	7	5	4	3	-
2002.....	4	4	8	4	7	6	11	6	8	-	1	-
2001.....	3	6	6	7	4	7	8	8	6	2	1	-
2000.....	5	1	5	8	5	7	2	5	3	-	-	-
1999.....	4	4	2	6	7	3	4	10	5	6	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Anemias (D50-D64)												
All races, both sexes												
2004.....	116	140	169	181	292	440	679	774	684	312	80	1
2003.....	104	121	155	184	300	447	689	784	721	342	72	-
2002.....	119	121	143	184	324	474	687	787	665	322	78	-
2001.....	120	119	130	186	306	525	701	808	659	294	90	-
2000.....	117	127	133	210	305	482	689	790	638	290	78	-
1999.....	98	105	126	175	311	497	667	775	647	316	82	-
All races, male												
2004.....	56	78	86	88	138	194	272	298	180	57	15	1
2003.....	48	67	89	104	147	203	274	270	196	75	8	-
2002.....	52	57	74	77	167	204	250	256	182	81	13	-
2001.....	58	53	58	91	146	220	276	276	177	52	15	-
2000.....	56	66	64	95	141	198	267	237	183	58	14	-
1999.....	48	43	62	92	134	217	267	261	172	73	12	-
All races, female												
2004.....	60	62	83	93	154	246	407	476	504	255	65	-
2003.....	56	54	66	80	153	244	415	514	525	267	64	-
2002.....	67	64	69	107	157	270	437	531	483	241	65	-
2001.....	62	66	72	95	160	305	425	532	482	242	75	-
2000.....	61	61	69	115	164	284	422	553	455	232	64	-
1999.....	50	62	64	83	177	280	400	514	475	243	70	-
White, both sexes												
2004.....	47	71	113	142	237	370	611	681	618	285	67	1
2003.....	44	56	97	135	239	379	606	718	638	306	65	-
2002.....	50	71	90	141	258	387	603	708	585	284	73	-
2001.....	61	69	99	146	247	446	613	728	599	266	69	-
2000.....	56	84	85	160	250	414	617	718	566	261	68	-
1999.....	41	46	76	135	254	428	604	692	583	277	68	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Anemias (D50-D64)												
White, male												
2004.....	26	35	60	70	116	159	243	265	168	52	14	1
2003.....	22	32	53	78	111	175	249	249	174	64	8	-
2002.....	26	36	47	61	128	173	216	231	157	68	12	-
2001.....	28	31	44	74	113	193	248	259	159	46	11	-
2000.....	27	45	42	81	118	169	238	212	156	52	13	-
1999.....	20	16	32	75	107	190	244	230	160	66	12	-
White, female												
2004.....	21	36	53	72	121	211	368	416	450	233	53	-
2003.....	22	24	44	57	128	204	357	469	464	242	57	-
2002.....	24	35	43	80	130	214	387	477	428	216	61	-
2001.....	33	38	55	72	134	253	365	469	440	220	58	-
2000.....	29	39	43	79	132	245	379	506	410	209	55	-
1999.....	21	30	44	60	147	238	360	462	423	211	56	-
Black, both sexes												
2004.....	67	65	51	36	47	63	54	79	59	22	11	-
2003.....	58	64	52	41	54	64	78	59	75	30	5	-
2002.....	65	47	47	41	59	80	74	70	68	34	4	-
2001.....	58	46	29	36	55	72	79	66	50	24	18	-
2000.....	59	40	43	43	46	58	64	64	61	28	10	-
1999.....	52	55	46	33	49	64	57	67	57	30	12	-
Black, male												
2004.....	30	40	24	17	19	30	20	30	9	3	1	-
2003.....	26	35	32	23	33	28	25	19	19	9	-	-
2002.....	25	20	26	16	35	28	31	21	22	9	1	-
2001.....	30	21	14	15	31	26	25	15	13	6	3	-
2000.....	29	18	19	13	18	23	23	20	22	5	1	-
1999.....	26	25	29	14	23	23	19	25	9	4	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Anemias (D50-D64)												
Black, female												
2004.....	37	25	27	19	28	33	34	49	50	19	10	-
2003.....	32	29	20	18	21	36	53	40	56	21	5	-
2002.....	40	27	21	25	24	52	43	49	46	25	3	-
2001.....	28	25	15	21	24	46	54	51	37	18	15	-
2000.....	30	22	24	30	28	35	41	44	39	23	9	-
1999.....	26	30	17	19	26	41	38	42	48	26	12	-
American Indian 1/, both sexes												
2004.....	-	1	-	1	2	1	2	2	2	-	-	-
2003.....	1	-	2	4	1	1	1	-	1	-	-	-
2002.....	2	2	-	-	-	1	1	1	1	1	1	-
2001.....	-	1	-	-	1	4	3	6	-	3	1	-
2000.....	-	1	-	2	2	6	2	1	3	-	-	-
1999.....	2	1	3	1	2	-	-	2	-	1	2	-
American Indian 1/, male												
2004.....	-	1	-	-	1	-	2	-	1	-	-	-
2003.....	-	-	1	1	1	-	-	-	-	-	-	-
2002.....	-	1	-	-	-	-	1	-	-	1	-	-
2001.....	-	1	-	-	1	1	-	2	-	-	-	-
2000.....	-	1	-	1	-	5	1	-	-	-	-	-
1999.....	-	1	1	-	-	-	-	1	-	1	-	-
American Indian 1/, female												
2004.....	-	-	-	1	1	1	-	2	1	-	-	-
2003.....	1	-	1	3	-	1	1	-	1	-	-	-
2002.....	2	1	-	-	-	1	-	1	1	-	1	-
2001.....	-	-	-	-	-	3	3	4	-	3	1	-
2000.....	-	-	-	1	2	1	1	1	3	-	-	-
1999.....	2	-	2	1	2	-	-	1	-	-	2	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Anemias (D50-D64)												
Asian or Pacific Islander 2/, both sexes												
2004.....	2	3	5	2	6	6	12	12	5	5	2	-
2003.....	1	1	4	4	6	3	4	7	7	6	2	-
2002.....	2	1	6	2	7	6	9	8	11	3	-	-
2001.....	1	3	2	4	3	3	6	8	10	1	2	-
2000.....	2	2	5	5	7	4	6	7	8	1	-	-
1999.....	3	3	1	6	6	5	6	14	7	8	-	-
Asian or Pacific Islander 2/, male												
2004.....	-	2	2	1	2	5	7	3	2	2	-	-
2003.....	-	-	3	2	2	-	-	2	3	2	-	-
2002.....	1	-	1	-	4	3	2	4	3	3	-	-
2001.....	-	-	-	2	1	-	3	-	5	-	1	-
2000.....	-	2	3	-	5	1	5	5	5	1	-	-
1999.....	2	1	-	3	4	4	4	5	3	2	-	-
Asian or Pacific Islander 2/, female												
2004.....	2	1	3	1	4	1	5	9	3	3	2	-
2003.....	1	1	1	2	4	3	4	5	4	4	2	-
2002.....	1	1	5	2	3	3	7	4	8	-	-	-
2001.....	1	3	2	2	2	3	3	8	5	1	1	-
2000.....	2	-	2	5	2	3	1	2	3	-	-	-
1999.....	1	2	1	3	2	1	2	9	4	6	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nutritional anemias (D50-D53)												
All races, both sexes												
2004.....	2	4	6	7	12	29	40	41	73	12	8	-
2003.....	3	6	8	8	24	27	40	60	60	35	10	-
2002.....	4	5	12	12	15	24	28	64	52	35	9	-
2001.....	2	5	5	11	17	36	43	62	51	34	11	-
2000.....	-	5	4	8	10	29	43	74	63	35	12	-
1999.....	2	3	5	5	20	28	56	52	64	37	12	-
All races, male												
2004.....	1	3	3	4	1	6	15	12	25	2	2	-
2003.....	2	2	6	4	13	18	14	19	15	7	-	-
2002.....	2	3	6	6	8	8	5	21	8	7	3	-
2001.....	2	3	1	9	8	14	20	23	10	6	2	-
2000.....	-	4	1	1	4	13	7	24	29	5	4	-
1999.....	1	1	4	1	9	16	19	23	16	7	1	-
All races, female												
2004.....	1	1	3	3	11	23	25	29	48	10	6	-
2003.....	1	4	2	4	11	9	26	41	45	28	10	-
2002.....	2	2	6	6	7	16	23	43	44	28	6	-
2001.....	-	2	4	2	9	22	23	39	41	28	9	-
2000.....	-	1	3	7	6	16	36	50	34	30	8	-
1999.....	1	2	1	4	11	12	37	29	48	30	11	-
White, both sexes												
2004.....	2	2	5	6	11	28	38	37	64	11	8	-
2003.....	2	3	4	6	15	25	34	54	57	33	9	-
2002.....	4	3	11	8	11	21	24	60	48	32	9	-
2001.....	2	2	2	9	15	30	38	52	51	32	11	-
2000.....	-	3	3	5	7	23	38	68	55	35	10	-
1999.....	-	2	3	4	17	22	49	48	55	33	10	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nutritional anemias (D50-D53)												
White, male												
2004.....	1	2	2	3	1	5	14	12	25	2	2	-
2003.....	2	1	2	4	7	16	14	18	14	6	-	-
2002.....	2	2	5	4	6	6	5	19	7	7	3	-
2001.....	2	2	-	7	6	11	17	21	10	6	2	-
2000.....	-	2	1	1	4	9	7	21	26	5	3	-
1999.....	-	1	2	1	7	12	16	22	15	6	1	-
White, female												
2004.....	1	-	3	3	10	23	24	25	39	9	6	-
2003.....	-	2	2	2	8	9	20	36	43	27	9	-
2002.....	2	1	6	4	5	15	19	41	41	25	6	-
2001.....	-	-	2	2	9	19	21	31	41	26	9	-
2000.....	-	1	2	4	3	14	31	47	29	30	7	-
1999.....	-	1	1	3	10	10	33	26	40	27	9	-
Black, both sexes												
2004.....	-	2	1	1	1	-	2	2	8	1	-	-
2003.....	1	3	4	2	8	2	6	5	1	-	-	-
2002.....	-	2	1	4	3	3	3	3	2	3	-	-
2001.....	-	3	3	1	2	6	4	8	-	1	-	-
2000.....	-	2	1	2	2	4	5	5	8	-	2	-
1999.....	1	1	1	-	3	6	7	4	8	3	2	-
Black, male												
2004.....	-	1	1	1	-	-	1	-	-	-	-	-
2003.....	-	1	4	-	6	2	-	1	-	-	-	-
2002.....	-	1	1	2	1	2	-	1	1	-	-	-
2001.....	-	1	1	1	2	3	3	1	-	-	-	-
2000.....	-	2	-	-	-	2	-	3	3	-	1	-
1999.....	1	-	1	-	2	4	3	1	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nutritional anemias (D50-D53)												
Asian or Pacific Islander 2/, both sexes												
2004.....	-	-	-	-	-	1	-	2	-	-	-	-
2003.....	-	-	-	-	1	-	-	1	2	2	1	-
2002.....	-	-	-	-	1	-	1	1	1	-	-	-
2001.....	-	-	-	1	-	-	1	1	-	-	-	-
2000.....	-	-	-	1	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	1	1	-	-
Asian or Pacific Islander 2/, male												
2004.....	-	-	-	-	-	1	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	1	1	-	-
2002.....	-	-	-	-	1	-	-	1	-	-	-	-
2001.....	-	-	-	1	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, female												
2004.....	-	-	-	-	-	-	-	2	-	-	-	-
2003.....	-	-	-	-	1	-	-	1	1	1	1	-
2002.....	-	-	-	-	-	-	1	-	1	-	-	-
2001.....	-	-	-	-	-	-	1	1	-	-	-	-
2000.....	-	-	-	1	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hemolytic anemias (D55-D59)												
All races, both sexes												
2004.....	62	44	37	27	26	40	44	56	24	11	-	-
2003.....	50	40	30	16	26	45	59	40	33	16	1	-
2002.....	62	30	21	34	37	49	52	60	28	10	3	-
2001.....	53	38	26	29	32	52	62	53	33	8	3	-
2000.....	64	35	25	29	36	48	60	63	33	5	-	-
1999.....	38	43	25	28	36	54	57	56	24	13	-	-
All races, male												
2004.....	26	26	16	7	9	26	22	23	7	2	-	-
2003.....	20	22	16	6	14	23	22	12	7	3	-	-
2002.....	26	12	10	12	17	15	14	17	9	5	-	-
2001.....	25	16	12	11	15	23	27	16	10	1	1	-
2000.....	30	15	9	11	16	19	22	19	12	-	-	-
1999.....	20	18	12	12	13	25	26	11	7	3	-	-
All races, female												
2004.....	36	18	21	20	17	14	22	33	17	9	-	-
2003.....	30	18	14	10	12	22	37	28	26	13	1	-
2002.....	36	18	11	22	20	34	38	43	19	5	3	-
2001.....	28	22	14	18	17	29	35	37	23	7	2	-
2000.....	34	20	16	18	20	29	38	44	21	5	-	-
1999.....	18	25	13	16	23	29	31	45	17	10	-	-
White, both sexes												
2004.....	9	6	17	17	15	34	43	51	23	10	-	-
2003.....	9	5	13	10	22	40	58	39	30	14	1	-
2002.....	15	5	10	25	26	41	51	56	27	9	2	-
2001.....	12	11	16	21	22	47	56	51	29	7	2	-
2000.....	17	13	7	17	26	45	57	62	30	5	-	-
1999.....	6	7	12	20	31	50	51	53	24	13	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hemolytic anemias (D55-D59)												
White, male												
2004.....	5	3	9	5	6	23	22	19	6	2	-	-
2003.....	3	3	7	4	11	20	22	12	5	2	-	-
2002.....	8	3	5	10	13	14	14	16	9	5	-	-
2001.....	5	5	7	8	11	22	26	16	9	1	-	-
2000.....	8	7	4	7	12	18	21	19	12	-	-	-
1999.....	4	3	5	7	10	24	24	9	7	3	-	-
White, female												
2004.....	4	3	8	12	9	11	21	32	17	8	-	-
2003.....	6	2	6	6	11	20	36	27	25	12	1	-
2002.....	7	2	5	15	13	27	37	40	18	4	2	-
2001.....	7	6	9	13	11	25	30	35	20	6	2	-
2000.....	9	6	3	10	14	27	36	43	18	5	-	-
1999.....	2	4	7	13	21	26	27	44	17	10	-	-
Black, both sexes												
2004.....	53	37	20	10	11	6	-	4	-	1	-	-
2003.....	41	35	17	6	4	5	1	1	3	2	-	-
2002.....	47	24	11	9	10	7	-	4	1	1	1	-
2001.....	41	27	10	8	10	4	4	2	2	-	1	-
2000.....	47	22	18	12	7	2	2	1	2	-	-	-
1999.....	32	35	12	8	4	4	6	3	-	-	-	-
Black, male												
2004.....	21	23	7	2	3	3	-	3	-	-	-	-
2003.....	17	19	9	2	3	3	-	-	2	1	-	-
2002.....	18	9	5	2	4	1	-	1	-	-	-	-
2001.....	20	11	5	3	4	1	1	-	1	-	1	-
2000.....	22	8	5	4	3	1	-	-	-	-	-	-
1999.....	16	15	7	5	2	1	2	2	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Aplastic anemias (D60-D61)												
All races, both sexes												
2004.....	21	44	62	61	101	114	143	138	94	35	7	-
2003.....	22	33	48	73	91	128	152	123	93	48	5	-
2002.....	26	37	44	60	94	134	168	132	73	37	-	-
2001.....	41	44	57	58	108	155	180	156	101	34	7	-
2000.....	29	40	54	78	100	136	177	175	90	39	10	-
1999.....	24	31	50	66	112	150	170	156	108	45	12	-
All races, male												
2004.....	11	19	32	31	55	53	68	56	29	8	1	-
2003.....	11	16	23	43	48	63	69	55	35	17	-	-
2002.....	13	14	25	18	50	70	74	57	31	9	-	-
2001.....	22	18	28	28	43	70	75	63	31	6	1	-
2000.....	13	22	25	43	48	56	80	55	24	10	4	-
1999.....	11	12	26	44	47	71	77	69	33	15	3	-
All races, female												
2004.....	10	25	30	30	46	61	75	82	65	27	6	-
2003.....	11	17	25	30	43	65	83	68	58	31	5	-
2002.....	13	23	19	42	44	64	94	75	42	28	-	-
2001.....	19	26	29	30	65	85	105	93	70	28	6	-
2000.....	16	18	29	35	52	80	97	120	66	29	6	-
1999.....	13	19	24	22	65	79	93	87	75	30	9	-
White, both sexes												
2004.....	17	37	52	54	85	100	133	122	86	32	6	-
2003.....	16	24	40	60	77	118	135	116	82	45	3	-
2002.....	15	27	31	53	78	128	152	119	64	34	-	-
2001.....	34	36	48	47	91	144	166	145	97	32	7	-
2000.....	22	35	43	70	90	121	164	156	82	35	10	-
1999.....	20	19	35	55	96	134	159	142	102	40	11	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Aplastic anemias (D60-D61)												
White, male												
2004.....	9	15	27	28	49	45	62	50	27	7	1	-
2003.....	9	12	19	35	40	62	65	53	32	14	-	-
2002.....	9	10	20	16	40	69	65	52	26	9	-	-
2001.....	18	13	23	25	32	66	71	62	31	6	1	-
2000.....	10	18	20	40	44	47	71	46	20	8	4	-
1999.....	8	7	17	39	38	65	72	64	31	13	3	-
White, female												
2004.....	8	22	25	26	36	55	71	72	59	25	5	-
2003.....	7	12	21	25	37	56	70	63	50	31	3	-
2002.....	6	17	11	37	38	59	87	67	38	25	-	-
2001.....	16	23	25	22	59	78	95	83	66	26	6	-
2000.....	12	17	23	30	46	74	93	110	62	27	6	-
1999.....	12	12	18	16	58	69	87	78	71	27	8	-
Black, both sexes												
2004.....	3	7	6	4	12	11	7	12	7	1	1	-
2003.....	4	8	4	11	10	9	16	7	10	3	1	-
2002.....	9	9	8	6	14	5	12	10	8	3	-	-
2001.....	6	5	7	8	13	9	12	6	2	2	-	-
2000.....	5	2	7	4	7	12	10	16	7	3	-	-
1999.....	2	10	13	8	13	11	8	11	5	4	1	-
Black, male												
2004.....	2	4	3	2	5	6	4	5	1	-	-	-
2003.....	2	4	1	7	5	1	4	2	2	3	-	-
2002.....	4	4	4	2	9	1	7	3	5	-	-	-
2001.....	4	4	5	2	9	4	2	1	-	-	-	-
2000.....	3	1	2	3	2	7	6	7	4	1	-	-
1999.....	2	4	9	3	8	2	3	4	1	2	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Aplastic anemias (D60-D61)												
Black, female												
2004.....	1	3	3	2	7	5	3	7	6	1	1	-
2003.....	2	4	3	4	5	8	12	5	8	-	1	-
2002.....	5	5	4	4	5	4	5	7	3	3	-	-
2001.....	2	1	2	6	4	5	10	5	2	2	-	-
2000.....	2	1	5	1	5	5	4	9	3	2	-	-
1999.....	-	6	4	5	5	9	5	7	4	2	1	-
American Indian 1/, both sexes												
2004.....	-	-	-	1	-	1	-	1	-	-	-	-
2003.....	1	-	1	-	1	-	-	-	-	-	-	-
2002.....	1	1	-	-	-	-	-	-	-	-	-	-
2001.....	-	1	-	-	1	1	-	4	-	-	-	-
2000.....	-	1	-	1	-	2	-	-	-	-	-	-
1999.....	-	-	2	-	1	-	-	-	-	-	-	-
American Indian 1/, male												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	1	-	1	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	1	-	-	1	-	-	-	-	-	-	-
2000.....	-	1	-	-	-	2	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, female												
2004.....	-	-	-	1	-	1	-	1	-	-	-	-
2003.....	1	-	-	-	-	-	-	-	-	-	-	-
2002.....	1	1	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	1	-	4	-	-	-	-
2000.....	-	-	-	1	-	-	-	-	-	-	-	-
1999.....	-	-	2	-	1	-	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Aplastic anemias (D60-D61)												
Asian or Pacific Islander 2/, both sexes												
2004.....	1	-	4	2	4	2	3	3	1	2	-	-
2003.....	1	1	3	2	3	1	1	-	1	-	1	-
2002.....	1	-	5	1	2	1	4	3	1	-	-	-
2001.....	1	2	2	3	3	1	2	1	2	-	-	-
2000.....	2	2	4	3	3	1	3	3	1	1	-	-
1999.....	2	2	-	3	2	5	3	3	1	1	-	-
Asian or Pacific Islander 2/, male												
2004.....	-	-	2	1	1	2	2	1	1	1	-	-
2003.....	-	-	2	1	2	-	-	-	1	-	-	-
2002.....	-	-	1	-	1	-	2	2	-	-	-	-
2001.....	-	-	-	1	1	-	2	-	-	-	-	-
2000.....	-	2	3	-	2	-	3	2	-	1	-	-
1999.....	1	1	-	2	1	4	2	1	1	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	1	-	2	1	3	-	1	2	-	1	-	-
2003.....	1	1	1	1	1	1	1	-	-	-	1	-
2002.....	1	-	4	1	1	1	2	1	1	-	-	-
2001.....	1	2	2	2	2	1	-	1	2	-	-	-
2000.....	2	-	1	3	1	1	-	1	1	-	-	-
1999.....	1	1	-	1	1	1	1	2	-	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute posthemorrhagic and other anemias (D62,D64)												
All races, both sexes												
2004.....	31	48	64	86	153	257	452	539	493	254	65	1
2003.....	29	42	69	87	159	247	438	561	535	243	56	-
2002.....	27	49	66	78	178	267	439	531	512	240	66	-
2001.....	24	32	42	88	149	282	416	537	474	218	69	-
2000.....	24	47	50	95	159	269	409	478	452	211	56	-
1999.....	34	28	46	76	143	265	384	511	451	221	58	-
All races, male												
2004.....	18	30	35	46	73	109	167	207	119	45	12	1
2003.....	15	27	44	51	72	99	169	184	139	48	8	-
2002.....	11	28	33	41	92	111	157	161	134	60	10	-
2001.....	9	16	17	43	80	113	154	174	126	39	11	-
2000.....	13	25	29	40	73	110	158	139	118	43	6	-
1999.....	16	12	20	35	65	105	145	158	116	48	8	-
All races, female												
2004.....	13	18	29	40	80	148	285	332	374	209	53	-
2003.....	14	15	25	36	87	148	269	377	396	195	48	-
2002.....	16	21	33	37	86	156	282	370	378	180	56	-
2001.....	15	16	25	45	69	169	262	363	348	179	58	-
2000.....	11	22	21	55	86	159	251	339	334	168	50	-
1999.....	18	16	26	41	78	160	239	353	335	173	50	-
White, both sexes												
2004.....	19	26	39	65	126	208	397	471	445	232	53	1
2003.....	17	24	40	59	125	196	379	509	469	214	52	-
2002.....	16	36	38	55	143	197	376	473	446	209	62	-
2001.....	13	20	33	69	119	225	353	480	422	195	49	-
2000.....	17	33	32	68	127	225	358	432	399	186	48	-
1999.....	15	18	26	56	110	222	345	449	402	191	47	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute posthemorrhagic and other anemias (D62,D64)												
White, male												
2004.....	11	15	22	34	60	86	145	184	110	41	11	1
2003.....	8	16	25	35	53	77	148	166	123	42	8	-
2002.....	7	21	17	31	69	84	132	144	115	47	9	-
2001.....	3	11	14	34	64	94	134	160	109	33	8	-
2000.....	9	18	17	33	58	95	139	126	98	39	6	-
1999.....	8	5	8	28	52	89	132	135	107	44	8	-
White, female												
2004.....	8	11	17	31	66	122	252	287	335	191	42	-
2003.....	9	8	15	24	72	119	231	343	346	172	44	-
2002.....	9	15	21	24	74	113	244	329	331	162	53	-
2001.....	10	9	19	35	55	131	219	320	313	162	41	-
2000.....	8	15	15	35	69	130	219	306	301	147	42	-
1999.....	7	13	18	28	58	133	213	314	295	147	39	-
Black, both sexes												
2004.....	11	19	24	21	23	46	45	61	44	19	10	-
2003.....	12	18	27	22	32	48	55	46	61	25	4	-
2002.....	9	12	27	22	32	65	59	53	57	27	3	-
2001.....	11	11	9	19	30	53	59	50	46	21	17	-
2000.....	7	14	17	25	30	40	47	42	44	25	8	-
1999.....	17	9	20	17	29	43	36	49	44	23	9	-
Black, male												
2004.....	7	12	13	12	11	21	15	22	8	3	1	-
2003.....	7	11	18	14	19	22	21	16	15	5	-	-
2002.....	3	6	16	10	21	24	24	16	16	9	1	-
2001.....	6	5	3	9	16	18	19	13	12	6	2	-
2000.....	4	7	12	6	13	13	17	10	15	4	-	-
1999.....	7	6	12	6	11	16	11	18	7	2	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute posthemorrhagic and other anemias (D62,D64)												
Black, female												
2004.....	4	7	11	9	12	25	30	39	36	16	9	-
2003.....	5	7	9	8	13	26	34	30	46	20	4	-
2002.....	6	6	11	12	11	41	35	37	41	18	2	-
2001.....	5	6	6	10	14	35	40	37	34	15	15	-
2000.....	3	7	5	19	17	27	30	32	29	21	8	-
1999.....	10	3	8	11	18	27	25	31	37	21	9	-
American Indian 1/, both sexes												
2004.....	-	1	-	-	2	-	2	1	1	-	-	-
2003.....	-	-	1	4	-	1	1	-	1	-	-	-
2002.....	1	1	-	-	-	-	1	1	-	1	1	-
2001.....	-	-	-	-	-	3	2	1	-	2	1	-
2000.....	-	-	-	1	-	2	2	-	3	-	-	-
1999.....	1	1	-	-	1	-	-	2	-	1	2	-
American Indian 1/, male												
2004.....	-	1	-	-	1	-	2	-	1	-	-	-
2003.....	-	-	-	1	-	-	-	-	-	-	-	-
2002.....	-	1	-	-	-	-	1	-	-	1	-	-
2001.....	-	-	-	-	-	1	-	1	-	-	-	-
2000.....	-	-	-	1	-	1	1	-	-	-	-	-
1999.....	-	1	-	-	-	-	-	1	-	1	-	-
American Indian 1/, female												
2004.....	-	-	-	-	1	-	-	1	-	-	-	-
2003.....	-	-	1	3	-	1	1	-	1	-	-	-
2002.....	1	-	-	-	-	-	-	1	-	-	1	-
2001.....	-	-	-	-	-	2	2	-	-	2	1	-
2000.....	-	-	-	-	-	1	1	-	3	-	-	-
1999.....	1	-	-	-	1	-	-	1	-	-	2	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute posthemorrhagic and other anemias (D62,D64)												
Asian or Pacific Islander 2/, both sexes												
2004.....	1	2	1	-	2	3	8	6	3	3	2	-
2003.....	-	-	1	2	2	2	3	6	4	4	-	-
2002.....	1	-	1	1	3	5	3	4	9	3	-	-
2001.....	-	1	-	-	-	1	2	6	6	-	2	-
2000.....	-	-	1	1	2	2	2	4	6	-	-	-
1999.....	1	-	-	3	3	-	3	11	5	6	-	-
Asian or Pacific Islander 2/, male												
2004.....	-	2	-	-	1	2	5	1	-	1	-	-
2003.....	-	-	1	1	-	-	-	2	1	1	-	-
2002.....	1	-	-	-	2	3	-	1	3	3	-	-
2001.....	-	-	-	-	-	-	1	-	5	-	1	-
2000.....	-	-	-	-	2	1	1	3	5	-	-	-
1999.....	1	-	-	1	2	-	2	4	2	1	-	-
Asian or Pacific Islander 2/, female												
2004.....	1	-	1	-	1	1	3	5	3	2	2	-
2003.....	-	-	-	1	2	2	3	4	3	3	-	-
2002.....	-	-	1	1	1	2	3	3	6	-	-	-
2001.....	-	1	-	-	-	1	1	6	1	-	1	-
2000.....	-	-	1	1	-	1	1	1	1	-	-	-
1999.....	-	-	-	2	1	-	1	7	3	5	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Coagulation defects, purpura and other hemorrhagic conditions (D65-D69)												
All races, both sexes												
2004.....	154	163	184	200	253	364	415	289	148	47	6	-
2003.....	167	163	156	221	285	374	379	292	154	46	4	-
2002.....	136	135	162	212	306	358	407	297	159	53	8	-
2001.....	127	151	165	215	276	366	392	280	144	53	10	-
2000.....	143	154	176	246	293	355	393	285	142	56	8	-
1999.....	119	129	165	208	327	394	336	264	151	54	7	-
All races, male												
2004.....	74	95	96	100	132	163	180	101	46	17	1	-
2003.....	80	81	75	105	145	170	163	111	48	13	-	-
2002.....	68	63	86	93	134	171	168	122	54	15	2	-
2001.....	63	68	83	101	128	147	186	111	39	10	-	-
2000.....	69	64	79	118	148	149	153	120	44	12	3	-
1999.....	63	64	65	95	161	174	123	106	50	12	3	-
All races, female												
2004.....	80	68	88	100	121	201	235	188	102	30	5	-
2003.....	87	82	81	116	140	204	216	181	106	33	4	-
2002.....	68	72	76	119	172	187	239	175	105	38	6	-
2001.....	64	83	82	114	148	219	206	169	105	43	10	-
2000.....	74	90	97	128	145	206	240	165	98	44	5	-
1999.....	56	65	100	113	166	220	213	158	101	42	4	-
White, both sexes												
2004.....	107	118	152	161	213	325	376	260	133	40	4	-
2003.....	115	121	121	177	247	316	338	258	134	44	3	-
2002.....	93	102	130	168	265	303	372	274	144	45	7	-
2001.....	88	124	126	182	239	320	346	249	137	45	9	-
2000.....	95	115	141	204	246	310	352	262	132	50	7	-
1999.....	83	88	137	159	274	341	299	238	138	49	6	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Coagulation defects, purpura and other hemorrhagic conditions (D65-D69)												
White, male												
2004.....	53	73	80	80	115	148	163	92	43	14	1	-
2003.....	52	60	58	81	128	147	148	96	45	13	-	-
2002.....	44	49	71	75	121	149	149	114	50	11	2	-
2001.....	44	56	65	85	109	127	168	100	37	10	-	-
2000.....	46	51	64	97	125	135	139	111	40	11	2	-
1999.....	47	45	52	72	137	148	104	97	48	11	2	-
White, female												
2004.....	54	45	72	81	98	177	213	168	90	26	3	-
2003.....	63	61	63	96	119	169	190	162	89	31	3	-
2002.....	49	53	59	93	144	154	223	160	94	34	5	-
2001.....	44	68	61	97	130	193	178	149	100	35	9	-
2000.....	49	64	77	107	121	175	213	151	92	39	5	-
1999.....	36	43	85	87	137	193	195	141	90	38	4	-
Black, both sexes												
2004.....	45	38	29	33	36	36	30	26	12	6	2	-
2003.....	45	35	27	37	36	44	31	27	18	2	-	-
2002.....	38	28	26	42	36	46	28	22	15	6	1	-
2001.....	36	22	32	28	31	37	38	26	6	7	1	-
2000.....	46	33	31	33	40	39	36	21	8	6	1	-
1999.....	32	37	25	42	40	46	30	25	12	5	1	-
Black, male												
2004.....	20	19	15	18	16	14	12	8	3	3	-	-
2003.....	23	18	13	18	17	18	10	11	2	-	-	-
2002.....	22	12	11	17	12	17	13	8	4	2	-	-
2001.....	17	10	13	13	14	16	14	6	1	-	-	-
2000.....	22	11	14	17	19	12	10	8	2	1	1	-
1999.....	14	16	10	21	16	21	15	9	2	1	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Coagulation defects, purpura and other hemorrhagic conditions (D65-D69)												
Black, female												
2004.....	25	19	14	15	20	22	18	18	9	3	2	-
2003.....	22	17	14	19	19	26	21	16	16	2	-	-
2002.....	16	16	15	25	24	29	15	14	11	4	1	-
2001.....	19	12	19	15	17	21	24	20	5	7	1	-
2000.....	24	22	17	16	21	27	26	13	6	5	-	-
1999.....	18	21	15	21	24	25	15	16	10	4	-	-
American Indian 1/, both sexes												
2004.....	-	1	1	3	-	-	-	-	1	-	-	-
2003.....	3	1	3	2	-	-	1	1	-	-	-	-
2002.....	2	2	1	1	1	1	-	-	-	-	-	-
2001.....	-	1	2	2	1	3	1	1	-	-	-	-
2000.....	1	4	-	3	-	1	-	-	-	-	-	-
1999.....	1	-	2	2	4	1	3	-	-	-	-	-
American Indian 1/, male												
2004.....	-	1	1	1	-	-	-	-	-	-	-	-
2003.....	3	-	1	1	-	-	-	-	-	-	-	-
2002.....	2	1	1	-	-	-	-	-	-	-	-	-
2001.....	-	-	1	2	1	1	-	1	-	-	-	-
2000.....	1	1	-	-	-	1	-	-	-	-	-	-
1999.....	1	-	2	-	1	1	1	-	-	-	-	-
American Indian 1/, female												
2004.....	-	-	-	2	-	-	-	-	1	-	-	-
2003.....	-	1	2	1	-	-	1	1	-	-	-	-
2002.....	-	1	-	1	1	1	-	-	-	-	-	-
2001.....	-	1	1	-	-	2	1	-	-	-	-	-
2000.....	-	3	-	3	-	-	-	-	-	-	-	-
1999.....	-	-	-	2	3	-	2	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Coagulation defects, purpura and other hemorrhagic conditions (D65-D69)												
Asian or Pacific Islander 2/, both sexes												
2004.....	2	6	2	3	4	3	9	3	2	1	-	-
2003.....	4	6	5	5	2	14	9	6	2	-	1	-
2002.....	3	3	5	1	4	8	7	1	-	2	-	-
2001.....	3	4	5	3	5	6	7	4	1	1	-	-
2000.....	1	2	4	6	7	5	5	2	2	-	-	-
1999.....	3	4	1	5	9	6	4	1	1	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	1	2	-	1	1	1	5	1	-	-	-	-
2003.....	2	3	3	5	-	5	5	4	1	-	-	-
2002.....	-	1	3	1	1	5	6	-	-	2	-	-
2001.....	2	2	4	1	4	3	4	4	1	-	-	-
2000.....	-	1	1	4	4	1	4	1	2	-	-	-
1999.....	1	3	1	2	7	4	3	-	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	1	4	2	2	3	2	4	2	2	1	-	-
2003.....	2	3	2	-	2	9	4	2	1	-	1	-
2002.....	3	2	2	-	3	3	1	1	-	-	-	-
2001.....	1	2	1	2	1	3	3	-	-	1	-	-
2000.....	1	1	3	2	3	4	1	1	-	-	-	-
1999.....	2	1	-	3	2	2	1	1	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other diseases of blood and blood-forming organs (D70-D76)

All races, both sexes

2004.....	43	43	52	63	92	94	108	77	45	21	3	-
2003.....	48	41	65	57	85	104	111	102	57	23	3	-
2002.....	50	53	63	65	94	118	137	96	43	14	3	-
2001.....	46	43	58	72	78	133	102	105	46	16	3	-
2000.....	45	36	57	77	89	116	109	88	39	16	1	-
1999.....	40	55	44	52	105	95	105	95	40	14	1	-

All races, male

2004.....	26	24	29	34	44	45	48	29	14	6	-	-
2003.....	28	18	31	25	45	52	50	38	15	5	-	-
2002.....	30	31	31	35	53	56	55	35	8	5	1	-
2001.....	26	22	31	35	32	68	42	33	12	5	1	-
2000.....	19	17	24	43	48	55	50	35	11	3	1	-
1999.....	14	28	24	22	59	53	49	31	12	5	-	-

All races, female

2004.....	17	19	23	29	48	49	60	48	31	15	3	-
2003.....	20	23	34	32	40	52	61	64	42	18	3	-
2002.....	20	22	32	30	41	62	82	61	35	9	2	-
2001.....	20	21	27	37	46	65	60	72	34	11	2	-
2000.....	26	19	33	34	41	61	59	53	28	13	-	-
1999.....	26	27	20	30	46	42	56	64	28	9	1	-

White, both sexes

2004.....	37	35	43	54	80	91	99	74	42	19	3	-
2003.....	39	32	58	49	80	93	103	97	52	23	2	-
2002.....	42	45	55	59	82	107	125	86	39	14	1	-
2001.....	35	36	49	56	71	125	92	99	41	13	3	-
2000.....	36	29	48	65	78	107	100	81	36	15	1	-
1999.....	31	42	36	45	94	91	97	89	36	13	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of blood and blood-forming organs (D70-D76)												
White, male												
2004.....	26	18	26	29	39	44	43	27	14	5	-	-
2003.....	24	12	29	23	41	45	47	38	14	5	-	-
2002.....	24	25	27	32	45	49	53	30	8	5	1	-
2001.....	19	18	27	27	30	65	37	30	12	5	1	-
2000.....	16	12	19	38	41	48	46	34	9	2	1	-
1999.....	10	23	22	18	55	51	44	28	10	5	-	-
White, female												
2004.....	11	17	17	25	41	47	56	47	28	14	3	-
2003.....	15	20	29	26	39	48	56	59	38	18	2	-
2002.....	18	20	28	27	37	58	72	56	31	9	-	-
2001.....	16	18	22	29	41	60	55	69	29	8	2	-
2000.....	20	17	29	27	37	59	54	47	27	13	-	-
1999.....	21	19	14	27	39	40	53	61	26	8	1	-
Black, both sexes												
2004.....	4	8	9	9	5	2	4	3	3	2	-	-
2003.....	6	5	6	7	4	8	7	5	5	-	1	-
2002.....	6	7	8	4	8	11	8	9	4	-	2	-
2001.....	9	5	7	13	5	6	8	4	4	2	-	-
2000.....	7	5	9	10	9	6	9	4	2	1	-	-
1999.....	7	13	8	7	9	3	7	4	3	1	-	-
Black, male												
2004.....	-	6	3	5	2	1	2	2	-	1	-	-
2003.....	3	4	2	1	3	5	2	-	1	-	-	-
2002.....	4	6	4	3	6	7	1	4	-	-	-	-
2001.....	7	2	4	7	1	2	4	1	-	-	-	-
2000.....	3	3	5	4	5	4	4	1	1	1	-	-
1999.....	3	5	2	4	3	1	4	1	2	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of blood and blood-forming organs (D70-D76)												
Black, female												
2004.....	4	2	6	4	3	1	2	1	3	1	-	-
2003.....	3	1	4	6	1	3	5	5	4	-	1	-
2002.....	2	1	4	1	2	4	7	5	4	-	2	-
2001.....	2	3	3	6	4	4	4	3	4	2	-	-
2000.....	4	2	4	6	4	2	5	3	1	-	-	-
1999.....	4	8	6	3	6	2	3	3	1	1	-	-
American Indian 1/, both sexes												
2004.....	1	-	-	-	1	-	-	-	-	-	-	-
2003.....	2	2	-	-	-	-	-	-	-	-	-	-
2002.....	1	-	-	-	1	-	-	-	-	-	-	-
2001.....	1	-	-	1	-	-	-	-	-	1	-	-
2000.....	-	-	-	-	1	2	-	1	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, male												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	1	-	-	-	-	-	-	-	-	-	-	-
2002.....	1	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	1	2	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, female												
2004.....	1	-	-	-	1	-	-	-	-	-	-	-
2003.....	1	2	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	1	-	-	-	-	-	-	-
2001.....	1	-	-	1	-	-	-	-	-	1	-	-
2000.....	-	-	-	-	-	-	-	1	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of blood and blood-forming organs (D70-D76)												
Asian or Pacific Islander 2/, both sexes												
2004.....	1	-	-	-	6	1	5	-	-	-	-	-
2003.....	1	2	1	1	1	3	1	-	-	-	-	-
2002.....	1	1	-	2	3	-	4	1	-	-	-	-
2001.....	1	2	2	2	2	2	2	2	1	-	-	-
2000.....	2	2	-	2	1	1	-	2	1	-	-	-
1999.....	2	-	-	-	2	1	1	2	-	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	-	-	-	-	3	-	3	-	-	-	-	-
2003.....	-	2	-	1	1	2	1	-	-	-	-	-
2002.....	1	-	-	-	2	-	1	1	-	-	-	-
2001.....	-	2	-	1	1	1	1	2	-	-	-	-
2000.....	-	2	-	1	1	1	-	-	1	-	-	-
1999.....	1	-	-	-	1	1	1	2	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	1	-	-	-	3	1	2	-	-	-	-	-
2003.....	1	-	1	-	-	1	-	-	-	-	-	-
2002.....	-	1	-	2	1	-	3	-	-	-	-	-
2001.....	1	-	2	1	1	1	1	-	1	-	-	-
2000.....	2	-	-	1	-	-	-	2	-	-	-	-
1999.....	1	-	-	-	1	-	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Certain disorders involving the immune mechanism (D80-D89)

All races, both sexes

2004.....	160	147	128	125	95	83	64	57	19	5	4	-
2003.....	169	152	123	105	114	90	69	46	16	8	4	-
2002.....	179	147	131	110	91	82	79	52	23	6	2	-
2001.....	154	119	120	104	106	90	67	41	21	4	2	-
2000.....	135	107	101	78	94	95	89	45	12	5	4	-
1999.....	103	87	80	82	79	76	50	36	17	5	2	-

All races, male

2004.....	76	66	50	56	25	20	22	14	6	2	-	-
2003.....	77	52	48	40	48	34	25	11	4	1	1	-
2002.....	84	71	58	39	38	24	31	13	7	1	-	-
2001.....	69	43	47	32	30	33	23	15	8	1	-	-
2000.....	63	44	48	32	44	31	31	13	5	2	-	-
1999.....	48	36	34	30	30	30	18	12	4	1	-	-

All races, female

2004.....	84	81	78	69	70	63	42	43	13	3	4	-
2003.....	92	100	75	65	66	56	44	35	12	7	3	-
2002.....	95	76	73	71	53	58	48	39	16	5	2	-
2001.....	85	76	73	72	76	57	44	26	13	3	2	-
2000.....	72	63	53	46	50	64	58	32	7	3	4	-
1999.....	55	51	46	52	49	46	32	24	13	4	2	-

White, both sexes

2004.....	57	65	62	77	63	65	48	53	16	4	4	-
2003.....	43	64	64	57	82	75	57	38	14	7	4	-
2002.....	62	72	70	70	68	70	67	47	21	5	1	-
2001.....	56	48	73	65	71	72	58	40	19	4	2	-
2000.....	50	45	54	45	78	80	71	41	12	3	3	-
1999.....	40	43	49	54	59	64	42	33	15	5	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Certain disorders involving the immune mechanism (D80-D89)												
White, male												
2004.....	33	31	29	36	15	19	18	13	5	2	-	-
2003.....	20	21	28	31	40	26	21	9	3	1	1	-
2002.....	28	39	37	25	33	22	29	11	7	1	-	-
2001.....	33	19	30	23	22	24	20	15	7	1	-	-
2000.....	26	27	29	23	38	28	25	13	5	1	-	-
1999.....	24	22	25	24	22	26	16	12	4	1	-	-
White, female												
2004.....	24	34	33	41	48	46	30	40	11	2	4	-
2003.....	23	43	36	26	42	49	36	29	11	6	3	-
2002.....	34	33	33	45	35	48	38	36	14	4	1	-
2001.....	23	29	43	42	49	48	38	25	12	3	2	-
2000.....	24	18	25	22	40	52	46	28	7	2	3	-
1999.....	16	21	24	30	37	38	26	21	11	4	1	-
Black, both sexes												
2004.....	101	80	65	48	29	16	15	4	3	1	-	-
2003.....	125	85	57	47	31	13	11	8	2	1	-	-
2002.....	116	74	58	39	23	12	12	4	2	1	-	-
2001.....	97	68	45	37	34	17	7	1	2	-	-	-
2000.....	85	61	46	32	15	15	18	4	-	2	1	-
1999.....	63	43	29	28	17	12	7	3	2	-	1	-
Black, male												
2004.....	42	35	21	20	10	1	3	1	1	-	-	-
2003.....	56	29	18	8	8	7	4	2	1	-	-	-
2002.....	56	32	20	13	5	2	2	2	-	-	-	-
2001.....	36	22	16	9	8	9	2	-	1	-	-	-
2000.....	37	17	19	8	5	3	6	-	-	1	-	-
1999.....	24	14	8	6	7	4	2	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Certain disorders involving the immune mechanism (D80-D89)

Asian or Pacific Islander 2/, both sexes

2004.....	-	2	1	-	3	1	1	-	-	-	-	-
2003.....	-	2	1	1	1	2	1	-	-	-	-	-
2002.....	-	-	2	1	-	-	-	1	-	-	1	-
2001.....	-	2	2	2	-	-	1	-	-	-	-	-
2000.....	-	-	-	1	1	-	-	-	-	-	-	-
1999.....	-	1	2	-	3	-	1	-	-	-	-	-

Asian or Pacific Islander 2/, male

2004.....	-	-	-	-	-	-	1	-	-	-	-	-
2003.....	-	1	1	1	-	1	-	-	-	-	-	-
2002.....	-	-	1	1	-	-	-	-	-	-	-	-
2001.....	-	1	1	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	1	1	-	-	-	-	-	-	-
1999.....	-	-	1	-	1	-	-	-	-	-	-	-

Asian or Pacific Islander 2/, female

2004.....	-	2	1	-	3	1	-	-	-	-	-	-
2003.....	-	1	-	-	1	1	1	-	-	-	-	-
2002.....	-	-	1	-	-	-	-	1	-	-	1	-
2001.....	-	1	1	2	-	-	1	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	1	1	-	2	-	1	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
IV. Endocrine, nutritional and metabolic diseases (E00-E88)												
All races, both sexes												
2004.....	4,580	6,254	7,435	8,746	11,080	13,919	15,495	12,340	7,449	2,571	510	5
2003.....	4,635	6,152	7,345	8,939	11,418	14,456	15,411	12,593	7,356	2,729	551	6
2002.....	4,536	5,861	6,877	8,743	11,558	14,625	15,345	12,652	7,331	2,672	530	4
2001.....	4,386	5,286	6,575	8,632	11,650	14,467	14,770	12,352	7,248	2,728	537	1
2000.....	4,021	5,019	6,369	8,524	11,680	14,515	14,168	11,870	7,070	2,586	563	1
1999.....	3,669	4,771	6,290	8,613	11,744	14,069	13,957	11,809	7,079	2,592	526	3
All races, male												
2004.....	2,729	3,563	4,209	4,757	5,637	6,613	6,690	4,516	2,135	521	81	2
2003.....	2,677	3,489	4,094	4,836	5,790	6,828	6,554	4,481	2,002	523	85	5
2002.....	2,638	3,372	3,806	4,533	5,704	6,680	6,307	4,484	1,934	520	73	4
2001.....	2,512	2,948	3,567	4,532	5,764	6,578	5,984	4,153	1,875	504	64	1
2000.....	2,276	2,766	3,419	4,333	5,698	6,548	5,722	3,934	1,841	497	77	1
1999.....	2,083	2,648	3,319	4,345	5,685	6,427	5,633	3,916	1,764	471	66	2
All races, female												
2004.....	1,851	2,691	3,226	3,989	5,443	7,306	8,805	7,824	5,314	2,050	429	3
2003.....	1,958	2,663	3,251	4,103	5,628	7,628	8,857	8,112	5,354	2,206	466	1
2002.....	1,898	2,489	3,071	4,210	5,854	7,945	9,038	8,168	5,397	2,152	457	-
2001.....	1,874	2,338	3,008	4,100	5,886	7,889	8,786	8,199	5,373	2,224	473	-
2000.....	1,745	2,253	2,950	4,191	5,982	7,967	8,446	7,936	5,229	2,089	486	-
1999.....	1,586	2,123	2,971	4,268	6,059	7,642	8,324	7,893	5,315	2,121	460	1
White, both sexes												
2004.....	3,357	4,736	5,704	6,788	8,792	11,479	13,220	10,715	6,497	2,242	411	3
2003.....	3,401	4,663	5,631	6,945	9,144	11,992	13,120	10,985	6,441	2,375	436	5
2002.....	3,330	4,426	5,225	6,673	9,260	12,154	13,148	11,002	6,467	2,314	443	3
2001.....	3,256	3,981	5,005	6,694	9,400	12,029	12,664	10,821	6,366	2,394	441	1
2000.....	2,964	3,798	4,895	6,625	9,443	12,098	12,116	10,387	6,226	2,259	449	1
1999.....	2,687	3,488	4,727	6,704	9,415	11,761	11,965	10,293	6,262	2,269	434	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
IV. Endocrine, nutritional and metabolic diseases (E00-E88)												
White, male												
2004.....	2,095	2,769	3,344	3,796	4,627	5,619	5,874	4,004	1,881	454	66	-
2003.....	2,041	2,743	3,237	3,884	4,791	5,843	5,725	4,013	1,774	466	73	4
2002.....	2,016	2,636	2,971	3,595	4,756	5,758	5,537	4,010	1,751	450	62	3
2001.....	1,921	2,296	2,798	3,660	4,847	5,622	5,220	3,747	1,650	438	51	1
2000.....	1,753	2,154	2,703	3,512	4,780	5,611	5,052	3,519	1,633	443	65	1
1999.....	1,596	2,015	2,597	3,495	4,763	5,532	4,961	3,461	1,536	399	51	1
White, female												
2004.....	1,262	1,967	2,360	2,992	4,165	5,860	7,346	6,711	4,616	1,788	345	3
2003.....	1,360	1,920	2,394	3,061	4,353	6,149	7,395	6,972	4,667	1,909	363	1
2002.....	1,314	1,790	2,254	3,078	4,504	6,396	7,611	6,992	4,716	1,864	381	-
2001.....	1,335	1,685	2,207	3,034	4,553	6,407	7,444	7,074	4,716	1,956	390	-
2000.....	1,211	1,644	2,192	3,113	4,663	6,487	7,064	6,868	4,593	1,816	384	-
1999.....	1,091	1,473	2,130	3,209	4,652	6,229	7,004	6,832	4,726	1,870	383	-
Black, both sexes												
2004.....	1,084	1,310	1,476	1,687	1,965	2,075	1,880	1,371	810	280	89	2
2003.....	1,082	1,312	1,466	1,709	1,931	2,081	1,907	1,347	776	298	105	1
2002.....	1,074	1,241	1,443	1,751	1,950	2,074	1,880	1,384	761	323	81	1
2001.....	1,008	1,151	1,355	1,671	1,954	2,094	1,783	1,321	784	290	90	-
2000.....	933	1,073	1,288	1,659	1,938	2,111	1,782	1,306	749	289	105	-
1999.....	882	1,111	1,369	1,664	1,982	1,960	1,721	1,312	723	276	84	2
Black, male												
2004.....	557	670	720	824	856	814	664	412	207	57	12	2
2003.....	545	651	718	803	828	815	674	372	170	43	9	1
2002.....	546	623	724	770	791	768	645	358	147	54	10	1
2001.....	509	568	655	746	789	808	637	330	187	50	13	-
2000.....	458	530	612	692	793	797	570	350	166	43	9	-
1999.....	430	548	615	744	772	732	557	379	191	55	12	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
IV. Endocrine, nutritional and metabolic diseases (E00-E88)												
Black, female												
2004.....	527	640	756	863	1,109	1,261	1,216	959	603	223	77	-
2003.....	537	661	748	906	1,103	1,266	1,233	975	606	255	96	-
2002.....	528	618	719	981	1,159	1,306	1,235	1,026	614	269	71	-
2001.....	499	583	700	925	1,165	1,286	1,146	991	597	240	77	-
2000.....	475	543	676	967	1,145	1,314	1,212	956	583	246	96	-
1999.....	452	563	754	920	1,210	1,228	1,164	933	532	221	72	1
American Indian 1/, both sexes												
2004.....	58	93	112	108	104	112	94	44	20	5	-	-
2003.....	77	80	98	114	108	102	94	60	24	17	3	-
2002.....	68	90	94	123	127	108	77	46	25	8	1	-
2001.....	63	69	89	87	106	99	83	52	24	6	1	-
2000.....	57	69	85	82	122	97	57	49	19	5	5	-
1999.....	57	81	95	113	120	101	68	61	21	8	2	-
American Indian 1/, male												
2004.....	32	51	63	48	42	63	30	13	4	-	-	-
2003.....	45	38	43	49	53	43	37	14	11	2	1	-
2002.....	34	46	48	52	58	40	24	16	7	3	-	-
2001.....	39	32	43	38	41	45	26	16	4	2	-	-
2000.....	24	31	48	37	42	43	16	17	7	1	2	-
1999.....	35	37	47	43	44	47	28	19	5	1	-	-
American Indian 1/, female												
2004.....	26	42	49	60	62	49	64	31	16	5	-	-
2003.....	32	42	55	65	55	59	57	46	13	15	2	-
2002.....	34	44	46	71	69	68	53	30	18	5	1	-
2001.....	24	37	46	49	65	54	57	36	20	4	1	-
2000.....	33	38	37	45	80	54	41	32	12	4	3	-
1999.....	22	44	48	70	76	54	40	42	16	7	2	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
IV. Endocrine, nutritional and metabolic diseases (E00-E88)												
Asian or Pacific Islander 2/, both sexes												
2004.....	81	115	143	163	219	253	301	210	122	44	10	-
2003.....	75	97	150	171	235	281	290	201	115	39	7	-
2002.....	64	104	115	196	221	289	240	220	78	27	5	-
2001.....	59	85	126	180	190	245	240	158	74	38	5	-
2000.....	67	79	101	158	177	209	213	128	76	33	4	-
1999.....	43	91	99	132	227	247	203	143	73	39	6	-
Asian or Pacific Islander 2/, male												
2004.....	45	73	82	89	112	117	122	87	43	10	3	-
2003.....	46	57	96	100	118	127	118	82	47	12	2	-
2002.....	42	67	63	116	99	114	101	100	29	13	1	-
2001.....	43	52	71	88	87	103	101	60	34	14	-	-
2000.....	41	51	56	92	83	97	84	48	35	10	1	-
1999.....	22	48	60	63	106	116	87	57	32	16	3	-
Asian or Pacific Islander 2/, female												
2004.....	36	42	61	74	107	136	179	123	79	34	7	-
2003.....	29	40	54	71	117	154	172	119	68	27	5	-
2002.....	22	37	52	80	122	175	139	120	49	14	4	-
2001.....	16	33	55	92	103	142	139	98	40	24	5	-
2000.....	26	28	45	66	94	112	129	80	41	23	3	-
1999.....	21	43	39	69	121	131	116	86	41	23	3	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Endocrine diseases (E00-E34)												
All races, both sexes												
2004.....	3,433	4,931	6,064	7,228	9,160	11,369	12,280	9,162	5,051	1,558	284	3
2003.....	3,487	4,921	6,041	7,432	9,537	11,905	12,195	9,271	4,981	1,630	272	4
2002.....	3,478	4,600	5,623	7,367	9,673	12,012	12,036	9,347	4,845	1,534	238	1
2001.....	3,374	4,267	5,486	7,301	9,732	11,907	11,693	8,938	4,716	1,542	230	1
2000.....	3,109	4,086	5,282	7,166	9,809	11,935	10,999	8,460	4,485	1,433	256	-
1999.....	2,878	3,947	5,317	7,344	9,909	11,577	10,899	8,354	4,363	1,331	226	3
All races, male												
2004.....	2,048	2,796	3,451	3,896	4,661	5,437	5,359	3,402	1,488	321	53	1
2003.....	2,018	2,763	3,365	3,976	4,824	5,665	5,242	3,365	1,382	328	46	3
2002.....	2,013	2,654	3,113	3,790	4,744	5,495	5,001	3,369	1,340	334	32	1
2001.....	1,961	2,393	2,923	3,779	4,781	5,370	4,779	3,088	1,275	304	38	1
2000.....	1,727	2,212	2,799	3,597	4,739	5,389	4,540	2,887	1,237	280	24	-
1999.....	1,641	2,153	2,774	3,677	4,723	5,297	4,480	2,911	1,136	237	33	2
All races, female												
2004.....	1,385	2,135	2,613	3,332	4,499	5,932	6,921	5,760	3,563	1,237	231	2
2003.....	1,469	2,158	2,676	3,456	4,713	6,240	6,953	5,906	3,599	1,302	226	1
2002.....	1,465	1,946	2,510	3,577	4,929	6,517	7,035	5,978	3,505	1,200	206	-
2001.....	1,413	1,874	2,563	3,522	4,951	6,537	6,914	5,850	3,441	1,238	192	-
2000.....	1,382	1,874	2,483	3,569	5,070	6,546	6,459	5,573	3,248	1,153	232	-
1999.....	1,237	1,794	2,543	3,667	5,186	6,280	6,419	5,443	3,227	1,094	193	1
White, both sexes												
2004.....	2,432	3,633	4,559	5,529	7,150	9,252	10,347	7,830	4,350	1,328	215	2
2003.....	2,479	3,641	4,527	5,669	7,522	9,779	10,288	7,969	4,308	1,398	220	4
2002.....	2,476	3,394	4,196	5,539	7,673	9,887	10,240	8,048	4,221	1,308	189	-
2001.....	2,440	3,130	4,107	5,585	7,746	9,801	9,962	7,746	4,117	1,321	174	1
2000.....	2,227	3,043	3,990	5,473	7,837	9,836	9,293	7,326	3,877	1,237	196	-
1999.....	2,052	2,808	3,920	5,649	7,861	9,582	9,261	7,175	3,769	1,149	179	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Endocrine diseases (E00-E34)												
White, male												
2004.....	1,525	2,112	2,687	3,063	3,774	4,584	4,672	2,990	1,293	276	43	-
2003.....	1,496	2,118	2,601	3,145	3,953	4,832	4,573	2,984	1,213	295	42	3
2002.....	1,484	2,033	2,400	2,967	3,926	4,714	4,388	3,009	1,219	291	27	-
2001.....	1,461	1,825	2,268	3,021	3,992	4,554	4,162	2,778	1,128	259	31	1
2000.....	1,291	1,691	2,188	2,878	3,944	4,594	3,992	2,561	1,085	247	20	-
1999.....	1,221	1,601	2,131	2,934	3,940	4,547	3,949	2,561	981	203	26	1
White, female												
2004.....	907	1,521	1,872	2,466	3,376	4,668	5,675	4,840	3,057	1,052	172	2
2003.....	983	1,523	1,926	2,524	3,569	4,947	5,715	4,985	3,095	1,103	178	1
2002.....	992	1,361	1,796	2,572	3,747	5,173	5,852	5,039	3,002	1,017	162	-
2001.....	979	1,305	1,839	2,564	3,754	5,247	5,800	4,968	2,989	1,062	143	-
2000.....	936	1,352	1,802	2,595	3,893	5,242	5,301	4,765	2,792	990	176	-
1999.....	831	1,207	1,789	2,715	3,921	5,035	5,312	4,614	2,788	946	153	-
Black, both sexes												
2004.....	888	1,109	1,271	1,458	1,715	1,799	1,593	1,117	592	197	63	1
2003.....	877	1,128	1,287	1,506	1,702	1,789	1,584	1,080	567	199	49	-
2002.....	884	1,032	1,238	1,538	1,690	1,777	1,531	1,085	547	207	46	1
2001.....	826	1,003	1,186	1,480	1,715	1,798	1,455	1,035	531	192	51	-
2000.....	774	911	1,131	1,471	1,709	1,829	1,472	996	534	173	57	-
1999.....	735	981	1,217	1,469	1,728	1,686	1,415	1,013	523	154	43	2
Black, male												
2004.....	459	572	632	710	750	694	557	329	156	37	7	1
2003.....	442	563	637	699	716	685	539	299	128	24	3	-
2002.....	461	521	612	671	683	652	512	264	95	35	5	1
2001.....	428	497	558	647	673	683	510	249	125	33	7	-
2000.....	380	448	526	603	684	672	462	269	118	22	2	-
1999.....	366	475	543	648	651	608	445	288	130	23	6	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Endocrine diseases (E00-E34)												
Black, female												
2004.....	429	537	639	748	965	1,105	1,036	788	436	160	56	-
2003.....	435	565	650	807	986	1,104	1,045	781	439	175	46	-
2002.....	423	511	626	867	1,007	1,125	1,019	821	452	172	41	-
2001.....	398	506	628	833	1,042	1,115	945	786	406	159	44	-
2000.....	394	463	605	868	1,025	1,157	1,010	727	416	151	55	-
1999.....	369	506	674	821	1,077	1,078	970	725	393	131	37	1
American Indian 1/, both sexes												
2004.....	46	89	105	104	95	96	81	34	16	3	-	-
2003.....	69	68	94	105	104	91	83	49	16	11	-	-
2002.....	61	83	86	113	112	96	69	33	19	2	-	-
2001.....	52	62	81	76	97	91	69	39	17	3	1	-
2000.....	53	60	75	77	111	83	53	37	13	2	2	-
1999.....	51	74	92	105	114	95	55	51	17	5	1	-
American Indian 1/, male												
2004.....	28	49	57	46	36	56	24	12	4	-	-	-
2003.....	42	33	41	44	51	39	33	14	10	2	-	-
2002.....	31	41	45	48	49	35	23	11	6	1	-	-
2001.....	31	28	37	33	36	42	22	14	3	-	-	-
2000.....	23	26	42	33	39	37	14	15	5	1	1	-
1999.....	33	33	45	40	42	43	20	16	4	1	-	-
American Indian 1/, female												
2004.....	18	40	48	58	59	40	57	22	12	3	-	-
2003.....	27	35	53	61	53	52	50	35	6	9	-	-
2002.....	30	42	41	65	63	61	46	22	13	1	-	-
2001.....	21	34	44	43	61	49	47	25	14	3	1	-
2000.....	30	34	33	44	72	46	39	22	8	1	1	-
1999.....	18	41	47	65	72	52	35	35	13	4	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Endocrine diseases (E00-E34)												
Asian or Pacific Islander 2/, both sexes												
2004.....	67	100	129	137	200	222	259	181	93	30	6	-
2003.....	62	84	133	152	209	246	240	173	90	22	3	-
2002.....	57	91	103	177	198	252	196	181	58	17	3	-
2001.....	56	72	112	160	174	217	207	118	51	26	4	-
2000.....	55	72	86	145	152	187	181	101	61	21	1	-
1999.....	40	84	88	121	206	214	168	115	54	23	3	-
Asian or Pacific Islander 2/, male												
2004.....	36	63	75	77	101	103	106	71	35	8	3	-
2003.....	38	49	86	88	104	109	97	68	31	7	1	-
2002.....	37	59	56	104	86	94	78	85	20	7	-	-
2001.....	41	43	60	78	80	91	85	47	19	12	-	-
2000.....	33	47	43	83	72	86	72	42	29	10	1	-
1999.....	21	44	55	55	90	99	66	46	21	10	1	-
Asian or Pacific Islander 2/, female												
2004.....	31	37	54	60	99	119	153	110	58	22	3	-
2003.....	24	35	47	64	105	137	143	105	59	15	2	-
2002.....	20	32	47	73	112	158	118	96	38	10	3	-
2001.....	15	29	52	82	94	126	122	71	32	14	4	-
2000.....	22	25	43	62	80	101	109	59	32	11	-	-
1999.....	19	40	33	66	116	115	102	69	33	13	2	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of thyroid gland (E00-E07)												
All races, both sexes												
2004.....	31	32	59	53	88	161	295	343	343	192	54	-
2003.....	35	48	50	66	103	215	328	403	342	188	62	-
2002.....	37	39	49	52	119	184	319	435	402	165	39	-
2001.....	41	37	46	57	98	226	299	450	376	168	29	-
2000.....	32	44	47	50	112	198	290	352	327	168	47	-
1999.....	32	41	35	61	116	184	304	363	369	153	47	-
All races, male												
2004.....	9	13	24	22	29	45	74	69	64	29	7	-
2003.....	7	16	10	25	22	60	59	76	49	19	7	-
2002.....	12	11	15	19	32	48	61	82	72	19	4	-
2001.....	14	11	16	16	29	70	54	85	65	29	4	-
2000.....	7	17	14	16	37	47	48	67	60	17	4	-
1999.....	13	12	13	20	33	45	80	68	55	26	5	-
All races, female												
2004.....	22	19	35	31	59	116	221	274	279	163	47	-
2003.....	28	32	40	41	81	155	269	327	293	169	55	-
2002.....	25	28	34	33	87	136	258	353	330	146	35	-
2001.....	27	26	30	41	69	156	245	365	311	139	25	-
2000.....	25	27	33	34	75	151	242	285	267	151	43	-
1999.....	19	29	22	41	83	139	224	295	314	127	42	-
White, both sexes												
2004.....	25	24	46	39	68	132	263	309	327	180	53	-
2003.....	25	43	42	55	88	197	295	362	326	176	58	-
2002.....	23	27	38	46	99	168	298	411	374	152	38	-
2001.....	29	33	34	47	82	198	258	411	357	158	26	-
2000.....	24	35	44	36	93	171	262	328	305	157	40	-
1999.....	22	26	28	50	100	162	269	340	345	143	40	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of thyroid gland (E00-E07)												
White, male												
2004.....	8	11	18	18	23	38	64	65	61	25	7	-
2003.....	5	15	8	20	20	52	54	69	46	18	7	-
2002.....	7	7	9	17	28	43	58	80	65	16	4	-
2001.....	13	10	13	14	24	58	46	79	62	26	4	-
2000.....	4	12	13	12	27	43	46	62	51	16	4	-
1999.....	10	5	11	12	30	40	67	64	50	25	4	-
White, female												
2004.....	17	13	28	21	45	94	199	244	266	155	46	-
2003.....	20	28	34	35	68	145	241	293	280	158	51	-
2002.....	16	20	29	29	71	125	240	331	309	136	34	-
2001.....	16	23	21	33	58	140	212	332	295	132	22	-
2000.....	20	23	31	24	66	128	216	266	254	141	36	-
1999.....	12	21	17	38	70	122	202	276	295	118	36	-
Black, both sexes												
2004.....	5	7	12	12	20	27	25	28	13	11	1	-
2003.....	10	5	8	11	13	13	27	37	15	10	4	-
2002.....	13	10	11	5	18	14	17	21	22	12	1	-
2001.....	11	4	10	8	15	23	32	37	17	7	2	-
2000.....	6	8	3	13	18	24	21	22	20	11	7	-
1999.....	8	15	7	9	15	18	29	22	22	9	6	-
Black, male												
2004.....	1	2	5	4	6	6	7	2	2	4	-	-
2003.....	2	1	2	5	1	7	5	7	3	-	-	-
2002.....	5	3	6	1	4	4	2	2	4	2	-	-
2001.....	1	1	3	1	5	10	8	6	3	2	-	-
2000.....	2	5	1	3	9	3	2	5	8	1	-	-
1999.....	2	7	2	6	3	4	10	3	5	-	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of thyroid gland (E00-E07)												
Black, female												
2004.....	4	5	7	8	14	21	18	26	11	7	1	-
2003.....	8	4	6	6	12	6	22	30	12	10	4	-
2002.....	8	7	5	4	14	10	15	19	18	10	1	-
2001.....	10	3	7	7	10	13	24	31	14	5	2	-
2000.....	4	3	2	10	9	21	19	17	12	10	7	-
1999.....	6	8	5	3	12	14	19	19	17	9	5	-
American Indian 1/, both sexes												
2004.....	1	-	-	1	-	-	3	2	1	-	-	-
2003.....	-	-	-	-	-	-	1	-	-	1	-	-
2002.....	-	-	-	-	1	2	1	1	3	-	-	-
2001.....	1	-	1	-	1	-	3	2	-	-	1	-
2000.....	-	1	-	-	1	1	-	1	-	-	-	-
1999.....	2	-	-	-	-	1	3	-	1	-	1	-
American Indian 1/, male												
2004.....	-	-	-	-	-	-	1	1	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	1	-	-	2	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	1	-	-	-	-	-	-	-
1999.....	1	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, female												
2004.....	1	-	-	1	-	-	2	1	1	-	-	-
2003.....	-	-	-	-	-	-	1	-	-	1	-	-
2002.....	-	-	-	-	1	1	1	1	1	-	-	-
2001.....	1	-	1	-	1	-	3	2	-	-	1	-
2000.....	-	1	-	-	-	1	-	1	-	-	-	-
1999.....	1	-	-	-	-	1	2	-	1	-	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of thyroid gland (E00-E07)												
Asian or Pacific Islander 2/, both sexes												
2004.....	-	1	1	1	-	2	4	4	2	1	-	-
2003.....	-	-	-	-	2	5	5	4	1	1	-	-
2002.....	1	2	-	1	1	-	3	2	3	1	-	-
2001.....	-	-	1	2	-	5	6	-	2	3	-	-
2000.....	2	-	-	1	-	2	7	1	2	-	-	-
1999.....	-	-	-	2	1	3	3	1	1	1	-	-
Asian or Pacific Islander 2/, male												
2004.....	-	-	1	-	-	1	2	1	1	-	-	-
2003.....	-	-	-	-	1	1	-	-	-	1	-	-
2002.....	-	1	-	1	-	-	1	-	1	1	-	-
2001.....	-	-	-	1	-	2	-	-	-	1	-	-
2000.....	1	-	-	1	-	1	-	-	1	-	-	-
1999.....	-	-	-	2	-	1	2	1	-	1	-	-
Asian or Pacific Islander 2/, female												
2004.....	-	1	-	1	-	1	2	3	1	1	-	-
2003.....	-	-	-	-	1	4	5	4	1	-	-	-
2002.....	1	1	-	-	1	-	2	2	2	-	-	-
2001.....	-	-	1	1	-	3	6	-	2	2	-	-
2000.....	1	-	-	-	-	1	7	1	1	-	-	-
1999.....	-	-	-	-	1	2	1	-	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diabetes mellitus (E10-E14)												
All races, both sexes												
2004.....	3,350	4,842	5,938	7,102	8,991	11,098	11,847	8,711	4,643	1,337	227	3
2003.....	3,413	4,806	5,925	7,310	9,346	11,574	11,725	8,758	4,572	1,424	210	3
2002.....	3,390	4,510	5,512	7,249	9,460	11,703	11,579	8,797	4,375	1,357	195	1
2001.....	3,280	4,188	5,382	7,195	9,536	11,546	11,259	8,366	4,259	1,350	196	1
2000.....	3,037	3,992	5,194	7,054	9,620	11,607	10,577	8,002	4,104	1,250	200	-
1999.....	2,793	3,865	5,232	7,215	9,693	11,269	10,488	7,893	3,943	1,165	177	3
All races, male												
2004.....	2,017	2,758	3,395	3,832	4,598	5,335	5,234	3,296	1,408	286	45	1
2003.....	1,987	2,715	3,329	3,929	4,758	5,547	5,132	3,260	1,318	306	39	3
2002.....	1,977	2,616	3,067	3,742	4,675	5,402	4,893	3,248	1,254	314	28	1
2001.....	1,920	2,362	2,879	3,734	4,703	5,244	4,674	2,966	1,188	269	34	1
2000.....	1,697	2,173	2,770	3,556	4,670	5,288	4,450	2,789	1,165	260	20	-
1999.....	1,602	2,120	2,736	3,630	4,649	5,190	4,349	2,816	1,071	210	28	2
All races, female												
2004.....	1,333	2,084	2,543	3,270	4,393	5,763	6,613	5,415	3,235	1,051	182	2
2003.....	1,426	2,091	2,596	3,381	4,588	6,027	6,593	5,498	3,254	1,118	171	-
2002.....	1,413	1,894	2,445	3,507	4,785	6,301	6,686	5,549	3,121	1,043	167	-
2001.....	1,360	1,826	2,503	3,461	4,833	6,302	6,585	5,400	3,071	1,081	162	-
2000.....	1,340	1,819	2,424	3,498	4,950	6,319	6,127	5,213	2,939	990	180	-
1999.....	1,191	1,745	2,496	3,585	5,044	6,079	6,139	5,077	2,872	955	149	1
White, both sexes												
2004.....	2,372	3,567	4,463	5,434	7,022	9,025	9,963	7,429	3,965	1,123	160	2
2003.....	2,420	3,552	4,435	5,570	7,364	9,481	9,877	7,507	3,926	1,208	162	3
2002.....	2,419	3,329	4,111	5,439	7,503	9,611	9,822	7,532	3,787	1,144	147	-
2001.....	2,368	3,066	4,027	5,501	7,584	9,488	9,582	7,224	3,685	1,145	143	1
2000.....	2,170	2,967	3,912	5,393	7,682	9,561	8,917	6,905	3,525	1,067	148	-
1999.....	1,988	2,754	3,855	5,547	7,683	9,315	8,902	6,746	3,379	995	137	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diabetes mellitus (E10-E14)												
White, male												
2004.....	1,503	2,081	2,647	3,014	3,727	4,497	4,564	2,895	1,218	247	36	-
2003.....	1,471	2,082	2,572	3,106	3,896	4,731	4,479	2,889	1,157	276	35	3
2002.....	1,462	2,004	2,367	2,928	3,872	4,630	4,290	2,891	1,141	274	23	-
2001.....	1,428	1,801	2,230	2,986	3,931	4,452	4,068	2,665	1,046	229	27	1
2000.....	1,269	1,662	2,161	2,849	3,890	4,508	3,911	2,473	1,023	228	16	-
1999.....	1,191	1,582	2,102	2,906	3,881	4,455	3,838	2,474	924	178	22	1
White, female												
2004.....	869	1,486	1,816	2,420	3,295	4,528	5,399	4,534	2,747	876	124	2
2003.....	949	1,470	1,863	2,464	3,468	4,750	5,398	4,618	2,769	932	127	-
2002.....	957	1,325	1,744	2,511	3,631	4,981	5,532	4,641	2,646	870	124	-
2001.....	940	1,265	1,797	2,515	3,653	5,036	5,514	4,559	2,639	916	116	-
2000.....	901	1,305	1,751	2,544	3,792	5,053	5,006	4,432	2,502	839	132	-
1999.....	797	1,172	1,753	2,641	3,802	4,860	5,064	4,272	2,455	817	115	-
Black, both sexes												
2004.....	866	1,088	1,246	1,431	1,676	1,758	1,552	1,073	572	182	61	1
2003.....	862	1,102	1,265	1,488	1,672	1,763	1,536	1,035	543	185	45	-
2002.....	854	1,010	1,214	1,522	1,653	1,748	1,501	1,054	517	195	45	1
2001.....	807	988	1,166	1,460	1,685	1,757	1,411	988	509	179	49	-
2000.....	762	894	1,121	1,441	1,677	1,781	1,434	962	507	160	49	-
1999.....	716	954	1,197	1,444	1,693	1,652	1,370	982	496	144	37	2
Black, male												
2004.....	450	565	619	696	736	681	543	320	152	31	6	1
2003.....	436	551	631	693	708	671	525	289	121	22	3	-
2002.....	447	514	600	663	669	644	506	261	90	33	5	1
2001.....	421	490	553	638	658	663	499	240	121	29	7	-
2000.....	374	438	524	593	671	659	454	260	109	21	2	-
1999.....	358	461	534	631	637	595	429	281	123	23	5	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diabetes mellitus (E10-E14)												
Black, female												
2004.....	416	523	627	735	940	1,077	1,009	753	420	151	55	-
2003.....	426	551	634	795	964	1,092	1,011	746	422	163	42	-
2002.....	407	496	614	859	984	1,104	995	793	427	162	40	-
2001.....	386	498	613	822	1,027	1,094	912	748	388	150	42	-
2000.....	388	456	597	848	1,006	1,122	980	702	398	139	47	-
1999.....	358	493	663	813	1,056	1,057	941	701	373	121	32	1
American Indian 1/, both sexes												
2004.....	45	88	105	103	94	96	77	32	15	3	-	-
2003.....	69	68	94	102	104	91	82	48	15	10	-	-
2002.....	61	83	84	113	110	94	68	32	16	2	-	-
2001.....	50	62	80	76	95	90	66	37	16	3	-	-
2000.....	53	59	75	77	109	81	52	36	13	2	2	-
1999.....	49	73	92	105	114	94	52	51	16	5	-	-
American Indian 1/, male												
2004.....	28	49	57	46	35	56	23	11	4	-	-	-
2003.....	42	33	41	44	51	39	33	14	9	2	-	-
2002.....	31	41	44	48	48	34	23	11	4	1	-	-
2001.....	30	28	37	33	35	41	22	14	2	-	-	-
2000.....	23	26	42	33	37	37	13	15	5	1	1	-
1999.....	32	33	45	40	42	43	19	16	4	1	-	-
American Indian 1/, female												
2004.....	17	39	48	57	59	40	54	21	11	3	-	-
2003.....	27	35	53	58	53	52	49	34	6	8	-	-
2002.....	30	42	40	65	62	60	45	21	12	1	-	-
2001.....	20	34	43	43	60	49	44	23	14	3	-	-
2000.....	30	33	33	44	72	44	39	21	8	1	1	-
1999.....	17	40	47	65	72	51	33	35	12	4	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diabetes mellitus (E10-E14)												
Asian or Pacific Islander 2/, both sexes												
2004.....	67	99	124	134	199	219	255	177	91	29	6	-
2003.....	62	84	131	150	206	239	230	168	88	21	3	-
2002.....	56	88	103	175	194	250	188	179	55	16	3	-
2001.....	55	72	109	158	172	211	200	117	49	23	4	-
2000.....	52	72	86	143	152	184	174	99	59	21	1	-
1999.....	40	84	88	119	203	208	164	114	52	21	3	-
Asian or Pacific Islander 2/, male												
2004.....	36	63	72	76	100	101	104	70	34	8	3	-
2003.....	38	49	85	86	103	106	95	68	31	6	1	-
2002.....	37	57	56	103	86	94	74	85	19	6	-	-
2001.....	41	43	59	77	79	88	85	47	19	11	-	-
2000.....	31	47	43	81	72	84	72	41	28	10	1	-
1999.....	21	44	55	53	89	97	63	45	20	8	1	-
Asian or Pacific Islander 2/, female												
2004.....	31	36	52	58	99	118	151	107	57	21	3	-
2003.....	24	35	46	64	103	133	135	100	57	15	2	-
2002.....	19	31	47	72	108	156	114	94	36	10	3	-
2001.....	14	29	50	81	93	123	115	70	30	12	4	-
2000.....	21	25	43	62	80	100	102	58	31	11	-	-
1999.....	19	40	33	66	114	111	101	69	32	13	2	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of pituitary gland (E22-E23)												
All races, both sexes												
2004.....	13	10	17	12	25	24	33	25	16	4	1	-
2003.....	7	13	16	9	18	26	32	23	12	7	-	-
2002.....	7	14	14	11	20	32	24	31	14	3	1	-
2001.....	13	11	17	9	16	29	33	26	12	4	1	-
2000.....	10	8	9	21	25	25	37	28	14	2	3	-
1999.....	5	12	15	20	22	35	25	28	14	2	-	-
All races, male												
2004.....	8	4	7	9	11	15	10	10	4	-	-	-
2003.....	2	9	5	7	11	16	14	7	2	-	-	-
2002.....	4	6	8	6	10	12	8	11	4	-	-	-
2001.....	7	7	10	5	9	12	13	4	5	-	-	-
2000.....	5	3	6	12	14	11	7	10	3	1	-	-
1999.....	2	4	8	7	8	22	13	6	3	-	-	-
All races, female												
2004.....	5	6	10	3	14	9	23	15	12	4	1	-
2003.....	5	4	11	2	7	10	18	16	10	7	-	-
2002.....	3	8	6	5	10	20	16	20	10	3	1	-
2001.....	6	4	7	4	7	17	20	22	7	4	1	-
2000.....	5	5	3	9	11	14	30	18	11	1	3	-
1999.....	3	8	7	13	14	13	12	22	11	2	-	-
White, both sexes												
2004.....	9	7	14	12	20	21	29	22	15	4	1	-
2003.....	7	10	12	7	17	23	26	22	8	7	-	-
2002.....	5	12	13	10	16	28	21	29	13	3	1	-
2001.....	10	8	15	7	16	27	32	25	11	3	1	-
2000.....	8	7	8	15	23	23	34	26	12	2	3	-
1999.....	4	9	10	14	18	25	21	26	14	2	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of pituitary gland (E22-E23)												
White, male												
2004.....	5	2	6	9	10	14	8	9	3	-	-	-
2003.....	2	7	5	6	11	14	12	6	-	-	-	-
2002.....	3	5	8	5	8	12	7	10	3	-	-	-
2001.....	5	5	10	4	9	10	12	4	4	-	-	-
2000.....	4	3	6	7	13	9	5	8	3	1	-	-
1999.....	2	3	6	3	6	16	10	6	3	-	-	-
White, female												
2004.....	4	5	8	3	10	7	21	13	12	4	1	-
2003.....	5	3	7	1	6	9	14	16	8	7	-	-
2002.....	2	7	5	5	8	16	14	19	10	3	1	-
2001.....	5	3	5	3	7	17	20	21	7	3	1	-
2000.....	4	4	2	8	10	14	29	18	9	1	3	-
1999.....	2	6	4	11	12	9	11	20	11	2	-	-
Black, both sexes												
2004.....	4	3	2	-	5	3	3	3	1	-	-	-
2003.....	-	3	3	1	1	3	4	1	2	-	-	-
2002.....	2	1	1	1	3	3	2	2	1	-	-	-
2001.....	1	3	2	2	-	2	1	1	1	1	-	-
2000.....	2	1	1	6	2	2	3	1	2	-	-	-
1999.....	1	2	5	6	2	8	3	2	-	-	-	-
Black, male												
2004.....	3	2	1	-	1	1	2	1	1	-	-	-
2003.....	-	2	-	-	-	2	1	1	1	-	-	-
2002.....	1	-	-	1	2	-	-	1	1	-	-	-
2001.....	1	2	-	1	-	2	1	-	1	-	-	-
2000.....	1	-	-	5	1	2	2	1	-	-	-	-
1999.....	-	1	2	4	1	5	2	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of pituitary gland (E22-E23)												
Asian or Pacific Islander 2/, both sexes												
2004.....	-	-	1	-	-	-	-	-	-	-	-	-
2003.....	-	-	1	1	-	-	2	-	1	-	-	-
2002.....	-	1	-	-	1	1	1	-	-	-	-	-
2001.....	1	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	1	-	-	-	-
1999.....	-	-	-	-	2	2	1	-	-	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	1	-	-	1	-	-	-	-	-
2002.....	-	1	-	-	-	-	1	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	1	-	-	-	-
1999.....	-	-	-	-	1	1	1	-	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	-	-	1	-	-	-	-	-	-	-	-	-
2003.....	-	-	1	-	-	-	1	-	1	-	-	-
2002.....	-	-	-	-	1	1	-	-	-	-	-	-
2001.....	1	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	1	1	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of adrenal glands (E24-E27)												
All races, both sexes												
2004.....	13	17	20	19	24	26	40	28	19	9	1	-
2003.....	13	23	20	22	27	43	43	26	24	5	-	-
2002.....	12	10	20	23	29	35	46	24	22	3	1	-
2001.....	18	11	13	17	41	52	50	43	28	5	3	-
2000.....	12	23	14	21	20	39	50	27	11	4	2	-
1999.....	20	8	12	25	27	40	21	23	13	3	2	-
All races, male												
2004.....	2	6	6	9	10	11	13	9	6	4	-	-
2003.....	7	14	4	6	10	20	12	8	4	1	-	-
2002.....	3	5	10	8	9	11	16	9	4	1	-	-
2001.....	13	3	7	12	16	20	24	20	7	-	-	-
2000.....	7	9	2	8	6	17	18	5	4	2	-	-
1999.....	10	4	6	8	6	16	12	7	3	-	-	-
All races, female												
2004.....	11	11	14	10	14	15	27	19	13	5	1	-
2003.....	6	9	16	16	17	23	31	18	20	4	-	-
2002.....	9	5	10	15	20	24	30	15	18	2	1	-
2001.....	5	8	6	5	25	32	26	23	21	5	3	-
2000.....	5	14	12	13	14	22	32	22	7	2	2	-
1999.....	10	4	6	17	21	24	9	16	10	3	2	-
White, both sexes												
2004.....	10	14	15	16	19	24	38	26	18	7	1	-
2003.....	11	17	17	19	26	37	41	24	22	3	-	-
2002.....	8	9	13	19	23	32	41	24	19	3	1	-
2001.....	14	10	8	16	40	47	46	41	26	3	3	-
2000.....	10	20	12	16	18	34	47	26	10	3	2	-
1999.....	20	4	11	22	22	38	20	21	11	3	2	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Disorders of adrenal glands (E24-E27)

White, male

2004.....	1	6	3	7	8	9	12	8	6	3	-	-
2003.....	6	9	4	6	9	18	10	8	4	-	-	-
2002.....	2	5	5	5	7	10	14	9	4	1	-	-
2001.....	9	3	6	11	16	17	22	19	6	-	-	-
2000.....	6	7	2	7	6	15	17	4	4	2	-	-
1999.....	10	2	6	7	5	15	11	6	2	-	-	-

White, female

2004.....	9	8	12	9	11	15	26	18	12	4	1	-
2003.....	5	8	13	13	17	19	31	16	18	3	-	-
2002.....	6	4	8	14	16	22	27	15	15	2	1	-
2001.....	5	7	2	5	24	30	24	22	20	3	3	-
2000.....	4	13	10	9	12	19	30	22	6	1	2	-
1999.....	10	2	5	15	17	23	9	15	9	3	2	-

Black, both sexes

2004.....	3	3	4	3	4	1	2	2	1	2	-	-
2003.....	2	6	3	1	1	5	2	-	2	2	-	-
2002.....	4	1	5	4	4	2	3	-	3	-	-	-
2001.....	4	1	4	1	1	4	4	2	1	2	-	-
2000.....	2	3	2	5	2	5	2	1	1	1	-	-
1999.....	-	4	1	3	5	2	1	2	2	-	-	-

Black, male

2004.....	1	-	2	2	1	1	1	1	-	1	-	-
2003.....	1	5	-	-	1	1	2	-	-	1	-	-
2002.....	1	-	4	3	1	1	1	-	-	-	-	-
2001.....	4	-	1	1	-	2	2	1	-	-	-	-
2000.....	1	2	-	1	-	2	-	1	-	-	-	-
1999.....	-	2	-	1	1	1	1	1	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of thymus (E32)												
All races, both sexes												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	1	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	1	-	-	-	-	-
2001.....	-	-	-	-	-	-	1	1	-	-	-	-
2000.....	-	-	1	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	1	-	-	-	-	-	-	-
All races, male												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	1	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	1	-	-	-	-	-
2000.....	-	-	1	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-
All races, female												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	1	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	1	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	1	-	-	-	-	-	-	-
White, both sexes												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	1	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	1	-	-	-	-	-
2001.....	-	-	-	-	-	-	1	1	-	-	-	-
2000.....	-	-	1	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	1	-	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other endocrine diseases (E15-E21,E28-E31,E34)												
All races, both sexes												
2004.....	26	30	30	42	32	60	65	55	30	16	1	-
2003.....	19	31	30	25	43	46	67	61	31	6	-	1
2002.....	32	27	28	32	45	58	67	60	32	6	2	-
2001.....	22	20	28	23	41	54	51	52	41	15	1	-
2000.....	18	19	17	20	32	66	45	51	29	9	4	-
1999.....	28	21	23	23	50	49	61	47	24	8	-	-
All races, male												
2004.....	12	15	19	24	13	31	28	18	6	2	1	-
2003.....	15	9	17	9	23	21	25	14	9	2	-	-
2002.....	17	16	13	15	18	22	23	19	6	-	-	-
2001.....	7	10	11	12	24	24	13	13	10	6	-	-
2000.....	11	10	6	5	12	26	17	16	5	-	-	-
1999.....	14	13	11	12	27	24	26	14	4	1	-	-
All races, female												
2004.....	14	15	11	18	19	29	37	37	24	14	-	-
2003.....	4	22	13	16	20	25	42	47	22	4	-	1
2002.....	15	11	15	17	27	36	44	41	26	6	2	-
2001.....	15	10	17	11	17	30	38	39	31	9	1	-
2000.....	7	9	11	15	20	40	28	35	24	9	4	-
1999.....	14	8	12	11	23	25	35	33	20	7	-	-
White, both sexes												
2004.....	16	21	21	28	21	50	54	44	25	14	-	-
2003.....	16	19	21	18	27	40	49	54	26	4	-	1
2002.....	21	17	21	25	32	48	57	52	28	6	2	-
2001.....	19	13	23	14	24	41	43	44	38	12	1	-
2000.....	15	14	13	13	21	47	33	41	25	8	3	-
1999.....	18	15	16	16	37	42	49	42	20	6	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other endocrine diseases (E15-E21,E28-E31,E34)

White, male

2004.....	8	12	13	15	6	26	24	13	5	1	-	-
2003.....	12	5	12	7	17	16	18	12	6	1	-	-
2002.....	10	12	11	12	11	19	19	19	6	-	-	-
2001.....	6	6	9	6	12	17	13	11	10	4	-	-
2000.....	8	7	5	3	8	19	13	14	4	-	-	-
1999.....	8	9	6	6	18	21	23	11	2	-	-	-

White, female

2004.....	8	9	8	13	15	24	30	31	20	13	-	-
2003.....	4	14	9	11	10	24	31	42	20	3	-	1
2002.....	11	5	10	13	21	29	38	33	22	6	2	-
2001.....	13	7	14	8	12	24	30	33	28	8	1	-
2000.....	7	7	8	10	13	28	20	27	21	8	3	-
1999.....	10	6	10	10	19	21	26	31	18	6	-	-

Black, both sexes

2004.....	10	8	7	12	10	10	11	11	5	2	1	-
2003.....	3	12	8	5	15	5	15	7	5	2	-	-
2002.....	11	10	7	6	12	10	8	8	4	-	-	-
2001.....	3	7	4	9	14	12	7	7	3	3	-	-
2000.....	2	5	4	6	10	17	12	10	4	1	1	-
1999.....	10	6	7	7	13	6	12	5	3	1	-	-

Black, male

2004.....	4	3	5	8	6	5	4	5	1	1	1	-
2003.....	3	4	4	1	6	4	6	2	3	1	-	-
2002.....	7	4	2	3	7	3	3	-	-	-	-	-
2001.....	1	4	1	6	10	6	-	2	-	2	-	-
2000.....	2	3	1	1	3	6	4	2	1	-	-	-
1999.....	6	4	5	6	9	3	3	3	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other endocrine diseases (E15-E21,E28-E31,E34)												
Asian or Pacific Islander 2/, both sexes												
2004.....	-	-	2	2	-	-	-	-	-	-	-	-
2003.....	-	-	1	1	1	1	3	-	-	-	-	-
2002.....	-	-	-	1	1	-	2	-	-	-	-	-
2001.....	-	-	1	-	2	1	1	1	-	-	-	-
2000.....	1	-	-	1	-	1	-	-	-	-	-	-
1999.....	-	-	-	-	-	1	-	-	1	1	-	-
Asian or Pacific Islander 2/, male												
2004.....	-	-	1	1	-	-	-	-	-	-	-	-
2003.....	-	-	1	1	-	1	1	-	-	-	-	-
2002.....	-	-	-	-	-	-	1	-	-	-	-	-
2001.....	-	-	1	-	1	1	-	-	-	-	-	-
2000.....	1	-	-	1	-	1	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	1	1	-	-
Asian or Pacific Islander 2/, female												
2004.....	-	-	1	1	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	1	-	2	-	-	-	-	-
2002.....	-	-	-	1	1	-	1	-	-	-	-	-
2001.....	-	-	-	-	1	-	1	1	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	1	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nutritional diseases (E40-E68)												
All races, both sexes												
2004.....	658	612	538	482	493	572	710	713	539	254	64	1
2003.....	588	588	501	492	496	619	697	793	589	310	74	2
2002.....	567	571	455	450	439	644	797	882	681	314	93	-
2001.....	518	442	393	368	477	604	749	851	696	327	85	-
2000.....	480	393	386	426	481	676	887	996	796	307	93	-
1999.....	394	363	315	362	496	631	876	986	836	383	94	-
All races, male												
2004.....	337	305	237	219	190	236	261	243	137	48	6	1
2003.....	286	299	225	229	198	234	264	252	142	50	11	2
2002.....	284	272	201	189	188	251	271	291	172	50	13	-
2001.....	238	197	178	148	205	212	278	235	172	55	3	-
2000.....	239	212	165	174	208	258	290	297	184	53	13	-
1999.....	188	199	151	156	224	252	308	302	199	71	9	-
All races, female												
2004.....	321	307	301	263	303	336	449	470	402	206	58	-
2003.....	302	289	276	263	298	385	433	541	447	260	63	-
2002.....	283	299	254	261	251	393	526	591	509	264	80	-
2001.....	280	245	215	220	272	392	471	616	524	272	82	-
2000.....	241	181	221	252	273	418	597	699	612	254	80	-
1999.....	206	164	164	206	272	379	568	684	637	312	85	-
White, both sexes												
2004.....	521	509	440	383	409	476	618	610	472	225	52	-
2003.....	477	492	414	431	416	534	597	705	513	265	55	1
2002.....	452	457	368	362	366	541	688	769	605	276	77	-
2001.....	400	364	322	297	390	503	635	765	614	289	78	-
2000.....	385	306	321	353	393	569	780	881	727	258	74	-
1999.....	319	288	253	296	386	527	745	869	765	329	80	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nutritional diseases (E40-E68)												
White, male												
2004.....	285	265	201	174	169	197	228	213	118	43	5	-
2003.....	240	261	195	207	163	197	225	226	123	43	7	1
2002.....	246	226	169	160	157	221	227	257	157	43	10	-
2001.....	192	169	144	119	171	181	236	210	146	48	2	-
2000.....	204	172	133	147	169	216	256	263	169	41	11	-
1999.....	162	159	124	128	175	207	260	266	178	55	7	-
White, female												
2004.....	236	244	239	209	240	279	390	397	354	182	47	-
2003.....	237	231	219	224	253	337	372	479	390	222	48	-
2002.....	206	231	199	202	209	320	461	512	448	233	67	-
2001.....	208	195	178	178	219	322	399	555	468	241	76	-
2000.....	181	134	188	206	224	353	524	618	558	217	63	-
1999.....	157	129	129	168	211	320	485	603	587	274	73	-
Black, both sexes												
2004.....	126	100	93	90	79	82	75	89	56	24	11	1
2003.....	104	88	81	53	75	79	90	77	70	37	15	1
2002.....	110	112	81	82	66	95	95	95	68	33	13	-
2001.....	110	76	64	63	79	89	98	75	71	30	7	-
2000.....	91	84	58	68	78	95	94	100	61	42	16	-
1999.....	71	69	59	60	99	91	119	102	64	47	14	-
Black, male												
2004.....	49	38	32	43	18	34	25	24	16	5	1	1
2003.....	44	32	28	20	32	36	34	22	16	5	3	1
2002.....	37	45	29	28	26	27	39	28	12	3	2	-
2001.....	41	27	30	25	31	29	33	24	21	5	1	-
2000.....	34	39	27	25	36	38	30	32	12	12	1	-
1999.....	25	36	25	24	44	38	41	30	19	13	2	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Nutritional diseases (E40-E68)

Black, female

2004.....	77	62	61	47	61	48	50	65	40	19	10	-
2003.....	60	56	53	33	43	43	56	55	54	32	12	-
2002.....	73	67	52	54	40	68	56	67	56	30	11	-
2001.....	69	49	34	38	48	60	65	51	50	25	6	-
2000.....	57	45	31	43	42	57	64	68	49	30	15	-
1999.....	46	33	34	36	55	53	78	72	45	34	12	-

American Indian 1/, both sexes

2004.....	5	3	3	2	2	7	6	4	3	-	-	-
2003.....	3	6	1	3	3	5	4	3	2	2	1	-
2002.....	3	1	3	3	5	2	6	6	3	3	1	-
2001.....	8	1	5	5	5	3	6	6	2	1	-	-
2000.....	2	3	3	2	5	5	3	8	3	1	3	-
1999.....	4	5	1	4	5	4	5	5	3	1	-	-

American Indian 1/, male

2004.....	1	2	3	1	1	2	3	1	-	-	-	-
2003.....	1	4	1	1	2	-	2	-	-	-	-	-
2002.....	-	1	1	-	4	-	1	2	-	2	-	-
2001.....	5	-	3	2	3	-	3	-	-	-	-	-
2000.....	-	1	1	1	1	2	1	-	1	-	1	-
1999.....	1	3	1	2	1	3	3	1	-	-	-	-

American Indian 1/, female

2004.....	4	1	-	1	1	5	3	3	3	-	-	-
2003.....	2	2	-	2	1	5	2	3	2	2	1	-
2002.....	3	-	2	3	1	2	5	4	3	1	1	-
2001.....	3	1	2	3	2	3	3	6	2	1	-	-
2000.....	2	2	2	1	4	3	2	8	2	1	2	-
1999.....	3	2	-	2	4	1	2	4	3	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nutritional diseases (E40-E68)												
Asian or Pacific Islander 2/, both sexes												
2004.....	6	-	2	7	3	7	11	10	8	5	1	-
2003.....	4	2	5	5	2	1	6	8	4	6	3	-
2002.....	2	1	3	3	2	6	8	12	5	2	2	-
2001.....	-	1	2	3	3	9	10	5	9	7	-	-
2000.....	2	-	4	3	5	7	10	7	5	6	-	-
1999.....	-	1	2	2	6	9	7	10	4	6	-	-
Asian or Pacific Islander 2/, male												
2004.....	2	-	1	1	2	3	5	5	3	-	-	-
2003.....	1	2	1	1	1	1	3	4	3	2	1	-
2002.....	1	-	2	1	1	3	4	4	3	2	1	-
2001.....	-	1	1	2	-	2	6	1	5	2	-	-
2000.....	1	-	4	1	2	2	3	2	2	-	-	-
1999.....	-	1	1	2	4	4	4	5	2	3	-	-
Asian or Pacific Islander 2/, female												
2004.....	4	-	1	6	1	4	6	5	5	5	1	-
2003.....	3	-	4	4	1	-	3	4	1	4	2	-
2002.....	1	1	1	2	1	3	4	8	2	-	1	-
2001.....	-	-	1	1	3	7	4	4	4	5	-	-
2000.....	1	-	-	2	3	5	7	5	3	6	-	-
1999.....	-	-	1	-	2	5	3	5	2	3	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malnutrition (E40-E46)												
All races, both sexes												
2004.....	73	68	84	116	172	328	536	613	471	227	62	1
2003.....	58	74	86	118	202	413	539	684	537	284	68	-
2002.....	55	73	83	136	218	419	648	778	616	283	87	-
2001.....	57	62	84	123	224	406	619	756	643	302	71	-
2000.....	46	61	97	178	245	478	721	888	712	280	84	-
1999.....	57	66	88	141	266	439	720	883	755	353	86	-
All races, male												
2004.....	32	40	41	58	83	148	194	212	120	43	6	1
2003.....	28	37	38	59	96	166	213	222	130	46	10	-
2002.....	24	39	45	64	110	176	230	267	163	45	13	-
2001.....	25	33	42	57	100	156	244	220	164	50	2	-
2000.....	18	39	47	79	128	194	240	275	165	46	13	-
1999.....	27	38	54	68	139	186	266	282	175	65	8	-
All races, female												
2004.....	41	28	43	58	89	180	342	401	351	184	56	-
2003.....	30	37	48	59	106	247	326	462	407	238	58	-
2002.....	31	34	38	72	108	243	418	511	453	238	74	-
2001.....	32	29	42	66	124	250	375	536	479	252	69	-
2000.....	28	22	50	99	117	284	481	613	547	234	71	-
1999.....	30	28	34	73	127	253	454	601	580	288	78	-
White, both sexes												
2004.....	51	55	63	89	137	268	466	521	409	200	50	-
2003.....	35	57	63	94	165	355	455	603	469	243	50	-
2002.....	42	51	61	105	180	356	561	675	544	247	72	-
2001.....	37	45	68	93	189	338	521	679	566	266	64	-
2000.....	31	45	74	146	189	405	634	781	650	235	67	-
1999.....	46	41	66	107	198	366	610	779	689	302	72	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malnutrition (E40-E46)												
White, male												
2004.....	23	30	29	39	68	120	166	183	102	38	5	-
2003.....	18	25	26	49	73	136	176	197	112	40	7	-
2002.....	17	24	29	51	91	153	192	235	148	38	10	-
2001.....	13	22	30	40	81	130	205	195	139	44	1	-
2000.....	11	26	32	63	96	162	209	242	152	35	11	-
1999.....	21	20	41	48	98	152	221	247	154	49	6	-
White, female												
2004.....	28	25	34	50	69	148	300	338	307	162	45	-
2003.....	17	32	37	45	92	219	279	406	357	203	43	-
2002.....	25	27	32	54	89	203	369	440	396	209	62	-
2001.....	24	23	38	53	108	208	316	484	427	222	63	-
2000.....	20	19	42	83	93	243	425	539	498	200	56	-
1999.....	25	21	25	59	100	214	389	532	535	253	66	-
Black, both sexes												
2004.....	20	12	19	24	32	48	55	78	51	23	11	1
2003.....	21	17	21	19	33	52	76	73	62	33	15	-
2002.....	12	22	22	28	33	59	76	86	65	31	12	-
2001.....	18	15	12	26	30	58	82	67	66	28	7	-
2000.....	15	16	21	28	49	63	74	94	57	39	14	-
1999.....	11	21	21	29	59	62	100	90	59	44	14	-
Black, male												
2004.....	9	9	11	18	13	23	20	23	15	5	1	1
2003.....	10	12	11	9	21	29	32	22	15	4	3	-
2002.....	7	15	16	12	15	20	33	26	12	3	2	-
2001.....	11	10	9	16	18	24	30	24	20	4	1	-
2000.....	7	13	13	14	29	30	27	31	10	11	1	-
1999.....	6	15	12	17	36	28	38	29	19	13	2	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malnutrition (E40-E46)												
Black, female												
2004.....	11	3	8	6	19	25	35	55	36	18	10	-
2003.....	11	5	10	10	12	23	44	51	47	29	12	-
2002.....	5	7	6	16	18	39	43	60	53	28	10	-
2001.....	7	5	3	10	12	34	52	43	46	24	6	-
2000.....	8	3	8	14	20	33	47	63	47	28	13	-
1999.....	5	6	9	12	23	34	62	61	40	31	12	-
American Indian 1/, both sexes												
2004.....	1	1	1	2	1	5	5	4	3	-	-	-
2003.....	2	-	-	1	3	5	2	2	2	2	1	-
2002.....	1	-	-	1	3	1	4	5	3	3	1	-
2001.....	2	1	2	3	3	3	6	5	2	1	-	-
2000.....	-	-	-	2	3	4	3	6	2	1	3	-
1999.....	-	3	1	3	3	3	4	4	3	1	-	-
American Indian 1/, male												
2004.....	-	1	1	1	-	2	3	1	-	-	-	-
2003.....	-	-	-	-	2	-	2	-	-	-	-	-
2002.....	-	-	-	-	3	-	1	2	-	2	-	-
2001.....	1	-	2	1	1	-	3	-	-	-	-	-
2000.....	-	-	-	1	1	1	1	-	1	-	1	-
1999.....	-	2	1	1	1	2	3	1	-	-	-	-
American Indian 1/, female												
2004.....	1	-	-	1	1	3	2	3	3	-	-	-
2003.....	2	-	-	1	1	5	-	2	2	2	1	-
2002.....	1	-	-	1	-	1	3	3	3	1	1	-
2001.....	1	1	-	2	2	3	3	5	2	1	-	-
2000.....	-	-	-	1	2	3	2	6	1	1	2	-
1999.....	-	1	-	2	2	1	1	3	3	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malnutrition (E40-E46)												
Asian or Pacific Islander 2/, both sexes												
2004.....	1	-	1	1	2	7	10	10	8	4	1	-
2003.....	-	-	2	4	1	1	6	6	4	6	2	-
2002.....	-	-	-	2	2	3	7	12	4	2	2	-
2001.....	-	1	2	1	2	7	10	5	9	7	-	-
2000.....	-	-	2	2	4	6	10	7	3	5	-	-
1999.....	-	1	-	2	6	8	6	10	4	6	-	-
Asian or Pacific Islander 2/, male												
2004.....	-	-	-	-	2	3	5	5	3	-	-	-
2003.....	-	-	1	1	-	1	3	3	3	2	-	-
2002.....	-	-	-	1	1	3	4	4	3	2	1	-
2001.....	-	1	1	-	-	2	6	1	5	2	-	-
2000.....	-	-	2	1	2	1	3	2	2	-	-	-
1999.....	-	1	-	2	4	4	4	5	2	3	-	-
Asian or Pacific Islander 2/, female												
2004.....	1	-	1	1	-	4	5	5	5	4	1	-
2003.....	-	-	1	3	1	-	3	3	1	4	2	-
2002.....	-	-	-	1	1	-	3	8	1	-	1	-
2001.....	-	-	1	1	2	5	4	4	4	5	-	-
2000.....	-	-	-	1	2	5	7	5	1	5	-	-
1999.....	-	-	-	-	2	4	2	5	2	3	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nutritional marasmus (E41)												
All races, both sexes												
2004.....	4	2	2	5	7	9	26	25	26	9	1	-
2003.....	2	2	3	8	6	20	31	29	28	14	3	-
2002.....	7	4	4	4	7	18	32	51	34	17	7	-
2001.....	4	3	5	6	10	18	33	44	38	26	3	-
2000.....	3	3	5	5	10	18	40	41	48	20	7	-
1999.....	2	4	2	6	19	20	38	38	38	24	2	-
All races, male												
2004.....	-	1	2	3	4	4	9	7	6	2	-	-
2003.....	2	2	1	6	1	9	10	8	9	-	-	-
2002.....	3	2	1	1	2	5	12	10	11	1	1	-
2001.....	2	1	4	2	3	9	15	16	11	2	-	-
2000.....	1	3	2	3	5	9	13	18	13	3	1	-
1999.....	-	3	2	3	11	6	18	16	13	7	-	-
All races, female												
2004.....	4	1	-	2	3	5	17	18	20	7	1	-
2003.....	-	-	2	2	5	11	21	21	19	14	3	-
2002.....	4	2	3	3	5	13	20	41	23	16	6	-
2001.....	2	2	1	4	7	9	18	28	27	24	3	-
2000.....	2	-	3	2	5	9	27	23	35	17	6	-
1999.....	2	1	-	3	8	14	20	22	25	17	2	-
White, both sexes												
2004.....	4	2	1	4	4	9	26	22	21	9	1	-
2003.....	1	2	1	6	5	16	25	28	26	12	2	-
2002.....	5	3	3	3	6	12	28	45	32	16	6	-
2001.....	3	2	2	4	8	18	27	38	35	24	3	-
2000.....	2	2	5	4	8	14	40	36	47	17	3	-
1999.....	1	3	1	4	15	12	36	35	33	22	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nutritional marasmus (E41)												
White, male												
2004.....	-	1	1	2	3	4	9	6	4	2	-	-
2003.....	1	2	-	4	1	7	9	8	8	-	-	-
2002.....	2	1	-	1	2	3	9	9	11	1	1	-
2001.....	1	-	2	2	2	9	13	13	10	2	-	-
2000.....	-	2	2	2	4	6	13	16	12	2	-	-
1999.....	-	2	1	2	9	5	16	14	12	5	-	-
White, female												
2004.....	4	1	-	2	1	5	17	16	17	7	1	-
2003.....	-	-	1	2	4	9	16	20	18	12	2	-
2002.....	3	2	3	2	4	9	19	36	21	15	5	-
2001.....	2	2	-	2	6	9	14	25	25	22	3	-
2000.....	2	-	3	2	4	8	27	20	35	15	3	-
1999.....	1	1	-	2	6	7	20	21	21	17	1	-
Black, both sexes												
2004.....	-	-	1	1	2	-	-	2	2	-	-	-
2003.....	1	-	2	2	1	3	5	-	2	1	-	-
2002.....	1	1	1	1	1	5	4	3	2	1	-	-
2001.....	1	1	2	1	1	-	5	4	2	2	-	-
2000.....	1	1	-	1	2	3	-	5	1	3	4	-
1999.....	1	1	1	1	4	8	1	3	5	1	1	-
Black, male												
2004.....	-	-	1	1	-	-	-	-	1	-	-	-
2003.....	1	-	1	2	-	2	1	-	1	-	-	-
2002.....	1	1	1	-	-	1	3	1	-	-	-	-
2001.....	1	1	1	-	1	-	1	2	-	-	-	-
2000.....	1	1	-	1	1	3	-	2	1	1	1	-
1999.....	-	1	1	-	2	1	1	2	1	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified malnutrition (E40,E42-E46)												
All races, both sexes												
2004.....	69	66	82	111	165	319	510	588	445	218	61	1
2003.....	56	72	83	110	196	393	508	655	509	270	65	-
2002.....	48	69	79	132	211	401	616	727	582	266	80	-
2001.....	53	59	79	117	214	388	586	712	605	276	68	-
2000.....	43	58	92	173	235	460	681	847	664	260	77	-
1999.....	55	62	86	135	247	419	682	845	717	329	84	-
All races, male												
2004.....	32	39	39	55	79	144	185	205	114	41	6	1
2003.....	26	35	37	53	95	157	203	214	121	46	10	-
2002.....	21	37	44	63	108	171	218	257	152	44	12	-
2001.....	23	32	38	55	97	147	229	204	153	48	2	-
2000.....	17	36	45	76	123	185	227	257	152	43	12	-
1999.....	27	35	52	65	128	180	248	266	162	58	8	-
All races, female												
2004.....	37	27	43	56	86	175	325	383	331	177	55	-
2003.....	30	37	46	57	101	236	305	441	388	224	55	-
2002.....	27	32	35	69	103	230	398	470	430	222	68	-
2001.....	30	27	41	62	117	241	357	508	452	228	66	-
2000.....	26	22	47	97	112	275	454	590	512	217	65	-
1999.....	28	27	34	70	119	239	434	579	555	271	76	-
White, both sexes												
2004.....	47	53	62	85	133	259	440	499	388	191	49	-
2003.....	34	55	62	88	160	339	430	575	443	231	48	-
2002.....	37	48	58	102	174	344	533	630	512	231	66	-
2001.....	34	43	66	89	181	320	494	641	531	242	61	-
2000.....	29	43	69	142	181	391	594	745	603	218	64	-
1999.....	45	38	65	103	183	354	574	744	656	280	71	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other and unspecified malnutrition (E40,E42-E46)

White, male

2004.....	23	29	28	37	65	116	157	177	98	36	5	-
2003.....	17	23	26	45	72	129	167	189	104	40	7	-
2002.....	15	23	29	50	89	150	183	226	137	37	9	-
2001.....	12	22	28	38	79	121	192	182	129	42	1	-
2000.....	11	24	30	61	92	156	196	226	140	33	11	-
1999.....	21	18	40	46	89	147	205	233	142	44	6	-

White, female

2004.....	24	24	34	48	68	143	283	322	290	155	44	-
2003.....	17	32	36	43	88	210	263	386	339	191	41	-
2002.....	22	25	29	52	85	194	350	404	375	194	57	-
2001.....	22	21	38	51	102	199	302	459	402	200	60	-
2000.....	18	19	39	81	89	235	398	519	463	185	53	-
1999.....	24	20	25	57	94	207	369	511	514	236	65	-

Black, both sexes

2004.....	20	12	18	23	30	48	55	76	49	23	11	1
2003.....	20	17	19	17	32	49	71	73	60	32	15	-
2002.....	11	21	21	27	32	54	72	83	63	30	12	-
2001.....	17	14	10	25	29	58	77	63	64	26	7	-
2000.....	14	15	21	27	47	60	74	89	56	36	10	-
1999.....	10	20	20	28	55	54	99	87	54	43	13	-

Black, male

2004.....	9	9	10	17	13	23	20	23	14	5	1	1
2003.....	9	12	10	7	21	27	31	22	14	4	3	-
2002.....	6	14	15	12	15	19	30	25	12	3	2	-
2001.....	10	9	8	16	17	24	29	22	20	4	1	-
2000.....	6	12	13	13	28	27	27	29	9	10	-	-
1999.....	6	14	11	17	34	27	37	27	18	12	2	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified malnutrition (E40,E42-E46)												
Black, female												
2004.....	11	3	8	6	17	25	35	53	35	18	10	-
2003.....	11	5	9	10	11	22	40	51	46	28	12	-
2002.....	5	7	6	15	17	35	42	58	51	27	10	-
2001.....	7	5	2	9	12	34	48	41	44	22	6	-
2000.....	8	3	8	14	19	33	47	60	47	26	10	-
1999.....	4	6	9	11	21	27	62	60	36	31	11	-
American Indian 1/, both sexes												
2004.....	1	1	1	2	1	5	5	4	2	-	-	-
2003.....	2	-	-	1	3	4	2	2	2	1	1	-
2002.....	-	-	-	1	3	1	4	4	3	3	-	-
2001.....	2	1	1	2	3	3	6	4	2	1	-	-
2000.....	-	-	-	2	3	3	3	6	2	1	3	-
1999.....	-	3	1	3	3	3	4	4	3	1	-	-
American Indian 1/, male												
2004.....	-	1	1	1	-	2	3	1	-	-	-	-
2003.....	-	-	-	-	2	-	2	-	-	-	-	-
2002.....	-	-	-	-	3	-	1	2	-	2	-	-
2001.....	1	-	1	1	1	-	3	-	-	-	-	-
2000.....	-	-	-	1	1	1	1	-	1	-	1	-
1999.....	-	2	1	1	1	2	3	1	-	-	-	-
American Indian 1/, female												
2004.....	1	-	-	1	1	3	2	3	2	-	-	-
2003.....	2	-	-	1	1	4	-	2	2	1	1	-
2002.....	-	-	-	1	-	1	3	2	3	1	-	-
2001.....	1	1	-	1	2	3	3	4	2	1	-	-
2000.....	-	-	-	1	2	2	2	6	1	1	2	-
1999.....	-	1	-	2	2	1	1	3	3	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified malnutrition (E40,E42-E46)												
Asian or Pacific Islander 2/, both sexes												
2004.....	1	-	1	1	1	7	10	9	6	4	1	-
2003.....	-	-	2	4	1	1	5	5	4	6	1	-
2002.....	-	-	-	2	2	2	7	10	4	2	2	-
2001.....	-	1	2	1	1	7	9	4	8	7	-	-
2000.....	-	-	2	2	4	6	10	7	3	5	-	-
1999.....	-	1	-	1	6	8	5	10	4	5	-	-
Asian or Pacific Islander 2/, male												
2004.....	-	-	-	-	1	3	5	4	2	-	-	-
2003.....	-	-	1	1	-	1	3	3	3	2	-	-
2002.....	-	-	-	1	1	2	4	4	3	2	1	-
2001.....	-	1	1	-	-	2	5	-	4	2	-	-
2000.....	-	-	2	1	2	1	3	2	2	-	-	-
1999.....	-	1	-	1	4	4	3	5	2	2	-	-
Asian or Pacific Islander 2/, female												
2004.....	1	-	1	1	-	4	5	5	4	4	1	-
2003.....	-	-	1	3	1	-	2	2	1	4	1	-
2002.....	-	-	-	1	1	-	3	6	1	-	1	-
2001.....	-	-	1	1	1	5	4	4	4	5	-	-
2000.....	-	-	-	1	2	5	7	5	1	5	-	-
1999.....	-	-	-	-	2	4	2	5	2	3	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other nutritional deficiencies (E50-E64)

All races, both sexes

2004.....	4	4	6	4	6	11	22	31	47	21	2	-
2003.....	3	3	5	7	13	15	25	43	33	17	6	-
2002.....	8	4	8	14	14	33	32	56	51	29	6	-
2001.....	5	2	5	8	19	33	42	42	41	22	12	-
2000.....	7	9	6	14	23	27	59	59	69	22	6	-
1999.....	4	10	6	8	14	33	56	61	67	27	8	-

All races, male

2004.....	2	2	4	2	2	5	10	13	14	5	-	-
2003.....	1	1	3	4	7	9	12	21	9	3	1	-
2002.....	4	4	5	6	6	17	11	16	6	4	-	-
2001.....	4	2	4	6	11	16	10	8	7	5	1	-
2000.....	3	6	5	8	6	13	24	14	17	6	-	-
1999.....	3	4	2	4	10	19	23	19	21	6	1	-

All races, female

2004.....	2	2	2	2	4	6	12	18	33	16	2	-
2003.....	2	2	2	3	6	6	13	22	24	14	5	-
2002.....	4	-	3	8	8	16	21	40	45	25	6	-
2001.....	1	-	1	2	8	17	32	34	34	17	11	-
2000.....	4	3	1	6	17	14	35	45	52	16	6	-
1999.....	1	6	4	4	4	14	33	42	46	21	7	-

White, both sexes

2004.....	4	3	4	2	6	11	20	30	45	20	2	-
2003.....	3	3	4	7	11	14	24	43	29	15	5	-
2002.....	7	3	6	9	9	31	28	51	47	27	5	-
2001.....	2	-	4	5	16	27	36	37	38	20	12	-
2000.....	5	7	4	13	22	20	53	55	62	20	4	-
1999.....	2	5	6	8	12	26	48	54	62	24	8	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other nutritional deficiencies (E50-E64)

White, male

2004.....	2	1	4	1	2	5	9	12	13	5	-	-
2003.....	1	1	2	4	6	8	11	21	8	2	-	-
2002.....	3	3	3	4	4	16	8	15	6	4	-	-
2001.....	1	-	3	3	9	15	9	8	6	4	1	-
2000.....	2	4	3	7	6	9	24	14	15	6	-	-
1999.....	2	1	2	4	8	13	21	18	21	6	1	-

White, female

2004.....	2	2	-	1	4	6	11	18	32	15	2	-
2003.....	2	2	2	3	5	6	13	22	21	13	5	-
2002.....	4	-	3	5	5	15	20	36	41	23	5	-
2001.....	1	-	1	2	7	12	27	29	32	16	11	-
2000.....	3	3	1	6	16	11	29	41	47	14	4	-
1999.....	-	4	4	4	4	13	27	36	41	18	7	-

Black, both sexes

2004.....	-	1	2	2	-	-	2	1	2	-	-	-
2003.....	-	-	1	-	2	1	1	-	4	2	-	-
2002.....	1	1	1	5	5	2	3	4	3	2	1	-
2001.....	3	2	1	2	3	4	6	5	3	2	-	-
2000.....	1	2	2	1	-	5	6	2	4	2	2	-
1999.....	1	5	-	-	2	7	7	7	5	3	-	-

Black, male

2004.....	-	1	-	1	-	-	1	1	1	-	-	-
2003.....	-	-	1	-	1	1	1	-	1	1	-	-
2002.....	1	1	1	2	2	1	3	1	-	-	-	-
2001.....	3	2	1	2	2	1	1	-	1	1	-	-
2000.....	1	2	2	1	-	2	-	-	2	-	-	-
1999.....	1	3	-	-	2	6	2	1	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other nutritional deficiencies (E50-E64)												
Asian or Pacific Islander 2/, both sexes												
2004.....	-	-	-	-	-	-	-	-	-	1	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	1	-
2002.....	-	-	1	-	-	-	-	-	1	-	-	-
2001.....	-	-	-	1	-	2	-	-	-	-	-	-
2000.....	-	-	-	-	-	1	-	-	2	-	-	-
1999.....	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	1	-
2002.....	-	-	1	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	1	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	1	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	-	-	-	-	-	-	-	-	-	1	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	1	-	-	-
2001.....	-	-	-	-	-	2	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	2	-	-	-
1999.....	-	-	-	-	-	-	1	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Obesity and other hyperalimentation (E65-E68)												
All races, both sexes												
2004.....	581	540	448	362	315	233	152	69	21	6	-	-
2003.....	527	511	410	367	281	191	133	66	19	9	-	2
2002.....	504	494	364	300	207	192	117	48	14	2	-	-
2001.....	456	378	304	237	234	165	88	53	12	3	2	-
2000.....	427	323	283	234	213	171	107	49	15	5	3	-
1999.....	333	287	221	213	216	159	100	42	14	3	-	-
All races, male												
2004.....	303	263	192	159	105	83	57	18	3	-	-	-
2003.....	257	261	184	166	95	59	39	9	3	1	-	2
2002.....	256	229	151	119	72	58	30	8	3	1	-	-
2001.....	209	162	132	85	94	40	24	7	1	-	-	-
2000.....	218	167	113	87	74	51	26	8	2	1	-	-
1999.....	158	157	95	84	75	47	19	1	3	-	-	-
All races, female												
2004.....	278	277	256	203	210	150	95	51	18	6	-	-
2003.....	270	250	226	201	186	132	94	57	16	8	-	-
2002.....	248	265	213	181	135	134	87	40	11	1	-	-
2001.....	247	216	172	152	140	125	64	46	11	3	2	-
2000.....	209	156	170	147	139	120	81	41	13	4	3	-
1999.....	175	130	126	129	141	112	81	41	11	3	-	-
White, both sexes												
2004.....	466	451	373	292	266	197	132	59	18	5	-	-
2003.....	439	432	347	330	240	165	118	59	15	7	-	1
2002.....	403	403	301	248	177	154	99	43	14	2	-	-
2001.....	361	319	250	199	185	138	78	49	10	3	2	-
2000.....	349	254	243	194	182	144	93	45	15	3	3	-
1999.....	271	242	181	181	176	135	87	36	14	3	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Obesity and other hyperalimentation (E65-E68)

White, male

2004.....	260	234	168	134	99	72	53	18	3	-	-	-
2003.....	221	235	167	154	84	53	38	8	3	1	-	1
2002.....	226	199	137	105	62	52	27	7	3	1	-	-
2001.....	178	147	111	76	81	36	22	7	1	-	-	-
2000.....	191	142	98	77	67	45	23	7	2	-	-	-
1999.....	139	138	81	76	69	42	18	1	3	-	-	-

White, female

2004.....	206	217	205	158	167	125	79	41	15	5	-	-
2003.....	218	197	180	176	156	112	80	51	12	6	-	-
2002.....	177	204	164	143	115	102	72	36	11	1	-	-
2001.....	183	172	139	123	104	102	56	42	9	3	2	-
2000.....	158	112	145	117	115	99	70	38	13	3	3	-
1999.....	132	104	100	105	107	93	69	35	11	3	-	-

Black, both sexes

2004.....	106	87	72	64	47	34	18	10	3	1	-	-
2003.....	83	71	59	34	40	26	13	4	4	2	-	1
2002.....	97	89	58	49	28	34	16	5	-	-	-	-
2001.....	89	59	51	35	46	27	10	3	2	-	-	-
2000.....	75	66	35	39	29	27	14	4	-	1	-	-
1999.....	59	43	38	31	38	22	12	5	-	-	-	-

Black, male

2004.....	40	28	21	24	5	11	4	-	-	-	-	-
2003.....	34	20	16	11	10	6	1	-	-	-	-	1
2002.....	29	29	12	14	9	6	3	1	-	-	-	-
2001.....	27	15	20	7	11	4	2	-	-	-	-	-
2000.....	26	24	12	10	7	6	3	1	-	1	-	-
1999.....	18	18	13	7	6	4	1	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Obesity and other hyperalimentation (E65-E68)

Black, female

2004.....	66	59	51	40	42	23	14	10	3	1	-	-
2003.....	49	51	43	23	30	20	12	4	4	2	-	-
2002.....	68	60	46	35	19	28	13	4	-	-	-	-
2001.....	62	44	31	28	35	23	8	3	2	-	-	-
2000.....	49	42	23	29	22	21	11	3	-	-	-	-
1999.....	41	25	25	24	32	18	11	5	-	-	-	-

American Indian 1/, both sexes

2004.....	4	2	2	-	1	2	1	-	-	-	-	-
2003.....	1	6	1	2	-	-	2	1	-	-	-	-
2002.....	2	1	3	2	2	1	1	-	-	-	-	-
2001.....	6	-	3	2	2	-	-	1	-	-	-	-
2000.....	1	3	3	-	1	-	-	-	-	-	-	-
1999.....	3	2	-	1	2	1	1	1	-	-	-	-

American Indian 1/, male

2004.....	1	1	2	-	1	-	-	-	-	-	-	-
2003.....	1	4	1	1	-	-	-	-	-	-	-	-
2002.....	-	1	1	-	1	-	-	-	-	-	-	-
2001.....	4	-	1	1	2	-	-	-	-	-	-	-
2000.....	-	1	1	-	-	-	-	-	-	-	-	-
1999.....	1	1	-	1	-	1	-	-	-	-	-	-

American Indian 1/, female

2004.....	3	1	-	-	-	2	1	-	-	-	-	-
2003.....	-	2	-	1	-	-	2	1	-	-	-	-
2002.....	2	-	2	2	1	1	1	-	-	-	-	-
2001.....	2	-	2	1	-	-	-	1	-	-	-	-
2000.....	1	2	2	-	1	-	-	-	-	-	-	-
1999.....	2	1	-	-	2	-	1	1	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Obesity and other hyperalimentation (E65-E68)												
Asian or Pacific Islander 2/, both sexes												
2004.....	5	-	1	6	1	-	1	-	-	-	-	-
2003.....	4	2	3	1	1	-	-	2	-	-	-	-
2002.....	2	1	2	1	-	3	1	-	-	-	-	-
2001.....	-	-	-	1	1	-	-	-	-	-	-	-
2000.....	2	-	2	1	1	-	-	-	-	1	-	-
1999.....	-	-	2	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	2	-	1	1	-	-	-	-	-	-	-	-
2003.....	1	2	-	-	1	-	-	1	-	-	-	-
2002.....	1	-	1	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	1	-	-	-	-	-	-	-	-
2000.....	1	-	2	-	-	-	-	-	-	-	-	-
1999.....	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	3	-	-	5	1	-	1	-	-	-	-	-
2003.....	3	-	3	1	-	-	-	1	-	-	-	-
2002.....	1	1	1	1	-	3	1	-	-	-	-	-
2001.....	-	-	-	-	1	-	-	-	-	-	-	-
2000.....	1	-	-	1	1	-	-	-	-	1	-	-
1999.....	-	-	1	-	-	1	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Metabolic disorders (E70-E88)												
All races, both sexes												
2004.....	489	711	833	1,036	1,427	1,978	2,505	2,465	1,859	759	162	1
2003.....	560	643	803	1,015	1,385	1,932	2,519	2,529	1,786	789	205	-
2002.....	491	690	799	926	1,446	1,969	2,512	2,423	1,805	824	199	3
2001.....	494	577	696	963	1,441	1,956	2,328	2,563	1,836	859	222	-
2000.....	432	540	701	932	1,390	1,904	2,282	2,414	1,789	846	214	1
1999.....	397	461	658	907	1,339	1,861	2,182	2,469	1,880	878	206	-
All races, male												
2004.....	344	462	521	642	786	940	1,070	871	510	152	22	-
2003.....	373	427	504	631	768	929	1,048	864	478	145	28	-
2002.....	341	446	492	554	772	934	1,035	824	422	136	28	3
2001.....	313	358	466	605	778	996	927	830	428	145	23	-
2000.....	310	342	455	562	751	901	892	750	420	164	40	1
1999.....	254	296	394	512	738	878	845	703	429	163	24	-
All races, female												
2004.....	145	249	312	394	641	1,038	1,435	1,594	1,349	607	140	1
2003.....	187	216	299	384	617	1,003	1,471	1,665	1,308	644	177	-
2002.....	150	244	307	372	674	1,035	1,477	1,599	1,383	688	171	-
2001.....	181	219	230	358	663	960	1,401	1,733	1,408	714	199	-
2000.....	122	198	246	370	639	1,003	1,390	1,664	1,369	682	174	-
1999.....	143	165	264	395	601	983	1,337	1,766	1,451	715	182	-
White, both sexes												
2004.....	404	594	705	876	1,233	1,751	2,255	2,275	1,675	689	144	1
2003.....	445	530	690	845	1,206	1,679	2,235	2,311	1,620	712	161	-
2002.....	402	575	661	772	1,221	1,726	2,220	2,185	1,641	730	177	3
2001.....	416	487	576	812	1,264	1,725	2,067	2,310	1,635	784	189	-
2000.....	352	449	584	799	1,213	1,693	2,043	2,180	1,622	764	179	1
1999.....	316	392	554	759	1,168	1,652	1,959	2,249	1,728	791	175	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Metabolic disorders (E70-E88)												
White, male												
2004.....	285	392	456	559	684	838	974	801	470	135	18	-
2003.....	305	364	441	532	675	814	927	803	438	128	24	-
2002.....	286	377	402	468	673	823	922	744	375	116	25	3
2001.....	268	302	386	520	684	887	822	759	376	131	18	-
2000.....	258	291	382	487	667	801	804	695	379	155	34	1
1999.....	213	255	342	433	648	778	752	634	377	141	18	-
White, female												
2004.....	119	202	249	317	549	913	1,281	1,474	1,205	554	126	1
2003.....	140	166	249	313	531	865	1,308	1,508	1,182	584	137	-
2002.....	116	198	259	304	548	903	1,298	1,441	1,266	614	152	-
2001.....	148	185	190	292	580	838	1,245	1,551	1,259	653	171	-
2000.....	94	158	202	312	546	892	1,239	1,485	1,243	609	145	-
1999.....	103	137	212	326	520	874	1,207	1,615	1,351	650	157	-
Black, both sexes												
2004.....	70	101	112	139	171	194	212	165	162	59	15	-
2003.....	101	96	98	150	154	213	233	190	139	62	41	-
2002.....	80	97	124	131	194	202	254	204	146	83	22	-
2001.....	72	72	105	128	160	207	230	211	182	68	32	-
2000.....	68	78	99	120	151	187	216	210	154	74	32	-
1999.....	76	61	93	135	155	183	187	197	136	75	27	-
Black, male												
2004.....	49	60	56	71	88	86	82	59	35	15	4	-
2003.....	59	56	53	84	80	94	101	51	26	14	3	-
2002.....	48	57	83	71	82	89	94	66	40	16	3	-
2001.....	40	44	67	74	85	96	94	57	41	12	5	-
2000.....	44	43	59	64	73	87	78	49	36	9	6	-
1999.....	39	37	47	72	77	86	71	61	42	19	4	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Metabolic disorders (E70-E88)												
Black, female												
2004.....	21	41	56	68	83	108	130	106	127	44	11	-
2003.....	42	40	45	66	74	119	132	139	113	48	38	-
2002.....	32	40	41	60	112	113	160	138	106	67	19	-
2001.....	32	28	38	54	75	111	136	154	141	56	27	-
2000.....	24	35	40	56	78	100	138	161	118	65	26	-
1999.....	37	24	46	63	78	97	116	136	94	56	23	-
American Indian 1/, both sexes												
2004.....	7	1	4	2	7	9	7	6	1	2	-	-
2003.....	5	6	3	6	1	6	7	8	6	4	2	-
2002.....	4	6	5	7	10	10	2	7	3	3	-	-
2001.....	3	6	3	6	4	5	8	7	5	2	-	-
2000.....	2	6	7	3	6	9	1	4	3	2	-	-
1999.....	2	2	2	4	1	2	8	5	1	2	1	-
American Indian 1/, male												
2004.....	3	-	3	1	5	5	3	-	-	-	-	-
2003.....	2	1	1	4	-	4	2	-	1	-	1	-
2002.....	3	4	2	4	5	5	-	3	1	-	-	-
2001.....	3	4	3	3	2	3	1	2	1	2	-	-
2000.....	1	4	5	3	2	4	1	2	1	-	-	-
1999.....	1	1	1	1	1	1	5	2	1	-	-	-
American Indian 1/, female												
2004.....	4	1	1	1	2	4	4	6	1	2	-	-
2003.....	3	5	2	2	1	2	5	8	5	4	1	-
2002.....	1	2	3	3	5	5	2	4	2	3	-	-
2001.....	-	2	-	3	2	2	7	5	4	-	-	-
2000.....	1	2	2	-	4	5	-	2	2	2	-	-
1999.....	1	1	1	3	-	1	3	3	-	2	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Metabolic disorders (E70-E88)												
Asian or Pacific Islander 2/, both sexes												
2004.....	8	15	12	19	16	24	31	19	21	9	3	-
2003.....	9	11	12	14	24	34	44	20	21	11	1	-
2002.....	5	12	9	16	21	31	36	27	15	8	-	-
2001.....	3	12	12	17	13	19	23	35	14	5	1	-
2000.....	10	7	11	10	20	15	22	20	10	6	3	-
1999.....	3	6	9	9	15	24	28	18	15	10	3	-
Asian or Pacific Islander 2/, male												
2004.....	7	10	6	11	9	11	11	11	5	2	-	-
2003.....	7	6	9	11	13	17	18	10	13	3	-	-
2002.....	4	8	5	11	12	17	19	11	6	4	-	-
2001.....	2	8	10	8	7	10	10	12	10	-	-	-
2000.....	7	4	9	8	9	9	9	4	4	-	-	-
1999.....	1	3	4	6	12	13	17	6	9	3	2	-
Asian or Pacific Islander 2/, female												
2004.....	1	5	6	8	7	13	20	8	16	7	3	-
2003.....	2	5	3	3	11	17	26	10	8	8	1	-
2002.....	1	4	4	5	9	14	17	16	9	4	-	-
2001.....	1	4	2	9	6	9	13	23	4	5	1	-
2000.....	3	3	2	2	11	6	13	16	6	6	3	-
1999.....	2	3	5	3	3	11	11	12	6	7	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cystic fibrosis (E84)												
All races, both sexes												
2004.....	13	7	6	4	3	2	3	1	1	-	-	-
2003.....	11	5	4	3	-	4	-	4	2	-	-	-
2002.....	9	6	6	2	4	4	2	1	1	-	-	-
2001.....	11	4	1	3	2	4	5	2	-	-	-	-
2000.....	7	2	2	3	2	-	-	2	1	1	-	-
1999.....	15	5	3	1	2	3	1	1	-	-	-	-
All races, male												
2004.....	5	4	3	2	3	1	-	1	-	-	-	-
2003.....	6	3	2	2	-	1	-	1	-	-	-	-
2002.....	4	6	3	1	3	1	2	-	-	-	-	-
2001.....	3	2	-	1	1	1	3	-	-	-	-	-
2000.....	4	2	2	2	1	-	-	-	-	-	-	-
1999.....	6	2	1	-	2	2	1	1	-	-	-	-
All races, female												
2004.....	8	3	3	2	-	1	3	-	1	-	-	-
2003.....	5	2	2	1	-	3	-	3	2	-	-	-
2002.....	5	-	3	1	1	3	-	1	1	-	-	-
2001.....	8	2	1	2	1	3	2	2	-	-	-	-
2000.....	3	-	-	1	1	-	-	2	1	1	-	-
1999.....	9	3	2	1	-	1	-	-	-	-	-	-
White, both sexes												
2004.....	13	6	6	4	3	2	3	1	1	-	-	-
2003.....	10	4	4	3	-	3	-	4	2	-	-	-
2002.....	8	5	4	2	3	4	2	1	1	-	-	-
2001.....	10	4	1	3	1	4	5	2	-	-	-	-
2000.....	5	1	2	3	1	-	-	2	1	1	-	-
1999.....	13	5	3	1	2	3	1	1	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Volume depletion, disorders of fluid, electrolyte and acid-base balance (E86-E87)												
All races, both sexes												
2004.....	107	143	158	212	380	674	1,098	1,381	1,299	616	142	1
2003.....	139	124	180	224	393	735	1,213	1,536	1,283	648	178	-
2002.....	106	146	185	235	454	729	1,310	1,564	1,377	703	176	3
2001.....	114	113	186	252	491	818	1,291	1,733	1,445	762	197	-
2000.....	99	141	183	282	460	840	1,315	1,706	1,463	743	191	1
1999.....	112	111	152	282	491	877	1,350	1,841	1,603	783	194	-
All races, male												
2004.....	68	84	76	113	183	278	416	417	340	125	20	-
2003.....	69	72	87	111	176	317	456	479	325	121	23	-
2002.....	54	69	96	116	201	295	484	498	308	115	22	3
2001.....	57	53	106	125	212	374	464	516	321	135	22	-
2000.....	57	78	97	149	226	366	500	498	349	150	37	1
1999.....	68	67	69	135	242	398	490	498	349	144	23	-
All races, female												
2004.....	39	59	82	99	197	396	682	964	959	491	122	1
2003.....	70	52	93	113	217	418	757	1,057	958	527	155	-
2002.....	52	77	89	119	253	434	826	1,066	1,069	588	154	-
2001.....	57	60	80	127	279	444	827	1,217	1,124	627	175	-
2000.....	42	63	86	133	234	474	815	1,208	1,114	593	154	-
1999.....	44	44	83	147	249	479	860	1,343	1,254	639	171	-
White, both sexes												
2004.....	87	100	117	164	303	583	970	1,261	1,159	554	126	1
2003.....	92	90	144	164	327	609	1,057	1,384	1,157	580	139	-
2002.....	75	112	142	179	361	607	1,128	1,395	1,242	627	157	3
2001.....	80	82	137	195	396	692	1,116	1,557	1,287	693	169	-
2000.....	69	107	141	220	374	727	1,161	1,531	1,319	672	162	1
1999.....	77	84	106	205	406	751	1,207	1,669	1,472	709	166	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Volume depletion, disorders of fluid, electrolyte and acid-base balance (E86-E87)

White, male

2004.....	55	54	52	91	149	239	370	369	308	110	16	-
2003.....	41	57	71	81	146	262	388	436	295	106	20	-
2002.....	36	49	71	84	163	245	414	444	268	99	20	3
2001.....	39	37	75	101	163	317	391	470	284	121	17	-
2000.....	41	57	72	117	184	307	440	457	311	142	31	1
1999.....	49	49	50	89	194	334	428	443	301	127	18	-

White, female

2004.....	32	46	65	73	154	344	600	892	851	444	110	1
2003.....	51	33	73	83	181	347	669	948	862	474	119	-
2002.....	39	63	71	95	198	362	714	951	974	528	137	-
2001.....	41	45	62	94	233	375	725	1,087	1,003	572	152	-
2000.....	28	50	69	103	190	420	721	1,074	1,008	530	131	-
1999.....	28	35	56	116	212	417	779	1,226	1,171	582	148	-

Black, both sexes

2004.....	16	40	37	44	71	82	113	106	128	51	14	-
2003.....	45	31	35	58	62	119	138	132	110	55	36	-
2002.....	28	30	36	51	85	110	164	146	120	66	19	-
2001.....	33	26	47	51	90	119	158	152	148	62	27	-
2000.....	29	29	37	60	76	104	145	160	134	63	26	-
1999.....	33	26	43	71	81	111	127	154	117	65	25	-

Black, male

2004.....	12	27	22	22	32	36	44	42	28	13	4	-
2003.....	27	14	15	30	29	51	65	37	23	12	2	-
2002.....	16	17	22	28	34	45	61	44	35	12	2	-
2001.....	17	14	29	23	47	52	68	37	32	12	5	-
2000.....	16	18	22	31	36	55	56	37	34	8	6	-
1999.....	18	18	19	44	44	56	51	49	39	16	3	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Volume depletion, disorders of fluid, electrolyte and acid-base balance (E86-E87)

Black, female

2004.....	4	13	15	22	39	46	69	64	100	38	10	-
2003.....	18	17	20	28	33	68	73	95	87	43	34	-
2002.....	12	13	14	23	51	65	103	102	85	54	17	-
2001.....	16	12	18	28	43	67	90	115	116	50	22	-
2000.....	13	11	15	29	40	49	89	123	100	55	20	-
1999.....	15	8	24	27	37	55	76	105	78	49	22	-

American Indian 1/, both sexes

2004.....	3	-	1	1	3	4	5	4	1	2	-	-
2003.....	2	2	-	1	-	2	4	8	5	3	2	-
2002.....	2	3	2	2	3	4	-	7	3	2	-	-
2001.....	1	3	1	2	4	2	5	6	2	2	-	-
2000.....	1	3	3	-	3	4	-	4	3	2	-	-
1999.....	-	-	1	2	-	1	4	5	1	2	1	-

American Indian 1/, male

2004.....	-	-	1	-	2	1	1	-	-	-	-	-
2003.....	1	-	-	-	-	1	-	-	1	-	1	-
2002.....	2	2	-	1	-	1	-	3	1	-	-	-
2001.....	1	2	1	-	2	1	-	2	-	2	-	-
2000.....	-	1	1	-	1	2	-	2	1	-	-	-
1999.....	-	-	-	-	-	1	2	2	1	-	-	-

American Indian 1/, female

2004.....	3	-	-	1	1	3	4	4	1	2	-	-
2003.....	1	2	-	1	-	1	4	8	4	3	1	-
2002.....	-	1	2	1	3	3	-	4	2	2	-	-
2001.....	-	1	-	2	2	1	5	4	2	-	-	-
2000.....	1	2	2	-	2	2	-	2	2	2	-	-
1999.....	-	-	1	2	-	-	2	3	-	2	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Volume depletion, disorders of fluid, electrolyte and acid-base balance (E86-E87)

Asian or Pacific Islander 2/, both sexes

2004.....	1	3	3	3	3	5	10	10	11	9	2	-
2003.....	-	1	1	1	4	5	14	12	11	10	1	-
2002.....	1	1	5	3	5	8	18	16	12	8	-	-
2001.....	-	2	1	4	1	5	12	18	8	5	1	-
2000.....	-	2	2	2	7	5	9	11	7	6	3	-
1999.....	2	1	2	4	4	14	12	13	13	7	2	-

Asian or Pacific Islander 2/, male

2004.....	1	3	1	-	-	2	1	6	4	2	-	-
2003.....	-	1	1	-	1	3	3	6	6	3	-	-
2002.....	-	1	3	3	4	4	9	7	4	4	-	-
2001.....	-	-	1	1	-	4	5	7	5	-	-	-
2000.....	-	2	2	1	5	2	4	2	3	-	-	-
1999.....	1	-	-	2	4	7	9	4	8	1	2	-

Asian or Pacific Islander 2/, female

2004.....	-	-	2	3	3	3	9	4	7	7	2	-
2003.....	-	-	-	1	3	2	11	6	5	7	1	-
2002.....	1	-	2	-	1	4	9	9	8	4	-	-
2001.....	-	2	-	3	1	1	7	11	3	5	1	-
2000.....	-	-	-	1	2	3	5	9	4	6	3	-
1999.....	1	1	2	2	-	7	3	9	5	6	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other metabolic disorders (E70-E83,E85,E88)												
All races, both sexes												
2004.....	369	561	669	820	1,044	1,302	1,404	1,083	559	143	20	-
2003.....	410	514	619	788	992	1,193	1,306	989	501	141	27	-
2002.....	376	538	608	689	988	1,236	1,200	858	427	121	23	-
2001.....	369	460	509	708	948	1,134	1,032	828	391	97	25	-
2000.....	326	397	516	647	928	1,064	967	706	325	102	23	-
1999.....	270	345	503	624	846	981	831	627	277	95	12	-
All races, male												
2004.....	271	374	442	527	600	661	654	453	170	27	2	-
2003.....	298	352	415	518	592	611	592	384	153	24	5	-
2002.....	283	371	393	437	568	638	549	326	114	21	6	-
2001.....	253	303	360	479	565	621	460	314	107	10	1	-
2000.....	249	262	356	411	524	535	392	252	71	14	3	-
1999.....	180	227	324	377	494	478	354	204	80	19	1	-
All races, female												
2004.....	98	187	227	293	444	641	750	630	389	116	18	-
2003.....	112	162	204	270	400	582	714	605	348	117	22	-
2002.....	93	167	215	252	420	598	651	532	313	100	17	-
2001.....	116	157	149	229	383	513	572	514	284	87	24	-
2000.....	77	135	160	236	404	529	575	454	254	88	20	-
1999.....	90	118	179	247	352	503	477	423	197	76	11	-
White, both sexes												
2004.....	304	488	582	708	927	1,166	1,282	1,013	515	135	18	-
2003.....	343	436	542	678	879	1,067	1,178	923	461	132	22	-
2002.....	319	458	515	591	857	1,115	1,090	789	398	103	20	-
2001.....	326	401	438	614	867	1,029	946	751	348	91	20	-
2000.....	278	341	441	576	838	966	882	647	302	91	17	-
1999.....	226	303	445	553	760	898	751	579	256	82	9	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other metabolic disorders (E70-E83,E85,E88)												
White, male												
2004.....	225	335	401	466	532	598	604	431	162	25	2	-
2003.....	258	305	368	449	529	551	539	366	143	22	4	-
2002.....	246	323	328	383	508	577	506	300	107	17	5	-
2001.....	226	263	311	418	520	569	428	289	92	10	1	-
2000.....	214	233	308	368	482	494	364	238	68	13	3	-
1999.....	159	204	291	344	452	442	323	190	76	14	-	-
White, female												
2004.....	79	153	181	242	395	568	678	582	353	110	16	-
2003.....	85	131	174	229	350	516	639	557	318	110	18	-
2002.....	73	135	187	208	349	538	584	489	291	86	15	-
2001.....	100	138	127	196	347	460	518	462	256	81	19	-
2000.....	64	108	133	208	356	472	518	409	234	78	14	-
1999.....	67	99	154	209	308	456	428	389	180	68	9	-
Black, both sexes												
2004.....	54	61	75	95	100	112	99	59	34	8	1	-
2003.....	55	64	63	92	92	93	95	58	29	7	5	-
2002.....	51	66	87	80	108	92	90	58	26	17	3	-
2001.....	39	46	58	77	70	88	72	59	34	6	5	-
2000.....	37	48	62	60	75	83	71	50	20	11	6	-
1999.....	41	35	50	64	74	72	60	43	19	10	2	-
Black, male												
2004.....	37	33	34	49	56	50	38	17	7	2	-	-
2003.....	32	41	38	54	51	43	36	14	3	2	1	-
2002.....	32	39	61	43	47	44	33	22	5	4	1	-
2001.....	23	30	38	51	38	44	26	20	9	-	-	-
2000.....	27	24	37	33	37	32	22	12	2	1	-	-
1999.....	20	19	28	28	33	30	20	12	3	3	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other metabolic disorders (E70-E83,E85,E88)

Black, female

2004.....	17	28	41	46	44	62	61	42	27	6	1	-
2003.....	23	23	25	38	41	50	59	44	26	5	4	-
2002.....	19	27	26	37	61	48	57	36	21	13	2	-
2001.....	16	16	20	26	32	44	46	39	25	6	5	-
2000.....	10	24	25	27	38	51	49	38	18	10	6	-
1999.....	21	16	22	36	41	42	40	31	16	7	1	-

American Indian 1/, both sexes

2004.....	4	1	3	1	4	5	2	2	-	-	-	-
2003.....	3	4	3	5	1	4	3	-	1	1	-	-
2002.....	2	3	3	5	7	6	2	-	-	1	-	-
2001.....	2	3	2	4	-	3	3	1	3	-	-	-
2000.....	1	3	4	3	3	5	1	-	-	-	-	-
1999.....	2	2	1	2	1	1	4	-	-	-	-	-

American Indian 1/, male

2004.....	3	-	2	1	3	4	2	-	-	-	-	-
2003.....	1	1	1	4	-	3	2	-	-	-	-	-
2002.....	1	2	2	3	5	4	-	-	-	-	-	-
2001.....	2	2	2	3	-	2	1	-	1	-	-	-
2000.....	1	3	4	3	1	2	1	-	-	-	-	-
1999.....	1	1	1	1	1	-	3	-	-	-	-	-

American Indian 1/, female

2004.....	1	1	1	-	1	1	-	2	-	-	-	-
2003.....	2	3	2	1	1	1	1	-	1	1	-	-
2002.....	1	1	1	2	2	2	2	-	-	1	-	-
2001.....	-	1	-	1	-	1	2	1	2	-	-	-
2000.....	-	-	-	-	2	3	-	-	-	-	-	-
1999.....	1	1	-	1	-	1	1	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other metabolic disorders (E70-E83,E85,E88)												
Asian or Pacific Islander 2/, both sexes												
2004.....	7	11	9	16	13	19	21	9	10	-	1	-
2003.....	9	10	11	13	20	29	30	8	10	1	-	-
2002.....	4	11	3	13	16	23	18	11	3	-	-	-
2001.....	2	10	11	13	11	14	11	17	6	-	-	-
2000.....	10	5	9	8	12	10	13	9	3	-	-	-
1999.....	1	5	7	5	11	10	16	5	2	3	1	-
Asian or Pacific Islander 2/, male												
2004.....	6	6	5	11	9	9	10	5	1	-	-	-
2003.....	7	5	8	11	12	14	15	4	7	-	-	-
2002.....	4	7	2	8	8	13	10	4	2	-	-	-
2001.....	2	8	9	7	7	6	5	5	5	-	-	-
2000.....	7	2	7	7	4	7	5	2	1	-	-	-
1999.....	-	3	4	4	8	6	8	2	1	2	-	-
Asian or Pacific Islander 2/, female												
2004.....	1	5	4	5	4	10	11	4	9	-	1	-
2003.....	2	5	3	2	8	15	15	4	3	1	-	-
2002.....	-	4	1	5	8	10	8	7	1	-	-	-
2001.....	-	2	2	6	4	8	6	12	1	-	-	-
2000.....	3	3	2	1	8	3	8	7	2	-	-	-
1999.....	1	2	3	1	3	4	8	3	1	1	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
V. Mental and behavioral disorders (F01-F99)												
All races, both sexes												
2004.....	1,852	1,525	1,319	1,376	2,431	5,157	10,031	13,568	12,048	5,752	1,259	8
2003.....	1,704	1,313	1,258	1,329	2,341	5,202	9,708	13,301	11,755	5,617	1,347	12
2002.....	1,553	1,248	1,106	1,248	2,225	4,761	8,937	12,267	11,115	5,159	1,164	5
2001.....	1,419	1,088	1,040	1,242	2,090	4,503	7,944	11,070	9,796	4,755	1,014	12
2000.....	1,257	1,051	925	1,118	1,997	4,012	7,106	9,868	8,791	4,173	977	8
1999.....	1,157	960	950	1,070	1,878	3,665	6,278	8,954	7,883	3,700	868	6
All races, male												
2004.....	1,421	1,158	949	866	1,292	2,268	3,698	4,099	2,812	938	138	8
2003.....	1,308	981	863	838	1,227	2,267	3,611	3,919	2,629	894	126	9
2002.....	1,235	937	794	818	1,167	2,094	3,338	3,657	2,458	776	141	5
2001.....	1,106	801	756	799	1,138	1,934	2,920	3,318	2,150	769	102	11
2000.....	982	776	666	725	1,092	1,807	2,578	2,917	1,876	643	98	8
1999.....	918	739	699	709	1,046	1,712	2,404	2,614	1,714	541	93	4
All races, female												
2004.....	431	367	370	510	1,139	2,889	6,333	9,469	9,236	4,814	1,121	-
2003.....	396	332	395	491	1,114	2,935	6,097	9,382	9,126	4,723	1,221	3
2002.....	318	311	312	430	1,058	2,667	5,599	8,610	8,657	4,383	1,023	-
2001.....	313	287	284	443	952	2,569	5,024	7,752	7,646	3,986	912	1
2000.....	275	275	259	393	905	2,205	4,528	6,951	6,915	3,530	879	-
1999.....	239	221	251	361	832	1,953	3,874	6,340	6,169	3,159	775	2
White, both sexes												
2004.....	1,455	1,244	1,083	1,111	2,079	4,533	9,157	12,574	11,130	5,327	1,104	4
2003.....	1,312	1,054	1,014	1,128	2,037	4,621	8,811	12,367	10,946	5,181	1,194	10
2002.....	1,202	988	875	1,034	1,930	4,241	8,173	11,386	10,400	4,757	1,016	3
2001.....	1,058	841	847	1,027	1,818	4,054	7,325	10,272	9,158	4,440	902	9
2000.....	977	831	728	926	1,724	3,595	6,508	9,188	8,219	3,891	868	6
1999.....	872	728	765	884	1,630	3,299	5,779	8,394	7,486	3,468	784	5

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
V. Mental and behavioral disorders (F01-F99)												
White, male												
2004.....	1,112	940	761	682	1,102	1,961	3,363	3,797	2,577	846	116	4
2003.....	995	791	668	709	1,049	1,997	3,261	3,640	2,455	806	101	7
2002.....	959	728	620	653	993	1,835	3,050	3,385	2,273	701	116	3
2001.....	829	602	603	647	980	1,736	2,679	3,076	1,999	702	85	8
2000.....	764	609	517	585	930	1,619	2,344	2,684	1,727	593	89	6
1999.....	699	551	562	577	897	1,519	2,187	2,432	1,617	499	84	3
White, female												
2004.....	343	304	322	429	977	2,572	5,794	8,777	8,553	4,481	988	-
2003.....	317	263	346	419	988	2,624	5,550	8,727	8,491	4,375	1,093	3
2002.....	243	260	255	381	937	2,406	5,123	8,001	8,127	4,056	900	-
2001.....	229	239	244	380	838	2,318	4,646	7,196	7,159	3,738	817	1
2000.....	213	222	211	341	794	1,976	4,164	6,504	6,492	3,298	779	-
1999.....	173	177	203	307	733	1,780	3,592	5,962	5,869	2,969	700	2
Black, both sexes												
2004.....	346	246	203	232	313	564	761	840	762	359	129	4
2003.....	336	227	214	172	262	510	780	792	679	376	137	2
2002.....	314	237	193	186	257	456	662	739	607	346	129	1
2001.....	325	218	178	192	249	406	546	684	532	276	99	3
2000.....	257	192	177	169	250	385	515	596	474	247	98	2
1999.....	246	217	165	170	228	324	444	470	350	203	72	1
Black, male												
2004.....	263	197	162	163	169	277	284	249	182	69	14	4
2003.....	268	165	171	111	157	233	293	224	143	74	22	2
2002.....	248	188	146	142	153	225	248	206	145	58	18	1
2001.....	252	175	143	136	142	182	214	194	115	51	13	3
2000.....	203	148	134	125	146	174	201	192	115	46	7	2
1999.....	188	174	125	121	137	168	193	151	77	31	5	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
V. Mental and behavioral disorders (F01-F99)												
Black, female												
2004.....	83	49	41	69	144	287	477	591	580	290	115	-
2003.....	68	62	43	61	105	277	487	568	536	302	115	-
2002.....	66	49	47	44	104	231	414	533	462	288	111	-
2001.....	73	43	35	56	107	224	332	490	417	225	86	-
2000.....	54	44	43	44	104	211	314	404	359	201	91	-
1999.....	58	43	40	49	91	156	251	319	273	172	67	-
American Indian 1/, both sexes												
2004.....	35	27	22	21	14	17	19	29	23	8	6	-
2003.....	44	26	18	19	15	17	21	26	19	14	3	-
2002.....	24	16	27	14	15	22	20	37	17	6	3	-
2001.....	26	18	12	16	10	12	16	34	18	4	5	-
2000.....	19	18	11	13	12	8	20	16	18	5	-	-
1999.....	30	14	15	8	11	17	15	20	11	7	4	-
American Indian 1/, male												
2004.....	30	15	19	15	10	8	10	9	9	2	3	-
2003.....	34	21	14	14	9	13	8	10	2	1	-	-
2002.....	18	14	20	11	10	10	9	17	4	3	1	-
2001.....	19	15	8	14	7	4	6	13	4	1	1	-
2000.....	11	12	9	10	10	3	7	9	6	1	-	-
1999.....	24	13	10	5	7	10	6	7	7	1	-	-
American Indian 1/, female												
2004.....	5	12	3	6	4	9	9	20	14	6	3	-
2003.....	10	5	4	5	6	4	13	16	17	13	3	-
2002.....	6	2	7	3	5	12	11	20	13	3	2	-
2001.....	7	3	4	2	3	8	10	21	14	3	4	-
2000.....	8	6	2	3	2	5	13	7	12	4	-	-
1999.....	6	1	5	3	4	7	9	13	4	6	4	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

V. Mental and behavioral disorders (F01-F99)

Asian or Pacific Islander 2/, both sexes

2004.....	16	8	11	12	25	43	94	125	133	58	20	-
2003.....	12	6	12	10	27	54	96	116	111	46	13	-
2002.....	13	7	11	14	23	42	82	105	91	50	16	1
2001.....	10	11	3	7	13	31	57	80	88	35	8	-
2000.....	4	10	9	10	11	24	63	68	80	30	11	-
1999.....	9	1	5	8	9	25	40	70	36	22	8	-

Asian or Pacific Islander 2/, male

2004.....	16	6	7	6	11	22	41	44	44	21	5	-
2003.....	11	4	10	4	12	24	49	45	29	13	3	-
2002.....	10	7	8	12	11	24	31	49	36	14	6	1
2001.....	6	9	2	2	9	12	21	35	32	15	3	-
2000.....	4	7	6	5	6	11	26	32	28	3	2	-
1999.....	7	1	2	6	5	15	18	24	13	10	4	-

Asian or Pacific Islander 2/, female

2004.....	-	2	4	6	14	21	53	81	89	37	15	-
2003.....	1	2	2	6	15	30	47	71	82	33	10	-
2002.....	3	-	3	2	12	18	51	56	55	36	10	-
2001.....	4	2	1	5	4	19	36	45	56	20	5	-
2000.....	-	3	3	5	5	13	37	36	52	27	9	-
1999.....	2	-	3	2	4	10	22	46	23	12	4	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Organic dementia (F01,F03)												
All races, both sexes												
2004.....	43	117	279	571	1,673	4,334	9,087	12,569	11,316	5,441	1,201	-
2003.....	41	94	272	579	1,592	4,365	8,810	12,286	10,948	5,279	1,261	1
2002.....	32	79	204	513	1,481	3,993	8,021	11,291	10,342	4,775	1,073	-
2001.....	32	81	198	526	1,393	3,679	6,992	10,066	9,022	4,383	928	-
2000.....	23	80	156	433	1,272	3,191	6,199	8,827	7,922	3,810	878	-
1999.....	28	59	149	378	1,155	2,772	5,278	7,843	6,974	3,288	769	-
All races, male												
2004.....	24	66	149	289	777	1,809	3,266	3,748	2,588	891	128	-
2003.....	21	52	140	305	769	1,796	3,201	3,571	2,427	828	116	-
2002.....	15	34	104	274	722	1,692	2,954	3,355	2,272	711	134	-
2001.....	16	36	108	273	682	1,520	2,524	2,999	1,960	700	88	-
2000.....	15	34	89	229	637	1,383	2,211	2,557	1,675	571	92	-
1999.....	18	41	86	216	594	1,219	1,978	2,263	1,488	479	86	-
All races, female												
2004.....	19	51	130	282	896	2,525	5,821	8,821	8,728	4,550	1,073	-
2003.....	20	42	132	274	823	2,569	5,609	8,715	8,521	4,451	1,145	1
2002.....	17	45	100	239	759	2,301	5,067	7,936	8,070	4,064	939	-
2001.....	16	45	90	253	711	2,159	4,468	7,067	7,062	3,683	840	-
2000.....	8	46	67	204	635	1,808	3,988	6,270	6,247	3,239	786	-
1999.....	10	18	63	162	561	1,553	3,300	5,580	5,486	2,809	683	-
White, both sexes												
2004.....	37	100	240	470	1,424	3,799	8,276	11,632	10,441	5,035	1,049	-
2003.....	32	80	231	487	1,380	3,863	7,986	11,410	10,183	4,867	1,120	1
2002.....	29	70	176	432	1,293	3,547	7,333	10,488	9,662	4,405	932	-
2001.....	27	67	170	444	1,209	3,313	6,438	9,347	8,416	4,091	825	-
2000.....	20	68	124	365	1,087	2,854	5,671	8,210	7,403	3,547	780	-
1999.....	26	51	128	318	1,012	2,501	4,857	7,355	6,620	3,077	694	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Organic dementia (F01,F03)												
White, male												
2004.....	21	56	127	236	666	1,560	2,953	3,472	2,375	804	107	-
2003.....	18	45	114	259	654	1,574	2,886	3,314	2,266	745	92	-
2002.....	12	32	90	223	625	1,480	2,699	3,111	2,093	644	109	-
2001.....	11	29	90	227	587	1,371	2,314	2,784	1,819	641	74	-
2000.....	13	27	72	191	538	1,241	2,009	2,349	1,540	529	84	-
1999.....	16	35	75	185	519	1,095	1,800	2,111	1,403	441	78	-
White, female												
2004.....	16	44	113	234	758	2,239	5,323	8,160	8,066	4,231	942	-
2003.....	14	35	117	228	726	2,289	5,100	8,096	7,917	4,122	1,028	1
2002.....	17	38	86	209	668	2,067	4,634	7,377	7,569	3,761	823	-
2001.....	16	38	80	217	622	1,942	4,124	6,563	6,597	3,450	751	-
2000.....	7	41	52	174	549	1,613	3,662	5,861	5,863	3,018	696	-
1999.....	10	16	53	133	493	1,406	3,057	5,244	5,217	2,636	616	-
Black, both sexes												
2004.....	5	16	35	93	224	492	708	791	726	342	128	-
2003.....	7	13	38	79	184	441	717	745	641	355	128	-
2002.....	2	9	25	73	167	389	601	677	580	315	122	-
2001.....	5	14	25	76	171	331	490	619	506	258	90	-
2000.....	3	11	27	58	169	311	452	544	429	228	88	-
1999.....	2	8	18	54	131	240	372	410	312	183	65	-
Black, male												
2004.....	2	9	21	52	101	229	268	228	165	65	14	-
2003.....	2	7	24	39	102	193	264	206	131	69	21	-
2002.....	2	2	12	44	86	183	222	186	142	51	18	-
2001.....	5	7	15	44	87	135	188	172	108	47	10	-
2000.....	2	6	15	35	88	130	173	175	104	38	6	-
1999.....	2	6	10	29	68	108	156	129	66	27	5	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Organic dementia (F01,F03)												
Black, female												
2004.....	3	7	14	41	123	263	440	563	561	277	114	-
2003.....	5	6	14	40	82	248	453	539	510	286	107	-
2002.....	-	7	13	29	81	206	379	491	438	264	104	-
2001.....	-	7	10	32	84	196	302	447	398	211	80	-
2000.....	1	5	12	23	81	181	279	369	325	190	82	-
1999.....	-	2	8	25	63	132	216	281	246	156	60	-
American Indian 1/, both sexes												
2004.....	-	-	-	2	5	11	17	28	22	8	6	-
2003.....	1	-	-	6	5	9	17	23	16	13	3	-
2002.....	-	-	-	3	6	20	16	31	15	6	3	-
2001.....	-	-	2	1	3	7	12	26	17	3	5	-
2000.....	-	-	-	3	7	8	16	12	16	5	-	-
1999.....	-	-	2	2	8	10	11	17	10	6	3	-
American Indian 1/, male												
2004.....	-	-	-	1	2	6	8	9	8	2	3	-
2003.....	-	-	-	5	3	6	5	8	2	1	-	-
2002.....	-	-	-	3	3	9	7	15	4	3	1	-
2001.....	-	-	2	1	1	4	3	11	3	-	1	-
2000.....	-	-	-	1	6	3	5	6	5	1	-	-
1999.....	-	-	1	-	5	4	5	4	6	1	-	-
American Indian 1/, female												
2004.....	-	-	-	1	3	5	9	19	14	6	3	-
2003.....	1	-	-	1	2	3	12	15	14	12	3	-
2002.....	-	-	-	-	3	11	9	16	11	3	2	-
2001.....	-	-	-	-	2	3	9	15	14	3	4	-
2000.....	-	-	-	2	1	5	11	6	11	4	-	-
1999.....	-	-	1	2	3	6	6	13	4	5	3	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Organic dementia (F01,F03)												
Asian or Pacific Islander 2/, both sexes												
2004.....	1	1	4	6	20	32	86	118	127	56	18	-
2003.....	1	1	3	7	23	52	90	108	108	44	10	-
2002.....	1	-	3	5	15	37	71	95	85	49	16	-
2001.....	-	-	1	5	10	28	52	74	83	31	8	-
2000.....	-	1	5	7	9	18	60	61	74	30	10	-
1999.....	-	-	1	4	4	21	38	61	32	22	7	-
Asian or Pacific Islander 2/, male												
2004.....	1	1	1	-	8	14	37	39	40	20	4	-
2003.....	1	-	2	2	10	23	46	43	28	13	3	-
2002.....	1	-	2	4	8	20	26	43	33	13	6	-
2001.....	-	-	1	1	7	10	19	32	30	12	3	-
2000.....	-	1	2	2	5	9	24	27	26	3	2	-
1999.....	-	-	-	2	2	12	17	19	13	10	3	-
Asian or Pacific Islander 2/, female												
2004.....	-	-	3	6	12	18	49	79	87	36	14	-
2003.....	-	1	1	5	13	29	44	65	80	31	7	-
2002.....	-	-	1	1	7	17	45	52	52	36	10	-
2001.....	-	-	-	4	3	18	33	42	53	19	5	-
2000.....	-	-	3	5	4	9	36	34	48	27	8	-
1999.....	-	-	1	2	2	9	21	42	19	12	4	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified organic mental disorders (F04-F09)												
All races, both sexes												
2004.....	13	14	28	38	66	163	331	485	394	174	30	-
2003.....	12	13	28	32	83	197	323	450	450	196	59	-
2002.....	6	21	23	44	102	178	348	481	455	236	56	-
2001.....	9	16	22	52	100	217	403	526	472	246	47	-
2000.....	9	22	28	44	109	239	373	595	516	251	65	-
1999.....	10	13	24	39	105	240	427	627	592	277	71	1
All races, male												
2004.....	7	10	14	21	38	83	135	156	120	16	1	-
2003.....	9	7	18	24	37	97	138	160	104	35	1	-
2002.....	4	10	14	30	51	74	138	144	96	34	5	-
2001.....	6	12	13	33	51	91	148	151	117	37	4	-
2000.....	4	14	14	26	55	102	118	189	108	47	4	-
1999.....	7	8	16	23	51	107	163	177	137	36	6	1
All races, female												
2004.....	6	4	14	17	28	80	196	329	274	158	29	-
2003.....	3	6	10	8	46	100	185	290	346	161	58	-
2002.....	2	11	9	14	51	104	210	337	359	202	51	-
2001.....	3	4	9	19	49	126	255	375	355	209	43	-
2000.....	5	8	14	18	54	137	255	406	408	204	61	-
1999.....	3	5	8	16	54	133	264	450	455	241	65	-
White, both sexes												
2004.....	11	12	22	30	63	149	306	459	370	162	29	-
2003.....	11	9	22	23	74	177	298	421	417	180	51	-
2002.....	4	18	17	37	86	164	323	441	439	211	52	-
2001.....	5	14	19	43	88	192	374	480	451	229	42	-
2000.....	8	20	22	40	98	220	338	553	487	237	56	-
1999.....	9	10	22	30	90	214	389	582	557	263	63	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified organic mental disorders (F04-F09)												
White, male												
2004.....	5	9	11	16	36	77	128	148	110	14	1	-
2003.....	8	4	16	16	31	87	126	152	94	31	1	-
2002.....	2	9	11	25	44	69	130	130	96	28	5	-
2001.....	2	11	11	29	45	80	137	138	111	33	3	-
2000.....	3	13	10	23	48	96	107	171	100	41	4	-
1999.....	6	8	14	18	45	91	146	159	128	33	5	-
White, female												
2004.....	6	3	11	14	27	72	178	311	260	148	28	-
2003.....	3	5	6	7	43	90	172	269	323	149	50	-
2002.....	2	9	6	12	42	95	193	311	343	183	47	-
2001.....	3	3	8	14	43	112	237	342	340	196	39	-
2000.....	5	7	12	17	50	124	231	382	387	196	52	-
1999.....	3	2	8	12	45	123	243	423	429	230	58	-
Black, both sexes												
2004.....	1	2	5	8	3	11	22	25	22	11	1	-
2003.....	1	4	6	7	9	19	23	20	29	14	7	-
2002.....	1	2	6	5	14	11	21	35	13	25	4	-
2001.....	4	2	3	9	11	24	27	41	18	15	5	-
2000.....	1	2	6	3	10	18	34	35	23	14	9	-
1999.....	1	3	2	9	13	24	36	39	32	13	6	1
Black, male												
2004.....	1	1	2	5	2	6	5	8	8	1	-	-
2003.....	1	3	2	7	6	9	10	5	10	4	-	-
2002.....	1	-	3	3	7	4	6	10	-	6	-	-
2001.....	4	1	2	4	5	11	10	12	5	2	1	-
2000.....	1	1	4	2	6	6	11	11	5	6	-	-
1999.....	1	-	2	5	6	15	16	14	8	3	-	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified organic mental disorders (F04-F09)												
Black, female												
2004.....	-	1	3	3	1	5	17	17	14	10	1	-
2003.....	-	1	4	-	3	10	13	15	19	10	7	-
2002.....	-	2	3	2	7	7	15	25	13	19	4	-
2001.....	-	1	1	5	6	13	17	29	13	13	4	-
2000.....	-	1	2	1	4	12	23	24	18	8	9	-
1999.....	-	3	-	4	7	9	20	25	24	10	6	-
American Indian 1/, both sexes												
2004.....	1	-	-	-	-	3	-	-	1	-	-	-
2003.....	-	-	-	-	-	1	-	3	3	1	-	-
2002.....	-	1	-	-	1	1	1	1	1	-	-	-
2001.....	-	-	-	-	-	-	-	3	-	1	-	-
2000.....	-	-	-	-	-	-	-	2	1	-	-	-
1999.....	-	-	-	-	-	1	2	1	1	1	1	-
American Indian 1/, male												
2004.....	1	-	-	-	-	-	-	-	1	-	-	-
2003.....	-	-	-	-	-	1	-	2	-	-	-	-
2002.....	-	1	-	-	-	-	1	1	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	1	-	-
2000.....	-	-	-	-	-	-	-	2	1	-	-	-
1999.....	-	-	-	-	-	-	1	1	1	-	-	-
American Indian 1/, female												
2004.....	-	-	-	-	-	3	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	1	3	1	-	-
2002.....	-	-	-	-	1	1	-	-	1	-	-	-
2001.....	-	-	-	-	-	-	-	3	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	1	1	-	-	1	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified organic mental disorders (F04-F09)												
Asian or Pacific Islander 2/, both sexes												
2004.....	-	-	1	-	-	-	3	1	1	1	-	-
2003.....	-	-	-	2	-	-	2	6	1	1	1	-
2002.....	1	-	-	2	1	2	3	4	2	-	-	-
2001.....	-	-	-	-	1	1	2	2	3	1	-	-
2000.....	-	-	-	1	1	1	1	5	5	-	-	-
1999.....	-	-	-	-	2	1	-	5	2	-	1	-
Asian or Pacific Islander 2/, male												
2004.....	-	-	1	-	-	-	2	-	1	1	-	-
2003.....	-	-	-	1	-	-	2	1	-	-	-	-
2002.....	1	-	-	2	-	1	1	3	-	-	-	-
2001.....	-	-	-	-	1	-	1	1	1	1	-	-
2000.....	-	-	-	1	1	-	-	5	2	-	-	-
1999.....	-	-	-	-	-	1	-	3	-	-	1	-
Asian or Pacific Islander 2/, female												
2004.....	-	-	-	-	-	-	1	1	-	-	-	-
2003.....	-	-	-	1	-	-	-	5	1	1	1	-
2002.....	-	-	-	-	1	1	2	1	2	-	-	-
2001.....	-	-	-	-	-	1	1	1	2	-	-	-
2000.....	-	-	-	-	-	1	1	-	3	-	-	-
1999.....	-	-	-	-	2	-	-	2	2	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Mental and behavioral disorders due to psychoactive substance use (F10-F19)												
All races, both sexes												
2004.....	1,678	1,272	905	667	520	404	294	142	62	13	8	8
2003.....	1,537	1,097	858	609	486	368	236	137	52	13	1	10
2002.....	1,426	1,040	776	581	462	351	203	120	40	9	-	5
2001.....	1,294	907	737	545	447	345	232	122	39	11	3	12
2000.....	1,143	864	653	553	451	351	225	116	50	5	3	8
1999.....	1,041	811	704	553	442	386	214	115	29	13	1	5
All races, male												
2004.....	1,315	1,025	734	502	384	280	187	80	34	7	3	8
2003.....	1,213	859	664	460	349	261	152	75	25	6	-	9
2002.....	1,160	837	621	460	312	241	124	67	25	5	-	5
2001.....	1,035	711	586	427	333	219	140	68	23	6	1	11
2000.....	921	679	521	429	323	243	145	61	25	3	1	8
1999.....	844	644	563	417	317	270	152	68	15	5	-	3
All races, female												
2004.....	363	247	171	165	136	124	107	62	28	6	5	-
2003.....	324	238	194	149	137	107	84	62	27	7	1	1
2002.....	266	203	155	121	150	110	79	53	15	4	-	-
2001.....	259	196	151	118	114	126	92	54	16	5	2	1
2000.....	222	185	132	124	128	108	80	55	25	2	2	-
1999.....	197	167	141	136	125	116	62	47	14	8	1	2
White, both sexes												
2004.....	1,305	1,025	725	530	431	357	275	133	54	11	8	4
2003.....	1,177	868	669	518	419	331	217	128	51	12	1	8
2002.....	1,093	806	587	474	390	313	182	102	37	9	-	3
2001.....	957	691	583	436	384	305	212	109	36	10	3	9
2000.....	875	664	503	441	389	310	207	108	44	4	3	6
1999.....	773	601	556	445	367	335	191	103	29	13	1	5

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Mental and behavioral disorders due to psychoactive substance use (F10-F19)												
White, male												
2004.....	1,022	825	578	385	311	242	177	75	28	5	3	4
2003.....	918	687	501	389	299	231	137	67	24	6	-	7
2002.....	896	640	468	363	254	212	108	56	24	5	-	3
2001.....	776	532	457	333	281	189	127	59	21	5	1	8
2000.....	709	521	399	335	274	209	131	57	20	2	1	6
1999.....	640	468	445	325	257	229	136	60	15	5	-	3
White, female												
2004.....	283	200	147	145	120	115	98	58	26	6	5	-
2003.....	259	181	168	129	120	100	80	61	27	6	1	1
2002.....	197	166	119	111	136	101	74	46	13	4	-	-
2001.....	181	159	126	103	103	116	85	50	15	5	2	1
2000.....	166	143	104	106	115	101	76	51	24	2	2	-
1999.....	133	133	111	120	110	106	55	43	14	8	1	2
Black, both sexes												
2004.....	326	214	153	113	76	35	16	8	7	2	-	4
2003.....	307	199	163	80	56	31	17	8	1	1	-	2
2002.....	299	213	155	90	62	34	19	11	1	-	-	1
2001.....	302	187	142	93	55	37	17	9	2	-	-	3
2000.....	245	174	136	100	56	39	15	7	6	1	-	2
1999.....	231	196	133	98	70	45	21	10	-	-	-	-
Black, male												
2004.....	251	180	133	98	62	30	8	4	5	2	-	4
2003.....	252	148	141	63	43	25	13	7	1	-	-	2
2002.....	239	178	128	83	50	25	15	8	-	-	-	1
2001.....	234	155	122	81	45	30	10	6	1	-	-	3
2000.....	197	140	110	83	45	33	11	3	5	1	-	2
1999.....	175	163	108	83	56	35	15	6	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Mental and behavioral disorders due to psychoactive substance use (F10-F19)

Black, female

2004.....	75	34	20	15	14	5	8	4	2	-	-	-
2003.....	55	51	22	17	13	6	4	1	-	1	-	-
2002.....	60	35	27	7	12	9	4	3	1	-	-	-
2001.....	68	32	20	12	10	7	7	3	1	-	-	-
2000.....	48	34	26	17	11	6	4	4	1	-	-	-
1999.....	56	33	25	15	14	10	6	4	-	-	-	-

American Indian 1/, both sexes

2004.....	33	27	22	18	9	2	1	-	-	-	-	-
2003.....	42	25	18	11	9	5	2	-	-	-	-	-
2002.....	24	14	26	10	7	1	2	4	-	-	-	-
2001.....	26	18	10	14	7	3	2	3	1	-	-	-
2000.....	19	17	11	10	5	-	2	1	-	-	-	-
1999.....	30	13	13	6	3	5	1	1	-	-	-	-

American Indian 1/, male

2004.....	28	15	19	13	8	1	1	-	-	-	-	-
2003.....	33	20	14	8	5	4	2	-	-	-	-	-
2002.....	18	12	19	8	6	1	1	1	-	-	-	-
2001.....	19	15	6	12	6	-	2	2	1	-	-	-
2000.....	11	12	9	9	4	-	2	1	-	-	-	-
1999.....	24	12	9	5	2	5	-	1	-	-	-	-

American Indian 1/, female

2004.....	5	12	3	5	1	1	-	-	-	-	-	-
2003.....	9	5	4	3	4	1	-	-	-	-	-	-
2002.....	6	2	7	2	1	-	1	3	-	-	-	-
2001.....	7	3	4	2	1	3	-	1	-	-	-	-
2000.....	8	5	2	1	1	-	-	-	-	-	-	-
1999.....	6	1	4	1	1	-	1	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Mental and behavioral disorders due to use of alcohol (F10)												
All races, both sexes												
2004.....	1,229	1,027	749	531	376	255	162	67	22	5	-	6
2003.....	1,167	903	731	475	352	232	140	70	17	7	-	8
2002.....	1,125	877	672	485	354	260	127	67	21	5	-	5
2001.....	1,028	780	654	483	356	247	155	61	25	7	1	11
2000.....	906	747	569	480	362	250	144	68	33	4	3	6
1999.....	838	697	612	469	339	281	143	69	18	8	1	3
All races, male												
2004.....	991	841	619	414	305	197	113	40	11	3	-	6
2003.....	936	723	575	369	273	174	110	41	7	3	-	7
2002.....	925	720	554	401	262	198	90	43	14	4	-	5
2001.....	827	619	531	387	284	177	104	39	17	4	1	10
2000.....	739	601	469	389	282	185	108	42	19	2	1	6
1999.....	695	562	492	368	259	217	111	47	9	2	-	2
All races, female												
2004.....	238	186	130	117	71	58	49	27	11	2	-	-
2003.....	231	180	156	106	79	58	30	29	10	4	-	1
2002.....	200	157	118	84	92	62	37	24	7	1	-	-
2001.....	201	161	123	96	72	70	51	22	8	3	-	1
2000.....	167	146	100	91	80	65	36	26	14	2	2	-
1999.....	143	135	120	101	80	64	32	22	9	6	1	1
White, both sexes												
2004.....	1,021	854	607	410	295	221	150	64	17	5	-	3
2003.....	943	743	579	397	294	200	123	63	17	6	-	6
2002.....	916	695	510	391	287	227	108	58	19	5	-	3
2001.....	790	599	519	380	301	215	140	51	23	7	1	8
2000.....	722	583	433	376	305	214	130	61	29	4	3	5
1999.....	647	532	476	372	273	236	122	60	18	8	1	3

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Mental and behavioral disorders due to use of alcohol (F10)												
White, male												
2004.....	819	699	495	311	239	169	107	39	8	3	-	3
2003.....	740	599	442	310	228	147	96	35	7	3	-	5
2002.....	760	563	423	316	207	173	76	34	13	4	-	3
2001.....	643	467	418	298	237	152	94	32	15	4	1	7
2000.....	589	468	354	300	235	153	96	38	16	2	1	5
1999.....	546	421	383	284	205	182	97	40	9	2	-	2
White, female												
2004.....	202	155	112	99	56	52	43	25	9	2	-	-
2003.....	203	144	137	87	66	53	27	28	10	3	-	1
2002.....	156	132	87	75	80	54	32	24	6	1	-	-
2001.....	147	132	101	82	64	63	46	19	8	3	-	1
2000.....	133	115	79	76	70	61	34	23	13	2	2	-
1999.....	101	111	93	88	68	54	25	20	9	6	1	1
Black, both sexes												
2004.....	174	146	117	97	69	27	9	3	4	-	-	3
2003.....	181	133	131	67	47	28	15	7	-	1	-	2
2002.....	177	162	129	78	59	29	17	7	1	-	-	1
2001.....	207	156	123	89	47	30	14	7	1	-	-	3
2000.....	163	139	122	92	51	34	12	6	4	-	-	1
1999.....	157	152	121	87	62	40	19	7	-	-	-	-
Black, male												
2004.....	143	127	103	84	55	23	4	1	2	-	-	3
2003.....	160	102	116	51	38	23	12	6	-	-	-	2
2002.....	141	139	106	72	49	21	13	7	-	-	-	1
2001.....	161	132	106	77	40	25	9	4	1	-	-	3
2000.....	135	115	103	78	43	31	10	3	3	-	-	1
1999.....	123	129	99	75	51	30	13	5	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Mental and behavioral disorders due to use of alcohol (F10)												
Black, female												
2004.....	31	19	14	13	14	4	5	2	2	-	-	-
2003.....	21	31	15	16	9	5	3	1	-	1	-	-
2002.....	36	23	23	6	10	8	4	-	1	-	-	-
2001.....	46	24	17	12	7	5	5	3	-	-	-	-
2000.....	28	24	19	14	8	3	2	3	1	-	-	-
1999.....	34	23	22	12	11	10	6	2	-	-	-	-
American Indian 1/, both sexes												
2004.....	27	23	20	18	8	1	1	-	-	-	-	-
2003.....	35	23	15	11	9	4	2	-	-	-	-	-
2002.....	24	13	25	10	6	1	2	1	-	-	-	-
2001.....	23	16	10	12	7	2	1	2	1	-	-	-
2000.....	17	16	11	10	5	-	2	1	-	-	-	-
1999.....	29	13	13	6	3	4	1	1	-	-	-	-
American Indian 1/, male												
2004.....	22	12	17	13	8	1	1	-	-	-	-	-
2003.....	28	19	11	8	5	4	2	-	-	-	-	-
2002.....	18	11	19	8	5	1	1	1	-	-	-	-
2001.....	17	13	6	11	6	-	1	2	1	-	-	-
2000.....	11	12	9	9	4	-	2	1	-	-	-	-
1999.....	23	12	9	5	2	4	-	1	-	-	-	-
American Indian 1/, female												
2004.....	5	11	3	5	-	-	-	-	-	-	-	-
2003.....	7	4	4	3	4	-	-	-	-	-	-	-
2002.....	6	2	6	2	1	-	1	-	-	-	-	-
2001.....	6	3	4	1	1	2	-	-	-	-	-	-
2000.....	6	4	2	1	1	-	-	-	-	-	-	-
1999.....	6	1	4	1	1	-	1	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Mental and behavioral disorders due to other psychoactive substance use (F11-F19)												
All races, both sexes												
2004.....	449	245	156	136	144	149	132	75	40	8	8	2
2003.....	370	194	127	134	134	136	96	67	35	6	1	2
2002.....	301	163	104	96	108	91	76	53	19	4	-	-
2001.....	266	127	83	62	91	98	77	61	14	4	2	1
2000.....	237	117	84	73	89	101	81	48	17	1	-	2
1999.....	203	114	92	84	103	105	71	46	11	5	-	2
All races, male												
2004.....	324	184	115	88	79	83	74	40	23	4	3	2
2003.....	277	136	89	91	76	87	42	34	18	3	-	2
2002.....	235	117	67	59	50	43	34	24	11	1	-	-
2001.....	208	92	55	40	49	42	36	29	6	2	-	1
2000.....	182	78	52	40	41	58	37	19	6	1	-	2
1999.....	149	82	71	49	58	53	41	21	6	3	-	1
All races, female												
2004.....	125	61	41	48	65	66	58	35	17	4	5	-
2003.....	93	58	38	43	58	49	54	33	17	3	1	-
2002.....	66	46	37	37	58	48	42	29	8	3	-	-
2001.....	58	35	28	22	42	56	41	32	8	2	2	-
2000.....	55	39	32	33	48	43	44	29	11	-	-	-
1999.....	54	32	21	35	45	52	30	25	5	2	-	1
White, both sexes												
2004.....	284	171	118	120	136	136	125	69	37	6	8	1
2003.....	234	125	90	121	125	131	94	65	34	6	1	2
2002.....	177	111	77	83	103	86	74	44	18	4	-	-
2001.....	167	92	64	56	83	90	72	58	13	3	2	1
2000.....	153	81	70	65	84	96	77	47	15	-	-	1
1999.....	126	69	80	73	94	99	69	43	11	5	-	2

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Mental and behavioral disorders due to other psychoactive substance use (F11-F19)

White, male

2004.....	203	126	83	74	72	73	70	36	20	2	3	1
2003.....	178	88	59	79	71	84	41	32	17	3	-	2
2002.....	136	77	45	47	47	39	32	22	11	1	-	-
2001.....	133	65	39	35	44	37	33	27	6	1	-	1
2000.....	120	53	45	35	39	56	35	19	4	-	-	1
1999.....	94	47	62	41	52	47	39	20	6	3	-	1

White, female

2004.....	81	45	35	46	64	63	55	33	17	4	5	-
2003.....	56	37	31	42	54	47	53	33	17	3	1	-
2002.....	41	34	32	36	56	47	42	22	7	3	-	-
2001.....	34	27	25	21	39	53	39	31	7	2	2	-
2000.....	33	28	25	30	45	40	42	28	11	-	-	-
1999.....	32	22	18	32	42	52	30	23	5	2	-	1

Black, both sexes

2004.....	152	68	36	16	7	8	7	5	3	2	-	1
2003.....	126	66	32	13	9	3	2	1	1	-	-	-
2002.....	122	51	26	12	3	5	2	4	-	-	-	-
2001.....	95	31	19	4	8	7	3	2	1	-	-	-
2000.....	82	35	14	8	5	5	3	1	2	1	-	1
1999.....	74	44	12	11	8	5	2	3	-	-	-	-

Black, male

2004.....	108	53	30	14	7	7	4	3	3	2	-	1
2003.....	92	46	25	12	5	2	1	1	1	-	-	-
2002.....	98	39	22	11	1	4	2	1	-	-	-	-
2001.....	73	23	16	4	5	5	1	2	-	-	-	-
2000.....	62	25	7	5	2	2	1	-	2	1	-	1
1999.....	52	34	9	8	5	5	2	1	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Schizophrenia, schizotypal and delusional disorders (F20-F29)												
All races, both sexes												
2004.....	30	34	27	35	52	68	72	72	43	14	1	-
2003.....	29	13	22	34	66	64	79	72	38	14	1	-
2002.....	18	29	17	30	57	73	70	80	42	23	7	-
2001.....	15	20	24	31	37	79	67	54	44	17	2	-
2000.....	19	17	21	31	63	67	86	72	48	19	3	-
1999.....	19	18	20	24	53	72	102	107	63	30	3	-
All races, male												
2004.....	19	20	15	17	25	28	23	23	10	2	-	-
2003.....	21	7	7	15	22	20	23	15	8	3	-	-
2002.....	13	15	12	14	26	27	17	17	7	3	-	-
2001.....	10	12	10	21	21	27	22	14	5	1	-	-
2000.....	9	13	11	15	29	22	30	18	7	4	-	-
1999.....	12	10	9	10	26	30	26	34	15	5	1	-
All races, female												
2004.....	11	14	12	18	27	40	49	49	33	12	1	-
2003.....	8	6	15	19	44	44	56	57	30	11	1	-
2002.....	5	14	5	16	31	46	53	63	35	20	7	-
2001.....	5	8	14	10	16	52	45	40	39	16	2	-
2000.....	10	4	10	16	34	45	56	54	41	15	3	-
1999.....	7	8	11	14	27	42	76	73	48	25	2	-
White, both sexes												
2004.....	24	30	24	26	47	59	66	63	39	13	1	-
2003.....	21	11	19	32	59	60	70	68	37	13	1	-
2002.....	14	21	17	24	49	65	62	76	37	20	7	-
2001.....	9	15	20	24	33	72	62	50	43	17	1	-
2000.....	17	16	16	29	57	61	81	68	43	17	3	-
1999.....	12	17	15	23	47	63	93	99	61	26	3	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Schizophrenia, schizotypal and delusional disorders (F20-F29)												
White, male												
2004.....	16	18	13	12	22	22	21	16	8	2	-	-
2003.....	14	6	6	15	20	19	21	14	8	3	-	-
2002.....	11	11	12	11	21	21	15	16	7	1	-	-
2001.....	7	8	9	18	19	24	20	13	5	1	-	-
2000.....	8	13	9	13	27	21	28	16	7	4	-	-
1999.....	6	10	7	10	23	23	23	32	14	4	1	-
White, female												
2004.....	8	12	11	14	25	37	45	47	31	11	1	-
2003.....	7	5	13	17	39	41	49	54	29	10	1	-
2002.....	3	10	5	13	28	44	47	60	30	19	7	-
2001.....	2	7	11	6	14	48	42	37	38	16	1	-
2000.....	9	3	7	16	30	40	53	52	36	13	3	-
1999.....	6	7	8	13	24	40	70	67	47	22	2	-
Black, both sexes												
2004.....	6	4	3	9	5	9	5	8	4	1	-	-
2003.....	8	2	2	2	6	4	7	4	1	1	-	-
2002.....	4	8	-	6	6	8	6	4	4	2	-	-
2001.....	5	5	4	7	4	7	3	4	1	-	1	-
2000.....	2	1	4	2	6	6	4	4	5	2	-	-
1999.....	5	1	4	1	6	7	9	7	2	4	-	-
Black, male												
2004.....	3	2	2	5	3	6	1	6	2	-	-	-
2003.....	7	1	1	-	2	1	2	1	-	-	-	-
2002.....	2	4	-	3	5	6	-	1	-	1	-	-
2001.....	3	4	1	3	2	3	1	1	-	-	-	-
2000.....	1	-	1	2	2	1	2	2	-	-	-	-
1999.....	4	-	1	-	3	5	3	1	1	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Mood (affective) disorders (F30-F39)												
All races, both sexes												
2004.....	17	20	13	22	56	101	146	200	149	54	7	-
2003.....	18	20	17	24	41	114	159	239	172	72	12	-
2002.....	15	12	25	23	57	98	185	187	144	71	15	-
2001.....	11	15	14	34	53	107	166	218	141	48	17	-
2000.....	14	17	16	25	47	92	142	177	168	57	14	-
1999.....	12	15	14	30	49	121	168	172	148	60	12	-
All races, male												
2004.....	11	6	4	12	27	32	44	58	36	12	3	-
2003.....	7	11	6	12	18	45	49	62	38	12	3	-
2002.....	7	4	12	10	21	27	53	37	37	17	1	-
2001.....	9	9	9	16	24	40	47	57	29	15	3	-
2000.....	6	11	8	9	20	26	43	59	39	14	-	-
1999.....	8	11	9	12	18	48	49	55	41	9	-	-
All races, female												
2004.....	6	14	9	10	29	69	102	142	113	42	4	-
2003.....	11	9	11	12	23	69	110	177	134	60	9	-
2002.....	8	8	13	13	36	71	132	150	107	54	14	-
2001.....	2	6	5	18	29	67	119	161	112	33	14	-
2000.....	8	6	8	16	27	66	99	118	129	43	14	-
1999.....	4	4	5	18	31	73	119	117	107	51	12	-
White, both sexes												
2004.....	15	16	13	22	53	93	138	195	145	54	7	-
2003.....	15	18	17	20	40	106	152	232	170	67	11	-
2002.....	13	12	21	20	54	95	171	182	141	70	12	-
2001.....	9	13	13	33	50	103	160	207	137	45	15	-
2000.....	13	17	15	23	44	86	136	172	162	57	14	-
1999.....	10	13	12	27	47	118	163	169	144	58	12	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Mood (affective) disorders (F30-F39)												
White, male												
2004.....	9	4	4	12	27	30	42	57	33	12	3	-
2003.....	6	9	6	10	18	43	48	61	37	11	3	-
2002.....	7	4	10	8	19	25	49	37	35	17	1	-
2001.....	7	8	9	15	23	39	45	55	28	13	2	-
2000.....	6	11	7	9	19	24	40	59	39	14	-	-
1999.....	6	9	8	12	18	46	47	54	39	9	-	-
White, female												
2004.....	6	12	9	10	26	63	96	138	112	42	4	-
2003.....	9	9	11	10	22	63	104	171	133	56	8	-
2002.....	6	8	11	12	35	70	122	145	106	53	11	-
2001.....	2	5	4	18	27	64	115	152	109	32	13	-
2000.....	7	6	8	14	25	62	96	113	123	43	14	-
1999.....	4	4	4	15	29	72	116	115	105	49	12	-
Black, both sexes												
2004.....	1	3	-	-	3	8	5	4	1	-	-	-
2003.....	3	1	-	2	1	7	6	6	1	4	-	-
2002.....	2	-	3	3	3	3	8	4	3	1	3	-
2001.....	2	2	1	1	3	3	4	8	3	1	2	-
2000.....	1	-	1	2	3	4	5	3	5	-	-	-
1999.....	2	1	2	3	2	3	3	2	4	2	-	-
Black, male												
2004.....	1	2	-	-	-	2	1	1	1	-	-	-
2003.....	1	1	-	1	-	1	1	1	-	1	-	-
2002.....	-	-	1	2	2	2	3	-	2	-	-	-
2001.....	2	1	-	1	1	1	2	2	-	1	1	-
2000.....	-	-	1	-	1	2	3	-	-	-	-	-
1999.....	2	1	1	-	-	2	2	1	2	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Mood (affective) disorders (F30-F39)												
Black, female												
2004.....	-	1	-	-	3	6	4	3	-	-	-	-
2003.....	2	-	-	1	1	6	5	5	1	3	-	-
2002.....	2	-	2	1	1	1	5	4	1	1	3	-
2001.....	-	1	1	-	2	2	2	6	3	-	1	-
2000.....	1	-	-	2	2	2	2	3	5	-	-	-
1999.....	-	-	1	3	2	1	1	1	2	2	-	-
American Indian 1/, both sexes												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	1	-	1	-	1	-	-	-	-	-	-
2002.....	-	-	1	-	-	-	1	-	-	-	-	-
2001.....	-	-	-	-	-	1	-	2	-	-	-	-
2000.....	-	-	-	-	-	-	1	1	1	-	-	-
1999.....	-	1	-	-	-	-	1	-	-	-	-	-
American Indian 1/, male												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	1	-	-	-	1	-	-	-	-	-	-
2002.....	-	-	1	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, female												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	1	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	1	-	-	-	-	-
2001.....	-	-	-	-	-	1	-	2	-	-	-	-
2000.....	-	-	-	-	-	-	1	1	1	-	-	-
1999.....	-	-	-	-	-	-	1	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Mood (affective) disorders (F30-F39)												
Asian or Pacific Islander 2/, both sexes												
2004.....	1	1	-	-	-	-	3	1	3	-	-	-
2003.....	-	-	-	1	-	-	1	1	1	1	1	-
2002.....	-	-	-	-	-	-	5	1	-	-	-	-
2001.....	-	-	-	-	-	-	2	1	1	2	-	-
2000.....	-	-	-	-	-	2	-	1	-	-	-	-
1999.....	-	-	-	-	-	-	1	1	-	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	1	-	-	-	-	-	1	-	2	-	-	-
2003.....	-	-	-	1	-	-	-	-	1	-	-	-
2002.....	-	-	-	-	-	-	1	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	1	1	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	-	1	-	-	-	-	2	1	1	-	-	-
2003.....	-	-	-	-	-	-	1	1	-	1	1	-
2002.....	-	-	-	-	-	-	4	1	-	-	-	-
2001.....	-	-	-	-	-	-	2	1	-	1	-	-
2000.....	-	-	-	-	-	2	-	1	-	-	-	-
1999.....	-	-	-	-	-	-	1	1	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Neurotic, stress-related and somatoform disorders (F40-F48)												
All races, both sexes												
2004.....	3	9	6	4	11	12	20	16	18	9	1	-
2003.....	6	6	4	6	15	10	16	23	13	3	2	1
2002.....	8	5	6	7	11	16	17	15	8	8	3	-
2001.....	3	4	3	9	9	10	8	11	11	5	1	-
2000.....	2	1	6	4	7	11	7	14	11	2	-	-
1999.....	4	4	3	3	11	17	14	14	14	1	2	-
All races, male												
2004.....	2	7	3	1	6	6	11	5	4	2	-	-
2003.....	2	5	1	4	5	7	9	4	2	-	1	-
2002.....	6	3	3	4	7	6	8	5	3	-	-	-
2001.....	3	2	1	6	3	5	5	2	5	1	1	-
2000.....	2	-	-	2	5	6	2	6	4	-	-	-
1999.....	4	3	-	2	4	8	7	3	3	-	-	-
All races, female												
2004.....	1	2	3	3	5	6	9	11	14	7	1	-
2003.....	4	1	3	2	10	3	7	19	11	3	1	1
2002.....	2	2	3	3	4	10	9	10	5	8	3	-
2001.....	-	2	2	3	6	5	3	9	6	4	-	-
2000.....	-	1	6	2	2	5	5	8	7	2	-	-
1999.....	-	1	3	1	7	9	7	11	11	1	2	-
White, both sexes												
2004.....	2	8	5	2	11	11	20	15	18	9	1	-
2003.....	5	6	4	6	15	9	14	21	13	3	2	1
2002.....	6	4	6	3	10	13	14	14	7	8	3	-
2001.....	3	3	3	7	8	10	8	9	11	4	1	-
2000.....	2	1	6	2	7	11	7	13	11	1	-	-
1999.....	3	3	3	3	11	15	14	13	13	1	2	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Neurotic, stress-related and somatoform disorders (F40-F48)												
White, male												
2004.....	1	7	2	-	6	5	11	5	4	2	-	-
2003.....	2	5	1	4	5	6	7	4	2	-	1	-
2002.....	5	2	3	1	7	5	7	5	2	-	-	-
2001.....	3	1	1	5	3	5	5	1	5	-	1	-
2000.....	2	-	-	-	5	6	2	6	4	-	-	-
1999.....	3	2	-	2	4	7	7	3	3	-	-	-
White, female												
2004.....	1	1	3	2	5	6	9	10	14	7	1	-
2003.....	3	1	3	2	10	3	7	17	11	3	1	1
2002.....	1	2	3	2	3	8	7	9	5	8	3	-
2001.....	-	2	2	2	5	5	3	8	6	4	-	-
2000.....	-	1	6	2	2	5	5	7	7	1	-	-
1999.....	-	1	3	1	7	8	7	10	10	1	2	-
Black, both sexes												
2004.....	-	1	1	2	-	-	-	1	-	-	-	-
2003.....	1	-	-	-	-	1	-	2	-	-	-	-
2002.....	2	1	-	4	1	3	2	-	1	-	-	-
2001.....	-	1	-	1	1	-	-	-	-	1	-	-
2000.....	-	-	-	2	-	-	-	1	-	1	-	-
1999.....	1	1	-	-	-	2	-	-	-	-	-	-
Black, male												
2004.....	-	-	1	1	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	1	-	-	-	-	-	-
2002.....	1	1	-	3	-	1	-	-	1	-	-	-
2001.....	-	1	-	-	-	-	-	-	-	1	-	-
2000.....	-	-	-	2	-	-	-	-	-	-	-	-
1999.....	1	1	-	-	-	1	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Neurotic, stress-related and somatoform disorders (F40-F48)												
Asian or Pacific Islander 2/, both sexes												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	1	-	-	-	-	-
2002.....	-	-	-	-	-	-	1	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	2	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	1	1	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	1	-	-	-	-	-
2002.....	-	-	-	-	-	-	1	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	1	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	1	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	1	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Mental retardation (F70-F79)												
All races, both sexes												
2004.....	56	40	49	31	37	39	23	12	6	-	-	-
2003.....	56	57	44	36	30	31	24	16	4	1	-	-
2002.....	33	54	45	34	32	23	28	13	9	1	-	-
2001.....	42	39	34	32	33	27	26	14	5	-	-	-
2000.....	36	44	31	21	29	30	16	16	5	1	-	-
1999.....	34	33	32	29	31	25	26	8	3	1	-	-
All races, male												
2004.....	39	18	28	20	26	19	8	3	1	-	-	-
2003.....	32	35	23	15	18	18	13	9	3	-	-	-
2002.....	25	33	24	18	18	13	12	5	2	-	-	-
2001.....	21	17	24	16	16	16	16	5	1	-	-	-
2000.....	21	22	18	12	16	17	6	8	2	-	-	-
1999.....	23	20	15	21	19	16	15	3	2	1	-	-
All races, female												
2004.....	17	22	21	11	11	20	15	9	5	-	-	-
2003.....	24	22	21	21	12	13	11	7	1	1	-	-
2002.....	8	21	21	16	14	10	16	8	7	1	-	-
2001.....	21	22	10	16	17	11	10	9	4	-	-	-
2000.....	15	22	13	9	13	13	10	8	3	1	-	-
1999.....	11	13	17	8	12	9	11	5	1	-	-	-
White, both sexes												
2004.....	50	36	45	27	36	35	22	11	5	-	-	-
2003.....	46	50	41	33	25	29	21	15	4	1	-	-
2002.....	29	51	42	31	29	18	27	13	9	1	-	-
2001.....	37	33	32	27	30	23	26	13	5	-	-	-
2000.....	32	41	29	20	25	28	15	16	4	1	-	-
1999.....	30	28	27	27	27	22	25	8	3	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Mental retardation (F70-F79)												
White, male												
2004.....	35	17	25	18	25	17	8	3	1	-	-	-
2003.....	26	31	20	13	15	17	12	8	3	-	-	-
2002.....	22	30	22	16	16	10	12	5	2	-	-	-
2001.....	18	12	22	13	15	13	16	5	1	-	-	-
2000.....	19	22	16	12	13	16	6	8	2	-	-	-
1999.....	20	18	13	19	15	14	15	3	2	1	-	-
White, female												
2004.....	15	19	20	9	11	18	14	8	4	-	-	-
2003.....	20	19	21	20	10	12	9	7	1	1	-	-
2002.....	7	21	20	15	13	8	15	8	7	1	-	-
2001.....	19	21	10	14	15	10	10	8	4	-	-	-
2000.....	13	19	13	8	12	12	9	8	2	1	-	-
1999.....	10	10	14	8	12	8	10	5	1	-	-	-
Black, both sexes												
2004.....	6	4	4	4	1	4	1	1	1	-	-	-
2003.....	9	7	3	2	5	2	3	1	-	-	-	-
2002.....	3	2	3	3	2	5	1	-	-	-	-	-
2001.....	5	6	2	5	3	2	-	1	-	-	-	-
2000.....	4	3	2	1	4	1	1	-	1	-	-	-
1999.....	4	5	4	2	4	3	1	-	-	-	-	-
Black, male												
2004.....	4	1	3	2	1	2	-	-	-	-	-	-
2003.....	5	4	3	1	3	1	1	1	-	-	-	-
2002.....	2	2	2	2	1	3	-	-	-	-	-	-
2001.....	3	5	2	3	1	2	-	-	-	-	-	-
2000.....	2	-	2	-	3	-	-	-	-	-	-	-
1999.....	3	2	2	2	4	2	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other and unspecified mental and behavioral disorders (F50-F69,F80-F99)

All races, both sexes

2004.....	12	19	12	8	16	36	58	72	60	47	11	-
2003.....	5	13	13	9	28	53	61	78	78	39	11	-
2002.....	15	8	10	16	23	29	65	80	75	36	10	-
2001.....	13	6	8	13	18	39	50	59	62	45	16	-
2000.....	11	6	14	7	19	31	58	51	71	28	14	-
1999.....	9	7	4	14	32	32	49	68	60	30	10	-

All races, male

2004.....	4	6	2	4	9	11	24	26	19	8	3	-
2003.....	3	5	4	3	9	23	26	23	22	10	5	-
2002.....	5	1	4	8	10	14	32	27	16	6	1	-
2001.....	6	2	5	7	8	16	18	22	10	9	5	-
2000.....	4	3	5	3	7	8	23	19	16	4	1	-
1999.....	2	2	1	8	17	14	14	11	13	6	-	-

All races, female

2004.....	8	13	10	4	7	25	34	46	41	39	8	-
2003.....	2	8	9	6	19	30	35	55	56	29	6	-
2002.....	10	7	6	8	13	15	33	53	59	30	9	-
2001.....	7	4	3	6	10	23	32	37	52	36	11	-
2000.....	7	3	9	4	12	23	35	32	55	24	13	-
1999.....	7	5	3	6	15	18	35	57	47	24	10	-

White, both sexes

2004.....	11	17	9	4	14	30	54	66	58	43	9	-
2003.....	5	12	11	9	25	46	53	72	71	38	8	-
2002.....	14	6	9	13	19	26	61	70	68	33	10	-
2001.....	11	5	7	13	16	36	45	57	59	44	15	-
2000.....	10	4	13	6	17	25	53	48	65	27	12	-
1999.....	9	5	2	11	29	31	47	65	59	29	9	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified mental and behavioral disorders (F50-F69,F80-F99)												
White, male												
2004.....	3	4	1	3	9	8	23	21	18	7	2	-
2003.....	3	4	4	3	7	20	24	20	21	10	4	-
2002.....	4	-	4	6	7	13	30	25	14	6	1	-
2001.....	5	1	4	7	7	15	15	21	9	9	4	-
2000.....	4	2	4	2	6	6	21	18	15	3	-	-
1999.....	2	1	-	6	16	14	13	10	13	6	-	-
White, female												
2004.....	8	13	8	1	5	22	31	45	40	36	7	-
2003.....	2	8	7	6	18	26	29	52	50	28	4	-
2002.....	10	6	5	7	12	13	31	45	54	27	9	-
2001.....	6	4	3	6	9	21	30	36	50	35	11	-
2000.....	6	2	9	4	11	19	32	30	50	24	12	-
1999.....	7	4	2	5	13	17	34	55	46	23	9	-
Black, both sexes												
2004.....	1	2	2	3	1	5	4	2	1	3	-	-
2003.....	-	1	2	-	1	5	7	6	6	1	2	-
2002.....	1	2	1	2	2	3	4	8	5	3	-	-
2001.....	2	1	1	-	1	2	5	2	2	1	1	-
2000.....	1	1	1	1	2	6	4	2	5	1	1	-
1999.....	-	2	2	3	2	-	2	2	-	1	1	-
Black, male												
2004.....	1	2	-	-	-	2	1	2	1	1	-	-
2003.....	-	1	-	-	1	2	2	3	1	-	1	-
2002.....	1	1	-	2	2	1	2	1	-	-	-	-
2001.....	1	1	1	-	1	-	3	1	1	-	1	-
2000.....	-	1	1	1	2	1	1	1	1	1	1	-
1999.....	-	1	1	2	-	-	1	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified mental and behavioral disorders (F50-F69,F80-F99)												
Asian or Pacific Islander 2/, both sexes												
2004.....	-	-	1	-	1	1	-	3	1	1	2	-
2003.....	-	-	-	-	1	1	1	-	1	-	1	-
2002.....	-	-	-	-	1	-	-	2	2	-	-	-
2001.....	-	-	-	-	1	1	-	-	1	-	-	-
2000.....	-	-	-	-	-	-	1	1	1	-	1	-
1999.....	-	-	-	-	1	1	-	1	1	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	-	-	1	-	-	1	-	3	-	-	1	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	1	2	-	-	-
2001.....	-	-	-	-	-	1	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	1	-	-	-	-	-
1999.....	-	-	-	-	1	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	-	-	-	-	1	-	-	-	1	1	1	-
2003.....	-	-	-	-	1	1	1	-	1	-	1	-
2002.....	-	-	-	-	1	-	-	1	-	-	-	-
2001.....	-	-	-	-	1	-	-	-	1	-	-	-
2000.....	-	-	-	-	-	-	-	1	1	-	1	-
1999.....	-	-	-	-	-	1	-	-	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VI. Diseases of the nervous system (G00-G98)												
All races, both sexes												
2004.....	2,144	2,692	3,146	4,215	7,098	14,141	22,539	24,306	17,278	6,548	1,178	1
2003.....	2,095	2,558	3,113	4,194	7,586	14,394	21,665	23,421	16,576	6,397	1,056	1
2002.....	1,979	2,350	2,890	4,088	7,235	13,741	20,663	22,024	14,850	5,852	954	1
2001.....	1,903	2,228	2,672	3,908	7,119	13,246	19,310	20,262	13,616	5,145	885	-
2000.....	1,938	2,104	2,567	3,836	7,041	12,820	17,898	18,587	12,693	4,617	773	3
1999.....	1,748	1,884	2,366	3,753	6,772	12,014	16,619	16,894	11,290	3,996	672	-
All races, male												
2004.....	1,140	1,403	1,612	2,178	3,622	6,550	9,360	8,297	4,241	1,088	106	1
2003.....	1,134	1,337	1,609	2,138	3,848	6,775	8,964	7,891	4,077	1,037	113	1
2002.....	1,026	1,195	1,515	2,142	3,696	6,449	8,484	7,378	3,635	944	98	1
2001.....	986	1,115	1,358	2,090	3,610	6,299	8,125	6,937	3,355	803	98	-
2000.....	1,016	1,074	1,330	2,045	3,622	6,038	7,477	6,223	3,107	732	76	2
1999.....	914	1,003	1,239	1,974	3,470	5,833	7,104	5,775	2,734	655	57	-
All races, female												
2004.....	1,004	1,289	1,534	2,037	3,476	7,591	13,179	16,009	13,037	5,460	1,072	-
2003.....	961	1,221	1,504	2,056	3,738	7,619	12,701	15,530	12,499	5,360	943	-
2002.....	953	1,155	1,375	1,946	3,539	7,292	12,179	14,646	11,215	4,908	856	-
2001.....	917	1,113	1,314	1,818	3,509	6,947	11,185	13,325	10,261	4,342	787	-
2000.....	922	1,030	1,237	1,791	3,419	6,782	10,421	12,364	9,586	3,885	697	1
1999.....	834	881	1,127	1,779	3,302	6,181	9,515	11,119	8,556	3,341	615	-
White, both sexes												
2004.....	1,770	2,284	2,732	3,695	6,432	12,966	20,945	22,780	16,155	6,072	1,033	-
2003.....	1,705	2,191	2,686	3,681	6,865	13,258	20,269	22,015	15,620	6,004	943	1
2002.....	1,604	2,016	2,524	3,607	6,564	12,698	19,434	20,755	13,997	5,486	853	1
2001.....	1,576	1,886	2,316	3,453	6,509	12,232	18,102	19,104	12,862	4,810	777	-
2000.....	1,623	1,799	2,230	3,409	6,393	11,922	16,826	17,512	12,027	4,332	697	2
1999.....	1,447	1,607	2,039	3,329	6,149	11,147	15,682	15,973	10,739	3,749	614	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

VI. Diseases of the nervous system (G00-G98)

White, male

2004.....	949	1,196	1,417	1,927	3,313	6,049	8,786	7,806	3,969	1,011	87	-
2003.....	932	1,161	1,383	1,884	3,521	6,297	8,409	7,447	3,838	955	101	1
2002.....	836	1,030	1,340	1,891	3,373	6,004	7,997	6,997	3,419	874	81	1
2001.....	817	954	1,190	1,861	3,321	5,868	7,652	6,549	3,154	758	82	-
2000.....	872	915	1,156	1,845	3,322	5,634	7,061	5,866	2,920	685	62	2
1999.....	760	868	1,076	1,765	3,195	5,444	6,767	5,467	2,582	604	49	-

White, female

2004.....	821	1,088	1,315	1,768	3,119	6,917	12,159	14,974	12,186	5,061	946	-
2003.....	773	1,030	1,303	1,797	3,344	6,961	11,860	14,568	11,782	5,049	842	-
2002.....	768	986	1,184	1,716	3,191	6,694	11,437	13,758	10,578	4,612	772	-
2001.....	759	932	1,126	1,592	3,188	6,364	10,450	12,555	9,708	4,052	695	-
2000.....	751	884	1,074	1,564	3,071	6,288	9,765	11,646	9,107	3,647	635	-
1999.....	687	739	963	1,564	2,954	5,703	8,915	10,506	8,157	3,145	565	-

Black, both sexes

2004.....	331	360	369	438	557	984	1,321	1,274	954	404	123	-
2003.....	349	324	364	443	594	937	1,163	1,149	814	334	100	-
2002.....	336	280	324	409	569	874	1,029	1,063	731	327	86	-
2001.....	300	302	305	391	517	875	1,010	992	649	298	96	-
2000.....	288	270	287	377	548	766	920	911	569	248	66	1
1999.....	276	244	282	363	545	737	782	780	470	218	54	-

Black, male

2004.....	174	180	170	203	259	403	457	386	205	59	16	-
2003.....	177	153	192	219	273	395	440	340	188	58	9	-
2002.....	165	138	155	209	281	366	397	294	163	61	14	-
2001.....	152	141	145	194	243	367	377	310	166	34	13	-
2000.....	132	145	148	179	260	338	339	272	140	43	11	-
1999.....	147	114	140	178	239	321	265	244	116	43	8	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

VI. Diseases of the nervous system (G00-G98)

Black, female

2004.....	157	180	199	235	298	581	864	888	749	345	107	-
2003.....	172	171	172	224	321	542	723	809	626	276	91	-
2002.....	171	142	169	200	288	508	632	769	568	266	72	-
2001.....	148	161	160	197	274	508	633	682	483	264	83	-
2000.....	156	125	139	198	288	428	581	639	429	205	55	1
1999.....	129	130	142	185	306	416	517	536	354	175	46	-

American Indian 1/, both sexes

2004.....	21	13	12	15	14	34	54	33	24	9	8	1
2003.....	16	7	17	19	28	36	54	47	29	7	3	-
2002.....	12	12	13	17	31	40	42	41	27	5	3	-
2001.....	11	15	16	19	17	27	42	28	16	10	-	-
2000.....	8	11	11	17	28	38	34	29	21	7	-	-
1999.....	8	11	15	16	20	25	30	26	17	7	2	-

American Indian 1/, male

2004.....	8	11	9	8	7	17	25	10	6	1	1	1
2003.....	11	2	10	8	13	12	21	16	4	-	-	-
2002.....	8	7	7	11	15	23	16	17	8	-	1	-
2001.....	7	9	6	10	10	13	17	8	1	1	-	-
2000.....	2	4	3	6	10	13	16	10	9	1	-	-
1999.....	3	9	8	9	9	12	16	3	6	-	-	-

American Indian 1/, female

2004.....	13	2	3	7	7	17	29	23	18	8	7	-
2003.....	5	5	7	11	15	24	33	31	25	7	3	-
2002.....	4	5	6	6	16	17	26	24	19	5	2	-
2001.....	4	6	10	9	7	14	25	20	15	9	-	-
2000.....	6	7	8	11	18	25	18	19	12	6	-	-
1999.....	5	2	7	7	11	13	14	23	11	7	2	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VI. Diseases of the nervous system (G00-G98)												
Asian or Pacific Islander 2/, both sexes												
2004.....	22	35	33	67	95	157	219	219	145	63	14	-
2003.....	25	36	46	51	99	163	179	210	113	52	10	-
2002.....	27	42	29	55	71	129	158	165	95	34	12	-
2001.....	16	25	35	45	76	112	156	138	89	27	12	-
2000.....	19	24	39	33	72	94	118	135	76	30	10	-
1999.....	17	22	30	45	58	105	125	115	64	22	2	-
Asian or Pacific Islander 2/, male												
2004.....	9	16	16	40	43	81	92	95	61	17	2	-
2003.....	14	21	24	27	41	71	94	88	47	24	3	-
2002.....	17	20	13	31	27	56	74	70	45	9	2	-
2001.....	10	11	17	25	36	51	79	70	34	10	3	-
2000.....	10	10	23	15	30	53	61	75	38	3	3	-
1999.....	4	12	15	22	27	56	56	61	30	8	-	-
Asian or Pacific Islander 2/, female												
2004.....	13	19	17	27	52	76	127	124	84	46	12	-
2003.....	11	15	22	24	58	92	85	122	66	28	7	-
2002.....	10	22	16	24	44	73	84	95	50	25	10	-
2001.....	6	14	18	20	40	61	77	68	55	17	9	-
2000.....	9	14	16	18	42	41	57	60	38	27	7	-
1999.....	13	10	15	23	31	49	69	54	34	14	2	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Meningitis (G00,G03)												
All races, both sexes												
2004.....	60	52	40	50	42	45	40	30	11	1	-	-
2003.....	58	55	53	57	39	55	35	33	18	1	-	1
2002.....	53	48	40	46	63	47	37	36	16	3	-	1
2001.....	50	58	55	46	48	46	50	25	12	4	-	-
2000.....	55	43	47	48	60	68	55	32	16	2	-	-
1999.....	46	41	49	49	66	65	57	40	20	2	-	-
All races, male												
2004.....	35	28	20	18	15	19	19	14	4	1	-	-
2003.....	38	30	31	27	18	26	14	13	5	1	-	1
2002.....	30	29	19	26	36	21	20	11	8	-	-	1
2001.....	30	30	31	24	27	26	21	11	3	1	-	-
2000.....	31	26	28	26	34	41	21	13	5	1	-	-
1999.....	29	24	21	28	32	26	23	13	7	-	-	-
All races, female												
2004.....	25	24	20	32	27	26	21	16	7	-	-	-
2003.....	20	25	22	30	21	29	21	20	13	-	-	-
2002.....	23	19	21	20	27	26	17	25	8	3	-	-
2001.....	20	28	24	22	21	20	29	14	9	3	-	-
2000.....	24	17	19	22	26	27	34	19	11	1	-	-
1999.....	17	17	28	21	34	39	34	27	13	2	-	-
White, both sexes												
2004.....	40	35	32	38	31	37	32	29	7	1	-	-
2003.....	37	36	36	46	30	48	30	28	13	1	-	1
2002.....	32	33	25	38	50	37	31	35	12	3	-	1
2001.....	35	40	44	35	43	40	42	20	9	2	-	-
2000.....	36	29	36	32	47	56	48	24	14	2	-	-
1999.....	33	32	39	38	51	47	48	36	12	2	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Meningitis (G00,G03)												
White, male												
2004.....	25	16	15	15	10	15	17	14	3	1	-	-
2003.....	23	22	21	23	14	25	12	11	4	1	-	1
2002.....	18	20	12	22	29	17	19	10	5	-	-	1
2001.....	21	21	24	17	24	21	18	7	2	-	-	-
2000.....	22	20	22	19	29	33	19	9	4	1	-	-
1999.....	19	20	17	21	25	17	21	12	4	-	-	-
White, female												
2004.....	15	19	17	23	21	22	15	15	4	-	-	-
2003.....	14	14	15	23	16	23	18	17	9	-	-	-
2002.....	14	13	13	16	21	20	12	25	7	3	-	-
2001.....	14	19	20	18	19	19	24	13	7	2	-	-
2000.....	14	9	14	13	18	23	29	15	10	1	-	-
1999.....	14	12	22	17	26	30	27	24	8	2	-	-
Black, both sexes												
2004.....	17	14	7	11	8	7	6	1	2	-	-	-
2003.....	17	18	17	10	8	5	4	4	5	-	-	-
2002.....	19	12	11	8	13	9	4	1	3	-	-	-
2001.....	13	16	9	9	5	5	6	4	2	2	-	-
2000.....	17	13	9	15	12	8	5	7	2	-	-	-
1999.....	13	8	10	9	12	17	8	4	8	-	-	-
Black, male												
2004.....	9	9	4	3	3	3	2	-	-	-	-	-
2003.....	11	8	10	4	4	-	2	1	1	-	-	-
2002.....	11	7	5	4	7	4	1	1	2	-	-	-
2001.....	8	8	6	6	3	4	2	4	1	1	-	-
2000.....	8	6	5	6	5	5	2	4	1	-	-	-
1999.....	10	4	4	6	6	8	2	1	3	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Meningitis (G00,G03)												
Asian or Pacific Islander 2/, both sexes												
2004.....	1	1	1	1	3	-	1	-	2	-	-	-
2003.....	2	-	-	-	1	2	1	1	-	-	-	-
2002.....	1	2	3	-	-	1	1	-	1	-	-	-
2001.....	-	1	2	-	-	1	1	1	1	-	-	-
2000.....	2	1	2	1	1	2	1	1	-	-	-	-
1999.....	-	1	-	1	3	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	-	1	1	-	2	-	-	-	1	-	-	-
2003.....	2	-	-	-	-	1	-	1	-	-	-	-
2002.....	-	1	1	-	-	-	-	-	1	-	-	-
2001.....	-	-	1	-	-	1	1	-	-	-	-	-
2000.....	1	-	1	1	-	2	-	-	-	-	-	-
1999.....	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	1	-	-	1	1	-	1	-	1	-	-	-
2003.....	-	-	-	-	1	1	1	-	-	-	-	-
2002.....	1	1	2	-	-	1	1	-	-	-	-	-
2001.....	-	1	1	-	-	-	-	1	1	-	-	-
2000.....	1	1	1	-	1	-	1	1	-	-	-	-
1999.....	-	1	-	1	2	-	1	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other inflammatory diseases of central nervous system (G04-G09)

All races, both sexes

2004.....	43	48	55	40	68	72	70	38	11	6	-	-
2003.....	43	51	51	67	59	75	81	31	14	5	-	-
2002.....	45	50	48	51	66	80	76	39	11	4	-	-
2001.....	40	43	44	50	70	68	55	30	11	2	1	-
2000.....	41	43	37	49	68	84	55	28	11	2	-	-
1999.....	36	38	35	45	61	51	51	30	9	5	1	-

All races, male

2004.....	21	31	32	27	38	38	39	19	3	1	-	-
2003.....	31	29	30	36	36	39	35	15	2	2	-	-
2002.....	25	28	28	34	37	40	41	12	4	2	-	-
2001.....	23	22	22	29	32	33	25	18	4	-	1	-
2000.....	29	20	22	25	40	50	26	9	4	-	-	-
1999.....	19	30	14	25	23	27	28	13	4	1	-	-

All races, female

2004.....	22	17	23	13	30	34	31	19	8	5	-	-
2003.....	12	22	21	31	23	36	46	16	12	3	-	-
2002.....	20	22	20	17	29	40	35	27	7	2	-	-
2001.....	17	21	22	21	38	35	30	12	7	2	-	-
2000.....	12	23	15	24	28	34	29	19	7	2	-	-
1999.....	17	8	21	20	38	24	23	17	5	4	1	-

White, both sexes

2004.....	33	40	46	36	59	64	62	33	11	6	-	-
2003.....	36	42	40	58	54	68	74	28	13	5	-	-
2002.....	32	44	39	42	53	68	68	39	11	4	-	-
2001.....	32	40	34	39	56	58	48	29	8	1	1	-
2000.....	35	33	34	39	56	74	46	25	10	2	-	-
1999.....	29	29	26	38	51	46	47	26	9	5	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other inflammatory diseases of central nervous system (G04-G09)

White, male

2004.....	20	27	29	23	33	34	34	15	3	1	-	-
2003.....	27	25	24	29	32	38	32	13	1	2	-	-
2002.....	17	25	23	27	31	35	37	12	4	2	-	-
2001.....	18	21	18	24	26	26	23	17	3	-	1	-
2000.....	25	15	21	24	35	46	23	9	4	-	-	-
1999.....	16	23	12	20	21	26	26	10	4	1	-	-

White, female

2004.....	13	13	17	13	26	30	28	18	8	5	-	-
2003.....	9	17	16	29	22	30	42	15	12	3	-	-
2002.....	15	19	16	15	22	33	31	27	7	2	-	-
2001.....	14	19	16	15	30	32	25	12	5	1	-	-
2000.....	10	18	13	15	21	28	23	16	6	2	-	-
1999.....	13	6	14	18	30	20	21	16	5	4	1	-

Black, both sexes

2004.....	8	8	8	1	9	8	8	5	-	-	-	-
2003.....	5	8	9	8	4	4	5	2	1	-	-	-
2002.....	12	4	9	7	11	10	4	-	-	-	-	-
2001.....	7	2	10	11	11	9	5	1	3	1	-	-
2000.....	6	7	2	9	8	7	7	3	1	-	-	-
1999.....	7	8	8	7	8	5	2	3	-	-	-	-

Black, male

2004.....	1	4	2	1	5	4	5	4	-	-	-	-
2003.....	2	4	5	6	3	1	2	2	1	-	-	-
2002.....	8	1	5	6	5	5	2	-	-	-	-	-
2001.....	4	1	4	5	5	6	1	1	1	-	-	-
2000.....	4	4	1	1	3	4	2	-	-	-	-	-
1999.....	3	6	2	5	2	1	1	2	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other inflammatory diseases of central nervous system (G04-G09)												
Black, female												
2004.....	7	4	6	-	4	4	3	1	-	-	-	-
2003.....	3	4	4	2	1	3	3	-	-	-	-	-
2002.....	4	3	4	1	6	5	2	-	-	-	-	-
2001.....	3	1	6	6	6	3	4	-	2	1	-	-
2000.....	2	3	1	8	5	3	5	3	1	-	-	-
1999.....	4	2	6	2	6	4	1	1	-	-	-	-
American Indian 1/, both sexes												
2004.....	-	-	1	-	-	-	-	-	-	-	-	-
2003.....	-	1	1	1	1	-	-	1	-	-	-	-
2002.....	-	1	-	-	-	-	1	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	1	1	-	1	1	-	-	-	-	-	-
1999.....	-	1	-	-	1	-	-	-	-	-	-	-
American Indian 1/, male												
2004.....	-	-	1	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	1	1	-	-	-	-	-	-	-
2002.....	-	1	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	1	-	-	-	-	-	-	-	-	-	-
1999.....	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, female												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	1	1	-	-	-	-	1	-	-	-	-
2002.....	-	-	-	-	-	-	1	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	1	-	1	1	-	-	-	-	-	-
1999.....	-	-	-	-	1	-	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other inflammatory diseases of central nervous system (G04-G09)												
Asian or Pacific Islander 2/, both sexes												
2004.....	2	-	-	3	-	-	-	-	-	-	-	-
2003.....	2	-	1	-	-	3	2	-	-	-	-	-
2002.....	1	1	-	2	2	2	3	-	-	-	-	-
2001.....	1	1	-	-	3	1	2	-	-	-	-	-
2000.....	-	2	-	1	3	2	2	-	-	-	-	-
1999.....	-	-	1	-	1	-	2	1	-	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	-	-	-	3	-	-	-	-	-	-	-	-
2003.....	2	-	1	-	-	-	1	-	-	-	-	-
2002.....	-	1	-	1	1	-	2	-	-	-	-	-
2001.....	1	-	-	-	1	1	1	-	-	-	-	-
2000.....	-	-	-	-	2	-	1	-	-	-	-	-
1999.....	-	-	-	-	-	-	1	1	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	2	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	3	1	-	-	-	-	-
2002.....	1	-	-	1	1	2	1	-	-	-	-	-
2001.....	-	1	-	-	2	-	1	-	-	-	-	-
2000.....	-	2	-	1	1	2	1	-	-	-	-	-
1999.....	-	-	1	-	1	-	1	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Parkinson's disease (G20-G21)												
All races, both sexes												
2004.....	48	103	246	640	1,574	3,526	5,226	4,174	1,937	441	46	-
2003.....	46	112	240	624	1,706	3,671	5,052	4,157	1,900	425	31	-
2002.....	39	95	223	626	1,600	3,368	4,771	3,959	1,781	440	32	-
2001.....	32	100	201	554	1,593	3,446	4,679	3,786	1,655	425	43	-
2000.....	33	82	188	626	1,482	3,322	4,330	3,567	1,627	369	27	1
1999.....	35	73	169	539	1,481	3,066	4,044	3,332	1,492	313	31	-
All races, male												
2004.....	37	71	156	421	1,036	2,189	3,117	2,260	888	160	10	-
2003.....	23	65	154	402	1,115	2,300	2,891	2,210	846	143	14	-
2002.....	26	57	136	433	1,054	2,065	2,746	2,109	793	147	10	-
2001.....	25	69	145	362	1,032	2,104	2,733	2,012	759	134	10	-
2000.....	19	55	115	398	940	2,052	2,478	1,879	726	115	8	1
1999.....	25	44	117	338	965	1,940	2,367	1,699	639	110	6	-
All races, female												
2004.....	11	32	90	219	538	1,337	2,109	1,914	1,049	281	36	-
2003.....	23	47	86	222	591	1,371	2,161	1,947	1,054	282	17	-
2002.....	13	38	87	193	546	1,303	2,025	1,850	988	293	22	-
2001.....	7	31	56	192	561	1,342	1,946	1,774	896	291	33	-
2000.....	14	27	73	228	542	1,270	1,852	1,688	901	254	19	-
1999.....	10	29	52	201	516	1,126	1,677	1,633	853	203	25	-
White, both sexes												
2004.....	40	97	224	582	1,493	3,325	4,988	3,975	1,839	416	37	-
2003.....	37	97	210	566	1,592	3,472	4,823	3,963	1,823	405	26	-
2002.....	38	82	208	567	1,494	3,193	4,587	3,795	1,708	415	30	-
2001.....	30	90	180	513	1,498	3,288	4,484	3,628	1,585	402	37	-
2000.....	29	75	176	585	1,397	3,163	4,143	3,418	1,560	350	26	1
1999.....	31	65	150	498	1,397	2,910	3,877	3,192	1,426	294	29	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Parkinson's disease (G20-G21)												
White, male												
2004.....	30	68	139	386	981	2,072	2,986	2,149	845	146	8	-
2003.....	19	56	133	368	1,043	2,179	2,767	2,097	806	133	12	-
2002.....	25	50	129	395	989	1,963	2,641	2,021	758	137	10	-
2001.....	23	62	135	339	972	2,019	2,622	1,934	725	127	9	-
2000.....	16	49	109	372	890	1,943	2,364	1,804	689	109	7	1
1999.....	24	41	106	314	922	1,842	2,281	1,625	606	102	6	-
White, female												
2004.....	10	29	85	196	512	1,253	2,002	1,826	994	270	29	-
2003.....	18	41	77	198	549	1,293	2,056	1,866	1,017	272	14	-
2002.....	13	32	79	172	505	1,230	1,946	1,774	950	278	20	-
2001.....	7	28	45	174	526	1,269	1,862	1,694	860	275	28	-
2000.....	13	26	67	213	507	1,220	1,779	1,614	871	241	19	-
1999.....	7	24	44	184	475	1,068	1,596	1,567	820	192	23	-
Black, both sexes												
2004.....	5	2	19	38	55	132	158	129	63	18	7	-
2003.....	7	11	22	39	81	134	151	113	53	11	4	-
2002.....	1	8	11	43	71	120	125	105	49	19	1	-
2001.....	2	5	15	25	66	109	129	116	54	18	4	-
2000.....	4	5	9	28	66	118	133	102	47	15	1	-
1999.....	4	7	15	33	64	109	112	94	52	15	2	-
Black, male												
2004.....	5	-	15	22	39	75	81	71	24	10	2	-
2003.....	3	7	14	22	51	81	80	62	25	5	1	-
2002.....	1	3	7	30	48	66	72	58	19	6	-	-
2001.....	2	4	9	15	44	55	72	55	29	4	-	-
2000.....	3	4	5	19	41	80	78	44	24	6	1	-
1999.....	1	3	10	22	37	67	56	47	22	5	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Parkinson's disease (G20-G21)												
Black, female												
2004.....	-	2	4	16	16	57	77	58	39	8	5	-
2003.....	4	4	8	17	30	53	71	51	28	6	3	-
2002.....	-	5	4	13	23	54	53	47	30	13	1	-
2001.....	-	1	6	10	22	54	57	61	25	14	4	-
2000.....	1	1	4	9	25	38	55	58	23	9	-	-
1999.....	3	4	5	11	27	42	56	47	30	10	2	-
American Indian 1/, both sexes												
2004.....	1	1	-	2	1	9	16	8	5	1	-	-
2003.....	-	-	4	3	10	9	17	11	5	1	-	-
2002.....	-	2	-	1	13	17	5	9	5	-	-	-
2001.....	-	-	3	5	5	10	4	7	1	1	-	-
2000.....	-	1	1	3	6	9	11	6	5	2	-	-
1999.....	-	-	-	1	3	5	11	3	2	1	-	-
American Indian 1/, male												
2004.....	1	1	-	1	1	6	11	4	2	-	-	-
2003.....	-	-	3	2	5	5	7	9	1	-	-	-
2002.....	-	2	-	1	9	12	4	6	3	-	-	-
2001.....	-	-	-	2	4	7	3	2	-	-	-	-
2000.....	-	1	-	1	3	5	7	4	4	-	-	-
1999.....	-	-	-	-	-	4	10	1	1	-	-	-
American Indian 1/, female												
2004.....	-	-	-	1	-	3	5	4	3	1	-	-
2003.....	-	-	1	1	5	4	10	2	4	1	-	-
2002.....	-	-	-	-	4	5	1	3	2	-	-	-
2001.....	-	-	3	3	1	3	1	5	1	1	-	-
2000.....	-	-	1	2	3	4	4	2	1	2	-	-
1999.....	-	-	-	1	3	1	1	2	1	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Parkinson's disease (G20-G21)												
Asian or Pacific Islander 2/, both sexes												
2004.....	2	3	3	18	25	60	64	62	30	6	2	-
2003.....	2	4	4	16	23	56	61	70	19	8	1	-
2002.....	-	3	4	15	22	38	54	50	19	6	1	-
2001.....	-	5	3	11	24	39	62	35	15	4	2	-
2000.....	-	1	2	10	13	32	43	41	15	2	-	-
1999.....	-	1	4	7	17	42	44	43	12	3	-	-
Asian or Pacific Islander 2/, male												
2004.....	1	2	2	12	15	36	39	36	17	4	-	-
2003.....	1	2	4	10	16	35	37	42	14	5	1	-
2002.....	-	2	-	7	8	24	29	24	13	4	-	-
2001.....	-	3	1	6	12	23	36	21	5	3	1	-
2000.....	-	1	1	6	6	24	29	27	9	-	-	-
1999.....	-	-	1	2	6	27	20	26	10	3	-	-
Asian or Pacific Islander 2/, female												
2004.....	1	1	1	6	10	24	25	26	13	2	2	-
2003.....	1	2	-	6	7	21	24	28	5	3	-	-
2002.....	-	1	4	8	14	14	25	26	6	2	1	-
2001.....	-	2	2	5	12	16	26	14	10	1	1	-
2000.....	-	-	1	4	7	8	14	14	6	2	-	-
1999.....	-	1	3	5	11	15	24	17	2	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Alzheimer's disease (G30)												
All races, both sexes												
2004.....	65	152	390	1,004	2,637	7,368	14,514	18,327	14,505	5,871	1,087	-
2003.....	57	152	402	979	2,857	7,348	13,809	17,368	13,774	5,704	975	-
2002.....	40	135	375	939	2,663	7,078	13,057	16,267	12,267	5,140	878	-
2001.....	56	165	364	858	2,564	6,568	11,974	14,843	11,168	4,474	796	-
2000.....	47	142	349	868	2,563	6,319	10,934	13,337	10,257	4,014	701	2
1999.....	40	125	323	830	2,374	5,769	10,067	11,862	9,041	3,473	604	-
All races, male												
2004.....	30	66	187	434	1,124	2,767	5,010	5,282	3,087	883	91	-
2003.....	25	76	167	456	1,228	2,855	4,751	4,916	2,933	824	91	-
2002.....	22	55	173	408	1,139	2,793	4,434	4,569	2,570	732	79	-
2001.....	27	61	167	397	1,075	2,616	4,179	4,225	2,312	608	79	-
2000.....	24	60	150	420	1,134	2,433	3,795	3,687	2,101	564	54	1
1999.....	18	68	154	389	1,048	2,332	3,572	3,382	1,861	505	48	-
All races, female												
2004.....	35	86	203	570	1,513	4,601	9,504	13,045	11,418	4,988	996	-
2003.....	32	76	235	523	1,629	4,493	9,058	12,452	10,841	4,880	884	-
2002.....	18	80	202	531	1,524	4,285	8,623	11,698	9,697	4,408	799	-
2001.....	29	104	197	461	1,489	3,952	7,795	10,618	8,856	3,866	717	-
2000.....	23	82	199	448	1,429	3,886	7,139	9,650	8,156	3,450	647	1
1999.....	22	57	169	441	1,326	3,437	6,495	8,480	7,180	2,968	556	-
White, both sexes												
2004.....	61	141	354	893	2,370	6,722	13,408	17,144	13,538	5,449	963	-
2003.....	55	142	376	861	2,591	6,745	12,880	16,313	12,967	5,351	873	-
2002.....	37	127	341	845	2,437	6,554	12,237	15,299	11,560	4,817	783	-
2001.....	54	158	341	767	2,369	6,032	11,194	13,961	10,565	4,182	707	-
2000.....	41	132	318	796	2,343	5,864	10,260	12,562	9,723	3,765	633	1
1999.....	35	113	300	757	2,169	5,363	9,470	11,216	8,611	3,267	552	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Alzheimer's disease (G30)												
White, male												
2004.....	29	63	169	390	1,010	2,541	4,667	4,957	2,871	821	75	-
2003.....	24	72	152	405	1,117	2,637	4,431	4,643	2,755	757	81	-
2002.....	19	51	159	363	1,046	2,600	4,146	4,321	2,418	676	64	-
2001.....	25	56	154	353	996	2,423	3,904	3,968	2,178	576	67	-
2000.....	21	55	138	389	1,043	2,271	3,583	3,466	1,974	525	45	1
1999.....	16	61	144	355	961	2,180	3,384	3,198	1,763	469	41	-
White, female												
2004.....	32	78	185	503	1,360	4,181	8,741	12,187	10,667	4,628	888	-
2003.....	31	70	224	456	1,474	4,108	8,449	11,670	10,212	4,594	792	-
2002.....	18	76	182	482	1,391	3,954	8,091	10,978	9,142	4,141	719	-
2001.....	29	102	187	414	1,373	3,609	7,290	9,993	8,387	3,606	640	-
2000.....	20	77	180	407	1,300	3,593	6,677	9,096	7,749	3,240	588	-
1999.....	19	52	156	402	1,208	3,183	6,086	8,018	6,848	2,798	511	-
Black, both sexes												
2004.....	4	10	27	100	234	567	956	1,022	841	364	106	-
2003.....	2	8	21	100	228	537	819	903	701	307	90	-
2002.....	3	3	29	80	203	473	714	843	615	292	81	-
2001.....	2	6	21	81	173	500	678	784	525	261	80	-
2000.....	4	7	23	67	194	400	608	682	464	219	58	1
1999.....	3	10	19	62	187	364	526	569	367	184	48	-
Black, male												
2004.....	1	2	13	39	99	197	294	270	170	48	13	-
2003.....	1	3	14	45	99	204	279	234	149	50	8	-
2002.....	3	2	12	38	88	179	248	202	119	52	12	-
2001.....	2	5	13	39	72	182	231	219	110	25	11	-
2000.....	2	5	11	30	83	137	185	181	99	36	6	-
1999.....	2	6	9	31	81	132	160	158	75	31	7	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Alzheimer's disease (G30)												
Black, female												
2004.....	3	8	14	61	135	370	662	752	671	316	93	-
2003.....	1	5	7	55	129	333	540	669	552	257	82	-
2002.....	-	1	17	42	115	294	466	641	496	240	69	-
2001.....	-	1	8	42	101	318	447	565	415	236	69	-
2000.....	2	2	12	37	111	263	423	501	365	183	52	1
1999.....	1	4	10	31	106	232	366	411	292	153	41	-
American Indian 1/, both sexes												
2004.....	-	-	2	2	9	13	30	23	18	8	7	-
2003.....	-	1	-	6	9	15	26	34	22	6	3	-
2002.....	-	-	1	7	7	13	29	28	22	5	3	-
2001.....	-	-	1	3	7	8	32	19	14	8	-	-
2000.....	-	1	2	2	12	21	18	19	15	5	-	-
1999.....	-	-	3	5	8	14	15	20	14	5	2	-
American Indian 1/, male												
2004.....	-	-	2	1	5	2	12	5	3	1	1	-
2003.....	-	-	-	1	3	1	8	7	2	-	-	-
2002.....	-	-	1	4	1	5	10	8	5	-	1	-
2001.....	-	-	-	2	2	3	14	6	1	-	-	-
2000.....	-	-	-	1	3	6	7	4	4	1	-	-
1999.....	-	-	1	1	4	6	5	1	5	-	-	-
American Indian 1/, female												
2004.....	-	-	-	1	4	11	18	18	15	7	6	-
2003.....	-	1	-	5	6	14	18	27	20	6	3	-
2002.....	-	-	-	3	6	8	19	20	17	5	2	-
2001.....	-	-	1	1	5	5	18	13	13	8	-	-
2000.....	-	1	2	1	9	15	11	15	11	4	-	-
1999.....	-	-	2	4	4	8	10	19	9	5	2	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Alzheimer's disease (G30)												
Asian or Pacific Islander 2/, both sexes												
2004.....	-	1	7	9	24	66	120	138	108	50	11	-
2003.....	-	1	5	12	29	51	84	118	84	40	9	-
2002.....	-	5	4	7	16	38	77	97	70	26	11	-
2001.....	-	1	1	7	15	28	70	79	64	23	9	-
2000.....	2	2	6	3	14	34	48	74	55	25	10	-
1999.....	2	2	1	6	10	28	56	57	49	17	2	-
Asian or Pacific Islander 2/, male												
2004.....	-	1	3	4	10	27	37	50	43	13	2	-
2003.....	-	1	1	5	9	13	33	32	27	17	2	-
2002.....	-	2	1	3	4	9	30	38	28	4	2	-
2001.....	-	-	-	3	5	8	30	32	23	7	1	-
2000.....	1	-	1	-	5	19	20	36	24	2	3	-
1999.....	-	1	-	2	2	14	23	25	18	5	-	-
Asian or Pacific Islander 2/, female												
2004.....	-	-	4	5	14	39	83	88	65	37	9	-
2003.....	-	-	4	7	20	38	51	86	57	23	7	-
2002.....	-	3	3	4	12	29	47	59	42	22	9	-
2001.....	-	1	1	4	10	20	40	47	41	16	8	-
2000.....	1	2	5	3	9	15	28	38	31	23	7	-
1999.....	2	1	1	4	8	14	33	32	31	12	2	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Multiple sclerosis (G35)												
All races, both sexes												
2004.....	447	526	446	379	318	286	177	98	27	3	-	-
2003.....	452	482	394	351	310	296	161	71	26	6	-	-
2002.....	436	452	401	321	302	252	175	79	21	2	1	-
2001.....	429	421	363	323	291	267	174	58	26	7	-	-
2000.....	433	405	337	294	272	225	155	62	12	-	-	-
1999.....	406	319	290	282	247	236	136	70	13	2	1	-
All races, male												
2004.....	144	189	146	136	110	87	50	21	4	-	-	-
2003.....	154	174	154	122	114	88	46	14	6	1	-	-
2002.....	149	159	171	105	100	84	48	23	4	-	-	-
2001.....	151	147	124	137	89	94	41	14	5	1	-	-
2000.....	157	143	120	103	95	72	41	17	5	-	-	-
1999.....	146	121	92	104	92	70	39	14	2	-	-	-
All races, female												
2004.....	303	337	300	243	208	199	127	77	23	3	-	-
2003.....	298	308	240	229	196	208	115	57	20	5	-	-
2002.....	287	293	230	216	202	168	127	56	17	2	1	-
2001.....	278	274	239	186	202	173	133	44	21	6	-	-
2000.....	276	262	217	191	177	153	114	45	7	-	-	-
1999.....	260	198	198	178	155	166	97	56	11	2	1	-
White, both sexes												
2004.....	397	471	392	347	305	273	173	98	25	2	-	-
2003.....	391	447	357	318	289	278	155	67	26	6	-	-
2002.....	369	411	363	293	288	236	170	76	20	2	1	-
2001.....	379	383	329	298	277	258	166	56	26	7	-	-
2000.....	384	363	297	274	257	219	150	62	11	-	-	-
1999.....	355	299	266	259	231	229	134	68	13	2	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Multiple sclerosis (G35)												
White, male												
2004.....	131	178	134	122	104	79	48	21	4	-	-	-
2003.....	132	165	139	112	109	84	44	12	6	1	-	-
2002.....	131	148	161	100	96	78	48	23	4	-	-	-
2001.....	138	136	115	129	86	91	39	14	5	1	-	-
2000.....	142	127	105	97	94	72	40	17	5	-	-	-
1999.....	128	114	83	100	87	67	39	14	2	-	-	-
White, female												
2004.....	266	293	258	225	201	194	125	77	21	2	-	-
2003.....	259	282	218	206	180	194	111	55	20	5	-	-
2002.....	238	263	202	193	192	158	122	53	16	2	1	-
2001.....	241	247	214	169	191	167	127	42	21	6	-	-
2000.....	242	236	192	177	163	147	110	45	6	-	-	-
1999.....	227	185	183	159	144	162	95	54	11	2	1	-
Black, both sexes												
2004.....	49	54	53	31	12	11	4	-	2	-	-	-
2003.....	58	35	37	32	20	18	6	3	-	-	-	-
2002.....	64	40	35	27	14	15	4	3	1	-	-	-
2001.....	48	37	33	23	14	8	8	2	-	-	-	-
2000.....	45	41	39	19	14	6	5	-	1	-	-	-
1999.....	45	20	24	23	16	6	1	2	-	-	-	-
Black, male												
2004.....	13	10	12	14	6	7	2	-	-	-	-	-
2003.....	22	9	15	10	5	4	2	2	-	-	-	-
2002.....	17	11	9	4	4	6	-	-	-	-	-	-
2001.....	12	10	8	8	3	3	2	-	-	-	-	-
2000.....	14	16	15	6	1	-	1	-	-	-	-	-
1999.....	18	7	9	4	5	3	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Multiple sclerosis (G35)												
Asian or Pacific Islander 2/, both sexes												
2004.....	1	-	-	-	1	2	-	-	-	1	-	-
2003.....	3	-	-	-	1	-	-	1	-	-	-	-
2002.....	1	-	1	1	-	1	1	-	-	-	-	-
2001.....	1	-	-	2	-	1	-	-	-	-	-	-
2000.....	1	1	1	-	1	-	-	-	-	-	-	-
1999.....	4	-	-	-	-	1	1	-	-	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	-	-	-	-	-	1	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	1	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	1	-	-	-	1	1	-	-	-	1	-	-
2003.....	3	-	-	-	1	-	-	1	-	-	-	-
2002.....	1	-	1	-	-	1	1	-	-	-	-	-
2001.....	1	-	-	2	-	1	-	-	-	-	-	-
2000.....	1	1	1	-	1	-	-	-	-	-	-	-
1999.....	4	-	-	-	-	1	1	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Epilepsy (G40-G41)												
All races, both sexes												
2004.....	114	104	73	67	51	84	63	43	27	5	-	-
2003.....	123	103	72	70	73	76	81	35	28	8	2	-
2002.....	108	71	75	71	76	96	75	51	22	7	1	-
2001.....	108	83	72	64	66	83	62	51	11	7	2	-
2000.....	107	88	66	64	89	73	80	53	28	4	1	-
1999.....	98	69	65	74	87	78	74	45	22	8	1	-
All races, male												
2004.....	81	56	28	34	17	37	20	20	9	-	-	-
2003.....	76	56	32	28	33	32	38	13	6	1	1	-
2002.....	69	32	41	35	34	38	30	25	5	2	-	-
2001.....	61	49	36	34	32	30	15	19	4	2	1	-
2000.....	60	52	39	33	40	29	18	20	8	1	1	-
1999.....	51	35	35	40	40	27	24	18	7	2	-	-
All races, female												
2004.....	33	48	45	33	34	47	43	23	18	5	-	-
2003.....	47	47	40	42	40	44	43	22	22	7	1	-
2002.....	39	39	34	36	42	58	45	26	17	5	1	-
2001.....	47	34	36	30	34	53	47	32	7	5	1	-
2000.....	47	36	27	31	49	44	62	33	20	3	-	-
1999.....	47	34	30	34	47	51	50	27	15	6	1	-
White, both sexes												
2004.....	88	72	51	52	40	61	55	34	24	2	-	-
2003.....	90	71	53	48	54	53	64	28	24	7	1	-
2002.....	71	54	58	53	54	71	62	45	18	2	1	-
2001.....	83	55	54	52	48	59	47	48	9	6	-	-
2000.....	82	64	46	49	66	50	66	45	18	2	-	-
1999.....	68	50	36	52	65	59	62	38	20	6	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Epilepsy (G40-G41)												
White, male												
2004.....	61	33	23	25	14	26	17	17	8	-	-	-
2003.....	53	38	24	14	22	21	27	11	4	1	1	-
2002.....	48	25	30	27	24	25	22	23	4	1	-	-
2001.....	47	33	28	27	24	20	12	18	3	2	-	-
2000.....	48	40	25	26	27	16	14	15	5	1	-	-
1999.....	35	22	19	28	33	19	21	17	6	2	-	-
White, female												
2004.....	27	39	28	27	26	35	38	17	16	2	-	-
2003.....	37	33	29	34	32	32	37	17	20	6	-	-
2002.....	23	29	28	26	30	46	40	22	14	1	1	-
2001.....	36	22	26	25	24	39	35	30	6	4	-	-
2000.....	34	24	21	23	39	34	52	30	13	1	-	-
1999.....	33	28	17	24	32	40	41	21	14	4	1	-
Black, both sexes												
2004.....	23	29	19	14	11	21	7	8	2	3	-	-
2003.....	30	32	17	20	17	20	14	7	4	1	1	-
2002.....	34	16	17	17	20	24	11	6	3	5	-	-
2001.....	23	25	16	11	16	21	15	3	1	1	2	-
2000.....	24	22	19	14	22	22	14	7	10	2	1	-
1999.....	30	17	25	22	22	19	12	7	2	2	-	-
Black, male												
2004.....	18	21	5	8	3	9	3	3	-	-	-	-
2003.....	22	18	7	12	11	10	8	2	2	-	-	-
2002.....	19	7	11	8	9	12	6	2	1	1	-	-
2001.....	13	15	7	6	7	9	3	1	1	-	1	-
2000.....	12	11	13	7	12	12	4	4	3	-	1	-
1999.....	16	11	15	12	7	8	3	1	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Transient cerebral ischemic attacks and related syndromes (G45)

All races, both sexes

2004.....	-	3	4	4	8	20	31	38	38	23	4	-
2003.....	-	2	6	4	13	22	36	76	60	36	6	-
2002.....	1	2	3	11	14	28	42	57	48	28	2	-
2001.....	4	2	5	13	11	31	35	70	60	26	11	-
2000.....	2	6	2	10	13	36	50	62	62	34	6	-
1999.....	5	1	7	12	16	30	50	78	76	32	4	-

All races, male

2004.....	-	-	3	1	6	9	6	8	12	3	1	-
2003.....	-	1	3	1	7	5	15	17	14	8	1	-
2002.....	-	1	2	7	7	14	21	15	8	2	-	-
2001.....	2	1	4	4	6	16	16	31	20	2	2	-
2000.....	2	1	2	4	4	21	16	17	13	4	2	-
1999.....	2	-	6	5	5	21	22	22	18	5	-	-

All races, female

2004.....	-	3	1	3	2	11	25	30	26	20	3	-
2003.....	-	1	3	3	6	17	21	59	46	28	5	-
2002.....	1	1	1	4	7	14	21	42	40	26	2	-
2001.....	2	1	1	9	5	15	19	39	40	24	9	-
2000.....	-	5	-	6	9	15	34	45	49	30	4	-
1999.....	3	1	1	7	11	9	28	56	58	27	4	-

White, both sexes

2004.....	-	3	4	4	7	18	28	36	35	21	3	-
2003.....	-	2	5	3	12	21	31	72	58	35	6	-
2002.....	1	2	1	8	14	26	40	53	44	27	2	-
2001.....	2	2	4	10	10	30	35	65	56	24	10	-
2000.....	2	4	2	8	12	34	48	61	56	31	6	-
1999.....	4	1	5	11	15	27	49	76	74	31	4	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Transient cerebral ischemic attacks and related syndromes (G45)

White, male

2004.....	-	-	3	1	5	8	6	8	12	3	1	-
2003.....	-	1	3	1	7	5	14	16	13	8	1	-
2002.....	-	1	1	5	7	13	19	14	7	2	-	-
2001.....	1	1	3	3	6	16	16	30	18	2	1	-
2000.....	2	-	2	2	4	20	16	17	10	4	2	-
1999.....	1	-	5	4	5	19	22	22	16	4	-	-

White, female

2004.....	-	3	1	3	2	10	22	28	23	18	2	-
2003.....	-	1	2	2	5	16	17	56	45	27	5	-
2002.....	1	1	-	3	7	13	21	39	37	25	2	-
2001.....	1	1	1	7	4	14	19	35	38	22	9	-
2000.....	-	4	-	6	8	14	32	44	46	27	4	-
1999.....	3	1	-	7	10	8	27	54	58	27	4	-

Black, both sexes

2004.....	-	-	-	-	1	1	2	1	3	1	1	-
2003.....	-	-	1	1	1	-	4	4	2	1	-	-
2002.....	-	-	1	3	-	2	2	3	4	1	-	-
2001.....	2	-	1	3	1	1	-	5	4	2	1	-
2000.....	-	2	-	2	1	2	2	1	4	3	-	-
1999.....	1	-	2	1	1	3	1	2	1	1	-	-

Black, male

2004.....	-	-	-	-	1	1	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	1	1	1	-	-	-
2002.....	-	-	-	2	-	1	2	1	1	-	-	-
2001.....	1	-	1	1	-	-	-	1	2	-	1	-
2000.....	-	1	-	2	-	1	-	-	1	-	-	-
1999.....	1	-	1	1	-	2	-	-	1	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Infantile cerebral palsy (G80)												
All races, both sexes												
2004.....	84	74	58	47	54	48	19	16	3	-	-	1
2003.....	83	64	60	34	40	35	32	17	2	-	-	-
2002.....	62	49	56	38	40	35	30	13	5	-	-	-
2001.....	60	56	39	30	51	42	20	12	2	4	-	-
2000.....	45	41	38	48	39	29	33	5	1	-	-	-
1999.....	39	32	31	33	35	30	22	17	2	1	-	-
All races, male												
2004.....	45	36	41	31	28	28	11	10	-	-	-	1
2003.....	51	33	34	19	23	22	16	8	2	-	-	-
2002.....	35	27	33	21	20	18	14	4	3	-	-	-
2001.....	36	25	22	18	35	23	11	6	2	2	-	-
2000.....	29	20	18	29	27	13	11	-	1	-	-	-
1999.....	26	20	22	22	17	16	10	5	1	-	-	-
All races, female												
2004.....	39	38	17	16	26	20	8	6	3	-	-	-
2003.....	32	31	26	15	17	13	16	9	-	-	-	-
2002.....	27	22	23	17	20	17	16	9	2	-	-	-
2001.....	24	31	17	12	16	19	9	6	-	2	-	-
2000.....	16	21	20	19	12	16	22	5	-	-	-	-
1999.....	13	12	9	11	18	14	12	12	1	1	-	-
White, both sexes												
2004.....	70	68	53	41	50	44	19	16	3	-	-	-
2003.....	68	57	55	33	37	34	29	17	2	-	-	-
2002.....	52	45	53	35	39	33	28	12	4	-	-	-
2001.....	51	51	37	27	48	41	20	12	2	4	-	-
2000.....	42	37	35	44	36	24	31	5	1	-	-	-
1999.....	32	30	29	31	32	29	22	17	2	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Infantile cerebral palsy (G80)												
White, male												
2004.....	38	35	36	27	26	24	11	10	-	-	-	-
2003.....	41	31	31	18	22	21	14	8	2	-	-	-
2002.....	30	24	30	21	20	17	13	4	2	-	-	-
2001.....	31	23	20	15	32	22	11	6	2	2	-	-
2000.....	28	19	17	27	24	9	11	-	1	-	-	-
1999.....	20	20	21	20	16	15	10	5	1	-	-	-
White, female												
2004.....	32	33	17	14	24	20	8	6	3	-	-	-
2003.....	27	26	24	15	15	13	15	9	-	-	-	-
2002.....	22	21	23	14	19	16	15	8	2	-	-	-
2001.....	20	28	17	12	16	19	9	6	-	2	-	-
2000.....	14	18	18	17	12	15	20	5	-	-	-	-
1999.....	12	10	8	11	16	14	12	12	1	1	-	-
Black, both sexes												
2004.....	14	5	5	6	4	4	-	-	-	-	-	-
2003.....	15	6	5	1	3	1	3	-	-	-	-	-
2002.....	9	3	3	3	1	2	2	1	1	-	-	-
2001.....	9	4	2	3	3	1	-	-	-	-	-	-
2000.....	3	4	3	4	3	5	2	-	-	-	-	-
1999.....	7	2	2	2	3	1	-	-	-	-	-	-
Black, male												
2004.....	7	1	5	4	2	4	-	-	-	-	-	-
2003.....	10	2	3	1	1	1	2	-	-	-	-	-
2002.....	5	2	3	-	-	1	1	-	1	-	-	-
2001.....	5	1	2	3	3	1	-	-	-	-	-	-
2000.....	1	1	1	2	3	4	-	-	-	4	-	-
1999.....	6	-	1	2	1	1	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

All other diseases of nervous system (G10-G12,G23-G25,G31,G36-G37,G43-G44,G47-G72,G81-G98)

All races, both sexes

2004.....	1,283	1,630	1,834	1,984	2,346	2,692	2,399	1,542	719	198	41	-
2003.....	1,233	1,537	1,835	2,008	2,489	2,816	2,378	1,633	754	212	42	-
2002.....	1,195	1,448	1,669	1,985	2,411	2,757	2,400	1,523	679	228	40	-
2001.....	1,124	1,300	1,529	1,970	2,425	2,695	2,261	1,387	671	196	32	-
2000.....	1,175	1,254	1,503	1,829	2,455	2,664	2,206	1,441	679	192	38	-
1999.....	1,043	1,186	1,397	1,889	2,405	2,689	2,118	1,420	615	160	30	-

All races, male

2004.....	747	926	999	1,076	1,248	1,376	1,088	663	234	40	4	-
2003.....	736	873	1,004	1,047	1,274	1,408	1,158	685	263	57	6	-
2002.....	670	807	912	1,073	1,269	1,376	1,130	610	240	59	9	-
2001.....	631	711	807	1,085	1,282	1,357	1,084	601	246	53	5	-
2000.....	665	697	836	1,007	1,308	1,327	1,071	581	244	47	11	-
1999.....	598	661	778	1,023	1,248	1,374	1,019	609	195	32	3	-

All races, female

2004.....	536	704	835	908	1,098	1,316	1,311	879	485	158	37	-
2003.....	497	664	831	961	1,215	1,408	1,220	948	491	155	36	-
2002.....	525	641	757	912	1,142	1,381	1,270	913	439	169	31	-
2001.....	493	589	722	885	1,143	1,338	1,177	786	425	143	27	-
2000.....	510	557	667	822	1,147	1,337	1,135	860	435	145	27	-
1999.....	445	525	619	866	1,157	1,315	1,099	811	420	128	27	-

White, both sexes

2004.....	1,041	1,357	1,576	1,702	2,077	2,422	2,180	1,415	673	175	30	-
2003.....	991	1,297	1,554	1,748	2,206	2,539	2,183	1,499	694	194	37	-
2002.....	972	1,218	1,436	1,726	2,135	2,480	2,211	1,401	620	216	36	-
2001.....	910	1,067	1,293	1,712	2,160	2,426	2,066	1,285	602	182	22	-
2000.....	972	1,062	1,286	1,582	2,179	2,438	2,034	1,310	634	180	32	-
1999.....	860	988	1,188	1,645	2,138	2,437	1,973	1,304	572	141	26	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

All other diseases of nervous system (G10-G12,G23-G25,G31,G36-G37,G43-G44,G47-G72,G81-G98)

White, male

2004.....	615	776	869	938	1,130	1,250	1,000	615	223	39	3	-
2003.....	613	751	856	914	1,155	1,287	1,068	636	247	52	6	-
2002.....	548	686	795	931	1,131	1,256	1,052	569	217	56	7	-
2001.....	513	601	693	954	1,155	1,230	1,007	555	218	48	4	-
2000.....	568	590	717	889	1,176	1,224	991	529	228	45	8	-
1999.....	501	567	669	903	1,125	1,259	963	564	180	26	2	-

White, female

2004.....	426	581	707	764	947	1,172	1,180	800	450	136	27	-
2003.....	378	546	698	834	1,051	1,252	1,115	863	447	142	31	-
2002.....	424	532	641	795	1,004	1,224	1,159	832	403	160	29	-
2001.....	397	466	600	758	1,005	1,196	1,059	730	384	134	18	-
2000.....	404	472	569	693	1,003	1,214	1,043	781	406	135	24	-
1999.....	359	421	519	742	1,013	1,178	1,010	740	392	115	24	-

Black, both sexes

2004.....	211	238	231	237	223	233	180	108	41	18	9	-
2003.....	215	206	235	232	232	218	157	113	48	14	5	-
2002.....	194	194	208	221	236	219	163	101	55	10	4	-
2001.....	194	207	198	225	228	221	169	77	60	13	9	-
2000.....	185	169	183	219	228	198	144	109	40	9	6	-
1999.....	166	172	177	204	232	213	120	99	40	16	4	-

Black, male

2004.....	120	133	114	112	101	103	70	38	11	1	1	-
2003.....	106	102	124	119	99	94	64	36	9	3	-	-
2002.....	101	105	103	117	120	92	65	30	20	2	2	-
2001.....	105	97	95	111	106	107	66	29	22	4	-	-
2000.....	88	97	97	106	112	95	67	39	12	1	3	-
1999.....	90	77	89	95	100	99	43	35	14	6	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of nervous system (G10-G12,G23-G25,G31,G36-G37,G43-G44,G47-G72,G81-G98)												
Black, female												
2004.....	91	105	117	125	122	130	110	70	30	17	8	-
2003.....	109	104	111	113	133	124	93	77	39	11	5	-
2002.....	93	89	105	104	116	127	98	71	35	8	2	-
2001.....	89	110	103	114	122	114	103	48	38	9	9	-
2000.....	97	72	86	113	116	103	77	70	28	8	3	-
1999.....	76	95	88	109	132	114	77	64	26	10	3	-
American Indian 1/, both sexes												
2004.....	17	7	7	9	4	9	6	2	-	-	1	-
2003.....	11	4	12	5	8	11	9	1	2	-	-	-
2002.....	7	6	9	9	10	10	6	4	-	-	-	-
2001.....	7	10	10	9	4	7	5	2	1	1	-	-
2000.....	4	7	7	10	9	5	4	4	-	-	-	-
1999.....	6	9	9	9	8	5	4	3	1	1	-	-
American Indian 1/, male												
2004.....	6	6	6	5	1	6	2	1	-	-	-	-
2003.....	8	2	7	2	4	5	5	-	1	-	-	-
2002.....	5	3	4	6	5	6	2	3	-	-	-	-
2001.....	5	6	4	5	3	2	-	-	-	1	-	-
2000.....	1	1	3	4	4	1	2	2	-	-	-	-
1999.....	3	7	6	7	5	1	1	1	-	-	-	-
American Indian 1/, female												
2004.....	11	1	1	4	3	3	4	1	-	-	1	-
2003.....	3	2	5	3	4	6	4	1	1	-	-	-
2002.....	2	3	5	3	5	4	4	1	-	-	-	-
2001.....	2	4	6	4	1	5	5	2	1	-	-	-
2000.....	3	6	4	6	5	4	2	2	-	-	-	-
1999.....	3	2	3	2	3	4	3	2	1	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

All other diseases of nervous system (G10-G12,G23-G25,G31,G36-G37,G43-G44,G47-G72,G81-G98)

Asian or Pacific Islander 2/, both sexes

2004.....	14	28	20	36	42	28	33	17	5	5	1	-
2003.....	16	30	34	23	43	48	29	20	10	4	-	-
2002.....	22	30	16	29	30	48	20	17	4	2	-	-
2001.....	13	16	28	24	33	41	21	23	8	-	1	-
2000.....	14	16	27	18	39	23	24	18	5	3	-	-
1999.....	11	17	23	31	27	34	21	14	2	2	-	-

Asian or Pacific Islander 2/, male

2004.....	6	11	10	21	16	17	16	9	-	-	-	-
2003.....	9	18	17	12	16	22	21	13	6	2	-	-
2002.....	16	13	10	19	13	22	11	8	3	1	-	-
2001.....	8	7	15	15	18	18	11	17	6	-	1	-
2000.....	8	9	19	8	16	7	11	11	4	1	-	-
1999.....	4	10	14	18	18	15	12	9	1	-	-	-

Asian or Pacific Islander 2/, female

2004.....	8	17	10	15	26	11	17	8	5	5	1	-
2003.....	7	12	17	11	27	26	8	7	4	2	-	-
2002.....	6	17	6	10	17	26	9	9	1	1	-	-
2001.....	5	9	13	9	15	23	10	6	2	-	-	-
2000.....	6	7	8	10	23	16	13	7	1	2	-	-
1999.....	7	7	9	13	9	19	9	5	1	2	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VII. Diseases of the eye and adnexa (H00-H57)												
All races, both sexes												
2004.....	2	-	2	2	3	6	3	6	3	6	2	-
2003.....	-	2	1	-	-	1	3	9	9	5	3	-
2002.....	-	2	3	1	4	2	6	7	3	3	-	-
2001.....	-	-	2	-	1	3	-	8	5	9	2	-
2000.....	2	1	-	5	5	4	7	9	11	4	1	-
1999.....	2	1	1	2	3	5	1	12	3	6	2	-
All races, male												
2004.....	1	-	-	2	2	3	-	-	1	1	1	-
2003.....	-	2	1	-	-	-	2	5	1	1	-	-
2002.....	-	1	2	1	2	2	2	1	-	-	-	-
2001.....	-	-	-	-	-	-	-	3	2	3	-	-
2000.....	1	1	-	4	5	3	2	4	3	1	-	-
1999.....	1	-	1	1	3	2	1	2	2	1	1	-
All races, female												
2004.....	1	-	2	-	1	3	3	6	2	5	1	-
2003.....	-	-	-	-	-	1	1	4	8	4	3	-
2002.....	-	1	1	-	2	-	4	6	3	3	-	-
2001.....	-	-	2	-	1	3	-	5	3	6	2	-
2000.....	1	-	-	1	-	1	5	5	8	3	1	-
1999.....	1	1	-	1	-	3	-	10	1	5	1	-
White, both sexes												
2004.....	2	-	1	2	2	6	2	6	3	6	2	-
2003.....	-	2	1	-	-	1	2	9	9	5	3	-
2002.....	-	2	2	1	3	1	6	7	3	3	-	-
2001.....	-	-	1	-	1	2	-	8	4	9	1	-
2000.....	2	1	-	5	4	3	4	6	9	3	1	-
1999.....	-	-	-	2	3	4	1	11	2	4	2	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VII. Diseases of the eye and adnexa (H00-H57)												
White, male												
2004.....	1	-	-	2	1	3	-	-	1	1	1	-
2003.....	-	2	1	-	-	-	1	5	1	1	-	-
2002.....	-	1	1	1	2	1	2	1	-	-	-	-
2001.....	-	-	-	-	-	-	-	3	2	3	-	-
2000.....	1	1	-	4	4	2	1	3	2	1	-	-
1999.....	-	-	-	1	3	2	1	2	1	-	1	-
White, female												
2004.....	1	-	1	-	1	3	2	6	2	5	1	-
2003.....	-	-	-	-	-	1	1	4	8	4	3	-
2002.....	-	1	1	-	1	-	4	6	3	3	-	-
2001.....	-	-	1	-	1	2	-	5	2	6	1	-
2000.....	1	-	-	1	-	1	3	3	7	2	1	-
1999.....	-	-	-	1	-	2	-	9	1	4	1	-
Black, both sexes												
2004.....	-	-	1	-	1	-	1	-	-	-	-	-
2003.....	-	-	-	-	-	-	1	-	-	-	-	-
2002.....	-	-	-	-	1	1	-	-	-	-	-	-
2001.....	-	-	1	-	-	1	-	-	1	-	1	-
2000.....	-	-	-	-	1	1	3	1	2	-	-	-
1999.....	1	1	1	-	-	1	-	1	1	2	-	-
Black, male												
2004.....	-	-	-	-	1	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	1	-	-	-	-	-
2002.....	-	-	-	-	-	1	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	1	1	1	-	1	-	-	-
1999.....	1	-	1	-	-	-	-	-	1	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VIII. Diseases of the ear and mastoid process (H60-H93)												
All races, both sexes												
2004.....	2	4	4	3	4	7	5	5	7	1	-	-
2003.....	3	2	10	4	8	2	7	3	3	1	2	-
2002.....	4	4	4	2	6	5	5	3	6	-	-	-
2001.....	1	7	5	8	4	5	6	5	4	2	3	-
2000.....	3	2	6	3	6	4	3	4	4	-	1	-
1999.....	4	6	3	4	3	8	4	4	2	1	1	-
All races, male												
2004.....	1	2	2	1	3	3	2	1	2	-	-	-
2003.....	1	-	3	-	3	1	5	1	1	-	-	-
2002.....	4	1	1	1	5	3	3	2	2	-	-	-
2001.....	-	1	3	3	2	2	2	3	1	1	-	-
2000.....	3	2	2	3	2	2	1	1	1	-	1	-
1999.....	3	1	1	2	3	3	4	1	2	-	-	-
All races, female												
2004.....	1	2	2	2	1	4	3	4	5	1	-	-
2003.....	2	2	7	4	5	1	2	2	2	1	2	-
2002.....	-	3	3	1	1	2	2	1	4	-	-	-
2001.....	1	6	2	5	2	3	4	2	3	1	3	-
2000.....	-	-	4	-	4	2	2	3	3	-	-	-
1999.....	1	5	2	2	-	5	-	3	-	1	1	-
White, both sexes												
2004.....	1	4	3	2	3	6	5	4	5	1	-	-
2003.....	2	2	9	2	5	2	6	3	3	1	2	-
2002.....	4	2	4	2	6	4	4	3	5	-	-	-
2001.....	1	7	5	6	4	4	5	4	4	2	3	-
2000.....	3	1	5	2	5	4	2	4	4	-	1	-
1999.....	4	3	3	3	3	5	4	4	1	-	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VIII. Diseases of the ear and mastoid process (H60-H93)												
White, male												
2004.....	-	2	2	-	2	2	2	1	2	-	-	-
2003.....	1	-	3	-	1	1	5	1	1	-	-	-
2002.....	4	1	1	1	5	3	2	2	2	-	-	-
2001.....	-	1	3	3	2	1	1	3	1	1	-	-
2000.....	3	1	1	2	2	2	-	1	1	-	1	-
1999.....	3	1	1	2	3	3	4	1	1	-	-	-
White, female												
2004.....	1	2	1	2	1	4	3	3	3	1	-	-
2003.....	1	2	6	2	4	1	1	2	2	1	2	-
2002.....	-	1	3	1	1	1	2	1	3	-	-	-
2001.....	1	6	2	3	2	3	4	1	3	1	3	-
2000.....	-	-	4	-	3	2	2	3	3	-	-	-
1999.....	1	2	2	1	-	2	-	3	-	-	1	-
Black, both sexes												
2004.....	1	-	1	1	-	1	-	1	2	-	-	-
2003.....	1	-	1	2	3	-	1	-	-	-	-	-
2002.....	-	2	-	-	-	-	1	-	1	-	-	-
2001.....	-	-	-	2	-	-	1	1	-	-	-	-
2000.....	-	1	1	1	-	-	1	-	-	-	-	-
1999.....	-	2	-	1	-	2	-	-	1	1	-	-
Black, male												
2004.....	1	-	-	1	-	1	-	-	-	-	-	-
2003.....	-	-	-	-	2	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	1	-	-	-	-	-
2001.....	-	-	-	-	-	-	1	-	-	-	-	-
2000.....	-	1	1	1	-	-	1	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

VIII. Diseases of the ear and mastoid process (H60-H93)

Asian or Pacific Islander 2/, both sexes

2004.....	-	-	-	-	1	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	1	-	-	-	-	-	-
2001.....	-	-	-	-	-	1	-	-	-	-	-	-
2000.....	-	-	-	-	1	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	1	-	-	-	-	-	-

Asian or Pacific Islander 2/, male

2004.....	-	-	-	-	1	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	1	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-

Asian or Pacific Islander 2/, female

2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	1	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	1	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	1	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
IX. Diseases of the circulatory system (I00-I99)												
All races, both sexes												
2004.....	27,415	35,122	43,485	53,829	75,659	112,752	152,450	153,778	113,145	45,667	10,087	45
2003.....	27,470	35,737	44,329	56,260	81,704	121,792	158,727	161,456	117,566	47,908	10,652	38
2002.....	27,364	35,306	43,719	58,053	86,715	126,712	161,956	163,997	118,216	47,754	10,565	51
2001.....	26,914	33,607	43,180	59,432	89,818	131,203	161,975	164,568	116,511	47,372	10,628	73
2000.....	26,064	33,685	44,313	62,095	94,887	135,869	161,494	165,884	117,435	47,576	10,639	62
1999.....	25,334	33,552	44,726	65,087	99,033	139,178	164,491	166,579	116,579	47,772	10,552	50
All races, male												
2004.....	19,103	23,878	28,487	33,268	43,239	59,144	69,851	58,637	33,212	9,739	1,516	38
2003.....	19,210	24,271	28,853	34,820	47,040	63,798	72,507	60,835	33,874	9,957	1,513	29
2002.....	19,077	23,889	28,457	35,763	49,879	65,967	73,286	61,517	33,525	9,885	1,522	36
2001.....	18,577	22,638	27,986	36,395	51,766	68,226	73,051	60,078	32,224	9,584	1,497	54
2000.....	18,141	22,943	28,744	37,891	55,009	70,441	72,828	60,554	32,176	9,662	1,539	43
1999.....	17,691	22,759	28,990	40,141	57,113	72,277	73,753	60,185	31,639	9,577	1,510	38
All races, female												
2004.....	8,312	11,244	14,998	20,561	32,420	53,608	82,599	95,141	79,933	35,928	8,571	7
2003.....	8,260	11,466	15,476	21,440	34,664	57,994	86,220	100,621	83,692	37,951	9,139	9
2002.....	8,287	11,417	15,262	22,290	36,836	60,745	88,670	102,480	84,691	37,869	9,043	15
2001.....	8,337	10,969	15,194	23,037	38,052	62,977	88,924	104,490	84,287	37,788	9,131	19
2000.....	7,923	10,742	15,569	24,204	39,878	65,428	88,666	105,330	85,259	37,914	9,100	19
1999.....	7,643	10,793	15,736	24,946	41,920	66,901	90,738	106,394	84,940	38,195	9,042	12
White, both sexes												
2004.....	20,018	26,795	34,387	43,330	63,329	97,649	136,485	140,514	103,684	41,706	8,828	30
2003.....	20,043	27,596	34,821	45,274	68,461	105,994	142,327	147,711	108,064	43,885	9,251	31
2002.....	20,244	27,382	34,509	47,058	73,411	110,427	145,548	150,167	108,604	43,692	9,225	35
2001.....	19,968	25,995	34,068	48,048	76,265	114,676	145,872	150,648	107,442	43,347	9,289	45
2000.....	19,631	26,210	35,058	50,934	80,795	119,172	145,815	151,833	108,448	43,459	9,378	45
1999.....	19,161	26,013	35,602	53,728	84,428	122,357	148,863	152,398	107,858	43,794	9,374	33

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
IX. Diseases of the circulatory system (I00-I99)												
White, male												
2004.....	14,500	18,704	22,988	27,355	36,933	52,086	63,338	54,100	30,513	8,842	1,290	26
2003.....	14,551	19,268	23,262	28,696	40,156	56,410	65,605	56,167	31,307	9,053	1,288	22
2002.....	14,636	19,020	23,001	29,653	43,113	58,468	66,580	56,853	30,785	8,999	1,275	26
2001.....	14,327	18,004	22,670	30,219	44,742	60,547	66,436	55,398	29,699	8,687	1,273	34
2000.....	14,128	18,298	23,363	31,878	47,804	62,685	66,430	55,789	29,650	8,661	1,320	34
1999.....	13,808	18,143	23,610	33,910	49,671	64,468	67,507	55,221	29,099	8,681	1,312	25
White, female												
2004.....	5,518	8,091	11,399	15,975	26,396	45,563	73,147	86,414	73,171	32,864	7,538	4
2003.....	5,492	8,328	11,559	16,578	28,305	49,584	76,722	91,544	76,757	34,832	7,963	9
2002.....	5,608	8,362	11,508	17,405	30,298	51,959	78,968	93,314	77,819	34,693	7,950	9
2001.....	5,641	7,991	11,398	17,829	31,523	54,129	79,436	95,250	77,743	34,660	8,016	11
2000.....	5,503	7,912	11,695	19,056	32,991	56,487	79,385	96,044	78,798	34,798	8,058	11
1999.....	5,353	7,870	11,992	19,818	34,757	57,889	81,356	97,177	78,759	35,113	8,062	8
Black, both sexes												
2004.....	6,622	7,353	8,010	8,990	10,573	12,672	13,055	10,580	7,735	3,288	1,083	14
2003.....	6,665	7,192	8,295	9,436	11,324	13,302	13,485	11,191	7,811	3,418	1,216	5
2002.....	6,407	7,041	8,097	9,488	11,319	13,719	13,673	11,397	7,953	3,437	1,161	15
2001.....	6,219	6,738	8,038	9,859	11,576	14,031	13,432	11,604	7,683	3,480	1,159	26
2000.....	5,735	6,650	8,124	9,691	12,235	14,129	13,224	11,911	7,581	3,507	1,133	17
1999.....	5,544	6,710	8,013	9,954	12,647	14,366	13,170	11,961	7,377	3,498	1,034	16
Black, male												
2004.....	4,067	4,531	4,799	5,020	5,368	5,801	5,149	3,428	2,007	652	177	11
2003.....	4,125	4,375	4,832	5,204	5,816	6,146	5,494	3,567	1,895	678	171	5
2002.....	3,943	4,290	4,750	5,216	5,741	6,131	5,393	3,587	2,055	661	190	9
2001.....	3,777	4,043	4,636	5,315	5,937	6,371	5,357	3,677	1,942	706	191	19
2000.....	3,535	4,103	4,640	5,140	6,130	6,376	5,233	3,792	1,950	762	188	9
1999.....	3,462	4,060	4,698	5,428	6,348	6,446	5,079	3,907	1,958	745	169	12

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
IX. Diseases of the circulatory system (I00-I99)												
Black, female												
2004.....	2,555	2,822	3,211	3,970	5,205	6,871	7,906	7,152	5,728	2,636	906	3
2003.....	2,540	2,817	3,463	4,232	5,508	7,156	7,991	7,624	5,916	2,740	1,045	-
2002.....	2,464	2,751	3,347	4,272	5,578	7,588	8,280	7,810	5,898	2,776	971	6
2001.....	2,442	2,695	3,402	4,544	5,639	7,660	8,075	7,927	5,741	2,774	968	7
2000.....	2,200	2,547	3,484	4,551	6,105	7,753	7,991	8,119	5,631	2,745	945	8
1999.....	2,082	2,650	3,315	4,526	6,299	7,920	8,091	8,054	5,419	2,753	865	4
American Indian 1/, both sexes												
2004.....	202	263	280	363	356	403	458	373	204	100	28	-
2003.....	219	250	345	367	393	423	420	337	215	68	34	2
2002.....	193	237	285	340	386	437	411	311	205	90	25	1
2001.....	183	232	262	351	395	394	391	319	174	75	26	-
2000.....	180	224	276	370	355	437	391	322	212	90	18	-
1999.....	161	214	259	330	401	449	381	324	188	66	29	-
American Indian 1/, male												
2004.....	123	174	177	218	195	212	208	151	58	24	5	-
2003.....	147	167	203	203	230	212	189	137	83	18	13	2
2002.....	139	147	186	194	217	227	171	143	58	21	8	1
2001.....	120	154	171	193	232	191	164	128	57	18	3	-
2000.....	118	142	166	217	210	221	178	123	60	23	2	-
1999.....	110	136	144	205	228	241	155	110	55	17	7	-
American Indian 1/, female												
2004.....	79	89	103	145	161	191	250	222	146	76	23	-
2003.....	72	83	142	164	163	211	231	200	132	50	21	-
2002.....	54	90	99	146	169	210	240	168	147	69	17	-
2001.....	63	78	91	158	163	203	227	191	117	57	23	-
2000.....	62	82	110	153	145	216	213	199	152	67	16	-
1999.....	51	78	115	125	173	208	226	214	133	49	22	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
IX. Diseases of the circulatory system (I00-I99)												
Asian or Pacific Islander 2/, both sexes												
2004.....	573	711	808	1,146	1,401	2,028	2,452	2,311	1,522	573	148	1
2003.....	543	699	868	1,183	1,526	2,073	2,495	2,217	1,476	537	151	-
2002.....	520	646	828	1,167	1,599	2,129	2,324	2,122	1,454	535	154	-
2001.....	544	642	812	1,174	1,582	2,102	2,280	1,997	1,212	470	154	2
2000.....	518	601	855	1,100	1,502	2,131	2,064	1,818	1,194	520	110	-
1999.....	468	615	852	1,075	1,557	2,006	2,077	1,896	1,156	414	115	1
Asian or Pacific Islander 2/, male												
2004.....	413	469	523	675	743	1,045	1,156	958	634	221	44	1
2003.....	387	461	556	717	838	1,030	1,219	964	589	208	41	-
2002.....	359	432	520	700	808	1,141	1,142	934	627	204	49	-
2001.....	353	437	509	668	855	1,117	1,094	875	526	173	30	1
2000.....	360	400	575	656	865	1,159	987	850	516	216	29	-
1999.....	311	420	538	598	866	1,122	1,012	947	527	134	22	1
Asian or Pacific Islander 2/, female												
2004.....	160	242	285	471	658	983	1,296	1,353	888	352	104	-
2003.....	156	238	312	466	688	1,043	1,276	1,253	887	329	110	-
2002.....	161	214	308	467	791	988	1,182	1,188	827	331	105	-
2001.....	191	205	303	506	727	985	1,186	1,122	686	297	124	1
2000.....	158	201	280	444	637	972	1,077	968	678	304	81	-
1999.....	157	195	314	477	691	884	1,065	949	629	280	93	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
All races, both sexes												
2004.....	122	146	175	256	356	538	608	461	278	76	14	-
2003.....	117	151	216	295	398	603	630	557	291	93	23	-
2002.....	140	190	243	272	431	620	607	462	263	82	12	-
2001.....	125	168	193	307	437	599	589	468	235	67	10	-
2000.....	133	183	237	327	492	601	575	416	231	54	11	-
1999.....	144	189	222	338	516	614	613	442	235	63	8	-
All races, male												
2004.....	49	54	53	95	107	178	202	104	53	16	1	-
2003.....	39	62	84	105	136	199	185	160	62	10	2	-
2002.....	55	73	83	84	161	177	164	100	55	8	1	-
2001.....	41	73	62	106	125	182	148	106	50	10	-	-
2000.....	55	72	94	98	139	177	144	96	40	13	1	-
1999.....	50	64	69	107	183	164	161	89	26	8	-	-
All races, female												
2004.....	73	92	122	161	249	360	406	357	225	60	13	-
2003.....	78	89	132	190	262	404	445	397	229	83	21	-
2002.....	85	117	160	188	270	443	443	362	208	74	11	-
2001.....	84	95	131	201	312	417	441	362	185	57	10	-
2000.....	78	111	143	229	353	424	431	320	191	41	10	-
1999.....	94	125	153	231	333	450	452	353	209	55	8	-
White, both sexes												
2004.....	96	105	152	219	323	498	580	445	265	75	13	-
2003.....	78	109	168	259	352	568	604	532	284	89	22	-
2002.....	110	149	199	230	400	570	582	438	256	79	11	-
2001.....	87	127	166	261	388	558	567	447	228	62	10	-
2000.....	107	148	202	293	441	575	554	400	216	50	11	-
1999.....	108	156	185	302	474	578	583	421	227	57	7	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)

White, male

2004.....	36	38	45	82	102	170	194	101	51	16	1	-
2003.....	27	43	64	91	125	193	182	154	60	10	2	-
2002.....	42	56	70	69	150	163	160	97	54	8	-	-
2001.....	32	61	54	88	114	172	141	101	48	8	-	-
2000.....	48	62	85	88	122	169	140	93	36	11	1	-
1999.....	40	53	58	100	170	151	155	87	26	7	-	-

White, female

2004.....	60	67	107	137	221	328	386	344	214	59	12	-
2003.....	51	66	104	168	227	375	422	378	224	79	20	-
2002.....	68	93	129	161	250	407	422	341	202	71	11	-
2001.....	55	66	112	173	274	386	426	346	180	54	10	-
2000.....	59	86	117	205	319	406	414	307	180	39	10	-
1999.....	68	103	127	202	304	427	428	334	201	50	7	-

Black, both sexes

2004.....	22	33	13	25	21	24	16	12	8	-	1	-
2003.....	31	34	38	30	36	23	17	18	4	3	1	-
2002.....	23	30	33	30	19	28	14	12	3	2	1	-
2001.....	33	31	22	31	38	28	12	13	4	5	-	-
2000.....	20	26	24	31	27	21	15	12	11	4	-	-
1999.....	29	29	23	25	31	24	16	16	5	4	1	-

Black, male

2004.....	11	14	5	10	3	4	4	2	1	-	-	-
2003.....	11	17	16	13	8	6	2	4	2	-	-	-
2002.....	10	12	10	13	8	8	3	2	-	-	1	-
2001.....	9	10	7	11	6	5	3	4	-	2	-	-
2000.....	5	7	7	8	7	7	3	3	3	2	-	-
1999.....	10	9	8	6	10	10	4	1	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Black, female												
2004.....	11	19	8	15	18	20	12	10	7	-	1	-
2003.....	20	17	22	17	28	17	15	14	2	3	1	-
2002.....	13	18	23	17	11	20	11	10	3	2	-	-
2001.....	24	21	15	20	32	23	9	9	4	3	-	-
2000.....	15	19	17	23	19	14	14	9	8	2	-	-
1999.....	19	20	15	19	21	14	12	15	5	4	1	-
American Indian 1/, both sexes												
2004.....	2	3	-	-	2	2	3	-	1	-	-	-
2003.....	1	-	3	-	4	1	-	1	-	-	-	-
2002.....	1	2	3	3	2	4	4	1	-	-	-	-
2001.....	1	2	2	4	3	2	2	1	-	-	-	-
2000.....	-	3	3	1	3	1	2	2	-	-	-	-
1999.....	1	1	4	2	4	1	2	-	1	-	-	-
American Indian 1/, male												
2004.....	2	1	-	-	-	-	2	-	-	-	-	-
2003.....	-	-	2	-	1	-	-	-	-	-	-	-
2002.....	1	-	2	-	-	3	1	1	-	-	-	-
2001.....	-	-	1	1	2	2	1	1	-	-	-	-
2000.....	-	1	-	-	1	-	1	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, female												
2004.....	-	2	-	-	2	2	1	-	1	-	-	-
2003.....	1	-	1	-	3	1	-	1	-	-	-	-
2002.....	-	2	1	3	2	1	3	-	-	-	-	-
2001.....	1	2	1	3	1	-	1	-	-	-	-	-
2000.....	-	2	3	1	2	1	1	2	-	-	-	-
1999.....	1	1	4	2	4	1	2	-	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Asian or Pacific Islander 2/, both sexes												
2004.....	2	5	10	12	10	14	9	4	4	1	-	-
2003.....	7	8	7	6	6	11	9	6	3	1	-	-
2002.....	6	9	8	9	10	18	7	11	4	1	-	-
2001.....	4	8	3	11	8	11	8	7	3	-	-	-
2000.....	6	6	8	2	21	4	4	2	4	-	-	-
1999.....	6	3	10	9	7	11	12	5	2	2	-	-
Asian or Pacific Islander 2/, male												
2004.....	-	1	3	3	2	4	2	1	1	-	-	-
2003.....	1	2	2	1	2	-	1	2	-	-	-	-
2002.....	2	5	1	2	3	3	-	-	1	-	-	-
2001.....	-	2	-	6	3	3	3	-	2	-	-	-
2000.....	2	2	2	2	8	1	2	-	1	-	-	-
1999.....	-	2	3	1	3	3	2	1	-	1	-	-
Asian or Pacific Islander 2/, female												
2004.....	2	4	7	9	8	10	7	3	3	1	-	-
2003.....	6	6	5	5	4	11	8	4	3	1	-	-
2002.....	4	4	7	7	7	15	7	11	3	1	-	-
2001.....	4	6	3	5	5	8	5	7	1	-	-	-
2000.....	4	4	6	-	13	3	2	2	3	-	-	-
1999.....	6	1	7	8	4	8	10	4	2	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever (I00-I02)												
All races, both sexes												
2004.....	2	2	4	4	4	3	7	1	4	-	-	-
2003.....	2	2	5	2	7	8	3	5	-	-	-	-
2002.....	4	2	3	1	3	12	5	9	2	2	-	-
2001.....	1	2	4	5	6	8	8	16	3	2	-	-
2000.....	2	1	5	4	3	5	2	6	-	-	-	-
1999.....	3	4	1	-	5	4	4	3	1	-	-	-
All races, male												
2004.....	-	1	2	2	2	-	2	1	1	-	-	-
2003.....	1	2	3	1	3	1	1	2	-	-	-	-
2002.....	1	1	-	-	1	3	1	3	-	2	-	-
2001.....	1	2	1	1	2	2	3	4	1	1	-	-
2000.....	-	1	4	1	3	3	-	1	-	-	-	-
1999.....	2	3	1	-	2	2	2	-	-	-	-	-
All races, female												
2004.....	2	1	2	2	2	3	5	-	3	-	-	-
2003.....	1	-	2	1	4	7	2	3	-	-	-	-
2002.....	3	1	3	1	2	9	4	6	2	-	-	-
2001.....	-	-	3	4	4	6	5	12	2	1	-	-
2000.....	2	-	1	3	-	2	2	5	-	-	-	-
1999.....	1	1	-	-	3	2	2	3	1	-	-	-
White, both sexes												
2004.....	2	1	2	3	4	3	6	1	4	-	-	-
2003.....	-	2	3	2	6	6	2	4	-	-	-	-
2002.....	4	2	3	1	3	10	5	8	1	2	-	-
2001.....	1	2	3	2	5	8	8	14	3	1	-	-
2000.....	1	-	5	3	3	5	2	5	-	-	-	-
1999.....	3	2	1	-	5	3	4	2	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever (I00-I02)												
White, male												
2004.....	-	-	1	2	2	-	2	1	1	-	-	-
2003.....	-	2	2	1	3	1	1	1	-	-	-	-
2002.....	1	1	-	-	1	3	1	3	-	2	-	-
2001.....	1	2	-	1	2	2	3	3	1	1	-	-
2000.....	-	-	4	1	3	3	-	1	-	-	-	-
1999.....	2	2	1	-	2	1	2	-	-	-	-	-
White, female												
2004.....	2	1	1	1	2	3	4	-	3	-	-	-
2003.....	-	-	1	1	3	5	1	3	-	-	-	-
2002.....	3	1	3	1	2	7	4	5	1	-	-	-
2001.....	-	-	3	1	3	6	5	11	2	-	-	-
2000.....	1	-	1	2	-	2	2	4	-	-	-	-
1999.....	1	-	-	-	3	2	2	2	-	-	-	-
Black, both sexes												
2004.....	-	1	2	1	-	-	1	-	-	-	-	-
2003.....	2	-	2	-	1	2	1	-	-	-	-	-
2002.....	-	-	-	-	-	2	-	-	1	-	-	-
2001.....	-	-	1	3	-	-	-	2	-	1	-	-
2000.....	1	1	-	1	-	-	-	1	-	-	-	-
1999.....	-	2	-	-	-	1	-	1	1	-	-	-
Black, male												
2004.....	-	1	1	-	-	-	-	-	-	-	-	-
2003.....	1	-	1	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	1	-	-	-	-	1	-	-	-	-
2000.....	-	1	-	-	-	-	-	-	-	-	-	-
1999.....	-	1	-	-	-	1	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic rheumatic heart diseases (I05-I09)												
All races, both sexes												
2004.....	120	144	171	252	352	535	601	460	274	76	14	-
2003.....	115	149	211	293	391	595	627	552	291	93	23	-
2002.....	136	188	240	271	428	608	602	453	261	80	12	-
2001.....	124	166	189	302	431	591	581	452	232	65	10	-
2000.....	131	182	232	323	489	596	573	410	231	54	11	-
1999.....	141	185	221	338	511	610	609	439	234	63	8	-
All races, male												
2004.....	49	53	51	93	105	178	200	103	52	16	1	-
2003.....	38	60	81	104	133	198	184	158	62	10	2	-
2002.....	54	72	83	84	160	174	163	97	55	6	1	-
2001.....	40	71	61	105	123	180	145	102	49	9	-	-
2000.....	55	71	90	97	136	174	144	95	40	13	1	-
1999.....	48	61	68	107	181	162	159	89	26	8	-	-
All races, female												
2004.....	71	91	120	159	247	357	401	357	222	60	13	-
2003.....	77	89	130	189	258	397	443	394	229	83	21	-
2002.....	82	116	157	187	268	434	439	356	206	74	11	-
2001.....	84	95	128	197	308	411	436	350	183	56	10	-
2000.....	76	111	142	226	353	422	429	315	191	41	10	-
1999.....	93	124	153	231	330	448	450	350	208	55	8	-
White, both sexes												
2004.....	94	104	150	216	319	495	574	444	261	75	13	-
2003.....	78	107	165	257	346	562	602	528	284	89	22	-
2002.....	106	147	196	229	397	560	577	430	255	77	11	-
2001.....	86	125	163	259	383	550	559	433	225	61	10	-
2000.....	106	148	197	290	438	570	552	395	216	50	11	-
1999.....	105	154	184	302	469	575	579	419	227	57	7	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic rheumatic heart diseases (I05-I09)												
White, male												
2004.....	36	38	44	80	100	170	192	100	50	16	1	-
2003.....	27	41	62	90	122	192	181	153	60	10	2	-
2002.....	41	55	70	69	149	160	159	94	54	6	-	-
2001.....	31	59	54	87	112	170	138	98	47	7	-	-
2000.....	48	62	81	87	119	166	140	92	36	11	1	-
1999.....	38	51	57	100	168	150	153	87	26	7	-	-
White, female												
2004.....	58	66	106	136	219	325	382	344	211	59	12	-
2003.....	51	66	103	167	224	370	421	375	224	79	20	-
2002.....	65	92	126	160	248	400	418	336	201	71	11	-
2001.....	55	66	109	172	271	380	421	335	178	54	10	-
2000.....	58	86	116	203	319	404	412	303	180	39	10	-
1999.....	67	103	127	202	301	425	426	332	201	50	7	-
Black, both sexes												
2004.....	22	32	11	24	21	24	15	12	8	-	1	-
2003.....	29	34	36	30	35	21	16	18	4	3	1	-
2002.....	23	30	33	30	19	26	14	12	2	2	1	-
2001.....	33	31	21	28	38	28	12	11	4	4	-	-
2000.....	19	25	24	30	27	21	15	11	11	4	-	-
1999.....	29	27	23	25	31	23	16	15	4	4	1	-
Black, male												
2004.....	11	13	4	10	3	4	4	2	1	-	-	-
2003.....	10	17	15	13	8	6	2	4	2	-	-	-
2002.....	10	12	10	13	8	8	3	2	-	-	1	-
2001.....	9	10	6	11	6	5	3	3	-	2	-	-
2000.....	5	6	7	8	8	7	3	3	3	2	-	-
1999.....	10	8	8	6	10	9	4	1	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic rheumatic heart diseases (I05-I09)												
Black, female												
2004.....	11	19	7	14	18	20	11	10	7	-	1	-
2003.....	19	17	21	17	27	15	14	14	2	3	1	-
2002.....	13	18	23	17	11	18	11	10	2	2	-	-
2001.....	24	21	15	17	32	23	9	8	4	2	-	-
2000.....	14	19	17	22	19	14	14	8	8	2	-	-
1999.....	19	19	15	19	21	14	12	14	4	4	1	-
American Indian 1/, both sexes												
2004.....	2	3	-	-	2	2	3	-	1	-	-	-
2003.....	1	-	3	-	4	1	-	1	-	-	-	-
2002.....	1	2	3	3	2	4	4	1	-	-	-	-
2001.....	1	2	2	4	2	2	2	1	-	-	-	-
2000.....	-	3	3	1	3	1	2	2	-	-	-	-
1999.....	1	1	4	2	4	1	2	-	1	-	-	-
American Indian 1/, male												
2004.....	2	1	-	-	-	-	2	-	-	-	-	-
2003.....	-	-	2	-	1	-	-	-	-	-	-	-
2002.....	1	-	2	-	-	3	1	1	-	-	-	-
2001.....	-	-	1	1	2	2	1	1	-	-	-	-
2000.....	-	1	-	-	1	-	1	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, female												
2004.....	-	2	-	-	2	2	1	-	1	-	-	-
2003.....	1	-	1	-	3	1	-	1	-	-	-	-
2002.....	-	2	1	3	2	1	3	-	-	-	-	-
2001.....	1	2	1	3	-	-	1	-	-	-	-	-
2000.....	-	2	3	1	2	1	1	2	-	-	-	-
1999.....	1	1	4	2	4	1	2	-	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic rheumatic heart diseases (I05-I09)												
Asian or Pacific Islander 2/, both sexes												
2004.....	2	5	10	12	10	14	9	4	4	1	-	-
2003.....	7	8	7	6	6	11	9	5	3	1	-	-
2002.....	6	9	8	9	10	18	7	10	4	1	-	-
2001.....	4	8	3	11	8	11	8	7	3	-	-	-
2000.....	6	6	8	2	21	4	4	2	4	-	-	-
1999.....	6	3	10	9	7	11	12	5	2	2	-	-
Asian or Pacific Islander 2/, male												
2004.....	-	1	3	3	2	4	2	1	1	-	-	-
2003.....	1	2	2	1	2	-	1	1	-	-	-	-
2002.....	2	5	1	2	3	3	-	-	1	-	-	-
2001.....	-	2	-	6	3	3	3	-	2	-	-	-
2000.....	2	2	2	2	8	1	2	-	1	-	-	-
1999.....	-	2	3	1	3	3	2	1	-	1	-	-
Asian or Pacific Islander 2/, female												
2004.....	2	4	7	9	8	10	7	3	3	1	-	-
2003.....	6	6	5	5	4	11	8	4	3	1	-	-
2002.....	4	4	7	7	7	15	7	10	3	1	-	-
2001.....	4	6	3	5	5	8	5	7	1	-	-	-
2000.....	4	4	6	-	13	3	2	2	3	-	-	-
1999.....	6	1	7	8	4	8	10	4	2	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic mitral valve diseases (I05)												
All races, both sexes												
2004.....	65	73	91	115	181	245	263	195	93	25	7	-
2003.....	68	70	123	162	198	301	304	220	103	29	6	-
2002.....	90	113	146	134	229	327	305	196	103	32	8	-
2001.....	84	99	109	170	246	331	309	237	107	27	3	-
2000.....	81	110	138	193	299	325	337	212	121	22	7	-
1999.....	93	111	128	203	285	361	334	232	112	28	2	-
All races, male												
2004.....	24	27	22	39	55	75	82	44	20	5	-	-
2003.....	20	31	58	62	61	95	87	57	24	4	-	-
2002.....	35	40	50	41	89	79	75	34	21	4	-	-
2001.....	25	38	28	60	69	89	72	57	21	3	-	-
2000.....	29	44	55	54	84	97	87	50	21	4	-	-
1999.....	32	37	36	64	94	97	79	41	14	3	-	-
All races, female												
2004.....	41	46	69	76	126	170	181	151	73	20	7	-
2003.....	48	39	65	100	137	206	217	163	79	25	6	-
2002.....	55	73	96	93	140	248	230	162	82	28	8	-
2001.....	59	61	81	110	177	242	237	180	86	24	3	-
2000.....	52	66	83	139	215	228	250	162	100	18	7	-
1999.....	61	74	92	139	191	264	255	191	98	25	2	-
White, both sexes												
2004.....	47	53	79	103	162	222	247	190	89	24	7	-
2003.....	45	44	93	144	181	282	289	209	101	28	5	-
2002.....	72	89	118	110	215	301	297	186	101	32	8	-
2001.....	62	74	95	142	219	309	301	227	105	26	3	-
2000.....	69	94	118	181	263	308	327	206	116	22	7	-
1999.....	64	91	106	179	259	342	319	222	111	24	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic mitral valve diseases (I05)												
White, male												
2004.....	16	21	19	35	52	70	79	44	19	5	-	-
2003.....	14	18	42	55	57	91	84	55	24	4	-	-
2002.....	27	31	42	35	84	72	74	32	21	4	-	-
2001.....	19	32	26	48	62	85	70	53	21	3	-	-
2000.....	28	40	49	53	72	90	85	48	19	4	-	-
1999.....	26	29	31	58	86	88	76	40	14	3	-	-
White, female												
2004.....	31	32	60	68	110	152	168	146	70	19	7	-
2003.....	31	26	51	89	124	191	205	154	77	24	5	-
2002.....	45	58	76	75	131	229	223	154	80	28	8	-
2001.....	43	42	69	94	157	224	231	174	84	23	3	-
2000.....	41	54	69	128	191	218	242	158	97	18	7	-
1999.....	38	62	75	121	173	254	243	182	97	21	1	-
Black, both sexes												
2004.....	15	15	5	10	11	14	9	5	3	-	-	-
2003.....	16	21	23	15	13	13	10	8	1	1	1	-
2002.....	14	18	23	15	8	15	4	6	-	-	-	-
2001.....	22	18	12	19	20	17	4	5	1	1	-	-
2000.....	8	13	13	11	18	14	6	4	3	-	-	-
1999.....	24	17	13	21	22	14	8	8	1	4	1	-
Black, male												
2004.....	7	5	1	4	1	3	1	-	1	-	-	-
2003.....	5	12	13	6	2	4	2	2	-	-	-	-
2002.....	6	7	7	4	3	5	-	1	-	-	-	-
2001.....	6	5	2	8	3	3	1	3	-	-	-	-
2000.....	1	2	4	1	6	6	-	2	2	-	-	-
1999.....	6	6	4	5	8	8	3	1	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic mitral valve diseases (I05)												
Black, female												
2004.....	8	10	4	6	10	11	8	5	2	-	-	-
2003.....	11	9	10	9	11	9	8	6	1	1	1	-
2002.....	8	11	16	11	5	10	4	5	-	-	-	-
2001.....	16	13	10	11	17	14	3	2	1	1	-	-
2000.....	7	11	9	10	12	8	6	2	1	-	-	-
1999.....	18	11	9	16	14	6	5	7	1	4	1	-
American Indian 1/, both sexes												
2004.....	1	2	-	-	-	1	2	-	1	-	-	-
2003.....	1	-	2	-	2	1	-	1	-	-	-	-
2002.....	1	1	1	3	1	-	2	1	-	-	-	-
2001.....	-	1	1	2	2	-	1	1	-	-	-	-
2000.....	-	1	2	1	1	-	1	1	-	-	-	-
1999.....	1	-	4	-	1	1	1	-	-	-	-	-
American Indian 1/, male												
2004.....	1	1	-	-	-	-	1	-	-	-	-	-
2003.....	-	-	2	-	1	-	-	-	-	-	-	-
2002.....	1	-	1	-	-	-	1	1	-	-	-	-
2001.....	-	-	-	-	2	-	-	1	-	-	-	-
2000.....	-	1	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, female												
2004.....	-	1	-	-	-	1	1	-	1	-	-	-
2003.....	1	-	-	-	1	1	-	1	-	-	-	-
2002.....	-	1	-	3	1	-	1	-	-	-	-	-
2001.....	-	1	1	2	-	-	1	-	-	-	-	-
2000.....	-	-	2	1	1	-	1	1	-	-	-	-
1999.....	1	-	4	-	1	1	1	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic mitral valve diseases (I05)												
Asian or Pacific Islander 2/, both sexes												
2004.....	2	3	7	2	8	8	5	-	-	1	-	-
2003.....	6	5	5	3	2	5	5	2	1	-	-	-
2002.....	3	5	4	6	5	11	2	3	2	-	-	-
2001.....	-	6	1	7	5	5	3	4	1	-	-	-
2000.....	4	2	5	-	17	3	3	1	2	-	-	-
1999.....	4	3	5	3	3	4	6	2	-	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	-	-	2	-	2	2	1	-	-	-	-	-
2003.....	1	1	1	1	1	-	1	-	-	-	-	-
2002.....	1	2	-	2	2	2	-	-	-	-	-	-
2001.....	-	1	-	4	2	1	1	-	-	-	-	-
2000.....	-	1	2	-	6	1	2	-	-	-	-	-
1999.....	-	2	1	1	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	2	3	5	2	6	6	4	-	-	1	-	-
2003.....	5	4	4	2	1	5	4	2	1	-	-	-
2002.....	2	3	4	4	3	9	2	3	2	-	-	-
2001.....	-	5	1	3	3	4	2	4	1	-	-	-
2000.....	4	1	3	-	11	2	1	1	2	-	-	-
1999.....	4	1	4	2	3	3	6	2	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Rheumatic aortic valve diseases (I06)

All races, both sexes

2004.....	1	7	6	5	8	17	23	15	13	5	1	-
2003.....	1	2	4	7	7	18	18	18	18	2	2	-
2002.....	4	8	3	8	8	19	27	25	21	2	-	-
2001.....	3	1	4	5	5	9	15	10	5	3	-	-
2000.....	3	3	2	3	2	3	3	7	3	1	-	-
1999.....	-	2	2	6	9	8	8	8	3	2	-	-

All races, male

2004.....	-	3	3	-	6	8	10	3	3	2	-	-
2003.....	-	1	-	1	2	8	9	6	6	-	-	-
2002.....	2	3	1	2	3	6	10	7	6	-	-	-
2001.....	1	1	3	2	-	6	7	1	1	-	-	-
2000.....	3	2	1	1	1	2	2	2	-	-	-	-
1999.....	-	2	1	2	1	2	5	2	-	-	-	-

All races, female

2004.....	1	4	3	5	2	9	13	12	10	3	1	-
2003.....	1	1	4	6	5	10	9	12	12	2	2	-
2002.....	2	5	2	6	5	13	17	18	15	2	-	-
2001.....	2	-	1	3	5	3	8	9	4	3	-	-
2000.....	-	1	1	2	1	1	1	5	3	1	-	-
1999.....	-	-	1	4	8	6	3	6	3	2	-	-

White, both sexes

2004.....	1	3	6	5	7	17	23	14	13	5	1	-
2003.....	1	2	2	7	6	18	18	18	18	2	2	-
2002.....	4	6	1	6	8	16	25	23	21	2	-	-
2001.....	2	1	4	5	4	9	14	8	5	3	-	-
2000.....	2	2	2	2	1	3	3	7	3	1	-	-
1999.....	-	2	2	5	9	8	7	8	3	2	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic aortic valve diseases (I06)												
White, male												
2004.....	-	1	3	-	6	8	10	2	3	2	-	-
2003.....	-	1	-	1	1	8	9	6	6	-	-	-
2002.....	2	3	1	1	3	6	10	7	6	-	-	-
2001.....	1	1	3	2	-	6	7	1	1	-	-	-
2000.....	2	1	1	-	-	2	2	2	-	-	-	-
1999.....	-	2	1	2	1	2	4	2	-	-	-	-
White, female												
2004.....	1	2	3	5	1	9	13	12	10	3	1	-
2003.....	1	1	2	6	5	10	9	12	12	2	2	-
2002.....	2	3	-	5	5	10	15	16	15	2	-	-
2001.....	1	-	1	3	4	3	7	7	4	3	-	-
2000.....	-	1	1	2	1	1	1	5	3	1	-	-
1999.....	-	-	1	3	8	6	3	6	3	2	-	-
Black, both sexes												
2004.....	-	3	-	-	-	-	-	1	-	-	-	-
2003.....	-	-	1	-	1	-	-	-	-	-	-	-
2002.....	-	1	1	2	-	1	1	-	-	-	-	-
2001.....	1	-	-	-	1	-	1	1	-	-	-	-
2000.....	-	1	-	1	-	-	-	-	-	-	-	-
1999.....	-	-	-	1	-	-	1	-	-	-	-	-
Black, male												
2004.....	-	1	-	-	-	-	-	1	-	-	-	-
2003.....	-	-	-	-	1	-	-	-	-	-	-	-
2002.....	-	-	-	1	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	1	-	1	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	1	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Disorders of both mitral and aortic valves (I08.0)

All races, both sexes

2004.....	19	24	26	49	63	108	143	143	82	25	5	-
2003.....	16	26	22	43	68	119	148	161	95	39	11	-
2002.....	13	6	18	33	58	96	98	82	59	19	3	-
2001.....	9	11	16	33	53	49	81	61	34	11	2	-
2000.....	9	15	20	23	44	56	50	44	30	9	3	-
1999.....	10	15	20	21	53	40	55	49	21	9	4	-

All races, male

2004.....	11	12	10	27	19	45	64	35	17	5	1	-
2003.....	7	11	6	18	35	50	50	59	20	6	1	-
2002.....	3	1	10	14	28	42	30	21	12	1	-	-
2001.....	5	5	7	12	24	19	31	19	9	2	-	-
2000.....	8	8	9	9	17	18	14	13	7	2	1	-
1999.....	3	6	9	10	24	15	18	16	2	1	-	-

All races, female

2004.....	8	12	16	22	44	63	79	108	65	20	4	-
2003.....	9	15	16	25	33	69	98	102	75	33	10	-
2002.....	10	5	8	19	30	54	68	61	47	18	3	-
2001.....	4	6	9	21	29	30	50	42	25	9	2	-
2000.....	1	7	11	14	27	38	36	31	23	7	2	-
1999.....	7	9	11	11	29	25	37	33	19	8	4	-

White, both sexes

2004.....	17	15	24	41	58	100	139	138	78	25	4	-
2003.....	12	21	18	33	59	111	144	157	91	38	11	-
2002.....	7	5	15	29	50	92	93	79	57	18	3	-
2001.....	8	10	14	30	45	47	77	58	33	9	2	-
2000.....	7	12	16	19	38	56	50	43	26	6	3	-
1999.....	8	13	17	18	48	39	51	47	19	9	4	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Disorders of both mitral and aortic valves (I08.0)

White, male

2004.....	9	7	9	24	18	43	61	33	17	5	1	-
2003.....	6	9	4	13	31	49	50	58	19	6	1	-
2002.....	2	1	8	12	23	39	28	21	12	1	-	-
2001.....	4	5	6	10	21	17	30	19	9	1	-	-
2000.....	6	6	8	7	14	18	14	13	6	1	1	-
1999.....	2	5	8	10	23	15	17	16	2	1	-	-

White, female

2004.....	8	8	15	17	40	57	78	105	61	20	3	-
2003.....	6	12	14	20	28	62	94	99	72	32	10	-
2002.....	5	4	7	17	27	53	65	58	45	17	3	-
2001.....	4	5	8	20	24	30	47	39	24	8	2	-
2000.....	1	6	8	12	24	38	36	30	20	5	2	-
1999.....	6	8	9	8	25	24	34	31	17	8	4	-

Black, both sexes

2004.....	2	8	2	6	5	3	2	4	2	-	1	-
2003.....	4	4	3	9	7	3	2	3	2	1	-	-
2002.....	4	1	3	4	6	2	4	2	1	1	-	-
2001.....	1	1	2	2	7	1	2	3	1	2	-	-
2000.....	1	3	3	3	3	-	-	1	3	3	-	-
1999.....	1	2	2	1	2	-	3	1	1	-	-	-

Black, male

2004.....	2	5	1	3	1	1	2	1	-	-	-	-
2003.....	1	2	1	5	3	1	-	1	1	-	-	-
2002.....	1	-	2	2	4	1	2	-	-	-	-	-
2001.....	1	-	1	1	2	1	1	-	-	1	-	-
2000.....	1	2	1	1	1	-	-	-	-	1	-	-
1999.....	1	1	1	-	1	-	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of both mitral and aortic valves (I08.0)												
Black, female												
2004.....	-	3	1	3	4	2	-	3	2	-	1	-
2003.....	3	2	2	4	4	2	2	2	1	1	-	-
2002.....	3	1	1	2	2	1	2	2	1	1	-	-
2001.....	-	1	1	1	5	-	1	3	1	1	-	-
2000.....	-	1	2	2	2	-	-	1	3	2	-	-
1999.....	-	1	1	1	1	-	3	1	1	-	-	-
American Indian 1/, both sexes												
2004.....	-	1	-	-	-	1	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	2	1	-	-	-	-	-
2001.....	-	-	-	1	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	1	-	-	-	-	-	-	-
1999.....	-	-	-	1	2	-	-	-	-	-	-	-
American Indian 1/, male												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	2	-	-	-	-	-	-
2001.....	-	-	-	1	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	1	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, female												
2004.....	-	1	-	-	-	1	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	1	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	1	2	-	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of both mitral and aortic valves (I08.0)												
Asian or Pacific Islander 2/, both sexes												
2004.....	-	-	-	2	-	4	2	1	2	-	-	-
2003.....	-	1	1	1	2	5	2	1	2	-	-	-
2002.....	2	-	-	-	2	-	-	1	1	-	-	-
2001.....	-	-	-	-	1	1	2	-	-	-	-	-
2000.....	1	-	1	1	2	-	-	-	1	-	-	-
1999.....	1	-	1	1	1	1	1	1	1	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	-	-	-	-	-	1	1	1	-	-	-	-
2003.....	-	-	1	-	1	-	-	-	-	-	-	-
2002.....	-	-	-	-	1	-	-	-	-	-	-	-
2001.....	-	-	-	-	1	1	-	-	-	-	-	-
2000.....	1	-	-	1	1	-	-	-	1	-	-	-
1999.....	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	-	-	-	2	-	3	1	-	2	-	-	-
2003.....	-	1	-	1	1	5	2	1	2	-	-	-
2002.....	2	-	-	-	1	-	-	1	1	-	-	-
2001.....	-	-	-	-	-	-	2	-	-	-	-	-
2000.....	-	-	1	-	1	-	-	-	-	-	-	-
1999.....	1	-	1	1	1	1	-	1	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)

All races, both sexes

2004.....	35	40	48	83	100	165	172	107	86	21	1	-
2003.....	30	51	62	81	118	157	157	153	75	23	4	-
2002.....	29	61	73	96	133	166	172	150	78	27	1	-
2001.....	28	55	60	94	127	202	176	144	86	24	5	-
2000.....	38	54	72	104	144	212	183	147	77	22	1	-
1999.....	38	57	71	108	164	201	212	150	98	24	2	-

All races, male

2004.....	14	11	16	27	25	50	44	21	12	4	-	-
2003.....	11	17	17	23	35	45	38	36	12	-	1	-
2002.....	14	28	22	27	40	47	48	35	16	1	1	-
2001.....	9	27	23	31	30	66	35	25	18	4	-	-
2000.....	15	17	25	33	34	57	41	30	12	7	-	-
1999.....	13	16	22	31	62	48	57	30	10	4	-	-

All races, female

2004.....	21	29	32	56	75	115	128	86	74	17	1	-
2003.....	19	34	45	58	83	112	119	117	63	23	3	-
2002.....	15	33	51	69	93	119	124	115	62	26	-	-
2001.....	19	28	37	63	97	136	141	119	68	20	5	-
2000.....	23	37	47	71	110	155	142	117	65	15	1	-
1999.....	25	41	49	77	102	153	155	120	88	20	2	-

White, both sexes

2004.....	29	33	41	67	92	156	165	102	81	21	1	-
2003.....	20	40	52	73	100	151	151	144	74	21	4	-
2002.....	23	47	62	84	124	151	162	142	76	25	-	-
2001.....	14	40	50	82	115	185	167	140	82	23	5	-
2000.....	28	40	61	88	136	203	172	139	71	21	1	-
1999.....	33	48	59	100	153	186	202	142	94	22	2	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)

White, male

2004.....	11	9	13	21	24	49	42	21	11	4	-	-
2003.....	7	13	16	21	33	44	38	34	11	-	1	-
2002.....	10	20	19	21	39	43	47	34	15	1	-	-
2001.....	7	21	19	27	29	62	31	25	16	3	-	-
2000.....	12	15	23	27	33	56	39	29	11	6	-	-
1999.....	10	15	17	30	58	45	56	29	10	3	-	-

White, female

2004.....	18	24	28	46	68	107	123	81	70	17	1	-
2003.....	13	27	36	52	67	107	113	110	63	21	3	-
2002.....	13	27	43	63	85	108	115	108	61	24	-	-
2001.....	7	19	31	55	86	123	136	115	66	20	5	-
2000.....	16	25	38	61	103	147	133	110	60	15	1	-
1999.....	23	33	42	70	95	141	146	113	84	19	2	-

Black, both sexes

2004.....	5	6	4	8	5	7	4	2	3	-	-	-
2003.....	9	9	9	6	14	5	4	7	1	1	-	-
2002.....	5	10	6	9	5	8	5	4	1	1	1	-
2001.....	9	12	7	7	10	10	5	2	2	1	-	-
2000.....	10	8	8	15	6	7	9	6	5	1	-	-
1999.....	4	8	8	2	7	9	4	6	2	-	-	-

Black, male

2004.....	2	2	2	3	1	-	1	-	-	-	-	-
2003.....	4	3	1	2	2	1	-	1	1	-	-	-
2002.....	3	5	1	6	1	2	1	1	-	-	1	-
2001.....	2	5	3	2	1	1	1	-	-	1	-	-
2000.....	3	1	2	5	1	1	1	1	1	1	-	-
1999.....	3	1	3	1	1	1	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Black, female												
2004.....	3	4	2	5	4	7	3	2	3	-	-	-
2003.....	5	6	8	4	12	4	4	6	-	1	-	-
2002.....	2	5	5	3	4	6	4	3	1	1	-	-
2001.....	7	7	4	5	9	9	4	2	2	-	-	-
2000.....	7	7	6	10	5	6	8	5	4	-	-	-
1999.....	1	7	5	1	6	8	4	6	2	-	-	-
American Indian 1/, both sexes												
2004.....	1	-	-	-	2	-	1	-	-	-	-	-
2003.....	-	-	-	-	2	-	-	-	-	-	-	-
2002.....	-	-	1	-	1	2	1	-	-	-	-	-
2001.....	1	1	1	1	-	2	1	-	-	-	-	-
2000.....	-	2	1	-	1	1	1	1	-	-	-	-
1999.....	-	1	-	1	1	-	1	-	1	-	-	-
American Indian 1/, male												
2004.....	1	-	-	-	-	-	1	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	1	-	-	1	-	-	-	-	-	-
2001.....	-	-	1	-	-	2	1	-	-	-	-	-
2000.....	-	-	-	-	-	-	1	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, female												
2004.....	-	-	-	-	2	-	-	-	-	-	-	-
2003.....	-	-	-	-	2	-	-	-	-	-	-	-
2002.....	-	-	-	-	1	1	1	-	-	-	-	-
2001.....	1	1	-	1	-	-	-	-	-	-	-	-
2000.....	-	2	1	-	1	1	-	1	-	-	-	-
1999.....	-	1	-	1	1	-	1	-	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Asian or Pacific Islander 2/, both sexes												
2004.....	-	1	3	8	1	2	2	3	2	-	-	-
2003.....	1	2	1	2	2	1	2	2	-	1	-	-
2002.....	1	4	4	3	3	5	4	4	1	1	-	-
2001.....	4	2	2	4	2	5	3	2	2	-	-	-
2000.....	-	4	2	1	1	1	1	1	1	-	-	-
1999.....	1	-	4	5	3	6	5	2	1	2	-	-
Asian or Pacific Islander 2/, male												
2004.....	-	-	1	3	-	1	-	-	1	-	-	-
2003.....	-	1	-	-	-	-	-	1	-	-	-	-
2002.....	1	3	1	-	-	1	-	-	1	-	-	-
2001.....	-	1	-	2	-	1	2	-	2	-	-	-
2000.....	-	1	-	1	-	-	-	-	-	-	-	-
1999.....	-	-	2	-	3	2	1	1	-	1	-	-
Asian or Pacific Islander 2/, female												
2004.....	-	1	2	5	1	1	2	3	1	-	-	-
2003.....	1	1	1	2	2	1	2	1	-	1	-	-
2002.....	-	1	3	3	3	4	4	4	-	1	-	-
2001.....	4	1	2	2	2	4	1	2	-	-	-	-
2000.....	-	3	2	-	1	1	1	1	1	-	-	-
1999.....	1	-	2	5	-	4	4	1	1	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive diseases (I10-I13)												
All races, both sexes												
2004.....	2,523	2,938	3,057	3,320	4,212	5,907	8,164	8,884	7,306	3,212	820	7
2003.....	2,529	2,773	2,900	3,318	4,232	6,195	7,922	8,485	6,974	3,105	738	-
2002.....	2,247	2,476	2,643	3,089	4,130	5,771	7,554	8,028	6,416	2,868	639	4
2001.....	2,042	2,119	2,359	2,893	3,975	5,731	7,236	7,763	5,879	2,689	612	11
2000.....	1,757	1,949	2,371	2,813	3,993	5,577	6,871	7,451	5,622	2,500	552	3
1999.....	1,587	1,914	2,203	2,900	3,886	5,559	6,682	7,047	5,329	2,292	537	1
All races, male												
2004.....	1,674	1,865	1,913	1,918	2,109	2,611	3,052	2,718	1,695	548	101	6
2003.....	1,657	1,765	1,691	1,892	2,096	2,663	2,868	2,513	1,570	491	84	-
2002.....	1,472	1,587	1,580	1,676	2,029	2,520	2,751	2,357	1,444	447	76	4
2001.....	1,314	1,294	1,365	1,603	1,943	2,446	2,589	2,260	1,277	424	58	9
2000.....	1,186	1,198	1,412	1,487	1,998	2,336	2,487	2,139	1,223	373	60	2
1999.....	1,009	1,182	1,256	1,546	1,865	2,402	2,377	1,995	1,143	362	48	1
All races, female												
2004.....	849	1,073	1,144	1,402	2,103	3,296	5,112	6,166	5,611	2,664	719	1
2003.....	872	1,008	1,209	1,426	2,136	3,532	5,054	5,972	5,404	2,614	654	-
2002.....	775	889	1,063	1,413	2,101	3,251	4,803	5,671	4,972	2,421	563	-
2001.....	728	825	994	1,290	2,032	3,285	4,647	5,503	4,602	2,265	554	2
2000.....	571	751	959	1,326	1,995	3,241	4,384	5,312	4,399	2,127	492	1
1999.....	578	732	947	1,354	2,021	3,157	4,305	5,052	4,186	1,930	489	-
White, both sexes												
2004.....	1,431	1,756	1,917	2,213	2,948	4,432	6,599	7,581	6,264	2,783	667	6
2003.....	1,429	1,635	1,850	2,127	2,920	4,663	6,512	7,224	6,051	2,717	596	-
2002.....	1,274	1,469	1,639	2,001	2,903	4,330	6,081	6,738	5,522	2,513	534	1
2001.....	1,125	1,244	1,448	1,839	2,773	4,358	5,865	6,526	5,149	2,355	489	6
2000.....	949	1,112	1,417	1,810	2,752	4,200	5,548	6,293	4,848	2,117	444	1
1999.....	856	1,079	1,324	1,853	2,671	4,138	5,369	5,926	4,626	1,943	438	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive diseases (I10-I13)												
White, male												
2004.....	1,009	1,162	1,229	1,308	1,516	1,972	2,489	2,343	1,457	456	82	5
2003.....	1,006	1,104	1,118	1,258	1,482	2,048	2,367	2,119	1,338	418	65	-
2002.....	889	1,001	1,007	1,109	1,463	1,917	2,210	1,993	1,228	378	55	1
2001.....	767	801	875	1,058	1,395	1,883	2,091	1,888	1,117	352	37	5
2000.....	683	722	876	994	1,423	1,798	2,006	1,802	1,029	298	44	1
1999.....	574	707	788	1,013	1,315	1,808	1,915	1,646	981	285	40	-
White, female												
2004.....	422	594	688	905	1,432	2,460	4,110	5,238	4,807	2,327	585	1
2003.....	423	531	732	869	1,438	2,615	4,145	5,105	4,713	2,299	531	-
2002.....	385	468	632	892	1,440	2,413	3,871	4,745	4,294	2,135	479	-
2001.....	358	443	573	781	1,378	2,475	3,774	4,638	4,032	2,003	452	1
2000.....	266	390	541	816	1,329	2,402	3,542	4,491	3,819	1,819	400	-
1999.....	282	372	536	840	1,356	2,330	3,454	4,280	3,645	1,658	398	-
Black, both sexes												
2004.....	1,038	1,094	1,050	1,015	1,150	1,313	1,363	1,084	896	365	132	1
2003.....	1,052	1,070	980	1,081	1,187	1,363	1,242	1,084	792	346	131	-
2002.....	934	955	932	995	1,122	1,304	1,306	1,128	771	314	95	2
2001.....	870	824	853	969	1,084	1,228	1,202	1,091	654	303	115	5
2000.....	759	804	899	934	1,142	1,244	1,156	1,031	693	333	97	2
1999.....	705	788	814	956	1,121	1,291	1,160	988	618	313	88	1
Black, male												
2004.....	626	653	627	563	535	569	479	301	188	71	16	1
2003.....	619	620	531	582	567	542	435	322	187	62	16	-
2002.....	557	554	526	509	512	539	464	310	178	56	18	2
2001.....	521	463	455	508	496	496	435	319	134	60	20	4
2000.....	467	459	501	459	519	474	415	290	168	57	11	1
1999.....	417	444	435	488	500	530	403	292	135	69	7	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive diseases (I10-I13)												
Black, female												
2004.....	412	441	423	452	615	744	884	783	708	294	116	-
2003.....	433	450	449	499	620	821	807	762	605	284	115	-
2002.....	377	401	406	486	610	765	842	818	593	258	77	-
2001.....	349	361	398	461	588	732	767	772	520	243	95	1
2000.....	292	345	398	475	623	770	741	741	525	276	86	1
1999.....	288	344	379	468	621	761	757	696	483	244	81	-
American Indian 1/, both sexes												
2004.....	13	32	20	18	19	22	32	15	15	7	2	-
2003.....	16	21	23	21	25	15	16	17	20	4	1	-
2002.....	6	7	14	21	11	16	15	15	10	8	1	1
2001.....	7	8	10	13	13	18	16	12	11	6	1	-
2000.....	13	10	16	11	14	15	19	23	6	4	2	-
1999.....	3	11	13	10	16	15	15	18	9	2	-	-
American Indian 1/, male												
2004.....	6	17	13	10	14	9	11	5	3	3	-	-
2003.....	8	15	15	9	11	10	6	1	7	-	1	-
2002.....	5	3	9	7	5	6	7	2	2	-	-	1
2001.....	4	4	7	7	8	6	4	4	3	-	-	-
2000.....	9	4	7	5	6	6	8	5	2	2	1	-
1999.....	3	6	4	6	9	7	3	5	3	2	-	-
American Indian 1/, female												
2004.....	7	15	7	8	5	13	21	10	12	4	2	-
2003.....	8	6	8	12	14	5	10	16	13	4	-	-
2002.....	1	4	5	14	6	10	8	13	8	8	1	-
2001.....	3	4	3	6	5	12	12	8	8	6	1	-
2000.....	4	6	9	6	8	9	11	18	4	2	1	-
1999.....	-	5	9	4	7	8	12	13	6	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive diseases (I10-I13)												
Asian or Pacific Islander 2/, both sexes												
2004.....	41	56	70	74	95	140	170	204	131	57	19	-
2003.....	32	47	47	89	100	154	152	160	111	38	10	-
2002.....	33	45	58	72	94	121	152	147	113	33	9	-
2001.....	40	43	48	72	105	127	153	134	65	25	7	-
2000.....	36	23	39	58	85	118	148	104	75	46	9	-
1999.....	23	36	52	81	78	115	138	115	76	34	11	-
Asian or Pacific Islander 2/, male												
2004.....	33	33	44	37	44	61	73	69	47	18	3	-
2003.....	24	26	27	43	36	63	60	71	38	11	2	-
2002.....	21	29	38	51	49	58	70	52	36	13	3	-
2001.....	22	26	28	30	44	61	59	49	23	12	1	-
2000.....	27	13	28	29	50	58	58	42	24	16	4	-
1999.....	15	25	29	39	41	57	56	52	24	6	1	-
Asian or Pacific Islander 2/, female												
2004.....	8	23	26	37	51	79	97	135	84	39	16	-
2003.....	8	21	20	46	64	91	92	89	73	27	8	-
2002.....	12	16	20	21	45	63	82	95	77	20	6	-
2001.....	18	17	20	42	61	66	94	85	42	13	6	-
2000.....	9	10	11	29	35	60	90	62	51	30	5	-
1999.....	8	11	23	42	37	58	82	63	52	28	10	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Essential (primary) hypertension (I10)												
All races, both sexes												
2004.....	391	459	584	690	984	1,538	2,163	2,544	2,225	1,118	304	2
2003.....	376	458	558	701	993	1,525	2,137	2,340	2,119	988	297	-
2002.....	307	375	488	629	951	1,371	1,972	2,155	1,852	902	233	-
2001.....	298	382	481	658	885	1,419	1,833	2,028	1,672	825	206	-
2000.....	304	350	466	579	920	1,346	1,637	1,861	1,528	697	200	-
1999.....	270	330	464	636	861	1,323	1,609	1,796	1,449	661	171	-
All races, male												
2004.....	262	259	353	390	451	641	748	693	423	160	28	1
2003.....	241	279	321	401	469	562	689	544	413	131	30	-
2002.....	189	237	274	323	458	539	644	553	346	115	17	-
2001.....	194	232	276	348	425	575	580	517	328	111	16	-
2000.....	197	218	275	284	456	530	542	481	270	88	24	-
1999.....	155	201	261	343	417	518	535	444	268	94	7	-
All races, female												
2004.....	129	200	231	300	533	897	1,415	1,851	1,802	958	276	1
2003.....	135	179	237	300	524	963	1,448	1,796	1,706	857	267	-
2002.....	118	138	214	306	493	832	1,328	1,602	1,506	787	216	-
2001.....	104	150	205	310	460	844	1,253	1,511	1,344	714	190	-
2000.....	107	132	191	295	464	816	1,095	1,380	1,258	609	176	-
1999.....	115	129	203	293	444	805	1,074	1,352	1,181	567	164	-
White, both sexes												
2004.....	258	302	429	508	766	1,241	1,843	2,225	1,937	999	249	1
2003.....	254	303	392	496	721	1,213	1,782	2,088	1,886	880	252	-
2002.....	203	265	367	467	723	1,086	1,681	1,865	1,637	809	196	-
2001.....	191	254	338	470	665	1,168	1,542	1,767	1,495	735	173	-
2000.....	183	240	327	435	723	1,080	1,397	1,615	1,354	613	161	-
1999.....	159	222	331	468	684	1,058	1,370	1,546	1,288	577	141	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Essential (primary) hypertension (I10)												
White, male												
2004.....	182	174	272	296	355	511	641	612	374	132	25	-
2003.....	170	195	226	298	354	443	551	478	357	113	23	-
2002.....	132	174	212	244	355	429	531	473	305	103	12	-
2001.....	126	154	202	261	319	466	472	439	289	97	11	-
2000.....	123	151	193	223	360	429	448	413	232	73	18	-
1999.....	102	144	191	259	332	417	445	363	234	74	6	-
White, female												
2004.....	76	128	157	212	411	730	1,202	1,613	1,563	867	224	1
2003.....	84	108	166	198	367	770	1,231	1,610	1,529	767	229	-
2002.....	71	91	155	223	368	657	1,150	1,392	1,332	706	184	-
2001.....	65	100	136	209	346	702	1,070	1,328	1,206	638	162	-
2000.....	60	89	134	212	363	651	949	1,202	1,122	540	143	-
1999.....	57	78	140	209	352	641	925	1,183	1,054	503	135	-
Black, both sexes												
2004.....	121	142	140	160	191	263	262	263	245	99	44	1
2003.....	114	141	153	187	244	276	311	211	193	96	40	-
2002.....	99	103	105	145	205	254	256	243	181	77	33	-
2001.....	98	118	133	169	192	227	246	224	154	82	31	-
2000.....	111	106	130	133	177	233	202	214	153	75	35	-
1999.....	106	105	120	146	164	237	204	212	140	72	25	-
Black, male												
2004.....	71	77	74	80	82	117	87	65	34	24	2	1
2003.....	67	75	89	99	100	106	118	52	41	16	6	-
2002.....	55	59	52	69	91	96	96	65	32	8	5	-
2001.....	64	72	67	80	93	99	90	64	35	13	5	-
2000.....	66	65	77	56	85	84	77	58	33	12	4	-
1999.....	51	56	64	73	78	87	76	64	25	16	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Essential (primary) hypertension (I10)												
Black, female												
2004.....	50	65	66	80	109	146	175	198	211	75	42	-
2003.....	47	66	64	88	144	170	193	159	152	80	34	-
2002.....	44	44	53	76	114	158	160	178	149	69	28	-
2001.....	34	46	66	89	99	128	156	160	119	69	26	-
2000.....	45	41	53	77	92	149	125	156	120	63	31	-
1999.....	55	49	56	73	86	150	128	148	115	56	24	-
American Indian 1/, both sexes												
2004.....	2	5	3	6	5	3	12	3	4	2	2	-
2003.....	2	6	3	2	8	5	3	6	8	1	-	-
2002.....	1	2	4	4	5	5	3	6	1	5	1	-
2001.....	1	1	-	3	6	1	4	5	2	2	1	-
2000.....	2	2	5	2	3	4	3	7	4	2	2	-
1999.....	1	1	3	2	2	2	2	4	4	2	-	-
American Indian 1/, male												
2004.....	-	2	2	4	5	-	4	-	1	1	-	-
2003.....	1	5	3	-	4	5	1	1	2	-	-	-
2002.....	1	-	2	-	3	1	1	1	-	-	-	-
2001.....	1	-	-	1	4	-	1	3	-	-	-	-
2000.....	2	1	3	-	1	2	1	2	1	1	1	-
1999.....	1	-	1	2	1	1	1	2	2	2	-	-
American Indian 1/, female												
2004.....	2	3	1	2	-	3	8	3	3	1	2	-
2003.....	1	1	-	2	4	-	2	5	6	1	-	-
2002.....	-	2	2	4	2	4	2	5	1	5	1	-
2001.....	-	1	-	2	2	1	3	2	2	2	1	-
2000.....	-	1	2	2	2	2	2	5	3	1	1	-
1999.....	-	1	2	-	1	1	1	2	2	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Essential (primary) hypertension (I10)												
Asian or Pacific Islander 2/, both sexes												
2004.....	10	10	12	16	22	31	46	53	39	18	9	-
2003.....	6	8	10	16	20	31	41	35	32	11	5	-
2002.....	4	5	12	13	18	26	32	41	33	11	3	-
2001.....	8	9	10	16	22	23	41	32	21	6	1	-
2000.....	8	2	4	9	17	29	35	25	17	7	2	-
1999.....	4	2	10	20	11	26	33	34	17	10	5	-
Asian or Pacific Islander 2/, male												
2004.....	9	6	5	10	9	13	16	16	14	3	1	-
2003.....	3	4	3	4	11	8	19	13	13	2	1	-
2002.....	1	4	8	10	9	13	16	14	9	4	-	-
2001.....	3	6	7	6	9	10	17	11	4	1	-	-
2000.....	6	1	2	5	10	15	16	8	4	2	1	-
1999.....	1	1	5	9	6	13	13	15	7	2	-	-
Asian or Pacific Islander 2/, female												
2004.....	1	4	7	6	13	18	30	37	25	15	8	-
2003.....	3	4	7	12	9	23	22	22	19	9	4	-
2002.....	3	1	4	3	9	13	16	27	24	7	3	-
2001.....	5	3	3	10	13	13	24	21	17	5	1	-
2000.....	2	1	2	4	7	14	19	17	13	5	1	-
1999.....	3	1	5	11	5	13	20	19	10	8	5	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart disease (I11)												
All races, both sexes												
2004.....	1,760	2,012	1,873	1,867	2,134	2,722	3,676	4,053	3,411	1,510	394	3
2003.....	1,784	1,852	1,799	1,892	2,153	2,945	3,620	3,976	3,285	1,590	338	-
2002.....	1,616	1,661	1,659	1,807	2,088	2,816	3,505	3,716	3,115	1,442	314	4
2001.....	1,414	1,380	1,389	1,557	2,062	2,719	3,400	3,658	2,958	1,398	331	11
2000.....	1,152	1,249	1,423	1,570	2,035	2,694	3,268	3,684	2,886	1,330	286	3
1999.....	1,041	1,260	1,277	1,553	1,964	2,672	3,158	3,498	2,710	1,188	283	1
All races, male												
2004.....	1,185	1,363	1,221	1,146	1,157	1,213	1,291	1,167	746	231	48	3
2003.....	1,195	1,222	1,086	1,141	1,123	1,326	1,301	1,143	669	206	36	-
2002.....	1,094	1,107	1,052	1,046	1,042	1,259	1,250	1,004	629	194	44	4
2001.....	931	879	870	906	1,055	1,188	1,226	957	565	197	27	9
2000.....	811	789	895	909	1,062	1,135	1,134	960	545	181	21	2
1999.....	673	800	760	858	958	1,185	1,067	912	499	160	23	1
All races, female												
2004.....	575	649	652	721	977	1,509	2,385	2,886	2,665	1,279	346	-
2003.....	589	630	713	751	1,030	1,619	2,319	2,833	2,616	1,384	302	-
2002.....	522	554	607	761	1,046	1,557	2,255	2,712	2,486	1,248	270	-
2001.....	483	501	519	651	1,007	1,531	2,174	2,701	2,393	1,201	304	2
2000.....	341	460	528	661	973	1,559	2,134	2,724	2,341	1,149	265	1
1999.....	368	460	517	695	1,006	1,487	2,091	2,586	2,211	1,028	260	-
White, both sexes												
2004.....	1,037	1,261	1,214	1,291	1,510	2,039	2,952	3,453	2,954	1,321	318	3
2003.....	1,043	1,150	1,185	1,235	1,518	2,199	2,982	3,362	2,859	1,399	269	-
2002.....	951	1,036	1,040	1,185	1,474	2,109	2,785	3,134	2,701	1,265	261	1
2001.....	810	843	870	999	1,435	2,051	2,754	3,073	2,611	1,225	262	6
2000.....	647	721	867	1,032	1,379	2,012	2,614	3,122	2,487	1,132	236	1
1999.....	590	728	801	997	1,329	1,976	2,541	2,983	2,371	1,017	231	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart disease (I11)												
White, male												
2004.....	748	880	797	802	836	927	1,039	992	634	193	37	3
2003.....	762	804	750	761	799	1,003	1,075	960	559	175	30	-
2002.....	690	730	671	698	754	952	998	844	533	161	31	1
2001.....	574	571	573	613	760	905	983	786	493	162	14	5
2000.....	480	485	564	616	749	855	893	813	456	147	18	1
1999.....	409	488	499	556	667	883	852	757	430	129	19	-
White, female												
2004.....	289	381	417	489	674	1,112	1,913	2,461	2,320	1,128	281	-
2003.....	281	346	435	474	719	1,196	1,907	2,402	2,300	1,224	239	-
2002.....	261	306	369	487	720	1,157	1,787	2,290	2,168	1,104	230	-
2001.....	236	272	297	386	675	1,146	1,771	2,287	2,118	1,063	248	1
2000.....	167	236	303	416	630	1,157	1,721	2,309	2,031	985	218	-
1999.....	181	240	302	441	662	1,093	1,689	2,226	1,941	888	212	-
Black, both sexes												
2004.....	697	698	610	537	582	609	652	494	389	164	69	-
2003.....	710	662	572	601	583	666	574	535	369	170	65	-
2002.....	637	592	581	576	571	642	641	515	357	160	47	2
2001.....	571	506	487	517	578	605	571	526	315	156	65	5
2000.....	475	507	524	494	612	625	584	509	360	172	46	2
1999.....	439	499	446	521	584	637	558	456	299	156	46	1
Black, male												
2004.....	417	448	387	323	302	252	221	139	91	29	10	-
2003.....	411	393	310	352	304	284	202	151	93	26	6	-
2002.....	384	357	359	317	269	273	219	139	76	24	10	2
2001.....	338	288	275	274	272	252	214	149	62	28	12	4
2000.....	308	291	308	269	287	248	212	134	77	24	2	1
1999.....	254	292	245	281	266	272	191	132	60	30	3	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart disease (I11)												
Black, female												
2004.....	280	250	223	214	280	357	431	355	298	135	59	-
2003.....	299	269	262	249	279	382	372	384	276	144	59	-
2002.....	253	235	222	259	302	369	422	376	281	136	37	-
2001.....	233	218	212	243	306	353	357	377	253	128	53	1
2000.....	167	216	216	225	325	377	372	375	283	148	44	1
1999.....	185	207	201	240	318	365	367	324	239	126	43	-
American Indian 1/, both sexes												
2004.....	8	20	11	5	7	13	7	10	8	3	-	-
2003.....	11	13	12	13	5	7	7	9	9	3	-	-
2002.....	3	3	7	10	3	8	5	5	6	-	-	1
2001.....	6	6	4	7	3	10	6	3	5	4	-	-
2000.....	10	5	6	6	5	5	11	6	2	2	-	-
1999.....	2	9	5	3	7	6	9	7	2	-	-	-
American Indian 1/, male												
2004.....	4	12	10	2	5	6	3	3	2	-	-	-
2003.....	5	9	8	6	2	4	2	-	3	-	-	-
2002.....	2	1	4	4	-	4	3	-	1	-	-	1
2001.....	3	3	3	5	2	4	2	-	1	-	-	-
2000.....	7	2	3	4	4	3	5	1	1	1	-	-
1999.....	2	5	-	2	5	3	2	1	1	-	-	-
American Indian 1/, female												
2004.....	4	8	1	3	2	7	4	7	6	3	-	-
2003.....	6	4	4	7	3	3	5	9	6	3	-	-
2002.....	1	2	3	6	3	4	2	5	5	-	-	-
2001.....	3	3	1	2	1	6	4	3	4	4	-	-
2000.....	3	3	3	2	1	2	6	5	1	1	-	-
1999.....	-	4	5	1	2	3	7	6	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart disease (I11)												
Asian or Pacific Islander 2/, both sexes												
2004.....	18	33	38	34	35	61	65	96	60	22	7	-
2003.....	20	27	30	43	47	73	57	70	48	18	4	-
2002.....	25	30	31	36	40	57	74	62	51	17	6	-
2001.....	27	25	28	34	46	53	69	56	27	13	4	-
2000.....	20	16	26	38	39	52	59	47	37	24	4	-
1999.....	10	24	25	32	44	53	50	52	38	15	6	-
Asian or Pacific Islander 2/, male												
2004.....	16	23	27	19	14	28	28	33	19	9	1	-
2003.....	17	16	18	22	18	35	22	32	14	5	-	-
2002.....	18	19	18	27	19	30	30	21	19	9	3	-
2001.....	16	17	19	14	21	27	27	22	9	7	1	-
2000.....	16	11	20	20	22	29	24	12	11	9	1	-
1999.....	8	15	16	19	20	27	22	22	8	1	1	-
Asian or Pacific Islander 2/, female												
2004.....	2	10	11	15	21	33	37	63	41	13	6	-
2003.....	3	11	12	21	29	38	35	38	34	13	4	-
2002.....	7	11	13	9	21	27	44	41	32	8	3	-
2001.....	11	8	9	20	25	26	42	34	18	6	3	-
2000.....	4	5	6	18	17	23	35	35	26	15	3	-
1999.....	2	9	9	13	24	26	28	30	30	14	5	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive renal disease (I12)												
All races, both sexes												
2004.....	267	356	440	602	876	1,317	1,801	1,739	1,209	424	85	2
2003.....	255	327	412	571	839	1,339	1,648	1,609	1,114	360	77	-
2002.....	234	306	357	497	845	1,221	1,574	1,646	1,063	374	60	-
2001.....	248	248	351	509	780	1,219	1,522	1,578	901	311	51	-
2000.....	222	247	377	515	770	1,161	1,479	1,412	847	320	42	-
1999.....	200	208	316	510	792	1,077	1,315	1,153	759	281	47	-
All races, male												
2004.....	164	188	236	301	393	600	793	670	387	127	15	2
2003.....	154	178	211	275	390	601	670	619	360	109	12	-
2002.....	133	161	178	240	398	564	658	629	361	100	11	-
2001.....	136	122	154	255	351	525	602	612	282	81	12	-
2000.....	134	134	191	220	348	519	613	510	299	67	8	-
1999.....	131	115	155	246	366	478	536	434	253	67	13	-
All races, female												
2004.....	103	168	204	301	483	717	1,008	1,069	822	297	70	-
2003.....	101	149	201	296	449	738	978	990	754	251	65	-
2002.....	101	145	179	257	447	657	916	1,017	702	274	49	-
2001.....	112	126	197	254	429	694	920	966	619	230	39	-
2000.....	88	113	186	295	422	642	866	902	548	253	34	-
1999.....	69	93	161	264	426	599	779	719	506	214	34	-
White, both sexes												
2004.....	102	150	198	334	540	922	1,399	1,443	987	335	67	2
2003.....	104	135	209	321	527	972	1,342	1,332	915	294	58	-
2002.....	99	124	169	276	561	881	1,214	1,338	865	312	53	-
2001.....	89	102	169	283	521	881	1,185	1,291	748	263	38	-
2000.....	86	105	179	271	477	845	1,176	1,157	702	243	35	-
1999.....	76	82	129	287	510	761	995	924	616	220	37	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive renal disease (I12)												
White, male												
2004.....	60	88	106	168	254	425	633	577	333	106	11	2
2003.....	56	75	104	156	254	467	569	520	307	91	10	-
2002.....	56	67	89	135	272	420	521	541	300	82	10	-
2001.....	46	52	68	134	248	398	484	517	245	64	10	-
2000.....	61	61	98	117	226	394	512	426	251	50	6	-
1999.....	44	49	64	144	247	349	419	359	209	51	10	-
White, female												
2004.....	42	62	92	166	286	497	766	866	654	229	56	-
2003.....	48	60	105	165	273	505	773	812	608	203	48	-
2002.....	43	57	80	141	289	461	693	797	565	230	43	-
2001.....	43	50	101	149	273	483	701	774	503	199	28	-
2000.....	25	44	81	154	251	451	664	731	451	193	29	-
1999.....	32	33	65	143	263	412	576	565	407	169	27	-
Black, both sexes												
2004.....	155	189	222	245	303	351	344	254	195	74	16	-
2003.....	146	183	190	222	277	324	262	238	176	59	19	-
2002.....	129	171	175	199	254	307	321	271	173	55	7	-
2001.....	154	138	169	205	226	290	301	257	134	44	12	-
2000.....	127	136	187	232	264	287	260	226	130	67	6	-
1999.....	120	118	170	198	259	295	276	206	127	53	10	-
Black, male												
2004.....	99	95	120	126	122	155	133	78	44	15	3	-
2003.....	95	98	99	104	127	118	82	84	46	16	2	-
2002.....	73	87	78	94	110	133	119	74	53	18	1	-
2001.....	87	67	82	112	91	106	107	84	26	14	2	-
2000.....	68	72	89	98	106	113	86	69	42	15	1	-
1999.....	84	59	82	93	105	121	102	67	37	14	3	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive renal disease (I12)												
Black, female												
2004.....	56	94	102	119	181	196	211	176	151	59	13	-
2003.....	51	85	91	118	150	206	180	154	130	43	17	-
2002.....	56	84	97	105	144	174	202	197	120	37	6	-
2001.....	67	71	87	93	135	184	194	173	108	30	10	-
2000.....	59	64	98	134	158	174	174	157	88	52	5	-
1999.....	36	59	88	105	154	174	174	139	90	39	7	-
American Indian 1/, both sexes												
2004.....	3	6	4	5	7	5	9	1	3	1	-	-
2003.....	3	1	7	6	11	2	5	2	2	-	-	-
2002.....	2	2	3	6	3	3	6	2	2	3	-	-
2001.....	-	1	4	3	4	6	5	2	4	-	-	-
2000.....	1	2	5	2	6	5	4	6	-	-	-	-
1999.....	-	-	5	3	3	3	4	6	2	-	-	-
American Indian 1/, male												
2004.....	2	2	1	2	4	3	2	1	-	1	-	-
2003.....	2	1	3	3	4	1	2	-	1	-	-	-
2002.....	2	2	3	2	2	1	3	-	1	-	-	-
2001.....	-	1	2	1	2	2	1	-	2	-	-	-
2000.....	-	-	1	1	1	1	1	1	-	-	-	-
1999.....	-	-	3	1	1	1	-	1	-	-	-	-
American Indian 1/, female												
2004.....	1	4	3	3	3	2	7	-	3	-	-	-
2003.....	1	-	4	3	7	1	3	2	1	-	-	-
2002.....	-	-	-	4	1	2	3	2	1	3	-	-
2001.....	-	-	2	2	2	4	4	2	2	-	-	-
2000.....	1	2	4	1	5	4	3	5	-	-	-	-
1999.....	-	-	2	2	2	2	4	5	2	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive renal disease (I12)												
Asian or Pacific Islander 2/, both sexes												
2004.....	7	11	16	18	26	39	49	41	24	14	2	-
2003.....	2	8	6	22	24	41	39	37	21	7	-	-
2002.....	4	9	10	16	27	30	33	35	23	4	-	-
2001.....	5	7	9	18	29	42	31	28	15	4	1	-
2000.....	8	4	6	10	23	24	39	23	15	10	1	-
1999.....	4	8	12	22	20	18	40	17	14	8	-	-
Asian or Pacific Islander 2/, male												
2004.....	3	3	9	5	13	17	25	14	10	5	1	-
2003.....	1	4	5	12	5	15	17	15	6	2	-	-
2002.....	2	5	8	9	14	10	15	14	7	-	-	-
2001.....	3	2	2	8	10	19	10	11	9	3	-	-
2000.....	5	1	3	4	15	11	14	14	6	2	1	-
1999.....	3	7	6	8	13	7	15	7	7	2	-	-
Asian or Pacific Islander 2/, female												
2004.....	4	8	7	13	13	22	24	27	14	9	1	-
2003.....	1	4	1	10	19	26	22	22	15	5	-	-
2002.....	2	4	2	7	13	20	18	21	16	4	-	-
2001.....	2	5	7	10	19	23	21	17	6	1	1	-
2000.....	3	3	3	6	8	13	25	9	9	8	-	-
1999.....	1	1	6	14	7	11	25	10	7	6	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart and renal disease (I13)												
All races, both sexes												
2004.....	105	111	160	161	218	330	524	548	461	160	37	-
2003.....	114	136	131	154	247	386	517	560	456	167	26	-
2002.....	90	134	139	156	246	363	503	511	386	150	32	-
2001.....	82	109	138	169	248	374	481	499	348	155	24	-
2000.....	79	103	105	149	268	376	487	494	361	153	24	-
1999.....	76	116	146	201	269	487	600	600	411	162	36	-
All races, male												
2004.....	63	55	103	81	108	157	220	188	139	30	10	-
2003.....	67	86	73	75	114	174	208	207	128	45	6	-
2002.....	56	82	76	67	131	158	199	171	108	38	4	-
2001.....	53	61	65	94	112	158	181	174	102	35	3	-
2000.....	44	57	51	74	132	152	198	188	109	37	7	-
1999.....	50	66	80	99	124	221	239	205	123	41	5	-
All races, female												
2004.....	42	56	57	80	110	173	304	360	322	130	27	-
2003.....	47	50	58	79	133	212	309	353	328	122	20	-
2002.....	34	52	63	89	115	205	304	340	278	112	28	-
2001.....	29	48	73	75	136	216	300	325	246	120	21	-
2000.....	35	46	54	75	136	224	289	306	252	116	17	-
1999.....	26	50	66	102	145	266	361	395	288	121	31	-
White, both sexes												
2004.....	34	43	76	80	132	230	405	460	386	128	33	-
2003.....	28	47	64	75	154	279	406	442	391	144	17	-
2002.....	21	44	63	73	145	254	401	401	319	127	24	-
2001.....	35	45	71	87	152	258	384	395	295	132	16	-
2000.....	33	46	44	72	173	263	361	399	305	129	12	-
1999.....	31	47	63	101	148	343	463	473	351	129	29	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart and renal disease (I13)												
White, male												
2004.....	19	20	54	42	71	109	176	162	116	25	9	-
2003.....	18	30	38	43	75	135	172	161	115	39	2	-
2002.....	11	30	35	32	82	116	160	135	90	32	2	-
2001.....	21	24	32	50	68	114	152	146	90	29	2	-
2000.....	19	25	21	38	88	120	153	150	90	28	2	-
1999.....	19	26	34	54	69	159	199	167	108	31	5	-
White, female												
2004.....	15	23	22	38	61	121	229	298	270	103	24	-
2003.....	10	17	26	32	79	144	234	281	276	105	15	-
2002.....	10	14	28	41	63	138	241	266	229	95	22	-
2001.....	14	21	39	37	84	144	232	249	205	103	14	-
2000.....	14	21	23	34	85	143	208	249	215	101	10	-
1999.....	12	21	29	47	79	184	264	306	243	98	24	-
Black, both sexes												
2004.....	65	65	78	73	74	90	105	73	67	28	3	-
2003.....	82	84	65	71	83	97	95	100	54	21	7	-
2002.....	69	89	71	75	92	101	88	99	60	22	8	-
2001.....	47	62	64	78	88	106	84	84	51	21	7	-
2000.....	46	55	58	75	89	99	110	82	50	19	10	-
1999.....	40	66	78	91	114	122	122	114	52	32	7	-
Black, male												
2004.....	39	33	46	34	29	45	38	19	19	3	1	-
2003.....	46	54	33	27	36	34	33	35	7	4	2	-
2002.....	45	51	37	29	42	37	30	32	17	6	2	-
2001.....	32	36	31	42	40	39	24	22	11	5	1	-
2000.....	25	31	27	36	41	29	40	29	16	6	4	-
1999.....	28	37	44	41	51	50	34	29	13	9	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart and renal disease (I13)												
Black, female												
2004.....	26	32	32	39	45	45	67	54	48	25	2	-
2003.....	36	30	32	44	47	63	62	65	47	17	5	-
2002.....	24	38	34	46	50	64	58	67	43	16	6	-
2001.....	15	26	33	36	48	67	60	62	40	16	6	-
2000.....	21	24	31	39	48	70	70	53	34	13	6	-
1999.....	12	29	34	50	63	72	88	85	39	23	7	-
American Indian 1/, both sexes												
2004.....	-	1	2	2	-	1	4	1	-	1	-	-
2003.....	-	1	1	-	1	1	1	-	1	-	1	-
2002.....	-	-	-	1	-	-	1	2	1	-	-	-
2001.....	-	-	2	-	-	1	1	2	-	-	-	-
2000.....	-	1	-	1	-	1	1	4	-	-	-	-
1999.....	-	1	-	2	4	4	-	1	1	-	-	-
American Indian 1/, male												
2004.....	-	1	-	2	-	-	2	1	-	1	-	-
2003.....	-	-	1	-	1	-	1	-	1	-	1	-
2002.....	-	-	-	1	-	-	-	1	-	-	-	-
2001.....	-	-	2	-	-	-	-	1	-	-	-	-
2000.....	-	1	-	-	-	-	1	1	-	-	-	-
1999.....	-	1	-	1	2	2	-	1	-	-	-	-
American Indian 1/, female												
2004.....	-	-	2	-	-	1	2	-	-	-	-	-
2003.....	-	1	-	-	-	1	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	1	1	1	-	-	-
2001.....	-	-	-	-	-	1	1	1	-	-	-	-
2000.....	-	-	-	1	-	1	-	3	-	-	-	-
1999.....	-	-	-	1	2	2	-	-	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart and renal disease (I13)												
Asian or Pacific Islander 2/, both sexes												
2004.....	6	2	4	6	12	9	10	14	8	3	1	-
2003.....	4	4	1	8	9	9	15	18	10	2	1	-
2002.....	-	1	5	7	9	8	13	9	6	1	-	-
2001.....	-	2	1	4	8	9	12	18	2	2	1	-
2000.....	-	1	3	1	6	13	15	9	6	5	2	-
1999.....	5	2	5	7	3	18	15	12	7	1	-	-
Asian or Pacific Islander 2/, male												
2004.....	5	1	3	3	8	3	4	6	4	1	-	-
2003.....	3	2	1	5	2	5	2	11	5	2	1	-
2002.....	-	1	4	5	7	5	9	3	1	-	-	-
2001.....	-	1	-	2	4	5	5	5	1	1	-	-
2000.....	-	-	3	-	3	3	4	8	3	3	1	-
1999.....	3	2	2	3	2	10	6	8	2	1	-	-
Asian or Pacific Islander 2/, female												
2004.....	1	1	1	3	4	6	6	8	4	2	1	-
2003.....	1	2	-	3	7	4	13	7	5	-	-	-
2002.....	-	-	1	2	2	3	4	6	5	1	-	-
2001.....	-	1	1	2	4	4	7	13	1	1	1	-
2000.....	-	1	-	1	3	10	11	1	3	2	1	-
1999.....	2	-	3	4	1	8	9	4	5	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Ischemic heart diseases (I20-I25)												
All races, both sexes												
2004.....	15,192	20,437	25,565	30,914	42,188	60,283	78,845	76,562	55,554	22,517	5,123	26
2003.....	15,372	20,907	26,470	32,648	46,232	66,062	83,308	81,998	58,794	23,939	5,433	29
2002.....	15,683	20,997	26,265	34,336	49,531	69,769	85,622	84,082	59,567	24,211	5,528	37
2001.....	15,612	20,270	26,304	35,502	51,961	73,082	86,772	84,880	59,558	24,206	5,665	48
2000.....	15,252	20,724	27,013	37,257	55,628	76,569	87,123	86,890	60,269	24,477	5,683	48
1999.....	15,251	20,575	27,802	40,080	58,760	79,379	89,917	88,158	60,502	24,983	5,733	39
All races, male												
2004.....	11,544	15,004	17,864	20,392	25,881	34,272	39,319	31,859	17,650	5,155	831	23
2003.....	11,820	15,342	18,425	21,547	28,452	37,191	41,382	33,563	18,269	5,232	808	22
2002.....	11,969	15,350	18,306	22,573	30,563	39,154	42,085	34,073	18,199	5,353	821	27
2001.....	11,818	14,794	18,212	23,181	31,891	40,894	42,059	33,466	17,702	5,124	814	33
2000.....	11,548	15,154	18,641	24,225	34,329	42,452	42,295	34,024	17,603	5,216	847	34
1999.....	11,614	15,021	19,102	26,184	36,003	44,041	43,264	34,168	17,472	5,245	841	31
All races, female												
2004.....	3,648	5,433	7,701	10,522	16,307	26,011	39,526	44,703	37,904	17,362	4,292	3
2003.....	3,552	5,565	8,045	11,101	17,780	28,871	41,926	48,435	40,525	18,707	4,625	7
2002.....	3,714	5,647	7,959	11,763	18,968	30,615	43,537	50,009	41,368	18,858	4,707	10
2001.....	3,794	5,476	8,092	12,321	20,070	32,188	44,713	51,414	41,856	19,082	4,851	15
2000.....	3,704	5,570	8,372	13,032	21,299	34,117	44,828	52,866	42,666	19,261	4,836	14
1999.....	3,637	5,554	8,700	13,896	22,757	35,338	46,653	53,990	43,030	19,738	4,892	8
White, both sexes												
2004.....	11,907	16,455	21,072	25,535	35,972	53,012	71,100	70,252	51,129	20,599	4,505	17
2003.....	12,085	17,028	21,621	27,015	39,524	58,115	75,130	75,347	54,228	21,994	4,730	23
2002.....	12,445	17,082	21,495	28,561	42,708	61,414	77,489	77,341	54,938	22,175	4,834	28
2001.....	12,441	16,436	21,493	29,456	44,934	64,552	78,643	78,104	55,084	22,157	4,976	30
2000.....	12,265	16,873	22,082	31,304	48,138	67,860	79,215	79,909	55,931	22,506	5,046	36
1999.....	12,272	16,733	22,933	33,894	50,959	70,631	81,999	81,066	56,125	22,985	5,122	27

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Ischemic heart diseases (I20-I25)												
White, male												
2004.....	9,321	12,364	15,034	17,216	22,506	30,687	35,987	29,562	16,329	4,703	706	15
2003.....	9,555	12,813	15,409	18,245	24,721	33,242	37,693	31,182	17,003	4,783	688	16
2002.....	9,771	12,781	15,313	19,219	26,909	35,116	38,536	31,656	16,828	4,882	701	21
2001.....	9,682	12,284	15,253	19,729	28,066	36,733	38,597	31,119	16,373	4,664	698	21
2000.....	9,559	12,586	15,635	20,885	30,317	38,199	38,901	31,567	16,346	4,726	729	27
1999.....	9,583	12,487	16,061	22,652	31,841	39,837	39,986	31,607	16,126	4,783	737	21
White, female												
2004.....	2,586	4,091	6,038	8,319	13,466	22,325	35,113	40,690	34,800	15,896	3,799	2
2003.....	2,530	4,215	6,212	8,770	14,803	24,873	37,437	44,165	37,225	17,211	4,042	7
2002.....	2,674	4,301	6,182	9,342	15,799	26,298	38,953	45,685	38,110	17,293	4,133	7
2001.....	2,759	4,152	6,240	9,727	16,868	27,819	40,046	46,985	38,711	17,493	4,278	9
2000.....	2,706	4,287	6,447	10,419	17,821	29,661	40,314	48,342	39,585	17,780	4,317	9
1999.....	2,689	4,246	6,872	11,242	19,118	30,794	42,013	49,459	39,999	18,202	4,385	6
Black, both sexes												
2004.....	2,911	3,480	3,916	4,567	5,286	6,090	6,280	4,966	3,586	1,595	542	8
2003.....	2,881	3,377	4,168	4,795	5,720	6,698	6,682	5,361	3,732	1,645	608	4
2002.....	2,866	3,441	4,166	4,920	5,815	6,962	6,757	5,534	3,808	1,717	602	9
2001.....	2,789	3,349	4,203	5,210	6,025	7,193	6,783	5,626	3,775	1,772	600	16
2000.....	2,624	3,383	4,302	5,124	6,495	7,298	6,668	5,891	3,644	1,676	568	12
1999.....	2,642	3,392	4,240	5,395	6,699	7,402	6,669	5,931	3,697	1,762	546	12
Black, male												
2004.....	1,938	2,252	2,434	2,647	2,851	2,929	2,594	1,710	979	332	100	7
2003.....	1,938	2,158	2,544	2,750	3,129	3,289	2,897	1,792	935	335	89	4
2002.....	1,896	2,219	2,574	2,821	3,070	3,255	2,841	1,850	1,022	357	88	6
2001.....	1,847	2,133	2,536	2,927	3,247	3,406	2,797	1,820	1,046	362	99	11
2000.....	1,701	2,227	2,561	2,796	3,409	3,451	2,748	1,939	957	369	104	7
1999.....	1,767	2,197	2,609	3,038	3,505	3,433	2,652	1,999	1,022	375	86	10

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Ischemic heart diseases (I20-I25)												
Black, female												
2004.....	973	1,228	1,482	1,920	2,435	3,161	3,686	3,256	2,607	1,263	442	1
2003.....	943	1,219	1,624	2,045	2,591	3,409	3,785	3,569	2,797	1,310	519	-
2002.....	970	1,222	1,592	2,099	2,745	3,707	3,916	3,684	2,786	1,360	514	3
2001.....	942	1,216	1,667	2,283	2,778	3,787	3,986	3,806	2,729	1,410	501	5
2000.....	923	1,156	1,741	2,328	3,086	3,847	3,920	3,952	2,687	1,307	464	5
1999.....	875	1,195	1,631	2,357	3,194	3,969	4,017	3,932	2,675	1,387	460	2
American Indian 1/, both sexes												
2004.....	116	146	166	209	201	218	242	200	112	49	11	-
2003.....	120	147	193	225	225	247	218	185	97	35	16	2
2002.....	112	144	179	205	221	254	211	154	101	45	10	-
2001.....	119	149	175	204	224	207	199	153	83	31	11	-
2000.....	101	135	167	229	185	246	192	156	113	45	7	-
1999.....	101	133	153	206	241	248	205	166	86	33	9	-
American Indian 1/, male												
2004.....	79	110	107	147	112	123	123	82	35	12	1	-
2003.....	93	104	125	138	143	133	115	80	38	11	6	2
2002.....	87	97	127	132	129	135	97	78	31	11	2	-
2001.....	82	111	122	121	138	112	90	67	25	10	2	-
2000.....	73	91	110	143	114	138	93	64	38	15	-	-
1999.....	73	95	96	134	141	140	94	59	30	6	2	-
American Indian 1/, female												
2004.....	37	36	59	62	89	95	119	118	77	37	10	-
2003.....	27	43	68	87	82	114	103	105	59	24	10	-
2002.....	25	47	52	73	92	119	114	76	70	34	8	-
2001.....	37	38	53	83	86	95	109	86	58	21	9	-
2000.....	28	44	57	86	71	108	99	92	75	30	7	-
1999.....	28	38	57	72	100	108	111	107	56	27	7	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Ischemic heart diseases (I20-I25)												
Asian or Pacific Islander 2/, both sexes												
2004.....	258	356	411	603	729	963	1,223	1,144	727	274	65	1
2003.....	286	355	488	613	763	1,002	1,278	1,105	737	265	79	-
2002.....	260	330	425	650	787	1,139	1,165	1,053	720	274	82	-
2001.....	263	336	433	632	778	1,130	1,147	997	616	246	78	2
2000.....	262	333	462	600	810	1,165	1,048	934	581	250	62	-
1999.....	236	317	476	585	861	1,098	1,044	995	594	203	56	-
Asian or Pacific Islander 2/, male												
2004.....	206	278	289	382	412	533	615	505	307	108	24	1
2003.....	234	267	347	414	459	527	677	509	293	103	25	-
2002.....	215	253	292	401	455	648	611	489	318	103	30	-
2001.....	207	266	301	404	440	643	575	460	258	88	15	1
2000.....	215	250	335	401	489	664	553	454	262	106	14	-
1999.....	191	242	336	360	516	631	532	503	294	81	16	-
Asian or Pacific Islander 2/, female												
2004.....	52	78	122	221	317	430	608	639	420	166	41	-
2003.....	52	88	141	199	304	475	601	596	444	162	54	-
2002.....	45	77	133	249	332	491	554	564	402	171	52	-
2001.....	56	70	132	228	338	487	572	537	358	158	63	1
2000.....	47	83	127	199	321	501	495	480	319	144	48	-
1999.....	45	75	140	225	345	467	512	492	300	122	40	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute myocardial infarction (I21-I22)												
All races, both sexes												
2004.....	6,167	8,346	10,412	12,357	15,970	21,573	26,464	24,512	16,524	5,939	1,277	6
2003.....	6,328	8,769	10,998	13,108	17,815	24,265	28,843	27,055	17,795	6,527	1,357	5
2002.....	6,564	8,758	11,175	14,230	19,773	26,154	30,436	28,423	18,352	6,623	1,320	5
2001.....	6,720	8,704	11,126	14,690	20,690	27,766	31,488	29,242	18,525	6,606	1,354	10
2000.....	6,647	9,030	11,983	15,898	22,561	30,059	32,151	29,922	18,701	6,755	1,367	8
1999.....	6,598	9,123	12,283	17,333	24,108	31,053	33,658	30,711	18,472	6,765	1,349	7
All races, male												
2004.....	4,643	6,107	7,235	8,017	9,641	11,987	12,906	9,983	5,286	1,393	233	4
2003.....	4,854	6,386	7,586	8,531	10,759	13,324	14,026	10,866	5,704	1,469	217	4
2002.....	4,929	6,287	7,679	9,214	12,050	14,316	14,603	11,478	5,741	1,559	213	4
2001.....	5,012	6,271	7,587	9,446	12,517	15,151	14,974	11,411	5,778	1,576	219	6
2000.....	4,946	6,490	8,148	10,177	13,646	16,295	15,336	11,765	5,701	1,539	238	5
1999.....	5,039	6,654	8,375	11,154	14,554	16,991	15,952	12,066	5,566	1,523	202	2
All races, female												
2004.....	1,524	2,239	3,177	4,340	6,329	9,586	13,558	14,529	11,238	4,546	1,044	2
2003.....	1,474	2,383	3,412	4,577	7,056	10,941	14,817	16,189	12,091	5,058	1,140	1
2002.....	1,635	2,471	3,496	5,016	7,723	11,838	15,833	16,945	12,611	5,064	1,107	1
2001.....	1,708	2,433	3,539	5,244	8,173	12,615	16,514	17,831	12,747	5,030	1,135	4
2000.....	1,701	2,540	3,835	5,721	8,915	13,764	16,815	18,157	13,000	5,216	1,129	3
1999.....	1,559	2,469	3,908	6,179	9,554	14,062	17,706	18,645	12,906	5,242	1,147	5
White, both sexes												
2004.....	4,964	6,913	8,829	10,405	13,719	19,121	23,898	22,418	15,122	5,360	1,085	4
2003.....	5,096	7,321	9,248	11,079	15,416	21,479	25,961	24,802	16,359	5,905	1,148	4
2002.....	5,374	7,341	9,380	12,067	17,184	23,108	27,457	25,992	16,780	5,985	1,116	4
2001.....	5,484	7,257	9,369	12,411	18,058	24,674	28,512	26,793	16,980	5,981	1,171	6
2000.....	5,458	7,574	10,085	13,575	19,710	26,733	29,233	27,383	17,270	6,112	1,174	7
1999.....	5,500	7,659	10,425	14,982	21,124	27,778	30,704	28,102	17,028	6,167	1,171	5

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute myocardial infarction (I21-I22)												
White, male												
2004.....	3,856	5,175	6,297	6,899	8,446	10,866	11,836	9,230	4,888	1,255	195	3
2003.....	4,019	5,478	6,526	7,397	9,464	11,962	12,739	10,067	5,274	1,330	177	3
2002.....	4,146	5,400	6,625	8,015	10,737	12,884	13,315	10,585	5,264	1,406	175	3
2001.....	4,225	5,373	6,564	8,205	11,150	13,690	13,744	10,533	5,289	1,410	187	5
2000.....	4,206	5,558	7,028	8,918	12,171	14,738	14,094	10,840	5,246	1,363	210	4
1999.....	4,319	5,690	7,277	9,830	13,039	15,479	14,755	11,087	5,116	1,375	166	1
White, female												
2004.....	1,108	1,738	2,532	3,506	5,273	8,255	12,062	13,188	10,234	4,105	890	1
2003.....	1,077	1,843	2,722	3,682	5,952	9,517	13,222	14,735	11,085	4,575	971	1
2002.....	1,228	1,941	2,755	4,052	6,447	10,224	14,142	15,407	11,516	4,579	941	1
2001.....	1,259	1,884	2,805	4,206	6,908	10,984	14,768	16,260	11,691	4,571	984	1
2000.....	1,252	2,016	3,057	4,657	7,539	11,995	15,139	16,543	12,024	4,749	964	3
1999.....	1,181	1,969	3,148	5,152	8,085	12,299	15,949	17,015	11,912	4,792	1,005	4
Black, both sexes												
2004.....	1,067	1,264	1,377	1,639	1,908	2,074	2,104	1,677	1,128	494	176	1
2003.....	1,071	1,254	1,471	1,719	2,042	2,359	2,352	1,828	1,197	545	183	1
2002.....	1,058	1,255	1,569	1,823	2,207	2,527	2,461	2,002	1,291	572	180	1
2001.....	1,068	1,265	1,518	1,955	2,261	2,604	2,478	2,028	1,309	544	157	4
2000.....	1,038	1,277	1,658	1,982	2,465	2,761	2,475	2,151	1,207	542	178	1
1999.....	971	1,271	1,611	2,027	2,569	2,773	2,525	2,183	1,254	540	159	2
Black, male												
2004.....	690	794	803	921	997	922	824	589	287	107	33	-
2003.....	702	771	871	935	1,076	1,131	997	603	325	106	34	1
2002.....	678	769	897	994	1,109	1,157	1,016	690	356	123	25	1
2001.....	658	768	858	1,042	1,148	1,181	987	679	377	134	28	1
2000.....	626	798	950	1,041	1,254	1,254	1,016	727	348	128	27	1
1999.....	624	819	937	1,120	1,284	1,246	984	765	356	128	30	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute myocardial infarction (I21-I22)												
Black, female												
2004.....	377	470	574	718	911	1,152	1,280	1,088	841	387	143	1
2003.....	369	483	600	784	966	1,228	1,355	1,225	872	439	149	-
2002.....	380	486	672	829	1,098	1,370	1,445	1,312	935	449	155	-
2001.....	410	497	660	913	1,113	1,423	1,491	1,349	932	410	129	3
2000.....	412	479	708	941	1,211	1,507	1,459	1,424	859	414	151	-
1999.....	347	452	674	907	1,285	1,527	1,541	1,418	898	412	129	1
American Indian 1/, both sexes												
2004.....	41	49	63	87	79	59	67	71	40	9	2	-
2003.....	45	58	94	92	80	89	83	56	25	10	2	-
2002.....	41	55	77	79	86	97	69	51	34	9	1	-
2001.....	57	53	72	76	86	69	70	62	28	15	5	-
2000.....	42	54	59	97	66	96	70	50	38	15	2	-
1999.....	39	61	55	90	93	92	85	64	24	11	4	-
American Indian 1/, male												
2004.....	27	40	36	60	49	30	35	21	13	1	-	-
2003.....	38	38	63	58	47	48	42	25	11	3	-	-
2002.....	34	42	57	49	41	44	29	25	11	3	1	-
2001.....	41	40	49	44	52	38	35	33	11	6	-	-
2000.....	30	38	35	64	37	54	33	22	12	7	-	-
1999.....	27	45	32	60	53	45	39	28	7	1	1	-
American Indian 1/, female												
2004.....	14	9	27	27	30	29	32	50	27	8	2	-
2003.....	7	20	31	34	33	41	41	31	14	7	2	-
2002.....	7	13	20	30	45	53	40	26	23	6	-	-
2001.....	16	13	23	32	34	31	35	29	17	9	5	-
2000.....	12	16	24	33	29	42	37	28	26	8	2	-
1999.....	12	16	23	30	40	47	46	36	17	10	3	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute myocardial infarction (I21-I22)												
Asian or Pacific Islander 2/, both sexes												
2004.....	95	120	143	226	264	319	395	346	234	76	14	1
2003.....	116	136	185	218	277	338	447	369	214	67	24	-
2002.....	91	107	149	261	296	422	449	378	247	57	23	-
2001.....	111	129	167	248	285	419	428	359	208	66	21	-
2000.....	109	125	181	244	320	469	373	338	186	86	13	-
1999.....	88	132	192	234	322	410	344	362	166	47	15	-
Asian or Pacific Islander 2/, male												
2004.....	70	98	99	137	149	169	211	143	98	30	5	1
2003.....	95	99	126	141	172	183	248	171	94	30	6	-
2002.....	71	76	100	156	163	231	243	178	110	27	12	-
2001.....	88	90	116	155	167	242	208	166	101	26	4	-
2000.....	84	96	135	154	184	249	193	176	95	41	1	-
1999.....	69	100	129	144	178	221	174	186	87	19	5	-
Asian or Pacific Islander 2/, female												
2004.....	25	22	44	89	115	150	184	203	136	46	9	-
2003.....	21	37	59	77	105	155	199	198	120	37	18	-
2002.....	20	31	49	105	133	191	206	200	137	30	11	-
2001.....	23	39	51	93	118	177	220	193	107	40	17	-
2000.....	25	29	46	90	136	220	180	162	91	45	12	-
1999.....	19	32	63	90	144	189	170	176	79	28	10	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute ischemic heart diseases (I24)												
All races, both sexes												
2004.....	136	184	175	179	220	308	362	327	246	111	33	1
2003.....	171	198	205	267	312	389	448	447	337	129	28	1
2002.....	226	210	250	289	342	420	478	452	339	153	27	-
2001.....	187	200	263	281	364	397	474	455	317	142	31	-
2000.....	189	219	265	294	350	440	466	439	318	136	35	1
1999.....	160	209	239	294	380	406	449	474	296	116	22	-
All races, male												
2004.....	96	132	119	124	130	154	175	143	76	21	6	1
2003.....	122	137	141	173	188	220	194	159	93	21	9	1
2002.....	162	153	167	193	197	216	228	172	87	35	4	-
2001.....	141	134	186	180	213	206	218	165	72	25	7	-
2000.....	141	163	178	194	204	232	214	153	68	28	5	1
1999.....	109	159	156	191	219	229	194	170	75	19	2	-
All races, female												
2004.....	40	52	56	55	90	154	187	184	170	90	27	-
2003.....	49	61	64	94	124	169	254	288	244	108	19	-
2002.....	64	57	83	96	145	204	250	280	252	118	23	-
2001.....	46	66	77	101	151	191	256	290	245	117	24	-
2000.....	48	56	87	100	146	208	252	286	250	108	30	-
1999.....	51	50	83	103	161	177	255	304	221	97	20	-
White, both sexes												
2004.....	91	132	136	138	179	257	314	299	224	104	29	1
2003.....	132	150	161	214	259	332	397	406	307	124	25	1
2002.....	171	172	195	237	287	369	427	409	312	143	23	-
2001.....	137	154	201	219	312	345	426	406	298	125	26	-
2000.....	145	160	207	236	294	373	417	400	288	130	32	-
1999.....	120	162	195	247	316	354	396	438	279	104	21	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute ischemic heart diseases (I24)												
White, male												
2004.....	65	96	93	92	102	127	158	127	68	20	5	1
2003.....	97	102	111	141	161	185	166	144	86	21	6	1
2002.....	123	126	133	159	164	193	203	162	78	32	4	-
2001.....	103	107	143	135	187	179	203	148	66	23	7	-
2000.....	107	124	142	154	171	199	193	142	60	27	4	-
1999.....	85	122	124	160	183	206	169	159	69	17	2	-
White, female												
2004.....	26	36	43	46	77	130	156	172	156	84	24	-
2003.....	35	48	50	73	98	147	231	262	221	103	19	-
2002.....	48	46	62	78	123	176	224	247	234	111	19	-
2001.....	34	47	58	84	125	166	223	258	232	102	19	-
2000.....	38	36	65	82	123	174	224	258	228	103	28	-
1999.....	35	40	71	87	133	148	227	279	210	87	19	-
Black, both sexes												
2004.....	42	49	37	38	33	48	40	23	21	7	3	-
2003.....	32	43	39	47	47	51	46	36	26	4	2	-
2002.....	51	32	47	47	53	45	45	38	21	8	4	-
2001.....	42	40	58	49	46	40	40	45	12	16	5	-
2000.....	40	51	52	47	46	56	43	35	26	6	3	1
1999.....	39	43	42	39	54	49	48	28	15	10	1	-
Black, male												
2004.....	28	34	25	29	22	26	16	14	7	1	1	-
2003.....	22	33	26	30	24	31	26	14	7	-	2	-
2002.....	35	21	29	32	32	20	21	9	8	3	-	-
2001.....	33	22	40	36	23	24	11	15	4	2	-	-
2000.....	31	35	32	32	27	28	18	9	7	1	1	1
1999.....	23	34	31	26	30	21	20	10	5	2	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other acute ischemic heart diseases (I24)

Black, female

2004.....	14	15	12	9	11	22	24	9	14	6	2	-
2003.....	10	10	13	17	23	20	20	22	19	4	-	-
2002.....	16	11	18	15	21	25	24	29	13	5	4	-
2001.....	9	18	18	13	23	16	29	30	8	14	5	-
2000.....	9	16	20	15	19	28	25	26	19	5	2	-
1999.....	16	9	11	13	24	28	28	18	10	8	1	-

American Indian 1/, both sexes

2004.....	1	2	1	2	2	1	4	4	1	-	1	-
2003.....	4	4	4	4	5	3	3	2	2	-	1	-
2002.....	3	3	5	3	2	4	4	1	5	2	-	-
2001.....	7	5	2	9	5	6	2	3	3	-	-	-
2000.....	4	4	5	8	7	4	4	1	2	-	-	-
1999.....	1	3	2	6	8	3	3	6	1	1	-	-

American Indian 1/, male

2004.....	1	1	1	2	1	-	-	1	1	-	-	-
2003.....	2	1	4	1	2	1	-	-	-	-	1	-
2002.....	3	3	4	1	1	2	3	-	1	-	-	-
2001.....	5	5	1	7	3	3	-	1	-	-	-	-
2000.....	3	2	3	6	4	3	1	1	1	-	-	-
1999.....	1	2	1	3	5	2	3	1	-	-	-	-

American Indian 1/, female

2004.....	-	1	-	-	1	1	4	3	-	-	1	-
2003.....	2	3	-	3	3	2	3	2	2	-	-	-
2002.....	-	-	1	2	1	2	1	1	4	2	-	-
2001.....	2	-	1	2	2	3	2	2	3	-	-	-
2000.....	1	2	2	2	3	1	3	-	1	-	-	-
1999.....	-	1	1	3	3	1	-	5	1	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute ischemic heart diseases (I24)												
Asian or Pacific Islander 2/, both sexes												
2004.....	2	1	1	1	6	2	4	1	-	-	-	-
2003.....	3	1	1	2	1	3	2	3	2	1	-	-
2002.....	1	3	3	2	-	2	2	4	1	-	-	-
2001.....	1	1	2	4	1	6	6	1	4	1	-	-
2000.....	-	4	1	3	3	7	2	3	2	-	-	-
1999.....	-	1	-	2	2	-	2	2	1	1	-	-
Asian or Pacific Islander 2/, male												
2004.....	2	1	-	1	5	1	1	1	-	-	-	-
2003.....	1	1	-	1	1	3	2	1	-	-	-	-
2002.....	1	3	1	1	-	1	1	1	-	-	-	-
2001.....	-	-	2	2	-	-	4	1	2	-	-	-
2000.....	-	2	1	2	2	2	2	1	-	-	-	-
1999.....	-	1	-	2	1	-	2	-	1	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	-	-	1	-	1	1	3	-	-	-	-	-
2003.....	2	-	1	1	-	-	-	2	2	1	-	-
2002.....	-	-	2	1	-	1	1	3	1	-	-	-
2001.....	1	1	-	2	1	6	2	-	2	1	-	-
2000.....	-	2	-	1	1	5	-	2	2	-	-	-
1999.....	-	-	-	-	1	-	-	2	-	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)												
All races, both sexes												
2004.....	8,889	11,907	14,978	18,378	25,998	38,402	52,019	51,723	38,784	16,467	3,813	19
2003.....	8,873	11,940	15,267	19,273	28,105	41,408	54,017	54,496	40,662	17,283	4,048	23
2002.....	8,893	12,029	14,840	19,817	29,416	43,195	54,708	55,207	40,876	17,435	4,181	32
2001.....	8,705	11,366	14,915	20,531	30,907	44,919	54,810	55,183	40,716	17,458	4,280	38
2000.....	8,416	11,475	14,765	21,065	32,717	46,070	54,506	56,529	41,250	17,586	4,281	39
1999.....	8,493	11,243	15,280	22,453	34,272	47,920	55,810	56,973	41,734	18,102	4,362	32
All races, male												
2004.....	6,805	8,765	10,510	12,251	16,110	22,131	26,238	21,733	12,288	3,741	592	18
2003.....	6,844	8,819	10,698	12,843	17,505	23,647	27,162	22,538	12,472	3,742	582	17
2002.....	6,878	8,910	10,460	13,166	18,316	24,622	27,254	22,423	12,371	3,759	604	23
2001.....	6,665	8,389	10,439	13,555	19,161	25,537	26,867	21,890	11,852	3,523	588	27
2000.....	6,461	8,501	10,315	13,854	20,479	25,925	26,745	22,106	11,834	3,649	604	28
1999.....	6,466	8,208	10,571	14,839	21,230	26,821	27,118	21,932	11,831	3,703	637	29
All races, female												
2004.....	2,084	3,142	4,468	6,127	9,888	16,271	25,781	29,990	26,496	12,726	3,221	1
2003.....	2,029	3,121	4,569	6,430	10,600	17,761	26,855	31,958	28,190	13,541	3,466	6
2002.....	2,015	3,119	4,380	6,651	11,100	18,573	27,454	32,784	28,505	13,676	3,577	9
2001.....	2,040	2,977	4,476	6,976	11,746	19,382	27,943	33,293	28,864	13,935	3,692	11
2000.....	1,955	2,974	4,450	7,211	12,238	20,145	27,761	34,423	29,416	13,937	3,677	11
1999.....	2,027	3,035	4,709	7,614	13,042	21,099	28,692	35,041	29,903	14,399	3,725	3
White, both sexes												
2004.....	6,852	9,410	12,107	14,992	22,074	33,634	46,888	47,535	35,783	15,135	3,391	12
2003.....	6,857	9,557	12,212	15,722	23,849	36,304	48,772	50,139	37,562	15,965	3,557	18
2002.....	6,900	9,569	11,920	16,257	25,237	37,937	49,605	50,940	37,846	16,047	3,695	24
2001.....	6,820	9,025	11,923	16,826	26,564	39,533	49,705	50,905	37,806	16,051	3,779	24
2000.....	6,662	9,139	11,790	17,493	28,134	40,754	49,565	52,126	38,373	16,264	3,840	29
1999.....	6,652	8,912	12,313	18,665	29,519	42,499	50,899	52,526	38,818	16,714	3,930	22

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)												
White, male												
2004.....	5,400	7,093	8,644	10,225	13,958	19,694	23,993	20,205	11,373	3,428	506	11
2003.....	5,439	7,233	8,772	10,707	15,096	21,095	24,788	20,971	11,643	3,432	505	12
2002.....	5,502	7,255	8,555	11,045	16,008	22,039	25,018	20,909	11,486	3,444	522	18
2001.....	5,354	6,804	8,546	11,389	16,729	22,864	24,650	20,438	11,018	3,231	504	16
2000.....	5,246	6,904	8,465	11,813	17,975	23,262	24,614	20,585	11,040	3,336	515	23
1999.....	5,179	6,675	8,660	12,662	18,619	24,152	25,062	20,361	10,941	3,391	569	20
White, female												
2004.....	1,452	2,317	3,463	4,767	8,116	13,940	22,895	27,330	24,410	11,707	2,885	1
2003.....	1,418	2,324	3,440	5,015	8,753	15,209	23,984	29,168	25,919	12,533	3,052	6
2002.....	1,398	2,314	3,365	5,212	9,229	15,898	24,587	30,031	26,360	12,603	3,173	6
2001.....	1,466	2,221	3,377	5,437	9,835	16,669	25,055	30,467	26,788	12,820	3,275	8
2000.....	1,416	2,235	3,325	5,680	10,159	17,492	24,951	31,541	27,333	12,928	3,325	6
1999.....	1,473	2,237	3,653	6,003	10,900	18,347	25,837	32,165	27,877	13,323	3,361	2
Black, both sexes												
2004.....	1,802	2,167	2,502	2,890	3,345	3,968	4,136	3,266	2,437	1,094	363	7
2003.....	1,778	2,080	2,658	3,029	3,631	4,288	4,284	3,497	2,509	1,096	423	3
2002.....	1,757	2,154	2,550	3,050	3,555	4,390	4,251	3,494	2,496	1,137	418	8
2001.....	1,679	2,044	2,627	3,206	3,718	4,549	4,265	3,553	2,454	1,212	438	12
2000.....	1,546	2,055	2,592	3,095	3,984	4,481	4,150	3,705	2,411	1,128	387	10
1999.....	1,632	2,078	2,587	3,329	4,076	4,580	4,096	3,720	2,428	1,212	386	10
Black, male												
2004.....	1,220	1,424	1,606	1,697	1,832	1,981	1,754	1,107	685	224	66	7
2003.....	1,214	1,354	1,647	1,785	2,029	2,127	1,874	1,175	603	229	53	3
2002.....	1,183	1,429	1,648	1,795	1,929	2,078	1,804	1,151	658	231	63	5
2001.....	1,156	1,343	1,638	1,849	2,076	2,201	1,799	1,126	665	226	71	10
2000.....	1,044	1,394	1,579	1,723	2,128	2,169	1,714	1,203	602	240	76	5
1999.....	1,120	1,344	1,641	1,892	2,191	2,166	1,648	1,224	661	245	56	9

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other forms of chronic ischemic heart disease (I20,I25)

Black, female

2004.....	582	743	896	1,193	1,513	1,987	2,382	2,159	1,752	870	297	-
2003.....	564	726	1,011	1,244	1,602	2,161	2,410	2,322	1,906	867	370	-
2002.....	574	725	902	1,255	1,626	2,312	2,447	2,343	1,838	906	355	3
2001.....	523	701	989	1,357	1,642	2,348	2,466	2,427	1,789	986	367	2
2000.....	502	661	1,013	1,372	1,856	2,312	2,436	2,502	1,809	888	311	5
1999.....	512	734	946	1,437	1,885	2,414	2,448	2,496	1,767	967	330	1

American Indian 1/, both sexes

2004.....	74	95	102	120	120	158	171	125	71	40	8	-
2003.....	71	85	95	129	140	155	132	127	70	25	13	2
2002.....	68	86	97	123	133	153	138	102	62	34	9	-
2001.....	55	91	101	119	133	132	127	88	52	16	6	-
2000.....	55	77	103	124	112	146	118	105	73	30	5	-
1999.....	61	69	96	110	140	153	117	96	61	21	5	-

American Indian 1/, male

2004.....	51	69	70	85	62	93	88	60	21	11	1	-
2003.....	53	65	58	79	94	84	73	55	27	8	5	2
2002.....	50	52	66	82	87	89	65	53	19	8	1	-
2001.....	36	66	72	70	83	71	55	33	14	4	2	-
2000.....	40	51	72	73	73	81	59	41	25	8	-	-
1999.....	45	48	63	71	83	93	52	30	23	5	1	-

American Indian 1/, female

2004.....	23	26	32	35	58	65	83	65	50	29	7	-
2003.....	18	20	37	50	46	71	59	72	43	17	8	-
2002.....	18	34	31	41	46	64	73	49	43	26	8	-
2001.....	19	25	29	49	50	61	72	55	38	12	4	-
2000.....	15	26	31	51	39	65	59	64	48	22	5	-
1999.....	16	21	33	39	57	60	65	66	38	16	4	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)												
Asian or Pacific Islander 2/, both sexes												
2004.....	161	235	267	376	459	642	824	797	493	198	51	-
2003.....	167	218	302	393	485	661	829	733	521	197	55	-
2002.....	168	220	273	387	491	715	714	671	472	217	59	-
2001.....	151	206	264	380	492	705	713	637	404	179	57	2
2000.....	153	204	280	353	487	689	673	593	393	164	49	-
1999.....	148	184	284	349	537	688	698	631	427	155	41	-
Asian or Pacific Islander 2/, male												
2004.....	134	179	190	244	258	363	403	361	209	78	19	-
2003.....	138	167	221	272	286	341	427	337	199	73	19	-
2002.....	143	174	191	244	292	416	367	310	208	76	18	-
2001.....	119	176	183	247	273	401	363	293	155	62	11	1
2000.....	131	152	199	245	303	413	358	277	167	65	13	-
1999.....	122	141	207	214	337	410	356	317	206	62	11	-
Asian or Pacific Islander 2/, female												
2004.....	27	56	77	132	201	279	421	436	284	120	32	-
2003.....	29	51	81	121	199	320	402	396	322	124	36	-
2002.....	25	46	82	143	199	299	347	361	264	141	41	-
2001.....	32	30	81	133	219	304	350	344	249	117	46	1
2000.....	22	52	81	108	184	276	315	316	226	99	36	-
1999.....	26	43	77	135	200	278	342	314	221	93	30	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)												
All races, both sexes												
2004.....	3,555	4,378	4,919	4,957	5,873	7,467	9,263	8,986	6,723	2,989	761	14
2003.....	3,533	4,208	4,902	5,194	6,300	8,253	10,086	9,600	7,316	3,135	823	14
2002.....	3,285	4,189	4,604	5,163	6,654	8,649	9,969	10,007	7,555	3,361	873	23
2001.....	3,209	3,924	4,514	5,272	6,733	9,041	10,155	9,984	7,556	3,323	873	25
2000.....	3,086	3,841	4,401	5,309	7,258	9,271	10,274	10,397	7,953	3,429	885	19
1999.....	2,945	3,747	4,636	5,691	7,704	9,666	10,655	10,745	8,204	3,591	863	24
All races, male												
2004.....	2,821	3,361	3,503	3,389	3,657	4,145	4,461	3,499	1,977	651	115	13
2003.....	2,837	3,203	3,491	3,490	3,932	4,543	4,790	3,759	2,106	700	112	12
2002.....	2,605	3,250	3,299	3,498	4,119	4,746	4,670	3,814	2,165	722	134	17
2001.....	2,549	2,978	3,267	3,483	4,185	4,897	4,774	3,630	2,167	616	108	19
2000.....	2,444	2,978	3,147	3,557	4,512	5,086	4,639	3,740	2,205	691	122	15
1999.....	2,314	2,842	3,281	3,842	4,704	5,200	4,763	3,895	2,196	692	122	22
All races, female												
2004.....	734	1,017	1,416	1,568	2,216	3,322	4,802	5,487	4,746	2,338	646	1
2003.....	696	1,005	1,411	1,704	2,368	3,710	5,296	5,841	5,210	2,435	711	2
2002.....	680	939	1,305	1,665	2,535	3,903	5,299	6,193	5,390	2,639	739	6
2001.....	660	946	1,247	1,789	2,548	4,144	5,381	6,354	5,389	2,707	765	6
2000.....	642	863	1,254	1,752	2,746	4,185	5,635	6,657	5,748	2,738	763	4
1999.....	631	905	1,355	1,849	3,000	4,466	5,892	6,850	6,008	2,899	741	2
White, both sexes												
2004.....	2,663	3,320	3,788	3,793	4,694	6,127	8,005	8,001	6,012	2,670	649	7
2003.....	2,653	3,255	3,672	3,976	5,000	6,822	8,750	8,563	6,593	2,837	707	10
2002.....	2,446	3,170	3,512	3,898	5,324	7,130	8,653	8,950	6,825	3,024	740	16
2001.....	2,413	2,897	3,335	3,941	5,375	7,447	8,869	8,927	6,867	2,978	745	16
2000.....	2,339	2,858	3,255	4,093	5,765	7,717	8,892	9,245	7,202	3,112	760	13
1999.....	2,183	2,765	3,446	4,342	6,130	8,014	9,303	9,568	7,430	3,248	749	15

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)												
White, male												
2004.....	2,191	2,594	2,720	2,654	2,984	3,437	3,889	3,127	1,769	574	94	6
2003.....	2,180	2,530	2,664	2,718	3,127	3,805	4,169	3,376	1,902	623	93	8
2002.....	1,999	2,504	2,563	2,678	3,362	3,952	4,048	3,427	1,953	646	111	12
2001.....	1,968	2,270	2,466	2,665	3,379	4,090	4,206	3,272	1,958	548	86	11
2000.....	1,903	2,252	2,408	2,811	3,636	4,249	4,037	3,348	1,997	611	93	12
1999.....	1,763	2,156	2,477	3,001	3,800	4,348	4,178	3,478	1,964	615	108	13
White, female												
2004.....	472	726	1,068	1,139	1,710	2,690	4,116	4,874	4,243	2,096	555	1
2003.....	473	725	1,008	1,258	1,873	3,017	4,581	5,187	4,691	2,214	614	2
2002.....	447	666	949	1,220	1,962	3,178	4,605	5,523	4,872	2,378	629	4
2001.....	445	627	869	1,276	1,996	3,357	4,663	5,655	4,909	2,430	659	5
2000.....	436	606	847	1,282	2,129	3,468	4,855	5,897	5,205	2,501	667	1
1999.....	420	609	969	1,341	2,330	3,666	5,125	6,090	5,466	2,633	641	2
Black, both sexes												
2004.....	795	939	995	1,018	1,062	1,173	1,084	814	603	277	97	7
2003.....	775	840	1,086	1,062	1,128	1,258	1,114	880	620	264	98	3
2002.....	754	901	984	1,125	1,175	1,350	1,148	903	626	291	118	7
2001.....	717	911	1,054	1,186	1,208	1,405	1,113	922	602	299	113	8
2000.....	681	882	1,029	1,083	1,325	1,349	1,204	1,010	663	280	114	6
1999.....	680	889	1,053	1,212	1,396	1,466	1,173	1,019	671	304	103	9
Black, male												
2004.....	551	669	684	632	609	611	489	288	168	61	18	7
2003.....	573	580	726	655	697	660	515	320	160	67	13	3
2002.....	530	652	659	722	660	699	546	308	170	63	18	5
2001.....	517	620	708	722	716	711	484	295	174	51	17	7
2000.....	489	646	662	648	769	716	505	340	175	67	26	3
1999.....	485	612	697	760	799	737	499	350	188	64	9	9

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)												
Black, female												
2004.....	244	270	311	386	453	562	595	526	435	216	79	-
2003.....	202	260	360	407	431	598	599	560	460	197	85	-
2002.....	224	249	325	403	515	651	602	595	456	228	100	2
2001.....	200	291	346	464	492	694	629	627	428	248	96	1
2000.....	192	236	367	435	556	633	699	670	488	213	88	3
1999.....	195	277	356	452	597	729	674	669	483	240	94	-
American Indian 1/, both sexes												
2004.....	30	37	33	39	27	34	37	31	18	10	-	-
2003.....	31	35	26	45	49	39	31	27	18	7	7	1
2002.....	27	37	23	38	43	32	34	23	22	7	3	-
2001.....	29	31	46	45	38	37	34	19	17	5	2	-
2000.....	20	27	25	31	24	33	25	30	15	5	1	-
1999.....	24	24	28	33	37	45	34	25	19	4	-	-
American Indian 1/, male												
2004.....	21	30	23	29	11	24	19	19	2	3	-	-
2003.....	24	26	16	29	37	17	20	10	9	2	3	1
2002.....	23	24	14	28	27	17	15	13	7	3	-	-
2001.....	22	18	37	26	24	17	17	7	5	-	2	-
2000.....	14	21	18	18	15	18	15	9	6	1	-	-
1999.....	17	18	22	23	21	29	15	9	6	1	-	-
American Indian 1/, female												
2004.....	9	7	10	10	16	10	18	12	16	7	-	-
2003.....	7	9	10	16	12	22	11	17	9	5	4	-
2002.....	4	13	9	10	16	15	19	10	15	4	3	-
2001.....	7	13	9	19	14	20	17	12	12	5	-	-
2000.....	6	6	7	13	9	15	10	21	9	4	1	-
1999.....	7	6	6	10	16	16	19	16	13	3	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)												
Asian or Pacific Islander 2/, both sexes												
2004.....	67	82	103	107	90	133	137	140	90	32	15	-
2003.....	74	78	118	111	123	134	191	130	85	27	11	-
2002.....	58	81	85	102	112	137	134	131	82	39	12	-
2001.....	50	85	79	100	112	152	139	116	70	41	13	1
2000.....	46	74	92	102	144	172	153	112	73	32	10	-
1999.....	58	69	109	104	141	141	145	133	84	35	11	-
Asian or Pacific Islander 2/, male												
2004.....	58	68	76	74	53	73	64	65	38	13	3	-
2003.....	60	67	85	88	71	61	86	53	35	8	3	-
2002.....	53	70	63	70	70	78	61	66	35	10	5	-
2001.....	42	70	56	70	66	79	67	56	30	17	3	1
2000.....	38	59	59	80	92	103	82	43	27	12	3	-
1999.....	49	56	85	58	84	86	71	58	38	12	5	-
Asian or Pacific Islander 2/, female												
2004.....	9	14	27	33	37	60	73	75	52	19	12	-
2003.....	14	11	33	23	52	73	105	77	50	19	8	-
2002.....	5	11	22	32	42	59	73	65	47	29	7	-
2001.....	8	15	23	30	46	73	72	60	40	24	10	-
2000.....	8	15	33	22	52	69	71	69	46	20	7	-
1999.....	9	13	24	46	57	55	74	75	46	23	6	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
All races, both sexes												
2004.....	5,334	7,529	10,059	13,421	20,125	30,935	42,756	42,737	32,061	13,478	3,052	5
2003.....	5,340	7,732	10,365	14,079	21,805	33,155	43,931	44,896	33,346	14,148	3,225	9
2002.....	5,608	7,840	10,236	14,654	22,762	34,546	44,739	45,200	33,321	14,074	3,308	9
2001.....	5,496	7,442	10,401	15,259	24,174	35,878	44,655	45,199	33,160	14,135	3,407	13
2000.....	5,330	7,634	10,364	15,756	25,459	36,799	44,232	46,132	33,297	14,157	3,396	20
1999.....	5,548	7,496	10,644	16,762	26,568	38,254	45,155	46,228	33,530	14,511	3,499	8
All races, male												
2004.....	3,984	5,404	7,007	8,862	12,453	17,986	21,777	18,234	10,311	3,090	477	5
2003.....	4,007	5,616	7,207	9,353	13,573	19,104	22,372	18,779	10,366	3,042	470	5
2002.....	4,273	5,660	7,161	9,668	14,197	19,876	22,584	18,609	10,206	3,037	470	6
2001.....	4,116	5,411	7,172	10,072	14,976	20,640	22,093	18,260	9,685	2,907	480	8
2000.....	4,017	5,523	7,168	10,297	15,967	20,839	22,106	18,366	9,629	2,958	482	13
1999.....	4,152	5,366	7,290	10,997	16,526	21,621	22,355	18,037	9,635	3,011	515	7
All races, female												
2004.....	1,350	2,125	3,052	4,559	7,672	12,949	20,979	24,503	21,750	10,388	2,575	-
2003.....	1,333	2,116	3,158	4,726	8,232	14,051	21,559	26,117	22,980	11,106	2,755	4
2002.....	1,335	2,180	3,075	4,986	8,565	14,670	22,155	26,591	23,115	11,037	2,838	3
2001.....	1,380	2,031	3,229	5,187	9,198	15,238	22,562	26,939	23,475	11,228	2,927	5
2000.....	1,313	2,111	3,196	5,459	9,492	15,960	22,126	27,766	23,668	11,199	2,914	7
1999.....	1,396	2,130	3,354	5,765	10,042	16,633	22,800	28,191	23,895	11,500	2,984	1
White, both sexes												
2004.....	4,189	6,090	8,319	11,199	17,380	27,507	38,883	39,534	29,771	12,465	2,742	5
2003.....	4,204	6,302	8,540	11,746	18,849	29,482	40,022	41,576	30,969	13,128	2,850	8
2002.....	4,454	6,399	8,408	12,359	19,913	30,807	40,952	41,990	31,021	13,023	2,955	8
2001.....	4,407	6,128	8,588	12,885	21,189	32,086	40,836	41,978	30,939	13,073	3,034	8
2000.....	4,323	6,281	8,535	13,400	22,369	33,037	40,673	42,881	31,171	13,152	3,080	16
1999.....	4,469	6,147	8,867	14,323	23,389	34,485	41,596	42,958	31,388	13,466	3,181	7

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)

White, male

2004.....	3,209	4,499	5,924	7,571	10,974	16,257	20,104	17,078	9,604	2,854	412	5
2003.....	3,259	4,703	6,108	7,989	11,969	17,290	20,619	17,595	9,741	2,809	412	4
2002.....	3,503	4,751	5,992	8,367	12,646	18,087	20,970	17,482	9,533	2,798	411	6
2001.....	3,386	4,534	6,080	8,724	13,350	18,774	20,444	17,166	9,060	2,683	418	5
2000.....	3,343	4,652	6,057	9,002	14,339	19,013	20,577	17,237	9,043	2,725	422	11
1999.....	3,416	4,519	6,183	9,661	14,819	19,804	20,884	16,883	8,977	2,776	461	7

White, female

2004.....	980	1,591	2,395	3,628	6,406	11,250	18,779	22,456	20,167	9,611	2,330	-
2003.....	945	1,599	2,432	3,757	6,880	12,192	19,403	23,981	21,228	10,319	2,438	4
2002.....	951	1,648	2,416	3,992	7,267	12,720	19,982	24,508	21,488	10,225	2,544	2
2001.....	1,021	1,594	2,508	4,161	7,839	13,312	20,392	24,812	21,879	10,390	2,616	3
2000.....	980	1,629	2,478	4,398	8,030	14,024	20,096	25,644	22,128	10,427	2,658	5
1999.....	1,053	1,628	2,684	4,662	8,570	14,681	20,712	26,075	22,411	10,690	2,720	-

Black, both sexes

2004.....	1,007	1,228	1,507	1,872	2,283	2,795	3,052	2,452	1,834	817	266	-
2003.....	1,003	1,240	1,572	1,967	2,503	3,030	3,170	2,617	1,889	832	325	-
2002.....	1,003	1,253	1,566	1,925	2,380	3,040	3,103	2,591	1,870	846	300	1
2001.....	962	1,133	1,573	2,020	2,510	3,144	3,152	2,631	1,852	913	325	4
2000.....	865	1,173	1,563	2,012	2,659	3,132	2,946	2,695	1,748	848	273	4
1999.....	952	1,189	1,534	2,117	2,680	3,114	2,923	2,701	1,757	908	283	1

Black, male

2004.....	669	755	922	1,065	1,223	1,370	1,265	819	517	163	48	-
2003.....	641	774	921	1,130	1,332	1,467	1,359	855	443	162	40	-
2002.....	653	777	989	1,073	1,269	1,379	1,258	843	488	168	45	-
2001.....	639	723	930	1,127	1,360	1,490	1,315	831	491	175	54	3
2000.....	555	748	917	1,075	1,359	1,453	1,209	863	427	173	50	2
1999.....	635	732	944	1,132	1,392	1,429	1,149	874	473	181	47	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)

Black, female

2004.....	338	473	585	807	1,060	1,425	1,787	1,633	1,317	654	218	-
2003.....	362	466	651	837	1,171	1,563	1,811	1,762	1,446	670	285	-
2002.....	350	476	577	852	1,111	1,661	1,845	1,748	1,382	678	255	1
2001.....	323	410	643	893	1,150	1,654	1,837	1,800	1,361	738	271	1
2000.....	310	425	646	937	1,300	1,679	1,737	1,832	1,321	675	223	2
1999.....	317	457	590	985	1,288	1,685	1,774	1,827	1,284	727	236	1

American Indian 1/, both sexes

2004.....	44	58	69	81	93	124	134	94	53	30	8	-
2003.....	40	50	69	84	91	116	101	100	52	18	6	1
2002.....	41	49	74	85	90	121	104	79	40	27	6	-
2001.....	26	60	55	74	95	95	93	69	35	11	4	-
2000.....	35	50	78	93	88	113	93	75	58	25	4	-
1999.....	37	45	68	77	103	108	83	71	42	17	5	-

American Indian 1/, male

2004.....	30	39	47	56	51	69	69	41	19	8	1	-
2003.....	29	39	42	50	57	67	53	45	18	6	2	1
2002.....	27	28	52	54	60	72	50	40	12	5	1	-
2001.....	14	48	35	44	59	54	38	26	9	4	-	-
2000.....	26	30	54	55	58	63	44	32	19	7	-	-
1999.....	28	30	41	48	62	64	37	21	17	4	1	-

American Indian 1/, female

2004.....	14	19	22	25	42	55	65	53	34	22	7	-
2003.....	11	11	27	34	34	49	48	55	34	12	4	-
2002.....	14	21	22	31	30	49	54	39	28	22	5	-
2001.....	12	12	20	30	36	41	55	43	26	7	4	-
2000.....	9	20	24	38	30	50	49	43	39	18	4	-
1999.....	9	15	27	29	41	44	46	50	25	13	4	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Asian or Pacific Islander 2/, both sexes												
2004.....	94	153	164	269	369	509	687	657	403	166	36	-
2003.....	93	140	184	282	362	527	638	603	436	170	44	-
2002.....	110	139	188	285	379	578	580	540	390	178	47	-
2001.....	101	121	185	280	380	553	574	521	334	138	44	1
2000.....	107	130	188	251	343	517	520	481	320	132	39	-
1999.....	90	115	175	245	396	547	553	498	343	120	30	-
Asian or Pacific Islander 2/, male												
2004.....	76	111	114	170	205	290	339	296	171	65	16	-
2003.....	78	100	136	184	215	280	341	284	164	65	16	-
2002.....	90	104	128	174	222	338	306	244	173	66	13	-
2001.....	77	106	127	177	207	322	296	237	125	45	8	-
2000.....	93	93	140	165	211	310	276	234	140	53	10	-
1999.....	73	85	122	156	253	324	285	259	168	50	6	-
Asian or Pacific Islander 2/, female												
2004.....	18	42	50	99	164	219	348	361	232	101	20	-
2003.....	15	40	48	98	147	247	297	319	272	105	28	-
2002.....	20	35	60	111	157	240	274	296	217	112	34	-
2001.....	24	15	58	103	173	231	278	284	209	93	36	1
2000.....	14	37	48	86	132	207	244	247	180	79	29	-
1999.....	17	30	53	89	143	223	268	239	175	70	24	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other heart diseases (I26-I51)												
All races, both sexes												
2004.....	4,987	5,890	7,244	8,989	12,916	19,868	27,985	30,758	23,870	9,971	2,166	7
2003.....	4,825	6,171	7,227	9,481	13,763	20,800	28,460	31,015	23,947	9,993	2,315	6
2002.....	4,862	5,913	7,033	9,377	14,303	21,276	28,500	30,794	23,517	9,779	2,210	6
2001.....	4,776	5,637	6,898	9,596	14,460	21,325	27,651	30,343	22,779	9,653	2,204	7
2000.....	4,551	5,417	6,945	9,875	14,804	21,524	26,843	29,519	22,677	9,591	2,206	8
1999.....	4,386	5,465	7,115	10,034	15,538	21,689	27,023	29,428	21,894	9,401	2,125	7
All races, male												
2004.....	3,131	3,712	4,359	5,157	6,876	9,799	12,161	11,330	6,977	2,142	325	6
2003.....	3,019	3,826	4,316	5,402	7,473	10,351	12,329	11,327	6,827	2,169	358	5
2002.....	3,082	3,587	4,184	5,275	7,716	10,463	12,308	11,167	6,558	2,070	338	3
2001.....	2,946	3,393	4,110	5,356	7,811	10,538	12,002	10,595	6,187	2,028	331	5
2000.....	2,873	3,384	4,148	5,584	8,037	10,505	11,501	10,488	6,178	1,983	359	5
1999.....	2,756	3,419	4,281	5,870	8,450	10,567	11,561	10,293	5,879	1,988	323	5
All races, female												
2004.....	1,856	2,178	2,885	3,832	6,040	10,069	15,824	19,428	16,893	7,829	1,841	1
2003.....	1,806	2,345	2,911	4,079	6,290	10,449	16,131	19,688	17,120	7,824	1,957	1
2002.....	1,780	2,326	2,849	4,102	6,587	10,813	16,192	19,627	16,959	7,709	1,872	3
2001.....	1,830	2,244	2,788	4,240	6,649	10,787	15,649	19,748	16,592	7,625	1,873	2
2000.....	1,678	2,033	2,797	4,291	6,767	11,019	15,342	19,031	16,499	7,608	1,847	3
1999.....	1,630	2,046	2,834	4,164	7,088	11,122	15,462	19,135	16,015	7,413	1,802	2
White, both sexes												
2004.....	3,537	4,378	5,693	7,237	10,846	17,269	25,348	28,422	22,126	9,237	1,931	4
2003.....	3,395	4,688	5,588	7,597	11,504	18,308	25,766	28,661	22,248	9,241	2,060	5
2002.....	3,481	4,482	5,499	7,614	12,102	18,674	25,844	28,481	21,808	9,026	1,946	3
2001.....	3,453	4,221	5,447	7,747	12,305	18,686	25,072	28,039	21,213	8,946	1,966	6
2000.....	3,294	4,139	5,459	8,050	12,567	18,934	24,445	27,240	21,096	8,849	1,974	7
1999.....	3,192	4,079	5,549	8,177	13,197	19,031	24,547	27,066	20,457	8,697	1,900	5

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other heart diseases (I26-I51)												
White, male												
2004.....	2,301	2,809	3,471	4,204	5,879	8,592	11,063	10,489	6,426	1,970	279	3
2003.....	2,161	2,934	3,409	4,390	6,369	9,229	11,227	10,524	6,355	1,995	304	4
2002.....	2,272	2,764	3,339	4,341	6,619	9,283	11,250	10,394	6,044	1,909	290	2
2001.....	2,211	2,576	3,296	4,419	6,708	9,306	10,903	9,804	5,733	1,859	286	5
2000.....	2,107	2,636	3,323	4,619	6,942	9,326	10,539	9,688	5,729	1,796	309	5
1999.....	2,041	2,598	3,391	4,853	7,285	9,361	10,556	9,430	5,445	1,810	276	4
White, female												
2004.....	1,236	1,569	2,222	3,033	4,967	8,677	14,285	17,933	15,700	7,267	1,652	1
2003.....	1,234	1,754	2,179	3,207	5,135	9,079	14,539	18,137	15,893	7,246	1,756	1
2002.....	1,209	1,718	2,160	3,273	5,483	9,391	14,594	18,087	15,764	7,117	1,656	1
2001.....	1,242	1,645	2,151	3,328	5,597	9,380	14,169	18,235	15,480	7,087	1,680	1
2000.....	1,187	1,503	2,136	3,431	5,625	9,608	13,906	17,552	15,367	7,053	1,665	2
1999.....	1,151	1,481	2,158	3,324	5,912	9,670	13,991	17,636	15,012	6,887	1,624	1
Black, both sexes												
2004.....	1,342	1,387	1,399	1,549	1,838	2,251	2,227	1,937	1,445	606	199	3
2003.....	1,320	1,347	1,474	1,679	2,007	2,144	2,265	1,968	1,404	656	216	1
2002.....	1,277	1,298	1,395	1,595	1,908	2,251	2,273	1,940	1,460	653	225	3
2001.....	1,224	1,297	1,327	1,664	1,891	2,320	2,222	1,994	1,354	607	206	1
2000.....	1,159	1,162	1,344	1,643	2,000	2,290	2,092	1,984	1,349	632	210	1
1999.....	1,100	1,270	1,432	1,679	2,101	2,338	2,150	2,054	1,227	615	190	2
Black, male												
2004.....	766	826	798	836	879	1,032	911	670	409	121	33	3
2003.....	788	809	821	908	968	951	905	638	353	138	42	1
2002.....	743	744	761	837	951	989	884	597	400	122	36	1
2001.....	681	749	745	838	947	1,065	926	663	350	136	40	-
2000.....	707	681	742	864	954	1,017	826	671	355	147	43	-
1999.....	665	749	814	916	1,041	1,030	857	703	351	147	40	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other heart diseases (I26-I51)												
Black, female												
2004.....	576	561	601	713	959	1,219	1,316	1,267	1,036	485	166	-
2003.....	532	538	653	771	1,039	1,193	1,360	1,330	1,051	518	174	-
2002.....	534	554	634	758	957	1,262	1,389	1,343	1,060	531	189	2
2001.....	543	548	582	826	944	1,255	1,296	1,331	1,004	471	166	1
2000.....	452	481	602	779	1,046	1,273	1,266	1,313	994	485	167	1
1999.....	435	521	618	763	1,060	1,308	1,293	1,351	876	468	150	1
American Indian 1/, both sexes												
2004.....	35	38	52	58	51	65	69	62	36	23	7	-
2003.....	41	51	71	61	58	61	80	62	44	16	11	-
2002.....	35	39	35	37	60	75	71	69	37	18	10	-
2001.....	24	40	36	45	64	69	71	65	33	18	6	-
2000.....	35	36	38	54	64	73	65	63	35	18	6	-
1999.....	31	32	47	55	60	79	62	51	45	16	11	-
American Indian 1/, male												
2004.....	16	24	31	32	29	32	22	30	13	4	2	-
2003.....	24	33	33	25	33	24	29	29	21	3	4	-
2002.....	25	22	24	21	37	41	30	31	10	5	6	-
2001.....	14	24	21	25	38	40	34	23	12	4	-	-
2000.....	22	23	21	28	42	39	26	25	7	2	-	-
1999.....	18	19	25	35	33	39	23	19	13	5	4	-
American Indian 1/, female												
2004.....	19	14	21	26	22	33	47	32	23	19	5	-
2003.....	17	18	38	36	25	37	51	33	23	13	7	-
2002.....	10	17	11	16	23	34	41	38	27	13	4	-
2001.....	10	16	15	20	26	29	37	42	21	14	6	-
2000.....	13	13	17	26	22	34	39	38	28	16	6	-
1999.....	13	13	22	20	27	40	39	32	32	11	7	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other heart diseases (I26-I51)												
Asian or Pacific Islander 2/, both sexes												
2004.....	73	87	100	145	181	283	341	337	263	105	29	-
2003.....	69	85	94	144	194	287	349	324	251	80	28	-
2002.....	69	94	104	131	233	276	312	304	212	82	29	-
2001.....	75	79	88	140	200	250	286	245	179	82	26	-
2000.....	63	80	104	128	173	227	241	232	197	92	16	-
1999.....	63	84	87	123	180	241	264	257	165	73	24	-
Asian or Pacific Islander 2/, male												
2004.....	48	53	59	85	89	143	165	141	129	47	11	-
2003.....	46	50	53	79	103	147	168	136	98	33	8	-
2002.....	42	57	60	76	109	150	144	145	104	34	6	-
2001.....	40	44	48	74	118	127	139	105	92	29	5	-
2000.....	37	44	62	73	99	123	110	104	87	38	7	-
1999.....	32	53	51	66	91	137	125	141	70	26	3	-
Asian or Pacific Islander 2/, female												
2004.....	25	34	41	60	92	140	176	196	134	58	18	-
2003.....	23	35	41	65	91	140	181	188	153	47	20	-
2002.....	27	37	44	55	124	126	168	159	108	48	23	-
2001.....	35	35	40	66	82	123	147	140	87	53	21	-
2000.....	26	36	42	55	74	104	131	128	110	54	9	-
1999.....	31	31	36	57	89	104	139	116	95	47	21	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
All races, both sexes												
2004.....	717	783	893	1,099	1,384	1,725	1,843	1,473	908	241	42	1
2003.....	756	850	977	1,135	1,505	1,921	1,996	1,586	848	247	53	-
2002.....	717	882	944	1,212	1,518	1,909	1,887	1,518	778	249	41	1
2001.....	716	834	914	1,148	1,521	1,795	1,790	1,477	773	247	33	-
2000.....	740	703	922	1,170	1,492	1,790	1,714	1,444	702	240	47	-
1999.....	659	733	859	1,134	1,590	1,886	1,780	1,405	772	254	43	2
All races, male												
2004.....	344	405	410	476	575	667	648	434	241	50	4	1
2003.....	368	388	446	516	668	820	719	504	218	58	6	-
2002.....	323	384	441	546	628	730	679	469	190	61	4	-
2001.....	305	351	434	517	686	706	684	459	206	54	3	-
2000.....	344	321	425	506	647	764	615	439	185	42	6	-
1999.....	301	336	436	528	722	769	634	437	221	53	9	2
All races, female												
2004.....	373	378	483	623	809	1,058	1,195	1,039	667	191	38	-
2003.....	388	462	531	619	837	1,101	1,277	1,082	630	189	47	-
2002.....	394	498	503	666	890	1,179	1,208	1,049	588	188	37	1
2001.....	411	483	480	631	835	1,089	1,106	1,018	567	193	30	-
2000.....	396	382	497	664	845	1,026	1,099	1,005	517	198	41	-
1999.....	358	397	423	606	868	1,117	1,146	968	551	201	34	-
White, both sexes												
2004.....	532	596	718	874	1,144	1,459	1,619	1,347	819	219	36	1
2003.....	567	662	772	897	1,209	1,685	1,777	1,433	754	228	45	-
2002.....	553	678	763	986	1,276	1,639	1,681	1,380	705	213	33	-
2001.....	554	645	732	920	1,291	1,539	1,609	1,332	689	212	29	-
2000.....	554	570	736	970	1,259	1,553	1,534	1,289	637	202	37	-
1999.....	498	554	662	912	1,311	1,628	1,565	1,254	701	235	40	2

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
White, male												
2004.....	264	310	339	396	486	570	547	389	215	45	4	1
2003.....	281	303	364	418	553	732	646	455	195	51	5	-
2002.....	273	300	378	454	545	641	603	435	164	50	4	-
2001.....	252	269	351	430	592	611	610	409	174	45	2	-
2000.....	265	264	347	435	565	675	551	385	166	33	3	-
1999.....	233	269	343	431	605	672	556	395	199	49	8	2
White, female												
2004.....	268	286	379	478	658	889	1,072	958	604	174	32	-
2003.....	286	359	408	479	656	953	1,131	978	559	177	40	-
2002.....	280	378	385	532	731	998	1,078	945	541	163	29	-
2001.....	302	376	381	490	699	928	999	923	515	167	27	-
2000.....	289	306	389	535	694	878	983	904	471	169	34	-
1999.....	265	285	319	481	706	956	1,009	859	502	186	32	-
Black, both sexes												
2004.....	170	171	162	203	222	253	189	113	80	21	6	-
2003.....	175	173	189	224	277	217	197	138	88	17	7	-
2002.....	153	191	173	213	226	254	189	124	66	34	7	1
2001.....	149	176	171	220	212	229	167	138	80	31	4	-
2000.....	173	120	179	189	219	221	166	139	61	35	10	-
1999.....	152	165	187	208	265	245	200	140	66	17	3	-
Black, male												
2004.....	73	86	63	73	77	92	87	37	23	5	-	-
2003.....	81	82	75	94	105	81	68	42	23	7	1	-
2002.....	47	80	59	87	75	82	67	29	22	10	-	-
2001.....	48	76	80	83	85	82	68	46	29	8	1	-
2000.....	73	56	77	66	79	82	60	50	18	9	3	-
1999.....	64	61	89	92	113	91	72	36	21	3	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Black, female												
2004.....	97	85	99	130	145	161	102	76	57	16	6	-
2003.....	94	91	114	130	172	136	129	96	65	10	6	-
2002.....	106	111	114	126	151	172	122	95	44	24	7	1
2001.....	101	100	91	137	127	147	99	92	51	23	3	-
2000.....	100	64	102	123	140	139	106	89	43	26	7	-
1999.....	88	104	98	116	152	154	128	104	45	14	2	-
American Indian 1/, both sexes												
2004.....	4	3	7	6	5	3	8	1	1	-	-	-
2003.....	4	9	7	5	7	2	9	5	-	1	-	-
2002.....	5	6	2	4	5	3	6	6	1	1	-	-
2001.....	4	5	1	4	5	6	2	-	1	-	-	-
2000.....	2	5	2	4	5	1	2	3	1	1	-	-
1999.....	3	4	5	5	5	4	2	4	2	1	-	-
American Indian 1/, male												
2004.....	2	2	6	3	4	-	3	-	1	-	-	-
2003.....	1	3	4	1	3	1	2	3	-	-	-	-
2002.....	2	-	1	2	2	2	3	1	-	-	-	-
2001.....	2	3	-	2	2	2	1	-	1	-	-	-
2000.....	-	1	-	2	3	1	-	-	-	-	-	-
1999.....	2	2	2	2	3	1	-	2	1	-	-	-
American Indian 1/, female												
2004.....	2	1	1	3	1	3	5	1	-	-	-	-
2003.....	3	6	3	4	4	1	7	2	-	1	-	-
2002.....	3	6	1	2	3	1	3	5	1	1	-	-
2001.....	2	2	1	2	3	4	1	-	-	-	-	-
2000.....	2	4	2	2	2	-	2	3	1	-	-	-
1999.....	1	2	3	3	2	3	2	2	1	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Asian or Pacific Islander 2/, both sexes												
2004.....	11	13	6	16	13	10	27	12	8	1	-	-
2003.....	10	6	9	9	12	17	13	10	6	1	1	-
2002.....	6	7	6	9	11	13	11	8	6	1	1	-
2001.....	9	8	10	4	13	21	12	7	3	4	-	-
2000.....	11	8	5	7	9	15	12	13	3	2	-	-
1999.....	6	10	5	9	9	9	13	7	3	1	-	-
Asian or Pacific Islander 2/, male												
2004.....	5	7	2	4	8	5	11	8	2	-	-	-
2003.....	5	-	3	3	7	6	3	4	-	-	-	-
2002.....	1	4	3	3	6	5	6	4	4	1	-	-
2001.....	3	3	3	2	7	11	5	4	2	1	-	-
2000.....	6	-	1	3	-	6	4	4	1	-	-	-
1999.....	2	4	2	3	1	5	6	4	-	1	-	-
Asian or Pacific Islander 2/, female												
2004.....	6	6	4	12	5	5	16	4	6	1	-	-
2003.....	5	6	6	6	5	11	10	6	6	1	1	-
2002.....	5	3	3	6	5	8	5	4	2	-	1	-
2001.....	6	5	7	2	6	10	7	3	1	3	-	-
2000.....	5	8	4	4	9	9	8	9	2	2	-	-
1999.....	4	6	3	6	8	4	7	3	3	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary embolism (I26)												
All races, both sexes												
2004.....	460	488	568	682	825	1,068	1,152	898	552	150	24	-
2003.....	526	560	620	682	922	1,168	1,258	957	531	151	32	-
2002.....	475	586	613	776	952	1,174	1,181	962	481	161	23	1
2001.....	489	585	601	746	998	1,165	1,146	982	511	154	18	-
2000.....	495	470	627	753	960	1,166	1,111	961	473	165	29	-
1999.....	452	516	601	753	1,115	1,315	1,260	1,006	533	174	33	2
All races, male												
2004.....	225	281	281	332	374	443	435	280	151	29	2	-
2003.....	264	288	311	349	437	524	484	314	142	33	4	-
2002.....	224	276	304	364	417	460	428	302	122	46	4	-
2001.....	218	250	297	371	457	493	454	316	140	37	1	-
2000.....	259	228	309	354	437	531	411	306	123	28	3	-
1999.....	213	256	318	367	533	545	437	317	141	36	7	2
All races, female												
2004.....	235	207	287	350	451	625	717	618	401	121	22	-
2003.....	262	272	309	333	485	644	774	643	389	118	28	-
2002.....	251	310	309	412	535	714	753	660	359	115	19	1
2001.....	271	335	304	375	541	672	692	666	371	117	17	-
2000.....	236	242	318	399	523	635	700	655	350	137	26	-
1999.....	239	260	283	386	582	770	823	689	392	138	26	-
White, both sexes												
2004.....	342	384	463	536	677	897	1,010	815	493	133	20	-
2003.....	393	438	495	531	737	1,026	1,109	858	461	138	26	-
2002.....	365	454	503	634	797	998	1,041	871	423	133	19	-
2001.....	383	466	486	610	845	992	1,029	882	447	130	16	-
2000.....	371	378	503	615	814	995	984	849	426	142	19	-
1999.....	343	393	465	608	912	1,121	1,113	903	474	162	30	2

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary embolism (I26)												
White, male												
2004.....	173	217	234	273	312	370	364	248	133	25	2	-
2003.....	203	225	250	280	361	461	427	279	125	26	4	-
2002.....	186	216	259	302	364	402	376	280	99	36	4	-
2001.....	184	193	238	312	398	423	402	280	113	31	1	-
2000.....	195	182	251	302	384	462	361	264	111	22	-	-
1999.....	162	200	253	295	445	468	378	282	125	33	6	2
White, female												
2004.....	169	167	229	263	365	527	646	567	360	108	18	-
2003.....	190	213	245	251	376	565	682	579	336	112	22	-
2002.....	179	238	244	332	433	596	665	591	324	97	15	-
2001.....	199	273	248	298	447	569	627	602	334	99	15	-
2000.....	176	196	252	313	430	533	623	585	315	120	19	-
1999.....	181	193	212	313	467	653	735	621	349	129	24	-
Black, both sexes												
2004.....	109	98	100	134	138	162	124	75	56	16	4	-
2003.....	125	115	114	146	178	135	137	91	66	13	5	-
2002.....	106	127	105	135	145	167	128	81	54	26	4	1
2001.....	100	112	111	132	143	157	112	97	60	23	2	-
2000.....	115	81	121	133	139	162	117	105	44	22	10	-
1999.....	106	115	130	142	195	187	141	97	55	12	3	-
Black, male												
2004.....	47	60	42	53	55	70	63	26	16	4	-	-
2003.....	57	61	54	68	69	60	54	31	17	7	-	-
2002.....	37	58	43	58	48	54	47	19	20	9	-	-
2001.....	31	54	59	56	54	62	50	34	24	6	-	-
2000.....	60	45	58	49	51	64	46	41	12	6	3	-
1999.....	48	52	63	72	86	75	57	32	16	3	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary embolism (I26)												
Black, female												
2004.....	62	38	58	81	83	92	61	49	40	12	4	-
2003.....	68	54	60	78	109	75	83	60	49	6	5	-
2002.....	69	69	62	77	97	113	81	62	34	17	4	1
2001.....	69	58	52	76	89	95	62	63	36	17	2	-
2000.....	55	36	63	84	88	98	71	64	32	16	7	-
1999.....	58	63	67	70	109	112	84	65	39	9	2	-
American Indian 1/, both sexes												
2004.....	1	1	4	4	4	2	5	-	1	-	-	-
2003.....	3	6	6	2	3	2	6	3	-	-	-	-
2002.....	3	3	2	3	5	3	4	4	1	1	-	-
2001.....	2	3	1	3	2	4	2	-	1	-	-	-
2000.....	2	5	-	3	2	-	1	2	1	-	-	-
1999.....	2	2	3	1	3	4	1	3	1	-	-	-
American Indian 1/, male												
2004.....	1	-	4	3	3	-	2	-	1	-	-	-
2003.....	1	2	4	-	3	1	1	2	-	-	-	-
2002.....	1	-	1	2	2	2	1	-	-	-	-	-
2001.....	1	2	-	2	-	1	1	-	1	-	-	-
2000.....	-	1	-	1	2	-	-	-	-	-	-	-
1999.....	2	1	1	-	2	1	-	1	-	-	-	-
American Indian 1/, female												
2004.....	-	1	-	1	1	2	3	-	-	-	-	-
2003.....	2	4	2	2	-	1	5	1	-	-	-	-
2002.....	2	3	1	1	3	1	3	4	1	1	-	-
2001.....	1	1	1	1	2	3	1	-	-	-	-	-
2000.....	2	4	-	2	-	-	1	2	1	-	-	-
1999.....	-	1	2	1	1	3	1	2	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary embolism (I26)												
Asian or Pacific Islander 2/, both sexes												
2004.....	8	5	1	8	6	7	13	8	2	1	-	-
2003.....	5	1	5	3	4	5	6	5	4	-	1	-
2002.....	1	2	3	4	5	6	8	6	3	1	-	-
2001.....	4	4	3	1	8	12	3	3	3	1	-	-
2000.....	7	6	3	2	5	9	9	5	2	1	-	-
1999.....	1	6	3	2	5	3	5	3	3	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	4	4	1	3	4	3	6	6	1	-	-	-
2003.....	3	-	3	1	4	2	2	2	-	-	-	-
2002.....	-	2	1	2	3	2	4	3	3	1	-	-
2001.....	2	1	-	1	5	7	1	2	2	-	-	-
2000.....	4	-	-	2	-	5	4	1	-	-	-	-
1999.....	1	3	1	-	-	1	2	2	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	4	1	-	5	2	4	7	2	1	1	-	-
2003.....	2	1	2	2	-	3	4	3	4	-	1	-
2002.....	1	-	2	2	2	4	4	3	-	-	-	-
2001.....	2	3	3	-	3	5	2	1	1	1	-	-
2000.....	3	6	3	-	5	4	5	4	2	1	-	-
1999.....	-	3	2	2	5	2	3	1	3	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
All races, both sexes												
2004.....	257	295	325	417	559	657	691	575	356	91	18	1
2003.....	230	290	357	453	583	753	738	629	317	96	21	-
2002.....	242	296	331	436	566	735	706	556	297	88	18	-
2001.....	227	249	313	402	523	630	644	495	262	93	15	-
2000.....	245	233	295	417	532	624	603	483	229	75	18	-
1999.....	207	217	258	381	475	571	520	399	239	80	10	-
All races, male												
2004.....	119	124	129	144	201	224	213	154	90	21	2	1
2003.....	104	100	135	167	231	296	235	190	76	25	2	-
2002.....	99	108	137	182	211	270	251	167	68	15	-	-
2001.....	87	101	137	146	229	213	230	143	66	17	2	-
2000.....	85	93	116	152	210	233	204	133	62	14	3	-
1999.....	88	80	118	161	189	224	197	120	80	17	2	-
All races, female												
2004.....	138	171	196	273	358	433	478	421	266	70	16	-
2003.....	126	190	222	286	352	457	503	439	241	71	19	-
2002.....	143	188	194	254	355	465	455	389	229	73	18	-
2001.....	140	148	176	256	294	417	414	352	196	76	13	-
2000.....	160	140	179	265	322	391	399	350	167	61	15	-
1999.....	119	137	140	220	286	347	323	279	159	63	8	-
White, both sexes												
2004.....	190	212	255	338	467	562	609	532	326	86	16	1
2003.....	174	224	277	366	472	659	668	575	293	90	19	-
2002.....	188	224	260	352	479	641	640	509	282	80	14	-
2001.....	171	179	246	310	446	547	580	450	242	82	13	-
2000.....	183	192	233	355	445	558	550	440	211	60	18	-
1999.....	155	161	197	304	399	507	452	351	227	73	10	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
White, male												
2004.....	91	93	105	123	174	200	183	141	82	20	2	1
2003.....	78	78	114	138	192	271	219	176	70	25	1	-
2002.....	87	84	119	152	181	239	227	155	65	14	-	-
2001.....	68	76	113	118	194	188	208	129	61	14	1	-
2000.....	70	82	96	133	181	213	190	121	55	11	3	-
1999.....	71	69	90	136	160	204	178	113	74	16	2	-
White, female												
2004.....	99	119	150	215	293	362	426	391	244	66	14	-
2003.....	96	146	163	228	280	388	449	399	223	65	18	-
2002.....	101	140	141	200	298	402	413	354	217	66	14	-
2001.....	103	103	133	192	252	359	372	321	181	68	12	-
2000.....	113	110	137	222	264	345	360	319	156	49	15	-
1999.....	84	92	107	168	239	303	274	238	153	57	8	-
Black, both sexes												
2004.....	61	73	62	69	84	91	65	38	24	5	2	-
2003.....	50	58	75	78	99	82	60	47	22	4	2	-
2002.....	47	64	68	78	81	87	61	43	12	8	3	-
2001.....	49	64	60	88	69	72	55	41	20	8	2	-
2000.....	58	39	58	56	80	59	49	34	17	13	-	-
1999.....	46	50	57	66	70	58	59	43	11	5	-	-
Black, male												
2004.....	26	26	21	20	22	22	24	11	7	1	-	-
2003.....	24	21	21	26	36	21	14	11	6	-	1	-
2002.....	10	22	16	29	27	28	20	10	2	1	-	-
2001.....	17	22	21	27	31	20	18	12	5	2	1	-
2000.....	13	11	19	17	28	18	14	9	6	3	-	-
1999.....	16	9	26	20	27	16	15	4	5	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)

Black, female

2004.....	35	47	41	49	62	69	41	27	17	4	2	-
2003.....	26	37	54	52	63	61	46	36	16	4	1	-
2002.....	37	42	52	49	54	59	41	33	10	7	3	-
2001.....	32	42	39	61	38	52	37	29	15	6	1	-
2000.....	45	28	39	39	52	41	35	25	11	10	-	-
1999.....	30	41	31	46	43	42	44	39	6	5	-	-

American Indian 1/, both sexes

2004.....	3	2	3	2	1	1	3	1	-	-	-	-
2003.....	1	3	1	3	4	-	3	2	-	1	-	-
2002.....	2	3	-	1	-	-	2	2	-	-	-	-
2001.....	2	2	-	1	3	2	-	-	-	-	-	-
2000.....	-	-	2	1	3	1	1	1	-	1	-	-
1999.....	1	2	2	4	2	-	1	1	1	1	-	-

American Indian 1/, male

2004.....	1	2	2	-	1	-	1	-	-	-	-	-
2003.....	-	1	-	1	-	-	1	1	-	-	-	-
2002.....	1	-	-	-	-	-	2	1	-	-	-	-
2001.....	1	1	-	-	2	1	-	-	-	-	-	-
2000.....	-	-	-	1	1	1	-	-	-	-	-	-
1999.....	-	1	1	2	1	-	-	1	1	-	-	-

American Indian 1/, female

2004.....	2	-	1	2	-	1	2	1	-	-	-	-
2003.....	1	2	1	2	4	-	2	1	-	1	-	-
2002.....	1	3	-	1	-	-	-	1	-	-	-	-
2001.....	1	1	-	1	1	1	-	-	-	-	-	-
2000.....	-	-	2	-	2	-	1	1	-	1	-	-
1999.....	1	1	1	2	1	-	1	-	-	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)

Asian or Pacific Islander 2/, both sexes

2004.....	3	8	5	8	7	3	14	4	6	-	-	-
2003.....	5	5	4	6	8	12	7	5	2	1	-	-
2002.....	5	5	3	5	6	7	3	2	3	-	1	-
2001.....	5	4	7	3	5	9	9	4	-	3	-	-
2000.....	4	2	2	5	4	6	3	8	1	1	-	-
1999.....	5	4	2	7	4	6	8	4	-	1	-	-

Asian or Pacific Islander 2/, male

2004.....	1	3	1	1	4	2	5	2	1	-	-	-
2003.....	2	-	-	2	3	4	1	2	-	-	-	-
2002.....	1	2	2	1	3	3	2	1	1	-	-	-
2001.....	1	2	3	1	2	4	4	2	-	1	-	-
2000.....	2	-	1	1	-	1	-	3	1	-	-	-
1999.....	1	1	1	3	1	4	4	2	-	1	-	-

Asian or Pacific Islander 2/, female

2004.....	2	5	4	7	3	1	9	2	5	-	-	-
2003.....	3	5	4	4	5	8	6	3	2	1	-	-
2002.....	4	3	1	4	3	4	1	1	2	-	1	-
2001.....	4	2	4	2	3	5	5	2	-	2	-	-
2000.....	2	2	1	4	4	5	3	5	-	1	-	-
1999.....	4	3	1	4	3	2	4	2	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other non-pulmonary forms of heart disease (I30-I51)												
All races, both sexes												
2004.....	4,270	5,107	6,351	7,890	11,532	18,143	26,142	29,285	22,962	9,730	2,124	6
2003.....	4,069	5,321	6,250	8,346	12,258	18,879	26,464	29,429	23,099	9,746	2,262	6
2002.....	4,145	5,031	6,089	8,165	12,785	19,367	26,613	29,276	22,739	9,530	2,169	5
2001.....	4,060	4,803	5,984	8,448	12,939	19,530	25,861	28,866	22,006	9,406	2,171	7
2000.....	3,811	4,714	6,023	8,705	13,312	19,734	25,129	28,075	21,975	9,351	2,159	8
1999.....	3,727	4,732	6,256	8,900	13,948	19,803	25,243	28,023	21,122	9,147	2,082	5
All races, male												
2004.....	2,787	3,307	3,949	4,681	6,301	9,132	11,513	10,896	6,736	2,092	321	5
2003.....	2,651	3,438	3,870	4,886	6,805	9,531	11,610	10,823	6,609	2,111	352	5
2002.....	2,759	3,203	3,743	4,729	7,088	9,733	11,629	10,698	6,368	2,009	334	3
2001.....	2,641	3,042	3,676	4,839	7,125	9,832	11,318	10,136	5,981	1,974	328	5
2000.....	2,529	3,063	3,723	5,078	7,390	9,741	10,886	10,049	5,993	1,941	353	5
1999.....	2,455	3,083	3,845	5,342	7,728	9,798	10,927	9,856	5,658	1,935	314	3
All races, female												
2004.....	1,483	1,800	2,402	3,209	5,231	9,011	14,629	18,389	16,226	7,638	1,803	1
2003.....	1,418	1,883	2,380	3,460	5,453	9,348	14,854	18,606	16,490	7,635	1,910	1
2002.....	1,386	1,828	2,346	3,436	5,697	9,634	14,984	18,578	16,371	7,521	1,835	2
2001.....	1,419	1,761	2,308	3,609	5,814	9,698	14,543	18,730	16,025	7,432	1,843	2
2000.....	1,282	1,651	2,300	3,627	5,922	9,993	14,243	18,026	15,982	7,410	1,806	3
1999.....	1,272	1,649	2,411	3,558	6,220	10,005	14,316	18,167	15,464	7,212	1,768	2
White, both sexes												
2004.....	3,005	3,782	4,975	6,363	9,702	15,810	23,729	27,075	21,307	9,018	1,895	3
2003.....	2,828	4,026	4,816	6,700	10,295	16,623	23,989	27,228	21,494	9,013	2,015	5
2002.....	2,928	3,804	4,736	6,628	10,826	17,035	24,163	27,101	21,103	8,813	1,913	3
2001.....	2,899	3,576	4,715	6,827	11,014	17,147	23,463	26,707	20,524	8,734	1,937	6
2000.....	2,740	3,569	4,723	7,080	11,308	17,381	22,911	25,951	20,459	8,647	1,937	7
1999.....	2,694	3,525	4,887	7,265	11,886	17,403	22,982	25,812	19,756	8,462	1,860	3

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other non-pulmonary forms of heart disease (I30-I51)												
White, male												
2004.....	2,037	2,499	3,132	3,808	5,393	8,022	10,516	10,100	6,211	1,925	275	2
2003.....	1,880	2,631	3,045	3,972	5,816	8,497	10,581	10,069	6,160	1,944	299	4
2002.....	1,999	2,464	2,961	3,887	6,074	8,642	10,647	9,959	5,880	1,859	286	2
2001.....	1,959	2,307	2,945	3,989	6,116	8,695	10,293	9,395	5,559	1,814	284	5
2000.....	1,842	2,372	2,976	4,184	6,377	8,651	9,988	9,303	5,563	1,763	306	5
1999.....	1,808	2,329	3,048	4,422	6,680	8,689	10,000	9,035	5,246	1,761	268	2
White, female												
2004.....	968	1,283	1,843	2,555	4,309	7,788	13,213	16,975	15,096	7,093	1,620	1
2003.....	948	1,395	1,771	2,728	4,479	8,126	13,408	17,159	15,334	7,069	1,716	1
2002.....	929	1,340	1,775	2,741	4,752	8,393	13,516	17,142	15,223	6,954	1,627	1
2001.....	940	1,269	1,770	2,838	4,898	8,452	13,170	17,312	14,965	6,920	1,653	1
2000.....	898	1,197	1,747	2,896	4,931	8,730	12,923	16,648	14,896	6,884	1,631	2
1999.....	886	1,196	1,839	2,843	5,206	8,714	12,982	16,777	14,510	6,701	1,592	1
Black, both sexes												
2004.....	1,172	1,216	1,237	1,346	1,616	1,998	2,038	1,824	1,365	585	193	3
2003.....	1,145	1,174	1,285	1,455	1,730	1,927	2,068	1,830	1,316	639	209	1
2002.....	1,124	1,107	1,222	1,382	1,682	1,997	2,084	1,816	1,394	619	218	2
2001.....	1,075	1,121	1,156	1,444	1,679	2,091	2,055	1,856	1,274	576	202	1
2000.....	986	1,042	1,165	1,454	1,781	2,069	1,926	1,845	1,288	597	200	1
1999.....	948	1,105	1,245	1,471	1,836	2,093	1,950	1,914	1,161	598	187	2
Black, male												
2004.....	693	740	735	763	802	940	824	633	386	116	33	3
2003.....	707	727	746	814	863	870	837	596	330	131	41	1
2002.....	696	664	702	750	876	907	817	568	378	112	36	1
2001.....	633	673	665	755	862	983	858	617	321	128	39	-
2000.....	634	625	665	798	875	935	766	621	337	138	40	-
1999.....	601	688	725	824	928	939	785	667	330	144	39	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other non-pulmonary forms of heart disease (I30-I51)

Black, female

2004.....	479	476	502	583	814	1,058	1,214	1,191	979	469	160	-
2003.....	438	447	539	641	867	1,057	1,231	1,234	986	508	168	-
2002.....	428	443	520	632	806	1,090	1,267	1,248	1,016	507	182	1
2001.....	442	448	491	689	817	1,108	1,197	1,239	953	448	163	1
2000.....	352	417	500	656	906	1,134	1,160	1,224	951	459	160	1
1999.....	347	417	520	647	908	1,154	1,165	1,247	831	454	148	1

American Indian 1/, both sexes

2004.....	31	35	45	52	46	62	61	61	35	23	7	-
2003.....	37	42	64	56	51	59	71	57	44	15	11	-
2002.....	30	33	33	33	55	72	65	63	36	17	10	-
2001.....	20	35	35	41	59	63	69	65	32	18	6	-
2000.....	33	31	36	50	59	72	63	60	34	17	6	-
1999.....	28	28	42	50	55	75	60	47	43	15	11	-

American Indian 1/, male

2004.....	14	22	25	29	25	32	19	30	12	4	2	-
2003.....	23	30	29	24	30	23	27	26	21	3	4	-
2002.....	23	22	23	19	35	39	27	30	10	5	6	-
2001.....	12	21	21	23	36	38	33	23	11	4	-	-
2000.....	22	22	21	26	39	38	26	25	7	2	-	-
1999.....	16	17	23	33	30	38	23	17	12	5	4	-

American Indian 1/, female

2004.....	17	13	20	23	21	30	42	31	23	19	5	-
2003.....	14	12	35	32	21	36	44	31	23	12	7	-
2002.....	7	11	10	14	20	33	38	33	26	12	4	-
2001.....	8	14	14	18	23	25	36	42	21	14	6	-
2000.....	11	9	15	24	20	34	37	35	27	15	6	-
1999.....	12	11	19	17	25	37	37	30	31	10	7	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other non-pulmonary forms of heart disease (I30-I51)

Asian or Pacific Islander 2/, both sexes

2004.....	62	74	94	129	168	273	314	325	255	104	29	-
2003.....	59	79	85	135	182	270	336	314	245	79	27	-
2002.....	63	87	98	122	222	263	301	296	206	81	28	-
2001.....	66	71	78	136	187	229	274	238	176	78	26	-
2000.....	52	72	99	121	164	212	229	219	194	90	16	-
1999.....	57	74	82	114	171	232	251	250	162	72	24	-

Asian or Pacific Islander 2/, male

2004.....	43	46	57	81	81	138	154	133	127	47	11	-
2003.....	41	50	50	76	96	141	165	132	98	33	8	-
2002.....	41	53	57	73	103	145	138	141	100	33	6	-
2001.....	37	41	45	72	111	116	134	101	90	28	5	-
2000.....	31	44	61	70	99	117	106	100	86	38	7	-
1999.....	30	49	49	63	90	132	119	137	70	25	3	-

Asian or Pacific Islander 2/, female

2004.....	19	28	37	48	87	135	160	192	128	57	18	-
2003.....	18	29	35	59	86	129	171	182	147	46	19	-
2002.....	22	34	41	49	119	118	163	155	106	48	22	-
2001.....	29	30	33	64	76	113	140	137	86	50	21	-
2000.....	21	28	38	51	65	95	123	119	108	52	9	-
1999.....	27	25	33	51	81	100	132	113	92	47	21	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute and subacute endocarditis (I33)												
All races, both sexes												
2004.....	100	81	84	103	131	157	161	122	21	11	1	1
2003.....	94	95	114	102	130	172	146	104	32	3	1	-
2002.....	90	73	95	107	131	161	145	76	36	8	1	-
2001.....	93	72	88	102	145	162	154	79	30	2	-	-
2000.....	97	88	77	105	119	166	136	83	30	5	1	-
1999.....	75	93	78	103	137	155	134	92	21	5	1	-
All races, male												
2004.....	64	52	46	67	52	72	90	58	10	4	-	1
2003.....	65	60	63	57	62	89	77	43	15	1	-	-
2002.....	62	42	49	66	67	78	75	31	15	2	1	-
2001.....	61	39	56	49	83	94	83	37	11	1	-	-
2000.....	61	52	44	59	63	95	74	40	9	2	-	-
1999.....	54	62	41	51	67	83	71	46	10	-	-	-
All races, female												
2004.....	36	29	38	36	79	85	71	64	11	7	1	-
2003.....	29	35	51	45	68	83	69	61	17	2	1	-
2002.....	28	31	46	41	64	83	70	45	21	6	-	-
2001.....	32	33	32	53	62	68	71	42	19	1	-	-
2000.....	36	36	33	46	56	71	62	43	21	3	1	-
1999.....	21	31	37	52	70	72	63	46	11	5	1	-
White, both sexes												
2004.....	59	60	66	83	111	132	147	113	19	8	-	-
2003.....	61	66	83	83	104	150	134	98	32	3	1	-
2002.....	60	46	79	82	110	145	138	70	33	8	1	-
2001.....	61	53	64	74	117	134	140	76	27	2	-	-
2000.....	65	59	63	89	102	148	122	74	26	5	-	-
1999.....	48	58	59	82	114	142	122	87	20	5	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute and subacute endocarditis (I33)												
White, male												
2004.....	39	41	38	54	47	62	83	54	10	3	-	-
2003.....	41	40	51	47	50	79	72	42	15	1	-	-
2002.....	45	25	43	52	58	74	71	28	14	2	1	-
2001.....	42	28	42	35	63	82	75	35	10	1	-	-
2000.....	44	38	38	47	57	87	66	37	8	2	-	-
1999.....	33	42	31	45	57	81	66	45	10	-	-	-
White, female												
2004.....	20	19	28	29	64	70	64	59	9	5	-	-
2003.....	20	26	32	36	54	71	62	56	17	2	1	-
2002.....	15	21	36	30	52	71	67	42	19	6	-	-
2001.....	19	25	22	39	54	52	65	41	17	1	-	-
2000.....	21	21	25	42	45	61	56	37	18	3	-	-
1999.....	15	16	28	37	57	61	56	42	10	5	-	-
Black, both sexes												
2004.....	38	18	17	18	18	18	9	9	2	3	1	1
2003.....	32	27	26	17	21	19	10	4	-	-	-	-
2002.....	28	25	16	23	16	14	5	5	3	-	-	-
2001.....	30	19	22	21	22	27	12	2	3	-	-	-
2000.....	27	26	13	15	16	16	10	6	4	-	1	-
1999.....	23	30	18	21	21	10	7	5	1	-	-	-
Black, male												
2004.....	23	10	8	11	5	7	3	4	-	1	-	1
2003.....	23	18	9	9	9	8	5	-	-	-	-	-
2002.....	15	16	6	12	8	3	3	3	1	-	-	-
2001.....	18	11	13	10	16	12	7	1	1	-	-	-
2000.....	15	12	6	11	5	6	5	1	1	-	-	-
1999.....	19	18	9	6	9	1	4	1	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute and subacute endocarditis (I33)												
Black, female												
2004.....	15	8	9	7	13	11	6	5	2	2	1	-
2003.....	9	9	17	8	12	11	5	4	-	-	-	-
2002.....	13	9	10	11	8	11	2	2	2	-	-	-
2001.....	12	8	9	11	6	15	5	1	2	-	-	-
2000.....	12	14	7	4	11	10	5	5	3	-	1	-
1999.....	4	12	9	15	12	9	3	4	1	-	-	-
American Indian 1/, both sexes												
2004.....	1	1	1	-	1	-	2	-	-	-	-	-
2003.....	-	-	2	1	1	1	-	1	-	-	-	-
2002.....	-	1	-	-	-	2	-	-	-	-	-	-
2001.....	1	-	-	2	-	-	-	-	-	-	-	-
2000.....	3	1	1	-	-	1	1	1	-	-	-	-
1999.....	1	1	-	-	-	-	2	-	-	-	-	-
American Indian 1/, male												
2004.....	1	-	-	-	-	-	1	-	-	-	-	-
2003.....	-	-	-	-	-	1	-	1	-	-	-	-
2002.....	-	-	-	-	-	1	-	-	-	-	-	-
2001.....	-	-	-	2	-	-	-	-	-	-	-	-
2000.....	2	1	-	-	-	1	1	-	-	-	-	-
1999.....	1	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, female												
2004.....	-	1	1	-	1	-	1	-	-	-	-	-
2003.....	-	-	2	1	1	-	-	-	-	-	-	-
2002.....	-	1	-	-	-	1	-	-	-	-	-	-
2001.....	1	-	-	-	-	-	-	-	-	-	-	-
2000.....	1	-	1	-	-	-	-	1	-	-	-	-
1999.....	-	1	-	-	-	-	1	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute and subacute endocarditis (I33)												
Asian or Pacific Islander 2/, both sexes												
2004.....	2	2	-	2	1	7	3	-	-	-	-	-
2003.....	1	2	3	1	4	2	2	1	-	-	-	-
2002.....	2	1	-	2	5	-	2	1	-	-	-	-
2001.....	1	-	2	5	6	1	2	1	-	-	-	-
2000.....	2	2	-	1	1	1	3	2	-	-	-	-
1999.....	3	4	1	-	2	3	3	-	-	-	1	-
Asian or Pacific Islander 2/, male												
2004.....	1	1	-	2	-	3	3	-	-	-	-	-
2003.....	1	2	3	1	3	1	-	-	-	-	-	-
2002.....	2	1	-	2	1	-	1	-	-	-	-	-
2001.....	1	-	1	2	4	-	1	1	-	-	-	-
2000.....	-	1	-	1	1	1	2	2	-	-	-	-
1999.....	1	2	1	-	1	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	1	1	-	-	1	4	-	-	-	-	-	-
2003.....	-	-	-	-	1	1	2	1	-	-	-	-
2002.....	-	-	-	-	4	-	1	1	-	-	-	-
2001.....	-	-	1	3	2	1	1	-	-	-	-	-
2000.....	2	1	-	-	-	-	1	-	-	-	-	-
1999.....	2	2	-	-	1	2	3	-	-	-	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
All races, both sexes												
2004.....	58	52	59	62	53	84	75	56	25	6	-	-
2003.....	48	59	45	80	71	70	83	45	27	5	-	-
2002.....	51	66	52	55	88	84	73	65	25	6	-	-
2001.....	54	54	52	47	73	87	67	49	28	7	1	1
2000.....	40	47	42	52	79	82	63	43	24	8	1	-
1999.....	46	63	62	69	77	88	72	56	25	4	-	-
All races, male												
2004.....	36	30	35	36	26	38	30	21	10	3	-	-
2003.....	30	32	25	44	40	34	31	13	8	1	-	-
2002.....	29	34	35	29	42	34	26	25	6	2	-	-
2001.....	29	28	33	28	31	39	22	14	8	2	-	1
2000.....	18	31	21	24	43	42	31	17	7	1	1	-
1999.....	25	40	30	34	35	38	27	19	11	-	-	-
All races, female												
2004.....	22	22	24	26	27	46	45	35	15	3	-	-
2003.....	18	27	20	36	31	36	52	32	19	4	-	-
2002.....	22	32	17	26	46	50	47	40	19	4	-	-
2001.....	25	26	19	19	42	48	45	35	20	5	1	-
2000.....	22	16	21	28	36	40	32	26	17	7	-	-
1999.....	21	23	32	35	42	50	45	37	14	4	-	-
White, both sexes												
2004.....	46	46	48	49	47	68	67	49	24	5	-	-
2003.....	38	51	39	71	62	63	73	41	26	5	-	-
2002.....	36	49	42	48	75	67	64	62	23	5	-	-
2001.....	38	42	43	36	68	80	61	43	27	7	-	1
2000.....	33	39	33	41	67	69	61	38	21	8	1	-
1999.....	35	53	53	57	65	77	65	50	22	4	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
White, male												
2004.....	28	26	30	26	22	33	26	18	10	2	-	-
2003.....	25	26	23	41	36	31	30	13	8	1	-	-
2002.....	21	26	29	25	38	30	22	25	5	1	-	-
2001.....	21	22	26	20	31	35	20	13	8	2	-	1
2000.....	15	28	17	15	37	34	31	15	6	1	1	-
1999.....	18	34	27	27	30	35	25	17	10	-	-	-
White, female												
2004.....	18	20	18	23	25	35	41	31	14	3	-	-
2003.....	13	25	16	30	26	32	43	28	18	4	-	-
2002.....	15	23	13	23	37	37	42	37	18	4	-	-
2001.....	17	20	17	16	37	45	41	30	19	5	-	-
2000.....	18	11	16	26	30	35	30	23	15	7	-	-
1999.....	17	19	26	30	35	42	40	33	12	4	-	-
Black, both sexes												
2004.....	11	6	10	11	6	14	8	5	1	-	-	-
2003.....	7	7	5	7	9	5	8	4	1	-	-	-
2002.....	14	14	8	7	12	14	6	3	2	-	-	-
2001.....	16	11	8	9	4	6	5	5	1	-	1	-
2000.....	7	7	8	10	8	8	2	5	2	-	-	-
1999.....	11	9	8	10	10	10	6	4	2	-	-	-
Black, male												
2004.....	7	4	4	9	4	5	4	1	-	-	-	-
2003.....	4	5	2	1	4	2	1	-	-	-	-	-
2002.....	7	8	5	4	3	4	3	-	1	-	-	-
2001.....	8	5	6	8	-	3	2	-	-	-	-	-
2000.....	3	2	4	8	3	5	-	2	-	-	-	-
1999.....	7	6	3	6	5	2	2	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Black, female												
2004.....	4	2	6	2	2	9	4	4	1	-	-	-
2003.....	3	2	3	6	5	3	7	4	1	-	-	-
2002.....	7	6	3	3	9	10	3	3	1	-	-	-
2001.....	8	6	2	1	4	3	3	5	1	-	1	-
2000.....	4	5	4	2	5	3	2	3	2	-	-	-
1999.....	4	3	5	4	5	8	4	4	2	-	-	-
American Indian 1/, both sexes												
2004.....	-	-	-	-	-	-	-	1	-	-	-	-
2003.....	1	1	-	1	-	-	-	-	-	-	-	-
2002.....	1	-	-	-	-	2	-	-	-	-	-	-
2001.....	-	1	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	1	2	-	-	-	-	-	-
1999.....	-	-	1	-	2	-	-	-	-	-	-	-
American Indian 1/, male												
2004.....	-	-	-	-	-	-	-	1	-	-	-	-
2003.....	1	1	-	1	-	-	-	-	-	-	-	-
2002.....	1	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	1	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	1	1	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, female												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	2	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	1	-	-	-	-	-	-
1999.....	-	-	1	-	2	-	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Asian or Pacific Islander 2/, both sexes												
2004.....	1	-	1	2	-	2	-	1	-	1	-	-
2003.....	2	-	1	1	-	2	2	-	-	-	-	-
2002.....	-	3	2	-	1	1	3	-	-	1	-	-
2001.....	-	-	1	2	1	1	1	1	-	-	-	-
2000.....	-	1	1	1	3	3	-	-	1	-	-	-
1999.....	-	1	-	2	-	1	1	2	1	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	1	-	1	1	-	-	-	1	-	1	-	-
2003.....	-	-	-	1	-	1	-	-	-	-	-	-
2002.....	-	-	1	-	1	-	1	-	-	1	-	-
2001.....	-	-	1	-	-	1	-	1	-	-	-	-
2000.....	-	1	-	1	2	2	-	-	1	-	-	-
1999.....	-	-	-	1	-	1	-	2	1	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	-	-	-	1	-	2	-	-	-	-	-	-
2003.....	2	-	1	-	-	1	2	-	-	-	-	-
2002.....	-	3	1	-	-	1	2	-	-	-	-	-
2001.....	-	-	-	2	1	-	1	-	-	-	-	-
2000.....	-	-	1	-	1	1	-	-	-	-	-	-
1999.....	-	1	-	1	-	-	1	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of endocardium (I34-I38)												
All races, both sexes												
2004.....	383	457	601	827	1,395	2,334	3,724	4,620	3,504	1,395	214	1
2003.....	361	489	543	885	1,366	2,412	3,723	4,539	3,492	1,257	229	2
2002.....	354	429	563	809	1,383	2,448	3,825	4,523	3,465	1,173	205	1
2001.....	380	403	523	858	1,521	2,517	3,619	4,419	3,237	1,142	195	-
2000.....	328	410	553	851	1,581	2,532	3,623	4,357	3,290	1,196	195	1
1999.....	340	383	544	881	1,559	2,583	3,739	4,373	3,055	1,120	167	-
All races, male												
2004.....	233	285	312	421	668	1,057	1,490	1,644	971	284	30	1
2003.....	196	297	282	435	664	1,113	1,519	1,576	962	245	39	1
2002.....	207	239	287	385	668	1,119	1,505	1,560	929	230	40	1
2001.....	240	229	281	429	735	1,180	1,454	1,493	840	242	34	-
2000.....	192	245	304	410	749	1,069	1,448	1,485	880	252	31	1
1999.....	196	219	296	457	762	1,159	1,555	1,446	751	241	25	-
All races, female												
2004.....	150	172	289	406	727	1,277	2,234	2,976	2,533	1,111	184	-
2003.....	165	192	261	450	702	1,299	2,204	2,963	2,530	1,012	190	1
2002.....	147	190	276	424	715	1,329	2,320	2,963	2,536	943	165	-
2001.....	140	174	242	429	786	1,337	2,165	2,926	2,397	900	161	-
2000.....	136	165	249	441	832	1,463	2,175	2,872	2,410	944	164	-
1999.....	144	164	248	424	797	1,424	2,184	2,927	2,304	879	142	-
White, both sexes												
2004.....	272	368	505	699	1,264	2,153	3,508	4,416	3,370	1,326	198	-
2003.....	262	388	436	759	1,217	2,216	3,522	4,341	3,360	1,205	219	2
2002.....	264	345	474	678	1,250	2,291	3,635	4,342	3,337	1,107	191	1
2001.....	297	310	442	725	1,378	2,333	3,447	4,239	3,112	1,097	180	-
2000.....	254	315	463	735	1,415	2,335	3,449	4,157	3,169	1,157	181	1
1999.....	266	307	448	753	1,368	2,400	3,557	4,180	2,920	1,082	157	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of endocardium (I34-I38)												
White, male												
2004.....	169	235	266	371	617	977	1,410	1,568	927	270	28	-
2003.....	140	231	237	381	604	1,036	1,442	1,520	926	234	36	1
2002.....	163	192	245	337	611	1,053	1,438	1,510	893	222	37	1
2001.....	189	182	236	367	672	1,099	1,399	1,426	805	231	31	-
2000.....	150	197	261	364	678	995	1,382	1,420	848	243	28	1
1999.....	161	180	246	404	675	1,086	1,484	1,384	716	231	24	-
White, female												
2004.....	103	133	239	328	647	1,176	2,098	2,848	2,443	1,056	170	-
2003.....	122	157	199	378	613	1,180	2,080	2,821	2,434	971	183	1
2002.....	101	153	229	341	639	1,238	2,197	2,832	2,444	885	154	-
2001.....	108	128	206	358	706	1,234	2,048	2,813	2,307	866	149	-
2000.....	104	118	202	371	737	1,340	2,067	2,737	2,321	914	153	-
1999.....	105	127	202	349	693	1,314	2,073	2,796	2,204	851	133	-
Black, both sexes												
2004.....	104	78	81	112	106	134	164	158	95	48	12	1
2003.....	94	87	90	97	122	153	148	135	102	46	9	-
2002.....	80	72	77	114	103	129	142	132	97	52	10	-
2001.....	78	83	71	113	112	148	141	135	97	34	12	-
2000.....	67	79	79	99	142	168	140	164	89	32	12	-
1999.....	66	69	84	103	161	145	145	142	105	34	6	-
Black, male												
2004.....	60	43	41	41	43	55	56	54	25	5	1	1
2003.....	54	60	39	43	46	59	54	36	22	8	2	-
2002.....	39	41	36	43	42	53	46	32	25	5	1	-
2001.....	47	43	39	51	47	64	41	44	22	8	2	-
2000.....	39	40	36	40	59	63	51	50	17	6	2	-
1999.....	33	32	44	43	74	57	54	34	23	9	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other diseases of endocardium (I34-I38)

Black, female

2004.....	44	35	40	71	63	79	108	104	70	43	11	-
2003.....	40	27	51	54	76	94	94	99	80	38	7	-
2002.....	41	31	41	71	61	76	96	100	72	47	9	-
2001.....	31	40	32	62	65	84	100	91	75	26	10	-
2000.....	28	39	43	59	83	105	89	114	72	26	10	-
1999.....	33	37	40	60	87	88	91	108	82	25	6	-

American Indian 1/, both sexes

2004.....	1	7	7	2	6	10	9	6	3	3	1	-
2003.....	2	3	5	9	4	6	9	9	3	2	-	-
2002.....	3	2	3	4	4	6	11	8	4	3	1	-
2001.....	2	3	2	2	7	4	4	7	2	3	-	-
2000.....	2	6	3	2	3	9	8	6	2	2	-	-
1999.....	2	-	4	6	5	8	7	9	5	-	1	-

American Indian 1/, male

2004.....	-	3	4	1	4	7	5	3	2	-	-	-
2003.....	1	3	2	4	3	3	3	4	3	1	-	-
2002.....	2	1	3	1	1	2	6	2	1	-	1	-
2001.....	2	1	1	1	3	2	2	4	-	1	-	-
2000.....	1	3	3	-	-	2	1	2	1	-	-	-
1999.....	1	-	2	4	3	2	4	4	2	-	-	-

American Indian 1/, female

2004.....	1	4	3	1	2	3	4	3	1	3	1	-
2003.....	1	-	3	5	1	3	6	5	-	1	-	-
2002.....	1	1	-	3	3	4	5	6	3	3	-	-
2001.....	-	2	1	1	4	2	2	3	2	2	-	-
2000.....	1	3	-	2	3	7	7	4	1	2	-	-
1999.....	1	-	2	2	2	6	3	5	3	-	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of endocardium (I34-I38)												
Asian or Pacific Islander 2/, both sexes												
2004.....	6	4	8	14	19	37	43	40	36	18	3	-
2003.....	3	11	12	20	23	37	44	54	27	4	1	-
2002.....	7	10	9	13	26	22	37	41	27	11	3	-
2001.....	3	7	8	18	24	32	27	38	26	8	3	-
2000.....	5	10	8	15	21	20	26	30	30	5	2	-
1999.....	6	7	8	19	25	30	30	42	25	4	3	-
Asian or Pacific Islander 2/, male												
2004.....	4	4	1	8	4	18	19	19	17	9	1	-
2003.....	1	3	4	7	11	15	20	16	11	2	1	-
2002.....	3	5	3	4	14	11	15	16	10	3	1	-
2001.....	2	3	5	10	13	15	12	19	13	2	1	-
2000.....	2	5	4	6	12	9	14	13	14	3	1	-
1999.....	1	7	4	6	10	14	13	24	10	1	1	-
Asian or Pacific Islander 2/, female												
2004.....	2	-	7	6	15	19	24	21	19	9	2	-
2003.....	2	8	8	13	12	22	24	38	16	2	-	-
2002.....	4	5	6	9	12	11	22	25	17	8	2	-
2001.....	1	4	3	8	11	17	15	19	13	6	2	-
2000.....	3	5	4	9	9	11	12	17	16	2	1	-
1999.....	5	-	4	13	15	16	17	18	15	3	2	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic mitral valve disorders (I34)												
All races, both sexes												
2004.....	54	72	85	126	211	352	466	509	340	113	17	-
2003.....	65	81	100	138	204	330	488	507	323	93	24	-
2002.....	57	72	118	146	242	373	523	554	333	116	14	-
2001.....	70	72	83	162	241	421	495	531	342	106	21	-
2000.....	55	73	103	148	262	429	524	529	360	119	23	-
1999.....	65	70	98	151	272	424	514	561	373	95	13	-
All races, male												
2004.....	28	35	33	52	84	136	165	146	83	21	1	-
2003.....	30	42	45	58	92	139	168	133	81	19	4	-
2002.....	26	31	49	56	94	128	164	196	74	20	2	-
2001.....	32	37	37	73	107	145	157	152	78	16	1	-
2000.....	22	30	43	69	102	169	180	164	87	29	1	-
1999.....	26	27	45	59	119	169	191	163	77	25	-	-
All races, female												
2004.....	26	37	52	74	127	216	301	363	257	92	16	-
2003.....	35	39	55	80	112	191	320	374	242	74	20	-
2002.....	31	41	69	90	148	245	359	358	259	96	12	-
2001.....	38	35	46	89	134	276	338	379	264	90	20	-
2000.....	33	43	60	79	160	260	344	365	273	90	22	-
1999.....	39	43	53	92	153	255	323	398	296	70	13	-
White, both sexes												
2004.....	42	60	68	109	188	322	436	478	326	106	14	-
2003.....	49	66	78	118	191	303	455	480	314	92	23	-
2002.....	40	57	104	124	220	342	497	537	328	109	14	-
2001.....	56	60	71	135	215	400	472	511	330	105	18	-
2000.....	44	59	90	130	238	399	506	495	347	114	21	-
1999.....	50	59	83	127	241	388	492	535	359	91	12	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic mitral valve disorders (I34)												
White, male												
2004.....	23	29	27	47	77	123	154	136	81	20	1	-
2003.....	22	33	39	50	87	128	158	129	78	19	4	-
2002.....	17	26	42	50	83	118	159	191	72	19	2	-
2001.....	29	33	33	63	96	140	153	147	77	16	1	-
2000.....	18	25	38	65	94	160	175	157	83	28	1	-
1999.....	22	24	41	52	104	156	188	158	73	25	-	-
White, female												
2004.....	19	31	41	62	111	199	282	342	245	86	13	-
2003.....	27	33	39	68	104	175	297	351	236	73	19	-
2002.....	23	31	62	74	137	224	338	346	256	90	12	-
2001.....	27	27	38	72	119	260	319	364	253	89	17	-
2000.....	26	34	52	65	144	239	331	338	264	86	20	-
1999.....	28	35	42	75	137	232	304	377	286	66	12	-
Black, both sexes												
2004.....	10	11	17	16	17	22	21	20	10	3	2	-
2003.....	15	12	18	15	11	20	21	18	7	1	1	-
2002.....	13	13	11	20	16	22	16	13	3	7	-	-
2001.....	13	12	10	23	21	14	19	13	9	1	2	-
2000.....	10	12	11	15	20	28	13	28	6	3	2	-
1999.....	11	10	14	17	25	25	15	18	10	3	1	-
Black, male												
2004.....	3	6	6	5	6	9	6	5	1	1	-	-
2003.....	7	7	5	6	5	7	6	2	3	-	-	-
2002.....	7	4	5	5	7	7	4	3	1	1	-	-
2001.....	2	4	4	9	8	3	3	2	-	-	-	-
2000.....	3	4	4	3	6	8	4	5	2	-	-	-
1999.....	2	2	4	4	14	8	1	2	2	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic mitral valve disorders (I34)												
Black, female												
2004.....	7	5	11	11	11	13	15	15	9	2	2	-
2003.....	8	5	13	9	6	13	15	16	4	1	1	-
2002.....	6	9	6	15	9	15	12	10	2	6	-	-
2001.....	11	8	6	14	13	11	16	11	9	1	2	-
2000.....	7	8	7	12	14	20	9	23	4	3	2	-
1999.....	9	8	10	13	11	17	14	16	8	3	1	-
American Indian 1/, both sexes												
2004.....	-	1	-	-	-	-	3	-	1	-	-	-
2003.....	-	-	-	1	1	-	3	3	-	-	-	-
2002.....	-	-	1	-	-	2	1	-	-	-	-	-
2001.....	-	-	1	1	1	1	-	-	1	-	-	-
2000.....	-	-	1	-	-	-	1	1	1	1	-	-
1999.....	1	-	1	2	-	1	1	1	1	-	-	-
American Indian 1/, male												
2004.....	-	-	-	-	-	-	1	-	-	-	-	-
2003.....	-	-	-	-	-	-	1	1	-	-	-	-
2002.....	-	-	1	-	-	-	1	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	1	-	-	-	-	-	-	-	-	-
1999.....	1	-	-	2	-	-	1	1	-	-	-	-
American Indian 1/, female												
2004.....	-	1	-	-	-	-	2	-	1	-	-	-
2003.....	-	-	-	1	1	-	2	2	-	-	-	-
2002.....	-	-	-	-	-	2	-	-	-	-	-	-
2001.....	-	-	1	1	1	1	-	-	1	-	-	-
2000.....	-	-	-	-	-	-	1	1	1	1	-	-
1999.....	-	-	1	-	-	1	-	-	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic mitral valve disorders (I34)												
Asian or Pacific Islander 2/, both sexes												
2004.....	2	-	-	1	6	8	6	11	3	4	1	-
2003.....	1	3	4	4	1	7	9	6	2	-	-	-
2002.....	4	2	2	2	6	7	9	4	2	-	-	-
2001.....	1	-	1	3	4	6	4	7	2	-	1	-
2000.....	1	2	1	3	4	2	4	5	6	1	-	-
1999.....	3	1	-	5	6	10	6	7	3	1	-	-
Asian or Pacific Islander 2/, male												
2004.....	2	-	-	-	1	4	4	5	1	-	-	-
2003.....	1	2	1	2	-	4	3	1	-	-	-	-
2002.....	2	1	1	1	4	3	-	2	1	-	-	-
2001.....	1	-	-	1	3	2	1	3	1	-	-	-
2000.....	1	1	-	1	2	1	1	2	2	1	-	-
1999.....	1	1	-	1	1	5	1	2	2	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	-	-	-	1	5	4	2	6	2	4	1	-
2003.....	-	1	3	2	1	3	6	5	2	-	-	-
2002.....	2	1	1	1	2	4	9	2	1	-	-	-
2001.....	-	-	1	2	1	4	3	4	1	-	1	-
2000.....	-	1	1	2	2	1	3	3	4	-	-	-
1999.....	2	-	-	4	5	5	5	5	1	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic aortic valve disorders (I35)												
All races, both sexes												
2004.....	148	206	276	375	756	1,363	2,372	3,197	2,517	1,053	157	-
2003.....	120	204	229	434	732	1,372	2,386	3,123	2,579	958	170	-
2002.....	143	176	261	382	761	1,428	2,412	3,090	2,531	876	154	1
2001.....	164	168	258	429	812	1,479	2,296	3,008	2,346	853	139	-
2000.....	141	192	251	426	878	1,485	2,329	3,060	2,344	880	131	-
1999.....	138	168	251	455	824	1,518	2,438	3,007	2,174	843	114	-
All races, male												
2004.....	96	136	160	213	390	650	975	1,178	714	225	23	-
2003.....	70	135	140	248	379	679	1,035	1,147	726	190	28	-
2002.....	89	111	146	200	416	713	1,033	1,098	714	171	31	1
2001.....	116	105	153	235	429	755	978	1,051	626	188	30	-
2000.....	101	135	160	229	456	656	970	1,086	650	183	22	-
1999.....	96	110	153	260	454	732	1,061	1,025	568	180	20	-
All races, female												
2004.....	52	70	116	162	366	713	1,397	2,019	1,803	828	134	-
2003.....	50	69	89	186	353	693	1,351	1,976	1,853	768	142	-
2002.....	54	65	115	182	345	715	1,379	1,992	1,817	705	123	-
2001.....	48	63	105	194	383	724	1,318	1,957	1,720	665	109	-
2000.....	40	57	91	197	422	829	1,359	1,974	1,694	697	109	-
1999.....	42	58	98	195	370	786	1,377	1,982	1,606	663	94	-
White, both sexes												
2004.....	115	179	246	330	700	1,254	2,247	3,072	2,419	1,000	148	-
2003.....	98	172	200	389	660	1,266	2,269	2,997	2,486	916	163	-
2002.....	117	156	231	324	698	1,344	2,298	2,960	2,431	828	143	1
2001.....	135	141	232	378	746	1,361	2,199	2,889	2,254	818	128	-
2000.....	120	158	215	367	796	1,392	2,219	2,934	2,260	852	121	-
1999.....	111	142	216	397	725	1,434	2,318	2,879	2,077	816	108	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic aortic valve disorders (I35)												
White, male												
2004.....	76	118	138	193	363	603	928	1,129	680	213	21	-
2003.....	58	113	126	226	347	632	993	1,110	702	180	26	-
2002.....	74	97	131	176	388	676	990	1,062	685	166	28	1
2001.....	93	91	135	209	395	693	943	1,005	595	178	27	-
2000.....	85	115	143	201	417	621	925	1,043	627	176	21	-
1999.....	79	94	132	235	402	696	1,007	984	540	173	19	-
White, female												
2004.....	39	61	108	137	337	651	1,319	1,943	1,739	787	127	-
2003.....	40	59	74	163	313	634	1,276	1,887	1,784	736	137	-
2002.....	43	59	100	148	310	668	1,308	1,898	1,746	662	115	-
2001.....	42	50	97	169	351	668	1,256	1,884	1,659	640	101	-
2000.....	35	43	72	166	379	771	1,294	1,891	1,633	676	100	-
1999.....	32	48	84	162	323	738	1,311	1,895	1,537	643	89	-
Black, both sexes												
2004.....	29	24	28	39	44	78	94	98	68	38	6	-
2003.....	20	27	21	31	59	83	83	82	74	38	7	-
2002.....	23	17	25	48	51	67	89	93	74	38	7	-
2001.....	27	24	22	41	51	94	79	88	70	27	9	-
2000.....	18	28	29	50	68	78	90	102	65	23	9	-
1999.....	25	22	30	45	79	67	93	100	75	24	3	-
Black, male												
2004.....	18	15	22	16	20	31	33	35	19	4	1	-
2003.....	11	20	12	15	24	36	27	21	15	8	2	-
2002.....	13	12	13	21	23	29	29	23	21	4	1	-
2001.....	21	11	14	19	24	49	24	30	21	7	2	-
2000.....	15	17	13	24	32	28	36	32	12	5	1	-
1999.....	17	12	17	22	41	28	42	25	19	6	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic aortic valve disorders (I35)												
Black, female												
2004.....	11	9	6	23	24	47	61	63	49	34	5	-
2003.....	9	7	9	16	35	47	56	61	59	30	5	-
2002.....	10	5	12	27	28	38	60	70	53	34	6	-
2001.....	6	13	8	22	27	45	55	58	49	20	7	-
2000.....	3	11	16	26	36	50	54	70	53	18	8	-
1999.....	8	10	13	23	38	39	51	75	56	18	3	-
American Indian 1/, both sexes												
2004.....	1	1	1	-	5	7	4	5	2	3	1	-
2003.....	1	2	3	6	2	4	5	5	3	2	-	-
2002.....	2	-	1	3	1	4	8	7	3	2	1	-
2001.....	2	1	-	-	5	3	1	7	1	2	-	-
2000.....	1	3	1	-	3	5	5	3	1	1	-	-
1999.....	1	-	1	2	4	2	5	6	3	-	1	-
American Indian 1/, male												
2004.....	-	1	-	-	4	5	2	3	2	-	-	-
2003.....	1	2	1	3	2	2	2	3	3	1	-	-
2002.....	1	-	1	1	-	2	4	2	1	-	1	-
2001.....	2	1	-	-	3	2	1	4	-	1	-	-
2000.....	1	1	1	-	-	1	1	1	1	-	-	-
1999.....	-	-	1	-	2	-	2	2	1	-	-	-
American Indian 1/, female												
2004.....	1	-	1	-	1	2	2	2	-	3	1	-
2003.....	-	-	2	3	-	2	3	2	-	1	-	-
2002.....	1	-	-	2	1	2	4	5	2	2	-	-
2001.....	-	-	-	-	2	1	-	3	1	1	-	-
2000.....	-	2	-	-	3	4	4	2	-	1	-	-
1999.....	1	-	-	2	2	2	3	4	2	-	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic aortic valve disorders (I35)												
Asian or Pacific Islander 2/, both sexes												
2004.....	3	2	1	6	7	24	27	22	28	12	2	-
2003.....	1	3	5	8	11	19	29	39	16	2	-	-
2002.....	1	3	4	7	11	13	17	30	23	8	3	-
2001.....	-	2	4	10	10	21	17	24	21	6	2	-
2000.....	2	3	6	9	11	10	15	21	18	4	1	-
1999.....	1	4	4	11	16	15	22	22	19	3	2	-
Asian or Pacific Islander 2/, male												
2004.....	2	2	-	4	3	11	12	11	13	8	1	-
2003.....	-	-	1	4	6	9	13	13	6	1	-	-
2002.....	1	2	1	2	5	6	10	11	7	1	1	-
2001.....	-	2	4	7	7	11	10	12	10	2	1	-
2000.....	-	2	3	4	7	6	8	10	10	2	-	-
1999.....	-	4	3	3	9	8	10	14	8	1	1	-
Asian or Pacific Islander 2/, female												
2004.....	1	-	1	2	4	13	15	11	15	4	1	-
2003.....	1	3	4	4	5	10	16	26	10	1	-	-
2002.....	-	1	3	5	6	7	7	19	16	7	2	-
2001.....	-	-	-	3	3	10	7	12	11	4	1	-
2000.....	2	1	3	5	4	4	7	11	8	2	1	-
1999.....	1	-	1	8	7	7	12	8	11	2	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of endocardium (I36-I38)												
All races, both sexes												
2004.....	181	179	240	326	428	619	886	914	647	229	40	1
2003.....	176	204	214	313	430	710	849	909	590	206	35	2
2002.....	154	181	184	281	380	647	890	879	601	181	37	-
2001.....	146	163	182	267	468	617	828	880	549	183	35	-
2000.....	132	145	199	277	441	618	770	768	586	197	41	1
1999.....	137	145	195	275	463	641	787	805	508	182	40	-
All races, male												
2004.....	109	114	119	156	194	271	350	320	174	38	6	1
2003.....	96	120	97	129	193	295	316	296	155	36	7	1
2002.....	92	97	92	129	158	278	308	266	141	39	7	-
2001.....	92	87	91	121	199	280	319	290	136	38	3	-
2000.....	69	80	101	112	191	244	298	235	143	40	8	1
1999.....	74	82	98	138	189	258	303	258	106	36	5	-
All races, female												
2004.....	72	65	121	170	234	348	536	594	473	191	34	-
2003.....	80	84	117	184	237	415	533	613	435	170	28	1
2002.....	62	84	92	152	222	369	582	613	460	142	30	-
2001.....	54	76	91	146	269	337	509	590	413	145	32	-
2000.....	63	65	98	165	250	374	472	533	443	157	33	-
1999.....	63	63	97	137	274	383	484	547	402	146	35	-
White, both sexes												
2004.....	115	129	191	260	376	577	825	866	625	220	36	-
2003.....	115	150	158	252	366	647	798	864	560	197	33	2
2002.....	107	132	139	230	332	605	840	845	578	170	34	-
2001.....	106	109	139	212	417	572	776	839	528	174	34	-
2000.....	90	98	158	238	381	544	724	728	562	191	39	1
1999.....	105	106	149	229	402	578	747	766	484	175	37	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

All other diseases of endocardium (I36-I38)

White, male

2004.....	70	88	101	131	177	251	328	303	166	37	6	-
2003.....	60	85	72	105	170	276	291	281	146	35	6	1
2002.....	72	69	72	111	140	259	289	257	136	37	7	-
2001.....	67	58	68	95	181	266	303	274	133	37	3	-
2000.....	47	57	80	98	167	214	282	220	138	39	6	1
1999.....	60	62	73	117	169	234	289	242	103	33	5	-

White, female

2004.....	45	41	90	129	199	326	497	563	459	183	30	-
2003.....	55	65	86	147	196	371	507	583	414	162	27	1
2002.....	35	63	67	119	192	346	551	588	442	133	27	-
2001.....	39	51	71	117	236	306	473	565	395	137	31	-
2000.....	43	41	78	140	214	330	442	508	424	152	33	-
1999.....	45	44	76	112	233	344	458	524	381	142	32	-

Black, both sexes

2004.....	65	43	36	57	45	34	49	40	17	7	4	1
2003.....	59	48	51	51	52	50	44	35	21	7	1	-
2002.....	44	42	41	46	36	40	37	26	20	7	3	-
2001.....	38	47	39	49	40	40	43	34	18	6	1	-
2000.....	39	39	39	34	54	62	37	34	18	6	1	-
1999.....	30	37	40	41	57	53	37	24	20	7	2	-

Black, male

2004.....	39	22	13	20	17	15	17	14	5	-	-	1
2003.....	36	33	22	22	17	16	21	13	4	-	-	-
2002.....	19	25	18	17	12	17	13	6	3	-	-	-
2001.....	24	28	21	23	15	12	14	12	1	1	-	-
2000.....	21	19	19	13	21	27	11	13	3	1	1	-
1999.....	14	18	23	17	19	21	11	7	2	3	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of endocardium (I36-I38)												
Black, female												
2004.....	26	21	23	37	28	19	32	26	12	7	4	-
2003.....	23	15	29	29	35	34	23	22	17	7	1	-
2002.....	25	17	23	29	24	23	24	20	17	7	3	-
2001.....	14	19	18	26	25	28	29	22	17	5	1	-
2000.....	18	20	20	21	33	35	26	21	15	5	-	-
1999.....	16	19	17	24	38	32	26	17	18	4	2	-
American Indian 1/, both sexes												
2004.....	-	5	6	2	1	3	2	1	-	-	-	-
2003.....	1	1	2	2	1	2	1	1	-	-	-	-
2002.....	1	2	1	1	3	-	2	1	1	1	-	-
2001.....	-	2	1	1	1	-	3	-	-	1	-	-
2000.....	1	3	1	2	-	4	2	2	-	-	-	-
1999.....	-	-	2	2	1	5	1	2	1	-	-	-
American Indian 1/, male												
2004.....	-	2	4	1	-	2	2	-	-	-	-	-
2003.....	-	1	1	1	1	1	-	-	-	-	-	-
2002.....	1	1	1	-	1	-	1	-	-	-	-	-
2001.....	-	-	1	1	-	-	1	-	-	-	-	-
2000.....	-	2	1	-	-	1	-	1	-	-	-	-
1999.....	-	-	1	2	1	2	1	1	1	-	-	-
American Indian 1/, female												
2004.....	-	3	2	1	1	1	-	1	-	-	-	-
2003.....	1	-	1	1	-	1	1	1	-	-	-	-
2002.....	-	1	-	1	2	-	1	1	1	1	-	-
2001.....	-	2	-	-	1	-	2	-	-	1	-	-
2000.....	1	1	-	2	-	3	2	1	-	-	-	-
1999.....	-	-	1	-	-	3	-	1	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

All other diseases of endocardium (I36-I38)

Asian or Pacific Islander 2/, both sexes

2004.....	1	2	7	7	6	5	10	7	5	2	-	-
2003.....	1	5	3	8	11	11	6	9	9	2	1	-
2002.....	2	5	3	4	9	2	11	7	2	3	-	-
2001.....	2	5	3	5	10	5	6	7	3	2	-	-
2000.....	2	5	1	3	6	8	7	4	6	-	1	-
1999.....	2	2	4	3	3	5	2	13	3	-	1	-

Asian or Pacific Islander 2/, male

2004.....	-	2	1	4	-	3	3	3	3	1	-	-
2003.....	-	1	2	1	5	2	4	2	5	1	1	-
2002.....	-	2	1	1	5	2	5	3	2	2	-	-
2001.....	1	1	1	2	3	2	1	4	2	-	-	-
2000.....	1	2	1	1	3	2	5	1	2	-	1	-
1999.....	-	2	1	2	-	1	2	8	-	-	-	-

Asian or Pacific Islander 2/, female

2004.....	1	-	6	3	6	2	7	4	2	1	-	-
2003.....	1	4	1	7	6	9	2	7	4	1	-	-
2002.....	2	3	2	3	4	-	6	4	-	1	-	-
2001.....	1	4	2	3	7	3	5	3	1	2	-	-
2000.....	1	3	-	2	3	6	2	3	4	-	-	-
1999.....	2	-	3	1	3	4	-	5	3	-	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiomyopathy (I42)												
All races, both sexes												
2004.....	1,353	1,486	1,638	1,826	2,383	3,247	3,833	3,327	2,058	691	123	2
2003.....	1,335	1,530	1,681	2,031	2,639	3,405	3,924	3,524	2,140	712	119	1
2002.....	1,347	1,575	1,602	2,020	2,919	3,670	4,106	3,564	2,063	733	163	1
2001.....	1,348	1,506	1,611	2,183	2,803	3,633	3,955	3,382	1,916	679	112	3
2000.....	1,411	1,456	1,704	2,298	2,981	3,822	3,767	3,308	1,957	652	114	-
1999.....	1,337	1,532	1,795	2,443	3,151	3,673	3,716	3,240	1,803	612	126	2
All races, male												
2004.....	922	1,002	1,112	1,177	1,488	1,933	2,042	1,600	808	200	24	2
2003.....	927	1,036	1,127	1,295	1,641	2,029	2,148	1,626	773	171	14	1
2002.....	955	1,069	1,070	1,285	1,849	2,206	2,227	1,683	727	199	28	-
2001.....	902	1,013	1,086	1,367	1,737	2,136	2,100	1,501	690	189	19	3
2000.....	992	1,011	1,111	1,456	1,868	2,264	2,009	1,527	652	198	19	-
1999.....	935	1,063	1,166	1,590	1,974	2,134	1,924	1,441	635	151	19	1
All races, female												
2004.....	431	484	526	649	895	1,314	1,791	1,727	1,250	491	99	-
2003.....	408	494	554	736	998	1,376	1,776	1,898	1,367	541	105	-
2002.....	392	506	532	735	1,070	1,464	1,879	1,881	1,336	534	135	1
2001.....	446	493	525	816	1,066	1,497	1,855	1,881	1,226	490	93	-
2000.....	419	445	593	842	1,113	1,558	1,758	1,781	1,305	454	95	-
1999.....	402	469	629	853	1,177	1,539	1,792	1,799	1,168	461	107	1
White, both sexes												
2004.....	913	1,072	1,216	1,393	1,937	2,773	3,431	3,036	1,901	622	106	1
2003.....	885	1,089	1,231	1,577	2,164	2,938	3,472	3,232	1,964	650	105	1
2002.....	919	1,136	1,209	1,605	2,421	3,154	3,670	3,220	1,863	655	147	1
2001.....	914	1,074	1,206	1,725	2,342	3,114	3,503	3,062	1,747	639	100	3
2000.....	950	1,038	1,294	1,785	2,478	3,266	3,348	2,995	1,794	576	98	-
1999.....	906	1,081	1,305	1,944	2,606	3,165	3,295	2,909	1,649	555	108	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiomyopathy (I42)												
White, male												
2004.....	652	725	836	910	1,246	1,687	1,853	1,479	748	182	18	1
2003.....	631	750	845	1,013	1,372	1,784	1,923	1,504	713	156	13	1
2002.....	661	787	829	1,030	1,552	1,938	2,018	1,540	655	180	25	-
2001.....	638	737	832	1,098	1,478	1,863	1,877	1,367	635	180	17	3
2000.....	678	730	861	1,140	1,583	1,957	1,808	1,397	609	166	17	-
1999.....	643	756	857	1,285	1,647	1,869	1,730	1,296	583	133	17	1
White, female												
2004.....	261	347	380	483	691	1,086	1,578	1,557	1,153	440	88	-
2003.....	254	339	386	564	792	1,154	1,549	1,728	1,251	494	92	-
2002.....	258	349	380	575	869	1,216	1,652	1,680	1,208	475	122	1
2001.....	276	337	374	627	864	1,251	1,626	1,695	1,112	459	83	-
2000.....	272	308	433	645	895	1,309	1,540	1,598	1,185	410	81	-
1999.....	263	325	448	659	959	1,296	1,565	1,613	1,066	422	91	-
Black, both sexes												
2004.....	406	387	387	387	391	408	345	237	136	53	15	1
2003.....	410	406	404	405	416	403	392	254	145	54	12	-
2002.....	386	402	350	372	429	450	368	284	175	68	15	-
2001.....	398	395	372	414	405	466	386	273	144	35	8	-
2000.....	425	386	353	463	447	481	371	275	132	62	14	-
1999.....	401	406	434	447	475	443	359	284	133	50	13	1
Black, male												
2004.....	248	256	249	240	214	211	157	97	47	10	5	1
2003.....	268	259	254	255	236	207	191	106	45	13	1	-
2002.....	264	259	212	224	256	226	171	113	62	13	3	-
2001.....	246	256	234	246	218	243	191	109	41	7	2	-
2000.....	285	260	215	283	246	261	177	110	32	24	2	-
1999.....	273	278	274	272	284	221	161	121	44	17	2	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiomyopathy (I42)												
Black, female												
2004.....	158	131	138	147	177	197	188	140	89	43	10	-
2003.....	142	147	150	150	180	196	201	148	100	41	11	-
2002.....	122	143	138	148	173	224	197	171	113	55	12	-
2001.....	152	139	138	168	187	223	195	164	103	28	6	-
2000.....	140	126	138	180	201	220	194	165	100	38	12	-
1999.....	128	128	160	175	191	222	198	163	89	33	11	1
American Indian 1/, both sexes												
2004.....	15	7	8	11	15	9	9	6	3	3	-	-
2003.....	19	11	20	12	16	13	10	3	3	1	-	-
2002.....	12	10	8	4	16	11	12	12	3	-	-	-
2001.....	4	9	8	10	11	7	15	9	8	1	-	-
2000.....	9	7	15	13	11	14	9	9	3	1	1	-
1999.....	9	12	14	11	18	13	9	7	3	2	-	-
American Indian 1/, male												
2004.....	9	5	4	6	9	6	2	4	-	-	-	-
2003.....	13	10	10	5	9	6	5	1	2	-	-	-
2002.....	9	7	5	3	12	7	7	7	-	-	-	-
2001.....	2	4	4	7	9	5	5	6	4	-	-	-
2000.....	8	6	9	8	8	7	6	6	-	1	-	-
1999.....	5	6	8	6	10	11	5	2	1	-	-	-
American Indian 1/, female												
2004.....	6	2	4	5	6	3	7	2	3	3	-	-
2003.....	6	1	10	7	7	7	5	2	1	1	-	-
2002.....	3	3	3	1	4	4	5	5	3	-	-	-
2001.....	2	5	4	3	2	2	10	3	4	1	-	-
2000.....	1	1	6	5	3	7	3	3	3	-	1	-
1999.....	4	6	6	5	8	2	4	5	2	2	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiomyopathy (I42)												
Asian or Pacific Islander 2/, both sexes												
2004.....	19	20	27	35	40	57	48	48	18	13	2	-
2003.....	21	24	26	37	43	51	50	35	28	7	2	-
2002.....	30	27	35	39	53	55	56	48	22	10	1	-
2001.....	32	28	25	34	45	46	51	38	17	4	4	-
2000.....	27	25	42	37	45	61	39	29	28	13	1	-
1999.....	21	33	42	41	52	52	53	40	18	5	5	-
Asian or Pacific Islander 2/, male												
2004.....	13	16	23	21	19	29	30	20	13	8	1	-
2003.....	15	17	18	22	24	32	29	15	13	2	-	-
2002.....	21	16	24	28	29	35	31	23	10	6	-	-
2001.....	16	16	16	16	32	25	27	19	10	2	-	-
2000.....	21	15	26	25	31	39	18	14	11	7	-	-
1999.....	14	23	27	27	33	33	28	22	7	1	-	-
Asian or Pacific Islander 2/, female												
2004.....	6	4	4	14	21	28	18	28	5	5	1	-
2003.....	6	7	8	15	19	19	21	20	15	5	2	-
2002.....	9	11	11	11	24	20	25	25	12	4	1	-
2001.....	16	12	9	18	13	21	24	19	7	2	4	-
2000.....	6	10	16	12	14	22	21	15	17	6	1	-
1999.....	7	10	15	14	19	19	25	18	11	4	5	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Conduction disorders and cardiac dysrhythmias (I44-I49)												
All races, both sexes												
2004.....	1,203	1,507	1,961	2,228	3,092	4,604	6,362	6,687	5,107	2,098	460	1
2003.....	1,169	1,616	1,870	2,366	3,382	4,968	6,611	6,879	5,121	2,229	559	2
2002.....	1,210	1,510	1,832	2,362	3,394	5,201	6,698	6,909	5,312	2,119	507	2
2001.....	1,168	1,414	1,770	2,413	3,560	5,147	6,474	6,715	5,044	2,160	535	1
2000.....	1,038	1,420	1,749	2,486	3,634	5,101	6,296	6,651	5,184	2,061	525	2
1999.....	1,099	1,412	2,039	2,605	3,975	5,317	6,559	6,786	5,150	2,176	579	1
All races, male												
2004.....	777	991	1,271	1,341	1,712	2,225	2,672	2,299	1,384	444	73	1
2003.....	781	1,057	1,176	1,427	1,921	2,394	2,755	2,317	1,371	465	76	2
2002.....	798	978	1,169	1,381	1,856	2,496	2,778	2,402	1,408	431	74	2
2001.....	767	900	1,104	1,367	1,932	2,514	2,717	2,252	1,293	424	79	1
2000.....	707	930	1,113	1,520	2,041	2,453	2,521	2,240	1,339	417	92	1
1999.....	718	928	1,264	1,618	2,147	2,557	2,779	2,292	1,371	476	78	-
All races, female												
2004.....	426	516	690	887	1,380	2,379	3,690	4,388	3,723	1,654	387	-
2003.....	388	559	694	939	1,461	2,574	3,856	4,562	3,750	1,764	483	-
2002.....	412	532	663	981	1,538	2,705	3,920	4,507	3,904	1,688	433	-
2001.....	401	514	666	1,046	1,628	2,633	3,757	4,463	3,751	1,736	456	-
2000.....	331	490	636	966	1,593	2,648	3,775	4,411	3,845	1,644	433	1
1999.....	381	484	775	987	1,828	2,760	3,780	4,494	3,779	1,700	501	1
White, both sexes												
2004.....	902	1,119	1,573	1,806	2,557	3,926	5,667	6,034	4,601	1,894	396	1
2003.....	857	1,260	1,486	1,861	2,806	4,307	5,889	6,239	4,658	2,006	484	2
2002.....	898	1,172	1,406	1,927	2,831	4,494	5,966	6,294	4,840	1,907	428	1
2001.....	869	1,076	1,402	1,947	2,985	4,438	5,760	6,105	4,622	1,945	468	1
2000.....	797	1,130	1,364	2,041	3,082	4,434	5,693	6,043	4,744	1,864	458	1
1999.....	852	1,100	1,644	2,160	3,388	4,617	5,901	6,176	4,757	1,973	497	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Conduction disorders and cardiac dysrhythmias (I44-I49)												
White, male												
2004.....	607	763	1,042	1,094	1,429	1,911	2,380	2,067	1,214	388	57	1
2003.....	589	841	950	1,144	1,611	2,078	2,463	2,102	1,225	418	62	2
2002.....	608	775	922	1,152	1,567	2,157	2,495	2,187	1,262	379	56	1
2001.....	589	690	887	1,141	1,626	2,167	2,410	2,050	1,165	368	65	1
2000.....	547	751	887	1,263	1,756	2,154	2,312	2,025	1,215	371	79	1
1999.....	570	725	1,042	1,357	1,857	2,212	2,506	2,068	1,239	417	63	-
White, female												
2004.....	295	356	531	712	1,128	2,015	3,287	3,967	3,387	1,506	339	-
2003.....	268	419	536	717	1,195	2,229	3,426	4,137	3,433	1,588	422	-
2002.....	290	397	484	775	1,264	2,337	3,471	4,107	3,578	1,528	372	-
2001.....	280	386	515	806	1,359	2,271	3,350	4,055	3,457	1,577	403	-
2000.....	250	379	477	778	1,326	2,280	3,381	4,018	3,529	1,493	379	-
1999.....	282	375	602	803	1,531	2,405	3,395	4,108	3,518	1,556	434	1
Black, both sexes												
2004.....	277	346	347	363	472	567	568	511	392	164	52	-
2003.....	287	309	341	443	500	558	572	514	355	190	63	-
2002.....	293	303	386	392	476	577	614	499	393	181	64	1
2001.....	274	299	333	414	488	606	600	518	358	172	60	-
2000.....	221	261	354	393	483	594	521	511	373	165	58	1
1999.....	226	285	360	404	522	613	589	534	346	173	73	-
Black, male												
2004.....	158	203	208	206	251	259	232	172	115	38	10	-
2003.....	174	186	198	245	269	265	218	154	96	35	10	-
2002.....	177	178	222	206	247	262	227	159	105	37	13	1
2001.....	163	182	199	199	256	292	247	170	92	41	12	-
2000.....	147	159	207	226	238	257	181	171	97	35	10	-
1999.....	134	183	205	235	253	285	247	182	108	47	14	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Black, female												
2004.....	119	143	139	157	221	308	336	339	277	126	42	-
2003.....	113	123	143	198	231	293	354	360	259	155	53	-
2002.....	116	125	164	186	229	315	387	340	288	144	51	-
2001.....	111	117	134	215	232	314	353	348	266	131	48	-
2000.....	74	102	147	167	245	337	340	340	276	130	48	1
1999.....	92	102	155	169	269	328	342	352	238	126	59	-
American Indian 1/, both sexes												
2004.....	6	13	10	12	8	9	13	13	11	5	1	-
2003.....	5	16	16	14	10	11	17	18	11	6	2	-
2002.....	4	8	11	6	10	21	13	11	7	4	4	-
2001.....	5	12	13	9	13	22	19	17	6	3	1	-
2000.....	9	10	7	15	17	11	13	15	14	5	2	-
1999.....	9	12	16	12	12	15	11	10	10	2	2	-
American Indian 1/, male												
2004.....	1	9	6	7	3	3	2	6	4	-	1	-
2003.....	3	9	8	7	7	6	6	8	7	1	1	-
2002.....	3	5	8	3	7	13	4	2	3	3	3	-
2001.....	3	9	7	3	8	12	13	5	3	-	-	-
2000.....	7	9	4	9	13	7	2	4	3	-	-	-
1999.....	6	10	9	9	9	8	2	5	4	2	1	-
American Indian 1/, female												
2004.....	5	4	4	5	5	6	11	7	7	5	-	-
2003.....	2	7	8	7	3	5	11	10	4	5	1	-
2002.....	1	3	3	3	3	8	9	9	4	1	1	-
2001.....	2	3	6	6	5	10	6	12	3	3	1	-
2000.....	2	1	3	6	4	4	11	11	11	5	2	-
1999.....	3	2	7	3	3	7	9	5	6	-	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Asian or Pacific Islander 2/, both sexes												
2004.....	18	29	31	47	55	102	114	129	103	35	11	-
2003.....	20	31	27	48	66	92	133	108	97	27	10	-
2002.....	15	27	29	37	77	109	105	105	72	27	11	-
2001.....	20	27	22	43	74	81	95	75	58	40	6	-
2000.....	11	19	24	37	52	62	69	82	53	27	7	-
1999.....	12	15	19	29	53	72	58	66	37	28	7	-
Asian or Pacific Islander 2/, male												
2004.....	11	16	15	34	29	52	58	54	51	18	5	-
2003.....	15	21	20	31	34	45	68	53	43	11	3	-
2002.....	10	20	17	20	35	64	52	54	38	12	2	-
2001.....	12	19	11	24	42	43	47	27	33	15	2	-
2000.....	6	11	15	22	34	35	26	40	24	11	3	-
1999.....	8	10	8	17	28	52	24	37	20	10	-	-
Asian or Pacific Islander 2/, female												
2004.....	7	13	16	13	26	50	56	75	52	17	6	-
2003.....	5	10	7	17	32	47	65	55	54	16	7	-
2002.....	5	7	12	17	42	45	53	51	34	15	9	-
2001.....	8	8	11	19	32	38	48	48	25	25	4	-
2000.....	5	8	9	15	18	27	43	42	29	16	4	-
1999.....	4	5	11	12	25	20	34	29	17	18	7	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Heart failure (I50)												
All races, both sexes												
2004.....	653	1,003	1,441	2,226	3,725	6,574	10,438	12,876	11,063	5,044	1,203	1
2003.....	623	1,007	1,462	2,302	3,836	6,697	10,439	12,844	11,082	5,071	1,222	-
2002.....	638	906	1,436	2,192	3,976	6,603	10,307	12,620	10,776	5,031	1,200	-
2001.....	601	926	1,448	2,270	4,004	6,879	10,255	12,793	10,735	4,971	1,209	1
2000.....	528	828	1,383	2,284	4,119	6,936	9,884	12,315	10,543	4,969	1,208	4
1999.....	484	853	1,254	2,240	4,284	6,926	9,845	12,169	10,197	4,850	1,111	1
All races, male												
2004.....	378	599	821	1,223	1,940	3,198	4,475	4,707	3,202	1,049	172	-
2003.....	364	605	849	1,281	2,032	3,262	4,396	4,668	3,115	1,111	206	-
2002.....	407	526	819	1,200	2,054	3,192	4,381	4,423	3,000	1,035	178	-
2001.....	361	556	791	1,235	2,109	3,300	4,372	4,320	2,886	1,020	172	-
2000.....	305	479	785	1,243	2,171	3,286	4,212	4,256	2,819	982	191	2
1999.....	300	506	722	1,232	2,294	3,264	4,054	4,155	2,636	991	177	1
All races, female												
2004.....	275	404	620	1,003	1,785	3,376	5,963	8,169	7,861	3,995	1,031	1
2003.....	259	402	613	1,021	1,804	3,435	6,043	8,176	7,967	3,960	1,016	-
2002.....	231	380	617	992	1,922	3,411	5,926	8,197	7,776	3,996	1,022	-
2001.....	240	370	657	1,035	1,895	3,579	5,883	8,473	7,849	3,951	1,037	1
2000.....	223	349	598	1,041	1,948	3,650	5,672	8,059	7,724	3,987	1,017	2
1999.....	184	347	532	1,008	1,990	3,662	5,791	8,014	7,561	3,859	934	-
White, both sexes												
2004.....	408	711	1,115	1,834	3,179	5,773	9,538	11,963	10,328	4,728	1,082	1
2003.....	406	750	1,109	1,873	3,245	5,926	9,516	11,935	10,361	4,723	1,094	-
2002.....	428	678	1,134	1,789	3,371	5,822	9,377	11,737	10,055	4,722	1,075	-
2001.....	418	693	1,149	1,849	3,428	6,108	9,358	11,903	10,063	4,645	1,092	1
2000.....	378	627	1,088	1,870	3,514	6,178	9,038	11,433	9,847	4,628	1,098	4
1999.....	326	626	1,006	1,827	3,684	6,102	8,979	11,241	9,583	4,490	1,019	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Heart failure (I50)												
White, male												
2004.....	248	429	634	1,012	1,694	2,822	4,125	4,387	2,993	982	152	-
2003.....	239	459	653	1,059	1,761	2,943	4,033	4,358	2,939	1,033	174	-
2002.....	280	402	652	984	1,762	2,852	4,034	4,138	2,800	977	156	-
2001.....	259	429	647	1,021	1,828	2,961	4,001	4,038	2,703	947	151	-
2000.....	224	383	624	1,049	1,892	2,959	3,872	3,961	2,618	905	166	2
1999.....	206	387	585	1,012	2,026	2,918	3,725	3,820	2,459	907	153	-
White, female												
2004.....	160	282	481	822	1,485	2,951	5,413	7,576	7,335	3,746	930	1
2003.....	167	291	456	814	1,484	2,983	5,483	7,577	7,422	3,690	920	-
2002.....	148	276	482	805	1,609	2,970	5,343	7,599	7,255	3,745	919	-
2001.....	159	264	502	828	1,600	3,147	5,357	7,865	7,360	3,698	941	1
2000.....	154	244	464	821	1,622	3,219	5,166	7,472	7,229	3,723	932	2
1999.....	120	239	421	815	1,658	3,184	5,254	7,421	7,124	3,583	866	-
Black, both sexes												
2004.....	233	273	291	346	489	710	786	783	632	277	104	-
2003.....	204	241	328	392	536	674	802	788	620	308	107	-
2002.....	195	207	283	363	535	690	823	772	623	274	109	-
2001.....	174	223	275	382	528	697	789	787	589	295	100	-
2000.....	143	189	266	373	544	674	742	783	608	292	102	-
1999.....	149	215	236	380	547	731	750	825	515	316	78	1
Black, male												
2004.....	125	159	164	188	214	326	307	271	165	55	16	-
2003.....	115	138	186	207	242	273	308	260	143	63	26	-
2002.....	116	111	158	191	262	298	310	227	156	47	17	-
2001.....	96	121	130	195	256	303	320	246	148	63	20	-
2000.....	78	90	143	175	248	285	288	255	167	60	22	-
1999.....	89	114	129	198	245	309	274	286	141	69	19	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Heart failure (I50)												
Black, female												
2004.....	108	114	127	158	275	384	479	512	467	222	88	-
2003.....	89	103	142	185	294	401	494	528	477	245	81	-
2002.....	79	96	125	172	273	392	513	545	467	227	92	-
2001.....	78	102	145	187	272	394	469	541	441	232	80	-
2000.....	65	99	123	198	296	389	454	528	441	232	80	-
1999.....	60	101	107	182	302	422	476	539	374	247	59	-
American Indian 1/, both sexes												
2004.....	7	4	17	22	12	32	24	32	17	11	5	-
2003.....	6	9	15	14	16	24	32	25	20	4	8	-
2002.....	9	9	7	19	20	28	24	27	19	9	5	-
2001.....	5	5	7	13	20	22	29	29	14	9	5	-
2000.....	5	4	10	17	25	29	29	28	14	8	2	-
1999.....	5	2	5	17	17	31	27	18	24	11	6	-
American Indian 1/, male												
2004.....	2	3	10	11	7	15	7	15	5	4	1	-
2003.....	4	5	8	5	9	6	12	12	6	1	2	-
2002.....	7	6	5	12	10	14	8	15	4	2	2	-
2001.....	4	4	4	6	11	13	13	6	4	3	-	-
2000.....	3	1	5	8	15	16	13	12	2	1	-	-
1999.....	2	-	3	12	7	12	9	5	5	3	3	-
American Indian 1/, female												
2004.....	5	1	7	11	5	17	17	17	12	7	4	-
2003.....	2	4	7	9	7	18	20	13	14	3	6	-
2002.....	2	3	2	7	10	14	16	12	15	7	3	-
2001.....	1	1	3	7	9	9	16	23	10	6	5	-
2000.....	2	3	5	9	10	13	16	16	12	7	2	-
1999.....	3	2	2	5	10	19	18	13	19	8	3	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Heart failure (I50)												
Asian or Pacific Islander 2/, both sexes												
2004.....	5	15	18	24	45	59	90	98	86	28	12	-
2003.....	7	7	10	23	39	73	89	96	81	36	13	-
2002.....	6	12	12	21	50	63	83	84	79	26	11	-
2001.....	4	5	17	26	28	52	79	74	69	22	12	-
2000.....	2	8	19	24	36	55	75	71	74	41	6	-
1999.....	4	10	7	16	36	62	89	85	75	33	8	-
Asian or Pacific Islander 2/, male												
2004.....	3	8	13	12	25	35	36	34	39	8	3	-
2003.....	6	3	2	10	20	40	43	38	27	14	4	-
2002.....	4	7	4	13	20	28	29	43	40	9	3	-
2001.....	2	2	10	13	14	23	38	30	31	7	1	-
2000.....	-	5	13	11	16	26	39	28	32	16	3	-
1999.....	3	5	5	10	16	25	46	44	31	12	2	-
Asian or Pacific Islander 2/, female												
2004.....	2	7	5	12	20	24	54	64	47	20	9	-
2003.....	1	4	8	13	19	33	46	58	54	22	9	-
2002.....	2	5	8	8	30	35	54	41	39	17	8	-
2001.....	2	3	7	13	14	29	41	44	38	15	11	-
2000.....	2	3	6	13	20	29	36	43	42	25	3	-
1999.....	1	5	2	6	20	37	43	41	44	21	6	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Congestive heart failure (I50.0)												
All races, both sexes												
2004.....	553	846	1,245	1,968	3,401	6,085	9,777	12,194	10,517	4,786	1,121	1
2003.....	525	861	1,280	2,070	3,491	6,177	9,833	12,165	10,541	4,825	1,138	-
2002.....	522	769	1,242	1,942	3,599	6,104	9,649	11,861	10,246	4,757	1,107	-
2001.....	512	766	1,259	2,045	3,646	6,372	9,545	12,065	10,158	4,681	1,108	1
2000.....	439	722	1,199	1,997	3,703	6,386	9,244	11,572	9,943	4,688	1,117	2
1999.....	395	694	1,072	1,996	3,873	6,362	9,198	11,452	9,639	4,566	1,040	1
All races, male												
2004.....	314	500	717	1,067	1,763	2,961	4,158	4,426	3,038	994	164	-
2003.....	300	514	729	1,153	1,840	3,002	4,124	4,403	2,956	1,042	191	-
2002.....	319	444	693	1,057	1,861	2,936	4,083	4,140	2,836	987	162	-
2001.....	300	456	678	1,098	1,907	3,019	4,022	4,067	2,733	954	161	-
2000.....	249	418	666	1,086	1,934	3,018	3,929	3,999	2,657	920	181	1
1999.....	242	408	602	1,077	2,072	2,987	3,774	3,893	2,503	937	163	1
All races, female												
2004.....	239	346	528	901	1,638	3,124	5,619	7,768	7,479	3,792	957	1
2003.....	225	347	551	917	1,651	3,175	5,709	7,762	7,585	3,783	947	-
2002.....	203	325	549	885	1,738	3,168	5,566	7,721	7,410	3,770	945	-
2001.....	212	310	581	947	1,739	3,353	5,523	7,998	7,425	3,727	947	1
2000.....	190	304	533	911	1,769	3,368	5,315	7,573	7,286	3,768	936	1
1999.....	153	286	470	919	1,801	3,375	5,424	7,559	7,136	3,629	877	-
White, both sexes												
2004.....	352	597	969	1,632	2,911	5,355	8,963	11,339	9,842	4,493	1,014	1
2003.....	353	646	970	1,698	2,964	5,480	8,987	11,324	9,890	4,511	1,027	-
2002.....	350	574	971	1,589	3,066	5,399	8,802	11,048	9,589	4,473	999	-
2001.....	359	570	1,002	1,660	3,132	5,683	8,732	11,243	9,536	4,377	1,006	1
2000.....	308	550	944	1,631	3,178	5,699	8,478	10,770	9,299	4,375	1,020	2
1999.....	258	508	860	1,630	3,332	5,630	8,399	10,606	9,068	4,227	957	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Congestive heart failure (I50.0)												
White, male												
2004.....	211	355	551	888	1,546	2,617	3,848	4,128	2,849	932	145	-
2003.....	204	391	561	961	1,596	2,718	3,795	4,118	2,798	972	164	-
2002.....	220	337	546	867	1,594	2,630	3,775	3,885	2,652	935	146	-
2001.....	214	354	554	906	1,660	2,726	3,696	3,807	2,564	885	142	-
2000.....	178	341	526	924	1,695	2,727	3,624	3,731	2,474	849	158	1
1999.....	163	313	486	886	1,831	2,678	3,468	3,590	2,334	860	142	-
White, female												
2004.....	141	242	418	744	1,365	2,738	5,115	7,211	6,993	3,561	869	1
2003.....	149	255	409	737	1,368	2,762	5,192	7,206	7,092	3,539	863	-
2002.....	130	237	425	722	1,472	2,769	5,027	7,163	6,937	3,538	853	-
2001.....	145	216	448	754	1,472	2,957	5,036	7,436	6,972	3,492	864	1
2000.....	130	209	418	707	1,483	2,972	4,854	7,039	6,825	3,526	862	1
1999.....	95	195	374	744	1,501	2,952	4,931	7,016	6,734	3,367	815	-
Black, both sexes												
2004.....	189	232	246	297	438	647	706	734	580	258	90	-
2003.....	164	203	288	341	480	605	732	729	558	278	95	-
2002.....	161	177	252	324	476	621	749	713	567	250	95	-
2001.....	146	188	234	348	469	625	717	726	547	275	88	-
2000.....	125	160	233	330	472	610	674	709	564	269	89	-
1999.....	130	178	201	337	492	652	693	752	476	300	70	1
Black, male												
2004.....	98	136	143	159	189	298	271	253	151	51	15	-
2003.....	90	117	159	181	218	242	276	239	128	57	23	-
2002.....	92	96	138	172	244	267	274	202	144	41	13	-
2001.....	81	97	111	174	224	265	280	226	137	60	18	-
2000.....	69	71	126	147	213	254	257	230	153	55	20	-
1999.....	75	91	109	173	220	277	257	259	134	64	16	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Congestive heart failure (I50.0)												
Black, female												
2004.....	91	96	103	138	249	349	435	481	429	207	75	-
2003.....	74	86	129	160	262	363	456	490	430	221	72	-
2002.....	69	81	114	152	232	354	475	511	423	209	82	-
2001.....	65	91	123	174	245	360	437	500	410	215	70	-
2000.....	56	89	107	183	259	356	417	479	411	214	69	-
1999.....	55	87	92	164	272	375	436	493	342	236	54	-
American Indian 1/, both sexes												
2004.....	7	4	14	18	11	28	23	32	16	9	5	-
2003.....	4	7	13	13	10	23	31	21	18	4	6	-
2002.....	8	7	7	14	14	27	23	24	18	9	3	-
2001.....	3	4	7	12	18	15	24	28	12	8	4	-
2000.....	4	4	8	15	22	26	24	25	12	8	2	-
1999.....	3	1	4	16	16	25	24	16	22	11	5	-
American Indian 1/, male												
2004.....	2	3	10	9	6	13	7	15	4	3	1	-
2003.....	3	4	7	4	6	5	12	9	5	1	1	-
2002.....	6	4	5	8	7	14	8	15	4	2	1	-
2001.....	3	3	4	6	9	7	11	6	4	2	-	-
2000.....	2	1	4	6	13	13	13	10	2	1	-	-
1999.....	1	-	2	11	6	9	7	5	4	3	3	-
American Indian 1/, female												
2004.....	5	1	4	9	5	15	16	17	12	6	4	-
2003.....	1	3	6	9	4	18	19	12	13	3	5	-
2002.....	2	3	2	6	7	13	15	9	14	7	2	-
2001.....	-	1	3	6	9	8	13	22	8	6	4	-
2000.....	2	3	4	9	9	13	11	15	10	7	2	-
1999.....	2	1	2	5	10	16	17	11	18	8	2	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Congestive heart failure (I50.0)												
Asian or Pacific Islander 2/, both sexes												
2004.....	5	13	16	21	41	55	85	89	79	26	12	-
2003.....	4	5	9	18	37	69	83	91	75	32	10	-
2002.....	3	11	12	15	43	57	75	76	72	25	10	-
2001.....	4	4	16	25	27	49	72	68	63	21	10	-
2000.....	2	8	14	21	31	51	68	68	68	36	6	-
1999.....	4	7	7	13	33	55	82	78	73	28	8	-
Asian or Pacific Islander 2/, male												
2004.....	3	6	13	11	22	33	32	30	34	8	3	-
2003.....	3	2	2	7	20	37	41	37	25	12	3	-
2002.....	1	7	4	10	16	25	26	38	36	9	2	-
2001.....	2	2	9	12	14	21	35	28	28	7	1	-
2000.....	-	5	10	9	13	24	35	28	28	15	3	-
1999.....	3	4	5	7	15	23	42	39	31	10	2	-
Asian or Pacific Islander 2/, female												
2004.....	2	7	3	10	19	22	53	59	45	18	9	-
2003.....	1	3	7	11	17	32	42	54	50	20	7	-
2002.....	2	4	8	5	27	32	49	38	36	16	8	-
2001.....	2	2	7	13	13	28	37	40	35	14	9	-
2000.....	2	3	4	12	18	27	33	40	40	21	3	-
1999.....	1	3	2	6	18	32	40	39	42	18	6	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified heart failure (I50.1-I50.9)												
All races, both sexes												
2004.....	100	157	196	258	324	489	661	682	546	258	82	-
2003.....	98	146	182	232	345	520	606	679	541	246	84	-
2002.....	116	137	194	250	377	499	658	759	530	274	93	-
2001.....	89	160	189	225	358	507	710	728	577	290	101	-
2000.....	89	106	184	287	416	550	640	743	600	281	91	2
1999.....	89	159	182	244	411	564	647	717	558	284	71	-
All races, male												
2004.....	64	99	104	156	177	237	317	281	164	55	8	-
2003.....	64	91	120	128	192	260	272	265	159	69	15	-
2002.....	88	82	126	143	193	256	298	283	164	48	16	-
2001.....	61	100	113	137	202	281	350	253	153	66	11	-
2000.....	56	61	119	157	237	268	283	257	162	62	10	1
1999.....	58	98	120	155	222	277	280	262	133	54	14	-
All races, female												
2004.....	36	58	92	102	147	252	344	401	382	203	74	-
2003.....	34	55	62	104	153	260	334	414	382	177	69	-
2002.....	28	55	68	107	184	243	360	476	366	226	77	-
2001.....	28	60	76	88	156	226	360	475	424	224	90	-
2000.....	33	45	65	130	179	282	357	486	438	219	81	1
1999.....	31	61	62	89	189	287	367	455	425	230	57	-
White, both sexes												
2004.....	56	114	146	202	268	418	575	624	486	235	68	-
2003.....	53	104	139	175	281	446	529	611	471	212	67	-
2002.....	78	104	163	200	305	423	575	689	466	249	76	-
2001.....	59	123	147	189	296	425	626	660	527	268	86	-
2000.....	70	77	144	239	336	479	560	663	548	253	78	2
1999.....	68	118	146	197	352	472	580	635	515	263	62	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified heart failure (I50.1-I50.9)												
White, male												
2004.....	37	74	83	124	148	205	277	259	144	50	7	-
2003.....	35	68	92	98	165	225	238	240	141	61	10	-
2002.....	60	65	106	117	168	222	259	253	148	42	10	-
2001.....	45	75	93	115	168	235	305	231	139	62	9	-
2000.....	46	42	98	125	197	232	248	230	144	56	8	1
1999.....	43	74	99	126	195	240	257	230	125	47	11	-
White, female												
2004.....	19	40	63	78	120	213	298	365	342	185	61	-
2003.....	18	36	47	77	116	221	291	371	330	151	57	-
2002.....	18	39	57	83	137	201	316	436	318	207	66	-
2001.....	14	48	54	74	128	190	321	429	388	206	77	-
2000.....	24	35	46	114	139	247	312	433	404	197	70	1
1999.....	25	44	47	71	157	232	323	405	390	216	51	-
Black, both sexes												
2004.....	44	41	45	49	51	63	80	49	52	19	14	-
2003.....	40	38	40	51	56	69	70	59	62	30	12	-
2002.....	34	30	31	39	59	69	74	59	56	24	14	-
2001.....	28	35	41	34	59	72	72	61	42	20	12	-
2000.....	18	29	33	43	72	64	68	74	44	23	13	-
1999.....	19	37	35	43	55	79	57	73	39	16	8	-
Black, male												
2004.....	27	23	21	29	25	28	36	18	14	4	1	-
2003.....	25	21	27	26	24	31	32	21	15	6	3	-
2002.....	24	15	20	19	18	31	36	25	12	6	4	-
2001.....	15	24	19	21	32	38	40	20	11	3	2	-
2000.....	9	19	17	28	35	31	31	25	14	5	2	-
1999.....	14	23	20	25	25	32	17	27	7	5	3	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified heart failure (I50.1-I50.9)												
Black, female												
2004.....	17	18	24	20	26	35	44	31	38	15	13	-
2003.....	15	17	13	25	32	38	38	38	47	24	9	-
2002.....	10	15	11	20	41	38	38	34	44	18	10	-
2001.....	13	11	22	13	27	34	32	41	31	17	10	-
2000.....	9	10	16	15	37	33	37	49	30	18	11	-
1999.....	5	14	15	18	30	47	40	46	32	11	5	-
American Indian 1/, both sexes												
2004.....	-	-	3	4	1	4	1	-	1	2	-	-
2003.....	2	2	2	1	6	1	1	4	2	-	2	-
2002.....	1	2	-	5	6	1	1	3	1	-	2	-
2001.....	2	1	-	1	2	7	5	1	2	1	1	-
2000.....	1	-	2	2	3	3	5	3	2	-	-	-
1999.....	2	1	1	1	1	6	3	2	2	-	1	-
American Indian 1/, male												
2004.....	-	-	-	2	1	2	-	-	1	1	-	-
2003.....	1	1	1	1	3	1	-	3	1	-	1	-
2002.....	1	2	-	4	3	-	-	-	-	-	1	-
2001.....	1	1	-	-	2	6	2	-	-	1	-	-
2000.....	1	-	1	2	2	3	-	2	-	-	-	-
1999.....	1	-	1	1	1	3	2	-	1	-	-	-
American Indian 1/, female												
2004.....	-	-	3	2	-	2	1	-	-	1	-	-
2003.....	1	1	1	-	3	-	1	1	1	-	1	-
2002.....	-	-	-	1	3	1	1	3	1	-	1	-
2001.....	1	-	-	1	-	1	3	1	2	-	1	-
2000.....	-	-	1	-	1	-	5	1	2	-	-	-
1999.....	1	1	-	-	-	3	1	2	1	-	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified heart failure (I50.1-I50.9)												
Asian or Pacific Islander 2/, both sexes												
2004.....	-	2	2	3	4	4	5	9	7	2	-	-
2003.....	3	2	1	5	2	4	6	5	6	4	3	-
2002.....	3	1	-	6	7	6	8	8	7	1	1	-
2001.....	-	1	1	1	1	3	7	6	6	1	2	-
2000.....	-	-	5	3	5	4	7	3	6	5	-	-
1999.....	-	3	-	3	3	7	7	7	2	5	-	-
Asian or Pacific Islander 2/, male												
2004.....	-	2	-	1	3	2	4	4	5	-	-	-
2003.....	3	1	-	3	-	3	2	1	2	2	1	-
2002.....	3	-	-	3	4	3	3	5	4	-	1	-
2001.....	-	-	1	1	-	2	3	2	3	-	-	-
2000.....	-	-	3	2	3	2	4	-	4	1	-	-
1999.....	-	1	-	3	1	2	4	5	-	2	-	-
Asian or Pacific Islander 2/, female												
2004.....	-	-	2	2	1	2	1	5	2	2	-	-
2003.....	-	1	1	2	2	1	4	4	4	2	2	-
2002.....	-	1	-	3	3	3	5	3	3	1	-	-
2001.....	-	1	-	-	1	1	4	4	3	1	2	-
2000.....	-	-	2	1	2	2	3	3	2	4	-	-
1999.....	-	2	-	-	2	5	3	2	2	3	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)

All races, both sexes

2004.....	47	34	19	27	22	25	23	40	29	20	10	-
2003.....	44	39	26	15	12	22	29	35	42	18	7	-
2002.....	53	56	26	35	55	61	56	64	54	22	8	1
2001.....	47	25	32	34	39	37	31	32	31	17	9	1
2000.....	26	47	29	17	32	31	37	42	31	23	7	-
1999.....	45	32	21	27	29	27	36	52	22	19	7	-

All races, male

2004.....	26	16	9	12	9	12	10	14	10	2	-	-
2003.....	20	24	12	6	7	12	16	15	13	6	3	-
2002.....	28	24	11	19	32	26	20	25	15	12	1	-
2001.....	28	10	19	22	21	24	15	13	10	3	3	-
2000.....	17	27	12	8	14	14	18	20	7	5	-	-
1999.....	24	14	13	17	14	13	17	14	6	5	-	-

All races, female

2004.....	21	18	10	15	13	13	13	26	19	18	10	-
2003.....	24	15	14	9	5	10	13	20	29	12	4	-
2002.....	25	32	15	16	23	35	36	39	39	10	7	1
2001.....	19	15	13	12	18	13	16	19	21	14	6	1
2000.....	9	20	17	9	18	17	19	22	24	18	7	-
1999.....	21	18	8	10	15	14	19	38	16	14	7	-

White, both sexes

2004.....	34	22	13	23	16	23	17	36	28	18	10	-
2003.....	32	32	23	14	12	21	27	34	38	17	6	-
2002.....	39	48	23	28	49	54	52	58	50	17	6	-
2001.....	33	21	27	27	30	33	28	31	29	15	7	-
2000.....	17	41	23	16	30	28	30	40	31	22	5	-
1999.....	34	28	14	23	27	23	35	47	21	19	6	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
White, male												
2004.....	17	12	6	8	8	12	8	13	10	2	-	-
2003.....	14	18	10	5	7	12	16	15	12	6	3	-
2002.....	19	22	9	12	30	23	18	24	15	10	1	-
2001.....	22	9	15	17	16	20	14	13	9	2	2	-
2000.....	9	24	8	7	14	13	15	18	7	5	-	-
1999.....	19	13	7	15	12	11	16	11	6	5	-	-
White, female												
2004.....	17	10	7	15	8	11	9	23	18	16	10	-
2003.....	18	14	13	9	5	9	11	19	26	11	3	-
2002.....	20	26	14	16	19	31	34	34	35	7	5	-
2001.....	11	12	12	10	14	13	14	18	20	13	5	-
2000.....	8	17	15	9	16	15	15	22	24	17	5	-
1999.....	15	15	7	8	15	12	19	36	15	14	6	-
Black, both sexes												
2004.....	13	11	4	4	4	2	5	3	1	2	-	-
2003.....	10	7	2	1	-	1	2	1	4	1	1	-
2002.....	14	8	2	6	3	7	3	5	4	5	1	1
2001.....	14	4	4	4	7	4	3	1	1	2	2	1
2000.....	7	6	4	1	1	3	6	2	-	1	1	-
1999.....	4	3	7	4	2	2	1	3	1	-	1	-
Black, male												
2004.....	9	4	2	4	1	-	1	-	-	-	-	-
2003.....	5	6	1	1	-	-	-	-	1	-	-	-
2002.....	9	2	2	6	2	3	1	-	-	2	-	-
2001.....	6	1	3	3	4	4	1	-	-	1	1	-
2000.....	6	3	3	1	-	1	2	2	-	-	-	-
1999.....	3	-	6	2	2	1	1	1	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Asian or Pacific Islander 2/, both sexes												
2004.....	-	-	2	-	1	-	1	1	-	-	-	-
2003.....	2	-	1	-	-	-	-	-	-	-	-	-
2002.....	-	-	1	1	3	-	1	1	-	-	1	-
2001.....	-	-	1	1	2	-	-	-	1	-	-	-
2000.....	2	-	2	-	1	-	1	-	-	-	-	-
1999.....	5	1	-	-	-	1	-	2	-	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	-	-	1	-	-	-	1	1	-	-	-	-
2003.....	1	-	1	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	1	-	-	1	1	-	-	-	-
2001.....	-	-	1	1	1	-	-	-	1	-	-	-
2000.....	2	-	1	-	-	-	1	-	-	-	-	-
1999.....	1	1	-	-	-	-	-	2	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	-	-	1	-	1	-	-	-	-	-	-	-
2003.....	1	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	1	-	3	-	-	-	-	-	1	-
2001.....	-	-	-	-	1	-	-	-	-	-	-	-
2000.....	-	-	1	-	1	-	-	-	-	-	-	-
1999.....	4	-	-	-	-	1	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

All other and ill-defined forms of heart disease (I50.0-I51.3,I51.6-I51.9)

All races, both sexes

2004.....	473	487	548	591	731	1,118	1,526	1,557	1,155	465	113	-
2003.....	395	486	509	565	822	1,133	1,509	1,459	1,163	451	125	1
2002.....	402	416	483	585	839	1,139	1,403	1,455	1,008	438	85	-
2001.....	369	403	460	541	794	1,068	1,306	1,397	985	428	110	-
2000.....	343	418	486	612	767	1,064	1,323	1,276	916	437	108	1
1999.....	301	364	463	532	736	1,034	1,142	1,255	849	361	91	1

All races, male

2004.....	351	332	343	404	406	597	704	553	341	106	22	-
2003.....	268	327	336	341	438	598	668	565	352	111	14	1
2002.....	273	291	303	364	520	582	617	549	268	98	12	-
2001.....	253	267	306	342	477	545	555	506	243	93	21	-
2000.....	237	288	333	358	441	518	573	464	280	84	19	1
1999.....	203	251	313	343	435	550	500	443	238	71	15	1

All races, female

2004.....	122	155	205	187	325	521	822	1,004	814	359	91	-
2003.....	127	159	173	224	384	535	841	894	811	340	111	-
2002.....	129	125	180	221	319	557	786	906	740	340	73	-
2001.....	116	136	154	199	317	523	751	891	742	335	89	-
2000.....	106	130	153	254	326	546	750	812	636	353	89	-
1999.....	98	113	150	189	301	484	642	812	611	290	76	-

White, both sexes

2004.....	371	384	439	476	591	962	1,354	1,428	1,036	417	103	-
2003.....	287	390	409	462	685	1,002	1,356	1,308	1,055	404	106	-
2002.....	284	330	369	471	719	1,008	1,261	1,318	902	392	65	-
2001.....	269	307	382	444	666	907	1,166	1,248	897	384	90	-
2000.....	246	320	395	503	620	923	1,170	1,171	827	387	96	1
1999.....	227	272	358	419	634	877	1,028	1,122	784	334	73	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and ill-defined forms of heart disease (I50.0-I51.3,I51.6-I51.9)												
White, male												
2004.....	277	268	280	333	330	518	631	514	299	96	20	-
2003.....	201	266	276	282	375	534	602	515	322	95	11	-
2002.....	202	235	232	295	456	515	551	507	236	88	10	-
2001.....	199	210	260	290	402	468	497	453	224	83	18	-
2000.....	175	221	280	299	360	452	502	430	252	70	15	1
1999.....	158	192	253	277	376	477	448	394	223	68	11	1
White, female												
2004.....	94	116	159	143	261	444	723	914	737	321	83	-
2003.....	86	124	133	180	310	468	754	793	733	309	95	-
2002.....	82	95	137	176	263	493	710	811	666	304	55	-
2001.....	70	97	122	154	264	439	669	795	673	301	72	-
2000.....	71	99	115	204	260	471	668	741	575	317	81	-
1999.....	69	80	105	142	258	400	580	728	561	266	62	-
Black, both sexes												
2004.....	90	97	100	105	130	145	153	118	106	38	9	-
2003.....	101	90	89	93	126	114	134	130	89	40	17	1
2002.....	114	76	100	105	108	116	123	116	97	39	19	-
2001.....	91	87	71	87	113	137	119	135	81	38	19	-
2000.....	89	88	88	100	140	125	134	99	80	45	12	-
1999.....	68	88	98	102	98	139	93	117	58	25	16	-
Black, male												
2004.....	63	61	59	64	70	77	64	34	34	7	1	-
2003.....	64	55	57	53	57	56	60	40	23	12	2	1
2002.....	69	49	61	64	56	58	56	34	28	8	2	-
2001.....	49	54	41	43	65	62	49	47	17	8	2	-
2000.....	61	59	51	54	76	57	62	30	23	13	4	-
1999.....	43	57	55	62	56	63	42	42	14	2	4	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and ill-defined forms of heart disease (I50.0-I51.3,I51.6-I51.9)												
Black, female												
2004.....	27	36	41	41	60	68	89	84	72	31	8	-
2003.....	37	35	32	40	69	58	74	90	66	28	15	-
2002.....	45	27	39	41	52	58	67	82	69	31	17	-
2001.....	42	33	30	44	48	75	70	88	64	30	17	-
2000.....	28	29	37	46	64	68	72	69	57	32	8	-
1999.....	25	31	43	40	42	76	51	75	44	23	12	-
American Indian 1/, both sexes												
2004.....	1	2	2	5	3	2	4	3	1	1	-	-
2003.....	4	2	6	5	4	4	3	1	7	2	1	-
2002.....	1	3	4	-	5	2	5	5	3	1	-	-
2001.....	3	5	5	3	8	8	2	3	2	2	-	-
2000.....	5	3	-	3	2	6	3	1	1	1	-	-
1999.....	-	1	2	4	1	7	4	3	1	-	2	-
American Indian 1/, male												
2004.....	1	2	1	4	2	1	2	1	1	-	-	-
2003.....	1	2	1	2	2	1	1	-	3	-	1	-
2002.....	1	3	2	-	5	2	2	4	2	-	-	-
2001.....	1	2	5	3	5	6	-	2	-	-	-	-
2000.....	1	2	-	1	2	4	3	1	1	-	-	-
1999.....	-	1	1	2	1	4	2	1	-	-	-	-
American Indian 1/, female												
2004.....	-	-	1	1	1	1	2	2	-	1	-	-
2003.....	3	-	5	3	2	3	2	1	4	2	-	-
2002.....	-	-	2	-	-	-	3	1	1	1	-	-
2001.....	2	3	-	-	3	2	2	1	2	2	-	-
2000.....	4	1	-	2	-	2	-	-	-	1	-	-
1999.....	-	-	1	2	-	3	2	2	1	-	2	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and ill-defined forms of heart disease (I50.0-I51.3,I51.6-I51.9)												
Asian or Pacific Islander 2/, both sexes												
2004.....	11	4	7	5	7	9	15	8	12	9	1	-
2003.....	3	4	5	5	7	13	16	20	12	5	1	-
2002.....	3	7	10	9	7	13	14	16	6	6	1	-
2001.....	6	4	2	7	7	16	19	11	5	4	1	-
2000.....	3	7	3	6	5	10	16	5	8	4	-	-
1999.....	6	3	5	7	3	11	17	13	6	2	-	-
Asian or Pacific Islander 2/, male												
2004.....	10	1	3	3	4	1	7	4	7	3	1	-
2003.....	2	4	2	4	4	7	5	10	4	4	-	-
2002.....	1	4	8	5	3	7	8	4	2	2	-	-
2001.....	4	1	-	6	5	9	9	4	2	2	1	-
2000.....	-	6	2	4	3	5	6	3	4	1	-	-
1999.....	2	1	4	2	2	6	8	6	1	1	-	-
Asian or Pacific Islander 2/, female												
2004.....	1	3	4	2	3	8	8	4	5	6	-	-
2003.....	1	-	3	1	3	6	11	10	8	1	1	-
2002.....	2	3	2	4	4	6	6	12	4	4	1	-
2001.....	2	3	2	1	2	7	10	7	3	2	-	-
2000.....	3	1	1	2	2	5	10	2	4	3	-	-
1999.....	4	2	1	5	1	5	9	7	5	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebrovascular diseases (I60-I69)												
All races, both sexes												
2004.....	3,559	4,368	5,598	7,806	12,095	20,504	29,588	30,162	21,179	7,752	1,452	5
2003.....	3,550	4,392	5,554	7,765	12,943	21,917	30,930	32,139	22,367	8,493	1,580	2
2002.....	3,440	4,293	5,604	8,026	13,966	22,911	31,978	33,197	23,110	8,521	1,584	3
2001.....	3,403	4,093	5,515	8,170	14,428	23,772	32,050	33,465	22,650	8,398	1,553	4
2000.....	3,386	4,145	5,811	8,671	14,978	24,678	32,342	34,030	23,205	8,530	1,611	2
1999.....	3,108	4,152	5,500	8,694	15,398	25,007	32,420	33,841	23,149	8,571	1,519	2
All races, male												
2004.....	2,015	2,395	3,128	4,180	6,027	9,333	11,692	9,872	5,346	1,449	192	3
2003.....	1,961	2,394	3,084	4,138	6,530	9,993	12,241	10,471	5,634	1,575	179	1
2002.....	1,876	2,373	3,083	4,305	6,902	10,283	12,372	10,884	5,713	1,510	209	1
2001.....	1,851	2,251	3,026	4,345	7,304	10,671	12,630	10,731	5,482	1,506	211	4
2000.....	1,834	2,294	3,202	4,513	7,529	11,192	12,754	10,864	5,588	1,564	196	1
1999.....	1,714	2,256	3,049	4,522	7,698	11,349	12,739	10,818	5,569	1,467	198	1
All races, female												
2004.....	1,544	1,973	2,470	3,626	6,068	11,171	17,896	20,290	15,833	6,303	1,260	2
2003.....	1,589	1,998	2,470	3,627	6,413	11,924	18,689	21,668	16,733	6,918	1,401	1
2002.....	1,564	1,920	2,521	3,721	7,064	12,628	19,606	22,313	17,397	7,011	1,375	2
2001.....	1,552	1,842	2,489	3,825	7,124	13,101	19,420	22,734	17,168	6,892	1,342	-
2000.....	1,552	1,851	2,609	4,158	7,449	13,486	19,588	23,166	17,617	6,966	1,415	1
1999.....	1,394	1,896	2,451	4,172	7,700	13,658	19,681	23,023	17,580	7,104	1,321	1
White, both sexes												
2004.....	2,287	3,049	4,054	5,982	9,817	17,397	26,275	27,442	19,337	7,059	1,263	3
2003.....	2,248	3,045	3,974	5,928	10,537	18,762	27,567	29,266	20,465	7,753	1,355	2
2002.....	2,203	3,052	4,094	6,112	11,460	19,741	28,557	30,288	21,155	7,769	1,388	2
2001.....	2,139	2,907	3,967	6,188	11,859	20,546	28,719	30,444	20,767	7,672	1,342	1
2000.....	2,287	2,930	4,309	6,775	12,449	21,385	28,982	30,998	21,356	7,711	1,397	1
1999.....	2,074	2,939	4,001	6,876	12,701	21,726	29,191	30,860	21,385	7,852	1,351	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebrovascular diseases (I60-I69)												
White, male												
2004.....	1,312	1,656	2,250	3,239	4,910	7,993	10,464	9,048	4,864	1,299	161	3
2003.....	1,257	1,633	2,222	3,213	5,358	8,619	10,969	9,587	5,144	1,406	158	1
2002.....	1,202	1,684	2,253	3,284	5,737	8,921	11,116	9,970	5,202	1,356	164	1
2001.....	1,179	1,594	2,196	3,337	6,056	9,291	11,373	9,749	5,019	1,351	184	1
2000.....	1,244	1,607	2,391	3,540	6,288	9,746	11,464	9,892	5,095	1,367	172	1
1999.....	1,147	1,613	2,230	3,601	6,399	9,885	11,528	9,838	5,084	1,336	177	-
White, female												
2004.....	975	1,393	1,804	2,743	4,907	9,404	15,811	18,394	14,473	5,760	1,102	-
2003.....	991	1,412	1,752	2,715	5,179	10,143	16,598	19,679	15,321	6,347	1,197	1
2002.....	1,001	1,368	1,841	2,828	5,723	10,820	17,441	20,318	15,953	6,413	1,224	1
2001.....	960	1,313	1,771	2,851	5,803	11,255	17,346	20,695	15,748	6,321	1,158	-
2000.....	1,043	1,323	1,918	3,235	6,161	11,639	17,518	21,106	16,261	6,344	1,225	-
1999.....	927	1,326	1,771	3,275	6,302	11,841	17,663	21,022	16,301	6,516	1,174	-
Black, both sexes												
2004.....	1,069	1,113	1,320	1,500	1,884	2,473	2,617	2,119	1,462	561	153	2
2003.....	1,140	1,143	1,338	1,501	1,937	2,546	2,667	2,288	1,538	601	196	-
2002.....	1,075	1,059	1,265	1,597	2,033	2,603	2,723	2,322	1,559	611	168	1
2001.....	1,104	1,002	1,308	1,651	2,063	2,650	2,645	2,412	1,538	609	172	3
2000.....	958	1,043	1,248	1,582	2,107	2,674	2,725	2,492	1,514	688	197	1
1999.....	896	1,028	1,267	1,538	2,275	2,733	2,618	2,453	1,461	623	144	1
Black, male												
2004.....	576	642	746	777	920	1,035	944	598	345	101	24	-
2003.....	620	646	735	751	936	1,102	986	659	345	113	15	-
2002.....	593	597	701	857	985	1,094	955	679	351	100	38	-
2001.....	590	560	705	860	999	1,129	957	733	320	116	20	3
2000.....	520	594	664	818	1,017	1,153	1,024	730	369	145	21	-
1999.....	498	547	704	787	1,081	1,177	939	743	365	110	18	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebrovascular diseases (I60-I69)												
Black, female												
2004.....	493	471	574	723	964	1,438	1,673	1,521	1,117	460	129	2
2003.....	520	497	603	750	1,001	1,444	1,681	1,629	1,193	488	181	-
2002.....	482	462	564	740	1,048	1,509	1,768	1,643	1,208	511	130	1
2001.....	514	442	603	791	1,064	1,521	1,688	1,679	1,218	493	152	-
2000.....	438	449	584	764	1,090	1,521	1,701	1,762	1,145	543	176	1
1999.....	398	481	563	751	1,194	1,556	1,679	1,710	1,096	513	126	1
American Indian 1/, both sexes												
2004.....	30	31	33	64	69	73	95	77	35	16	7	-
2003.....	32	23	40	46	67	75	92	54	45	7	3	-
2002.....	27	34	37	49	67	69	89	59	47	15	4	-
2001.....	23	26	28	67	74	82	78	68	39	17	7	-
2000.....	25	32	47	44	64	86	92	66	47	16	-	-
1999.....	19	31	32	43	62	90	76	72	39	11	5	-
American Indian 1/, male												
2004.....	18	14	20	25	36	34	44	27	5	3	2	-
2003.....	14	13	20	20	34	34	32	19	15	2	-	-
2002.....	14	19	16	20	29	33	28	25	14	4	-	-
2001.....	13	11	12	29	35	25	26	24	13	4	1	-
2000.....	12	17	26	27	38	30	39	24	12	3	-	-
1999.....	11	12	14	23	36	48	27	20	9	3	1	-
American Indian 1/, female												
2004.....	12	17	13	39	33	39	51	50	30	13	5	-
2003.....	18	10	20	26	33	41	60	35	30	5	3	-
2002.....	13	15	21	29	38	36	61	34	33	11	4	-
2001.....	10	15	16	38	39	57	52	44	26	13	6	-
2000.....	13	15	21	17	26	56	53	42	35	13	-	-
1999.....	8	19	18	20	26	42	49	52	30	8	4	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebrovascular diseases (I60-I69)												
Asian or Pacific Islander 2/, both sexes												
2004.....	173	175	191	260	325	561	601	524	345	116	29	-
2003.....	130	181	202	290	402	534	604	531	319	132	26	-
2002.....	135	148	208	268	406	498	609	528	349	126	24	-
2001.....	137	158	212	264	432	494	608	541	306	100	32	-
2000.....	116	140	207	270	358	533	543	474	288	115	17	-
1999.....	119	154	200	237	360	458	535	456	264	85	19	1
Asian or Pacific Islander 2/, male												
2004.....	109	83	112	139	161	271	240	199	132	46	5	-
2003.....	70	102	107	154	202	238	254	206	130	54	6	-
2002.....	67	73	113	144	151	235	273	210	146	50	7	-
2001.....	69	86	113	119	214	226	274	225	130	35	6	-
2000.....	58	76	121	128	186	263	227	218	112	49	3	-
1999.....	58	84	101	111	182	239	245	217	111	18	2	1
Asian or Pacific Islander 2/, female												
2004.....	64	92	79	121	164	290	361	325	213	70	24	-
2003.....	60	79	95	136	200	296	350	325	189	78	20	-
2002.....	68	75	95	124	255	263	336	318	203	76	17	-
2001.....	68	72	99	145	218	268	334	316	176	65	26	-
2000.....	58	64	86	142	172	270	316	256	176	66	14	-
1999.....	61	70	99	126	178	219	290	239	153	67	17	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Subarachnoid hemorrhage (I60)												
All races, both sexes												
2004.....	654	607	540	472	505	571	512	361	166	58	8	-
2003.....	687	594	518	483	496	573	548	346	152	40	4	-
2002.....	585	574	552	524	535	587	548	343	171	40	8	2
2001.....	624	600	473	502	555	596	506	365	162	49	5	-
2000.....	667	639	545	495	566	641	527	372	174	44	9	1
1999.....	620	579	543	541	599	615	533	345	150	45	4	1
All races, male												
2004.....	285	223	198	148	158	176	137	98	33	9	-	-
2003.....	267	214	197	180	154	168	153	93	31	7	1	-
2002.....	201	217	209	187	179	171	123	78	35	10	3	1
2001.....	255	218	166	154	203	173	144	88	30	4	1	-
2000.....	249	219	184	150	201	191	142	85	40	11	2	1
1999.....	242	209	184	186	173	183	139	74	25	8	-	1
All races, female												
2004.....	369	384	342	324	347	395	375	263	133	49	8	-
2003.....	420	380	321	303	342	405	395	253	121	33	3	-
2002.....	384	357	343	337	356	416	425	265	136	30	5	1
2001.....	369	382	307	348	352	423	362	277	132	45	4	-
2000.....	418	420	361	345	365	450	385	287	134	33	7	-
1999.....	378	370	359	355	426	432	394	271	125	37	4	-
White, both sexes												
2004.....	493	493	429	394	429	485	456	327	155	54	7	-
2003.....	526	483	433	388	422	485	474	300	134	37	2	-
2002.....	444	452	456	437	436	511	485	301	156	35	6	2
2001.....	456	463	403	395	471	511	461	315	148	40	5	-
2000.....	523	534	442	394	473	561	468	326	157	34	5	1
1999.....	482	469	434	465	514	546	474	308	131	33	3	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Subarachnoid hemorrhage (I60)												
White, male												
2004.....	216	181	156	126	140	154	124	94	32	9	-	-
2003.....	215	169	161	146	143	141	133	80	27	6	-	-
2002.....	158	172	171	157	149	154	108	70	33	8	3	1
2001.....	192	164	143	128	174	151	138	76	27	4	1	-
2000.....	196	188	145	123	168	172	131	82	37	10	2	1
1999.....	193	174	152	155	152	171	125	70	23	4	-	-
White, female												
2004.....	277	312	273	268	289	331	332	233	123	45	7	-
2003.....	311	314	272	242	279	344	341	220	107	31	2	-
2002.....	286	280	285	280	287	357	377	231	123	27	3	1
2001.....	264	299	260	267	297	360	323	239	121	36	4	-
2000.....	327	346	297	271	305	389	337	244	120	24	3	-
1999.....	289	295	282	310	362	375	349	238	108	29	3	-
Black, both sexes												
2004.....	119	88	83	47	44	47	41	22	7	2	1	-
2003.....	123	84	57	63	41	51	46	31	15	3	1	-
2002.....	116	89	65	58	50	48	35	29	12	2	-	-
2001.....	131	102	54	76	47	56	25	31	9	5	-	-
2000.....	116	83	68	67	65	62	37	34	9	8	4	-
1999.....	112	88	72	50	64	42	43	23	14	8	1	-
Black, male												
2004.....	47	34	32	15	12	11	9	2	-	-	-	-
2003.....	39	36	30	23	6	13	13	9	3	1	-	-
2002.....	33	35	26	23	20	12	8	5	-	-	-	-
2001.....	49	37	20	18	15	17	4	4	2	-	-	-
2000.....	40	23	23	18	25	17	6	1	1	-	-	-
1999.....	43	30	23	22	16	6	13	2	1	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Subarachnoid hemorrhage (I60)												
Black, female												
2004.....	72	54	51	32	32	36	32	20	7	2	1	-
2003.....	84	48	27	40	35	38	33	22	12	2	1	-
2002.....	83	54	39	35	30	36	27	24	12	2	-	-
2001.....	82	65	34	58	32	39	21	27	7	5	-	-
2000.....	76	60	45	49	40	45	31	33	8	8	4	-
1999.....	69	58	49	28	48	36	30	21	13	7	1	-
American Indian 1/, both sexes												
2004.....	8	7	2	4	6	3	2	-	-	-	-	-
2003.....	8	3	3	5	3	4	4	1	1	-	-	-
2002.....	2	5	4	3	4	3	2	-	1	-	1	-
2001.....	4	3	2	7	3	2	2	3	-	1	-	-
2000.....	6	6	8	2	-	1	-	2	-	-	-	-
1999.....	3	4	5	-	3	2	-	-	1	-	-	-
American Indian 1/, male												
2004.....	3	2	-	2	1	1	-	-	-	-	-	-
2003.....	2	1	-	2	-	-	1	-	1	-	-	-
2002.....	2	2	2	-	3	1	-	-	-	-	-	-
2001.....	2	3	1	2	2	-	-	-	-	-	-	-
2000.....	3	2	4	-	-	-	-	1	-	-	-	-
1999.....	-	-	-	-	1	1	-	-	-	-	-	-
American Indian 1/, female												
2004.....	5	5	2	2	5	2	2	-	-	-	-	-
2003.....	6	2	3	3	3	4	3	1	-	-	-	-
2002.....	-	3	2	3	1	2	2	-	1	-	1	-
2001.....	2	-	1	5	1	2	2	3	-	1	-	-
2000.....	3	4	4	2	-	1	-	1	-	-	-	-
1999.....	3	4	5	-	2	1	-	-	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Subarachnoid hemorrhage (I60)												
Asian or Pacific Islander 2/, both sexes												
2004.....	34	19	26	27	26	36	13	12	4	2	-	-
2003.....	30	24	25	27	30	33	24	14	2	-	1	-
2002.....	23	28	27	26	45	25	26	13	2	3	1	-
2001.....	33	32	14	24	34	27	18	16	5	3	-	-
2000.....	22	16	27	32	28	17	22	10	8	2	-	-
1999.....	23	18	32	26	18	25	16	14	4	4	-	1
Asian or Pacific Islander 2/, male												
2004.....	19	6	10	5	5	10	4	2	1	-	-	-
2003.....	11	8	6	9	5	14	6	4	-	-	1	-
2002.....	8	8	10	7	7	4	7	3	2	2	-	-
2001.....	12	14	2	6	12	5	2	8	1	-	-	-
2000.....	10	6	12	9	8	2	5	1	2	1	-	-
1999.....	6	5	9	9	4	5	1	2	1	3	-	1
Asian or Pacific Islander 2/, female												
2004.....	15	13	16	22	21	26	9	10	3	2	-	-
2003.....	19	16	19	18	25	19	18	10	2	-	-	-
2002.....	15	20	17	19	38	21	19	10	-	1	1	-
2001.....	21	18	12	18	22	22	16	8	4	3	-	-
2000.....	12	10	15	23	20	15	17	9	6	1	-	-
1999.....	17	13	23	17	14	20	15	12	3	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intracerebral and other intracranial hemorrhage (I61-I62)												
All races, both sexes												
2004.....	1,471	1,614	1,753	2,113	2,898	4,262	4,981	3,696	1,857	448	40	1
2003.....	1,440	1,499	1,685	1,990	3,000	4,300	4,850	3,755	1,723	433	56	-
2002.....	1,354	1,451	1,664	2,001	3,133	4,386	4,817	3,703	1,689	395	55	-
2001.....	1,371	1,345	1,651	2,025	3,092	4,323	4,693	3,578	1,609	401	43	4
2000.....	1,339	1,427	1,716	2,239	3,145	4,459	4,702	3,622	1,737	437	51	-
1999.....	1,197	1,399	1,528	1,971	3,233	4,367	4,420	3,161	1,500	372	46	-
All races, male												
2004.....	866	936	1,010	1,155	1,430	1,959	2,081	1,340	576	103	8	1
2003.....	872	845	946	1,066	1,527	1,959	2,051	1,366	494	117	8	-
2002.....	805	802	953	1,095	1,584	1,946	1,978	1,314	467	84	11	-
2001.....	789	772	914	1,044	1,614	2,003	1,919	1,226	409	100	9	4
2000.....	768	875	977	1,190	1,560	2,007	1,846	1,228	481	119	8	-
1999.....	711	803	877	1,033	1,622	1,959	1,697	1,053	419	79	9	-
All races, female												
2004.....	605	678	743	958	1,468	2,303	2,900	2,356	1,281	345	32	-
2003.....	568	654	739	924	1,473	2,341	2,799	2,389	1,229	316	48	-
2002.....	549	649	711	906	1,549	2,440	2,839	2,389	1,222	311	44	-
2001.....	582	573	737	981	1,478	2,320	2,774	2,352	1,200	301	34	-
2000.....	571	552	739	1,049	1,585	2,452	2,856	2,394	1,256	318	43	-
1999.....	486	596	651	938	1,611	2,408	2,723	2,108	1,081	293	37	-
White, both sexes												
2004.....	867	1,071	1,287	1,659	2,437	3,722	4,483	3,408	1,689	417	29	1
2003.....	800	979	1,176	1,549	2,526	3,782	4,358	3,442	1,557	386	52	-
2002.....	787	1,003	1,243	1,552	2,663	3,879	4,375	3,411	1,547	361	45	-
2001.....	784	946	1,187	1,572	2,605	3,835	4,266	3,265	1,474	359	38	1
2000.....	834	981	1,267	1,794	2,715	3,952	4,280	3,331	1,588	389	42	-
1999.....	734	959	1,117	1,607	2,786	3,887	4,022	2,879	1,387	333	39	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intracerebral and other intracranial hemorrhage (I61-I62)												
White, male												
2004.....	531	617	734	920	1,206	1,730	1,897	1,251	524	93	5	1
2003.....	497	550	657	852	1,283	1,744	1,858	1,268	439	101	7	-
2002.....	471	548	721	858	1,371	1,751	1,812	1,216	426	71	8	-
2001.....	459	545	680	815	1,375	1,790	1,760	1,114	377	87	9	1
2000.....	487	600	717	947	1,356	1,800	1,692	1,131	433	97	7	-
1999.....	439	553	654	859	1,419	1,742	1,542	958	386	72	6	-
White, female												
2004.....	336	454	553	739	1,231	1,992	2,586	2,157	1,165	324	24	-
2003.....	303	429	519	697	1,243	2,038	2,500	2,174	1,118	285	45	-
2002.....	316	455	522	694	1,292	2,128	2,563	2,195	1,121	290	37	-
2001.....	325	401	507	757	1,230	2,045	2,506	2,151	1,097	272	29	-
2000.....	347	381	550	847	1,359	2,152	2,588	2,200	1,155	292	35	-
1999.....	295	406	463	748	1,367	2,145	2,480	1,921	1,001	261	33	-
Black, both sexes												
2004.....	510	438	383	345	341	361	357	193	115	21	10	-
2003.....	557	432	403	322	328	377	339	209	124	32	3	-
2002.....	481	359	323	328	333	359	310	195	102	27	7	-
2001.....	512	320	357	343	325	341	307	206	105	31	5	3
2000.....	438	371	354	327	318	356	277	209	116	41	8	-
1999.....	394	349	317	276	328	350	268	186	81	32	5	-
Black, male												
2004.....	277	266	225	176	164	141	134	62	32	5	2	-
2003.....	326	249	229	153	166	165	134	64	39	8	1	-
2002.....	289	207	182	174	157	134	106	59	26	8	2	-
2001.....	289	186	179	181	161	135	107	71	20	8	-	3
2000.....	245	233	210	178	141	148	93	64	32	16	1	-
1999.....	232	196	178	131	142	149	93	59	25	6	3	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intracerebral and other intracranial hemorrhage (I61-I62)												
Black, female												
2004.....	233	172	158	169	177	220	223	131	83	16	8	-
2003.....	231	183	174	169	162	212	205	145	85	24	2	-
2002.....	192	152	141	154	176	225	204	136	76	19	5	-
2001.....	223	134	178	162	164	206	200	135	85	23	5	-
2000.....	193	138	144	149	177	208	184	145	84	25	7	-
1999.....	162	153	139	145	186	201	175	127	56	26	2	-
American Indian 1/, both sexes												
2004.....	8	13	11	13	15	13	14	9	3	2	-	-
2003.....	16	8	15	8	11	13	21	7	3	-	-	-
2002.....	15	16	10	12	13	8	8	9	3	1	1	-
2001.....	8	8	10	16	19	12	11	11	2	-	-	-
2000.....	11	9	10	9	13	13	17	10	5	1	-	-
1999.....	8	11	12	5	12	13	10	12	2	-	1	-
American Indian 1/, male												
2004.....	4	7	8	3	8	5	9	3	1	1	-	-
2003.....	8	4	8	3	5	3	8	2	2	-	-	-
2002.....	9	12	4	3	2	-	3	7	-	1	-	-
2001.....	6	3	4	7	10	5	2	4	-	-	-	-
2000.....	6	5	6	7	6	-	7	3	2	1	-	-
1999.....	5	5	6	3	4	4	2	3	-	-	-	-
American Indian 1/, female												
2004.....	4	6	3	10	7	8	5	6	2	1	-	-
2003.....	8	4	7	5	6	10	13	5	1	-	-	-
2002.....	6	4	6	9	11	8	5	2	3	-	1	-
2001.....	2	5	6	9	9	7	9	7	2	-	-	-
2000.....	5	4	4	2	7	13	10	7	3	-	-	-
1999.....	3	6	6	2	8	9	8	9	2	-	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intracerebral and other intracranial hemorrhage (I61-I62)												
Asian or Pacific Islander 2/, both sexes												
2004.....	86	92	72	96	105	166	127	86	50	8	1	-
2003.....	67	80	91	111	135	128	132	97	39	15	1	-
2002.....	71	73	88	109	124	140	124	88	37	6	2	-
2001.....	67	71	97	94	143	135	109	96	28	11	-	-
2000.....	56	66	85	109	99	138	128	72	28	6	1	-
1999.....	61	80	82	83	107	117	120	84	30	7	1	-
Asian or Pacific Islander 2/, male												
2004.....	54	46	43	56	52	83	41	24	19	4	1	-
2003.....	41	42	52	58	73	47	51	32	14	8	-	-
2002.....	36	35	46	60	54	61	57	32	15	4	1	-
2001.....	35	38	51	41	68	73	50	37	12	5	-	-
2000.....	30	37	44	58	57	59	54	30	14	5	-	-
1999.....	35	49	39	40	57	64	60	33	8	1	-	-
Asian or Pacific Islander 2/, female												
2004.....	32	46	29	40	53	83	86	62	31	4	-	-
2003.....	26	38	39	53	62	81	81	65	25	7	1	-
2002.....	35	38	42	49	70	79	67	56	22	2	1	-
2001.....	32	33	46	53	75	62	59	59	16	6	-	-
2000.....	26	29	41	51	42	79	74	42	14	1	1	-
1999.....	26	31	43	43	50	53	60	51	22	6	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebral infarction (I63)												
All races, both sexes												
2004.....	193	289	387	543	875	1,504	2,104	2,416	1,614	609	110	2
2003.....	208	290	366	609	933	1,612	2,412	2,565	1,890	712	122	-
2002.....	236	249	408	558	1,008	1,679	2,422	2,665	1,894	754	119	-
2001.....	197	268	347	578	1,138	1,806	2,451	2,751	1,867	763	114	-
2000.....	215	270	435	633	1,130	2,015	2,694	2,973	2,066	794	146	1
1999.....	206	309	430	769	1,321	2,183	2,857	3,218	2,187	787	168	-
All races, male												
2004.....	129	182	244	315	440	727	842	801	415	107	11	1
2003.....	123	165	231	344	504	724	981	873	448	111	17	-
2002.....	149	164	233	321	477	793	1,006	876	479	138	18	-
2001.....	112	176	199	341	578	855	1,028	841	457	134	13	-
2000.....	118	160	265	343	600	949	1,116	990	487	143	16	-
1999.....	124	185	244	442	677	1,025	1,141	1,066	511	119	29	-
All races, female												
2004.....	64	107	143	228	435	777	1,262	1,615	1,199	502	99	1
2003.....	85	125	135	265	429	888	1,431	1,692	1,442	601	105	-
2002.....	87	85	175	237	531	886	1,416	1,789	1,415	616	101	-
2001.....	85	92	148	237	560	951	1,423	1,910	1,410	629	101	-
2000.....	97	110	170	290	530	1,066	1,578	1,983	1,579	651	130	1
1999.....	82	124	186	327	644	1,158	1,716	2,152	1,676	668	139	-
White, both sexes												
2004.....	137	212	286	431	708	1,281	1,874	2,171	1,457	552	97	1
2003.....	143	206	259	491	760	1,389	2,163	2,337	1,739	655	109	-
2002.....	165	191	317	436	835	1,459	2,149	2,416	1,733	695	101	-
2001.....	147	197	256	447	954	1,565	2,210	2,503	1,723	699	97	-
2000.....	146	183	338	500	941	1,751	2,417	2,688	1,900	719	130	-
1999.....	149	230	323	620	1,084	1,883	2,598	2,927	2,017	719	150	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebral infarction (I63)												
White, male												
2004.....	92	137	173	253	353	623	760	716	374	97	10	1
2003.....	91	117	168	282	411	622	902	791	416	102	17	-
2002.....	102	128	179	252	400	697	898	790	439	122	14	-
2001.....	87	130	154	273	497	737	928	748	421	122	9	-
2000.....	76	111	207	269	508	820	999	896	441	124	15	-
1999.....	93	144	183	368	553	880	1,045	959	459	109	27	-
White, female												
2004.....	45	75	113	178	355	658	1,114	1,455	1,083	455	87	-
2003.....	52	89	91	209	349	767	1,261	1,546	1,323	553	92	-
2002.....	63	63	138	184	435	762	1,251	1,626	1,294	573	87	-
2001.....	60	67	102	174	457	828	1,282	1,755	1,302	577	88	-
2000.....	70	72	131	231	433	931	1,418	1,792	1,459	595	115	-
1999.....	56	86	140	252	531	1,003	1,553	1,968	1,558	610	123	-
Black, both sexes												
2004.....	49	68	90	92	146	196	177	185	129	43	10	1
2003.....	57	75	91	101	142	190	204	186	125	46	12	-
2002.....	65	53	77	104	149	188	217	197	128	42	14	-
2001.....	45	63	81	114	161	201	197	202	122	57	14	-
2000.....	58	73	85	118	158	212	239	234	138	69	15	1
1999.....	48	65	93	130	205	263	217	243	152	61	17	-
Black, male												
2004.....	32	39	60	56	70	92	64	61	30	6	1	-
2003.....	28	43	56	53	76	85	68	63	24	6	-	-
2002.....	42	33	45	59	68	78	83	65	31	8	3	-
2001.....	24	42	41	62	69	102	81	72	26	12	2	-
2000.....	39	43	49	68	74	99	99	70	33	15	1	-
1999.....	25	37	51	66	111	127	79	88	41	8	2	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebral infarction (I63)												
Black, female												
2004.....	17	29	30	36	76	104	113	124	99	37	9	1
2003.....	29	32	35	48	66	105	136	123	101	40	12	-
2002.....	23	20	32	45	81	110	134	132	97	34	11	-
2001.....	21	21	40	52	92	99	116	130	96	45	12	-
2000.....	19	30	36	50	84	113	140	164	105	54	14	1
1999.....	23	28	42	64	94	136	138	155	111	53	15	-
American Indian 1/, both sexes												
2004.....	1	2	-	6	3	6	5	6	4	1	-	-
2003.....	2	3	1	2	3	3	10	5	3	-	-	-
2002.....	2	2	7	3	4	10	11	6	1	3	-	-
2001.....	1	3	3	3	5	1	3	7	2	2	-	-
2000.....	2	4	1	2	9	7	6	7	3	-	-	-
1999.....	2	2	1	3	6	6	5	7	3	1	-	-
American Indian 1/, male												
2004.....	1	2	-	1	2	3	2	5	1	-	-	-
2003.....	1	2	-	1	2	3	2	1	-	-	-	-
2002.....	1	1	4	-	1	7	4	3	-	1	-	-
2001.....	-	1	1	2	3	1	-	3	-	-	-	-
2000.....	-	2	1	1	6	2	2	2	1	-	-	-
1999.....	1	-	1	2	3	3	1	1	2	-	-	-
American Indian 1/, female												
2004.....	-	-	-	5	1	3	3	1	3	1	-	-
2003.....	1	1	1	1	1	-	8	4	3	-	-	-
2002.....	1	1	3	3	3	3	7	3	1	2	-	-
2001.....	1	2	2	1	2	-	3	4	2	2	-	-
2000.....	2	2	-	1	3	5	4	5	2	-	-	-
1999.....	1	2	-	1	3	3	4	6	1	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebral infarction (I63)												
Asian or Pacific Islander 2/, both sexes												
2004.....	6	7	11	14	18	21	48	54	24	13	3	-
2003.....	6	6	15	15	28	30	35	37	23	11	1	-
2002.....	4	3	7	15	20	22	45	46	32	14	4	-
2001.....	4	5	7	14	18	39	41	39	20	5	3	-
2000.....	9	10	11	13	22	45	32	44	25	6	1	-
1999.....	7	12	13	16	26	31	37	41	15	6	1	-
Asian or Pacific Islander 2/, male												
2004.....	4	4	11	5	15	9	16	19	10	4	-	-
2003.....	3	3	7	8	15	14	9	18	8	3	-	-
2002.....	4	2	5	10	8	11	21	18	9	7	1	-
2001.....	1	3	3	4	9	15	19	18	10	-	2	-
2000.....	3	4	8	5	12	28	16	22	12	4	-	-
1999.....	5	4	9	6	10	15	16	18	9	2	-	-
Asian or Pacific Islander 2/, female												
2004.....	2	3	-	9	3	12	32	35	14	9	3	-
2003.....	3	3	8	7	13	16	26	19	15	8	1	-
2002.....	-	1	2	5	12	11	24	28	23	7	3	-
2001.....	3	2	4	10	9	24	22	21	10	5	1	-
2000.....	6	6	3	8	10	17	16	22	13	2	1	-
1999.....	2	8	4	10	16	16	21	23	6	4	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Stroke, not specified as hemorrhage or infarction (I64)												
All races, both sexes												
2004.....	965	1,422	2,259	3,580	5,967	10,505	16,017	16,975	12,329	4,493	870	1
2003.....	965	1,570	2,365	3,600	6,448	11,610	17,085	18,425	13,283	5,037	949	1
2002.....	989	1,608	2,375	3,861	7,187	12,299	17,964	19,474	13,922	5,118	957	1
2001.....	961	1,495	2,442	3,974	7,529	13,066	18,299	19,929	13,867	5,156	956	-
2000.....	911	1,435	2,487	4,253	7,955	13,429	18,408	20,219	14,096	5,203	957	-
1999.....	854	1,481	2,400	4,301	7,953	13,667	18,598	20,387	14,254	5,265	905	1
All races, male												
2004.....	585	803	1,285	1,927	3,034	4,712	6,194	5,353	2,971	837	120	1
2003.....	569	924	1,345	1,952	3,270	5,324	6,602	5,810	3,267	922	101	1
2002.....	562	961	1,344	2,110	3,597	5,525	6,722	6,259	3,341	883	118	-
2001.....	562	866	1,403	2,178	3,807	5,762	7,020	6,289	3,310	920	130	-
2000.....	562	835	1,429	2,243	3,993	6,032	7,172	6,240	3,311	915	125	-
1999.....	514	830	1,381	2,273	4,019	6,202	7,294	6,361	3,334	890	112	-
All races, female												
2004.....	380	619	974	1,653	2,933	5,793	9,823	11,622	9,358	3,656	750	-
2003.....	396	646	1,020	1,648	3,178	6,286	10,483	12,615	10,016	4,115	848	-
2002.....	427	647	1,031	1,751	3,590	6,774	11,242	13,215	10,581	4,235	839	1
2001.....	399	629	1,039	1,796	3,722	7,304	11,279	13,640	10,557	4,236	826	-
2000.....	349	600	1,058	2,010	3,962	7,397	11,236	13,979	10,785	4,288	832	-
1999.....	340	651	1,019	2,028	3,934	7,465	11,304	14,026	10,920	4,375	793	1
White, both sexes												
2004.....	599	970	1,588	2,657	4,783	8,827	14,191	15,446	11,294	4,094	749	1
2003.....	614	1,072	1,665	2,693	5,179	9,848	15,225	16,798	12,154	4,604	803	1
2002.....	624	1,104	1,650	2,878	5,813	10,537	15,990	17,756	12,748	4,667	837	-
2001.....	603	1,028	1,690	2,947	6,101	11,252	16,325	18,122	12,745	4,724	819	-
2000.....	609	969	1,795	3,271	6,508	11,598	16,434	18,438	12,995	4,716	827	-
1999.....	549	1,010	1,695	3,309	6,479	11,803	16,719	18,604	13,183	4,820	804	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Stroke, not specified as hemorrhage or infarction (I64)												
White, male												
2004.....	376	545	912	1,449	2,447	4,005	5,525	4,908	2,720	745	99	1
2003.....	373	629	974	1,490	2,647	4,565	5,902	5,323	2,971	817	88	1
2002.....	364	665	937	1,565	2,933	4,761	6,007	5,721	3,035	795	90	-
2001.....	361	595	967	1,628	3,103	5,000	6,268	5,717	3,026	828	114	-
2000.....	388	569	1,048	1,744	3,284	5,215	6,429	5,668	3,014	802	108	-
1999.....	341	583	982	1,753	3,321	5,371	6,606	5,787	3,048	811	101	-
White, female												
2004.....	223	425	676	1,208	2,336	4,822	8,666	10,538	8,574	3,349	650	-
2003.....	241	443	691	1,203	2,532	5,283	9,323	11,475	9,183	3,787	715	-
2002.....	260	439	713	1,313	2,880	5,776	9,983	12,035	9,713	3,872	747	-
2001.....	242	433	723	1,319	2,998	6,252	10,057	12,405	9,719	3,896	705	-
2000.....	221	400	747	1,527	3,224	6,383	10,005	12,770	9,981	3,914	719	-
1999.....	208	427	713	1,556	3,158	6,432	10,113	12,817	10,135	4,009	703	-
Black, both sexes												
2004.....	321	404	592	795	1,010	1,398	1,486	1,253	826	329	96	-
2003.....	322	439	622	781	1,080	1,472	1,522	1,339	912	356	127	-
2002.....	329	461	640	864	1,169	1,512	1,625	1,401	949	373	108	1
2001.....	327	415	664	896	1,212	1,548	1,606	1,478	920	366	113	-
2000.....	275	418	604	881	1,258	1,551	1,675	1,498	913	408	120	-
1999.....	283	426	642	880	1,289	1,608	1,574	1,518	886	388	86	1
Black, male												
2004.....	176	234	329	407	493	566	524	333	179	64	17	-
2003.....	181	258	327	388	520	625	556	370	205	70	9	-
2002.....	181	269	359	475	583	640	561	412	213	60	25	-
2001.....	183	242	383	483	593	652	586	443	194	70	13	-
2000.....	160	237	322	450	609	681	616	448	232	83	15	-
1999.....	161	223	364	463	596	692	561	453	216	70	10	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Stroke, not specified as hemorrhage or infarction (I64)												
Black, female												
2004.....	145	170	263	388	517	832	962	920	647	265	79	-
2003.....	141	181	295	393	560	847	966	969	707	286	118	-
2002.....	148	192	281	389	586	872	1,064	989	736	313	83	1
2001.....	144	173	281	413	619	896	1,020	1,035	726	296	100	-
2000.....	115	181	282	431	649	870	1,059	1,050	681	325	105	-
1999.....	122	203	278	417	693	916	1,013	1,065	670	318	76	1
American Indian 1/, both sexes												
2004.....	12	6	16	34	40	41	51	41	21	10	5	-
2003.....	6	9	18	24	34	44	44	25	30	3	2	-
2002.....	5	9	12	29	35	35	55	37	30	7	1	-
2001.....	7	12	11	33	37	53	47	37	22	11	5	-
2000.....	6	11	22	18	29	50	49	41	31	10	-	-
1999.....	5	13	10	28	28	52	41	46	29	9	2	-
American Indian 1/, male												
2004.....	9	2	8	17	22	21	26	14	3	1	1	-
2003.....	3	6	9	13	18	22	16	11	8	1	-	-
2002.....	1	4	4	17	17	20	18	12	10	2	-	-
2001.....	3	4	5	13	18	16	17	13	9	3	-	-
2000.....	3	6	11	11	21	20	21	16	4	2	-	-
1999.....	4	6	4	16	20	29	16	14	7	2	-	-
American Indian 1/, female												
2004.....	3	4	8	17	18	20	25	27	18	9	4	-
2003.....	3	3	9	11	16	22	28	14	22	2	2	-
2002.....	4	5	8	12	18	15	37	25	20	5	1	-
2001.....	4	8	6	20	19	37	30	24	13	8	5	-
2000.....	3	5	11	7	8	30	28	25	27	8	-	-
1999.....	1	7	6	12	8	23	25	32	22	7	2	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Stroke, not specified as hemorrhage or infarction (I64)												
Asian or Pacific Islander 2/, both sexes												
2004.....	33	42	63	94	134	239	289	235	188	60	20	-
2003.....	23	50	60	102	155	246	294	263	187	74	17	-
2002.....	31	34	73	90	170	215	294	280	195	71	11	-
2001.....	24	40	77	98	179	213	321	292	180	55	19	-
2000.....	21	37	66	83	160	230	250	242	157	69	10	-
1999.....	17	32	53	84	157	204	264	219	156	48	13	-
Asian or Pacific Islander 2/, male												
2004.....	24	22	36	54	72	120	119	98	69	27	3	-
2003.....	12	31	35	61	85	112	128	106	83	34	4	-
2002.....	16	23	44	53	64	104	136	114	83	26	3	-
2001.....	15	25	48	54	93	94	149	116	81	19	3	-
2000.....	11	23	48	38	79	116	106	108	61	28	2	-
1999.....	8	18	31	41	82	110	111	107	63	7	1	-
Asian or Pacific Islander 2/, female												
2004.....	9	20	27	40	62	119	170	137	119	33	17	-
2003.....	11	19	25	41	70	134	166	157	104	40	13	-
2002.....	15	11	29	37	106	111	158	166	112	45	8	-
2001.....	9	15	29	44	86	119	172	176	99	36	16	-
2000.....	10	14	18	45	81	114	144	134	96	41	8	-
1999.....	9	14	22	43	75	94	153	112	93	41	12	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other cerebrovascular diseases and their sequelae (I67,I69)												
All races, both sexes												
2004.....	276	436	659	1,098	1,850	3,662	5,974	6,714	5,213	2,144	424	1
2003.....	250	439	620	1,083	2,066	3,822	6,035	7,048	5,319	2,271	449	1
2002.....	276	411	605	1,082	2,103	3,960	6,227	7,012	5,434	2,214	445	-
2001.....	250	385	602	1,091	2,114	3,981	6,101	6,842	5,145	2,029	435	-
2000.....	254	374	628	1,051	2,182	4,134	6,011	6,844	5,132	2,052	448	-
1999.....	231	384	599	1,112	2,292	4,175	6,012	6,730	5,058	2,102	396	-
All races, male												
2004.....	150	251	391	635	965	1,759	2,438	2,280	1,351	393	53	-
2003.....	130	246	365	596	1,075	1,818	2,454	2,329	1,394	418	52	-
2002.....	159	229	344	592	1,065	1,848	2,543	2,357	1,391	395	59	-
2001.....	133	219	344	628	1,102	1,878	2,519	2,287	1,276	348	58	-
2000.....	137	205	347	587	1,175	2,013	2,478	2,321	1,269	376	45	-
1999.....	123	229	363	588	1,207	1,980	2,468	2,264	1,280	371	48	-
All races, female												
2004.....	126	185	268	463	885	1,903	3,536	4,434	3,862	1,751	371	1
2003.....	120	193	255	487	991	2,004	3,581	4,719	3,925	1,853	397	1
2002.....	117	182	261	490	1,038	2,112	3,684	4,655	4,043	1,819	386	-
2001.....	117	166	258	463	1,012	2,103	3,582	4,555	3,869	1,681	377	-
2000.....	117	169	281	464	1,007	2,121	3,533	4,523	3,863	1,676	403	-
1999.....	108	155	236	524	1,085	2,195	3,544	4,466	3,778	1,731	348	-
White, both sexes												
2004.....	191	303	464	841	1,460	3,082	5,271	6,090	4,742	1,942	381	-
2003.....	165	305	441	807	1,650	3,258	5,347	6,389	4,881	2,071	389	1
2002.....	183	302	428	809	1,713	3,355	5,558	6,404	4,971	2,011	399	-
2001.....	149	273	431	827	1,728	3,383	5,457	6,239	4,677	1,850	383	-
2000.....	175	263	467	816	1,812	3,523	5,383	6,215	4,716	1,853	393	-
1999.....	160	271	432	875	1,838	3,607	5,378	6,142	4,667	1,947	355	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other cerebrovascular diseases and their sequelae (I67,I69)												
White, male												
2004.....	97	176	275	491	764	1,481	2,158	2,079	1,214	355	47	-
2003.....	81	168	262	443	874	1,547	2,174	2,125	1,291	380	46	-
2002.....	107	171	245	452	884	1,558	2,291	2,173	1,269	360	49	-
2001.....	80	160	252	493	907	1,613	2,279	2,094	1,168	310	51	-
2000.....	97	139	274	457	972	1,739	2,213	2,115	1,170	334	40	-
1999.....	81	159	259	466	954	1,721	2,210	2,064	1,168	340	43	-
White, female												
2004.....	94	127	189	350	696	1,601	3,113	4,011	3,528	1,587	334	-
2003.....	84	137	179	364	776	1,711	3,173	4,264	3,590	1,691	343	1
2002.....	76	131	183	357	829	1,797	3,267	4,231	3,702	1,651	350	-
2001.....	69	113	179	334	821	1,770	3,178	4,145	3,509	1,540	332	-
2000.....	78	124	193	359	840	1,784	3,170	4,100	3,546	1,519	353	-
1999.....	79	112	173	409	884	1,886	3,168	4,078	3,499	1,607	312	-
Black, both sexes												
2004.....	70	115	172	221	343	471	556	466	385	166	36	1
2003.....	81	113	165	234	346	456	556	523	362	164	53	-
2002.....	84	97	160	243	332	496	536	500	368	167	39	-
2001.....	89	102	152	222	318	504	510	495	382	150	40	-
2000.....	71	98	137	189	308	493	497	517	338	162	50	-
1999.....	59	100	143	202	389	470	516	483	328	134	35	-
Black, male												
2004.....	44	69	100	123	181	225	213	140	104	26	4	-
2003.....	46	60	93	134	168	214	215	153	74	28	5	-
2002.....	48	53	89	126	157	230	197	138	81	24	8	-
2001.....	45	53	82	116	161	223	179	143	78	26	5	-
2000.....	36	58	60	104	168	208	210	147	71	31	4	-
1999.....	37	61	88	105	216	203	193	141	82	25	3	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other cerebrovascular diseases and their sequelae (I67,I69)

Black, female

2004.....	26	46	72	98	162	246	343	326	281	140	32	1
2003.....	35	53	72	100	178	242	341	370	288	136	48	-
2002.....	36	44	71	117	175	266	339	362	287	143	31	-
2001.....	44	49	70	106	157	281	331	352	304	124	35	-
2000.....	35	40	77	85	140	285	287	370	267	131	46	-
1999.....	22	39	55	97	173	267	323	342	246	109	32	-

American Indian 1/, both sexes

2004.....	1	3	4	7	5	10	23	21	7	3	2	-
2003.....	-	-	3	7	16	11	13	16	8	4	1	-
2002.....	3	2	4	2	11	13	13	7	12	4	1	-
2001.....	3	-	2	8	10	14	15	10	13	3	2	-
2000.....	-	2	6	13	13	15	20	6	8	5	-	-
1999.....	1	1	4	7	13	17	20	7	4	1	2	-

American Indian 1/, male

2004.....	1	1	4	2	3	4	7	5	-	1	1	-
2003.....	-	-	3	1	9	6	5	5	4	1	-	-
2002.....	1	-	2	-	6	5	3	3	4	-	-	-
2001.....	2	-	1	5	2	3	7	4	4	1	1	-
2000.....	-	2	4	8	5	8	9	2	5	-	-	-
1999.....	1	1	3	2	8	11	8	2	-	1	1	-

American Indian 1/, female

2004.....	-	2	-	5	2	6	16	16	7	2	1	-
2003.....	-	-	-	6	7	5	8	11	4	3	1	-
2002.....	2	2	2	2	5	8	10	4	8	4	1	-
2001.....	1	-	1	3	8	11	8	6	9	2	1	-
2000.....	-	-	2	5	8	7	11	4	3	5	-	-
1999.....	-	-	1	5	5	6	12	5	4	-	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other cerebrovascular diseases and their sequelae (I67,I69)												
Asian or Pacific Islander 2/, both sexes												
2004.....	14	15	19	29	42	99	124	137	79	33	5	-
2003.....	4	21	11	35	54	97	119	120	68	32	6	-
2002.....	6	10	13	28	47	96	120	101	83	32	6	-
2001.....	9	10	17	34	58	80	119	98	73	26	10	-
2000.....	8	11	18	33	49	103	111	106	70	32	5	-
1999.....	11	12	20	28	52	81	98	98	59	20	4	-
Asian or Pacific Islander 2/, male												
2004.....	8	5	12	19	17	49	60	56	33	11	1	-
2003.....	3	18	7	18	24	51	60	46	25	9	1	-
2002.....	3	5	8	14	18	55	52	43	37	11	2	-
2001.....	6	6	9	14	32	39	54	46	26	11	1	-
2000.....	4	6	9	18	30	58	46	57	23	11	1	-
1999.....	4	8	13	15	29	45	57	57	30	5	1	-
Asian or Pacific Islander 2/, female												
2004.....	6	10	7	10	25	50	64	81	46	22	4	-
2003.....	1	3	4	17	30	46	59	74	43	23	5	-
2002.....	3	5	5	14	29	41	68	58	46	21	4	-
2001.....	3	4	8	20	26	41	65	52	47	15	9	-
2000.....	4	5	9	15	19	45	65	49	47	21	4	-
1999.....	7	4	7	13	23	36	41	41	29	15	3	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of arteries, arterioles and capillaries (I70-I78)												
All races, both sexes												
2004.....	755	1,029	1,537	2,235	3,529	5,124	6,623	6,385	4,566	1,990	486	-
2003.....	795	1,071	1,665	2,434	3,711	5,697	6,834	6,685	4,811	2,137	537	1
2002.....	712	1,135	1,638	2,631	3,979	5,813	6,994	6,858	4,953	2,159	569	1
2001.....	724	1,048	1,620	2,598	4,082	6,122	7,066	7,081	4,996	2,219	557	2
2000.....	719	1,036	1,651	2,826	4,523	6,347	7,152	7,007	5,088	2,275	555	1
1999.....	682	1,049	1,647	2,729	4,555	6,432	7,217	7,154	5,148	2,346	605	-
All races, male												
2004.....	518	674	993	1,372	2,065	2,705	3,166	2,583	1,395	402	64	-
2003.....	549	716	1,099	1,576	2,165	3,187	3,279	2,638	1,430	453	78	1
2002.....	476	751	1,080	1,690	2,345	3,154	3,339	2,763	1,470	472	76	1
2001.....	472	684	1,056	1,630	2,461	3,241	3,389	2,756	1,429	473	77	2
2000.....	495	715	1,109	1,814	2,757	3,534	3,420	2,768	1,472	487	76	1
1999.....	452	711	1,115	1,749	2,727	3,529	3,413	2,667	1,484	488	92	-
All races, female												
2004.....	237	355	544	863	1,464	2,419	3,457	3,802	3,171	1,588	422	-
2003.....	246	355	566	858	1,546	2,510	3,555	4,047	3,381	1,684	459	-
2002.....	236	384	558	941	1,634	2,659	3,655	4,095	3,483	1,687	493	-
2001.....	252	364	564	968	1,621	2,881	3,677	4,325	3,567	1,746	480	-
2000.....	224	321	542	1,012	1,766	2,813	3,732	4,239	3,616	1,788	479	-
1999.....	230	338	532	980	1,828	2,903	3,804	4,487	3,664	1,858	513	-
White, both sexes												
2004.....	566	806	1,255	1,890	3,132	4,594	6,017	5,861	4,213	1,822	425	-
2003.....	587	879	1,390	2,093	3,274	5,149	6,184	6,167	4,449	1,961	467	1
2002.....	517	910	1,364	2,280	3,523	5,226	6,386	6,366	4,571	2,010	491	1
2001.....	546	847	1,328	2,263	3,608	5,496	6,455	6,569	4,632	2,028	483	2
2000.....	526	833	1,367	2,460	4,053	5,730	6,542	6,477	4,692	2,094	485	-
1999.....	528	850	1,420	2,383	4,101	5,824	6,625	6,608	4,748	2,153	532	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of arteries, arterioles and capillaries (I70-I78)												
White, male												
2004.....	401	531	818	1,181	1,875	2,460	2,914	2,398	1,302	375	59	-
2003.....	418	602	927	1,373	1,945	2,901	2,978	2,458	1,334	418	68	1
2002.....	345	604	910	1,504	2,098	2,882	3,082	2,589	1,355	445	64	1
2001.....	354	570	883	1,448	2,210	2,947	3,120	2,588	1,322	438	64	2
2000.....	370	591	946	1,621	2,520	3,233	3,175	2,589	1,351	441	65	-
1999.....	354	593	982	1,567	2,498	3,235	3,157	2,475	1,378	442	74	-
White, female												
2004.....	165	275	437	709	1,257	2,134	3,103	3,463	2,911	1,447	366	-
2003.....	169	277	463	720	1,329	2,248	3,206	3,709	3,115	1,543	399	-
2002.....	172	306	454	776	1,425	2,344	3,304	3,777	3,216	1,565	427	-
2001.....	192	277	445	815	1,398	2,549	3,335	3,981	3,310	1,590	419	-
2000.....	156	242	421	839	1,533	2,497	3,367	3,888	3,341	1,653	420	-
1999.....	174	257	438	816	1,603	2,589	3,468	4,133	3,370	1,711	458	-
Black, both sexes												
2004.....	160	185	251	287	326	445	493	417	299	145	54	-
2003.....	186	164	235	291	367	447	542	419	300	151	59	-
2002.....	171	196	238	291	368	496	515	407	319	128	68	-
2001.....	151	180	257	270	408	531	513	426	318	171	63	-
2000.....	155	179	251	298	396	520	518	450	340	159	61	1
1999.....	131	176	195	299	373	514	496	468	340	173	64	-
Black, male												
2004.....	99	117	156	163	153	201	194	137	73	24	4	-
2003.....	115	99	147	169	177	222	241	134	65	26	8	-
2002.....	116	128	148	148	192	219	209	134	93	22	9	-
2001.....	100	98	146	140	211	235	220	126	84	26	10	-
2000.....	103	106	138	159	197	246	203	145	90	38	9	1
1999.....	81	103	111	157	191	235	200	156	80	43	18	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of arteries, arterioles and capillaries (I70-I78)												
Black, female												
2004.....	61	68	95	124	173	244	299	280	226	121	50	-
2003.....	71	65	88	122	190	225	301	285	235	125	51	-
2002.....	55	68	90	143	176	277	306	273	226	106	59	-
2001.....	51	82	111	130	197	296	293	300	234	145	53	-
2000.....	52	73	113	139	199	274	315	305	250	121	52	-
1999.....	50	73	84	142	182	279	296	312	260	130	46	-
American Indian 1/, both sexes												
2004.....	5	10	8	11	14	20	17	18	5	3	1	-
2003.....	7	6	14	13	14	21	13	17	9	5	3	-
2002.....	10	9	14	24	23	19	18	11	9	3	-	-
2001.....	5	5	10	16	13	14	25	19	7	3	1	-
2000.....	5	8	3	28	21	15	18	12	10	7	3	-
1999.....	4	6	8	13	18	15	16	17	8	4	4	-
American Indian 1/, male												
2004.....	2	7	5	2	4	12	6	7	2	1	-	-
2003.....	6	2	7	11	8	10	6	8	2	2	2	-
2002.....	5	4	8	13	15	9	6	5	1	1	-	-
2001.....	4	4	8	9	7	6	9	8	4	-	-	-
2000.....	2	6	1	12	8	7	9	5	1	1	1	-
1999.....	3	4	5	6	9	7	7	7	-	1	-	-
American Indian 1/, female												
2004.....	3	3	3	9	10	8	11	11	3	2	1	-
2003.....	1	4	7	2	6	11	7	9	7	3	1	-
2002.....	5	5	6	11	8	10	12	6	8	2	-	-
2001.....	1	1	2	7	6	8	16	11	3	3	1	-
2000.....	3	2	2	16	13	8	9	7	9	6	2	-
1999.....	1	2	3	7	9	8	9	10	8	3	4	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of arteries, arterioles and capillaries (I70-I78)												
Asian or Pacific Islander 2/, both sexes												
2004.....	24	28	23	47	57	65	96	89	49	20	6	-
2003.....	15	22	26	37	56	80	95	82	53	20	8	-
2002.....	14	20	22	36	65	72	75	74	54	18	10	-
2001.....	22	16	25	49	53	81	73	67	39	17	10	-
2000.....	33	16	30	40	53	82	74	68	46	15	6	-
1999.....	19	17	24	34	63	79	80	61	52	16	5	-
Asian or Pacific Islander 2/, male												
2004.....	16	19	14	26	33	32	52	41	18	2	1	-
2003.....	10	13	18	23	35	54	54	38	29	7	-	-
2002.....	10	15	14	25	40	44	42	35	21	4	3	-
2001.....	14	12	19	33	33	53	40	34	19	9	3	-
2000.....	20	12	24	22	32	48	33	29	30	7	1	-
1999.....	14	11	17	19	29	52	49	29	26	2	-	-
Asian or Pacific Islander 2/, female												
2004.....	8	9	9	21	24	33	44	48	31	18	5	-
2003.....	5	9	8	14	21	26	41	44	24	13	8	-
2002.....	4	5	8	11	25	28	33	39	33	14	7	-
2001.....	8	4	6	16	20	28	33	33	20	8	7	-
2000.....	13	4	6	18	21	34	41	39	16	8	5	-
1999.....	5	6	7	15	34	27	31	32	26	14	5	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerosis (I70)												
All races, both sexes												
2004.....	93	170	289	409	723	1,294	2,112	2,682	2,382	1,246	349	-
2003.....	120	171	295	436	775	1,482	2,355	2,946	2,582	1,375	385	1
2002.....	121	207	280	494	806	1,521	2,541	3,090	2,802	1,416	415	1
2001.....	121	198	275	471	864	1,624	2,532	3,214	2,786	1,471	417	1
2000.....	105	170	281	495	928	1,685	2,592	3,188	2,870	1,518	429	-
1999.....	129	172	314	485	991	1,770	2,615	3,362	2,987	1,581	460	-
All races, male												
2004.....	62	98	177	229	390	626	923	996	652	240	45	-
2003.....	77	113	185	245	404	742	992	1,009	668	272	53	1
2002.....	75	127	178	282	446	748	1,055	1,088	755	281	53	1
2001.....	75	118	168	273	470	765	1,026	1,123	710	285	54	1
2000.....	71	108	186	290	510	838	1,097	1,090	738	283	55	-
1999.....	84	114	193	277	518	845	1,082	1,100	773	316	66	-
All races, female												
2004.....	31	72	112	180	333	668	1,189	1,686	1,730	1,006	304	-
2003.....	43	58	110	191	371	740	1,363	1,937	1,914	1,103	332	-
2002.....	46	80	102	212	360	773	1,486	2,002	2,047	1,135	362	-
2001.....	46	80	107	198	394	859	1,506	2,091	2,076	1,186	363	-
2000.....	34	62	95	205	418	847	1,495	2,098	2,132	1,235	374	-
1999.....	45	58	121	208	473	925	1,533	2,262	2,214	1,265	394	-
White, both sexes												
2004.....	73	138	248	343	632	1,171	1,963	2,508	2,233	1,166	312	-
2003.....	90	139	246	365	666	1,338	2,146	2,755	2,405	1,287	342	1
2002.....	85	170	227	435	697	1,346	2,320	2,915	2,637	1,331	371	1
2001.....	97	166	216	406	760	1,475	2,339	3,034	2,624	1,374	377	1
2000.....	73	140	226	431	815	1,535	2,375	2,984	2,693	1,431	383	-
1999.....	103	140	262	413	887	1,596	2,423	3,139	2,796	1,475	413	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerosis (I70)												
White, male												
2004.....	53	81	151	197	348	568	857	946	614	228	41	-
2003.....	59	93	156	206	343	663	899	943	625	257	47	1
2002.....	50	106	144	251	389	663	967	1,030	710	266	46	1
2001.....	59	101	136	237	418	692	946	1,067	666	268	44	1
2000.....	50	93	154	252	461	775	1,008	1,015	687	261	48	-
1999.....	70	90	164	238	463	755	1,000	1,022	722	291	56	-
White, female												
2004.....	20	57	97	146	284	603	1,106	1,562	1,619	938	271	-
2003.....	31	46	90	159	323	675	1,247	1,812	1,780	1,030	295	-
2002.....	35	64	83	184	308	683	1,353	1,885	1,927	1,065	325	-
2001.....	38	65	80	169	342	783	1,393	1,967	1,958	1,106	333	-
2000.....	23	47	72	179	354	760	1,367	1,969	2,006	1,170	335	-
1999.....	33	50	98	175	424	841	1,423	2,117	2,074	1,184	357	-
Black, both sexes												
2004.....	18	28	39	52	76	109	129	138	125	68	34	-
2003.....	28	30	42	62	96	122	179	153	149	78	35	-
2002.....	31	29	48	49	86	156	196	151	137	68	38	-
2001.....	20	28	56	58	90	132	158	151	139	87	32	-
2000.....	28	28	51	51	97	132	191	173	153	76	40	-
1999.....	23	27	50	63	88	154	162	190	162	92	40	-
Black, male												
2004.....	8	13	26	25	37	51	57	35	30	10	4	-
2003.....	17	19	25	33	55	63	80	50	33	12	5	-
2002.....	22	17	32	25	46	76	78	49	36	13	5	-
2001.....	13	14	29	30	45	65	61	45	35	13	7	-
2000.....	17	14	29	33	42	54	79	63	41	19	5	-
1999.....	11	20	27	34	46	78	65	68	39	23	10	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerosis (I70)												
Black, female												
2004.....	10	15	13	27	39	58	72	103	95	58	30	-
2003.....	11	11	17	29	41	59	99	103	116	66	30	-
2002.....	9	12	16	24	40	80	118	102	101	55	33	-
2001.....	7	14	27	28	45	67	97	106	104	74	25	-
2000.....	11	14	22	18	55	78	112	110	112	57	35	-
1999.....	12	7	23	29	42	76	97	122	123	69	30	-
American Indian 1/, both sexes												
2004.....	-	3	-	4	6	4	6	8	2	2	-	-
2003.....	2	1	4	2	6	2	4	3	5	1	2	-
2002.....	4	4	3	8	9	7	9	2	4	2	-	-
2001.....	2	3	1	2	6	4	10	8	6	1	1	-
2000.....	-	2	1	8	8	3	10	8	3	4	2	-
1999.....	2	-	2	5	6	4	8	5	5	3	3	-
American Indian 1/, male												
2004.....	-	3	-	1	2	4	3	2	-	1	-	-
2003.....	1	-	2	2	3	1	3	2	-	-	1	-
2002.....	2	2	2	4	3	1	4	1	1	-	-	-
2001.....	2	2	1	2	3	1	5	2	3	-	-	-
2000.....	-	1	1	3	3	1	5	3	-	-	1	-
1999.....	2	-	2	3	3	3	3	1	-	1	-	-
American Indian 1/, female												
2004.....	-	-	-	3	4	-	3	6	2	1	-	-
2003.....	1	1	2	-	3	1	1	1	5	1	1	-
2002.....	2	2	1	4	6	6	5	1	3	2	-	-
2001.....	-	1	-	-	3	3	5	6	3	1	1	-
2000.....	-	1	-	5	5	2	5	5	3	4	1	-
1999.....	-	-	-	2	3	1	5	4	5	2	3	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerosis (I70)												
Asian or Pacific Islander 2/, both sexes												
2004.....	2	1	2	10	9	10	14	28	22	10	3	-
2003.....	-	1	3	7	7	20	26	35	23	9	6	-
2002.....	1	4	2	2	14	12	16	22	24	15	6	-
2001.....	2	1	2	5	8	13	25	21	17	9	7	-
2000.....	4	-	3	5	8	15	16	23	21	7	4	-
1999.....	1	5	-	4	10	16	22	28	24	11	4	-
Asian or Pacific Islander 2/, male												
2004.....	1	1	-	6	3	3	6	13	8	1	-	-
2003.....	-	1	2	4	3	15	10	14	10	3	-	-
2002.....	1	2	-	2	8	8	6	8	8	2	2	-
2001.....	1	1	2	4	4	7	14	9	6	4	3	-
2000.....	4	-	2	2	4	8	5	9	10	3	1	-
1999.....	1	4	-	2	6	9	14	9	12	1	-	-
Asian or Pacific Islander 2/, female												
2004.....	1	-	2	4	6	7	8	15	14	9	3	-
2003.....	-	-	1	3	4	5	16	21	13	6	6	-
2002.....	-	2	2	-	6	4	10	14	16	13	4	-
2001.....	1	-	-	1	4	6	11	12	11	5	4	-
2000.....	-	-	1	3	4	7	11	14	11	4	3	-
1999.....	-	1	-	2	4	7	8	19	12	10	4	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Aortic aneurysm and dissection (I71)												
All races, both sexes												
2004.....	423	516	802	1,137	1,795	2,416	2,715	1,949	950	224	24	-
2003.....	452	590	887	1,317	1,954	2,726	2,745	2,059	965	245	29	-
2002.....	383	598	862	1,404	2,078	2,779	2,739	2,013	914	239	21	-
2001.....	374	532	862	1,430	2,087	2,899	2,852	2,122	1,022	242	26	1
2000.....	394	585	908	1,552	2,460	3,043	2,792	2,134	985	213	25	-
1999.....	355	570	867	1,578	2,423	3,095	2,841	2,118	935	257	27	-
All races, male												
2004.....	319	390	582	786	1,160	1,369	1,490	932	423	73	4	-
2003.....	347	448	659	912	1,262	1,667	1,525	1,033	411	74	7	-
2002.....	291	458	634	1,020	1,349	1,688	1,497	1,034	368	74	5	-
2001.....	287	398	633	981	1,401	1,722	1,591	1,000	406	78	7	1
2000.....	301	438	673	1,097	1,633	1,885	1,558	1,063	425	68	6	-
1999.....	270	430	658	1,114	1,598	1,905	1,596	1,008	398	83	6	-
All races, female												
2004.....	104	126	220	351	635	1,047	1,225	1,017	527	151	20	-
2003.....	105	142	228	405	692	1,059	1,220	1,026	554	171	22	-
2002.....	92	140	228	384	729	1,091	1,242	979	546	165	16	-
2001.....	87	134	229	449	686	1,177	1,261	1,122	616	164	19	-
2000.....	93	147	235	455	827	1,158	1,234	1,071	560	145	19	-
1999.....	85	140	209	464	825	1,190	1,245	1,110	537	174	21	-
White, both sexes												
2004.....	315	396	664	1,002	1,635	2,212	2,486	1,798	898	210	20	-
2003.....	334	482	761	1,176	1,773	2,521	2,542	1,921	915	219	25	-
2002.....	272	482	749	1,261	1,891	2,580	2,548	1,877	843	219	16	-
2001.....	270	431	736	1,276	1,905	2,639	2,654	1,987	943	221	24	1
2000.....	303	475	782	1,381	2,270	2,795	2,603	1,988	900	188	23	-
1999.....	273	473	777	1,420	2,232	2,871	2,658	1,971	871	233	22	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Aortic aneurysm and dissection (I71)												
White, male												
2004.....	244	307	482	698	1,072	1,272	1,380	865	405	70	3	-
2003.....	266	379	570	827	1,170	1,561	1,421	977	388	67	6	-
2002.....	210	370	556	936	1,238	1,585	1,415	985	345	72	4	-
2001.....	212	333	544	892	1,293	1,588	1,499	950	385	73	7	1
2000.....	237	364	592	1,006	1,527	1,747	1,480	1,015	389	62	6	-
1999.....	208	369	600	1,025	1,500	1,805	1,506	946	376	74	5	-
White, female												
2004.....	71	89	182	304	563	940	1,106	933	493	140	17	-
2003.....	68	103	191	349	603	960	1,121	944	527	152	19	-
2002.....	62	112	193	325	653	995	1,133	892	498	147	12	-
2001.....	58	98	192	384	612	1,051	1,155	1,037	558	148	17	-
2000.....	66	111	190	375	743	1,048	1,123	973	511	126	17	-
1999.....	65	104	177	395	732	1,066	1,152	1,025	495	159	17	-
Black, both sexes												
2004.....	90	97	122	112	114	154	164	105	35	13	2	-
2003.....	102	90	107	112	135	152	142	99	36	22	2	-
2002.....	99	102	95	109	137	145	143	96	49	20	5	-
2001.....	88	88	104	107	137	198	157	98	64	16	2	-
2000.....	69	98	102	129	152	190	143	114	64	20	1	-
1999.....	71	85	70	128	146	166	140	118	44	22	5	-
Black, male												
2004.....	61	67	85	74	57	65	67	44	11	2	-	-
2003.....	70	57	76	66	62	72	67	33	11	5	-	-
2002.....	74	76	66	59	74	68	54	28	15	2	1	-
2001.....	64	55	72	59	79	93	68	28	13	2	-	-
2000.....	52	63	62	68	83	101	54	35	21	4	-	-
1999.....	52	53	44	74	75	63	62	44	12	8	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Aortic aneurysm and dissection (I71)

Black, female

2004.....	29	30	37	38	57	89	97	61	24	11	2	-
2003.....	32	33	31	46	73	80	75	66	25	17	2	-
2002.....	25	26	29	50	63	77	89	68	34	18	4	-
2001.....	24	33	32	48	58	105	89	70	51	14	2	-
2000.....	17	35	40	61	69	89	89	79	43	16	1	-
1999.....	19	32	26	54	71	103	78	74	32	14	4	-

American Indian 1/, both sexes

2004.....	1	2	3	2	4	10	5	4	1	-	1	-
2003.....	2	2	4	4	4	12	4	5	-	1	1	-
2002.....	2	1	5	10	9	6	7	3	2	-	-	-
2001.....	1	2	6	9	3	7	8	6	-	1	-	-
2000.....	-	1	2	13	8	9	3	-	3	1	-	-
1999.....	-	3	2	4	8	8	3	6	1	-	-	-

American Indian 1/, male

2004.....	1	2	3	-	2	6	3	3	-	-	-	-
2003.....	2	1	3	3	2	6	1	2	-	1	1	-
2002.....	1	-	2	7	8	5	2	2	-	-	-	-
2001.....	1	2	4	4	3	5	3	4	-	-	-	-
2000.....	-	1	-	7	4	5	2	-	1	-	-	-
1999.....	-	2	1	1	5	4	2	3	-	-	-	-

American Indian 1/, female

2004.....	-	-	-	2	2	4	2	1	1	-	1	-
2003.....	-	1	1	1	2	6	3	3	-	-	-	-
2002.....	1	1	3	3	1	1	5	1	2	-	-	-
2001.....	-	-	2	5	-	2	5	2	-	1	-	-
2000.....	-	-	2	6	4	4	1	-	2	1	-	-
1999.....	-	1	1	3	3	4	1	3	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Aortic aneurysm and dissection (I71)												
Asian or Pacific Islander 2/, both sexes												
2004.....	17	21	13	21	42	40	60	42	16	1	1	-
2003.....	14	16	15	25	42	41	57	34	14	3	1	-
2002.....	10	13	13	24	41	48	41	37	20	-	-	-
2001.....	15	11	16	38	42	55	33	31	15	4	-	-
2000.....	22	11	22	29	30	49	43	32	18	4	1	-
1999.....	11	9	18	26	37	50	40	23	19	2	-	-
Asian or Pacific Islander 2/, male												
2004.....	13	14	12	14	29	26	40	20	7	1	1	-
2003.....	9	11	10	16	28	28	36	21	12	1	-	-
2002.....	6	12	10	18	29	30	26	19	8	-	-	-
2001.....	10	8	13	26	26	36	21	18	8	3	-	-
2000.....	12	10	19	16	19	32	22	13	14	2	-	-
1999.....	10	6	13	14	18	33	26	15	10	1	-	-
Asian or Pacific Islander 2/, female												
2004.....	4	7	1	7	13	14	20	22	9	-	-	-
2003.....	5	5	5	9	14	13	21	13	2	2	1	-
2002.....	4	1	3	6	12	18	15	18	12	-	-	-
2001.....	5	3	3	12	16	19	12	13	7	1	-	-
2000.....	10	1	3	13	11	17	21	19	4	2	1	-
1999.....	1	3	5	12	19	17	14	8	9	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)												
All races, both sexes												
2004.....	239	343	446	689	1,011	1,414	1,796	1,754	1,234	520	113	-
2003.....	223	310	483	681	982	1,489	1,734	1,680	1,264	517	123	-
2002.....	208	330	496	733	1,095	1,513	1,714	1,755	1,237	504	133	-
2001.....	229	318	483	697	1,131	1,599	1,682	1,745	1,188	506	114	-
2000.....	220	281	462	779	1,135	1,619	1,768	1,685	1,233	544	101	1
1999.....	198	307	466	666	1,141	1,567	1,761	1,674	1,226	508	118	-
All races, male												
2004.....	137	186	234	357	515	710	753	655	320	89	15	-
2003.....	125	155	255	419	499	778	762	596	351	107	18	-
2002.....	110	166	268	388	550	718	787	641	347	117	18	-
2001.....	110	168	255	376	590	754	772	633	313	110	16	-
2000.....	123	169	250	427	614	811	765	615	309	136	15	1
1999.....	98	167	264	358	611	779	735	559	313	89	20	-
All races, female												
2004.....	102	157	212	332	496	704	1,043	1,099	914	431	98	-
2003.....	98	155	228	262	483	711	972	1,084	913	410	105	-
2002.....	98	164	228	345	545	795	927	1,114	890	387	115	-
2001.....	119	150	228	321	541	845	910	1,112	875	396	98	-
2000.....	97	112	212	352	521	808	1,003	1,070	924	408	86	-
1999.....	100	140	202	308	530	788	1,026	1,115	913	419	98	-
White, both sexes												
2004.....	178	272	343	545	865	1,211	1,568	1,555	1,082	446	93	-
2003.....	163	258	383	552	835	1,290	1,496	1,491	1,129	455	100	-
2002.....	160	258	388	584	935	1,300	1,518	1,574	1,091	460	104	-
2001.....	179	250	376	581	943	1,382	1,462	1,548	1,065	433	82	-
2000.....	150	218	359	648	968	1,400	1,564	1,505	1,099	475	79	-
1999.....	152	237	381	550	982	1,357	1,544	1,498	1,081	445	97	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)												
White, male												
2004.....	104	143	185	286	455	620	677	587	283	77	15	-
2003.....	93	130	201	340	432	677	658	538	321	94	15	-
2002.....	85	128	210	317	471	634	700	574	300	107	14	-
2001.....	83	136	203	319	499	667	675	571	271	97	13	-
2000.....	83	134	200	363	532	711	687	559	275	118	11	-
1999.....	76	134	218	304	535	675	651	507	280	77	13	-
White, female												
2004.....	74	129	158	259	410	591	891	968	799	369	78	-
2003.....	70	128	182	212	403	613	838	953	808	361	85	-
2002.....	75	130	178	267	464	666	818	1,000	791	353	90	-
2001.....	96	114	173	262	444	715	787	977	794	336	69	-
2000.....	67	84	159	285	436	689	877	946	824	357	68	-
1999.....	76	103	163	246	447	682	893	991	801	368	84	-
Black, both sexes												
2004.....	52	60	90	123	136	182	200	174	139	64	18	-
2003.....	56	44	86	117	136	173	221	167	115	51	22	-
2002.....	41	65	95	133	145	195	176	160	133	40	25	-
2001.....	43	64	97	105	181	201	198	177	115	68	29	-
2000.....	58	53	98	118	147	198	184	163	123	63	20	1
1999.....	37	64	75	108	139	194	194	160	134	59	19	-
Black, male												
2004.....	30	37	45	64	59	85	70	58	32	12	-	-
2003.....	28	23	46	70	60	87	94	51	21	9	3	-
2002.....	20	35	50	64	72	75	77	57	42	7	3	-
2001.....	23	29	45	51	87	77	91	53	36	11	3	-
2000.....	34	29	47	58	72	91	70	47	28	15	4	1
1999.....	18	30	40	49	70	94	73	44	29	12	7	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other diseases of arteries, arterioles and capillaries (I72-I78)

Black, female

2004.....	22	23	45	59	77	97	130	116	107	52	18	-
2003.....	28	21	40	47	76	86	127	116	94	42	19	-
2002.....	21	30	45	69	73	120	99	103	91	33	22	-
2001.....	20	35	52	54	94	124	107	124	79	57	26	-
2000.....	24	24	51	60	75	107	114	116	95	48	16	-
1999.....	19	34	35	59	69	100	121	116	105	47	12	-

American Indian 1/, both sexes

2004.....	4	5	5	5	4	6	6	6	2	1	-	-
2003.....	3	3	6	7	4	7	5	9	4	3	-	-
2002.....	4	4	6	6	5	6	2	6	3	1	-	-
2001.....	2	-	3	5	4	3	7	5	1	1	-	-
2000.....	5	5	-	7	5	3	5	4	4	2	1	-
1999.....	2	3	4	4	4	3	5	6	2	1	1	-

American Indian 1/, male

2004.....	1	2	2	1	-	2	-	2	2	-	-	-
2003.....	3	1	2	6	3	3	2	4	2	1	-	-
2002.....	2	2	4	2	4	3	-	2	-	1	-	-
2001.....	1	-	3	3	1	-	1	2	1	-	-	-
2000.....	2	4	-	2	1	1	2	2	-	1	-	-
1999.....	1	2	2	2	1	-	2	3	-	-	-	-

American Indian 1/, female

2004.....	3	3	3	4	4	4	6	4	-	1	-	-
2003.....	-	2	4	1	1	4	3	5	2	2	-	-
2002.....	2	2	2	4	1	3	2	4	3	-	-	-
2001.....	1	-	-	2	3	3	6	3	-	1	-	-
2000.....	3	1	-	5	4	2	3	2	4	1	1	-
1999.....	1	1	2	2	3	3	3	3	2	1	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other diseases of arteries, arterioles and capillaries (I72-I78)

Asian or Pacific Islander 2/, both sexes

2004.....	5	6	8	16	6	15	22	19	11	9	2	-
2003.....	1	5	8	5	7	19	12	13	16	8	1	-
2002.....	3	3	7	10	10	12	18	15	10	3	4	-
2001.....	5	4	7	6	3	13	15	15	7	4	3	-
2000.....	7	5	5	6	15	18	15	13	7	4	1	-
1999.....	7	3	6	4	16	13	18	10	9	3	1	-

Asian or Pacific Islander 2/, male

2004.....	2	4	2	6	1	3	6	8	3	-	-	-
2003.....	1	1	6	3	4	11	8	3	7	3	-	-
2002.....	3	1	4	5	3	6	10	8	5	2	1	-
2001.....	3	3	4	3	3	10	5	7	5	2	-	-
2000.....	4	2	3	4	9	8	6	7	6	2	-	-
1999.....	3	1	4	3	5	10	9	5	4	-	-	-

Asian or Pacific Islander 2/, female

2004.....	3	2	6	10	5	12	16	11	8	9	2	-
2003.....	-	4	2	2	3	8	4	10	9	5	1	-
2002.....	-	2	3	5	7	6	8	7	5	1	3	-
2001.....	2	1	3	3	-	3	10	8	2	2	3	-
2000.....	3	3	2	2	6	10	9	6	1	2	1	-
1999.....	4	2	2	1	11	3	9	5	5	3	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of circulatory system (I80-I99)												
All races, both sexes												
2004.....	277	314	309	309	363	528	637	566	392	149	26	-
2003.....	282	272	297	319	425	518	643	577	382	148	26	-
2002.....	280	302	293	322	375	552	701	576	390	134	23	-
2001.....	232	272	291	366	475	572	611	568	414	140	27	1
2000.....	266	231	285	326	469	573	588	571	343	149	21	-
1999.....	176	208	237	312	380	498	619	509	322	116	25	1
All races, male												
2004.....	172	174	177	154	174	246	259	171	96	27	2	-
2003.....	165	166	154	160	188	214	223	163	82	27	4	-
2002.....	147	168	141	160	163	216	267	173	86	25	1	-
2001.....	135	149	155	174	231	254	234	164	97	19	6	1
2000.....	150	126	138	170	220	245	227	175	72	26	-	-
1999.....	96	106	118	163	187	225	238	155	66	19	8	-
All races, female												
2004.....	105	140	132	155	189	282	378	395	296	122	24	-
2003.....	117	106	143	159	237	304	420	414	300	121	22	-
2002.....	133	134	152	162	212	336	434	403	304	109	22	-
2001.....	97	123	136	192	244	318	377	404	317	121	21	-
2000.....	116	105	147	156	249	328	361	396	271	123	21	-
1999.....	80	102	119	149	193	273	381	354	256	97	17	1
White, both sexes												
2004.....	194	246	244	254	291	447	566	511	350	131	24	-
2003.....	221	212	230	255	350	429	564	514	339	130	21	-
2002.....	214	238	219	260	315	472	609	515	354	120	21	-
2001.....	177	213	219	294	398	480	551	519	369	127	23	-
2000.....	203	175	222	242	395	488	529	516	309	132	21	-
1999.....	131	177	190	243	325	429	549	451	290	107	24	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other disorders of circulatory system (I80-I99)

White, male

2004.....	120	144	141	125	145	212	227	159	84	23	2	-
2003.....	127	139	113	126	156	178	189	143	73	23	3	-
2002.....	115	130	109	127	137	186	226	154	74	21	1	-
2001.....	102	118	113	140	193	215	211	149	87	15	4	-
2000.....	117	94	107	131	192	214	205	158	64	22	-	-
1999.....	69	92	100	124	163	191	210	138	59	18	8	-

White, female

2004.....	74	102	103	129	146	235	339	352	266	108	22	-
2003.....	94	73	117	129	194	251	375	371	266	107	18	-
2002.....	99	108	110	133	178	286	383	361	280	99	20	-
2001.....	75	95	106	154	205	265	340	370	282	112	19	-
2000.....	86	81	115	111	203	274	324	358	245	110	21	-
1999.....	62	85	90	119	162	238	339	313	231	89	16	1

Black, both sexes

2004.....	80	61	61	47	68	76	59	45	39	16	2	-
2003.....	55	57	62	59	70	81	70	53	41	16	5	-
2002.....	61	62	68	60	54	75	85	54	33	12	2	-
2001.....	48	55	68	64	67	81	55	42	40	13	3	1
2000.....	60	53	56	79	68	82	50	51	30	15	-	-
1999.....	41	27	42	62	47	64	61	51	29	8	1	-

Black, male

2004.....	51	27	33	24	27	31	23	10	12	3	-	-
2003.....	34	26	38	31	31	34	28	18	8	4	1	-
2002.....	28	36	30	31	23	27	37	15	11	4	-	-
2001.....	29	30	42	31	31	35	19	12	8	4	2	1
2000.....	32	29	27	36	26	28	16	14	8	4	-	-
1999.....	24	11	17	36	20	31	24	13	5	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other disorders of circulatory system (I80-I99)

Black, female

2004.....	29	34	28	23	41	45	36	35	27	13	2	-
2003.....	21	31	24	28	39	47	42	35	33	12	4	-
2002.....	33	26	38	29	31	48	48	39	22	8	2	-
2001.....	19	25	26	33	36	46	36	30	32	9	1	-
2000.....	28	24	29	43	42	54	34	37	22	11	-	-
1999.....	17	16	25	26	27	33	37	38	24	7	1	-

American Indian 1/, both sexes

2004.....	1	3	1	3	-	3	-	1	-	2	-	-
2003.....	2	2	1	1	-	3	1	1	-	1	-	-
2002.....	2	2	3	1	2	-	3	2	1	1	-	-
2001.....	4	2	1	2	4	2	-	1	1	-	-	-
2000.....	1	-	2	3	4	1	3	-	1	-	-	-
1999.....	2	-	2	1	-	1	5	-	-	-	-	-

American Indian 1/, male

2004.....	-	1	1	2	-	2	-	-	-	1	-	-
2003.....	2	-	1	-	-	1	1	-	-	-	-	-
2002.....	2	2	-	1	2	-	2	1	-	-	-	-
2001.....	3	-	-	1	4	-	-	1	-	-	-	-
2000.....	-	-	1	2	1	1	2	-	-	-	-	-
1999.....	2	-	-	1	-	-	1	-	-	-	-	-

American Indian 1/, female

2004.....	1	2	-	1	-	1	-	1	-	1	-	-
2003.....	-	2	-	1	-	2	-	1	-	1	-	-
2002.....	-	-	3	-	-	-	1	1	1	1	-	-
2001.....	1	2	1	1	-	2	-	-	1	-	-	-
2000.....	1	-	1	1	3	-	1	-	1	-	-	-
1999.....	-	-	2	-	-	1	4	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of circulatory system (I80-I99)												
Asian or Pacific Islander 2/, both sexes												
2004.....	2	4	3	5	4	2	12	9	3	-	-	-
2003.....	4	1	4	4	5	5	8	9	2	1	-	-
2002.....	3	-	3	1	4	5	4	5	2	1	-	-
2001.....	3	2	3	6	6	9	5	6	4	-	1	-
2000.....	2	3	5	2	2	2	6	4	3	2	-	-
1999.....	2	4	3	6	8	4	4	7	3	1	-	-
Asian or Pacific Islander 2/, male												
2004.....	1	2	2	3	2	1	9	2	-	-	-	-
2003.....	2	1	2	3	1	1	5	2	1	-	-	-
2002.....	2	-	2	1	1	3	2	3	1	-	-	-
2001.....	1	1	-	2	3	4	4	2	2	-	-	-
2000.....	1	3	3	1	1	2	4	3	-	-	-	-
1999.....	1	3	1	2	4	3	3	4	2	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	1	2	1	2	2	1	3	7	3	-	-	-
2003.....	2	-	2	1	4	4	3	7	1	1	-	-
2002.....	1	-	1	-	3	2	2	2	1	1	-	-
2001.....	2	1	3	4	3	5	1	4	2	-	1	-
2000.....	1	-	2	1	1	-	2	1	3	2	-	-
1999.....	1	1	2	4	4	1	1	3	1	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
All races, both sexes												
2004.....	193	228	222	215	264	401	488	424	295	101	19	-
2003.....	207	204	224	254	321	384	490	436	284	122	13	-
2002.....	217	239	217	251	268	399	536	452	297	90	15	-
2001.....	167	202	225	269	356	440	457	430	306	104	23	1
2000.....	191	171	192	242	349	431	435	427	245	107	15	-
1999.....	116	140	150	206	245	314	391	324	195	73	14	1
All races, male												
2004.....	112	113	122	101	124	176	190	123	74	16	2	-
2003.....	111	116	117	123	134	148	161	127	51	20	1	-
2002.....	104	126	95	118	113	144	192	133	62	10	1	-
2001.....	91	104	112	120	172	181	160	113	65	15	5	1
2000.....	96	84	82	125	149	173	163	115	50	19	-	-
1999.....	52	63	63	102	119	132	143	81	35	14	4	-
All races, female												
2004.....	81	115	100	114	140	225	298	301	221	85	17	-
2003.....	96	88	107	131	187	236	329	309	233	102	12	-
2002.....	113	113	122	133	155	255	344	319	235	80	14	-
2001.....	76	98	113	149	184	259	297	317	241	89	18	-
2000.....	95	87	110	117	200	258	272	312	195	88	15	-
1999.....	64	77	87	104	126	182	248	243	160	59	10	1
White, both sexes												
2004.....	135	175	166	179	209	338	434	386	259	89	17	-
2003.....	163	156	168	206	263	316	431	391	251	107	8	-
2002.....	162	190	163	204	223	343	461	404	267	82	13	-
2001.....	129	158	164	216	294	372	412	393	273	93	21	-
2000.....	142	131	147	177	292	364	394	388	223	95	15	-
1999.....	87	117	121	162	209	273	345	286	173	67	14	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
White, male												
2004.....	78	93	93	84	100	152	167	115	64	13	2	-
2003.....	86	98	83	98	110	120	139	112	44	18	-	-
2002.....	79	96	72	94	96	129	160	117	54	9	1	-
2001.....	67	82	78	100	143	156	146	102	59	12	4	-
2000.....	75	64	62	98	126	153	150	104	45	16	-	-
1999.....	38	53	52	76	105	114	127	73	33	13	4	-
White, female												
2004.....	57	82	73	95	109	186	267	271	195	76	15	-
2003.....	77	58	85	108	153	196	292	279	207	89	8	-
2002.....	83	94	91	110	127	214	301	287	213	73	12	-
2001.....	62	76	86	116	151	216	266	291	214	81	17	-
2000.....	67	67	85	79	166	211	244	284	178	79	15	-
1999.....	49	64	69	86	104	159	218	213	140	54	10	1
Black, both sexes												
2004.....	57	50	53	31	52	60	45	33	34	12	2	-
2003.....	40	46	53	46	56	62	54	39	33	13	5	-
2002.....	52	47	51	46	42	54	70	43	28	7	2	-
2001.....	35	41	59	49	58	60	43	32	31	11	2	1
2000.....	47	39	43	62	53	66	33	38	21	11	-	-
1999.....	27	21	26	43	33	40	44	34	20	5	-	-
Black, male												
2004.....	34	20	27	13	23	22	17	7	10	3	-	-
2003.....	23	17	32	24	23	26	19	14	7	2	1	-
2002.....	23	28	22	23	16	14	29	13	7	1	-	-
2001.....	22	21	34	20	26	24	13	10	5	3	1	1
2000.....	21	19	19	25	21	19	8	10	5	3	-	-
1999.....	13	9	11	26	13	18	15	7	1	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Asian or Pacific Islander 2/, both sexes												
2004.....	-	2	3	4	3	-	9	5	2	-	-	-
2003.....	3	1	3	1	2	4	4	5	-	1	-	-
2002.....	2	-	1	1	2	2	4	3	1	-	-	-
2001.....	2	2	1	4	1	6	2	4	2	-	-	-
2000.....	1	1	2	2	1	-	5	1	-	1	-	-
1999.....	2	2	2	1	3	1	1	4	2	1	-	-
Asian or Pacific Islander 2/, male												
2004.....	-	-	2	3	1	-	6	1	-	-	-	-
2003.....	1	1	2	1	1	1	2	1	-	-	-	-
2002.....	1	-	1	1	-	1	2	2	1	-	-	-
2001.....	1	1	-	-	-	1	1	-	1	-	-	-
2000.....	-	1	1	1	1	-	3	1	-	-	-	-
1999.....	1	1	-	-	1	-	-	1	1	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	-	2	1	1	2	-	3	4	2	-	-	-
2003.....	2	-	1	-	1	3	2	4	-	1	-	-
2002.....	1	-	-	-	2	1	2	1	-	-	-	-
2001.....	1	1	1	4	1	5	1	4	1	-	-	-
2000.....	1	-	1	1	-	-	2	-	-	1	-	-
1999.....	1	1	2	1	2	1	1	3	1	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

All other and unspecified disorders of circulatory system (I83-I99)

All races, both sexes

2004.....	84	86	87	94	99	127	149	142	97	48	7	-
2003.....	75	68	73	65	104	134	153	141	98	26	13	-
2002.....	63	63	76	71	107	153	165	124	93	44	8	-
2001.....	65	70	66	97	119	132	154	138	108	36	4	-
2000.....	75	60	93	84	120	142	153	144	98	42	6	-
1999.....	60	68	87	106	135	184	228	185	127	43	11	-

All races, male

2004.....	60	61	55	53	50	70	69	48	22	11	-	-
2003.....	54	50	37	37	54	66	62	36	31	7	3	-
2002.....	43	42	46	42	50	72	75	40	24	15	-	-
2001.....	44	45	43	54	59	73	74	51	32	4	1	-
2000.....	54	42	56	45	71	72	64	60	22	7	-	-
1999.....	44	43	55	61	68	93	95	74	31	5	4	-

All races, female

2004.....	24	25	32	41	49	57	80	94	75	37	7	-
2003.....	21	18	36	28	50	68	91	105	67	19	10	-
2002.....	20	21	30	29	57	81	90	84	69	29	8	-
2001.....	21	25	23	43	60	59	80	87	76	32	3	-
2000.....	21	18	37	39	49	70	89	84	76	35	6	-
1999.....	16	25	32	45	67	91	133	111	96	38	7	-

White, both sexes

2004.....	59	71	78	75	82	109	132	125	91	42	7	-
2003.....	58	56	62	49	87	113	133	123	88	23	13	-
2002.....	52	48	56	56	92	129	148	111	87	38	8	-
2001.....	48	55	55	78	104	108	139	126	96	34	2	-
2000.....	61	44	75	65	103	124	135	128	86	37	6	-
1999.....	44	60	69	81	116	156	204	165	117	40	10	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

All other and unspecified disorders of circulatory system (I83-I99)

White, male

2004.....	42	51	48	41	45	60	60	44	20	10	-	-
2003.....	41	41	30	28	46	58	50	31	29	5	3	-
2002.....	36	34	37	33	41	57	66	37	20	12	-	-
2001.....	35	36	35	40	50	59	65	47	28	3	-	-
2000.....	42	30	45	33	66	61	55	54	19	6	-	-
1999.....	31	39	48	48	58	77	83	65	26	5	4	-

White, female

2004.....	17	20	30	34	37	49	72	81	71	32	7	-
2003.....	17	15	32	21	41	55	83	92	59	18	10	-
2002.....	16	14	19	23	51	72	82	74	67	26	8	-
2001.....	13	19	20	38	54	49	74	79	68	31	2	-
2000.....	19	14	30	32	37	63	80	74	67	31	6	-
1999.....	13	21	21	33	58	79	121	100	91	35	6	-

Black, both sexes

2004.....	23	11	8	16	16	16	14	12	5	4	-	-
2003.....	15	11	9	13	14	19	16	14	8	3	-	-
2002.....	9	15	17	14	12	21	15	11	5	5	-	-
2001.....	13	14	9	15	9	21	12	10	9	2	1	-
2000.....	13	14	13	17	15	16	17	13	9	4	-	-
1999.....	14	6	16	19	14	24	17	17	9	3	1	-

Black, male

2004.....	17	7	6	11	4	9	6	3	2	-	-	-
2003.....	11	9	6	7	8	8	9	4	1	2	-	-
2002.....	5	8	8	8	7	13	8	2	4	3	-	-
2001.....	7	9	8	11	5	11	6	2	3	1	1	-
2000.....	11	10	8	11	5	9	8	4	3	1	-	-
1999.....	11	2	6	10	7	13	9	6	4	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and unspecified disorders of circulatory system (I83-I99)												
Black, female												
2004.....	6	4	2	5	12	7	8	9	3	4	-	-
2003.....	4	2	3	6	6	11	7	10	7	1	-	-
2002.....	4	7	9	6	5	8	7	9	1	2	-	-
2001.....	6	5	1	4	4	10	6	8	6	1	-	-
2000.....	2	4	5	6	10	7	9	9	6	3	-	-
1999.....	3	4	10	9	7	11	8	11	5	3	1	-
American Indian 1/, both sexes												
2004.....	-	2	1	2	-	-	-	1	-	2	-	-
2003.....	1	1	1	-	-	1	-	-	-	-	-	-
2002.....	1	-	1	1	1	-	2	-	-	-	-	-
2001.....	3	1	-	2	1	-	-	-	1	-	-	-
2000.....	-	-	2	2	1	-	-	-	-	-	-	-
1999.....	2	-	1	1	-	1	4	-	-	-	-	-
American Indian 1/, male												
2004.....	-	1	1	1	-	-	-	-	-	1	-	-
2003.....	1	-	1	-	-	-	-	-	-	-	-	-
2002.....	1	-	-	1	1	-	1	-	-	-	-	-
2001.....	2	-	-	1	1	-	-	-	-	-	-	-
2000.....	-	-	1	1	-	-	-	-	-	-	-	-
1999.....	2	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, female												
2004.....	-	1	-	1	-	-	-	1	-	1	-	-
2003.....	-	1	-	-	-	1	-	-	-	-	-	-
2002.....	-	-	1	-	-	-	1	-	-	-	-	-
2001.....	1	1	-	1	-	-	-	-	1	-	-	-
2000.....	-	-	1	1	1	-	-	-	-	-	-	-
1999.....	-	-	1	-	-	1	4	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and unspecified disorders of circulatory system (I83-I99)												
Asian or Pacific Islander 2/, both sexes												
2004.....	2	2	-	1	1	2	3	4	1	-	-	-
2003.....	1	-	1	3	3	1	4	4	2	-	-	-
2002.....	1	-	2	-	2	3	-	2	1	1	-	-
2001.....	1	-	2	2	5	3	3	2	2	-	1	-
2000.....	1	2	3	-	1	2	1	3	3	1	-	-
1999.....	-	2	1	5	5	3	3	3	1	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	1	2	-	-	1	1	3	1	-	-	-	-
2003.....	1	-	-	2	-	-	3	1	1	-	-	-
2002.....	1	-	1	-	1	2	-	1	-	-	-	-
2001.....	-	-	-	2	3	3	3	2	1	-	-	-
2000.....	1	2	2	-	-	2	1	2	-	-	-	-
1999.....	-	2	1	2	3	3	3	3	1	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	1	-	-	1	-	1	-	3	1	-	-	-
2003.....	-	-	1	1	3	1	1	3	1	-	-	-
2002.....	-	-	1	-	1	1	-	1	1	1	-	-
2001.....	1	-	2	-	2	-	-	-	1	-	1	-
2000.....	-	-	1	-	1	-	-	1	3	1	-	-
1999.....	-	-	-	3	2	-	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
X. Diseases of the respiratory system (J00-J98)												
All races, both sexes												
2004.....	4,388	7,125	11,379	16,542	25,750	37,305	44,018	38,111	23,867	8,859	1,918	1
2003.....	4,471	7,028	11,485	17,157	26,766	39,063	44,831	39,735	25,089	9,659	2,130	5
2002.....	4,278	6,854	10,617	16,513	27,408	39,212	45,117	39,780	25,019	9,504	2,113	10
2001.....	4,256	6,651	10,390	17,025	28,010	39,077	42,897	38,757	23,793	9,025	2,049	9
2000.....	4,093	6,511	10,183	17,296	28,721	39,256	43,258	38,353	23,927	9,137	2,097	8
1999.....	3,799	6,467	10,379	17,666	29,311	39,486	42,627	37,537	23,220	9,097	2,134	6
All races, male												
2004.....	2,462	3,806	6,080	8,918	13,597	19,216	21,452	16,877	8,854	2,690	459	1
2003.....	2,443	3,792	6,139	9,170	14,202	20,039	21,633	17,245	9,213	2,782	485	3
2002.....	2,344	3,645	5,683	8,767	14,661	20,325	21,990	17,473	9,301	2,687	420	7
2001.....	2,319	3,562	5,501	9,036	14,975	20,260	20,837	16,683	8,737	2,557	437	6
2000.....	2,212	3,528	5,451	9,271	15,395	20,295	21,249	16,900	8,711	2,697	458	6
1999.....	2,028	3,421	5,644	9,707	16,037	20,813	21,090	16,621	8,377	2,647	474	4
All races, female												
2004.....	1,926	3,319	5,299	7,624	12,153	18,089	22,566	21,234	15,013	6,169	1,459	-
2003.....	2,028	3,236	5,346	7,987	12,564	19,024	23,198	22,490	15,876	6,877	1,645	2
2002.....	1,934	3,209	4,934	7,746	12,747	18,887	23,127	22,307	15,718	6,817	1,693	3
2001.....	1,937	3,089	4,889	7,989	13,035	18,817	22,060	22,074	15,056	6,468	1,612	3
2000.....	1,881	2,983	4,732	8,025	13,326	18,961	22,009	21,453	15,216	6,440	1,639	2
1999.....	1,771	3,046	4,735	7,959	13,274	18,673	21,537	20,916	14,843	6,450	1,660	2
White, both sexes												
2004.....	3,476	6,011	9,960	14,779	23,438	34,260	40,844	35,501	22,156	8,177	1,704	1
2003.....	3,557	5,918	10,044	15,342	24,297	35,870	41,691	37,215	23,331	8,930	1,881	4
2002.....	3,400	5,811	9,250	14,785	24,940	36,084	41,868	37,137	23,325	8,769	1,906	8
2001.....	3,365	5,665	9,084	15,210	25,525	35,984	39,859	36,178	22,043	8,283	1,785	5
2000.....	3,249	5,509	8,848	15,339	26,118	36,261	40,264	35,873	22,261	8,440	1,869	3
1999.....	3,045	5,467	9,068	15,836	26,637	36,376	39,770	35,033	21,623	8,396	1,926	5

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
X. Diseases of the respiratory system (J00-J98)												
White, male												
2004.....	1,960	3,168	5,262	7,884	12,278	17,564	19,813	15,642	8,197	2,455	397	1
2003.....	1,958	3,167	5,291	8,135	12,773	18,215	19,927	16,089	8,489	2,525	426	2
2002.....	1,862	3,061	4,888	7,805	13,262	18,551	20,304	16,253	8,617	2,436	379	6
2001.....	1,812	3,036	4,736	8,010	13,564	18,489	19,259	15,504	8,034	2,320	375	4
2000.....	1,731	2,958	4,701	8,117	13,885	18,595	19,709	15,741	8,025	2,465	400	3
1999.....	1,600	2,862	4,883	8,643	14,482	19,042	19,564	15,441	7,716	2,420	428	3
White, female												
2004.....	1,516	2,843	4,698	6,895	11,160	16,696	21,031	19,859	13,959	5,722	1,307	-
2003.....	1,599	2,751	4,753	7,207	11,524	17,655	21,764	21,126	14,842	6,405	1,455	2
2002.....	1,538	2,750	4,362	6,980	11,678	17,533	21,564	20,884	14,708	6,333	1,527	2
2001.....	1,553	2,629	4,348	7,200	11,961	17,495	20,600	20,674	14,009	5,963	1,410	1
2000.....	1,518	2,551	4,147	7,222	12,233	17,666	20,555	20,132	14,236	5,975	1,469	-
1999.....	1,445	2,605	4,185	7,193	12,155	17,334	20,206	19,592	13,907	5,976	1,498	2
Black, both sexes												
2004.....	820	968	1,223	1,495	1,871	2,391	2,406	1,921	1,232	518	167	-
2003.....	818	953	1,215	1,528	1,996	2,514	2,391	1,852	1,310	539	191	1
2002.....	768	922	1,180	1,454	2,023	2,469	2,562	2,004	1,298	585	169	1
2001.....	789	861	1,105	1,505	2,046	2,480	2,389	1,952	1,338	591	216	2
2000.....	756	881	1,161	1,659	2,163	2,406	2,347	1,894	1,315	566	187	4
1999.....	676	892	1,131	1,566	2,232	2,557	2,269	2,028	1,290	578	175	1
Black, male												
2004.....	452	562	702	888	1,065	1,287	1,217	870	444	153	44	-
2003.....	428	542	713	876	1,153	1,403	1,270	807	498	162	46	1
2002.....	421	523	686	819	1,135	1,382	1,291	861	488	190	33	1
2001.....	441	460	642	860	1,160	1,407	1,223	849	500	170	46	1
2000.....	419	498	647	979	1,254	1,343	1,163	838	505	183	42	2
1999.....	385	495	665	915	1,292	1,438	1,186	914	488	180	40	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
X. Diseases of the respiratory system (J00-J98)												
Black, female												
2004.....	368	406	521	607	806	1,104	1,189	1,051	788	365	123	-
2003.....	390	411	502	652	843	1,111	1,121	1,045	812	377	145	-
2002.....	347	399	494	635	888	1,087	1,271	1,143	810	395	136	-
2001.....	348	401	463	645	886	1,073	1,166	1,103	838	421	170	1
2000.....	337	383	514	680	909	1,063	1,184	1,056	810	383	145	2
1999.....	291	397	466	651	940	1,119	1,083	1,114	802	398	135	-
American Indian 1/, both sexes												
2004.....	45	65	83	94	135	159	157	87	55	24	9	-
2003.....	48	70	104	108	132	162	152	126	77	19	16	-
2002.....	43	53	76	96	115	139	137	115	72	24	8	-
2001.....	39	47	80	93	119	151	130	110	61	27	6	1
2000.....	32	52	74	97	120	126	122	104	66	14	8	-
1999.....	35	39	76	79	119	155	113	110	61	23	8	-
American Indian 1/, male												
2004.....	28	32	42	47	69	77	76	36	24	10	5	-
2003.....	26	35	55	50	71	89	77	46	30	14	3	-
2002.....	19	27	39	46	57	78	66	55	27	7	2	-
2001.....	21	22	40	40	64	73	64	48	20	11	2	1
2000.....	25	27	48	49	52	72	62	44	31	7	3	-
1999.....	20	22	34	42	62	81	48	50	23	7	1	-
American Indian 1/, female												
2004.....	17	33	41	47	66	82	81	51	31	14	4	-
2003.....	22	35	49	58	61	73	75	80	47	5	13	-
2002.....	24	26	37	50	58	61	71	60	45	17	6	-
2001.....	18	25	40	53	55	78	66	62	41	16	4	-
2000.....	7	25	26	48	68	54	60	60	35	7	5	-
1999.....	15	17	42	37	57	74	65	60	38	16	7	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
X. Diseases of the respiratory system (J00-J98)												
Asian or Pacific Islander 2/, both sexes												
2004.....	47	81	113	174	306	495	611	602	424	140	38	-
2003.....	48	87	122	179	341	517	597	542	371	171	42	-
2002.....	67	68	111	178	330	520	550	524	324	126	30	1
2001.....	63	78	121	217	320	462	519	517	351	124	42	1
2000.....	56	69	100	201	320	463	525	482	285	117	33	1
1999.....	43	69	104	185	323	398	475	366	246	100	25	-
Asian or Pacific Islander 2/, male												
2004.....	22	44	74	99	185	288	346	329	189	72	13	-
2003.....	31	48	80	109	205	332	359	303	196	81	10	-
2002.....	42	34	70	97	207	314	329	304	169	54	6	-
2001.....	45	44	83	126	187	291	291	282	183	56	14	-
2000.....	37	45	55	126	204	285	315	277	150	42	13	1
1999.....	23	42	62	107	201	252	292	216	150	40	5	-
Asian or Pacific Islander 2/, female												
2004.....	25	37	39	75	121	207	265	273	235	68	25	-
2003.....	17	39	42	70	136	185	238	239	175	90	32	-
2002.....	25	34	41	81	123	206	221	220	155	72	24	1
2001.....	18	34	38	91	133	171	228	235	168	68	28	1
2000.....	19	24	45	75	116	178	210	205	135	75	20	-
1999.....	20	27	42	78	122	146	183	150	96	60	20	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Acute upper respiratory infections (J00-J06)

All races, both sexes

2004.....	5	7	7	6	5	11	15	11	11	10	8	-
2003.....	11	6	12	4	5	4	17	29	19	10	11	-
2002.....	11	11	3	6	12	10	10	11	19	16	4	-
2001.....	3	4	8	6	5	12	14	16	24	9	8	-
2000.....	9	3	3	10	14	17	20	27	31	7	9	-
1999.....	7	4	8	5	10	16	26	28	21	15	8	-

All races, male

2004.....	2	6	1	1	4	4	6	5	2	2	1	-
2003.....	7	2	9	4	2	2	7	10	7	5	2	-
2002.....	9	5	-	3	5	3	4	2	6	2	-	-
2001.....	1	1	3	2	-	3	5	3	1	4	-	-
2000.....	6	1	3	5	7	5	7	7	7	-	-	-
1999.....	2	2	4	4	5	8	11	9	2	2	-	-

All races, female

2004.....	3	1	6	5	1	7	9	6	9	8	7	-
2003.....	4	4	3	-	3	2	10	19	12	5	9	-
2002.....	2	6	3	3	7	7	6	9	13	14	4	-
2001.....	2	3	5	4	5	9	9	13	23	5	8	-
2000.....	3	2	-	5	7	12	13	20	24	7	9	-
1999.....	5	2	4	1	5	8	15	19	19	13	8	-

White, both sexes

2004.....	4	5	7	5	4	11	14	10	11	8	6	-
2003.....	6	6	8	2	5	4	17	27	17	7	11	-
2002.....	9	10	3	4	10	10	8	10	17	16	4	-
2001.....	2	3	6	6	3	11	14	16	23	9	6	-
2000.....	8	3	2	8	14	17	16	24	30	7	9	-
1999.....	5	3	6	4	7	16	24	26	20	13	8	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute upper respiratory infections (J00-J06)												
White, male												
2004.....	1	4	1	1	3	4	5	4	2	1	1	-
2003.....	4	2	6	2	2	2	7	9	5	3	2	-
2002.....	7	5	-	2	4	3	3	2	4	2	-	-
2001.....	1	-	2	2	-	3	5	3	1	4	-	-
2000.....	6	1	2	3	7	5	5	6	6	-	-	-
1999.....	2	2	2	3	4	8	10	7	1	1	-	-
White, female												
2004.....	3	1	6	4	1	7	9	6	9	7	5	-
2003.....	2	4	2	-	3	2	10	18	12	4	9	-
2002.....	2	5	3	2	6	7	5	8	13	14	4	-
2001.....	1	3	4	4	3	8	9	13	22	5	6	-
2000.....	2	2	-	5	7	12	11	18	24	7	9	-
1999.....	3	1	4	1	3	8	14	19	19	12	8	-
Black, both sexes												
2004.....	1	1	-	1	1	-	1	1	-	1	1	-
2003.....	4	-	3	2	-	-	-	2	1	2	-	-
2002.....	2	1	-	2	-	-	2	-	-	-	-	-
2001.....	1	1	2	-	1	1	-	-	1	-	2	-
2000.....	1	-	-	2	-	-	1	1	1	-	-	-
1999.....	1	1	2	-	3	-	1	2	1	2	-	-
Black, male												
2004.....	1	1	-	-	1	-	1	1	-	-	-	-
2003.....	2	-	3	2	-	-	-	1	1	1	-	-
2002.....	2	-	-	1	-	-	1	-	-	-	-	-
2001.....	-	1	1	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	2	-	-	1	1	1	-	-	-
1999.....	-	-	2	-	1	-	-	2	1	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Acute pharyngitis and tonsillitis (J02-J03)

All races, both sexes

2004.....	2	1	3	-	1	1	-	2	-	-	-	-
2003.....	2	1	1	-	2	-	-	-	-	2	-	-
2002.....	3	4	-	-	1	-	2	-	1	1	-	-
2001.....	-	-	2	3	-	3	2	1	2	1	1	-
2000.....	2	1	1	3	2	2	-	1	2	-	-	-
1999.....	-	-	1	-	-	2	2	2	1	-	-	-

All races, male

2004.....	-	1	-	-	1	1	-	1	-	-	-	-
2003.....	2	-	-	-	-	-	-	-	-	1	-	-
2002.....	3	1	-	-	-	-	2	-	-	-	-	-
2001.....	-	-	1	1	-	2	-	-	-	1	-	-
2000.....	-	1	1	1	2	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	1	-	2	-	-	-	-

All races, female

2004.....	2	-	3	-	-	-	-	1	-	-	-	-
2003.....	-	1	1	-	2	-	-	-	-	1	-	-
2002.....	-	3	-	-	1	-	-	-	1	1	-	-
2001.....	-	-	1	2	-	1	2	1	2	-	1	-
2000.....	2	-	-	2	-	2	-	1	2	-	-	-
1999.....	-	-	1	-	-	1	2	-	1	-	-	-

White, both sexes

2004.....	2	-	3	-	1	1	-	2	-	-	-	-
2003.....	1	1	-	-	2	-	-	-	-	1	-	-
2002.....	2	3	-	-	1	-	1	-	1	1	-	-
2001.....	-	-	2	3	-	3	2	1	2	1	1	-
2000.....	1	1	1	3	2	2	-	1	2	-	-	-
1999.....	-	-	1	-	-	2	2	-	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute pharyngitis and tonsillitis (J02-J03)												
White, male												
2004.....	-	-	-	-	1	1	-	1	-	-	-	-
2003.....	1	-	-	-	-	-	-	-	-	-	-	-
2002.....	2	1	-	-	-	-	1	-	-	-	-	-
2001.....	-	-	1	1	-	2	-	-	-	1	-	-
2000.....	-	1	1	1	2	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	1	-	-	-	-	-	-
White, female												
2004.....	2	-	3	-	-	-	-	1	-	-	-	-
2003.....	-	1	-	-	2	-	-	-	-	1	-	-
2002.....	-	2	-	-	1	-	-	-	1	1	-	-
2001.....	-	-	1	2	-	1	2	1	2	-	1	-
2000.....	1	-	-	2	-	2	-	1	2	-	-	-
1999.....	-	-	1	-	-	1	2	-	1	-	-	-
Black, both sexes												
2004.....	-	1	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	1	-	-
2002.....	1	1	-	-	-	-	1	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	1	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	2	-	-	-	-
Black, male												
2004.....	-	1	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	1	-	-
2002.....	1	-	-	-	-	-	1	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	2	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute laryngitis and tracheitis (J04)												
All races, both sexes												
2004.....	-	1	2	-	2	2	3	1	-	-	-	-
2003.....	3	2	4	-	-	-	1	-	-	-	-	-
2002.....	1	-	1	1	2	-	-	-	-	-	-	-
2001.....	1	-	-	1	-	-	3	2	-	-	-	-
2000.....	1	-	1	1	-	1	1	-	-	-	-	-
1999.....	1	-	1	-	1	1	1	-	1	1	-	-
All races, male												
2004.....	-	1	-	-	1	1	3	1	-	-	-	-
2003.....	2	1	3	-	-	-	-	-	-	-	-	-
2002.....	1	-	-	-	1	-	-	-	-	-	-	-
2001.....	1	-	-	-	-	-	2	-	-	-	-	-
2000.....	1	-	1	1	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	1	-	-	-	-	-	-
All races, female												
2004.....	-	-	2	-	1	1	-	-	-	-	-	-
2003.....	1	1	1	-	-	-	1	-	-	-	-	-
2002.....	-	-	1	1	1	-	-	-	-	-	-	-
2001.....	-	-	-	1	-	-	1	2	-	-	-	-
2000.....	-	-	-	-	-	1	1	-	-	-	-	-
1999.....	1	-	1	-	1	-	1	-	1	1	-	-
White, both sexes												
2004.....	-	1	2	-	2	2	2	-	-	-	-	-
2003.....	1	2	3	-	-	-	1	-	-	-	-	-
2002.....	1	-	1	1	2	-	-	-	-	-	-	-
2001.....	1	-	-	1	-	-	3	2	-	-	-	-
2000.....	1	-	-	-	-	1	-	-	-	-	-	-
1999.....	1	-	1	-	1	1	1	-	1	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified acute upper respiratory infections (J00-J01,J05-J06)												
All races, both sexes												
2004.....	3	5	2	6	2	8	12	8	11	10	8	-
2003.....	6	3	7	4	3	4	16	29	19	8	11	-
2002.....	7	7	2	5	9	10	8	11	18	15	4	-
2001.....	2	4	6	2	5	9	9	13	22	8	7	-
2000.....	6	2	1	6	12	14	19	26	29	7	9	-
1999.....	6	4	6	5	9	13	23	26	19	14	8	-
All races, male												
2004.....	2	4	1	1	2	2	3	3	2	2	1	-
2003.....	3	1	6	4	2	2	7	10	7	4	2	-
2002.....	5	4	-	3	4	3	2	2	6	2	-	-
2001.....	-	1	2	1	-	1	3	3	1	3	-	-
2000.....	5	-	1	3	5	5	7	7	7	-	-	-
1999.....	2	2	4	4	5	6	11	7	2	2	-	-
All races, female												
2004.....	1	1	1	5	-	6	9	5	9	8	7	-
2003.....	3	2	1	-	1	2	9	19	12	4	9	-
2002.....	2	3	2	2	5	7	6	9	12	13	4	-
2001.....	2	3	4	1	5	8	6	10	21	5	7	-
2000.....	1	2	-	3	7	9	12	19	22	7	9	-
1999.....	4	2	2	1	4	7	12	19	17	12	8	-
White, both sexes												
2004.....	2	4	2	5	1	8	12	8	11	8	6	-
2003.....	4	3	5	2	3	4	16	27	17	6	11	-
2002.....	6	7	2	3	7	10	7	10	16	15	4	-
2001.....	1	3	4	2	3	8	9	13	21	8	5	-
2000.....	6	2	1	5	12	14	16	23	28	7	9	-
1999.....	4	3	4	4	6	13	21	26	18	12	8	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified acute upper respiratory infections (J00-J01,J05-J06)												
White, male												
2004.....	1	3	1	1	1	2	3	3	2	1	1	-
2003.....	2	1	4	2	2	2	7	9	5	3	2	-
2002.....	4	4	-	2	3	3	2	2	4	2	-	-
2001.....	-	-	1	1	-	1	3	3	1	3	-	-
2000.....	5	-	1	2	5	5	5	6	6	-	-	-
1999.....	2	2	2	3	4	6	10	7	1	1	-	-
White, female												
2004.....	1	1	1	4	-	6	9	5	9	7	5	-
2003.....	2	2	1	-	1	2	9	18	12	3	9	-
2002.....	2	3	2	1	4	7	5	8	12	13	4	-
2001.....	1	3	3	1	3	7	6	10	20	5	5	-
2000.....	1	2	-	3	7	9	11	17	22	7	9	-
1999.....	2	1	2	1	2	7	11	19	17	11	8	-
Black, both sexes												
2004.....	1	-	-	1	1	-	-	-	-	1	1	-
2003.....	2	-	2	2	-	-	-	2	1	1	-	-
2002.....	1	-	-	2	-	-	1	-	-	-	-	-
2001.....	1	1	2	-	1	1	-	-	1	-	2	-
2000.....	-	-	-	1	-	-	1	1	1	-	-	-
1999.....	1	1	2	-	3	-	1	-	1	2	-	-
Black, male												
2004.....	1	-	-	-	1	-	-	-	-	-	-	-
2003.....	1	-	2	2	-	-	-	1	1	-	-	-
2002.....	1	-	-	1	-	-	-	-	-	-	-	-
2001.....	-	1	1	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	1	-	-	1	1	1	-	-	-
1999.....	-	-	2	-	1	-	-	-	1	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of the respiratory system (J10-J98)												
All races, both sexes												
2004.....	4,383	7,118	11,372	16,536	25,745	37,294	44,003	38,100	23,856	8,849	1,910	1
2003.....	4,460	7,022	11,473	17,153	26,761	39,059	44,814	39,706	25,070	9,649	2,119	5
2002.....	4,267	6,843	10,614	16,507	27,396	39,202	45,107	39,769	25,000	9,488	2,109	10
2001.....	4,253	6,647	10,382	17,019	28,005	39,065	42,883	38,741	23,769	9,016	2,041	9
2000.....	4,084	6,508	10,180	17,286	28,707	39,239	43,238	38,326	23,896	9,130	2,088	8
1999.....	3,792	6,463	10,371	17,661	29,301	39,470	42,601	37,509	23,199	9,082	2,126	6
All races, male												
2004.....	2,460	3,800	6,079	8,917	13,593	19,212	21,446	16,872	8,852	2,688	458	1
2003.....	2,436	3,790	6,130	9,166	14,200	20,037	21,626	17,235	9,206	2,777	483	3
2002.....	2,335	3,640	5,683	8,764	14,656	20,322	21,986	17,471	9,295	2,685	420	7
2001.....	2,318	3,561	5,498	9,034	14,975	20,257	20,832	16,680	8,736	2,553	437	6
2000.....	2,206	3,527	5,448	9,266	15,388	20,290	21,242	16,893	8,704	2,697	458	6
1999.....	2,026	3,419	5,640	9,703	16,032	20,805	21,079	16,612	8,375	2,645	474	4
All races, female												
2004.....	1,923	3,318	5,293	7,619	12,152	18,082	22,557	21,228	15,004	6,161	1,452	-
2003.....	2,024	3,232	5,343	7,987	12,561	19,022	23,188	22,471	15,864	6,872	1,636	2
2002.....	1,932	3,203	4,931	7,743	12,740	18,880	23,121	22,298	15,705	6,803	1,689	3
2001.....	1,935	3,086	4,884	7,985	13,030	18,808	22,051	22,061	15,033	6,463	1,604	3
2000.....	1,878	2,981	4,732	8,020	13,319	18,949	21,996	21,433	15,192	6,433	1,630	2
1999.....	1,766	3,044	4,731	7,958	13,269	18,665	21,522	20,897	14,824	6,437	1,652	2
White, both sexes												
2004.....	3,472	6,006	9,953	14,774	23,434	34,249	40,830	35,491	22,145	8,169	1,698	1
2003.....	3,551	5,912	10,036	15,340	24,292	35,866	41,674	37,188	23,314	8,923	1,870	4
2002.....	3,391	5,801	9,247	14,781	24,930	36,074	41,860	37,127	23,308	8,753	1,902	8
2001.....	3,363	5,662	9,078	15,204	25,522	35,973	39,845	36,162	22,020	8,274	1,779	5
2000.....	3,241	5,506	8,846	15,331	26,104	36,244	40,248	35,849	22,231	8,433	1,860	3
1999.....	3,040	5,464	9,062	15,832	26,630	36,360	39,746	35,007	21,603	8,383	1,918	5

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of the respiratory system (J10-J98)												
White, male												
2004.....	1,959	3,164	5,261	7,883	12,275	17,560	19,808	15,638	8,195	2,454	396	1
2003.....	1,954	3,165	5,285	8,133	12,771	18,213	19,920	16,080	8,484	2,522	424	2
2002.....	1,855	3,056	4,888	7,803	13,258	18,548	20,301	16,251	8,613	2,434	379	6
2001.....	1,811	3,036	4,734	8,008	13,564	18,486	19,254	15,501	8,033	2,316	375	4
2000.....	1,725	2,957	4,699	8,114	13,878	18,590	19,704	15,735	8,019	2,465	400	3
1999.....	1,598	2,860	4,881	8,640	14,478	19,034	19,554	15,434	7,715	2,419	428	3
White, female												
2004.....	1,513	2,842	4,692	6,891	11,159	16,689	21,022	19,853	13,950	5,715	1,302	-
2003.....	1,597	2,747	4,751	7,207	11,521	17,653	21,754	21,108	14,830	6,401	1,446	2
2002.....	1,536	2,745	4,359	6,978	11,672	17,526	21,559	20,876	14,695	6,319	1,523	2
2001.....	1,552	2,626	4,344	7,196	11,958	17,487	20,591	20,661	13,987	5,958	1,404	1
2000.....	1,516	2,549	4,147	7,217	12,226	17,654	20,544	20,114	14,212	5,968	1,460	-
1999.....	1,442	2,604	4,181	7,192	12,152	17,326	20,192	19,573	13,888	5,964	1,490	2
Black, both sexes												
2004.....	819	967	1,223	1,494	1,870	2,391	2,405	1,920	1,232	517	166	-
2003.....	814	953	1,212	1,526	1,996	2,514	2,391	1,850	1,309	537	191	1
2002.....	766	921	1,180	1,452	2,023	2,469	2,560	2,004	1,298	585	169	1
2001.....	788	860	1,103	1,505	2,045	2,479	2,389	1,952	1,337	591	214	2
2000.....	755	881	1,161	1,657	2,163	2,406	2,346	1,893	1,314	566	187	4
1999.....	675	891	1,129	1,566	2,229	2,557	2,268	2,026	1,289	576	175	1
Black, male												
2004.....	451	561	702	888	1,064	1,287	1,216	869	444	153	44	-
2003.....	426	542	710	874	1,153	1,403	1,270	806	497	161	46	1
2002.....	419	523	686	818	1,135	1,382	1,290	861	488	190	33	1
2001.....	441	459	641	860	1,160	1,407	1,223	849	500	170	46	1
2000.....	419	498	647	977	1,254	1,343	1,162	837	504	183	42	2
1999.....	385	495	663	915	1,291	1,438	1,186	912	487	179	40	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of the respiratory system (J10-J98)												
Black, female												
2004.....	368	406	521	606	806	1,104	1,189	1,051	788	364	122	-
2003.....	388	411	502	652	843	1,111	1,121	1,044	812	376	145	-
2002.....	347	398	494	634	888	1,087	1,270	1,143	810	395	136	-
2001.....	347	401	462	645	885	1,072	1,166	1,103	837	421	168	1
2000.....	336	383	514	680	909	1,063	1,184	1,056	810	383	145	2
1999.....	290	396	466	651	938	1,119	1,082	1,114	802	397	135	-
American Indian 1/, both sexes												
2004.....	45	64	83	94	135	159	157	87	55	24	9	-
2003.....	48	70	103	108	132	162	152	126	77	19	16	-
2002.....	43	53	76	96	115	139	137	115	72	24	8	-
2001.....	39	47	80	93	119	151	130	110	61	27	6	1
2000.....	32	52	73	97	120	126	120	104	66	14	8	-
1999.....	34	39	76	79	119	155	113	110	61	23	8	-
American Indian 1/, male												
2004.....	28	31	42	47	69	77	76	36	24	10	5	-
2003.....	26	35	55	50	71	89	77	46	30	14	3	-
2002.....	19	27	39	46	57	78	66	55	27	7	2	-
2001.....	21	22	40	40	64	73	64	48	20	11	2	1
2000.....	25	27	47	49	52	72	61	44	31	7	3	-
1999.....	20	22	34	42	62	81	48	50	23	7	1	-
American Indian 1/, female												
2004.....	17	33	41	47	66	82	81	51	31	14	4	-
2003.....	22	35	48	58	61	73	75	80	47	5	13	-
2002.....	24	26	37	50	58	61	71	60	45	17	6	-
2001.....	18	25	40	53	55	78	66	62	41	16	4	-
2000.....	7	25	26	48	68	54	59	60	35	7	5	-
1999.....	14	17	42	37	57	74	65	60	38	16	7	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of the respiratory system (J10-J98)												
Asian or Pacific Islander 2/, both sexes												
2004.....	47	81	113	174	306	495	611	602	424	139	37	-
2003.....	47	87	122	179	341	517	597	542	370	170	42	-
2002.....	67	68	111	178	328	520	550	523	322	126	30	1
2001.....	63	78	121	217	319	462	519	517	351	124	42	1
2000.....	56	69	100	201	320	463	524	480	285	117	33	1
1999.....	43	69	104	184	323	398	474	366	246	100	25	-
Asian or Pacific Islander 2/, male												
2004.....	22	44	74	99	185	288	346	329	189	71	13	-
2003.....	30	48	80	109	205	332	359	303	195	80	10	-
2002.....	42	34	70	97	206	314	329	304	167	54	6	-
2001.....	45	44	83	126	187	291	291	282	183	56	14	-
2000.....	37	45	55	126	204	285	315	277	150	42	13	1
1999.....	23	42	62	106	201	252	291	216	150	40	5	-
Asian or Pacific Islander 2/, female												
2004.....	25	37	39	75	121	207	265	273	235	68	24	-
2003.....	17	39	42	70	136	185	238	239	175	90	32	-
2002.....	25	34	41	81	122	206	221	219	155	72	24	1
2001.....	18	34	38	91	132	171	228	235	168	68	28	1
2000.....	19	24	45	75	116	178	209	203	135	75	20	-
1999.....	20	27	42	78	122	146	183	150	96	60	20	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Influenza (J10-J11)												
All races, both sexes												
2004.....	11	23	32	31	68	115	176	238	212	84	20	-
2003.....	32	47	38	72	111	200	257	341	303	128	31	-
2002.....	9	13	17	19	37	59	118	174	144	59	27	-
2001.....	7	8	13	10	11	18	38	37	39	14	6	-
2000.....	24	39	54	76	123	189	303	372	305	130	48	-
1999.....	13	29	42	60	104	154	297	368	303	162	48	-
All races, male												
2004.....	5	11	14	18	40	56	69	92	52	22	2	-
2003.....	18	26	16	38	51	89	100	132	84	26	3	-
2002.....	5	7	13	13	19	28	53	66	45	10	3	-
2001.....	4	4	5	3	5	6	14	15	8	4	2	-
2000.....	14	20	24	43	63	81	115	127	86	35	6	-
1999.....	7	13	24	27	47	71	131	127	70	35	6	-
All races, female												
2004.....	6	12	18	13	28	59	107	146	160	62	18	-
2003.....	14	21	22	34	60	111	157	209	219	102	28	-
2002.....	4	6	4	6	18	31	65	108	99	49	24	-
2001.....	3	4	8	7	6	12	24	22	31	10	4	-
2000.....	10	19	30	33	60	108	188	245	219	95	42	-
1999.....	6	16	18	33	57	83	166	241	233	127	42	-
White, both sexes												
2004.....	9	22	28	28	64	100	171	231	208	81	20	-
2003.....	27	41	35	64	106	191	248	330	292	123	29	-
2002.....	8	11	14	13	34	57	115	172	136	58	26	-
2001.....	7	7	10	8	11	18	34	34	38	13	6	-
2000.....	22	37	50	70	110	180	289	353	295	125	47	-
1999.....	10	26	39	55	93	146	283	356	298	157	46	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Influenza (J10-J11)												
White, male												
2004.....	5	10	13	17	37	46	68	89	52	22	2	-
2003.....	16	22	14	35	49	85	97	129	83	24	3	-
2002.....	5	5	11	9	19	27	51	65	44	10	3	-
2001.....	4	3	3	1	5	6	14	14	7	4	2	-
2000.....	12	18	23	40	59	75	112	119	83	32	6	-
1999.....	6	13	23	24	43	70	124	124	70	34	6	-
White, female												
2004.....	4	12	15	11	27	54	103	142	156	59	18	-
2003.....	11	19	21	29	57	106	151	201	209	99	26	-
2002.....	3	6	3	4	15	30	64	107	92	48	23	-
2001.....	3	4	7	7	6	12	20	20	31	9	4	-
2000.....	10	19	27	30	51	105	177	234	212	93	41	-
1999.....	4	13	16	31	50	76	159	232	228	123	40	-
Black, both sexes												
2004.....	1	-	4	3	4	12	4	3	4	2	-	-
2003.....	5	5	2	6	4	5	8	8	8	3	1	-
2002.....	-	1	3	4	2	1	3	1	5	1	1	-
2001.....	-	1	2	2	-	-	4	3	1	1	-	-
2000.....	2	2	4	5	11	5	8	17	8	4	1	-
1999.....	2	3	2	3	10	5	10	8	4	4	2	-
Black, male												
2004.....	-	-	1	1	3	9	1	1	-	-	-	-
2003.....	2	4	1	3	1	2	2	2	-	1	-	-
2002.....	-	1	2	3	-	-	2	1	1	-	-	-
2001.....	-	1	2	2	-	-	-	1	1	-	-	-
2000.....	2	2	1	2	3	4	1	7	1	3	-	-
1999.....	1	-	1	2	4	1	5	2	-	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Influenza (J10-J11)												
Black, female												
2004.....	1	-	3	2	1	3	3	2	4	2	-	-
2003.....	3	1	1	3	3	3	6	6	8	2	1	-
2002.....	-	-	1	1	2	1	1	-	4	1	1	-
2001.....	-	-	-	-	-	-	4	2	-	1	-	-
2000.....	-	-	3	3	8	1	7	10	7	1	1	-
1999.....	1	3	1	1	6	4	5	6	4	3	2	-
American Indian 1/, both sexes												
2004.....	-	-	-	-	-	1	-	1	-	-	-	-
2003.....	-	-	-	-	1	2	1	1	2	1	1	-
2002.....	-	1	-	1	1	1	-	-	1	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	1	-	1	1	-	-	-
1999.....	-	-	-	-	1	1	1	-	-	-	-	-
American Indian 1/, male												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	1	-	1	-	-	1	-	-
2002.....	-	1	-	1	-	1	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	1	-	-	-
1999.....	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, female												
2004.....	-	-	-	-	-	1	-	1	-	-	-	-
2003.....	-	-	-	-	-	2	-	1	2	-	1	-
2002.....	-	-	-	-	1	-	-	-	1	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	1	-	1	-	-	-	-
1999.....	-	-	-	-	1	1	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Influenza (J10-J11)												
Asian or Pacific Islander 2/, both sexes												
2004.....	1	1	-	-	-	2	1	3	-	1	-	-
2003.....	-	1	1	2	-	2	-	2	1	1	-	-
2002.....	1	-	-	1	-	-	-	1	2	-	-	-
2001.....	-	-	1	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	1	2	3	6	1	1	1	-	-
1999.....	1	-	1	2	-	2	3	4	1	1	-	-
Asian or Pacific Islander 2/, male												
2004.....	-	1	-	-	-	1	-	2	-	-	-	-
2003.....	-	-	1	-	-	2	-	1	1	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	1	1	2	2	1	1	-	-	-
1999.....	-	-	-	1	-	-	1	1	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	1	-	-	-	-	1	1	1	-	1	-	-
2003.....	-	1	-	2	-	-	-	1	-	1	-	-
2002.....	1	-	-	1	-	-	-	1	2	-	-	-
2001.....	-	-	1	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	1	1	4	-	-	1	-	-
1999.....	1	-	1	1	-	2	2	3	1	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonia (J12-J18)												
All races, both sexes												
2004.....	1,083	1,347	1,752	2,339	3,944	6,902	10,873	12,111	9,980	4,477	1,190	-
2003.....	1,190	1,245	1,800	2,516	4,132	7,498	11,487	13,312	10,897	5,044	1,341	2
2002.....	1,022	1,258	1,699	2,401	4,390	7,846	11,961	13,803	11,346	5,113	1,329	3
2001.....	984	1,147	1,536	2,313	4,316	7,492	11,129	13,345	10,653	4,822	1,275	5
2000.....	967	1,159	1,627	2,457	4,533	7,723	11,606	13,380	11,013	4,986	1,313	3
1999.....	886	1,069	1,485	2,322	4,375	7,622	11,119	13,131	10,848	4,997	1,372	2
All races, male												
2004.....	671	780	1,018	1,362	2,185	3,609	5,279	5,116	3,412	1,250	266	-
2003.....	733	713	1,016	1,468	2,325	3,943	5,497	5,460	3,631	1,334	280	1
2002.....	614	735	983	1,371	2,447	4,086	5,728	5,650	3,830	1,342	253	2
2001.....	598	682	896	1,335	2,389	3,989	5,285	5,358	3,579	1,278	258	4
2000.....	588	690	946	1,404	2,503	4,003	5,552	5,448	3,638	1,338	267	2
1999.....	552	598	827	1,367	2,452	3,949	5,307	5,374	3,525	1,260	279	1
All races, female												
2004.....	412	567	734	977	1,759	3,293	5,594	6,995	6,568	3,227	924	-
2003.....	457	532	784	1,048	1,807	3,555	5,990	7,852	7,266	3,710	1,061	1
2002.....	408	523	716	1,030	1,943	3,760	6,233	8,153	7,516	3,771	1,076	1
2001.....	386	465	640	978	1,927	3,503	5,844	7,987	7,074	3,544	1,017	1
2000.....	379	469	681	1,053	2,030	3,720	6,054	7,932	7,375	3,648	1,046	1
1999.....	334	471	658	955	1,923	3,673	5,812	7,757	7,323	3,737	1,093	1
White, both sexes												
2004.....	800	1,050	1,410	1,881	3,357	6,013	9,730	11,028	9,132	4,117	1,060	-
2003.....	885	955	1,421	2,043	3,490	6,593	10,390	12,281	10,027	4,652	1,184	1
2002.....	775	970	1,362	2,005	3,778	6,912	10,814	12,667	10,471	4,712	1,198	3
2001.....	738	899	1,225	1,913	3,712	6,525	10,044	12,275	9,814	4,411	1,108	4
2000.....	714	895	1,276	1,960	3,829	6,848	10,494	12,324	10,185	4,603	1,169	1
1999.....	656	833	1,161	1,939	3,745	6,816	10,160	12,061	10,014	4,614	1,239	2

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonia (J12-J18)												
White, male												
2004.....	496	616	815	1,073	1,858	3,134	4,731	4,670	3,131	1,139	226	-
2003.....	548	540	771	1,176	1,952	3,450	4,945	5,040	3,304	1,203	238	-
2002.....	465	563	779	1,141	2,122	3,597	5,168	5,180	3,517	1,212	229	2
2001.....	430	533	700	1,089	2,049	3,446	4,759	4,917	3,256	1,156	222	4
2000.....	422	518	743	1,105	2,110	3,548	5,007	5,009	3,332	1,229	235	1
1999.....	398	458	643	1,140	2,081	3,528	4,844	4,914	3,214	1,158	255	1
White, female												
2004.....	304	434	595	808	1,499	2,879	4,999	6,358	6,001	2,978	834	-
2003.....	337	415	650	867	1,538	3,143	5,445	7,241	6,723	3,449	946	1
2002.....	310	407	583	864	1,656	3,315	5,646	7,487	6,954	3,500	969	1
2001.....	308	366	525	824	1,663	3,079	5,285	7,358	6,558	3,255	886	-
2000.....	292	377	533	855	1,719	3,300	5,487	7,315	6,853	3,374	934	-
1999.....	258	375	518	799	1,664	3,288	5,316	7,147	6,800	3,456	984	1
Black, both sexes												
2004.....	257	261	287	383	473	683	865	801	597	283	101	-
2003.....	267	243	319	391	526	715	802	750	641	277	113	1
2002.....	211	256	300	330	512	717	882	859	651	312	107	-
2001.....	215	219	258	316	488	770	823	792	636	321	139	-
2000.....	226	238	294	411	570	695	861	810	649	311	121	2
1999.....	206	210	283	330	546	660	751	876	675	309	113	-
Black, male												
2004.....	161	145	176	252	267	364	414	315	185	71	30	-
2003.....	163	148	205	247	305	387	384	284	223	75	30	1
2002.....	124	158	180	190	269	377	419	320	209	93	19	-
2001.....	145	132	163	195	282	438	385	291	227	83	27	-
2000.....	146	155	169	247	319	361	407	318	218	83	25	1
1999.....	139	128	160	198	319	336	353	356	227	80	20	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonia (J12-J18)												
Black, female												
2004.....	96	116	111	131	206	319	451	486	412	212	71	-
2003.....	104	95	114	144	221	328	418	466	418	202	83	-
2002.....	87	98	120	140	243	340	463	539	442	219	88	-
2001.....	70	87	95	121	206	332	438	501	409	238	112	-
2000.....	80	83	125	164	251	334	454	492	431	228	96	1
1999.....	67	82	123	132	227	324	398	520	448	229	93	-
American Indian 1/, both sexes												
2004.....	7	14	18	21	17	41	49	29	29	9	6	-
2003.....	17	21	28	30	25	29	50	50	40	10	12	-
2002.....	13	17	7	18	15	34	46	37	38	17	6	-
2001.....	14	9	16	19	23	37	48	41	31	17	2	-
2000.....	13	6	18	20	25	36	34	35	35	9	3	-
1999.....	12	12	18	15	21	36	34	46	38	18	4	-
American Indian 1/, male												
2004.....	5	8	5	10	8	21	25	13	13	4	3	-
2003.....	9	10	17	15	15	14	25	13	14	7	2	-
2002.....	7	7	5	10	8	18	14	16	14	4	2	-
2001.....	9	4	9	11	12	18	24	19	10	8	1	-
2000.....	11	3	13	15	14	18	15	11	15	4	-	-
1999.....	9	6	7	6	10	24	14	16	14	6	-	-
American Indian 1/, female												
2004.....	2	6	13	11	9	20	24	16	16	5	3	-
2003.....	8	11	11	15	10	15	25	37	26	3	10	-
2002.....	6	10	2	8	7	16	32	21	24	13	4	-
2001.....	5	5	7	8	11	19	24	22	21	9	1	-
2000.....	2	3	5	5	11	18	19	24	20	5	3	-
1999.....	3	6	11	9	11	12	20	30	24	12	4	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonia (J12-J18)												
Asian or Pacific Islander 2/, both sexes												
2004.....	19	22	37	54	97	165	229	253	222	68	23	-
2003.....	21	26	32	52	91	161	245	231	189	105	32	-
2002.....	23	15	30	48	85	183	219	240	186	72	18	-
2001.....	17	20	37	65	93	160	214	237	172	73	26	1
2000.....	14	20	39	66	109	144	217	211	144	63	20	-
1999.....	12	14	23	38	63	110	174	148	121	56	16	-
Asian or Pacific Islander 2/, male												
2004.....	9	11	22	27	52	90	109	118	83	36	7	-
2003.....	13	15	23	30	53	92	143	123	90	49	10	-
2002.....	18	7	19	30	48	94	127	134	90	33	3	-
2001.....	14	13	24	40	46	87	117	131	86	31	8	-
2000.....	9	14	21	37	60	76	123	110	73	22	7	-
1999.....	6	6	17	23	42	61	96	88	70	16	4	-
Asian or Pacific Islander 2/, female												
2004.....	10	11	15	27	45	75	120	135	139	32	16	-
2003.....	8	11	9	22	38	69	102	108	99	56	22	-
2002.....	5	8	11	18	37	89	92	106	96	39	15	-
2001.....	3	7	13	25	47	73	97	106	86	42	18	1
2000.....	5	6	18	29	49	68	94	101	71	41	13	-
1999.....	6	8	6	15	21	49	78	60	51	40	12	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Viral pneumonia, not elsewhere classified (J12)												
All races, both sexes												
2004.....	5	3	4	11	8	20	35	31	12	11	8	-
2003.....	5	8	11	8	18	28	33	44	33	17	4	-
2002.....	6	9	7	8	22	31	39	32	27	14	5	-
2001.....	6	9	6	14	9	24	37	25	26	9	3	-
2000.....	11	14	12	23	24	41	50	53	41	23	4	-
1999.....	5	10	13	23	21	35	59	32	41	19	6	-
All races, male												
2004.....	3	3	-	4	4	4	18	10	4	2	-	-
2003.....	2	4	4	5	9	14	16	19	6	4	3	-
2002.....	3	4	3	4	11	14	20	18	4	5	-	-
2001.....	4	5	2	9	6	15	25	13	9	3	-	-
2000.....	7	9	5	12	8	20	18	15	9	6	-	-
1999.....	1	5	6	8	9	28	31	10	9	2	1	-
All races, female												
2004.....	2	-	4	7	4	16	17	21	8	9	8	-
2003.....	3	4	7	3	9	14	17	25	27	13	1	-
2002.....	3	5	4	4	11	17	19	14	23	9	5	-
2001.....	2	4	4	5	3	9	12	12	17	6	3	-
2000.....	4	5	7	11	16	21	32	38	32	17	4	-
1999.....	4	5	7	15	12	7	28	22	32	17	5	-
White, both sexes												
2004.....	5	2	2	10	8	20	35	29	11	11	8	-
2003.....	4	6	9	8	17	27	30	42	33	17	3	-
2002.....	6	8	7	7	21	29	38	30	27	12	4	-
2001.....	5	9	6	11	8	22	35	24	25	9	3	-
2000.....	9	13	9	22	21	40	46	50	41	23	3	-
1999.....	5	10	12	20	20	30	56	30	40	17	6	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Viral pneumonia, not elsewhere classified (J12)												
White, male												
2004.....	3	2	-	4	4	4	18	9	3	2	-	-
2003.....	2	3	2	5	8	13	14	19	6	4	2	-
2002.....	3	3	3	4	11	12	19	17	4	4	-	-
2001.....	3	5	2	8	5	14	24	12	8	3	-	-
2000.....	5	8	4	12	6	20	18	15	9	6	-	-
1999.....	1	5	6	8	9	23	29	9	9	2	1	-
White, female												
2004.....	2	-	2	6	4	16	17	20	8	9	8	-
2003.....	2	3	7	3	9	14	16	23	27	13	1	-
2002.....	3	5	4	3	10	17	19	13	23	8	4	-
2001.....	2	4	4	3	3	8	11	12	17	6	3	-
2000.....	4	5	5	10	15	20	28	35	32	17	3	-
1999.....	4	5	6	12	11	7	27	21	31	15	5	-
Black, both sexes												
2004.....	-	1	1	1	-	-	-	1	-	-	-	-
2003.....	1	1	2	-	-	-	2	1	-	-	1	-
2002.....	-	-	-	-	1	1	1	1	-	2	-	-
2001.....	-	-	-	2	1	-	1	1	-	-	-	-
2000.....	1	1	2	1	2	-	1	-	-	-	1	-
1999.....	-	-	-	2	-	3	2	2	1	1	-	-
Black, male												
2004.....	-	1	-	-	-	-	-	1	-	-	-	-
2003.....	-	-	2	-	-	-	2	-	-	-	1	-
2002.....	-	-	-	-	-	1	1	1	-	1	-	-
2001.....	-	-	-	1	1	-	-	1	-	-	-	-
2000.....	1	1	1	-	1	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	3	2	1	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Viral pneumonia, not elsewhere classified (J12)												
Black, female												
2004.....	-	-	1	1	-	-	-	-	-	-	-	-
2003.....	1	1	-	-	-	-	-	1	-	-	-	-
2002.....	-	-	-	-	1	-	-	-	-	1	-	-
2001.....	-	-	-	1	-	-	1	-	-	-	-	-
2000.....	-	-	1	1	1	-	1	-	-	-	1	-
1999.....	-	-	-	2	-	-	-	1	1	1	-	-
American Indian 1/, both sexes												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	1	-	-	-	-	-	1	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	1	-	-	1	2	-	-	-	-	-
1999.....	-	-	1	-	1	-	1	-	-	-	-	-
American Indian 1/, male												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	1	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, female												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	1	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	1	-	-	1	2	-	-	-	-	-
1999.....	-	-	1	-	1	-	1	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Viral pneumonia, not elsewhere classified (J12)												
Asian or Pacific Islander 2/, both sexes												
2004.....	-	-	1	-	-	-	-	1	1	-	-	-
2003.....	-	1	-	-	1	1	1	1	-	-	-	-
2002.....	-	-	-	1	-	1	-	-	-	-	1	-
2001.....	1	-	-	1	-	2	1	-	1	-	-	-
2000.....	1	-	-	-	1	-	1	3	-	-	-	-
1999.....	-	-	-	1	-	2	-	-	-	1	-	-
Asian or Pacific Islander 2/, male												
2004.....	-	-	-	-	-	-	-	-	1	-	-	-
2003.....	-	1	-	-	1	1	-	-	-	-	-	-
2002.....	-	-	-	-	-	1	-	-	-	-	-	-
2001.....	1	-	-	-	-	1	1	-	1	-	-	-
2000.....	1	-	-	-	1	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	2	-	-	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	-	-	1	-	-	-	-	1	-	-	-	-
2003.....	-	-	-	-	-	-	1	1	-	-	-	-
2002.....	-	-	-	1	-	-	-	-	-	-	1	-
2001.....	-	-	-	1	-	1	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	1	3	-	-	-	-
1999.....	-	-	-	1	-	-	-	-	-	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bacterial pneumonia (J13-J15)												
All races, both sexes												
2004.....	160	194	220	259	363	583	775	732	491	202	38	-
2003.....	193	172	215	285	402	610	821	799	572	229	46	-
2002.....	138	173	183	250	412	617	824	784	558	238	62	-
2001.....	138	194	182	253	383	627	761	752	524	214	44	-
2000.....	137	170	209	255	433	600	759	763	501	213	38	-
1999.....	99	139	165	225	381	528	664	639	421	153	39	1
All races, male												
2004.....	96	100	113	152	201	317	421	345	194	71	10	-
2003.....	122	99	126	171	242	327	422	379	211	64	6	-
2002.....	76	104	100	132	237	346	421	355	206	73	16	-
2001.....	80	111	104	158	219	358	397	331	199	70	8	-
2000.....	81	96	118	145	251	362	399	354	208	73	7	-
1999.....	67	78	82	131	216	272	331	284	141	48	9	-
All races, female												
2004.....	64	94	107	107	162	266	354	387	297	131	28	-
2003.....	71	73	89	114	160	283	399	420	361	165	40	-
2002.....	62	69	83	118	175	271	403	429	352	165	46	-
2001.....	58	83	78	95	164	269	364	421	325	144	36	-
2000.....	56	74	91	110	182	238	360	409	293	140	31	-
1999.....	32	61	83	94	165	256	333	355	280	105	30	1
White, both sexes												
2004.....	115	150	173	210	309	490	690	659	439	181	33	-
2003.....	149	129	166	241	349	530	743	720	519	205	43	-
2002.....	108	131	153	212	358	548	734	723	509	221	53	-
2001.....	111	165	141	213	326	536	689	674	488	198	35	-
2000.....	108	137	172	204	369	527	672	690	446	198	31	-
1999.....	78	109	131	193	323	468	604	586	380	139	35	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bacterial pneumonia (J13-J15)												
White, male												
2004.....	69	71	86	118	168	268	376	316	176	66	9	-
2003.....	93	70	91	145	208	285	387	343	194	57	6	-
2002.....	58	76	83	111	204	307	367	331	189	66	16	-
2001.....	59	94	81	134	182	300	360	295	180	65	8	-
2000.....	64	76	95	117	214	314	362	313	180	68	6	-
1999.....	50	60	67	112	183	241	298	255	124	45	9	-
White, female												
2004.....	46	79	87	92	141	222	314	343	263	115	24	-
2003.....	56	59	75	96	141	245	356	377	325	148	37	-
2002.....	50	55	70	101	154	241	367	392	320	155	37	-
2001.....	52	71	60	79	144	236	329	379	308	133	27	-
2000.....	44	61	77	87	155	213	310	377	266	130	25	-
1999.....	28	49	64	81	140	227	306	331	256	94	26	1
Black, both sexes												
2004.....	38	36	34	42	41	70	63	59	39	21	5	-
2003.....	35	34	40	32	37	63	51	60	41	18	1	-
2002.....	20	34	25	27	40	49	61	40	31	17	8	-
2001.....	24	23	34	31	45	71	52	62	26	10	8	-
2000.....	24	27	31	42	53	50	62	47	44	13	7	-
1999.....	18	28	28	25	51	46	42	40	29	7	3	-
Black, male												
2004.....	24	25	21	29	25	36	31	25	11	5	1	-
2003.....	22	23	28	18	23	35	19	24	13	3	-	-
2002.....	11	23	15	12	23	29	34	15	8	7	-	-
2001.....	18	13	19	20	30	46	23	27	14	3	-	-
2000.....	13	17	20	21	32	34	25	23	23	5	1	-
1999.....	14	17	11	15	29	23	24	19	12	2	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bacterial pneumonia (J13-J15)												
Black, female												
2004.....	14	11	13	13	16	34	32	34	28	16	4	-
2003.....	13	11	12	14	14	28	32	36	28	15	1	-
2002.....	9	11	10	15	17	20	27	25	23	10	8	-
2001.....	6	10	15	11	15	25	29	35	12	7	8	-
2000.....	11	10	11	21	21	16	37	24	21	8	6	-
1999.....	4	11	17	10	22	23	18	21	17	5	3	-
American Indian 1/, both sexes												
2004.....	3	5	7	3	1	4	4	1	2	-	-	-
2003.....	5	6	2	3	3	3	7	3	2	1	1	-
2002.....	3	4	1	2	2	1	5	2	4	-	-	-
2001.....	1	2	1	1	1	4	2	1	1	1	-	-
2000.....	3	2	2	2	2	6	3	5	2	-	-	-
1999.....	1	2	3	2	2	2	2	3	6	1	1	-
American Indian 1/, male												
2004.....	1	2	2	2	1	3	2	-	1	-	-	-
2003.....	4	4	2	2	1	1	3	2	2	1	-	-
2002.....	2	3	1	1	1	-	2	1	1	-	-	-
2001.....	1	1	-	1	1	2	1	1	1	-	-	-
2000.....	2	-	1	2	1	5	-	3	1	-	-	-
1999.....	1	1	1	-	1	1	-	2	3	-	-	-
American Indian 1/, female												
2004.....	2	3	5	1	-	1	2	1	1	-	-	-
2003.....	1	2	-	1	2	2	4	1	-	-	1	-
2002.....	1	1	-	1	1	1	3	1	3	-	-	-
2001.....	-	1	1	-	-	2	1	-	-	1	-	-
2000.....	1	2	1	-	1	1	3	2	1	-	-	-
1999.....	-	1	2	2	1	1	2	1	3	1	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bacterial pneumonia (J13-J15)												
Asian or Pacific Islander 2/, both sexes												
2004.....	4	3	6	4	12	19	18	13	11	-	-	-
2003.....	4	3	7	9	13	14	20	16	10	5	1	-
2002.....	7	4	4	9	12	19	24	19	14	-	1	-
2001.....	2	4	6	8	11	16	18	15	9	5	1	-
2000.....	2	4	4	7	9	17	22	21	9	2	-	-
1999.....	2	-	3	5	5	12	16	10	6	6	-	-
Asian or Pacific Islander 2/, male												
2004.....	2	2	4	3	7	10	12	4	6	-	-	-
2003.....	3	2	5	6	10	6	13	10	2	3	-	-
2002.....	5	2	1	8	9	10	18	8	8	-	-	-
2001.....	2	3	4	3	6	10	13	8	4	2	-	-
2000.....	2	3	2	5	4	9	12	15	4	-	-	-
1999.....	2	-	3	4	3	7	9	8	2	1	-	-
Asian or Pacific Islander 2/, female												
2004.....	2	1	2	1	5	9	6	9	5	-	-	-
2003.....	1	1	2	3	3	8	7	6	8	2	1	-
2002.....	2	2	3	1	3	9	6	11	6	-	1	-
2001.....	-	1	2	5	5	6	5	7	5	3	1	-
2000.....	-	1	2	2	5	8	10	6	5	2	-	-
1999.....	-	-	-	1	2	5	7	2	4	5	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonia due to other or unspecified organisms (J16,J18)												
All races, both sexes												
2004.....	918	1,150	1,528	2,069	3,573	6,299	10,063	11,348	9,477	4,264	1,144	-
2003.....	992	1,065	1,574	2,223	3,712	6,860	10,633	12,469	10,292	4,798	1,291	2
2002.....	878	1,076	1,509	2,143	3,956	7,198	11,098	12,987	10,761	4,861	1,262	3
2001.....	840	944	1,348	2,046	3,924	6,841	10,331	12,568	10,103	4,599	1,228	5
2000.....	819	975	1,406	2,179	4,076	7,082	10,797	12,564	10,471	4,750	1,271	3
1999.....	782	920	1,307	2,074	3,973	7,059	10,396	12,460	10,386	4,825	1,327	1
All races, male												
2004.....	572	677	905	1,206	1,980	3,288	4,840	4,761	3,214	1,177	256	-
2003.....	609	610	886	1,292	2,074	3,602	5,059	5,062	3,414	1,266	271	1
2002.....	535	627	880	1,235	2,199	3,726	5,287	5,277	3,620	1,264	237	2
2001.....	514	566	790	1,168	2,164	3,616	4,863	5,014	3,371	1,205	250	4
2000.....	500	585	823	1,247	2,244	3,621	5,135	5,079	3,421	1,259	260	2
1999.....	484	515	739	1,228	2,227	3,649	4,945	5,080	3,375	1,210	269	1
All races, female												
2004.....	346	473	623	863	1,593	3,011	5,223	6,587	6,263	3,087	888	-
2003.....	383	455	688	931	1,638	3,258	5,574	7,407	6,878	3,532	1,020	1
2002.....	343	449	629	908	1,757	3,472	5,811	7,710	7,141	3,597	1,025	1
2001.....	326	378	558	878	1,760	3,225	5,468	7,554	6,732	3,394	978	1
2000.....	319	390	583	932	1,832	3,461	5,662	7,485	7,050	3,491	1,011	1
1999.....	298	405	568	846	1,746	3,410	5,451	7,380	7,011	3,615	1,058	-
White, both sexes												
2004.....	680	898	1,235	1,661	3,040	5,503	9,005	10,340	8,682	3,925	1,019	-
2003.....	732	820	1,246	1,794	3,124	6,036	9,617	11,519	9,475	4,430	1,138	1
2002.....	661	831	1,202	1,786	3,399	6,335	10,042	11,914	9,935	4,479	1,141	3
2001.....	622	725	1,078	1,689	3,378	5,967	9,320	11,577	9,301	4,204	1,070	4
2000.....	597	745	1,095	1,734	3,439	6,281	9,776	11,584	9,698	4,382	1,135	1
1999.....	573	714	1,018	1,726	3,402	6,318	9,500	11,445	9,594	4,458	1,198	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonia due to other or unspecified organisms (J16,J18)												
White, male												
2004.....	424	543	729	951	1,686	2,862	4,337	4,345	2,952	1,071	217	-
2003.....	453	467	678	1,026	1,736	3,152	4,544	4,678	3,104	1,142	230	-
2002.....	404	484	693	1,026	1,907	3,278	4,782	4,832	3,324	1,142	213	2
2001.....	368	434	617	947	1,862	3,132	4,375	4,610	3,068	1,088	214	4
2000.....	353	434	644	976	1,890	3,214	4,627	4,681	3,143	1,155	229	1
1999.....	347	393	570	1,020	1,889	3,264	4,517	4,650	3,081	1,111	245	1
White, female												
2004.....	256	355	506	710	1,354	2,641	4,668	5,995	5,730	2,854	802	-
2003.....	279	353	568	768	1,388	2,884	5,073	6,841	6,371	3,288	908	1
2002.....	257	347	509	760	1,492	3,057	5,260	7,082	6,611	3,337	928	1
2001.....	254	291	461	742	1,516	2,835	4,945	6,967	6,233	3,116	856	-
2000.....	244	311	451	758	1,549	3,067	5,149	6,903	6,555	3,227	906	-
1999.....	226	321	448	706	1,513	3,054	4,983	6,795	6,513	3,347	953	-
Black, both sexes												
2004.....	219	224	252	340	432	613	802	741	558	262	96	-
2003.....	231	208	277	359	489	652	749	689	600	259	111	1
2002.....	191	222	275	303	471	667	820	818	620	293	99	-
2001.....	191	196	224	283	442	699	770	729	610	311	131	-
2000.....	201	210	261	368	515	645	798	763	605	298	113	2
1999.....	188	182	255	303	495	611	707	834	645	301	110	-
Black, male												
2004.....	137	119	155	223	242	328	383	289	174	66	29	-
2003.....	141	125	175	229	282	352	363	260	210	72	29	1
2002.....	113	135	165	178	246	347	384	304	201	85	19	-
2001.....	127	119	144	174	251	392	362	263	213	80	27	-
2000.....	132	137	148	226	286	327	382	295	195	78	24	1
1999.....	125	111	149	183	290	310	327	336	215	78	20	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonia due to other or unspecified organisms (J16,J18)												
Black, female												
2004.....	82	105	97	117	190	285	419	452	384	196	67	-
2003.....	90	83	102	130	207	300	386	429	390	187	82	-
2002.....	78	87	110	125	225	320	436	514	419	208	80	-
2001.....	64	77	80	109	191	307	408	466	397	231	104	-
2000.....	69	73	113	142	229	318	416	468	410	220	89	1
1999.....	63	71	106	120	205	301	380	498	430	223	90	-
American Indian 1/, both sexes												
2004.....	4	9	11	18	16	37	45	28	27	9	6	-
2003.....	12	15	26	27	22	26	43	47	38	9	11	-
2002.....	10	12	6	16	13	33	41	34	34	17	6	-
2001.....	13	7	15	18	22	33	46	40	30	16	2	-
2000.....	10	4	15	18	23	29	29	30	33	9	3	-
1999.....	11	10	14	13	18	34	31	43	32	17	3	-
American Indian 1/, male												
2004.....	4	6	3	8	7	18	23	13	12	4	3	-
2003.....	5	6	15	13	14	13	22	11	12	6	2	-
2002.....	5	3	4	9	7	18	12	15	13	4	2	-
2001.....	8	3	9	10	11	16	23	18	9	8	1	-
2000.....	9	3	12	13	13	13	15	8	14	4	-	-
1999.....	8	5	6	6	9	23	14	14	11	6	-	-
American Indian 1/, female												
2004.....	-	3	8	10	9	19	22	15	15	5	3	-
2003.....	7	9	11	14	8	13	21	36	26	3	9	-
2002.....	5	9	2	7	6	15	29	19	21	13	4	-
2001.....	5	4	6	8	11	17	23	22	21	8	1	-
2000.....	1	1	3	5	10	16	14	22	19	5	3	-
1999.....	3	5	8	7	9	11	17	29	21	11	3	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonia due to other or unspecified organisms (J16,J18)												
Asian or Pacific Islander 2/, both sexes												
2004.....	15	19	30	50	85	146	211	239	210	68	23	-
2003.....	17	22	25	43	77	146	224	214	179	100	31	-
2002.....	16	11	26	38	73	163	195	221	172	72	16	-
2001.....	14	16	31	56	82	142	195	222	162	68	25	1
2000.....	11	16	35	59	99	127	194	187	135	61	20	-
1999.....	10	14	20	32	58	96	158	138	115	49	16	-
Asian or Pacific Islander 2/, male												
2004.....	7	9	18	24	45	80	97	114	76	36	7	-
2003.....	10	12	18	24	42	85	130	113	88	46	10	-
2002.....	13	5	18	22	39	83	109	126	82	33	3	-
2001.....	11	10	20	37	40	76	103	123	81	29	8	-
2000.....	6	11	19	32	55	67	111	95	69	22	7	-
1999.....	4	6	14	19	39	52	87	80	68	15	4	-
Asian or Pacific Islander 2/, female												
2004.....	8	10	12	26	40	66	114	125	134	32	16	-
2003.....	7	10	7	19	35	61	94	101	91	54	21	-
2002.....	3	6	8	16	34	80	86	95	90	39	13	-
2001.....	3	6	11	19	42	66	92	99	81	39	17	1
2000.....	5	5	16	27	44	60	83	92	66	39	13	-
1999.....	6	8	6	13	19	44	71	58	47	34	12	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute lower respiratory infections (J20-J22)												
All races, both sexes												
2004.....	5	6	8	9	12	21	33	42	54	23	8	-
2003.....	12	11	10	13	17	35	45	57	60	32	15	-
2002.....	12	11	7	10	23	21	45	70	42	27	13	-
2001.....	10	4	12	10	18	24	34	60	55	25	5	-
2000.....	4	6	4	19	24	41	41	80	77	23	12	-
1999.....	8	8	13	14	21	44	63	86	96	46	21	-
All races, male												
2004.....	5	2	4	6	6	8	10	15	18	4	1	-
2003.....	6	6	6	11	6	17	18	14	15	4	2	-
2002.....	9	3	3	4	9	11	16	21	11	5	2	-
2001.....	6	1	5	3	7	10	10	15	10	4	1	-
2000.....	2	2	1	12	10	21	14	26	15	6	-	-
1999.....	3	4	6	6	11	22	29	19	18	6	5	-
All races, female												
2004.....	-	4	4	3	6	13	23	27	36	19	7	-
2003.....	6	5	4	2	11	18	27	43	45	28	13	-
2002.....	3	8	4	6	14	10	29	49	31	22	11	-
2001.....	4	3	7	7	11	14	24	45	45	21	4	-
2000.....	2	4	3	7	14	20	27	54	62	17	12	-
1999.....	5	4	7	8	10	22	34	67	78	40	16	-
White, both sexes												
2004.....	3	5	8	8	11	16	33	39	53	22	8	-
2003.....	11	9	9	11	15	34	42	56	59	30	14	-
2002.....	10	9	5	9	19	19	40	64	41	27	12	-
2001.....	6	3	11	9	13	21	33	58	52	23	5	-
2000.....	4	6	4	18	23	38	39	75	72	22	12	-
1999.....	5	6	11	11	16	41	58	81	93	44	19	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute lower respiratory infections (J20-J22)												
White, male												
2004.....	3	2	4	5	6	7	10	13	18	4	1	-
2003.....	6	5	5	9	6	16	17	14	15	4	2	-
2002.....	8	3	1	3	9	11	14	19	11	5	2	-
2001.....	3	1	4	3	5	8	10	14	9	4	1	-
2000.....	2	2	1	11	9	21	12	24	12	6	-	-
1999.....	3	3	6	4	8	21	27	16	17	6	4	-
White, female												
2004.....	-	3	4	3	5	9	23	26	35	18	7	-
2003.....	5	4	4	2	9	18	25	42	44	26	12	-
2002.....	2	6	4	6	10	8	26	45	30	22	10	-
2001.....	3	2	7	6	8	13	23	44	43	19	4	-
2000.....	2	4	3	7	14	17	27	51	60	16	12	-
1999.....	2	3	5	7	8	20	31	65	76	38	15	-
Black, both sexes												
2004.....	1	-	-	1	1	5	-	2	1	-	-	-
2003.....	1	2	-	2	2	1	2	1	1	2	1	-
2002.....	2	1	2	1	4	2	5	4	-	-	1	-
2001.....	4	1	-	-	4	-	-	2	3	2	-	-
2000.....	-	-	-	1	1	3	2	5	4	1	-	-
1999.....	3	1	1	3	5	2	4	4	2	2	2	-
Black, male												
2004.....	1	-	-	1	-	1	-	1	-	-	-	-
2003.....	-	1	-	2	-	1	1	-	-	-	-	-
2002.....	1	-	2	1	-	-	2	2	-	-	-	-
2001.....	3	-	-	-	1	-	-	1	1	-	-	-
2000.....	-	-	-	1	1	-	2	2	2	-	-	-
1999.....	-	1	-	2	3	-	1	2	-	-	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute bronchitis and bronchiolitis (J20-J21)												
All races, both sexes												
2004.....	4	6	4	8	10	14	18	25	29	13	2	-
2003.....	9	9	8	11	11	19	26	38	36	19	8	-
2002.....	11	8	5	10	13	15	24	39	31	15	9	-
2001.....	8	3	11	9	14	18	21	38	36	13	4	-
2000.....	1	4	3	15	16	30	23	45	44	15	7	-
1999.....	7	6	12	13	16	28	41	54	55	30	11	-
All races, male												
2004.....	4	2	4	5	5	4	3	11	12	4	-	-
2003.....	5	5	6	9	3	7	10	6	9	3	1	-
2002.....	8	3	2	4	6	8	10	12	7	3	1	-
2001.....	5	1	5	3	5	6	4	9	5	2	1	-
2000.....	1	2	1	9	5	18	9	14	10	5	-	-
1999.....	2	3	5	6	9	15	19	12	12	5	2	-
All races, female												
2004.....	-	4	-	3	5	10	15	14	17	9	2	-
2003.....	4	4	2	2	8	12	16	32	27	16	7	-
2002.....	3	5	3	6	7	7	14	27	24	12	8	-
2001.....	3	2	6	6	9	12	17	29	31	11	3	-
2000.....	-	2	2	6	11	12	14	31	34	10	7	-
1999.....	5	3	7	7	7	13	22	42	43	25	9	-
White, both sexes												
2004.....	3	5	4	7	9	10	18	22	29	12	2	-
2003.....	9	7	7	9	9	19	24	37	35	18	8	-
2002.....	9	6	4	9	12	14	22	34	30	15	8	-
2001.....	4	2	10	8	10	15	21	37	33	11	4	-
2000.....	1	4	3	14	15	27	22	42	42	14	7	-
1999.....	4	4	10	10	11	25	38	51	53	29	11	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute bronchitis and bronchiolitis (J20-J21)												
White, male												
2004.....	3	2	4	4	5	3	3	9	12	4	-	-
2003.....	5	4	5	7	3	7	9	6	9	3	1	-
2002.....	7	3	1	3	6	8	9	10	7	3	1	-
2001.....	2	1	4	3	3	4	4	9	4	2	1	-
2000.....	1	2	1	8	4	18	8	14	8	5	-	-
1999.....	2	2	5	4	6	14	18	10	11	5	2	-
White, female												
2004.....	-	3	-	3	4	7	15	13	17	8	2	-
2003.....	4	3	2	2	6	12	15	31	26	15	7	-
2002.....	2	3	3	6	6	6	13	24	23	12	7	-
2001.....	2	1	6	5	7	11	17	28	29	9	3	-
2000.....	-	2	2	6	11	9	14	28	34	9	7	-
1999.....	2	2	5	6	5	11	20	41	42	24	9	-
Black, both sexes												
2004.....	1	-	-	1	1	4	-	2	-	-	-	-
2003.....	-	2	-	2	2	-	1	1	1	1	-	-
2002.....	2	1	1	1	1	1	2	4	-	-	1	-
2001.....	4	1	-	-	3	-	-	1	3	2	-	-
2000.....	-	-	-	1	1	3	1	3	2	1	-	-
1999.....	3	1	1	3	5	2	2	3	1	1	-	-
Black, male												
2004.....	1	-	-	1	-	1	-	1	-	-	-	-
2003.....	-	1	-	2	-	-	1	-	-	-	-	-
2002.....	1	-	1	1	-	-	1	2	-	-	-	-
2001.....	3	-	-	-	1	-	-	-	1	-	-	-
2000.....	-	-	-	1	1	-	1	-	2	-	-	-
1999.....	-	1	-	2	3	-	-	2	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Unspecified acute lower respiratory infection (J22)												
All races, both sexes												
2004.....	1	-	4	1	2	7	15	17	25	10	6	-
2003.....	3	2	2	2	6	16	19	19	24	13	7	-
2002.....	1	3	2	-	10	6	21	31	11	12	4	-
2001.....	2	1	1	1	4	6	13	22	19	12	1	-
2000.....	3	2	1	4	8	11	18	35	33	8	5	-
1999.....	1	2	1	1	5	16	22	32	41	16	10	-
All races, male												
2004.....	1	-	-	1	1	4	7	4	6	-	1	-
2003.....	1	1	-	2	3	10	8	8	6	1	1	-
2002.....	1	-	1	-	3	3	6	9	4	2	1	-
2001.....	1	-	-	-	2	4	6	6	5	2	-	-
2000.....	1	-	-	3	5	3	5	12	5	1	-	-
1999.....	1	1	1	-	2	7	10	7	6	1	3	-
All races, female												
2004.....	-	-	4	-	1	3	8	13	19	10	5	-
2003.....	2	1	2	-	3	6	11	11	18	12	6	-
2002.....	-	3	1	-	7	3	15	22	7	10	3	-
2001.....	1	1	1	1	2	2	7	16	14	10	1	-
2000.....	2	2	1	1	3	8	13	23	28	7	5	-
1999.....	-	1	-	1	3	9	12	25	35	15	7	-
White, both sexes												
2004.....	-	-	4	1	2	6	15	17	24	10	6	-
2003.....	2	2	2	2	6	15	18	19	24	12	6	-
2002.....	1	3	1	-	7	5	18	30	11	12	4	-
2001.....	2	1	1	1	3	6	12	21	19	12	1	-
2000.....	3	2	1	4	8	11	17	33	30	8	5	-
1999.....	1	2	1	1	5	16	20	30	40	15	8	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Unspecified acute lower respiratory infection (J22)												
White, male												
2004.....	-	-	-	1	1	4	7	4	6	-	1	-
2003.....	1	1	-	2	3	9	8	8	6	1	1	-
2002.....	1	-	-	-	3	3	5	9	4	2	1	-
2001.....	1	-	-	-	2	4	6	5	5	2	-	-
2000.....	1	-	-	3	5	3	4	10	4	1	-	-
1999.....	1	1	1	-	2	7	9	6	6	1	2	-
White, female												
2004.....	-	-	4	-	1	2	8	13	18	10	5	-
2003.....	1	1	2	-	3	6	10	11	18	11	5	-
2002.....	-	3	1	-	4	2	13	21	7	10	3	-
2001.....	1	1	1	1	1	2	6	16	14	10	1	-
2000.....	2	2	1	1	3	8	13	23	26	7	5	-
1999.....	-	1	-	1	3	9	11	24	34	14	6	-
Black, both sexes												
2004.....	-	-	-	-	-	1	-	-	1	-	-	-
2003.....	1	-	-	-	-	1	1	-	-	1	1	-
2002.....	-	-	1	-	3	1	3	-	-	-	-	-
2001.....	-	-	-	-	1	-	-	1	-	-	-	-
2000.....	-	-	-	-	-	-	1	2	2	-	-	-
1999.....	-	-	-	-	-	-	2	1	1	1	2	-
Black, male												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	1	-	-	-	-	-	-
2002.....	-	-	1	-	-	-	1	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	1	-	-	-	-
2000.....	-	-	-	-	-	-	1	2	-	-	-	-
1999.....	-	-	-	-	-	-	1	-	-	-	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of upper respiratory tract (J30-J39)												
All races, both sexes												
2004.....	22	24	22	22	24	50	59	58	32	9	1	-
2003.....	15	24	33	31	33	51	54	60	34	12	2	-
2002.....	17	32	21	26	35	45	37	46	36	10	3	-
2001.....	23	13	26	27	33	47	69	51	41	8	4	-
2000.....	26	20	24	33	46	49	59	53	28	6	3	-
1999.....	19	19	29	37	38	63	55	54	31	15	-	-
All races, male												
2004.....	9	16	15	12	12	24	31	28	16	3	-	-
2003.....	7	13	18	20	19	29	31	25	13	5	1	-
2002.....	9	15	9	11	18	22	16	19	14	1	1	-
2001.....	11	4	16	11	14	21	37	23	19	1	1	-
2000.....	11	9	10	18	23	22	34	22	12	4	-	-
1999.....	13	10	17	22	20	32	24	26	11	5	-	-
All races, female												
2004.....	13	8	7	10	12	26	28	30	16	6	1	-
2003.....	8	11	15	11	14	22	23	35	21	7	1	-
2002.....	8	17	12	15	17	23	21	27	22	9	2	-
2001.....	12	9	10	16	19	26	32	28	22	7	3	-
2000.....	15	11	14	15	23	27	25	31	16	2	3	-
1999.....	6	9	12	15	18	31	31	28	20	10	-	-
White, both sexes												
2004.....	17	18	18	18	17	42	54	55	28	8	1	-
2003.....	12	18	28	27	27	43	53	54	31	11	2	-
2002.....	13	23	18	20	33	41	31	44	34	9	3	-
2001.....	18	9	23	23	22	41	63	47	34	8	3	-
2000.....	21	18	20	23	37	42	56	47	27	6	3	-
1999.....	16	15	23	28	34	55	49	47	29	13	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of upper respiratory tract (J30-J39)												
White, male												
2004.....	7	13	13	10	9	19	28	27	14	3	-	-
2003.....	6	10	18	18	14	25	31	24	12	5	1	-
2002.....	7	10	8	9	17	20	12	18	14	1	1	-
2001.....	8	1	13	9	12	16	33	21	15	1	1	-
2000.....	10	8	8	14	21	19	32	18	11	4	-	-
1999.....	11	7	15	16	19	29	24	22	9	4	-	-
White, female												
2004.....	10	5	5	8	8	23	26	28	14	5	1	-
2003.....	6	8	10	9	13	18	22	30	19	6	1	-
2002.....	6	13	10	11	16	21	19	26	20	8	2	-
2001.....	10	8	10	14	10	25	30	26	19	7	2	-
2000.....	11	10	12	9	16	23	24	29	16	2	3	-
1999.....	5	8	8	12	15	26	25	25	20	9	-	-
Black, both sexes												
2004.....	5	5	4	4	6	7	4	3	3	1	-	-
2003.....	3	6	5	3	5	7	1	3	2	1	-	-
2002.....	3	8	3	5	2	3	5	1	2	1	-	-
2001.....	4	3	2	2	10	6	6	4	7	-	1	-
2000.....	3	2	4	9	8	5	3	4	1	-	-	-
1999.....	3	4	6	8	4	5	5	6	1	2	-	-
Black, male												
2004.....	2	2	2	2	2	4	3	1	1	-	-	-
2003.....	1	3	-	1	4	3	-	-	-	-	-	-
2002.....	2	5	1	2	1	2	4	1	-	-	-	-
2001.....	2	2	2	1	1	5	4	2	4	-	-	-
2000.....	-	1	2	3	2	1	2	2	1	-	-	-
1999.....	2	3	2	5	1	2	-	3	1	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of upper respiratory tract (J30-J39)												
Black, female												
2004.....	3	3	2	2	4	3	1	2	2	1	-	-
2003.....	2	3	5	2	1	4	1	3	2	1	-	-
2002.....	1	3	2	3	1	1	1	-	2	1	-	-
2001.....	2	1	-	1	9	1	2	2	3	-	1	-
2000.....	3	1	2	6	6	4	1	2	-	-	-	-
1999.....	1	1	4	3	3	3	5	3	-	1	-	-
American Indian 1/, both sexes												
2004.....	-	-	-	-	1	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	1	1	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	1	-	-	-	-	-	-	-	-
2000.....	1	-	-	-	1	1	-	1	-	-	-	-
1999.....	-	-	-	1	-	-	1	-	-	-	-	-
American Indian 1/, male												
2004.....	-	-	-	-	1	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	1	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	1	-	1	-	-	-	-
1999.....	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, female												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	1	1	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	1	-	-	-	1	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	1	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of upper respiratory tract (J30-J39)												
Asian or Pacific Islander 2/, both sexes												
2004.....	-	1	-	-	-	1	1	-	1	-	-	-
2003.....	-	-	-	1	1	1	-	3	1	-	-	-
2002.....	-	-	-	1	-	1	1	1	-	-	-	-
2001.....	1	1	1	1	1	-	-	-	-	-	-	-
2000.....	1	-	-	1	-	1	-	1	-	-	-	-
1999.....	-	-	-	-	-	3	-	1	1	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	-	1	-	-	-	1	-	-	1	-	-	-
2003.....	-	-	-	1	1	1	-	1	1	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	1	1	1	-	1	-	-	-	-	-	-	-
2000.....	1	-	-	1	-	1	-	1	-	-	-	-
1999.....	-	-	-	-	-	1	-	1	1	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	-	-	-	-	-	-	1	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	2	-	-	-	-
2002.....	-	-	-	1	-	1	1	1	-	-	-	-
2001.....	-	-	-	1	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	2	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic lower respiratory diseases (J40-J47)												
All races, both sexes												
2004.....	2,279	4,205	7,549	11,323	17,067	23,398	24,170	17,842	8,601	2,459	337	1
2003.....	2,261	4,344	7,733	11,805	18,114	24,593	24,693	18,193	8,874	2,516	351	2
2002.....	2,249	4,220	7,060	11,405	18,383	24,495	24,746	17,971	8,508	2,430	375	7
2001.....	2,197	4,176	6,990	11,838	18,913	24,442	23,320	17,542	8,125	2,345	379	1
2000.....	2,077	3,985	6,754	11,833	19,324	24,309	23,413	17,085	7,823	2,238	350	2
1999.....	2,031	4,179	7,118	12,441	20,203	24,978	23,657	16,831	7,510	2,154	338	4
All races, male												
2004.....	1,193	2,155	3,874	5,843	8,660	11,737	11,518	7,964	3,345	832	90	1
2003.....	1,145	2,252	3,991	6,001	9,332	12,266	11,697	8,055	3,547	812	98	2
2002.....	1,178	2,141	3,642	5,838	9,501	12,457	11,975	8,140	3,448	762	86	5
2001.....	1,111	2,159	3,543	6,046	9,831	12,433	11,258	7,806	3,279	733	94	1
2000.....	1,044	2,072	3,492	6,149	10,061	12,305	11,533	7,843	3,165	776	92	2
1999.....	1,004	2,169	3,789	6,642	10,835	13,021	11,684	7,853	3,154	779	108	3
All races, female												
2004.....	1,086	2,050	3,675	5,480	8,407	11,661	12,652	9,878	5,256	1,627	247	-
2003.....	1,116	2,092	3,742	5,804	8,782	12,327	12,996	10,138	5,327	1,704	253	-
2002.....	1,071	2,079	3,418	5,567	8,882	12,038	12,771	9,831	5,060	1,668	289	2
2001.....	1,086	2,017	3,447	5,792	9,082	12,009	12,062	9,736	4,846	1,612	285	-
2000.....	1,033	1,913	3,262	5,684	9,263	12,004	11,880	9,242	4,658	1,462	258	-
1999.....	1,027	2,010	3,329	5,799	9,368	11,957	11,973	8,978	4,356	1,375	230	1
White, both sexes												
2004.....	1,884	3,661	6,822	10,403	15,870	21,891	22,851	16,883	8,123	2,299	297	1
2003.....	1,861	3,777	6,993	10,843	16,763	22,999	23,332	17,255	8,377	2,345	312	2
2002.....	1,842	3,709	6,344	10,447	17,054	22,937	23,321	17,020	8,047	2,258	337	5
2001.....	1,821	3,666	6,301	10,809	17,576	22,961	22,035	16,625	7,617	2,193	337	-
2000.....	1,702	3,469	6,063	10,793	17,972	22,832	22,193	16,186	7,360	2,070	310	1
1999.....	1,697	3,656	6,400	11,383	18,728	23,314	22,384	15,941	7,073	2,022	306	3

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic lower respiratory diseases (J40-J47)												
White, male												
2004.....	985	1,830	3,454	5,318	7,967	10,901	10,780	7,445	3,120	761	76	1
2003.....	946	1,942	3,569	5,471	8,531	11,312	10,880	7,556	3,301	744	86	2
2002.....	964	1,856	3,224	5,292	8,723	11,514	11,167	7,615	3,232	691	73	4
2001.....	911	1,887	3,156	5,465	9,034	11,524	10,518	7,317	3,039	666	76	-
2000.....	849	1,791	3,102	5,523	9,263	11,418	10,836	7,340	2,938	702	78	1
1999.....	827	1,885	3,362	6,017	9,956	12,010	10,934	7,376	2,922	716	97	2
White, female												
2004.....	899	1,831	3,368	5,085	7,903	10,990	12,071	9,438	5,003	1,538	221	-
2003.....	915	1,835	3,424	5,372	8,232	11,687	12,452	9,699	5,076	1,601	226	-
2002.....	878	1,853	3,120	5,155	8,331	11,423	12,154	9,405	4,815	1,567	264	1
2001.....	910	1,779	3,145	5,344	8,542	11,437	11,517	9,308	4,578	1,527	261	-
2000.....	853	1,678	2,961	5,270	8,709	11,414	11,357	8,846	4,422	1,368	232	-
1999.....	870	1,771	3,038	5,366	8,772	11,304	11,450	8,565	4,151	1,306	209	1
Black, both sexes												
2004.....	364	469	637	791	979	1,209	1,007	700	342	112	30	-
2003.....	360	494	624	825	1,109	1,245	1,071	681	365	129	35	-
2002.....	366	461	615	811	1,105	1,257	1,141	717	352	140	31	1
2001.....	334	448	583	873	1,112	1,191	1,021	700	377	117	30	-
2000.....	340	445	628	881	1,133	1,191	961	671	360	133	31	-
1999.....	304	473	618	905	1,207	1,358	1,022	701	343	111	25	1
Black, male												
2004.....	193	286	362	450	563	661	547	365	154	42	9	-
2003.....	175	271	351	455	647	716	639	349	171	45	12	-
2002.....	192	256	359	466	641	742	625	379	158	55	10	1
2001.....	174	237	326	500	660	710	584	363	166	51	12	-
2000.....	169	240	355	532	677	691	532	360	173	60	8	-
1999.....	160	255	374	537	718	815	598	365	172	52	10	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic lower respiratory diseases (J40-J47)												
Black, female												
2004.....	171	183	275	341	416	548	460	335	188	70	21	-
2003.....	185	223	273	370	462	529	432	332	194	84	23	-
2002.....	174	205	256	345	464	515	516	338	194	85	21	-
2001.....	160	211	257	373	452	481	437	337	211	66	18	-
2000.....	171	205	273	349	456	500	429	311	187	73	23	-
1999.....	144	218	244	368	489	543	424	336	171	59	15	-
American Indian 1/, both sexes												
2004.....	18	41	48	53	83	83	78	35	13	5	2	-
2003.....	23	38	54	59	71	99	64	50	16	6	-	-
2002.....	15	22	50	64	71	75	64	47	20	2	1	-
2001.....	10	28	44	54	68	82	47	44	19	3	2	1
2000.....	7	36	35	58	66	66	60	49	15	3	4	-
1999.....	11	18	45	44	79	88	51	36	10	1	2	-
American Indian 1/, male												
2004.....	9	18	28	26	42	39	36	12	7	4	1	-
2003.....	13	20	27	26	41	57	34	25	5	4	-	-
2002.....	7	11	24	30	34	40	37	20	8	-	-	-
2001.....	5	16	18	20	36	42	21	21	8	1	1	1
2000.....	5	19	22	24	26	39	38	24	8	1	3	-
1999.....	5	7	21	25	43	39	24	18	4	-	-	-
American Indian 1/, female												
2004.....	9	23	20	27	41	44	42	23	6	1	1	-
2003.....	10	18	27	33	30	42	30	25	11	2	-	-
2002.....	8	11	26	34	37	35	27	27	12	2	1	-
2001.....	5	12	26	34	32	40	26	23	11	2	1	-
2000.....	2	17	13	34	40	27	22	25	7	2	1	-
1999.....	6	11	24	19	36	49	27	18	6	1	2	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic lower respiratory diseases (J40-J47)												
Asian or Pacific Islander 2/, both sexes												
2004.....	13	34	42	76	135	215	234	224	123	43	8	-
2003.....	17	35	62	78	171	250	226	207	116	36	4	-
2002.....	26	28	51	83	153	226	220	187	89	30	6	1
2001.....	32	34	62	102	157	208	217	173	112	32	10	-
2000.....	28	35	28	101	153	220	199	179	88	32	5	1
1999.....	19	32	55	109	189	218	200	153	84	20	5	-
Asian or Pacific Islander 2/, male												
2004.....	6	21	30	49	88	136	155	142	64	25	4	-
2003.....	11	19	44	49	113	181	144	125	70	19	-	-
2002.....	15	18	35	50	103	161	146	126	50	16	3	-
2001.....	21	19	43	61	101	157	135	105	66	15	5	-
2000.....	21	22	13	70	95	157	127	119	46	13	3	1
1999.....	12	22	32	63	118	157	128	94	56	11	1	-
Asian or Pacific Islander 2/, female												
2004.....	7	13	12	27	47	79	79	82	59	18	4	-
2003.....	6	16	18	29	58	69	82	82	46	17	4	-
2002.....	11	10	16	33	50	65	74	61	39	14	3	1
2001.....	11	15	19	41	56	51	82	68	46	17	5	-
2000.....	7	13	15	31	58	63	72	60	42	19	2	-
1999.....	7	10	23	46	71	61	72	59	28	9	4	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)												
All races, both sexes												
2004.....	2,258	4,184	7,519	11,276	16,968	23,213	23,950	17,660	8,502	2,434	335	1
2003.....	2,252	4,316	7,697	11,752	18,021	24,421	24,469	18,031	8,781	2,491	349	2
2002.....	2,233	4,198	7,008	11,344	18,279	24,311	24,501	17,805	8,431	2,408	373	7
2001.....	2,179	4,153	6,945	11,767	18,809	24,270	23,120	17,354	8,034	2,320	374	1
2000.....	2,067	3,951	6,719	11,768	19,215	24,132	23,185	16,922	7,745	2,221	348	2
1999.....	2,013	4,148	7,078	12,365	20,107	24,789	23,428	16,657	7,445	2,139	335	4
All races, male												
2004.....	1,182	2,144	3,859	5,828	8,631	11,678	11,469	7,921	3,324	828	90	1
2003.....	1,139	2,247	3,974	5,990	9,303	12,215	11,631	8,016	3,532	806	98	2
2002.....	1,171	2,133	3,623	5,822	9,466	12,407	11,913	8,097	3,428	761	85	5
2001.....	1,105	2,147	3,520	6,021	9,798	12,383	11,216	7,765	3,259	732	93	1
2000.....	1,043	2,058	3,479	6,128	10,021	12,257	11,475	7,816	3,146	770	91	2
1999.....	1,001	2,159	3,774	6,625	10,797	12,945	11,611	7,801	3,132	775	107	3
All races, female												
2004.....	1,076	2,040	3,660	5,448	8,337	11,535	12,481	9,739	5,178	1,606	245	-
2003.....	1,113	2,069	3,723	5,762	8,718	12,206	12,838	10,015	5,249	1,685	251	-
2002.....	1,062	2,065	3,385	5,522	8,813	11,904	12,588	9,708	5,003	1,647	288	2
2001.....	1,074	2,006	3,425	5,746	9,011	11,887	11,904	9,589	4,775	1,588	281	-
2000.....	1,024	1,893	3,240	5,640	9,194	11,875	11,710	9,106	4,599	1,451	257	-
1999.....	1,012	1,989	3,304	5,740	9,310	11,844	11,817	8,856	4,313	1,364	228	1
White, both sexes												
2004.....	1,864	3,645	6,795	10,359	15,779	21,718	22,637	16,711	8,027	2,275	295	1
2003.....	1,856	3,757	6,960	10,800	16,679	22,838	23,121	17,098	8,287	2,320	310	2
2002.....	1,832	3,689	6,294	10,397	16,957	22,760	23,087	16,863	7,975	2,237	335	5
2001.....	1,805	3,649	6,262	10,747	17,481	22,801	21,846	16,444	7,531	2,169	332	-
2000.....	1,693	3,442	6,032	10,737	17,880	22,666	21,975	16,028	7,283	2,054	309	1
1999.....	1,681	3,630	6,364	11,316	18,641	23,143	22,171	15,770	7,010	2,008	303	3

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)												
White, male												
2004.....	974	1,821	3,441	5,304	7,939	10,847	10,732	7,403	3,100	757	76	1
2003.....	942	1,939	3,553	5,463	8,506	11,265	10,819	7,518	3,286	738	86	2
2002.....	960	1,849	3,206	5,280	8,694	11,466	11,108	7,577	3,213	690	72	4
2001.....	906	1,880	3,135	5,446	9,006	11,482	10,482	7,279	3,022	665	75	-
2000.....	848	1,778	3,091	5,509	9,228	11,376	10,778	7,313	2,920	696	78	1
1999.....	825	1,878	3,349	6,001	9,921	11,944	10,866	7,326	2,902	712	96	2
White, female												
2004.....	890	1,824	3,354	5,055	7,840	10,871	11,905	9,308	4,927	1,518	219	-
2003.....	914	1,818	3,407	5,337	8,173	11,573	12,302	9,580	5,001	1,582	224	-
2002.....	872	1,840	3,088	5,117	8,263	11,294	11,979	9,286	4,762	1,547	263	1
2001.....	899	1,769	3,127	5,301	8,475	11,319	11,364	9,165	4,509	1,504	257	-
2000.....	845	1,664	2,941	5,228	8,652	11,290	11,197	8,715	4,363	1,358	231	-
1999.....	856	1,752	3,015	5,315	8,720	11,199	11,305	8,444	4,108	1,296	207	1
Black, both sexes												
2004.....	364	466	635	789	974	1,205	1,004	696	341	112	30	-
2003.....	357	488	623	824	1,106	1,242	1,067	681	364	129	35	-
2002.....	362	460	613	806	1,101	1,254	1,135	715	351	139	31	1
2001.....	333	447	580	871	1,109	1,186	1,016	697	376	117	30	-
2000.....	339	440	625	874	1,127	1,188	958	669	360	133	31	-
1999.....	302	470	615	902	1,201	1,350	1,016	700	342	110	25	1
Black, male												
2004.....	193	284	361	450	562	661	547	364	153	42	9	-
2003.....	173	269	351	455	644	715	637	349	171	45	12	-
2002.....	190	255	358	464	637	740	623	379	158	55	10	1
2001.....	174	236	325	498	658	707	580	362	165	51	12	-
2000.....	169	240	353	527	673	688	532	360	173	60	8	-
1999.....	159	253	372	537	716	811	597	365	171	52	10	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)												
Black, female												
2004.....	171	182	274	339	412	544	457	332	188	70	21	-
2003.....	184	219	272	369	462	527	430	332	193	84	23	-
2002.....	172	205	255	342	464	514	512	336	193	84	21	-
2001.....	159	211	255	373	451	479	436	335	211	66	18	-
2000.....	170	200	272	347	454	500	426	309	187	73	23	-
1999.....	143	217	243	365	485	539	419	335	171	58	15	-
American Indian 1/, both sexes												
2004.....	17	41	48	53	83	82	78	35	13	5	2	-
2003.....	23	38	54	57	69	99	63	49	16	6	-	-
2002.....	15	21	50	62	70	74	64	47	18	2	1	-
2001.....	10	26	43	52	67	81	46	44	18	3	2	1
2000.....	7	35	35	58	66	65	59	49	15	3	4	-
1999.....	11	18	44	43	79	87	49	35	10	1	2	-
American Indian 1/, male												
2004.....	9	18	28	26	42	38	36	12	7	4	1	-
2003.....	13	20	27	26	40	57	34	25	5	4	-	-
2002.....	7	11	24	28	33	40	37	20	7	-	-	-
2001.....	5	15	17	18	35	41	21	21	8	1	1	1
2000.....	5	18	22	24	26	39	38	24	8	1	3	-
1999.....	5	7	21	25	43	38	24	17	4	-	-	-
American Indian 1/, female												
2004.....	8	23	20	27	41	44	42	23	6	1	1	-
2003.....	10	18	27	31	29	42	29	24	11	2	-	-
2002.....	8	10	26	34	37	34	27	27	11	2	1	-
2001.....	5	11	26	34	32	40	25	23	10	2	1	-
2000.....	2	17	13	34	40	26	21	25	7	2	1	-
1999.....	6	11	23	18	36	49	25	18	6	1	2	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)												
Asian or Pacific Islander 2/, both sexes												
2004.....	13	32	41	75	132	208	231	218	121	42	8	-
2003.....	16	33	60	71	167	242	218	203	114	36	4	-
2002.....	24	28	51	79	151	223	215	180	87	30	6	1
2001.....	31	31	60	97	152	202	212	169	109	31	10	-
2000.....	28	34	27	99	142	213	193	176	87	31	4	1
1999.....	19	30	55	104	186	209	192	152	83	20	5	-
Asian or Pacific Islander 2/, male												
2004.....	6	21	29	48	88	132	154	142	64	25	4	-
2003.....	11	19	43	46	113	178	141	124	70	19	-	-
2002.....	14	18	35	50	102	161	145	121	50	16	3	-
2001.....	20	16	43	59	99	153	133	103	64	15	5	-
2000.....	21	22	13	68	94	154	127	119	45	13	2	1
1999.....	12	21	32	62	117	152	124	93	55	11	1	-
Asian or Pacific Islander 2/, female												
2004.....	7	11	12	27	44	76	77	76	57	17	4	-
2003.....	5	14	17	25	54	64	77	79	44	17	4	-
2002.....	10	10	16	29	49	62	70	59	37	14	3	1
2001.....	11	15	17	38	53	49	79	66	45	16	5	-
2000.....	7	12	14	31	48	59	66	57	42	18	2	-
1999.....	7	9	23	42	69	57	68	59	28	9	4	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bronchitis, not specified as acute or chronic (J40)												
All races, both sexes												
2004.....	13	13	16	20	23	36	53	66	69	39	10	-
2003.....	4	12	13	15	28	52	58	88	80	37	12	-
2002.....	9	15	12	17	38	54	75	92	79	45	24	-
2001.....	13	12	15	23	38	72	68	104	74	39	13	-
2000.....	13	13	25	27	38	84	98	129	93	47	20	-
1999.....	13	13	28	33	60	70	106	109	83	49	16	-
All races, male												
2004.....	8	6	9	14	16	18	18	24	26	8	1	-
2003.....	1	5	8	7	15	25	23	29	20	7	1	-
2002.....	6	6	6	10	15	24	25	31	20	10	5	-
2001.....	6	9	9	11	19	32	24	35	20	10	1	-
2000.....	5	7	14	16	15	41	55	42	26	10	1	-
1999.....	7	8	13	15	29	35	42	37	22	10	2	-
All races, female												
2004.....	5	7	7	6	7	18	35	42	43	31	9	-
2003.....	3	7	5	8	13	27	35	59	60	30	11	-
2002.....	3	9	6	7	23	30	50	61	59	35	19	-
2001.....	7	3	6	12	19	40	44	69	54	29	12	-
2000.....	8	6	11	11	23	43	43	87	67	37	19	-
1999.....	6	5	15	18	31	35	64	72	61	39	14	-
White, both sexes												
2004.....	10	11	16	19	23	34	52	60	64	39	9	-
2003.....	3	7	12	14	26	51	52	81	75	32	11	-
2002.....	7	14	11	16	38	53	70	87	75	43	23	-
2001.....	12	12	12	19	34	65	64	102	71	36	12	-
2000.....	10	11	22	23	33	78	93	126	90	45	20	-
1999.....	8	11	25	28	52	58	98	100	80	49	13	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bronchitis, not specified as acute or chronic (J40)												
White, male												
2004.....	6	5	9	13	16	16	18	23	24	8	1	-
2003.....	1	4	7	7	13	24	23	25	19	7	1	-
2002.....	4	6	6	10	15	23	24	31	19	10	4	-
2001.....	5	9	7	8	16	27	21	35	19	7	1	-
2000.....	5	7	11	13	14	38	55	41	26	9	1	-
1999.....	5	7	11	13	25	29	36	36	21	10	1	-
White, female												
2004.....	4	6	7	6	7	18	34	37	40	31	8	-
2003.....	2	3	5	7	13	27	29	56	56	25	10	-
2002.....	3	8	5	6	23	30	46	56	56	33	19	-
2001.....	7	3	5	11	18	38	43	67	52	29	11	-
2000.....	5	4	11	10	19	40	38	85	64	36	19	-
1999.....	3	4	14	15	27	29	62	64	59	39	12	-
Black, both sexes												
2004.....	3	1	-	1	-	2	-	3	4	-	-	-
2003.....	1	4	1	1	-	-	4	6	4	4	1	-
2002.....	2	1	1	1	-	-	5	5	3	2	1	-
2001.....	1	-	3	4	4	6	2	2	1	3	-	-
2000.....	3	1	3	3	5	5	4	3	2	2	-	-
1999.....	4	2	2	4	6	8	8	8	1	-	2	-
Black, male												
2004.....	2	-	-	1	-	2	-	-	1	-	-	-
2003.....	-	1	1	-	-	-	-	3	-	-	-	-
2002.....	2	-	-	-	-	-	1	-	1	-	1	-
2001.....	1	-	2	3	3	4	2	-	-	3	-	-
2000.....	-	-	3	2	1	2	-	1	-	1	-	-
1999.....	2	1	1	1	3	4	6	1	-	-	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bronchitis, not specified as acute or chronic (J40)												
Black, female												
2004.....	1	1	-	-	-	-	-	3	3	-	-	-
2003.....	1	3	-	1	-	-	4	3	4	4	1	-
2002.....	-	1	1	1	-	-	4	5	2	2	-	-
2001.....	-	-	1	1	1	2	-	2	1	-	-	-
2000.....	3	1	-	1	4	3	4	2	2	1	-	-
1999.....	2	1	1	3	3	4	2	7	1	-	1	-
American Indian 1/, both sexes												
2004.....	-	-	-	-	-	-	-	1	-	-	-	-
2003.....	-	1	-	-	-	-	1	1	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	1	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	1	-	-	1	-	1	-	1	-	-	1	-
American Indian 1/, male												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	1	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	1	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, female												
2004.....	-	-	-	-	-	-	-	1	-	-	-	-
2003.....	-	1	-	-	-	-	1	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	1	-	-	-	-	1	-	1	-	-	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bronchitis, not specified as acute or chronic (J40)												
Asian or Pacific Islander 2/, both sexes												
2004.....	-	1	-	-	-	-	1	2	1	-	1	-
2003.....	-	-	-	-	2	1	1	-	1	1	-	-
2002.....	-	-	-	-	-	1	-	-	1	-	-	-
2001.....	-	-	-	-	-	-	2	-	2	-	1	-
2000.....	-	1	-	1	-	1	1	-	1	-	-	-
1999.....	-	-	1	-	2	3	-	-	2	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	-	1	-	-	-	-	-	1	1	-	-	-
2003.....	-	-	-	-	2	1	-	-	1	-	-	-
2002.....	-	-	-	-	-	1	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	1	-	1	-	-	-
2000.....	-	-	-	1	-	1	-	-	-	-	-	-
1999.....	-	-	1	-	1	2	-	-	1	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	-	-	-	-	-	-	1	1	-	-	1	-
2003.....	-	-	-	-	-	-	1	-	-	1	-	-
2002.....	-	-	-	-	-	-	-	-	1	-	-	-
2001.....	-	-	-	-	-	-	1	-	1	-	1	-
2000.....	-	1	-	-	-	-	1	-	1	-	-	-
1999.....	-	-	-	-	1	1	-	-	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic bronchitis (J41-J42)												
All races, both sexes												
2004.....	7	21	17	20	34	51	59	61	39	29	5	-
2003.....	14	9	13	25	36	55	81	63	49	14	2	-
2002.....	11	7	20	34	46	77	80	70	35	24	3	-
2001.....	8	13	21	33	39	57	64	84	49	28	10	-
2000.....	6	7	22	40	53	72	79	107	68	33	4	-
1999.....	11	22	34	30	50	69	94	84	65	33	7	-
All races, male												
2004.....	4	12	8	10	14	19	28	21	12	10	4	-
2003.....	9	7	6	10	14	34	31	22	15	6	1	-
2002.....	8	4	12	22	27	45	33	31	15	10	-	-
2001.....	3	7	11	24	22	35	24	35	13	4	1	-
2000.....	3	2	9	25	23	43	35	50	20	9	-	-
1999.....	3	16	14	21	29	38	42	34	23	10	4	-
All races, female												
2004.....	3	9	9	10	20	32	31	40	27	19	1	-
2003.....	5	2	7	15	22	21	50	41	34	8	1	-
2002.....	3	3	8	12	19	32	47	39	20	14	3	-
2001.....	5	6	10	9	17	22	40	49	36	24	9	-
2000.....	3	5	13	15	30	29	44	57	48	24	4	-
1999.....	8	6	20	9	21	31	52	50	42	23	3	-
White, both sexes												
2004.....	3	17	12	19	30	46	54	58	33	26	4	-
2003.....	10	7	12	24	30	49	73	58	46	14	2	-
2002.....	10	4	17	30	43	72	72	69	33	23	3	-
2001.....	8	12	21	30	33	53	60	80	45	27	10	-
2000.....	5	6	22	35	49	66	73	103	67	32	3	-
1999.....	7	20	32	30	45	65	85	81	63	30	7	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic bronchitis (J41-J42)												
White, male												
2004.....	-	11	5	10	12	18	27	18	12	8	3	-
2003.....	7	6	6	10	10	30	25	21	15	6	1	-
2002.....	7	2	9	18	25	42	29	30	13	10	-	-
2001.....	3	7	11	23	20	33	22	32	12	4	1	-
2000.....	2	1	9	24	20	40	32	47	20	9	-	-
1999.....	1	14	13	21	26	36	37	31	22	9	4	-
White, female												
2004.....	3	6	7	9	18	28	27	40	21	18	1	-
2003.....	3	1	6	14	20	19	48	37	31	8	1	-
2002.....	3	2	8	12	18	30	43	39	20	13	3	-
2001.....	5	5	10	7	13	20	38	48	33	23	9	-
2000.....	3	5	13	11	29	26	41	56	47	23	3	-
1999.....	6	6	19	9	19	29	48	50	41	21	3	-
Black, both sexes												
2004.....	4	3	5	-	4	3	3	3	5	2	-	-
2003.....	4	2	1	1	4	3	5	3	1	-	-	-
2002.....	1	2	1	3	2	4	4	1	2	-	-	-
2001.....	-	1	-	3	4	2	1	3	2	-	-	-
2000.....	1	-	-	4	4	5	6	2	1	1	1	-
1999.....	4	2	2	-	5	4	6	-	2	1	-	-
Black, male												
2004.....	4	-	3	-	2	-	-	3	-	1	-	-
2003.....	2	1	-	-	2	3	4	-	-	-	-	-
2002.....	1	1	1	3	1	3	1	1	2	-	-	-
2001.....	-	-	-	1	1	-	-	2	-	-	-	-
2000.....	1	-	-	-	3	2	3	1	-	-	-	-
1999.....	2	2	1	-	3	2	4	-	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic bronchitis (J41-J42)												
Asian or Pacific Islander 2/, both sexes												
2004.....	-	1	-	1	-	2	1	-	1	1	1	-
2003.....	-	-	-	-	2	2	3	2	2	-	-	-
2002.....	-	-	2	1	1	1	4	-	-	1	-	-
2001.....	-	-	-	-	2	2	3	1	2	1	-	-
2000.....	-	1	-	1	-	1	-	2	-	-	-	-
1999.....	-	-	-	-	-	-	3	3	-	2	-	-
Asian or Pacific Islander 2/, male												
2004.....	-	1	-	-	-	1	1	-	-	1	1	-
2003.....	-	-	-	-	2	1	2	1	-	-	-	-
2002.....	-	-	2	1	1	-	3	-	-	-	-	-
2001.....	-	-	-	-	1	2	2	1	1	-	-	-
2000.....	-	1	-	1	-	1	-	2	-	-	-	-
1999.....	-	-	-	-	-	-	1	3	-	1	-	-
Asian or Pacific Islander 2/, female												
2004.....	-	-	-	1	-	1	-	-	1	-	-	-
2003.....	-	-	-	-	-	1	1	1	2	-	-	-
2002.....	-	-	-	-	-	1	1	-	-	1	-	-
2001.....	-	-	-	-	1	-	1	-	1	1	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	2	-	-	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Emphysema (J43)												
All races, both sexes												
2004.....	283	564	1,019	1,571	2,222	2,782	2,542	1,629	629	152	17	-
2003.....	293	630	1,125	1,618	2,445	3,124	2,796	1,740	672	143	19	2
2002.....	298	643	1,060	1,665	2,723	3,202	2,907	1,826	726	157	15	1
2001.....	307	705	1,124	1,892	2,912	3,481	2,856	1,827	727	161	21	-
2000.....	312	680	1,127	1,976	3,144	3,522	2,989	1,832	685	150	18	-
1999.....	332	723	1,296	2,246	3,410	3,768	3,187	1,785	622	146	27	-
All races, male												
2004.....	157	304	569	851	1,200	1,442	1,283	779	285	53	5	-
2003.....	174	339	593	827	1,314	1,608	1,387	792	300	48	5	2
2002.....	172	334	591	875	1,460	1,723	1,505	866	329	52	4	1
2001.....	177	395	571	988	1,566	1,809	1,446	929	326	54	6	-
2000.....	179	366	577	1,077	1,661	1,839	1,534	898	317	58	5	-
1999.....	183	394	691	1,230	1,906	2,027	1,650	886	295	68	9	-
All races, female												
2004.....	126	260	450	720	1,022	1,340	1,259	850	344	99	12	-
2003.....	119	291	532	791	1,131	1,516	1,409	948	372	95	14	-
2002.....	126	309	469	790	1,263	1,479	1,402	960	397	105	11	-
2001.....	130	310	553	904	1,346	1,672	1,410	898	401	107	15	-
2000.....	133	314	550	899	1,483	1,683	1,455	934	368	92	13	-
1999.....	149	329	605	1,016	1,504	1,741	1,537	899	327	78	18	-
White, both sexes												
2004.....	232	501	930	1,468	2,091	2,634	2,418	1,546	604	137	13	-
2003.....	258	569	1,026	1,508	2,287	2,934	2,640	1,657	622	129	17	2
2002.....	261	586	974	1,536	2,545	3,031	2,750	1,731	689	153	12	1
2001.....	256	624	1,039	1,771	2,726	3,302	2,732	1,744	681	150	20	-
2000.....	262	609	1,036	1,819	2,964	3,342	2,851	1,731	644	134	16	-
1999.....	286	654	1,197	2,104	3,216	3,547	3,050	1,692	589	135	22	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Emphysema (J43)												
White, male												
2004.....	125	261	511	781	1,115	1,347	1,210	729	270	47	3	-
2003.....	148	300	531	765	1,215	1,480	1,294	741	264	42	4	2
2002.....	145	299	529	780	1,354	1,606	1,410	804	313	50	2	1
2001.....	146	344	515	920	1,442	1,698	1,368	874	305	51	6	-
2000.....	146	325	522	966	1,553	1,735	1,453	827	293	47	3	-
1999.....	148	348	627	1,135	1,765	1,895	1,554	836	273	63	7	-
White, female												
2004.....	107	240	419	687	976	1,287	1,208	817	334	90	10	-
2003.....	110	269	495	743	1,072	1,454	1,346	916	358	87	13	-
2002.....	116	287	445	756	1,191	1,425	1,340	927	376	103	10	-
2001.....	110	280	524	851	1,284	1,604	1,364	870	376	99	14	-
2000.....	116	284	514	853	1,411	1,607	1,398	904	351	87	13	-
1999.....	138	306	570	969	1,451	1,652	1,496	856	316	72	15	-
Black, both sexes												
2004.....	47	52	79	81	105	122	98	66	19	10	3	-
2003.....	29	57	85	97	124	148	120	57	37	11	1	-
2002.....	31	49	68	112	156	137	133	69	32	4	3	-
2001.....	46	70	72	107	149	126	97	65	40	9	1	-
2000.....	47	63	84	135	146	138	109	74	31	14	-	-
1999.....	43	60	86	127	154	176	108	76	28	11	3	-
Black, male												
2004.....	31	37	52	57	66	76	55	40	11	4	2	-
2003.....	20	37	53	54	75	95	68	36	27	4	1	-
2002.....	23	30	47	82	92	90	80	42	14	2	2	-
2001.....	27	44	47	62	94	74	61	40	18	2	-	-
2000.....	30	36	52	97	90	74	63	47	20	9	-	-
1999.....	33	38	56	88	109	103	75	38	17	5	2	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Emphysema (J43)												
Black, female												
2004.....	16	15	27	24	39	46	43	26	8	6	1	-
2003.....	9	20	32	43	49	53	52	21	10	7	-	-
2002.....	8	19	21	30	64	47	53	27	18	2	1	-
2001.....	19	26	25	45	55	52	36	25	22	7	1	-
2000.....	17	27	32	38	56	64	46	27	11	5	-	-
1999.....	10	22	30	39	45	73	33	38	11	6	1	-
American Indian 1/, both sexes												
2004.....	3	5	7	7	6	5	5	4	1	-	-	-
2003.....	3	2	6	2	7	11	7	5	2	-	-	-
2002.....	2	5	5	6	8	9	4	6	1	-	-	-
2001.....	-	5	5	7	12	15	4	3	1	-	-	-
2000.....	1	6	5	8	11	13	5	5	1	-	1	-
1999.....	2	4	3	7	9	9	4	4	-	-	-	-
American Indian 1/, male												
2004.....	1	2	4	3	4	5	1	1	1	-	-	-
2003.....	3	1	4	-	4	10	3	2	-	-	-	-
2002.....	-	3	4	4	4	7	4	4	-	-	-	-
2001.....	-	5	2	2	10	6	1	3	-	-	-	-
2000.....	1	3	2	3	3	7	2	4	1	-	1	-
1999.....	1	3	1	2	6	4	2	2	-	-	-	-
American Indian 1/, female												
2004.....	2	3	3	4	2	-	4	3	-	-	-	-
2003.....	-	1	2	2	3	1	4	3	2	-	-	-
2002.....	2	2	1	2	4	2	-	2	1	-	-	-
2001.....	-	-	3	5	2	9	3	-	1	-	-	-
2000.....	-	3	3	5	8	6	3	1	-	-	-	-
1999.....	1	1	2	5	3	5	2	2	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Emphysema (J43)												
Asian or Pacific Islander 2/, both sexes												
2004.....	1	6	3	15	20	21	21	13	5	5	1	-
2003.....	3	2	8	11	27	31	29	21	11	3	1	-
2002.....	4	3	13	11	14	25	20	20	4	-	-	-
2001.....	5	6	8	7	25	38	23	15	5	2	-	-
2000.....	2	2	2	14	23	29	24	22	9	2	1	-
1999.....	1	5	10	8	31	36	25	13	5	-	2	-
Asian or Pacific Islander 2/, male												
2004.....	-	4	2	10	15	14	17	9	3	2	-	-
2003.....	3	1	5	8	20	23	22	13	9	2	-	-
2002.....	4	2	11	9	10	20	11	16	2	-	-	-
2001.....	4	2	7	4	20	31	16	12	3	1	-	-
2000.....	2	2	1	11	15	23	16	20	3	2	1	-
1999.....	1	5	7	5	26	25	19	10	5	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	1	2	1	5	5	7	4	4	2	3	1	-
2003.....	-	1	3	3	7	8	7	8	2	1	1	-
2002.....	-	1	2	2	4	5	9	4	2	-	-	-
2001.....	1	4	1	3	5	7	7	3	2	1	-	-
2000.....	-	-	1	3	8	6	8	2	6	-	-	-
1999.....	-	-	3	3	5	11	6	3	-	-	2	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic obstructive pulmonary disease (J44)												
All races, both sexes												
2004.....	1,686	3,309	6,224	9,417	14,433	20,036	20,931	15,553	7,540	2,119	290	1
2003.....	1,626	3,366	6,283	9,832	15,242	20,826	21,146	15,805	7,744	2,222	291	-
2002.....	1,610	3,245	5,668	9,371	15,146	20,577	21,028	15,472	7,366	2,096	312	5
2001.....	1,527	3,158	5,497	9,531	15,474	20,276	19,714	14,968	6,945	2,015	308	1
2000.....	1,440	2,981	5,275	9,397	15,574	20,034	19,590	14,466	6,672	1,910	295	2
1999.....	1,362	3,047	5,480	9,703	16,167	20,428	19,570	14,298	6,451	1,814	267	4
All races, male												
2004.....	922	1,713	3,195	4,861	7,315	10,112	10,046	7,001	2,948	741	78	1
2003.....	846	1,779	3,270	5,066	7,867	10,432	10,072	7,095	3,140	731	86	-
2002.....	865	1,681	2,920	4,826	7,855	10,484	10,225	7,072	3,023	679	72	4
2001.....	803	1,652	2,831	4,894	8,075	10,391	9,613	6,680	2,843	653	82	1
2000.....	746	1,577	2,785	4,889	8,190	10,214	9,728	6,722	2,744	678	85	2
1999.....	707	1,622	2,968	5,244	8,681	10,701	9,748	6,763	2,740	670	90	3
All races, female												
2004.....	764	1,596	3,029	4,556	7,118	9,924	10,885	8,552	4,592	1,378	212	-
2003.....	780	1,587	3,013	4,766	7,375	10,394	11,074	8,710	4,604	1,491	205	-
2002.....	745	1,564	2,748	4,545	7,291	10,093	10,803	8,400	4,343	1,417	240	1
2001.....	724	1,506	2,666	4,637	7,399	9,885	10,101	8,288	4,102	1,362	226	-
2000.....	694	1,404	2,490	4,508	7,384	9,820	9,862	7,744	3,928	1,232	210	-
1999.....	655	1,425	2,512	4,459	7,486	9,727	9,822	7,535	3,711	1,144	177	1
White, both sexes												
2004.....	1,451	2,934	5,681	8,672	13,430	18,767	19,804	14,742	7,126	1,988	259	1
2003.....	1,405	2,978	5,727	9,060	14,141	19,513	20,028	14,999	7,333	2,074	258	-
2002.....	1,380	2,911	5,115	8,633	14,089	19,264	19,834	14,660	6,974	1,943	283	4
2001.....	1,318	2,829	5,013	8,712	14,426	19,069	18,629	14,193	6,521	1,885	272	-
2000.....	1,234	2,654	4,767	8,629	14,518	18,832	18,580	13,730	6,287	1,774	262	1
1999.....	1,197	2,716	4,949	8,900	15,012	19,091	18,522	13,570	6,084	1,706	244	3

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic obstructive pulmonary disease (J44)												
White, male												
2004.....	786	1,479	2,875	4,435	6,732	9,406	9,404	6,551	2,749	682	68	1
2003.....	725	1,553	2,949	4,621	7,201	9,648	9,376	6,662	2,936	670	75	-
2002.....	733	1,479	2,597	4,414	7,221	9,689	9,539	6,625	2,830	613	64	3
2001.....	683	1,463	2,548	4,419	7,444	9,633	8,982	6,262	2,639	594	66	-
2000.....	624	1,388	2,487	4,427	7,535	9,461	9,133	6,311	2,551	622	74	1
1999.....	608	1,428	2,643	4,755	7,999	9,869	9,123	6,362	2,546	616	82	2
White, female												
2004.....	665	1,455	2,806	4,237	6,698	9,361	10,400	8,191	4,377	1,306	191	-
2003.....	680	1,425	2,778	4,439	6,940	9,865	10,652	8,337	4,397	1,404	183	-
2002.....	647	1,432	2,518	4,219	6,868	9,575	10,295	8,035	4,144	1,330	219	1
2001.....	635	1,366	2,465	4,293	6,982	9,436	9,647	7,931	3,882	1,291	206	-
2000.....	610	1,266	2,280	4,202	6,983	9,371	9,447	7,419	3,736	1,152	188	-
1999.....	589	1,288	2,306	4,145	7,013	9,222	9,399	7,208	3,538	1,090	162	1
Black, both sexes												
2004.....	218	324	474	652	822	1,025	865	593	298	93	25	-
2003.....	200	343	477	662	921	1,043	897	597	305	110	31	-
2002.....	209	302	479	633	868	1,070	959	619	304	126	25	1
2001.....	187	292	417	699	882	993	875	593	320	102	26	-
2000.....	188	287	459	652	899	980	804	558	299	107	27	-
1999.....	149	310	459	695	954	1,107	855	579	289	93	20	1
Black, male												
2004.....	126	211	275	372	474	566	478	314	138	35	7	-
2003.....	107	203	272	381	549	597	554	307	141	40	11	-
2002.....	120	184	283	354	518	633	531	329	140	51	7	1
2001.....	104	169	247	414	531	611	505	312	144	45	11	-
2000.....	108	166	269	395	558	600	455	302	145	45	8	-
1999.....	88	182	287	422	565	682	505	318	146	46	7	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic obstructive pulmonary disease (J44)												
Black, female												
2004.....	92	113	199	280	348	459	387	279	160	58	18	-
2003.....	93	140	205	281	372	446	343	290	164	70	20	-
2002.....	89	118	196	279	350	437	428	290	164	75	18	-
2001.....	83	123	170	285	351	382	370	281	176	57	15	-
2000.....	80	121	190	257	341	380	349	256	154	62	19	-
1999.....	61	128	172	273	389	425	350	261	143	47	13	-
American Indian 1/, both sexes												
2004.....	12	35	39	45	77	75	72	30	12	5	2	-
2003.....	16	32	43	54	60	83	52	42	13	6	-	-
2002.....	13	14	43	51	60	63	58	41	16	2	1	-
2001.....	8	16	36	44	52	64	41	39	16	2	2	1
2000.....	6	23	29	47	52	51	53	42	14	3	3	-
1999.....	6	9	39	33	66	73	45	29	9	1	1	-
American Indian 1/, male												
2004.....	8	15	23	23	38	33	35	11	6	4	1	-
2003.....	10	17	20	26	36	45	29	21	5	4	-	-
2002.....	7	7	20	22	28	32	31	16	7	-	-	-
2001.....	5	8	13	15	25	34	20	18	7	1	1	1
2000.....	4	12	20	20	23	31	35	19	7	1	2	-
1999.....	4	2	20	21	35	32	22	14	3	-	-	-
American Indian 1/, female												
2004.....	4	20	16	22	39	42	37	19	6	1	1	-
2003.....	6	15	23	28	24	38	23	21	8	2	-	-
2002.....	6	7	23	29	32	31	27	25	9	2	1	-
2001.....	3	8	23	29	27	30	21	21	9	1	1	-
2000.....	2	11	9	27	29	20	18	23	7	2	1	-
1999.....	2	7	19	12	31	41	23	15	6	1	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic obstructive pulmonary disease (J44)												
Asian or Pacific Islander 2/, both sexes												
2004.....	5	16	30	48	104	169	190	188	104	33	4	-
2003.....	5	13	36	56	120	187	169	167	93	32	2	-
2002.....	8	18	31	54	129	180	177	152	72	25	3	-
2001.....	14	21	31	76	114	150	169	143	88	26	8	-
2000.....	12	17	20	69	105	171	153	136	72	26	3	1
1999.....	10	12	33	75	135	157	148	120	69	14	2	-
Asian or Pacific Islander 2/, male												
2004.....	2	8	22	31	71	107	129	125	55	20	2	-
2003.....	4	6	29	38	81	142	113	105	58	17	-	-
2002.....	5	11	20	36	88	130	124	102	46	15	1	-
2001.....	11	12	23	46	75	113	106	88	53	13	4	-
2000.....	10	11	9	47	74	122	105	90	41	10	1	1
1999.....	7	10	18	46	82	118	98	69	45	8	1	-
Asian or Pacific Islander 2/, female												
2004.....	3	8	8	17	33	62	61	63	49	13	2	-
2003.....	1	7	7	18	39	45	56	62	35	15	2	-
2002.....	3	7	11	18	41	50	53	50	26	10	2	-
2001.....	3	9	8	30	39	37	63	55	35	13	4	-
2000.....	2	6	11	22	31	49	48	46	31	16	2	-
1999.....	3	2	15	29	53	39	50	51	24	6	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Asthma (J45-J46)												
All races, both sexes												
2004.....	269	277	243	248	256	308	365	351	225	95	13	-
2003.....	315	299	263	262	270	364	388	335	236	75	25	-
2002.....	305	288	248	257	326	401	411	345	225	86	19	1
2001.....	324	265	288	288	346	384	418	371	239	77	22	-
2000.....	296	270	270	328	406	420	429	388	227	81	11	-
1999.....	295	343	240	353	420	454	471	381	224	97	18	-
All races, male												
2004.....	91	109	78	92	86	87	94	96	53	16	2	-
2003.....	109	117	97	80	93	116	118	78	57	14	5	-
2002.....	120	108	94	89	109	131	125	97	41	10	4	-
2001.....	116	84	98	104	116	116	109	86	57	11	3	-
2000.....	110	106	94	121	132	120	123	104	39	15	-	-
1999.....	101	119	88	115	152	144	129	81	52	17	2	-
All races, female												
2004.....	178	168	165	156	170	221	271	255	172	79	11	-
2003.....	206	182	166	182	177	248	270	257	179	61	20	-
2002.....	185	180	154	168	217	270	286	248	184	76	15	1
2001.....	208	181	190	184	230	268	309	285	182	66	19	-
2000.....	186	164	176	207	274	300	306	284	188	66	11	-
1999.....	194	224	152	238	268	310	342	300	172	80	16	-
White, both sexes												
2004.....	168	182	156	181	205	237	309	305	200	85	10	-
2003.....	180	196	183	194	195	291	328	303	211	71	22	-
2002.....	174	174	177	182	242	340	361	316	204	75	14	-
2001.....	211	172	177	215	262	312	361	325	213	71	18	-
2000.....	182	162	185	231	316	348	378	338	195	69	8	-
1999.....	183	229	161	254	316	382	416	327	194	88	17	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Asthma (J45-J46)												
White, male												
2004.....	57	65	41	65	64	60	73	82	45	12	1	-
2003.....	61	76	60	60	67	83	101	69	52	13	5	-
2002.....	71	63	65	58	79	106	106	87	38	7	2	-
2001.....	69	57	54	76	84	91	89	76	47	9	1	-
2000.....	71	57	62	79	106	102	105	87	30	9	-	-
1999.....	63	81	55	77	106	115	116	61	40	14	2	-
White, female												
2004.....	111	117	115	116	141	177	236	223	155	73	9	-
2003.....	119	120	123	134	128	208	227	234	159	58	17	-
2002.....	103	111	112	124	163	234	255	229	166	68	12	-
2001.....	142	115	123	139	178	221	272	249	166	62	17	-
2000.....	111	105	123	152	210	246	273	251	165	60	8	-
1999.....	120	148	106	177	210	267	300	266	154	74	15	-
Black, both sexes												
2004.....	92	86	77	55	43	53	38	31	15	7	2	-
2003.....	123	82	59	63	57	48	41	18	17	4	2	-
2002.....	119	106	64	57	75	43	34	21	10	7	2	-
2001.....	99	84	88	58	70	59	41	34	13	3	3	-
2000.....	100	89	79	80	73	60	35	32	27	9	3	-
1999.....	102	96	66	76	82	55	39	37	22	5	-	-
Black, male												
2004.....	30	36	31	20	20	17	14	7	3	2	-	-
2003.....	44	27	25	20	18	20	11	3	3	1	-	-
2002.....	44	40	27	25	26	14	10	7	1	2	-	-
2001.....	42	23	29	18	29	18	12	8	3	1	1	-
2000.....	30	38	29	33	21	10	11	9	8	5	-	-
1999.....	34	30	27	26	36	20	7	8	7	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Asthma (J45-J46)												
Black, female												
2004.....	62	50	46	35	23	36	24	24	12	5	2	-
2003.....	79	55	34	43	39	28	30	15	14	3	2	-
2002.....	75	66	37	32	49	29	24	14	9	5	2	-
2001.....	57	61	59	40	41	41	29	26	10	2	2	-
2000.....	70	51	50	47	52	50	24	23	19	4	3	-
1999.....	68	66	39	50	46	35	32	29	15	4	-	-
American Indian 1/, both sexes												
2004.....	2	1	2	1	-	2	-	-	-	-	-	-
2003.....	4	3	5	1	2	4	3	1	1	-	-	-
2002.....	-	1	2	5	2	2	2	-	1	-	-	-
2001.....	2	5	2	1	3	1	1	2	1	1	-	-
2000.....	-	6	1	3	3	1	1	2	-	-	-	-
1999.....	2	5	2	2	4	4	-	1	1	-	-	-
American Indian 1/, male												
2004.....	-	1	1	-	-	-	-	-	-	-	-	-
2003.....	-	2	3	-	-	2	2	1	-	-	-	-
2002.....	-	-	-	2	1	1	2	-	-	-	-	-
2001.....	-	2	2	1	-	-	-	-	1	-	-	-
2000.....	-	3	-	1	-	1	1	1	-	-	-	-
1999.....	-	2	-	1	2	2	-	1	1	-	-	-
American Indian 1/, female												
2004.....	2	-	1	1	-	2	-	-	-	-	-	-
2003.....	4	1	2	1	2	2	1	-	1	-	-	-
2002.....	-	1	2	3	1	1	-	-	1	-	-	-
2001.....	2	3	-	-	3	1	1	2	-	1	-	-
2000.....	-	3	-	2	3	-	-	1	-	-	-	-
1999.....	2	3	2	1	2	2	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Asthma (J45-J46)												
Asian or Pacific Islander 2/, both sexes												
2004.....	7	8	8	11	8	16	18	15	10	3	1	-
2003.....	8	18	16	4	16	21	16	13	7	-	1	-
2002.....	12	7	5	13	7	16	14	8	10	4	3	1
2001.....	12	4	21	14	11	12	15	10	12	2	1	-
2000.....	14	13	5	14	14	11	15	16	5	3	-	-
1999.....	8	13	11	21	18	13	16	16	7	4	1	-
Asian or Pacific Islander 2/, male												
2004.....	4	7	5	7	2	10	7	7	5	2	1	-
2003.....	4	12	9	-	8	11	4	5	2	-	-	-
2002.....	5	5	2	4	3	10	7	3	2	1	2	-
2001.....	5	2	13	9	3	7	8	2	6	1	1	-
2000.....	9	8	3	8	5	7	6	7	1	1	-	-
1999.....	4	6	6	11	8	7	6	11	4	2	-	-
Asian or Pacific Islander 2/, female												
2004.....	3	1	3	4	6	6	11	8	5	1	-	-
2003.....	4	6	7	4	8	10	12	8	5	-	1	-
2002.....	7	2	3	9	4	6	7	5	8	3	1	1
2001.....	7	2	8	5	8	5	7	8	6	1	-	-
2000.....	5	5	2	6	9	4	9	9	4	2	-	-
1999.....	4	7	5	10	10	6	10	5	3	2	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bronchiectasis (J47)												
All races, both sexes												
2004.....	21	21	30	47	99	185	220	182	99	25	2	-
2003.....	9	28	36	53	93	172	224	162	93	25	2	-
2002.....	16	22	52	61	104	184	245	166	77	22	2	-
2001.....	18	23	45	71	104	172	200	188	91	25	5	-
2000.....	10	34	35	65	109	177	228	163	78	17	2	-
1999.....	18	31	40	76	96	189	229	174	65	15	3	-
All races, male												
2004.....	11	11	15	15	29	59	49	43	21	4	-	-
2003.....	6	5	17	11	29	51	66	39	15	6	-	-
2002.....	7	8	19	16	35	50	62	43	20	1	1	-
2001.....	6	12	23	25	33	50	42	41	20	1	1	-
2000.....	1	14	13	21	40	48	58	27	19	6	1	-
1999.....	3	10	15	17	38	76	73	52	22	4	1	-
All races, female												
2004.....	10	10	15	32	70	126	171	139	78	21	2	-
2003.....	3	23	19	42	64	121	158	123	78	19	2	-
2002.....	9	14	33	45	69	134	183	123	57	21	1	-
2001.....	12	11	22	46	71	122	158	147	71	24	4	-
2000.....	9	20	22	44	69	129	170	136	59	11	1	-
1999.....	15	21	25	59	58	113	156	122	43	11	2	-
White, both sexes												
2004.....	20	16	27	44	91	173	214	172	96	24	2	-
2003.....	5	20	33	43	84	161	211	157	90	25	2	-
2002.....	10	20	50	50	97	177	234	157	72	21	2	-
2001.....	16	17	39	62	95	160	189	181	86	24	5	-
2000.....	9	27	31	56	92	166	218	158	77	16	1	-
1999.....	16	26	36	67	87	171	213	171	63	14	3	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bronchiectasis (J47)												
White, male												
2004.....	11	9	13	14	28	54	48	42	20	4	-	-
2003.....	4	3	16	8	25	47	61	38	15	6	-	-
2002.....	4	7	18	12	29	48	59	38	19	1	1	-
2001.....	5	7	21	19	28	42	36	38	17	1	1	-
2000.....	1	13	11	14	35	42	58	27	18	6	-	-
1999.....	2	7	13	16	35	66	68	50	20	4	1	-
White, female												
2004.....	9	7	14	30	63	119	166	130	76	20	2	-
2003.....	1	17	17	35	59	114	150	119	75	19	2	-
2002.....	6	13	32	38	68	129	175	119	53	20	1	-
2001.....	11	10	18	43	67	118	153	143	69	23	4	-
2000.....	8	14	20	42	57	124	160	131	59	10	1	-
1999.....	14	19	23	51	52	105	145	121	43	10	2	-
Black, both sexes												
2004.....	-	3	2	2	5	4	3	4	1	-	-	-
2003.....	3	6	1	1	3	3	4	-	1	-	-	-
2002.....	4	1	2	5	4	3	6	2	1	1	-	-
2001.....	1	1	3	2	3	5	5	3	1	-	-	-
2000.....	1	5	3	7	6	3	3	2	-	-	-	-
1999.....	2	3	3	3	6	8	6	1	1	1	-	-
Black, male												
2004.....	-	2	1	-	1	-	-	1	1	-	-	-
2003.....	2	2	-	-	3	1	2	-	-	-	-	-
2002.....	2	1	1	2	4	2	2	-	-	-	-	-
2001.....	-	1	1	2	2	3	4	1	1	-	-	-
2000.....	-	-	2	5	4	3	-	-	-	-	-	-
1999.....	1	2	2	-	2	4	1	-	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bronchiectasis (J47)												
Black, female												
2004.....	-	1	1	2	4	4	3	3	-	-	-	-
2003.....	1	4	1	1	-	2	2	-	1	-	-	-
2002.....	2	-	1	3	-	1	4	2	1	1	-	-
2001.....	1	-	2	-	1	2	1	2	-	-	-	-
2000.....	1	5	1	2	2	-	3	2	-	-	-	-
1999.....	1	1	1	3	4	4	5	1	-	1	-	-
American Indian 1/, both sexes												
2004.....	1	-	-	-	-	1	-	-	-	-	-	-
2003.....	-	-	-	2	2	-	1	1	-	-	-	-
2002.....	-	1	-	2	1	1	-	-	2	-	-	-
2001.....	-	2	1	2	1	1	1	-	1	-	-	-
2000.....	-	1	-	-	-	1	1	-	-	-	-	-
1999.....	-	-	1	1	-	1	2	1	-	-	-	-
American Indian 1/, male												
2004.....	-	-	-	-	-	1	-	-	-	-	-	-
2003.....	-	-	-	-	1	-	-	-	-	-	-	-
2002.....	-	-	-	2	1	-	-	-	1	-	-	-
2001.....	-	1	1	2	1	1	-	-	-	-	-	-
2000.....	-	1	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	1	-	1	-	-	-	-
American Indian 1/, female												
2004.....	1	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	2	1	-	1	1	-	-	-	-
2002.....	-	1	-	-	-	1	-	-	1	-	-	-
2001.....	-	1	-	-	-	-	1	-	1	-	-	-
2000.....	-	-	-	-	-	1	1	-	-	-	-	-
1999.....	-	-	1	1	-	-	2	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bronchiectasis (J47)												
Asian or Pacific Islander 2/, both sexes												
2004.....	-	2	1	1	3	7	3	6	2	1	-	-
2003.....	1	2	2	7	4	8	8	4	2	-	-	-
2002.....	2	-	-	4	2	3	5	7	2	-	-	-
2001.....	1	3	2	5	5	6	5	4	3	1	-	-
2000.....	-	1	1	2	11	7	6	3	1	1	1	-
1999.....	-	2	-	5	3	9	8	1	1	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	-	-	1	1	-	4	1	-	-	-	-	-
2003.....	-	-	1	3	-	3	3	1	-	-	-	-
2002.....	1	-	-	-	1	-	1	5	-	-	-	-
2001.....	1	3	-	2	2	4	2	2	2	-	-	-
2000.....	-	-	-	2	1	3	-	-	1	-	1	-
1999.....	-	1	-	1	1	5	4	1	1	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	-	2	-	-	3	3	2	6	2	1	-	-
2003.....	1	2	1	4	4	5	5	3	2	-	-	-
2002.....	1	-	-	4	1	3	4	2	2	-	-	-
2001.....	-	-	2	3	3	2	3	2	1	1	-	-
2000.....	-	1	1	-	10	4	6	3	-	1	-	-
1999.....	-	1	-	4	2	4	4	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Lung diseases due to external agents (J60-J70)												
All races, both sexes												
2004.....	246	341	483	693	1,314	2,341	3,697	4,042	2,904	1,128	241	-
2003.....	220	349	511	707	1,301	2,484	3,824	4,115	3,032	1,216	252	-
2002.....	227	305	456	723	1,324	2,592	3,811	4,319	3,072	1,236	231	-
2001.....	248	293	441	694	1,426	2,528	3,775	4,157	3,027	1,177	266	-
2000.....	219	282	379	737	1,463	2,520	3,489	4,048	2,864	1,131	239	-
1999.....	173	281	374	693	1,356	2,340	3,228	3,627	2,641	1,071	213	-
All races, male												
2004.....	157	210	317	465	841	1,425	2,078	2,080	1,302	404	75	-
2003.....	140	223	320	483	823	1,513	2,158	2,055	1,304	419	69	-
2002.....	150	190	279	464	848	1,584	2,128	2,202	1,342	432	56	-
2001.....	160	189	264	459	921	1,502	2,154	2,083	1,244	396	61	-
2000.....	134	173	238	491	982	1,541	1,978	2,050	1,199	386	65	-
1999.....	97	171	257	471	892	1,505	1,873	1,834	1,060	389	47	-
All races, female												
2004.....	89	131	166	228	473	916	1,619	1,962	1,602	724	166	-
2003.....	80	126	191	224	478	971	1,666	2,060	1,728	797	183	-
2002.....	77	115	177	259	476	1,008	1,683	2,117	1,730	804	175	-
2001.....	88	104	177	235	505	1,026	1,621	2,074	1,783	781	205	-
2000.....	85	109	141	246	481	979	1,511	1,998	1,665	745	174	-
1999.....	76	110	117	222	464	835	1,355	1,793	1,581	682	166	-
White, both sexes												
2004.....	196	286	377	593	1,127	2,085	3,366	3,733	2,665	1,022	214	-
2003.....	189	296	399	587	1,133	2,190	3,432	3,792	2,780	1,113	219	-
2002.....	176	260	375	598	1,151	2,294	3,488	3,997	2,846	1,119	207	-
2001.....	193	252	364	590	1,226	2,264	3,429	3,808	2,772	1,071	224	-
2000.....	175	238	307	620	1,243	2,251	3,171	3,748	2,642	1,044	211	-
1999.....	135	225	305	581	1,156	2,077	2,930	3,347	2,444	962	192	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Lung diseases due to external agents (J60-J70)												
White, male												
2004.....	129	174	244	396	722	1,278	1,905	1,925	1,199	362	71	-
2003.....	122	190	254	412	718	1,341	1,951	1,910	1,204	380	64	-
2002.....	115	161	236	390	740	1,409	1,964	2,059	1,235	393	54	-
2001.....	126	165	216	400	805	1,359	1,978	1,924	1,153	367	55	-
2000.....	106	151	195	410	833	1,373	1,826	1,924	1,106	358	60	-
1999.....	73	137	205	394	763	1,339	1,706	1,696	986	350	44	-
White, female												
2004.....	67	112	133	197	405	807	1,461	1,808	1,466	660	143	-
2003.....	67	106	145	175	415	849	1,481	1,882	1,576	733	155	-
2002.....	61	99	139	208	411	885	1,524	1,938	1,611	726	153	-
2001.....	67	87	148	190	421	905	1,451	1,884	1,619	704	169	-
2000.....	69	87	112	210	410	878	1,345	1,824	1,536	686	151	-
1999.....	62	88	100	187	393	738	1,224	1,651	1,458	612	148	-
Black, both sexes												
2004.....	39	52	93	92	164	209	275	242	181	86	22	-
2003.....	28	44	102	111	135	248	311	258	209	78	24	-
2002.....	45	38	68	113	148	256	270	264	191	97	18	-
2001.....	51	37	72	91	174	230	294	284	205	93	34	-
2000.....	42	40	65	107	201	234	263	243	184	75	21	-
1999.....	35	52	63	99	183	239	254	247	177	98	18	-
Black, male												
2004.....	23	34	63	63	104	117	142	115	71	33	3	-
2003.....	16	28	60	65	85	144	156	106	75	27	4	-
2002.....	31	25	37	70	92	147	130	109	87	33	2	-
2001.....	32	23	45	53	100	129	148	128	71	23	5	-
2000.....	28	20	37	72	135	146	123	97	71	22	4	-
1999.....	22	31	47	67	118	147	133	115	59	33	3	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Lung diseases due to external agents (J60-J70)												
Black, female												
2004.....	16	18	30	29	60	92	133	127	110	53	19	-
2003.....	12	16	42	46	50	104	155	152	134	51	20	-
2002.....	14	13	31	43	56	109	140	155	104	64	16	-
2001.....	19	14	27	38	74	101	146	156	134	70	29	-
2000.....	14	20	28	35	66	88	140	146	113	53	17	-
1999.....	13	21	16	32	65	92	121	132	118	65	15	-
American Indian 1/, both sexes												
2004.....	7	-	5	3	7	11	10	8	6	4	-	-
2003.....	-	3	4	4	10	8	18	11	7	2	3	-
2002.....	4	2	7	2	6	13	14	11	5	5	1	-
2001.....	3	-	2	4	7	9	14	8	7	1	2	-
2000.....	1	4	3	4	5	6	8	8	6	1	1	-
1999.....	1	3	4	6	4	6	5	13	3	2	-	-
American Indian 1/, male												
2004.....	3	-	3	2	2	7	5	4	2	1	-	-
2003.....	-	1	2	2	4	5	10	5	5	2	1	-
2002.....	2	2	3	-	5	9	8	8	2	3	-	-
2001.....	1	-	2	2	3	2	8	3	1	-	-	-
2000.....	-	2	2	4	3	5	3	4	3	1	-	-
1999.....	1	3	3	5	-	6	2	9	2	-	-	-
American Indian 1/, female												
2004.....	4	-	2	1	5	4	5	4	4	3	-	-
2003.....	-	2	2	2	6	3	8	6	2	-	2	-
2002.....	2	-	4	2	1	4	6	3	3	2	1	-
2001.....	2	-	-	2	4	7	6	5	6	1	2	-
2000.....	1	2	1	-	2	1	5	4	3	-	1	-
1999.....	-	-	1	1	4	-	3	4	1	2	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Lung diseases due to external agents (J60-J70)												
Asian or Pacific Islander 2/, both sexes												
2004.....	4	3	8	5	16	36	46	59	52	16	5	-
2003.....	3	6	6	5	23	38	63	54	36	23	6	-
2002.....	2	5	6	10	19	29	39	47	30	15	5	-
2001.....	1	4	3	9	19	25	38	57	43	12	6	-
2000.....	1	-	4	6	14	29	47	49	32	11	6	-
1999.....	2	1	2	7	13	18	39	20	17	9	3	-
Asian or Pacific Islander 2/, male												
2004.....	2	2	7	4	13	23	26	36	30	8	1	-
2003.....	2	4	4	4	16	23	41	34	20	10	-	-
2002.....	2	2	3	4	11	19	26	26	18	3	-	-
2001.....	1	1	1	4	13	12	20	28	19	6	1	-
2000.....	-	-	4	5	11	17	26	25	19	5	1	-
1999.....	1	-	2	5	11	13	32	14	13	6	-	-
Asian or Pacific Islander 2/, female												
2004.....	2	1	1	1	3	13	20	23	22	8	4	-
2003.....	1	2	2	1	7	15	22	20	16	13	6	-
2002.....	-	3	3	6	8	10	13	21	12	12	5	-
2001.....	-	3	2	5	6	13	18	29	24	6	5	-
2000.....	1	-	-	1	3	12	21	24	13	6	5	-
1999.....	1	1	-	2	2	5	7	6	4	3	3	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumoconioses and chemical effects (J60-J66,J68)												
All races, both sexes												
2004.....	13	25	45	82	127	211	231	185	92	11	2	-
2003.....	10	27	43	91	142	220	267	203	84	12	2	-
2002.....	12	26	49	80	143	256	224	217	76	17	2	-
2001.....	8	27	37	91	175	252	264	188	66	16	2	-
2000.....	7	21	36	85	196	249	235	217	80	20	-	-
1999.....	12	19	38	104	170	251	252	180	54	18	1	-
All races, male												
2004.....	12	25	43	80	124	204	221	178	87	10	2	-
2003.....	9	26	41	90	132	213	256	192	80	12	2	-
2002.....	9	24	45	77	137	245	212	207	75	17	1	-
2001.....	7	26	36	90	171	239	255	181	63	16	2	-
2000.....	7	19	35	79	186	241	226	208	79	19	-	-
1999.....	12	19	38	99	170	243	243	172	53	17	1	-
All races, female												
2004.....	1	-	2	2	3	7	10	7	5	1	-	-
2003.....	1	1	2	1	10	7	11	11	4	-	-	-
2002.....	3	2	4	3	6	11	12	10	1	-	1	-
2001.....	1	1	1	1	4	13	9	7	3	-	-	-
2000.....	-	2	1	6	10	8	9	9	1	1	-	-
1999.....	-	-	-	5	-	8	9	8	1	1	-	-
White, both sexes												
2004.....	12	24	41	75	119	207	219	176	83	10	2	-
2003.....	9	25	42	85	136	208	252	198	80	12	2	-
2002.....	11	24	43	77	136	239	212	212	71	14	1	-
2001.....	6	24	31	87	165	244	249	177	64	16	2	-
2000.....	6	21	34	80	186	234	222	210	76	18	-	-
1999.....	8	19	28	95	160	239	245	168	53	15	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumoconioses and chemical effects (J60-J66,J68)												
White, male												
2004.....	11	24	39	73	116	200	209	170	78	9	2	-
2003.....	8	24	40	84	126	201	242	188	76	12	2	-
2002.....	8	24	40	74	130	229	201	203	70	14	-	-
2001.....	6	23	30	86	161	231	240	170	61	16	2	-
2000.....	6	19	33	74	176	227	214	201	75	17	-	-
1999.....	8	19	28	91	160	231	236	161	52	14	-	-
White, female												
2004.....	1	-	2	2	3	7	10	6	5	1	-	-
2003.....	1	1	2	1	10	7	10	10	4	-	-	-
2002.....	3	-	3	3	6	10	11	9	1	-	1	-
2001.....	-	1	1	1	4	13	9	7	3	-	-	-
2000.....	-	2	1	6	10	7	8	9	1	1	-	-
1999.....	-	-	-	4	-	8	9	7	1	1	-	-
Black, both sexes												
2004.....	1	1	4	7	7	4	12	8	9	1	-	-
2003.....	1	1	-	6	5	12	11	5	4	-	-	-
2002.....	1	2	4	3	7	12	11	5	5	3	1	-
2001.....	2	3	5	4	8	8	13	10	2	-	-	-
2000.....	1	-	2	5	10	13	10	7	3	1	-	-
1999.....	4	-	8	8	8	9	7	11	1	3	1	-
Black, male												
2004.....	1	1	4	7	7	4	12	8	9	1	-	-
2003.....	1	1	-	6	5	12	10	4	4	-	-	-
2002.....	1	-	4	3	7	11	10	4	5	3	1	-
2001.....	1	3	5	4	8	8	13	10	2	-	-	-
2000.....	1	-	2	5	10	12	9	7	3	1	-	-
1999.....	4	-	8	7	8	9	7	10	1	3	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonitis due to solids and liquids (J69)												
All races, both sexes												
2004.....	226	312	431	598	1,169	2,107	3,457	3,850	2,808	1,117	239	-
2003.....	204	317	456	610	1,144	2,245	3,544	3,908	2,945	1,204	250	-
2002.....	208	271	400	627	1,166	2,323	3,577	4,100	2,995	1,219	229	-
2001.....	236	259	400	589	1,241	2,259	3,500	3,958	2,960	1,160	264	-
2000.....	209	257	341	642	1,257	2,258	3,247	3,825	2,779	1,111	239	-
1999.....	155	259	327	582	1,172	2,068	2,962	3,443	2,586	1,053	212	-
All races, male												
2004.....	143	183	268	376	709	1,207	1,852	1,898	1,211	394	73	-
2003.....	129	196	275	388	681	1,285	1,896	1,861	1,223	407	67	-
2002.....	136	162	231	380	703	1,332	1,911	1,994	1,267	415	55	-
2001.....	151	160	224	362	744	1,253	1,892	1,893	1,180	379	59	-
2000.....	126	153	201	405	790	1,291	1,747	1,839	1,116	367	65	-
1999.....	82	151	214	367	716	1,252	1,623	1,659	1,006	372	46	-
All races, female												
2004.....	83	129	163	222	460	900	1,605	1,952	1,597	723	166	-
2003.....	75	121	181	222	463	960	1,648	2,047	1,722	797	183	-
2002.....	72	109	169	247	463	991	1,666	2,106	1,728	804	174	-
2001.....	85	99	176	227	497	1,006	1,608	2,065	1,780	781	205	-
2000.....	83	104	140	237	467	967	1,500	1,986	1,663	744	174	-
1999.....	73	108	113	215	456	816	1,339	1,784	1,580	681	166	-
White, both sexes												
2004.....	178	258	330	505	992	1,855	3,139	3,550	2,578	1,012	212	-
2003.....	174	268	347	496	982	1,968	3,167	3,590	2,697	1,101	217	-
2002.....	158	228	326	505	1,001	2,042	3,266	3,783	2,774	1,105	206	-
2001.....	184	221	329	490	1,051	2,003	3,169	3,620	2,707	1,054	222	-
2000.....	166	214	271	530	1,050	2,006	2,942	3,532	2,562	1,026	211	-
1999.....	121	204	269	479	982	1,820	2,672	3,175	2,390	947	192	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonitis due to solids and liquids (J69)												
White, male												
2004.....	116	148	200	314	598	1,064	1,691	1,751	1,117	353	69	-
2003.....	112	165	210	323	582	1,129	1,703	1,720	1,127	368	62	-
2002.....	102	133	194	309	602	1,173	1,758	1,855	1,165	379	54	-
2001.....	118	139	182	307	638	1,118	1,731	1,745	1,091	350	53	-
2000.....	99	131	160	329	653	1,139	1,607	1,720	1,028	341	60	-
1999.....	62	118	173	298	597	1,099	1,463	1,532	933	336	44	-
White, female												
2004.....	62	110	130	191	394	791	1,448	1,799	1,461	659	143	-
2003.....	62	103	137	173	400	839	1,464	1,870	1,570	733	155	-
2002.....	56	95	132	196	399	869	1,508	1,928	1,609	726	152	-
2001.....	66	82	147	183	413	885	1,438	1,875	1,616	704	169	-
2000.....	67	83	111	201	397	867	1,335	1,812	1,534	685	151	-
1999.....	59	86	96	181	385	721	1,209	1,643	1,457	611	148	-
Black, both sexes												
2004.....	37	51	89	85	156	205	262	234	172	85	22	-
2003.....	27	42	100	105	130	234	300	253	205	78	24	-
2002.....	44	36	63	110	140	244	259	259	186	94	17	-
2001.....	48	34	67	86	166	222	281	274	203	93	34	-
2000.....	41	39	63	102	188	219	253	236	180	74	21	-
1999.....	31	51	54	91	175	227	246	236	176	95	17	-
Black, male												
2004.....	22	33	59	56	97	113	130	107	62	32	3	-
2003.....	15	27	60	59	80	130	146	102	71	27	4	-
2002.....	30	25	32	67	85	136	120	105	82	30	1	-
2001.....	31	20	40	49	92	121	135	118	69	23	5	-
2000.....	27	20	35	67	123	132	114	90	67	21	4	-
1999.....	18	30	38	60	110	137	126	105	58	30	2	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonitis due to solids and liquids (J69)												
Black, female												
2004.....	15	18	30	29	59	92	132	127	110	53	19	-
2003.....	12	15	40	46	50	104	154	151	134	51	20	-
2002.....	14	11	31	43	55	108	139	154	104	64	16	-
2001.....	17	14	27	37	74	101	146	156	134	70	29	-
2000.....	14	19	28	35	65	87	139	146	113	53	17	-
1999.....	13	21	16	31	65	90	120	131	118	65	15	-
American Indian 1/, both sexes												
2004.....	7	-	5	3	6	11	10	7	6	4	-	-
2003.....	-	3	4	4	9	7	15	11	7	2	3	-
2002.....	4	2	6	2	6	10	13	11	5	5	1	-
2001.....	3	-	1	4	6	9	12	8	7	1	2	-
2000.....	1	4	3	4	5	6	8	8	5	1	1	-
1999.....	1	3	2	5	4	3	5	12	3	2	-	-
American Indian 1/, male												
2004.....	3	-	3	2	2	7	5	4	2	1	-	-
2003.....	-	1	2	2	3	4	7	5	5	2	1	-
2002.....	2	2	2	-	5	6	7	8	2	3	-	-
2001.....	1	-	1	2	2	2	6	3	1	-	-	-
2000.....	-	2	2	4	3	5	3	4	2	1	-	-
1999.....	1	3	1	4	-	3	2	8	2	-	-	-
American Indian 1/, female												
2004.....	4	-	2	1	4	4	5	3	4	3	-	-
2003.....	-	2	2	2	6	3	8	6	2	-	2	-
2002.....	2	-	4	2	1	4	6	3	3	2	1	-
2001.....	2	-	-	2	4	7	6	5	6	1	2	-
2000.....	1	2	1	-	2	1	5	4	3	-	1	-
1999.....	-	-	1	1	4	-	3	4	1	2	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonitis due to solids and liquids (J69)												
Asian or Pacific Islander 2/, both sexes												
2004.....	4	3	7	5	15	36	46	59	52	16	5	-
2003.....	3	4	5	5	23	36	62	54	36	23	6	-
2002.....	2	5	5	10	19	27	39	47	30	15	5	-
2001.....	1	4	3	9	18	25	38	56	43	12	6	-
2000.....	1	-	4	6	14	27	44	49	32	10	6	-
1999.....	2	1	2	7	11	18	39	20	17	9	3	-
Asian or Pacific Islander 2/, male												
2004.....	2	2	6	4	12	23	26	36	30	8	1	-
2003.....	2	3	3	4	16	22	40	34	20	10	-	-
2002.....	2	2	3	4	11	17	26	26	18	3	-	-
2001.....	1	1	1	4	12	12	20	27	19	6	1	-
2000.....	-	-	4	5	11	15	23	25	19	4	1	-
1999.....	1	-	2	5	9	13	32	14	13	6	-	-
Asian or Pacific Islander 2/, female												
2004.....	2	1	1	1	3	13	20	23	22	8	4	-
2003.....	1	1	2	1	7	14	22	20	16	13	6	-
2002.....	-	3	2	6	8	10	13	21	12	12	5	-
2001.....	-	3	2	5	6	13	18	29	24	6	5	-
2000.....	1	-	-	1	3	12	21	24	13	6	5	-
1999.....	1	1	-	2	2	5	7	6	4	3	3	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonitis due to food and vomit (J69.0)												
All races, both sexes												
2004.....	226	312	431	598	1,169	2,106	3,456	3,849	2,808	1,117	239	-
2003.....	203	317	456	610	1,144	2,244	3,542	3,907	2,945	1,204	250	-
2002.....	208	271	400	625	1,166	2,322	3,574	4,100	2,995	1,219	229	-
2001.....	236	259	400	588	1,241	2,259	3,498	3,957	2,958	1,160	264	-
2000.....	209	256	340	642	1,257	2,257	3,247	3,825	2,778	1,111	239	-
1999.....	155	259	326	579	1,172	2,065	2,960	3,442	2,586	1,053	212	-
All races, male												
2004.....	143	183	268	376	709	1,206	1,851	1,898	1,211	394	73	-
2003.....	129	196	275	388	681	1,285	1,895	1,861	1,223	407	67	-
2002.....	136	162	231	379	703	1,332	1,911	1,994	1,267	415	55	-
2001.....	151	160	224	361	744	1,253	1,890	1,893	1,179	379	59	-
2000.....	126	152	201	405	790	1,291	1,747	1,839	1,115	367	65	-
1999.....	82	151	214	364	716	1,252	1,623	1,659	1,006	372	46	-
All races, female												
2004.....	83	129	163	222	460	900	1,605	1,951	1,597	723	166	-
2003.....	74	121	181	222	463	959	1,647	2,046	1,722	797	183	-
2002.....	72	109	169	246	463	990	1,663	2,106	1,728	804	174	-
2001.....	85	99	176	227	497	1,006	1,608	2,064	1,779	781	205	-
2000.....	83	104	139	237	467	966	1,500	1,986	1,663	744	174	-
1999.....	73	108	112	215	456	813	1,337	1,783	1,580	681	166	-
White, both sexes												
2004.....	178	258	330	505	992	1,854	3,139	3,549	2,578	1,012	212	-
2003.....	173	268	347	496	982	1,967	3,165	3,590	2,697	1,101	217	-
2002.....	158	228	326	503	1,001	2,041	3,264	3,783	2,774	1,105	206	-
2001.....	184	221	329	489	1,051	2,003	3,167	3,619	2,705	1,054	222	-
2000.....	166	213	270	530	1,050	2,005	2,942	3,532	2,561	1,026	211	-
1999.....	121	204	268	477	982	1,818	2,670	3,174	2,390	947	192	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonitis due to food and vomit (J69.0)												
White, male												
2004.....	116	148	200	314	598	1,063	1,691	1,751	1,117	353	69	-
2003.....	112	165	210	323	582	1,129	1,702	1,720	1,127	368	62	-
2002.....	102	133	194	308	602	1,173	1,758	1,855	1,165	379	54	-
2001.....	118	139	182	306	638	1,118	1,729	1,745	1,090	350	53	-
2000.....	99	130	160	329	653	1,139	1,607	1,720	1,027	341	60	-
1999.....	62	118	173	296	597	1,099	1,463	1,532	933	336	44	-
White, female												
2004.....	62	110	130	191	394	791	1,448	1,798	1,461	659	143	-
2003.....	61	103	137	173	400	838	1,463	1,870	1,570	733	155	-
2002.....	56	95	132	195	399	868	1,506	1,928	1,609	726	152	-
2001.....	66	82	147	183	413	885	1,438	1,874	1,615	704	169	-
2000.....	67	83	110	201	397	866	1,335	1,812	1,534	685	151	-
1999.....	59	86	95	181	385	719	1,207	1,642	1,457	611	148	-
Black, both sexes												
2004.....	37	51	89	85	156	205	261	234	172	85	22	-
2003.....	27	42	100	105	130	234	300	252	205	78	24	-
2002.....	44	36	63	110	140	244	258	259	186	94	17	-
2001.....	48	34	67	86	166	222	281	274	203	93	34	-
2000.....	41	39	63	102	188	219	253	236	180	74	21	-
1999.....	31	51	54	90	175	226	246	236	176	95	17	-
Black, male												
2004.....	22	33	59	56	97	113	129	107	62	32	3	-
2003.....	15	27	60	59	80	130	146	102	71	27	4	-
2002.....	30	25	32	67	85	136	120	105	82	30	1	-
2001.....	31	20	40	49	92	121	135	118	69	23	5	-
2000.....	27	20	35	67	123	132	114	90	67	21	4	-
1999.....	18	30	38	59	110	137	126	105	58	30	2	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonitis due to food and vomit (J69.0)												
Black, female												
2004.....	15	18	30	29	59	92	132	127	110	53	19	-
2003.....	12	15	40	46	50	104	154	150	134	51	20	-
2002.....	14	11	31	43	55	108	138	154	104	64	16	-
2001.....	17	14	27	37	74	101	146	156	134	70	29	-
2000.....	14	19	28	35	65	87	139	146	113	53	17	-
1999.....	13	21	16	31	65	89	120	131	118	65	15	-
American Indian 1/, both sexes												
2004.....	7	-	5	3	6	11	10	7	6	4	-	-
2003.....	-	3	4	4	9	7	15	11	7	2	3	-
2002.....	4	2	6	2	6	10	13	11	5	5	1	-
2001.....	3	-	1	4	6	9	12	8	7	1	2	-
2000.....	1	4	3	4	5	6	8	8	5	1	1	-
1999.....	1	3	2	5	4	3	5	12	3	2	-	-
American Indian 1/, male												
2004.....	3	-	3	2	2	7	5	4	2	1	-	-
2003.....	-	1	2	2	3	4	7	5	5	2	1	-
2002.....	2	2	2	-	5	6	7	8	2	3	-	-
2001.....	1	-	1	2	2	2	6	3	1	-	-	-
2000.....	-	2	2	4	3	5	3	4	2	1	-	-
1999.....	1	3	1	4	-	3	2	8	2	-	-	-
American Indian 1/, female												
2004.....	4	-	2	1	4	4	5	3	4	3	-	-
2003.....	-	2	2	2	6	3	8	6	2	-	2	-
2002.....	2	-	4	2	1	4	6	3	3	2	1	-
2001.....	2	-	-	2	4	7	6	5	6	1	2	-
2000.....	1	2	1	-	2	1	5	4	3	-	1	-
1999.....	-	-	1	1	4	-	3	4	1	2	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonitis due to food and vomit (J69.0)												
Asian or Pacific Islander 2/, both sexes												
2004.....	4	3	7	5	15	36	46	59	52	16	5	-
2003.....	3	4	5	5	23	36	62	54	36	23	6	-
2002.....	2	5	5	10	19	27	39	47	30	15	5	-
2001.....	1	4	3	9	18	25	38	56	43	12	6	-
2000.....	1	-	4	6	14	27	44	49	32	10	6	-
1999.....	2	1	2	7	11	18	39	20	17	9	3	-
Asian or Pacific Islander 2/, male												
2004.....	2	2	6	4	12	23	26	36	30	8	1	-
2003.....	2	3	3	4	16	22	40	34	20	10	-	-
2002.....	2	2	3	4	11	17	26	26	18	3	-	-
2001.....	1	1	1	4	12	12	20	27	19	6	1	-
2000.....	-	-	4	5	11	15	23	25	19	4	1	-
1999.....	1	-	2	5	9	13	32	14	13	6	-	-
Asian or Pacific Islander 2/, female												
2004.....	2	1	1	1	3	13	20	23	22	8	4	-
2003.....	1	1	2	1	7	14	22	20	16	13	6	-
2002.....	-	3	2	6	8	10	13	21	12	12	5	-
2001.....	-	3	2	5	6	13	18	29	24	6	5	-
2000.....	1	-	-	1	3	12	21	24	13	6	5	-
1999.....	1	1	-	2	2	5	7	6	4	3	3	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pneumonitis due to solids and liquids (J69.1-J69.8)												
All races, both sexes												
2004.....	-	-	-	-	-	1	1	1	-	-	-	-
2003.....	1	-	-	-	-	1	2	1	-	-	-	-
2002.....	-	-	-	2	-	1	3	-	-	-	-	-
2001.....	-	-	-	1	-	-	2	1	2	-	-	-
2000.....	-	1	1	-	-	1	-	-	1	-	-	-
1999.....	-	-	1	3	-	3	2	1	-	-	-	-
All races, male												
2004.....	-	-	-	-	-	1	1	-	-	-	-	-
2003.....	-	-	-	-	-	-	1	-	-	-	-	-
2002.....	-	-	-	1	-	-	-	-	-	-	-	-
2001.....	-	-	-	1	-	-	2	-	1	-	-	-
2000.....	-	1	-	-	-	-	-	-	1	-	-	-
1999.....	-	-	-	3	-	-	-	-	-	-	-	-
All races, female												
2004.....	-	-	-	-	-	-	-	1	-	-	-	-
2003.....	1	-	-	-	-	1	1	1	-	-	-	-
2002.....	-	-	-	1	-	1	3	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	1	1	-	-	-
2000.....	-	-	1	-	-	1	-	-	-	-	-	-
1999.....	-	-	1	-	-	3	2	1	-	-	-	-
White, both sexes												
2004.....	-	-	-	-	-	1	-	1	-	-	-	-
2003.....	1	-	-	-	-	1	2	-	-	-	-	-
2002.....	-	-	-	2	-	1	2	-	-	-	-	-
2001.....	-	-	-	1	-	-	2	1	2	-	-	-
2000.....	-	1	-	-	-	1	-	-	1	-	-	-
1999.....	-	-	1	2	-	2	2	1	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other lung diseases due to external agents (J67,J70)												
All races, both sexes												
2004.....	7	4	7	13	18	23	9	7	4	-	-	-
2003.....	6	5	12	6	15	19	13	4	3	-	-	-
2002.....	7	8	7	16	15	13	10	2	1	-	-	-
2001.....	4	7	4	14	10	17	11	11	1	1	-	-
2000.....	3	4	2	10	10	13	7	6	5	-	-	-
1999.....	6	3	9	7	14	21	14	4	1	-	-	-
All races, male												
2004.....	2	2	6	9	8	14	5	4	4	-	-	-
2003.....	2	1	4	5	10	15	6	2	1	-	-	-
2002.....	5	4	3	7	8	7	5	1	-	-	-	-
2001.....	2	3	4	7	6	10	7	9	1	1	-	-
2000.....	1	1	2	7	6	9	5	3	4	-	-	-
1999.....	3	1	5	5	6	10	7	3	1	-	-	-
All races, female												
2004.....	5	2	1	4	10	9	4	3	-	-	-	-
2003.....	4	4	8	1	5	4	7	2	2	-	-	-
2002.....	2	4	4	9	7	6	5	1	1	-	-	-
2001.....	2	4	-	7	4	7	4	2	-	-	-	-
2000.....	2	3	-	3	4	4	2	3	1	-	-	-
1999.....	3	2	4	2	8	11	7	1	-	-	-	-
White, both sexes												
2004.....	6	4	6	13	16	23	8	7	4	-	-	-
2003.....	6	3	10	6	15	14	13	4	3	-	-	-
2002.....	7	8	6	16	14	13	10	2	1	-	-	-
2001.....	3	7	4	13	10	17	11	11	1	1	-	-
2000.....	3	3	2	10	7	11	7	6	4	-	-	-
1999.....	6	2	8	7	14	18	13	4	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other lung diseases due to external agents (J67,J70)												
White, male												
2004.....	2	2	5	9	8	14	5	4	4	-	-	-
2003.....	2	1	4	5	10	11	6	2	1	-	-	-
2002.....	5	4	2	7	8	7	5	1	-	-	-	-
2001.....	2	3	4	7	6	10	7	9	1	1	-	-
2000.....	1	1	2	7	4	7	5	3	3	-	-	-
1999.....	3	-	4	5	6	9	7	3	1	-	-	-
White, female												
2004.....	4	2	1	4	8	9	3	3	-	-	-	-
2003.....	4	2	6	1	5	3	7	2	2	-	-	-
2002.....	2	4	4	9	6	6	5	1	1	-	-	-
2001.....	1	4	-	6	4	7	4	2	-	-	-	-
2000.....	2	2	-	3	3	4	2	3	1	-	-	-
1999.....	3	2	4	2	8	9	6	1	-	-	-	-
Black, both sexes												
2004.....	1	-	-	-	1	-	1	-	-	-	-	-
2003.....	-	1	2	-	-	2	-	-	-	-	-	-
2002.....	-	-	1	-	1	-	-	-	-	-	-	-
2001.....	1	-	-	1	-	-	-	-	-	-	-	-
2000.....	-	1	-	-	3	2	-	-	1	-	-	-
1999.....	-	1	1	-	-	3	1	-	-	-	-	-
Black, male												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	2	-	-	-	-	-	-
2002.....	-	-	1	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	2	2	-	-	1	-	-	-
1999.....	-	1	1	-	-	1	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Suppurative and necrotic conditions of lower respiratory tract (J85-J86)												
All races, both sexes												
2004.....	52	53	49	52	68	74	88	73	31	3	1	-
2003.....	56	59	57	59	81	82	91	63	25	9	1	-
2002.....	51	44	56	59	83	95	99	68	26	6	5	-
2001.....	61	52	56	66	81	115	94	85	38	8	2	-
2000.....	51	39	51	59	95	108	81	82	32	6	3	-
1999.....	59	53	53	92	89	99	81	66	36	12	1	-
All races, male												
2004.....	39	36	32	37	45	34	57	42	13	1	1	-
2003.....	41	39	34	41	48	38	46	31	13	4	-	-
2002.....	36	31	43	38	52	57	61	38	6	1	2	-
2001.....	47	31	38	38	49	68	49	41	17	2	1	-
2000.....	35	31	31	42	54	67	39	51	16	1	3	-
1999.....	43	34	36	70	57	52	50	28	14	5	1	-
All races, female												
2004.....	13	17	17	15	23	40	31	31	18	2	-	-
2003.....	15	20	23	18	33	44	45	32	12	5	1	-
2002.....	15	13	13	21	31	38	38	30	20	5	3	-
2001.....	14	21	18	28	32	47	45	44	21	6	1	-
2000.....	16	8	20	17	41	41	42	31	16	5	-	-
1999.....	16	19	17	22	32	47	31	38	22	7	-	-
White, both sexes												
2004.....	38	43	41	44	59	61	77	62	29	3	1	-
2003.....	45	46	44	51	70	69	75	60	20	9	1	-
2002.....	38	35	43	52	69	74	87	58	23	6	3	-
2001.....	42	49	45	55	69	101	84	75	35	7	2	-
2000.....	45	29	41	46	80	98	73	68	28	6	1	-
1999.....	50	30	42	72	78	89	74	55	30	11	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Suppurative and necrotic conditions of lower respiratory tract (J85-J86)												
White, male												
2004.....	32	29	26	31	40	29	49	37	13	1	1	-
2003.....	32	33	27	34	42	35	34	29	9	4	-	-
2002.....	29	25	32	35	44	43	55	33	6	1	-	-
2001.....	33	30	30	28	39	58	42	37	15	2	1	-
2000.....	30	24	23	31	49	61	34	44	15	1	1	-
1999.....	38	17	29	56	51	45	48	21	11	4	-	-
White, female												
2004.....	6	14	15	13	19	32	28	25	16	2	-	-
2003.....	13	13	17	17	28	34	41	31	11	5	1	-
2002.....	9	10	11	17	25	31	32	25	17	5	3	-
2001.....	9	19	15	27	30	43	42	38	20	5	1	-
2000.....	15	5	18	15	31	37	39	24	13	5	-	-
1999.....	12	13	13	16	27	44	26	34	19	7	-	-
Black, both sexes												
2004.....	12	9	8	6	6	12	4	9	2	-	-	-
2003.....	11	11	9	7	10	10	13	3	5	-	-	-
2002.....	10	9	10	6	10	19	8	8	3	-	2	-
2001.....	15	2	11	9	8	11	8	6	3	1	-	-
2000.....	6	10	8	12	13	9	6	12	4	-	2	-
1999.....	6	22	10	15	9	8	7	8	5	-	1	-
Black, male												
2004.....	6	6	6	4	4	5	3	4	-	-	-	-
2003.....	9	6	5	6	6	2	9	2	4	-	-	-
2002.....	7	6	8	3	5	13	4	3	-	-	2	-
2001.....	10	1	8	8	6	7	6	2	2	-	-	-
2000.....	5	7	7	10	4	5	3	6	1	-	2	-
1999.....	4	16	7	11	5	6	2	5	2	-	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Suppurative and necrotic conditions of lower respiratory tract (J85-J86)												
Asian or Pacific Islander 2/, both sexes												
2004.....	1	1	-	1	3	1	6	2	-	-	-	-
2003.....	-	2	4	1	1	2	3	-	-	-	-	-
2002.....	1	-	2	1	1	-	3	1	-	-	-	-
2001.....	1	1	-	2	3	1	2	3	-	-	-	-
2000.....	-	-	-	-	2	1	2	2	-	-	-	-
1999.....	2	1	1	4	1	2	-	3	1	1	-	-
Asian or Pacific Islander 2/, male												
2004.....	-	1	-	1	1	-	4	1	-	-	-	-
2003.....	-	-	2	1	-	1	3	-	-	-	-	-
2002.....	-	-	2	-	1	-	2	1	-	-	-	-
2001.....	1	-	-	2	3	1	1	2	-	-	-	-
2000.....	-	-	-	-	1	1	2	1	-	-	-	-
1999.....	1	1	-	3	-	1	-	2	1	1	-	-
Asian or Pacific Islander 2/, female												
2004.....	1	-	-	-	2	1	2	1	-	-	-	-
2003.....	-	2	2	-	1	1	-	-	-	-	-	-
2002.....	1	-	-	1	-	-	1	-	-	-	-	-
2001.....	-	1	-	-	-	-	1	1	-	-	-	-
2000.....	-	-	-	-	1	-	-	1	-	-	-	-
1999.....	1	-	1	1	1	1	-	1	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pleural effusion and plaque (J90,J92)												
All races, both sexes												
2004.....	11	27	31	48	77	125	198	205	144	75	4	-
2003.....	12	21	28	45	76	122	190	222	139	45	10	-
2002.....	14	14	29	46	75	116	198	223	154	60	17	-
2001.....	16	11	33	39	81	122	198	236	141	54	11	-
2000.....	12	17	22	43	68	116	176	173	132	60	12	-
1999.....	10	21	26	44	75	124	166	183	132	49	9	-
All races, male												
2004.....	7	21	11	31	35	67	92	92	55	19	1	-
2003.....	4	13	14	24	33	56	88	95	52	15	1	-
2002.....	6	9	19	17	36	54	95	83	57	10	3	-
2001.....	7	4	17	24	46	49	92	96	42	17	3	-
2000.....	7	10	10	23	40	60	77	69	42	18	4	-
1999.....	7	9	14	27	42	67	76	73	47	16	1	-
All races, female												
2004.....	4	6	20	17	42	58	106	113	89	56	3	-
2003.....	8	8	14	21	43	66	102	127	87	30	9	-
2002.....	8	5	10	29	39	62	103	140	97	50	14	-
2001.....	9	7	16	15	35	73	106	140	99	37	8	-
2000.....	5	7	12	20	28	56	99	104	90	42	8	-
1999.....	3	12	12	17	33	57	90	110	85	33	8	-
White, both sexes												
2004.....	8	23	29	39	67	113	180	185	138	67	3	-
2003.....	7	18	21	38	70	97	177	208	135	39	8	-
2002.....	13	11	24	40	60	104	180	207	148	56	13	-
2001.....	9	10	27	33	68	106	182	219	125	43	10	-
2000.....	10	15	18	33	60	107	158	159	125	54	9	-
1999.....	7	15	23	40	66	112	153	163	125	44	6	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pleural effusion and plaque (J90,J92)												
White, male												
2004.....	5	17	10	27	32	62	85	86	53	17	1	-
2003.....	2	10	8	21	30	41	83	91	50	14	1	-
2002.....	5	7	16	14	30	49	87	79	54	10	3	-
2001.....	3	3	15	22	40	41	85	90	39	15	3	-
2000.....	7	8	9	17	32	55	71	63	38	18	2	-
1999.....	5	5	12	25	39	60	70	66	46	13	-	-
White, female												
2004.....	3	6	19	12	35	51	95	99	85	50	2	-
2003.....	5	8	13	17	40	56	94	117	85	25	7	-
2002.....	8	4	8	26	30	55	93	128	94	46	10	-
2001.....	6	7	12	11	28	65	97	129	86	28	7	-
2000.....	3	7	9	16	28	52	87	96	87	36	7	-
1999.....	2	10	11	15	27	52	83	97	79	31	6	-
Black, both sexes												
2004.....	3	4	2	8	7	9	14	16	5	6	1	-
2003.....	5	3	6	5	5	19	11	11	4	6	2	-
2002.....	1	2	5	5	14	9	17	16	6	3	3	-
2001.....	7	-	5	6	11	14	15	15	12	11	1	-
2000.....	2	2	2	9	7	9	15	11	6	6	2	-
1999.....	3	6	3	4	8	10	11	15	6	3	3	-
Black, male												
2004.....	2	4	1	4	2	4	7	5	1	2	-	-
2003.....	2	3	5	2	2	13	3	3	2	1	-	-
2002.....	1	2	3	2	6	3	8	4	3	-	-	-
2001.....	4	-	2	2	5	8	7	5	3	2	-	-
2000.....	-	2	-	6	7	5	4	5	3	-	1	-
1999.....	2	4	2	2	2	6	4	6	-	2	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pleural effusion and plaque (J90,J92)												
Asian or Pacific Islander 2/, both sexes												
2004.....	-	-	-	1	3	1	3	3	1	2	-	-
2003.....	-	-	-	1	1	2	2	2	-	-	-	-
2002.....	-	1	-	1	-	3	-	-	-	1	1	-
2001.....	-	1	-	-	1	2	1	2	4	-	-	-
2000.....	-	-	2	1	1	-	2	2	1	-	1	-
1999.....	-	-	-	-	1	2	2	5	1	2	-	-
Asian or Pacific Islander 2/, male												
2004.....	-	-	-	-	1	-	-	1	1	-	-	-
2003.....	-	-	-	1	1	1	2	1	-	-	-	-
2002.....	-	-	-	1	-	2	-	-	-	-	-	-
2001.....	-	1	-	-	-	-	-	1	-	-	-	-
2000.....	-	-	1	-	1	-	2	1	1	-	1	-
1999.....	-	-	-	-	1	1	2	1	1	1	-	-
Asian or Pacific Islander 2/, female												
2004.....	-	-	-	1	2	1	3	2	-	2	-	-
2003.....	-	-	-	-	-	1	-	1	-	-	-	-
2002.....	-	1	-	-	-	1	-	-	-	1	1	-
2001.....	-	-	-	-	1	2	1	1	4	-	-	-
2000.....	-	-	1	1	-	-	-	1	-	-	-	-
1999.....	-	-	-	-	-	1	-	4	-	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of respiratory system (J80-J84,J93-J98)												
All races, both sexes												
2004.....	674	1,092	1,446	2,019	3,171	4,268	4,709	3,489	1,898	591	108	-
2003.....	662	922	1,263	1,905	2,896	3,994	4,173	3,343	1,706	647	116	1
2002.....	666	946	1,269	1,818	3,046	3,933	4,092	3,095	1,672	547	109	-
2001.....	707	943	1,275	2,022	3,126	4,277	4,226	3,228	1,650	563	93	3
2000.....	704	961	1,265	2,029	3,031	4,184	4,070	3,053	1,622	550	108	3
1999.....	593	804	1,231	1,958	3,040	4,046	3,935	3,163	1,602	576	124	-
All races, male												
2004.....	374	569	794	1,143	1,769	2,252	2,312	1,443	639	153	22	-
2003.....	342	505	715	1,080	1,563	2,086	1,991	1,368	547	158	29	-
2002.....	328	509	692	1,008	1,726	2,023	1,914	1,252	542	122	14	-
2001.....	374	487	714	1,115	1,713	2,179	1,933	1,243	538	118	16	1
2000.....	371	520	696	1,084	1,652	2,190	1,900	1,257	531	133	21	2
1999.....	300	411	670	1,071	1,676	2,086	1,905	1,278	476	150	27	-
All races, female												
2004.....	300	523	652	876	1,402	2,016	2,397	2,046	1,259	438	86	-
2003.....	320	417	548	825	1,333	1,908	2,182	1,975	1,159	489	87	1
2002.....	338	437	577	810	1,320	1,910	2,178	1,843	1,130	425	95	-
2001.....	333	456	561	907	1,413	2,098	2,293	1,985	1,112	445	77	2
2000.....	333	441	569	945	1,379	1,994	2,170	1,796	1,091	417	87	1
1999.....	293	393	561	887	1,364	1,960	2,030	1,885	1,126	426	97	-
White, both sexes												
2004.....	517	898	1,220	1,760	2,862	3,928	4,368	3,275	1,769	550	94	-
2003.....	514	752	1,086	1,676	2,618	3,650	3,925	3,152	1,593	601	101	1
2002.....	516	773	1,062	1,597	2,732	3,636	3,784	2,898	1,562	508	103	-
2001.....	529	767	1,072	1,764	2,825	3,936	3,941	3,021	1,533	505	84	1
2000.....	548	799	1,067	1,768	2,750	3,848	3,775	2,889	1,497	503	98	1
1999.....	464	658	1,058	1,723	2,714	3,710	3,655	2,956	1,497	516	110	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of respiratory system (J80-J84,J93-J98)												
White, male												
2004.....	297	473	682	1,006	1,604	2,084	2,152	1,346	595	145	18	-
2003.....	276	413	619	957	1,429	1,908	1,882	1,287	506	144	29	-
2002.....	257	426	581	910	1,554	1,878	1,783	1,183	500	111	14	-
2001.....	293	413	597	991	1,575	2,028	1,815	1,167	500	101	14	-
2000.....	287	437	595	963	1,502	2,020	1,774	1,194	484	115	18	1
1999.....	237	335	586	964	1,518	1,932	1,777	1,199	440	134	22	-
White, female												
2004.....	220	425	538	754	1,258	1,844	2,216	1,929	1,174	405	76	-
2003.....	238	339	467	719	1,189	1,742	2,043	1,865	1,087	457	72	1
2002.....	259	347	481	687	1,178	1,758	2,001	1,715	1,062	397	89	-
2001.....	236	354	475	773	1,250	1,908	2,126	1,854	1,033	404	70	1
2000.....	261	362	472	805	1,248	1,828	2,001	1,695	1,013	388	80	-
1999.....	227	323	472	759	1,196	1,778	1,878	1,757	1,057	382	88	-
Black, both sexes												
2004.....	137	167	188	206	230	245	232	144	97	27	12	-
2003.....	134	145	145	176	200	264	172	135	74	41	15	-
2002.....	128	145	174	177	226	205	229	134	88	31	6	-
2001.....	158	149	170	206	238	257	218	146	93	45	9	2
2000.....	134	142	156	222	219	255	227	120	98	36	9	2
1999.....	113	120	143	199	257	270	204	161	76	47	11	-
Black, male												
2004.....	63	84	91	111	119	122	99	62	32	5	2	-
2003.....	58	78	83	93	103	135	76	60	22	12	-	-
2002.....	61	70	94	81	121	98	96	42	30	9	-	-
2001.....	71	63	93	99	105	110	89	56	25	11	2	1
2000.....	69	71	76	104	106	130	88	40	34	15	2	1
1999.....	55	57	70	91	121	125	90	58	26	10	4	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of respiratory system (J80-J84,J93-J98)												
Black, female												
2004.....	74	83	97	95	111	123	133	82	65	22	10	-
2003.....	76	67	62	83	97	129	96	75	52	29	15	-
2002.....	67	75	80	96	105	107	133	92	58	22	6	-
2001.....	87	86	77	107	133	147	129	90	68	34	7	1
2000.....	65	71	80	118	113	125	139	80	64	21	7	1
1999.....	58	63	73	108	136	145	114	103	50	37	7	-
American Indian 1/, both sexes												
2004.....	11	9	12	16	27	21	18	13	7	6	1	-
2003.....	8	8	16	14	25	19	18	13	12	-	-	-
2002.....	8	9	11	11	18	14	11	19	8	-	-	-
2001.....	9	10	16	15	19	21	20	16	4	6	-	-
2000.....	10	6	15	14	23	16	17	9	8	1	-	-
1999.....	9	6	8	12	13	24	21	15	10	2	2	-
American Indian 1/, male												
2004.....	9	5	6	8	16	9	9	7	2	1	1	-
2003.....	4	4	8	7	10	12	7	3	6	-	-	-
2002.....	3	6	6	5	8	9	7	10	3	-	-	-
2001.....	3	2	10	6	11	9	11	5	1	2	-	-
2000.....	9	3	9	5	9	9	5	4	3	1	-	-
1999.....	5	6	3	5	8	12	7	7	3	1	1	-
American Indian 1/, female												
2004.....	2	4	6	8	11	12	9	6	5	5	-	-
2003.....	4	4	8	7	15	7	11	10	6	-	-	-
2002.....	5	3	5	6	10	5	4	9	5	-	-	-
2001.....	6	8	6	9	8	12	9	11	3	4	-	-
2000.....	1	3	6	9	14	7	12	5	5	-	-	-
1999.....	4	-	5	7	5	12	14	8	7	1	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of respiratory system (J80-J84,J93-J98)												
Asian or Pacific Islander 2/, both sexes												
2004.....	9	18	26	37	52	74	91	57	25	8	1	-
2003.....	6	17	16	39	53	61	58	43	27	5	-	-
2002.....	14	19	22	33	70	78	68	44	14	8	-	-
2001.....	11	17	17	37	44	63	47	45	20	7	-	-
2000.....	12	14	27	25	39	65	51	35	19	10	1	-
1999.....	7	20	22	24	56	42	55	31	19	11	1	-
Asian or Pacific Islander 2/, male												
2004.....	5	7	15	18	30	37	52	28	10	2	1	-
2003.....	4	10	5	23	21	31	26	18	13	2	-	-
2002.....	7	7	11	12	43	38	28	17	9	2	-	-
2001.....	7	9	14	19	22	32	18	15	12	4	-	-
2000.....	6	9	16	12	35	31	33	19	10	2	1	-
1999.....	3	13	11	11	29	17	31	14	7	5	-	-
Asian or Pacific Islander 2/, female												
2004.....	4	11	11	19	22	37	39	29	15	6	-	-
2003.....	2	7	11	16	32	30	32	25	14	3	-	-
2002.....	7	12	11	21	27	40	40	27	5	6	-	-
2001.....	4	8	3	18	22	31	29	30	8	3	-	-
2000.....	6	5	11	13	4	34	18	16	9	8	-	-
1999.....	4	7	11	13	27	25	24	17	12	6	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XI. Diseases of the digestive system (K00-K92)												
All races, both sexes												
2004.....	6,843	6,605	6,331	6,539	7,937	9,936	11,136	9,756	6,363	2,310	404	2
2003.....	6,760	6,393	6,278	6,868	8,259	10,225	11,282	9,926	6,369	2,299	441	5
2002.....	6,349	6,025	5,859	6,574	8,349	10,329	11,314	10,063	6,363	2,236	411	7
2001.....	6,079	5,537	5,912	6,809	8,543	10,261	10,958	10,009	6,256	2,300	432	12
2000.....	5,576	5,359	5,720	6,572	8,530	10,432	10,629	9,986	6,091	2,179	427	4
1999.....	5,249	5,177	5,719	6,895	8,789	10,166	10,492	9,546	6,020	2,130	433	7
All races, male												
2004.....	4,800	4,378	3,828	3,668	4,050	4,689	4,502	3,298	1,721	446	68	-
2003.....	4,741	4,201	3,925	3,871	4,278	4,705	4,657	3,385	1,703	452	80	5
2002.....	4,474	3,933	3,605	3,723	4,317	4,779	4,576	3,445	1,647	459	69	5
2001.....	4,216	3,643	3,621	3,832	4,427	4,679	4,386	3,322	1,659	503	79	12
2000.....	3,845	3,470	3,528	3,725	4,385	4,888	4,207	3,334	1,579	448	61	4
1999.....	3,675	3,428	3,487	3,916	4,569	4,649	4,293	3,168	1,590	402	73	4
All races, female												
2004.....	2,043	2,227	2,503	2,871	3,887	5,247	6,634	6,458	4,642	1,864	336	2
2003.....	2,019	2,192	2,353	2,997	3,981	5,520	6,625	6,541	4,666	1,847	361	-
2002.....	1,875	2,092	2,254	2,851	4,032	5,550	6,738	6,618	4,716	1,777	342	2
2001.....	1,863	1,894	2,291	2,977	4,116	5,582	6,572	6,687	4,597	1,797	353	-
2000.....	1,731	1,889	2,192	2,847	4,145	5,544	6,422	6,652	4,512	1,731	366	-
1999.....	1,574	1,749	2,232	2,979	4,220	5,517	6,199	6,378	4,430	1,728	360	3
White, both sexes												
2004.....	5,615	5,449	5,329	5,630	6,945	8,938	10,122	8,996	5,883	2,115	368	2
2003.....	5,437	5,299	5,257	5,894	7,286	9,209	10,325	9,181	5,892	2,105	387	4
2002.....	5,159	5,039	4,927	5,685	7,364	9,325	10,323	9,339	5,929	2,068	364	5
2001.....	4,865	4,493	4,976	5,848	7,564	9,178	10,052	9,254	5,847	2,108	390	8
2000.....	4,447	4,406	4,767	5,623	7,555	9,399	9,723	9,188	5,665	2,007	382	3
1999.....	4,271	4,245	4,777	5,905	7,770	9,131	9,632	8,786	5,607	1,983	390	4

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XI. Diseases of the digestive system (K00-K92)												
White, male												
2004.....	4,007	3,624	3,235	3,152	3,552	4,220	4,080	3,044	1,592	412	58	-
2003.....	3,857	3,481	3,315	3,336	3,770	4,247	4,236	3,131	1,571	409	63	4
2002.....	3,689	3,313	3,044	3,237	3,820	4,331	4,166	3,187	1,525	419	56	5
2001.....	3,416	2,968	3,062	3,291	3,935	4,179	4,042	3,050	1,540	453	69	8
2000.....	3,093	2,893	2,966	3,221	3,894	4,401	3,855	3,054	1,465	397	52	3
1999.....	3,018	2,832	2,956	3,393	4,081	4,198	3,958	2,880	1,472	363	68	3
White, female												
2004.....	1,608	1,825	2,094	2,478	3,393	4,718	6,042	5,952	4,291	1,703	310	2
2003.....	1,580	1,818	1,942	2,558	3,516	4,962	6,089	6,050	4,321	1,696	324	-
2002.....	1,470	1,726	1,883	2,448	3,544	4,994	6,157	6,152	4,404	1,649	308	-
2001.....	1,449	1,525	1,914	2,557	3,629	4,999	6,010	6,204	4,307	1,655	321	-
2000.....	1,354	1,513	1,801	2,402	3,661	4,998	5,868	6,134	4,200	1,610	330	-
1999.....	1,253	1,413	1,821	2,512	3,689	4,933	5,674	5,906	4,135	1,620	322	1
Black, both sexes												
2004.....	1,028	951	816	729	804	771	806	594	389	154	32	-
2003.....	1,115	917	828	790	776	808	738	594	383	164	45	1
2002.....	1,014	807	750	723	787	798	795	588	351	141	42	2
2001.....	1,027	883	766	812	803	874	742	603	340	163	37	4
2000.....	953	792	791	781	817	830	733	664	377	141	39	1
1999.....	824	775	797	817	833	840	718	627	355	127	34	3
Black, male												
2004.....	664	632	491	422	415	358	320	192	102	22	9	-
2003.....	758	609	500	436	414	356	322	197	94	30	10	1
2002.....	668	516	451	399	399	344	316	204	92	25	11	-
2001.....	696	576	452	467	410	396	271	210	84	39	8	4
2000.....	635	492	470	419	415	384	286	211	87	35	7	1
1999.....	556	499	448	433	398	360	275	222	89	29	2	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XI. Diseases of the digestive system (K00-K92)												
Black, female												
2004.....	364	319	325	307	389	413	486	402	287	132	23	-
2003.....	357	308	328	354	362	452	416	397	289	134	35	-
2002.....	346	291	299	324	388	454	479	384	259	116	31	2
2001.....	331	307	314	345	393	478	471	393	256	124	29	-
2000.....	318	300	321	362	402	446	447	453	290	106	32	-
1999.....	268	276	349	384	435	480	443	405	266	98	32	2
American Indian 1/, both sexes												
2004.....	132	127	96	83	61	62	43	35	15	7	1	-
2003.....	137	107	105	73	50	57	46	34	28	4	1	-
2002.....	113	98	91	73	67	39	40	31	14	5	-	-
2001.....	121	87	92	59	59	57	30	39	11	6	1	-
2000.....	121	92	74	79	47	53	38	27	9	5	2	-
1999.....	101	97	79	71	56	50	23	20	10	7	3	-
American Indian 1/, male												
2004.....	71	71	49	39	23	34	14	9	2	1	-	-
2003.....	81	60	53	40	23	19	19	10	9	-	1	-
2002.....	74	54	52	41	31	19	15	10	4	2	-	-
2001.....	68	51	60	31	28	25	10	11	6	2	1	-
2000.....	79	49	40	37	14	23	15	7	4	1	1	-
1999.....	64	54	41	30	25	20	8	5	3	3	-	-
American Indian 1/, female												
2004.....	61	56	47	44	38	28	29	26	13	6	1	-
2003.....	56	47	52	33	27	38	27	24	19	4	-	-
2002.....	39	44	39	32	36	20	25	21	10	3	-	-
2001.....	53	36	32	28	31	32	20	28	5	4	-	-
2000.....	42	43	34	42	33	30	23	20	5	4	1	-
1999.....	37	43	38	41	31	30	15	15	7	4	3	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XI. Diseases of the digestive system (K00-K92)												
Asian or Pacific Islander 2/, both sexes												
2004.....	68	78	90	97	127	165	165	131	76	34	3	-
2003.....	71	70	88	111	147	151	173	117	66	26	8	-
2002.....	63	81	91	93	131	167	156	105	69	22	5	-
2001.....	66	74	78	90	117	152	134	113	58	23	4	-
2000.....	55	69	88	89	111	150	135	107	40	26	4	-
1999.....	53	60	66	102	130	145	119	113	48	13	6	-
Asian or Pacific Islander 2/, male												
2004.....	58	51	53	55	60	77	88	53	25	11	1	-
2003.....	45	51	57	59	71	83	80	47	29	13	6	-
2002.....	43	50	58	46	67	85	79	44	26	13	2	-
2001.....	36	48	47	43	54	79	63	51	29	9	1	-
2000.....	38	36	52	48	62	80	51	62	23	15	1	-
1999.....	37	43	42	60	65	71	52	61	26	7	3	-
Asian or Pacific Islander 2/, female												
2004.....	10	27	37	42	67	88	77	78	51	23	2	-
2003.....	26	19	31	52	76	68	93	70	37	13	2	-
2002.....	20	31	33	47	64	82	77	61	43	9	3	-
2001.....	30	26	31	47	63	73	71	62	29	14	3	-
2000.....	17	33	36	41	49	70	84	45	17	11	3	-
1999.....	16	17	24	42	65	74	67	52	22	6	3	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of oral cavity, salivary glands and jaws (K00-K14)												
All races, both sexes												
2004.....	8	9	14	13	13	20	36	49	29	11	2	-
2003.....	4	4	9	6	8	12	19	27	22	6	2	-
2002.....	3	6	6	7	13	14	22	31	22	11	4	-
2001.....	2	2	3	2	4	21	31	22	21	6	3	-
2000.....	4	-	2	4	8	9	20	28	14	6	1	-
1999.....	3	1	2	8	12	14	13	18	15	7	3	-
All races, male												
2004.....	5	4	6	5	3	14	14	16	5	1	-	-
2003.....	3	3	4	2	3	6	6	8	4	1	1	-
2002.....	3	4	5	5	8	8	7	9	6	1	1	-
2001.....	2	2	3	-	2	8	13	9	6	1	-	-
2000.....	1	-	2	3	5	4	5	10	3	1	-	-
1999.....	3	1	1	3	8	4	3	7	3	3	1	-
All races, female												
2004.....	3	5	8	8	10	6	22	33	24	10	2	-
2003.....	1	1	5	4	5	6	13	19	18	5	1	-
2002.....	-	2	1	2	5	6	15	22	16	10	3	-
2001.....	-	-	-	2	2	13	18	13	15	5	3	-
2000.....	3	-	-	1	3	5	15	18	11	5	1	-
1999.....	-	-	1	5	4	10	10	11	12	4	2	-
White, both sexes												
2004.....	5	9	13	10	11	19	34	47	28	11	2	-
2003.....	4	2	6	6	6	12	19	27	22	6	2	-
2002.....	1	5	6	6	13	13	21	29	20	11	4	-
2001.....	2	2	3	2	4	16	27	20	20	6	3	-
2000.....	4	-	2	3	7	9	17	27	14	4	1	-
1999.....	3	1	1	8	9	12	11	17	14	6	3	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of oral cavity, salivary glands and jaws (K00-K14)												
White, male												
2004.....	3	4	6	4	2	13	13	16	5	1	-	-
2003.....	3	2	3	2	2	6	6	8	4	1	1	-
2002.....	1	4	5	4	8	8	6	8	6	1	1	-
2001.....	2	2	3	-	2	6	10	8	6	1	-	-
2000.....	1	-	2	3	4	4	5	10	3	1	-	-
1999.....	3	1	-	3	7	4	2	7	3	2	1	-
White, female												
2004.....	2	5	7	6	9	6	21	31	23	10	2	-
2003.....	1	-	3	4	4	6	13	19	18	5	1	-
2002.....	-	1	1	2	5	5	15	21	14	10	3	-
2001.....	-	-	-	2	2	10	17	12	14	5	3	-
2000.....	3	-	-	-	3	5	12	17	11	3	1	-
1999.....	-	-	1	5	2	8	9	10	11	4	2	-
Black, both sexes												
2004.....	3	-	1	3	2	1	1	-	1	-	-	-
2003.....	-	1	2	-	2	-	-	-	-	-	-	-
2002.....	1	1	-	1	-	1	1	1	2	-	-	-
2001.....	-	-	-	-	-	5	3	2	1	-	-	-
2000.....	-	-	-	1	1	-	3	-	-	1	-	-
1999.....	-	-	1	-	1	2	2	1	1	-	-	-
Black, male												
2004.....	2	-	-	1	1	1	-	-	-	-	-	-
2003.....	-	1	-	-	1	-	-	-	-	-	-	-
2002.....	1	-	-	1	-	-	1	-	-	-	-	-
2001.....	-	-	-	-	-	2	2	1	-	-	-	-
2000.....	-	-	-	-	1	-	-	-	-	-	-	-
1999.....	-	-	1	-	-	-	1	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of oral cavity, salivary glands and jaws (K00-K14)												
Asian or Pacific Islander 2/, both sexes												
2004.....	-	-	-	-	-	-	-	2	-	-	-	-
2003.....	-	1	1	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	1	-	-	-	-
2001.....	-	-	-	-	-	-	1	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	1	-	1	-	-
1999.....	-	-	-	-	2	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, male												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	1	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	1	-	-	-	-
2001.....	-	-	-	-	-	-	1	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	1	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, female												
2004.....	-	-	-	-	-	-	-	2	-	-	-	-
2003.....	-	1	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	1	-	1	-	-
1999.....	-	-	-	-	1	-	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of other parts of digestive system (K20-K92)												
All races, both sexes												
2004.....	6,835	6,596	6,317	6,526	7,924	9,916	11,100	9,707	6,334	2,299	402	2
2003.....	6,756	6,389	6,269	6,862	8,251	10,213	11,263	9,899	6,347	2,293	439	5
2002.....	6,346	6,019	5,853	6,567	8,336	10,315	11,292	10,032	6,341	2,225	407	7
2001.....	6,077	5,535	5,909	6,807	8,539	10,240	10,927	9,987	6,235	2,294	429	12
2000.....	5,572	5,359	5,718	6,568	8,522	10,423	10,609	9,958	6,077	2,173	426	4
1999.....	5,246	5,176	5,717	6,887	8,777	10,152	10,479	9,528	6,005	2,123	430	7
All races, male												
2004.....	4,795	4,374	3,822	3,663	4,047	4,675	4,488	3,282	1,716	445	68	-
2003.....	4,738	4,198	3,921	3,869	4,275	4,699	4,651	3,377	1,699	451	79	5
2002.....	4,471	3,929	3,600	3,718	4,309	4,771	4,569	3,436	1,641	458	68	5
2001.....	4,214	3,641	3,618	3,832	4,425	4,671	4,373	3,313	1,653	502	79	12
2000.....	3,844	3,470	3,526	3,722	4,380	4,884	4,202	3,324	1,576	447	61	4
1999.....	3,672	3,427	3,486	3,913	4,561	4,645	4,290	3,161	1,587	399	72	4
All races, female												
2004.....	2,040	2,222	2,495	2,863	3,877	5,241	6,612	6,425	4,618	1,854	334	2
2003.....	2,018	2,191	2,348	2,993	3,976	5,514	6,612	6,522	4,648	1,842	360	-
2002.....	1,875	2,090	2,253	2,849	4,027	5,544	6,723	6,596	4,700	1,767	339	2
2001.....	1,863	1,894	2,291	2,975	4,114	5,569	6,554	6,674	4,582	1,792	350	-
2000.....	1,728	1,889	2,192	2,846	4,142	5,539	6,407	6,634	4,501	1,726	365	-
1999.....	1,574	1,749	2,231	2,974	4,216	5,507	6,189	6,367	4,418	1,724	358	3
White, both sexes												
2004.....	5,610	5,440	5,316	5,620	6,934	8,919	10,088	8,949	5,855	2,104	366	2
2003.....	5,433	5,297	5,251	5,888	7,280	9,197	10,306	9,154	5,870	2,099	385	4
2002.....	5,158	5,034	4,921	5,679	7,351	9,312	10,302	9,310	5,909	2,057	360	5
2001.....	4,863	4,491	4,973	5,846	7,560	9,162	10,025	9,234	5,827	2,102	387	8
2000.....	4,443	4,406	4,765	5,620	7,548	9,390	9,706	9,161	5,651	2,003	381	3
1999.....	4,268	4,244	4,776	5,897	7,761	9,119	9,621	8,769	5,593	1,977	387	4

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of other parts of digestive system (K20-K92)												
White, male												
2004.....	4,004	3,620	3,229	3,148	3,550	4,207	4,067	3,028	1,587	411	58	-
2003.....	3,854	3,479	3,312	3,334	3,768	4,241	4,230	3,123	1,567	408	62	4
2002.....	3,688	3,309	3,039	3,233	3,812	4,323	4,160	3,179	1,519	418	55	5
2001.....	3,414	2,966	3,059	3,291	3,933	4,173	4,032	3,042	1,534	452	69	8
2000.....	3,092	2,893	2,964	3,218	3,890	4,397	3,850	3,044	1,462	396	52	3
1999.....	3,015	2,831	2,956	3,390	4,074	4,194	3,956	2,873	1,469	361	67	3
White, female												
2004.....	1,606	1,820	2,087	2,472	3,384	4,712	6,021	5,921	4,268	1,693	308	2
2003.....	1,579	1,818	1,939	2,554	3,512	4,956	6,076	6,031	4,303	1,691	323	-
2002.....	1,470	1,725	1,882	2,446	3,539	4,989	6,142	6,131	4,390	1,639	305	-
2001.....	1,449	1,525	1,914	2,555	3,627	4,989	5,993	6,192	4,293	1,650	318	-
2000.....	1,351	1,513	1,801	2,402	3,658	4,993	5,856	6,117	4,189	1,607	329	-
1999.....	1,253	1,413	1,820	2,507	3,687	4,925	5,665	5,896	4,124	1,616	320	1
Black, both sexes												
2004.....	1,025	951	815	726	802	770	805	594	388	154	32	-
2003.....	1,115	916	826	790	774	808	738	594	383	164	45	1
2002.....	1,013	806	750	722	787	797	794	587	349	141	42	2
2001.....	1,027	883	766	812	803	869	739	601	339	163	37	4
2000.....	953	792	791	780	816	830	730	664	377	140	39	1
1999.....	824	775	796	817	832	838	716	626	354	127	34	3
Black, male												
2004.....	662	632	491	421	414	357	320	192	102	22	9	-
2003.....	758	608	500	436	413	356	322	197	94	30	10	1
2002.....	667	516	451	398	399	344	315	204	92	25	11	-
2001.....	696	576	452	467	410	394	269	209	84	39	8	4
2000.....	635	492	470	419	414	384	286	211	87	35	7	1
1999.....	556	499	447	433	398	360	274	222	89	29	2	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of other parts of digestive system (K20-K92)												
Black, female												
2004.....	363	319	324	305	388	413	485	402	286	132	23	-
2003.....	357	308	326	354	361	452	416	397	289	134	35	-
2002.....	346	290	299	324	388	453	479	383	257	116	31	2
2001.....	331	307	314	345	393	475	470	392	255	124	29	-
2000.....	318	300	321	361	402	446	444	453	290	105	32	-
1999.....	268	276	349	384	434	478	442	404	265	98	32	2
American Indian 1/, both sexes												
2004.....	132	127	96	83	61	62	42	35	15	7	1	-
2003.....	137	107	105	73	50	57	46	34	28	4	1	-
2002.....	112	98	91	73	67	39	40	31	14	5	-	-
2001.....	121	87	92	59	59	57	30	39	11	6	1	-
2000.....	121	92	74	79	47	53	38	27	9	5	2	-
1999.....	101	97	79	71	56	50	23	20	10	7	3	-
American Indian 1/, male												
2004.....	71	71	49	39	23	34	13	9	2	1	-	-
2003.....	81	60	53	40	23	19	19	10	9	-	1	-
2002.....	73	54	52	41	31	19	15	10	4	2	-	-
2001.....	68	51	60	31	28	25	10	11	6	2	1	-
2000.....	79	49	40	37	14	23	15	7	4	1	1	-
1999.....	64	54	41	30	25	20	8	5	3	3	-	-
American Indian 1/, female												
2004.....	61	56	47	44	38	28	29	26	13	6	1	-
2003.....	56	47	52	33	27	38	27	24	19	4	-	-
2002.....	39	44	39	32	36	20	25	21	10	3	-	-
2001.....	53	36	32	28	31	32	20	28	5	4	-	-
2000.....	42	43	34	42	33	30	23	20	5	4	1	-
1999.....	37	43	38	41	31	30	15	15	7	4	3	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of other parts of digestive system (K20-K92)												
Asian or Pacific Islander 2/, both sexes												
2004.....	68	78	90	97	127	165	165	129	76	34	3	-
2003.....	71	69	87	111	147	151	173	117	66	26	8	-
2002.....	63	81	91	93	131	167	156	104	69	22	5	-
2001.....	66	74	78	90	117	152	133	113	58	23	4	-
2000.....	55	69	88	89	111	150	135	106	40	25	4	-
1999.....	53	60	66	102	128	145	119	113	48	12	6	-
Asian or Pacific Islander 2/, male												
2004.....	58	51	53	55	60	77	88	53	25	11	1	-
2003.....	45	51	56	59	71	83	80	47	29	13	6	-
2002.....	43	50	58	46	67	85	79	43	26	13	2	-
2001.....	36	48	47	43	54	79	62	51	29	9	1	-
2000.....	38	36	52	48	62	80	51	62	23	15	1	-
1999.....	37	43	42	60	64	71	52	61	26	6	3	-
Asian or Pacific Islander 2/, female												
2004.....	10	27	37	42	67	88	77	76	51	23	2	-
2003.....	26	18	31	52	76	68	93	70	37	13	2	-
2002.....	20	31	33	47	64	82	77	61	43	9	3	-
2001.....	30	26	31	47	63	73	71	62	29	14	3	-
2000.....	17	33	36	41	49	70	84	44	17	10	3	-
1999.....	16	17	24	42	64	74	67	52	22	6	3	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of esophagus (K20-K22)												
All races, both sexes												
2004.....	47	52	75	95	125	209	325	407	304	132	26	-
2003.....	53	57	68	91	137	232	333	428	318	134	30	-
2002.....	50	43	55	91	148	261	361	449	340	140	22	-
2001.....	41	52	70	97	156	257	383	434	342	122	36	-
2000.....	49	52	68	105	160	264	381	449	374	140	25	-
1999.....	44	53	65	108	171	241	374	431	379	129	37	-
All races, male												
2004.....	36	33	47	53	61	118	143	154	86	37	11	-
2003.....	33	37	46	57	80	112	143	177	108	46	10	-
2002.....	36	22	31	61	73	129	167	180	126	46	1	-
2001.....	31	36	47	56	83	124	170	158	105	32	11	-
2000.....	37	35	51	68	83	140	170	185	123	37	6	-
1999.....	24	28	42	65	104	134	178	175	119	34	10	-
All races, female												
2004.....	11	19	28	42	64	91	182	253	218	95	15	-
2003.....	20	20	22	34	57	120	190	251	210	88	20	-
2002.....	14	21	24	30	75	132	194	269	214	94	21	-
2001.....	10	16	23	41	73	133	213	276	237	90	25	-
2000.....	12	17	17	37	77	124	211	264	251	103	19	-
1999.....	20	25	23	43	67	107	196	256	260	95	27	-
White, both sexes												
2004.....	41	44	57	81	108	187	304	391	285	122	24	-
2003.....	41	48	58	80	126	211	313	404	299	126	29	-
2002.....	44	34	49	71	132	228	336	428	325	127	21	-
2001.....	33	45	59	86	135	236	355	404	330	118	34	-
2000.....	39	43	56	92	138	244	362	424	354	135	23	-
1999.....	32	47	53	94	153	224	350	400	364	123	35	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of esophagus (K20-K22)												
White, male												
2004.....	33	27	36	42	53	105	134	149	80	34	10	-
2003.....	24	30	37	49	77	102	136	169	102	44	10	-
2002.....	30	16	27	46	67	113	154	174	117	44	1	-
2001.....	25	30	38	48	70	110	157	147	102	31	11	-
2000.....	28	29	40	59	70	127	161	176	118	34	5	-
1999.....	17	26	31	58	91	123	164	165	113	34	10	-
White, female												
2004.....	8	17	21	39	55	82	170	242	205	88	14	-
2003.....	17	18	21	31	49	109	177	235	197	82	19	-
2002.....	14	18	22	25	65	115	182	254	208	83	20	-
2001.....	8	15	21	38	65	126	198	257	228	87	23	-
2000.....	11	14	16	33	68	117	201	248	236	101	18	-
1999.....	15	21	22	36	62	101	186	235	251	89	25	-
Black, both sexes												
2004.....	6	8	15	13	17	19	16	13	13	8	2	-
2003.....	10	7	9	10	8	19	17	17	16	6	1	-
2002.....	5	9	4	19	14	25	19	19	11	10	-	-
2001.....	7	7	8	10	16	16	23	26	10	4	2	-
2000.....	10	8	11	12	18	19	18	23	16	5	2	-
1999.....	10	5	11	13	15	15	18	27	14	6	1	-
Black, male												
2004.....	3	6	9	10	8	11	8	5	4	2	1	-
2003.....	7	5	8	7	3	9	4	6	5	1	-	-
2002.....	5	6	3	14	6	11	9	5	7	1	-	-
2001.....	5	6	7	7	9	11	11	9	3	1	-	-
2000.....	9	5	10	9	10	13	8	7	3	3	1	-
1999.....	5	1	10	7	12	10	10	8	5	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of esophagus (K20-K22)												
Black, female												
2004.....	3	2	6	3	9	8	8	8	9	6	1	-
2003.....	3	2	1	3	5	10	13	11	11	5	1	-
2002.....	-	3	1	5	8	14	10	14	4	9	-	-
2001.....	2	1	1	3	7	5	12	17	7	3	2	-
2000.....	1	3	1	3	8	6	10	16	13	2	1	-
1999.....	5	4	1	6	3	5	8	19	9	6	1	-
American Indian 1/, both sexes												
2004.....	-	-	-	-	-	-	2	2	1	1	-	-
2003.....	1	1	-	-	1	1	-	2	-	-	-	-
2002.....	1	-	1	-	1	1	-	-	-	1	-	-
2001.....	1	-	2	-	2	2	2	2	1	-	-	-
2000.....	-	-	1	1	2	-	1	1	2	-	-	-
1999.....	1	1	-	1	1	-	2	2	1	-	-	-
American Indian 1/, male												
2004.....	-	-	-	-	-	-	-	-	-	1	-	-
2003.....	1	1	-	-	-	-	-	1	-	-	-	-
2002.....	1	-	1	-	-	1	-	-	-	-	-	-
2001.....	1	-	2	-	2	1	1	-	-	-	-	-
2000.....	-	-	1	-	2	-	1	1	-	-	-	-
1999.....	1	1	-	-	1	-	2	-	1	-	-	-
American Indian 1/, female												
2004.....	-	-	-	-	-	-	2	2	1	-	-	-
2003.....	-	-	-	-	1	1	-	1	-	-	-	-
2002.....	-	-	-	-	1	-	-	-	-	1	-	-
2001.....	-	-	-	-	-	1	1	2	1	-	-	-
2000.....	-	-	-	1	-	-	-	-	2	-	-	-
1999.....	-	-	-	1	-	-	-	2	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of esophagus (K20-K22)												
Asian or Pacific Islander 2/, both sexes												
2004.....	-	-	3	1	-	3	3	1	5	1	-	-
2003.....	1	1	1	1	2	1	3	5	3	2	-	-
2002.....	-	-	1	1	1	7	6	2	4	2	1	-
2001.....	-	-	1	1	3	3	3	2	1	-	-	-
2000.....	-	1	-	-	2	1	-	1	2	-	-	-
1999.....	1	-	1	-	2	2	4	2	-	-	1	-
Asian or Pacific Islander 2/, male												
2004.....	-	-	2	1	-	2	1	-	2	-	-	-
2003.....	1	1	1	1	-	1	3	1	1	1	-	-
2002.....	-	-	-	1	-	4	4	1	2	1	-	-
2001.....	-	-	-	1	2	2	1	2	-	-	-	-
2000.....	-	1	-	-	1	-	-	1	2	-	-	-
1999.....	1	-	1	-	-	1	2	2	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	-	-	1	-	-	1	2	1	3	1	-	-
2003.....	-	-	-	-	2	-	-	4	2	1	-	-
2002.....	-	-	1	-	1	3	2	1	2	1	1	-
2001.....	-	-	1	-	1	1	2	-	1	-	-	-
2000.....	-	-	-	-	1	1	-	-	-	-	-	-
1999.....	-	-	-	-	2	1	2	-	-	-	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Peptic ulcer (K25-K28)												
All races, both sexes												
2004.....	150	159	215	250	332	505	643	606	413	141	25	-
2003.....	159	183	200	228	359	545	734	685	391	138	30	-
2002.....	154	209	202	254	411	592	699	714	407	148	33	-
2001.....	147	183	235	294	424	615	823	788	458	184	40	1
2000.....	177	200	243	299	449	642	773	801	498	177	33	-
1999.....	142	186	236	314	522	665	835	771	470	164	34	2
All races, male												
2004.....	97	98	130	133	171	257	273	209	120	35	7	-
2003.....	110	105	132	141	205	273	333	262	126	25	5	-
2002.....	106	129	118	142	236	287	335	279	116	39	5	-
2001.....	95	123	150	179	218	315	334	277	125	47	5	1
2000.....	131	134	150	172	270	341	326	291	150	49	9	-
1999.....	102	111	154	196	308	331	381	279	138	29	2	1
All races, female												
2004.....	53	61	85	117	161	248	370	397	293	106	18	-
2003.....	49	78	68	87	154	272	401	423	265	113	25	-
2002.....	48	80	84	112	175	305	364	435	291	109	28	-
2001.....	52	60	85	115	206	300	489	511	333	137	35	-
2000.....	46	66	93	127	179	301	447	510	348	128	24	-
1999.....	40	75	82	118	214	334	454	492	332	135	32	1
White, both sexes												
2004.....	118	129	180	212	281	447	583	562	386	129	24	-
2003.....	116	148	164	197	317	475	655	630	365	127	24	-
2002.....	118	164	161	213	363	524	652	661	380	138	28	-
2001.....	112	138	193	245	374	548	755	731	431	171	39	1
2000.....	132	162	208	250	397	569	717	731	468	164	29	-
1999.....	106	152	198	259	457	600	760	713	439	150	28	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Peptic ulcer (K25-K28)												
White, male												
2004.....	79	79	108	117	145	226	242	195	112	34	6	-
2003.....	76	83	108	120	179	238	290	240	115	21	3	-
2002.....	77	99	95	119	205	253	311	260	108	36	5	-
2001.....	68	91	122	148	184	277	303	257	115	43	5	1
2000.....	96	110	130	146	234	300	298	263	138	42	8	-
1999.....	76	88	127	160	272	302	348	255	133	24	2	1
White, female												
2004.....	39	50	72	95	136	221	341	367	274	95	18	-
2003.....	40	65	56	77	138	237	365	390	250	106	21	-
2002.....	41	65	66	94	158	271	341	401	272	102	23	-
2001.....	44	47	71	97	190	271	452	474	316	128	34	-
2000.....	36	52	78	104	163	269	419	468	330	122	21	-
1999.....	30	64	71	99	185	298	412	458	306	126	26	-
Black, both sexes												
2004.....	29	27	32	34	39	39	38	28	20	8	1	-
2003.....	39	33	28	20	34	51	57	34	16	4	4	-
2002.....	33	35	33	30	39	51	26	38	14	8	4	-
2001.....	32	35	34	33	41	45	41	38	20	9	1	-
2000.....	38	35	30	37	43	53	39	50	23	8	4	-
1999.....	32	32	34	45	48	43	56	41	24	11	4	1
Black, male												
2004.....	15	17	21	14	19	22	18	9	5	-	1	-
2003.....	31	22	20	15	22	25	32	13	5	1	-	-
2002.....	27	24	21	17	24	26	14	14	3	2	-	-
2001.....	25	26	22	21	27	23	19	15	5	1	-	-
2000.....	28	21	19	18	30	27	19	18	5	3	1	-
1999.....	23	22	25	29	26	19	21	18	3	3	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Peptic ulcer (K25-K28)												
Black, female												
2004.....	14	10	11	20	20	17	20	19	15	8	-	-
2003.....	8	11	8	5	12	26	25	21	11	3	4	-
2002.....	6	11	12	13	15	25	12	24	11	6	4	-
2001.....	7	9	12	12	14	22	22	23	15	8	1	-
2000.....	10	14	11	19	13	26	20	32	18	5	3	-
1999.....	9	10	9	16	22	24	35	23	21	8	4	1
American Indian 1/, both sexes												
2004.....	-	1	-	1	3	2	1	1	1	-	-	-
2003.....	2	1	-	2	-	5	-	1	-	1	-	-
2002.....	-	4	3	2	2	2	1	2	1	-	-	-
2001.....	1	4	2	4	3	4	2	-	1	-	-	-
2000.....	3	1	3	4	1	-	3	1	-	-	-	-
1999.....	1	-	-	1	1	3	-	-	-	-	1	-
American Indian 1/, male												
2004.....	-	-	-	-	1	1	-	-	-	-	-	-
2003.....	1	-	-	2	-	1	-	-	-	-	-	-
2002.....	-	3	-	2	1	2	1	1	-	-	-	-
2001.....	1	1	2	2	3	3	-	-	1	-	-	-
2000.....	3	1	1	3	-	-	2	-	-	-	-	-
1999.....	1	-	-	-	1	2	-	-	-	-	-	-
American Indian 1/, female												
2004.....	-	1	-	1	2	1	1	1	1	-	-	-
2003.....	1	1	-	-	-	4	-	1	-	1	-	-
2002.....	-	1	3	-	1	-	-	1	1	-	-	-
2001.....	-	3	-	2	-	1	2	-	-	-	-	-
2000.....	-	-	2	1	1	-	1	1	-	-	-	-
1999.....	-	-	-	1	-	1	-	-	-	-	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Peptic ulcer (K25-K28)												
Asian or Pacific Islander 2/, both sexes												
2004.....	3	2	3	3	9	17	21	15	6	4	-	-
2003.....	2	1	8	9	8	14	22	20	10	6	2	-
2002.....	3	6	5	9	7	15	20	13	12	2	1	-
2001.....	2	6	6	12	6	18	25	19	6	4	-	-
2000.....	4	2	2	8	8	20	14	19	7	5	-	-
1999.....	3	2	4	9	16	19	19	17	7	3	1	-
Asian or Pacific Islander 2/, male												
2004.....	3	2	1	2	6	8	13	5	3	1	-	-
2003.....	2	-	4	4	4	9	11	9	6	3	2	-
2002.....	2	3	2	4	6	6	9	4	5	1	-	-
2001.....	1	5	4	8	4	12	12	5	4	3	-	-
2000.....	4	2	-	5	6	14	7	10	7	4	-	-
1999.....	2	1	2	7	9	8	12	6	2	2	-	-
Asian or Pacific Islander 2/, female												
2004.....	-	-	2	1	3	9	8	10	3	3	-	-
2003.....	-	1	4	5	4	5	11	11	4	3	-	-
2002.....	1	3	3	5	1	9	11	9	7	1	1	-
2001.....	1	1	2	4	2	6	13	14	2	1	-	-
2000.....	-	-	2	3	2	6	7	9	-	1	-	-
1999.....	1	1	2	2	7	11	7	11	5	1	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gastric ulcer (K25)												
All races, both sexes												
2004.....	60	50	70	78	96	157	190	173	116	31	6	-
2003.....	62	74	68	72	97	155	205	191	106	28	6	-
2002.....	50	77	61	81	126	155	203	195	106	42	11	-
2001.....	48	51	72	95	136	178	238	196	106	42	7	1
2000.....	64	61	81	99	135	178	198	227	132	38	6	-
1999.....	58	68	64	83	159	197	235	212	109	39	5	-
All races, male												
2004.....	35	29	39	37	44	66	85	54	33	7	1	-
2003.....	44	39	41	42	43	64	81	60	29	2	1	-
2002.....	31	44	42	43	71	67	79	73	28	10	2	-
2001.....	26	36	39	51	69	82	89	56	18	3	1	1
2000.....	44	35	41	53	77	91	69	78	30	12	1	-
1999.....	42	38	37	42	85	82	98	76	22	6	-	-
All races, female												
2004.....	25	21	31	41	52	91	105	119	83	24	5	-
2003.....	18	35	27	30	54	91	124	131	77	26	5	-
2002.....	19	33	19	38	55	88	124	122	78	32	9	-
2001.....	22	15	33	44	67	96	149	140	88	39	6	-
2000.....	20	26	40	46	58	87	129	149	102	26	5	-
1999.....	16	30	27	41	74	115	137	136	87	33	5	-
White, both sexes												
2004.....	45	42	58	61	79	136	170	158	108	28	6	-
2003.....	38	61	57	60	85	131	181	175	93	27	4	-
2002.....	39	63	46	66	112	130	181	174	97	41	10	-
2001.....	38	38	57	71	115	157	214	177	98	35	7	1
2000.....	48	44	67	81	116	156	181	200	124	32	5	-
1999.....	40	50	52	62	139	178	212	183	101	37	4	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gastric ulcer (K25)												
White, male												
2004.....	29	23	31	32	36	55	73	51	32	7	1	-
2003.....	24	32	33	32	36	54	71	53	25	2	-	-
2002.....	22	32	30	33	62	57	70	66	27	9	2	-
2001.....	20	26	29	36	55	73	79	51	15	2	1	1
2000.....	34	23	31	41	62	78	60	67	26	9	1	-
1999.....	28	26	29	31	72	73	88	61	21	5	-	-
White, female												
2004.....	16	19	27	29	43	81	97	107	76	21	5	-
2003.....	14	29	24	28	49	77	110	122	68	25	4	-
2002.....	17	31	16	33	50	73	111	108	70	32	8	-
2001.....	18	12	28	35	60	84	135	126	83	33	6	-
2000.....	14	21	36	40	54	78	121	133	98	23	4	-
1999.....	12	24	23	31	67	105	124	122	80	32	4	-
Black, both sexes												
2004.....	14	8	10	16	14	14	14	7	6	1	-	-
2003.....	22	13	9	8	10	20	19	10	10	-	1	-
2002.....	11	10	12	9	9	20	13	15	6	1	-	-
2001.....	10	10	12	14	18	15	10	13	6	6	-	-
2000.....	14	16	13	15	17	17	13	17	7	4	1	-
1999.....	17	17	11	20	16	12	19	21	6	2	1	-
Black, male												
2004.....	5	6	8	5	6	7	8	1	1	-	-	-
2003.....	19	7	6	8	5	6	8	5	2	-	-	-
2002.....	9	10	10	6	5	6	5	4	-	1	-	-
2001.....	6	7	7	10	13	6	5	4	1	-	-	-
2000.....	8	11	9	11	13	10	8	8	3	1	-	-
1999.....	13	12	7	10	10	7	6	12	1	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gastric ulcer (K25)												
Black, female												
2004.....	9	2	2	11	8	7	6	6	5	1	-	-
2003.....	3	6	3	-	5	14	11	5	8	-	1	-
2002.....	2	-	2	3	4	14	8	11	6	-	-	-
2001.....	4	3	5	4	5	9	5	9	5	6	-	-
2000.....	6	5	4	4	4	7	5	9	4	3	1	-
1999.....	4	5	4	10	6	5	13	9	5	1	1	-
American Indian 1/, both sexes												
2004.....	-	-	-	1	-	2	-	1	-	-	-	-
2003.....	1	-	-	2	-	1	-	1	-	1	-	-
2002.....	-	3	1	2	1	1	-	1	1	-	-	-
2001.....	-	1	1	4	-	-	2	-	1	-	-	-
2000.....	1	-	1	2	-	-	1	1	-	-	-	-
1999.....	1	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, male												
2004.....	-	-	-	-	-	1	-	-	-	-	-	-
2003.....	-	-	-	2	-	1	-	-	-	-	-	-
2002.....	-	2	-	2	1	1	-	-	-	-	-	-
2001.....	-	1	1	2	-	-	-	-	1	-	-	-
2000.....	1	-	1	1	-	-	-	-	-	-	-	-
1999.....	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, female												
2004.....	-	-	-	1	-	1	-	1	-	-	-	-
2003.....	1	-	-	-	-	-	-	1	-	1	-	-
2002.....	-	1	1	-	-	-	-	1	1	-	-	-
2001.....	-	-	-	2	-	-	2	-	-	-	-	-
2000.....	-	-	-	1	-	-	1	1	-	-	-	-
1999.....	-	-	-	-	-	1	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gastric ulcer (K25)												
Asian or Pacific Islander 2/, both sexes												
2004.....	1	-	2	-	3	5	6	7	2	2	-	-
2003.....	1	-	2	2	2	3	5	5	3	-	1	-
2002.....	-	1	2	4	4	4	9	5	2	-	1	-
2001.....	-	2	2	6	3	6	12	6	1	1	-	-
2000.....	1	1	-	1	2	5	3	9	1	2	-	-
1999.....	-	1	1	1	4	6	4	8	2	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	1	-	-	-	2	3	4	2	-	-	-	-
2003.....	1	-	2	-	2	3	2	2	2	-	1	-
2002.....	-	-	2	2	3	3	4	3	1	-	-	-
2001.....	-	2	2	3	1	3	5	1	1	1	-	-
2000.....	1	1	-	-	2	3	1	3	1	2	-	-
1999.....	-	-	1	1	3	2	4	3	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	-	-	2	-	1	2	2	5	2	2	-	-
2003.....	-	-	-	2	-	-	3	3	1	-	-	-
2002.....	-	1	-	2	1	1	5	2	1	-	1	-
2001.....	-	-	-	3	2	3	7	5	-	-	-	-
2000.....	-	-	-	1	-	2	2	6	-	-	-	-
1999.....	-	1	-	-	1	4	-	5	2	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Duodenal ulcer (K26)												
All races, both sexes												
2004.....	48	59	80	87	126	184	216	200	128	44	3	-
2003.....	55	58	63	76	132	183	247	230	123	44	6	-
2002.....	48	82	78	88	149	230	230	235	132	40	5	-
2001.....	59	62	86	102	140	213	277	290	133	50	10	-
2000.....	73	77	94	98	153	224	276	235	138	49	6	-
1999.....	53	56	102	114	171	216	241	239	130	44	3	1
All races, male												
2004.....	32	37	54	49	74	102	88	77	47	13	1	-
2003.....	37	32	44	51	84	104	125	97	40	9	1	-
2002.....	37	55	47	55	84	110	121	92	32	15	2	-
2001.....	43	39	60	63	77	110	123	118	48	11	2	-
2000.....	59	54	57	60	91	116	118	85	50	14	2	-
1999.....	39	34	71	83	113	123	121	87	44	10	1	1
All races, female												
2004.....	16	22	26	38	52	82	128	123	81	31	2	-
2003.....	18	26	19	25	48	79	122	133	83	35	5	-
2002.....	11	27	31	33	65	120	109	143	100	25	3	-
2001.....	16	23	26	39	63	103	154	172	85	39	8	-
2000.....	14	23	37	38	62	108	158	150	88	35	4	-
1999.....	14	22	31	31	58	93	120	152	86	34	2	-
White, both sexes												
2004.....	41	50	67	77	112	169	199	187	121	43	3	-
2003.....	47	45	49	65	118	159	224	212	118	41	5	-
2002.....	34	64	66	74	134	210	223	221	130	38	5	-
2001.....	44	49	73	91	126	192	262	277	130	47	10	-
2000.....	54	66	83	83	136	203	260	219	130	47	6	-
1999.....	41	49	92	99	153	196	224	225	121	37	3	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Duodenal ulcer (K26)												
White, male												
2004.....	26	32	47	43	64	94	81	71	43	13	1	-
2003.....	31	23	36	44	76	92	114	87	37	8	1	-
2002.....	27	45	41	48	75	98	118	89	31	15	2	-
2001.....	30	32	51	58	65	97	115	113	46	10	2	-
2000.....	42	46	53	53	81	104	111	79	46	14	2	-
1999.....	29	30	63	71	104	114	114	83	42	9	1	1
White, female												
2004.....	15	18	20	34	48	75	118	116	78	30	2	-
2003.....	16	22	13	21	42	67	110	125	81	33	4	-
2002.....	7	19	25	26	59	112	105	132	99	23	3	-
2001.....	14	17	22	33	61	95	147	164	84	37	8	-
2000.....	12	20	30	30	55	99	149	140	84	33	4	-
1999.....	12	19	29	28	49	82	110	142	79	28	2	-
Black, both sexes												
2004.....	6	8	12	10	9	9	11	7	5	1	-	-
2003.....	7	13	9	6	10	16	16	13	2	1	1	-
2002.....	12	15	10	11	13	14	4	9	-	2	-	-
2001.....	14	9	12	9	10	12	11	7	2	2	-	-
2000.....	16	9	8	10	15	15	10	12	4	1	-	-
1999.....	10	7	9	11	10	13	14	8	7	5	-	-
Black, male												
2004.....	5	5	6	6	6	6	4	3	2	-	-	-
2003.....	5	9	6	4	7	9	7	7	1	-	-	-
2002.....	9	8	6	7	8	9	3	3	-	-	-	-
2001.....	12	5	8	3	8	6	5	4	1	-	-	-
2000.....	14	6	4	4	9	9	4	3	-	-	-	-
1999.....	8	4	7	9	4	6	5	3	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Duodenal ulcer (K26)												
Asian or Pacific Islander 2/, both sexes												
2004.....	1	-	1	-	4	6	5	6	2	-	-	-
2003.....	-	-	5	5	4	6	7	5	3	2	-	-
2002.....	2	3	2	3	1	5	3	5	2	-	-	-
2001.....	-	3	1	2	3	7	4	6	1	1	-	-
2000.....	3	1	1	4	1	6	5	4	4	1	-	-
1999.....	2	-	1	4	8	7	3	6	2	2	-	-
Asian or Pacific Islander 2/, male												
2004.....	1	-	1	-	3	2	3	3	2	-	-	-
2003.....	-	-	2	3	1	3	4	3	2	1	-	-
2002.....	1	2	-	-	1	2	-	-	1	-	-	-
2001.....	-	2	1	2	3	5	3	1	1	1	-	-
2000.....	3	1	-	2	1	3	2	3	4	-	-	-
1999.....	2	-	1	3	5	3	2	1	1	1	-	-
Asian or Pacific Islander 2/, female												
2004.....	-	-	-	-	1	4	2	3	-	-	-	-
2003.....	-	-	3	2	3	3	3	2	1	1	-	-
2002.....	1	1	2	3	-	3	3	5	1	-	-	-
2001.....	-	1	-	-	-	2	1	5	-	-	-	-
2000.....	-	-	1	2	-	3	3	1	-	1	-	-
1999.....	-	-	-	1	3	4	1	5	1	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Peptic ulcer, site unspecified (K27)												
All races, both sexes												
2004.....	40	49	61	80	105	159	226	227	163	65	16	-
2003.....	41	50	69	76	121	198	271	259	154	64	18	-
2002.....	56	47	62	80	131	198	258	277	163	64	17	-
2001.....	39	69	73	93	145	218	301	301	212	91	21	-
2000.....	37	62	67	97	155	232	290	331	220	88	20	-
1999.....	31	61	69	117	185	246	352	315	226	79	26	1
All races, male												
2004.....	28	31	35	44	50	87	96	77	38	15	5	-
2003.....	29	33	47	46	70	102	120	102	53	14	3	-
2002.....	38	27	29	43	77	104	131	112	51	14	1	-
2001.....	26	47	48	63	70	120	120	102	56	33	2	-
2000.....	27	45	51	57	98	130	132	126	69	22	6	-
1999.....	21	38	46	71	107	122	158	113	70	12	1	-
All races, female												
2004.....	12	18	26	36	55	72	130	150	125	50	11	-
2003.....	12	17	22	30	51	96	151	157	101	50	15	-
2002.....	18	20	33	37	54	94	127	165	112	50	16	-
2001.....	13	22	25	30	75	98	181	199	156	58	19	-
2000.....	10	17	16	40	57	102	158	205	151	66	14	-
1999.....	10	23	23	46	78	124	194	202	156	67	25	1
White, both sexes												
2004.....	30	37	51	70	87	137	205	211	152	57	15	-
2003.....	30	41	58	68	107	177	240	239	146	57	15	-
2002.....	45	34	49	69	113	175	241	259	148	58	13	-
2001.....	29	50	59	79	130	193	272	276	197	88	21	-
2000.....	28	52	57	83	140	203	270	304	207	84	17	-
1999.....	25	52	54	98	159	222	318	300	213	75	21	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Peptic ulcer, site unspecified (K27)												
White, male												
2004.....	22	24	28	39	43	75	84	72	35	14	4	-
2003.....	21	27	39	42	60	90	99	98	49	11	2	-
2002.....	28	19	24	37	65	92	119	103	46	12	1	-
2001.....	18	32	39	52	62	104	107	92	52	31	2	-
2000.....	19	41	45	50	88	114	123	115	65	19	5	-
1999.....	19	31	35	58	93	112	143	108	69	10	1	-
White, female												
2004.....	8	13	23	31	44	62	121	139	117	43	11	-
2003.....	9	14	19	26	47	87	141	141	97	46	13	-
2002.....	17	15	25	32	48	83	122	156	102	46	12	-
2001.....	11	18	20	27	68	89	165	184	145	57	19	-
2000.....	9	11	12	33	52	89	147	189	142	65	12	-
1999.....	6	21	19	40	66	110	175	192	144	65	20	-
Black, both sexes												
2004.....	9	10	10	7	14	16	11	14	8	6	1	-
2003.....	10	7	10	6	12	14	22	11	4	3	2	-
2002.....	10	10	11	9	16	17	8	14	7	5	4	-
2001.....	8	16	10	10	13	18	20	18	12	1	-	-
2000.....	7	10	9	10	10	20	14	21	11	3	3	-
1999.....	5	8	13	14	22	17	23	12	11	4	3	1
Black, male												
2004.....	5	5	7	3	6	9	6	5	2	-	1	-
2003.....	7	6	8	3	9	9	17	1	2	1	-	-
2002.....	9	6	5	4	10	11	6	7	2	1	-	-
2001.....	7	14	7	8	6	11	9	7	3	1	-	-
2000.....	6	4	6	3	7	8	5	7	2	2	1	-
1999.....	2	6	11	10	12	6	10	3	1	2	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Peptic ulcer, site unspecified (K27)												
Asian or Pacific Islander 2/, both sexes												
2004.....	1	2	-	3	2	6	10	2	2	2	-	-
2003.....	1	1	1	2	2	5	9	9	4	4	1	-
2002.....	1	2	1	2	2	6	8	3	8	1	-	-
2001.....	2	1	3	4	-	5	9	7	3	2	-	-
2000.....	-	-	1	3	5	9	5	6	2	1	-	-
1999.....	1	1	2	4	3	6	11	3	2	-	1	-
Asian or Pacific Islander 2/, male												
2004.....	1	2	-	2	1	3	6	-	1	1	-	-
2003.....	1	-	-	1	1	3	4	3	2	2	1	-
2002.....	1	1	-	2	2	1	5	1	3	1	-	-
2001.....	1	1	1	3	-	4	4	3	1	1	-	-
2000.....	-	-	-	3	3	8	3	4	2	1	-	-
1999.....	-	1	-	3	1	3	5	2	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	-	-	-	1	1	3	4	2	1	1	-	-
2003.....	-	1	1	1	1	2	5	6	2	2	-	-
2002.....	-	1	1	-	-	5	3	2	5	-	-	-
2001.....	1	-	2	1	-	1	5	4	2	1	-	-
2000.....	-	-	1	-	2	1	2	2	-	-	-	-
1999.....	1	-	2	1	2	3	6	1	2	-	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gastrojejunal ulcer (K28)												
All races, both sexes												
2004.....	2	1	4	5	5	5	11	6	6	1	-	-
2003.....	1	1	-	4	9	9	11	5	8	2	-	-
2002.....	-	3	1	5	5	9	8	7	6	2	-	-
2001.....	1	1	4	4	3	6	7	1	7	1	2	-
2000.....	3	-	1	5	6	8	9	8	8	2	1	-
1999.....	-	1	1	-	7	6	7	5	5	2	-	-
All races, male												
2004.....	2	1	2	3	3	2	4	1	2	-	-	-
2003.....	-	1	-	2	8	3	7	3	4	-	-	-
2002.....	-	3	-	1	4	6	4	2	5	-	-	-
2001.....	-	1	3	2	2	3	2	1	3	-	-	-
2000.....	1	-	1	2	4	4	7	2	1	1	-	-
1999.....	-	1	-	-	3	4	4	3	2	1	-	-
All races, female												
2004.....	-	-	2	2	2	3	7	5	4	1	-	-
2003.....	1	-	-	2	1	6	4	2	4	2	-	-
2002.....	-	-	1	4	1	3	4	5	1	2	-	-
2001.....	1	-	1	2	1	3	5	-	4	1	2	-
2000.....	2	-	-	3	2	4	2	6	7	1	1	-
1999.....	-	-	1	-	4	2	3	2	3	1	-	-
White, both sexes												
2004.....	2	-	4	4	3	5	9	6	5	1	-	-
2003.....	1	1	-	4	7	8	10	4	8	2	-	-
2002.....	-	3	-	4	4	9	7	7	5	1	-	-
2001.....	1	1	4	4	3	6	7	1	6	1	1	-
2000.....	2	-	1	3	5	7	6	8	7	1	1	-
1999.....	-	1	-	-	6	4	6	5	4	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gastrojejunal ulcer (K28)												
Asian or Pacific Islander 2/, both sexes												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	1	1	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	1	-	-
2001.....	-	-	-	-	-	-	-	-	1	-	-	-
2000.....	-	-	-	-	-	-	1	-	-	1	-	-
1999.....	-	-	-	-	1	-	1	-	1	1	-	-
Asian or Pacific Islander 2/, male												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	1	1	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	1	-	-	-
2000.....	-	-	-	-	-	-	1	-	-	1	-	-
1999.....	-	-	-	-	-	-	1	-	1	1	-	-
Asian or Pacific Islander 2/, female												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	1	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	1	-	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gastritis and duodenitis (K29)												
All races, both sexes												
2004.....	8	27	25	18	24	49	53	54	48	20	4	-
2003.....	28	14	21	12	36	36	55	71	39	25	3	-
2002.....	15	18	17	14	26	46	57	59	41	26	8	-
2001.....	9	15	13	22	29	42	49	84	58	22	2	-
2000.....	12	25	22	21	29	42	60	73	48	18	9	-
1999.....	17	16	11	22	27	41	72	72	62	16	8	-
All races, male												
2004.....	6	17	20	11	13	23	23	24	20	1	-	-
2003.....	21	7	15	7	17	23	24	25	10	4	1	-
2002.....	9	15	10	7	15	25	21	23	11	6	-	-
2001.....	6	13	6	11	14	18	17	37	17	8	-	-
2000.....	9	15	15	15	15	16	22	24	14	3	-	-
1999.....	15	10	4	10	13	25	30	23	19	1	1	-
All races, female												
2004.....	2	10	5	7	11	26	30	30	28	19	4	-
2003.....	7	7	6	5	19	13	31	46	29	21	2	-
2002.....	6	3	7	7	11	21	36	36	30	20	8	-
2001.....	3	2	7	11	15	24	32	47	41	14	2	-
2000.....	3	10	7	6	14	26	38	49	34	15	9	-
1999.....	2	6	7	12	14	16	42	49	43	15	7	-
White, both sexes												
2004.....	5	26	17	15	21	40	42	52	42	17	4	-
2003.....	23	12	19	10	31	29	46	64	36	20	3	-
2002.....	11	13	15	11	23	43	50	53	37	24	7	-
2001.....	6	13	10	18	22	39	42	80	57	21	2	-
2000.....	9	21	16	19	27	37	58	61	47	17	9	-
1999.....	16	14	11	15	22	31	57	66	59	14	8	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gastritis and duodenitis (K29)												
White, male												
2004.....	3	17	13	11	11	21	18	22	19	1	-	-
2003.....	17	6	14	5	12	18	21	24	8	4	1	-
2002.....	6	11	9	7	13	25	21	20	11	6	-	-
2001.....	3	11	5	8	11	18	15	35	17	7	-	-
2000.....	6	13	12	14	14	13	21	20	14	3	-	-
1999.....	14	8	4	7	9	20	22	22	17	1	1	-
White, female												
2004.....	2	9	4	4	10	19	24	30	23	16	4	-
2003.....	6	6	5	5	19	11	25	40	28	16	2	-
2002.....	5	2	6	4	10	18	29	33	26	18	7	-
2001.....	3	2	5	10	11	21	27	45	40	14	2	-
2000.....	3	8	4	5	13	24	37	41	33	14	9	-
1999.....	2	6	7	8	13	11	35	44	42	13	7	-
Black, both sexes												
2004.....	2	1	6	3	2	7	7	1	5	1	-	-
2003.....	3	2	2	2	2	5	5	5	2	5	-	-
2002.....	4	4	-	3	2	3	5	5	2	2	1	-
2001.....	3	2	3	4	6	2	4	2	1	1	-	-
2000.....	1	4	6	2	2	4	2	11	-	1	-	-
1999.....	1	2	-	6	5	6	13	6	2	2	-	-
Black, male												
2004.....	2	-	5	-	1	1	3	1	1	-	-	-
2003.....	3	1	1	2	2	4	1	-	2	-	-	-
2002.....	3	3	-	-	2	-	-	3	-	-	-	-
2001.....	3	2	1	3	2	-	2	1	-	1	-	-
2000.....	1	2	3	1	1	2	1	3	-	-	-	-
1999.....	1	2	-	2	4	3	6	1	2	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gastritis and duodenitis (K29)												
Black, female												
2004.....	-	1	1	3	1	6	4	-	4	1	-	-
2003.....	-	1	1	-	-	1	4	5	-	5	-	-
2002.....	1	1	-	3	-	3	5	2	2	2	1	-
2001.....	-	-	2	1	4	2	2	1	1	-	-	-
2000.....	-	2	3	1	1	2	1	8	-	1	-	-
1999.....	-	-	-	4	1	3	7	5	-	2	-	-
American Indian 1/, both sexes												
2004.....	1	-	1	-	-	2	-	-	-	-	-	-
2003.....	-	-	-	-	1	-	-	1	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	1	-	-	-	-
2000.....	1	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	1	-	1	-	-	1	-	-	-
American Indian 1/, male												
2004.....	1	-	1	-	-	1	-	-	-	-	-	-
2003.....	-	-	-	-	1	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	1	-	-	-	-
2000.....	1	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	1	-	1	-	-	-	-	-	-
American Indian 1/, female												
2004.....	-	-	-	-	-	1	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	1	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gastritis and duodenitis (K29)												
Asian or Pacific Islander 2/, both sexes												
2004.....	-	-	1	-	1	-	4	1	1	2	-	-
2003.....	2	-	-	-	2	2	4	1	1	-	-	-
2002.....	-	1	2	-	1	-	2	1	2	-	-	-
2001.....	-	-	-	-	1	1	3	1	-	-	-	-
2000.....	1	-	-	-	-	1	-	1	1	-	-	-
1999.....	-	-	-	-	-	3	2	-	-	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	-	-	1	-	1	-	2	1	-	-	-	-
2003.....	1	-	-	-	2	1	2	1	-	-	-	-
2002.....	-	1	1	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	1	-	-	-	-	-	-	-
2000.....	1	-	-	-	-	1	-	1	-	-	-	-
1999.....	-	-	-	-	-	1	2	-	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	-	-	-	-	-	-	2	-	1	2	-	-
2003.....	1	-	-	-	-	1	2	-	1	-	-	-
2002.....	-	-	1	-	1	-	2	1	2	-	-	-
2001.....	-	-	-	-	-	1	3	1	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	1	-	-	-
1999.....	-	-	-	-	-	2	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Dyspepsia and other diseases of stomach and duodenum (K30-K31)												
All races, both sexes												
2004.....	38	33	39	41	47	85	95	75	49	21	2	-
2003.....	27	26	29	43	62	75	102	83	63	16	4	-
2002.....	28	30	30	43	52	85	92	84	45	16	4	-
2001.....	28	16	24	40	51	71	90	89	48	19	2	-
2000.....	27	28	27	43	62	86	78	89	59	15	6	-
1999.....	14	19	33	31	54	74	94	98	55	18	3	-
All races, male												
2004.....	16	18	16	24	22	37	43	31	20	3	-	-
2003.....	13	8	12	19	33	33	41	23	12	4	-	-
2002.....	11	9	13	25	26	39	32	24	13	2	1	-
2001.....	11	7	13	22	23	26	42	20	16	6	1	-
2000.....	13	11	9	30	34	37	32	32	16	4	1	-
1999.....	5	12	12	19	25	37	29	32	21	4	-	-
All races, female												
2004.....	22	15	23	17	25	48	52	44	29	18	2	-
2003.....	14	18	17	24	29	42	61	60	51	12	4	-
2002.....	17	21	17	18	26	46	60	60	32	14	3	-
2001.....	17	9	11	18	28	45	48	69	32	13	1	-
2000.....	14	17	18	13	28	49	46	57	43	11	5	-
1999.....	9	7	21	12	29	37	65	66	34	14	3	-
White, both sexes												
2004.....	25	29	32	31	38	76	89	70	41	20	2	-
2003.....	21	21	25	36	57	62	90	74	60	15	4	-
2002.....	22	26	27	36	46	77	77	77	39	15	4	-
2001.....	26	13	19	35	44	58	83	81	48	18	2	-
2000.....	25	25	23	32	56	76	72	85	52	14	6	-
1999.....	12	19	27	25	44	71	85	88	51	18	3	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Dyspepsia and other diseases of stomach and duodenum (K30-K31)												
White, male												
2004.....	10	16	11	16	17	34	39	28	16	3	-	-
2003.....	10	5	12	15	31	29	35	23	11	4	-	-
2002.....	9	8	12	21	22	36	27	22	10	2	1	-
2001.....	9	6	10	18	19	21	40	19	16	5	1	-
2000.....	11	9	7	22	30	32	31	30	14	4	1	-
1999.....	3	12	11	14	19	37	26	30	19	4	-	-
White, female												
2004.....	15	13	21	15	21	42	50	42	25	17	2	-
2003.....	11	16	13	21	26	33	55	51	49	11	4	-
2002.....	13	18	15	15	24	41	50	55	29	13	3	-
2001.....	17	7	9	17	25	37	43	62	32	13	1	-
2000.....	14	16	16	10	26	44	41	55	38	10	5	-
1999.....	9	7	16	11	25	34	59	58	32	14	3	-
Black, both sexes												
2004.....	13	4	6	10	9	9	6	5	7	1	-	-
2003.....	5	4	4	7	3	9	9	6	2	1	-	-
2002.....	6	2	3	3	5	8	11	6	6	1	-	-
2001.....	2	3	5	5	6	12	6	8	-	1	-	-
2000.....	2	3	4	10	5	9	5	2	7	1	-	-
1999.....	2	-	6	4	9	2	6	10	3	-	-	-
Black, male												
2004.....	6	2	4	8	5	3	4	3	4	-	-	-
2003.....	3	2	-	4	2	1	4	-	-	-	-	-
2002.....	2	-	1	1	3	3	5	1	3	-	-	-
2001.....	2	1	3	4	3	5	2	1	-	1	-	-
2000.....	2	2	2	8	3	4	1	-	2	-	-	-
1999.....	2	-	1	3	5	-	1	2	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Dyspepsia and other diseases of stomach and duodenum (K30-K31)												
Asian or Pacific Islander 2/, both sexes												
2004.....	-	-	-	-	-	-	-	-	1	-	-	-
2003.....	1	1	-	-	1	3	2	2	1	-	-	-
2002.....	-	1	-	1	-	-	2	1	-	-	-	-
2001.....	-	-	-	-	-	1	1	-	-	-	-	-
2000.....	-	-	-	1	1	1	1	2	-	-	-	-
1999.....	-	-	-	2	1	1	3	-	1	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	1	-	-	-	3	1	-	1	-	-	-
2002.....	-	1	-	1	-	-	-	1	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	1	1	-	2	-	-	-	-
1999.....	-	-	-	2	1	-	2	-	1	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	-	-	-	-	-	-	-	-	1	-	-	-
2003.....	1	-	-	-	1	-	1	2	-	-	-	-
2002.....	-	-	-	-	-	-	2	-	-	-	-	-
2001.....	-	-	-	-	-	1	1	-	-	-	-	-
2000.....	-	-	-	1	-	-	1	-	-	-	-	-
1999.....	-	-	-	-	-	1	1	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of appendix (K35-K38)												
All races, both sexes												
2004.....	31	24	28	33	48	58	80	46	30	8	1	-
2003.....	25	18	25	29	35	56	76	42	28	7	1	-
2002.....	30	28	36	32	57	61	59	56	25	10	3	-
2001.....	15	22	23	34	55	37	53	51	30	11	1	-
2000.....	25	28	32	26	45	53	68	39	24	4	1	-
1999.....	24	22	26	30	39	43	53	40	18	3	1	-
All races, male												
2004.....	22	10	19	17	26	33	37	26	8	3	-	-
2003.....	16	13	17	20	26	28	47	21	10	3	-	-
2002.....	20	18	27	24	32	35	32	22	6	1	2	-
2001.....	11	13	17	23	34	22	30	20	12	4	1	-
2000.....	17	20	23	15	25	30	30	15	9	-	-	-
1999.....	19	17	17	18	20	28	19	23	9	2	1	-
All races, female												
2004.....	9	14	9	16	22	25	43	20	22	5	1	-
2003.....	9	5	8	9	9	28	29	21	18	4	1	-
2002.....	10	10	9	8	25	26	27	34	19	9	1	-
2001.....	4	9	6	11	21	15	23	31	18	7	-	-
2000.....	8	8	9	11	20	23	38	24	15	4	1	-
1999.....	5	5	9	12	19	15	34	17	9	1	-	-
White, both sexes												
2004.....	24	22	22	26	43	52	65	42	29	6	1	-
2003.....	19	15	17	23	30	46	71	39	25	7	1	-
2002.....	23	25	31	26	45	56	56	52	22	10	2	-
2001.....	14	13	16	28	52	33	44	44	28	10	1	-
2000.....	19	21	29	20	34	42	58	38	22	4	1	-
1999.....	22	15	15	25	32	38	46	37	17	3	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of appendix (K35-K38)												
White, male												
2004.....	17	10	16	13	23	30	33	22	7	2	-	-
2003.....	13	11	12	16	22	23	44	20	10	3	-	-
2002.....	16	17	23	19	25	32	30	19	4	1	1	-
2001.....	10	10	12	19	33	20	26	14	11	3	1	-
2000.....	12	14	20	13	18	26	25	14	8	-	-	-
1999.....	17	12	12	15	15	24	15	21	9	2	1	-
White, female												
2004.....	7	12	6	13	20	22	32	20	22	4	1	-
2003.....	6	4	5	7	8	23	27	19	15	4	1	-
2002.....	7	8	8	7	20	24	26	33	18	9	1	-
2001.....	4	3	4	9	19	13	18	30	17	7	-	-
2000.....	7	7	9	7	16	16	33	24	14	4	1	-
1999.....	5	3	3	10	17	14	31	16	8	1	-	-
Black, both sexes												
2004.....	6	1	6	6	4	4	12	4	1	-	-	-
2003.....	3	3	7	4	5	9	2	3	2	-	-	-
2002.....	6	3	4	5	9	3	2	4	3	-	1	-
2001.....	1	9	7	6	1	3	8	3	2	1	-	-
2000.....	4	7	3	6	10	9	7	1	2	-	-	-
1999.....	2	7	10	3	7	5	6	2	1	-	-	-
Black, male												
2004.....	5	-	3	4	3	3	3	4	1	-	-	-
2003.....	1	2	4	3	4	4	1	1	-	-	-	-
2002.....	4	1	3	4	5	2	2	3	2	-	1	-
2001.....	1	3	5	4	-	2	3	3	1	1	-	-
2000.....	3	6	3	2	7	3	3	1	1	-	-	-
1999.....	2	5	4	2	5	4	4	2	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of appendix (K35-K38)												
Asian or Pacific Islander 2/, both sexes												
2004.....	-	-	-	1	1	2	3	-	-	2	-	-
2003.....	3	-	-	2	-	1	3	-	1	-	-	-
2002.....	-	-	1	1	3	2	1	-	-	-	-	-
2001.....	-	-	-	-	2	1	1	4	-	-	-	-
2000.....	2	-	-	-	1	1	3	-	-	-	-	-
1999.....	-	-	-	2	-	-	1	1	-	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	-	-	-	-	-	-	1	-	-	1	-	-
2003.....	2	-	-	1	-	1	2	-	-	-	-	-
2002.....	-	-	1	1	2	1	-	-	-	-	-	-
2001.....	-	-	-	-	1	-	1	3	-	-	-	-
2000.....	2	-	-	-	-	1	2	-	-	-	-	-
1999.....	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	-	-	-	1	1	2	2	-	-	1	-	-
2003.....	1	-	-	1	-	-	1	-	1	-	-	-
2002.....	-	-	-	-	1	1	1	-	-	-	-	-
2001.....	-	-	-	-	1	1	-	1	-	-	-	-
2000.....	-	-	-	-	1	-	1	-	-	-	-	-
1999.....	-	-	-	1	-	-	1	1	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hernia (K40-K46)												
All races, both sexes												
2004.....	45	70	60	90	149	217	292	296	166	61	10	-
2003.....	60	76	85	101	136	203	259	296	202	56	9	-
2002.....	44	55	71	90	135	208	281	320	172	85	11	-
2001.....	58	57	74	81	132	185	258	269	190	88	6	-
2000.....	58	51	62	84	126	224	242	294	188	63	10	-
1999.....	39	36	71	77	138	186	265	269	196	56	8	-
All races, male												
2004.....	25	35	27	38	82	99	99	101	54	14	1	-
2003.....	35	30	42	48	63	93	106	95	62	17	4	-
2002.....	19	27	29	40	63	84	105	97	45	20	-	-
2001.....	31	23	41	40	54	71	102	88	48	19	1	-
2000.....	32	20	29	39	53	91	104	96	42	12	4	-
1999.....	20	21	39	34	59	75	117	96	51	8	2	-
All races, female												
2004.....	20	35	33	52	67	118	193	195	112	47	9	-
2003.....	25	46	43	53	73	110	153	201	140	39	5	-
2002.....	25	28	42	50	72	124	176	223	127	65	11	-
2001.....	27	34	33	41	78	114	156	181	142	69	5	-
2000.....	26	31	33	45	73	133	138	198	146	51	6	-
1999.....	19	15	32	43	79	111	148	173	145	48	6	-
White, both sexes												
2004.....	39	57	50	78	130	201	268	272	156	55	9	-
2003.....	48	60	71	88	120	185	230	276	184	55	7	-
2002.....	34	47	55	71	121	197	258	296	167	83	10	-
2001.....	47	45	62	69	116	164	235	251	179	77	4	-
2000.....	47	45	54	68	117	198	223	270	176	57	9	-
1999.....	30	32	62	60	123	163	245	248	183	54	6	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hernia (K40-K46)												
White, male												
2004.....	22	29	24	30	73	92	90	92	49	13	1	-
2003.....	30	27	35	42	54	86	94	87	57	17	2	-
2002.....	17	24	23	31	55	80	98	89	42	19	-	-
2001.....	23	16	34	36	49	64	91	75	45	18	1	-
2000.....	25	18	25	31	51	85	95	88	39	10	4	-
1999.....	17	21	36	29	57	68	109	86	44	7	2	-
White, female												
2004.....	17	28	26	48	57	109	178	180	107	42	8	-
2003.....	18	33	36	46	66	99	136	189	127	38	5	-
2002.....	17	23	32	40	66	117	160	207	125	64	10	-
2001.....	24	29	28	33	67	100	144	176	134	59	3	-
2000.....	22	27	29	37	66	113	128	182	137	47	5	-
1999.....	13	11	26	31	66	95	136	162	139	47	4	-
Black, both sexes												
2004.....	6	12	8	10	18	11	20	18	8	6	1	-
2003.....	11	16	12	12	12	17	22	14	15	1	1	-
2002.....	8	8	13	15	12	10	20	19	4	2	1	-
2001.....	11	12	11	10	15	17	17	16	9	9	2	-
2000.....	8	5	8	14	7	22	16	24	12	3	1	-
1999.....	7	4	8	12	12	20	18	15	12	2	2	-
Black, male												
2004.....	3	6	3	7	9	4	7	7	3	1	-	-
2003.....	5	3	7	6	7	6	9	6	4	-	1	-
2002.....	2	3	4	8	7	4	7	6	2	1	-	-
2001.....	8	7	6	2	5	6	9	11	3	1	-	-
2000.....	5	1	4	7	2	4	8	8	3	-	-	-
1999.....	3	-	2	5	2	5	8	7	6	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hernia (K40-K46)												
Black, female												
2004.....	3	6	5	3	9	7	13	11	5	5	1	-
2003.....	6	13	5	6	5	11	13	8	11	1	-	-
2002.....	6	5	9	7	5	6	13	13	2	1	1	-
2001.....	3	5	5	8	10	11	8	5	6	8	2	-
2000.....	3	4	4	7	5	18	8	16	9	3	1	-
1999.....	4	4	6	7	10	15	10	8	6	1	2	-
American Indian 1/, both sexes												
2004.....	-	1	1	2	1	1	2	3	-	-	-	-
2003.....	1	-	1	1	3	-	1	3	2	-	1	-
2002.....	2	-	1	3	-	1	3	1	-	-	-	-
2001.....	-	-	1	1	-	-	1	1	1	-	-	-
2000.....	2	1	-	2	1	1	1	-	-	-	-	-
1999.....	1	-	1	1	1	-	-	2	-	-	-	-
American Indian 1/, male												
2004.....	-	-	-	1	-	1	-	2	-	-	-	-
2003.....	-	-	-	-	1	-	-	2	1	-	1	-
2002.....	-	-	1	1	-	-	-	-	-	-	-	-
2001.....	-	-	1	1	-	-	-	1	-	-	-	-
2000.....	1	1	-	1	-	-	-	-	-	-	-	-
1999.....	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, female												
2004.....	-	1	1	1	1	-	2	1	-	-	-	-
2003.....	1	-	1	1	2	-	1	1	1	-	-	-
2002.....	2	-	-	2	-	1	3	1	-	-	-	-
2001.....	-	-	-	-	-	-	1	-	1	-	-	-
2000.....	1	-	-	1	1	1	1	-	-	-	-	-
1999.....	1	-	-	1	1	-	-	2	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hernia (K40-K46)												
Asian or Pacific Islander 2/, both sexes												
2004.....	-	-	1	-	-	4	2	3	2	-	-	-
2003.....	-	-	1	-	1	1	6	3	1	-	-	-
2002.....	-	-	2	1	2	-	-	4	1	-	-	-
2001.....	-	-	-	1	1	4	5	1	1	2	-	-
2000.....	1	-	-	-	1	3	2	-	-	3	-	-
1999.....	1	-	-	4	2	3	2	4	1	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	-	-	-	-	-	2	2	-	2	-	-	-
2003.....	-	-	-	-	1	1	3	-	-	-	-	-
2002.....	-	-	1	-	1	-	-	2	1	-	-	-
2001.....	-	-	-	1	-	1	2	1	-	-	-	-
2000.....	1	-	-	-	-	2	1	-	-	2	-	-
1999.....	-	-	-	-	-	2	-	3	1	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	-	-	1	-	-	2	-	3	-	-	-	-
2003.....	-	-	1	-	-	-	3	3	1	-	-	-
2002.....	-	-	1	1	1	-	-	2	-	-	-	-
2001.....	-	-	-	-	1	3	3	-	1	2	-	-
2000.....	-	-	-	-	1	1	1	-	-	1	-	-
1999.....	1	-	-	4	2	1	2	1	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Crohn's disease and ulcerative colitis (K50-K51)												
All races, both sexes												
2004.....	60	70	74	70	91	134	136	94	52	14	2	-
2003.....	49	60	63	76	91	119	124	93	42	8	1	-
2002.....	49	39	48	70	86	126	124	97	38	6	1	-
2001.....	41	66	51	64	96	116	123	109	46	10	1	-
2000.....	39	49	41	60	99	119	111	94	40	4	1	-
1999.....	49	45	42	55	95	118	124	88	35	12	1	-
All races, male												
2004.....	29	34	41	31	41	57	47	26	10	4	1	-
2003.....	23	33	29	30	41	53	48	34	15	2	-	-
2002.....	19	18	18	36	40	51	47	27	9	2	-	-
2001.....	21	32	26	33	51	49	44	34	10	1	-	-
2000.....	15	21	17	25	41	50	30	23	13	-	-	-
1999.....	21	22	16	30	42	48	45	22	9	1	-	-
All races, female												
2004.....	31	36	33	39	50	77	89	68	42	10	1	-
2003.....	26	27	34	46	50	66	76	59	27	6	1	-
2002.....	30	21	30	34	46	75	77	70	29	4	1	-
2001.....	20	34	25	31	45	67	79	75	36	9	1	-
2000.....	24	28	24	35	58	69	81	71	27	4	1	-
1999.....	28	23	26	25	53	70	79	66	26	11	1	-
White, both sexes												
2004.....	55	60	68	64	84	129	132	91	51	14	2	-
2003.....	44	56	56	71	83	116	119	90	40	8	1	-
2002.....	44	36	45	67	80	120	118	95	38	6	1	-
2001.....	35	58	49	60	92	111	118	99	45	9	1	-
2000.....	37	44	39	58	95	113	108	92	36	4	1	-
1999.....	44	37	40	55	92	108	122	85	34	11	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Crohn's disease and ulcerative colitis (K50-K51)												
White, male												
2004.....	25	26	37	28	39	54	44	24	10	4	1	-
2003.....	21	31	25	28	38	51	47	33	14	2	-	-
2002.....	17	15	17	33	36	48	45	26	9	2	-	-
2001.....	17	27	24	30	49	48	43	31	9	1	-	-
2000.....	14	20	17	24	40	46	30	23	12	-	-	-
1999.....	18	20	15	30	41	43	44	22	9	1	-	-
White, female												
2004.....	30	34	31	36	45	75	88	67	41	10	1	-
2003.....	23	25	31	43	45	65	72	57	26	6	1	-
2002.....	27	21	28	34	44	72	73	69	29	4	1	-
2001.....	18	31	25	30	43	63	75	68	36	8	1	-
2000.....	23	24	22	34	55	67	78	69	24	4	1	-
1999.....	26	17	25	25	51	65	78	63	25	10	1	-
Black, both sexes												
2004.....	5	10	6	4	7	4	4	3	1	-	-	-
2003.....	4	4	5	4	7	3	4	2	2	-	-	-
2002.....	4	2	3	2	4	5	6	1	-	-	-	-
2001.....	5	8	1	4	2	4	5	9	-	1	-	-
2000.....	1	4	2	1	3	5	3	2	4	-	-	-
1999.....	5	7	2	-	2	9	1	2	1	1	-	-
Black, male												
2004.....	4	8	4	2	2	3	3	2	-	-	-	-
2003.....	2	2	3	1	3	2	-	-	1	-	-	-
2002.....	2	2	1	2	4	2	2	1	-	-	-	-
2001.....	4	5	1	3	-	1	1	2	-	-	-	-
2000.....	-	1	-	1	-	4	-	1	1	-	-	-
1999.....	3	2	1	-	1	4	1	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Crohn's disease and ulcerative colitis (K50-K51)												
Black, female												
2004.....	1	2	2	2	5	1	1	1	1	-	-	-
2003.....	2	2	2	3	4	1	4	2	1	-	-	-
2002.....	2	-	2	-	-	3	4	-	-	-	-	-
2001.....	1	3	-	1	2	3	4	7	-	1	-	-
2000.....	1	3	2	-	3	1	3	2	3	-	-	-
1999.....	2	5	1	-	1	5	-	2	1	1	-	-
American Indian 1/, both sexes												
2004.....	-	-	-	1	-	1	-	-	-	-	-	-
2003.....	-	-	1	-	-	-	-	-	-	-	-	-
2002.....	-	1	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	2	-	-	-	-	-	-	-
2000.....	1	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	1	-	-	1	-	-	-	-	-	-	-
American Indian 1/, male												
2004.....	-	-	-	1	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	1	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	2	-	-	-	-	-	-	-
2000.....	1	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, female												
2004.....	-	-	-	-	-	1	-	-	-	-	-	-
2003.....	-	-	1	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	1	-	-	1	-	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Crohn's disease and ulcerative colitis (K50-K51)												
Asian or Pacific Islander 2/, both sexes												
2004.....	-	-	-	1	-	-	-	-	-	-	-	-
2003.....	1	-	1	1	1	-	1	1	-	-	-	-
2002.....	1	-	-	1	2	1	-	1	-	-	-	-
2001.....	1	-	1	-	-	1	-	1	1	-	-	-
2000.....	-	1	-	1	1	1	-	-	-	-	-	-
1999.....	-	-	-	-	-	1	1	1	-	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	1	1	-	-	1	1	-	-	-	-
2002.....	-	-	-	1	-	1	-	-	-	-	-	-
2001.....	-	-	1	-	-	-	-	1	1	-	-	-
2000.....	-	-	-	-	1	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	-	-	-	1	-	-	-	-	-	-	-	-
2003.....	1	-	-	-	1	-	-	-	-	-	-	-
2002.....	1	-	-	-	2	-	-	1	-	-	-	-
2001.....	1	-	-	-	-	1	-	-	-	-	-	-
2000.....	-	1	-	1	-	1	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	1	1	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Vascular disorders and obstruction of intestine without hernia (K55-K56)												
All races, both sexes												
2004.....	285	421	606	868	1,369	2,183	2,777	2,657	1,768	664	108	-
2003.....	305	455	579	946	1,492	2,215	2,832	2,685	1,823	634	113	-
2002.....	269	413	558	863	1,480	2,199	2,817	2,716	1,796	612	80	-
2001.....	290	380	596	964	1,566	2,207	2,684	2,661	1,649	585	103	-
2000.....	296	372	569	907	1,549	2,334	2,617	2,710	1,585	540	93	-
1999.....	256	384	568	964	1,519	2,196	2,564	2,561	1,605	564	96	1
All races, male												
2004.....	142	191	271	364	554	869	955	791	415	116	17	-
2003.....	150	235	258	400	621	832	1,002	784	429	103	22	-
2002.....	138	178	262	358	609	823	998	793	421	112	18	-
2001.....	140	161	270	402	685	831	925	767	382	115	16	-
2000.....	138	176	265	393	646	914	884	771	340	84	14	-
1999.....	116	181	248	384	649	839	877	708	336	81	16	-
All races, female												
2004.....	143	230	335	504	815	1,314	1,822	1,866	1,353	548	91	-
2003.....	155	220	321	546	871	1,383	1,830	1,901	1,394	531	91	-
2002.....	131	235	296	505	871	1,376	1,819	1,923	1,375	500	62	-
2001.....	150	219	326	562	881	1,376	1,759	1,894	1,267	470	87	-
2000.....	158	196	304	514	903	1,420	1,733	1,939	1,245	456	79	-
1999.....	140	203	320	580	870	1,357	1,687	1,853	1,269	483	80	1
White, both sexes												
2004.....	222	329	502	743	1,181	1,915	2,506	2,431	1,614	610	100	-
2003.....	219	355	463	799	1,295	1,950	2,574	2,479	1,701	567	94	-
2002.....	198	334	456	720	1,256	1,969	2,540	2,509	1,675	559	70	-
2001.....	205	293	473	801	1,368	1,929	2,461	2,465	1,542	536	88	-
2000.....	218	296	446	757	1,354	2,068	2,370	2,483	1,469	492	85	-
1999.....	195	320	484	803	1,297	1,929	2,345	2,329	1,498	529	89	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Vascular disorders and obstruction of intestine without hernia (K55-K56)

White, male

2004.....	113	149	217	304	462	748	844	710	377	104	15	-
2003.....	107	174	201	332	515	722	892	716	400	85	13	-
2002.....	99	142	202	302	512	737	887	718	392	101	15	-
2001.....	99	123	218	323	599	722	846	696	352	104	11	-
2000.....	102	133	198	318	563	798	786	695	318	72	11	-
1999.....	86	151	208	317	559	743	788	626	311	72	14	-

White, female

2004.....	109	180	285	439	719	1,167	1,662	1,721	1,237	506	85	-
2003.....	112	181	262	467	780	1,228	1,682	1,763	1,301	482	81	-
2002.....	99	192	254	418	744	1,232	1,653	1,791	1,283	458	55	-
2001.....	106	170	255	478	769	1,207	1,615	1,769	1,190	432	77	-
2000.....	116	163	248	439	791	1,270	1,584	1,788	1,151	420	74	-
1999.....	109	169	276	486	738	1,186	1,557	1,703	1,187	457	75	1

Black, both sexes

2004.....	56	84	91	102	159	223	229	185	137	47	7	-
2003.....	76	85	97	123	169	221	214	174	103	58	15	-
2002.....	64	69	83	125	189	188	242	175	105	47	9	-
2001.....	78	73	114	143	178	231	198	164	95	43	13	-
2000.....	75	73	114	129	163	229	209	204	108	40	6	-
1999.....	56	57	79	144	193	237	190	208	97	32	4	-

Black, male

2004.....	25	38	50	51	77	98	92	62	34	10	2	-
2003.....	41	52	49	58	91	90	92	56	21	14	6	-
2002.....	34	32	48	50	82	66	94	64	25	6	3	-
2001.....	37	33	47	74	80	92	71	65	27	10	4	-
2000.....	35	42	63	63	70	97	84	63	21	10	2	-
1999.....	27	27	39	59	74	84	78	67	19	8	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Vascular disorders and obstruction of intestine without hernia (K55-K56)												
Black, female												
2004.....	31	46	41	51	82	125	137	123	103	37	5	-
2003.....	35	33	48	65	78	131	122	118	82	44	9	-
2002.....	30	37	35	75	107	122	148	111	80	41	6	-
2001.....	41	40	67	69	98	139	127	99	68	33	9	-
2000.....	40	31	51	66	93	132	125	141	87	30	4	-
1999.....	29	30	40	85	119	153	112	141	78	24	4	-
American Indian 1/, both sexes												
2004.....	2	5	3	9	12	11	12	7	6	2	-	-
2003.....	7	7	9	7	11	12	13	4	5	2	-	-
2002.....	2	5	8	5	13	10	7	12	4	-	-	-
2001.....	5	6	5	10	8	15	9	11	3	1	1	-
2000.....	3	3	1	8	12	10	11	4	1	1	1	-
1999.....	3	5	3	6	10	7	8	5	-	1	1	-
American Indian 1/, male												
2004.....	-	1	-	2	4	5	3	-	1	-	-	-
2003.....	-	3	4	3	6	3	4	1	3	-	-	-
2002.....	1	3	4	1	4	4	2	4	2	-	-	-
2001.....	3	2	3	2	1	5	3	1	-	-	1	-
2000.....	1	1	-	6	3	4	5	2	-	-	1	-
1999.....	1	2	1	2	3	1	2	2	-	-	-	-
American Indian 1/, female												
2004.....	2	4	3	7	8	6	9	7	5	2	-	-
2003.....	7	4	5	4	5	9	9	3	2	2	-	-
2002.....	1	2	4	4	9	6	5	8	2	-	-	-
2001.....	2	4	2	8	7	10	6	10	3	1	-	-
2000.....	2	2	1	2	9	6	6	2	1	1	-	-
1999.....	2	3	2	4	7	6	6	3	-	1	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Vascular disorders and obstruction of intestine without hernia (K55-K56)												
Asian or Pacific Islander 2/, both sexes												
2004.....	5	3	10	14	17	34	30	34	11	5	1	-
2003.....	3	8	10	17	17	32	31	28	14	7	4	-
2002.....	5	5	11	13	22	32	28	20	12	6	1	-
2001.....	2	8	4	10	12	32	16	21	9	5	1	-
2000.....	-	-	8	13	20	27	27	19	7	7	1	-
1999.....	2	2	2	11	19	23	21	19	10	2	2	-
Asian or Pacific Islander 2/, male												
2004.....	4	3	4	7	11	18	16	19	3	2	-	-
2003.....	2	6	4	7	9	17	14	11	5	4	3	-
2002.....	4	1	8	5	11	16	15	7	2	5	-	-
2001.....	1	3	2	3	5	12	5	5	3	1	-	-
2000.....	-	-	4	6	10	15	9	11	1	2	-	-
1999.....	2	1	-	6	13	11	9	13	6	1	2	-
Asian or Pacific Islander 2/, female												
2004.....	1	-	6	7	6	16	14	15	8	3	1	-
2003.....	1	2	6	10	8	15	17	17	9	3	1	-
2002.....	1	4	3	8	11	16	13	13	10	1	1	-
2001.....	1	5	2	7	7	20	11	16	6	4	1	-
2000.....	-	-	4	7	10	12	18	8	6	5	1	-
1999.....	-	1	2	5	6	12	12	6	4	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diverticular disease of intestine (K57)												
All races, both sexes												
2004.....	60	90	138	183	327	484	721	668	455	163	24	-
2003.....	60	96	137	190	311	542	718	728	443	164	29	-
2002.....	56	79	106	179	306	543	712	662	464	123	21	-
2001.....	48	86	143	206	318	520	661	711	470	173	27	-
2000.....	53	58	116	213	330	481	658	683	472	136	18	-
1999.....	49	93	118	214	362	510	623	736	479	144	28	-
All races, male												
2004.....	36	44	67	75	125	180	226	171	88	21	4	-
2003.....	29	39	63	78	121	178	212	192	97	28	5	-
2002.....	40	41	48	76	121	199	210	169	84	21	2	-
2001.....	26	43	60	70	125	202	207	175	92	33	4	-
2000.....	27	36	49	92	134	158	199	162	93	19	-	-
1999.....	20	36	60	82	146	167	188	170	111	24	3	-
All races, female												
2004.....	24	46	71	108	202	304	495	497	367	142	20	-
2003.....	31	57	74	112	190	364	506	536	346	136	24	-
2002.....	16	38	58	103	185	344	502	493	380	102	19	-
2001.....	22	43	83	136	193	318	454	536	378	140	23	-
2000.....	26	22	67	121	196	323	459	521	379	117	18	-
1999.....	29	57	58	132	216	343	435	566	368	120	25	-
White, both sexes												
2004.....	49	84	118	154	296	448	668	624	434	147	19	-
2003.....	54	74	115	168	276	500	667	690	416	156	27	-
2002.....	47	68	90	157	276	492	655	623	427	113	20	-
2001.....	39	75	121	168	284	478	630	656	449	163	23	-
2000.....	45	47	91	185	294	447	614	640	456	127	15	-
1999.....	44	74	99	185	327	476	582	701	445	134	24	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diverticular disease of intestine (K57)												
White, male												
2004.....	33	41	57	62	115	166	204	159	83	17	4	-
2003.....	28	34	55	70	107	169	195	181	94	26	5	-
2002.....	35	34	43	66	114	173	188	158	80	19	1	-
2001.....	22	40	51	58	114	183	195	164	87	32	3	-
2000.....	25	32	37	81	118	149	189	153	91	19	-	-
1999.....	19	29	50	76	132	155	176	162	103	23	3	-
White, female												
2004.....	16	43	61	92	181	282	464	465	351	130	15	-
2003.....	26	40	60	98	169	331	472	509	322	130	22	-
2002.....	12	34	47	91	162	319	467	465	347	94	19	-
2001.....	17	35	70	110	170	295	435	492	362	131	20	-
2000.....	20	15	54	104	176	298	425	487	365	108	15	-
1999.....	25	45	49	109	195	321	406	539	342	111	21	-
Black, both sexes												
2004.....	11	3	15	24	28	27	45	40	20	12	5	-
2003.....	6	21	20	16	28	35	43	30	25	6	2	-
2002.....	8	8	15	20	23	43	51	32	31	9	1	-
2001.....	8	10	16	35	31	38	26	51	16	10	2	-
2000.....	6	9	23	26	35	28	38	36	16	9	3	-
1999.....	4	18	17	27	30	29	37	30	30	10	4	-
Black, male												
2004.....	3	1	9	11	8	10	18	11	5	2	-	-
2003.....	1	5	8	4	11	7	15	10	2	1	-	-
2002.....	5	5	5	8	5	21	18	8	2	2	1	-
2001.....	3	3	7	10	10	16	8	9	2	1	1	-
2000.....	1	4	11	9	16	7	9	5	2	-	-	-
1999.....	1	6	9	5	12	11	10	6	5	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diverticular disease of intestine (K57)												
Black, female												
2004.....	8	2	6	13	20	17	27	29	15	10	5	-
2003.....	5	16	12	12	17	28	28	20	23	5	2	-
2002.....	3	3	10	12	18	22	33	24	29	7	-	-
2001.....	5	7	9	25	21	22	18	42	14	9	1	-
2000.....	5	5	12	17	19	21	29	31	14	9	3	-
1999.....	3	12	8	22	18	18	27	24	25	9	4	-
American Indian 1/, both sexes												
2004.....	-	1	3	2	1	-	2	-	-	-	-	-
2003.....	-	-	-	3	1	3	3	1	-	-	-	-
2002.....	1	2	-	1	1	-	1	1	1	-	-	-
2001.....	-	-	3	1	-	-	-	3	-	-	-	-
2000.....	1	-	2	1	1	-	1	2	-	-	-	-
1999.....	1	-	2	1	1	3	-	1	-	-	-	-
American Indian 1/, male												
2004.....	-	1	-	1	1	-	1	-	-	-	-	-
2003.....	-	-	-	3	1	-	1	-	-	-	-	-
2002.....	-	1	-	1	-	-	-	-	-	-	-	-
2001.....	-	-	-	1	-	-	-	1	-	-	-	-
2000.....	-	-	1	1	-	-	-	-	-	-	-	-
1999.....	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, female												
2004.....	-	-	3	1	-	-	1	-	-	-	-	-
2003.....	-	-	-	-	-	3	2	1	-	-	-	-
2002.....	1	1	-	-	1	-	1	1	1	-	-	-
2001.....	-	-	3	-	-	-	-	2	-	-	-	-
2000.....	1	-	1	-	1	-	1	2	-	-	-	-
1999.....	1	-	1	1	1	3	-	1	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diverticular disease of intestine (K57)												
Asian or Pacific Islander 2/, both sexes												
2004.....	-	2	2	3	2	9	6	4	1	4	-	-
2003.....	-	1	2	3	6	4	5	7	2	2	-	-
2002.....	-	1	1	1	6	8	5	6	5	1	-	-
2001.....	1	1	3	2	3	4	5	1	5	-	2	-
2000.....	1	2	-	1	-	6	5	5	-	-	-	-
1999.....	-	1	-	1	4	2	4	4	4	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	-	1	1	1	1	4	3	1	-	2	-	-
2003.....	-	-	-	1	2	2	1	1	1	1	-	-
2002.....	-	1	-	1	2	5	4	3	2	-	-	-
2001.....	1	-	2	1	1	3	4	1	3	-	-	-
2000.....	1	-	-	1	-	2	1	4	-	-	-	-
1999.....	-	1	-	1	2	1	2	2	3	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	-	1	1	2	1	5	3	3	1	2	-	-
2003.....	-	1	2	2	4	2	4	6	1	1	-	-
2002.....	-	-	1	-	4	3	1	3	3	1	-	-
2001.....	-	1	1	1	2	1	1	-	2	-	2	-
2000.....	-	2	-	-	-	4	4	1	-	-	-	-
1999.....	-	-	-	-	2	1	2	2	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of intestines and peritoneum (K52,K58-K63)												
All races, both sexes												
2004.....	134	173	239	316	489	695	961	857	530	199	32	-
2003.....	141	185	249	326	470	731	892	772	534	197	41	-
2002.....	135	183	248	335	455	734	863	799	499	169	41	-
2001.....	136	152	212	330	490	749	864	821	492	169	19	3
2000.....	110	165	214	286	470	701	835	726	493	161	32	-
1999.....	114	157	207	320	498	713	768	714	422	160	32	-
All races, male												
2004.....	68	80	92	136	207	304	350	237	119	38	4	-
2003.....	72	92	124	137	211	313	322	231	120	33	8	-
2002.....	64	87	111	153	211	324	277	250	117	32	9	-
2001.....	63	72	98	139	209	304	301	232	121	31	5	3
2000.....	41	86	110	119	217	278	277	196	120	33	3	-
1999.....	60	78	89	136	214	295	280	205	103	32	5	-
All races, female												
2004.....	66	93	147	180	282	391	611	620	411	161	28	-
2003.....	69	93	125	189	259	418	570	541	414	164	33	-
2002.....	71	96	137	182	244	410	586	549	382	137	32	-
2001.....	73	80	114	191	281	445	563	589	371	138	14	-
2000.....	69	79	104	167	253	423	558	530	373	128	29	-
1999.....	54	79	118	184	284	418	488	509	319	128	27	-
White, both sexes												
2004.....	111	145	197	266	432	628	890	807	490	183	27	-
2003.....	105	154	203	269	407	660	828	718	495	186	39	-
2002.....	114	154	204	299	389	670	799	755	466	159	37	-
2001.....	110	114	173	289	438	666	780	769	459	155	18	1
2000.....	85	137	166	248	420	642	771	666	450	150	29	-
1999.....	92	134	175	277	444	647	704	669	396	149	30	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of intestines and peritoneum (K52,K58-K63)												
White, male												
2004.....	54	64	77	106	183	271	322	219	108	36	3	-
2003.....	54	72	104	117	181	282	296	214	107	31	8	-
2002.....	56	73	91	131	176	296	250	240	103	30	8	-
2001.....	51	52	78	116	178	269	275	218	115	29	5	1
2000.....	31	70	81	105	198	250	254	172	106	31	2	-
1999.....	49	66	75	119	196	262	258	182	93	26	4	-
White, female												
2004.....	57	81	120	160	249	357	568	588	382	147	24	-
2003.....	51	82	99	152	226	378	532	504	388	155	31	-
2002.....	58	81	113	168	213	374	549	515	363	129	29	-
2001.....	59	62	95	173	260	397	505	551	344	126	13	-
2000.....	54	67	85	143	222	392	517	494	344	119	27	-
1999.....	43	68	100	158	248	385	446	487	303	123	26	-
Black, both sexes												
2004.....	16	25	38	42	50	54	60	39	29	13	4	-
2003.....	32	28	42	52	50	61	58	45	31	10	2	-
2002.....	21	25	38	31	58	52	53	39	27	8	4	-
2001.....	22	32	30	38	45	75	73	46	27	14	1	2
2000.....	21	24	38	31	46	45	53	44	38	10	3	-
1999.....	21	20	25	39	47	50	56	38	26	9	1	-
Black, male												
2004.....	11	13	14	25	23	26	22	14	8	2	1	-
2003.....	15	18	19	20	24	26	22	12	10	2	-	-
2002.....	8	13	16	20	32	23	21	10	12	-	1	-
2001.....	11	19	15	23	26	34	21	12	4	2	-	2
2000.....	8	14	20	10	17	23	21	15	9	1	1	-
1999.....	11	10	10	17	15	22	20	20	10	4	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of intestines and peritoneum (K52,K58-K63)												
Black, female												
2004.....	5	12	24	17	27	28	38	25	21	11	3	-
2003.....	17	10	23	32	26	35	36	33	21	8	2	-
2002.....	13	12	22	11	26	29	32	29	15	8	3	-
2001.....	11	13	15	15	19	41	52	34	23	12	1	-
2000.....	13	10	18	21	29	22	32	29	29	9	2	-
1999.....	10	10	15	22	32	28	36	18	16	5	1	-
American Indian 1/, both sexes												
2004.....	4	3	2	6	2	2	4	4	2	1	-	-
2003.....	1	2	1	2	3	4	3	2	5	-	-	-
2002.....	-	2	4	2	3	1	2	2	3	1	-	-
2001.....	2	4	7	1	3	3	2	3	1	-	-	-
2000.....	2	1	1	4	-	5	1	5	1	-	-	-
1999.....	-	2	3	2	4	5	2	1	-	1	-	-
American Indian 1/, male												
2004.....	1	3	1	4	1	1	-	1	1	-	-	-
2003.....	1	1	1	-	2	2	3	1	-	-	-	-
2002.....	-	-	3	-	-	1	-	-	1	1	-	-
2001.....	1	1	4	-	3	-	1	1	1	-	-	-
2000.....	2	1	1	2	-	2	-	1	1	-	-	-
1999.....	-	1	2	-	1	2	-	-	-	1	-	-
American Indian 1/, female												
2004.....	3	-	1	2	1	1	4	3	1	1	-	-
2003.....	-	1	-	2	1	2	-	1	5	-	-	-
2002.....	-	2	1	2	3	-	2	2	2	-	-	-
2001.....	1	3	3	1	-	3	1	2	-	-	-	-
2000.....	-	-	-	2	-	3	1	4	-	-	-	-
1999.....	-	1	1	2	3	3	2	1	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of intestines and peritoneum (K52,K58-K63)												
Asian or Pacific Islander 2/, both sexes												
2004.....	3	-	2	2	5	11	7	7	9	2	1	-
2003.....	3	1	3	3	10	6	3	7	3	1	-	-
2002.....	-	2	2	3	5	11	9	3	3	1	-	-
2001.....	2	2	2	2	4	5	9	3	5	-	-	-
2000.....	2	3	9	3	4	9	10	11	4	1	-	-
1999.....	1	1	4	2	3	11	6	6	-	1	1	-
Asian or Pacific Islander 2/, male												
2004.....	2	-	-	1	-	6	6	3	2	-	-	-
2003.....	2	1	-	-	4	3	1	4	3	-	-	-
2002.....	-	1	1	2	3	4	6	-	1	1	-	-
2001.....	-	-	1	-	2	1	4	1	1	-	-	-
2000.....	-	1	8	2	2	3	2	8	4	1	-	-
1999.....	-	1	2	-	2	9	2	3	-	1	1	-
Asian or Pacific Islander 2/, female												
2004.....	1	-	2	1	5	5	1	4	7	2	1	-
2003.....	1	-	3	3	6	3	2	3	-	1	-	-
2002.....	-	1	1	1	2	7	3	3	2	-	-	-
2001.....	2	2	1	2	2	4	5	2	4	-	-	-
2000.....	2	2	1	1	2	6	8	3	-	-	-	-
1999.....	1	-	2	2	1	2	4	3	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of peritoneum (K65-K66)												
All races, both sexes												
2004.....	85	107	108	140	180	250	286	242	121	46	9	-
2003.....	109	90	128	161	190	257	274	242	126	60	5	-
2002.....	107	105	117	123	197	278	328	267	150	53	2	-
2001.....	95	108	103	135	198	265	292	276	128	50	3	-
2000.....	91	122	129	138	235	287	260	259	136	51	5	-
1999.....	93	104	113	152	247	272	270	205	138	51	10	-
All races, male												
2004.....	53	52	55	78	89	104	126	73	33	8	2	-
2003.....	58	50	68	78	87	115	118	74	38	14	-	-
2002.....	63	57	59	58	76	115	102	86	34	5	1	-
2001.....	50	64	54	77	86	110	119	86	33	12	-	-
2000.....	43	62	71	66	104	130	95	88	37	9	1	-
1999.....	55	53	64	74	122	122	113	66	40	12	2	-
All races, female												
2004.....	32	55	53	62	91	146	160	169	88	38	7	-
2003.....	51	40	60	83	103	142	156	168	88	46	5	-
2002.....	44	48	58	65	121	163	226	181	116	48	1	-
2001.....	45	44	49	58	112	155	173	190	95	38	3	-
2000.....	48	60	58	72	131	157	165	171	99	42	4	-
1999.....	38	51	49	78	125	150	157	139	98	39	8	-
White, both sexes												
2004.....	66	83	89	118	145	228	251	224	115	39	8	-
2003.....	77	65	92	134	161	234	248	219	113	52	4	-
2002.....	80	89	87	101	166	250	293	247	140	48	1	-
2001.....	70	77	71	110	167	229	259	251	116	45	3	-
2000.....	67	92	102	108	203	252	231	235	123	44	4	-
1999.....	65	73	89	118	209	233	248	179	124	49	8	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of peritoneum (K65-K66)												
White, male												
2004.....	40	40	46	65	73	95	112	68	31	7	1	-
2003.....	41	37	51	62	78	106	106	68	36	11	-	-
2002.....	50	49	43	51	66	99	91	80	31	5	1	-
2001.....	40	43	40	65	73	96	109	77	27	10	-	-
2000.....	32	49	56	50	88	115	83	76	34	9	1	-
1999.....	39	37	48	58	106	106	104	59	37	12	2	-
White, female												
2004.....	26	43	43	53	72	133	139	156	84	32	7	-
2003.....	36	28	41	72	83	128	142	151	77	41	4	-
2002.....	30	40	44	50	100	151	202	167	109	43	-	-
2001.....	30	34	31	45	94	133	150	174	89	35	3	-
2000.....	35	43	46	58	115	137	148	159	89	35	3	-
1999.....	26	36	41	60	103	127	144	120	87	37	6	-
Black, both sexes												
2004.....	18	19	17	20	27	19	32	15	5	7	1	-
2003.....	27	23	30	19	27	21	22	21	13	6	1	-
2002.....	24	14	25	20	27	24	28	19	10	5	1	-
2001.....	23	25	26	24	24	28	29	23	11	4	-	-
2000.....	23	29	23	26	25	30	28	20	12	5	-	-
1999.....	27	30	19	28	35	33	21	21	13	2	2	-
Black, male												
2004.....	12	9	8	11	11	6	12	4	2	1	1	-
2003.....	14	11	13	11	9	9	10	6	2	2	-	-
2002.....	13	7	12	7	8	13	8	5	3	-	-	-
2001.....	10	18	9	12	11	10	9	7	5	1	-	-
2000.....	11	13	13	15	11	13	12	10	2	-	-	-
1999.....	16	15	12	13	15	14	9	6	3	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of peritoneum (K65-K66)												
Black, female												
2004.....	6	10	9	9	16	13	20	11	3	6	-	-
2003.....	13	12	17	8	18	12	12	15	11	4	1	-
2002.....	11	7	13	13	19	11	20	14	7	5	1	-
2001.....	13	7	17	12	13	18	20	16	6	3	-	-
2000.....	12	16	10	11	14	17	16	10	10	5	-	-
1999.....	11	15	7	15	20	19	12	15	10	2	2	-
American Indian 1/, both sexes												
2004.....	-	2	2	-	4	-	1	-	-	-	-	-
2003.....	3	-	3	1	-	-	1	1	-	1	-	-
2002.....	2	-	2	-	1	1	1	-	-	-	-	-
2001.....	1	2	1	-	-	-	1	-	-	-	-	-
2000.....	-	-	2	3	1	2	-	1	-	2	-	-
1999.....	1	-	1	-	-	-	1	-	1	-	-	-
American Indian 1/, male												
2004.....	-	1	1	-	2	-	1	-	-	-	-	-
2003.....	2	-	1	-	-	-	1	-	-	-	-	-
2002.....	-	-	1	-	1	1	-	-	-	-	-	-
2001.....	-	1	1	-	-	-	1	-	-	-	-	-
2000.....	-	-	-	-	1	1	-	-	-	-	-	-
1999.....	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, female												
2004.....	-	1	1	-	2	-	-	-	-	-	-	-
2003.....	1	-	2	1	-	-	-	1	-	1	-	-
2002.....	2	-	1	-	-	-	1	-	-	-	-	-
2001.....	1	1	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	2	3	-	1	-	1	-	2	-	-
1999.....	1	-	-	-	-	-	1	-	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of peritoneum (K65-K66)												
Asian or Pacific Islander 2/, both sexes												
2004.....	1	3	-	2	4	3	2	3	1	-	-	-
2003.....	2	2	3	7	2	2	3	1	-	1	-	-
2002.....	1	2	3	2	3	3	6	1	-	-	-	-
2001.....	1	4	5	1	7	8	3	2	1	1	-	-
2000.....	1	1	2	1	6	3	1	3	1	-	1	-
1999.....	-	1	4	6	3	6	-	5	-	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	1	2	-	2	3	3	1	1	-	-	-	-
2003.....	1	2	3	5	-	-	1	-	-	1	-	-
2002.....	-	1	3	-	1	2	3	1	-	-	-	-
2001.....	-	2	4	-	2	4	-	2	1	1	-	-
2000.....	-	-	2	1	4	1	-	2	1	-	-	-
1999.....	-	1	3	3	1	2	-	1	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	-	1	-	-	1	-	1	2	1	-	-	-
2003.....	1	-	-	2	2	2	2	1	-	-	-	-
2002.....	1	1	-	2	2	1	3	-	-	-	-	-
2001.....	1	2	1	1	5	4	3	-	-	-	-	-
2000.....	1	1	-	-	2	2	1	1	-	-	1	-
1999.....	-	-	1	3	2	4	-	4	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of liver (K70-K76)												
All races, both sexes												
2004.....	5,280	4,650	3,912	3,437	3,367	3,067	2,126	1,117	389	79	11	1
2003.....	5,168	4,494	3,886	3,633	3,468	3,095	2,179	1,121	376	76	11	4
2002.....	4,828	4,117	3,623	3,449	3,507	3,112	2,225	1,130	417	83	10	4
2001.....	4,601	3,753	3,528	3,488	3,485	3,132	2,070	1,105	368	104	11	8
2000.....	4,106	3,620	3,495	3,461	3,518	3,074	2,014	1,101	366	78	10	4
1999.....	3,897	3,476	3,516	3,630	3,635	3,112	2,044	1,050	388	92	16	3
All races, male												
2004.....	3,857	3,282	2,555	2,100	1,907	1,591	992	437	145	28	3	-
2003.....	3,796	3,127	2,632	2,245	1,957	1,549	1,009	471	143	24	3	4
2002.....	3,550	2,885	2,383	2,120	1,993	1,567	1,062	493	132	29	3	4
2001.....	3,351	2,629	2,346	2,180	1,983	1,565	972	466	136	36	4	8
2000.....	2,997	2,484	2,320	2,129	1,994	1,607	883	473	139	26	2	4
1999.....	2,891	2,487	2,311	2,305	2,047	1,540	965	449	143	27	5	3
All races, female												
2004.....	1,423	1,368	1,357	1,337	1,460	1,476	1,134	680	244	51	8	1
2003.....	1,372	1,367	1,254	1,388	1,511	1,546	1,170	650	233	52	8	-
2002.....	1,278	1,232	1,240	1,329	1,514	1,545	1,163	637	285	54	7	-
2001.....	1,250	1,124	1,182	1,308	1,502	1,567	1,098	639	232	68	7	-
2000.....	1,109	1,136	1,175	1,332	1,524	1,467	1,131	628	227	52	8	-
1999.....	1,006	989	1,205	1,325	1,588	1,572	1,079	601	245	65	11	-
White, both sexes												
2004.....	4,409	3,886	3,353	3,017	3,034	2,838	1,964	1,025	353	76	10	1
2003.....	4,245	3,794	3,335	3,194	3,140	2,876	2,033	1,045	337	68	8	4
2002.....	3,983	3,482	3,120	3,049	3,197	2,881	2,055	1,048	385	80	7	4
2001.....	3,756	3,112	3,058	3,081	3,144	2,881	1,925	1,024	347	93	11	6
2000.....	3,320	3,027	2,989	3,044	3,183	2,852	1,865	1,022	336	73	8	3
1999.....	3,228	2,873	2,971	3,189	3,278	2,860	1,916	965	356	84	14	2

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of liver (K70-K76)												
White, male												
2004.....	3,274	2,757	2,207	1,847	1,726	1,476	925	415	135	28	2	-
2003.....	3,145	2,641	2,275	1,979	1,782	1,448	940	441	124	22	2	4
2002.....	2,968	2,461	2,063	1,883	1,828	1,461	982	458	122	29	1	4
2001.....	2,770	2,189	2,033	1,933	1,803	1,438	908	437	126	31	4	6
2000.....	2,448	2,107	2,011	1,897	1,813	1,493	832	437	129	23	2	3
1999.....	2,423	2,081	1,990	2,038	1,861	1,422	917	420	133	24	4	2
White, female												
2004.....	1,135	1,129	1,146	1,170	1,308	1,362	1,039	610	218	48	8	1
2003.....	1,100	1,153	1,060	1,215	1,358	1,428	1,093	604	213	46	6	-
2002.....	1,015	1,021	1,057	1,166	1,369	1,420	1,073	590	263	51	6	-
2001.....	986	923	1,025	1,148	1,341	1,443	1,017	587	221	62	7	-
2000.....	872	920	978	1,147	1,370	1,359	1,033	585	207	50	6	-
1999.....	805	792	981	1,151	1,417	1,438	999	545	223	60	10	-
Black, both sexes												
2004.....	704	601	427	319	242	141	110	59	24	2	1	-
2003.....	764	563	424	350	245	144	93	56	27	7	3	-
2002.....	702	506	390	307	227	175	110	62	26	2	3	-
2001.....	691	533	368	333	253	190	107	47	17	7	-	2
2000.....	650	465	394	332	267	164	100	53	26	3	1	1
1999.....	547	484	446	345	270	193	97	65	25	6	2	1
Black, male												
2004.....	474	428	272	196	144	73	47	13	6	-	1	-
2003.....	550	399	283	218	136	68	47	24	13	1	1	-
2002.....	483	346	252	189	125	80	51	25	7	-	2	-
2001.....	493	363	249	210	143	95	47	15	7	2	-	2
2000.....	460	308	244	196	151	87	35	22	8	1	-	1
1999.....	387	329	266	215	149	89	38	24	6	1	1	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of liver (K70-K76)												
Black, female												
2004.....	230	173	155	123	98	68	63	46	18	2	-	-
2003.....	214	164	141	132	109	76	46	32	14	6	2	-
2002.....	219	160	138	118	102	95	59	37	19	2	1	-
2001.....	198	170	119	123	110	95	60	32	10	5	-	-
2000.....	190	157	150	136	116	77	65	31	18	2	1	-
1999.....	160	155	180	130	121	104	59	41	19	5	1	-
American Indian 1/, both sexes												
2004.....	118	106	72	49	34	33	10	8	1	-	-	-
2003.....	115	90	81	44	23	23	11	4	7	-	-	-
2002.....	95	75	66	47	35	15	16	3	-	-	-	-
2001.....	106	65	61	35	35	19	10	8	1	2	-	-
2000.....	101	80	59	45	24	22	12	3	-	-	1	-
1999.....	86	81	59	44	32	15	5	7	-	1	-	-
American Indian 1/, male												
2004.....	66	61	38	23	12	19	5	2	-	-	-	-
2003.....	72	52	40	24	9	8	3	-	5	-	-	-
2002.....	67	43	39	26	18	8	8	1	-	-	-	-
2001.....	59	42	42	21	14	10	3	1	1	1	-	-
2000.....	66	42	33	16	7	12	3	2	-	-	-	-
1999.....	55	49	29	21	16	7	3	2	-	1	-	-
American Indian 1/, female												
2004.....	52	45	34	26	22	14	5	6	1	-	-	-
2003.....	43	38	41	20	14	15	8	4	2	-	-	-
2002.....	28	32	27	21	17	7	8	2	-	-	-	-
2001.....	47	23	19	14	21	9	7	7	-	1	-	-
2000.....	35	38	26	29	17	10	9	1	-	-	1	-
1999.....	31	32	30	23	16	8	2	5	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of liver (K70-K76)												
Asian or Pacific Islander 2/, both sexes												
2004.....	49	57	60	52	57	55	42	25	11	1	-	-
2003.....	44	47	46	45	60	52	42	16	5	1	-	-
2002.....	48	54	47	46	48	41	44	17	6	1	-	-
2001.....	48	43	41	39	53	42	28	26	3	2	-	-
2000.....	35	48	53	40	44	36	37	23	4	2	-	-
1999.....	36	38	40	52	55	44	26	13	7	1	-	-
Asian or Pacific Islander 2/, male												
2004.....	43	36	38	34	25	23	15	7	4	-	-	-
2003.....	29	35	34	24	30	25	19	6	1	1	-	-
2002.....	32	35	29	22	22	18	21	9	3	-	-	-
2001.....	29	35	22	16	23	22	14	13	2	2	-	-
2000.....	23	27	32	20	23	15	13	12	2	2	-	-
1999.....	26	28	26	31	21	22	7	3	4	1	-	-
Asian or Pacific Islander 2/, female												
2004.....	6	21	22	18	32	32	27	18	7	1	-	-
2003.....	15	12	12	21	30	27	23	10	4	-	-	-
2002.....	16	19	18	24	26	23	23	8	3	1	-	-
2001.....	19	8	19	23	30	20	14	13	1	-	-	-
2000.....	12	21	21	20	21	21	24	11	2	-	-	-
1999.....	10	10	14	21	34	22	19	10	3	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Alcoholic liver disease (K70)												
All races, both sexes												
2004.....	2,369	1,940	1,453	1,078	763	481	228	86	15	1	-	1
2003.....	2,238	1,837	1,410	1,075	713	491	238	67	19	-	-	2
2002.....	2,139	1,712	1,371	1,023	826	495	235	79	16	3	-	2
2001.....	2,082	1,582	1,331	1,062	786	515	202	86	12	6	-	5
2000.....	1,888	1,575	1,394	1,085	806	533	218	76	17	2	-	2
1999.....	1,837	1,511	1,348	1,104	831	533	256	94	21	3	-	2
All races, male												
2004.....	1,772	1,453	1,096	824	568	357	170	56	12	1	-	-
2003.....	1,704	1,386	1,073	800	543	349	169	40	12	-	-	2
2002.....	1,624	1,277	1,060	771	596	334	161	54	10	1	-	2
2001.....	1,536	1,235	1,007	789	590	370	135	60	8	4	-	5
2000.....	1,458	1,172	1,044	821	612	419	142	52	12	1	-	2
1999.....	1,426	1,179	1,015	867	630	364	174	59	13	1	-	2
All races, female												
2004.....	597	487	357	254	195	124	58	30	3	-	-	1
2003.....	534	451	337	275	170	142	69	27	7	-	-	-
2002.....	515	435	311	252	230	161	74	25	6	2	-	-
2001.....	546	347	324	273	196	145	67	26	4	2	-	-
2000.....	430	403	350	264	194	114	76	24	5	1	-	-
1999.....	411	332	333	237	201	169	82	35	8	2	-	-
White, both sexes												
2004.....	2,048	1,654	1,260	947	691	448	206	81	13	1	-	1
2003.....	1,876	1,590	1,207	941	645	462	220	62	17	-	-	2
2002.....	1,808	1,466	1,190	903	749	450	215	71	14	3	-	2
2001.....	1,709	1,332	1,166	918	711	472	188	74	10	6	-	5
2000.....	1,576	1,329	1,204	961	728	487	194	70	16	2	-	1
1999.....	1,547	1,261	1,144	958	740	475	233	81	18	3	-	2

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Alcoholic liver disease (K70)												
White, male												
2004.....	1,561	1,245	958	726	512	333	154	53	11	1	-	-
2003.....	1,444	1,209	934	704	494	329	153	37	11	-	-	2
2002.....	1,392	1,120	927	684	542	306	149	48	8	1	-	2
2001.....	1,287	1,045	886	683	533	343	126	53	6	4	-	5
2000.....	1,237	1,013	917	735	553	387	130	48	11	1	-	1
1999.....	1,217	993	883	753	564	325	161	50	12	1	-	2
White, female												
2004.....	487	409	302	221	179	115	52	28	2	-	-	1
2003.....	432	381	273	237	151	133	67	25	6	-	-	-
2002.....	416	346	263	219	207	144	66	23	6	2	-	-
2001.....	422	287	280	235	178	129	62	21	4	2	-	-
2000.....	339	316	287	226	175	100	64	22	5	1	-	-
1999.....	330	268	261	205	176	150	72	31	6	2	-	-
Black, both sexes												
2004.....	241	216	146	105	50	16	15	1	2	-	-	-
2003.....	274	192	153	107	52	18	15	4	1	-	-	-
2002.....	255	186	138	90	55	36	15	7	1	-	-	-
2001.....	295	200	128	120	58	33	11	8	1	-	-	-
2000.....	249	187	138	107	63	35	15	4	1	-	-	1
1999.....	227	194	167	117	77	45	19	11	3	-	-	-
Black, male												
2004.....	159	158	104	80	42	9	10	1	1	-	-	-
2003.....	204	139	108	76	38	11	13	3	1	-	-	-
2002.....	180	117	103	66	39	23	9	5	1	-	-	-
2001.....	206	150	92	90	45	19	7	4	1	-	-	-
2000.....	177	124	87	74	50	24	9	3	1	-	-	1
1999.....	164	147	110	93	55	30	11	7	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Alcoholic liver disease (K70)												
Black, female												
2004.....	82	58	42	25	8	7	5	-	1	-	-	-
2003.....	70	53	45	31	14	7	2	1	-	-	-	-
2002.....	75	69	35	24	16	13	6	2	-	-	-	-
2001.....	89	50	36	30	13	14	4	4	-	-	-	-
2000.....	72	63	51	33	13	11	6	1	-	-	-	-
1999.....	63	47	57	24	22	15	8	4	2	-	-	-
American Indian 1/, both sexes												
2004.....	65	54	33	18	9	11	2	2	-	-	-	-
2003.....	75	43	38	18	9	7	-	1	-	-	-	-
2002.....	55	43	31	20	14	5	1	-	-	-	-	-
2001.....	61	34	27	18	10	3	1	1	1	-	-	-
2000.....	50	44	35	15	8	8	2	-	-	-	-	-
1999.....	53	43	32	19	8	7	1	1	-	-	-	-
American Indian 1/, male												
2004.....	39	37	23	11	3	9	2	1	-	-	-	-
2003.....	45	29	22	13	4	5	-	-	-	-	-	-
2002.....	38	26	19	13	10	3	-	-	-	-	-	-
2001.....	32	26	21	11	6	2	-	1	1	-	-	-
2000.....	33	25	26	11	3	5	1	-	-	-	-	-
1999.....	39	27	18	13	6	4	1	1	-	-	-	-
American Indian 1/, female												
2004.....	26	17	10	7	6	2	-	1	-	-	-	-
2003.....	30	14	16	5	5	2	-	1	-	-	-	-
2002.....	17	17	12	7	4	2	1	-	-	-	-	-
2001.....	29	8	6	7	4	1	1	-	-	-	-	-
2000.....	17	19	9	4	5	3	1	-	-	-	-	-
1999.....	14	16	14	6	2	3	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Alcoholic liver disease (K70)												
Asian or Pacific Islander 2/, both sexes												
2004.....	15	16	14	8	13	6	5	2	-	-	-	-
2003.....	13	12	12	9	7	4	3	-	1	-	-	-
2002.....	21	17	12	10	8	4	4	1	1	-	-	-
2001.....	17	16	10	6	7	7	2	3	-	-	-	-
2000.....	13	15	17	2	7	3	7	2	-	-	-	-
1999.....	10	13	5	10	6	6	3	1	-	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	13	13	11	7	11	6	4	1	-	-	-	-
2003.....	11	9	9	7	7	4	3	-	-	-	-	-
2002.....	14	14	11	8	5	2	3	1	1	-	-	-
2001.....	11	14	8	5	6	6	2	2	-	-	-	-
2000.....	11	10	14	1	6	3	2	1	-	-	-	-
1999.....	6	12	4	8	5	5	1	1	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	2	3	3	1	2	-	1	1	-	-	-	-
2003.....	2	3	3	2	-	-	-	-	1	-	-	-
2002.....	7	3	1	2	3	2	1	-	-	-	-	-
2001.....	6	2	2	1	1	1	-	1	-	-	-	-
2000.....	2	5	3	1	1	-	5	1	-	-	-	-
1999.....	4	1	1	2	1	1	2	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic hepatitis, not elsewhere classified (K73)												
All races, both sexes												
2004.....	19	5	10	9	13	12	11	9	-	-	-	-
2003.....	21	13	12	21	12	14	11	7	3	-	-	-
2002.....	35	35	48	65	70	77	51	28	8	-	-	-
2001.....	37	43	36	54	66	62	42	23	7	3	-	-
2000.....	32	36	36	68	53	64	39	26	8	1	-	-
1999.....	35	37	46	40	64	58	46	21	4	1	-	-
All races, male												
2004.....	11	4	6	4	9	3	3	1	-	-	-	-
2003.....	14	7	6	10	6	5	3	3	-	-	-	-
2002.....	16	5	10	13	19	19	9	7	1	-	-	-
2001.....	16	11	5	10	18	12	10	7	3	-	-	-
2000.....	13	9	4	12	7	10	8	6	2	-	-	-
1999.....	18	10	11	7	19	15	10	5	-	-	-	-
All races, female												
2004.....	8	1	4	5	4	9	8	8	-	-	-	-
2003.....	7	6	6	11	6	9	8	4	3	-	-	-
2002.....	19	30	38	52	51	58	42	21	7	-	-	-
2001.....	21	32	31	44	48	50	32	16	4	3	-	-
2000.....	19	27	32	56	46	54	31	20	6	1	-	-
1999.....	17	27	35	33	45	43	36	16	4	1	-	-
White, both sexes												
2004.....	16	4	9	9	12	12	9	9	-	-	-	-
2003.....	17	12	12	18	9	13	11	6	2	-	-	-
2002.....	26	31	42	56	65	76	45	26	8	-	-	-
2001.....	34	40	33	49	55	54	40	22	7	3	-	-
2000.....	24	31	34	54	49	61	36	24	8	1	-	-
1999.....	29	29	36	33	54	51	42	19	4	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic hepatitis, not elsewhere classified (K73)												
White, male												
2004.....	8	3	6	4	9	3	2	1	-	-	-	-
2003.....	14	6	6	9	4	5	3	3	-	-	-	-
2002.....	12	5	10	12	18	19	8	7	1	-	-	-
2001.....	15	10	4	8	18	12	9	7	3	-	-	-
2000.....	10	8	4	9	6	10	8	6	2	-	-	-
1999.....	16	10	11	7	13	12	8	5	-	-	-	-
White, female												
2004.....	8	1	3	5	3	9	7	8	-	-	-	-
2003.....	3	6	6	9	5	8	8	3	2	-	-	-
2002.....	14	26	32	44	47	57	37	19	7	-	-	-
2001.....	19	30	29	41	37	42	31	15	4	3	-	-
2000.....	14	23	30	45	43	51	28	18	6	1	-	-
1999.....	13	19	25	26	41	39	34	14	4	1	-	-
Black, both sexes												
2004.....	2	1	-	-	1	-	-	-	-	-	-	-
2003.....	4	1	-	3	2	-	-	-	-	-	-	-
2002.....	8	2	5	7	2	-	4	2	-	-	-	-
2001.....	3	2	2	4	5	7	1	1	-	-	-	-
2000.....	6	3	2	8	3	1	2	2	-	-	-	-
1999.....	5	4	7	6	6	3	3	2	-	-	-	-
Black, male												
2004.....	2	1	-	-	-	-	-	-	-	-	-	-
2003.....	-	1	-	1	1	-	-	-	-	-	-	-
2002.....	3	-	-	1	-	-	1	-	-	-	-	-
2001.....	1	-	1	2	-	-	-	-	-	-	-	-
2000.....	3	1	-	1	-	-	-	-	-	-	-	-
1999.....	2	-	-	-	2	-	2	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic hepatitis, not elsewhere classified (K73)												
Asian or Pacific Islander 2/, both sexes												
2004.....	1	-	-	-	-	-	1	-	-	-	-	-
2003.....	-	-	-	-	-	1	-	1	1	-	-	-
2002.....	-	2	1	2	2	-	2	-	-	-	-	-
2001.....	-	1	-	1	5	1	1	-	-	-	-	-
2000.....	-	2	-	4	1	2	1	-	-	-	-	-
1999.....	-	3	2	1	4	4	1	-	-	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	1	-	-	-	-	-	1	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	1	-	-	-	-	-	-	-
2001.....	-	1	-	-	-	-	1	-	-	-	-	-
2000.....	-	-	-	2	1	-	-	-	-	-	-	-
1999.....	-	-	-	-	4	3	-	-	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	1	-	1	1	-	-	-
2002.....	-	2	1	2	1	-	2	-	-	-	-	-
2001.....	-	-	-	1	5	1	-	-	-	-	-	-
2000.....	-	2	-	2	-	2	1	-	-	-	-	-
1999.....	-	3	2	1	-	1	1	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Fibrosis and cirrhosis of liver (K74)												
All races, both sexes												
2004.....	1,757	1,635	1,526	1,521	1,735	1,739	1,262	634	183	27	2	-
2003.....	1,750	1,626	1,530	1,633	1,952	1,809	1,295	638	182	29	1	1
2002.....	1,638	1,511	1,420	1,582	1,815	1,840	1,302	621	206	21	3	2
2001.....	1,583	1,374	1,384	1,636	1,882	1,748	1,232	605	202	34	1	1
2000.....	1,360	1,336	1,397	1,598	1,872	1,770	1,205	640	178	27	4	1
1999.....	1,268	1,255	1,440	1,689	1,914	1,805	1,203	587	201	29	4	-
All races, male												
2004.....	1,255	1,124	904	824	893	836	548	230	74	9	2	-
2003.....	1,286	1,092	973	926	980	827	548	261	68	13	-	1
2002.....	1,203	1,041	835	868	930	858	572	249	66	7	-	2
2001.....	1,162	915	855	950	978	769	537	240	78	15	1	1
2000.....	989	885	891	896	969	811	484	267	69	11	1	1
1999.....	939	862	866	963	968	829	513	229	73	13	2	-
All races, female												
2004.....	502	511	622	697	842	903	714	404	109	18	-	-
2003.....	464	534	557	707	972	982	747	377	114	16	1	-
2002.....	435	470	585	714	885	982	730	372	140	14	3	-
2001.....	421	459	529	686	904	979	695	365	124	19	-	-
2000.....	371	451	506	702	903	959	721	373	109	16	3	-
1999.....	329	393	574	726	946	976	690	358	128	16	2	-
White, both sexes												
2004.....	1,446	1,369	1,323	1,357	1,597	1,631	1,188	588	169	26	1	-
2003.....	1,438	1,358	1,340	1,464	1,806	1,707	1,222	602	172	26	1	1
2002.....	1,344	1,298	1,232	1,424	1,680	1,735	1,223	588	193	20	3	2
2001.....	1,288	1,141	1,212	1,475	1,731	1,639	1,168	579	191	31	1	-
2000.....	1,098	1,137	1,213	1,419	1,714	1,675	1,138	603	164	26	4	1
1999.....	1,058	1,048	1,242	1,518	1,770	1,694	1,145	554	186	26	4	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Fibrosis and cirrhosis of liver (K74)												
White, male												
2004.....	1,040	949	789	732	824	789	521	219	69	9	1	-
2003.....	1,058	907	852	837	914	780	516	246	63	11	-	1
2002.....	996	889	721	778	867	823	538	238	62	7	-	2
2001.....	945	759	738	855	906	721	517	233	73	14	1	-
2000.....	798	754	779	809	894	765	462	251	63	11	1	1
1999.....	786	722	753	873	904	781	494	220	67	12	2	-
White, female												
2004.....	406	420	534	625	773	842	667	369	100	17	-	-
2003.....	380	451	488	627	892	927	706	356	109	15	1	-
2002.....	348	409	511	646	813	912	685	350	131	13	3	-
2001.....	343	382	474	620	825	918	651	346	118	17	-	-
2000.....	300	383	434	610	820	910	676	352	101	15	3	-
1999.....	272	326	489	645	866	913	651	334	119	14	2	-
Black, both sexes												
2004.....	270	214	156	114	104	66	46	30	8	1	1	-
2003.....	276	224	143	132	110	65	43	27	8	2	-	-
2002.....	261	179	150	119	92	81	42	20	11	-	-	-
2001.....	241	191	134	125	105	81	42	13	9	2	-	1
2000.....	229	159	144	134	119	67	45	24	12	1	-	-
1999.....	174	172	155	128	104	82	43	25	11	2	-	-
Black, male												
2004.....	189	151	91	65	57	35	20	9	2	-	1	-
2003.....	207	161	94	72	49	32	17	13	3	1	-	-
2002.....	185	131	94	73	48	27	18	6	3	-	-	-
2001.....	183	130	96	78	55	39	11	3	3	-	-	1
2000.....	170	109	93	71	62	34	14	9	4	-	-	-
1999.....	131	113	90	74	53	37	16	7	3	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Fibrosis and cirrhosis of liver (K74)												
Black, female												
2004.....	81	63	65	49	47	31	26	21	6	1	-	-
2003.....	69	63	49	60	61	33	26	14	5	1	-	-
2002.....	76	48	56	46	44	54	24	14	8	-	-	-
2001.....	58	61	38	47	50	42	31	10	6	2	-	-
2000.....	59	50	51	63	57	33	31	15	8	1	-	-
1999.....	43	59	65	54	51	45	27	18	8	2	-	-
American Indian 1/, both sexes												
2004.....	26	28	28	25	12	10	4	3	-	-	-	-
2003.....	20	24	29	14	8	10	6	1	2	-	-	-
2002.....	17	20	18	16	16	5	13	3	-	-	-	-
2001.....	31	22	16	13	16	8	6	2	-	-	-	-
2000.....	23	25	19	22	12	11	4	2	-	-	-	-
1999.....	16	24	22	19	12	6	1	1	-	-	-	-
American Indian 1/, male												
2004.....	12	12	12	11	5	3	1	-	-	-	-	-
2003.....	12	9	12	5	3	2	2	-	2	-	-	-
2002.....	12	11	9	8	7	2	7	1	-	-	-	-
2001.....	19	10	11	8	4	5	1	-	-	-	-	-
2000.....	16	14	6	4	3	6	-	1	-	-	-	-
1999.....	6	16	9	5	4	2	-	-	-	-	-	-
American Indian 1/, female												
2004.....	14	16	16	14	7	7	3	3	-	-	-	-
2003.....	8	15	17	9	5	8	4	1	-	-	-	-
2002.....	5	9	9	8	9	3	6	2	-	-	-	-
2001.....	12	12	5	5	12	3	5	2	-	-	-	-
2000.....	7	11	13	18	9	5	4	1	-	-	-	-
1999.....	10	8	13	14	8	4	1	1	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Fibrosis and cirrhosis of liver (K74)												
Asian or Pacific Islander 2/, both sexes												
2004.....	15	24	19	25	22	32	24	13	6	-	-	-
2003.....	16	20	18	23	28	27	24	8	-	1	-	-
2002.....	16	14	20	23	27	19	24	10	2	1	-	-
2001.....	23	20	22	23	30	20	16	11	2	1	-	-
2000.....	10	15	21	23	27	17	18	11	2	-	-	-
1999.....	20	11	21	24	28	23	14	7	4	1	-	-
Asian or Pacific Islander 2/, male												
2004.....	14	12	12	16	7	9	6	2	3	-	-	-
2003.....	9	15	15	12	14	13	13	2	-	1	-	-
2002.....	10	10	11	9	8	6	9	4	1	-	-	-
2001.....	15	16	10	9	13	4	8	4	2	1	-	-
2000.....	5	8	13	12	10	6	8	6	2	-	-	-
1999.....	16	11	14	11	7	9	3	2	3	1	-	-
Asian or Pacific Islander 2/, female												
2004.....	1	12	7	9	15	23	18	11	3	-	-	-
2003.....	7	5	3	11	14	14	11	6	-	-	-	-
2002.....	6	4	9	14	19	13	15	6	1	1	-	-
2001.....	8	4	12	14	17	16	8	7	-	-	-	-
2000.....	5	7	8	11	17	11	10	5	-	-	-	-
1999.....	4	-	7	13	21	14	11	5	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of liver (K71-K72,K75-K76)												
All races, both sexes												
2004.....	1,135	1,070	923	829	856	835	625	388	191	51	9	-
2003.....	1,159	1,018	934	904	791	781	635	409	172	47	10	1
2002.....	1,016	859	784	779	796	700	637	402	187	59	7	-
2001.....	899	754	777	736	751	807	594	391	147	61	10	2
2000.....	826	673	668	710	787	707	552	359	163	48	6	1
1999.....	757	673	682	797	826	716	539	348	162	59	12	1
All races, male												
2004.....	819	701	549	448	437	395	271	150	59	18	1	-
2003.....	792	642	580	509	428	368	289	167	63	11	3	1
2002.....	707	562	478	468	448	356	320	183	55	21	3	-
2001.....	637	468	479	431	397	414	290	159	47	17	3	2
2000.....	537	418	381	400	406	367	249	148	56	14	1	1
1999.....	508	436	419	468	430	332	268	156	57	13	3	1
All races, female												
2004.....	316	369	374	381	419	440	354	238	132	33	8	-
2003.....	367	376	354	395	363	413	346	242	109	36	7	-
2002.....	309	297	306	311	348	344	317	219	132	38	4	-
2001.....	262	286	298	305	354	393	304	232	100	44	7	-
2000.....	289	255	287	310	381	340	303	211	107	34	5	-
1999.....	249	237	263	329	396	384	271	192	105	46	9	-
White, both sexes												
2004.....	899	859	761	704	734	747	561	347	171	49	9	-
2003.....	914	834	776	771	680	694	580	375	146	42	7	1
2002.....	805	687	656	666	703	620	572	363	170	57	4	-
2001.....	725	599	647	639	647	716	529	349	139	53	10	1
2000.....	622	530	538	610	692	629	497	325	148	44	4	1
1999.....	594	535	549	680	714	640	496	311	148	54	10	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of liver (K71-K72,K75-K76)												
White, male												
2004.....	665	560	454	385	381	351	248	142	55	18	1	-
2003.....	629	519	483	429	370	334	268	155	50	11	2	1
2002.....	568	447	405	409	401	313	287	165	51	21	1	-
2001.....	523	375	405	387	346	362	256	144	44	13	3	1
2000.....	403	332	311	344	360	331	232	132	53	11	1	1
1999.....	404	356	343	405	380	304	254	145	54	11	2	-
White, female												
2004.....	234	299	307	319	353	396	313	205	116	31	8	-
2003.....	285	315	293	342	310	360	312	220	96	31	5	-
2002.....	237	240	251	257	302	307	285	198	119	36	3	-
2001.....	202	224	242	252	301	354	273	205	95	40	7	-
2000.....	219	198	227	266	332	298	265	193	95	33	3	-
1999.....	190	179	206	275	334	336	242	166	94	43	8	-
Black, both sexes												
2004.....	191	170	125	100	87	59	49	28	14	1	-	-
2003.....	210	146	128	108	81	61	35	25	18	5	3	-
2002.....	178	139	97	91	78	58	49	33	14	2	3	-
2001.....	152	140	104	84	85	69	53	25	7	5	-	1
2000.....	166	116	110	83	82	61	38	23	13	2	1	-
1999.....	141	114	117	94	83	63	32	27	11	4	2	1
Black, male												
2004.....	124	118	77	51	45	29	17	3	3	-	-	-
2003.....	139	98	81	69	48	25	17	8	9	-	1	-
2002.....	115	98	55	49	38	30	23	14	3	-	2	-
2001.....	103	83	60	40	43	37	29	8	3	2	-	1
2000.....	110	74	64	50	39	29	12	10	3	1	-	-
1999.....	90	69	66	48	39	22	9	10	2	1	1	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of liver (K71-K72,K75-K76)												
Black, female												
2004.....	67	52	48	49	42	30	32	25	11	1	-	-
2003.....	71	48	47	39	33	36	18	17	9	5	2	-
2002.....	63	41	42	42	40	28	26	19	11	2	1	-
2001.....	49	57	44	44	42	32	24	17	4	3	-	-
2000.....	56	42	46	33	43	32	26	13	10	1	1	-
1999.....	51	45	51	46	44	41	23	17	9	3	1	-
American Indian 1/, both sexes												
2004.....	27	24	10	6	13	12	3	3	1	-	-	-
2003.....	20	23	14	12	5	6	5	2	5	-	-	-
2002.....	22	12	17	11	4	4	2	-	-	-	-	-
2001.....	14	9	17	4	8	8	3	5	-	2	-	-
2000.....	26	11	5	6	4	3	6	1	-	-	1	-
1999.....	16	13	4	6	12	2	3	5	-	1	-	-
American Indian 1/, male												
2004.....	15	12	3	1	4	7	2	1	-	-	-	-
2003.....	15	14	6	6	1	1	1	-	3	-	-	-
2002.....	16	6	11	5	1	3	1	-	-	-	-	-
2001.....	8	6	10	2	4	3	2	-	-	1	-	-
2000.....	17	3	1	1	1	1	2	1	-	-	-	-
1999.....	10	6	2	3	6	1	2	1	-	1	-	-
American Indian 1/, female												
2004.....	12	12	7	5	9	5	1	2	1	-	-	-
2003.....	5	9	8	6	4	5	4	2	2	-	-	-
2002.....	6	6	6	6	3	1	1	-	-	-	-	-
2001.....	6	3	7	2	4	5	1	5	-	1	-	-
2000.....	9	8	4	5	3	2	4	-	-	-	1	-
1999.....	6	7	2	3	6	1	1	4	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of liver (K71-K72,K75-K76)												
Asian or Pacific Islander 2/, both sexes												
2004.....	18	17	27	19	22	17	12	10	5	1	-	-
2003.....	15	15	16	13	25	20	15	7	3	-	-	-
2002.....	11	21	14	11	11	18	14	6	3	-	-	-
2001.....	8	6	9	9	11	14	9	12	1	1	-	-
2000.....	12	16	15	11	9	14	11	10	2	2	-	-
1999.....	6	11	12	17	17	11	8	5	3	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	15	11	15	11	7	8	4	4	1	-	-	-
2003.....	9	11	10	5	9	8	3	4	1	-	-	-
2002.....	8	11	7	5	8	10	9	4	1	-	-	-
2001.....	3	4	4	2	4	12	3	7	-	1	-	-
2000.....	7	9	5	5	6	6	3	5	-	2	-	-
1999.....	4	5	8	12	5	5	3	-	1	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	3	6	12	8	15	9	8	6	4	1	-	-
2003.....	6	4	6	8	16	12	12	3	2	-	-	-
2002.....	3	10	7	6	3	8	5	2	2	-	-	-
2001.....	5	2	5	7	7	2	6	5	1	-	-	-
2000.....	5	7	10	6	3	8	8	5	2	-	-	-
1999.....	2	6	4	5	12	6	5	5	2	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cholelithiasis and other disorders of gallbladder (K80-K82)												
All races, both sexes												
2004.....	77	102	145	184	274	432	576	547	425	155	28	-
2003.....	80	77	105	193	296	440	614	498	371	133	25	-
2002.....	77	93	123	194	310	420	572	525	351	150	28	-
2001.....	68	94	148	193	333	432	513	542	364	106	31	-
2000.....	66	80	109	184	266	419	507	528	358	143	24	-
1999.....	69	95	108	171	304	417	531	521	324	120	27	-
All races, male												
2004.....	41	56	82	104	144	217	266	219	123	36	7	-
2003.....	44	45	61	101	164	233	286	199	99	27	3	-
2002.....	48	59	63	103	155	229	271	195	112	26	2	-
2001.....	40	61	81	96	173	231	233	209	111	21	8	-
2000.....	30	48	62	100	142	224	237	204	105	27	3	-
1999.....	30	54	66	89	163	208	239	203	101	22	4	-
All races, female												
2004.....	36	46	63	80	130	215	310	328	302	119	21	-
2003.....	36	32	44	92	132	207	328	299	272	106	22	-
2002.....	29	34	60	91	155	191	301	330	239	124	26	-
2001.....	28	33	67	97	160	201	280	333	253	85	23	-
2000.....	36	32	47	84	124	195	270	324	253	116	21	-
1999.....	39	41	42	82	141	209	292	318	223	98	23	-
White, both sexes												
2004.....	52	75	117	152	222	374	514	496	397	143	26	-
2003.....	52	59	83	154	267	390	552	455	335	121	22	-
2002.....	55	77	100	166	265	361	510	483	327	142	24	-
2001.....	50	72	119	150	283	379	456	493	335	97	29	-
2000.....	49	64	87	151	215	373	454	489	335	135	23	-
1999.....	54	72	81	134	264	378	486	481	299	113	25	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cholelithiasis and other disorders of gallbladder (K80-K82)												
White, male												
2004.....	27	42	71	88	123	192	243	204	117	35	7	-
2003.....	31	35	47	82	150	208	258	184	90	25	2	-
2002.....	35	50	51	90	136	204	243	179	105	25	2	-
2001.....	31	48	64	73	152	209	211	187	101	19	7	-
2000.....	23	41	51	87	119	203	220	193	99	26	2	-
1999.....	24	38	53	72	145	191	227	184	93	22	4	-
White, female												
2004.....	25	33	46	64	99	182	271	292	280	108	19	-
2003.....	21	24	36	72	117	182	294	271	245	96	20	-
2002.....	20	27	49	76	129	157	267	304	222	117	22	-
2001.....	19	24	55	77	131	170	245	306	234	78	22	-
2000.....	26	23	36	64	96	170	234	296	236	109	21	-
1999.....	30	34	28	62	119	187	259	297	206	91	21	-
Black, both sexes												
2004.....	22	22	24	22	39	42	49	38	20	10	1	-
2003.....	22	15	18	29	21	35	43	35	21	12	3	-
2002.....	19	12	17	22	34	45	49	26	14	6	4	-
2001.....	17	20	22	32	46	35	43	33	21	8	2	-
2000.....	14	13	16	28	37	37	39	33	18	6	-	-
1999.....	13	14	21	29	30	26	39	25	15	6	2	-
Black, male												
2004.....	13	13	8	12	14	17	15	11	3	-	-	-
2003.....	11	8	10	15	8	18	19	11	6	2	1	-
2002.....	10	6	11	9	10	17	21	8	3	-	-	-
2001.....	9	12	12	16	18	10	11	15	6	2	1	-
2000.....	6	5	9	10	15	15	12	9	3	-	-	-
1999.....	4	10	9	12	13	12	11	9	4	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cholelithiasis and other disorders of gallbladder (K80-K82)												
Black, female												
2004.....	9	9	16	10	25	25	34	27	17	10	1	-
2003.....	11	7	8	14	13	17	24	24	15	10	2	-
2002.....	9	6	6	13	24	28	28	18	11	6	4	-
2001.....	8	8	10	16	28	25	32	18	15	6	1	-
2000.....	8	8	7	18	22	22	27	24	15	6	-	-
1999.....	9	4	12	17	17	14	28	16	11	6	2	-
American Indian 1/, both sexes												
2004.....	-	2	3	3	2	5	2	4	2	1	1	-
2003.....	2	1	3	4	2	4	5	2	5	-	-	-
2002.....	2	1	3	3	3	3	2	4	2	-	-	-
2001.....	-	1	1	1	-	5	1	4	-	-	-	-
2000.....	1	1	1	-	2	5	3	3	1	1	-	-
1999.....	2	1	2	4	2	4	-	1	4	1	-	-
American Indian 1/, male												
2004.....	-	1	2	1	1	2	1	3	-	-	-	-
2003.....	-	1	3	2	2	2	2	-	-	-	-	-
2002.....	2	1	-	2	3	1	1	1	-	-	-	-
2001.....	-	1	1	1	-	3	-	3	-	-	-	-
2000.....	-	1	-	-	1	2	1	-	1	-	-	-
1999.....	2	-	2	2	1	1	-	-	1	-	-	-
American Indian 1/, female												
2004.....	-	1	1	2	1	3	1	1	2	1	1	-
2003.....	2	-	-	2	-	2	3	2	5	-	-	-
2002.....	-	-	3	1	-	2	1	3	2	-	-	-
2001.....	-	-	-	-	-	2	1	1	-	-	-	-
2000.....	1	-	1	-	1	3	2	3	-	1	-	-
1999.....	-	1	-	2	1	3	-	1	3	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cholelithiasis and other disorders of gallbladder (K80-K82)												
Asian or Pacific Islander 2/, both sexes												
2004.....	3	3	1	7	11	11	11	9	6	1	-	-
2003.....	4	2	1	6	6	11	14	6	10	-	-	-
2002.....	1	3	3	3	8	11	11	12	8	2	-	-
2001.....	1	1	6	10	4	13	13	12	8	1	-	-
2000.....	2	2	5	5	12	4	11	3	4	1	1	-
1999.....	-	8	4	4	8	9	6	14	6	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	1	-	1	3	6	6	7	1	3	1	-	-
2003.....	2	1	1	2	4	5	7	4	3	-	-	-
2002.....	1	2	1	2	6	7	6	7	4	1	-	-
2001.....	-	-	4	6	3	9	11	4	4	-	-	-
2000.....	1	1	2	3	7	4	4	2	2	1	1	-
1999.....	-	6	2	3	4	4	1	10	3	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	2	3	-	4	5	5	4	8	3	-	-	-
2003.....	2	1	-	4	2	6	7	2	7	-	-	-
2002.....	-	1	2	1	2	4	5	5	4	1	-	-
2001.....	1	1	2	4	1	4	2	8	4	1	-	-
2000.....	1	1	3	2	5	-	7	1	2	-	-	-
1999.....	-	2	2	1	4	5	5	4	3	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of biliary tract and pancreas (K83-K86)												
All races, both sexes												
2004.....	306	324	319	387	477	582	638	503	369	125	15	1
2003.....	288	293	348	376	509	621	664	552	334	127	19	1
2002.....	287	351	315	381	502	603	636	560	312	110	16	1
2001.....	313	297	345	398	532	563	614	508	357	120	25	-
2000.....	270	306	292	354	487	586	570	517	303	92	21	-
1999.....	269	267	303	347	480	530	504	517	285	104	15	-
All races, male												
2004.....	207	218	204	237	247	288	267	190	108	19	1	-
2003.....	197	202	215	232	280	312	296	190	79	24	6	1
2002.....	198	219	220	242	287	317	288	197	76	20	2	-
2001.....	198	194	205	227	311	268	241	180	81	28	2	-
2000.....	179	190	161	228	261	312	244	173	80	24	4	-
1999.....	176	174	181	204	261	274	210	189	70	17	-	-
All races, female												
2004.....	99	106	115	150	230	294	371	313	261	106	14	1
2003.....	91	91	133	144	229	309	368	362	255	103	13	-
2002.....	89	132	95	139	215	286	348	363	236	90	14	1
2001.....	115	103	140	171	221	295	373	328	276	92	23	-
2000.....	91	116	131	126	226	274	326	344	223	68	17	-
1999.....	93	93	122	143	219	256	294	328	215	87	15	-
White, both sexes												
2004.....	222	242	248	331	399	512	572	454	335	116	14	1
2003.....	207	226	277	308	419	564	605	499	305	113	18	-
2002.....	211	288	244	324	436	545	579	512	291	95	15	-
2001.....	222	233	285	334	463	504	560	464	319	111	23	-
2000.....	211	236	234	280	423	521	514	468	274	81	18	-
1999.....	206	207	238	291	430	467	452	461	262	90	14	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of biliary tract and pancreas (K83-K86)												
White, male												
2004.....	149	163	160	206	209	255	244	176	97	17	1	-
2003.....	143	155	168	194	236	282	270	171	71	21	5	-
2002.....	148	178	173	211	246	290	265	180	74	11	2	-
2001.....	144	153	169	191	278	239	226	163	71	25	2	-
2000.....	136	150	129	182	229	280	221	153	68	19	4	-
1999.....	128	131	147	176	238	246	192	160	64	13	-	-
White, female												
2004.....	73	79	88	125	190	257	328	278	238	99	13	1
2003.....	64	71	109	114	183	282	335	328	234	92	13	-
2002.....	63	110	71	113	190	255	314	332	217	84	13	-
2001.....	78	80	116	143	185	265	334	301	248	86	21	-
2000.....	75	86	105	98	194	241	293	315	206	62	14	-
1999.....	78	76	91	115	192	221	260	301	198	77	14	-
Black, both sexes												
2004.....	78	75	65	42	67	59	46	41	23	7	1	-
2003.....	73	62	63	56	70	46	41	44	25	10	1	1
2002.....	67	56	61	47	53	39	44	40	18	13	1	1
2001.....	83	55	49	56	56	46	43	34	30	6	2	-
2000.....	53	61	50	61	58	49	43	42	22	8	3	-
1999.....	58	54	59	51	39	47	43	42	19	11	-	-
Black, male												
2004.....	53	48	41	25	36	30	13	11	7	1	-	-
2003.....	48	44	40	29	36	22	20	15	7	2	1	1
2002.....	45	38	37	24	35	19	15	13	2	9	-	-
2001.....	51	38	29	33	27	24	10	11	5	3	-	-
2000.....	38	38	27	40	29	23	17	16	9	3	-	-
1999.....	44	39	30	26	16	22	13	20	5	3	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of biliary tract and pancreas (K83-K86)												
Black, female												
2004.....	25	27	24	17	31	29	33	30	16	6	1	-
2003.....	25	18	23	27	34	24	21	29	18	8	-	-
2002.....	22	18	24	23	18	20	29	27	16	4	1	1
2001.....	32	17	20	23	29	22	33	23	25	3	2	-
2000.....	15	23	23	21	29	26	26	26	13	5	3	-
1999.....	14	15	29	25	23	25	30	22	14	8	-	-
American Indian 1/, both sexes												
2004.....	3	2	1	5	1	2	5	1	-	-	-	-
2003.....	3	1	2	3	3	1	2	1	1	-	-	-
2002.....	6	4	1	2	4	2	4	-	1	1	-	-
2001.....	4	4	5	3	1	6	1	2	2	-	-	-
2000.....	3	4	1	5	1	1	2	1	1	-	-	-
1999.....	2	2	3	3	2	7	1	-	-	1	-	-
American Indian 1/, male												
2004.....	2	2	-	3	1	1	1	-	-	-	-	-
2003.....	3	-	2	2	1	1	1	-	-	-	-	-
2002.....	2	1	1	2	-	-	2	-	-	-	-	-
2001.....	2	2	2	1	-	2	1	-	2	-	-	-
2000.....	2	1	1	3	-	-	1	-	1	-	-	-
1999.....	2	1	1	1	1	2	1	-	-	1	-	-
American Indian 1/, female												
2004.....	1	-	1	2	-	1	4	1	-	-	-	-
2003.....	-	1	-	1	2	-	1	1	1	-	-	-
2002.....	4	3	-	-	4	2	2	-	1	1	-	-
2001.....	2	2	3	2	1	4	-	2	-	-	-	-
2000.....	1	3	-	2	1	1	1	-	-	-	-	-
1999.....	-	1	2	2	1	5	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of biliary tract and pancreas (K83-K86)												
Asian or Pacific Islander 2/, both sexes												
2004.....	3	5	5	9	10	9	15	7	11	2	-	-
2003.....	5	4	6	9	17	10	16	8	3	4	-	-
2002.....	3	3	9	8	9	17	9	8	2	1	-	-
2001.....	4	5	6	5	12	7	10	8	6	3	-	-
2000.....	3	5	7	8	5	15	11	6	6	3	-	-
1999.....	3	4	3	2	9	9	8	14	4	2	1	-
Asian or Pacific Islander 2/, male												
2004.....	3	5	3	3	1	2	9	3	4	1	-	-
2003.....	3	3	5	7	7	7	5	4	1	1	-	-
2002.....	3	2	9	5	6	8	6	4	-	-	-	-
2001.....	1	1	5	2	6	3	4	6	3	-	-	-
2000.....	3	1	4	3	3	9	5	4	2	2	-	-
1999.....	2	3	3	1	6	4	4	9	1	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	-	-	2	6	9	7	6	4	7	1	-	-
2003.....	2	1	1	2	10	3	11	4	2	3	-	-
2002.....	-	1	-	3	3	9	3	4	2	1	-	-
2001.....	3	4	1	3	6	4	6	2	3	3	-	-
2000.....	-	4	3	5	2	6	6	2	4	1	-	-
1999.....	1	1	-	1	3	5	4	5	3	2	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pancreatitis (K85,K86.0-K86.1)												
All races, both sexes												
2004.....	260	272	247	288	344	374	391	293	193	61	6	1
2003.....	241	233	272	277	355	426	430	307	168	69	10	1
2002.....	244	291	250	280	346	420	413	327	165	49	5	1
2001.....	258	253	279	307	397	400	399	299	201	70	12	-
2000.....	224	240	244	271	354	397	378	309	161	36	8	-
1999.....	230	205	235	280	358	380	334	319	147	56	9	-
All races, male												
2004.....	184	189	159	180	184	190	153	101	48	7	-	-
2003.....	163	164	177	180	195	207	188	104	35	10	4	1
2002.....	172	184	175	186	197	217	177	113	35	8	-	-
2001.....	162	170	166	171	236	187	150	108	43	14	2	-
2000.....	148	154	135	180	197	206	153	95	38	10	3	-
1999.....	149	136	147	167	198	194	135	111	34	6	-	-
All races, female												
2004.....	76	83	88	108	160	184	238	192	145	54	6	1
2003.....	78	69	95	97	160	219	242	203	133	59	6	-
2002.....	72	107	75	94	149	203	236	214	130	41	5	1
2001.....	96	83	113	136	161	213	249	191	158	56	10	-
2000.....	76	86	109	91	157	191	225	214	123	26	5	-
1999.....	81	69	88	113	160	186	199	208	113	50	9	-
White, both sexes												
2004.....	185	199	193	246	286	331	347	262	175	55	6	1
2003.....	173	181	214	233	287	388	389	273	154	59	9	-
2002.....	177	242	190	243	309	384	378	300	155	42	5	-
2001.....	177	203	228	261	349	364	360	269	184	66	10	-
2000.....	175	181	194	213	309	355	343	279	150	31	8	-
1999.....	170	159	184	237	317	333	300	286	135	52	8	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pancreatitis (K85,K86.0-K86.1)												
White, male												
2004.....	132	139	126	152	155	168	135	93	43	6	-	-
2003.....	116	124	136	150	160	187	169	92	31	9	3	-
2002.....	127	151	134	164	176	198	163	101	34	3	-	-
2001.....	114	136	133	147	215	168	137	95	41	14	2	-
2000.....	112	119	107	142	172	183	138	83	33	8	3	-
1999.....	104	103	120	143	180	176	122	94	30	5	-	-
White, female												
2004.....	53	60	67	94	131	163	212	169	132	49	6	1
2003.....	57	57	78	83	127	201	220	181	123	50	6	-
2002.....	50	91	56	79	133	186	215	199	121	39	5	-
2001.....	63	67	95	114	134	196	223	174	143	52	8	-
2000.....	63	62	87	71	137	172	205	196	117	23	5	-
1999.....	66	56	64	94	137	157	178	192	105	47	8	-
Black, both sexes												
2004.....	70	67	50	32	49	38	33	26	15	6	-	-
2003.....	62	50	51	38	53	30	27	28	12	7	1	1
2002.....	59	45	54	33	29	25	25	24	7	7	-	1
2001.....	76	44	42	39	38	29	32	23	16	2	2	-
2000.....	45	53	42	47	42	30	27	28	6	4	-	-
1999.....	55	43	46	41	31	36	28	25	11	4	-	-
Black, male												
2004.....	47	44	31	22	27	21	10	7	5	1	-	-
2003.....	42	39	35	24	28	14	14	9	4	1	1	1
2002.....	40	30	35	18	18	14	9	9	1	5	-	-
2001.....	45	32	27	22	17	16	9	8	2	-	-	-
2000.....	33	33	23	32	23	15	12	11	3	1	-	-
1999.....	41	32	23	23	12	15	10	11	4	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pancreatitis (K85,K86.0-K86.1)												
Black, female												
2004.....	23	23	19	10	22	17	23	19	10	5	-	-
2003.....	20	11	16	14	25	16	13	19	8	6	-	-
2002.....	19	15	19	15	11	11	16	15	6	2	-	1
2001.....	31	12	15	17	21	13	23	15	14	2	2	-
2000.....	12	20	19	15	19	15	15	17	3	3	-	-
1999.....	14	11	23	18	19	21	18	14	7	3	-	-
American Indian 1/, both sexes												
2004.....	2	2	1	5	1	-	3	1	-	-	-	-
2003.....	3	1	2	2	2	1	2	1	-	-	-	-
2002.....	5	2	-	1	3	2	3	-	1	-	-	-
2001.....	2	3	4	3	-	4	1	1	-	-	-	-
2000.....	2	3	1	5	1	-	1	-	-	-	-	-
1999.....	2	2	2	1	2	6	1	-	-	-	-	-
American Indian 1/, male												
2004.....	2	2	-	3	1	-	1	-	-	-	-	-
2003.....	3	-	2	2	1	1	1	-	-	-	-	-
2002.....	2	1	-	1	-	-	1	-	-	-	-	-
2001.....	2	2	2	1	-	1	1	-	-	-	-	-
2000.....	1	1	1	3	-	-	-	-	-	-	-	-
1999.....	2	1	1	-	1	2	1	-	-	-	-	-
American Indian 1/, female												
2004.....	-	-	1	2	-	-	2	1	-	-	-	-
2003.....	-	1	-	-	1	-	1	1	-	-	-	-
2002.....	3	1	-	-	3	2	2	-	1	-	-	-
2001.....	-	1	2	2	-	3	-	1	-	-	-	-
2000.....	1	2	-	2	1	-	1	-	-	-	-	-
1999.....	-	1	1	1	1	4	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pancreatitis (K85,K86.0-K86.1)												
Asian or Pacific Islander 2/, both sexes												
2004.....	3	4	3	5	8	5	8	4	3	-	-	-
2003.....	3	1	5	4	13	7	12	5	2	3	-	-
2002.....	3	2	6	3	5	9	7	3	2	-	-	-
2001.....	3	3	5	4	10	3	6	6	1	2	-	-
2000.....	2	3	7	6	2	12	7	2	5	1	-	-
1999.....	3	1	3	1	8	5	5	8	1	-	1	-
Asian or Pacific Islander 2/, male												
2004.....	3	4	2	3	1	1	7	1	-	-	-	-
2003.....	2	1	4	4	6	5	4	3	-	-	-	-
2002.....	3	2	6	3	3	5	4	3	-	-	-	-
2001.....	1	-	4	1	4	2	3	5	-	-	-	-
2000.....	2	1	4	3	2	8	3	1	2	1	-	-
1999.....	2	-	3	1	5	1	2	6	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	-	-	1	2	7	4	1	3	3	-	-	-
2003.....	1	-	1	-	7	2	8	2	2	3	-	-
2002.....	-	-	-	-	2	4	3	-	2	-	-	-
2001.....	2	3	1	3	6	1	3	1	1	2	-	-
2000.....	-	2	3	3	-	4	4	1	3	-	-	-
1999.....	1	1	-	-	3	4	3	2	1	-	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of biliary tract and pancreas (K83,K86.2-K86.9)												
All races, both sexes												
2004.....	46	52	72	99	133	208	247	210	176	64	9	-
2003.....	47	60	76	99	154	195	234	245	166	58	9	-
2002.....	43	60	65	101	156	183	223	233	147	61	11	-
2001.....	55	44	66	91	135	163	215	209	156	50	13	-
2000.....	46	66	48	83	133	189	192	208	142	56	13	-
1999.....	39	62	68	67	122	150	170	198	138	48	6	-
All races, male												
2004.....	23	29	45	57	63	98	114	89	60	12	1	-
2003.....	34	38	38	52	85	105	108	86	44	14	2	-
2002.....	26	35	45	56	90	100	111	84	41	12	2	-
2001.....	36	24	39	56	75	81	91	72	38	14	-	-
2000.....	31	36	26	48	64	106	91	78	42	14	1	-
1999.....	27	38	34	37	63	80	75	78	36	11	-	-
All races, female												
2004.....	23	23	27	42	70	110	133	121	116	52	8	-
2003.....	13	22	38	47	69	90	126	159	122	44	7	-
2002.....	17	25	20	45	66	83	112	149	106	49	9	-
2001.....	19	20	27	35	60	82	124	137	118	36	13	-
2000.....	15	30	22	35	69	83	101	130	100	42	12	-
1999.....	12	24	34	30	59	70	95	120	102	37	6	-
White, both sexes												
2004.....	37	43	55	85	113	181	225	192	160	61	8	-
2003.....	34	45	63	75	132	176	216	226	151	54	9	-
2002.....	34	46	54	81	127	161	201	212	136	53	10	-
2001.....	45	30	57	73	114	140	200	195	135	45	13	-
2000.....	36	55	40	67	114	166	171	189	124	50	10	-
1999.....	36	48	54	54	113	134	152	175	127	38	6	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of biliary tract and pancreas (K83,K86.2-K86.9)												
White, male												
2004.....	17	24	34	54	54	87	109	83	54	11	1	-
2003.....	27	31	32	44	76	95	101	79	40	12	2	-
2002.....	21	27	39	47	70	92	102	79	40	8	2	-
2001.....	30	17	36	44	63	71	89	68	30	11	-	-
2000.....	24	31	22	40	57	97	83	70	35	11	1	-
1999.....	24	28	27	33	58	70	70	66	34	8	-	-
White, female												
2004.....	20	19	21	31	59	94	116	109	106	50	7	-
2003.....	7	14	31	31	56	81	115	147	111	42	7	-
2002.....	13	19	15	34	57	69	99	133	96	45	8	-
2001.....	15	13	21	29	51	69	111	127	105	34	13	-
2000.....	12	24	18	27	57	69	88	119	89	39	9	-
1999.....	12	20	27	21	55	64	82	109	93	30	6	-
Black, both sexes												
2004.....	8	8	15	10	18	21	13	15	8	1	1	-
2003.....	11	12	12	18	17	16	14	16	13	3	-	-
2002.....	8	11	7	14	24	14	19	16	11	6	1	-
2001.....	7	11	7	17	18	17	11	11	14	4	-	-
2000.....	8	8	8	14	16	19	16	14	16	4	3	-
1999.....	3	11	13	10	8	11	15	17	8	7	-	-
Black, male												
2004.....	6	4	10	3	9	9	3	4	2	-	-	-
2003.....	6	5	5	5	8	8	6	6	3	1	-	-
2002.....	5	8	2	6	17	5	6	4	1	4	-	-
2001.....	6	6	2	11	10	8	1	3	3	3	-	-
2000.....	5	5	4	8	6	8	5	5	6	2	-	-
1999.....	3	7	7	3	4	7	3	9	1	2	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of biliary tract and pancreas (K83,K86.2-K86.9)												
Black, female												
2004.....	2	4	5	7	9	12	10	11	6	1	1	-
2003.....	5	7	7	13	9	8	8	10	10	2	-	-
2002.....	3	3	5	8	7	9	13	12	10	2	1	-
2001.....	1	5	5	6	8	9	10	8	11	1	-	-
2000.....	3	3	4	6	10	11	11	9	10	2	3	-
1999.....	-	4	6	7	4	4	12	8	7	5	-	-
American Indian 1/, both sexes												
2004.....	1	-	-	-	-	2	2	-	-	-	-	-
2003.....	-	-	-	1	1	-	-	-	1	-	-	-
2002.....	1	2	1	1	1	-	1	-	-	1	-	-
2001.....	2	1	1	-	1	2	-	1	2	-	-	-
2000.....	1	1	-	-	-	1	1	1	1	-	-	-
1999.....	-	-	1	2	-	1	-	-	-	1	-	-
American Indian 1/, male												
2004.....	-	-	-	-	-	1	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	1	1	-	-	1	-	-	-	-	-
2001.....	-	-	-	-	-	1	-	-	2	-	-	-
2000.....	1	-	-	-	-	-	1	-	1	-	-	-
1999.....	-	-	-	1	-	-	-	-	-	1	-	-
American Indian 1/, female												
2004.....	1	-	-	-	-	1	2	-	-	-	-	-
2003.....	-	-	-	1	1	-	-	-	1	-	-	-
2002.....	1	2	-	-	1	-	-	-	-	1	-	-
2001.....	2	1	1	-	1	1	-	1	-	-	-	-
2000.....	-	1	-	-	-	1	-	1	-	-	-	-
1999.....	-	-	1	1	-	1	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of biliary tract and pancreas (K83,K86.2-K86.9)												
Asian or Pacific Islander 2/, both sexes												
2004.....	-	1	2	4	2	4	7	3	8	2	-	-
2003.....	2	3	1	5	4	3	4	3	1	1	-	-
2002.....	-	1	3	5	4	8	2	5	-	1	-	-
2001.....	1	2	1	1	2	4	4	2	5	1	-	-
2000.....	1	2	-	2	3	3	4	4	1	2	-	-
1999.....	-	3	-	1	1	4	3	6	3	2	-	-
Asian or Pacific Islander 2/, male												
2004.....	-	1	1	-	-	1	2	2	4	1	-	-
2003.....	1	2	1	3	1	2	1	1	1	1	-	-
2002.....	-	-	3	2	3	3	2	1	-	-	-	-
2001.....	-	1	1	1	2	1	1	1	3	-	-	-
2000.....	1	-	-	-	1	1	2	3	-	1	-	-
1999.....	-	3	-	-	1	3	2	3	1	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	-	-	1	4	2	3	5	1	4	1	-	-
2003.....	1	1	-	2	3	1	3	2	-	-	-	-
2002.....	-	1	-	3	1	5	-	4	-	1	-	-
2001.....	1	1	-	-	-	3	3	1	2	1	-	-
2000.....	-	2	-	2	2	2	2	1	1	1	-	-
1999.....	-	-	-	1	-	1	1	3	2	2	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of digestive system (K90,K92)												
All races, both sexes												
2004.....	229	294	334	414	625	966	1,391	1,538	1,215	471	105	-
2003.....	204	265	346	457	659	1,046	1,407	1,603	1,257	518	118	-
2002.....	217	256	304	449	664	1,047	1,466	1,594	1,284	494	127	2
2001.....	187	254	344	461	674	1,049	1,450	1,539	1,235	531	122	-
2000.....	193	203	299	387	697	1,111	1,435	1,595	1,133	551	138	-
1999.....	170	223	300	452	686	1,034	1,358	1,455	1,149	490	114	1
All races, male												
2004.....	160	206	196	262	358	498	641	593	367	82	10	-
2003.....	141	175	207	276	369	552	664	599	351	97	12	-
2002.....	150	165	208	273	372	547	622	601	339	97	22	1
2001.....	140	170	204	277	376	535	636	564	364	109	21	-
2000.....	135	132	194	231	361	556	669	591	295	120	14	-
1999.....	118	143	183	267	388	522	619	521	317	105	21	-
All races, female												
2004.....	69	88	138	152	267	468	750	945	848	389	95	-
2003.....	63	90	139	181	290	494	743	1,004	906	421	106	-
2002.....	67	91	96	176	292	500	844	993	945	397	105	1
2001.....	47	84	140	184	298	514	814	975	871	422	101	-
2000.....	58	71	105	156	336	555	766	1,004	838	431	124	-
1999.....	52	80	117	185	298	512	739	934	832	385	93	1
White, both sexes												
2004.....	172	229	266	332	520	844	1,240	1,408	1,127	427	96	-
2003.....	162	210	273	357	551	899	1,275	1,472	1,159	478	104	-
2002.....	174	197	237	368	556	899	1,324	1,471	1,190	458	113	1
2001.....	138	190	265	372	578	907	1,322	1,422	1,142	478	109	-
2000.....	140	146	225	308	592	956	1,289	1,457	1,053	506	121	-
1999.....	122	175	233	367	589	894	1,223	1,347	1,066	456	101	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of digestive system (K90,K92)												
White, male												
2004.....	125	160	149	213	298	442	573	545	346	76	7	-
2003.....	114	138	168	223	306	477	606	552	328	92	11	-
2002.....	125	132	167	223	311	476	568	556	311	88	17	1
2001.....	102	127	161	225	321	459	587	522	340	94	18	-
2000.....	103	98	150	189	305	480	604	551	274	104	12	-
1999.....	85	111	149	221	333	452	566	479	291	96	20	-
White, female												
2004.....	47	69	117	119	222	402	667	863	781	351	89	-
2003.....	48	72	105	134	245	422	669	920	831	386	93	-
2002.....	49	65	70	145	245	423	756	915	879	370	96	-
2001.....	36	63	104	147	257	448	735	900	802	384	91	-
2000.....	37	48	75	119	287	476	685	906	779	402	109	-
1999.....	37	64	84	146	256	442	657	868	775	360	81	-
Black, both sexes												
2004.....	53	59	59	75	94	112	131	105	75	32	8	-
2003.....	40	50	65	86	93	132	108	108	83	38	12	-
2002.....	42	53	61	73	91	126	128	102	78	28	12	1
2001.....	44	59	72	79	83	127	116	101	80	45	12	-
2000.....	47	52	69	65	97	127	130	119	73	41	16	-
1999.....	39	41	59	71	90	123	115	94	72	29	12	1
Black, male												
2004.....	33	43	40	45	54	50	55	35	19	3	2	-
2003.....	26	34	35	43	55	65	46	37	16	4	-	-
2002.....	24	30	37	45	51	57	48	38	21	4	3	-
2001.....	34	40	39	45	49	65	45	33	16	12	2	-
2000.....	28	30	42	30	52	62	56	34	18	14	2	-
1999.....	27	31	29	38	49	61	44	32	20	8	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of digestive system (K90,K92)												
Black, female												
2004.....	20	16	19	30	40	62	76	70	56	29	6	-
2003.....	14	16	30	43	38	67	62	71	67	34	12	-
2002.....	18	23	24	28	40	69	80	64	57	24	9	1
2001.....	10	19	33	34	34	62	71	68	64	33	10	-
2000.....	19	22	27	35	45	65	74	85	55	27	14	-
1999.....	12	10	30	33	41	62	71	62	52	21	11	1
American Indian 1/, both sexes												
2004.....	3	3	7	5	1	3	1	5	2	2	-	-
2003.....	2	4	3	6	1	3	6	11	3	-	-	-
2002.....	-	3	2	5	3	3	1	6	2	2	-	-
2001.....	1	1	4	3	4	3	1	4	1	3	-	-
2000.....	3	1	3	6	2	6	3	6	3	1	-	-
1999.....	3	4	4	7	1	5	4	1	3	2	1	-
American Indian 1/, male												
2004.....	1	1	5	3	-	3	1	1	-	-	-	-
2003.....	1	2	1	4	-	2	3	5	-	-	-	-
2002.....	-	1	2	4	3	1	1	3	1	1	-	-
2001.....	1	1	2	2	2	1	-	2	1	1	-	-
2000.....	2	1	2	5	-	2	2	1	1	1	-	-
1999.....	2	-	2	3	1	4	-	1	1	-	-	-
American Indian 1/, female												
2004.....	2	2	2	2	1	-	-	4	2	2	-	-
2003.....	1	2	2	2	1	1	3	6	3	-	-	-
2002.....	-	2	-	1	-	2	-	3	1	1	-	-
2001.....	-	-	2	1	2	2	1	2	-	2	-	-
2000.....	1	-	1	1	2	4	1	5	2	-	-	-
1999.....	1	4	2	4	-	1	4	-	2	2	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of digestive system (K90,K92)												
Asian or Pacific Islander 2/, both sexes												
2004.....	1	3	2	2	10	7	19	20	11	10	1	-
2003.....	-	1	5	8	14	12	18	12	12	2	2	-
2002.....	1	3	4	3	14	19	13	15	14	6	2	-
2001.....	4	4	3	7	9	12	11	12	12	5	1	-
2000.....	3	4	2	8	6	22	13	13	4	3	1	-
1999.....	6	3	4	7	6	12	16	13	8	3	-	-
Asian or Pacific Islander 2/, male												
2004.....	1	2	2	1	6	3	12	12	2	3	1	-
2003.....	-	1	3	6	8	8	9	5	7	1	1	-
2002.....	1	2	2	1	7	13	5	4	6	4	2	-
2001.....	3	2	2	5	4	10	4	7	7	2	1	-
2000.....	2	3	-	7	4	12	7	5	2	1	-	-
1999.....	4	1	3	5	5	5	9	9	5	1	-	-
Asian or Pacific Islander 2/, female												
2004.....	-	1	-	1	4	4	7	8	9	7	-	-
2003.....	-	-	2	2	6	4	9	7	5	1	1	-
2002.....	-	1	2	2	7	6	8	11	8	2	-	-
2001.....	1	2	1	2	5	2	7	5	5	3	-	-
2000.....	1	1	2	1	2	10	6	8	2	2	1	-
1999.....	2	2	1	2	1	7	7	4	3	2	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XII. Diseases of the skin and subcutaneous tissue (L00-L98)												
All races, both sexes												
2004.....	145	197	216	266	363	524	708	653	465	235	47	-
2003.....	153	183	205	237	372	503	677	659	500	229	50	-
2002.....	129	167	157	196	362	484	625	735	500	206	67	-
2001.....	120	140	160	207	309	484	616	694	450	208	64	-
2000.....	85	125	158	197	350	473	629	665	502	251	53	1
1999.....	97	107	134	214	315	486	618	677	499	222	54	-
All races, male												
2004.....	71	85	100	103	154	218	253	200	127	49	5	-
2003.....	76	82	106	104	168	195	240	214	114	47	9	-
2002.....	62	79	64	89	173	199	239	239	122	27	9	-
2001.....	60	68	72	111	141	204	208	194	96	26	7	-
2000.....	37	63	83	72	158	185	226	199	114	44	7	1
1999.....	46	52	58	100	135	184	220	211	119	43	5	-
All races, female												
2004.....	74	112	116	163	209	306	455	453	338	186	42	-
2003.....	77	101	99	133	204	308	437	445	386	182	41	-
2002.....	67	88	93	107	189	285	386	496	378	179	58	-
2001.....	60	72	88	96	168	280	408	500	354	182	57	-
2000.....	48	62	75	125	192	288	403	466	388	207	46	-
1999.....	51	55	76	114	180	302	398	466	380	179	49	-
White, both sexes												
2004.....	106	136	162	195	255	418	565	533	378	181	36	-
2003.....	102	136	150	177	259	393	540	548	401	182	29	-
2002.....	84	117	102	140	255	371	480	571	396	159	51	-
2001.....	86	99	112	151	233	369	484	556	357	166	41	-
2000.....	60	88	114	140	245	349	493	524	393	184	37	-
1999.....	59	68	88	147	227	347	477	526	388	157	34	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XII. Diseases of the skin and subcutaneous tissue (L00-L98)												
White, male												
2004.....	53	62	79	75	110	180	206	169	109	33	3	-
2003.....	55	63	77	72	111	158	188	176	96	36	6	-
2002.....	41	60	45	61	123	158	195	193	105	23	7	-
2001.....	41	52	56	84	114	150	164	156	78	20	2	-
2000.....	28	47	63	48	108	136	174	163	88	29	6	-
1999.....	29	34	40	72	97	129	171	169	97	32	2	-
White, female												
2004.....	53	74	83	120	145	238	359	364	269	148	33	-
2003.....	47	73	73	105	148	235	352	372	305	146	23	-
2002.....	43	57	57	79	132	213	285	378	291	136	44	-
2001.....	45	47	56	67	119	219	320	400	279	146	39	-
2000.....	32	41	51	92	137	213	319	361	305	155	31	-
1999.....	30	34	48	75	130	218	306	357	291	125	32	-
Black, both sexes												
2004.....	37	56	48	64	102	99	134	108	81	51	11	-
2003.....	47	45	52	54	108	106	127	96	90	44	21	-
2002.....	42	47	51	48	93	111	135	154	101	44	14	-
2001.....	31	36	42	55	67	110	125	134	92	40	22	-
2000.....	20	34	40	54	94	112	128	136	98	61	16	1
1999.....	35	38	45	61	85	130	134	141	105	64	20	-
Black, male												
2004.....	16	20	18	25	40	36	43	26	16	14	2	-
2003.....	18	18	29	27	54	36	47	34	16	10	3	-
2002.....	21	17	18	24	46	40	43	41	15	2	1	-
2001.....	18	13	16	26	24	53	43	35	17	6	5	-
2000.....	7	14	18	24	45	43	49	34	22	14	1	1
1999.....	16	17	17	26	36	51	45	39	20	11	3	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XII. Diseases of the skin and subcutaneous tissue (L00-L98)												
Black, female												
2004.....	21	36	30	39	62	63	91	82	65	37	9	-
2003.....	29	27	23	27	54	70	80	62	74	34	18	-
2002.....	21	30	33	24	47	71	92	113	86	42	13	-
2001.....	13	23	26	29	43	57	82	99	75	34	17	-
2000.....	13	20	22	30	49	69	79	102	76	47	15	-
1999.....	19	21	28	35	49	79	89	102	85	53	17	-
American Indian 1/, both sexes												
2004.....	-	1	4	1	1	1	2	1	1	1	-	-
2003.....	3	1	2	1	1	3	4	2	4	-	-	-
2002.....	2	1	1	2	5	-	3	2	-	2	-	-
2001.....	-	2	3	-	2	2	1	-	-	1	1	-
2000.....	3	-	2	1	5	4	3	-	3	4	-	-
1999.....	1	-	-	4	1	3	1	4	2	1	-	-
American Indian 1/, male												
2004.....	-	-	2	-	1	-	1	-	-	-	-	-
2003.....	2	-	-	-	1	1	1	-	1	-	-	-
2002.....	-	-	-	1	3	-	-	-	-	1	-	-
2001.....	-	1	-	-	1	-	-	-	-	-	-	-
2000.....	1	-	1	-	4	1	1	-	2	-	-	-
1999.....	-	-	-	1	-	1	1	-	-	-	-	-
American Indian 1/, female												
2004.....	-	1	2	1	-	1	1	1	1	1	-	-
2003.....	1	1	2	1	-	2	3	2	3	-	-	-
2002.....	2	1	1	1	2	-	3	2	-	1	-	-
2001.....	-	1	3	-	1	2	1	-	-	1	1	-
2000.....	2	-	1	1	1	3	2	-	1	4	-	-
1999.....	1	-	-	3	1	2	-	4	2	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XII. Diseases of the skin and subcutaneous tissue (L00-L98)												
Asian or Pacific Islander 2/, both sexes												
2004.....	2	4	2	6	5	6	7	11	5	2	-	-
2003.....	1	1	1	5	4	1	6	13	5	3	-	-
2002.....	1	2	3	6	9	2	7	8	3	1	2	-
2001.....	3	3	3	1	7	3	6	4	1	1	-	-
2000.....	2	3	2	2	6	8	5	5	8	2	-	-
1999.....	2	1	1	2	2	6	6	6	4	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	2	3	1	3	3	2	3	5	2	2	-	-
2003.....	1	1	-	5	2	-	4	4	1	1	-	-
2002.....	-	2	1	3	1	1	1	5	2	1	1	-
2001.....	1	2	-	1	2	1	1	3	1	-	-	-
2000.....	1	2	1	-	1	5	2	2	2	1	-	-
1999.....	1	1	1	1	2	3	3	3	2	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	-	1	1	3	2	4	4	6	3	-	-	-
2003.....	-	-	1	-	2	1	2	9	4	2	-	-
2002.....	1	-	2	3	8	1	6	3	1	-	1	-
2001.....	2	1	3	-	5	2	5	1	-	1	-	-
2000.....	1	1	1	2	5	3	3	3	6	1	-	-
1999.....	1	-	-	1	-	3	3	3	2	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Infections of skin and subcutaneous tissue (L00-L08)												
All races, both sexes												
2004.....	73	91	104	121	145	222	275	240	173	81	11	-
2003.....	80	92	94	101	144	200	245	234	171	77	19	-
2002.....	58	80	67	78	115	163	189	196	157	59	16	-
2001.....	58	60	74	85	104	151	176	210	128	63	15	-
2000.....	32	60	68	68	121	146	168	180	142	62	11	1
1999.....	36	44	52	67	95	108	143	153	122	54	16	-
All races, male												
2004.....	43	46	47	50	68	102	106	84	58	12	2	-
2003.....	51	51	51	49	67	82	98	83	45	16	6	-
2002.....	38	46	34	38	56	67	83	73	40	10	4	-
2001.....	32	34	41	48	54	66	55	66	32	7	-	-
2000.....	19	39	37	32	63	64	73	66	34	8	1	1
1999.....	22	27	22	31	40	39	60	55	34	18	1	-
All races, female												
2004.....	30	45	57	71	77	120	169	156	115	69	9	-
2003.....	29	41	43	52	77	118	147	151	126	61	13	-
2002.....	20	34	33	40	59	96	106	123	117	49	12	-
2001.....	26	26	33	37	50	85	121	144	96	56	15	-
2000.....	13	21	31	36	58	82	95	114	108	54	10	-
1999.....	14	17	30	36	55	69	83	98	88	36	15	-
White, both sexes												
2004.....	59	71	86	100	121	203	244	214	161	70	11	-
2003.....	61	80	78	83	117	180	221	216	160	72	12	-
2002.....	45	62	56	59	93	141	171	189	143	55	14	-
2001.....	47	45	58	72	95	136	157	198	116	56	14	-
2000.....	27	49	58	55	104	127	152	164	130	53	10	-
1999.....	26	31	40	56	83	99	134	142	112	47	12	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Infections of skin and subcutaneous tissue (L00-L08)												
White, male												
2004.....	33	36	40	39	58	93	97	80	56	9	2	-
2003.....	38	44	43	38	58	73	89	76	42	16	4	-
2002.....	29	37	31	29	47	60	79	70	36	9	3	-
2001.....	24	28	36	43	51	58	52	62	31	6	-	-
2000.....	16	33	34	25	54	56	66	61	29	6	1	-
1999.....	16	19	19	30	36	32	55	50	30	15	1	-
White, female												
2004.....	26	35	46	61	63	110	147	134	105	61	9	-
2003.....	23	36	35	45	59	107	132	140	118	56	8	-
2002.....	16	25	25	30	46	81	92	119	107	46	11	-
2001.....	23	17	22	29	44	78	105	136	85	50	14	-
2000.....	11	16	24	30	50	71	86	103	101	47	9	-
1999.....	10	12	21	26	47	67	79	92	82	32	11	-
Black, both sexes												
2004.....	13	16	15	17	21	17	26	23	12	10	-	-
2003.....	15	10	15	15	24	19	19	13	8	5	7	-
2002.....	12	15	9	17	16	22	14	6	14	3	1	-
2001.....	11	12	13	13	6	13	17	11	12	6	1	-
2000.....	4	10	8	12	12	15	14	14	9	8	1	1
1999.....	9	13	12	8	10	7	8	10	7	7	4	-
Black, male												
2004.....	9	8	5	9	7	8	8	4	2	3	-	-
2003.....	10	6	8	8	8	8	6	6	2	-	2	-
2002.....	9	7	2	8	8	7	4	2	4	-	-	-
2001.....	8	4	5	5	2	7	3	3	1	1	-	-
2000.....	2	5	2	7	7	5	6	3	3	2	-	1
1999.....	6	8	3	-	3	6	4	4	3	3	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Infections of skin and subcutaneous tissue (L00-L08)												
Black, female												
2004.....	4	8	10	8	14	9	18	19	10	7	-	-
2003.....	5	4	7	7	16	11	13	7	6	5	5	-
2002.....	3	8	7	9	8	15	10	4	10	3	1	-
2001.....	3	8	8	8	4	6	14	8	11	5	1	-
2000.....	2	5	6	5	5	10	8	11	6	6	1	-
1999.....	3	5	9	8	7	1	4	6	4	4	4	-
American Indian 1/, both sexes												
2004.....	-	1	2	-	-	1	1	1	-	1	-	-
2003.....	3	1	1	-	-	1	-	1	1	-	-	-
2002.....	1	1	1	-	2	-	2	-	-	-	-	-
2001.....	-	2	1	-	1	1	-	-	-	1	-	-
2000.....	1	-	2	1	2	2	-	-	1	1	-	-
1999.....	1	-	-	3	1	1	-	-	1	-	-	-
American Indian 1/, male												
2004.....	-	-	1	-	-	-	1	-	-	-	-	-
2003.....	2	-	-	-	-	1	-	-	-	-	-	-
2002.....	-	-	-	-	1	-	-	-	-	-	-	-
2001.....	-	1	-	-	1	-	-	-	-	-	-	-
2000.....	1	-	1	-	2	1	-	-	1	-	-	-
1999.....	-	-	-	1	-	1	-	-	-	-	-	-
American Indian 1/, female												
2004.....	-	1	1	-	-	1	-	1	-	1	-	-
2003.....	1	1	1	-	-	-	-	1	1	-	-	-
2002.....	1	1	1	-	1	-	2	-	-	-	-	-
2001.....	-	1	1	-	-	1	-	-	-	1	-	-
2000.....	-	-	1	1	-	1	-	-	-	1	-	-
1999.....	1	-	-	2	1	-	-	-	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Infections of skin and subcutaneous tissue (L00-L08)												
Asian or Pacific Islander 2/, both sexes												
2004.....	1	3	1	4	3	1	4	2	-	-	-	-
2003.....	1	1	-	3	3	-	5	4	2	-	-	-
2002.....	-	2	1	2	4	-	2	1	-	1	1	-
2001.....	-	1	2	-	2	1	2	1	-	-	-	-
2000.....	-	1	-	-	3	2	2	2	2	-	-	-
1999.....	-	-	-	-	1	1	1	1	2	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	1	2	1	2	3	1	-	-	-	-	-	-
2003.....	1	1	-	3	1	-	3	1	1	-	-	-
2002.....	-	2	1	1	-	-	-	1	-	1	1	-
2001.....	-	1	-	-	-	1	-	1	-	-	-	-
2000.....	-	1	-	-	-	2	1	2	1	-	-	-
1999.....	-	-	-	-	1	-	1	1	1	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	-	1	-	2	-	-	4	2	-	-	-	-
2003.....	-	-	-	-	2	-	2	3	1	-	-	-
2002.....	-	-	-	1	4	-	2	-	-	-	-	-
2001.....	-	-	2	-	2	-	2	-	-	-	-	-
2000.....	-	-	-	-	3	-	1	-	1	-	-	-
1999.....	-	-	-	-	-	1	-	-	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified diseases of skin and subcutaneous tissue (L10-L98)												
All races, both sexes												
2004.....	72	106	112	145	218	302	433	413	292	154	36	-
2003.....	73	91	111	136	228	303	432	425	329	152	31	-
2002.....	71	87	90	118	247	321	436	539	343	147	51	-
2001.....	62	80	86	122	205	333	440	484	322	145	49	-
2000.....	53	65	90	129	229	327	461	485	360	189	42	-
1999.....	61	63	82	147	220	378	475	524	377	168	38	-
All races, male												
2004.....	28	39	53	53	86	116	147	116	69	37	3	-
2003.....	25	31	55	55	101	113	142	131	69	31	3	-
2002.....	24	33	30	51	117	132	156	166	82	17	5	-
2001.....	28	34	31	63	87	138	153	128	64	19	7	-
2000.....	18	24	46	40	95	121	153	133	80	36	6	-
1999.....	24	25	36	69	95	145	160	156	85	25	4	-
All races, female												
2004.....	44	67	59	92	132	186	286	297	223	117	33	-
2003.....	48	60	56	81	127	190	290	294	260	121	28	-
2002.....	47	54	60	67	130	189	280	373	261	130	46	-
2001.....	34	46	55	59	118	195	287	356	258	126	42	-
2000.....	35	41	44	89	134	206	308	352	280	153	36	-
1999.....	37	38	46	78	125	233	315	368	292	143	34	-
White, both sexes												
2004.....	47	65	76	95	134	215	321	319	217	111	25	-
2003.....	41	56	72	94	142	213	319	332	241	110	17	-
2002.....	39	55	46	81	162	230	309	382	253	104	37	-
2001.....	39	54	54	79	138	233	327	358	241	110	27	-
2000.....	33	39	56	85	141	222	341	360	263	131	27	-
1999.....	33	37	48	91	144	248	343	384	276	110	22	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified diseases of skin and subcutaneous tissue (L10-L98)												
White, male												
2004.....	20	26	39	36	52	87	109	89	53	24	1	-
2003.....	17	19	34	34	53	85	99	100	54	20	2	-
2002.....	12	23	14	32	76	98	116	123	69	14	4	-
2001.....	17	24	20	41	63	92	112	94	47	14	2	-
2000.....	12	14	29	23	54	80	108	102	59	23	5	-
1999.....	13	15	21	42	61	97	116	119	67	17	1	-
White, female												
2004.....	27	39	37	59	82	128	212	230	164	87	24	-
2003.....	24	37	38	60	89	128	220	232	187	90	15	-
2002.....	27	32	32	49	86	132	193	259	184	90	33	-
2001.....	22	30	34	38	75	141	215	264	194	96	25	-
2000.....	21	25	27	62	87	142	233	258	204	108	22	-
1999.....	20	22	27	49	83	151	227	265	209	93	21	-
Black, both sexes												
2004.....	24	40	33	47	81	82	108	85	69	41	11	-
2003.....	32	35	37	39	84	87	108	83	82	39	14	-
2002.....	30	32	42	31	77	89	121	148	87	41	13	-
2001.....	20	24	29	42	61	97	108	123	80	34	21	-
2000.....	16	24	32	42	82	97	114	122	89	53	15	-
1999.....	26	25	33	53	75	123	126	131	98	57	16	-
Black, male												
2004.....	7	12	13	16	33	28	35	22	14	11	2	-
2003.....	8	12	21	19	46	28	41	28	14	10	1	-
2002.....	12	10	16	16	38	33	39	39	11	2	1	-
2001.....	10	9	11	21	22	46	40	32	16	5	5	-
2000.....	5	9	16	17	38	38	43	31	19	12	1	-
1999.....	10	9	14	26	33	45	41	35	17	8	3	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified diseases of skin and subcutaneous tissue (L10-L98)												
Black, female												
2004.....	17	28	20	31	48	54	73	63	55	30	9	-
2003.....	24	23	16	20	38	59	67	55	68	29	13	-
2002.....	18	22	26	15	39	56	82	109	76	39	12	-
2001.....	10	15	18	21	39	51	68	91	64	29	16	-
2000.....	11	15	16	25	44	59	71	91	70	41	14	-
1999.....	16	16	19	27	42	78	85	96	81	49	13	-
American Indian 1/, both sexes												
2004.....	-	-	2	1	1	-	1	-	1	-	-	-
2003.....	-	-	1	1	1	2	4	1	3	-	-	-
2002.....	1	-	-	2	3	-	1	2	-	2	-	-
2001.....	-	-	2	-	1	1	1	-	-	-	1	-
2000.....	2	-	-	-	3	2	3	-	2	3	-	-
1999.....	-	-	-	1	-	2	1	4	1	1	-	-
American Indian 1/, male												
2004.....	-	-	1	-	1	-	-	-	-	-	-	-
2003.....	-	-	-	-	1	-	1	-	1	-	-	-
2002.....	-	-	-	1	2	-	-	-	-	1	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	2	-	1	-	1	-	-	-
1999.....	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, female												
2004.....	-	-	1	1	-	-	1	-	1	-	-	-
2003.....	-	-	1	1	-	2	3	1	2	-	-	-
2002.....	1	-	-	1	1	-	1	2	-	1	-	-
2001.....	-	-	2	-	1	1	1	-	-	-	1	-
2000.....	2	-	-	-	1	2	2	-	1	3	-	-
1999.....	-	-	-	1	-	2	-	4	1	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified diseases of skin and subcutaneous tissue (L10-L98)												
Asian or Pacific Islander 2/, both sexes												
2004.....	1	1	1	2	2	5	3	9	5	2	-	-
2003.....	-	-	1	2	1	1	1	9	3	3	-	-
2002.....	1	-	2	4	5	2	5	7	3	-	1	-
2001.....	3	2	1	1	5	2	4	3	1	1	-	-
2000.....	2	2	2	2	3	6	3	3	6	2	-	-
1999.....	2	1	1	2	1	5	5	5	2	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	1	1	-	1	-	1	3	5	2	2	-	-
2003.....	-	-	-	2	1	-	1	3	-	1	-	-
2002.....	-	-	-	2	1	1	1	4	2	-	-	-
2001.....	1	1	-	1	2	-	1	2	1	-	-	-
2000.....	1	1	1	-	1	3	1	-	1	1	-	-
1999.....	1	1	1	1	1	3	2	2	1	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	-	-	1	1	2	4	-	4	3	-	-	-
2003.....	-	-	1	-	-	1	-	6	3	2	-	-
2002.....	1	-	2	2	4	1	4	3	1	-	1	-
2001.....	2	1	1	-	3	2	3	1	-	1	-	-
2000.....	1	1	1	2	2	3	2	3	5	1	-	-
1999.....	1	-	-	1	-	2	3	3	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XIII. Diseases of the musculoskeletal system and connective tissue (M00-M99)												
All races, both sexes												
2004.....	637	757	825	950	1,276	1,803	2,273	2,014	1,432	585	145	-
2003.....	629	701	848	957	1,365	1,883	2,192	2,031	1,422	636	171	-
2002.....	571	652	817	971	1,382	1,903	2,136	2,080	1,505	615	157	1
2001.....	571	598	735	966	1,385	1,885	2,164	1,975	1,433	582	122	-
2000.....	522	585	720	980	1,345	1,920	2,065	1,876	1,419	539	151	-
1999.....	533	567	717	962	1,380	1,876	1,921	1,827	1,307	519	137	-
All races, male												
2004.....	253	253	323	332	428	605	657	515	278	87	11	-
2003.....	220	251	293	381	482	644	619	506	265	76	19	-
2002.....	209	219	294	345	501	595	630	496	284	88	15	1
2001.....	219	217	251	314	479	614	626	463	254	91	13	-
2000.....	154	222	244	341	471	626	605	461	272	82	13	-
1999.....	177	194	241	304	485	608	557	437	239	69	15	-
All races, female												
2004.....	384	504	502	618	848	1,198	1,616	1,499	1,154	498	134	-
2003.....	409	450	555	576	883	1,239	1,573	1,525	1,157	560	152	-
2002.....	362	433	523	626	881	1,308	1,506	1,584	1,221	527	142	-
2001.....	352	381	484	652	906	1,271	1,538	1,512	1,179	491	109	-
2000.....	368	363	476	639	874	1,294	1,460	1,415	1,147	457	138	-
1999.....	356	373	476	658	895	1,268	1,364	1,390	1,068	450	122	-
White, both sexes												
2004.....	436	564	665	784	1,113	1,604	2,085	1,863	1,350	547	128	-
2003.....	445	541	683	794	1,178	1,686	1,993	1,893	1,322	586	152	-
2002.....	418	499	660	815	1,210	1,713	1,950	1,931	1,422	583	141	-
2001.....	404	468	576	815	1,199	1,709	2,003	1,830	1,337	546	105	-
2000.....	374	440	579	846	1,173	1,732	1,902	1,762	1,317	507	131	-
1999.....	372	427	584	802	1,231	1,699	1,798	1,717	1,232	483	124	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XIII. Diseases of the musculoskeletal system and connective tissue (M00-M99)												
White, male												
2004.....	180	196	268	277	363	535	601	467	262	75	10	-
2003.....	168	192	236	308	407	562	553	466	237	67	15	-
2002.....	164	172	240	281	435	535	564	455	261	79	13	-
2001.....	162	167	199	262	413	558	573	432	231	84	9	-
2000.....	115	179	198	292	395	556	549	414	244	77	11	-
1999.....	135	146	201	254	438	527	525	403	219	60	13	-
White, female												
2004.....	256	368	397	507	750	1,069	1,484	1,396	1,088	472	118	-
2003.....	277	349	447	486	771	1,124	1,440	1,427	1,085	519	137	-
2002.....	254	327	420	534	775	1,178	1,386	1,476	1,161	504	128	-
2001.....	242	301	377	553	786	1,151	1,430	1,398	1,106	462	96	-
2000.....	259	261	381	554	778	1,176	1,353	1,348	1,073	430	120	-
1999.....	237	281	383	548	793	1,172	1,273	1,314	1,013	423	111	-
Black, both sexes												
2004.....	183	158	137	140	135	160	147	115	67	33	15	-
2003.....	166	134	140	135	153	160	153	97	83	40	13	-
2002.....	137	132	131	127	140	149	149	118	71	26	12	1
2001.....	150	113	125	131	146	144	133	121	80	28	17	-
2000.....	129	120	121	112	140	151	133	96	84	22	16	-
1999.....	130	117	113	138	129	151	94	88	62	29	10	-
Black, male												
2004.....	66	44	49	48	57	61	43	36	15	9	1	-
2003.....	49	48	52	64	64	67	52	27	22	6	1	-
2002.....	41	45	46	50	56	47	52	32	17	6	2	1
2001.....	53	42	45	46	51	49	46	26	18	4	4	-
2000.....	32	38	40	46	65	54	47	43	20	3	2	-
1999.....	34	38	32	47	42	71	24	30	15	8	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XIII. Diseases of the musculoskeletal system and connective tissue (M00-M99)												
Black, female												
2004.....	117	114	88	92	78	99	104	79	52	24	14	-
2003.....	117	86	88	71	89	93	101	70	61	34	12	-
2002.....	96	87	85	77	84	102	97	86	54	20	10	-
2001.....	97	71	80	85	95	95	87	95	62	24	13	-
2000.....	97	82	81	66	75	97	86	53	64	19	14	-
1999.....	96	79	81	91	87	80	70	58	47	21	9	-
American Indian 1/, both sexes												
2004.....	8	13	11	9	9	6	12	6	4	1	-	-
2003.....	9	10	10	11	12	11	11	10	2	3	1	-
2002.....	10	7	10	13	6	9	8	3	3	2	-	-
2001.....	5	4	11	8	15	9	10	10	3	1	-	-
2000.....	6	10	8	8	9	10	11	3	3	1	-	-
1999.....	12	14	4	7	7	8	4	6	3	3	1	-
American Indian 1/, male												
2004.....	3	7	4	4	3	2	4	1	1	-	-	-
2003.....	2	2	4	3	4	5	2	3	1	-	1	-
2002.....	3	-	3	7	-	4	-	-	1	-	-	-
2001.....	1	1	2	3	6	2	1	3	2	1	-	-
2000.....	1	2	2	1	3	4	5	1	1	-	-	-
1999.....	4	6	2	-	2	2	-	-	1	1	1	-
American Indian 1/, female												
2004.....	5	6	7	5	6	4	8	5	3	1	-	-
2003.....	7	8	6	8	8	6	9	7	1	3	-	-
2002.....	7	7	7	6	6	5	8	3	2	2	-	-
2001.....	4	3	9	5	9	7	9	7	1	-	-	-
2000.....	5	8	6	7	6	6	6	2	2	1	-	-
1999.....	8	8	2	7	5	6	4	6	2	2	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XIII. Diseases of the musculoskeletal system and connective tissue (M00-M99)												
Asian or Pacific Islander 2/, both sexes												
2004.....	10	22	12	17	19	33	29	30	11	4	2	-
2003.....	9	16	15	17	22	26	35	31	15	7	5	-
2002.....	6	14	16	16	26	32	29	28	9	4	4	-
2001.....	12	13	23	12	25	23	18	14	13	7	-	-
2000.....	13	15	12	14	23	27	19	15	15	9	4	-
1999.....	19	9	16	15	13	18	25	16	10	4	2	-
Asian or Pacific Islander 2/, male												
2004.....	4	6	2	3	5	7	9	11	-	3	-	-
2003.....	1	9	1	6	7	10	12	10	5	3	2	-
2002.....	1	2	5	7	10	9	14	9	5	3	-	-
2001.....	3	7	5	3	9	5	6	2	3	2	-	-
2000.....	6	3	4	2	8	12	4	3	7	2	-	-
1999.....	4	4	6	3	3	8	8	4	4	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	6	16	10	14	14	26	20	19	11	1	2	-
2003.....	8	7	14	11	15	16	23	21	10	4	3	-
2002.....	5	12	11	9	16	23	15	19	4	1	4	-
2001.....	9	6	18	9	16	18	12	12	10	5	-	-
2000.....	7	12	8	12	15	15	15	12	8	7	4	-
1999.....	15	5	10	12	10	10	17	12	6	4	2	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatoid arthritis and related inflammatory polyarthropathies (M05-M08)												
All races, both sexes												
2004.....	57	110	144	226	346	503	533	371	152	40	13	-
2003.....	55	91	159	211	363	524	591	383	163	55	4	-
2002.....	55	107	160	241	361	591	565	416	184	46	6	-
2001.....	61	105	152	234	374	566	539	354	154	45	6	-
2000.....	57	85	155	256	390	556	576	346	170	26	4	-
1999.....	48	101	170	239	398	640	517	386	147	39	7	-
All races, male												
2004.....	19	32	51	72	84	128	97	69	22	5	2	-
2003.....	12	31	60	60	100	141	116	69	21	3	1	-
2002.....	18	37	53	77	100	137	132	71	30	5	-	-
2001.....	17	39	47	67	103	142	112	51	16	4	-	-
2000.....	12	29	45	74	121	133	121	67	16	2	-	-
1999.....	19	33	57	69	120	152	100	64	20	2	-	-
All races, female												
2004.....	38	78	93	154	262	375	436	302	130	35	11	-
2003.....	43	60	99	151	263	383	475	314	142	52	3	-
2002.....	37	70	107	164	261	454	433	345	154	41	6	-
2001.....	44	66	105	167	271	424	427	303	138	41	6	-
2000.....	45	56	110	182	269	423	455	279	154	24	4	-
1999.....	29	68	113	170	278	488	417	322	127	37	7	-
White, both sexes												
2004.....	45	98	129	199	316	458	488	346	140	38	8	-
2003.....	51	83	145	188	335	484	539	351	158	48	3	-
2002.....	46	91	138	218	330	545	524	387	176	45	5	-
2001.....	50	91	133	219	344	530	508	326	140	40	5	-
2000.....	50	72	139	235	364	517	546	330	158	22	3	-
1999.....	42	87	159	219	379	596	488	370	144	39	7	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatoid arthritis and related inflammatory polyarthropathies (M05-M08)												
White, male												
2004.....	16	29	48	63	77	120	92	61	22	5	2	-
2003.....	12	26	56	57	91	127	106	64	19	3	1	-
2002.....	17	32	48	64	94	129	121	67	28	5	-	-
2001.....	14	37	42	63	94	134	101	46	16	3	-	-
2000.....	11	27	42	69	111	123	115	65	15	1	-	-
1999.....	16	29	55	62	115	138	96	62	20	2	-	-
White, female												
2004.....	29	69	81	136	239	338	396	285	118	33	6	-
2003.....	39	57	89	131	244	357	433	287	139	45	2	-
2002.....	29	59	90	154	236	416	403	320	148	40	5	-
2001.....	36	54	91	156	250	396	407	280	124	37	5	-
2000.....	39	45	97	166	253	394	431	265	143	21	3	-
1999.....	26	58	104	157	264	458	392	308	124	37	7	-
Black, both sexes												
2004.....	9	8	13	20	23	33	30	21	6	2	4	-
2003.....	3	6	10	16	22	29	38	22	4	6	1	-
2002.....	6	11	11	16	25	32	31	19	6	1	1	-
2001.....	9	13	10	13	24	23	24	22	12	4	1	-
2000.....	7	9	13	15	18	26	21	12	10	2	1	-
1999.....	3	12	8	17	15	36	19	14	3	-	-	-
Black, male												
2004.....	2	1	3	9	6	8	3	5	-	-	-	-
2003.....	-	3	4	2	8	9	8	3	2	-	-	-
2002.....	-	4	3	9	6	6	8	2	1	-	-	-
2001.....	3	2	3	3	7	6	9	4	-	-	-	-
2000.....	1	2	2	5	9	5	5	2	1	-	-	-
1999.....	1	4	1	7	4	14	2	2	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatoid arthritis and related inflammatory polyarthropathies (M05-M08)												
Black, female												
2004.....	7	7	10	11	17	25	27	16	6	2	4	-
2003.....	3	3	6	14	14	20	30	19	2	6	1	-
2002.....	6	7	8	7	19	26	23	17	5	1	1	-
2001.....	6	11	7	10	17	17	15	18	12	4	1	-
2000.....	6	7	11	10	9	21	16	10	9	2	1	-
1999.....	2	8	7	10	11	22	17	12	3	-	-	-
American Indian 1/, both sexes												
2004.....	2	4	1	2	3	2	5	1	2	-	-	-
2003.....	1	2	1	4	3	5	3	3	-	1	-	-
2002.....	3	2	5	4	4	4	2	-	1	-	-	-
2001.....	1	-	3	1	3	2	3	3	1	-	-	-
2000.....	-	1	3	4	2	4	2	-	1	-	-	-
1999.....	2	2	1	3	1	4	2	1	-	-	-	-
American Indian 1/, male												
2004.....	-	2	-	-	1	-	2	-	-	-	-	-
2003.....	-	2	-	1	1	1	1	1	-	-	-	-
2002.....	1	-	1	2	-	1	-	-	1	-	-	-
2001.....	-	-	1	1	1	1	-	1	-	-	-	-
2000.....	-	-	1	-	1	3	-	-	-	-	-	-
1999.....	2	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, female												
2004.....	2	2	1	2	2	2	3	1	2	-	-	-
2003.....	1	-	1	3	2	4	2	2	-	1	-	-
2002.....	2	2	4	2	4	3	2	-	-	-	-	-
2001.....	1	-	2	-	2	1	3	2	1	-	-	-
2000.....	-	1	2	4	1	1	2	-	1	-	-	-
1999.....	-	2	1	3	-	4	2	1	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatoid arthritis and related inflammatory polyarthropathies (M05-M08)												
Asian or Pacific Islander 2/, both sexes												
2004.....	1	-	1	5	4	10	10	3	4	-	1	-
2003.....	-	-	3	3	3	6	11	7	1	-	-	-
2002.....	-	3	6	3	2	10	8	10	1	-	-	-
2001.....	1	1	6	1	3	11	4	3	1	1	-	-
2000.....	-	3	-	2	6	9	7	4	1	2	-	-
1999.....	1	-	2	-	3	4	8	1	-	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	1	-	-	-	-	-	-	3	-	-	-	-
2003.....	-	-	-	-	-	4	1	1	-	-	-	-
2002.....	-	1	1	2	-	1	3	2	-	-	-	-
2001.....	-	-	1	-	1	1	2	-	-	1	-	-
2000.....	-	-	-	-	-	2	1	-	-	1	-	-
1999.....	-	-	1	-	-	-	2	-	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	-	-	1	5	4	10	10	-	4	-	1	-
2003.....	-	-	3	3	3	2	10	6	1	-	-	-
2002.....	-	2	5	1	2	9	5	8	1	-	-	-
2001.....	1	1	5	1	2	10	2	3	1	-	-	-
2000.....	-	3	-	2	6	7	6	4	1	1	-	-
1999.....	1	-	1	-	3	4	6	1	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Systemic lupus erythematosus (M32)												
All races, both sexes												
2004.....	148	131	102	72	82	87	83	39	10	4	1	-
2003.....	137	117	112	79	81	104	79	36	11	3	-	-
2002.....	104	104	112	84	91	93	75	43	18	-	1	-
2001.....	126	97	96	85	104	107	84	36	18	1	-	-
2000.....	117	111	104	93	76	106	76	37	11	3	-	-
1999.....	117	92	87	98	108	98	64	40	15	3	2	-
All races, male												
2004.....	28	17	17	9	18	14	14	8	2	1	-	-
2003.....	20	13	19	18	14	9	10	8	1	-	-	-
2002.....	17	11	11	12	18	18	11	3	3	-	-	-
2001.....	21	13	11	10	18	12	19	12	3	-	-	-
2000.....	11	19	18	15	18	17	15	5	2	1	-	-
1999.....	17	13	11	11	17	18	13	7	4	1	-	-
All races, female												
2004.....	120	114	85	63	64	73	69	31	8	3	1	-
2003.....	117	104	93	61	67	95	69	28	10	3	-	-
2002.....	87	93	101	72	73	75	64	40	15	-	1	-
2001.....	105	84	85	75	86	95	65	24	15	1	-	-
2000.....	106	92	86	78	58	89	61	32	9	2	-	-
1999.....	100	79	76	87	91	80	51	33	11	2	2	-
White, both sexes												
2004.....	84	84	62	50	63	71	71	37	10	3	1	-
2003.....	82	74	87	57	64	93	71	35	9	3	-	-
2002.....	66	70	74	54	73	83	66	40	16	-	1	-
2001.....	79	68	64	66	87	93	77	30	16	1	-	-
2000.....	77	72	76	72	61	92	71	37	10	3	-	-
1999.....	71	63	67	67	87	89	59	39	12	3	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Systemic lupus erythematosus (M32)												
White, male												
2004.....	17	12	13	5	15	12	11	8	2	-	-	-
2003.....	16	7	15	15	13	8	10	8	1	-	-	-
2002.....	12	11	9	9	16	17	9	3	3	-	-	-
2001.....	14	8	6	9	16	11	18	12	3	-	-	-
2000.....	8	12	15	13	16	16	13	5	2	1	-	-
1999.....	13	9	9	8	15	17	13	7	2	1	-	-
White, female												
2004.....	67	72	49	45	48	59	60	29	8	3	1	-
2003.....	66	67	72	42	51	85	61	27	8	3	-	-
2002.....	54	59	65	45	57	66	57	37	13	-	1	-
2001.....	65	60	58	57	71	82	59	18	13	1	-	-
2000.....	69	60	61	59	45	76	58	32	8	2	-	-
1999.....	58	54	58	59	72	72	46	32	10	2	1	-
Black, both sexes												
2004.....	59	44	35	19	16	14	9	2	-	1	-	-
2003.....	49	35	23	20	15	9	5	1	2	-	-	-
2002.....	36	29	35	27	15	10	8	2	2	-	-	-
2001.....	41	26	27	16	12	12	5	5	2	-	-	-
2000.....	37	34	25	19	11	10	5	-	1	-	-	-
1999.....	34	27	17	28	19	7	3	1	2	-	1	-
Black, male												
2004.....	11	5	4	3	2	2	2	-	-	1	-	-
2003.....	4	4	4	2	-	1	-	-	-	-	-	-
2002.....	5	-	2	3	2	1	1	-	-	-	-	-
2001.....	6	5	5	1	2	1	1	-	-	-	-	-
2000.....	3	7	2	2	1	1	2	-	-	-	-	-
1999.....	2	4	1	3	2	-	-	-	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Systemic lupus erythematosus (M32)												
Black, female												
2004.....	48	39	31	16	14	12	7	2	-	-	-	-
2003.....	45	31	19	18	15	8	5	1	2	-	-	-
2002.....	31	29	33	24	13	9	7	2	2	-	-	-
2001.....	35	21	22	15	10	11	4	5	2	-	-	-
2000.....	34	27	23	17	10	9	3	-	1	-	-	-
1999.....	32	23	16	25	17	7	3	1	1	-	1	-
American Indian 1/, both sexes												
2004.....	2	-	2	2	2	-	2	-	-	-	-	-
2003.....	3	3	-	-	-	1	2	-	-	-	-	-
2002.....	1	1	1	2	2	-	-	-	-	-	-	-
2001.....	1	1	1	1	3	-	2	-	-	-	-	-
2000.....	1	1	-	1	1	1	-	-	-	-	-	-
1999.....	3	1	1	-	-	1	1	-	-	-	-	-
American Indian 1/, male												
2004.....	-	-	-	1	-	-	1	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, female												
2004.....	2	-	2	1	2	-	1	-	-	-	-	-
2003.....	3	3	-	-	-	1	2	-	-	-	-	-
2002.....	1	1	1	2	2	-	-	-	-	-	-	-
2001.....	1	1	1	1	3	-	2	-	-	-	-	-
2000.....	1	1	-	1	1	1	-	-	-	-	-	-
1999.....	3	1	1	-	-	1	1	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Systemic lupus erythematosus (M32)												
Asian or Pacific Islander 2/, both sexes												
2004.....	3	3	3	1	1	2	1	-	-	-	-	-
2003.....	3	5	2	2	2	1	1	-	-	-	-	-
2002.....	1	4	2	1	1	-	1	1	-	-	-	-
2001.....	5	2	4	2	2	2	-	1	-	-	-	-
2000.....	2	4	3	1	3	3	-	-	-	-	-	-
1999.....	9	1	2	3	2	1	1	-	1	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	-	-	-	-	1	-	-	-	-	-	-	-
2003.....	-	2	-	1	1	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	1	-	-	-	-	-
2001.....	1	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	1	-	1	-	-	-	-	-	-	-
1999.....	2	-	1	-	-	1	-	-	1	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	3	3	3	1	-	2	1	-	-	-	-	-
2003.....	3	3	2	1	1	1	1	-	-	-	-	-
2002.....	1	4	2	1	1	-	-	1	-	-	-	-
2001.....	4	2	4	2	2	2	-	1	-	-	-	-
2000.....	2	4	2	1	2	3	-	-	-	-	-	-
1999.....	7	1	1	3	2	-	1	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other arthropathies and related disorders (M00,M02,M10-M31,M33-M35)												
All races, both sexes												
2004.....	262	307	327	374	429	567	671	538	401	207	47	-
2003.....	248	285	341	367	478	617	590	553	399	202	68	-
2002.....	263	258	334	376	487	602	633	565	432	190	63	1
2001.....	234	238	297	390	520	607	669	570	413	182	54	-
2000.....	227	245	299	385	530	644	649	528	429	177	54	-
1999.....	251	234	313	398	530	579	579	525	400	169	53	-
All races, male												
2004.....	97	94	118	114	128	178	200	140	85	36	3	-
2003.....	81	97	102	131	159	224	173	154	92	28	6	-
2002.....	82	91	113	112	161	195	192	136	90	25	7	1
2001.....	90	76	104	103	178	198	204	146	74	39	7	-
2000.....	70	94	98	122	157	223	205	133	89	33	4	-
1999.....	87	79	99	122	171	196	180	138	88	33	9	-
All races, female												
2004.....	165	213	209	260	301	389	471	398	316	171	44	-
2003.....	167	188	239	236	319	393	417	399	307	174	62	-
2002.....	181	167	221	264	326	407	441	429	342	165	56	-
2001.....	144	162	193	287	342	409	465	424	339	143	47	-
2000.....	157	151	201	263	373	421	444	395	340	144	50	-
1999.....	164	155	214	276	359	383	399	387	312	136	44	-
White, both sexes												
2004.....	182	223	268	312	377	509	626	507	378	190	43	-
2003.....	166	216	271	307	410	553	539	521	361	181	59	-
2002.....	190	199	279	326	422	549	581	520	404	181	59	-
2001.....	156	190	240	324	455	562	623	538	388	170	43	-
2000.....	154	184	240	336	459	588	598	497	390	166	52	-
1999.....	163	177	241	339	467	530	542	496	376	154	44	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other arthropathies and related disorders (M00,M02,M10-M31,M33-M35)												
White, male												
2004.....	69	70	97	97	112	162	187	132	80	31	2	-
2003.....	57	71	82	105	134	196	152	147	82	25	6	-
2002.....	60	66	89	95	137	180	176	128	82	23	7	-
2001.....	60	59	89	84	156	183	190	141	67	36	4	-
2000.....	49	73	75	104	133	199	185	114	78	32	4	-
1999.....	63	58	78	105	152	174	171	131	82	31	8	-
White, female												
2004.....	113	153	171	215	265	347	439	375	298	159	41	-
2003.....	109	145	189	202	276	357	387	374	279	156	53	-
2002.....	130	133	190	231	285	369	405	392	322	158	52	-
2001.....	96	131	151	240	299	379	433	397	321	134	39	-
2000.....	105	111	165	232	326	389	413	383	312	134	48	-
1999.....	100	119	163	234	315	356	371	365	294	123	36	-
Black, both sexes												
2004.....	74	63	51	51	41	41	36	21	20	14	4	-
2003.....	77	57	60	50	52	51	39	17	29	17	7	-
2002.....	64	53	47	41	52	43	41	37	25	7	4	1
2001.....	73	39	41	56	45	37	43	31	22	9	11	-
2000.....	61	48	48	36	59	42	43	26	34	9	1	-
1999.....	73	49	63	48	52	41	30	23	19	14	8	-
Black, male												
2004.....	25	18	19	14	14	11	10	6	5	2	1	-
2003.....	24	22	19	23	19	23	14	4	6	2	-	-
2002.....	21	24	21	15	19	11	11	7	5	1	-	1
2001.....	27	13	13	17	13	13	12	4	6	2	3	-
2000.....	16	16	20	15	19	17	18	16	8	-	-	-
1999.....	20	18	19	15	16	20	6	6	4	2	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other arthropathies and related disorders (M00,M02,M10-M31,M33-M35)												
Black, female												
2004.....	49	45	32	37	27	30	26	15	15	12	3	-
2003.....	53	35	41	27	33	28	25	13	23	15	7	-
2002.....	43	29	26	26	33	32	30	30	20	6	4	-
2001.....	46	26	28	39	32	24	31	27	16	7	8	-
2000.....	45	32	28	21	40	25	25	10	26	9	1	-
1999.....	53	31	44	33	36	21	24	17	15	12	7	-
American Indian 1/, both sexes												
2004.....	3	5	4	3	2	2	3	1	1	-	-	-
2003.....	2	4	5	3	4	1	2	2	-	1	-	-
2002.....	5	1	2	3	-	1	1	3	-	1	-	-
2001.....	2	3	5	3	5	2	2	1	-	-	-	-
2000.....	4	6	3	3	4	4	4	-	-	-	-	-
1999.....	7	2	1	2	3	2	1	3	2	-	-	-
American Indian 1/, male												
2004.....	2	3	1	2	1	-	-	-	-	-	-	-
2003.....	-	-	1	-	1	1	1	-	-	-	-	-
2002.....	1	-	-	2	-	1	-	-	-	-	-	-
2001.....	1	1	-	-	3	-	1	1	-	-	-	-
2000.....	1	2	1	1	1	1	2	-	-	-	-	-
1999.....	2	1	1	-	-	1	-	-	-	-	-	-
American Indian 1/, female												
2004.....	1	2	3	1	1	2	3	1	1	-	-	-
2003.....	2	4	4	3	3	-	1	2	-	1	-	-
2002.....	4	1	2	1	-	-	1	3	-	1	-	-
2001.....	1	2	5	3	2	2	1	-	-	-	-	-
2000.....	3	4	2	2	3	3	2	-	-	-	-	-
1999.....	5	1	-	2	3	1	1	3	2	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other arthropathies and related disorders (M00,M02,M10-M31,M33-M35)												
Asian or Pacific Islander 2/, both sexes												
2004.....	3	16	4	8	9	15	6	9	2	3	-	-
2003.....	3	8	5	7	12	12	10	13	9	3	2	-
2002.....	4	5	6	6	13	9	10	5	3	1	-	-
2001.....	3	6	11	7	15	6	1	-	3	3	-	-
2000.....	8	7	8	10	8	10	4	5	5	2	1	-
1999.....	8	6	8	9	8	6	6	3	3	1	1	-
Asian or Pacific Islander 2/, male												
2004.....	1	3	1	1	1	5	3	2	-	3	-	-
2003.....	-	4	-	3	5	4	6	3	4	1	-	-
2002.....	-	1	3	-	5	3	5	1	3	1	-	-
2001.....	2	3	2	2	6	2	1	-	1	1	-	-
2000.....	4	3	2	2	4	6	-	3	3	1	-	-
1999.....	2	2	1	2	3	1	3	1	2	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	2	13	3	7	8	10	3	7	2	-	-	-
2003.....	3	4	5	4	7	8	4	10	5	2	2	-
2002.....	4	4	3	6	8	6	5	4	-	-	-	-
2001.....	1	3	9	5	9	4	-	-	2	2	-	-
2000.....	4	4	6	8	4	4	4	2	2	1	1	-
1999.....	6	4	7	7	5	5	3	2	1	1	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Dorsopathies (M40-M54)												
All races, both sexes												
2004.....	37	47	61	55	103	145	188	183	134	22	6	-
2003.....	41	48	55	68	97	161	183	153	76	32	6	-
2002.....	33	48	46	62	102	127	154	157	94	21	3	-
2001.....	33	37	37	51	88	129	169	168	133	24	-	-
2000.....	23	29	41	63	78	127	164	141	85	27	7	-
1999.....	34	26	37	56	83	111	124	127	73	18	6	-
All races, male												
2004.....	20	23	37	27	49	70	71	57	36	4	2	-
2003.....	31	27	25	40	42	73	73	46	16	6	1	-
2002.....	23	22	24	24	50	60	63	54	16	6	-	-
2001.....	19	20	20	26	36	57	67	47	32	4	-	-
2000.....	16	14	23	35	45	46	69	46	28	5	1	-
1999.....	13	10	23	31	39	55	51	41	22	6	1	-
All races, female												
2004.....	17	24	24	28	54	75	117	126	98	18	4	-
2003.....	10	21	30	28	55	88	110	107	60	26	5	-
2002.....	10	26	22	38	52	67	91	103	78	15	3	-
2001.....	14	17	17	25	52	72	102	121	101	20	-	-
2000.....	7	15	18	28	33	81	95	95	57	22	6	-
1999.....	21	16	14	25	44	56	73	86	51	12	5	-
White, both sexes												
2004.....	29	41	51	41	89	135	177	171	128	20	6	-
2003.....	33	39	42	58	74	139	170	142	72	31	5	-
2002.....	27	39	36	54	91	112	145	150	87	20	2	-
2001.....	29	27	29	41	76	111	154	158	128	24	-	-
2000.....	19	25	38	55	66	114	149	136	79	24	6	-
1999.....	31	21	30	45	74	98	118	120	65	17	6	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Dorsopathies (M40-M54)												
White, male												
2004.....	16	19	29	20	40	62	64	54	34	3	2	-
2003.....	24	19	18	33	31	64	67	41	15	6	1	-
2002.....	20	16	18	19	45	51	59	49	12	6	-	-
2001.....	16	14	15	21	31	50	64	44	30	4	-	-
2000.....	15	11	21	28	37	39	64	45	27	4	1	-
1999.....	12	9	17	27	34	47	49	38	18	6	1	-
White, female												
2004.....	13	22	22	21	49	73	113	117	94	17	4	-
2003.....	9	20	24	25	43	75	103	101	57	25	4	-
2002.....	7	23	18	35	46	61	86	101	75	14	2	-
2001.....	13	13	14	20	45	61	90	114	98	20	-	-
2000.....	4	14	17	27	29	75	85	91	52	20	5	-
1999.....	19	12	13	18	40	51	69	82	47	11	5	-
Black, both sexes												
2004.....	7	4	9	11	13	8	7	7	6	1	-	-
2003.....	7	7	12	8	19	20	9	7	3	1	1	-
2002.....	6	8	9	6	6	12	7	5	6	-	-	-
2001.....	4	9	7	9	10	16	10	8	4	-	-	-
2000.....	3	4	3	8	12	13	10	4	4	2	1	-
1999.....	3	4	5	10	9	13	3	5	6	-	-	-
Black, male												
2004.....	3	2	8	6	9	7	3	1	2	1	-	-
2003.....	6	6	6	6	11	9	4	3	1	-	-	-
2002.....	3	6	5	3	2	8	2	3	4	-	-	-
2001.....	3	5	4	4	5	6	2	2	1	-	-	-
2000.....	1	3	2	7	8	7	2	1	-	1	-	-
1999.....	1	-	4	4	5	8	1	2	3	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Dorsopathies (M40-M54)												
Black, female												
2004.....	4	2	1	5	4	1	4	6	4	-	-	-
2003.....	1	1	6	2	8	11	5	4	2	1	1	-
2002.....	3	2	4	3	4	4	5	2	2	-	-	-
2001.....	1	4	3	5	5	10	8	6	3	-	-	-
2000.....	2	1	1	1	4	6	8	3	4	1	1	-
1999.....	2	4	1	6	4	5	2	3	3	-	-	-
American Indian 1/, both sexes												
2004.....	1	1	-	1	-	1	1	1	-	-	-	-
2003.....	-	-	1	1	2	-	1	-	-	-	-	-
2002.....	-	-	1	2	-	-	-	-	1	1	-	-
2001.....	-	-	1	1	1	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	2	1	-	-	-	-
1999.....	-	1	1	-	-	-	-	1	1	1	-	-
American Indian 1/, male												
2004.....	1	1	-	-	-	1	1	-	-	-	-	-
2003.....	-	-	1	-	-	-	-	-	-	-	-	-
2002.....	-	-	1	2	-	-	-	-	-	-	-	-
2001.....	-	-	1	1	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	1	-	-	-	-	-
1999.....	-	1	1	-	-	-	-	-	1	-	-	-
American Indian 1/, female												
2004.....	-	-	-	1	-	-	-	1	-	-	-	-
2003.....	-	-	-	1	2	-	1	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	1	1	-	-
2001.....	-	-	-	-	1	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	1	1	-	-	-	-
1999.....	-	-	-	-	-	-	-	1	-	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Dorsopathies (M40-M54)												
Asian or Pacific Islander 2/, both sexes												
2004.....	-	1	1	2	1	1	3	4	-	1	-	-
2003.....	1	2	-	1	2	2	3	4	1	-	-	-
2002.....	-	1	-	-	5	3	2	2	-	-	1	-
2001.....	-	1	-	-	1	2	5	2	1	-	-	-
2000.....	1	-	-	-	-	-	3	-	2	1	-	-
1999.....	-	-	1	1	-	-	3	1	1	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	-	1	-	1	-	-	3	2	-	-	-	-
2003.....	1	2	-	1	-	-	2	2	-	-	-	-
2002.....	-	-	-	-	3	1	2	2	-	-	-	-
2001.....	-	1	-	-	-	1	1	1	1	-	-	-
2000.....	-	-	-	-	-	-	2	-	1	-	-	-
1999.....	-	-	1	-	-	-	1	1	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	-	-	1	1	1	1	-	2	-	1	-	-
2003.....	-	-	-	-	2	2	1	2	1	-	-	-
2002.....	-	1	-	-	2	2	-	-	-	-	1	-
2001.....	-	-	-	-	1	1	4	1	-	-	-	-
2000.....	1	-	-	-	-	-	1	-	1	1	-	-
1999.....	-	-	-	1	-	-	2	-	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Soft tissue disorders (M60-M79)												
All races, both sexes												
2004.....	88	94	89	93	126	156	189	144	75	27	3	-
2003.....	98	82	91	103	129	154	178	139	85	25	5	-
2002.....	73	73	73	91	113	140	167	133	80	29	2	-
2001.....	82	72	74	83	108	146	145	107	60	19	4	-
2000.....	68	70	61	80	108	149	116	97	64	11	6	-
1999.....	53	66	55	77	103	127	133	105	57	12	4	-
All races, male												
2004.....	55	54	49	44	59	82	95	74	30	11	2	-
2003.....	49	32	43	50	73	81	93	62	32	7	2	-
2002.....	40	30	42	58	60	60	87	66	31	11	1	-
2001.....	50	38	31	50	63	84	76	48	23	7	2	-
2000.....	29	39	28	40	61	78	64	42	34	4	2	-
1999.....	28	37	27	37	67	75	72	50	26	5	-	-
All races, female												
2004.....	33	40	40	49	67	74	94	70	45	16	1	-
2003.....	49	50	48	53	56	73	85	77	53	18	3	-
2002.....	33	43	31	33	53	80	80	67	49	18	1	-
2001.....	32	34	43	33	45	62	69	59	37	12	2	-
2000.....	39	31	33	40	47	71	52	55	30	7	4	-
1999.....	25	29	28	40	36	52	61	55	31	7	4	-
White, both sexes												
2004.....	62	64	69	71	109	134	167	121	66	21	2	-
2003.....	73	69	65	84	109	128	158	125	77	23	4	-
2002.....	56	55	58	75	90	122	148	111	68	28	2	-
2001.....	61	55	47	64	86	117	127	93	52	18	3	-
2000.....	51	52	43	66	85	115	101	85	55	9	4	-
1999.....	41	43	38	58	87	109	116	87	47	12	4	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Soft tissue disorders (M60-M79)												
White, male												
2004.....	36	38	39	36	47	71	86	61	26	9	2	-
2003.....	37	28	33	37	64	69	85	58	28	6	1	-
2002.....	33	23	35	47	47	52	77	57	27	11	1	-
2001.....	39	28	19	40	52	75	66	43	20	6	1	-
2000.....	20	35	23	35	46	65	59	38	28	3	1	-
1999.....	21	27	19	29	58	62	61	40	23	5	-	-
White, female												
2004.....	26	26	30	35	62	63	81	60	40	12	-	-
2003.....	36	41	32	47	45	59	73	67	49	17	3	-
2002.....	23	32	23	28	43	70	71	54	41	17	1	-
2001.....	22	27	28	24	34	42	61	50	32	12	2	-
2000.....	31	17	20	31	39	50	42	47	27	6	3	-
1999.....	20	16	19	29	29	47	55	47	24	7	4	-
Black, both sexes												
2004.....	24	26	16	21	15	20	18	18	8	6	1	-
2003.....	20	11	19	15	20	22	19	14	8	1	1	-
2002.....	16	15	13	12	20	15	15	19	11	1	-	-
2001.....	17	16	25	17	20	27	16	10	7	-	1	-
2000.....	15	15	16	14	19	30	15	11	8	2	2	-
1999.....	12	17	14	16	16	16	14	16	10	-	-	-
Black, male												
2004.....	17	14	7	7	11	10	8	10	3	2	-	-
2003.....	10	3	8	11	9	9	7	4	4	-	1	-
2002.....	6	7	6	8	11	7	9	6	4	-	-	-
2001.....	11	9	10	9	10	9	10	4	2	-	1	-
2000.....	8	4	5	5	13	11	5	4	5	1	1	-
1999.....	7	7	6	7	9	11	10	9	3	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Soft tissue disorders (M60-M79)												
Asian or Pacific Islander 2/, both sexes												
2004.....	2	2	2	-	2	2	3	3	-	-	-	-
2003.....	2	1	5	3	-	2	1	-	-	1	-	-
2002.....	1	1	2	2	3	2	1	3	1	-	-	-
2001.....	3	1	2	1	2	-	1	2	-	-	-	-
2000.....	1	1	-	-	3	3	-	1	1	-	-	-
1999.....	-	2	3	2	-	2	3	2	-	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	2	2	1	-	1	1	1	2	-	-	-	-
2003.....	-	1	1	1	-	1	1	-	-	1	-	-
2002.....	1	-	1	2	2	1	1	3	-	-	-	-
2001.....	-	1	2	1	1	-	-	1	-	-	-	-
2000.....	1	-	-	-	2	2	-	-	1	-	-	-
1999.....	-	2	2	1	-	2	1	1	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	-	-	1	-	1	1	2	1	-	-	-	-
2003.....	2	-	4	2	-	1	-	-	-	-	-	-
2002.....	-	1	1	-	1	1	-	-	1	-	-	-
2001.....	3	-	-	-	1	-	1	1	-	-	-	-
2000.....	-	1	-	-	1	1	-	1	-	-	-	-
1999.....	-	-	1	1	-	-	2	1	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Osteopathies, chondropathies and other disorders of musculoskeletal system and connective tissue (M80-M99)												
All races, both sexes												
2004.....	45	68	102	130	190	345	609	739	660	285	75	-
2003.....	50	78	90	129	217	323	571	767	688	319	88	-
2002.....	43	62	92	117	228	350	542	766	697	329	82	-
2001.....	35	49	79	123	191	330	558	740	655	311	58	-
2000.....	30	45	60	103	163	338	484	727	660	295	80	-
1999.....	30	48	55	94	158	321	504	644	615	278	65	-
All races, male												
2004.....	34	33	51	66	90	133	180	167	103	30	2	-
2003.....	27	51	44	82	94	116	154	167	103	32	9	-
2002.....	29	28	51	62	112	125	145	166	114	41	7	-
2001.....	22	31	38	58	81	121	148	159	106	37	4	-
2000.....	16	27	32	55	69	129	131	168	103	37	6	-
1999.....	13	22	24	34	71	112	141	137	79	22	5	-
All races, female												
2004.....	11	35	51	64	100	212	429	572	557	255	73	-
2003.....	23	27	46	47	123	207	417	600	585	287	79	-
2002.....	14	34	41	55	116	225	397	600	583	288	75	-
2001.....	13	18	41	65	110	209	410	581	549	274	54	-
2000.....	14	18	28	48	94	209	353	559	557	258	74	-
1999.....	17	26	31	60	87	209	363	507	536	256	60	-
White, both sexes												
2004.....	34	54	86	111	159	297	556	681	628	275	68	-
2003.....	40	60	73	100	186	289	516	719	645	300	81	-
2002.....	33	45	75	88	204	302	486	723	671	309	72	-
2001.....	29	37	63	101	151	296	514	685	613	293	54	-
2000.....	23	35	43	82	138	306	437	677	625	283	66	-
1999.....	24	36	49	74	137	277	475	605	588	258	62	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Osteopathies, chondropathies and other disorders of musculoskeletal system and connective tissue (M80-M99)												
White, male												
2004.....	26	28	42	56	72	108	161	151	98	27	2	-
2003.....	22	41	32	61	74	98	133	148	92	27	6	-
2002.....	22	24	41	47	96	106	122	151	109	34	5	-
2001.....	19	21	28	45	64	105	134	146	95	35	4	-
2000.....	12	21	22	43	52	114	113	147	94	36	5	-
1999.....	10	14	23	23	64	89	135	125	74	15	4	-
White, female												
2004.....	8	26	44	55	87	189	395	530	530	248	66	-
2003.....	18	19	41	39	112	191	383	571	553	273	75	-
2002.....	11	21	34	41	108	196	364	572	562	275	67	-
2001.....	10	16	35	56	87	191	380	539	518	258	50	-
2000.....	11	14	21	39	86	192	324	530	531	247	61	-
1999.....	14	22	26	51	73	188	340	480	514	243	58	-
Black, both sexes												
2004.....	10	13	13	18	27	44	47	46	27	9	6	-
2003.....	10	18	16	26	25	29	43	36	37	15	3	-
2002.....	9	16	16	25	22	37	47	36	21	17	7	-
2001.....	6	10	15	20	35	29	35	45	33	15	4	-
2000.....	6	10	16	20	21	30	39	43	27	7	11	-
1999.....	5	8	6	19	18	38	25	29	22	15	1	-
Black, male												
2004.....	8	4	8	9	15	23	17	14	5	3	-	-
2003.....	5	10	11	20	17	16	19	13	9	4	-	-
2002.....	6	4	9	12	16	14	21	14	3	5	2	-
2001.....	3	8	10	12	14	14	12	12	9	2	-	-
2000.....	3	6	9	12	15	13	15	20	6	1	1	-
1999.....	3	5	1	11	6	18	5	11	4	6	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Osteopathies, chondropathies and other disorders of musculoskeletal system and connective tissue (M80-M99)												
Black, female												
2004.....	2	9	5	9	12	21	30	32	22	6	6	-
2003.....	5	8	5	6	8	13	24	23	28	11	3	-
2002.....	3	12	7	13	6	23	26	22	18	12	5	-
2001.....	3	2	5	8	21	15	23	33	24	13	4	-
2000.....	3	4	7	8	6	17	24	23	21	6	10	-
1999.....	2	3	5	8	12	20	20	18	18	9	1	-
American Indian 1/, both sexes												
2004.....	-	1	2	-	2	1	-	1	-	1	-	-
2003.....	-	-	1	2	3	2	3	5	2	1	1	-
2002.....	1	1	1	-	-	3	2	-	1	-	-	-
2001.....	-	-	1	1	3	3	2	4	1	-	-	-
2000.....	-	-	-	-	1	-	3	2	2	1	-	-
1999.....	-	4	-	1	3	1	-	1	-	2	1	-
American Indian 1/, male												
2004.....	-	1	1	-	1	1	-	-	-	-	-	-
2003.....	-	-	1	1	2	1	-	2	1	-	1	-
2002.....	1	-	1	-	-	2	-	-	-	-	-	-
2001.....	-	-	-	1	2	1	-	1	1	-	-	-
2000.....	-	-	-	-	1	-	2	1	1	-	-	-
1999.....	-	3	-	-	1	1	-	-	-	1	1	-
American Indian 1/, female												
2004.....	-	-	1	-	1	-	-	1	-	1	-	-
2003.....	-	-	-	1	1	1	3	3	1	1	-	-
2002.....	-	1	-	-	-	1	2	-	1	-	-	-
2001.....	-	-	1	-	1	2	2	3	-	-	-	-
2000.....	-	-	-	-	-	-	1	1	1	1	-	-
1999.....	-	1	-	1	2	-	-	1	-	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Osteopathies, chondropathies and other disorders of musculoskeletal system and connective tissue (M80-M99)												
Asian or Pacific Islander 2/, both sexes												
2004.....	1	-	1	1	2	3	6	11	5	-	1	-
2003.....	-	-	-	1	3	3	9	7	4	3	3	-
2002.....	-	-	-	4	2	8	7	7	4	3	3	-
2001.....	-	2	-	1	2	2	7	6	8	3	-	-
2000.....	1	-	1	1	3	2	5	5	6	4	3	-
1999.....	1	-	-	-	-	5	4	9	5	3	1	-
Asian or Pacific Islander 2/, male												
2004.....	-	-	-	1	2	1	2	2	-	-	-	-
2003.....	-	-	-	-	1	1	2	4	1	1	2	-
2002.....	-	-	-	3	-	3	2	1	2	2	-	-
2001.....	-	2	-	-	1	1	2	-	1	-	-	-
2000.....	1	-	1	-	1	2	1	-	2	-	-	-
1999.....	-	-	-	-	-	4	1	1	1	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	1	-	1	-	-	2	4	9	5	-	1	-
2003.....	-	-	-	1	2	2	7	3	3	2	1	-
2002.....	-	-	-	1	2	5	5	6	2	1	3	-
2001.....	-	-	-	1	1	1	5	6	7	3	-	-
2000.....	-	-	-	1	2	-	4	5	4	4	3	-
1999.....	1	-	-	-	-	1	3	8	4	3	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Osteoporosis (M80-M81)												
All races, both sexes												
2004.....	2	9	11	15	47	125	293	440	468	223	55	-
2003.....	4	6	6	14	63	123	295	482	488	239	57	-
2002.....	4	6	9	21	56	144	287	516	512	243	59	-
2001.....	2	7	12	28	40	125	286	460	458	225	37	-
2000.....	3	2	7	27	38	153	273	471	452	220	64	-
1999.....	1	5	8	28	42	147	281	427	443	204	50	-
All races, male												
2004.....	2	5	4	4	11	22	40	52	42	13	1	-
2003.....	3	2	1	4	12	14	38	43	49	20	2	-
2002.....	2	-	3	4	10	20	35	53	62	20	3	-
2001.....	1	2	3	5	11	19	30	51	44	24	2	-
2000.....	2	1	-	4	7	27	35	58	42	17	5	-
1999.....	1	1	-	5	6	22	39	43	28	9	4	-
All races, female												
2004.....	-	4	7	11	36	103	253	388	426	210	54	-
2003.....	1	4	5	10	51	109	257	439	439	219	55	-
2002.....	2	6	6	17	46	124	252	463	450	223	56	-
2001.....	1	5	9	23	29	106	256	409	414	201	35	-
2000.....	1	1	7	23	31	126	238	413	410	203	59	-
1999.....	-	4	8	23	36	125	242	384	415	195	46	-
White, both sexes												
2004.....	2	8	10	15	46	122	284	425	457	220	52	-
2003.....	3	6	6	13	62	118	283	475	476	233	55	-
2002.....	4	6	9	19	54	134	277	506	504	236	53	-
2001.....	2	6	11	27	37	121	279	447	444	219	37	-
2000.....	3	2	5	27	34	150	265	464	443	213	57	-
1999.....	1	5	7	27	42	143	275	416	435	196	48	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Osteoporosis (M80-M81)												
White, male												
2004.....	2	5	3	4	11	21	37	51	42	12	1	-
2003.....	2	2	1	4	12	14	35	42	45	19	1	-
2002.....	2	-	3	4	10	18	34	52	61	18	2	-
2001.....	1	1	3	5	11	19	30	51	42	23	2	-
2000.....	2	1	-	4	6	24	35	56	38	16	5	-
1999.....	1	1	-	4	6	21	38	43	28	6	3	-
White, female												
2004.....	-	3	7	11	35	101	247	374	415	208	51	-
2003.....	1	4	5	9	50	104	248	433	431	214	54	-
2002.....	2	6	6	15	44	116	243	454	443	218	51	-
2001.....	1	5	8	22	26	102	249	396	402	196	35	-
2000.....	1	1	5	23	28	126	230	408	405	197	52	-
1999.....	-	4	7	23	36	122	237	373	407	190	45	-
Black, both sexes												
2004.....	-	1	-	-	1	-	5	6	6	2	3	-
2003.....	1	-	-	-	-	3	2	1	6	4	-	-
2002.....	-	-	-	1	1	5	5	3	5	4	3	-
2001.....	-	1	-	1	1	3	1	6	8	4	-	-
2000.....	-	-	2	-	2	2	3	2	3	3	4	-
1999.....	-	-	1	1	-	4	3	5	4	3	-	-
Black, male												
2004.....	-	-	-	-	-	-	2	-	-	1	-	-
2003.....	1	-	-	-	-	-	2	-	2	1	-	-
2002.....	-	-	-	-	-	1	-	-	-	-	1	-
2001.....	-	1	-	-	-	-	-	-	1	1	-	-
2000.....	-	-	-	-	1	2	-	2	1	1	-	-
1999.....	-	-	-	1	-	1	-	-	-	2	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Osteoporosis (M80-M81)												
Black, female												
2004.....	-	1	-	-	1	-	3	6	6	1	3	-
2003.....	-	-	-	-	-	3	-	1	4	3	-	-
2002.....	-	-	-	1	1	4	5	3	5	4	2	-
2001.....	-	-	-	1	1	3	1	6	7	3	-	-
2000.....	-	-	2	-	1	-	3	-	2	2	4	-
1999.....	-	-	1	-	-	3	3	5	4	1	-	-
American Indian 1/, both sexes												
2004.....	-	-	1	-	-	1	-	1	-	1	-	-
2003.....	-	-	-	1	1	1	3	3	2	1	-	-
2002.....	-	-	-	-	-	1	-	-	1	-	-	-
2001.....	-	-	1	-	1	1	2	2	-	-	-	-
2000.....	-	-	-	-	-	-	1	1	1	-	-	-
1999.....	-	-	-	-	-	-	-	1	-	2	1	-
American Indian 1/, male												
2004.....	-	-	1	-	-	1	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	1	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	1	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	1	1	-
American Indian 1/, female												
2004.....	-	-	-	-	-	-	-	1	-	1	-	-
2003.....	-	-	-	1	1	1	3	3	1	1	-	-
2002.....	-	-	-	-	-	1	-	-	1	-	-	-
2001.....	-	-	1	-	1	1	2	2	-	-	-	-
2000.....	-	-	-	-	-	-	1	1	-	-	-	-
1999.....	-	-	-	-	-	-	-	1	-	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Osteoporosis (M80-M81)												
Asian or Pacific Islander 2/, both sexes												
2004.....	-	-	-	-	-	2	4	8	5	-	-	-
2003.....	-	-	-	-	-	1	7	3	4	1	2	-
2002.....	-	-	-	1	1	4	5	7	2	3	3	-
2001.....	-	-	-	-	1	-	4	5	6	2	-	-
2000.....	-	-	-	-	2	1	4	4	5	4	3	-
1999.....	-	-	-	-	-	-	3	5	4	3	1	-
Asian or Pacific Islander 2/, male												
2004.....	-	-	-	-	-	-	1	1	-	-	-	-
2003.....	-	-	-	-	-	-	1	1	1	-	1	-
2002.....	-	-	-	-	-	1	1	1	1	2	-	-
2001.....	-	-	-	-	-	-	-	-	1	-	-	-
2000.....	-	-	-	-	-	1	-	-	2	-	-	-
1999.....	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	-	-	-	-	-	2	3	7	5	-	-	-
2003.....	-	-	-	-	-	1	6	2	3	1	1	-
2002.....	-	-	-	1	1	3	4	6	1	1	3	-
2001.....	-	-	-	-	1	-	4	5	5	2	-	-
2000.....	-	-	-	-	2	-	4	4	3	4	3	-
1999.....	-	-	-	-	-	-	2	5	4	3	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other osteopathies, chondropathies and disorders of musculoskeletal system and connective tissue (M83-M99)												
All races, both sexes												
2004.....	43	59	91	115	143	220	316	299	192	62	20	-
2003.....	46	72	84	115	154	200	276	285	200	80	31	-
2002.....	39	56	83	96	172	206	255	250	185	86	23	-
2001.....	33	42	67	95	151	205	272	280	197	86	21	-
2000.....	27	43	53	76	125	185	211	256	208	75	16	-
1999.....	29	43	47	66	116	174	223	217	172	74	15	-
All races, male												
2004.....	32	28	47	62	79	111	140	115	61	17	1	-
2003.....	24	49	43	78	82	102	116	124	54	12	7	-
2002.....	27	28	48	58	102	105	110	113	52	21	4	-
2001.....	21	29	35	53	70	102	118	108	62	13	2	-
2000.....	14	26	32	51	62	102	96	110	61	20	1	-
1999.....	12	21	24	29	65	90	102	94	51	13	1	-
All races, female												
2004.....	11	31	44	53	64	109	176	184	131	45	19	-
2003.....	22	23	41	37	72	98	160	161	146	68	24	-
2002.....	12	28	35	38	70	101	145	137	133	65	19	-
2001.....	12	13	32	42	81	103	154	172	135	73	19	-
2000.....	13	17	21	25	63	83	115	146	147	55	15	-
1999.....	17	22	23	37	51	84	121	123	121	61	14	-
White, both sexes												
2004.....	32	46	76	96	113	175	272	256	171	55	16	-
2003.....	37	54	67	87	124	171	233	244	169	67	26	-
2002.....	29	39	66	69	150	168	209	217	167	73	19	-
2001.....	27	31	52	74	114	175	235	238	169	74	17	-
2000.....	20	33	38	55	104	156	172	213	182	70	9	-
1999.....	23	31	42	47	95	134	200	189	153	62	14	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other osteopathies, chondropathies and disorders of musculoskeletal system and connective tissue (M83-M99)												
White, male												
2004.....	24	23	39	52	61	87	124	100	56	15	1	-
2003.....	20	39	31	57	62	84	98	106	47	8	5	-
2002.....	20	24	38	43	86	88	88	99	48	16	3	-
2001.....	18	20	25	40	53	86	104	95	53	12	2	-
2000.....	10	20	22	39	46	90	78	91	56	20	-	-
1999.....	9	13	23	19	58	68	97	82	46	9	1	-
White, female												
2004.....	8	23	37	44	52	88	148	156	115	40	15	-
2003.....	17	15	36	30	62	87	135	138	122	59	21	-
2002.....	9	15	28	26	64	80	121	118	119	57	16	-
2001.....	9	11	27	34	61	89	131	143	116	62	15	-
2000.....	10	13	16	16	58	66	94	122	126	50	9	-
1999.....	14	18	19	28	37	66	103	107	107	53	13	-
Black, both sexes												
2004.....	10	12	13	18	26	44	42	40	21	7	3	-
2003.....	9	18	16	26	25	26	41	35	31	11	3	-
2002.....	9	16	16	24	21	32	42	33	16	13	4	-
2001.....	6	9	15	19	34	26	34	39	25	11	4	-
2000.....	6	10	14	20	19	28	36	41	24	4	7	-
1999.....	5	8	5	18	18	34	22	24	18	12	1	-
Black, male												
2004.....	8	4	8	9	15	23	15	14	5	2	-	-
2003.....	4	10	11	20	17	16	17	13	7	3	-	-
2002.....	6	4	9	12	16	13	21	14	3	5	1	-
2001.....	3	7	10	12	14	14	12	12	8	1	-	-
2000.....	3	6	9	12	14	11	15	18	5	-	1	-
1999.....	3	5	1	10	6	17	5	11	4	4	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other osteopathies, chondropathies and disorders of musculoskeletal system and connective tissue (M83-M99)												
Black, female												
2004.....	2	8	5	9	11	21	27	26	16	5	3	-
2003.....	5	8	5	6	8	10	24	22	24	8	3	-
2002.....	3	12	7	12	5	19	21	19	13	8	3	-
2001.....	3	2	5	7	20	12	22	27	17	10	4	-
2000.....	3	4	5	8	5	17	21	23	19	4	6	-
1999.....	2	3	4	8	12	17	17	13	14	8	1	-
American Indian 1/, both sexes												
2004.....	-	1	1	-	2	-	-	-	-	-	-	-
2003.....	-	-	1	1	2	1	-	2	-	-	1	-
2002.....	1	1	1	-	-	2	2	-	-	-	-	-
2001.....	-	-	-	1	2	2	-	2	1	-	-	-
2000.....	-	-	-	-	1	-	2	1	1	1	-	-
1999.....	-	4	-	1	3	1	-	-	-	-	-	-
American Indian 1/, male												
2004.....	-	1	-	-	1	-	-	-	-	-	-	-
2003.....	-	-	1	1	2	1	-	2	-	-	1	-
2002.....	1	-	1	-	-	2	-	-	-	-	-	-
2001.....	-	-	-	1	2	1	-	1	1	-	-	-
2000.....	-	-	-	-	1	-	2	1	-	-	-	-
1999.....	-	3	-	-	1	1	-	-	-	-	-	-
American Indian 1/, female												
2004.....	-	-	1	-	1	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	1	-	-	-	-	2	-	-	-	-	-
2001.....	-	-	-	-	-	1	-	1	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	1	1	-	-
1999.....	-	1	-	1	2	-	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other osteopathies, chondropathies and disorders of musculoskeletal system and connective tissue (M83-M99)												
Asian or Pacific Islander 2/, both sexes												
2004.....	1	-	1	1	2	1	2	3	-	-	1	-
2003.....	-	-	-	1	3	2	2	4	-	2	1	-
2002.....	-	-	-	3	1	4	2	-	2	-	-	-
2001.....	-	2	-	1	1	2	3	1	2	1	-	-
2000.....	1	-	1	1	1	1	1	1	1	-	-	-
1999.....	1	-	-	-	-	5	1	4	1	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	-	-	-	1	2	1	1	1	-	-	-	-
2003.....	-	-	-	-	1	1	1	3	-	1	1	-
2002.....	-	-	-	3	-	2	1	-	1	-	-	-
2001.....	-	2	-	-	1	1	2	-	-	-	-	-
2000.....	1	-	1	-	1	1	1	-	-	-	-	-
1999.....	-	-	-	-	-	4	-	1	1	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	1	-	1	-	-	-	1	2	-	-	1	-
2003.....	-	-	-	1	2	1	1	1	-	1	-	-
2002.....	-	-	-	-	1	2	1	-	1	-	-	-
2001.....	-	-	-	1	-	1	1	1	2	1	-	-
2000.....	-	-	-	1	-	-	-	1	1	-	-	-
1999.....	1	-	-	-	-	1	1	3	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XIV. Diseases of the genitourinary system (N00-N98)												
All races, both sexes												
2004.....	1,537	2,138	2,672	3,710	5,378	8,555	11,347	11,019	7,769	2,922	569	3
2003.....	1,443	1,989	2,683	3,697	5,591	8,503	11,294	11,187	7,542	2,866	536	2
2002.....	1,339	1,831	2,384	3,497	5,585	8,552	10,995	10,851	7,460	2,703	548	2
2001.....	1,303	1,716	2,294	3,734	5,651	8,441	10,401	10,567	7,166	2,677	568	1
2000.....	1,191	1,546	2,235	3,474	5,362	8,293	10,032	9,935	6,863	2,639	548	3
1999.....	1,082	1,470	2,120	3,389	5,475	8,041	9,652	9,740	6,701	2,616	514	1
All races, male												
2004.....	857	1,157	1,391	1,889	2,646	4,119	5,220	4,488	2,666	773	124	1
2003.....	827	1,084	1,374	1,869	2,750	4,118	5,218	4,494	2,577	734	112	2
2002.....	741	937	1,231	1,793	2,846	4,117	4,960	4,309	2,493	745	139	2
2001.....	707	884	1,150	1,941	2,782	3,970	4,683	4,194	2,354	677	104	-
2000.....	655	821	1,135	1,742	2,625	3,989	4,461	3,869	2,267	708	119	3
1999.....	571	766	1,117	1,687	2,737	3,926	4,358	3,778	2,095	674	115	-
All races, female												
2004.....	680	981	1,281	1,821	2,732	4,436	6,127	6,531	5,103	2,149	445	2
2003.....	616	905	1,309	1,828	2,841	4,385	6,076	6,693	4,965	2,132	424	-
2002.....	598	894	1,153	1,704	2,739	4,435	6,035	6,542	4,967	1,958	409	-
2001.....	596	832	1,144	1,793	2,869	4,471	5,718	6,373	4,812	2,000	464	1
2000.....	536	725	1,100	1,732	2,737	4,304	5,571	6,066	4,596	1,931	429	-
1999.....	511	704	1,003	1,702	2,738	4,115	5,294	5,962	4,606	1,942	399	1
White, both sexes												
2004.....	934	1,371	1,826	2,695	4,167	7,028	9,799	9,789	6,867	2,588	465	3
2003.....	865	1,312	1,843	2,675	4,290	7,033	9,732	9,987	6,680	2,533	439	1
2002.....	791	1,201	1,629	2,545	4,335	7,063	9,535	9,601	6,647	2,341	448	1
2001.....	797	1,116	1,540	2,678	4,425	6,984	8,947	9,332	6,369	2,368	467	1
2000.....	728	1,000	1,531	2,495	4,131	6,916	8,665	8,790	6,083	2,336	464	2
1999.....	654	983	1,450	2,433	4,253	6,664	8,326	8,510	5,959	2,282	421	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XIV. Diseases of the genitourinary system (N00-N98)												
White, male												
2004.....	515	768	966	1,380	2,106	3,459	4,608	4,023	2,384	668	103	1
2003.....	478	716	956	1,379	2,149	3,486	4,539	4,099	2,310	649	95	1
2002.....	439	613	835	1,339	2,259	3,461	4,377	3,859	2,221	644	107	1
2001.....	443	582	801	1,427	2,244	3,372	4,101	3,752	2,108	598	90	-
2000.....	397	534	793	1,301	2,105	3,379	3,923	3,493	2,013	614	102	2
1999.....	347	541	773	1,237	2,169	3,283	3,799	3,302	1,858	605	94	-
White, female												
2004.....	419	603	860	1,315	2,061	3,569	5,191	5,766	4,483	1,920	362	2
2003.....	387	596	887	1,296	2,141	3,547	5,193	5,888	4,370	1,884	344	-
2002.....	352	588	794	1,206	2,076	3,602	5,158	5,742	4,426	1,697	341	-
2001.....	354	534	739	1,251	2,181	3,612	4,846	5,580	4,261	1,770	377	1
2000.....	331	466	738	1,194	2,026	3,537	4,742	5,297	4,070	1,722	362	-
1999.....	307	442	677	1,196	2,084	3,381	4,527	5,208	4,101	1,677	327	-
Black, both sexes												
2004.....	557	699	770	895	1,077	1,314	1,337	1,060	796	303	91	-
2003.....	533	625	763	918	1,158	1,287	1,366	1,066	769	294	83	1
2002.....	506	565	676	865	1,120	1,312	1,270	1,093	720	327	91	1
2001.....	461	524	673	949	1,087	1,287	1,283	1,091	698	280	95	-
2000.....	423	494	636	874	1,094	1,235	1,215	1,005	688	273	72	1
1999.....	401	429	603	855	1,089	1,232	1,167	1,084	658	298	90	1
Black, male												
2004.....	319	357	378	442	482	572	511	389	227	90	17	-
2003.....	330	340	382	440	551	547	592	341	226	71	16	1
2002.....	279	298	354	408	531	572	500	391	236	86	28	1
2001.....	239	274	311	466	471	520	504	377	209	67	13	-
2000.....	235	260	305	393	466	549	473	320	206	82	13	1
1999.....	214	194	310	393	508	570	497	408	200	57	20	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XIV. Diseases of the genitourinary system (N00-N98)												
Black, female												
2004.....	238	342	392	453	595	742	826	671	569	213	74	-
2003.....	203	285	381	478	607	740	774	725	543	223	67	-
2002.....	227	267	322	457	589	740	770	702	484	241	63	-
2001.....	222	250	362	483	616	767	779	714	489	213	82	-
2000.....	188	234	331	481	628	686	742	685	482	191	59	-
1999.....	187	235	293	462	581	662	670	676	458	241	70	1
American Indian 1/, both sexes												
2004.....	24	23	25	40	36	47	35	31	15	6	1	-
2003.....	18	24	28	32	28	58	43	28	20	13	3	-
2002.....	18	25	25	34	40	34	34	27	15	5	4	-
2001.....	18	32	30	34	39	45	34	25	18	6	2	-
2000.....	19	19	26	39	33	27	38	24	21	4	3	-
1999.....	12	20	21	34	29	34	22	23	12	8	3	-
American Indian 1/, male												
2004.....	14	10	13	26	9	15	17	11	7	3	-	-
2003.....	9	14	10	15	10	31	15	11	5	5	-	-
2002.....	10	11	10	14	12	10	18	7	5	1	3	-
2001.....	11	9	12	11	18	19	14	12	6	1	-	-
2000.....	10	10	14	19	12	8	14	10	9	2	1	-
1999.....	5	9	8	14	13	16	4	7	1	2	1	-
American Indian 1/, female												
2004.....	10	13	12	14	27	32	18	20	8	3	1	-
2003.....	9	10	18	17	18	27	28	17	15	8	3	-
2002.....	8	14	15	20	28	24	16	20	10	4	1	-
2001.....	7	23	18	23	21	26	20	13	12	5	2	-
2000.....	9	9	12	20	21	19	24	14	12	2	2	-
1999.....	7	11	13	20	16	18	18	16	11	6	2	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XIV. Diseases of the genitourinary system (N00-N98)												
Asian or Pacific Islander 2/, both sexes												
2004.....	22	45	51	80	98	166	176	139	91	25	12	-
2003.....	27	28	49	72	115	125	153	106	73	26	11	-
2002.....	24	40	54	53	90	143	156	130	78	30	5	-
2001.....	27	44	51	73	100	125	137	119	81	23	4	-
2000.....	21	33	42	66	104	115	114	116	71	26	9	-
1999.....	15	38	46	67	104	111	137	123	72	28	-	-
Asian or Pacific Islander 2/, male												
2004.....	9	22	34	41	49	73	84	65	48	12	4	-
2003.....	10	14	26	35	40	54	72	43	36	9	1	-
2002.....	13	15	32	32	44	74	65	52	31	14	1	-
2001.....	14	19	26	37	49	59	64	53	31	11	1	-
2000.....	13	17	23	29	42	53	51	46	39	10	3	-
1999.....	5	22	26	43	47	57	58	61	36	10	-	-
Asian or Pacific Islander 2/, female												
2004.....	13	23	17	39	49	93	92	74	43	13	8	-
2003.....	17	14	23	37	75	71	81	63	37	17	10	-
2002.....	11	25	22	21	46	69	91	78	47	16	4	-
2001.....	13	25	25	36	51	66	73	66	50	12	3	-
2000.....	8	16	19	37	62	62	63	70	32	16	6	-
1999.....	10	16	20	24	57	54	79	62	36	18	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of urinary system (N00-N39)												
All races, both sexes												
2004.....	1,506	2,100	2,628	3,655	5,294	8,434	11,130	10,806	7,600	2,850	561	3
2003.....	1,419	1,960	2,647	3,638	5,524	8,374	11,116	10,981	7,391	2,800	532	2
2002.....	1,320	1,811	2,354	3,459	5,518	8,453	10,807	10,666	7,317	2,647	533	1
2001.....	1,275	1,691	2,265	3,691	5,558	8,315	10,239	10,378	7,021	2,624	559	1
2000.....	1,167	1,526	2,209	3,427	5,285	8,174	9,858	9,752	6,734	2,586	535	2
1999.....	1,061	1,444	2,084	3,342	5,399	7,913	9,477	9,560	6,590	2,568	498	1
All races, male												
2004.....	843	1,136	1,362	1,854	2,592	4,032	5,061	4,356	2,543	724	121	1
2003.....	822	1,067	1,357	1,835	2,708	4,038	5,095	4,359	2,472	693	110	2
2002.....	738	932	1,211	1,769	2,796	4,053	4,821	4,183	2,397	711	128	1
2001.....	697	872	1,135	1,918	2,728	3,890	4,576	4,063	2,251	641	98	-
2000.....	649	809	1,116	1,718	2,577	3,912	4,345	3,743	2,178	676	114	2
1999.....	560	755	1,097	1,656	2,686	3,847	4,228	3,644	2,014	645	105	-
All races, female												
2004.....	663	964	1,266	1,801	2,702	4,402	6,069	6,450	5,057	2,126	440	2
2003.....	597	893	1,290	1,803	2,816	4,336	6,021	6,622	4,919	2,107	422	-
2002.....	582	879	1,143	1,690	2,722	4,400	5,986	6,483	4,920	1,936	405	-
2001.....	578	819	1,130	1,773	2,830	4,425	5,663	6,315	4,770	1,983	461	1
2000.....	518	717	1,093	1,709	2,708	4,262	5,513	6,009	4,556	1,910	421	-
1999.....	501	689	987	1,686	2,713	4,066	5,249	5,916	4,576	1,923	393	1
White, both sexes												
2004.....	914	1,342	1,788	2,652	4,103	6,925	9,606	9,595	6,713	2,521	457	3
2003.....	851	1,291	1,818	2,630	4,231	6,918	9,567	9,793	6,539	2,469	435	1
2002.....	778	1,183	1,608	2,517	4,283	6,982	9,363	9,439	6,515	2,291	434	1
2001.....	779	1,097	1,524	2,640	4,345	6,875	8,801	9,162	6,234	2,319	459	1
2000.....	708	987	1,510	2,458	4,066	6,813	8,507	8,626	5,966	2,288	452	2
1999.....	641	963	1,423	2,397	4,195	6,555	8,168	8,349	5,856	2,237	408	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of urinary system (N00-N39)												
White, male												
2004.....	507	754	942	1,352	2,062	3,388	4,466	3,900	2,272	622	100	1
2003.....	477	705	944	1,352	2,110	3,417	4,426	3,974	2,210	610	93	1
2002.....	438	608	821	1,321	2,220	3,413	4,252	3,747	2,129	613	97	1
2001.....	437	573	793	1,406	2,197	3,300	4,005	3,632	2,010	565	85	-
2000.....	394	527	779	1,281	2,065	3,315	3,816	3,380	1,933	585	97	2
1999.....	342	534	759	1,212	2,129	3,209	3,683	3,180	1,784	577	86	-
White, female												
2004.....	407	588	846	1,300	2,041	3,537	5,140	5,695	4,441	1,899	357	2
2003.....	374	586	874	1,278	2,121	3,501	5,141	5,819	4,329	1,859	342	-
2002.....	340	575	787	1,196	2,063	3,569	5,111	5,692	4,386	1,678	337	-
2001.....	342	524	731	1,234	2,148	3,575	4,796	5,530	4,224	1,754	374	1
2000.....	314	460	731	1,177	2,001	3,498	4,691	5,246	4,033	1,703	355	-
1999.....	299	429	664	1,185	2,066	3,346	4,485	5,169	4,072	1,660	322	-
Black, both sexes												
2004.....	547	692	765	884	1,059	1,298	1,321	1,045	783	299	91	-
2003.....	524	620	752	907	1,151	1,274	1,358	1,054	760	293	83	1
2002.....	500	563	667	856	1,105	1,297	1,257	1,073	710	323	90	-
2001.....	451	520	660	944	1,077	1,270	1,267	1,075	688	277	94	-
2000.....	420	487	631	865	1,083	1,221	1,201	990	678	270	71	-
1999.....	394	423	594	844	1,072	1,214	1,150	1,066	651	295	87	1
Black, male												
2004.....	313	352	374	436	474	558	501	383	218	88	17	-
2003.....	326	336	377	434	549	536	586	331	222	70	16	1
2002.....	277	298	348	403	520	559	489	379	233	85	27	-
2001.....	235	272	304	464	465	512	493	368	204	65	12	-
2000.....	232	255	300	389	459	538	465	311	199	80	13	-
1999.....	209	190	304	387	498	565	483	397	194	56	18	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of urinary system (N00-N39)												
Black, female												
2004.....	234	340	391	448	585	740	820	662	565	211	74	-
2003.....	198	284	375	473	602	738	772	723	538	223	67	-
2002.....	223	265	319	453	585	738	768	694	477	238	63	-
2001.....	216	248	356	480	612	758	774	707	484	212	82	-
2000.....	188	232	331	476	624	683	736	679	479	190	58	-
1999.....	185	233	290	457	574	649	667	669	457	239	69	1
American Indian 1/, both sexes												
2004.....	24	22	25	39	36	46	34	30	15	6	1	-
2003.....	18	23	28	31	28	58	43	28	19	12	3	-
2002.....	18	25	25	33	40	34	33	27	15	5	4	-
2001.....	18	31	30	34	38	45	34	24	18	6	2	-
2000.....	19	19	26	39	32	27	37	23	21	3	3	-
1999.....	11	20	21	34	28	33	22	23	12	8	3	-
American Indian 1/, male												
2004.....	14	9	13	25	9	14	16	11	7	3	-	-
2003.....	9	13	10	15	10	31	15	11	4	4	-	-
2002.....	10	11	10	13	12	10	17	7	5	1	3	-
2001.....	11	9	12	11	17	19	14	12	6	1	-	-
2000.....	10	10	14	19	11	8	13	9	9	2	1	-
1999.....	4	9	8	14	12	16	4	7	1	2	1	-
American Indian 1/, female												
2004.....	10	13	12	14	27	32	18	19	8	3	1	-
2003.....	9	10	18	16	18	27	28	17	15	8	3	-
2002.....	8	14	15	20	28	24	16	20	10	4	1	-
2001.....	7	22	18	23	21	26	20	12	12	5	2	-
2000.....	9	9	12	20	21	19	24	14	12	1	2	-
1999.....	7	11	13	20	16	17	18	16	11	6	2	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of urinary system (N00-N39)												
Asian or Pacific Islander 2/, both sexes												
2004.....	21	44	50	80	96	165	169	136	89	24	12	-
2003.....	26	26	49	70	114	124	148	106	73	26	11	-
2002.....	24	40	54	53	90	140	154	127	77	28	5	-
2001.....	27	43	51	73	98	125	137	117	81	22	4	-
2000.....	20	33	42	65	104	113	113	113	69	25	9	-
1999.....	15	38	46	67	104	111	137	122	71	28	-	-
Asian or Pacific Islander 2/, male												
2004.....	9	21	33	41	47	72	78	62	46	11	4	-
2003.....	10	13	26	34	39	54	68	43	36	9	1	-
2002.....	13	15	32	32	44	71	63	50	30	12	1	-
2001.....	14	18	26	37	49	59	64	51	31	10	1	-
2000.....	13	17	23	29	42	51	51	43	37	9	3	-
1999.....	5	22	26	43	47	57	58	60	35	10	-	-
Asian or Pacific Islander 2/, female												
2004.....	12	23	17	39	49	93	91	74	43	13	8	-
2003.....	16	13	23	36	75	70	80	63	37	17	10	-
2002.....	11	25	22	21	46	69	91	77	47	16	4	-
2001.....	13	25	25	36	49	66	73	66	50	12	3	-
2000.....	7	16	19	36	62	62	62	70	32	16	6	-
1999.....	10	16	20	24	57	54	79	62	36	18	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Glomerular and renal tubulo-interstitial diseases (N00-N15)												
All races, both sexes												
2004.....	66	82	75	107	158	261	354	308	225	65	14	-
2003.....	73	74	78	122	160	240	373	342	199	72	13	-
2002.....	60	70	75	109	172	289	332	337	229	61	10	-
2001.....	64	58	68	115	173	281	298	304	214	80	7	-
2000.....	60	68	77	113	171	274	323	331	220	91	22	-
1999.....	78	75	82	120	199	309	350	366	220	85	11	-
All races, male												
2004.....	27	48	43	52	75	118	151	136	89	27	3	-
2003.....	40	30	35	51	69	109	159	145	90	21	1	-
2002.....	32	39	33	54	90	144	135	147	87	13	5	-
2001.....	28	27	29	56	95	118	147	121	71	29	2	-
2000.....	35	30	37	53	64	131	132	126	87	30	3	-
1999.....	35	35	43	54	76	145	176	154	78	34	4	-
All races, female												
2004.....	39	34	32	55	83	143	203	172	136	38	11	-
2003.....	33	44	43	71	91	131	214	197	109	51	12	-
2002.....	28	31	42	55	82	145	197	190	142	48	5	-
2001.....	36	31	39	59	78	163	151	183	143	51	5	-
2000.....	25	38	40	60	107	143	191	205	133	61	19	-
1999.....	43	40	39	66	123	164	174	212	142	51	7	-
White, both sexes												
2004.....	51	62	58	88	138	225	319	279	198	57	11	-
2003.....	53	60	57	105	132	206	321	313	173	66	8	-
2002.....	44	54	59	97	158	249	298	304	209	55	8	-
2001.....	43	46	49	88	153	252	258	277	193	77	6	-
2000.....	33	56	58	89	144	234	281	295	190	81	19	-
1999.....	57	60	58	90	162	274	315	330	197	77	10	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Glomerular and renal tubulo-interstitial diseases (N00-N15)												
White, male												
2004.....	23	36	32	43	63	101	139	120	76	25	3	-
2003.....	29	26	24	44	59	92	133	136	78	20	1	-
2002.....	24	32	28	44	82	119	122	135	81	12	4	-
2001.....	18	21	19	38	83	103	130	109	67	28	2	-
2000.....	17	26	28	39	55	114	119	115	72	26	3	-
1999.....	26	28	31	41	57	130	158	136	67	31	4	-
White, female												
2004.....	28	26	26	45	75	124	180	159	122	32	8	-
2003.....	24	34	33	61	73	114	188	177	95	46	7	-
2002.....	20	22	31	53	76	130	176	169	128	43	4	-
2001.....	25	25	30	50	70	149	128	168	126	49	4	-
2000.....	16	30	30	50	89	120	162	180	118	55	16	-
1999.....	31	32	27	49	105	144	157	194	130	46	6	-
Black, both sexes												
2004.....	12	18	13	17	17	30	27	22	20	7	3	-
2003.....	17	11	16	13	19	27	40	20	24	3	4	-
2002.....	12	12	11	11	12	31	23	29	15	6	2	-
2001.....	17	8	15	24	17	24	31	22	15	2	1	-
2000.....	26	9	15	22	21	34	32	25	23	9	1	-
1999.....	19	12	18	23	33	28	26	34	20	7	1	-
Black, male												
2004.....	3	11	9	8	10	15	7	13	8	1	-	-
2003.....	10	3	8	6	7	14	22	5	11	1	-	-
2002.....	6	6	4	9	7	18	9	10	5	1	1	-
2001.....	9	5	9	16	10	10	12	10	2	-	-	-
2000.....	17	3	9	13	7	14	10	7	11	3	-	-
1999.....	9	5	10	10	18	12	12	17	10	3	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Glomerular and renal tubulo-interstitial diseases (N00-N15)												
Black, female												
2004.....	9	7	4	9	7	15	20	9	12	6	3	-
2003.....	7	8	8	7	12	13	18	15	13	2	4	-
2002.....	6	6	7	2	5	13	14	19	10	5	1	-
2001.....	8	3	6	8	7	14	19	12	13	2	1	-
2000.....	9	6	6	9	14	20	22	18	12	6	1	-
1999.....	10	7	8	13	15	16	14	17	10	4	1	-
American Indian 1/, both sexes												
2004.....	2	1	2	1	1	2	1	-	1	-	-	-
2003.....	1	2	1	-	2	3	-	1	-	1	-	-
2002.....	1	1	1	-	-	2	3	-	2	-	-	-
2001.....	1	2	-	-	1	2	3	-	1	1	-	-
2000.....	1	1	-	1	1	1	2	1	1	-	2	-
1999.....	-	2	1	3	-	1	-	-	1	-	-	-
American Indian 1/, male												
2004.....	1	-	1	1	-	-	1	-	-	-	-	-
2003.....	1	-	-	-	2	2	-	-	-	-	-	-
2002.....	-	1	-	-	-	2	1	-	1	-	-	-
2001.....	-	1	-	-	1	2	3	-	-	1	-	-
2000.....	1	1	-	-	-	1	1	1	-	-	-	-
1999.....	-	2	-	1	-	-	-	-	-	-	-	-
American Indian 1/, female												
2004.....	1	1	1	-	1	2	-	-	1	-	-	-
2003.....	-	2	1	-	-	1	-	1	-	1	-	-
2002.....	1	-	1	-	-	-	2	-	1	-	-	-
2001.....	1	1	-	-	-	-	-	-	1	-	-	-
2000.....	-	-	-	1	1	-	1	-	1	-	2	-
1999.....	-	-	1	2	-	1	-	-	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Glomerular and renal tubulo-interstitial diseases (N00-N15)												
Asian or Pacific Islander 2/, both sexes												
2004.....	1	1	2	1	2	4	7	7	6	1	-	-
2003.....	2	1	4	4	7	4	12	8	2	2	1	-
2002.....	3	3	4	1	2	7	8	4	3	-	-	-
2001.....	3	2	4	3	2	3	6	5	5	-	-	-
2000.....	-	2	4	1	5	5	8	10	6	1	-	-
1999.....	2	1	5	4	4	6	9	2	2	1	-	-
Asian or Pacific Islander 2/, male												
2004.....	-	1	1	-	2	2	4	3	5	1	-	-
2003.....	-	1	3	1	1	1	4	4	1	-	-	-
2002.....	2	-	1	1	1	5	3	2	-	-	-	-
2001.....	1	-	1	2	1	3	2	2	2	-	-	-
2000.....	-	-	-	1	2	2	2	3	4	1	-	-
1999.....	-	-	2	2	1	3	6	1	1	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	1	-	1	1	-	2	3	4	1	-	-	-
2003.....	2	-	1	3	6	3	8	4	1	2	1	-
2002.....	1	3	3	-	1	2	5	2	3	-	-	-
2001.....	2	2	3	1	1	-	4	3	3	-	-	-
2000.....	-	2	4	-	3	3	6	7	2	-	-	-
1999.....	2	1	3	2	3	3	3	1	1	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Glomerular diseases (N00-N07)												
All races, both sexes												
2004.....	34	31	38	48	60	93	109	89	53	16	3	-
2003.....	32	24	34	51	72	84	115	84	37	15	6	-
2002.....	28	37	27	39	64	98	116	87	44	17	2	-
2001.....	31	24	26	60	76	92	93	89	44	12	2	-
2000.....	32	32	34	54	72	87	120	97	34	20	3	-
1999.....	32	30	43	46	82	127	113	101	39	13	1	-
All races, male												
2004.....	18	17	24	23	32	46	53	38	20	5	1	-
2003.....	22	13	15	30	29	39	52	32	19	5	1	-
2002.....	16	23	13	22	34	50	49	39	18	2	1	-
2001.....	19	14	17	29	44	47	51	40	14	6	-	-
2000.....	23	17	21	30	33	53	47	47	13	9	-	-
1999.....	17	14	23	21	33	66	61	43	16	8	-	-
All races, female												
2004.....	16	14	14	25	28	47	56	51	33	11	2	-
2003.....	10	11	19	21	43	45	63	52	18	10	5	-
2002.....	12	14	14	17	30	48	67	48	26	15	1	-
2001.....	12	10	9	31	32	45	42	49	30	6	2	-
2000.....	9	15	13	24	39	34	73	50	21	11	3	-
1999.....	15	16	20	25	49	61	52	58	23	5	1	-
White, both sexes												
2004.....	28	24	29	37	51	78	95	81	42	14	2	-
2003.....	21	22	26	43	54	68	96	73	28	14	3	-
2002.....	17	27	22	35	57	81	98	68	38	16	1	-
2001.....	17	19	17	44	68	82	81	81	39	11	1	-
2000.....	15	25	25	40	64	74	107	79	27	17	2	-
1999.....	21	20	30	33	70	111	94	85	32	11	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Glomerular diseases (N00-N07)												
White, male												
2004.....	16	13	17	19	27	38	47	33	15	4	1	-
2003.....	15	12	10	25	22	29	42	28	14	5	1	-
2002.....	9	18	13	19	30	40	43	33	16	2	1	-
2001.....	11	10	11	20	40	43	47	37	13	6	-	-
2000.....	10	14	16	22	31	46	44	38	9	7	-	-
1999.....	12	9	17	16	27	60	50	35	13	7	-	-
White, female												
2004.....	12	11	12	18	24	40	48	48	27	10	1	-
2003.....	6	10	16	18	32	39	54	45	14	9	2	-
2002.....	8	9	9	16	27	41	55	35	22	14	-	-
2001.....	6	9	6	24	28	39	34	44	26	5	1	-
2000.....	5	11	9	18	33	28	63	41	18	10	2	-
1999.....	9	11	13	17	43	51	44	50	19	4	1	-
Black, both sexes												
2004.....	4	6	8	10	9	11	12	5	9	1	1	-
2003.....	9	1	5	6	12	12	12	4	8	-	2	-
2002.....	10	6	3	3	6	13	12	16	4	1	1	-
2001.....	10	3	7	14	6	9	9	5	4	1	1	-
2000.....	16	5	7	12	6	10	12	11	4	2	-	-
1999.....	10	7	8	10	9	12	12	15	5	2	-	-
Black, male												
2004.....	1	3	6	3	5	6	4	3	3	-	-	-
2003.....	6	-	3	4	5	7	7	1	4	-	-	-
2002.....	6	4	-	2	4	8	4	5	1	-	-	-
2001.....	7	3	5	8	3	3	2	2	-	-	-	-
2000.....	12	2	5	7	1	4	3	5	2	1	-	-
1999.....	5	3	5	3	5	5	6	7	2	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Glomerular diseases (N00-N07)												
Black, female												
2004.....	3	3	2	7	4	5	8	2	6	1	1	-
2003.....	3	1	2	2	7	5	5	3	4	-	2	-
2002.....	4	2	3	1	2	5	8	11	3	1	1	-
2001.....	3	-	2	6	3	6	7	3	4	1	1	-
2000.....	4	3	2	5	5	6	9	6	2	1	-	-
1999.....	5	4	3	7	4	7	6	8	3	1	-	-
American Indian 1/, both sexes												
2004.....	2	-	1	1	-	1	-	-	-	-	-	-
2003.....	1	-	-	-	2	2	-	1	-	-	-	-
2002.....	-	1	-	-	-	1	2	-	1	-	-	-
2001.....	1	2	-	-	-	1	1	-	-	-	-	-
2000.....	1	1	-	1	-	1	-	1	-	-	1	-
1999.....	-	2	1	1	-	1	-	-	-	-	-	-
American Indian 1/, male												
2004.....	1	-	1	1	-	-	-	-	-	-	-	-
2003.....	1	-	-	-	2	2	-	-	-	-	-	-
2002.....	-	1	-	-	-	1	-	-	1	-	-	-
2001.....	-	1	-	-	-	1	1	-	-	-	-	-
2000.....	1	1	-	-	-	1	-	1	-	-	-	-
1999.....	-	2	-	1	-	-	-	-	-	-	-	-
American Indian 1/, female												
2004.....	1	-	-	-	-	1	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	1	-	-	-	-
2002.....	-	-	-	-	-	-	2	-	-	-	-	-
2001.....	1	1	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	1	-	-	-	-	-	-	1	-
1999.....	-	-	1	-	-	1	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Glomerular diseases (N00-N07)												
Asian or Pacific Islander 2/, both sexes												
2004.....	-	1	-	-	-	3	2	3	2	1	-	-
2003.....	1	1	3	2	4	2	7	6	1	1	1	-
2002.....	1	3	2	1	1	3	4	3	1	-	-	-
2001.....	3	-	2	2	2	-	2	3	1	-	-	-
2000.....	-	1	2	1	2	2	1	6	3	1	-	-
1999.....	1	1	4	2	3	3	7	1	2	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	-	1	-	-	-	2	2	2	2	1	-	-
2003.....	-	1	2	1	-	1	3	3	1	-	-	-
2002.....	1	-	-	1	-	1	2	1	-	-	-	-
2001.....	1	-	1	1	1	-	1	1	1	-	-	-
2000.....	-	-	-	1	1	2	-	3	2	1	-	-
1999.....	-	-	1	1	1	1	5	1	1	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	-	-	-	-	-	1	-	1	-	-	-	-
2003.....	1	-	1	1	4	1	4	3	-	1	1	-
2002.....	-	3	2	-	1	2	2	2	1	-	-	-
2001.....	2	-	1	1	1	-	1	2	-	-	-	-
2000.....	-	1	2	-	1	-	1	3	1	-	-	-
1999.....	1	1	3	1	2	2	2	-	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute and rapidly progressive nephritic syndrome (N00-N01)												
All races, both sexes												
2004.....	3	-	1	3	1	5	5	6	1	-	1	-
2003.....	2	-	-	2	4	2	5	2	-	-	-	-
2002.....	1	-	1	4	3	2	2	5	3	2	-	-
2001.....	-	1	-	2	1	4	4	4	4	-	-	-
2000.....	1	1	-	3	2	2	6	4	2	2	-	-
1999.....	1	1	2	4	3	5	3	5	1	-	-	-
All races, male												
2004.....	1	-	-	1	1	3	4	1	1	-	-	-
2003.....	1	-	-	1	3	1	1	-	-	-	-	-
2002.....	-	-	1	2	1	-	1	3	-	1	-	-
2001.....	-	1	-	2	1	1	2	-	1	-	-	-
2000.....	1	1	-	1	1	1	5	2	1	1	-	-
1999.....	-	1	1	4	1	3	1	1	1	-	-	-
All races, female												
2004.....	2	-	1	2	-	2	1	5	-	-	1	-
2003.....	1	-	-	1	1	1	4	2	-	-	-	-
2002.....	1	-	-	2	2	2	1	2	3	1	-	-
2001.....	-	-	-	-	-	3	2	4	3	-	-	-
2000.....	-	-	-	2	1	1	1	2	1	1	-	-
1999.....	1	-	1	-	2	2	2	4	-	-	-	-
White, both sexes												
2004.....	3	-	1	2	1	4	4	6	1	-	-	-
2003.....	2	-	-	1	4	2	4	2	-	-	-	-
2002.....	1	-	1	3	2	2	2	4	2	2	-	-
2001.....	-	-	-	2	1	4	4	3	3	-	-	-
2000.....	-	1	-	2	2	1	6	4	2	2	-	-
1999.....	1	-	1	3	3	5	3	4	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute and rapidly progressive nephritic syndrome (N00-N01)												
White, male												
2004.....	1	-	-	1	1	2	3	1	1	-	-	-
2003.....	1	-	-	-	3	1	1	-	-	-	-	-
2002.....	-	-	1	1	1	-	1	2	-	1	-	-
2001.....	-	-	-	2	1	1	2	-	1	-	-	-
2000.....	-	1	-	1	1	1	5	2	1	1	-	-
1999.....	-	-	-	3	1	3	1	-	1	-	-	-
White, female												
2004.....	2	-	1	1	-	2	1	5	-	-	-	-
2003.....	1	-	-	1	1	1	3	2	-	-	-	-
2002.....	1	-	-	2	1	2	1	2	2	1	-	-
2001.....	-	-	-	-	-	3	2	3	2	-	-	-
2000.....	-	-	-	1	1	-	1	2	1	1	-	-
1999.....	1	-	1	-	2	2	2	4	-	-	-	-
Black, both sexes												
2004.....	-	-	-	1	-	1	1	-	-	-	1	-
2003.....	-	-	-	1	-	-	-	-	-	-	-	-
2002.....	-	-	-	1	1	-	-	1	1	-	-	-
2001.....	-	-	-	-	-	-	-	1	1	-	-	-
2000.....	1	-	-	1	-	1	-	-	-	-	-	-
1999.....	-	-	1	-	-	-	-	1	-	-	-	-
Black, male												
2004.....	-	-	-	-	-	1	1	-	-	-	-	-
2003.....	-	-	-	1	-	-	-	-	-	-	-	-
2002.....	-	-	-	1	-	-	-	1	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	1	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	1	-	-	-	-	1	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nephrotic syndrome (N04)												
All races, both sexes												
2004.....	7	3	5	10	13	19	20	18	11	1	-	-
2003.....	2	5	9	9	15	16	27	15	5	2	2	-
2002.....	3	6	9	9	11	17	30	18	10	3	1	-
2001.....	5	8	4	11	13	24	16	20	8	1	1	-
2000.....	10	1	8	10	14	14	30	23	11	6	1	-
1999.....	4	4	9	6	24	32	22	17	6	5	-	-
All races, male												
2004.....	5	-	4	4	8	8	9	6	3	-	-	-
2003.....	-	3	4	5	6	6	12	2	3	-	-	-
2002.....	1	5	5	7	4	5	11	7	2	-	1	-
2001.....	4	5	3	4	3	9	7	10	4	-	-	-
2000.....	8	1	6	6	4	10	10	11	4	3	-	-
1999.....	3	1	3	1	11	12	16	8	-	4	-	-
All races, female												
2004.....	2	3	1	6	5	11	11	12	8	1	-	-
2003.....	2	2	5	4	9	10	15	13	2	2	2	-
2002.....	2	1	4	2	7	12	19	11	8	3	-	-
2001.....	1	3	1	7	10	15	9	10	4	1	1	-
2000.....	2	-	2	4	10	4	20	12	7	3	1	-
1999.....	1	3	6	5	13	20	6	9	6	1	-	-
White, both sexes												
2004.....	7	3	4	7	12	17	18	15	9	1	-	-
2003.....	2	3	6	8	13	14	22	12	5	2	-	-
2002.....	1	5	8	6	10	15	23	13	8	3	1	-
2001.....	4	7	4	7	11	21	14	17	6	1	1	-
2000.....	4	1	8	6	12	11	28	19	10	6	1	-
1999.....	2	3	5	6	20	27	17	13	6	5	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nephrotic syndrome (N04)												
White, male												
2004.....	5	-	3	4	8	7	8	5	2	-	-	-
2003.....	-	2	1	4	5	4	8	2	3	-	-	-
2002.....	1	4	5	5	4	4	9	6	1	-	1	-
2001.....	4	4	3	1	2	9	7	8	3	-	-	-
2000.....	3	1	6	3	4	8	9	8	3	3	-	-
1999.....	2	1	2	1	9	10	12	5	-	4	-	-
White, female												
2004.....	2	3	1	3	4	10	10	10	7	1	-	-
2003.....	2	1	5	4	8	10	14	10	2	2	-	-
2002.....	-	1	3	1	6	11	14	7	7	3	-	-
2001.....	-	3	1	6	9	12	7	9	3	1	1	-
2000.....	1	-	2	3	8	3	19	11	7	3	1	-
1999.....	-	2	3	5	11	17	5	8	6	1	-	-
Black, both sexes												
2004.....	-	-	1	3	1	2	1	2	1	-	-	-
2003.....	-	1	2	1	2	2	4	1	-	-	1	-
2002.....	2	1	-	2	1	2	4	4	1	-	-	-
2001.....	1	1	-	4	2	3	2	1	1	-	-	-
2000.....	6	-	-	3	2	2	2	2	-	-	-	-
1999.....	2	1	3	-	3	3	3	3	-	-	-	-
Black, male												
2004.....	-	-	1	-	-	1	-	1	-	-	-	-
2003.....	-	-	2	1	1	2	3	-	-	-	-	-
2002.....	-	1	-	1	-	1	1	1	-	-	-	-
2001.....	-	1	-	3	1	-	-	1	-	-	-	-
2000.....	5	-	-	2	-	1	1	1	-	-	-	-
1999.....	1	-	1	-	2	1	2	2	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nephrotic syndrome (N04)												
Black, female												
2004.....	-	-	-	3	1	1	1	1	1	-	-	-
2003.....	-	1	-	-	1	-	1	1	-	-	1	-
2002.....	2	-	-	1	1	1	3	3	1	-	-	-
2001.....	1	-	-	1	1	3	2	-	1	-	-	-
2000.....	1	-	-	1	2	1	1	1	-	-	-	-
1999.....	1	1	2	-	1	2	1	1	-	-	-	-
American Indian 1/, both sexes												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	1	-	-	-	-
2002.....	-	-	-	-	-	-	1	-	1	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	1	-	-	-	-
1999.....	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, male												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	1	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	1	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, female												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	1	-	-	-	-
2002.....	-	-	-	-	-	-	1	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	1	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nephrotic syndrome (N04)												
Asian or Pacific Islander 2/, both sexes												
2004.....	-	-	-	-	-	-	1	1	1	-	-	-
2003.....	-	1	1	-	-	-	1	1	-	-	1	-
2002.....	-	-	1	1	-	-	2	1	-	-	-	-
2001.....	-	-	-	-	-	-	-	2	1	-	-	-
2000.....	-	-	-	1	-	1	-	1	1	-	-	-
1999.....	-	-	1	-	1	1	2	1	-	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	-	-	-	-	-	-	1	-	1	-	-	-
2003.....	-	1	1	-	-	-	1	-	-	-	-	-
2002.....	-	-	-	1	-	-	1	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	1	1	-	-	-
2000.....	-	-	-	1	-	1	-	1	1	-	-	-
1999.....	-	-	-	-	-	1	2	1	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	-	-	-	-	-	-	-	1	-	-	-	-
2003.....	-	-	-	-	-	-	-	1	-	-	1	-
2002.....	-	-	1	-	-	-	1	1	-	-	-	-
2001.....	-	-	-	-	-	-	-	1	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	1	-	1	-	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified glomerular diseases (N02-N03,N05-N07)												
All races, both sexes												
2004.....	24	28	32	35	46	69	84	65	41	15	2	-
2003.....	28	19	25	40	53	66	83	67	32	13	4	-
2002.....	24	31	17	26	50	79	84	64	31	12	1	-
2001.....	26	15	22	47	62	64	73	65	32	11	1	-
2000.....	21	30	26	41	56	71	84	70	21	12	2	-
1999.....	27	25	32	36	55	90	88	79	32	8	1	-
All races, male												
2004.....	12	17	20	18	23	35	40	31	16	5	1	-
2003.....	21	10	11	24	20	32	39	30	16	5	1	-
2002.....	15	18	7	13	29	45	37	29	16	1	-	-
2001.....	15	8	14	23	40	37	42	30	9	6	-	-
2000.....	14	15	15	23	28	42	32	34	8	5	-	-
1999.....	14	12	19	16	21	51	44	34	15	4	-	-
All races, female												
2004.....	12	11	12	17	23	34	44	34	25	10	1	-
2003.....	7	9	14	16	33	34	44	37	16	8	3	-
2002.....	9	13	10	13	21	34	47	35	15	11	1	-
2001.....	11	7	8	24	22	27	31	35	23	5	1	-
2000.....	7	15	11	18	28	29	52	36	13	7	2	-
1999.....	13	13	13	20	34	39	44	45	17	4	1	-
White, both sexes												
2004.....	18	21	24	28	38	57	73	60	32	13	2	-
2003.....	17	19	20	34	37	52	70	59	23	12	3	-
2002.....	15	22	13	26	45	64	73	51	28	11	-	-
2001.....	13	12	13	35	56	57	63	61	30	10	-	-
2000.....	11	23	17	32	50	62	73	56	15	9	1	-
1999.....	18	17	24	24	47	79	74	68	25	6	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified glomerular diseases (N02-N03,N05-N07)												
White, male												
2004.....	10	13	14	14	18	29	36	27	12	4	1	-
2003.....	14	10	9	21	14	24	33	26	11	5	1	-
2002.....	8	14	7	13	25	36	33	25	15	1	-	-
2001.....	7	6	8	17	37	33	38	29	9	6	-	-
2000.....	7	12	10	18	26	37	30	28	5	3	-	-
1999.....	10	8	15	12	17	47	37	30	12	3	-	-
White, female												
2004.....	8	8	10	14	20	28	37	33	20	9	1	-
2003.....	3	9	11	13	23	28	37	33	12	7	2	-
2002.....	7	8	6	13	20	28	40	26	13	10	-	-
2001.....	6	6	5	18	19	24	25	32	21	4	-	-
2000.....	4	11	7	14	24	25	43	28	10	6	1	-
1999.....	8	9	9	12	30	32	37	38	13	3	1	-
Black, both sexes												
2004.....	4	6	7	6	8	8	10	3	8	1	-	-
2003.....	9	-	3	4	10	10	8	3	8	-	1	-
2002.....	8	5	3	-	4	11	8	11	2	1	1	-
2001.....	9	2	7	10	4	6	7	3	2	1	1	-
2000.....	9	5	7	8	4	7	10	9	4	2	-	-
1999.....	8	6	4	10	6	9	9	11	5	2	-	-
Black, male												
2004.....	1	3	5	3	5	4	3	2	3	-	-	-
2003.....	6	-	1	2	4	5	4	1	4	-	-	-
2002.....	6	3	-	-	4	7	3	3	1	-	-	-
2001.....	7	2	5	5	2	3	2	1	-	-	-	-
2000.....	6	2	5	5	1	3	2	4	2	1	-	-
1999.....	4	3	3	3	3	4	4	4	2	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified glomerular diseases (N02-N03,N05-N07)												
Asian or Pacific Islander 2/, both sexes												
2004.....	-	1	-	-	-	3	1	2	1	1	-	-
2003.....	1	-	2	2	4	2	5	5	1	1	-	-
2002.....	1	3	1	-	1	3	2	2	1	-	-	-
2001.....	3	-	2	2	2	-	2	1	-	-	-	-
2000.....	-	1	2	-	2	1	1	5	2	1	-	-
1999.....	1	1	3	1	2	2	5	-	2	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	-	1	-	-	-	2	1	2	1	1	-	-
2003.....	-	-	1	1	-	1	2	3	1	-	-	-
2002.....	1	-	-	-	-	1	1	1	-	-	-	-
2001.....	1	-	1	1	1	-	1	-	-	-	-	-
2000.....	-	-	-	-	1	1	-	2	1	1	-	-
1999.....	-	-	1	-	1	-	3	-	1	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	-	-	-	-	-	1	-	-	-	-	-	-
2003.....	1	-	1	1	4	1	3	2	-	1	-	-
2002.....	-	3	1	-	1	2	1	1	1	-	-	-
2001.....	2	-	1	1	1	-	1	1	-	-	-	-
2000.....	-	1	2	-	1	-	1	3	1	-	-	-
1999.....	1	1	2	1	1	2	2	-	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Renal tubulo-interstitial diseases (N10-N15)												
All races, both sexes												
2004.....	32	51	37	59	98	168	245	219	172	49	11	-
2003.....	41	50	44	71	88	156	258	258	162	57	7	-
2002.....	32	33	48	70	108	191	216	250	185	44	8	-
2001.....	33	34	42	55	97	189	205	215	170	68	5	-
2000.....	28	36	43	59	99	187	203	234	186	71	19	-
1999.....	46	45	39	74	117	182	237	265	181	72	10	-
All races, male												
2004.....	9	31	19	29	43	72	98	98	69	22	2	-
2003.....	18	17	20	21	40	70	107	113	71	16	-	-
2002.....	16	16	20	32	56	94	86	108	69	11	4	-
2001.....	9	13	12	27	51	71	96	81	57	23	2	-
2000.....	12	13	16	23	31	78	85	79	74	21	3	-
1999.....	18	21	20	33	43	79	115	111	62	26	4	-
All races, female												
2004.....	23	20	18	30	55	96	147	121	103	27	9	-
2003.....	23	33	24	50	48	86	151	145	91	41	7	-
2002.....	16	17	28	38	52	97	130	142	116	33	4	-
2001.....	24	21	30	28	46	118	109	134	113	45	3	-
2000.....	16	23	27	36	68	109	118	155	112	50	16	-
1999.....	28	24	19	41	74	103	122	154	119	46	6	-
White, both sexes												
2004.....	23	38	29	51	87	147	224	198	156	43	9	-
2003.....	32	38	31	62	78	138	225	240	145	52	5	-
2002.....	27	27	37	62	101	168	200	236	171	39	7	-
2001.....	26	27	32	44	85	170	177	196	154	66	5	-
2000.....	18	31	33	49	80	160	174	216	163	64	17	-
1999.....	36	40	28	57	92	163	221	245	165	66	9	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Renal tubulo-interstitial diseases (N10-N15)												
White, male												
2004.....	7	23	15	24	36	63	92	87	61	21	2	-
2003.....	14	14	14	19	37	63	91	108	64	15	-	-
2002.....	15	14	15	25	52	79	79	102	65	10	3	-
2001.....	7	11	8	18	43	60	83	72	54	22	2	-
2000.....	7	12	12	17	24	68	75	77	63	19	3	-
1999.....	14	19	14	25	30	70	108	101	54	24	4	-
White, female												
2004.....	16	15	14	27	51	84	132	111	95	22	7	-
2003.....	18	24	17	43	41	75	134	132	81	37	5	-
2002.....	12	13	22	37	49	89	121	134	106	29	4	-
2001.....	19	16	24	26	42	110	94	124	100	44	3	-
2000.....	11	19	21	32	56	92	99	139	100	45	14	-
1999.....	22	21	14	32	62	93	113	144	111	42	5	-
Black, both sexes												
2004.....	8	12	5	7	8	19	15	17	11	6	2	-
2003.....	8	10	11	7	7	15	28	16	16	3	2	-
2002.....	2	6	8	8	6	18	11	13	11	5	1	-
2001.....	7	5	8	10	11	15	22	17	11	1	-	-
2000.....	10	4	8	10	15	24	20	14	19	7	1	-
1999.....	9	5	10	13	24	16	14	19	15	5	1	-
Black, male												
2004.....	2	8	3	5	5	9	3	10	5	1	-	-
2003.....	4	3	5	2	2	7	15	4	7	1	-	-
2002.....	-	2	4	7	3	10	5	5	4	1	1	-
2001.....	2	2	4	8	7	7	10	8	2	-	-	-
2000.....	5	1	4	6	6	10	7	2	9	2	-	-
1999.....	4	2	5	7	13	7	6	10	8	2	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Renal tubulo-interstitial diseases (N10-N15)												
Black, female												
2004.....	6	4	2	2	3	10	12	7	6	5	2	-
2003.....	4	7	6	5	5	8	13	12	9	2	2	-
2002.....	2	4	4	1	3	8	6	8	7	4	-	-
2001.....	5	3	4	2	4	8	12	9	9	1	-	-
2000.....	5	3	4	4	9	14	13	12	10	5	1	-
1999.....	5	3	5	6	11	9	8	9	7	3	1	-
American Indian 1/, both sexes												
2004.....	-	1	1	-	1	1	1	-	1	-	-	-
2003.....	-	2	1	-	-	1	-	-	-	1	-	-
2002.....	1	-	1	-	-	1	1	-	1	-	-	-
2001.....	-	-	-	-	1	1	2	-	1	1	-	-
2000.....	-	-	-	-	1	-	2	-	1	-	1	-
1999.....	-	-	-	2	-	-	-	-	1	-	-	-
American Indian 1/, male												
2004.....	-	-	-	-	-	-	1	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	1	1	-	-	-	-	-
2001.....	-	-	-	-	1	1	2	-	-	1	-	-
2000.....	-	-	-	-	-	-	1	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, female												
2004.....	-	1	1	-	1	1	-	-	1	-	-	-
2003.....	-	2	1	-	-	1	-	-	-	1	-	-
2002.....	1	-	1	-	-	-	-	-	1	-	-	-
2001.....	-	-	-	-	-	-	-	-	1	-	-	-
2000.....	-	-	-	-	1	-	1	-	1	-	1	-
1999.....	-	-	-	2	-	-	-	-	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Renal tubulo-interstitial diseases (N10-N15)												
Asian or Pacific Islander 2/, both sexes												
2004.....	1	-	2	1	2	1	5	4	4	-	-	-
2003.....	1	-	1	2	3	2	5	2	1	1	-	-
2002.....	2	-	2	-	1	4	4	1	2	-	-	-
2001.....	-	2	2	1	-	3	4	2	4	-	-	-
2000.....	-	1	2	-	3	3	7	4	3	-	-	-
1999.....	1	-	1	2	1	3	2	1	-	1	-	-
Asian or Pacific Islander 2/, male												
2004.....	-	-	1	-	2	-	2	1	3	-	-	-
2003.....	-	-	1	-	1	-	1	1	-	-	-	-
2002.....	1	-	1	-	1	4	1	1	-	-	-	-
2001.....	-	-	-	1	-	3	1	1	1	-	-	-
2000.....	-	-	-	-	1	-	2	-	2	-	-	-
1999.....	-	-	1	1	-	2	1	-	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	1	-	1	1	-	1	3	3	1	-	-	-
2003.....	1	-	-	2	2	2	4	1	1	1	-	-
2002.....	1	-	1	-	-	-	3	-	2	-	-	-
2001.....	-	2	2	-	-	-	3	1	3	-	-	-
2000.....	-	1	2	-	2	3	5	4	1	-	-	-
1999.....	1	-	-	1	1	1	1	1	-	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Renal failure (N17-N19)												
All races, both sexes												
2004.....	1,208	1,705	2,178	2,894	4,103	6,222	7,608	7,062	4,676	1,695	317	2
2003.....	1,131	1,611	2,130	2,939	4,267	6,198	7,672	7,184	4,541	1,621	307	2
2002.....	1,072	1,464	1,919	2,774	4,272	6,228	7,429	6,856	4,398	1,530	296	-
2001.....	1,041	1,376	1,850	2,963	4,248	6,007	6,891	6,517	4,174	1,514	300	1
2000.....	932	1,234	1,789	2,754	4,094	5,794	6,429	6,040	3,878	1,420	265	2
1999.....	836	1,144	1,640	2,641	4,056	5,560	6,092	5,718	3,656	1,372	252	-
All races, male												
2004.....	710	925	1,147	1,518	2,098	3,113	3,782	3,157	1,793	507	83	1
2003.....	686	923	1,113	1,532	2,187	3,140	3,807	3,198	1,771	486	80	2
2002.....	615	772	1,014	1,468	2,271	3,190	3,618	3,046	1,672	492	81	-
2001.....	588	735	945	1,581	2,171	3,025	3,356	2,953	1,604	429	64	-
2000.....	526	678	929	1,412	2,080	2,981	3,121	2,617	1,458	443	73	2
1999.....	462	612	880	1,340	2,100	2,912	2,976	2,535	1,334	415	67	-
All races, female												
2004.....	498	780	1,031	1,376	2,005	3,109	3,826	3,905	2,883	1,188	234	1
2003.....	445	688	1,017	1,407	2,080	3,058	3,865	3,986	2,770	1,135	227	-
2002.....	457	692	905	1,306	2,001	3,038	3,811	3,810	2,726	1,038	215	-
2001.....	453	641	905	1,382	2,077	2,982	3,535	3,564	2,570	1,085	236	1
2000.....	406	556	860	1,342	2,014	2,813	3,308	3,423	2,420	977	192	-
1999.....	374	532	760	1,301	1,956	2,648	3,116	3,183	2,322	957	185	-
White, both sexes												
2004.....	696	1,058	1,432	2,038	3,115	5,044	6,499	6,191	4,096	1,488	257	2
2003.....	653	1,030	1,435	2,067	3,203	5,058	6,524	6,324	3,994	1,411	254	1
2002.....	601	912	1,279	1,966	3,231	5,076	6,385	6,013	3,863	1,311	242	-
2001.....	619	860	1,199	2,077	3,260	4,898	5,865	5,717	3,651	1,324	250	1
2000.....	545	761	1,188	1,925	3,103	4,774	5,491	5,299	3,417	1,238	214	2
1999.....	482	722	1,093	1,870	3,102	4,551	5,150	4,900	3,212	1,178	198	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Renal failure (N17-N19)												
White, male												
2004.....	408	604	772	1,088	1,653	2,609	3,323	2,818	1,605	441	67	1
2003.....	385	592	764	1,116	1,681	2,657	3,295	2,903	1,571	417	66	1
2002.....	349	483	670	1,076	1,782	2,685	3,193	2,723	1,477	426	61	-
2001.....	363	477	644	1,153	1,736	2,559	2,912	2,644	1,434	381	56	-
2000.....	311	428	643	1,048	1,661	2,517	2,736	2,352	1,309	379	66	2
1999.....	269	415	601	986	1,675	2,426	2,570	2,204	1,196	376	55	-
White, female												
2004.....	288	454	660	950	1,462	2,435	3,176	3,373	2,491	1,047	190	1
2003.....	268	438	671	951	1,522	2,401	3,229	3,421	2,423	994	188	-
2002.....	252	429	609	890	1,449	2,391	3,192	3,290	2,386	885	181	-
2001.....	256	383	555	924	1,524	2,339	2,953	3,073	2,217	943	194	1
2000.....	234	333	545	877	1,442	2,257	2,755	2,947	2,108	859	148	-
1999.....	213	307	492	884	1,427	2,125	2,580	2,696	2,016	802	143	-
Black, both sexes												
2004.....	475	590	685	756	877	1,024	968	749	519	191	53	-
2003.....	444	540	634	790	953	992	1,014	772	488	190	46	1
2002.....	442	496	577	742	938	1,021	910	746	468	197	51	-
2001.....	387	457	586	807	884	982	908	708	455	175	48	-
2000.....	353	431	548	743	889	918	846	659	409	169	43	-
1999.....	335	374	494	692	855	902	836	716	392	172	54	-
Black, male												
2004.....	282	296	337	376	395	432	394	284	155	59	12	-
2003.....	284	309	321	376	468	412	445	253	168	58	13	1
2002.....	251	265	309	356	444	443	364	277	167	55	19	-
2001.....	204	238	268	390	386	404	383	259	143	39	7	-
2000.....	195	225	257	326	380	416	341	229	124	59	5	-
1999.....	184	169	252	310	380	431	368	285	112	29	12	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Renal failure (N17-N19)												
Black, female												
2004.....	193	294	348	380	482	592	574	465	364	132	41	-
2003.....	160	231	313	414	485	580	569	519	320	132	33	-
2002.....	191	231	268	386	494	578	546	469	301	142	32	-
2001.....	183	219	318	417	498	578	525	449	312	136	41	-
2000.....	158	206	291	417	509	502	505	430	285	110	38	-
1999.....	151	205	242	382	475	471	468	431	280	143	42	-
American Indian 1/, both sexes												
2004.....	19	19	19	30	29	34	24	24	8	4	1	-
2003.....	13	18	24	24	20	47	27	22	14	5	-	-
2002.....	12	21	21	24	32	22	21	14	11	2	2	-
2001.....	14	24	25	24	30	30	19	15	10	2	1	-
2000.....	16	16	21	35	29	21	24	10	13	2	-	-
1999.....	7	15	18	30	24	25	16	19	6	7	-	-
American Indian 1/, male												
2004.....	11	9	9	18	9	13	9	10	4	2	-	-
2003.....	7	11	10	11	6	27	14	10	4	3	-	-
2002.....	7	9	10	10	9	6	13	4	2	1	1	-
2001.....	10	5	10	7	12	15	6	10	4	-	-	-
2000.....	8	8	11	18	10	6	9	4	6	2	-	-
1999.....	4	7	8	12	10	11	1	5	1	2	-	-
American Indian 1/, female												
2004.....	8	10	10	12	20	21	15	14	4	2	1	-
2003.....	6	7	14	13	14	20	13	12	10	2	-	-
2002.....	5	12	11	14	23	16	8	10	9	1	1	-
2001.....	4	19	15	17	18	15	13	5	6	2	1	-
2000.....	8	8	10	17	19	15	15	6	7	-	-	-
1999.....	3	8	10	18	14	14	15	14	5	5	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Renal failure (N17-N19)												
Asian or Pacific Islander 2/, both sexes												
2004.....	18	38	42	70	82	120	117	98	53	12	6	-
2003.....	21	23	37	58	91	101	107	66	45	15	7	-
2002.....	17	35	42	42	71	109	113	83	56	20	1	-
2001.....	21	35	40	55	74	97	99	77	58	13	1	-
2000.....	18	26	32	51	73	81	68	72	39	11	8	-
1999.....	12	33	35	49	75	82	90	83	46	15	-	-
Asian or Pacific Islander 2/, male												
2004.....	9	16	29	36	41	59	56	45	29	5	4	-
2003.....	10	11	18	29	32	44	53	32	28	8	1	-
2002.....	8	15	25	26	36	56	48	42	26	10	-	-
2001.....	11	15	23	31	37	47	55	40	23	9	1	-
2000.....	12	17	18	20	29	42	35	32	19	3	2	-
1999.....	5	21	19	32	35	44	37	41	25	8	-	-
Asian or Pacific Islander 2/, female												
2004.....	9	22	13	34	41	61	61	53	24	7	2	-
2003.....	11	12	19	29	59	57	54	34	17	7	6	-
2002.....	9	20	17	16	35	53	65	41	30	10	1	-
2001.....	10	20	17	24	37	50	44	37	35	4	-	-
2000.....	6	9	14	31	44	39	33	40	20	8	6	-
1999.....	7	12	16	17	40	38	53	42	21	7	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Urolithiasis (N20-N23)												
All races, both sexes												
2004.....	11	14	15	28	25	45	55	48	31	8	1	-
2003.....	9	12	19	22	23	43	47	65	27	13	1	-
2002.....	8	13	11	22	23	38	48	49	17	15	2	-
2001.....	13	6	19	19	28	44	38	36	31	11	2	-
2000.....	11	10	8	16	21	50	45	43	27	6	-	-
1999.....	7	10	20	15	25	25	44	42	26	10	-	-
All races, male												
2004.....	5	9	7	7	10	23	19	17	8	3	-	-
2003.....	5	6	10	10	13	17	19	27	8	2	-	-
2002.....	5	3	5	11	9	17	18	21	2	5	-	-
2001.....	6	3	12	12	14	12	18	11	10	5	-	-
2000.....	9	3	3	4	11	15	21	12	7	2	-	-
1999.....	6	8	10	10	8	12	18	14	3	1	-	-
All races, female												
2004.....	6	5	8	21	15	22	36	31	23	5	1	-
2003.....	4	6	9	12	10	26	28	38	19	11	1	-
2002.....	3	10	6	11	14	21	30	28	15	10	2	-
2001.....	7	3	7	7	14	32	20	25	21	6	2	-
2000.....	2	7	5	12	10	35	24	31	20	4	-	-
1999.....	1	2	10	5	17	13	26	28	23	9	-	-
White, both sexes												
2004.....	9	10	14	27	21	40	50	44	29	7	1	-
2003.....	6	10	16	18	22	39	44	63	24	12	1	-
2002.....	6	11	9	20	22	34	44	46	16	14	2	-
2001.....	10	6	17	16	22	40	35	35	28	11	2	-
2000.....	10	6	8	15	18	45	44	38	25	5	-	-
1999.....	5	9	19	11	22	21	43	37	25	8	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Urolithiasis (N20-N23)												
White, male												
2004.....	5	7	7	7	9	20	16	17	8	2	-	-
2003.....	2	5	7	7	13	14	17	27	8	2	-	-
2002.....	4	3	3	11	8	16	15	19	2	5	-	-
2001.....	4	3	11	9	13	11	17	11	9	5	-	-
2000.....	8	2	3	3	9	15	21	12	7	2	-	-
1999.....	4	7	10	7	8	10	17	11	3	1	-	-
White, female												
2004.....	4	3	7	20	12	20	34	27	21	5	1	-
2003.....	4	5	9	11	9	25	27	36	16	10	1	-
2002.....	2	8	6	9	14	18	29	27	14	9	2	-
2001.....	6	3	6	7	9	29	18	24	19	6	2	-
2000.....	2	4	5	12	9	30	23	26	18	3	-	-
1999.....	1	2	9	4	14	11	26	26	22	7	-	-
Black, both sexes												
2004.....	1	3	-	-	4	5	3	4	2	-	-	-
2003.....	3	2	2	2	1	4	2	2	2	1	-	-
2002.....	1	1	2	2	1	3	4	3	1	1	-	-
2001.....	3	-	1	3	4	4	3	1	3	-	-	-
2000.....	1	3	-	-	3	5	-	1	2	1	-	-
1999.....	2	-	1	3	2	3	-	5	1	2	-	-
Black, male												
2004.....	-	1	-	-	1	3	2	-	-	-	-	-
2003.....	3	1	2	1	-	3	2	-	-	-	-	-
2002.....	1	-	2	-	1	1	3	2	-	-	-	-
2001.....	2	-	1	3	1	1	1	-	1	-	-	-
2000.....	1	1	-	-	2	-	-	-	-	-	-	-
1999.....	2	-	-	2	-	2	-	3	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Urolithiasis (N20-N23)												
Asian or Pacific Islander 2/, both sexes												
2004.....	1	1	1	1	-	-	1	-	-	1	-	-
2003.....	-	-	1	2	-	-	1	-	1	-	-	-
2002.....	1	1	-	-	-	1	-	-	-	-	-	-
2001.....	-	-	1	-	2	-	-	-	-	-	-	-
2000.....	-	1	-	1	-	-	1	4	-	-	-	-
1999.....	-	1	-	1	1	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	-	1	-	-	-	-	-	-	-	1	-	-
2003.....	-	-	1	2	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	1	-	-	-	-	-	-	-	-
1999.....	-	1	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	1	-	1	1	-	-	1	-	-	-	-	-
2003.....	-	-	-	-	-	-	1	-	1	-	-	-
2002.....	1	1	-	-	-	1	-	-	-	-	-	-
2001.....	-	-	1	-	2	-	-	-	-	-	-	-
2000.....	-	1	-	-	-	-	1	4	-	-	-	-
1999.....	-	-	-	-	1	1	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified disorders of kidney (N25-N27)												
All races, both sexes												
2004.....	3	6	5	7	7	14	12	11	4	-	2	-
2003.....	3	3	4	6	10	16	8	9	6	5	1	-
2002.....	3	4	4	7	8	12	13	16	8	2	-	-
2001.....	1	4	4	3	6	6	8	6	7	2	1	-
2000.....	-	3	8	3	13	12	16	13	3	3	1	-
1999.....	4	2	5	7	9	20	15	6	8	3	1	-
All races, male												
2004.....	2	5	2	3	3	8	5	5	2	-	1	-
2003.....	3	-	1	3	4	9	2	2	5	1	-	-
2002.....	1	2	-	1	6	8	5	3	3	1	-	-
2001.....	1	2	1	-	1	2	4	3	1	-	-	-
2000.....	-	2	4	2	7	5	6	8	2	1	-	-
1999.....	2	1	-	5	5	8	8	6	3	1	-	-
All races, female												
2004.....	1	1	3	4	4	6	7	6	2	-	1	-
2003.....	-	3	3	3	6	7	6	7	1	4	1	-
2002.....	2	2	4	6	2	4	8	13	5	1	-	-
2001.....	-	2	3	3	5	4	4	3	6	2	1	-
2000.....	-	1	4	1	6	7	10	5	1	2	1	-
1999.....	2	1	5	2	4	12	7	-	5	2	1	-
White, both sexes												
2004.....	2	4	5	6	7	12	11	11	4	-	2	-
2003.....	3	2	2	3	8	15	7	9	6	5	1	-
2002.....	1	4	4	6	8	8	11	14	8	1	-	-
2001.....	1	2	4	3	6	4	6	5	7	2	-	-
2000.....	-	3	7	1	11	9	15	12	3	3	-	-
1999.....	2	1	2	5	7	17	11	4	5	3	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified disorders of kidney (N25-N27)												
White, male												
2004.....	1	3	2	2	3	7	5	5	2	-	1	-
2003.....	3	-	-	1	3	8	2	2	5	1	-	-
2002.....	-	2	-	1	6	6	5	3	3	1	-	-
2001.....	1	1	1	-	1	1	3	2	1	-	-	-
2000.....	-	2	3	1	6	4	6	8	2	1	-	-
1999.....	1	1	-	3	4	6	7	4	3	1	-	-
White, female												
2004.....	1	1	3	4	4	5	6	6	2	-	1	-
2003.....	-	2	2	2	5	7	5	7	1	4	1	-
2002.....	1	2	4	5	2	2	6	11	5	-	-	-
2001.....	-	1	3	3	5	3	3	3	6	2	-	-
2000.....	-	1	4	-	5	5	9	4	1	2	-	-
1999.....	1	-	2	2	3	11	4	-	2	2	1	-
Black, both sexes												
2004.....	1	2	-	1	-	1	1	-	-	-	-	-
2003.....	-	1	2	3	1	1	1	-	-	-	-	-
2002.....	2	-	-	1	-	2	1	2	-	1	-	-
2001.....	-	1	-	-	-	1	1	1	-	-	1	-
2000.....	-	-	-	1	2	3	-	1	-	-	1	-
1999.....	2	1	3	2	1	3	4	1	2	-	-	-
Black, male												
2004.....	1	2	-	1	-	-	-	-	-	-	-	-
2003.....	-	-	1	2	1	1	-	-	-	-	-	-
2002.....	1	-	-	-	-	1	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	1	1	-	-	-	-
2000.....	-	-	-	1	1	1	-	-	-	-	-	-
1999.....	1	-	-	2	-	2	1	1	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified disorders of kidney (N25-N27)												
Asian or Pacific Islander 2/, both sexes												
2004.....	-	-	-	-	-	1	-	-	-	-	-	-
2003.....	-	-	-	-	1	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	2	1	-	-	-	-	-
2001.....	-	1	-	-	-	-	1	-	-	-	-	-
2000.....	-	-	-	1	-	-	1	-	-	-	-	-
1999.....	-	-	-	-	1	-	-	1	1	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	-	-	-	-	-	1	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	1	-	-	-	-	-	-
2001.....	-	1	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	1	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	1	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	1	1	-	-	-	-	-
2001.....	-	-	-	-	-	-	1	-	-	-	-	-
2000.....	-	-	-	1	-	-	1	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Urinary tract infection, site not specified (N39.0)												
All races, both sexes												
2004.....	171	229	279	527	887	1,735	2,888	3,181	2,536	1,038	218	1
2003.....	149	203	331	481	962	1,738	2,826	3,201	2,501	1,053	207	-
2002.....	124	215	281	462	941	1,760	2,776	3,228	2,542	1,003	219	1
2001.....	109	192	260	505	1,000	1,821	2,826	3,357	2,497	972	238	-
2000.....	115	155	267	463	878	1,896	2,853	3,145	2,512	1,027	243	-
1999.....	100	160	275	483	992	1,866	2,792	3,278	2,576	1,061	230	-
All races, male												
2004.....	68	106	118	226	343	687	990	957	608	177	33	-
2003.....	57	76	148	202	376	693	1,014	904	552	172	28	-
2002.....	50	89	120	191	367	629	947	892	595	189	40	1
2001.....	44	79	113	220	394	659	966	911	527	165	31	-
2000.....	52	66	110	200	358	711	962	902	592	185	36	-
1999.....	35	69	132	194	426	703	965	875	557	178	31	-
All races, female												
2004.....	103	123	161	301	544	1,048	1,898	2,224	1,928	861	185	1
2003.....	92	127	183	279	586	1,045	1,812	2,297	1,949	881	179	-
2002.....	74	126	161	271	574	1,131	1,829	2,336	1,947	814	179	-
2001.....	65	113	147	285	606	1,162	1,860	2,446	1,970	807	207	-
2000.....	63	89	157	263	520	1,185	1,891	2,243	1,920	842	207	-
1999.....	65	91	143	289	566	1,163	1,827	2,403	2,019	883	199	-
White, both sexes												
2004.....	128	162	222	420	736	1,471	2,542	2,897	2,266	930	178	1
2003.....	104	152	253	382	782	1,474	2,495	2,924	2,231	943	168	-
2002.....	89	170	216	359	780	1,507	2,442	2,903	2,304	879	177	1
2001.....	75	148	203	394	824	1,544	2,478	2,981	2,268	863	192	-
2000.....	80	123	200	370	709	1,623	2,508	2,819	2,251	927	216	-
1999.....	71	132	204	365	807	1,583	2,498	2,944	2,319	938	195	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Urinary tract infection, site not specified (N39.0)												
White, male												
2004.....	51	75	90	173	285	579	875	864	540	145	28	-
2003.....	41	58	113	154	304	583	896	832	505	161	25	-
2002.....	35	68	95	151	296	531	830	802	531	159	31	1
2001.....	32	55	89	171	320	560	864	803	468	139	26	-
2000.....	34	49	78	154	289	607	839	822	516	164	26	-
1999.....	29	60	95	139	328	581	857	770	479	153	24	-
White, female												
2004.....	77	87	132	247	451	892	1,667	2,033	1,726	785	150	1
2003.....	63	94	140	228	478	891	1,599	2,092	1,726	782	143	-
2002.....	54	102	121	208	484	976	1,612	2,101	1,773	720	146	-
2001.....	43	93	114	223	504	984	1,614	2,178	1,800	724	166	-
2000.....	46	74	122	216	420	1,016	1,669	1,997	1,735	763	190	-
1999.....	42	72	109	226	479	1,002	1,641	2,174	1,840	785	171	-
Black, both sexes												
2004.....	40	62	51	91	134	220	296	249	234	96	34	-
2003.....	39	48	71	88	160	237	289	243	241	95	33	-
2002.....	30	41	57	85	139	225	294	277	218	113	37	-
2001.....	29	36	49	89	151	240	309	332	208	97	42	-
2000.....	32	29	61	82	145	244	304	288	230	86	25	-
1999.....	25	24	64	108	160	256	257	294	231	111	32	-
Black, male												
2004.....	16	29	24	42	55	98	92	79	53	27	5	-
2003.....	15	16	32	43	64	99	107	67	41	9	3	-
2002.....	11	20	21	33	61	89	102	83	58	28	7	-
2001.....	10	20	22	42	60	89	90	97	53	25	5	-
2000.....	17	17	29	38	60	96	108	69	59	16	8	-
1999.....	6	9	33	50	88	109	92	86	69	23	6	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Urinary tract infection, site not specified (N39.0)												
Black, female												
2004.....	24	33	27	49	79	122	204	170	181	69	29	-
2003.....	24	32	39	45	96	138	182	176	200	86	30	-
2002.....	19	21	36	52	78	136	192	194	160	85	30	-
2001.....	19	16	27	47	91	151	219	235	155	72	37	-
2000.....	15	12	32	44	85	148	196	219	171	70	17	-
1999.....	19	15	31	58	72	147	165	208	162	88	26	-
American Indian 1/, both sexes												
2004.....	2	2	2	8	6	10	8	6	6	2	-	-
2003.....	4	2	2	6	5	8	16	4	5	6	3	-
2002.....	3	3	2	8	8	8	9	12	2	3	1	-
2001.....	3	5	4	8	7	12	12	9	5	3	1	-
2000.....	2	1	3	1	2	4	11	12	7	1	1	-
1999.....	4	3	2	-	3	6	5	4	4	1	3	-
American Indian 1/, male												
2004.....	1	-	1	6	-	1	5	1	3	1	-	-
2003.....	1	2	-	3	2	2	1	-	-	1	-	-
2002.....	2	1	-	2	3	1	3	2	2	-	1	-
2001.....	1	3	1	3	4	1	5	2	1	-	-	-
2000.....	1	-	1	1	1	1	3	4	3	-	1	-
1999.....	-	-	-	-	1	4	2	2	-	-	1	-
American Indian 1/, female												
2004.....	1	2	1	2	6	9	3	5	3	1	-	-
2003.....	3	-	2	3	3	6	15	4	5	5	3	-
2002.....	1	2	2	6	5	7	6	10	-	3	-	-
2001.....	2	2	3	5	3	11	7	7	4	3	1	-
2000.....	1	1	2	-	1	3	8	8	4	1	-	-
1999.....	4	3	2	-	2	2	3	2	4	1	2	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Urinary tract infection, site not specified (N39.0)												
Asian or Pacific Islander 2/, both sexes												
2004.....	1	3	4	8	11	34	42	29	30	10	6	-
2003.....	2	1	5	5	15	19	26	30	24	9	3	-
2002.....	2	1	6	10	14	20	31	36	18	8	4	-
2001.....	2	3	4	14	18	25	27	35	16	9	3	-
2000.....	1	2	3	10	22	25	30	26	24	13	1	-
1999.....	-	1	5	10	22	21	32	36	22	11	-	-
Asian or Pacific Islander 2/, male												
2004.....	-	2	3	5	3	9	18	13	12	4	-	-
2003.....	-	-	3	2	6	9	10	5	6	1	-	-
2002.....	2	-	4	5	7	8	12	5	4	2	1	-
2001.....	1	1	1	4	10	9	7	9	5	1	-	-
2000.....	-	-	2	7	8	7	12	7	14	5	1	-
1999.....	-	-	4	5	9	9	14	17	9	2	-	-
Asian or Pacific Islander 2/, female												
2004.....	1	1	1	3	8	25	24	16	18	6	6	-
2003.....	2	1	2	3	9	10	16	25	18	8	3	-
2002.....	-	1	2	5	7	12	19	31	14	6	3	-
2001.....	1	2	3	10	8	16	20	26	11	8	3	-
2000.....	1	2	1	3	14	18	18	19	10	8	-	-
1999.....	-	1	1	5	13	12	18	19	13	9	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of urinary system (N28-N36,N39.1-N39.9)												
All races, both sexes												
2004.....	47	64	76	92	114	157	213	196	128	44	9	-
2003.....	54	57	85	68	102	139	190	180	117	36	3	-
2002.....	53	45	64	85	102	126	209	180	123	36	6	-
2001.....	47	55	64	86	103	156	178	158	98	45	11	-
2000.....	49	56	60	78	108	148	192	180	94	39	4	-
1999.....	36	53	62	76	118	133	184	150	104	37	4	1
All races, male												
2004.....	31	43	45	48	63	83	114	84	43	10	1	-
2003.....	31	32	50	37	59	70	94	83	46	11	1	-
2002.....	35	27	39	44	53	65	98	74	38	11	2	-
2001.....	30	26	35	49	53	74	85	64	38	13	1	-
2000.....	27	30	33	47	57	69	103	78	32	15	2	-
1999.....	20	30	32	53	71	67	85	60	39	16	3	-
All races, female												
2004.....	16	21	31	44	51	74	99	112	85	34	8	-
2003.....	23	25	35	31	43	69	96	97	71	25	2	-
2002.....	18	18	25	41	49	61	111	106	85	25	4	-
2001.....	17	29	29	37	50	82	93	94	60	32	10	-
2000.....	22	26	27	31	51	79	89	102	62	24	2	-
1999.....	16	23	30	23	47	66	99	90	65	21	1	1
White, both sexes												
2004.....	28	46	57	73	86	133	185	173	120	39	8	-
2003.....	32	37	55	55	84	126	176	160	111	32	3	-
2002.....	37	32	41	69	84	108	183	159	115	31	5	-
2001.....	31	35	52	62	80	137	159	147	87	42	9	-
2000.....	40	38	49	58	81	128	168	163	80	34	3	-
1999.....	24	39	47	56	95	109	151	134	98	33	4	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of urinary system (N28-N36,N39.1-N39.9)												
White, male												
2004.....	19	29	39	39	49	72	108	76	41	9	1	-
2003.....	17	24	36	30	50	63	83	74	43	9	1	-
2002.....	26	20	25	38	46	56	87	65	35	10	1	-
2001.....	19	16	29	35	44	66	79	63	31	12	1	-
2000.....	24	20	24	36	45	58	95	71	27	13	2	-
1999.....	13	23	22	36	57	56	74	55	36	15	3	-
White, female												
2004.....	9	17	18	34	37	61	77	97	79	30	7	-
2003.....	15	13	19	25	34	63	93	86	68	23	2	-
2002.....	11	12	16	31	38	52	96	94	80	21	4	-
2001.....	12	19	23	27	36	71	80	84	56	30	8	-
2000.....	16	18	25	22	36	70	73	92	53	21	1	-
1999.....	11	16	25	20	38	53	77	79	62	18	1	-
Black, both sexes												
2004.....	18	17	16	19	27	18	26	21	8	5	1	-
2003.....	21	18	27	11	17	13	12	17	5	4	-	-
2002.....	13	13	20	15	15	15	25	16	8	5	-	-
2001.....	15	18	9	21	21	19	15	11	7	3	2	-
2000.....	8	15	7	17	23	17	19	16	14	5	1	-
1999.....	11	12	14	16	21	22	27	16	5	3	-	1
Black, male												
2004.....	11	13	4	9	13	10	6	7	2	1	-	-
2003.....	14	7	13	6	9	7	10	6	2	2	-	-
2002.....	7	7	12	5	7	7	11	7	3	1	-	-
2001.....	10	9	4	13	8	8	6	1	5	1	-	-
2000.....	2	9	5	11	9	11	6	6	5	2	-	-
1999.....	7	7	9	13	12	9	10	5	3	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of urinary system (N28-N36,N39.1-N39.9)												
Black, female												
2004.....	7	4	12	10	14	8	20	14	6	4	1	-
2003.....	7	11	14	5	8	6	2	11	3	2	-	-
2002.....	6	6	8	10	8	8	14	9	5	4	-	-
2001.....	5	9	5	8	13	11	9	10	2	2	2	-
2000.....	6	6	2	6	14	6	13	10	9	3	1	-
1999.....	4	5	5	3	9	13	17	11	2	2	-	1
American Indian 1/, both sexes												
2004.....	1	-	2	-	-	-	-	-	-	-	-	-
2003.....	-	1	1	1	1	-	-	1	-	-	-	-
2002.....	2	-	1	1	-	2	-	1	-	-	1	-
2001.....	-	-	1	2	-	-	-	-	2	-	-	-
2000.....	-	1	1	2	-	1	-	-	-	-	-	-
1999.....	-	-	-	1	1	1	-	-	1	-	-	-
American Indian 1/, male												
2004.....	1	-	2	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	1	-	-	-	1	-	-	-	-
2002.....	1	-	-	1	-	1	-	1	-	-	1	-
2001.....	-	-	1	1	-	-	-	-	1	-	-	-
2000.....	-	1	1	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	1	1	1	-	-	-	-	-	-
American Indian 1/, female												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	1	1	-	1	-	-	-	-	-	-	-
2002.....	1	-	1	-	-	1	-	-	-	-	-	-
2001.....	-	-	-	1	-	-	-	-	1	-	-	-
2000.....	-	-	-	2	-	1	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of urinary system (N28-N36,N39.1-N39.9)												
Asian or Pacific Islander 2/, both sexes												
2004.....	-	1	1	-	1	6	2	2	-	-	-	-
2003.....	1	1	2	1	-	-	2	2	1	-	-	-
2002.....	1	-	2	-	3	1	1	4	-	-	-	-
2001.....	1	2	2	1	2	-	4	-	2	-	-	-
2000.....	1	2	3	1	4	2	5	1	-	-	-	-
1999.....	1	2	1	3	1	1	6	-	-	1	-	-
Asian or Pacific Islander 2/, male												
2004.....	-	1	-	-	1	1	-	1	-	-	-	-
2003.....	-	1	1	-	-	-	1	2	1	-	-	-
2002.....	1	-	2	-	-	1	-	1	-	-	-	-
2001.....	1	1	1	-	1	-	-	-	1	-	-	-
2000.....	1	-	3	-	3	-	2	1	-	-	-	-
1999.....	-	-	1	3	1	1	1	-	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	-	-	1	-	-	5	2	1	-	-	-	-
2003.....	1	-	1	1	-	-	1	-	-	-	-	-
2002.....	-	-	-	-	3	-	1	3	-	-	-	-
2001.....	-	1	1	1	1	-	4	-	1	-	-	-
2000.....	-	2	-	1	1	2	3	-	-	-	-	-
1999.....	1	2	-	-	-	-	5	-	-	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of male genital organs (N40-N50)												
All races, both sexes												
2004.....	14	21	29	35	53	86	159	132	123	48	3	-
2003.....	5	17	17	34	42	80	121	135	105	41	2	-
2002.....	3	5	20	23	49	63	139	126	96	34	11	1
2001.....	10	12	15	22	54	80	107	131	103	36	6	-
2000.....	6	12	19	24	48	76	116	126	89	32	5	1
1999.....	10	11	20	31	51	79	130	134	80	29	10	-
All races, male												
2004.....	14	21	29	35	53	86	159	132	123	48	3	-
2003.....	5	17	17	34	42	80	121	135	105	41	2	-
2002.....	3	5	20	23	49	63	139	126	96	34	11	1
2001.....	10	12	15	22	54	80	107	131	103	36	6	-
2000.....	6	12	19	24	48	76	116	126	89	32	5	1
1999.....	10	11	20	31	51	79	130	134	80	29	10	-
All races, female												
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....
White, both sexes												
2004.....	8	14	24	28	43	71	142	123	112	45	3	-
2003.....	1	11	12	27	39	69	111	125	100	39	2	-
2002.....	1	5	14	18	38	47	125	112	92	31	10	-
2001.....	6	9	8	20	47	72	96	120	98	33	5	-
2000.....	3	7	14	20	40	63	107	113	80	29	5	-
1999.....	4	7	14	25	40	74	116	122	73	28	8	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of male genital organs (N40-N50)												
White, male												
2004.....	8	14	24	28	43	71	142	123	112	45	3	-
2003.....	1	11	12	27	39	69	111	125	100	39	2	-
2002.....	1	5	14	18	38	47	125	112	92	31	10	-
2001.....	6	9	8	20	47	72	96	120	98	33	5	-
2000.....	3	7	14	20	40	63	107	113	80	29	5	-
1999.....	4	7	14	25	40	74	116	122	73	28	8	-
White, female												
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....
Black, both sexes												
2004.....	6	5	4	6	8	13	10	6	9	2	-	-
2003.....	4	4	5	6	2	11	6	10	4	1	-	-
2002.....	2	-	6	4	11	13	11	12	3	1	1	1
2001.....	4	2	7	2	6	8	11	9	5	2	1	-
2000.....	3	5	5	4	7	11	8	9	7	2	-	1
1999.....	5	4	6	6	10	5	14	11	6	1	2	-
Black, male												
2004.....	6	5	4	6	8	13	10	6	9	2	-	-
2003.....	4	4	5	6	2	11	6	10	4	1	-	-
2002.....	2	-	6	4	11	13	11	12	3	1	1	1
2001.....	4	2	7	2	6	8	11	9	5	2	1	-
2000.....	3	5	5	4	7	11	8	9	7	2	-	1
1999.....	5	4	6	6	10	5	14	11	6	1	2	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hyperplasia of prostate (N40)												
All races, both sexes												
2004.....	3	3	10	14	31	58	119	115	109	41	3	-
2003.....	-	2	10	15	35	58	107	112	96	38	2	-
2002.....	-	1	9	9	32	48	116	93	85	31	11	1
2001.....	2	3	6	14	31	60	81	110	94	33	5	-
2000.....	-	4	7	10	28	64	95	106	83	29	4	1
1999.....	1	4	9	15	32	59	101	109	67	24	9	-
All races, male												
2004.....	3	3	10	14	31	58	119	115	109	41	3	-
2003.....	-	2	10	15	35	58	107	112	96	38	2	-
2002.....	-	1	9	9	32	48	116	93	85	31	11	1
2001.....	2	3	6	14	31	60	81	110	94	33	5	-
2000.....	-	4	7	10	28	64	95	106	83	29	4	1
1999.....	1	4	9	15	32	59	101	109	67	24	9	-
All races, female												
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....
White, both sexes												
2004.....	1	3	9	13	26	51	105	108	100	38	3	-
2003.....	-	1	8	10	34	52	98	105	92	36	2	-
2002.....	-	1	6	8	26	38	107	83	82	28	10	-
2001.....	1	2	3	12	28	55	74	102	90	31	5	-
2000.....	-	2	5	10	25	57	88	96	74	27	4	-
1999.....	1	3	6	12	28	56	91	99	63	24	7	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hyperplasia of prostate (N40)												
White, male												
2004.....	1	3	9	13	26	51	105	108	100	38	3	-
2003.....	-	1	8	10	34	52	98	105	92	36	2	-
2002.....	-	1	6	8	26	38	107	83	82	28	10	-
2001.....	1	2	3	12	28	55	74	102	90	31	5	-
2000.....	-	2	5	10	25	57	88	96	74	27	4	-
1999.....	1	3	6	12	28	56	91	99	63	24	7	-
White, female												
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....
Black, both sexes												
2004.....	2	-	1	1	5	5	8	4	7	2	-	-
2003.....	-	1	2	4	1	6	5	7	3	1	-	-
2002.....	-	-	3	1	6	7	7	9	2	1	1	1
2001.....	1	1	3	2	2	5	7	6	4	1	-	-
2000.....	-	2	2	-	2	7	7	7	7	1	-	1
1999.....	-	1	3	3	3	3	10	9	4	-	2	-
Black, male												
2004.....	2	-	1	1	5	5	8	4	7	2	-	-
2003.....	-	1	2	4	1	6	5	7	3	1	-	-
2002.....	-	-	3	1	6	7	7	9	2	1	1	1
2001.....	1	1	3	2	2	5	7	6	4	1	-	-
2000.....	-	2	2	-	2	7	7	7	7	1	-	1
1999.....	-	1	3	3	3	3	10	9	4	-	2	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of male genital organs (N41-N50)												
All races, both sexes												
2004.....	11	18	19	21	22	28	40	17	14	7	-	-
2003.....	5	15	7	19	7	22	14	23	9	3	-	-
2002.....	3	4	11	14	17	15	23	33	11	3	-	-
2001.....	8	9	9	8	23	20	26	21	9	3	1	-
2000.....	6	8	12	14	20	12	21	20	6	3	1	-
1999.....	9	7	11	16	19	20	29	25	13	5	1	-
All races, male												
2004.....	11	18	19	21	22	28	40	17	14	7	-	-
2003.....	5	15	7	19	7	22	14	23	9	3	-	-
2002.....	3	4	11	14	17	15	23	33	11	3	-	-
2001.....	8	9	9	8	23	20	26	21	9	3	1	-
2000.....	6	8	12	14	20	12	21	20	6	3	1	-
1999.....	9	7	11	16	19	20	29	25	13	5	1	-
All races, female												
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....
White, both sexes												
2004.....	7	11	15	15	17	20	37	15	12	7	-	-
2003.....	1	10	4	17	5	17	13	20	8	3	-	-
2002.....	1	4	8	10	12	9	18	29	10	3	-	-
2001.....	5	7	5	8	19	17	22	18	8	2	-	-
2000.....	3	5	9	10	15	6	19	17	6	2	1	-
1999.....	3	4	8	13	12	18	25	23	10	4	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of male genital organs (N41-N50)												
White, male												
2004.....	7	11	15	15	17	20	37	15	12	7	-	-
2003.....	1	10	4	17	5	17	13	20	8	3	-	-
2002.....	1	4	8	10	12	9	18	29	10	3	-	-
2001.....	5	7	5	8	19	17	22	18	8	2	-	-
2000.....	3	5	9	10	15	6	19	17	6	2	1	-
1999.....	3	4	8	13	12	18	25	23	10	4	1	-
White, female												
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....
Black, both sexes												
2004.....	4	5	3	5	3	8	2	2	2	-	-	-
2003.....	4	3	3	2	1	5	1	3	1	-	-	-
2002.....	2	-	3	3	5	6	4	3	1	-	-	-
2001.....	3	1	4	-	4	3	4	3	1	1	1	-
2000.....	3	3	3	4	5	4	1	2	-	1	-	-
1999.....	5	3	3	3	7	2	4	2	2	1	-	-
Black, male												
2004.....	4	5	3	5	3	8	2	2	2	-	-	-
2003.....	4	3	3	2	1	5	1	3	1	-	-	-
2002.....	2	-	3	3	5	6	4	3	1	-	-	-
2001.....	3	1	4	-	4	3	4	3	1	1	1	-
2000.....	3	3	3	4	5	4	1	2	-	1	-	-
1999.....	5	3	3	3	7	2	4	2	2	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of breast (N60-N64)												
All races, both sexes												
2004.....	3	4	4	2	1	4	7	13	12	6	2	-
2003.....	2	3	2	5	4	3	5	10	8	5	-	-
2002.....	3	4	2	3	4	3	3	6	9	7	1	-
2001.....	-	3	3	4	8	8	8	13	11	1	2	-
2000.....	3	1	-	2	2	4	8	7	4	6	2	-
1999.....	1	1	2	-	3	6	5	8	5	4	-	-
All races, male												
2004.....	-	-	-	-	1	1	-	-	-	1	-	-
2003.....	-	-	-	-	-	-	2	-	-	-	-	-
2002.....	-	-	-	1	1	1	-	-	-	-	-	-
2001.....	-	-	-	1	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	1	-	-	-	-	-	-
1999.....	1	-	-	-	-	-	-	-	1	-	-	-
All races, female												
2004.....	3	4	4	2	-	3	7	13	12	5	2	-
2003.....	2	3	2	5	4	3	3	10	8	5	-	-
2002.....	3	4	2	2	3	2	3	6	9	7	1	-
2001.....	-	3	3	3	8	8	8	13	11	1	2	-
2000.....	3	1	-	2	2	3	8	7	4	6	2	-
1999.....	-	1	2	-	3	6	5	8	4	4	-	-
White, both sexes												
2004.....	1	4	4	2	1	2	6	11	12	6	2	-
2003.....	1	2	1	2	3	3	5	9	5	5	-	-
2002.....	3	3	2	-	4	3	3	5	7	6	1	-
2001.....	-	3	1	4	8	6	8	12	9	1	2	-
2000.....	3	1	-	-	2	4	8	4	4	5	2	-
1999.....	1	1	2	-	1	3	4	8	4	3	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of breast (N60-N64)												
Black, female												
2004.....	2	-	-	-	-	1	1	2	-	-	-	-
2003.....	1	-	1	2	1	-	-	1	3	-	-	-
2002.....	-	1	-	2	-	-	-	1	2	1	-	-
2001.....	-	-	2	-	-	2	-	1	2	-	-	-
2000.....	-	-	-	2	-	-	-	3	-	1	-	-
1999.....	-	-	-	-	2	2	1	-	1	1	-	-
American Indian 1/, both sexes												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, male												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, female												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	1	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of female genital organs (N70-N98)												
All races, both sexes												
2004.....	14	13	11	18	30	31	51	68	34	18	3	-
2003.....	17	9	17	20	21	46	52	61	38	20	2	-
2002.....	13	11	8	12	14	33	46	53	38	15	3	-
2001.....	18	10	11	17	31	38	47	45	31	16	1	-
2000.....	15	7	7	21	27	39	50	50	36	15	6	-
1999.....	10	14	14	16	22	43	40	38	26	15	6	-
All races, male												
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....
All races, female												
2004.....	14	13	11	18	30	31	51	68	34	18	3	-
2003.....	17	9	17	20	21	46	52	61	38	20	2	-
2002.....	13	11	8	12	14	33	46	53	38	15	3	-
2001.....	18	10	11	17	31	38	47	45	31	16	1	-
2000.....	15	7	7	21	27	39	50	50	36	15	6	-
1999.....	10	14	14	16	22	43	40	38	26	15	6	-
White, both sexes												
2004.....	11	11	10	13	20	30	45	60	30	16	3	-
2003.....	12	8	12	16	17	43	49	60	36	20	2	-
2002.....	9	10	5	10	10	31	44	45	33	13	3	-
2001.....	12	7	7	14	25	31	42	38	28	15	1	-
2000.....	14	5	7	17	23	36	43	47	33	14	5	-
1999.....	8	12	11	11	17	32	38	31	26	14	5	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Inflammatory diseases of female pelvic organs (N70-N76)												
All races, both sexes												
2004.....	3	8	7	7	9	10	11	17	11	6	-	-
2003.....	7	3	7	6	10	17	12	18	9	4	-	-
2002.....	7	4	2	4	4	17	12	22	11	3	-	-
2001.....	3	4	8	8	10	13	18	23	11	4	-	-
2000.....	3	4	4	11	10	17	12	11	12	3	1	-
1999.....	2	6	7	4	11	15	17	6	6	5	1	-
All races, male												
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....
All races, female												
2004.....	3	8	7	7	9	10	11	17	11	6	-	-
2003.....	7	3	7	6	10	17	12	18	9	4	-	-
2002.....	7	4	2	4	4	17	12	22	11	3	-	-
2001.....	3	4	8	8	10	13	18	23	11	4	-	-
2000.....	3	4	4	11	10	17	12	11	12	3	1	-
1999.....	2	6	7	4	11	15	17	6	6	5	1	-
White, both sexes												
2004.....	2	6	7	6	8	9	7	16	10	5	-	-
2003.....	4	2	6	5	8	15	10	17	9	4	-	-
2002.....	5	3	2	4	3	16	12	19	10	3	-	-
2001.....	2	4	4	6	8	11	16	22	8	4	-	-
2000.....	3	2	4	9	8	15	11	9	12	2	1	-
1999.....	2	6	6	2	8	10	16	6	6	5	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Noninflammatory disorders of female genital tract (N80-N98)												
All races, both sexes												
2004.....	11	5	4	11	21	21	40	51	23	12	3	-
2003.....	10	6	10	14	11	29	40	43	29	16	2	-
2002.....	6	7	6	8	10	16	34	31	27	12	3	-
2001.....	15	6	3	9	21	25	29	22	20	12	1	-
2000.....	12	3	3	10	17	22	38	39	24	12	5	-
1999.....	8	8	7	12	11	28	23	32	20	10	5	-
All races, male												
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....
All races, female												
2004.....	11	5	4	11	21	21	40	51	23	12	3	-
2003.....	10	6	10	14	11	29	40	43	29	16	2	-
2002.....	6	7	6	8	10	16	34	31	27	12	3	-
2001.....	15	6	3	9	21	25	29	22	20	12	1	-
2000.....	12	3	3	10	17	22	38	39	24	12	5	-
1999.....	8	8	7	12	11	28	23	32	20	10	5	-
White, both sexes												
2004.....	9	5	3	7	12	21	38	44	20	11	3	-
2003.....	8	6	6	11	9	28	39	43	27	16	2	-
2002.....	4	7	3	6	7	15	32	26	23	10	3	-
2001.....	10	3	3	8	17	20	26	16	20	11	1	-
2000.....	11	3	3	8	15	21	32	38	21	12	4	-
1999.....	6	6	5	9	9	22	22	25	20	9	5	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XVI. Certain conditions originating in the perinatal period (P00-P96)												
All races, both sexes												
2004.....	1	1	2	1	-	1	-	1	-	-	-	-
2003.....	1	-	-	-	2	-	-	-	-	-	-	-
2002.....	-	1	-	1	2	-	2	-	-	1	-	2
2001.....	2	2	-	2	1	1	-	1	-	-	-	1
2000.....	1	1	1	-	-	-	-	-	-	-	-	1
1999.....	-	1	2	1	1	1	-	-	1	-	-	5
All races, male												
2004.....	-	1	2	1	-	1	-	-	-	-	-	-
2003.....	1	-	-	-	2	-	-	-	-	-	-	-
2002.....	-	1	-	-	1	-	1	-	-	-	-	2
2001.....	2	1	-	1	1	1	-	-	-	-	-	1
2000.....	1	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	1	-	1	1	-	-	1	-	-	4
All races, female												
2004.....	1	-	-	-	-	-	-	1	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	1	1	-	1	-	-	1	-	-
2001.....	-	1	-	1	-	-	-	1	-	-	-	-
2000.....	-	1	1	-	-	-	-	-	-	-	-	1
1999.....	-	1	1	1	-	-	-	-	-	-	-	1
White, both sexes												
2004.....	1	1	2	-	-	1	-	1	-	-	-	-
2003.....	-	-	-	-	1	-	-	-	-	-	-	-
2002.....	-	1	-	1	1	-	2	-	-	1	-	1
2001.....	2	1	-	2	1	1	-	1	-	-	-	1
2000.....	1	1	1	-	-	-	-	-	-	-	-	1
1999.....	-	1	2	1	1	1	-	-	1	-	-	3

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intrauterine hypoxia and birth asphyxia (P20-P21)												
All races, both sexes												
2004.....	1	-	1	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	2	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	1	-	1	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	1	-	-	-	-	-	1
All races, male												
2004.....	-	-	1	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	2	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	1	-	1	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	1	-	-	-	-	-	1
All races, female												
2004.....	1	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes												
2004.....	1	-	1	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	1	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	1	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	1	-	-	-	-	-	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other and unspecified conditions originating in the perinatal period (P05,P08,P29,P50-P96)

All races, both sexes

2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	1	-	1	-	-	1	-	-	1	-	-
2001.....	1	-	-	1	-	1	-	1	-	-	-	-
2000.....	1	-	1	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	1	-	-	-	1	-	-	-

All races, male

2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	1	-	-	-	-	-	-	-	-	-	-
2001.....	1	-	-	-	-	1	-	-	-	-	-	-
2000.....	1	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	1	-	-	-	1	-	-	-

All races, female

2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	1	-	-	1	-	-	1	-	-
2001.....	-	-	-	1	-	-	-	1	-	-	-	-
2000.....	-	-	1	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-

White, both sexes

2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	1	-	1	-	-	1	-	-	1	-	-
2001.....	1	-	-	1	-	1	-	1	-	-	-	-
2000.....	1	-	1	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	1	-	-	-	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XVII. Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)												
All races, both sexes												
2004.....	358	360	274	230	198	232	240	169	90	28	9	-
2003.....	343	376	290	223	190	224	240	170	100	40	9	-
2002.....	363	351	283	224	216	232	257	184	86	33	2	1
2001.....	333	298	283	214	234	244	221	182	87	38	10	-
2000.....	325	296	273	248	216	242	246	178	97	39	6	1
1999.....	305	304	253	251	218	249	242	187	100	34	5	2
All races, male												
2004.....	186	187	133	106	102	114	81	69	27	5	1	-
2003.....	172	189	147	108	90	86	92	57	27	12	1	-
2002.....	186	170	151	107	106	105	96	65	22	9	1	1
2001.....	167	146	130	109	119	110	93	70	17	9	4	-
2000.....	190	130	134	120	102	99	98	74	41	13	2	-
1999.....	162	157	126	106	103	102	102	60	35	6	1	1
All races, female												
2004.....	172	173	141	124	96	118	159	100	63	23	8	-
2003.....	171	187	143	115	100	138	148	113	73	28	8	-
2002.....	177	181	132	117	110	127	161	119	64	24	1	-
2001.....	166	152	153	105	115	134	128	112	70	29	6	-
2000.....	135	166	139	128	114	143	148	104	56	26	4	1
1999.....	143	147	127	145	115	147	140	127	65	28	4	1
White, both sexes												
2004.....	316	324	250	208	184	205	219	158	85	28	7	-
2003.....	305	335	261	191	169	204	220	160	90	36	9	-
2002.....	314	304	248	202	191	212	232	170	83	30	2	1
2001.....	293	267	260	193	210	226	203	172	81	34	6	-
2000.....	294	255	241	219	191	225	232	168	94	34	5	1
1999.....	265	275	225	225	191	221	224	170	92	34	5	2

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XVII. Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)												
White, male												
2004.....	164	166	122	96	96	99	75	66	27	5	-	-
2003.....	155	169	129	92	77	78	79	55	24	10	1	-
2002.....	157	146	132	99	91	95	83	61	22	8	1	1
2001.....	148	130	119	103	110	106	81	64	14	9	1	-
2000.....	168	114	123	99	89	94	94	70	40	12	1	-
1999.....	140	143	114	95	91	83	96	53	33	6	1	1
White, female												
2004.....	152	158	128	112	88	106	144	92	58	23	7	-
2003.....	150	166	132	99	92	126	141	105	66	26	8	-
2002.....	157	158	116	103	100	117	149	109	61	22	1	-
2001.....	145	137	141	90	100	120	122	108	67	25	5	-
2000.....	126	141	118	120	102	131	138	98	54	22	4	1
1999.....	125	132	111	130	100	138	128	117	59	28	4	1
Black, both sexes												
2004.....	33	24	23	17	11	22	17	9	3	-	2	-
2003.....	28	34	26	24	16	15	15	8	8	4	-	-
2002.....	40	41	28	20	20	14	18	13	2	3	-	-
2001.....	33	27	21	15	21	17	11	7	5	4	4	-
2000.....	28	32	27	25	18	14	11	7	3	5	1	-
1999.....	33	25	21	21	25	22	16	16	8	-	-	-
Black, male												
2004.....	17	14	11	9	5	13	5	3	-	-	1	-
2003.....	10	18	15	14	11	5	11	2	2	2	-	-
2002.....	22	22	18	8	13	7	9	4	-	1	-	-
2001.....	17	13	10	4	8	4	8	4	2	-	3	-
2000.....	19	11	7	18	8	5	4	3	1	1	1	-
1999.....	17	12	10	6	10	14	5	7	2	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XVII. Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)												
Asian or Pacific Islander 2/, both sexes												
2004.....	8	9	1	5	3	5	4	2	2	-	-	-
2003.....	8	5	2	6	4	5	5	-	2	-	-	-
2002.....	9	4	5	2	5	5	3	1	1	-	-	-
2001.....	6	3	2	6	2	1	5	3	1	-	-	-
2000.....	2	7	2	2	4	2	2	3	-	-	-	-
1999.....	4	1	5	4	2	5	2	1	-	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	5	6	-	1	1	2	1	-	-	-	-	-
2003.....	6	1	2	2	2	3	2	-	1	-	-	-
2002.....	7	1	1	-	2	3	2	-	-	-	-	-
2001.....	2	3	1	2	-	-	4	2	1	-	-	-
2000.....	2	4	1	2	3	-	-	1	-	-	-	-
1999.....	2	-	2	4	2	4	1	-	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	3	3	1	4	2	3	3	2	2	-	-	-
2003.....	2	4	-	4	2	2	3	-	1	-	-	-
2002.....	2	3	4	2	3	2	1	1	1	-	-	-
2001.....	4	-	1	4	2	1	1	1	-	-	-	-
2000.....	-	3	1	-	1	2	2	2	-	-	-	-
1999.....	2	1	3	-	-	1	1	1	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Spina bifida (Q05)												
All races, both sexes												
2004.....	5	3	2	3	2	1	1	2	-	-	-	-
2003.....	7	2	3	1	2	-	2	3	-	-	-	-
2002.....	11	6	1	1	3	2	1	-	-	-	-	-
2001.....	7	1	4	-	1	1	-	1	-	-	-	-
2000.....	5	2	1	1	3	1	1	-	-	-	-	-
1999.....	4	4	1	-	-	2	-	1	1	-	-	-
All races, male												
2004.....	2	1	-	1	1	1	-	-	-	-	-	-
2003.....	4	1	1	-	1	-	1	-	-	-	-	-
2002.....	7	1	-	1	2	1	-	-	-	-	-	-
2001.....	4	-	2	-	-	-	-	-	-	-	-	-
2000.....	2	2	1	1	2	-	1	-	-	-	-	-
1999.....	2	1	1	-	-	1	-	1	-	-	-	-
All races, female												
2004.....	3	2	2	2	1	-	1	2	-	-	-	-
2003.....	3	1	2	1	1	-	1	3	-	-	-	-
2002.....	4	5	1	-	1	1	1	-	-	-	-	-
2001.....	3	1	2	-	1	1	-	1	-	-	-	-
2000.....	3	-	-	-	1	1	-	-	-	-	-	-
1999.....	2	3	-	-	-	1	-	-	1	-	-	-
White, both sexes												
2004.....	5	3	1	3	2	1	1	2	-	-	-	-
2003.....	7	2	3	1	2	-	2	3	-	-	-	-
2002.....	10	6	1	1	2	2	1	-	-	-	-	-
2001.....	7	1	4	-	1	1	-	1	-	-	-	-
2000.....	5	2	1	1	3	1	1	-	-	-	-	-
1999.....	4	4	1	-	-	2	-	1	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other congenital malformations of nervous system (Q01-Q04,Q06-Q07)												
All races, both sexes												
2004.....	6	11	9	9	5	3	10	1	1	-	-	-
2003.....	13	7	5	9	6	4	9	2	1	1	1	-
2002.....	11	9	8	8	4	5	7	1	-	-	-	-
2001.....	11	6	9	8	8	9	5	5	-	-	-	-
2000.....	6	3	7	7	5	7	4	4	-	-	-	-
1999.....	13	7	8	8	3	7	6	3	-	-	-	-
All races, male												
2004.....	3	5	3	2	3	2	3	1	-	-	-	-
2003.....	6	3	2	2	2	-	5	1	-	-	1	-
2002.....	4	4	5	6	1	4	4	-	-	-	-	-
2001.....	7	4	3	3	5	3	3	1	-	-	-	-
2000.....	4	2	4	7	2	3	1	3	-	-	-	-
1999.....	7	3	5	6	2	1	4	1	-	-	-	-
All races, female												
2004.....	3	6	6	7	2	1	7	-	1	-	-	-
2003.....	7	4	3	7	4	4	4	1	1	1	-	-
2002.....	7	5	3	2	3	1	3	1	-	-	-	-
2001.....	4	2	6	5	3	6	2	4	-	-	-	-
2000.....	2	1	3	-	3	4	3	1	-	-	-	-
1999.....	6	4	3	2	1	6	2	2	-	-	-	-
White, both sexes												
2004.....	5	10	6	7	4	3	8	1	-	-	-	-
2003.....	12	6	3	8	6	4	8	2	1	1	1	-
2002.....	10	8	7	8	3	5	7	1	-	-	-	-
2001.....	7	6	9	7	8	8	5	5	-	-	-	-
2000.....	6	3	5	6	4	6	4	4	-	-	-	-
1999.....	9	5	7	7	3	7	5	3	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Congenital malformations of heart (Q20-Q24)												
All races, both sexes												
2004.....	105	83	56	48	49	64	72	45	17	9	4	-
2003.....	86	80	59	69	51	63	82	49	35	11	2	-
2002.....	97	86	53	55	64	75	84	48	42	14	-	-
2001.....	111	65	55	58	63	82	71	62	30	16	6	-
2000.....	93	85	63	65	68	78	97	54	35	14	2	-
1999.....	95	85	70	83	78	89	85	84	40	13	2	-
All races, male												
2004.....	57	47	27	16	29	27	21	13	8	1	-	-
2003.....	37	43	26	34	18	18	27	9	8	2	-	-
2002.....	44	43	27	23	26	31	26	9	7	4	-	-
2001.....	52	36	24	29	37	30	30	16	1	3	2	-
2000.....	50	37	33	37	30	35	37	18	12	2	1	-
1999.....	55	40	36	32	36	31	32	21	13	3	-	-
All races, female												
2004.....	48	36	29	32	20	37	51	32	9	8	4	-
2003.....	49	37	33	35	33	45	55	40	27	9	2	-
2002.....	53	43	26	32	38	44	58	39	35	10	-	-
2001.....	59	29	31	29	26	52	41	46	29	13	4	-
2000.....	43	48	30	28	38	43	60	36	23	12	1	-
1999.....	40	45	34	51	42	58	53	63	27	10	2	-
White, both sexes												
2004.....	94	76	53	42	43	61	68	41	16	9	4	-
2003.....	77	71	51	57	50	56	78	49	32	11	2	-
2002.....	85	76	42	48	58	66	77	44	40	14	-	-
2001.....	96	54	48	53	57	77	68	57	29	15	4	-
2000.....	86	72	54	55	58	74	93	53	35	12	2	-
1999.....	80	77	59	73	68	83	81	74	37	13	2	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Congenital malformations of heart (Q20-Q24)												
White, male												
2004.....	51	45	25	15	25	27	21	13	8	1	-	-
2003.....	33	38	21	30	18	16	24	9	7	2	-	-
2002.....	37	39	21	22	24	27	24	9	7	4	-	-
2001.....	45	32	21	27	35	30	28	13	1	3	-	-
2000.....	45	31	31	29	24	34	36	17	12	2	1	-
1999.....	44	38	31	27	32	25	30	17	12	3	-	-
White, female												
2004.....	43	31	28	27	18	34	47	28	8	8	4	-
2003.....	44	33	30	27	32	40	54	40	25	9	2	-
2002.....	48	37	21	26	34	39	53	35	33	10	-	-
2001.....	51	22	27	26	22	47	40	44	28	12	4	-
2000.....	41	41	23	26	34	40	57	36	23	10	1	-
1999.....	36	39	28	46	36	58	51	57	25	10	2	-
Black, both sexes												
2004.....	8	5	2	3	6	3	3	3	1	-	-	-
2003.....	6	8	7	8	-	4	3	-	2	-	-	-
2002.....	9	8	8	6	3	5	4	4	1	-	-	-
2001.....	13	10	7	3	5	5	1	3	1	1	2	-
2000.....	6	10	7	9	7	2	2	1	-	2	-	-
1999.....	10	8	9	8	9	4	4	10	3	-	-	-
Black, male												
2004.....	4	1	2	1	4	-	-	-	-	-	-	-
2003.....	2	5	4	3	-	-	2	-	1	-	-	-
2002.....	4	3	6	1	2	2	1	-	-	-	-	-
2001.....	6	3	3	1	2	-	1	1	-	-	2	-
2000.....	4	5	-	8	3	1	1	1	-	-	-	-
1999.....	8	2	4	3	3	4	2	4	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other congenital malformations of circulatory system (Q25-Q28)												
All races, both sexes												
2004.....	42	43	31	33	36	64	56	39	31	6	2	-
2003.....	48	37	47	36	45	49	51	41	14	13	2	-
2002.....	45	45	34	43	51	63	69	48	17	5	1	-
2001.....	32	39	31	39	66	48	46	39	26	11	1	-
2000.....	43	35	44	31	51	54	59	56	23	9	-	-
1999.....	49	42	35	57	41	50	60	37	25	7	2	-
All races, male												
2004.....	26	20	17	17	17	35	21	20	6	1	1	-
2003.....	27	18	28	18	23	26	26	17	2	5	-	-
2002.....	22	24	22	20	30	23	27	19	3	2	1	-
2001.....	16	25	15	19	30	29	23	16	9	2	-	-
2000.....	31	14	24	12	31	23	25	23	8	6	-	-
1999.....	29	27	14	25	15	27	26	14	10	1	1	-
All races, female												
2004.....	16	23	14	16	19	29	35	19	25	5	1	-
2003.....	21	19	19	18	22	23	25	24	12	8	2	-
2002.....	23	21	12	23	21	40	42	29	14	3	-	-
2001.....	16	14	16	20	36	19	23	23	17	9	1	-
2000.....	12	21	20	19	20	31	34	33	15	3	-	-
1999.....	20	15	21	32	26	23	34	23	15	6	1	-
White, both sexes												
2004.....	33	33	26	29	34	52	53	38	31	6	1	-
2003.....	38	32	40	30	36	44	45	38	12	11	2	-
2002.....	31	32	28	38	43	55	61	46	16	4	1	-
2001.....	26	34	27	34	55	46	45	35	23	10	1	-
2000.....	32	26	35	27	45	51	55	54	21	8	-	-
1999.....	40	34	28	49	34	40	54	34	22	7	2	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other congenital malformations of circulatory system (Q25-Q28)												
White, male												
2004.....	21	14	14	14	16	27	20	20	6	1	-	-
2003.....	21	17	23	14	15	23	23	15	2	5	-	-
2002.....	14	18	18	17	23	19	22	19	3	1	1	-
2001.....	13	21	13	18	26	27	23	14	7	2	-	-
2000.....	21	10	21	9	28	23	25	23	8	5	-	-
1999.....	26	20	12	21	13	21	23	12	10	1	1	-
White, female												
2004.....	12	19	12	15	18	25	33	18	25	5	1	-
2003.....	17	15	17	16	21	21	22	23	10	6	2	-
2002.....	17	14	10	21	20	36	39	27	13	3	-	-
2001.....	13	13	14	16	29	19	22	21	16	8	1	-
2000.....	11	16	14	18	17	28	30	31	13	3	-	-
1999.....	14	14	16	28	21	19	31	22	12	6	1	-
Black, both sexes												
2004.....	7	5	5	4	1	11	3	-	-	-	1	-
2003.....	6	4	6	5	8	5	4	3	2	2	-	-
2002.....	12	11	4	5	7	6	7	2	1	1	-	-
2001.....	5	4	4	2	9	2	1	3	3	1	-	-
2000.....	10	7	8	2	4	2	3	1	2	1	-	-
1999.....	8	7	4	6	7	8	5	3	3	-	-	-
Black, male												
2004.....	4	2	3	3	-	8	1	-	-	-	1	-
2003.....	3	1	4	3	7	3	3	2	-	-	-	-
2002.....	6	5	3	3	6	3	4	-	-	1	-	-
2001.....	3	4	2	-	3	2	-	2	2	-	-	-
2000.....	9	3	2	1	2	-	-	-	-	1	-	-
1999.....	2	6	2	2	2	5	2	2	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other congenital malformations of circulatory system (Q25-Q28)												
Asian or Pacific Islander 2/, both sexes												
2004.....	1	4	-	-	1	1	-	1	-	-	-	-
2003.....	3	1	-	1	1	-	2	-	-	-	-	-
2002.....	2	1	2	-	1	2	-	-	-	-	-	-
2001.....	1	-	-	3	1	-	-	1	-	-	-	-
2000.....	1	2	-	1	2	1	-	1	-	-	-	-
1999.....	-	-	2	2	-	2	1	-	-	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	1	3	-	-	1	-	-	-	-	-	-	-
2003.....	2	-	-	1	1	-	-	-	-	-	-	-
2002.....	2	-	1	-	1	1	-	-	-	-	-	-
2001.....	-	-	-	1	-	-	-	-	-	-	-	-
2000.....	1	1	-	1	1	-	-	-	-	-	-	-
1999.....	-	-	-	2	-	1	1	-	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	-	1	-	-	-	1	-	1	-	-	-	-
2003.....	1	1	-	-	-	-	2	-	-	-	-	-
2002.....	-	1	1	-	-	1	-	-	-	-	-	-
2001.....	1	-	-	2	1	-	-	1	-	-	-	-
2000.....	-	1	-	-	1	1	-	1	-	-	-	-
1999.....	-	-	2	-	-	1	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Congenital malformations of respiratory system (Q30-Q34)												
All races, both sexes												
2004.....	5	-	2	2	3	4	4	1	-	2	-	-
2003.....	2	2	4	1	-	1	1	4	-	1	-	-
2002.....	-	-	2	3	-	3	2	2	-	-	-	-
2001.....	1	2	4	-	4	1	3	3	3	-	-	-
2000.....	2	1	3	7	-	3	2	2	-	-	-	-
1999.....	1	3	1	-	4	3	2	-	-	1	-	-
All races, male												
2004.....	2	-	1	-	2	4	1	-	-	1	-	-
2003.....	2	1	1	1	-	1	-	3	-	1	-	-
2002.....	-	-	-	1	-	1	-	1	-	-	-	-
2001.....	1	1	2	-	2	1	-	1	-	-	-	-
2000.....	1	-	3	2	-	2	1	-	-	-	-	-
1999.....	-	2	1	-	2	1	2	-	-	1	-	-
All races, female												
2004.....	3	-	1	2	1	-	3	1	-	1	-	-
2003.....	-	1	3	-	-	-	1	1	-	-	-	-
2002.....	-	-	2	2	-	2	2	1	-	-	-	-
2001.....	-	1	2	-	2	-	3	2	3	-	-	-
2000.....	1	1	-	5	-	1	1	2	-	-	-	-
1999.....	1	1	-	-	2	2	-	-	-	-	-	-
White, both sexes												
2004.....	3	-	2	1	3	3	4	1	-	2	-	-
2003.....	2	2	3	1	-	1	1	4	-	-	-	-
2002.....	-	-	2	3	-	3	2	2	-	-	-	-
2001.....	1	2	3	-	4	1	3	3	3	-	-	-
2000.....	2	1	3	6	-	3	1	2	-	-	-	-
1999.....	1	3	1	-	3	3	2	-	-	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Congenital malformations of respiratory system (Q30-Q34)												
White, male												
2004.....	1	-	1	-	2	3	1	-	-	1	-	-
2003.....	2	1	1	1	-	1	-	3	-	-	-	-
2002.....	-	-	-	1	-	1	-	1	-	-	-	-
2001.....	1	1	1	-	2	1	-	1	-	-	-	-
2000.....	1	-	3	2	-	2	-	-	-	-	-	-
1999.....	-	2	1	-	1	1	2	-	-	1	-	-
White, female												
2004.....	2	-	1	1	1	-	3	1	-	1	-	-
2003.....	-	1	2	-	-	-	1	1	-	-	-	-
2002.....	-	-	2	2	-	2	2	1	-	-	-	-
2001.....	-	1	2	-	2	-	3	2	3	-	-	-
2000.....	1	1	-	4	-	1	1	2	-	-	-	-
1999.....	1	1	-	-	2	2	-	-	-	-	-	-
Black, both sexes												
2004.....	2	-	-	1	-	-	-	-	-	-	-	-
2003.....	-	-	1	-	-	-	-	-	-	1	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	1	-	-	1	-	-	-	-	-
1999.....	-	-	-	-	1	-	-	-	-	-	-	-
Black, male												
2004.....	1	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	1	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	1	-	-	-	-	-
1999.....	-	-	-	-	1	-	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other congenital malformations and deformations (Q10-Q18,Q35-Q89)												
All races, both sexes												
2004.....	63	58	72	78	76	86	95	75	39	11	3	-
2003.....	62	78	55	63	67	99	93	70	47	14	3	-
2002.....	75	68	70	58	83	79	91	83	26	14	1	1
2001.....	51	54	68	64	77	97	91	71	26	11	2	-
2000.....	55	67	65	79	75	93	80	61	38	16	4	-
1999.....	52	58	59	69	81	93	88	62	34	13	-	2
All races, male												
2004.....	34	31	34	41	34	38	33	34	12	2	-	-
2003.....	40	41	24	32	36	40	32	27	17	4	-	-
2002.....	44	32	37	30	45	42	36	36	11	3	-	1
2001.....	30	28	31	34	38	43	34	36	7	4	1	-
2000.....	36	23	31	38	32	35	33	30	20	5	1	-
1999.....	32	34	27	30	44	41	37	23	12	1	-	1
All races, female												
2004.....	29	27	38	37	42	48	62	41	27	9	3	-
2003.....	22	37	31	31	31	59	61	43	30	10	3	-
2002.....	31	36	33	28	38	37	55	47	15	11	1	-
2001.....	21	26	37	30	39	54	57	35	19	7	1	-
2000.....	19	44	34	41	43	58	47	31	18	11	3	-
1999.....	20	24	32	39	37	52	51	39	22	12	-	1
White, both sexes												
2004.....	54	46	64	72	71	75	83	69	36	11	2	-
2003.....	53	65	52	53	58	91	84	63	42	13	3	-
2002.....	66	60	59	51	74	76	81	75	26	12	1	1
2001.....	44	46	61	57	72	87	78	70	24	9	-	-
2000.....	48	57	60	69	67	84	75	54	37	14	3	-
1999.....	49	51	53	66	74	81	81	58	32	13	-	2

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other congenital malformations and deformations (Q10-Q18,Q35-Q89)												
White, male												
2004.....	28	23	31	37	33	32	28	31	12	2	-	-
2003.....	35	34	21	28	32	37	26	27	15	3	-	-
2002.....	38	28	32	27	40	40	30	32	11	3	-	1
2001.....	25	23	28	32	36	41	25	35	6	4	-	-
2000.....	31	20	30	32	28	32	31	27	19	5	-	-
1999.....	29	30	23	29	41	34	36	22	11	1	-	1
White, female												
2004.....	26	23	33	35	38	43	55	38	24	9	2	-
2003.....	18	31	31	25	26	54	58	36	27	10	3	-
2002.....	28	32	27	24	34	36	51	43	15	9	1	-
2001.....	19	23	33	25	36	46	53	35	18	5	-	-
2000.....	17	37	30	37	39	52	44	27	18	9	3	-
1999.....	20	21	30	37	33	47	45	36	21	12	-	1
Black, both sexes												
2004.....	7	8	8	6	3	8	10	6	2	-	1	-
2003.....	6	8	3	7	7	6	7	5	4	1	-	-
2002.....	7	7	9	7	8	3	7	7	-	2	-	-
2001.....	7	6	6	6	5	9	8	1	1	2	2	-
2000.....	7	6	4	10	7	9	5	5	1	2	1	-
1999.....	3	6	5	3	7	10	6	3	2	-	-	-
Black, male												
2004.....	5	7	3	4	1	5	4	3	-	-	-	-
2003.....	3	5	3	4	3	2	5	-	1	1	-	-
2002.....	5	4	5	3	4	2	4	4	-	-	-	-
2001.....	5	3	3	2	2	2	6	1	-	-	1	-
2000.....	5	-	1	6	3	3	2	2	1	-	1	-
1999.....	3	3	3	1	3	5	1	1	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other congenital malformations and deformations (Q10-Q18,Q35-Q89)												
Asian or Pacific Islander 2/, both sexes												
2004.....	2	3	-	-	2	3	2	-	1	-	-	-
2003.....	3	3	-	1	2	2	2	-	1	-	-	-
2002.....	2	1	1	-	1	-	1	1	-	-	-	-
2001.....	-	2	1	1	-	1	4	-	1	-	-	-
2000.....	-	2	1	-	1	-	-	2	-	-	-	-
1999.....	-	-	1	-	-	2	1	1	-	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	1	1	-	-	-	1	1	-	-	-	-	-
2003.....	2	1	-	-	1	1	1	-	1	-	-	-
2002.....	1	-	-	-	1	-	1	-	-	-	-	-
2001.....	-	2	-	-	-	-	3	-	1	-	-	-
2000.....	-	2	-	-	1	-	-	1	-	-	-	-
1999.....	-	-	1	-	-	2	-	-	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	1	2	-	-	2	2	1	-	1	-	-	-
2003.....	1	2	-	1	1	1	1	-	-	-	-	-
2002.....	1	1	1	-	-	-	-	1	-	-	-	-
2001.....	-	-	1	1	-	1	1	-	-	-	-	-
2000.....	-	-	1	-	-	-	-	1	-	-	-	-
1999.....	-	-	-	-	-	-	1	1	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Down's syndrome (Q90)												
All races, both sexes												
2004.....	130	160	102	56	25	7	2	5	2	-	-	-
2003.....	123	168	112	42	17	5	1	1	3	-	1	-
2002.....	123	136	115	55	11	2	2	2	1	-	-	-
2001.....	119	128	108	42	14	6	3	1	-	-	-	-
2000.....	119	103	89	57	14	6	2	1	-	-	-	-
1999.....	86	102	77	33	11	4	-	-	-	-	1	-
All races, male												
2004.....	61	82	51	28	15	4	2	1	1	-	-	-
2003.....	55	82	63	20	9	-	-	-	-	-	-	-
2002.....	64	66	60	26	2	1	2	-	1	-	-	-
2001.....	56	52	52	23	7	4	1	-	-	-	-	-
2000.....	65	52	38	22	5	1	-	-	-	-	-	-
1999.....	35	50	42	13	4	-	-	-	-	-	-	-
All races, female												
2004.....	69	78	51	28	10	3	-	4	1	-	-	-
2003.....	68	86	49	22	8	5	1	1	3	-	1	-
2002.....	59	70	55	29	9	1	-	2	-	-	-	-
2001.....	63	76	56	19	7	2	2	1	-	-	-	-
2000.....	54	51	51	35	9	5	2	1	-	-	-	-
1999.....	51	52	35	20	7	4	-	-	-	-	1	-
White, both sexes												
2004.....	120	154	98	53	25	7	2	5	2	-	-	-
2003.....	114	155	104	39	15	5	1	1	3	-	1	-
2002.....	111	122	109	52	11	2	2	2	1	-	-	-
2001.....	111	121	104	40	12	6	3	1	-	-	-	-
2000.....	114	94	82	54	14	6	2	1	-	-	-	-
1999.....	78	98	74	29	9	4	-	-	-	-	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Down's syndrome (Q90)												
White, male												
2004.....	57	78	49	26	15	4	2	1	1	-	-	-
2003.....	53	75	58	17	8	-	-	-	-	-	-	-
2002.....	57	57	57	25	2	1	2	-	1	-	-	-
2001.....	54	49	50	22	6	4	1	-	-	-	-	-
2000.....	63	49	35	19	5	1	-	-	-	-	-	-
1999.....	33	49	41	13	2	-	-	-	-	-	-	-
White, female												
2004.....	63	76	49	27	10	3	-	4	1	-	-	-
2003.....	61	80	46	22	7	5	1	1	3	-	1	-
2002.....	54	65	52	27	9	1	-	2	-	-	-	-
2001.....	57	72	54	18	6	2	2	1	-	-	-	-
2000.....	51	45	47	35	9	5	2	1	-	-	-	-
1999.....	45	49	33	16	7	4	-	-	-	-	1	-
Black, both sexes												
2004.....	8	6	4	2	-	-	-	-	-	-	-	-
2003.....	9	13	7	3	1	-	-	-	-	-	-	-
2002.....	10	14	6	2	-	-	-	-	-	-	-	-
2001.....	5	7	4	2	2	-	-	-	-	-	-	-
2000.....	4	9	6	2	-	-	-	-	-	-	-	-
1999.....	8	2	2	4	1	-	-	-	-	-	-	-
Black, male												
2004.....	3	4	2	1	-	-	-	-	-	-	-	-
2003.....	2	7	4	3	1	-	-	-	-	-	-	-
2002.....	6	9	3	1	-	-	-	-	-	-	-	-
2001.....	1	3	2	1	1	-	-	-	-	-	-	-
2000.....	1	3	2	2	-	-	-	-	-	-	-	-
1999.....	2	1	1	-	1	-	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other chromosomal abnormalities, not elsewhere classified (Q91-Q99)												
All races, both sexes												
2004.....	2	2	-	1	2	3	-	1	-	-	-	-
2003.....	2	2	5	1	2	3	-	-	-	-	-	-
2002.....	1	1	-	1	-	3	1	-	-	-	-	-
2001.....	-	3	3	2	1	-	2	-	2	-	1	-
2000.....	2	-	1	1	-	-	1	-	1	-	-	1
1999.....	5	3	2	1	-	1	1	-	-	-	-	-
All races, male												
2004.....	1	1	-	1	1	3	-	-	-	-	-	-
2003.....	1	-	2	1	1	1	-	-	-	-	-	-
2002.....	1	-	-	-	-	2	1	-	-	-	-	-
2001.....	-	-	-	1	-	-	2	-	-	-	1	-
2000.....	1	-	-	1	-	-	-	-	1	-	-	-
1999.....	2	-	-	-	-	-	1	-	-	-	-	-
All races, female												
2004.....	1	1	-	-	1	-	-	1	-	-	-	-
2003.....	1	2	3	-	1	2	-	-	-	-	-	-
2002.....	-	1	-	1	-	1	-	-	-	-	-	-
2001.....	-	3	3	1	1	-	-	-	2	-	-	-
2000.....	1	-	1	-	-	-	1	-	-	-	-	1
1999.....	3	3	2	1	-	1	-	-	-	-	-	-
White, both sexes												
2004.....	2	2	-	1	2	3	-	1	-	-	-	-
2003.....	2	2	5	1	2	3	-	-	-	-	-	-
2002.....	1	-	-	1	-	3	1	-	-	-	-	-
2001.....	-	3	3	2	1	-	1	-	2	-	1	-
2000.....	1	-	1	1	-	-	1	-	1	-	-	1
1999.....	4	3	2	1	-	1	1	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XVIII. Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)												
All races, both sexes												
2004.....	1,132	1,047	994	1,014	1,309	2,075	3,084	3,844	3,806	2,183	929	119
2003.....	1,296	1,077	990	1,033	1,489	1,989	3,017	3,723	3,702	2,225	915	112
2002.....	1,304	1,042	960	995	1,403	2,103	2,833	3,350	3,406	1,944	854	109
2001.....	1,622	1,192	1,100	1,207	1,567	2,209	2,794	3,239	3,220	1,882	748	115
2000.....	1,415	1,152	1,101	1,353	1,687	2,235	2,794	3,286	3,014	1,798	726	115
1999.....	983	908	952	1,140	1,558	1,945	2,468	2,818	2,799	1,592	589	75
All races, male												
2004.....	733	663	605	585	679	949	1,232	1,272	982	390	124	86
2003.....	829	682	612	618	813	967	1,193	1,148	919	385	118	83
2002.....	828	652	621	563	761	1,010	1,218	1,103	857	379	100	77
2001.....	1,066	758	699	685	812	1,014	1,170	1,017	760	392	105	78
2000.....	966	743	686	832	876	1,100	1,144	1,078	733	331	100	90
1999.....	650	578	603	665	848	949	981	894	699	306	85	56
All races, female												
2004.....	399	384	389	429	630	1,126	1,852	2,572	2,824	1,793	805	33
2003.....	467	395	378	415	676	1,022	1,824	2,575	2,783	1,840	797	29
2002.....	476	390	339	432	642	1,093	1,615	2,247	2,549	1,565	754	32
2001.....	556	434	401	522	755	1,195	1,624	2,222	2,460	1,490	643	37
2000.....	449	409	415	521	811	1,135	1,650	2,208	2,281	1,467	626	25
1999.....	333	330	349	475	710	996	1,487	1,924	2,100	1,286	504	19
White, both sexes												
2004.....	869	815	797	838	1,088	1,814	2,752	3,517	3,479	2,025	808	85
2003.....	985	855	790	846	1,249	1,755	2,733	3,444	3,401	2,063	811	96
2002.....	954	815	783	806	1,212	1,844	2,556	3,077	3,153	1,801	755	83
2001.....	1,286	956	892	975	1,329	1,956	2,509	2,985	2,978	1,732	673	88
2000.....	1,087	907	876	1,072	1,446	1,965	2,509	3,015	2,802	1,680	647	83
1999.....	723	694	753	919	1,335	1,680	2,193	2,582	2,596	1,466	520	61

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XVIII. Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)												
White, male												
2004.....	557	513	484	480	580	835	1,105	1,165	912	353	102	66
2003.....	644	535	486	504	690	861	1,074	1,057	840	354	101	71
2002.....	602	507	509	458	659	871	1,094	1,025	792	347	77	58
2001.....	857	600	558	552	704	914	1,063	933	696	358	95	61
2000.....	744	574	551	658	750	974	1,036	988	687	308	88	66
1999.....	478	437	482	538	731	824	865	807	640	280	79	47
White, female												
2004.....	312	302	313	358	508	979	1,647	2,352	2,567	1,672	706	19
2003.....	341	320	304	342	559	894	1,659	2,387	2,561	1,709	710	25
2002.....	352	308	274	348	553	973	1,462	2,052	2,361	1,454	678	25
2001.....	429	356	334	423	625	1,042	1,446	2,052	2,282	1,374	578	27
2000.....	343	333	325	414	696	991	1,473	2,027	2,115	1,372	559	17
1999.....	245	257	271	381	604	856	1,328	1,775	1,956	1,186	441	14
Black, both sexes												
2004.....	246	209	178	156	198	236	291	271	278	131	103	29
2003.....	290	205	186	171	229	205	240	242	257	135	89	11
2002.....	329	205	157	168	172	230	248	237	214	121	88	21
2001.....	294	203	172	201	207	222	246	214	214	131	65	22
2000.....	298	216	208	256	223	249	260	239	184	95	78	28
1999.....	236	196	182	202	201	239	251	210	183	114	65	11
Black, male												
2004.....	166	138	108	92	87	99	113	85	54	27	17	16
2003.....	176	132	118	105	118	91	95	80	69	23	14	8
2002.....	213	131	101	92	93	124	114	67	53	28	21	14
2001.....	175	139	125	116	94	88	88	71	54	28	10	13
2000.....	203	152	123	159	114	116	99	76	39	14	11	21
1999.....	155	132	113	116	105	114	101	74	54	23	6	7

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XVIII. Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)												
Black, female												
2004.....	80	71	70	64	111	137	178	186	224	104	86	13
2003.....	114	73	68	66	111	114	145	162	188	112	75	3
2002.....	116	74	56	76	79	106	134	170	161	93	67	7
2001.....	119	64	47	85	113	134	158	143	160	103	55	9
2000.....	95	64	85	97	109	133	161	163	145	81	67	7
1999.....	81	64	69	86	96	125	150	136	129	91	59	4
American Indian 1/, both sexes												
2004.....	5	5	8	6	8	7	7	9	13	6	1	3
2003.....	13	8	8	4	1	5	9	8	12	5	5	2
2002.....	11	8	13	8	11	7	6	4	10	7	3	3
2001.....	13	14	12	15	7	6	10	7	9	5	3	4
2000.....	13	13	6	7	4	7	6	11	8	8	-	1
1999.....	11	7	7	8	12	11	16	9	8	6	-	1
American Indian 1/, male												
2004.....	3	2	5	4	4	4	1	3	3	1	-	3
2003.....	3	6	5	2	-	4	5	3	1	1	1	2
2002.....	5	4	8	5	6	4	2	1	1	1	-	3
2001.....	12	8	3	8	4	3	6	1	1	1	-	3
2000.....	9	8	5	4	3	4	2	4	2	3	-	1
1999.....	10	4	3	5	5	5	10	2	3	2	-	1
American Indian 1/, female												
2004.....	2	3	3	2	4	3	6	6	10	5	1	-
2003.....	10	2	3	2	1	1	4	5	11	4	4	-
2002.....	6	4	5	3	5	3	4	3	9	6	3	-
2001.....	1	6	9	7	3	3	4	6	8	4	3	1
2000.....	4	5	1	3	1	3	4	7	6	5	-	-
1999.....	1	3	4	3	7	6	6	7	5	4	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XVIII. Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)												
Asian or Pacific Islander 2/, both sexes												
2004.....	12	18	11	14	15	18	34	47	36	21	17	2
2003.....	8	9	6	12	10	24	35	29	32	22	10	3
2002.....	10	14	7	13	8	22	23	32	29	15	8	2
2001.....	29	19	24	16	24	25	29	33	19	14	7	1
2000.....	17	16	11	18	14	14	19	21	20	15	1	3
1999.....	13	11	10	11	10	15	8	17	12	6	4	2
Asian or Pacific Islander 2/, male												
2004.....	7	10	8	9	8	11	13	19	13	9	5	1
2003.....	6	9	3	7	5	11	19	8	9	7	2	2
2002.....	8	10	3	8	3	11	8	10	11	3	2	2
2001.....	22	11	13	9	10	9	13	12	9	5	-	1
2000.....	10	9	7	11	9	6	7	10	5	6	1	2
1999.....	7	5	5	6	7	6	5	11	2	1	-	1
Asian or Pacific Islander 2/, female												
2004.....	5	8	3	5	7	7	21	28	23	12	12	1
2003.....	2	-	3	5	5	13	16	21	23	15	8	1
2002.....	2	4	4	5	5	11	15	22	18	12	6	-
2001.....	7	8	11	7	14	16	16	21	10	9	7	-
2000.....	7	7	4	7	5	8	12	11	15	9	-	1
1999.....	6	6	5	5	3	9	3	6	10	5	4	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Senility (R54)												
All races, both sexes												
2004.....	6	4	21	42	100	305	621	999	1,302	918	488	-
2003.....	3	4	20	40	120	264	608	960	1,247	931	478	-
2002.....	3	5	12	37	104	275	521	871	1,162	791	485	-
2001.....	3	6	13	36	118	245	554	834	1,152	846	381	-
2000.....	-	2	9	35	112	266	510	777	975	720	379	-
1999.....	3	3	13	32	98	215	438	643	847	569	288	-
All races, male												
2004.....	5	3	12	27	48	133	246	306	321	163	72	-
2003.....	2	2	7	23	60	114	225	282	314	146	62	-
2002.....	1	1	7	21	45	111	207	280	297	156	52	-
2001.....	3	4	6	21	57	99	220	251	260	179	53	-
2000.....	-	-	4	18	52	117	193	226	219	141	48	-
1999.....	3	2	8	18	60	92	160	192	208	102	45	-
All races, female												
2004.....	1	1	9	15	52	172	375	693	981	755	416	-
2003.....	1	2	13	17	60	150	383	678	933	785	416	-
2002.....	2	4	5	16	59	164	314	591	865	635	433	-
2001.....	-	2	7	15	61	146	334	583	892	667	328	-
2000.....	-	2	5	17	60	149	317	551	756	579	331	-
1999.....	-	1	5	14	38	123	278	451	639	467	243	-
White, both sexes												
2004.....	5	3	15	34	90	276	561	946	1,211	865	438	-
2003.....	2	3	17	32	112	242	575	903	1,158	880	435	-
2002.....	3	5	12	32	91	250	485	807	1,097	743	433	-
2001.....	-	4	9	23	108	221	512	793	1,072	783	346	-
2000.....	-	2	7	27	95	236	456	711	919	678	339	-
1999.....	2	3	9	30	87	187	391	612	805	533	263	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Senility (R54)												
White, male												
2004.....	4	2	8	23	43	122	219	294	309	151	61	-
2003.....	2	2	6	16	56	105	209	262	297	135	59	-
2002.....	1	1	7	19	38	100	198	262	281	146	40	-
2001.....	-	3	4	16	54	92	205	235	238	167	46	-
2000.....	-	-	3	14	48	103	176	206	213	134	44	-
1999.....	2	2	5	17	53	80	141	179	195	95	43	-
White, female												
2004.....	1	1	7	11	47	154	342	652	902	714	377	-
2003.....	-	1	11	16	56	137	366	641	861	745	376	-
2002.....	2	4	5	13	53	150	287	545	816	597	393	-
2001.....	-	1	5	7	54	129	307	558	834	616	300	-
2000.....	-	2	4	13	47	133	280	505	706	544	295	-
1999.....	-	1	4	13	34	107	250	433	610	438	220	-
Black, both sexes												
2004.....	1	1	6	7	10	27	52	45	74	38	42	-
2003.....	1	1	3	8	8	20	25	45	72	38	36	-
2002.....	-	-	-	5	13	21	29	52	50	40	45	-
2001.....	3	2	3	12	7	23	40	35	67	54	29	-
2000.....	-	-	2	8	17	28	50	59	47	32	40	-
1999.....	1	-	4	1	11	25	46	27	38	35	25	-
Black, male												
2004.....	1	1	4	4	5	11	25	9	9	7	8	-
2003.....	-	-	1	7	4	8	11	16	13	5	2	-
2002.....	-	-	-	2	7	10	7	14	13	8	11	-
2001.....	3	1	2	4	1	7	14	15	18	11	7	-
2000.....	-	-	1	4	4	13	16	18	5	3	4	-
1999.....	1	-	3	-	7	11	19	11	11	7	2	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Senility (R54)												
Black, female												
2004.....	-	-	2	3	5	16	27	36	65	31	34	-
2003.....	1	1	2	1	4	12	14	29	59	33	34	-
2002.....	-	-	-	3	6	11	22	38	37	32	34	-
2001.....	-	1	1	8	6	16	26	20	49	43	22	-
2000.....	-	-	1	4	13	15	34	41	42	29	36	-
1999.....	-	-	1	1	4	14	27	16	27	28	23	-
American Indian 1/, both sexes												
2004.....	-	-	-	-	-	-	1	-	6	4	-	-
2003.....	-	-	-	-	-	-	2	4	9	3	3	-
2002.....	-	-	-	-	-	-	1	2	3	2	1	-
2001.....	-	-	1	1	1	-	1	2	4	4	3	-
2000.....	-	-	-	-	-	1	-	6	4	3	-	-
1999.....	-	-	-	1	-	3	1	2	3	1	-	-
American Indian 1/, male												
2004.....	-	-	-	-	-	-	-	-	1	1	-	-
2003.....	-	-	-	-	-	-	2	2	1	1	1	-
2002.....	-	-	-	-	-	-	1	-	-	-	-	-
2001.....	-	-	-	1	1	-	1	-	-	1	-	-
2000.....	-	-	-	-	-	-	-	2	-	2	-	-
1999.....	-	-	-	1	-	1	-	1	2	-	-	-
American Indian 1/, female												
2004.....	-	-	-	-	-	-	1	-	5	3	-	-
2003.....	-	-	-	-	-	-	-	2	8	2	2	-
2002.....	-	-	-	-	-	-	-	2	3	2	1	-
2001.....	-	-	1	-	-	-	-	2	4	3	3	-
2000.....	-	-	-	-	-	1	-	4	4	1	-	-
1999.....	-	-	-	-	-	2	1	1	1	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Senility (R54)												
Asian or Pacific Islander 2/, both sexes												
2004.....	-	-	-	1	-	2	7	8	11	11	8	-
2003.....	-	-	-	-	-	2	6	8	8	10	4	-
2002.....	-	-	-	-	-	4	6	10	12	6	6	-
2001.....	-	-	-	-	2	1	1	4	9	5	3	-
2000.....	-	-	-	-	-	1	4	1	5	7	-	-
1999.....	-	-	-	-	-	-	-	2	1	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	-	-	-	-	-	-	2	3	2	4	3	-
2003.....	-	-	-	-	-	1	3	2	3	5	-	-
2002.....	-	-	-	-	-	1	1	4	3	2	1	-
2001.....	-	-	-	-	1	-	-	1	4	-	-	-
2000.....	-	-	-	-	-	1	1	-	1	2	-	-
1999.....	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	-	-	-	1	-	2	5	5	9	7	5	-
2003.....	-	-	-	-	-	1	3	6	5	5	4	-
2002.....	-	-	-	-	-	3	5	6	9	4	5	-
2001.....	-	-	-	-	1	1	1	3	5	5	3	-
2000.....	-	-	-	-	-	-	3	1	4	5	-	-
1999.....	-	-	-	-	-	-	-	1	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Ill-defined and unknown causes of mortality (R95-R99)												
All races, both sexes												
2004.....	872	769	648	603	644	851	1,009	1,022	915	445	166	118
2003.....	1,048	833	674	651	779	877	1,067	1,156	982	496	172	110
2002.....	1,025	821	679	633	758	995	1,103	1,113	928	485	152	108
2001.....	1,392	972	845	843	911	1,103	1,177	1,179	930	458	168	115
2000.....	1,193	933	826	958	1,056	1,200	1,329	1,321	1,021	577	174	114
1999.....	781	680	673	755	931	994	1,119	1,160	942	506	163	74
All races, male												
2004.....	570	504	417	356	373	439	431	409	258	81	16	85
2003.....	677	541	438	427	468	466	481	393	268	93	21	82
2002.....	677	531	449	375	440	504	512	413	243	89	19	77
2001.....	922	622	542	496	513	554	533	411	213	105	26	78
2000.....	823	607	535	607	558	616	572	466	286	110	23	89
1999.....	521	438	449	462	537	511	475	392	244	106	26	55
All races, female												
2004.....	302	265	231	247	271	412	578	613	657	364	150	33
2003.....	371	292	236	224	311	411	586	763	714	403	151	28
2002.....	348	290	230	258	318	491	591	700	685	396	133	31
2001.....	470	350	303	347	398	549	644	768	717	353	142	37
2000.....	370	326	291	351	498	584	757	855	735	467	151	25
1999.....	260	242	224	293	394	483	644	768	698	400	137	19
White, both sexes												
2004.....	687	615	539	505	534	761	904	927	837	415	145	84
2003.....	822	672	545	553	660	786	968	1,070	912	461	151	95
2002.....	760	655	562	516	663	879	998	1,026	850	447	138	82
2001.....	1,132	800	702	708	785	980	1,051	1,091	863	420	145	88
2000.....	933	754	675	769	929	1,064	1,212	1,223	943	536	156	83
1999.....	583	522	538	614	811	872	994	1,049	872	458	143	60

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Ill-defined and unknown causes of mortality (R95-R99)												
White, male												
2004.....	443	403	343	300	322	397	387	372	242	74	12	65
2003.....	544	427	349	355	395	417	438	368	244	86	17	70
2002.....	500	422	372	305	388	441	454	389	225	78	15	58
2001.....	761	506	450	416	449	497	476	378	198	96	25	61
2000.....	649	491	439	483	486	547	524	425	265	102	19	66
1999.....	386	333	361	375	467	450	416	344	220	95	22	46
White, female												
2004.....	244	212	196	205	212	364	517	555	595	341	133	19
2003.....	278	245	196	198	265	369	530	702	668	375	134	25
2002.....	260	233	190	211	275	438	544	637	625	369	123	24
2001.....	371	294	252	292	336	483	575	713	665	324	120	27
2000.....	284	263	236	286	443	517	688	798	678	434	137	17
1999.....	197	189	177	239	344	422	578	705	652	363	121	14
Black, both sexes												
2004.....	173	138	97	86	97	82	86	82	68	27	18	29
2003.....	207	146	120	88	113	79	84	78	61	32	16	11
2002.....	249	148	105	103	82	104	93	77	66	35	11	21
2001.....	225	147	114	110	104	107	104	68	58	33	22	22
2000.....	236	156	138	168	114	121	108	84	74	36	17	27
1999.....	179	144	123	124	102	108	112	100	63	42	19	11
Black, male												
2004.....	121	91	65	47	45	36	37	32	11	6	4	16
2003.....	125	100	84	64	70	43	35	24	22	7	4	8
2002.....	167	98	71	62	48	56	53	21	16	10	3	14
2001.....	133	101	79	67	55	49	44	24	12	7	1	13
2000.....	161	103	88	110	64	62	46	33	19	6	3	20
1999.....	122	97	82	77	60	54	48	41	22	9	4	7

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Ill-defined and unknown causes of mortality (R95-R99)												
Black, female												
2004.....	52	47	32	39	52	46	49	50	57	21	14	13
2003.....	82	46	36	24	43	36	49	54	39	25	12	3
2002.....	82	50	34	41	34	48	40	56	50	25	8	7
2001.....	92	46	35	43	49	58	60	44	46	26	21	9
2000.....	75	53	50	58	50	59	62	51	55	30	14	7
1999.....	57	47	41	47	42	54	64	59	41	33	15	4
American Indian 1/, both sexes												
2004.....	5	3	7	4	2	4	4	5	3	-	1	3
2003.....	11	7	4	3	1	5	4	2	1	1	1	2
2002.....	10	6	9	4	9	3	1	2	6	1	1	3
2001.....	9	10	8	12	3	4	4	3	1	1	-	4
2000.....	11	10	5	6	2	5	1	4	-	3	-	1
1999.....	9	6	6	7	10	6	10	2	3	3	-	1
American Indian 1/, male												
2004.....	3	1	4	3	-	2	1	1	-	-	-	3
2003.....	2	6	3	2	-	4	2	1	-	-	-	2
2002.....	5	3	5	2	4	3	-	1	-	-	-	3
2001.....	9	7	2	6	1	3	4	1	-	-	-	3
2000.....	7	6	4	4	2	3	-	1	-	1	-	1
1999.....	8	4	3	4	4	3	9	-	1	1	-	1
American Indian 1/, female												
2004.....	2	2	3	1	2	2	3	4	3	-	1	-
2003.....	9	1	1	1	1	1	2	1	1	1	1	-
2002.....	5	3	4	2	5	-	1	1	6	1	1	-
2001.....	-	3	6	6	2	1	-	2	1	1	-	1
2000.....	4	4	1	2	-	2	1	3	-	2	-	-
1999.....	1	2	3	3	6	3	1	2	2	2	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Ill-defined and unknown causes of mortality (R95-R99)												
Asian or Pacific Islander 2/, both sexes												
2004.....	7	13	5	8	11	4	15	8	7	3	2	2
2003.....	8	8	5	7	5	7	11	6	8	2	4	2
2002.....	6	12	3	10	4	9	11	8	6	2	2	2
2001.....	26	15	21	13	19	12	18	17	8	4	1	1
2000.....	13	13	8	15	11	10	8	10	4	2	1	3
1999.....	10	8	6	10	8	8	3	9	4	3	1	2
Asian or Pacific Islander 2/, male												
2004.....	3	9	5	6	6	4	6	4	5	1	-	1
2003.....	6	8	2	6	3	2	6	-	2	-	-	2
2002.....	5	8	1	6	-	4	5	2	2	1	1	2
2001.....	19	8	11	7	8	5	9	8	3	2	-	1
2000.....	6	7	4	10	6	4	2	7	2	1	1	2
1999.....	5	4	3	6	6	4	2	7	1	1	-	1
Asian or Pacific Islander 2/, female												
2004.....	4	4	-	2	5	-	9	4	2	2	2	1
2003.....	2	-	3	1	2	5	5	6	6	2	4	-
2002.....	1	4	2	4	4	5	6	6	4	1	1	-
2001.....	7	7	10	6	11	7	9	9	5	2	1	-
2000.....	7	6	4	5	5	6	6	3	2	1	-	1
1999.....	5	4	3	4	2	4	1	2	3	2	1	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other ill-defined and unknown causes of mortality (R96-R99)												
All races, both sexes												
2004.....	872	769	648	603	644	851	1,009	1,022	915	445	166	118
2003.....	1,048	833	674	651	779	877	1,067	1,156	982	496	172	110
2002.....	1,025	821	679	633	758	995	1,103	1,113	928	485	152	108
2001.....	1,392	972	845	843	911	1,103	1,177	1,179	930	458	168	115
2000.....	1,193	933	826	958	1,056	1,200	1,329	1,321	1,021	577	174	114
1999.....	781	680	673	755	931	994	1,119	1,160	942	506	163	74
All races, male												
2004.....	570	504	417	356	373	439	431	409	258	81	16	85
2003.....	677	541	438	427	468	466	481	393	268	93	21	82
2002.....	677	531	449	375	440	504	512	413	243	89	19	77
2001.....	922	622	542	496	513	554	533	411	213	105	26	78
2000.....	823	607	535	607	558	616	572	466	286	110	23	89
1999.....	521	438	449	462	537	511	475	392	244	106	26	55
All races, female												
2004.....	302	265	231	247	271	412	578	613	657	364	150	33
2003.....	371	292	236	224	311	411	586	763	714	403	151	28
2002.....	348	290	230	258	318	491	591	700	685	396	133	31
2001.....	470	350	303	347	398	549	644	768	717	353	142	37
2000.....	370	326	291	351	498	584	757	855	735	467	151	25
1999.....	260	242	224	293	394	483	644	768	698	400	137	19
White, both sexes												
2004.....	687	615	539	505	534	761	904	927	837	415	145	84
2003.....	822	672	545	553	660	786	968	1,070	912	461	151	95
2002.....	760	655	562	516	663	879	998	1,026	850	447	138	82
2001.....	1,132	800	702	708	785	980	1,051	1,091	863	420	145	88
2000.....	933	754	675	769	929	1,064	1,212	1,223	943	536	156	83
1999.....	583	522	538	614	811	872	994	1,049	872	458	143	60

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other ill-defined and unknown causes of mortality (R96-R99)												
White, male												
2004.....	443	403	343	300	322	397	387	372	242	74	12	65
2003.....	544	427	349	355	395	417	438	368	244	86	17	70
2002.....	500	422	372	305	388	441	454	389	225	78	15	58
2001.....	761	506	450	416	449	497	476	378	198	96	25	61
2000.....	649	491	439	483	486	547	524	425	265	102	19	66
1999.....	386	333	361	375	467	450	416	344	220	95	22	46
White, female												
2004.....	244	212	196	205	212	364	517	555	595	341	133	19
2003.....	278	245	196	198	265	369	530	702	668	375	134	25
2002.....	260	233	190	211	275	438	544	637	625	369	123	24
2001.....	371	294	252	292	336	483	575	713	665	324	120	27
2000.....	284	263	236	286	443	517	688	798	678	434	137	17
1999.....	197	189	177	239	344	422	578	705	652	363	121	14
Black, both sexes												
2004.....	173	138	97	86	97	82	86	82	68	27	18	29
2003.....	207	146	120	88	113	79	84	78	61	32	16	11
2002.....	249	148	105	103	82	104	93	77	66	35	11	21
2001.....	225	147	114	110	104	107	104	68	58	33	22	22
2000.....	236	156	138	168	114	121	108	84	74	36	17	27
1999.....	179	144	123	124	102	108	112	100	63	42	19	11
Black, male												
2004.....	121	91	65	47	45	36	37	32	11	6	4	16
2003.....	125	100	84	64	70	43	35	24	22	7	4	8
2002.....	167	98	71	62	48	56	53	21	16	10	3	14
2001.....	133	101	79	67	55	49	44	24	12	7	1	13
2000.....	161	103	88	110	64	62	46	33	19	6	3	20
1999.....	122	97	82	77	60	54	48	41	22	9	4	7

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other ill-defined and unknown causes of mortality (R96-R99)												
Black, female												
2004.....	52	47	32	39	52	46	49	50	57	21	14	13
2003.....	82	46	36	24	43	36	49	54	39	25	12	3
2002.....	82	50	34	41	34	48	40	56	50	25	8	7
2001.....	92	46	35	43	49	58	60	44	46	26	21	9
2000.....	75	53	50	58	50	59	62	51	55	30	14	7
1999.....	57	47	41	47	42	54	64	59	41	33	15	4
American Indian 1/, both sexes												
2004.....	5	3	7	4	2	4	4	5	3	-	1	3
2003.....	11	7	4	3	1	5	4	2	1	1	1	2
2002.....	10	6	9	4	9	3	1	2	6	1	1	3
2001.....	9	10	8	12	3	4	4	3	1	1	-	4
2000.....	11	10	5	6	2	5	1	4	-	3	-	1
1999.....	9	6	6	7	10	6	10	2	3	3	-	1
American Indian 1/, male												
2004.....	3	1	4	3	-	2	1	1	-	-	-	3
2003.....	2	6	3	2	-	4	2	1	-	-	-	2
2002.....	5	3	5	2	4	3	-	1	-	-	-	3
2001.....	9	7	2	6	1	3	4	1	-	-	-	3
2000.....	7	6	4	4	2	3	-	1	-	1	-	1
1999.....	8	4	3	4	4	3	9	-	1	1	-	1
American Indian 1/, female												
2004.....	2	2	3	1	2	2	3	4	3	-	1	-
2003.....	9	1	1	1	1	1	2	1	1	1	1	-
2002.....	5	3	4	2	5	-	1	1	6	1	1	-
2001.....	-	3	6	6	2	1	-	2	1	1	-	1
2000.....	4	4	1	2	-	2	1	3	-	2	-	-
1999.....	1	2	3	3	6	3	1	2	2	2	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other ill-defined and unknown causes of mortality (R96-R99)												
Asian or Pacific Islander 2/, both sexes												
2004.....	7	13	5	8	11	4	15	8	7	3	2	2
2003.....	8	8	5	7	5	7	11	6	8	2	4	2
2002.....	6	12	3	10	4	9	11	8	6	2	2	2
2001.....	26	15	21	13	19	12	18	17	8	4	1	1
2000.....	13	13	8	15	11	10	8	10	4	2	1	3
1999.....	10	8	6	10	8	8	3	9	4	3	1	2
Asian or Pacific Islander 2/, male												
2004.....	3	9	5	6	6	4	6	4	5	1	-	1
2003.....	6	8	2	6	3	2	6	-	2	-	-	2
2002.....	5	8	1	6	-	4	5	2	2	1	1	2
2001.....	19	8	11	7	8	5	9	8	3	2	-	1
2000.....	6	7	4	10	6	4	2	7	2	1	1	2
1999.....	5	4	3	6	6	4	2	7	1	1	-	1
Asian or Pacific Islander 2/, female												
2004.....	4	4	-	2	5	-	9	4	2	2	2	1
2003.....	2	-	3	1	2	5	5	6	6	2	4	-
2002.....	1	4	2	4	4	5	6	6	4	1	1	-
2001.....	7	7	10	6	11	7	9	9	5	2	1	-
2000.....	7	6	4	5	5	6	6	3	2	1	-	1
1999.....	5	4	3	4	2	4	1	2	3	2	1	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R53,R55-R94)												
All races, both sexes												
2004.....	254	274	325	369	565	919	1,454	1,823	1,589	820	275	1
2003.....	245	240	296	342	590	848	1,342	1,607	1,473	798	265	2
2002.....	276	216	269	325	541	833	1,209	1,366	1,316	668	217	1
2001.....	227	214	242	328	538	861	1,063	1,226	1,138	578	199	-
2000.....	222	217	266	360	519	769	955	1,188	1,018	501	173	1
1999.....	199	225	266	353	529	736	911	1,015	1,010	517	138	1
All races, male												
2004.....	158	156	176	202	258	377	555	557	403	146	36	1
2003.....	150	139	167	168	285	387	487	473	337	146	35	1
2002.....	150	120	165	167	276	395	499	410	317	134	29	-
2001.....	141	132	151	168	242	361	417	355	287	108	26	-
2000.....	143	136	147	207	266	367	379	386	228	80	29	1
1999.....	126	138	146	185	251	346	346	310	247	98	14	1
All races, female												
2004.....	96	118	149	167	307	542	899	1,266	1,186	674	239	-
2003.....	95	101	129	174	305	461	855	1,134	1,136	652	230	1
2002.....	126	96	104	158	265	438	710	956	999	534	188	1
2001.....	86	82	91	160	296	500	646	871	851	470	173	-
2000.....	79	81	119	153	253	402	576	802	790	421	144	-
1999.....	73	87	120	168	278	390	565	705	763	419	124	-
White, both sexes												
2004.....	177	197	243	299	464	777	1,287	1,644	1,431	745	225	1
2003.....	161	180	228	261	477	727	1,190	1,471	1,331	722	225	1
2002.....	191	155	209	258	458	715	1,073	1,244	1,206	611	184	1
2001.....	154	152	181	244	436	755	946	1,101	1,043	529	182	-
2000.....	154	151	194	276	422	665	841	1,081	940	466	152	-
1999.....	138	169	206	275	437	621	808	921	919	475	114	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R53,R55-R94)												
White, male												
2004.....	110	108	133	157	215	316	499	499	361	128	29	1
2003.....	98	106	131	133	239	339	427	427	299	133	25	1
2002.....	101	84	130	134	233	330	442	374	286	123	22	-
2001.....	96	91	104	120	201	325	382	320	260	95	24	-
2000.....	95	83	109	161	216	324	336	357	209	72	25	-
1999.....	90	102	116	146	211	294	308	284	225	90	14	1
White, female												
2004.....	67	89	110	142	249	461	788	1,145	1,070	617	196	-
2003.....	63	74	97	128	238	388	763	1,044	1,032	589	200	-
2002.....	90	71	79	124	225	385	631	870	920	488	162	1
2001.....	58	61	77	124	235	430	564	781	783	434	158	-
2000.....	59	68	85	115	206	341	505	724	731	394	127	-
1999.....	48	67	90	129	226	327	500	637	694	385	100	-
Black, both sexes												
2004.....	72	70	75	63	91	127	153	144	136	66	43	-
2003.....	82	58	63	75	108	106	131	119	124	65	37	-
2002.....	80	57	52	60	77	105	126	108	98	46	32	-
2001.....	66	54	55	79	96	92	102	111	89	44	14	-
2000.....	62	60	68	80	92	100	102	96	63	27	21	1
1999.....	56	52	55	77	88	106	93	83	82	37	21	-
Black, male												
2004.....	44	46	39	41	37	52	51	44	34	14	5	-
2003.....	51	32	33	34	44	40	49	40	34	11	8	-
2002.....	46	33	30	28	38	58	54	32	24	10	7	-
2001.....	39	37	44	45	38	32	30	32	24	10	2	-
2000.....	42	49	34	45	46	41	37	25	15	5	4	1
1999.....	32	35	28	39	38	49	34	22	21	7	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R53,R55-R94)												
Black, female												
2004.....	28	24	36	22	54	75	102	100	102	52	38	-
2003.....	31	26	30	41	64	66	82	79	90	54	29	-
2002.....	34	24	22	32	39	47	72	76	74	36	25	-
2001.....	27	17	11	34	58	60	72	79	65	34	12	-
2000.....	20	11	34	35	46	59	65	71	48	22	17	-
1999.....	24	17	27	38	50	57	59	61	61	30	21	-
American Indian 1/, both sexes												
2004.....	-	2	1	2	6	3	2	4	4	2	-	-
2003.....	2	1	4	1	-	-	3	2	2	1	1	-
2002.....	1	2	4	4	2	4	4	-	1	4	1	-
2001.....	4	4	3	2	3	2	5	2	4	-	-	-
2000.....	2	3	1	1	2	1	5	1	4	2	-	-
1999.....	2	1	1	-	2	2	5	5	2	2	-	-
American Indian 1/, male												
2004.....	-	1	1	1	4	2	-	2	2	-	-	-
2003.....	1	-	2	-	-	-	1	-	-	-	-	-
2002.....	-	1	3	3	2	1	1	-	1	1	-	-
2001.....	3	1	1	1	2	-	1	-	1	-	-	-
2000.....	2	2	1	-	1	1	2	1	2	-	-	-
1999.....	2	-	-	-	1	1	1	1	-	1	-	-
American Indian 1/, female												
2004.....	-	1	-	1	2	1	2	2	2	2	-	-
2003.....	1	1	2	1	-	-	2	2	2	1	1	-
2002.....	1	1	1	1	-	3	3	-	-	3	1	-
2001.....	1	3	2	1	1	2	4	2	3	-	-	-
2000.....	-	1	-	1	1	-	3	-	2	2	-	-
1999.....	-	1	1	-	1	1	4	4	2	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

All other symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R53,R55-R94)

Asian or Pacific Islander 2/, both sexes

2004.....	5	5	6	5	4	12	12	31	18	7	7	-
2003.....	-	1	1	5	5	15	18	15	16	10	2	1
2002.....	4	2	4	3	4	9	6	14	11	7	-	-
2001.....	3	4	3	3	3	12	10	12	2	5	3	-
2000.....	4	3	3	3	3	3	7	10	11	6	-	-
1999.....	3	3	4	1	2	7	5	6	7	3	3	-

Asian or Pacific Islander 2/, male

2004.....	4	1	3	3	2	7	5	12	6	4	2	-
2003.....	-	1	1	1	2	8	10	6	4	2	2	-
2002.....	3	2	2	2	3	6	2	4	6	-	-	-
2001.....	3	3	2	2	1	4	4	3	2	3	-	-
2000.....	4	2	3	1	3	1	4	3	2	3	-	-
1999.....	2	1	2	-	1	2	3	3	1	-	-	-

Asian or Pacific Islander 2/, female

2004.....	1	4	3	2	2	5	7	19	12	3	5	-
2003.....	-	-	-	4	3	7	8	9	12	8	-	1
2002.....	1	-	2	1	1	3	4	10	5	7	-	-
2001.....	-	1	1	1	2	8	6	9	-	2	3	-
2000.....	-	1	-	2	-	2	3	7	9	3	-	-
1999.....	1	2	2	1	1	5	2	3	6	3	3	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XX. External causes of mortality (*U01-*U03,V01-Y89)												
All races, both sexes												
2004.....	12,203	8,773	6,586	5,565	5,997	7,677	9,114	7,742	5,100	1,791	311	144
2003.....	11,407	8,430	6,141	5,395	6,124	7,668	8,769	7,817	4,743	1,721	349	143
2002.....	10,624	7,649	5,845	5,409	6,269	7,768	8,589	7,472	4,674	1,709	304	150
2001 3/.....	10,092	7,117	5,509	5,337	6,138	7,876	8,300	7,214	4,500	1,616	325	186
2000.....	9,038	6,501	5,302	5,002	6,190	7,470	7,684	6,956	4,368	1,520	321	133
1999.....	8,298	6,326	5,135	5,443	6,419	7,694	8,025	7,134	4,332	1,483	270	190
All races, male												
2004.....	8,777	6,221	4,532	3,737	3,836	4,521	4,920	3,603	1,946	514	76	123
2003.....	8,141	6,006	4,313	3,621	3,874	4,550	4,692	3,651	1,836	479	70	121
2002.....	7,636	5,573	4,050	3,658	4,051	4,616	4,544	3,450	1,719	539	57	129
2001 3/.....	7,307	5,078	3,800	3,565	3,897	4,670	4,473	3,364	1,722	422	66	157
2000.....	6,612	4,636	3,642	3,298	3,917	4,432	4,130	3,245	1,622	443	67	116
1999.....	6,059	4,541	3,604	3,644	4,064	4,499	4,341	3,326	1,609	429	63	156
All races, female												
2004.....	3,426	2,552	2,054	1,828	2,161	3,156	4,194	4,139	3,154	1,277	235	21
2003.....	3,266	2,424	1,828	1,774	2,250	3,118	4,077	4,166	2,907	1,242	279	22
2002.....	2,988	2,076	1,795	1,751	2,218	3,152	4,045	4,022	2,955	1,170	247	21
2001 3/.....	2,785	2,039	1,709	1,772	2,241	3,206	3,827	3,850	2,778	1,194	259	29
2000.....	2,426	1,865	1,660	1,704	2,273	3,038	3,554	3,711	2,746	1,077	254	17
1999.....	2,239	1,785	1,531	1,799	2,355	3,195	3,684	3,808	2,723	1,054	207	34
White, both sexes												
2004.....	10,174	7,426	5,662	4,804	5,251	6,910	8,429	7,304	4,785	1,706	285	127
2003.....	9,482	7,186	5,280	4,617	5,391	6,940	8,080	7,353	4,457	1,636	319	125
2002.....	8,840	6,529	5,015	4,612	5,582	7,015	7,968	7,023	4,374	1,599	276	136
2001 3/.....	8,427	6,060	4,657	4,600	5,389	7,140	7,663	6,727	4,207	1,507	294	157
2000.....	7,438	5,488	4,531	4,284	5,429	6,755	7,102	6,522	4,085	1,402	285	118
1999.....	6,870	5,319	4,279	4,686	5,622	6,897	7,384	6,611	4,069	1,383	243	150

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XX. External causes of mortality (*U01-*U03,V01-Y89)												
White, male												
2004.....	7,288	5,218	3,896	3,247	3,386	4,114	4,555	3,400	1,811	490	68	109
2003.....	6,780	5,110	3,726	3,123	3,461	4,147	4,331	3,442	1,719	453	63	105
2002.....	6,337	4,729	3,490	3,113	3,630	4,176	4,222	3,237	1,591	499	48	117
2001 3/.....	6,085	4,298	3,214	3,095	3,446	4,247	4,142	3,140	1,606	386	57	133
2000.....	5,416	3,889	3,101	2,808	3,461	4,017	3,820	3,047	1,507	398	56	104
1999.....	5,011	3,787	3,023	3,148	3,580	4,043	4,018	3,077	1,494	401	55	124
White, female												
2004.....	2,886	2,208	1,766	1,557	1,865	2,796	3,874	3,904	2,974	1,216	217	18
2003.....	2,702	2,076	1,554	1,494	1,930	2,793	3,749	3,911	2,738	1,183	256	20
2002.....	2,503	1,800	1,525	1,499	1,952	2,839	3,746	3,786	2,783	1,100	228	19
2001 3/.....	2,342	1,762	1,443	1,505	1,943	2,893	3,521	3,587	2,601	1,121	237	24
2000.....	2,022	1,599	1,430	1,476	1,968	2,738	3,282	3,475	2,578	1,004	229	14
1999.....	1,859	1,532	1,256	1,538	2,042	2,854	3,366	3,534	2,575	982	188	26
Black, both sexes												
2004.....	1,665	1,104	713	578	547	587	505	290	243	67	24	11
2003.....	1,573	982	676	595	549	523	492	335	209	66	22	16
2002.....	1,467	890	660	581	526	574	453	340	224	88	20	8
2001 3/.....	1,391	861	679	588	593	568	477	363	233	83	27	24
2000.....	1,319	817	616	581	615	547	450	334	221	93	31	12
1999.....	1,164	809	665	618	634	644	512	414	223	92	22	37
Black, male												
2004.....	1,245	826	500	387	348	329	277	118	105	18	8	10
2003.....	1,134	727	475	399	307	300	248	146	77	18	5	14
2002.....	1,084	686	465	404	332	342	239	159	91	32	5	7
2001 3/.....	1,034	641	476	385	358	325	249	149	89	24	8	21
2000.....	998	618	441	411	370	318	237	146	88	34	8	9
1999.....	882	614	460	416	395	369	247	193	90	25	6	30

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XX. External causes of mortality (*U01-*U03,V01-Y89)												
Black, female												
2004.....	420	278	213	191	199	258	228	172	138	49	16	1
2003.....	439	255	201	196	242	223	244	189	132	48	17	2
2002.....	383	204	195	177	194	232	214	181	133	56	15	1
2001 3/.....	357	220	203	203	235	243	228	214	144	59	19	3
2000.....	321	199	175	170	245	229	213	188	133	59	23	3
1999.....	282	195	205	202	239	275	265	221	133	67	16	7
American Indian 1/, both sexes												
2004.....	137	77	65	59	54	33	38	24	12	2	-	6
2003.....	142	89	67	58	41	42	29	28	16	3	2	1
2002.....	135	88	55	64	34	38	31	9	17	5	-	3
2001 3/.....	119	69	55	43	34	46	30	21	16	7	-	1
2000.....	118	76	46	40	41	33	24	18	6	3	2	2
1999.....	99	72	62	56	47	40	32	18	6	4	3	-
American Indian 1/, male												
2004.....	102	62	48	34	29	16	14	13	6	-	-	4
2003.....	92	57	44	35	27	19	13	13	7	-	1	1
2002.....	98	62	28	47	21	23	14	5	6	1	-	2
2001 3/.....	92	53	36	21	20	27	17	9	4	1	-	1
2000.....	94	53	28	21	27	23	13	12	2	1	-	2
1999.....	68	57	40	37	32	20	20	8	1	1	2	-
American Indian 1/, female												
2004.....	35	15	17	25	25	17	24	11	6	2	-	2
2003.....	50	32	23	23	14	23	16	15	9	3	1	-
2002.....	37	26	27	17	13	15	17	4	11	4	-	1
2001 3/.....	27	16	19	22	14	19	13	12	12	6	-	-
2000.....	24	23	18	19	14	10	11	6	4	2	2	-
1999.....	31	15	22	19	15	20	12	10	5	3	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
XX. External causes of mortality (*U01-*U03,V01-Y89)												
Asian or Pacific Islander 2/, both sexes												
2004.....	227	166	146	124	145	147	142	124	60	16	2	-
2003.....	210	173	118	125	143	163	168	101	61	16	6	1
2002.....	182	142	115	152	127	141	137	100	59	17	8	3
2001 3/.....	155	127	118	106	122	122	130	103	44	19	4	4
2000.....	163	120	109	97	105	135	108	82	56	22	3	1
1999.....	165	126	129	83	116	113	97	91	34	4	2	3
Asian or Pacific Islander 2/, male												
2004.....	142	115	88	69	73	62	74	72	24	6	-	-
2003.....	135	112	68	64	79	84	100	50	33	8	1	1
2002.....	117	96	67	94	68	75	69	49	31	7	4	3
2001 3/.....	96	86	74	64	73	71	65	66	23	11	1	2
2000.....	104	76	72	58	59	74	60	40	25	10	3	1
1999.....	98	83	81	43	57	67	56	48	24	2	-	2
Asian or Pacific Islander 2/, female												
2004.....	85	51	58	55	72	85	68	52	36	10	2	-
2003.....	75	61	50	61	64	79	68	51	28	8	5	-
2002.....	65	46	48	58	59	66	68	51	28	10	4	-
2001 3/.....	59	41	44	42	49	51	65	37	21	8	3	2
2000.....	59	44	37	39	46	61	48	42	31	12	-	-
1999.....	67	43	48	40	59	46	41	43	10	2	2	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Accidents (unintentional injuries) (V01-X59,Y85-Y86)												
All races, both sexes												
2004.....	7,516	5,408	4,243	3,819	4,297	5,900	7,557	6,780	4,666	1,709	292	88
2003.....	6,972	5,186	3,984	3,711	4,370	5,875	7,233	6,839	4,352	1,620	335	76
2002.....	6,434	4,557	3,788	3,672	4,414	5,938	6,966	6,507	4,255	1,595	294	85
2001.....	5,875	4,159	3,499	3,506	4,329	5,972	6,716	6,266	4,074	1,519	312	120
2000.....	5,408	4,021	3,484	3,375	4,323	5,596	6,162	5,933	3,939	1,419	304	85
1999.....	4,943	3,944	3,341	3,692	4,516	5,772	6,510	6,138	3,954	1,380	257	106
All races, male												
2004.....	5,386	3,752	2,810	2,448	2,544	3,222	3,808	2,962	1,705	476	67	74
2003.....	4,871	3,613	2,711	2,365	2,556	3,251	3,600	2,991	1,595	440	67	70
2002.....	4,552	3,255	2,526	2,369	2,662	3,268	3,419	2,795	1,488	480	54	74
2001.....	4,224	2,872	2,356	2,200	2,551	3,310	3,342	2,730	1,486	381	60	104
2000.....	3,960	2,804	2,329	2,117	2,522	3,039	3,024	2,563	1,391	397	62	83
1999.....	3,606	2,753	2,262	2,336	2,656	3,096	3,276	2,647	1,397	376	60	94
All races, female												
2004.....	2,130	1,656	1,433	1,371	1,753	2,678	3,749	3,818	2,961	1,233	225	14
2003.....	2,101	1,573	1,273	1,346	1,814	2,624	3,633	3,848	2,757	1,180	268	6
2002.....	1,882	1,302	1,262	1,303	1,752	2,670	3,547	3,712	2,767	1,115	240	11
2001.....	1,651	1,287	1,143	1,306	1,778	2,662	3,374	3,536	2,588	1,138	252	16
2000.....	1,448	1,217	1,155	1,258	1,801	2,557	3,138	3,370	2,548	1,022	242	2
1999.....	1,337	1,191	1,079	1,356	1,860	2,676	3,234	3,491	2,557	1,004	197	12
White, both sexes												
2004.....	6,158	4,485	3,593	3,269	3,737	5,294	7,000	6,397	4,402	1,632	269	79
2003.....	5,667	4,328	3,373	3,126	3,826	5,311	6,649	6,451	4,088	1,544	307	70
2002.....	5,285	3,789	3,196	3,073	3,894	5,361	6,455	6,120	3,991	1,499	267	78
2001.....	4,814	3,464	2,903	2,995	3,783	5,387	6,201	5,858	3,819	1,419	283	103
2000.....	4,347	3,334	2,941	2,855	3,762	5,039	5,701	5,570	3,693	1,317	272	79
1999.....	3,999	3,248	2,714	3,130	3,897	5,143	5,987	5,696	3,716	1,292	234	89

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Accidents (unintentional injuries) (V01-X59,Y85-Y86)												
White, male												
2004.....	4,391	3,079	2,359	2,107	2,220	2,904	3,519	2,786	1,591	453	62	68
2003.....	3,973	3,009	2,299	1,997	2,262	2,940	3,295	2,819	1,484	416	60	64
2002.....	3,715	2,681	2,126	1,965	2,351	2,939	3,151	2,617	1,374	448	45	67
2001.....	3,454	2,371	1,948	1,880	2,239	2,978	3,072	2,542	1,386	348	53	91
2000.....	3,179	2,297	1,945	1,770	2,207	2,712	2,781	2,400	1,290	357	53	77
1999.....	2,917	2,248	1,852	1,975	2,292	2,740	3,018	2,443	1,292	351	54	79
White, female												
2004.....	1,767	1,406	1,234	1,162	1,517	2,390	3,481	3,611	2,811	1,179	207	11
2003.....	1,694	1,319	1,074	1,129	1,564	2,371	3,354	3,632	2,604	1,128	247	6
2002.....	1,570	1,108	1,070	1,108	1,543	2,422	3,304	3,503	2,617	1,051	222	11
2001.....	1,360	1,093	955	1,115	1,544	2,409	3,129	3,316	2,433	1,071	230	12
2000.....	1,168	1,037	996	1,085	1,555	2,327	2,920	3,170	2,403	960	219	2
1999.....	1,082	1,000	862	1,155	1,605	2,403	2,969	3,253	2,424	941	180	10
Black, both sexes												
2004.....	1,131	758	510	424	411	460	406	256	204	59	21	4
2003.....	1,061	691	478	455	407	398	414	280	190	59	20	5
2002.....	947	626	467	445	387	436	370	291	197	76	20	4
2001.....	900	576	488	421	431	450	386	299	200	74	25	14
2000.....	879	566	431	417	451	424	354	280	191	80	28	5
1999.....	786	569	497	459	492	506	430	351	200	81	19	16
Black, male												
2004.....	839	555	357	278	254	255	219	103	90	17	5	3
2003.....	745	500	336	302	226	230	207	122	72	17	5	5
2002.....	699	478	330	304	235	256	196	131	82	25	5	4
2001.....	663	421	343	272	245	257	205	124	74	21	6	12
2000.....	658	434	316	285	258	253	185	121	79	32	7	5
1999.....	594	422	333	300	300	285	203	163	80	22	5	14

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Accidents (unintentional injuries) (V01-X59,Y85-Y86)												
Black, female												
2004.....	292	203	153	146	157	205	187	153	114	42	16	1
2003.....	316	191	142	153	181	168	207	158	118	42	15	-
2002.....	248	148	137	141	152	180	174	160	115	51	15	-
2001.....	237	155	145	149	186	193	181	175	126	53	19	2
2000.....	221	132	115	132	193	171	169	159	112	48	21	-
1999.....	192	147	164	159	192	221	227	188	120	59	14	2
American Indian 1/, both sexes												
2004.....	104	67	50	40	49	29	36	24	11	2	-	5
2003.....	114	74	54	49	35	35	27	24	15	3	2	1
2002.....	98	66	46	47	32	31	26	7	17	5	-	1
2001.....	80	57	41	33	28	40	23	21	14	7	-	-
2000.....	89	55	39	31	32	29	22	13	6	3	2	1
1999.....	67	52	53	46	36	36	23	16	5	3	2	-
American Indian 1/, male												
2004.....	82	56	38	20	26	13	14	13	6	-	-	3
2003.....	72	46	36	29	22	16	13	10	7	-	1	1
2002.....	71	50	22	36	20	17	13	4	6	1	-	1
2001.....	60	42	26	17	16	22	12	9	4	1	-	-
2000.....	70	34	22	17	19	20	12	9	2	1	-	1
1999.....	47	42	33	29	22	18	11	7	1	1	1	-
American Indian 1/, female												
2004.....	22	11	12	20	23	16	22	11	5	2	-	2
2003.....	42	28	18	20	13	19	14	14	8	3	1	-
2002.....	27	16	24	11	12	14	13	3	11	4	-	-
2001.....	20	15	15	16	12	18	11	12	10	6	-	-
2000.....	19	21	17	14	13	9	10	4	4	2	2	-
1999.....	20	10	20	17	14	18	12	9	4	2	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Accidents (unintentional injuries) (V01-X59,Y85-Y86)												
Asian or Pacific Islander 2/, both sexes												
2004.....	123	98	90	86	100	117	115	103	49	16	2	-
2003.....	130	93	79	81	102	131	143	84	59	14	6	-
2002.....	104	76	79	107	101	110	115	89	50	15	7	2
2001.....	81	62	67	57	87	95	106	88	41	19	4	3
2000.....	93	66	73	72	78	104	85	70	49	19	2	-
1999.....	91	75	77	57	91	87	70	75	33	4	2	1
Asian or Pacific Islander 2/, male												
2004.....	74	62	56	43	44	50	56	60	18	6	-	-
2003.....	81	58	40	37	46	65	85	40	32	7	1	-
2002.....	67	46	48	64	56	56	59	43	26	6	4	2
2001.....	47	38	39	31	51	53	53	55	22	11	1	1
2000.....	53	39	46	45	38	54	46	33	20	7	2	-
1999.....	48	41	44	32	42	53	44	34	24	2	-	1
Asian or Pacific Islander 2/, female												
2004.....	49	36	34	43	56	67	59	43	31	10	2	-
2003.....	49	35	39	44	56	66	58	44	27	7	5	-
2002.....	37	30	31	43	45	54	56	46	24	9	3	-
2001.....	34	24	28	26	36	42	53	33	19	8	3	2
2000.....	40	27	27	27	40	50	39	37	29	12	-	-
1999.....	43	34	33	25	49	34	26	41	9	2	2	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Transport accidents (V01-V99,Y85)												
All races, both sexes												
2004.....	3,178	2,510	2,000	1,689	1,540	1,749	1,561	930	328	58	4	27
2003.....	3,098	2,507	1,934	1,578	1,632	1,782	1,595	994	350	55	2	20
2002.....	2,970	2,229	1,936	1,597	1,754	1,855	1,578	936	330	47	9	39
2001.....	2,872	2,152	1,709	1,567	1,715	1,943	1,564	903	304	63	2	43
2000.....	2,717	2,115	1,768	1,599	1,716	1,860	1,482	985	298	49	3	33
1999.....	2,524	2,131	1,681	1,727	1,852	1,902	1,566	964	311	50	5	37
All races, male												
2004.....	2,325	1,773	1,341	1,077	921	1,001	904	501	171	27	2	21
2003.....	2,198	1,778	1,309	1,019	956	1,019	866	534	187	30	-	18
2002.....	2,089	1,615	1,262	1,039	1,062	1,060	857	527	180	26	4	36
2001.....	2,032	1,467	1,164	974	1,041	1,106	877	492	163	28	1	37
2000.....	1,922	1,456	1,161	996	1,025	1,050	808	539	167	27	1	33
1999.....	1,782	1,454	1,107	1,091	1,071	1,089	870	551	179	25	3	34
All races, female												
2004.....	853	737	659	612	619	748	657	429	157	31	2	6
2003.....	900	729	625	559	676	763	729	460	163	25	2	2
2002.....	881	614	674	558	692	795	721	409	150	21	5	3
2001.....	840	685	545	593	674	837	687	411	141	35	1	6
2000.....	795	659	607	603	691	810	674	446	131	22	2	-
1999.....	742	677	574	636	781	813	696	413	132	25	2	3
White, both sexes												
2004.....	2,601	2,135	1,719	1,458	1,347	1,548	1,447	881	305	55	4	25
2003.....	2,564	2,109	1,644	1,338	1,439	1,586	1,462	924	322	51	1	19
2002.....	2,494	1,900	1,658	1,356	1,565	1,684	1,453	870	309	44	6	34
2001.....	2,432	1,834	1,443	1,346	1,511	1,767	1,450	843	287	58	2	35
2000.....	2,245	1,796	1,523	1,375	1,503	1,683	1,377	914	275	44	3	31
1999.....	2,102	1,802	1,412	1,497	1,637	1,716	1,438	888	281	45	4	28

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Transport accidents (V01-V99,Y85)												
White, male												
2004.....	1,906	1,506	1,143	933	804	893	827	476	157	27	2	20
2003.....	1,832	1,504	1,129	873	860	908	792	495	168	27	-	17
2002.....	1,753	1,371	1,087	875	943	949	780	484	163	25	4	31
2001.....	1,735	1,246	984	836	921	997	817	453	155	27	1	31
2000.....	1,598	1,226	987	845	899	942	743	503	154	25	1	31
1999.....	1,494	1,230	931	950	946	977	796	503	163	20	2	26
White, female												
2004.....	695	629	576	525	543	655	620	405	148	28	2	5
2003.....	732	605	515	465	579	678	670	429	154	24	1	2
2002.....	741	529	571	481	622	735	673	386	146	19	2	3
2001.....	697	588	459	510	590	770	633	390	132	31	1	4
2000.....	647	570	536	530	604	741	634	411	121	19	2	-
1999.....	608	572	481	547	691	739	642	385	118	25	2	2
Black, both sexes												
2004.....	443	289	213	161	124	136	79	32	17	3	-	-
2003.....	394	299	211	176	135	134	86	48	22	2	-	1
2002.....	346	251	200	163	126	117	92	44	13	2	3	3
2001.....	338	252	208	172	141	127	73	38	14	5	-	8
2000.....	361	244	180	165	160	109	74	50	16	3	-	1
1999.....	323	246	190	163	155	138	98	59	22	5	1	9
Black, male												
2004.....	333	205	153	112	81	81	57	17	12	-	-	-
2003.....	286	216	143	116	73	82	51	27	13	1	-	1
2002.....	253	195	138	115	85	83	58	32	10	1	-	3
2001.....	237	183	152	111	78	84	39	23	7	1	-	6
2000.....	258	188	133	120	100	73	47	27	10	2	-	1
1999.....	236	173	134	105	93	85	56	41	11	5	1	8

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Transport accidents (V01-V99,Y85)												
Black, female												
2004.....	110	84	60	49	43	55	22	15	5	3	-	-
2003.....	108	83	68	60	62	52	35	21	9	1	-	-
2002.....	93	56	62	48	41	34	34	12	3	1	3	-
2001.....	101	69	56	61	63	43	34	15	7	4	-	2
2000.....	103	56	47	45	60	36	27	23	6	1	-	-
1999.....	87	73	56	58	62	53	42	18	11	-	-	1
American Indian 1/, both sexes												
2004.....	56	33	18	22	26	16	10	1	1	-	-	2
2003.....	56	38	27	19	17	9	7	8	1	-	-	-
2002.....	63	31	32	20	15	11	7	2	2	1	-	-
2001.....	44	34	16	20	11	16	9	4	1	-	-	-
2000.....	43	26	24	19	15	14	5	3	2	1	-	1
1999.....	39	36	23	27	17	14	9	1	-	-	-	-
American Indian 1/, male												
2004.....	45	30	16	10	16	8	5	-	1	-	-	1
2003.....	34	23	14	10	7	3	2	3	1	-	-	-
2002.....	46	23	14	15	8	8	6	1	2	-	-	-
2001.....	32	23	9	11	7	9	6	2	-	-	-	-
2000.....	32	12	13	9	10	9	4	2	1	-	-	1
1999.....	27	28	14	14	12	9	5	-	-	-	-	-
American Indian 1/, female												
2004.....	11	3	2	12	10	8	5	1	-	-	-	1
2003.....	22	15	13	9	10	6	5	5	-	-	-	-
2002.....	17	8	18	5	7	3	1	1	-	1	-	-
2001.....	12	11	7	9	4	7	3	2	1	-	-	-
2000.....	11	14	11	10	5	5	1	1	1	1	-	-
1999.....	12	8	9	13	5	5	4	1	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Transport accidents (V01-V99,Y85)												
Asian or Pacific Islander 2/, both sexes												
2004.....	78	53	50	48	43	49	25	16	5	-	-	-
2003.....	84	61	52	45	41	53	40	14	5	2	1	-
2002.....	67	47	46	58	48	43	26	20	6	-	-	2
2001.....	58	32	42	29	52	33	32	18	2	-	-	-
2000.....	68	49	41	40	38	54	26	18	5	1	-	-
1999.....	60	47	56	40	43	34	21	16	8	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	41	32	29	22	20	19	15	8	1	-	-	-
2003.....	46	35	23	20	16	26	21	9	5	2	-	-
2002.....	37	26	23	34	26	20	13	10	5	-	-	2
2001.....	28	15	19	16	35	16	15	14	1	-	-	-
2000.....	34	30	28	22	16	26	14	7	2	-	-	-
1999.....	25	23	28	22	20	18	13	7	5	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	37	21	21	26	23	30	10	8	4	-	-	-
2003.....	38	26	29	25	25	27	19	5	-	-	1	-
2002.....	30	21	23	24	22	23	13	10	1	-	-	-
2001.....	30	17	23	13	17	17	17	4	1	-	-	-
2000.....	34	19	13	18	22	28	12	11	3	1	-	-
1999.....	35	24	28	18	23	16	8	9	3	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Railway accidents (V05,V15,V80.6,V81.2-V81.9)												
All races, both sexes												
2004.....	39	22	15	9	12	10	7	7	2	-	-	-
2003.....	33	26	10	8	5	8	6	1	-	-	-	1
2002.....	38	24	13	9	6	4	6	3	1	-	-	3
2001.....	39	21	18	7	11	11	7	3	-	1	-	8
2000.....	33	16	15	8	7	5	6	4	2	-	-	6
1999.....	27	16	14	6	8	7	6	2	1	-	-	6
All races, male												
2004.....	35	21	12	5	8	8	6	5	1	-	-	-
2003.....	27	22	8	8	3	4	5	2	-	-	-	1
2002.....	34	23	13	6	5	2	4	2	1	-	-	2
2001.....	33	19	13	6	8	10	2	-	-	-	-	8
2000.....	28	15	13	7	6	4	4	2	1	-	-	6
1999.....	21	15	11	2	7	4	4	2	1	-	-	6
All races, female												
2004.....	4	1	3	4	4	2	1	2	1	-	-	-
2003.....	6	4	2	-	2	4	1	1	-	-	-	-
2002.....	4	1	-	3	1	2	2	1	-	-	-	1
2001.....	6	2	5	1	3	1	5	3	-	1	-	-
2000.....	5	1	2	1	1	1	2	2	1	-	-	-
1999.....	6	1	3	4	1	3	2	-	-	-	-	-
White, both sexes												
2004.....	25	19	6	5	12	6	7	7	2	-	-	-
2003.....	29	22	10	6	4	6	4	1	-	-	-	1
2002.....	29	15	10	6	5	4	4	1	1	-	-	3
2001.....	34	18	12	6	9	10	5	3	-	1	-	6
2000.....	22	11	14	6	7	5	4	3	2	-	-	6
1999.....	21	12	11	5	7	6	5	1	1	-	-	5

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Railway accidents (V05,V15,V80.6,V81.2-V81.9)												
White, male												
2004.....	23	18	6	4	8	4	6	5	1	-	-	-
2003.....	24	18	8	6	3	3	3	-	-	-	-	1
2002.....	26	15	10	4	4	2	2	1	1	-	-	2
2001.....	28	16	10	5	6	9	-	-	-	-	-	6
2000.....	19	10	12	5	6	4	3	2	1	-	-	6
1999.....	17	11	9	1	6	3	3	1	1	-	-	5
White, female												
2004.....	2	1	-	1	4	2	1	2	1	-	-	-
2003.....	5	4	2	-	1	3	1	1	-	-	-	-
2002.....	3	-	-	2	1	2	2	-	-	-	-	1
2001.....	6	2	2	1	3	1	5	3	-	1	-	-
2000.....	3	1	2	1	1	1	1	1	1	-	-	-
1999.....	4	1	2	4	1	3	2	-	-	-	-	-
Black, both sexes												
2004.....	9	3	8	3	-	4	-	-	-	-	-	-
2003.....	4	3	-	2	1	2	2	-	-	-	-	-
2002.....	8	7	3	2	1	-	2	-	-	-	-	-
2001.....	4	3	6	1	2	1	-	-	-	-	-	2
2000.....	8	4	1	2	-	-	2	1	-	-	-	-
1999.....	6	4	2	1	1	-	-	1	-	-	-	1
Black, male												
2004.....	8	3	5	1	-	4	-	-	-	-	-	-
2003.....	3	3	-	2	-	1	2	-	-	-	-	-
2002.....	7	6	3	2	1	-	2	-	-	-	-	-
2001.....	4	3	3	1	2	1	-	-	-	-	-	2
2000.....	6	4	1	2	-	-	1	-	-	-	-	-
1999.....	4	4	1	1	1	-	-	1	-	-	-	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)												
All races, both sexes												
2004.....	2,854	2,293	1,800	1,547	1,427	1,683	1,497	887	319	57	4	26
2003.....	2,788	2,232	1,726	1,444	1,523	1,680	1,530	961	343	53	1	16
2002.....	2,697	2,004	1,760	1,469	1,644	1,763	1,508	909	321	47	8	35
2001.....	2,553	1,939	1,526	1,409	1,581	1,823	1,489	868	294	59	2	33
2000.....	2,425	1,917	1,589	1,436	1,602	1,764	1,409	953	285	47	3	26
1999.....	2,225	1,885	1,485	1,565	1,711	1,772	1,471	920	291	46	4	30
All races, male												
2004.....	2,055	1,596	1,174	965	834	954	857	472	167	26	2	20
2003.....	1,937	1,553	1,130	908	870	944	823	511	182	29	-	14
2002.....	1,861	1,423	1,117	923	965	992	801	512	176	26	4	33
2001.....	1,774	1,296	1,013	846	940	1,027	832	478	157	28	1	27
2000.....	1,689	1,293	1,013	870	932	980	759	517	161	27	1	26
1999.....	1,537	1,250	950	969	962	1,012	815	524	164	24	2	27
All races, female												
2004.....	799	697	626	582	593	729	640	415	152	31	2	6
2003.....	851	679	596	536	653	736	707	450	161	24	1	2
2002.....	836	581	643	546	679	771	707	397	145	21	4	2
2001.....	779	643	513	563	641	796	657	390	137	31	1	6
2000.....	736	624	576	566	670	784	650	436	124	20	2	-
1999.....	688	635	535	596	749	760	656	396	127	22	2	3
White, both sexes												
2004.....	2,327	1,941	1,547	1,332	1,246	1,495	1,385	841	296	54	4	24
2003.....	2,295	1,861	1,458	1,226	1,338	1,497	1,404	892	316	49	-	15
2002.....	2,256	1,708	1,501	1,248	1,466	1,605	1,391	846	300	44	5	30
2001.....	2,153	1,654	1,282	1,213	1,386	1,661	1,382	813	279	55	2	29
2000.....	1,989	1,622	1,362	1,234	1,402	1,598	1,308	885	262	43	3	24
1999.....	1,847	1,587	1,244	1,356	1,512	1,603	1,355	849	263	41	4	22

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)

White, male

2004.....	1,678	1,349	998	831	725	854	781	449	153	26	2	19
2003.....	1,607	1,302	968	781	779	841	755	473	164	26	-	13
2002.....	1,557	1,210	957	777	857	894	730	471	159	25	4	28
2001.....	1,509	1,106	846	730	828	925	776	441	150	27	1	25
2000.....	1,392	1,085	852	736	816	879	696	482	148	25	1	24
1999.....	1,284	1,051	798	841	851	907	748	480	150	19	2	20

White, female

2004.....	649	592	549	501	521	641	604	392	143	28	2	5
2003.....	688	559	490	445	559	656	649	419	152	23	-	2
2002.....	699	498	544	471	609	711	661	375	141	19	1	2
2001.....	644	548	436	483	558	736	606	372	129	28	1	4
2000.....	597	537	510	498	586	719	612	403	114	18	2	-
1999.....	563	536	446	515	661	696	607	369	113	22	2	2

Black, both sexes

2004.....	409	267	189	151	117	123	78	32	17	3	-	-
2003.....	364	277	194	161	130	125	81	48	21	2	-	1
2002.....	318	224	187	150	118	107	86	43	13	2	3	3
2001.....	308	226	189	153	136	116	70	37	13	4	-	4
2000.....	342	225	169	150	151	102	71	47	16	2	-	1
1999.....	288	224	169	150	145	128	90	54	21	5	-	8

Black, male

2004.....	303	186	134	105	75	73	56	17	12	-	-	-
2003.....	258	197	127	102	70	77	47	27	12	1	-	1
2002.....	227	170	126	102	77	73	53	31	10	1	-	3
2001.....	214	158	140	94	73	78	38	23	7	1	-	2
2000.....	242	171	124	107	94	68	46	26	10	2	-	1
1999.....	207	155	116	96	84	82	52	37	10	5	-	7

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)

Black, female

2004.....	106	81	55	46	42	50	22	15	5	3	-	-
2003.....	106	80	67	59	60	48	34	21	9	1	-	-
2002.....	91	54	61	48	41	34	33	12	3	1	3	-
2001.....	94	68	49	59	63	38	32	14	6	3	-	2
2000.....	100	54	45	43	57	34	25	21	6	-	-	-
1999.....	81	69	53	54	61	46	38	17	11	-	-	1

American Indian 1/, both sexes

2004.....	44	33	16	18	25	16	10	1	1	-	-	2
2003.....	53	35	26	17	15	9	7	7	1	-	-	-
2002.....	62	30	32	18	13	11	7	2	2	1	-	-
2001.....	38	29	15	18	9	16	7	4	1	-	-	-
2000.....	37	24	23	17	14	13	5	3	2	1	-	1
1999.....	35	30	21	23	14	14	9	1	-	-	-	-

American Indian 1/, male

2004.....	36	30	14	8	15	8	5	-	1	-	-	1
2003.....	33	21	14	8	5	3	2	2	1	-	-	-
2002.....	45	22	14	13	6	8	6	1	2	-	-	-
2001.....	27	18	8	9	6	9	4	2	-	-	-	-
2000.....	26	10	13	9	9	8	4	2	1	-	-	1
1999.....	25	23	13	12	9	9	5	-	-	-	-	-

American Indian 1/, female

2004.....	8	3	2	10	10	8	5	1	-	-	-	1
2003.....	20	14	12	9	10	6	5	5	-	-	-	-
2002.....	17	8	18	5	7	3	1	1	-	1	-	-
2001.....	11	11	7	9	3	7	3	2	1	-	-	-
2000.....	11	14	10	8	5	5	1	1	1	-	-	-
1999.....	10	7	8	11	5	5	4	1	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)

Asian or Pacific Islander 2/, both sexes

2004.....	74	52	48	46	39	49	24	13	5	-	-	-
2003.....	76	59	48	40	40	49	38	14	5	2	1	-
2002.....	61	42	40	53	47	40	24	18	6	-	-	2
2001.....	54	30	40	25	50	30	30	14	1	-	-	-
2000.....	57	46	35	35	35	51	25	18	5	1	-	-
1999.....	55	44	51	36	40	27	17	16	7	-	-	-

Asian or Pacific Islander 2/, male

2004.....	38	31	28	21	19	19	15	6	1	-	-	-
2003.....	39	33	21	17	16	23	19	9	5	2	-	-
2002.....	32	21	20	31	25	17	12	9	5	-	-	2
2001.....	24	14	19	13	33	15	14	12	-	-	-	-
2000.....	29	27	24	18	13	25	13	7	2	-	-	-
1999.....	21	21	23	20	18	14	10	7	4	-	-	-

Asian or Pacific Islander 2/, female

2004.....	36	21	20	25	20	30	9	7	4	-	-	-
2003.....	37	26	27	23	24	26	19	5	-	-	1	-
2002.....	29	21	20	22	22	23	12	9	1	-	-	-
2001.....	30	16	21	12	17	15	16	2	1	-	-	-
2000.....	28	19	11	17	22	26	12	11	3	1	-	-
1999.....	34	23	28	16	22	13	7	9	3	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pedestrian involved in collision with motor vehicle (V02-V04)												
All races, both sexes												
2004.....	273	229	192	132	151	165	157	90	33	9	-	6
2003.....	244	227	182	158	136	174	167	95	37	7	-	4
2002.....	251	208	166	180	184	177	148	102	39	10	1	11
2001.....	249	197	148	111	173	177	195	92	34	6	-	14
2000.....	230	201	158	142	154	184	162	110	39	5	-	6
1999.....	208	206	158	134	171	168	137	118	35	1	1	8
All races, male												
2004.....	219	157	131	85	81	90	89	53	16	5	-	4
2003.....	184	173	127	103	89	84	84	51	26	6	-	4
2002.....	183	147	107	115	110	92	79	50	24	6	1	10
2001.....	173	140	107	62	111	98	108	54	19	3	-	11
2000.....	160	147	106	93	88	103	83	60	22	4	-	6
1999.....	161	148	107	88	99	95	85	59	18	-	1	7
All races, female												
2004.....	54	72	61	47	70	75	68	37	17	4	-	2
2003.....	60	54	55	55	47	90	83	44	11	1	-	-
2002.....	68	61	59	65	74	85	69	52	15	4	-	1
2001.....	76	57	41	49	62	79	87	38	15	3	-	3
2000.....	70	54	52	49	66	81	79	50	17	1	-	-
1999.....	47	58	51	46	72	73	52	59	17	1	-	1
White, both sexes												
2004.....	200	185	148	105	127	136	140	77	28	8	-	6
2003.....	181	188	152	121	109	142	143	83	34	6	-	3
2002.....	191	170	132	135	156	146	125	87	34	9	1	7
2001.....	185	162	111	91	139	147	165	78	33	6	-	12
2000.....	177	155	128	112	128	155	143	93	33	4	-	6
1999.....	158	154	127	108	134	140	119	105	27	1	1	4

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pedestrian involved in collision with motor vehicle (V02-V04)												
White, male												
2004.....	158	122	102	68	66	75	77	44	14	5	-	4
2003.....	139	147	107	78	71	67	72	42	23	5	-	3
2002.....	138	120	87	88	91	75	65	42	20	5	1	6
2001.....	130	113	77	50	90	80	96	43	19	3	-	10
2000.....	126	111	86	69	72	83	69	48	18	3	-	6
1999.....	122	108	88	75	74	77	72	49	15	-	1	4
White, female												
2004.....	42	63	46	37	61	61	63	33	14	3	-	2
2003.....	42	41	45	43	38	75	71	41	11	1	-	-
2002.....	53	50	45	47	65	71	60	45	14	4	-	1
2001.....	55	49	34	41	49	67	69	35	14	3	-	2
2000.....	51	44	42	43	56	72	74	45	15	1	-	-
1999.....	36	46	39	33	60	63	47	56	12	1	-	-
Black, both sexes												
2004.....	60	37	29	13	15	16	9	6	3	1	-	-
2003.....	47	32	20	24	17	18	9	5	1	-	-	1
2002.....	48	29	25	25	15	14	15	6	2	1	-	3
2001.....	53	29	28	14	19	15	12	6	1	-	-	2
2000.....	41	40	21	19	17	18	7	9	2	1	-	-
1999.....	38	45	19	14	23	15	9	9	4	-	-	4
Black, male												
2004.....	53	30	23	10	10	10	6	5	2	-	-	-
2003.....	37	23	15	20	15	12	6	4	1	-	-	1
2002.....	36	23	18	16	11	11	11	3	1	1	-	3
2001.....	36	23	25	9	13	11	4	4	-	-	-	1
2000.....	28	32	14	16	13	14	6	8	2	1	-	-
1999.....	32	35	15	9	17	9	5	8	1	-	-	3

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pedestrian involved in collision with motor vehicle (V02-V04)												
Black, female												
2004.....	7	7	6	3	5	6	3	1	1	1	-	-
2003.....	10	9	5	4	2	6	3	1	-	-	-	-
2002.....	12	6	7	9	4	3	4	3	1	-	-	-
2001.....	17	6	3	5	6	4	8	2	1	-	-	1
2000.....	13	8	7	3	4	4	1	1	-	-	-	-
1999.....	6	10	4	5	6	6	4	1	3	-	-	1
American Indian 1/, both sexes												
2004.....	4	4	1	4	1	1	-	-	-	-	-	-
2003.....	7	2	3	1	1	1	-	-	-	-	-	-
2002.....	7	3	3	2	2	1	1	1	1	-	-	-
2001.....	4	3	2	1	3	-	-	-	-	-	-	-
2000.....	4	2	2	4	1	-	1	-	2	-	-	-
1999.....	6	5	1	4	5	4	2	-	-	-	-	-
American Indian 1/, male												
2004.....	3	4	1	3	1	1	-	-	-	-	-	-
2003.....	4	1	3	1	1	-	-	-	-	-	-	-
2002.....	6	2	1	2	2	1	1	-	1	-	-	-
2001.....	4	2	2	1	3	-	-	-	-	-	-	-
2000.....	3	2	1	4	1	-	1	-	1	-	-	-
1999.....	4	4	-	1	4	3	2	-	-	-	-	-
American Indian 1/, female												
2004.....	1	-	-	1	-	-	-	-	-	-	-	-
2003.....	3	1	-	-	-	1	-	-	-	-	-	-
2002.....	1	1	2	-	-	-	-	1	-	-	-	-
2001.....	-	1	-	-	-	-	-	-	-	-	-	-
2000.....	1	-	-	-	-	-	-	-	1	-	-	-
1999.....	2	1	1	3	1	1	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pedestrian involved in collision with motor vehicle (V02-V04)												
Asian or Pacific Islander 2/, both sexes												
2004.....	9	3	14	10	8	12	8	7	2	-	-	-
2003.....	9	5	7	12	9	13	15	7	2	1	-	-
2002.....	5	6	6	18	11	16	7	8	2	-	-	1
2001.....	7	3	7	5	12	15	18	8	-	-	-	-
2000.....	8	4	7	7	8	11	11	8	2	-	-	-
1999.....	6	2	11	8	9	9	7	4	4	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	5	1	5	4	4	4	6	4	-	-	-	-
2003.....	4	2	2	4	2	5	6	5	2	1	-	-
2002.....	3	2	1	9	6	5	2	5	2	-	-	1
2001.....	3	2	3	2	5	7	8	7	-	-	-	-
2000.....	3	2	5	4	2	6	7	4	1	-	-	-
1999.....	3	1	4	3	4	6	6	2	2	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	4	2	9	6	4	8	2	3	2	-	-	-
2003.....	5	3	5	8	7	8	9	2	-	-	-	-
2002.....	2	4	5	9	5	11	5	3	-	-	-	-
2001.....	4	1	4	3	7	8	10	1	-	-	-	-
2000.....	5	2	2	3	6	5	4	4	1	-	-	-
1999.....	3	1	7	5	5	3	1	2	2	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pedalcyclist involved in collision with motor vehicle (V12-V14,V19.0-V19.2,V19.4-V19.6)												
All races, both sexes												
2004.....	55	38	25	27	22	20	8	6	1	-	-	-
2003.....	54	36	20	12	23	13	7	1	1	-	-	-
2002.....	34	26	20	15	13	13	7	4	-	-	-	1
2001.....	44	34	21	22	14	13	6	3	1	1	1	3
2000.....	35	36	17	19	14	13	9	5	-	-	-	1
1999.....	33	21	12	21	16	6	4	7	-	1	-	1
All races, male												
2004.....	48	32	23	26	21	20	8	5	-	-	-	-
2003.....	48	32	20	10	21	12	7	1	-	-	-	-
2002.....	28	24	18	15	11	11	7	4	-	-	-	1
2001.....	41	31	21	22	14	13	6	3	1	1	1	3
2000.....	32	32	16	17	13	13	7	4	-	-	-	1
1999.....	30	18	9	20	16	6	4	7	-	1	-	1
All races, female												
2004.....	7	6	2	1	1	-	-	1	1	-	-	-
2003.....	6	4	-	2	2	1	-	-	1	-	-	-
2002.....	6	2	2	-	2	2	-	-	-	-	-	-
2001.....	3	3	-	-	-	-	-	-	-	-	-	-
2000.....	3	4	1	2	1	-	2	1	-	-	-	-
1999.....	3	3	3	1	-	-	-	-	-	-	-	-
White, both sexes												
2004.....	47	32	23	24	18	18	8	6	1	-	-	-
2003.....	42	29	19	10	22	11	6	1	1	-	-	-
2002.....	30	24	16	12	10	11	7	4	-	-	-	1
2001.....	33	29	20	14	10	13	5	2	1	1	1	2
2000.....	24	33	15	13	14	11	8	3	-	-	-	1
1999.....	28	19	11	14	14	5	4	7	-	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pedalcyclist involved in collision with motor vehicle (V12-V14,V19.0-V19.2,V19.4-V19.6)												
White, male												
2004.....	40	27	21	23	17	18	8	5	-	-	-	-
2003.....	37	25	19	9	20	10	6	1	-	-	-	-
2002.....	24	22	14	12	8	9	7	4	-	-	-	1
2001.....	30	26	20	14	10	13	5	2	1	1	1	2
2000.....	22	29	14	11	13	11	6	2	-	-	-	1
1999.....	25	16	8	13	14	5	4	7	-	1	-	-
White, female												
2004.....	7	5	2	1	1	-	-	1	1	-	-	-
2003.....	5	4	-	1	2	1	-	-	1	-	-	-
2002.....	6	2	2	-	2	2	-	-	-	-	-	-
2001.....	3	3	-	-	-	-	-	-	-	-	-	-
2000.....	2	4	1	2	1	-	2	1	-	-	-	-
1999.....	3	3	3	1	-	-	-	-	-	-	-	-
Black, both sexes												
2004.....	8	5	1	1	1	-	-	-	-	-	-	-
2003.....	10	6	-	-	-	1	-	-	-	-	-	-
2002.....	3	2	2	3	1	1	-	-	-	-	-	-
2001.....	11	5	1	6	2	-	-	-	-	-	-	1
2000.....	10	3	2	5	-	-	-	2	-	-	-	-
1999.....	4	1	1	3	1	-	-	-	-	-	-	1
Black, male												
2004.....	8	4	1	1	1	-	-	-	-	-	-	-
2003.....	9	6	-	-	-	1	-	-	-	-	-	-
2002.....	3	2	2	3	1	1	-	-	-	-	-	-
2001.....	11	5	1	6	2	-	-	-	-	-	-	1
2000.....	10	3	2	5	-	-	-	2	-	-	-	-
1999.....	4	1	1	3	1	-	-	-	-	-	-	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Motorcyclist involved in any accident except collision with railway train (V20-V24,V26-V34,V36-V39)												
All races, both sexes												
2004.....	398	243	125	71	31	24	10	1	1	1	-	-
2003.....	337	233	98	57	39	12	14	1	3	-	-	-
2002.....	300	175	92	49	26	14	10	8	1	1	-	-
2001.....	237	117	72	37	27	9	12	4	-	1	-	-
2000.....	192	115	68	43	29	11	1	4	2	-	-	1
1999.....	153	109	61	32	16	12	1	3	-	-	-	-
All races, male												
2004.....	351	223	110	68	30	19	8	1	1	-	-	-
2003.....	300	210	89	54	37	10	12	-	3	-	-	-
2002.....	267	154	87	45	25	13	8	7	-	1	-	-
2001.....	209	107	66	35	24	9	11	4	-	1	-	-
2000.....	174	101	58	38	27	10	1	4	2	-	-	1
1999.....	134	96	58	31	15	11	1	2	-	-	-	-
All races, female												
2004.....	47	20	15	3	1	5	2	-	-	1	-	-
2003.....	37	23	9	3	2	2	2	1	-	-	-	-
2002.....	33	21	5	4	1	1	2	1	1	-	-	-
2001.....	28	10	6	2	3	-	1	-	-	-	-	-
2000.....	18	14	10	5	2	1	-	-	-	-	-	-
1999.....	19	13	3	1	1	1	-	1	-	-	-	-
White, both sexes												
2004.....	374	231	121	69	31	23	9	1	1	1	-	-
2003.....	322	224	95	57	36	12	14	1	3	-	-	-
2002.....	291	169	89	45	24	14	8	8	1	1	-	-
2001.....	229	111	67	35	23	8	12	3	-	1	-	-
2000.....	179	112	65	40	26	9	1	3	2	-	-	1
1999.....	146	102	57	30	16	12	1	3	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Motor vehicle accident involving collision with railway train (V25,V35,V45,V55,V65,V75,V81.0-V81.1,V87.6,V88.6)												
All races, both sexes												
2004.....	13	12	7	8	7	6	10	5	3	-	-	-
2003.....	11	17	10	7	12	15	8	2	1	1	-	-
2002.....	15	15	10	14	5	6	11	7	2	2	-	-
2001.....	13	7	10	7	11	14	5	2	-	-	-	-
2000.....	17	13	9	9	12	12	9	6	1	-	-	-
1999.....	11	15	12	5	13	12	9	8	2	1	-	-
All races, male												
2004.....	11	10	6	6	4	4	10	4	2	-	-	-
2003.....	7	15	7	5	9	9	7	1	1	1	-	-
2002.....	12	10	5	7	2	4	5	3	1	1	-	-
2001.....	10	5	8	3	8	6	3	1	-	-	-	-
2000.....	15	10	6	6	10	10	4	4	1	-	-	-
1999.....	7	9	6	3	8	11	5	4	1	-	-	-
All races, female												
2004.....	2	2	1	2	3	2	-	1	1	-	-	-
2003.....	4	2	3	2	3	6	1	1	-	-	-	-
2002.....	3	5	5	7	3	2	6	4	1	1	-	-
2001.....	3	2	2	4	3	8	2	1	-	-	-	-
2000.....	2	3	3	3	2	2	5	2	-	-	-	-
1999.....	4	6	6	2	5	1	4	4	1	1	-	-
White, both sexes												
2004.....	10	10	4	6	6	6	9	5	3	-	-	-
2003.....	9	16	9	7	10	12	7	2	1	1	-	-
2002.....	13	15	8	12	5	6	11	7	1	2	-	-
2001.....	10	6	10	7	11	13	5	2	-	-	-	-
2000.....	16	10	8	7	11	10	8	6	1	-	-	-
1999.....	10	14	11	4	12	12	9	6	1	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Motor vehicle accident involving collision with railway train (V25,V35,V45,V55,V65,V75,V81.0-V81.1,V87.6,V88.6)												
White, male												
2004.....	8	9	3	4	3	4	9	4	2	-	-	-
2003.....	6	14	7	5	9	7	6	1	1	1	-	-
2002.....	10	10	4	6	2	4	5	3	-	1	-	-
2001.....	9	4	8	3	8	6	3	1	-	-	-	-
2000.....	14	9	5	4	9	9	3	4	1	-	-	-
1999.....	7	8	5	3	8	11	5	3	-	-	-	-
White, female												
2004.....	2	1	1	2	3	2	-	1	1	-	-	-
2003.....	3	2	2	2	1	5	1	1	-	-	-	-
2002.....	3	5	4	6	3	2	6	4	1	1	-	-
2001.....	1	2	2	4	3	7	2	1	-	-	-	-
2000.....	2	1	3	3	2	1	5	2	-	-	-	-
1999.....	3	6	6	1	4	1	4	3	1	1	-	-
Black, both sexes												
2004.....	3	2	3	2	1	-	1	-	-	-	-	-
2003.....	1	1	-	-	1	3	1	-	-	-	-	-
2002.....	2	-	2	1	-	-	-	-	1	-	-	-
2001.....	3	-	-	-	-	1	-	-	-	-	-	-
2000.....	1	3	1	2	1	2	1	-	-	-	-	-
1999.....	1	1	1	1	1	-	-	2	1	-	-	-
Black, male												
2004.....	3	1	3	2	1	-	1	-	-	-	-	-
2003.....	1	1	-	-	-	2	1	-	-	-	-	-
2002.....	2	-	1	-	-	-	-	-	1	-	-	-
2001.....	1	-	-	-	-	-	-	-	-	-	-	-
2000.....	1	1	1	2	1	1	1	-	-	-	-	-
1999.....	-	1	1	-	-	-	-	1	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other motor vehicle accident involving collision with railway train (V45,V55,V65,V75,V81.0-V81.1,V87.6,V88.6)												
All races, both sexes												
2004.....	13	12	7	8	7	6	10	5	3	-	-	-
2003.....	10	17	9	7	12	15	8	2	-	1	-	-
2002.....	15	15	10	14	5	6	11	7	2	2	-	-
2001.....	13	7	10	7	11	14	5	2	-	-	-	-
2000.....	16	13	9	9	12	12	9	6	1	-	-	-
1999.....	11	15	12	5	13	12	9	8	2	1	-	-
All races, male												
2004.....	11	10	6	6	4	4	10	4	2	-	-	-
2003.....	6	15	6	5	9	9	7	1	-	1	-	-
2002.....	12	10	5	7	2	4	5	3	1	1	-	-
2001.....	10	5	8	3	8	6	3	1	-	-	-	-
2000.....	14	10	6	6	10	10	4	4	1	-	-	-
1999.....	7	9	6	3	8	11	5	4	1	-	-	-
All races, female												
2004.....	2	2	1	2	3	2	-	1	1	-	-	-
2003.....	4	2	3	2	3	6	1	1	-	-	-	-
2002.....	3	5	5	7	3	2	6	4	1	1	-	-
2001.....	3	2	2	4	3	8	2	1	-	-	-	-
2000.....	2	3	3	3	2	2	5	2	-	-	-	-
1999.....	4	6	6	2	5	1	4	4	1	1	-	-
White, both sexes												
2004.....	10	10	4	6	6	6	9	5	3	-	-	-
2003.....	8	16	8	7	10	12	7	2	-	1	-	-
2002.....	13	15	8	12	5	6	11	7	1	2	-	-
2001.....	10	6	10	7	11	13	5	2	-	-	-	-
2000.....	15	10	8	7	11	10	8	6	1	-	-	-
1999.....	10	14	11	4	12	12	9	6	1	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other motor vehicle accident involving collision with railway train (V45,V55,V65,V75,V81.0-V81.1,V87.6,V88.6)												
White, male												
2004.....	8	9	3	4	3	4	9	4	2	-	-	-
2003.....	5	14	6	5	9	7	6	1	-	1	-	-
2002.....	10	10	4	6	2	4	5	3	-	1	-	-
2001.....	9	4	8	3	8	6	3	1	-	-	-	-
2000.....	13	9	5	4	9	9	3	4	1	-	-	-
1999.....	7	8	5	3	8	11	5	3	-	-	-	-
White, female												
2004.....	2	1	1	2	3	2	-	1	1	-	-	-
2003.....	3	2	2	2	1	5	1	1	-	-	-	-
2002.....	3	5	4	6	3	2	6	4	1	1	-	-
2001.....	1	2	2	4	3	7	2	1	-	-	-	-
2000.....	2	1	3	3	2	1	5	2	-	-	-	-
1999.....	3	6	6	1	4	1	4	3	1	1	-	-
Black, both sexes												
2004.....	3	2	3	2	1	-	1	-	-	-	-	-
2003.....	1	1	-	-	1	3	1	-	-	-	-	-
2002.....	2	-	2	1	-	-	-	-	1	-	-	-
2001.....	3	-	-	-	-	1	-	-	-	-	-	-
2000.....	1	3	1	2	1	2	1	-	-	-	-	-
1999.....	1	1	1	1	1	-	-	2	1	-	-	-
Black, male												
2004.....	3	1	3	2	1	-	1	-	-	-	-	-
2003.....	1	1	-	-	-	2	1	-	-	-	-	-
2002.....	2	-	1	-	-	-	-	-	1	-	-	-
2001.....	1	-	-	-	-	-	-	-	-	-	-	-
2000.....	1	1	1	2	1	1	1	-	-	-	-	-
1999.....	-	1	1	-	-	-	-	1	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Occupant of motor vehicle involved in collision with other (non-motorized) road vehicle, streetcar, animal or pedestrian (V40-V41,V46-V47,V50-V51,V56-V57, V60-V61,V66-V67,V70-V71,V76-V77)												
All races, both sexes												
2004.....	217	180	151	117	113	107	100	40	14	2	2	-
2003.....	258	172	131	107	136	115	90	64	22	3	-	1
2002.....	235	175	163	114	108	112	91	52	15	3	1	-
2001.....	203	130	107	107	95	126	80	56	13	8	-	1
2000.....	173	124	113	99	119	96	73	46	23	5	-	-
1999.....	166	133	110	122	100	105	80	35	11	4	-	1
All races, male												
2004.....	157	134	103	77	71	75	56	20	8	-	1	-
2003.....	180	132	93	72	85	68	52	42	10	2	-	1
2002.....	165	128	125	74	71	71	42	34	9	1	1	-
2001.....	146	108	79	70	57	76	50	27	8	5	-	1
2000.....	125	86	78	67	69	61	41	25	8	4	-	-
1999.....	131	87	78	83	63	62	51	24	5	1	-	-
All races, female												
2004.....	60	46	48	40	42	32	44	20	6	2	1	-
2003.....	78	40	38	35	51	47	38	22	12	1	-	-
2002.....	70	47	38	40	37	41	49	18	6	2	-	-
2001.....	57	22	28	37	38	50	30	29	5	3	-	-
2000.....	48	38	35	32	50	35	32	21	15	1	-	-
1999.....	35	46	32	39	37	43	29	11	6	3	-	1
White, both sexes												
2004.....	174	142	123	102	97	97	95	36	14	2	2	-
2003.....	204	144	109	85	118	103	86	61	21	3	-	1
2002.....	187	137	135	94	94	104	85	49	14	3	1	-
2001.....	166	114	88	93	83	113	72	55	13	7	-	1
2000.....	134	98	97	80	100	83	66	45	20	5	-	-
1999.....	129	106	88	106	87	89	73	30	11	4	-	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Occupant of motor vehicle involved in collision with other (non-motorized) road vehicle, streetcar, animal or pedestrian (V40-V41,V46-V47,V50-V51,V56-V57, V60-V61,V66-V67,V70-V71,V76-V77)												
White, male												
2004.....	128	100	87	66	60	68	52	19	8	-	1	-
2003.....	146	109	76	58	75	60	50	40	9	2	-	1
2002.....	133	99	104	56	59	65	38	31	8	1	1	-
2001.....	120	95	66	59	50	65	46	26	8	4	-	1
2000.....	98	65	66	53	57	50	36	24	8	4	-	-
1999.....	108	71	64	73	53	54	47	21	5	1	-	-
White, female												
2004.....	46	42	36	36	37	29	43	17	6	2	1	-
2003.....	58	35	33	27	43	43	36	21	12	1	-	-
2002.....	54	38	31	38	35	39	47	18	6	2	-	-
2001.....	46	19	22	34	33	48	26	29	5	3	-	-
2000.....	36	33	31	27	43	33	30	21	12	1	-	-
1999.....	21	35	24	33	34	35	26	9	6	3	-	1
Black, both sexes												
2004.....	39	34	23	12	12	9	3	4	-	-	-	-
2003.....	45	23	19	20	15	12	3	1	1	-	-	-
2002.....	34	35	25	17	12	7	4	2	1	-	-	-
2001.....	28	14	14	10	8	12	8	1	-	1	-	-
2000.....	32	22	13	16	17	8	6	-	3	-	-	-
1999.....	29	22	16	12	11	16	5	3	-	-	-	-
Black, male												
2004.....	27	31	13	11	7	7	2	1	-	-	-	-
2003.....	29	19	15	12	7	8	2	1	1	-	-	-
2002.....	26	26	21	15	10	5	3	2	1	-	-	-
2001.....	21	11	11	8	5	11	4	1	-	1	-	-
2000.....	24	17	11	12	11	8	5	-	-	-	-	-
1999.....	19	14	13	7	9	8	3	3	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Occupant of motor vehicle involved in collision with other (non-motorized) road vehicle, streetcar, animal or pedestrian (V40-V41,V46-V47,V50-V51,V56-V57, V60-V61,V66-V67,V70-V71,V76-V77)												
Black, female												
2004.....	12	3	10	1	5	2	1	3	-	-	-	-
2003.....	16	4	4	8	8	4	1	-	-	-	-	-
2002.....	8	9	4	2	2	2	1	-	-	-	-	-
2001.....	7	3	3	2	3	1	4	-	-	-	-	-
2000.....	8	5	2	4	6	-	1	-	3	-	-	-
1999.....	10	8	3	5	2	8	2	-	-	-	-	-
American Indian 1/, both sexes												
2004.....	2	3	3	-	1	-	1	-	-	-	-	-
2003.....	3	2	1	1	-	-	-	-	-	-	-	-
2002.....	7	2	2	-	-	1	-	-	-	-	-	-
2001.....	3	1	4	2	-	1	-	-	-	-	-	-
2000.....	1	-	-	-	2	-	-	-	-	-	-	-
1999.....	5	1	3	1	-	-	2	1	-	-	-	-
American Indian 1/, male												
2004.....	2	2	2	-	1	-	1	-	-	-	-	-
2003.....	3	2	1	1	-	-	-	-	-	-	-	-
2002.....	3	2	-	-	-	1	-	-	-	-	-	-
2001.....	2	1	2	1	-	-	-	-	-	-	-	-
2000.....	1	-	-	-	1	-	-	-	-	-	-	-
1999.....	3	1	1	1	-	-	1	-	-	-	-	-
American Indian 1/, female												
2004.....	-	1	1	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	4	-	2	-	-	-	-	-	-	-	-	-
2001.....	1	-	2	1	-	1	-	-	-	-	-	-
2000.....	-	-	-	-	1	-	-	-	-	-	-	-
1999.....	2	-	2	-	-	-	1	1	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Occupant of motor vehicle involved in collision with other (non-motorized) road vehicle, streetcar, animal or pedestrian (V40-V41,V46-V47,V50-V51,V56-V57, V60-V61,V66-V67,V70-V71,V76-V77)												
Asian or Pacific Islander 2/, both sexes												
2004.....	2	1	2	3	3	1	1	-	-	-	-	-
2003.....	6	3	2	1	3	-	1	2	-	-	-	-
2002.....	7	1	1	3	2	-	2	1	-	-	-	-
2001.....	6	1	1	2	4	-	-	-	-	-	-	-
2000.....	6	4	3	3	-	5	1	1	-	-	-	-
1999.....	3	4	3	3	2	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	-	1	1	-	3	-	1	-	-	-	-	-
2003.....	2	2	1	1	3	-	-	1	-	-	-	-
2002.....	3	1	-	3	2	-	1	1	-	-	-	-
2001.....	3	1	-	2	2	-	-	-	-	-	-	-
2000.....	2	4	1	2	-	3	-	1	-	-	-	-
1999.....	1	1	-	2	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	2	-	1	3	-	1	-	-	-	-	-	-
2003.....	4	1	1	-	-	-	1	1	-	-	-	-
2002.....	4	-	1	-	-	-	1	-	-	-	-	-
2001.....	3	-	1	-	2	-	-	-	-	-	-	-
2000.....	4	-	2	1	-	2	1	-	-	-	-	-
1999.....	2	3	3	1	1	-	-	1	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Occupant of car, pickup truck or van involved in collision with other motor vehicle (V42-V44,V49,V52-V54,V59)												
All races, both sexes												
2004.....	615	565	450	482	438	560	514	328	117	19	1	1
2003.....	643	521	450	401	495	575	543	349	120	11	1	1
2002.....	643	503	512	405	537	619	553	323	115	12	3	8
2001.....	569	496	401	437	502	579	471	306	121	17	1	2
2000.....	548	486	433	398	523	588	520	316	98	14	1	2
1999.....	539	498	400	434	543	568	508	283	81	18	1	6
All races, male												
2004.....	363	324	242	259	226	286	285	168	62	9	1	1
2003.....	388	303	255	214	240	302	276	177	60	3	-	1
2002.....	388	324	273	219	280	328	292	167	59	8	1	7
2001.....	332	281	221	241	266	295	250	169	60	6	-	2
2000.....	335	289	231	202	281	288	276	158	59	6	-	2
1999.....	319	265	224	228	285	291	269	152	51	9	1	6
All races, female												
2004.....	252	241	208	223	212	274	229	160	55	10	-	-
2003.....	255	218	195	187	255	273	267	172	60	8	1	-
2002.....	255	179	239	186	257	291	261	156	56	4	2	1
2001.....	237	215	180	196	236	284	221	137	61	11	1	-
2000.....	213	197	202	196	242	300	244	158	39	8	1	-
1999.....	220	233	176	206	258	277	239	131	30	9	-	-
White, both sexes												
2004.....	506	472	400	425	396	506	477	313	110	18	1	1
2003.....	532	423	378	337	440	523	512	328	114	10	-	1
2002.....	537	430	439	353	488	573	523	308	111	11	1	8
2001.....	478	440	336	379	442	541	444	293	115	15	1	2
2000.....	450	419	367	356	472	547	490	300	95	12	1	2
1999.....	452	429	341	380	486	528	475	265	72	16	1	4

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Occupant of car, pickup truck or van involved in collision with other motor vehicle (V42-V44,V49,V52-V54,V59)												
White, male												
2004.....	305	270	207	232	202	257	259	160	56	9	1	1
2003.....	319	241	217	182	217	273	256	166	57	3	-	1
2002.....	326	277	234	189	250	299	273	161	55	8	1	7
2001.....	278	253	184	208	238	272	235	160	56	6	-	2
2000.....	275	247	183	180	256	264	260	151	57	6	-	2
1999.....	265	225	188	194	254	268	250	142	45	7	1	4
White, female												
2004.....	201	202	193	193	194	249	218	153	54	9	-	-
2003.....	213	182	161	155	223	250	256	162	57	7	-	-
2002.....	211	153	205	164	238	274	250	147	56	3	-	1
2001.....	200	187	152	171	204	269	209	133	59	9	1	-
2000.....	175	172	184	176	216	283	230	149	38	6	1	-
1999.....	187	204	153	186	232	260	225	123	27	9	-	-
Black, both sexes												
2004.....	73	73	35	44	32	37	30	13	6	1	-	-
2003.....	84	78	52	49	37	38	26	16	5	1	-	-
2002.....	74	49	49	32	36	32	23	12	2	1	2	-
2001.....	73	44	53	50	48	25	23	11	6	2	-	-
2000.....	83	47	51	31	41	27	24	13	3	-	-	-
1999.....	65	54	42	41	42	29	27	15	7	2	-	2
Black, male												
2004.....	39	40	25	23	18	20	23	7	5	-	-	-
2003.....	54	48	29	27	16	21	18	9	2	-	-	-
2002.....	47	34	28	18	21	20	14	6	2	-	-	-
2001.....	45	24	30	30	19	15	14	8	4	-	-	-
2000.....	50	34	39	19	19	16	12	6	2	-	-	-
1999.....	41	32	25	26	26	20	15	8	4	2	-	2

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Occupant of car, pickup truck or van involved in collision with other motor vehicle (V42-V44,V49,V52-V54,V59)												
Black, female												
2004.....	34	33	10	21	14	17	7	6	1	1	-	-
2003.....	30	30	23	22	21	17	8	7	3	1	-	-
2002.....	27	15	21	14	15	12	9	6	-	1	2	-
2001.....	28	20	23	20	29	10	9	3	2	2	-	-
2000.....	33	13	12	12	22	11	12	7	1	-	-	-
1999.....	24	22	17	15	16	9	12	7	3	-	-	-
American Indian 1/, both sexes												
2004.....	10	8	2	5	4	7	5	-	-	-	-	-
2003.....	6	6	8	6	6	4	1	4	-	-	-	-
2002.....	13	5	10	5	5	5	2	-	1	-	-	-
2001.....	4	6	2	4	2	6	2	1	-	-	-	-
2000.....	5	10	5	4	5	3	2	2	-	1	-	-
1999.....	6	5	8	6	4	1	2	-	-	-	-	-
American Indian 1/, male												
2004.....	7	7	2	2	2	4	3	-	-	-	-	-
2003.....	4	5	3	2	2	1	-	1	-	-	-	-
2002.....	8	3	4	3	3	4	1	-	1	-	-	-
2001.....	2	2	1	1	1	4	1	-	-	-	-	-
2000.....	5	3	2	1	3	2	2	1	-	-	-	-
1999.....	5	3	6	3	1	-	1	-	-	-	-	-
American Indian 1/, female												
2004.....	3	1	-	3	2	3	2	-	-	-	-	-
2003.....	2	1	5	4	4	3	1	3	-	-	-	-
2002.....	5	2	6	2	2	1	1	-	-	-	-	-
2001.....	2	4	1	3	1	2	1	1	-	-	-	-
2000.....	-	7	3	3	2	1	-	1	-	1	-	-
1999.....	1	2	2	3	3	1	1	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Occupant of car, pickup truck or van involved in collision with other motor vehicle (V42-V44,V49,V52-V54,V59)												
Asian or Pacific Islander 2/, both sexes												
2004.....	26	12	13	8	6	10	2	2	1	-	-	-
2003.....	21	14	12	9	12	10	4	1	1	-	1	-
2002.....	19	19	14	15	8	9	5	3	1	-	-	-
2001.....	14	6	10	4	10	7	2	1	-	-	-	-
2000.....	10	10	10	7	5	11	4	1	-	1	-	-
1999.....	16	10	9	7	11	10	4	3	2	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	12	7	8	2	4	5	-	1	1	-	-	-
2003.....	11	9	6	3	5	7	2	1	1	-	-	-
2002.....	7	10	7	9	6	5	4	-	1	-	-	-
2001.....	7	2	6	2	8	4	-	1	-	-	-	-
2000.....	5	5	7	2	3	6	2	-	-	-	-	-
1999.....	8	5	5	5	4	3	3	2	2	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	14	5	5	6	2	5	2	1	-	-	-	-
2003.....	10	5	6	6	7	3	2	-	-	-	1	-
2002.....	12	9	7	6	2	4	1	3	-	-	-	-
2001.....	7	4	4	2	2	3	2	-	-	-	-	-
2000.....	5	5	3	5	2	5	2	1	-	1	-	-
1999.....	8	5	4	2	7	7	1	1	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Occupant of heavy transport vehicle or bus involved in collision with other motor vehicle (V62-V64,V69,V72-V74,V79)												
All races, both sexes												
2004.....	38	21	17	6	6	4	2	1	-	-	-	-
2003.....	25	25	17	11	9	7	5	1	-	-	-	-
2002.....	25	15	11	6	8	4	3	5	3	-	-	-
2001.....	18	8	16	7	5	2	1	1	-	-	-	-
2000.....	21	11	6	5	5	5	-	-	-	-	-	-
1999.....	19	19	10	5	11	8	11	1	3	-	-	-
All races, male												
2004.....	35	18	16	4	5	2	1	-	-	-	-	-
2003.....	21	25	14	8	2	4	2	-	-	-	-	-
2002.....	24	14	10	5	5	3	3	3	1	-	-	-
2001.....	16	7	15	7	4	-	-	-	-	-	-	-
2000.....	17	9	5	5	5	5	-	-	-	-	-	-
1999.....	18	17	7	2	6	3	5	-	-	-	-	-
All races, female												
2004.....	3	3	1	2	1	2	1	1	-	-	-	-
2003.....	4	-	3	3	7	3	3	1	-	-	-	-
2002.....	1	1	1	1	3	1	-	2	2	-	-	-
2001.....	2	1	1	-	1	2	1	1	-	-	-	-
2000.....	4	2	1	-	-	-	-	-	-	-	-	-
1999.....	1	2	3	3	5	5	6	1	3	-	-	-
White, both sexes												
2004.....	32	17	14	5	6	3	2	-	-	-	-	-
2003.....	20	22	17	10	9	5	5	1	-	-	-	-
2002.....	21	11	9	5	6	4	2	3	3	-	-	-
2001.....	18	8	14	5	5	2	1	1	-	-	-	-
2000.....	18	9	5	5	5	4	-	-	-	-	-	-
1999.....	16	15	9	4	11	8	9	1	2	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Occupant of heavy transport vehicle or bus involved in collision with other motor vehicle (V62-V64,V69,V72-V74,V79)

Asian or Pacific Islander 2/, both sexes

2004.....	-	-	1	-	-	-	-	-	-	-	-	-
2003.....	2	-	-	1	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	1	-	-	-	-	-

Asian or Pacific Islander 2/, male

2004.....	-	-	1	-	-	-	-	-	-	-	-	-
2003.....	1	-	-	1	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-

Asian or Pacific Islander 2/, female

2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	1	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	1	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Occupant of motor vehicle involved in noncollision accident (V48,V58,V68,V78)												
All races, both sexes												
2004.....	230	177	136	105	73	69	42	26	17	5	-	3
2003.....	247	191	132	101	87	78	53	43	14	4	-	1
2002.....	221	159	149	90	90	75	53	30	7	3	-	6
2001.....	227	144	130	88	86	80	51	27	4	1	-	2
2000.....	188	123	103	80	68	67	45	28	11	3	-	3
1999.....	167	142	112	101	70	60	37	31	6	2	1	1
All races, male												
2004.....	173	136	92	71	49	42	23	18	7	1	-	2
2003.....	166	123	89	66	58	48	34	19	8	2	-	1
2002.....	156	113	96	60	59	45	32	18	6	1	-	6
2001.....	162	102	89	59	58	47	27	15	1	1	-	1
2000.....	142	84	71	48	52	43	23	18	7	2	-	3
1999.....	127	103	74	70	35	39	18	21	2	1	-	1
All races, female												
2004.....	57	41	44	34	24	27	19	8	10	4	-	1
2003.....	81	68	43	35	29	30	19	24	6	2	-	-
2002.....	65	46	53	30	31	30	21	12	1	2	-	-
2001.....	65	42	41	29	28	33	24	12	3	-	-	1
2000.....	46	39	32	32	16	24	22	10	4	1	-	-
1999.....	40	39	38	31	35	21	19	10	4	1	1	-
White, both sexes												
2004.....	190	160	118	91	57	57	39	26	17	5	-	3
2003.....	203	160	107	85	77	64	48	39	13	4	-	1
2002.....	182	138	128	82	79	69	48	25	7	3	-	5
2001.....	200	112	108	79	80	76	48	23	3	1	-	2
2000.....	152	94	87	73	59	59	41	23	11	3	-	3
1999.....	138	119	93	89	61	54	34	29	5	1	1	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Occupant of motor vehicle involved in noncollision accident (V48,V58,V68,V78)												
White, male												
2004.....	140	124	80	62	37	34	22	18	7	1	-	2
2003.....	137	101	70	55	54	41	32	18	8	2	-	1
2002.....	122	97	83	55	51	44	28	14	6	1	-	5
2001.....	142	80	70	53	56	44	26	12	1	1	-	1
2000.....	115	63	61	44	47	39	21	15	7	2	-	3
1999.....	103	88	63	61	29	36	17	19	2	-	-	1
White, female												
2004.....	50	36	38	29	20	23	17	8	10	4	-	1
2003.....	66	59	37	30	23	23	16	21	5	2	-	-
2002.....	60	41	45	27	28	25	20	11	1	2	-	-
2001.....	58	32	38	26	24	32	22	11	2	-	-	1
2000.....	37	31	26	29	12	20	20	8	4	1	-	-
1999.....	35	31	30	28	32	18	17	10	3	1	1	-
Black, both sexes												
2004.....	31	13	15	13	10	7	2	-	-	-	-	-
2003.....	29	24	16	11	7	11	2	2	1	-	-	-
2002.....	24	15	14	5	7	4	4	5	-	-	-	-
2001.....	18	26	18	6	4	2	1	2	1	-	-	-
2000.....	25	26	13	6	5	4	3	3	-	-	-	-
1999.....	21	15	9	7	5	5	2	2	-	1	-	-
Black, male												
2004.....	26	9	9	9	8	7	1	-	-	-	-	-
2003.....	22	17	13	7	4	5	2	1	-	-	-	-
2002.....	22	13	9	3	6	1	3	4	-	-	-	-
2001.....	15	18	17	5	1	2	1	1	-	-	-	-
2000.....	20	19	8	4	3	3	2	2	-	-	-	-
1999.....	18	11	5	6	4	3	1	2	-	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Occupant of motor vehicle involved in noncollision accident (V48,V58,V68,V78)												
Black, female												
2004.....	5	4	6	4	2	-	1	-	-	-	-	-
2003.....	7	7	3	4	3	6	-	1	1	-	-	-
2002.....	2	2	5	2	1	3	1	1	-	-	-	-
2001.....	3	8	1	1	3	-	-	1	1	-	-	-
2000.....	5	7	5	2	2	1	1	1	-	-	-	-
1999.....	3	4	4	1	1	2	1	-	-	-	-	-
American Indian 1/, both sexes												
2004.....	6	2	1	-	4	3	-	-	-	-	-	-
2003.....	10	3	4	1	1	-	2	2	-	-	-	-
2002.....	9	5	4	2	1	1	1	-	-	-	-	-
2001.....	5	4	2	3	1	2	-	1	-	-	-	-
2000.....	7	2	2	-	1	1	1	-	-	-	-	-
1999.....	5	3	2	4	1	1	-	-	-	-	-	-
American Indian 1/, male												
2004.....	6	2	1	-	3	1	-	-	-	-	-	-
2003.....	6	2	2	1	-	-	-	-	-	-	-	-
2002.....	9	3	3	1	1	-	1	-	-	-	-	-
2001.....	3	4	1	1	-	1	-	1	-	-	-	-
2000.....	6	2	1	-	1	-	-	-	-	-	-	-
1999.....	5	3	2	3	1	-	-	-	-	-	-	-
American Indian 1/, female												
2004.....	-	-	-	-	1	2	-	-	-	-	-	-
2003.....	4	1	2	-	1	-	2	2	-	-	-	-
2002.....	-	2	1	1	-	1	-	-	-	-	-	-
2001.....	2	-	1	2	1	1	-	-	-	-	-	-
2000.....	1	-	1	-	-	1	1	-	-	-	-	-
1999.....	-	-	-	1	-	1	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Occupant of motor vehicle involved in noncollision accident (V48,V58,V68,V78)												
Asian or Pacific Islander 2/, both sexes												
2004.....	3	2	2	1	2	2	1	-	-	-	-	-
2003.....	5	4	5	4	2	3	1	-	-	-	-	-
2002.....	6	1	3	1	3	1	-	-	-	-	-	1
2001.....	4	2	2	-	1	-	2	1	-	-	-	-
2000.....	4	1	1	1	3	3	-	2	-	-	-	-
1999.....	3	5	8	1	3	-	1	-	1	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	1	1	2	-	1	-	-	-	-	-	-	-
2003.....	1	3	4	3	-	2	-	-	-	-	-	-
2002.....	3	-	1	1	1	-	-	-	-	-	-	1
2001.....	2	-	1	-	1	-	-	1	-	-	-	-
2000.....	1	-	1	-	1	1	-	1	-	-	-	-
1999.....	1	1	4	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	2	1	-	1	1	2	1	-	-	-	-	-
2003.....	4	1	1	1	2	1	1	-	-	-	-	-
2002.....	3	1	2	-	2	1	-	-	-	-	-	-
2001.....	2	2	1	-	-	-	2	-	-	-	-	-
2000.....	3	1	-	1	2	2	-	1	-	-	-	-
1999.....	2	4	4	1	2	-	1	-	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Occupant of special-use motor vehicle involved in any accident (V83-V86)												
All races, both sexes												
2004.....	56	45	43	29	27	26	11	14	3	1	-	-
2003.....	69	61	35	40	25	35	20	13	3	1	-	-
2002.....	44	38	38	31	40	32	26	9	2	-	1	-
2001.....	47	50	40	41	36	48	28	7	2	-	-	-
2000.....	47	58	56	48	34	34	20	12	4	-	-	-
1999.....	62	46	50	61	49	42	27	24	4	-	-	-
All races, male												
2004.....	49	43	40	28	26	25	11	12	2	1	-	-
2003.....	56	55	33	38	24	33	18	12	2	1	-	-
2002.....	38	35	35	29	34	27	25	7	1	-	1	-
2001.....	44	47	40	40	32	45	26	6	2	-	-	-
2000.....	42	54	54	45	32	34	20	11	4	-	-	-
1999.....	56	43	47	59	48	42	24	23	3	-	-	-
All races, female												
2004.....	7	2	3	1	1	1	-	2	1	-	-	-
2003.....	13	6	2	2	1	2	2	1	1	-	-	-
2002.....	6	3	3	2	6	5	1	2	1	-	-	-
2001.....	3	3	-	1	4	3	2	1	-	-	-	-
2000.....	5	4	2	3	2	-	-	1	-	-	-	-
1999.....	6	3	3	2	1	-	3	1	1	-	-	-
White, both sexes												
2004.....	49	41	38	26	25	24	9	14	3	1	-	-
2003.....	62	55	33	39	25	35	19	12	3	1	-	-
2002.....	42	37	37	27	40	32	26	9	2	-	1	-
2001.....	41	48	39	41	34	46	27	7	2	-	-	-
2000.....	44	57	48	47	31	32	20	12	4	-	-	-
1999.....	59	43	45	56	48	41	27	23	4	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Occupant of special-use motor vehicle involved in any accident (V83-V86)												
White, male												
2004.....	43	39	35	25	24	23	9	12	2	1	-	-
2003.....	49	49	32	38	24	33	17	11	2	1	-	-
2002.....	36	34	34	26	34	27	25	7	1	-	1	-
2001.....	38	45	39	40	30	44	25	6	2	-	-	-
2000.....	39	53	46	44	29	32	20	11	4	-	-	-
1999.....	53	40	42	54	47	41	24	22	3	-	-	-
White, female												
2004.....	6	2	3	1	1	1	-	2	1	-	-	-
2003.....	13	6	1	1	1	2	2	1	1	-	-	-
2002.....	6	3	3	1	6	5	1	2	1	-	-	-
2001.....	3	3	-	1	4	2	2	1	-	-	-	-
2000.....	5	4	2	3	2	-	-	1	-	-	-	-
1999.....	6	3	3	2	1	-	3	1	1	-	-	-
Black, both sexes												
2004.....	5	1	4	2	-	2	1	-	-	-	-	-
2003.....	3	5	1	-	-	-	1	1	-	-	-	-
2002.....	1	-	1	2	-	-	-	-	-	-	-	-
2001.....	4	1	1	-	2	1	-	-	-	-	-	-
2000.....	3	1	5	1	3	-	-	-	-	-	-	-
1999.....	2	-	4	4	1	-	-	1	-	-	-	-
Black, male												
2004.....	5	1	4	2	-	2	1	-	-	-	-	-
2003.....	3	5	1	-	-	-	1	1	-	-	-	-
2002.....	1	-	1	1	-	-	-	-	-	-	-	-
2001.....	4	1	1	-	2	-	-	-	-	-	-	-
2000.....	3	1	5	1	3	-	-	-	-	-	-	-
1999.....	2	-	4	4	1	-	-	1	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other and unspecified motor vehicle accidents (V09.0,V09.2,V80.3-V80.5,V82.0-V82.1,V87.0-V87.5,V87.7-V87.8,V88.0-V88.5,V88.7-V88.8,V89.0,V89.2)

All races, both sexes

2004.....	959	783	654	570	559	702	643	376	130	20	1	16
2003.....	900	749	651	550	561	656	623	392	142	26	-	9
2002.....	929	690	599	565	633	711	606	369	137	16	2	9
2001.....	946	756	581	552	632	775	640	370	119	25	-	11
2000.....	974	750	626	593	644	754	570	426	107	20	2	13
1999.....	867	696	560	650	722	791	657	410	149	19	1	13

All races, male

2004.....	649	519	411	341	321	391	366	191	69	10	-	13
2003.....	587	485	403	338	305	374	331	208	72	14	-	7
2002.....	600	474	361	354	368	398	308	219	75	8	-	9
2001.....	641	468	367	307	366	438	351	199	66	11	-	9
2000.....	647	481	388	349	355	413	304	233	58	11	1	13
1999.....	554	464	340	385	387	452	353	232	84	12	-	12

All races, female

2004.....	310	264	243	229	238	311	277	185	61	10	1	3
2003.....	313	264	248	212	256	282	292	184	70	12	-	2
2002.....	329	216	238	211	265	313	298	150	62	8	2	-
2001.....	305	288	214	245	266	337	289	171	53	14	-	2
2000.....	327	269	238	244	289	341	266	193	49	9	1	-
1999.....	313	232	220	265	335	339	304	178	65	7	1	1

White, both sexes

2004.....	745	651	558	479	483	625	597	363	119	19	1	14
2003.....	720	600	539	475	492	590	564	364	126	24	-	9
2002.....	762	577	508	483	564	646	556	346	127	15	1	9
2001.....	793	624	489	469	559	702	603	349	112	24	-	10
2000.....	795	635	542	501	556	688	531	400	96	19	2	11
1999.....	711	586	462	565	643	714	604	380	141	17	1	12

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other and unspecified motor vehicle accidents (V09.0,V09.2,V80.3-V80.5,V82.0-V82.1,V87.0-V87.5,V87.7-V87.8,V88.0-V88.5,V88.7-V88.8,V89.0,V89.2)

White, male

2004.....	497	431	343	281	281	355	337	186	63	10	-	12
2003.....	469	392	340	295	273	338	302	194	61	12	-	7
2002.....	490	391	305	300	334	355	281	200	68	8	-	9
2001.....	545	382	308	265	321	393	329	188	63	11	-	9
2000.....	527	402	332	291	303	379	280	224	51	10	1	11
1999.....	458	392	279	337	351	401	323	215	80	10	-	11

White, female

2004.....	248	220	215	198	202	270	260	177	56	9	1	2
2003.....	251	208	199	180	219	252	262	170	65	12	-	2
2002.....	272	186	203	183	230	291	275	146	59	7	1	-
2001.....	248	242	181	204	238	309	274	161	49	13	-	1
2000.....	268	233	210	210	253	309	251	176	45	9	1	-
1999.....	253	194	183	228	292	313	281	165	61	7	1	1

Black, both sexes

2004.....	170	92	74	62	46	50	31	8	8	1	-	-
2003.....	133	97	83	57	51	40	39	23	13	1	-	-
2002.....	121	85	64	60	45	49	37	16	7	-	1	-
2001.....	113	102	68	64	51	60	26	17	5	1	-	1
2000.....	136	80	59	67	65	41	30	19	8	1	-	1
1999.....	119	78	74	65	61	63	46	22	8	2	-	1

Black, male

2004.....	123	61	51	46	30	26	21	4	5	-	-	-
2003.....	92	68	51	36	26	26	17	11	8	1	-	-
2002.....	79	64	41	41	27	35	19	15	5	-	-	-
2001.....	76	71	49	33	30	39	15	9	3	-	-	-
2000.....	96	61	40	45	43	24	20	7	6	1	-	1
1999.....	82	53	50	39	26	42	28	14	4	2	-	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified motor vehicle accidents (V09.0,V09.2,V80.3-V80.5,V82.0-V82.1,V87.0-V87.5,V87.7-V87.8,V88.0-V88.5,V88.7-V88.8,V89.0,V89.2)												
Black, female												
2004.....	47	31	23	16	16	24	10	4	3	1	-	-
2003.....	41	29	32	21	25	14	22	12	5	-	-	-
2002.....	42	21	23	19	18	14	18	1	2	-	1	-
2001.....	37	31	19	31	21	21	11	8	2	1	-	1
2000.....	40	19	19	22	22	17	10	12	2	-	-	-
1999.....	37	25	24	26	35	21	18	8	4	-	-	-
American Indian 1/, both sexes												
2004.....	16	11	8	9	14	5	3	1	1	-	-	2
2003.....	20	21	9	7	5	4	4	1	1	-	-	-
2002.....	25	13	13	7	5	3	3	1	-	1	-	-
2001.....	17	13	5	7	3	5	5	2	1	-	-	-
2000.....	18	8	11	9	5	6	1	1	-	-	-	1
1999.....	11	12	5	6	4	7	3	-	-	-	-	-
American Indian 1/, male												
2004.....	13	10	7	3	7	2	-	-	1	-	-	1
2003.....	11	10	5	3	1	2	2	1	1	-	-	-
2002.....	18	11	6	5	-	2	3	1	-	-	-	-
2001.....	11	7	2	4	2	2	3	1	-	-	-	-
2000.....	9	2	6	4	3	3	1	1	-	-	-	1
1999.....	6	8	2	2	3	5	1	-	-	-	-	-
American Indian 1/, female												
2004.....	3	1	1	6	7	3	3	1	-	-	-	1
2003.....	9	11	4	4	4	2	2	-	-	-	-	-
2002.....	7	2	7	2	5	1	-	-	-	1	-	-
2001.....	6	6	3	3	1	3	2	1	1	-	-	-
2000.....	9	6	5	5	2	3	-	-	-	-	-	-
1999.....	5	4	3	4	1	2	2	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified motor vehicle accidents (V09.0,V09.2,V80.3-V80.5,V82.0-V82.1,V87.0-V87.5,V87.7-V87.8,V88.0-V88.5,V88.7-V88.8,V89.0,V89.2)												
Asian or Pacific Islander 2/, both sexes												
2004.....	28	29	14	20	16	22	12	4	2	-	-	-
2003.....	27	31	20	11	13	22	16	4	2	1	-	-
2002.....	21	15	14	15	19	13	10	6	3	-	-	-
2001.....	23	17	19	12	19	8	6	2	1	-	-	-
2000.....	25	27	14	16	18	19	8	6	3	-	-	-
1999.....	26	20	19	14	14	7	4	8	-	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	16	17	10	11	3	8	8	1	-	-	-	-
2003.....	15	15	7	4	5	8	10	2	2	1	-	-
2002.....	13	8	9	8	7	6	5	3	2	-	-	-
2001.....	9	8	8	5	13	4	4	1	-	-	-	-
2000.....	15	16	10	9	6	7	3	1	1	-	-	-
1999.....	8	11	9	7	7	4	1	3	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	12	12	4	9	13	14	4	3	2	-	-	-
2003.....	12	16	13	7	8	14	6	2	-	-	-	-
2002.....	8	7	5	7	12	7	5	3	1	-	-	-
2001.....	14	9	11	7	6	4	2	1	1	-	-	-
2000.....	10	11	4	7	12	12	5	5	2	-	-	-
1999.....	18	9	10	7	7	3	3	5	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified land transport accidents (V01,V06,V09.1,V09.3,V09.9,V10-V11,V16-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.7-V80.9,V87.9,V88.9,V89.1,V89.3,V89.9)												
All races, both sexes												
2004.....	57	52	43	34	37	19	27	14	3	1	-	1
2003.....	64	61	50	34	34	34	32	17	5	1	1	3
2002.....	52	37	35	23	31	30	25	8	3	-	1	1
2001.....	66	46	48	57	41	48	31	17	7	1	-	2
2000.....	76	48	43	44	41	45	34	18	10	1	-	1
1999.....	98	85	66	73	58	85	61	32	15	3	1	1
All races, male												
2004.....	46	37	32	22	23	12	20	8	1	1	-	1
2003.....	51	46	37	27	25	21	22	11	4	1	-	3
2002.....	44	30	26	20	25	23	21	4	1	-	-	1
2001.....	51	37	35	44	26	28	15	6	4	-	-	2
2000.....	46	33	37	31	31	28	20	12	4	-	-	1
1999.....	78	62	47	50	38	47	33	18	12	1	1	1
All races, female												
2004.....	11	15	11	12	14	7	7	6	2	-	-	-
2003.....	13	15	13	7	9	13	10	6	1	-	1	-
2002.....	8	7	9	3	6	7	4	4	2	-	1	-
2001.....	15	9	13	13	15	20	16	11	3	1	-	-
2000.....	30	15	6	13	10	17	14	6	6	1	-	-
1999.....	20	23	19	23	20	38	28	14	3	2	-	-
White, both sexes												
2004.....	45	44	38	31	30	14	27	12	3	1	-	1
2003.....	48	52	42	28	29	31	29	16	4	1	1	3
2002.....	47	30	28	18	27	23	24	8	3	-	1	1
2001.....	54	38	39	42	39	42	28	14	5	1	-	-
2000.....	70	39	37	37	35	38	32	17	10	-	-	1
1999.....	83	78	59	61	50	73	55	30	13	3	-	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other and unspecified land transport accidents (V01,V06,V09.1,V09.3,V09.9,V10-V11,V16-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.7-V80.9,V87.9,V88.9,V89.1,V89.3,V89.9)

White, male

2004.....	35	31	28	20	19	10	20	7	1	1	-	1
2003.....	38	39	30	22	21	19	20	10	3	1	-	3
2002.....	39	24	22	16	21	16	21	4	1	-	-	1
2001.....	44	30	32	32	24	26	14	4	3	-	-	-
2000.....	43	26	34	25	26	25	19	12	4	-	-	1
1999.....	65	58	41	45	32	42	30	17	10	1	-	1

White, female

2004.....	10	13	10	11	11	4	7	5	2	-	-	-
2003.....	10	13	12	6	8	12	9	6	1	-	1	-
2002.....	8	6	6	2	6	7	3	4	2	-	1	-
2001.....	10	8	7	10	15	16	14	10	2	1	-	-
2000.....	27	13	3	12	9	13	13	5	6	-	-	-
1999.....	18	20	18	16	18	31	25	13	3	2	-	-

Black, both sexes

2004.....	6	8	4	2	3	5	-	-	-	-	-	-
2003.....	13	9	7	4	3	2	1	-	1	-	-	-
2002.....	5	5	3	3	3	5	1	-	-	-	-	-
2001.....	9	6	7	11	1	4	1	-	1	-	-	2
2000.....	2	7	3	5	3	4	1	1	-	1	-	-
1999.....	12	7	4	7	5	8	3	2	1	-	1	-

Black, male

2004.....	5	6	3	2	3	2	-	-	-	-	-	-
2003.....	12	7	7	4	3	1	-	-	1	-	-	-
2002.....	5	4	3	3	3	5	-	-	-	-	-	-
2001.....	5	5	3	9	1	2	-	-	-	-	-	2
2000.....	1	5	2	5	2	2	-	-	-	-	-	-
1999.....	11	4	3	3	4	2	1	1	1	-	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified land transport accidents (V01,V06,V09.1,V09.3,V09.9,V10-V11,V16-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.7-V80.9,V87.9,V88.9,V89.1,V89.3,V89.9)												
Asian or Pacific Islander 2/, both sexes												
2004.....	3	-	-	1	4	-	-	2	-	-	-	-
2003.....	2	-	1	2	1	1	2	-	-	-	-	-
2002.....	-	2	4	1	-	2	-	-	-	-	-	-
2001.....	2	-	2	3	1	2	1	3	1	-	-	-
2000.....	3	2	2	1	3	3	1	-	-	-	-	-
1999.....	2	-	2	2	2	4	3	-	1	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	3	-	-	-	1	-	-	1	-	-	-	-
2003.....	1	-	-	1	-	1	2	-	-	-	-	-
2002.....	-	2	1	-	-	2	-	-	-	-	-	-
2001.....	2	-	-	2	1	-	-	2	1	-	-	-
2000.....	1	2	1	1	3	1	1	-	-	-	-	-
1999.....	2	-	2	-	1	3	2	-	1	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	-	-	-	1	3	-	-	1	-	-	-	-
2003.....	1	-	1	1	1	-	-	-	-	-	-	-
2002.....	-	-	3	1	-	-	-	-	-	-	-	-
2001.....	-	-	2	1	-	2	1	1	-	-	-	-
2000.....	2	-	1	-	-	2	-	-	-	-	-	-
1999.....	-	-	-	2	1	1	1	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Water transport accidents (V90-V94)												
All races, both sexes												
2004.....	63	40	25	25	14	10	10	6	1	-	-	-
2003.....	60	27	36	19	11	19	3	2	-	-	-	-
2002.....	51	39	27	34	24	13	10	1	-	-	-	-
2001.....	52	39	31	22	23	6	8	2	1	-	-	-
2000.....	54	34	24	30	20	7	7	1	-	-	-	-
1999.....	47	38	32	28	16	5	8	1	-	-	-	-
All races, male												
2004.....	51	35	24	23	14	10	10	6	1	-	-	-
2003.....	54	24	35	18	11	19	3	2	-	-	-	-
2002.....	47	37	25	33	24	11	10	1	-	-	-	-
2001.....	48	33	29	22	22	6	8	2	1	-	-	-
2000.....	50	32	22	26	19	7	5	-	-	-	-	-
1999.....	44	36	30	28	15	3	7	-	-	-	-	-
All races, female												
2004.....	12	5	1	2	-	-	-	-	-	-	-	-
2003.....	6	3	1	1	-	-	-	-	-	-	-	-
2002.....	4	2	2	1	-	2	-	-	-	-	-	-
2001.....	4	6	2	-	1	-	-	-	-	-	-	-
2000.....	4	2	2	4	1	-	2	1	-	-	-	-
1999.....	3	2	2	-	1	2	1	1	-	-	-	-
White, both sexes												
2004.....	53	36	20	24	12	9	10	6	1	-	-	-
2003.....	51	23	32	13	10	17	2	2	-	-	-	-
2002.....	39	29	26	28	20	13	8	1	-	-	-	-
2001.....	45	30	27	16	20	6	8	2	1	-	-	-
2000.....	47	31	20	25	20	6	7	1	-	-	-	-
1999.....	37	31	25	23	13	4	7	1	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Water transport accidents (V90-V94)												
White, male												
2004.....	44	32	19	22	12	9	10	6	1	-	-	-
2003.....	45	20	32	12	10	17	2	2	-	-	-	-
2002.....	35	27	24	27	20	11	8	1	-	-	-	-
2001.....	41	24	25	16	19	6	8	2	1	-	-	-
2000.....	43	29	19	22	19	6	5	-	-	-	-	-
1999.....	35	30	23	23	12	3	6	-	-	-	-	-
White, female												
2004.....	9	4	1	2	-	-	-	-	-	-	-	-
2003.....	6	3	-	1	-	-	-	-	-	-	-	-
2002.....	4	2	2	1	-	2	-	-	-	-	-	-
2001.....	4	6	2	-	1	-	-	-	-	-	-	-
2000.....	4	2	1	3	1	-	2	1	-	-	-	-
1999.....	2	1	2	-	1	1	1	1	-	-	-	-
Black, both sexes												
2004.....	8	4	5	-	1	1	-	-	-	-	-	-
2003.....	6	3	3	5	-	2	1	-	-	-	-	-
2002.....	9	10	1	4	3	-	1	-	-	-	-	-
2001.....	5	9	4	5	1	-	-	-	-	-	-	-
2000.....	5	2	3	4	-	1	-	-	-	-	-	-
1999.....	6	3	7	4	2	-	1	-	-	-	-	-
Black, male												
2004.....	7	3	5	-	1	1	-	-	-	-	-	-
2003.....	6	3	3	5	-	2	1	-	-	-	-	-
2002.....	9	10	1	4	3	-	1	-	-	-	-	-
2001.....	5	9	4	5	1	-	-	-	-	-	-	-
2000.....	5	2	3	3	-	1	-	-	-	-	-	-
1999.....	6	3	7	4	2	-	1	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Water transport accidents (V90-V94)												
Asian or Pacific Islander 2/, both sexes												
2004.....	1	-	-	1	-	-	-	-	-	-	-	-
2003.....	2	1	-	-	-	-	-	-	-	-	-	-
2002.....	3	-	-	1	-	-	1	-	-	-	-	-
2001.....	1	-	-	-	1	-	-	-	-	-	-	-
2000.....	1	-	1	1	-	-	-	-	-	-	-	-
1999.....	1	1	-	1	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	-	-	-	1	-	-	-	-	-	-	-	-
2003.....	2	1	-	-	-	-	-	-	-	-	-	-
2002.....	3	-	-	1	-	-	1	-	-	-	-	-
2001.....	1	-	-	-	1	-	-	-	-	-	-	-
2000.....	1	-	-	1	-	-	-	-	-	-	-	-
1999.....	1	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	1	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	1	-	-	-	-	-	-	-	-	-
1999.....	-	1	-	-	-	1	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Air and space transport accidents (V95-V97)												
All races, both sexes												
2004.....	98	52	72	41	26	12	4	1	-	-	-	-
2003.....	90	95	69	39	28	20	7	3	-	-	-	-
2002.....	77	75	60	36	31	17	4	2	-	-	-	-
2001.....	100	72	56	50	33	21	8	3	-	-	-	-
2000.....	87	74	61	48	21	19	8	1	1	-	-	-
1999.....	74	70	50	36	36	14	6	2	1	-	-	-
All races, male												
2004.....	90	43	63	37	24	12	2	1	-	-	-	-
2003.....	80	82	63	34	25	17	5	3	-	-	-	-
2002.....	62	65	50	34	30	14	4	1	-	-	-	-
2001.....	77	56	47	40	26	15	8	2	-	-	-	-
2000.....	76	64	47	41	16	17	7	1	1	-	-	-
1999.....	61	62	48	27	31	10	5	1	1	-	-	-
All races, female												
2004.....	8	9	9	4	2	-	2	-	-	-	-	-
2003.....	10	13	6	5	3	3	2	-	-	-	-	-
2002.....	15	10	10	2	1	3	-	1	-	-	-	-
2001.....	23	16	9	10	7	6	-	1	-	-	-	-
2000.....	11	10	14	7	5	2	1	-	-	-	-	-
1999.....	13	8	2	9	5	4	1	1	-	-	-	-
White, both sexes												
2004.....	96	52	70	40	26	12	4	1	-	-	-	-
2003.....	89	94	66	37	28	20	7	3	-	-	-	-
2002.....	75	75	59	35	31	17	4	2	-	-	-	-
2001.....	99	68	55	50	32	20	8	3	-	-	-	-
2000.....	82	73	60	44	20	18	8	1	1	-	-	-
1999.....	71	69	49	35	35	14	6	2	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified transport accidents and their sequelae (V98-V99,Y85)												
All races, both sexes												
2004.....	67	51	45	33	24	15	16	15	3	-	-	-
2003.....	63	66	43	34	31	21	17	10	2	1	-	-
2002.....	55	50	41	25	18	27	25	13	4	-	-	-
2001.....	62	35	30	22	25	34	20	10	2	2	-	-
2000.....	42	26	36	33	25	19	18	8	-	1	-	-
1999.....	53	37	33	19	23	19	14	7	3	1	-	-
All races, male												
2004.....	48	41	36	25	18	5	9	9	1	-	-	-
2003.....	49	51	36	24	22	14	8	7	1	-	-	-
2002.....	41	37	31	23	13	18	17	7	2	-	-	-
2001.....	49	26	27	16	19	20	11	4	1	-	-	-
2000.....	33	19	29	21	21	13	13	7	-	-	-	-
1999.....	41	29	21	15	18	13	6	6	1	-	-	-
All races, female												
2004.....	19	10	9	8	6	10	7	6	2	-	-	-
2003.....	14	15	7	10	9	7	9	3	1	1	-	-
2002.....	14	13	10	2	5	9	8	6	2	-	-	-
2001.....	13	9	3	6	6	14	9	6	1	2	-	-
2000.....	9	7	7	12	4	6	5	1	-	1	-	-
1999.....	12	8	12	4	5	6	8	1	2	1	-	-
White, both sexes												
2004.....	55	43	38	26	21	12	14	14	3	-	-	-
2003.....	52	57	36	28	30	15	16	10	2	1	-	-
2002.....	48	43	34	20	16	21	22	12	4	-	-	-
2001.....	47	26	28	19	24	28	18	8	2	1	-	-
2000.....	35	20	30	29	19	17	18	7	-	1	-	-
1999.....	43	25	23	17	20	16	10	5	3	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified transport accidents and their sequelae (V98-V99,Y85)												
White, male												
2004.....	38	33	31	20	16	4	8	8	1	-	-	-
2003.....	39	44	31	19	22	11	7	7	1	-	-	-
2002.....	35	30	25	18	11	12	15	6	2	-	-	-
2001.....	36	18	25	13	18	17	10	4	1	-	-	-
2000.....	28	13	24	18	17	11	13	6	-	-	-	-
1999.....	35	19	13	13	15	12	4	4	1	-	-	-
White, female												
2004.....	17	10	7	6	5	8	6	6	2	-	-	-
2003.....	13	13	5	9	8	4	9	3	1	1	-	-
2002.....	13	13	9	2	5	9	7	6	2	-	-	-
2001.....	11	8	3	6	6	11	8	4	1	1	-	-
2000.....	7	7	6	11	2	6	5	1	-	1	-	-
1999.....	8	6	10	4	5	4	6	1	2	1	-	-
Black, both sexes												
2004.....	10	7	6	5	3	3	1	-	-	-	-	-
2003.....	7	6	6	4	1	3	1	-	-	-	-	-
2002.....	6	5	6	3	1	5	2	1	-	-	-	-
2001.....	11	7	2	2	1	5	2	1	-	1	-	-
2000.....	3	5	3	4	6	2	-	1	-	-	-	-
1999.....	8	7	8	1	2	2	4	2	-	-	-	-
Black, male												
2004.....	9	7	5	4	2	1	1	-	-	-	-	-
2003.....	7	5	5	3	-	1	1	-	-	-	-	-
2002.....	5	5	5	3	1	5	2	1	-	-	-	-
2001.....	9	7	2	2	1	2	1	-	-	-	-	-
2000.....	3	5	2	3	4	2	-	1	-	-	-	-
1999.....	5	6	7	1	2	1	2	2	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified transport accidents and their sequelae (V98-V99,Y85)												
Asian or Pacific Islander 2/, both sexes												
2004.....	-	1	1	-	-	-	1	1	-	-	-	-
2003.....	3	1	1	2	-	3	-	-	-	-	-	-
2002.....	-	2	1	2	1	1	1	-	-	-	-	-
2001.....	1	1	-	1	-	1	-	1	-	-	-	-
2000.....	2	1	3	-	-	-	-	-	-	-	-	-
1999.....	2	2	1	1	1	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	-	1	-	-	-	-	-	1	-	-	-	-
2003.....	3	1	-	2	-	2	-	-	-	-	-	-
2002.....	-	2	1	2	1	1	-	-	-	-	-	-
2001.....	1	-	-	1	-	1	-	-	-	-	-	-
2000.....	-	1	3	-	-	-	-	-	-	-	-	-
1999.....	1	2	1	1	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	-	-	1	-	-	-	1	-	-	-	-	-
2003.....	-	-	1	-	-	1	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	1	-	-	-	-	-
2001.....	-	1	-	-	-	-	-	1	-	-	-	-
2000.....	2	-	-	-	-	-	-	-	-	-	-	-
1999.....	1	-	-	-	-	1	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Falls (W00-W19)												
All races, both sexes												
2004.....	617	669	724	885	1,370	2,246	3,436	3,447	2,433	934	148	4
2003.....	535	579	641	787	1,261	2,136	3,113	3,220	2,221	806	157	1
2002.....	530	515	574	759	1,208	2,011	2,869	3,032	2,045	784	129	1
2001.....	504	463	541	747	1,086	1,873	2,567	2,647	1,850	718	135	4
2000.....	462	448	501	595	1,065	1,641	2,200	2,352	1,682	618	120	-
1999.....	377	398	489	663	1,000	1,636	2,220	2,308	1,625	533	112	4
All races, male												
2004.....	465	499	488	575	819	1,189	1,669	1,465	891	239	35	3
2003.....	409	408	419	485	724	1,145	1,516	1,359	796	206	25	1
2002.....	394	379	392	474	728	1,079	1,365	1,262	691	231	24	-
2001.....	395	344	374	469	621	1,041	1,236	1,149	655	188	31	3
2000.....	358	326	354	378	613	895	1,052	992	592	173	27	-
1999.....	295	296	354	433	594	840	1,110	964	568	140	29	4
All races, female												
2004.....	152	170	236	310	551	1,057	1,767	1,982	1,542	695	113	1
2003.....	126	171	222	302	537	991	1,597	1,861	1,425	600	132	-
2002.....	136	136	182	285	480	932	1,504	1,770	1,354	553	105	1
2001.....	109	119	167	278	465	832	1,331	1,498	1,195	530	104	1
2000.....	104	122	147	217	452	746	1,148	1,360	1,090	445	93	-
1999.....	82	102	135	230	406	796	1,110	1,344	1,057	393	83	-
White, both sexes												
2004.....	516	566	624	786	1,236	2,101	3,263	3,309	2,338	897	141	4
2003.....	429	493	558	687	1,139	1,994	2,924	3,092	2,120	784	150	-
2002.....	448	437	507	669	1,096	1,874	2,739	2,910	1,942	754	122	1
2001.....	431	391	468	660	993	1,742	2,439	2,521	1,767	687	125	4
2000.....	377	374	428	514	966	1,539	2,088	2,262	1,613	584	111	-
1999.....	301	324	398	575	894	1,518	2,114	2,206	1,551	506	103	4

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Falls (W00-W19)												
White, male												
2004.....	387	418	410	523	736	1,120	1,582	1,396	845	227	33	3
2003.....	326	349	364	416	659	1,073	1,420	1,306	757	200	25	-
2002.....	329	321	345	416	663	1,004	1,297	1,205	646	220	22	-
2001.....	334	292	322	414	571	967	1,167	1,088	622	172	27	3
2000.....	289	269	305	326	563	833	997	951	559	159	25	-
1999.....	237	239	291	372	532	771	1,057	915	535	131	28	4
White, female												
2004.....	129	148	214	263	500	981	1,681	1,913	1,493	670	108	1
2003.....	103	144	194	271	480	921	1,504	1,786	1,363	584	125	-
2002.....	119	116	162	253	433	870	1,442	1,705	1,296	534	100	1
2001.....	97	99	146	246	422	775	1,272	1,433	1,145	515	98	1
2000.....	88	105	123	188	403	706	1,091	1,311	1,054	425	86	-
1999.....	64	85	107	203	362	747	1,057	1,291	1,016	375	75	-
Black, both sexes												
2004.....	78	85	77	77	98	99	103	71	65	23	6	-
2003.....	77	68	65	72	80	79	111	74	59	14	4	-
2002.....	68	57	54	57	81	86	69	72	67	22	5	-
2001.....	66	56	55	73	72	94	81	71	51	16	7	-
2000.....	67	60	56	64	73	65	72	50	46	21	7	-
1999.....	64	59	72	81	70	75	80	59	55	21	7	-
Black, male												
2004.....	61	67	63	40	64	52	53	29	34	7	2	-
2003.....	57	46	42	52	37	40	48	32	19	3	-	-
2002.....	54	42	37	37	46	48	37	34	29	8	1	-
2001.....	55	41	38	47	40	52	45	30	19	8	3	-
2000.....	54	47	42	41	37	39	31	20	23	9	1	-
1999.....	49	45	48	57	43	43	36	23	20	6	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Falls (W00-W19)												
Black, female												
2004.....	17	18	14	37	34	47	50	42	31	16	4	-
2003.....	20	22	23	20	43	39	63	42	40	11	4	-
2002.....	14	15	17	20	35	38	32	38	38	14	4	-
2001.....	11	15	17	26	32	42	36	41	32	8	4	-
2000.....	13	13	14	23	36	26	41	30	23	12	6	-
1999.....	15	14	24	24	27	32	44	36	35	15	6	-
American Indian 1/, both sexes												
2004.....	9	6	12	4	8	3	11	13	4	2	-	-
2003.....	15	8	8	8	8	11	10	9	8	2	1	1
2002.....	6	8	3	7	4	9	12	5	7	1	-	-
2001.....	2	6	6	3	5	8	4	10	7	4	-	-
2000.....	11	7	5	-	6	9	3	6	2	2	1	-
1999.....	5	3	9	1	9	5	3	8	3	3	1	-
American Indian 1/, male												
2004.....	8	6	8	2	3	2	3	10	1	-	-	-
2003.....	12	5	8	7	7	4	7	4	3	-	-	1
2002.....	3	7	2	4	2	3	4	3	-	-	-	-
2001.....	2	5	5	1	2	4	2	2	2	1	-	-
2000.....	10	6	1	-	2	8	2	5	1	1	-	-
1999.....	3	2	6	-	4	1	-	4	1	1	-	-
American Indian 1/, female												
2004.....	1	-	4	2	5	1	8	3	3	2	-	-
2003.....	3	3	-	1	1	7	3	5	5	2	1	-
2002.....	3	1	1	3	2	6	8	2	7	1	-	-
2001.....	-	1	1	2	3	4	2	8	5	3	-	-
2000.....	1	1	4	-	4	1	1	1	1	1	1	-
1999.....	2	1	3	1	5	4	3	4	2	2	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Falls (W00-W19)												
Asian or Pacific Islander 2/, both sexes												
2004.....	14	12	11	18	28	43	59	54	26	12	1	-
2003.....	14	10	10	20	34	52	68	45	34	6	2	-
2002.....	8	13	10	26	27	42	49	45	29	7	2	-
2001.....	5	10	12	11	16	29	43	45	25	11	3	-
2000.....	7	7	12	17	20	28	37	34	21	11	1	-
1999.....	7	12	10	6	27	38	23	35	16	3	1	-
Asian or Pacific Islander 2/, male												
2004.....	9	8	7	10	16	15	31	30	11	5	-	-
2003.....	14	8	5	10	21	28	41	17	17	3	-	-
2002.....	8	9	8	17	17	24	27	20	16	3	1	-
2001.....	4	6	9	7	8	18	22	29	12	7	1	-
2000.....	5	4	6	11	11	15	22	16	9	4	1	-
1999.....	6	10	9	4	15	25	17	22	12	2	-	-
Asian or Pacific Islander 2/, female												
2004.....	5	4	4	8	12	28	28	24	15	7	1	-
2003.....	-	2	5	10	13	24	27	28	17	3	2	-
2002.....	-	4	2	9	10	18	22	25	13	4	1	-
2001.....	1	4	3	4	8	11	21	16	13	4	2	-
2000.....	2	3	6	6	9	13	15	18	12	7	-	-
1999.....	1	2	1	2	12	13	6	13	4	1	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Fall on same level (W00-W09,W18)												
All races, both sexes												
2004.....	146	157	225	265	431	753	1,204	1,253	930	359	63	1
2003.....	143	160	177	223	381	710	1,113	1,174	847	296	53	1
2002.....	132	136	150	208	364	657	947	1,132	832	318	49	-
2001.....	104	107	137	181	287	502	739	775	580	243	43	1
2000.....	92	86	106	129	238	405	588	665	498	202	36	-
1999.....	56	46	56	108	170	311	399	426	334	100	31	-
All races, male												
2004.....	114	105	143	168	238	401	569	513	349	89	16	1
2003.....	102	107	110	120	210	377	551	486	301	74	8	1
2002.....	90	84	95	127	222	339	434	465	284	89	7	-
2001.....	69	70	89	107	145	276	357	318	203	59	9	-
2000.....	65	54	61	74	131	200	278	259	168	51	8	-
1999.....	39	29	32	66	98	160	190	172	118	20	7	-
All races, female												
2004.....	32	52	82	97	193	352	635	740	581	270	47	-
2003.....	41	53	67	103	171	333	562	688	546	222	45	-
2002.....	42	52	55	81	142	318	513	667	548	229	42	-
2001.....	35	37	48	74	142	226	382	457	377	184	34	1
2000.....	27	32	45	55	107	205	310	406	330	151	28	-
1999.....	17	17	24	42	72	151	209	254	216	80	24	-
White, both sexes												
2004.....	124	132	186	240	384	688	1,145	1,211	903	346	60	1
2003.....	117	128	150	205	346	674	1,054	1,130	813	287	49	-
2002.....	117	118	141	189	332	610	902	1,084	791	309	46	-
2001.....	91	91	120	167	260	460	704	737	552	234	42	1
2000.....	73	69	89	113	221	386	559	641	478	191	33	-
1999.....	40	34	38	88	155	286	374	409	319	97	29	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Fall on same level (W00-W09,W18)												
White, male												
2004.....	96	90	116	159	212	367	537	493	335	85	16	1
2003.....	80	86	91	108	195	355	523	465	285	72	8	-
2002.....	80	72	90	115	202	311	407	443	262	85	6	-
2001.....	58	60	78	99	134	255	337	304	193	55	9	-
2000.....	52	43	52	63	125	187	263	249	158	49	8	-
1999.....	31	22	23	54	92	147	178	165	111	19	7	-
White, female												
2004.....	28	42	70	81	172	321	608	718	568	261	44	-
2003.....	37	42	59	97	151	319	531	665	528	215	41	-
2002.....	37	46	51	74	130	299	495	641	529	224	40	-
2001.....	33	31	42	68	126	205	367	433	359	179	33	1
2000.....	21	26	37	50	96	199	296	392	320	142	25	-
1999.....	9	12	15	34	63	139	196	244	208	78	22	-
Black, both sexes												
2004.....	17	20	28	18	35	50	33	17	18	8	2	-
2003.....	16	21	17	7	13	16	23	19	13	4	2	-
2002.....	13	10	8	11	21	26	18	24	25	6	1	-
2001.....	13	13	13	11	19	24	18	16	18	5	-	-
2000.....	12	13	13	9	12	9	17	13	10	5	2	-
1999.....	14	11	14	18	11	13	18	9	10	1	2	-
Black, male												
2004.....	15	12	20	6	20	27	17	4	10	2	-	-
2003.....	13	14	13	5	3	12	10	7	4	1	-	-
2002.....	8	5	4	8	13	19	12	10	12	3	-	-
2001.....	11	9	8	7	7	11	11	5	5	3	-	-
2000.....	7	8	8	7	4	5	6	6	5	1	-	-
1999.....	7	7	7	11	6	6	8	2	4	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Fall on same level (W00-W09,W18)												
Black, female												
2004.....	2	8	8	12	15	23	16	13	8	6	2	-
2003.....	3	7	4	2	10	4	13	12	9	3	2	-
2002.....	5	5	4	3	8	7	6	14	13	3	1	-
2001.....	2	4	5	4	12	13	7	11	13	2	-	-
2000.....	5	5	5	2	8	4	11	7	5	4	2	-
1999.....	7	4	7	7	5	7	10	7	6	1	2	-
American Indian 1/, both sexes												
2004.....	2	1	5	1	1	1	3	5	3	2	-	-
2003.....	6	3	5	4	3	6	6	5	3	2	1	1
2002.....	1	2	-	3	2	7	5	2	4	-	-	-
2001.....	-	1	2	1	1	5	2	3	3	2	-	-
2000.....	4	2	2	-	2	2	1	2	1	-	1	-
1999.....	1	1	1	1	-	3	2	2	2	-	-	-
American Indian 1/, male												
2004.....	1	1	3	-	-	1	1	4	1	-	-	-
2003.....	5	1	5	4	3	3	5	2	1	-	-	1
2002.....	1	2	-	2	1	3	2	1	-	-	-	-
2001.....	-	1	2	-	1	3	1	-	2	-	-	-
2000.....	4	2	1	-	1	2	1	1	1	-	-	-
1999.....	1	-	-	-	-	-	-	1	1	-	-	-
American Indian 1/, female												
2004.....	1	-	2	1	1	-	2	1	2	2	-	-
2003.....	1	2	-	-	-	3	1	3	2	2	1	-
2002.....	-	-	-	1	1	4	3	1	4	-	-	-
2001.....	-	-	-	1	-	2	1	3	1	2	-	-
2000.....	-	-	1	-	1	-	-	1	-	-	1	-
1999.....	-	1	1	1	-	3	2	1	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Fall on same level (W00-W09,W18)												
Asian or Pacific Islander 2/, both sexes												
2004.....	3	4	6	6	11	14	23	20	6	3	1	-
2003.....	4	8	5	7	19	14	30	20	18	3	1	-
2002.....	1	6	1	5	9	14	22	22	12	3	2	-
2001.....	-	2	2	2	7	13	15	19	7	2	1	-
2000.....	3	2	2	7	3	8	11	9	9	6	-	-
1999.....	1	-	3	1	4	9	5	6	3	2	-	-
Asian or Pacific Islander 2/, male												
2004.....	2	2	4	3	6	6	14	12	3	2	-	-
2003.....	4	6	1	3	9	7	13	12	11	1	-	-
2002.....	1	5	1	2	6	6	13	11	10	1	1	-
2001.....	-	-	1	1	3	7	8	9	3	1	-	-
2000.....	2	1	-	4	1	6	8	3	4	1	-	-
1999.....	-	-	2	1	-	7	4	4	2	1	-	-
Asian or Pacific Islander 2/, female												
2004.....	1	2	2	3	5	8	9	8	3	1	1	-
2003.....	-	2	4	4	10	7	17	8	7	2	1	-
2002.....	-	1	-	3	3	8	9	11	2	2	1	-
2001.....	-	2	1	1	4	6	7	10	4	1	1	-
2000.....	1	1	2	3	2	2	3	6	5	5	-	-
1999.....	1	-	1	-	4	2	1	2	1	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Fall from one level to another (W10-W17)												
All races, both sexes												
2004.....	221	237	226	218	290	346	408	311	159	37	6	2
2003.....	192	178	207	227	273	346	409	304	178	50	6	-
2002.....	177	188	194	218	295	381	400	321	135	37	7	-
2001.....	199	178	168	221	267	378	390	311	146	29	10	1
2000.....	179	168	158	178	265	310	320	233	110	25	4	-
1999.....	149	168	199	223	254	354	358	264	137	27	3	2
All races, male												
2004.....	190	194	178	165	208	207	237	159	74	11	3	2
2003.....	162	144	154	163	196	225	233	156	91	13	1	-
2002.....	140	160	153	157	203	252	242	171	61	10	-	-
2001.....	173	149	137	171	180	256	219	164	60	11	2	1
2000.....	156	136	125	128	191	212	199	139	54	8	-	-
1999.....	128	135	159	176	164	207	228	141	61	10	1	2
All races, female												
2004.....	31	43	48	53	82	139	171	152	85	26	3	-
2003.....	30	34	53	64	77	121	176	148	87	37	5	-
2002.....	37	28	41	61	92	129	158	150	74	27	7	-
2001.....	26	29	31	50	87	122	171	147	86	18	8	-
2000.....	23	32	33	50	74	98	121	94	56	17	4	-
1999.....	21	33	40	47	90	147	130	123	76	17	2	-
White, both sexes												
2004.....	188	205	203	203	267	328	383	295	152	37	5	2
2003.....	155	162	184	192	246	322	383	288	170	49	6	-
2002.....	159	162	180	197	272	355	379	299	125	34	7	-
2001.....	167	155	144	199	246	352	375	294	138	26	9	1
2000.....	148	146	143	152	244	296	306	222	104	22	4	-
1999.....	121	147	184	203	233	330	341	251	133	25	3	2

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Fall from one level to another (W10-W17)												
White, male												
2004.....	163	166	156	157	189	201	224	155	70	11	2	2
2003.....	130	132	139	133	177	209	221	150	89	13	1	-
2002.....	127	138	143	143	191	234	233	160	58	9	-	-
2001.....	144	129	116	152	165	237	211	157	56	9	1	1
2000.....	128	117	114	111	174	201	190	132	49	7	-	-
1999.....	104	116	145	159	146	190	216	136	61	9	1	2
White, female												
2004.....	25	39	47	46	78	127	159	140	82	26	3	-
2003.....	25	30	45	59	69	113	162	138	81	36	5	-
2002.....	32	24	37	54	81	121	146	139	67	25	7	-
2001.....	23	26	28	47	81	115	164	137	82	17	8	-
2000.....	20	29	29	41	70	95	116	90	55	15	4	-
1999.....	17	31	39	44	87	140	125	115	72	16	2	-
Black, both sexes												
2004.....	25	25	19	11	15	10	14	9	3	-	1	-
2003.....	24	15	18	28	18	14	19	8	6	1	-	-
2002.....	14	20	11	12	18	16	11	14	8	3	-	-
2001.....	28	17	14	18	18	19	12	11	6	3	-	-
2000.....	27	19	12	21	14	11	8	5	5	2	-	-
1999.....	21	14	10	19	12	14	10	9	4	2	-	-
Black, male												
2004.....	20	22	18	7	13	4	7	3	3	-	1	-
2003.....	19	11	11	26	10	11	6	5	2	-	-	-
2002.....	10	17	7	8	9	11	6	7	3	1	-	-
2001.....	25	15	12	16	14	14	8	2	2	2	-	-
2000.....	24	17	9	14	11	9	3	3	4	1	-	-
1999.....	18	12	9	17	10	10	6	3	-	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Fall from one level to another (W10-W17)												
Black, female												
2004.....	5	3	1	4	2	6	7	6	-	-	-	-
2003.....	5	4	7	2	8	3	13	3	4	1	-	-
2002.....	4	3	4	4	9	5	5	7	5	2	-	-
2001.....	3	2	2	2	4	5	4	9	4	1	-	-
2000.....	3	2	3	7	3	2	5	2	1	1	-	-
1999.....	3	2	1	2	2	4	4	6	4	1	-	-
American Indian 1/, both sexes												
2004.....	4	2	2	-	3	-	2	-	-	-	-	-
2003.....	5	1	2	2	3	2	-	-	-	-	-	-
2002.....	1	2	1	-	-	-	2	-	1	-	-	-
2001.....	-	2	2	2	1	1	-	1	-	-	-	-
2000.....	2	1	-	-	-	1	1	1	-	-	-	-
1999.....	3	1	2	-	3	2	-	-	-	-	-	-
American Indian 1/, male												
2004.....	4	2	2	-	3	-	1	-	-	-	-	-
2003.....	5	1	2	1	3	1	-	-	-	-	-	-
2002.....	-	2	1	-	-	-	1	-	-	-	-	-
2001.....	-	1	2	1	-	-	-	1	-	-	-	-
2000.....	2	1	-	-	-	1	1	1	-	-	-	-
1999.....	2	1	2	-	3	1	-	-	-	-	-	-
American Indian 1/, female												
2004.....	-	-	-	-	-	-	1	-	-	-	-	-
2003.....	-	-	-	1	-	1	-	-	-	-	-	-
2002.....	1	-	-	-	-	-	1	-	1	-	-	-
2001.....	-	1	-	1	1	1	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	1	-	-	-	-	1	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Fall from one level to another (W10-W17)												
Asian or Pacific Islander 2/, both sexes												
2004.....	4	5	2	4	5	8	9	7	4	-	-	-
2003.....	8	-	3	5	6	8	7	8	2	-	-	-
2002.....	3	4	2	9	5	10	8	8	1	-	-	-
2001.....	4	4	8	2	2	6	3	5	2	-	1	-
2000.....	2	2	3	5	7	2	5	5	1	1	-	-
1999.....	4	6	3	1	6	8	7	4	-	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	3	4	2	1	3	2	5	1	1	-	-	-
2003.....	8	-	2	3	6	4	6	1	-	-	-	-
2002.....	3	3	2	6	3	7	2	4	-	-	-	-
2001.....	4	4	7	2	1	5	-	4	2	-	1	-
2000.....	2	1	2	3	6	1	5	3	1	-	-	-
1999.....	4	6	3	-	5	6	6	2	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	1	1	-	3	2	6	4	6	3	-	-	-
2003.....	-	-	1	2	-	4	1	7	2	-	-	-
2002.....	-	1	-	3	2	3	6	4	1	-	-	-
2001.....	-	-	1	-	1	1	3	1	-	-	-	-
2000.....	-	1	1	2	1	1	-	2	-	1	-	-
1999.....	-	-	-	1	1	2	1	2	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Unspecified fall (W19)												
All races, both sexes												
2004.....	250	275	273	402	649	1,147	1,824	1,883	1,344	538	79	1
2003.....	200	241	257	337	607	1,080	1,591	1,742	1,196	460	98	-
2002.....	221	191	230	333	549	973	1,522	1,579	1,078	429	73	1
2001.....	201	178	236	345	532	993	1,438	1,561	1,124	446	82	2
2000.....	191	194	237	288	562	926	1,292	1,454	1,074	391	80	-
1999.....	172	184	234	332	576	971	1,463	1,618	1,154	406	78	2
All races, male												
2004.....	161	200	167	242	373	581	863	793	468	139	16	-
2003.....	145	157	155	202	318	543	732	717	404	119	16	-
2002.....	164	135	144	190	303	488	689	626	346	132	17	-
2001.....	153	125	148	191	296	509	660	667	392	118	20	2
2000.....	137	136	168	176	291	483	575	594	370	114	19	-
1999.....	128	132	163	191	332	473	692	651	389	110	21	2
All races, female												
2004.....	89	75	106	160	276	566	961	1,090	876	399	63	1
2003.....	55	84	102	135	289	537	859	1,025	792	341	82	-
2002.....	57	56	86	143	246	485	833	953	732	297	56	1
2001.....	48	53	88	154	236	484	778	894	732	328	62	-
2000.....	54	58	69	112	271	443	717	860	704	277	61	-
1999.....	44	52	71	141	244	498	771	967	765	296	57	-
White, both sexes												
2004.....	204	229	235	343	585	1,085	1,735	1,803	1,283	514	76	1
2003.....	157	203	224	290	547	998	1,487	1,674	1,137	448	95	-
2002.....	172	157	186	283	492	909	1,458	1,527	1,026	411	69	1
2001.....	173	145	204	294	487	930	1,360	1,490	1,077	427	74	2
2000.....	156	159	196	249	501	857	1,223	1,399	1,031	371	74	-
1999.....	140	143	176	284	506	902	1,399	1,546	1,099	384	71	2

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Unspecified fall (W19)												
White, male												
2004.....	128	162	138	207	335	552	821	748	440	131	15	-
2003.....	116	131	134	175	287	509	676	691	383	115	16	-
2002.....	122	111	112	158	270	459	657	602	326	126	16	-
2001.....	132	103	128	163	272	475	619	627	373	108	17	2
2000.....	109	109	139	152	264	445	544	570	352	103	17	-
1999.....	102	101	123	159	294	434	663	614	363	103	20	2
White, female												
2004.....	76	67	97	136	250	533	914	1,055	843	383	61	1
2003.....	41	72	90	115	260	489	811	983	754	333	79	-
2002.....	50	46	74	125	222	450	801	925	700	285	53	1
2001.....	41	42	76	131	215	455	741	863	704	319	57	-
2000.....	47	50	57	97	237	412	679	829	679	268	57	-
1999.....	38	42	53	125	212	468	736	932	736	281	51	-
Black, both sexes												
2004.....	36	40	30	48	48	39	56	45	44	15	3	-
2003.....	37	32	30	37	49	49	69	47	40	9	2	-
2002.....	41	27	35	34	42	44	40	34	34	13	4	-
2001.....	25	26	28	44	35	51	51	44	27	8	7	-
2000.....	28	28	31	34	47	45	47	32	31	14	5	-
1999.....	29	34	48	44	47	48	52	41	41	18	5	-
Black, male												
2004.....	26	33	25	27	31	21	29	22	21	5	1	-
2003.....	25	21	18	21	24	17	32	20	13	2	-	-
2002.....	36	20	26	21	24	18	19	17	14	4	1	-
2001.....	19	17	18	24	19	27	26	23	12	3	3	-
2000.....	23	22	25	20	22	25	22	11	14	7	1	-
1999.....	24	26	32	29	27	27	22	18	16	5	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Unspecified fall (W19)												
Black, female												
2004.....	10	7	5	21	17	18	27	23	23	10	2	-
2003.....	12	11	12	16	25	32	37	27	27	7	2	-
2002.....	5	7	9	13	18	26	21	17	20	9	3	-
2001.....	6	9	10	20	16	24	25	21	15	5	4	-
2000.....	5	6	6	14	25	20	25	21	17	7	4	-
1999.....	5	8	16	15	20	21	30	23	25	13	4	-
American Indian 1/, both sexes												
2004.....	3	3	5	3	4	2	6	8	1	-	-	-
2003.....	4	4	1	2	2	3	4	4	5	-	-	-
2002.....	4	4	2	4	2	2	5	3	2	1	-	-
2001.....	2	3	2	-	3	2	2	6	4	2	-	-
2000.....	5	4	3	-	4	6	1	3	1	2	-	-
1999.....	1	1	6	-	6	-	1	6	1	3	1	-
American Indian 1/, male												
2004.....	3	3	3	2	-	1	1	6	-	-	-	-
2003.....	2	3	1	2	1	-	2	2	2	-	-	-
2002.....	2	3	1	2	1	-	1	2	-	-	-	-
2001.....	2	3	1	-	1	1	1	1	-	1	-	-
2000.....	4	3	-	-	1	5	-	3	-	1	-	-
1999.....	-	1	4	-	1	-	-	3	-	1	-	-
American Indian 1/, female												
2004.....	-	-	2	1	4	1	5	2	1	-	-	-
2003.....	2	1	-	-	1	3	2	2	3	-	-	-
2002.....	2	1	1	2	1	2	4	1	2	1	-	-
2001.....	-	-	1	-	2	1	1	5	4	1	-	-
2000.....	1	1	3	-	3	1	1	-	1	1	-	-
1999.....	1	-	2	-	5	-	1	3	1	2	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Unspecified fall (W19)												
Asian or Pacific Islander 2/, both sexes												
2004.....	7	3	3	8	12	21	27	27	16	9	-	-
2003.....	2	2	2	8	9	30	31	17	14	3	1	-
2002.....	4	3	7	12	13	18	19	15	16	4	-	-
2001.....	1	4	2	7	7	10	25	21	16	9	1	-
2000.....	2	3	7	5	10	18	21	20	11	4	1	-
1999.....	2	6	4	4	17	21	11	25	13	1	1	-
Asian or Pacific Islander 2/, male												
2004.....	4	2	1	6	7	7	12	17	7	3	-	-
2003.....	2	2	2	4	6	17	22	4	6	2	-	-
2002.....	4	1	5	9	8	11	12	5	6	2	-	-
2001.....	-	2	1	4	4	6	14	16	7	6	-	-
2000.....	1	2	4	4	4	8	9	10	4	3	1	-
1999.....	2	4	4	3	10	12	7	16	10	1	-	-
Asian or Pacific Islander 2/, female												
2004.....	3	1	2	2	5	14	15	10	9	6	-	-
2003.....	-	-	-	4	3	13	9	13	8	1	1	-
2002.....	-	2	2	3	5	7	7	10	10	2	-	-
2001.....	1	2	1	3	3	4	11	5	9	3	1	-
2000.....	1	1	3	1	6	10	12	10	7	1	-	-
1999.....	-	2	-	1	7	9	4	9	3	-	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Accidental discharge of firearms (W32-W34)												
All races, both sexes												
2004.....	65	25	10	20	20	15	13	5	3	-	-	-
2003.....	44	36	26	23	16	16	21	3	3	-	-	-
2002.....	43	36	20	17	19	18	14	5	1	1	-	-
2001.....	42	44	25	19	14	24	17	6	2	-	-	2
2000.....	40	36	14	24	17	17	7	7	4	-	-	-
1999.....	41	25	23	15	18	15	8	8	2	-	-	-
All races, male												
2004.....	56	20	9	15	18	12	11	5	3	-	-	-
2003.....	35	35	24	20	10	15	18	3	3	-	-	-
2002.....	39	32	18	15	17	15	11	4	1	1	-	-
2001.....	36	34	21	16	10	22	14	5	2	-	-	1
2000.....	38	30	14	20	14	16	7	7	4	-	-	-
1999.....	32	22	19	13	18	13	7	5	2	-	-	-
All races, female												
2004.....	9	5	1	5	2	3	2	-	-	-	-	-
2003.....	9	1	2	3	6	1	3	-	-	-	-	-
2002.....	4	4	2	2	2	3	3	1	-	-	-	-
2001.....	6	10	4	3	4	2	3	1	-	-	-	1
2000.....	2	6	-	4	3	1	-	-	-	-	-	-
1999.....	9	3	4	2	-	2	1	3	-	-	-	-
White, both sexes												
2004.....	61	23	8	19	17	13	11	4	3	-	-	-
2003.....	41	34	25	22	12	16	17	2	2	-	-	-
2002.....	37	32	18	16	19	14	14	5	1	1	-	-
2001.....	41	38	24	18	12	23	15	6	2	-	-	2
2000.....	37	34	13	22	15	16	6	7	4	-	-	-
1999.....	36	21	20	13	17	13	7	8	2	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Accidental discharge of firearms (W32-W34)												
White, male												
2004.....	54	19	7	14	16	10	9	4	3	-	-	-
2003.....	33	34	23	19	8	15	14	2	2	-	-	-
2002.....	34	29	17	14	17	11	11	4	1	1	-	-
2001.....	36	29	20	15	8	21	12	5	2	-	-	1
2000.....	35	28	13	19	12	15	6	7	4	-	-	-
1999.....	27	19	18	11	17	11	6	5	2	-	-	-
White, female												
2004.....	7	4	1	5	1	3	2	-	-	-	-	-
2003.....	8	-	2	3	4	1	3	-	-	-	-	-
2002.....	3	3	1	2	2	3	3	1	-	-	-	-
2001.....	5	9	4	3	4	2	3	1	-	-	-	1
2000.....	2	6	-	3	3	1	-	-	-	-	-	-
1999.....	9	2	2	2	-	2	1	3	-	-	-	-
Black, both sexes												
2004.....	4	2	2	1	3	2	2	1	-	-	-	-
2003.....	3	2	1	1	4	-	3	1	1	-	-	-
2002.....	5	4	2	-	-	4	-	-	-	-	-	-
2001.....	1	5	1	1	2	1	2	-	-	-	-	-
2000.....	3	2	1	1	2	-	1	-	-	-	-	-
1999.....	5	4	3	2	1	2	1	-	-	-	-	-
Black, male												
2004.....	2	1	2	1	2	2	2	1	-	-	-	-
2003.....	2	1	1	1	2	-	3	1	1	-	-	-
2002.....	4	3	1	-	-	4	-	-	-	-	-	-
2001.....	-	4	1	1	2	1	2	-	-	-	-	-
2000.....	3	2	1	-	2	-	1	-	-	-	-	-
1999.....	5	3	1	2	1	2	1	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Accidental explosion (W39-W40)												
All races, both sexes												
2004.....	12	9	8	11	6	5	6	3	-	-	-	-
2003.....	13	9	10	5	10	7	2	4	4	-	-	-
2002.....	11	11	8	3	8	9	2	2	-	-	-	-
2001.....	18	6	8	9	9	10	9	2	-	-	-	-
2000.....	7	10	6	7	13	4	8	4	1	-	-	-
1999.....	15	13	6	5	9	5	5	7	1	-	-	-
All races, male												
2004.....	11	8	8	7	3	2	4	2	-	-	-	-
2003.....	11	8	8	5	8	5	2	2	3	-	-	-
2002.....	9	10	8	1	6	6	1	2	-	-	-	-
2001.....	16	6	7	8	7	4	5	-	-	-	-	-
2000.....	5	10	4	5	11	1	5	4	-	-	-	-
1999.....	14	11	3	2	8	4	5	4	1	-	-	-
All races, female												
2004.....	1	1	-	4	3	3	2	1	-	-	-	-
2003.....	2	1	2	-	2	2	-	2	1	-	-	-
2002.....	2	1	-	2	2	3	1	-	-	-	-	-
2001.....	2	-	1	1	2	6	4	2	-	-	-	-
2000.....	2	-	2	2	2	3	3	-	1	-	-	-
1999.....	1	2	3	3	1	1	-	3	-	-	-	-
White, both sexes												
2004.....	10	9	7	11	6	5	5	3	-	-	-	-
2003.....	13	5	10	5	10	7	2	3	-	-	-	-
2002.....	11	9	7	3	6	9	2	2	-	-	-	-
2001.....	16	6	6	7	7	8	9	2	-	-	-	-
2000.....	6	9	6	7	10	3	6	4	-	-	-	-
1999.....	15	10	5	5	8	5	5	6	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Accidental explosion (W39-W40)												
White, male												
2004.....	9	8	7	7	3	2	4	2	-	-	-	-
2003.....	11	5	8	5	8	5	2	2	-	-	-	-
2002.....	9	8	7	1	4	6	1	2	-	-	-	-
2001.....	14	6	5	6	5	3	5	-	-	-	-	-
2000.....	5	9	4	5	8	-	3	4	-	-	-	-
1999.....	14	9	3	2	7	4	5	4	-	-	-	-
White, female												
2004.....	1	1	-	4	3	3	1	1	-	-	-	-
2003.....	2	-	2	-	2	2	-	1	-	-	-	-
2002.....	2	1	-	2	2	3	1	-	-	-	-	-
2001.....	2	-	1	1	2	5	4	2	-	-	-	-
2000.....	1	-	2	2	2	3	3	-	-	-	-	-
1999.....	1	1	2	3	1	1	-	2	-	-	-	-
Black, both sexes												
2004.....	2	-	1	-	-	-	1	-	-	-	-	-
2003.....	-	3	-	-	-	-	-	-	4	-	-	-
2002.....	-	2	-	-	2	-	-	-	-	-	-	-
2001.....	2	-	2	1	2	2	-	-	-	-	-	-
2000.....	1	1	-	-	3	1	2	-	1	-	-	-
1999.....	-	3	-	-	1	-	-	-	-	-	-	-
Black, male												
2004.....	2	-	1	-	-	-	-	-	-	-	-	-
2003.....	-	3	-	-	-	-	-	-	3	-	-	-
2002.....	-	2	-	-	2	-	-	-	-	-	-	-
2001.....	2	-	2	1	2	1	-	-	-	-	-	-
2000.....	-	1	-	-	3	1	2	-	-	-	-	-
1999.....	-	2	-	-	1	-	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Accidental explosion (W39-W40)												
Black, female												
2004.....	-	-	-	-	-	-	1	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	1	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	1	-	-	-	-	-	-
2000.....	1	-	-	-	-	-	-	-	1	-	-	-
1999.....	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	1	-	-	-	-	-	1	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	1	-	-	-	-	1	-	-	-	-
American Indian 1/, male												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, female												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	1	-	-	-	-	-	1	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	1	-	-	-	-	1	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Accidental drowning and submersion (W65-W74)												
All races, both sexes												
2004.....	204	152	119	105	83	76	82	51	23	3	1	14
2003.....	186	153	112	102	100	94	68	58	20	5	-	30
2002.....	169	118	94	81	88	88	79	48	25	5	1	20
2001.....	145	109	97	78	81	85	72	56	15	4	-	34
2000.....	172	120	97	80	99	71	85	37	19	8	-	24
1999.....	172	122	97	84	101	88	70	45	16	7	-	34
All races, male												
2004.....	158	119	89	81	59	54	59	27	13	2	1	14
2003.....	147	125	86	73	72	57	47	31	15	2	-	30
2002.....	137	93	68	64	68	56	50	31	15	4	-	18
2001.....	117	83	72	61	63	55	47	37	8	-	-	29
2000.....	143	102	77	63	70	38	50	22	7	5	-	23
1999.....	145	95	75	64	65	63	48	19	6	3	-	30
All races, female												
2004.....	46	33	30	24	24	22	23	24	10	1	-	-
2003.....	39	28	26	29	28	37	21	27	5	3	-	-
2002.....	32	25	26	17	20	32	29	17	10	1	1	2
2001.....	28	26	25	17	18	30	25	19	7	4	-	5
2000.....	29	18	20	17	29	33	35	15	12	3	-	1
1999.....	27	27	22	20	36	25	22	26	10	4	-	4
White, both sexes												
2004.....	159	120	99	92	65	63	71	43	21	2	-	14
2003.....	153	131	93	92	86	80	65	49	17	4	-	29
2002.....	133	103	78	68	71	75	68	41	22	3	-	20
2001.....	125	85	81	66	73	76	62	50	14	4	-	28
2000.....	130	105	82	70	86	58	79	34	16	6	-	23
1999.....	142	99	82	72	82	74	64	39	16	5	-	32

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Accidental drowning and submersion (W65-W74)												
White, male												
2004.....	125	95	73	73	47	45	54	23	13	2	-	14
2003.....	119	105	70	66	63	51	45	29	12	2	-	29
2002.....	110	81	55	54	56	50	44	29	14	3	-	18
2001.....	99	63	58	52	57	50	41	33	8	-	-	25
2000.....	107	89	70	55	59	31	50	20	7	3	-	22
1999.....	122	76	63	53	53	53	44	17	6	3	-	28
White, female												
2004.....	34	25	26	19	18	18	17	20	8	-	-	-
2003.....	34	26	23	26	23	29	20	20	5	2	-	-
2002.....	23	22	23	14	15	25	24	12	8	-	-	2
2001.....	26	22	23	14	16	26	21	17	6	4	-	3
2000.....	23	16	12	15	27	27	29	14	9	3	-	1
1999.....	20	23	19	19	29	21	20	22	10	2	-	4
Black, both sexes												
2004.....	34	19	10	8	12	5	5	3	2	1	1	-
2003.....	24	19	15	5	11	6	2	4	1	-	-	1
2002.....	24	12	10	9	9	10	7	6	2	2	1	-
2001.....	17	21	11	7	6	5	8	3	1	-	-	3
2000.....	29	9	8	7	8	9	3	3	2	2	-	1
1999.....	20	19	10	9	12	11	4	3	-	2	-	1
Black, male												
2004.....	23	15	7	5	10	4	3	-	-	-	1	-
2003.....	22	17	13	3	7	3	2	2	1	-	-	1
2002.....	19	11	8	8	7	5	4	2	1	1	-	-
2001.....	15	17	9	7	5	4	5	3	-	-	-	3
2000.....	24	8	5	6	7	4	-	2	-	2	-	1
1999.....	18	16	7	8	9	8	3	2	-	-	-	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Accidental drowning and submersion (W65-W74)												
Asian or Pacific Islander 2/, both sexes												
2004.....	7	11	8	4	6	8	5	5	-	-	-	-
2003.....	6	2	4	4	3	7	1	5	2	1	-	-
2002.....	9	2	5	4	8	3	3	1	1	-	-	-
2001.....	1	1	4	5	2	4	2	3	-	-	-	3
2000.....	7	3	6	2	4	4	3	-	1	-	-	-
1999.....	5	2	3	2	5	2	2	3	-	-	-	1
Asian or Pacific Islander 2/, male												
2004.....	7	7	7	2	2	5	2	4	-	-	-	-
2003.....	5	2	3	3	2	2	-	-	2	-	-	-
2002.....	6	1	4	2	5	1	2	-	-	-	-	-
2001.....	1	1	4	2	1	1	1	1	-	-	-	1
2000.....	7	2	1	1	3	3	-	-	-	-	-	-
1999.....	3	1	3	2	1	1	1	-	-	-	-	1
Asian or Pacific Islander 2/, female												
2004.....	-	4	1	2	4	3	3	1	-	-	-	-
2003.....	1	-	1	1	1	5	1	5	-	1	-	-
2002.....	3	1	1	2	3	2	1	1	1	-	-	-
2001.....	-	-	-	3	1	3	1	2	-	-	-	2
2000.....	-	1	5	1	1	1	3	-	1	-	-	-
1999.....	2	1	-	-	4	1	1	3	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Accidental inhalation and ingestion of food or other objects causing obstruction of respiratory tract (W78-W80)												
All races, both sexes												
2004.....	183	187	199	207	331	523	748	672	529	211	41	-
2003.....	165	182	204	233	335	494	706	659	425	167	42	-
2002.....	143	170	152	251	313	540	670	637	439	186	41	-
2001.....	180	167	163	221	318	535	658	669	430	181	43	-
2000.....	151	156	173	219	350	573	662	667	488	178	48	-
1999.....	134	127	145	212	354	481	591	630	435	158	35	-
All races, male												
2004.....	99	116	112	122	178	293	360	313	193	71	11	-
2003.....	87	105	135	128	177	278	368	303	177	52	10	-
2002.....	87	107	92	149	171	295	321	280	168	58	8	-
2001.....	104	107	87	127	167	280	338	300	164	47	6	-
2000.....	101	94	96	125	176	296	347	274	173	46	10	-
1999.....	97	72	80	111	204	238	286	264	154	46	4	-
All races, female												
2004.....	84	71	87	85	153	230	388	359	336	140	30	-
2003.....	78	77	69	105	158	216	338	356	248	115	32	-
2002.....	56	63	60	102	142	245	349	357	271	128	33	-
2001.....	76	60	76	94	151	255	320	369	266	134	37	-
2000.....	50	62	77	94	174	277	315	393	315	132	38	-
1999.....	37	55	65	101	150	243	305	366	281	112	31	-
White, both sexes												
2004.....	153	158	164	166	267	455	668	605	484	195	36	-
2003.....	130	152	164	192	280	440	627	598	396	156	36	-
2002.....	112	137	123	190	259	472	600	575	401	166	34	-
2001.....	154	130	131	186	258	453	576	608	376	162	38	-
2000.....	120	127	142	171	289	507	594	605	442	155	45	-
1999.....	112	96	109	176	292	406	520	569	400	144	33	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Accidental inhalation and ingestion of food or other objects causing obstruction of respiratory tract (W78-W80)												
White, male												
2004.....	82	99	95	97	146	264	321	282	175	67	10	-
2003.....	70	89	106	111	150	248	326	275	165	49	10	-
2002.....	65	86	77	111	147	258	284	255	157	50	6	-
2001.....	86	80	70	109	136	237	287	273	141	42	6	-
2000.....	78	76	78	97	147	262	308	241	160	36	9	-
1999.....	80	53	59	93	172	199	254	241	142	42	3	-
White, female												
2004.....	71	59	69	69	121	191	347	323	309	128	26	-
2003.....	60	63	58	81	130	192	301	323	231	107	26	-
2002.....	47	51	46	79	112	214	316	320	244	116	28	-
2001.....	68	50	61	77	122	216	289	335	235	120	32	-
2000.....	42	51	64	74	142	245	286	364	282	119	36	-
1999.....	32	43	50	83	120	207	266	328	258	102	30	-
Black, both sexes												
2004.....	28	25	29	39	51	61	70	54	37	13	4	-
2003.....	31	23	34	35	49	39	65	51	19	9	5	-
2002.....	29	29	25	55	53	64	61	56	35	15	5	-
2001.....	21	32	31	33	56	66	69	54	46	16	4	-
2000.....	28	25	28	45	54	61	59	54	38	18	3	-
1999.....	21	27	32	33	57	69	63	55	33	14	2	-
Black, male												
2004.....	15	16	13	23	27	25	36	23	15	4	1	-
2003.....	15	12	23	15	24	23	35	20	7	2	-	-
2002.....	21	18	13	35	24	36	31	22	10	6	1	-
2001.....	13	23	17	18	29	34	45	23	19	3	-	-
2000.....	21	15	16	25	25	33	33	27	9	8	1	-
1999.....	16	16	19	16	31	37	29	22	10	4	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Accidental inhalation and ingestion of food or other objects causing obstruction of respiratory tract (W78-W80)												
Black, female												
2004.....	13	9	16	16	24	36	34	31	22	9	3	-
2003.....	16	11	11	20	25	16	30	31	12	7	5	-
2002.....	8	11	12	20	29	28	30	34	25	9	4	-
2001.....	8	9	14	15	27	32	24	31	27	13	4	-
2000.....	7	10	12	20	29	28	26	27	29	10	2	-
1999.....	5	11	13	17	26	32	34	33	23	10	1	-
American Indian 1/, both sexes												
2004.....	1	2	4	-	3	3	2	3	1	-	-	-
2003.....	3	2	3	2	1	5	2	1	1	-	-	-
2002.....	2	2	2	3	-	1	1	-	-	1	-	-
2001.....	3	1	1	-	1	6	3	2	3	1	-	-
2000.....	3	3	1	-	2	1	5	1	2	-	-	-
1999.....	1	1	4	1	1	2	1	1	-	-	-	-
American Indian 1/, male												
2004.....	1	1	3	-	1	2	1	1	1	-	-	-
2003.....	1	1	3	1	1	2	-	1	1	-	-	-
2002.....	1	1	1	1	-	-	-	-	-	-	-	-
2001.....	3	1	-	-	-	4	-	2	-	-	-	-
2000.....	2	3	1	-	2	1	3	1	-	-	-	-
1999.....	1	1	2	1	-	-	-	-	-	-	-	-
American Indian 1/, female												
2004.....	-	1	1	-	2	1	1	2	-	-	-	-
2003.....	2	1	-	1	-	3	2	-	-	-	-	-
2002.....	1	1	1	2	-	1	1	-	-	1	-	-
2001.....	-	-	1	-	1	2	3	-	3	1	-	-
2000.....	1	-	-	-	-	-	2	-	2	-	-	-
1999.....	-	-	2	-	1	2	1	1	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Accidental inhalation and ingestion of food or other objects causing obstruction of respiratory tract (W78-W80)												
Asian or Pacific Islander 2/, both sexes												
2004.....	1	2	2	2	10	4	8	10	7	3	1	-
2003.....	1	5	3	4	5	10	12	9	9	2	1	-
2002.....	-	2	2	3	1	3	8	6	3	4	2	-
2001.....	2	4	-	2	3	10	10	5	5	2	1	-
2000.....	-	1	2	3	5	4	4	7	6	5	-	-
1999.....	-	3	-	2	4	4	7	5	2	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	1	-	1	2	4	2	2	7	2	-	-	-
2003.....	1	3	3	1	2	5	7	7	4	1	-	-
2002.....	-	2	1	2	-	1	6	3	1	2	1	-
2001.....	2	3	-	-	2	5	6	2	4	2	-	-
2000.....	-	-	1	3	2	-	3	5	4	2	-	-
1999.....	-	2	-	1	1	2	3	1	2	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	-	2	1	-	6	2	6	3	5	3	1	-
2003.....	-	2	-	3	3	5	5	2	5	1	1	-
2002.....	-	-	1	1	1	2	2	3	2	2	1	-
2001.....	-	1	-	2	1	5	4	3	1	-	1	-
2000.....	-	1	1	-	3	4	1	2	2	3	-	-
1999.....	-	1	-	1	3	2	4	4	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other accidental and unspecified threats to breathing (W75-W77,W81-W84)												
All races, both sexes												
2004.....	62	32	25	24	18	20	17	20	4	2	2	-
2003.....	55	35	24	15	25	16	22	21	11	4	-	1
2002.....	53	40	25	27	20	28	17	28	14	6	-	-
2001.....	45	30	21	22	34	32	23	18	15	5	-	-
2000.....	58	51	32	25	32	34	26	23	22	6	-	-
1999.....	59	44	32	43	72	102	86	110	62	19	6	-
All races, male												
2004.....	52	24	20	15	10	10	11	4	2	1	1	-
2003.....	45	27	20	11	16	8	7	8	3	1	-	1
2002.....	47	29	18	18	11	19	8	9	5	-	-	-
2001.....	40	22	12	15	19	16	9	4	6	-	-	-
2000.....	48	38	26	18	19	17	12	9	8	2	-	-
1999.....	46	36	19	30	41	56	33	39	18	5	2	-
All races, female												
2004.....	10	8	5	9	8	10	6	16	2	1	1	-
2003.....	10	8	4	4	9	8	15	13	8	3	-	-
2002.....	6	11	7	9	9	9	9	19	9	6	-	-
2001.....	5	8	9	7	15	16	14	14	9	5	-	-
2000.....	10	13	6	7	13	17	14	14	14	4	-	-
1999.....	13	8	13	13	31	46	53	71	44	14	4	-
White, both sexes												
2004.....	58	30	24	20	18	17	16	19	4	2	2	-
2003.....	51	32	22	12	24	13	20	20	10	3	-	1
2002.....	50	33	25	23	18	25	17	23	12	6	-	-
2001.....	39	26	17	21	30	30	20	17	13	5	-	-
2000.....	55	46	31	22	25	31	25	22	17	6	-	-
1999.....	52	34	27	39	61	86	78	100	61	18	5	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other accidental and unspecified threats to breathing (W75-W77,W81-W84)												
White, male												
2004.....	48	22	19	13	10	7	11	4	2	1	1	-
2003.....	41	27	18	8	16	5	6	7	2	1	-	1
2002.....	44	23	18	15	11	16	8	6	4	-	-	-
2001.....	36	20	10	15	18	15	8	4	5	-	-	-
2000.....	47	36	25	15	14	15	11	8	6	2	-	-
1999.....	42	27	16	26	32	48	30	36	18	5	2	-
White, female												
2004.....	10	8	5	7	8	10	5	15	2	1	1	-
2003.....	10	5	4	4	8	8	14	13	8	2	-	-
2002.....	6	10	7	8	7	9	9	17	8	6	-	-
2001.....	3	6	7	6	12	15	12	13	8	5	-	-
2000.....	8	10	6	7	11	16	14	14	11	4	-	-
1999.....	10	7	11	13	29	38	48	64	43	13	3	-
Black, both sexes												
2004.....	4	2	-	2	-	3	-	1	-	-	-	-
2003.....	4	2	1	2	-	3	2	-	1	1	-	-
2002.....	2	5	-	4	2	3	-	5	2	-	-	-
2001.....	4	4	1	-	3	2	3	1	2	-	-	-
2000.....	1	4	1	2	7	3	1	1	4	-	-	-
1999.....	5	9	4	3	11	15	8	9	1	1	1	-
Black, male												
2004.....	4	2	-	1	-	3	-	-	-	-	-	-
2003.....	4	-	1	2	-	3	1	-	1	-	-	-
2002.....	2	4	-	3	-	3	-	3	1	-	-	-
2001.....	3	2	1	-	1	1	1	-	1	-	-	-
2000.....	-	2	1	2	5	2	1	1	1	-	-	-
1999.....	2	8	2	3	9	7	3	2	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Accidental exposure to electric current (W85-W87)												
All races, both sexes												
2004.....	31	20	18	4	4	2	3	-	-	-	-	-
2003.....	22	24	7	9	8	1	1	3	-	-	-	-
2002.....	29	13	10	6	4	4	3	-	-	-	-	-
2001.....	18	19	13	11	2	7	4	5	-	-	-	-
2000.....	22	18	15	4	3	1	2	-	-	-	-	-
1999.....	17	28	12	11	10	5	4	2	-	-	-	-
All races, male												
2004.....	31	17	18	3	4	2	2	-	-	-	-	-
2003.....	19	23	7	9	8	1	1	3	-	-	-	-
2002.....	28	12	10	6	4	2	2	-	-	-	-	-
2001.....	16	17	13	11	2	6	3	5	-	-	-	-
2000.....	21	18	14	4	2	1	1	-	-	-	-	-
1999.....	17	27	12	11	9	4	4	1	-	-	-	-
All races, female												
2004.....	-	3	-	1	-	-	1	-	-	-	-	-
2003.....	3	1	-	-	-	-	-	-	-	-	-	-
2002.....	1	1	-	-	-	2	1	-	-	-	-	-
2001.....	2	2	-	-	-	1	1	-	-	-	-	-
2000.....	1	-	1	-	1	-	1	-	-	-	-	-
1999.....	-	1	-	-	1	1	-	1	-	-	-	-
White, both sexes												
2004.....	27	19	17	3	4	2	3	-	-	-	-	-
2003.....	20	21	6	9	7	1	1	3	-	-	-	-
2002.....	25	11	9	5	3	3	2	-	-	-	-	-
2001.....	15	15	12	10	2	7	3	5	-	-	-	-
2000.....	19	17	11	3	3	1	2	-	-	-	-	-
1999.....	14	25	11	9	10	5	4	1	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Accidental exposure to smoke, fire and flames (X00-X09)												
All races, both sexes												
2004.....	238	205	207	183	166	214	245	163	81	22	7	3
2003.....	256	204	188	198	221	226	234	157	67	24	5	4
2002.....	235	183	157	174	211	211	221	129	70	18	4	3
2001.....	206	215	163	190	221	225	198	159	72	18	2	3
2000.....	225	187	182	188	213	257	215	158	78	24	3	5
1999.....	170	157	162	193	223	258	229	150	74	18	3	4
All races, male												
2004.....	169	127	124	121	89	108	126	70	31	5	3	2
2003.....	148	138	116	113	131	124	104	72	27	8	-	3
2002.....	146	111	108	117	125	108	123	55	31	9	2	3
2001.....	137	142	94	112	117	106	98	69	30	8	1	3
2000.....	158	126	115	107	125	136	108	69	34	11	1	5
1999.....	110	98	96	111	133	128	119	67	34	9	1	4
All races, female												
2004.....	69	78	83	62	77	106	119	93	50	17	4	1
2003.....	108	66	72	85	90	102	130	85	40	16	5	1
2002.....	89	72	49	57	86	103	98	74	39	9	2	-
2001.....	69	73	69	78	104	119	100	90	42	10	1	-
2000.....	67	61	67	81	88	121	107	89	44	13	2	-
1999.....	60	59	66	82	90	130	110	83	40	9	2	-
White, both sexes												
2004.....	181	154	153	138	116	163	193	133	59	21	3	2
2003.....	188	156	147	152	168	175	181	119	59	19	2	4
2002.....	174	140	107	130	159	164	178	104	49	11	3	3
2001.....	149	155	122	138	163	170	151	126	53	10	1	3
2000.....	175	129	131	139	158	189	156	129	51	17	-	5
1999.....	127	123	112	136	157	187	173	108	53	13	2	3

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Accidental exposure to smoke, fire and flames (X00-X09)												
White, male												
2004.....	126	92	98	89	64	81	104	58	22	5	3	1
2003.....	115	103	93	87	102	90	78	58	23	6	-	3
2002.....	105	86	75	86	93	85	105	47	20	5	1	3
2001.....	95	105	67	80	83	76	78	57	21	4	-	3
2000.....	122	89	80	76	93	100	81	61	22	8	-	5
1999.....	82	74	75	77	92	89	95	52	26	7	1	3
White, female												
2004.....	55	62	55	49	52	82	89	75	37	16	-	1
2003.....	73	53	54	65	66	85	103	61	36	13	2	1
2002.....	69	54	32	44	66	79	73	57	29	6	2	-
2001.....	54	50	55	58	80	94	73	69	32	6	1	-
2000.....	53	40	51	63	65	89	75	68	29	9	-	-
1999.....	45	49	37	59	65	98	78	56	27	6	1	-
Black, both sexes												
2004.....	50	47	51	39	45	47	49	26	22	1	4	1
2003.....	63	45	34	42	48	51	49	36	8	4	3	-
2002.....	58	39	48	38	47	42	41	25	20	7	1	-
2001.....	51	53	37	48	54	49	43	31	18	8	1	-
2000.....	48	53	49	45	53	64	57	29	26	7	3	-
1999.....	41	34	46	52	61	67	53	41	21	5	1	1
Black, male												
2004.....	38	32	25	30	24	27	21	10	9	-	-	1
2003.....	32	33	20	26	27	34	23	13	4	1	-	-
2002.....	39	23	32	27	28	20	18	8	11	4	1	-
2001.....	39	32	24	29	33	26	18	11	8	4	1	-
2000.....	34	35	34	29	31	35	26	8	12	3	1	-
1999.....	27	24	20	30	39	36	23	15	8	2	-	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Accidental exposure to smoke, fire and flames (X00-X09)												
Black, female												
2004.....	12	15	26	9	21	20	28	16	13	1	4	-
2003.....	31	12	14	16	21	17	26	23	4	3	3	-
2002.....	19	16	16	11	19	22	23	17	9	3	-	-
2001.....	12	21	13	19	21	23	25	20	10	4	-	-
2000.....	14	18	15	16	22	29	31	21	14	4	2	-
1999.....	14	10	26	22	22	31	30	26	13	3	1	-
American Indian 1/, both sexes												
2004.....	3	3	2	4	1	2	2	1	-	-	-	-
2003.....	3	1	3	2	3	-	-	-	-	-	-	-
2002.....	1	2	1	4	4	1	-	-	1	-	-	-
2001.....	5	5	2	3	2	2	-	1	-	-	-	-
2000.....	1	3	2	1	1	1	-	-	-	-	-	-
1999.....	1	-	3	5	1	3	3	-	-	-	-	-
American Indian 1/, male												
2004.....	3	2	1	2	-	-	1	-	-	-	-	-
2003.....	1	-	1	-	1	-	-	-	-	-	-	-
2002.....	1	2	-	4	3	1	-	-	-	-	-	-
2001.....	2	4	2	2	1	1	-	1	-	-	-	-
2000.....	1	1	1	-	1	1	-	-	-	-	-	-
1999.....	1	-	-	4	1	2	1	-	-	-	-	-
American Indian 1/, female												
2004.....	-	1	1	2	1	2	1	1	-	-	-	-
2003.....	2	1	2	2	2	-	-	-	-	-	-	-
2002.....	-	-	1	-	1	-	-	-	1	-	-	-
2001.....	3	1	-	1	1	1	-	-	-	-	-	-
2000.....	-	2	1	1	-	-	-	-	-	-	-	-
1999.....	-	-	3	1	-	1	2	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Accidental exposure to smoke, fire and flames (X00-X09)												
Asian or Pacific Islander 2/, both sexes												
2004.....	4	1	1	2	4	2	1	3	-	-	-	-
2003.....	2	2	4	2	2	-	4	2	-	1	-	-
2002.....	2	2	1	2	1	4	2	-	-	-	-	-
2001.....	1	2	2	1	2	4	4	1	1	-	-	-
2000.....	1	2	-	3	1	3	2	-	1	-	-	-
1999.....	1	-	1	-	4	1	-	1	-	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	2	1	-	-	1	-	-	2	-	-	-	-
2003.....	-	2	2	-	1	-	3	1	-	1	-	-
2002.....	1	-	1	-	1	2	-	-	-	-	-	-
2001.....	1	1	1	1	-	3	2	-	1	-	-	-
2000.....	1	1	-	2	-	-	1	-	-	-	-	-
1999.....	-	-	1	-	1	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	2	-	1	2	3	2	1	1	-	-	-	-
2003.....	2	-	2	2	1	-	1	1	-	-	-	-
2002.....	1	2	-	2	-	2	2	-	-	-	-	-
2001.....	-	1	1	-	2	1	2	1	-	-	-	-
2000.....	-	1	-	1	1	3	1	-	1	-	-	-
1999.....	1	-	-	-	3	-	-	1	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Accidental contact with heat and hot substances (X10-X19)												
All races, both sexes												
2004.....	9	10	5	4	4	6	10	8	8	4	-	-
2003.....	5	4	4	7	3	6	14	12	7	1	1	-
2002.....	3	3	8	8	8	15	10	10	4	2	-	-
2001.....	10	10	7	8	8	8	16	12	7	3	-	-
2000.....	2	2	9	5	13	9	16	16	6	3	1	-
1999.....	3	9	6	9	9	7	19	16	9	3	-	-
All races, male												
2004.....	8	9	2	2	2	5	3	3	3	1	-	-
2003.....	2	2	3	3	1	3	9	5	3	-	-	-
2002.....	2	1	2	4	4	8	4	3	3	1	-	-
2001.....	9	4	1	4	3	5	7	4	2	1	-	-
2000.....	1	2	8	4	6	4	8	6	-	1	-	-
1999.....	2	7	3	5	4	5	8	2	3	-	-	-
All races, female												
2004.....	1	1	3	2	2	1	7	5	5	3	-	-
2003.....	3	2	1	4	2	3	5	7	4	1	1	-
2002.....	1	2	6	4	4	7	6	7	1	1	-	-
2001.....	1	6	6	4	5	3	9	8	5	2	-	-
2000.....	1	-	1	1	7	5	8	10	6	2	1	-
1999.....	1	2	3	4	5	2	11	14	6	3	-	-
White, both sexes												
2004.....	8	6	4	3	2	4	9	5	6	4	-	-
2003.....	3	3	1	5	3	3	11	10	4	1	1	-
2002.....	2	3	4	6	4	11	8	7	4	-	-	-
2001.....	8	7	6	7	4	7	12	12	7	3	-	-
2000.....	2	1	6	4	10	4	12	11	5	3	1	-
1999.....	1	6	3	5	7	6	15	11	9	2	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Accidental contact with heat and hot substances (X10-X19)												
White, male												
2004.....	7	5	2	2	1	4	3	2	1	1	-	-
2003.....	1	1	1	3	1	2	6	3	2	-	-	-
2002.....	1	1	-	2	2	6	3	2	3	-	-	-
2001.....	7	2	1	4	1	4	5	4	2	1	-	-
2000.....	1	1	6	4	5	-	6	5	-	1	-	-
1999.....	-	4	1	3	3	4	6	1	3	-	-	-
White, female												
2004.....	1	1	2	1	1	-	6	3	5	3	-	-
2003.....	2	2	-	2	2	1	5	7	2	1	1	-
2002.....	1	2	4	4	2	5	5	5	1	-	-	-
2001.....	1	5	5	3	3	3	7	8	5	2	-	-
2000.....	1	-	-	-	5	4	6	6	5	2	1	-
1999.....	1	2	2	2	4	2	9	10	6	2	-	-
Black, both sexes												
2004.....	-	4	1	1	2	2	1	3	1	-	-	-
2003.....	2	1	3	2	-	3	1	2	2	-	-	-
2002.....	1	-	3	2	2	3	1	2	-	2	-	-
2001.....	2	3	1	1	4	-	4	-	-	-	-	-
2000.....	-	1	3	1	2	5	2	5	1	-	-	-
1999.....	2	3	3	4	2	1	2	2	-	1	-	-
Black, male												
2004.....	-	4	-	-	1	1	-	1	1	-	-	-
2003.....	1	1	2	-	-	1	1	2	1	-	-	-
2002.....	1	-	2	2	2	1	-	1	-	1	-	-
2001.....	2	2	-	-	2	-	2	-	-	-	-	-
2000.....	-	1	2	-	1	4	1	1	-	4	-	-
1999.....	2	3	2	2	1	1	1	1	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Accidental contact with heat and hot substances (X10-X19)												
Asian or Pacific Islander 2/, both sexes												
2004.....	1	-	-	-	-	-	-	-	1	-	-	-
2003.....	-	-	-	-	-	-	2	-	-	-	-	-
2002.....	-	-	1	-	2	1	1	1	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	1	-	2	-	-	-	-	-
1999.....	-	-	-	-	-	-	2	3	-	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	1	-	-	-	-	-	-	-	1	-	-	-
2003.....	-	-	-	-	-	-	2	-	-	-	-	-
2002.....	-	-	-	-	-	1	1	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	1	-	-	-	-	-
1999.....	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	1	-	2	-	-	1	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	1	-	1	-	-	-	-	-
1999.....	-	-	-	-	-	-	1	3	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Earthquake, volcanic eruption, avalanche, landslide and other earth movements (X34-X36)												
All races, both sexes												
2004.....	1	4	-	-	-	-	-	-	-	-	-	-
2003.....	3	3	-	-	-	-	-	1	-	-	-	-
2002.....	-	-	2	-	-	-	-	-	-	-	-	-
2001.....	1	2	-	-	-	-	-	-	-	-	-	-
2000.....	5	-	1	-	-	-	-	-	-	-	-	-
1999.....	1	1	1	-	-	1	-	-	-	-	-	-
All races, male												
2004.....	1	3	-	-	-	-	-	-	-	-	-	-
2003.....	2	2	-	-	-	-	-	1	-	-	-	-
2002.....	-	-	2	-	-	-	-	-	-	-	-	-
2001.....	-	1	-	-	-	-	-	-	-	-	-	-
2000.....	5	-	-	-	-	-	-	-	-	-	-	-
1999.....	1	1	1	-	-	-	-	-	-	-	-	-
All races, female												
2004.....	-	1	-	-	-	-	-	-	-	-	-	-
2003.....	1	1	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	1	1	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	1	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	1	-	-	-	-	-	-
White, both sexes												
2004.....	1	4	-	-	-	-	-	-	-	-	-	-
2003.....	2	3	-	-	-	-	-	1	-	-	-	-
2002.....	-	-	2	-	-	-	-	-	-	-	-	-
2001.....	1	2	-	-	-	-	-	-	-	-	-	-
2000.....	4	-	1	-	-	-	-	-	-	-	-	-
1999.....	1	1	1	-	-	1	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cataclysmic storm and flood (X37-X38)												
All races, both sexes												
2004.....	7	8	9	3	8	4	7	3	-	-	-	-
2003.....	6	4	4	3	6	6	3	4	1	-	-	-
2002.....	8	2	4	4	6	5	3	1	1	-	-	-
2001.....	4	6	7	3	12	4	1	1	1	-	-	-
2000.....	8	3	4	3	1	2	4	1	1	-	-	-
1999.....	15	10	8	10	4	9	7	4	1	1	-	-
All races, male												
2004.....	4	8	2	2	4	1	5	1	-	-	-	-
2003.....	5	4	1	2	4	2	2	3	1	-	-	-
2002.....	7	2	2	1	3	2	2	1	-	-	-	-
2001.....	3	3	4	2	6	4	-	1	-	-	-	-
2000.....	2	1	2	3	1	2	2	1	-	-	-	-
1999.....	9	4	6	4	3	6	6	1	-	1	-	-
All races, female												
2004.....	3	-	7	1	4	3	2	2	-	-	-	-
2003.....	1	-	3	1	2	4	1	1	-	-	-	-
2002.....	1	-	2	3	3	3	1	-	1	-	-	-
2001.....	1	3	3	1	6	-	1	-	1	-	-	-
2000.....	6	2	2	-	-	-	2	-	1	-	-	-
1999.....	6	6	2	6	1	3	1	3	1	-	-	-
White, both sexes												
2004.....	6	7	9	3	7	4	7	3	-	-	-	-
2003.....	5	3	3	2	6	6	3	4	1	-	-	-
2002.....	6	2	4	4	6	5	3	1	1	-	-	-
2001.....	4	5	3	2	9	3	1	1	1	-	-	-
2000.....	5	-	4	3	1	2	4	1	1	-	-	-
1999.....	10	9	7	9	3	7	7	3	1	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cataclysmic storm and flood (X37-X38)												
White, male												
2004.....	3	7	2	2	4	1	5	1	-	-	-	-
2003.....	4	3	-	1	4	2	2	3	1	-	-	-
2002.....	5	2	2	1	3	2	2	1	-	-	-	-
2001.....	3	3	2	1	4	3	-	1	-	-	-	-
2000.....	2	-	2	3	1	2	2	1	-	-	-	-
1999.....	5	3	5	3	2	4	6	1	-	1	-	-
White, female												
2004.....	3	-	7	1	3	3	2	2	-	-	-	-
2003.....	1	-	3	1	2	4	1	1	-	-	-	-
2002.....	1	-	2	3	3	3	1	-	1	-	-	-
2001.....	1	2	1	1	5	-	1	-	1	-	-	-
2000.....	3	-	2	-	-	-	2	-	1	-	-	-
1999.....	5	6	2	6	1	3	1	2	1	-	-	-
Black, both sexes												
2004.....	1	-	-	-	1	-	-	-	-	-	-	-
2003.....	1	1	1	1	-	-	-	-	-	-	-	-
2002.....	2	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	1	4	1	3	-	-	-	-	-	-	-
2000.....	3	2	-	-	-	-	-	-	-	-	-	-
1999.....	4	1	1	-	1	2	-	1	-	-	-	-
Black, male												
2004.....	1	-	-	-	-	-	-	-	-	-	-	-
2003.....	1	1	1	1	-	-	-	-	-	-	-	-
2002.....	2	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	2	1	2	-	-	-	-	-	-	-
2000.....	-	1	-	-	-	-	-	-	-	-	-	-
1999.....	3	1	1	-	1	2	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Accidental poisoning and exposure to noxious substances (X40-X49)												
All races, both sexes												
2004.....	2,337	1,111	466	240	181	175	140	107	49	9	-	9
2003.....	2,083	957	413	229	174	146	138	105	47	10	4	6
2002.....	1,736	754	344	218	165	154	141	103	64	13	1	9
2001.....	1,281	530	268	161	130	153	126	91	47	12	2	10
2000.....	1,069	447	241	146	132	118	127	109	60	14	1	7
1999.....	918	449	219	158	142	138	127	100	31	12	-	12
All races, male												
2004.....	1,550	676	253	127	95	81	62	44	19	1	-	4
2003.....	1,368	584	257	137	86	71	71	34	19	4	1	5
2002.....	1,176	491	199	126	83	71	65	38	25	8	-	7
2001.....	893	341	159	88	82	83	54	28	18	2	1	10
2000.....	768	260	141	79	68	49	53	51	23	2	-	6
1999.....	650	298	150	98	74	63	53	38	10	5	-	8
All races, female												
2004.....	787	435	213	113	86	94	78	63	30	8	-	5
2003.....	715	373	156	92	88	75	67	71	28	6	3	1
2002.....	560	263	145	92	82	83	76	65	39	5	1	2
2001.....	388	189	109	73	48	70	72	63	29	10	1	-
2000.....	301	187	100	67	64	69	74	58	37	12	1	1
1999.....	268	151	69	60	68	75	74	62	21	7	-	4
White, both sexes												
2004.....	1,904	873	379	194	154	154	119	101	42	8	-	6
2003.....	1,676	787	350	167	149	133	130	94	41	9	4	4
2002.....	1,375	585	270	183	144	137	127	94	58	11	1	7
2001.....	952	416	200	135	110	131	106	80	41	10	-	9
2000.....	790	349	194	122	111	111	118	92	54	11	-	6
1999.....	685	353	157	127	113	128	117	84	29	10	-	9

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Accidental poisoning and exposure to noxious substances (X40-X49)

White, male

2004.....	1,234	503	189	93	80	67	52	42	17	-	-	3
2003.....	1,088	461	211	90	73	64	67	29	15	3	1	3
2002.....	911	361	139	101	69	64	54	34	25	7	-	5
2001.....	651	252	107	73	69	66	47	25	17	1	-	9
2000.....	558	187	105	60	55	46	49	43	21	1	-	5
1999.....	469	221	100	77	61	58	49	35	10	4	-	6

White, female

2004.....	670	370	190	101	74	87	67	59	25	8	-	3
2003.....	588	326	139	77	76	69	63	65	26	6	3	1
2002.....	464	224	131	82	75	73	73	60	33	4	1	2
2001.....	301	164	93	62	41	65	59	55	24	9	-	-
2000.....	232	162	89	62	56	65	69	49	33	10	-	1
1999.....	216	132	57	50	52	70	68	49	19	6	-	3

Black, both sexes

2004.....	400	211	73	40	24	18	17	6	6	1	-	2
2003.....	366	152	57	52	18	13	7	10	6	1	-	2
2002.....	339	154	69	28	17	15	12	6	5	1	-	1
2001.....	310	105	60	24	17	17	17	11	6	2	2	1
2000.....	266	92	43	24	19	7	8	16	6	3	1	1
1999.....	217	85	59	26	24	10	8	14	2	2	-	3

Black, male

2004.....	293	155	54	30	15	12	8	2	2	1	-	1
2003.....	255	113	45	42	12	7	3	5	4	1	-	2
2002.....	247	119	57	22	13	7	9	1	-	-	-	1
2001.....	229	81	47	15	12	14	6	3	1	1	1	1
2000.....	199	70	34	19	12	3	4	8	2	1	-	1
1999.....	170	70	47	18	12	5	4	3	-	1	-	2

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Accidental poisoning and exposure to noxious substances (X40-X49)												
Black, female												
2004.....	107	56	19	10	9	6	9	4	4	-	-	1
2003.....	111	39	12	10	6	6	4	5	2	-	-	-
2002.....	92	35	12	6	4	8	3	5	5	1	-	-
2001.....	81	24	13	9	5	3	11	8	5	1	1	-
2000.....	67	22	9	5	7	4	4	8	4	2	1	-
1999.....	47	15	12	8	12	5	4	11	2	1	-	1
American Indian 1/, both sexes												
2004.....	20	12	6	4	1	1	-	-	-	-	-	1
2003.....	25	10	4	8	1	-	-	-	-	-	-	-
2002.....	14	12	2	1	-	-	-	-	1	-	-	1
2001.....	11	5	6	1	2	-	1	-	-	-	-	-
2000.....	8	3	-	-	1	-	-	1	-	-	-	-
1999.....	6	5	1	3	1	-	-	1	-	-	-	-
American Indian 1/, male												
2004.....	13	7	4	2	-	-	-	-	-	-	-	-
2003.....	15	5	1	4	1	-	-	-	-	-	-	-
2002.....	11	9	1	1	-	-	-	-	-	-	-	1
2001.....	7	4	4	-	1	-	1	-	-	-	-	-
2000.....	8	2	-	-	-	-	-	-	-	-	-	-
1999.....	3	5	1	2	-	-	-	-	-	-	-	-
American Indian 1/, female												
2004.....	7	5	2	2	1	1	-	-	-	-	-	1
2003.....	10	5	3	4	-	-	-	-	-	-	-	-
2002.....	3	3	1	-	-	-	-	-	1	-	-	-
2001.....	4	1	2	1	1	-	-	-	-	-	-	-
2000.....	-	1	-	-	1	-	-	1	-	-	-	-
1999.....	3	-	-	1	1	-	-	1	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Accidental poisoning and exposure to noxious substances (X40-X49)												
Asian or Pacific Islander 2/, both sexes												
2004.....	13	15	8	2	2	2	4	-	1	-	-	-
2003.....	16	8	2	2	6	-	1	1	-	-	-	-
2002.....	8	3	3	6	4	2	2	3	-	1	-	-
2001.....	8	4	2	1	1	5	2	-	-	-	-	-
2000.....	5	3	4	-	1	-	1	-	-	-	-	-
1999.....	10	6	2	2	4	-	2	1	-	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	10	11	6	2	-	2	2	-	-	-	-	-
2003.....	10	5	-	1	-	-	1	-	-	-	-	-
2002.....	7	2	2	2	1	-	2	3	-	1	-	-
2001.....	6	4	1	-	-	3	-	-	-	-	-	-
2000.....	3	1	2	-	1	-	-	-	-	-	-	-
1999.....	8	2	2	1	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	3	4	2	-	2	-	2	-	1	-	-	-
2003.....	6	3	2	1	6	-	-	1	-	-	-	-
2002.....	1	1	1	4	3	2	-	-	-	-	-	-
2001.....	2	-	1	1	1	2	2	-	-	-	-	-
2000.....	2	2	2	-	-	-	1	-	-	-	-	-
1999.....	2	4	-	1	3	-	2	1	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Accidental poisoning by and exposure to drugs and other biological substances (X40-X44)												
All races, both sexes												
2004.....	2,217	1,031	421	201	133	135	98	78	36	8	-	9
2003.....	1,982	872	353	189	123	108	112	85	38	7	3	6
2002.....	1,634	667	287	169	117	111	100	80	53	9	1	7
2001.....	1,183	468	225	131	101	108	87	59	38	9	2	9
2000.....	989	374	181	117	90	91	85	80	48	11	-	7
1999.....	836	380	174	122	99	100	92	79	25	11	-	9
All races, male												
2004.....	1,462	621	220	98	66	59	36	25	13	1	-	4
2003.....	1,291	519	209	109	55	52	55	21	16	1	1	5
2002.....	1,100	423	158	91	53	48	42	27	20	5	-	5
2001.....	819	300	127	67	59	54	32	10	12	1	1	9
2000.....	705	207	103	64	46	35	27	36	16	2	-	6
1999.....	589	251	113	70	51	40	31	29	6	4	-	5
All races, female												
2004.....	755	410	201	103	67	76	62	53	23	7	-	5
2003.....	691	353	144	80	68	56	57	64	22	6	2	1
2002.....	534	244	129	78	64	63	58	53	33	4	1	2
2001.....	364	168	98	64	42	54	55	49	26	8	1	-
2000.....	284	167	78	53	44	56	58	44	32	9	-	1
1999.....	247	129	61	52	48	60	61	50	19	7	-	4
White, both sexes												
2004.....	1,802	805	344	162	116	120	86	75	31	7	-	6
2003.....	1,594	715	297	141	105	98	104	76	35	6	3	4
2002.....	1,293	524	221	140	104	98	92	74	49	7	1	5
2001.....	881	362	168	109	85	95	80	52	34	7	-	8
2000.....	726	292	141	96	77	86	78	66	44	8	-	6
1999.....	621	299	122	98	80	93	85	65	24	9	-	6

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Accidental poisoning by and exposure to drugs and other biological substances (X40-X44)

White, male

2004.....	1,162	456	164	68	57	49	30	25	12	-	-	3
2003.....	1,025	407	170	73	47	46	51	18	13	-	1	3
2002.....	851	316	104	71	45	42	34	24	20	4	-	3
2001.....	596	217	84	55	48	43	32	8	12	-	-	8
2000.....	505	146	72	48	36	34	24	30	15	1	-	5
1999.....	421	185	71	54	43	37	28	26	6	3	-	3

White, female

2004.....	640	349	180	94	59	71	56	50	19	7	-	3
2003.....	569	308	127	68	58	52	53	58	22	6	2	1
2002.....	442	208	117	69	59	56	58	50	29	3	1	2
2001.....	285	145	84	54	37	52	48	44	22	7	-	-
2000.....	221	146	69	48	41	52	54	36	29	7	-	1
1999.....	200	114	51	44	37	56	57	39	18	6	-	3

Black, both sexes

2004.....	387	201	66	34	16	13	9	3	5	1	-	2
2003.....	353	142	50	43	12	10	7	8	3	1	-	2
2002.....	323	135	63	23	10	11	6	4	4	1	-	1
2001.....	287	100	52	20	13	11	6	7	4	2	2	1
2000.....	253	79	37	21	12	5	6	13	4	3	-	1
1999.....	202	73	50	22	15	7	6	12	1	2	-	3

Black, male

2004.....	282	149	48	27	9	9	5	-	1	1	-	1
2003.....	244	105	38	34	7	6	3	3	3	1	-	2
2002.....	235	103	53	17	7	6	6	1	-	-	-	1
2001.....	213	78	40	12	10	9	-	2	-	1	1	1
2000.....	192	60	30	16	9	1	3	6	1	1	-	1
1999.....	158	61	40	15	8	3	3	3	-	1	-	2

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Accidental poisoning by and exposure to drugs and other biological substances (X40-X44)												
Black, female												
2004.....	105	52	18	7	7	4	4	3	4	-	-	1
2003.....	109	37	12	9	5	4	4	5	-	-	-	-
2002.....	88	32	10	6	3	5	-	3	4	1	-	-
2001.....	74	22	12	8	3	2	6	5	4	1	1	-
2000.....	61	19	7	5	3	4	3	7	3	2	-	-
1999.....	44	12	10	7	7	4	3	9	1	1	-	1
American Indian 1/, both sexes												
2004.....	18	12	5	3	-	1	-	-	-	-	-	1
2003.....	22	8	4	4	1	-	-	-	-	-	-	-
2002.....	12	5	1	1	-	-	-	-	-	-	-	1
2001.....	7	3	3	1	2	-	-	-	-	-	-	-
2000.....	6	2	-	-	-	-	-	1	-	-	-	-
1999.....	5	4	-	-	1	-	-	1	-	-	-	-
American Indian 1/, male												
2004.....	11	7	3	1	-	-	-	-	-	-	-	-
2003.....	14	3	1	1	1	-	-	-	-	-	-	-
2002.....	9	2	-	1	-	-	-	-	-	-	-	1
2001.....	4	2	2	-	1	-	-	-	-	-	-	-
2000.....	6	1	-	-	-	-	-	-	-	-	-	-
1999.....	2	4	-	-	-	-	-	-	-	-	-	-
American Indian 1/, female												
2004.....	7	5	2	2	-	1	-	-	-	-	-	1
2003.....	8	5	3	3	-	-	-	-	-	-	-	-
2002.....	3	3	1	-	-	-	-	-	-	-	-	-
2001.....	3	1	1	1	1	-	-	-	-	-	-	-
2000.....	-	1	-	-	-	-	-	1	-	-	-	-
1999.....	3	-	-	-	1	-	-	1	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Accidental poisoning by and exposure to drugs and other biological substances (X40-X44)												
Asian or Pacific Islander 2/, both sexes												
2004.....	10	13	6	2	1	1	3	-	-	-	-	-
2003.....	13	7	2	1	5	-	1	1	-	-	-	-
2002.....	6	3	2	5	3	2	2	2	-	1	-	-
2001.....	8	3	2	1	1	2	1	-	-	-	-	-
2000.....	4	1	3	-	1	-	1	-	-	-	-	-
1999.....	8	4	2	2	3	-	1	1	-	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	7	9	5	2	-	1	1	-	-	-	-	-
2003.....	8	4	-	1	-	-	1	-	-	-	-	-
2002.....	5	2	1	2	1	-	2	2	-	1	-	-
2001.....	6	3	1	-	-	2	-	-	-	-	-	-
2000.....	2	-	1	-	1	-	-	-	-	-	-	-
1999.....	8	1	2	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	3	4	1	-	1	-	2	-	-	-	-	-
2003.....	5	3	2	-	5	-	-	1	-	-	-	-
2002.....	1	1	1	3	2	2	-	-	-	-	-	-
2001.....	2	-	1	1	1	-	1	-	-	-	-	-
2000.....	2	1	2	-	-	-	1	-	-	-	-	-
1999.....	-	3	-	1	3	-	1	1	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Accidental poisoning by and exposure to other and unspecified solid or liquids substances (X45-X46,X48-X49)

All races, both sexes

2004.....	72	37	19	15	12	16	14	5	4	-	-	-
2003.....	56	43	23	11	9	6	7	5	1	1	-	-
2002.....	55	46	22	10	16	9	12	4	5	-	-	-
2001.....	58	29	21	11	9	8	6	3	1	1	-	-
2000.....	41	36	24	10	14	6	7	6	4	2	-	-
1999.....	41	33	21	16	17	13	10	7	1	-	-	3

All races, male

2004.....	51	22	14	11	7	9	8	2	3	-	-	-
2003.....	43	32	21	8	3	2	4	2	-	1	-	-
2002.....	42	39	20	7	8	6	3	1	2	-	-	-
2001.....	41	19	15	7	8	6	3	2	1	-	-	-
2000.....	32	24	15	4	6	2	4	5	2	-	-	-
1999.....	29	21	16	13	8	6	6	2	-	-	-	3

All races, female

2004.....	21	15	5	4	5	7	6	3	1	-	-	-
2003.....	13	11	2	3	6	4	3	3	1	-	-	-
2002.....	13	7	2	3	8	3	9	3	3	-	-	-
2001.....	17	10	6	4	1	2	3	1	-	1	-	-
2000.....	9	12	9	6	8	4	3	1	2	2	-	-
1999.....	12	12	5	3	9	7	4	5	1	-	-	-

White, both sexes

2004.....	62	32	13	13	9	14	11	4	4	-	-	-
2003.....	46	36	20	7	8	5	7	5	1	1	-	-
2002.....	42	31	17	9	11	8	10	2	3	-	-	-
2001.....	43	25	15	9	8	8	5	3	1	1	-	-
2000.....	32	29	21	9	10	6	7	5	4	2	-	-
1999.....	33	24	16	10	14	13	9	7	1	-	-	3

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Accidental poisoning by and exposure to other and unspecified solid or liquids substances (X45-X46,X48-X49)												
White, male												
2004.....	42	19	9	10	6	7	6	2	3	-	-	-
2003.....	35	25	18	4	3	2	4	2	-	1	-	-
2002.....	32	25	16	6	4	6	3	1	2	-	-	-
2001.....	31	17	10	6	7	6	3	2	1	-	-	-
2000.....	26	18	12	3	5	2	4	4	2	-	-	-
1999.....	24	14	12	8	8	6	5	2	-	-	-	3
White, female												
2004.....	20	13	4	3	3	7	5	2	1	-	-	-
2003.....	11	11	2	3	5	3	3	3	1	-	-	-
2002.....	10	6	1	3	7	2	7	1	1	-	-	-
2001.....	12	8	5	3	1	2	2	1	-	1	-	-
2000.....	6	11	9	6	5	4	3	1	2	2	-	-
1999.....	9	10	4	2	6	7	4	5	1	-	-	-
Black, both sexes												
2004.....	8	4	5	2	1	2	2	1	-	-	-	-
2003.....	7	6	3	2	-	1	-	-	-	-	-	-
2002.....	10	10	3	1	5	1	2	2	1	-	-	-
2001.....	12	2	3	2	1	-	-	-	-	-	-	-
2000.....	7	7	3	1	3	-	-	1	-	-	-	-
1999.....	8	8	4	4	3	-	1	-	-	-	-	-
Black, male												
2004.....	7	2	4	1	1	2	1	-	-	-	-	-
2003.....	6	6	3	2	-	-	-	-	-	-	-	-
2002.....	7	9	2	1	4	-	-	-	-	-	-	-
2001.....	8	-	3	1	1	-	-	-	-	-	-	-
2000.....	4	6	3	1	6	-	-	1	-	-	-	-
1999.....	5	6	3	3	-	-	1	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Accidental poisoning by and exposure to other gases and vapors (X47)												
All races, both sexes												
2004.....	48	43	26	24	36	24	28	24	9	1	-	-
2003.....	45	42	37	29	42	32	19	15	8	2	1	-
2002.....	47	41	35	39	32	34	29	19	6	4	-	2
2001.....	40	33	22	19	20	37	33	29	8	2	-	1
2000.....	39	37	36	19	28	21	35	23	8	1	1	-
1999.....	41	36	24	20	26	25	25	14	5	1	-	-
All races, male												
2004.....	37	33	19	18	22	13	18	17	3	-	-	-
2003.....	34	33	27	20	28	17	12	11	3	2	-	-
2002.....	34	29	21	28	22	17	20	10	3	3	-	2
2001.....	33	22	17	14	15	23	19	16	5	1	-	1
2000.....	31	29	23	11	16	12	22	10	5	-	-	-
1999.....	32	26	21	15	15	17	16	7	4	1	-	-
All races, female												
2004.....	11	10	7	6	14	11	10	7	6	1	-	-
2003.....	11	9	10	9	14	15	7	4	5	-	1	-
2002.....	13	12	14	11	10	17	9	9	3	1	-	-
2001.....	7	11	5	5	5	14	14	13	3	1	-	-
2000.....	8	8	13	8	12	9	13	13	3	1	1	-
1999.....	9	10	3	5	11	8	9	7	1	-	-	-
White, both sexes												
2004.....	40	36	22	19	29	20	22	22	7	1	-	-
2003.....	36	36	33	19	36	30	19	13	5	2	1	-
2002.....	40	30	32	34	29	31	25	18	6	4	-	2
2001.....	28	29	17	17	17	28	21	25	6	2	-	1
2000.....	32	28	32	17	24	19	33	21	6	1	-	-
1999.....	31	30	19	19	19	22	23	12	4	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Accidental poisoning by and exposure to other gases and vapors (X47)												
White, male												
2004.....	30	28	16	15	17	11	16	15	2	-	-	-
2003.....	28	29	23	13	23	16	12	9	2	2	-	-
2002.....	28	20	19	24	20	16	17	9	3	3	-	2
2001.....	24	18	13	12	14	17	12	15	4	1	-	1
2000.....	27	23	21	9	14	10	21	9	4	-	-	-
1999.....	24	22	17	15	10	15	16	7	4	1	-	-
White, female												
2004.....	10	8	6	4	12	9	6	7	5	1	-	-
2003.....	8	7	10	6	13	14	7	4	3	-	1	-
2002.....	12	10	13	10	9	15	8	9	3	1	-	-
2001.....	4	11	4	5	3	11	9	10	2	1	-	-
2000.....	5	5	11	8	10	9	12	12	2	1	-	-
1999.....	7	8	2	4	9	7	7	5	-	-	-	-
Black, both sexes												
2004.....	5	6	2	4	7	3	6	2	1	-	-	-
2003.....	6	4	4	7	6	2	-	2	3	-	-	-
2002.....	6	9	3	4	2	3	4	-	-	-	-	-
2001.....	11	3	5	2	3	6	11	4	2	-	-	-
2000.....	6	6	3	2	4	2	2	2	2	-	1	-
1999.....	7	4	5	-	6	3	1	2	1	-	-	-
Black, male												
2004.....	4	4	2	2	5	1	2	2	1	-	-	-
2003.....	5	2	4	6	5	1	-	2	1	-	-	-
2002.....	5	7	2	4	2	1	3	-	-	-	-	-
2001.....	8	3	4	2	1	5	6	1	1	-	-	-
2000.....	3	4	1	2	2	2	1	1	1	-	-	-
1999.....	7	3	4	-	4	2	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Accidental poisoning by and exposure to other gases and vapors (X47)												
Asian or Pacific Islander 2/, both sexes												
2004.....	3	1	2	-	-	1	-	-	1	-	-	-
2003.....	2	1	-	1	-	-	-	-	-	-	-	-
2002.....	1	-	-	1	1	-	-	1	-	-	-	-
2001.....	-	1	-	-	-	3	-	-	-	-	-	-
2000.....	1	2	1	-	-	-	-	-	-	-	-	-
1999.....	2	2	-	-	1	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	3	1	1	-	-	1	-	-	-	-	-	-
2003.....	1	1	-	-	-	-	-	-	-	-	-	-
2002.....	1	-	-	-	-	-	-	1	-	-	-	-
2001.....	-	1	-	-	-	1	-	-	-	-	-	-
2000.....	1	1	1	-	-	-	-	-	-	-	-	-
1999.....	-	1	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	-	-	1	-	-	-	-	-	1	-	-	-
2003.....	1	-	-	1	-	-	-	-	-	-	-	-
2002.....	-	-	-	1	1	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	2	-	-	-	-	-	-
2000.....	-	1	-	-	-	-	-	-	-	-	-	-
1999.....	2	1	-	-	-	-	1	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and unspecified accidents and their sequelae (W20-W31,W35-W38,W41-W64,W92-W99,X20-X32,X39,X50-X59,Y86)												
All races, both sexes												
2004.....	571	465	452	443	565	864	1,289	1,371	1,208	466	89	31
2003.....	496	487	414	521	578	945	1,316	1,598	1,196	548	124	14
2002.....	499	480	448	524	607	1,000	1,358	1,576	1,262	533	109	13
2001.....	543	406	477	469	699	1,072	1,461	1,697	1,331	515	128	24
2000.....	467	424	439	478	669	1,009	1,328	1,574	1,280	519	128	16
1999.....	492	425	460	560	722	1,125	1,578	1,794	1,387	579	96	15
All races, male												
2004.....	456	353	343	301	341	463	592	527	379	129	14	30
2003.....	390	372	324	359	362	523	589	633	361	137	31	12
2002.....	387	370	339	352	379	547	609	583	369	142	16	10
2001.....	421	301	348	312	413	581	654	636	438	107	20	21
2000.....	387	337	315	313	392	534	571	589	383	130	23	16
1999.....	402	327	337	361	432	587	727	692	422	142	21	14
All races, female												
2004.....	115	112	109	142	224	401	697	844	829	337	75	1
2003.....	106	115	90	162	216	422	727	965	835	411	93	2
2002.....	112	110	109	172	228	453	749	993	893	391	93	3
2001.....	122	105	129	157	286	491	807	1,061	893	408	108	3
2000.....	80	87	124	165	277	475	757	985	897	389	105	-
1999.....	90	98	123	199	290	538	851	1,102	965	437	75	1
White, both sexes												
2004.....	472	380	385	375	497	765	1,188	1,291	1,140	448	83	28
2003.....	387	397	347	442	502	857	1,206	1,532	1,116	517	113	13
2002.....	414	395	379	417	541	888	1,243	1,488	1,192	503	101	13
2001.....	441	354	390	398	611	969	1,357	1,587	1,258	480	117	22
2000.....	381	344	367	401	585	895	1,234	1,489	1,215	491	112	14
1999.....	397	341	370	465	616	991	1,445	1,673	1,313	548	87	13

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and unspecified accidents and their sequelae (W20-W31,W35-W38,W41-W64,W92-W99,X20-X32,X39,X50-X59,Y86)												
White, male												
2004.....	381	286	296	259	304	408	545	496	356	123	13	27
2003.....	310	303	268	308	310	476	536	606	337	128	24	11
2002.....	322	300	288	281	339	486	559	548	341	137	12	10
2001.....	339	259	290	264	364	532	603	594	413	101	19	19
2000.....	314	267	258	260	349	465	524	556	357	122	18	14
1999.....	327	263	278	297	366	518	666	632	387	138	18	12
White, female												
2004.....	91	94	89	116	193	357	643	795	784	325	70	1
2003.....	77	94	79	134	192	381	670	926	779	389	89	2
2002.....	92	95	91	136	202	402	684	940	851	366	89	3
2001.....	102	95	100	134	247	437	754	993	845	379	98	3
2000.....	67	77	109	141	236	430	710	933	858	369	94	-
1999.....	70	78	92	168	250	473	779	1,041	926	410	69	1
Black, both sexes												
2004.....	84	73	52	55	51	86	79	59	54	17	6	1
2003.....	94	73	55	67	61	70	88	54	67	28	8	1
2002.....	69	70	54	88	47	91	87	75	53	25	5	-
2001.....	86	40	76	59	71	87	85	90	62	27	11	2
2000.....	68	72	59	62	70	100	75	72	51	26	14	2
1999.....	81	75	76	85	97	116	113	107	66	30	7	2
Black, male												
2004.....	64	57	38	35	30	47	39	20	17	5	1	1
2003.....	68	55	44	44	43	37	40	20	18	9	5	1
2002.....	53	58	40	54	27	49	39	28	20	4	2	-
2001.....	67	32	49	41	39	40	41	31	19	4	1	2
2000.....	61	63	45	42	35	59	39	27	22	7	4	2
1999.....	63	58	51	58	60	59	47	54	31	4	2	2

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and unspecified accidents and their sequelae (W20-W31,W35-W38,W41-W64,W92-W99,X20-X32,X39,X50-X59,Y86)												
Black, female												
2004.....	20	16	14	20	21	39	40	39	37	12	5	-
2003.....	26	18	11	23	18	33	48	34	49	19	3	-
2002.....	16	12	14	34	20	42	48	47	33	21	3	-
2001.....	19	8	27	18	32	47	44	59	43	23	10	-
2000.....	7	9	14	20	35	41	36	45	29	19	10	-
1999.....	18	17	25	27	37	57	66	53	35	26	5	-
American Indian 1/, both sexes												
2004.....	11	8	6	4	10	4	9	6	5	-	-	2
2003.....	8	13	9	8	5	9	7	5	4	1	1	-
2002.....	7	10	5	11	9	9	4	-	6	2	-	-
2001.....	10	4	7	5	7	6	6	4	3	2	-	-
2000.....	16	7	6	9	6	4	9	2	-	-	1	-
1999.....	7	5	9	5	5	11	7	4	2	-	1	-
American Indian 1/, male												
2004.....	9	7	4	2	6	1	4	2	3	-	-	2
2003.....	7	11	9	5	5	6	3	2	2	-	1	-
2002.....	5	8	3	10	7	5	3	-	4	1	-	-
2001.....	10	3	5	3	5	2	3	2	2	-	-	-
2000.....	11	6	5	6	3	1	3	1	-	-	-	-
1999.....	7	4	7	4	3	5	5	3	-	-	1	-
American Indian 1/, female												
2004.....	2	1	2	2	4	3	5	4	2	-	-	-
2003.....	1	2	-	3	-	3	4	3	2	1	-	-
2002.....	2	2	2	1	2	4	1	-	2	1	-	-
2001.....	-	1	2	2	2	4	3	2	1	2	-	-
2000.....	5	1	1	3	3	3	6	1	-	-	1	-
1999.....	-	1	2	1	2	6	2	1	2	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

All other and unspecified accidents and their sequelae (W20-W31,W35-W38,W41-W64,W92-W99,X20-X32,X39,X50-X59,Y86)

Asian or Pacific Islander 2/, both sexes

2004.....	4	4	9	9	7	9	13	15	9	1	-	-
2003.....	7	4	3	4	10	9	15	7	9	2	2	-
2002.....	9	5	10	8	10	12	24	13	11	3	3	-
2001.....	6	8	4	7	10	10	13	16	8	6	-	-
2000.....	2	1	7	6	8	10	10	11	14	2	1	-
1999.....	7	4	5	5	4	7	13	10	6	1	1	-

Asian or Pacific Islander 2/, male

2004.....	2	3	5	5	1	7	4	9	3	1	-	-
2003.....	5	3	3	2	4	4	10	5	4	-	1	-
2002.....	7	4	8	7	6	7	8	7	4	-	2	-
2001.....	5	7	4	4	5	7	7	9	4	2	-	-
2000.....	1	1	7	5	5	9	5	5	4	1	1	-
1999.....	5	2	1	2	3	5	9	3	4	-	-	-

Asian or Pacific Islander 2/, female

2004.....	2	1	4	4	6	2	9	6	6	-	-	-
2003.....	2	1	-	2	6	5	5	2	5	2	1	-
2002.....	2	1	2	1	4	5	16	6	7	3	1	-
2001.....	1	1	-	3	5	3	6	7	4	4	-	-
2000.....	1	-	-	1	3	1	5	6	10	1	-	-
1999.....	2	2	4	3	1	2	4	7	2	1	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)												
All races, both sexes												
2004.....	3,163	2,349	1,662	1,133	1,146	1,189	931	552	207	35	5	11
2003.....	2,996	2,314	1,529	1,167	1,168	1,159	956	558	205	34	1	7
2002.....	2,835	2,186	1,432	1,197	1,266	1,231	1,028	573	211	41	1	10
2001 3/.....	2,682	1,985	1,332	1,212	1,220	1,219	973	538	200	28	3	15
2000.....	2,450	1,708	1,237	1,073	1,219	1,227	954	590	205	34	4	7
1999.....	2,264	1,678	1,218	1,187	1,284	1,280	937	580	182	39	-	16
All races, male												
2004.....	2,329	1,787	1,301	937	964	1,005	812	475	172	27	5	11
2003.....	2,271	1,771	1,216	958	996	985	824	484	178	28	-	7
2002.....	2,188	1,708	1,129	975	1,078	1,054	884	492	179	33	-	10
2001 3/.....	2,020	1,534	1,029	1,001	1,039	1,033	852	468	168	25	3	14
2000.....	1,842	1,306	959	870	1,018	1,062	822	507	168	26	4	7
1999.....	1,702	1,284	971	972	1,079	1,073	795	501	150	30	-	16
All races, female												
2004.....	834	562	361	196	182	184	119	77	35	8	-	-
2003.....	725	543	313	209	172	174	132	74	27	6	1	-
2002.....	647	478	303	222	188	177	144	81	32	8	1	-
2001 3/.....	662	451	303	211	181	186	121	70	32	3	-	1
2000.....	608	402	278	203	201	165	132	83	37	8	-	-
1999.....	562	394	247	215	205	207	142	79	32	9	-	-
White, both sexes												
2004.....	2,967	2,196	1,555	1,064	1,073	1,135	884	536	192	35	5	9
2003.....	2,810	2,181	1,437	1,092	1,106	1,107	922	534	204	31	1	7
2002.....	2,640	2,064	1,350	1,130	1,199	1,163	1,000	551	203	38	1	10
2001 3/.....	2,510	1,859	1,247	1,132	1,150	1,169	931	518	195	27	3	14
2000.....	2,273	1,604	1,171	1,017	1,151	1,174	916	574	198	32	2	6
1999.....	2,121	1,567	1,131	1,104	1,226	1,226	903	563	178	38	-	13

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)												
White, male												
2004.....	2,191	1,657	1,228	878	905	965	773	464	163	27	5	9
2003.....	2,141	1,673	1,143	901	944	944	797	468	177	27	-	7
2002.....	2,054	1,610	1,074	919	1,034	1,001	860	475	173	30	-	10
2001 3/.....	1,893	1,438	958	942	979	990	820	455	164	25	3	13
2000.....	1,706	1,225	908	823	961	1,020	797	495	162	24	2	6
1999.....	1,586	1,196	903	911	1,029	1,031	770	487	148	29	-	13
White, female												
2004.....	776	539	327	186	168	170	111	72	29	8	-	-
2003.....	669	508	294	191	162	163	125	66	27	4	1	-
2002.....	586	454	276	211	165	162	140	76	30	8	1	-
2001 3/.....	617	421	289	190	171	179	111	63	31	2	-	1
2000.....	567	379	263	194	190	154	119	79	36	8	-	-
1999.....	535	371	228	193	197	195	133	76	30	9	-	-
Black, both sexes												
2004.....	114	107	59	34	42	33	27	5	7	-	-	1
2003.....	128	76	61	43	34	26	18	11	-	1	-	-
2002.....	123	69	58	33	49	39	14	16	2	2	-	-
2001 3/.....	118	82	49	43	48	30	24	8	4	1	-	-
2000.....	119	62	41	39	46	30	23	9	2	-	1	-
1999.....	84	68	50	58	38	35	17	10	4	1	-	2
Black, male												
2004.....	88	93	47	32	39	30	24	5	4	-	-	1
2003.....	93	57	50	32	27	24	16	8	-	-	-	-
2002.....	89	58	41	31	37	34	14	14	2	2	-	-
2001 3/.....	89	65	41	32	42	26	20	5	4	-	-	-
2000.....	90	51	31	36	40	25	15	8	2	-	1	-
1999.....	71	55	42	47	33	30	15	9	2	1	-	2

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)												
Asian or Pacific Islander 2/, both sexes												
2004.....	64	43	39	27	28	19	20	11	8	-	-	-
2003.....	47	52	24	28	26	22	16	12	1	2	-	-
2002.....	52	44	20	31	18	24	14	6	6	1	-	-
2001 3/.....	34	37	26	34	19	18	15	12	1	-	-	1
2000.....	43	29	22	14	17	22	15	7	5	2	1	1
1999.....	48	35	33	21	15	19	15	7	-	-	-	1
Asian or Pacific Islander 2/, male												
2004.....	39	34	20	20	18	8	15	6	5	-	-	-
2003.....	29	37	18	22	23	14	11	7	1	1	-	-
2002.....	30	33	11	22	7	14	10	3	4	1	-	-
2001 3/.....	19	25	23	26	15	15	9	8	-	-	-	1
2000.....	33	18	17	8	12	16	10	4	4	2	1	1
1999.....	38	28	24	10	12	12	8	5	-	-	-	1
Asian or Pacific Islander 2/, female												
2004.....	25	9	19	7	10	11	5	5	3	-	-	-
2003.....	18	15	6	6	3	8	5	5	-	1	-	-
2002.....	22	11	9	9	11	10	4	3	2	-	-	-
2001 3/.....	15	12	3	8	4	3	6	4	1	-	-	-
2000.....	10	11	5	6	5	6	5	3	1	-	-	-
1999.....	10	7	9	11	3	7	7	2	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Intentional self-poisoning (suicide) by and exposure to drugs and other biological substances (X60-X64)

All races, both sexes

2004.....	606	382	202	85	77	73	47	29	21	2	-	-
2003.....	525	329	169	109	72	68	48	35	8	-	-	-
2002.....	481	300	164	96	85	74	57	26	12	-	-	-
2001.....	434	236	131	97	79	65	31	25	11	3	-	3
2000.....	398	239	113	62	84	61	45	30	9	3	-	-
1999.....	370	212	117	96	81	78	53	18	12	2	-	1

All races, male

2004.....	275	165	94	30	33	30	18	17	7	-	-	-
2003.....	245	145	67	40	30	21	15	19	5	-	-	-
2002.....	241	126	68	39	33	33	26	11	5	-	-	-
2001.....	178	97	44	53	31	25	11	12	5	2	-	3
2000.....	176	105	38	24	30	27	24	12	3	1	-	-
1999.....	164	88	49	50	35	22	22	7	6	1	-	1

All races, female

2004.....	331	217	108	55	44	43	29	12	14	2	-	-
2003.....	280	184	102	69	42	47	33	16	3	-	-	-
2002.....	240	174	96	57	52	41	31	15	7	-	-	-
2001.....	256	139	87	44	48	40	20	13	6	1	-	-
2000.....	222	134	75	38	54	34	21	18	6	2	-	-
1999.....	206	124	68	46	46	56	31	11	6	1	-	-

White, both sexes

2004.....	573	365	191	80	74	73	46	28	18	2	-	-
2003.....	506	315	162	103	69	68	47	34	8	-	-	-
2002.....	450	294	153	91	80	70	55	26	12	-	-	-
2001.....	423	226	123	90	76	64	31	24	11	2	-	3
2000.....	373	232	112	59	83	60	43	30	9	3	-	-
1999.....	355	201	112	91	81	74	53	18	12	2	-	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intentional self-poisoning (suicide) by and exposure to drugs and other biological substances (X60-X64)												
Asian or Pacific Islander 2/, both sexes												
2004.....	11	4	2	3	1	-	1	1	2	-	-	-
2003.....	4	6	3	2	2	-	1	-	-	-	-	-
2002.....	5	3	2	3	2	3	1	-	-	-	-	-
2001.....	1	2	1	4	1	-	-	1	-	-	-	-
2000.....	3	1	1	1	-	-	-	-	-	-	-	-
1999.....	3	4	3	3	-	3	-	-	-	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	5	2	-	3	-	-	-	1	-	-	-	-
2003.....	2	2	2	1	2	-	-	-	-	-	-	-
2002.....	2	1	-	2	-	2	1	-	-	-	-	-
2001.....	-	1	1	2	1	-	-	-	-	-	-	-
2000.....	2	-	1	-	-	-	-	-	-	-	-	-
1999.....	2	-	3	2	-	2	-	-	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	6	2	2	-	1	-	1	-	2	-	-	-
2003.....	2	4	1	1	-	-	1	-	-	-	-	-
2002.....	3	2	2	1	2	1	-	-	-	-	-	-
2001.....	1	1	-	2	-	-	-	1	-	-	-	-
2000.....	1	1	-	1	-	-	-	-	-	-	-	-
1999.....	1	4	-	1	-	1	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intentional self-poisoning (suicide) by and exposure to other and unspecified solid or liquid substances and their vapors (X65-X66,X68-X69)												
All races, both sexes												
2004.....	28	34	13	7	7	6	7	2	1	-	-	-
2003.....	30	13	16	6	3	6	4	2	2	-	-	-
2002.....	19	14	3	6	9	6	4	3	-	-	-	-
2001.....	13	18	6	6	7	1	3	6	-	-	-	-
2000.....	20	9	7	7	6	3	5	1	2	-	-	-
1999.....	19	17	14	4	4	2	4	1	-	1	-	-
All races, male												
2004.....	15	21	7	3	7	5	7	1	1	-	-	-
2003.....	16	7	11	3	-	4	3	-	-	-	-	-
2002.....	13	10	2	3	5	4	2	1	-	-	-	-
2001.....	11	13	4	4	5	1	2	4	-	-	-	-
2000.....	12	5	6	6	4	2	2	1	2	-	-	-
1999.....	13	11	9	2	3	1	3	-	-	1	-	-
All races, female												
2004.....	13	13	6	4	-	1	-	1	-	-	-	-
2003.....	14	6	5	3	3	2	1	2	2	-	-	-
2002.....	6	4	1	3	4	2	2	2	-	-	-	-
2001.....	2	5	2	2	2	-	1	2	-	-	-	-
2000.....	8	4	1	1	2	1	3	-	-	-	-	-
1999.....	6	6	5	2	1	1	1	1	-	-	-	-
White, both sexes												
2004.....	26	29	11	7	7	6	6	2	1	-	-	-
2003.....	25	11	14	6	3	6	4	1	2	-	-	-
2002.....	18	13	3	6	8	6	4	3	-	-	-	-
2001.....	11	10	5	2	6	1	2	6	-	-	-	-
2000.....	18	7	6	5	4	2	3	1	2	-	-	-
1999.....	18	16	14	3	3	2	4	1	-	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intentional self-poisoning (suicide) by and exposure to other gases and vapors (X67)												
All races, both sexes												
2004.....	184	102	68	25	34	40	28	25	5	-	-	-
2003.....	153	121	63	34	43	31	33	24	7	1	-	-
2002.....	160	116	44	41	35	46	38	28	8	1	-	-
2001.....	153	132	55	46	36	45	35	25	9	-	-	-
2000.....	144	111	52	38	32	37	37	25	7	1	-	-
1999.....	148	90	61	33	34	56	42	38	11	1	-	-
All races, male												
2004.....	144	79	54	19	23	25	22	16	3	-	-	-
2003.....	113	93	49	28	33	20	21	17	5	1	-	-
2002.....	123	91	36	29	24	33	30	21	8	1	-	-
2001.....	112	91	39	34	26	32	29	19	6	-	-	-
2000.....	101	84	38	25	23	30	29	20	7	1	-	-
1999.....	103	65	41	26	23	40	30	35	11	1	-	-
All races, female												
2004.....	40	23	14	6	11	15	6	9	2	-	-	-
2003.....	40	28	14	6	10	11	12	7	2	-	-	-
2002.....	37	25	8	12	11	13	8	7	-	-	-	-
2001.....	41	41	16	12	10	13	6	6	3	-	-	-
2000.....	43	27	14	13	9	7	8	5	-	-	-	-
1999.....	45	25	20	7	11	16	12	3	-	-	-	-
White, both sexes												
2004.....	178	99	64	25	34	39	27	25	5	-	-	-
2003.....	148	119	56	32	43	31	33	24	7	1	-	-
2002.....	156	111	44	40	35	44	38	27	8	1	-	-
2001.....	148	126	53	46	35	44	35	25	9	-	-	-
2000.....	141	109	52	36	32	36	36	25	7	1	-	-
1999.....	145	89	60	32	34	56	42	38	11	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Intentional self-harm (suicide) by hanging, strangulation and suffocation (X70)

All races, both sexes

2004.....	497	369	206	131	98	109	97	63	34	11	1	5
2003.....	460	327	168	120	107	109	103	65	29	5	1	1
2002.....	444	279	168	135	115	109	94	71	25	8	1	6
2001.....	388	241	151	127	111	103	109	62	29	2	-	-
2000.....	336	203	140	105	106	127	99	76	35	9	3	-
1999.....	314	189	139	136	113	129	99	75	34	9	-	4

All races, male

2004.....	391	292	163	104	76	79	72	39	25	7	1	5
2003.....	370	269	128	95	88	81	77	49	21	2	-	1
2002.....	346	236	132	106	86	79	60	50	18	4	-	6
2001.....	325	196	113	97	93	71	82	43	22	2	-	-
2000.....	277	153	103	64	81	92	69	55	25	5	3	-
1999.....	250	159	107	94	86	87	64	52	22	5	-	4

All races, female

2004.....	106	77	43	27	22	30	25	24	9	4	-	-
2003.....	90	58	40	25	19	28	26	16	8	3	1	-
2002.....	98	43	36	29	29	30	34	21	7	4	1	-
2001.....	63	45	38	30	18	32	27	19	7	-	-	-
2000.....	59	50	37	41	25	35	30	21	10	4	-	-
1999.....	64	30	32	42	27	42	35	23	12	4	-	-

White, both sexes

2004.....	441	332	174	112	81	95	88	59	29	11	1	5
2003.....	417	278	145	94	90	97	95	57	29	3	1	1
2002.....	404	250	155	111	103	96	87	68	19	7	1	6
2001.....	344	215	127	107	96	93	99	54	28	2	-	-
2000.....	294	185	126	92	92	109	90	72	32	8	1	-
1999.....	276	167	117	115	102	112	92	68	34	9	-	4

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intentional self-harm (suicide) by hanging, strangulation and suffocation (X70)												
White, male												
2004.....	353	261	141	90	62	73	65	38	21	7	1	5
2003.....	339	229	111	74	74	74	70	44	21	1	-	1
2002.....	320	215	125	87	82	73	57	48	14	3	-	6
2001.....	291	178	92	81	81	64	77	38	22	2	-	-
2000.....	242	141	95	55	71	79	64	52	22	4	1	-
1999.....	219	141	93	82	78	75	61	47	22	5	-	4
White, female												
2004.....	88	71	33	22	19	22	23	21	8	4	-	-
2003.....	78	49	34	20	16	23	25	13	8	2	1	-
2002.....	84	35	30	24	21	23	30	20	5	4	1	-
2001.....	53	37	35	26	15	29	22	16	6	-	-	-
2000.....	52	44	31	37	21	30	26	20	10	4	-	-
1999.....	57	26	24	33	24	37	31	21	12	4	-	-
Black, both sexes												
2004.....	27	21	10	2	5	1	1	-	2	-	-	-
2003.....	22	20	12	4	5	1	2	1	-	-	-	-
2002.....	17	11	5	7	5	2	1	-	-	-	-	-
2001.....	17	12	7	8	3	1	1	-	-	-	-	-
2000.....	17	7	4	2	4	6	1	1	-	-	1	-
1999.....	14	5	5	9	2	3	-	1	-	-	-	-
Black, male												
2004.....	21	18	9	2	5	1	1	-	2	-	-	-
2003.....	19	18	9	3	3	1	2	1	-	-	-	-
2002.....	15	9	4	7	2	2	1	-	-	-	-	-
2001.....	15	9	7	7	3	1	1	-	-	-	-	-
2000.....	17	4	1	2	2	5	1	1	-	-	1	-
1999.....	13	4	3	8	2	3	-	1	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intentional self-harm (suicide) by hanging, strangulation and suffocation (X70)												
Asian or Pacific Islander 2/, both sexes												
2004.....	27	16	22	16	12	13	8	4	3	-	-	-
2003.....	17	29	11	21	12	11	6	7	-	2	-	-
2002.....	21	16	8	17	7	11	6	3	6	1	-	-
2001.....	17	13	15	11	12	9	8	8	1	-	-	-
2000.....	22	9	10	11	9	12	8	3	3	1	1	-
1999.....	23	13	17	12	8	14	7	6	-	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	15	13	13	11	9	5	6	1	2	-	-	-
2003.....	9	22	8	17	11	6	5	4	-	1	-	-
2002.....	9	11	3	12	2	4	2	2	4	1	-	-
2001.....	9	8	12	8	9	6	3	5	-	-	-	-
2000.....	15	6	7	7	7	8	4	2	3	1	1	-
1999.....	17	11	11	4	5	9	3	4	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	12	3	9	5	3	8	2	3	1	-	-	-
2003.....	8	7	3	4	1	5	1	3	-	1	-	-
2002.....	12	5	5	5	5	7	4	1	2	-	-	-
2001.....	8	5	3	3	3	3	5	3	1	-	-	-
2000.....	7	3	3	4	2	4	4	1	-	-	-	-
1999.....	6	2	6	8	3	5	4	2	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Intentional self-harm (suicide) by discharge of firearms (X72-X74)

All races, both sexes

2004.....	1,565	1,283	1,045	794	837	903	685	387	128	18	4	3
2003.....	1,574	1,329	988	833	867	886	709	399	136	24	-	-
2002.....	1,484	1,295	939	846	930	918	762	392	133	25	-	2
2001.....	1,471	1,195	888	857	901	920	728	382	135	18	2	6
2000.....	1,367	1,001	832	791	909	917	694	410	129	18	1	5
1999.....	1,215	1,058	802	827	964	946	660	399	102	23	-	5

All races, male

2004.....	1,296	1,101	893	717	758	828	651	371	126	17	4	3
2003.....	1,349	1,122	879	747	791	815	665	375	132	22	-	-
2002.....	1,283	1,115	815	753	867	854	722	374	127	24	-	2
2001.....	1,242	1,020	751	761	831	846	685	360	126	18	2	6
2000.....	1,146	854	714	698	830	856	649	388	119	18	1	5
1999.....	1,030	882	712	742	873	875	624	376	99	21	-	5

All races, female

2004.....	269	182	152	77	79	75	34	16	2	1	-	-
2003.....	225	207	109	86	76	71	44	24	4	2	-	-
2002.....	201	180	124	93	63	64	40	18	6	1	-	-
2001.....	229	175	137	96	70	74	43	22	9	-	-	-
2000.....	221	147	118	93	79	61	45	22	10	-	-	-
1999.....	185	176	90	85	91	71	36	23	3	2	-	-

White, both sexes

2004.....	1,497	1,215	1,005	755	800	871	663	381	125	18	4	2
2003.....	1,487	1,276	945	797	835	856	687	391	136	23	-	-
2002.....	1,395	1,238	897	817	894	878	747	380	131	23	-	2
2001.....	1,397	1,137	851	820	860	889	700	374	131	18	2	5
2000.....	1,290	941	791	759	867	894	679	403	126	17	1	4
1999.....	1,154	998	756	785	929	917	640	390	100	22	-	5

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intentional self-harm (suicide) by discharge of firearms (X72-X74)												
White, male												
2004.....	1,242	1,040	861	680	723	797	630	365	124	17	4	2
2003.....	1,275	1,079	839	717	762	787	646	368	132	22	-	-
2002.....	1,208	1,062	778	726	836	816	707	362	125	22	-	2
2001.....	1,181	969	717	731	794	818	662	353	122	18	2	5
2000.....	1,083	800	679	667	790	836	637	381	116	17	1	4
1999.....	974	827	672	707	838	849	605	368	97	20	-	5
White, female												
2004.....	255	175	144	75	77	74	33	16	1	1	-	-
2003.....	212	197	106	80	73	69	41	23	4	1	-	-
2002.....	187	176	119	91	58	62	40	18	6	1	-	-
2001.....	216	168	134	89	66	71	38	21	9	-	-	-
2000.....	207	141	112	92	77	58	42	22	10	-	-	-
1999.....	180	171	84	78	91	68	35	22	3	2	-	-
Black, both sexes												
2004.....	48	55	32	30	31	29	19	5	3	-	-	-
2003.....	69	39	36	32	24	23	16	6	-	1	-	-
2002.....	60	40	33	22	34	31	11	12	2	2	-	-
2001.....	60	43	27	27	35	26	23	6	4	-	-	-
2000.....	61	39	33	28	34	19	13	6	2	-	-	-
1999.....	43	46	36	36	29	28	15	9	2	1	-	-
Black, male												
2004.....	41	49	26	29	31	28	18	5	2	-	-	-
2003.....	59	30	33	26	21	21	14	6	-	-	-	-
2002.....	51	37	28	20	30	29	11	12	2	2	-	-
2001.....	50	39	25	22	31	23	19	5	4	-	-	-
2000.....	48	35	27	27	33	17	10	6	2	-	-	-
1999.....	40	42	32	30	29	25	14	8	2	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Intentional self-harm (suicide) by jumping from a high place (X80)

All races, both sexes

2004.....	59	43	33	24	25	19	21	17	7	-	-	1
2003.....	60	54	32	19	22	18	18	14	12	2	-	1
2002.....	58	46	29	16	24	19	25	14	12	1	-	-
2001.....	61	44	29	23	27	29	18	8	6	1	-	1
2000.....	51	28	22	13	22	19	24	16	6	2	-	-
1999.....	46	30	19	25	27	24	27	17	4	1	-	2

All races, male

2004.....	41	33	24	18	19	11	12	9	2	-	-	1
2003.....	37	35	20	13	15	11	12	10	6	2	-	1
2002.....	35	33	21	7	17	11	14	9	7	1	-	-
2001.....	35	32	21	13	16	22	8	7	3	-	-	-
2000.....	38	17	12	10	12	11	16	8	5	1	-	-
1999.....	31	18	9	16	21	14	18	11	3	-	-	2

All races, female

2004.....	18	10	9	6	6	8	9	8	5	-	-	-
2003.....	23	19	12	6	7	7	6	4	6	-	-	-
2002.....	23	13	8	9	7	8	11	5	5	-	-	-
2001.....	26	12	8	10	11	7	10	1	3	1	-	1
2000.....	13	11	10	3	10	8	8	8	1	1	-	-
1999.....	15	12	10	9	6	10	9	6	1	1	-	-

White, both sexes

2004.....	52	39	28	22	19	17	17	14	5	-	-	1
2003.....	53	48	31	17	19	14	17	10	11	2	-	1
2002.....	52	39	23	14	21	14	25	12	12	1	-	-
2001.....	52	37	26	22	24	25	17	7	6	1	-	1
2000.....	44	24	20	13	19	15	21	13	6	2	-	-
1999.....	40	25	16	22	24	22	22	16	4	1	-	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intentional self-harm (suicide) by jumping from a high place (X80)												
White, male												
2004.....	36	29	20	17	14	10	10	8	1	-	-	1
2003.....	34	31	19	11	12	9	12	7	5	2	-	1
2002.....	33	27	18	6	14	9	14	9	7	1	-	-
2001.....	31	28	18	13	13	18	7	6	3	-	-	-
2000.....	32	16	11	10	10	8	15	6	5	1	-	-
1999.....	26	15	7	15	19	14	16	10	3	-	-	1
White, female												
2004.....	16	10	8	5	5	7	7	6	4	-	-	-
2003.....	19	17	12	6	7	5	5	3	6	-	-	-
2002.....	19	12	5	8	7	5	11	3	5	-	-	-
2001.....	21	9	8	9	11	7	10	1	3	1	-	1
2000.....	12	8	9	3	9	7	6	7	1	1	-	-
1999.....	14	10	9	7	5	8	6	6	1	1	-	-
Black, both sexes												
2004.....	4	2	3	-	3	-	1	-	1	-	-	-
2003.....	4	3	-	-	1	2	-	-	-	-	-	-
2002.....	2	1	5	-	-	4	-	-	-	-	-	-
2001.....	7	5	3	-	3	1	-	-	-	-	-	-
2000.....	3	2	1	-	1	2	2	2	-	-	-	-
1999.....	4	3	2	1	2	1	1	-	-	-	-	-
Black, male												
2004.....	3	2	3	-	2	-	1	-	-	-	-	-
2003.....	1	2	-	-	1	2	-	-	-	-	-	-
2002.....	1	1	3	-	-	2	-	-	-	-	-	-
2001.....	4	4	3	-	3	1	-	-	-	-	-	-
2000.....	3	1	1	-	-	1	-	1	-	-	-	-
1999.....	3	1	2	-	1	-	1	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intentional self-harm (suicide) by jumping from a high place (X80)												
Asian or Pacific Islander 2/, both sexes												
2004.....	3	2	2	2	3	2	3	3	1	-	-	-
2003.....	3	3	-	2	2	2	1	4	1	-	-	-
2002.....	4	5	1	2	3	1	-	2	-	-	-	-
2001.....	2	2	-	1	-	3	1	1	-	-	-	-
2000.....	3	2	1	-	2	2	1	1	-	-	-	-
1999.....	2	2	1	2	1	1	4	1	-	-	-	1
Asian or Pacific Islander 2/, male												
2004.....	2	2	1	1	3	1	1	1	1	-	-	-
2003.....	2	2	-	2	2	-	-	3	1	-	-	-
2002.....	1	4	-	1	3	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	3	1	1	-	-	-	-
2000.....	2	-	-	-	2	2	1	1	-	-	-	-
1999.....	2	2	-	1	1	-	1	1	-	-	-	1
Asian or Pacific Islander 2/, female												
2004.....	1	-	1	1	-	1	2	2	-	-	-	-
2003.....	1	1	-	-	-	2	1	1	-	-	-	-
2002.....	3	1	1	1	-	1	-	2	-	-	-	-
2001.....	2	2	-	1	-	-	-	-	-	-	-	-
2000.....	1	2	1	-	-	-	-	-	-	-	-	-
1999.....	-	-	1	1	-	1	3	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Intentional self-harm (suicide) by all other and unspecified means and their sequelae (*U03,X71,X75-X79,X81-X84,X87.0)

All races, both sexes

2004.....	224	136	95	67	68	39	46	29	11	4	-	2
2003.....	194	141	93	46	54	41	41	19	11	2	-	5
2002.....	189	136	85	57	68	59	48	39	21	6	-	2
2001 3/.....	162	119	72	56	59	56	49	30	10	4	1	5
2000.....	134	117	71	57	60	63	50	32	17	1	-	2
1999.....	152	82	66	66	61	45	52	32	19	2	-	4

All races, male

2004.....	167	96	66	46	48	27	30	22	8	3	-	2
2003.....	141	100	62	32	39	33	31	14	9	1	-	5
2002.....	147	97	55	38	46	40	30	26	14	3	-	2
2001 3/.....	117	85	57	39	37	36	35	23	6	3	1	5
2000.....	92	88	48	43	38	44	33	23	7	-	-	2
1999.....	111	61	44	42	38	34	34	20	9	1	-	4

All races, female

2004.....	57	40	29	21	20	12	16	7	3	1	-	-
2003.....	53	41	31	14	15	8	10	5	2	1	-	-
2002.....	42	39	30	19	22	19	18	13	7	3	-	-
2001 3/.....	45	34	15	17	22	20	14	7	4	1	-	-
2000.....	42	29	23	14	22	19	17	9	10	1	-	-
1999.....	41	21	22	24	23	11	18	12	10	1	-	-

White, both sexes

2004.....	200	117	82	63	58	34	37	27	9	4	-	1
2003.....	174	134	84	43	47	35	39	17	11	2	-	5
2002.....	165	119	75	51	58	55	44	35	21	6	-	2
2001 3/.....	135	108	62	45	53	53	47	28	10	4	1	5
2000.....	113	106	64	53	54	58	44	30	16	1	-	2
1999.....	133	71	56	56	53	43	50	32	17	2	-	2

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intentional self-harm (suicide) by all other and unspecified means and their sequelae (*U03,X71,X75-X79,X81-X84,X87.0)												
White, male												
2004.....	149	79	58	43	45	25	23	20	6	3	-	1
2003.....	129	97	59	31	36	29	30	14	9	1	-	5
2002.....	130	86	51	34	42	37	26	24	14	3	-	2
2001 3/.....	99	78	50	31	34	34	33	23	6	3	1	5
2000.....	75	80	43	40	34	40	30	23	7	-	-	2
1999.....	98	54	38	34	34	32	33	20	9	1	-	2
White, female												
2004.....	51	38	24	20	13	9	14	7	3	1	-	-
2003.....	45	37	25	12	11	6	9	3	2	1	-	-
2002.....	35	33	24	17	16	18	18	11	7	3	-	-
2001 3/.....	36	30	12	14	19	19	14	5	4	1	-	-
2000.....	38	26	21	13	20	18	14	7	9	1	-	-
1999.....	35	17	18	22	19	11	17	12	8	1	-	-
Black, both sexes												
2004.....	16	13	5	1	2	2	4	-	-	-	-	1
2003.....	14	5	7	2	3	-	-	2	-	-	-	-
2002.....	20	12	7	2	6	1	1	3	-	-	-	-
2001 3/.....	22	8	6	4	5	1	-	2	-	-	-	-
2000.....	12	7	3	3	5	-	3	-	-	-	-	-
1999.....	11	9	4	8	5	2	1	-	2	-	-	2
Black, male												
2004.....	13	13	4	1	-	1	2	-	-	-	-	1
2003.....	8	3	3	1	1	-	-	-	-	-	-	-
2002.....	15	8	3	2	3	1	1	1	-	-	-	-
2001 3/.....	15	4	3	2	3	-	-	-	-	-	-	-
2000.....	9	6	2	3	4	-	1	-	-	-	-	-
1999.....	8	5	2	6	1	2	-	-	-	-	-	2

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Assault (homicide) (*U01-*U02,X85-Y09,X87.1)												
All races, both sexes												
2004.....	826	523	356	260	186	164	125	73	26	4	1	36
2003.....	798	490	296	227	205	187	130	77	22	5	-	41
2002.....	760	523	318	229	192	169	127	64	24	6	1	41
2001 3/.....	1,004	628	390	310	222	183	129	71	29	4	1	32
2000.....	696	455	283	233	210	172	129	68	23	8	1	30
1999.....	670	420	302	259	214	185	119	65	24	10	-	49
All races, male												
2004.....	641	389	248	177	125	81	57	29	11	1	-	29
2003.....	595	352	216	156	129	110	61	32	13	1	-	29
2002.....	553	383	231	152	125	100	58	32	6	3	-	32
2001 3/.....	747	461	262	204	128	98	57	33	7	1	1	23
2000.....	511	338	199	147	132	85	69	25	10	5	-	15
1999.....	483	323	218	183	121	101	61	28	8	5	-	33
All races, female												
2004.....	185	134	108	83	61	83	68	44	15	3	1	7
2003.....	203	138	80	71	76	77	69	45	9	4	-	12
2002.....	207	140	87	77	67	69	69	32	18	3	1	9
2001 3/.....	257	167	128	106	94	85	72	38	22	3	-	9
2000.....	185	117	84	86	78	87	60	43	13	3	1	15
1999.....	187	97	84	76	93	84	58	37	16	5	-	16
White, both sexes												
2004.....	487	339	243	192	131	117	102	63	16	4	1	34
2003.....	493	327	203	160	144	143	101	65	18	3	-	36
2002.....	453	366	221	160	146	119	99	53	18	5	-	36
2001 3/.....	681	461	283	225	167	139	101	63	22	3	-	24
2000.....	432	309	188	150	153	138	89	57	18	7	1	25
1999.....	429	287	209	193	160	133	92	51	21	7	-	34

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Assault (homicide) (*U01-*U02,X85-Y09,X87.1)												
White, male												
2004.....	369	244	165	119	86	56	43	26	4	1	-	27
2003.....	354	217	147	105	89	82	46	27	11	1	-	25
2002.....	311	254	155	105	88	66	42	28	3	3	-	29
2001 3/.....	496	328	190	142	91	73	45	28	5	-	-	16
2000.....	295	224	126	86	90	67	42	20	7	4	-	13
1999.....	308	210	146	130	86	70	47	19	6	4	-	23
White, female												
2004.....	118	95	78	73	45	61	59	37	12	3	1	7
2003.....	139	110	56	55	55	61	55	38	7	2	-	11
2002.....	142	112	66	55	58	53	57	25	15	2	-	7
2001 3/.....	185	133	93	83	76	66	56	35	17	3	-	8
2000.....	137	85	62	64	63	71	47	37	11	3	1	12
1999.....	121	77	63	63	74	63	45	32	15	3	-	11
Black, both sexes												
2004.....	299	158	96	55	49	42	21	9	9	-	-	2
2003.....	274	143	80	54	52	39	26	11	4	2	-	4
2002.....	273	132	86	55	42	48	23	11	5	1	-	3
2001 3/.....	274	141	84	71	48	40	23	7	6	1	1	8
2000.....	231	123	81	74	51	30	34	8	5	1	-	4
1999.....	198	116	79	60	44	46	21	13	3	3	-	14
Black, male												
2004.....	243	125	70	48	35	23	14	3	7	-	-	2
2003.....	216	123	59	44	35	27	14	5	2	-	-	3
2002.....	217	113	68	40	34	33	15	4	2	-	-	2
2001 3/.....	213	111	59	52	35	25	9	4	2	1	1	7
2000.....	193	97	63	58	37	16	23	3	3	1	-	1
1999.....	150	100	60	50	30	29	10	8	2	1	-	10

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Assault (homicide) (*U01-*U02,X85-Y09,X87.1)												
Black, female												
2004.....	56	33	26	7	14	19	7	6	2	-	-	-
2003.....	58	20	21	10	17	12	12	6	2	2	-	1
2002.....	56	19	18	15	8	15	8	7	3	1	-	1
2001 3/.....	61	30	25	19	13	15	14	3	4	-	-	1
2000.....	38	26	18	16	14	14	11	5	2	-	-	3
1999.....	48	16	19	10	14	17	11	5	1	2	-	4
American Indian 1/, both sexes												
2004.....	7	4	4	6	-	2	1	-	-	-	-	-
2003.....	9	3	1	3	2	1	-	1	-	-	-	-
2002.....	14	7	2	6	1	-	3	-	-	-	-	1
2001 3/.....	13	3	2	5	1	-	2	-	1	-	-	-
2000.....	9	2	4	2	2	2	-	2	-	-	-	1
1999.....	18	8	3	3	4	2	3	-	-	-	-	-
American Indian 1/, male												
2004.....	6	2	3	5	-	1	-	-	-	-	-	-
2003.....	8	1	1	2	2	-	-	-	-	-	-	-
2002.....	10	3	2	3	1	-	1	-	-	-	-	-
2001 3/.....	10	3	2	3	1	-	1	-	-	-	-	-
2000.....	7	1	3	1	2	1	-	2	-	-	-	1
1999.....	13	6	3	3	3	1	3	-	-	-	-	-
American Indian 1/, female												
2004.....	1	2	1	1	-	1	1	-	-	-	-	-
2003.....	1	2	-	1	-	1	-	1	-	-	-	-
2002.....	4	4	-	3	-	-	2	-	-	-	-	1
2001 3/.....	3	-	-	2	-	-	1	-	1	-	-	-
2000.....	2	1	1	1	-	1	-	-	-	-	-	-
1999.....	5	2	-	-	1	1	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Assault (homicide) (*U01-*U02,X85-Y09,X87.1)												
Asian or Pacific Islander 2/, both sexes												
2004.....	33	22	13	7	6	3	1	1	1	-	-	-
2003.....	22	17	12	10	7	4	3	-	-	-	-	1
2002.....	20	18	9	8	3	2	2	-	1	-	1	1
2001 3/.....	36	23	21	9	6	4	3	1	-	-	-	-
2000.....	24	21	10	7	4	2	6	1	-	-	-	-
1999.....	25	9	11	3	6	4	3	1	-	-	-	1
Asian or Pacific Islander 2/, male												
2004.....	23	18	10	5	4	1	-	-	-	-	-	-
2003.....	17	11	9	5	3	1	1	-	-	-	-	1
2002.....	15	13	6	4	2	1	-	-	1	-	-	1
2001 3/.....	28	19	11	7	1	-	2	1	-	-	-	-
2000.....	16	16	7	2	3	1	4	-	-	-	-	-
1999.....	12	7	9	-	2	1	1	1	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	10	4	3	2	2	2	1	1	1	-	-	-
2003.....	5	6	3	5	4	3	2	-	-	-	-	-
2002.....	5	5	3	4	1	1	2	-	-	-	1	-
2001 3/.....	8	4	10	2	5	4	1	-	-	-	-	-
2000.....	8	5	3	5	1	1	2	1	-	-	-	-
1999.....	13	2	2	3	4	3	2	-	-	-	-	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Assault (homicide) by drugs, biological substances and other and unspecified noxious substances (*U01.6-*U01.7,X85-X90)

All races, both sexes

2004.....	-	2	4	2	-	2	1	2	2	-	-	-
2003.....	4	1	-	1	1	2	3	2	-	1	-	-
2002.....	5	4	1	2	-	1	3	3	-	-	-	-
2001.....	3	3	2	3	2	-	-	1	2	-	-	-
2000.....	3	1	3	2	2	2	1	1	-	-	-	-
1999.....	4	1	1	2	3	-	2	-	1	-	-	-

All races, male

2004.....	-	2	4	2	-	-	-	1	-	-	-	-
2003.....	3	1	-	-	1	2	2	-	-	-	-	-
2002.....	4	3	1	-	-	1	-	2	-	-	-	-
2001.....	3	2	1	1	1	-	-	1	-	-	-	-
2000.....	2	1	1	-	-	-	-	-	-	-	-	-
1999.....	3	1	-	-	1	-	-	-	-	-	-	-

All races, female

2004.....	-	-	-	-	-	2	1	1	2	-	-	-
2003.....	1	-	-	1	-	-	1	2	-	1	-	-
2002.....	1	1	-	2	-	-	3	1	-	-	-	-
2001.....	-	1	1	2	1	-	-	-	2	-	-	-
2000.....	1	-	2	2	2	2	1	1	-	-	-	-
1999.....	1	-	1	2	2	-	2	-	1	-	-	-

White, both sexes

2004.....	-	2	4	1	-	1	1	-	2	-	-	-
2003.....	1	-	-	1	1	1	2	1	-	1	-	-
2002.....	4	4	1	2	-	1	2	3	-	-	-	-
2001.....	3	2	2	3	2	-	-	1	2	-	-	-
2000.....	2	1	1	2	2	2	1	1	-	-	-	-
1999.....	3	1	1	2	3	-	2	-	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Assault (homicide) by hanging, strangulation and suffocation (X91)												
All races, both sexes												
2004.....	33	21	17	16	16	12	13	4	-	-	-	-
2003.....	39	23	15	17	15	16	18	5	1	-	-	4
2002.....	36	19	20	14	14	7	7	5	2	1	-	2
2001.....	31	21	24	20	27	16	13	4	7	-	-	4
2000.....	25	23	15	19	13	15	9	6	4	-	-	3
1999.....	28	17	9	29	16	19	8	9	3	-	-	7
All races, male												
2004.....	17	8	9	3	7	5	4	1	-	-	-	-
2003.....	14	9	4	5	6	8	8	2	-	-	-	2
2002.....	12	8	13	9	5	3	-	3	-	-	-	2
2001.....	12	14	10	11	12	7	1	1	3	-	-	1
2000.....	8	14	6	7	6	5	3	2	-	-	-	1
1999.....	12	11	2	18	4	5	2	4	1	-	-	3
All races, female												
2004.....	16	13	8	13	9	7	9	3	-	-	-	-
2003.....	25	14	11	12	9	8	10	3	1	-	-	2
2002.....	24	11	7	5	9	4	7	2	2	1	-	-
2001.....	19	7	14	9	15	9	12	3	4	-	-	3
2000.....	17	9	9	12	7	10	6	4	4	-	-	2
1999.....	16	6	7	11	12	14	6	5	2	-	-	4
White, both sexes												
2004.....	22	13	14	14	12	10	11	4	-	-	-	-
2003.....	24	19	10	13	7	13	12	4	-	-	-	4
2002.....	27	13	17	11	11	4	6	4	2	-	-	2
2001.....	24	15	15	16	20	9	10	4	5	-	-	3
2000.....	20	15	12	12	8	13	6	5	3	-	-	1
1999.....	20	12	4	22	12	15	6	8	3	-	-	5

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Assault (homicide) by hanging, strangulation and suffocation (X91)												
White, male												
2004.....	13	6	7	2	6	4	3	1	-	-	-	-
2003.....	8	7	3	4	3	6	4	2	-	-	-	2
2002.....	9	6	11	8	4	-	-	3	-	-	-	2
2001.....	11	10	9	9	8	5	1	1	2	-	-	-
2000.....	7	9	6	4	4	5	3	2	-	-	-	1
1999.....	10	8	2	13	4	4	2	3	1	-	-	2
White, female												
2004.....	9	7	7	12	6	6	8	3	-	-	-	-
2003.....	16	12	7	9	4	7	8	2	-	-	-	2
2002.....	18	7	6	3	7	4	6	1	2	-	-	-
2001.....	13	5	6	7	12	4	9	3	3	-	-	3
2000.....	13	6	6	8	4	8	3	3	3	-	-	-
1999.....	10	4	2	9	8	11	4	5	2	-	-	3
Black, both sexes												
2004.....	8	7	3	2	4	2	2	-	-	-	-	-
2003.....	15	3	3	3	5	3	5	-	1	-	-	-
2002.....	8	5	2	3	3	3	1	1	-	1	-	-
2001.....	7	6	7	3	7	6	2	-	1	-	-	1
2000.....	3	7	2	7	5	1	3	1	1	-	-	2
1999.....	8	5	5	7	3	3	2	1	-	-	-	2
Black, male												
2004.....	3	2	2	1	1	1	1	-	-	-	-	-
2003.....	6	2	1	1	1	2	3	-	-	-	-	-
2002.....	3	2	2	1	1	3	-	-	-	-	-	-
2001.....	1	4	1	1	4	2	-	-	1	-	-	1
2000.....	-	4	-	3	2	-	-	-	-	-	-	-
1999.....	2	3	-	5	-	1	-	1	-	-	-	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Assault (homicide) by hanging, strangulation and suffocation (X91)												
Asian or Pacific Islander 2/, both sexes												
2004.....	3	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	1	2	1	2	-	1	-	-	-	-	-
2002.....	1	-	1	-	-	-	-	-	-	-	-	-
2001.....	-	-	2	1	-	1	-	-	-	-	-	-
2000.....	2	1	1	-	-	1	-	-	-	-	-	-
1999.....	-	-	-	-	1	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	1	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	1	-	1	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	1	-	-	-	-	-	-	-	-
2000.....	1	1	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	2	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	1	2	1	1	-	-	-	-	-	-	-
2002.....	1	-	1	-	-	-	-	-	-	-	-	-
2001.....	-	-	2	-	-	1	-	-	-	-	-	-
2000.....	1	-	1	-	-	1	-	-	-	-	-	-
1999.....	-	-	-	-	1	1	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)												
All races, both sexes												
2004.....	424	279	189	122	71	62	38	22	5	2	-	21
2003.....	427	250	144	83	78	55	33	17	5	1	-	18
2002.....	421	263	146	92	69	72	43	14	4	1	-	13
2001.....	382	213	151	114	78	61	38	12	2	2	-	11
2000.....	377	235	132	99	85	54	27	23	4	1	-	12
1999.....	342	206	167	102	91	61	34	17	4	2	-	15
All races, male												
2004.....	336	216	131	83	55	31	10	11	3	1	-	20
2003.....	326	186	106	63	52	27	8	9	1	-	-	16
2002.....	316	197	104	61	40	44	20	8	-	1	-	12
2001.....	292	159	105	79	44	38	16	6	-	1	-	11
2000.....	285	172	94	58	50	23	14	6	2	1	-	7
1999.....	250	167	127	68	56	34	14	7	2	1	-	14
All races, female												
2004.....	88	63	58	39	16	31	28	11	2	1	-	1
2003.....	101	64	38	20	26	28	25	8	4	1	-	2
2002.....	105	66	42	31	29	28	23	6	4	-	-	1
2001.....	90	54	46	35	34	23	22	6	2	1	-	-
2000.....	92	63	38	41	35	31	13	17	2	-	-	5
1999.....	92	39	40	34	35	27	20	10	2	1	-	1
White, both sexes												
2004.....	245	177	134	93	51	49	35	20	2	2	-	20
2003.....	265	168	100	58	58	41	27	17	3	-	-	14
2002.....	253	190	94	62	54	55	36	13	3	1	-	11
2001.....	236	142	113	84	57	47	31	12	2	1	-	9
2000.....	229	165	95	70	64	47	17	19	4	1	-	10
1999.....	213	139	113	79	69	43	29	16	4	1	-	10

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)												
White, male												
2004.....	182	129	90	56	35	25	8	10	-	1	-	19
2003.....	196	116	72	43	38	20	5	9	-	-	-	13
2002.....	183	134	66	38	27	32	15	8	-	1	-	10
2001.....	173	94	76	51	29	27	10	6	-	-	-	9
2000.....	160	117	60	37	33	20	5	5	2	1	-	5
1999.....	162	106	80	50	40	20	11	6	2	1	-	9
White, female												
2004.....	63	48	44	37	16	24	27	10	2	1	-	1
2003.....	69	52	28	15	20	21	22	8	3	-	-	1
2002.....	70	56	28	24	27	23	21	5	3	-	-	1
2001.....	63	48	37	33	28	20	21	6	2	1	-	-
2000.....	69	48	35	33	31	27	12	14	2	-	-	5
1999.....	51	33	33	29	29	23	18	10	2	-	-	1
Black, both sexes												
2004.....	161	87	51	25	16	11	3	2	3	-	-	1
2003.....	147	72	37	23	19	11	6	-	2	1	-	3
2002.....	150	63	46	26	14	15	6	1	1	-	-	2
2001.....	130	61	33	27	18	13	5	-	-	1	-	2
2000.....	131	56	29	25	20	7	10	4	-	-	-	1
1999.....	104	59	45	20	19	17	4	1	-	1	-	5
Black, male												
2004.....	140	73	38	23	16	5	2	1	3	-	-	1
2003.....	118	65	27	19	14	7	3	-	1	-	-	2
2002.....	118	56	33	21	12	11	5	-	-	-	-	2
2001.....	109	55	25	25	14	11	4	-	-	1	-	2
2000.....	113	43	26	20	17	3	9	1	-	-	-	1
1999.....	77	55	39	17	14	14	2	1	-	-	-	5

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)												
Asian or Pacific Islander 2/, both sexes												
2004.....	17	14	3	4	4	1	-	-	-	-	-	-
2003.....	14	8	6	2	1	2	-	-	-	-	-	1
2002.....	13	9	5	3	-	2	1	-	-	-	-	-
2001.....	14	8	4	1	2	1	2	-	-	-	-	-
2000.....	15	14	6	3	1	-	-	-	-	-	-	-
1999.....	17	6	7	2	1	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	13	13	2	4	4	-	-	-	-	-	-	-
2003.....	11	5	6	1	-	-	-	-	-	-	-	1
2002.....	10	6	4	2	-	1	-	-	-	-	-	-
2001.....	9	8	3	1	-	-	2	-	-	-	-	-
2000.....	11	12	6	1	-	-	-	-	-	-	-	-
1999.....	6	4	6	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	4	1	1	-	-	1	-	-	-	-	-	-
2003.....	3	3	-	1	1	2	-	-	-	-	-	-
2002.....	3	3	1	1	-	1	1	-	-	-	-	-
2001.....	5	-	1	-	2	1	-	-	-	-	-	-
2000.....	4	2	-	2	1	-	-	-	-	-	-	-
1999.....	11	2	1	2	1	1	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Assault (homicide) by sharp object (X99)												
All races, both sexes												
2004.....	129	75	55	48	29	31	25	12	5	-	-	2
2003.....	128	72	44	36	42	34	20	13	1	-	-	5
2002.....	120	96	42	37	45	24	22	13	2	2	-	5
2001.....	106	74	38	56	23	35	17	6	5	1	-	3
2000.....	86	62	39	35	27	33	26	7	3	2	-	4
1999.....	101	65	42	33	30	28	13	8	4	1	-	4
All races, male												
2004.....	100	54	37	35	18	17	14	2	2	-	-	2
2003.....	94	45	31	17	27	21	9	3	-	-	-	4
2002.....	88	69	28	24	30	12	8	9	-	1	-	3
2001.....	78	54	18	30	11	13	7	4	2	-	-	3
2000.....	54	52	25	22	22	20	16	2	3	2	-	3
1999.....	72	47	28	23	15	17	6	4	2	1	-	2
All races, female												
2004.....	29	21	18	13	11	14	11	10	3	-	-	-
2003.....	34	27	13	19	15	13	11	10	1	-	-	1
2002.....	32	27	14	13	15	12	14	4	2	1	-	2
2001.....	28	20	20	26	12	22	10	2	3	1	-	-
2000.....	32	10	14	13	5	13	10	5	-	-	-	1
1999.....	29	18	14	10	15	11	7	4	2	-	-	2
White, both sexes												
2004.....	66	45	32	31	16	14	18	8	1	-	-	2
2003.....	71	45	27	27	26	21	14	8	1	-	-	5
2002.....	60	53	30	27	32	15	14	11	1	2	-	5
2001.....	66	48	24	38	17	26	12	6	3	1	-	1
2000.....	54	37	24	19	17	23	19	7	2	1	-	4
1999.....	62	40	30	22	20	14	9	4	3	1	-	3

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Assault (homicide) by sharp object (X99)												
White, male												
2004.....	52	29	21	21	12	6	11	1	-	-	-	2
2003.....	49	25	19	12	15	12	8	1	-	-	-	4
2002.....	41	35	18	19	19	10	5	8	-	1	-	3
2001.....	44	34	13	20	9	10	7	4	1	-	-	1
2000.....	31	33	16	11	15	13	12	2	2	1	-	3
1999.....	45	24	19	15	9	7	5	2	1	1	-	1
White, female												
2004.....	14	16	11	10	4	8	7	7	1	-	-	-
2003.....	22	20	8	15	11	9	6	7	1	-	-	1
2002.....	19	18	12	8	13	5	9	3	1	1	-	2
2001.....	22	14	11	18	8	16	5	2	2	1	-	-
2000.....	23	4	8	8	2	10	7	5	-	-	-	1
1999.....	17	16	11	7	11	7	4	2	2	-	-	2
Black, both sexes												
2004.....	58	27	19	15	12	17	7	4	4	-	-	-
2003.....	50	25	14	7	14	12	4	5	-	-	-	-
2002.....	54	36	11	9	12	9	8	2	1	-	-	-
2001.....	35	24	9	15	5	9	5	-	2	-	-	2
2000.....	29	24	14	16	8	7	4	-	1	1	-	-
1999.....	34	21	12	11	7	13	3	4	1	-	-	1
Black, male												
2004.....	43	23	13	13	6	11	3	1	2	-	-	-
2003.....	41	20	9	5	10	9	1	2	-	-	-	-
2002.....	45	29	9	5	10	2	3	1	-	-	-	-
2001.....	30	18	4	9	2	3	-	-	1	-	-	2
2000.....	21	18	9	11	5	5	2	-	1	1	-	-
1999.....	25	19	9	8	5	9	-	2	1	-	-	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Assault (homicide) by sharp object (X99)												
Black, female												
2004.....	15	4	6	2	6	6	4	3	2	-	-	-
2003.....	9	5	5	2	4	3	3	3	-	-	-	-
2002.....	9	7	2	4	2	7	5	1	1	-	-	-
2001.....	5	6	5	6	3	6	5	-	1	-	-	-
2000.....	8	6	5	5	3	2	2	-	-	-	-	-
1999.....	9	2	3	3	2	4	3	2	-	-	-	-
American Indian 1/, both sexes												
2004.....	2	-	-	1	-	-	-	-	-	-	-	-
2003.....	2	-	-	-	1	-	-	-	-	-	-	-
2002.....	4	2	-	-	-	-	-	-	-	-	-	-
2001.....	2	-	1	1	-	-	-	-	-	-	-	-
2000.....	2	-	1	-	1	2	-	-	-	-	-	-
1999.....	2	2	-	-	1	-	-	-	-	-	-	-
American Indian 1/, male												
2004.....	2	-	-	1	-	-	-	-	-	-	-	-
2003.....	1	-	-	-	1	-	-	-	-	-	-	-
2002.....	1	1	-	-	-	-	-	-	-	-	-	-
2001.....	2	-	1	-	-	-	-	-	-	-	-	-
2000.....	1	-	-	-	1	1	-	-	-	-	-	-
1999.....	1	2	-	-	-	-	-	-	-	-	-	-
American Indian 1/, female												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	1	-	-	-	-	-	-	-	-	-	-	-
2002.....	3	1	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	1	-	-	-	-	-	-	-	-
2000.....	1	-	1	-	-	1	-	-	-	-	-	-
1999.....	1	-	-	-	1	-	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Assault (homicide) by sharp object (X99)												
Asian or Pacific Islander 2/, both sexes												
2004.....	3	3	4	1	1	-	-	-	-	-	-	-
2003.....	5	2	3	2	1	1	2	-	-	-	-	-
2002.....	2	5	1	1	1	-	-	-	-	-	-	-
2001.....	3	2	4	2	1	-	-	-	-	-	-	-
2000.....	1	1	-	-	1	1	3	-	-	-	-	-
1999.....	3	2	-	-	2	1	1	-	-	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	3	2	3	-	-	-	-	-	-	-	-	-
2003.....	3	-	3	-	1	-	-	-	-	-	-	-
2002.....	1	4	1	-	1	-	-	-	-	-	-	-
2001.....	2	2	-	1	-	-	-	-	-	-	-	-
2000.....	1	1	-	-	1	1	2	-	-	-	-	-
1999.....	1	2	-	-	1	1	1	-	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	-	1	1	1	1	-	-	-	-	-	-	-
2003.....	2	2	-	2	-	1	2	-	-	-	-	-
2002.....	1	1	-	1	-	-	-	-	-	-	-	-
2001.....	1	-	4	1	1	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	1	-	-	-	-	-
1999.....	2	-	-	-	1	-	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Assault (homicide) by blunt object (Y00)												
All races, both sexes												
2004.....	11	7	5	4	1	3	5	2	-	-	-	1
2003.....	10	9	5	12	4	4	4	1	-	-	-	-
2002.....	14	10	5	5	3	5	5	2	-	-	-	1
2001.....	18	11	9	8	7	3	3	2	1	-	-	1
2000.....	20	10	9	10	8	2	3	1	-	-	-	-
1999.....	15	16	10	12	9	3	8	1	-	-	-	1
All races, male												
2004.....	6	6	2	4	-	1	2	2	-	-	-	1
2003.....	8	6	2	9	4	3	3	-	-	-	-	-
2002.....	12	6	3	4	3	2	4	1	-	-	-	1
2001.....	15	8	8	5	3	2	3	1	-	-	-	1
2000.....	14	8	6	8	3	2	-	1	-	-	-	-
1999.....	11	11	7	9	7	2	6	-	-	-	-	-
All races, female												
2004.....	5	1	3	-	1	2	3	-	-	-	-	-
2003.....	2	3	3	3	-	1	1	1	-	-	-	-
2002.....	2	4	2	1	-	3	1	1	-	-	-	-
2001.....	3	3	1	3	4	1	-	1	1	-	-	-
2000.....	6	2	3	2	5	-	3	-	-	-	-	-
1999.....	4	5	3	3	2	1	2	1	-	-	-	1
White, both sexes												
2004.....	5	5	3	4	-	3	3	2	-	-	-	1
2003.....	7	3	5	10	3	4	4	1	-	-	-	-
2002.....	10	8	4	4	3	2	5	2	-	-	-	1
2001.....	9	6	5	4	7	3	3	2	1	-	-	1
2000.....	13	5	4	4	7	2	2	1	-	-	-	-
1999.....	13	13	7	10	9	3	5	1	-	-	-	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Assault (homicide) by blunt object (Y00)												
White, male												
2004.....	3	4	2	4	-	1	1	2	-	-	-	1
2003.....	5	1	2	7	3	3	3	-	-	-	-	-
2002.....	8	4	2	3	3	-	4	1	-	-	-	1
2001.....	7	4	4	1	3	2	3	1	-	-	-	1
2000.....	11	4	3	3	2	2	-	1	-	-	-	-
1999.....	9	8	5	7	7	2	4	-	-	-	-	-
White, female												
2004.....	2	1	1	-	-	2	2	-	-	-	-	-
2003.....	2	2	3	3	-	1	1	1	-	-	-	-
2002.....	2	4	2	1	-	2	1	1	-	-	-	-
2001.....	2	2	1	3	4	1	-	1	1	-	-	-
2000.....	2	1	1	1	5	-	2	-	-	-	-	-
1999.....	4	5	2	3	2	1	1	1	-	-	-	1
Black, both sexes												
2004.....	6	2	2	-	1	-	1	-	-	-	-	-
2003.....	2	5	-	2	-	-	-	-	-	-	-	-
2002.....	4	2	1	1	-	3	-	-	-	-	-	-
2001.....	7	4	3	2	-	-	-	-	-	-	-	-
2000.....	6	4	4	5	1	-	1	-	-	-	-	-
1999.....	1	3	3	2	-	-	2	-	-	-	-	-
Black, male												
2004.....	3	2	-	-	-	-	1	-	-	-	-	-
2003.....	2	4	-	2	-	-	-	-	-	-	-	-
2002.....	4	2	1	1	-	2	-	-	-	-	-	-
2001.....	7	3	3	2	-	-	-	-	-	-	-	-
2000.....	3	4	2	5	1	-	-	-	-	-	-	-
1999.....	1	3	2	2	-	-	2	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Assault (homicide) by blunt object (Y00)												
Asian or Pacific Islander 2/, both sexes												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	1	-	-	1	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	1	1	1	2	-	-	-	-	-	-	-	-
2000.....	1	-	1	1	-	-	-	-	-	-	-	-
1999.....	1	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	1	-	-	1	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	1	1	2	-	-	-	-	-	-	-	-
2000.....	-	-	1	-	-	-	-	-	-	-	-	-
1999.....	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	1	-	-	-	-	-	-	-	-	-	-	-
2000.....	1	-	-	1	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	1	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Assault (homicide) by bodily force (Y04)												
All races, both sexes												
2004.....	7	2	-	2	3	1	-	1	-	-	-	-
2003.....	9	6	3	4	-	4	-	1	-	1	-	-
2002.....	10	3	3	2	2	1	1	-	1	-	-	-
2001.....	10	4	4	1	5	5	2	3	1	-	-	-
2000.....	9	7	2	1	3	1	1	1	1	-	-	-
1999.....	13	5	2	5	3	2	-	1	1	-	-	1
All races, male												
2004.....	7	2	-	1	2	-	-	1	-	-	-	-
2003.....	8	5	3	3	-	3	-	1	-	-	-	-
2002.....	9	2	2	2	2	1	1	-	1	-	-	-
2001.....	8	4	3	1	4	3	2	2	-	-	-	-
2000.....	9	6	1	1	-	1	-	-	-	-	-	-
1999.....	11	5	2	4	3	1	-	-	-	-	-	1
All races, female												
2004.....	-	-	-	1	1	1	-	-	-	-	-	-
2003.....	1	1	-	1	-	1	-	-	-	1	-	-
2002.....	1	1	1	-	-	-	-	-	-	-	-	-
2001.....	2	-	1	-	1	2	-	1	1	-	-	-
2000.....	-	1	1	-	3	-	1	1	1	-	-	-
1999.....	2	-	-	1	-	1	-	1	1	-	-	-
White, both sexes												
2004.....	5	2	-	1	1	1	-	1	-	-	-	-
2003.....	5	3	3	2	-	4	-	1	-	-	-	-
2002.....	7	3	3	1	1	-	1	-	-	-	-	-
2001.....	7	3	4	1	4	4	2	2	1	-	-	-
2000.....	5	6	2	-	2	1	1	1	1	-	-	-
1999.....	10	2	2	4	2	2	-	-	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Neglect, abandonment and other maltreatment syndromes (Y06-Y07)												
All races, both sexes												
2004.....	1	2	1	-	1	1	2	3	1	-	1	1
2003.....	3	5	1	-	5	3	1	2	1	-	-	1
2002.....	2	2	5	2	2	3	7	3	-	1	1	-
2001.....	2	1	1	2	5	4	2	4	2	-	-	-
2000.....	5	1	2	-	3	1	-	5	2	1	-	1
1999.....	1	1	-	1	-	3	4	2	-	-	-	-
All races, male												
2004.....	1	2	-	-	1	1	-	1	-	-	-	-
2003.....	2	3	-	-	2	1	-	1	-	-	-	-
2002.....	1	-	-	-	1	1	3	-	-	-	-	-
2001.....	-	-	-	1	-	1	1	-	-	-	-	-
2000.....	2	1	1	-	1	1	-	-	2	1	-	1
1999.....	-	1	-	1	-	-	-	1	-	-	-	-
All races, female												
2004.....	-	-	1	-	-	-	2	2	1	-	1	1
2003.....	1	2	1	-	3	2	1	1	1	-	-	1
2002.....	1	2	5	2	1	2	4	3	-	1	1	-
2001.....	2	1	1	1	5	3	1	4	2	-	-	-
2000.....	3	-	1	-	2	-	-	5	-	-	-	-
1999.....	1	-	-	-	-	3	4	1	-	-	-	-
White, both sexes												
2004.....	1	2	1	-	1	1	1	3	-	-	1	1
2003.....	3	2	-	-	3	2	1	2	1	-	-	1
2002.....	2	2	5	1	2	3	7	3	-	1	-	-
2001.....	2	1	1	2	4	3	2	3	1	-	-	-
2000.....	3	-	2	-	3	1	-	4	-	1	-	1
1999.....	1	-	-	-	-	2	3	1	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Neglect, abandonment and other maltreatment syndromes (Y06-Y07)												
Black, female												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	2	-	-	1	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	1	1	-	1	1	-	-	-
2000.....	1	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	1	1	-	-	-	-
American Indian 1/, both sexes												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	1	-	1	-	-	-	-	-	-
American Indian 1/, male												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, female												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	1	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Assault (homicide) by all other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5,*U01.8-*U01.9,*U02,X92,X96-X98,Y01-Y03,Y08-Y09,Y87.1)

All races, both sexes

2004.....	220	135	85	66	65	52	41	27	13	2	-	11
2003.....	178	124	84	74	60	69	51	36	14	2	-	13
2002.....	152	126	96	74	57	56	39	24	15	1	-	20
2001 3/.....	452	301	161	106	75	59	54	39	9	1	1	13
2000.....	171	116	80	67	69	64	62	24	9	4	1	10
1999.....	166	109	71	75	62	68	50	27	11	7	-	21

All races, male

2004.....	174	99	65	49	42	26	27	10	6	-	-	6
2003.....	140	97	70	59	37	45	31	16	12	1	-	7
2002.....	111	98	80	52	44	36	22	9	5	1	-	14
2001 3/.....	339	220	117	76	53	34	27	18	2	-	1	7
2000.....	137	84	65	51	50	33	36	14	3	1	-	3
1999.....	124	80	52	60	35	42	33	12	3	3	-	13

All races, female

2004.....	46	36	20	17	23	26	14	17	7	2	-	5
2003.....	38	27	14	15	23	24	20	20	2	1	-	6
2002.....	41	28	16	22	13	20	17	15	10	-	-	6
2001 3/.....	113	81	44	30	22	25	27	21	7	1	-	6
2000.....	34	32	15	16	19	31	26	10	6	3	1	7
1999.....	42	29	19	15	27	26	17	15	8	4	-	8

White, both sexes

2004.....	142	93	55	48	50	38	33	25	11	2	-	10
2003.....	117	87	58	49	46	57	41	31	13	2	-	12
2002.....	90	93	67	51	43	39	28	17	12	1	-	17
2001 3/.....	334	244	119	77	56	47	41	33	7	1	-	10
2000.....	106	80	48	43	50	49	43	19	8	4	1	9
1999.....	107	80	52	54	45	54	38	21	9	5	-	15

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Assault (homicide) by all other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5,*U01.8-*U01.9,*U02,X92,X96-X98,Y01-Y03,Y08-Y09,Y87.1)

White, male

2004.....	113	70	41	35	31	19	20	10	4	-	-	5
2003.....	88	64	48	38	27	37	25	13	11	1	-	6
2002.....	60	70	55	36	33	22	14	6	3	1	-	13
2001 3/.....	252	181	84	58	38	26	21	14	2	-	-	5
2000.....	78	55	38	31	35	25	22	10	3	1	-	3
1999.....	72	61	38	42	23	36	25	7	2	2	-	11

White, female

2004.....	29	23	14	13	19	19	13	15	7	2	-	5
2003.....	29	23	10	11	19	20	16	18	2	1	-	6
2002.....	30	23	12	15	10	17	14	11	9	-	-	4
2001 3/.....	82	63	35	19	18	21	20	19	5	1	-	5
2000.....	28	25	10	12	15	24	21	9	5	3	1	6
1999.....	35	19	14	12	22	18	13	14	7	3	-	4

Black, both sexes

2004.....	64	35	21	11	14	12	8	2	2	-	-	1
2003.....	54	32	26	17	13	11	10	5	1	-	-	1
2002.....	53	26	26	15	12	17	8	7	3	-	-	1
2001 3/.....	93	44	32	24	16	10	11	5	2	-	1	3
2000.....	58	30	30	20	16	15	16	3	1	-	-	1
1999.....	48	24	14	19	15	12	9	5	2	2	-	5

Black, male

2004.....	52	25	17	9	11	6	7	-	2	-	-	1
2003.....	45	28	22	15	10	7	6	3	1	-	-	1
2002.....	43	24	23	11	10	14	7	3	2	-	-	-
2001 3/.....	65	30	26	15	14	8	5	3	-	-	1	2
2000.....	53	26	26	18	12	8	12	2	-	-	-	-
1999.....	42	16	10	17	11	5	6	4	1	1	-	2

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Assault (homicide) by all other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5,*U01.8-*U01.9,*U02,X92,X96-X98,Y01-Y03,Y08-Y09,Y87.1)												
Asian or Pacific Islander 2/, both sexes												
2004.....	10	5	6	2	1	2	-	-	-	-	-	-
2003.....	3	4	-	5	1	1	-	-	-	-	-	-
2002.....	4	4	2	3	2	-	1	-	-	-	-	1
2001 3/.....	17	12	10	3	3	2	1	1	-	-	-	-
2000.....	4	5	1	3	2	-	3	-	-	-	-	-
1999.....	3	1	4	1	1	1	1	1	-	-	-	1
Asian or Pacific Islander 2/, male												
2004.....	6	3	5	1	-	1	-	-	-	-	-	-
2003.....	3	4	-	4	-	1	-	-	-	-	-	-
2002.....	4	3	1	2	1	-	-	-	-	-	-	1
2001 3/.....	16	8	7	2	1	-	-	1	-	-	-	-
2000.....	3	2	-	1	2	-	2	-	-	-	-	-
1999.....	3	1	3	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	4	2	1	1	1	1	-	-	-	-	-	-
2003.....	-	-	-	1	1	-	-	-	-	-	-	-
2002.....	-	1	1	1	1	-	1	-	-	-	-	-
2001 3/.....	1	4	3	1	2	2	1	-	-	-	-	-
2000.....	1	3	1	2	-	-	1	-	-	-	-	-
1999.....	-	-	1	1	1	1	1	-	-	-	-	1

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Event of undetermined intent (Y10-Y34,Y87.2,Y89.9)												
All races, both sexes												
2004.....	512	275	124	91	65	49	70	44	26	9	1	9
2003.....	490	245	123	74	57	45	51	36	12	4	-	18
2002.....	431	207	113	80	59	44	51	39	17	8	1	14
2001.....	371	152	73	58	41	53	55	22	11	9	2	18
2000.....	298	154	74	55	49	43	36	26	25	5	1	11
1999.....	265	136	86	72	42	51	45	37	18	2	1	19
All races, male												
2004.....	313	163	68	56	35	24	28	18	8	2	-	9
2003.....	318	153	69	37	28	26	23	15	4	2	-	14
2002.....	264	131	72	46	37	24	22	20	6	5	-	13
2001.....	229	105	41	40	24	28	29	12	6	2	-	16
2000.....	196	105	47	38	32	29	20	14	5	2	-	11
1999.....	175	94	54	47	25	31	24	20	4	1	-	13
All races, female												
2004.....	199	112	56	35	30	25	42	26	18	7	1	-
2003.....	172	92	54	37	29	19	28	21	8	2	-	4
2002.....	167	76	41	34	22	20	29	19	11	3	1	1
2001.....	142	47	32	18	17	25	26	10	5	7	2	2
2000.....	102	49	27	17	17	14	16	12	20	3	1	-
1999.....	90	42	32	25	17	20	21	17	14	1	1	6
White, both sexes												
2004.....	420	239	103	67	56	41	65	39	24	8	1	5
2003.....	403	203	96	59	44	40	45	34	12	4	-	11
2002.....	344	179	93	65	50	37	42	37	16	6	1	12
2001.....	297	129	60	46	37	48	48	20	11	8	2	15
2000.....	240	116	61	45	39	38	34	22	21	5	-	8
1999.....	203	111	75	63	34	47	42	33	16	2	1	14

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Event of undetermined intent (Y10-Y34,Y87.2,Y89.9)												
White, male												
2004.....	252	138	53	41	31	19	27	16	8	2	-	5
2003.....	251	120	51	28	23	24	21	14	4	2	-	8
2002.....	195	110	59	34	28	19	20	19	6	4	-	11
2001.....	175	85	32	32	20	26	27	11	6	2	-	13
2000.....	154	74	38	29	24	27	18	11	4	2	-	8
1999.....	128	70	43	42	19	28	21	18	4	1	-	9
White, female												
2004.....	168	101	50	26	25	22	38	23	16	6	1	-
2003.....	152	83	45	31	21	16	24	20	8	2	-	3
2002.....	149	69	34	31	22	18	22	18	10	2	1	1
2001.....	122	44	28	14	17	22	21	9	5	6	2	2
2000.....	86	42	23	16	15	11	16	11	17	3	-	-
1999.....	75	41	32	21	15	19	21	15	12	1	1	5
Black, both sexes												
2004.....	81	32	18	19	3	7	3	2	2	1	-	4
2003.....	75	31	22	9	12	3	4	2	-	-	-	7
2002.....	80	23	16	10	7	7	9	1	1	2	-	1
2001.....	67	21	11	10	2	2	3	2	-	1	-	2
2000.....	53	31	12	8	7	4	2	3	3	-	1	3
1999.....	59	20	8	8	6	4	1	3	-	-	-	5
Black, male												
2004.....	55	24	14	13	1	4	1	-	-	-	-	4
2003.....	58	25	15	8	4	1	1	1	-	-	-	6
2002.....	63	17	13	9	7	5	2	-	-	1	-	1
2001.....	50	19	8	8	2	-	1	1	-	-	-	2
2000.....	38	24	9	7	6	2	2	2	1	-	-	3
1999.....	46	19	8	5	5	3	1	1	-	-	-	4

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Event of undetermined intent (Y10-Y34,Y87.2,Y89.9)												
Black, female												
2004.....	26	8	4	6	2	3	2	2	2	1	-	-
2003.....	17	6	7	1	8	2	3	1	-	-	-	1
2002.....	17	6	3	1	-	2	7	1	1	1	-	-
2001.....	17	2	3	2	-	2	2	1	-	1	-	-
2000.....	15	7	3	1	1	2	-	1	2	-	1	-
1999.....	13	1	-	3	1	1	-	2	-	-	-	1
American Indian 1/, both sexes												
2004.....	7	3	1	2	1	-	1	-	-	-	-	-
2003.....	6	5	2	1	-	1	-	-	-	-	-	-
2002.....	2	3	1	2	-	-	-	-	-	-	-	1
2001.....	4	-	1	1	-	1	1	-	-	-	-	1
2000.....	3	6	-	-	1	-	-	-	-	-	-	-
1999.....	2	2	1	1	1	-	1	-	1	-	-	-
American Indian 1/, male												
2004.....	2	1	-	1	1	-	-	-	-	-	-	-
2003.....	3	5	2	1	-	-	-	-	-	-	-	-
2002.....	1	2	-	1	-	-	-	-	-	-	-	1
2001.....	2	-	1	-	-	1	1	-	-	-	-	1
2000.....	2	6	-	-	-	-	-	-	-	-	-	-
1999.....	1	2	1	-	1	-	1	-	-	-	-	-
American Indian 1/, female												
2004.....	5	2	1	1	-	-	1	-	-	-	-	-
2003.....	3	-	-	-	-	1	-	-	-	-	-	-
2002.....	1	1	1	1	-	-	-	-	-	-	-	-
2001.....	2	-	-	1	-	-	-	-	-	-	-	-
2000.....	1	-	-	-	1	-	-	-	-	-	-	-
1999.....	1	-	-	1	-	-	-	-	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Event of undetermined intent (Y10-Y34,Y87.2,Y89.9)												
Asian or Pacific Islander 2/, both sexes												
2004.....	4	1	2	3	5	1	1	3	-	-	-	-
2003.....	6	6	3	5	1	1	2	-	-	-	-	-
2002.....	5	2	3	3	2	-	-	1	-	-	-	-
2001.....	3	2	1	1	2	2	3	-	-	-	-	-
2000.....	2	1	1	2	2	1	-	1	1	-	-	-
1999.....	1	3	2	-	1	-	1	1	1	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	4	-	1	1	2	1	-	2	-	-	-	-
2003.....	6	3	1	-	1	1	1	-	-	-	-	-
2002.....	5	2	-	2	2	-	-	1	-	-	-	-
2001.....	2	1	-	-	2	1	-	-	-	-	-	-
2000.....	2	1	-	2	2	-	-	1	-	-	-	-
1999.....	-	3	2	-	-	-	1	1	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	-	1	1	2	3	-	1	1	-	-	-	-
2003.....	-	3	2	5	-	-	1	-	-	-	-	-
2002.....	-	-	3	1	-	-	-	-	-	-	-	-
2001.....	1	1	1	1	-	1	3	-	-	-	-	-
2000.....	-	-	1	-	-	1	-	-	1	-	-	-
1999.....	1	-	-	-	1	-	-	-	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Poisoning by and exposure to drugs and biological substances, undetermined intent (Y10-Y14)												
All races, both sexes												
2004.....	361	180	63	34	19	10	8	4	3	-	-	1
2003.....	353	161	59	29	11	9	6	4	2	-	-	4
2002.....	319	119	61	27	13	9	5	5	3	2	-	3
2001.....	259	87	37	19	9	11	9	4	5	1	-	3
2000.....	199	72	31	13	4	8	5	4	3	-	-	5
1999.....	175	72	35	24	12	9	8	4	4	-	1	6
All races, male												
2004.....	195	86	29	16	6	3	-	2	1	-	-	1
2003.....	209	89	28	10	3	3	2	1	-	-	-	4
2002.....	178	65	34	9	6	4	2	3	1	1	-	2
2001.....	153	49	17	10	2	5	3	1	1	-	-	3
2000.....	129	47	15	6	2	4	3	2	-	-	-	5
1999.....	111	43	16	11	5	4	2	-	-	-	-	4
All races, female												
2004.....	166	94	34	18	13	7	8	2	2	-	-	-
2003.....	144	72	31	19	8	6	4	3	2	-	-	-
2002.....	141	54	27	18	7	5	3	2	2	1	-	1
2001.....	106	38	20	9	7	6	6	3	4	1	-	-
2000.....	70	25	16	7	2	4	2	2	3	-	-	-
1999.....	64	29	19	13	7	5	6	4	4	-	1	2
White, both sexes												
2004.....	306	158	52	28	18	9	6	3	3	-	-	-
2003.....	291	139	50	23	8	9	5	4	2	-	-	1
2002.....	259	100	54	26	12	8	5	5	3	2	-	3
2001.....	211	77	33	14	9	10	8	3	5	1	-	3
2000.....	163	58	29	11	4	8	5	4	3	-	-	3
1999.....	132	60	29	23	11	9	8	4	4	-	1	4

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Poisoning by and exposure to drugs and biological substances, undetermined intent (Y10-Y14)												
White, male												
2004.....	162	72	21	12	6	2	-	2	1	-	-	-
2003.....	163	73	22	5	1	3	2	1	-	-	-	1
2002.....	132	51	30	8	5	4	2	3	1	1	-	2
2001.....	119	41	13	6	2	4	3	-	1	-	-	3
2000.....	104	35	13	5	2	4	3	2	-	-	-	3
1999.....	80	32	10	10	4	4	2	-	-	-	-	2
White, female												
2004.....	144	86	31	16	12	7	6	1	2	-	-	-
2003.....	128	66	28	18	7	6	3	3	2	-	-	-
2002.....	127	49	24	18	7	4	3	2	2	1	-	1
2001.....	92	36	20	8	7	6	5	3	4	1	-	-
2000.....	59	23	16	6	2	4	2	2	3	-	-	-
1999.....	52	28	19	13	7	5	6	4	4	-	1	2
Black, both sexes												
2004.....	48	20	9	6	-	1	1	-	-	-	-	1
2003.....	55	18	9	5	3	-	1	-	-	-	-	3
2002.....	60	17	7	1	1	1	-	-	-	-	-	-
2001.....	48	9	4	4	-	-	-	1	-	-	-	-
2000.....	35	12	2	2	-	-	-	-	-	-	-	2
1999.....	40	9	6	1	1	-	-	-	-	-	-	2
Black, male												
2004.....	30	14	8	4	-	1	-	-	-	-	-	1
2003.....	41	14	6	5	2	-	-	-	-	-	-	3
2002.....	46	12	4	1	1	-	-	-	-	-	-	-
2001.....	34	8	4	4	-	-	-	1	-	-	-	-
2000.....	24	10	2	1	-	-	-	-	-	-	-	2
1999.....	30	8	6	1	1	-	-	-	-	-	-	2

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Poisoning by and exposure to other gases and vapors, undetermined intent (Y17)												
All races, both sexes												
2004.....	8	7	7	1	3	2	1	-	1	1	-	-
2003.....	7	3	7	6	2	3	3	1	-	-	-	-
2002.....	5	7	2	3	3	-	3	1	-	-	-	-
2001.....	3	6	2	2	4	3	2	-	-	-	-	-
2000.....	5	2	1	2	4	2	2	1	3	-	-	-
1999.....	6	1	7	3	3	4	6	4	-	-	-	-
All races, male												
2004.....	5	6	6	1	3	2	1	-	1	-	-	-
2003.....	6	2	4	4	2	1	1	1	-	-	-	-
2002.....	3	5	1	2	2	-	1	1	-	-	-	-
2001.....	1	6	1	2	4	2	-	-	-	-	-	-
2000.....	4	2	1	2	2	2	1	1	1	-	-	-
1999.....	4	-	6	3	2	3	3	3	-	-	-	-
All races, female												
2004.....	3	1	1	-	-	-	-	-	-	1	-	-
2003.....	1	1	3	2	-	2	2	-	-	-	-	-
2002.....	2	2	1	1	1	-	2	-	-	-	-	-
2001.....	2	-	1	-	-	1	2	-	-	-	-	-
2000.....	1	-	-	-	2	-	1	-	2	-	-	-
1999.....	2	1	1	-	1	1	3	1	-	-	-	-
White, both sexes												
2004.....	6	7	7	1	3	2	1	-	1	1	-	-
2003.....	5	3	5	4	2	3	3	1	-	-	-	-
2002.....	5	7	2	3	3	-	1	1	-	-	-	-
2001.....	3	5	2	2	4	3	1	-	-	-	-	-
2000.....	5	1	1	1	4	2	2	1	3	-	-	-
1999.....	6	1	6	1	3	4	6	4	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Discharge of firearms, undetermined intent (Y22-Y24)												
All races, both sexes												
2004.....	17	11	9	11	5	2	6	-	2	-	-	3
2003.....	12	14	6	4	4	4	5	1	-	-	-	-
2002.....	10	9	3	8	7	3	2	-	-	-	-	-
2001.....	13	9	3	6	4	6	5	1	-	-	-	-
2000.....	14	12	3	7	4	5	1	2	-	-	-	1
1999.....	10	6	8	9	6	6	7	4	-	-	-	-
All races, male												
2004.....	13	8	6	9	4	1	5	-	1	-	-	3
2003.....	11	12	4	3	4	4	4	1	-	-	-	-
2002.....	9	5	2	4	6	3	1	-	-	-	-	-
2001.....	9	8	3	6	2	6	3	1	-	-	-	-
2000.....	12	10	1	5	4	4	1	2	-	-	-	1
1999.....	6	5	6	8	6	5	7	4	-	-	-	-
All races, female												
2004.....	4	3	3	2	1	1	1	-	1	-	-	-
2003.....	1	2	2	1	-	-	1	-	-	-	-	-
2002.....	1	4	1	4	1	-	1	-	-	-	-	-
2001.....	4	1	-	-	2	-	2	-	-	-	-	-
2000.....	2	2	2	2	-	1	-	-	-	-	-	-
1999.....	4	1	2	1	-	1	-	-	-	-	-	-
White, both sexes												
2004.....	14	10	9	10	5	2	6	-	2	-	-	2
2003.....	11	12	5	4	4	4	5	1	-	-	-	-
2002.....	9	8	2	8	6	2	2	-	-	-	-	-
2001.....	11	6	3	6	4	6	4	1	-	-	-	-
2000.....	12	8	3	6	3	5	1	2	-	-	-	1
1999.....	10	5	7	9	4	6	6	4	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Discharge of firearms, undetermined intent (Y22-Y24)												
White, male												
2004.....	10	8	6	8	4	1	5	-	1	-	-	2
2003.....	10	11	4	3	4	4	4	1	-	-	-	-
2002.....	8	5	2	4	5	2	1	-	-	-	-	-
2001.....	7	5	3	6	2	6	2	1	-	-	-	-
2000.....	10	6	1	4	3	4	1	2	-	-	-	1
1999.....	6	4	5	8	4	5	6	4	-	-	-	-
White, female												
2004.....	4	2	3	2	1	1	1	-	1	-	-	-
2003.....	1	1	1	1	-	-	1	-	-	-	-	-
2002.....	1	3	-	4	1	-	1	-	-	-	-	-
2001.....	4	1	-	-	2	-	2	-	-	-	-	-
2000.....	2	2	2	2	-	1	-	-	-	-	-	-
1999.....	4	1	2	1	-	1	-	-	-	-	-	-
Black, both sexes												
2004.....	3	1	-	1	-	-	-	-	-	-	-	1
2003.....	1	1	-	-	-	-	-	-	-	-	-	-
2002.....	1	1	-	-	1	1	-	-	-	-	-	-
2001.....	2	3	-	-	-	-	1	-	-	-	-	-
2000.....	1	3	-	1	1	-	-	-	-	-	-	-
1999.....	-	1	-	-	2	-	1	-	-	-	-	-
Black, male												
2004.....	3	-	-	1	-	-	-	-	-	-	-	1
2003.....	1	1	-	-	-	-	-	-	-	-	-	-
2002.....	1	-	-	-	1	1	-	-	-	-	-	-
2001.....	2	3	-	-	-	-	1	-	-	-	-	-
2000.....	1	3	-	1	1	-	-	-	-	-	-	-
1999.....	-	1	-	-	2	-	1	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Falling, jumping or pushed from a high place, undetermined intent (Y30)												
All races, both sexes												
2004.....	2	3	2	3	5	-	-	3	-	2	-	-
2003.....	6	-	4	4	1	2	3	2	2	-	-	-
2002.....	7	6	2	1	2	1	6	1	2	-	-	-
2001.....	6	1	3	-	3	3	2	1	-	-	-	-
2000.....	4	3	4	1	2	1	1	1	2	1	-	-
1999.....	2	1	1	-	1	3	-	-	1	-	-	-
All races, male												
2004.....	1	2	1	2	4	-	-	1	-	1	-	-
2003.....	4	-	1	3	-	1	2	1	1	-	-	-
2002.....	5	6	2	1	2	-	1	1	-	-	-	-
2001.....	4	-	2	-	3	1	1	1	-	-	-	-
2000.....	3	1	3	1	1	1	1	1	1	-	-	-
1999.....	1	1	1	-	-	1	-	-	1	-	-	-
All races, female												
2004.....	1	1	1	1	1	-	-	2	-	1	-	-
2003.....	2	-	3	1	1	1	1	1	1	-	-	-
2002.....	2	-	-	-	-	1	5	-	2	-	-	-
2001.....	2	1	1	-	-	2	1	-	-	-	-	-
2000.....	1	2	1	-	1	-	-	-	1	1	-	-
1999.....	1	-	-	-	1	2	-	-	-	-	-	-
White, both sexes												
2004.....	2	3	2	2	5	-	-	3	-	2	-	-
2003.....	6	-	3	2	1	2	1	2	2	-	-	-
2002.....	7	5	1	1	2	1	5	-	2	-	-	-
2001.....	3	1	2	-	3	3	2	1	-	-	-	-
2000.....	2	3	2	1	2	1	1	1	2	1	-	-
1999.....	2	-	1	-	1	3	-	-	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and unspecified events of undetermined intent and their sequelae (Y20-Y21,Y25-Y29,Y31-Y34,Y87.2,Y89.9)												
All races, both sexes												
2004.....	111	68	43	41	33	35	54	37	19	6	1	5
2003.....	102	65	45	30	37	26	32	27	8	4	-	14
2002.....	80	62	43	41	33	31	34	32	12	6	1	10
2001.....	77	46	26	30	21	30	35	16	6	7	2	15
2000.....	72	61	32	31	33	25	27	18	17	4	1	5
1999.....	65	53	34	34	20	29	24	25	13	2	-	13
All races, male												
2004.....	86	57	26	27	18	18	22	15	5	1	-	5
2003.....	81	50	30	16	17	17	13	10	3	2	-	10
2002.....	61	48	33	30	21	17	16	15	5	4	-	10
2001.....	52	40	17	21	13	14	21	9	5	2	-	13
2000.....	48	42	26	23	21	18	14	8	3	2	-	5
1999.....	51	43	25	24	12	18	12	13	3	1	-	9
All races, female												
2004.....	25	11	17	14	15	17	32	22	14	5	1	-
2003.....	21	15	15	14	20	9	19	17	5	2	-	4
2002.....	19	14	10	11	12	14	18	17	7	2	1	-
2001.....	25	6	9	9	8	16	14	7	1	5	2	2
2000.....	24	19	6	8	12	7	13	10	14	2	1	-
1999.....	14	10	9	10	8	11	12	12	10	1	-	4
White, both sexes												
2004.....	79	55	33	26	25	28	51	33	17	5	1	3
2003.....	81	47	33	25	27	21	29	25	8	4	-	10
2002.....	57	55	32	27	26	26	28	31	11	4	1	8
2001.....	56	37	19	24	17	26	31	15	6	6	2	12
2000.....	55	44	23	26	25	20	25	14	13	4	-	4
1999.....	47	42	31	29	15	25	22	21	11	2	-	10

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and unspecified events of undetermined intent and their sequelae (Y20-Y21,Y25-Y29,Y31-Y34,Y87.2,Y89.9)												
White, male												
2004.....	63	46	19	19	14	14	21	13	5	1	-	3
2003.....	64	34	21	15	14	15	12	9	3	2	-	7
2002.....	41	42	25	19	14	13	14	15	5	3	-	8
2001.....	37	32	13	18	9	13	20	9	5	2	-	10
2000.....	35	29	21	18	15	16	12	5	2	2	-	4
1999.....	36	32	22	22	9	15	10	11	3	1	-	7
White, female												
2004.....	16	9	14	7	11	14	30	20	12	4	1	-
2003.....	17	13	12	10	13	6	17	16	5	2	-	3
2002.....	16	13	7	8	12	13	14	16	6	1	1	-
2001.....	19	5	6	6	8	13	11	6	1	4	2	2
2000.....	20	15	2	8	10	4	13	9	11	2	-	-
1999.....	11	10	9	7	6	10	12	10	8	1	-	3
Black, both sexes												
2004.....	28	11	9	11	3	6	2	2	2	1	-	2
2003.....	17	12	9	2	9	3	2	2	-	-	-	4
2002.....	18	5	8	9	5	5	6	1	1	2	-	1
2001.....	15	9	6	5	2	2	1	1	-	1	-	2
2000.....	15	14	8	4	6	4	2	3	3	-	1	1
1999.....	18	9	1	5	3	4	-	3	-	-	-	3
Black, male												
2004.....	20	10	6	7	1	3	1	-	-	-	-	2
2003.....	14	10	7	1	2	1	-	1	-	-	-	3
2002.....	15	5	8	8	5	4	2	-	-	1	-	1
2001.....	12	8	4	3	2	-	-	-	-	-	-	2
2000.....	11	10	5	4	5	2	2	2	1	-	-	1
1999.....	15	9	1	2	2	3	-	1	-	-	-	2

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

All other and unspecified events of undetermined intent and their sequelae (Y20-Y21,Y25-Y29,Y31-Y34,Y87.2,Y89.9)

Black, female

2004.....	8	1	3	4	2	3	1	2	2	1	-	-
2003.....	3	2	2	1	7	2	2	1	-	-	-	1
2002.....	3	-	-	1	-	1	4	1	1	1	-	-
2001.....	3	1	2	2	-	2	1	1	-	1	-	-
2000.....	4	4	3	-	1	2	-	1	2	-	1	-
1999.....	3	-	-	3	1	1	-	2	-	-	-	1

American Indian 1/, both sexes

2004.....	2	2	-	1	1	-	1	-	-	-	-	-
2003.....	2	3	1	-	-	1	-	-	-	-	-	-
2002.....	1	2	-	2	-	-	-	-	-	-	-	1
2001.....	3	-	-	-	-	-	1	-	-	-	-	1
2000.....	1	3	-	-	1	-	-	-	-	-	-	-
1999.....	-	-	1	-	1	-	1	-	1	-	-	-

American Indian 1/, male

2004.....	1	1	-	-	1	-	-	-	-	-	-	-
2003.....	1	3	1	-	-	-	-	-	-	-	-	-
2002.....	1	1	-	1	-	-	-	-	-	-	-	1
2001.....	1	-	-	-	-	-	1	-	-	-	-	1
2000.....	1	3	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	1	-	1	-	1	-	-	-	-	-

American Indian 1/, female

2004.....	1	1	-	1	-	-	1	-	-	-	-	-
2003.....	1	-	-	-	-	1	-	-	-	-	-	-
2002.....	-	1	-	1	-	-	-	-	-	-	-	-
2001.....	2	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	1	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

All other and unspecified events of undetermined intent and their sequelae (Y20-Y21,Y25-Y29,Y31-Y34,Y87.2,Y89.9)

Asian or Pacific Islander 2/, both sexes

2004.....	2	-	1	3	4	1	-	2	-	-	-	-
2003.....	2	3	2	3	1	1	1	-	-	-	-	-
2002.....	4	-	3	3	2	-	-	-	-	-	-	-
2001.....	3	-	1	1	2	2	2	-	-	-	-	-
2000.....	1	-	1	1	1	1	-	1	1	-	-	-
1999.....	-	2	1	-	1	-	1	1	1	-	-	-

Asian or Pacific Islander 2/, male

2004.....	2	-	1	1	2	1	-	2	-	-	-	-
2003.....	2	3	1	-	1	1	1	-	-	-	-	-
2002.....	4	-	-	2	2	-	-	-	-	-	-	-
2001.....	2	-	-	-	2	1	-	-	-	-	-	-
2000.....	1	-	-	1	1	-	-	1	-	-	-	-
1999.....	-	2	1	-	-	-	1	1	-	-	-	-

Asian or Pacific Islander 2/, female

2004.....	-	-	-	2	2	-	-	-	-	-	-	-
2003.....	-	-	1	3	-	-	-	-	-	-	-	-
2002.....	-	-	3	1	-	-	-	-	-	-	-	-
2001.....	1	-	1	1	-	1	2	-	-	-	-	-
2000.....	-	-	1	-	-	1	-	-	1	-	-	-
1999.....	-	-	-	-	1	-	-	-	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Legal intervention (Y35,Y89.0)												
All races, both sexes												
2004.....	27	9	3	4	3	3	2	-	-	-	-	-
2003.....	22	12	7	2	2	2	-	-	-	-	-	-
2002.....	13	10	6	4	2	-	1	-	-	-	-	-
2001.....	18	15	9	4	2	3	1	1	-	-	-	-
2000.....	17	6	4	2	3	1	-	1	-	-	-	-
1999.....	31	9	11	2	-	1	-	-	-	-	-	-
All races, male												
2004.....	27	8	3	3	3	3	2	-	-	-	-	-
2003.....	19	11	7	2	2	2	-	-	-	-	-	-
2002.....	12	10	6	4	2	-	1	-	-	-	-	-
2001.....	18	15	7	4	2	3	1	1	-	-	-	-
2000.....	17	6	3	2	3	1	-	1	-	-	-	-
1999.....	30	9	11	2	-	1	-	-	-	-	-	-
All races, female												
2004.....	-	1	-	1	-	-	-	-	-	-	-	-
2003.....	3	1	-	-	-	-	-	-	-	-	-	-
2002.....	1	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	2	-	-	-	-	-	-	-	-	-
2000.....	-	-	1	-	-	-	-	-	-	-	-	-
1999.....	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes												
2004.....	18	5	3	2	3	3	2	-	-	-	-	-
2003.....	19	10	7	2	2	1	-	-	-	-	-	-
2002.....	13	8	5	4	1	-	1	-	-	-	-	-
2001.....	17	10	8	4	2	3	1	-	-	-	-	-
2000.....	11	5	4	1	2	1	-	1	-	-	-	-
1999.....	25	7	8	2	-	1	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other legal intervention and their sequelae (Y35.0-Y35.4,Y35.6-Y35.7,Y89.0)												
All races, both sexes												
2004.....	21	5	3	4	2	3	2	-	-	-	-	-
2003.....	19	8	3	2	2	2	-	-	-	-	-	-
2002.....	6	6	3	3	2	-	1	-	-	-	-	-
2001.....	11	10	6	4	2	3	1	1	-	-	-	-
2000.....	8	5	2	2	3	1	-	1	-	-	-	-
1999.....	19	5	7	2	-	1	-	-	-	-	-	-
All races, male												
2004.....	21	4	3	3	2	3	2	-	-	-	-	-
2003.....	16	7	3	2	2	2	-	-	-	-	-	-
2002.....	6	6	3	3	2	-	1	-	-	-	-	-
2001.....	11	10	5	4	2	3	1	1	-	-	-	-
2000.....	8	5	2	2	3	1	-	1	-	-	-	-
1999.....	18	5	7	2	-	1	-	-	-	-	-	-
All races, female												
2004.....	-	1	-	1	-	-	-	-	-	-	-	-
2003.....	3	1	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	1	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes												
2004.....	16	2	3	2	2	3	2	-	-	-	-	-
2003.....	17	7	3	2	2	1	-	-	-	-	-	-
2002.....	6	4	3	3	1	-	1	-	-	-	-	-
2001.....	11	5	6	4	2	3	1	-	-	-	-	-
2000.....	5	4	2	1	2	1	-	1	-	-	-	-
1999.....	17	3	6	2	-	1	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Operations of war and their sequelae (Y36,Y89.1)												
All races, both sexes												
2004.....	-	3	1	1	-	5	9	1	-	-	-	-
2003.....	1	5	-	-	2	2	3	-	-	-	-	-
2002.....	-	2	1	2	1	4	6	1	-	-	-	-
2001.....	1	-	-	-	4	7	1	1	-	-	-	-
2000.....	2	-	1	1	2	4	5	1	-	-	-	-
1999.....	2	1	-	2	6	4	4	2	1	-	-	-
All races, male												
2004.....	-	3	1	1	-	5	9	1	-	-	-	-
2003.....	1	5	-	-	2	2	3	-	-	-	-	-
2002.....	-	2	1	2	1	4	6	1	-	-	-	-
2001.....	1	-	-	-	4	6	1	1	-	-	-	-
2000.....	2	-	1	1	2	4	5	1	-	-	-	-
1999.....	2	1	-	2	6	4	3	2	1	-	-	-
All races, female												
2004.....	-	-	-	-	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	1	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes												
2004.....	-	3	1	1	-	5	9	1	-	-	-	-
2003.....	1	4	-	-	2	2	3	-	-	-	-	-
2002.....	-	1	1	2	-	4	5	1	-	-	-	-
2001.....	1	-	-	-	3	7	1	1	-	-	-	-
2000.....	1	-	1	1	2	4	5	1	-	-	-	-
1999.....	2	1	-	1	6	4	4	2	1	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Complications of medical and surgical care (Y40-Y84,Y88)												
All races, both sexes												
2004.....	159	206	197	257	300	367	420	292	175	34	12	-
2003.....	128	178	202	214	320	398	396	307	152	58	13	1
2002.....	151	164	187	225	335	382	410	288	167	59	7	-
2001.....	141	178	206	247	320	439	425	315	186	56	7	1
2000.....	167	157	219	263	384	427	398	337	176	54	11	-
1999.....	123	138	177	229	357	401	410	312	153	52	12	-
All races, male												
2004.....	81	119	101	115	165	181	204	118	50	8	4	-
2003.....	66	101	94	103	161	174	181	129	46	8	3	1
2002.....	67	84	85	110	146	166	154	110	40	18	3	-
2001.....	68	91	105	116	149	192	191	119	55	13	2	-
2000.....	84	77	104	123	208	212	190	134	48	13	1	-
1999.....	61	77	88	102	177	193	182	128	49	17	3	-
All races, female												
2004.....	78	87	96	142	135	186	216	174	125	26	8	-
2003.....	62	77	108	111	159	224	215	178	106	50	10	-
2002.....	84	80	102	115	189	216	256	178	127	41	4	-
2001.....	73	87	101	131	171	247	234	196	131	43	5	1
2000.....	83	80	115	140	176	215	208	203	128	41	10	-
1999.....	62	61	89	127	180	208	228	184	104	35	9	-
White, both sexes												
2004.....	124	159	164	209	251	315	367	268	151	27	9	-
2003.....	89	133	164	178	267	336	360	269	135	54	11	1
2002.....	105	122	149	178	292	331	366	261	146	51	7	-
2001.....	107	137	156	198	247	387	380	267	160	50	6	1
2000.....	134	120	165	215	320	361	357	297	155	41	10	-
1999.....	91	98	142	193	299	343	356	266	137	44	8	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Complications of medical and surgical care (Y40-Y84,Y88)												
White, male												
2004.....	67	92	87	100	141	162	182	107	45	7	1	-
2003.....	43	78	79	90	139	154	169	114	43	7	3	1
2002.....	50	65	70	84	128	147	143	97	35	14	3	-
2001.....	49	66	79	95	112	171	176	103	45	11	1	-
2000.....	70	64	80	98	175	186	177	119	44	11	1	-
1999.....	46	55	71	87	148	169	159	108	43	16	1	-
White, female												
2004.....	57	67	77	109	110	153	185	161	106	20	8	-
2003.....	46	55	85	88	128	182	191	155	92	47	8	-
2002.....	55	57	79	94	164	184	223	164	111	37	4	-
2001.....	58	71	77	103	135	216	204	164	115	39	5	1
2000.....	64	56	85	117	145	175	180	178	111	30	9	-
1999.....	45	43	71	106	151	174	197	158	94	28	7	-
Black, both sexes												
2004.....	32	45	30	44	42	45	48	18	21	7	3	-
2003.....	34	39	35	34	44	56	30	31	15	4	2	-
2002.....	44	38	32	38	39	44	36	21	19	7	-	-
2001.....	31	38	47	43	63	46	41	46	23	6	1	-
2000.....	31	35	51	42	59	59	37	34	20	12	1	-
1999.....	31	34	28	32	54	53	43	37	16	7	3	-
Black, male												
2004.....	12	26	12	14	19	17	19	7	4	1	3	-
2003.....	21	20	15	13	15	17	10	10	3	1	-	-
2002.....	16	18	12	20	17	14	11	10	5	4	-	-
2001.....	18	22	25	21	33	17	14	14	9	2	1	-
2000.....	13	12	22	24	28	22	12	12	3	1	-	-
1999.....	15	16	14	13	27	22	18	12	6	1	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Complications of medical and surgical care (Y40-Y84,Y88)												
Black, female												
2004.....	20	19	18	30	23	28	29	11	17	6	-	-
2003.....	13	19	20	21	29	39	20	21	12	3	2	-
2002.....	28	20	20	18	22	30	25	11	14	3	-	-
2001.....	13	16	22	22	30	29	27	32	14	4	-	-
2000.....	18	23	29	18	31	37	25	22	17	11	1	-
1999.....	16	18	14	19	27	31	25	25	10	6	2	-
American Indian 1/, both sexes												
2004.....	1	-	1	3	1	-	-	-	1	-	-	-
2003.....	1	1	3	1	2	1	2	2	1	-	-	-
2002.....	1	3	2	6	1	2	2	2	-	-	-	-
2001.....	2	1	-	1	2	3	1	-	1	-	-	-
2000.....	1	-	-	4	1	1	2	3	-	-	-	-
1999.....	1	2	1	2	1	2	3	2	-	1	1	-
American Indian 1/, male												
2004.....	1	-	1	1	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	1	-	-	2	-	-	-	-
2002.....	1	-	1	4	-	1	-	1	-	-	-	-
2001.....	1	1	-	-	-	2	-	-	-	-	-	-
2000.....	1	-	-	-	1	1	1	1	-	-	-	-
1999.....	-	2	1	1	1	1	3	1	-	-	1	-
American Indian 1/, female												
2004.....	-	-	-	2	1	-	-	-	1	-	-	-
2003.....	1	1	3	1	1	1	2	-	1	-	-	-
2002.....	-	3	1	2	1	1	2	1	-	-	-	-
2001.....	1	-	-	1	2	1	1	-	1	-	-	-
2000.....	-	-	-	4	-	-	1	2	-	-	-	-
1999.....	1	-	-	1	-	1	-	1	-	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Complications of medical and surgical care (Y40-Y84,Y88)												
Asian or Pacific Islander 2/, both sexes												
2004.....	2	2	2	1	6	7	5	6	2	-	-	-
2003.....	4	5	-	1	7	5	4	5	1	-	-	-
2002.....	1	1	4	3	3	5	6	4	2	1	-	-
2001.....	1	2	3	5	8	3	3	2	2	-	-	-
2000.....	1	2	3	2	4	6	2	3	1	1	-	-
1999.....	-	4	6	2	3	3	8	7	-	-	-	-
Asian or Pacific Islander 2/, male												
2004.....	1	1	1	-	5	2	3	4	1	-	-	-
2003.....	2	3	-	-	6	3	2	3	-	-	-	-
2002.....	-	1	2	2	1	4	-	2	-	-	-	-
2001.....	-	2	1	-	4	2	1	2	1	-	-	-
2000.....	-	1	2	1	4	3	-	2	1	1	-	-
1999.....	-	4	2	1	1	1	2	7	-	-	-	-
Asian or Pacific Islander 2/, female												
2004.....	1	1	1	1	1	5	2	2	1	-	-	-
2003.....	2	2	-	1	1	2	2	2	1	-	-	-
2002.....	1	-	2	1	2	1	6	2	2	1	-	-
2001.....	1	-	2	5	4	1	2	-	1	-	-	-
2000.....	1	1	1	1	-	3	2	1	-	-	-	-
1999.....	-	-	4	1	2	2	6	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Drugs, medicaments and biological substances causing adverse effects in therapeutic use, and their sequelae (Y40-Y59,Y88.0)												
All races, both sexes												
2004.....	17	24	31	28	25	42	38	28	17	3	-	-
2003.....	18	20	23	31	24	29	26	19	16	1	-	-
2002.....	13	22	12	19	29	18	27	10	9	2	-	-
2001.....	15	10	15	18	22	28	24	26	12	2	-	1
2000.....	18	16	15	15	20	28	35	19	12	1	1	-
1999.....	16	15	17	17	21	17	18	12	4	2	-	-
All races, male												
2004.....	9	16	11	11	16	24	14	6	7	1	-	-
2003.....	8	11	14	12	16	15	8	6	4	-	-	-
2002.....	5	8	6	10	13	7	8	2	2	-	-	-
2001.....	6	3	9	8	9	15	9	4	3	-	-	-
2000.....	9	7	10	5	10	13	18	8	5	-	-	-
1999.....	7	10	8	8	11	7	8	3	1	-	-	-
All races, female												
2004.....	8	8	20	17	9	18	24	22	10	2	-	-
2003.....	10	9	9	19	8	14	18	13	12	1	-	-
2002.....	8	14	6	9	16	11	19	8	7	2	-	-
2001.....	9	7	6	10	13	13	15	22	9	2	-	1
2000.....	9	9	5	10	10	15	17	11	7	1	1	-
1999.....	9	5	9	9	10	10	10	9	3	2	-	-
White, both sexes												
2004.....	12	21	23	22	21	38	32	27	17	2	-	-
2003.....	15	15	17	29	21	26	22	18	16	1	-	-
2002.....	10	16	8	18	25	15	26	10	9	2	-	-
2001.....	12	9	12	16	15	23	22	24	12	1	-	1
2000.....	11	14	9	13	16	27	31	16	12	1	1	-
1999.....	14	12	12	13	19	13	17	12	4	2	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Drugs, medicaments and biological substances causing adverse effects in therapeutic use, and their sequelae (Y40-Y59,Y88.0)

White, male

2004.....	6	14	10	8	13	22	13	6	7	1	-	-
2003.....	6	7	10	11	13	14	7	6	4	-	-	-
2002.....	4	7	6	9	12	6	8	2	2	-	-	-
2001.....	5	3	9	7	7	13	9	4	3	-	-	-
2000.....	7	6	7	5	8	13	17	7	5	-	-	-
1999.....	6	8	4	7	10	6	8	3	1	-	-	-

White, female

2004.....	6	7	13	14	8	16	19	21	10	1	-	-
2003.....	9	8	7	18	8	12	15	12	12	1	-	-
2002.....	6	9	2	9	13	9	18	8	7	2	-	-
2001.....	7	6	3	9	8	10	13	20	9	1	-	1
2000.....	4	8	2	8	8	14	14	9	7	1	1	-
1999.....	8	4	8	6	9	7	9	9	3	2	-	-

Black, both sexes

2004.....	5	3	7	6	3	2	6	1	-	1	-	-
2003.....	1	4	6	2	3	3	2	1	-	-	-	-
2002.....	3	5	4	1	3	2	1	-	-	-	-	-
2001.....	3	1	3	2	5	4	2	2	-	1	-	-
2000.....	7	2	6	1	3	1	4	2	-	-	-	-
1999.....	2	3	1	4	2	4	1	-	-	-	-	-

Black, male

2004.....	3	2	1	3	2	1	1	-	-	-	-	-
2003.....	1	3	4	1	3	1	1	-	-	-	-	-
2002.....	1	1	-	1	1	-	-	-	-	-	-	-
2001.....	1	-	-	1	2	2	-	-	-	-	-	-
2000.....	2	1	3	-	1	-	1	-	-	-	-	-
1999.....	1	2	1	1	1	1	-	-	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Misadventures to patients during medical and surgical care and their sequelae (Y60-Y69,Y88.1)												
All races, both sexes												
2004.....	12	17	15	13	12	19	35	11	1	-	1	-
2003.....	11	10	16	10	12	19	18	21	5	2	-	-
2002.....	13	17	9	12	19	20	21	14	7	-	-	-
2001.....	14	5	12	21	17	28	22	12	7	4	-	-
2000.....	12	14	9	20	25	25	16	15	9	2	-	-
1999.....	9	12	9	18	24	24	26	15	4	-	-	-
All races, male												
2004.....	4	7	6	4	4	10	13	5	-	-	-	-
2003.....	6	5	8	3	4	7	6	8	1	1	-	-
2002.....	6	9	5	5	7	8	6	3	2	-	-	-
2001.....	4	5	9	6	7	8	12	5	4	1	-	-
2000.....	5	7	1	8	13	16	8	6	1	1	-	-
1999.....	3	6	3	7	9	12	10	6	3	-	-	-
All races, female												
2004.....	8	10	9	9	8	9	22	6	1	-	1	-
2003.....	5	5	8	7	8	12	12	13	4	1	-	-
2002.....	7	8	4	7	12	12	15	11	5	-	-	-
2001.....	10	-	3	15	10	20	10	7	3	3	-	-
2000.....	7	7	8	12	12	9	8	9	8	1	-	-
1999.....	6	6	6	11	15	12	16	9	1	-	-	-
White, both sexes												
2004.....	11	11	11	11	11	17	30	10	1	-	1	-
2003.....	7	8	12	6	8	17	16	20	4	1	-	-
2002.....	8	11	7	10	14	17	20	13	6	-	-	-
2001.....	11	4	8	14	13	23	21	11	6	4	-	-
2000.....	10	13	3	14	19	21	14	14	7	1	-	-
1999.....	7	10	9	15	16	21	22	11	4	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Misadventures to patients during medical and surgical care and their sequelae (Y60-Y69,Y88.1)												
White, male												
2004.....	4	2	3	4	3	9	9	4	-	-	-	-
2003.....	4	5	6	2	3	7	6	7	-	-	-	-
2002.....	6	7	3	4	5	7	6	3	1	-	-	-
2001.....	4	4	7	5	4	7	11	4	3	1	-	-
2000.....	4	7	-	3	9	14	8	6	1	1	-	-
1999.....	3	6	3	5	7	10	8	4	3	-	-	-
White, female												
2004.....	7	9	8	7	8	8	21	6	1	-	1	-
2003.....	3	3	6	4	5	10	10	13	4	1	-	-
2002.....	2	4	4	6	9	10	14	10	5	-	-	-
2001.....	7	-	1	9	9	16	10	7	3	3	-	-
2000.....	6	6	3	11	10	7	6	8	6	-	-	-
1999.....	4	4	6	10	9	11	14	7	1	-	-	-
Black, both sexes												
2004.....	1	5	2	2	-	2	4	1	-	-	-	-
2003.....	4	2	4	4	2	2	2	1	1	1	-	-
2002.....	5	6	2	1	5	3	-	1	1	-	-	-
2001.....	2	1	4	6	4	5	1	1	-	-	-	-
2000.....	2	1	6	5	6	3	1	1	2	1	-	-
1999.....	2	2	-	2	8	2	4	3	-	-	-	-
Black, male												
2004.....	-	4	1	-	-	1	3	1	-	-	-	-
2003.....	2	-	2	1	-	-	-	1	1	1	-	-
2002.....	-	2	2	-	2	1	-	-	1	-	-	-
2001.....	-	1	2	1	3	1	1	1	-	-	-	-
2000.....	1	-	1	4	4	1	-	-	-	-	-	-
1999.....	-	-	-	1	2	1	2	1	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other complications of medical and surgical care and their sequelae (Y70-Y84,Y88.2-Y88.3)												
All races, both sexes												
2004.....	130	165	151	216	263	306	347	253	157	31	11	-
2003.....	99	148	163	173	284	350	352	267	131	55	13	1
2002.....	125	125	166	194	287	344	362	264	151	57	7	-
2001.....	112	163	179	208	281	383	379	277	167	50	7	-
2000.....	137	127	195	228	339	374	347	303	155	51	10	-
1999.....	98	111	151	194	312	360	366	285	145	50	12	-
All races, male												
2004.....	68	96	84	100	145	147	177	107	43	7	4	-
2003.....	52	85	72	88	141	152	167	115	41	7	3	1
2002.....	56	67	74	95	126	151	140	105	36	18	3	-
2001.....	58	83	87	102	133	169	170	110	48	12	2	-
2000.....	70	63	93	110	185	183	164	120	42	12	1	-
1999.....	51	61	77	87	157	174	164	119	45	17	3	-
All races, female												
2004.....	62	69	67	116	118	159	170	146	114	24	7	-
2003.....	47	63	91	85	143	198	185	152	90	48	10	-
2002.....	69	58	92	99	161	193	222	159	115	39	4	-
2001.....	54	80	92	106	148	214	209	167	119	38	5	-
2000.....	67	64	102	118	154	191	183	183	113	39	9	-
1999.....	47	50	74	107	155	186	202	166	100	33	9	-
White, both sexes												
2004.....	101	127	130	176	219	260	305	231	133	25	8	-
2003.....	67	110	135	143	238	293	322	231	115	52	11	1
2002.....	87	95	134	150	253	299	320	238	131	49	7	-
2001.....	84	124	136	168	219	341	337	232	142	45	6	-
2000.....	113	93	153	188	285	313	312	267	136	39	9	-
1999.....	70	76	121	165	264	309	317	243	129	42	8	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other complications of medical and surgical care and their sequelae (Y70-Y84,Y88.2-Y88.3)												
White, male												
2004.....	57	76	74	88	125	131	160	97	38	6	1	-
2003.....	33	66	63	77	123	133	156	101	39	7	3	1
2002.....	40	51	61	71	111	134	129	92	32	14	3	-
2001.....	40	59	63	83	101	151	156	95	39	10	1	-
2000.....	59	51	73	90	158	159	152	106	38	10	1	-
1999.....	37	41	64	75	131	153	143	101	39	16	1	-
White, female												
2004.....	44	51	56	88	94	129	145	134	95	19	7	-
2003.....	34	44	72	66	115	160	166	130	76	45	8	-
2002.....	47	44	73	79	142	165	191	146	99	35	4	-
2001.....	44	65	73	85	118	190	181	137	103	35	5	-
2000.....	54	42	80	98	127	154	160	161	98	29	8	-
1999.....	33	35	57	90	133	156	174	142	90	26	7	-
Black, both sexes												
2004.....	26	37	21	36	39	41	38	16	21	6	3	-
2003.....	29	33	25	28	39	51	26	29	14	3	2	-
2002.....	36	27	26	36	31	39	35	20	18	7	-	-
2001.....	26	36	40	35	54	37	38	43	23	5	1	-
2000.....	22	32	39	36	50	55	32	31	18	11	1	-
1999.....	27	29	27	26	44	47	38	34	16	7	3	-
Black, male												
2004.....	9	20	10	11	17	15	15	6	4	1	3	-
2003.....	18	17	9	11	12	16	9	9	2	-	-	-
2002.....	15	15	10	19	14	13	11	10	4	4	-	-
2001.....	17	21	23	19	28	14	13	13	9	2	1	-
2000.....	10	11	18	20	23	21	11	12	3	1	-	-
1999.....	14	14	13	11	24	20	16	11	6	1	1	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other complications of medical and surgical care and their sequelae (Y70-Y84,Y88.2-Y88.3)												
Black, female												
2004.....	17	17	11	25	22	26	23	10	17	5	-	-
2003.....	11	16	16	17	27	35	17	20	12	3	2	-
2002.....	21	12	16	17	17	26	24	10	14	3	-	-
2001.....	9	15	17	16	26	23	25	30	14	3	-	-
2000.....	12	21	21	16	27	34	21	19	15	10	1	-
1999.....	13	15	14	15	20	27	22	23	10	6	2	-
American Indian 1/, both sexes												
2004.....	1	-	-	3	1	-	-	-	1	-	-	-
2003.....	1	1	3	1	2	1	1	2	1	-	-	-
2002.....	1	2	2	5	1	2	2	2	-	-	-	-
2001.....	2	1	-	1	2	3	1	-	1	-	-	-
2000.....	1	-	-	3	1	-	2	3	-	-	-	-
1999.....	1	2	-	2	1	1	3	2	-	1	1	-
American Indian 1/, male												
2004.....	1	-	-	1	-	-	-	-	-	-	-	-
2003.....	-	-	-	-	1	-	-	2	-	-	-	-
2002.....	1	-	1	3	-	1	-	1	-	-	-	-
2001.....	1	1	-	-	-	2	-	-	-	-	-	-
2000.....	1	-	-	-	1	-	1	1	-	-	-	-
1999.....	-	2	-	1	1	-	3	1	-	-	1	-
American Indian 1/, female												
2004.....	-	-	-	2	1	-	-	-	1	-	-	-
2003.....	1	1	3	1	1	1	1	-	1	-	-	-
2002.....	-	2	1	2	1	1	2	1	-	-	-	-
2001.....	1	-	-	1	2	1	1	-	1	-	-	-
2000.....	-	-	-	3	-	-	1	2	-	-	-	-
1999.....	1	-	-	1	-	1	-	1	-	1	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other complications of medical and surgical care and their sequelae (Y70-Y84,Y88.2-Y88.3)

Asian or Pacific Islander 2/, both sexes

2004.....	2	1	-	1	4	5	4	6	2	-	-	-
2003.....	2	4	-	1	5	5	3	5	1	-	-	-
2002.....	1	1	4	3	2	4	5	4	2	1	-	-
2001.....	-	2	3	4	6	2	3	2	1	-	-	-
2000.....	1	2	3	1	3	6	1	2	1	1	-	-
1999.....	-	4	3	1	3	3	8	6	-	-	-	-

Asian or Pacific Islander 2/, male

2004.....	1	-	-	-	3	1	2	4	1	-	-	-
2003.....	1	2	-	-	5	3	2	3	-	-	-	-
2002.....	-	1	2	2	1	3	-	2	-	-	-	-
2001.....	-	2	1	-	4	2	1	2	-	-	-	-
2000.....	-	1	2	-	3	3	-	1	1	1	-	-
1999.....	-	4	-	-	1	1	2	6	-	-	-	-

Worktable 292F. Deaths from 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Montana, New York, and Wisconsin in 2003. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white or black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other complications of medical and surgical care and their sequelae (Y70-Y84,Y88.2-Y88.3)												
Asian or Pacific Islander 2/, female												
2004.....	1	1	-	1	1	4	2	2	1	-	-	-
2003.....	1	2	-	1	-	2	1	2	1	-	-	-
2002.....	1	-	2	1	1	1	5	2	2	1	-	-
2001.....	-	-	2	4	2	-	2	-	1	-	-	-
2000.....	1	1	1	1	-	3	1	1	-	-	-	-
1999.....	-	-	3	1	2	2	6	-	-	-	-	-

- Quantity zero.

... Category not applicable.

1/ Includes Aleuts and Eskimos.

2/ Includes Chinese, Filipino, Hawaiian, Japanese, and Other Asian or Pacific Islander.

3/ Figures include September 11, 2001 related deaths for which death certificates were filed as of 10/24/02; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001."

Source: CDC/NCHS, National Vital Statistics System, Mortality.