

Worktable Orig13. Age-adjusted death rates for 358 selected causes, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), and year	All origins 1/			Hispanic			Non-Hispanic 2/			Non-Hispanic white			Non-Hispanic black		
	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female
All causes															
2004.....	800.8	955.7	679.2	586.7	706.8	485.9	814.1	971.1	691.4	797.1	949.0	677.5	1,044.7	1,291.5	869.4
2003.....	832.7	994.3	706.2	621.2	748.1	515.8	844.5	1,008.0	717.2	826.1	984.0	702.1	1,083.2	1,341.1	899.8
2002.....	845.3	1,013.7	715.2	629.3	766.7	518.3	856.5	1,026.5	725.8	837.5	1,002.2	709.9	1,099.2	1,360.6	915.3
2001.....	854.5	1,029.1	721.8	658.7	802.5	544.2	864.0	1,039.8	730.9	842.9	1,012.8	713.5	1,116.5	1,393.7	925.5
2000.....	869.0	1,053.8	731.4	665.7	818.1	546.0	877.9	1,063.8	740.0	855.5	1,035.4	721.5	1,137.0	1,422.0	941.2
1999.....	875.6	1,067.0	734.0	676.4	830.5	555.9	883.9	1,076.4	741.9	859.8	1,045.5	722.3	1,150.1	1,449.4	946.0
I. Certain infectious and parasitic diseases (A00-B99)															
2004.....	21.7	26.3	17.6	21.0	26.5	15.8	21.6	26.0	17.8	17.8	21.3	14.8	51.5	66.3	39.8
2003.....	22.1	26.7	17.9	21.3	27.2	15.6	22.0	26.4	18.0	17.9	21.4	14.9	53.4	69.6	40.7
2002.....	22.2	27.2	17.7	21.6	27.4	16.0	22.0	26.9	17.8	17.8	21.5	14.6	55.2	72.1	41.8
2001.....	21.3	26.3	16.9	21.7	27.8	15.9	21.1	25.9	16.9	16.9	20.6	13.7	54.2	71.2	41.0
2000.....	21.3	26.3	16.8	21.4	28.3	15.1	21.0	25.8	16.8	16.6	20.2	13.5	55.7	73.8	41.6
1999.....	21.8	27.2	17.2	23.1	29.5	17.0	21.5	26.7	17.1	17.0	20.8	13.7	57.7	76.7	42.7
Intestinal infectious diseases (A00-A09)															
2004.....	1.5	1.4	1.5	0.9	0.8	0.9	1.5	1.5	1.5	1.6	1.5	1.6	1.0	1.0	0.9
2003.....	1.1	1.0	1.1	0.7	0.8	0.6	1.1	1.0	1.1	1.1	1.1	1.2	0.8	0.7	0.8
2002.....	0.9	0.9	0.9	0.5	0.5	0.5	0.9	0.9	0.9	0.9	0.9	0.9	0.6	0.7	0.6
2001.....	0.5	0.5	0.6	0.4	0.4	0.4	0.6	0.6	0.6	0.6	0.6	0.6	0.4	0.5	0.4
2000.....	0.5	0.5	0.5	0.3	0.4	0.3	0.5	0.5	0.5	0.5	0.5	0.5	0.4	0.4	0.3
1999.....	0.4	0.4	0.4	0.3	0.3	0.2	0.4	0.4	0.4	0.4	0.5	0.4	0.3	0.3	0.3
Cholera (A00)															
2004.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Other intestinal infectious diseases (A01-A08)															
2004.....	1.4	1.4	1.4	0.8	0.8	0.9	1.5	1.4	1.5	1.5	1.5	1.6	0.9	1.0	0.9
2003.....	1.0	1.0	1.0	0.6	0.7	0.6	1.0	1.0	1.1	1.1	1.0	1.1	0.8	0.7	0.8
2002.....	0.8	0.8	0.8	0.5	0.5	0.5	0.8	0.8	0.8	0.9	0.9	0.9	0.6	0.7	0.6
2001.....	0.5	0.5	0.5	0.4	0.4	0.4	0.5	0.5	0.5	0.6	0.6	0.6	0.4	0.5	0.4
2000.....	0.5	0.5	0.5	0.3	0.4	0.3	0.5	0.5	0.5	0.5	0.5	0.5	0.3	0.4	0.3
1999.....	0.4	0.4	0.4	0.2	0.3	*	0.4	0.4	0.4	0.4	0.4	0.4	0.3	0.3	0.3
Typhoid fever (A01.0)															
2004.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Paratyphoid fevers and other salmonella infections (A01.1-A01.4,A02)															
2004.....	0.0	0.0	*	*	*	*	0.0	*	*	*	*	*	*	*	*
2003.....	0.0	0.0	0.0	*	*	*	0.0	0.0	*	0.0	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2001.....	0.0	0.0	*	*	*	*	0.0	*	*	0.0	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	0.0	*	*	0.0	*	*	*	*	*
1999.....	0.0	0.0	*	*	*	*	0.0	*	*	0.0	*	*	*	*	*

Worktable Orig13. Age-adjusted death rates for 358 selected causes, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), and year	All origins 1/			Hispanic			Non-Hispanic 2/			Non-Hispanic white			Non-Hispanic black		
	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female
Shigellosis (A03)															
2004.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Other bacterial food-borne intoxications (A05)															
2004.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Amebiasis (A06)															
2004.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Intestinal infections due to other specified organisms (A04,A07-A08)															
2004.....	1.4	1.4	1.4	0.8	0.8	0.9	1.4	1.4	1.5	1.5	1.5	1.6	0.9	0.9	0.9
2003.....	1.0	1.0	1.0	0.6	0.7	0.5	1.0	1.0	1.1	1.1	1.0	1.1	0.7	0.7	0.7
2002.....	0.8	0.8	0.8	0.5	0.5	0.5	0.8	0.8	0.8	0.9	0.9	0.9	0.6	0.7	0.6
2001.....	0.5	0.5	0.5	0.4	*	0.4	0.5	0.5	0.5	0.5	0.5	0.5	0.4	0.4	0.4
2000.....	0.5	0.5	0.4	0.3	0.4	0.3	0.5	0.5	0.4	0.5	0.5	0.5	0.3	0.4	0.3
1999.....	0.4	0.4	0.4	0.2	*	*	0.4	0.4	0.4	0.4	0.4	0.4	0.3	0.3	0.3
Diarrhea and gastroenteritis of infectious origin (A09)															
2004.....	0.0	0.0	0.0	*	*	*	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2003.....	0.0	0.0	0.0	*	*	*	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2002.....	0.0	0.0	0.0	*	*	*	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2001.....	0.0	0.0	0.0	*	*	*	0.0	*	0.0	0.0	*	0.0	*	*	*
2000.....	0.0	0.0	0.0	*	*	*	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
1999.....	0.0	0.0	0.0	*	*	*	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
Tuberculosis (A16-A19)															
2004.....	0.2	0.3	0.1	0.5	0.8	0.3	0.2	0.3	0.1	0.1	0.2	0.1	0.7	1.0	0.4
2003.....	0.2	0.3	0.1	0.4	0.6	0.3	0.2	0.3	0.1	0.1	0.2	0.1	0.6	1.0	0.3
2002.....	0.3	0.4	0.2	0.5	0.6	0.4	0.3	0.4	0.2	0.1	0.2	0.1	0.8	1.1	0.5
2001.....	0.3	0.4	0.2	0.5	0.8	0.3	0.2	0.3	0.2	0.1	0.2	0.1	0.8	1.2	0.5
2000.....	0.3	0.4	0.2	0.5	0.7	0.4	0.2	0.4	0.2	0.1	0.2	0.1	0.9	1.5	0.6
1999.....	0.3	0.5	0.2	0.8	1.1	0.6	0.3	0.5	0.2	0.2	0.3	0.1	1.0	1.5	0.6
Respiratory tuberculosis (A16)															
2004.....	0.2	0.2	0.1	0.4	0.6	0.3	0.1	0.2	0.1	0.1	0.1	0.1	0.5	0.8	0.3
2003.....	0.2	0.3	0.1	0.4	0.6	0.2	0.2	0.2	0.1	0.1	0.1	0.1	0.5	0.8	0.3
2002.....	0.2	0.3	0.1	0.4	0.5	0.3	0.2	0.3	0.1	0.1	0.2	0.1	0.6	0.9	0.4
2001.....	0.2	0.3	0.1	0.4	0.7	0.2	0.2	0.3	0.1	0.1	0.2	0.1	0.6	1.0	0.3
2000.....	0.2	0.3	0.1	0.4	0.7	0.2	0.2	0.3	0.1	0.1	0.2	0.1	0.8	1.3	0.4
1999.....	0.3	0.4	0.2	0.6	0.8	0.5	0.3	0.4	0.1	0.2	0.3	0.1	0.8	1.2	0.5

Worktable Orig13. Age-adjusted death rates for 358 selected causes, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), and year	All origins 1/			Hispanic			Non-Hispanic 2/			Non-Hispanic white			Non-Hispanic black		
	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female
Herpes-viral (herpes simplex) infections (B00)															
2004.....	0.1	0.1	0.1	*	*	*	0.1	0.1	0.0	0.1	0.1	0.1	*	*	*
2003.....	0.1	0.0	0.1	*	*	*	0.1	0.1	0.1	0.1	0.1	0.1	*	*	*
2002.....	0.0	0.1	0.1	*	*	*	0.1	0.1	0.1	0.0	0.0	0.1	0.1	*	*
2001.....	0.1	0.1	0.1	*	*	*	0.1	0.1	0.1	0.1	0.1	0.1	0.0	*	*
2000.....	0.0	0.0	0.0	*	*	*	0.0	0.0	0.0	0.0	0.0	0.0	0.1	*	*
1999.....	0.0	0.0	0.0	*	*	*	0.0	0.0	0.0	0.0	0.0	0.0	0.1	*	*
Zoster (herpes zoster)(B02)															
2004.....	0.0	0.1	0.0	*	*	*	0.1	0.0	0.0	0.1	0.0	0.1	*	*	*
2003.....	0.0	0.0	0.0	*	*	*	0.0	0.0	0.0	0.1	0.0	0.1	*	*	*
2002.....	0.0	0.0	0.0	*	*	*	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2001.....	0.0	0.0	0.0	*	*	*	0.0	0.0	0.0	0.0	0.0	0.1	*	*	*
2000.....	0.0	0.0	0.0	*	*	*	0.0	0.0	0.1	0.0	0.0	0.0	*	*	*
1999.....	0.1	0.0	0.1	*	*	*	0.1	0.0	0.1	0.1	0.0	0.1	*	*	*
Smallpox (B03)															
2004.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Measles (B05)															
2004.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Rubella (German measles) (B06)															
2004.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Viral hepatitis (B15-B19)															
2004.....	1.8	2.4	1.2	2.8	3.5	2.0	1.7	2.3	1.1	1.5	2.1	1.0	2.7	3.8	1.7
2003.....	1.8	2.5	1.2	2.9	3.8	2.1	1.7	2.4	1.1	1.6	2.2	1.0	2.7	3.8	1.8
2002.....	2.0	2.7	1.3	3.3	4.2	2.4	1.9	2.6	1.2	1.7	2.4	1.1	2.8	3.9	1.9
2001.....	2.0	2.6	1.3	3.7	4.5	2.8	1.8	2.5	1.2	1.7	2.3	1.0	2.8	3.5	2.1
2000.....	1.9	2.6	1.3	3.6	4.6	2.6	1.8	2.4	1.2	1.6	2.2	1.1	2.7	3.8	1.8
1999.....	1.8	2.4	1.2	3.5	4.3	2.7	1.6	2.2	1.1	1.5	2.0	1.0	2.6	3.5	1.9
Human immunodeficiency virus (HIV) disease (B20-B24)															
2004.....	4.5	6.6	2.4	5.3	8.2	2.4	4.4	6.4	2.5	1.9	3.1	0.6	20.9	29.8	13.3
2003.....	4.7	7.1	2.4	5.9	9.2	2.7	4.6	6.8	2.4	2.0	3.4	0.6	21.7	32.0	13.1
2002.....	4.9	7.4	2.5	5.8	9.1	2.6	4.8	7.2	2.5	2.1	3.5	0.6	23.0	33.9	13.8
2001.....	5.0	7.5	2.5	6.2	9.7	2.7	4.8	7.3	2.5	2.1	3.6	0.6	23.2	34.4	13.8
2000.....	5.2	7.9	2.5	6.7	10.6	2.9	5.0	7.5	2.5	2.2	3.8	0.7	23.6	35.5	13.5
1999.....	5.3	8.2	2.5	6.9	10.9	3.0	5.1	7.8	2.5	2.3	4.0	0.7	24.0	36.5	13.4

Worktable Orig13. Age-adjusted death rates for 358 selected causes, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), and year	All origins 1/			Hispanic			Non-Hispanic 2/			Non-Hispanic white			Non-Hispanic black		
	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female
Human immunodeficiency virus (HIV) disease resulting in infectious and parasitic diseases (B20)															
2004.....	1.8	2.5	1.0	2.3	3.6	1.1	1.7	2.4	1.0	0.7	1.2	0.3	7.9	11.1	5.3
2003.....	1.8	2.7	0.9	2.6	4.0	1.2	1.7	2.6	0.9	0.8	1.3	0.3	7.8	11.7	4.6
2002.....	1.9	2.9	1.0	2.6	4.0	1.2	1.9	2.7	1.0	0.8	1.4	0.3	8.8	13.1	5.2
2001.....	1.9	2.9	1.0	2.8	4.3	1.3	1.8	2.7	1.0	0.8	1.4	0.3	8.7	12.8	5.3
2000.....	2.0	3.1	1.0	3.0	4.6	1.4	1.9	2.9	1.0	0.9	1.4	0.3	9.1	13.5	5.3
1999.....	2.1	3.3	1.1	3.0	4.6	1.5	2.0	3.1	1.0	0.9	1.5	0.3	9.5	14.3	5.3
Human immunodeficiency virus (HIV) disease resulting in malignant neoplasms (B21)															
2004.....	0.3	0.5	0.1	0.3	0.6	0.1	0.3	0.5	0.1	0.2	0.3	0.0	1.1	1.9	0.5
2003.....	0.3	0.5	0.1	0.4	0.7	0.1	0.3	0.5	0.1	0.2	0.3	0.0	1.1	1.6	0.6
2002.....	0.3	0.5	0.1	0.4	0.6	*	0.3	0.5	0.1	0.2	0.4	0.0	1.2	1.9	0.6
2001.....	0.3	0.6	0.1	0.4	0.7	0.1	0.3	0.5	0.1	0.2	0.4	0.1	1.2	2.0	0.6
2000.....	0.4	0.6	0.1	0.5	0.8	0.1	0.3	0.6	0.1	0.2	0.4	0.0	1.3	2.1	0.5
1999.....	0.4	0.7	0.1	0.6	1.0	0.2	0.4	0.6	0.1	0.2	0.4	0.0	1.3	2.2	0.5
Human immunodeficiency virus (HIV) disease resulting in other specified diseases (B22)															
2004.....	0.4	0.7	0.2	0.6	0.9	0.2	0.4	0.6	0.2	0.2	0.4	0.1	1.8	2.7	1.1
2003.....	0.4	0.7	0.2	0.6	1.0	0.2	0.4	0.7	0.2	0.2	0.4	0.1	1.9	3.0	1.0
2002.....	0.5	0.7	0.2	0.6	1.0	0.1	0.4	0.7	0.2	0.2	0.4	0.0	1.9	2.9	1.0
2001.....	0.5	0.8	0.2	0.7	1.2	0.2	0.5	0.8	0.2	0.2	0.4	0.1	2.1	3.2	1.2
2000.....	0.6	0.9	0.2	0.7	1.2	0.2	0.5	0.9	0.2	0.3	0.5	0.1	2.4	3.7	1.3
1999.....	0.6	0.9	0.2	0.7	1.2	0.2	0.5	0.9	0.2	0.3	0.5	0.1	2.3	3.7	1.2
Human immunodeficiency virus (HIV) disease resulting in other conditions (B23)															
2004.....	0.7	1.1	0.4	0.7	1.1	0.3	0.7	1.1	0.4	0.3	0.4	0.1	3.9	5.8	2.4
2003.....	1.1	1.6	0.6	1.3	2.0	0.7	1.1	1.6	0.7	0.4	0.6	0.1	5.9	8.4	3.8
2002.....	1.2	1.7	0.7	1.4	2.1	0.7	1.1	1.7	0.7	0.4	0.7	0.1	6.4	9.3	3.9
2001.....	1.2	1.8	0.7	1.5	2.2	0.8	1.2	1.7	0.7	0.4	0.7	0.1	6.3	9.2	3.9
2000.....	1.2	1.8	0.7	1.5	2.4	0.7	1.2	1.7	0.7	0.4	0.7	0.2	6.1	9.0	3.7
1999.....	1.2	1.8	0.7	1.5	2.2	0.8	1.2	1.8	0.7	0.4	0.7	0.2	6.5	9.6	3.8
Unspecified human immunodeficiency virus (HIV) disease (B24)															
2004.....	1.2	1.7	0.7	1.3	2.0	0.6	1.2	1.7	0.7	0.5	0.8	0.2	6.0	8.3	4.0
2003.....	1.0	1.5	0.6	1.0	1.4	0.5	1.0	1.5	0.5	0.4	0.7	0.1	5.0	7.3	3.0
2002.....	1.0	1.4	0.5	0.8	1.3	0.4	1.0	1.5	0.5	0.4	0.7	0.1	4.8	6.9	2.9
2001.....	1.0	1.5	0.5	0.8	1.3	0.3	1.0	1.5	0.5	0.4	0.8	0.1	4.8	7.1	2.8
2000.....	1.0	1.5	0.5	1.0	1.6	0.5	1.0	1.5	0.5	0.4	0.7	0.1	4.8	7.2	2.7
1999.....	1.0	1.5	0.5	1.2	1.9	0.4	0.9	1.4	0.5	0.4	0.7	0.1	4.5	6.7	2.6
All other and unspecified viral diseases (A81,A85.0-A85.1,A85.8,A86-A89,B01,B04,B07-B09,B25-B34)															
2004.....	0.3	0.4	0.3	0.3	0.4	0.2	0.3	0.4	0.3	0.3	0.4	0.3	0.3	0.3	0.3
2003.....	0.3	0.4	0.3	0.3	0.4	0.2	0.4	0.4	0.3	0.4	0.4	0.3	0.3	0.4	0.3
2002.....	0.4	0.4	0.3	0.2	0.2	0.2	0.4	0.4	0.3	0.4	0.4	0.3	0.3	0.4	0.3
2001.....	0.3	0.3	0.3	0.3	0.3	0.2	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3
2000.....	0.4	0.4	0.3	0.3	0.4	0.3	0.4	0.4	0.3	0.4	0.4	0.3	0.4	0.4	0.4
1999.....	0.4	0.4	0.4	0.3	0.4	0.3	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4
Other and unspecified infectious and parasitic diseases and their sequelae (B35-B99)															
2004.....	1.5	1.9	1.1	2.1	2.8	1.6	1.4	1.8	1.1	1.4	1.7	1.0	2.0	2.6	1.5
2003.....	1.5	1.8	1.2	2.1	2.7	1.5	1.4	1.7	1.1	1.3	1.6	1.0	2.0	2.5	1.6
2002.....	1.3	1.7	0.9	1.8	2.4	1.2	1.2	1.6	0.9	1.1	1.5	0.8	1.8	2.5	1.3
2001.....	1.1	1.3	0.9	1.2	1.5	1.0	1.1	1.3	0.9	1.0	1.2	0.8	1.7	2.1	1.4
2000.....	1.1	1.3	0.8	1.3	1.8	0.9	1.1	1.3	0.8	0.9	1.2	0.8	1.7	2.3	1.3
1999.....	1.5	1.8	1.2	1.6	1.8	1.5	1.5	1.8	1.2	1.2	1.6	1.0	2.9	3.4	2.6

Worktable Orig13. Age-adjusted death rates for 358 selected causes, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), and year	All origins 1/			Hispanic			Non-Hispanic 2/			Non-Hispanic white			Non-Hispanic black		
	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female
Helminthiasis (B65-B83)															
2004.....	0.0	0.0	*	*	*	*	0.0	*	*	*	*	*	*	*	*
2003.....	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000.....	0.0	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Schistosomiasis (bilharziasis) (B65)															
2004.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Other and unspecified helminthiasis (B66-B83)															
2004.....	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2003.....	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000.....	0.0	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Sequelae of tuberculosis (B90)															
2004.....	0.0	0.0	0.0	*	*	*	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2003.....	0.0	0.0	0.0	*	*	*	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2002.....	0.0	0.0	0.0	*	*	*	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2001.....	0.0	0.0	0.0	*	*	*	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2000.....	0.0	0.1	0.0	*	*	*	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
1999.....	0.1	0.1	0.1	0.1	*	*	0.1	0.1	0.1	0.0	0.1	0.0	0.1	*	*
Sequelae of poliomyelitis (B91)															
2004.....	0.1	0.1	0.1	*	*	*	0.1	0.1	0.1	0.1	0.1	0.1	*	*	*
2003.....	0.1	0.1	0.1	*	*	*	0.1	0.1	0.1	0.1	0.1	0.1	*	*	*
2002.....	0.1	0.1	0.1	*	*	*	0.1	0.1	0.1	0.1	0.1	0.1	*	*	*
2001.....	0.1	0.1	0.1	*	*	*	0.1	0.2	0.1	0.1	0.1	0.1	*	*	*
2000.....	0.1	0.1	0.1	*	*	*	0.1	0.1	0.1	0.1	0.1	0.1	*	*	*
1999.....	0.1	0.1	0.1	*	*	*	0.1	0.1	0.1	0.1	0.1	0.1	*	*	*
All other and unspecified infectious and parasitic diseases and their sequelae (B85-B89,B92-B99)															
2004.....	0.9	1.2	0.7	1.6	2.0	1.2	0.9	1.1	0.7	0.9	1.1	0.7	1.0	1.4	0.8
2003.....	0.9	1.1	0.7	1.5	1.9	1.1	0.8	1.0	0.6	0.8	1.0	0.6	1.1	1.4	0.8
2002.....	0.6	0.8	0.4	1.1	1.5	0.8	0.6	0.7	0.4	0.5	0.7	0.4	0.7	1.0	0.6
2001.....	0.4	0.5	0.4	0.6	0.7	0.5	0.4	0.5	0.3	0.4	0.5	0.3	0.6	0.6	0.5
2000.....	0.3	0.4	0.3	0.5	0.6	0.4	0.3	0.4	0.3	0.3	0.4	0.2	0.5	0.7	0.4
1999.....	0.5	0.6	0.5	0.5	0.5	0.4	0.5	0.6	0.5	0.4	0.5	0.3	1.2	1.4	1.1
II. Neoplasms (C00-D48)															
2004.....	190.4	233.3	161.2	125.0	154.8	104.2	194.8	238.4	165.1	193.4	235.1	164.8	235.6	311.6	189.6
2003.....	194.7	239.1	164.8	129.8	160.3	108.8	198.8	243.9	168.4	197.2	240.6	167.8	241.6	319.3	194.6
2002.....	198.1	244.6	167.0	131.5	165.1	108.8	202.1	249.2	170.6	200.4	245.5	170.0	247.0	329.7	197.5
2001.....	200.9	249.8	168.8	135.6	172.1	111.4	204.6	254.0	172.2	202.4	249.4	171.4	251.3	341.0	198.5
2000.....	204.5	255.0	171.8	138.4	175.5	114.1	208.1	259.2	175.0	205.6	253.9	174.3	256.7	350.3	201.0
1999.....	205.6	258.0	171.7	138.1	174.1	114.0	209.1	262.2	174.7	206.2	256.7	173.5	260.8	355.7	204.2

Worktable Orig13. Age-adjusted death rates for 358 selected causes, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), and year	All origins 1/			Hispanic			Non-Hispanic 2/			Non-Hispanic white			Non-Hispanic black		
	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female
Malignant neoplasms (C00-C97)															
2004.....	185.8	227.7	157.4	121.9	151.2	101.4	190.1	232.7	161.2	188.6	229.2	160.9	231.1	306.4	185.7
2003.....	190.1	233.3	160.9	126.6	156.5	105.9	194.1	238.1	164.4	192.4	234.6	163.8	237.3	314.2	190.8
2002.....	193.5	238.9	163.1	128.4	161.4	106.1	197.4	243.4	166.6	195.6	239.6	165.9	242.5	324.4	193.4
2001.....	196.0	243.7	164.7	132.3	168.2	108.6	199.7	247.9	168.0	197.4	243.1	167.2	246.7	335.3	194.4
2000.....	199.6	248.9	167.6	134.9	171.7	110.8	203.1	253.0	170.7	200.6	247.7	170.0	252.0	344.8	196.9
1999.....	200.8	251.9	167.6	134.8	170.6	110.9	204.2	256.0	170.6	201.2	250.4	169.4	255.9	350.3	199.8
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)															
2004.....	2.6	4.0	1.5	1.6	2.6	0.8	2.7	4.1	1.5	2.5	3.8	1.5	3.9	7.0	1.7
2003.....	2.6	4.0	1.5	1.7	2.7	0.9	2.7	4.2	1.5	2.6	3.9	1.5	3.9	7.0	1.7
2002.....	2.7	4.1	1.5	1.5	2.6	0.7	2.8	4.1	1.6	2.6	3.9	1.5	3.7	6.4	1.7
2001.....	2.7	4.1	1.6	1.8	2.9	1.0	2.8	4.2	1.6	2.6	3.9	1.6	3.9	6.8	1.8
2000.....	2.7	4.1	1.6	1.8	3.1	0.8	2.8	4.1	1.6	2.6	3.8	1.6	4.2	7.4	1.8
1999.....	2.7	4.1	1.6	1.9	3.0	1.0	2.8	4.2	1.6	2.6	3.9	1.6	4.4	7.6	2.1
Of lip (C00)															
2004.....	0.0	0.0	0.0	*	*	*	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2003.....	0.0	0.0	0.0	*	*	*	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2002.....	0.0	0.0	0.0	*	*	*	0.0	0.0	*	0.0	0.0	*	*	*	*
2001.....	0.0	0.0	*	*	*	*	0.0	0.0	*	0.0	0.0	*	*	*	*
2000.....	0.0	0.0	*	*	*	*	0.0	0.0	*	0.0	0.0	*	*	*	*
1999.....	0.0	0.0	*	*	*	*	0.0	0.0	*	0.0	0.0	*	*	*	*
Of tongue (C01-C02)															
2004.....	0.6	0.9	0.4	0.4	0.6	0.2	0.6	0.9	0.4	0.6	0.9	0.4	0.7	1.2	0.3
2003.....	0.6	0.9	0.4	0.5	0.7	0.3	0.7	1.0	0.4	0.7	1.0	0.4	0.8	1.3	0.4
2002.....	0.7	0.9	0.4	0.4	0.7	0.2	0.7	1.0	0.4	0.7	0.9	0.4	0.8	1.3	0.3
2001.....	0.6	0.9	0.4	0.4	0.6	0.4	0.7	1.0	0.4	0.6	0.9	0.4	0.7	1.2	0.3
2000.....	0.6	0.9	0.4	0.5	0.8	*	0.6	0.9	0.4	0.6	0.9	0.4	0.8	1.4	0.3
1999.....	0.6	0.9	0.4	0.5	0.8	0.2	0.7	1.0	0.4	0.6	0.9	0.4	0.9	1.4	0.4
Of pharynx (C10-C13,C14.0)															
2004.....	1.1	1.8	0.5	0.7	1.2	0.3	1.1	1.8	0.5	1.0	1.6	0.5	2.2	3.9	0.9
2003.....	1.1	1.8	0.5	0.7	1.4	0.3	1.1	1.8	0.5	1.0	1.6	0.5	2.1	3.9	0.7
2002.....	1.1	1.7	0.6	0.6	1.2	0.2	1.1	1.8	0.6	1.0	1.6	0.5	1.9	3.3	0.9
2001.....	1.1	1.8	0.6	0.8	1.2	0.4	1.2	1.9	0.6	1.0	1.6	0.5	2.0	3.6	0.9
2000.....	1.1	1.8	0.6	0.7	1.2	0.3	1.1	1.9	0.6	1.0	1.5	0.5	2.3	4.1	0.9
1999.....	1.1	1.8	0.6	0.8	1.3	0.3	1.1	1.9	0.6	1.0	1.6	0.5	2.3	4.1	1.0
Of other and unspecified sites within the lip, oral cavity and pharynx (C03-C09,C14.2-C14.8)															
2004.....	0.9	1.2	0.5	0.5	0.7	0.3	0.9	1.3	0.6	0.9	1.3	0.6	1.0	1.8	0.4
2003.....	0.9	1.3	0.5	0.4	0.6	0.3	0.9	1.3	0.5	0.9	1.3	0.6	1.1	1.8	0.6
2002.....	0.9	1.3	0.6	0.5	0.8	0.3	0.9	1.4	0.6	0.9	1.4	0.6	1.1	1.8	0.5
2001.....	0.9	1.3	0.6	0.6	1.0	0.3	0.9	1.3	0.6	0.9	1.3	0.6	1.2	1.9	0.6
2000.....	0.9	1.3	0.6	0.6	1.0	0.2	0.9	1.3	0.6	0.9	1.3	0.6	1.1	1.9	0.6
1999.....	0.9	1.3	0.6	0.6	0.9	0.4	1.0	1.3	0.7	0.9	1.3	0.7	1.3	2.1	0.7
Malignant neoplasms of digestive organs (C15-C26)															
2004.....	43.9	56.4	34.2	39.1	50.1	30.5	44.2	56.7	34.4	42.4	54.5	32.8	61.3	79.7	48.4
2003.....	45.1	57.7	35.3	40.2	51.0	31.7	45.3	58.0	35.4	43.3	55.7	33.7	63.4	82.3	50.3
2002.....	45.7	58.8	35.6	40.2	51.0	31.7	46.0	59.1	35.8	44.0	56.8	34.1	64.4	84.1	50.9
2001.....	46.4	59.4	36.5	41.7	52.9	33.4	46.6	59.6	36.6	44.6	57.3	34.8	65.7	84.4	52.8
2000.....	47.2	60.5	37.2	42.1	54.6	32.9	47.4	60.7	37.3	45.3	58.2	35.6	67.2	87.5	53.4
1999.....	47.3	60.7	37.3	41.9	53.0	33.5	47.5	61.0	37.4	45.4	58.4	35.7	67.5	88.2	53.4

Worktable Orig13. Age-adjusted death rates for 358 selected causes, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), and year	All origins 1/			Hispanic			Non-Hispanic 2/			Non-Hispanic white			Non-Hispanic black		
	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female
Of liver and intrahepatic bile ducts (C22)															
2004.....	5.1	7.4	3.2	7.6	10.9	4.9	4.9	7.1	3.1	4.4	6.3	2.8	7.0	10.9	4.2
2003.....	5.0	7.4	3.1	7.8	10.8	5.2	4.8	7.1	3.0	4.3	6.3	2.7	6.7	10.3	4.0
2002.....	4.9	7.1	3.0	7.5	10.7	4.9	4.7	6.9	2.9	4.2	6.2	2.7	6.4	10.0	3.7
2001.....	4.7	6.9	3.0	7.3	10.4	4.9	4.6	6.7	2.9	4.1	6.0	2.6	6.3	9.2	4.1
2000.....	4.7	6.8	3.0	7.5	10.7	4.9	4.5	6.6	2.9	4.0	5.8	2.6	6.2	9.7	3.6
1999.....	4.5	6.5	3.0	7.2	10.0	5.0	4.4	6.3	2.8	3.9	5.6	2.6	6.0	8.8	3.9
Liver (C22.0,C22.2-C22.9)															
2004.....	3.9	6.1	2.2	6.3	9.5	3.6	3.8	5.8	2.1	3.2	5.0	1.8	6.0	9.8	3.2
2003.....	3.9	6.0	2.1	6.4	9.5	3.8	3.7	5.7	2.0	3.2	5.0	1.7	5.6	9.1	3.1
2002.....	3.8	5.9	2.1	6.4	9.6	3.9	3.7	5.7	2.0	3.2	5.0	1.8	5.4	8.8	2.9
2001.....	3.7	5.7	2.1	6.2	9.3	3.8	3.5	5.5	2.0	3.1	4.9	1.7	5.4	8.2	3.3
2000.....	3.7	5.7	2.1	6.3	9.6	3.8	3.5	5.4	2.0	3.0	4.7	1.7	5.3	8.7	2.9
1999.....	3.6	5.4	2.1	6.1	9.0	3.8	3.4	5.2	2.0	3.0	4.6	1.8	5.2	8.0	3.2
Intrahepatic bile duct carcinoma (C22.1)															
2004.....	1.2	1.3	1.1	1.3	1.5	1.2	1.2	1.3	1.0	1.1	1.3	1.0	1.0	1.1	1.0
2003.....	1.1	1.3	1.0	1.3	1.4	1.3	1.1	1.3	0.9	1.1	1.3	0.9	1.1	1.3	0.9
2002.....	1.0	1.2	0.9	1.1	1.2	1.0	1.1	1.2	0.9	1.0	1.2	0.9	0.9	1.2	0.8
2001.....	1.0	1.2	0.9	1.2	1.1	1.2	1.0	1.2	0.9	1.0	1.2	0.8	0.9	1.1	0.8
2000.....	1.0	1.1	0.9	1.1	1.2	1.1	1.0	1.1	0.8	1.0	1.2	0.8	0.8	0.9	0.8
1999.....	0.9	1.0	0.9	1.1	1.0	1.2	0.9	1.0	0.8	0.9	1.0	0.8	0.8	0.9	0.7
Of gallbladder and extrahepatic bile ducts (C23-C24)															
2004.....	1.1	1.0	1.2	1.4	1.2	1.5	1.1	1.0	1.2	1.1	1.0	1.2	1.2	1.0	1.4
2003.....	1.2	1.1	1.2	1.4	1.1	1.6	1.1	1.1	1.2	1.1	1.1	1.2	1.1	0.9	1.3
2002.....	1.2	1.1	1.3	1.6	1.4	1.9	1.1	1.1	1.2	1.1	1.1	1.2	1.3	1.1	1.5
2001.....	1.3	1.2	1.4	1.9	1.5	2.1	1.2	1.1	1.3	1.2	1.1	1.3	1.3	1.1	1.5
2000.....	1.3	1.2	1.4	1.9	1.5	2.2	1.3	1.2	1.4	1.3	1.2	1.3	1.2	0.9	1.4
1999.....	1.3	1.2	1.4	2.0	1.3	2.6	1.3	1.2	1.3	1.2	1.2	1.3	1.3	1.0	1.4
Of pancreas (C25)															
2004.....	10.6	12.3	9.3	8.2	9.1	7.4	10.8	12.5	9.4	10.6	12.3	9.2	13.8	15.7	12.2
2003.....	10.5	12.0	9.3	8.2	9.2	7.4	10.6	12.2	9.4	10.4	12.0	9.1	13.9	15.6	12.6
2002.....	10.5	12.1	9.2	7.9	8.4	7.4	10.7	12.3	9.3	10.5	12.2	9.0	13.8	15.5	12.5
2001.....	10.5	12.1	9.3	8.4	9.0	7.8	10.7	12.3	9.4	10.5	12.2	9.2	13.8	15.1	12.6
2000.....	10.6	12.2	9.3	8.4	9.4	7.5	10.7	12.3	9.4	10.5	12.2	9.1	14.3	16.0	12.9
1999.....	10.6	12.3	9.3	7.9	8.8	7.2	10.8	12.4	9.4	10.5	12.2	9.2	14.3	16.7	12.5
Of other and ill-defined digestive organs (C26)															
2004.....	0.4	0.4	0.3	0.3	0.4	0.2	0.4	0.4	0.3	0.4	0.4	0.3	0.4	0.5	0.4
2003.....	0.5	0.5	0.4	0.4	0.4	0.4	0.5	0.6	0.4	0.4	0.5	0.4	0.6	0.8	0.5
2002.....	0.5	0.5	0.4	0.4	0.4	0.3	0.5	0.6	0.4	0.5	0.5	0.4	0.6	0.9	0.5
2001.....	0.5	0.6	0.4	0.3	0.5	0.3	0.5	0.6	0.4	0.5	0.6	0.4	0.7	0.9	0.6
2000.....	0.5	0.6	0.4	0.4	0.6	0.3	0.5	0.6	0.4	0.5	0.6	0.4	0.7	0.7	0.6
1999.....	0.6	0.7	0.5	0.5	0.7	0.4	0.6	0.7	0.5	0.6	0.7	0.5	0.7	0.9	0.5
Malignant neoplasms of respiratory and intrathoracic organs (C30-C39)															
2004.....	54.8	72.7	41.6	23.6	35.9	14.6	56.9	75.2	43.5	57.4	74.5	44.8	63.5	96.8	41.6
2003.....	55.7	74.5	42.0	24.3	36.7	15.2	57.8	76.9	43.8	58.3	76.2	45.1	64.9	100.0	41.9
2002.....	56.5	76.0	42.3	24.9	38.2	15.2	58.5	78.4	44.1	59.0	77.6	45.4	65.8	102.0	41.9
2001.....	57.0	78.1	41.8	25.1	39.2	15.1	59.0	80.4	43.5	59.3	79.3	44.7	66.4	106.1	40.6
2000.....	57.8	79.7	42.1	25.9	41.2	15.1	59.7	81.9	43.7	59.8	80.7	44.8	68.2	108.7	41.7
1999.....	57.2	79.8	41.1	26.5	42.4	15.1	58.9	81.8	42.5	58.9	80.4	43.4	68.7	109.9	41.7

Worktable Orig13. Age-adjusted death rates for 358 selected causes, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), and year	All origins 1/			Hispanic			Non-Hispanic 2/			Non-Hispanic white			Non-Hispanic black		
	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female
Of nasal cavity, middle ear and accessory sinuses (C30-C31)															
2004.....	0.2	0.2	0.1	*	*	*	0.2	0.2	0.1	0.2	0.2	0.1	0.2	0.3	*
2003.....	0.1	0.2	0.1	0.1	0.2	*	0.2	0.2	0.1	0.2	0.2	0.1	0.2	0.3	*
2002.....	0.2	0.2	0.1	0.2	*	*	0.2	0.2	0.1	0.2	0.2	0.1	0.1	0.2	*
2001.....	0.2	0.2	0.1	0.1	*	*	0.2	0.2	0.1	0.2	0.2	0.1	0.2	0.3	*
2000.....	0.1	0.2	0.1	0.1	*	*	0.1	0.2	0.1	0.1	0.2	0.1	0.2	0.3	0.1
1999.....	0.2	0.2	0.1	*	*	*	0.2	0.2	0.1	0.2	0.2	0.1	0.2	0.2	*
Of larynx (C32)															
2004.....	1.2	2.2	0.4	1.0	2.1	*	1.2	2.2	0.5	1.2	2.1	0.4	2.3	4.6	0.7
2003.....	1.3	2.4	0.5	0.9	1.9	*	1.3	2.4	0.5	1.2	2.2	0.5	2.5	5.2	0.7
2002.....	1.3	2.3	0.5	0.9	1.7	0.3	1.3	2.4	0.5	1.2	2.2	0.5	2.5	4.9	0.9
2001.....	1.3	2.5	0.5	1.0	1.9	0.2	1.4	2.5	0.5	1.2	2.3	0.5	2.6	5.0	0.9
2000.....	1.4	2.5	0.5	0.9	1.9	*	1.4	2.6	0.5	1.3	2.3	0.5	2.8	5.5	1.0
1999.....	1.4	2.5	0.5	1.2	2.5	0.3	1.4	2.5	0.5	1.3	2.3	0.5	2.6	5.2	0.8
Of trachea, bronchus and lung (C33-C34)															
2004.....	53.2	70.1	40.9	22.4	33.4	14.3	55.4	72.6	42.8	56.0	72.1	44.1	60.9	91.7	40.6
2003.....	54.1	71.7	41.3	23.2	34.5	14.8	56.2	74.1	43.1	56.7	73.6	44.3	62.0	94.2	40.9
2002.....	54.9	73.2	41.6	23.7	36.2	14.6	56.9	75.6	43.3	57.5	75.0	44.6	62.9	96.5	40.8
2001.....	55.3	75.2	41.0	23.8	36.8	14.5	57.2	77.4	42.7	57.7	76.6	44.0	63.4	100.5	39.4
2000.....	56.1	76.7	41.3	24.8	39.0	14.7	57.9	78.9	42.9	58.2	77.9	44.1	64.9	102.6	40.4
1999.....	55.5	76.9	40.2	25.0	39.6	14.6	57.2	78.8	41.7	57.2	77.6	42.7	65.7	104.3	40.6
Of thymus, heart, mediastinum and pleura (C37-C38)															
2004.....	0.2	0.2	0.1	0.2	0.2	*	0.2	0.2	0.1	0.1	0.2	0.1	0.2	0.2	0.2
2003.....	0.2	0.2	0.1	0.1	*	*	0.2	0.2	0.1	0.1	0.2	0.1	0.2	0.2	0.1
2002.....	0.2	0.2	0.1	0.2	*	*	0.2	0.2	0.1	0.2	0.2	0.1	0.2	0.2	0.1
2001.....	0.2	0.2	0.1	0.2	0.3	*	0.2	0.2	0.1	0.2	0.2	0.1	0.2	0.2	0.2
2000.....	0.1	0.2	0.1	0.1	*	*	0.1	0.2	0.1	0.1	0.2	0.1	0.2	0.3	0.2
1999.....	0.1	0.2	0.1	0.1	*	*	0.2	0.2	0.1	0.1	0.2	0.1	0.2	0.2	0.2
Of other and ill-defined sites in respiratory system and intrathoracic organs (C39)															
2004.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2003.....	0.0	0.0	0.0	*	*	*	0.0	0.0	*	0.0	0.0	*	*	*	*
2002.....	0.0	0.0	0.0	*	*	*	0.0	0.0	0.0	0.0	0.0	*	*	*	*
2001.....	0.0	0.0	*	*	*	*	0.0	0.0	*	0.0	0.0	*	*	*	*
2000.....	0.0	0.0	*	*	*	*	0.0	0.0	*	0.0	0.0	*	*	*	*
1999.....	0.0	0.0	*	*	*	*	0.0	0.0	*	0.0	0.0	*	*	*	*
Malignant neoplasms of bone and articular cartilage (C40-C41)															
2004.....	0.4	0.5	0.4	0.4	0.3	0.4	0.4	0.5	0.4	0.5	0.5	0.3	0.5	0.6	0.5
2003.....	0.4	0.5	0.3	0.4	0.5	0.2	0.4	0.5	0.3	0.4	0.6	0.4	0.4	0.5	0.4
2002.....	0.4	0.5	0.3	0.4	0.5	0.2	0.4	0.5	0.3	0.4	0.5	0.3	0.4	0.5	0.3
2001.....	0.5	0.6	0.4	0.4	0.5	0.4	0.5	0.6	0.4	0.5	0.6	0.4	0.5	0.6	0.4
2000.....	0.4	0.6	0.3	0.4	0.4	0.4	0.4	0.6	0.3	0.4	0.6	0.3	0.4	0.6	0.3
1999.....	0.4	0.5	0.4	0.4	0.7	0.3	0.4	0.5	0.4	0.4	0.5	0.4	0.4	0.5	0.3
Melanoma and other malignant neoplasms of skin (C43-C44)															
2004.....	3.5	5.3	2.1	0.9	1.2	0.7	3.6	5.6	2.2	4.1	6.2	2.5	0.9	1.4	0.6
2003.....	3.5	5.2	2.2	1.2	1.7	0.8	3.6	5.5	2.3	4.1	6.2	2.6	0.9	1.2	0.7
2002.....	3.4	5.1	2.1	1.2	1.5	0.9	3.6	5.4	2.2	4.0	6.0	2.6	1.0	1.5	0.6
2001.....	3.5	5.3	2.2	1.2	1.6	0.9	3.7	5.6	2.3	4.2	6.2	2.6	1.0	1.4	0.7
2000.....	3.5	5.2	2.2	1.2	1.9	0.8	3.6	5.4	2.3	4.1	6.0	2.6	1.1	1.5	0.9
1999.....	3.5	5.2	2.2	1.2	1.6	0.9	3.6	5.5	2.3	4.0	6.0	2.6	1.0	1.6	0.5

Worktable Orig13. Age-adjusted death rates for 358 selected causes, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), and year	All origins 1/			Hispanic			Non-Hispanic 2/			Non-Hispanic white			Non-Hispanic black		
	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female
Malignant melanoma of skin (C43)															
2004.....	2.7	3.9	1.7	0.6	0.7	0.5	2.8	4.2	1.8	3.3	4.8	2.1	0.4	0.5	0.4
2003.....	2.7	3.9	1.8	0.8	1.0	0.6	2.8	4.1	1.8	3.2	4.7	2.1	0.5	0.4	0.4
2002.....	2.6	3.8	1.7	0.8	0.8	0.7	2.7	4.0	1.8	3.1	4.5	2.1	0.4	0.5	0.3
2001.....	2.7	3.9	1.8	0.8	0.9	0.7	2.8	4.1	1.8	3.2	4.7	2.1	0.4	0.3	0.5
2000.....	2.7	3.8	1.8	0.8	1.1	0.5	2.8	4.0	1.9	3.2	4.5	2.2	0.6	0.6	0.5
1999.....	2.6	3.8	1.7	0.8	1.0	0.6	2.8	4.0	1.8	3.1	4.5	2.1	0.4	0.5	0.3
Other malignant neoplasms of skin (C44)															
2004.....	0.8	1.3	0.4	0.3	0.4	0.2	0.8	1.4	0.4	0.9	1.5	0.4	0.5	0.8	0.2
2003.....	0.8	1.4	0.4	0.4	0.7	0.2	0.8	1.4	0.4	0.9	1.5	0.5	0.5	0.7	0.3
2002.....	0.8	1.4	0.4	0.4	0.7	*	0.8	1.4	0.4	0.9	1.5	0.5	0.6	1.0	0.3
2001.....	0.9	1.4	0.5	0.4	0.7	*	0.9	1.5	0.5	0.9	1.5	0.5	0.6	1.0	0.2
2000.....	0.8	1.4	0.4	0.4	0.7	0.2	0.8	1.4	0.4	0.9	1.5	0.4	0.6	0.9	0.4
1999.....	0.8	1.4	0.4	0.4	0.6	0.2	0.9	1.5	0.4	0.9	1.6	0.5	0.6	1.1	0.2
Malignant neoplasms of mesothelial and soft tissue (C45-C49)															
2004.....	2.4	3.2	1.8	1.4	1.9	1.1	2.5	3.3	1.9	2.6	3.5	1.9	2.0	2.3	1.8
2003.....	2.4	3.2	1.8	1.7	2.1	1.4	2.4	3.2	1.8	2.5	3.4	1.9	2.0	2.5	1.7
2002.....	2.4	3.2	1.8	1.7	2.3	1.3	2.4	3.3	1.8	2.5	3.4	1.8	2.0	2.4	1.8
2001.....	2.4	3.2	1.8	1.6	2.1	1.3	2.4	3.3	1.8	2.5	3.5	1.9	1.9	2.2	1.7
2000.....	2.5	3.3	1.8	1.5	2.0	1.2	2.5	3.4	1.9	2.6	3.6	1.9	2.1	2.6	1.9
1999.....	2.4	3.3	1.8	1.7	1.9	1.5	2.5	3.4	1.8	2.5	3.5	1.8	2.2	2.6	1.9
Mesothelioma (C45)															
2004.....	0.8	1.6	0.3	0.4	0.7	*	0.9	1.7	0.3	0.9	1.8	0.3	0.3	0.8	*
2003.....	0.8	1.6	0.3	0.6	1.1	0.2	0.9	1.7	0.3	0.9	1.8	0.3	0.4	0.9	*
2002.....	0.9	1.7	0.2	0.5	1.0	*	0.9	1.7	0.2	0.9	1.9	0.3	0.4	0.8	*
2001.....	0.8	1.6	0.3	0.4	0.8	*	0.9	1.7	0.3	1.0	1.8	0.3	0.2	0.5	*
2000.....	0.9	1.7	0.3	0.4	0.7	*	0.9	1.7	0.3	0.9	1.9	0.3	0.4	0.8	0.1
1999.....	0.9	1.6	0.3	0.4	0.7	0.2	0.9	1.7	0.3	0.9	1.8	0.3	0.4	0.8	0.2
Kaposi's sarcoma (C46)															
2004.....	0.0	0.0	*	*	*	*	0.0	0.0	*	0.0	0.0	*	*	*	*
2003.....	0.0	0.0	0.0	*	*	*	0.0	0.0	*	0.0	0.0	*	*	*	*
2002.....	0.0	0.0	0.0	*	*	*	0.0	0.0	*	0.0	*	*	*	*	*
2001.....	0.0	0.0	*	*	*	*	0.0	0.0	*	0.0	*	*	*	*	*
2000.....	0.0	0.0	*	*	*	*	0.0	0.0	*	0.0	*	*	*	*	*
1999.....	0.0	0.0	*	*	*	*	0.0	0.0	*	0.0	0.0	*	*	*	*
Of other mesothelial and soft tissue (C47-C49)															
2004.....	1.5	1.6	1.5	1.0	1.0	1.0	1.6	1.6	1.6	1.6	1.6	1.6	1.6	1.6	1.7
2003.....	1.5	1.5	1.5	1.1	1.0	1.1	1.5	1.5	1.5	1.5	1.6	1.5	1.6	1.6	1.7
2002.....	1.5	1.5	1.5	1.1	1.2	1.1	1.5	1.5	1.5	1.5	1.6	1.5	1.7	1.6	1.6
2001.....	1.5	1.5	1.5	1.1	1.2	1.1	1.5	1.6	1.5	1.6	1.6	1.5	1.6	1.7	1.6
2000.....	1.6	1.6	1.5	1.1	1.2	1.0	1.6	1.7	1.6	1.6	1.7	1.6	1.6	1.6	1.7
1999.....	1.6	1.6	1.5	1.2	1.3	1.2	1.6	1.7	1.5	1.6	1.7	1.5	1.8	1.8	1.7
Malignant neoplasm of breast (C50)															
2004.....	13.8	0.3	24.4	8.7	*	15.6	14.1	0.3	25.1	13.7	0.3	24.5	19.4	0.5	32.9
2003.....	14.2	0.3	25.3	8.9	*	16.1	14.6	0.3	25.9	14.2	0.3	25.2	20.5	0.5	34.6
2002.....	14.5	0.3	25.6	8.6	*	15.5	14.8	0.3	26.2	14.4	0.3	25.6	20.5	0.6	34.6
2001.....	14.8	0.3	26.0	9.1	*	16.3	15.1	0.4	26.6	14.7	0.3	26.0	20.8	0.6	35.1
2000.....	15.3	0.4	26.8	9.5	*	16.9	15.6	0.4	27.3	15.2	0.4	26.8	20.9	0.6	35.1
1999.....	15.2	0.3	26.6	9.2	*	16.4	15.5	0.3	27.2	15.1	0.3	26.6	21.2	0.7	35.7

Worktable Orig13. Age-adjusted death rates for 358 selected causes, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), and year	All origins 1/			Hispanic			Non-Hispanic 2/			Non-Hispanic white			Non-Hispanic black		
	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female
Malignant neoplasms of female genital organs (C51-C58)															
2004.....	9.0	...	16.2	7.0	...	12.6	9.2	...	16.4	9.0	...	16.2	12.0	...	20.3
2003.....	9.2	...	16.4	7.0	...	12.6	9.3	...	16.6	9.2	...	16.4	12.1	...	20.3
2002.....	9.4	...	16.7	7.3	...	13.1	9.5	...	16.9	9.3	...	16.7	12.4	...	20.8
2001.....	9.5	...	16.8	7.6	...	13.5	9.6	...	17.0	9.4	...	16.7	12.3	...	20.6
2000.....	9.5	...	16.8	8.1	...	14.3	9.6	...	16.9	9.4	...	16.7	12.7	...	21.2
1999.....	9.4	...	16.7	7.3	...	13.0	9.5	...	16.8	9.4	...	16.7	12.4	...	20.8
Of vulva and vagina (C51-C52)															
2004.....	0.4	...	0.7	0.3	...	0.5	0.4	...	0.7	0.4	...	0.7	0.4	...	0.7
2003.....	0.4	...	0.7	0.3	...	0.5	0.4	...	0.7	0.4	...	0.7	0.4	...	0.6
2002.....	0.4	...	0.7	0.3	...	0.5	0.4	...	0.7	0.4	...	0.7	0.4	...	0.6
2001.....	0.4	...	0.7	0.3	...	0.5	0.4	...	0.7	0.4	...	0.7	0.5	...	0.8
2000.....	0.4	...	0.7	0.4	...	0.6	0.4	...	0.7	0.4	...	0.7	0.4	...	0.7
1999.....	0.4	...	0.7	0.4	...	0.6	0.4	...	0.7	0.4	...	0.7	0.4	...	0.7
Of cervix uteri (C53)															
2004.....	1.3	...	2.4	1.7	...	3.1	1.3	...	2.4	1.1	...	2.1	2.7	...	4.6
2003.....	1.3	...	2.5	1.6	...	3.0	1.3	...	2.4	1.1	...	2.2	2.7	...	4.8
2002.....	1.3	...	2.6	1.7	...	3.1	1.3	...	2.5	1.1	...	2.2	2.9	...	5.0
2001.....	1.4	...	2.7	1.8	...	3.4	1.4	...	2.6	1.2	...	2.3	2.8	...	4.9
2000.....	1.5	...	2.8	2.2	...	4.0	1.5	...	2.7	1.2	...	2.3	3.2	...	5.5
1999.....	1.5	...	2.8	1.9	...	3.6	1.5	...	2.8	1.3	...	2.5	3.2	...	5.4
Of corpus uteri and uterus, part unspecified (C54-C55)															
2004.....	2.3	...	4.1	1.6	...	2.9	2.4	...	4.2	2.2	...	3.9	4.2	...	7.0
2003.....	2.3	...	4.1	1.7	...	3.0	2.4	...	4.2	2.2	...	3.9	4.3	...	7.1
2002.....	2.4	...	4.2	1.9	...	3.3	2.4	...	4.2	2.2	...	3.9	4.5	...	7.4
2001.....	2.4	...	4.2	1.9	...	3.3	2.4	...	4.2	2.3	...	4.0	4.3	...	7.0
2000.....	2.4	...	4.1	1.8	...	3.1	2.4	...	4.2	2.2	...	3.9	4.5	...	7.4
1999.....	2.4	...	4.1	1.8	...	3.1	2.4	...	4.1	2.3	...	3.9	4.2	...	6.9
Of ovary (C56)															
2004.....	4.9	...	8.8	3.3	...	5.9	5.1	...	9.0	5.2	...	9.3	4.6	...	7.8
2003.....	5.0	...	8.8	3.3	...	5.9	5.1	...	9.1	5.3	...	9.4	4.5	...	7.5
2002.....	5.1	...	9.0	3.3	...	6.0	5.2	...	9.2	5.4	...	9.6	4.5	...	7.5
2001.....	5.1	...	9.0	3.5	...	6.1	5.2	...	9.2	5.4	...	9.5	4.6	...	7.6
2000.....	5.1	...	8.9	3.7	...	6.4	5.2	...	9.1	5.3	...	9.5	4.4	...	7.3
1999.....	5.0	...	8.8	3.1	...	5.5	5.1	...	9.0	5.2	...	9.3	4.4	...	7.4
Of other and unspecified female genital organs (C57)															
2004.....	0.1	...	0.1	*	...	*	0.1	...	0.1	0.1	...	0.1	0.1	...	0.1
2003.....	0.1	...	0.2	*	...	*	0.1	...	0.2	0.1	...	0.2	0.2	...	0.3
2002.....	0.1	...	0.3	*	...	*	0.1	...	0.3	0.1	...	0.3	0.2	...	0.3
2001.....	0.1	...	0.2	*	...	*	0.1	...	0.2	0.1	...	0.3	0.1	...	0.2
2000.....	0.1	...	0.2	*	...	*	0.1	...	0.2	0.1	...	0.2	0.2	...	0.3
1999.....	0.1	...	0.3	*	...	*	0.1	...	0.3	0.1	...	0.3	0.2	...	0.3
Of placenta (C58)															
2004.....	0.0	...	0.0	*	...	*	*	...	*	*	...	*	*	...	*
2003.....	*	...	*	*	...	*	*	...	*	*	...	*	*	...	*
2002.....	*	...	*	*	...	*	*	...	*	*	...	*	*	...	*
2001.....	*	...	*	*	...	*	*	...	*	*	...	*	*	...	*
2000.....	*	...	*	*	...	*	*	...	*	*	...	*	*	...	*
1999.....	*	...	*	*	...	*	*	...	*	*	...	*	*	...	*

Worktable Orig13. Age-adjusted death rates for 358 selected causes, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), and year	All origins 1/			Hispanic			Non-Hispanic 2/			Non-Hispanic white			Non-Hispanic black		
	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female
Malignant neoplasms of male genital organs (C60-C63)															
2004.....	10.0	25.9	...	7.9	19.7	...	10.1	26.2	...	9.3	24.1	...	20.2	56.8	...
2003.....	10.3	26.9	...	8.2	20.7	...	10.4	27.2	...	9.6	25.1	...	20.9	58.8	...
2002.....	10.8	28.4	...	8.7	22.2	...	10.9	28.7	...	10.0	26.3	...	22.4	63.2	...
2001.....	11.1	29.6	...	9.4	24.1	...	11.2	29.8	...	10.2	27.1	...	23.5	67.2	...
2000.....	11.5	30.8	...	8.6	22.2	...	11.6	31.2	...	10.6	28.5	...	24.3	69.2	...
1999.....	11.9	31.8	...	9.1	23.6	...	12.0	32.1	...	11.0	29.4	...	24.7	70.1	...
Of prostate (C61)															
2004.....	9.8	25.4	...	7.6	19.1	...	9.9	25.7	...	9.1	23.6	...	20.0	56.4	...
2003.....	10.1	26.5	...	7.9	20.2	...	10.2	26.8	...	9.4	24.6	...	20.7	58.2	...
2002.....	10.6	27.9	...	8.4	21.6	...	10.7	28.2	...	9.8	25.8	...	22.2	62.8	...
2001.....	10.9	29.1	...	9.1	23.5	...	11.0	29.3	...	10.0	26.7	...	23.3	66.8	...
2000.....	11.3	30.4	...	8.3	21.6	...	11.4	30.7	...	10.3	28.0	...	24.1	68.8	...
1999.....	11.6	31.3	...	8.8	23.0	...	11.7	31.7	...	10.7	28.9	...	24.5	69.7	...
Of testis (C62)															
2004.....	0.1	0.3	...	0.1	0.2	...	0.1	0.3	...	0.1	0.3	...	0.1	0.2	...
2003.....	0.1	0.2	...	0.1	0.2	...	0.1	0.2	...	0.1	0.3	...	0.1	0.2	...
2002.....	0.1	0.3	...	0.1	0.3	...	0.1	0.3	...	0.2	0.3	...	0.1	0.2	...
2001.....	0.1	0.2	...	0.1	0.3	...	0.1	0.2	...	0.1	0.3	...	0.1	0.2	...
2000.....	0.1	0.3	...	0.1	0.2	...	0.1	0.3	...	0.1	0.3	...	0.1	0.2	...
1999.....	0.1	0.3	...	0.1	0.2	...	0.1	0.3	...	0.2	0.3	...	0.1	0.1	...
Of penis and other and unspecified male genital organs (C60,C63)															
2004.....	0.1	0.2	...	0.2	0.4	...	0.1	0.2	...	0.1	0.2	...	0.1	0.2	...
2003.....	0.1	0.2	...	0.1	0.3	...	0.1	0.2	...	0.1	0.2	...	0.1	0.4	...
2002.....	0.1	0.2	...	0.1	0.4	...	0.1	0.2	...	0.1	0.2	...	0.1	0.2	...
2001.....	0.1	0.2	...	0.1	0.3	...	0.1	0.2	...	0.1	0.2	...	0.1	0.3	...
2000.....	0.1	0.2	...	0.2	0.3	...	0.1	0.2	...	0.1	0.2	...	0.1	0.3	...
1999.....	0.1	0.2	...	0.2	0.4	...	0.1	0.2	...	0.1	0.2	...	0.1	0.3	...
Malignant neoplasms of urinary tract (C64-C68)															
2004.....	8.7	13.7	5.1	6.0	9.0	3.8	8.8	14.0	5.2	9.2	14.6	5.3	7.9	11.1	5.7
2003.....	8.7	13.7	5.1	6.2	9.1	4.0	8.8	13.9	5.2	9.1	14.5	5.3	8.1	11.8	5.8
2002.....	8.9	13.9	5.3	6.2	9.6	3.8	9.0	14.1	5.4	9.2	14.6	5.4	8.3	11.7	6.2
2001.....	8.8	14.1	5.2	6.2	9.8	3.7	9.0	14.3	5.3	9.2	14.8	5.3	8.3	11.9	6.1
2000.....	8.8	14.1	5.3	5.9	9.4	3.5	8.9	14.3	5.3	9.2	14.7	5.4	8.1	12.2	5.7
1999.....	8.7	13.8	5.2	6.0	9.4	3.5	8.8	14.0	5.2	8.9	14.4	5.2	8.4	12.1	6.1
Of kidney and renal pelvis (C64-C65)															
2004.....	4.1	5.9	2.7	3.7	5.5	2.2	4.1	5.9	2.8	4.2	6.1	2.8	4.1	5.9	2.8
2003.....	4.2	6.0	2.7	3.7	5.0	2.6	4.2	6.1	2.7	4.3	6.2	2.8	4.1	6.1	2.8
2002.....	4.2	6.1	2.8	3.6	5.3	2.3	4.2	6.2	2.8	4.4	6.3	2.9	4.1	5.9	2.9
2001.....	4.3	6.2	2.8	3.7	5.6	2.3	4.3	6.2	2.8	4.4	6.3	2.9	4.3	6.5	2.9
2000.....	4.2	6.2	2.8	3.5	5.3	2.2	4.3	6.2	2.8	4.3	6.3	2.9	4.2	6.3	2.8
1999.....	4.1	5.9	2.7	3.6	5.1	2.4	4.1	5.9	2.7	4.1	6.0	2.7	4.2	6.2	2.8
Of bladder (C67)															
2004.....	4.4	7.6	2.2	2.2	3.5	1.4	4.5	7.8	2.3	4.7	8.2	2.3	3.6	5.0	2.8
2003.....	4.3	7.3	2.2	2.4	4.0	1.4	4.4	7.5	2.3	4.5	7.9	2.3	3.8	5.5	2.7
2002.....	4.4	7.5	2.4	2.5	4.2	1.5	4.5	7.7	2.4	4.6	8.0	2.4	4.0	5.5	3.1
2001.....	4.3	7.5	2.3	2.4	4.0	1.4	4.4	7.7	2.3	4.6	8.1	2.3	3.8	5.1	2.9
2000.....	4.3	7.6	2.3	2.3	4.1	1.2	4.4	7.8	2.3	4.6	8.1	2.3	3.8	5.6	2.7
1999.....	4.4	7.6	2.3	2.3	4.1	1.1	4.4	7.8	2.3	4.6	8.1	2.3	4.0	5.8	3.0

Worktable Orig13. Age-adjusted death rates for 358 selected causes, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), and year	All origins 1/			Hispanic			Non-Hispanic 2/			Non-Hispanic white			Non-Hispanic black		
	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female
Of other and unspecified sites within the urinary tract (C66,C68)															
2004.....	0.2	0.3	0.2	*	*	*	0.2	0.3	0.2	0.2	0.3	0.1	0.2	*	0.2
2003.....	0.2	0.3	0.2	*	*	*	0.2	0.3	0.2	0.2	0.3	0.2	0.2	*	0.2
2002.....	0.2	0.3	0.2	*	*	*	0.2	0.3	0.2	0.2	0.3	0.2	0.2	*	0.2
2001.....	0.2	0.3	0.2	*	*	*	0.2	0.3	0.2	0.2	0.3	0.2	0.2	0.2	0.2
2000.....	0.2	0.3	0.2	*	*	*	0.2	0.3	0.2	0.2	0.3	0.2	0.2	*	0.2
1999.....	0.2	0.3	0.2	*	*	*	0.2	0.3	0.2	0.2	0.3	0.2	0.2	*	0.2
Malignant neoplasm of eye and adnexa (C69)															
2004.....	0.1	0.1	0.0	*	*	*	0.1	0.1	0.1	0.1	0.1	0.1	*	*	*
2003.....	0.1	0.1	0.1	*	*	*	0.1	0.1	0.1	0.1	0.1	0.1	*	*	*
2002.....	0.1	0.1	0.1	*	*	*	0.1	0.1	0.1	0.1	0.1	0.1	*	*	*
2001.....	0.1	0.1	0.0	*	*	*	0.1	0.1	0.1	0.1	0.1	0.1	*	*	*
2000.....	0.1	0.1	0.0	*	*	*	0.1	0.1	0.1	0.1	0.1	0.1	*	*	*
1999.....	0.1	0.1	0.1	*	*	*	0.1	0.1	0.1	0.1	0.1	0.1	*	*	*
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)															
2004.....	4.3	5.2	3.5	2.7	3.1	2.3	4.4	5.4	3.6	4.8	5.8	3.9	2.5	3.1	2.0
2003.....	4.4	5.3	3.6	2.8	3.3	2.4	4.5	5.5	3.7	4.9	5.9	4.1	2.7	3.4	2.2
2002.....	4.4	5.5	3.6	2.9	3.4	2.4	4.6	5.6	3.7	4.9	6.1	4.0	2.6	3.2	2.2
2001.....	4.5	5.5	3.6	2.9	3.4	2.4	4.6	5.7	3.7	4.9	6.1	4.0	2.6	3.2	2.2
2000.....	4.5	5.6	3.7	3.0	3.7	2.6	4.6	5.7	3.8	5.0	6.2	4.0	2.7	3.3	2.3
1999.....	4.6	5.6	3.8	2.9	3.2	2.6	4.8	5.8	3.9	5.1	6.3	4.2	2.8	3.3	2.3
Of brain (C71)															
2004.....	4.2	5.1	3.4	2.6	3.0	2.3	4.4	5.3	3.6	4.7	5.7	3.9	2.4	3.0	2.0
2003.....	4.3	5.3	3.5	2.7	3.2	2.3	4.5	5.5	3.6	4.8	5.9	4.0	2.6	3.4	2.1
2002.....	4.3	5.3	3.5	2.8	3.4	2.3	4.5	5.5	3.6	4.9	5.9	4.0	2.5	3.0	2.2
2001.....	4.4	5.4	3.5	2.8	3.3	2.4	4.5	5.5	3.6	4.9	6.0	3.9	2.6	3.1	2.2
2000.....	4.5	5.5	3.6	3.0	3.6	2.5	4.6	5.7	3.7	4.9	6.0	4.0	2.7	3.2	2.2
1999.....	4.5	5.5	3.7	2.8	3.1	2.6	4.7	5.7	3.8	5.0	6.1	4.1	2.6	3.2	2.2
Of other parts of central nervous system (C70,C72)															
2004.....	0.1	0.1	0.1	*	*	*	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2003.....	0.1	0.1	0.1	*	*	*	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2002.....	0.1	0.1	0.1	*	*	*	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2001.....	0.1	0.1	0.1	*	*	*	0.1	0.1	0.1	0.1	0.1	0.1	*	*	*
2000.....	0.1	0.1	0.1	*	*	*	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
1999.....	0.1	0.1	0.1	*	*	*	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
Malignant neoplasms of thyroid and other endocrine glands (C73-C75)															
2004.....	0.7	0.7	0.7	0.7	0.6	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.6	0.7
2003.....	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.6	0.5	0.6
2002.....	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.6	0.8
2001.....	0.8	0.8	0.7	0.7	0.6	0.8	0.8	0.8	0.7	0.8	0.8	0.7	0.7	0.7	0.8
2000.....	0.8	0.8	0.7	0.8	0.7	0.8	0.8	0.8	0.7	0.7	0.8	0.7	0.8	0.6	0.9
1999.....	0.7	0.8	0.7	0.9	0.8	0.9	0.7	0.8	0.7	0.7	0.8	0.7	0.8	0.7	0.7
Other malignant neoplasms of other and unspecified sites (C76-C80,C97)															
2004.....	13.3	16.1	11.2	8.7	10.4	7.5	13.6	16.5	11.4	13.5	16.2	11.4	17.1	22.5	13.4
2003.....	13.7	16.6	11.6	9.6	11.6	8.0	14.0	16.9	11.8	13.8	16.7	11.8	17.5	22.3	14.2
2002.....	14.1	17.2	11.9	9.6	11.7	8.1	14.4	17.6	12.1	14.1	17.2	11.9	18.5	23.4	15.3
2001.....	14.2	17.2	12.0	9.7	11.7	8.2	14.4	17.5	12.2	14.1	17.0	12.1	18.6	24.6	14.7
2000.....	14.7	17.8	12.5	10.5	12.8	8.9	14.9	18.1	12.7	14.7	17.6	12.6	19.0	25.1	14.9
1999.....	16.0	19.5	13.6	10.5	13.3	8.6	16.3	19.8	13.8	16.0	19.4	13.6	20.9	26.8	17.0

Worktable Orig13. Age-adjusted death rates for 358 selected causes, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), and year	All origins 1/			Hispanic			Non-Hispanic 2/			Non-Hispanic white			Non-Hispanic black		
	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)															
2004.....	18.4	23.6	14.6	13.2	16.2	10.7	18.7	24.0	14.8	18.9	24.4	14.9	19.2	23.9	16.0
2003.....	19.1	24.5	15.1	13.8	16.3	11.9	19.3	24.9	15.2	19.6	25.5	15.3	19.2	23.4	16.4
2002.....	19.5	25.1	15.5	14.5	17.4	12.4	19.8	25.5	15.7	20.1	26.0	15.8	19.6	24.8	16.2
2001.....	20.0	25.5	16.0	14.9	19.3	11.6	20.2	25.8	16.2	20.4	26.1	16.3	20.3	25.7	16.8
2000.....	20.4	26.0	16.5	15.5	19.3	12.7	20.6	26.3	16.6	20.9	26.7	16.8	20.2	25.4	16.9
1999.....	20.6	26.1	16.7	15.4	17.6	13.5	20.8	26.5	16.8	21.0	26.8	16.9	20.6	26.0	17.0
Hodgkin's disease (C81)															
2004.....	0.4	0.5	0.3	0.4	0.5	0.3	0.4	0.6	0.4	0.4	0.6	0.4	0.4	0.5	0.4
2003.....	0.5	0.5	0.4	0.4	0.5	0.3	0.5	0.6	0.4	0.5	0.6	0.4	0.4	0.5	0.3
2002.....	0.5	0.6	0.4	0.4	0.5	0.4	0.5	0.6	0.4	0.5	0.6	0.4	0.4	0.6	0.3
2001.....	0.5	0.6	0.4	0.5	0.7	0.3	0.5	0.6	0.4	0.5	0.6	0.4	0.4	0.5	0.4
2000.....	0.5	0.6	0.4	0.5	0.7	0.3	0.5	0.5	0.4	0.5	0.6	0.4	0.3	0.4	0.3
1999.....	0.5	0.6	0.4	0.6	0.8	0.5	0.5	0.6	0.4	0.5	0.7	0.4	0.4	0.6	0.3
Of other lymphoid, hematopoietic and related tissue (C82-C85,C96)															
2004.....	7.1	8.9	5.7	5.2	6.4	4.2	7.2	9.0	5.8	7.5	9.4	6.1	5.1	6.3	4.2
2003.....	7.3	9.3	5.9	5.4	6.4	4.7	7.4	9.5	6.0	7.8	9.9	6.2	5.0	5.9	4.3
2002.....	7.6	9.6	6.2	5.8	6.6	5.2	7.7	9.7	6.2	8.1	10.1	6.5	5.5	6.9	4.4
2001.....	7.9	10.0	6.4	5.9	7.6	4.6	8.0	10.1	6.5	8.3	10.4	6.8	5.7	7.4	4.5
2000.....	8.3	10.3	6.8	6.4	7.8	5.3	8.3	10.4	6.8	8.7	10.8	7.2	5.3	6.8	4.3
1999.....	8.4	10.4	6.9	6.2	7.1	5.5	8.5	10.5	7.0	8.8	10.9	7.3	5.7	7.6	4.5
Non-Hodgkin's lymphoma (C82-C85)															
2004.....	7.0	8.8	5.7	5.2	6.4	4.2	7.1	9.0	5.8	7.5	9.3	6.1	5.0	6.2	4.1
2003.....	7.3	9.3	5.9	5.4	6.4	4.7	7.4	9.4	6.0	7.8	9.9	6.2	5.0	5.9	4.3
2002.....	7.6	9.6	6.2	5.8	6.6	5.2	7.7	9.7	6.2	8.0	10.1	6.5	5.4	6.9	4.4
2001.....	7.9	9.9	6.4	5.9	7.5	4.6	8.0	10.0	6.5	8.3	10.4	6.8	5.7	7.4	4.5
2000.....	8.2	10.2	6.7	6.3	7.7	5.3	8.3	10.3	6.8	8.7	10.8	7.1	5.3	6.7	4.3
1999.....	8.3	10.3	6.9	6.2	7.1	5.5	8.4	10.5	6.9	8.8	10.8	7.3	5.7	7.6	4.4
Of other and unspecified lymphoid, hematopoietic and related tissue (C96)															
2004.....	0.0	0.0	0.0	*	*	*	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2003.....	0.0	0.0	0.0	*	*	*	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2002.....	0.0	0.0	0.0	*	*	*	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2001.....	0.0	0.0	0.0	*	*	*	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2000.....	0.0	0.0	0.0	*	*	*	0.0	0.1	0.0	0.0	0.0	0.0	*	*	*
1999.....	0.0	0.0	0.0	*	*	*	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
Malignant immunoproliferative diseases (C88)															
2004.....	0.1	0.1	0.1	*	*	*	0.1	0.2	0.1	0.1	0.2	0.1	*	*	*
2003.....	0.1	0.2	0.1	*	*	*	0.1	0.2	0.1	0.1	0.2	0.1	*	*	*
2002.....	0.1	0.2	0.1	*	*	*	0.1	0.2	0.1	0.2	0.2	0.1	*	*	*
2001.....	0.1	0.2	0.1	*	*	*	0.1	0.2	0.1	0.1	0.2	0.1	*	*	*
2000.....	0.1	0.2	0.1	*	*	*	0.1	0.2	0.1	0.1	0.2	0.1	*	*	*
1999.....	0.2	0.2	0.1	*	*	*	0.2	0.2	0.1	0.2	0.2	0.1	0.1	*	*
Multiple myeloma and malignant plasma cell neoplasms (C90)															
2004.....	3.6	4.4	3.0	2.9	3.3	2.5	3.6	4.5	3.0	3.4	4.2	2.7	7.0	8.4	6.1
2003.....	3.7	4.6	3.1	2.9	3.2	2.7	3.8	4.7	3.1	3.5	4.4	2.8	7.2	8.5	6.4
2002.....	3.8	4.8	3.2	3.3	4.2	2.7	3.8	4.8	3.2	3.6	4.5	2.9	6.9	8.4	6.1
2001.....	3.8	4.7	3.2	3.2	4.3	2.5	3.8	4.7	3.2	3.6	4.4	2.9	7.3	8.8	6.4
2000.....	3.9	4.7	3.3	3.2	3.7	2.8	3.9	4.7	3.3	3.6	4.4	3.0	7.7	8.9	6.9
1999.....	3.9	4.7	3.3	3.2	3.4	3.0	3.9	4.8	3.3	3.5	4.5	2.9	7.5	8.9	6.7

Worktable Orig13. Age-adjusted death rates for 358 selected causes, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), and year	All origins 1/			Hispanic			Non-Hispanic 2/			Non-Hispanic white			Non-Hispanic black		
	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female
Carcinoma in situ of other and unspecified sites (D00-D04,D09)															
2004.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	0.0	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Benign neoplasms (D10-D36)															
2004.....	0.5	0.5	0.6	0.4	0.5	0.4	0.5	0.4	0.6	0.5	0.4	0.5	0.9	0.9	0.9
2003.....	0.5	0.5	0.5	0.5	0.3	0.6	0.5	0.5	0.5	0.5	0.5	0.5	0.8	0.9	0.7
2002.....	0.5	0.5	0.6	0.5	0.5	0.4	0.6	0.5	0.6	0.5	0.5	0.6	0.9	0.9	0.9
2001.....	0.6	0.6	0.6	0.4	0.3	0.5	0.5	0.6	0.6	0.5	0.5	0.5	1.0	1.1	0.9
2000.....	0.6	0.5	0.6	0.5	0.4	0.6	0.6	0.5	0.6	0.5	0.5	0.6	0.9	0.8	1.0
1999.....	0.6	0.6	0.6	0.4	0.3	0.4	0.6	0.6	0.6	0.5	0.5	0.5	1.0	1.0	1.0
Of female genital organs (D25-D28)															
2004.....	0.0	...	0.0	*	...	*	0.0	...	0.0	*	...	*	*	...	*
2003.....	0.0	...	0.0	*	...	*	0.0	...	0.0	*	...	*	*	...	*
2002.....	0.0	...	0.0	*	...	*	0.0	...	0.0	0.0	...	0.0	0.1	...	0.1
2001.....	0.0	...	0.0	*	...	*	0.0	...	0.0	0.0	...	0.0	*	...	*
2000.....	0.0	...	0.0	*	...	*	0.0	...	0.0	0.0	...	0.0	0.1	...	0.2
1999.....	0.0	...	0.0	*	...	*	0.0	...	0.0	*	...	*	*	...	*
Of eye, brain and other parts of central nervous system (D31-D33)															
2004.....	0.3	0.3	0.3	0.2	0.2	0.2	0.3	0.3	0.3	0.3	0.3	0.3	0.5	0.5	0.6
2003.....	0.3	0.3	0.3	0.3	*	0.4	0.3	0.3	0.3	0.3	0.3	0.3	0.5	0.4	0.5
2002.....	0.3	0.3	0.4	0.2	0.2	0.2	0.3	0.3	0.4	0.3	0.3	0.4	0.5	0.4	0.5
2001.....	0.3	0.3	0.3	0.2	*	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.6	0.6	0.6
2000.....	0.3	0.3	0.4	0.3	*	0.3	0.4	0.3	0.4	0.3	0.3	0.4	0.5	0.4	0.5
1999.....	0.3	0.3	0.3	0.2	*	0.2	0.4	0.3	0.4	0.3	0.3	0.3	0.6	0.5	0.6
Of other and unspecified sites (D10-D24,D29-D30,D34-D36)															
2004.....	0.2	0.2	0.2	0.2	0.2	*	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.3	0.3
2003.....	0.2	0.2	0.2	0.2	*	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.4	0.2
2002.....	0.2	0.2	0.2	0.2	0.3	*	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.4	0.3
2001.....	0.2	0.3	0.2	0.2	*	0.2	0.2	0.3	0.2	0.2	0.2	0.2	0.3	0.5	0.2
2000.....	0.2	0.2	0.2	0.2	*	0.3	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.4	0.3
1999.....	0.2	0.2	0.2	0.2	0.2	*	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.5	0.3
Neoplasms of uncertain or unknown behavior of specified sites (D37-D47)															
2004.....	3.8	4.9	3.0	2.5	2.9	2.2	3.9	5.0	3.1	4.0	5.2	3.1	3.3	4.1	2.8
2003.....	3.9	5.0	3.2	2.7	3.3	2.2	4.0	5.1	3.2	4.1	5.3	3.3	3.3	4.1	2.9
2002.....	3.8	5.0	3.1	2.6	3.1	2.2	3.9	5.1	3.1	4.0	5.2	3.2	3.4	4.2	2.9
2001.....	4.1	5.3	3.3	2.7	3.4	2.2	4.1	5.4	3.4	4.3	5.5	3.4	3.5	4.4	3.0
2000.....	4.1	5.3	3.3	2.8	3.3	2.5	4.2	5.4	3.4	4.3	5.5	3.5	3.5	4.4	2.9
1999.....	4.1	5.3	3.3	2.7	3.0	2.6	4.1	5.4	3.3	4.2	5.6	3.4	3.6	4.2	3.2
Neoplasms of uncertain or unknown behavior of unspecified sites (D48)															
2004.....	0.3	0.2	0.2	0.2	0.2	*	0.3	0.3	0.2	0.3	0.2	0.2	0.3	0.3	0.3
2003.....	0.2	0.2	0.2	0.1	*	*	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
2002.....	0.2	0.2	0.3	0.1	*	*	0.2	0.2	0.3	0.2	0.2	0.2	0.2	0.2	0.2
2001.....	0.2	0.2	0.2	0.2	0.2	*	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
2000.....	0.2	0.2	0.2	0.2	*	*	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.2
1999.....	0.2	0.2	0.2	0.1	*	*	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.3	0.3

Worktable Orig13. Age-adjusted death rates for 358 selected causes, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), and year	All origins 1/			Hispanic			Non-Hispanic 2/			Non-Hispanic white			Non-Hispanic black		
	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female
III. Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)															
2004.....	3.1	3.2	3.1	2.1	2.1	2.0	3.2	3.3	3.1	2.7	2.9	2.6	7.1	7.3	7.0
2003.....	3.3	3.3	3.2	2.3	2.4	2.2	3.3	3.4	3.3	2.8	2.9	2.7	7.6	7.9	7.4
2002.....	3.4	3.4	3.3	2.7	3.2	2.3	3.4	3.4	3.4	2.8	2.9	2.8	7.7	7.9	7.6
2001.....	3.4	3.4	3.3	2.6	2.8	2.5	3.4	3.4	3.4	2.9	2.9	2.8	7.6	7.4	7.6
2000.....	3.4	3.4	3.3	2.3	2.3	2.3	3.4	3.5	3.4	2.9	3.0	2.8	7.5	7.3	7.7
1999.....	3.3	3.4	3.2	2.6	2.9	2.3	3.4	3.5	3.3	2.8	3.0	2.7	7.4	7.5	7.3
Anemias (D50-D64)															
2004.....	1.5	1.5	1.5	1.0	0.9	1.0	1.6	1.6	1.5	1.3	1.3	1.3	3.4	3.6	3.2
2003.....	1.6	1.6	1.5	1.1	1.2	1.1	1.6	1.6	1.5	1.3	1.3	1.3	3.6	4.0	3.2
2002.....	1.6	1.6	1.6	1.3	1.6	1.1	1.6	1.6	1.6	1.3	1.3	1.4	3.7	4.0	3.5
2001.....	1.6	1.6	1.6	1.2	1.1	1.2	1.7	1.6	1.7	1.4	1.4	1.4	3.6	3.7	3.4
2000.....	1.6	1.6	1.6	0.9	0.9	1.0	1.7	1.6	1.7	1.4	1.4	1.4	3.5	3.5	3.5
1999.....	1.6	1.7	1.6	1.2	1.4	1.1	1.7	1.7	1.6	1.4	1.4	1.4	3.7	3.8	3.5
Nutritional anemias (D50-D53)															
2004.....	0.1	0.1	0.1	*	*	*	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2003.....	0.1	0.1	0.1	*	*	*	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	0.1
2002.....	0.1	0.1	0.1	*	*	*	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2001.....	0.1	0.1	0.1	*	*	*	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2000.....	0.1	0.1	0.1	*	*	*	0.1	0.1	0.1	0.1	0.1	0.1	0.2	*	0.2
1999.....	0.1	0.1	0.1	*	*	*	0.1	0.1	0.1	0.1	0.1	0.1	0.2	*	0.2
Hemolytic anemias (D55-D59)															
2004.....	0.3	0.3	0.3	0.1	*	*	0.3	0.4	0.3	0.1	0.1	0.1	1.5	1.7	1.4
2003.....	0.3	0.3	0.3	0.1	*	*	0.3	0.3	0.3	0.1	0.1	0.1	1.4	1.5	1.3
2002.....	0.3	0.3	0.3	0.1	*	0.1	0.3	0.3	0.3	0.1	0.1	0.1	1.5	1.5	1.5
2001.....	0.3	0.3	0.3	0.1	*	*	0.3	0.3	0.3	0.1	0.1	0.1	1.5	1.6	1.4
2000.....	0.3	0.3	0.3	0.1	*	0.2	0.3	0.3	0.3	0.1	0.1	0.1	1.5	1.3	1.5
1999.....	0.3	0.3	0.3	0.1	*	*	0.3	0.4	0.3	0.1	0.1	0.1	1.6	1.8	1.4
Aplastic anemias (D60-D61)															
2004.....	0.3	0.4	0.3	0.3	0.2	0.3	0.3	0.4	0.3	0.3	0.3	0.3	0.4	0.4	0.3
2003.....	0.3	0.4	0.3	0.2	0.3	0.2	0.3	0.4	0.3	0.3	0.4	0.3	0.4	0.5	0.4
2002.....	0.3	0.4	0.3	0.3	0.5	0.3	0.3	0.4	0.3	0.3	0.4	0.3	0.4	0.5	0.4
2001.....	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.3
2000.....	0.4	0.4	0.4	0.3	0.2	0.2	0.4	0.4	0.4	0.4	0.4	0.4	0.5	0.6	0.4
1999.....	0.4	0.5	0.4	0.3	0.3	0.3	0.4	0.5	0.4	0.4	0.4	0.4	0.5	0.5	0.4
Acute posthemorrhagic and other anemias (D62,D64)															
2004.....	0.8	0.8	0.8	0.6	0.6	0.6	0.8	0.8	0.8	0.8	0.7	0.8	1.4	1.5	1.3
2003.....	0.8	0.8	0.8	0.7	0.7	0.8	0.8	0.8	0.8	0.8	0.7	0.8	1.6	1.8	1.4
2002.....	0.8	0.8	0.9	0.7	0.9	0.6	0.9	0.8	0.9	0.8	0.7	0.8	1.7	1.9	1.5
2001.....	0.9	0.8	0.9	0.6	0.6	0.6	0.9	0.8	0.9	0.8	0.7	0.8	1.6	1.5	1.6
2000.....	0.8	0.8	0.8	0.5	0.5	0.5	0.8	0.8	0.8	0.8	0.8	0.8	1.4	1.4	1.4
1999.....	0.8	0.8	0.9	0.8	0.9	0.7	0.8	0.8	0.9	0.8	0.7	0.8	1.5	1.3	1.5
Coagulation defects, purpura and other hemorrhagic conditions (D65-D69)															
2004.....	0.9	0.9	0.8	0.7	0.7	0.6	0.9	1.0	0.9	0.8	0.9	0.8	1.4	1.5	1.4
2003.....	0.9	1.0	0.9	0.7	0.7	0.7	0.9	1.0	0.9	0.9	0.9	0.8	1.5	1.5	1.4
2002.....	0.9	1.0	0.9	0.9	1.0	0.8	0.9	1.0	0.9	0.9	0.9	0.8	1.4	1.4	1.4
2001.....	0.9	1.0	0.9	0.9	1.0	0.9	0.9	1.0	0.9	0.9	0.9	0.8	1.5	1.5	1.5
2000.....	1.0	1.0	0.9	0.8	0.8	0.9	0.9	1.0	0.9	0.9	1.0	0.9	1.5	1.5	1.5
1999.....	1.0	1.0	0.9	0.9	0.9	0.9	1.0	1.0	0.9	0.9	0.9	0.8	1.6	1.7	1.6

Worktable Orig13. Age-adjusted death rates for 358 selected causes, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), and year	All origins 1/			Hispanic			Non-Hispanic 2/			Non-Hispanic white			Non-Hispanic black		
	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female
Other diseases of blood and blood-forming organs (D70-D76)															
2004.....	0.3	0.3	0.3	0.1	0.1	0.1	0.3	0.3	0.3	0.3	0.3	0.3	0.2	0.3	0.2
2003.....	0.3	0.3	0.3	0.2	0.2	*	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3
2002.....	0.3	0.4	0.3	0.2	0.3	0.2	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.5	0.3
2001.....	0.3	0.3	0.3	0.3	0.3	0.2	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3
2000.....	0.3	0.4	0.3	0.2	0.3	0.2	0.3	0.4	0.3	0.3	0.3	0.3	0.3	0.4	0.3
1999.....	0.3	0.3	0.3	0.2	0.3	*	0.3	0.3	0.3	0.3	0.3	0.2	0.3	0.4	0.3
Certain disorders involving the immune mechanism (D80-D89)															
2004.....	0.5	0.5	0.5	0.3	0.4	0.2	0.5	0.4	0.5	0.3	0.3	0.2	2.1	1.9	2.2
2003.....	0.5	0.5	0.5	0.3	0.4	0.2	0.5	0.5	0.6	0.3	0.3	0.3	2.3	2.1	2.5
2002.....	0.5	0.5	0.5	0.2	0.3	0.2	0.5	0.5	0.5	0.3	0.3	0.3	2.2	2.0	2.4
2001.....	0.5	0.5	0.5	0.3	0.3	0.2	0.5	0.5	0.5	0.3	0.3	0.3	2.2	2.0	2.4
2000.....	0.5	0.5	0.5	0.3	0.4	0.3	0.5	0.5	0.5	0.3	0.3	0.3	2.1	1.9	2.3
1999.....	0.4	0.4	0.4	0.3	0.3	0.2	0.4	0.4	0.4	0.3	0.3	0.2	1.8	1.6	1.9
IV. Endocrine, nutritional and metabolic diseases (E00-E88)															
2004.....	33.1	37.2	29.8	37.3	39.4	35.2	32.9	37.0	29.5	30.2	34.7	26.5	60.8	64.6	57.7
2003.....	34.2	38.0	31.1	40.6	43.8	37.9	33.8	37.6	30.6	31.0	35.3	27.5	62.3	64.2	60.2
2002.....	34.4	37.7	31.7	41.4	44.0	39.2	34.1	37.3	31.3	31.2	35.0	28.0	63.3	63.4	62.2
2001.....	34.3	37.1	31.8	42.5	43.9	41.0	33.8	36.7	31.3	31.0	34.4	28.2	62.5	63.8	60.9
2000.....	34.1	36.9	31.8	42.6	45.1	40.4	33.6	36.4	31.3	30.8	34.2	28.1	63.0	62.2	62.7
1999.....	34.0	36.6	31.8	43.8	45.1	42.4	33.5	36.2	31.3	30.5	33.7	27.9	63.0	63.8	62.2
Endocrine diseases (E00-E34)															
2004.....	25.4	28.9	22.7	32.7	34.7	30.8	24.9	28.5	22.2	22.3	26.2	19.4	50.2	53.4	47.4
2003.....	26.3	29.6	23.7	35.7	38.8	33.3	25.7	29.0	23.1	23.0	26.7	20.1	51.4	52.9	49.7
2002.....	26.4	29.3	24.1	36.3	38.6	34.4	25.8	28.7	23.5	23.2	26.6	20.5	51.6	51.2	50.8
2001.....	26.3	28.9	24.2	37.5	38.3	36.3	25.6	28.3	23.6	23.1	26.1	20.7	51.2	51.8	50.1
2000.....	26.0	28.4	24.1	37.5	40.0	35.4	25.4	27.8	23.5	22.8	25.7	20.5	51.7	50.6	51.8
1999.....	26.0	28.5	24.1	38.6	39.7	37.5	25.4	27.9	23.4	22.6	25.6	20.3	51.9	51.5	51.7
Disorders of thyroid gland (E00-E07)															
2004.....	0.6	0.4	0.7	0.4	0.3	0.5	0.6	0.4	0.7	0.6	0.4	0.7	0.8	0.6	0.9
2003.....	0.7	0.4	0.8	0.5	0.3	0.7	0.7	0.4	0.8	0.7	0.3	0.9	0.7	0.5	0.9
2002.....	0.7	0.4	0.8	0.5	0.4	0.6	0.7	0.4	0.8	0.6	0.4	0.9	0.7	0.4	0.9
2001.....	0.7	0.4	0.8	0.5	0.4	0.5	0.7	0.4	0.8	0.7	0.4	0.8	0.8	0.6	1.0
2000.....	0.6	0.4	0.8	0.4	*	0.5	0.7	0.4	0.8	0.6	0.3	0.8	0.8	0.6	1.0
1999.....	0.7	0.4	0.8	0.5	*	0.6	0.7	0.4	0.8	0.7	0.4	0.8	0.9	0.6	1.0
Diabetes mellitus (E10-E14)															
2004.....	24.5	28.2	21.7	32.1	34.3	30.2	24.0	27.7	21.1	21.5	25.5	18.4	48.7	52.1	46.0
2003.....	25.3	28.9	22.5	35.0	38.3	32.4	24.7	28.3	21.9	22.1	26.1	19.0	50.0	51.7	48.3
2002.....	25.4	28.6	23.0	35.6	38.1	33.6	24.8	28.0	22.3	22.2	25.9	19.3	50.3	50.3	49.4
2001.....	25.3	28.1	23.1	36.7	37.8	35.5	24.6	27.5	22.4	22.1	25.4	19.5	49.8	50.5	48.7
2000.....	25.0	27.8	23.0	36.9	39.6	34.6	24.4	27.1	22.4	21.8	25.1	19.4	50.2	49.4	50.2
1999.....	25.0	27.8	23.0	37.8	39.1	36.6	24.3	27.1	22.3	21.6	25.0	19.2	50.4	50.0	50.1
Disorders of pituitary gland (E22-E23)															
2004.....	0.1	0.1	0.1	*	*	*	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2003.....	0.1	0.1	0.1	*	*	*	0.1	0.1	0.1	0.1	0.1	0.0	0.1	*	*
2002.....	0.1	0.1	0.1	*	*	*	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2001.....	0.1	0.1	0.1	*	*	*	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2000.....	0.1	0.1	0.1	*	*	*	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
1999.....	0.1	0.1	0.1	*	*	*	0.1	0.1	0.1	0.1	0.1	0.1	0.2	0.2	0.1

Worktable Orig13. Age-adjusted death rates for 358 selected causes, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), and year	All origins 1/			Hispanic			Non-Hispanic 2/			Non-Hispanic white			Non-Hispanic black		
	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female
Disorders of adrenal glands (E24-E27)															
2004.....	0.1	0.1	0.1	*	*	*	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	0.1
2003.....	0.1	0.1	0.1	*	*	*	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2002.....	0.1	0.1	0.1	*	*	*	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2001.....	0.1	0.1	0.1	*	*	*	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2000.....	0.1	0.1	0.1	*	*	*	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	0.2
1999.....	0.1	0.1	0.1	*	*	*	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
Diseases of thymus (E32)															
2004.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Other endocrine diseases (E15-E21,E28-E31,E34)															
2004.....	0.1	0.2	0.1	*	*	*	0.2	0.2	0.1	0.1	0.1	0.1	0.4	0.5	0.3
2003.....	0.2	0.1	0.1	0.1	*	*	0.2	0.2	0.2	0.1	0.1	0.1	0.4	0.5	0.3
2002.....	0.1	0.1	0.1	*	*	*	0.1	0.1	0.2	0.1	0.1	0.1	0.3	0.3	0.4
2001.....	0.2	0.2	0.1	0.1	*	*	0.2	0.2	0.2	0.1	0.1	0.1	0.3	0.4	0.3
2000.....	0.1	0.1	0.2	*	*	*	0.1	0.1	0.2	0.1	0.1	0.1	0.4	0.3	0.4
1999.....	0.2	0.2	0.1	0.2	*	*	0.2	0.2	0.1	0.1	0.1	0.1	0.4	0.5	0.3
Nutritional diseases (E40-E68)															
2004.....	2.4	2.3	2.4	1.4	1.3	1.5	2.5	2.4	2.5	2.3	2.3	2.3	4.3	3.9	4.6
2003.....	2.5	2.4	2.5	1.7	1.5	1.7	2.6	2.4	2.6	2.5	2.4	2.4	4.1	3.8	4.3
2002.....	2.6	2.5	2.6	1.7	1.6	1.7	2.6	2.5	2.7	2.5	2.4	2.4	4.5	4.0	4.8
2001.....	2.4	2.2	2.5	1.8	1.8	1.8	2.4	2.2	2.6	2.3	2.1	2.4	4.2	4.0	4.4
2000.....	2.6	2.5	2.6	1.9	1.7	1.9	2.7	2.5	2.7	2.5	2.4	2.5	4.4	4.3	4.5
1999.....	2.5	2.5	2.5	1.8	1.9	1.7	2.6	2.5	2.6	2.4	2.3	2.4	4.4	4.5	4.4
Malnutrition (E40-E46)															
2004.....	0.9	0.9	1.0	0.6	0.5	0.7	1.0	0.9	1.0	0.9	0.8	0.9	1.6	1.8	1.5
2003.....	1.1	1.0	1.1	0.8	0.7	0.8	1.1	1.0	1.1	1.0	0.9	1.1	1.9	2.1	1.7
2002.....	1.2	1.1	1.2	1.0	1.0	0.9	1.2	1.2	1.2	1.1	1.1	1.2	2.0	2.0	2.0
2001.....	1.2	1.1	1.3	1.1	1.1	1.1	1.2	1.1	1.3	1.2	1.0	1.2	1.9	2.2	1.7
2000.....	1.4	1.3	1.5	1.1	1.1	1.2	1.4	1.3	1.5	1.4	1.2	1.4	2.3	2.5	2.1
1999.....	1.5	1.4	1.5	1.2	1.2	1.2	1.5	1.4	1.5	1.4	1.3	1.4	2.5	3.0	2.2
Nutritional marasmus (E41)															
2004.....	0.0	0.0	0.0	*	*	*	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2003.....	0.0	0.0	0.1	*	*	*	0.0	0.0	0.1	0.0	0.0	0.1	0.1	*	*
2002.....	0.1	0.0	0.1	*	*	*	0.1	0.0	0.1	0.1	0.0	0.1	0.1	*	*
2001.....	0.1	0.1	0.1	*	*	*	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2000.....	0.1	0.1	0.1	*	*	*	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
1999.....	0.1	0.1	0.1	*	*	*	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
Other and unspecified malnutrition (E40,E42-E46)															
2004.....	0.9	0.9	0.9	0.6	0.5	0.6	0.9	0.9	0.9	0.9	0.8	0.9	1.6	1.8	1.5
2003.....	1.0	0.9	1.0	0.7	0.7	0.7	1.0	1.0	1.1	1.0	0.9	1.0	1.8	2.0	1.6
2002.....	1.1	1.1	1.1	0.9	0.9	0.9	1.1	1.1	1.1	1.1	1.0	1.1	1.9	1.9	1.9
2001.....	1.1	1.0	1.2	1.0	1.0	1.0	1.2	1.0	1.2	1.1	1.0	1.2	1.8	2.2	1.6
2000.....	1.3	1.2	1.4	1.1	1.0	1.2	1.3	1.2	1.4	1.3	1.2	1.3	2.2	2.4	2.0
1999.....	1.4	1.3	1.4	1.2	1.1	1.2	1.4	1.3	1.4	1.3	1.2	1.4	2.4	2.8	2.1

Worktable Orig13. Age-adjusted death rates for 358 selected causes, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), and year	All origins 1/			Hispanic			Non-Hispanic 2/			Non-Hispanic white			Non-Hispanic black		
	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female
Other nutritional deficiencies (E50-E64)															
2004.....	0.1	0.0	0.0	*	*	*	0.1	0.1	0.0	0.1	0.0	0.1	*	*	*
2003.....	0.1	0.1	0.0	*	*	*	0.1	0.1	0.0	0.1	0.1	0.0	*	*	*
2002.....	0.1	0.1	0.1	*	*	*	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2001.....	0.1	0.1	0.1	*	*	*	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2000.....	0.1	0.1	0.1	*	*	*	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	0.1
1999.....	0.1	0.1	0.1	*	*	*	0.1	0.1	0.1	0.1	0.1	0.1	0.2	*	0.2
Obesity and other hyperalimentation (E65-E68)															
2004.....	1.4	1.4	1.4	0.8	0.7	0.8	1.4	1.4	1.5	1.4	1.4	1.3	2.7	2.0	3.1
2003.....	1.3	1.3	1.4	0.8	0.7	0.9	1.4	1.4	1.4	1.4	1.4	1.3	2.2	1.7	2.6
2002.....	1.3	1.2	1.3	0.7	0.5	0.8	1.3	1.3	1.3	1.2	1.3	1.2	2.4	1.8	2.7
2001.....	1.1	1.0	1.2	0.7	0.6	0.7	1.1	1.0	1.2	1.1	1.0	1.1	2.2	1.6	2.6
2000.....	1.1	1.1	1.1	0.6	0.5	0.7	1.1	1.1	1.1	1.0	1.1	1.0	2.0	1.6	2.3
1999.....	0.9	0.9	1.0	0.5	0.6	0.5	1.0	0.9	1.0	0.9	0.9	0.9	1.7	1.4	2.0
Metabolic disorders (E70-E88)															
2004.....	5.3	6.0	4.7	3.2	3.4	2.9	5.4	6.1	4.8	5.5	6.2	4.9	6.3	7.2	5.6
2003.....	5.4	6.0	4.8	3.2	3.4	2.9	5.6	6.2	4.9	5.6	6.2	4.9	6.8	7.5	6.2
2002.....	5.4	5.9	5.0	3.4	3.8	3.1	5.6	6.1	5.1	5.5	6.0	5.0	7.2	8.1	6.6
2001.....	5.5	6.0	5.0	3.3	3.9	2.9	5.7	6.2	5.2	5.7	6.1	5.2	7.1	8.0	6.4
2000.....	5.5	5.9	5.0	3.3	3.4	3.1	5.6	6.1	5.1	5.6	6.1	5.0	6.9	7.4	6.3
1999.....	5.5	5.7	5.1	3.4	3.5	3.2	5.6	5.8	5.2	5.6	5.7	5.2	6.7	7.8	6.1
Cystic fibrosis (E84)															
2004.....	0.2	0.2	0.2	*	*	*	0.2	0.2	0.2	0.2	0.2	0.2	*	*	*
2003.....	0.2	0.2	0.1	*	*	*	0.2	0.2	0.2	0.2	0.2	0.2	0.0	*	*
2002.....	0.2	0.2	0.2	0.0	*	*	0.2	0.2	0.2	0.2	0.2	0.2	0.1	*	*
2001.....	0.2	0.2	0.2	0.1	*	*	0.2	0.2	0.2	0.2	0.2	0.2	*	*	*
2000.....	0.2	0.2	0.2	*	*	*	0.2	0.2	0.2	0.2	0.2	0.3	*	*	*
1999.....	0.2	0.2	0.2	*	*	*	0.2	0.2	0.2	0.2	0.2	0.3	*	*	*
Volume depletion, disorders of fluid, electrolyte and acid-base balance (E86-E87)															
2004.....	2.2	2.0	2.2	1.3	1.3	1.2	2.2	2.1	2.3	2.1	2.0	2.2	3.2	3.6	2.9
2003.....	2.4	2.2	2.4	1.2	1.4	1.1	2.4	2.2	2.5	2.3	2.1	2.4	3.8	4.0	3.6
2002.....	2.5	2.3	2.6	1.3	1.3	1.3	2.6	2.3	2.7	2.5	2.2	2.6	4.1	4.3	3.9
2001.....	2.7	2.5	2.8	1.4	1.4	1.3	2.8	2.5	2.9	2.7	2.4	2.8	4.4	4.6	4.2
2000.....	2.8	2.7	2.8	1.5	1.4	1.5	2.9	2.8	2.9	2.8	2.6	2.8	4.3	4.5	4.0
1999.....	3.0	2.8	3.1	1.7	1.6	1.7	3.1	2.8	3.1	3.0	2.6	3.1	4.3	5.2	3.8
Other metabolic disorders (E70-E83,E85,E88)															
2004.....	3.0	3.8	2.3	1.8	2.1	1.6	3.1	3.9	2.4	3.1	4.0	2.5	3.1	3.6	2.6
2003.....	2.9	3.7	2.3	1.9	2.1	1.8	3.0	3.8	2.3	3.0	3.9	2.3	2.9	3.5	2.5
2002.....	2.8	3.5	2.2	2.1	2.5	1.8	2.8	3.6	2.2	2.8	3.6	2.2	3.1	3.8	2.6
2001.....	2.6	3.4	2.0	1.9	2.4	1.5	2.7	3.4	2.0	2.7	3.5	2.1	2.7	3.4	2.1
2000.....	2.5	3.1	2.0	1.8	2.0	1.6	2.5	3.1	2.0	2.6	3.2	2.0	2.5	2.8	2.3
1999.....	2.3	2.8	1.9	1.6	1.8	1.5	2.3	2.8	1.9	2.4	2.9	1.9	2.4	2.6	2.2
V. Mental and behavioral disorders (F01-F99)															
2004.....	20.3	19.9	19.5	12.0	13.1	10.5	20.7	20.2	20.0	20.9	20.0	20.4	22.0	25.6	18.9
2003.....	20.2	19.6	19.6	12.0	13.3	10.3	20.6	19.8	20.0	20.8	19.7	20.4	21.5	24.3	18.8
2002.....	19.2	18.8	18.4	11.4	14.2	8.9	19.4	18.9	18.8	19.6	18.6	19.2	20.4	23.8	17.2
2001.....	18.0	17.8	17.1	11.7	13.4	9.9	18.2	18.0	17.3	18.2	17.7	17.7	19.1	23.0	15.7
2000.....	16.7	16.8	15.7	10.5	12.7	8.3	16.9	16.8	16.0	16.9	16.4	16.3	18.1	22.3	14.6
1999.....	15.5	16.0	14.3	11.2	13.7	8.7	15.6	16.0	14.5	15.6	15.7	14.7	16.1	20.6	12.5

Worktable Orig13. Age-adjusted death rates for 358 selected causes, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), and year	All origins 1/			Hispanic			Non-Hispanic 2/			Non-Hispanic white			Non-Hispanic black		
	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female
Organic dementia (F01,F03)															
2004.....	15.3	12.9	16.4	8.4	7.3	8.9	15.6	13.2	16.8	15.9	13.3	17.2	15.7	15.4	15.6
2003.....	15.3	12.8	16.4	7.9	7.0	8.3	15.6	13.1	16.8	15.9	13.2	17.2	15.1	14.4	15.2
2002.....	14.3	12.2	15.3	7.2	7.3	7.1	14.6	12.4	15.6	14.9	12.5	16.0	13.7	13.4	13.6
2001.....	13.2	11.4	14.0	7.7	7.1	8.0	13.4	11.5	14.2	13.7	11.7	14.6	12.2	11.8	12.1
2000.....	12.0	10.3	12.6	6.4	6.4	6.3	12.2	10.4	12.8	12.4	10.5	13.2	11.2	11.5	10.9
1999.....	10.6	9.3	11.1	6.5	6.4	6.6	10.8	9.4	11.2	11.1	9.6	11.6	8.7	8.9	8.5
Other and unspecified organic mental disorders (F04-F09)															
2004.....	0.6	0.6	0.6	0.4	0.3	0.5	0.6	0.6	0.6	0.6	0.6	0.6	0.5	0.5	0.5
2003.....	0.6	0.6	0.6	0.5	0.4	0.5	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.7	0.5
2002.....	0.7	0.6	0.7	0.7	0.7	0.7	0.7	0.6	0.7	0.7	0.6	0.7	0.6	0.6	0.7
2001.....	0.7	0.7	0.8	0.6	0.6	0.6	0.7	0.7	0.8	0.8	0.7	0.8	0.8	0.8	0.7
2000.....	0.8	0.7	0.9	0.7	0.5	0.7	0.8	0.7	0.9	0.8	0.7	0.9	0.8	0.8	0.7
1999.....	0.9	0.8	0.9	0.8	0.9	0.8	0.9	0.8	0.9	0.9	0.8	1.0	0.9	1.1	0.8
Mental and behavioral disorders due to psychoactive substance use (F10-F19)															
2004.....	3.6	5.6	1.7	2.9	5.0	0.9	3.6	5.6	1.8	3.5	5.4	1.7	5.0	8.7	2.2
2003.....	3.4	5.4	1.6	3.2	5.6	1.0	3.4	5.3	1.7	3.3	5.0	1.6	5.0	8.4	2.3
2002.....	3.3	5.2	1.5	3.2	5.6	0.9	3.3	5.1	1.6	3.1	4.7	1.5	5.3	9.0	2.3
2001.....	3.2	5.0	1.5	3.0	5.2	0.9	3.2	4.9	1.5	2.9	4.5	1.4	5.5	9.5	2.3
2000.....	3.1	5.0	1.4	3.1	5.5	0.9	3.1	4.9	1.5	2.8	4.4	1.3	5.5	9.4	2.4
1999.....	3.1	5.1	1.4	3.3	6.0	0.8	3.1	4.9	1.4	2.7	4.3	1.3	5.8	9.8	2.7
Mental and behavioral disorders due to use of alcohol (F10)															
2004.....	2.4	3.9	1.0	1.8	3.3	0.4	2.4	4.0	1.0	2.4	3.8	1.1	3.0	5.6	1.0
2003.....	2.4	3.8	1.0	2.1	3.8	0.5	2.4	3.8	1.1	2.3	3.7	1.1	3.0	5.5	1.1
2002.....	2.3	3.9	1.0	2.2	4.1	0.5	2.3	3.8	1.0	2.3	3.6	1.0	3.4	6.2	1.2
2001.....	2.3	3.8	1.0	2.1	3.9	0.5	2.3	3.8	1.0	2.2	3.4	1.0	3.7	6.8	1.3
2000.....	2.2	3.7	0.9	2.3	4.2	0.5	2.2	3.7	0.9	2.1	3.4	0.9	3.6	6.7	1.3
1999.....	2.2	3.7	0.9	2.3	4.2	0.5	2.2	3.6	0.9	2.0	3.3	0.9	3.8	6.8	1.5
Mental and behavioral disorders due to other psychoactive substance use (F11-F19)															
2004.....	1.2	1.6	0.7	1.1	1.7	0.5	1.2	1.7	0.7	1.1	1.5	0.7	2.0	3.1	1.2
2003.....	1.1	1.5	0.6	1.1	1.8	0.5	1.0	1.5	0.7	0.9	1.3	0.6	1.9	2.8	1.2
2002.....	0.9	1.3	0.6	0.9	1.5	0.4	0.9	1.2	0.6	0.8	1.1	0.5	1.9	2.8	1.2
2001.....	0.9	1.2	0.5	0.8	1.3	0.4	0.9	1.2	0.5	0.7	1.0	0.5	1.8	2.7	1.1
2000.....	0.9	1.2	0.6	0.9	1.3	0.4	0.9	1.2	0.5	0.7	1.0	0.5	1.8	2.7	1.1
1999.....	0.9	1.3	0.5	1.1	1.8	0.3	0.8	1.2	0.5	0.7	1.0	0.4	1.9	2.9	1.1
Schizophrenia, schizotypal and delusional disorders (F20-F29)															
2004.....	0.2	0.2	0.2	*	*	*	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.4	0.2
2003.....	0.2	0.1	0.2	*	*	*	0.2	0.1	0.2	0.2	0.2	0.2	0.2	0.2	0.2
2002.....	0.2	0.2	0.2	*	*	*	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.3	0.2
2001.....	0.2	0.1	0.2	*	*	*	0.2	0.2	0.2	0.2	0.1	0.2	0.2	0.2	0.2
2000.....	0.2	0.2	0.2	*	*	*	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
1999.....	0.2	0.2	0.2	*	*	*	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.2
Mood (affective) disorders (F30-F39)															
2004.....	0.3	0.2	0.3	0.1	*	*	0.3	0.2	0.3	0.3	0.2	0.3	0.1	*	0.1
2003.....	0.3	0.3	0.3	*	*	*	0.3	0.3	0.3	0.3	0.3	0.4	0.2	*	0.2
2002.....	0.3	0.2	0.3	*	*	*	0.3	0.2	0.4	0.3	0.2	0.4	0.1	*	0.1
2001.....	0.3	0.3	0.3	*	*	*	0.3	0.3	0.3	0.3	0.3	0.3	0.1	*	0.1
2000.....	0.3	0.3	0.3	*	*	*	0.3	0.3	0.3	0.3	0.3	0.3	0.1	*	0.1
1999.....	0.3	0.3	0.3	*	*	*	0.3	0.3	0.3	0.4	0.3	0.3	0.1	*	*

Worktable Orig13. Age-adjusted death rates for 358 selected causes, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), and year	All origins 1/			Hispanic			Non-Hispanic 2/			Non-Hispanic white			Non-Hispanic black		
	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female
Neurotic, stress-related and somatoform disorders (F40-F48)															
2004.....	0.0	0.0	0.0	*	*	*	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2003.....	0.0	0.0	0.0	*	*	*	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2002.....	0.0	0.0	0.0	*	*	*	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2001.....	0.0	0.0	0.0	*	*	*	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2000.....	0.0	0.0	0.0	*	*	*	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
1999.....	0.0	0.0	0.0	*	*	*	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
Mental retardation (F70-F79)															
2004.....	0.2	0.2	0.1	0.1	*	*	0.2	0.2	0.1	0.2	0.2	0.1	0.2	0.2	0.2
2003.....	0.2	0.2	0.1	0.1	*	*	0.2	0.2	0.1	0.2	0.2	0.1	0.2	0.2	0.1
2002.....	0.2	0.2	0.1	0.1	*	*	0.2	0.2	0.1	0.2	0.2	0.2	0.2	0.2	*
2001.....	0.1	0.1	0.1	*	*	*	0.1	0.2	0.1	0.1	0.2	0.1	0.2	0.2	0.1
2000.....	0.1	0.2	0.1	0.1	*	*	0.1	0.2	0.1	0.2	0.2	0.1	0.1	0.2	0.1
1999.....	0.2	0.2	0.1	0.1	*	*	0.2	0.2	0.1	0.2	0.2	0.1	0.2	0.2	0.2
Other and unspecified mental and behavioral disorders (F50-F69,F80-F99)															
2004.....	0.2	0.1	0.2	*	*	*	0.2	0.1	0.2	0.2	0.1	0.2	0.1	*	*
2003.....	0.2	0.1	0.2	0.1	*	*	0.2	0.1	0.2	0.2	0.1	0.2	0.2	*	0.2
2002.....	0.2	0.1	0.2	*	*	*	0.2	0.1	0.2	0.2	0.1	0.2	0.1	*	0.2
2001.....	0.1	0.1	0.2	*	*	*	0.1	0.1	0.2	0.2	0.1	0.2	0.1	*	*
2000.....	0.1	0.1	0.2	*	*	*	0.1	0.1	0.2	0.2	0.1	0.2	0.1	*	*
1999.....	0.1	0.1	0.2	*	*	*	0.2	0.1	0.2	0.2	0.1	0.2	0.1	*	*
VI. Diseases of the nervous system (G00-G98)															
2004.....	37.1	37.5	36.3	22.7	22.9	22.1	38.0	38.4	37.1	39.1	39.6	38.1	33.3	32.8	33.0
2003.....	37.2	37.8	36.1	23.4	25.0	22.0	37.9	38.5	36.9	39.1	39.7	38.1	31.7	32.8	30.5
2002.....	35.6	36.5	34.4	21.1	21.4	20.5	36.3	37.2	35.1	37.5	38.5	36.3	29.6	30.7	28.4
2001.....	34.2	35.8	32.7	19.9	21.3	18.8	34.9	36.5	33.4	35.9	37.6	34.4	29.1	30.7	27.8
2000.....	33.0	34.8	31.5	19.9	21.9	18.2	33.6	35.4	32.1	34.6	36.4	33.0	27.4	29.2	26.1
1999.....	30.9	33.2	29.1	18.7	19.8	17.7	31.5	33.9	29.6	32.4	34.9	30.4	25.3	27.0	24.0
Meningitis (G00,G03)															
2004.....	0.3	0.3	0.2	0.3	0.3	0.2	0.3	0.3	0.2	0.2	0.2	0.2	0.5	0.5	0.4
2003.....	0.3	0.3	0.2	0.2	0.2	0.2	0.3	0.3	0.2	0.2	0.2	0.2	0.6	0.6	0.6
2002.....	0.2	0.3	0.2	0.3	0.3	0.2	0.2	0.3	0.2	0.2	0.2	0.2	0.6	0.8	0.4
2001.....	0.3	0.3	0.2	0.2	0.3	0.2	0.3	0.3	0.2	0.2	0.2	0.2	0.6	0.8	0.4
2000.....	0.3	0.3	0.2	0.2	0.3	0.1	0.3	0.3	0.2	0.2	0.3	0.2	0.6	0.8	0.6
1999.....	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.2	0.7	0.9	0.6
Other inflammatory diseases of central nervous system (G04-G09)															
2004.....	0.2	0.3	0.2	0.2	0.3	*	0.2	0.3	0.2	0.2	0.3	0.2	0.3	0.4	0.3
2003.....	0.3	0.3	0.2	0.3	0.3	0.2	0.3	0.3	0.2	0.2	0.3	0.2	0.3	0.4	0.3
2002.....	0.2	0.3	0.2	0.2	0.2	0.2	0.3	0.3	0.2	0.2	0.3	0.2	0.4	0.5	0.3
2001.....	0.2	0.2	0.2	0.1	0.2	*	0.2	0.3	0.2	0.2	0.2	0.2	0.4	0.4	0.4
2000.....	0.2	0.3	0.2	0.2	0.4	*	0.2	0.3	0.2	0.2	0.2	0.2	0.4	0.4	0.3
1999.....	0.2	0.3	0.2	0.2	0.2	*	0.2	0.3	0.2	0.2	0.2	0.2	0.3	0.4	0.3
Parkinson's disease (G20-G21)															
2004.....	6.1	9.3	4.1	3.8	5.7	2.6	6.2	9.5	4.2	6.6	10.1	4.4	2.7	4.3	1.9
2003.....	6.2	9.3	4.2	3.6	5.5	2.4	6.3	9.5	4.3	6.7	10.1	4.6	2.8	4.5	1.9
2002.....	5.9	9.0	4.1	3.5	5.1	2.6	6.0	9.2	4.1	6.4	9.7	4.4	2.5	4.0	1.7
2001.....	5.9	9.1	4.0	3.8	5.6	2.7	5.9	9.3	4.1	6.3	9.9	4.3	2.5	4.0	1.8
2000.....	5.7	8.8	3.9	3.6	5.4	2.6	5.8	9.0	4.0	6.1	9.5	4.3	2.5	4.2	1.6
1999.....	5.4	8.4	3.7	3.5	5.1	2.5	5.4	8.5	3.7	5.8	9.0	3.9	2.4	3.8	1.7

Worktable Orig13. Age-adjusted death rates for 358 selected causes, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), and year	All origins 1/			Hispanic			Non-Hispanic 2/			Non-Hispanic white			Non-Hispanic black		
	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female
Alzheimer's disease (G30)															
2004.....	21.8	17.7	23.8	13.0	10.8	14.1	22.2	18.1	24.3	23.0	18.7	25.2	18.7	15.5	20.1
2003.....	21.4	17.5	23.3	13.4	12.2	14.1	21.7	17.8	23.7	22.6	18.4	24.7	16.9	14.9	17.6
2002.....	20.2	16.6	22.0	11.6	9.7	12.6	20.6	17.0	22.4	21.5	17.7	23.4	15.3	13.2	16.1
2001.....	19.1	16.1	20.5	10.7	9.2	11.5	19.4	16.4	20.8	20.2	17.0	21.7	14.8	13.3	15.3
2000.....	18.1	15.2	19.3	10.4	9.1	10.9	18.3	15.5	19.6	19.1	16.1	20.4	13.2	11.6	13.9
1999.....	16.5	14.4	17.4	9.6	8.1	10.3	16.7	14.6	17.6	17.4	15.2	18.4	11.4	10.3	11.8
Multiple sclerosis (G35)															
2004.....	1.1	0.8	1.4	0.2	0.2	0.3	1.2	0.8	1.5	1.2	0.9	1.5	1.1	0.8	1.3
2003.....	1.1	0.8	1.3	0.3	0.2	0.4	1.2	0.9	1.4	1.2	0.9	1.4	1.1	0.8	1.4
2002.....	1.1	0.8	1.3	0.3	0.2	0.3	1.1	0.8	1.4	1.2	0.9	1.4	1.1	0.7	1.5
2001.....	1.1	0.8	1.3	0.3	0.2	0.4	1.1	0.8	1.4	1.2	0.9	1.4	1.0	0.6	1.3
2000.....	1.0	0.8	1.2	0.3	0.3	0.4	1.1	0.8	1.3	1.1	0.8	1.4	1.0	0.7	1.3
1999.....	1.0	0.7	1.2	0.3	*	0.4	1.0	0.7	1.3	1.1	0.8	1.3	0.9	0.6	1.1
Epilepsy (G40-G41)															
2004.....	0.5	0.5	0.4	0.4	0.4	0.3	0.5	0.5	0.4	0.4	0.5	0.3	0.9	1.1	0.7
2003.....	0.5	0.5	0.4	0.4	0.4	0.3	0.5	0.5	0.4	0.4	0.5	0.4	1.0	1.4	0.8
2002.....	0.5	0.6	0.4	0.4	0.6	0.3	0.5	0.6	0.4	0.4	0.5	0.4	1.0	1.3	0.8
2001.....	0.4	0.5	0.4	0.4	0.5	0.3	0.4	0.5	0.4	0.4	0.5	0.4	1.0	1.2	0.8
2000.....	0.5	0.5	0.4	0.4	0.5	0.3	0.5	0.5	0.4	0.4	0.5	0.4	1.0	1.3	0.9
1999.....	0.5	0.6	0.4	0.4	0.4	0.3	0.5	0.6	0.4	0.4	0.5	0.4	1.1	1.4	0.9
Transient cerebral ischemic attacks and related syndromes (G45)															
2004.....	0.1	0.0	0.1	*	*	*	0.1	0.0	0.1	0.1	0.0	0.1	*	*	*
2003.....	0.1	0.1	0.1	*	*	*	0.1	0.1	0.1	0.1	0.1	0.1	*	*	*
2002.....	0.1	0.1	0.1	*	*	*	0.1	0.1	0.1	0.1	0.1	0.1	*	*	*
2001.....	0.1	0.1	0.1	*	*	*	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2000.....	0.1	0.1	0.1	*	*	*	0.1	0.1	0.1	0.1	0.1	0.1	*	*	*
1999.....	0.1	0.1	0.1	*	*	*	0.1	0.1	0.1	0.1	0.1	0.1	*	*	*
Infantile cerebral palsy (G80)															
2004.....	0.5	0.5	0.4	0.3	0.3	0.2	0.5	0.6	0.4	0.5	0.6	0.4	0.7	0.8	0.6
2003.....	0.5	0.5	0.4	0.3	0.3	0.3	0.5	0.6	0.4	0.4	0.5	0.4	0.7	0.8	0.6
2002.....	0.4	0.5	0.4	0.3	0.3	0.3	0.4	0.5	0.4	0.4	0.5	0.3	0.6	0.7	0.6
2001.....	0.4	0.5	0.3	0.2	0.3	0.2	0.4	0.5	0.4	0.4	0.5	0.3	0.5	0.6	0.5
2000.....	0.4	0.4	0.3	0.2	0.3	0.2	0.4	0.4	0.4	0.4	0.4	0.3	0.6	0.6	0.5
1999.....	0.4	0.4	0.3	0.3	0.3	0.3	0.4	0.4	0.3	0.4	0.4	0.3	0.5	0.6	0.4
All other diseases of nervous system (G10-G12,G23-G25,G31,G36-G37,G43-G44,G47-G72,G81-G98)															
2004.....	6.8	8.1	5.7	4.5	5.0	4.0	7.0	8.3	5.9	7.0	8.4	5.8	8.2	9.3	7.5
2003.....	7.0	8.4	5.9	4.8	5.7	4.0	7.2	8.6	6.1	7.2	8.7	6.0	8.2	9.3	7.4
2002.....	6.9	8.3	5.8	4.5	5.0	3.9	7.1	8.5	5.9	7.1	8.6	5.9	8.1	9.5	7.0
2001.....	6.7	8.1	5.7	4.1	5.0	3.3	6.9	8.4	5.8	6.9	8.3	5.8	8.2	9.7	7.3
2000.....	6.8	8.3	5.7	4.5	5.7	3.6	7.0	8.4	5.8	7.0	8.5	5.8	8.0	9.6	6.9
1999.....	6.6	8.1	5.6	4.2	5.2	3.5	6.8	8.3	5.7	6.8	8.4	5.6	7.8	8.9	7.2
VII. Diseases of the eye and adnexa (H00-H57)															
2004.....	0.0	*	0.0	*	*	*	0.0	*	0.0	0.0	*	0.0	*	*	*
2003.....	0.0	*	0.0	*	*	*	0.0	*	0.0	0.0	*	0.0	*	*	*
2002.....	0.0	*	0.0	*	*	*	0.0	*	0.0	0.0	*	0.0	*	*	*
2001.....	0.0	*	0.0	*	*	*	0.0	*	0.0	0.0	*	*	*	*	*
2000.....	0.0	0.0	0.0	*	*	*	0.0	0.0	0.0	0.0	*	0.0	*	*	*
1999.....	0.0	*	0.0	*	*	*	0.0	*	0.0	0.0	*	0.0	*	*	*

Worktable Orig13. Age-adjusted death rates for 358 selected causes, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), and year	All origins 1/			Hispanic			Non-Hispanic 2/			Non-Hispanic white			Non-Hispanic black		
	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female
VIII. Diseases of the ear and mastoid process (H60-H93)															
2004.....	0.0	0.0	0.0	*	*	*	0.0	0.0	0.0	0.0	*	0.0	*	*	*
2003.....	0.0	0.0	0.0	*	*	*	0.0	0.0	0.0	0.0	*	0.0	*	*	*
2002.....	0.0	0.0	0.0	*	*	*	0.0	0.0	0.0	0.0	0.0	*	*	*	*
2001.....	0.0	0.0	0.0	*	*	*	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2000.....	0.0	0.0	0.0	*	*	*	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
1999.....	0.0	0.0	0.0	*	*	*	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
IX. Diseases of the circulatory system (I00-I99)															
2004.....	288.0	341.7	245.3	211.2	251.9	178.3	292.3	347.0	248.9	284.8	339.6	240.9	390.6	461.2	338.8
2003.....	307.7	365.0	262.3	229.3	267.5	197.8	311.7	370.2	265.4	303.7	362.4	256.9	414.8	489.2	360.1
2002.....	319.0	378.8	272.1	237.5	281.1	202.9	322.8	383.6	275.1	315.0	376.3	266.5	424.3	496.8	371.1
2001.....	328.2	389.8	280.4	253.4	299.5	217.6	331.4	393.9	283.0	323.2	386.0	274.0	435.3	514.1	379.5
2000.....	341.3	409.0	290.0	259.6	308.5	221.7	344.6	413.1	292.5	336.5	405.7	283.5	446.8	526.5	390.9
1999.....	350.8	420.5	297.9	269.0	321.9	228.8	353.7	424.3	300.3	345.6	416.3	291.4	454.5	539.6	395.2
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)															
2004.....	1.1	0.8	1.3	0.8	0.7	0.9	1.1	0.8	1.3	1.1	0.8	1.3	0.8	0.6	1.0
2003.....	1.2	1.0	1.4	0.8	0.6	1.0	1.2	1.0	1.4	1.3	1.0	1.4	1.1	0.9	1.2
2002.....	1.2	0.9	1.5	1.1	0.6	1.4	1.2	0.9	1.5	1.3	0.9	1.5	0.9	0.8	1.0
2001.....	1.2	0.9	1.5	0.9	0.7	1.1	1.2	0.9	1.5	1.2	0.9	1.4	1.1	0.7	1.4
2000.....	1.3	1.0	1.5	0.9	0.7	1.1	1.3	1.0	1.6	1.3	1.0	1.6	1.0	0.7	1.2
1999.....	1.3	0.9	1.6	1.1	0.4	1.5	1.3	0.9	1.6	1.4	1.0	1.6	1.0	0.8	1.2
Acute rheumatic fever (I00-I02)															
2004.....	0.0	*	0.0	*	*	*	0.0	*	*	0.0	*	*	*	*	*
2003.....	0.0	*	0.0	*	*	*	0.0	*	0.0	0.0	*	*	*	*	*
2002.....	0.0	*	0.0	*	*	*	0.0	*	0.0	0.0	*	0.0	*	*	*
2001.....	0.0	0.0	0.0	*	*	*	0.0	0.0	0.0	0.0	*	0.0	*	*	*
2000.....	0.0	*	0.0	*	*	*	0.0	*	0.0	0.0	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	0.0	*	*	0.0	*	*	*	*	*
Chronic rheumatic heart diseases (I05-I09)															
2004.....	1.1	0.8	1.3	0.8	0.7	0.9	1.1	0.8	1.3	1.1	0.8	1.3	0.8	0.6	1.0
2003.....	1.2	1.0	1.4	0.8	0.6	1.0	1.2	1.0	1.4	1.2	1.0	1.4	1.1	0.9	1.2
2002.....	1.2	0.9	1.5	1.1	0.6	1.4	1.2	0.9	1.5	1.3	0.9	1.5	0.9	0.8	1.0
2001.....	1.2	0.9	1.4	0.9	0.7	1.1	1.2	0.9	1.4	1.2	0.9	1.4	1.1	0.7	1.3
2000.....	1.3	0.9	1.5	0.9	0.6	1.1	1.3	1.0	1.5	1.3	1.0	1.6	1.0	0.7	1.2
1999.....	1.3	0.9	1.6	1.1	0.4	1.5	1.3	0.9	1.6	1.4	0.9	1.6	1.0	0.7	1.2
Rheumatic mitral valve diseases (I05)															
2004.....	0.5	0.4	0.6	0.4	0.3	0.4	0.5	0.4	0.6	0.5	0.4	0.6	0.4	0.3	0.5
2003.....	0.6	0.5	0.7	0.4	0.4	0.4	0.6	0.5	0.7	0.6	0.5	0.7	0.5	0.5	0.6
2002.....	0.6	0.4	0.8	0.5	0.3	0.7	0.6	0.4	0.7	0.6	0.4	0.8	0.5	0.4	0.6
2001.....	0.7	0.4	0.8	0.5	0.3	0.6	0.7	0.5	0.8	0.7	0.5	0.8	0.6	0.4	0.8
2000.....	0.7	0.5	0.9	0.5	0.4	0.6	0.7	0.5	0.9	0.8	0.6	0.9	0.5	0.4	0.6
1999.....	0.7	0.5	0.9	0.6	0.3	0.9	0.7	0.5	0.9	0.8	0.5	0.9	0.6	0.5	0.7
Rheumatic aortic valve diseases (I06)															
2004.....	0.0	0.0	0.0	*	*	*	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2003.....	0.0	0.0	0.0	*	*	*	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2002.....	0.0	0.0	0.1	*	*	*	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2001.....	0.0	0.0	0.0	*	*	*	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2000.....	0.0	*	*	*	*	*	0.0	*	*	0.0	*	*	*	*	*
1999.....	0.0	*	0.0	*	*	*	0.0	*	0.0	0.0	*	0.0	*	*	*

Worktable Orig13. Age-adjusted death rates for 358 selected causes, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), and year	All origins 1/			Hispanic			Non-Hispanic 2/			Non-Hispanic white			Non-Hispanic black		
	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female
Disorders of both mitral and aortic valves (I08.0)															
2004.....	0.2	0.2	0.2	0.2	*	*	0.2	0.2	0.3	0.3	0.2	0.3	0.2	0.2	0.2
2003.....	0.3	0.2	0.3	0.2	*	*	0.3	0.2	0.3	0.3	0.3	0.3	0.2	0.2	0.2
2002.....	0.2	0.2	0.2	0.2	*	*	0.2	0.1	0.2	0.2	0.2	0.2	0.1	*	*
2001.....	0.1	0.1	0.1	*	*	*	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2000.....	0.1	0.1	0.1	*	*	*	0.1	0.1	0.1	0.1	0.1	0.1	*	*	*
1999.....	0.1	0.1	0.1	*	*	*	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)															
2004.....	0.3	0.2	0.4	0.2	*	0.3	0.3	0.2	0.4	0.3	0.2	0.4	0.2	*	0.3
2003.....	0.3	0.2	0.4	0.2	*	0.4	0.3	0.2	0.4	0.3	0.2	0.4	0.3	0.2	0.4
2002.....	0.4	0.2	0.4	0.4	*	0.5	0.4	0.3	0.4	0.4	0.3	0.4	0.3	0.2	0.3
2001.....	0.4	0.3	0.5	0.3	*	0.3	0.4	0.3	0.5	0.4	0.2	0.5	0.3	0.2	0.4
2000.....	0.4	0.3	0.5	0.3	*	0.4	0.4	0.3	0.5	0.4	0.3	0.5	0.4	0.2	0.5
1999.....	0.4	0.3	0.5	0.3	*	0.5	0.5	0.3	0.5	0.4	0.3	0.5	0.3	0.2	0.3
Hypertensive diseases (I10-I13)															
2004.....	18.1	18.7	17.0	15.5	17.0	13.9	18.3	18.8	17.2	15.3	15.5	14.4	46.5	51.8	41.7
2003.....	18.1	18.3	17.2	16.5	17.7	15.0	18.1	18.3	17.3	15.1	15.0	14.5	46.8	51.9	42.3
2002.....	17.1	17.4	16.3	15.8	17.0	14.5	17.1	17.4	16.3	14.2	14.1	13.6	45.4	50.2	41.2
2001.....	16.5	16.6	15.8	15.3	16.2	14.2	16.5	16.6	15.9	13.7	13.5	13.3	43.5	48.3	39.5
2000.....	16.2	16.3	15.5	14.2	15.4	13.0	16.2	16.3	15.5	13.3	13.2	12.8	43.9	47.7	40.5
1999.....	15.8	15.9	15.2	14.1	14.7	13.3	15.8	15.9	15.2	12.9	12.7	12.5	43.6	48.0	39.9
Essential (primary) hypertension (I10)															
2004.....	4.4	4.0	4.6	3.2	2.8	3.3	4.5	4.0	4.7	4.1	3.6	4.3	8.4	8.1	8.2
2003.....	4.4	3.7	4.6	3.2	2.5	3.5	4.4	3.8	4.7	4.0	3.3	4.3	8.6	8.7	8.2
2002.....	4.0	3.5	4.2	2.8	2.3	3.2	4.1	3.5	4.2	3.7	3.2	3.9	7.8	7.7	7.6
2001.....	3.9	3.5	4.0	3.2	2.7	3.3	4.0	3.5	4.1	3.6	3.1	3.7	7.7	8.3	7.2
2000.....	3.7	3.4	3.8	3.0	3.2	2.8	3.8	3.4	3.8	3.4	2.9	3.5	7.4	7.7	7.1
1999.....	3.6	3.3	3.7	2.8	2.6	2.8	3.7	3.3	3.8	3.3	2.9	3.4	7.3	7.5	7.1
Hypertensive heart disease (I11)															
2004.....	9.5	10.1	8.5	7.8	9.1	6.7	9.6	10.2	8.6	7.9	8.3	7.2	24.6	28.8	21.0
2003.....	9.6	10.1	8.8	8.9	10.3	7.5	9.6	10.0	8.8	7.9	8.1	7.3	25.4	29.1	22.2
2002.....	9.1	9.5	8.3	8.4	9.7	7.1	9.2	9.5	8.4	7.5	7.6	7.0	24.7	28.2	21.5
2001.....	8.7	8.9	8.1	7.7	8.8	6.7	8.7	8.9	8.2	7.2	7.1	6.8	23.4	26.5	20.7
2000.....	8.6	8.7	8.1	7.0	7.5	6.3	8.7	8.7	8.1	7.1	7.0	6.7	24.0	26.6	21.5
1999.....	8.3	8.3	7.9	6.9	7.1	6.6	8.4	8.4	7.9	6.8	6.7	6.5	23.2	26.0	20.7
Hypertensive renal disease (I12)															
2004.....	3.2	3.5	3.0	3.4	3.9	3.0	3.2	3.5	3.0	2.4	2.8	2.2	10.2	11.2	9.4
2003.....	3.1	3.4	2.8	3.5	3.7	3.3	3.0	3.3	2.8	2.4	2.7	2.2	9.3	10.2	8.7
2002.....	3.0	3.3	2.8	3.4	3.7	3.2	2.9	3.3	2.7	2.3	2.6	2.1	9.4	10.4	8.8
2001.....	2.9	3.2	2.7	3.3	3.5	3.2	2.9	3.1	2.7	2.2	2.5	2.1	9.1	9.8	8.5
2000.....	2.8	3.1	2.6	3.2	3.5	3.0	2.8	3.1	2.6	2.1	2.4	1.9	9.2	9.6	8.8
1999.....	2.6	2.9	2.4	3.2	3.8	2.9	2.5	2.8	2.4	1.9	2.2	1.7	8.9	9.9	8.2
Hypertensive heart and renal disease (I13)															
2004.....	1.0	1.1	0.9	1.0	1.2	0.9	1.0	1.1	0.9	0.7	0.8	0.7	3.3	3.7	3.0
2003.....	1.1	1.1	1.0	1.0	1.2	0.9	1.1	1.2	1.0	0.8	0.9	0.7	3.5	3.8	3.2
2002.....	1.0	1.0	0.9	1.1	1.2	1.0	1.0	1.0	0.9	0.7	0.7	0.7	3.6	3.9	3.2
2001.....	1.0	1.1	1.0	1.1	1.3	1.0	1.0	1.0	0.9	0.7	0.8	0.7	3.4	3.6	3.1
2000.....	1.0	1.1	0.9	1.0	1.2	0.9	1.0	1.1	0.9	0.8	0.8	0.7	3.4	3.7	3.1
1999.....	1.2	1.4	1.1	1.1	1.2	1.0	1.2	1.4	1.1	0.9	1.0	0.8	4.2	4.5	3.9

Worktable Orig13. Age-adjusted death rates for 358 selected causes, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), and year	All origins 1/			Hispanic			Non-Hispanic 2/			Non-Hispanic white			Non-Hispanic black		
	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female
Ischemic heart disease (I20-I25)															
2004.....	150.2	194.1	116.7	119.2	149.6	95.3	151.9	196.8	117.8	150.9	196.8	115.6	182.0	226.7	150.5
2003.....	162.9	209.9	127.2	130.0	158.0	107.4	164.6	212.7	128.1	163.3	212.3	125.7	197.2	246.1	162.7
2002.....	170.8	220.4	133.6	138.3	172.2	112.2	172.3	222.7	134.4	171.0	222.7	131.7	205.1	253.2	171.4
2001.....	177.8	228.5	139.9	149.9	185.8	122.6	179.0	230.5	140.5	177.5	230.2	137.7	213.5	264.4	178.4
2000.....	186.8	241.4	146.5	153.2	190.1	125.1	188.0	243.5	147.2	186.6	243.5	144.3	220.4	271.9	185.0
1999.....	194.6	251.2	152.9	162.2	200.1	133.1	195.7	253.1	153.5	194.3	253.0	150.8	226.6	280.3	189.7
Acute myocardial infarction (I21-I22)															
2004.....	52.2	67.8	40.2	40.9	51.2	32.7	53.0	68.9	40.7	52.9	69.4	39.9	62.9	76.0	53.5
2003.....	57.9	75.0	44.8	45.4	54.8	37.9	58.7	76.3	45.2	58.5	76.6	44.3	69.6	85.5	58.4
2002.....	62.1	80.6	48.1	49.5	61.6	40.2	62.8	81.7	48.5	62.5	82.1	47.4	75.0	90.9	64.1
2001.....	65.4	84.9	50.9	52.8	66.2	43.0	66.1	86.0	51.2	65.8	86.5	50.1	77.8	94.3	66.8
2000.....	69.9	91.3	54.1	54.1	67.6	43.9	70.6	92.5	54.5	70.4	93.0	53.4	82.0	100.1	70.0
1999.....	73.2	96.1	56.3	57.2	71.1	46.6	73.9	97.3	56.7	73.8	98.0	55.7	84.1	103.3	71.3
Other acute ischemic heart diseases (I24)															
2004.....	0.8	1.0	0.6	0.2	0.2	*	0.9	1.1	0.7	0.8	1.0	0.6	1.4	2.1	1.0
2003.....	1.1	1.3	0.9	0.3	0.2	0.3	1.1	1.4	0.9	1.1	1.3	0.9	1.7	2.4	1.2
2002.....	1.2	1.5	0.9	0.4	0.5	0.3	1.2	1.5	1.0	1.2	1.5	0.9	1.7	2.3	1.3
2001.....	1.2	1.5	0.9	0.4	0.7	0.2	1.2	1.5	1.0	1.2	1.5	0.9	1.8	2.4	1.4
2000.....	1.2	1.5	0.9	0.5	0.7	0.4	1.2	1.6	1.0	1.2	1.5	0.9	1.8	2.5	1.4
1999.....	1.2	1.5	0.9	0.4	0.4	0.4	1.2	1.6	1.0	1.2	1.5	0.9	1.8	2.5	1.3
Other forms of chronic ischemic heart disease (I20,I25)															
2004.....	97.1	125.3	75.8	78.1	98.2	62.4	98.0	126.8	76.4	97.2	126.4	75.1	117.6	148.6	96.0
2003.....	103.8	133.5	81.5	84.3	102.9	69.3	104.7	135.0	82.0	103.7	134.4	80.5	125.9	158.1	103.1
2002.....	107.6	138.3	84.5	88.4	110.0	71.8	108.3	139.5	84.9	107.3	139.1	83.4	128.5	160.0	106.1
2001.....	111.2	142.1	88.1	96.6	118.9	79.4	111.7	143.0	88.3	110.5	142.3	86.7	134.0	167.8	110.2
2000.....	115.7	148.6	91.5	98.7	121.9	80.8	116.1	149.4	91.7	115.1	149.0	90.0	136.5	169.3	113.7
1999.....	120.2	153.6	95.6	104.6	128.5	86.2	120.5	154.2	95.8	119.4	153.5	94.1	140.7	174.5	117.0
Atherosclerotic cardiovascular disease, so described (I25.0)															
2004.....	21.1	27.5	15.7	17.5	24.6	11.8	21.2	27.6	15.9	19.9	25.9	14.8	36.5	49.5	27.3
2003.....	22.7	29.6	17.1	19.1	26.7	13.0	22.9	29.6	17.3	21.5	27.6	16.2	38.6	53.1	28.4
2002.....	23.5	30.4	17.8	19.3	26.2	13.9	23.6	30.4	17.9	22.1	28.4	16.8	40.6	55.2	30.3
2001.....	24.1	30.8	18.5	21.0	28.5	15.1	24.1	30.8	18.6	22.5	28.6	17.3	42.1	56.4	32.0
2000.....	25.2	32.3	19.4	21.3	29.0	15.4	25.2	32.2	19.5	23.5	30.0	18.2	43.9	59.2	33.3
1999.....	26.4	33.5	20.6	23.2	31.1	17.0	26.4	33.4	20.7	24.6	30.9	19.3	46.0	61.5	35.1
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)															
2004.....	76.0	97.8	60.1	60.6	73.6	50.6	76.8	99.2	60.6	77.3	100.5	60.2	81.2	99.1	68.7
2003.....	81.1	104.0	64.4	65.2	76.3	56.3	81.9	105.4	64.7	82.2	106.8	64.2	87.3	105.0	74.7
2002.....	84.1	108.0	66.7	69.1	83.9	57.9	84.7	109.1	66.9	85.2	110.7	66.6	87.9	104.8	75.8
2001.....	87.1	111.3	69.6	75.6	90.5	64.3	87.6	112.2	69.7	88.0	113.7	69.4	91.8	111.4	78.2
2000.....	90.5	116.3	72.0	77.3	92.9	65.4	90.9	117.2	72.2	91.5	119.0	71.8	92.6	110.1	80.4
1999.....	93.8	120.1	75.0	81.4	97.4	69.2	94.1	120.9	75.1	94.8	122.6	74.8	94.7	113.0	81.9
Other heart diseases (I26-I51)															
2004.....	55.2	61.7	49.9	29.5	33.3	26.3	56.8	63.5	51.3	55.6	61.9	50.3	74.0	87.4	64.5
2003.....	57.6	64.5	52.0	32.4	36.6	29.0	59.0	66.2	53.2	57.7	64.6	52.0	77.0	89.3	67.9
2002.....	58.5	65.5	52.8	31.5	36.1	27.9	60.0	67.2	54.1	58.6	65.7	52.9	77.8	89.3	69.2
2001.....	59.1	66.1	53.4	32.6	36.2	29.6	60.4	67.6	54.6	58.9	65.8	53.3	79.1	93.3	69.3
2000.....	60.0	67.9	53.8	33.9	38.7	30.2	61.2	69.3	54.9	59.7	67.6	53.5	79.7	93.7	70.1
1999.....	60.9	69.2	54.5	34.5	40.4	30.0	62.1	70.5	55.5	60.4	68.4	54.1	82.3	98.8	71.1

Worktable Orig13. Age-adjusted death rates for 358 selected causes, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), and year	All origins 1/			Hispanic			Non-Hispanic 2/			Non-Hispanic white			Non-Hispanic black		
	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)															
2004.....	4.4	4.0	4.7	2.3	1.9	2.5	4.6	4.2	4.8	4.3	4.0	4.5	7.7	7.5	8.0
2003.....	4.8	4.6	5.0	2.3	1.9	2.7	4.9	4.7	5.1	4.6	4.5	4.8	8.4	8.3	8.5
2002.....	4.8	4.4	5.0	2.5	2.1	2.8	4.9	4.6	5.2	4.6	4.4	4.8	8.2	7.3	8.9
2001.....	4.7	4.5	4.9	2.4	2.6	2.4	4.8	4.6	5.1	4.6	4.3	4.7	8.1	8.0	8.4
2000.....	4.7	4.5	4.9	2.4	2.0	2.6	4.8	4.6	5.0	4.6	4.4	4.7	8.1	7.8	8.5
1999.....	4.8	4.7	5.0	2.7	2.5	2.8	5.0	4.8	5.1	4.6	4.5	4.6	8.8	8.5	9.1
Pulmonary embolism (I26)															
2004.....	2.7	2.7	2.7	1.3	1.1	1.4	2.8	2.8	2.9	2.7	2.6	2.7	5.0	5.3	4.8
2003.....	3.0	3.0	2.9	1.2	1.2	1.2	3.1	3.1	3.1	2.9	2.9	2.9	5.5	6.0	5.3
2002.....	3.0	2.9	3.1	1.4	1.3	1.5	3.1	3.0	3.2	3.0	2.9	3.0	5.3	5.1	5.5
2001.....	3.1	3.0	3.1	1.4	1.6	1.3	3.2	3.1	3.2	3.0	2.9	3.0	5.4	5.7	5.3
2000.....	3.0	3.0	3.0	1.2	1.2	1.2	3.2	3.2	3.1	3.0	3.0	2.9	5.5	5.8	5.5
1999.....	3.3	3.3	3.3	1.7	1.6	1.7	3.4	3.4	3.4	3.1	3.2	3.1	6.2	6.6	6.0
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)															
2004.....	1.7	1.4	1.9	1.0	0.8	1.2	1.7	1.4	2.0	1.6	1.4	1.8	2.8	2.2	3.2
2003.....	1.8	1.6	2.0	1.1	0.7	1.5	1.9	1.6	2.1	1.7	1.6	1.9	2.8	2.3	3.2
2002.....	1.8	1.5	1.9	1.1	0.8	1.3	1.8	1.6	2.0	1.7	1.5	1.9	2.9	2.3	3.4
2001.....	1.7	1.5	1.8	1.0	0.9	1.1	1.7	1.5	1.9	1.6	1.4	1.7	2.7	2.3	3.1
2000.....	1.7	1.5	1.8	1.1	0.8	1.4	1.7	1.5	1.9	1.6	1.4	1.7	2.6	2.0	3.0
1999.....	1.5	1.4	1.6	1.0	0.9	1.1	1.6	1.5	1.7	1.5	1.4	1.5	2.6	2.0	3.1
Other non-pulmonary forms of heart disease (I30-I51)															
2004.....	50.8	57.7	45.3	27.3	31.4	23.7	52.2	59.3	46.5	51.3	57.9	45.8	66.3	79.9	56.5
2003.....	52.8	60.0	47.0	30.1	34.7	26.3	54.1	61.4	48.1	53.0	60.1	47.2	68.6	81.0	59.4
2002.....	53.8	61.1	47.8	29.0	34.0	25.1	55.1	62.7	48.9	54.0	61.3	48.0	69.6	82.0	60.3
2001.....	54.4	61.6	48.5	30.2	33.6	27.1	55.5	63.0	49.5	54.4	61.5	48.6	70.9	85.3	60.9
2000.....	55.2	63.4	48.9	31.5	36.8	27.6	56.3	64.7	49.8	55.1	63.2	48.9	71.6	85.9	61.6
1999.....	56.1	64.5	49.6	31.8	37.9	27.2	57.2	65.7	50.5	55.8	63.8	49.4	73.5	90.2	62.1
Acute and subacute endocarditis (I33)															
2004.....	0.4	0.5	0.3	0.4	0.6	0.2	0.4	0.5	0.3	0.4	0.5	0.3	0.7	0.8	0.6
2003.....	0.4	0.5	0.3	0.4	0.4	0.3	0.4	0.5	0.3	0.4	0.5	0.3	0.8	0.9	0.6
2002.....	0.4	0.5	0.3	0.3	0.5	0.2	0.4	0.5	0.3	0.4	0.5	0.3	0.7	0.8	0.6
2001.....	0.4	0.5	0.3	0.3	0.3	0.2	0.4	0.6	0.3	0.4	0.5	0.3	0.8	1.1	0.6
2000.....	0.4	0.5	0.3	0.4	0.4	0.3	0.4	0.5	0.3	0.4	0.5	0.3	0.8	0.9	0.7
1999.....	0.4	0.5	0.3	0.3	0.6	0.2	0.4	0.5	0.3	0.4	0.5	0.3	0.8	0.9	0.6
Diseases of pericardium and acute myocarditis (I30-I31,I40)															
2004.....	0.3	0.3	0.2	0.2	0.2	0.2	0.3	0.3	0.2	0.2	0.2	0.2	0.5	0.6	0.4
2003.....	0.3	0.3	0.3	0.2	0.2	0.2	0.3	0.4	0.3	0.3	0.3	0.2	0.5	0.5	0.5
2002.....	0.3	0.3	0.3	0.3	0.3	0.2	0.3	0.3	0.3	0.3	0.3	0.2	0.5	0.5	0.5
2001.....	0.3	0.3	0.3	0.3	0.2	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.5	0.5	0.5
2000.....	0.3	0.3	0.2	0.2	0.3	0.2	0.3	0.3	0.2	0.2	0.3	0.2	0.4	0.5	0.4
1999.....	0.3	0.4	0.3	0.3	0.2	0.3	0.3	0.4	0.3	0.3	0.3	0.3	0.5	0.6	0.5
Other diseases of endocardium (I34-I38)															
2004.....	6.7	6.9	6.5	3.6	3.9	3.4	6.9	7.1	6.7	7.1	7.3	6.9	4.9	5.2	4.7
2003.....	6.8	7.0	6.6	3.7	3.8	3.7	6.9	7.2	6.7	7.2	7.4	6.9	5.0	5.1	4.8
2002.....	6.9	7.1	6.7	3.7	4.1	3.4	7.0	7.2	6.9	7.3	7.5	7.1	4.9	4.6	4.9
2001.....	6.9	7.2	6.6	3.9	4.1	3.7	7.1	7.4	6.8	7.3	7.6	7.0	5.0	5.3	4.7
2000.....	7.2	7.5	6.9	4.4	4.7	4.1	7.3	7.6	7.0	7.5	7.8	7.3	5.3	5.5	5.1
1999.....	7.2	7.6	6.9	4.2	4.8	3.8	7.3	7.7	7.0	7.6	7.9	7.2	5.4	5.4	5.2

Worktable Orig13. Age-adjusted death rates for 358 selected causes, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), and year	All origins 1/			Hispanic			Non-Hispanic 2/			Non-Hispanic white			Non-Hispanic black		
	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female
Nonrheumatic mitral valve disorders (I34)															
2004.....	0.8	0.8	0.9	0.4	0.4	0.5	0.9	0.8	0.9	0.9	0.8	1.0	0.7	0.6	0.8
2003.....	0.9	0.8	0.9	0.5	0.4	0.5	0.9	0.8	0.9	0.9	0.8	1.0	0.7	0.6	0.7
2002.....	1.0	0.8	1.0	0.5	0.4	0.6	1.0	0.9	1.1	1.0	0.9	1.1	0.7	0.6	0.7
2001.....	1.0	0.9	1.1	0.6	0.5	0.7	1.0	0.9	1.1	1.0	0.9	1.1	0.7	0.5	0.8
2000.....	1.0	1.0	1.1	0.5	0.7	0.5	1.1	1.0	1.1	1.1	1.0	1.2	0.8	0.6	0.9
1999.....	1.1	1.0	1.1	0.5	*	0.5	1.1	1.0	1.1	1.1	1.0	1.2	0.8	0.6	0.9
Nonrheumatic aortic valve disorders (I35)															
2004.....	4.2	4.4	4.0	2.1	2.4	2.0	4.3	4.5	4.1	4.5	4.8	4.3	2.4	2.6	2.2
2003.....	4.2	4.6	4.0	2.3	2.4	2.2	4.3	4.7	4.0	4.6	4.9	4.3	2.4	2.5	2.3
2002.....	4.3	4.6	4.0	2.3	2.8	2.0	4.4	4.7	4.1	4.6	5.0	4.3	2.5	2.5	2.4
2001.....	4.3	4.7	4.0	2.4	2.6	2.2	4.4	4.8	4.1	4.6	5.0	4.3	2.5	3.0	2.2
2000.....	4.5	4.9	4.2	2.8	3.3	2.5	4.5	5.0	4.3	4.8	5.2	4.5	2.7	2.9	2.5
1999.....	4.5	5.0	4.2	2.8	3.3	2.3	4.6	5.0	4.2	4.8	5.3	4.4	2.8	3.1	2.5
All other diseases of endocardium (I36-I38)															
2004.....	1.7	1.7	1.6	1.0	1.1	0.9	1.7	1.7	1.7	1.7	1.7	1.7	1.8	1.9	1.7
2003.....	1.7	1.6	1.7	1.0	1.0	1.0	1.7	1.7	1.7	1.7	1.7	1.7	1.9	2.0	1.8
2002.....	1.6	1.6	1.7	0.8	0.9	0.8	1.7	1.6	1.7	1.7	1.6	1.7	1.7	1.6	1.7
2001.....	1.6	1.7	1.6	0.9	1.0	0.8	1.6	1.7	1.6	1.6	1.7	1.6	1.7	1.8	1.7
2000.....	1.6	1.6	1.6	1.0	0.7	1.1	1.7	1.6	1.6	1.6	1.6	1.6	1.8	2.0	1.7
1999.....	1.6	1.6	1.6	1.0	1.1	0.9	1.7	1.7	1.7	1.7	1.6	1.7	1.8	1.8	1.8
Cardiomyopathy (I42)															
2004.....	8.5	11.9	6.1	6.1	8.4	4.3	8.7	12.2	6.2	8.0	11.3	5.6	15.2	21.0	11.0
2003.....	9.1	12.6	6.5	7.2	9.9	5.1	9.3	12.8	6.7	8.5	11.9	6.1	16.2	22.5	11.8
2002.....	9.6	13.4	6.8	7.0	9.8	4.9	9.8	13.6	6.9	9.0	12.7	6.3	16.9	23.4	12.3
2001.....	9.5	13.1	6.8	7.5	9.6	5.7	9.6	13.3	6.9	8.8	12.3	6.2	17.3	23.8	12.7
2000.....	9.8	13.8	7.0	7.7	10.3	5.8	10.0	14.0	7.0	9.0	12.8	6.3	18.1	25.3	13.1
1999.....	10.0	13.9	7.1	7.4	10.2	5.2	10.1	14.0	7.2	9.2	12.8	6.5	18.6	26.3	13.1
Conduction disorders and cardiac dysrhythmias (I44-I49)															
2004.....	12.5	13.8	11.3	4.5	4.9	4.2	13.0	14.4	11.7	12.6	13.9	11.4	18.0	21.6	15.6
2003.....	13.2	14.6	12.0	5.3	5.9	4.7	13.7	15.2	12.3	13.3	14.7	12.0	18.7	21.7	16.5
2002.....	13.6	15.1	12.2	5.0	5.7	4.4	14.1	15.7	12.6	13.7	15.2	12.3	19.1	22.0	17.0
2001.....	13.6	15.1	12.3	5.4	5.8	5.1	14.1	15.6	12.7	13.6	15.0	12.3	19.4	23.2	16.8
2000.....	13.9	15.6	12.4	5.7	6.6	5.1	14.3	16.1	12.8	13.9	15.7	12.4	19.1	22.1	16.9
1999.....	14.7	16.7	13.0	6.2	7.5	5.3	15.1	17.2	13.4	14.7	16.7	13.1	20.3	24.8	17.3
Heart failure (I50)															
2004.....	18.9	19.9	18.0	9.7	10.2	9.2	19.4	20.5	18.4	19.5	20.5	18.6	21.3	23.7	19.7
2003.....	19.4	20.6	18.4	10.8	11.7	10.1	19.8	21.0	18.8	19.8	21.1	18.8	22.2	23.9	20.9
2002.....	19.5	20.4	18.6	10.2	10.7	9.8	19.9	20.9	18.9	19.9	21.0	19.0	22.1	23.7	20.7
2001.....	20.2	21.2	19.2	10.7	10.9	10.4	20.5	21.6	19.6	20.6	21.7	19.7	22.4	24.7	20.9
2000.....	20.3	21.5	19.2	10.9	11.6	10.4	20.6	22.0	19.6	20.7	22.0	19.6	22.4	24.6	21.0
1999.....	20.3	21.4	19.3	11.1	11.7	10.6	20.6	21.8	19.6	20.6	21.7	19.6	22.8	25.5	21.1
Congestive heart failure (I50.0)															
2004.....	17.6	18.5	16.8	9.0	9.3	8.6	18.0	18.9	17.2	18.2	19.1	17.4	19.1	21.1	17.7
2003.....	18.1	19.1	17.3	10.1	10.8	9.5	18.5	19.5	17.6	18.6	19.7	17.7	19.9	21.2	18.8
2002.....	18.0	18.8	17.3	9.4	9.6	9.1	18.4	19.3	17.7	18.5	19.4	17.8	19.9	21.1	18.8
2001.....	18.7	19.5	18.0	9.9	9.8	9.8	19.1	19.9	18.3	19.2	20.1	18.4	20.2	22.0	19.1
2000.....	18.8	19.8	17.9	10.0	10.4	9.7	19.1	20.2	18.2	19.2	20.4	18.3	20.0	21.7	19.0
1999.....	18.8	19.8	17.9	10.4	10.9	10.0	19.1	20.1	18.2	19.1	20.1	18.3	20.5	23.0	19.1

Worktable Orig13. Age-adjusted death rates for 358 selected causes, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), and year	All origins 1/			Hispanic			Non-Hispanic 2/			Non-Hispanic white			Non-Hispanic black		
	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female
Other and unspecified heart failure (I50.1-I50.9)															
2004.....	1.3	1.5	1.1	0.7	0.8	0.6	1.3	1.5	1.2	1.3	1.4	1.1	2.2	2.7	1.9
2003.....	1.3	1.5	1.1	0.7	0.9	0.6	1.3	1.6	1.2	1.2	1.5	1.1	2.3	2.7	2.0
2002.....	1.4	1.6	1.3	0.8	1.0	0.7	1.4	1.6	1.3	1.3	1.6	1.2	2.2	2.6	2.0
2001.....	1.5	1.7	1.3	0.8	1.1	0.7	1.5	1.7	1.3	1.4	1.6	1.2	2.2	2.8	1.8
2000.....	1.5	1.7	1.4	0.8	1.2	0.6	1.5	1.7	1.4	1.5	1.6	1.3	2.3	2.9	2.0
1999.....	1.5	1.7	1.3	0.7	0.8	0.6	1.5	1.7	1.4	1.5	1.7	1.3	2.2	2.6	2.0
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)															
2004.....	0.2	0.2	0.2	0.1	0.1	0.1	0.2	0.2	0.2	0.2	0.2	0.2	0.4	0.5	0.4
2003.....	0.2	0.2	0.2	0.2	0.2	0.1	0.2	0.2	0.2	0.2	0.2	0.2	0.4	0.4	0.3
2002.....	0.3	0.3	0.3	0.2	0.2	0.2	0.3	0.3	0.3	0.3	0.3	0.3	0.5	0.6	0.4
2001.....	0.3	0.3	0.2	0.1	0.2	*	0.3	0.3	0.3	0.2	0.2	0.2	0.5	0.6	0.4
2000.....	0.3	0.3	0.2	0.2	0.2	*	0.3	0.3	0.2	0.2	0.2	0.2	0.4	0.4	0.4
1999.....	0.3	0.3	0.2	0.2	0.2	0.2	0.3	0.3	0.2	0.2	0.3	0.2	0.4	0.6	0.3
All other and ill-defined forms of heart disease (I50.0-I51.3,I51.6-I51.9)															
2004.....	3.3	4.0	2.7	2.5	3.1	2.0	3.4	4.1	2.7	3.2	4.0	2.6	5.2	6.6	4.2
2003.....	3.3	4.0	2.7	2.4	2.6	2.1	3.4	4.1	2.8	3.3	4.0	2.7	5.0	6.1	4.1
2002.....	3.2	4.0	2.6	2.3	2.9	1.8	3.3	4.0	2.7	3.2	3.9	2.6	5.0	6.2	4.0
2001.....	3.2	3.8	2.6	2.0	2.4	1.7	3.2	3.9	2.6	3.1	3.8	2.5	5.0	6.0	4.3
2000.....	3.2	3.9	2.6	2.1	2.7	1.7	3.2	3.9	2.6	3.1	3.8	2.5	5.2	6.6	4.1
1999.....	3.0	3.7	2.4	2.0	2.7	1.5	3.0	3.7	2.4	2.8	3.5	2.3	4.8	6.1	3.8
Cerebrovascular diseases (I60-I69)															
2004.....	50.0	50.4	48.9	38.2	41.5	35.4	50.5	50.7	49.5	48.3	48.2	47.7	71.1	76.3	66.6
2003.....	53.5	54.1	52.3	40.5	43.0	38.1	54.1	54.5	52.9	51.7	51.9	50.8	75.5	81.0	70.9
2002.....	56.2	56.5	55.2	41.3	44.3	38.6	56.8	57.0	55.9	54.6	54.4	53.9	77.5	83.0	72.8
2001.....	57.9	59.0	56.4	44.9	48.9	41.6	58.4	59.2	56.9	56.0	56.5	54.8	79.9	86.5	74.8
2000.....	60.9	62.4	59.1	46.4	50.5	43.0	61.3	62.7	59.6	59.0	59.9	57.6	82.9	90.8	77.2
1999.....	61.6	63.2	59.8	46.6	52.6	42.2	62.1	63.5	60.4	59.8	60.8	58.4	82.8	90.6	77.1
Subarachnoid hemorrhage (I60)															
2004.....	2.0	1.6	2.4	2.4	2.1	2.7	2.0	1.5	2.4	1.8	1.4	2.2	2.8	2.1	3.3
2003.....	2.0	1.6	2.4	2.3	1.8	2.6	2.0	1.6	2.4	1.9	1.4	2.2	2.9	2.3	3.4
2002.....	2.1	1.6	2.6	2.3	1.8	2.6	2.1	1.6	2.6	2.0	1.5	2.3	3.0	2.2	3.6
2001.....	2.2	1.7	2.5	2.5	1.9	2.9	2.1	1.6	2.5	1.9	1.5	2.3	3.2	2.4	3.8
2000.....	2.4	1.8	2.8	2.6	2.0	3.1	2.3	1.8	2.8	2.2	1.7	2.5	3.4	2.3	4.3
1999.....	2.4	1.8	2.8	2.8	2.1	3.4	2.3	1.7	2.8	2.1	1.7	2.5	3.3	2.4	4.0
Intracerebral and other intracranial hemorrhage (I61-I62)															
2004.....	9.2	10.3	8.4	8.4	9.9	7.2	9.2	10.3	8.4	8.6	9.5	7.8	13.8	16.1	12.0
2003.....	9.3	10.4	8.4	7.9	9.0	6.9	9.3	10.4	8.5	8.5	9.5	7.8	14.4	17.0	12.4
2002.....	9.4	10.4	8.5	8.3	9.4	7.3	9.4	10.4	8.6	8.7	9.6	8.0	13.4	15.5	11.7
2001.....	9.4	10.5	8.6	8.6	10.0	7.5	9.4	10.5	8.6	8.7	9.6	7.9	14.0	16.1	12.3
2000.....	9.9	11.1	9.0	8.9	10.4	7.6	9.9	11.0	9.1	9.2	10.1	8.5	14.5	17.2	12.4
1999.....	9.3	10.3	8.5	9.0	10.8	7.4	9.3	10.2	8.6	8.6	9.4	8.0	13.5	15.6	11.8
Cerebral infarction (I63)															
2004.....	3.6	3.8	3.5	2.4	2.4	2.3	3.7	3.8	3.5	3.6	3.7	3.5	5.2	5.9	4.6
2003.....	4.1	4.2	3.9	2.5	2.4	2.4	4.2	4.3	4.0	4.1	4.2	3.9	5.5	6.0	5.1
2002.....	4.3	4.4	4.1	2.7	2.9	2.6	4.3	4.5	4.2	4.2	4.3	4.1	5.6	6.2	5.1
2001.....	4.5	4.6	4.3	3.3	3.4	3.1	4.5	4.7	4.3	4.4	4.5	4.2	5.9	6.9	5.3
2000.....	5.0	5.3	4.7	3.4	4.1	3.0	5.0	5.3	4.8	4.9	5.1	4.7	6.7	7.6	6.0
1999.....	5.4	5.7	5.2	3.7	4.0	3.5	5.5	5.8	5.2	5.4	5.6	5.1	7.2	8.4	6.4

Worktable Orig13. Age-adjusted death rates for 358 selected causes, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), and year	All origins 1/			Hispanic			Non-Hispanic 2/			Non-Hispanic white			Non-Hispanic black		
	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female
Stroke, not specified as hemorrhage or infarction (I64)															
2004.....	25.4	24.9	25.3	18.0	19.6	16.6	25.8	25.2	25.7	24.9	24.1	24.9	36.4	37.7	34.8
2003.....	28.0	27.7	27.6	20.3	21.8	19.0	28.3	27.9	28.0	27.4	26.8	27.2	39.3	41.3	37.3
2002.....	30.0	29.6	29.8	20.5	21.9	19.3	30.4	29.9	30.2	29.4	28.6	29.4	41.9	44.9	39.3
2001.....	31.4	31.4	30.9	22.5	25.0	20.8	31.7	31.6	31.3	30.7	30.3	30.4	43.2	47.0	40.3
2000.....	32.8	33.0	32.2	23.6	25.6	21.9	33.2	33.3	32.5	32.2	32.0	31.7	44.7	49.2	41.6
1999.....	33.6	34.0	32.8	23.3	26.4	21.1	33.9	34.2	33.1	32.9	33.0	32.3	45.1	49.0	42.3
Other cerebrovascular diseases and their sequelae (I67,I69)															
2004.....	9.6	9.7	9.4	7.0	7.4	6.6	9.7	9.9	9.5	9.5	9.5	9.3	13.0	14.5	11.9
2003.....	10.1	10.2	9.9	7.5	7.9	7.2	10.2	10.3	10.0	9.9	9.9	9.8	13.5	14.4	12.7
2002.....	10.4	10.6	10.2	7.5	8.3	6.9	10.6	10.7	10.3	10.3	10.4	10.1	13.7	14.2	13.1
2001.....	10.4	10.7	10.1	7.9	8.6	7.3	10.5	10.8	10.2	10.2	10.5	9.9	13.7	14.2	13.1
2000.....	10.8	11.3	10.3	7.9	8.4	7.4	10.8	11.4	10.4	10.6	11.1	10.1	13.6	14.5	12.9
1999.....	10.9	11.4	10.5	7.7	9.3	6.8	11.0	11.5	10.6	10.8	11.1	10.4	13.7	15.2	12.6
Diseases of arteries, arterioles and capillaries (I70-I78)															
2004.....	11.9	14.3	10.1	6.9	8.8	5.5	12.1	14.6	10.3	12.2	14.7	10.2	13.4	15.3	11.9
2003.....	12.8	15.7	10.7	7.9	10.4	6.1	13.1	16.0	10.9	13.1	16.1	10.9	14.2	16.8	12.3
2002.....	13.4	16.4	11.2	8.3	9.9	7.1	13.6	16.8	11.4	13.7	17.0	11.3	14.6	17.2	12.7
2001.....	14.0	17.1	11.7	8.5	10.2	7.2	14.2	17.4	11.9	14.3	17.5	11.9	15.3	17.6	13.6
2000.....	14.7	18.4	12.0	9.4	11.5	7.9	14.9	18.7	12.2	14.9	18.9	12.1	15.9	18.6	14.1
1999.....	15.0	18.5	12.5	9.4	12.3	7.5	15.2	18.8	12.7	15.3	19.0	12.6	15.6	18.4	13.8
Atherosclerosis (I70)															
2004.....	3.9	4.1	3.7	2.4	2.9	2.1	4.0	4.1	3.8	4.1	4.3	3.9	3.5	3.5	3.4
2003.....	4.4	4.5	4.3	2.6	3.2	2.3	4.5	4.5	4.3	4.6	4.6	4.4	4.3	4.8	3.9
2002.....	4.7	4.9	4.6	3.1	3.3	3.0	4.8	5.0	4.6	4.9	5.1	4.7	4.5	5.0	4.1
2001.....	5.0	5.1	4.8	3.5	3.7	3.3	5.0	5.1	4.9	5.2	5.3	5.0	4.3	4.7	4.1
2000.....	5.2	5.5	5.0	3.7	3.8	3.5	5.3	5.6	5.0	5.4	5.7	5.1	4.9	5.5	4.5
1999.....	5.5	5.7	5.3	3.8	4.4	3.4	5.6	5.8	5.4	5.7	5.8	5.5	5.0	5.9	4.5
Aortic aneurysm and dissection (I71)															
2004.....	4.6	6.7	3.1	2.4	3.5	1.5	4.8	6.9	3.3	4.8	7.0	3.3	4.5	6.2	3.4
2003.....	5.1	7.5	3.3	3.0	4.6	1.9	5.2	7.7	3.4	5.3	7.9	3.4	4.7	6.2	3.5
2002.....	5.2	7.7	3.3	2.5	3.8	1.6	5.3	7.9	3.4	5.4	8.1	3.5	4.7	6.1	3.6
2001.....	5.4	8.0	3.6	2.5	4.1	1.5	5.5	8.2	3.7	5.6	8.4	3.7	5.0	6.5	4.0
2000.....	5.7	8.7	3.7	3.2	4.8	2.1	5.8	8.9	3.7	5.9	9.2	3.7	5.3	6.9	4.2
1999.....	5.8	8.8	3.7	2.8	4.5	1.7	5.9	9.0	3.8	6.1	9.3	3.8	5.0	6.5	3.9
Other diseases of arteries, arterioles and capillaries (I72-I78)															
2004.....	3.3	3.5	3.1	2.2	2.4	2.0	3.4	3.6	3.2	3.2	3.5	3.0	5.3	5.7	5.1
2003.....	3.3	3.7	3.1	2.2	2.6	2.0	3.4	3.7	3.2	3.3	3.6	3.0	5.3	5.8	4.9
2002.....	3.5	3.8	3.3	2.7	2.9	2.6	3.5	3.9	3.3	3.4	3.7	3.2	5.5	6.1	5.1
2001.....	3.6	3.9	3.3	2.4	2.4	2.4	3.6	4.0	3.3	3.4	3.8	3.2	5.9	6.5	5.6
2000.....	3.7	4.2	3.4	2.6	2.9	2.4	3.7	4.2	3.4	3.6	4.1	3.2	5.8	6.3	5.4
1999.....	3.7	4.0	3.4	2.8	3.4	2.4	3.7	4.0	3.4	3.6	3.9	3.3	5.7	6.0	5.4
Other disorders of circulatory system (I80-I99)															
2004.....	1.5	1.6	1.5	1.0	0.9	1.0	1.6	1.7	1.5	1.5	1.6	1.4	2.8	2.9	2.6
2003.....	1.6	1.6	1.6	1.1	1.2	1.1	1.6	1.6	1.6	1.5	1.5	1.5	3.0	3.2	2.8
2002.....	1.6	1.6	1.6	1.1	1.0	1.2	1.7	1.6	1.6	1.6	1.5	1.6	2.9	3.2	2.7
2001.....	1.6	1.7	1.6	1.4	1.5	1.3	1.7	1.7	1.6	1.6	1.6	1.5	2.8	3.1	2.5
2000.....	1.7	1.7	1.6	1.4	1.6	1.3	1.7	1.7	1.6	1.6	1.6	1.5	2.9	3.1	2.8
1999.....	1.5	1.5	1.4	1.2	1.3	1.2	1.5	1.5	1.4	1.4	1.4	1.3	2.5	2.7	2.2

Worktable Orig13. Age-adjusted death rates for 358 selected causes, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), and year	All origins 1/			Hispanic			Non-Hispanic 2/			Non-Hispanic white			Non-Hispanic black		
	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)															
2004.....	1.2	1.2	1.2	0.7	0.5	0.8	1.2	1.2	1.2	1.1	1.1	1.1	2.2	2.3	2.2
2003.....	1.2	1.1	1.3	0.8	0.8	0.9	1.3	1.2	1.3	1.2	1.1	1.2	2.4	2.5	2.3
2002.....	1.2	1.1	1.3	0.8	0.5	1.0	1.3	1.2	1.3	1.2	1.1	1.3	2.3	2.4	2.3
2001.....	1.2	1.2	1.3	1.1	1.0	1.1	1.3	1.2	1.3	1.2	1.1	1.2	2.2	2.3	2.1
2000.....	1.2	1.2	1.2	1.0	1.1	1.0	1.2	1.2	1.3	1.2	1.1	1.2	2.2	2.2	2.3
1999.....	1.0	0.9	1.0	0.8	0.9	0.8	1.0	0.9	1.0	0.9	0.8	0.9	1.7	1.7	1.7
All other and unspecified disorders of circulatory system (I83-I99)															
2004.....	0.4	0.5	0.3	0.3	0.4	0.2	0.4	0.5	0.3	0.4	0.4	0.3	0.5	0.7	0.5
2003.....	0.4	0.4	0.3	0.3	0.4	0.2	0.4	0.4	0.3	0.4	0.4	0.3	0.6	0.8	0.4
2002.....	0.4	0.4	0.3	0.3	0.5	0.2	0.4	0.4	0.3	0.4	0.4	0.3	0.6	0.9	0.4
2001.....	0.4	0.5	0.3	0.3	0.5	*	0.4	0.5	0.3	0.4	0.5	0.3	0.5	0.8	0.4
2000.....	0.4	0.5	0.3	0.4	0.5	0.3	0.4	0.5	0.3	0.4	0.5	0.3	0.7	0.9	0.5
1999.....	0.5	0.6	0.4	0.4	0.4	0.4	0.5	0.6	0.4	0.5	0.6	0.4	0.7	1.0	0.5
X. Diseases of the respiratory system (J00-J98)															
2004.....	76.1	93.0	65.5	47.2	57.2	40.4	77.8	95.0	67.0	80.0	96.5	69.7	66.4	91.1	51.9
2003.....	80.5	98.0	69.5	49.9	62.0	41.8	82.2	99.9	71.1	84.4	101.1	73.9	69.5	96.4	53.9
2002.....	81.4	100.4	69.7	51.5	64.7	42.7	83.0	102.2	71.2	85.0	103.6	73.8	71.4	98.3	56.0
2001.....	81.6	101.2	69.8	54.1	68.5	45.0	83.0	102.8	71.1	84.7	103.9	73.3	72.6	100.5	56.9
2000.....	83.7	105.7	70.9	54.5	70.1	44.4	85.1	107.3	72.1	86.8	108.3	74.3	74.4	104.0	57.6
1999.....	84.3	107.7	70.6	54.3	70.7	44.2	85.6	109.3	71.8	87.2	110.2	73.8	76.3	107.9	58.5
Acute upper respiratory infections (J00-J06)															
2004.....	0.0	0.0	0.0	*	*	*	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2003.....	0.1	0.1	0.0	*	*	*	0.1	0.1	0.0	0.1	0.1	0.0	0.1	*	*
2002.....	0.0	0.0	0.1	*	*	*	0.0	0.0	0.1	0.0	0.0	0.1	*	*	*
2001.....	0.0	0.0	0.1	*	*	*	0.0	0.0	0.0	0.0	0.0	0.1	0.1	*	*
2000.....	0.0	0.0	0.1	*	*	*	0.1	0.0	0.1	0.1	0.1	0.1	*	*	*
1999.....	0.1	0.1	0.1	*	*	*	0.1	0.1	0.1	0.1	0.0	0.1	0.1	*	*
Acute pharyngitis and tonsillitis (J02-J03)															
2004.....	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2003.....	0.0	*	*	*	*	*	0.0	*	*	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	0.0	*	*	0.0	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Acute laryngitis and tracheitis (J04)															
2004.....	0.0	*	*	*	*	*	0.0	*	*	*	*	*	*	*	*
2003.....	0.0	*	*	*	*	*	0.0	*	*	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	0.0	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	0.0	*	*	*	*	*	*	*	*
Other and unspecified acute upper respiratory infections (J00-J01,J05-J06)															
2004.....	0.0	0.0	0.0	*	*	*	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2003.....	0.0	0.1	0.0	*	*	*	0.0	0.1	0.0	0.0	0.1	0.0	*	*	*
2002.....	0.0	0.0	0.0	*	*	*	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2001.....	0.0	0.0	0.0	*	*	*	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2000.....	0.0	0.0	0.0	*	*	*	0.0	0.0	0.0	0.0	0.0	0.1	*	*	*
1999.....	0.0	0.1	0.0	*	*	*	0.0	0.1	0.1	0.0	0.0	0.0	*	*	*

Worktable Orig13. Age-adjusted death rates for 358 selected causes, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), and year	All origins 1/			Hispanic			Non-Hispanic 2/			Non-Hispanic white			Non-Hispanic black		
	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female
Acute bronchitis and bronchiolitis (J20-J21)															
2004.....	0.1	0.1	0.1	0.1	*	*	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2003.....	0.1	0.1	0.1	0.1	0.1	*	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*
2002.....	0.1	0.1	0.1	*	*	*	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*
2001.....	0.1	0.1	0.1	*	*	*	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2000.....	0.1	0.1	0.1	0.1	*	*	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	0.1
1999.....	0.1	0.1	0.1	0.1	*	*	0.1	0.1	0.1	0.1	0.1	0.1	0.2	0.2	0.2
Unspecified acute lower respiratory infection (J22)															
2004.....	0.0	0.0	0.0	*	*	*	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2003.....	0.0	0.0	0.0	*	*	*	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2002.....	0.0	0.0	0.0	*	*	*	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2001.....	0.0	0.0	0.0	*	*	*	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2000.....	0.0	0.0	0.0	*	*	*	0.0	0.0	0.0	0.1	0.0	0.0	*	*	*
1999.....	0.1	0.0	0.1	*	*	*	0.1	0.0	0.1	0.1	0.0	0.1	*	*	*
Other diseases of upper respiratory tract (J30-J39)															
2004.....	0.1	0.2	0.1	0.1	*	*	0.1	0.2	0.1	0.1	0.2	0.1	0.2	0.3	0.2
2003.....	0.1	0.2	0.1	0.1	*	*	0.1	0.2	0.1	0.1	0.2	0.1	0.2	0.2	0.2
2002.....	0.1	0.1	0.1	*	*	*	0.1	0.1	0.1	0.1	0.1	0.1	0.2	0.3	0.1
2001.....	0.2	0.2	0.1	0.2	*	*	0.2	0.2	0.1	0.1	0.2	0.1	0.3	0.4	0.2
2000.....	0.2	0.2	0.1	0.1	*	*	0.2	0.2	0.1	0.1	0.2	0.1	0.2	0.3	0.2
1999.....	0.2	0.2	0.1	0.1	*	*	0.2	0.2	0.1	0.1	0.2	0.1	0.3	0.3	0.2
Chronic lower respiratory diseases (J40-J47)															
2004.....	41.1	49.5	36.0	18.4	23.8	14.9	42.6	51.0	37.3	44.9	52.8	40.0	28.7	41.6	21.2
2003.....	43.3	52.3	37.8	20.2	27.1	15.8	44.7	53.7	39.1	47.0	55.4	41.8	30.5	45.1	22.3
2002.....	43.5	53.5	37.4	20.6	27.2	16.2	44.8	54.9	38.6	46.9	56.5	41.2	31.5	46.8	22.9
2001.....	43.7	54.0	37.6	20.7	27.6	16.5	44.9	55.4	38.7	47.0	56.9	41.1	31.3	47.0	22.6
2000.....	44.2	55.8	37.4	21.1	28.8	16.3	45.3	57.1	38.5	47.2	58.5	40.7	32.0	47.9	23.1
1999.....	45.4	58.7	37.7	23.4	33.0	17.7	46.5	59.9	38.6	48.3	61.3	40.8	33.9	52.0	23.7
Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)															
2004.....	40.8	49.2	35.6	18.2	23.6	14.7	42.2	50.8	36.9	44.5	52.5	39.6	28.6	41.5	21.1
2003.....	43.0	52.1	37.4	20.1	27.0	15.7	44.4	53.5	38.7	46.6	55.1	41.4	30.5	45.0	22.2
2002.....	43.1	53.2	37.0	20.3	27.0	16.0	44.4	54.6	38.2	46.5	56.2	40.7	31.4	46.7	22.7
2001.....	43.3	53.8	37.2	20.5	27.4	16.2	44.6	55.1	38.3	46.6	56.7	40.7	31.1	46.8	22.5
2000.....	43.8	55.5	37.0	20.8	28.5	15.9	45.0	56.8	38.1	46.8	58.3	40.3	31.8	47.8	22.9
1999.....	45.1	58.4	37.3	23.1	32.8	17.3	46.1	59.6	38.3	47.9	61.0	40.4	33.7	51.8	23.5
Bronchitis, not specified as acute or chronic (J40)															
2004.....	0.1	0.2	0.1	0.1	*	*	0.1	0.1	0.1	0.1	0.2	0.1	0.1	*	*
2003.....	0.1	0.1	0.1	0.2	*	*	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	0.2
2002.....	0.2	0.2	0.2	0.2	*	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.1	*	0.1
2001.....	0.2	0.2	0.2	0.1	*	*	0.2	0.2	0.2	0.2	0.2	0.2	0.1	0.3	*
2000.....	0.2	0.2	0.2	*	*	*	0.2	0.3	0.2	0.2	0.3	0.2	0.2	0.2	0.2
1999.....	0.2	0.2	0.2	0.2	*	*	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.2
Chronic bronchitis (J41-J42)															
2004.....	0.1	0.1	0.1	*	*	*	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*
2003.....	0.1	0.2	0.1	*	*	*	0.1	0.2	0.1	0.1	0.2	0.1	0.1	*	0.1
2002.....	0.2	0.2	0.1	0.1	*	*	0.2	0.2	0.1	0.2	0.2	0.1	0.1	0.2	*
2001.....	0.1	0.2	0.1	*	*	*	0.1	0.2	0.1	0.2	0.2	0.1	0.1	*	*
2000.....	0.2	0.2	0.2	0.3	*	*	0.2	0.2	0.2	0.2	0.2	0.2	0.1	*	*
1999.....	0.2	0.2	0.2	0.2	*	0.3	0.2	0.2	0.2	0.2	0.2	0.2	0.1	*	*

Worktable Orig13. Age-adjusted death rates for 358 selected causes, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), and year	All origins 1/			Hispanic			Non-Hispanic 2/			Non-Hispanic white			Non-Hispanic black		
	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female
Emphysema (J43)															
2004.....	4.6	5.8	3.8	1.6	2.2	1.2	4.8	6.1	4.0	5.1	6.3	4.3	2.8	4.7	1.7
2003.....	5.1	6.3	4.3	2.0	2.9	1.4	5.3	6.6	4.5	5.6	6.8	4.9	3.3	5.4	2.1
2002.....	5.4	7.0	4.4	2.0	2.8	1.5	5.6	7.2	4.6	6.0	7.5	5.0	3.5	5.9	2.1
2001.....	5.8	7.4	4.7	2.1	3.3	1.4	6.0	7.7	4.9	6.4	8.0	5.3	3.4	5.4	2.2
2000.....	6.0	7.8	4.9	2.1	3.2	1.3	6.2	8.1	5.1	6.6	8.3	5.5	3.8	6.2	2.4
1999.....	6.5	8.6	5.2	2.4	3.8	1.5	6.7	8.8	5.3	7.1	9.2	5.8	4.0	6.8	2.3
Other chronic obstructive pulmonary disease (J44)															
2004.....	34.7	42.0	30.1	15.2	20.3	11.9	35.8	43.3	31.3	38.1	45.1	33.8	22.4	33.4	16.1
2003.....	36.2	44.2	31.3	16.5	22.6	12.7	37.3	45.5	32.4	39.6	47.2	35.0	23.6	36.4	16.5
2002.....	35.9	44.7	30.7	16.7	22.9	12.6	37.0	45.8	31.7	39.0	47.4	34.0	24.2	37.2	17.0
2001.....	35.7	44.8	30.3	16.7	22.9	12.9	36.7	45.9	31.3	38.7	47.4	33.6	23.9	37.8	16.2
2000.....	35.8	45.9	29.9	16.8	23.4	12.6	36.7	46.9	30.7	38.6	48.4	32.9	23.8	37.7	16.0
1999.....	36.5	47.9	29.8	18.5	27.4	13.2	37.3	48.8	30.5	39.0	50.3	32.6	25.3	40.9	16.7
Asthma (J45-J46)															
2004.....	1.3	1.0	1.4	1.1	0.9	1.4	1.3	1.1	1.5	1.0	0.8	1.2	3.1	3.1	3.1
2003.....	1.4	1.1	1.6	1.3	1.1	1.4	1.4	1.2	1.6	1.1	0.9	1.3	3.2	3.0	3.4
2002.....	1.5	1.2	1.7	1.3	1.0	1.5	1.5	1.2	1.7	1.2	0.9	1.4	3.5	3.4	3.5
2001.....	1.5	1.2	1.7	1.4	1.0	1.7	1.5	1.2	1.8	1.2	0.9	1.4	3.6	3.3	3.9
2000.....	1.6	1.3	1.8	1.6	1.5	1.7	1.6	1.3	1.8	1.3	1.0	1.5	3.9	3.5	4.2
1999.....	1.7	1.3	2.0	1.8	1.3	2.2	1.7	1.3	1.9	1.4	1.0	1.6	4.0	3.6	4.2
Bronchiectasis (J47)															
2004.....	0.3	0.2	0.4	0.2	*	0.2	0.3	0.2	0.4	0.3	0.2	0.4	0.1	*	0.1
2003.....	0.3	0.2	0.4	0.2	*	*	0.3	0.2	0.4	0.3	0.2	0.4	0.1	*	*
2002.....	0.3	0.2	0.4	0.2	*	0.3	0.3	0.2	0.4	0.3	0.2	0.4	0.1	*	*
2001.....	0.3	0.2	0.4	0.2	*	0.2	0.3	0.2	0.4	0.4	0.2	0.4	0.1	*	*
2000.....	0.3	0.2	0.4	0.3	*	0.4	0.3	0.2	0.4	0.4	0.2	0.4	0.1	*	*
1999.....	0.3	0.3	0.4	0.3	*	0.3	0.3	0.3	0.4	0.4	0.3	0.4	0.2	*	0.2
Lung diseases due to external agents (J60-J70)															
2004.....	6.0	8.6	4.3	3.5	4.5	2.9	6.1	8.9	4.4	6.1	8.9	4.4	6.4	9.5	4.7
2003.....	6.3	9.0	4.7	3.6	4.6	3.0	6.4	9.2	4.7	6.4	9.3	4.7	7.0	9.9	5.5
2002.....	6.5	9.4	4.8	4.2	5.8	3.2	6.6	9.6	4.9	6.6	9.7	4.9	6.9	10.0	5.3
2001.....	6.6	9.5	4.9	4.6	6.4	3.5	6.6	9.7	4.9	6.7	9.8	4.8	7.3	10.1	5.8
2000.....	6.5	9.7	4.7	4.3	5.9	3.3	6.6	9.9	4.7	6.6	9.9	4.8	7.1	10.3	5.3
1999.....	6.1	9.2	4.4	3.8	4.9	3.2	6.1	9.3	4.4	6.1	9.3	4.4	7.2	10.7	5.2
Pneumoconioses and chemical effects (J60-J66,J68)															
2004.....	0.3	0.9	0.0	0.1	0.4	*	0.4	0.9	0.0	0.4	1.0	0.0	0.2	0.7	*
2003.....	0.4	1.0	0.0	0.1	*	*	0.4	1.0	0.0	0.4	1.0	0.0	0.2	0.5	*
2002.....	0.4	1.0	0.0	0.2	0.5	*	0.4	1.0	0.0	0.4	1.0	0.0	0.2	0.6	*
2001.....	0.4	1.0	0.0	0.2	0.4	*	0.4	1.0	0.0	0.4	1.1	0.0	0.2	0.7	*
2000.....	0.4	1.1	0.0	0.2	0.5	*	0.4	1.1	0.0	0.4	1.2	0.0	0.3	0.7	*
1999.....	0.4	1.0	0.0	*	*	*	0.4	1.1	0.0	0.4	1.1	0.0	0.3	0.8	*
Pneumonitis due to solids and liquids (J69)															
2004.....	5.6	7.7	4.3	3.3	4.1	2.8	5.7	7.9	4.4	5.7	7.9	4.4	6.2	8.8	4.7
2003.....	5.9	8.0	4.6	3.5	4.3	2.9	6.0	8.2	4.7	6.0	8.2	4.7	6.8	9.3	5.4
2002.....	6.1	8.4	4.7	3.9	5.3	3.1	6.2	8.6	4.8	6.2	8.7	4.8	6.7	9.4	5.2
2001.....	6.1	8.5	4.8	4.4	6.0	3.5	6.2	8.6	4.9	6.2	8.7	4.8	7.1	9.5	5.8
2000.....	6.1	8.6	4.6	4.1	5.5	3.3	6.1	8.7	4.7	6.1	8.7	4.7	6.8	9.5	5.3
1999.....	5.6	8.1	4.3	3.7	4.7	3.1	5.7	8.2	4.3	5.7	8.1	4.3	6.8	10.0	5.2

Worktable Orig13. Age-adjusted death rates for 358 selected causes, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), and year	All origins 1/			Hispanic			Non-Hispanic 2/			Non-Hispanic white			Non-Hispanic black		
	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female
Pneumonitis due to food and vomit (J69.0)															
2004.....	5.6	7.7	4.3	3.3	4.1	2.8	5.7	7.9	4.4	5.7	7.9	4.4	6.2	8.8	4.7
2003.....	5.9	8.0	4.6	3.5	4.3	2.9	6.0	8.2	4.7	6.0	8.2	4.7	6.8	9.3	5.4
2002.....	6.1	8.4	4.7	3.9	5.3	3.1	6.2	8.6	4.8	6.2	8.7	4.8	6.7	9.4	5.2
2001.....	6.1	8.5	4.8	4.4	6.0	3.5	6.2	8.6	4.9	6.2	8.6	4.8	7.1	9.5	5.8
2000.....	6.1	8.6	4.6	4.1	5.5	3.3	6.1	8.7	4.7	6.1	8.7	4.7	6.8	9.5	5.3
1999.....	5.6	8.1	4.3	3.7	4.7	3.1	5.7	8.2	4.3	5.7	8.1	4.3	6.8	10.0	5.1
Other pneumonitis due to solids and liquids (J69.1-J69.8)															
2004.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Other lung diseases due to external agents (J67,J70)															
2004.....	0.0	0.0	0.0	*	*	*	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2003.....	0.0	0.0	0.0	*	*	*	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2002.....	0.0	0.0	0.0	*	*	*	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2001.....	0.0	0.0	0.0	*	*	*	0.0	0.0	0.0	0.0	0.1	0.0	*	*	*
2000.....	0.0	0.0	0.0	*	*	*	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
1999.....	0.0	0.0	0.0	*	*	*	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
Suppurative and necrotic conditions of lower respiratory tract (J85-J86)															
2004.....	0.2	0.3	0.2	0.1	*	*	0.2	0.3	0.2	0.2	0.3	0.2	0.3	0.4	0.2
2003.....	0.2	0.3	0.2	0.2	0.2	*	0.2	0.3	0.2	0.2	0.3	0.2	0.4	0.6	0.2
2002.....	0.2	0.4	0.2	0.2	0.4	*	0.2	0.4	0.2	0.2	0.3	0.2	0.4	0.7	0.3
2001.....	0.3	0.4	0.2	0.2	0.3	*	0.3	0.4	0.2	0.3	0.3	0.2	0.4	0.6	0.2
2000.....	0.2	0.4	0.2	0.2	*	*	0.3	0.4	0.2	0.2	0.4	0.2	0.5	0.7	0.3
1999.....	0.3	0.4	0.2	0.2	0.3	*	0.3	0.4	0.2	0.3	0.4	0.2	0.5	0.8	0.3
Pleural effusion and plaque (J90,J92)															
2004.....	0.3	0.4	0.3	0.2	*	0.2	0.3	0.4	0.3	0.3	0.4	0.3	0.3	0.4	0.3
2003.....	0.3	0.4	0.3	0.2	*	*	0.3	0.4	0.3	0.3	0.4	0.3	0.4	0.4	0.3
2002.....	0.3	0.4	0.3	0.2	*	*	0.3	0.4	0.3	0.4	0.4	0.3	0.4	0.4	0.3
2001.....	0.3	0.4	0.3	0.2	*	*	0.4	0.4	0.3	0.4	0.4	0.3	0.5	0.5	0.4
2000.....	0.3	0.4	0.3	0.3	0.4	*	0.3	0.4	0.3	0.3	0.4	0.3	0.4	0.5	0.3
1999.....	0.3	0.4	0.3	0.2	*	*	0.3	0.4	0.3	0.3	0.4	0.3	0.3	0.4	0.3
All other diseases of respiratory system (J80-J84,J93-J98)															
2004.....	8.4	10.1	7.2	7.6	8.5	6.9	8.5	10.2	7.3	8.6	10.4	7.3	7.8	9.4	6.8
2003.....	7.9	9.5	6.9	7.1	8.1	6.3	8.0	9.5	6.9	8.0	9.7	7.0	7.2	8.8	6.2
2002.....	7.9	9.4	6.9	6.9	7.7	6.2	8.0	9.5	6.9	8.0	9.6	7.0	7.5	8.7	6.7
2001.....	8.4	10.0	7.3	7.6	9.0	6.7	8.5	10.0	7.4	8.5	10.1	7.3	8.3	9.3	7.8
2000.....	8.4	10.2	7.2	7.7	8.7	6.9	8.4	10.2	7.3	8.5	10.3	7.3	8.2	9.8	7.3
1999.....	8.3	10.0	7.2	7.6	9.0	6.7	8.3	10.1	7.2	8.3	10.1	7.1	8.3	9.6	7.6
XI. Diseases of the digestive system (K00-K92)															
2004.....	28.8	33.4	24.5	29.3	36.2	22.7	28.6	32.9	24.5	28.4	32.5	24.5	32.2	40.4	26.0
2003.....	29.8	34.8	25.1	31.7	39.7	24.2	29.5	34.2	25.2	29.3	33.8	25.1	33.3	41.9	26.7
2002.....	29.9	35.0	25.3	31.4	40.9	22.8	29.7	34.3	25.4	29.5	33.9	25.4	32.8	40.6	26.9
2001.....	30.3	35.5	25.7	33.0	42.6	24.5	29.9	34.7	25.7	29.6	34.0	25.6	35.1	44.7	27.8
2000.....	30.4	35.8	25.5	33.7	44.2	24.5	30.0	34.9	25.5	29.4	34.1	25.1	35.8	44.4	29.2
1999.....	30.3	36.0	25.3	34.5	44.7	25.4	29.9	35.2	25.3	29.4	34.3	24.9	35.7	44.6	29.1

Worktable Orig13. Age-adjusted death rates for 358 selected causes, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), and year	All origins 1/			Hispanic			Non-Hispanic 2/			Non-Hispanic white			Non-Hispanic black			
	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	
Diseases of oral cavity, salivary glands and jaws (K00-K14)																
2004.....	0.1	0.1	0.1	*	*	*	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*	*
2003.....	0.0	0.0	0.0	*	*	*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2002.....	0.0	0.1	0.0	*	*	*	0.0	0.1	0.0	0.0	0.1	0.0	*	*	*	
2001.....	0.0	0.0	0.0	*	*	*	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	
2000.....	0.0	0.0	0.0	*	*	*	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	
1999.....	0.0	0.0	0.0	*	*	*	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	
Diseases of other parts of digestive system (K20-K92)																
2004.....	28.7	33.3	24.4	29.2	36.2	22.6	28.5	32.8	24.5	28.3	32.4	24.4	32.2	40.3	25.9	
2003.....	29.8	34.8	25.1	31.6	39.6	24.1	29.4	34.2	25.1	29.3	33.7	25.1	33.3	41.9	26.7	
2002.....	29.9	35.0	25.3	31.4	40.8	22.8	29.6	34.2	25.4	29.5	33.9	25.3	32.8	40.6	26.9	
2001.....	30.2	35.4	25.6	33.0	42.5	24.5	29.9	34.7	25.6	29.5	33.9	25.5	35.0	44.6	27.8	
2000.....	30.3	35.7	25.5	33.7	44.2	24.5	29.9	34.9	25.5	29.4	34.1	25.1	35.7	44.4	29.2	
1999.....	30.3	35.9	25.3	34.4	44.7	25.4	29.9	35.1	25.2	29.3	34.3	24.8	35.7	44.5	29.0	
Diseases of esophagus (K20-K22)																
2004.....	0.6	0.7	0.6	0.3	0.5	0.2	0.6	0.7	0.6	0.7	0.7	0.6	0.6	0.8	0.4	
2003.....	0.7	0.8	0.6	0.5	0.7	0.3	0.7	0.8	0.6	0.7	0.9	0.6	0.6	0.7	0.5	
2002.....	0.7	0.9	0.6	0.5	0.7	0.3	0.7	0.9	0.6	0.8	0.9	0.7	0.7	0.9	0.5	
2001.....	0.7	0.9	0.7	0.4	0.5	0.4	0.8	0.9	0.7	0.8	0.9	0.7	0.6	0.9	0.5	
2000.....	0.8	1.0	0.7	0.5	0.8	0.3	0.8	1.0	0.7	0.8	1.0	0.7	0.7	1.0	0.5	
1999.....	0.8	1.0	0.7	0.7	0.9	0.6	0.8	1.0	0.7	0.8	1.0	0.7	0.7	0.9	0.5	
Peptic ulcer (K25-K28)																
2004.....	1.2	1.4	1.1	0.9	1.1	0.7	1.2	1.4	1.1	1.2	1.4	1.1	1.3	1.7	1.1	
2003.....	1.3	1.6	1.1	1.1	1.4	0.9	1.3	1.6	1.1	1.3	1.5	1.2	1.5	2.1	1.0	
2002.....	1.4	1.7	1.2	1.2	1.6	0.8	1.4	1.7	1.2	1.4	1.7	1.2	1.4	2.0	1.1	
2001.....	1.6	1.8	1.4	1.2	1.6	0.8	1.6	1.8	1.4	1.6	1.8	1.4	1.6	2.3	1.1	
2000.....	1.6	2.0	1.4	1.4	1.6	1.2	1.7	2.1	1.4	1.6	2.0	1.4	1.7	2.4	1.3	
1999.....	1.7	2.0	1.4	1.3	1.8	0.9	1.7	2.1	1.4	1.7	2.0	1.4	1.9	2.5	1.4	
Gastric ulcer (K25)																
2004.....	0.4	0.4	0.3	0.3	0.3	0.2	0.4	0.4	0.4	0.4	0.4	0.4	0.5	0.6	0.4	
2003.....	0.4	0.4	0.4	0.3	0.3	0.2	0.4	0.4	0.4	0.4	0.4	0.4	0.5	0.7	0.4	
2002.....	0.4	0.5	0.4	0.4	0.6	0.3	0.4	0.4	0.4	0.4	0.4	0.4	0.5	0.6	0.4	
2001.....	0.5	0.4	0.4	0.4	0.5	0.2	0.5	0.5	0.4	0.4	0.4	0.4	0.6	0.8	0.4	
2000.....	0.5	0.5	0.4	0.5	0.5	0.5	0.5	0.5	0.4	0.5	0.5	0.4	0.7	1.0	0.4	
1999.....	0.5	0.5	0.4	0.4	0.6	0.3	0.5	0.5	0.4	0.5	0.5	0.5	0.7	1.0	0.5	
Duodenal ulcer (K26)																
2004.....	0.4	0.5	0.3	0.3	0.5	*	0.4	0.5	0.3	0.4	0.5	0.4	0.4	0.5	0.2	
2003.....	0.4	0.6	0.4	0.3	0.3	0.3	0.5	0.6	0.3	0.5	0.6	0.4	0.5	0.7	0.3	
2002.....	0.5	0.6	0.4	0.3	0.4	0.2	0.5	0.6	0.4	0.5	0.6	0.4	0.4	0.6	0.3	
2001.....	0.5	0.7	0.4	0.3	0.5	*	0.6	0.7	0.5	0.6	0.7	0.5	0.5	0.6	0.3	
2000.....	0.5	0.7	0.4	0.3	0.5	*	0.6	0.7	0.4	0.6	0.7	0.4	0.5	0.6	0.4	
1999.....	0.5	0.7	0.4	0.3	0.4	*	0.6	0.7	0.4	0.5	0.7	0.4	0.5	0.6	0.4	
Peptic ulcer, site unspecified (K27)																
2004.....	0.4	0.5	0.4	0.3	0.3	0.3	0.4	0.5	0.4	0.4	0.4	0.4	0.4	0.5	0.4	
2003.....	0.5	0.6	0.4	0.5	0.6	0.3	0.5	0.6	0.4	0.5	0.5	0.4	0.4	0.7	0.3	
2002.....	0.5	0.6	0.4	0.4	0.6	0.3	0.5	0.6	0.4	0.5	0.6	0.4	0.5	0.7	0.4	
2001.....	0.6	0.7	0.5	0.5	0.6	0.4	0.6	0.7	0.5	0.6	0.6	0.5	0.6	0.8	0.4	
2000.....	0.6	0.8	0.5	0.5	0.5	0.5	0.6	0.8	0.5	0.6	0.8	0.5	0.6	0.7	0.5	
1999.....	0.6	0.8	0.6	0.5	0.8	0.4	0.6	0.8	0.6	0.6	0.8	0.6	0.6	0.8	0.5	

Worktable Orig13. Age-adjusted death rates for 358 selected causes, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Table with columns: Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), and year; All origins 1/; Hispanic; Non-Hispanic 2/; Non-Hispanic white; Non-Hispanic black. Rows include Gastrojejunal ulcer (K28), Gastritis and duodenitis (K29), Dyspepsia and other diseases of stomach and duodenum (K30-K31), Diseases of appendix (K35-K38), Hernia (K40-K46), Crohn's disease and ulcerative colitis (K50-K51), and Vascular disorders and obstruction of intestine without hernia (K55-K56).

Worktable Orig13. Age-adjusted death rates for 358 selected causes, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), and year	All origins 1/			Hispanic			Non-Hispanic 2/			Non-Hispanic white			Non-Hispanic black		
	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female
Diverticular disease of intestine (K57)															
2004.....	1.1	0.9	1.3	0.8	0.7	0.8	1.1	0.9	1.2	1.2	0.9	1.3	1.0	0.9	1.0
2003.....	1.2	1.0	1.3	1.0	0.7	1.1	1.2	1.0	1.3	1.2	1.0	1.4	1.0	0.8	1.1
2002.....	1.2	0.9	1.3	0.9	0.7	1.0	1.2	1.0	1.3	1.2	1.0	1.3	1.1	1.0	1.1
2001.....	1.2	1.0	1.3	1.0	0.8	1.0	1.2	1.0	1.4	1.3	1.0	1.4	1.1	0.9	1.2
2000.....	1.2	1.0	1.3	0.9	0.7	1.1	1.2	1.0	1.3	1.2	1.0	1.4	1.1	0.8	1.2
1999.....	1.3	1.0	1.4	1.2	1.1	1.2	1.3	1.0	1.4	1.3	1.0	1.4	1.1	0.9	1.2
Other diseases of intestines and peritoneum (K52,K58-K63)															
2004.....	1.7	1.5	1.7	1.1	1.1	1.0	1.7	1.5	1.8	1.7	1.5	1.9	1.7	1.9	1.5
2003.....	1.7	1.6	1.7	1.1	1.2	1.1	1.7	1.6	1.7	1.7	1.6	1.7	1.9	2.0	1.8
2002.....	1.7	1.6	1.7	1.0	1.0	0.9	1.7	1.6	1.8	1.7	1.6	1.8	1.7	1.9	1.5
2001.....	1.7	1.6	1.7	1.3	1.4	1.2	1.7	1.6	1.8	1.7	1.6	1.8	2.0	2.1	1.9
2000.....	1.6	1.5	1.7	1.1	1.2	1.1	1.7	1.5	1.7	1.6	1.5	1.7	1.8	1.9	1.7
1999.....	1.6	1.6	1.7	1.2	1.1	1.3	1.6	1.6	1.7	1.6	1.5	1.7	1.7	2.0	1.5
Diseases of peritoneum (K65-K66)															
2004.....	0.6	0.6	0.6	0.5	0.5	0.5	0.6	0.6	0.6	0.6	0.6	0.5	0.9	0.9	0.8
2003.....	0.6	0.7	0.6	0.6	0.6	0.7	0.6	0.7	0.6	0.6	0.6	0.6	1.0	1.0	1.0
2002.....	0.7	0.6	0.7	0.4	0.4	0.3	0.7	0.6	0.7	0.6	0.6	0.7	1.0	1.0	1.0
2001.....	0.7	0.7	0.6	0.5	0.5	0.5	0.7	0.7	0.6	0.6	0.6	0.6	1.1	1.2	1.0
2000.....	0.7	0.7	0.7	0.5	0.7	0.4	0.7	0.7	0.7	0.7	0.7	0.6	1.2	1.3	1.1
1999.....	0.7	0.8	0.6	0.6	0.7	0.6	0.7	0.8	0.6	0.7	0.7	0.6	1.1	1.2	1.0
Diseases of liver (K70-K76)															
2004.....	12.0	16.2	8.2	17.3	23.8	11.3	11.5	15.4	8.0	11.5	15.4	8.0	12.0	17.0	8.1
2003.....	12.4	16.8	8.4	18.7	25.9	12.1	11.8	16.0	8.2	11.9	16.0	8.2	12.4	17.8	8.3
2002.....	12.3	16.7	8.4	18.7	26.8	11.3	11.8	15.8	8.2	11.8	15.8	8.2	12.4	17.2	8.7
2001.....	12.3	16.7	8.3	19.2	28.0	11.3	11.7	15.8	8.1	11.6	15.5	8.1	13.2	19.1	8.6
2000.....	12.2	16.7	8.2	19.7	28.8	11.6	11.6	15.7	8.0	11.4	15.5	7.7	13.4	18.7	9.3
1999.....	12.3	17.0	8.1	19.6	28.6	11.6	11.7	16.1	7.9	11.4	15.7	7.6	14.1	19.7	9.6
Alcoholic liver disease (K70)															
2004.....	4.2	6.3	2.2	6.3	10.9	2.0	3.9	5.9	2.2	4.0	6.0	2.2	3.6	5.7	2.0
2003.....	4.2	6.4	2.1	6.7	11.6	2.2	3.9	5.9	2.1	4.0	6.0	2.1	3.9	6.1	2.1
2002.....	4.2	6.3	2.2	6.9	12.1	2.1	3.9	5.8	2.2	3.9	5.8	2.2	3.9	5.9	2.3
2001.....	4.3	6.6	2.2	7.6	13.6	2.3	4.0	6.0	2.2	3.9	5.9	2.2	4.5	7.1	2.4
2000.....	4.3	6.8	2.2	7.8	14.4	2.1	4.1	6.2	2.1	4.0	6.1	2.0	4.7	7.1	2.9
1999.....	4.3	6.9	2.1	7.6	13.5	2.4	4.1	6.3	2.1	4.0	6.2	2.0	5.2	8.2	2.8
Chronic hepatitis, not elsewhere classified (K73)															
2004.....	0.0	0.0	0.0	*	*	*	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2003.....	0.0	0.1	0.0	*	*	*	0.0	0.1	0.0	0.0	0.1	0.0	*	*	*
2002.....	0.2	0.1	0.2	0.4	*	0.6	0.2	0.1	0.2	0.2	0.1	0.2	0.2	*	0.2
2001.....	0.2	0.1	0.2	0.3	*	0.4	0.2	0.1	0.2	0.2	0.1	0.2	0.2	*	0.2
2000.....	0.2	0.1	0.2	0.3	*	0.5	0.1	0.1	0.2	0.1	0.1	0.2	0.2	*	0.2
1999.....	0.2	0.1	0.2	0.2	*	0.3	0.2	0.1	0.2	0.1	0.1	0.2	0.2	*	0.3
Fibrosis and cirrhosis of liver (K74)															
2004.....	4.8	6.1	3.6	7.6	8.7	6.5	4.6	5.8	3.4	4.6	5.9	3.5	4.4	6.2	3.0
2003.....	5.1	6.5	3.8	8.0	9.2	6.7	4.9	6.3	3.7	5.0	6.4	3.8	4.6	6.4	3.2
2002.....	5.0	6.4	3.8	8.1	10.1	6.3	4.8	6.1	3.7	4.9	6.2	3.7	4.5	6.1	3.2
2001.....	5.1	6.5	3.8	7.9	10.0	6.0	4.9	6.2	3.7	4.9	6.2	3.8	4.8	6.9	3.0
2000.....	5.1	6.5	3.8	8.4	10.2	6.7	4.8	6.2	3.6	4.8	6.3	3.6	4.7	6.5	3.3
1999.....	5.0	6.6	3.7	8.2	10.2	6.4	4.8	6.3	3.6	4.9	6.3	3.6	4.8	6.7	3.3

Worktable Orig13. Age-adjusted death rates for 358 selected causes, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories in the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), and year	All origins 1/			Hispanic			Non-Hispanic 2/			Non-Hispanic white			Non-Hispanic black		
	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female
Other diseases of liver (K71-K72,K75-K76)															
2004.....	3.0	3.7	2.4	3.4	4.1	2.7	3.0	3.6	2.4	2.8	3.5	2.2	4.0	5.0	3.1
2003.....	3.0	3.8	2.4	4.0	5.0	3.1	3.0	3.7	2.3	2.9	3.6	2.3	4.0	5.3	3.0
2002.....	2.9	3.8	2.1	3.3	4.4	2.3	2.9	3.7	2.1	2.8	3.6	2.0	3.8	5.0	3.0
2001.....	2.7	3.5	2.1	3.4	4.3	2.6	2.7	3.5	2.0	2.6	3.3	2.0	3.8	5.0	2.9
2000.....	2.6	3.3	2.0	3.2	4.1	2.4	2.6	3.2	2.0	2.4	3.0	1.9	3.9	5.0	2.9
1999.....	2.7	3.4	2.1	3.6	4.8	2.5	2.6	3.3	2.0	2.4	3.2	1.8	3.9	4.7	3.3
Cholelithiasis and other disorders of gallbladder (K80-K82)															
2004.....	1.0	1.2	0.9	1.2	1.3	1.2	1.0	1.1	0.9	1.0	1.2	0.9	1.3	1.2	1.2
2003.....	1.0	1.2	0.9	1.1	1.3	0.9	1.0	1.1	0.9	1.0	1.1	0.9	1.1	1.3	1.0
2002.....	1.0	1.2	0.9	1.2	1.3	1.1	1.0	1.2	0.9	1.0	1.2	0.9	1.1	1.1	1.1
2001.....	1.0	1.2	0.9	1.3	1.5	1.2	1.0	1.2	0.9	1.0	1.2	0.9	1.3	1.4	1.2
2000.....	1.0	1.2	0.9	1.3	1.7	1.1	1.0	1.2	0.9	1.0	1.1	0.9	1.2	1.0	1.2
1999.....	1.0	1.2	0.9	1.5	1.6	1.3	1.0	1.2	0.9	1.0	1.2	0.9	1.1	1.1	1.1
Disorders of biliary tract and pancreas (K83-K86)															
2004.....	1.6	2.0	1.3	1.4	1.7	1.1	1.6	2.0	1.3	1.6	1.9	1.3	2.4	3.1	1.9
2003.....	1.7	2.0	1.4	1.4	1.9	1.1	1.7	2.0	1.4	1.6	1.9	1.3	2.5	3.3	1.9
2002.....	1.7	2.1	1.3	1.5	1.9	1.2	1.7	2.1	1.3	1.7	2.0	1.3	2.3	3.1	1.7
2001.....	1.8	2.1	1.5	1.5	1.5	1.5	1.8	2.1	1.5	1.7	2.0	1.4	2.6	3.2	2.0
2000.....	1.7	2.1	1.4	1.7	2.2	1.3	1.7	2.1	1.4	1.6	1.9	1.3	2.5	3.4	1.9
1999.....	1.6	2.0	1.3	1.7	2.1	1.4	1.6	2.0	1.3	1.5	1.8	1.3	2.4	3.3	1.8
Pancreatitis (K85,K86.0-K86.1)															
2004.....	1.2	1.4	0.9	1.0	1.2	0.7	1.2	1.5	0.9	1.1	1.4	0.9	1.9	2.5	1.4
2003.....	1.2	1.5	0.9	1.0	1.4	0.7	1.2	1.5	1.0	1.1	1.3	0.9	1.9	2.7	1.3
2002.....	1.2	1.5	0.9	1.0	1.4	0.7	1.2	1.6	1.0	1.2	1.5	0.9	1.7	2.4	1.1
2001.....	1.3	1.5	1.1	1.0	1.0	1.0	1.3	1.6	1.1	1.2	1.5	1.0	2.0	2.5	1.5
2000.....	1.2	1.5	1.0	1.1	1.6	0.8	1.2	1.5	1.0	1.2	1.4	0.9	1.9	2.6	1.4
1999.....	1.2	1.5	1.0	1.1	1.5	0.8	1.2	1.5	1.0	1.1	1.4	0.9	1.9	2.7	1.4
Other disorders of biliary tract and pancreas (K83,K86.2-K86.9)															
2004.....	0.5	0.5	0.4	0.4	0.4	0.4	0.5	0.5	0.4	0.5	0.5	0.4	0.5	0.6	0.5
2003.....	0.5	0.6	0.4	0.4	0.5	0.4	0.5	0.6	0.4	0.5	0.6	0.4	0.6	0.7	0.6
2002.....	0.5	0.6	0.4	0.4	0.4	0.4	0.5	0.6	0.4	0.5	0.6	0.4	0.6	0.7	0.6
2001.....	0.5	0.5	0.4	0.5	0.5	0.5	0.5	0.5	0.4	0.4	0.5	0.4	0.6	0.6	0.5
2000.....	0.5	0.5	0.4	0.6	0.6	0.6	0.4	0.6	0.4	0.4	0.5	0.4	0.6	0.7	0.6
1999.....	0.4	0.5	0.4	0.6	0.6	0.6	0.4	0.5	0.4	0.4	0.5	0.3	0.5	0.6	0.4
All other diseases of digestive system (K90,K92)															
2004.....	2.6	3.1	2.3	1.4	1.7	1.2	2.7	3.1	2.3	2.7	3.1	2.3	3.4	4.2	2.9
2003.....	2.8	3.2	2.4	1.5	1.7	1.4	2.9	3.3	2.5	2.8	3.2	2.5	3.6	4.3	3.1
2002.....	2.9	3.3	2.5	1.5	1.8	1.2	2.9	3.4	2.6	2.9	3.3	2.5	3.7	4.5	3.1
2001.....	2.9	3.4	2.5	1.6	2.0	1.3	3.0	3.5	2.6	2.9	3.4	2.6	3.8	4.7	3.2
2000.....	2.9	3.4	2.6	1.6	1.8	1.4	3.0	3.5	2.6	2.9	3.4	2.5	4.0	4.8	3.5
1999.....	2.9	3.3	2.5	1.5	1.9	1.3	2.9	3.4	2.5	2.8	3.3	2.5	3.6	4.4	3.1
XII. Diseases of the skin and subcutaneous tissue (L00-L98)															
2004.....	1.4	1.3	1.5	1.1	1.1	1.1	1.4	1.3	1.5	1.2	1.1	1.2	3.6	3.2	3.8
2003.....	1.4	1.3	1.5	1.2	1.2	1.2	1.4	1.3	1.5	1.2	1.1	1.2	3.7	3.6	3.8
2002.....	1.4	1.3	1.4	1.0	1.1	1.0	1.4	1.3	1.4	1.1	1.1	1.1	4.0	3.3	4.3
2001.....	1.3	1.2	1.4	1.1	1.1	1.1	1.3	1.2	1.4	1.1	1.0	1.2	3.7	3.5	3.7
2000.....	1.4	1.2	1.4	1.1	1.0	1.2	1.3	1.2	1.4	1.1	1.0	1.2	4.0	3.8	4.0
1999.....	1.3	1.3	1.4	1.0	0.9	1.1	1.4	1.3	1.4	1.1	1.0	1.1	4.3	4.0	4.4

Worktable Orig13. Age-adjusted death rates for 358 selected causes, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), and year	All origins 1/			Hispanic			Non-Hispanic 2/			Non-Hispanic white			Non-Hispanic black		
	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female
Soft tissue disorders (M60-M79)															
2004.....	0.4	0.5	0.4	0.4	0.5	0.3	0.4	0.5	0.3	0.4	0.5	0.3	0.8	1.0	0.6
2003.....	0.4	0.5	0.4	0.3	0.5	0.2	0.4	0.5	0.4	0.4	0.5	0.4	0.7	0.8	0.7
2002.....	0.4	0.5	0.3	0.3	0.4	0.3	0.4	0.5	0.3	0.4	0.5	0.3	0.7	0.9	0.5
2001.....	0.4	0.5	0.3	0.4	0.4	0.3	0.4	0.5	0.3	0.3	0.4	0.3	0.7	0.9	0.6
2000.....	0.4	0.5	0.3	0.3	0.4	0.3	0.4	0.5	0.3	0.3	0.4	0.2	0.8	1.0	0.7
1999.....	0.4	0.5	0.3	0.4	0.5	0.3	0.4	0.5	0.3	0.3	0.4	0.2	0.7	1.0	0.6
Osteopathies, chondropathies and other disorders of musculoskeletal system and connective tissue (M80-M99)															
2004.....	1.1	0.8	1.2	0.7	0.5	0.8	1.1	0.8	1.3	1.1	0.8	1.3	1.2	1.3	1.0
2003.....	1.1	0.8	1.3	0.6	0.7	0.5	1.2	0.8	1.3	1.2	0.8	1.4	1.2	1.5	0.9
2002.....	1.2	0.8	1.3	0.7	0.6	0.7	1.2	0.8	1.3	1.2	0.8	1.4	1.1	1.3	1.0
2001.....	1.1	0.8	1.3	0.6	0.6	0.6	1.2	0.8	1.3	1.2	0.8	1.3	1.2	1.2	1.0
2000.....	1.1	0.8	1.2	0.7	*	0.9	1.1	0.8	1.2	1.1	0.8	1.3	1.1	1.4	0.9
1999.....	1.1	0.7	1.2	0.6	0.4	0.7	1.1	0.7	1.2	1.1	0.7	1.3	0.9	1.0	0.8
Osteoporosis (M80-M81)															
2004.....	0.6	0.2	0.7	0.2	*	0.3	0.6	0.2	0.8	0.6	0.2	0.8	0.1	*	0.1
2003.....	0.6	0.2	0.8	0.2	*	0.3	0.6	0.2	0.8	0.7	0.2	0.9	*	*	*
2002.....	0.6	0.2	0.8	0.2	*	0.3	0.7	0.2	0.9	0.7	0.2	0.9	0.1	*	0.2
2001.....	0.6	0.2	0.8	0.2	*	0.3	0.6	0.2	0.8	0.7	0.2	0.9	0.1	*	0.2
2000.....	0.6	0.2	0.8	0.3	*	0.5	0.6	0.2	0.8	0.7	0.2	0.9	0.1	*	*
1999.....	0.6	0.2	0.8	0.3	*	0.4	0.6	0.2	0.8	0.7	0.2	0.9	0.1	*	*
Other osteopathies, chondropathies and disorders of musculoskeletal system and connective tissue (M83-M99)															
2004.....	0.5	0.6	0.5	0.5	0.4	0.5	0.5	0.6	0.5	0.5	0.6	0.5	1.1	1.3	0.9
2003.....	0.6	0.6	0.5	0.4	0.6	0.2	0.6	0.6	0.5	0.5	0.6	0.5	1.1	1.4	0.8
2002.....	0.5	0.6	0.4	0.5	0.5	0.4	0.5	0.6	0.4	0.5	0.6	0.4	1.0	1.2	0.8
2001.....	0.5	0.6	0.5	0.4	0.6	0.3	0.6	0.6	0.5	0.5	0.6	0.5	1.0	1.2	0.9
2000.....	0.5	0.6	0.4	0.4	*	0.4	0.5	0.6	0.4	0.4	0.6	0.4	1.0	1.3	0.8
1999.....	0.4	0.5	0.4	0.3	*	0.3	0.4	0.5	0.4	0.4	0.5	0.4	0.8	0.9	0.7
XIV. Diseases of the genitourinary system (N00-N98)															
2004.....	20.1	23.0	18.3	16.5	18.6	15.1	20.3	23.2	18.4	18.5	21.5	16.7	38.5	43.8	35.3
2003.....	20.4	23.4	18.5	16.9	18.9	15.5	20.5	23.6	18.6	18.7	21.8	16.9	39.5	45.1	35.9
2002.....	20.2	23.2	18.3	16.0	17.5	14.9	20.3	23.5	18.5	18.6	21.6	16.7	38.7	44.9	35.1
2001.....	20.2	23.2	18.5	17.3	18.7	16.5	20.3	23.3	18.5	18.4	21.6	16.6	39.2	43.8	36.5
2000.....	19.8	23.0	18.0	16.9	18.5	15.9	19.9	23.1	18.0	18.1	21.4	16.2	38.2	43.1	35.3
1999.....	19.5	22.7	17.7	17.2	20.0	15.5	19.6	22.7	17.8	17.7	20.8	16.0	38.4	45.2	34.6
Diseases of urinary system (N00-N39)															
2004.....	19.8	22.4	18.1	16.2	18.2	14.9	19.9	22.6	18.2	18.2	20.8	16.5	38.0	42.9	35.0
2003.....	20.1	22.9	18.3	16.6	18.3	15.4	20.2	23.1	18.4	18.4	21.3	16.6	39.1	44.5	35.5
2002.....	19.9	22.6	18.1	15.7	17.1	14.7	20.0	22.9	18.3	18.3	21.1	16.5	38.2	44.1	34.7
2001.....	19.9	22.6	18.2	17.1	18.4	16.2	19.9	22.7	18.3	18.1	21.0	16.4	38.7	43.0	36.1
2000.....	19.5	22.4	17.7	16.6	18.0	15.7	19.5	22.5	17.8	17.8	20.8	16.0	37.8	42.4	35.0
1999.....	19.2	22.0	17.5	17.0	19.7	15.3	19.2	22.1	17.5	17.4	20.1	15.8	37.8	44.2	34.1
Glomerular and renal tubulo-interstitial diseases (N00-N15)															
2004.....	0.6	0.7	0.6	0.6	0.6	0.6	0.6	0.7	0.6	0.6	0.7	0.6	0.9	1.1	0.8
2003.....	0.7	0.7	0.6	0.6	0.7	0.6	0.7	0.7	0.6	0.6	0.7	0.6	0.9	1.1	0.9
2002.....	0.7	0.8	0.6	0.6	0.8	0.6	0.7	0.8	0.6	0.6	0.7	0.6	0.8	1.0	0.7
2001.....	0.7	0.8	0.6	0.6	0.7	0.6	0.7	0.7	0.6	0.6	0.7	0.6	0.9	1.0	0.8
2000.....	0.7	0.8	0.7	0.6	0.7	0.6	0.7	0.8	0.7	0.6	0.7	0.6	1.1	1.2	1.0
1999.....	0.7	0.9	0.7	0.8	0.8	0.8	0.7	0.9	0.7	0.7	0.8	0.6	1.1	1.5	0.9

Worktable Orig13. Age-adjusted death rates for 358 selected causes, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Table with 17 columns: Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), and year; All origins 1/ (Both Sexes, Male, Female); Hispanic (Both Sexes, Male, Female); Non-Hispanic 2/ (Both Sexes, Male, Female); Non-Hispanic white (Both Sexes, Male, Female); Non-Hispanic black (Both Sexes, Male, Female). Rows include categories like Glomerular diseases (N00-N07), Acute and rapidly progressive nephritic syndrome (N00-N01), Nephrotic syndrome (N04), Other and unspecified glomerular diseases (N02-N03,N05-N07), Renal tubulo-interstitial diseases (N10-N15), Renal failure (N17-N19), and Urolithiasis (N20-N23).

Worktable Orig13. Age-adjusted death rates for 358 selected causes, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), and year	All origins 1/			Hispanic			Non-Hispanic 2/			Non-Hispanic white			Non-Hispanic black		
	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female
Other and unspecified disorders of kidney (N25-N27)															
2004.....	0.0	0.0	0.0	*	*	*	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2003.....	0.0	0.0	0.0	*	*	*	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2002.....	0.0	0.0	0.0	*	*	*	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2001.....	0.0	*	0.0	*	*	*	0.0	*	0.0	0.0	*	0.0	*	*	*
2000.....	0.0	0.0	0.0	*	*	*	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
1999.....	0.0	0.0	0.0	*	*	*	0.0	0.0	0.0	0.0	0.0	0.0	0.1	*	*
Urinary tract infection, site not specified (N39.0)															
2004.....	4.6	4.0	5.0	3.5	3.2	3.8	4.7	4.0	5.0	4.6	3.8	4.9	6.6	6.5	6.6
2003.....	4.7	3.9	5.1	3.3	2.8	3.5	4.7	4.0	5.1	4.6	3.9	5.0	6.8	6.2	7.1
2002.....	4.7	4.0	5.2	3.3	2.3	3.9	4.8	4.0	5.2	4.6	3.9	5.1	6.9	6.8	7.0
2001.....	5.0	4.1	5.4	3.8	2.9	4.4	5.0	4.2	5.4	4.8	4.0	5.2	7.4	7.1	7.5
2000.....	5.0	4.4	5.3	4.0	3.1	4.5	5.0	4.4	5.4	4.9	4.2	5.2	7.2	7.1	7.2
1999.....	5.2	4.4	5.5	4.0	3.4	4.4	5.2	4.5	5.6	5.0	4.2	5.5	7.6	8.1	7.2
Other diseases of urinary system (N28-N36,N39.1-N39.9)															
2004.....	0.4	0.5	0.3	0.4	0.5	0.3	0.4	0.5	0.4	0.4	0.5	0.3	0.8	0.9	0.7
2003.....	0.4	0.5	0.3	0.2	0.2	*	0.4	0.5	0.3	0.4	0.5	0.3	0.6	0.8	0.5
2002.....	0.4	0.5	0.3	0.3	0.4	0.2	0.4	0.5	0.3	0.4	0.4	0.3	0.7	0.9	0.6
2001.....	0.4	0.4	0.3	0.4	0.5	0.4	0.4	0.5	0.3	0.4	0.4	0.3	0.7	0.8	0.6
2000.....	0.4	0.5	0.3	0.3	0.4	0.2	0.4	0.5	0.3	0.4	0.5	0.3	0.7	0.9	0.6
1999.....	0.4	0.5	0.3	0.3	0.3	0.2	0.4	0.5	0.3	0.3	0.4	0.3	0.7	1.0	0.5
Diseases of male genital organs (N40-N50)															
2004.....	0.2	0.6	...	0.2	0.5	...	0.2	0.6	...	0.2	0.6	...	0.3	0.8	...
2003.....	0.2	0.6	...	0.2	0.6	...	0.2	0.6	...	0.2	0.6	...	0.2	0.6	...
2002.....	0.2	0.6	...	0.2	0.4	...	0.2	0.5	...	0.2	0.5	...	0.3	0.8	...
2001.....	0.2	0.6	...	*	*	...	0.2	0.6	...	0.2	0.6	...	0.3	0.8	...
2000.....	0.2	0.6	...	0.2	0.6	...	0.2	0.6	...	0.2	0.6	...	0.3	0.8	...
1999.....	0.2	0.6	...	*	*	...	0.2	0.6	...	0.2	0.6	...	0.3	0.9	...
Hyperplasia of prostate (N40)															
2004.....	0.2	0.5	...	*	*	...	0.2	0.5	...	0.2	0.5	...	0.1	0.4	...
2003.....	0.2	0.5	...	0.2	0.4	...	0.2	0.5	...	0.2	0.5	...	0.1	0.4	...
2002.....	0.1	0.4	...	*	*	...	0.1	0.4	...	0.1	0.4	...	0.2	0.5	...
2001.....	0.1	0.5	...	*	*	...	0.2	0.5	...	0.2	0.5	...	0.1	0.4	...
2000.....	0.2	0.5	...	*	*	...	0.2	0.5	...	0.2	0.5	...	0.2	0.5	...
1999.....	0.2	0.5	...	*	*	...	0.2	0.5	...	0.2	0.5	...	0.2	0.6	...
Other diseases of male genital organs (N41-N50)															
2004.....	0.1	0.2	...	*	*	...	0.1	0.2	...	0.1	0.1	...	0.1	0.3	...
2003.....	0.0	0.1	...	*	*	...	0.0	0.1	...	0.0	0.1	...	0.1	0.2	...
2002.....	0.1	0.1	...	*	*	...	0.1	0.1	...	0.0	0.1	...	0.1	0.3	...
2001.....	0.1	0.1	...	*	*	...	0.1	0.1	...	0.1	0.1	...	0.1	0.3	...
2000.....	0.0	0.1	...	*	*	...	0.0	0.1	...	0.0	0.1	...	0.1	0.3	...
1999.....	0.1	0.2	...	*	*	...	0.1	0.2	...	0.0	0.1	...	0.1	0.4	...
Disorders of breast (N60-N64)															
2004.....	0.0	*	0.0	*	*	*	0.0	*	0.0	0.0	*	0.0	*	*	*
2003.....	0.0	*	0.0	*	*	*	0.0	*	0.0	0.0	*	0.0	*	*	*
2002.....	0.0	*	0.0	*	*	*	0.0	*	0.0	0.0	*	0.0	*	*	*
2001.....	0.0	*	0.0	*	*	*	0.0	*	0.0	0.0	*	0.0	*	*	*
2000.....	0.0	*	0.0	*	*	*	0.0	*	0.0	0.0	*	0.0	*	*	*
1999.....	0.0	*	0.0	*	*	*	0.0	*	0.0	0.0	*	0.0	*	*	*

Worktable Orig13. Age-adjusted death rates for 358 selected causes, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), and year	All origins 1/			Hispanic			Non-Hispanic 2/			Non-Hispanic white			Non-Hispanic black		
	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female
Disorders of female genital organs (N70-N98)															
2004.....	0.1	...	0.2	0.1	...	0.2	0.1	...	0.2	0.1	...	0.2	0.2	...	0.3
2003.....	0.1	...	0.2	0.1	...	0.2	0.1	...	0.2	0.1	...	0.2	0.2	...	0.3
2002.....	0.1	...	0.2	0.1	...	0.2	0.1	...	0.2	0.1	...	0.2	0.2	...	0.3
2001.....	0.1	...	0.2	0.1	...	0.2	0.1	...	0.2	0.1	...	0.2	0.2	...	0.4
2000.....	0.1	...	0.2	*	...	*	0.1	...	0.2	0.1	...	0.2	0.2	...	0.3
1999.....	0.1	...	0.2	*	...	*	0.1	...	0.2	0.1	...	0.2	0.3	...	0.4
Inflammatory diseases of female pelvic organs (N70-N76)															
2004.....	0.0	...	0.1	*	...	*	0.0	...	0.1	0.0	...	0.0	*	...	*
2003.....	0.0	...	0.1	*	...	*	0.0	...	0.1	0.0	...	0.1	*	...	*
2002.....	0.0	...	0.1	*	...	*	0.0	...	0.1	0.0	...	0.0	0.1	...	0.1
2001.....	0.0	...	0.1	*	...	*	0.0	...	0.1	0.0	...	0.1	0.1	...	0.1
2000.....	0.0	...	0.1	*	...	*	0.0	...	0.1	0.0	...	0.1	0.1	...	0.1
1999.....	0.0	...	0.1	*	...	*	0.0	...	0.1	0.0	...	0.0	0.1	...	0.2
Noninflammatory disorders of female genital tract (N80-N98)															
2004.....	0.1	...	0.1	*	...	*	0.1	...	0.1	0.1	...	0.1	0.1	...	0.2
2003.....	0.1	...	0.1	*	...	*	0.1	...	0.1	0.1	...	0.1	0.1	...	0.2
2002.....	0.1	...	0.1	*	...	*	0.1	...	0.1	0.0	...	0.1	0.1	...	0.2
2001.....	0.1	...	0.1	*	...	*	0.1	...	0.1	0.0	...	0.1	0.1	...	0.2
2000.....	0.1	...	0.1	*	...	*	0.1	...	0.1	0.1	...	0.1	0.1	...	0.2
1999.....	0.1	...	0.1	*	...	*	0.1	...	0.1	0.1	...	0.1	0.2	...	0.3
XV. Pregnancy, childbirth and the puerperium (000-099)															
2004.....	0.2	...	0.5	0.2	...	0.5	0.2	...	0.5	0.1	...	0.3	0.6	...	1.2
2003.....	0.2	...	0.4	0.2	...	0.5	0.2	...	0.3	0.1	...	0.2	0.5	...	1.0
2002.....	0.1	...	0.3	0.2	...	0.3	0.1	...	0.2	0.1	...	0.2	0.4	...	0.8
2001.....	0.1	...	0.3	0.2	...	0.4	0.1	...	0.3	0.1	...	0.2	0.4	...	0.8
2000.....	0.1	...	0.3	0.2	...	0.4	0.1	...	0.3	0.1	...	0.2	0.4	...	0.7
1999.....	0.2	...	0.3	0.2	...	0.4	0.1	...	0.3	0.1	...	0.2	0.4	...	0.8
Pregnancy with abortive outcome (000-007)															
2004.....	0.0	...	0.0	*	...	*	0.0	...	0.0	*	...	*	*	...	*
2003.....	0.0	...	0.0	*	...	*	0.0	...	0.0	*	...	*	*	...	*
2002.....	0.0	...	0.0	*	...	*	0.0	...	0.0	*	...	*	*	...	*
2001.....	0.0	...	0.0	*	...	*	0.0	...	0.0	*	...	*	*	...	*
2000.....	0.0	...	0.0	*	...	*	0.0	...	0.0	*	...	*	*	...	*
1999.....	0.0	...	0.0	*	...	*	0.0	...	0.0	*	...	*	*	...	*
Ectopic pregnancy (000)															
2004.....	*	...	*	*	...	*	*	...	*	*	...	*	*	...	*
2003.....	0.0	...	0.0	*	...	*	*	...	*	*	...	*	*	...	*
2002.....	*	...	*	*	...	*	*	...	*	*	...	*	*	...	*
2001.....	0.0	...	0.0	*	...	*	0.0	...	0.0	*	...	*	*	...	*
2000.....	0.0	...	0.0	*	...	*	0.0	...	0.0	*	...	*	*	...	*
1999.....	*	...	*	*	...	*	*	...	*	*	...	*	*	...	*
Spontaneous abortion (003)															
2004.....	*	...	*	*	...	*	*	...	*	*	...	*	*	...	*
2003.....	*	...	*	*	...	*	*	...	*	*	...	*	*	...	*
2002.....	*	...	*	*	...	*	*	...	*	*	...	*	*	...	*
2001.....	*	...	*	*	...	*	*	...	*	*	...	*	*	...	*
2000.....	*	...	*	*	...	*	*	...	*	*	...	*	*	...	*
1999.....	*	...	*	*	...	*	*	...	*	*	...	*	*	...	*

Worktable Orig13. Age-adjusted death rates for 358 selected causes, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), and year	All origins 1/			Hispanic			Non-Hispanic 2/			Non-Hispanic white			Non-Hispanic black		
	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female
Medical abortion (O04)															
2004.....	*	...	*	*	...	*	*	...	*	*	...	*	*	...	*
2003.....	*	...	*	*	...	*	*	...	*	*	...	*	*	...	*
2002.....	*	...	*	*	...	*	*	...	*	*	...	*	*	...	*
2001.....	*	...	*	*	...	*	*	...	*	*	...	*	*	...	*
2000.....	*	...	*	*	...	*	*	...	*	*	...	*	*	...	*
1999.....	*	...	*	*	...	*	*	...	*	*	...	*	*	...	*
Other abortion (O05)															
2004.....	*	...	*	*	...	*	*	...	*	*	...	*	*	...	*
2003.....	*	...	*	*	...	*	*	...	*	*	...	*	*	...	*
2002.....	*	...	*	*	...	*	*	...	*	*	...	*	*	...	*
2001.....	*	...	*	*	...	*	*	...	*	*	...	*	*	...	*
2000.....	*	...	*	*	...	*	*	...	*	*	...	*	*	...	*
1999.....	*	...	*	*	...	*	*	...	*	*	...	*	*	...	*
Other and unspecified pregnancy with abortive outcome (O01-O02,O06-O07)															
2004.....	*	...	*	*	...	*	*	...	*	*	...	*	*	...	*
2003.....	*	...	*	*	...	*	*	...	*	*	...	*	*	...	*
2002.....	*	...	*	*	...	*	*	...	*	*	...	*	*	...	*
2001.....	*	...	*	*	...	*	*	...	*	*	...	*	*	...	*
2000.....	*	...	*	*	...	*	*	...	*	*	...	*	*	...	*
1999.....	*	...	*	*	...	*	*	...	*	*	...	*	*	...	*
Other direct obstetric deaths (O10-O92)															
2004.....	0.1	...	0.2	0.1	...	0.2	0.1	...	0.2	0.1	...	0.1	0.3	...	0.7
2003.....	0.1	...	0.2	0.2	...	0.4	0.1	...	0.2	0.1	...	0.1	0.3	...	0.7
2002.....	0.1	...	0.2	0.1	...	0.3	0.1	...	0.2	0.0	...	0.1	0.3	...	0.6
2001.....	0.1	...	0.2	0.2	...	0.4	0.1	...	0.2	0.1	...	0.1	0.3	...	0.6
2000.....	0.1	...	0.2	0.2	...	0.3	0.1	...	0.2	0.1	...	0.1	0.3	...	0.5
1999.....	0.1	...	0.2	0.1	...	0.3	0.1	...	0.2	0.1	...	0.1	0.3	...	0.6
Eclampsia and pre-eclampsia (O11,O13-O16)															
2004.....	0.0	...	0.0	*	...	*	0.0	...	0.0	*	...	*	0.1	...	0.1
2003.....	0.0	...	0.0	*	...	*	0.0	...	0.0	*	...	*	0.1	...	0.1
2002.....	0.0	...	0.0	*	...	*	0.0	...	0.0	*	...	*	0.1	...	0.1
2001.....	0.0	...	0.1	0.1	...	0.1	0.0	...	0.0	*	...	*	0.1	...	0.1
2000.....	0.0	...	0.1	0.1	...	0.1	0.0	...	0.1	0.0	...	0.0	0.1	...	0.1
1999.....	0.0	...	0.1	*	...	*	0.0	...	0.1	0.0	...	0.0	0.1	...	0.2
Hemorrhage of pregnancy and childbirth and placenta previa (O20,O44-O46,O67,O72)															
2004.....	0.0	...	0.0	*	...	*	0.0	...	0.0	*	...	*	*	...	*
2003.....	0.0	...	0.0	*	...	*	0.0	...	0.0	*	...	*	*	...	*
2002.....	0.0	...	0.0	*	...	*	0.0	...	0.0	*	...	*	*	...	*
2001.....	0.0	...	0.0	*	...	*	0.0	...	0.0	*	...	*	*	...	*
2000.....	0.0	...	0.0	*	...	*	0.0	...	0.0	*	...	*	*	...	*
1999.....	0.0	...	0.0	*	...	*	0.0	...	0.0	*	...	*	*	...	*
Complications predominately related to the puerperium (O85-O92)															
2004.....	0.0	...	0.1	*	...	*	0.0	...	0.1	0.0	...	0.0	0.1	...	0.2
2003.....	0.0	...	0.1	*	...	*	0.0	...	0.1	0.0	...	0.0	0.1	...	0.2
2002.....	0.0	...	0.1	*	...	*	0.0	...	0.1	0.0	...	0.0	0.1	...	0.2
2001.....	0.0	...	0.1	*	...	*	0.0	...	0.1	0.0	...	0.1	0.1	...	0.2
2000.....	0.0	...	0.1	*	...	*	0.0	...	0.1	0.0	...	0.1	0.1	...	0.2
1999.....	0.0	...	0.1	*	...	*	0.0	...	0.1	0.0	...	0.1	0.1	...	0.2

Worktable Orig13. Age-adjusted death rates for 358 selected causes, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), and year	All origins 1/			Hispanic			Non-Hispanic 2/			Non-Hispanic white			Non-Hispanic black		
	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female
Obstetric embolism (088)															
2004.....	0.0	...	0.0	*	...	*	0.0	...	0.0	0.0	...	0.0	*	...	*
2003.....	0.0	...	0.0	*	...	*	0.0	...	0.0	*	...	*	*	...	*
2002.....	0.0	...	0.0	*	...	*	0.0	...	0.0	0.0	...	0.0	*	...	*
2001.....	0.0	...	0.0	*	...	*	0.0	...	0.0	0.0	...	0.0	0.1	...	0.1
2000.....	0.0	...	0.0	*	...	*	0.0	...	0.0	0.0	...	0.0	*	...	*
1999.....	0.0	...	0.0	*	...	*	0.0	...	0.0	0.0	...	0.0	*	...	*
Other complications predominately related to the puerperium (085-087,089-092)															
2004.....	0.0	...	0.0	*	...	*	0.0	...	0.0	0.0	...	0.0	0.1	...	0.1
2003.....	0.0	...	0.0	*	...	*	0.0	...	0.0	*	...	*	*	...	*
2002.....	0.0	...	0.0	*	...	*	0.0	...	0.0	*	...	*	0.1	...	0.1
2001.....	0.0	...	0.0	*	...	*	0.0	...	0.0	*	...	*	*	...	*
2000.....	0.0	...	0.0	*	...	*	0.0	...	0.0	*	...	*	*	...	*
1999.....	0.0	...	0.0	*	...	*	0.0	...	0.0	*	...	*	*	...	*
All other direct obstetric causes (010,012,021-043,047-066,068-071,073-075)															
2004.....	0.1	...	0.1	*	...	*	0.1	...	0.1	0.0	...	0.1	0.2	...	0.3
2003.....	0.0	...	0.1	0.1	...	0.2	0.0	...	0.1	0.0	...	0.0	0.1	...	0.3
2002.....	0.0	...	0.1	0.1	...	0.1	0.1	...	0.1	0.0	...	0.0	0.1	...	0.2
2001.....	0.0	...	0.1	0.0	...	0.1	0.0	...	0.1	0.0	...	0.0	0.1	...	0.2
2000.....	0.0	...	0.1	0.1	...	0.1	0.0	...	0.1	0.0	...	0.0	0.1	...	0.2
1999.....	0.0	...	0.1	*	...	*	0.0	...	0.1	0.0	...	0.0	0.1	...	0.2
Obstetric death of unspecified cause (095)															
2004.....	0.0	...	0.0	*	...	*	0.0	...	0.0	*	...	*	*	...	*
2003.....	0.0	...	0.0	*	...	*	0.0	...	0.0	*	...	*	*	...	*
2002.....	*	...	*	*	...	*	*	...	*	*	...	*	*	...	*
2001.....	*	...	*	*	...	*	*	...	*	*	...	*	*	...	*
2000.....	*	...	*	*	...	*	*	...	*	*	...	*	*	...	*
1999.....	*	...	*	*	...	*	*	...	*	*	...	*	*	...	*
Other deaths related to pregnancy, childbirth and the puerperium (096-097)															
2004.....	0.1	...	0.1	0.1	...	0.2	0.1	...	0.1	0.0	...	0.1	0.1	...	0.2
2003.....	0.0	...	0.0	*	...	*	0.0	...	0.0	0.0	...	0.0	*	...	*
2002.....	0.0	...	0.0	*	...	*	*	...	*	*	...	*	*	...	*
2001.....	*	...	*	*	...	*	*	...	*	*	...	*	*	...	*
2000.....	*	...	*	*	...	*	*	...	*	*	...	*	*	...	*
1999.....	*	...	*	*	...	*	*	...	*	*	...	*	*	...	*
Indirect obstetric deaths (098-099)															
2004.....	0.0	...	0.1	0.0	...	0.1	0.1	...	0.1	0.0	...	0.1	0.1	...	0.2
2003.....	0.0	...	0.1	*	...	*	0.0	...	0.1	0.0	...	0.1	0.1	...	0.1
2002.....	0.0	...	0.0	*	...	*	0.0	...	0.0	0.0	...	0.0	*	...	*
2001.....	0.0	...	0.0	*	...	*	0.0	...	0.0	0.0	...	0.0	0.0	...	0.1
2000.....	0.0	...	0.0	*	...	*	0.0	...	0.0	0.0	...	0.0	*	...	*
1999.....	0.0	...	0.0	*	...	*	0.0	...	0.0	0.0	...	0.0	*	...	*
XVI. Certain conditions originating in the perinatal period (P00-P96)															
2004.....	4.8	5.4	4.2	4.1	4.6	3.6	4.9	5.5	4.4	3.6	4.1	3.2	10.3	11.4	9.3
2003.....	4.9	5.5	4.3	4.2	4.6	3.8	5.1	5.7	4.4	3.8	4.3	3.3	10.3	11.5	9.0
2002.....	4.9	5.4	4.3	4.0	4.4	3.6	5.0	5.6	4.5	3.8	4.2	3.3	10.2	11.1	9.3
2001.....	4.7	5.3	4.2	3.8	4.2	3.3	4.9	5.4	4.4	3.6	4.0	3.2	10.7	11.8	9.5
2000.....	5.1	5.6	4.5	3.8	4.2	3.5	5.3	5.9	4.7	3.8	4.3	3.4	11.7	13.0	10.3
1999.....	5.2	5.7	4.6	3.9	4.3	3.6	5.4	5.9	4.8	3.8	4.3	3.4	12.1	13.5	10.8

Worktable Orig13. Age-adjusted death rates for 358 selected causes, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), and year	All origins 1/			Hispanic			Non-Hispanic 2/			Non-Hispanic white			Non-Hispanic black		
	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female
Newborn affected by maternal factors and by complications of pregnancy, labor and delivery (P00-P04)															
2004.....	1.1	1.1	1.0	0.8	0.9	0.7	1.1	1.2	1.0	0.8	0.9	0.8	2.3	2.4	2.1
2003.....	1.1	1.2	1.0	0.9	1.0	0.8	1.1	1.2	1.0	0.9	1.0	0.8	2.1	2.3	2.0
2002.....	1.0	1.2	0.9	0.8	0.9	0.7	1.1	1.2	1.0	0.9	0.9	0.8	2.1	2.3	2.0
2001.....	1.0	1.1	0.9	0.7	0.8	0.5	1.0	1.1	0.9	0.8	0.9	0.7	2.1	2.2	1.9
2000.....	1.0	1.1	0.9	0.7	0.7	0.6	1.1	1.2	1.0	0.8	0.9	0.8	2.1	2.3	2.0
1999.....	1.0	1.1	0.9	0.6	0.6	0.7	1.1	1.1	1.0	0.8	0.9	0.7	2.3	2.5	2.1
Disorders related to short gestation and low birth weight, not elsewhere classified (P07)															
2004.....	1.6	1.8	1.4	1.2	1.4	1.1	1.6	1.8	1.5	1.1	1.2	1.0	3.8	4.3	3.4
2003.....	1.7	1.8	1.5	1.4	1.5	1.3	1.7	1.9	1.5	1.1	1.3	1.0	4.1	4.5	3.6
2002.....	1.6	1.7	1.4	1.3	1.4	1.2	1.6	1.8	1.5	1.1	1.2	1.0	3.9	4.3	3.5
2001.....	1.5	1.6	1.4	1.1	1.2	1.1	1.6	1.7	1.5	1.0	1.1	1.0	4.0	4.4	3.7
2000.....	1.6	1.7	1.5	1.1	1.2	1.1	1.7	1.8	1.6	1.0	1.1	1.0	4.4	4.7	4.0
1999.....	1.6	1.7	1.4	1.2	1.3	1.1	1.7	1.8	1.5	1.0	1.1	0.9	4.5	4.9	4.0
Birth trauma (P10-P15)															
2004.....	0.0	*	*	*	*	*	0.0	*	*	*	*	*	*	*	*
2003.....	0.0	*	*	*	*	*	0.0	*	*	*	*	*	*	*	*
2002.....	0.1	0.1	0.1	0.1	0.2	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.2	0.2	0.1
2001.....	0.0	0.1	0.0	0.0	*	*	0.0	0.1	0.0	0.0	0.0	0.0	0.1	0.1	*
2000.....	0.0	0.0	0.0	*	*	*	0.0	*	0.0	0.0	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	0.0	*	*	*	*	*	*	*	*
Intrauterine hypoxia and birth asphyxia (P20-P21)															
2004.....	0.2	0.2	0.2	0.2	0.2	0.1	0.2	0.2	0.2	0.2	0.2	0.1	0.3	0.3	0.3
2003.....	0.2	0.2	0.2	0.2	0.2	0.1	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.3	0.3
2002.....	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.3	0.3
2001.....	0.2	0.2	0.2	0.2	0.2	0.1	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.3	0.3
2000.....	0.2	0.3	0.2	0.2	0.2	0.2	0.2	0.3	0.2	0.2	0.2	0.2	0.4	0.4	0.3
1999.....	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.2	0.2	0.2	0.2	0.4	0.4	0.3
Respiratory distress of newborn (P22)															
2004.....	0.3	0.3	0.3	0.3	0.3	0.2	0.3	0.3	0.3	0.2	0.3	0.2	0.6	0.7	0.5
2003.....	0.3	0.3	0.2	0.2	0.3	0.2	0.3	0.3	0.2	0.2	0.3	0.2	0.5	0.6	0.5
2002.....	0.3	0.4	0.3	0.3	0.3	0.2	0.3	0.4	0.3	0.3	0.3	0.2	0.7	0.7	0.7
2001.....	0.3	0.4	0.3	0.3	0.3	0.3	0.4	0.4	0.3	0.3	0.3	0.2	0.8	0.9	0.6
2000.....	0.4	0.5	0.3	0.3	0.3	0.3	0.4	0.5	0.3	0.3	0.4	0.2	0.8	1.0	0.7
1999.....	0.4	0.5	0.3	0.3	0.4	0.3	0.4	0.5	0.3	0.3	0.3	0.3	0.9	1.1	0.7
Other respiratory conditions originating in the perinatal period (P23-P28)															
2004.....	0.4	0.5	0.4	0.4	0.4	0.3	0.4	0.5	0.4	0.3	0.4	0.3	0.9	0.9	0.8
2003.....	0.4	0.5	0.3	0.4	0.4	0.3	0.4	0.5	0.4	0.4	0.4	0.3	0.8	1.0	0.7
2002.....	0.4	0.5	0.4	0.3	0.4	0.3	0.5	0.5	0.4	0.4	0.4	0.3	0.9	1.0	0.8
2001.....	0.5	0.6	0.5	0.4	0.4	0.3	0.6	0.6	0.5	0.4	0.5	0.4	1.1	1.3	1.0
2000.....	0.6	0.6	0.5	0.5	0.5	0.4	0.6	0.7	0.5	0.4	0.5	0.4	1.2	1.4	1.0
1999.....	0.6	0.7	0.5	0.5	0.6	0.4	0.7	0.7	0.6	0.5	0.6	0.4	1.3	1.4	1.2
Infections specific to the perinatal period (P35-P39)															
2004.....	0.4	0.4	0.3	0.3	0.4	0.3	0.3	0.4	0.3	0.3	0.3	0.2	0.7	0.8	0.6
2003.....	0.3	0.4	0.3	0.3	0.3	0.3	0.3	0.4	0.3	0.3	0.3	0.2	0.7	0.8	0.6
2002.....	0.3	0.4	0.3	0.3	0.3	0.2	0.3	0.3	0.3	0.3	0.3	0.2	0.6	0.6	0.6
2001.....	0.3	0.3	0.3	0.3	0.3	0.2	0.3	0.3	0.3	0.2	0.3	0.2	0.6	0.7	0.6
2000.....	0.3	0.4	0.3	0.3	0.4	0.2	0.3	0.4	0.3	0.3	0.3	0.2	0.7	0.8	0.6
1999.....	0.3	0.4	0.3	0.2	0.3	0.2	0.3	0.4	0.3	0.2	0.3	0.2	0.7	0.9	0.6

Worktable Orig13. Age-adjusted death rates for 358 selected causes, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), and year	All origins 1/			Hispanic			Non-Hispanic 2/			Non-Hispanic white			Non-Hispanic black		
	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female
Other and unspecified conditions originating in the perinatal period (P05,P08,P29,P50-P96)															
2004.....	0.9	1.0	0.8	0.9	1.0	0.8	0.9	1.0	0.8	0.7	0.8	0.6	1.7	1.9	1.5
2003.....	0.9	1.0	0.8	0.8	0.9	0.8	0.9	1.0	0.8	0.7	0.8	0.6	1.7	2.0	1.4
2002.....	0.8	0.9	0.7	0.8	0.8	0.7	0.8	0.9	0.7	0.6	0.7	0.5	1.5	1.6	1.3
2001.....	0.8	1.0	0.7	0.8	0.9	0.7	0.8	1.0	0.7	0.6	0.7	0.5	1.6	1.8	1.3
2000.....	1.0	1.1	0.8	0.8	0.9	0.7	1.0	1.1	0.8	0.7	0.8	0.7	2.0	2.3	1.6
1999.....	1.0	1.1	0.8	0.8	0.9	0.8	1.0	1.1	0.8	0.7	0.8	0.6	2.0	2.2	1.9
XVII. Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)															
2004.....	3.6	3.7	3.4	3.2	3.3	3.1	3.6	3.8	3.4	3.5	3.7	3.4	4.1	4.5	3.7
2003.....	3.6	3.8	3.4	3.4	3.5	3.2	3.6	3.8	3.4	3.5	3.6	3.4	4.0	4.5	3.7
2002.....	3.7	3.8	3.5	3.4	3.6	3.2	3.7	3.8	3.5	3.6	3.8	3.4	4.2	4.5	4.0
2001.....	3.6	3.8	3.4	3.4	3.8	3.0	3.6	3.8	3.4	3.5	3.8	3.3	4.2	4.4	4.1
2000.....	3.8	4.1	3.6	3.6	3.9	3.3	3.8	4.1	3.6	3.7	3.9	3.5	4.5	4.9	4.2
1999.....	3.8	3.9	3.6	3.4	3.4	3.4	3.8	4.0	3.6	3.7	3.9	3.5	4.5	4.8	4.2
Anencephaly and similar malformations (Q00)															
2004.....	0.1	0.1	0.1	0.1	0.1	0.2	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2003.....	0.1	0.1	0.1	0.2	0.2	0.2	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	0.1
2002.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	0.1
2001.....	0.1	0.1	0.1	0.1	0.1	0.2	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	0.1
2000.....	0.1	0.1	0.1	0.1	0.1	0.2	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	0.1
1999.....	0.1	0.1	0.1	0.2	0.1	0.2	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	0.1
Spina bifida (Q05)															
2004.....	0.0	0.0	0.0	0.1	*	*	0.0	0.0	0.0	0.0	0.0	0.1	*	*	*
2003.....	0.0	0.0	0.0	*	*	*	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2002.....	0.1	0.1	0.1	*	*	*	0.1	0.1	0.0	0.1	0.1	0.1	*	*	*
2001.....	0.0	0.0	0.1	*	*	*	0.0	0.1	0.1	0.1	0.1	0.0	*	*	*
2000.....	0.0	0.0	0.0	0.0	*	*	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
1999.....	0.0	0.0	0.0	*	*	*	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
All other congenital malformations of nervous system (Q01-Q04,Q06-Q07)															
2004.....	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.2	0.3	0.4	0.3	0.4
2003.....	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.4	0.3
2002.....	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3
2001.....	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.2	0.3	0.3	0.4
2000.....	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.2	0.3	0.2	0.4	0.4	0.4
1999.....	0.3	0.3	0.3	0.2	0.2	0.3	0.3	0.3	0.3	0.3	0.3	0.2	0.4	0.3	0.4
Congenital malformations of heart (Q20-Q24)															
2004.....	1.0	1.1	0.9	0.9	1.0	0.8	1.0	1.1	0.9	1.0	1.1	0.9	1.2	1.3	1.1
2003.....	1.1	1.2	1.0	0.9	1.0	0.8	1.1	1.2	1.0	1.1	1.1	1.0	1.3	1.4	1.2
2002.....	1.1	1.2	1.0	1.1	1.2	0.9	1.1	1.2	1.0	1.1	1.2	1.0	1.3	1.4	1.2
2001.....	1.1	1.2	1.0	0.9	1.1	0.8	1.2	1.3	1.1	1.1	1.2	1.0	1.5	1.5	1.4
2000.....	1.2	1.3	1.2	1.1	1.2	1.0	1.2	1.3	1.1	1.2	1.3	1.1	1.5	1.6	1.3
1999.....	1.3	1.4	1.2	1.1	1.2	1.1	1.3	1.4	1.2	1.3	1.4	1.1	1.5	1.7	1.4
Other congenital malformations of circulatory system (Q25-Q28)															
2004.....	0.3	0.3	0.3	0.3	0.2	0.3	0.3	0.4	0.3	0.3	0.3	0.3	0.4	0.5	0.3
2003.....	0.3	0.3	0.3	0.3	0.3	0.2	0.3	0.3	0.3	0.3	0.3	0.3	0.4	0.4	0.3
2002.....	0.3	0.4	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.5	0.6	0.4
2001.....	0.3	0.3	0.3	0.4	0.6	0.3	0.3	0.3	0.3	0.3	0.3	0.2	0.3	0.4	0.3
2000.....	0.3	0.4	0.3	0.4	0.4	0.3	0.3	0.4	0.3	0.3	0.3	0.3	0.4	0.5	0.4
1999.....	0.3	0.4	0.3	0.3	0.4	0.3	0.3	0.4	0.3	0.3	0.3	0.3	0.5	0.5	0.4

Worktable Orig13. Age-adjusted death rates for 358 selected causes, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), and year	All origins 1/			Hispanic			Non-Hispanic 2/			Non-Hispanic white			Non-Hispanic black		
	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female
Congenital malformations of respiratory system (Q30-Q34)															
2004.....	0.2	0.3	0.2	0.2	0.3	0.2	0.2	0.3	0.2	0.2	0.2	0.2	0.3	0.4	0.3
2003.....	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.2	0.2	0.2	0.2	0.3	0.4	0.2
2002.....	0.2	0.2	0.2	0.2	0.3	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.3	0.3
2001.....	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.4	0.2
2000.....	0.3	0.3	0.2	0.2	0.3	0.2	0.3	0.3	0.2	0.2	0.3	0.2	0.4	0.4	0.3
1999.....	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.2	0.2	0.2	0.2	0.3	0.4	0.2
Other congenital malformations and deformations (Q10-Q18,Q35-Q89)															
2004.....	0.9	1.0	0.8	0.8	0.8	0.7	0.9	1.0	0.8	0.9	1.0	0.8	1.0	1.2	0.8
2003.....	0.9	1.0	0.8	0.9	1.0	0.8	0.9	1.0	0.8	0.9	1.0	0.8	1.1	1.3	0.9
2002.....	0.9	1.0	0.8	0.8	0.8	0.8	0.9	1.0	0.8	0.9	1.0	0.8	1.0	1.2	0.9
2001.....	0.9	1.0	0.8	0.8	1.0	0.7	0.9	1.0	0.8	0.9	1.0	0.8	1.1	1.2	0.9
2000.....	0.9	1.1	0.8	0.9	1.1	0.7	0.9	1.0	0.8	0.9	1.0	0.8	1.1	1.3	0.9
1999.....	0.9	1.1	0.9	0.9	0.9	0.8	1.0	1.0	0.8	0.9	1.0	0.8	1.1	1.2	0.9
Down's syndrome (Q90)															
2004.....	0.3	0.3	0.2	0.1	0.2	0.1	0.3	0.3	0.3	0.3	0.3	0.3	0.2	0.3	0.2
2003.....	0.3	0.3	0.3	0.2	0.2	0.2	0.3	0.3	0.3	0.3	0.3	0.3	0.2	0.3	0.2
2002.....	0.3	0.3	0.3	0.2	0.2	0.1	0.3	0.3	0.3	0.3	0.3	0.3	0.2	0.3	0.2
2001.....	0.3	0.3	0.3	0.1	0.2	0.1	0.3	0.3	0.3	0.3	0.3	0.3	0.2	0.2	0.2
2000.....	0.3	0.3	0.3	0.2	0.2	0.2	0.2	0.3	0.2	0.2	0.3	0.3	0.2	0.2	0.3
1999.....	0.2	0.2	0.2	0.1	*	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
All other chromosomal abnormalities, not elsewhere classified (Q91-Q99)															
2004.....	0.4	0.3	0.4	0.3	0.3	0.4	0.4	0.3	0.4	0.3	0.3	0.4	0.4	0.4	0.5
2003.....	0.4	0.3	0.4	0.4	0.3	0.4	0.4	0.3	0.4	0.3	0.3	0.4	0.4	0.3	0.5
2002.....	0.4	0.3	0.4	0.3	0.3	0.4	0.4	0.3	0.4	0.4	0.3	0.4	0.4	0.3	0.5
2001.....	0.3	0.3	0.4	0.4	0.4	0.4	0.3	0.3	0.4	0.3	0.3	0.4	0.4	0.3	0.4
2000.....	0.4	0.3	0.4	0.3	0.3	0.4	0.4	0.3	0.4	0.3	0.3	0.4	0.4	0.4	0.5
1999.....	0.3	0.3	0.4	0.3	0.3	0.4	0.4	0.3	0.4	0.3	0.3	0.4	0.4	0.3	0.5
XVIII. Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)															
2004.....	10.0	10.7	9.0	5.3	6.1	4.5	10.3	11.0	9.4	9.9	10.5	9.0	14.2	16.1	12.3
2003.....	10.6	11.4	9.5	5.6	6.6	4.5	11.0	11.8	9.9	10.6	11.3	9.6	15.0	17.6	12.8
2002.....	10.3	11.4	9.0	5.5	6.6	4.5	10.7	11.8	9.3	10.2	11.1	9.0	14.8	18.2	12.0
2001.....	11.3	12.7	9.8	7.9	9.9	5.8	11.5	12.8	10.0	11.0	12.2	9.7	15.4	18.0	13.1
2000.....	11.4	13.1	9.7	7.8	10.3	5.5	11.6	13.2	10.0	11.0	12.5	9.4	16.7	20.0	13.9
1999.....	9.7	11.0	8.4	5.2	6.2	4.3	10.0	11.3	8.7	9.3	10.4	8.1	15.6	19.3	12.7
Senility (R54)															
2004.....	1.6	1.3	1.7	0.7	0.5	0.8	1.6	1.3	1.7	1.7	1.4	1.8	1.3	1.1	1.4
2003.....	1.6	1.2	1.7	0.8	0.4	0.9	1.6	1.3	1.7	1.7	1.3	1.8	1.2	0.9	1.2
2002.....	1.5	1.2	1.6	0.5	0.5	0.5	1.5	1.2	1.6	1.5	1.3	1.7	1.2	1.1	1.2
2001.....	1.5	1.2	1.6	0.7	0.6	0.7	1.5	1.3	1.6	1.6	1.3	1.6	1.3	1.3	1.3
2000.....	1.4	1.1	1.5	0.6	*	0.6	1.4	1.2	1.5	1.4	1.2	1.5	1.4	1.0	1.5
1999.....	1.2	1.0	1.2	0.5	*	0.7	1.2	1.0	1.2	1.2	1.1	1.3	1.0	1.1	1.0
Ill-defined and unknown causes of mortality (R95-R99)															
2004.....	5.2	6.3	4.1	3.3	4.2	2.3	5.4	6.5	4.3	5.1	6.1	4.0	7.8	9.5	6.3
2003.....	5.9	7.2	4.7	3.4	4.5	2.3	6.2	7.4	4.9	5.9	7.1	4.7	8.9	11.3	7.0
2002.....	5.9	7.2	4.6	3.6	4.7	2.6	6.1	7.5	4.8	5.8	7.0	4.6	9.1	11.8	6.8
2001.....	7.1	8.7	5.6	5.9	8.1	3.8	7.2	8.6	5.7	6.8	8.2	5.4	9.7	11.8	7.9
2000.....	7.4	9.1	5.8	6.0	8.3	3.8	7.4	9.1	5.9	7.0	8.5	5.5	11.0	14.0	8.6
1999.....	6.0	7.3	4.8	3.5	4.6	2.5	6.2	7.5	5.0	5.6	6.7	4.5	10.3	13.4	7.8

Worktable Orig13. Age-adjusted death rates for 358 selected causes, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), and year	All origins 1/			Hispanic			Non-Hispanic 2/			Non-Hispanic white			Non-Hispanic black		
	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female
Sudden infant death syndrome (R95)															
2004.....	0.8	0.9	0.6	0.4	0.5	0.4	0.9	1.0	0.7	0.7	0.8	0.6	1.6	1.8	1.3
2003.....	0.7	0.9	0.6	0.5	0.6	0.4	0.8	0.9	0.7	0.7	0.8	0.5	1.5	1.5	1.4
2002.....	0.8	0.9	0.7	0.5	0.6	0.4	0.9	1.0	0.7	0.7	0.8	0.6	1.5	1.6	1.3
2001.....	0.8	0.9	0.6	0.4	0.5	0.4	0.8	1.0	0.7	0.7	0.8	0.6	1.7	1.8	1.5
2000.....	0.9	1.1	0.8	0.5	0.6	0.4	1.0	1.2	0.8	0.8	0.9	0.7	1.9	2.2	1.6
1999.....	1.0	1.1	0.8	0.6	0.6	0.5	1.1	1.2	0.9	0.8	1.0	0.7	2.0	2.2	1.8
Other ill-defined and unknown causes of mortality (R96-R99)															
2004.....	4.4	5.4	3.5	2.8	3.7	1.9	4.5	5.5	3.6	4.4	5.3	3.4	6.2	7.7	5.0
2003.....	5.2	6.3	4.0	2.9	3.9	1.9	5.3	6.5	4.2	5.2	6.3	4.1	7.4	9.7	5.6
2002.....	5.1	6.3	4.0	3.1	4.1	2.3	5.3	6.5	4.1	5.0	6.1	4.0	7.6	10.1	5.5
2001.....	6.3	7.8	4.9	5.5	7.6	3.4	6.3	7.6	5.0	6.1	7.4	4.8	8.1	10.0	6.5
2000.....	6.5	8.1	5.0	5.5	7.7	3.4	6.4	7.9	5.0	6.2	7.6	4.8	9.1	11.8	7.0
1999.....	5.1	6.2	4.0	2.9	3.9	2.0	5.2	6.3	4.1	4.8	5.8	3.8	8.3	11.1	6.1
All other symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R53,R55-R94)															
2004.....	3.2	3.1	3.2	1.4	1.4	1.3	3.3	3.2	3.3	3.2	3.0	3.3	5.1	5.6	4.6
2003.....	3.2	3.0	3.1	1.5	1.6	1.4	3.3	3.1	3.2	3.1	2.9	3.1	5.0	5.4	4.6
2002.....	2.9	3.0	2.8	1.3	1.3	1.3	3.0	3.1	2.9	2.9	2.9	2.8	4.6	5.3	3.9
2001.....	2.8	2.8	2.7	1.3	1.2	1.3	2.9	2.9	2.7	2.7	2.7	2.6	4.4	4.9	3.8
2000.....	2.7	2.8	2.5	1.2	1.5	1.0	2.8	2.9	2.6	2.6	2.7	2.4	4.3	4.9	3.8
1999.....	2.6	2.7	2.4	1.2	1.3	1.1	2.6	2.7	2.5	2.5	2.6	2.3	4.3	4.8	3.8
XX. External causes of mortality (*U01-*U03,V01-Y89)															
2004.....	57.3	82.7	33.7	44.6	66.6	22.0	58.6	84.4	35.0	58.3	82.5	35.8	67.1	106.3	33.3
2003.....	57.0	82.9	33.0	45.6	69.2	21.8	57.9	84.0	34.1	57.1	81.5	34.7	67.7	107.5	33.2
2002.....	56.7	82.7	32.6	45.4	69.2	21.6	57.6	83.8	33.7	56.5	80.9	34.0	68.7	110.0	33.4
2001 3/.....	56.1	82.6	31.8	46.4	70.2	22.7	56.8	83.4	32.7	55.4	80.3	32.7	69.6	110.7	34.4
2000.....	53.9	79.5	30.5	45.1	68.8	21.7	54.5	80.2	31.4	52.6	76.3	31.2	69.4	111.4	33.7
1999.....	54.3	80.1	31.0	45.6	70.5	21.3	54.8	80.5	31.8	52.7	76.3	31.4	71.3	114.2	35.2
Accidents (unintentional injuries) (V01-X59,Y85-Y86)															
2004.....	37.7	52.1	24.5	29.8	43.0	16.4	38.5	53.0	25.4	39.7	54.2	26.3	37.1	55.5	22.2
2003.....	37.3	51.8	24.1	30.6	44.9	16.3	37.8	52.2	24.8	38.8	53.1	25.5	36.8	55.0	22.1
2002.....	36.9	51.5	23.5	30.7	45.6	16.1	37.3	51.8	24.2	38.0	52.3	24.8	37.6	57.2	21.8
2001.....	35.7	50.2	22.5	30.7	44.7	16.8	35.9	50.4	23.0	36.2	50.4	23.3	38.3	57.9	22.4
2000.....	34.9	49.3	22.0	30.1	44.5	16.1	35.2	49.5	22.5	35.3	49.2	22.7	38.4	59.2	21.8
1999.....	35.3	49.8	22.3	30.6	45.8	16.0	35.5	49.9	22.8	35.3	49.1	22.9	40.8	62.4	23.5
Transport accidents (V01-V99,Y85)															
2004.....	16.2	23.1	9.7	15.1	22.1	7.9	16.3	23.2	9.9	16.8	23.6	10.3	16.1	24.9	8.6
2003.....	16.4	23.4	9.8	15.9	23.2	8.4	16.4	23.3	10.0	16.8	23.6	10.3	16.2	25.1	8.8
2002.....	16.7	23.9	9.9	16.0	23.4	8.4	16.7	23.8	10.1	17.1	24.1	10.5	16.2	25.4	8.6
2001.....	16.6	23.7	9.8	16.7	24.6	8.9	16.5	23.6	10.0	16.7	23.7	10.1	16.9	26.1	9.1
2000.....	16.6	23.7	10.0	15.7	23.1	8.3	16.7	23.7	10.2	16.9	23.8	10.5	17.1	26.7	9.0
1999.....	16.6	23.6	10.2	15.4	22.5	8.4	16.7	23.7	10.5	16.9	23.6	10.6	17.5	27.1	9.7
Railway accidents (V05,V15,V80.6,V81.2-V81.9)															
2004.....	0.1	0.3	0.1	0.2	0.4	*	0.1	0.2	0.1	0.1	0.2	0.1	0.2	0.4	*
2003.....	0.1	0.2	0.0	0.2	0.3	*	0.1	0.2	0.0	0.1	0.2	0.0	0.2	0.4	*
2002.....	0.2	0.3	0.0	0.2	0.3	*	0.2	0.3	0.0	0.1	0.3	0.0	0.3	0.5	*
2001.....	0.2	0.3	0.1	0.3	0.5	*	0.2	0.3	0.0	0.2	0.2	0.0	0.2	0.4	*
2000.....	0.2	0.3	0.1	0.2	0.4	*	0.2	0.3	0.0	0.2	0.3	0.0	0.3	0.5	*
1999.....	0.2	0.3	0.0	0.2	0.4	*	0.2	0.3	0.1	0.2	0.3	0.1	0.2	0.4	*

Worktable Orig13. Age-adjusted death rates for 358 selected causes, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), and year	All origins 1/			Hispanic			Non-Hispanic 2/			Non-Hispanic white			Non-Hispanic black		
	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female
Occupant of motor vehicle involved in collision with other (non-motorized) road vehicle, streetcar, animal or pedestrian (V40-V41,V46-V47,V50-V51,V56-V57,V60-V61,V66-V67,V70-V71,V76-V77)															
2004.....	1.6	2.3	0.8	1.4	2.2	0.6	1.6	2.3	0.9	1.6	2.4	0.9	1.7	2.7	0.9
2003.....	1.7	2.5	0.9	1.5	2.3	0.6	1.7	2.5	0.9	1.8	2.6	1.0	1.7	2.7	0.7
2002.....	1.7	2.6	0.9	1.6	2.4	0.7	1.7	2.6	1.0	1.8	2.6	1.0	1.9	3.2	0.9
2001.....	1.5	2.2	0.8	1.5	2.3	0.7	1.5	2.2	0.8	1.4	2.2	0.8	1.8	2.9	0.8
2000.....	1.4	2.0	0.7	1.2	1.7	0.6	1.4	2.1	0.8	1.4	2.0	0.8	1.7	2.7	0.8
1999.....	1.3	2.0	0.7	1.2	1.8	0.5	1.4	2.0	0.8	1.4	2.0	0.7	1.7	2.7	0.9
Occupant of car, pickup truck or van involved in collision with other motor vehicle (V42-V44,V49,V52-V54,V59)															
2004.....	3.6	4.5	2.8	3.6	4.6	2.5	3.5	4.4	2.8	3.6	4.5	2.9	3.3	4.5	2.4
2003.....	3.7	4.6	2.8	3.7	5.0	2.3	3.6	4.5	2.9	3.7	4.5	3.0	3.5	4.7	2.6
2002.....	3.9	5.0	2.9	4.1	5.7	2.4	3.8	4.8	3.0	3.9	4.8	3.1	3.6	4.9	2.5
2001.....	3.7	4.7	2.8	3.8	5.3	2.4	3.6	4.6	2.8	3.7	4.6	3.0	3.6	5.1	2.5
2000.....	3.7	4.6	2.9	3.4	4.7	2.2	3.7	4.6	2.9	3.8	4.7	3.0	3.6	5.0	2.4
1999.....	3.7	4.6	2.9	3.2	4.2	2.2	3.7	4.6	2.9	3.8	4.6	3.1	3.7	5.2	2.5
Occupant of heavy transport vehicle or bus involved in collision with other motor vehicle (V62-V64,V69,V72-V74,V79)															
2004.....	0.1	0.1	0.0	0.1	0.1	*	0.1	0.1	0.0	0.1	0.1	*	0.1	0.2	*
2003.....	0.1	0.1	0.0	0.1	0.1	*	0.1	0.1	0.0	0.1	0.1	0.0	0.1	0.2	*
2002.....	0.1	0.1	0.0	0.1	0.2	*	0.1	0.1	0.0	0.1	0.1	*	0.1	0.2	*
2001.....	0.1	0.1	0.0	*	*	*	0.1	0.1	0.0	0.1	0.1	0.0	*	*	*
2000.....	0.1	0.1	0.0	*	*	*	0.1	0.1	0.0	0.1	0.1	*	0.1	0.2	*
1999.....	0.1	0.1	0.0	*	*	*	0.1	0.1	0.0	0.1	0.1	0.0	0.1	*	*
Occupant of motor vehicle involved in noncollision accident (V48,V58,V68,V78)															
2004.....	1.4	1.9	0.8	1.5	2.2	0.7	1.3	1.9	0.8	1.4	1.9	0.8	1.2	1.8	0.7
2003.....	1.5	2.0	0.9	1.6	2.2	1.0	1.5	2.0	0.9	1.5	2.1	0.9	1.2	1.8	0.7
2002.....	1.5	2.1	0.9	1.6	2.3	0.9	1.5	2.1	0.9	1.5	2.2	0.9	1.2	1.9	0.6
2001.....	1.4	1.9	0.8	1.6	2.2	0.8	1.3	1.9	0.8	1.4	1.9	0.8	1.1	1.8	0.6
2000.....	1.3	1.9	0.7	1.4	2.0	0.7	1.3	1.9	0.8	1.4	1.9	0.8	1.2	1.8	0.6
1999.....	1.3	1.9	0.8	1.2	1.8	0.6	1.3	1.9	0.8	1.3	1.9	0.8	1.2	1.8	0.7
Occupant of special-use motor vehicle involved in any accident (V83-V86)															
2004.....	0.4	0.7	0.1	0.1	0.2	*	0.4	0.7	0.1	0.5	0.8	0.1	0.2	0.3	*
2003.....	0.4	0.7	0.1	0.1	0.3	*	0.4	0.8	0.1	0.5	0.9	0.1	0.1	0.2	*
2002.....	0.3	0.6	0.1	0.2	0.3	*	0.4	0.7	0.1	0.4	0.8	0.1	0.1	0.1	*
2001.....	0.4	0.7	0.1	0.1	0.2	*	0.4	0.7	0.1	0.5	0.8	0.1	0.1	0.2	*
2000.....	0.4	0.7	0.1	0.2	0.3	*	0.4	0.7	0.1	0.5	0.8	0.1	0.1	0.3	*
1999.....	0.4	0.7	0.1	0.1	0.2	*	0.4	0.7	0.1	0.5	0.9	0.1	0.1	0.3	*
Other and unspecified motor vehicle accidents (V09.0,V09.2,V80.3-V80.5,V82.0-V82.1,V87.0-V87.5,V87.7-V87.8,V88.0-V88.5,V88.7-V88.8,V89.0,V89.2)															
2004.....	5.4	7.3	3.7	4.8	6.8	2.6	5.5	7.4	3.8	5.6	7.4	3.9	5.7	8.5	3.4
2003.....	5.2	7.1	3.5	5.0	7.0	2.8	5.3	7.1	3.6	5.3	7.1	3.6	5.6	8.2	3.4
2002.....	5.4	7.4	3.6	4.6	6.3	2.8	5.6	7.5	3.8	5.7	7.6	3.9	5.5	8.2	3.3
2001.....	5.7	7.9	3.8	4.8	6.9	2.7	5.8	8.0	3.9	5.9	8.0	4.0	6.0	8.9	3.5
2000.....	6.0	8.2	4.1	5.5	7.8	3.2	6.0	8.2	4.1	6.1	8.2	4.2	6.4	9.5	3.8
1999.....	6.0	8.1	4.1	5.7	8.0	3.5	6.0	8.1	4.2	6.1	8.1	4.3	6.3	9.3	4.0
Streetcar accidents (V82.2-V82.9)															
2004.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig13. Age-adjusted death rates for 358 selected causes, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Table with 17 columns: Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), and year; All origins 1/ (Both Sexes, Male, Female); Hispanic (Both Sexes, Male, Female); Non-Hispanic 2/ (Both Sexes, Male, Female); Non-Hispanic white (Both Sexes, Male, Female); Non-Hispanic black (Both Sexes, Male, Female). Rows include categories like Other and unspecified land transport accidents, Water transport accidents, Air and space transport accidents, Other and unspecified transport accidents and their sequelae, Falls, Fall on same level, and Fall from one level to another.

Worktable Orig13. Age-adjusted death rates for 358 selected causes, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), and year	All origins 1/			Hispanic			Non-Hispanic 2/			Non-Hispanic white			Non-Hispanic black		
	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female
Unspecified fall (W19)															
2004.....	3.1	3.8	2.5	2.2	2.7	1.8	3.1	3.9	2.6	3.3	4.0	2.8	1.8	2.8	1.2
2003.....	2.8	3.4	2.4	1.8	2.3	1.3	2.9	3.5	2.4	3.0	3.6	2.5	1.8	2.3	1.5
2002.....	2.6	3.2	2.2	1.9	2.4	1.5	2.7	3.2	2.3	2.8	3.3	2.4	1.6	2.4	1.1
2001.....	2.7	3.5	2.2	1.8	2.1	1.5	2.7	3.5	2.2	2.8	3.6	2.4	1.7	2.6	1.2
2000.....	2.6	3.3	2.1	2.1	2.7	1.6	2.6	3.3	2.1	2.7	3.4	2.2	1.7	2.5	1.2
1999.....	2.9	3.7	2.3	2.4	3.4	1.6	2.9	3.7	2.3	3.0	3.7	2.4	2.1	3.1	1.4
Accidental discharge of firearms (W32-W34)															
2004.....	0.2	0.4	0.0	0.1	0.2	*	0.2	0.4	0.1	0.2	0.4	0.1	0.3	0.6	*
2003.....	0.2	0.5	0.1	0.2	0.3	*	0.3	0.5	0.1	0.3	0.5	0.1	0.3	0.6	*
2002.....	0.3	0.4	0.1	0.2	0.2	*	0.3	0.5	0.1	0.3	0.5	0.1	0.4	0.8	*
2001.....	0.3	0.5	0.1	0.2	0.4	*	0.3	0.5	0.1	0.3	0.5	0.1	0.3	0.6	*
2000.....	0.3	0.5	0.1	0.2	0.2	*	0.3	0.5	0.1	0.3	0.5	0.1	0.3	0.7	0.1
1999.....	0.3	0.5	0.1	0.2	0.3	*	0.3	0.5	0.1	0.3	0.5	0.1	0.4	0.8	0.1
Accidental explosion (W39-W40)															
2004.....	0.1	0.1	0.0	*	*	*	0.1	0.1	0.0	0.1	0.1	0.0	*	*	*
2003.....	0.1	0.1	0.0	*	*	*	0.1	0.1	0.0	0.1	0.1	*	*	*	*
2002.....	0.0	0.1	*	*	*	*	0.0	0.1	*	0.1	0.1	*	*	*	*
2001.....	0.1	0.1	0.0	*	*	*	0.1	0.1	0.0	0.1	0.1	0.0	*	*	*
2000.....	0.1	0.1	0.0	*	*	*	0.1	0.1	0.0	0.1	0.1	0.0	*	*	*
1999.....	0.1	0.1	0.0	*	*	*	0.1	0.1	0.0	0.1	0.1	0.0	*	*	*
Accidental drowning and submersion (W65-W74)															
2004.....	1.1	1.8	0.5	1.0	1.6	0.3	1.1	1.8	0.5	1.1	1.7	0.5	1.4	2.3	0.5
2003.....	1.1	1.8	0.5	1.1	1.8	0.3	1.1	1.8	0.5	1.1	1.7	0.5	1.2	2.2	0.4
2002.....	1.2	2.0	0.5	1.1	1.9	0.3	1.2	1.9	0.5	1.1	1.7	0.5	1.6	2.8	0.5
2001.....	1.1	1.8	0.5	1.0	1.6	0.3	1.2	1.8	0.5	1.1	1.7	0.5	1.5	2.7	0.5
2000.....	1.2	2.0	0.5	1.1	1.7	0.4	1.3	2.0	0.5	1.2	1.8	0.5	1.6	2.7	0.6
1999.....	1.3	2.0	0.5	1.2	2.0	0.3	1.2	2.0	0.5	1.1	1.8	0.5	1.7	3.0	0.5
Accidental inhalation and ingestion of food or other objects causing obstruction of respiratory tract (W78-W80)															
2004.....	1.5	1.9	1.2	0.9	1.2	0.7	1.5	1.9	1.3	1.5	1.9	1.3	1.9	2.5	1.6
2003.....	1.5	1.9	1.2	0.8	1.2	0.6	1.5	1.9	1.2	1.5	1.9	1.2	1.8	2.4	1.4
2002.....	1.4	1.8	1.2	0.8	1.1	0.6	1.5	1.9	1.2	1.4	1.8	1.1	2.1	2.8	1.7
2001.....	1.5	1.9	1.2	0.9	1.2	0.7	1.5	1.9	1.2	1.5	1.8	1.2	2.2	3.2	1.6
2000.....	1.6	2.0	1.3	1.0	1.4	0.7	1.6	2.0	1.3	1.5	1.9	1.3	2.2	3.0	1.7
1999.....	1.4	1.8	1.2	1.0	1.2	0.8	1.5	1.8	1.2	1.4	1.7	1.2	2.2	2.9	1.7
Other accidental and unspecified threats to breathing (W75-W77,W81-W84)															
2004.....	0.5	0.6	0.3	0.3	0.4	0.1	0.5	0.7	0.4	0.5	0.7	0.3	0.6	0.8	0.5
2003.....	0.4	0.6	0.3	0.3	0.4	0.2	0.5	0.7	0.3	0.5	0.7	0.3	0.6	0.8	0.4
2002.....	0.5	0.7	0.3	0.2	0.3	0.1	0.5	0.7	0.3	0.5	0.7	0.3	0.7	0.8	0.6
2001.....	0.5	0.6	0.3	0.2	0.3	0.2	0.5	0.7	0.3	0.5	0.7	0.3	0.6	0.8	0.5
2000.....	0.5	0.7	0.3	0.3	0.4	0.2	0.5	0.7	0.3	0.5	0.7	0.3	0.6	0.9	0.5
1999.....	0.6	0.8	0.4	0.4	0.5	0.2	0.6	0.9	0.4	0.6	0.8	0.4	0.8	1.0	0.5
Accidental exposure to electric current (W85-W87)															
2004.....	0.1	0.2	0.0	0.1	0.3	*	0.1	0.2	*	0.1	0.3	*	0.1	0.2	*
2003.....	0.1	0.2	*	0.1	0.2	*	0.1	0.2	*	0.1	0.3	*	0.1	0.2	*
2002.....	0.2	0.3	0.0	0.2	0.3	*	0.1	0.3	0.0	0.1	0.3	0.0	0.1	0.2	*
2001.....	0.1	0.3	*	0.2	0.4	*	0.1	0.3	*	0.2	0.3	*	0.1	0.2	*
2000.....	0.1	0.3	0.0	0.1	0.2	*	0.1	0.3	*	0.2	0.3	*	0.1	0.2	*
1999.....	0.2	0.3	0.0	0.2	0.3	*	0.2	0.3	*	0.2	0.3	*	0.1	0.2	*

Worktable Orig13. Age-adjusted death rates for 358 selected causes, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Table with columns for Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), and year; All origins 1/; Hispanic; Non-Hispanic 2/; Non-Hispanic white; Non-Hispanic black. Rows include Accidental exposure to radiation (W88-W91), Accidental exposure to smoke, fire and flames (X00-X09), Accidental contact with heat and hot substances (X10-X19), Victim of lightning (X33), Earthquake, volcanic eruption, avalanche, landslide and other earth movements (X34-X36), Cataclysmic storm and flood (X37-X38), and Accidental poisoning and exposure to noxious substances (X40-X49).

Worktable Orig13. Age-adjusted death rates for 358 selected causes, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Table with 17 columns: Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), and year; All origins 1/ (Both Sexes, Male, Female); Hispanic (Both Sexes, Male, Female); Non-Hispanic 2/ (Both Sexes, Male, Female); Non-Hispanic white (Both Sexes, Male, Female); Non-Hispanic black (Both Sexes, Male, Female). Rows include categories like 'Accidental poisoning by and exposure to drugs and other biological substances (X40-X44)', 'Accidental poisoning by and exposure to other and unspecified solid or liquids substances (X45-X46, X48-X49)', 'Accidental poisoning by and exposure to other gases and vapors (X47)', 'All other and unspecified accidents and their sequelae (W20-W31, W35-W38, W41-W64, W92-W99, X20-X32, X39, X50-X59, Y86)', 'Intentional self-harm (suicide) (*U03, X60-X84, Y87.0)', 'Intentional self-poisoning (suicide) by and exposure to drugs and other biological substances (X60-X64)', and 'Intentional self-poisoning (suicide) by and exposure to other and unspecified solid or liquid substances and their vapors (X65-X66, X68-X69)'. Values are death rates per 100,000 population, with asterisks indicating statistical significance.

Worktable Orig13. Age-adjusted death rates for 358 selected causes, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), and year	All origins 1/			Hispanic			Non-Hispanic 2/			Non-Hispanic white			Non-Hispanic black		
	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female
Intentional self-poisoning (suicide) by and exposure to other gases and vapors (X67)															
2004.....	0.5	0.7	0.2	0.1	0.2	*	0.5	0.8	0.2	0.6	1.0	0.2	0.1	0.2	*
2003.....	0.5	0.8	0.2	0.1	0.2	*	0.5	0.9	0.2	0.6	1.0	0.3	0.1	0.2	*
2002.....	0.5	0.8	0.2	0.1	0.2	*	0.5	0.9	0.2	0.6	1.1	0.3	0.1	0.2	*
2001.....	0.5	0.8	0.2	0.1	0.2	*	0.6	0.9	0.2	0.6	1.0	0.3	0.1	0.2	*
2000.....	0.5	0.8	0.2	0.1	0.2	*	0.6	0.9	0.2	0.6	1.0	0.3	0.1	0.2	*
1999.....	0.5	0.9	0.3	0.1	0.2	*	0.6	1.0	0.3	0.7	1.1	0.3	0.1	0.2	*
Intentional self-harm (suicide) by hanging, strangulation and suffocation (X70)															
2004.....	2.5	4.1	0.9	2.1	3.6	0.7	2.5	4.2	0.9	2.7	4.5	1.0	1.4	2.4	0.4
2003.....	2.3	3.9	0.8	1.9	3.3	0.5	2.3	4.0	0.8	2.5	4.2	0.8	1.3	2.3	0.4
2002.....	2.2	3.7	0.8	1.9	3.2	0.5	2.3	3.9	0.8	2.5	4.2	0.8	1.1	2.0	0.3
2001.....	2.2	3.7	0.7	2.0	3.6	0.5	2.2	3.8	0.7	2.4	4.1	0.7	1.1	2.0	0.3
2000.....	2.0	3.4	0.7	1.9	3.4	0.5	2.0	3.5	0.7	2.1	3.7	0.7	1.2	2.2	0.3
1999.....	1.9	3.3	0.6	1.8	3.3	0.4	2.0	3.3	0.7	2.0	3.5	0.7	1.0	2.0	0.2
Intentional self-harm (suicide) by discharge of firearms (X72-X74)															
2004.....	5.6	10.3	1.5	2.4	4.4	0.5	6.0	11.1	1.6	6.8	12.3	1.9	2.8	5.6	0.5
2003.....	5.8	10.7	1.4	2.4	4.5	0.3	6.2	11.5	1.5	6.9	12.8	1.8	2.8	5.5	0.6
2002.....	5.9	11.1	1.4	2.5	4.8	0.4	6.3	11.8	1.5	7.0	13.0	1.8	3.0	6.0	0.5
2001.....	5.9	11.0	1.4	2.5	4.8	0.4	6.3	11.7	1.6	7.0	12.9	1.8	3.1	6.1	0.7
2000.....	5.9	11.0	1.5	2.7	5.1	0.5	6.3	11.6	1.6	6.9	12.8	1.8	3.2	6.1	0.7
1999.....	6.0	11.1	1.5	2.8	5.3	0.6	6.3	11.8	1.6	6.9	12.8	1.8	3.3	6.7	0.6
Intentional self-harm (suicide) by jumping from a high place (X80)															
2004.....	0.2	0.3	0.1	0.2	0.3	*	0.2	0.3	0.1	0.2	0.3	0.1	0.2	0.3	*
2003.....	0.2	0.3	0.1	0.2	0.3	*	0.2	0.4	0.1	0.2	0.4	0.1	0.2	0.3	*
2002.....	0.2	0.4	0.1	0.2	0.3	*	0.3	0.4	0.1	0.3	0.4	0.2	0.2	0.3	*
2001.....	0.2	0.3	0.1	0.2	0.3	*	0.2	0.3	0.1	0.2	0.3	0.1	0.2	0.3	*
2000.....	0.2	0.3	0.1	0.2	0.3	*	0.2	0.3	0.1	0.2	0.3	0.1	0.2	0.3	*
1999.....	0.3	0.4	0.1	0.2	0.3	*	0.3	0.4	0.1	0.3	0.4	0.1	0.2	0.4	*
Intentional self-harm (suicide) by all other and unspecified means and their sequelae (*U03,X71,X75-X79,X81-X84,X87.0)															
2004.....	0.6	1.0	0.3	0.5	0.8	0.1	0.7	1.0	0.3	0.7	1.0	0.3	0.5	0.8	0.2
2003.....	0.6	0.9	0.3	0.4	0.7	*	0.6	1.0	0.3	0.6	1.0	0.3	0.5	0.8	0.2
2002.....	0.6	1.0	0.3	0.5	0.8	0.2	0.6	1.0	0.3	0.7	1.0	0.3	0.5	0.9	0.2
2001 3/.....	0.6	0.9	0.3	0.4	0.8	0.2	0.6	1.0	0.3	0.6	1.0	0.3	0.5	0.7	0.3
2000.....	0.6	0.9	0.3	0.5	0.8	*	0.6	0.9	0.3	0.6	0.9	0.3	0.5	0.8	0.2
1999.....	0.6	0.9	0.3	0.4	0.6	0.2	0.6	0.9	0.3	0.6	0.9	0.3	0.5	0.8	0.3
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)															
2004.....	5.9	9.2	2.5	7.2	11.5	2.3	5.6	8.6	2.6	2.7	3.6	1.8	20.7	36.2	6.5
2003.....	6.0	9.4	2.6	7.7	12.1	2.7	5.7	8.9	2.6	2.7	3.6	1.8	21.7	37.9	6.6
2002.....	6.1	9.4	2.8	7.3	11.6	2.5	5.8	8.9	2.8	2.8	3.7	1.9	21.6	37.4	7.0
2001 3/.....	7.1	10.8	3.3	8.3	12.9	3.1	6.8	10.3	3.3	4.0	5.6	2.5	21.7	37.2	7.5
2000.....	5.9	9.0	2.8	7.5	11.8	2.8	5.6	8.5	2.7	2.8	3.6	1.9	21.1	36.4	7.2
1999.....	6.0	9.1	2.9	7.6	12.2	2.6	5.7	8.5	2.9	2.9	3.8	2.0	20.7	35.2	7.5
Assault (homicide) by drugs, biological substances and other and unspecified noxious substances (*U01.6-*U01.7,X85-X90)															
2004.....	0.0	0.0	0.0	*	*	*	0.0	0.0	0.0	0.0	0.0	0.0	0.1	*	*
2003.....	0.0	0.0	0.0	*	*	*	0.0	0.0	0.0	0.0	0.0	*	0.1	*	*
2002.....	0.0	0.0	0.0	*	*	*	0.0	0.0	0.0	0.0	0.0	*	*	*	*
2001.....	0.0	0.0	0.0	*	*	*	0.0	0.0	0.0	0.0	0.0	*	*	*	*
2000.....	0.0	0.0	0.0	*	*	*	0.0	0.0	0.0	0.0	*	0.0	*	*	*
1999.....	0.0	0.0	0.0	*	*	*	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*

Worktable Orig13. Age-adjusted death rates for 358 selected causes, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), and year	All origins 1/			Hispanic			Non-Hispanic 2/			Non-Hispanic white			Non-Hispanic black		
	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female
Assault (homicide) by hanging, strangulation and suffocation (X91)															
2004.....	0.2	0.1	0.3	0.2	0.2	0.3	0.2	0.1	0.3	0.2	0.1	0.2	0.6	0.4	0.7
2003.....	0.2	0.1	0.3	0.3	0.3	0.3	0.2	0.1	0.3	0.2	0.1	0.2	0.5	0.4	0.7
2002.....	0.2	0.2	0.3	0.2	0.2	0.2	0.2	0.2	0.3	0.2	0.1	0.2	0.6	0.4	0.7
2001.....	0.2	0.2	0.3	0.2	0.2	0.3	0.2	0.2	0.3	0.2	0.1	0.2	0.6	0.4	0.8
2000.....	0.2	0.2	0.3	0.2	0.2	0.3	0.2	0.2	0.3	0.2	0.2	0.2	0.6	0.3	0.8
1999.....	0.2	0.2	0.3	0.3	0.3	0.2	0.2	0.2	0.3	0.2	0.1	0.3	0.6	0.4	0.8
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)															
2004.....	4.0	6.6	1.1	4.8	8.1	1.0	3.7	6.3	1.2	1.5	2.1	0.8	15.6	29.0	3.0
2003.....	4.1	6.8	1.2	5.1	8.5	1.2	3.9	6.5	1.2	1.5	2.1	0.8	16.3	30.1	3.3
2002.....	4.1	6.8	1.3	4.8	8.1	1.2	3.9	6.5	1.3	1.6	2.2	0.9	16.1	29.5	3.6
2001.....	3.9	6.6	1.3	4.9	8.2	1.2	3.7	6.2	1.3	1.6	2.2	0.9	15.2	28.2	3.1
2000.....	3.8	6.3	1.2	4.8	8.0	1.3	3.6	6.0	1.3	1.5	2.0	0.9	15.0	27.8	3.2
1999.....	3.8	6.3	1.3	5.0	8.2	1.4	3.6	6.0	1.3	1.5	2.2	0.9	14.8	26.9	3.6
Assault (homicide) by sharp object (X99)															
2004.....	0.7	1.0	0.4	1.0	1.5	0.5	0.6	0.9	0.4	0.4	0.5	0.3	2.2	3.4	1.2
2003.....	0.7	1.0	0.4	1.1	1.5	0.5	0.7	0.9	0.4	0.3	0.4	0.2	2.3	3.6	1.2
2002.....	0.7	1.0	0.4	1.1	1.6	0.5	0.7	0.9	0.4	0.4	0.5	0.2	2.3	3.4	1.2
2001.....	0.7	1.0	0.4	1.1	1.6	0.5	0.6	0.8	0.4	0.4	0.5	0.3	2.1	3.2	1.2
2000.....	0.6	0.9	0.4	1.1	1.7	0.5	0.6	0.8	0.3	0.3	0.4	0.2	2.0	3.0	1.1
1999.....	0.7	1.0	0.4	1.0	1.5	0.4	0.6	0.9	0.4	0.4	0.5	0.2	2.1	3.3	1.0
Assault (homicide) by blunt object (Y00)															
2004.....	0.0	0.0	0.0	*	*	*	0.0	0.1	0.0	0.0	0.0	*	0.1	*	*
2003.....	0.0	0.1	0.0	*	*	*	0.0	0.1	0.0	0.0	0.0	0.0	0.1	0.2	*
2002.....	0.1	0.1	0.0	0.1	0.1	*	0.1	0.1	0.0	0.1	0.1	0.0	0.1	0.2	*
2001.....	0.1	0.1	0.1	0.1	*	*	0.1	0.1	0.1	0.0	0.1	0.0	0.2	0.3	*
2000.....	0.1	0.1	0.1	0.1	*	*	0.1	0.1	0.1	0.1	0.1	0.0	0.2	0.3	0.1
1999.....	0.1	0.1	0.0	*	*	*	0.1	0.1	0.0	0.1	0.1	0.0	0.2	0.3	0.1
Assault (homicide) by bodily force (Y04)															
2004.....	0.0	0.0	*	*	*	*	0.0	0.0	*	0.0	0.0	*	0.1	*	*
2003.....	0.0	0.1	*	0.1	0.1	*	0.0	0.1	*	0.0	0.0	*	0.1	0.2	*
2002.....	0.0	0.1	*	*	*	*	0.0	0.1	*	0.0	0.0	*	0.1	0.2	*
2001.....	0.1	0.1	0.0	*	*	*	0.1	0.1	0.0	0.0	0.1	0.0	0.1	0.2	*
2000.....	0.1	0.1	*	*	*	*	0.1	0.1	*	0.0	0.1	*	0.1	0.2	*
1999.....	0.1	0.1	0.0	*	*	*	0.1	0.1	0.0	0.0	0.1	*	0.2	0.3	*
Sexual assault (homicide) by bodily force (Y05)															
2004.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Neglect, abandonment and other maltreatment syndromes (Y06-Y07)															
2004.....	0.1	0.1	0.1	0.1	0.1	*	0.1	0.1	0.0	0.0	0.0	0.0	0.1	0.1	0.1
2003.....	0.1	0.1	0.1	0.1	0.1	*	0.1	0.1	0.1	0.0	0.1	0.0	0.2	0.2	0.1
2002.....	0.1	0.1	0.1	0.1	0.1	*	0.1	0.1	0.1	0.0	0.0	0.1	0.2	0.2	0.2
2001.....	0.1	0.1	0.1	0.1	0.1	*	0.1	0.1	0.1	0.1	0.1	0.0	0.2	0.2	0.2
2000.....	0.1	0.1	0.1	0.1	*	*	0.1	0.1	0.1	0.1	0.1	0.0	0.2	0.2	0.2
1999.....	0.1	0.1	0.1	0.0	*	*	0.1	0.1	0.1	0.0	0.1	0.0	0.2	0.2	0.2

Worktable Orig13. Age-adjusted death rates for 358 selected causes, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), and year	All origins 1/			Hispanic			Non-Hispanic 2/			Non-Hispanic white			Non-Hispanic black		
	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female
Assault (homicide) by all other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5,*U01.8-*U01.9,*U02.X92,X96-X98,Y01-Y03,Y08-Y09,Y87.1)															
2004.....	0.9	1.1	0.6	1.0	1.5	0.5	0.8	1.1	0.6	0.6	0.8	0.4	2.0	3.0	1.3
2003.....	0.9	1.2	0.6	1.1	1.6	0.6	0.8	1.2	0.6	0.6	0.8	0.4	2.2	3.3	1.2
2002.....	0.8	1.1	0.5	0.9	1.4	0.5	0.8	1.1	0.5	0.6	0.8	0.4	2.2	3.4	1.2
2001 3/.....	2.0	2.8	1.1	1.8	2.7	0.9	2.0	2.8	1.1	1.7	2.5	0.9	3.3	4.6	2.1
2000.....	1.0	1.4	0.7	1.1	1.6	0.6	1.0	1.3	0.7	0.6	0.8	0.5	3.0	4.5	1.7
1999.....	1.0	1.3	0.7	1.2	1.9	0.6	1.0	1.2	0.7	0.7	0.8	0.5	2.5	3.7	1.6
Event of undetermined intent (Y10-Y34,Y87.2,Y89.9)															
2004.....	1.7	2.2	1.2	0.9	1.2	0.5	1.8	2.3	1.3	1.9	2.4	1.4	1.9	2.8	1.2
2003.....	1.7	2.3	1.2	0.9	1.4	0.4	1.8	2.4	1.3	2.0	2.5	1.4	1.9	2.9	1.0
2002.....	1.7	2.2	1.2	0.8	1.3	0.4	1.8	2.3	1.3	1.8	2.3	1.3	2.1	3.1	1.3
2001.....	1.5	2.0	0.9	0.9	1.3	0.4	1.5	2.1	1.0	1.6	2.1	1.1	1.8	2.9	0.9
2000.....	1.3	1.9	0.8	0.6	0.9	0.3	1.4	2.0	0.9	1.4	1.9	0.9	1.9	3.1	1.0
1999.....	1.4	2.0	0.9	0.7	1.2	0.3	1.5	2.0	0.9	1.4	1.9	0.9	2.0	3.3	1.0
Poisoning by and exposure to drugs and biological substances, undetermined intent (Y10-Y14)															
2004.....	1.1	1.4	0.9	0.4	0.5	0.3	1.2	1.5	1.0	1.3	1.6	1.1	1.1	1.6	0.7
2003.....	1.2	1.5	0.9	0.4	0.6	0.2	1.3	1.7	1.0	1.5	1.8	1.1	1.2	1.8	0.7
2002.....	1.1	1.3	0.8	0.4	0.5	0.3	1.2	1.5	0.9	1.3	1.6	1.0	1.3	1.8	0.9
2001.....	1.0	1.3	0.7	0.4	0.5	0.2	1.0	1.4	0.7	1.1	1.4	0.8	1.1	1.9	0.5
2000.....	0.8	1.2	0.6	0.3	0.4	0.2	0.9	1.3	0.6	1.0	1.3	0.6	1.1	1.7	0.7
1999.....	0.9	1.2	0.6	0.3	0.5	0.2	0.9	1.2	0.6	0.9	1.2	0.7	1.2	2.0	0.6
Poisoning by and exposure to other and unspecified solid or liquid substance, undetermined intent (Y15-Y16,Y18-Y19)															
2004.....	0.0	0.0	*	*	*	*	0.0	0.0	*	0.0	0.0	*	*	*	*
2003.....	0.0	0.0	0.0	*	*	*	0.0	0.0	*	0.0	0.0	*	*	*	*
2002.....	0.0	0.0	*	*	*	*	0.0	0.0	*	0.0	0.0	*	*	*	*
2001.....	0.0	0.0	*	*	*	*	0.0	0.0	*	0.0	0.0	*	*	*	*
2000.....	0.0	0.0	*	*	*	*	0.0	0.0	*	0.0	0.0	*	*	*	*
1999.....	0.0	0.0	*	*	*	*	0.0	0.0	*	0.0	0.0	*	*	*	*
Poisoning by and exposure to other gases and vapors, undetermined intent (Y17)															
2004.....	0.0	0.0	*	*	*	*	0.0	0.0	*	0.0	0.0	*	*	*	*
2003.....	0.0	0.1	0.0	*	*	*	0.0	0.1	0.0	0.0	0.1	0.0	*	*	*
2002.....	0.0	0.0	*	*	*	*	0.0	0.0	*	0.0	0.0	*	*	*	*
2001.....	0.0	0.1	*	*	*	*	0.0	0.0	*	0.0	0.0	*	*	*	*
2000.....	0.0	0.0	0.0	*	*	*	0.0	0.0	*	0.0	0.0	*	*	*	*
1999.....	0.0	0.0	0.0	*	*	*	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
Discharge of firearms, undetermined intent (Y22-Y24)															
2004.....	0.1	0.1	0.0	0.0	*	*	0.1	0.1	0.0	0.1	0.1	0.0	0.1	0.1	*
2003.....	0.1	0.1	0.0	0.1	0.1	*	0.1	0.1	0.0	0.1	0.1	0.0	0.1	0.2	*
2002.....	0.1	0.1	0.0	0.1	0.1	*	0.1	0.1	0.0	0.1	0.1	0.0	0.1	0.2	*
2001.....	0.1	0.1	0.0	0.1	*	*	0.1	0.1	0.0	0.1	0.1	0.0	0.1	0.2	*
2000.....	0.1	0.1	0.0	0.0	*	*	0.1	0.1	0.0	0.1	0.1	0.0	0.1	0.3	*
1999.....	0.1	0.2	0.0	0.1	0.2	*	0.1	0.2	0.0	0.1	0.2	0.0	0.2	0.3	*
Falling, jumping or pushed from a high place, undetermined intent (Y30)															
2004.....	0.0	0.0	0.0	*	*	*	0.0	0.0	0.0	0.0	0.0	*	*	*	*
2003.....	0.0	0.0	0.0	*	*	*	0.0	0.0	0.0	0.0	0.0	*	*	*	*
2002.....	0.0	0.0	0.0	*	*	*	0.0	0.0	0.0	0.0	0.0	*	*	*	*
2001.....	0.0	0.0	0.0	*	*	*	0.0	0.0	0.0	0.0	0.0	*	*	*	*
2000.....	0.0	0.0	*	*	*	*	0.0	0.0	*	0.0	0.0	*	*	*	*
1999.....	0.0	0.0	*	*	*	*	0.0	0.0	*	0.0	0.0	*	*	*	*

Worktable Orig13. Age-adjusted death rates for 358 selected causes, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2004

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States: see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States, 2004." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), and year	All origins 1/			Hispanic			Non-Hispanic 2/			Non-Hispanic white			Non-Hispanic black		
	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female	Both Sexes	Male	Female
Misadventures to patients during medical and surgical care and their sequelae (Y60-Y69,Y88.1)															
2004.....	0.1	0.0	0.1	*	*	*	0.1	0.0	0.1	0.1	0.0	0.1	0.1	*	*
2003.....	0.1	0.0	0.1	*	*	*	0.1	0.0	0.1	0.0	0.0	0.1	0.1	*	0.1
2002.....	0.0	0.0	0.1	*	*	*	0.0	0.0	0.1	0.0	0.0	0.0	0.1	*	0.1
2001.....	0.1	0.1	0.1	*	*	*	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2000.....	0.1	0.1	0.1	*	*	*	0.1	0.1	0.1	0.0	0.1	0.0	0.1	*	0.2
1999.....	0.1	0.1	0.1	*	*	*	0.1	0.1	0.1	0.0	0.1	0.1	0.1	*	0.2
Other complications of medical and surgical care and their sequelae (Y70-Y84,Y88.2-Y88.3)															
2004.....	0.8	0.9	0.7	0.6	0.7	0.5	0.8	0.9	0.7	0.8	0.9	0.6	1.3	1.3	1.3
2003.....	0.8	0.9	0.8	0.6	0.7	0.6	0.8	0.9	0.8	0.8	0.9	0.7	1.3	1.2	1.3
2002.....	0.8	0.9	0.8	0.5	0.4	0.6	0.9	0.9	0.8	0.8	0.9	0.8	1.4	1.5	1.4
2001.....	0.9	1.0	0.9	0.7	0.8	0.6	0.9	1.0	0.9	0.8	0.9	0.8	1.7	2.0	1.5
2000.....	0.9	1.1	0.9	0.6	0.8	0.6	1.0	1.1	0.9	0.9	1.0	0.8	1.6	1.7	1.6
1999.....	0.9	1.0	0.8	0.5	0.5	0.4	0.9	1.0	0.8	0.8	1.0	0.8	1.5	1.7	1.4

* Figure does not meet standards of reliability or precision.

0.0 Quantity more than zero but less than 0.05.

... Category not applicable.

1/ Figures for origin not stated are included in "All origins" but not distributed among specified origins.

2/ Includes races other than white and black.

3/ Figures include September 11, 2001 related deaths for which death certificates were filed as of 10/24/02; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001."

Source: CDC/NCHS, National Vital Statistics System, Mortality.