

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All causes												
All races 3/, both sexes												
2004.....	816.5	685.2	29.9	16.8	80.1	102.1	193.5	427.0	910.3	2,164.6	5,275.1	13,823.5
2003.....	841.9	700.0	31.5	17.0	81.5	103.6	201.6	433.2	940.9	2,255.0	5,463.1	14,593.3
2002.....	847.3	695.0	31.2	17.4	81.4	103.6	202.9	430.1	952.4	2,314.7	5,556.9	14,828.3
2001.....	848.5	683.4	33.3	17.3	80.7	105.2	203.6	428.9	964.6	2,353.3	5,582.4	15,112.8
2000.....	854.0	736.7	32.4	18.0	79.9	101.4	198.9	425.6	992.2	2,399.1	5,666.5	15,524.4
1999.....	857.0	736.0	34.2	18.6	79.3	102.2	198.0	418.2	1,005.0	2,457.3	5,714.5	15,554.6
All races 3/, male												
2004.....	817.6	753.7	32.4	19.2	114.7	139.5	243.6	543.5	1,128.8	2,644.8	6,394.3	15,031.1
2003.....	840.3	777.4	35.1	19.8	116.5	141.4	255.0	552.2	1,165.5	2,771.7	6,641.8	15,794.0
2002.....	846.6	761.5	35.2	20.0	117.3	142.2	257.5	547.5	1,184.0	2,855.3	6,760.5	16,254.5
2001.....	846.4	749.8	37.0	19.8	117.0	143.7	259.6	545.1	1,192.7	2,911.5	6,833.0	16,744.8
2000.....	853.0	806.5	35.9	20.9	114.9	138.6	255.2	542.8	1,230.7	2,979.6	6,972.6	17,501.4
1999.....	859.2	805.0	37.9	21.5	113.1	139.7	254.9	533.1	1,252.0	3,073.7	7,083.3	17,597.2
All races 3/, female												
2004.....	815.4	613.4	27.3	14.3	43.6	63.5	143.5	314.3	707.4	1,761.4	4,521.8	13,280.3
2003.....	843.4	619.1	27.8	14.0	44.4	64.6	148.5	318.4	732.7	1,823.0	4,675.5	14,062.5
2002.....	848.0	625.3	27.0	14.7	43.7	64.0	148.8	316.9	738.0	1,864.7	4,757.9	14,209.6
2001.....	850.4	613.9	29.5	14.6	42.6	66.0	148.2	316.8	754.0	1,890.8	4,760.5	14,429.9
2000.....	855.0	663.4	28.7	15.0	43.1	63.5	143.2	312.5	772.2	1,921.2	4,814.7	14,719.2
1999.....	854.9	663.6	30.3	15.6	43.7	64.1	141.8	307.6	777.6	1,952.3	4,825.4	14,731.3
White, both sexes												
2004.....	863.2	574.0	27.0	15.5	76.4	92.9	179.8	394.0	863.5	2,120.4	5,272.8	14,010.1
2003.....	890.1	591.3	28.5	15.8	77.1	94.3	185.7	397.5	892.2	2,207.0	5,457.8	14,792.3
2002.....	895.7	586.7	28.1	16.1	77.1	93.5	186.6	395.3	902.5	2,267.6	5,544.2	15,015.9
2001.....	895.1	570.9	30.7	16.2	75.5	95.3	185.6	393.2	913.5	2,298.9	5,560.9	15,287.5
2000.....	900.2	610.6	29.2	17.0	74.5	90.3	179.9	388.3	940.2	2,341.5	5,637.7	15,707.0
1999.....	901.4	606.1	30.8	17.2	73.2	91.4	177.3	380.4	948.1	2,396.2	5,679.5	15,718.0
White, male												
2004.....	854.2	631.6	29.4	17.9	108.3	127.0	229.1	504.0	1,065.9	2,584.0	6,384.8	15,250.7
2003.....	877.6	658.7	31.5	18.4	108.9	128.8	237.6	509.2	1,102.3	2,707.3	6,620.6	16,037.9
2002.....	884.0	650.9	31.5	18.4	109.7	128.3	239.3	505.4	1,118.6	2,795.4	6,738.8	16,473.2
2001.....	881.9	627.6	34.2	18.4	108.0	130.3	239.7	501.3	1,127.5	2,842.3	6,799.7	16,935.4
2000.....	887.8	667.6	32.6	19.8	105.8	124.1	233.6	496.9	1,163.3	2,905.7	6,933.1	17,716.4
1999.....	892.1	667.0	33.9	19.7	102.8	125.4	230.8	484.6	1,179.7	2,998.7	7,040.1	17,752.9

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All causes												
White, female												
2004.....	871.9	513.6	24.4	13.0	42.4	56.8	129.5	285.4	671.8	1,723.6	4,514.4	13,450.9
2003.....	902.3	520.7	25.5	13.1	43.2	58.0	132.9	287.2	693.7	1,780.6	4,671.6	14,240.6
2002.....	907.0	519.4	24.5	13.7	42.4	56.9	133.2	286.8	698.7	1,819.7	4,742.5	14,382.8
2001.....	907.9	511.5	27.1	13.9	40.8	58.6	131.0	286.8	712.2	1,840.2	4,738.5	14,597.6
2000.....	912.3	550.5	25.5	14.1	41.1	55.1	125.7	281.4	730.9	1,868.3	4,785.3	14,890.7
1999.....	910.4	542.0	27.5	14.6	41.5	56.2	123.2	277.9	731.0	1,893.9	4,787.8	14,900.6
Black, both sexes												
2004.....	744.3	1,284.7	44.8	23.6	109.4	179.3	322.3	744.8	1,499.6	2,982.6	6,170.4	13,338.5
2003.....	763.6	1,273.8	46.8	22.9	113.1	181.6	343.6	771.4	1,549.9	3,108.4	6,353.9	13,976.7
2002.....	768.4	1,263.6	47.1	24.5	113.7	186.8	348.6	765.8	1,565.2	3,159.3	6,518.9	14,260.8
2001.....	773.5	1,304.5	47.5	23.3	117.7	185.0	356.5	771.9	1,576.8	3,257.6	6,585.6	14,550.8
2000.....	781.1	1,426.1	49.9	24.2	119.4	187.8	356.9	786.9	1,607.8	3,335.6	6,701.3	14,714.3
1999.....	788.1	1,457.3	52.6	26.5	121.2	184.4	368.4	782.4	1,655.6	3,413.6	6,753.3	14,757.9
Black, male												
2004.....	792.6	1,414.2	48.6	26.0	164.3	252.3	397.0	954.9	1,960.8	3,818.3	7,710.3	14,452.5
2003.....	813.7	1,410.0	53.7	26.8	171.3	256.2	426.7	991.4	2,011.2	3,981.3	8,067.0	14,903.4
2002.....	816.7	1,351.5	54.4	28.9	172.6	264.5	434.7	983.0	2,039.2	4,024.5	8,169.6	15,635.5
2001.....	823.9	1,426.4	53.2	27.7	180.7	259.2	445.0	996.1	2,025.3	4,166.6	8,355.9	16,439.9
2000.....	834.1	1,567.6	54.5	28.2	181.4	261.0	453.0	1,017.7	2,080.1	4,253.5	8,486.0	16,791.0
1999.....	847.4	1,592.8	59.0	32.0	184.6	258.6	469.2	1,030.7	2,145.6	4,352.3	8,559.1	17,304.5
Black, female												
2004.....	700.3	1,149.9	40.9	21.1	53.7	112.3	256.0	564.1	1,128.6	2,386.1	5,300.0	12,896.9
2003.....	717.9	1,132.2	39.7	18.9	54.0	113.5	270.0	582.3	1,178.8	2,487.2	5,385.5	13,616.7
2002.....	724.4	1,172.0	39.5	19.9	54.4	116.4	272.3	579.4	1,184.2	2,545.0	5,584.4	13,734.2
2001.....	727.7	1,176.3	41.7	18.7	54.9	117.9	278.0	579.1	1,215.5	2,616.1	5,591.6	13,849.9
2000.....	733.0	1,279.8	45.3	20.0	58.3	121.8	271.9	588.3	1,227.2	2,689.6	5,696.5	13,941.3
1999.....	734.3	1,317.4	46.1	20.9	58.6	117.6	279.4	569.8	1,262.7	2,751.5	5,742.4	13,805.9

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
I. Certain infectious and parasitic diseases (A00-B99)												
All races 3/, both sexes												
2004.....	22.0	11.8	1.2	0.4	1.1	5.1	15.5	24.8	28.0	49.5	113.5	249.5
2003.....	22.2	12.3	1.2	0.5	1.2	5.4	16.7	25.1	27.6	49.3	114.4	255.2
2002.....	22.2	14.4	1.1	0.6	1.2	6.2	17.5	24.9	26.1	50.7	113.4	252.8
2001.....	21.2	14.0	1.5	0.5	1.3	6.7	17.6	22.9	25.4	48.0	104.9	246.3
2000.....	21.0	13.4	1.4	0.5	1.2	7.5	17.7	22.9	24.4	45.8	103.7	256.0
1999.....	21.5	14.8	1.5	0.6	1.2	8.4	18.7	22.1	24.1	48.0	105.1	267.9
All races 3/, male												
2004.....	23.7	13.8	1.1	0.4	1.3	6.0	21.0	34.9	35.8	56.6	128.1	273.1
2003.....	24.0	12.6	1.4	0.4	1.2	6.7	23.2	35.8	34.9	56.2	128.8	269.1
2002.....	24.2	14.8	1.2	0.6	1.2	7.6	24.6	36.1	33.0	58.0	130.8	270.9
2001.....	23.3	14.7	1.6	0.5	1.4	8.5	25.2	33.1	31.9	54.5	122.3	269.7
2000.....	23.0	14.1	1.5	0.5	1.3	9.5	25.4	33.7	30.8	52.7	120.5	273.0
1999.....	23.7	17.0	1.8	0.7	1.3	11.2	27.1	32.2	30.4	55.8	123.3	290.1
All races 3/, female												
2004.....	20.3	9.6	1.2	0.4	1.0	4.2	9.9	15.1	20.7	43.5	103.7	238.8
2003.....	20.6	12.1	1.1	0.5	1.1	4.0	10.3	14.7	20.8	43.5	104.8	249.1
2002.....	20.2	14.0	0.9	0.5	1.1	4.7	10.4	14.1	19.7	44.6	101.9	245.0
2001.....	19.2	13.2	1.3	0.5	1.3	4.8	10.2	13.0	19.4	42.5	93.5	236.5
2000.....	19.0	12.7	1.4	0.5	1.0	5.4	10.1	12.5	18.4	40.1	92.7	249.1
1999.....	19.3	12.5	1.2	0.6	1.2	5.6	10.3	12.4	18.4	41.7	93.3	258.9
White, both sexes												
2004.....	19.9	9.9	1.1	0.3	0.7	2.8	10.2	18.4	22.7	44.3	108.0	242.1
2003.....	20.0	9.0	1.0	0.4	0.8	3.1	11.0	18.6	22.2	43.9	108.7	245.0
2002.....	19.7	12.4	1.0	0.5	0.8	3.5	11.2	18.1	20.8	45.1	106.9	242.3
2001.....	18.6	10.6	1.3	0.4	0.8	3.7	11.3	16.3	20.2	42.5	98.8	235.4
2000.....	18.2	10.6	1.2	0.4	0.8	4.1	11.5	16.1	19.2	40.2	96.0	242.9
1999.....	18.6	11.9	1.3	0.5	0.8	5.0	12.0	15.3	18.9	41.9	97.1	251.9
White, male												
2004.....	20.9	11.6	1.1	0.4	0.8	3.6	14.7	26.3	28.3	50.1	121.5	266.6
2003.....	20.9	8.6	1.2	0.3	0.9	4.2	16.1	27.1	27.5	49.4	122.0	259.4
2002.....	20.9	12.7	1.2	0.6	0.8	4.7	16.8	26.5	26.0	51.2	122.0	260.3
2001.....	19.8	11.3	1.5	0.5	0.9	5.1	17.3	24.0	25.0	47.8	115.1	256.7
2000.....	19.3	11.2	1.2	0.4	1.0	5.8	17.4	23.8	23.9	46.1	111.0	258.5
1999.....	19.8	14.3	1.6	0.5	1.0	7.3	18.5	22.3	23.1	48.3	114.0	270.4

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
I. Certain infectious and parasitic diseases (A00-B99)												
White, female												
2004.....	19.0	8.1	1.1	0.3	0.6	1.9	5.7	10.6	17.3	39.3	98.8	231.1
2003.....	19.1	9.4	0.9	0.4	0.8	2.0	5.7	10.2	17.2	39.3	99.7	238.6
2002.....	18.6	12.1	0.8	0.4	0.8	2.1	5.6	9.8	15.9	39.9	96.8	234.5
2001.....	17.4	9.9	1.2	0.4	0.8	2.2	5.3	8.8	15.7	38.0	88.0	226.4
2000.....	17.2	10.0	1.2	0.4	0.6	2.4	5.6	8.4	14.8	35.3	86.2	236.5
1999.....	17.4	9.4	1.1	0.5	0.7	2.7	5.5	8.5	14.9	36.6	86.0	244.5
Black, both sexes												
2004.....	40.4	21.8	1.5	0.6	3.4	20.2	52.7	75.4	78.0	102.1	193.5	375.3
2003.....	41.7	28.2	2.3	0.9	3.1	20.2	58.1	77.5	78.1	102.7	194.9	416.6
2002.....	42.9	24.6	1.7	0.8	3.0	24.0	61.8	79.8	74.7	107.1	204.2	413.1
2001.....	42.3	31.2	2.2	0.9	4.1	26.4	62.4	76.5	73.9	102.9	189.8	404.7
2000.....	43.2	29.9	2.4	0.8	3.4	29.8	62.6	79.4	70.0	103.2	206.8	436.7
1999.....	44.5	31.1	2.5	1.2	3.5	31.1	66.8	78.2	70.7	110.4	207.6	469.0
Black, male												
2004.....	47.8	24.3	*	*	3.6	22.6	69.2	108.1	112.2	129.0	233.9	408.3
2003.....	50.0	31.8	2.5	0.8	2.9	24.3	78.1	112.5	111.3	132.0	235.8	456.7
2002.....	51.7	25.3	1.6	0.7	2.9	28.1	84.8	119.0	103.0	135.3	257.0	448.1
2001.....	51.2	33.2	2.6	0.9	3.9	32.8	86.3	113.1	102.1	128.6	231.3	470.9
2000.....	52.7	31.3	2.8	0.8	3.4	36.2	87.9	121.1	98.2	128.3	258.6	486.7
1999.....	54.6	33.2	3.2	1.3	3.1	40.1	94.7	119.8	100.0	140.0	253.0	537.8
Black, female												
2004.....	33.7	19.1	*	0.7	3.3	18.0	38.1	47.3	50.5	83.0	170.7	362.3
2003.....	34.2	24.4	2.0	0.9	3.2	16.4	40.3	47.5	51.3	81.8	171.7	401.0
2002.....	35.0	23.9	1.7	1.0	3.1	20.3	41.5	46.2	51.9	87.0	174.3	399.6
2001.....	34.2	29.0	1.8	0.8	4.2	20.6	41.3	45.1	51.1	84.7	166.6	380.2
2000.....	34.5	28.4	2.1	0.8	3.4	24.0	40.1	43.5	47.2	85.6	177.6	418.1
1999.....	35.3	28.9	1.9	1.1	4.0	23.0	42.2	42.6	47.2	89.5	182.1	443.3

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Intestinal infectious diseases (A00-A09)												
All races 3/, both sexes												
2004.....	1.5	*	*	*	*	*	0.1	0.2	0.9	3.3	12.9	34.4
2003.....	1.1	*	*	*	*	*	0.0	0.2	0.7	2.5	9.4	25.0
2002.....	0.9	*	*	*	*	*	0.0	0.2	0.5	2.0	7.1	21.0
2001.....	0.6	*	*	*	*	*	*	0.1	0.3	1.3	4.6	13.1
2000.....	0.5	*	*	*	*	*	0.0	0.1	0.3	1.3	3.9	11.9
1999.....	0.4	0.6	*	*	*	*	*	0.1	0.2	1.0	3.3	10.6
All races 3/, male												
2004.....	1.1	*	*	*	*	*	0.1	0.2	0.9	3.2	12.9	32.2
2003.....	0.8	*	*	*	*	*	*	0.2	0.6	2.3	9.4	21.9
2002.....	0.7	*	*	*	*	*	*	0.2	0.6	2.3	7.1	20.4
2001.....	0.4	*	*	*	*	*	*	0.1	0.3	1.4	4.7	12.2
2000.....	0.4	*	*	*	*	*	*	0.1	0.4	1.3	4.1	11.5
1999.....	0.3	*	*	*	*	*	*	*	0.3	1.0	3.6	9.7
All races 3/, female												
2004.....	1.8	*	*	*	*	*	*	0.2	0.8	3.5	12.9	35.4
2003.....	1.4	*	*	*	*	*	*	0.2	0.7	2.6	9.3	26.4
2002.....	1.0	*	*	*	*	*	*	0.1	0.5	1.8	7.0	21.3
2001.....	0.7	*	*	*	*	*	*	0.1	0.4	1.3	4.6	13.4
2000.....	0.6	*	*	*	*	*	*	*	0.2	1.2	3.7	12.0
1999.....	0.5	*	*	*	*	*	*	*	0.2	1.0	3.1	10.9
White, both sexes												
2004.....	1.7	*	*	*	*	*	0.1	0.2	0.9	3.5	13.6	36.7
2003.....	1.2	*	*	*	*	*	*	0.2	0.7	2.6	9.8	26.3
2002.....	1.0	*	*	*	*	*	*	0.2	0.5	2.1	7.5	22.1
2001.....	0.6	*	*	*	*	*	*	0.1	0.3	1.3	4.8	13.8
2000.....	0.6	*	*	*	*	*	*	0.1	0.3	1.3	4.1	12.3
1999.....	0.5	*	*	*	*	*	*	0.1	0.2	1.1	3.4	11.2
White, male												
2004.....	1.3	*	*	*	*	*	*	0.2	0.9	3.3	13.5	34.2
2003.....	0.9	*	*	*	*	*	*	0.2	0.6	2.4	9.9	23.2
2002.....	0.8	*	*	*	*	*	*	0.2	0.5	2.3	7.5	21.5
2001.....	0.5	*	*	*	*	*	*	*	0.3	1.4	4.8	12.8
2000.....	0.4	*	*	*	*	*	*	*	0.4	1.4	4.3	12.0
1999.....	0.4	*	*	*	*	*	*	*	0.2	1.1	3.8	10.2

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other intestinal infectious diseases (A01-A08)												
All races 3/, both sexes												
2004.....	1.5	*	*	*	*	*	0.1	0.2	0.9	3.3	12.5	33.6
2003.....	1.0	*	*	*	*	*	0.0	0.2	0.6	2.4	9.1	24.2
2002.....	0.8	*	*	*	*	*	0.0	0.1	0.5	1.9	6.8	20.2
2001.....	0.5	*	*	*	*	*	*	0.1	0.3	1.3	4.5	12.6
2000.....	0.5	*	*	*	*	*	*	0.1	0.3	1.2	3.7	11.3
1999.....	0.4	0.6	*	*	*	*	*	0.1	0.2	0.9	3.0	9.9
All races 3/, male												
2004.....	1.1	*	*	*	*	*	0.1	0.2	0.9	3.1	12.6	31.7
2003.....	0.8	*	*	*	*	*	*	0.2	0.6	2.2	9.1	21.1
2002.....	0.7	*	*	*	*	*	*	0.2	0.5	2.2	6.9	19.7
2001.....	0.4	*	*	*	*	*	*	0.1	0.3	1.3	4.5	11.9
2000.....	0.4	*	*	*	*	*	*	0.1	0.4	1.3	4.0	10.6
1999.....	0.3	*	*	*	*	*	*	*	0.2	0.9	3.4	9.3
All races 3/, female												
2004.....	1.8	*	*	*	*	*	*	0.2	0.8	3.4	12.5	34.5
2003.....	1.3	*	*	*	*	*	*	0.1	0.7	2.5	9.1	25.5
2002.....	1.0	*	*	*	*	*	*	0.1	0.5	1.7	6.8	20.4
2001.....	0.7	*	*	*	*	*	*	0.1	0.3	1.2	4.4	13.0
2000.....	0.6	*	*	*	*	*	*	*	0.2	1.1	3.6	11.5
1999.....	0.5	*	*	*	*	*	*	*	0.2	0.9	2.8	10.2
White, both sexes												
2004.....	1.7	*	*	*	*	*	0.1	0.2	0.9	3.4	13.2	35.9
2003.....	1.2	*	*	*	*	*	*	0.2	0.6	2.6	9.5	25.4
2002.....	1.0	*	*	*	*	*	*	0.2	0.5	2.0	7.3	21.1
2001.....	0.6	*	*	*	*	*	*	0.1	0.3	1.3	4.6	13.3
2000.....	0.5	*	*	*	*	*	*	0.1	0.3	1.2	4.0	11.7
1999.....	0.4	*	*	*	*	*	*	0.1	0.2	0.9	3.1	10.5
White, male												
2004.....	1.3	*	*	*	*	*	*	0.2	0.9	3.2	13.2	33.7
2003.....	0.9	*	*	*	*	*	*	0.2	0.6	2.4	9.6	22.4
2002.....	0.7	*	*	*	*	*	*	0.2	0.5	2.2	7.3	20.7
2001.....	0.5	*	*	*	*	*	*	*	0.3	1.3	4.6	12.4
2000.....	0.4	*	*	*	*	*	*	*	0.4	1.3	4.1	11.1
1999.....	0.3	*	*	*	*	*	*	*	0.2	0.9	3.6	9.7

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Intestinal infections due to other specified organisms (A04,A07-A08)

All races 3/, both sexes

2004.....	1.4	*	*	*	*	*	0.1	0.2	0.8	3.2	12.5	33.5
2003.....	1.0	*	*	*	*	*	*	0.2	0.6	2.4	9.0	23.9
2002.....	0.8	*	*	*	*	*	*	0.1	0.5	1.9	6.8	20.2
2001.....	0.5	*	*	*	*	*	*	0.1	0.3	1.2	4.3	12.4
2000.....	0.4	*	*	*	*	*	*	0.1	0.3	1.1	3.6	11.1
1999.....	0.4	*	*	*	*	*	*	0.1	0.2	0.8	2.9	9.7

All races 3/, male

2004.....	1.1	*	*	*	*	*	*	0.2	0.9	3.1	12.5	31.4
2003.....	0.7	*	*	*	*	*	*	0.2	0.6	2.2	9.0	20.6
2002.....	0.6	*	*	*	*	*	*	0.1	0.5	2.1	6.8	19.6
2001.....	0.4	*	*	*	*	*	*	*	0.3	1.3	4.4	11.8
2000.....	0.4	*	*	*	*	*	*	*	0.3	1.2	3.9	10.5
1999.....	0.3	*	*	*	*	*	*	*	0.2	0.8	3.3	9.1

All races 3/, female

2004.....	1.8	*	*	*	*	*	*	0.2	0.8	3.3	12.5	34.4
2003.....	1.3	*	*	*	*	*	*	0.1	0.6	2.5	9.0	25.4
2002.....	1.0	*	*	*	*	*	*	0.1	0.5	1.7	6.7	20.4
2001.....	0.6	*	*	*	*	*	*	0.1	0.3	1.2	4.3	12.7
2000.....	0.5	*	*	*	*	*	*	*	0.2	1.1	3.4	11.4
1999.....	0.4	*	*	*	*	*	*	*	0.2	0.9	2.7	9.9

White, both sexes

2004.....	1.7	*	*	*	*	*	0.1	0.2	0.9	3.4	13.2	35.8
2003.....	1.2	*	*	*	*	*	*	0.1	0.6	2.5	9.4	25.2
2002.....	0.9	*	*	*	*	*	*	0.1	0.5	2.0	7.2	21.1
2001.....	0.6	*	*	*	*	*	*	0.1	0.3	1.3	4.5	13.1
2000.....	0.5	*	*	*	*	*	*	0.1	0.3	1.2	3.8	11.5
1999.....	0.4	*	*	*	*	*	*	*	0.2	0.9	3.0	10.2

White, male

2004.....	1.3	*	*	*	*	*	*	0.2	0.9	3.2	13.1	33.5
2003.....	0.9	*	*	*	*	*	*	0.2	0.6	2.3	9.5	22.0
2002.....	0.7	*	*	*	*	*	*	0.2	0.5	2.2	7.2	20.6
2001.....	0.4	*	*	*	*	*	*	*	0.3	1.3	4.4	12.3
2000.....	0.4	*	*	*	*	*	*	*	0.3	1.2	4.0	11.0
1999.....	0.3	*	*	*	*	*	*	*	0.2	0.8	3.5	9.5

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Tuberculosis (A16-A19)												
All races 3/, both sexes												
2004.....	0.2	*	*	*	*	*	0.1	0.2	0.3	0.7	1.3	2.4
2003.....	0.2	*	*	*	*	*	0.1	0.3	0.4	0.6	1.7	2.3
2002.....	0.3	*	*	*	*	*	0.1	0.2	0.4	0.9	1.8	2.6
2001.....	0.3	*	*	*	*	0.1	0.1	0.2	0.5	0.8	1.6	2.8
2000.....	0.3	*	*	*	*	0.1	0.1	0.2	0.3	0.8	2.0	2.7
1999.....	0.3	*	*	*	*	*	0.1	0.3	0.5	0.9	2.1	4.2
All races 3/, male												
2004.....	0.3	*	*	*	*	*	0.2	0.3	0.4	1.1	1.6	3.8
2003.....	0.3	*	*	*	*	*	0.1	0.4	0.5	0.9	2.3	3.1
2002.....	0.3	*	*	*	*	*	0.2	0.3	0.6	1.2	2.4	4.0
2001.....	0.3	*	*	*	*	*	0.1	0.3	0.7	1.2	2.0	4.4
2000.....	0.3	*	*	*	*	*	0.1	0.4	0.5	1.0	2.8	4.6
1999.....	0.4	*	*	*	*	*	0.2	0.5	0.8	1.4	2.8	7.1
All races 3/, female												
2004.....	0.2	*	*	*	*	*	*	0.1	*	0.4	1.0	1.7
2003.....	0.2	*	*	*	*	*	*	0.1	0.2	0.4	1.3	1.9
2002.....	0.2	*	*	*	*	*	*	0.1	0.2	0.6	1.5	2.1
2001.....	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.4	2.2
2000.....	0.2	*	*	*	*	*	*	*	0.2	0.7	1.5	2.0
1999.....	0.2	*	*	*	*	*	*	*	0.3	0.6	1.6	3.0
White, both sexes												
2004.....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.5	1.0	1.7
2003.....	0.2	*	*	*	*	*	*	0.1	0.3	0.5	1.3	1.8
2002.....	0.2	*	*	*	*	*	0.1	0.1	0.3	0.6	1.4	2.1
2001.....	0.2	*	*	*	*	*	0.1	0.1	0.3	0.6	1.2	2.2
2000.....	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.4	2.2
1999.....	0.2	*	*	*	*	*	0.1	0.2	0.4	0.7	1.6	3.2
White, male												
2004.....	0.2	*	*	*	*	*	*	0.2	0.3	0.7	1.2	2.8
2003.....	0.2	*	*	*	*	*	*	0.2	0.4	0.6	1.7	2.4
2002.....	0.2	*	*	*	*	*	*	0.2	0.3	0.9	1.8	3.2
2001.....	0.2	*	*	*	*	*	*	0.2	0.5	0.7	1.5	3.1
2000.....	0.2	*	*	*	*	*	*	0.2	0.2	0.7	1.8	3.8
1999.....	0.3	*	*	*	*	*	*	0.3	0.5	1.0	2.1	5.4

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Respiratory tuberculosis (A16)												
All races 3/, both sexes												
2004.....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.6	0.9	2.2
2003.....	0.2	*	*	*	*	*	0.1	0.2	0.3	0.5	1.4	1.8
2002.....	0.2	*	*	*	*	*	0.1	0.2	0.3	0.7	1.5	2.3
2001.....	0.2	*	*	*	*	*	0.1	0.2	0.4	0.6	1.3	2.4
2000.....	0.2	*	*	*	*	*	0.1	0.2	0.2	0.6	1.7	2.3
1999.....	0.3	*	*	*	*	*	0.1	0.2	0.5	0.8	1.8	3.7
All races 3/, male												
2004.....	0.2	*	*	*	*	*	0.1	0.2	0.3	0.9	1.2	3.7
2003.....	0.2	*	*	*	*	*	*	0.3	0.4	0.7	1.9	2.5
2002.....	0.3	*	*	*	*	*	0.1	0.2	0.5	1.0	1.9	3.5
2001.....	0.3	*	*	*	*	*	*	0.3	0.6	0.9	1.7	3.8
2000.....	0.3	*	*	*	*	*	0.1	0.3	0.4	0.9	2.4	4.1
1999.....	0.4	*	*	*	*	*	0.1	0.4	0.7	1.2	2.5	6.4
All races 3/, female												
2004.....	0.1	*	*	*	*	*	*	*	*	0.3	0.7	1.5
2003.....	0.1	*	*	*	*	*	*	0.1	0.2	0.3	1.0	1.5
2002.....	0.2	*	*	*	*	*	*	0.1	0.2	0.4	1.3	1.8
2001.....	0.2	*	*	*	*	*	*	*	*	0.4	1.1	1.8
2000.....	0.2	*	*	*	*	*	*	*	*	0.5	1.2	1.6
1999.....	0.2	*	*	*	*	*	*	*	0.2	0.4	1.3	2.6
White, both sexes												
2004.....	0.1	*	*	*	*	*	0.1	0.1	0.1	0.4	0.7	1.6
2003.....	0.1	*	*	*	*	*	*	0.1	0.2	0.4	1.1	1.4
2002.....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.4	1.1	1.8
2001.....	0.2	*	*	*	*	*	*	0.1	0.3	0.4	1.0	1.8
2000.....	0.2	*	*	*	*	*	*	0.1	0.1	0.4	1.2	1.9
1999.....	0.2	*	*	*	*	*	0.1	0.1	0.3	0.5	1.3	3.0
White, male												
2004.....	0.2	*	*	*	*	*	*	*	0.2	0.6	1.0	2.6
2003.....	0.2	*	*	*	*	*	*	0.2	0.3	0.5	1.4	1.8
2002.....	0.2	*	*	*	*	*	*	0.1	0.3	0.7	1.4	2.9
2001.....	0.2	*	*	*	*	*	*	0.1	0.5	0.6	1.3	2.6
2000.....	0.2	*	*	*	*	*	*	0.2	0.2	0.6	1.5	3.4
1999.....	0.3	*	*	*	*	*	*	0.2	0.5	0.8	1.8	5.2

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Zoonotic and other bacterial diseases (A20-A49)												
All races 3/, both sexes												
2004.....	12.0	7.5	0.7	0.2	0.4	0.9	2.1	5.7	13.7	33.9	85.6	194.8
2003.....	12.3	7.9	0.7	0.2	0.5	0.9	2.2	5.6	13.9	34.2	88.7	209.5
2002.....	12.3	8.6	0.6	0.3	0.4	0.9	2.1	5.5	13.3	36.2	90.2	210.5
2001.....	11.9	8.9	0.9	0.3	0.5	0.8	2.0	5.4	13.0	34.3	85.9	213.7
2000.....	11.6	8.6	0.8	0.2	0.4	0.8	2.1	5.3	12.6	32.3	83.8	222.5
1999.....	11.7	9.1	0.8	0.3	0.5	0.8	2.0	5.0	12.1	33.2	83.9	231.2
All races 3/, male												
2004.....	11.0	8.9	0.7	0.2	0.4	0.9	2.1	6.3	14.7	37.7	97.8	214.7
2003.....	11.1	8.4	0.8	0.2	0.5	0.9	2.2	6.2	15.1	38.2	99.4	223.2
2002.....	11.1	8.6	0.7	0.3	0.4	0.9	2.2	6.1	14.4	39.7	103.7	222.5
2001.....	10.8	9.5	0.9	0.3	0.5	0.8	2.2	5.9	14.1	38.4	99.9	233.5
2000.....	10.4	8.9	0.9	0.2	0.5	0.9	2.2	6.1	13.7	36.1	96.0	233.1
1999.....	10.5	10.4	0.9	0.3	0.5	0.9	2.3	5.6	13.3	37.6	97.5	246.7
All races 3/, female												
2004.....	12.9	6.1	0.7	0.2	0.4	0.9	2.0	5.2	12.7	30.7	77.5	185.8
2003.....	13.5	7.5	0.5	0.3	0.5	0.8	2.2	5.1	12.8	30.8	81.5	203.5
2002.....	13.5	8.5	0.6	0.3	0.4	0.8	2.0	4.9	12.3	33.3	81.3	205.3
2001.....	12.9	8.3	0.8	0.2	0.5	0.8	1.8	4.9	12.0	30.8	76.6	205.4
2000.....	12.9	8.3	0.7	0.2	0.4	0.7	2.0	4.5	11.5	29.2	75.9	218.2
1999.....	12.9	7.7	0.7	0.3	0.4	0.8	1.8	4.5	11.1	29.7	75.0	225.0
White, both sexes												
2004.....	11.8	6.1	0.6	0.2	0.3	0.7	1.7	4.7	11.7	30.6	80.5	186.1
2003.....	12.1	5.6	0.5	0.2	0.4	0.7	1.8	4.5	11.9	30.5	83.9	198.2
2002.....	12.1	7.3	0.6	0.3	0.4	0.7	1.7	4.5	11.2	32.5	84.4	199.4
2001.....	11.7	6.8	0.8	0.3	0.4	0.6	1.6	4.3	10.9	30.7	80.6	202.9
2000.....	11.3	7.1	0.7	0.2	0.4	0.6	1.7	4.2	10.5	28.7	77.2	209.6
1999.....	11.3	7.2	0.8	0.2	0.4	0.7	1.6	4.0	10.1	29.2	76.9	216.1
White, male												
2004.....	10.8	7.3	0.6	0.2	0.4	0.7	1.8	5.2	12.5	33.9	91.7	207.4
2003.....	10.8	5.6	0.7	0.2	0.4	0.7	1.8	5.0	12.8	33.9	94.0	212.7
2002.....	10.9	7.3	0.7	0.4	0.4	0.8	1.8	5.1	12.3	35.7	96.1	211.9
2001.....	10.5	7.5	0.9	0.3	0.4	0.6	1.8	4.7	11.8	34.1	93.8	221.0
2000.....	10.0	7.2	0.8	0.2	0.5	0.6	1.8	4.8	11.6	32.3	88.6	219.5
1999.....	10.0	8.5	0.8	0.2	0.4	0.7	1.8	4.4	10.8	32.8	89.5	230.1

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Zoonotic and other bacterial diseases (A20-A49)												
White, female												
2004.....	12.8	4.9	0.6	0.2	0.3	0.7	1.6	4.2	11.0	27.7	72.8	176.5
2003.....	13.4	5.6	0.4	0.2	0.4	0.7	1.8	4.1	11.1	27.5	77.0	191.9
2002.....	13.4	7.4	0.5	0.2	0.4	0.7	1.6	4.0	10.1	29.9	76.5	194.0
2001.....	12.8	6.1	0.7	0.2	0.4	0.6	1.4	4.0	10.1	27.8	71.8	195.4
2000.....	12.6	7.0	0.7	0.2	0.3	0.6	1.6	3.6	9.5	25.6	69.7	205.6
1999.....	12.6	5.9	0.7	0.2	0.3	0.6	1.4	3.6	9.5	26.1	68.6	210.5
Black, both sexes												
2004.....	16.2	15.1	0.9	*	0.8	2.1	4.8	14.2	33.0	72.1	162.1	328.8
2003.....	16.9	19.6	1.2	0.3	0.8	2.0	5.3	14.5	33.8	75.1	161.9	379.6
2002.....	16.9	14.8	0.8	0.5	0.5	2.0	4.9	13.6	34.4	77.1	175.8	376.3
2001.....	16.5	20.3	1.1	*	1.0	2.0	4.9	14.1	33.8	74.3	165.0	373.1
2000.....	16.8	18.4	1.2	0.3	0.7	2.1	5.1	14.3	32.5	74.2	177.9	402.3
1999.....	17.0	19.0	0.9	0.3	0.8	1.9	5.4	13.8	31.6	77.9	177.2	428.2
Black, male												
2004.....	15.3	16.9	*	*	0.9	2.3	4.9	15.9	38.0	85.1	196.8	352.7
2003.....	16.0	22.0	*	*	0.7	2.3	5.4	16.6	40.0	89.6	194.1	416.0
2002.....	15.7	15.7	*	*	*	2.0	5.4	15.5	37.5	89.1	218.9	402.0
2001.....	15.6	20.7	*	*	1.1	2.0	5.0	15.8	38.9	88.9	200.4	435.7
2000.....	15.6	19.8	*	*	0.8	2.5	5.1	17.5	36.6	84.3	212.6	441.7
1999.....	16.3	20.5	*	*	0.7	2.1	6.3	16.2	38.8	94.9	215.2	472.2
Black, female												
2004.....	17.1	13.3	*	*	0.8	1.9	4.7	12.8	28.9	62.9	142.4	319.4
2003.....	17.7	17.0	*	*	0.9	1.7	5.1	12.8	28.9	64.8	143.7	365.4
2002.....	18.0	13.9	*	0.6	*	2.1	4.5	11.9	31.9	68.6	151.4	366.4
2001.....	17.3	19.8	*	*	0.8	2.0	4.7	12.6	29.8	64.0	145.1	349.8
2000.....	18.0	16.9	*	*	*	1.8	5.1	11.7	29.2	67.0	158.4	387.6
1999.....	17.7	17.5	*	*	0.8	1.8	4.6	11.8	25.8	66.0	155.9	411.8

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Septicemia (A40-A41)												
All races 3/, both sexes												
2004.....	11.4	6.6	0.5	0.2	0.3	0.8	1.9	5.4	12.9	32.4	81.6	186.7
2003.....	11.7	6.9	0.5	0.2	0.4	0.8	2.1	5.3	13.1	32.6	85.0	202.5
2002.....	11.7	7.3	0.5	0.2	0.3	0.8	1.9	5.2	12.6	34.7	86.5	203.0
2001.....	11.3	7.7	0.7	0.2	0.3	0.7	1.8	5.0	12.3	32.8	82.3	205.9
2000.....	11.1	7.2	0.6	0.2	0.3	0.7	1.9	4.9	11.9	31.0	80.4	215.7
1999.....	11.0	7.5	0.6	0.2	0.3	0.7	1.8	4.6	11.4	31.2	79.4	220.7
All races 3/, male												
2004.....	10.4	8.0	0.5	0.1	0.3	0.8	2.0	5.9	13.8	35.8	93.3	203.9
2003.....	10.5	7.4	0.7	0.2	0.4	0.8	2.1	5.8	14.1	36.2	95.0	213.8
2002.....	10.6	7.7	0.5	0.2	0.3	0.8	2.0	5.7	13.7	38.0	99.0	213.9
2001.....	10.2	8.4	0.8	0.2	0.4	0.7	2.0	5.4	13.2	36.7	95.5	224.3
2000.....	9.8	7.1	0.7	0.2	0.3	0.8	2.0	5.6	13.0	34.5	91.9	223.0
1999.....	9.8	9.0	0.6	0.2	0.3	0.7	2.1	5.1	12.4	35.2	91.9	233.5
All races 3/, female												
2004.....	12.3	5.2	0.6	0.2	0.3	0.8	1.9	4.9	12.0	29.6	73.7	179.0
2003.....	12.9	6.5	0.4	0.2	0.4	0.8	2.0	4.8	12.2	29.5	78.3	197.5
2002.....	12.9	7.0	0.5	0.2	0.3	0.8	1.8	4.7	11.7	31.9	78.2	198.3
2001.....	12.4	7.0	0.6	0.1	0.3	0.7	1.7	4.7	11.4	29.5	73.7	198.3
2000.....	12.3	7.3	0.6	0.1	0.2	0.6	1.9	4.2	11.0	28.1	72.9	212.7
1999.....	12.2	5.8	0.5	0.2	0.3	0.6	1.6	4.1	10.5	28.0	71.3	215.5
White, both sexes												
2004.....	11.2	5.2	0.5	0.2	0.3	0.7	1.6	4.4	11.0	29.1	76.6	178.1
2003.....	11.5	4.6	0.4	0.2	0.3	0.6	1.7	4.3	11.2	28.9	80.1	191.1
2002.....	11.5	6.1	0.5	0.2	0.3	0.6	1.6	4.2	10.6	31.1	80.7	191.9
2001.....	11.1	5.4	0.7	0.2	0.3	0.5	1.4	4.0	10.3	29.3	77.1	195.3
2000.....	10.8	5.7	0.6	0.2	0.2	0.5	1.6	3.9	9.9	27.4	73.9	203.0
1999.....	10.6	5.5	0.5	0.2	0.3	0.6	1.4	3.6	9.4	27.3	72.8	205.9
White, male												
2004.....	10.2	6.3	0.5	0.2	0.3	0.6	1.6	4.8	11.7	32.2	87.6	196.5
2003.....	10.2	4.6	0.6	0.2	0.3	0.6	1.7	4.7	11.9	32.1	89.5	203.3
2002.....	10.3	6.2	0.5	0.3	0.3	0.7	1.6	4.7	11.7	34.1	91.5	202.9
2001.....	9.9	6.2	0.8	0.2	0.3	0.5	1.7	4.3	11.1	32.5	89.4	211.8
2000.....	9.4	5.8	0.6	0.2	0.2	0.6	1.7	4.4	10.9	30.8	84.6	209.8
1999.....	9.3	6.9	0.5	0.2	0.3	0.6	1.6	4.0	10.0	30.6	84.6	217.4

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Septicemia (A40-A41)												
White, female												
2004.....	12.2	4.0	0.5	0.2	0.2	0.7	1.5	4.0	10.3	26.5	69.2	169.9
2003.....	12.8	4.5	*	0.2	0.3	0.6	1.7	3.8	10.5	26.2	73.8	185.8
2002.....	12.8	6.0	0.4	0.2	0.3	0.6	1.5	3.8	9.6	28.6	73.4	187.1
2001.....	12.3	4.6	0.6	0.1	0.2	0.5	1.2	3.7	9.7	26.6	69.0	188.3
2000.....	12.1	5.6	0.5	*	0.1	0.4	1.5	3.3	9.0	24.6	66.8	200.2
1999.....	11.9	4.0	0.5	0.2	0.2	0.5	1.2	3.2	9.0	24.5	65.1	201.4
Black, both sexes												
2004.....	15.6	14.5	*	*	0.7	1.9	4.5	13.5	31.6	69.9	155.6	320.3
2003.....	16.3	18.6	0.9	*	0.6	1.9	4.9	13.7	32.2	73.0	157.5	372.0
2002.....	16.3	13.5	*	0.3	0.4	1.9	4.4	12.9	33.2	74.7	171.2	367.0
2001.....	15.8	19.3	0.9	*	0.7	1.8	4.6	13.4	32.1	71.8	160.2	364.8
2000.....	16.2	16.4	1.0	*	0.6	2.0	4.7	13.6	31.4	71.9	173.4	392.6
1999.....	16.1	17.7	*	*	0.6	1.5	4.9	12.9	30.3	74.4	168.3	412.3
Black, male												
2004.....	14.6	16.3	*	*	0.7	2.0	4.7	15.1	36.2	81.1	189.0	341.7
2003.....	15.3	21.1	*	*	*	2.1	5.0	15.3	37.8	85.7	189.0	404.5
2002.....	15.0	14.8	*	*	*	1.9	4.9	14.7	35.9	86.1	212.3	393.5
2001.....	14.9	20.1	*	*	0.8	1.8	4.7	14.9	36.9	85.6	195.1	424.4
2000.....	14.9	16.0	*	*	*	2.3	4.6	16.3	35.5	81.2	207.1	427.8
1999.....	15.3	19.8	*	*	*	1.6	5.8	15.1	37.1	90.3	201.7	453.5
Black, female												
2004.....	16.4	12.7	*	*	0.8	1.8	4.3	12.0	27.9	61.9	136.7	311.9
2003.....	17.2	16.1	*	*	0.7	1.6	4.9	12.3	27.8	63.9	139.7	359.3
2002.....	17.4	12.1	*	*	*	1.9	4.1	11.4	30.9	66.6	148.0	356.9
2001.....	16.7	18.6	*	*	*	1.8	4.6	12.1	28.3	62.0	140.6	342.7
2000.....	17.4	16.9	*	*	*	1.7	4.8	11.3	28.2	65.3	154.4	379.5
1999.....	16.9	15.4	*	*	*	1.5	4.2	11.1	24.8	63.2	149.6	396.9

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other zoonotic and bacterial diseases (A21-A22,A24-A32,A42-A44,A48-A49)

All races 3/, both sexes

2004.....	0.6	*	*	*	*	0.1	0.1	0.3	0.8	1.5	4.0	7.8
2003.....	0.5	*	*	*	*	*	0.1	0.3	0.8	1.6	3.7	7.0
2002.....	0.5	*	*	*	*	*	0.2	0.3	0.6	1.5	3.7	7.4
2001.....	0.5	*	*	*	*	0.1	0.1	0.3	0.6	1.4	3.5	7.5
2000.....	0.5	0.5	*	*	*	*	0.1	0.3	0.6	1.3	3.3	6.7
1999.....	0.6	0.6	*	*	*	0.1	0.2	0.4	0.7	2.0	4.3	10.3

All races 3/, male

2004.....	0.5	*	*	*	*	*	0.1	0.3	0.8	1.9	4.4	10.8
2003.....	0.6	*	*	*	*	*	0.1	0.4	0.9	2.0	4.5	9.3
2002.....	0.5	*	*	*	*	*	0.2	0.4	0.7	1.7	4.6	8.5
2001.....	0.5	*	*	*	*	*	0.1	0.4	0.8	1.7	4.3	8.9
2000.....	0.5	*	*	*	*	*	0.1	0.4	0.7	1.6	4.0	9.9
1999.....	0.6	*	*	*	*	0.1	0.2	0.4	0.8	2.4	5.4	13.1

All races 3/, female

2004.....	0.6	*	*	*	*	*	0.1	0.3	0.7	1.1	3.7	6.4
2003.....	0.5	*	*	*	*	*	*	0.3	0.6	1.3	3.1	5.9
2002.....	0.5	*	*	*	*	*	0.1	0.2	0.5	1.3	3.0	6.9
2001.....	0.5	*	*	*	*	*	0.1	0.2	0.5	1.2	2.9	6.9
2000.....	0.4	*	*	*	*	*	0.1	0.2	0.5	1.0	2.8	5.3
1999.....	0.6	*	*	*	*	*	0.2	0.3	0.6	1.6	3.6	9.2

White, both sexes

2004.....	0.6	*	*	*	*	*	0.1	0.2	0.7	1.4	3.8	7.7
2003.....	0.5	*	*	*	*	*	0.1	0.3	0.7	1.5	3.7	7.0
2002.....	0.5	*	*	*	*	*	0.1	0.3	0.5	1.4	3.6	7.3
2001.....	0.5	*	*	*	*	*	0.1	0.3	0.5	1.3	3.4	7.4
2000.....	0.5	*	*	*	*	*	0.1	0.3	0.5	1.2	3.2	6.4
1999.....	0.6	*	*	*	*	0.1	0.2	0.3	0.7	1.8	3.9	9.9

White, male

2004.....	0.5	*	*	*	*	*	*	0.3	0.8	1.7	4.1	10.9
2003.....	0.6	*	*	*	*	*	0.1	0.3	0.8	1.8	4.5	9.3
2002.....	0.5	*	*	*	*	*	0.2	0.3	0.6	1.6	4.5	8.7
2001.....	0.5	*	*	*	*	*	0.1	0.4	0.7	1.5	4.3	8.9
2000.....	0.5	*	*	*	*	*	*	0.4	0.6	1.4	3.9	9.5
1999.....	0.6	*	*	*	*	*	0.2	0.4	0.8	2.2	4.8	12.6

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other zoonotic and bacterial diseases (A21-A22,A24-A32,A42-A44,A48-A49)

White, female

2004.....	0.6	*	*	*	*	*	*	0.2	0.7	1.1	3.6	6.3
2003.....	0.5	*	*	*	*	*	*	0.2	0.6	1.3	3.1	6.0
2002.....	0.5	*	*	*	*	*	*	0.2	0.5	1.2	3.0	6.7
2001.....	0.5	*	*	*	*	*	*	0.2	0.4	1.1	2.7	6.8
2000.....	0.5	*	*	*	*	*	*	0.2	0.4	1.0	2.7	5.1
1999.....	0.6	*	*	*	*	*	0.2	0.3	0.5	1.5	3.3	8.8

Black, both sexes

2004.....	0.6	*	*	*	*	*	*	0.7	1.3	2.2	6.4	8.2
2003.....	0.5	*	*	*	*	*	*	0.7	1.4	2.2	4.4	7.6
2002.....	0.6	*	*	*	*	*	0.4	0.6	1.2	2.4	4.6	9.2
2001.....	0.6	*	*	*	*	*	*	0.6	1.5	2.5	4.7	8.0
2000.....	0.5	*	*	*	*	*	*	0.5	1.0	2.2	4.5	9.7
1999.....	0.8	*	*	*	*	*	*	0.8	1.3	3.5	8.8	15.9

Black, male

2004.....	0.7	*	*	*	*	*	*	*	1.7	3.9	7.8	*
2003.....	0.7	*	*	*	*	*	*	1.1	1.9	3.9	*	*
2002.....	0.6	*	*	*	*	*	*	*	*	3.0	6.6	*
2001.....	0.6	*	*	*	*	*	*	*	1.9	3.3	*	*
2000.....	0.6	*	*	*	*	*	*	*	*	3.1	*	*
1999.....	0.9	*	*	*	*	*	*	*	*	4.6	13.2	*

Black, female

2004.....	0.6	*	*	*	*	*	*	*	*	*	5.7	*
2003.....	0.4	*	*	*	*	*	*	*	*	*	4.0	*
2002.....	0.6	*	*	*	*	*	*	*	*	2.0	3.4	9.4
2001.....	0.6	*	*	*	*	*	*	*	*	*	4.5	*
2000.....	0.5	*	*	*	*	*	*	*	*	*	3.9	*
1999.....	0.7	*	*	*	*	*	*	*	*	2.7	6.4	14.9

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Viral diseases (A80-B34)												
All races 3/, both sexes												
2004.....	6.8	2.9	0.3	0.1	0.6	3.9	12.5	16.2	10.3	7.0	6.5	7.8
2003.....	7.1	2.9	0.4	0.2	0.6	4.2	13.7	16.7	10.0	7.7	7.5	9.1
2002.....	7.4	4.1	0.3	0.2	0.6	5.0	14.6	17.3	9.6	7.8	7.8	10.1
2001.....	7.4	3.0	0.4	0.2	0.7	5.6	15.0	16.0	9.7	7.9	7.2	8.3
2000.....	7.5	2.9	0.4	0.2	0.6	6.4	15.0	16.3	9.4	8.0	8.1	9.6
1999.....	7.6	3.5	0.5	0.3	0.6	7.1	15.6	15.1	8.8	8.1	8.3	11.6
All races 3/, male												
2004.....	9.5	3.2	0.4	0.1	0.7	4.8	17.7	24.6	15.9	9.4	7.3	9.0
2003.....	10.1	2.5	0.4	0.2	0.6	5.4	19.8	25.8	15.2	10.1	9.3	10.2
2002.....	10.6	4.6	0.4	0.1	0.6	6.3	21.3	26.9	14.5	10.0	9.6	11.7
2001.....	10.5	3.1	0.5	0.2	0.7	7.4	22.1	25.2	14.5	9.6	8.5	9.3
2000.....	10.8	2.9	0.4	0.2	0.7	8.3	22.3	25.8	14.3	9.9	10.4	10.8
1999.....	10.9	4.0	0.6	0.3	0.6	9.8	23.5	24.1	13.3	10.3	9.9	13.3
All races 3/, female												
2004.....	4.1	2.6	0.3	0.1	0.6	3.0	7.3	8.2	5.0	5.0	6.0	7.2
2003.....	4.2	3.3	0.4	0.2	0.5	3.0	7.6	7.9	5.1	5.7	6.3	8.7
2002.....	4.4	3.6	*	0.2	0.6	3.6	7.9	7.9	5.1	6.0	6.5	9.3
2001.....	4.3	2.8	0.3	0.2	0.8	3.7	7.9	7.1	5.3	6.4	6.4	7.9
2000.....	4.4	2.9	0.4	0.2	0.6	4.5	7.7	7.1	5.0	6.4	6.6	9.1
1999.....	4.3	3.1	0.4	0.2	0.6	4.4	7.8	6.5	4.7	6.4	7.2	10.9
White, both sexes												
2004.....	4.6	2.6	0.3	0.1	0.3	1.8	7.7	11.0	7.2	5.5	6.0	7.8
2003.....	4.9	2.3	0.4	0.1	0.3	2.2	8.4	11.5	6.9	6.1	7.0	9.3
2002.....	5.1	3.7	0.3	0.1	0.3	2.5	8.9	11.7	6.8	6.2	7.2	10.3
2001.....	5.0	2.3	0.3	0.1	0.3	2.8	9.2	10.7	6.9	6.4	6.7	8.4
2000.....	5.1	2.1	0.3	0.2	0.3	3.3	9.4	10.7	6.6	6.5	7.7	9.6
1999.....	5.2	2.8	0.4	0.2	0.3	4.1	9.7	9.8	6.1	6.6	7.9	11.6
White, male												
2004.....	6.8	2.8	0.3	*	0.3	2.6	11.9	17.3	11.1	7.2	6.8	8.8
2003.....	7.2	1.9	0.3	*	0.4	3.3	13.2	18.4	10.5	7.8	8.4	10.3
2002.....	7.5	4.1	0.4	*	0.4	3.7	14.1	18.7	10.2	7.7	8.8	11.7
2001.....	7.4	2.4	0.4	0.1	0.4	4.2	14.8	17.5	10.3	7.8	7.9	9.7
2000.....	7.5	2.2	*	0.1	0.4	4.9	14.9	17.5	9.9	7.6	9.7	10.9
1999.....	7.6	3.4	0.6	0.2	0.4	6.2	15.7	15.9	9.2	8.2	9.5	12.7

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Viral diseases (A80-B34)												
White, female												
2004.....	2.5	2.4	*	*	0.2	0.9	3.4	4.7	3.4	4.1	5.5	7.3
2003.....	2.7	2.7	0.4	*	0.3	1.1	3.5	4.6	3.5	4.7	5.9	8.9
2002.....	2.8	3.3	*	0.1	0.3	1.3	3.6	4.7	3.7	4.9	6.1	9.7
2001.....	2.7	2.2	*	0.1	0.3	1.4	3.6	3.9	3.8	5.2	5.9	7.9
2000.....	2.8	2.1	*	0.2	0.2	1.7	3.8	4.0	3.6	5.6	6.3	9.1
1999.....	2.8	2.1	*	0.2	0.3	1.8	3.7	3.8	3.3	5.3	6.8	11.1
Black, both sexes												
2004.....	21.5	4.2	*	0.4	2.4	17.5	46.3	57.8	38.9	19.3	10.4	7.3
2003.....	22.3	5.0	*	0.5	2.1	17.6	51.3	58.5	38.0	20.5	12.9	7.9
2002.....	23.5	5.8	*	0.3	2.2	21.2	55.1	62.4	33.6	21.6	13.6	7.4
2001.....	23.7	6.6	0.8	0.5	2.9	23.6	56.1	59.4	34.7	19.2	11.7	6.1
2000.....	24.1	7.3	0.9	0.4	2.5	26.9	55.6	61.7	32.7	20.2	12.8	8.8
1999.....	24.3	7.3	1.1	0.8	2.4	27.7	58.5	59.2	32.0	21.2	11.3	11.5
Black, male												
2004.....	29.4	*	*	*	2.5	19.6	62.0	87.1	64.7	31.3	12.5	*
2003.....	31.1	*	*	*	2.0	21.2	70.9	89.8	63.0	33.1	17.8	*
2002.....	33.0	6.4	*	*	2.2	25.1	76.9	98.0	55.7	34.5	20.1	*
2001.....	33.1	7.5	*	*	2.6	30.1	79.1	92.6	56.4	27.4	15.9	*
2000.....	34.3	7.0	*	*	2.4	32.9	80.5	98.6	54.8	32.5	19.4	*
1999.....	35.0	7.5	*	0.9	2.1	36.4	85.2	97.1	52.8	31.3	13.5	*
Black, female												
2004.....	14.2	*	*	*	2.3	15.6	32.3	32.6	18.0	10.7	9.1	*
2003.....	14.2	*	*	*	2.1	14.3	33.9	31.6	18.0	11.6	10.2	*
2002.....	14.9	*	*	*	2.2	17.7	35.9	31.9	15.9	12.5	9.9	*
2001.....	15.0	*	*	*	3.3	17.7	35.8	30.8	17.2	13.4	9.4	*
2000.....	14.8	7.6	*	*	2.6	21.5	33.6	30.0	14.9	11.6	9.0	8.6
1999.....	14.6	7.0	*	0.6	2.8	19.9	34.9	26.7	15.3	14.1	10.0	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Viral hepatitis (B15-B19)												
All races 3/, both sexes												
2004.....	1.8	*	*	*	*	0.2	1.4	5.3	4.3	3.4	3.6	2.7
2003.....	1.9	*	*	*	*	0.2	1.5	5.5	3.9	3.8	4.2	2.4
2002.....	2.0	*	*	*	0.1	0.2	1.6	5.8	4.0	4.2	4.6	3.9
2001.....	2.0	*	*	*	0.1	0.2	1.8	5.3	4.0	4.6	4.9	3.8
2000.....	1.9	*	*	*	*	0.2	1.7	5.0	3.7	4.6	5.3	4.0
1999.....	1.7	*	*	*	*	0.2	1.6	4.1	3.3	4.8	5.3	4.7
All races 3/, male												
2004.....	2.4	*	*	*	*	0.2	1.8	8.0	6.1	4.0	3.6	3.0
2003.....	2.5	*	*	*	*	0.2	2.1	8.3	5.4	4.2	4.8	2.8
2002.....	2.7	*	*	*	*	0.3	2.3	8.9	5.5	4.5	5.3	5.0
2001.....	2.5	*	*	*	*	0.2	2.5	8.2	5.3	4.8	5.3	4.4
2000.....	2.4	*	*	*	*	0.2	2.4	7.7	5.0	4.8	6.4	5.7
1999.....	2.2	*	*	*	*	0.3	2.3	6.3	4.3	5.1	6.6	6.9
All races 3/, female												
2004.....	1.3	*	*	*	*	0.1	1.0	2.7	2.5	3.0	3.5	2.5
2003.....	1.3	*	*	*	*	0.1	0.9	2.9	2.5	3.5	3.8	2.3
2002.....	1.4	*	*	*	*	0.2	1.0	2.8	2.7	3.9	4.1	3.3
2001.....	1.4	*	*	*	*	0.2	1.1	2.5	2.8	4.4	4.6	3.5
2000.....	1.4	*	*	*	*	0.2	1.1	2.4	2.6	4.5	4.7	3.3
1999.....	1.3	*	*	*	*	0.2	0.9	2.0	2.4	4.5	4.5	3.9
White, both sexes												
2004.....	1.8	*	*	*	*	0.1	1.4	5.2	3.8	3.1	3.2	2.5
2003.....	1.8	*	*	*	*	0.2	1.5	5.4	3.3	3.4	3.9	2.3
2002.....	2.0	*	*	*	0.1	0.2	1.6	5.6	3.5	3.7	4.4	3.7
2001.....	1.9	*	*	*	*	0.2	1.8	5.0	3.6	4.1	4.7	3.6
2000.....	1.8	*	*	*	*	0.2	1.7	4.6	3.2	4.2	5.1	3.9
1999.....	1.7	*	*	*	*	0.2	1.5	3.8	2.9	4.3	5.1	4.5
White, male												
2004.....	2.4	*	*	*	*	0.1	1.9	7.8	5.3	3.6	3.3	2.7
2003.....	2.4	*	*	*	*	0.2	2.1	8.2	4.7	3.8	4.4	2.8
2002.....	2.6	*	*	*	*	0.3	2.3	8.7	4.8	4.1	5.2	4.9
2001.....	2.5	*	*	*	*	0.2	2.5	7.8	4.8	4.5	5.4	4.5
2000.....	2.4	*	*	*	*	0.2	2.4	7.2	4.3	4.3	6.2	5.7
1999.....	2.2	*	*	*	*	0.3	2.3	5.7	3.9	4.8	6.5	6.5

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Viral hepatitis (B15-B19)												
White, female												
2004.....	1.2	*	*	*	*	*	0.9	2.6	2.3	2.7	3.2	2.4
2003.....	1.2	*	*	*	*	*	0.9	2.7	2.0	3.0	3.5	2.1
2002.....	1.3	*	*	*	*	*	0.9	2.7	2.4	3.3	3.9	3.2
2001.....	1.3	*	*	*	*	0.1	1.0	2.2	2.4	3.7	4.2	3.2
2000.....	1.3	*	*	*	*	0.1	1.0	2.1	2.2	4.1	4.5	3.2
1999.....	1.2	*	*	*	*	0.2	0.8	1.9	2.0	3.9	4.1	3.7
Black, both sexes												
2004.....	2.2	*	*	*	*	*	1.4	7.5	8.7	4.6	5.0	*
2003.....	2.2	*	*	*	*	*	1.7	7.4	8.0	5.4	6.0	*
2002.....	2.3	*	*	*	*	0.4	1.9	8.0	7.1	6.1	5.2	*
2001.....	2.2	*	*	*	*	*	2.1	7.9	6.7	5.6	5.7	*
2000.....	2.2	*	*	*	*	0.4	2.0	7.9	5.9	5.7	6.2	*
1999.....	2.0	*	*	*	*	*	2.1	6.9	5.0	6.9	6.1	6.4
Black, male												
2004.....	3.1	*	*	*	*	*	1.5	11.7	13.8	6.7	*	*
2003.....	3.0	*	*	*	*	*	2.4	11.5	11.8	6.1	6.2	*
2002.....	3.0	*	*	*	*	*	2.5	12.4	10.5	7.1	*	*
2001.....	2.7	*	*	*	*	*	2.9	11.8	8.9	4.9	*	*
2000.....	2.9	*	*	*	*	*	2.9	12.3	8.1	7.0	7.6	*
1999.....	2.6	*	*	*	*	*	3.0	11.6	6.2	5.9	*	*
Black, female												
2004.....	1.5	*	*	*	*	*	1.4	3.8	4.5	3.2	5.2	*
2003.....	1.6	*	*	*	*	*	1.1	4.0	5.0	4.9	5.9	*
2002.....	1.6	*	*	*	*	*	1.4	4.3	4.3	5.4	5.4	*
2001.....	1.8	*	*	*	*	*	1.4	4.7	4.9	6.0	6.8	*
2000.....	1.5	*	*	*	*	*	1.3	4.2	4.1	4.7	5.5	*
1999.....	1.5	*	*	*	*	*	1.3	2.8	4.1	7.6	6.5	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Human immunodeficiency virus (HIV) disease (B20-B24)												
All races 3/, both sexes												
2004.....	4.4	*	*	0.1	0.5	3.7	10.9	10.6	5.4	2.4	0.8	*
2003.....	4.7	*	*	0.1	0.4	4.0	12.0	10.9	5.4	2.4	0.7	*
2002.....	4.9	*	*	0.1	0.4	4.6	12.7	11.2	5.1	2.2	0.8	*
2001.....	5.0	*	*	0.1	0.6	5.3	13.0	10.5	5.2	2.1	0.7	*
2000.....	5.1	*	*	0.1	0.5	6.1	13.1	11.0	5.1	2.2	0.7	*
1999.....	5.3	*	0.2	0.2	0.5	6.8	13.8	10.7	4.8	2.2	0.6	*
All races 3/, male												
2004.....	6.6	*	*	*	0.5	4.5	15.7	16.3	9.0	4.0	1.4	*
2003.....	7.1	*	*	*	0.4	5.1	17.5	17.2	9.1	4.0	1.5	*
2002.....	7.4	*	*	*	0.4	5.9	18.8	17.7	8.5	3.9	1.4	*
2001.....	7.6	*	*	0.1	0.5	7.1	19.5	16.8	8.6	3.5	1.5	*
2000.....	7.9	*	*	0.1	0.5	8.0	19.8	17.8	8.7	3.8	1.3	*
1999.....	8.2	*	*	0.2	0.5	9.5	21.0	17.5	8.3	3.8	1.0	*
All races 3/, female												
2004.....	2.4	*	*	*	0.4	2.8	6.2	5.2	2.0	1.0	0.5	*
2003.....	2.4	*	*	*	0.4	2.8	6.5	4.8	2.1	1.0	0.3	*
2002.....	2.5	*	*	*	0.4	3.3	6.7	4.8	1.9	0.8	0.3	*
2001.....	2.5	*	*	*	0.6	3.5	6.7	4.4	2.0	0.9	*	*
2000.....	2.5	*	*	0.1	0.4	4.2	6.5	4.4	1.8	0.8	0.3	*
1999.....	2.5	*	*	0.2	0.5	4.1	6.7	4.1	1.6	0.8	0.3	*
White, both sexes												
2004.....	2.4	*	*	*	0.1	1.6	6.2	5.5	2.8	1.2	0.5	*
2003.....	2.5	*	*	*	0.2	2.0	6.8	5.8	2.9	1.2	0.4	*
2002.....	2.6	*	*	*	0.2	2.2	7.0	5.8	2.8	1.0	0.4	*
2001.....	2.7	*	*	*	0.2	2.6	7.3	5.5	2.8	1.0	0.3	*
2000.....	2.8	*	*	0.1	0.2	3.1	7.5	5.9	2.8	1.0	0.4	*
1999.....	3.0	*	*	0.1	0.2	3.7	8.0	5.7	2.5	1.1	0.3	*
White, male												
2004.....	3.9	*	*	*	0.2	2.4	9.9	9.2	5.0	2.2	0.9	*
2003.....	4.2	*	*	*	0.2	3.0	11.0	10.0	5.0	2.1	0.9	*
2002.....	4.3	*	*	*	0.2	3.3	11.6	9.8	4.9	1.8	0.6	*
2001.....	4.5	*	*	*	0.2	3.9	12.1	9.5	4.8	1.9	0.7	*
2000.....	4.7	*	*	*	0.2	4.6	12.4	10.1	5.0	1.9	0.7	*
1999.....	5.0	*	*	*	0.3	5.8	13.2	9.9	4.7	2.0	0.7	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Human immunodeficiency virus (HIV) disease (B20-B24)												
White, female												
2004.....	0.8	*	*	*	*	0.7	2.4	1.9	0.7	0.4	*	*
2003.....	0.9	*	*	*	0.1	0.9	2.5	1.7	0.9	0.4	*	*
2002.....	0.9	*	*	*	0.1	1.0	2.5	1.8	0.8	*	*	*
2001.....	0.9	*	*	*	0.2	1.2	2.4	1.5	0.8	0.3	*	*
2000.....	1.0	*	*	*	*	1.5	2.6	1.7	0.8	0.3	*	*
1999.....	1.0	*	*	*	0.2	1.6	2.7	1.6	0.6	0.3	*	*
Black, both sexes												
2004.....	18.8	*	*	*	2.2	17.1	44.6	50.0	29.7	13.6	4.5	*
2003.....	19.6	*	*	0.5	1.9	17.3	49.2	50.6	29.6	14.5	4.7	*
2002.....	20.8	*	*	*	2.0	20.6	52.8	53.9	26.1	14.4	5.6	*
2001.....	21.1	*	*	0.4	2.7	23.2	53.7	51.2	27.7	12.8	5.3	*
2000.....	21.4	*	*	0.3	2.1	26.3	53.5	53.4	26.0	13.5	4.8	*
1999.....	21.8	*	*	0.6	2.3	27.3	56.1	51.8	26.2	13.5	3.6	*
Black, male												
2004.....	25.9	*	*	*	2.2	19.2	60.3	74.9	50.3	23.4	7.5	*
2003.....	27.7	*	*	*	1.9	20.8	68.0	77.8	50.7	25.8	9.0	*
2002.....	29.5	*	*	*	1.9	24.5	74.0	85.0	44.4	26.3	12.0	*
2001.....	30.1	*	*	*	2.3	29.8	75.9	80.7	47.1	21.7	11.5	*
2000.....	31.0	*	*	*	2.0	32.3	77.5	86.0	46.0	24.6	9.4	*
1999.....	31.9	*	*	0.7	1.9	36.0	81.9	85.0	45.5	24.7	6.5	*
Black, female												
2004.....	12.4	*	*	*	2.2	15.1	30.6	28.6	13.1	6.5	*	*
2003.....	12.3	*	*	*	1.9	14.2	32.5	27.3	12.5	6.4	*	*
2002.....	12.8	*	*	*	2.1	17.1	34.0	27.2	11.3	6.0	*	*
2001.....	12.9	*	*	*	3.1	17.3	34.1	25.9	12.1	6.5	*	*
2000.....	12.8	*	*	*	2.2	20.8	32.2	25.3	10.0	5.8	*	*
1999.....	12.6	*	*	*	2.7	19.5	33.4	23.4	10.7	5.6	*	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Human immunodeficiency virus (HIV) disease resulting in infectious and parasitic diseases (B20)

All races 3/, both sexes

2004.....	1.7	*	*	*	0.2	1.5	4.4	4.2	1.9	0.9	0.3	*
2003.....	1.8	*	*	*	0.2	1.6	4.5	4.3	2.0	0.8	0.2	*
2002.....	1.9	*	*	*	0.2	1.9	5.1	4.5	1.9	0.7	0.2	*
2001.....	1.9	*	*	*	0.2	2.1	5.1	4.2	1.9	0.7	0.2	*
2000.....	2.0	*	*	*	0.2	2.4	5.3	4.4	1.8	0.8	0.3	*
1999.....	2.1	*	*	0.1	0.2	2.8	5.6	4.4	1.6	0.7	0.3	*

All races 3/, male

2004.....	2.6	*	*	*	0.2	1.9	6.3	6.2	3.2	1.5	0.5	*
2003.....	2.7	*	*	*	0.2	2.1	6.5	6.9	3.3	1.3	0.4	*
2002.....	2.9	*	*	*	0.2	2.3	7.5	7.2	3.0	1.3	0.5	*
2001.....	2.9	*	*	*	0.3	2.8	7.6	6.7	3.1	1.2	0.5	*
2000.....	3.1	*	*	*	0.2	3.2	7.8	7.1	2.9	1.4	0.5	*
1999.....	3.3	*	*	*	0.2	3.8	8.4	7.2	2.7	1.3	0.6	*

All races 3/, female

2004.....	1.0	*	*	*	0.2	1.2	2.4	2.2	0.7	0.4	*	*
2003.....	0.9	*	*	*	0.1	1.1	2.5	1.8	0.9	0.4	*	*
2002.....	1.0	*	*	*	0.2	1.4	2.7	1.9	0.8	0.3	*	*
2001.....	1.0	*	*	*	0.2	1.5	2.6	1.8	0.8	0.4	*	*
2000.....	1.0	*	*	*	0.2	1.6	2.8	1.7	0.7	0.2	*	*
1999.....	1.0	*	*	*	0.3	1.8	2.7	1.8	0.5	0.3	*	*

White, both sexes

2004.....	1.0	*	*	*	*	0.7	2.6	2.3	0.9	0.5	0.2	*
2003.....	1.0	*	*	*	0.1	0.8	2.7	2.4	1.1	0.4	*	*
2002.....	1.1	*	*	*	0.1	0.9	3.0	2.3	1.1	0.4	*	*
2001.....	1.1	*	*	*	0.1	1.1	3.0	2.2	1.1	0.4	*	*
2000.....	1.1	*	*	*	0.1	1.3	3.1	2.4	1.0	0.4	*	*
1999.....	1.2	*	*	*	0.1	1.6	3.2	2.4	0.9	0.4	*	*

White, male

2004.....	1.6	*	*	*	*	1.0	4.2	3.7	1.7	0.8	*	*
2003.....	1.7	*	*	*	*	1.3	4.4	4.0	1.8	0.7	*	*
2002.....	1.7	*	*	*	0.1	1.3	4.7	3.9	1.9	0.7	*	*
2001.....	1.8	*	*	*	*	1.7	4.8	3.8	1.8	0.6	*	*
2000.....	1.9	*	*	*	0.1	1.9	4.9	4.2	1.7	0.7	*	*
1999.....	2.0	*	*	*	*	2.4	5.2	4.0	1.6	0.7	*	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Human immunodeficiency virus (HIV) disease resulting in infectious and parasitic diseases (B20)

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
White, female												
2004.....	0.4	*	*	*	*	0.3	1.0	0.9	0.2	*	*	*
2003.....	0.4	*	*	*	*	0.4	1.1	0.8	0.4	*	*	*
2002.....	0.4	*	*	*	*	0.5	1.2	0.8	0.4	*	*	*
2001.....	0.4	*	*	*	*	0.5	1.1	0.7	0.4	*	*	*
2000.....	0.4	*	*	*	*	0.6	1.2	0.7	0.4	*	*	*
1999.....	0.4	*	*	*	*	0.7	1.2	0.7	0.2	*	*	*
Black, both sexes												
2004.....	7.2	*	*	*	1.0	7.0	16.9	18.8	10.9	5.4	*	*
2003.....	7.1	*	*	*	0.7	6.7	17.1	19.1	10.8	4.6	*	*
2002.....	8.0	*	*	*	0.8	8.0	20.1	21.8	9.0	4.6	*	*
2001.....	8.0	*	*	*	1.1	8.9	20.0	20.1	9.7	4.5	*	*
2000.....	8.3	*	*	*	0.9	10.2	21.3	20.4	8.8	4.8	2.2	*
1999.....	8.6	*	*	*	1.1	11.0	22.2	21.5	8.1	4.5	*	*
Black, male												
2004.....	9.7	*	*	*	0.9	7.6	23.0	27.4	17.8	8.9	*	*
2003.....	10.2	*	*	*	0.7	8.2	23.4	30.4	18.3	8.0	*	*
2002.....	11.4	*	*	*	0.8	9.5	28.6	35.0	14.9	7.7	*	*
2001.....	11.3	*	*	*	1.2	10.8	28.8	31.1	16.4	7.5	*	*
2000.....	11.9	*	*	*	0.8	13.0	30.4	33.0	15.0	9.2	*	*
1999.....	12.6	*	*	*	0.8	14.0	32.9	35.0	14.1	8.4	*	*
Black, female												
2004.....	4.9	*	*	*	1.1	6.3	11.6	11.4	5.3	2.9	*	*
2003.....	4.3	*	*	*	0.7	5.3	11.6	9.3	4.7	2.2	*	*
2002.....	4.9	*	*	*	0.8	6.7	12.7	10.5	4.3	2.3	*	*
2001.....	4.9	*	*	*	1.1	7.2	12.2	10.6	4.4	2.4	*	*
2000.....	5.0	*	*	*	1.0	7.7	13.1	9.6	3.8	*	*	*
1999.....	5.0	*	*	*	1.4	8.3	12.7	9.9	3.3	*	*	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Human immunodeficiency virus (HIV) disease resulting in malignant neoplasms (B21)

All races 3/, both sexes

2004.....	0.3	*	*	*	*	0.2	0.7	0.8	0.5	0.2	*	*
2003.....	0.3	*	*	*	*	0.2	0.8	0.7	0.5	0.2	*	*
2002.....	0.3	*	*	*	*	0.2	0.8	0.8	0.4	0.2	*	*
2001.....	0.3	*	*	*	*	0.3	0.9	0.7	0.5	0.2	*	*
2000.....	0.4	*	*	*	*	0.4	0.9	0.8	0.4	0.2	*	*
1999.....	0.4	*	*	*	*	0.5	1.0	0.8	0.5	0.2	*	*

All races 3/, male

2004.....	0.5	*	*	*	*	0.3	1.2	1.3	0.9	0.4	*	*
2003.....	0.5	*	*	*	*	0.3	1.3	1.1	0.8	0.4	*	*
2002.....	0.5	*	*	*	*	0.3	1.3	1.4	0.6	0.4	*	*
2001.....	0.6	*	*	*	*	0.5	1.5	1.2	0.9	0.3	*	*
2000.....	0.6	*	*	*	*	0.5	1.6	1.4	0.8	0.3	*	*
1999.....	0.7	*	*	*	*	0.9	1.8	1.4	0.9	0.3	*	*

All races 3/, female

2004.....	0.1	*	*	*	*	*	0.2	0.2	*	*	*	*
2003.....	0.1	*	*	*	*	*	0.2	0.3	0.2	*	*	*
2002.....	0.1	*	*	*	*	*	0.3	0.2	*	*	*	*
2001.....	0.1	*	*	*	*	0.2	0.2	0.2	0.2	*	*	*
2000.....	0.1	*	*	*	*	0.2	0.3	0.2	*	*	*	*
1999.....	0.1	*	*	*	*	0.2	0.3	0.2	*	*	*	*

White, both sexes

2004.....	0.2	*	*	*	*	0.1	0.5	0.5	0.3	*	*	*
2003.....	0.2	*	*	*	*	0.1	0.6	0.5	0.3	0.1	*	*
2002.....	0.2	*	*	*	*	0.2	0.5	0.5	0.3	0.1	*	*
2001.....	0.2	*	*	*	*	0.2	0.6	0.5	0.3	*	*	*
2000.....	0.3	*	*	*	*	0.2	0.7	0.6	0.3	*	*	*
1999.....	0.3	*	*	*	*	0.4	0.8	0.6	0.3	*	*	*

White, male

2004.....	0.4	*	*	*	*	0.2	0.9	1.0	0.6	*	*	*
2003.....	0.4	*	*	*	*	0.2	1.0	0.8	0.5	*	*	*
2002.....	0.4	*	*	*	*	0.3	1.0	1.0	0.5	*	*	*
2001.....	0.4	*	*	*	*	0.3	1.2	0.9	0.5	*	*	*
2000.....	0.5	*	*	*	*	0.4	1.2	1.1	0.6	*	*	*
1999.....	0.5	*	*	*	*	0.7	1.4	1.0	0.6	*	*	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Human immunodeficiency virus (HIV) disease resulting in malignant neoplasms (B21)

White, female

2004.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2003.....	0.1	*	*	*	*	*	*	0.1	*	*	*	*
2002.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	0.1	*	*	*	*	*
1999.....	0.1	*	*	*	*	*	0.2	0.1	*	*	*	*

Black, both sexes

2004.....	1.0	*	*	*	*	0.7	2.3	2.8	2.0	1.2	*	*
2003.....	1.0	*	*	*	*	0.7	2.2	2.6	2.1	*	*	*
2002.....	1.1	*	*	*	*	0.7	2.6	3.1	1.4	*	*	*
2001.....	1.1	*	*	*	*	1.1	2.6	2.5	2.5	*	*	*
2000.....	1.2	*	*	*	*	1.1	3.0	2.9	1.6	*	*	*
1999.....	1.2	*	*	*	*	1.5	3.0	2.8	1.9	*	*	*

Black, male

2004.....	1.6	*	*	*	*	0.9	3.5	4.5	3.8	*	*	*
2003.....	1.4	*	*	*	*	1.0	3.4	3.5	3.4	*	*	*
2002.....	1.6	*	*	*	*	0.9	4.0	4.9	2.4	*	*	*
2001.....	1.7	*	*	*	*	1.7	4.3	3.9	4.4	*	*	*
2000.....	1.9	*	*	*	*	1.6	5.0	4.9	3.2	*	*	*
1999.....	1.9	*	*	*	*	2.3	4.9	4.8	3.3	*	*	*

Black, female

2004.....	0.5	*	*	*	*	*	1.2	1.4	*	*	*	*
2003.....	0.6	*	*	*	*	*	1.1	1.8	*	*	*	*
2002.....	0.6	*	*	*	*	*	1.4	1.5	*	*	*	*
2001.....	0.5	*	*	*	*	*	1.1	1.2	*	*	*	*
2000.....	0.5	*	*	*	*	0.7	1.3	1.2	*	*	*	*
1999.....	0.5	*	*	*	*	0.8	1.2	1.0	*	*	*	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Human immunodeficiency virus (HIV) disease resulting in other specified diseases (B22)

All races 3/, both sexes

2004.....	0.4	*	*	*	*	0.3	1.0	1.1	0.6	0.2	*	*
2003.....	0.5	*	*	*	*	0.3	1.2	1.0	0.5	0.2	*	*
2002.....	0.5	*	*	*	*	0.5	1.2	1.0	0.6	0.2	*	*
2001.....	0.5	*	*	*	*	0.6	1.3	1.0	0.6	0.3	*	*
2000.....	0.6	*	*	*	*	0.7	1.5	1.2	0.6	0.3	*	*
1999.....	0.6	*	*	*	*	0.7	1.6	1.1	0.5	0.2	*	*

All races 3/, male

2004.....	0.7	*	*	*	*	0.4	1.6	1.8	1.0	0.3	*	*
2003.....	0.7	*	*	*	*	0.5	1.9	1.7	1.0	0.5	*	*
2002.....	0.8	*	*	*	*	0.7	1.9	1.6	1.0	0.4	*	*
2001.....	0.8	*	*	*	*	0.8	2.0	1.6	1.0	0.4	*	*
2000.....	0.9	*	*	*	*	1.0	2.4	2.0	1.1	0.4	*	*
1999.....	0.9	*	*	*	*	1.1	2.5	1.8	0.9	0.4	*	*

All races 3/, female

2004.....	0.2	*	*	*	*	0.3	0.5	0.5	0.2	*	*	*
2003.....	0.2	*	*	*	*	0.2	0.5	0.4	0.1	*	*	*
2002.....	0.2	*	*	*	*	0.3	0.5	0.4	*	*	*	*
2001.....	0.2	*	*	*	*	0.3	0.5	0.3	0.2	*	*	*
2000.....	0.2	*	*	*	*	0.4	0.6	0.4	*	*	*	*
1999.....	0.2	*	*	*	*	0.4	0.6	0.3	*	*	*	*

White, both sexes

2004.....	0.3	*	*	*	*	0.2	0.6	0.7	0.3	*	*	*
2003.....	0.3	*	*	*	*	0.2	0.8	0.6	0.3	0.2	*	*
2002.....	0.3	*	*	*	*	0.3	0.8	0.6	0.3	*	*	*
2001.....	0.3	*	*	*	*	0.3	0.8	0.6	0.4	0.1	*	*
2000.....	0.4	*	*	*	*	0.4	1.0	0.7	0.3	0.1	*	*
1999.....	0.4	*	*	*	*	0.5	1.0	0.7	0.3	0.1	*	*

White, male

2004.....	0.5	*	*	*	*	0.3	1.1	1.2	0.5	*	*	*
2003.....	0.5	*	*	*	*	0.3	1.3	1.2	0.5	0.3	*	*
2002.....	0.5	*	*	*	*	0.5	1.4	1.1	0.6	*	*	*
2001.....	0.5	*	*	*	*	0.5	1.3	1.1	0.6	*	*	*
2000.....	0.6	*	*	*	*	0.7	1.7	1.2	0.6	*	*	*
1999.....	0.6	*	*	*	*	0.7	1.8	1.3	0.5	*	*	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Human immunodeficiency virus (HIV) disease resulting in other specified diseases (B22)

White, female

2004.....	0.1	*	*	*	*	*	0.2	0.2	*	*	*	*
2003.....	0.1	*	*	*	*	*	0.2	*	*	*	*	*
2002.....	0.1	*	*	*	*	*	0.2	0.1	*	*	*	*
2001.....	0.1	*	*	*	*	*	0.2	*	*	*	*	*
2000.....	0.1	*	*	*	*	0.1	0.2	0.1	*	*	*	*
1999.....	0.1	*	*	*	*	0.2	0.2	0.1	*	*	*	*

Black, both sexes

2004.....	1.7	*	*	*	*	1.5	3.8	4.4	2.9	1.2	*	*
2003.....	1.7	*	*	*	*	1.3	4.3	4.2	3.1	*	*	*
2002.....	1.7	*	*	*	*	2.0	4.3	3.7	2.7	1.5	*	*
2001.....	1.9	*	*	*	*	2.2	4.9	4.0	2.9	1.7	*	*
2000.....	2.1	*	*	*	*	2.8	5.2	5.2	2.9	1.4	*	*
1999.....	2.1	*	*	*	*	2.7	5.8	4.0	3.1	1.5	*	*

Black, male

2004.....	2.3	*	*	*	*	1.5	5.3	6.8	5.2	*	*	*
2003.....	2.5	*	*	*	*	1.7	6.5	6.5	5.7	*	*	*
2002.....	2.5	*	*	*	*	2.4	6.3	5.8	5.0	2.9	*	*
2001.....	2.8	*	*	*	*	3.0	7.4	6.3	4.5	*	*	*
2000.....	3.2	*	*	*	*	3.6	7.6	8.8	5.5	*	*	*
1999.....	3.2	*	*	*	*	3.9	8.7	6.6	5.7	*	*	*

Black, female

2004.....	1.1	*	*	*	*	1.4	2.6	2.3	*	*	*	*
2003.....	0.9	*	*	*	*	1.0	2.4	2.1	*	*	*	*
2002.....	1.0	*	*	*	*	1.7	2.5	1.9	*	*	*	*
2001.....	1.1	*	*	*	*	1.5	2.7	2.0	1.6	*	*	*
2000.....	1.2	*	*	*	*	2.1	3.0	2.2	*	*	*	*
1999.....	1.1	*	*	*	*	1.6	3.2	1.8	*	*	*	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Human immunodeficiency virus (HIV) disease resulting in other conditions (B23)

All races 3/, both sexes

2004.....	0.7	*	*	*	0.1	0.5	1.7	1.8	1.0	0.5	0.2	*
2003.....	1.1	*	*	*	0.1	0.9	2.7	2.7	1.4	0.7	0.2	*
2002.....	1.2	*	*	*	0.1	1.0	3.0	2.8	1.4	0.6	0.3	*
2001.....	1.2	*	*	*	0.2	1.2	3.1	2.6	1.4	0.5	0.2	*
2000.....	1.2	*	*	*	0.1	1.3	2.9	2.6	1.5	0.6	0.2	*
1999.....	1.2	*	*	0.1	0.1	1.4	3.0	2.6	1.4	0.7	*	*

All races 3/, male

2004.....	1.1	*	*	*	*	0.7	2.4	2.7	1.7	0.8	*	*
2003.....	1.6	*	*	*	*	1.0	3.8	4.1	2.2	1.1	0.4	*
2002.....	1.7	*	*	*	*	1.3	4.2	4.2	2.3	1.0	0.4	*
2001.....	1.8	*	*	*	0.1	1.5	4.4	4.1	2.3	0.9	0.4	*
2000.....	1.8	*	*	*	*	1.6	4.2	4.0	2.4	0.9	0.5	*
1999.....	1.8	*	*	*	0.1	1.9	4.3	4.1	2.3	1.2	*	*

All races 3/, female

2004.....	0.4	*	*	*	*	0.4	1.1	0.8	0.4	*	*	*
2003.....	0.6	*	*	*	0.1	0.7	1.7	1.4	0.5	0.3	*	*
2002.....	0.7	*	*	*	*	0.7	1.8	1.4	0.5	0.3	*	*
2001.....	0.7	*	*	*	0.2	0.9	1.9	1.3	0.5	*	*	*
2000.....	0.7	*	*	*	0.1	1.1	1.6	1.2	0.6	0.3	*	*
1999.....	0.7	*	*	*	0.1	0.9	1.8	1.1	0.5	0.2	*	*

White, both sexes

2004.....	0.3	*	*	*	*	0.2	0.8	0.7	0.5	0.3	*	*
2003.....	0.5	*	*	*	*	0.3	1.3	1.2	0.7	0.3	*	*
2002.....	0.5	*	*	*	*	0.4	1.4	1.2	0.6	0.2	*	*
2001.....	0.6	*	*	*	*	0.5	1.5	1.2	0.6	0.2	*	*
2000.....	0.6	*	*	*	*	0.6	1.5	1.2	0.7	0.2	*	*
1999.....	0.6	*	*	*	*	0.6	1.5	1.1	0.6	0.3	*	*

White, male

2004.....	0.5	*	*	*	*	0.3	1.2	1.2	0.9	0.5	*	*
2003.....	0.8	*	*	*	*	0.4	2.0	2.0	1.1	0.5	*	*
2002.....	0.9	*	*	*	*	0.6	2.3	1.9	1.1	0.4	*	*
2001.....	0.9	*	*	*	*	0.7	2.4	1.9	1.2	0.4	*	*
2000.....	0.9	*	*	*	*	0.7	2.4	1.9	1.2	0.4	*	*
1999.....	0.9	*	*	*	*	0.9	2.3	1.8	1.2	0.5	*	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Human immunodeficiency virus (HIV) disease resulting in other conditions (B23)

White, female

2004.....	0.1	*	*	*	*	*	0.4	0.3	*	*	*	*
2003.....	0.2	*	*	*	*	0.2	0.6	0.4	0.2	*	*	*
2002.....	0.2	*	*	*	*	0.2	0.6	0.4	*	*	*	*
2001.....	0.2	*	*	*	*	0.3	0.7	0.4	*	*	*	*
2000.....	0.2	*	*	*	*	0.4	0.6	0.4	0.2	*	*	*
1999.....	0.2	*	*	*	*	0.3	0.6	0.4	*	*	*	*

Black, both sexes

2004.....	3.5	*	*	*	0.3	2.9	8.2	9.6	6.3	2.5	*	*
2003.....	5.3	*	*	*	0.6	4.4	12.9	14.3	8.0	4.6	*	*
2002.....	5.7	*	*	*	0.4	5.1	13.7	15.4	8.4	4.7	*	*
2001.....	5.7	*	*	*	0.7	5.9	14.2	14.4	7.9	3.4	2.1	*
2000.....	5.5	*	*	*	0.4	6.5	13.1	14.1	8.2	3.9	*	*
1999.....	5.8	*	*	*	0.6	6.7	14.4	14.5	8.3	4.6	*	*

Black, male

2004.....	5.0	*	*	*	*	3.5	10.9	15.1	10.3	4.7	*	*
2003.....	7.3	*	*	*	*	4.9	17.3	21.1	13.6	7.7	*	*
2002.....	8.0	*	*	*	*	6.3	18.6	23.6	14.2	8.6	*	*
2001.....	8.0	*	*	*	*	7.6	19.3	22.3	13.4	6.2	*	*
2000.....	7.8	*	*	*	*	7.6	18.3	22.2	13.6	6.9	*	*
1999.....	8.3	*	*	*	*	8.7	19.8	23.4	14.0	8.8	*	*

Black, female

2004.....	2.2	*	*	*	*	2.3	5.7	4.9	3.2	*	*	*
2003.....	3.6	*	*	*	0.7	4.0	9.0	8.4	3.5	2.4	*	*
2002.....	3.6	*	*	*	*	4.0	9.5	8.3	3.8	*	*	*
2001.....	3.6	*	*	*	1.0	4.4	9.8	7.5	3.4	*	*	*
2000.....	3.5	*	*	*	*	5.5	8.4	7.2	3.9	*	*	*
1999.....	3.6	*	*	*	0.7	4.9	9.6	6.8	3.7	*	*	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Unspecified human immunodeficiency virus (HIV) disease (B24)												
All races 3/, both sexes												
2004.....	1.2	*	*	*	0.1	1.0	3.1	2.8	1.4	0.5	0.2	*
2003.....	1.0	*	*	*	0.1	1.0	2.8	2.2	1.1	0.5	0.2	*
2002.....	1.0	*	*	*	0.1	1.0	2.7	2.1	0.9	0.4	*	*
2001.....	1.0	*	*	*	0.1	1.1	2.7	2.0	0.8	0.4	*	*
2000.....	1.0	*	*	*	0.1	1.3	2.5	2.1	0.9	0.4	*	*
1999.....	1.0	*	*	*	0.1	1.3	2.6	1.8	0.8	0.4	*	*
All races 3/, male												
2004.....	1.7	*	*	*	0.1	1.2	4.3	4.2	2.3	0.9	*	*
2003.....	1.5	*	*	*	*	1.3	4.0	3.4	1.8	0.8	*	*
2002.....	1.5	*	*	*	*	1.2	3.9	3.3	1.5	0.7	*	*
2001.....	1.5	*	*	*	*	1.5	4.0	3.4	1.4	0.7	*	*
2000.....	1.5	*	*	*	0.1	1.6	3.8	3.3	1.6	0.7	*	*
1999.....	1.5	*	*	*	*	1.8	4.0	3.0	1.5	0.7	*	*
All races 3/, female												
2004.....	0.7	*	*	*	0.1	0.9	1.9	1.4	0.5	0.2	*	*
2003.....	0.5	*	*	*	0.1	0.7	1.6	1.0	0.4	0.2	*	*
2002.....	0.5	*	*	*	0.1	0.8	1.4	0.9	0.3	*	*	*
2001.....	0.5	*	*	*	0.1	0.7	1.4	0.7	0.3	*	*	*
2000.....	0.5	*	*	*	*	0.9	1.2	0.8	0.2	*	*	*
1999.....	0.5	*	*	*	*	0.8	1.3	0.7	0.3	*	*	*
White, both sexes												
2004.....	0.6	*	*	*	*	0.4	1.6	1.3	0.7	0.2	*	*
2003.....	0.5	*	*	*	*	0.5	1.4	1.1	0.6	0.2	*	*
2002.....	0.5	*	*	*	*	0.5	1.3	1.1	0.5	0.2	*	*
2001.....	0.5	*	*	*	*	0.5	1.4	1.0	0.4	0.2	*	*
2000.....	0.5	*	*	*	*	0.6	1.3	1.0	0.5	0.2	*	*
1999.....	0.5	*	*	*	*	0.7	1.5	1.0	0.4	0.2	*	*
White, male												
2004.....	1.0	*	*	*	*	0.6	2.5	2.2	1.3	0.5	*	*
2003.....	0.9	*	*	*	*	0.7	2.2	1.9	1.1	0.3	*	*
2002.....	0.8	*	*	*	*	0.7	2.2	1.8	0.8	0.3	*	*
2001.....	0.8	*	*	*	*	0.7	2.4	1.8	0.7	0.5	*	*
2000.....	0.8	*	*	*	*	0.9	2.2	1.8	0.9	0.4	*	*
1999.....	0.9	*	*	*	*	1.1	2.5	1.7	0.8	0.4	*	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Unspecified human immunodeficiency virus (HIV) disease (B24)

White, female

2004.....	0.2	*	*	*	*	0.2	0.7	0.4	0.2	*	*	*
2003.....	0.2	*	*	*	*	0.2	0.5	0.3	*	*	*	*
2002.....	0.2	*	*	*	*	0.2	0.4	0.4	*	*	*	*
2001.....	0.1	*	*	*	*	0.2	0.4	0.2	*	*	*	*
2000.....	0.2	*	*	*	*	0.3	0.5	0.3	*	*	*	*
1999.....	0.2	*	*	*	*	0.3	0.5	0.3	*	*	*	*

Black, both sexes

2004.....	5.4	*	*	*	0.6	5.1	13.4	14.4	7.6	3.3	*	*
2003.....	4.5	*	*	*	0.4	4.2	12.6	10.5	5.6	3.1	*	*
2002.....	4.3	*	*	*	0.5	4.7	12.0	10.0	4.5	2.8	*	*
2001.....	4.4	*	*	*	0.6	5.1	12.0	10.3	4.7	2.3	*	*
2000.....	4.3	*	*	*	0.5	5.7	11.0	10.7	4.5	2.5	*	*
1999.....	4.1	*	*	*	0.4	5.5	10.8	9.1	4.8	2.0	*	*

Black, male

2004.....	7.3	*	*	*	*	5.7	17.6	21.1	13.2	5.7	*	*
2003.....	6.3	*	*	*	*	5.0	17.4	16.2	9.7	6.1	*	*
2002.....	6.0	*	*	*	*	5.3	16.5	15.8	8.0	5.1	*	*
2001.....	6.3	*	*	*	*	6.8	16.1	17.0	8.4	3.9	*	*
2000.....	6.3	*	*	*	*	6.6	16.2	17.1	8.7	4.5	*	*
1999.....	5.9	*	*	*	*	7.1	15.6	15.1	8.4	3.4	*	*

Black, female

2004.....	3.7	*	*	*	0.6	4.6	9.6	8.6	3.0	*	*	*
2003.....	2.8	*	*	*	*	3.4	8.4	5.6	2.3	*	*	*
2002.....	2.8	*	*	*	*	4.2	8.0	5.1	1.7	*	*	*
2001.....	2.7	*	*	*	0.7	3.5	8.3	4.5	1.8	*	*	*
2000.....	2.6	*	*	*	*	4.9	6.4	5.1	*	*	*	*
1999.....	2.4	*	*	*	*	4.0	6.6	3.9	2.0	*	*	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

All other and unspecified viral diseases (A81,A85.0-A85.1,A85.8,A86-A89,B01,B04,B07-B09,B25-B34)

All races 3/, both sexes

2004.....	0.3	2.3	0.3	0.1	0.1	0.1	0.1	0.2	0.5	0.8	1.2	2.0
2003.....	0.3	2.2	0.4	0.1	0.1	0.1	0.1	0.2	0.5	1.0	1.3	2.8
2002.....	0.4	3.2	0.3	0.1	0.1	0.1	0.2	0.2	0.4	0.9	1.3	2.9
2001.....	0.3	2.1	0.3	0.1	0.1	0.1	0.1	0.2	0.4	0.9	1.0	2.2
2000.....	0.4	1.8	0.3	0.1	0.1	0.1	0.1	0.2	0.5	1.0	1.4	3.3
1999.....	0.4	2.7	0.3	0.1	0.1	0.1	0.2	0.3	0.6	0.9	1.7	4.0

All races 3/, male

2004.....	0.3	2.7	0.3	*	0.1	*	0.1	0.2	0.6	0.9	1.5	2.3
2003.....	0.3	1.8	0.4	*	0.1	*	0.2	0.2	0.6	1.1	1.5	3.0
2002.....	0.4	3.6	0.4	*	0.1	0.1	0.2	0.3	0.4	0.9	1.6	2.8
2001.....	0.3	2.1	0.4	*	0.1	*	0.2	0.2	0.5	0.9	1.2	2.4
2000.....	0.4	1.9	0.3	*	0.1	0.1	0.1	0.2	0.5	1.2	1.8	3.1
1999.....	0.4	2.8	0.4	*	*	*	0.2	0.3	0.6	1.1	1.6	3.9

All races 3/, female

2004.....	0.3	1.9	0.3	*	0.1	*	0.1	0.2	0.4	0.7	1.0	1.8
2003.....	0.3	2.6	0.4	*	*	*	0.1	0.2	0.4	0.8	1.2	2.7
2002.....	0.4	2.8	*	0.1	*	0.1	0.2	0.2	0.4	0.9	1.2	2.9
2001.....	0.3	2.0	0.3	0.1	*	*	0.1	0.2	0.4	0.9	0.9	2.1
2000.....	0.4	1.7	*	*	*	*	0.1	0.2	0.5	0.9	1.2	3.4
1999.....	0.4	2.5	*	*	*	0.1	0.1	0.3	0.6	0.8	1.8	4.0

White, both sexes

2004.....	0.3	2.1	0.2	0.1	0.1	0.1	0.1	0.2	0.5	0.8	1.3	2.0
2003.....	0.4	1.8	0.3	0.1	0.1	0.1	0.1	0.2	0.5	1.0	1.4	3.0
2002.....	0.4	3.1	0.3	0.1	0.1	0.1	0.2	0.2	0.4	1.0	1.4	3.1
2001.....	0.3	1.7	0.3	0.1	0.1	*	0.1	0.2	0.5	1.0	1.1	2.3
2000.....	0.4	1.5	0.3	0.1	0.1	0.1	0.1	0.2	0.5	1.1	1.4	3.4
1999.....	0.4	2.3	0.3	0.1	0.1	0.1	0.2	0.3	0.6	1.0	1.8	4.2

White, male

2004.....	0.3	2.3	*	*	*	*	0.1	0.2	0.6	1.0	1.7	2.4
2003.....	0.4	1.4	*	*	0.1	*	0.1	0.2	0.6	1.2	1.5	3.2
2002.....	0.4	3.2	0.3	*	*	*	0.2	0.2	0.4	1.0	1.6	2.9
2001.....	0.3	1.8	0.4	*	*	*	0.1	0.2	0.5	1.0	1.3	2.6
2000.....	0.4	1.6	*	*	0.1	*	0.1	0.2	0.6	1.2	1.8	3.3
1999.....	0.4	2.6	0.4	*	*	*	0.2	0.3	0.6	1.1	1.7	3.9

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other and unspecified infectious and parasitic diseases and their sequelae (B35-B99)

All races 3/, both sexes

2004.....	1.5	1.0	*	*	0.1	0.2	0.7	2.4	2.8	4.4	7.1	9.7
2003.....	1.5	1.1	*	*	0.1	0.2	0.7	2.3	2.7	4.2	7.1	9.0
2002.....	1.3	1.3	*	0.1	0.1	0.3	0.6	1.7	2.2	3.7	6.4	8.3
2001.....	1.1	1.6	*	0.0	0.1	0.2	0.5	1.2	1.9	3.6	5.4	8.2
2000.....	1.0	1.3	*	0.1	0.1	0.2	0.5	1.0	1.7	3.3	5.7	9.0
1999.....	1.4	1.6	0.1	0.1	0.1	0.4	0.8	1.6	2.4	4.7	7.4	10.0

All races 3/, male

2004.....	1.8	1.3	*	*	0.1	0.2	0.9	3.5	3.8	5.1	8.4	12.9
2003.....	1.7	1.3	*	*	*	0.3	0.9	3.2	3.5	4.7	8.3	10.4
2002.....	1.5	1.5	*	*	*	0.3	0.9	2.5	2.9	4.7	7.9	11.9
2001.....	1.2	1.6	*	*	0.1	0.3	0.7	1.6	2.3	3.8	7.1	10.1
2000.....	1.1	1.5	*	*	*	0.2	0.7	1.3	2.0	4.3	6.9	12.6
1999.....	1.6	1.9	*	*	0.1	0.4	1.0	1.9	2.7	5.4	9.4	13.1

All races 3/, female

2004.....	1.3	*	*	*	*	0.2	0.5	1.4	2.0	3.8	6.2	8.3
2003.....	1.3	*	*	*	*	0.2	0.5	1.4	1.9	3.9	6.3	8.3
2002.....	1.0	1.2	*	*	*	0.2	0.4	0.9	1.6	2.9	5.5	6.8
2001.....	1.0	1.6	*	*	*	0.2	0.4	0.8	1.5	3.5	4.4	7.4
2000.....	0.9	1.1	*	*	*	0.2	0.3	0.7	1.5	2.5	4.9	7.6
1999.....	1.3	1.2	*	*	*	0.4	0.6	1.3	2.1	4.0	6.2	8.8

White, both sexes

2004.....	1.6	0.8	*	*	0.1	0.2	0.7	2.5	2.7	4.2	6.8	9.6
2003.....	1.5	0.9	*	*	*	0.2	0.7	2.3	2.4	4.2	6.7	8.9
2002.....	1.3	1.1	*	0.1	0.1	0.2	0.6	1.6	2.0	3.6	6.4	8.3
2001.....	1.1	1.2	*	*	0.1	0.2	0.5	1.1	1.7	3.5	5.4	7.9
2000.....	1.0	1.0	*	*	0.1	0.1	0.4	0.9	1.5	3.2	5.6	8.9
1999.....	1.3	1.2	*	0.1	0.1	0.3	0.6	1.2	2.0	4.4	7.2	9.6

White, male

2004.....	1.8	*	*	*	0.1	0.2	0.8	3.5	3.5	4.9	8.1	13.2
2003.....	1.7	*	*	*	*	0.2	0.9	3.2	3.2	4.5	7.9	10.5
2002.....	1.5	*	*	*	*	0.2	0.8	2.4	2.6	4.6	7.7	11.8
2001.....	1.2	*	*	*	*	0.2	0.6	1.4	2.1	3.7	6.9	9.9
2000.....	1.1	1.3	*	*	*	0.2	0.6	1.1	1.8	4.1	6.5	12.0
1999.....	1.5	1.6	*	*	*	0.3	0.9	1.6	2.4	5.1	9.0	12.0

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other and unspecified infectious and parasitic diseases and their sequelae (B35-B99)

White, female

2004.....	1.3	*	*	*	*	0.1	0.5	1.4	1.9	3.5	5.9	7.9
2003.....	1.3	*	*	*	*	*	0.4	1.3	1.7	3.9	5.9	8.2
2002.....	1.0	*	*	*	*	0.2	0.3	0.8	1.4	2.8	5.5	6.7
2001.....	1.0	1.4	*	*	*	*	0.3	0.7	1.3	3.3	4.3	7.1
2000.....	0.9	*	*	*	*	*	0.3	0.6	1.3	2.4	4.9	7.6
1999.....	1.2	*	*	*	*	0.2	0.4	0.9	1.7	3.7	6.0	8.6

Black, both sexes

2004.....	1.5	*	*	*	*	0.4	1.2	2.4	4.3	5.4	9.2	13.3
2003.....	1.5	*	*	*	*	0.5	1.1	3.2	4.2	4.0	9.6	9.0
2002.....	1.5	*	*	*	*	0.6	1.3	2.9	4.1	3.7	6.4	9.5
2001.....	1.3	3.7	*	*	*	0.6	1.1	2.0	3.3	5.1	5.7	11.4
2000.....	1.3	3.3	*	*	*	0.6	1.3	2.0	3.0	4.5	7.1	10.9
1999.....	2.2	3.8	*	*	*	1.3	2.5	4.1	5.1	7.5	10.0	14.3

Black, male

2004.....	1.8	*	*	*	*	*	1.4	3.7	6.6	5.5	11.7	*
2003.....	1.8	*	*	*	*	*	1.3	4.1	5.5	5.3	11.0	*
2002.....	1.8	*	*	*	*	0.8	1.8	4.2	6.3	4.9	7.2	*
2001.....	1.5	*	*	*	*	*	1.6	3.0	4.1	5.9	6.8	*
2000.....	1.5	*	*	*	*	*	1.7	2.6	3.8	6.3	10.3	*
1999.....	2.3	*	*	*	*	1.4	2.5	4.6	5.1	8.7	13.5	*

Black, female

2004.....	1.3	*	*	*	*	*	0.9	1.2	2.4	5.3	7.9	13.0
2003.....	1.4	*	*	*	*	*	1.0	2.4	3.1	3.1	8.8	8.9
2002.....	1.1	*	*	*	*	*	0.9	1.8	2.4	2.9	6.0	*
2001.....	1.1	*	*	*	*	*	0.7	1.2	2.7	4.5	5.1	11.8
2000.....	1.1	*	*	*	*	*	0.9	1.5	2.4	3.3	5.3	9.0
1999.....	2.2	*	*	*	*	1.1	2.4	3.7	5.1	6.7	8.1	11.4

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Mycoses (B35-B49)												
All races 3/, both sexes												
2004.....	0.4	0.9	*	*	*	0.1	0.2	0.3	0.7	1.1	2.0	2.5
2003.....	0.4	1.0	*	*	*	0.1	0.2	0.3	0.6	1.3	2.1	2.4
2002.....	0.4	1.2	*	*	*	0.1	0.2	0.4	0.7	1.4	2.5	2.5
2001.....	0.4	1.4	*	*	0.1	0.1	0.2	0.3	0.7	1.3	2.2	3.0
2000.....	0.4	1.3	*	*	*	0.1	0.2	0.4	0.8	1.5	2.2	3.2
1999.....	0.7	1.4	*	*	0.1	0.2	0.3	0.7	1.2	2.3	3.3	3.9
All races 3/, male												
2004.....	0.4	1.0	*	*	*	0.1	0.2	0.4	0.8	1.3	2.7	3.8
2003.....	0.4	1.1	*	*	*	0.1	0.2	0.3	0.7	1.3	2.9	3.3
2002.....	0.5	1.2	*	*	*	0.2	0.3	0.5	1.0	2.0	3.5	4.2
2001.....	0.4	1.4	*	*	*	*	0.2	0.4	0.8	1.4	3.0	3.8
2000.....	0.5	1.5	*	*	*	0.1	0.3	0.4	0.9	2.0	2.9	4.9
1999.....	0.8	1.6	*	*	*	0.2	0.4	0.7	1.4	2.9	4.6	6.6
All races 3/, female												
2004.....	0.3	*	*	*	*	*	0.1	0.2	0.5	1.0	1.5	1.9
2003.....	0.3	*	*	*	*	*	0.1	0.3	0.5	1.2	1.6	2.0
2002.....	0.3	1.1	*	*	*	*	0.1	0.2	0.5	0.9	1.7	1.7
2001.....	0.4	1.5	*	*	*	*	0.1	0.2	0.5	1.2	1.6	2.6
2000.....	0.4	*	*	*	*	*	0.1	0.3	0.7	1.1	1.8	2.6
1999.....	0.6	1.1	*	*	*	0.2	0.2	0.6	1.0	1.9	2.4	2.9
White, both sexes												
2004.....	0.3	0.7	*	*	*	*	0.1	0.2	0.6	1.0	1.8	2.4
2003.....	0.4	0.7	*	*	*	*	0.1	0.3	0.5	1.2	2.0	2.3
2002.....	0.4	0.9	*	*	*	0.1	0.1	0.3	0.6	1.3	2.4	2.4
2001.....	0.4	1.0	*	*	*	0.1	0.1	0.3	0.6	1.1	2.0	2.6
2000.....	0.4	1.0	*	*	*	*	0.2	0.3	0.7	1.4	2.1	3.0
1999.....	0.6	1.0	*	*	*	0.1	0.2	0.6	1.0	2.2	3.1	3.5
White, male												
2004.....	0.4	*	*	*	*	*	0.2	0.3	0.7	1.2	2.5	4.0
2003.....	0.4	*	*	*	*	*	0.2	0.3	0.6	1.3	2.8	3.4
2002.....	0.5	*	*	*	*	*	0.2	0.4	0.8	1.8	3.4	4.1
2001.....	0.4	*	*	*	*	*	0.2	0.4	0.7	1.2	2.9	3.4
2000.....	0.5	*	*	*	*	*	0.2	0.4	0.9	1.8	2.7	4.4
1999.....	0.7	1.4	*	*	*	0.2	0.3	0.6	1.3	2.7	4.4	5.7

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Mycoses (B35-B49)												
White, female												
2004.....	0.3	*	*	*	*	*	*	0.2	0.5	0.9	1.3	1.7
2003.....	0.3	*	*	*	*	*	*	0.2	0.4	1.2	1.5	1.7
2002.....	0.3	*	*	*	*	*	*	0.2	0.4	0.8	1.7	1.7
2001.....	0.3	*	*	*	*	*	0.1	0.2	0.5	1.1	1.5	2.2
2000.....	0.4	*	*	*	*	*	*	0.3	0.6	1.0	1.8	2.4
1999.....	0.5	*	*	*	*	0.1	0.1	0.5	0.8	1.7	2.2	2.6
Black, both sexes												
2004.....	0.5	*	*	*	*	*	0.4	0.6	1.4	1.7	4.1	*
2003.....	0.6	*	*	*	*	*	0.4	0.7	1.5	1.6	3.5	*
2002.....	0.6	*	*	*	*	0.4	0.5	1.0	1.6	2.2	2.9	*
2001.....	0.6	3.5	*	*	*	*	0.4	0.7	1.6	2.8	3.5	7.1
2000.....	0.6	*	*	*	*	*	0.5	0.8	1.5	2.5	3.1	6.6
1999.....	1.0	3.5	*	*	*	0.6	0.8	1.4	2.2	4.1	5.1	8.9
Black, male												
2004.....	0.6	*	*	*	*	*	*	0.9	2.2	*	*	*
2003.....	0.6	*	*	*	*	*	*	*	2.0	*	*	*
2002.....	0.9	*	*	*	*	*	*	1.7	2.4	3.4	*	*
2001.....	0.8	*	*	*	*	*	*	*	1.9	3.9	*	*
2000.....	0.7	*	*	*	*	*	*	1.0	*	3.2	*	*
1999.....	1.1	*	*	*	*	*	1.0	1.8	2.3	4.9	*	*
Black, female												
2004.....	0.5	*	*	*	*	*	*	*	*	*	3.6	*
2003.....	0.5	*	*	*	*	*	*	*	*	*	*	*
2002.....	0.5	*	*	*	*	*	*	*	*	*	*	*
2001.....	0.5	*	*	*	*	*	*	*	*	*	*	*
2000.....	0.6	*	*	*	*	*	*	*	1.5	2.1	*	*
1999.....	0.9	*	*	*	*	*	*	1.1	2.2	3.6	4.6	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Protozoal diseases (B50-B64)												
All races 3/, both sexes												
2004.....	0.1	*	*	*	*	*	0.1	0.1	0.1	0.2	0.2	*
2003.....	0.1	*	*	*	*	*	0.1	0.1	0.1	0.1	0.2	*
2002.....	0.1	*	*	*	*	0.1	0.1	0.1	0.1	0.2	0.2	*
2001.....	0.1	*	*	*	*	0.1	0.1	0.1	0.2	0.3	0.2	*
2000.....	0.1	*	*	*	*	*	0.1	0.1	0.2	0.2	0.2	*
1999.....	0.1	*	*	*	*	0.1	0.2	0.1	0.2	0.3	0.3	*
All races 3/, male												
2004.....	0.1	*	*	*	*	*	0.1	0.1	*	0.3	*	*
2003.....	0.1	*	*	*	*	*	*	0.1	0.2	*	*	*
2002.....	0.1	*	*	*	*	*	0.1	0.1	0.2	*	*	*
2001.....	0.1	*	*	*	*	*	0.1	0.1	0.3	0.4	*	*
2000.....	0.1	*	*	*	*	*	0.2	0.1	0.3	0.4	*	*
1999.....	0.1	*	*	*	*	*	0.2	0.2	0.2	0.3	*	*
All races 3/, female												
2004.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2003.....	0.1	*	*	*	*	*	*	*	*	*	*	*
2002.....	0.1	*	*	*	*	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*	*	0.2	0.3	*	*
2000.....	0.1	*	*	*	*	*	*	*	*	*	*	*
1999.....	0.1	*	*	*	*	*	0.1	*	*	0.2	0.3	*
White, both sexes												
2004.....	0.1	*	*	*	*	*	*	0.1	*	0.2	0.2	*
2003.....	0.1	*	*	*	*	*	*	0.1	0.1	0.1	*	*
2002.....	0.1	*	*	*	*	*	0.1	0.1	0.1	0.2	0.2	*
2001.....	0.1	*	*	*	*	*	0.1	0.1	0.2	0.4	0.2	*
2000.....	0.1	*	*	*	*	*	0.1	*	0.2	0.2	0.2	*
1999.....	0.1	*	*	*	*	*	0.1	0.1	0.1	0.3	0.3	*
White, male												
2004.....	0.1	*	*	*	*	*	*	0.1	*	0.3	*	*
2003.....	0.1	*	*	*	*	*	*	*	*	*	*	*
2002.....	0.1	*	*	*	*	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	0.1	*	0.2	0.4	*	*
2000.....	0.1	*	*	*	*	*	0.1	*	0.3	0.4	*	*
1999.....	0.1	*	*	*	*	*	0.2	0.1	*	0.3	*	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pneumocystosis (B59)												
All races 3/, both sexes												
2004.....	0.1	*	*	*	*	*	0.1	0.1	0.1	0.1	0.2	*
2003.....	0.1	*	*	*	*	*	0.1	0.1	0.1	0.1	0.2	*
2002.....	0.1	*	*	*	*	0.1	0.1	0.1	0.1	0.2	*	*
2001.....	0.1	*	*	*	*	0.1	0.1	0.1	0.2	0.3	0.2	*
2000.....	0.1	*	*	*	*	*	0.1	0.1	0.2	0.2	0.2	*
1999.....	0.1	*	*	*	*	0.1	0.1	0.1	0.1	0.2	0.2	*
All races 3/, male												
2004.....	0.1	*	*	*	*	*	0.1	*	*	*	*	*
2003.....	0.1	*	*	*	*	*	*	0.1	*	*	*	*
2002.....	0.1	*	*	*	*	*	0.1	0.1	0.2	*	*	*
2001.....	0.1	*	*	*	*	*	0.1	0.1	0.2	0.3	*	*
2000.....	0.1	*	*	*	*	*	0.1	0.1	0.2	0.3	*	*
1999.....	0.1	*	*	*	*	*	0.1	0.2	*	0.3	*	*
All races 3/, female												
2004.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2003.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2002.....	0.1	*	*	*	*	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*	*	*	0.3	*	*
2000.....	0.1	*	*	*	*	*	*	*	*	*	*	*
1999.....	0.1	*	*	*	*	*	0.1	*	*	0.2	*	*
White, both sexes												
2004.....	0.0	*	*	*	*	*	*	0.1	*	0.1	0.2	*
2003.....	0.0	*	*	*	*	*	*	*	0.1	*	*	*
2002.....	0.1	*	*	*	*	*	*	*	0.1	0.2	*	*
2001.....	0.1	*	*	*	*	*	0.1	0.1	0.2	0.3	0.2	*
2000.....	0.1	*	*	*	*	*	0.1	*	0.2	0.2	0.2	*
1999.....	0.1	*	*	*	*	*	0.1	0.1	0.1	0.3	0.2	*
White, male												
2004.....	0.1	*	*	*	*	*	*	*	*	*	*	*
2003.....	0.1	*	*	*	*	*	*	*	*	*	*	*
2002.....	0.1	*	*	*	*	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*	*	0.2	0.3	*	*
2000.....	0.1	*	*	*	*	*	*	*	0.2	0.3	*	*
1999.....	0.1	*	*	*	*	*	0.1	*	*	0.3	*	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Sequelae of poliomyelitis (B91)												
All races 3/, both sexes												
2004.....	0.1	...	*	*	*	*	*	*	0.1	0.5	0.9	1.4
2003.....	0.1	...	*	*	*	*	*	*	0.2	0.4	0.7	1.2
2002.....	0.1	...	*	*	*	*	*	*	0.2	0.5	0.9	1.3
2001.....	0.1	...	*	*	*	*	*	0.1	0.2	0.5	0.9	1.2
2000.....	0.1	...	*	*	*	*	*	0.1	0.2	0.4	0.8	1.2
1999.....	0.1	...	*	*	*	*	*	0.1	0.1	0.4	0.6	1.3
All races 3/, male												
2004.....	0.1	...	*	*	*	*	*	*	*	0.5	1.1	1.9
2003.....	0.1	...	*	*	*	*	*	*	0.2	0.4	0.8	1.5
2002.....	0.1	...	*	*	*	*	*	*	0.2	0.5	0.9	1.9
2001.....	0.1	...	*	*	*	*	*	*	0.2	0.6	1.1	1.8
2000.....	0.1	...	*	*	*	*	*	*	0.2	0.5	0.8	*
1999.....	0.1	...	*	*	*	*	*	*	*	0.5	0.7	*
All races 3/, female												
2004.....	0.1	...	*	*	*	*	*	*	*	0.4	0.7	1.1
2003.....	0.1	...	*	*	*	*	*	*	0.2	0.4	0.7	1.1
2002.....	0.1	...	*	*	*	*	*	*	*	0.5	0.9	1.0
2001.....	0.1	...	*	*	*	*	*	*	0.2	0.5	0.7	1.0
2000.....	0.1	...	*	*	*	*	*	*	0.2	0.4	0.8	1.1
1999.....	0.1	...	*	*	*	*	*	*	*	0.3	0.6	1.2
White, both sexes												
2004.....	0.1	...	*	*	*	*	*	*	0.1	0.5	1.0	1.5
2003.....	0.1	...	*	*	*	*	*	*	0.2	0.5	0.8	1.3
2002.....	0.1	...	*	*	*	*	*	*	0.2	0.5	1.0	1.4
2001.....	0.1	...	*	*	*	*	*	*	0.2	0.6	1.0	1.4
2000.....	0.1	...	*	*	*	*	*	*	0.2	0.5	0.9	1.4
1999.....	0.1	...	*	*	*	*	*	0.1	0.2	0.4	0.7	1.4
White, male												
2004.....	0.1	...	*	*	*	*	*	*	*	0.6	1.2	2.1
2003.....	0.1	...	*	*	*	*	*	*	0.2	0.5	0.9	1.5
2002.....	0.1	...	*	*	*	*	*	*	0.2	0.6	1.0	2.1
2001.....	0.1	...	*	*	*	*	*	*	0.2	0.6	1.2	2.0
2000.....	0.1	...	*	*	*	*	*	*	0.2	0.6	0.9	*
1999.....	0.1	...	*	*	*	*	*	*	*	0.5	0.8	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

All other and unspecified infectious and parasitic diseases and their sequelae (B85-B89,B92-B99)

All races 3/, both sexes

2004.....	1.0	*	*	*	*	0.1	0.5	2.0	1.9	2.5	3.8	5.3
2003.....	0.9	*	*	*	*	0.1	0.5	1.9	1.7	2.3	3.8	4.8
2002.....	0.6	*	*	*	*	0.1	0.4	1.2	1.2	1.6	2.6	3.6
2001.....	0.4	*	*	*	*	0.1	0.3	0.6	0.7	1.3	1.8	3.2
2000.....	0.3	*	*	*	*	*	0.2	0.4	0.5	1.0	2.0	3.7
1999.....	0.5	*	*	*	*	0.1	0.4	0.7	0.8	1.5	2.6	3.9

All races 3/, male

2004.....	1.2	*	*	*	*	0.1	0.5	2.9	2.7	2.8	4.0	6.3
2003.....	1.1	*	*	*	*	*	0.6	2.7	2.3	2.6	4.2	4.8
2002.....	0.7	*	*	*	*	*	0.5	1.8	1.6	1.9	3.0	4.7
2001.....	0.5	*	*	*	*	*	0.4	0.9	0.9	1.3	2.2	3.4
2000.....	0.4	*	*	*	*	*	0.2	0.6	0.5	1.2	2.4	4.3
1999.....	0.5	*	*	*	*	0.1	0.4	0.9	0.9	1.6	3.2	3.8

All races 3/, female

2004.....	0.8	*	*	*	*	*	0.4	1.1	1.2	2.2	3.6	4.8
2003.....	0.8	*	*	*	*	*	0.3	1.0	1.2	2.1	3.6	4.8
2002.....	0.5	*	*	*	*	*	0.2	0.6	0.8	1.4	2.4	3.1
2001.....	0.4	*	*	*	*	*	0.2	0.4	0.5	1.3	1.6	3.2
2000.....	0.3	*	*	*	*	*	0.1	0.3	0.4	0.8	1.7	3.4
1999.....	0.5	*	*	*	*	0.1	0.3	0.5	0.8	1.4	2.3	3.9

White, both sexes

2004.....	1.0	*	*	*	*	0.1	0.5	2.1	1.9	2.4	3.7	5.1
2003.....	0.9	*	*	*	*	0.1	0.5	1.9	1.6	2.2	3.5	4.8
2002.....	0.6	*	*	*	*	*	0.3	1.2	1.1	1.6	2.6	3.5
2001.....	0.4	*	*	*	*	*	0.2	0.6	0.7	1.3	1.8	3.3
2000.....	0.4	*	*	*	*	*	0.2	0.4	0.4	1.0	1.9	3.6
1999.....	0.5	*	*	*	*	0.1	0.2	0.5	0.6	1.4	2.5	3.8

White, male

2004.....	1.2	*	*	*	*	*	0.6	3.1	2.5	2.8	4.0	6.2
2003.....	1.1	*	*	*	*	*	0.7	2.8	2.2	2.4	3.8	4.8
2002.....	0.8	*	*	*	*	*	0.5	1.8	1.4	1.9	2.9	4.7
2001.....	0.5	*	*	*	*	*	0.3	0.9	0.8	1.4	2.2	3.5
2000.....	0.4	*	*	*	*	*	0.2	0.6	0.5	1.1	2.4	4.1
1999.....	0.5	*	*	*	*	*	0.3	0.7	0.7	1.5	3.0	3.7

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

All other and unspecified infectious and parasitic diseases and their sequelae (B85-B89,B92-B99)

White, female

2004.....	0.8	*	*	*	*	*	0.4	1.2	1.2	2.0	3.5	4.6
2003.....	0.8	*	*	*	*	*	0.3	1.0	1.0	2.1	3.3	4.8
2002.....	0.5	*	*	*	*	*	0.2	0.5	0.7	1.3	2.4	2.9
2001.....	0.4	*	*	*	*	*	0.1	0.3	0.5	1.2	1.6	3.2
2000.....	0.3	*	*	*	*	*	*	0.3	0.3	0.8	1.6	3.4
1999.....	0.4	*	*	*	*	*	0.1	0.3	0.6	1.3	2.3	3.8

Black, both sexes

2004.....	0.8	*	*	*	*	*	0.4	1.4	2.3	3.2	4.3	8.2
2003.....	0.8	*	*	*	*	*	0.4	2.1	2.5	2.0	5.5	*
2002.....	0.6	*	*	*	*	*	0.4	1.4	1.9	*	2.9	*
2001.....	0.4	*	*	*	*	*	0.4	0.8	1.0	2.0	*	*
2000.....	0.4	*	*	*	*	*	*	0.6	1.1	1.2	2.9	*
1999.....	0.9	*	*	*	*	0.4	1.1	2.1	2.4	2.8	4.3	*

Black, male

2004.....	0.9	*	*	*	*	*	*	2.5	3.6	3.3	*	*
2003.....	1.0	*	*	*	*	*	*	3.0	3.2	2.8	6.5	*
2002.....	0.7	*	*	*	*	*	0.7	1.8	3.1	*	*	*
2001.....	0.5	*	*	*	*	*	*	1.3	*	*	*	*
2000.....	0.4	*	*	*	*	*	*	*	*	*	*	*
1999.....	0.9	*	*	*	*	*	1.0	2.0	2.5	3.2	7.1	*

Black, female

2004.....	0.6	*	*	*	*	*	*	*	1.3	3.1	3.8	7.9
2003.....	0.7	*	*	*	*	*	*	1.4	1.9	*	4.9	*
2002.....	0.5	*	*	*	*	*	*	1.0	*	*	*	*
2001.....	0.4	*	*	*	*	*	*	*	*	2.4	*	*
2000.....	0.3	*	*	*	*	*	*	*	*	*	*	*
1999.....	0.9	*	*	*	*	*	1.2	2.1	2.3	2.5	*	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
II. Neoplasms (C00-D48)												
All races 3/, both sexes												
2004.....	193.2	3.6	2.8	2.7	4.4	9.5	34.1	120.7	337.9	767.8	1,316.2	1,731.0
2003.....	196.2	3.4	2.8	2.8	4.2	9.8	35.7	123.8	347.4	783.1	1,339.0	1,778.8
2002.....	197.9	3.5	3.0	2.8	4.5	10.1	36.6	125.6	355.6	805.0	1,348.0	1,802.6
2001.....	199.2	2.9	3.1	2.7	4.5	10.5	37.6	128.3	361.4	816.9	1,352.8	1,848.3
2000.....	201.3	4.0	3.1	2.7	4.6	10.2	37.4	129.3	371.7	830.2	1,373.2	1,904.3
1999.....	201.8	3.3	3.1	2.7	4.7	10.4	37.9	129.4	379.3	841.6	1,368.0	1,888.7
All races 3/, male												
2004.....	203.1	3.5	2.9	3.0	5.1	9.1	29.9	126.2	382.1	923.3	1,707.7	2,448.4
2003.....	206.0	3.2	3.2	3.0	4.9	9.4	31.7	129.3	391.9	947.7	1,741.5	2,515.6
2002.....	208.5	3.5	3.1	3.1	5.1	9.7	32.4	129.9	405.1	980.6	1,757.2	2,592.2
2001.....	210.1	2.8	3.3	2.8	5.3	9.7	33.4	132.4	410.7	1,002.0	1,775.3	2,721.3
2000.....	212.0	4.2	3.4	2.9	5.4	9.6	33.6	133.0	421.5	1,018.9	1,808.1	2,819.6
1999.....	213.7	3.4	3.3	2.9	5.4	10.2	34.2	133.7	434.1	1,037.8	1,813.1	2,831.4
All races 3/, female												
2004.....	183.7	3.7	2.7	2.4	3.6	10.0	38.3	115.3	296.9	637.2	1,052.8	1,408.2
2003.....	186.6	3.6	2.4	2.6	3.6	10.3	39.7	118.6	306.2	645.5	1,070.0	1,453.0
2002.....	187.6	3.6	2.8	2.6	3.8	10.6	40.8	121.5	309.7	658.8	1,076.3	1,460.0
2001.....	188.7	3.0	2.9	2.6	3.7	11.3	41.8	124.4	315.9	663.5	1,075.1	1,483.0
2000.....	191.1	3.8	2.8	2.5	3.8	10.8	41.2	125.8	325.8	674.8	1,089.5	1,531.6
1999.....	190.3	3.2	2.9	2.6	4.0	10.6	41.5	125.3	328.9	681.0	1,079.0	1,508.8
White, both sexes												
2004.....	205.7	3.8	2.8	2.7	4.3	9.2	32.6	114.9	331.3	766.2	1,321.0	1,739.9
2003.....	208.9	3.6	2.9	2.9	4.3	9.6	34.3	117.6	341.1	782.6	1,341.1	1,784.4
2002.....	210.6	3.7	2.9	2.9	4.6	9.9	35.3	120.2	347.8	803.3	1,348.0	1,806.4
2001.....	211.7	2.9	3.4	2.7	4.6	10.2	36.1	123.1	354.0	812.1	1,349.1	1,849.1
2000.....	213.9	4.1	3.1	2.8	4.5	10.1	35.3	123.6	364.5	824.9	1,369.0	1,903.1
1999.....	213.9	3.2	3.3	2.7	4.6	10.2	36.0	123.6	369.9	835.5	1,361.0	1,884.1
White, male												
2004.....	215.0	3.8	2.9	3.0	5.0	9.1	28.9	119.3	370.2	912.4	1,699.4	2,452.1
2003.....	218.3	3.4	3.2	3.1	5.0	9.3	30.8	121.7	380.6	939.1	1,731.4	2,517.5
2002.....	220.6	3.9	3.0	3.0	5.3	9.5	31.4	123.6	391.2	971.1	1,742.3	2,590.9
2001.....	221.7	2.7	3.5	2.7	5.3	9.7	32.2	125.7	397.6	987.3	1,754.1	2,707.4
2000.....	223.3	4.4	3.5	3.0	5.3	9.7	31.7	125.5	407.5	1,001.3	1,785.1	2,806.3
1999.....	224.8	3.4	3.5	2.9	5.2	9.9	32.8	125.5	418.5	1,020.9	1,789.0	2,810.7

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
II. Neoplasms (C00-D48)												
White, female												
2004.....	196.7	3.9	2.8	2.4	3.5	9.4	36.5	110.6	294.4	641.1	1,062.9	1,418.8
2003.....	199.7	3.8	2.5	2.6	3.6	9.8	37.9	113.6	303.7	649.2	1,077.2	1,459.6
2002.....	200.9	3.5	2.7	2.7	3.9	10.2	39.2	116.9	306.8	660.9	1,083.3	1,465.6
2001.....	201.9	3.1	3.2	2.7	3.8	10.8	40.0	120.6	313.0	664.2	1,080.2	1,489.8
2000.....	204.8	3.9	2.7	2.6	3.7	10.5	39.0	121.6	324.3	676.9	1,095.2	1,536.2
1999.....	203.3	2.9	3.2	2.5	3.9	10.4	39.3	121.7	324.4	680.9	1,080.5	1,511.8
Black, both sexes												
2004.....	165.0	*	2.8	2.8	5.1	12.7	48.0	179.8	464.7	928.7	1,477.5	1,831.0
2003.....	167.2	*	2.6	2.8	4.2	12.7	50.1	188.1	476.3	938.4	1,517.5	1,910.9
2002.....	168.9	3.1	3.5	2.8	4.3	12.9	51.2	184.1	491.1	975.2	1,552.0	1,949.3
2001.....	170.2	*	2.1	3.0	4.7	13.2	52.4	186.4	491.9	1,005.6	1,575.5	2,005.1
2000.....	172.4	3.3	3.2	2.5	5.4	12.5	55.1	195.3	501.6	1,020.3	1,597.2	2,063.0
1999.....	174.5	4.3	2.5	3.0	5.9	12.8	55.1	195.5	527.7	1,035.4	1,612.3	2,070.3
Black, male												
2004.....	179.2	*	3.2	2.8	5.7	10.6	39.6	199.5	576.9	1,225.1	2,126.4	2,810.4
2003.....	181.2	*	3.0	3.0	4.7	11.0	42.8	209.5	590.5	1,237.7	2,180.9	2,892.3
2002.....	184.3	*	3.3	3.3	4.8	11.5	44.0	199.5	617.4	1,288.8	2,261.7	3,055.5
2001.....	187.5	*	2.6	3.3	5.5	10.9	45.7	206.7	610.7	1,352.6	2,331.5	3,298.4
2000.....	191.5	*	3.3	2.8	6.4	10.6	49.6	217.0	633.3	1,383.0	2,387.8	3,340.9
1999.....	194.1	*	2.8	3.1	6.9	12.2	48.4	223.8	666.0	1,388.8	2,397.8	3,404.0
Black, female												
2004.....	152.1	*	2.4	2.9	4.5	14.7	55.4	162.9	374.4	717.2	1,110.7	1,442.8
2003.....	154.4	*	2.1	2.6	3.6	14.3	56.5	169.6	384.3	725.4	1,142.4	1,529.6
2002.....	155.0	*	3.7	2.2	3.8	14.1	57.6	170.9	389.6	752.6	1,150.2	1,525.5
2001.....	154.6	*	1.6	2.6	3.8	15.3	58.4	168.9	396.3	760.7	1,150.9	1,525.3
2000.....	155.0	*	3.0	2.2	4.3	14.3	59.9	176.5	395.4	765.2	1,152.2	1,587.4
1999.....	156.8	*	2.1	2.9	4.9	13.3	61.0	171.3	416.7	786.1	1,172.5	1,571.7

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms (C00-C97)												
All races 3/, both sexes												
2004.....	188.6	1.8	2.5	2.5	4.1	9.1	33.4	119.0	333.4	755.1	1,280.4	1,653.3
2003.....	191.5	1.9	2.5	2.6	4.0	9.4	35.0	122.2	343.0	770.3	1,302.5	1,698.2
2002.....	193.2	1.8	2.6	2.6	4.3	9.7	35.8	123.8	351.1	792.1	1,311.9	1,723.9
2001.....	194.4	1.6	2.7	2.5	4.3	10.1	36.8	126.5	356.5	802.8	1,315.8	1,765.6
2000.....	196.5	2.4	2.7	2.5	4.4	9.8	36.6	127.5	366.7	816.3	1,335.6	1,819.4
1999.....	197.0	1.8	2.7	2.5	4.5	10.0	37.1	127.6	374.6	827.1	1,331.5	1,805.8
All races 3/, male												
2004.....	198.4	1.8	2.6	2.7	4.8	8.6	29.1	124.3	376.7	907.6	1,662.1	2,349.5
2003.....	201.3	1.7	2.8	2.8	4.6	8.9	30.8	127.4	386.8	931.7	1,695.4	2,413.8
2002.....	203.8	2.0	2.7	2.9	4.9	9.2	31.5	128.0	399.8	964.8	1,711.3	2,491.1
2001.....	205.3	1.5	2.9	2.5	5.0	9.3	32.6	130.3	405.2	984.6	1,727.1	2,613.6
2000.....	207.2	2.6	3.0	2.7	5.1	9.2	32.7	130.9	415.8	1,001.9	1,760.6	2,710.7
1999.....	208.9	1.9	2.9	2.7	5.2	9.7	33.3	131.7	428.4	1,019.4	1,767.0	2,722.2
All races 3/, female												
2004.....	179.1	1.9	2.4	2.2	3.4	9.6	37.7	113.8	293.2	627.1	1,023.5	1,340.1
2003.....	182.0	2.1	2.1	2.4	3.4	9.9	39.1	117.1	302.3	635.3	1,040.1	1,381.9
2002.....	183.0	1.6	2.4	2.4	3.6	10.2	40.0	119.8	306.0	648.5	1,046.7	1,391.1
2001.....	183.9	1.8	2.5	2.4	3.5	10.9	41.0	122.7	311.5	652.2	1,045.4	1,410.7
2000.....	186.2	2.3	2.5	2.2	3.6	10.4	40.4	124.2	321.3	663.6	1,058.5	1,456.4
1999.....	185.6	1.7	2.5	2.3	3.7	10.3	40.9	123.7	325.0	669.5	1,048.5	1,436.5
White, both sexes												
2004.....	200.7	2.0	2.5	2.5	4.0	8.8	32.0	113.3	326.9	753.3	1,284.3	1,659.7
2003.....	203.7	2.1	2.6	2.7	4.1	9.1	33.6	116.0	336.6	769.5	1,303.4	1,701.4
2002.....	205.5	1.8	2.5	2.7	4.4	9.5	34.5	118.5	343.3	790.1	1,311.0	1,725.7
2001.....	206.4	1.7	3.0	2.4	4.3	9.8	35.3	121.2	349.1	797.8	1,311.1	1,765.2
2000.....	208.6	2.5	2.8	2.6	4.3	9.7	34.5	121.8	359.4	811.0	1,330.3	1,815.7
1999.....	208.7	1.8	2.9	2.4	4.3	9.8	35.3	121.9	365.2	820.7	1,323.7	1,798.8
White, male												
2004.....	209.9	1.9	2.5	2.8	4.7	8.6	28.2	117.5	364.9	896.3	1,652.7	2,348.9
2003.....	213.1	1.8	2.9	2.9	4.7	8.9	29.9	119.9	375.6	922.7	1,683.6	2,412.1
2002.....	215.5	2.2	2.7	2.8	5.0	9.1	30.5	121.8	386.0	954.8	1,695.3	2,486.8
2001.....	216.4	1.5	3.1	2.4	5.1	9.3	31.3	123.6	392.1	969.4	1,704.6	2,597.6
2000.....	218.1	2.8	3.1	2.8	5.0	9.2	30.9	123.5	401.9	984.3	1,736.0	2,693.7
1999.....	219.5	2.0	3.2	2.7	5.0	9.5	31.9	123.5	412.7	1,002.3	1,741.8	2,696.3

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms (C00-C97)												
White, female												
2004.....	191.7	2.1	2.5	2.2	3.3	9.0	35.8	109.2	290.8	630.8	1,033.1	1,348.9
2003.....	194.6	2.3	2.3	2.5	3.4	9.4	37.3	112.1	299.8	638.9	1,046.3	1,386.5
2002.....	195.8	1.4	2.3	2.5	3.7	9.9	38.5	115.3	303.1	650.4	1,053.1	1,395.1
2001.....	196.7	2.0	2.8	2.5	3.5	10.4	39.3	118.9	308.6	652.9	1,049.8	1,416.7
2000.....	199.4	2.3	2.4	2.3	3.6	10.1	38.2	120.1	319.7	665.6	1,063.4	1,459.1
1999.....	198.2	1.5	2.7	2.2	3.7	10.1	38.7	120.2	320.7	669.4	1,049.6	1,438.2
Black, both sexes												
2004.....	161.9	*	2.5	2.5	4.8	12.0	46.8	177.6	458.5	915.9	1,446.0	1,774.0
2003.....	164.3	*	2.2	2.6	3.9	12.2	48.9	185.6	471.0	925.4	1,487.9	1,853.1
2002.....	165.9	*	2.9	2.5	4.0	12.3	50.0	181.7	485.6	962.6	1,520.7	1,885.7
2001.....	167.1	*	1.9	2.6	4.3	12.9	51.4	184.4	486.1	991.2	1,543.9	1,931.7
2000.....	169.3	*	2.8	2.2	5.1	11.9	54.0	192.6	496.0	1,005.3	1,566.3	1,998.9
1999.....	171.3	*	2.0	2.7	5.5	12.2	53.9	192.9	521.4	1,019.7	1,579.3	2,008.4
Black, male												
2004.....	176.2	*	2.8	2.5	5.4	10.0	38.4	197.0	569.2	1,209.7	2,087.2	2,748.8
2003.....	178.3	*	2.4	2.8	4.5	10.3	41.7	207.0	583.8	1,221.5	2,144.2	2,825.5
2002.....	181.5	*	2.6	3.1	4.5	11.2	43.0	197.3	610.3	1,274.7	2,223.0	2,976.1
2001.....	184.5	*	2.4	2.9	5.0	10.5	44.6	204.8	604.2	1,335.3	2,290.0	3,209.9
2000.....	188.5	*	2.7	2.5	6.1	10.1	48.4	214.2	626.4	1,363.8	2,351.8	3,264.8
1999.....	191.0	*	2.1	2.8	6.4	11.4	47.3	221.3	659.0	1,369.8	2,359.4	3,340.7
Black, female												
2004.....	148.9	*	2.2	2.6	4.2	13.9	54.2	160.9	369.4	706.2	1,083.6	1,387.7
2003.....	151.4	*	2.1	2.3	3.3	13.9	55.4	167.2	380.4	714.6	1,116.9	1,475.3
2002.....	151.7	*	3.2	2.0	3.6	13.3	56.2	168.2	385.4	741.1	1,123.1	1,468.0
2001.....	151.3	*	*	2.2	3.6	15.0	57.3	166.8	390.9	748.4	1,125.0	1,457.5
2000.....	151.8	*	2.9	1.9	4.1	13.5	58.9	173.9	391.0	753.1	1,124.0	1,527.7
1999.....	153.3	*	1.9	2.6	4.6	13.0	59.8	168.5	411.1	772.7	1,142.6	1,510.4

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)

All races 3/, both sexes

2004.....	2.7	*	*	*	0.1	0.1	0.6	2.9	6.1	10.1	13.7	18.0
2003.....	2.7	*	*	*	0.1	0.1	0.6	2.8	6.5	10.2	13.1	18.6
2002.....	2.7	*	*	*	0.1	0.1	0.6	2.7	6.8	10.4	13.5	18.7
2001.....	2.7	*	*	*	0.1	0.1	0.6	2.7	6.9	10.4	14.2	18.5
2000.....	2.7	*	*	*	0.1	0.1	0.6	2.8	6.8	10.7	13.2	19.4
1999.....	2.7	*	*	*	0.1	0.1	0.6	2.9	6.8	10.3	14.3	19.3

All races 3/, male

2004.....	3.7	*	*	*	*	0.2	0.8	4.6	10.0	15.6	20.3	24.9
2003.....	3.7	*	*	*	0.1	0.2	0.9	4.6	10.6	15.6	19.4	26.9
2002.....	3.7	*	*	*	*	0.2	0.8	4.4	10.9	15.9	19.6	26.9
2001.....	3.7	*	*	*	*	0.2	0.9	4.4	11.3	16.0	20.6	25.8
2000.....	3.6	*	*	*	*	0.1	1.0	4.5	10.9	16.0	19.2	27.1
1999.....	3.6	*	*	*	0.1	0.1	0.9	4.6	10.8	15.8	20.9	27.1

All races 3/, female

2004.....	1.7	*	*	*	*	*	0.4	1.2	2.5	5.6	9.2	14.9
2003.....	1.7	*	*	*	*	0.1	0.4	1.0	2.8	5.7	9.0	15.0
2002.....	1.7	*	*	*	*	*	0.4	1.1	3.0	5.8	9.5	15.1
2001.....	1.8	*	*	*	*	0.1	0.4	1.2	2.8	5.8	10.0	15.4
2000.....	1.8	*	*	*	*	0.1	0.3	1.3	2.9	6.3	9.2	16.3
1999.....	1.8	*	*	*	*	*	0.4	1.3	3.1	5.8	9.9	16.1

White, both sexes

2004.....	2.7	*	*	*	*	0.1	0.5	2.5	5.6	9.9	13.7	18.6
2003.....	2.7	*	*	*	*	0.1	0.5	2.4	6.2	9.8	12.9	19.0
2002.....	2.7	*	*	*	*	0.1	0.5	2.4	6.3	10.2	13.7	18.9
2001.....	2.7	*	*	*	0.1	0.1	0.5	2.4	6.3	10.0	14.4	19.0
2000.....	2.7	*	*	*	*	0.1	0.5	2.4	6.0	10.2	13.1	19.7
1999.....	2.7	*	*	*	*	0.1	0.5	2.3	6.1	9.9	14.1	19.8

White, male

2004.....	3.6	*	*	*	*	0.2	0.6	3.9	9.1	14.8	19.9	24.8
2003.....	3.7	*	*	*	*	0.2	0.8	4.0	9.7	14.6	18.6	27.2
2002.....	3.7	*	*	*	*	0.2	0.7	3.8	9.9	15.3	19.7	27.0
2001.....	3.6	*	*	*	*	0.1	0.7	3.9	10.1	15.1	20.4	26.4
2000.....	3.5	*	*	*	*	0.1	0.7	3.7	9.6	14.9	18.7	27.6
1999.....	3.5	*	*	*	*	*	0.7	3.6	9.6	15.1	20.3	27.3

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)												
White, female												
2004.....	1.8	*	*	*	*	*	0.3	1.1	2.4	5.8	9.4	15.7
2003.....	1.8	*	*	*	*	*	0.2	0.9	2.8	5.7	9.1	15.4
2002.....	1.8	*	*	*	*	*	0.3	1.0	2.8	5.9	9.6	15.3
2001.....	1.9	*	*	*	*	*	0.3	1.0	2.7	5.8	10.3	15.9
2000.....	1.9	*	*	*	*	*	0.3	1.2	2.7	6.3	9.4	16.5
1999.....	1.9	*	*	*	*	*	0.3	1.0	2.8	5.6	10.1	16.7
Black, both sexes												
2004.....	3.0	*	*	*	*	*	1.3	6.0	11.1	13.8	15.3	12.4
2003.....	2.9	*	*	*	*	*	1.1	5.1	11.2	15.4	15.3	15.5
2002.....	2.8	*	*	*	*	*	1.1	5.5	12.1	12.9	11.8	16.0
2001.....	2.9	*	*	*	*	*	1.2	5.1	12.6	15.5	13.9	12.9
2000.....	3.0	*	*	*	*	*	1.1	6.0	13.5	15.5	15.1	15.3
1999.....	3.2	*	*	*	*	*	1.2	7.1	13.5	15.2	16.9	14.3
Black, male												
2004.....	4.7	*	*	*	*	*	1.9	10.4	19.8	26.6	26.4	26.8
2003.....	4.6	*	*	*	*	*	1.2	9.0	21.1	28.7	27.9	25.0
2002.....	4.3	*	*	*	*	*	1.6	9.5	21.9	23.0	18.4	26.8
2001.....	4.4	*	*	*	*	*	1.7	8.7	23.3	27.1	23.5	*
2000.....	4.8	*	*	*	*	*	1.7	10.5	24.3	28.2	28.8	*
1999.....	4.8	*	*	*	*	*	1.8	12.1	23.2	25.1	31.7	24.6
Black, female												
2004.....	1.4	*	*	*	*	*	0.8	2.3	4.2	4.6	9.0	*
2003.....	1.4	*	*	*	*	*	1.0	1.7	3.2	6.0	8.1	11.7
2002.....	1.4	*	*	*	*	*	0.7	2.0	4.2	5.8	8.1	11.9
2001.....	1.5	*	*	*	*	*	0.8	2.1	3.9	7.4	8.4	10.5
2000.....	1.4	*	*	*	*	*	*	2.0	4.8	6.5	7.3	14.2
1999.....	1.7	*	*	*	*	*	0.7	2.9	5.7	8.2	8.6	10.5

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of tongue (C01-C02)												
All races 3/, both sexes												
2004.....	0.6	*	*	*	*	*	0.2	0.7	1.5	2.5	2.9	4.1
2003.....	0.6	*	*	*	*	0.1	0.2	0.7	1.6	2.5	3.0	3.7
2002.....	0.7	*	*	*	*	0.1	0.2	0.7	1.7	2.4	3.3	4.0
2001.....	0.6	*	*	*	*	0.1	0.2	0.7	1.7	2.3	3.3	3.8
2000.....	0.6	*	*	*	*	0.1	0.2	0.7	1.7	2.3	3.0	4.0
1999.....	0.6	*	*	*	*	*	0.2	0.7	1.5	2.4	3.1	4.0
All races 3/, male												
2004.....	0.8	*	*	*	*	*	0.2	1.1	2.5	3.6	4.0	4.8
2003.....	0.9	*	*	*	*	*	0.2	1.2	2.5	3.6	4.2	5.0
2002.....	0.9	*	*	*	*	*	0.3	1.0	2.7	3.6	4.4	5.2
2001.....	0.9	*	*	*	*	*	0.2	1.1	2.8	3.5	4.4	4.1
2000.....	0.8	*	*	*	*	*	0.2	1.1	2.7	3.4	3.6	5.1
1999.....	0.8	*	*	*	*	*	0.3	1.1	2.4	3.7	4.5	5.4
All races 3/, female												
2004.....	0.4	*	*	*	*	*	0.2	0.4	0.6	1.5	2.2	3.8
2003.....	0.4	*	*	*	*	*	0.1	0.3	0.7	1.5	2.2	3.2
2002.....	0.5	*	*	*	*	*	*	0.3	0.9	1.5	2.5	3.5
2001.....	0.4	*	*	*	*	*	0.1	0.3	0.7	1.2	2.5	3.6
2000.....	0.5	*	*	*	*	*	0.1	0.4	0.8	1.4	2.5	3.6
1999.....	0.4	*	*	*	*	*	0.2	0.3	0.8	1.3	2.2	3.5
White, both sexes												
2004.....	0.7	*	*	*	*	*	0.2	0.7	1.5	2.5	3.0	4.2
2003.....	0.7	*	*	*	*	*	0.2	0.7	1.6	2.4	3.1	3.8
2002.....	0.7	*	*	*	*	*	0.2	0.6	1.7	2.4	3.4	4.1
2001.....	0.7	*	*	*	*	0.1	0.1	0.7	1.6	2.3	3.3	3.9
2000.....	0.7	*	*	*	*	0.1	0.2	0.7	1.6	2.3	3.0	4.1
1999.....	0.6	*	*	*	*	*	0.2	0.6	1.4	2.4	3.2	4.1
White, male												
2004.....	0.9	*	*	*	*	*	0.2	1.1	2.5	3.6	4.1	4.8
2003.....	0.9	*	*	*	*	*	0.2	1.1	2.5	3.5	4.2	5.1
2002.....	0.9	*	*	*	*	*	0.3	1.0	2.5	3.5	4.5	5.2
2001.....	0.9	*	*	*	*	*	0.2	1.1	2.7	3.5	4.4	4.3
2000.....	0.8	*	*	*	*	*	0.2	1.0	2.6	3.4	3.6	5.1
1999.....	0.8	*	*	*	*	*	0.2	1.0	2.2	3.7	4.6	5.5

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of pharynx (C10-C13,C14.0)												
All races 3/, both sexes												
2004.....	1.1	*	*	*	*	*	0.2	1.3	2.8	4.5	5.6	5.1
2003.....	1.1	*	*	*	*	0.1	0.2	1.1	2.9	4.6	5.1	6.0
2002.....	1.1	*	*	*	*	*	0.2	1.2	3.0	4.5	5.1	5.4
2001.....	1.1	*	*	*	*	*	0.3	1.2	3.1	4.5	5.5	5.9
2000.....	1.1	*	*	*	*	*	0.2	1.2	3.1	4.7	5.3	5.6
1999.....	1.1	*	*	*	*	*	0.3	1.3	3.1	4.6	5.6	5.2
All races 3/, male												
2004.....	1.7	*	*	*	*	*	0.3	2.1	4.7	7.4	9.1	8.7
2003.....	1.6	*	*	*	*	*	0.4	1.9	4.7	7.4	8.4	10.8
2002.....	1.6	*	*	*	*	*	0.3	1.9	4.8	7.1	8.2	9.8
2001.....	1.6	*	*	*	*	*	0.4	2.0	5.1	7.2	8.9	10.0
2000.....	1.6	*	*	*	*	*	0.4	1.9	5.2	7.5	8.6	9.9
1999.....	1.6	*	*	*	*	*	0.4	2.1	5.1	7.2	9.0	9.9
All races 3/, female												
2004.....	0.6	*	*	*	*	*	0.1	0.5	1.1	2.1	3.3	3.5
2003.....	0.6	*	*	*	*	*	0.1	0.4	1.2	2.3	2.8	3.9
2002.....	0.6	*	*	*	*	*	0.2	0.5	1.3	2.3	3.0	3.5
2001.....	0.6	*	*	*	*	*	0.1	0.5	1.2	2.3	3.3	4.2
2000.....	0.6	*	*	*	*	*	*	0.5	1.3	2.5	3.1	3.9
1999.....	0.6	*	*	*	*	*	0.1	0.5	1.3	2.5	3.3	3.4
White, both sexes												
2004.....	1.1	*	*	*	*	*	0.2	1.0	2.4	4.2	5.4	5.1
2003.....	1.0	*	*	*	*	*	0.2	0.9	2.5	4.3	4.7	5.8
2002.....	1.0	*	*	*	*	*	0.2	0.9	2.5	4.3	4.8	5.2
2001.....	1.1	*	*	*	*	*	0.2	0.9	2.7	4.2	5.4	6.0
2000.....	1.0	*	*	*	*	*	0.1	0.9	2.6	4.3	5.0	5.3
1999.....	1.0	*	*	*	*	*	0.2	0.9	2.7	4.2	5.2	5.3
White, male												
2004.....	1.5	*	*	*	*	*	0.2	1.6	3.9	6.7	8.6	8.2
2003.....	1.5	*	*	*	*	*	0.3	1.5	4.0	6.6	7.7	10.6
2002.....	1.5	*	*	*	*	*	0.2	1.5	4.1	6.7	7.8	9.5
2001.....	1.5	*	*	*	*	*	0.3	1.5	4.4	6.4	8.5	9.9
2000.....	1.4	*	*	*	*	*	0.2	1.4	4.2	6.6	8.0	9.5
1999.....	1.5	*	*	*	*	*	0.2	1.4	4.3	6.6	8.3	9.9

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of pharynx (C10-C13,C14.0)												
White, female												
2004.....	0.6	*	*	*	*	*	*	0.4	0.9	2.1	3.2	3.7
2003.....	0.6	*	*	*	*	*	*	0.3	1.1	2.3	2.7	3.7
2002.....	0.6	*	*	*	*	*	*	0.4	1.1	2.3	2.8	3.4
2001.....	0.6	*	*	*	*	*	*	0.3	1.1	2.2	3.3	4.3
2000.....	0.6	*	*	*	*	*	*	0.4	1.1	2.3	3.1	3.7
1999.....	0.6	*	*	*	*	*	0.1	0.4	1.1	2.3	3.2	3.5
Black, both sexes												
2004.....	1.7	*	*	*	*	*	0.6	3.2	7.0	8.0	8.1	5.6
2003.....	1.5	*	*	*	*	*	0.5	2.6	6.0	8.5	7.8	7.3
2002.....	1.4	*	*	*	*	*	0.6	2.8	6.4	6.0	6.2	7.4
2001.....	1.5	*	*	*	*	*	0.6	2.8	6.5	7.6	7.4	*
2000.....	1.6	*	*	*	*	*	0.6	2.9	7.8	8.6	8.0	7.5
1999.....	1.6	*	*	*	*	*	0.5	3.7	6.9	8.2	9.7	*
Black, male												
2004.....	2.7	*	*	*	*	*	0.9	5.6	12.4	15.4	13.9	*
2003.....	2.6	*	*	*	*	*	*	4.7	11.5	16.7	15.8	*
2002.....	2.2	*	*	*	*	*	0.8	5.1	11.8	10.6	10.3	*
2001.....	2.3	*	*	*	*	*	1.0	4.8	11.7	14.5	12.9	*
2000.....	2.6	*	*	*	*	*	1.0	5.1	14.1	15.7	15.4	*
1999.....	2.6	*	*	*	*	*	0.9	6.6	12.0	13.7	19.1	*
Black, female												
2004.....	0.7	*	*	*	*	*	*	1.1	2.7	2.8	4.9	*
2003.....	0.6	*	*	*	*	*	*	0.8	1.6	2.7	3.3	*
2002.....	0.7	*	*	*	*	*	*	0.8	2.0	2.8	3.9	*
2001.....	0.7	*	*	*	*	*	*	1.2	2.3	2.8	4.3	*
2000.....	0.7	*	*	*	*	*	*	1.0	2.7	3.6	3.8	*
1999.....	0.8	*	*	*	*	*	*	1.2	2.9	4.3	4.5	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Of other and unspecified sites within the lip, oral cavity and pharynx (C03-C09,C14.2-C14.8)

All races 3/, both sexes

2004.....	0.9	*	*	*	*	*	0.2	0.8	1.8	3.1	4.9	8.4
2003.....	0.9	*	*	*	*	*	0.2	0.9	2.0	3.0	4.8	8.7
2002.....	0.9	*	*	*	*	*	0.2	0.9	2.1	3.4	5.0	8.7
2001.....	0.9	*	*	*	*	*	0.2	0.8	2.0	3.6	5.3	8.3
2000.....	0.9	*	*	*	*	*	0.2	0.9	1.9	3.5	4.7	9.5
1999.....	0.9	*	*	*	*	*	0.2	0.9	2.1	3.3	5.4	9.7

All races 3/, male

2004.....	1.1	*	*	*	*	*	0.2	1.3	2.8	4.5	6.8	10.8
2003.....	1.2	*	*	*	*	*	0.3	1.5	3.3	4.4	6.3	10.7
2002.....	1.2	*	*	*	*	*	0.2	1.4	3.4	5.1	6.7	11.1
2001.....	1.2	*	*	*	*	*	0.2	1.2	3.3	5.1	7.0	10.6
2000.....	1.1	*	*	*	*	*	0.3	1.4	2.9	5.0	6.7	11.4
1999.....	1.1	*	*	*	*	*	0.3	1.4	3.2	4.7	7.2	11.4

All races 3/, female

2004.....	0.6	*	*	*	*	*	*	0.3	0.8	2.0	3.7	7.3
2003.....	0.7	*	*	*	*	*	0.1	0.3	0.9	1.8	3.8	7.7
2002.....	0.7	*	*	*	*	*	0.1	0.3	0.9	2.0	3.9	7.7
2001.....	0.7	*	*	*	*	*	0.1	0.4	0.9	2.3	4.1	7.3
2000.....	0.7	*	*	*	*	*	0.1	0.4	0.9	2.3	3.4	8.8
1999.....	0.7	*	*	*	*	*	0.1	0.4	1.0	2.1	4.3	9.0

White, both sexes

2004.....	0.9	*	*	*	*	*	0.1	0.8	1.7	3.2	5.1	8.8
2003.....	1.0	*	*	*	*	*	0.2	0.8	2.0	3.0	4.9	9.0
2002.....	1.0	*	*	*	*	*	0.2	0.8	2.0	3.5	5.2	9.1
2001.....	1.0	*	*	*	*	*	0.1	0.8	1.9	3.5	5.5	8.5
2000.....	1.0	*	*	*	*	*	0.2	0.8	1.7	3.5	4.8	9.9
1999.....	1.0	*	*	*	*	*	0.1	0.8	2.0	3.2	5.6	10.0

White, male

2004.....	1.2	*	*	*	*	*	0.2	1.2	2.7	4.4	6.8	11.2
2003.....	1.2	*	*	*	*	*	0.3	1.4	3.1	4.3	6.3	11.1
2002.....	1.3	*	*	*	*	*	0.2	1.3	3.3	5.1	7.0	11.5
2001.....	1.2	*	*	*	*	*	0.2	1.2	3.0	5.0	7.2	11.0
2000.....	1.2	*	*	*	*	*	0.3	1.3	2.7	4.8	6.7	12.2
1999.....	1.2	*	*	*	*	*	0.2	1.2	3.0	4.6	7.2	11.5

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Of other and unspecified sites within the lip, oral cavity and pharynx (C03-C09,C14.2-C14.8)

White, female

2004.....	0.7	*	*	*	*	*	*	0.3	0.8	2.1	3.8	7.7
2003.....	0.7	*	*	*	*	*	*	0.3	0.9	1.9	4.0	8.1
2002.....	0.7	*	*	*	*	*	0.1	0.3	0.9	2.1	4.0	8.0
2001.....	0.8	*	*	*	*	*	*	0.3	0.9	2.3	4.4	7.5
2000.....	0.8	*	*	*	*	*	0.1	0.4	0.9	2.5	3.6	8.9
1999.....	0.8	*	*	*	*	*	*	0.3	1.0	2.1	4.5	9.5

Black, both sexes

2004.....	0.8	*	*	*	*	*	0.3	1.7	2.6	3.1	4.2	*
2003.....	0.8	*	*	*	*	*	0.4	1.4	3.0	3.7	4.9	*
2002.....	0.8	*	*	*	*	*	*	1.6	3.1	3.7	3.9	6.2
2001.....	0.8	*	*	*	*	*	*	1.2	3.7	5.4	3.8	*
2000.....	0.8	*	*	*	*	*	*	1.9	3.2	4.1	4.7	*
1999.....	0.9	*	*	*	*	*	0.4	1.9	3.9	4.2	4.5	6.4

Black, male

2004.....	1.2	*	*	*	*	*	*	2.9	4.3	5.7	7.5	*
2003.....	1.2	*	*	*	*	*	*	2.5	5.8	6.5	7.1	*
2002.....	1.2	*	*	*	*	*	*	2.7	5.7	6.7	*	*
2001.....	1.2	*	*	*	*	*	*	1.9	6.9	8.4	6.5	*
2000.....	1.2	*	*	*	*	*	*	3.1	5.8	7.6	8.5	*
1999.....	1.3	*	*	*	*	*	*	2.9	6.6	6.8	7.7	*

Black, female

2004.....	0.4	*	*	*	*	*	*	*	*	*	*	*
2003.....	0.4	*	*	*	*	*	*	*	*	*	3.7	*
2002.....	0.4	*	*	*	*	*	*	*	*	*	3.4	*
2001.....	0.5	*	*	*	*	*	*	*	*	3.3	*	*
2000.....	0.4	*	*	*	*	*	*	*	*	*	*	*
1999.....	0.5	*	*	*	*	*	*	1.0	1.7	2.4	*	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of digestive organs (C15-C26)												
All races 3/, both sexes												
2004.....	44.7	*	0.1	*	0.3	1.5	7.1	29.3	78.5	170.4	302.9	445.3
2003.....	45.5	*	0.1	0.1	0.3	1.6	7.0	30.0	79.1	176.7	309.8	461.2
2002.....	45.7	*	0.2	0.1	0.3	1.5	7.4	29.7	80.6	181.0	311.9	467.4
2001.....	46.0	*	*	*	0.3	1.5	7.4	30.1	80.4	183.8	315.0	487.9
2000.....	46.4	*	0.2	0.1	0.2	1.5	7.2	29.6	83.1	184.4	323.9	504.9
1999.....	46.4	*	0.2	*	0.3	1.5	7.4	29.3	84.0	187.0	320.2	507.4
All races 3/, male												
2004.....	49.8	*	*	*	0.4	1.7	8.5	39.5	104.7	222.9	385.0	531.7
2003.....	50.4	*	*	*	0.3	1.7	8.5	39.9	105.2	230.8	395.9	546.7
2002.....	50.7	*	0.3	*	0.3	1.8	9.1	39.7	108.1	237.5	397.9	557.9
2001.....	50.5	*	*	*	0.3	1.7	8.8	39.7	106.9	238.4	402.1	593.2
2000.....	50.8	*	*	*	0.3	1.8	8.9	39.1	109.6	239.8	413.7	614.1
1999.....	50.7	*	0.3	*	0.3	1.8	8.9	38.9	110.0	243.0	411.2	620.8
All races 3/, female												
2004.....	39.8	*	*	*	0.3	1.3	5.6	19.5	54.2	126.4	247.7	406.4
2003.....	40.7	*	*	*	0.3	1.4	5.6	20.5	54.8	131.5	252.4	423.4
2002.....	40.8	*	*	*	0.3	1.3	5.7	20.1	55.2	133.9	254.9	428.1
2001.....	41.7	*	*	*	0.2	1.3	6.0	20.9	56.0	138.5	257.7	443.9
2000.....	42.3	*	*	*	0.2	1.2	5.6	20.5	58.6	138.9	265.3	460.4
1999.....	42.3	*	*	*	0.3	1.2	5.9	20.1	60.0	141.1	261.1	461.7
White, both sexes												
2004.....	46.3	*	*	*	0.3	1.4	6.5	26.9	73.9	163.8	297.5	440.0
2003.....	47.0	*	*	*	0.3	1.4	6.3	27.0	74.5	171.0	303.7	453.8
2002.....	47.3	*	0.2	*	0.3	1.4	6.8	27.3	75.9	174.6	304.8	460.4
2001.....	47.6	*	*	*	0.2	1.4	6.7	27.6	75.7	176.8	307.3	481.6
2000.....	48.0	*	*	*	0.2	1.3	6.4	27.1	78.1	177.8	315.6	497.0
1999.....	48.0	*	0.2	*	0.2	1.3	6.6	26.9	78.5	180.6	312.0	499.7
White, male												
2004.....	51.3	*	*	*	0.3	1.5	7.9	36.5	98.5	213.9	378.6	528.5
2003.....	52.0	*	*	*	0.3	1.5	7.7	35.9	99.3	224.0	389.3	539.4
2002.....	52.3	*	*	*	0.3	1.5	8.3	36.5	101.9	230.0	389.6	550.5
2001.....	52.1	*	*	*	0.3	1.5	8.0	36.4	100.8	230.1	394.0	588.1
2000.....	52.3	*	*	*	0.3	1.6	7.9	35.7	102.6	231.5	404.4	605.3
1999.....	52.2	*	*	*	0.3	1.5	8.0	35.4	102.5	235.6	401.1	614.4

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of digestive organs (C15-C26)												
White, female												
2004.....	41.3	*	*	*	0.2	1.1	5.0	17.5	50.6	120.9	242.1	400.0
2003.....	42.2	*	*	*	0.2	1.3	4.9	18.3	51.0	125.9	245.9	415.8
2002.....	42.4	*	*	*	0.3	1.3	5.2	18.2	51.5	127.6	247.9	421.3
2001.....	43.2	*	*	*	0.2	1.2	5.4	19.0	52.1	131.8	249.7	437.0
2000.....	43.9	*	*	*	0.2	1.1	4.9	18.7	55.0	132.7	257.2	453.1
1999.....	43.9	*	*	*	0.2	1.1	5.1	18.5	56.0	134.7	253.6	453.7
Black, both sexes												
2004.....	42.7	*	*	*	0.6	2.3	10.2	47.1	123.4	242.7	379.3	524.2
2003.....	43.7	*	*	*	0.3	2.1	11.2	52.0	125.1	239.2	395.7	567.1
2002.....	43.6	*	*	*	0.4	2.1	11.1	48.2	126.6	252.1	407.8	566.6
2001.....	44.1	*	*	*	*	2.2	11.6	49.8	126.5	257.2	417.3	585.6
2000.....	44.5	*	*	*	*	2.3	11.7	51.2	129.3	257.9	429.8	606.9
1999.....	44.6	*	*	*	0.5	2.4	11.9	49.2	134.3	260.0	429.7	613.1
Black, male												
2004.....	47.7	*	*	*	0.8	2.5	12.1	63.5	169.2	330.8	488.9	575.2
2003.....	48.4	*	*	*	*	2.5	13.6	72.2	170.6	319.2	498.1	654.8
2002.....	48.5	*	*	*	*	2.7	13.9	65.2	173.0	333.8	531.0	671.1
2001.....	48.1	*	*	*	*	2.6	14.2	66.5	169.6	340.1	531.2	676.3
2000.....	49.2	*	*	*	*	2.7	14.9	69.1	178.7	339.5	552.0	728.8
1999.....	49.4	*	*	*	*	3.0	14.3	68.6	184.7	338.3	569.1	715.9
Black, female												
2004.....	38.2	*	*	*	*	2.0	8.6	33.0	86.6	179.9	317.4	504.0
2003.....	39.3	*	*	*	*	1.8	9.1	34.5	88.5	182.4	337.7	533.1
2002.....	39.2	*	*	*	*	1.5	8.6	33.6	89.3	194.0	338.0	526.6
2001.....	40.5	*	*	*	*	1.8	9.4	35.4	91.8	198.7	353.3	551.9
2000.....	40.3	*	*	*	*	2.0	8.9	35.8	89.5	200.6	361.0	561.5
1999.....	40.4	*	*	*	*	1.9	9.8	32.7	93.9	204.9	351.6	574.7

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of esophagus (C15)												
All races 3/, both sexes												
2004.....	4.4	*	*	*	*	0.1	0.7	3.4	10.3	19.2	26.7	25.8
2003.....	4.4	*	*	*	*	0.1	0.6	3.4	10.5	19.7	26.4	25.0
2002.....	4.4	*	*	*	*	*	0.7	3.5	10.4	20.2	26.6	24.6
2001.....	4.4	*	*	*	*	0.1	0.7	3.6	10.1	20.6	26.3	25.8
2000.....	4.3	*	*	*	*	0.1	0.7	3.6	10.8	20.0	25.6	25.9
1999.....	4.3	*	*	*	*	0.1	0.7	3.5	10.6	20.1	24.4	26.8
All races 3/, male												
2004.....	7.0	*	*	*	*	0.1	1.1	5.9	17.9	33.4	48.8	48.5
2003.....	6.9	*	*	*	*	0.1	1.1	6.0	18.5	34.1	47.8	45.3
2002.....	6.9	*	*	*	*	*	1.1	6.0	18.2	35.6	48.3	43.4
2001.....	6.9	*	*	*	*	0.1	1.1	6.3	17.6	36.0	47.4	48.1
2000.....	6.7	*	*	*	*	0.2	1.2	6.3	18.6	34.1	45.8	47.8
1999.....	6.6	*	*	*	*	0.1	1.1	6.1	18.3	35.3	44.4	48.0
All races 3/, female												
2004.....	2.0	*	*	*	*	*	0.2	1.0	3.2	7.3	11.9	15.7
2003.....	2.0	*	*	*	*	*	0.2	1.0	3.2	7.7	12.0	16.1
2002.....	2.0	*	*	*	*	*	0.2	1.0	3.2	7.4	12.2	16.5
2001.....	2.0	*	*	*	*	*	0.2	1.0	3.2	7.9	12.4	16.5
2000.....	2.1	*	*	*	*	*	0.1	1.0	3.5	8.3	12.4	17.1
1999.....	2.0	*	*	*	*	*	0.2	0.9	3.6	7.6	11.5	18.3
White, both sexes												
2004.....	4.7	*	*	*	*	0.1	0.7	3.3	10.0	19.2	27.1	26.2
2003.....	4.7	*	*	*	*	0.1	0.7	3.2	10.2	19.6	26.7	25.8
2002.....	4.6	*	*	*	*	*	0.7	3.4	9.9	19.8	26.7	25.1
2001.....	4.6	*	*	*	*	0.1	0.7	3.4	9.4	19.8	26.5	26.3
2000.....	4.5	*	*	*	*	0.1	0.7	3.3	10.0	19.5	25.3	26.6
1999.....	4.4	*	*	*	*	0.1	0.6	3.2	9.8	19.4	24.2	27.2
White, male												
2004.....	7.5	*	*	*	*	0.1	1.2	5.8	17.6	33.5	49.7	49.2
2003.....	7.4	*	*	*	*	0.1	1.1	5.7	18.0	33.9	48.8	46.7
2002.....	7.3	*	*	*	*	*	1.2	6.0	17.5	35.2	48.8	44.6
2001.....	7.2	*	*	*	*	0.1	1.1	5.9	16.4	34.9	47.9	48.8
2000.....	7.0	*	*	*	*	0.2	1.2	5.9	17.5	33.6	45.6	49.3
1999.....	6.9	*	*	*	*	0.1	1.1	5.7	17.0	34.4	44.0	48.0

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of esophagus (C15)												
White, female												
2004.....	2.0	*	*	*	*	*	0.2	0.8	2.8	6.9	11.7	15.9
2003.....	2.0	*	*	*	*	*	0.2	0.8	2.7	7.5	11.7	16.5
2002.....	2.0	*	*	*	*	*	0.2	0.8	2.8	6.8	11.9	16.7
2001.....	2.1	*	*	*	*	*	0.2	0.9	2.7	7.1	12.3	16.8
2000.....	2.1	*	*	*	*	*	0.1	0.8	2.9	7.8	11.9	17.3
1999.....	2.0	*	*	*	*	*	0.1	0.7	3.1	6.9	11.2	18.9
Black, both sexes												
2004.....	4.0	*	*	*	*	*	0.7	5.2	15.6	24.7	27.2	21.4
2003.....	4.2	*	*	*	*	*	0.8	6.0	16.9	25.6	27.9	21.3
2002.....	4.2	*	*	*	*	*	1.0	5.1	17.3	28.5	29.6	20.8
2001.....	4.5	*	*	*	*	*	0.9	5.9	18.5	32.3	29.5	23.6
2000.....	4.5	*	*	*	*	*	0.8	6.6	19.8	28.2	33.7	22.2
1999.....	4.6	*	*	*	*	*	0.9	6.5	20.0	29.9	31.1	23.9
Black, male												
2004.....	5.8	*	*	*	*	*	1.0	8.3	25.5	41.3	47.9	38.7
2003.....	6.2	*	*	*	*	*	1.3	10.2	28.7	44.3	46.0	38.6
2002.....	6.3	*	*	*	*	*	1.4	8.3	30.1	49.2	51.9	37.5
2001.....	7.0	*	*	*	*	*	1.5	10.2	32.5	55.3	54.7	46.5
2000.....	6.7	*	*	*	*	*	1.2	11.2	34.0	47.0	58.4	38.1
1999.....	7.0	*	*	*	*	*	1.7	11.3	34.8	50.6	56.9	51.6
Black, female												
2004.....	2.3	*	*	*	*	*	*	2.5	7.5	12.9	15.6	14.6
2003.....	2.3	*	*	*	*	*	*	2.5	7.4	12.2	17.7	14.6
2002.....	2.3	*	*	*	*	*	*	2.4	7.0	13.8	16.9	14.4
2001.....	2.3	*	*	*	*	*	*	2.2	7.3	16.1	15.4	15.2
2000.....	2.5	*	*	*	*	*	*	2.7	8.3	14.9	19.8	16.3
1999.....	2.3	*	*	*	*	*	*	2.4	8.2	15.2	16.7	13.6

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of stomach (C16)												
All races 3/, both sexes												
2004.....	4.0	*	*	*	*	0.3	1.1	2.7	6.5	14.9	26.3	42.2
2003.....	4.2	*	*	*	*	0.3	1.1	3.0	6.7	15.2	27.6	43.6
2002.....	4.2	*	*	*	0.1	0.3	1.0	2.8	6.9	15.4	28.7	47.6
2001.....	4.3	*	*	*	0.1	0.3	1.0	2.9	7.0	16.5	30.2	45.4
2000.....	4.5	*	*	*	*	0.3	1.1	2.9	7.4	16.6	31.0	52.7
1999.....	4.6	*	*	*	*	0.3	1.1	3.0	7.8	17.1	31.6	50.1
All races 3/, male												
2004.....	4.8	*	*	*	*	0.3	1.2	3.5	9.1	20.6	37.2	62.9
2003.....	4.9	*	*	*	*	0.3	1.2	3.8	9.5	21.9	38.4	60.0
2002.....	5.0	*	*	*	*	0.4	1.1	3.7	9.5	22.5	40.6	67.2
2001.....	5.1	*	*	*	*	0.4	1.1	3.8	10.2	23.2	43.1	62.6
2000.....	5.3	*	*	*	*	0.3	1.3	3.8	10.5	23.7	43.9	75.6
1999.....	5.5	*	*	*	*	0.4	1.3	4.2	11.0	24.2	46.5	71.3
All races 3/, female												
2004.....	3.3	*	*	*	*	0.3	0.9	2.0	4.1	10.1	19.0	32.9
2003.....	3.4	*	*	*	*	0.3	1.0	2.1	4.1	9.7	20.3	36.4
2002.....	3.5	*	*	*	0.1	0.3	0.9	1.9	4.5	9.5	20.8	39.1
2001.....	3.5	*	*	*	*	0.3	0.9	2.0	4.1	10.9	21.7	38.3
2000.....	3.7	*	*	*	*	0.3	0.9	2.1	4.6	10.8	22.6	43.4
1999.....	3.7	*	*	*	*	0.3	1.0	1.9	4.9	11.3	21.9	41.6
White, both sexes												
2004.....	3.8	*	*	*	*	0.3	0.9	2.3	5.5	12.9	23.7	38.8
2003.....	3.9	*	*	*	*	0.3	0.9	2.4	5.9	13.5	24.6	38.7
2002.....	4.0	*	*	*	0.1	0.3	0.8	2.3	5.9	13.4	25.4	42.2
2001.....	4.1	*	*	*	*	0.3	0.8	2.4	6.1	14.5	26.9	41.1
2000.....	4.2	*	*	*	*	0.2	0.9	2.5	6.5	14.5	27.3	47.3
1999.....	4.3	*	*	*	*	0.3	0.9	2.5	6.6	15.2	28.1	44.8
White, male												
2004.....	4.5	*	*	*	*	0.3	1.1	3.0	7.8	17.5	33.2	58.4
2003.....	4.6	*	*	*	*	0.3	1.0	3.1	8.3	19.2	34.3	53.5
2002.....	4.8	*	*	*	*	0.4	0.9	3.1	8.3	19.7	36.6	60.5
2001.....	4.9	*	*	*	*	0.3	1.0	3.2	8.8	20.2	38.7	57.7
2000.....	5.0	*	*	*	*	0.3	1.0	3.2	9.3	20.9	39.0	68.9
1999.....	5.1	*	*	*	*	0.3	1.0	3.5	9.3	21.7	41.5	63.4

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of stomach (C16)												
White, female												
2004.....	3.1	*	*	*	*	0.2	0.8	1.6	3.4	8.8	17.1	29.9
2003.....	3.2	*	*	*	*	0.3	0.8	1.7	3.6	8.7	17.9	32.1
2002.....	3.2	*	*	*	*	0.2	0.7	1.5	3.7	8.0	17.9	34.2
2001.....	3.3	*	*	*	*	0.2	0.7	1.7	3.5	9.7	19.0	34.1
2000.....	3.5	*	*	*	*	0.2	0.7	1.7	3.9	9.2	19.6	38.6
1999.....	3.4	*	*	*	*	0.2	0.8	1.5	4.0	9.8	19.3	37.3
Black, both sexes												
2004.....	5.3	*	*	*	*	0.5	1.6	5.3	12.8	30.2	50.4	71.3
2003.....	5.4	*	*	*	*	0.4	1.8	5.7	12.7	27.1	54.3	85.8
2002.....	5.6	*	*	*	*	0.6	1.6	5.4	12.9	30.9	56.8	100.4
2001.....	5.6	*	*	*	*	0.5	2.1	5.4	12.9	30.5	58.4	88.1
2000.....	5.9	*	*	*	*	0.7	2.0	5.7	13.8	32.0	63.9	101.7
1999.....	6.1	*	*	*	*	0.7	2.2	5.9	15.9	31.1	64.6	102.4
Black, male												
2004.....	6.5	*	*	*	*	*	1.9	7.2	19.4	47.3	75.2	100.3
2003.....	6.5	*	*	*	*	*	2.0	8.3	19.6	44.0	79.5	106.3
2002.....	6.6	*	*	*	*	0.8	1.8	7.7	18.9	46.7	79.8	137.2
2001.....	6.7	*	*	*	*	*	2.3	7.9	19.7	48.2	87.1	104.4
2000.....	7.0	*	*	*	*	0.8	2.9	7.9	20.6	46.7	91.1	136.1
1999.....	7.3	*	*	*	*	*	2.7	8.5	23.1	46.0	100.5	147.6
Black, female												
2004.....	4.2	*	*	*	*	*	1.4	3.6	7.5	18.1	36.3	59.9
2003.....	4.3	*	*	*	*	*	1.6	3.5	7.1	15.0	40.0	77.8
2002.....	4.7	*	*	*	*	*	1.4	3.4	8.0	19.8	43.9	86.3
2001.....	4.5	*	*	*	*	*	1.9	3.2	7.5	18.0	42.3	82.1
2000.....	5.0	*	*	*	*	*	1.3	3.7	8.4	21.6	48.6	88.9
1999.....	5.0	*	*	*	*	*	1.8	3.7	10.1	20.6	44.4	85.4

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of small intestine (C17)												
All races 3/, both sexes												
2004.....	0.4	*	*	*	*	*	0.1	0.3	0.7	1.5	2.4	3.1
2003.....	0.4	*	*	*	*	*	0.1	0.3	0.7	1.3	2.5	3.0
2002.....	0.4	*	*	*	*	*	0.1	0.3	0.7	1.4	2.3	2.8
2001.....	0.4	*	*	*	*	*	0.1	0.3	0.8	1.4	2.5	3.2
2000.....	0.4	*	*	*	*	*	0.1	0.3	0.8	1.5	2.4	3.0
1999.....	0.4	*	*	*	*	*	0.1	0.4	0.6	1.4	2.4	3.2
All races 3/, male												
2004.....	0.4	*	*	*	*	*	0.1	0.3	0.9	1.9	3.0	4.3
2003.....	0.4	*	*	*	*	*	*	0.3	0.9	1.7	2.7	3.5
2002.....	0.4	*	*	*	*	*	0.1	0.3	0.8	1.6	2.8	3.7
2001.....	0.4	*	*	*	*	*	0.1	0.4	0.9	1.9	3.0	4.1
2000.....	0.4	*	*	*	*	*	0.1	0.4	0.9	1.8	3.0	4.2
1999.....	0.4	*	*	*	*	*	0.1	0.5	0.9	1.7	3.2	4.4
All races 3/, female												
2004.....	0.4	*	*	*	*	*	0.1	0.3	0.6	1.1	1.9	2.5
2003.....	0.4	*	*	*	*	*	*	0.3	0.6	1.0	2.3	2.7
2002.....	0.3	*	*	*	*	*	*	0.2	0.5	1.2	2.0	2.4
2001.....	0.4	*	*	*	*	*	0.1	0.2	0.7	1.1	2.1	2.8
2000.....	0.4	*	*	*	*	*	0.1	0.3	0.6	1.2	2.1	2.5
1999.....	0.3	*	*	*	*	*	*	0.3	0.4	1.2	1.8	2.7
White, both sexes												
2004.....	0.4	*	*	*	*	*	0.1	0.3	0.6	1.4	2.3	2.9
2003.....	0.4	*	*	*	*	*	0.1	0.2	0.7	1.3	2.4	2.9
2002.....	0.4	*	*	*	*	*	0.1	0.3	0.6	1.4	2.2	2.6
2001.....	0.4	*	*	*	*	*	0.1	0.3	0.7	1.4	2.4	3.0
2000.....	0.4	*	*	*	*	*	0.1	0.3	0.7	1.4	2.4	3.1
1999.....	0.4	*	*	*	*	*	0.1	0.4	0.6	1.4	2.2	3.0
White, male												
2004.....	0.4	*	*	*	*	*	*	0.3	0.8	1.8	2.9	4.1
2003.....	0.4	*	*	*	*	*	*	0.3	0.9	1.6	2.6	3.3
2002.....	0.4	*	*	*	*	*	*	0.3	0.7	1.6	2.7	3.6
2001.....	0.4	*	*	*	*	*	0.1	0.3	0.8	1.8	2.9	3.7
2000.....	0.4	*	*	*	*	*	*	0.3	0.8	1.8	2.9	4.3
1999.....	0.4	*	*	*	*	*	0.1	0.5	0.9	1.7	3.0	4.4

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of colon, rectum and anus (C18-C21)												
All races 3/, both sexes												
2004.....	18.3	*	*	*	0.1	0.7	3.0	10.3	28.2	64.7	128.9	225.6
2003.....	19.2	*	*	*	0.1	0.7	3.0	10.9	29.8	70.5	134.7	239.7
2002.....	19.7	*	*	*	0.1	0.7	3.2	11.4	31.7	73.3	137.4	241.4
2001.....	20.0	*	*	*	0.1	0.8	3.2	11.6	31.5	75.3	139.4	255.7
2000.....	20.4	*	*	*	0.1	0.7	3.0	11.2	32.7	76.7	146.0	270.4
1999.....	20.5	*	*	*	0.1	0.7	3.0	11.1	33.5	77.4	146.7	271.7
All races 3/, male												
2004.....	18.6	*	*	*	0.2	0.7	3.2	11.9	34.2	80.6	156.9	254.7
2003.....	19.6	*	*	*	0.1	0.8	3.2	12.5	36.0	88.0	164.4	273.3
2002.....	20.1	*	*	*	0.2	0.8	3.6	13.0	39.4	91.3	167.4	275.1
2001.....	20.2	*	*	*	0.1	0.8	3.5	13.1	38.3	93.7	169.5	298.3
2000.....	20.6	*	*	*	0.1	0.8	3.4	12.8	39.6	95.4	179.5	315.6
1999.....	20.6	*	*	*	0.1	0.8	3.0	12.5	40.5	95.1	181.9	324.0
All races 3/, female												
2004.....	18.0	*	*	*	0.1	0.6	2.9	8.7	22.7	51.4	110.0	212.5
2003.....	18.9	*	*	*	*	0.7	2.7	9.3	24.0	55.9	114.8	224.8
2002.....	19.2	*	*	*	*	0.7	2.8	9.8	24.5	58.3	117.5	226.8
2001.....	19.8	*	*	*	0.1	0.7	2.9	10.2	25.3	60.1	119.6	237.8
2000.....	20.2	*	*	*	*	0.6	2.7	9.8	26.4	61.2	124.2	251.9
1999.....	20.3	*	*	*	0.1	0.6	2.9	9.7	27.0	62.9	123.8	250.6
White, both sexes												
2004.....	19.2	*	*	*	0.1	0.7	2.8	9.6	26.9	62.8	127.4	225.1
2003.....	20.2	*	*	*	0.1	0.7	2.7	10.0	28.2	68.7	133.3	238.5
2002.....	20.7	*	*	*	0.1	0.7	2.9	10.7	30.1	71.8	135.2	240.7
2001.....	21.0	*	*	*	0.1	0.7	3.0	10.8	30.1	73.4	136.9	255.3
2000.....	21.5	*	*	*	0.1	0.6	2.7	10.4	31.2	75.1	144.0	269.2
1999.....	21.6	*	*	*	0.1	0.7	2.8	10.4	32.0	75.6	144.5	270.9
White, male												
2004.....	19.5	*	*	*	0.1	0.7	3.0	11.2	32.8	78.2	154.6	255.4
2003.....	20.5	*	*	*	0.1	0.7	3.0	11.5	34.3	86.2	162.8	272.9
2002.....	21.1	*	*	*	0.1	0.8	3.3	12.2	37.6	89.8	164.2	273.6
2001.....	21.2	*	*	*	*	0.7	3.2	12.3	36.8	91.8	167.0	297.3
2000.....	21.6	*	*	*	*	0.7	3.1	12.0	37.6	93.4	176.9	313.3
1999.....	21.8	*	*	*	*	0.7	2.9	11.9	38.8	93.7	179.3	324.5

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of colon, rectum and anus (C18-C21)												
White, female												
2004.....	18.9	*	*	*	*	0.6	2.6	7.9	21.3	49.6	108.9	211.4
2003.....	19.8	*	*	*	*	0.6	2.4	8.4	22.5	53.8	113.4	223.2
2002.....	20.3	*	*	*	*	0.7	2.6	9.1	23.1	56.4	115.7	226.4
2001.....	20.7	*	*	*	*	0.7	2.7	9.3	23.7	57.9	116.8	237.7
2000.....	21.4	*	*	*	*	0.5	2.4	9.0	25.2	59.7	122.3	251.3
1999.....	21.4	*	*	*	*	0.6	2.6	8.9	25.7	60.5	121.7	249.3
Black, both sexes												
2004.....	17.2	*	*	*	*	1.0	4.4	16.6	44.8	94.6	165.1	260.8
2003.....	18.2	*	*	*	*	0.9	5.0	18.7	48.1	100.4	172.0	281.5
2002.....	18.2	*	*	*	*	0.7	4.9	18.6	49.5	99.2	183.6	274.7
2001.....	18.6	*	*	*	*	1.0	4.9	19.0	49.8	104.0	187.6	286.5
2000.....	18.8	*	*	*	*	1.0	5.1	19.5	50.1	104.7	190.1	310.0
1999.....	18.7	*	*	*	*	1.0	4.8	17.7	52.7	106.2	193.1	306.1
Black, male												
2004.....	17.4	*	*	*	*	1.0	4.5	19.1	53.4	120.4	214.6	278.2
2003.....	18.3	*	*	*	*	1.0	5.2	22.0	58.9	125.8	213.0	317.0
2002.....	18.6	*	*	*	*	1.0	5.8	21.2	61.2	121.6	237.5	328.1
2001.....	18.4	*	*	*	*	1.2	5.7	20.8	58.4	128.1	228.4	347.2
2000.....	19.1	*	*	*	*	1.0	6.0	22.0	62.5	131.5	241.0	377.1
1999.....	18.6	*	*	*	*	1.2	4.7	20.1	65.8	126.6	244.4	350.4
Black, female												
2004.....	17.0	*	*	*	*	0.9	4.3	14.4	37.8	76.2	137.0	254.0
2003.....	18.1	*	*	*	*	0.9	4.8	16.0	39.4	82.3	148.8	267.8
2002.....	17.8	*	*	*	*	*	4.1	16.3	40.2	83.3	153.0	254.2
2001.....	18.8	*	*	*	*	0.9	4.3	17.5	42.8	87.0	164.7	264.0
2000.....	18.5	*	*	*	*	1.0	4.3	17.2	40.2	85.8	161.4	285.0
1999.....	18.9	*	*	*	*	0.9	4.9	15.6	42.2	91.8	164.4	289.5

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Colon (C18)												
All races 3/, both sexes												
2004.....	15.2	*	*	*	0.1	0.5	2.1	7.8	22.3	53.8	109.8	195.3
2003.....	16.1	*	*	*	0.1	0.6	2.2	8.5	23.6	59.1	115.7	208.4
2002.....	16.5	*	*	*	0.1	0.5	2.4	8.9	25.3	61.0	118.5	211.3
2001.....	16.8	*	*	*	0.1	0.6	2.4	8.9	25.7	63.5	120.1	221.6
2000.....	17.3	*	*	*	0.1	0.5	2.3	8.9	26.5	64.3	126.5	236.6
1999.....	17.4	*	*	*	0.1	0.5	2.3	8.8	27.4	65.1	127.0	238.1
All races 3/, male												
2004.....	15.1	*	*	*	0.1	0.5	2.2	9.1	26.3	65.7	131.7	219.9
2003.....	16.1	*	*	*	0.1	0.6	2.4	9.6	28.0	72.9	140.1	236.1
2002.....	16.6	*	*	*	0.1	0.6	2.6	9.9	30.9	75.5	142.6	240.2
2001.....	16.7	*	*	*	*	0.6	2.6	9.9	30.6	77.7	144.4	256.7
2000.....	17.1	*	*	*	*	0.6	2.6	9.9	31.4	78.6	154.0	275.8
1999.....	17.2	*	*	*	0.1	0.6	2.3	9.8	32.7	79.2	156.1	282.2
All races 3/, female												
2004.....	15.2	*	*	*	*	0.5	2.1	6.6	18.5	43.8	95.1	184.2
2003.....	16.1	*	*	*	*	0.5	2.1	7.4	19.6	47.5	99.4	196.1
2002.....	16.4	*	*	*	*	0.5	2.1	7.9	20.2	49.0	102.6	198.7
2001.....	16.9	*	*	*	0.1	0.6	2.3	8.0	21.2	51.8	104.2	206.9
2000.....	17.4	*	*	*	*	0.5	2.1	7.9	22.0	52.5	108.5	220.6
1999.....	17.5	*	*	*	*	0.5	2.3	7.9	22.5	53.5	108.2	220.4
White, both sexes												
2004.....	15.9	*	*	*	0.1	0.5	2.0	7.2	21.0	51.7	108.1	194.8
2003.....	16.9	*	*	*	0.1	0.5	2.0	7.6	22.2	57.3	114.4	206.9
2002.....	17.3	*	*	*	0.1	0.5	2.2	8.2	23.9	59.4	116.5	210.5
2001.....	17.6	*	*	*	0.1	0.5	2.2	8.2	24.2	61.6	117.6	221.2
2000.....	18.1	*	*	*	0.1	0.5	2.1	8.1	25.1	62.8	124.3	235.0
1999.....	18.2	*	*	*	0.1	0.5	2.1	8.2	25.9	63.4	124.9	237.1
White, male												
2004.....	15.8	*	*	*	*	0.5	2.1	8.5	25.0	63.3	129.1	220.7
2003.....	16.9	*	*	*	*	0.6	2.2	8.8	26.5	71.1	138.5	236.2
2002.....	17.3	*	*	*	*	0.6	2.4	9.2	29.3	73.9	139.7	238.7
2001.....	17.5	*	*	*	*	0.6	2.4	9.1	29.3	75.9	141.6	255.5
2000.....	17.9	*	*	*	*	0.6	2.3	9.2	29.7	77.0	151.1	272.5
1999.....	18.2	*	*	*	*	0.5	2.2	9.3	31.1	78.0	153.6	282.9

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Colon (C18)												
White, female												
2004.....	16.0	*	*	*	*	0.5	1.8	6.0	17.2	41.8	93.8	183.1
2003.....	16.8	*	*	*	*	0.5	1.8	6.5	18.2	45.5	98.1	193.9
2002.....	17.3	*	*	*	*	0.5	2.0	7.2	18.8	47.1	100.9	198.2
2001.....	17.7	*	*	*	*	0.5	2.1	7.3	19.5	49.5	101.6	206.9
2000.....	18.3	*	*	*	*	0.4	1.9	7.1	20.8	50.9	106.6	219.8
1999.....	18.3	*	*	*	*	0.5	2.0	7.1	21.1	51.2	106.1	218.7
Black, both sexes												
2004.....	14.7	*	*	*	*	0.6	3.1	13.2	37.8	82.9	146.7	227.3
2003.....	15.6	*	*	*	*	0.7	4.0	15.4	40.1	87.2	151.4	249.4
2002.....	15.7	*	*	*	*	0.6	3.8	15.5	41.7	86.0	161.6	244.1
2001.....	16.1	*	*	*	*	0.8	4.0	15.7	42.9	91.6	165.7	249.9
2000.....	16.3	*	*	*	*	0.7	4.0	16.3	42.5	89.6	171.0	279.7
1999.....	16.3	*	*	*	*	0.9	4.0	15.0	45.0	92.2	171.6	273.9
Black, male												
2004.....	14.5	*	*	*	*	*	2.7	15.1	44.4	103.2	187.4	243.4
2003.....	15.4	*	*	*	*	*	3.9	17.4	48.4	108.2	188.1	270.0
2002.....	15.8	*	*	*	*	*	4.3	17.0	50.9	104.9	206.3	292.7
2001.....	15.6	*	*	*	*	0.8	4.3	17.3	48.4	109.5	201.9	307.5
2000.....	16.1	*	*	*	*	*	4.5	17.9	51.6	108.4	215.6	346.0
1999.....	15.8	*	*	*	*	0.9	3.7	16.3	54.7	108.5	214.3	304.7
Black, female												
2004.....	14.8	*	*	*	*	*	3.4	11.6	32.5	68.4	123.7	220.9
2003.....	15.8	*	*	*	*	*	4.0	13.6	33.5	72.3	130.6	241.4
2002.....	15.6	*	*	*	*	*	3.3	14.2	34.3	72.7	136.3	225.5
2001.....	16.5	*	*	*	*	0.9	3.8	14.3	38.5	79.0	145.4	228.6
2000.....	16.4	*	*	*	*	0.7	3.6	14.9	35.2	76.4	145.9	255.0
1999.....	16.8	*	*	*	*	0.8	4.3	13.9	37.2	80.7	147.7	262.4

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Rectosigmoid junction and rectum (C19-C20)												
All races 3/, both sexes												
2004.....	2.9	*	*	*	*	0.2	0.8	2.1	5.5	10.3	18.1	29.1
2003.....	2.9	*	*	*	*	0.1	0.6	2.2	5.7	10.9	18.0	29.9
2002.....	3.0	*	*	*	*	0.2	0.7	2.2	5.9	11.7	18.0	29.0
2001.....	3.0	*	*	*	*	0.2	0.7	2.5	5.5	11.2	18.3	32.6
2000.....	3.0	*	*	*	*	0.1	0.6	2.1	5.8	11.8	18.8	32.6
1999.....	3.0	*	*	*	*	0.2	0.6	2.0	5.8	11.7	18.8	32.1
All races 3/, male												
2004.....	3.3	*	*	*	*	0.2	0.9	2.6	7.4	14.3	24.5	33.5
2003.....	3.3	*	*	*	*	0.2	0.7	2.7	7.6	14.6	23.7	36.4
2002.....	3.4	*	*	*	*	0.2	0.8	2.9	8.1	15.3	24.2	33.6
2001.....	3.4	*	*	*	*	0.2	0.8	3.0	7.4	15.5	24.4	40.7
2000.....	3.4	*	*	*	*	0.2	0.7	2.6	7.8	16.3	24.8	38.9
1999.....	3.3	*	*	*	*	0.2	0.6	2.5	7.6	15.4	25.1	40.7
All races 3/, female												
2004.....	2.5	*	*	*	*	0.2	0.7	1.7	3.8	7.0	13.7	27.2
2003.....	2.6	*	*	*	*	0.1	0.5	1.6	3.9	7.8	14.2	27.0
2002.....	2.6	*	*	*	*	0.2	0.5	1.7	3.8	8.7	13.9	27.0
2001.....	2.6	*	*	*	*	0.1	0.5	1.9	3.7	7.6	14.4	29.2
2000.....	2.6	*	*	*	*	0.1	0.5	1.6	4.0	8.1	14.8	30.1
1999.....	2.7	*	*	*	*	0.1	0.5	1.6	4.1	8.7	14.7	28.7
White, both sexes												
2004.....	3.1	*	*	*	*	0.2	0.7	2.1	5.5	10.5	18.3	29.1
2003.....	3.1	*	*	*	*	0.1	0.6	2.1	5.6	10.8	18.0	30.1
2002.....	3.2	*	*	*	*	0.2	0.6	2.2	5.8	11.8	17.8	29.0
2001.....	3.2	*	*	*	*	0.2	0.6	2.4	5.4	11.2	18.4	32.6
2000.....	3.2	*	*	*	*	0.1	0.5	2.1	5.7	11.6	18.9	33.0
1999.....	3.2	*	*	*	*	0.2	0.6	2.0	5.7	11.7	18.8	32.2
White, male												
2004.....	3.5	*	*	*	*	0.2	0.8	2.6	7.4	14.4	24.7	33.6
2003.....	3.5	*	*	*	*	0.2	0.7	2.6	7.5	14.6	23.9	36.0
2002.....	3.6	*	*	*	*	0.2	0.7	2.8	8.0	15.4	23.9	33.7
2001.....	3.6	*	*	*	*	0.2	0.8	3.0	7.3	15.4	24.7	41.0
2000.....	3.5	*	*	*	*	0.1	0.7	2.6	7.6	15.9	25.0	39.8
1999.....	3.5	*	*	*	*	0.2	0.6	2.4	7.4	15.3	25.0	40.6

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Rectosigmoid junction and rectum (C19-C20)

White, female

2004.....	2.7	*	*	*	*	0.2	0.6	1.6	3.7	7.1	13.9	27.1
2003.....	2.7	*	*	*	*	*	0.5	1.6	3.7	7.6	14.1	27.5
2002.....	2.8	*	*	*	*	0.2	0.5	1.6	3.7	8.6	13.8	27.0
2001.....	2.8	*	*	*	*	0.1	0.5	1.8	3.7	7.7	14.1	29.1
2000.....	2.8	*	*	*	*	*	0.4	1.6	3.9	8.1	14.9	30.3
1999.....	2.9	*	*	*	*	*	0.6	1.6	4.1	8.7	14.7	28.9

Black, both sexes

2004.....	2.4	*	*	*	*	*	1.2	3.1	6.7	11.1	17.8	31.9
2003.....	2.4	*	*	*	*	*	0.9	3.1	7.5	12.3	19.5	31.5
2002.....	2.4	*	*	*	*	*	0.9	2.8	7.3	12.6	21.4	29.7
2001.....	2.4	*	*	*	*	*	0.8	3.1	6.7	12.0	21.0	35.6
2000.....	2.4	*	*	*	*	*	0.9	2.9	7.3	14.5	18.2	28.8
1999.....	2.3	*	*	*	*	*	0.6	2.4	7.5	13.2	20.7	31.9

Black, male

2004.....	2.7	*	*	*	*	*	1.5	3.7	8.7	16.4	27.0	31.8
2003.....	2.8	*	*	*	*	*	1.0	4.2	10.3	16.7	24.0	45.9
2002.....	2.7	*	*	*	*	*	1.2	3.9	9.9	16.3	30.4	34.3
2001.....	2.7	*	*	*	*	*	1.3	3.4	9.7	18.2	25.6	38.6
2000.....	2.8	*	*	*	*	*	1.2	3.8	10.4	22.5	24.2	30.0
1999.....	2.7	*	*	*	*	*	0.8	3.4	10.8	17.6	29.8	45.7

Black, female

2004.....	2.1	*	*	*	*	*	0.8	2.5	5.0	7.3	12.6	31.9
2003.....	2.1	*	*	*	*	*	0.8	2.2	5.3	9.2	16.9	25.9
2002.....	2.1	*	*	*	*	*	0.7	2.0	5.3	10.0	16.2	27.9
2001.....	2.1	*	*	*	*	*	*	2.8	4.2	7.6	18.3	34.5
2000.....	2.0	*	*	*	*	*	*	2.2	4.8	8.9	14.8	28.3
1999.....	1.9	*	*	*	*	*	*	1.6	4.8	10.1	15.7	26.7

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Anus and anus canal (C21)												
All races 3/, both sexes												
2004.....	0.2	*	*	*	*	*	0.1	0.3	0.4	0.6	1.0	1.2
2003.....	0.2	*	*	*	*	*	0.1	0.2	0.4	0.6	0.9	1.4
2002.....	0.2	*	*	*	*	*	0.1	0.3	0.4	0.6	0.9	1.2
2001.....	0.2	*	*	*	*	*	0.1	0.2	0.4	0.6	0.9	1.4
2000.....	0.2	*	*	*	*	*	0.1	0.2	0.4	0.6	0.8	1.2
1999.....	0.2	*	*	*	*	*	0.1	0.2	0.3	0.6	0.8	1.4
All races 3/, male												
2004.....	0.2	*	*	*	*	*	0.1	0.2	0.4	0.6	0.7	*
2003.....	0.1	*	*	*	*	*	0.1	0.2	0.3	0.5	0.5	*
2002.....	0.2	*	*	*	*	*	0.1	0.3	0.4	0.5	0.6	*
2001.....	0.1	*	*	*	*	*	0.1	0.2	0.3	0.6	0.7	*
2000.....	0.1	*	*	*	*	*	0.1	0.2	0.4	0.5	0.8	*
1999.....	0.1	*	*	*	*	*	*	0.2	0.3	0.4	0.6	*
All races 3/, female												
2004.....	0.2	*	*	*	*	*	0.1	0.3	0.4	0.7	1.1	1.1
2003.....	0.2	*	*	*	*	*	*	0.3	0.5	0.6	1.2	1.6
2002.....	0.2	*	*	*	*	*	0.1	0.3	0.5	0.6	1.0	1.2
2001.....	0.2	*	*	*	*	*	0.1	0.3	0.4	0.6	1.1	1.7
2000.....	0.2	*	*	*	*	*	0.1	0.3	0.4	0.6	0.9	1.3
1999.....	0.2	*	*	*	*	*	*	0.2	0.4	0.7	0.9	1.6
White, both sexes												
2004.....	0.2	*	*	*	*	*	0.1	0.3	0.5	0.6	1.0	1.1
2003.....	0.2	*	*	*	*	*	0.1	0.2	0.4	0.6	0.9	1.5
2002.....	0.2	*	*	*	*	*	0.1	0.3	0.4	0.6	0.9	1.2
2001.....	0.2	*	*	*	*	*	0.1	0.2	0.4	0.6	0.9	1.5
2000.....	0.2	*	*	*	*	*	0.1	0.3	0.4	0.6	0.8	1.1
1999.....	0.2	*	*	*	*	*	0.1	0.2	0.4	0.5	0.8	1.5
White, male												
2004.....	0.2	*	*	*	*	*	0.1	0.2	0.5	0.5	0.7	*
2003.....	0.1	*	*	*	*	*	*	0.1	0.4	0.5	0.5	*
2002.....	0.2	*	*	*	*	*	*	0.3	0.4	0.5	0.6	*
2001.....	0.1	*	*	*	*	*	*	0.2	0.3	0.6	0.7	*
2000.....	0.1	*	*	*	*	*	*	0.2	0.4	0.5	0.8	*
1999.....	0.1	*	*	*	*	*	*	0.2	0.3	0.4	0.7	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Of liver and intrahepatic bile ducts (C22)

All races 3/, both sexes

2004.....	5.2	*	0.1	*	0.1	0.2	0.9	5.6	10.7	19.9	31.3	32.4
2003.....	5.1	*	0.1	*	0.1	0.2	0.9	5.4	9.9	19.8	30.7	33.4
2002.....	4.9	*	0.2	*	0.1	0.3	0.9	4.9	9.6	19.6	29.3	35.0
2001.....	4.7	*	*	*	0.1	0.2	0.9	4.8	9.3	18.8	29.0	34.1
2000.....	4.6	*	0.1	0.0	0.1	0.2	1.0	4.5	9.3	18.7	28.8	32.7
1999.....	4.4	*	0.2	*	0.1	0.2	1.0	4.1	8.8	19.0	28.0	30.6

All races 3/, male

2004.....	6.8	*	*	*	0.1	0.3	1.3	9.2	16.6	28.3	42.7	41.5
2003.....	6.6	*	*	*	0.1	0.3	1.3	8.9	14.9	27.9	44.6	47.5
2002.....	6.4	*	*	*	*	0.3	1.4	8.2	14.5	28.5	41.6	46.2
2001.....	6.1	*	*	*	*	0.2	1.3	7.7	13.9	26.8	41.0	50.3
2000.....	5.9	*	*	*	0.1	0.3	1.3	7.2	13.8	27.0	42.1	44.7
1999.....	5.6	*	0.3	*	0.1	0.2	1.4	6.5	12.9	27.5	39.0	40.1

All races 3/, female

2004.....	3.7	*	*	*	*	0.1	0.4	2.2	5.3	12.8	23.6	28.3
2003.....	3.5	*	*	*	*	0.2	0.5	2.1	5.3	13.0	21.4	27.2
2002.....	3.4	*	*	*	*	0.2	0.5	1.8	5.1	12.2	21.1	30.2
2001.....	3.4	*	*	*	*	0.2	0.4	1.9	5.0	12.1	21.1	27.2
2000.....	3.3	*	*	*	*	0.1	0.6	1.9	5.1	11.8	20.1	27.8
1999.....	3.3	*	*	*	*	0.2	0.6	1.8	5.1	11.9	20.9	26.8

White, both sexes

2004.....	5.1	*	*	*	0.1	0.1	0.7	4.8	9.4	18.3	30.0	31.2
2003.....	4.9	*	*	*	0.1	0.1	0.7	4.6	8.7	18.3	29.5	32.1
2002.....	4.8	*	0.2	*	*	0.2	0.8	4.1	8.5	18.3	28.4	33.8
2001.....	4.6	*	*	*	0.1	0.2	0.6	4.0	8.2	17.6	27.9	33.3
2000.....	4.5	*	*	*	0.1	0.1	0.7	3.7	8.1	17.3	27.8	32.0
1999.....	4.4	*	0.2	*	0.1	0.1	0.8	3.4	7.6	17.8	27.2	29.9

White, male

2004.....	6.5	*	*	*	*	0.2	1.0	7.9	14.2	25.9	41.4	39.9
2003.....	6.4	*	*	*	*	0.1	1.0	7.4	13.1	26.0	43.0	45.6
2002.....	6.1	*	*	*	*	0.2	1.1	6.8	12.6	26.5	40.0	45.1
2001.....	5.9	*	*	*	*	0.1	0.9	6.4	12.2	25.4	39.6	49.7
2000.....	5.7	*	*	*	*	0.2	1.0	5.7	11.8	25.0	41.1	43.1
1999.....	5.5	*	*	*	*	0.1	1.1	5.2	11.1	25.8	38.0	40.6

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Of liver and intrahepatic bile ducts (C22)

White, female

2004.....	3.7	*	*	*	*	*	0.4	1.9	4.9	11.8	22.3	27.3
2003.....	3.5	*	*	*	*	0.1	0.5	1.8	4.6	11.7	20.3	26.2
2002.....	3.5	*	*	*	*	0.2	0.4	1.5	4.5	11.4	20.6	29.0
2001.....	3.3	*	*	*	*	0.2	0.3	1.6	4.4	10.9	20.1	26.5
2000.....	3.3	*	*	*	*	*	0.4	1.7	4.6	10.9	19.0	27.4
1999.....	3.3	*	*	*	*	*	0.5	1.6	4.4	11.0	20.1	25.6

Black, both sexes

2004.....	5.2	*	*	*	*	0.4	1.4	9.6	17.4	25.6	34.4	36.1
2003.....	4.9	*	*	*	*	0.5	1.4	9.8	16.3	22.9	31.3	41.7
2002.....	4.6	*	*	*	*	0.5	1.4	8.7	14.8	23.0	30.9	38.0
2001.....	4.5	*	*	*	*	*	1.7	8.7	14.8	21.7	32.4	35.0
2000.....	4.4	*	*	*	*	0.4	1.8	8.6	14.1	23.6	28.7	31.6
1999.....	4.2	*	*	*	*	0.4	1.6	7.2	13.2	24.2	29.4	34.4

Black, male

2004.....	7.3	*	*	*	*	*	2.2	16.7	29.9	40.3	44.0	46.7
2003.....	6.8	*	*	*	*	*	2.2	17.1	25.9	33.5	44.3	63.6
2002.....	6.5	*	*	*	*	*	2.2	15.1	24.2	37.7	47.0	41.8
2001.....	5.9	*	*	*	*	*	2.6	14.8	23.6	30.0	43.0	42.0
2000.....	6.1	*	*	*	*	*	2.4	14.8	23.5	37.3	39.7	48.4
1999.....	5.5	*	*	*	*	*	2.4	12.8	20.0	36.5	41.2	30.5

Black, female

2004.....	3.3	*	*	*	*	*	*	3.5	7.3	15.0	29.0	31.9
2003.....	3.2	*	*	*	*	*	*	3.5	8.6	15.3	23.9	33.2
2002.....	2.9	*	*	*	*	*	0.7	3.2	7.3	12.6	21.8	36.6
2001.....	3.2	*	*	*	*	*	0.8	3.5	7.7	15.9	26.4	32.4
2000.....	2.8	*	*	*	*	*	1.2	3.3	6.5	14.0	22.5	25.3
1999.....	3.0	*	*	*	*	*	0.9	2.4	7.8	15.6	22.7	35.9

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Liver (C22.0,C22.2-C22.9)												
All races 3/, both sexes												
2004.....	4.0	*	0.1	*	0.1	0.2	0.6	4.9	8.5	15.2	22.9	23.2
2003.....	3.9	*	0.1	*	0.1	0.2	0.7	4.6	7.8	15.3	23.0	24.3
2002.....	3.8	*	0.2	*	0.1	0.2	0.7	4.3	7.6	15.7	22.2	24.7
2001.....	3.7	*	*	*	0.1	0.2	0.7	4.2	7.5	14.7	21.9	24.2
2000.....	3.6	*	0.1	0.0	0.1	0.2	0.8	3.8	7.5	14.8	22.1	23.4
1999.....	3.5	*	0.2	*	0.1	0.2	0.8	3.5	7.0	15.2	21.5	22.5
All races 3/, male												
2004.....	5.6	*	*	*	0.1	0.2	1.1	8.4	13.9	23.2	33.1	30.6
2003.....	5.5	*	*	*	*	0.2	1.1	8.1	12.4	22.5	35.4	35.2
2002.....	5.3	*	*	*	*	0.3	1.2	7.4	12.3	24.0	32.7	34.6
2001.....	5.1	*	*	*	*	0.2	1.1	7.1	12.0	22.0	32.2	38.5
2000.....	5.0	*	*	*	*	0.2	1.2	6.4	11.9	22.4	33.7	33.9
1999.....	4.7	*	0.3	*	0.1	0.2	1.2	5.9	10.9	23.1	31.5	31.2
All races 3/, female												
2004.....	2.5	*	*	*	*	*	0.2	1.5	3.4	8.6	16.1	19.8
2003.....	2.4	*	*	*	*	0.1	0.3	1.3	3.5	9.3	14.7	19.5
2002.....	2.4	*	*	*	*	0.1	0.3	1.3	3.2	8.7	15.2	20.4
2001.....	2.4	*	*	*	*	0.1	0.3	1.4	3.4	8.6	15.1	18.2
2000.....	2.3	*	*	*	*	0.1	0.4	1.3	3.5	8.5	14.4	19.1
1999.....	2.4	*	*	*	*	0.1	0.4	1.2	3.5	8.6	15.0	18.9
White, both sexes												
2004.....	3.8	*	*	*	0.1	0.1	0.5	4.1	7.1	13.7	21.7	21.9
2003.....	3.7	*	*	*	0.1	0.1	0.5	3.8	6.6	13.9	21.7	23.0
2002.....	3.7	*	0.2	*	*	0.1	0.6	3.5	6.5	14.4	21.4	23.7
2001.....	3.5	*	*	*	*	0.1	0.5	3.4	6.5	13.5	20.8	23.5
2000.....	3.4	*	*	*	*	0.1	0.6	3.0	6.4	13.3	21.0	22.5
1999.....	3.4	*	0.2	*	0.1	0.1	0.6	2.8	5.9	13.9	20.7	21.7
White, male												
2004.....	5.3	*	*	*	*	0.1	0.8	7.0	11.6	20.8	31.6	28.8
2003.....	5.2	*	*	*	*	*	0.8	6.6	10.6	20.8	33.7	33.1
2002.....	5.0	*	*	*	*	0.1	0.9	6.1	10.5	22.0	31.4	33.5
2001.....	4.8	*	*	*	*	*	0.7	5.8	10.4	20.7	30.9	37.8
2000.....	4.6	*	*	*	*	*	0.8	5.0	10.0	20.3	32.6	31.8
1999.....	4.5	*	*	*	*	*	0.9	4.7	9.2	21.4	30.6	31.4

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Liver (C22.0,C22.2-C22.9)												
White, female												
2004.....	2.4	*	*	*	*	*	0.2	1.2	2.9	7.6	14.9	18.8
2003.....	2.3	*	*	*	*	*	0.2	1.1	2.9	8.0	13.6	18.5
2002.....	2.4	*	*	*	*	*	0.3	1.0	2.8	7.9	14.7	19.4
2001.....	2.3	*	*	*	*	0.1	0.2	1.1	2.8	7.4	14.1	17.5
2000.....	2.3	*	*	*	*	*	0.3	1.0	3.0	7.5	13.4	18.7
1999.....	2.3	*	*	*	*	*	0.3	1.0	2.9	7.7	14.2	17.8
Black, both sexes												
2004.....	4.5	*	*	*	*	0.4	1.1	8.8	15.7	21.2	27.7	29.0
2003.....	4.2	*	*	*	*	0.4	1.3	8.7	14.2	18.6	25.3	33.0
2002.....	4.0	*	*	*	*	0.4	1.3	8.1	13.0	19.7	24.5	28.8
2001.....	3.9	*	*	*	*	*	1.5	8.0	12.8	18.4	26.1	27.0
2000.....	3.8	*	*	*	*	*	1.6	8.0	12.1	20.4	24.0	24.7
1999.....	3.6	*	*	*	*	*	1.3	6.9	11.6	20.9	24.3	28.4
Black, male												
2004.....	6.7	*	*	*	*	*	2.0	15.8	27.9	35.2	38.1	36.8
2003.....	6.0	*	*	*	*	*	2.1	16.0	23.5	27.7	37.2	51.1
2002.....	5.8	*	*	*	*	*	2.1	14.5	22.3	32.8	38.2	35.4
2001.....	5.3	*	*	*	*	*	2.5	14.1	20.9	25.6	35.3	34.0
2000.....	5.5	*	*	*	*	*	2.2	14.2	21.5	33.2	34.2	42.7
1999.....	5.0	*	*	*	*	*	2.2	12.5	18.8	32.4	34.7	24.6
Black, female												
2004.....	2.5	*	*	*	*	*	*	2.8	5.8	11.2	21.9	26.0
2003.....	2.5	*	*	*	*	*	*	2.4	6.8	12.1	18.5	25.9
2002.....	2.3	*	*	*	*	*	*	2.6	5.5	10.4	16.7	26.3
2001.....	2.6	*	*	*	*	*	*	2.7	6.3	13.3	21.0	24.4
2000.....	2.2	*	*	*	*	*	1.0	2.6	4.6	11.3	18.2	18.0
1999.....	2.4	*	*	*	*	*	*	2.1	5.9	12.9	18.4	29.8

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Intrahepatic bile duct carcinoma (C22.1)												
All races 3/, both sexes												
2004.....	1.2	*	*	*	*	0.1	0.2	0.8	2.3	4.7	8.3	9.2
2003.....	1.1	*	*	*	*	0.1	0.2	0.8	2.1	4.5	7.7	9.1
2002.....	1.0	*	*	*	*	0.1	0.2	0.6	2.0	3.9	7.1	10.3
2001.....	1.0	*	*	*	*	*	0.2	0.6	1.8	4.1	7.1	9.9
2000.....	1.0	*	*	*	*	*	0.2	0.7	1.8	3.9	6.7	9.3
1999.....	0.9	*	*	*	*	*	0.2	0.6	1.7	3.8	6.5	8.1
All races 3/, male												
2004.....	1.2	*	*	*	*	*	0.2	0.9	2.6	5.1	9.5	10.9
2003.....	1.2	*	*	*	*	*	0.2	0.8	2.5	5.4	9.2	12.4
2002.....	1.0	*	*	*	*	*	0.2	0.8	2.2	4.5	8.8	11.6
2001.....	1.0	*	*	*	*	*	0.2	0.7	1.9	4.8	8.8	11.9
2000.....	1.0	*	*	*	*	*	0.2	0.7	1.9	4.6	8.3	10.8
1999.....	0.9	*	*	*	*	*	0.2	0.6	1.9	4.4	7.5	8.9
All races 3/, female												
2004.....	1.2	*	*	*	*	*	0.2	0.7	1.9	4.3	7.5	8.5
2003.....	1.1	*	*	*	*	*	0.2	0.8	1.8	3.7	6.6	7.6
2002.....	1.0	*	*	*	*	*	0.2	0.5	1.8	3.4	5.9	9.8
2001.....	1.0	*	*	*	*	*	0.1	0.5	1.6	3.6	6.0	9.0
2000.....	1.0	*	*	*	*	*	0.1	0.6	1.7	3.3	5.7	8.8
1999.....	0.9	*	*	*	*	*	0.2	0.6	1.6	3.3	5.8	7.8
White, both sexes												
2004.....	1.3	*	*	*	*	0.1	0.2	0.7	2.3	4.6	8.3	9.3
2003.....	1.2	*	*	*	*	0.1	0.2	0.8	2.1	4.4	7.8	9.2
2002.....	1.1	*	*	*	*	*	0.2	0.6	2.0	3.9	7.0	10.2
2001.....	1.1	*	*	*	*	*	0.2	0.6	1.7	4.1	7.0	9.8
2000.....	1.0	*	*	*	*	*	0.2	0.7	1.7	4.0	6.7	9.5
1999.....	1.0	*	*	*	*	*	0.2	0.6	1.7	3.9	6.5	8.3
White, male												
2004.....	1.3	*	*	*	*	*	0.3	0.8	2.6	5.0	9.8	11.1
2003.....	1.2	*	*	*	*	*	0.2	0.7	2.4	5.2	9.4	12.5
2002.....	1.1	*	*	*	*	*	0.2	0.7	2.2	4.5	8.6	11.6
2001.....	1.1	*	*	*	*	*	0.2	0.7	1.8	4.7	8.7	11.8
2000.....	1.0	*	*	*	*	*	0.2	0.7	1.8	4.7	8.4	11.3
1999.....	0.9	*	*	*	*	*	0.2	0.6	1.9	4.5	7.4	9.2

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Intrahepatic bile duct carcinoma (C22.1)												
White, female												
2004.....	1.3	*	*	*	*	*	0.2	0.6	2.0	4.2	7.4	8.5
2003.....	1.2	*	*	*	*	*	0.2	0.8	1.7	3.7	6.7	7.7
2002.....	1.1	*	*	*	*	*	0.2	0.5	1.7	3.5	5.9	9.6
2001.....	1.1	*	*	*	*	*	0.1	0.5	1.6	3.6	5.9	9.0
2000.....	1.0	*	*	*	*	*	0.2	0.6	1.6	3.4	5.6	8.8
1999.....	1.0	*	*	*	*	*	0.2	0.6	1.5	3.4	5.9	7.9
Black, both sexes												
2004.....	0.7	*	*	*	*	*	*	0.8	1.7	4.4	6.6	7.0
2003.....	0.7	*	*	*	*	*	*	1.1	2.1	4.3	6.0	8.8
2002.....	0.6	*	*	*	*	*	*	0.6	1.9	3.3	6.4	9.2
2001.....	0.6	*	*	*	*	*	*	0.7	2.0	3.4	6.2	8.0
2000.....	0.6	*	*	*	*	*	*	0.7	2.0	3.3	4.7	6.9
1999.....	0.5	*	*	*	*	*	*	*	1.6	3.3	5.1	*
Black, male												
2004.....	0.7	*	*	*	*	*	*	0.9	2.0	5.1	5.8	*
2003.....	0.7	*	*	*	*	*	*	1.1	2.4	5.8	7.1	*
2002.....	0.6	*	*	*	*	*	*	*	1.9	4.9	8.9	*
2001.....	0.6	*	*	*	*	*	*	*	2.7	4.3	7.7	*
2000.....	0.5	*	*	*	*	*	*	*	2.0	4.1	*	*
1999.....	0.5	*	*	*	*	*	*	*	*	4.1	6.5	*
Black, female												
2004.....	0.8	*	*	*	*	*	*	0.8	1.4	3.9	7.1	*
2003.....	0.7	*	*	*	*	*	*	1.1	1.8	3.2	5.4	*
2002.....	0.6	*	*	*	*	*	*	*	1.8	2.1	5.0	10.3
2001.....	0.6	*	*	*	*	*	*	*	*	2.7	5.5	*
2000.....	0.6	*	*	*	*	*	*	*	1.9	2.7	4.3	*
1999.....	0.6	*	*	*	*	*	*	*	1.9	2.7	4.3	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Of gallbladder and extrahepatic bile ducts (C23-C24)

All races 3/, both sexes

2004.....	1.2	*	*	*	*	*	0.1	0.4	1.8	4.7	8.7	13.3
2003.....	1.2	*	*	*	*	*	0.1	0.6	1.7	4.2	9.3	14.3
2002.....	1.2	*	*	*	*	*	0.1	0.6	1.7	4.6	8.9	14.5
2001.....	1.3	*	*	*	*	*	0.2	0.6	1.8	4.8	9.9	15.8
2000.....	1.3	*	*	*	*	*	0.1	0.6	2.0	5.3	10.0	16.0
1999.....	1.3	*	*	*	*	*	0.1	0.6	2.0	5.1	10.0	16.2

All races 3/, male

2004.....	0.9	*	*	*	*	*	*	0.3	1.4	4.1	8.6	12.9
2003.....	0.9	*	*	*	*	*	0.1	0.5	1.5	4.2	8.5	14.1
2002.....	0.9	*	*	*	*	*	0.1	0.4	1.4	4.3	8.5	15.0
2001.....	1.0	*	*	*	*	*	0.1	0.5	1.4	4.0	9.9	15.8
2000.....	1.0	*	*	*	*	*	0.1	0.4	1.6	5.0	9.5	15.8
1999.....	1.0	*	*	*	*	*	0.1	0.5	1.6	4.5	9.3	15.2

All races 3/, female

2004.....	1.4	*	*	*	*	*	0.1	0.5	2.1	5.2	8.8	13.5
2003.....	1.4	*	*	*	*	*	0.1	0.7	2.0	4.3	9.8	14.4
2002.....	1.4	*	*	*	*	*	0.2	0.8	1.9	4.9	9.2	14.3
2001.....	1.6	*	*	*	*	*	0.2	0.7	2.1	5.5	9.9	15.8
2000.....	1.6	*	*	*	*	*	0.1	0.8	2.3	5.5	10.4	16.0
1999.....	1.6	*	*	*	*	*	0.2	0.7	2.4	5.5	10.5	16.7

White, both sexes

2004.....	1.2	*	*	*	*	*	0.1	0.4	1.7	4.5	8.6	13.4
2003.....	1.3	*	*	*	*	*	0.1	0.5	1.7	4.2	9.2	14.4
2002.....	1.3	*	*	*	*	*	0.1	0.5	1.6	4.6	8.8	14.5
2001.....	1.4	*	*	*	*	*	0.2	0.6	1.8	4.7	9.9	15.8
2000.....	1.4	*	*	*	*	*	0.1	0.6	1.9	5.2	10.2	16.3
1999.....	1.4	*	*	*	*	*	0.1	0.5	2.0	5.1	9.9	16.4

White, male

2004.....	0.9	*	*	*	*	*	*	0.4	1.4	3.9	8.5	13.2
2003.....	1.0	*	*	*	*	*	*	0.5	1.4	4.3	8.4	14.5
2002.....	1.0	*	*	*	*	*	*	0.4	1.4	4.3	8.4	15.3
2001.....	1.0	*	*	*	*	*	0.1	0.5	1.4	4.0	9.8	15.5
2000.....	1.1	*	*	*	*	*	0.1	0.5	1.6	5.0	9.6	16.2
1999.....	1.0	*	*	*	*	*	0.1	0.4	1.6	4.4	9.3	15.8

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Of gallbladder and extrahepatic bile ducts (C23-C24)

White, female

2004.....	1.5	*	*	*	*	*	0.1	0.5	2.0	5.0	8.7	13.5
2003.....	1.5	*	*	*	*	*	*	0.6	2.0	4.2	9.7	14.4
2002.....	1.5	*	*	*	*	*	0.2	0.7	1.9	4.9	9.0	14.2
2001.....	1.7	*	*	*	*	*	0.2	0.7	2.1	5.3	10.0	15.9
2000.....	1.7	*	*	*	*	*	0.2	0.8	2.2	5.4	10.5	16.3
1999.....	1.7	*	*	*	*	*	0.1	0.7	2.4	5.6	10.2	16.6

Black, both sexes

2004.....	0.8	*	*	*	*	*	*	0.4	2.2	5.5	8.7	11.3
2003.....	0.7	*	*	*	*	*	*	0.8	1.5	3.8	9.1	12.3
2002.....	0.9	*	*	*	*	*	*	0.9	2.3	5.2	9.7	14.3
2001.....	0.9	*	*	*	*	*	*	0.7	2.1	5.8	8.8	15.7
2000.....	0.8	*	*	*	*	*	*	0.8	1.9	5.8	7.9	11.6
1999.....	0.8	*	*	*	*	*	*	0.7	2.2	5.2	9.9	12.1

Black, male

2004.....	0.5	*	*	*	*	*	*	*	1.6	4.4	9.2	*
2003.....	0.5	*	*	*	*	*	*	*	*	3.2	7.6	*
2002.....	0.6	*	*	*	*	*	*	*	1.9	4.4	8.6	*
2001.....	0.6	*	*	*	*	*	*	*	*	4.8	9.1	*
2000.....	0.5	*	*	*	*	*	*	*	*	4.8	7.0	*
1999.....	0.6	*	*	*	*	*	*	*	*	5.3	8.9	*

Black, female

2004.....	1.1	*	*	*	*	*	*	*	2.7	6.2	8.5	14.2
2003.....	1.0	*	*	*	*	*	*	1.0	1.6	4.2	9.9	14.2
2002.....	1.2	*	*	*	*	*	*	1.2	2.7	5.8	10.4	16.4
2001.....	1.1	*	*	*	*	*	*	1.0	2.6	6.4	8.6	16.0
2000.....	1.1	*	*	*	*	*	*	1.3	2.4	6.5	8.4	12.0
1999.....	1.1	*	*	*	*	*	*	*	2.8	5.2	10.5	14.9

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of pancreas (C25)												
All races 3/, both sexes												
2004.....	10.8	*	*	*	*	0.2	1.2	6.3	19.8	44.3	76.0	97.7
2003.....	10.6	*	*	*	*	0.2	1.2	6.2	19.1	44.2	75.4	95.2
2002.....	10.5	*	*	*	*	0.1	1.3	6.1	19.2	44.8	75.0	94.5
2001.....	10.5	*	*	*	*	0.1	1.3	6.1	19.2	44.5	74.1	100.1
2000.....	10.4	*	*	*	*	0.1	1.2	6.3	19.4	44.1	76.5	96.4
1999.....	10.4	*	*	*	*	0.1	1.3	6.4	19.8	45.0	73.1	99.7
All races 3/, male												
2004.....	10.9	*	*	*	*	0.2	1.5	8.1	24.1	52.6	84.6	102.0
2003.....	10.5	*	*	*	*	0.2	1.5	7.7	23.1	50.9	85.4	95.6
2002.....	10.5	*	*	*	*	0.1	1.6	7.8	23.6	51.4	84.0	100.9
2001.....	10.3	*	*	*	*	0.1	1.4	7.6	23.6	50.8	84.0	105.1
2000.....	10.3	*	*	*	*	0.2	1.4	8.0	23.6	50.4	85.9	102.9
1999.....	10.4	*	*	*	*	0.2	1.6	8.3	23.9	52.4	81.8	107.5
All races 3/, female												
2004.....	10.7	*	*	*	*	0.1	0.9	4.7	15.8	37.3	70.1	95.8
2003.....	10.6	*	*	*	*	0.1	1.0	4.8	15.3	38.6	68.8	95.0
2002.....	10.5	*	*	*	*	*	1.0	4.4	15.1	39.3	69.1	91.7
2001.....	10.6	*	*	*	*	0.1	1.1	4.7	15.2	39.4	67.6	98.1
2000.....	10.5	*	*	*	*	0.1	1.0	4.6	15.5	38.8	70.4	93.8
1999.....	10.5	*	*	*	*	0.1	1.0	4.7	16.0	39.0	67.4	96.6
White, both sexes												
2004.....	11.4	*	*	*	*	0.1	1.1	6.1	19.1	43.5	75.6	97.1
2003.....	11.2	*	*	*	*	0.2	1.2	5.9	18.4	43.7	74.8	94.4
2002.....	11.1	*	*	*	*	0.1	1.3	5.8	18.7	43.7	74.3	94.3
2001.....	11.1	*	*	*	*	0.1	1.3	5.9	19.0	43.8	73.3	99.0
2000.....	11.0	*	*	*	*	0.1	1.2	6.1	18.9	43.1	75.3	94.7
1999.....	11.0	*	*	*	*	0.1	1.3	6.2	19.1	44.2	71.7	98.6
White, male												
2004.....	11.5	*	*	*	*	0.2	1.4	7.7	23.4	51.6	85.1	102.9
2003.....	11.2	*	*	*	*	0.2	1.4	7.2	22.5	50.8	85.4	96.0
2002.....	11.1	*	*	*	*	*	1.6	7.5	23.0	50.7	84.2	101.6
2001.....	11.0	*	*	*	*	0.2	1.4	7.4	23.5	50.0	83.8	106.5
2000.....	10.9	*	*	*	*	0.2	1.4	7.8	23.0	49.8	85.2	102.4
1999.....	10.9	*	*	*	*	0.2	1.6	7.9	22.9	51.6	80.8	107.4

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of pancreas (C25)												
White, female												
2004.....	11.3	*	*	*	*	*	0.9	4.4	15.1	36.7	69.1	94.5
2003.....	11.2	*	*	*	*	0.1	1.0	4.5	14.6	37.6	67.7	93.8
2002.....	11.0	*	*	*	*	*	1.0	4.2	14.5	37.8	67.7	91.1
2001.....	11.2	*	*	*	*	*	1.1	4.5	14.7	38.5	66.3	95.9
2000.....	11.1	*	*	*	*	*	0.9	4.4	15.0	37.5	68.8	91.5
1999.....	11.0	*	*	*	*	*	0.9	4.6	15.5	37.9	65.8	95.0
Black, both sexes												
2004.....	9.5	*	*	*	*	*	1.6	9.3	28.4	57.9	88.0	113.6
2003.....	9.5	*	*	*	*	*	1.8	10.0	27.3	56.3	93.1	112.9
2002.....	9.3	*	*	*	*	*	1.8	8.7	27.6	60.7	89.9	106.6
2001.....	9.1	*	*	*	*	*	1.6	9.0	25.5	58.4	92.7	121.3
2000.....	9.3	*	*	*	*	*	1.7	9.0	27.5	59.1	97.8	119.8
1999.....	9.4	*	*	*	*	*	2.0	9.6	28.2	59.5	94.0	118.6
Black, male												
2004.....	9.4	*	*	*	*	*	2.0	11.4	36.1	72.3	89.6	100.3
2003.....	9.2	*	*	*	*	*	2.4	13.3	33.2	63.2	97.9	104.3
2002.....	9.0	*	*	*	*	*	2.1	11.4	34.1	68.4	96.1	100.8
2001.....	8.6	*	*	*	*	*	1.7	11.1	30.3	68.1	100.3	99.8
2000.....	9.0	*	*	*	*	*	2.0	11.8	34.2	65.8	106.9	109.6
1999.....	9.4	*	*	*	*	*	2.4	13.1	36.5	67.9	106.7	114.8
Black, female												
2004.....	9.6	*	*	*	*	*	1.2	7.5	22.2	47.7	87.0	118.9
2003.....	9.7	*	*	*	*	*	1.3	7.2	22.5	51.5	90.4	116.3
2002.....	9.5	*	*	*	*	*	1.4	6.4	22.5	55.3	86.4	108.8
2001.....	9.6	*	*	*	*	*	1.4	7.1	21.6	51.6	88.4	129.2
2000.....	9.7	*	*	*	*	*	1.4	6.6	22.1	54.4	92.7	123.6
1999.....	9.4	*	*	*	*	*	1.6	6.6	21.6	53.6	86.9	120.0

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Of other and ill-defined digestive organs (C26)

All races 3/, both sexes

2004.....	0.4	*	*	*	*	*	0.1	0.2	0.5	1.2	2.6	5.1
2003.....	0.5	*	*	*	*	*	0.0	0.2	0.6	1.7	3.4	7.0
2002.....	0.5	*	*	*	*	*	0.1	0.2	0.6	1.7	3.7	6.9
2001.....	0.5	*	*	*	*	*	0.1	0.2	0.7	1.7	3.6	7.8
2000.....	0.5	*	*	*	*	*	0.1	0.2	0.7	1.7	3.5	7.8
1999.....	0.6	*	*	*	*	*	0.1	0.3	0.8	1.9	4.1	9.1

All races 3/, male

2004.....	0.3	*	*	*	*	*	*	0.2	0.6	1.4	3.1	5.0
2003.....	0.5	*	*	*	*	*	*	0.2	0.8	2.1	4.1	7.3
2002.....	0.5	*	*	*	*	*	*	0.2	0.7	2.3	4.7	6.4
2001.....	0.5	*	*	*	*	*	*	0.3	1.0	2.0	4.2	8.9
2000.....	0.5	*	*	*	*	*	*	0.2	1.0	2.3	4.1	7.5
1999.....	0.6	*	*	*	*	*	0.1	0.4	1.0	2.2	5.2	10.4

All races 3/, female

2004.....	0.4	*	*	*	*	*	*	0.2	0.4	1.0	2.3	5.2
2003.....	0.5	*	*	*	*	*	*	0.2	0.4	1.3	2.9	6.8
2002.....	0.5	*	*	*	*	*	*	0.1	0.5	1.1	3.0	7.2
2001.....	0.5	*	*	*	*	*	*	0.1	0.5	1.5	3.2	7.3
2000.....	0.5	*	*	*	*	*	*	0.2	0.6	1.2	3.1	8.0
1999.....	0.6	*	*	*	*	*	*	0.2	0.6	1.7	3.4	8.5

White, both sexes

2004.....	0.4	*	*	*	*	*	*	0.2	0.5	1.2	2.7	5.3
2003.....	0.5	*	*	*	*	*	*	0.2	0.6	1.7	3.3	7.0
2002.....	0.5	*	*	*	*	*	*	0.2	0.6	1.6	3.7	7.0
2001.....	0.5	*	*	*	*	*	0.1	0.2	0.6	1.7	3.5	7.7
2000.....	0.5	*	*	*	*	*	0.1	0.2	0.7	1.6	3.4	7.9
1999.....	0.6	*	*	*	*	*	0.1	0.2	0.7	2.0	4.2	9.0

White, male

2004.....	0.4	*	*	*	*	*	*	0.2	0.6	1.5	3.1	5.3
2003.....	0.5	*	*	*	*	*	*	0.2	0.7	2.1	4.0	7.0
2002.....	0.5	*	*	*	*	*	*	0.2	0.7	2.2	4.7	6.2
2001.....	0.5	*	*	*	*	*	*	0.3	0.9	2.0	4.2	8.8
2000.....	0.5	*	*	*	*	*	*	0.2	1.0	2.1	4.2	7.7
1999.....	0.6	*	*	*	*	*	0.1	0.3	0.9	2.2	5.2	10.3

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms of respiratory and intrathoracic organs (C30-C39)

All races 3/, both sexes

2004.....	55.4	*	*	*	0.1	0.4	5.8	31.3	110.7	272.3	382.0	306.3
2003.....	56.0	*	*	*	0.1	0.4	5.9	31.6	115.0	276.6	386.7	307.9
2002.....	56.3	*	*	*	0.1	0.5	6.3	31.7	119.3	282.1	386.9	305.1
2001.....	56.5	*	*	*	0.1	0.5	6.6	32.2	121.9	287.3	380.6	312.2
2000.....	57.0	*	*	*	0.1	0.6	6.4	33.0	126.7	291.9	380.7	310.8
1999.....	56.2	*	*	*	0.1	0.5	6.4	32.7	127.9	289.4	371.2	306.6

All races 3/, male

2004.....	64.4	*	*	*	0.1	0.5	6.2	37.9	137.8	349.2	532.5	485.7
2003.....	65.4	*	*	*	0.1	0.5	6.5	38.6	143.3	358.8	541.5	493.3
2002.....	66.1	*	*	*	0.2	0.6	6.6	38.8	150.7	367.5	544.9	498.8
2001.....	67.2	*	*	*	0.1	0.4	7.2	39.7	154.0	382.6	546.1	534.1
2000.....	68.1	*	*	*	0.1	0.6	7.3	40.7	161.1	390.8	550.4	541.7
1999.....	68.0	*	*	*	*	0.6	7.1	40.6	165.3	390.0	544.0	550.6

All races 3/, female

2004.....	46.7	*	*	*	*	0.3	5.4	24.9	85.5	207.8	280.7	225.6
2003.....	46.9	*	*	*	*	0.4	5.3	24.9	88.8	207.9	283.2	225.9
2002.....	46.8	*	*	*	*	0.4	6.0	24.9	90.3	211.1	282.0	221.1
2001.....	46.1	*	*	*	*	0.5	5.9	25.0	92.3	208.3	271.8	219.3
2000.....	46.2	*	*	*	*	0.5	5.5	25.6	94.9	210.4	270.0	216.8
1999.....	45.0	*	*	*	*	0.5	5.7	25.1	93.4	207.0	258.9	208.3

White, both sexes

2004.....	59.7	*	*	*	0.1	0.4	5.6	29.9	109.8	276.8	386.7	310.7
2003.....	60.3	*	*	*	0.1	0.5	5.7	30.1	114.6	281.2	391.1	311.2
2002.....	60.7	*	*	*	0.1	0.5	6.1	30.4	118.1	286.8	392.0	309.0
2001.....	60.8	*	*	*	0.1	0.5	6.3	30.8	121.7	290.5	383.6	314.8
2000.....	61.2	*	*	*	0.1	0.6	5.9	31.4	126.2	294.2	384.3	313.4
1999.....	60.2	*	*	*	*	0.5	6.0	31.0	126.9	291.1	372.9	308.1

White, male

2004.....	68.3	*	*	*	*	0.5	5.8	35.2	133.1	348.9	532.1	490.5
2003.....	69.4	*	*	*	*	0.5	6.2	35.9	139.1	358.3	540.0	496.2
2002.....	70.1	*	*	*	0.1	0.5	6.2	36.1	145.3	367.4	544.1	503.6
2001.....	71.1	*	*	*	*	0.5	6.8	36.8	150.1	379.8	542.1	535.5
2000.....	72.0	*	*	*	*	0.6	6.5	37.5	156.8	386.9	547.6	543.3
1999.....	71.6	*	*	*	*	0.6	6.5	37.2	160.4	385.5	539.7	551.9

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms of respiratory and intrathoracic organs (C30-C39)

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
White, female												
2004.....	51.2	*	*	*	*	0.3	5.3	24.7	87.7	215.1	287.5	229.6
2003.....	51.5	*	*	*	*	0.4	5.3	24.3	91.4	215.4	290.4	229.3
2002.....	51.5	*	*	*	*	0.4	6.0	24.7	92.5	218.5	289.9	224.5
2001.....	50.7	*	*	*	*	0.5	5.9	24.9	94.9	215.2	278.3	222.3
2000.....	50.7	*	*	*	*	0.5	5.3	25.3	97.5	216.6	276.9	220.1
1999.....	49.1	*	*	*	*	0.5	5.5	24.8	95.5	212.4	263.6	210.1
Black, both sexes												
2004.....	44.6	*	*	*	*	0.5	8.1	48.1	148.9	291.4	397.6	287.6
2003.....	45.0	*	*	*	*	0.5	8.1	50.2	152.1	301.4	397.3	293.2
2002.....	45.3	*	*	*	*	0.6	9.4	49.9	160.6	307.1	396.1	278.0
2001.....	45.2	*	*	*	*	*	9.6	49.7	155.8	319.7	399.1	290.2
2000.....	46.2	*	*	*	*	0.7	10.9	53.8	162.6	329.8	391.0	295.9
1999.....	46.2	*	*	*	*	0.5	10.1	54.0	167.7	326.1	399.2	300.4
Black, male												
2004.....	57.5	*	*	*	*	*	9.3	67.2	220.4	432.8	638.4	508.6
2003.....	58.4	*	*	*	*	*	9.9	68.5	224.5	453.3	654.7	530.7
2002.....	59.4	*	*	*	*	0.8	11.6	69.6	239.1	466.0	657.5	498.5
2001.....	60.6	*	*	*	*	*	11.8	71.7	231.0	501.1	689.2	571.9
2000.....	62.0	*	*	*	*	*	14.4	76.4	242.2	515.0	672.8	584.7
1999.....	62.4	*	*	*	*	*	12.6	78.5	254.9	507.7	685.3	599.9
Black, female												
2004.....	32.9	*	*	*	*	*	7.1	31.7	91.3	190.5	261.5	200.0
2003.....	32.8	*	*	*	*	*	6.6	34.5	93.9	193.4	251.8	200.9
2002.....	32.4	*	*	*	*	*	7.4	33.0	97.4	194.2	248.2	193.5
2001.....	31.2	*	*	*	*	*	7.7	30.8	95.3	191.7	236.1	185.7
2000.....	31.8	*	*	*	*	*	7.8	34.3	98.4	199.5	232.3	188.4
1999.....	31.6	*	*	*	*	*	7.9	33.0	97.8	198.0	239.1	188.4

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Of nasal cavity, middle ear and accessory sinuses (C30-C31)

All races 3/, both sexes

2004.....	0.2	*	*	*	*	*	0.1	0.1	0.3	0.6	0.9	1.4
2003.....	0.2	*	*	*	*	*	0.1	0.1	0.3	0.5	0.9	1.2
2002.....	0.2	*	*	*	*	*	0.1	0.2	0.3	0.6	0.8	1.2
2001.....	0.2	*	*	*	*	*	0.1	0.2	0.3	0.5	1.1	1.7
2000.....	0.1	*	*	*	*	*	0.1	0.1	0.3	0.5	0.8	1.2
1999.....	0.2	*	*	*	*	*	0.1	0.2	0.3	0.6	0.9	1.5

All races 3/, male

2004.....	0.2	*	*	*	*	*	*	0.2	0.4	0.7	1.2	1.8
2003.....	0.2	*	*	*	*	*	*	0.2	0.4	0.8	1.4	*
2002.....	0.2	*	*	*	*	*	*	0.3	0.4	0.8	1.0	1.7
2001.....	0.2	*	*	*	*	*	*	0.2	0.5	0.6	1.3	1.8
2000.....	0.2	*	*	*	*	*	0.1	0.2	0.5	0.8	1.0	*
1999.....	0.2	*	*	*	*	*	0.1	0.3	0.4	0.7	1.2	1.8

All races 3/, female

2004.....	0.1	*	*	*	*	*	*	0.1	0.2	0.4	0.7	1.2
2003.....	0.1	*	*	*	*	*	*	*	0.2	0.3	0.6	1.1
2002.....	0.1	*	*	*	*	*	*	*	0.2	0.4	0.7	1.0
2001.....	0.2	*	*	*	*	*	*	*	0.2	0.4	0.9	1.6
2000.....	0.1	*	*	*	*	*	*	*	0.2	0.4	0.6	1.1
1999.....	0.1	*	*	*	*	*	*	*	0.2	0.5	0.7	1.4

White, both sexes

2004.....	0.2	*	*	*	*	*	0.1	0.1	0.3	0.6	0.9	1.4
2003.....	0.2	*	*	*	*	*	0.1	0.2	0.3	0.5	0.9	1.2
2002.....	0.2	*	*	*	*	*	*	0.2	0.3	0.6	0.9	1.3
2001.....	0.2	*	*	*	*	*	0.1	0.2	0.3	0.5	1.1	1.7
2000.....	0.2	*	*	*	*	*	0.1	0.1	0.3	0.5	0.8	1.3
1999.....	0.2	*	*	*	*	*	0.1	0.2	0.3	0.6	0.9	1.5

White, male

2004.....	0.2	*	*	*	*	*	*	0.1	0.4	0.7	1.2	1.8
2003.....	0.2	*	*	*	*	*	*	0.2	0.4	0.8	1.3	*
2002.....	0.2	*	*	*	*	*	*	0.3	0.5	0.8	1.1	1.7
2001.....	0.2	*	*	*	*	*	*	0.2	0.4	0.6	1.3	1.8
2000.....	0.2	*	*	*	*	*	*	0.2	0.4	0.8	1.1	*
1999.....	0.2	*	*	*	*	*	*	0.2	0.4	0.7	1.2	1.9

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of larynx (C32)												
All races 3/, both sexes												
2004.....	1.2	*	*	*	*	*	0.1	1.1	3.2	5.7	6.6	6.5
2003.....	1.3	*	*	*	*	*	0.2	1.0	3.4	6.2	6.9	6.5
2002.....	1.3	*	*	*	*	*	0.2	1.1	3.4	5.9	7.6	5.5
2001.....	1.3	*	*	*	*	*	0.2	1.1	3.6	6.4	7.2	6.7
2000.....	1.4	*	*	*	*	*	0.1	1.1	3.6	6.6	8.1	6.5
1999.....	1.4	*	*	*	*	*	0.2	1.1	3.8	6.7	7.3	7.0
All races 3/, male												
2004.....	2.0	*	*	*	*	*	0.2	1.8	5.4	10.1	13.0	15.6
2003.....	2.1	*	*	*	*	*	0.3	1.8	5.8	10.8	13.7	15.3
2002.....	2.1	*	*	*	*	*	0.3	1.8	5.8	10.0	14.9	13.2
2001.....	2.2	*	*	*	*	*	0.3	1.8	6.1	11.1	14.2	18.3
2000.....	2.2	*	*	*	*	*	0.2	1.8	6.2	11.4	15.7	17.4
1999.....	2.2	*	*	*	*	*	0.3	1.9	6.4	11.4	14.1	17.8
All races 3/, female												
2004.....	0.5	*	*	*	*	*	*	0.4	1.2	2.0	2.3	2.4
2003.....	0.5	*	*	*	*	*	*	0.3	1.2	2.4	2.4	2.7
2002.....	0.5	*	*	*	*	*	*	0.4	1.2	2.5	2.8	2.2
2001.....	0.5	*	*	*	*	*	0.1	0.4	1.3	2.5	2.6	1.9
2000.....	0.6	*	*	*	*	*	*	0.4	1.2	2.7	3.1	2.1
1999.....	0.6	*	*	*	*	*	*	0.3	1.4	2.9	2.8	2.6
White, both sexes												
2004.....	1.3	*	*	*	*	*	0.1	0.9	3.0	5.4	6.4	6.3
2003.....	1.3	*	*	*	*	*	0.2	0.9	3.1	5.8	6.6	6.2
2002.....	1.3	*	*	*	*	*	0.1	0.9	3.1	5.7	7.3	5.6
2001.....	1.3	*	*	*	*	*	0.2	0.9	3.2	6.0	6.9	6.6
2000.....	1.3	*	*	*	*	*	0.1	0.9	3.1	6.1	7.7	6.6
1999.....	1.4	*	*	*	*	*	0.1	0.9	3.4	6.4	7.2	6.7
White, male												
2004.....	2.0	*	*	*	*	*	0.2	1.5	4.8	9.5	12.4	15.0
2003.....	2.1	*	*	*	*	*	0.3	1.5	5.2	9.8	12.9	14.5
2002.....	2.0	*	*	*	*	*	0.2	1.5	5.1	9.5	14.1	13.3
2001.....	2.1	*	*	*	*	*	0.3	1.4	5.4	10.3	13.3	18.3
2000.....	2.1	*	*	*	*	*	0.2	1.5	5.3	10.4	15.0	17.6
1999.....	2.1	*	*	*	*	*	0.2	1.6	5.7	10.7	13.8	16.8

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of larynx (C32)												
White, female												
2004.....	0.5	*	*	*	*	*	*	0.4	1.2	1.9	2.3	2.5
2003.....	0.5	*	*	*	*	*	*	0.3	1.1	2.4	2.4	2.5
2002.....	0.6	*	*	*	*	*	*	0.3	1.2	2.5	2.6	2.3
2001.....	0.5	*	*	*	*	*	*	0.3	1.2	2.3	2.7	1.7
2000.....	0.6	*	*	*	*	*	*	0.4	1.1	2.6	2.9	2.1
1999.....	0.6	*	*	*	*	*	*	0.3	1.2	2.8	2.9	2.7
Black, both sexes												
2004.....	1.7	*	*	*	*	*	*	2.3	6.6	10.3	11.2	8.2
2003.....	1.8	*	*	*	*	*	*	2.5	7.5	11.5	11.7	11.1
2002.....	1.8	*	*	*	*	*	0.5	2.6	7.5	10.3	13.7	*
2001.....	1.8	*	*	*	*	*	0.4	2.8	7.9	11.4	11.7	8.3
2000.....	2.0	*	*	*	*	*	*	2.9	9.0	12.8	14.1	6.9
1999.....	1.8	*	*	*	*	*	0.6	2.9	8.6	11.6	10.0	8.3
Black, male												
2004.....	2.9	*	*	*	*	*	*	4.2	12.6	20.4	24.8	23.8
2003.....	3.1	*	*	*	*	*	*	4.5	13.6	23.8	27.9	27.1
2002.....	3.0	*	*	*	*	*	0.7	4.5	14.6	19.3	29.3	*
2001.....	3.1	*	*	*	*	*	*	5.1	14.2	21.3	27.7	*
2000.....	3.3	*	*	*	*	*	*	5.1	17.2	24.3	28.5	*
1999.....	3.1	*	*	*	*	*	0.9	5.2	15.9	21.8	22.8	24.6
Black, female												
2004.....	0.5	*	*	*	*	*	*	*	1.7	3.2	3.5	*
2003.....	0.6	*	*	*	*	*	*	0.8	2.5	2.8	*	*
2002.....	0.7	*	*	*	*	*	*	1.1	1.8	4.0	4.9	*
2001.....	0.7	*	*	*	*	*	*	*	2.8	4.5	*	*
2000.....	0.7	*	*	*	*	*	*	1.0	2.4	4.6	6.0	*
1999.....	0.7	*	*	*	*	*	*	0.9	2.7	4.4	*	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of trachea, bronchus and lung (C33-C34)												
All races 3/, both sexes												
2004.....	53.8	*	*	*	0.1	0.4	5.5	29.9	106.9	265.5	373.6	297.5
2003.....	54.4	*	*	*	0.1	0.4	5.6	30.3	111.0	269.3	377.8	298.9
2002.....	54.7	*	*	*	0.1	0.4	6.0	30.3	115.3	275.0	377.6	297.2
2001.....	54.8	*	*	*	0.1	0.4	6.2	30.7	117.7	279.7	371.4	302.7
2000.....	55.3	*	*	*	0.1	0.5	6.1	31.6	122.4	284.2	370.8	302.1
1999.....	54.5	*	*	*	0.1	0.4	6.0	31.3	123.4	281.5	362.1	297.0
All races 3/, male												
2004.....	62.0	*	*	*	*	0.4	5.8	35.8	131.7	337.6	517.3	466.8
2003.....	62.9	*	*	*	*	0.4	6.1	36.5	136.7	346.6	525.1	475.1
2002.....	63.7	*	*	*	*	0.4	6.2	36.6	144.0	355.9	527.9	482.2
2001.....	64.7	*	*	*	*	0.4	6.7	37.4	147.1	370.0	529.5	512.4
2000.....	65.5	*	*	*	*	0.5	6.9	38.5	154.0	377.9	532.2	521.2
1999.....	65.4	*	*	*	*	0.5	6.6	38.2	158.0	377.2	527.6	529.2
All races 3/, female												
2004.....	45.9	*	*	*	*	0.3	5.2	24.2	83.9	205.0	277.0	221.3
2003.....	46.1	*	*	*	*	0.4	5.1	24.4	87.1	204.8	279.4	221.0
2002.....	46.0	*	*	*	*	0.4	5.8	24.3	88.8	207.7	277.8	217.0
2001.....	45.3	*	*	*	*	0.4	5.7	24.3	90.5	204.9	267.5	215.0
2000.....	45.4	*	*	*	*	0.5	5.3	25.0	93.3	206.9	265.6	212.8
1999.....	44.1	*	*	*	*	0.4	5.5	24.6	91.5	203.1	254.5	203.5
White, both sexes												
2004.....	58.1	*	*	*	*	0.3	5.3	28.7	106.3	270.4	378.6	302.0
2003.....	58.7	*	*	*	0.1	0.4	5.4	28.9	110.8	274.2	382.6	302.4
2002.....	59.1	*	*	*	*	0.4	5.8	29.1	114.5	280.0	382.9	300.9
2001.....	59.1	*	*	*	*	0.4	6.0	29.6	117.8	283.4	374.7	305.4
2000.....	59.5	*	*	*	*	0.5	5.7	30.2	122.5	287.0	374.7	304.5
1999.....	58.5	*	*	*	*	0.5	5.8	29.7	122.9	283.5	363.8	298.8
White, male												
2004.....	65.9	*	*	*	*	0.4	5.5	33.3	127.6	338.1	517.5	472.2
2003.....	66.9	*	*	*	*	0.4	5.7	34.1	133.1	347.1	524.6	478.5
2002.....	67.7	*	*	*	*	0.4	5.8	34.2	139.3	356.4	527.8	486.6
2001.....	68.6	*	*	*	*	0.4	6.3	34.9	143.8	368.1	526.4	513.9
2000.....	69.4	*	*	*	*	0.5	6.1	35.7	150.8	374.9	529.9	522.4
1999.....	69.1	*	*	*	*	0.5	6.1	35.2	153.9	373.4	523.5	531.5

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of trachea, bronchus and lung (C33-C34)												
White, female												
2004.....	50.4	*	*	*	*	0.3	5.2	24.1	86.1	212.5	283.8	225.3
2003.....	50.7	*	*	*	*	0.4	5.1	23.8	89.8	212.2	286.6	224.4
2002.....	50.7	*	*	*	*	0.4	5.8	24.2	91.0	215.1	285.8	220.2
2001.....	49.8	*	*	*	*	0.4	5.7	24.3	93.3	211.9	274.0	218.1
2000.....	49.9	*	*	*	*	0.5	5.2	24.8	96.1	213.2	272.7	215.9
1999.....	48.2	*	*	*	*	0.4	5.4	24.4	93.8	208.6	259.1	205.3
Black, both sexes												
2004.....	42.7	*	*	*	*	0.5	7.7	45.4	141.6	279.6	384.4	277.5
2003.....	42.9	*	*	*	*	*	7.7	47.4	144.0	288.6	383.1	280.9
2002.....	43.2	*	*	*	*	0.4	8.7	46.9	152.4	295.3	381.3	272.3
2001.....	43.1	*	*	*	*	*	8.9	46.5	147.0	306.5	385.4	280.7
2000.....	43.9	*	*	*	*	0.6	10.1	50.4	152.5	315.8	375.0	288.1
1999.....	44.1	*	*	*	*	0.4	9.3	50.6	158.1	313.1	387.9	290.5
Black, male												
2004.....	54.3	*	*	*	*	*	8.8	62.5	206.9	410.4	611.1	481.8
2003.....	54.9	*	*	*	*	*	9.3	63.6	210.0	427.1	622.2	502.5
2002.....	56.0	*	*	*	*	*	10.6	64.6	223.6	444.6	626.2	484.6
2001.....	57.2	*	*	*	*	*	11.0	66.2	215.2	476.4	659.8	549.2
2000.....	58.3	*	*	*	*	*	13.4	70.7	223.5	488.8	642.5	562.8
1999.....	58.9	*	*	*	*	*	11.3	72.6	237.6	484.4	661.0	574.2
Black, female												
2004.....	32.1	*	*	*	*	*	6.8	30.7	89.0	186.2	256.3	196.5
2003.....	32.0	*	*	*	*	*	6.2	33.4	91.0	190.0	248.0	194.8
2002.....	31.6	*	*	*	*	*	7.0	31.7	95.3	189.3	242.6	191.0
2001.....	30.2	*	*	*	*	*	7.1	29.5	92.0	186.7	231.3	181.0
2000.....	30.8	*	*	*	*	*	7.3	32.9	95.3	194.1	224.3	185.9
1999.....	30.7	*	*	*	*	*	7.6	31.9	94.4	192.3	234.9	184.4

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Of thymus, heart, mediastinum and pleura (C37-C38)

All races 3/, both sexes

2004.....	0.1	*	*	*	*	*	0.1	0.2	0.3	0.5	0.8	0.9
2003.....	0.2	*	*	*	*	0.1	0.1	0.1	0.3	0.5	0.9	1.3
2002.....	0.2	*	*	*	*	0.1	0.1	0.2	0.2	0.6	0.8	1.1
2001.....	0.2	*	*	*	*	*	0.1	0.2	0.3	0.6	0.8	1.0
2000.....	0.2	*	*	*	*	*	0.1	0.1	0.3	0.5	1.0	0.9
1999.....	0.2	*	*	*	*	*	0.1	0.1	0.3	0.5	0.9	1.0

All races 3/, male

2004.....	0.2	*	*	*	*	*	*	0.2	0.3	0.7	1.0	1.5
2003.....	0.2	*	*	*	*	*	0.1	0.1	0.3	0.6	1.1	1.7
2002.....	0.2	*	*	*	*	*	0.1	0.2	0.4	0.6	0.9	1.7
2001.....	0.2	*	*	*	*	*	0.1	0.2	0.4	0.8	0.9	1.5
2000.....	0.2	*	*	*	*	*	0.1	0.2	0.3	0.6	1.4	*
1999.....	0.2	*	*	*	*	*	0.1	0.2	0.4	0.6	1.0	*

All races 3/, female

2004.....	0.1	*	*	*	*	*	*	0.1	0.2	0.4	0.6	0.7
2003.....	0.1	*	*	*	*	*	*	0.1	0.2	0.4	0.7	1.1
2002.....	0.1	*	*	*	*	*	*	0.1	*	0.5	0.7	0.9
2001.....	0.2	*	*	*	*	*	*	0.2	0.2	0.4	0.7	0.8
2000.....	0.1	*	*	*	*	*	*	0.1	0.2	0.4	0.7	0.7
1999.....	0.1	*	*	*	*	*	*	*	0.3	0.5	0.8	0.8

White, both sexes

2004.....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.5	0.8	1.0
2003.....	0.2	*	*	*	*	*	0.1	0.1	0.3	0.5	0.8	1.4
2002.....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.6	0.8	1.3
2001.....	0.2	*	*	*	*	*	0.1	0.2	0.3	0.6	0.8	1.0
2000.....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.5	1.0	0.9
1999.....	0.2	*	*	*	*	*	0.1	0.1	0.3	0.5	0.9	1.0

White, male

2004.....	0.2	*	*	*	*	*	*	0.2	0.3	0.7	1.1	1.6
2003.....	0.2	*	*	*	*	*	*	0.1	0.3	0.6	1.0	1.8
2002.....	0.2	*	*	*	*	*	*	0.2	0.3	0.7	1.0	1.9
2001.....	0.2	*	*	*	*	*	*	0.2	0.4	0.7	1.0	*
2000.....	0.2	*	*	*	*	*	0.1	0.2	0.3	0.7	1.4	*
1999.....	0.2	*	*	*	*	*	0.1	0.2	0.4	0.5	1.0	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of bone and articular cartilage (C40-C41)												
All races 3/, both sexes												
2004.....	0.4	*	*	0.2	0.5	0.2	0.2	0.3	0.6	1.0	1.8	2.7
2003.....	0.4	*	*	0.2	0.5	0.2	0.2	0.3	0.5	0.9	1.9	2.8
2002.....	0.4	*	*	0.2	0.4	0.2	0.2	0.3	0.6	0.8	1.6	2.9
2001.....	0.5	*	*	0.2	0.5	0.2	0.2	0.3	0.5	1.1	1.9	3.5
2000.....	0.4	*	*	0.2	0.4	0.2	0.2	0.3	0.6	1.0	1.9	3.4
1999.....	0.4	*	*	0.2	0.5	0.2	0.2	0.4	0.4	1.0	1.8	3.0
All races 3/, male												
2004.....	0.5	*	*	0.2	0.6	0.3	0.2	0.4	0.7	1.1	2.1	3.2
2003.....	0.5	*	*	0.2	0.6	0.3	0.3	0.4	0.6	1.1	2.4	3.5
2002.....	0.5	*	*	0.2	0.5	0.3	0.3	0.4	0.7	0.9	2.0	3.7
2001.....	0.5	*	*	0.2	0.7	0.2	0.3	0.4	0.6	1.4	2.5	3.6
2000.....	0.5	*	*	0.2	0.6	0.3	0.2	0.4	0.7	1.2	2.4	4.2
1999.....	0.5	*	*	0.2	0.6	0.3	0.2	0.4	0.6	1.2	2.0	4.5
All races 3/, female												
2004.....	0.4	*	*	0.2	0.3	0.2	0.2	0.2	0.5	1.0	1.6	2.5
2003.....	0.4	*	*	0.2	0.3	0.2	0.1	0.2	0.3	0.7	1.5	2.5
2002.....	0.4	*	*	0.2	0.3	0.2	0.1	0.2	0.5	0.8	1.3	2.5
2001.....	0.4	*	*	0.2	0.3	0.2	0.2	0.2	0.5	0.8	1.5	3.4
2000.....	0.4	*	*	0.2	0.3	0.1	0.1	0.2	0.4	0.8	1.6	3.1
1999.....	0.4	*	*	0.2	0.4	0.2	0.2	0.3	0.3	0.8	1.7	2.4
White, both sexes												
2004.....	0.5	*	*	0.2	0.5	0.2	0.2	0.3	0.6	1.1	1.8	2.8
2003.....	0.5	*	*	0.2	0.5	0.2	0.3	0.3	0.5	0.9	1.9	2.9
2002.....	0.4	*	*	0.2	0.5	0.3	0.2	0.3	0.6	0.8	1.6	3.0
2001.....	0.5	*	*	0.2	0.5	0.2	0.2	0.3	0.5	1.0	2.0	3.6
2000.....	0.5	*	*	0.2	0.5	0.2	0.2	0.3	0.5	1.0	1.9	3.5
1999.....	0.5	*	*	0.2	0.5	0.2	0.2	0.4	0.4	1.0	1.9	3.1
White, male												
2004.....	0.5	*	*	0.2	0.6	0.3	0.2	0.4	0.6	1.2	2.1	3.2
2003.....	0.5	*	*	0.2	0.6	0.3	0.3	0.4	0.6	1.1	2.5	3.5
2002.....	0.5	*	*	0.2	0.5	0.3	0.3	0.4	0.7	0.9	2.1	4.0
2001.....	0.5	*	*	0.2	0.7	0.3	0.3	0.3	0.6	1.4	2.6	3.6
2000.....	0.5	*	*	0.2	0.5	0.3	0.2	0.4	0.7	1.2	2.4	4.5
1999.....	0.5	*	*	0.2	0.6	0.3	0.2	0.4	0.5	1.3	2.0	4.8

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Melanoma and other malignant neoplasms of skin (C43-C44)												
All races 3/, both sexes												
2004.....	3.5	*	*	*	0.1	0.5	1.3	3.3	6.7	11.8	19.8	30.2
2003.....	3.5	*	*	*	0.1	0.5	1.4	3.4	6.8	11.3	19.7	30.8
2002.....	3.4	*	*	*	0.1	0.5	1.5	3.3	6.4	11.5	19.5	29.8
2001.....	3.5	*	*	*	0.1	0.5	1.5	3.6	6.8	12.1	18.7	31.2
2000.....	3.4	*	*	*	0.1	0.5	1.5	3.7	6.7	11.9	18.0	31.8
1999.....	3.4	*	*	*	0.1	0.5	1.5	3.6	6.7	11.8	18.8	29.2
All races 3/, male												
2004.....	4.7	*	*	*	*	0.6	1.6	4.4	9.8	17.9	32.4	53.9
2003.....	4.6	*	*	*	0.1	0.6	1.7	4.6	9.9	16.9	32.0	53.5
2002.....	4.5	*	*	*	0.1	0.6	1.8	4.6	9.3	17.4	31.1	50.2
2001.....	4.6	*	*	*	0.1	0.6	1.8	4.8	9.9	18.7	30.0	54.0
2000.....	4.5	*	*	*	0.1	0.6	1.7	4.9	9.5	17.8	29.6	55.4
1999.....	4.5	*	*	*	0.1	0.6	2.0	4.9	9.8	17.9	30.0	50.6
All races 3/, female												
2004.....	2.4	*	*	*	0.1	0.4	1.0	2.2	3.7	6.6	11.2	19.5
2003.....	2.4	*	*	*	*	0.4	1.1	2.3	4.0	6.7	11.5	20.7
2002.....	2.4	*	*	*	0.1	0.4	1.1	2.1	3.7	6.6	11.8	21.0
2001.....	2.5	*	*	*	*	0.4	1.1	2.4	4.0	6.7	11.3	21.7
2000.....	2.5	*	*	*	0.1	0.4	1.2	2.5	4.1	7.0	10.5	22.2
1999.....	2.4	*	*	*	*	0.4	1.1	2.4	3.9	6.7	11.5	20.6
White, both sexes												
2004.....	4.2	*	*	*	0.1	0.6	1.5	3.8	7.5	13.2	21.6	32.7
2003.....	4.2	*	*	*	0.1	0.6	1.7	3.9	7.7	12.7	21.5	33.3
2002.....	4.1	*	*	*	0.1	0.6	1.7	3.8	7.1	12.7	21.3	32.2
2001.....	4.2	*	*	*	0.1	0.6	1.8	4.2	7.6	13.3	20.3	33.7
2000.....	4.0	*	*	*	0.1	0.6	1.7	4.2	7.4	13.0	19.5	34.3
1999.....	4.0	*	*	*	0.1	0.6	1.8	4.1	7.5	13.0	20.3	31.5
White, male												
2004.....	5.6	*	*	*	*	0.7	1.8	5.0	10.9	20.1	35.3	58.9
2003.....	5.5	*	*	*	0.2	0.8	2.0	5.2	11.1	18.9	34.9	58.2
2002.....	5.3	*	*	*	*	0.8	2.1	5.2	10.3	19.0	34.0	54.5
2001.....	5.4	*	*	*	0.2	0.7	2.2	5.4	11.0	20.4	32.6	58.7
2000.....	5.2	*	*	*	0.1	0.7	2.0	5.6	10.5	19.4	31.9	60.6
1999.....	5.3	*	*	*	0.1	0.7	2.3	5.5	10.9	19.7	32.2	55.0

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant melanoma of skin (C43)												
All races 3/, both sexes												
2004.....	2.7	*	*	*	0.1	0.5	1.2	2.8	5.3	9.3	14.7	17.7
2003.....	2.7	*	*	*	0.1	0.5	1.3	2.9	5.5	8.7	14.6	17.6
2002.....	2.6	*	*	*	0.1	0.5	1.3	2.8	5.1	8.8	13.7	17.9
2001.....	2.6	*	*	*	0.1	0.5	1.4	3.1	5.3	9.2	13.3	17.8
2000.....	2.6	*	*	*	0.1	0.5	1.4	3.2	5.4	8.9	13.1	18.9
1999.....	2.6	*	*	*	0.1	0.5	1.4	3.1	5.3	9.0	13.4	16.4
All races 3/, male												
2004.....	3.5	*	*	*	*	0.6	1.4	3.7	7.6	13.7	23.7	31.4
2003.....	3.5	*	*	*	0.1	0.6	1.5	3.8	7.8	12.6	23.3	30.2
2002.....	3.3	*	*	*	0.1	0.6	1.6	3.8	7.2	13.0	21.2	29.2
2001.....	3.4	*	*	*	0.1	0.5	1.7	3.9	7.3	13.9	21.3	29.9
2000.....	3.3	*	*	*	0.1	0.5	1.6	4.2	7.4	12.9	20.8	31.8
1999.....	3.3	*	*	*	0.1	0.5	1.8	4.0	7.5	13.2	20.9	27.5
All races 3/, female												
2004.....	1.9	*	*	*	0.1	0.4	1.0	2.0	3.2	5.5	8.7	11.5
2003.....	1.9	*	*	*	*	0.4	1.0	2.1	3.5	5.4	8.7	12.1
2002.....	1.9	*	*	*	0.1	0.4	1.1	1.9	3.2	5.4	8.8	12.9
2001.....	1.9	*	*	*	*	0.4	1.1	2.3	3.3	5.3	8.1	12.8
2000.....	2.0	*	*	*	0.1	0.4	1.2	2.2	3.5	5.7	8.2	13.7
1999.....	1.9	*	*	*	*	0.4	1.0	2.2	3.3	5.5	8.5	11.9
White, both sexes												
2004.....	3.3	*	*	*	0.1	0.6	1.4	3.3	6.1	10.5	16.2	19.2
2003.....	3.2	*	*	*	0.1	0.6	1.5	3.4	6.3	9.8	16.0	19.1
2002.....	3.1	*	*	*	0.1	0.6	1.6	3.3	5.8	9.9	15.1	19.2
2001.....	3.2	*	*	*	0.1	0.6	1.7	3.6	6.1	10.3	14.6	19.3
2000.....	3.1	*	*	*	0.1	0.6	1.6	3.7	6.1	9.9	14.3	20.3
1999.....	3.1	*	*	*	0.1	0.6	1.7	3.6	6.1	10.0	14.5	17.6
White, male												
2004.....	4.3	*	*	*	*	0.7	1.7	4.3	8.7	15.6	26.0	34.2
2003.....	4.2	*	*	*	0.1	0.7	1.8	4.4	8.8	14.2	25.5	32.9
2002.....	4.0	*	*	*	*	0.7	1.9	4.4	8.1	14.4	23.3	31.7
2001.....	4.1	*	*	*	0.1	0.6	2.0	4.6	8.4	15.5	23.3	32.7
2000.....	4.0	*	*	*	0.1	0.7	1.9	4.8	8.5	14.3	22.5	34.7
1999.....	4.0	*	*	*	0.1	0.6	2.1	4.6	8.5	14.7	22.6	29.7

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other malignant neoplasms of skin (C44)												
All races 3/, both sexes												
2004.....	0.8	*	*	*	*	*	0.1	0.5	1.3	2.5	5.0	12.5
2003.....	0.8	*	*	*	*	*	0.1	0.5	1.3	2.7	5.1	13.1
2002.....	0.8	*	*	*	*	*	0.1	0.5	1.3	2.7	5.8	12.0
2001.....	0.9	*	*	*	*	*	0.1	0.5	1.5	2.9	5.4	13.4
2000.....	0.8	*	*	*	*	*	0.1	0.5	1.3	3.0	4.9	12.9
1999.....	0.8	*	*	*	*	*	0.1	0.6	1.4	2.8	5.4	12.8
All races 3/, male												
2004.....	1.1	*	*	*	*	*	0.1	0.7	2.3	4.2	8.7	22.5
2003.....	1.2	*	*	*	*	*	0.2	0.8	2.1	4.4	8.7	23.3
2002.....	1.2	*	*	*	*	*	0.2	0.8	2.2	4.4	9.9	21.0
2001.....	1.2	*	*	*	*	*	0.1	0.9	2.5	4.8	8.7	24.1
2000.....	1.1	*	*	*	*	*	0.2	0.8	2.1	4.9	8.8	23.6
1999.....	1.2	*	*	*	*	*	0.2	0.9	2.4	4.7	9.1	23.1
All races 3/, female												
2004.....	0.5	*	*	*	*	*	*	0.2	0.5	1.1	2.6	8.0
2003.....	0.5	*	*	*	*	*	*	0.2	0.5	1.2	2.8	8.6
2002.....	0.5	*	*	*	*	*	*	0.2	0.6	1.2	3.1	8.1
2001.....	0.5	*	*	*	*	*	*	0.2	0.6	1.3	3.2	8.9
2000.....	0.5	*	*	*	*	*	*	0.2	0.6	1.3	2.4	8.5
1999.....	0.5	*	*	*	*	*	*	0.2	0.5	1.3	3.0	8.6
White, both sexes												
2004.....	0.9	*	*	*	*	*	0.1	0.5	1.4	2.7	5.4	13.6
2003.....	0.9	*	*	*	*	*	0.1	0.5	1.4	2.9	5.5	14.2
2002.....	0.9	*	*	*	*	*	0.1	0.5	1.3	2.8	6.3	13.0
2001.....	1.0	*	*	*	*	*	0.1	0.5	1.6	3.0	5.8	14.3
2000.....	0.9	*	*	*	*	*	0.1	0.5	1.3	3.1	5.2	14.0
1999.....	0.9	*	*	*	*	*	0.1	0.5	1.4	3.0	5.7	13.9
White, male												
2004.....	1.3	*	*	*	*	*	0.1	0.7	2.3	4.5	9.3	24.7
2003.....	1.3	*	*	*	*	*	0.2	0.8	2.2	4.7	9.4	25.3
2002.....	1.3	*	*	*	*	*	0.2	0.8	2.2	4.6	10.7	22.7
2001.....	1.3	*	*	*	*	*	0.1	0.8	2.5	4.9	9.3	25.9
2000.....	1.3	*	*	*	*	*	0.1	0.7	2.1	5.2	9.3	25.9
1999.....	1.3	*	*	*	*	*	0.2	0.8	2.4	4.9	9.5	25.3

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of mesothelial and soft tissue (C45-C49)												
All races 3/, both sexes												
2004.....	2.4	*	0.2	0.2	0.3	0.4	0.8	1.6	4.2	8.7	16.1	15.7
2003.....	2.4	*	*	0.2	0.4	0.5	0.7	1.7	4.1	8.8	15.7	14.9
2002.....	2.4	*	0.1	0.2	0.3	0.4	0.8	1.6	4.3	9.1	14.9	15.7
2001.....	2.4	*	*	0.1	0.3	0.5	0.8	1.8	4.2	9.1	15.1	14.9
2000.....	2.4	*	0.1	0.1	0.4	0.5	0.8	1.8	4.4	9.6	15.1	15.7
1999.....	2.4	*	*	0.2	0.4	0.5	0.9	1.8	4.4	9.3	14.8	14.8
All races 3/, male												
2004.....	2.8	*	*	0.2	0.4	0.5	0.9	1.6	4.8	11.3	24.7	27.9
2003.....	2.8	*	*	0.2	0.4	0.5	0.8	1.8	4.6	11.6	24.3	25.1
2002.....	2.8	*	*	0.2	0.3	0.5	0.9	1.6	5.0	12.3	22.5	28.1
2001.....	2.8	*	*	0.1	0.4	0.6	0.8	2.0	4.8	12.5	22.7	25.9
2000.....	2.9	*	*	0.1	0.4	0.6	0.8	1.8	5.3	12.9	23.5	28.4
1999.....	2.8	*	*	0.2	0.4	0.6	1.1	1.9	5.5	12.8	22.6	24.6
All races 3/, female												
2004.....	2.0	*	*	0.2	0.3	0.3	0.7	1.6	3.6	6.5	10.3	10.2
2003.....	2.0	*	*	0.2	0.3	0.4	0.7	1.6	3.7	6.4	9.9	10.3
2002.....	1.9	*	*	0.2	0.3	0.3	0.7	1.5	3.7	6.4	9.8	10.4
2001.....	2.0	*	*	0.1	0.3	0.4	0.7	1.6	3.7	6.3	10.1	10.3
2000.....	2.0	*	*	0.1	0.3	0.5	0.7	1.7	3.6	6.9	9.6	10.5
1999.....	1.9	*	*	0.2	0.3	0.4	0.7	1.7	3.5	6.4	9.6	10.8
White, both sexes												
2004.....	2.7	*	*	0.2	0.3	0.4	0.8	1.6	4.4	9.3	17.0	16.5
2003.....	2.6	*	*	0.2	0.4	0.4	0.7	1.7	4.3	9.3	16.4	15.6
2002.....	2.6	*	*	0.2	0.3	0.4	0.8	1.6	4.4	9.5	15.6	16.5
2001.....	2.6	*	*	0.1	0.3	0.5	0.8	1.8	4.4	9.7	15.8	15.4
2000.....	2.7	*	*	0.1	0.3	0.5	0.7	1.8	4.6	10.0	15.9	16.2
1999.....	2.6	*	*	0.2	0.3	0.4	0.8	1.9	4.6	9.7	15.2	15.3
White, male												
2004.....	3.2	*	*	0.2	0.3	0.6	0.8	1.6	5.1	12.1	26.1	29.5
2003.....	3.1	*	*	0.2	0.4	0.5	0.8	1.8	4.7	12.3	25.8	26.1
2002.....	3.1	*	*	0.2	0.4	0.5	0.9	1.7	5.2	13.0	23.8	29.8
2001.....	3.1	*	*	0.1	0.4	0.6	0.8	2.0	5.0	13.4	23.9	27.0
2000.....	3.2	*	*	*	0.4	0.6	0.8	1.9	5.5	13.4	24.8	29.5
1999.....	3.1	*	*	0.2	0.4	0.5	1.0	2.0	5.7	13.5	23.6	26.2

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms of mesothelial and soft tissue (C45-C49)

White, female

2004.....	2.2	*	*	0.2	0.3	0.3	0.7	1.6	3.6	6.9	10.8	10.6
2003.....	2.2	*	*	0.2	0.3	0.3	0.7	1.6	3.8	6.7	10.1	10.9
2002.....	2.1	*	*	0.2	0.3	0.4	0.7	1.5	3.7	6.5	10.1	10.7
2001.....	2.1	*	*	0.1	0.3	0.4	0.7	1.5	3.8	6.5	10.5	10.5
2000.....	2.2	*	*	0.1	0.3	0.5	0.7	1.6	3.7	7.1	10.1	10.7
1999.....	2.1	*	*	0.1	0.3	0.4	0.6	1.7	3.5	6.6	9.7	10.9

Black, both sexes

2004.....	1.5	*	*	*	0.5	0.6	1.1	2.1	3.4	6.0	10.5	9.3
2003.....	1.6	*	*	*	0.3	0.8	0.9	2.3	3.9	6.3	10.1	9.9
2002.....	1.5	*	*	*	0.4	0.6	1.1	1.8	4.2	7.0	10.2	7.7
2001.....	1.4	*	*	*	0.4	0.6	0.8	2.3	3.7	5.9	8.5	11.1
2000.....	1.6	*	*	*	0.5	0.6	1.2	2.2	4.3	8.2	8.3	10.9
1999.....	1.6	*	*	*	0.5	0.7	1.3	1.9	4.3	6.6	11.5	10.5

Black, male

2004.....	1.5	*	*	*	*	*	1.1	2.1	3.5	7.1	15.0	*
2003.....	1.7	*	*	*	*	0.8	1.1	2.8	4.6	6.9	12.7	*
2002.....	1.6	*	*	*	*	0.8	1.3	2.1	4.3	8.0	12.9	*
2001.....	1.5	*	*	*	0.7	0.8	1.1	2.2	3.9	6.4	11.2	*
2000.....	1.6	*	*	*	*	*	1.4	1.6	5.0	10.1	12.4	*
1999.....	1.7	*	*	*	0.8	0.8	1.6	1.8	5.2	7.9	14.5	*

Black, female

2004.....	1.5	*	*	*	*	*	1.1	2.1	3.4	5.2	7.9	*
2003.....	1.4	*	*	*	*	0.8	0.8	1.8	3.4	5.9	8.6	*
2002.....	1.4	*	*	*	*	*	0.8	1.6	4.0	6.3	8.6	*
2001.....	1.4	*	*	*	*	*	0.7	2.3	3.6	5.5	6.9	8.8
2000.....	1.5	*	*	*	*	*	1.0	2.7	3.7	6.9	6.0	8.6
1999.....	1.5	*	*	*	*	*	1.0	1.9	3.6	5.6	9.8	10.1

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Mesothelioma (C45)												
All races 3/, both sexes												
2004.....	0.9	*	*	*	*	*	0.1	0.3	1.3	3.5	7.9	5.7
2003.....	0.9	*	*	*	*	*	0.1	0.3	1.3	3.7	7.5	5.7
2002.....	0.8	*	*	*	*	*	0.1	0.3	1.4	4.0	7.2	5.9
2001.....	0.8	*	*	*	*	*	0.1	0.4	1.3	3.9	7.1	5.5
2000.....	0.8	*	*	*	*	*	0.1	0.3	1.5	4.2	6.9	5.7
1999.....	0.8	*	*	*	*	*	0.1	0.4	1.6	4.2	6.8	4.8
All races 3/, male												
2004.....	1.4	*	*	*	*	*	0.1	0.4	2.2	6.2	16.3	14.4
2003.....	1.4	*	*	*	*	*	0.1	0.5	2.2	6.9	15.4	14.2
2002.....	1.4	*	*	*	*	*	0.1	0.4	2.4	7.6	14.7	15.9
2001.....	1.4	*	*	*	*	*	*	0.5	2.2	7.3	14.4	14.5
2000.....	1.4	*	*	*	*	*	*	0.5	2.5	7.8	14.2	15.5
1999.....	1.4	*	*	*	*	*	0.1	0.6	2.5	7.7	14.0	12.7
All races 3/, female												
2004.....	0.3	*	*	*	*	*	*	0.2	0.5	1.2	2.3	1.8
2003.....	0.3	*	*	*	*	*	*	0.2	0.5	1.1	2.2	1.9
2002.....	0.3	*	*	*	*	*	*	0.1	0.5	1.0	2.2	1.6
2001.....	0.3	*	*	*	*	*	*	0.2	0.5	1.1	2.3	1.7
2000.....	0.3	*	*	*	*	*	*	0.2	0.5	1.3	2.1	1.8
1999.....	0.3	*	*	*	*	*	*	0.2	0.6	1.3	2.1	1.6
White, both sexes												
2004.....	1.0	*	*	*	*	*	0.1	0.3	1.5	3.9	8.6	6.1
2003.....	1.0	*	*	*	*	*	0.1	0.4	1.4	4.1	8.1	6.0
2002.....	1.0	*	*	*	*	*	0.1	0.3	1.5	4.4	7.7	6.4
2001.....	1.0	*	*	*	*	*	0.1	0.4	1.5	4.3	7.7	5.9
2000.....	1.0	*	*	*	*	*	0.1	0.3	1.6	4.5	7.4	6.1
1999.....	1.0	*	*	*	*	*	0.1	0.4	1.7	4.6	7.2	5.0
White, male												
2004.....	1.6	*	*	*	*	*	*	0.4	2.4	6.8	17.3	15.6
2003.....	1.6	*	*	*	*	*	*	0.5	2.4	7.5	16.6	14.9
2002.....	1.6	*	*	*	*	*	0.1	0.4	2.6	8.2	15.6	17.3
2001.....	1.6	*	*	*	*	*	*	0.6	2.5	8.0	15.4	15.6
2000.....	1.6	*	*	*	*	*	*	0.5	2.7	8.2	15.2	16.6
1999.....	1.6	*	*	*	*	*	*	0.6	2.7	8.4	14.8	13.6

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of other mesothelial and soft tissue (C47-C49)												
All races 3/, both sexes												
2004.....	1.6	*	0.2	0.2	0.3	0.4	0.7	1.3	2.8	5.2	8.1	9.6
2003.....	1.5	*	*	0.2	0.4	0.4	0.6	1.3	2.8	5.0	8.1	8.8
2002.....	1.5	*	0.1	0.2	0.3	0.4	0.7	1.3	2.9	5.1	7.6	9.3
2001.....	1.5	*	*	0.1	0.3	0.5	0.7	1.4	2.8	5.2	8.0	9.1
2000.....	1.6	*	0.1	0.1	0.4	0.5	0.7	1.4	2.9	5.3	8.1	9.6
1999.....	1.5	*	*	0.2	0.4	0.5	0.8	1.5	2.9	5.1	7.9	9.6
All races 3/, male												
2004.....	1.4	*	*	0.2	0.3	0.5	0.7	1.2	2.6	5.1	8.3	12.7
2003.....	1.4	*	*	0.2	0.4	0.5	0.7	1.3	2.4	4.7	8.7	10.7
2002.....	1.4	*	*	0.2	0.3	0.5	0.8	1.3	2.6	4.7	7.7	11.7
2001.....	1.4	*	*	0.1	0.4	0.5	0.7	1.4	2.5	5.2	8.2	10.7
2000.....	1.4	*	*	0.1	0.4	0.5	0.7	1.4	2.7	5.0	9.2	12.5
1999.....	1.5	*	*	0.2	0.4	0.5	0.9	1.4	2.9	5.1	8.5	11.3
All races 3/, female												
2004.....	1.7	*	*	0.2	0.3	0.3	0.7	1.4	3.1	5.3	8.0	8.2
2003.....	1.6	*	*	0.2	0.3	0.4	0.6	1.4	3.2	5.3	7.6	8.0
2002.....	1.6	*	*	0.2	0.3	0.3	0.6	1.3	3.2	5.4	7.6	8.3
2001.....	1.6	*	*	0.1	0.2	0.4	0.7	1.4	3.2	5.1	7.8	8.4
2000.....	1.7	*	*	0.1	0.3	0.5	0.7	1.5	3.1	5.6	7.4	8.4
1999.....	1.6	*	*	0.2	0.3	0.4	0.6	1.5	2.8	5.1	7.5	8.9
White, both sexes												
2004.....	1.7	*	*	0.2	0.3	0.4	0.7	1.3	2.9	5.4	8.4	9.9
2003.....	1.6	*	*	0.2	0.4	0.4	0.6	1.3	2.8	5.1	8.3	9.1
2002.....	1.6	*	*	0.2	0.3	0.4	0.7	1.3	2.9	5.1	7.9	9.6
2001.....	1.6	*	*	0.1	0.3	0.5	0.7	1.4	2.9	5.3	8.1	9.1
2000.....	1.7	*	*	0.1	0.3	0.5	0.7	1.4	3.0	5.4	8.4	9.7
1999.....	1.6	*	*	0.2	0.3	0.4	0.7	1.5	2.9	5.1	8.0	9.8
White, male												
2004.....	1.5	*	*	0.2	0.3	0.5	0.7	1.1	2.7	5.2	8.6	13.0
2003.....	1.4	*	*	0.2	0.4	0.5	0.7	1.2	2.3	4.8	9.1	10.9
2002.....	1.4	*	*	0.2	0.4	0.4	0.8	1.2	2.6	4.8	8.1	12.0
2001.....	1.5	*	*	0.1	0.4	0.5	0.7	1.4	2.5	5.4	8.4	10.6
2000.....	1.5	*	*	*	0.4	0.6	0.7	1.4	2.8	5.1	9.4	12.4
1999.....	1.5	*	*	0.2	0.4	0.5	0.9	1.4	2.9	5.0	8.6	12.0

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Of other mesothelial and soft tissue (C47-C49)

White, female

2004.....	1.8	*	*	0.2	0.3	0.3	0.6	1.4	3.1	5.5	8.2	8.5
2003.....	1.8	*	*	0.2	0.3	0.3	0.6	1.4	3.3	5.4	7.7	8.3
2002.....	1.8	*	*	0.2	0.3	0.3	0.6	1.4	3.1	5.4	7.7	8.5
2001.....	1.7	*	*	0.1	0.3	0.4	0.6	1.4	3.2	5.2	8.0	8.4
2000.....	1.8	*	*	0.1	0.3	0.5	0.7	1.4	3.1	5.7	7.7	8.5
1999.....	1.7	*	*	0.1	0.3	0.4	0.5	1.5	2.8	5.2	7.5	8.9

Black, both sexes

2004.....	1.3	*	*	*	0.5	0.5	1.0	1.8	2.9	4.7	7.4	7.0
2003.....	1.3	*	*	*	0.3	0.7	0.9	1.9	3.2	5.0	7.1	7.0
2002.....	1.2	*	*	*	0.4	0.5	0.9	1.7	3.6	5.4	6.9	5.9
2001.....	1.2	*	*	*	0.4	0.6	0.8	1.9	3.2	4.8	6.8	9.8
2000.....	1.3	*	*	*	0.5	0.4	1.1	1.9	3.3	5.5	5.8	9.1
1999.....	1.3	*	*	*	0.5	0.6	1.2	1.6	3.4	5.2	7.8	8.3

Black, male

2004.....	1.1	*	*	*	*	*	0.9	1.6	2.4	4.8	6.7	*
2003.....	1.2	*	*	*	*	*	1.0	2.1	3.2	4.1	6.2	*
2002.....	1.1	*	*	*	*	*	1.1	1.8	3.2	4.7	5.7	*
2001.....	1.2	*	*	*	0.7	*	1.0	1.6	2.9	4.3	7.4	*
2000.....	1.1	*	*	*	*	*	1.2	1.1	3.1	4.7	6.7	*
1999.....	1.3	*	*	*	0.8	*	1.5	1.4	3.4	5.6	7.7	*

Black, female

2004.....	1.5	*	*	*	*	*	1.0	2.0	3.3	4.6	7.9	*
2003.....	1.4	*	*	*	*	0.8	0.8	1.7	3.2	5.7	7.7	*
2002.....	1.3	*	*	*	*	*	0.8	1.6	3.8	5.9	7.6	*
2001.....	1.3	*	*	*	*	*	0.7	2.1	3.5	5.1	6.4	8.8
2000.....	1.4	*	*	*	*	*	1.0	2.6	3.4	6.1	5.3	*
1999.....	1.4	*	*	*	*	*	1.0	1.7	3.5	4.9	7.9	9.2

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of breast (C50)												
All races 3/, both sexes												
2004.....	14.1	*	*	*	*	1.0	5.7	15.0	29.1	44.9	72.2	124.2
2003.....	14.4	*	*	*	*	1.0	6.1	15.6	29.6	45.5	74.9	132.5
2002.....	14.5	*	*	*	*	1.1	6.1	16.1	29.5	46.5	76.5	134.6
2001.....	14.7	*	*	*	*	1.2	6.3	16.8	30.2	47.6	76.8	134.2
2000.....	15.0	*	*	*	*	1.2	6.3	16.9	31.2	49.1	78.9	147.4
1999.....	14.9	*	*	*	*	1.1	6.1	16.9	31.1	49.5	78.9	144.4
All races 3/, male												
2004.....	0.3	*	*	*	*	*	*	0.2	0.5	1.1	2.1	3.3
2003.....	0.3	*	*	*	*	*	*	0.2	0.5	1.1	2.0	3.9
2002.....	0.3	*	*	*	*	*	*	0.2	0.5	1.0	2.2	3.4
2001.....	0.3	*	*	*	*	*	*	0.2	0.6	1.4	2.3	3.5
2000.....	0.3	*	*	*	*	*	*	0.2	0.7	1.4	2.2	4.2
1999.....	0.3	*	*	*	*	*	*	0.2	0.6	1.2	2.0	4.5
All races 3/, female												
2004.....	27.5	*	*	*	*	2.0	11.3	29.3	55.8	81.6	119.5	178.6
2003.....	28.2	*	*	*	*	2.1	12.2	30.4	56.6	82.6	123.7	189.4
2002.....	28.3	*	*	*	*	2.1	12.0	31.4	56.2	84.4	125.9	191.5
2001.....	28.6	*	*	*	*	2.4	12.4	32.8	57.5	85.8	125.8	188.9
2000.....	29.2	*	*	*	*	2.3	12.4	33.0	59.3	88.3	128.9	205.7
1999.....	28.9	*	*	*	*	2.2	12.0	32.9	59.2	88.9	128.9	200.8
White, both sexes												
2004.....	14.6	*	*	*	*	0.8	5.0	13.6	28.1	44.8	73.0	124.8
2003.....	15.0	*	*	*	*	0.9	5.5	14.4	28.4	45.1	75.1	132.5
2002.....	15.1	*	*	*	*	0.9	5.3	14.9	28.5	46.2	76.6	135.3
2001.....	15.2	*	*	*	*	1.0	5.6	15.7	29.1	47.0	77.1	134.8
2000.....	15.7	*	*	*	*	1.0	5.7	15.8	30.2	49.2	79.4	147.3
1999.....	15.5	*	*	*	*	0.9	5.4	15.7	29.9	49.0	79.0	146.0
White, male												
2004.....	0.3	*	*	*	*	*	*	0.1	0.4	1.0	2.1	3.0
2003.....	0.3	*	*	*	*	*	*	0.2	0.5	1.0	2.0	3.9
2002.....	0.3	*	*	*	*	*	*	0.2	0.5	0.9	2.1	3.3
2001.....	0.3	*	*	*	*	*	*	0.2	0.6	1.4	2.2	3.3
2000.....	0.3	*	*	*	*	*	*	0.2	0.7	1.4	2.2	3.9
1999.....	0.3	*	*	*	*	*	*	0.2	0.5	1.2	2.0	4.4

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of breast (C50)												
White, female												
2004.....	28.6	*	*	*	*	1.7	10.1	27.0	54.3	82.2	121.3	179.7
2003.....	29.3	*	*	*	*	1.8	11.1	28.4	54.9	82.6	124.6	189.4
2002.....	29.5	*	*	*	*	1.8	10.7	29.4	55.0	84.6	126.5	192.6
2001.....	29.8	*	*	*	*	2.0	11.3	31.0	56.0	85.6	126.9	189.9
2000.....	30.7	*	*	*	*	2.1	11.3	31.2	57.9	89.3	130.2	205.5
1999.....	30.2	*	*	*	*	1.9	10.8	30.9	57.5	88.9	129.5	202.9
Black, both sexes												
2004.....	14.5	*	*	*	*	2.1	10.9	26.0	43.7	56.3	82.3	143.5
2003.....	15.1	*	*	*	*	2.2	11.1	26.5	45.4	59.3	90.8	161.6
2002.....	14.9	*	*	*	*	2.3	11.7	27.0	43.1	60.1	94.0	152.6
2001.....	15.0	*	*	*	*	2.6	11.1	27.3	44.6	63.5	92.3	152.0
2000.....	14.8	*	*	*	*	2.3	11.2	27.8	45.4	58.6	90.5	176.4
1999.....	15.0	*	*	*	*	2.3	11.6	28.7	46.3	63.3	94.5	147.3
Black, male												
2004.....	0.3	*	*	*	*	*	*	*	*	*	*	*
2003.....	0.3	*	*	*	*	*	*	*	*	*	*	*
2002.....	0.4	*	*	*	*	*	*	*	*	*	*	*
2001.....	0.4	*	*	*	*	*	*	*	*	*	*	*
2000.....	0.4	*	*	*	*	*	*	*	*	*	*	*
1999.....	0.4	*	*	*	*	*	*	*	*	*	*	*
Black, female												
2004.....	27.5	*	*	*	*	4.0	20.5	48.0	78.2	95.1	127.4	196.9
2003.....	28.5	*	*	*	*	4.1	20.8	48.9	81.1	100.1	141.0	222.4
2002.....	28.2	*	*	*	*	4.3	22.0	49.8	76.6	101.1	145.0	209.1
2001.....	28.3	*	*	*	*	5.0	20.8	50.3	79.8	106.6	141.8	205.9
2000.....	27.9	*	*	*	*	4.2	20.9	51.5	80.9	98.6	139.8	238.7
1999.....	28.2	*	*	*	*	4.3	21.7	53.0	82.2	106.2	145.4	199.3

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms of female genital organs (C51-C58)

White, female

2004.....	19.2	*	*	*	0.2	1.6	5.8	14.8	33.5	62.4	92.6	105.9
2003.....	19.3	*	*	*	0.2	1.5	5.6	15.2	35.5	62.0	92.7	108.6
2002.....	19.5	*	*	*	0.3	1.7	5.8	16.1	35.0	62.7	94.2	110.3
2001.....	19.6	*	*	*	0.2	1.8	5.9	16.4	34.5	62.3	96.3	113.0
2000.....	19.4	*	*	*	0.2	1.8	5.7	16.2	35.3	61.8	94.9	115.5
1999.....	19.2	*	*	*	0.2	1.6	6.0	16.6	35.2	62.1	93.1	110.5

Black, both sexes

2004.....	8.6	*	*	*	*	1.3	3.9	10.0	22.9	52.2	65.0	80.9
2003.....	8.5	*	*	*	*	1.3	3.9	9.8	24.0	47.8	69.7	89.3
2002.....	8.6	*	*	*	*	1.2	4.1	9.8	23.4	49.8	71.0	98.3
2001.....	8.5	*	*	*	*	1.3	4.1	9.4	24.0	49.2	71.3	94.9
2000.....	8.6	*	*	*	*	1.3	4.3	9.6	23.8	51.9	73.1	96.7
1999.....	8.4	*	*	*	*	1.0	4.0	8.8	25.1	52.7	71.9	91.5

Black, male

2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

Black, female

2004.....	16.4	*	*	*	*	2.4	7.3	18.5	41.4	89.5	101.8	113.0
2003.....	16.3	*	*	*	*	2.6	7.3	18.2	43.3	81.8	109.1	124.0
2002.....	16.4	*	*	*	*	2.3	7.7	18.3	42.3	85.2	111.3	136.0
2001.....	16.2	*	*	*	*	2.5	7.7	17.6	43.4	84.0	111.4	130.1
2000.....	16.4	*	*	*	*	2.4	8.1	17.8	42.9	88.4	114.2	132.6
1999.....	16.1	*	*	*	*	1.9	7.5	16.4	45.3	89.8	112.2	125.7

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Of vulva and vagina (C51-C52)

White, female

2004.....	0.9	*	*	*	*	*	0.2	0.4	1.0	2.3	5.0	10.4
2003.....	0.9	*	*	*	*	*	0.1	0.4	0.9	2.2	5.0	10.8
2002.....	0.9	*	*	*	*	*	0.2	0.4	0.9	2.1	5.0	11.4
2001.....	0.9	*	*	*	*	*	0.1	0.4	0.7	2.1	5.2	10.4
2000.....	0.9	*	*	*	*	*	0.1	0.4	0.9	2.5	5.0	11.1
1999.....	0.9	*	*	*	*	*	0.1	0.4	0.8	2.1	5.1	12.6

Black, both sexes

2004.....	0.3	*	*	*	*	*	*	*	0.8	1.3	2.3	*
2003.....	0.3	*	*	*	*	*	*	*	*	*	2.1	6.1
2002.....	0.2	*	*	*	*	*	*	*	*	*	2.4	*
2001.....	0.3	*	*	*	*	*	*	*	0.9	1.5	3.4	6.4
2000.....	0.3	*	*	*	*	*	*	*	*	*	2.6	7.2
1999.....	0.3	*	*	*	*	*	*	*	*	1.3	3.1	*

Black, male

2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

Black, female

2004.....	0.6	*	*	*	*	*	*	*	1.5	2.3	3.6	*
2003.....	0.5	*	*	*	*	*	*	*	*	*	3.3	8.5
2002.....	0.5	*	*	*	*	*	*	*	*	*	3.7	*
2001.....	0.6	*	*	*	*	*	*	*	1.6	2.6	5.3	8.8
2000.....	0.5	*	*	*	*	*	*	*	*	*	4.1	9.9
1999.....	0.5	*	*	*	*	*	*	*	*	2.3	4.8	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of cervix uteri (C53)												
White, female												
2004.....	2.4	*	*	*	*	1.0	2.6	3.7	5.0	5.1	5.5	5.9
2003.....	2.5	*	*	*	*	0.9	2.7	4.0	5.0	4.8	6.0	7.1
2002.....	2.5	*	*	*	*	1.0	2.8	4.0	4.7	5.3	6.0	7.3
2001.....	2.7	*	*	*	*	1.1	3.0	4.5	5.0	5.0	6.6	8.8
2000.....	2.7	*	*	*	*	1.1	2.8	4.3	5.2	5.7	7.1	8.8
1999.....	2.8	*	*	*	*	1.1	2.9	4.7	5.4	5.7	6.7	9.5
Black, both sexes												
2004.....	2.0	*	*	*	*	0.8	2.2	3.7	5.3	7.6	8.8	16.1
2003.....	2.1	*	*	*	*	0.9	2.2	3.6	5.5	7.4	11.3	14.6
2002.....	2.2	*	*	*	*	0.8	2.5	4.1	5.0	9.0	10.8	17.5
2001.....	2.1	*	*	*	*	0.8	2.4	3.8	5.2	7.8	11.9	19.0
2000.....	2.4	*	*	*	*	0.9	2.6	4.3	6.3	9.5	11.9	19.7
1999.....	2.3	*	*	*	*	0.7	2.5	3.4	7.1	10.7	11.3	21.7
Black, male												
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....
Black, female												
2004.....	3.9	*	*	*	*	1.6	4.2	6.9	9.6	13.0	13.8	22.4
2003.....	4.0	*	*	*	*	1.7	4.2	6.8	10.0	12.6	17.7	20.3
2002.....	4.2	*	*	*	*	1.5	4.7	7.6	8.9	15.4	16.9	24.2
2001.....	4.0	*	*	*	*	1.5	4.5	7.0	9.3	13.3	18.5	26.1
2000.....	4.5	*	*	*	*	1.8	4.9	7.9	11.3	16.3	18.6	27.0
1999.....	4.4	*	*	*	*	1.2	4.7	6.4	12.8	18.3	17.6	29.8

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of corpus uteri and uterus, part unspecified (C54-C55)												
White, female												
2004.....	4.7	*	*	*	*	*	0.7	2.5	7.9	16.4	25.1	32.6
2003.....	4.7	*	*	*	*	*	0.6	2.5	8.3	16.9	24.1	32.8
2002.....	4.7	*	*	*	*	*	0.6	2.6	8.0	16.5	24.1	34.2
2001.....	4.8	*	*	*	*	0.2	0.6	2.7	7.8	16.2	26.2	34.7
2000.....	4.6	*	*	*	*	0.1	0.6	2.6	8.3	15.7	24.6	34.3
1999.....	4.6	*	*	*	*	*	0.6	2.5	7.7	16.3	26.1	32.8
Black, both sexes												
2004.....	2.9	*	*	*	*	*	0.5	1.9	8.0	23.0	26.1	26.2
2003.....	2.9	*	*	*	*	*	0.5	1.8	9.0	21.0	27.8	30.0
2002.....	3.0	*	*	*	*	*	0.4	1.9	8.4	22.3	28.3	38.9
2001.....	2.8	*	*	*	*	*	0.4	2.1	7.7	21.2	28.4	32.2
2000.....	2.9	*	*	*	*	*	0.4	1.6	8.3	22.7	30.5	33.2
1999.....	2.7	*	*	*	*	*	0.5	1.4	7.9	21.3	27.3	35.4
Black, male												
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....
Black, female												
2004.....	5.5	*	*	*	*	*	1.0	3.6	14.4	39.4	40.9	36.6
2003.....	5.5	*	*	*	*	*	1.0	3.3	16.2	36.0	43.5	41.7
2002.....	5.7	*	*	*	*	*	0.7	3.6	15.2	38.1	44.3	53.8
2001.....	5.4	*	*	*	*	*	0.7	3.9	13.9	36.2	44.3	44.2
2000.....	5.5	*	*	*	*	*	0.7	3.0	14.9	38.6	47.7	45.5
1999.....	5.2	*	*	*	*	*	0.9	2.6	14.3	36.3	42.5	48.6

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of ovary (C56)												
White, female												
2004.....	10.9	*	*	*	0.1	0.5	2.1	8.0	19.3	38.0	56.0	56.0
2003.....	11.0	*	*	*	0.1	0.5	2.1	8.1	21.0	37.1	56.3	56.4
2002.....	11.1	*	*	*	0.2	0.5	2.2	8.9	20.8	37.7	57.4	55.9
2001.....	11.0	*	*	*	*	0.5	2.1	8.6	20.5	38.0	56.8	57.3
2000.....	10.9	*	*	*	*	0.5	2.2	8.7	20.4	37.0	56.8	59.0
1999.....	10.6	*	*	*	0.1	0.4	2.3	8.7	20.5	36.9	53.6	54.3
Black, both sexes												
2004.....	3.3	*	*	*	*	*	0.9	3.9	8.6	20.1	27.3	32.7
2003.....	3.1	*	*	*	*	*	1.0	3.8	8.6	17.7	27.5	36.8
2002.....	3.1	*	*	*	*	*	1.0	3.4	9.1	17.1	28.8	34.7
2001.....	3.1	*	*	*	*	*	1.1	3.0	10.0	18.3	26.9	36.5
2000.....	2.9	*	*	*	*	*	1.0	3.3	8.4	18.1	26.6	34.4
1999.....	3.0	*	*	*	*	*	0.8	3.5	9.1	18.5	28.7	28.7
Black, male												
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....
Black, female												
2004.....	6.2	*	*	*	*	*	1.7	7.2	15.6	34.5	42.8	45.7
2003.....	6.0	*	*	*	*	*	1.9	7.1	15.5	30.2	43.1	51.0
2002.....	5.9	*	*	*	*	*	1.9	6.4	16.4	29.3	45.2	48.1
2001.....	5.9	*	*	*	*	*	2.1	5.7	18.0	31.2	42.0	50.1
2000.....	5.6	*	*	*	*	*	2.0	6.2	15.2	30.8	41.6	47.2
1999.....	5.7	*	*	*	*	*	1.5	6.6	16.4	31.5	44.8	39.4

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms of male genital organs (C60-C63)

All races 3/, both sexes

2004.....	10.1	*	*	*	0.1	0.2	0.3	1.3	7.3	31.4	93.1	181.7
2003.....	10.4	*	*	*	0.1	0.2	0.3	1.3	7.7	33.3	96.2	185.9
2002.....	10.8	*	*	*	0.1	0.2	0.3	1.2	7.9	35.0	100.8	195.9
2001.....	11.0	*	*	*	0.1	0.2	0.3	1.3	7.8	36.2	103.8	201.9
2000.....	11.3	*	*	*	0.1	0.2	0.3	1.2	8.2	37.4	106.5	210.1
1999.....	11.6	*	*	*	0.1	0.3	0.4	1.2	8.5	40.2	110.6	208.7

All races 3/, male

2004.....	20.5	*	*	*	0.2	0.4	0.6	2.6	15.1	68.8	231.5	585.7
2003.....	21.1	*	*	*	0.2	0.4	0.6	2.6	16.0	73.2	240.2	606.5
2002.....	21.9	*	*	*	0.2	0.5	0.6	2.4	16.4	77.0	252.6	647.3
2001.....	22.4	*	*	*	0.2	0.4	0.6	2.7	16.2	79.8	261.7	684.2
2000.....	22.9	*	*	*	0.2	0.4	0.6	2.4	17.1	82.9	269.8	726.1
1999.....	23.6	*	*	*	0.2	0.5	0.7	2.5	17.6	89.2	280.8	726.5

All races 3/, female

2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

White, both sexes

2004.....	10.2	*	*	*	0.1	0.2	0.3	1.1	6.4	28.2	86.5	173.1
2003.....	10.5	*	*	*	0.1	0.2	0.3	1.1	6.6	30.0	89.2	178.3
2002.....	10.9	*	*	*	0.1	0.3	0.4	1.0	6.7	31.1	93.3	187.6
2001.....	11.0	*	*	*	0.1	0.2	0.3	1.1	6.7	32.3	95.6	190.3
2000.....	11.2	*	*	*	0.1	0.3	0.3	1.0	7.1	32.9	97.8	199.5
1999.....	11.6	*	*	*	0.1	0.3	0.4	1.1	7.1	35.8	102.6	196.3

White, male

2004.....	20.6	*	*	*	0.2	0.4	0.6	2.2	13.1	61.1	213.4	557.1
2003.....	21.2	*	*	*	0.2	0.5	0.7	2.2	13.5	65.2	221.1	581.0
2002.....	22.0	*	*	*	0.2	0.5	0.7	2.0	13.9	67.7	232.3	619.3
2001.....	22.3	*	*	*	0.2	0.5	0.6	2.2	13.8	70.5	239.7	645.0
2000.....	22.8	*	*	*	0.2	0.5	0.6	2.1	14.6	72.2	246.3	690.4
1999.....	23.6	*	*	*	0.2	0.6	0.8	2.2	14.8	78.7	259.1	685.1

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of prostate (C61)												
All races 3/, both sexes												
2004.....	9.9	*	*	*	*	*	0.1	1.0	7.0	30.9	92.5	180.9
2003.....	10.2	*	*	*	*	*	0.1	1.0	7.4	32.9	95.5	184.9
2002.....	10.6	*	*	*	*	*	0.1	1.0	7.6	34.5	100.1	195.0
2001.....	10.8	*	*	*	*	*	0.1	1.1	7.5	35.7	103.2	200.9
2000.....	11.0	*	*	*	*	*	0.1	1.0	7.9	37.0	105.7	209.3
1999.....	11.4	*	*	*	*	*	0.1	1.0	8.2	39.7	109.9	207.5
All races 3/, male												
2004.....	20.1	*	*	*	*	*	0.1	2.1	14.6	67.8	229.8	583.1
2003.....	20.7	*	*	*	*	*	0.1	2.1	15.4	72.3	238.3	603.3
2002.....	21.5	*	*	*	*	*	0.1	1.9	15.8	75.9	251.0	644.5
2001.....	22.0	*	*	*	*	*	0.1	2.1	15.7	78.9	260.2	681.2
2000.....	22.5	*	*	*	*	*	0.1	2.0	16.6	82.0	267.9	723.1
1999.....	23.2	*	*	*	*	*	0.1	2.1	17.1	88.2	279.1	722.4
All races 3/, female												
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....
White, both sexes												
2004.....	10.0	*	*	*	*	*	*	0.8	6.1	27.7	85.8	172.3
2003.....	10.3	*	*	*	*	*	*	0.9	6.3	29.5	88.5	177.5
2002.....	10.6	*	*	*	*	*	0.1	0.8	6.4	30.6	92.6	186.7
2001.....	10.8	*	*	*	*	*	*	0.9	6.4	31.9	95.1	189.4
2000.....	11.0	*	*	*	*	*	*	0.8	6.8	32.5	97.0	198.6
1999.....	11.4	*	*	*	*	*	0.1	0.9	6.9	35.3	101.9	195.2
White, male												
2004.....	20.1	*	*	*	*	*	*	1.7	12.6	60.2	211.7	554.4
2003.....	20.7	*	*	*	*	*	*	1.7	13.0	64.2	219.4	578.2
2002.....	21.5	*	*	*	*	*	0.1	1.5	13.2	66.7	230.7	616.3
2001.....	21.9	*	*	*	*	*	*	1.8	13.3	69.6	238.2	641.8
2000.....	22.3	*	*	*	*	*	*	1.7	13.9	71.2	244.5	687.5
1999.....	23.1	*	*	*	*	*	0.1	1.7	14.2	77.7	257.4	681.0

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Of testis (C62)

All races 3/, both sexes

2004.....	0.1	*	*	*	0.1	0.2	0.2	0.2	0.1	*	*	*
2003.....	0.1	*	*	*	0.1	0.2	0.2	0.1	0.1	*	0.2	*
2002.....	0.1	*	*	*	0.1	0.2	0.2	0.2	0.1	0.1	*	*
2001.....	0.1	*	*	*	0.1	0.2	0.2	0.2	0.1	0.1	*	*
2000.....	0.1	*	*	*	0.1	0.2	0.2	0.1	0.1	*	*	*
1999.....	0.1	*	*	*	0.1	0.2	0.3	0.2	0.1	0.1	*	*

All races 3/, male

2004.....	0.2	*	*	*	0.2	0.4	0.4	0.4	0.2	*	*	*
2003.....	0.2	*	*	*	0.2	0.4	0.4	0.3	0.2	*	0.4	*
2002.....	0.3	*	*	*	0.2	0.5	0.4	0.3	0.3	0.3	*	*
2001.....	0.2	*	*	*	0.2	0.4	0.4	0.3	0.2	0.3	*	*
2000.....	0.2	*	*	*	0.2	0.4	0.5	0.3	0.2	*	*	*
1999.....	0.3	*	*	*	0.2	0.5	0.5	0.3	0.2	0.3	*	*

All races 3/, female

2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

White, both sexes

2004.....	0.1	*	*	*	0.1	0.2	0.2	0.2	0.1	*	*	*
2003.....	0.1	*	*	*	0.1	0.2	0.2	0.2	0.1	*	*	*
2002.....	0.2	*	*	*	0.1	0.3	0.3	0.2	0.1	0.2	*	*
2001.....	0.1	*	*	*	0.1	0.2	0.2	0.2	*	*	*	*
2000.....	0.1	*	*	*	0.1	0.2	0.3	0.1	0.1	*	*	*
1999.....	0.2	*	*	*	0.1	0.3	0.3	0.2	0.1	0.1	*	*

White, male

2004.....	0.3	*	*	*	0.2	0.4	0.5	0.4	0.2	*	*	*
2003.....	0.3	*	*	*	0.2	0.5	0.5	0.3	0.2	*	*	*
2002.....	0.3	*	*	*	0.2	0.5	0.5	0.4	0.3	0.3	*	*
2001.....	0.3	*	*	*	0.2	0.5	0.5	0.4	*	*	*	*
2000.....	0.3	*	*	*	0.2	0.5	0.5	0.3	0.2	*	*	*
1999.....	0.3	*	*	*	0.2	0.6	0.6	0.4	0.3	0.3	*	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Of penis and other and unspecified male genital organs (C60,C63)

All races 3/, both sexes

2004.....	0.1	*	*	*	*	*	*	0.1	0.2	0.4	0.5	0.6
2003.....	0.1	*	*	*	*	*	*	0.1	0.2	0.3	0.6	0.8
2002.....	0.1	*	*	*	*	*	*	0.1	0.2	0.4	0.5	0.6
2001.....	0.1	*	*	*	*	*	*	0.1	0.2	0.3	0.5	0.8
2000.....	0.1	*	*	*	*	*	*	0.1	0.2	0.3	0.7	0.7
1999.....	0.1	*	*	*	*	*	*	*	0.2	0.3	0.6	1.1

All races 3/, male

2004.....	0.2	*	*	*	*	*	*	0.2	0.3	0.9	1.3	2.1
2003.....	0.2	*	*	*	*	*	*	0.2	0.4	0.7	1.4	2.6
2002.....	0.2	*	*	*	*	*	*	0.1	0.4	0.8	1.3	2.0
2001.....	0.2	*	*	*	*	*	*	0.2	0.3	0.7	1.2	2.8
2000.....	0.2	*	*	*	*	*	*	0.1	0.4	0.7	1.7	2.4
1999.....	0.2	*	*	*	*	*	*	*	0.3	0.7	1.4	3.7

All races 3/, female

2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

White, both sexes

2004.....	0.1	*	*	*	*	*	*	0.1	0.2	0.4	0.6	0.6
2003.....	0.1	*	*	*	*	*	*	0.1	0.2	0.3	0.6	0.7
2002.....	0.1	*	*	*	*	*	*	0.1	0.2	0.3	0.5	0.7
2001.....	0.1	*	*	*	*	*	*	0.1	0.2	0.3	0.5	0.9
2000.....	0.1	*	*	*	*	*	*	0.1	0.2	0.3	0.6	0.7
1999.....	0.1	*	*	*	*	*	*	*	0.1	0.3	0.6	1.0

White, male

2004.....	0.2	*	*	*	*	*	*	0.2	0.3	0.8	1.4	2.0
2003.....	0.2	*	*	*	*	*	*	0.2	0.4	0.7	1.4	2.1
2002.....	0.2	*	*	*	*	*	*	0.1	0.3	0.8	1.3	2.1
2001.....	0.2	*	*	*	*	*	*	0.1	0.3	0.7	1.2	2.9
2000.....	0.2	*	*	*	*	*	*	0.2	0.4	0.7	1.6	2.3
1999.....	0.2	*	*	*	*	*	*	*	0.3	0.7	1.4	3.6

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of urinary tract (C64-C68)												
All races 3/, both sexes												
2004.....	8.8	*	*	0.1	*	0.2	0.9	4.4	13.9	32.3	65.9	102.7
2003.....	8.7	*	*	0.1	0.1	0.2	1.0	4.3	14.2	32.5	65.3	101.4
2002.....	8.8	*	*	0.1	0.1	0.2	1.0	4.6	14.0	34.6	65.1	102.0
2001.....	8.7	*	0.1	0.1	0.1	0.2	1.1	4.6	14.4	33.8	63.8	103.9
2000.....	8.7	*	*	0.1	0.1	0.1	1.0	4.5	14.3	33.5	64.0	106.0
1999.....	8.5	*	*	0.1	*	0.2	1.0	4.5	14.4	33.5	61.8	103.0
All races 3/, male												
2004.....	11.7	*	*	*	*	0.2	1.2	6.4	20.9	49.0	105.3	182.8
2003.....	11.6	*	*	*	*	0.2	1.3	6.4	21.4	48.5	105.0	178.8
2002.....	11.6	*	*	0.1	*	0.2	1.3	6.7	21.1	51.4	103.3	183.6
2001.....	11.6	*	*	*	*	0.2	1.4	6.8	21.6	51.5	102.5	192.4
2000.....	11.4	*	*	*	*	0.1	1.3	6.6	21.3	50.7	103.3	198.9
1999.....	11.2	*	*	*	*	0.2	1.4	6.6	21.5	50.6	100.8	188.4
All races 3/, female												
2004.....	6.0	*	*	*	*	0.1	0.6	2.4	7.5	18.4	39.4	66.7
2003.....	6.0	*	*	0.1	*	0.1	0.6	2.3	7.5	19.2	38.7	67.2
2002.....	6.1	*	*	*	*	0.1	0.6	2.6	7.5	20.5	39.7	66.6
2001.....	6.0	*	*	0.1	*	0.2	0.8	2.6	7.9	19.2	38.3	66.9
2000.....	6.0	*	*	*	*	0.1	0.7	2.6	7.9	19.4	38.5	68.1
1999.....	5.9	*	*	*	*	0.1	0.7	2.5	7.9	19.5	36.5	68.6
White, both sexes												
2004.....	9.9	*	*	0.1	*	0.1	0.9	4.5	14.6	33.7	68.4	105.5
2003.....	9.7	*	*	0.1	*	0.1	0.9	4.4	14.6	33.9	67.2	104.1
2002.....	9.8	*	*	0.1	*	0.1	1.0	4.7	14.5	35.8	66.7	103.9
2001.....	9.7	*	*	0.1	*	0.1	1.0	4.7	14.8	35.2	65.2	106.4
2000.....	9.6	*	*	0.1	*	0.1	1.0	4.7	14.9	34.6	65.3	109.2
1999.....	9.3	*	*	0.1	*	0.1	1.0	4.5	14.8	34.3	63.2	104.6
White, male												
2004.....	13.2	*	*	*	*	0.1	1.1	6.6	21.9	51.3	109.9	190.9
2003.....	12.9	*	*	*	*	0.1	1.3	6.5	22.1	50.4	108.5	186.6
2002.....	13.0	*	*	*	*	0.2	1.3	7.0	21.9	53.6	106.3	190.7
2001.....	12.9	*	*	*	*	0.1	1.4	7.0	22.1	53.8	105.4	200.5
2000.....	12.7	*	*	*	*	*	1.4	6.8	22.1	52.3	106.0	206.8
1999.....	12.4	*	*	*	*	0.2	1.3	6.5	22.1	52.2	103.8	194.9

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of urinary tract (C64-C68)												
White, female												
2004.....	6.6	*	*	*	*	*	0.6	2.4	7.7	18.6	40.1	67.0
2003.....	6.5	*	*	*	*	*	0.5	2.2	7.5	19.9	39.2	67.6
2002.....	6.7	*	*	*	*	*	0.6	2.6	7.5	20.8	40.1	66.2
2001.....	6.5	*	*	*	*	0.1	0.7	2.5	7.9	19.5	38.5	67.0
2000.....	6.6	*	*	*	*	0.1	0.7	2.6	8.1	19.7	38.6	69.6
1999.....	6.4	*	*	*	*	*	0.6	2.5	7.9	19.5	36.6	68.4
Black, both sexes												
2004.....	5.3	*	*	*	*	0.4	1.2	4.8	12.6	28.3	53.9	88.8
2003.....	5.5	*	*	*	*	*	1.5	5.0	13.6	27.7	57.8	89.0
2002.....	5.5	*	*	*	*	0.4	1.0	4.8	13.8	30.3	58.4	93.2
2001.....	5.5	*	*	*	*	0.4	1.7	5.0	14.9	27.8	58.5	86.6
2000.....	5.3	*	*	*	*	*	1.1	4.6	13.1	30.7	61.4	80.4
1999.....	5.5	*	*	*	*	0.5	1.4	5.5	14.7	31.9	53.6	93.1
Black, male												
2004.....	6.2	*	*	*	*	*	1.7	6.8	18.3	39.6	77.4	120.2
2003.....	6.6	*	*	*	*	*	2.1	7.1	19.2	40.7	87.4	126.2
2002.....	6.3	*	*	*	*	*	1.4	6.1	20.0	40.0	85.2	135.1
2001.....	6.5	*	*	*	*	*	1.9	7.2	21.6	39.2	85.6	127.1
2000.....	6.4	*	*	*	*	*	1.2	6.3	19.1	44.8	93.0	140.7
1999.....	6.6	*	*	*	*	*	2.1	8.5	21.0	43.1	79.0	138.3
Black, female												
2004.....	4.5	*	*	*	*	*	0.8	3.1	7.9	20.2	40.6	76.4
2003.....	4.4	*	*	*	*	*	1.0	3.2	9.1	18.5	41.0	74.5
2002.....	4.7	*	*	*	*	*	0.6	3.7	8.9	23.4	43.2	77.2
2001.....	4.6	*	*	*	*	*	1.4	3.2	9.6	19.7	43.3	71.6
2000.....	4.3	*	*	*	*	*	1.1	3.0	8.4	20.8	43.6	57.9
1999.....	4.6	*	*	*	*	*	0.9	3.0	9.6	24.1	39.4	76.2

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Of kidney and renal pelvis (C64-C65)

All races 3/, both sexes

2004.....	4.2	*	*	0.1	*	0.1	0.6	3.0	8.7	16.7	26.8	32.7
2003.....	4.2	*	*	0.1	0.1	0.1	0.7	2.9	9.0	16.9	27.0	33.2
2002.....	4.2	*	*	0.1	0.0	0.1	0.7	3.3	8.8	17.7	26.3	32.5
2001.....	4.2	*	0.1	0.1	0.1	0.1	0.8	3.3	9.2	17.6	26.3	32.3
2000.....	4.2	*	*	0.1	0.1	0.1	0.7	3.3	9.0	17.1	26.6	33.8
1999.....	4.0	*	*	0.1	*	0.1	0.8	3.1	8.9	16.7	24.6	31.4

All races 3/, male

2004.....	5.3	*	*	*	*	0.2	0.8	4.3	12.8	23.9	38.2	47.0
2003.....	5.4	*	*	*	*	0.2	1.0	4.3	13.3	23.7	39.5	48.7
2002.....	5.4	*	*	*	*	0.2	1.0	4.8	13.0	25.1	37.9	50.1
2001.....	5.4	*	*	*	*	0.1	1.1	4.9	13.4	25.2	37.4	52.9
2000.....	5.3	*	*	*	*	0.1	1.0	4.8	13.1	24.2	38.2	55.1
1999.....	5.0	*	*	*	*	0.2	1.1	4.5	12.9	23.6	35.7	49.4

All races 3/, female

2004.....	3.1	*	*	*	*	*	0.4	1.7	5.0	10.6	19.1	26.3
2003.....	3.1	*	*	0.1	*	0.1	0.4	1.6	4.9	11.2	18.7	26.4
2002.....	3.1	*	*	*	*	*	0.4	1.8	4.9	11.6	18.7	24.9
2001.....	3.1	*	*	0.1	*	0.1	0.5	1.9	5.3	11.3	19.0	23.7
2000.....	3.1	*	*	*	*	*	0.5	1.9	5.3	11.3	18.9	25.1
1999.....	3.0	*	*	*	*	*	0.5	1.8	5.1	11.0	17.4	24.1

White, both sexes

2004.....	4.6	*	*	0.1	*	0.1	0.6	3.1	9.1	17.1	27.3	33.7
2003.....	4.6	*	*	0.1	*	0.1	0.7	3.0	9.1	17.5	27.7	33.8
2002.....	4.6	*	*	0.1	*	0.1	0.7	3.3	9.1	18.3	26.8	33.2
2001.....	4.6	*	*	0.1	*	0.1	0.7	3.4	9.4	18.2	26.7	32.8
2000.....	4.6	*	*	*	*	0.1	0.7	3.4	9.3	17.5	27.0	34.5
1999.....	4.3	*	*	0.1	*	0.1	0.7	3.1	9.1	17.1	24.9	31.5

White, male

2004.....	5.8	*	*	*	*	*	0.8	4.5	13.2	24.4	39.0	48.6
2003.....	5.9	*	*	*	*	0.1	1.0	4.4	13.6	24.3	40.4	50.0
2002.....	5.9	*	*	*	*	*	1.0	4.9	13.3	25.9	38.6	51.9
2001.....	5.8	*	*	*	*	*	1.0	5.0	13.6	26.0	37.7	54.4
2000.....	5.7	*	*	*	*	*	1.0	4.9	13.4	24.4	38.7	56.6
1999.....	5.4	*	*	*	*	0.1	1.0	4.5	13.3	23.9	36.3	49.8

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of kidney and renal pelvis (C64-C65)												
White, female												
2004.....	3.4	*	*	*	*	*	0.4	1.8	5.1	10.9	19.4	27.0
2003.....	3.4	*	*	*	*	*	0.4	1.7	4.9	11.6	19.1	26.6
2002.....	3.4	*	*	*	*	*	0.4	1.8	5.1	11.9	18.9	25.1
2001.....	3.4	*	*	*	*	*	0.4	1.8	5.4	11.6	19.4	23.8
2000.....	3.4	*	*	*	*	*	0.4	1.9	5.4	11.6	19.2	25.5
1999.....	3.2	*	*	*	*	*	0.4	1.8	5.3	11.3	17.5	24.2
Black, both sexes												
2004.....	2.8	*	*	*	*	0.4	0.8	3.1	8.0	15.8	25.8	28.2
2003.....	2.9	*	*	*	*	*	1.1	3.2	9.5	14.6	23.8	33.8
2002.....	2.8	*	*	*	*	*	0.8	3.5	8.6	15.4	26.1	28.2
2001.....	3.0	*	*	*	*	*	1.2	3.7	9.5	15.0	25.6	29.8
2000.....	2.9	*	*	*	*	*	0.9	3.4	8.8	16.5	25.6	29.7
1999.....	2.9	*	*	*	*	0.4	1.1	3.8	8.6	16.1	23.5	31.2
Black, male												
2004.....	3.6	*	*	*	*	*	1.1	4.5	12.1	24.0	37.3	38.7
2003.....	3.7	*	*	*	*	*	1.6	4.9	13.8	22.2	37.0	43.8
2002.....	3.5	*	*	*	*	*	1.2	4.8	13.9	21.6	36.1	39.7
2001.....	3.8	*	*	*	*	*	1.5	5.4	14.9	22.3	40.3	46.5
2000.....	3.6	*	*	*	*	*	1.0	4.9	13.1	26.3	38.2	49.6
1999.....	3.6	*	*	*	*	*	1.6	5.8	13.2	23.5	32.0	49.2
Black, female												
2004.....	2.2	*	*	*	*	*	*	1.8	4.7	10.0	19.4	24.0
2003.....	2.2	*	*	*	*	*	0.7	1.8	6.0	9.2	16.4	30.0
2002.....	2.2	*	*	*	*	*	*	2.4	4.2	11.0	20.5	23.8
2001.....	2.2	*	*	*	*	*	1.0	2.2	5.2	9.9	17.4	23.6
2000.....	2.1	*	*	*	*	*	0.8	2.2	5.2	9.6	18.6	22.3
1999.....	2.2	*	*	*	*	*	*	2.1	5.0	10.9	18.8	24.5

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of bladder (C67)												
All races 3/, both sexes												
2004.....	4.4	*	*	*	*	*	0.3	1.3	5.0	14.9	37.4	67.9
2003.....	4.3	*	*	*	*	*	0.2	1.3	5.0	14.9	36.4	64.8
2002.....	4.4	*	*	*	*	*	0.3	1.3	4.9	15.9	37.0	66.6
2001.....	4.3	*	*	*	*	*	0.3	1.2	5.0	15.4	35.7	68.9
2000.....	4.3	*	*	*	*	*	0.2	1.2	5.0	15.5	35.7	69.6
1999.....	4.3	*	*	*	*	*	0.2	1.3	5.2	15.9	35.3	68.5
All races 3/, male												
2004.....	6.2	*	*	*	*	*	0.3	2.0	7.9	24.0	64.7	133.0
2003.....	6.0	*	*	*	*	*	0.3	2.0	7.8	23.9	63.1	125.1
2002.....	6.0	*	*	*	*	*	0.3	1.8	7.7	25.2	63.0	128.9
2001.....	6.0	*	*	*	*	*	0.3	1.8	7.9	25.2	62.5	134.8
2000.....	5.9	*	*	*	*	*	0.3	1.7	7.9	25.2	62.6	139.4
1999.....	5.9	*	*	*	*	*	0.3	2.0	8.2	25.7	62.5	134.8
All races 3/, female												
2004.....	2.7	*	*	*	*	*	0.2	0.6	2.3	7.3	19.0	38.6
2003.....	2.6	*	*	*	*	*	0.2	0.6	2.4	7.4	18.5	38.1
2002.....	2.8	*	*	*	*	*	0.2	0.7	2.4	8.2	19.7	39.6
2001.....	2.7	*	*	*	*	*	0.3	0.6	2.3	7.2	18.1	41.4
2000.....	2.7	*	*	*	*	*	0.2	0.6	2.4	7.5	18.1	41.2
1999.....	2.7	*	*	*	*	*	0.2	0.6	2.5	7.8	17.7	41.8
White, both sexes												
2004.....	5.0	*	*	*	*	*	0.2	1.3	5.3	15.8	39.1	69.6
2003.....	4.8	*	*	*	*	*	0.2	1.3	5.3	15.7	37.5	66.6
2002.....	4.9	*	*	*	*	*	0.3	1.3	5.1	16.5	38.1	67.7
2001.....	4.8	*	*	*	*	*	0.3	1.2	5.2	16.1	36.7	70.8
2000.....	4.8	*	*	*	*	*	0.3	1.2	5.3	16.2	36.4	72.0
1999.....	4.8	*	*	*	*	*	0.2	1.3	5.4	16.3	36.3	69.8
White, male												
2004.....	7.1	*	*	*	*	*	0.3	2.1	8.4	25.7	68.3	139.3
2003.....	6.8	*	*	*	*	*	0.3	2.0	8.3	25.3	65.7	131.0
2002.....	6.8	*	*	*	*	*	0.4	1.9	8.2	26.5	65.3	134.1
2001.....	6.8	*	*	*	*	*	0.3	1.9	8.3	26.6	65.0	141.1
2000.....	6.7	*	*	*	*	*	0.3	1.9	8.4	26.5	64.7	145.7
1999.....	6.7	*	*	*	*	*	0.3	1.9	8.5	26.9	64.8	140.7

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of bladder (C67)												
White, female												
2004.....	2.9	*	*	*	*	*	0.2	0.6	2.4	7.3	19.2	38.1
2003.....	2.9	*	*	*	*	*	0.1	0.5	2.4	7.6	18.4	38.1
2002.....	3.0	*	*	*	*	*	0.2	0.7	2.3	8.1	19.8	38.9
2001.....	2.9	*	*	*	*	*	0.3	0.6	2.2	7.2	17.8	41.3
2000.....	2.9	*	*	*	*	*	0.2	0.6	2.5	7.5	17.8	42.1
1999.....	2.9	*	*	*	*	*	0.2	0.6	2.4	7.6	17.7	41.4
Black, both sexes												
2004.....	2.4	*	*	*	*	*	0.5	1.6	4.1	11.7	27.2	60.1
2003.....	2.4	*	*	*	*	*	0.4	1.6	3.9	12.0	32.3	53.4
2002.....	2.5	*	*	*	*	*	*	1.1	4.5	14.3	31.0	63.3
2001.....	2.4	*	*	*	*	*	0.4	1.3	4.9	11.9	31.2	55.0
2000.....	2.3	*	*	*	*	*	*	1.0	4.1	13.3	34.7	49.7
1999.....	2.5	*	*	*	*	*	*	1.5	5.6	15.0	29.1	60.3
Black, male												
2004.....	2.6	*	*	*	*	*	*	2.3	5.9	14.5	39.5	81.5
2003.....	2.8	*	*	*	*	*	*	2.2	5.4	17.5	48.8	80.3
2002.....	2.7	*	*	*	*	*	*	1.2	5.6	18.3	47.6	91.1
2001.....	2.5	*	*	*	*	*	*	1.7	6.4	16.1	43.0	78.3
2000.....	2.7	*	*	*	*	*	*	1.4	5.8	17.6	52.7	88.8
1999.....	2.9	*	*	*	*	*	*	2.6	7.4	19.1	46.1	86.7
Black, female												
2004.....	2.2	*	*	*	*	*	*	0.9	2.6	9.7	20.3	51.6
2003.....	2.1	*	*	*	*	*	*	1.2	2.8	8.1	23.0	42.9
2002.....	2.3	*	*	*	*	*	*	0.9	3.7	11.6	21.6	52.6
2001.....	2.2	*	*	*	*	*	*	0.8	3.8	8.9	24.6	46.3
2000.....	2.0	*	*	*	*	*	*	*	2.7	10.3	24.5	35.2
1999.....	2.2	*	*	*	*	*	*	*	4.2	12.1	19.6	50.4

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Of other and unspecified sites within the urinary tract (C66,C68)

All races 3/, both sexes

2004.....	0.2	*	*	*	*	*	*	0.1	0.2	0.8	1.8	2.1
2003.....	0.2	*	*	*	*	*	*	0.1	0.2	0.8	1.9	3.4
2002.....	0.2	*	*	*	*	*	*	0.1	0.3	0.9	1.8	2.9
2001.....	0.2	*	*	*	*	*	*	0.1	0.2	0.9	1.8	2.7
2000.....	0.2	*	*	*	*	*	*	0.1	0.3	0.9	1.8	2.6
1999.....	0.2	*	*	*	*	*	*	0.1	0.3	0.9	1.9	3.1

All races 3/, male

2004.....	0.2	*	*	*	*	*	*	*	0.3	1.1	2.4	2.8
2003.....	0.2	*	*	*	*	*	*	0.1	0.2	0.9	2.4	5.1
2002.....	0.3	*	*	*	*	*	*	0.1	0.4	1.1	2.4	4.7
2001.....	0.2	*	*	*	*	*	*	0.1	0.3	1.1	2.6	4.7
2000.....	0.2	*	*	*	*	*	*	*	0.3	1.3	2.5	4.4
1999.....	0.2	*	*	*	*	*	*	*	0.4	1.3	2.6	4.2

All races 3/, female

2004.....	0.2	*	*	*	*	*	*	*	0.2	0.5	1.4	1.8
2003.....	0.2	*	*	*	*	*	*	*	0.2	0.7	1.6	2.7
2002.....	0.2	*	*	*	*	*	*	*	0.2	0.8	1.3	2.2
2001.....	0.2	*	*	*	*	*	*	*	0.2	0.6	1.3	1.8
2000.....	0.2	*	*	*	*	*	*	*	0.2	0.6	1.4	1.8
1999.....	0.2	*	*	*	*	*	*	*	0.3	0.6	1.4	2.7

White, both sexes

2004.....	0.2	*	*	*	*	*	*	*	0.2	0.8	1.9	2.3
2003.....	0.2	*	*	*	*	*	*	0.1	0.2	0.7	2.0	3.6
2002.....	0.2	*	*	*	*	*	*	0.1	0.3	1.0	1.8	3.0
2001.....	0.2	*	*	*	*	*	*	0.1	0.2	0.9	1.9	2.8
2000.....	0.2	*	*	*	*	*	*	0.1	0.2	0.9	1.9	2.7
1999.....	0.3	*	*	*	*	*	*	0.1	0.3	0.9	1.9	3.3

White, male

2004.....	0.3	*	*	*	*	*	*	*	0.3	1.2	2.7	3.0
2003.....	0.3	*	*	*	*	*	*	0.1	0.3	0.8	2.5	5.5
2002.....	0.3	*	*	*	*	*	*	*	0.4	1.2	2.5	4.7
2001.....	0.3	*	*	*	*	*	*	*	0.2	1.2	2.7	4.9
2000.....	0.3	*	*	*	*	*	*	*	0.3	1.3	2.6	4.6
1999.....	0.3	*	*	*	*	*	*	*	0.4	1.3	2.7	4.4

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasm of eye and adnexa (C69)

All races 3/, both sexes

2004.....	0.1	*	*	*	*	*	*	0.1	0.1	0.3	0.3	0.5
2003.....	0.1	*	*	*	*	*	*	0.1	0.2	0.2	0.4	0.7
2002.....	0.1	*	*	*	*	*	*	0.1	0.2	0.2	0.5	0.7
2001.....	0.1	*	*	*	*	*	*	0.1	0.1	0.3	0.5	0.6
2000.....	0.1	*	*	*	*	*	*	0.1	0.2	0.3	0.4	0.8
1999.....	0.1	*	*	*	*	*	*	0.1	0.1	0.3	0.5	0.8

All races 3/, male

2004.....	0.1	*	*	*	*	*	*	*	0.2	0.3	0.5	*
2003.....	0.1	*	*	*	*	*	*	*	0.2	0.3	0.6	*
2002.....	0.1	*	*	*	*	*	*	*	0.2	0.3	0.6	*
2001.....	0.1	*	*	*	*	*	*	*	*	0.3	0.6	*
2000.....	0.1	*	*	*	*	*	*	*	0.2	0.4	0.6	*
1999.....	0.1	*	*	*	*	*	*	*	*	0.4	0.4	*

All races 3/, female

2004.....	0.1	*	*	*	*	*	*	*	*	0.2	*	*
2003.....	0.1	*	*	*	*	*	*	*	0.2	0.2	0.3	*
2002.....	0.1	*	*	*	*	*	*	*	0.2	0.2	0.4	0.7
2001.....	0.1	*	*	*	*	*	*	*	*	0.2	0.4	0.6
2000.....	0.1	*	*	*	*	*	*	*	*	0.2	0.3	0.7
1999.....	0.1	*	*	*	*	*	*	*	*	*	0.5	0.8

White, both sexes

2004.....	0.1	*	*	*	*	*	*	0.1	0.2	0.3	0.4	0.5
2003.....	0.1	*	*	*	*	*	*	0.1	0.2	0.3	0.5	0.7
2002.....	0.1	*	*	*	*	*	*	0.1	0.2	0.3	0.5	0.8
2001.....	0.1	*	*	*	*	*	*	0.1	0.1	0.3	0.5	0.7
2000.....	0.1	*	*	*	*	*	*	0.1	0.2	0.3	0.5	0.8
1999.....	0.1	*	*	*	*	*	*	0.1	0.1	0.3	0.5	0.9

White, male

2004.....	0.1	*	*	*	*	*	*	*	0.2	0.4	0.6	*
2003.....	0.1	*	*	*	*	*	*	*	0.2	0.3	0.7	*
2002.....	0.1	*	*	*	*	*	*	*	0.3	0.3	0.6	*
2001.....	0.1	*	*	*	*	*	*	*	*	0.4	0.7	*
2000.....	0.1	*	*	*	*	*	*	*	0.2	0.4	0.6	*
1999.....	0.1	*	*	*	*	*	*	*	*	0.4	0.5	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)												
All races 3/, both sexes												
2004.....	4.4	*	0.6	0.8	0.5	0.9	2.0	4.6	9.5	15.7	19.6	16.6
2003.....	4.4	*	0.7	0.8	0.6	1.0	2.4	4.8	9.7	15.5	19.5	15.2
2002.....	4.4	*	0.7	0.8	0.5	1.0	2.2	5.1	9.6	16.2	19.1	16.4
2001.....	4.4	*	0.7	0.7	0.6	1.1	2.3	5.0	9.7	15.4	20.4	16.4
2000.....	4.5	0.5	0.7	0.8	0.6	1.0	2.4	5.1	9.6	16.5	20.0	16.9
1999.....	4.6	*	0.7	0.8	0.6	1.1	2.4	5.1	10.3	16.9	20.4	16.0
All races 3/, male												
2004.....	4.9	*	0.7	0.8	0.6	1.1	2.5	5.8	11.8	18.4	23.7	19.4
2003.....	5.0	*	0.7	0.8	0.6	1.1	2.9	6.2	11.4	18.6	24.6	20.0
2002.....	5.0	*	0.8	0.9	0.7	1.1	2.8	6.2	11.7	20.2	23.8	20.1
2001.....	5.0	*	0.7	0.7	0.6	1.3	2.8	6.1	12.1	19.2	25.8	20.1
2000.....	5.1	*	0.7	0.8	0.7	1.3	3.1	6.2	11.8	20.2	24.3	22.4
1999.....	5.1	*	0.7	0.7	0.7	1.4	2.8	6.2	12.4	20.8	25.5	20.7
All races 3/, female												
2004.....	3.9	*	0.6	0.8	0.4	0.7	1.6	3.4	7.3	13.4	16.8	15.4
2003.....	3.9	*	0.6	0.8	0.5	0.9	2.0	3.5	8.1	12.8	16.1	13.1
2002.....	3.9	*	0.7	0.7	0.4	0.8	1.7	4.1	7.7	12.8	16.0	14.8
2001.....	3.9	*	0.6	0.8	0.5	0.8	1.7	3.9	7.6	12.3	16.8	14.8
2000.....	3.9	*	0.6	0.7	0.5	0.7	1.7	4.0	7.6	13.4	17.3	14.7
1999.....	4.1	*	0.8	0.8	0.5	0.8	1.9	4.1	8.4	13.7	17.1	14.1
White, both sexes												
2004.....	4.9	*	0.7	0.8	0.6	1.0	2.3	5.0	10.3	17.0	20.8	17.3
2003.....	5.0	*	0.7	0.9	0.6	1.1	2.7	5.2	10.5	16.6	20.7	15.7
2002.....	5.0	*	0.7	0.8	0.5	1.1	2.5	5.6	10.4	17.4	20.1	17.1
2001.....	5.0	*	0.7	0.8	0.6	1.2	2.6	5.4	10.5	16.5	21.6	17.0
2000.....	5.1	*	0.7	0.8	0.6	1.1	2.6	5.6	10.4	17.8	21.0	17.5
1999.....	5.2	*	0.8	0.8	0.6	1.3	2.6	5.6	11.1	18.2	21.4	16.6
White, male												
2004.....	5.5	*	0.7	0.8	0.7	1.1	2.7	6.2	12.8	20.0	24.8	20.6
2003.....	5.6	*	0.8	0.9	0.7	1.3	3.2	6.7	12.3	19.8	25.8	20.7
2002.....	5.7	*	0.7	0.9	0.7	1.2	3.1	6.8	12.4	21.6	24.8	21.2
2001.....	5.7	*	0.8	0.7	0.7	1.5	3.2	6.7	12.9	20.5	27.1	20.5
2000.....	5.7	*	0.7	0.9	0.7	1.4	3.4	6.8	12.8	21.7	25.5	23.5
1999.....	5.8	*	0.8	0.8	0.6	1.5	3.1	6.7	13.3	22.3	26.8	21.8

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)												
White, female												
2004.....	4.4	*	0.6	0.7	0.4	0.8	1.8	3.7	8.0	14.4	18.1	15.8
2003.....	4.4	*	0.6	0.9	0.5	1.0	2.3	3.8	8.7	13.9	17.2	13.4
2002.....	4.4	*	0.6	0.8	0.4	1.0	1.9	4.5	8.5	13.8	16.9	15.4
2001.....	4.4	*	0.7	0.8	0.5	0.9	1.9	4.3	8.2	13.0	18.0	15.6
2000.....	4.4	*	0.6	0.7	0.5	0.9	1.8	4.5	8.2	14.5	18.0	15.0
1999.....	4.6	*	0.8	0.8	0.5	1.0	2.1	4.5	9.1	14.8	17.9	14.5
Black, both sexes												
2004.....	2.0	*	*	0.7	0.5	0.7	1.3	2.8	4.9	8.0	10.0	10.4
2003.....	2.1	*	*	0.7	0.5	0.7	1.4	3.1	5.5	8.9	10.5	10.5
2002.....	2.1	*	1.1	0.8	0.5	0.6	1.2	2.7	5.6	8.7	11.8	9.5
2001.....	2.0	*	*	0.6	0.5	0.5	1.1	3.0	5.8	9.2	9.9	11.7
2000.....	2.1	*	*	0.8	0.6	0.7	1.6	2.9	5.7	8.1	11.9	10.6
1999.....	2.1	*	*	0.8	0.6	0.7	1.3	3.2	6.0	8.2	11.8	9.9
Black, male												
2004.....	2.2	*	*	0.7	*	0.8	1.9	3.9	6.7	7.9	14.2	*
2003.....	2.5	*	*	0.8	0.7	*	2.1	4.0	6.7	12.0	15.2	*
2002.....	2.3	*	*	1.0	*	0.8	1.3	2.7	8.1	10.4	16.4	*
2001.....	2.2	*	*	0.7	*	*	1.2	3.6	7.9	10.1	12.4	*
2000.....	2.4	*	*	0.9	0.7	1.1	2.0	3.7	7.1	9.8	13.0	*
1999.....	2.3	*	*	0.7	0.8	0.9	1.8	4.4	7.2	10.0	12.6	*
Black, female												
2004.....	1.8	*	*	0.8	*	*	0.7	1.9	3.4	8.1	7.6	11.4
2003.....	1.8	*	*	0.6	*	*	0.9	2.2	4.6	6.7	7.8	11.7
2002.....	1.9	*	*	*	*	*	1.1	2.8	3.7	7.4	9.3	10.3
2001.....	1.8	*	*	*	*	*	1.0	2.4	4.2	8.5	8.6	9.7
2000.....	1.9	*	*	0.6	*	*	1.2	2.2	4.6	6.9	11.3	10.7
1999.....	1.9	*	*	0.9	*	*	1.0	2.1	5.0	6.9	11.4	9.2

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of brain (C71)												
All races 3/, both sexes												
2004.....	4.3	*	0.6	0.8	0.5	0.9	2.0	4.5	9.3	15.5	19.3	16.1
2003.....	4.4	*	0.6	0.8	0.6	1.0	2.4	4.7	9.5	15.2	19.1	14.6
2002.....	4.4	*	0.7	0.8	0.5	0.9	2.2	5.0	9.5	15.8	18.7	16.0
2001.....	4.3	*	0.6	0.7	0.5	1.0	2.3	4.9	9.6	15.1	20.0	16.0
2000.....	4.4	0.5	0.7	0.8	0.6	1.0	2.4	5.0	9.5	16.2	19.7	16.3
1999.....	4.5	*	0.7	0.7	0.5	1.1	2.3	5.0	10.1	16.6	19.9	15.5
All races 3/, male												
2004.....	4.8	*	0.7	0.8	0.6	1.1	2.5	5.7	11.6	18.2	23.3	19.1
2003.....	4.9	*	0.7	0.8	0.6	1.1	2.9	6.1	11.2	18.4	24.3	19.2
2002.....	4.9	*	0.8	0.9	0.6	1.1	2.7	6.1	11.5	19.7	23.1	19.7
2001.....	4.9	*	0.6	0.7	0.6	1.3	2.8	6.0	11.9	18.8	25.3	19.7
2000.....	5.0	*	0.7	0.8	0.6	1.3	3.1	6.1	11.7	19.9	23.9	21.8
1999.....	5.0	*	0.7	0.7	0.6	1.3	2.7	6.1	12.2	20.5	25.0	20.1
All races 3/, female												
2004.....	3.8	*	0.5	0.7	0.4	0.7	1.5	3.3	7.2	13.2	16.6	14.7
2003.....	3.8	*	0.6	0.8	0.5	0.8	1.9	3.5	8.0	12.6	15.6	12.6
2002.....	3.8	*	0.6	0.7	0.4	0.8	1.7	4.0	7.6	12.5	15.8	14.4
2001.....	3.8	*	0.6	0.8	0.5	0.8	1.7	3.8	7.5	12.1	16.4	14.4
2000.....	3.9	*	0.6	0.7	0.5	0.7	1.7	4.0	7.4	13.2	16.9	14.1
1999.....	4.0	*	0.7	0.7	0.4	0.8	1.9	4.0	8.2	13.5	16.6	13.6
White, both sexes												
2004.....	4.9	*	0.6	0.8	0.5	1.0	2.2	4.9	10.2	16.8	20.5	16.7
2003.....	4.9	*	0.7	0.8	0.6	1.1	2.7	5.2	10.3	16.4	20.3	15.0
2002.....	4.9	*	0.6	0.8	0.5	1.0	2.5	5.6	10.2	17.0	19.7	16.7
2001.....	4.9	*	0.7	0.7	0.6	1.2	2.5	5.4	10.3	16.1	21.1	16.6
2000.....	5.0	*	0.7	0.8	0.6	1.1	2.6	5.5	10.2	17.5	20.7	16.9
1999.....	5.1	*	0.8	0.7	0.5	1.2	2.5	5.5	11.0	17.9	21.0	16.2
White, male												
2004.....	5.4	*	0.7	0.8	0.7	1.1	2.7	6.2	12.6	19.8	24.5	20.4
2003.....	5.5	*	0.7	0.8	0.7	1.3	3.1	6.6	12.1	19.6	25.5	19.8
2002.....	5.5	*	0.7	0.9	0.7	1.1	3.1	6.8	12.2	21.1	24.1	20.8
2001.....	5.6	*	0.7	0.7	0.6	1.5	3.2	6.6	12.7	20.1	26.6	20.1
2000.....	5.6	*	0.7	0.8	0.7	1.3	3.3	6.6	12.6	21.4	25.2	23.0
1999.....	5.6	*	0.8	0.7	0.6	1.5	3.0	6.6	13.1	21.9	26.3	21.2

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of brain (C71)												
White, female												
2004.....	4.3	*	0.6	0.7	0.4	0.8	1.8	3.7	7.9	14.2	17.8	15.1
2003.....	4.3	*	0.6	0.8	0.5	0.9	2.2	3.8	8.7	13.6	16.7	12.9
2002.....	4.3	*	0.5	0.8	0.3	0.9	1.9	4.4	8.3	13.6	16.7	14.9
2001.....	4.3	*	0.6	0.8	0.5	0.9	1.9	4.2	8.1	12.8	17.5	15.1
2000.....	4.4	*	0.6	0.7	0.5	0.8	1.8	4.4	8.0	14.2	17.7	14.5
1999.....	4.5	*	0.8	0.7	0.4	1.0	2.1	4.4	9.0	14.5	17.5	14.1
Black, both sexes												
2004.....	1.9	*	*	0.7	0.5	0.7	1.2	2.7	4.7	7.8	9.8	10.4
2003.....	2.1	*	*	0.7	0.5	0.7	1.4	3.0	5.3	8.5	10.1	10.2
2002.....	2.0	*	1.0	0.8	0.4	0.6	1.2	2.6	5.3	8.2	11.3	9.2
2001.....	2.0	*	*	0.6	0.5	0.5	1.0	2.9	5.7	9.1	9.7	11.7
2000.....	2.0	*	*	0.8	0.6	0.6	1.5	2.8	5.5	7.9	11.5	10.0
1999.....	2.0	*	*	0.8	0.6	0.6	1.3	3.1	5.7	8.0	11.1	9.2
Black, male												
2004.....	2.2	*	*	0.7	*	*	1.8	3.8	6.4	7.6	13.6	*
2003.....	2.4	*	*	0.7	0.6	*	2.1	4.0	6.4	11.5	15.2	*
2002.....	2.2	*	*	1.0	*	0.8	1.3	2.5	7.6	9.7	15.5	*
2001.....	2.1	*	*	0.7	*	*	1.2	3.5	7.6	10.1	12.4	*
2000.....	2.3	*	*	0.9	0.7	1.0	2.0	3.7	6.9	9.7	12.1	*
1999.....	2.3	*	*	0.7	0.8	0.8	1.7	4.2	7.0	9.6	12.0	*
Black, female												
2004.....	1.7	*	*	0.7	*	*	0.7	1.7	3.3	8.0	7.6	11.4
2003.....	1.8	*	*	0.6	*	*	0.9	2.2	4.4	6.3	7.2	11.3
2002.....	1.8	*	*	*	*	*	1.1	2.7	3.4	7.1	8.9	10.3
2001.....	1.8	*	*	*	*	*	0.9	2.4	4.1	8.4	8.3	9.7
2000.....	1.8	*	*	0.6	*	*	1.1	2.1	4.4	6.7	11.1	10.3
1999.....	1.8	*	*	0.9	*	*	0.9	2.1	4.6	6.9	10.7	8.8

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Of other parts of central nervous system (C70,C72)

All races 3/, both sexes

2004.....	0.1	*	*	*	*	*	*	0.1	0.2	0.2	0.3	0.5
2003.....	0.1	*	*	*	*	*	*	0.1	0.1	0.2	0.4	0.6
2002.....	0.1	*	*	*	*	*	*	0.1	0.2	0.4	0.4	*
2001.....	0.1	*	*	*	*	*	*	0.1	0.2	0.3	0.5	*
2000.....	0.1	*	*	*	*	*	0.0	0.1	0.2	0.3	0.3	0.6
1999.....	0.1	*	*	*	*	*	0.1	0.1	0.2	0.3	0.5	*

All races 3/, male

2004.....	0.1	*	*	*	*	*	*	*	0.2	*	0.4	*
2003.....	0.1	*	*	*	*	*	*	*	0.2	0.2	*	*
2002.....	0.1	*	*	*	*	*	*	*	0.2	0.5	0.7	*
2001.....	0.1	*	*	*	*	*	*	*	0.2	0.4	0.5	*
2000.....	0.1	*	*	*	*	*	*	0.1	*	0.3	*	*
1999.....	0.1	*	*	*	*	*	0.1	0.1	0.2	0.4	0.5	*

All races 3/, female

2004.....	0.1	*	*	*	*	*	*	*	*	*	0.3	0.7
2003.....	0.1	*	*	*	*	*	*	*	*	0.2	0.5	*
2002.....	0.1	*	*	*	*	*	*	0.1	0.1	0.3	*	*
2001.....	0.1	*	*	*	*	*	*	*	*	0.2	0.4	*
2000.....	0.1	*	*	*	*	*	*	*	0.2	0.3	0.3	*
1999.....	0.1	*	*	*	*	*	*	0.1	0.2	0.2	0.5	*

White, both sexes

2004.....	0.1	*	*	*	*	*	*	0.1	0.2	0.2	0.3	0.6
2003.....	0.1	*	*	*	*	*	*	0.1	0.1	0.2	0.4	0.7
2002.....	0.1	*	*	*	*	*	*	0.1	0.1	0.4	0.4	*
2001.....	0.1	*	*	*	*	*	*	0.1	0.2	0.3	0.5	*
2000.....	0.1	*	*	*	*	*	*	0.1	0.2	0.3	0.3	0.5
1999.....	0.1	*	*	*	*	*	0.1	0.1	0.2	0.3	0.5	*

White, male

2004.....	0.1	*	*	*	*	*	*	*	0.2	*	*	*
2003.....	0.1	*	*	*	*	*	*	*	0.2	*	*	*
2002.....	0.1	*	*	*	*	*	*	*	*	0.5	0.7	*
2001.....	0.1	*	*	*	*	*	*	*	0.2	0.4	0.5	*
2000.....	0.1	*	*	*	*	*	*	0.1	*	0.3	*	*
1999.....	0.1	*	*	*	*	*	0.1	*	*	0.4	0.5	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of thyroid and other endocrine glands (C73-C75)												
All races 3/, both sexes												
2004.....	0.7	*	0.5	0.2	0.1	0.1	0.2	0.4	1.1	2.5	3.9	6.0
2003.....	0.7	*	0.4	0.3	*	0.1	0.2	0.5	1.2	2.2	3.8	5.1
2002.....	0.7	*	0.5	0.2	0.1	0.1	0.2	0.5	1.2	2.3	4.3	5.3
2001.....	0.7	*	0.5	0.2	0.1	0.1	0.2	0.6	1.3	2.6	4.1	5.3
2000.....	0.7	*	0.5	0.2	0.1	0.1	0.2	0.5	1.3	2.6	4.1	6.2
1999.....	0.7	*	0.5	0.2	0.1	0.1	0.2	0.5	1.3	2.3	4.1	5.8
All races 3/, male												
2004.....	0.6	*	0.5	0.3	0.1	*	0.2	0.4	1.2	2.7	3.7	5.5
2003.....	0.6	*	0.5	0.3	*	0.1	0.2	0.6	1.1	2.2	3.6	4.5
2002.....	0.6	*	0.4	0.3	0.1	*	0.2	0.6	1.3	2.3	4.3	4.2
2001.....	0.7	*	0.5	0.2	*	0.1	0.2	0.6	1.4	2.9	3.9	4.5
2000.....	0.7	*	0.5	0.2	*	*	0.2	0.5	1.3	2.7	4.3	5.3
1999.....	0.7	*	0.5	0.3	*	*	0.2	0.6	1.3	2.6	4.2	5.1
All races 3/, female												
2004.....	0.8	*	0.5	0.2	*	*	0.1	0.5	1.1	2.2	4.1	6.3
2003.....	0.7	*	0.4	0.2	*	0.1	0.2	0.5	1.2	2.1	4.0	5.4
2002.....	0.8	*	0.5	0.2	*	0.1	0.2	0.5	1.1	2.3	4.2	5.8
2001.....	0.8	*	0.4	0.2	*	0.1	0.2	0.5	1.2	2.4	4.3	5.7
2000.....	0.8	*	0.5	0.2	*	*	0.2	0.4	1.3	2.4	4.0	6.5
1999.....	0.8	*	0.4	0.2	*	*	0.2	0.5	1.3	2.1	4.0	6.1
White, both sexes												
2004.....	0.8	*	0.5	0.2	0.1	0.1	0.2	0.4	1.2	2.5	4.0	5.9
2003.....	0.7	*	0.4	0.2	*	0.1	0.2	0.5	1.2	2.3	3.8	5.2
2002.....	0.8	*	0.4	0.2	*	0.1	0.2	0.5	1.2	2.3	4.3	5.3
2001.....	0.8	*	0.5	0.2	0.1	0.1	0.2	0.6	1.3	2.8	4.0	5.3
2000.....	0.8	*	0.5	0.2	0.1	0.1	0.2	0.4	1.3	2.5	4.2	6.2
1999.....	0.8	*	0.5	0.2	*	0.1	0.2	0.5	1.3	2.3	4.1	5.8
White, male												
2004.....	0.7	*	0.5	0.3	*	*	0.2	0.4	1.2	2.8	3.8	5.3
2003.....	0.7	*	0.5	0.3	*	*	0.2	0.6	1.2	2.4	3.6	4.6
2002.....	0.7	*	0.4	0.3	*	*	0.2	0.6	1.3	2.3	4.5	4.1
2001.....	0.7	*	0.6	0.2	*	*	0.2	0.6	1.4	3.1	4.0	4.5
2000.....	0.7	*	0.6	0.2	*	*	0.2	0.5	1.3	2.8	4.5	5.6
1999.....	0.7	*	0.6	0.3	*	*	0.2	0.6	1.3	2.6	4.4	4.9

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms of thyroid and other endocrine glands (C73-C75)

White, female

2004.....	0.8	*	0.5	0.2	*	*	0.2	0.5	1.2	2.2	4.2	6.2
2003.....	0.8	*	0.4	0.2	*	*	0.2	0.5	1.2	2.2	3.8	5.5
2002.....	0.8	*	0.4	0.2	*	*	0.2	0.5	1.1	2.3	4.2	5.9
2001.....	0.8	*	0.5	0.2	*	*	0.2	0.5	1.2	2.4	4.1	5.7
2000.....	0.8	*	0.4	0.2	*	*	0.2	0.4	1.2	2.3	3.9	6.5
1999.....	0.8	*	0.4	0.2	*	*	0.2	0.5	1.3	2.0	3.9	6.2

Black, both sexes

2004.....	0.5	*	*	0.3	*	*	*	*	1.1	2.3	2.9	7.0
2003.....	0.4	*	*	0.4	*	*	*	*	0.9	1.6	4.3	*
2002.....	0.5	*	*	*	*	*	*	0.5	1.1	1.8	4.6	*
2001.....	0.6	*	*	*	*	*	*	0.7	1.4	1.8	5.0	*
2000.....	0.6	*	*	*	*	*	*	0.8	1.4	2.7	3.7	*
1999.....	0.6	*	*	0.3	*	*	0.4	*	1.6	2.4	3.3	*

Black, male

2004.....	0.4	*	*	*	*	*	*	*	*	*	*	*
2003.....	0.4	*	*	*	*	*	*	*	*	*	*	*
2002.....	0.5	*	*	*	*	*	*	*	*	*	*	*
2001.....	0.5	*	*	*	*	*	*	*	*	*	*	*
2000.....	0.4	*	*	*	*	*	*	*	*	*	*	*
1999.....	0.5	*	*	*	*	*	*	*	*	*	*	*

Black, female

2004.....	0.6	*	*	*	*	*	*	*	*	2.8	3.3	*
2003.....	0.5	*	*	*	*	*	*	*	*	2.1	4.9	*
2002.....	0.6	*	*	*	*	*	*	*	*	*	5.8	*
2001.....	0.6	*	*	*	*	*	*	1.0	*	2.1	5.9	*
2000.....	0.7	*	*	*	*	*	*	*	*	3.0	4.6	*
1999.....	0.6	*	*	*	*	*	*	*	1.5	2.3	3.8	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other malignant neoplasms of other and unspecified sites (C76-C80,C97)

All races 3/, both sexes

2004.....	13.5	*	*	0.1	0.3	0.6	2.2	8.1	22.3	50.7	94.4	141.2
2003.....	13.8	*	*	0.1	0.2	0.5	2.2	8.7	23.1	51.9	96.4	147.9
2002.....	14.1	*	0.1	0.1	0.3	0.6	2.4	8.7	24.1	53.7	96.7	154.5
2001.....	14.1	*	*	0.1	0.2	0.6	2.4	8.6	24.5	53.9	97.4	155.2
2000.....	14.5	*	0.1	0.1	0.2	0.6	2.5	9.0	25.1	56.4	101.9	159.0
1999.....	15.7	*	*	0.1	0.3	0.7	2.8	9.4	28.3	63.0	109.3	168.4

All races 3/, male

2004.....	14.1	*	*	0.1	0.3	0.6	2.2	9.5	27.0	61.8	117.1	170.3
2003.....	14.3	*	*	*	0.3	0.6	2.2	10.1	28.1	62.6	119.2	181.3
2002.....	14.7	*	*	*	0.4	0.7	2.4	10.4	29.0	66.0	118.5	196.9
2001.....	14.5	*	*	0.1	0.3	0.7	2.5	10.3	29.4	65.1	119.1	196.5
2000.....	14.8	*	*	0.1	0.2	0.6	2.6	10.5	30.0	68.4	124.6	200.1
1999.....	16.2	*	*	0.1	0.3	0.9	2.8	11.1	33.9	76.6	133.5	218.9

All races 3/, female

2004.....	13.0	*	*	*	0.2	0.6	2.1	6.7	17.9	41.3	79.1	128.1
2003.....	13.4	*	*	*	0.2	0.5	2.2	7.4	18.5	43.0	81.2	133.2
2002.....	13.5	*	*	*	0.2	0.6	2.4	7.1	19.6	43.4	82.3	136.0
2001.....	13.6	*	*	*	0.2	0.6	2.3	7.1	19.9	44.6	83.1	137.9
2000.....	14.1	*	*	*	0.2	0.5	2.4	7.5	20.6	46.6	87.0	142.3
1999.....	15.3	*	*	*	0.2	0.6	2.7	7.9	23.1	51.8	93.6	148.1

White, both sexes

2004.....	14.4	*	*	0.1	0.3	0.6	2.1	7.5	21.5	50.2	94.6	143.3
2003.....	14.7	*	*	0.1	0.2	0.5	2.1	8.2	22.3	51.5	97.2	149.3
2002.....	14.9	*	*	0.1	0.3	0.6	2.2	8.2	23.4	53.0	96.4	154.9
2001.....	14.8	*	*	0.1	0.2	0.6	2.3	8.1	23.5	52.7	97.0	157.1
2000.....	15.3	*	*	0.1	0.3	0.6	2.3	8.3	24.3	55.8	101.2	160.3
1999.....	16.6	*	*	0.1	0.3	0.7	2.6	8.8	27.0	62.2	108.9	168.0

White, male

2004.....	14.8	*	*	*	0.3	0.7	2.1	8.6	25.6	60.5	117.7	172.2
2003.....	15.1	*	*	*	0.3	0.6	2.1	9.3	26.8	61.6	119.7	184.1
2002.....	15.5	*	*	*	0.4	0.7	2.1	9.8	27.8	65.4	117.7	198.4
2001.....	15.1	*	*	*	0.3	0.7	2.3	9.4	27.6	63.3	118.2	199.0
2000.....	15.5	*	*	*	0.2	0.6	2.3	9.6	28.3	67.4	122.8	201.3
1999.....	17.0	*	*	*	0.3	0.8	2.6	10.1	32.2	75.2	132.9	218.7

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other malignant neoplasms of other and unspecified sites (C76-C80,C97)

White, female

2004.....	13.9	*	*	*	0.2	0.6	2.1	6.4	17.5	41.4	78.8	130.3
2003.....	14.4	*	*	*	0.2	0.5	2.1	7.1	18.0	42.8	82.0	133.9
2002.....	14.4	*	*	*	0.2	0.5	2.4	6.7	19.2	42.5	82.1	136.0
2001.....	14.6	*	*	*	0.1	0.5	2.2	6.8	19.5	43.8	83.0	139.6
2000.....	15.2	*	*	*	0.3	0.5	2.3	7.1	20.6	46.0	87.0	143.7
1999.....	16.2	*	*	*	0.2	0.6	2.5	7.6	22.2	51.4	93.1	147.7

Black, both sexes

2004.....	11.9	*	*	*	*	0.7	3.3	13.5	33.7	66.0	109.0	135.9
2003.....	12.1	*	*	*	*	0.7	3.4	14.4	35.6	66.4	105.9	150.0
2002.....	12.6	*	*	*	*	0.9	3.7	13.5	36.2	70.3	115.9	168.4
2001.....	12.6	*	*	*	0.3	0.9	3.6	14.6	37.4	73.8	114.5	152.9
2000.....	12.7	*	*	*	*	0.6	4.3	15.2	37.0	71.6	121.8	155.2
1999.....	13.9	*	*	*	*	0.7	4.3	15.7	44.0	79.2	127.5	184.3

Black, male

2004.....	13.4	*	*	*	*	*	3.7	18.3	45.6	90.2	137.3	178.8
2003.....	13.2	*	*	*	*	*	3.6	18.3	46.8	86.8	140.2	174.1
2002.....	13.4	*	*	*	*	0.8	4.3	16.7	47.5	87.2	151.8	212.3
2001.....	14.0	*	*	*	*	0.9	3.9	19.3	50.8	97.3	150.1	208.8
2000.....	14.2	*	*	*	*	*	5.0	19.9	51.5	91.2	166.2	204.1
1999.....	15.0	*	*	*	*	0.8	4.3	20.6	57.9	104.4	158.6	241.4

Black, female

2004.....	10.6	*	*	*	*	0.8	2.9	9.4	24.0	48.7	93.0	118.9
2003.....	11.2	*	*	*	*	0.8	3.3	11.1	26.6	51.9	86.6	140.6
2002.....	11.8	*	*	*	*	1.0	3.1	10.7	27.0	58.3	95.7	151.6
2001.....	11.4	*	*	*	*	0.9	3.3	10.5	26.6	57.2	94.5	132.2
2000.....	11.4	*	*	*	*	*	3.6	11.2	25.4	57.9	96.8	136.9
1999.....	12.9	*	*	*	*	*	4.3	11.6	32.8	61.4	110.0	163.0

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)												
All races 3/, both sexes												
2004.....	18.6	0.6	0.8	0.9	1.5	2.1	3.4	9.0	25.8	68.2	139.8	189.4
2003.....	19.1	0.8	0.9	1.0	1.5	2.2	3.9	9.3	26.8	70.3	143.3	198.0
2002.....	19.5	0.7	0.8	0.9	1.7	2.3	3.9	9.9	28.2	73.9	143.6	197.5
2001.....	19.8	0.7	1.0	0.9	1.8	2.5	4.1	10.4	29.5	74.6	145.2	199.9
2000.....	20.1	1.0	0.9	0.9	1.8	2.3	4.1	10.7	30.0	76.3	149.2	204.6
1999.....	20.2	0.8	0.9	0.9	1.9	2.4	4.2	10.8	31.8	77.8	148.0	199.4
All races 3/, male												
2004.....	20.4	*	0.9	1.0	1.9	2.4	4.2	11.0	32.1	87.4	181.3	254.9
2003.....	21.0	*	1.0	1.1	1.8	2.6	4.7	11.4	33.9	90.4	184.8	268.7
2002.....	21.4	*	0.8	1.0	2.0	2.7	4.6	11.9	35.0	95.2	188.0	269.1
2001.....	21.5	*	1.0	1.0	2.1	2.9	5.1	12.7	36.3	94.7	187.0	275.3
2000.....	21.6	1.1	1.0	1.1	2.3	2.7	4.8	13.0	36.1	96.8	192.6	281.8
1999.....	21.7	*	1.0	1.0	2.2	2.8	5.1	13.2	38.8	97.3	189.0	279.0
All races 3/, female												
2004.....	16.8	*	0.6	0.7	1.2	1.7	2.7	7.0	19.9	52.0	111.8	160.0
2003.....	17.3	1.1	0.8	0.8	1.2	1.7	3.1	7.2	20.1	53.4	115.6	166.8
2002.....	17.7	*	0.7	0.8	1.4	1.9	3.2	7.9	21.9	56.2	114.1	166.5
2001.....	18.1	*	1.0	0.7	1.4	2.1	3.2	8.1	23.1	57.9	117.7	168.4
2000.....	18.6	*	0.9	0.7	1.4	1.9	3.4	8.5	24.4	59.5	120.9	173.1
1999.....	18.7	*	0.8	0.7	1.5	2.0	3.4	8.5	25.3	61.9	121.3	167.3
White, both sexes												
2004.....	20.2	0.6	0.8	0.9	1.6	2.0	3.4	8.7	25.7	68.9	143.4	194.9
2003.....	20.9	0.9	0.9	1.0	1.5	2.1	3.8	9.0	27.0	71.5	146.9	204.5
2002.....	21.3	*	0.8	1.0	1.8	2.2	3.9	9.6	28.0	75.4	147.6	204.0
2001.....	21.5	0.7	1.1	0.9	1.8	2.4	4.0	10.2	29.2	75.8	148.7	205.9
2000.....	21.9	1.2	1.0	1.0	1.8	2.2	4.0	10.4	30.1	78.0	153.3	208.7
1999.....	21.9	0.8	1.0	0.9	1.8	2.3	4.2	10.7	31.5	79.3	151.3	204.1
White, male												
2004.....	22.2	*	0.8	1.1	1.9	2.4	4.1	10.7	32.2	88.2	186.3	264.0
2003.....	23.1	*	1.1	1.1	1.8	2.6	4.6	11.1	34.2	92.8	191.0	279.5
2002.....	23.4	*	0.9	1.0	2.1	2.6	4.5	11.6	34.8	97.3	193.7	279.4
2001.....	23.3	*	1.1	1.0	2.1	2.8	4.8	12.6	35.9	96.3	191.8	284.8
2000.....	23.6	*	1.0	1.1	2.2	2.6	4.8	12.6	36.1	98.7	198.2	290.2
1999.....	23.6	*	1.0	1.0	2.2	2.6	5.0	13.0	38.6	99.1	193.4	285.8

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)												
White, female												
2004.....	18.3	*	0.7	0.7	1.2	1.6	2.6	6.8	19.5	52.4	114.2	163.8
2003.....	18.8	1.4	0.8	0.8	1.2	1.6	3.0	7.0	20.1	53.4	117.2	171.3
2002.....	19.2	*	0.8	0.9	1.4	1.8	3.2	7.5	21.5	56.8	116.7	171.2
2001.....	19.7	*	1.1	0.8	1.4	2.0	3.2	7.9	22.8	58.5	120.1	172.8
2000.....	20.3	*	1.0	0.8	1.3	1.8	3.2	8.3	24.4	60.6	123.8	175.5
1999.....	20.4	*	0.9	0.7	1.5	2.0	3.3	8.4	24.9	62.8	123.7	171.3
Black, both sexes												
2004.....	13.6	*	*	0.7	1.6	3.0	4.5	12.6	32.4	75.1	123.4	154.0
2003.....	13.4	*	*	0.8	1.4	2.8	5.2	13.0	31.1	73.1	126.1	150.5
2002.....	13.6	*	*	0.8	1.5	2.9	4.8	13.6	35.9	76.0	122.3	152.3
2001.....	14.1	*	*	1.0	1.8	3.4	5.6	13.2	36.7	78.0	126.9	155.7
2000.....	13.9	*	0.8	0.6	2.2	2.9	5.3	14.7	35.8	75.0	121.0	175.5
1999.....	14.1	*	*	0.9	2.3	2.9	5.5	13.8	38.9	76.2	126.8	162.0
Black, male												
2004.....	14.4	*	*	0.8	1.9	3.3	5.2	15.8	39.7	98.5	150.6	193.7
2003.....	13.9	*	*	0.9	1.6	3.4	6.4	16.6	40.0	87.0	148.9	184.5
2002.....	14.6	*	*	1.0	1.4	3.6	6.2	16.1	44.4	97.8	152.6	202.6
2001.....	15.1	*	*	1.3	2.0	3.9	7.3	16.0	46.0	99.2	159.5	206.5
2000.....	14.7	*	*	0.8	2.7	3.5	5.7	17.9	44.5	96.0	155.0	224.9
1999.....	15.1	*	*	1.1	2.6	3.8	6.8	17.3	47.4	95.4	161.1	220.3
Black, female												
2004.....	12.8	*	*	0.7	1.3	2.7	3.9	9.8	26.6	58.3	108.1	138.2
2003.....	12.9	*	*	0.8	1.1	2.4	4.2	9.9	23.8	63.3	113.2	137.3
2002.....	12.8	*	*	0.6	1.5	2.2	3.5	11.4	29.1	60.5	105.1	133.1
2001.....	13.1	*	*	0.6	1.6	2.9	4.0	10.8	29.2	63.0	108.6	136.8
2000.....	13.1	*	*	*	1.8	2.4	5.0	12.0	28.8	60.2	101.8	157.1
1999.....	13.2	*	*	0.7	2.0	2.1	4.4	10.8	32.0	62.6	107.6	140.2

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hodgkin's disease (C81)												
All races 3/, both sexes												
2004.....	0.4	*	*	*	0.2	0.3	0.3	0.4	0.7	1.2	1.9	2.2
2003.....	0.5	*	*	*	0.2	0.4	0.3	0.4	0.7	1.2	2.2	2.4
2002.....	0.5	*	*	*	0.2	0.4	0.4	0.4	0.6	1.3	1.9	2.7
2001.....	0.5	*	*	*	0.2	0.4	0.4	0.5	0.6	1.2	2.2	2.2
2000.....	0.5	*	*	*	0.2	0.4	0.3	0.4	0.7	1.3	2.1	2.5
1999.....	0.5	*	*	*	0.2	0.4	0.4	0.5	0.7	1.5	2.1	2.4
All races 3/, male												
2004.....	0.5	*	*	*	0.2	0.3	0.3	0.5	0.9	1.6	2.5	3.2
2003.....	0.5	*	*	*	0.1	0.4	0.4	0.5	0.9	1.5	2.6	2.6
2002.....	0.5	*	*	*	0.2	0.4	0.4	0.5	0.8	1.8	2.5	3.6
2001.....	0.5	*	*	*	0.2	0.4	0.5	0.6	0.7	1.6	2.6	2.6
2000.....	0.5	*	*	*	0.2	0.4	0.4	0.5	0.7	1.6	2.5	3.7
1999.....	0.6	*	*	*	0.3	0.4	0.5	0.5	1.0	1.9	2.5	3.8
All races 3/, female												
2004.....	0.4	*	*	*	0.1	0.3	0.2	0.3	0.5	0.9	1.5	1.7
2003.....	0.4	*	*	*	0.2	0.4	0.3	0.3	0.5	0.9	1.9	2.3
2002.....	0.4	*	*	*	0.2	0.4	0.3	0.4	0.4	0.9	1.6	2.3
2001.....	0.4	*	*	*	0.2	0.4	0.3	0.4	0.4	0.8	2.0	2.0
2000.....	0.4	*	*	*	0.2	0.3	0.3	0.3	0.6	1.0	1.8	2.0
1999.....	0.4	*	*	*	0.2	0.3	0.3	0.4	0.5	1.2	1.9	1.8
White, both sexes												
2004.....	0.5	*	*	*	0.2	0.3	0.3	0.4	0.7	1.3	2.0	2.3
2003.....	0.5	*	*	*	0.2	0.4	0.4	0.4	0.7	1.2	2.2	2.5
2002.....	0.5	*	*	*	0.2	0.4	0.4	0.5	0.6	1.3	2.1	2.9
2001.....	0.5	*	*	*	0.2	0.4	0.4	0.5	0.6	1.3	2.3	2.4
2000.....	0.5	*	*	*	0.2	0.4	0.3	0.4	0.7	1.3	2.2	2.6
1999.....	0.6	*	*	*	0.2	0.4	0.4	0.5	0.8	1.6	2.3	2.4
White, male												
2004.....	0.5	*	*	*	0.2	0.3	0.3	0.5	1.0	1.7	2.6	3.5
2003.....	0.5	*	*	*	0.1	0.5	0.5	0.5	0.9	1.5	2.7	2.7
2002.....	0.6	*	*	*	0.2	0.4	0.4	0.5	0.8	1.9	2.6	3.9
2001.....	0.6	*	*	*	0.2	0.4	0.5	0.6	0.7	1.7	2.7	2.7
2000.....	0.5	*	*	*	0.2	0.4	0.4	0.6	0.7	1.7	2.7	3.9
1999.....	0.6	*	*	*	0.3	0.4	0.5	0.5	1.1	2.0	2.7	3.8

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Of other lymphoid, hematopoietic and related tissue (C82-C85,C96)

All races 3/, both sexes

2004.....	7.2	*	*	0.1	0.3	0.6	1.2	3.7	9.9	26.2	54.8	75.3
2003.....	7.4	*	*	0.1	0.3	0.6	1.4	3.8	10.8	26.9	56.2	79.0
2002.....	7.6	*	*	0.1	0.4	0.7	1.5	4.0	11.5	28.9	57.4	77.6
2001.....	7.9	*	*	0.1	0.4	0.8	1.7	4.4	12.3	29.4	58.8	80.1
2000.....	8.1	*	*	0.1	0.5	0.7	1.7	4.7	13.1	30.6	61.2	80.7
1999.....	8.2	*	*	0.1	0.4	0.7	1.8	4.8	13.3	31.8	62.0	78.0

All races 3/, male

2004.....	7.7	*	*	0.1	0.4	0.8	1.7	4.8	12.4	32.5	67.5	96.3
2003.....	8.0	*	*	0.1	0.4	0.8	1.8	4.9	13.8	34.0	69.3	104.0
2002.....	8.2	*	*	0.2	0.5	0.9	2.0	5.0	14.4	36.2	71.0	100.2
2001.....	8.4	*	*	0.1	0.5	0.9	2.2	5.8	15.1	36.7	71.6	107.1
2000.....	8.6	*	*	0.1	0.6	0.9	2.2	6.0	16.0	37.5	74.8	106.6
1999.....	8.6	*	*	0.1	0.5	0.9	2.3	6.2	16.3	38.4	75.0	104.7

All races 3/, female

2004.....	6.7	*	*	*	0.3	0.5	0.8	2.6	7.6	20.8	46.2	65.8
2003.....	6.8	*	*	0.1	0.3	0.4	1.0	2.8	8.0	21.0	47.4	68.0
2002.....	7.1	*	*	*	0.2	0.5	1.1	3.0	8.8	22.9	48.4	67.8
2001.....	7.3	*	*	*	0.3	0.6	1.1	3.1	9.7	23.3	50.4	68.8
2000.....	7.6	*	*	*	0.3	0.6	1.2	3.3	10.5	24.8	52.3	70.2
1999.....	7.8	*	*	*	0.3	0.6	1.2	3.4	10.6	26.5	53.6	67.3

White, both sexes

2004.....	8.0	*	*	0.1	0.3	0.6	1.2	3.7	10.2	27.5	57.8	79.0
2003.....	8.3	*	*	0.1	0.3	0.6	1.4	3.8	11.2	28.3	59.2	83.6
2002.....	8.5	*	*	0.1	0.3	0.6	1.5	3.9	11.9	30.3	60.4	82.4
2001.....	8.8	*	*	0.1	0.3	0.7	1.6	4.5	12.6	30.8	61.8	83.8
2000.....	9.1	*	*	0.1	0.4	0.7	1.7	4.7	13.8	32.2	64.5	84.8
1999.....	9.2	*	*	0.1	0.4	0.7	1.7	4.9	13.9	33.6	65.1	81.5

White, male

2004.....	8.5	*	*	0.1	0.4	0.8	1.7	4.8	12.7	33.9	71.0	101.7
2003.....	9.0	*	*	*	0.4	0.7	1.8	4.8	14.4	36.0	73.1	110.1
2002.....	9.1	*	*	0.2	0.5	0.8	1.9	5.0	14.8	37.9	74.6	106.1
2001.....	9.3	*	*	0.1	0.4	0.9	2.2	5.8	15.4	38.3	75.0	111.9
2000.....	9.6	*	*	0.2	0.6	0.8	2.2	6.0	16.5	39.3	78.8	112.5
1999.....	9.6	*	*	0.1	0.5	0.8	2.2	6.3	16.7	40.3	78.3	108.3

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of other lymphoid, hematopoietic and related tissue (C82-C85,C96)												
White, female												
2004.....	7.5	*	*	*	0.3	0.4	0.8	2.6	7.9	22.0	48.8	68.8
2003.....	7.7	*	*	*	0.3	0.4	1.0	2.8	8.3	21.7	49.8	71.8
2002.....	8.0	*	*	*	0.2	0.5	1.1	2.9	9.1	23.9	50.9	72.1
2001.....	8.3	*	*	*	0.3	0.5	1.1	3.1	10.1	24.6	53.0	72.1
2000.....	8.7	*	*	*	0.2	0.6	1.1	3.4	11.2	26.2	55.1	73.5
1999.....	8.8	*	*	*	0.3	0.5	1.2	3.5	11.2	28.0	56.5	70.7
Black, both sexes												
2004.....	3.7	*	*	*	0.5	1.1	1.5	4.8	9.1	18.5	28.2	39.2
2003.....	3.6	*	*	*	0.4	1.1	1.9	4.8	8.9	18.3	28.3	31.5
2002.....	3.9	*	*	*	0.4	1.2	2.0	4.7	10.8	21.3	30.5	31.2
2001.....	4.1	*	*	*	0.5	1.2	2.2	5.0	11.0	21.1	30.9	39.6
2000.....	3.8	*	*	*	0.6	1.0	2.1	5.0	10.6	19.3	27.2	36.9
1999.....	4.0	*	*	*	0.5	1.1	2.4	4.8	11.1	20.1	31.9	40.8
Black, male												
2004.....	4.0	*	*	*	0.6	1.4	1.9	6.5	11.8	23.4	32.9	42.7
2003.....	3.8	*	*	*	*	1.3	2.3	6.5	11.2	20.4	31.0	34.4
2002.....	4.3	*	*	*	*	1.5	2.9	5.5	13.7	28.0	35.6	42.9
2001.....	4.6	*	*	*	0.7	1.3	3.3	6.7	13.7	27.8	37.4	55.6
2000.....	4.2	*	*	*	0.7	1.2	2.6	7.0	14.6	24.9	31.5	46.1
1999.....	4.6	*	*	*	*	1.5	3.2	6.6	14.7	25.6	40.9	59.8
Black, female												
2004.....	3.4	*	*	*	*	0.8	1.2	3.3	6.9	14.9	25.6	37.8
2003.....	3.4	*	*	*	*	0.8	1.5	3.3	7.1	16.7	26.8	30.4
2002.....	3.6	*	*	*	*	0.8	1.3	4.0	8.5	16.5	27.6	26.7
2001.....	3.6	*	*	*	*	1.1	1.3	3.6	8.8	16.5	27.3	33.7
2000.....	3.4	*	*	*	*	0.9	1.6	3.3	7.3	15.4	24.7	33.5
1999.....	3.5	*	*	*	*	*	1.7	3.2	8.2	16.3	26.9	33.7

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Non-Hodgkin's lymphoma (C82-C85)												
All races 3/, both sexes												
2004.....	7.1	*	*	0.1	0.3	0.6	1.2	3.7	9.9	26.1	54.6	75.0
2003.....	7.4	*	*	0.1	0.3	0.6	1.4	3.8	10.8	26.8	56.1	78.8
2002.....	7.6	*	*	0.1	0.3	0.7	1.5	3.9	11.5	28.8	57.2	77.3
2001.....	7.8	*	*	0.1	0.4	0.8	1.7	4.4	12.2	29.3	58.6	79.6
2000.....	8.1	*	*	0.1	0.5	0.7	1.7	4.6	13.1	30.4	61.0	80.2
1999.....	8.2	*	*	0.1	0.4	0.7	1.8	4.8	13.3	31.7	61.8	77.7
All races 3/, male												
2004.....	7.6	*	*	0.1	0.4	0.8	1.7	4.8	12.3	32.5	67.3	95.8
2003.....	8.0	*	*	0.1	0.4	0.8	1.8	4.8	13.8	33.9	69.1	103.6
2002.....	8.1	*	*	0.2	0.4	0.9	2.0	5.0	14.4	36.0	70.7	99.7
2001.....	8.4	*	*	0.1	0.5	0.9	2.2	5.8	15.0	36.5	71.3	106.6
2000.....	8.6	*	*	0.1	0.6	0.9	2.1	6.0	15.9	37.4	74.5	105.9
1999.....	8.6	*	*	0.1	0.5	0.9	2.3	6.2	16.2	38.3	74.7	104.4
All races 3/, female												
2004.....	6.6	*	*	*	0.3	0.5	0.8	2.6	7.6	20.7	46.0	65.7
2003.....	6.8	*	*	*	0.3	0.4	1.0	2.8	8.0	20.9	47.4	67.8
2002.....	7.1	*	*	*	0.2	0.5	1.1	3.0	8.8	22.8	48.2	67.6
2001.....	7.3	*	*	*	0.3	0.6	1.1	3.1	9.7	23.3	50.2	68.3
2000.....	7.6	*	*	*	0.3	0.6	1.2	3.3	10.5	24.7	52.1	69.8
1999.....	7.7	*	*	*	0.3	0.6	1.2	3.4	10.6	26.4	53.4	66.9
White, both sexes												
2004.....	8.0	*	*	0.1	0.3	0.6	1.2	3.6	10.2	27.4	57.6	78.8
2003.....	8.3	*	*	0.1	0.3	0.6	1.4	3.8	11.2	28.2	59.1	83.3
2002.....	8.5	*	*	0.1	0.3	0.6	1.5	3.9	11.9	30.2	60.3	82.1
2001.....	8.8	*	*	0.1	0.3	0.7	1.6	4.5	12.6	30.7	61.6	83.4
2000.....	9.1	*	*	0.1	0.4	0.7	1.7	4.7	13.7	32.1	64.3	84.3
1999.....	9.2	*	*	0.1	0.4	0.7	1.7	4.9	13.8	33.5	64.9	81.2
White, male												
2004.....	8.5	*	*	0.1	0.4	0.8	1.7	4.7	12.6	33.9	70.8	101.2
2003.....	8.9	*	*	*	0.4	0.7	1.8	4.8	14.4	35.9	73.0	109.7
2002.....	9.1	*	*	0.2	0.5	0.8	1.9	5.0	14.8	37.7	74.4	105.6
2001.....	9.3	*	*	0.1	0.4	0.9	2.1	5.8	15.3	38.0	74.7	111.5
2000.....	9.6	*	*	0.1	0.6	0.8	2.2	6.0	16.5	39.2	78.4	111.8
1999.....	9.6	*	*	0.1	0.5	0.8	2.2	6.3	16.6	40.2	78.0	108.0

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Non-Hodgkin's lymphoma (C82-C85)												
White, female												
2004.....	7.5	*	*	*	0.3	0.4	0.8	2.6	7.9	21.9	48.6	68.7
2003.....	7.7	*	*	*	0.3	0.4	1.0	2.8	8.3	21.7	49.7	71.6
2002.....	8.0	*	*	*	0.2	0.5	1.1	2.9	9.1	23.9	50.8	71.8
2001.....	8.2	*	*	*	0.3	0.5	1.1	3.1	10.1	24.5	52.9	71.6
2000.....	8.6	*	*	*	0.2	0.6	1.1	3.4	11.1	26.1	55.0	73.1
1999.....	8.8	*	*	*	0.3	0.5	1.2	3.5	11.1	27.9	56.3	70.4
Black, both sexes												
2004.....	3.7	*	*	*	0.5	1.1	1.5	4.8	9.0	18.5	27.9	38.6
2003.....	3.6	*	*	*	0.4	1.1	1.9	4.8	8.9	18.2	28.3	31.2
2002.....	3.9	*	*	*	0.4	1.2	2.0	4.7	10.8	21.2	30.2	30.6
2001.....	4.0	*	*	*	0.5	1.2	2.2	5.0	10.9	21.0	30.5	39.3
2000.....	3.7	*	*	*	0.6	1.0	2.1	5.0	10.5	19.3	26.8	36.3
1999.....	4.0	*	*	*	0.5	1.1	2.4	4.7	11.1	20.1	31.8	40.5
Black, male												
2004.....	4.0	*	*	*	0.6	1.4	1.9	6.5	11.8	23.4	32.9	41.7
2003.....	3.8	*	*	*	*	1.3	2.3	6.5	11.2	20.4	31.0	34.4
2002.....	4.3	*	*	*	*	1.5	2.9	5.5	13.7	27.8	35.3	42.9
2001.....	4.5	*	*	*	0.7	1.3	3.2	6.7	13.6	27.7	36.8	54.5
2000.....	4.2	*	*	*	*	1.2	2.6	6.9	14.5	24.9	31.2	43.8
1999.....	4.6	*	*	*	*	1.5	3.2	6.6	14.6	25.6	40.6	59.8
Black, female												
2004.....	3.3	*	*	*	*	0.8	1.2	3.3	6.8	14.9	25.2	37.4
2003.....	3.4	*	*	*	*	0.8	1.5	3.3	7.1	16.6	26.8	30.0
2002.....	3.5	*	*	*	*	0.8	1.3	4.0	8.5	16.5	27.3	25.9
2001.....	3.6	*	*	*	*	1.1	1.3	3.6	8.7	16.3	26.9	33.7
2000.....	3.3	*	*	*	*	0.9	1.6	3.3	7.3	15.4	24.4	33.5
1999.....	3.5	*	*	*	*	*	1.7	3.2	8.2	16.3	26.9	33.3

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant immunoproliferative diseases (C88)												
All races 3/, both sexes												
2004.....	0.1	*	*	*	*	*	*	*	0.1	0.4	1.0	1.5
2003.....	0.1	*	*	*	*	*	*	*	0.1	0.5	1.2	2.2
2002.....	0.1	*	*	*	*	*	*	*	0.2	0.4	1.4	2.0
2001.....	0.1	*	*	*	*	*	*	*	0.2	0.5	1.1	2.0
2000.....	0.1	*	*	*	*	*	*	*	0.1	0.4	1.2	2.1
1999.....	0.2	*	*	*	*	*	*	*	0.2	0.5	1.4	2.7
All races 3/, male												
2004.....	0.1	*	*	*	*	*	*	*	*	0.6	1.3	2.5
2003.....	0.2	*	*	*	*	*	*	*	0.1	0.7	1.7	3.6
2002.....	0.2	*	*	*	*	*	*	*	0.2	0.6	2.2	2.8
2001.....	0.1	*	*	*	*	*	*	*	0.2	0.6	1.6	2.9
2000.....	0.1	*	*	*	*	*	*	*	0.2	0.6	1.7	3.4
1999.....	0.2	*	*	*	*	*	*	*	0.3	0.8	1.9	3.9
All races 3/, female												
2004.....	0.1	*	*	*	*	*	*	*	*	0.3	0.7	1.1
2003.....	0.1	*	*	*	*	*	*	*	*	0.3	0.8	1.6
2002.....	0.1	*	*	*	*	*	*	*	*	0.2	0.8	1.6
2001.....	0.1	*	*	*	*	*	*	*	*	0.4	0.9	1.7
2000.....	0.1	*	*	*	*	*	*	*	*	0.3	0.9	1.6
1999.....	0.1	*	*	*	*	*	*	*	0.2	0.3	1.1	2.2
White, both sexes												
2004.....	0.1	*	*	*	*	*	*	*	0.1	0.4	1.1	1.7
2003.....	0.2	*	*	*	*	*	*	*	0.1	0.5	1.2	2.4
2002.....	0.2	*	*	*	*	*	*	*	0.2	0.4	1.5	2.1
2001.....	0.2	*	*	*	*	*	*	*	0.2	0.5	1.2	2.2
2000.....	0.2	*	*	*	*	*	*	*	0.2	0.5	1.3	2.2
1999.....	0.2	*	*	*	*	*	*	*	0.2	0.6	1.4	2.9
White, male												
2004.....	0.1	*	*	*	*	*	*	*	*	0.6	1.4	2.8
2003.....	0.2	*	*	*	*	*	*	*	*	0.8	1.9	4.0
2002.....	0.2	*	*	*	*	*	*	*	0.2	0.7	2.3	2.9
2001.....	0.2	*	*	*	*	*	*	*	0.2	0.7	1.7	3.1
2000.....	0.2	*	*	*	*	*	*	*	0.2	0.7	1.8	3.5
1999.....	0.2	*	*	*	*	*	*	*	0.3	0.8	1.9	4.3

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Multiple myeloma and malignant plasma cell neoplasms (C90)

All races 3/, both sexes

2004.....	3.6	*	*	*	*	*	0.3	1.6	5.8	15.0	29.7	31.6
2003.....	3.7	*	*	*	*	*	0.3	1.7	5.8	16.0	29.9	34.1
2002.....	3.8	*	*	*	*	*	0.3	1.8	6.1	17.0	29.8	34.7
2001.....	3.8	*	*	*	*	*	0.3	1.7	6.4	16.9	29.8	34.2
2000.....	3.8	*	*	*	*	*	0.3	1.8	6.2	16.9	30.3	35.7
1999.....	3.8	*	*	*	*	0.1	0.3	1.8	6.9	16.7	29.6	33.9

All races 3/, male

2004.....	3.8	*	*	*	*	*	0.3	1.8	6.8	18.3	36.8	42.1
2003.....	3.9	*	*	*	*	*	0.4	2.1	7.3	19.1	36.0	46.8
2002.....	4.0	*	*	*	*	*	0.3	2.2	7.3	20.4	37.5	48.0
2001.....	4.0	*	*	*	*	*	0.4	2.1	7.6	20.2	37.1	45.1
2000.....	3.9	*	*	*	*	*	0.4	2.1	6.8	19.8	37.9	48.3
1999.....	3.9	*	*	*	*	*	0.4	2.1	8.2	19.4	36.5	47.4

All races 3/, female

2004.....	3.4	*	*	*	*	*	0.3	1.3	4.9	12.2	24.9	26.9
2003.....	3.6	*	*	*	*	*	0.2	1.4	4.5	13.5	25.9	28.5
2002.....	3.6	*	*	*	*	*	0.3	1.4	5.0	14.1	24.7	29.0
2001.....	3.6	*	*	*	*	*	0.3	1.3	5.4	14.2	25.0	29.7
2000.....	3.7	*	*	*	*	*	0.3	1.6	5.6	14.4	25.4	30.6
1999.....	3.7	*	*	*	*	*	0.3	1.5	5.7	14.5	25.2	28.4

White, both sexes

2004.....	3.6	*	*	*	*	*	0.3	1.4	5.2	13.7	28.3	30.3
2003.....	3.7	*	*	*	*	*	0.3	1.4	5.3	14.9	28.3	32.6
2002.....	3.9	*	*	*	*	*	0.2	1.5	5.5	15.8	28.7	33.6
2001.....	3.8	*	*	*	*	*	0.3	1.5	5.6	15.7	28.3	32.8
2000.....	3.8	*	*	*	*	*	0.3	1.6	5.4	15.7	28.7	33.0
1999.....	3.8	*	*	*	*	*	0.3	1.5	5.9	15.5	28.0	31.9

White, male

2004.....	3.9	*	*	*	*	*	0.3	1.6	6.3	17.0	35.4	40.7
2003.....	4.0	*	*	*	*	*	0.4	1.7	6.6	18.1	34.6	45.8
2002.....	4.1	*	*	*	*	*	0.3	1.9	6.6	19.4	36.7	47.3
2001.....	4.0	*	*	*	*	*	0.4	1.8	6.6	19.0	35.7	44.0
2000.....	3.9	*	*	*	*	*	0.3	1.8	6.1	18.6	36.4	46.7
1999.....	3.9	*	*	*	*	*	0.3	1.9	7.3	18.1	34.9	45.4

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Multiple myeloma and malignant plasma cell neoplasms (C90)

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
White, female												
2004.....	3.4	*	*	*	*	*	0.2	1.1	4.3	10.8	23.4	25.6
2003.....	3.5	*	*	*	*	*	0.2	1.1	4.0	12.1	24.0	26.7
2002.....	3.6	*	*	*	*	*	0.2	1.1	4.3	12.8	23.4	27.7
2001.....	3.6	*	*	*	*	*	0.2	1.1	4.6	13.0	23.4	28.1
2000.....	3.7	*	*	*	*	*	0.2	1.3	4.8	13.3	23.7	27.4
1999.....	3.6	*	*	*	*	*	0.2	1.2	4.6	13.3	23.4	26.5
Black, both sexes												
2004.....	4.7	*	*	*	*	*	0.7	3.4	12.3	30.3	52.6	53.9
2003.....	4.8	*	*	*	*	*	0.7	4.1	11.7	30.6	54.6	59.8
2002.....	4.6	*	*	*	*	*	0.7	4.1	13.0	31.3	48.1	54.3
2001.....	4.8	*	*	*	*	*	0.9	3.6	14.9	31.1	52.8	57.1
2000.....	5.0	*	*	*	*	*	0.8	4.4	13.8	31.9	54.4	70.1
1999.....	4.9	*	*	*	*	*	0.9	4.2	16.3	30.1	52.4	62.5
Black, male												
2004.....	4.6	*	*	*	*	*	0.7	3.9	14.1	36.3	62.9	71.5
2003.....	4.7	*	*	*	*	*	0.9	5.1	15.2	33.5	62.9	73.0
2002.....	4.6	*	*	*	*	*	0.8	4.6	15.5	36.7	58.2	70.8
2001.....	4.8	*	*	*	*	*	1.0	4.3	18.2	36.3	65.3	69.2
2000.....	4.8	*	*	*	*	*	*	4.9	15.0	38.5	66.6	76.1
1999.....	4.8	*	*	*	*	*	1.1	4.3	18.1	34.9	63.6	80.9
Black, female												
2004.....	4.7	*	*	*	*	*	0.7	3.1	11.0	26.0	46.7	46.9
2003.....	4.9	*	*	*	*	*	*	3.3	8.9	28.5	49.9	54.7
2002.....	4.7	*	*	*	*	*	*	3.7	11.0	27.6	42.4	48.1
2001.....	4.8	*	*	*	*	*	0.8	3.0	12.3	27.4	45.8	52.6
2000.....	5.2	*	*	*	*	*	0.9	4.0	12.9	27.3	47.6	67.8
1999.....	5.1	*	*	*	*	*	*	4.1	14.9	26.8	46.1	55.6

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Leukemia (C91-C95)												
All races 3/, both sexes												
2004.....	7.3	0.6	0.8	0.8	1.0	1.0	1.6	3.3	9.2	25.4	52.4	78.8
2003.....	7.4	0.7	0.8	0.8	1.0	1.1	1.8	3.3	9.3	25.7	53.9	80.3
2002.....	7.5	0.5	0.7	0.8	1.2	1.2	1.7	3.6	9.8	26.4	53.1	80.5
2001.....	7.5	0.6	1.0	0.8	1.2	1.3	1.7	3.8	10.0	26.7	53.3	81.4
2000.....	7.6	1.0	0.9	0.8	1.2	1.2	1.8	3.7	9.9	27.2	54.5	83.6
1999.....	7.5	0.7	0.8	0.7	1.2	1.2	1.8	3.7	10.6	27.2	52.8	82.4
All races 3/, male												
2004.....	8.3	*	0.9	0.9	1.3	1.2	1.8	3.8	11.9	34.4	73.2	110.7
2003.....	8.4	*	0.9	0.9	1.2	1.4	2.0	4.0	11.9	35.1	75.3	111.8
2002.....	8.5	*	0.7	0.8	1.4	1.4	1.9	4.2	12.2	36.3	74.9	114.5
2001.....	8.5	*	1.0	0.9	1.4	1.5	1.9	4.2	12.8	35.7	74.2	117.5
2000.....	8.5	1.1	0.9	0.9	1.5	1.4	1.9	4.3	12.5	37.2	75.7	119.8
1999.....	8.4	*	0.8	0.8	1.5	1.4	1.9	4.3	13.1	36.8	73.2	119.2
All races 3/, female												
2004.....	6.3	*	0.6	0.7	0.8	0.9	1.4	2.8	6.8	17.8	38.4	64.5
2003.....	6.4	1.1	0.8	0.7	0.7	0.8	1.6	2.7	7.0	17.8	39.6	66.4
2002.....	6.5	*	0.6	0.7	1.0	1.0	1.5	3.1	7.6	18.1	38.7	65.7
2001.....	6.6	*	1.0	0.7	1.0	1.1	1.5	3.3	7.5	19.2	39.5	66.3
2000.....	6.7	*	0.9	0.6	0.9	1.0	1.7	3.2	7.6	19.0	40.6	68.8
1999.....	6.7	*	0.7	0.7	1.0	1.1	1.6	3.2	8.3	19.3	39.6	67.6
White, both sexes												
2004.....	8.0	*	0.7	0.8	1.1	1.0	1.6	3.3	9.4	26.0	54.2	81.6
2003.....	8.2	0.9	0.9	0.9	1.0	1.1	1.8	3.4	9.6	26.6	56.0	83.4
2002.....	8.2	*	0.7	0.8	1.2	1.2	1.8	3.6	9.9	27.5	54.9	83.0
2001.....	8.3	0.6	1.1	0.8	1.2	1.3	1.7	3.7	10.2	27.5	55.1	84.7
2000.....	8.3	1.2	1.0	0.8	1.2	1.1	1.8	3.7	10.0	28.3	56.6	86.0
1999.....	8.3	0.8	0.9	0.7	1.2	1.2	1.8	3.7	10.8	28.0	54.5	85.4
White, male												
2004.....	9.1	*	0.8	0.9	1.3	1.2	1.8	3.8	12.2	35.0	75.8	115.3
2003.....	9.4	*	1.0	1.0	1.2	1.3	1.9	4.1	12.2	36.4	78.7	116.9
2002.....	9.3	*	0.8	0.8	1.5	1.4	2.0	4.1	12.3	37.5	77.4	119.1
2001.....	9.3	*	1.1	0.8	1.5	1.5	1.9	4.3	13.0	36.7	76.8	123.1
2000.....	9.4	*	0.9	0.9	1.4	1.3	1.9	4.2	12.6	38.5	78.5	123.6
1999.....	9.2	*	0.9	0.8	1.5	1.3	1.9	4.2	13.3	37.9	75.5	124.1

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Leukemia (C91-C95)												
White, female												
2004.....	6.9	*	0.7	0.7	0.8	0.9	1.4	2.8	6.8	18.4	39.5	66.4
2003.....	7.0	1.4	0.8	0.7	0.7	0.9	1.6	2.7	7.1	18.2	40.6	68.6
2002.....	7.1	*	0.6	0.8	1.0	0.9	1.6	3.1	7.6	18.9	39.8	67.3
2001.....	7.3	*	1.1	0.7	1.0	1.1	1.6	3.2	7.6	19.7	40.8	68.6
2000.....	7.4	*	1.0	0.7	0.9	0.9	1.6	3.2	7.7	19.8	42.1	70.8
1999.....	7.3	*	0.8	0.7	1.0	1.2	1.6	3.2	8.4	19.8	40.6	69.9
Black, both sexes												
2004.....	4.8	*	*	0.6	1.0	1.3	2.0	4.0	10.3	25.2	41.1	60.1
2003.....	4.6	*	*	0.7	0.9	1.2	2.3	3.7	9.6	23.1	41.6	59.2
2002.....	4.7	*	*	0.6	0.9	1.2	1.6	4.3	11.2	22.3	42.1	64.7
2001.....	4.7	*	*	0.9	1.1	1.4	2.1	4.2	9.8	24.9	41.6	57.1
2000.....	4.8	*	*	0.5	1.4	1.6	2.0	4.8	10.7	22.7	38.2	66.0
1999.....	4.7	*	*	0.8	1.4	1.2	1.9	4.3	10.6	24.8	41.2	55.8
Black, male												
2004.....	5.3	*	*	0.7	1.1	1.4	2.2	4.9	12.8	37.0	52.3	79.5
2003.....	5.0	*	*	0.7	1.1	1.6	2.8	4.5	12.3	31.1	53.3	77.2
2002.....	5.1	*	*	0.7	1.0	1.4	1.8	5.2	13.7	31.5	56.8	86.8
2001.....	5.2	*	*	1.1	1.2	1.7	2.5	4.5	13.0	34.0	55.0	77.2
2000.....	5.3	*	*	0.6	1.9	1.9	2.0	5.5	13.8	31.6	54.8	98.0
1999.....	5.2	*	*	1.0	1.6	1.5	2.0	5.5	13.4	32.9	54.7	77.3
Black, female												
2004.....	4.4	*	*	*	0.8	1.2	1.8	3.2	8.3	16.7	34.8	52.4
2003.....	4.2	*	*	0.7	0.6	0.8	1.9	3.0	7.4	17.4	34.9	52.3
2002.....	4.3	*	*	*	0.7	1.1	1.4	3.6	9.2	15.7	33.8	56.3
2001.....	4.3	*	*	0.6	1.0	1.1	1.7	3.9	7.3	18.4	34.0	49.7
2000.....	4.2	*	*	*	0.9	1.3	2.1	4.3	8.1	16.4	28.8	54.1
1999.....	4.3	*	*	0.6	1.3	1.0	1.7	3.3	8.3	19.0	33.6	47.7

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Lymphoid leukemia (C91)												
All races 3/, both sexes												
2004.....	2.1	*	0.3	0.4	0.4	0.3	0.4	0.8	2.4	6.5	14.1	28.9
2003.....	2.2	*	0.4	0.4	0.4	0.3	0.4	0.7	2.3	6.8	15.4	29.6
2002.....	2.2	*	0.2	0.4	0.5	0.3	0.4	0.9	2.5	6.7	14.9	30.1
2001.....	2.2	*	0.3	0.4	0.5	0.4	0.4	0.8	2.5	7.0	15.1	31.2
2000.....	2.2	*	0.3	0.4	0.5	0.3	0.3	0.8	2.4	7.1	15.7	32.0
1999.....	2.2	*	0.3	0.4	0.5	0.3	0.3	0.8	2.8	7.6	15.9	30.8
All races 3/, male												
2004.....	2.5	*	0.4	0.5	0.6	0.4	0.4	1.0	3.5	9.1	20.1	41.7
2003.....	2.6	*	0.4	0.6	0.6	0.4	0.5	1.0	3.3	10.2	22.1	41.7
2002.....	2.6	*	*	0.5	0.7	0.4	0.6	1.1	3.3	10.0	21.3	43.0
2001.....	2.5	*	0.3	0.5	0.7	0.4	0.5	1.1	3.6	10.0	21.2	43.8
2000.....	2.6	*	0.3	0.5	0.7	0.4	0.4	1.0	3.5	10.4	22.4	43.8
1999.....	2.6	*	0.3	0.4	0.7	0.4	0.4	1.1	3.9	10.7	22.7	45.6
All races 3/, female												
2004.....	1.7	*	*	0.3	0.3	0.2	0.3	0.6	1.3	4.3	10.1	23.1
2003.....	1.8	*	0.4	0.3	0.3	0.2	0.3	0.5	1.4	3.9	10.8	24.3
2002.....	1.8	*	*	0.4	0.4	0.3	0.3	0.6	1.7	4.0	10.6	24.5
2001.....	1.8	*	0.3	0.3	0.3	0.3	0.3	0.6	1.5	4.4	11.0	25.9
2000.....	1.8	*	0.3	0.3	0.3	0.2	0.2	0.6	1.4	4.5	11.2	27.3
1999.....	1.9	*	0.3	0.3	0.3	0.2	0.3	0.5	1.7	5.0	11.5	24.8
White, both sexes												
2004.....	2.3	*	0.3	0.4	0.5	0.3	0.3	0.8	2.4	6.7	14.6	30.2
2003.....	2.4	*	0.4	0.5	0.5	0.3	0.4	0.7	2.4	7.0	16.1	31.0
2002.....	2.4	*	0.2	0.5	0.6	0.4	0.5	0.9	2.4	7.0	15.3	31.6
2001.....	2.4	*	0.3	0.4	0.6	0.4	0.4	0.8	2.5	7.1	15.6	32.8
2000.....	2.4	*	0.4	0.5	0.5	0.3	0.3	0.8	2.5	7.4	16.3	33.1
1999.....	2.5	*	0.4	0.4	0.5	0.4	0.3	0.8	2.8	7.8	16.5	32.0
White, male												
2004.....	2.7	*	0.3	0.5	0.6	0.4	0.4	1.0	3.6	9.1	20.8	43.7
2003.....	2.9	*	0.5	0.6	0.6	0.4	0.5	1.0	3.5	10.5	23.2	43.5
2002.....	2.8	*	*	0.5	0.7	0.4	0.6	1.2	3.3	10.4	22.0	45.1
2001.....	2.8	*	*	0.5	0.8	0.4	0.4	1.0	3.6	10.2	22.0	46.2
2000.....	2.8	*	0.3	0.6	0.7	0.4	0.5	1.0	3.6	10.6	23.3	45.0
1999.....	2.9	*	0.4	0.4	0.7	0.5	0.4	1.1	3.9	11.0	23.6	47.7

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Lymphoid leukemia (C91)												
White, female												
2004.....	1.9	*	*	0.4	0.4	0.2	0.3	0.6	1.3	4.5	10.3	24.1
2003.....	2.0	*	0.4	0.3	0.3	0.2	0.3	0.5	1.4	3.9	11.2	25.4
2002.....	2.0	*	*	0.5	0.4	0.3	0.3	0.6	1.7	4.1	10.9	25.8
2001.....	2.0	*	*	0.4	0.3	0.3	0.3	0.5	1.5	4.4	11.3	27.3
2000.....	2.1	*	0.4	0.3	0.3	0.2	0.2	0.6	1.4	4.6	11.7	28.3
1999.....	2.1	*	*	0.3	0.4	0.3	0.3	0.5	1.7	5.2	11.8	25.7
Black, both sexes												
2004.....	1.4	*	*	*	*	*	0.4	0.9	2.9	7.3	13.0	19.7
2003.....	1.3	*	*	*	0.3	*	0.4	0.8	2.4	7.3	11.6	19.8
2002.....	1.3	*	*	*	0.4	*	*	0.8	3.3	6.4	13.8	18.4
2001.....	1.5	*	*	0.4	0.4	*	*	1.3	3.1	7.7	13.2	19.0
2000.....	1.3	*	*	*	0.4	*	*	1.1	2.4	6.6	11.7	25.0
1999.....	1.4	*	*	0.3	0.5	*	0.4	0.9	3.4	7.4	12.4	22.0
Black, male												
2004.....	1.8	*	*	*	*	*	*	1.3	4.3	12.5	18.7	30.8
2003.....	1.6	*	*	*	*	*	*	1.4	3.4	9.9	15.8	26.1
2002.....	1.6	*	*	*	*	*	*	1.0	4.3	9.0	20.4	31.1
2001.....	1.7	*	*	*	*	*	*	1.5	4.6	11.0	18.8	28.4
2000.....	1.6	*	*	*	*	*	*	1.4	3.9	10.1	18.2	40.4
1999.....	1.7	*	*	*	*	*	*	1.3	4.6	11.5	17.5	32.8
Black, female												
2004.....	1.0	*	*	*	*	*	*	*	1.8	3.7	9.8	15.4
2003.....	1.1	*	*	*	*	*	*	*	1.5	5.5	9.2	17.4
2002.....	1.1	*	*	*	*	*	*	*	2.5	4.6	10.1	13.6
2001.....	1.2	*	*	*	*	*	*	1.1	1.9	5.3	10.1	15.6
2000.....	1.0	*	*	*	*	*	*	*	*	4.1	8.0	19.3
1999.....	1.1	*	*	*	*	*	*	*	2.3	4.6	9.5	18.0

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Myeloid leukemia (C92)												
All races 3/, both sexes												
2004.....	3.4	*	0.4	0.2	0.4	0.6	0.9	1.9	5.0	12.9	23.9	25.6
2003.....	3.4	*	0.3	0.3	0.4	0.6	1.1	2.0	5.1	13.0	23.7	26.1
2002.....	3.4	*	0.3	0.2	0.4	0.6	1.0	2.0	5.4	13.4	23.8	25.7
2001.....	3.5	*	0.5	0.2	0.5	0.7	1.1	2.3	5.4	13.6	23.6	25.1
2000.....	3.5	*	0.4	0.2	0.5	0.7	1.1	2.2	5.4	13.4	23.8	25.3
1999.....	3.3	*	0.4	0.2	0.5	0.7	1.1	2.2	5.5	12.9	21.5	25.6
All races 3/, male												
2004.....	3.8	*	0.4	0.3	0.5	0.5	1.0	2.1	5.9	17.3	32.7	36.1
2003.....	3.8	*	0.3	0.2	0.4	0.7	1.1	2.2	6.2	17.1	32.9	37.9
2002.....	3.9	*	0.4	0.2	0.4	0.7	1.0	2.2	6.6	17.8	33.7	37.6
2001.....	3.9	*	0.5	0.2	0.5	0.8	1.1	2.4	6.7	17.5	32.6	37.2
2000.....	3.9	*	0.4	0.2	0.6	0.8	1.1	2.5	6.4	17.8	33.2	39.1
1999.....	3.7	*	0.4	0.2	0.5	0.7	1.1	2.4	6.4	17.0	29.4	37.7
All races 3/, female												
2004.....	3.0	*	0.3	0.2	0.4	0.6	0.9	1.8	4.0	9.2	18.0	20.8
2003.....	3.0	*	*	0.3	0.3	0.5	1.0	1.7	4.0	9.6	17.6	20.9
2002.....	3.0	*	*	0.2	0.4	0.5	0.9	1.8	4.3	9.7	17.3	20.5
2001.....	3.1	*	0.5	0.2	0.5	0.6	1.0	2.2	4.3	10.4	17.6	20.1
2000.....	3.1	*	0.3	0.2	0.4	0.6	1.2	1.9	4.5	9.9	17.7	19.7
1999.....	3.0	*	0.3	0.2	0.5	0.6	1.0	2.0	4.6	9.5	16.3	20.7
White, both sexes												
2004.....	3.7	*	0.3	0.2	0.4	0.6	0.9	1.9	5.1	13.4	24.8	26.7
2003.....	3.7	*	0.3	0.3	0.4	0.6	1.0	2.0	5.2	13.6	24.7	26.9
2002.....	3.8	*	0.3	0.2	0.4	0.6	1.0	2.0	5.4	14.0	24.8	26.5
2001.....	3.8	*	0.5	0.2	0.5	0.7	1.1	2.3	5.6	14.1	24.4	26.0
2000.....	3.8	*	0.4	0.2	0.4	0.6	1.1	2.2	5.4	14.1	24.9	25.9
1999.....	3.6	*	0.4	0.2	0.4	0.6	1.1	2.2	5.6	13.3	22.2	26.7
White, male												
2004.....	4.2	*	0.3	0.3	0.5	0.5	1.0	2.1	6.2	18.0	33.8	38.3
2003.....	4.3	*	*	0.2	0.4	0.7	1.1	2.3	6.4	17.8	34.6	39.6
2002.....	4.3	*	0.4	0.2	0.4	0.7	1.0	2.2	6.6	18.6	35.0	38.7
2001.....	4.3	*	0.5	0.2	0.5	0.8	1.1	2.4	6.9	18.1	33.9	39.1
2000.....	4.3	*	0.4	0.2	0.5	0.7	1.1	2.5	6.4	18.5	34.8	41.0
1999.....	4.0	*	0.3	0.2	0.5	0.6	1.1	2.4	6.6	17.5	30.4	39.5

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Myeloid leukemia (C92)												
White, female												
2004.....	3.3	*	0.3	0.2	0.4	0.6	0.8	1.8	4.1	9.6	18.7	21.5
2003.....	3.3	*	*	0.3	0.3	0.5	1.0	1.7	4.2	10.0	18.1	21.3
2002.....	3.3	*	*	0.2	0.4	0.5	1.0	1.8	4.3	10.2	17.9	21.2
2001.....	3.4	*	0.6	0.2	0.5	0.6	1.0	2.2	4.4	10.8	18.1	20.5
2000.....	3.3	*	*	0.2	0.4	0.5	1.1	1.9	4.5	10.4	18.4	19.8
1999.....	3.3	*	0.4	0.2	0.4	0.7	1.0	2.1	4.7	9.8	16.8	21.6
Black, both sexes												
2004.....	2.2	*	*	*	0.5	0.7	1.1	2.3	5.0	10.2	17.0	17.8
2003.....	2.1	*	*	0.3	0.5	0.6	1.3	2.1	4.7	10.4	16.6	21.0
2002.....	2.1	*	*	*	0.4	0.8	0.9	2.1	5.6	10.2	16.3	19.3
2001.....	2.1	*	*	*	0.4	0.7	1.4	2.1	4.4	11.4	16.6	17.2
2000.....	2.3	*	*	*	0.7	1.1	1.4	2.8	6.0	9.8	14.4	20.3
1999.....	2.1	*	*	0.3	0.7	0.7	1.2	2.4	4.9	10.9	15.6	14.3
Black, male												
2004.....	2.2	*	*	*	*	0.8	1.2	2.7	5.6	13.6	20.6	19.9
2003.....	2.3	*	*	*	*	0.8	1.6	2.2	5.8	14.2	21.4	31.3
2002.....	2.3	*	*	*	*	0.9	1.1	2.5	6.8	13.8	20.9	32.2
2001.....	2.3	*	*	*	*	1.0	1.6	2.2	5.7	15.6	20.3	*
2000.....	2.4	*	*	*	1.0	1.3	1.4	3.1	7.0	12.9	18.2	24.2
1999.....	2.3	*	*	*	0.8	0.9	1.2	3.0	6.3	14.3	20.3	*
Black, female												
2004.....	2.1	*	*	*	*	0.7	1.0	2.0	4.5	7.7	14.9	16.9
2003.....	2.0	*	*	*	*	*	1.1	2.1	3.9	7.6	13.9	17.0
2002.....	1.9	*	*	*	*	0.7	0.8	1.8	4.6	7.6	13.6	14.4
2001.....	2.0	*	*	*	*	*	1.2	2.1	3.3	8.5	14.5	16.4
2000.....	2.2	*	*	*	*	0.8	1.5	2.6	5.2	7.6	12.3	18.9
1999.....	1.9	*	*	*	0.7	*	1.2	1.8	3.7	8.5	12.9	12.3

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Monocytic leukemia (C93)												
All races 3/, both sexes												
2004.....	0.0	*	*	*	*	*	*	*	*	0.1	0.4	0.6
2003.....	0.0	*	*	*	*	*	*	*	*	0.1	0.4	0.6
2002.....	0.1	*	*	*	*	*	*	*	*	0.2	0.4	0.6
2001.....	0.1	*	*	*	*	*	*	*	*	0.2	0.5	0.7
2000.....	0.1	*	*	*	*	*	*	*	0.1	0.2	0.4	0.7
1999.....	0.1	*	*	*	*	*	*	*	*	0.2	0.5	0.9
All races 3/, male												
2004.....	0.1	*	*	*	*	*	*	*	*	*	0.7	*
2003.....	0.1	*	*	*	*	*	*	*	*	*	0.6	*
2002.....	0.1	*	*	*	*	*	*	*	*	0.3	0.5	*
2001.....	0.1	*	*	*	*	*	*	*	*	*	0.6	*
2000.....	0.1	*	*	*	*	*	*	*	*	0.3	0.6	*
1999.....	0.1	*	*	*	*	*	*	*	*	0.3	0.8	*
All races 3/, female												
2004.....	0.0	*	*	*	*	*	*	*	*	*	0.3	*
2003.....	0.0	*	*	*	*	*	*	*	*	*	0.3	*
2002.....	0.0	*	*	*	*	*	*	*	*	*	0.3	*
2001.....	0.1	*	*	*	*	*	*	*	*	*	0.4	*
2000.....	0.0	*	*	*	*	*	*	*	*	*	*	*
1999.....	0.1	*	*	*	*	*	*	*	*	0.2	0.3	0.7
White, both sexes												
2004.....	0.0	*	*	*	*	*	*	*	*	0.1	0.5	0.6
2003.....	0.1	*	*	*	*	*	*	*	*	0.2	0.5	0.7
2002.....	0.1	*	*	*	*	*	*	*	*	0.2	0.4	0.6
2001.....	0.1	*	*	*	*	*	*	*	*	0.2	0.5	0.8
2000.....	0.1	*	*	*	*	*	*	*	*	0.2	0.4	0.7
1999.....	0.1	*	*	*	*	*	*	*	*	0.3	0.5	1.0
White, male												
2004.....	0.1	*	*	*	*	*	*	*	*	*	0.7	*
2003.....	0.1	*	*	*	*	*	*	*	*	*	0.6	*
2002.....	0.1	*	*	*	*	*	*	*	*	0.3	0.6	*
2001.....	0.1	*	*	*	*	*	*	*	*	*	0.6	*
2000.....	0.1	*	*	*	*	*	*	*	*	0.3	0.6	*
1999.....	0.1	*	*	*	*	*	*	*	*	0.3	0.8	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified leukemia (C94-C95)												
All races 3/, both sexes												
2004.....	1.8	*	0.1	0.1	0.2	0.2	0.3	0.6	1.9	5.8	13.9	23.8
2003.....	1.8	*	0.2	0.1	0.1	0.2	0.3	0.6	1.9	5.7	14.4	24.0
2002.....	1.8	*	*	0.1	0.2	0.2	0.3	0.7	1.9	6.1	14.0	24.1
2001.....	1.8	*	0.2	0.2	0.2	0.2	0.3	0.7	2.0	5.9	14.2	24.4
2000.....	1.9	*	0.2	0.1	0.2	0.2	0.3	0.7	2.0	6.5	14.6	25.6
1999.....	1.9	*	*	0.2	0.3	0.2	0.3	0.7	2.3	6.5	14.9	25.2
All races 3/, male												
2004.....	2.0	*	*	0.2	0.2	0.2	0.4	0.7	2.4	7.8	19.7	31.7
2003.....	1.9	*	*	0.1	0.2	0.2	0.4	0.8	2.3	7.6	19.6	31.1
2002.....	2.0	*	*	0.2	0.3	0.3	0.3	0.8	2.3	8.2	19.3	32.6
2001.....	2.0	*	0.3	0.2	0.2	0.3	0.3	0.8	2.4	8.0	19.8	35.3
2000.....	2.0	*	*	0.1	0.2	0.3	0.4	0.8	2.5	8.8	19.5	35.7
1999.....	2.1	*	*	0.2	0.3	0.2	0.4	0.8	2.7	8.8	20.3	34.6
All races 3/, female												
2004.....	1.6	*	*	0.1	0.1	0.1	0.3	0.4	1.4	4.2	10.0	20.3
2003.....	1.6	*	*	0.1	0.1	0.1	0.3	0.5	1.5	4.2	10.9	20.8
2002.....	1.6	*	*	0.1	0.2	0.2	0.3	0.6	1.6	4.4	10.5	20.4
2001.....	1.6	*	*	0.1	0.2	0.2	0.2	0.5	1.7	4.3	10.5	19.8
2000.....	1.7	*	*	0.1	0.2	0.2	0.3	0.7	1.6	4.5	11.4	21.4
1999.....	1.8	*	*	0.1	0.2	0.2	0.3	0.6	1.9	4.6	11.5	21.3
White, both sexes												
2004.....	1.9	*	*	0.1	0.2	0.2	0.3	0.6	1.9	5.8	14.4	24.1
2003.....	2.0	*	0.2	0.1	0.2	0.2	0.3	0.6	1.9	5.9	14.7	24.8
2002.....	2.0	*	*	0.1	0.2	0.2	0.3	0.6	1.9	6.3	14.4	24.2
2001.....	2.0	*	0.3	0.2	0.2	0.2	0.3	0.7	2.0	6.0	14.6	25.1
2000.....	2.1	*	0.2	0.1	0.2	0.2	0.3	0.7	2.0	6.6	15.0	26.3
1999.....	2.1	*	*	0.2	0.3	0.2	0.3	0.7	2.3	6.6	15.3	25.7
White, male												
2004.....	2.2	*	*	0.2	0.2	0.2	0.3	0.7	2.4	7.7	20.5	32.1
2003.....	2.1	*	*	0.1	0.2	0.2	0.4	0.8	2.3	7.8	20.2	32.7
2002.....	2.2	*	*	0.1	0.3	0.3	0.3	0.7	2.3	8.3	19.9	34.0
2001.....	2.2	*	*	0.2	0.2	0.3	0.3	0.8	2.4	8.1	20.3	36.4
2000.....	2.2	*	*	0.1	0.2	0.3	0.4	0.7	2.5	9.0	19.9	36.4
1999.....	2.3	*	*	0.2	0.3	0.2	0.4	0.7	2.8	9.1	20.8	35.4

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified leukemia (C94-C95)												
White, female												
2004.....	1.7	*	*	*	*	*	0.3	0.4	1.4	4.2	10.2	20.5
2003.....	1.8	*	*	0.1	*	0.1	0.3	0.5	1.5	4.2	11.0	21.4
2002.....	1.8	*	*	*	0.2	0.2	0.3	0.6	1.6	4.6	10.7	20.0
2001.....	1.8	*	*	0.2	0.2	*	0.2	0.5	1.6	4.3	10.9	20.3
2000.....	1.9	*	*	*	0.2	0.2	0.3	0.7	1.6	4.6	11.8	22.2
1999.....	1.9	*	*	0.1	0.2	0.2	0.3	0.6	1.9	4.6	11.7	21.8
Black, both sexes												
2004.....	1.3	*	*	*	*	*	0.5	0.7	2.3	7.6	11.1	22.3
2003.....	1.2	*	*	*	*	*	0.5	0.7	2.4	5.3	13.2	18.4
2002.....	1.2	*	*	*	*	*	*	1.3	2.3	5.6	11.8	26.1
2001.....	1.2	*	*	*	*	*	0.4	0.8	2.3	5.8	11.5	20.6
2000.....	1.2	*	*	*	*	*	0.4	0.9	2.3	6.2	12.0	20.0
1999.....	1.2	*	*	*	*	*	*	1.0	2.3	6.2	12.8	19.4
Black, male												
2004.....	1.3	*	*	*	*	*	*	0.9	2.9	10.9	12.8	27.8
2003.....	1.2	*	*	*	*	*	*	1.0	3.1	6.8	16.1	*
2002.....	1.2	*	*	*	*	*	*	1.5	2.5	8.7	15.2	22.5
2001.....	1.3	*	*	*	*	*	*	*	2.7	7.4	15.6	29.5
2000.....	1.3	*	*	*	*	*	*	*	2.8	8.6	18.2	32.3
1999.....	1.2	*	*	*	*	*	*	1.1	2.5	7.1	16.3	24.6
Black, female												
2004.....	1.2	*	*	*	*	*	*	*	1.9	5.2	10.1	20.1
2003.....	1.2	*	*	*	*	*	*	*	1.9	4.2	11.6	17.8
2002.....	1.2	*	*	*	*	*	*	1.0	2.0	3.4	9.9	27.5
2001.....	1.1	*	*	*	*	*	*	*	2.0	4.6	9.3	17.3
2000.....	1.0	*	*	*	*	*	*	*	1.8	4.4	8.5	15.5
1999.....	1.2	*	*	*	*	*	*	1.0	2.2	5.6	10.8	17.5

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Benign neoplasms (D10-D36)												
All races 3/, both sexes												
2004.....	0.5	*	*	*	*	0.1	0.1	0.3	0.6	1.3	3.6	7.9
2003.....	0.5	*	*	*	*	0.1	0.2	0.3	0.6	1.4	3.7	7.4
2002.....	0.6	*	*	*	*	0.1	0.2	0.4	0.6	1.6	3.6	7.8
2001.....	0.5	*	*	*	0.1	0.1	0.1	0.3	0.7	1.5	3.6	8.7
2000.....	0.6	0.6	*	*	*	0.1	0.2	0.3	0.6	1.5	4.2	9.0
1999.....	0.6	*	*	*	*	0.1	0.1	0.3	0.7	1.6	4.0	8.4
All races 3/, male												
2004.....	0.4	*	*	*	*	0.1	0.1	0.2	0.5	1.2	3.4	7.4
2003.....	0.4	*	*	*	*	*	0.1	0.3	0.6	1.4	3.6	7.3
2002.....	0.4	*	*	*	*	0.1	0.2	0.3	0.6	1.6	3.4	6.3
2001.....	0.5	*	*	*	*	*	0.1	0.3	0.8	1.4	3.9	8.9
2000.....	0.4	*	*	*	*	*	0.1	0.3	0.4	1.5	3.8	8.7
1999.....	0.4	*	*	*	*	*	0.2	0.3	0.7	1.6	3.6	9.2
All races 3/, female												
2004.....	0.7	*	*	*	*	0.1	0.2	0.4	0.7	1.4	3.8	8.1
2003.....	0.6	*	*	*	*	0.1	0.2	0.4	0.6	1.3	3.7	7.4
2002.....	0.7	*	*	*	*	*	0.2	0.4	0.6	1.6	3.8	8.4
2001.....	0.6	*	*	*	*	0.1	0.2	0.3	0.7	1.5	3.4	8.6
2000.....	0.7	*	*	*	*	0.1	0.3	0.3	0.7	1.5	4.4	9.1
1999.....	0.7	*	*	*	*	*	0.1	0.3	0.8	1.6	4.2	8.1
White, both sexes												
2004.....	0.5	*	*	*	*	0.1	0.1	0.3	0.6	1.1	3.5	7.8
2003.....	0.5	*	*	*	*	0.1	0.1	0.3	0.6	1.2	3.7	7.4
2002.....	0.6	*	*	*	*	0.1	0.2	0.3	0.6	1.5	3.5	7.4
2001.....	0.5	*	*	*	*	0.1	0.1	0.3	0.6	1.4	3.4	8.2
2000.....	0.6	*	*	*	*	*	0.1	0.3	0.5	1.4	4.1	8.9
1999.....	0.6	*	*	*	*	*	0.1	0.2	0.6	1.5	3.8	8.4
White, male												
2004.....	0.4	*	*	*	*	*	*	0.2	0.5	1.1	3.2	7.3
2003.....	0.4	*	*	*	*	*	0.1	0.3	0.6	1.3	3.5	7.3
2002.....	0.4	*	*	*	*	*	0.2	0.3	0.6	1.6	3.2	5.7
2001.....	0.5	*	*	*	*	*	*	0.3	0.6	1.3	3.8	8.4
2000.....	0.4	*	*	*	*	*	*	0.3	0.4	1.3	3.9	8.7
1999.....	0.4	*	*	*	*	*	0.1	0.2	0.6	1.5	3.4	9.3

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Benign neoplasms (D10-D36)												
White, female												
2004.....	0.7	*	*	*	*	*	0.1	0.3	0.7	1.2	3.7	8.0
2003.....	0.7	*	*	*	*	*	0.2	0.3	0.6	1.2	3.8	7.4
2002.....	0.7	*	*	*	*	*	0.2	0.4	0.6	1.5	3.7	8.1
2001.....	0.6	*	*	*	*	*	0.1	0.3	0.6	1.4	3.2	8.0
2000.....	0.7	*	*	*	*	*	0.2	0.2	0.7	1.5	4.3	9.0
1999.....	0.7	*	*	*	*	*	*	0.2	0.7	1.4	4.1	8.1
Black, both sexes												
2004.....	0.6	*	*	*	*	*	*	0.5	1.1	2.7	5.8	10.2
2003.....	0.6	*	*	*	*	*	0.5	0.6	0.9	2.9	4.1	7.9
2002.....	0.6	*	*	*	*	*	*	0.7	1.0	2.6	5.5	11.3
2001.....	0.7	*	*	*	*	*	*	0.6	1.8	2.6	5.5	15.4
2000.....	0.6	*	*	*	*	*	0.5	0.7	0.8	2.7	5.6	10.3
1999.....	0.7	*	*	*	*	*	*	0.7	1.8	2.8	6.6	8.6
Black, male												
2004.....	0.5	*	*	*	*	*	*	*	*	*	7.5	*
2003.....	0.5	*	*	*	*	*	*	*	*	3.5	*	*
2002.....	0.5	*	*	*	*	*	*	*	*	*	5.7	*
2001.....	0.6	*	*	*	*	*	*	*	2.1	*	6.2	*
2000.....	0.4	*	*	*	*	*	*	*	*	2.9	*	*
1999.....	0.6	*	*	*	*	*	*	*	2.1	*	7.4	*
Black, female												
2004.....	0.7	*	*	*	*	*	*	*	*	3.2	4.9	9.8
2003.....	0.6	*	*	*	*	*	*	*	*	2.4	3.5	8.5
2002.....	0.7	*	*	*	*	*	*	1.0	*	2.7	5.4	11.1
2001.....	0.7	*	*	*	*	*	*	*	1.5	2.5	5.1	14.7
2000.....	0.8	*	*	*	*	*	0.7	*	*	2.6	6.3	10.3
1999.....	0.7	*	*	*	*	*	*	*	1.5	3.2	6.2	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Of eye, brain and other parts of central nervous system (D31-D33)

All races 3/, both sexes

2004.....	0.3	*	*	*	*	*	0.0	0.2	0.4	0.9	2.5	5.6
2003.....	0.3	*	*	*	*	*	0.1	0.2	0.4	0.9	2.5	5.4
2002.....	0.3	*	*	*	*	*	0.1	0.2	0.3	1.0	2.4	5.1
2001.....	0.3	*	*	*	*	*	0.0	0.1	0.4	1.0	2.3	5.9
2000.....	0.3	*	*	*	*	*	0.1	0.1	0.3	1.0	2.9	6.2
1999.....	0.3	*	*	*	*	*	0.1	0.1	0.5	1.0	2.7	5.5

All races 3/, male

2004.....	0.2	*	*	*	*	*	*	0.2	0.2	0.7	2.2	5.2
2003.....	0.3	*	*	*	*	*	*	0.2	0.3	0.9	2.3	5.2
2002.....	0.2	*	*	*	*	*	*	0.2	0.3	1.0	2.0	3.2
2001.....	0.2	*	*	*	*	*	*	*	0.4	0.9	2.2	5.4
2000.....	0.3	*	*	*	*	*	*	0.2	0.2	0.9	2.5	5.9
1999.....	0.2	*	*	*	*	*	*	0.1	0.4	1.0	2.4	5.5

All races 3/, female

2004.....	0.4	*	*	*	*	*	*	0.2	0.5	1.0	2.6	5.8
2003.....	0.4	*	*	*	*	*	*	0.2	0.4	1.0	2.6	5.5
2002.....	0.4	*	*	*	*	*	0.1	0.2	0.4	1.1	2.7	5.9
2001.....	0.4	*	*	*	*	*	*	0.2	0.4	1.0	2.4	6.2
2000.....	0.4	*	*	*	*	*	*	0.1	0.5	1.0	3.1	6.4
1999.....	0.4	*	*	*	*	*	*	0.1	0.5	1.1	3.0	5.5

White, both sexes

2004.....	0.3	*	*	*	*	*	*	0.2	0.3	0.7	2.3	5.7
2003.....	0.3	*	*	*	*	*	*	0.2	0.3	0.8	2.5	5.4
2002.....	0.3	*	*	*	*	*	0.1	0.2	0.3	1.0	2.3	4.9
2001.....	0.3	*	*	*	*	*	*	0.1	0.3	0.9	2.2	5.5
2000.....	0.4	*	*	*	*	*	0.1	0.1	0.3	0.9	2.8	6.3
1999.....	0.3	*	*	*	*	*	*	0.1	0.4	1.0	2.6	5.6

White, male

2004.....	0.2	*	*	*	*	*	*	0.2	0.2	0.6	2.0	5.2
2003.....	0.3	*	*	*	*	*	*	0.1	0.3	0.8	2.3	5.3
2002.....	0.2	*	*	*	*	*	*	0.2	0.3	1.0	1.9	2.9
2001.....	0.2	*	*	*	*	*	*	*	0.2	0.9	2.1	4.9
2000.....	0.3	*	*	*	*	*	*	0.2	*	0.8	2.5	6.0
1999.....	0.3	*	*	*	*	*	*	*	0.3	1.0	2.2	5.5

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Of eye, brain and other parts of central nervous system (D31-D33)

White, female

2004.....	0.4	*	*	*	*	*	*	0.2	0.5	0.8	2.6	5.9
2003.....	0.4	*	*	*	*	*	*	0.2	0.4	0.8	2.6	5.4
2002.....	0.4	*	*	*	*	*	*	0.2	0.4	1.0	2.6	5.8
2001.....	0.4	*	*	*	*	*	*	0.1	0.3	0.9	2.3	5.8
2000.....	0.5	*	*	*	*	*	*	*	0.5	1.0	3.0	6.4
1999.....	0.4	*	*	*	*	*	*	*	0.5	1.0	2.9	5.6

Black, both sexes

2004.....	0.4	*	*	*	*	*	*	*	0.7	2.1	3.9	5.9
2003.....	0.3	*	*	*	*	*	*	*	*	2.2	2.6	6.1
2002.....	0.3	*	*	*	*	*	*	*	*	1.5	3.9	5.9
2001.....	0.4	*	*	*	*	*	*	*	1.2	1.8	3.6	11.1
2000.....	0.3	*	*	*	*	*	*	*	*	1.3	3.9	*
1999.....	0.4	*	*	*	*	*	*	*	1.1	1.9	4.4	*

Black, male

2004.....	0.3	*	*	*	*	*	*	*	*	*	*	*
2003.....	0.2	*	*	*	*	*	*	*	*	*	*	*
2002.....	0.2	*	*	*	*	*	*	*	*	*	*	*
2001.....	0.3	*	*	*	*	*	*	*	*	*	*	*
2000.....	0.2	*	*	*	*	*	*	*	*	*	*	*
1999.....	0.3	*	*	*	*	*	*	*	*	*	*	*

Black, female

2004.....	0.4	*	*	*	*	*	*	*	*	2.5	3.5	*
2003.....	0.4	*	*	*	*	*	*	*	*	2.0	*	*
2002.....	0.4	*	*	*	*	*	*	*	*	*	4.1	*
2001.....	0.5	*	*	*	*	*	*	*	*	*	3.8	10.9
2000.....	0.4	*	*	*	*	*	*	*	*	*	4.8	*
1999.....	0.5	*	*	*	*	*	*	*	*	2.7	4.1	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Of other and unspecified sites (D10-D24,D29-D30,D34-D36)

All races 3/, both sexes

2004.....	0.2	*	*	*	*	0.1	0.1	0.1	0.3	0.4	1.1	2.2
2003.....	0.2	*	*	*	*	0.1	0.1	0.1	0.3	0.4	1.1	1.9
2002.....	0.2	*	*	*	*	0.1	0.1	0.1	0.3	0.6	1.2	2.6
2001.....	0.2	*	*	*	*	*	0.1	0.1	0.3	0.5	1.2	2.5
2000.....	0.2	0.5	*	*	*	*	0.1	0.1	0.2	0.5	1.3	2.6
1999.....	0.2	*	*	*	*	*	0.1	0.1	0.3	0.5	1.2	2.8

All races 3/, male

2004.....	0.2	*	*	*	*	*	*	*	0.3	0.5	1.3	2.3
2003.....	0.2	*	*	*	*	*	0.1	0.1	0.3	0.5	1.3	2.1
2002.....	0.2	*	*	*	*	*	0.1	0.1	0.3	0.7	1.4	3.1
2001.....	0.2	*	*	*	*	*	*	0.2	0.4	0.5	1.7	3.5
2000.....	0.2	*	*	*	*	*	*	0.2	0.2	0.6	1.3	2.9
1999.....	0.2	*	*	*	*	*	0.1	0.1	0.4	0.6	1.3	3.7

All races 3/, female

2004.....	0.2	*	*	*	*	*	*	0.1	0.1	0.2	0.3	1.0	2.2
2003.....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.3	1.0	1.8	
2002.....	0.2	*	*	*	*	*	0.1	0.2	0.2	0.5	1.0	2.4	
2001.....	0.2	*	*	*	*	*	*	0.1	0.3	0.4	0.9	2.1	
2000.....	0.2	*	*	*	*	*	0.1	*	0.2	0.5	1.2	2.6	
1999.....	0.2	*	*	*	*	*	*	0.1	0.2	0.4	1.2	2.4	

White, both sexes

2004.....	0.2	*	*	*	*	*	0.1	0.1	0.3	0.4	1.1	2.0
2003.....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.4	1.1	1.9
2002.....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.5	1.2	2.3
2001.....	0.2	*	*	*	*	*	0.1	0.1	0.3	0.4	1.2	2.5
2000.....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.5	1.3	2.5
1999.....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.5	1.2	2.8

White, male

2004.....	0.2	*	*	*	*	*	*	*	0.3	0.5	1.2	2.1
2003.....	0.2	*	*	*	*	*	*	0.1	0.3	0.5	1.2	2.0
2002.....	0.2	*	*	*	*	*	*	0.1	0.3	0.6	1.4	2.8
2001.....	0.2	*	*	*	*	*	*	0.2	0.4	0.5	1.7	3.5
2000.....	0.2	*	*	*	*	*	*	0.2	0.2	0.5	1.3	2.7
1999.....	0.2	*	*	*	*	*	*	*	0.3	0.5	1.2	3.8

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Neoplasms of uncertain or unknown behavior of specified sites (D37-D47)

All races 3/, both sexes

2004.....	3.8	0.8	0.3	0.2	0.2	0.3	0.5	1.2	3.5	10.8	30.8	65.6
2003.....	3.9	0.9	0.2	0.2	0.2	0.3	0.5	1.2	3.6	11.0	31.6	69.6
2002.....	3.8	0.9	0.3	0.2	0.2	0.3	0.5	1.3	3.6	10.8	31.1	67.0
2001.....	4.0	0.5	0.3	0.2	0.2	0.2	0.6	1.4	4.0	12.0	32.2	70.1
2000.....	4.0	0.7	0.3	0.2	0.2	0.3	0.6	1.4	4.2	11.8	32.0	72.0
1999.....	4.0	0.8	0.4	0.2	0.2	0.3	0.6	1.4	3.8	12.5	31.4	70.8

All races 3/, male

2004.....	4.1	*	0.3	0.2	0.2	0.3	0.6	1.4	4.5	13.9	40.8	87.9
2003.....	4.1	*	0.3	0.2	0.2	0.3	0.7	1.4	4.3	14.1	41.3	91.6
2002.....	4.0	*	0.3	0.2	0.2	0.3	0.6	1.4	4.5	13.7	41.3	91.9
2001.....	4.2	*	0.3	0.2	0.1	0.3	0.7	1.6	4.6	15.3	43.1	96.6
2000.....	4.2	*	0.4	0.2	0.2	0.3	0.7	1.6	4.9	15.1	42.4	97.6
1999.....	4.2	*	0.4	0.2	0.2	0.4	0.7	1.6	4.8	16.2	41.4	97.8

All races 3/, female

2004.....	3.6	*	*	0.2	0.2	0.2	0.4	1.0	2.7	8.2	24.0	55.5
2003.....	3.8	*	*	0.1	0.1	0.2	0.4	1.1	3.0	8.4	25.0	59.9
2002.....	3.7	*	0.3	0.2	0.2	0.3	0.5	1.1	2.8	8.3	24.3	56.2
2001.....	3.9	*	0.3	0.2	0.2	0.2	0.6	1.2	3.4	9.3	25.0	59.0
2000.....	3.9	*	*	0.2	0.1	0.3	0.5	1.2	3.5	9.2	25.2	61.5
1999.....	3.8	*	0.3	0.2	0.2	0.3	0.5	1.2	2.9	9.5	24.9	60.0

White, both sexes

2004.....	4.3	0.8	0.3	0.2	0.2	0.3	0.5	1.2	3.5	11.2	31.8	68.2
2003.....	4.4	0.8	0.2	0.1	0.2	0.3	0.5	1.2	3.6	11.4	32.7	72.1
2002.....	4.3	1.0	0.3	0.2	0.2	0.3	0.5	1.3	3.7	11.2	32.1	69.3
2001.....	4.5	*	0.3	0.2	0.1	0.3	0.6	1.4	4.1	12.5	33.3	71.6
2000.....	4.5	0.7	0.3	0.2	0.2	0.3	0.6	1.4	4.3	12.1	33.2	74.3
1999.....	4.4	0.7	0.3	0.2	0.2	0.3	0.6	1.4	3.8	12.8	32.4	73.4

White, male

2004.....	4.5	*	0.3	0.2	0.2	0.3	0.6	1.4	4.4	14.4	42.3	92.1
2003.....	4.5	*	*	0.2	0.2	0.3	0.7	1.4	4.2	14.6	43.0	95.6
2002.....	4.5	*	*	0.2	0.2	0.3	0.6	1.5	4.5	14.3	42.6	95.4
2001.....	4.7	*	*	0.2	0.1	0.3	0.7	1.7	4.7	16.0	44.4	99.2
2000.....	4.6	*	0.4	0.2	0.2	0.3	0.7	1.6	5.0	15.3	43.9	101.1
1999.....	4.6	*	0.3	0.2	0.2	0.3	0.7	1.7	4.9	16.6	42.8	102.7

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Neoplasms of uncertain or unknown behavior of specified sites (D37-D47)

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
White, female												
2004.....	4.0	*	*	0.2	0.2	0.2	0.4	1.0	2.7	8.5	24.7	57.4
2003.....	4.2	*	*	0.1	0.1	0.2	0.4	1.0	3.1	8.6	25.8	61.7
2002.....	4.1	*	*	0.2	0.2	0.2	0.4	1.1	2.9	8.6	25.1	57.9
2001.....	4.3	*	*	0.2	0.1	0.2	0.5	1.2	3.5	9.5	25.9	60.1
2000.....	4.4	*	*	0.2	*	0.3	0.5	1.1	3.6	9.4	26.1	63.4
1999.....	4.2	*	0.4	0.2	0.2	0.3	0.5	1.2	2.9	9.7	25.5	61.6
Black, both sexes												
2004.....	2.3	*	*	*	*	0.4	0.7	1.5	4.7	9.5	24.1	42.3
2003.....	2.2	*	*	*	*	0.4	0.6	1.8	4.1	9.6	24.4	46.7
2002.....	2.2	*	*	*	*	0.4	0.7	1.5	4.2	9.4	24.8	49.9
2001.....	2.3	*	*	0.3	*	*	0.8	1.3	3.8	10.4	25.1	55.6
2000.....	2.3	*	*	*	*	0.4	0.5	1.9	4.2	11.4	23.3	51.6
1999.....	2.4	*	*	*	*	*	0.8	1.7	4.1	12.2	24.9	48.8
Black, male												
2004.....	2.3	*	*	*	*	*	0.8	1.9	6.3	12.6	29.8	48.7
2003.....	2.2	*	*	*	*	*	0.8	1.8	5.3	12.4	30.5	56.3
2002.....	2.2	*	*	*	*	*	*	1.6	5.4	11.1	32.4	66.5
2001.....	2.2	*	*	*	*	*	0.8	1.3	4.2	12.9	34.4	71.5
2000.....	2.4	*	*	*	*	*	0.8	2.1	5.4	14.9	29.7	64.6
1999.....	2.3	*	*	*	*	*	0.7	1.7	4.6	15.9	29.2	52.7
Black, female												
2004.....	2.2	*	*	*	*	*	0.7	1.1	3.4	7.2	20.9	39.8
2003.....	2.2	*	*	*	*	*	*	1.7	3.1	7.6	21.0	42.9
2002.....	2.3	*	*	*	*	*	0.9	1.4	3.3	8.1	20.5	43.5
2001.....	2.3	*	*	*	*	*	0.7	1.3	3.5	8.7	19.8	49.7
2000.....	2.2	*	*	*	*	*	*	1.7	3.3	9.0	19.8	46.8
1999.....	2.5	*	*	*	*	*	0.8	1.7	3.7	9.5	22.5	47.3

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Neoplasms of uncertain or unknown behavior of unspecified sites (D48)

All races 3/, both sexes

2004.....	0.3	0.5	*	*	0.1	*	0.1	0.2	0.3	0.6	1.4	4.1
2003.....	0.2	*	*	*	*	0.1	0.0	0.1	0.2	0.5	1.2	3.4
2002.....	0.2	*	*	*	*	0.1	0.1	0.1	0.2	0.4	1.4	3.8
2001.....	0.2	*	*	*	*	*	0.0	0.1	0.2	0.5	1.2	3.8
2000.....	0.2	*	*	*	*	*	0.1	0.1	0.3	0.4	1.4	3.9
1999.....	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.1	3.5

All races 3/, male

2004.....	0.2	*	*	*	*	*	*	0.2	0.3	0.6	1.3	3.6
2003.....	0.2	*	*	*	*	*	*	*	0.2	0.4	1.2	2.6
2002.....	0.2	*	*	*	*	*	0.1	*	0.2	0.5	1.1	3.0
2001.....	0.2	*	*	*	*	*	*	0.2	0.2	0.6	1.1	1.8
2000.....	0.2	*	*	*	*	*	*	*	0.3	0.4	1.3	2.4
1999.....	0.1	*	*	*	*	*	*	*	0.2	0.5	1.0	2.1

All races 3/, female

2004.....	0.3	*	*	*	*	*	*	0.1	0.3	0.5	1.4	4.4
2003.....	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.1	3.8
2002.....	0.3	*	*	*	*	*	*	0.2	0.3	0.4	1.5	4.2
2001.....	0.3	*	*	*	*	*	*	0.1	0.3	0.5	1.2	4.6
2000.....	0.3	*	*	*	*	*	*	0.1	0.3	0.5	1.5	4.4
1999.....	0.2	*	*	*	*	*	*	0.1	0.2	0.4	1.2	4.1

White, both sexes

2004.....	0.3	*	*	*	*	*	0.1	0.2	0.3	0.6	1.4	4.2
2003.....	0.2	*	*	*	*	0.1	*	0.1	0.2	0.5	1.2	3.5
2002.....	0.2	*	*	*	*	*	0.1	0.1	0.3	0.4	1.4	3.9
2001.....	0.2	*	*	*	*	*	*	0.2	0.2	0.5	1.2	4.0
2000.....	0.2	*	*	*	*	*	0.1	0.1	0.3	0.4	1.4	4.1
1999.....	0.2	*	*	*	*	*	*	0.1	0.2	0.4	1.1	3.4

White, male

2004.....	0.2	*	*	*	*	*	*	0.2	0.3	0.6	1.3	3.7
2003.....	0.2	*	*	*	*	*	*	*	0.2	0.5	1.3	2.5
2002.....	0.2	*	*	*	*	*	*	*	0.2	0.5	1.2	3.1
2001.....	0.2	*	*	*	*	*	*	0.2	0.2	0.5	1.2	2.0
2000.....	0.2	*	*	*	*	*	*	*	0.2	0.4	1.2	2.6
1999.....	0.2	*	*	*	*	*	*	*	0.2	0.5	1.0	2.2

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
III. Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)												
All races 3/, both sexes												
2004.....	3.2	2.3	0.4	0.2	0.5	0.8	1.4	2.2	3.5	7.0	17.3	52.9
2003.....	3.3	2.6	0.5	0.2	0.4	0.8	1.4	2.3	3.5	7.4	17.6	56.7
2002.....	3.4	2.0	0.5	0.2	0.5	0.8	1.5	2.4	3.6	7.6	18.4	56.8
2001.....	3.3	2.6	0.5	0.2	0.4	0.8	1.5	2.3	3.6	7.3	18.9	58.5
2000.....	3.3	1.8	0.5	0.2	0.4	0.8	1.4	2.3	3.7	7.6	18.8	58.9
1999.....	3.2	2.5	0.5	0.2	0.5	0.8	1.4	2.1	3.3	7.3	18.2	60.3
All races 3/, male												
2004.....	2.8	2.8	0.5	0.2	0.5	0.8	1.5	2.2	3.7	7.3	18.1	52.1
2003.....	2.9	2.9	0.5	0.2	0.4	0.8	1.5	2.4	3.4	7.9	18.8	55.1
2002.....	2.9	2.5	0.5	0.2	0.5	0.9	1.4	2.3	3.7	7.7	18.9	57.2
2001.....	2.8	2.9	0.6	0.2	0.4	0.9	1.5	2.2	3.3	7.2	20.0	58.1
2000.....	2.8	2.1	0.5	0.3	0.4	0.8	1.4	2.3	3.5	8.1	19.1	60.4
1999.....	2.8	3.0	0.6	0.2	0.5	0.9	1.5	2.1	3.1	7.5	19.3	63.2
All races 3/, female												
2004.....	3.6	1.8	0.4	0.2	0.4	0.8	1.3	2.1	3.3	6.8	16.8	53.3
2003.....	3.7	2.3	0.4	0.2	0.4	0.8	1.3	2.2	3.6	6.9	16.7	57.5
2002.....	3.8	1.5	0.5	0.2	0.4	0.7	1.5	2.5	3.5	7.5	18.1	56.6
2001.....	3.9	2.4	0.3	0.2	0.4	0.7	1.5	2.3	3.8	7.5	18.2	58.7
2000.....	3.8	1.6	0.6	0.2	0.4	0.8	1.5	2.2	3.8	7.2	18.6	58.3
1999.....	3.7	1.9	0.4	0.2	0.5	0.8	1.4	2.1	3.5	7.1	17.4	59.2
White, both sexes												
2004.....	3.0	2.0	0.3	0.2	0.2	0.4	0.7	1.4	2.7	6.4	17.2	52.6
2003.....	3.0	2.3	0.4	0.2	0.2	0.4	0.7	1.3	2.6	6.7	17.1	56.4
2002.....	3.1	1.9	0.4	0.1	0.3	0.4	0.7	1.4	2.8	7.0	17.8	56.2
2001.....	3.1	2.3	0.4	0.2	0.2	0.4	0.7	1.3	2.9	6.7	18.4	58.3
2000.....	3.1	1.7	0.4	0.1	0.2	0.3	0.7	1.4	2.9	7.0	18.4	58.8
1999.....	3.0	1.8	0.4	0.2	0.3	0.3	0.7	1.3	2.5	6.6	17.8	59.6
White, male												
2004.....	2.6	2.2	0.3	0.2	0.3	0.4	0.8	1.5	2.9	6.8	17.9	52.5
2003.....	2.6	2.8	0.3	0.2	0.2	0.4	0.8	1.4	2.5	7.3	18.5	55.1
2002.....	2.6	2.5	0.5	0.1	0.3	0.5	0.8	1.5	3.0	7.1	18.3	56.3
2001.....	2.6	2.4	0.6	0.2	0.2	0.4	0.7	1.4	2.8	6.6	19.6	58.9
2000.....	2.5	1.7	0.4	0.2	0.2	0.3	0.8	1.4	2.9	7.6	18.7	59.7
1999.....	2.5	2.2	0.5	0.2	0.3	0.4	0.8	1.4	2.4	6.9	18.8	63.6

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
III. Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)												
White, female												
2004.....	3.3	1.7	*	0.1	0.2	0.3	0.6	1.2	2.4	6.1	16.7	52.6
2003.....	3.5	1.7	0.4	0.1	0.2	0.4	0.6	1.2	2.6	6.2	16.2	57.0
2002.....	3.6	*	0.4	0.2	0.2	0.4	0.7	1.4	2.6	6.9	17.5	56.2
2001.....	3.6	2.1	*	*	0.2	0.4	0.7	1.2	3.0	6.9	17.5	58.0
2000.....	3.6	1.7	0.4	*	0.2	0.3	0.7	1.3	2.9	6.5	18.2	58.4
1999.....	3.4	1.4	*	0.2	0.3	0.3	0.6	1.2	2.7	6.4	17.2	58.0
Black, both sexes												
2004.....	5.7	3.5	0.9	0.5	1.6	3.5	6.0	8.7	11.9	14.0	22.1	65.7
2003.....	6.0	3.0	0.9	0.5	1.5	3.4	6.4	9.6	12.0	15.0	26.1	69.4
2002.....	6.0	*	0.8	0.5	1.5	3.5	6.5	9.8	11.2	14.9	28.1	71.9
2001.....	5.9	3.5	*	0.4	1.6	3.6	6.9	10.1	10.0	14.3	27.9	64.8
2000.....	5.8	*	1.2	0.6	1.4	3.9	6.3	9.8	11.0	13.8	26.7	66.6
1999.....	5.7	5.5	1.4	0.6	1.8	4.0	6.6	8.5	10.7	13.7	24.9	71.1
Black, male												
2004.....	5.3	*	*	*	1.7	3.8	6.3	8.3	12.6	14.8	23.1	61.6
2003.....	5.6	*	*	*	1.5	3.4	6.7	10.0	12.3	15.6	27.9	66.7
2002.....	5.5	*	*	*	1.9	4.2	6.1	9.4	11.1	15.3	29.0	78.3
2001.....	5.2	*	*	*	1.7	4.3	6.7	9.7	9.0	14.2	28.8	52.2
2000.....	5.1	*	*	0.6	1.3	3.8	6.0	9.9	9.7	13.0	25.7	73.8
1999.....	5.3	6.5	*	0.6	2.0	4.4	6.6	8.2	10.3	13.8	27.4	63.3
Black, female												
2004.....	6.0	*	*	*	1.6	3.2	5.8	9.0	11.3	13.5	21.6	67.3
2003.....	6.3	*	*	*	1.5	3.4	6.2	9.2	11.8	14.6	25.0	70.5
2002.....	6.4	*	*	*	1.1	2.8	6.9	10.1	11.2	14.7	27.6	69.4
2001.....	6.4	*	*	*	1.4	2.9	7.1	10.5	10.8	14.4	27.4	69.5
2000.....	6.4	*	*	*	1.4	4.0	6.6	9.7	12.0	14.3	27.3	64.0
1999.....	6.2	*	*	*	1.5	3.6	6.6	8.7	11.1	13.6	23.6	74.0

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Anemias (D50-D64)												
All races 3/, both sexes												
2004.....	1.6	*	0.2	0.1	0.3	0.4	0.5	0.6	1.1	2.6	8.6	38.1
2003.....	1.6	0.5	0.2	0.1	0.3	0.4	0.4	0.6	1.0	2.6	8.8	40.7
2002.....	1.6	*	0.1	0.1	0.3	0.5	0.4	0.6	1.0	2.8	9.1	40.3
2001.....	1.6	*	0.1	0.1	0.3	0.4	0.5	0.6	1.0	2.7	9.8	42.0
2000.....	1.6	*	0.2	0.1	0.2	0.4	0.4	0.6	1.1	2.8	9.5	42.4
1999.....	1.6	*	0.2	0.1	0.3	0.4	0.4	0.6	1.0	2.6	9.5	43.8
All races 3/, male												
2004.....	1.3	*	*	0.1	0.3	0.4	0.5	0.5	1.2	2.7	8.9	36.5
2003.....	1.3	*	*	0.1	0.2	0.4	0.4	0.6	1.2	3.0	9.3	38.0
2002.....	1.3	*	*	0.1	0.3	0.5	0.4	0.6	1.0	2.9	8.9	38.3
2001.....	1.3	*	*	0.1	0.3	0.5	0.4	0.6	0.9	2.9	9.9	40.0
2000.....	1.2	*	*	0.1	0.2	0.3	0.3	0.6	1.1	2.8	9.5	40.1
1999.....	1.3	*	*	0.1	0.3	0.4	0.5	0.6	0.9	2.7	10.1	43.4
All races 3/, female												
2004.....	1.8	*	*	0.1	0.3	0.4	0.4	0.6	1.0	2.5	8.4	38.8
2003.....	1.9	*	*	0.1	0.3	0.4	0.4	0.6	0.8	2.3	8.5	41.9
2002.....	1.9	*	*	0.1	0.2	0.4	0.4	0.7	1.0	2.6	9.2	41.2
2001.....	2.0	*	*	0.1	0.3	0.3	0.5	0.6	1.0	2.5	9.6	42.9
2000.....	2.0	*	*	0.1	0.2	0.4	0.5	0.6	1.0	2.8	9.4	43.3
1999.....	1.9	*	*	0.2	0.3	0.4	0.4	0.6	1.0	2.6	9.2	44.0
White, both sexes												
2004.....	1.5	*	*	0.1	0.1	0.1	0.2	0.3	0.7	2.4	8.5	37.7
2003.....	1.5	*	*	0.1	0.1	0.1	0.1	0.3	0.6	2.4	8.6	40.6
2002.....	1.5	*	*	0.1	0.1	0.1	0.1	0.3	0.7	2.5	8.7	39.7
2001.....	1.5	*	*	0.1	0.1	0.1	0.1	0.3	0.8	2.5	9.4	41.7
2000.....	1.5	*	*	0.1	0.1	0.1	0.1	0.3	0.8	2.5	9.3	42.0
1999.....	1.5	*	*	0.1	0.1	*	0.1	0.3	0.6	2.4	9.4	43.0
White, male												
2004.....	1.1	*	*	*	*	*	0.2	0.3	0.8	2.5	8.6	36.7
2003.....	1.1	*	*	*	*	0.1	0.1	0.3	0.7	2.6	9.1	37.9
2002.....	1.1	*	*	*	*	0.1	0.1	0.3	0.8	2.6	8.5	37.2
2001.....	1.1	*	*	*	*	0.1	0.1	0.2	0.7	2.6	9.8	40.4
2000.....	1.1	*	*	*	*	*	0.1	0.3	0.9	2.7	9.2	39.1
1999.....	1.1	*	*	*	*	*	0.1	0.3	0.5	2.5	9.9	43.4

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Anemias (D50-D64)												
White, female												
2004.....	1.8	*	*	*	*	*	0.1	0.2	0.7	2.2	8.4	38.2
2003.....	1.8	*	*	*	*	*	0.1	0.2	0.6	2.2	8.2	41.8
2002.....	1.8	*	*	*	*	*	*	0.3	0.7	2.4	8.8	40.8
2001.....	1.9	*	*	*	*	*	0.2	0.3	0.8	2.4	9.1	42.3
2000.....	1.9	*	*	*	*	*	0.1	0.3	0.8	2.4	9.3	43.2
1999.....	1.9	*	*	*	0.1	*	0.1	0.3	0.7	2.3	9.0	42.9
Black, both sexes												
2004.....	2.7	*	*	0.3	1.4	2.3	2.4	2.7	4.0	4.8	11.8	48.2
2003.....	2.7	*	*	0.4	1.3	2.1	2.3	3.0	4.2	5.6	14.5	49.3
2002.....	2.8	*	*	0.4	1.3	2.4	2.5	3.1	3.6	5.9	16.0	52.3
2001.....	2.7	*	*	0.4	1.2	2.2	2.7	3.1	3.0	5.5	16.0	48.5
2000.....	2.6	*	*	0.5	1.1	2.3	2.4	3.0	3.4	5.4	13.3	51.0
1999.....	2.8	*	0.8	0.5	1.4	2.6	2.5	3.1	4.2	5.0	13.4	52.9
Black, male												
2004.....	2.6	*	*	*	1.5	2.7	2.7	2.5	4.9	5.0	13.9	42.7
2003.....	2.7	*	*	*	1.3	2.2	2.6	3.3	5.4	7.9	14.9	49.0
2002.....	2.7	*	*	*	1.6	2.8	2.5	2.6	3.9	7.3	16.9	56.8
2001.....	2.5	*	*	*	1.4	2.9	2.4	3.6	3.1	6.7	15.0	42.0
2000.....	2.3	*	*	*	1.0	2.1	2.0	3.3	3.4	4.5	13.9	55.4
1999.....	2.7	*	*	*	1.7	2.8	2.7	3.3	5.1	5.4	12.9	44.5
Black, female												
2004.....	2.8	*	*	*	1.3	2.0	2.2	2.9	3.2	4.6	10.5	50.4
2003.....	2.8	*	*	*	1.2	2.0	2.1	2.8	3.2	3.9	14.2	49.4
2002.....	3.0	*	*	*	0.9	2.0	2.5	3.6	3.3	5.0	15.4	50.5
2001.....	2.8	*	*	*	1.1	1.6	2.9	2.6	2.8	4.6	16.5	50.9
2000.....	3.0	*	*	*	1.1	2.4	2.8	2.8	3.4	6.0	13.0	49.4
1999.....	3.0	*	*	*	1.2	2.3	2.3	3.0	3.6	4.7	13.6	56.1

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Nutritional anemias (D50-D53)												
All races 3/, both sexes												
2004.....	0.1	*	*	*	*	*	*	*	*	*	0.5	2.8
2003.....	0.1	*	*	*	*	*	*	*	*	0.2	0.5	3.5
2002.....	0.1	*	*	*	*	*	*	*	*	0.1	0.4	3.5
2001.....	0.1	*	*	*	*	*	*	*	*	0.2	0.6	3.6
2000.....	0.1	*	*	*	*	*	*	*	*	*	0.6	4.3
1999.....	0.1	*	*	*	*	*	*	*	*	0.1	0.7	4.0
All races 3/, male												
2004.....	0.1	*	*	*	*	*	*	*	*	*	0.4	2.7
2003.....	0.1	*	*	*	*	*	*	*	*	*	0.6	2.8
2002.....	0.1	*	*	*	*	*	*	*	*	*	*	2.8
2001.....	0.1	*	*	*	*	*	*	*	*	*	0.7	3.2
2000.....	0.1	*	*	*	*	*	*	*	*	*	0.4	5.1
1999.....	0.1	*	*	*	*	*	*	*	*	*	0.7	3.9
All races 3/, female												
2004.....	0.1	*	*	*	*	*	*	*	*	*	0.6	2.8
2003.....	0.1	*	*	*	*	*	*	*	*	*	0.5	3.8
2002.....	0.1	*	*	*	*	*	*	*	*	*	0.5	3.8
2001.....	0.1	*	*	*	*	*	*	*	*	*	0.6	3.8
2000.....	0.1	*	*	*	*	*	*	*	*	*	0.7	4.0
1999.....	0.1	*	*	*	*	*	*	*	*	*	0.7	4.0
White, both sexes												
2004.....	0.1	*	*	*	*	*	*	*	*	*	0.6	2.7
2003.....	0.1	*	*	*	*	*	*	*	*	0.1	0.5	3.6
2002.....	0.1	*	*	*	*	*	*	*	*	*	0.4	3.6
2001.....	0.1	*	*	*	*	*	*	*	*	0.2	0.6	3.7
2000.....	0.1	*	*	*	*	*	*	*	*	*	0.5	4.4
1999.....	0.1	*	*	*	*	*	*	*	*	0.1	0.6	3.9
White, male												
2004.....	0.1	*	*	*	*	*	*	*	*	*	*	3.0
2003.....	0.1	*	*	*	*	*	*	*	*	*	0.6	2.9
2002.....	0.1	*	*	*	*	*	*	*	*	*	*	2.9
2001.....	0.1	*	*	*	*	*	*	*	*	*	0.6	3.3
2000.....	0.1	*	*	*	*	*	*	*	*	*	*	5.0
1999.....	0.1	*	*	*	*	*	*	*	*	*	0.6	4.1

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hemolytic anemias (D55-D59)												
White, female												
2004.....	0.1	*	*	*	*	*	*	*	*	0.2	0.5	1.9
2003.....	0.2	*	*	*	*	*	*	*	*	*	0.8	2.2
2002.....	0.2	*	*	*	*	*	*	*	*	0.3	0.9	2.2
2001.....	0.2	*	*	*	*	*	*	*	*	0.3	0.8	2.2
2000.....	0.2	*	*	*	*	*	*	*	*	0.3	0.9	2.4
1999.....	0.2	*	*	*	*	*	*	*	*	0.4	0.8	2.6
Black, both sexes												
2004.....	1.5	*	*	*	1.3	2.3	2.1	2.1	2.0	1.2	*	*
2003.....	1.3	*	*	0.3	1.1	2.0	2.1	2.3	1.9	*	*	*
2002.....	1.4	*	*	0.3	1.1	2.3	2.2	2.5	1.3	*	*	*
2001.....	1.4	*	*	*	1.0	2.0	2.3	2.4	1.5	*	*	*
2000.....	1.4	*	*	0.4	0.9	2.0	2.1	2.3	1.6	*	*	*
1999.....	1.5	*	0.8	0.4	1.2	2.4	2.2	2.2	2.0	*	*	*
Black, male												
2004.....	1.5	*	*	*	1.4	2.6	2.4	1.9	2.3	*	*	*
2003.....	1.4	*	*	*	1.0	2.1	2.3	2.3	2.3	*	*	*
2002.....	1.4	*	*	*	1.4	2.6	2.3	2.0	*	*	*	*
2001.....	1.4	*	*	*	1.2	2.6	2.1	2.7	*	*	*	*
2000.....	1.3	*	*	*	0.9	1.8	1.9	2.4	*	*	*	*
1999.....	1.7	*	*	*	1.7	2.7	2.5	2.2	2.1	*	*	*
Black, female												
2004.....	1.4	*	*	*	1.2	1.9	1.9	2.3	1.7	*	*	*
2003.....	1.3	*	*	*	1.2	1.9	1.9	2.2	1.6	*	*	*
2002.....	1.4	*	*	*	0.9	1.9	2.2	2.8	1.4	*	*	*
2001.....	1.3	*	*	*	0.8	1.4	2.4	2.2	1.5	*	*	*
2000.....	1.4	*	*	*	0.9	2.1	2.3	2.3	2.0	*	*	*
1999.....	1.3	*	*	*	0.8	2.1	1.9	2.2	1.9	*	*	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Aplastic anemias (D60-D61)												
All races 3/, both sexes												
2004.....	0.3	*	*	*	0.1	*	0.1	0.1	0.4	0.9	2.0	5.6
2003.....	0.3	*	*	*	0.1	0.1	0.1	0.1	0.3	0.9	2.2	5.7
2002.....	0.3	*	*	0.1	0.1	0.1	0.1	0.1	0.3	0.8	2.4	5.3
2001.....	0.4	*	*	0.1	0.1	0.1	0.1	0.2	0.4	0.9	2.7	6.8
2000.....	0.4	*	*	*	0.1	0.1	0.1	0.2	0.4	1.0	2.5	7.4
1999.....	0.4	*	*	0.1	0.1	*	0.1	0.2	0.3	1.0	2.6	7.7
All races 3/, male												
2004.....	0.3	*	*	*	*	*	0.1	0.1	0.4	1.0	2.3	6.2
2003.....	0.3	*	*	*	*	0.1	*	0.1	0.3	1.1	2.6	7.4
2002.....	0.3	*	*	*	*	*	*	0.1	0.3	0.8	2.8	7.0
2001.....	0.3	*	*	*	*	*	*	0.2	0.4	0.9	2.9	7.8
2000.....	0.3	*	*	*	*	*	*	0.1	0.4	1.1	2.8	7.6
1999.....	0.4	*	*	*	*	*	*	0.2	0.3	1.1	3.1	10.1
All races 3/, female												
2004.....	0.4	*	*	*	*	*	*	0.1	0.4	0.8	1.8	5.4
2003.....	0.3	*	*	*	*	*	*	0.1	0.3	0.7	1.9	5.0
2002.....	0.3	*	*	*	*	*	*	0.1	0.3	0.9	2.1	4.5
2001.....	0.4	*	*	*	*	*	*	0.2	0.4	0.9	2.5	6.3
2000.....	0.4	*	*	*	*	*	0.1	0.2	0.4	0.9	2.4	7.3
1999.....	0.4	*	*	*	*	*	*	0.2	0.3	0.9	2.3	6.8
White, both sexes												
2004.....	0.4	*	*	*	0.1	*	0.1	0.1	0.4	0.9	2.0	5.6
2003.....	0.3	*	*	*	*	0.1	0.1	0.1	0.3	0.9	2.2	5.8
2002.....	0.3	*	*	*	*	*	0.1	0.1	0.3	0.8	2.5	5.2
2001.....	0.4	*	*	0.1	0.1	*	0.1	0.2	0.4	0.9	2.7	7.1
2000.....	0.4	*	*	*	*	*	0.1	0.1	0.4	1.0	2.6	7.4
1999.....	0.4	*	*	*	0.1	*	0.1	0.1	0.3	0.9	2.7	7.8
White, male												
2004.....	0.3	*	*	*	*	*	*	0.1	0.3	1.0	2.3	6.2
2003.....	0.3	*	*	*	*	*	*	0.1	0.3	1.0	2.7	7.6
2002.....	0.3	*	*	*	*	*	*	*	0.3	0.8	2.9	6.9
2001.....	0.3	*	*	*	*	*	*	0.2	0.3	0.8	3.0	8.5
2000.....	0.3	*	*	*	*	*	*	0.1	0.4	1.1	2.7	7.0
1999.....	0.4	*	*	*	*	*	*	0.1	0.2	1.0	3.1	10.3

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Acute posthemorrhagic and other anemias (D62,D64)

All races 3/, both sexes

2004.....	0.9	*	*	*	*	*	0.1	0.1	0.4	1.3	5.5	27.8
2003.....	0.9	*	*	*	*	*	*	0.1	0.4	1.3	5.3	29.6
2002.....	0.9	*	*	*	*	*	*	0.1	0.4	1.4	5.5	29.4
2001.....	0.8	*	*	*	*	*	0.1	0.1	0.3	1.3	5.6	29.5
2000.....	0.8	*	*	*	*	*	*	0.1	0.4	1.4	5.5	28.2
1999.....	0.8	*	*	*	*	*	0.0	0.2	0.3	1.2	5.3	29.9

All races 3/, male

2004.....	0.6	*	*	*	*	*	*	0.2	0.5	1.4	5.3	25.4
2003.....	0.6	*	*	*	*	*	*	0.2	0.5	1.5	5.2	26.2
2002.....	0.6	*	*	*	*	*	*	0.1	0.5	1.6	5.3	26.3
2001.....	0.6	*	*	*	*	*	*	*	0.3	1.5	5.4	26.9
2000.....	0.6	*	*	*	*	*	*	0.1	0.5	1.4	5.5	24.9
1999.....	0.6	*	*	*	*	*	*	0.2	0.3	1.2	5.2	27.7

All races 3/, female

2004.....	1.1	*	*	*	*	*	*	0.1	0.3	1.2	5.6	28.9
2003.....	1.1	*	*	*	*	*	*	0.1	0.3	1.2	5.4	31.1
2002.....	1.1	*	*	*	*	*	*	0.1	0.4	1.2	5.7	30.7
2001.....	1.1	*	*	*	*	*	*	0.1	0.3	1.1	5.7	30.5
2000.....	1.1	*	*	*	*	*	*	0.1	0.3	1.4	5.5	29.6
1999.....	1.1	*	*	*	*	*	*	0.2	0.3	1.2	5.4	30.8

White, both sexes

2004.....	0.9	*	*	*	*	*	*	0.1	0.3	1.2	5.2	27.4
2003.....	0.9	*	*	*	*	*	*	0.1	0.3	1.2	5.0	29.2
2002.....	0.9	*	*	*	*	*	*	0.1	0.3	1.2	5.0	28.7
2001.....	0.9	*	*	*	*	*	*	*	0.2	1.2	5.1	28.8
2000.....	0.8	*	*	*	*	*	*	0.1	0.3	1.2	5.2	27.8
1999.....	0.8	*	*	*	*	*	*	0.1	0.2	1.0	5.1	28.9

White, male

2004.....	0.6	*	*	*	*	*	*	0.1	0.3	1.3	4.9	25.4
2003.....	0.6	*	*	*	*	*	*	*	0.4	1.2	4.9	25.9
2002.....	0.6	*	*	*	*	*	*	*	0.3	1.4	4.7	25.1
2001.....	0.6	*	*	*	*	*	*	*	0.2	1.3	5.1	26.4
2000.....	0.6	*	*	*	*	*	*	*	0.3	1.2	5.3	24.3
1999.....	0.6	*	*	*	*	*	*	*	*	1.1	5.1	27.2

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute posthemorrhagic and other anemias (D62,D64)												
White, female												
2004.....	1.1	*	*	*	*	*	*	*	0.2	1.1	5.4	28.3
2003.....	1.2	*	*	*	*	*	*	*	0.2	1.1	5.1	30.7
2002.....	1.2	*	*	*	*	*	*	*	0.3	1.1	5.2	30.2
2001.....	1.1	*	*	*	*	*	*	*	0.3	1.0	5.2	29.8
2000.....	1.1	*	*	*	*	*	*	*	0.3	1.2	5.2	29.2
1999.....	1.1	*	*	*	*	*	*	*	0.3	1.0	5.2	29.6
Black, both sexes												
2004.....	0.9	*	*	*	*	*	*	0.5	1.5	2.5	9.1	37.8
2003.....	1.0	*	*	*	*	*	*	0.5	1.6	3.2	10.5	39.7
2002.....	1.0	*	*	*	*	*	*	*	1.5	3.2	12.9	41.6
2001.....	1.0	*	*	*	*	*	*	0.5	0.8	2.9	11.9	41.1
2000.....	0.9	*	*	*	*	*	*	*	1.3	3.3	9.5	37.2
1999.....	0.9	*	*	*	*	*	*	0.7	1.2	2.8	8.7	39.9
Black, male												
2004.....	0.7	*	*	*	*	*	*	*	1.9	3.2	10.0	33.8
2003.....	0.9	*	*	*	*	*	*	*	2.4	4.6	12.1	37.5
2002.....	0.9	*	*	*	*	*	*	*	1.9	4.4	13.8	45.0
2001.....	0.7	*	*	*	*	*	*	*	*	3.6	10.9	37.4
2000.....	0.7	*	*	*	*	*	*	*	*	*	9.1	33.4
1999.....	0.6	*	*	*	*	*	*	*	*	*	8.3	31.6
Black, female												
2004.....	1.0	*	*	*	*	*	*	*	*	2.1	8.7	39.4
2003.....	1.1	*	*	*	*	*	*	*	*	2.1	9.6	40.5
2002.....	1.2	*	*	*	*	*	*	*	*	2.3	12.3	40.3
2001.....	1.2	*	*	*	*	*	*	*	*	2.5	12.4	42.5
2000.....	1.1	*	*	*	*	*	*	*	*	3.7	9.7	38.6
1999.....	1.1	*	*	*	*	*	*	*	*	3.0	9.0	42.9

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Coagulation defects, purpura and other hemorrhagic conditions (D65-D69)

All races 3/, both sexes

2004.....	0.9	0.9	*	*	0.1	0.2	0.3	0.7	1.2	2.5	6.0	10.1
2003.....	0.9	1.0	*	*	0.1	0.2	0.3	0.7	1.1	2.8	5.9	10.5
2002.....	0.9	0.6	*	*	0.1	0.2	0.3	0.7	1.1	2.8	6.0	11.3
2001.....	0.9	0.8	*	*	0.1	0.2	0.4	0.7	1.2	2.7	6.0	11.1
2000.....	0.9	0.6	*	*	0.1	0.1	0.3	0.7	1.4	2.9	6.1	11.6
1999.....	0.9	1.1	*	0.1	0.1	0.1	0.4	0.6	1.2	2.9	6.0	11.5

All races 3/, male

2004.....	0.8	1.2	*	*	*	0.1	0.3	0.7	1.4	2.8	6.6	10.9
2003.....	0.8	1.1	*	*	*	0.1	0.3	0.7	1.2	3.0	6.5	11.9
2002.....	0.8	*	*	*	*	0.2	0.3	0.6	1.2	2.7	6.7	13.9
2001.....	0.8	*	*	*	*	0.1	0.3	0.7	1.2	2.8	6.7	12.3
2000.....	0.8	*	*	*	*	0.1	0.3	0.7	1.2	3.2	6.2	14.6
1999.....	0.8	1.4	*	*	*	0.1	0.4	0.7	1.1	3.1	6.2	14.3

All races 3/, female

2004.....	1.0	*	*	*	*	0.2	0.4	0.7	1.0	2.2	5.6	9.7
2003.....	1.0	*	*	*	*	0.2	0.3	0.7	1.1	2.6	5.4	9.9
2002.....	1.0	*	*	*	0.1	0.1	0.3	0.7	1.1	2.9	5.6	10.1
2001.....	1.0	*	*	*	*	0.2	0.4	0.7	1.3	2.6	5.6	10.5
2000.....	1.0	*	*	*	0.1	0.2	0.3	0.6	1.5	2.7	6.0	10.4
1999.....	1.0	*	*	*	*	0.2	0.4	0.6	1.3	2.8	5.8	10.3

White, both sexes

2004.....	0.9	0.8	*	*	*	0.1	0.3	0.6	1.1	2.3	6.1	10.0
2003.....	0.9	0.9	*	*	0.1	0.1	0.3	0.6	1.0	2.7	5.7	10.3
2002.....	1.0	*	*	*	0.1	0.2	0.3	0.6	1.0	2.7	5.9	11.3
2001.....	0.9	0.7	*	*	*	0.1	0.3	0.5	1.2	2.6	5.9	11.0
2000.....	0.9	*	*	*	0.1	0.1	0.3	0.6	1.2	2.8	5.9	11.8
1999.....	0.9	0.9	*	*	0.1	0.1	0.3	0.5	1.1	2.7	5.8	11.4

White, male

2004.....	0.9	*	*	*	*	0.1	0.3	0.6	1.3	2.7	6.6	11.0
2003.....	0.8	*	*	*	*	*	0.3	0.6	1.0	2.9	6.4	11.8
2002.....	0.9	*	*	*	*	0.2	0.3	0.5	1.1	2.7	6.5	14.1
2001.....	0.8	*	*	*	*	0.1	0.3	0.5	1.2	2.7	6.5	12.5
2000.....	0.9	*	*	*	*	*	0.3	0.6	1.1	3.0	6.2	14.8
1999.....	0.8	*	*	*	*	*	0.3	0.6	1.0	2.8	5.8	14.6

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Coagulation defects, purpura and other hemorrhagic conditions (D65-D69)

White, female

2004.....	1.0	*	*	*	*	0.1	0.3	0.6	0.9	2.1	5.7	9.5
2003.....	1.0	*	*	*	*	0.1	0.2	0.6	1.0	2.5	5.2	9.7
2002.....	1.0	*	*	*	*	0.1	0.3	0.6	1.0	2.8	5.5	10.1
2001.....	1.0	*	*	*	*	0.2	0.3	0.5	1.2	2.6	5.5	10.4
2000.....	1.0	*	*	*	*	*	0.3	0.5	1.3	2.6	5.8	10.5
1999.....	1.0	*	*	*	*	*	0.2	0.5	1.2	2.5	5.8	10.2

Black, both sexes

2004.....	1.1	*	*	*	*	*	0.9	1.5	2.3	4.0	6.6	13.0
2003.....	1.1	*	*	*	*	*	0.6	1.8	2.2	4.3	7.6	13.7
2002.....	1.0	*	*	*	*	*	0.4	1.6	2.0	4.6	7.7	13.1
2001.....	1.1	*	*	*	*	0.5	0.9	1.8	2.1	3.5	7.9	12.3
2000.....	1.1	*	*	*	*	0.4	0.6	1.7	2.6	4.4	8.2	11.3
1999.....	1.2	*	*	*	*	*	1.0	1.4	2.6	5.0	8.4	13.7

Black, male

2004.....	1.0	*	*	*	*	*	*	1.6	2.6	4.7	7.2	*
2003.....	1.0	*	*	*	*	*	*	1.8	2.5	4.9	7.9	*
2002.....	0.9	*	*	*	*	*	*	1.6	2.0	4.1	8.6	*
2001.....	0.9	*	*	*	*	*	0.8	1.9	2.0	3.9	8.8	*
2000.....	0.9	*	*	*	*	*	*	1.8	2.3	5.3	6.7	*
1999.....	1.1	*	*	*	*	*	0.8	1.4	2.5	5.4	11.1	*

Black, female

2004.....	1.1	*	*	*	*	*	1.1	1.4	2.1	3.5	6.3	12.6
2003.....	1.2	*	*	*	*	*	0.7	1.8	2.0	3.8	7.5	13.8
2002.....	1.1	*	*	*	*	*	*	1.6	2.1	5.0	7.1	12.3
2001.....	1.2	*	*	*	*	*	0.9	1.8	2.2	3.3	7.4	13.9
2000.....	1.2	*	*	*	*	*	0.7	1.6	2.9	3.8	9.0	10.3
1999.....	1.3	*	*	*	*	*	1.1	1.5	2.7	4.7	6.9	13.1

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other diseases of blood and blood-forming organs (D70-D76)

All races 3/, both sexes

2004.....	0.3	0.6	0.1	*	*	*	0.1	0.2	0.3	0.8	1.6	3.0
2003.....	0.3	0.5	0.1	0.1	*	0.1	0.1	0.2	0.4	0.8	1.7	3.9
2002.....	0.3	*	0.2	*	*	*	0.1	0.2	0.4	0.9	2.0	3.4
2001.....	0.3	0.6	0.1	*	*	*	0.1	0.2	0.4	0.8	1.9	3.9
2000.....	0.3	*	*	*	0.1	0.1	0.1	0.2	0.4	0.9	1.8	3.4
1999.....	0.3	0.6	*	*	0.1	0.1	0.1	0.2	0.4	0.9	1.6	3.6

All races 3/, male

2004.....	0.3	*	*	*	*	*	0.1	0.2	0.4	0.9	1.8	3.2
2003.....	0.3	*	*	*	*	*	0.1	0.2	0.4	0.8	2.0	4.0
2002.....	0.3	*	0.3	*	*	*	0.1	0.2	0.5	1.1	2.2	3.5
2001.....	0.3	*	*	*	*	*	*	0.2	0.4	0.8	2.2	3.9
2000.....	0.3	*	*	*	*	*	0.1	0.2	0.4	1.1	2.2	4.1
1999.....	0.3	*	*	*	*	*	0.1	0.1	0.5	1.0	2.1	4.0

All races 3/, female

2004.....	0.3	*	*	*	*	*	*	0.2	0.3	0.8	1.4	2.9
2003.....	0.3	*	*	*	*	*	*	0.1	0.4	0.7	1.5	3.9
2002.....	0.3	*	*	*	*	*	0.1	0.2	0.4	0.7	1.9	3.3
2001.....	0.3	*	*	*	*	*	0.1	0.1	0.4	0.8	1.6	3.8
2000.....	0.3	*	*	*	*	*	*	0.2	0.4	0.7	1.6	3.1
1999.....	0.3	*	*	*	*	*	0.1	0.2	0.4	0.8	1.3	3.4

White, both sexes

2004.....	0.3	*	*	*	*	*	0.1	0.2	0.3	0.8	1.6	3.2
2003.....	0.3	*	*	*	*	*	0.1	0.2	0.4	0.8	1.7	4.1
2002.....	0.3	*	0.2	*	*	*	0.1	0.2	0.4	0.9	2.0	3.4
2001.....	0.3	*	*	*	*	*	0.1	0.2	0.4	0.8	1.9	3.9
2000.....	0.3	*	*	*	*	*	0.1	0.2	0.4	0.9	1.9	3.5
1999.....	0.3	*	*	*	*	*	0.1	0.2	0.4	0.9	1.7	3.7

White, male

2004.....	0.3	*	*	*	*	*	*	0.2	0.4	0.9	1.9	3.4
2003.....	0.3	*	*	*	*	*	*	0.2	0.4	0.9	2.0	4.4
2002.....	0.3	*	*	*	*	*	*	0.2	0.5	1.1	2.2	3.5
2001.....	0.3	*	*	*	*	*	*	0.2	0.4	0.8	2.3	4.1
2000.....	0.3	*	*	*	*	*	0.1	0.2	0.3	1.1	2.1	4.1
1999.....	0.3	*	*	*	*	*	0.1	*	0.5	1.0	2.2	4.0

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Certain disorders involving the immune mechanism (D80-D89)

All races 3/, both sexes

2004.....	0.5	*	*	*	0.1	0.2	0.5	0.8	0.9	1.2	1.1	1.7
2003.....	0.5	0.5	*	*	0.1	0.2	0.6	0.8	1.0	1.2	1.2	1.6
2002.....	0.5	0.5	*	*	*	0.2	0.6	0.9	1.0	1.1	1.3	1.8
2001.....	0.5	0.8	*	*	*	0.2	0.6	0.9	0.9	1.1	1.2	1.5
2000.....	0.5	*	*	*	*	0.2	0.6	0.8	0.9	0.9	1.5	1.6
1999.....	0.4	*	0.2	*	*	0.2	0.5	0.7	0.7	0.9	1.0	1.4

All races 3/, male

2004.....	0.4	*	*	*	*	0.2	0.6	0.8	0.8	1.0	0.8	1.5
2003.....	0.5	*	*	*	*	0.2	0.7	0.8	0.7	1.1	1.1	*
2002.....	0.5	*	*	*	*	0.2	0.6	0.9	1.0	0.9	1.1	1.5
2001.....	0.4	*	*	*	*	0.3	0.6	0.8	0.7	0.7	1.1	1.8
2000.....	0.5	*	*	*	*	0.2	0.6	0.8	0.8	0.9	1.3	1.6
1999.....	0.4	*	*	*	*	0.3	0.5	0.7	0.6	0.7	1.0	*

All races 3/, female

2004.....	0.5	*	*	*	*	0.2	0.5	0.7	1.1	1.4	1.4	1.9
2003.....	0.6	*	*	*	*	0.2	0.6	0.8	1.2	1.3	1.3	1.7
2002.....	0.6	*	*	*	*	0.1	0.6	0.9	1.1	1.2	1.4	1.9
2001.....	0.6	*	*	*	*	0.1	0.5	1.0	1.1	1.5	1.3	1.4
2000.....	0.5	*	*	*	*	0.2	0.6	0.8	0.9	1.0	1.6	1.5
1999.....	0.4	*	*	*	*	0.2	0.5	0.6	0.8	1.0	1.1	1.5

White, both sexes

2004.....	0.3	*	*	*	*	0.1	0.2	0.3	0.5	0.9	1.0	1.8
2003.....	0.3	*	*	*	*	0.1	0.2	0.3	0.5	0.9	1.1	1.5
2002.....	0.3	*	*	*	*	0.1	0.2	0.4	0.6	0.9	1.2	1.8
2001.....	0.3	0.8	*	*	*	0.1	0.2	0.3	0.6	0.9	1.2	1.6
2000.....	0.3	*	*	*	*	0.1	0.2	0.3	0.5	0.8	1.4	1.5
1999.....	0.3	*	*	*	*	0.1	0.2	0.3	0.5	0.7	1.0	1.4

White, male

2004.....	0.3	*	*	*	*	*	0.3	0.4	0.5	0.7	0.8	1.5
2003.....	0.3	*	*	*	*	*	0.3	0.3	0.4	1.0	1.0	*
2002.....	0.3	*	*	*	*	*	0.3	0.4	0.7	0.8	1.1	*
2001.....	0.3	*	*	*	*	0.2	0.2	0.4	0.5	0.6	1.0	2.0
2000.....	0.3	*	*	*	*	*	0.3	0.4	0.6	0.8	1.2	*
1999.....	0.3	*	*	*	*	0.2	0.2	0.4	0.5	0.6	1.0	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Certain disorders involving the immune mechanism (D80-D89)												
White, female												
2004.....	0.3	*	*	*	*	*	0.1	0.2	0.5	1.0	1.1	1.9
2003.....	0.3	*	*	*	*	*	0.1	0.3	0.6	0.8	1.2	1.7
2002.....	0.3	*	*	*	*	*	0.1	0.3	0.6	0.9	1.3	1.9
2001.....	0.3	*	*	*	*	*	0.1	0.3	0.6	1.1	1.3	1.5
2000.....	0.3	*	*	*	*	*	0.2	0.3	0.4	0.7	1.5	1.5
1999.....	0.3	*	*	*	*	*	*	0.2	0.4	0.8	1.0	1.4
Black, both sexes												
2004.....	1.7	*	*	*	*	0.9	2.6	4.2	5.0	4.4	3.1	*
2003.....	1.9	*	*	*	*	0.9	3.4	4.5	5.1	4.6	2.4	*
2002.....	1.9	*	*	*	*	0.7	3.5	4.9	5.0	3.7	2.5	*
2001.....	1.8	*	*	*	*	0.9	3.3	4.9	4.5	4.3	2.5	*
2000.....	1.8	*	*	*	*	1.0	3.0	4.8	4.4	2.8	3.6	*
1999.....	1.5	*	*	*	*	1.0	3.0	3.6	3.0	2.7	*	*
Black, male												
2004.....	1.6	*	*	*	*	0.9	2.7	4.0	4.3	4.2	*	*
2003.....	1.7	*	*	*	*	0.8	3.6	4.6	3.8	*	*	*
2002.....	1.7	*	*	*	*	0.9	3.1	5.0	4.4	*	*	*
2001.....	1.6	*	*	*	*	1.1	3.4	3.8	3.4	*	*	*
2000.....	1.6	*	*	*	*	1.1	3.1	4.5	3.3	*	*	*
1999.....	1.3	*	*	*	*	1.1	2.8	3.3	2.1	*	*	*
Black, female												
2004.....	1.9	*	*	*	*	0.8	2.4	4.4	5.5	4.6	4.2	*
2003.....	2.2	*	*	*	*	1.0	3.3	4.5	6.2	6.2	*	*
2002.....	2.1	*	*	*	*	*	3.8	4.8	5.5	4.5	3.2	*
2001.....	2.1	*	*	*	*	*	3.2	5.9	5.3	5.5	*	*
2000.....	1.9	*	*	*	*	1.0	2.9	5.0	5.2	3.5	4.1	*
1999.....	1.7	*	*	*	*	0.9	3.1	4.0	3.8	3.3	*	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
IV. Endocrine, nutritional and metabolic diseases (E00-E88)												
All races 3/, both sexes												
2004.....	33.7	5.7	0.9	0.5	1.2	3.0	7.1	18.8	47.1	107.4	226.8	470.6
2003.....	34.5	6.2	1.2	0.5	1.3	3.1	7.2	19.2	48.4	111.0	232.1	492.8
2002.....	34.5	6.6	0.9	0.5	1.3	3.1	7.4	18.7	47.9	111.1	235.3	504.8
2001.....	34.0	7.2	1.2	0.5	1.2	2.8	6.7	18.6	46.9	110.8	232.5	519.2
2000.....	33.5	6.8	1.0	0.5	1.3	2.9	6.6	17.8	46.9	109.9	232.0	521.0
1999.....	33.3	6.9	1.0	0.5	1.4	2.6	6.5	17.2	46.5	110.5	229.3	529.8
All races 3/, male												
2004.....	32.2	7.1	1.0	0.6	1.3	3.2	8.4	22.7	55.5	123.3	254.9	481.0
2003.....	32.5	6.8	1.4	0.6	1.4	3.6	8.5	22.8	56.5	127.3	259.6	490.8
2002.....	31.9	7.4	0.8	0.6	1.4	3.6	8.9	22.1	56.1	123.3	255.6	504.5
2001.....	30.9	8.0	1.2	0.6	1.2	3.0	7.8	22.0	53.6	124.1	251.9	507.7
2000.....	30.2	7.4	1.1	0.6	1.4	3.4	7.7	20.8	53.1	120.8	251.5	517.4
1999.....	29.8	7.3	1.1	0.5	1.3	2.9	7.6	20.3	52.3	120.9	250.5	521.0
All races 3/, female												
2004.....	35.2	4.2	0.8	0.5	1.2	2.7	5.8	14.9	39.2	94.0	207.8	465.9
2003.....	36.5	5.5	1.1	0.4	1.2	2.7	5.9	15.8	40.9	97.4	213.7	493.7
2002.....	37.0	5.8	1.0	0.4	1.2	2.6	5.9	15.5	40.2	100.9	221.9	504.9
2001.....	37.0	6.4	1.2	0.5	1.3	2.5	5.5	15.3	40.6	99.7	219.8	524.0
2000.....	36.7	6.2	1.0	0.5	1.2	2.3	5.6	14.8	41.2	100.8	219.4	522.5
1999.....	36.6	6.5	0.9	0.5	1.4	2.4	5.4	14.2	41.2	102.0	215.4	533.3
White, both sexes												
2004.....	33.7	5.8	0.9	0.5	1.2	2.6	6.2	16.7	42.1	97.7	213.6	453.5
2003.....	34.5	5.4	1.2	0.6	1.3	2.7	6.4	17.0	43.2	101.3	218.4	475.4
2002.....	34.4	6.2	1.0	0.6	1.3	2.8	6.7	16.5	42.4	100.3	221.8	487.1
2001.....	34.0	6.8	1.2	0.5	1.2	2.4	6.0	16.4	41.5	100.8	218.7	502.4
2000.....	33.5	6.3	1.1	0.6	1.3	2.5	6.1	15.5	41.8	99.8	217.5	503.5
1999.....	33.1	6.0	1.1	0.5	1.4	2.4	5.7	14.8	40.2	99.7	215.1	511.4
White, male												
2004.....	32.5	7.2	1.0	0.5	1.3	2.8	7.5	20.8	50.7	114.5	245.1	470.5
2003.....	32.9	6.3	1.4	0.7	1.4	3.1	7.8	20.6	51.6	118.7	249.4	484.1
2002.....	32.2	7.6	0.8	0.7	1.3	3.2	8.2	19.9	50.8	114.6	246.6	498.9
2001.....	31.2	7.2	1.3	0.6	1.2	2.6	7.1	19.6	48.5	116.4	240.7	500.5
2000.....	30.6	6.8	1.2	0.6	1.3	3.0	7.1	18.8	48.2	112.9	241.3	510.5
1999.....	29.9	6.3	1.2	0.5	1.3	2.6	6.8	18.1	46.7	112.3	240.3	504.7

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
IV. Endocrine, nutritional and metabolic diseases (E00-E88)												
White, female												
2004.....	34.9	4.3	0.9	0.5	1.2	2.4	4.9	12.7	34.0	83.3	192.1	445.8
2003.....	36.1	4.5	1.1	0.4	1.2	2.4	5.1	13.4	35.2	86.5	197.5	471.5
2002.....	36.6	4.7	1.1	0.4	1.3	2.4	5.2	13.1	34.5	88.3	205.2	482.1
2001.....	36.8	6.3	1.2	0.5	1.2	2.2	4.8	13.1	34.9	87.6	204.2	503.2
2000.....	36.3	5.7	1.0	0.5	1.2	1.9	5.0	12.3	35.7	88.8	201.8	500.6
1999.....	36.2	5.7	0.9	0.5	1.4	2.2	4.6	11.7	34.2	89.2	198.6	514.0
Black, both sexes												
2004.....	41.8	5.9	*	0.5	1.7	6.0	14.4	36.9	96.1	210.8	397.6	719.1
2003.....	42.3	9.9	1.0	0.4	1.7	6.3	13.8	38.6	100.5	213.0	406.3	736.9
2002.....	42.6	8.3	0.8	0.6	2.0	6.2	14.0	38.6	101.6	219.3	410.0	757.0
2001.....	41.6	10.3	0.8	0.5	1.8	5.5	12.8	38.3	98.8	217.2	410.3	763.1
2000.....	41.4	8.9	0.8	0.4	1.8	6.0	12.3	37.8	96.6	217.4	424.6	766.1
1999.....	41.4	11.4	0.8	0.6	1.6	5.0	13.1	37.9	104.0	221.8	406.2	763.6
Black, male												
2004.....	37.7	7.7	*	0.6	1.8	6.8	16.7	41.8	107.5	232.8	411.5	683.5
2003.....	37.1	8.9	*	*	1.6	7.6	15.3	42.8	111.2	229.5	420.0	619.3
2002.....	36.6	7.0	*	*	2.3	7.5	16.3	43.1	114.5	222.8	405.4	610.0
2001.....	35.9	11.7	*	*	1.7	6.5	14.4	43.3	108.1	222.3	425.2	658.1
2000.....	34.6	9.9	*	*	1.9	6.8	13.6	41.1	104.7	217.3	413.9	655.0
1999.....	35.2	12.7	*	0.6	1.6	5.6	14.7	41.9	109.6	223.0	396.3	746.4
Black, female												
2004.....	45.5	*	*	*	1.6	5.2	12.3	32.7	86.9	195.0	389.7	733.2
2003.....	47.0	10.8	*	*	1.8	5.2	12.4	35.0	92.0	201.2	398.5	782.6
2002.....	48.0	9.7	*	0.7	1.6	5.1	12.0	34.8	91.3	216.8	412.7	813.3
2001.....	46.8	8.8	*	*	1.8	4.6	11.4	34.1	91.4	213.5	401.9	802.0
2000.....	47.6	7.9	*	*	1.6	5.4	11.2	35.0	90.1	217.4	430.6	807.4
1999.....	47.1	10.1	*	*	1.6	4.5	11.7	34.5	99.5	220.9	411.7	770.1

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Endocrine diseases (E00-E34)												
All races 3/, both sexes												
2004.....	25.8	0.6	0.1	0.1	0.5	1.6	4.8	13.8	37.8	88.8	182.3	330.4
2003.....	26.5	0.7	0.1	0.1	0.5	1.8	4.8	14.2	39.3	92.5	187.3	342.7
2002.....	26.4	1.0	*	0.1	0.5	1.7	5.0	14.1	38.4	93.2	188.8	347.6
2001.....	26.1	1.1	*	0.1	0.4	1.6	4.6	14.0	38.5	93.0	187.7	350.3
2000.....	25.6	1.2	0.1	0.1	0.5	1.7	4.5	13.5	38.6	92.3	185.5	345.2
1999.....	25.5	1.0	0.1	0.1	0.5	1.6	4.5	13.4	39.0	93.7	183.9	343.6
All races 3/, male												
2004.....	25.0	*	*	0.1	0.5	1.8	5.9	16.8	44.6	101.5	206.9	349.1
2003.....	25.3	*	*	0.1	0.5	2.0	5.9	17.0	45.6	105.4	211.6	354.4
2002.....	24.8	1.1	*	0.1	0.5	2.0	6.1	16.7	45.1	102.8	206.6	365.2
2001.....	24.1	1.3	*	0.1	0.5	1.8	5.5	16.7	43.7	103.2	203.5	362.2
2000.....	23.4	1.3	*	*	0.5	2.0	5.3	15.5	43.0	100.4	203.5	360.9
1999.....	23.3	*	*	*	0.5	1.7	5.3	15.9	43.2	101.3	203.1	361.8
All races 3/, female												
2004.....	26.7	*	*	0.1	0.5	1.4	3.7	10.8	31.5	78.0	165.8	322.0
2003.....	27.7	*	*	0.1	0.4	1.6	3.8	11.6	33.4	81.8	171.0	337.6
2002.....	28.0	*	*	*	0.5	1.4	3.9	11.6	32.2	85.3	177.1	339.9
2001.....	28.0	*	*	*	0.4	1.4	3.7	11.4	33.7	84.6	177.3	345.3
2000.....	27.7	1.1	*	0.1	0.4	1.4	3.7	11.5	34.5	85.6	173.8	338.8
1999.....	27.5	1.3	*	0.1	0.5	1.5	3.8	11.0	35.0	87.4	171.3	336.3
White, both sexes												
2004.....	25.3	0.7	*	0.1	0.4	1.4	4.2	11.9	33.0	79.5	169.5	313.3
2003.....	26.1	*	*	0.1	0.4	1.5	4.2	12.2	34.2	83.0	174.6	326.4
2002.....	26.0	1.1	*	0.1	0.4	1.5	4.4	12.0	33.3	83.2	176.5	331.6
2001.....	25.7	1.1	*	0.1	0.3	1.3	4.0	12.1	33.4	83.5	175.1	335.2
2000.....	25.1	1.0	*	0.1	0.4	1.4	4.0	11.4	33.8	82.7	171.8	329.3
1999.....	24.9	1.0	*	0.1	0.4	1.4	3.9	11.3	33.0	83.6	170.8	325.9
White, male												
2004.....	24.9	*	*	*	0.4	1.5	5.2	14.9	39.8	92.9	197.4	338.1
2003.....	25.3	*	*	*	0.5	1.6	5.2	14.9	40.7	97.1	202.8	347.0
2002.....	24.8	1.3	*	0.1	0.4	1.8	5.4	14.5	40.1	94.6	198.7	361.5
2001.....	24.1	*	*	0.1	0.4	1.4	4.9	14.6	39.0	95.9	193.5	356.8
2000.....	23.5	*	*	*	0.5	1.7	4.8	13.5	38.5	92.9	194.3	353.0
1999.....	23.2	*	*	*	0.4	1.5	4.6	13.8	37.8	93.5	194.5	349.4

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Endocrine diseases (E00-E34)												
White, female												
2004.....	25.8	*	*	*	0.4	1.2	3.1	9.0	26.6	68.0	150.5	302.1
2003.....	26.8	*	*	*	0.3	1.3	3.2	9.4	28.1	71.1	155.5	317.3
2002.....	27.1	*	*	*	0.4	1.2	3.4	9.6	26.9	73.6	161.5	318.5
2001.....	27.2	*	*	*	0.2	1.2	3.1	9.6	28.2	73.0	162.8	326.1
2000.....	26.8	*	*	*	0.3	1.1	3.2	9.4	29.4	74.1	157.0	319.7
1999.....	26.5	1.4	*	*	0.4	1.2	3.2	8.8	28.4	75.3	155.3	316.4
Black, both sexes												
2004.....	34.2	*	*	0.3	1.1	3.6	10.0	29.0	82.1	183.1	341.0	555.3
2003.....	34.7	*	*	*	0.9	4.1	9.8	30.9	87.4	187.7	343.6	552.8
2002.....	34.5	*	*	*	1.1	3.9	10.1	31.1	86.0	191.3	343.0	559.8
2001.....	34.0	*	*	*	1.0	3.9	9.3	30.4	86.3	191.4	344.2	555.5
2000.....	33.8	*	*	*	1.0	3.9	8.7	30.8	83.6	192.2	360.0	550.6
1999.....	34.0	*	*	*	0.9	3.3	9.4	31.5	92.2	194.5	342.2	552.6
Black, male												
2004.....	31.0	*	*	*	1.1	4.3	12.2	34.0	93.1	202.3	348.3	525.5
2003.....	30.5	*	*	*	1.0	4.9	11.7	34.6	97.4	199.1	345.2	473.4
2002.....	29.5	*	*	*	1.2	4.7	12.1	35.7	96.3	193.2	333.9	427.8
2001.....	29.2	*	*	*	1.0	4.9	10.8	35.2	93.3	191.1	351.1	469.7
2000.....	28.1	*	*	*	1.0	4.3	9.8	33.9	89.3	188.3	343.4	474.0
1999.....	28.5	*	*	*	0.7	3.5	10.8	35.6	95.9	191.0	323.7	523.8
Black, female												
2004.....	37.2	*	*	*	1.1	3.0	8.0	24.7	73.2	169.4	336.9	567.0
2003.....	38.6	*	*	*	0.8	3.4	8.2	27.7	79.3	179.6	342.7	583.7
2002.....	39.0	*	*	*	0.9	3.1	8.4	27.1	77.7	189.9	348.2	610.4
2001.....	38.3	*	*	*	1.1	3.0	8.0	26.2	80.7	191.6	340.4	587.3
2000.....	39.1	*	*	*	1.0	3.6	7.7	28.1	78.9	194.9	369.4	579.1
1999.....	39.0	*	*	*	1.0	3.1	8.2	27.9	89.2	196.9	352.5	563.3

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Disorders of thyroid gland (E00-E07)												
All races 3/, both sexes												
2004.....	0.6	*	*	*	*	0.1	0.1	0.2	0.3	0.8	3.5	19.2
2003.....	0.7	*	*	*	*	0.1	0.1	0.2	0.4	0.9	4.2	21.1
2002.....	0.7	*	*	*	*	0.1	0.1	0.2	0.3	0.9	3.9	22.7
2001.....	0.7	*	*	*	*	*	0.1	0.2	0.3	0.8	4.2	23.2
2000.....	0.6	*	*	*	*	0.1	0.1	0.1	0.4	0.9	3.9	21.1
1999.....	0.6	*	*	*	*	0.1	0.1	0.2	0.3	1.0	4.0	22.4
All races 3/, male												
2004.....	0.3	*	*	*	*	*	*	0.1	0.3	0.6	2.3	11.2
2003.....	0.3	*	*	*	*	*	*	*	0.2	0.6	2.3	10.5
2002.....	0.3	*	*	*	*	*	*	0.1	0.2	0.6	2.1	12.7
2001.....	0.3	*	*	*	*	*	*	0.1	0.2	0.5	2.5	14.1
2000.....	0.3	*	*	*	*	*	*	*	0.3	0.6	1.9	12.1
1999.....	0.3	*	*	*	*	*	*	0.1	0.2	0.6	2.6	12.9
All races 3/, female												
2004.....	0.9	*	*	*	*	*	0.1	0.2	0.4	0.9	4.3	22.8
2003.....	1.1	*	*	*	*	*	0.1	0.3	0.5	1.2	5.5	25.8
2002.....	1.0	*	*	*	*	*	0.1	0.2	0.4	1.2	5.1	27.0
2001.....	1.0	*	*	*	*	*	0.1	0.2	0.4	1.1	5.3	27.1
2000.....	1.0	*	*	*	*	*	0.2	0.2	0.5	1.1	5.3	24.8
1999.....	1.0	*	*	*	*	*	0.1	0.2	0.4	1.2	4.9	26.3
White, both sexes												
2004.....	0.6	*	*	*	*	*	0.1	0.1	0.3	0.7	3.4	19.8
2003.....	0.7	*	*	*	*	*	0.1	0.1	0.4	0.9	4.3	21.7
2002.....	0.7	*	*	*	*	*	*	0.1	0.3	0.9	4.1	23.5
2001.....	0.7	*	*	*	*	*	0.1	0.1	0.3	0.8	4.0	23.9
2000.....	0.7	*	*	*	*	*	0.1	0.1	0.4	0.8	3.9	21.6
1999.....	0.7	*	*	*	*	*	0.1	0.1	0.3	0.9	3.9	23.0
White, male												
2004.....	0.3	*	*	*	*	*	*	*	0.2	0.6	2.2	11.6
2003.....	0.3	*	*	*	*	*	*	*	0.2	0.5	2.3	10.7
2002.....	0.3	*	*	*	*	*	*	*	*	0.6	2.2	13.1
2001.....	0.3	*	*	*	*	*	*	0.1	0.2	0.5	2.3	14.5
2000.....	0.3	*	*	*	*	*	*	*	0.2	0.5	2.0	12.0
1999.....	0.3	*	*	*	*	*	*	*	*	0.6	2.5	13.2

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Disorders of thyroid gland (E00-E07)												
White, female												
2004.....	1.0	*	*	*	*	*	*	0.2	0.3	0.8	4.3	23.5
2003.....	1.2	*	*	*	*	*	*	0.2	0.5	1.2	5.6	26.5
2002.....	1.2	*	*	*	*	*	*	0.2	0.4	1.2	5.3	28.0
2001.....	1.1	*	*	*	*	*	*	0.2	0.4	1.1	5.2	27.8
2000.....	1.1	*	*	*	*	*	*	0.2	0.5	1.0	5.1	25.5
1999.....	1.1	*	*	*	*	*	*	0.2	0.4	1.2	4.9	27.0
Black, both sexes												
2004.....	0.5	*	*	*	*	*	*	*	*	1.8	5.2	14.9
2003.....	0.5	*	*	*	*	*	*	0.4	*	1.4	4.1	19.3
2002.....	0.5	*	*	*	*	*	*	0.5	0.8	1.4	3.2	16.6
2001.....	0.5	*	*	*	*	*	*	*	*	1.4	5.8	19.3
2000.....	0.6	*	*	*	*	*	0.5	*	*	1.9	4.9	18.8
1999.....	0.6	*	*	*	*	*	*	0.6	0.9	1.5	5.2	18.8
Black, male												
2004.....	0.3	*	*	*	*	*	*	*	*	*	*	*
2003.....	0.3	*	*	*	*	*	*	*	*	*	*	*
2002.....	0.2	*	*	*	*	*	*	*	*	*	*	*
2001.....	0.3	*	*	*	*	*	*	*	*	*	*	*
2000.....	0.3	*	*	*	*	*	*	*	*	*	*	*
1999.....	0.3	*	*	*	*	*	*	*	*	*	*	*
Black, female												
2004.....	0.7	*	*	*	*	*	*	*	*	2.2	6.1	17.7
2003.....	0.7	*	*	*	*	*	*	*	*	*	4.5	22.7
2002.....	0.7	*	*	*	*	*	*	*	*	*	4.1	19.7
2001.....	0.7	*	*	*	*	*	*	*	*	*	6.1	21.9
2000.....	0.8	*	*	*	*	*	*	*	*	*	6.8	19.7
1999.....	0.8	*	*	*	*	*	*	*	*	*	5.7	21.9

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diabetes mellitus (E10-E14)												
All races 3/, both sexes												
2004.....	24.9	*	*	0.1	0.4	1.5	4.6	13.4	37.1	87.2	176.9	307.0
2003.....	25.5	*	*	0.1	0.4	1.6	4.6	13.9	38.5	90.8	181.1	317.5
2002.....	25.4	*	*	0.1	0.4	1.6	4.8	13.7	37.7	91.4	182.8	320.6
2001.....	25.1	*	*	0.1	0.4	1.5	4.3	13.6	37.8	91.4	181.4	321.8
2000.....	24.6	*	*	0.1	0.4	1.6	4.3	13.1	37.8	90.7	179.5	319.7
1999.....	24.5	*	*	0.1	0.4	1.4	4.3	12.9	38.3	91.8	178.0	317.2
All races 3/, male												
2004.....	24.4	*	*	*	0.4	1.7	5.7	16.5	44.0	100.0	202.5	333.9
2003.....	24.8	*	*	*	0.5	1.9	5.7	16.7	45.0	104.0	207.2	340.7
2002.....	24.2	*	*	0.1	0.5	1.9	5.9	16.4	44.5	101.4	202.6	348.5
2001.....	23.5	*	*	0.1	0.4	1.7	5.3	16.4	43.1	101.7	198.9	343.1
2000.....	22.9	*	*	*	0.5	1.9	5.1	15.3	42.4	99.1	199.6	345.1
1999.....	22.8	*	*	*	0.4	1.6	5.1	15.6	42.6	99.8	198.2	345.7
All races 3/, female												
2004.....	25.4	*	*	0.1	0.4	1.3	3.5	10.4	30.7	76.4	159.6	294.9
2003.....	26.2	*	*	*	0.3	1.4	3.5	11.2	32.4	79.8	163.6	307.2
2002.....	26.5	*	*	*	0.4	1.3	3.7	11.1	31.4	83.1	169.7	308.4
2001.....	26.6	*	*	*	0.3	1.3	3.4	11.0	32.9	82.8	169.9	312.9
2000.....	26.3	*	*	*	0.4	1.2	3.4	11.1	33.6	83.7	166.3	309.4
1999.....	26.2	*	*	*	0.4	1.3	3.5	10.4	34.3	85.2	164.9	305.8
White, both sexes												
2004.....	24.4	*	*	*	0.3	1.3	4.0	11.6	32.4	78.1	164.2	289.4
2003.....	25.0	*	*	0.1	0.3	1.4	4.0	11.8	33.5	81.4	168.4	300.8
2002.....	24.9	*	*	0.1	0.3	1.4	4.3	11.7	32.7	81.5	170.4	303.7
2001.....	24.6	*	*	0.1	0.3	1.2	3.8	11.7	32.7	81.9	168.9	306.0
2000.....	24.1	*	*	*	0.3	1.3	3.8	11.1	33.1	81.2	166.0	303.5
1999.....	23.9	*	*	0.1	0.4	1.3	3.7	10.9	32.4	81.8	165.1	298.9
White, male												
2004.....	24.3	*	*	*	0.3	1.4	5.0	14.7	39.2	91.6	193.3	323.0
2003.....	24.8	*	*	*	0.4	1.5	5.1	14.7	40.2	95.8	198.6	333.4
2002.....	24.2	*	*	*	0.4	1.6	5.3	14.3	39.6	93.3	194.7	344.3
2001.....	23.5	*	*	*	0.3	1.3	4.7	14.3	38.4	94.6	189.2	337.3
2000.....	22.9	*	*	*	0.4	1.6	4.7	13.3	38.0	91.8	190.5	337.4
1999.....	22.7	*	*	*	0.4	1.4	4.5	13.5	37.3	92.3	189.9	333.4

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diabetes mellitus (E10-E14)												
White, female												
2004.....	24.5	*	*	*	0.3	1.1	2.9	8.6	25.9	66.5	144.4	274.3
2003.....	25.2	*	*	*	0.3	1.2	2.9	9.0	27.2	69.2	148.0	286.3
2002.....	25.6	*	*	*	0.3	1.1	3.2	9.2	26.2	71.5	154.0	286.1
2001.....	25.7	*	*	*	0.2	1.1	2.9	9.2	27.4	71.2	155.5	293.0
2000.....	25.3	*	*	*	0.3	1.0	3.0	9.0	28.5	72.4	149.8	289.7
1999.....	25.0	*	*	*	0.3	1.2	3.0	8.4	27.8	73.1	148.9	285.1
Black, both sexes												
2004.....	33.2	*	*	*	0.9	3.3	9.5	28.1	80.5	179.3	332.8	532.4
2003.....	33.8	*	*	*	0.7	3.8	9.5	30.1	85.7	184.9	336.1	527.5
2002.....	33.6	*	*	*	1.0	3.6	9.6	30.1	84.2	188.1	336.9	537.8
2001.....	33.1	*	*	*	1.0	3.7	8.8	29.6	85.0	188.4	335.2	529.7
2000.....	32.8	*	*	*	0.9	3.5	8.0	30.2	82.5	188.4	350.6	524.9
1999.....	33.0	*	*	*	0.8	3.0	8.9	30.3	90.2	190.8	333.5	529.0
Black, male												
2004.....	30.2	*	*	*	1.0	3.9	11.8	33.1	91.6	198.4	340.8	505.7
2003.....	29.8	*	*	*	0.8	4.7	11.5	33.7	96.0	197.2	337.3	453.5
2002.....	29.0	*	*	*	1.2	4.4	11.7	34.8	94.7	190.1	329.9	417.0
2001.....	28.5	*	*	*	1.0	4.8	10.2	34.5	92.2	187.7	341.9	450.5
2000.....	27.4	*	*	*	1.0	3.8	9.4	33.4	88.2	184.9	337.0	452.1
1999.....	27.7	*	*	*	*	3.3	10.3	34.7	93.8	186.5	314.8	506.2
Black, female												
2004.....	36.0	*	*	*	0.9	2.7	7.6	23.8	71.6	165.7	328.2	543.0
2003.....	37.4	*	*	*	*	3.0	7.8	27.0	77.4	176.1	335.4	556.2
2002.....	37.8	*	*	*	0.7	2.9	7.8	26.0	75.8	186.7	340.9	584.1
2001.....	37.2	*	*	*	1.0	2.8	7.6	25.4	79.1	188.9	331.5	559.1
2000.....	37.8	*	*	*	0.9	3.1	6.8	27.5	77.8	190.9	358.3	552.0
1999.....	37.8	*	*	*	0.9	2.6	7.6	26.6	87.4	193.9	343.9	537.5

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other endocrine diseases (E15-E21,E28-E31,E34)

All races 3/, both sexes

2004.....	0.2	*	*	*	*	*	0.0	0.1	0.2	0.4	1.0	2.1
2003.....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.4	0.9	2.1
2002.....	0.2	0.7	*	*	*	*	*	0.1	0.2	0.4	1.0	2.2
2001.....	0.2	0.7	*	*	*	*	0.1	0.1	0.2	0.3	0.8	2.5
2000.....	0.1	0.7	*	*	*	*	0.1	0.1	0.1	0.3	0.9	2.2
1999.....	0.2	0.7	*	*	*	*	0.1	0.1	0.2	0.4	0.9	1.9

All races 3/, male

2004.....	0.1	*	*	*	*	*	*	0.1	0.2	0.4	1.1	1.8
2003.....	0.1	*	*	*	*	*	*	0.1	0.2	0.4	0.9	1.7
2002.....	0.1	*	*	*	*	*	*	0.1	0.2	0.4	0.9	1.8
2001.....	0.1	*	*	*	*	*	*	*	0.2	0.4	0.7	2.2
2000.....	0.1	*	*	*	*	*	*	0.1	*	*	0.9	1.7
1999.....	0.1	*	*	*	*	*	*	0.1	0.2	0.5	1.0	*

All races 3/, female

2004.....	0.2	*	*	*	*	*	*	0.1	0.2	0.4	0.9	2.2
2003.....	0.2	*	*	*	*	*	*	*	0.2	0.4	0.9	2.2
2002.....	0.2	*	*	*	*	*	*	0.1	0.2	0.4	1.0	2.3
2001.....	0.2	*	*	*	*	*	*	*	0.2	0.3	0.9	2.6
2000.....	0.2	*	*	*	*	*	*	*	0.2	0.3	0.9	2.4
1999.....	0.2	*	*	*	*	*	*	0.1	0.2	0.3	0.8	2.0

White, both sexes

2004.....	0.1	*	*	*	*	*	*	0.1	0.2	0.3	0.9	1.9
2003.....	0.1	*	*	*	*	*	0.1	0.1	0.2	0.3	0.8	2.0
2002.....	0.2	0.8	*	*	*	*	*	0.1	0.2	0.4	0.9	2.1
2001.....	0.1	0.7	*	*	*	*	*	0.1	0.2	0.2	0.7	2.4
2000.....	0.1	*	*	*	*	*	*	0.1	0.1	0.2	0.7	2.0
1999.....	0.1	*	*	*	*	*	*	0.1	0.2	0.3	0.8	1.8

White, male

2004.....	0.1	*	*	*	*	*	*	*	0.2	0.3	1.1	*
2003.....	0.1	*	*	*	*	*	*	0.1	*	0.3	0.7	*
2002.....	0.1	*	*	*	*	*	*	*	0.2	0.3	0.8	2.0
2001.....	0.1	*	*	*	*	*	*	*	*	*	0.7	2.1
2000.....	0.1	*	*	*	*	*	*	*	*	*	0.7	*
1999.....	0.1	*	*	*	*	*	*	*	*	0.3	1.0	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Nutritional diseases (E40-E68)												
All races 3/, both sexes												
2004.....	2.5	*	*	*	0.1	0.8	1.5	2.9	4.0	5.3	9.9	32.3
2003.....	2.5	*	*	*	0.2	0.7	1.5	2.8	3.9	5.4	10.2	37.5
2002.....	2.6	*	*	*	0.2	0.8	1.6	2.6	3.9	4.9	11.3	42.9
2001.....	2.4	*	*	*	0.2	0.6	1.3	2.4	3.3	4.6	10.8	44.5
2000.....	2.6	*	*	*	0.1	0.6	1.3	2.3	3.2	4.9	12.6	51.7
1999.....	2.5	*	*	*	0.2	0.5	1.2	2.0	2.9	4.7	12.3	55.3
All races 3/, male												
2004.....	2.1	*	*	*	0.2	0.8	1.6	3.0	3.9	4.9	9.5	28.8
2003.....	2.1	*	*	*	0.2	0.9	1.5	2.8	3.9	5.1	9.7	31.5
2002.....	2.2	*	*	*	0.2	0.9	1.8	2.6	3.7	4.5	10.3	37.8
2001.....	1.9	*	*	*	0.2	0.7	1.4	2.4	3.1	4.3	9.8	35.8
2000.....	2.0	*	*	*	0.2	0.7	1.5	2.4	3.2	4.6	11.2	44.6
1999.....	2.0	*	*	*	0.2	0.6	1.4	2.0	3.1	4.6	11.6	48.7
All races 3/, female												
2004.....	2.8	*	*	*	*	0.7	1.4	2.9	4.0	5.6	10.1	33.9
2003.....	2.9	*	*	*	0.2	0.5	1.5	2.8	3.9	5.6	10.6	40.1
2002.....	3.0	*	*	*	0.2	0.6	1.4	2.6	4.0	5.1	12.0	45.1
2001.....	2.9	*	*	*	0.2	0.5	1.2	2.5	3.5	4.9	11.4	48.1
2000.....	3.1	*	*	*	0.1	0.4	1.2	2.2	3.2	5.2	13.6	54.6
1999.....	2.9	*	*	*	0.2	0.4	1.0	2.0	2.6	4.7	12.8	58.0
White, both sexes												
2004.....	2.4	*	*	*	0.1	0.6	1.3	2.7	3.8	5.0	9.5	31.0
2003.....	2.6	*	*	*	0.2	0.6	1.4	2.7	3.8	5.3	9.8	36.1
2002.....	2.6	*	*	*	0.1	0.6	1.4	2.5	3.6	4.6	10.8	41.6
2001.....	2.4	*	*	*	0.1	0.5	1.1	2.2	3.2	4.3	10.1	43.8
2000.....	2.6	*	*	*	0.1	0.4	1.2	2.2	3.0	4.6	12.1	50.6
1999.....	2.5	*	*	*	0.1	0.4	1.1	1.9	2.6	4.2	11.5	54.2
White, male												
2004.....	2.1	*	*	*	0.1	0.6	1.5	2.9	3.9	4.7	9.1	27.8
2003.....	2.2	*	*	*	0.2	0.8	1.5	2.8	3.9	5.1	9.1	30.5
2002.....	2.2	*	*	*	0.2	0.7	1.7	2.7	3.6	4.3	9.8	37.1
2001.....	1.9	*	*	*	0.1	0.6	1.2	2.2	3.0	4.0	9.3	34.5
2000.....	2.1	*	*	*	0.1	0.6	1.4	2.5	3.0	4.3	10.7	43.7
1999.....	2.0	*	*	*	*	0.5	1.3	2.0	2.9	4.1	10.7	46.9

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Nutritional diseases (E40-E68)												
White, female												
2004.....	2.8	*	*	*	*	0.6	1.1	2.5	3.8	5.2	9.7	32.5
2003.....	3.0	*	*	*	0.1	0.4	1.3	2.6	3.7	5.6	10.3	38.6
2002.....	3.0	*	*	*	*	0.5	1.2	2.3	3.7	4.8	11.4	43.5
2001.....	2.9	*	*	*	*	0.4	1.0	2.2	3.3	4.6	10.6	47.7
2000.....	3.1	*	*	*	*	0.3	1.1	1.9	3.0	4.9	13.1	53.4
1999.....	3.0	*	*	*	*	0.3	0.9	1.7	2.4	4.3	12.1	57.2
Black, both sexes												
2004.....	3.3	*	*	*	0.3	1.9	3.2	5.1	6.7	9.8	15.8	50.8
2003.....	3.0	*	*	*	0.4	1.7	2.8	4.4	6.1	7.5	17.2	58.1
2002.....	3.3	*	*	*	0.5	1.9	2.9	4.4	7.3	8.8	19.7	62.1
2001.....	3.0	*	*	*	0.5	1.2	2.8	4.8	5.5	8.5	19.8	56.2
2000.....	3.1	*	*	*	0.3	1.5	2.5	4.3	5.8	8.8	20.6	68.5
1999.....	3.0	*	*	*	0.4	1.3	2.4	3.5	5.4	9.7	23.2	72.4
Black, male												
2004.....	2.6	*	*	*	*	2.0	3.0	4.0	5.4	8.5	16.4	45.7
2003.....	2.4	*	*	*	*	2.0	2.3	4.0	4.9	7.3	19.7	48.0
2002.....	2.6	*	*	*	*	2.3	3.0	3.2	6.3	7.7	18.9	48.2
2001.....	2.5	*	*	*	*	1.3	2.9	4.2	5.0	8.1	18.2	57.9
2000.....	2.5	*	*	*	*	1.7	2.5	3.3	6.1	8.9	20.6	65.7
1999.....	2.6	*	*	*	*	1.5	2.4	2.8	5.7	10.0	24.3	75.0
Black, female												
2004.....	3.9	*	*	*	*	1.8	3.4	6.1	7.7	10.7	15.4	52.8
2003.....	3.6	*	*	*	*	1.5	3.1	4.8	7.1	7.6	15.8	62.0
2002.....	4.0	*	*	*	*	1.6	2.7	5.4	8.1	9.5	20.1	67.4
2001.....	3.6	*	*	*	*	1.2	2.8	5.4	5.9	8.8	20.7	55.6
2000.....	3.6	*	*	*	*	1.3	2.6	5.1	5.6	8.8	20.6	69.5
1999.....	3.5	*	*	*	*	1.1	2.3	4.2	5.1	9.4	22.5	71.4

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malnutrition (E40-E46)												
All races 3/, both sexes												
2004.....	1.0	*	*	*	*	*	0.1	0.3	0.5	1.6	6.7	28.3
2003.....	1.1	*	*	*	*	*	0.1	0.2	0.6	1.7	7.4	33.4
2002.....	1.2	*	*	*	*	*	0.1	0.2	0.6	1.9	8.4	38.4
2001.....	1.2	*	*	*	*	*	0.1	0.3	0.6	1.9	8.2	40.2
2000.....	1.4	*	*	*	*	*	0.1	0.2	0.7	2.3	9.7	46.3
1999.....	1.4	*	*	*	*	*	0.1	0.3	0.6	2.2	9.5	50.0
All races 3/, male												
2004.....	0.7	*	*	*	*	*	*	0.2	0.6	1.7	6.6	25.3
2003.....	0.8	*	*	*	*	*	*	0.2	0.6	1.9	7.4	28.2
2002.....	0.9	*	*	*	*	*	0.1	0.2	0.7	2.1	8.0	35.1
2001.....	0.8	*	*	*	*	*	*	0.2	0.6	1.9	8.0	33.6
2000.....	0.9	*	*	*	*	*	*	0.2	0.7	2.5	8.9	40.7
1999.....	1.0	*	*	*	*	*	0.1	0.3	0.8	2.5	9.4	44.4
All races 3/, female												
2004.....	1.2	*	*	*	*	*	0.1	0.3	0.5	1.5	6.7	29.6
2003.....	1.4	*	*	*	*	*	*	0.2	0.6	1.7	7.4	35.6
2002.....	1.6	*	*	*	*	*	0.1	0.2	0.5	1.8	8.6	39.8
2001.....	1.6	*	*	*	*	*	0.1	0.3	0.5	1.9	8.2	43.0
2000.....	1.8	*	*	*	*	*	0.2	0.3	0.6	2.1	10.2	48.6
1999.....	1.8	*	*	*	*	*	0.1	0.3	0.5	2.0	9.5	52.3
White, both sexes												
2004.....	1.0	*	*	*	*	*	0.1	0.2	0.5	1.4	6.3	26.9
2003.....	1.1	*	*	*	*	*	0.1	0.2	0.5	1.6	7.0	32.1
2002.....	1.3	*	*	*	*	*	0.1	0.2	0.5	1.8	8.0	37.0
2001.....	1.3	*	*	*	*	*	0.1	0.2	0.5	1.8	7.6	39.5
2000.....	1.5	*	*	*	*	*	0.1	0.2	0.6	2.1	9.3	45.2
1999.....	1.5	*	*	*	*	*	0.1	0.2	0.5	1.9	8.8	48.9
White, male												
2004.....	0.7	*	*	*	*	*	*	0.2	0.5	1.5	6.1	24.1
2003.....	0.8	*	*	*	*	*	*	0.2	0.4	1.7	6.7	27.2
2002.....	0.9	*	*	*	*	*	0.1	0.2	0.5	1.9	7.5	34.3
2001.....	0.8	*	*	*	*	*	*	0.1	0.5	1.7	7.4	32.2
2000.....	1.0	*	*	*	*	*	*	0.2	0.6	2.2	8.4	39.7
1999.....	1.0	*	*	*	*	*	*	0.2	0.6	2.0	8.5	42.2

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malnutrition (E40-E46)												
White, female												
2004.....	1.3	*	*	*	*	*	*	0.3	0.5	1.4	6.5	28.2
2003.....	1.5	*	*	*	*	*	*	0.2	0.6	1.6	7.3	34.2
2002.....	1.6	*	*	*	*	*	*	0.3	0.5	1.7	8.4	38.2
2001.....	1.7	*	*	*	*	*	*	0.3	0.5	1.9	7.7	42.6
2000.....	1.9	*	*	*	*	*	0.1	0.2	0.6	2.0	9.9	47.4
1999.....	2.0	*	*	*	*	*	0.2	0.2	0.4	1.8	9.0	51.6
Black, both sexes												
2004.....	1.0	*	*	*	*	*	*	0.6	1.1	3.2	10.4	46.0
2003.....	1.2	*	*	*	*	*	*	0.7	1.4	3.0	13.0	53.4
2002.....	1.3	*	*	*	*	*	*	*	1.7	3.6	14.0	57.6
2001.....	1.2	*	*	*	*	*	*	0.6	1.1	3.4	14.8	51.6
2000.....	1.4	*	*	*	*	*	*	0.5	1.5	4.7	14.9	63.8
1999.....	1.5	*	*	*	*	*	*	0.6	1.8	5.4	17.9	66.0
Black, male												
2004.....	0.9	*	*	*	*	*	*	*	1.5	4.3	12.0	43.7
2003.....	1.0	*	*	*	*	*	*	*	1.9	4.2	17.2	45.9
2002.....	1.0	*	*	*	*	*	*	*	2.6	3.9	15.2	46.1
2001.....	1.1	*	*	*	*	*	*	1.0	*	4.9	15.9	55.6
2000.....	1.2	*	*	*	*	*	*	*	2.4	6.3	17.3	61.1
1999.....	1.4	*	*	*	*	*	*	*	2.5	7.8	20.3	73.8
Black, female												
2004.....	1.2	*	*	*	*	*	*	*	*	2.5	9.4	46.9
2003.....	1.3	*	*	*	*	*	*	*	*	2.2	10.7	56.3
2002.....	1.5	*	*	*	*	*	*	*	*	3.4	13.3	62.0
2001.....	1.3	*	*	*	*	*	*	*	*	2.2	14.2	50.1
2000.....	1.6	*	*	*	*	*	*	*	*	3.5	13.6	64.8
1999.....	1.6	*	*	*	*	*	*	*	*	3.6	16.5	63.1

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified malnutrition (E40,E42-E46)												
All races 3/, both sexes												
2004.....	0.9	*	*	*	*	*	0.1	0.2	0.5	1.5	6.4	27.0
2003.....	1.0	*	*	*	*	*	0.1	0.2	0.6	1.7	7.0	31.8
2002.....	1.2	*	*	*	*	*	0.1	0.2	0.6	1.9	8.0	36.0
2001.....	1.1	*	*	*	*	*	0.1	0.2	0.5	1.8	7.7	37.7
2000.....	1.3	*	*	*	*	*	0.1	0.2	0.6	2.2	9.2	43.6
1999.....	1.4	*	*	*	*	*	0.1	0.3	0.6	2.1	9.0	47.5
All races 3/, male												
2004.....	0.7	*	*	*	*	*	*	0.2	0.6	1.6	6.3	24.3
2003.....	0.7	*	*	*	*	*	*	0.2	0.5	1.8	7.0	27.1
2002.....	0.8	*	*	*	*	*	0.1	0.2	0.6	2.1	7.7	33.5
2001.....	0.8	*	*	*	*	*	*	0.2	0.6	1.8	7.5	31.3
2000.....	0.9	*	*	*	*	*	*	0.2	0.7	2.4	8.4	37.8
1999.....	0.9	*	*	*	*	*	0.1	0.3	0.8	2.3	8.9	41.4
All races 3/, female												
2004.....	1.2	*	*	*	*	*	0.1	0.3	0.5	1.4	6.4	28.2
2003.....	1.3	*	*	*	*	*	*	0.2	0.6	1.6	7.0	33.9
2002.....	1.5	*	*	*	*	*	0.1	0.2	0.5	1.7	8.2	37.1
2001.....	1.5	*	*	*	*	*	*	0.3	0.5	1.8	7.9	40.4
2000.....	1.7	*	*	*	*	*	0.2	0.2	0.5	2.1	9.7	45.9
1999.....	1.8	*	*	*	*	*	0.1	0.3	0.5	1.9	9.1	50.0
White, both sexes												
2004.....	1.0	*	*	*	*	*	0.1	0.2	0.5	1.4	6.0	25.7
2003.....	1.1	*	*	*	*	*	0.1	0.2	0.5	1.6	6.7	30.5
2002.....	1.2	*	*	*	*	*	0.1	0.2	0.5	1.7	7.7	34.7
2001.....	1.2	*	*	*	*	*	0.1	0.2	0.5	1.7	7.2	37.0
2000.....	1.4	*	*	*	*	*	0.1	0.2	0.5	2.0	8.8	42.5
1999.....	1.4	*	*	*	*	*	0.1	0.2	0.5	1.8	8.4	46.5
White, male												
2004.....	0.7	*	*	*	*	*	*	0.2	0.5	1.4	5.8	23.2
2003.....	0.7	*	*	*	*	*	*	0.2	0.4	1.6	6.4	26.0
2002.....	0.9	*	*	*	*	*	*	0.2	0.5	1.9	7.3	32.5
2001.....	0.8	*	*	*	*	*	*	0.1	0.5	1.6	6.9	30.1
2000.....	0.9	*	*	*	*	*	*	0.2	0.5	2.1	8.0	37.0
1999.....	0.9	*	*	*	*	*	*	0.2	0.6	1.8	8.1	39.4

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified malnutrition (E40,E42-E46)												
White, female												
2004.....	1.2	*	*	*	*	*	*	0.2	0.5	1.3	6.2	26.9
2003.....	1.4	*	*	*	*	*	*	0.2	0.6	1.5	6.9	32.4
2002.....	1.5	*	*	*	*	*	*	0.2	0.5	1.6	8.0	35.6
2001.....	1.6	*	*	*	*	*	*	0.2	0.5	1.8	7.4	39.9
2000.....	1.8	*	*	*	*	*	0.1	0.2	0.5	1.9	9.4	44.7
1999.....	1.9	*	*	*	*	*	0.1	0.2	0.4	1.7	8.6	49.4
Black, both sexes												
2004.....	1.0	*	*	*	*	*	*	0.6	1.0	3.1	10.4	44.8
2003.....	1.1	*	*	*	*	*	*	0.6	1.3	2.9	12.2	52.5
2002.....	1.2	*	*	*	*	*	*	*	1.6	3.5	13.1	55.8
2001.....	1.1	*	*	*	*	*	*	0.6	0.9	3.2	14.3	49.1
2000.....	1.3	*	*	*	*	*	*	0.5	1.5	4.5	14.6	59.7
1999.....	1.4	*	*	*	*	*	*	0.5	1.7	5.0	16.9	62.8
Black, male												
2004.....	0.9	*	*	*	*	*	*	*	*	4.2	12.0	42.7
2003.....	0.9	*	*	*	*	*	*	*	1.8	3.9	16.4	44.8
2002.....	0.9	*	*	*	*	*	*	*	2.5	3.9	14.1	45.0
2001.....	1.0	*	*	*	*	*	*	*	*	4.8	15.6	53.3
2000.....	1.1	*	*	*	*	*	*	*	2.3	6.0	16.4	55.4
1999.....	1.3	*	*	*	*	*	*	*	2.4	7.5	19.7	69.1
Black, female												
2004.....	1.1	*	*	*	*	*	*	*	*	2.3	9.4	45.7
2003.....	1.3	*	*	*	*	*	*	*	*	2.1	9.9	55.5
2002.....	1.4	*	*	*	*	*	*	*	*	3.2	12.5	60.0
2001.....	1.2	*	*	*	*	*	*	*	*	2.1	13.6	47.6
2000.....	1.5	*	*	*	*	*	*	*	*	3.4	13.6	61.4
1999.....	1.5	*	*	*	*	*	*	*	*	3.3	15.3	60.5

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Obesity and other hyperalimmentation (E65-E68)												
All races 3/, both sexes												
2004.....	1.4	*	*	*	0.1	0.7	1.4	2.6	3.4	3.7	3.0	2.0
2003.....	1.4	*	*	*	0.2	0.7	1.4	2.5	3.3	3.5	2.5	2.0
2002.....	1.3	*	*	*	0.2	0.7	1.5	2.3	3.2	2.8	2.4	1.4
2001.....	1.1	*	*	*	0.2	0.6	1.2	2.2	2.7	2.6	2.0	1.6
2000.....	1.1	*	*	*	0.1	0.5	1.2	2.1	2.5	2.4	2.2	1.7
1999.....	0.9	*	*	*	0.1	0.5	1.1	1.7	2.1	2.3	2.1	1.4
All races 3/, male												
2004.....	1.4	*	*	*	0.2	0.7	1.5	2.7	3.3	3.1	2.7	1.4
2003.....	1.3	*	*	*	0.2	0.9	1.4	2.6	3.3	3.1	1.9	*
2002.....	1.2	*	*	*	0.2	0.8	1.7	2.4	3.0	2.3	1.7	*
2001.....	1.0	*	*	*	0.1	0.6	1.3	2.1	2.4	2.2	1.3	*
2000.....	1.0	*	*	*	0.2	0.7	1.4	2.2	2.4	1.9	1.6	*
1999.....	0.9	*	*	*	0.1	0.6	1.3	1.7	2.2	1.9	1.4	*
All races 3/, female												
2004.....	1.5	*	*	*	*	0.7	1.3	2.6	3.5	4.1	3.2	2.2
2003.....	1.5	*	*	*	0.2	0.5	1.4	2.5	3.3	3.9	2.9	2.5
2002.....	1.3	*	*	*	0.1	0.6	1.2	2.3	3.5	3.2	2.9	1.6
2001.....	1.2	*	*	*	0.2	0.5	1.1	2.2	2.9	2.9	2.5	2.0
2000.....	1.1	*	*	*	*	0.4	1.0	1.9	2.6	2.8	2.7	2.0
1999.....	1.0	*	*	*	0.1	0.4	0.9	1.7	2.1	2.7	2.6	1.9
White, both sexes												
2004.....	1.4	*	*	*	0.1	0.6	1.2	2.5	3.3	3.5	2.8	1.9
2003.....	1.4	*	*	*	0.1	0.6	1.3	2.5	3.3	3.6	2.5	1.9
2002.....	1.2	*	*	*	0.1	0.6	1.3	2.2	3.1	2.7	2.2	1.4
2001.....	1.0	*	*	*	0.1	0.5	1.0	2.0	2.6	2.4	1.9	1.6
2000.....	1.0	*	*	*	0.1	0.4	1.1	1.9	2.4	2.3	2.1	1.7
1999.....	0.9	*	*	*	0.1	0.4	1.0	1.6	2.1	2.2	2.0	1.4
White, male												
2004.....	1.4	*	*	*	0.1	0.6	1.4	2.8	3.3	3.2	2.7	1.5
2003.....	1.4	*	*	*	0.2	0.8	1.4	2.6	3.5	3.3	2.0	*
2002.....	1.2	*	*	*	0.1	0.6	1.6	2.5	3.0	2.3	1.7	*
2001.....	1.0	*	*	*	*	0.6	1.1	2.1	2.5	2.1	1.3	*
2000.....	1.0	*	*	*	0.1	0.6	1.3	2.2	2.4	2.0	1.5	*
1999.....	0.9	*	*	*	*	0.5	1.2	1.7	2.2	2.0	1.4	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Obesity and other hyperalimentation (E65-E68)												
White, female												
2004.....	1.4	*	*	*	*	0.6	1.0	2.2	3.3	3.8	3.0	2.0
2003.....	1.4	*	*	*	*	0.4	1.2	2.4	3.1	3.9	2.8	2.3
2002.....	1.2	*	*	*	*	0.5	1.1	2.0	3.1	3.0	2.5	1.7
2001.....	1.1	*	*	*	*	0.4	0.9	1.9	2.8	2.6	2.3	2.0
2000.....	1.0	*	*	*	*	0.3	0.9	1.6	2.4	2.6	2.5	2.1
1999.....	0.9	*	*	*	*	0.3	0.7	1.5	1.9	2.4	2.4	1.8
Black, both sexes												
2004.....	2.2	*	*	*	0.3	1.8	3.0	4.5	5.5	6.4	5.2	*
2003.....	1.9	*	*	*	0.3	1.7	2.7	3.8	4.7	4.3	4.0	*
2002.....	2.0	*	*	*	0.5	1.9	2.6	4.0	5.6	4.6	5.2	*
2001.....	1.8	*	*	*	0.5	1.2	2.6	4.1	4.3	4.9	3.9	*
2000.....	1.6	*	*	*	*	1.3	2.3	3.7	4.1	4.1	4.5	*
1999.....	1.4	*	*	*	0.3	1.2	2.1	2.9	3.4	4.2	3.8	*
Black, male												
2004.....	1.7	*	*	*	*	2.0	2.7	3.3	3.8	4.0	*	*
2003.....	1.4	*	*	*	*	2.0	2.3	3.2	2.9	3.0	*	*
2002.....	1.5	*	*	*	*	2.2	2.8	2.5	3.5	3.3	*	*
2001.....	1.3	*	*	*	*	1.2	2.5	3.0	3.1	*	*	*
2000.....	1.3	*	*	*	*	1.5	2.3	2.7	3.3	*	*	*
1999.....	1.1	*	*	*	*	1.5	2.1	2.2	2.9	*	*	*
Black, female												
2004.....	2.7	*	*	*	*	1.7	3.3	5.5	6.8	8.1	5.8	*
2003.....	2.2	*	*	*	*	1.4	3.1	4.2	6.1	5.3	5.1	*
2002.....	2.4	*	*	*	*	1.6	2.4	5.2	7.2	5.5	6.7	*
2001.....	2.2	*	*	*	*	1.1	2.6	5.0	5.3	6.4	5.1	*
2000.....	1.9	*	*	*	*	1.2	2.2	4.6	4.8	5.3	5.5	*
1999.....	1.7	*	*	*	*	1.0	2.1	3.5	3.8	5.8	5.0	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Metabolic disorders (E70-E88)												
All races 3/, both sexes												
2004.....	5.4	4.9	0.8	0.4	0.6	0.6	0.8	2.1	5.3	13.3	34.6	107.9
2003.....	5.5	5.2	1.1	0.4	0.7	0.6	0.9	2.2	5.2	13.1	34.6	112.6
2002.....	5.5	5.5	0.8	0.4	0.6	0.6	0.8	2.0	5.6	13.0	35.2	114.3
2001.....	5.5	6.0	1.0	0.4	0.6	0.6	0.8	2.1	5.0	13.1	34.1	124.4
2000.....	5.4	5.3	0.9	0.4	0.7	0.6	0.8	2.0	5.1	12.6	33.9	124.1
1999.....	5.3	5.6	0.8	0.4	0.7	0.6	0.8	1.8	4.7	12.2	33.1	130.8
All races 3/, male												
2004.....	5.1	6.0	0.8	0.4	0.6	0.6	0.9	3.0	7.0	16.9	38.5	103.1
2003.....	5.1	5.8	1.1	0.5	0.7	0.6	1.1	3.0	6.9	16.8	38.4	104.8
2002.....	4.9	6.1	0.8	0.5	0.6	0.7	1.0	2.8	7.3	16.0	38.8	101.5
2001.....	4.9	6.5	1.1	0.4	0.6	0.6	1.0	2.8	6.8	16.7	38.6	109.8
2000.....	4.8	5.5	0.9	0.5	0.7	0.7	0.9	2.9	6.8	15.8	36.7	112.0
1999.....	4.5	6.3	0.9	0.4	0.7	0.6	0.9	2.4	6.1	15.1	35.8	110.5
All races 3/, female												
2004.....	5.8	3.8	0.7	0.4	0.6	0.5	0.7	1.3	3.7	10.3	31.9	110.1
2003.....	5.9	4.5	1.0	0.3	0.6	0.5	0.7	1.5	3.6	10.0	32.1	116.1
2002.....	6.0	4.9	0.9	0.4	0.6	0.5	0.7	1.3	4.0	10.5	32.8	119.9
2001.....	6.1	5.5	1.0	0.4	0.7	0.5	0.6	1.5	3.4	10.2	31.1	130.6
2000.....	6.0	5.0	0.8	0.3	0.7	0.5	0.7	1.1	3.5	10.0	32.0	129.1
1999.....	6.1	4.9	0.7	0.4	0.8	0.6	0.6	1.2	3.5	9.8	31.3	139.0
White, both sexes												
2004.....	5.9	5.0	0.8	0.4	0.7	0.6	0.8	2.1	5.2	13.2	34.6	109.2
2003.....	5.9	4.6	1.1	0.4	0.7	0.6	0.9	2.1	5.1	12.9	34.0	112.9
2002.....	5.9	5.0	0.9	0.5	0.7	0.7	0.8	2.0	5.4	12.6	34.6	114.0
2001.....	5.9	5.6	1.1	0.4	0.7	0.6	0.9	2.1	4.9	13.0	33.6	123.4
2000.....	5.8	5.0	0.9	0.5	0.8	0.6	0.8	1.9	5.0	12.5	33.6	123.6
1999.....	5.7	4.9	0.9	0.4	0.8	0.6	0.7	1.7	4.6	11.9	32.7	131.3
White, male												
2004.....	5.5	6.2	0.9	0.4	0.7	0.7	0.9	2.9	7.0	16.9	38.6	104.6
2003.....	5.4	5.6	1.1	0.6	0.7	0.7	1.1	2.9	6.9	16.5	37.5	106.6
2002.....	5.2	6.2	0.8	0.6	0.7	0.8	1.0	2.7	7.1	15.7	38.1	100.2
2001.....	5.2	5.9	1.1	0.4	0.7	0.6	1.0	2.8	6.6	16.5	37.9	109.2
2000.....	5.1	5.2	1.0	0.5	0.7	0.7	0.9	2.8	6.7	15.7	36.3	113.9
1999.....	4.7	5.6	1.0	0.5	0.7	0.6	0.8	2.3	6.0	14.7	35.0	108.4

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Metabolic disorders (E70-E88)												
White, female												
2004.....	6.3	3.7	0.8	0.4	0.8	0.6	0.7	1.2	3.5	10.1	31.9	111.2
2003.....	6.3	3.7	1.0	0.3	0.7	0.6	0.7	1.4	3.4	9.8	31.7	115.6
2002.....	6.5	3.9	1.0	0.4	0.8	0.6	0.6	1.3	3.9	9.9	32.2	120.0
2001.....	6.6	5.3	1.1	0.5	0.8	0.6	0.7	1.4	3.4	10.1	30.7	129.4
2000.....	6.4	4.8	0.8	0.4	0.8	0.6	0.7	1.0	3.4	9.8	31.7	127.6
1999.....	6.7	4.1	0.8	0.4	0.9	0.7	0.6	1.1	3.3	9.6	31.2	140.4
Black, both sexes												
2004.....	4.2	5.1	*	*	*	0.4	1.2	2.8	7.3	17.9	40.8	113.1
2003.....	4.6	7.9	0.8	*	0.4	0.5	1.1	3.3	7.0	17.8	45.4	126.0
2002.....	4.8	7.6	0.8	0.3	0.4	0.4	1.1	3.1	8.4	19.3	47.3	135.1
2001.....	4.6	8.7	*	*	*	0.4	0.7	3.2	7.0	17.3	46.2	151.4
2000.....	4.4	6.2	*	*	0.4	0.6	1.1	2.8	7.2	16.4	44.0	147.0
1999.....	4.4	9.3	*	0.4	0.3	0.4	1.3	2.9	6.5	17.6	40.8	138.7
Black, male												
2004.....	4.0	6.2	*	*	*	*	1.6	3.9	9.0	22.0	46.8	112.3
2003.....	4.2	6.8	*	*	*	*	1.3	4.2	8.9	23.1	55.0	98.0
2002.....	4.4	6.7	*	*	0.6	*	1.2	4.1	11.9	21.8	52.5	134.0
2001.....	4.2	9.0	*	*	*	*	0.8	4.0	9.8	23.0	55.9	130.5
2000.....	4.0	7.4	*	*	*	0.8	1.3	3.9	9.4	20.0	50.0	115.3
1999.....	4.1	10.4	*	*	*	*	1.5	3.5	7.9	21.9	48.3	147.6
Black, female												
2004.....	4.4	*	*	*	*	*	0.9	1.8	6.0	14.9	37.4	113.4
2003.....	4.9	9.0	*	*	*	*	1.0	2.4	5.6	14.0	40.0	136.9
2002.....	5.1	8.5	*	*	*	*	0.9	2.2	5.5	17.4	44.3	135.5
2001.....	4.9	8.5	*	*	*	*	*	2.5	4.7	13.2	40.8	159.1
2000.....	4.9	*	*	*	*	*	0.9	1.8	5.5	13.8	40.6	158.8
1999.....	4.6	8.1	*	*	*	*	1.2	2.4	5.3	14.6	36.7	135.4

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Cystic fibrosis (E84)

All races 3/, both sexes

2004.....	0.2	*	*	0.1	0.4	0.3	0.2	0.1	*	*	*	*
2003.....	0.2	*	*	0.1	0.4	0.3	0.2	0.1	*	*	*	*
2002.....	0.2	*	*	0.1	0.4	0.3	0.2	0.1	*	*	*	*
2001.....	0.2	*	*	0.1	0.4	0.3	0.2	0.1	*	*	*	*
2000.....	0.2	*	*	0.1	0.4	0.3	0.2	0.1	*	*	*	*
1999.....	0.2	*	*	0.1	0.5	0.3	0.1	0.1	*	*	*	*

All races 3/, male

2004.....	0.2	*	*	*	0.4	0.3	0.1	*	*	*	*	*
2003.....	0.2	*	*	*	0.3	0.4	0.2	*	*	*	*	*
2002.....	0.2	*	*	*	0.3	0.4	0.2	*	*	*	*	*
2001.....	0.2	*	*	*	0.4	0.3	0.2	0.1	*	*	*	*
2000.....	0.2	*	*	0.1	0.4	0.3	0.2	*	*	*	*	*
1999.....	0.2	*	*	0.1	0.4	0.4	0.2	*	*	*	*	*

All races 3/, female

2004.....	0.2	*	*	*	0.4	0.3	0.2	*	*	*	*	*
2003.....	0.2	*	*	*	0.4	0.3	0.2	*	*	*	*	*
2002.....	0.2	*	*	0.1	0.4	0.3	0.2	*	*	*	*	*
2001.....	0.2	*	*	0.1	0.5	0.3	0.1	0.1	*	*	*	*
2000.....	0.2	*	*	0.1	0.5	0.4	0.2	*	*	*	*	*
1999.....	0.2	*	*	0.1	0.5	0.3	0.1	*	*	*	*	*

White, both sexes

2004.....	0.2	*	*	0.1	0.5	0.4	0.2	0.1	*	*	*	*
2003.....	0.2	*	*	0.1	0.5	0.4	0.2	0.1	*	*	*	*
2002.....	0.2	*	*	0.1	0.5	0.4	0.2	0.1	*	*	*	*
2001.....	0.2	*	*	0.1	0.5	0.4	0.2	0.1	*	*	*	*
2000.....	0.2	*	*	0.1	0.5	0.4	0.3	0.1	*	*	*	*
1999.....	0.2	*	*	0.1	0.6	0.4	0.2	0.1	*	*	*	*

White, male

2004.....	0.2	*	*	*	0.5	0.4	0.2	*	*	*	*	*
2003.....	0.2	*	*	*	0.4	0.4	0.2	*	*	*	*	*
2002.....	0.2	*	*	*	0.4	0.5	0.2	*	*	*	*	*
2001.....	0.2	*	*	*	0.4	0.4	0.3	0.1	*	*	*	*
2000.....	0.2	*	*	0.1	0.5	0.4	0.2	*	*	*	*	*
1999.....	0.2	*	*	0.1	0.5	0.4	0.2	*	*	*	*	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Volume depletion, disorders of fluid, electrolyte and acid-base balance (E86-E87)

All races 3/, both sexes

2004.....	2.2	1.6	*	*	0.0	0.1	0.2	0.5	1.0	3.2	13.7	70.7
2003.....	2.4	1.9	0.2	*	0.1	0.1	0.2	0.6	1.1	3.4	15.1	77.3
2002.....	2.5	1.8	*	*	0.1	0.1	0.2	0.5	1.2	3.8	16.0	83.2
2001.....	2.7	2.2	0.2	*	*	0.1	0.2	0.5	1.2	4.1	16.8	93.9
2000.....	2.8	1.9	*	*	0.1	0.1	0.3	0.5	1.3	4.0	17.4	96.8
1999.....	2.9	1.9	0.1	*	0.1	0.1	0.3	0.5	1.1	4.2	18.2	106.4

All races 3/, male

2004.....	1.6	1.9	*	*	*	0.1	0.2	0.6	1.1	3.5	13.3	59.8
2003.....	1.7	2.2	*	*	*	0.1	0.2	0.6	1.2	3.4	15.0	65.6
2002.....	1.7	2.1	*	*	*	*	0.2	0.6	1.3	3.8	15.3	67.9
2001.....	1.8	2.2	*	*	*	*	0.2	0.6	1.3	4.1	16.8	76.5
2000.....	2.0	1.8	*	*	*	*	0.3	0.6	1.5	4.5	17.7	84.3
1999.....	2.0	2.3	*	*	*	*	0.3	0.7	1.2	4.5	18.4	85.0

All races 3/, female

2004.....	2.8	1.3	*	*	*	*	0.2	0.4	0.9	2.9	13.9	75.7
2003.....	3.1	1.7	*	*	*	*	0.2	0.5	1.0	3.3	15.2	82.5
2002.....	3.3	1.5	*	*	*	0.1	0.2	0.5	1.2	3.7	16.5	89.8
2001.....	3.6	2.2	*	*	*	0.1	0.2	0.5	1.1	4.1	16.8	101.2
2000.....	3.5	2.0	*	*	*	*	0.2	0.4	1.2	3.6	17.2	101.9
1999.....	3.8	1.5	*	*	*	*	0.2	0.4	1.0	3.9	18.1	115.1

White, both sexes

2004.....	2.4	1.4	*	*	*	0.1	0.1	0.5	0.9	2.9	13.4	70.8
2003.....	2.5	1.4	*	*	*	0.1	0.2	0.5	1.0	3.1	14.5	76.6
2002.....	2.7	1.4	*	*	*	0.1	0.2	0.4	1.1	3.4	15.2	82.4
2001.....	2.9	1.7	0.2	*	*	0.1	0.2	0.4	1.0	3.7	16.0	93.0
2000.....	2.9	1.5	*	*	*	0.1	0.2	0.4	1.2	3.7	17.0	96.0
1999.....	3.1	1.3	*	*	*	0.1	0.2	0.4	0.9	3.8	17.8	106.6

White, male

2004.....	1.6	1.5	*	*	*	*	0.2	0.6	0.9	3.3	13.0	59.0
2003.....	1.7	1.6	*	*	*	*	0.2	0.5	1.1	3.1	14.0	65.6
2002.....	1.7	1.7	*	*	*	*	0.2	0.4	1.1	3.4	14.4	66.1
2001.....	1.9	1.6	*	*	*	*	0.2	0.4	1.1	3.6	15.7	75.8
2000.....	2.0	1.4	*	*	*	*	0.2	0.5	1.3	4.1	16.9	84.9
1999.....	2.0	1.6	*	*	*	*	0.2	0.6	1.0	3.8	17.4	82.4

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Volume depletion, disorders of fluid, electrolyte and acid-base balance (E86-E87)

White, female

2004.....	3.1	*	*	*	*	*	0.1	0.4	0.9	2.6	13.7	76.1
2003.....	3.3	*	*	*	*	*	0.2	0.5	0.9	3.1	14.8	81.5
2002.....	3.6	*	*	*	*	*	0.2	0.4	1.1	3.4	15.8	89.5
2001.....	3.8	1.8	*	*	*	*	0.2	0.4	1.0	3.8	16.2	100.2
2000.....	3.8	1.6	*	*	*	*	0.2	0.3	1.1	3.3	17.0	100.5
1999.....	4.2	*	*	*	*	*	0.2	0.3	0.9	3.7	17.9	116.4

Black, both sexes

2004.....	2.1	*	*	*	*	*	0.6	0.9	2.7	6.6	19.6	84.3
2003.....	2.4	4.5	*	*	*	*	0.5	1.4	2.4	7.0	26.2	97.1
2002.....	2.6	3.6	*	*	*	*	0.6	1.3	2.5	8.1	28.4	104.2
2001.....	2.8	5.1	*	*	*	*	0.4	1.5	2.9	8.4	29.3	119.4
2000.....	2.7	3.3	*	*	*	*	0.6	1.4	2.7	8.2	27.2	119.8
1999.....	2.7	4.8	*	*	*	*	0.8	1.4	2.9	9.2	26.3	115.1

Black, male

2004.....	1.8	*	*	*	*	*	*	1.1	3.8	7.5	22.3	86.4
2003.....	2.0	*	*	*	*	*	*	1.8	2.4	8.3	32.7	77.2
2002.....	2.1	*	*	*	*	*	*	1.7	3.3	8.8	30.4	99.7
2001.....	2.2	*	*	*	*	*	*	1.7	3.8	10.1	35.3	97.6
2000.....	2.2	*	*	*	*	*	0.8	1.8	3.7	9.8	33.6	98.0
1999.....	2.5	*	*	*	*	*	1.2	1.6	3.5	12.9	32.9	125.4

Black, female

2004.....	2.3	*	*	*	*	*	*	*	1.7	6.0	18.1	83.5
2003.....	2.8	*	*	*	*	*	*	1.0	2.4	6.1	22.5	104.9
2002.....	2.9	*	*	*	*	*	*	0.9	1.8	7.5	27.3	106.0
2001.....	3.2	*	*	*	*	*	*	1.3	2.1	7.3	25.9	127.6
2000.....	3.1	*	*	*	*	*	*	1.1	1.9	7.1	23.5	127.9
1999.....	2.9	*	*	*	*	*	*	1.2	2.4	6.6	22.5	111.3

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other metabolic disorders (E70-E83,E85,E88)												
All races 3/, both sexes												
2004.....	3.0	3.1	0.7	0.3	0.2	0.2	0.5	1.5	4.2	10.1	20.9	37.1
2003.....	2.9	3.1	0.8	0.3	0.2	0.2	0.5	1.6	4.1	9.7	19.4	35.2
2002.....	2.8	3.4	0.7	0.3	0.2	0.2	0.5	1.5	4.3	9.2	19.1	31.1
2001.....	2.6	3.7	0.8	0.3	0.2	0.1	0.4	1.5	3.8	9.0	17.2	30.5
2000.....	2.4	3.2	0.8	0.3	0.2	0.2	0.3	1.4	3.8	8.6	16.4	27.3
1999.....	2.2	3.4	0.7	0.3	0.2	0.1	0.4	1.2	3.6	8.0	14.8	24.3
All races 3/, male												
2004.....	3.3	3.8	0.7	0.3	0.2	0.2	0.6	2.2	5.8	13.4	25.2	43.2
2003.....	3.2	3.5	1.0	0.4	0.3	0.2	0.7	2.3	5.7	13.3	23.3	39.2
2002.....	3.0	3.6	0.7	0.4	0.3	0.2	0.6	2.2	6.0	12.1	23.4	33.6
2001.....	2.9	4.1	0.9	0.3	0.2	0.2	0.5	2.2	5.5	12.6	21.7	33.3
2000.....	2.6	3.6	0.8	0.3	0.2	0.3	0.4	2.2	5.3	11.3	19.0	27.7
1999.....	2.4	3.7	0.8	0.3	0.2	0.1	0.5	1.7	4.8	10.5	17.3	25.5
All races 3/, female												
2004.....	2.8	2.4	0.6	0.2	0.2	0.1	0.3	0.8	2.7	7.3	17.9	34.4
2003.....	2.6	2.8	0.7	0.2	0.2	0.2	0.3	0.9	2.5	6.7	16.8	33.4
2002.....	2.5	3.2	0.7	0.2	0.1	0.1	0.3	0.8	2.8	6.7	16.3	30.0
2001.....	2.3	3.2	0.7	0.3	0.1	0.1	0.3	0.9	2.3	6.1	14.3	29.3
2000.....	2.3	2.8	0.7	0.2	0.2	*	0.2	0.6	2.3	6.3	14.8	27.1
1999.....	2.1	3.1	0.6	0.3	0.2	0.1	0.3	0.8	2.4	5.9	13.2	23.9
White, both sexes												
2004.....	3.3	3.4	0.7	0.3	0.2	0.2	0.5	1.5	4.3	10.2	21.2	38.4
2003.....	3.2	3.1	0.9	0.3	0.2	0.2	0.5	1.6	4.1	9.8	19.5	36.1
2002.....	3.0	3.3	0.7	0.3	0.2	0.2	0.5	1.5	4.3	9.1	19.3	31.6
2001.....	2.8	3.7	0.8	0.3	0.2	0.1	0.4	1.5	3.9	9.3	17.5	30.4
2000.....	2.7	3.4	0.8	0.3	0.2	0.2	0.4	1.4	3.8	8.8	16.6	27.5
1999.....	2.4	3.1	0.8	0.3	0.2	0.1	0.4	1.2	3.7	8.1	14.9	24.6
White, male												
2004.....	3.6	4.2	0.8	0.3	0.2	0.2	0.6	2.2	6.1	13.6	25.6	45.5
2003.....	3.5	3.8	1.0	0.5	0.3	0.2	0.7	2.4	5.8	13.4	23.5	40.9
2002.....	3.3	4.0	0.7	0.4	0.3	0.2	0.7	2.2	5.9	12.2	23.6	34.1
2001.....	3.1	4.0	0.9	0.3	0.2	0.2	0.6	2.2	5.5	12.8	22.1	33.3
2000.....	2.9	3.8	0.9	0.4	0.2	0.2	0.5	2.3	5.4	11.6	19.4	29.0
1999.....	2.6	3.5	0.9	0.3	0.2	0.1	0.5	1.7	5.0	10.8	17.5	25.9

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other metabolic disorders (E70-E83,E85,E88)												
White, female												
2004.....	3.0	2.5	0.7	0.3	0.2	0.2	0.3	0.8	2.6	7.4	18.1	35.1
2003.....	2.9	2.4	0.8	0.2	0.2	0.2	0.3	0.9	2.5	6.8	16.8	34.0
2002.....	2.7	2.6	0.8	0.2	0.1	0.1	0.3	0.8	2.7	6.5	16.4	30.4
2001.....	2.5	3.3	0.8	0.3	0.2	*	0.3	0.9	2.4	6.3	14.4	29.1
2000.....	2.4	3.0	0.7	0.2	0.2	*	0.2	0.7	2.2	6.4	14.7	26.9
1999.....	2.3	2.7	0.7	0.3	0.2	0.2	0.2	0.7	2.4	5.9	13.3	24.0
Black, both sexes												
2004.....	2.1	*	*	*	*	*	0.6	1.8	4.7	11.3	21.2	28.8
2003.....	2.0	3.3	*	*	*	*	0.6	1.8	4.6	10.8	19.2	28.9
2002.....	2.2	3.9	*	*	*	*	0.5	1.8	5.8	11.1	18.9	30.9
2001.....	1.8	3.7	*	*	*	*	*	1.7	4.1	8.8	16.9	31.9
2000.....	1.7	*	*	*	*	*	0.4	1.3	4.5	8.2	16.8	27.2
1999.....	1.7	4.5	*	0.3	*	*	0.5	1.5	3.6	8.4	14.6	23.6
Black, male												
2004.....	2.2	*	*	*	*	*	0.8	2.7	5.2	14.5	24.5	25.8
2003.....	2.1	*	*	*	*	*	0.8	2.3	6.4	14.8	22.3	20.9
2002.....	2.3	*	*	*	*	*	*	2.4	8.5	12.8	22.1	34.3
2001.....	1.9	*	*	*	*	*	*	2.3	6.0	12.9	20.6	32.9
2000.....	1.7	*	*	*	*	*	*	2.0	5.6	10.2	16.4	*
1999.....	1.5	*	*	*	*	*	*	1.9	4.4	9.0	15.4	*
Black, female												
2004.....	2.1	*	*	*	*	*	*	1.1	4.3	8.9	19.4	29.9
2003.....	2.0	*	*	*	*	*	*	1.4	3.1	7.9	17.4	32.0
2002.....	2.1	*	*	*	*	*	*	1.3	3.6	9.9	17.1	29.6
2001.....	1.7	*	*	*	*	*	*	1.2	2.6	5.9	14.9	31.6
2000.....	1.7	*	*	*	*	*	*	*	3.6	6.7	17.0	30.9
1999.....	1.8	*	*	*	*	*	0.7	1.2	2.9	8.0	14.1	24.1

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

V. Mental and behavioral disorders (F01-F99)

All races 3/, both sexes

2004.....	21.0	*	*	0.1	0.7	1.9	5.0	9.0	9.8	20.6	117.1	671.4
2003.....	20.7	*	*	0.1	0.7	1.8	5.2	8.7	9.2	20.0	115.9	679.3
2002.....	19.3	*	*	0.1	0.6	1.7	5.3	8.1	8.8	19.0	107.6	646.7
2001.....	17.8	*	*	0.1	0.6	1.8	4.9	7.8	8.4	18.2	99.0	604.8
2000.....	16.4	*	*	0.1	0.6	1.8	4.9	7.3	8.1	16.9	89.9	561.6
1999.....	15.1	*	*	0.1	0.6	1.8	5.1	7.1	8.0	16.0	81.3	515.3

All races 3/, male

2004.....	16.2	*	*	*	0.9	2.7	7.2	13.6	15.1	25.6	114.3	529.7
2003.....	15.7	*	*	*	0.9	2.6	7.5	13.3	13.7	24.7	114.0	523.8
2002.....	14.8	*	*	*	0.8	2.3	7.5	12.7	13.5	23.9	106.9	506.0
2001.....	13.8	*	*	*	0.8	2.6	7.0	12.2	12.8	23.3	97.3	488.0
2000.....	12.8	*	*	*	0.9	2.6	7.2	11.5	12.4	21.9	89.9	451.0
1999.....	12.3	*	*	0.1	0.8	2.6	7.7	11.3	12.6	21.2	85.5	415.8

All races 3/, female

2004.....	25.6	*	*	*	0.5	1.0	2.7	4.6	4.9	16.4	119.0	735.1
2003.....	25.5	*	*	*	0.4	1.0	2.9	4.2	5.0	16.1	117.1	748.1
2002.....	23.7	*	*	*	0.4	1.0	3.0	3.7	4.5	14.9	108.0	707.8
2001.....	21.7	*	*	*	0.3	1.0	2.8	3.6	4.3	13.9	100.1	653.7
2000.....	19.8	*	*	*	0.4	1.0	2.5	3.3	4.2	12.9	90.0	606.6
1999.....	17.8	*	*	*	0.3	1.0	2.5	3.1	3.8	11.8	78.6	555.4

White, both sexes

2004.....	23.1	*	*	0.1	0.8	1.9	5.0	8.7	9.4	20.0	118.4	687.9
2003.....	22.8	*	*	0.1	0.7	1.8	5.0	8.2	8.7	19.9	116.8	697.4
2002.....	21.3	*	*	0.1	0.6	1.7	5.1	7.6	8.2	18.7	108.8	663.8
2001.....	19.6	*	*	0.1	0.6	1.8	4.6	7.0	7.8	17.8	100.8	621.6
2000.....	17.9	*	*	0.1	0.7	1.8	4.6	6.6	7.5	16.5	90.7	577.6
1999.....	16.5	*	*	0.1	0.6	1.9	4.6	6.3	7.3	15.5	82.3	534.6

White, male

2004.....	17.2	*	*	*	1.1	2.6	7.3	13.0	14.1	24.2	113.6	538.9
2003.....	16.7	*	*	*	1.0	2.7	7.3	12.4	12.6	24.1	113.4	535.8
2002.....	15.7	*	*	*	0.9	2.3	7.4	11.8	12.2	22.6	106.7	514.9
2001.....	14.5	*	*	*	0.9	2.6	6.6	10.9	11.5	22.3	98.0	498.4
2000.....	13.4	*	*	*	0.9	2.6	6.8	10.3	11.2	20.6	89.7	459.4
1999.....	12.8	*	*	*	0.8	2.8	7.1	10.0	11.3	20.1	84.9	429.2

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
V. Mental and behavioral disorders (F01-F99)												
White, female												
2004.....	28.8	*	*	*	0.5	1.1	2.6	4.3	4.9	16.4	121.7	755.1
2003.....	28.7	*	*	*	0.4	1.0	2.7	4.0	5.0	16.4	119.2	769.0
2002.....	26.7	*	*	*	0.4	1.0	2.8	3.4	4.4	15.3	110.3	728.4
2001.....	24.4	*	*	*	0.3	1.0	2.5	3.2	4.3	14.1	102.6	673.1
2000.....	22.2	*	*	*	0.4	1.0	2.3	3.0	4.0	13.0	91.4	625.6
1999.....	20.1	*	*	*	0.3	1.0	2.1	2.6	3.6	11.8	80.6	576.9
Black, both sexes												
2004.....	14.4	*	*	*	0.4	2.1	5.5	13.9	15.5	31.5	133.2	589.4
2003.....	14.0	*	*	*	0.5	1.9	6.9	13.7	16.0	25.4	131.4	578.8
2002.....	13.2	*	*	*	0.4	1.8	7.2	13.7	16.3	26.3	115.9	540.8
2001.....	12.6	*	*	*	0.6	1.9	7.6	15.7	15.6	26.4	100.7	488.5
2000.....	11.8	*	*	*	0.7	2.4	7.3	14.6	15.1	25.3	98.2	442.6
1999.....	10.7	*	*	*	0.5	1.9	8.6	14.6	16.0	24.2	84.7	349.1
Black, male												
2004.....	14.0	*	*	*	*	3.5	7.9	21.2	27.8	46.0	156.2	510.6
2003.....	13.4	*	*	*	0.7	2.5	9.5	22.5	27.3	37.7	148.4	482.7
2002.....	13.1	*	*	*	*	2.4	9.8	22.3	28.4	42.1	135.7	457.8
2001.....	13.0	*	*	*	0.9	3.0	11.0	25.7	28.1	40.3	116.5	423.2
2000.....	12.5	*	*	*	1.1	3.4	10.7	24.1	25.9	39.7	113.5	415.2
1999.....	11.9	*	*	*	0.8	2.5	12.5	23.1	28.2	37.9	111.0	309.3
Black, female												
2004.....	14.8	*	*	*	*	0.7	3.3	7.6	5.6	21.1	120.2	620.6
2003.....	14.5	*	*	*	*	1.4	4.6	6.0	6.9	16.6	121.8	616.1
2002.....	13.3	*	*	*	*	1.3	4.9	6.3	6.6	15.0	104.8	572.6
2001.....	12.1	*	*	*	*	0.9	4.6	7.1	5.6	16.7	91.9	512.8
2000.....	11.2	*	*	*	*	1.4	4.2	6.4	6.4	15.2	89.5	452.9
1999.....	9.6	*	*	*	*	1.2	5.1	7.2	6.3	14.5	70.1	364.0

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Organic dementia (F01,F03)												
All races 3/, both sexes												
2004.....	15.9	*	*	*	*	*	*	0.2	1.4	12.2	103.5	628.2
2003.....	15.7	*	*	*	*	*	*	0.1	1.3	11.8	102.4	631.7
2002.....	14.5	*	*	*	*	*	*	0.1	1.1	10.9	94.3	598.3
2001.....	13.1	*	*	*	*	*	*	0.1	1.1	10.5	84.9	554.0
2000.....	11.7	*	*	*	*	*	*	0.1	1.0	9.3	76.0	505.6
1999.....	10.3	*	*	*	*	*	*	0.1	0.9	8.3	65.8	454.4
All races 3/, male												
2004.....	9.5	*	*	*	*	*	*	0.2	1.5	12.6	97.3	487.8
2003.....	9.3	*	*	*	*	*	*	0.1	1.4	12.9	96.9	480.4
2002.....	8.7	*	*	*	*	*	*	0.1	1.1	12.0	91.4	465.7
2001.....	7.8	*	*	*	*	*	*	0.2	1.2	11.5	81.1	442.4
2000.....	6.9	*	*	*	*	*	*	0.1	1.1	10.4	73.7	398.9
1999.....	6.2	*	*	*	*	*	*	0.1	1.1	9.8	66.4	361.7
All races 3/, female												
2004.....	22.1	*	*	*	*	*	*	0.1	1.2	11.7	107.7	691.3
2003.....	21.9	*	*	*	*	*	*	0.1	1.2	11.0	106.0	698.5
2002.....	20.1	*	*	*	*	*	*	0.1	1.0	10.0	96.3	655.9
2001.....	18.2	*	*	*	*	*	*	0.1	1.0	9.6	87.3	600.8
2000.....	16.3	*	*	*	*	*	*	0.1	0.9	8.3	77.5	549.1
1999.....	14.2	*	*	*	*	*	*	*	0.7	7.1	65.5	491.7
White, both sexes												
2004.....	17.9	*	*	*	*	*	*	0.2	1.4	11.9	104.4	642.8
2003.....	17.6	*	*	*	*	*	*	0.1	1.3	11.8	103.1	647.9
2002.....	16.4	*	*	*	*	*	*	0.1	1.1	10.9	95.4	613.9
2001.....	14.8	*	*	*	*	*	*	0.1	1.1	10.4	86.4	569.1
2000.....	13.1	*	*	*	*	*	*	0.1	0.9	9.0	76.6	519.6
1999.....	11.7	*	*	*	*	*	*	0.1	0.9	8.2	66.7	471.2
White, male												
2004.....	10.5	*	*	*	*	*	*	0.2	1.5	12.3	96.3	496.5
2003.....	10.2	*	*	*	*	*	*	0.1	1.4	12.5	96.2	491.1
2002.....	9.6	*	*	*	*	*	*	*	1.1	11.6	91.2	473.7
2001.....	8.7	*	*	*	*	*	*	0.1	1.1	11.1	81.8	452.2
2000.....	7.6	*	*	*	*	*	*	*	1.0	9.9	73.5	406.1
1999.....	6.9	*	*	*	*	*	*	*	1.1	9.6	66.3	373.7

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Organic dementia (F01,F03)												
White, female												
2004.....	25.0	*	*	*	*	*	*	0.1	1.2	11.5	110.0	708.7
2003.....	24.8	*	*	*	*	*	*	*	1.2	11.1	107.7	717.4
2002.....	23.0	*	*	*	*	*	*	0.2	1.1	10.2	98.2	674.7
2001.....	20.7	*	*	*	*	*	*	0.1	1.1	9.7	89.4	618.0
2000.....	18.5	*	*	*	*	*	*	*	0.9	8.3	78.5	565.7
1999.....	16.3	*	*	*	*	*	*	*	0.7	7.1	67.0	510.4
Black, both sexes												
2004.....	9.2	*	*	*	*	*	*	*	1.8	18.3	120.6	560.3
2003.....	8.8	*	*	*	*	*	*	*	1.8	15.4	118.0	545.3
2002.....	7.9	*	*	*	*	*	*	*	1.3	14.2	102.7	503.1
2001.....	7.0	*	*	*	*	*	*	*	1.5	14.8	86.9	452.3
2000.....	6.4	*	*	*	*	*	*	*	1.6	13.7	83.2	403.2
1999.....	5.0	*	*	*	*	*	*	*	1.1	11.3	67.5	309.3
Black, male												
2004.....	6.3	*	*	*	*	*	*	*	2.3	21.2	138.4	468.9
2003.....	5.8	*	*	*	*	*	*	*	2.5	19.8	128.9	445.2
2002.....	5.3	*	*	*	*	*	*	*	*	18.6	116.2	425.6
2001.....	4.6	*	*	*	*	*	*	*	1.9	19.0	95.1	382.4
2000.....	4.5	*	*	*	*	*	*	*	1.9	18.0	91.7	372.5
1999.....	3.5	*	*	*	*	*	*	*	*	14.3	81.2	266.0
Black, female												
2004.....	11.9	*	*	*	*	*	*	*	1.3	16.2	110.6	596.6
2003.....	11.5	*	*	*	*	*	*	*	1.3	12.2	111.8	584.1
2002.....	10.2	*	*	*	*	*	*	*	1.4	11.1	95.0	532.7
2001.....	9.1	*	*	*	*	*	*	*	*	11.9	82.3	478.2
2000.....	8.1	*	*	*	*	*	*	*	*	10.7	78.4	414.7
1999.....	6.3	*	*	*	*	*	*	*	*	9.1	59.9	325.5

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other and unspecified organic mental disorders (F04-F09)

All races 3/, both sexes

2004.....	0.6	*	*	*	*	*	*	0.1	0.1	0.6	3.8	22.3
2003.....	0.6	*	*	*	*	*	*	0.1	0.1	0.6	4.0	24.5
2002.....	0.7	*	*	*	*	*	*	0.1	0.2	0.8	4.1	26.7
2001.....	0.8	*	*	*	*	*	*	*	0.2	0.8	4.9	29.3
2000.....	0.8	*	*	*	*	*	*	*	0.2	0.8	5.0	33.7
1999.....	0.9	*	*	*	*	*	*	*	0.2	0.8	5.5	37.7

All races 3/, male

2004.....	0.4	*	*	*	*	*	*	*	0.2	0.7	4.2	19.4
2003.....	0.5	*	*	*	*	*	*	*	0.2	0.7	4.6	20.8
2002.....	0.4	*	*	*	*	*	*	*	0.2	1.0	4.2	20.1
2001.....	0.5	*	*	*	*	*	*	*	0.2	1.0	4.8	23.8
2000.....	0.5	*	*	*	*	*	*	*	0.2	1.0	4.5	28.4
1999.....	0.6	*	*	*	*	*	*	*	0.2	0.9	5.6	29.8

All races 3/, female

2004.....	0.8	*	*	*	*	*	*	*	*	0.4	3.6	23.6
2003.....	0.8	*	*	*	*	*	*	*	*	0.5	3.7	26.2
2002.....	0.9	*	*	*	*	*	*	*	0.1	0.7	4.1	29.6
2001.....	1.0	*	*	*	*	*	*	*	*	0.7	5.0	31.6
2000.....	1.1	*	*	*	*	*	*	*	0.2	0.7	5.2	35.8
1999.....	1.2	*	*	*	*	*	*	*	*	0.7	5.4	40.9

White, both sexes

2004.....	0.7	*	*	*	*	*	*	*	0.1	0.6	3.9	23.3
2003.....	0.7	*	*	*	*	*	*	*	0.1	0.6	4.1	25.1
2002.....	0.8	*	*	*	*	*	*	*	0.2	0.8	4.3	27.5
2001.....	0.8	*	*	*	*	*	*	*	0.2	0.8	5.0	30.2
2000.....	0.9	*	*	*	*	*	*	*	0.2	0.9	5.0	34.7
1999.....	1.0	*	*	*	*	*	*	*	0.2	0.7	5.5	38.9

White, male

2004.....	0.5	*	*	*	*	*	*	*	0.2	0.7	4.4	20.1
2003.....	0.5	*	*	*	*	*	*	*	0.2	0.6	4.6	21.3
2002.....	0.5	*	*	*	*	*	*	*	0.2	0.9	4.3	20.6
2001.....	0.5	*	*	*	*	*	*	*	0.2	1.0	4.8	24.2
2000.....	0.6	*	*	*	*	*	*	*	0.2	1.0	4.6	28.5
1999.....	0.6	*	*	*	*	*	*	*	0.2	0.9	5.4	30.1

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other and unspecified organic mental disorders (F04-F09)

White, female

2004.....	0.9	*	*	*	*	*	*	*	*	0.5	3.6	24.7
2003.....	0.9	*	*	*	*	*	*	*	*	0.6	3.8	26.8
2002.....	1.1	*	*	*	*	*	*	*	*	0.6	4.2	30.5
2001.....	1.1	*	*	*	*	*	*	*	*	0.7	5.1	32.6
2000.....	1.3	*	*	*	*	*	*	*	*	0.8	5.3	37.3
1999.....	1.4	*	*	*	*	*	*	*	*	0.6	5.5	42.4

Black, both sexes

2004.....	0.3	*	*	*	*	*	*	*	*	*	3.3	16.6
2003.....	0.4	*	*	*	*	*	*	*	*	*	4.3	20.4
2002.....	0.4	*	*	*	*	*	*	*	*	*	3.3	22.9
2001.....	0.5	*	*	*	*	*	*	*	*	1.2	5.4	24.3
2000.....	0.5	*	*	*	*	*	*	*	*	*	5.7	25.3
1999.....	0.5	*	*	*	*	*	*	*	*	1.3	6.6	28.7

Black, male

2004.....	0.3	*	*	*	*	*	*	*	*	*	*	*
2003.....	0.4	*	*	*	*	*	*	*	*	*	*	*
2002.....	0.2	*	*	*	*	*	*	*	*	*	*	*
2001.....	0.4	*	*	*	*	*	*	*	*	*	6.2	22.7
2000.....	0.4	*	*	*	*	*	*	*	*	*	*	25.4
1999.....	0.5	*	*	*	*	*	*	*	*	*	9.5	29.3

Black, female

2004.....	0.4	*	*	*	*	*	*	*	*	*	3.5	16.5
2003.....	0.4	*	*	*	*	*	*	*	*	*	3.7	20.7
2002.....	0.5	*	*	*	*	*	*	*	*	*	3.6	25.1
2001.....	0.5	*	*	*	*	*	*	*	*	*	5.0	24.8
2000.....	0.5	*	*	*	*	*	*	*	*	*	6.0	25.3
1999.....	0.6	*	*	*	*	*	*	*	*	*	5.0	28.5

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Mental and behavioral disorders due to psychoactive substance use (F10-F19)

All races 3/, both sexes

2004.....	3.6	*	*	*	0.6	1.7	4.6	8.3	7.5	6.4	5.4	4.6
2003.....	3.5	*	*	*	0.5	1.6	4.8	7.9	7.0	6.0	4.7	4.3
2002.....	3.3	*	*	*	0.5	1.5	4.9	7.4	6.8	5.7	4.4	3.7
2001.....	3.2	*	*	*	0.5	1.6	4.5	7.2	6.5	5.4	4.6	4.0
2000.....	3.1	*	*	*	0.5	1.6	4.6	6.8	6.2	5.5	4.7	4.1
1999.....	3.1	*	*	*	0.4	1.6	4.7	6.6	6.4	5.4	4.9	3.8

All races 3/, male

2004.....	5.6	*	*	*	0.8	2.5	6.8	12.7	12.6	10.5	8.9	8.2
2003.....	5.3	*	*	*	0.8	2.4	7.1	12.5	11.3	9.7	8.0	7.3
2002.....	5.1	*	*	*	0.7	2.1	7.1	11.9	11.4	9.3	7.2	7.0
2001.....	4.9	*	*	*	0.7	2.4	6.6	11.5	10.7	9.2	7.2	7.5
2000.....	4.8	*	*	*	0.7	2.4	6.9	10.9	10.3	9.1	8.0	7.3
1999.....	4.8	*	*	*	0.7	2.4	7.3	10.6	10.6	8.9	8.8	7.4

All races 3/, female

2004.....	1.8	*	*	*	0.3	0.8	2.3	4.0	2.8	3.0	3.0	3.0
2003.....	1.7	*	*	*	0.2	0.8	2.6	3.5	3.0	2.9	2.5	3.0
2002.....	1.6	*	*	*	0.2	0.8	2.7	3.1	2.6	2.7	2.5	2.2
2001.....	1.5	*	*	*	0.2	0.8	2.4	3.1	2.6	2.3	2.9	2.5
2000.....	1.5	*	*	*	0.2	0.8	2.2	2.8	2.5	2.5	2.5	2.8
1999.....	1.4	*	*	*	0.2	0.8	2.2	2.7	2.5	2.6	2.4	2.4

White, both sexes

2004.....	3.6	*	*	*	0.7	1.7	4.6	7.8	7.1	6.0	5.5	4.7
2003.....	3.4	*	*	*	0.5	1.6	4.7	7.5	6.4	5.9	4.8	4.5
2002.....	3.2	*	*	*	0.5	1.5	4.7	6.9	6.1	5.4	4.3	3.6
2001.....	3.0	*	*	*	0.5	1.6	4.2	6.4	5.9	5.1	4.6	4.0
2000.....	3.0	*	*	*	0.5	1.6	4.3	6.0	5.6	5.2	4.6	4.1
1999.....	2.9	*	*	*	0.5	1.6	4.3	5.7	5.7	5.0	4.8	3.9

White, male

2004.....	5.5	*	*	*	1.0	2.5	6.9	12.1	11.6	9.5	8.9	8.2
2003.....	5.2	*	*	*	0.8	2.5	7.0	11.6	10.3	9.4	7.9	7.4
2002.....	4.9	*	*	*	0.7	2.2	7.0	11.0	10.0	8.5	7.0	6.8
2001.....	4.6	*	*	*	0.8	2.4	6.2	10.3	9.4	8.4	7.0	7.3
2000.....	4.6	*	*	*	0.7	2.3	6.5	9.7	9.1	8.3	7.7	7.2
1999.....	4.6	*	*	*	0.7	2.4	6.7	9.4	9.2	7.9	8.4	7.4

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Mental and behavioral disorders due to psychoactive substance use (F10-F19)

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
White, female												
2004.....	1.8	*	*	*	0.3	0.9	2.3	3.7	2.7	3.1	3.1	3.1
2003.....	1.7	*	*	*	0.2	0.7	2.4	3.4	2.8	2.9	2.6	3.2
2002.....	1.5	*	*	*	0.2	0.8	2.4	2.8	2.4	2.9	2.6	2.2
2001.....	1.5	*	*	*	0.2	0.8	2.2	2.7	2.6	2.4	3.0	2.6
2000.....	1.4	*	*	*	0.3	0.8	2.0	2.4	2.3	2.5	2.6	2.9
1999.....	1.3	*	*	*	0.2	0.7	1.8	2.2	2.3	2.6	2.4	2.5
Black, both sexes												
2004.....	4.3	*	*	*	*	1.7	4.9	13.1	12.7	10.9	5.1	*
2003.....	4.3	*	*	*	*	1.7	6.4	12.5	13.1	8.0	4.9	*
2002.....	4.4	*	*	*	*	1.6	6.7	12.9	13.9	9.0	5.5	*
2001.....	4.6	*	*	*	0.4	1.6	7.1	14.7	13.0	8.9	5.7	*
2000.....	4.6	*	*	*	0.5	2.1	6.8	13.9	12.7	9.4	5.9	*
1999.....	4.7	*	*	*	0.4	1.6	8.1	13.8	13.8	10.2	7.3	*
Black, male												
2004.....	6.9	*	*	*	*	2.9	7.3	20.2	24.2	22.2	10.6	*
2003.....	6.7	*	*	*	*	2.2	8.9	21.2	23.5	14.9	10.7	*
2002.....	7.0	*	*	*	*	2.2	9.1	21.3	26.0	19.0	11.5	*
2001.....	7.4	*	*	*	0.7	2.5	10.3	24.1	24.5	18.2	11.8	*
2000.....	7.3	*	*	*	0.9	3.1	9.9	23.3	22.9	18.7	13.3	*
1999.....	7.4	*	*	*	*	2.3	11.9	22.0	25.5	20.4	15.4	*
Black, female												
2004.....	2.0	*	*	*	*	*	2.8	6.9	3.4	2.9	*	*
2003.....	2.0	*	*	*	*	1.2	4.2	5.0	4.8	3.0	*	*
2002.....	2.1	*	*	*	*	1.0	4.5	5.8	4.2	*	*	*
2001.....	2.1	*	*	*	*	0.9	4.2	6.7	3.7	2.2	*	*
2000.....	2.1	*	*	*	*	1.2	4.1	5.9	4.4	2.9	*	*
1999.....	2.3	*	*	*	*	1.1	4.8	6.8	4.4	3.0	*	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Mental and behavioral disorders due to use of alcohol (F10)

All races 3/, both sexes

2004.....	2.5	*	*	*	0.2	0.7	2.8	5.8	6.1	4.9	3.2	1.9
2003.....	2.4	*	*	*	0.2	0.7	3.0	5.8	5.9	4.5	2.9	2.0
2002.....	2.4	*	*	*	0.2	0.6	3.1	5.6	5.8	4.6	3.0	2.0
2001.....	2.3	*	*	*	0.2	0.8	2.9	5.5	5.7	4.6	3.2	2.1
2000.....	2.2	*	*	*	0.1	0.7	2.9	5.1	5.4	4.6	3.2	2.5
1999.....	2.2	*	*	*	0.2	0.7	2.9	5.0	5.5	4.4	3.5	2.3

All races 3/, male

2004.....	3.9	*	*	*	0.3	1.1	4.3	9.4	10.4	8.5	5.9	3.6
2003.....	3.8	*	*	*	0.3	1.1	4.6	9.3	9.7	7.7	5.5	3.5
2002.....	3.8	*	*	*	0.2	1.0	4.7	9.1	10.0	8.0	5.7	4.4
2001.....	3.7	*	*	*	0.3	1.1	4.5	8.9	9.5	8.1	5.6	4.7
2000.....	3.6	*	*	*	0.2	1.1	4.6	8.4	9.2	8.1	6.0	5.2
1999.....	3.6	*	*	*	0.3	1.1	4.6	8.3	9.2	7.6	6.8	4.9

All races 3/, female

2004.....	1.0	*	*	*	0.1	0.4	1.2	2.5	2.1	1.9	1.4	1.2
2003.....	1.1	*	*	*	*	0.3	1.5	2.5	2.3	1.9	1.1	1.3
2002.....	1.0	*	*	*	*	0.3	1.5	2.2	2.0	1.8	1.3	1.0
2001.....	1.0	*	*	*	*	0.4	1.4	2.2	2.2	1.7	1.6	1.1
2000.....	0.9	*	*	*	*	0.3	1.2	1.9	1.9	1.7	1.3	1.5
1999.....	0.9	*	*	*	*	0.4	1.3	1.9	2.1	1.8	1.3	1.3

White, both sexes

2004.....	2.5	*	*	*	0.2	0.7	2.9	5.9	5.9	4.4	3.2	2.0
2003.....	2.4	*	*	*	0.2	0.7	3.1	5.8	5.5	4.4	2.8	2.0
2002.....	2.4	*	*	*	0.2	0.7	3.1	5.5	5.3	4.3	2.9	2.0
2001.....	2.3	*	*	*	0.2	0.7	2.8	5.1	5.2	4.3	3.1	2.1
2000.....	2.2	*	*	*	0.2	0.7	2.9	4.8	4.9	4.2	3.1	2.5
1999.....	2.1	*	*	*	0.2	0.8	2.8	4.6	4.9	4.0	3.2	2.3

White, male

2004.....	4.0	*	*	*	0.3	1.1	4.5	9.3	9.9	7.5	5.9	3.7
2003.....	3.8	*	*	*	0.3	1.1	4.7	9.0	9.0	7.4	5.2	3.4
2002.....	3.8	*	*	*	0.3	1.0	4.8	9.0	8.9	7.2	5.4	4.1
2001.....	3.6	*	*	*	0.3	1.1	4.3	8.3	8.4	7.3	5.5	4.4
2000.....	3.5	*	*	*	0.2	1.1	4.6	7.8	8.2	7.3	5.6	5.1
1999.....	3.4	*	*	*	0.3	1.1	4.4	7.6	8.1	6.7	6.4	4.7

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Mental and behavioral disorders due to use of alcohol (F10)												
White, female												
2004.....	1.1	*	*	*	*	0.3	1.3	2.5	2.1	1.8	1.4	1.2
2003.....	1.1	*	*	*	*	0.3	1.4	2.6	2.3	1.8	1.2	1.4
2002.....	1.0	*	*	*	*	0.3	1.4	2.2	1.9	1.8	1.3	1.1
2001.....	1.0	*	*	*	*	0.4	1.4	2.0	2.1	1.7	1.6	1.1
2000.....	0.9	*	*	*	*	0.3	1.1	1.8	1.8	1.7	1.4	1.5
1999.....	0.9	*	*	*	*	0.4	1.1	1.7	1.9	1.8	1.2	1.3
Black, both sexes												
2004.....	2.5	*	*	*	*	0.7	2.1	6.6	9.1	9.6	3.6	*
2003.....	2.5	*	*	*	*	0.6	2.8	7.1	9.6	6.7	4.4	*
2002.....	2.7	*	*	*	*	0.6	3.2	7.2	11.0	8.1	4.8	*
2001.....	3.0	*	*	*	*	0.6	3.6	9.0	11.0	8.1	4.7	*
2000.....	2.9	*	*	*	*	0.9	3.2	8.7	10.7	8.6	5.0	*
1999.....	3.0	*	*	*	*	0.5	4.0	8.4	11.4	9.1	6.5	*
Black, male												
2004.....	4.3	*	*	*	*	1.2	3.5	11.5	17.8	19.3	7.5	*
2003.....	4.3	*	*	*	*	1.0	4.3	12.8	17.7	12.5	9.9	*
2002.....	4.6	*	*	*	*	1.1	4.7	12.3	20.8	17.3	9.8	*
2001.....	5.2	*	*	*	*	1.0	6.0	15.2	21.0	16.9	10.0	*
2000.....	4.9	*	*	*	*	1.5	4.9	15.0	20.0	17.7	12.4	*
1999.....	4.9	*	*	*	*	0.8	6.1	13.9	21.5	18.5	13.2	*
Black, female												
2004.....	0.9	*	*	*	*	*	0.9	2.5	2.1	2.7	*	*
2003.....	0.9	*	*	*	*	*	1.5	2.1	3.0	2.5	*	*
2002.....	1.0	*	*	*	*	*	1.8	2.8	3.1	*	*	*
2001.....	1.1	*	*	*	*	*	1.6	3.6	2.9	*	*	*
2000.....	1.1	*	*	*	*	*	1.7	3.3	3.2	2.3	*	*
1999.....	1.3	*	*	*	*	*	2.2	3.6	3.4	2.4	*	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Mental and behavioral disorders due to other psychoactive substance use (F11-F19)

All races 3/, both sexes

2004.....	1.2	*	*	*	0.4	1.0	1.8	2.4	1.4	1.5	2.2	2.7
2003.....	1.1	*	*	*	0.3	0.9	1.8	2.1	1.2	1.5	1.8	2.3
2002.....	0.9	*	*	*	0.3	0.8	1.8	1.8	1.0	1.1	1.3	1.7
2001.....	0.9	*	*	*	0.3	0.8	1.6	1.8	0.8	0.8	1.4	1.8
2000.....	0.9	*	*	*	0.3	0.9	1.6	1.6	0.8	0.9	1.5	1.6
1999.....	0.9	*	*	*	0.3	0.9	1.8	1.6	0.9	1.0	1.4	1.5

All races 3/, male

2004.....	1.6	*	*	*	0.6	1.4	2.5	3.3	2.1	2.0	3.0	4.6
2003.....	1.5	*	*	*	0.5	1.3	2.5	3.2	1.7	2.0	2.5	3.8
2002.....	1.3	*	*	*	0.4	1.1	2.4	2.8	1.4	1.3	1.5	2.6
2001.....	1.2	*	*	*	0.4	1.2	2.1	2.7	1.2	1.1	1.6	2.8
2000.....	1.2	*	*	*	0.5	1.2	2.3	2.5	1.1	1.0	1.9	2.1
1999.....	1.3	*	*	*	0.4	1.3	2.7	2.3	1.3	1.3	2.0	2.5

All races 3/, female

2004.....	0.7	*	*	*	0.2	0.5	1.1	1.5	0.7	1.1	1.6	1.8
2003.....	0.7	*	*	*	0.2	0.5	1.1	1.1	0.7	1.0	1.3	1.7
2002.....	0.6	*	*	*	0.2	0.5	1.2	0.9	0.6	1.0	1.2	1.2
2001.....	0.5	*	*	*	0.1	0.4	1.0	0.9	0.5	0.6	1.3	1.4
2000.....	0.6	*	*	*	0.2	0.6	1.0	0.8	0.6	0.8	1.2	1.3
1999.....	0.5	*	*	*	0.1	0.4	0.9	0.8	0.4	0.8	1.1	1.1

White, both sexes

2004.....	1.1	*	*	*	0.4	1.0	1.7	1.9	1.2	1.6	2.3	2.7
2003.....	1.0	*	*	*	0.4	0.9	1.6	1.7	0.9	1.5	2.0	2.5
2002.....	0.9	*	*	*	0.3	0.9	1.6	1.4	0.8	1.2	1.4	1.6
2001.....	0.8	*	*	*	0.3	0.9	1.3	1.3	0.7	0.9	1.4	1.9
2000.....	0.8	*	*	*	0.4	0.9	1.4	1.2	0.7	0.9	1.6	1.6
1999.....	0.8	*	*	*	0.3	0.9	1.5	1.1	0.7	1.0	1.5	1.6

White, male

2004.....	1.5	*	*	*	0.6	1.5	2.4	2.7	1.7	2.0	3.1	4.5
2003.....	1.4	*	*	*	0.5	1.3	2.3	2.6	1.3	2.1	2.7	4.0
2002.....	1.2	*	*	*	0.5	1.2	2.2	2.1	1.1	1.3	1.6	2.7
2001.....	1.1	*	*	*	0.5	1.3	1.9	2.0	1.0	1.1	1.6	2.9
2000.....	1.1	*	*	*	0.5	1.2	2.0	1.9	1.0	1.0	2.1	2.1
1999.....	1.1	*	*	*	0.4	1.3	2.3	1.7	1.1	1.3	2.0	2.7

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Mental and behavioral disorders due to other psychoactive substance use (F11-F19)

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
White, female												
2004.....	0.7	*	*	*	0.2	0.5	1.0	1.2	0.6	1.3	1.7	2.0
2003.....	0.6	*	*	*	0.2	0.4	0.9	0.9	0.6	1.1	1.5	1.8
2002.....	0.6	*	*	*	0.2	0.5	1.0	0.6	0.6	1.1	1.3	1.1
2001.....	0.5	*	*	*	0.2	0.4	0.8	0.6	0.5	0.7	1.4	1.5
2000.....	0.5	*	*	*	0.2	0.5	0.8	0.6	0.5	0.9	1.2	1.4
1999.....	0.4	*	*	*	0.1	0.4	0.7	0.5	0.4	0.8	1.2	1.1
Black, both sexes												
2004.....	1.8	*	*	*	*	1.0	2.8	6.5	3.6	1.3	*	*
2003.....	1.7	*	*	*	*	1.1	3.6	5.5	3.5	1.3	*	*
2002.....	1.7	*	*	*	*	1.0	3.5	5.8	2.9	*	*	*
2001.....	1.6	*	*	*	*	1.1	3.4	5.8	2.0	*	*	*
2000.....	1.6	*	*	*	0.4	1.3	3.6	5.2	2.0	*	*	*
1999.....	1.7	*	*	*	*	1.1	4.1	5.4	2.3	*	*	*
Black, male												
2004.....	2.6	*	*	*	*	1.7	3.8	8.8	6.4	2.9	*	*
2003.....	2.4	*	*	*	*	1.2	4.6	8.4	5.8	*	*	*
2002.....	2.4	*	*	*	*	1.1	4.4	9.0	5.2	*	*	*
2001.....	2.3	*	*	*	*	1.5	4.3	8.9	3.4	*	*	*
2000.....	2.3	*	*	*	*	1.7	5.0	8.3	2.9	*	*	*
1999.....	2.5	*	*	*	*	1.5	5.9	8.1	4.1	*	*	*
Black, female												
2004.....	1.1	*	*	*	*	*	1.9	4.5	1.3	*	*	*
2003.....	1.1	*	*	*	*	1.0	2.7	2.9	1.8	*	*	*
2002.....	1.1	*	*	*	*	0.8	2.8	3.0	*	*	*	*
2001.....	1.0	*	*	*	*	*	2.6	3.1	*	*	*	*
2000.....	1.0	*	*	*	*	0.9	2.4	2.6	*	*	*	*
1999.....	1.0	*	*	*	*	0.7	2.6	3.2	*	*	*	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Schizophrenia, schizotypal and delusional disorders (F20-F29)												
All races 3/, both sexes												
2004.....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.5	1.1	2.7
2003.....	0.2	*	*	*	*	*	0.1	0.1	0.1	0.5	1.1	2.7
2002.....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.5	1.1	3.3
2001.....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.4	1.2	2.7
2000.....	0.2	*	*	*	*	*	0.0	0.1	0.2	0.5	1.2	3.3
1999.....	0.2	*	*	*	*	*	0.0	0.1	0.2	0.4	1.4	4.9
All races 3/, male												
2004.....	0.2	*	*	*	*	*	*	0.2	0.3	0.5	1.0	2.3
2003.....	0.1	*	*	*	*	*	*	0.2	*	0.4	0.8	1.8
2002.....	0.1	*	*	*	*	*	*	0.1	0.2	0.5	0.9	1.9
2001.....	0.1	*	*	*	*	*	0.1	0.1	0.2	0.5	1.0	1.5
2000.....	0.1	*	*	*	*	*	*	0.1	0.2	0.5	1.1	2.4
1999.....	0.2	*	*	*	*	*	*	0.1	*	0.4	1.2	4.6
All races 3/, female												
2004.....	0.2	*	*	*	*	*	*	*	0.2	0.4	1.1	2.8
2003.....	0.2	*	*	*	*	*	*	*	0.1	0.6	1.3	3.0
2002.....	0.2	*	*	*	*	*	*	*	*	0.5	1.3	3.9
2001.....	0.2	*	*	*	*	*	*	*	0.2	0.3	1.3	3.1
2000.....	0.2	*	*	*	*	*	*	*	*	0.5	1.3	3.8
1999.....	0.2	*	*	*	*	*	*	*	*	0.4	1.6	5.0
White, both sexes												
2004.....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.5	1.1	2.6
2003.....	0.2	*	*	*	*	*	*	0.1	0.1	0.6	1.1	2.8
2002.....	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.1	3.4
2001.....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.4	1.2	2.8
2000.....	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.3	3.4
1999.....	0.2	*	*	*	*	*	*	0.1	0.2	0.4	1.4	5.0
White, male												
2004.....	0.2	*	*	*	*	*	*	0.2	0.3	0.5	0.9	1.9
2003.....	0.1	*	*	*	*	*	*	0.2	*	0.5	0.9	1.9
2002.....	0.1	*	*	*	*	*	*	0.1	0.2	0.4	0.8	1.9
2001.....	0.1	*	*	*	*	*	*	*	*	0.5	1.0	*
2000.....	0.2	*	*	*	*	*	*	*	0.2	0.5	1.1	2.4
1999.....	0.2	*	*	*	*	*	*	*	*	0.4	1.1	4.7

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Mood (affective) disorders (F30-F39)												
All races 3/, both sexes												
2004.....	0.3	*	*	*	*	*	*	0.1	0.1	0.4	1.9	8.4
2003.....	0.3	*	*	*	*	*	*	0.1	0.1	0.4	2.1	10.5
2002.....	0.3	*	*	*	*	*	*	0.1	0.1	0.4	2.2	9.1
2001.....	0.3	*	*	*	*	*	0.1	0.1	0.1	0.5	2.2	9.6
2000.....	0.3	*	*	*	*	*	*	0.1	0.1	0.4	1.9	9.8
1999.....	0.3	*	*	*	*	*	*	0.1	0.1	0.4	2.4	9.4
All races 3/, male												
2004.....	0.2	*	*	*	*	*	*	*	*	0.5	1.5	7.2
2003.....	0.2	*	*	*	*	*	*	*	*	0.4	1.8	8.0
2002.....	0.2	*	*	*	*	*	*	*	*	0.4	1.6	6.6
2001.....	0.2	*	*	*	*	*	*	0.1	*	0.5	1.7	8.0
2000.....	0.2	*	*	*	*	*	*	*	*	0.3	1.4	9.1
1999.....	0.2	*	*	*	*	*	*	*	0.2	0.4	2.0	8.8
All races 3/, female												
2004.....	0.4	*	*	*	*	*	*	*	0.2	0.4	2.2	9.0
2003.....	0.4	*	*	*	*	*	*	*	0.1	0.4	2.3	11.6
2002.....	0.4	*	*	*	*	*	*	*	0.2	0.5	2.7	10.1
2001.....	0.4	*	*	*	*	*	*	*	*	0.5	2.5	10.3
2000.....	0.4	*	*	*	*	*	*	*	*	0.4	2.2	10.1
1999.....	0.4	*	*	*	*	*	*	*	*	0.5	2.6	9.7
White, both sexes												
2004.....	0.3	*	*	*	*	*	*	0.1	0.1	0.5	2.0	9.2
2003.....	0.4	*	*	*	*	*	*	0.1	0.1	0.4	2.2	11.3
2002.....	0.4	*	*	*	*	*	*	0.1	0.1	0.5	2.3	9.8
2001.....	0.4	*	*	*	*	*	0.1	*	0.1	0.5	2.3	10.1
2000.....	0.3	*	*	*	*	*	*	0.1	0.2	0.4	2.0	10.6
1999.....	0.4	*	*	*	*	*	*	*	0.1	0.5	2.5	10.2
White, male												
2004.....	0.2	*	*	*	*	*	*	*	*	0.5	1.5	7.7
2003.....	0.2	*	*	*	*	*	*	*	*	0.4	2.0	8.6
2002.....	0.2	*	*	*	*	*	*	*	*	0.4	1.6	7.2
2001.....	0.2	*	*	*	*	*	*	*	*	0.5	1.9	8.3
2000.....	0.2	*	*	*	*	*	*	*	*	0.4	1.4	10.1
1999.....	0.3	*	*	*	*	*	*	*	*	0.4	2.1	9.5

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Mental retardation (F70-F79)												
All races 3/, both sexes												
2004.....	0.2	*	*	*	0.1	0.1	0.1	0.3	0.3	0.4	0.5	*
2003.....	0.2	*	*	*	0.1	0.1	0.1	0.3	0.4	0.4	0.4	0.4
2002.....	0.2	*	*	*	0.1	0.1	0.2	0.2	0.4	0.4	0.4	0.5
2001.....	0.1	*	*	*	0.1	0.1	0.1	0.2	0.3	0.4	0.4	*
2000.....	0.1	*	*	*	0.1	0.1	0.1	0.2	0.3	0.3	0.4	0.5
1999.....	0.2	*	*	0.1	0.1	0.1	0.2	0.2	0.3	0.3	0.4	*
All races 3/, male												
2004.....	0.2	*	*	*	*	0.1	0.2	0.4	0.3	0.5	0.5	*
2003.....	0.2	*	*	*	*	0.1	0.2	0.3	0.4	0.4	0.6	*
2002.....	0.2	*	*	*	*	*	0.2	0.3	0.4	0.4	0.5	*
2001.....	0.2	*	*	*	*	0.1	0.1	0.2	0.3	0.4	0.6	*
2000.....	0.2	*	*	*	0.1	0.1	0.2	0.2	0.3	0.3	0.5	*
1999.....	0.2	*	*	*	*	0.1	0.2	0.3	0.3	0.5	0.6	*
All races 3/, female												
2004.....	0.1	*	*	*	*	*	0.1	0.2	0.3	0.2	0.5	*
2003.....	0.1	*	*	*	*	*	0.1	0.2	0.3	0.3	0.3	*
2002.....	0.1	*	*	*	*	*	0.1	0.1	0.3	0.3	0.3	*
2001.....	0.1	*	*	*	*	*	0.1	0.2	0.2	0.3	0.3	*
2000.....	0.1	*	*	*	*	*	0.1	0.2	0.3	0.2	0.3	*
1999.....	0.1	*	*	*	*	0.1	0.2	0.1	0.2	0.2	0.3	*
White, both sexes												
2004.....	0.2	*	*	*	*	0.1	0.1	0.3	0.3	0.4	0.5	*
2003.....	0.2	*	*	*	0.1	0.1	0.1	0.3	0.4	0.4	0.4	0.5
2002.....	0.2	*	*	*	*	0.1	0.2	0.2	0.4	0.4	0.4	0.6
2001.....	0.2	*	*	*	*	0.1	0.1	0.2	0.3	0.4	0.4	*
2000.....	0.2	*	*	*	0.1	0.1	0.1	0.2	0.3	0.3	0.4	0.5
1999.....	0.2	*	*	*	0.1	0.1	0.2	0.2	0.3	0.3	0.4	*
White, male												
2004.....	0.2	*	*	*	*	*	0.2	0.4	0.3	0.6	0.5	*
2003.....	0.2	*	*	*	*	*	0.2	0.4	0.4	0.4	0.6	*
2002.....	0.2	*	*	*	*	*	0.2	0.3	0.5	0.4	0.5	*
2001.....	0.2	*	*	*	*	*	0.1	0.2	0.3	0.4	0.6	*
2000.....	0.2	*	*	*	*	*	0.2	0.2	0.4	0.3	0.5	*
1999.....	0.2	*	*	*	*	0.1	0.2	0.3	0.3	0.5	0.7	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other and unspecified mental and behavioral disorders (F50-F69,F80-F99)

All races 3/, both sexes

2004.....	0.2	*	*	*	0.1	*	0.1	0.1	0.1	0.1	0.7	3.9
2003.....	0.2	*	*	*	0.1	0.1	0.1	0.1	0.1	0.2	0.9	4.4
2002.....	0.2	*	*	*	*	0.1	0.1	0.1	*	0.2	0.7	4.4
2001.....	0.1	*	*	*	*	0.1	*	*	*	0.2	0.7	4.1
2000.....	0.1	*	*	*	*	*	*	*	0.1	0.1	0.7	3.9
1999.....	0.1	*	*	*	*	*	0.1	0.1	*	0.2	0.7	4.0

All races 3/, male

2004.....	0.1	*	*	*	*	*	*	*	*	*	0.7	3.7
2003.....	0.1	*	*	*	*	*	*	*	*	*	1.0	4.2
2002.....	0.1	*	*	*	*	*	*	*	*	*	0.9	3.6
2001.....	0.1	*	*	*	*	*	*	*	*	*	0.7	3.5
2000.....	0.1	*	*	*	*	*	*	*	*	*	0.6	3.3
1999.....	0.1	*	*	*	*	*	*	*	*	0.3	0.6	2.5

All races 3/, female

2004.....	0.2	*	*	*	0.1	*	0.1	0.1	0.2	*	0.8	4.0
2003.....	0.2	*	*	*	0.1	0.1	0.1	*	*	0.3	0.8	4.5
2002.....	0.2	*	*	*	*	0.1	0.1	0.1	*	0.2	0.6	4.7
2001.....	0.2	*	*	*	*	*	*	*	*	*	0.7	4.4
2000.....	0.2	*	*	*	*	*	*	*	*	*	0.8	4.1
1999.....	0.2	*	*	*	*	*	*	*	*	0.2	0.7	4.7

White, both sexes

2004.....	0.2	*	*	*	0.1	*	0.1	0.1	0.1	*	0.7	4.0
2003.....	0.2	*	*	*	0.1	*	0.1	0.1	0.1	0.2	0.9	4.4
2002.....	0.2	*	*	*	*	0.1	0.1	0.1	*	0.2	0.8	4.4
2001.....	0.2	*	*	*	*	*	*	*	*	0.2	0.7	4.4
2000.....	0.1	*	*	*	*	*	*	*	*	0.1	0.7	4.0
1999.....	0.2	*	*	*	*	*	0.1	*	*	0.2	0.7	4.3

White, male

2004.....	0.1	*	*	*	*	*	*	*	*	*	0.7	3.5
2003.....	0.1	*	*	*	*	*	*	*	*	*	0.9	4.2
2002.....	0.1	*	*	*	*	*	*	*	*	*	0.9	3.7
2001.....	0.1	*	*	*	*	*	*	*	*	*	0.7	3.7
2000.....	0.1	*	*	*	*	*	*	*	*	*	0.6	3.2
1999.....	0.1	*	*	*	*	*	*	*	*	0.3	0.6	2.7

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
VI. Diseases of the nervous system (G00-G98)												
All races 3/, both sexes												
2004.....	38.1	9.2	2.0	1.3	2.0	2.3	4.1	9.2	20.1	61.3	282.8	1,014.7
2003.....	37.7	10.0	1.8	1.3	2.2	2.3	4.2	9.3	20.3	64.2	280.2	1,006.7
2002.....	35.7	9.5	2.1	1.2	2.0	2.2	4.2	9.1	19.7	62.0	270.2	951.0
2001.....	33.9	8.9	2.1	1.2	2.1	2.2	4.1	8.7	19.4	60.2	258.9	906.2
2000.....	32.3	10.2	2.1	1.3	1.9	2.1	4.1	9.1	19.2	59.1	248.5	864.9
1999.....	30.1	11.6	2.4	1.3	2.0	2.1	3.9	8.5	17.9	57.1	234.2	790.8
All races 3/, male												
2004.....	30.0	10.5	2.0	1.5	2.6	2.7	4.6	10.0	21.5	68.8	304.9	910.7
2003.....	29.9	11.0	1.8	1.5	2.8	2.7	4.8	10.3	21.9	71.7	305.4	907.9
2002.....	28.5	11.0	2.4	1.4	2.6	2.7	4.9	9.8	21.2	70.3	293.9	867.4
2001.....	27.4	10.0	2.2	1.3	2.8	2.4	4.7	9.2	20.3	68.7	289.2	861.6
2000.....	26.2	11.5	2.1	1.3	2.4	2.5	4.6	9.9	20.6	68.3	277.0	826.2
1999.....	25.0	12.2	2.6	1.4	2.8	2.4	4.4	9.0	19.7	65.6	268.8	772.7
All races 3/, female												
2004.....	45.9	8.0	2.0	1.1	1.4	1.9	3.6	8.3	18.7	54.9	267.9	1,061.5
2003.....	45.3	9.0	1.8	1.1	1.5	2.0	3.5	8.3	18.8	58.0	263.4	1,050.4
2002.....	42.8	7.8	1.7	1.1	1.4	1.8	3.5	8.4	18.3	55.0	254.4	987.3
2001.....	40.2	7.8	1.9	1.1	1.4	1.9	3.5	8.2	18.4	53.2	239.0	924.9
2000.....	38.2	8.8	2.0	1.2	1.3	1.8	3.6	8.2	18.0	51.6	229.9	880.7
1999.....	35.1	11.0	2.1	1.1	1.3	1.7	3.4	8.1	16.2	50.2	211.8	798.2
White, both sexes												
2004.....	42.7	9.1	1.8	1.2	1.9	2.1	4.0	9.0	20.2	63.5	293.3	1,051.0
2003.....	42.4	9.2	1.6	1.2	2.1	2.2	4.0	9.0	20.4	66.4	291.6	1,047.3
2002.....	40.2	9.3	1.8	1.1	1.9	2.1	4.0	8.8	19.9	64.1	281.7	989.7
2001.....	38.0	7.9	1.8	1.1	2.0	2.0	3.9	8.5	19.4	62.4	268.7	942.3
2000.....	36.3	9.4	1.8	1.2	1.8	2.0	3.9	8.8	19.4	60.9	258.2	900.8
1999.....	33.7	10.4	2.2	1.2	1.9	2.0	3.7	8.2	17.9	58.6	243.2	825.1
White, male												
2004.....	33.4	10.6	1.8	1.4	2.4	2.6	4.6	9.9	21.7	71.2	316.4	945.7
2003.....	33.3	11.1	1.5	1.4	2.9	2.5	4.6	10.0	22.0	74.0	317.1	944.4
2002.....	31.7	11.0	2.1	1.3	2.4	2.6	4.7	9.4	21.5	72.2	305.7	904.3
2001.....	30.5	8.7	2.1	1.2	2.7	2.1	4.5	9.0	20.4	70.9	300.2	896.4
2000.....	29.1	10.3	2.0	1.3	2.4	2.4	4.3	9.8	20.6	70.4	287.3	859.9
1999.....	27.7	11.8	2.5	1.2	2.7	2.3	4.2	8.7	19.7	67.5	279.6	806.2

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
VI. Diseases of the nervous system (G00-G98)												
White, female												
2004.....	51.8	7.5	1.7	1.0	1.3	1.6	3.4	8.2	18.9	56.9	277.5	1,098.4
2003.....	51.4	7.1	1.7	1.0	1.4	1.9	3.3	8.0	19.0	59.9	274.4	1,092.9
2002.....	48.6	7.5	1.5	1.0	1.2	1.6	3.3	8.2	18.5	57.1	265.6	1,026.8
2001.....	45.4	7.1	1.6	1.0	1.2	1.8	3.4	8.1	18.4	55.2	247.8	961.5
2000.....	43.2	8.4	1.6	1.1	1.2	1.6	3.5	7.8	18.2	52.9	239.0	917.4
1999.....	39.5	9.0	1.8	1.1	1.1	1.6	3.3	7.8	16.1	51.2	219.4	832.7
Black, both sexes												
2004.....	21.9	10.4	3.2	1.8	3.0	4.1	6.0	12.8	25.1	57.4	231.7	776.9
2003.....	20.8	14.6	2.8	2.0	2.6	3.9	6.7	13.8	24.9	60.7	213.9	699.3
2002.....	19.5	11.6	3.3	1.6	2.9	3.8	6.6	13.7	22.9	58.0	197.3	655.4
2001.....	19.0	14.4	3.1	1.6	3.0	4.1	6.4	12.6	23.9	54.4	199.5	624.9
2000.....	17.9	14.3	3.4	1.8	2.5	3.5	6.6	13.5	22.8	55.9	183.9	561.2
1999.....	16.7	17.8	3.3	1.9	3.0	3.1	6.2	13.4	22.1	55.2	167.6	485.3
Black, male												
2004.....	17.3	10.4	3.2	1.7	3.7	3.8	6.3	14.4	27.1	64.0	239.4	661.6
2003.....	17.5	12.5	3.2	2.1	3.1	4.6	7.4	15.2	28.0	69.2	235.5	620.4
2002.....	16.7	11.9	4.0	1.9	3.6	4.5	7.6	15.1	24.9	69.9	218.9	570.4
2001.....	16.1	17.1	2.9	1.4	3.8	4.7	7.5	13.7	25.3	63.3	218.9	593.4
2000.....	15.4	17.9	3.1	1.9	2.6	3.7	7.7	13.6	26.9	64.2	205.0	537.4
1999.....	14.6	15.9	3.2	2.3	3.9	3.6	7.2	14.8	23.9	61.3	180.2	481.6
Black, female												
2004.....	26.1	10.5	3.3	1.9	2.3	4.3	5.7	11.5	23.6	52.7	227.4	822.6
2003.....	23.9	16.7	2.3	1.8	2.2	3.2	6.2	12.6	22.4	54.6	201.7	730.0
2002.....	22.2	11.2	2.7	1.4	2.2	3.2	5.7	12.4	21.2	49.4	185.1	688.0
2001.....	21.5	11.7	3.4	1.8	2.2	3.4	5.4	11.6	22.9	48.1	188.5	636.5
2000.....	20.2	10.6	3.6	1.8	2.4	3.3	5.7	13.5	19.5	50.0	172.0	570.1
1999.....	18.5	19.8	3.4	1.4	2.0	2.7	5.3	12.2	20.6	50.9	160.6	486.7

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Meningitis (G00,G03)												
All races 3/, both sexes												
2004.....	0.2	1.6	0.2	0.1	0.1	0.1	0.2	0.3	0.3	0.5	0.7	0.9
2003.....	0.3	1.9	0.2	*	0.1	0.1	0.2	0.3	0.4	0.5	0.7	1.1
2002.....	0.2	1.8	*	0.1	0.1	0.1	0.1	0.3	0.3	0.6	0.7	1.2
2001.....	0.3	1.9	0.2	0.1	0.1	0.1	0.2	0.3	0.4	0.5	0.8	0.9
2000.....	0.3	2.4	0.2	0.1	*	0.1	0.2	0.3	0.4	0.6	1.0	1.2
1999.....	0.3	3.1	0.3	0.1	0.1	0.1	0.2	0.3	0.4	0.6	1.0	1.5
All races 3/, male												
2004.....	0.2	1.5	*	*	*	0.1	0.2	0.4	0.3	0.4	0.7	*
2003.....	0.3	1.9	*	*	0.1	0.1	0.2	0.4	0.5	0.5	0.8	*
2002.....	0.3	2.0	*	*	*	*	0.2	0.4	0.4	0.7	0.8	*
2001.....	0.3	1.9	*	*	0.1	0.1	0.2	0.3	0.5	0.6	0.9	*
2000.....	0.3	2.5	0.3	*	*	0.1	0.2	0.3	0.5	0.7	1.3	*
1999.....	0.3	3.1	0.3	*	0.1	0.1	0.2	0.4	0.4	0.7	1.0	1.7
All races 3/, female												
2004.....	0.2	1.7	*	*	*	*	0.1	0.2	0.3	0.6	0.6	0.7
2003.....	0.2	2.0	*	*	*	*	0.1	0.2	0.3	0.5	0.6	1.0
2002.....	0.2	1.7	*	*	*	*	*	0.2	0.3	0.5	0.6	1.1
2001.....	0.2	2.0	*	*	*	*	0.1	0.2	0.4	0.4	0.6	0.8
2000.....	0.2	2.3	*	*	*	0.1	0.1	0.2	0.3	0.5	0.8	1.0
1999.....	0.3	3.0	*	*	*	0.1	0.1	0.2	0.4	0.5	1.0	1.4
White, both sexes												
2004.....	0.2	1.4	*	*	*	*	0.1	0.2	0.3	0.4	0.6	0.8
2003.....	0.2	1.7	*	*	0.1	0.1	0.1	0.2	0.3	0.5	0.7	1.0
2002.....	0.2	1.7	*	0.1	*	0.1	0.1	0.2	0.3	0.6	0.6	1.2
2001.....	0.2	1.5	*	*	0.1	0.1	0.1	0.2	0.4	0.5	0.7	0.8
2000.....	0.2	1.8	0.2	0.1	*	0.1	0.1	0.2	0.3	0.5	0.9	1.0
1999.....	0.3	2.3	0.3	*	0.1	0.1	0.1	0.2	0.3	0.6	0.9	1.3
White, male												
2004.....	0.2	1.5	*	*	*	*	0.2	0.3	0.3	0.3	0.7	*
2003.....	0.3	2.0	*	*	*	*	0.1	0.3	0.4	0.5	0.8	*
2002.....	0.2	1.8	*	*	*	*	0.1	0.3	0.3	0.7	0.8	*
2001.....	0.2	1.4	*	*	*	*	0.1	0.3	0.4	0.6	0.9	*
2000.....	0.3	2.0	*	*	*	*	0.1	0.3	0.4	0.7	1.2	*
1999.....	0.3	3.0	0.3	*	*	*	0.1	0.3	0.4	0.6	0.9	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other inflammatory diseases of central nervous system (G04-G09)

All races 3/, both sexes

2004.....	0.2	*	*	0.1	0.1	0.1	0.1	0.2	0.4	0.6	1.1	1.1
2003.....	0.2	*	*	0.1	0.1	0.1	0.2	0.2	0.4	0.7	1.2	1.1
2002.....	0.3	*	*	0.1	0.1	0.1	0.1	0.2	0.4	0.6	1.2	1.2
2001.....	0.2	*	*	*	0.1	0.1	0.1	0.2	0.3	0.7	1.0	1.0
2000.....	0.2	*	*	0.1	0.1	0.1	0.1	0.2	0.3	0.6	1.1	1.0
1999.....	0.2	*	*	0.1	0.1	0.1	0.1	0.2	0.3	0.6	0.8	1.1

All races 3/, male

2004.....	0.3	*	*	*	*	*	0.1	0.2	0.5	0.8	1.5	1.5
2003.....	0.3	*	*	*	*	*	0.2	0.3	0.4	0.9	1.4	*
2002.....	0.3	*	*	*	0.1	0.1	0.2	0.3	0.4	0.9	1.6	*
2001.....	0.2	*	*	*	*	*	0.1	0.2	0.4	0.7	1.2	1.8
2000.....	0.2	*	*	*	*	*	0.2	0.2	0.4	0.8	1.6	*
1999.....	0.2	*	*	0.1	*	*	0.2	0.2	0.4	0.6	1.1	*

All races 3/, female

2004.....	0.2	*	*	*	*	0.1	0.1	0.2	0.3	0.4	0.8	1.0
2003.....	0.2	*	*	*	*	0.1	0.1	0.1	0.3	0.5	1.1	0.9
2002.....	0.2	*	*	*	*	0.1	0.1	0.2	0.3	0.5	1.0	1.1
2001.....	0.2	*	*	*	*	0.2	0.1	0.2	0.3	0.6	0.9	0.7
2000.....	0.2	*	*	*	*	0.1	0.1	0.1	0.3	0.5	0.8	0.9
1999.....	0.2	*	*	*	*	*	0.1	0.2	0.2	0.6	0.6	0.9

White, both sexes

2004.....	0.2	*	*	*	*	0.1	0.1	0.2	0.3	0.6	1.1	1.1
2003.....	0.2	*	*	*	*	0.1	0.1	0.2	0.3	0.7	1.2	1.1
2002.....	0.2	*	*	*	0.1	0.1	0.1	0.2	0.4	0.6	1.2	1.3
2001.....	0.2	*	*	*	*	0.1	0.1	0.2	0.3	0.6	0.9	1.0
2000.....	0.2	*	*	0.1	*	*	0.1	0.2	0.3	0.6	1.1	1.0
1999.....	0.2	*	*	0.1	0.1	0.1	0.1	0.2	0.3	0.6	0.8	1.1

White, male

2004.....	0.3	*	*	*	*	*	0.1	0.2	0.5	0.8	1.5	*
2003.....	0.3	*	*	*	*	*	0.2	0.3	0.4	0.8	1.5	*
2002.....	0.3	*	*	*	*	0.1	0.2	0.2	0.4	0.8	1.6	*
2001.....	0.2	*	*	*	*	*	0.1	0.2	0.4	0.7	1.1	1.8
2000.....	0.2	*	*	*	*	*	0.2	0.2	0.4	0.8	1.6	*
1999.....	0.2	*	*	*	*	*	0.1	0.2	0.4	0.6	1.2	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Parkinson's disease (G20-G21)												
All races 3/, both sexes												
2004.....	6.1	*	*	*	*	*	*	0.2	1.2	12.0	67.5	135.8
2003.....	6.2	*	*	*	*	*	*	0.2	1.3	12.7	67.8	138.2
2002.....	5.9	*	*	*	*	*	*	0.1	1.2	12.2	63.9	135.2
2001.....	5.8	*	*	*	*	*	*	0.1	1.2	11.7	64.6	134.2
2000.....	5.6	*	*	*	*	*	*	0.1	1.1	11.5	61.9	131.9
1999.....	5.2	*	*	*	*	*	*	0.1	1.0	11.0	58.2	124.4
All races 3/, male												
2004.....	7.2	*	*	*	*	*	*	0.2	1.6	17.3	101.7	220.0
2003.....	7.1	*	*	*	*	*	*	0.2	1.6	18.2	100.7	222.4
2002.....	6.8	*	*	*	*	*	*	0.2	1.5	17.9	94.7	220.1
2001.....	6.7	*	*	*	*	*	*	0.2	1.8	16.8	97.0	224.4
2000.....	6.4	*	*	*	*	*	*	0.2	1.5	16.1	92.8	222.3
1999.....	6.0	*	*	*	*	*	*	0.2	1.4	15.7	89.5	205.6
All races 3/, female												
2004.....	5.1	*	*	*	*	*	*	*	0.8	7.5	44.4	97.9
2003.....	5.3	*	*	*	*	*	*	0.1	0.9	8.1	45.8	101.0
2002.....	5.0	*	*	*	*	*	*	*	0.9	7.4	43.5	98.4
2001.....	4.9	*	*	*	*	*	*	*	0.7	7.5	43.3	96.4
2000.....	4.8	*	*	*	*	*	*	0.1	0.8	7.6	41.7	95.0
1999.....	4.5	*	*	*	*	*	*	*	0.7	7.1	37.8	91.7
White, both sexes												
2004.....	7.2	*	*	*	*	*	*	0.2	1.3	13.0	71.9	143.1
2003.....	7.2	*	*	*	*	*	*	0.2	1.3	13.6	72.2	146.0
2002.....	6.9	*	*	*	*	*	*	0.2	1.3	13.0	68.2	143.3
2001.....	6.8	*	*	*	*	*	*	0.1	1.2	12.6	68.8	141.8
2000.....	6.5	*	*	*	*	*	*	0.1	1.2	12.3	65.6	139.5
1999.....	6.1	*	*	*	*	*	*	0.1	1.1	11.7	61.5	131.2
White, male												
2004.....	8.3	*	*	*	*	*	*	0.2	1.7	18.6	107.9	231.3
2003.....	8.2	*	*	*	*	*	*	0.2	1.6	19.3	106.6	233.3
2002.....	7.9	*	*	*	*	*	*	0.2	1.6	19.0	100.5	232.7
2001.....	7.8	*	*	*	*	*	*	0.2	1.9	17.9	103.0	237.6
2000.....	7.4	*	*	*	*	*	*	0.2	1.6	17.2	97.5	235.3
1999.....	7.0	*	*	*	*	*	*	0.2	1.5	16.8	94.4	216.7

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Parkinson's disease (G20-G21)												
White, female												
2004.....	6.0	*	*	*	*	*	*	*	0.9	8.2	47.4	103.3
2003.....	6.2	*	*	*	*	*	*	0.2	1.0	8.7	48.8	107.4
2002.....	5.9	*	*	*	*	*	*	*	0.9	7.9	46.5	104.4
2001.....	5.8	*	*	*	*	*	*	*	0.7	8.1	46.2	101.7
2000.....	5.6	*	*	*	*	*	*	*	0.9	8.2	44.7	100.6
1999.....	5.2	*	*	*	*	*	*	*	0.6	7.5	40.0	96.8
Black, both sexes												
2004.....	1.6	*	*	*	*	*	*	*	0.7	5.4	29.2	61.2
2003.....	1.6	*	*	*	*	*	*	*	1.2	7.0	29.0	52.8
2002.....	1.5	*	*	*	*	*	*	*	*	6.8	25.4	51.7
2001.....	1.5	*	*	*	*	*	*	*	0.8	5.5	25.2	59.0
2000.....	1.5	*	*	*	*	*	*	*	*	5.7	27.4	51.6
1999.....	1.4	*	*	*	*	*	*	*	0.9	5.9	24.4	52.0
Black, male												
2004.....	1.9	*	*	*	*	*	*	*	*	8.5	43.4	106.3
2003.....	1.9	*	*	*	*	*	*	*	1.7	10.3	45.4	97.0
2002.....	1.7	*	*	*	*	*	*	*	*	11.1	39.6	89.0
2001.....	1.7	*	*	*	*	*	*	*	*	8.5	37.4	99.8
2000.....	1.8	*	*	*	*	*	*	*	*	8.8	47.8	86.5
1999.....	1.6	*	*	*	*	*	*	*	*	8.7	37.8	86.7
Black, female												
2004.....	1.4	*	*	*	*	*	*	*	*	3.2	21.1	43.3
2003.....	1.4	*	*	*	*	*	*	*	*	4.7	19.8	35.6
2002.....	1.2	*	*	*	*	*	*	*	*	3.6	17.4	37.4
2001.....	1.3	*	*	*	*	*	*	*	*	3.3	18.3	43.8
2000.....	1.2	*	*	*	*	*	*	*	*	3.5	15.9	38.6
1999.....	1.2	*	*	*	*	*	*	*	*	3.9	16.9	39.0

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Alzheimer's disease (G30)												
All races 3/, both sexes												
2004.....	22.5	*	*	*	*	*	*	0.2	1.9	19.7	168.7	818.8
2003.....	21.8	*	*	*	*	*	*	0.2	2.0	20.9	164.4	802.4
2002.....	20.4	*	*	*	*	*	*	0.1	1.9	19.7	158.1	752.3
2001.....	18.9	*	*	*	*	*	*	0.2	2.1	18.7	147.5	710.3
2000.....	17.6	*	*	*	*	*	*	0.2	2.0	18.7	139.6	667.7
1999.....	16.0	*	*	*	*	*	*	0.2	1.9	17.4	129.5	601.3
All races 3/, male												
2004.....	13.1	*	*	*	*	*	*	0.2	1.8	18.5	149.0	619.6
2003.....	12.8	*	*	*	*	*	*	0.2	1.8	20.2	147.6	606.5
2002.....	12.0	*	*	*	*	*	*	0.1	1.8	18.6	142.2	572.0
2001.....	11.3	*	*	*	*	*	*	0.2	1.9	17.7	136.3	556.1
2000.....	10.5	*	*	*	*	*	*	0.2	1.8	18.7	127.6	522.1
1999.....	9.8	*	*	*	*	*	*	0.2	1.9	17.3	122.6	485.7
All races 3/, female												
2004.....	31.5	*	*	*	*	*	*	0.3	1.9	20.8	181.9	908.4
2003.....	30.5	*	*	*	*	*	*	0.2	2.1	21.5	175.7	889.0
2002.....	28.5	*	*	*	*	*	*	0.1	2.0	20.6	168.7	830.5
2001.....	26.3	*	*	*	*	*	*	0.2	2.3	19.5	154.8	774.9
2000.....	24.5	*	*	*	*	*	*	0.2	2.2	18.6	147.4	727.0
1999.....	21.9	*	*	*	*	*	*	0.2	1.8	17.5	134.0	648.0
White, both sexes												
2004.....	25.6	*	*	*	*	*	*	0.3	2.0	20.5	174.1	846.8
2003.....	25.0	*	*	*	*	*	*	0.2	2.2	21.7	170.7	834.0
2002.....	23.5	*	*	*	*	*	*	0.1	2.1	20.7	164.8	781.8
2001.....	21.7	*	*	*	*	*	*	0.2	2.3	19.6	152.6	738.1
2000.....	20.2	*	*	*	*	*	*	0.2	2.2	19.5	144.8	695.3
1999.....	18.3	*	*	*	*	*	*	0.2	2.0	18.1	134.5	627.9
White, male												
2004.....	14.9	*	*	*	*	*	*	0.2	1.9	19.0	153.7	640.9
2003.....	14.6	*	*	*	*	*	*	0.2	1.9	20.8	152.4	630.3
2002.....	13.7	*	*	*	*	*	*	0.1	1.9	19.3	147.3	594.8
2001.....	12.8	*	*	*	*	*	*	0.2	2.0	18.5	140.5	577.2
2000.....	11.9	*	*	*	*	*	*	0.2	1.9	19.5	132.5	542.1
1999.....	11.2	*	*	*	*	*	*	0.2	2.1	17.9	127.4	506.9

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Alzheimer's disease (G30)												
White, female												
2004.....	36.1	*	*	*	*	*	*	0.3	2.1	21.7	188.0	939.6
2003.....	35.2	*	*	*	*	*	*	0.2	2.4	22.5	183.1	924.3
2002.....	33.0	*	*	*	*	*	*	0.1	2.2	21.8	176.5	863.0
2001.....	30.3	*	*	*	*	*	*	0.2	2.6	20.6	160.6	805.4
2000.....	28.2	*	*	*	*	*	*	0.2	2.4	19.5	152.9	757.6
1999.....	25.3	*	*	*	*	*	*	0.2	2.0	18.3	139.1	676.5
Black, both sexes												
2004.....	11.0	*	*	*	*	*	*	*	1.3	19.3	153.1	657.9
2003.....	9.7	*	*	*	*	*	*	*	1.0	19.2	138.1	583.8
2002.....	8.9	*	*	*	*	*	*	*	1.2	16.8	123.1	543.7
2001.....	8.4	*	*	*	*	*	*	*	1.1	15.2	124.7	506.7
2000.....	7.5	*	*	*	*	*	*	*	1.2	15.8	109.9	445.1
1999.....	6.5	*	*	*	*	*	*	*	1.2	15.1	98.2	372.4
Black, male												
2004.....	6.2	*	*	*	*	*	*	*	*	19.1	136.7	497.7
2003.....	6.0	*	*	*	*	*	*	*	*	20.3	136.2	459.8
2002.....	5.3	*	*	*	*	*	*	*	*	18.0	122.5	412.8
2001.....	5.1	*	*	*	*	*	*	*	*	16.1	121.5	414.1
2000.....	4.5	*	*	*	*	*	*	*	*	16.5	97.5	371.3
1999.....	4.0	*	*	*	*	*	*	*	*	16.5	89.8	317.5
Black, female												
2004.....	15.3	*	*	*	*	*	*	*	1.4	19.4	162.4	721.4
2003.....	13.2	*	*	*	*	*	*	*	*	18.4	139.2	631.9
2002.....	12.0	*	*	*	*	*	*	*	*	15.9	123.4	593.9
2001.....	11.3	*	*	*	*	*	*	*	*	14.6	126.4	541.0
2000.....	10.2	*	*	*	*	*	*	*	*	15.2	116.9	472.6
1999.....	8.7	*	*	*	*	*	*	*	*	14.2	102.9	392.9

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Multiple sclerosis (G35)												
All races 3/, both sexes												
2004.....	1.1	*	*	*	*	0.1	0.6	1.8	3.3	3.8	3.6	2.6
2003.....	1.1	*	*	*	*	0.1	0.6	1.9	3.1	3.6	3.6	2.2
2002.....	1.1	*	*	*	*	0.1	0.6	2.0	3.2	3.4	3.4	2.2
2001.....	1.1	*	*	*	*	0.1	0.7	1.9	3.1	3.4	3.5	2.1
2000.....	1.0	*	*	*	*	0.1	0.7	1.9	3.1	3.1	3.1	1.7
1999.....	1.0	*	*	*	*	0.1	0.7	1.9	2.6	2.9	3.0	2.1
All races 3/, male												
2004.....	0.8	*	*	*	*	*	0.4	1.3	2.4	2.9	2.6	1.7
2003.....	0.8	*	*	*	*	0.1	0.5	1.4	2.4	2.8	2.6	1.5
2002.....	0.8	*	*	*	*	*	0.4	1.3	2.6	2.5	2.6	1.9
2001.....	0.7	*	*	*	*	*	0.5	1.4	2.2	2.7	2.7	1.5
2000.....	0.7	*	*	*	*	*	0.5	1.4	2.3	2.4	2.3	1.8
1999.....	0.6	*	*	*	*	*	0.5	1.3	1.9	2.4	2.3	*
All races 3/, female												
2004.....	1.5	*	*	*	*	0.1	0.7	2.4	4.2	4.5	4.2	3.1
2003.....	1.4	*	*	*	*	0.1	0.8	2.5	3.8	4.3	4.2	2.5
2002.....	1.4	*	*	*	*	0.1	0.8	2.6	3.8	4.2	3.9	2.4
2001.....	1.4	*	*	*	*	0.2	0.9	2.5	3.9	3.9	4.0	2.3
2000.....	1.3	*	*	*	*	0.2	0.8	2.4	3.8	3.6	3.6	1.7
1999.....	1.2	*	*	*	*	0.2	0.9	2.5	3.2	3.3	3.5	2.4
White, both sexes												
2004.....	1.2	*	*	*	*	0.1	0.5	1.9	3.5	4.1	3.9	2.9
2003.....	1.2	*	*	*	*	0.1	0.6	2.0	3.4	3.8	3.8	2.3
2002.....	1.2	*	*	*	*	0.1	0.6	2.0	3.4	3.7	3.6	2.4
2001.....	1.2	*	*	*	*	0.1	0.6	2.0	3.3	3.6	3.8	2.2
2000.....	1.1	*	*	*	*	0.1	0.6	2.0	3.2	3.3	3.3	1.9
1999.....	1.0	*	*	*	*	0.1	0.7	2.0	2.8	3.0	3.3	2.2
White, male												
2004.....	0.8	*	*	*	*	*	0.4	1.3	2.6	3.1	2.7	1.8
2003.....	0.9	*	*	*	*	*	0.5	1.4	2.6	3.0	2.8	*
2002.....	0.8	*	*	*	*	*	0.4	1.4	2.8	2.7	2.8	2.1
2001.....	0.8	*	*	*	*	*	0.4	1.5	2.4	2.9	2.9	1.7
2000.....	0.8	*	*	*	*	*	0.5	1.5	2.3	2.6	2.5	2.0
1999.....	0.7	*	*	*	*	*	0.5	1.3	2.0	2.5	2.4	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Multiple sclerosis (G35)												
White, female												
2004.....	1.6	*	*	*	*	*	0.7	2.5	4.3	5.0	4.6	3.3
2003.....	1.5	*	*	*	*	*	0.7	2.5	4.1	4.5	4.4	2.7
2002.....	1.5	*	*	*	*	*	0.8	2.6	4.0	4.5	4.1	2.5
2001.....	1.5	*	*	*	*	0.2	0.8	2.5	4.1	4.2	4.3	2.5
2000.....	1.4	*	*	*	*	0.2	0.8	2.4	4.0	3.9	3.8	1.9
1999.....	1.4	*	*	*	*	0.1	0.8	2.6	3.5	3.4	3.9	2.5
Black, both sexes												
2004.....	0.9	*	*	*	*	*	0.9	1.9	3.7	2.5	*	*
2003.....	0.9	*	*	*	*	0.4	1.0	2.4	2.6	3.0	2.4	*
2002.....	0.9	*	*	*	*	*	1.1	2.6	2.8	2.4	*	*
2001.....	0.8	*	*	*	*	*	1.1	2.2	2.8	2.2	*	*
2000.....	0.8	*	*	*	*	*	1.2	2.2	3.3	2.0	*	*
1999.....	0.7	*	*	*	*	*	1.0	1.9	1.8	2.4	*	*
Black, male												
2004.....	0.5	*	*	*	*	*	*	1.2	1.7	2.8	*	*
2003.....	0.6	*	*	*	*	*	*	1.8	1.9	*	*	*
2002.....	0.5	*	*	*	*	*	*	1.5	1.7	*	*	*
2001.....	0.5	*	*	*	*	*	*	1.4	*	*	*	*
2000.....	0.5	*	*	*	*	*	1.0	1.3	2.8	*	*	*
1999.....	0.4	*	*	*	*	*	*	1.5	*	*	*	*
Black, female												
2004.....	1.2	*	*	*	*	*	1.2	2.6	5.3	2.3	*	*
2003.....	1.2	*	*	*	*	*	1.4	3.0	3.1	3.7	*	*
2002.....	1.3	*	*	*	*	*	1.5	3.5	3.8	3.3	*	*
2001.....	1.1	*	*	*	*	*	1.5	2.9	3.7	2.7	*	*
2000.....	1.1	*	*	*	*	*	1.4	3.1	3.6	2.7	*	*
1999.....	0.9	*	*	*	*	*	1.4	2.2	2.1	3.1	*	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Epilepsy (G40-G41)												
All races 3/, both sexes												
2004.....	0.5	*	0.2	0.1	0.2	0.5	0.5	0.6	0.6	0.6	1.1	1.5
2003.....	0.5	*	0.2	0.1	0.3	0.4	0.5	0.7	0.6	0.8	1.2	1.5
2002.....	0.5	*	0.1	0.1	0.3	0.4	0.6	0.6	0.5	0.8	1.3	1.8
2001.....	0.5	*	*	0.1	0.2	0.4	0.6	0.5	0.6	0.7	1.2	1.6
2000.....	0.5	*	0.2	0.1	0.2	0.4	0.5	0.6	0.6	0.8	1.2	2.0
1999.....	0.5	*	0.2	0.1	0.3	0.4	0.5	0.6	0.6	0.9	1.2	1.8
All races 3/, male												
2004.....	0.5	*	*	*	0.3	0.6	0.7	0.8	0.6	0.6	1.1	1.9
2003.....	0.5	*	*	0.1	0.3	0.5	0.6	0.8	0.7	0.7	1.4	1.5
2002.....	0.5	*	*	0.1	0.3	0.5	0.7	0.8	0.6	0.8	1.3	2.3
2001.....	0.5	*	*	0.1	0.3	0.5	0.7	0.6	0.7	0.8	0.9	2.0
2000.....	0.5	*	*	0.1	0.2	0.5	0.6	0.7	0.8	0.9	1.0	2.4
1999.....	0.5	*	0.3	0.1	0.3	0.5	0.7	0.7	0.6	1.0	1.1	2.3
All races 3/, female												
2004.....	0.4	*	*	*	0.2	0.4	0.4	0.4	0.6	0.7	1.2	1.4
2003.....	0.4	*	*	*	0.2	0.3	0.4	0.5	0.6	0.8	1.1	1.6
2002.....	0.4	*	*	*	0.3	0.3	0.4	0.4	0.5	0.8	1.3	1.5
2001.....	0.4	*	*	*	0.2	0.3	0.4	0.4	0.5	0.6	1.3	1.4
2000.....	0.4	*	*	*	0.2	0.3	0.4	0.5	0.5	0.8	1.4	1.9
1999.....	0.4	*	*	0.1	0.2	0.3	0.4	0.5	0.5	0.8	1.4	1.7
White, both sexes												
2004.....	0.4	*	0.2	0.1	0.2	0.4	0.5	0.5	0.5	0.6	1.0	1.4
2003.....	0.4	*	0.2	0.1	0.3	0.4	0.5	0.6	0.5	0.6	1.0	1.4
2002.....	0.4	*	*	0.1	0.2	0.4	0.5	0.5	0.5	0.7	1.2	1.6
2001.....	0.4	*	*	0.1	0.2	0.4	0.5	0.5	0.5	0.6	0.9	1.6
2000.....	0.4	*	0.2	0.1	0.2	0.4	0.5	0.5	0.5	0.7	1.0	1.7
1999.....	0.4	*	0.2	0.1	0.2	0.4	0.5	0.5	0.4	0.7	1.1	1.7
White, male												
2004.....	0.5	*	*	*	0.2	0.5	0.6	0.7	0.5	0.5	0.9	1.8
2003.....	0.5	*	*	0.1	0.3	0.5	0.6	0.7	0.5	0.5	1.0	*
2002.....	0.5	*	*	*	0.2	0.4	0.6	0.7	0.5	0.7	1.0	2.2
2001.....	0.5	*	*	0.1	0.3	0.4	0.6	0.6	0.6	0.7	0.7	2.0
2000.....	0.5	*	*	*	0.2	0.4	0.6	0.7	0.6	0.7	0.7	1.9
1999.....	0.5	*	0.3	*	0.3	0.5	0.6	0.6	0.4	0.8	0.9	2.3

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Epilepsy (G40-G41)												
White, female												
2004.....	0.4	*	*	*	0.2	0.3	0.3	0.3	0.5	0.6	1.1	1.2
2003.....	0.4	*	*	*	0.2	0.3	0.4	0.5	0.5	0.8	1.0	1.5
2002.....	0.4	*	*	*	0.2	0.3	0.4	0.4	0.5	0.7	1.3	1.3
2001.....	0.4	*	*	*	0.2	0.3	0.4	0.4	0.4	0.6	1.1	1.4
2000.....	0.4	*	*	*	0.2	0.3	0.4	0.4	0.4	0.7	1.3	1.6
1999.....	0.4	*	*	*	0.2	0.3	0.4	0.4	0.4	0.6	1.2	1.5
Black, both sexes												
2004.....	0.8	*	*	*	0.4	0.8	0.9	1.0	1.7	1.4	2.8	*
2003.....	0.9	*	*	*	0.3	0.7	0.9	1.4	1.8	2.2	3.5	*
2002.....	0.8	*	*	*	0.4	0.6	0.9	1.5	1.2	2.2	3.6	*
2001.....	0.8	*	*	*	0.4	0.7	1.2	0.9	1.6	1.6	3.8	*
2000.....	0.8	*	*	*	0.5	0.6	0.8	1.3	1.7	2.2	3.9	6.3
1999.....	0.9	*	*	*	0.4	0.8	1.1	1.4	1.8	2.7	3.4	*
Black, male												
2004.....	0.9	*	*	*	0.6	0.9	1.1	1.5	2.0	*	*	*
2003.....	1.0	*	*	*	*	0.8	1.1	2.1	2.0	3.2	*	*
2002.....	1.0	*	*	*	*	1.0	1.2	2.0	*	*	*	*
2001.....	0.9	*	*	*	*	1.0	1.7	1.2	1.9	*	*	*
2000.....	0.9	*	*	*	*	*	1.0	1.4	2.2	*	*	*
1999.....	1.1	*	*	*	*	1.0	1.6	1.8	2.5	*	*	*
Black, female												
2004.....	0.7	*	*	*	*	0.8	0.7	*	1.4	*	*	*
2003.....	0.7	*	*	*	*	*	0.8	0.9	1.6	*	*	*
2002.....	0.7	*	*	*	*	*	*	1.1	*	2.0	*	*
2001.....	0.6	*	*	*	*	*	0.7	*	*	*	4.0	*
2000.....	0.7	*	*	*	*	*	*	1.2	*	*	3.4	*
1999.....	0.8	*	*	*	*	*	*	1.0	*	2.6	3.4	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Transient cerebral ischemic attacks and related syndromes (G45)												
All races 3/, both sexes												
2004.....	0.1	*	*	*	*	*	*	*	*	*	0.4	2.1
2003.....	0.1	*	*	*	*	*	*	*	*	*	0.5	3.8
2002.....	0.1	*	*	*	*	*	*	*	*	0.1	0.5	2.9
2001.....	0.1	*	*	*	*	*	*	*	*	0.1	0.5	3.8
2000.....	0.1	*	*	*	*	*	*	*	*	0.1	0.7	3.9
1999.....	0.1	*	*	*	*	*	*	*	*	0.2	0.7	4.6
All races 3/, male												
2004.....	0.0	*	*	*	*	*	*	*	*	*	*	1.6
2003.....	0.1	*	*	*	*	*	*	*	*	*	0.4	2.8
2002.....	0.1	*	*	*	*	*	*	*	*	*	0.7	1.8
2001.....	0.1	*	*	*	*	*	*	*	*	*	0.6	4.2
2000.....	0.1	*	*	*	*	*	*	*	*	*	0.8	2.9
1999.....	0.1	*	*	*	*	*	*	*	*	*	0.9	3.8
All races 3/, female												
2004.....	0.1	*	*	*	*	*	*	*	*	*	0.5	2.4
2003.....	0.1	*	*	*	*	*	*	*	*	*	0.5	4.2
2002.....	0.1	*	*	*	*	*	*	*	*	*	0.5	3.4
2001.....	0.1	*	*	*	*	*	*	*	*	*	0.4	3.6
2000.....	0.1	*	*	*	*	*	*	*	*	*	0.7	4.2
1999.....	0.1	*	*	*	*	*	*	*	*	*	0.5	4.9
White, both sexes												
2004.....	0.1	*	*	*	*	*	*	*	*	*	0.4	2.2
2003.....	0.1	*	*	*	*	*	*	*	*	*	0.5	4.0
2002.....	0.1	*	*	*	*	*	*	*	*	0.1	0.6	3.0
2001.....	0.1	*	*	*	*	*	*	*	*	0.1	0.6	3.9
2000.....	0.1	*	*	*	*	*	*	*	*	0.1	0.7	4.0
1999.....	0.1	*	*	*	*	*	*	*	*	0.2	0.7	4.9
White, male												
2004.....	0.0	*	*	*	*	*	*	*	*	*	*	1.8
2003.....	0.1	*	*	*	*	*	*	*	*	*	*	2.9
2002.....	0.1	*	*	*	*	*	*	*	*	*	0.7	1.8
2001.....	0.1	*	*	*	*	*	*	*	*	*	0.7	4.3
2000.....	0.1	*	*	*	*	*	*	*	*	*	0.8	3.0
1999.....	0.1	*	*	*	*	*	*	*	*	*	0.9	3.9

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Infantile cerebral palsy (G80)												
All races 3/, both sexes												
2004.....	0.5	*	0.5	0.6	0.8	0.5	0.3	0.4	0.5	0.5	0.5	*
2003.....	0.5	*	0.4	0.6	0.7	0.5	0.2	0.4	0.4	0.4	0.5	*
2002.....	0.4	*	0.6	0.5	0.7	0.4	0.2	0.3	0.4	0.4	0.5	*
2001.....	0.4	*	0.5	0.5	0.7	0.3	0.2	0.3	0.4	0.4	0.5	*
2000.....	0.4	*	0.4	0.6	0.6	0.3	0.2	0.3	0.3	0.5	0.5	*
1999.....	0.4	*	0.5	0.5	0.5	0.3	0.2	0.2	0.3	0.4	0.4	0.5
All races 3/, male												
2004.....	0.6	*	0.4	0.6	0.9	0.5	0.3	0.4	0.6	0.7	0.7	*
2003.....	0.5	*	0.5	0.7	0.8	0.5	0.3	0.5	0.5	0.5	0.7	*
2002.....	0.5	*	0.7	0.5	0.8	0.4	0.3	0.3	0.5	0.5	0.6	*
2001.....	0.5	*	0.5	0.5	0.8	0.4	0.3	0.3	0.4	0.6	0.7	*
2000.....	0.4	*	0.3	0.6	0.7	0.4	0.2	0.3	0.3	0.7	0.5	*
1999.....	0.4	*	0.6	0.5	0.6	0.4	0.2	0.3	0.4	0.5	0.5	*
All races 3/, female												
2004.....	0.4	*	0.5	0.5	0.6	0.4	0.3	0.4	0.4	0.4	0.4	*
2003.....	0.4	*	0.4	0.5	0.6	0.4	0.2	0.3	0.4	0.3	0.4	*
2002.....	0.4	*	0.5	0.5	0.5	0.3	0.2	0.3	0.3	0.4	0.4	*
2001.....	0.3	*	0.4	0.5	0.5	0.3	0.2	0.2	0.4	0.3	0.4	*
2000.....	0.3	*	0.4	0.5	0.5	0.3	0.2	0.2	0.3	0.3	0.5	*
1999.....	0.3	*	0.4	0.4	0.4	0.3	0.2	0.2	0.2	0.3	0.4	*
White, both sexes												
2004.....	0.5	*	0.3	0.5	0.7	0.5	0.3	0.4	0.5	0.6	0.5	*
2003.....	0.4	*	0.3	0.5	0.6	0.4	0.2	0.4	0.5	0.4	0.5	*
2002.....	0.4	*	0.5	0.4	0.6	0.3	0.2	0.3	0.4	0.5	0.5	*
2001.....	0.4	*	0.4	0.4	0.6	0.3	0.2	0.3	0.4	0.5	0.5	*
2000.....	0.4	*	0.3	0.5	0.6	0.3	0.2	0.3	0.3	0.5	0.5	*
1999.....	0.3	*	0.5	0.4	0.4	0.4	0.2	0.2	0.3	0.4	0.5	0.5
White, male												
2004.....	0.5	*	*	0.6	0.8	0.5	0.3	0.4	0.6	0.7	0.7	*
2003.....	0.5	*	0.4	0.6	0.7	0.4	0.3	0.4	0.5	0.5	0.8	*
2002.....	0.4	*	0.6	0.5	0.7	0.3	0.3	0.3	0.5	0.6	0.7	*
2001.....	0.5	*	0.5	0.4	0.7	0.3	0.3	0.3	0.4	0.6	0.7	*
2000.....	0.4	*	*	0.6	0.7	0.4	0.2	0.3	0.4	0.7	0.5	*
1999.....	0.4	*	0.6	0.4	0.5	0.4	0.2	0.3	0.4	0.5	0.6	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other diseases of nervous system (G10-G12,G23-G25,G31,G36-G37,G43-G44,G47-G72,G81-G98)												
All races 3/, both sexes												
2004.....	6.9	6.7	1.1	0.5	0.9	1.1	2.4	5.5	11.9	23.5	39.2	51.4
2003.....	7.1	7.3	0.9	0.5	1.0	1.1	2.4	5.4	12.1	24.5	40.4	56.0
2002.....	6.9	6.9	1.1	0.5	0.9	1.2	2.5	5.4	11.7	24.1	40.5	53.8
2001.....	6.7	6.1	1.2	0.5	1.0	1.1	2.3	5.2	11.2	24.0	39.4	51.9
2000.....	6.7	7.1	1.2	0.5	0.9	1.1	2.4	5.5	11.4	23.3	39.4	55.4
1999.....	6.5	7.7	1.2	0.6	1.1	1.0	2.1	5.0	10.9	23.3	39.3	53.6
All races 3/, male												
2004.....	7.3	7.8	1.1	0.6	1.3	1.3	2.9	6.6	13.8	27.6	47.2	62.4
2003.....	7.5	8.3	0.9	0.6	1.5	1.4	3.0	6.6	14.0	27.8	49.8	70.0
2002.....	7.3	8.4	1.3	0.6	1.3	1.5	3.0	6.4	13.4	28.2	49.3	66.1
2001.....	7.1	7.1	1.2	0.6	1.5	1.2	2.9	5.9	12.5	28.5	49.0	69.7
2000.....	7.2	8.1	1.3	0.5	1.3	1.4	2.8	6.5	13.2	27.9	49.1	72.0
1999.....	7.0	8.6	1.2	0.6	1.6	1.2	2.6	5.8	12.6	27.4	49.7	70.3
All races 3/, female												
2004.....	6.5	5.6	1.1	0.4	0.4	0.8	2.0	4.5	10.2	20.0	33.9	46.5
2003.....	6.6	6.2	1.0	0.4	0.5	0.9	1.8	4.3	10.3	21.8	34.1	49.9
2002.....	6.5	5.4	0.9	0.3	0.5	0.8	1.9	4.4	10.1	20.6	34.6	48.5
2001.....	6.3	5.0	1.2	0.5	0.5	0.9	1.8	4.4	10.0	20.2	33.1	44.5
2000.....	6.3	6.0	1.1	0.5	0.5	0.8	1.9	4.6	9.7	19.5	33.0	48.7
1999.....	6.1	6.8	1.2	0.5	0.5	0.7	1.7	4.3	9.2	20.0	32.6	46.8
White, both sexes												
2004.....	7.3	6.8	1.0	0.5	0.9	1.0	2.5	5.3	11.8	23.7	39.8	52.3
2003.....	7.5	6.8	0.9	0.5	1.1	1.1	2.3	5.3	12.0	24.9	41.1	56.9
2002.....	7.3	6.9	1.0	0.5	0.9	1.2	2.4	5.3	11.7	24.3	41.1	54.7
2001.....	7.0	5.8	1.1	0.5	1.0	1.0	2.3	5.0	10.9	24.2	39.8	52.5
2000.....	7.1	6.9	1.1	0.5	1.0	1.1	2.2	5.4	11.3	23.4	40.2	56.2
1999.....	6.9	7.5	1.1	0.5	1.1	1.0	2.1	4.8	10.7	23.4	40.0	54.2
White, male												
2004.....	7.8	8.0	1.0	0.6	1.3	1.3	3.0	6.5	13.6	28.1	48.0	64.7
2003.....	8.0	8.2	0.7	0.6	1.6	1.4	3.0	6.5	13.9	28.3	50.8	72.0
2002.....	7.8	8.5	1.1	0.6	1.3	1.6	3.0	6.3	13.4	28.3	50.4	67.5
2001.....	7.5	6.6	1.1	0.6	1.6	1.1	2.8	5.7	12.3	28.8	49.7	70.1
2000.....	7.6	7.4	1.2	0.5	1.4	1.4	2.7	6.5	13.0	28.1	50.1	73.1
1999.....	7.4	8.4	1.1	0.5	1.7	1.2	2.5	5.7	12.5	27.6	50.9	71.5

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

All other diseases of nervous system (G10-G12,G23-G25,G31,G36-G37,G43-G44,G47-G72,G81-G98)

White, female

2004.....	6.8	5.6	1.0	0.4	0.4	0.7	1.9	4.2	10.1	19.9	34.2	46.8
2003.....	7.0	5.2	1.0	0.4	0.5	0.9	1.7	4.0	10.1	22.0	34.5	50.3
2002.....	6.9	5.2	0.8	0.3	0.4	0.7	1.8	4.3	10.0	20.9	34.9	49.2
2001.....	6.6	4.9	1.0	0.4	0.4	0.8	1.7	4.3	9.5	20.4	33.2	45.1
2000.....	6.6	6.3	1.0	0.4	0.5	0.7	1.8	4.3	9.7	19.4	33.6	49.3
1999.....	6.4	6.5	1.1	0.5	0.4	0.7	1.6	3.9	8.9	19.9	32.8	47.3

Black, both sexes

2004.....	6.2	6.8	1.7	0.5	0.9	1.8	3.2	8.2	16.2	26.5	41.5	49.6
2003.....	6.2	9.9	1.5	0.7	0.8	1.5	3.7	8.0	16.0	27.1	38.2	52.5
2002.....	6.0	7.9	1.7	0.5	1.0	1.6	3.5	7.7	15.2	27.1	39.6	50.5
2001.....	6.1	8.0	2.0	0.6	1.2	1.9	3.2	7.9	16.0	27.1	41.3	48.8
2000.....	5.9	7.5	1.6	0.5	1.0	1.7	3.6	8.5	14.4	27.0	37.3	51.3
1999.....	5.7	9.1	1.4	0.8	1.1	1.3	3.0	8.4	14.6	26.5	36.7	50.7

Black, male

2004.....	6.2	7.4	1.9	*	1.2	1.7	3.3	9.8	19.1	29.5	48.2	50.7
2003.....	6.3	9.8	1.8	0.8	1.2	1.8	4.4	9.2	18.3	30.7	44.6	50.0
2002.....	6.3	7.8	2.0	0.6	1.4	1.9	4.3	9.1	17.7	33.8	45.0	57.9
2001.....	6.3	10.2	1.9	*	1.6	2.4	4.0	9.2	17.0	31.4	50.9	62.4
2000.....	6.2	10.9	1.6	*	1.0	2.0	4.3	9.1	17.8	31.9	49.1	63.4
1999.....	5.7	10.4	*	1.0	1.6	1.5	3.5	9.1	15.6	28.7	43.7	65.6

Black, female

2004.....	6.2	6.2	1.6	*	0.7	1.8	3.1	6.9	13.8	24.4	37.8	49.2
2003.....	6.0	9.9	*	0.6	*	1.3	3.0	7.1	14.0	24.6	34.6	53.5
2002.....	5.7	7.9	*	*	0.7	1.3	2.7	6.5	13.3	22.3	36.5	47.6
2001.....	5.9	*	2.0	0.8	0.9	1.5	2.5	6.7	15.2	24.1	35.9	43.8
2000.....	5.5	*	1.6	*	0.9	1.4	3.0	8.0	11.7	23.6	30.7	46.8
1999.....	5.7	7.7	*	0.7	*	1.1	2.6	7.8	13.8	25.0	32.9	45.1

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
IX. Diseases of the circulatory system (I00-I99)												
All races 3/, both sexes												
2004.....	294.9	14.5	1.5	0.8	3.3	10.3	37.3	112.2	270.3	701.3	2,044.6	6,639.9
2003.....	311.9	14.8	1.5	0.9	3.6	10.6	38.9	114.8	287.0	752.4	2,179.9	7,162.1
2002.....	320.2	16.5	1.5	0.9	3.2	10.3	38.4	115.6	297.1	792.2	2,266.8	7,414.0
2001.....	325.5	15.4	1.9	0.9	3.3	10.5	37.7	114.6	303.4	815.0	2,331.6	7,699.8
2000.....	334.6	17.4	1.5	0.9	3.4	9.8	37.5	117.0	321.3	853.6	2,405.6	8,055.8
1999.....	342.0	17.6	1.5	0.9	3.5	9.9	38.3	117.3	329.2	891.0	2,484.0	8,220.5
All races 3/, male												
2004.....	282.7	15.6	1.5	0.8	4.1	13.1	49.8	158.3	374.1	907.8	2,472.0	6,837.7
2003.....	297.7	16.2	1.5	1.0	4.3	13.2	51.8	162.5	395.7	980.4	2,644.5	7,348.4
2002.....	304.7	17.3	1.5	0.9	4.2	13.1	51.5	163.7	409.5	1,031.7	2,740.6	7,659.3
2001.....	307.6	15.8	2.0	0.9	4.0	13.0	50.6	161.8	416.5	1,062.5	2,833.1	7,958.0
2000.....	317.2	18.3	1.7	1.0	4.0	12.1	50.3	166.7	443.8	1,118.8	2,936.2	8,470.3
1999.....	324.2	18.5	1.6	1.0	4.1	12.3	51.9	166.9	454.0	1,172.7	3,033.6	8,624.0
All races 3/, female												
2004.....	306.7	13.5	1.5	0.8	2.4	7.5	24.8	67.7	174.0	527.9	1,756.9	6,551.0
2003.....	325.8	13.3	1.5	0.7	2.9	7.9	26.1	68.7	186.1	561.7	1,869.4	7,079.7
2002.....	335.2	15.7	1.4	0.8	2.3	7.5	25.5	69.2	193.1	592.8	1,952.2	7,307.6
2001.....	342.8	15.0	1.8	0.9	2.5	7.9	24.9	69.1	198.9	609.9	2,002.1	7,591.7
2000.....	351.3	16.5	1.4	0.8	2.9	7.5	24.9	69.2	208.3	635.2	2,059.6	7,887.0
1999.....	359.1	16.6	1.4	0.8	2.8	7.4	24.9	69.5	214.3	660.4	2,127.0	8,057.9
White, both sexes												
2004.....	313.3	12.6	1.2	0.7	2.8	8.4	32.6	98.3	246.7	668.6	2,025.0	6,728.0
2003.....	331.7	11.9	1.2	0.7	3.0	8.7	33.5	100.2	261.7	716.1	2,160.0	7,256.7
2002.....	341.0	14.1	1.3	0.8	2.8	8.4	33.5	101.7	271.9	758.7	2,244.3	7,507.0
2001.....	346.4	12.7	1.6	0.8	2.7	8.8	32.2	100.6	277.3	778.4	2,308.0	7,796.7
2000.....	356.4	15.3	1.4	0.8	2.8	8.1	32.2	103.2	294.6	818.2	2,379.9	8,159.4
1999.....	364.2	15.2	1.3	0.8	2.9	8.3	32.6	103.6	301.8	854.5	2,458.8	8,322.3
White, male												
2004.....	298.1	13.7	1.3	0.7	3.5	11.0	45.0	142.7	345.6	873.8	2,461.9	6,960.5
2003.....	314.0	12.5	1.1	0.9	3.6	11.1	46.3	146.0	367.0	942.1	2,630.9	7,485.4
2002.....	322.2	15.2	1.5	0.8	3.6	10.8	46.3	148.3	380.4	998.4	2,730.2	7,786.5
2001.....	324.9	13.7	1.6	0.8	3.4	11.1	44.7	146.4	387.5	1,025.3	2,819.1	8,082.3
2000.....	335.6	16.1	1.6	0.9	3.4	10.3	44.9	151.6	413.8	1,085.1	2,921.9	8,607.0
1999.....	342.5	15.9	1.5	0.9	3.4	10.5	45.7	151.4	422.7	1,137.0	3,022.0	8,738.2

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
IX. Diseases of the circulatory system (I00-I99)												
White, female												
2004.....	328.3	11.4	1.1	0.7	2.0	5.7	20.0	54.5	153.0	493.0	1,726.9	6,623.2
2003.....	348.9	11.2	1.4	0.6	2.4	6.3	20.5	55.1	162.2	523.4	1,841.6	7,155.4
2002.....	359.5	13.0	1.2	0.7	1.9	5.8	20.6	55.8	169.6	555.3	1,918.2	7,385.6
2001.....	367.4	11.7	1.5	0.8	2.1	6.3	19.5	55.4	173.7	570.0	1,968.7	7,677.1
2000.....	376.6	14.4	1.1	0.6	2.3	5.8	19.5	55.6	182.7	594.3	2,023.3	7,977.5
1999.....	385.2	14.4	1.2	0.7	2.4	5.9	19.4	56.5	188.5	618.9	2,089.6	8,155.3
Black, both sexes												
2004.....	262.4	23.0	2.9	1.5	6.0	23.6	76.2	234.0	529.7	1,129.0	2,586.3	6,397.4
2003.....	275.3	27.1	2.8	1.4	7.2	23.7	80.7	241.4	560.5	1,214.6	2,728.8	6,895.6
2002.....	278.2	28.6	2.0	1.4	6.1	24.3	77.7	240.0	573.2	1,233.0	2,840.6	7,111.7
2001.....	282.4	29.3	3.6	1.3	6.8	23.4	79.8	240.5	582.8	1,284.1	2,906.2	7,346.8
2000.....	287.0	28.6	2.6	1.5	7.0	22.4	78.9	244.0	604.7	1,325.0	2,983.2	7,548.9
1999.....	291.1	29.9	2.5	1.4	7.0	21.2	83.0	244.3	617.5	1,374.7	3,038.5	7,610.9
Black, male												
2004.....	259.1	24.6	2.8	1.4	7.9	28.8	94.5	309.1	721.6	1,439.5	3,048.5	6,222.9
2003.....	270.6	32.7	3.3	1.5	8.8	28.9	99.4	320.5	747.6	1,550.8	3,283.2	6,580.1
2002.....	270.5	28.8	2.0	1.5	7.5	30.0	96.5	316.1	768.1	1,563.8	3,305.9	6,961.1
2001.....	274.3	29.0	3.8	1.2	8.2	28.6	99.1	314.9	767.4	1,629.3	3,451.2	7,393.4
2000.....	277.6	29.7	2.2	1.4	7.7	26.3	97.5	320.9	801.9	1,649.0	3,515.1	7,717.4
1999.....	284.2	31.6	2.6	1.5	8.6	25.9	104.5	326.1	825.3	1,731.9	3,543.4	7,943.3
Black, female												
2004.....	265.3	21.3	3.0	1.6	4.2	18.8	59.9	169.4	375.4	907.4	2,325.1	6,466.6
2003.....	279.5	21.3	2.4	1.3	5.6	19.0	64.1	173.3	410.1	975.4	2,415.4	7,018.2
2002.....	285.2	28.5	2.1	1.3	4.6	19.1	61.0	174.8	416.5	998.1	2,577.2	7,169.4
2001.....	289.9	29.6	3.4	1.4	5.4	18.6	62.7	176.5	434.1	1,040.5	2,600.1	7,329.5
2000.....	295.6	27.4	2.9	1.5	6.3	18.9	62.4	177.9	445.8	1,097.1	2,683.7	7,486.2
1999.....	297.4	28.2	2.4	1.2	5.5	16.9	64.0	174.3	450.8	1,122.8	2,755.9	7,486.6

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
All races 3/, both sexes												
2004.....	1.1	*	*	*	*	0.1	0.2	0.5	1.1	3.3	8.8	17.1
2003.....	1.2	*	*	*	*	0.1	0.2	0.5	1.3	3.8	9.6	20.5
2002.....	1.2	*	*	*	*	0.1	0.2	0.6	1.6	3.8	9.6	17.8
2001.....	1.2	*	*	*	*	0.1	0.3	0.6	1.4	4.1	9.4	17.7
2000.....	1.3	*	*	*	*	0.1	0.3	0.7	1.7	4.5	9.5	16.8
1999.....	1.3	*	*	*	*	0.1	0.3	0.7	1.7	4.6	10.0	18.0
All races 3/, male												
2004.....	0.7	*	*	*	*	*	0.2	0.4	0.8	2.4	7.3	11.5
2003.....	0.8	*	*	*	*	*	0.2	0.4	1.1	2.9	7.5	16.2
2002.....	0.8	*	*	*	*	*	0.2	0.5	1.2	3.0	6.7	11.8
2001.....	0.7	*	*	*	*	*	0.2	0.5	1.1	2.8	6.6	12.8
2000.....	0.8	*	*	*	*	*	0.3	0.6	1.4	2.9	6.6	12.2
1999.....	0.8	*	*	*	*	*	0.2	0.5	1.2	3.5	6.8	10.3
All races 3/, female												
2004.....	1.5	*	*	*	*	*	0.3	0.6	1.4	4.1	9.9	19.5
2003.....	1.7	*	*	*	*	*	0.2	0.7	1.5	4.5	11.0	22.3
2002.....	1.7	*	*	*	*	*	0.2	0.7	2.0	4.6	11.6	20.4
2001.....	1.7	*	*	*	*	*	0.3	0.7	1.7	5.1	11.3	19.8
2000.....	1.8	*	*	*	*	0.1	0.4	0.8	2.0	5.8	11.4	18.7
1999.....	1.9	*	*	*	*	0.1	0.3	0.8	2.2	5.6	12.2	21.1
White, both sexes												
2004.....	1.2	*	*	*	*	*	0.2	0.4	1.0	3.4	9.3	18.2
2003.....	1.4	*	*	*	*	*	0.2	0.4	1.2	3.8	10.2	21.8
2002.....	1.4	*	*	*	*	0.1	0.2	0.5	1.5	4.0	10.1	18.9
2001.....	1.3	*	*	*	*	*	0.2	0.5	1.4	4.1	10.0	18.7
2000.....	1.4	*	*	*	*	0.1	0.3	0.6	1.7	4.6	10.1	17.6
1999.....	1.4	*	*	*	*	0.1	0.2	0.6	1.7	4.8	10.5	18.9
White, male												
2004.....	0.8	*	*	*	*	*	0.1	0.4	0.7	2.5	7.8	12.4
2003.....	0.9	*	*	*	*	*	0.1	0.3	0.9	3.0	8.1	17.3
2002.....	0.8	*	*	*	*	*	0.2	0.4	1.1	3.0	7.1	12.6
2001.....	0.8	*	*	*	*	*	0.2	0.4	1.1	2.8	6.9	13.3
2000.....	0.9	*	*	*	*	*	0.2	0.6	1.5	2.9	7.0	12.7
1999.....	0.8	*	*	*	*	*	0.2	0.5	1.1	3.7	7.0	11.1

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)

White, female

2004.....	1.7	*	*	*	*	*	0.2	0.5	1.4	4.2	10.4	20.8
2003.....	1.8	*	*	*	*	*	0.2	0.5	1.4	4.6	11.6	23.8
2002.....	1.9	*	*	*	*	*	0.2	0.6	1.9	4.8	12.1	21.6
2001.....	1.8	*	*	*	*	*	0.2	0.6	1.6	5.2	12.0	21.0
2000.....	1.9	*	*	*	*	*	0.3	0.7	1.9	6.0	12.2	19.6
1999.....	2.0	*	*	*	*	*	0.3	0.7	2.2	5.7	12.8	22.0

Black, both sexes

2004.....	0.6	*	*	*	*	*	0.6	1.0	1.6	2.7	4.0	5.9
2003.....	0.8	*	*	*	*	*	0.5	1.3	2.6	3.9	4.1	7.6
2002.....	0.7	*	*	*	*	*	0.3	0.9	2.4	2.9	4.4	*
2001.....	0.8	*	*	*	*	*	0.7	1.3	2.1	4.1	4.2	6.8
2000.....	0.7	*	*	*	*	*	0.6	1.0	2.0	3.5	3.9	8.4
1999.....	0.8	*	*	*	*	*	0.6	1.2	2.2	3.4	4.4	8.3

Black, male

2004.....	0.5	*	*	*	*	*	*	0.9	*	*	*	*
2003.....	0.6	*	*	*	*	*	*	0.9	2.7	3.0	*	*
2002.....	0.5	*	*	*	*	*	*	*	1.9	3.0	*	*
2001.....	0.5	*	*	*	*	*	*	1.0	*	*	*	*
2000.....	0.5	*	*	*	*	*	*	*	*	*	*	*
1999.....	0.5	*	*	*	*	*	*	*	*	*	*	*

Black, female

2004.....	0.8	*	*	*	*	*	*	1.1	1.7	3.3	5.0	*
2003.....	1.0	*	*	*	*	*	*	1.7	2.5	4.5	5.1	8.1
2002.....	0.8	*	*	*	*	*	*	1.1	2.8	2.8	5.0	*
2001.....	1.1	*	*	*	*	*	0.9	1.7	2.6	5.3	5.3	*
2000.....	1.0	*	*	*	*	*	0.8	1.3	2.7	4.3	4.8	*
1999.....	1.0	*	*	*	*	*	0.7	1.4	2.6	4.1	4.5	11.0

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Chronic rheumatic heart diseases (I05-I09)												
All races 3/, both sexes												
2004.....	1.1	*	*	*	*	0.1	0.2	0.5	1.1	3.3	8.8	17.0
2003.....	1.2	*	*	*	*	0.1	0.2	0.5	1.3	3.7	9.5	20.3
2002.....	1.2	*	*	*	*	0.1	0.2	0.6	1.6	3.8	9.5	17.5
2001.....	1.2	*	*	*	*	0.1	0.3	0.6	1.4	4.0	9.3	17.2
2000.....	1.3	*	*	*	*	0.1	0.3	0.7	1.7	4.4	9.5	16.7
1999.....	1.3	*	*	*	*	0.1	0.3	0.7	1.7	4.6	10.0	17.9
All races 3/, male												
2004.....	0.7	*	*	*	*	*	0.2	0.4	0.7	2.3	7.2	11.4
2003.....	0.8	*	*	*	*	*	0.2	0.4	1.1	2.8	7.4	16.1
2002.....	0.7	*	*	*	*	*	0.2	0.5	1.2	2.9	6.6	11.4
2001.....	0.7	*	*	*	*	*	0.2	0.4	1.1	2.7	6.5	12.3
2000.....	0.8	*	*	*	*	*	0.3	0.6	1.4	2.8	6.5	12.1
1999.....	0.8	*	*	*	*	*	0.2	0.5	1.1	3.5	6.7	10.3
All races 3/, female												
2004.....	1.5	*	*	*	*	*	0.3	0.6	1.4	4.0	9.8	19.5
2003.....	1.7	*	*	*	*	*	0.2	0.7	1.5	4.5	10.9	22.2
2002.....	1.7	*	*	*	*	*	0.2	0.7	2.0	4.6	11.4	20.2
2001.....	1.7	*	*	*	*	*	0.3	0.7	1.7	5.0	11.2	19.3
2000.....	1.7	*	*	*	*	0.1	0.3	0.7	2.0	5.7	11.4	18.5
1999.....	1.8	*	*	*	*	0.1	0.3	0.8	2.2	5.5	12.1	21.0
White, both sexes												
2004.....	1.2	*	*	*	*	*	0.2	0.4	1.0	3.4	9.2	18.1
2003.....	1.4	*	*	*	*	*	0.2	0.4	1.1	3.8	10.1	21.7
2002.....	1.3	*	*	*	*	0.1	0.2	0.5	1.5	3.9	10.0	18.6
2001.....	1.3	*	*	*	*	*	0.2	0.5	1.3	4.0	9.8	18.3
2000.....	1.4	*	*	*	*	0.1	0.3	0.6	1.7	4.5	10.1	17.5
1999.....	1.4	*	*	*	*	0.1	0.2	0.6	1.7	4.8	10.5	18.9
White, male												
2004.....	0.8	*	*	*	*	*	0.1	0.4	0.7	2.4	7.7	12.3
2003.....	0.9	*	*	*	*	*	0.1	0.3	0.9	2.9	8.0	17.2
2002.....	0.8	*	*	*	*	*	0.2	0.4	1.1	3.0	7.0	12.2
2001.....	0.8	*	*	*	*	*	0.2	0.4	1.1	2.7	6.8	12.9
2000.....	0.8	*	*	*	*	*	0.2	0.5	1.4	2.8	6.9	12.6
1999.....	0.8	*	*	*	*	*	0.2	0.5	1.1	3.6	6.9	11.1

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Chronic rheumatic heart diseases (I05-I09)												
White, female												
2004.....	1.7	*	*	*	*	*	0.2	0.5	1.3	4.1	10.3	20.7
2003.....	1.8	*	*	*	*	*	0.2	0.5	1.4	4.6	11.5	23.7
2002.....	1.9	*	*	*	*	*	0.2	0.6	1.9	4.7	12.0	21.4
2001.....	1.8	*	*	*	*	*	0.2	0.6	1.6	5.1	11.8	20.5
2000.....	1.9	*	*	*	*	*	0.3	0.7	1.9	6.0	12.2	19.5
1999.....	2.0	*	*	*	*	*	0.3	0.7	2.2	5.7	12.8	22.0
Black, both sexes												
2004.....	0.6	*	*	*	*	*	0.6	1.0	1.5	2.6	3.9	5.9
2003.....	0.8	*	*	*	*	*	0.5	1.3	2.5	3.8	3.8	7.6
2002.....	0.7	*	*	*	*	*	*	0.9	2.4	2.9	4.1	*
2001.....	0.8	*	*	*	*	*	0.7	1.3	2.1	4.0	4.2	*
2000.....	0.7	*	*	*	*	*	0.6	1.0	2.0	3.4	3.9	8.1
1999.....	0.7	*	*	*	*	*	0.5	1.2	2.1	3.4	4.3	7.7
Black, male												
2004.....	0.4	*	*	*	*	*	*	0.9	*	*	*	*
2003.....	0.6	*	*	*	*	*	*	*	2.6	3.0	*	*
2002.....	0.5	*	*	*	*	*	*	*	1.9	3.0	*	*
2001.....	0.5	*	*	*	*	*	*	1.0	*	*	*	*
2000.....	0.4	*	*	*	*	*	*	*	*	*	*	*
1999.....	0.5	*	*	*	*	*	*	*	*	*	*	*
Black, female												
2004.....	0.8	*	*	*	*	*	*	1.1	1.6	3.2	4.9	*
2003.....	1.0	*	*	*	*	*	*	1.6	2.5	4.4	4.6	8.1
2002.....	0.8	*	*	*	*	*	*	1.1	2.8	2.8	4.7	*
2001.....	1.1	*	*	*	*	*	0.9	1.7	2.6	5.0	5.3	*
2000.....	0.9	*	*	*	*	*	0.8	1.3	2.7	4.2	4.8	*
1999.....	1.0	*	*	*	*	*	*	1.4	2.6	4.1	4.5	10.1

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Rheumatic mitral valve diseases (I05)												
All races 3/, both sexes												
2004.....	0.5	*	*	*	*	*	0.1	0.3	0.6	1.6	3.9	6.6
2003.....	0.6	*	*	*	*	*	0.1	0.3	0.7	2.0	4.7	7.6
2002.....	0.6	*	*	*	*	0.1	0.1	0.4	1.0	2.0	5.0	7.4
2001.....	0.7	*	*	*	*	0.1	0.2	0.4	0.8	2.3	5.1	8.5
2000.....	0.7	*	*	*	*	0.1	0.2	0.4	1.0	2.7	5.4	8.5
1999.....	0.7	*	*	*	*	*	0.1	0.4	1.0	2.6	5.7	9.0
All races 3/, male												
2004.....	0.3	*	*	*	*	*	0.1	0.2	0.4	1.1	3.0	4.6
2003.....	0.4	*	*	*	*	*	*	0.2	0.7	1.5	3.5	5.9
2002.....	0.4	*	*	*	*	*	0.1	0.3	0.7	1.6	3.0	4.2
2001.....	0.4	*	*	*	*	*	0.1	0.3	0.5	1.6	3.2	6.2
2000.....	0.4	*	*	*	*	*	0.2	0.3	0.9	1.7	3.8	6.1
1999.....	0.4	*	*	*	*	*	0.1	0.3	0.6	1.9	3.7	4.9
All races 3/, female												
2004.....	0.7	*	*	*	*	*	0.2	0.3	0.8	2.0	4.5	7.5
2003.....	0.8	*	*	*	*	*	0.1	0.4	0.7	2.4	5.5	8.4
2002.....	0.9	*	*	*	*	*	0.1	0.4	1.2	2.3	6.2	8.7
2001.....	0.9	*	*	*	*	*	0.2	0.5	1.1	2.9	6.3	9.4
2000.....	1.0	*	*	*	*	*	0.2	0.5	1.2	3.5	6.4	9.5
1999.....	1.0	*	*	*	*	*	0.2	0.5	1.3	3.3	7.0	10.7
White, both sexes												
2004.....	0.5	*	*	*	*	*	0.1	0.2	0.5	1.7	4.1	7.1
2003.....	0.6	*	*	*	*	*	0.1	0.3	0.6	2.0	5.0	8.1
2002.....	0.7	*	*	*	*	*	0.1	0.3	0.9	2.0	5.2	7.9
2001.....	0.7	*	*	*	*	*	0.2	0.3	0.8	2.3	5.4	9.1
2000.....	0.8	*	*	*	*	*	0.2	0.4	1.0	2.8	5.7	9.1
1999.....	0.8	*	*	*	*	*	0.1	0.4	1.0	2.7	6.0	9.5
White, male												
2004.....	0.3	*	*	*	*	*	*	0.2	0.3	1.2	3.2	5.0
2003.....	0.4	*	*	*	*	*	*	0.2	0.5	1.5	3.8	6.4
2002.....	0.4	*	*	*	*	*	*	0.2	0.7	1.6	3.2	4.5
2001.....	0.4	*	*	*	*	*	0.2	0.2	0.6	1.5	3.4	6.5
2000.....	0.5	*	*	*	*	*	0.2	0.3	0.9	1.7	4.0	6.4
1999.....	0.4	*	*	*	*	*	*	0.3	0.6	2.0	3.8	5.3

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Rheumatic mitral valve diseases (I05)												
White, female												
2004.....	0.8	*	*	*	*	*	0.1	0.3	0.7	2.1	4.7	8.0
2003.....	0.9	*	*	*	*	*	0.1	0.3	0.6	2.5	5.8	8.8
2002.....	1.0	*	*	*	*	*	0.1	0.4	1.1	2.4	6.6	9.3
2001.....	1.0	*	*	*	*	*	0.2	0.4	1.0	2.9	6.7	10.1
2000.....	1.1	*	*	*	*	*	0.2	0.5	1.1	3.6	6.8	10.3
1999.....	1.2	*	*	*	*	*	0.1	0.4	1.3	3.3	7.5	11.2
Black, both sexes												
2004.....	0.3	*	*	*	*	*	0.4	0.6	0.7	1.2	2.3	*
2003.....	0.4	*	*	*	*	*	*	0.8	1.6	1.6	2.3	*
2002.....	0.4	*	*	*	*	*	*	0.6	1.6	1.4	*	*
2001.....	0.4	*	*	*	*	*	0.3	0.9	1.2	2.3	2.2	*
2000.....	0.4	*	*	*	*	*	*	*	1.1	1.8	2.2	*
1999.....	0.5	*	*	*	*	*	*	0.9	1.3	2.6	2.4	*
Black, male												
2004.....	0.2	*	*	*	*	*	*	*	*	*	*	*
2003.....	0.3	*	*	*	*	*	*	*	2.0	*	*	*
2002.....	0.3	*	*	*	*	*	*	*	*	*	*	*
2001.....	0.2	*	*	*	*	*	*	*	*	*	*	*
2000.....	0.2	*	*	*	*	*	*	*	*	*	*	*
1999.....	0.3	*	*	*	*	*	*	*	*	*	*	*
Black, female												
2004.....	0.4	*	*	*	*	*	*	*	*	*	*	*
2003.....	0.5	*	*	*	*	*	*	1.0	*	2.0	*	*
2002.....	0.4	*	*	*	*	*	*	*	1.8	*	*	*
2001.....	0.6	*	*	*	*	*	*	1.1	1.6	2.9	*	*
2000.....	0.5	*	*	*	*	*	*	*	1.5	2.3	*	*
1999.....	0.6	*	*	*	*	*	*	1.1	1.5	3.1	*	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Disorders of both mitral and aortic valves (I08.0)												
All races 3/, both sexes												
2004.....	0.2	*	*	*	*	*	*	0.1	0.2	0.6	1.9	5.2
2003.....	0.3	*	*	*	*	*	*	0.1	0.2	0.6	2.1	6.5
2002.....	0.2	*	*	*	*	*	*	0.1	0.1	0.5	1.5	3.5
2001.....	0.1	*	*	*	*	*	*	*	0.1	0.5	1.0	2.5
2000.....	0.1	*	*	*	*	*	*	*	0.1	0.4	0.9	2.0
1999.....	0.1	*	*	*	*	*	*	*	0.1	0.4	0.8	2.0
All races 3/, male												
2004.....	0.2	*	*	*	*	*	*	*	0.2	0.5	2.1	3.8
2003.....	0.2	*	*	*	*	*	*	*	*	0.6	1.9	6.0
2002.....	0.1	*	*	*	*	*	*	*	*	0.5	1.4	2.4
2001.....	0.1	*	*	*	*	*	*	*	*	0.4	1.0	2.3
2000.....	0.1	*	*	*	*	*	*	*	*	0.3	0.7	1.9
1999.....	0.1	*	*	*	*	*	*	*	*	0.4	0.7	*
All races 3/, female												
2004.....	0.3	*	*	*	*	*	*	*	0.2	0.7	1.8	5.9
2003.....	0.3	*	*	*	*	*	*	*	0.2	0.6	2.2	6.7
2002.....	0.2	*	*	*	*	*	*	*	*	0.5	1.6	4.0
2001.....	0.2	*	*	*	*	*	*	*	*	0.5	1.1	2.5
2000.....	0.1	*	*	*	*	*	*	*	*	0.4	1.0	2.1
1999.....	0.1	*	*	*	*	*	*	*	0.2	0.4	0.8	2.2
White, both sexes												
2004.....	0.3	*	*	*	*	*	*	0.1	0.2	0.6	2.1	5.6
2003.....	0.3	*	*	*	*	*	*	*	0.2	0.6	2.2	7.0
2002.....	0.2	*	*	*	*	*	*	*	0.1	0.5	1.6	3.8
2001.....	0.2	*	*	*	*	*	*	*	0.1	0.5	1.1	2.6
2000.....	0.1	*	*	*	*	*	*	*	0.1	0.4	1.0	2.0
1999.....	0.1	*	*	*	*	*	*	*	0.1	0.4	0.8	2.1
White, male												
2004.....	0.2	*	*	*	*	*	*	*	*	0.6	2.2	4.1
2003.....	0.2	*	*	*	*	*	*	*	*	0.6	2.1	6.4
2002.....	0.1	*	*	*	*	*	*	*	*	0.5	1.5	2.7
2001.....	0.1	*	*	*	*	*	*	*	*	0.4	1.0	2.5
2000.....	0.1	*	*	*	*	*	*	*	*	0.3	0.7	1.9
1999.....	0.1	*	*	*	*	*	*	*	*	0.4	0.7	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)

All races 3/, both sexes

2004.....	0.3	*	*	*	*	*	0.1	0.1	0.3	1.0	2.6	4.4
2003.....	0.3	*	*	*	*	*	0.1	0.1	0.4	1.1	2.4	5.4
2002.....	0.4	*	*	*	*	*	0.1	0.1	0.5	1.3	2.7	5.6
2001.....	0.4	*	*	*	*	*	0.1	0.1	0.5	1.2	3.0	5.9
2000.....	0.4	*	*	*	*	*	0.1	0.2	0.5	1.3	3.2	5.8
1999.....	0.4	*	*	*	*	*	0.1	0.2	0.5	1.5	3.4	6.6

All races 3/, male

2004.....	0.2	*	*	*	*	*	*	0.1	0.2	0.6	1.8	2.5
2003.....	0.2	*	*	*	*	*	*	0.1	0.3	0.7	1.6	3.4
2002.....	0.2	*	*	*	*	*	*	0.1	0.4	0.8	1.9	3.8
2001.....	0.2	*	*	*	*	*	*	0.1	0.4	0.7	2.0	3.6
2000.....	0.2	*	*	*	*	*	*	0.1	0.4	0.8	2.0	4.0
1999.....	0.2	*	*	*	*	*	*	0.1	0.3	1.1	2.2	3.7

All races 3/, female

2004.....	0.5	*	*	*	*	*	*	0.2	0.4	1.3	3.1	5.3
2003.....	0.5	*	*	*	*	*	*	0.2	0.5	1.4	3.0	6.3
2002.....	0.5	*	*	*	*	*	*	0.2	0.6	1.6	3.2	6.3
2001.....	0.5	*	*	*	*	*	*	0.2	0.5	1.6	3.7	6.8
2000.....	0.6	*	*	*	*	*	0.1	0.2	0.7	1.8	4.0	6.6
1999.....	0.6	*	*	*	*	*	0.1	0.2	0.7	1.8	4.2	7.8

White, both sexes

2004.....	0.3	*	*	*	*	*	*	0.1	0.3	1.0	2.8	4.7
2003.....	0.4	*	*	*	*	*	*	0.1	0.4	1.1	2.6	5.7
2002.....	0.4	*	*	*	*	*	*	0.1	0.5	1.3	2.7	5.9
2001.....	0.4	*	*	*	*	*	*	0.1	0.4	1.2	3.1	6.3
2000.....	0.4	*	*	*	*	*	0.1	0.2	0.5	1.4	3.4	6.0
1999.....	0.5	*	*	*	*	*	0.1	0.2	0.5	1.6	3.5	6.9

White, male

2004.....	0.2	*	*	*	*	*	*	*	0.2	0.6	1.9	2.6
2003.....	0.2	*	*	*	*	*	*	*	0.3	0.7	1.8	3.5
2002.....	0.2	*	*	*	*	*	*	*	0.4	0.8	2.0	4.0
2001.....	0.2	*	*	*	*	*	*	*	0.4	0.8	2.1	3.7
2000.....	0.3	*	*	*	*	*	*	0.1	0.4	0.8	2.1	4.1
1999.....	0.3	*	*	*	*	*	*	*	0.3	1.2	2.3	3.9

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hypertensive diseases (I10-I13)												
All races 3/, both sexes												
2004.....	18.6	*	*	*	0.2	1.1	4.0	11.1	20.6	40.8	108.5	416.1
2003.....	18.4	*	*	*	0.2	1.1	4.0	10.9	20.3	41.2	109.7	409.5
2002.....	17.2	*	*	*	0.1	1.0	3.7	9.9	19.2	39.5	104.6	390.8
2001.....	16.4	*	*	*	0.1	1.0	3.4	9.0	17.7	37.5	103.1	384.7
2000.....	15.9	*	*	*	0.1	0.8	3.1	8.3	17.8	37.0	100.7	380.3
1999.....	15.4	*	*	*	0.1	0.8	3.1	7.9	17.3	36.8	100.1	366.0
All races 3/, male												
2004.....	16.0	*	*	*	0.2	1.5	5.4	14.8	27.0	47.8	108.5	335.7
2003.....	15.4	*	*	*	0.3	1.4	5.2	14.7	25.7	47.8	107.3	322.4
2002.....	14.5	*	*	*	0.2	1.4	5.0	13.2	24.8	44.6	103.7	311.1
2001.....	13.5	*	*	*	0.2	1.2	4.5	12.0	21.9	42.7	101.0	309.4
2000.....	13.0	*	*	*	0.2	1.0	4.0	11.2	22.4	42.0	98.8	309.3
1999.....	12.6	*	*	*	0.2	1.1	4.1	10.3	21.4	41.1	99.3	297.3
All races 3/, female												
2004.....	21.2	*	*	*	0.1	0.7	2.6	7.5	14.7	34.9	108.5	452.3
2003.....	21.2	*	*	*	0.1	0.8	2.8	7.3	15.3	35.7	111.3	448.0
2002.....	19.9	*	*	*	*	0.7	2.4	6.7	14.1	35.2	105.2	425.4
2001.....	19.2	*	*	*	*	0.7	2.3	6.1	13.8	33.2	104.5	416.3
2000.....	18.6	*	*	*	0.1	0.6	2.2	5.5	13.5	32.9	101.9	409.3
1999.....	18.1	*	*	*	*	0.5	2.1	5.6	13.6	33.3	100.7	393.7
White, both sexes												
2004.....	17.2	*	*	*	0.1	0.6	2.7	7.5	14.8	32.4	95.4	394.8
2003.....	16.9	*	*	*	0.1	0.7	2.6	7.2	14.6	31.8	97.2	389.7
2002.....	15.8	*	*	*	0.1	0.6	2.3	6.7	13.7	30.9	91.3	368.7
2001.....	15.0	*	*	*	0.1	0.6	2.0	5.8	12.4	28.9	90.6	364.3
2000.....	14.4	*	*	*	0.1	0.4	2.0	5.3	12.2	28.3	87.5	357.1
1999.....	13.9	*	*	*	*	0.5	1.8	4.9	11.8	28.0	86.2	343.4
White, male												
2004.....	14.2	*	*	*	0.2	1.0	3.8	10.6	19.8	38.4	95.2	318.7
2003.....	13.6	*	*	*	0.2	0.9	3.8	10.4	19.2	37.5	95.2	301.5
2002.....	12.7	*	*	*	0.1	0.8	3.4	9.4	18.2	35.3	90.1	290.6
2001.....	11.8	*	*	*	*	0.8	3.0	8.1	16.0	33.6	88.2	288.6
2000.....	11.3	*	*	*	*	0.6	2.8	7.6	15.9	32.9	86.1	286.2
1999.....	10.8	*	*	*	*	0.8	2.6	6.8	15.1	31.7	85.3	273.5

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hypertensive diseases (I10-I13)												
White, female												
2004.....	20.1	*	*	*	*	0.3	1.5	4.4	10.1	27.2	95.6	429.1
2003.....	20.1	*	*	*	*	0.4	1.4	4.1	10.3	26.9	98.6	428.7
2002.....	18.8	*	*	*	*	0.3	1.3	4.0	9.4	27.1	92.1	402.6
2001.....	18.2	*	*	*	*	0.3	1.1	3.6	9.1	24.9	92.1	396.0
2000.....	17.4	*	*	*	*	0.2	1.2	2.9	8.7	24.5	88.5	385.8
1999.....	16.9	*	*	*	*	0.2	0.9	3.1	8.6	24.9	86.8	371.5
Black, both sexes												
2004.....	32.2	*	*	*	0.5	3.9	13.5	39.2	73.9	124.9	269.0	698.5
2003.....	32.1	*	*	*	0.6	3.8	13.7	40.2	74.2	132.7	265.4	686.5
2002.....	30.7	*	*	*	0.4	3.9	12.8	35.9	71.5	125.4	270.7	685.4
2001.....	29.1	*	*	*	0.4	3.5	12.5	35.0	66.1	123.0	257.1	664.2
2000.....	28.9	*	*	*	0.5	3.1	11.3	33.0	69.7	125.5	261.8	673.8
1999.....	28.6	*	*	*	0.5	2.7	12.5	32.4	67.2	126.3	270.5	639.9
Black, male												
2004.....	31.3	*	*	*	4.9	17.5	50.0	99.0	152.1	291.8	572.2	
2003.....	30.8	*	*	*	0.8	4.7	16.3	51.7	93.5	161.7	275.6	612.0
2002.....	29.3	*	*	*	5.3	16.7	46.1	91.8	145.7	287.7	602.5	
2001.....	27.7	*	*	*	4.3	15.7	46.3	81.2	145.4	274.0	604.8	
2000.....	26.8	*	*	*	3.6	13.9	42.5	88.1	143.1	269.2	606.6	
1999.....	26.8	*	*	*	0.8	3.2	15.6	41.4	82.8	145.3	286.9	589.4
Black, female												
2004.....	33.0	*	*	*	3.1	10.0	30.0	53.8	105.5	256.2	748.6	
2003.....	33.3	*	*	*	3.0	11.4	30.3	58.7	112.1	259.6	715.4	
2002.....	31.9	*	*	*	2.6	9.4	27.2	55.1	111.1	261.0	717.1	
2001.....	30.5	*	*	*	2.8	9.7	25.3	54.0	107.2	247.7	686.2	
2000.....	30.8	*	*	*	2.6	9.0	24.8	54.9	113.0	257.6	698.8	
1999.....	30.3	*	*	*	2.3	9.7	24.7	54.6	113.0	261.3	658.8	

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Essential (primary) hypertension (I10)												
All races 3/, both sexes												
2004.....	4.6	*	*	*	*	0.1	0.3	1.6	3.6	9.1	28.5	127.4
2003.....	4.4	*	*	*	*	0.1	0.4	1.4	3.6	9.2	28.5	121.9
2002.....	4.0	*	*	*	*	0.1	0.4	1.3	3.2	8.6	26.3	112.0
2001.....	3.9	*	*	*	*	0.1	0.3	1.3	3.4	8.4	25.9	107.4
2000.....	3.7	*	*	*	*	0.1	0.3	1.4	3.4	8.2	24.1	101.1
1999.....	3.6	*	*	*	*	0.1	0.3	1.2	3.3	8.1	24.0	98.1
All races 3/, male												
2004.....	3.2	*	*	*	*	*	0.5	2.1	4.4	10.0	26.6	86.5
2003.....	3.0	*	*	*	*	*	0.4	1.9	4.5	10.4	24.3	77.4
2002.....	2.8	*	*	*	*	*	0.5	1.7	4.0	9.4	23.3	74.2
2001.....	2.8	*	*	*	*	0.1	0.4	1.7	4.2	9.3	23.2	74.8
2000.....	2.6	*	*	*	*	*	0.4	1.9	4.2	8.9	22.0	70.3
1999.....	2.5	*	*	*	*	*	0.4	1.5	4.1	9.2	21.9	68.1
All races 3/, female												
2004.....	5.9	*	*	*	*	*	0.2	1.1	2.9	8.3	29.8	145.8
2003.....	5.8	*	*	*	*	*	0.4	1.0	2.9	8.3	31.3	141.5
2002.....	5.2	*	*	*	*	*	0.3	1.0	2.5	8.0	28.2	128.3
2001.....	5.0	*	*	*	*	*	0.3	0.9	2.7	7.7	27.6	121.1
2000.....	4.6	*	*	*	*	*	0.3	0.9	2.6	7.5	25.5	113.6
1999.....	4.5	*	*	*	*	*	0.2	1.0	2.7	7.3	25.4	110.2
White, both sexes												
2004.....	4.6	*	*	*	*	*	0.3	1.2	2.9	8.0	26.7	123.5
2003.....	4.5	*	*	*	*	*	0.3	1.1	2.9	7.7	26.1	119.9
2002.....	4.1	*	*	*	*	*	0.3	1.0	2.8	7.5	24.3	108.6
2001.....	3.9	*	*	*	*	0.1	0.2	0.9	2.7	7.1	24.0	104.6
2000.....	3.6	*	*	*	*	*	0.3	1.0	2.7	7.2	22.2	97.5
1999.....	3.5	*	*	*	*	*	0.2	0.9	2.7	7.1	22.0	94.3
White, male												
2004.....	3.2	*	*	*	*	*	0.4	1.7	3.7	8.8	24.6	84.0
2003.....	2.9	*	*	*	*	*	0.4	1.6	3.6	8.9	21.4	74.3
2002.....	2.7	*	*	*	*	*	0.3	1.3	3.5	8.2	21.0	71.0
2001.....	2.6	*	*	*	*	*	0.3	1.3	3.4	7.9	20.8	71.1
2000.....	2.5	*	*	*	*	*	0.4	1.4	3.4	7.9	19.8	66.4
1999.....	2.4	*	*	*	*	*	0.3	1.2	3.4	8.0	19.7	62.7

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Essential (primary) hypertension (I10)												
White, female												
2004.....	6.0	*	*	*	*	*	0.2	0.7	2.2	7.2	28.1	141.3
2003.....	6.0	*	*	*	*	*	0.3	0.7	2.2	6.6	29.2	140.2
2002.....	5.4	*	*	*	*	*	0.2	0.6	2.1	6.9	26.5	124.9
2001.....	5.1	*	*	*	*	*	0.1	0.6	2.1	6.4	26.1	118.7
2000.....	4.7	*	*	*	*	*	0.2	0.6	2.1	6.6	23.8	110.2
1999.....	4.6	*	*	*	*	*	*	0.6	2.1	6.4	23.5	107.0
Black, both sexes												
2004.....	5.5	*	*	*	*	*	1.0	4.5	9.7	20.3	52.8	183.6
2003.....	5.5	*	*	*	*	*	1.1	3.8	10.6	25.2	59.8	157.5
2002.....	5.0	*	*	*	*	*	1.2	4.0	7.9	20.7	52.9	158.6
2001.....	4.9	*	*	*	*	*	1.1	4.0	9.9	21.6	50.1	150.8
2000.....	4.7	*	*	*	*	*	1.0	4.4	9.7	18.7	47.4	149.2
1999.....	4.6	*	*	*	*	*	1.1	3.9	9.4	18.9	48.7	143.2
Black, male												
2004.....	4.4	*	*	*	*	*	1.3	5.2	11.7	22.4	56.8	124.2
2003.....	4.7	*	*	*	*	*	1.1	5.1	13.3	28.0	63.2	119.9
2002.....	4.0	*	*	*	*	*	1.4	4.8	9.4	22.8	55.1	117.9
2001.....	4.3	*	*	*	*	*	1.2	5.2	12.3	25.1	55.6	132.8
2000.....	4.0	*	*	*	*	*	0.9	5.9	13.0	20.6	48.7	123.4
1999.....	3.8	*	*	*	*	*	1.2	4.2	11.3	22.2	50.1	124.2
Black, female												
2004.....	6.4	*	*	*	*	*	0.7	3.9	8.2	18.7	50.5	207.1
2003.....	6.3	*	*	*	*	*	1.1	2.8	8.5	23.2	57.9	172.2
2002.....	5.8	*	*	*	*	*	0.9	3.4	6.6	19.3	51.6	174.2
2001.....	5.5	*	*	*	*	*	0.9	3.0	8.0	19.2	46.9	157.5
2000.....	5.3	*	*	*	*	*	1.0	3.2	6.9	17.4	46.7	158.8
1999.....	5.3	*	*	*	*	*	1.0	3.7	7.9	16.5	47.9	150.2

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hypertensive heart disease (I11)												
All races 3/, both sexes												
2004.....	9.7	*	*	*	0.1	0.8	3.0	7.8	13.4	21.7	49.3	192.8
2003.....	9.7	*	*	*	0.1	0.8	2.9	7.9	13.1	22.1	51.0	195.0
2002.....	9.2	*	*	*	0.1	0.7	2.7	7.2	12.5	21.3	49.6	187.0
2001.....	8.7	*	*	*	0.1	0.6	2.4	6.3	10.9	19.8	48.7	189.5
2000.....	8.4	*	*	*	0.1	0.5	2.2	5.6	11.0	19.6	48.2	193.1
1999.....	8.1	*	*	*	0.1	0.5	2.1	5.3	10.7	19.1	47.7	184.9
All races 3/, male												
2004.....	8.9	*	*	*	0.1	1.1	4.1	10.7	18.5	27.3	48.0	145.4
2003.....	8.8	*	*	*	0.2	1.0	4.0	10.8	17.2	27.1	51.0	142.2
2002.....	8.2	*	*	*	0.1	1.1	3.8	9.8	16.9	25.2	49.4	134.6
2001.....	7.5	*	*	*	*	0.8	3.3	8.6	14.4	23.6	48.4	134.4
2000.....	7.2	*	*	*	0.1	0.7	2.9	7.8	14.5	23.7	46.5	139.1
1999.....	6.8	*	*	*	0.1	0.7	2.9	7.0	13.7	21.9	46.8	133.6
All races 3/, female												
2004.....	10.5	*	*	*	*	0.4	1.9	5.1	8.6	16.9	50.2	214.1
2003.....	10.7	*	*	*	*	0.5	1.9	5.2	9.3	17.8	51.0	218.3
2002.....	10.2	*	*	*	*	0.4	1.6	4.6	8.4	18.1	49.8	209.7
2001.....	9.8	*	*	*	*	0.4	1.6	4.1	7.8	16.6	48.8	212.6
2000.....	9.7	*	*	*	*	0.4	1.5	3.5	7.8	16.2	49.4	215.1
1999.....	9.4	*	*	*	*	0.3	1.3	3.6	7.9	16.8	48.3	205.5
White, both sexes												
2004.....	8.9	*	*	*	0.1	0.5	2.1	5.6	10.0	17.6	43.2	183.7
2003.....	8.8	*	*	*	0.1	0.5	2.0	5.4	9.8	17.3	45.1	185.3
2002.....	8.3	*	*	*	*	0.4	1.9	5.1	9.1	16.7	42.9	177.3
2001.....	7.8	*	*	*	*	0.4	1.6	4.2	7.9	15.2	42.6	179.9
2000.....	7.6	*	*	*	*	0.3	1.5	3.7	7.6	15.0	41.5	181.8
1999.....	7.3	*	*	*	*	0.3	1.3	3.4	7.5	14.4	40.9	175.3
White, male												
2004.....	7.8	*	*	*	*	0.7	3.1	8.1	13.9	22.3	41.9	136.4
2003.....	7.7	*	*	*	0.1	0.7	3.1	7.9	13.4	21.3	44.8	131.9
2002.....	7.1	*	*	*	*	0.7	2.8	7.4	12.7	19.9	42.6	124.8
2001.....	6.4	*	*	*	*	0.6	2.4	6.1	10.9	18.8	41.9	123.7
2000.....	6.1	*	*	*	*	0.4	2.1	5.5	10.4	18.6	39.6	129.3
1999.....	5.8	*	*	*	*	0.6	2.0	4.9	10.0	16.6	39.7	123.7

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hypertensive heart disease (I11)												
White, female												
2004.....	9.9	*	*	*	*	0.2	1.1	3.1	6.3	13.5	44.0	205.0
2003.....	10.0	*	*	*	*	0.2	1.0	2.9	6.4	13.9	45.2	209.0
2002.....	9.5	*	*	*	*	0.2	0.9	2.8	5.8	14.1	43.1	200.1
2001.....	9.2	*	*	*	*	0.2	0.8	2.4	5.1	12.3	43.0	203.5
2000.....	9.0	*	*	*	*	0.2	0.9	1.9	5.0	11.9	42.9	203.1
1999.....	8.7	*	*	*	*	*	0.7	2.0	5.1	12.5	41.7	196.0
Black, both sexes												
2004.....	17.6	*	*	*	0.3	2.7	9.2	26.2	45.1	64.6	126.8	314.7
2003.....	17.9	*	*	*	0.4	2.5	9.4	27.9	44.7	69.3	126.3	332.3
2002.....	17.2	*	*	*	*	2.8	8.6	24.3	44.4	68.0	133.0	320.4
2001.....	16.0	*	*	*	*	2.1	8.3	23.0	39.2	65.6	124.4	326.1
2000.....	16.1	*	*	*	*	2.1	7.4	21.2	42.2	66.8	131.9	340.0
1999.....	15.5	*	*	*	0.3	1.8	7.9	20.9	39.6	67.2	131.9	305.1
Black, male												
2004.....	18.4	*	*	*	*	3.5	11.7	33.6	64.6	86.6	131.7	267.2
2003.....	18.1	*	*	*	*	3.0	11.2	35.0	57.1	92.3	137.1	287.8
2002.....	17.5	*	*	*	*	3.9	11.6	31.3	60.8	83.6	141.1	267.0
2001.....	15.9	*	*	*	*	2.6	10.3	30.2	49.8	79.1	137.1	284.8
2000.....	15.6	*	*	*	*	2.6	9.2	27.8	54.9	81.4	139.3	273.3
1999.....	15.2	*	*	*	*	2.1	10.1	26.3	50.6	80.4	142.3	263.6
Black, female												
2004.....	16.9	*	*	*	*	1.9	7.0	19.8	29.4	48.9	124.0	333.5
2003.....	17.8	*	*	*	*	2.0	7.9	21.8	34.7	52.9	120.2	349.6
2002.....	16.9	*	*	*	*	1.7	6.0	18.3	31.2	56.8	128.5	340.9
2001.....	16.2	*	*	*	*	1.7	6.5	16.8	30.6	56.1	117.3	341.4
2000.....	16.5	*	*	*	*	1.7	5.8	15.5	31.9	56.6	127.7	364.9
1999.....	15.9	*	*	*	*	1.5	5.9	16.2	30.8	57.9	126.0	320.6

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hypertensive renal disease (I12)												
All races 3/, both sexes												
2004.....	3.3	*	*	*	*	0.2	0.4	1.2	2.7	8.0	24.0	71.1
2003.....	3.1	*	*	*	*	0.2	0.4	1.1	2.6	7.7	23.2	67.0
2002.....	3.0	*	*	*	*	0.2	0.4	1.0	2.5	7.3	21.9	68.4
2001.....	2.9	*	*	*	*	0.2	0.4	1.1	2.4	7.0	21.8	64.5
2000.....	2.8	*	*	*	*	0.1	0.4	1.0	2.6	7.0	21.4	61.8
1999.....	2.5	*	*	*	*	0.1	0.4	1.0	2.2	7.1	19.6	53.9
All races 3/, male												
2004.....	2.9	*	*	*	*	0.2	0.5	1.4	3.0	8.2	26.7	79.5
2003.....	2.7	*	*	*	*	0.2	0.5	1.4	2.9	8.0	24.7	76.1
2002.....	2.6	*	*	*	*	0.2	0.5	1.2	2.7	7.7	24.1	79.2
2001.....	2.4	*	*	*	*	0.2	0.5	1.3	2.3	7.3	22.6	76.0
2000.....	2.4	*	*	*	*	0.1	0.5	1.1	2.8	6.8	23.2	72.0
1999.....	2.2	*	*	*	*	0.2	0.5	1.2	2.4	7.4	21.1	64.3
All races 3/, female												
2004.....	3.6	*	*	*	*	0.2	0.3	0.9	2.5	7.8	22.3	67.4
2003.....	3.5	*	*	*	*	0.1	0.3	0.8	2.4	7.5	22.2	63.0
2002.....	3.4	*	*	*	*	0.1	0.3	0.8	2.3	7.1	20.6	63.7
2001.....	3.3	*	*	*	*	0.2	0.3	0.9	2.5	6.8	21.3	59.7
2000.....	3.1	*	*	*	*	0.1	0.3	0.8	2.4	7.1	20.2	57.7
1999.....	2.8	*	*	*	*	*	0.3	0.7	2.1	6.8	18.6	49.8
White, both sexes												
2004.....	2.8	*	*	*	*	0.1	0.2	0.5	1.4	5.5	20.1	64.6
2003.....	2.7	*	*	*	*	0.1	0.2	0.5	1.4	5.3	20.1	61.1
2002.....	2.6	*	*	*	*	0.1	0.2	0.5	1.3	5.3	18.4	61.9
2001.....	2.5	*	*	*	*	0.1	0.2	0.5	1.3	5.0	18.3	58.7
2000.....	2.3	*	*	*	*	*	0.2	0.4	1.4	4.6	18.2	55.7
1999.....	2.1	*	*	*	*	0.1	0.2	0.4	1.0	4.9	15.9	47.7
White, male												
2004.....	2.4	*	*	*	*	0.1	0.3	0.6	1.6	5.7	22.6	75.4
2003.....	2.3	*	*	*	*	*	0.3	0.6	1.5	5.6	22.3	71.0
2002.....	2.2	*	*	*	*	*	0.2	0.5	1.4	5.6	20.5	74.2
2001.....	2.1	*	*	*	*	*	0.2	0.5	1.1	5.2	19.6	71.1
2000.....	2.0	*	*	*	*	*	0.2	0.5	1.6	4.7	20.5	66.1
1999.....	1.8	*	*	*	*	*	0.2	0.5	1.1	5.3	17.6	58.3

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hypertensive renal disease (I12)												
White, female												
2004.....	3.1	*	*	*	*	*	0.1	0.4	1.2	5.3	18.4	59.8
2003.....	3.1	*	*	*	*	*	0.1	0.4	1.3	5.1	18.6	56.6
2002.....	2.9	*	*	*	*	*	0.1	0.4	1.2	5.0	16.9	56.5
2001.....	2.9	*	*	*	*	*	0.1	0.4	1.4	4.9	17.5	53.5
2000.....	2.7	*	*	*	*	*	*	0.3	1.2	4.6	16.6	51.4
1999.....	2.4	*	*	*	*	*	0.1	0.4	0.9	4.6	14.8	43.5
Black, both sexes												
2004.....	6.8	*	*	*	*	0.7	2.1	5.6	14.2	31.6	69.9	152.0
2003.....	6.2	*	*	*	*	0.7	2.0	5.5	13.5	29.2	59.7	143.5
2002.....	6.2	*	*	*	*	0.6	2.1	4.9	13.1	26.8	65.1	150.3
2001.....	5.9	*	*	*	*	0.8	2.0	5.9	12.1	25.8	62.5	137.3
2000.....	5.9	*	*	*	*	0.4	2.2	5.2	13.2	30.0	59.7	134.2
1999.....	5.7	*	*	*	*	0.4	2.0	5.2	12.1	27.8	63.0	126.3
Black, male												
2004.....	6.3	*	*	*	*	0.8	2.6	7.6	16.6	34.4	80.2	139.1
2003.....	5.7	*	*	*	*	1.0	2.4	7.7	16.0	32.5	56.4	154.3
2002.....	5.5	*	*	*	*	*	2.7	6.2	14.0	29.1	72.3	156.5
2001.....	5.3	*	*	*	*	0.9	2.6	7.6	13.2	29.4	62.7	143.0
2000.....	5.1	*	*	*	*	*	2.8	6.0	14.8	29.8	60.3	146.5
1999.....	5.3	*	*	*	*	*	2.4	7.4	13.3	29.1	68.6	141.8
Black, female												
2004.....	7.3	*	*	*	*	*	1.7	3.9	12.2	29.7	64.0	157.1
2003.....	6.7	*	*	*	*	*	1.6	3.6	11.5	26.8	61.6	139.4
2002.....	6.7	*	*	*	*	*	1.6	3.8	12.4	25.2	61.1	147.9
2001.....	6.4	*	*	*	*	0.7	1.5	4.3	11.3	23.3	62.5	135.1
2000.....	6.7	*	*	*	*	*	1.6	4.6	12.0	30.1	59.3	129.6
1999.....	6.1	*	*	*	*	*	1.6	3.3	11.1	26.9	59.9	120.5

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hypertensive heart and renal disease (I13)												
All races 3/, both sexes												
2004.....	1.0	*	*	*	*	0.1	0.2	0.5	0.9	2.1	6.6	24.8
2003.....	1.1	*	*	*	*	0.1	0.2	0.5	1.0	2.2	7.0	25.6
2002.....	1.0	*	*	*	*	0.1	0.2	0.4	1.0	2.2	6.8	23.5
2001.....	1.0	*	*	*	*	0.1	0.2	0.4	1.0	2.3	6.8	23.3
2000.....	1.0	*	*	*	*	0.1	0.2	0.4	0.9	2.3	7.0	24.3
1999.....	1.2	*	*	*	*	0.1	0.3	0.4	1.1	2.6	8.9	29.1
All races 3/, male												
2004.....	0.9	*	*	*	*	*	0.3	0.6	1.1	2.2	7.2	24.3
2003.....	0.9	*	*	*	*	*	0.3	0.6	1.2	2.3	7.4	26.7
2002.....	0.9	*	*	*	*	*	0.2	0.5	1.2	2.4	7.0	23.1
2001.....	0.8	*	*	*	*	*	0.3	0.5	1.0	2.5	6.8	24.2
2000.....	0.8	*	*	*	*	*	0.2	0.5	0.9	2.5	7.2	27.8
1999.....	1.0	*	*	*	*	*	0.3	0.6	1.3	2.7	9.6	31.3
All races 3/, female												
2004.....	1.2	*	*	*	*	*	0.1	0.4	0.7	1.9	6.2	25.0
2003.....	1.2	*	*	*	*	*	0.1	0.3	0.7	2.1	6.8	25.2
2002.....	1.2	*	*	*	*	*	0.1	0.3	0.8	2.0	6.7	23.7
2001.....	1.1	*	*	*	*	*	0.2	0.2	0.9	2.1	6.8	22.9
2000.....	1.1	*	*	*	*	*	0.1	0.3	0.8	2.1	6.9	22.9
1999.....	1.4	*	*	*	*	*	0.2	0.3	0.9	2.4	8.5	28.2
White, both sexes												
2004.....	0.9	*	*	*	*	*	0.1	0.2	0.5	1.3	5.5	23.0
2003.....	0.9	*	*	*	*	*	0.1	0.2	0.5	1.4	6.0	23.4
2002.....	0.8	*	*	*	*	*	0.1	0.1	0.5	1.4	5.7	21.0
2001.....	0.8	*	*	*	*	*	0.1	0.2	0.5	1.5	5.7	21.0
2000.....	0.8	*	*	*	*	*	0.1	0.2	0.4	1.5	5.6	22.0
1999.....	1.0	*	*	*	*	*	0.1	0.2	0.5	1.5	7.3	26.1
White, male												
2004.....	0.7	*	*	*	*	*	*	0.2	0.6	1.5	6.1	22.9
2003.....	0.7	*	*	*	*	*	*	0.2	0.6	1.6	6.6	24.3
2002.....	0.7	*	*	*	*	*	*	0.1	0.6	1.6	6.0	20.6
2001.....	0.7	*	*	*	*	*	*	0.2	0.5	1.6	5.9	22.7
2000.....	0.7	*	*	*	*	*	0.1	0.2	0.5	1.7	6.2	24.4
1999.....	0.8	*	*	*	*	*	*	0.2	0.6	1.7	8.2	28.8

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hypertensive heart and renal disease (I13)												
White, female												
2004.....	1.0	*	*	*	*	*	*	0.1	0.4	1.2	5.1	23.0
2003.....	1.0	*	*	*	*	*	*	*	0.4	1.3	5.5	22.9
2002.....	1.0	*	*	*	*	*	*	*	0.4	1.2	5.6	21.1
2001.....	1.0	*	*	*	*	*	*	0.1	0.5	1.4	5.5	20.3
2000.....	1.0	*	*	*	*	*	*	0.1	0.4	1.4	5.2	21.1
1999.....	1.1	*	*	*	*	*	*	0.1	0.5	1.4	6.7	25.0
Black, both sexes												
2004.....	2.3	*	*	*	*	*	1.2	2.8	4.9	8.5	19.6	48.2
2003.....	2.4	*	*	*	*	0.4	1.2	3.0	5.4	9.0	19.6	53.1
2002.....	2.4	*	*	*	*	*	0.9	2.7	6.1	9.9	19.6	56.1
2001.....	2.2	*	*	*	*	0.4	1.2	2.1	5.0	9.9	20.1	50.1
2000.....	2.2	*	*	*	*	*	0.8	2.2	4.6	9.9	22.8	50.4
1999.....	2.8	*	*	*	*	*	1.5	2.4	6.0	12.5	26.9	65.4
Black, male												
2004.....	2.2	*	*	*	*	*	1.9	3.5	6.1	8.7	23.1	41.7
2003.....	2.3	*	*	*	*	*	1.7	4.0	7.1	8.9	18.9	50.0
2002.....	2.3	*	*	*	*	*	1.0	3.8	7.5	10.1	19.2	61.1
2001.....	2.1	*	*	*	*	*	1.6	3.2	5.9	11.9	18.5	44.3
2000.....	2.0	*	*	*	*	*	1.0	2.8	5.3	11.3	20.9	63.4
1999.....	2.5	*	*	*	*	*	1.8	3.5	7.6	13.5	25.8	59.8
Black, female												
2004.....	2.4	*	*	*	*	*	*	2.3	4.0	8.3	17.6	50.8
2003.....	2.5	*	*	*	*	*	0.8	2.1	4.0	9.1	19.9	54.3
2002.....	2.5	*	*	*	*	*	0.9	1.8	4.9	9.7	19.8	54.2
2001.....	2.3	*	*	*	*	*	0.8	1.1	4.2	8.6	21.0	52.2
2000.....	2.4	*	*	*	*	*	*	1.6	4.1	9.0	23.9	45.5
1999.....	3.0	*	*	*	*	*	1.2	1.5	4.8	11.7	27.5	67.5

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Ischemic heart diseases (I20-I25)												
All races 3/, both sexes												
2004.....	153.7	*	*	*	0.3	2.7	16.2	59.9	158.2	395.9	1,072.6	3,287.4
2003.....	165.1	0.5	*	0.1	0.4	2.7	17.0	62.2	169.8	430.2	1,160.7	3,610.2
2002.....	171.4	0.6	*	0.0	0.3	2.5	17.1	63.8	177.7	458.9	1,220.2	3,775.0
2001.....	176.3	0.6	*	0.0	0.4	2.8	16.7	64.0	184.0	477.6	1,271.3	3,958.2
2000.....	183.1	0.8	*	*	0.3	2.3	16.7	66.1	196.7	505.1	1,324.2	4,182.5
1999.....	189.8	0.7	*	0.1	0.4	2.5	17.2	67.7	203.5	536.6	1,384.8	4,318.1
All races 3/, male												
2004.....	161.6	*	*	*	0.4	4.0	24.4	92.8	234.8	549.1	1,410.3	3,680.3
2003.....	172.2	*	*	*	0.5	4.0	25.7	97.2	251.5	598.8	1,524.4	4,005.2
2002.....	178.4	*	*	*	0.4	3.7	26.1	99.4	263.3	640.1	1,598.9	4,205.3
2001.....	181.7	*	*	*	0.6	4.1	25.5	99.5	271.6	663.7	1,663.5	4,395.8
2000.....	188.7	*	*	*	0.4	3.5	25.8	103.5	290.2	705.2	1,736.8	4,701.7
1999.....	195.4	*	*	*	0.5	3.7	26.6	106.2	299.3	749.8	1,813.7	4,837.5
All races 3/, female												
2004.....	146.1	*	*	*	0.1	1.4	8.0	28.0	87.1	267.3	845.3	3,110.6
2003.....	158.1	*	*	*	0.3	1.4	8.4	28.3	94.0	289.2	917.7	3,435.5
2002.....	164.7	*	*	*	0.2	1.3	8.2	29.5	98.5	308.1	968.9	3,588.3
2001.....	171.2	*	*	*	0.2	1.5	8.1	29.7	103.1	323.4	1,013.6	3,775.1
2000.....	177.6	*	*	*	0.2	1.2	7.6	29.9	110.4	340.3	1,055.2	3,971.0
1999.....	184.5	*	*	*	0.2	1.4	7.9	30.7	115.1	362.0	1,106.3	4,108.8
White, both sexes												
2004.....	166.1	*	*	*	0.2	2.5	15.7	56.8	151.3	385.6	1,073.4	3,343.9
2003.....	178.3	*	*	*	0.3	2.5	16.4	58.8	162.0	418.9	1,159.0	3,671.7
2002.....	185.4	*	*	*	0.3	2.3	16.5	60.7	169.5	448.8	1,217.8	3,836.4
2001.....	190.5	*	*	*	0.3	2.6	15.9	60.7	175.1	465.8	1,268.5	4,022.7
2000.....	197.9	0.8	*	*	0.3	2.1	15.8	62.8	187.3	493.4	1,320.9	4,257.7
1999.....	205.1	*	*	*	0.3	2.3	16.2	64.4	194.3	524.8	1,383.7	4,389.2
White, male												
2004.....	174.3	*	*	*	0.3	3.7	24.1	89.6	227.1	539.9	1,422.1	3,768.8
2003.....	185.5	*	*	*	0.5	3.8	25.1	93.5	243.5	587.9	1,529.5	4,106.1
2002.....	192.5	*	*	*	0.4	3.5	25.6	96.3	254.3	632.9	1,608.0	4,299.7
2001.....	195.8	*	*	*	0.5	3.9	24.7	96.3	262.3	653.8	1,672.4	4,494.0
2000.....	203.6	*	*	*	0.4	3.3	25.1	100.5	280.3	697.3	1,744.8	4,813.9
1999.....	210.5	*	*	*	0.4	3.4	25.5	102.8	289.0	741.3	1,827.8	4,933.9

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Ischemic heart diseases (I20-I25)												
White, female												
2004.....	158.1	*	*	*	*	1.2	7.2	24.4	79.5	253.5	835.6	3,152.3
2003.....	171.4	*	*	*	0.2	1.2	7.5	24.6	85.0	274.9	908.5	3,479.3
2002.....	178.4	*	*	*	0.2	1.1	7.3	25.5	89.5	292.7	956.0	3,635.2
2001.....	185.3	*	*	*	0.2	1.3	7.0	25.6	93.1	307.2	1,000.3	3,825.5
2000.....	192.4	*	*	*	0.2	0.9	6.5	25.6	100.0	322.5	1,042.0	4,031.8
1999.....	200.0	*	*	*	0.1	1.3	6.8	26.7	105.5	344.3	1,092.6	4,170.3
Black, both sexes												
2004.....	121.0	*	*	*	0.7	4.7	23.2	96.0	255.0	568.6	1,243.5	3,014.3
2003.....	129.2	*	*	*	0.7	4.3	24.7	99.3	273.1	615.2	1,363.0	3,310.1
2002.....	132.9	*	*	*	0.5	3.9	25.1	101.6	288.0	636.1	1,422.7	3,462.9
2001.....	136.9	*	*	*	0.9	4.6	25.6	102.3	297.9	673.0	1,479.0	3,615.1
2000.....	140.1	*	*	*	0.4	4.3	25.6	105.6	314.6	702.2	1,523.2	3,684.7
1999.....	143.7	*	*	*	0.8	3.9	27.5	108.5	320.1	735.6	1,552.7	3,805.8
Black, male												
2004.....	125.2	*	*	*	1.1	6.5	31.9	138.4	362.4	761.9	1,537.6	3,100.5
2003.....	133.4	*	*	*	0.9	6.2	35.0	144.5	381.8	827.3	1,744.8	3,285.3
2002.....	135.3	*	*	*	*	5.1	35.2	144.3	407.3	840.8	1,748.8	3,556.2
2001.....	138.9	*	*	*	1.2	6.6	36.0	145.7	412.8	894.0	1,825.4	3,775.0
2000.....	141.5	*	*	*	*	5.9	35.6	149.4	439.2	907.9	1,877.0	3,885.2
1999.....	146.0	*	*	*	0.9	5.7	39.5	156.9	452.9	962.3	1,870.8	4,080.1
Black, female												
2004.....	117.1	*	*	*	*	3.0	15.5	59.6	168.6	430.7	1,077.3	2,980.1
2003.....	125.3	*	*	*	*	2.5	15.6	60.5	185.6	464.3	1,147.2	3,319.7
2002.....	130.8	*	*	*	*	2.8	16.3	64.9	192.2	490.8	1,238.1	3,427.2
2001.....	135.2	*	*	*	*	2.8	16.4	65.0	205.3	517.1	1,284.5	3,555.7
2000.....	138.8	*	*	*	*	2.8	16.7	68.0	214.1	557.4	1,324.0	3,610.0
1999.....	141.7	*	*	*	0.7	2.3	16.9	67.0	213.6	575.8	1,374.6	3,703.2

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute myocardial infarction (I21-I22)												
All races 3/, both sexes												
2004.....	53.4	*	*	*	0.1	1.1	6.4	24.2	64.5	153.4	370.3	992.9
2003.....	58.7	*	*	*	0.2	1.0	7.0	25.5	70.9	168.6	412.7	1,118.8
2002.....	62.3	*	*	*	0.2	1.0	7.0	26.5	74.9	186.1	444.4	1,191.3
2001.....	64.9	*	*	*	0.2	1.2	7.1	27.6	78.4	193.2	471.2	1,265.4
2000.....	68.5	0.5	*	*	0.1	0.9	6.9	28.9	86.6	209.1	503.3	1,338.5
1999.....	71.5	*	*	*	0.2	1.1	7.2	29.6	90.0	225.0	529.3	1,379.3
All races 3/, male												
2004.....	57.4	*	*	*	0.2	1.6	9.6	37.3	95.3	209.5	477.0	1,120.5
2003.....	62.6	*	*	*	0.2	1.5	10.4	39.8	104.1	231.0	530.6	1,263.5
2002.....	66.2	*	*	*	0.2	1.4	10.5	40.6	109.2	256.2	569.2	1,366.4
2001.....	68.6	*	*	*	0.2	1.7	10.6	42.4	114.0	264.7	604.1	1,461.3
2000.....	72.7	*	*	*	0.1	1.3	10.6	44.7	125.7	286.9	648.3	1,568.3
1999.....	76.1	*	*	*	0.2	1.5	11.0	46.3	131.8	310.0	684.4	1,622.1
All races 3/, female												
2004.....	49.6	*	*	*	*	0.6	3.3	11.6	35.9	106.3	298.5	935.5
2003.....	54.8	*	*	*	0.1	0.6	3.6	11.7	40.0	116.5	333.9	1,054.8
2002.....	58.4	*	*	*	0.1	0.5	3.6	12.9	43.2	127.7	361.5	1,115.3
2001.....	61.3	*	*	*	*	0.7	3.5	13.3	45.4	134.0	383.9	1,183.5
2000.....	64.6	*	*	*	0.1	0.5	3.3	13.6	50.5	145.1	408.7	1,244.8
1999.....	67.0	*	*	*	0.1	0.6	3.4	13.5	51.5	155.4	428.6	1,281.5
White, both sexes												
2004.....	57.8	*	*	*	0.1	1.1	6.4	23.6	63.5	151.2	372.1	1,004.1
2003.....	63.4	*	*	*	0.2	1.0	6.8	24.7	69.5	166.8	412.7	1,132.6
2002.....	67.3	*	*	*	0.1	1.0	6.9	25.9	73.5	184.2	443.3	1,201.2
2001.....	70.1	*	*	*	0.1	1.1	6.8	26.9	76.8	190.8	471.1	1,277.8
2000.....	74.1	*	*	*	0.1	0.9	6.8	28.2	84.9	206.7	502.6	1,353.5
1999.....	77.4	*	*	*	0.1	1.0	7.0	29.1	88.6	223.3	530.2	1,393.2
White, male												
2004.....	62.3	*	*	*	0.1	1.6	9.7	37.0	95.1	208.6	484.2	1,143.7
2003.....	67.7	*	*	*	0.2	1.4	10.3	39.2	103.6	230.7	532.6	1,289.3
2002.....	71.8	*	*	*	0.2	1.4	10.5	40.4	108.9	257.3	572.0	1,386.1
2001.....	74.4	*	*	*	0.2	1.6	10.6	42.4	113.7	264.7	609.1	1,481.1
2000.....	78.8	*	*	*	0.2	1.3	10.6	44.7	125.0	287.2	652.5	1,592.9
1999.....	82.7	*	*	*	0.2	1.4	10.9	46.4	131.3	311.1	692.3	1,644.0

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute myocardial infarction (I21-I22)												
White, female												
2004.....	53.3	*	*	*	*	0.5	3.0	10.4	33.5	102.1	295.6	941.1
2003.....	59.2	*	*	*	*	0.5	3.2	10.4	37.2	112.3	331.5	1,063.2
2002.....	62.8	*	*	*	*	0.5	3.3	11.5	40.1	122.2	357.0	1,120.8
2001.....	66.0	*	*	*	*	0.7	3.1	11.7	42.0	128.4	379.6	1,192.7
2000.....	69.6	*	*	*	*	0.4	2.9	11.9	47.3	139.3	404.1	1,256.2
1999.....	72.3	*	*	*	*	0.6	3.1	12.1	48.6	150.1	423.9	1,292.4
Black, both sexes												
2004.....	42.0	*	*	*	0.3	1.7	8.8	35.4	91.1	204.7	420.0	979.9
2003.....	45.7	*	*	*	*	1.6	10.1	37.5	98.6	220.0	479.9	1,094.9
2002.....	48.5	*	*	*	*	1.6	9.7	37.9	106.9	238.8	517.3	1,201.2
2001.....	50.0	*	*	*	0.3	1.8	10.0	39.2	109.8	252.6	537.8	1,239.9
2000.....	52.2	*	*	*	*	1.7	9.9	41.1	120.1	268.7	571.1	1,275.7
1999.....	53.3	*	*	*	*	1.6	10.3	40.9	120.9	279.5	584.6	1,318.8
Black, male												
2004.....	42.4	*	*	*	*	2.0	11.6	50.3	123.5	265.8	486.1	1,009.3
2003.....	46.6	*	*	*	*	2.3	13.8	53.0	133.3	283.0	600.2	1,113.5
2002.....	48.3	*	*	*	*	2.1	13.3	52.0	141.6	300.1	623.4	1,280.1
2001.....	49.2	*	*	*	*	2.5	13.2	51.9	143.8	317.1	638.0	1,382.0
2000.....	52.0	*	*	*	*	2.2	13.4	54.9	160.3	335.8	687.3	1,418.5
1999.....	53.4	*	*	*	*	2.3	14.2	56.6	165.5	353.6	685.6	1,498.7
Black, female												
2004.....	41.6	*	*	*	*	1.4	6.2	22.5	65.0	161.1	382.7	968.3
2003.....	44.9	*	*	*	*	1.0	6.7	24.1	70.7	175.2	411.9	1,087.7
2002.....	48.8	*	*	*	*	1.2	6.5	25.9	79.1	195.3	457.2	1,171.0
2001.....	50.6	*	*	*	*	1.2	7.1	28.2	82.4	207.0	481.5	1,187.2
2000.....	52.5	*	*	*	*	1.3	6.8	29.1	87.7	221.6	505.6	1,222.5
1999.....	53.2	*	*	*	*	0.9	6.9	27.4	85.1	227.4	528.1	1,251.5

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other acute ischemic heart diseases (I24)												
All races 3/, both sexes												
2004.....	0.8	*	*	*	*	*	0.2	0.5	1.2	2.2	5.2	14.8
2003.....	1.1	*	*	*	*	*	0.2	0.7	1.4	3.2	6.5	20.0
2002.....	1.2	*	*	*	*	*	0.2	0.9	1.7	3.5	7.1	21.1
2001.....	1.2	*	*	*	*	*	0.2	0.8	1.8	3.5	6.9	21.5
2000.....	1.2	*	*	*	*	*	0.2	0.7	2.0	3.5	7.3	21.9
1999.....	1.2	*	*	*	*	*	0.2	0.7	1.9	3.7	7.0	21.9
All races 3/, male												
2004.....	0.9	*	*	*	*	*	0.3	0.7	1.8	3.0	6.3	16.3
2003.....	1.1	*	*	*	*	*	0.3	1.0	2.1	4.3	8.0	19.5
2002.....	1.2	*	*	*	*	*	0.2	1.3	2.5	4.7	8.7	21.4
2001.....	1.2	*	*	*	*	*	0.3	1.2	2.6	4.7	8.5	20.7
2000.....	1.3	*	*	*	*	*	0.3	1.2	2.9	4.8	9.1	20.7
1999.....	1.2	*	*	*	*	*	0.3	1.0	2.8	4.9	8.8	22.3
All races 3/, female												
2004.....	0.8	*	*	*	*	*	0.1	0.3	0.7	1.4	4.4	14.1
2003.....	1.0	*	*	*	*	*	0.1	0.4	0.9	2.2	5.5	20.2
2002.....	1.1	*	*	*	*	*	0.1	0.5	1.0	2.4	5.9	21.0
2001.....	1.1	*	*	*	*	*	0.1	0.4	1.1	2.5	5.9	21.8
2000.....	1.1	*	*	*	*	*	0.1	0.3	1.1	2.4	6.1	22.4
1999.....	1.1	*	*	*	*	*	*	0.4	1.1	2.6	5.8	21.7
White, both sexes												
2004.....	0.8	*	*	*	*	*	0.2	0.4	1.1	2.0	4.9	15.0
2003.....	1.1	*	*	*	*	*	0.2	0.6	1.3	3.0	6.3	20.2
2002.....	1.2	*	*	*	*	*	0.2	0.8	1.6	3.3	7.0	21.4
2001.....	1.2	*	*	*	*	*	0.2	0.7	1.6	3.3	6.8	21.5
2000.....	1.2	*	*	*	*	*	0.2	0.7	1.8	3.3	7.1	22.1
1999.....	1.2	*	*	*	*	*	0.1	0.6	1.7	3.5	6.8	22.4
White, male												
2004.....	0.9	*	*	*	*	*	0.3	0.6	1.6	2.6	6.1	16.2
2003.....	1.1	*	*	*	*	*	0.3	0.9	1.8	4.1	7.6	19.7
2002.....	1.3	*	*	*	*	*	0.2	1.1	2.3	4.4	8.6	21.9
2001.....	1.2	*	*	*	*	*	0.2	1.1	2.4	4.4	8.5	20.7
2000.....	1.3	*	*	*	*	*	0.3	1.1	2.6	4.4	8.9	21.0
1999.....	1.2	*	*	*	*	*	0.2	0.8	2.5	4.7	8.6	22.9

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other acute ischemic heart diseases (I24)												
White, female												
2004.....	0.8	*	*	*	*	*	*	0.2	0.6	1.4	4.2	14.4
2003.....	1.1	*	*	*	*	*	*	0.3	0.8	2.0	5.5	20.5
2002.....	1.2	*	*	*	*	*	*	0.4	0.9	2.3	5.9	21.1
2001.....	1.2	*	*	*	*	*	*	0.3	0.9	2.4	5.7	21.7
2000.....	1.2	*	*	*	*	*	*	0.3	0.9	2.3	5.9	22.6
1999.....	1.2	*	*	*	*	*	*	0.3	1.1	2.5	5.6	22.1
Black, both sexes												
2004.....	1.0	*	*	*	*	*	*	1.4	3.0	4.1	8.8	15.2
2003.....	1.1	*	*	*	*	*	0.5	1.3	3.0	5.5	9.9	19.8
2002.....	1.2	*	*	*	*	*	0.3	1.7	3.0	5.9	9.3	21.1
2001.....	1.2	*	*	*	*	*	0.5	1.5	3.9	5.7	8.5	24.0
2000.....	1.2	*	*	*	*	*	0.4	1.5	4.2	5.6	10.8	21.9
1999.....	1.2	*	*	*	*	*	0.5	1.8	3.6	5.7	10.7	17.2
Black, male												
2004.....	1.3	*	*	*	*	*	*	1.9	4.6	7.1	11.7	22.8
2003.....	1.4	*	*	*	*	*	*	1.9	4.8	7.6	16.1	24.0
2002.....	1.4	*	*	*	*	*	*	2.7	4.2	9.1	11.8	21.4
2001.....	1.4	*	*	*	*	*	*	2.4	5.5	8.5	10.3	23.8
2000.....	1.5	*	*	*	*	*	*	2.4	6.1	8.6	13.9	*
1999.....	1.4	*	*	*	*	*	0.7	2.6	6.1	8.2	12.6	*
Black, female												
2004.....	0.8	*	*	*	*	*	*	0.9	1.7	2.0	7.2	12.2
2003.....	0.9	*	*	*	*	*	*	*	1.5	4.0	6.4	18.2
2002.....	1.0	*	*	*	*	*	*	0.8	2.0	3.6	8.0	20.9
2001.....	1.1	*	*	*	*	*	*	*	2.6	3.7	7.4	24.0
2000.....	1.0	*	*	*	*	*	*	*	2.7	3.5	9.0	22.3
1999.....	1.0	*	*	*	*	*	*	1.1	1.5	3.8	9.6	16.2

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other forms of chronic ischemic heart disease (I20,I25)

All races 3/, both sexes

2004.....	99.5	*	*	*	0.2	1.6	9.6	35.1	92.5	240.3	697.1	2,279.7
2003.....	105.3	*	*	*	0.2	1.7	9.8	35.9	97.5	258.4	741.5	2,471.4
2002.....	108.0	*	*	*	0.2	1.4	9.9	36.5	101.0	269.4	768.8	2,562.5
2001.....	110.3	*	*	*	0.2	1.6	9.5	35.6	103.8	280.9	793.1	2,671.3
2000.....	113.3	*	*	*	0.2	1.4	9.5	36.4	108.1	292.4	813.6	2,822.1
1999.....	117.2	*	*	*	0.2	1.4	9.8	37.5	111.5	308.0	848.5	2,917.0

All races 3/, male

2004.....	103.3	*	*	*	0.2	2.4	14.5	54.8	137.7	336.5	926.9	2,543.6
2003.....	108.5	*	*	*	0.3	2.5	14.9	56.4	145.4	363.5	985.8	2,722.2
2002.....	110.9	*	*	*	0.2	2.2	15.4	57.5	151.5	379.3	1,021.0	2,817.4
2001.....	111.9	*	*	*	0.3	2.4	14.6	55.9	154.9	394.3	1,050.9	2,913.8
2000.....	114.8	*	*	*	0.2	2.1	14.8	57.7	161.6	413.5	1,079.4	3,112.7
1999.....	118.0	*	*	*	0.3	2.1	15.3	59.0	164.7	434.9	1,120.5	3,193.1

All races 3/, female

2004.....	95.7	*	*	*	*	0.7	4.6	16.2	50.5	159.6	542.4	2,161.0
2003.....	102.3	*	*	*	0.1	0.8	4.7	16.2	53.1	170.5	578.3	2,360.5
2002.....	105.2	*	*	*	*	0.7	4.4	16.2	54.3	178.0	601.4	2,451.9
2001.....	108.8	*	*	*	0.1	0.7	4.4	16.0	56.7	186.9	623.7	2,569.8
2000.....	111.9	*	*	*	*	0.6	4.3	16.0	58.8	192.8	640.3	2,703.8
1999.....	116.4	*	*	*	0.1	0.7	4.3	16.7	62.6	204.0	671.8	2,805.7

White, both sexes

2004.....	107.5	*	*	*	0.1	1.4	9.2	32.8	86.7	232.4	696.4	2,324.8
2003.....	113.8	*	*	*	0.2	1.5	9.4	33.5	91.3	249.1	740.0	2,518.8
2002.....	116.9	*	*	*	0.2	1.4	9.4	34.0	94.4	261.3	767.5	2,613.9
2001.....	119.1	*	*	*	0.2	1.4	8.9	33.1	96.7	271.7	790.5	2,723.5
2000.....	122.6	*	*	*	0.2	1.2	8.9	33.9	100.6	283.4	811.2	2,882.1
1999.....	126.5	*	*	*	0.1	1.3	9.0	34.7	104.0	298.0	846.7	2,973.6

White, male

2004.....	111.1	*	*	*	0.2	2.2	14.2	52.1	130.5	328.7	931.8	2,608.9
2003.....	116.6	*	*	*	0.2	2.3	14.4	53.4	138.1	353.1	989.3	2,797.1
2002.....	119.4	*	*	*	0.2	2.1	14.9	54.8	143.1	371.2	1,027.4	2,891.6
2001.....	120.1	*	*	*	0.3	2.2	13.9	52.9	146.2	384.6	1,054.9	2,992.2
2000.....	123.5	*	*	*	0.2	1.9	14.2	54.8	152.7	405.6	1,083.4	3,200.0
1999.....	126.5	*	*	*	0.2	2.0	14.4	55.5	155.2	425.5	1,126.9	3,267.1

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other forms of chronic ischemic heart disease (I20,I25)												
White, female												
2004.....	103.9	*	*	*	*	0.6	4.1	13.8	45.4	149.9	535.9	2,196.8
2003.....	111.0	*	*	*	*	0.7	4.2	13.9	47.0	160.6	571.4	2,395.5
2002.....	114.4	*	*	*	*	0.6	3.9	13.6	48.5	168.1	593.1	2,493.2
2001.....	118.1	*	*	*	*	0.6	3.9	13.6	50.2	176.4	615.0	2,611.0
2000.....	121.6	*	*	*	*	0.5	3.5	13.4	51.8	180.9	632.0	2,753.0
1999.....	126.5	*	*	*	*	0.6	3.6	14.3	55.9	191.7	663.1	2,855.7
Black, both sexes												
2004.....	78.0	*	*	*	0.4	2.9	14.2	59.2	161.0	359.8	814.7	2,019.1
2003.....	82.3	*	*	*	0.4	2.6	14.2	60.6	171.5	389.7	873.2	2,195.3
2002.....	83.2	*	*	*	*	2.2	15.1	61.9	178.1	391.4	896.1	2,240.6
2001.....	85.8	*	*	*	0.5	2.7	15.1	61.6	184.2	414.8	932.7	2,351.2
2000.....	86.6	*	*	*	*	2.5	15.3	63.1	190.2	427.8	941.3	2,387.1
1999.....	89.3	*	*	*	0.5	2.3	16.8	65.8	195.7	450.4	957.4	2,469.8
Black, male												
2004.....	81.5	*	*	*	*	4.3	19.8	86.2	234.3	489.0	1,039.8	2,068.3
2003.....	85.5	*	*	*	*	3.9	20.6	89.6	243.7	536.7	1,128.5	2,147.8
2002.....	85.7	*	*	*	*	2.9	21.7	89.5	261.4	531.5	1,113.6	2,254.6
2001.....	88.2	*	*	*	0.7	4.2	22.2	91.3	263.6	568.3	1,177.1	2,369.1
2000.....	88.0	*	*	*	*	3.6	21.6	92.1	272.7	563.5	1,175.7	2,446.0
1999.....	91.1	*	*	*	*	3.3	24.6	97.7	281.3	600.5	1,172.6	2,561.5
Black, female												
2004.....	74.7	*	*	*	*	1.5	9.1	36.1	102.0	267.6	687.4	1,999.6
2003.....	79.4	*	*	*	*	1.4	8.5	35.7	113.4	285.0	728.9	2,213.8
2002.....	81.0	*	*	*	*	1.5	9.4	38.2	111.1	291.9	772.9	2,235.2
2001.....	83.5	*	*	*	*	1.5	8.9	36.1	120.3	306.4	795.5	2,344.5
2000.....	85.3	*	*	*	*	1.5	9.7	38.1	123.7	332.3	809.4	2,365.2
1999.....	87.6	*	*	*	*	1.3	9.9	38.5	127.0	344.6	836.9	2,435.5

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Atherosclerotic cardiovascular disease, so described (I25.0)												
All races 3/, both sexes												
2004.....	21.7	*	*	*	*	0.5	3.3	13.6	32.0	58.7	129.0	400.4
2003.....	23.1	*	*	*	*	0.5	3.4	13.9	32.7	62.7	142.5	442.9
2002.....	23.6	*	*	*	*	0.5	3.4	13.4	33.1	64.7	146.2	474.5
2001.....	23.9	*	*	*	0.1	0.5	3.2	12.9	33.3	65.6	152.7	493.6
2000.....	24.7	*	*	*	*	0.4	3.0	13.2	34.0	68.3	158.1	534.6
1999.....	25.8	*	*	*	*	0.4	3.3	13.0	35.3	72.7	166.2	563.4
All races 3/, male												
2004.....	23.9	*	*	*	*	0.8	5.2	22.0	49.0	83.6	164.9	414.0
2003.....	25.2	*	*	*	*	0.9	5.2	22.7	49.9	88.9	181.1	462.1
2002.....	25.4	*	*	*	*	0.7	5.4	21.5	51.2	91.8	185.3	491.8
2001.....	25.3	*	*	*	*	0.7	5.0	21.0	51.4	92.4	193.9	502.0
2000.....	26.0	*	*	*	*	0.6	4.9	21.5	52.6	97.2	199.3	550.8
1999.....	26.8	*	*	*	*	0.6	5.3	20.9	53.7	103.0	207.0	578.6
All races 3/, female												
2004.....	19.5	*	*	*	*	0.2	1.4	5.5	16.1	37.7	104.8	394.3
2003.....	21.1	*	*	*	*	0.2	1.5	5.5	16.7	40.8	116.7	434.4
2002.....	21.9	*	*	*	*	0.2	1.3	5.5	16.2	42.1	120.2	467.1
2001.....	22.5	*	*	*	*	0.2	1.4	5.1	16.7	43.3	125.5	490.1
2000.....	23.5	*	*	*	*	0.1	1.2	5.2	16.8	44.6	131.3	528.0
1999.....	24.8	*	*	*	*	0.2	1.3	5.3	18.3	47.9	139.8	557.2
White, both sexes												
2004.....	22.1	*	*	*	*	0.4	3.2	12.3	28.7	53.2	122.2	395.6
2003.....	23.6	*	*	*	*	0.5	3.2	12.6	29.0	56.5	135.5	439.3
2002.....	24.1	*	*	*	*	0.4	3.1	12.1	29.4	58.1	138.4	470.6
2001.....	24.3	*	*	*	*	0.4	3.0	11.6	28.8	58.3	144.5	489.7
2000.....	25.1	*	*	*	*	0.3	2.8	11.8	29.4	61.2	149.2	529.5
1999.....	26.1	*	*	*	*	0.3	3.0	11.3	30.4	64.8	157.0	557.5
White, male												
2004.....	24.1	*	*	*	*	0.7	5.1	20.4	44.1	76.6	156.3	408.8
2003.....	25.3	*	*	*	*	0.8	5.0	20.8	44.8	80.0	171.9	458.7
2002.....	25.5	*	*	*	*	0.7	5.1	19.9	45.9	82.9	174.7	488.0
2001.....	25.3	*	*	*	*	0.6	4.7	19.2	45.1	82.7	184.2	498.6
2000.....	26.0	*	*	*	*	0.5	4.6	19.6	46.3	87.8	187.5	545.6
1999.....	26.6	*	*	*	*	0.5	4.9	18.6	46.9	92.5	195.2	571.2

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Atherosclerotic cardiovascular disease, so described (I25.0)												
White, female												
2004.....	20.1	*	*	*	*	0.2	1.3	4.3	14.1	33.1	99.0	389.7
2003.....	21.9	*	*	*	*	0.2	1.4	4.5	14.1	36.5	110.8	430.7
2002.....	22.7	*	*	*	*	0.1	1.1	4.5	13.8	37.0	114.0	463.0
2001.....	23.3	*	*	*	*	0.2	1.2	4.1	13.4	37.8	118.2	486.0
2000.....	24.3	*	*	*	*	*	1.0	4.2	13.5	39.0	123.9	522.9
1999.....	25.6	*	*	*	*	0.2	1.0	4.2	15.0	41.6	131.9	552.0
Black, both sexes												
2004.....	25.0	*	*	*	*	0.9	4.8	25.7	66.7	120.0	226.9	505.1
2003.....	26.0	*	*	*	*	0.9	4.9	25.7	69.7	128.1	241.6	543.2
2002.....	27.0	*	*	*	*	0.8	5.8	25.5	71.4	136.3	259.0	575.5
2001.....	27.7	*	*	*	*	0.7	5.6	25.3	77.5	143.4	266.5	594.5
2000.....	28.5	*	*	*	*	0.7	5.1	26.9	78.2	145.5	278.4	646.6
1999.....	29.9	*	*	*	*	0.7	6.3	27.5	81.4	158.6	291.2	668.6
Black, male												
2004.....	28.9	*	*	*	*	1.5	6.9	38.1	104.6	172.0	306.2	531.5
2003.....	30.5	*	*	*	*	1.5	7.7	40.4	106.0	190.3	331.4	583.9
2002.....	31.1	*	*	*	*	1.1	8.6	38.7	111.4	197.2	357.2	599.3
2001.....	31.5	*	*	*	*	0.9	8.5	39.2	117.4	208.2	351.7	609.3
2000.....	32.3	*	*	*	*	1.0	7.8	42.0	120.0	207.3	369.7	701.2
1999.....	33.8	*	*	*	*	1.0	9.6	42.8	123.3	229.3	380.0	715.9
Black, female												
2004.....	21.5	*	*	*	*	*	3.0	15.1	36.2	83.0	182.0	494.6
2003.....	21.9	*	*	*	*	*	2.6	13.1	40.5	83.9	190.9	527.4
2002.....	23.3	*	*	*	*	*	3.3	14.2	39.2	93.0	203.5	566.4
2001.....	24.3	*	*	*	*	*	3.1	13.3	45.4	97.7	218.6	589.0
2000.....	25.0	*	*	*	*	*	2.8	14.0	44.6	102.0	227.1	626.3
1999.....	26.4	*	*	*	*	*	3.4	14.4	47.8	108.8	241.5	650.9

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
All races 3/, both sexes												
2004.....	77.8	*	*	*	0.1	1.1	6.2	21.5	60.5	181.7	568.1	1,879.3
2003.....	82.2	*	*	*	0.2	1.1	6.5	22.0	64.9	195.7	599.0	2,028.5
2002.....	84.4	*	*	*	0.1	1.0	6.5	23.1	68.0	204.7	622.6	2,088.0
2001.....	86.4	*	*	*	0.2	1.1	6.2	22.7	70.5	215.3	640.5	2,177.7
2000.....	88.6	*	*	*	0.1	1.0	6.5	23.2	74.1	224.1	655.5	2,287.5
1999.....	91.4	*	*	*	0.2	1.0	6.5	24.5	76.3	235.2	682.3	2,353.6
All races 3/, male												
2004.....	79.4	*	*	*	0.2	1.7	9.3	32.8	88.7	252.9	762.0	2,129.6
2003.....	83.3	*	*	*	0.2	1.6	9.7	33.7	95.5	274.6	804.7	2,260.1
2002.....	85.5	*	*	*	0.2	1.5	10.0	35.9	100.3	287.5	835.7	2,325.6
2001.....	86.5	*	*	*	0.2	1.8	9.5	34.9	103.5	301.9	856.9	2,411.8
2000.....	88.8	*	*	*	0.2	1.5	9.9	36.2	109.0	316.3	880.1	2,561.9
1999.....	91.2	*	*	*	0.2	1.5	10.0	38.1	111.0	331.9	913.6	2,614.4
All races 3/, female												
2004.....	76.2	*	*	*	*	0.5	3.2	10.6	34.3	121.9	437.6	1,766.7
2003.....	81.1	*	*	*	0.1	0.6	3.2	10.8	36.4	129.7	461.6	1,926.2
2002.....	83.3	*	*	*	*	0.5	3.1	10.7	38.0	135.9	481.1	1,984.9
2001.....	86.2	*	*	*	0.1	0.5	3.0	11.0	40.0	143.6	498.2	2,079.8
2000.....	88.4	*	*	*	*	0.5	3.0	10.7	42.0	148.2	509.0	2,175.8
1999.....	91.6	*	*	*	0.1	0.5	3.0	11.4	44.3	156.1	532.1	2,248.4
White, both sexes												
2004.....	85.4	*	*	*	0.1	1.0	6.0	20.5	58.1	179.1	574.2	1,929.2
2003.....	90.2	*	*	*	0.1	1.0	6.2	20.9	62.2	192.6	604.6	2,079.5
2002.....	92.8	*	*	*	0.1	0.9	6.3	21.9	65.1	203.2	629.1	2,143.3
2001.....	94.8	*	*	*	0.1	1.0	5.9	21.5	67.9	213.4	646.0	2,233.8
2000.....	97.4	*	*	*	0.1	0.9	6.1	22.1	71.2	222.2	662.0	2,352.7
1999.....	100.4	*	*	*	0.1	1.0	6.1	23.4	73.5	233.2	689.7	2,416.1
White, male												
2004.....	87.0	*	*	*	0.2	1.5	9.1	31.6	86.4	252.0	775.6	2,200.2
2003.....	91.3	*	*	*	0.2	1.5	9.4	32.5	93.3	273.1	817.4	2,338.4
2002.....	93.9	*	*	*	0.1	1.4	9.8	34.9	97.3	288.3	852.7	2,403.6
2001.....	94.8	*	*	*	0.2	1.6	9.2	33.7	101.1	301.9	870.7	2,493.6
2000.....	97.5	*	*	*	0.1	1.4	9.6	35.2	106.4	317.8	895.9	2,654.4
1999.....	99.9	*	*	*	0.2	1.4	9.5	36.9	108.3	333.0	931.7	2,695.9

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)

White, female

2004.....	83.8	*	*	*	*	0.5	2.9	9.4	31.3	116.7	436.9	1,807.0
2003.....	89.1	*	*	*	*	0.5	2.9	9.4	32.9	124.0	460.7	1,964.9
2002.....	91.7	*	*	*	*	0.4	2.8	9.1	34.7	131.1	479.1	2,030.2
2001.....	94.9	*	*	*	*	0.4	2.6	9.6	36.7	138.6	496.8	2,125.0
2000.....	97.3	*	*	*	*	0.4	2.5	9.3	38.3	141.9	508.1	2,230.1
1999.....	100.8	*	*	*	*	0.5	2.6	10.1	40.9	150.1	531.1	2,303.8

Black, both sexes

2004.....	52.9	*	*	*	0.3	2.0	9.3	33.5	94.3	239.8	587.8	1,514.0
2003.....	56.3	*	*	*	0.3	1.8	9.3	34.9	101.8	261.5	631.6	1,652.1
2002.....	56.2	*	*	*	*	1.4	9.4	36.4	106.7	255.1	637.0	1,665.1
2001.....	58.0	*	*	*	0.4	2.0	9.5	36.3	106.7	271.4	666.3	1,756.7
2000.....	58.1	*	*	*	*	1.8	10.2	36.1	112.0	282.3	662.9	1,740.5
1999.....	59.3	*	*	*	0.3	1.5	10.4	38.3	114.2	291.8	666.2	1,801.2

Black, male

2004.....	52.6	*	*	*	*	2.8	12.9	48.1	129.7	317.0	733.6	1,536.8
2003.....	55.0	*	*	*	*	2.4	13.0	49.1	137.6	346.5	797.1	1,563.9
2002.....	54.5	*	*	*	*	1.8	13.1	50.8	150.1	334.3	756.5	1,655.3
2001.....	56.7	*	*	*	*	3.2	13.7	52.1	146.2	360.1	825.4	1,759.8
2000.....	55.8	*	*	*	*	2.6	13.9	50.2	152.7	356.1	806.0	1,744.8
1999.....	57.3	*	*	*	*	2.3	15.0	54.9	157.9	371.2	792.6	1,845.5

Black, female

2004.....	53.2	*	*	*	*	1.2	6.2	21.0	65.8	184.6	505.4	1,505.0
2003.....	57.5	*	*	*	*	1.2	6.0	22.6	72.9	201.1	538.0	1,686.4
2002.....	57.7	*	*	*	*	1.1	6.1	24.0	71.9	198.9	569.4	1,668.8
2001.....	59.2	*	*	*	*	1.0	5.9	22.8	75.0	208.7	576.9	1,755.5
2000.....	60.3	*	*	*	*	1.1	6.9	24.1	79.2	230.3	582.3	1,738.9
1999.....	61.2	*	*	*	*	0.8	6.4	24.1	79.1	235.8	595.4	1,784.6

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other heart diseases (I26-I51)												
All races 3/, both sexes												
2004.....	56.6	10.0	1.1	0.5	2.1	4.3	9.7	21.5	45.2	118.6	368.9	1,373.9
2003.....	58.5	10.4	1.1	0.6	2.2	4.5	10.3	21.4	48.0	126.8	382.8	1,427.2
2002.....	58.8	11.7	1.0	0.6	2.1	4.5	10.3	21.7	48.7	129.6	390.9	1,443.5
2001.....	58.6	11.1	1.4	0.6	2.0	4.4	9.9	21.6	49.5	131.4	389.5	1,475.5
2000.....	58.8	12.0	1.1	0.6	2.2	4.4	9.9	21.5	50.9	134.2	391.3	1,509.4
1999.....	59.4	12.9	1.2	0.6	2.3	4.4	10.3	21.6	52.9	138.8	398.5	1,512.9
All races 3/, male												
2004.....	50.6	10.5	1.1	0.5	2.6	5.3	12.0	27.7	57.7	142.8	420.8	1,377.7
2003.....	52.3	11.4	1.0	0.7	2.6	5.3	12.7	27.2	60.7	154.2	440.0	1,431.3
2002.....	52.4	12.1	1.0	0.6	2.7	5.6	12.8	28.2	60.8	156.5	448.2	1,448.6
2001.....	51.8	10.8	1.5	0.6	2.4	5.2	12.4	27.6	61.7	158.7	452.0	1,473.4
2000.....	52.2	12.2	1.2	0.7	2.6	5.2	12.2	27.8	64.7	164.0	451.0	1,549.1
1999.....	53.1	13.1	1.3	0.7	2.7	5.3	12.9	27.7	67.5	172.7	459.7	1,548.9
All races 3/, female												
2004.....	62.4	9.4	1.1	0.6	1.5	3.3	7.4	15.5	33.6	98.4	334.0	1,372.1
2003.....	64.5	9.3	1.2	0.5	1.7	3.7	7.9	15.7	36.3	103.8	344.5	1,425.4
2002.....	65.0	11.4	1.0	0.5	1.4	3.4	7.8	15.5	37.5	107.2	352.8	1,441.3
2001.....	65.2	11.5	1.3	0.6	1.5	3.6	7.5	15.9	38.3	108.7	348.4	1,476.4
2000.....	65.1	11.8	0.9	0.5	1.8	3.5	7.5	15.3	38.2	109.6	352.3	1,493.2
1999.....	65.5	12.7	1.0	0.6	1.8	3.6	7.7	15.6	39.4	111.1	358.7	1,498.5
White, both sexes												
2004.....	60.4	8.7	0.9	0.5	1.8	3.5	8.1	18.2	40.6	113.4	368.6	1,408.8
2003.....	62.2	8.2	0.9	0.5	1.8	3.6	8.6	18.1	43.1	120.3	383.4	1,461.4
2002.....	62.6	10.0	0.9	0.5	1.7	3.6	8.7	18.4	43.9	124.2	390.3	1,475.5
2001.....	62.3	9.3	1.1	0.5	1.6	3.6	8.1	18.4	44.6	125.6	387.6	1,509.6
2000.....	62.5	10.6	1.0	0.5	1.7	3.6	8.1	18.2	46.1	128.1	389.6	1,541.6
1999.....	62.9	11.0	1.0	0.5	1.9	3.6	8.4	18.1	47.2	132.2	395.1	1,543.3
White, male												
2004.....	53.0	9.1	0.9	0.5	2.2	4.3	10.3	23.9	52.1	137.0	419.2	1,407.9
2003.....	54.7	8.8	0.8	0.6	2.2	4.4	10.8	23.3	54.7	147.2	441.1	1,467.6
2002.....	54.9	10.5	1.0	0.6	2.3	4.5	11.1	24.3	55.3	150.4	448.3	1,482.1
2001.....	54.1	9.6	1.2	0.5	2.0	4.2	10.4	24.0	55.9	152.2	448.7	1,503.4
2000.....	54.6	10.8	1.2	0.6	2.1	4.3	10.3	23.9	59.2	157.4	449.5	1,580.5
1999.....	55.1	11.4	1.1	0.6	2.2	4.3	10.9	23.7	60.6	165.1	456.1	1,571.5

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other heart diseases (I26-I51)												
White, female												
2004.....	67.6	8.3	0.8	0.5	1.3	2.6	5.9	12.7	29.8	93.1	334.0	1,409.2
2003.....	69.6	7.6	1.0	0.4	1.5	2.8	6.3	13.0	32.1	97.3	344.4	1,458.6
2002.....	70.2	9.6	0.8	0.5	1.1	2.6	6.4	12.6	33.1	101.9	351.4	1,472.6
2001.....	70.4	8.9	1.1	0.5	1.3	2.9	5.7	12.8	34.0	103.1	347.1	1,512.2
2000.....	70.2	10.4	0.7	0.4	1.4	2.8	6.0	12.5	33.9	103.4	350.1	1,525.8
1999.....	70.5	10.7	0.8	0.5	1.6	2.8	6.0	12.6	34.5	104.7	355.1	1,531.9
Black, both sexes												
2004.....	50.8	14.8	2.3	1.1	3.7	10.0	22.2	50.0	96.1	195.5	450.2	1,180.7
2003.....	52.5	19.6	2.0	1.0	4.4	10.9	23.6	49.9	102.1	215.7	449.2	1,238.1
2002.....	52.5	20.2	1.5	0.8	4.1	11.3	22.8	51.4	102.0	207.6	469.1	1,270.4
2001.....	52.8	20.1	2.9	1.0	4.2	10.5	23.7	51.1	103.5	213.0	480.6	1,277.7
2000.....	52.7	19.0	1.9	1.1	4.7	10.3	23.0	51.6	102.6	220.2	477.9	1,306.0
1999.....	54.4	22.3	2.0	1.0	4.5	10.5	24.7	52.4	113.3	229.9	495.2	1,302.8
Black, male												
2004.....	50.4	15.7	2.3	1.0	4.7	12.2	26.7	63.6	125.6	237.6	540.9	1,224.9
2003.....	51.5	23.2	2.2	1.1	5.7	12.4	28.6	63.8	132.3	264.0	523.5	1,220.9
2002.....	50.8	20.9	*	0.9	5.1	14.2	27.4	65.9	127.9	255.2	537.3	1,238.3
2001.....	51.7	17.7	3.2	0.9	5.3	12.8	28.3	62.5	132.1	258.5	585.9	1,349.1
2000.....	51.5	18.9	1.9	1.2	5.4	12.2	28.1	66.5	130.5	266.0	558.0	1,402.3
1999.....	54.3	21.8	2.1	1.1	5.8	12.7	30.1	67.3	147.3	287.8	580.2	1,454.1
Black, female												
2004.....	51.1	13.9	2.3	1.1	2.7	8.0	18.3	38.4	72.3	165.4	398.9	1,163.2
2003.....	53.4	15.8	1.8	0.8	3.2	9.4	19.1	38.0	77.8	181.3	407.1	1,244.8
2002.....	54.0	19.4	1.8	0.8	3.1	8.7	18.7	39.0	81.1	173.8	430.6	1,282.7
2001.....	53.8	22.7	2.5	1.0	3.0	8.5	19.6	41.3	80.5	180.9	421.5	1,251.2
2000.....	53.9	19.2	2.0	1.0	4.1	8.7	18.6	38.7	80.1	187.9	432.8	1,270.2
1999.....	54.5	22.8	1.8	0.8	3.3	8.5	19.9	39.7	86.1	189.1	447.7	1,246.2

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
All races 3/, both sexes												
2004.....	4.5	3.8	0.2	0.1	0.3	0.8	1.7	3.3	5.8	13.4	27.5	54.8
2003.....	4.8	3.3	0.2	0.1	0.3	0.9	1.9	3.3	6.5	14.4	30.4	58.0
2002.....	4.8	4.6	0.2	0.1	0.3	0.8	1.7	3.4	6.9	14.9	29.8	56.3
2001.....	4.7	4.2	0.1	0.1	0.3	0.9	1.7	3.4	6.9	14.6	28.5	57.5
2000.....	4.6	4.9	0.2	0.1	0.3	0.9	1.7	3.5	6.7	14.5	28.3	57.4
1999.....	4.7	5.7	*	0.1	0.4	0.9	1.8	3.3	6.7	14.8	30.0	59.6
All races 3/, male												
2004.....	3.6	4.1	*	*	0.2	0.6	1.4	3.1	5.8	12.5	25.2	48.3
2003.....	3.9	3.7	*	*	0.2	0.6	1.6	3.3	6.2	14.2	29.9	54.4
2002.....	3.8	4.7	*	*	0.2	0.7	1.4	3.1	6.5	14.1	27.7	52.1
2001.....	3.8	3.9	*	*	0.2	0.6	1.5	3.0	6.5	14.5	27.9	55.6
2000.....	3.8	5.1	*	*	0.2	0.7	1.4	3.3	6.4	13.9	28.3	54.8
1999.....	3.9	5.9	*	*	0.3	0.6	1.5	3.0	6.8	15.1	29.1	60.3
All races 3/, female												
2004.....	5.3	3.5	*	*	0.4	0.9	1.9	3.4	5.7	14.3	29.1	57.7
2003.....	5.7	2.9	*	*	0.5	1.1	2.2	3.4	6.9	14.6	30.8	59.6
2002.....	5.7	4.5	*	*	0.4	1.0	1.9	3.6	7.2	15.6	31.2	58.1
2001.....	5.5	4.6	*	*	0.4	1.1	2.0	3.7	7.3	14.6	28.9	58.2
2000.....	5.5	4.6	*	*	0.5	1.0	2.1	3.7	7.0	15.0	28.4	58.5
1999.....	5.5	5.6	*	*	0.5	1.1	2.0	3.6	6.6	14.6	30.5	59.2
White, both sexes												
2004.....	4.5	3.2	*	0.1	0.2	0.6	1.4	2.9	5.3	12.6	26.6	55.3
2003.....	4.8	2.5	*	0.1	0.3	0.7	1.5	3.0	6.0	13.3	30.1	57.8
2002.....	4.8	3.9	*	0.1	0.2	0.6	1.5	3.0	6.3	14.2	29.1	56.1
2001.....	4.7	3.6	*	0.1	0.3	0.7	1.5	3.0	6.4	13.8	27.9	56.8
2000.....	4.7	4.1	0.2	0.1	0.3	0.7	1.5	3.1	6.3	13.8	27.7	56.4
1999.....	4.7	5.2	*	0.1	0.4	0.7	1.5	2.9	6.0	13.7	28.9	59.2
White, male												
2004.....	3.6	3.4	*	*	0.2	0.5	1.2	2.9	5.4	12.0	23.8	48.0
2003.....	3.9	2.9	*	*	0.2	0.5	1.3	2.9	5.8	13.3	29.7	54.0
2002.....	3.9	4.1	*	*	0.2	0.5	1.3	2.9	6.1	13.7	27.2	51.9
2001.....	3.8	3.4	*	*	0.2	0.5	1.3	2.8	5.9	14.0	27.1	53.6
2000.....	3.8	4.3	*	*	0.2	0.6	1.2	3.0	6.1	13.6	27.7	52.9
1999.....	3.9	5.4	*	*	0.2	0.5	1.4	2.8	6.2	14.1	28.1	60.3

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
White, female												
2004.....	5.4	3.0	*	*	0.3	0.8	1.6	2.9	5.2	13.2	28.5	58.6
2003.....	5.7	2.0	*	*	0.4	1.0	1.8	3.1	6.3	13.2	30.4	59.5
2002.....	5.8	3.7	*	*	0.3	0.7	1.7	3.1	6.5	14.7	30.4	58.0
2001.....	5.6	3.9	*	*	0.3	0.9	1.6	3.2	6.8	13.7	28.4	58.1
2000.....	5.5	3.9	*	*	0.4	0.8	1.7	3.2	6.5	14.0	27.7	57.8
1999.....	5.5	4.9	*	*	0.5	0.8	1.6	3.1	5.7	13.5	29.5	58.8
Black, both sexes												
2004.....	5.6	5.9	*	*	0.7	1.9	3.8	6.7	11.5	24.5	44.4	62.0
2003.....	6.1	6.5	*	*	0.7	2.0	4.5	6.7	13.1	29.3	42.2	72.9
2002.....	5.9	8.2	*	*	0.6	2.4	3.5	6.8	13.8	26.0	45.9	68.6
2001.....	5.8	6.8	*	*	0.6	2.1	3.8	6.8	13.7	25.9	41.9	77.7
2000.....	5.7	8.0	*	*	0.7	2.1	4.0	7.4	12.2	24.7	42.2	76.6
1999.....	6.2	9.4	*	*	0.7	2.2	4.0	7.3	14.8	28.8	49.1	72.1
Black, male												
2004.....	4.6	6.5	*	*	*	1.5	3.3	5.9	11.5	20.8	49.8	64.6
2003.....	5.2	7.1	*	*	0.7	1.6	4.1	6.9	12.7	28.0	42.0	76.1
2002.....	4.5	9.0	*	*	*	2.0	2.9	5.6	11.8	23.1	42.7	65.4
2001.....	4.8	6.9	*	*	*	1.6	3.3	5.4	13.8	24.3	44.1	95.3
2000.....	4.6	7.7	*	*	*	1.7	3.2	6.4	12.2	21.2	43.0	92.3
1999.....	5.1	8.8	*	*	*	1.9	3.1	6.2	14.1	30.1	50.1	71.5
Black, female												
2004.....	6.5	*	*	*	0.8	2.2	4.3	7.4	11.4	27.2	41.4	61.0
2003.....	6.9	*	*	*	0.8	2.3	4.9	6.6	13.4	30.2	42.3	71.7
2002.....	7.2	7.3	*	*	0.7	2.7	4.1	7.9	15.4	28.1	47.7	69.8
2001.....	6.8	6.6	*	*	*	2.7	4.4	8.1	13.6	27.0	40.7	71.1
2000.....	6.8	8.3	*	*	1.1	2.4	4.6	8.1	12.3	27.1	41.8	70.8
1999.....	7.2	10.1	*	*	0.7	2.5	4.8	8.1	15.3	27.8	48.5	72.3

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pulmonary embolism (I26)												
All races 3/, both sexes												
2004.....	2.8	*	*	*	0.2	0.5	1.1	2.2	3.6	8.2	17.1	33.4
2003.....	3.0	*	*	*	0.2	0.6	1.3	2.3	4.2	8.7	18.9	35.5
2002.....	3.0	*	*	*	0.2	0.5	1.2	2.2	4.5	9.5	18.5	35.4
2001.....	3.0	*	*	*	0.2	0.6	1.2	2.3	4.7	9.5	18.4	37.8
2000.....	3.0	*	*	*	0.2	0.5	1.2	2.3	4.5	9.3	18.4	38.4
1999.....	3.2	*	*	*	0.2	0.6	1.2	2.3	4.7	10.1	21.1	42.0
All races 3/, male												
2004.....	2.3	*	*	*	0.1	0.5	1.0	2.1	4.0	8.4	16.8	30.6
2003.....	2.6	*	*	*	0.2	0.4	1.1	2.4	4.5	9.4	19.6	34.1
2002.....	2.5	*	*	*	0.1	0.5	1.0	2.2	4.5	9.4	17.5	34.1
2001.....	2.6	*	*	*	0.1	0.5	1.1	2.1	4.5	10.0	19.0	38.0
2000.....	2.6	*	*	*	0.1	0.5	1.0	2.4	4.6	9.5	19.3	37.5
1999.....	2.7	*	*	*	0.2	0.5	1.1	2.2	5.0	10.9	20.4	42.0
All races 3/, female												
2004.....	3.2	*	*	*	0.2	0.6	1.3	2.2	3.3	8.0	17.3	34.7
2003.....	3.4	*	*	*	0.3	0.7	1.4	2.2	4.0	8.2	18.4	36.0
2002.....	3.5	*	*	*	0.2	0.6	1.3	2.2	4.5	9.5	19.2	36.0
2001.....	3.5	*	*	*	0.2	0.7	1.3	2.5	4.9	9.1	18.0	37.7
2000.....	3.4	*	*	*	0.3	0.6	1.3	2.2	4.4	9.1	17.8	38.8
1999.....	3.7	*	*	*	0.3	0.7	1.3	2.4	4.4	9.6	21.5	42.1
White, both sexes												
2004.....	2.8	*	*	*	0.2	0.5	0.9	1.9	3.4	7.6	16.5	33.4
2003.....	3.0	*	*	*	0.2	0.4	1.0	2.1	3.9	8.0	18.6	34.8
2002.....	3.0	*	*	*	0.2	0.4	1.0	2.0	4.2	9.0	17.9	34.8
2001.....	3.1	*	*	*	0.2	0.5	1.0	2.0	4.4	9.1	17.9	37.0
2000.....	3.0	*	*	*	0.2	0.5	1.0	2.1	4.2	8.9	17.8	37.4
1999.....	3.2	*	*	*	0.2	0.4	1.0	2.0	4.2	9.4	20.3	41.7
White, male												
2004.....	2.3	*	*	*	0.1	0.4	0.8	2.0	3.7	8.0	15.7	30.0
2003.....	2.6	*	*	*	*	0.3	0.9	2.1	4.1	8.8	19.1	33.2
2002.....	2.5	*	*	*	*	0.3	0.9	2.1	4.3	9.1	17.0	33.3
2001.....	2.6	*	*	*	*	0.3	0.9	2.0	4.1	9.7	18.3	36.1
2000.....	2.6	*	*	*	*	0.4	0.8	2.2	4.3	9.3	18.6	35.8
1999.....	2.7	*	*	*	*	0.3	1.0	2.0	4.6	10.1	19.4	41.3

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pulmonary embolism (I26)												
White, female												
2004.....	3.2	*	*	*	0.2	0.6	1.1	1.9	3.1	7.3	17.1	34.9
2003.....	3.4	*	*	*	0.2	0.6	1.2	2.0	3.7	7.3	18.2	35.6
2002.....	3.5	*	*	*	0.2	0.5	1.2	2.0	4.1	8.9	18.5	35.5
2001.....	3.6	*	*	*	0.2	0.6	1.0	2.1	4.7	8.6	17.6	37.4
2000.....	3.4	*	*	*	0.2	0.5	1.1	1.9	4.2	8.5	17.2	38.1
1999.....	3.7	*	*	*	0.3	0.5	1.0	2.1	3.8	8.8	20.8	41.8
Black, both sexes												
2004.....	3.6	*	*	*	0.4	1.2	2.7	4.6	6.8	15.7	28.8	42.6
2003.....	4.0	*	*	*	0.6	1.5	3.1	4.6	8.3	19.0	27.7	51.1
2002.....	3.8	*	*	*	0.4	1.5	2.3	4.5	8.8	16.6	30.6	49.0
2001.....	3.8	*	*	*	*	1.6	2.7	4.7	8.8	16.5	28.5	55.9
2000.....	3.9	*	*	*	0.6	1.2	2.8	5.1	8.3	16.4	30.4	56.6
1999.....	4.3	*	*	*	0.5	1.6	2.6	5.1	10.3	20.5	36.2	53.2
Black, male												
2004.....	3.2	*	*	*	*	1.2	2.4	4.0	7.9	15.0	37.0	45.7
2003.....	3.6	*	*	*	*	1.3	2.8	5.1	9.3	19.3	32.2	57.3
2002.....	3.0	*	*	*	*	1.5	1.9	4.3	8.6	15.1	29.0	51.5
2001.....	3.3	*	*	*	*	1.3	2.5	3.7	10.0	15.9	33.0	72.6
2000.....	3.3	*	*	*	*	0.9	2.5	5.3	9.4	14.6	33.3	71.5
1999.....	3.8	*	*	*	*	1.5	2.0	4.7	10.8	23.2	40.6	60.9
Black, female												
2004.....	3.9	*	*	*	*	1.1	2.9	5.1	6.0	16.2	24.1	41.3
2003.....	4.3	*	*	*	0.6	1.6	3.4	4.2	7.4	18.7	25.2	48.6
2002.....	4.4	*	*	*	*	1.5	2.6	4.8	8.9	17.6	31.5	48.1
2001.....	4.3	*	*	*	*	1.8	3.0	5.5	7.8	16.9	25.9	49.7
2000.....	4.4	*	*	*	0.8	1.5	3.0	5.0	7.3	17.7	28.8	51.1
1999.....	4.8	*	*	*	*	1.8	3.1	5.4	9.8	18.6	33.7	50.4

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)

All races 3/, both sexes

2004.....	1.7	3.7	0.1	*	0.1	0.2	0.5	1.1	2.1	5.3	10.4	21.4
2003.....	1.8	3.2	0.1	0.1	0.1	0.3	0.6	1.0	2.3	5.6	11.6	22.6
2002.....	1.8	4.6	0.2	0.1	0.1	0.3	0.5	1.1	2.4	5.5	11.3	20.9
2001.....	1.6	4.2	0.1	0.1	0.1	0.3	0.6	1.1	2.2	5.1	10.1	19.6
2000.....	1.6	4.7	0.2	0.1	0.1	0.3	0.6	1.1	2.2	5.2	9.9	19.0
1999.....	1.5	5.7	*	0.1	0.2	0.3	0.6	1.0	2.0	4.6	8.9	17.5

All races 3/, male

2004.....	1.2	4.0	*	*	*	0.2	0.5	1.0	1.8	4.1	8.4	17.7
2003.....	1.3	3.6	*	*	*	0.2	0.5	0.9	1.8	4.8	10.3	20.3
2002.....	1.3	4.7	*	*	*	0.2	0.4	0.9	1.9	4.7	10.3	18.0
2001.....	1.2	3.9	*	*	0.1	0.2	0.4	0.9	2.0	4.5	8.9	17.6
2000.....	1.2	4.9	*	*	*	0.2	0.5	0.9	1.8	4.4	9.0	17.3
1999.....	1.2	5.9	*	*	0.1	0.2	0.5	0.8	1.7	4.2	8.7	18.4

All races 3/, female

2004.....	2.2	3.5	*	*	0.2	0.3	0.6	1.1	2.4	6.3	11.8	23.1
2003.....	2.3	2.8	*	*	0.2	0.4	0.8	1.2	2.8	6.4	12.4	23.6
2002.....	2.2	4.4	*	*	0.1	0.3	0.6	1.3	2.8	6.1	12.0	22.1
2001.....	2.0	4.5	*	*	0.1	0.4	0.7	1.3	2.5	5.5	11.0	20.5
2000.....	2.1	4.5	*	*	0.2	0.4	0.7	1.4	2.5	5.8	10.6	19.7
1999.....	1.8	5.5	*	*	0.2	0.4	0.7	1.2	2.2	5.0	9.0	17.2

White, both sexes

2004.....	1.7	3.2	*	*	0.1	0.2	0.5	1.0	1.9	5.0	10.1	21.9
2003.....	1.8	2.5	*	*	0.1	0.3	0.5	0.9	2.1	5.3	11.5	23.0
2002.....	1.8	3.8	*	*	0.1	0.2	0.4	1.0	2.1	5.2	11.2	21.3
2001.....	1.7	3.6	*	*	0.1	0.3	0.5	1.0	2.0	4.7	10.0	19.7
2000.....	1.7	4.0	0.2	*	0.1	0.2	0.5	1.0	2.0	5.0	10.0	19.0
1999.....	1.5	5.1	*	0.1	0.2	0.2	0.5	0.9	1.8	4.3	8.7	17.6

White, male

2004.....	1.2	3.3	*	*	*	0.1	0.4	0.9	1.6	4.0	8.2	18.0
2003.....	1.4	2.9	*	*	*	0.1	0.4	0.8	1.7	4.5	10.6	20.8
2002.....	1.3	4.0	*	*	*	0.2	0.4	0.9	1.8	4.6	10.2	18.6
2001.....	1.2	3.4	*	*	*	0.2	0.4	0.8	1.8	4.3	8.8	17.4
2000.....	1.2	4.2	*	*	*	0.2	0.4	0.8	1.8	4.3	9.1	17.1
1999.....	1.2	5.4	*	*	0.1	0.1	0.4	0.8	1.6	4.0	8.7	19.0

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)

White, female

2004.....	2.2	3.0	*	*	*	0.2	0.5	1.0	2.1	5.9	11.5	23.7
2003.....	2.3	2.0	*	*	0.2	0.4	0.6	1.0	2.5	5.9	12.2	23.9
2002.....	2.2	3.6	*	*	*	0.2	0.5	1.1	2.4	5.8	11.9	22.5
2001.....	2.1	3.8	*	*	*	0.3	0.6	1.1	2.1	5.1	10.8	20.7
2000.....	2.1	3.8	*	*	0.2	0.3	0.6	1.2	2.3	5.5	10.5	19.8
1999.....	1.8	4.8	*	*	0.2	0.3	0.6	1.0	1.9	4.6	8.7	17.0

Black, both sexes

2004.....	2.0	5.7	*	*	0.3	0.7	1.2	2.1	4.7	8.8	15.7	19.5
2003.....	2.1	6.2	*	*	*	0.5	1.4	2.1	4.8	10.4	14.5	21.9
2002.....	2.2	8.2	*	*	*	0.8	1.3	2.3	5.0	9.4	15.3	19.6
2001.....	2.0	6.8	*	*	*	0.6	1.1	2.1	4.9	9.4	13.4	21.8
2000.....	1.9	7.6	*	*	*	0.8	1.2	2.2	4.0	8.2	11.8	20.0
1999.....	1.9	9.3	*	*	*	0.6	1.5	2.2	4.5	8.3	12.9	18.8

Black, male

2004.....	1.4	6.5	*	*	*	*	0.9	1.8	3.6	5.8	12.8	*
2003.....	1.5	6.8	*	*	*	*	1.2	1.8	3.4	8.7	9.9	*
2002.....	1.5	9.0	*	*	*	*	1.0	1.3	3.2	8.0	13.8	*
2001.....	1.5	6.9	*	*	*	*	0.8	1.6	3.8	8.4	11.2	22.7
2000.....	1.3	7.4	*	*	*	*	0.7	1.2	2.8	6.6	9.7	*
1999.....	1.3	8.8	*	*	*	*	1.2	1.5	3.3	6.9	9.5	*

Black, female

2004.....	2.6	*	*	*	*	1.1	1.4	2.3	5.5	11.0	17.3	19.7
2003.....	2.6	*	*	*	*	*	1.6	2.4	5.9	11.5	17.1	23.1
2002.....	2.8	7.3	*	*	*	1.1	1.5	3.1	6.4	10.4	16.2	21.8
2001.....	2.5	6.6	*	*	*	0.8	1.4	2.6	5.8	10.1	14.7	21.5
2000.....	2.4	7.9	*	*	*	0.9	1.6	3.1	5.0	9.4	13.0	19.7
1999.....	2.5	9.7	*	*	*	0.8	1.7	2.7	5.4	9.2	14.8	21.9

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other non-pulmonary forms of heart disease (I30-I51)												
All races 3/, both sexes												
2004.....	52.2	6.2	0.9	0.5	1.7	3.5	8.1	18.2	39.4	105.2	341.4	1,319.1
2003.....	53.6	7.1	0.9	0.5	1.8	3.7	8.4	18.0	41.5	112.4	352.4	1,369.2
2002.....	54.0	7.1	0.8	0.5	1.8	3.7	8.6	18.4	41.8	114.6	361.1	1,387.2
2001.....	53.9	6.9	1.3	0.5	1.7	3.5	8.2	18.3	42.6	116.8	361.0	1,418.1
2000.....	54.1	7.1	0.9	0.5	1.8	3.5	8.1	18.0	44.2	119.7	362.9	1,452.0
1999.....	54.7	7.2	1.0	0.5	1.9	3.6	8.5	18.2	46.2	124.0	368.5	1,453.4
All races 3/, male												
2004.....	47.1	6.4	1.0	0.5	2.4	4.7	10.6	24.5	51.8	130.3	395.6	1,329.4
2003.....	48.3	7.7	0.8	0.6	2.4	4.7	11.1	24.0	54.4	140.0	410.2	1,376.9
2002.....	48.6	7.3	0.9	0.5	2.5	5.0	11.4	25.1	54.3	142.4	420.4	1,396.5
2001.....	48.0	6.9	1.3	0.5	2.2	4.5	10.9	24.6	55.3	144.2	424.1	1,417.8
2000.....	48.5	7.1	1.0	0.6	2.4	4.5	10.8	24.5	58.3	150.2	422.7	1,494.4
1999.....	49.1	7.3	1.1	0.6	2.4	4.6	11.4	24.7	60.8	157.6	430.5	1,488.5
All races 3/, female												
2004.....	57.1	5.9	0.9	0.5	1.1	2.3	5.5	12.2	27.9	84.1	304.9	1,314.4
2003.....	58.8	6.4	1.0	0.4	1.3	2.6	5.7	12.3	29.4	89.2	313.7	1,365.8
2002.....	59.3	6.9	0.8	0.4	1.1	2.4	5.9	11.9	30.2	91.6	321.7	1,383.1
2001.....	59.7	7.0	1.2	0.6	1.1	2.4	5.5	12.2	30.9	94.1	319.5	1,418.2
2000.....	59.6	7.2	0.7	0.4	1.3	2.5	5.5	11.7	31.3	94.7	323.9	1,434.8
1999.....	60.0	7.1	0.9	0.5	1.3	2.5	5.7	12.0	32.8	96.6	328.2	1,439.2
White, both sexes												
2004.....	55.9	5.4	0.7	0.4	1.5	2.8	6.7	15.3	35.3	100.7	342.0	1,353.6
2003.....	57.4	5.7	0.8	0.4	1.5	2.9	7.0	15.1	37.1	107.0	353.3	1,403.6
2002.....	57.8	6.2	0.8	0.4	1.5	3.0	7.2	15.4	37.5	109.9	361.2	1,419.3
2001.....	57.6	5.7	1.0	0.5	1.4	2.8	6.6	15.4	38.3	111.7	359.7	1,452.9
2000.....	57.8	6.5	0.8	0.4	1.5	2.9	6.7	15.1	39.9	114.2	361.9	1,485.2
1999.....	58.1	5.9	0.9	0.5	1.6	2.9	7.0	15.2	41.2	118.4	366.1	1,484.0
White, male												
2004.....	49.4	5.7	0.8	0.4	2.1	3.8	9.1	21.0	46.7	125.1	395.4	1,359.9
2003.....	50.8	5.8	0.7	0.5	2.0	3.9	9.5	20.4	49.0	133.9	411.4	1,413.6
2002.....	51.1	6.4	0.9	0.5	2.1	3.9	9.8	21.3	49.1	136.7	421.1	1,430.2
2001.....	50.3	6.3	1.0	0.4	1.8	3.7	9.1	21.2	50.0	138.2	421.6	1,449.9
2000.....	50.8	6.4	1.0	0.5	1.9	3.7	9.0	20.9	53.1	143.8	421.8	1,527.6
1999.....	51.1	6.0	1.0	0.5	2.0	3.8	9.5	21.0	54.4	151.0	427.9	1,511.1

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other non-pulmonary forms of heart disease (I30-I51)

White, female

2004.....	62.2	5.2	0.7	0.4	0.9	1.8	4.4	9.7	24.5	79.9	305.5	1,350.7
2003.....	63.9	5.6	0.9	0.3	1.1	1.9	4.5	9.9	25.8	84.0	314.0	1,399.2
2002.....	64.4	5.9	0.7	0.4	0.8	1.9	4.7	9.5	26.6	87.2	321.0	1,414.6
2001.....	64.8	5.0	1.0	0.5	0.9	1.9	4.1	9.6	27.2	89.3	318.7	1,454.1
2000.....	64.6	6.6	0.6	0.3	1.0	2.0	4.3	9.3	27.4	89.4	322.4	1,468.0
1999.....	64.9	5.8	0.8	0.4	1.1	2.0	4.4	9.6	28.8	91.3	325.6	1,473.2

Black, both sexes

2004.....	45.2	8.9	2.1	1.0	3.1	8.1	18.4	43.4	84.6	170.9	405.7	1,118.7
2003.....	46.4	13.0	1.7	0.9	3.7	8.9	19.1	43.2	89.0	186.3	407.0	1,165.2
2002.....	46.6	12.0	1.1	0.7	3.5	8.9	19.2	44.6	88.2	181.6	423.2	1,201.8
2001.....	47.0	13.4	2.7	0.9	3.6	8.4	19.9	44.3	89.8	187.1	438.7	1,200.0
2000.....	47.0	11.1	1.6	1.0	4.0	8.3	19.1	44.2	90.3	195.5	435.7	1,229.4
1999.....	48.2	12.9	1.8	0.9	3.9	8.3	20.6	45.2	98.6	201.1	446.1	1,230.8

Black, male

2004.....	45.8	9.2	2.2	0.9	4.2	10.6	23.4	57.7	114.1	216.9	491.1	1,160.3
2003.....	46.4	16.1	1.8	1.0	5.0	10.8	24.6	56.9	119.6	236.0	481.5	1,144.8
2002.....	46.3	11.9	*	0.8	4.6	12.2	24.5	60.3	116.1	232.1	494.6	1,172.9
2001.....	46.9	10.8	3.0	0.8	4.7	11.3	25.1	57.2	118.3	234.1	541.8	1,253.8
2000.....	46.9	11.2	*	1.0	5.1	10.5	24.8	60.1	118.3	244.8	515.0	1,310.1
1999.....	49.2	13.0	2.0	1.1	5.2	10.8	26.9	61.0	133.1	257.7	530.0	1,382.7

Black, female

2004.....	44.6	8.6	2.0	1.1	1.9	5.8	13.9	31.0	60.9	138.2	357.5	1,102.2
2003.....	46.5	9.9	1.6	0.8	2.4	7.1	14.2	31.5	64.4	151.0	364.9	1,173.1
2002.....	46.8	12.1	*	0.7	2.3	6.0	14.6	31.1	65.8	145.7	382.8	1,212.9
2001.....	47.1	16.1	2.3	0.9	2.5	5.8	15.3	33.2	66.9	153.9	380.9	1,180.0
2000.....	47.1	10.9	1.6	0.9	2.9	6.3	13.9	30.6	67.8	160.8	391.0	1,199.3
1999.....	47.3	12.8	1.6	0.8	2.6	6.0	15.1	31.6	70.8	161.3	399.2	1,174.0

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute and subacute endocarditis (I33)												
All races 3/, both sexes												
2004.....	0.4	*	*	*	*	0.1	0.2	0.5	0.6	1.3	2.5	3.2
2003.....	0.4	*	*	*	*	0.1	0.3	0.4	0.7	1.3	2.5	3.0
2002.....	0.4	*	*	*	*	0.1	0.2	0.4	0.6	1.3	2.4	2.6
2001.....	0.4	*	*	*	*	0.1	0.3	0.5	0.6	1.3	2.5	2.5
2000.....	0.4	*	*	*	*	0.1	0.3	0.5	0.7	1.2	2.4	2.8
1999.....	0.4	*	*	*	*	0.1	0.3	0.4	0.7	1.3	2.4	2.9
All races 3/, male												
2004.....	0.5	*	*	*	*	0.1	0.2	0.6	0.7	1.4	3.1	4.8
2003.....	0.5	*	*	*	*	*	0.3	0.6	0.9	1.4	3.2	4.1
2002.....	0.5	*	*	*	*	0.2	0.3	0.6	0.7	1.6	3.0	3.5
2001.....	0.5	*	*	*	*	*	0.3	0.7	0.8	1.6	3.5	3.8
2000.....	0.5	*	*	*	*	0.1	0.3	0.6	0.8	1.5	3.5	4.2
1999.....	0.5	*	*	*	*	0.1	0.3	0.6	0.9	1.4	3.2	4.7
All races 3/, female												
2004.....	0.4	*	*	*	*	*	0.1	0.3	0.4	1.1	2.0	2.5
2003.....	0.4	*	*	*	*	*	0.2	0.3	0.6	1.1	2.0	2.5
2002.....	0.3	*	*	*	*	*	0.1	0.3	0.6	1.1	2.0	2.2
2001.....	0.4	*	*	*	*	*	0.2	0.3	0.5	1.1	1.8	2.0
2000.....	0.4	*	*	*	*	*	0.2	0.3	0.5	1.0	1.8	2.3
1999.....	0.3	*	*	*	*	*	0.2	0.2	0.5	1.2	1.8	2.1
White, both sexes												
2004.....	0.4	*	*	*	*	0.1	0.2	0.3	0.5	1.2	2.4	3.2
2003.....	0.4	*	*	*	*	0.1	0.2	0.3	0.6	1.2	2.5	3.1
2002.....	0.4	*	*	*	*	0.1	0.2	0.3	0.5	1.2	2.5	2.7
2001.....	0.4	*	*	*	*	0.1	0.2	0.4	0.5	1.2	2.4	2.6
2000.....	0.4	*	*	*	*	0.1	0.2	0.4	0.6	1.2	2.4	2.7
1999.....	0.4	*	*	*	*	0.1	0.2	0.3	0.6	1.2	2.4	3.0
White, male												
2004.....	0.5	*	*	*	*	*	0.3	0.4	0.7	1.4	3.1	4.9
2003.....	0.5	*	*	*	*	*	0.3	0.5	0.8	1.3	3.3	4.4
2002.....	0.5	*	*	*	*	0.2	0.2	0.5	0.6	1.5	3.2	3.6
2001.....	0.5	*	*	*	*	*	0.2	0.5	0.7	1.3	3.5	3.9
2000.....	0.5	*	*	*	*	*	0.3	0.5	0.8	1.4	3.5	4.2
1999.....	0.5	*	*	*	*	*	0.2	0.4	0.7	1.4	3.4	5.1

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute and subacute endocarditis (I33)												
White, female												
2004.....	0.3	*	*	*	*	*	*	0.2	0.4	1.1	1.9	2.4
2003.....	0.4	*	*	*	*	*	0.2	0.2	0.5	1.0	1.9	2.6
2002.....	0.3	*	*	*	*	*	0.1	0.2	0.5	1.0	2.0	2.3
2001.....	0.3	*	*	*	*	*	0.2	0.2	0.4	1.1	1.7	2.1
2000.....	0.3	*	*	*	*	*	0.2	0.2	0.4	1.0	1.7	2.1
1999.....	0.3	*	*	*	*	*	0.1	0.2	0.4	1.1	1.8	2.1
Black, both sexes												
2004.....	0.5	*	*	*	*	*	*	1.4	1.2	2.1	2.7	*
2003.....	0.6	*	*	*	*	*	0.6	1.3	1.9	2.2	3.0	*
2002.....	0.5	*	*	*	*	*	0.4	1.2	1.6	2.3	*	*
2001.....	0.6	*	*	*	*	*	0.7	1.3	1.6	2.6	4.1	*
2000.....	0.6	*	*	*	*	*	0.5	1.3	1.6	1.9	2.8	*
1999.....	0.6	*	*	*	*	*	0.7	1.2	2.0	2.6	*	*
Black, male												
2004.....	0.6	*	*	*	*	*	*	1.8	*	*	*	*
2003.....	0.7	*	*	*	*	*	*	2.0	2.2	*	*	*
2002.....	0.6	*	*	*	*	*	*	1.5	1.9	2.9	*	*
2001.....	0.8	*	*	*	*	*	0.7	1.9	2.1	3.8	*	*
2000.....	0.6	*	*	*	*	*	0.8	1.8	*	*	*	*
1999.....	0.7	*	*	*	*	*	1.0	1.9	2.5	*	*	*
Black, female												
2004.....	0.5	*	*	*	*	*	*	1.0	*	2.0	*	*
2003.....	0.5	*	*	*	*	*	*	*	1.7	2.0	*	*
2002.....	0.5	*	*	*	*	*	*	1.0	*	*	*	*
2001.....	0.5	*	*	*	*	*	0.7	*	*	*	3.3	*
2000.....	0.5	*	*	*	*	*	*	1.0	1.6	*	*	*
1999.....	0.5	*	*	*	*	*	*	*	1.6	2.8	*	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Diseases of pericardium and acute myocarditis (I30-I31,I40)

All races 3/, both sexes

2004.....	0.3	0.5	0.2	*	0.1	0.1	0.2	0.3	0.4	0.6	1.2	1.8
2003.....	0.3	0.8	0.1	0.1	0.1	0.1	0.2	0.3	0.4	0.8	1.2	1.6
2002.....	0.3	0.6	0.1	0.1	0.1	0.1	0.2	0.3	0.4	0.8	1.2	2.1
2001.....	0.3	0.8	0.2	0.1	0.1	0.1	0.2	0.3	0.4	0.7	1.2	1.9
2000.....	0.3	0.5	0.2	0.0	0.1	0.1	0.2	0.2	0.4	0.7	1.2	1.8
1999.....	0.3	0.7	0.2	0.1	0.1	0.1	0.3	0.3	0.5	0.8	1.3	2.0

All races 3/, male

2004.....	0.3	*	*	*	*	*	0.2	0.3	0.5	0.7	1.3	2.3
2003.....	0.3	*	*	*	0.2	0.1	0.2	0.4	0.4	1.0	1.3	1.5
2002.....	0.3	*	*	*	0.1	0.2	0.2	0.4	0.5	0.9	1.2	2.4
2001.....	0.3	*	*	*	0.1	0.2	0.2	0.3	0.5	0.7	1.2	1.8
2000.....	0.3	*	*	*	0.1	0.1	0.3	0.2	0.4	0.8	1.5	2.1
1999.....	0.3	*	*	*	0.1	0.2	0.3	0.3	0.6	0.8	1.4	2.5

All races 3/, female

2004.....	0.3	*	*	*	*	0.1	0.1	0.2	0.3	0.5	1.2	1.6
2003.....	0.3	*	*	*	0.1	*	0.2	0.2	0.3	0.7	1.1	1.7
2002.....	0.3	*	*	*	*	*	0.1	0.2	0.4	0.7	1.3	2.0
2001.....	0.3	*	*	*	*	*	0.2	0.3	0.3	0.6	1.2	2.0
2000.....	0.3	*	*	*	*	0.1	0.1	0.2	0.3	0.6	1.0	1.7
1999.....	0.3	*	*	*	*	*	0.2	0.3	0.4	0.8	1.3	1.9

White, both sexes

2004.....	0.2	*	*	*	*	0.1	0.1	0.2	0.4	0.6	1.2	1.8
2003.....	0.3	*	*	*	0.1	0.1	0.2	0.2	0.4	0.8	1.2	1.7
2002.....	0.3	*	*	*	0.1	0.1	0.2	0.2	0.4	0.8	1.1	2.2
2001.....	0.3	*	*	*	0.1	0.1	0.2	0.2	0.4	0.7	1.2	1.9
2000.....	0.3	*	*	*	0.1	0.1	0.2	0.2	0.3	0.7	1.2	1.8
1999.....	0.3	*	*	*	0.1	0.1	0.2	0.3	0.5	0.8	1.3	2.0

White, male

2004.....	0.3	*	*	*	*	*	0.1	0.3	0.5	0.7	1.3	2.2
2003.....	0.3	*	*	*	0.1	*	0.2	0.3	0.4	1.1	1.3	1.7
2002.....	0.3	*	*	*	*	*	0.1	0.2	0.3	0.5	0.9	2.5
2001.....	0.3	*	*	*	*	*	0.2	0.2	0.3	0.5	0.7	2.0
2000.....	0.3	*	*	*	*	*	0.2	0.2	0.4	0.7	1.5	2.1
1999.....	0.3	*	*	*	*	0.1	0.3	0.3	0.6	0.8	1.4	2.5

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other diseases of endocardium (I34-I38)												
All races 3/, both sexes												
2004.....	6.9	*	*	*	0.1	0.3	0.7	1.7	3.6	12.0	46.7	200.3
2003.....	6.9	*	*	*	0.1	0.3	0.7	1.5	3.7	12.3	47.7	201.9
2002.....	6.9	*	*	*	0.1	0.3	0.8	1.6	3.7	12.0	49.3	203.9
2001.....	6.9	*	*	*	0.1	0.3	0.7	1.6	3.7	13.0	48.8	204.2
2000.....	7.0	*	*	*	0.1	0.3	0.8	1.6	4.0	13.2	49.8	213.2
1999.....	7.0	*	*	*	0.2	0.4	0.8	1.6	3.9	13.2	51.7	209.8
All races 3/, male												
2004.....	5.5	*	*	*	0.1	0.3	0.8	2.2	4.3	12.9	48.8	194.2
2003.....	5.4	*	*	*	0.1	0.3	0.9	1.7	4.3	13.2	51.1	195.3
2002.....	5.4	*	*	*	0.1	0.3	0.9	2.0	4.1	12.7	51.6	198.5
2001.....	5.4	*	*	*	*	0.3	0.8	2.1	4.2	14.0	52.8	200.8
2000.....	5.5	*	*	*	0.1	0.3	0.9	2.1	4.7	14.0	51.6	215.8
1999.....	5.5	*	*	*	0.2	0.3	1.0	1.9	4.5	14.7	56.4	206.4
All races 3/, female												
2004.....	8.3	*	*	*	0.1	0.3	0.6	1.2	3.1	11.3	45.3	203.0
2003.....	8.3	*	*	*	*	0.3	0.6	1.3	3.1	11.5	45.4	204.8
2002.....	8.4	*	*	*	*	0.3	0.6	1.2	3.4	11.4	47.7	206.3
2001.....	8.2	*	*	*	0.1	0.3	0.5	1.2	3.2	12.1	46.2	205.6
2000.....	8.5	*	*	*	0.1	0.3	0.7	1.2	3.3	12.6	48.6	212.1
1999.....	8.5	*	*	*	0.2	0.4	0.7	1.4	3.3	12.1	48.7	211.2
White, both sexes												
2004.....	7.8	*	*	*	0.1	0.3	0.6	1.5	3.5	12.3	49.0	212.5
2003.....	7.8	*	*	*	0.1	0.3	0.6	1.3	3.5	12.4	49.9	214.4
2002.....	7.9	*	*	*	0.1	0.3	0.7	1.4	3.6	12.1	52.0	216.2
2001.....	7.8	*	*	*	0.1	0.3	0.6	1.5	3.5	13.2	51.2	216.5
2000.....	7.9	*	*	*	0.1	0.3	0.7	1.5	3.7	13.4	51.9	225.8
1999.....	7.9	*	*	*	0.2	0.3	0.8	1.4	3.7	13.1	54.0	221.4
White, male												
2004.....	6.1	*	*	*	*	0.3	0.7	1.9	4.2	13.4	50.9	205.2
2003.....	6.1	*	*	*	0.1	0.3	0.8	1.5	4.0	13.5	53.4	207.8
2002.....	6.1	*	*	*	*	0.3	0.9	1.7	4.0	13.0	54.4	211.7
2001.....	6.1	*	*	*	*	0.3	0.7	1.9	4.0	14.2	55.5	212.0
2000.....	6.1	*	*	*	*	0.3	0.8	1.9	4.5	14.2	53.8	229.0
1999.....	6.2	*	*	*	0.2	0.3	0.9	1.7	4.3	14.7	58.8	218.2

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other diseases of endocardium (I34-I38)												
White, female												
2004.....	9.5	*	*	*	*	0.2	0.5	1.0	2.9	11.3	47.6	215.8
2003.....	9.5	*	*	*	*	0.3	0.5	1.2	2.9	11.6	47.5	217.2
2002.....	9.6	*	*	*	*	0.3	0.5	1.0	3.3	11.4	50.3	218.2
2001.....	9.5	*	*	*	*	0.3	0.5	1.0	3.0	12.3	48.4	218.4
2000.....	9.7	*	*	*	*	0.3	0.5	1.1	3.0	12.7	50.7	224.5
1999.....	9.6	*	*	*	0.1	0.4	0.6	1.2	3.1	11.8	50.8	222.7
Black, both sexes												
2004.....	3.3	*	*	*	*	0.5	1.3	3.8	5.5	12.6	30.0	88.3
2003.....	3.4	*	*	*	*	0.6	1.4	3.4	6.4	12.8	30.7	85.2
2002.....	3.2	*	*	*	*	0.5	1.4	3.7	5.6	12.9	28.1	86.4
2001.....	3.3	*	*	*	*	0.6	1.2	3.3	6.1	13.5	30.6	85.4
2000.....	3.5	*	*	*	*	0.6	1.5	3.2	6.5	14.6	33.6	92.9
1999.....	3.5	*	*	*	*	0.6	1.6	3.4	6.4	16.1	32.0	91.5
Black, male												
2004.....	2.9	*	*	*	*	*	1.5	5.0	6.5	11.6	30.9	84.4
2003.....	2.9	*	*	*	*	*	1.7	4.3	8.0	12.5	31.9	70.9
2002.....	2.6	*	*	*	*	*	1.3	4.3	6.5	12.1	28.4	67.5
2001.....	2.9	*	*	*	*	*	1.5	4.4	7.3	14.2	30.9	86.2
2000.....	3.0	*	*	*	*	*	1.6	4.3	7.0	14.5	34.5	86.5
1999.....	3.0	*	*	*	*	*	1.8	3.9	7.2	17.2	34.1	77.3
Black, female												
2004.....	3.7	*	*	*	*	*	1.0	2.8	4.7	13.3	29.4	89.8
2003.....	3.8	*	*	*	*	0.7	1.1	2.7	5.1	13.0	30.0	90.7
2002.....	3.8	*	*	*	*	*	1.5	3.2	4.9	13.4	27.9	93.6
2001.....	3.6	*	*	*	*	*	1.0	2.5	5.1	13.0	30.4	85.0
2000.....	4.0	*	*	*	*	*	1.4	2.3	6.1	14.6	33.1	95.3
1999.....	4.0	*	*	*	*	*	1.4	3.0	5.8	15.2	30.8	96.8

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Nonrheumatic mitral valve disorders (I34)												
All races 3/, both sexes												
2004.....	0.9	*	*	*	*	0.1	0.2	0.3	0.5	1.8	6.3	20.1
2003.....	0.9	*	*	*	*	0.1	0.2	0.3	0.6	1.9	6.4	20.1
2002.....	1.0	*	*	*	0.1	0.1	0.2	0.3	0.7	2.1	7.0	22.1
2001.....	1.0	*	*	*	0.1	0.1	0.2	0.3	0.6	2.2	7.3	22.7
2000.....	1.0	*	*	*	0.1	0.1	0.2	0.3	0.7	2.2	7.7	24.3
1999.....	1.0	*	*	*	0.1	0.1	0.2	0.4	0.7	2.3	7.7	25.1
All races 3/, male												
2004.....	0.6	*	*	*	*	*	0.2	0.3	0.5	1.6	5.8	16.6
2003.....	0.6	*	*	*	*	*	0.2	0.2	0.6	1.8	6.0	16.4
2002.....	0.7	*	*	*	*	*	0.2	0.2	0.6	1.8	5.7	21.0
2001.....	0.7	*	*	*	*	0.1	0.2	0.3	0.6	2.2	6.1	19.0
2000.....	0.7	*	*	*	*	*	0.2	0.3	0.6	2.1	7.2	22.9
1999.....	0.7	*	*	*	*	*	0.2	0.3	0.6	2.1	7.5	22.2
All races 3/, female												
2004.....	1.1	*	*	*	*	0.1	0.2	0.3	0.6	2.0	6.7	21.7
2003.....	1.1	*	*	*	*	0.1	0.2	0.3	0.6	1.9	6.6	21.7
2002.....	1.2	*	*	*	*	0.1	0.2	0.3	0.8	2.4	7.9	22.6
2001.....	1.3	*	*	*	*	0.2	0.2	0.3	0.6	2.2	8.1	24.3
2000.....	1.3	*	*	*	*	0.1	0.3	0.3	0.8	2.4	8.1	24.9
1999.....	1.3	*	*	*	*	0.2	0.2	0.4	0.8	2.4	7.8	26.2
White, both sexes												
2004.....	1.0	*	*	*	*	0.1	0.2	0.3	0.5	1.9	6.6	21.1
2003.....	1.0	*	*	*	*	0.1	0.2	0.3	0.6	1.9	6.6	21.4
2002.....	1.1	*	*	*	*	0.1	0.2	0.2	0.7	2.2	7.4	23.8
2001.....	1.1	*	*	*	*	0.1	0.2	0.3	0.6	2.2	7.7	24.2
2000.....	1.1	*	*	*	*	0.1	0.2	0.3	0.7	2.3	8.1	25.5
1999.....	1.2	*	*	*	*	0.1	0.3	0.3	0.7	2.3	8.0	26.5
White, male												
2004.....	0.7	*	*	*	*	*	0.2	0.3	0.5	1.7	5.9	17.5
2003.....	0.7	*	*	*	*	*	0.1	0.2	0.6	1.9	6.2	17.6
2002.....	0.7	*	*	*	*	*	0.2	0.2	0.6	1.8	6.0	22.6
2001.....	0.8	*	*	*	*	*	0.2	0.4	0.6	2.2	6.5	20.5
2000.....	0.8	*	*	*	*	*	0.2	0.3	0.6	2.2	7.6	24.3
1999.....	0.8	*	*	*	*	*	0.3	0.3	0.7	2.1	7.9	23.7

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Nonrheumatic mitral valve disorders (I34)												
White, female												
2004.....	1.3	*	*	*	*	*	0.2	0.2	0.6	2.0	7.0	22.7
2003.....	1.3	*	*	*	*	*	0.2	0.3	0.6	2.0	6.9	23.0
2002.....	1.4	*	*	*	*	0.1	0.2	0.3	0.8	2.5	8.2	24.3
2001.....	1.4	*	*	*	*	0.2	0.2	0.3	0.6	2.2	8.5	25.7
2000.....	1.5	*	*	*	*	0.1	0.3	0.3	0.8	2.4	8.5	25.9
1999.....	1.5	*	*	*	*	0.2	0.3	0.4	0.7	2.4	8.0	27.6
Black, both sexes												
2004.....	0.5	*	*	*	*	*	*	0.5	1.0	1.9	4.3	9.9
2003.....	0.5	*	*	*	*	*	*	0.5	1.1	1.5	4.2	7.9
2002.....	0.5	*	*	*	*	*	*	0.5	0.9	2.1	3.9	6.8
2001.....	0.5	*	*	*	*	*	*	0.6	0.9	2.6	3.5	7.7
2000.....	0.5	*	*	*	*	*	*	0.6	0.9	2.1	4.5	12.2
1999.....	0.5	*	*	*	*	*	*	0.6	1.0	2.6	4.4	10.2
Black, male												
2004.....	0.4	*	*	*	*	*	*	*	*	*	*	*
2003.....	0.4	*	*	*	*	*	*	*	*	*	*	*
2002.....	0.3	*	*	*	*	*	*	*	*	*	*	*
2001.....	0.3	*	*	*	*	*	*	*	*	*	*	*
2000.....	0.3	*	*	*	*	*	*	*	*	*	*	*
1999.....	0.3	*	*	*	*	*	*	*	*	*	*	*
Black, female												
2004.....	0.6	*	*	*	*	*	*	*	*	2.2	4.4	11.0
2003.....	0.6	*	*	*	*	*	*	*	*	*	4.5	8.9
2002.....	0.6	*	*	*	*	*	*	*	*	2.4	4.4	*
2001.....	0.7	*	*	*	*	*	*	*	*	2.8	4.5	9.7
2000.....	0.7	*	*	*	*	*	*	*	*	2.7	4.9	13.7
1999.....	0.7	*	*	*	*	*	*	*	*	2.5	5.3	12.3

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Nonrheumatic aortic valve disorders (I35)												
All races 3/, both sexes												
2004.....	4.3	*	*	*	*	0.1	0.2	0.6	1.7	6.1	28.8	142.5
2003.....	4.3	*	*	*	*	0.1	0.2	0.5	1.6	6.4	29.2	144.9
2002.....	4.3	*	*	*	*	0.1	0.2	0.6	1.6	6.3	30.2	144.8
2001.....	4.3	*	*	*	*	0.1	0.2	0.7	1.7	6.8	30.0	144.1
2000.....	4.4	*	*	*	*	0.1	0.2	0.6	1.8	7.1	30.9	151.3
1999.....	4.4	*	*	*	*	0.1	0.3	0.7	1.8	6.9	32.4	147.8
All races 3/, male												
2004.....	3.4	*	*	*	*	0.1	0.3	0.8	2.1	7.2	31.1	141.9
2003.....	3.5	*	*	*	*	0.1	0.3	0.7	2.0	7.5	33.3	144.7
2002.....	3.5	*	*	*	*	0.1	0.3	0.8	2.0	7.4	34.4	144.9
2001.....	3.5	*	*	*	*	0.1	0.3	1.0	2.1	8.0	34.8	145.9
2000.....	3.5	*	*	*	*	0.1	0.3	1.0	2.5	8.2	33.3	158.2
1999.....	3.5	*	*	*	*	0.1	0.4	0.9	2.3	8.6	37.2	150.3
All races 3/, female												
2004.....	5.2	*	*	*	*	*	0.1	0.4	1.2	5.3	27.2	142.7
2003.....	5.1	*	*	*	*	*	0.1	0.4	1.1	5.4	26.5	145.0
2002.....	5.2	*	*	*	*	*	0.2	0.4	1.3	5.3	27.4	144.8
2001.....	5.1	*	*	*	*	*	*	0.4	1.3	5.8	26.9	143.4
2000.....	5.3	*	*	*	*	*	0.1	0.3	1.2	6.1	29.2	148.5
1999.....	5.2	*	*	*	*	*	0.2	0.4	1.3	5.6	29.2	146.8
White, both sexes												
2004.....	5.0	*	*	*	*	0.1	0.2	0.6	1.7	6.5	30.3	151.6
2003.....	5.0	*	*	*	*	0.1	0.2	0.5	1.6	6.6	30.7	154.1
2002.....	5.0	*	*	*	*	0.1	0.2	0.6	1.7	6.4	31.9	153.2
2001.....	4.9	*	*	*	*	0.1	0.2	0.7	1.7	7.0	31.5	152.8
2000.....	5.1	*	*	*	*	0.1	0.2	0.6	1.8	7.2	32.4	160.7
1999.....	5.0	*	*	*	*	0.1	0.3	0.6	1.8	6.9	34.0	156.1
White, male												
2004.....	3.9	*	*	*	*	*	0.3	0.8	2.1	7.6	32.7	150.1
2003.....	4.0	*	*	*	*	*	0.3	0.6	2.1	7.8	35.0	154.4
2002.....	4.0	*	*	*	*	*	0.3	0.8	2.1	7.7	36.4	154.4
2001.....	3.9	*	*	*	*	*	0.3	1.0	2.2	8.3	36.3	153.5
2000.....	4.0	*	*	*	*	*	0.3	0.9	2.6	8.4	35.0	168.4
1999.....	4.0	*	*	*	*	*	0.3	0.8	2.3	8.7	39.0	159.0

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Nonrheumatic aortic valve disorders (I35)												
White, female												
2004.....	6.1	*	*	*	*	*	*	0.3	1.3	5.5	28.7	152.2
2003.....	6.0	*	*	*	*	*	0.1	0.4	1.1	5.6	27.8	154.0
2002.....	6.0	*	*	*	*	*	0.1	0.4	1.4	5.3	29.0	152.7
2001.....	5.9	*	*	*	*	*	*	0.4	1.3	6.0	28.4	152.5
2000.....	6.1	*	*	*	*	*	0.1	0.3	1.1	6.2	30.7	157.6
1999.....	6.0	*	*	*	*	*	0.2	0.4	1.3	5.5	30.7	155.0
Black, both sexes												
2004.....	1.5	*	*	*	*	*	*	0.9	1.8	4.8	17.3	59.2
2003.....	1.5	*	*	*	*	*	0.4	0.8	1.7	5.3	16.9	58.6
2002.....	1.5	*	*	*	*	*	0.4	1.0	1.6	5.9	16.2	63.0
2001.....	1.5	*	*	*	*	*	*	1.0	1.8	5.5	18.3	59.6
2000.....	1.7	*	*	*	*	*	0.4	0.9	2.3	7.1	18.3	62.3
1999.....	1.7	*	*	*	*	*	0.4	1.2	2.2	7.5	17.7	64.4
Black, male												
2004.....	1.3	*	*	*	*	*	*	1.2	2.9	5.0	17.8	58.6
2003.....	1.2	*	*	*	*	*	*	1.1	2.6	5.5	17.8	48.0
2002.....	1.2	*	*	*	*	*	*	1.3	2.1	6.3	16.6	52.5
2001.....	1.5	*	*	*	*	*	*	1.7	2.2	6.2	21.5	68.1
2000.....	1.4	*	*	*	*	*	*	1.5	2.8	8.2	19.4	57.7
1999.....	1.6	*	*	*	*	*	*	1.7	2.7	9.3	21.5	58.6
Black, female												
2004.....	1.7	*	*	*	*	*	*	*	*	4.6	17.0	59.5
2003.....	1.7	*	*	*	*	*	*	*	*	5.1	16.4	62.8
2002.....	1.8	*	*	*	*	*	*	*	*	5.6	15.9	66.9
2001.....	1.6	*	*	*	*	*	*	*	1.5	5.0	16.5	56.4
2000.....	1.9	*	*	*	*	*	*	*	2.0	6.4	17.7	64.0
1999.....	1.9	*	*	*	*	*	*	*	1.7	6.3	15.5	66.6

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other diseases of endocardium (I36-I38)												
All races 3/, both sexes												
2004.....	1.7	*	*	*	0.0	0.1	0.3	0.8	1.4	4.1	11.6	37.7
2003.....	1.7	*	*	*	*	0.1	0.3	0.7	1.5	4.1	12.1	36.9
2002.....	1.7	*	*	*	*	0.1	0.3	0.7	1.4	3.6	12.1	37.0
2001.....	1.6	*	*	*	*	0.1	0.3	0.6	1.4	4.0	11.5	37.4
2000.....	1.6	*	*	*	*	0.1	0.3	0.7	1.4	3.9	11.2	37.6
1999.....	1.6	*	*	*	0.1	0.1	0.3	0.6	1.4	4.0	11.7	37.0
All races 3/, male												
2004.....	1.4	*	*	*	*	0.1	0.3	1.1	1.7	4.2	11.9	35.7
2003.....	1.3	*	*	*	*	0.1	0.4	0.8	1.6	3.9	11.9	34.2
2002.....	1.3	*	*	*	*	0.1	0.4	0.9	1.5	3.5	11.5	32.6
2001.....	1.3	*	*	*	*	0.1	0.3	0.8	1.5	3.9	12.0	35.9
2000.....	1.2	*	*	*	*	*	0.4	0.8	1.6	3.6	11.1	34.7
1999.....	1.3	*	*	*	*	0.1	0.4	0.7	1.6	3.9	11.7	33.9
All races 3/, female												
2004.....	2.0	*	*	*	*	*	0.2	0.6	1.2	4.0	11.4	38.5
2003.....	2.1	*	*	*	*	0.1	0.2	0.7	1.4	4.2	12.3	38.1
2002.....	2.0	*	*	*	*	0.1	0.3	0.5	1.3	3.8	12.4	38.9
2001.....	1.9	*	*	*	*	*	0.2	0.4	1.3	4.1	11.2	38.0
2000.....	1.9	*	*	*	*	0.1	0.2	0.6	1.3	4.1	11.3	38.7
1999.....	2.0	*	*	*	*	0.1	0.3	0.6	1.3	4.1	11.7	38.2
White, both sexes												
2004.....	1.9	*	*	*	*	0.1	0.2	0.6	1.3	4.0	12.1	39.9
2003.....	1.8	*	*	*	*	0.1	0.2	0.6	1.3	3.9	12.6	38.9
2002.....	1.8	*	*	*	*	0.1	0.3	0.5	1.2	3.5	12.7	39.2
2001.....	1.7	*	*	*	*	*	0.2	0.5	1.1	3.9	11.9	39.5
2000.....	1.7	*	*	*	*	0.1	0.2	0.6	1.2	3.8	11.4	39.6
1999.....	1.7	*	*	*	0.1	0.1	0.3	0.5	1.2	3.9	12.0	38.8
White, male												
2004.....	1.5	*	*	*	*	0.1	0.3	0.8	1.6	4.2	12.3	37.6
2003.....	1.4	*	*	*	*	*	0.3	0.6	1.4	3.8	12.2	35.8
2002.....	1.4	*	*	*	*	0.1	0.3	0.7	1.3	3.4	12.0	34.8
2001.....	1.4	*	*	*	*	*	0.3	0.6	1.2	3.8	12.6	38.0
2000.....	1.3	*	*	*	*	*	0.3	0.6	1.4	3.6	11.2	36.4
1999.....	1.3	*	*	*	*	*	0.3	0.6	1.4	3.9	12.0	35.5

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

All other diseases of endocardium (I36-I38)

White, female

2004.....	2.2	*	*	*	*	*	0.2	0.4	1.0	3.8	12.0	40.9
2003.....	2.3	*	*	*	*	*	0.2	0.5	1.2	4.0	12.8	40.2
2002.....	2.2	*	*	*	*	*	0.2	0.4	1.1	3.6	13.1	41.1
2001.....	2.1	*	*	*	*	*	0.2	0.3	1.1	4.1	11.5	40.2
2000.....	2.1	*	*	*	*	*	0.1	0.5	1.1	4.0	11.5	40.9
1999.....	2.1	*	*	*	*	0.1	0.2	0.5	1.1	3.9	12.0	40.2

Black, both sexes

2004.....	1.3	*	*	*	*	*	0.7	2.5	2.7	5.9	8.3	19.2
2003.....	1.4	*	*	*	*	*	0.7	2.2	3.6	6.0	9.6	18.7
2002.....	1.2	*	*	*	*	*	0.8	2.2	3.1	4.9	8.0	16.6
2001.....	1.2	*	*	*	*	*	0.6	1.7	3.4	5.3	8.8	18.1
2000.....	1.3	*	*	*	*	*	0.8	1.8	3.2	5.3	10.8	18.5
1999.....	1.2	*	*	*	*	*	0.9	1.7	3.2	6.0	9.9	16.9

Black, male

2004.....	1.3	*	*	*	*	*	0.9	3.4	2.7	5.1	8.9	*
2003.....	1.3	*	*	*	*	*	0.9	2.8	4.5	5.5	10.4	*
2002.....	1.1	*	*	*	*	*	0.8	2.5	3.7	4.1	8.6	*
2001.....	1.1	*	*	*	*	*	*	2.2	4.3	5.5	7.7	*
2000.....	1.2	*	*	*	*	*	0.8	2.1	3.5	5.0	11.5	*
1999.....	1.1	*	*	*	*	*	0.9	1.8	3.9	5.3	9.8	*

Black, female

2004.....	1.4	*	*	*	*	*	*	1.7	2.7	6.4	8.0	19.3
2003.....	1.4	*	*	*	*	*	*	1.7	2.9	6.4	9.1	19.0
2002.....	1.4	*	*	*	*	*	0.8	1.9	2.7	5.4	7.6	19.3
2001.....	1.3	*	*	*	*	*	*	1.3	2.6	5.2	9.4	18.9
2000.....	1.4	*	*	*	*	*	0.8	1.5	3.0	5.6	10.4	17.6
1999.....	1.4	*	*	*	*	*	0.8	1.6	2.7	6.4	10.0	18.0

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Cardiomyopathy (I42)												
All races 3/, both sexes												
2004.....	8.7	2.8	0.2	0.2	0.6	1.4	3.1	6.1	10.7	22.8	54.6	127.6
2003.....	9.2	3.1	0.4	0.1	0.7	1.5	3.3	6.3	11.5	25.5	57.0	137.8
2002.....	9.6	2.9	0.2	0.2	0.7	1.6	3.5	6.5	11.9	27.0	61.1	142.0
2001.....	9.4	3.4	0.4	0.2	0.7	1.4	3.3	6.3	12.3	27.2	60.3	138.3
2000.....	9.7	3.1	0.3	0.2	0.7	1.5	3.3	6.7	13.0	28.7	61.4	142.3
1999.....	9.8	3.5	0.2	0.2	0.7	1.5	3.6	6.7	14.0	30.4	60.4	139.2
All races 3/, male												
2004.....	10.3	2.7	0.3	0.2	0.9	2.1	4.4	8.6	15.1	31.6	76.2	174.6
2003.....	10.8	3.4	0.3	0.2	0.9	2.1	4.7	8.9	16.1	35.2	81.0	178.8
2002.....	11.3	2.7	*	0.2	1.0	2.3	4.8	9.4	16.7	37.8	87.2	189.7
2001.....	11.0	2.9	0.5	0.2	0.9	2.0	4.8	8.9	17.3	37.4	84.9	184.7
2000.....	11.4	2.8	0.3	0.3	0.9	2.1	4.8	9.6	18.2	40.0	87.6	195.3
1999.....	11.5	3.1	*	0.2	0.9	2.0	5.2	9.6	19.6	43.0	84.3	188.2
All races 3/, female												
2004.....	7.2	3.0	*	0.2	0.3	0.7	1.9	3.6	6.7	15.4	40.1	106.4
2003.....	7.7	2.9	0.4	*	0.5	0.9	1.9	3.7	7.2	17.4	40.9	119.7
2002.....	8.0	3.1	*	0.1	0.4	0.9	2.2	3.6	7.5	18.1	43.7	121.3
2001.....	8.0	3.9	0.3	0.2	0.4	0.8	1.9	3.9	7.7	18.8	44.2	118.9
2000.....	8.0	3.4	*	0.1	0.4	1.0	1.9	4.0	8.2	19.4	44.3	120.7
1999.....	8.1	3.8	*	0.1	0.4	1.0	2.0	4.0	8.9	20.0	44.9	119.4
White, both sexes												
2004.....	8.7	2.7	0.2	0.2	0.5	1.0	2.5	4.8	9.2	20.9	53.7	129.3
2003.....	9.2	3.0	0.3	0.1	0.5	1.1	2.6	5.0	9.7	23.6	55.8	139.8
2002.....	9.6	2.8	0.2	0.2	0.5	1.2	2.7	5.2	10.3	25.4	59.8	141.7
2001.....	9.3	2.8	0.3	0.1	0.5	1.0	2.5	5.0	10.5	25.5	58.6	139.2
2000.....	9.5	3.0	0.2	0.2	0.5	1.1	2.5	5.3	11.2	26.5	59.4	142.4
1999.....	9.6	3.1	0.2	0.1	0.5	1.2	2.7	5.3	11.7	28.1	58.6	138.6
White, male												
2004.....	10.2	2.6	*	0.1	0.8	1.6	3.5	7.1	12.9	29.3	75.5	178.3
2003.....	10.7	2.9	*	0.2	0.7	1.6	3.8	7.3	13.8	32.6	79.9	182.6
2002.....	11.2	2.4	*	0.2	0.8	1.7	3.8	7.7	14.6	35.4	86.4	190.9
2001.....	10.8	2.7	0.4	*	0.7	1.5	3.7	7.2	14.9	35.2	83.0	187.0
2000.....	11.1	2.7	*	0.2	0.7	1.6	3.7	7.7	15.8	37.1	85.2	197.5
1999.....	11.1	2.6	*	0.2	0.6	1.6	4.0	7.6	16.3	39.9	82.4	188.0

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Cardiomyopathy (I42)												
White, female												
2004.....	7.2	2.8	*	0.2	0.3	0.5	1.4	2.6	5.7	13.7	38.8	107.2
2003.....	7.8	3.1	*	*	0.3	0.6	1.4	2.8	5.9	15.8	39.4	120.8
2002.....	8.0	3.2	*	0.1	0.2	0.6	1.6	2.8	6.2	16.8	42.0	120.4
2001.....	7.9	2.9	*	0.1	0.3	0.6	1.3	2.8	6.4	17.2	42.4	119.2
2000.....	8.0	3.4	*	*	0.3	0.7	1.4	2.9	6.9	17.6	42.4	120.0
1999.....	8.1	3.5	*	0.1	0.3	0.7	1.4	3.0	7.3	18.3	42.9	118.8
Black, both sexes												
2004.....	11.2	*	*	*	1.3	3.7	8.1	15.8	26.7	44.9	75.7	124.4
2003.....	11.9	3.5	*	*	1.5	4.2	8.4	16.3	29.3	48.0	81.0	135.7
2002.....	12.2	3.4	*	*	1.6	4.6	9.2	16.7	28.5	47.5	84.8	161.0
2001.....	12.4	6.8	0.9	0.3	1.7	3.9	9.4	17.4	30.3	49.1	90.2	141.2
2000.....	12.9	3.6	*	0.4	1.8	4.2	8.9	19.3	30.2	55.0	92.9	151.1
1999.....	13.2	5.1	*	0.4	1.8	3.8	10.0	19.2	35.2	56.1	88.5	153.0
Black, male												
2004.....	13.7	*	*	*	2.0	5.3	11.5	22.3	39.1	62.9	102.5	158.0
2003.....	14.5	*	*	*	1.8	5.8	11.7	23.0	41.7	69.1	112.3	172.0
2002.....	14.9	*	*	*	2.1	6.8	12.6	24.6	40.0	68.5	113.9	204.8
2001.....	14.9	*	*	*	2.3	5.4	13.1	23.7	43.3	67.2	127.7	180.4
2000.....	15.8	*	*	*	2.5	5.8	13.0	26.9	43.6	77.4	132.6	193.7
1999.....	16.4	*	*	*	2.6	5.2	14.3	27.9	52.0	81.8	117.4	215.6
Black, female												
2004.....	8.9	*	*	*	0.6	2.3	5.2	10.2	16.7	32.0	60.6	111.0
2003.....	9.5	*	*	*	1.1	2.8	5.5	10.5	19.4	33.0	63.3	121.5
2002.....	9.8	*	*	*	1.1	2.7	6.2	9.8	19.2	32.5	68.4	144.2
2001.....	10.2	8.8	*	*	1.1	2.5	6.1	12.0	19.7	36.3	69.1	126.7
2000.....	10.3	*	*	*	1.2	2.8	5.3	12.8	19.5	39.2	70.6	135.2
1999.....	10.3	*	*	*	1.0	2.5	6.2	11.8	21.8	38.0	72.3	129.7

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Conduction disorders and cardiac dysrhythmias (I44-I49)												
All races 3/, both sexes												
2004.....	12.8	0.8	0.1	0.1	0.4	0.8	1.9	5.0	11.9	28.8	84.5	295.3
2003.....	13.4	0.8	0.1	0.1	0.4	0.8	1.9	5.0	12.5	31.3	90.0	313.7
2002.....	13.6	1.4	0.1	0.1	0.5	0.8	2.0	4.9	12.6	31.5	93.4	323.2
2001.....	13.5	0.9	0.2	0.1	0.4	0.7	1.7	5.0	12.6	32.6	92.4	328.2
2000.....	13.6	1.0	0.2	0.1	0.5	0.7	1.9	4.8	13.1	33.3	92.2	340.2
1999.....	14.3	1.0	0.3	0.1	0.5	0.8	1.9	5.2	14.5	35.7	97.1	353.7
All races 3/, male												
2004.....	11.5	*	*	0.1	0.5	0.9	2.3	6.5	16.2	36.2	93.8	278.5
2003.....	12.0	*	*	0.2	0.5	0.9	2.4	6.6	16.6	40.1	99.9	292.7
2002.....	12.2	1.6	*	0.1	0.6	0.9	2.5	6.6	16.8	39.0	103.8	310.5
2001.....	12.0	*	*	0.1	0.6	0.9	2.2	6.7	16.5	39.8	104.9	311.6
2000.....	12.1	1.1	*	0.1	0.5	0.8	2.3	6.6	17.5	42.9	101.9	333.2
1999.....	12.9	1.2	0.3	0.2	0.6	1.0	2.3	7.1	19.2	45.4	110.8	353.4
All races 3/, female												
2004.....	14.1	*	*	0.1	0.3	0.7	1.4	3.5	8.0	22.6	78.3	302.9
2003.....	14.8	*	*	0.3	0.3	0.7	1.5	3.5	8.7	24.0	83.3	323.0
2002.....	15.1	1.2	*	0.3	0.6	1.4	3.3	3.3	8.6	25.3	86.6	328.8
2001.....	15.1	*	*	0.1	0.2	0.6	1.3	3.4	9.0	26.7	84.2	335.2
2000.....	15.1	*	*	0.4	0.4	0.6	1.5	3.0	8.9	25.4	85.8	343.0
1999.....	15.7	*	*	0.1	0.4	0.6	1.4	3.4	10.2	27.8	88.2	353.8
White, both sexes												
2004.....	13.5	0.7	*	0.1	0.4	0.7	1.6	4.4	10.9	27.3	83.0	295.0
2003.....	14.2	*	*	0.1	0.4	0.7	1.8	4.4	11.5	29.4	88.7	314.5
2002.....	14.4	1.1	*	0.1	0.4	0.6	1.8	4.4	11.3	30.0	91.7	324.4
2001.....	14.3	0.9	0.2	0.1	0.3	0.6	1.5	4.5	11.4	30.9	90.3	329.7
2000.....	14.4	1.0	0.2	0.1	0.4	0.6	1.7	4.3	12.0	31.8	91.0	341.6
1999.....	15.2	0.8	0.3	0.1	0.4	0.7	1.7	4.7	13.4	34.3	95.4	355.9
White, male												
2004.....	11.9	*	*	*	0.5	0.8	2.1	5.9	15.0	34.3	91.5	273.7
2003.....	12.5	*	*	0.1	0.5	0.8	2.2	6.0	15.5	37.7	97.9	291.3
2002.....	12.7	1.6	*	0.1	0.6	0.8	2.4	6.0	15.4	37.3	101.6	308.9
2001.....	12.4	*	*	0.1	0.5	0.7	2.0	6.1	15.0	37.8	101.6	310.2
2000.....	12.7	*	*	*	0.5	0.8	2.1	5.9	16.3	41.1	101.1	332.8
1999.....	13.5	*	0.3	0.2	0.5	0.9	2.2	6.6	17.9	43.7	108.0	350.9

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Conduction disorders and cardiac dysrhythmias (I44-I49)												
White, female												
2004.....	15.0	*	*	*	0.3	0.5	1.1	3.0	7.0	21.4	77.1	304.7
2003.....	15.9	*	*	*	0.3	0.5	1.3	2.9	7.8	22.3	82.5	324.7
2002.....	16.1	*	*	*	0.2	0.5	1.2	2.8	7.5	23.7	85.1	331.1
2001.....	16.1	*	*	*	0.2	0.5	1.0	2.9	8.1	25.0	82.9	337.9
2000.....	16.1	*	*	*	0.3	0.4	1.2	2.6	8.0	24.0	84.3	345.2
1999.....	16.9	*	*	*	0.4	0.5	1.2	2.9	9.3	26.5	87.0	357.9
Black, both sexes												
2004.....	12.1	*	*	*	0.6	1.6	4.1	9.9	23.9	48.2	114.1	315.6
2003.....	12.4	*	*	*	0.7	1.7	3.6	10.5	23.5	55.2	115.1	327.3
2002.....	12.5	*	*	*	0.6	1.6	3.7	10.3	26.1	51.4	123.5	337.6
2001.....	12.6	*	*	*	0.7	1.5	3.6	10.5	24.9	54.0	127.6	340.2
2000.....	12.3	*	*	*	0.7	1.4	3.6	9.9	25.2	52.9	121.6	346.3
1999.....	13.0	*	*	*	0.8	1.7	3.6	10.6	27.1	56.3	132.6	359.0
Black, male												
2004.....	11.9	*	*	*	0.8	1.7	4.5	12.4	31.8	63.3	136.7	332.8
2003.....	12.0	*	*	*	0.9	1.6	4.0	13.3	31.2	72.3	136.2	307.6
2002.....	11.9	*	*	*	0.8	1.9	4.3	13.4	34.0	64.7	140.3	336.6
2001.....	12.2	*	*	*	1.0	1.8	4.1	13.1	33.7	65.9	158.6	357.4
2000.....	11.7	*	*	*	*	1.5	4.2	13.5	33.6	67.9	132.6	361.0
1999.....	12.9	*	*	*	1.0	1.9	4.0	13.5	36.6	71.8	163.6	411.3
Black, female												
2004.....	12.3	*	*	*	*	1.6	3.7	7.7	17.5	37.4	101.3	308.7
2003.....	12.8	*	*	*	*	1.7	3.2	8.1	17.4	43.0	103.2	335.0
2002.....	13.1	*	*	*	*	1.4	3.1	7.6	19.7	42.1	114.0	338.0
2001.....	12.9	*	*	*	*	1.1	3.2	8.3	17.9	45.7	110.2	333.8
2000.....	12.8	*	*	*	0.8	1.2	3.1	6.7	18.4	42.4	115.4	340.8
1999.....	13.1	*	*	*	*	1.5	3.3	8.1	19.4	45.4	115.3	339.5

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Heart failure (I50)												
All races 3/, both sexes												
2004.....	19.5	0.7	*	*	0.1	0.2	0.7	2.5	8.4	32.2	131.2	621.2
2003.....	19.8	0.9	*	*	0.1	0.2	0.7	2.4	8.8	33.5	133.2	641.1
2002.....	19.6	0.6	*	*	0.1	0.3	0.6	2.5	8.8	33.8	132.8	645.0
2001.....	20.0	0.6	*	*	0.1	0.2	0.7	2.4	9.4	34.3	136.3	674.6
2000.....	19.8	0.8	*	0.1	0.1	0.2	0.6	2.2	9.1	34.8	136.1	684.9
1999.....	19.7	*	*	*	0.1	0.2	0.6	2.1	8.9	35.4	137.2	681.9
All races 3/, male												
2004.....	15.4	*	*	*	*	0.3	0.8	3.0	10.1	37.5	147.0	605.5
2003.....	15.7	1.4	*	*	0.1	0.3	0.9	2.9	10.8	39.7	148.6	629.8
2002.....	15.3	*	*	*	*	0.3	0.8	3.1	10.5	39.2	149.0	621.4
2001.....	15.5	*	*	*	*	0.3	0.8	3.1	11.1	40.3	153.8	646.4
2000.....	15.3	1.1	*	*	0.1	0.2	0.7	2.7	10.9	41.1	153.7	672.2
1999.....	15.2	*	*	*	0.1	0.2	0.7	2.7	10.8	42.5	152.0	667.0
All races 3/, female												
2004.....	23.4	*	*	*	*	0.2	0.6	2.0	6.8	27.8	120.5	628.2
2003.....	23.7	*	*	*	*	0.2	0.5	1.9	7.0	28.3	122.9	646.1
2002.....	23.7	*	*	*	*	0.2	0.5	1.8	7.2	29.2	122.0	655.3
2001.....	24.3	*	*	*	*	0.2	0.6	1.8	7.8	29.3	124.7	686.4
2000.....	24.1	*	*	*	*	0.1	0.5	1.6	7.5	29.6	124.6	690.0
1999.....	24.0	*	*	*	*	0.2	0.5	1.6	7.1	29.6	127.6	687.9
White, both sexes												
2004.....	21.5	0.6	*	*	0.1	0.2	0.5	1.9	7.4	31.4	132.4	641.5
2003.....	21.8	0.6	*	*	0.1	0.2	0.5	1.9	7.8	32.2	134.3	660.4
2002.....	21.6	*	*	*	*	0.2	0.5	1.9	8.0	32.5	133.3	664.5
2001.....	22.1	*	*	*	*	0.2	0.5	2.0	8.5	33.0	137.0	695.1
2000.....	21.8	*	*	*	*	0.1	0.4	1.8	8.2	33.4	136.7	703.7
1999.....	21.6	*	*	*	*	0.1	0.5	1.7	8.0	34.1	136.7	699.2
White, male												
2004.....	16.8	*	*	*	*	0.2	0.7	2.3	8.8	36.8	148.2	625.5
2003.....	17.1	*	*	*	*	0.2	0.6	2.3	9.6	38.6	150.4	650.8
2002.....	16.7	*	*	*	*	0.2	0.6	2.5	9.5	37.7	150.3	641.9
2001.....	16.9	*	*	*	*	0.2	0.6	2.6	10.3	39.0	154.6	666.5
2000.....	16.7	*	*	*	*	0.1	0.6	2.2	10.0	40.0	154.6	690.0
1999.....	16.4	*	*	*	*	*	0.6	2.1	9.8	41.3	152.1	680.0

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Heart failure (I50)												
White, female												
2004.....	26.1	*	*	*	*	0.1	0.4	1.5	6.0	26.8	121.7	648.7
2003.....	26.3	*	*	*	*	0.1	0.3	1.4	6.1	26.8	123.4	664.7
2002.....	26.4	*	*	*	*	0.1	0.3	1.4	6.5	28.1	121.8	674.3
2001.....	27.1	*	*	*	*	0.2	0.4	1.4	6.9	28.0	125.3	707.1
2000.....	26.7	*	*	*	*	*	0.3	1.3	6.6	27.9	124.9	709.3
1999.....	26.6	*	*	*	*	*	0.3	1.2	6.3	28.0	126.6	707.0
Black, both sexes												
2004.....	13.5	*	*	*	*	0.7	1.9	7.5	19.4	48.2	150.4	506.5
2003.....	14.0	*	*	*	*	0.7	2.3	6.8	20.6	54.3	150.4	531.8
2002.....	13.7	*	*	*	*	0.8	1.8	7.2	18.6	53.2	156.9	528.0
2001.....	13.8	*	*	*	*	0.6	1.9	6.4	19.6	54.5	157.3	543.8
2000.....	13.6	*	*	*	*	0.4	1.6	5.7	18.6	55.4	154.4	558.4
1999.....	13.8	*	*	*	*	0.5	1.6	6.4	18.9	56.4	163.4	552.9
Black, male												
2004.....	11.8	*	*	*	*	0.9	2.2	9.3	25.0	55.7	176.2	503.7
2003.....	11.8	*	*	*	*	0.8	2.7	8.3	26.3	63.2	163.9	513.0
2002.....	11.5	*	*	*	*	1.0	2.2	9.0	22.9	64.7	174.4	479.2
2001.....	11.6	*	*	*	*	0.9	2.2	7.4	22.2	65.3	183.3	541.2
2000.....	11.2	*	*	*	*	*	1.7	7.1	21.4	61.9	173.5	581.2
1999.....	11.8	*	*	*	*	*	1.9	8.4	22.9	65.2	179.2	603.5
Black, female												
2004.....	15.1	*	*	*	*	*	1.7	5.9	15.0	42.8	135.8	507.6
2003.....	15.9	*	*	*	*	*	1.9	5.6	16.0	48.0	142.7	539.2
2002.....	15.8	*	*	*	*	*	1.5	5.6	15.1	45.1	147.0	546.7
2001.....	15.8	*	*	*	*	*	1.7	5.5	17.6	46.9	142.6	544.8
2000.....	15.7	*	*	*	*	*	1.5	4.5	16.4	50.9	143.7	549.9
1999.....	15.7	*	*	*	*	*	1.4	4.7	15.7	50.2	154.6	534.0

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Congestive heart failure (I50.0)												
All races 3/, both sexes												
2004.....	18.1	*	*	*	0.0	0.2	0.6	2.1	7.2	29.1	122.3	588.9
2003.....	18.4	0.6	*	*	0.1	0.2	0.6	2.0	7.7	30.3	124.4	608.2
2002.....	18.2	*	*	*	*	0.2	0.5	2.0	7.6	30.3	123.7	609.0
2001.....	18.5	*	*	*	0.1	0.2	0.5	2.1	8.0	31.1	126.6	636.1
2000.....	18.3	*	*	*	0.1	0.1	0.5	1.8	7.9	31.0	126.4	644.4
1999.....	18.2	*	*	*	0.1	0.1	0.5	1.7	7.4	31.9	127.3	642.7
All races 3/, male												
2004.....	14.2	*	*	*	*	0.2	0.7	2.5	8.7	33.6	136.4	571.8
2003.....	14.5	*	*	*	*	0.2	0.7	2.4	9.3	35.8	138.3	594.6
2002.....	14.0	*	*	*	*	0.3	0.6	2.5	8.9	35.2	138.1	584.6
2001.....	14.2	*	*	*	*	0.2	0.6	2.6	9.3	36.2	141.2	609.3
2000.....	14.0	*	*	*	*	0.1	0.5	2.2	9.3	36.4	142.4	632.2
1999.....	13.9	*	*	*	*	0.1	0.6	2.2	8.9	38.0	140.5	628.2
All races 3/, female												
2004.....	21.9	*	*	*	*	0.1	0.5	1.7	5.8	25.3	112.8	596.6
2003.....	22.3	*	*	*	*	0.2	0.5	1.6	6.2	25.7	115.2	614.2
2002.....	22.2	*	*	*	*	0.2	0.4	1.5	6.3	26.3	114.1	619.6
2001.....	22.8	*	*	*	*	0.1	0.4	1.6	6.8	26.8	117.0	647.3
2000.....	22.4	*	*	*	*	0.1	0.4	1.4	6.6	26.6	116.1	649.4
1999.....	22.4	*	*	*	*	0.1	0.4	1.3	6.1	26.9	118.7	648.5
White, both sexes												
2004.....	20.1	*	*	*	*	0.1	0.4	1.6	6.3	28.5	123.8	609.2
2003.....	20.4	*	*	*	*	0.1	0.4	1.6	6.8	29.4	125.8	628.4
2002.....	20.1	*	*	*	*	0.1	0.4	1.5	6.8	29.3	124.5	628.8
2001.....	20.5	*	*	*	*	0.1	0.4	1.7	7.3	30.0	127.7	656.5
2000.....	20.2	*	*	*	*	0.1	0.3	1.4	7.2	29.9	127.3	663.6
1999.....	20.0	*	*	*	*	0.1	0.4	1.3	6.7	30.7	127.2	660.1
White, male												
2004.....	15.6	*	*	*	*	0.2	0.5	2.0	7.5	33.1	137.9	591.7
2003.....	15.9	*	*	*	*	0.2	0.5	1.9	8.2	35.0	140.4	616.2
2002.....	15.4	*	*	*	*	0.2	0.5	1.9	8.0	33.8	139.8	605.8
2001.....	15.5	*	*	*	*	0.2	0.5	2.2	8.7	35.1	142.6	629.0
2000.....	15.4	*	*	*	*	*	0.4	1.8	8.6	35.7	143.7	650.5
1999.....	15.0	*	*	*	*	*	0.5	1.6	8.1	37.0	140.7	641.7

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Congestive heart failure (I50.0)												
White, female												
2004.....	24.6	*	*	*	*	*	0.3	1.3	5.2	24.5	114.2	617.1
2003.....	24.9	*	*	*	*	*	0.3	1.2	5.4	24.5	116.0	633.9
2002.....	24.8	*	*	*	*	*	0.3	1.1	5.7	25.5	114.2	638.8
2001.....	25.4	*	*	*	*	0.1	0.3	1.2	5.9	25.7	117.8	667.9
2000.....	25.0	*	*	*	*	*	0.3	1.1	5.8	25.0	116.5	668.8
1999.....	24.9	*	*	*	*	*	0.3	0.9	5.4	25.5	118.3	667.4
Black, both sexes												
2004.....	12.1	*	*	*	*	0.5	1.5	6.1	16.5	42.4	136.0	468.7
2003.....	12.4	*	*	*	*	0.6	1.9	5.7	17.8	48.0	136.2	484.3
2002.....	12.3	*	*	*	*	0.8	1.6	6.0	16.2	47.4	142.1	482.6
2001.....	12.4	*	*	*	*	0.5	1.6	5.5	16.6	48.9	142.0	502.4
2000.....	12.1	*	*	*	*	*	1.3	4.9	16.1	48.5	140.0	510.2
1999.....	12.4	*	*	*	*	0.5	1.3	5.6	15.9	50.4	148.4	509.5
Black, male												
2004.....	10.3	*	*	*	*	0.8	1.6	7.3	21.6	48.2	158.4	466.9
2003.....	10.4	*	*	*	*	*	2.3	6.7	22.4	56.2	146.1	466.1
2002.....	10.2	*	*	*	*	1.0	1.8	7.4	19.9	59.4	155.2	428.8
2001.....	10.2	*	*	*	*	*	1.9	6.4	18.4	57.6	160.4	500.4
2000.....	9.8	*	*	*	*	*	1.3	6.2	18.1	52.7	154.7	528.2
1999.....	10.5	*	*	*	*	*	1.6	7.2	18.8	57.8	164.2	554.2
Black, female												
2004.....	13.6	*	*	*	*	*	1.4	5.1	12.4	38.3	123.4	469.4
2003.....	14.4	*	*	*	*	*	1.7	4.8	14.0	42.3	130.6	491.4
2002.....	14.2	*	*	*	*	*	1.4	4.8	13.3	38.9	134.6	503.2
2001.....	14.4	*	*	*	*	*	1.4	4.7	15.2	42.8	131.7	503.1
2000.....	14.2	*	*	*	*	*	1.2	3.9	14.5	45.5	131.8	503.5
1999.....	14.2	*	*	*	*	*	1.0	4.2	13.5	45.2	139.6	492.8

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified heart failure (I50.1-I50.9)												
All races 3/, both sexes												
2004.....	1.3	*	*	*	*	*	0.1	0.4	1.2	3.2	8.9	32.3
2003.....	1.3	*	*	*	*	*	0.1	0.4	1.2	3.1	8.7	32.9
2002.....	1.4	*	*	*	*	*	0.1	0.5	1.2	3.4	9.1	36.1
2001.....	1.4	*	*	*	*	0.1	0.1	0.4	1.4	3.2	9.7	38.5
2000.....	1.5	0.6	*	*	*	*	0.1	0.4	1.2	3.8	9.6	40.5
1999.....	1.5	*	*	*	*	*	0.1	0.4	1.4	3.6	9.9	39.2
All races 3/, male												
2004.....	1.2	*	*	*	*	*	0.2	0.5	1.5	4.0	10.6	33.7
2003.....	1.2	*	*	*	*	*	0.2	0.5	1.6	3.8	10.3	35.2
2002.....	1.3	*	*	*	*	*	0.2	0.7	1.6	4.0	10.9	36.8
2001.....	1.3	*	*	*	*	*	0.2	0.5	1.8	4.1	12.7	37.2
2000.....	1.3	*	*	*	*	*	0.2	0.5	1.5	4.7	11.3	40.0
1999.....	1.3	*	*	*	*	*	0.2	0.5	1.9	4.5	11.6	38.8
All races 3/, female												
2004.....	1.4	*	*	*	*	*	0.1	0.3	1.0	2.5	7.7	31.6
2003.....	1.4	*	*	*	*	*	*	0.3	0.8	2.6	7.7	31.9
2002.....	1.5	*	*	*	*	*	*	0.3	0.9	2.9	7.9	35.7
2001.....	1.6	*	*	*	*	*	0.1	0.2	1.0	2.4	7.7	39.1
2000.....	1.7	*	*	*	*	*	*	0.3	0.9	3.1	8.5	40.6
1999.....	1.6	*	*	*	*	*	0.1	0.3	1.0	2.7	8.8	39.4
White, both sexes												
2004.....	1.4	*	*	*	*	*	0.1	0.3	1.0	2.9	8.6	32.3
2003.....	1.4	*	*	*	*	*	0.1	0.3	1.0	2.9	8.5	32.0
2002.....	1.5	*	*	*	*	*	0.1	0.4	1.2	3.2	8.7	35.6
2001.....	1.5	*	*	*	*	*	0.1	0.3	1.2	3.0	9.3	38.7
2000.....	1.5	*	*	*	*	*	0.1	0.3	1.1	3.6	9.3	40.2
1999.....	1.5	*	*	*	*	*	0.1	0.4	1.3	3.4	9.5	39.2
White, male												
2004.....	1.2	*	*	*	*	*	0.1	0.4	1.3	3.7	10.3	33.8
2003.....	1.2	*	*	*	*	*	0.2	0.4	1.4	3.6	10.0	34.6
2002.....	1.3	*	*	*	*	*	0.1	0.5	1.5	3.9	10.5	36.0
2001.....	1.4	*	*	*	*	*	0.2	0.4	1.6	3.9	12.0	37.5
2000.....	1.3	*	*	*	*	*	0.2	0.5	1.4	4.4	10.9	39.5
1999.....	1.3	*	*	*	*	*	0.1	0.5	1.8	4.4	11.4	38.3

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other and unspecified heart failure (I50.1-I50.9)

White, female

2004.....	1.5	*	*	*	*	*	*	0.2	0.8	2.3	7.4	31.6
2003.....	1.5	*	*	*	*	*	*	0.2	0.7	2.3	7.5	30.8
2002.....	1.6	*	*	*	*	*	*	0.2	0.8	2.6	7.6	35.5
2001.....	1.7	*	*	*	*	*	*	0.2	0.9	2.3	7.5	39.2
2000.....	1.8	*	*	*	*	*	*	0.2	0.8	2.9	8.3	40.5
1999.....	1.7	*	*	*	*	*	*	0.3	0.9	2.6	8.3	39.5

Black, both sexes

2004.....	1.5	*	*	*	*	*	0.4	1.3	3.0	5.8	14.4	37.8
2003.....	1.5	*	*	*	*	*	*	1.1	2.8	6.3	14.2	47.6
2002.....	1.4	*	*	*	*	*	*	1.2	2.3	5.8	14.8	45.4
2001.....	1.4	*	*	*	*	*	*	0.9	3.0	5.6	15.2	41.5
2000.....	1.5	*	*	*	*	*	0.3	0.8	2.5	6.9	14.4	48.2
1999.....	1.4	*	*	*	*	*	0.4	0.8	3.0	6.0	15.0	43.4

Black, male

2004.....	1.5	*	*	*	*	*	*	1.9	3.4	7.5	17.8	36.8
2003.....	1.5	*	*	*	*	*	*	1.5	3.9	7.0	17.8	46.9
2002.....	1.3	*	*	*	*	*	*	1.7	3.0	5.3	19.2	50.4
2001.....	1.4	*	*	*	*	*	*	1.0	3.8	7.7	23.0	40.8
2000.....	1.4	*	*	*	*	*	*	*	3.3	9.2	18.8	53.0
1999.....	1.3	*	*	*	*	*	*	1.2	4.1	7.4	15.1	49.2

Black, female

2004.....	1.5	*	*	*	*	*	*	0.8	2.6	4.5	12.4	38.2
2003.....	1.6	*	*	*	*	*	*	0.8	2.0	5.7	12.1	47.8
2002.....	1.5	*	*	*	*	*	*	0.8	1.8	6.2	12.3	43.5
2001.....	1.4	*	*	*	*	*	*	0.8	2.3	4.1	10.9	41.7
2000.....	1.5	*	*	*	*	*	*	*	1.9	5.4	11.9	46.4
1999.....	1.5	*	*	*	*	*	*	*	2.2	5.0	15.0	41.2

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
All races 3/, both sexes												
2004.....	0.2	0.6	0.2	0.1	0.1	0.1	0.2	0.3	0.2	0.3	0.4	2.0
2003.....	0.2	0.7	0.2	0.1	0.1	0.2	0.2	0.2	0.2	0.1	0.4	2.2
2002.....	0.3	0.8	0.1	*	0.2	0.2	0.3	0.3	0.3	0.5	0.9	3.2
2001.....	0.3	0.6	0.3	0.1	0.1	0.2	0.3	0.3	0.2	0.4	0.5	2.0
2000.....	0.2	0.6	0.1	0.1	0.1	0.2	0.3	0.2	0.3	0.3	0.6	2.4
1999.....	0.3	0.7	0.2	0.1	0.2	0.2	0.3	0.3	0.2	0.3	0.5	2.4
All races 3/, male												
2004.....	0.2	*	*	*	0.2	0.1	0.3	0.3	0.2	0.2	0.4	1.7
2003.....	0.2	*	*	0.1	0.2	0.2	0.2	0.2	0.3	*	0.5	2.6
2002.....	0.3	*	*	*	0.1	0.2	0.3	0.3	0.3	0.6	0.9	3.8
2001.....	0.3	*	0.3	*	0.1	0.2	0.3	0.3	0.2	0.5	0.8	2.2
2000.....	0.2	*	*	*	0.1	0.2	0.3	0.2	0.3	0.3	0.7	2.6
1999.....	0.3	*	*	*	0.2	0.3	0.3	0.3	0.2	0.4	0.6	2.1
All races 3/, female												
2004.....	0.2	*	*	*	*	0.1	0.2	0.3	0.2	0.3	0.3	2.2
2003.....	0.2	*	*	*	*	0.1	0.2	0.3	0.2	*	0.3	2.0
2002.....	0.3	*	*	*	0.2	0.1	0.3	0.3	0.3	0.4	0.9	3.0
2001.....	0.2	*	*	*	0.1	0.2	0.2	0.2	0.2	0.3	0.4	1.9
2000.....	0.2	*	*	*	0.1	0.2	0.2	0.2	0.3	0.3	0.5	2.4
1999.....	0.2	*	*	*	0.2	0.1	0.2	0.3	0.2	0.2	0.4	2.5
White, both sexes												
2004.....	0.2	*	*	*	0.1	0.1	0.2	0.2	0.1	0.2	0.3	2.1
2003.....	0.2	*	*	0.1	0.1	0.1	0.2	0.2	0.2	0.2	0.4	2.2
2002.....	0.3	*	*	*	0.1	0.2	0.2	0.3	0.3	0.5	0.9	3.2
2001.....	0.2	*	0.2	*	0.1	0.1	0.3	0.2	0.2	0.4	0.5	2.1
2000.....	0.2	*	*	*	0.1	0.2	0.2	0.2	0.3	0.3	0.5	2.6
1999.....	0.2	*	*	*	0.2	0.2	0.2	0.2	0.2	0.3	0.5	2.5
White, male												
2004.....	0.2	*	*	*	0.2	0.2	0.2	0.2	*	*	0.4	1.8
2003.....	0.2	*	*	*	0.1	0.2	0.2	0.2	0.2	*	0.6	2.8
2002.....	0.3	*	*	*	*	0.2	0.3	0.3	0.3	0.6	0.9	4.0
2001.....	0.2	*	*	*	*	0.1	0.3	0.2	0.2	0.5	0.8	2.2
2000.....	0.2	*	*	*	0.1	0.2	0.3	0.2	0.3	0.3	0.6	2.7
1999.....	0.2	*	*	*	0.2	0.2	0.3	0.2	0.2	0.4	0.6	2.0

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

All other and ill-defined forms of heart disease (I50.0-I51.3,I51.6-I51.9)

All races 3/, both sexes

2004.....	3.4	0.6	*	0.1	0.2	0.5	1.0	2.1	3.6	7.2	20.4	67.7
2003.....	3.4	0.5	*	0.0	0.2	0.4	1.0	1.9	3.6	7.6	20.5	67.8
2002.....	3.3	0.6	*	*	0.2	0.4	1.1	1.9	3.4	7.8	20.0	65.0
2001.....	3.2	*	*	0.1	0.2	0.4	1.0	1.8	3.4	7.3	18.9	66.3
2000.....	3.1	0.8	*	*	0.2	0.4	0.9	1.8	3.7	7.5	19.3	64.6
1999.....	2.9	0.8	*	*	0.2	0.4	0.9	1.6	3.5	6.9	17.8	61.5

All races 3/, male

2004.....	3.5	*	*	*	0.4	0.7	1.4	3.0	4.8	9.6	24.9	67.8
2003.....	3.4	*	*	*	0.4	0.7	1.5	2.5	4.9	9.3	24.6	72.1
2002.....	3.3	*	*	*	0.4	0.6	1.5	2.7	4.6	10.6	23.6	66.7
2001.....	3.2	*	*	*	0.3	0.7	1.5	2.6	4.7	9.9	22.1	66.4
2000.....	3.1	*	*	*	0.4	0.6	1.3	2.5	5.3	9.6	22.4	69.0
1999.....	3.0	*	*	*	0.3	0.6	1.3	2.3	4.9	9.4	21.8	64.3

All races 3/, female

2004.....	3.3	*	*	*	0.1	0.2	0.7	1.1	2.4	5.1	17.3	67.7
2003.....	3.3	*	*	*	0.1	0.2	0.5	1.2	2.3	6.1	17.8	66.0
2002.....	3.2	*	*	*	*	0.2	0.6	1.2	2.2	5.4	17.5	64.3
2001.....	3.1	*	*	*	*	0.2	0.5	1.1	2.2	5.2	16.8	66.3
2000.....	3.1	*	*	*	*	0.2	0.5	1.0	2.2	5.7	17.3	62.7
1999.....	2.8	*	*	*	*	0.2	0.5	0.9	2.1	4.8	15.2	60.4

White, both sexes

2004.....	3.5	*	*	*	0.2	0.4	1.0	1.9	3.3	6.7	20.0	68.1
2003.....	3.5	*	*	*	0.2	0.4	0.9	1.6	3.3	7.2	20.5	67.5
2002.....	3.4	*	*	*	0.2	0.3	1.0	1.6	3.1	7.5	19.9	64.5
2001.....	3.2	*	*	*	0.2	0.4	0.9	1.6	3.2	7.0	18.4	65.7
2000.....	3.2	0.7	*	*	0.2	0.4	0.7	1.6	3.4	7.0	18.8	64.7
1999.....	3.0	0.7	*	*	0.2	0.3	0.7	1.4	3.1	6.5	17.3	61.4

White, male

2004.....	3.6	*	*	*	0.3	0.6	1.4	2.8	4.5	9.0	24.5	68.2
2003.....	3.5	*	*	*	0.3	0.6	1.4	2.3	4.7	9.0	24.5	72.2
2002.....	3.4	*	*	*	0.3	0.5	1.4	2.4	4.2	10.3	23.3	66.9
2001.....	3.2	*	*	*	0.3	0.6	1.4	2.3	4.5	9.5	21.4	66.2
2000.....	3.2	*	*	*	0.3	0.6	1.1	2.3	5.0	9.0	21.6	69.2
1999.....	3.0	*	*	*	0.3	0.5	1.1	2.1	4.5	8.9	21.2	64.5

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

All other and ill-defined forms of heart disease (I50.0-I51.3,I51.6-I51.9)

White, female

2004.....	3.5	*	*	*	*	0.2	0.6	1.0	2.2	4.7	17.0	68.1
2003.....	3.5	*	*	*	*	*	0.5	1.0	2.1	5.7	17.8	65.4
2002.....	3.4	*	*	*	*	0.1	0.6	0.9	2.0	5.1	17.6	63.4
2001.....	3.3	*	*	*	*	0.2	0.4	0.8	2.0	4.8	16.3	65.5
2000.....	3.2	*	*	*	*	0.1	0.3	0.8	2.0	5.3	17.0	62.8
1999.....	2.9	*	*	*	*	*	0.4	0.7	1.8	4.5	14.7	60.2

Black, both sexes

2004.....	3.6	*	*	*	0.4	1.0	1.9	3.8	6.8	13.6	30.0	76.4
2003.....	3.5	*	*	*	0.5	1.0	1.9	3.9	6.5	12.8	25.3	80.5
2002.....	3.4	*	*	*	0.5	0.8	1.8	4.4	6.7	12.6	24.8	80.5
2001.....	3.4	*	*	*	0.4	0.9	2.1	4.1	6.2	12.0	27.1	83.8
2000.....	3.5	*	*	*	0.6	0.9	1.9	3.7	7.2	14.5	28.2	73.8
1999.....	3.3	*	*	*	0.4	0.9	2.0	3.4	7.8	12.2	25.6	68.9

Black, male

2004.....	4.0	*	*	*	*	1.6	2.4	5.2	9.3	18.6	39.3	75.5
2003.....	3.7	*	*	*	0.9	1.2	2.9	5.0	9.1	15.5	32.7	80.3
2002.....	3.8	*	*	*	0.9	1.2	2.5	5.8	9.3	17.1	32.7	77.2
2001.....	3.5	*	*	*	*	1.4	2.6	5.2	8.4	15.6	32.7	84.0
2000.....	3.9	*	*	*	1.0	1.3	2.5	5.4	10.1	19.0	36.0	80.7
1999.....	3.6	*	*	*	*	1.4	2.8	4.4	10.6	17.4	32.3	72.6

Black, female

2004.....	3.3	*	*	*	*	*	1.4	2.6	4.8	10.0	24.7	76.8
2003.....	3.3	*	*	*	*	0.7	1.0	2.9	4.4	10.9	21.0	80.6
2002.....	3.2	*	*	*	*	*	1.3	3.2	4.5	9.4	20.3	81.7
2001.....	3.3	*	*	*	*	*	1.6	3.1	4.5	9.4	24.0	83.8
2000.....	3.2	*	*	*	*	*	1.4	2.2	4.9	11.3	23.9	71.3
1999.....	3.0	*	*	*	*	*	1.3	2.6	5.6	8.5	21.9	67.5

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Cerebrovascular diseases (I60-I69)												
All races 3/, both sexes												
2004.....	51.1	3.1	0.3	0.2	0.5	1.4	5.4	14.9	34.3	107.8	386.2	1,245.9
2003.....	54.2	2.5	0.3	0.2	0.5	1.5	5.5	15.0	35.6	112.9	410.7	1,370.1
2002.....	56.4	2.9	0.3	0.2	0.4	1.4	5.4	15.1	37.2	120.3	431.0	1,445.9
2001.....	57.4	2.7	0.4	0.2	0.5	1.5	5.5	15.1	38.0	123.4	443.9	1,500.2
2000.....	59.6	3.3	0.3	0.2	0.5	1.5	5.8	16.0	41.0	128.6	461.3	1,589.2
1999.....	60.0	2.7	0.3	0.2	0.5	1.4	5.7	15.2	40.6	130.8	469.8	1,614.8
All races 3/, male												
2004.....	40.7	3.4	0.3	0.2	0.5	1.4	5.6	16.7	39.5	121.1	402.9	1,118.1
2003.....	42.9	2.8	0.3	0.2	0.5	1.6	5.8	16.7	40.8	127.8	431.4	1,236.0
2002.....	44.2	3.2	0.4	0.2	0.5	1.4	5.3	16.7	42.7	135.0	445.9	1,317.9
2001.....	45.2	3.1	0.3	0.2	0.5	1.6	5.7	16.7	43.4	140.4	467.3	1,380.2
2000.....	46.9	3.8	*	0.2	0.5	1.5	5.8	17.5	47.2	145.0	490.8	1,484.3
1999.....	47.1	3.3	0.3	0.2	0.5	1.5	5.8	16.7	46.5	147.3	500.4	1,512.8
All races 3/, female												
2004.....	61.2	2.8	*	0.2	0.5	1.4	5.1	13.1	29.5	96.6	374.9	1,303.4
2003.....	65.1	2.2	0.3	0.1	0.5	1.4	5.3	13.4	30.9	100.5	396.8	1,429.4
2002.....	68.2	2.5	0.3	0.2	0.3	1.4	5.5	13.6	32.1	108.1	421.2	1,501.5
2001.....	69.2	2.3	0.4	0.2	0.5	1.5	5.4	13.6	32.9	109.3	428.6	1,550.4
2000.....	71.8	2.7	0.4	0.2	0.5	1.5	5.7	14.5	35.3	115.1	442.1	1,632.0
1999.....	72.3	2.1	0.3	0.2	0.5	1.4	5.6	13.8	35.1	117.2	449.8	1,656.0
White, both sexes												
2004.....	53.7	2.5	0.2	0.2	0.5	1.2	4.4	11.4	28.6	99.0	377.7	1,257.8
2003.....	57.0	2.0	0.2	0.2	0.5	1.2	4.2	11.4	29.4	103.7	403.0	1,382.2
2002.....	59.5	2.3	0.3	0.2	0.4	1.2	4.2	11.6	31.4	110.7	423.5	1,459.5
2001.....	60.5	1.9	0.3	0.2	0.4	1.3	4.4	11.4	31.7	113.0	436.4	1,511.2
2000.....	62.8	2.5	0.3	0.2	0.5	1.3	4.5	12.4	34.8	119.4	452.4	1,601.6
1999.....	63.3	2.3	0.3	0.2	0.4	1.3	4.4	11.7	34.0	121.1	461.6	1,631.6
White, male												
2004.....	41.8	2.8	*	0.2	0.5	1.2	4.7	12.8	32.4	110.8	393.7	1,129.3
2003.....	44.2	2.0	*	0.2	0.5	1.3	4.4	12.9	33.3	117.3	422.4	1,247.0
2002.....	45.7	2.6	0.4	0.2	0.5	1.2	4.2	12.9	35.6	123.8	437.5	1,327.4
2001.....	46.6	2.2	*	0.2	0.5	1.4	4.5	12.7	36.1	128.5	458.8	1,386.2
2000.....	48.4	3.0	*	0.2	0.5	1.4	4.5	13.6	39.7	133.8	480.0	1,490.7
1999.....	48.7	2.8	*	0.2	0.5	1.3	4.6	12.8	38.9	136.0	490.3	1,522.7

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Cerebrovascular diseases (I60-I69)												
White, female												
2004.....	65.3	2.2	*	0.2	0.4	1.1	4.1	10.1	25.1	89.0	366.8	1,315.7
2003.....	69.5	2.0	*	0.1	0.4	1.2	4.0	10.0	25.8	92.1	389.9	1,442.1
2002.....	73.0	2.0	*	0.2	0.3	1.2	4.3	10.4	27.4	99.5	414.1	1,516.9
2001.....	74.0	1.6	0.4	0.2	0.4	1.3	4.2	10.2	27.6	99.9	421.6	1,563.5
2000.....	76.9	2.0	*	0.2	0.4	1.3	4.4	11.2	30.2	107.3	434.2	1,646.7
1999.....	77.6	1.9	*	0.1	0.4	1.2	4.3	10.7	29.4	108.6	442.8	1,675.4
Black, both sexes												
2004.....	46.9	6.0	*	*	0.8	3.1	12.3	39.0	83.9	195.3	511.7	1,211.2
2003.....	49.3	4.9	*	*	0.9	3.1	13.8	41.0	89.8	201.1	531.1	1,348.7
2002.....	50.0	5.5	*	0.4	0.6	3.1	12.8	40.7	88.0	215.1	552.3	1,383.9
2001.....	51.1	6.6	*	*	0.8	2.9	13.2	42.6	91.1	222.5	560.3	1,452.7
2000.....	52.5	7.5	*	*	0.7	2.9	14.5	43.4	93.8	222.9	588.8	1,530.0
1999.....	52.2	4.6	*	*	0.7	2.7	14.3	41.5	96.3	231.9	590.5	1,492.5
Black, male												
2004.....	41.5	6.8	*	*	1.0	2.9	12.8	44.8	107.4	235.2	551.0	1,061.0
2003.....	43.2	5.9	*	*	0.8	3.3	14.3	46.9	112.1	237.4	588.9	1,180.3
2002.....	43.5	5.8	*	*	0.7	2.9	12.2	46.5	110.3	262.9	587.8	1,252.2
2001.....	44.6	8.1	*	*	0.7	3.1	13.2	48.8	111.9	269.2	613.9	1,349.1
2000.....	46.1	8.6	*	*	*	2.7	14.7	49.5	115.4	268.5	659.2	1,458.8
1999.....	45.9	*	*	*	*	2.8	14.7	48.6	117.9	274.7	650.6	1,448.3
Black, female												
2004.....	51.9	*	*	*	*	3.3	11.8	33.9	65.0	166.8	489.5	1,270.7
2003.....	54.8	*	*	*	1.1	2.8	13.3	36.0	71.8	175.3	498.3	1,414.2
2002.....	55.8	*	*	*	*	3.3	13.4	35.7	70.1	181.2	532.2	1,434.3
2001.....	56.9	*	*	*	1.0	2.8	13.1	37.3	74.4	189.5	530.3	1,491.2
2000.....	58.3	*	*	*	0.8	3.1	14.4	38.1	76.4	190.9	549.2	1,556.5
1999.....	57.9	*	*	*	0.8	2.7	13.9	35.3	78.9	201.7	556.8	1,509.1

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Subarachnoid hemorrhage (I60)												
All races 3/, both sexes												
2004.....	2.1	*	*	0.0	0.2	0.5	1.7	3.1	3.9	5.3	8.3	12.2
2003.....	2.1	*	*	*	0.1	0.5	1.7	3.1	4.0	5.3	8.7	11.5
2002.....	2.1	*	*	0.0	0.1	0.5	1.8	3.1	4.2	5.8	8.9	12.2
2001.....	2.2	*	*	*	0.2	0.5	1.8	3.2	4.2	5.8	8.8	13.2
2000.....	2.3	*	*	0.1	0.2	0.6	1.9	3.5	4.9	5.8	9.4	14.1
1999.....	2.3	*	*	*	0.2	0.6	2.1	3.5	4.7	6.2	9.4	13.1
All races 3/, male												
2004.....	1.5	*	*	*	0.2	0.5	1.3	2.7	3.0	3.6	6.0	9.3
2003.....	1.5	*	*	*	0.2	0.5	1.4	2.4	3.1	4.0	6.2	9.1
2002.....	1.5	*	*	*	0.2	0.4	1.4	2.3	3.3	4.4	5.8	9.1
2001.....	1.5	*	*	*	0.2	0.5	1.4	2.6	3.2	4.3	6.4	9.5
2000.....	1.6	*	*	*	0.2	0.6	1.5	2.7	3.5	4.2	6.8	11.2
1999.....	1.6	*	*	*	0.2	0.6	1.8	2.8	3.4	4.3	6.7	9.0
All races 3/, female												
2004.....	2.6	*	*	*	0.2	0.5	2.0	3.5	4.8	6.7	9.9	13.5
2003.....	2.6	*	*	*	0.1	0.5	1.9	3.7	4.8	6.5	10.4	12.5
2002.....	2.8	*	*	*	0.1	0.6	2.1	3.8	5.1	6.9	11.0	13.6
2001.....	2.8	*	*	*	0.2	0.5	2.1	3.7	5.2	7.0	10.3	14.8
2000.....	3.0	*	*	*	0.2	0.6	2.3	4.3	6.2	7.0	11.2	15.3
1999.....	3.0	*	*	*	0.2	0.6	2.4	4.1	5.9	7.7	11.1	14.8
White, both sexes												
2004.....	2.1	*	*	*	0.1	0.4	1.5	2.8	3.7	5.2	8.1	12.4
2003.....	2.0	*	*	*	0.1	0.5	1.4	2.8	3.8	5.1	8.3	11.1
2002.....	2.1	*	*	*	0.1	0.4	1.6	2.8	4.0	5.5	8.7	12.0
2001.....	2.1	*	*	*	0.1	0.5	1.5	2.8	4.0	5.4	8.6	12.7
2000.....	2.3	*	*	*	0.2	0.6	1.7	3.2	4.7	5.4	9.2	13.6
1999.....	2.3	*	*	*	0.2	0.5	1.8	3.1	4.4	6.1	9.2	12.6
White, male												
2004.....	1.5	*	*	*	0.1	0.4	1.2	2.4	2.8	3.6	5.9	9.9
2003.....	1.5	*	*	*	0.2	0.5	1.2	2.3	2.8	4.0	5.9	8.6
2002.....	1.5	*	*	*	0.2	0.4	1.3	2.2	3.1	4.2	5.7	9.1
2001.....	1.5	*	*	*	0.2	0.5	1.3	2.3	2.9	4.1	6.4	9.2
2000.....	1.6	*	*	*	0.2	0.6	1.4	2.5	3.3	4.0	6.9	11.8
1999.....	1.6	*	*	*	0.2	0.6	1.6	2.6	3.3	4.2	6.8	9.0

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Subarachnoid hemorrhage (I60)												
White, female												
2004.....	2.6	*	*	*	*	0.4	1.8	3.2	4.6	6.5	9.6	13.5
2003.....	2.6	*	*	*	*	0.5	1.6	3.3	4.8	6.1	10.0	12.2
2002.....	2.7	*	*	*	*	0.5	1.8	3.3	4.8	6.6	10.8	13.3
2001.....	2.7	*	*	*	*	0.5	1.8	3.3	5.0	6.5	10.1	14.2
2000.....	2.9	*	*	*	0.2	0.6	1.9	3.9	6.0	6.6	10.8	14.3
1999.....	2.9	*	*	*	0.2	0.5	2.1	3.6	5.5	7.6	10.9	14.1
Black, both sexes												
2004.....	2.2	*	*	*	*	0.9	2.9	5.0	5.9	5.3	8.8	9.0
2003.....	2.3	*	*	*	*	0.9	3.2	5.0	5.1	6.1	9.9	14.6
2002.....	2.4	*	*	*	*	1.0	3.2	5.4	5.8	6.4	8.6	12.8
2001.....	2.5	*	*	*	*	0.8	3.4	5.7	6.2	7.4	8.6	13.8
2000.....	2.7	*	*	*	*	0.9	3.6	6.1	6.2	8.0	10.8	17.2
1999.....	2.6	*	*	*	*	0.9	3.8	6.2	6.7	6.9	9.4	14.7
Black, male												
2004.....	1.6	*	*	*	*	*	2.3	4.1	5.1	3.7	5.6	*
2003.....	1.6	*	*	*	*	*	2.5	3.3	5.4	4.1	7.3	*
2002.....	1.6	*	*	*	*	0.8	2.1	3.3	5.2	6.1	5.7	*
2001.....	1.7	*	*	*	*	0.8	2.5	4.5	5.0	4.8	6.2	*
2000.....	1.7	*	*	*	*	*	2.6	4.4	4.2	6.3	7.0	*
1999.....	1.8	*	*	*	*	0.8	2.8	4.9	5.0	5.6	*	*
Black, female												
2004.....	2.8	*	*	*	*	1.1	3.5	5.7	6.5	6.3	10.7	11.8
2003.....	2.9	*	*	*	*	1.0	3.9	6.4	4.9	7.5	11.3	15.0
2002.....	3.1	*	*	*	*	1.2	4.1	7.3	6.4	6.6	10.2	15.6
2001.....	3.2	*	*	*	*	0.9	4.1	6.8	7.0	9.2	9.9	16.4
2000.....	3.5	*	*	*	*	1.2	4.4	7.7	7.8	9.2	13.0	22.8
1999.....	3.3	*	*	*	*	1.0	4.7	7.3	8.1	7.9	11.4	18.4

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Intracerebral and other intracranial hemorrhage (I61-I62)												
All races 3/, both sexes												
2004.....	9.4	1.3	*	*	0.2	0.5	2.1	6.2	11.6	27.1	71.3	124.3
2003.....	9.4	0.8	*	0.1	0.2	0.5	2.2	6.3	11.4	27.2	71.1	126.6
2002.....	9.4	1.0	*	0.1	0.2	0.5	2.1	6.0	11.7	28.1	72.3	127.2
2001.....	9.4	1.0	*	0.1	0.2	0.6	2.3	6.2	11.8	27.9	71.7	127.9
2000.....	9.7	1.1	*	0.1	0.2	0.5	2.3	6.5	12.9	29.3	74.1	137.9
1999.....	9.1	1.1	*	0.1	0.2	0.5	2.2	5.9	12.3	28.3	71.9	122.3
All races 3/, male												
2004.....	8.9	1.4	*	*	0.2	0.6	2.6	7.4	13.9	30.7	77.4	134.4
2003.....	8.9	*	*	*	0.2	0.5	2.5	7.6	13.3	31.1	77.8	137.4
2002.....	8.7	1.1	*	*	0.2	0.5	2.3	7.3	13.7	32.3	77.2	135.0
2001.....	8.8	1.2	*	*	0.3	0.6	2.6	7.5	13.9	32.0	78.6	134.2
2000.....	9.1	1.3	*	*	0.2	0.6	2.7	7.8	15.9	33.1	79.0	149.6
1999.....	8.5	1.3	*	*	0.2	0.5	2.6	7.0	14.7	32.0	75.9	130.7
All races 3/, female												
2004.....	9.8	1.3	*	*	0.2	0.5	1.6	5.1	9.4	24.2	67.1	119.8
2003.....	9.8	*	*	*	0.2	0.5	1.9	5.1	9.6	24.0	66.6	121.8
2002.....	10.0	1.0	*	*	0.1	0.4	1.8	4.8	9.8	24.6	69.0	123.8
2001.....	9.9	*	*	*	0.2	0.5	1.9	5.1	10.0	24.6	67.1	125.2
2000.....	10.4	*	*	*	0.2	0.4	2.0	5.3	10.2	26.1	70.9	133.1
1999.....	9.8	*	*	*	0.1	0.4	1.8	4.9	10.1	25.2	69.2	118.9
White, both sexes												
2004.....	9.4	1.0	*	*	0.2	0.4	1.6	4.4	9.5	25.7	71.0	126.5
2003.....	9.3	*	*	0.1	0.2	0.4	1.6	4.2	9.0	25.7	70.8	127.7
2002.....	9.5	0.9	*	0.1	0.1	0.4	1.4	4.2	9.9	26.5	72.4	129.2
2001.....	9.4	0.7	*	0.1	0.2	0.5	1.7	4.3	9.8	26.2	71.8	128.9
2000.....	9.8	0.7	*	*	0.2	0.4	1.6	4.6	10.8	28.0	73.9	139.4
1999.....	9.2	0.8	*	*	0.1	0.4	1.5	4.1	10.2	27.2	71.7	123.2
White, male												
2004.....	8.8	*	*	*	0.2	0.5	2.0	5.3	11.2	28.9	77.4	137.6
2003.....	8.6	*	*	*	0.2	0.4	1.8	5.3	10.4	29.2	77.7	138.9
2002.....	8.7	*	*	*	0.2	0.4	1.6	5.1	11.5	30.6	77.8	136.9
2001.....	8.6	*	*	*	0.3	0.5	1.9	5.2	11.7	30.0	78.8	134.9
2000.....	8.9	*	*	*	0.2	0.5	1.9	5.6	13.1	31.4	79.0	150.5
1999.....	8.3	*	*	*	0.2	0.4	1.7	4.9	12.2	31.0	75.2	131.7

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Intracerebral and other intracranial hemorrhage (I61-I62)												
White, female												
2004.....	10.1	*	*	*	0.2	0.3	1.1	3.4	7.9	22.9	66.6	121.5
2003.....	10.0	*	*	*	0.2	0.4	1.3	3.2	7.7	22.6	66.2	122.8
2002.....	10.2	*	*	*	*	0.3	1.2	3.3	8.3	23.1	68.7	125.9
2001.....	10.1	*	*	*	0.1	0.4	1.4	3.3	8.1	22.9	67.1	126.3
2000.....	10.7	*	*	*	0.2	0.4	1.4	3.6	8.7	25.2	70.6	134.9
1999.....	10.0	*	*	*	*	0.3	1.2	3.4	8.2	24.0	69.4	119.7
Black, both sexes												
2004.....	10.2	*	*	*	0.3	1.3	5.5	19.1	28.3	39.6	72.2	95.6
2003.....	10.5	*	*	*	0.4	1.2	6.1	20.6	30.2	38.0	72.9	107.4
2002.....	9.7	*	*	*	*	1.2	6.0	18.9	25.8	39.2	69.4	98.3
2001.....	10.0	*	*	*	0.4	1.4	5.9	20.6	26.7	40.0	68.6	106.6
2000.....	10.3	*	*	*	*	1.2	7.1	21.0	29.7	39.0	69.0	117.0
1999.....	9.5	*	*	*	*	1.1	6.8	18.8	27.9	36.7	68.2	96.9
Black, male												
2004.....	10.7	*	*	*	*	1.3	6.6	22.6	38.0	47.1	76.6	100.3
2003.....	11.0	*	*	*	*	1.4	6.6	25.1	38.8	44.9	84.3	116.8
2002.....	10.1	*	*	*	*	1.3	6.8	24.1	33.1	47.2	68.8	101.8
2001.....	10.3	*	*	*	*	1.5	6.7	25.2	32.3	49.5	71.2	112.3
2000.....	10.9	*	*	*	*	1.4	8.3	25.7	40.6	46.7	73.0	130.3
1999.....	9.8	*	*	*	*	1.2	8.2	23.3	35.2	40.1	74.4	109.0
Black, female												
2004.....	9.8	*	*	*	*	1.3	4.5	16.0	20.5	34.2	69.7	93.7
2003.....	10.1	*	*	*	*	1.0	5.6	16.8	23.3	33.1	66.5	103.7
2002.....	9.4	*	*	*	*	1.1	5.3	14.5	20.0	33.4	69.7	96.9
2001.....	9.8	*	*	*	*	1.2	5.2	16.6	22.2	33.3	67.1	104.4
2000.....	9.8	*	*	*	*	1.0	6.1	16.9	20.8	33.6	66.8	112.0
1999.....	9.2	*	*	*	*	0.9	5.6	14.9	22.1	34.3	64.7	92.4

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Cerebral infarction (I63)												
All races 3/, both sexes												
2004.....	3.7	*	*	*	*	0.1	0.3	0.8	2.3	7.7	27.8	97.7
2003.....	4.2	*	*	*	*	0.1	0.3	0.9	2.4	8.4	31.3	112.2
2002.....	4.3	*	*	*	*	0.1	0.2	1.0	2.5	8.6	32.2	118.3
2001.....	4.4	*	*	*	*	0.1	0.3	0.9	2.4	9.4	33.9	124.8
2000.....	4.9	*	*	*	*	0.1	0.3	0.9	2.9	9.6	38.1	141.0
1999.....	5.3	*	*	*	*	0.1	0.3	0.9	3.1	11.3	41.2	153.1
All races 3/, male												
2004.....	3.0	*	*	*	*	0.1	0.3	1.0	3.0	9.0	30.1	88.5
2003.....	3.3	*	*	*	*	*	0.4	1.2	2.9	10.2	33.1	100.3
2002.....	3.4	*	*	*	*	0.1	0.2	1.2	3.1	9.6	35.4	108.7
2001.....	3.5	*	*	*	*	0.1	0.3	1.1	3.1	11.1	37.8	111.2
2000.....	3.9	*	*	*	*	*	0.4	1.1	3.6	11.4	42.3	133.3
1999.....	4.2	*	*	*	*	0.1	0.4	1.2	3.8	13.5	45.0	144.6
All races 3/, female												
2004.....	4.4	*	*	*	*	0.1	0.3	0.6	1.7	6.6	26.3	101.9
2003.....	5.0	*	*	*	*	*	0.2	0.7	1.8	6.9	30.1	117.5
2002.....	5.1	*	*	*	*	0.1	0.2	0.7	1.9	7.7	30.1	122.4
2001.....	5.3	*	*	*	*	*	0.3	0.8	1.8	8.0	31.3	130.4
2000.....	5.8	*	*	*	*	0.1	0.3	0.8	2.2	8.1	35.3	144.2
1999.....	6.3	*	*	*	*	*	0.3	0.7	2.5	9.6	38.8	156.6
White, both sexes												
2004.....	4.0	*	*	*	*	0.1	0.3	0.7	2.0	7.1	27.3	97.6
2003.....	4.4	*	*	*	*	0.1	0.2	0.8	1.9	7.9	30.9	113.7
2002.....	4.6	*	*	*	*	0.1	0.2	0.8	2.2	8.0	31.6	119.1
2001.....	4.8	*	*	*	*	0.1	0.2	0.8	2.1	8.8	33.4	126.0
2000.....	5.2	*	*	*	*	0.1	0.3	0.8	2.5	9.0	37.4	141.7
1999.....	5.7	*	*	*	*	0.1	0.3	0.8	2.7	10.5	40.6	154.4
White, male												
2004.....	3.2	*	*	*	*	*	0.3	0.8	2.6	8.2	29.5	87.9
2003.....	3.5	*	*	*	*	*	0.3	1.0	2.5	9.5	32.9	101.5
2002.....	3.6	*	*	*	*	*	0.2	1.1	2.8	8.9	34.8	108.6
2001.....	3.7	*	*	*	*	0.1	0.3	0.9	2.7	10.5	37.0	110.5
2000.....	4.1	*	*	*	*	*	0.3	0.9	3.2	10.6	41.2	133.1
1999.....	4.4	*	*	*	*	*	0.3	1.0	3.3	12.5	44.1	144.0

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Cerebral infarction (I63)												
White, female												
2004.....	4.8	*	*	*	*	*	0.2	0.5	1.5	6.2	25.8	102.0
2003.....	5.4	*	*	*	*	*	0.2	0.5	1.5	6.5	29.6	119.1
2002.....	5.5	*	*	*	*	*	0.2	0.6	1.7	7.2	29.5	123.7
2001.....	5.8	*	*	*	*	*	0.2	0.6	1.5	7.3	31.1	132.5
2000.....	6.3	*	*	*	*	*	0.2	0.7	1.9	7.6	35.0	145.2
1999.....	6.9	*	*	*	*	*	0.2	0.6	2.1	8.9	38.4	158.5
Black, both sexes												
2004.....	3.3	*	*	*	*	*	0.7	1.8	5.4	13.7	37.5	103.5
2003.....	3.5	*	*	*	*	*	0.8	2.1	6.0	14.2	40.1	107.7
2002.....	3.5	*	*	*	*	*	0.5	2.1	4.9	15.0	42.0	113.1
2001.....	3.7	*	*	*	*	*	0.7	2.1	5.7	16.5	42.1	121.3
2000.....	4.1	*	*	*	*	*	0.8	2.1	6.5	16.7	49.2	142.6
1999.....	4.4	*	*	*	*	*	0.7	2.2	6.6	20.4	53.0	150.8
Black, male												
2004.....	3.1	*	*	*	*	*	*	2.5	7.7	17.5	43.4	97.4
2003.....	3.1	*	*	*	*	*	0.9	2.4	8.0	18.2	43.2	97.0
2002.....	3.1	*	*	*	*	*	*	2.7	6.6	18.1	46.2	114.7
2001.....	3.3	*	*	*	*	*	*	2.5	7.3	19.0	53.9	127.1
2000.....	3.7	*	*	*	*	*	0.8	2.9	8.4	20.8	60.0	137.2
1999.....	4.0	*	*	*	*	*	*	2.9	8.3	26.0	63.3	162.9
Black, female												
2004.....	3.6	*	*	*	*	*	0.7	1.2	3.7	11.1	34.1	105.9
2003.....	3.9	*	*	*	*	*	0.7	1.7	4.4	11.4	38.4	111.8
2002.....	3.9	*	*	*	*	*	*	1.6	3.6	12.8	39.6	112.5
2001.....	4.0	*	*	*	*	*	0.8	1.8	4.3	14.7	35.5	119.1
2000.....	4.5	*	*	*	*	*	0.9	1.4	4.9	13.8	43.1	144.7
1999.....	4.7	*	*	*	*	*	0.8	1.6	5.3	16.4	47.2	146.3

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Stroke, not specified as hemorrhage or infarction (I64)												
All races 3/, both sexes												
2004.....	26.0	*	*	*	*	0.2	0.9	3.7	12.7	51.7	204.5	713.4
2003.....	28.4	*	*	*	0.1	0.3	1.0	3.7	14.1	54.8	223.0	799.7
2002.....	30.1	*	*	*	*	0.2	0.9	4.0	15.0	60.5	237.6	859.4
2001.....	31.1	*	*	*	*	0.2	0.9	3.8	15.6	62.8	249.4	906.2
2000.....	32.1	*	*	*	*	0.2	0.9	3.8	16.2	66.4	257.6	954.7
1999.....	32.6	*	*	*	*	0.2	0.8	3.8	16.3	66.5	263.9	982.4
All races 3/, male												
2004.....	19.7	*	*	*	*	0.2	1.1	4.4	14.9	58.9	209.0	615.5
2003.....	21.5	*	*	*	*	0.3	1.2	4.4	16.9	62.5	231.4	699.0
2002.....	22.6	*	*	*	*	0.2	1.0	4.6	18.0	68.8	241.0	762.8
2001.....	23.5	*	*	*	*	0.1	0.9	4.5	18.7	72.1	256.3	819.7
2000.....	24.2	*	*	*	*	0.2	0.9	4.7	19.4	75.1	270.6	863.2
1999.....	24.7	*	*	*	*	0.2	0.8	4.6	19.4	75.9	280.4	896.4
All races 3/, female												
2004.....	32.2	*	*	*	*	0.2	0.8	2.9	10.6	45.7	201.4	757.4
2003.....	35.0	*	*	*	*	0.2	0.8	3.0	11.5	48.3	217.4	844.2
2002.....	37.4	*	*	*	*	0.2	0.9	3.3	12.1	53.6	235.4	901.3
2001.....	38.5	*	*	*	*	0.2	0.8	3.0	12.7	55.1	244.9	942.4
2000.....	39.7	*	*	*	*	0.2	0.8	3.0	13.1	59.2	249.0	992.0
1999.....	40.3	*	*	*	*	0.2	0.7	3.0	13.5	58.9	253.3	1,017.1
White, both sexes												
2004.....	27.6	*	*	*	*	0.1	0.7	2.8	10.3	46.6	199.1	721.0
2003.....	30.2	*	*	*	*	0.2	0.7	2.8	11.5	49.6	218.1	807.1
2002.....	32.1	*	*	*	*	0.2	0.7	3.0	12.1	54.7	232.6	867.3
2001.....	33.1	*	*	*	*	0.1	0.7	2.8	12.5	56.7	244.3	913.6
2000.....	34.2	*	*	*	*	0.2	0.7	2.9	13.3	60.7	251.8	963.5
1999.....	34.8	*	*	*	*	0.2	0.6	2.8	13.2	60.5	258.6	993.4
White, male												
2004.....	20.5	*	*	*	*	*	0.8	3.4	12.1	53.0	203.3	622.4
2003.....	22.4	*	*	*	*	0.2	0.9	3.4	13.8	56.6	225.7	704.0
2002.....	23.5	*	*	*	*	0.2	0.8	3.5	14.5	61.7	235.1	766.7
2001.....	24.4	*	*	*	*	*	0.7	3.5	14.9	64.7	250.2	823.5
2000.....	25.2	*	*	*	*	0.2	0.7	3.7	16.1	68.5	263.5	865.2
1999.....	25.8	*	*	*	*	*	0.7	3.5	15.8	69.0	274.3	903.1

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Stroke, not specified as hemorrhage or infarction (I64)												
White, female												
2004.....	34.7	*	*	*	*	0.2	0.6	2.1	8.6	41.2	196.2	765.4
2003.....	37.8	*	*	*	*	0.2	0.6	2.2	9.2	43.6	213.0	852.8
2002.....	40.4	*	*	*	*	0.2	0.6	2.4	9.8	48.8	230.9	910.9
2001.....	41.6	*	*	*	*	0.2	0.6	2.2	10.4	49.9	240.4	951.3
2000.....	43.0	*	*	*	*	0.1	0.6	2.2	10.7	54.3	244.0	1,003.5
1999.....	43.6	*	*	*	*	0.2	0.5	2.2	10.8	53.5	248.3	1,029.7
Black, both sexes												
2004.....	23.1	*	*	*	*	0.6	2.6	10.6	34.3	104.2	289.9	706.1
2003.....	24.6	*	*	*	*	0.8	3.1	10.6	38.4	108.9	305.0	797.6
2002.....	26.0	*	*	*	*	0.4	2.5	11.4	41.7	120.5	325.3	840.7
2001.....	26.6	*	*	*	*	0.4	2.5	11.2	42.6	126.3	333.8	883.4
2000.....	27.2	*	*	*	*	0.5	2.3	11.3	41.8	129.3	351.8	919.4
1999.....	27.5	*	*	*	*	0.4	2.2	11.8	44.8	131.9	351.1	917.6
Black, male												
2004.....	19.1	*	*	*	*	*	2.8	12.4	43.5	124.7	303.5	589.1
2003.....	20.6	*	*	*	*	1.0	3.7	12.6	47.5	127.8	333.1	681.9
2002.....	22.0	*	*	*	*	*	2.3	13.3	53.4	151.0	344.5	761.2
2001.....	22.7	*	*	*	*	*	2.6	13.4	55.3	155.8	364.3	816.9
2000.....	23.2	*	*	*	*	*	2.2	13.8	51.3	154.9	392.7	897.2
1999.....	23.2	*	*	*	*	*	2.3	14.3	55.3	155.7	385.2	877.6
Black, female												
2004.....	26.6	*	*	*	*	*	2.5	9.0	26.9	89.5	282.3	752.5
2003.....	28.3	*	*	*	*	*	2.5	8.9	31.1	95.4	289.1	842.6
2002.....	29.6	*	*	*	*	*	2.7	9.8	32.3	98.8	314.4	871.2
2001.....	30.1	*	*	*	*	*	2.4	9.3	32.3	105.4	316.6	908.1
2000.....	30.9	*	*	*	*	*	2.3	9.1	34.2	111.2	328.8	927.6
1999.....	31.3	*	*	*	*	*	2.1	9.6	36.4	115.1	332.0	932.6

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other cerebrovascular diseases and their sequelae (I67,I69)

All races 3/, both sexes

2004.....	9.9	1.4	0.2	0.1	0.1	0.1	0.4	1.1	3.8	16.0	74.3	298.3
2003.....	10.3	1.2	0.1	0.1	0.1	0.1	0.3	1.0	3.8	17.2	76.6	320.1
2002.....	10.5	1.4	0.2	0.1	0.0	0.1	0.4	1.1	3.8	17.4	80.0	328.9
2001.....	10.3	1.2	0.2	0.1	*	0.1	0.3	1.0	3.9	17.5	80.2	328.2
2000.....	10.5	1.6	0.2	*	0.1	0.1	0.3	1.1	4.1	17.6	82.1	341.4
1999.....	10.6	1.1	0.1	0.1	0.1	0.1	0.3	1.1	4.1	18.5	83.3	343.9

All races 3/, male

2004.....	7.6	1.5	*	*	*	0.1	0.4	1.1	4.6	19.0	80.4	270.4
2003.....	7.8	1.2	*	*	*	0.1	0.3	1.1	4.6	20.0	82.9	290.2
2002.....	7.9	1.7	0.3	0.1	*	0.1	0.3	1.2	4.5	20.0	86.4	302.3
2001.....	7.9	1.5	*	*	*	0.1	0.3	1.1	4.6	20.8	88.2	305.5
2000.....	8.1	1.8	*	*	*	*	0.3	1.2	4.7	21.2	92.0	326.9
1999.....	8.1	1.4	*	*	*	*	0.3	1.1	5.2	21.6	92.4	332.1

All races 3/, female

2004.....	12.1	1.4	*	*	*	0.1	0.3	1.0	3.0	13.4	70.2	310.8
2003.....	12.7	1.2	*	*	*	*	0.4	0.9	3.1	14.8	72.4	333.3
2002.....	13.0	1.1	*	*	*	*	0.4	1.0	3.2	15.3	75.7	340.4
2001.....	12.7	*	*	*	*	0.1	0.3	1.0	3.2	14.7	74.9	337.6
2000.....	12.8	1.4	*	*	*	0.1	0.3	1.1	3.6	14.6	75.6	347.4
1999.....	12.9	*	*	*	*	0.1	0.4	1.0	3.2	15.9	77.4	348.7

White, both sexes

2004.....	10.5	1.1	*	0.1	0.1	0.1	0.3	0.8	3.1	14.4	72.2	300.3
2003.....	11.0	1.0	*	*	0.1	0.1	0.3	0.8	3.1	15.5	74.8	322.5
2002.....	11.3	0.9	0.2	*	*	0.1	0.3	0.9	3.2	15.9	78.1	332.0
2001.....	11.1	0.9	0.2	*	*	0.1	0.3	0.8	3.3	16.0	78.3	329.9
2000.....	11.3	1.2	*	*	*	0.1	0.3	0.9	3.5	16.3	80.0	343.4
1999.....	11.4	0.9	*	*	0.1	0.1	0.3	0.9	3.4	16.8	81.5	348.1

White, male

2004.....	7.9	*	*	*	*	*	0.4	0.8	3.7	17.1	77.6	271.5
2003.....	8.2	*	*	*	*	0.1	0.2	0.8	3.7	18.0	80.2	294.0
2002.....	8.4	*	*	*	*	0.1	0.3	1.0	3.8	18.3	84.0	306.3
2001.....	8.3	*	*	*	*	*	0.3	0.8	3.9	19.1	86.4	308.1
2000.....	8.6	1.6	*	*	*	*	0.3	1.0	4.1	19.5	89.4	330.0
1999.....	8.5	1.3	*	*	*	*	0.2	0.9	4.2	19.3	90.0	334.9

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other cerebrovascular diseases and their sequelae (I67,I69)

White, female

2004.....	13.1	*	*	*	*	*	0.3	0.8	2.5	12.2	68.6	313.3
2003.....	13.8	*	*	*	*	*	0.3	0.8	2.6	13.3	71.2	335.2
2002.....	14.1	*	*	*	*	*	0.4	0.8	2.7	13.8	74.2	343.2
2001.....	13.7	*	*	*	*	0.1	0.3	0.7	2.6	13.3	72.9	339.1
2000.....	13.9	*	*	*	*	*	0.3	0.8	3.0	13.7	73.8	348.8
1999.....	14.1	*	*	*	*	*	0.3	0.9	2.7	14.7	75.8	353.4

Black, both sexes

2004.....	8.1	3.2	*	*	*	*	0.5	2.5	9.9	32.5	103.2	296.9
2003.....	8.3	*	*	*	*	*	0.6	2.8	10.1	33.9	103.1	321.5
2002.....	8.4	3.4	*	*	*	*	0.7	2.8	9.7	34.1	107.0	318.9
2001.....	8.3	3.1	*	*	*	*	0.7	3.0	10.0	32.3	107.3	327.6
2000.....	8.2	3.9	*	*	*	*	0.7	2.9	9.6	30.0	108.0	333.8
1999.....	8.3	*	*	*	*	*	0.8	2.6	10.2	35.9	108.8	312.5

Black, male

2004.....	7.0	*	*	*	*	*	*	3.1	13.1	42.1	121.9	272.2
2003.....	6.9	*	*	*	*	*	*	3.4	12.4	42.5	121.0	271.1
2002.....	6.8	*	*	*	*	*	*	3.1	12.1	40.4	122.5	269.1
2001.....	6.6	*	*	*	*	*	0.8	3.1	11.9	40.1	118.3	285.9
2000.....	6.6	*	*	*	*	*	0.8	2.8	10.8	39.8	126.6	291.8
1999.....	7.1	*	*	*	*	*	0.8	3.3	14.0	47.2	121.8	294.1

Black, female

2004.....	9.1	*	*	*	*	*	*	2.0	7.3	25.7	92.7	306.8
2003.....	9.6	*	*	*	*	*	0.7	2.2	8.2	27.8	93.0	341.1
2002.....	9.8	*	*	*	*	*	0.8	2.5	7.9	29.6	98.3	338.0
2001.....	9.9	*	*	*	*	*	*	2.8	8.5	26.9	101.1	343.1
2000.....	9.6	*	*	*	*	*	0.7	3.0	8.6	23.2	97.5	349.4
1999.....	9.3	*	*	*	*	*	0.7	2.0	7.1	28.0	101.6	319.3

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diseases of arteries, arterioles and capillaries (I70-I78)												
All races 3/, both sexes												
2004.....	12.1	*	*	*	0.2	0.4	1.1	3.1	8.8	31.2	90.6	276.3
2003.....	13.0	*	*	*	0.2	0.4	1.1	3.4	9.8	33.5	97.4	300.6
2002.....	13.4	*	*	*	0.2	0.4	1.1	3.1	10.4	36.2	100.6	316.5
2001.....	13.8	*	*	*	0.2	0.4	1.2	3.1	10.5	36.5	104.9	337.3
2000.....	14.4	*	*	*	0.2	0.4	1.1	3.2	11.1	40.0	109.2	352.0
1999.....	14.6	*	*	*	0.1	0.4	1.2	3.2	11.3	39.5	111.6	367.2
All races 3/, male												
2004.....	11.6	*	*	*	0.2	0.6	1.5	4.3	11.9	40.8	112.5	294.7
2003.....	12.6	*	*	*	0.2	0.5	1.5	4.7	13.5	44.8	125.5	318.3
2002.....	13.0	*	*	*	0.2	0.5	1.4	4.3	14.3	48.6	127.8	344.0
2001.....	13.3	*	*	*	0.2	0.6	1.6	4.2	14.3	49.3	133.0	364.5
2000.....	14.1	*	*	*	0.2	0.5	1.4	4.4	15.7	55.1	142.5	391.4
1999.....	14.1	*	*	*	0.2	0.5	1.5	4.3	16.0	54.0	144.2	396.5
All races 3/, female												
2004.....	12.6	*	*	*	*	0.3	0.8	1.9	6.0	23.2	75.8	268.0
2003.....	13.3	*	*	*	0.1	0.3	0.7	2.1	6.4	24.1	78.6	292.8
2002.....	13.8	*	*	*	0.1	0.3	0.8	2.0	6.8	25.8	82.5	304.6
2001.....	14.4	*	*	*	0.1	0.3	0.8	2.0	7.1	25.8	86.4	325.9
2000.....	14.6	*	*	*	0.1	0.3	0.8	2.1	6.8	27.5	87.5	336.0
1999.....	15.2	*	*	*	*	0.2	0.9	2.0	7.0	27.7	90.5	355.4
White, both sexes												
2004.....	13.2	*	*	*	0.2	0.4	1.0	2.8	8.3	31.5	91.8	281.3
2003.....	14.2	*	*	*	0.2	0.4	1.0	3.0	9.5	33.8	98.6	306.4
2002.....	14.7	*	*	*	0.2	0.4	1.0	2.7	10.0	36.5	101.8	323.7
2001.....	15.1	*	*	*	0.1	0.4	1.0	2.8	10.0	36.8	105.9	344.1
2000.....	15.7	*	*	*	0.2	0.4	0.9	2.8	10.6	40.5	110.2	358.3
1999.....	16.0	*	*	*	0.1	0.3	1.1	2.9	11.1	40.1	112.9	372.8
White, male												
2004.....	12.7	*	*	*	0.2	0.5	1.3	4.0	11.2	41.5	114.6	303.7
2003.....	13.8	*	*	*	0.2	0.5	1.4	4.2	13.2	45.4	126.8	327.4
2002.....	14.2	*	*	*	0.2	0.4	1.3	3.7	13.7	49.4	130.2	354.1
2001.....	14.5	*	*	*	0.2	0.6	1.3	3.7	13.8	50.0	134.7	375.1
2000.....	15.4	*	*	*	0.2	0.5	1.2	3.9	15.3	56.4	145.0	401.0
1999.....	15.4	*	*	*	0.2	0.5	1.3	4.0	15.9	55.3	146.4	404.8

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diseases of arteries, arterioles and capillaries (I70-I78)												
White, female												
2004.....	13.8	*	*	*	*	0.2	0.6	1.6	5.6	22.9	76.2	271.1
2003.....	14.6	*	*	*	*	0.2	0.6	1.7	6.0	23.9	79.5	297.1
2002.....	15.2	*	*	*	*	0.3	0.7	1.6	6.5	25.6	82.7	310.4
2001.....	15.8	*	*	*	*	0.3	0.7	1.8	6.5	25.6	86.7	331.0
2000.....	16.0	*	*	*	*	0.2	0.7	1.7	6.2	27.1	87.3	340.9
1999.....	16.7	*	*	*	*	0.2	0.8	1.8	6.6	27.4	90.9	360.0
Black, both sexes												
2004.....	8.8	*	*	*	*	0.8	2.4	5.8	15.0	35.4	94.3	258.0
2003.....	9.2	*	*	*	*	0.7	2.4	7.0	14.4	38.5	100.8	271.0
2002.....	9.4	*	*	*	*	0.9	2.1	6.9	16.4	39.1	104.8	273.8
2001.....	9.7	*	*	*	*	0.7	2.6	5.8	17.2	40.6	110.5	300.3
2000.....	10.0	*	*	*	*	0.7	2.4	6.6	17.6	41.9	113.2	315.9
1999.....	9.7	*	*	*	*	0.6	2.1	5.9	15.6	40.9	111.4	333.2
Black, male												
2004.....	8.3	*	*	*	*	1.2	3.2	8.0	21.1	43.8	110.0	236.4
2003.....	8.9	*	*	*	*	1.1	3.2	9.4	20.0	48.7	130.6	242.9
2002.....	9.0	*	*	*	*	1.2	2.8	9.5	23.5	48.5	122.8	276.6
2001.....	9.0	*	*	*	*	0.9	3.6	8.3	21.6	50.8	133.9	279.1
2000.....	9.3	*	*	*	*	0.9	3.1	8.9	22.4	52.1	136.0	325.2
1999.....	9.0	*	*	*	*	0.9	2.5	8.0	20.2	51.2	133.7	348.0
Black, female												
2004.....	9.2	*	*	*	*	*	1.8	3.9	10.1	29.4	85.4	266.6
2003.....	9.5	*	*	*	*	*	1.7	5.0	10.0	31.2	83.9	281.9
2002.....	9.7	*	*	*	*	*	1.5	4.7	10.8	32.3	94.7	272.7
2001.....	10.3	*	*	*	*	*	1.6	3.7	13.7	33.4	97.3	308.2
2000.....	10.6	*	*	*	*	*	1.7	4.7	13.7	34.8	100.4	312.5
1999.....	10.3	*	*	*	*	*	1.8	4.2	11.9	33.6	99.0	327.7

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Atherosclerosis (I70)												
All races 3/, both sexes												
2004.....	4.0	*	*	*	*	*	0.1	0.4	1.6	6.1	26.3	137.0
2003.....	4.5	*	*	*	*	*	0.1	0.5	1.7	6.6	29.8	154.6
2002.....	4.8	*	*	*	*	*	0.1	0.5	1.8	7.1	31.9	168.1
2001.....	4.9	*	*	*	*	*	0.1	0.5	1.9	7.3	33.1	179.1
2000.....	5.1	*	*	*	*	*	0.1	0.5	1.9	7.7	34.6	188.8
1999.....	5.4	*	*	*	*	*	0.1	0.5	2.0	8.0	35.9	202.0
All races 3/, male												
2004.....	3.1	*	*	*	*	*	0.1	0.5	2.0	7.3	29.7	128.2
2003.....	3.4	*	*	*	*	*	0.2	0.7	2.2	7.8	33.6	138.6
2002.....	3.7	*	*	*	*	*	0.1	0.6	2.4	8.8	35.5	156.6
2001.....	3.7	*	*	*	*	*	0.1	0.6	2.4	9.0	35.9	167.2
2000.....	3.9	*	*	*	*	*	0.2	0.6	2.5	9.6	39.7	176.5
1999.....	4.0	*	*	*	*	*	0.1	0.7	2.7	9.6	40.0	189.0
All races 3/, female												
2004.....	4.9	*	*	*	*	*	*	0.2	1.2	5.1	24.0	141.0
2003.....	5.5	*	*	*	*	*	*	0.3	1.2	5.6	27.3	161.7
2002.....	5.9	*	*	*	*	*	0.1	0.3	1.3	5.7	29.5	173.1
2001.....	6.2	*	*	*	*	*	*	0.3	1.4	5.9	31.2	184.1
2000.....	6.3	*	*	*	*	*	*	0.3	1.2	6.2	31.3	193.8
1999.....	6.7	*	*	*	*	*	*	0.3	1.4	6.7	33.2	207.2
White, both sexes												
2004.....	4.6	*	*	*	*	*	0.1	0.4	1.6	6.1	27.1	142.0
2003.....	5.0	*	*	*	*	*	0.1	0.4	1.6	6.5	30.3	159.5
2002.....	5.4	*	*	*	*	*	0.1	0.4	1.7	7.1	32.1	174.7
2001.....	5.6	*	*	*	*	*	0.1	0.5	1.8	7.3	33.8	185.9
2000.....	5.7	*	*	*	*	*	0.1	0.4	1.8	7.7	35.1	195.2
1999.....	6.0	*	*	*	*	*	0.1	0.5	2.0	8.0	36.4	207.7
White, male												
2004.....	3.5	*	*	*	*	*	*	0.5	1.9	7.4	30.4	134.4
2003.....	3.7	*	*	*	*	*	0.2	0.6	2.1	7.5	33.7	143.3
2002.....	4.0	*	*	*	*	*	0.1	0.5	2.3	8.8	35.6	163.2
2001.....	4.1	*	*	*	*	*	0.1	0.6	2.3	9.0	36.4	173.9
2000.....	4.3	*	*	*	*	*	0.2	0.5	2.5	9.7	40.3	181.4
1999.....	4.4	*	*	*	*	*	0.1	0.7	2.6	9.5	40.2	193.7

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Atherosclerosis (I70)												
White, female												
2004.....	5.6	*	*	*	*	*	*	0.2	1.2	5.0	24.9	145.4
2003.....	6.3	*	*	*	*	*	*	0.3	1.1	5.6	28.0	166.7
2002.....	6.7	*	*	*	*	*	*	0.3	1.3	5.7	29.8	179.7
2001.....	7.0	*	*	*	*	*	*	0.3	1.3	5.9	32.1	190.9
2000.....	7.1	*	*	*	*	*	*	0.2	1.1	6.1	31.7	200.8
1999.....	7.6	*	*	*	*	*	*	0.3	1.4	6.8	34.0	213.3
Black, both sexes												
2004.....	2.2	*	*	*	*	*	*	0.5	2.3	7.4	23.9	102.9
2003.....	2.6	*	*	*	*	*	*	0.8	2.6	9.2	30.7	121.1
2002.....	2.7	*	*	*	*	*	*	1.0	2.9	8.0	36.5	117.0
2001.....	2.6	*	*	*	*	*	*	0.7	3.3	8.9	30.7	125.6
2000.....	2.9	*	*	*	*	*	*	1.1	3.2	8.9	35.2	138.3
1999.....	3.0	*	*	*	*	*	*	0.7	3.2	9.2	34.9	154.3
Black, male												
2004.....	1.7	*	*	*	*	*	*	*	3.0	8.6	30.1	78.5
2003.....	2.3	*	*	*	*	*	*	1.2	3.6	12.4	40.3	104.3
2002.....	2.3	*	*	*	*	*	*	1.4	4.2	10.1	44.2	110.4
2001.....	2.1	*	*	*	*	*	*	1.0	3.8	10.9	37.1	113.5
2000.....	2.4	*	*	*	*	*	*	1.4	3.9	11.0	40.3	147.6
1999.....	2.5	*	*	*	*	*	*	*	4.4	11.8	44.0	164.0
Black, female												
2004.....	2.6	*	*	*	*	*	*	*	1.7	6.5	20.5	112.6
2003.....	2.9	*	*	*	*	*	*	*	1.8	7.0	25.2	127.6
2002.....	3.1	*	*	*	*	*	*	*	1.9	6.5	32.2	119.5
2001.....	3.1	*	*	*	*	*	*	*	2.9	7.5	27.1	130.1
2000.....	3.3	*	*	*	*	*	*	*	2.7	7.5	32.4	134.8
1999.....	3.4	*	*	*	*	*	*	*	2.3	7.4	29.8	150.7

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Aortic aneurysm and dissection (I71)												
All races 3/, both sexes												
2004.....	4.7	*	*	*	0.1	0.3	0.7	1.8	4.5	15.9	39.6	64.8
2003.....	5.1	*	*	*	0.1	0.3	0.7	2.0	5.3	17.8	42.5	70.0
2002.....	5.1	*	*	*	0.1	0.3	0.7	1.8	5.5	19.1	43.3	69.4
2001.....	5.3	*	*	*	0.1	0.3	0.7	1.7	5.5	19.2	45.7	77.5
2000.....	5.6	*	*	*	0.1	0.3	0.7	1.8	6.2	21.8	47.2	79.2
1999.....	5.7	*	*	*	0.1	0.2	0.7	1.7	6.0	21.7	48.6	80.3
All races 3/, male												
2004.....	5.6	*	*	*	0.2	0.5	1.1	2.7	6.9	23.1	54.8	95.0
2003.....	6.3	*	*	*	0.1	0.4	1.1	3.1	8.2	26.0	61.9	105.5
2002.....	6.4	*	*	*	0.2	0.4	1.0	2.7	8.5	28.5	62.7	106.6
2001.....	6.5	*	*	*	0.2	0.4	1.2	2.6	8.5	28.7	66.4	114.8
2000.....	7.0	*	*	*	0.1	0.4	1.0	2.7	9.5	32.9	70.6	127.3
1999.....	7.0	*	*	*	0.1	0.4	1.0	2.7	9.5	32.7	72.7	125.3
All races 3/, female												
2004.....	3.8	*	*	*	*	0.2	0.3	0.8	2.3	9.8	29.3	51.2
2003.....	3.9	*	*	*	*	0.1	0.3	1.0	2.6	11.0	29.5	54.2
2002.....	4.0	*	*	*	*	0.1	0.3	0.8	2.7	11.2	30.5	53.3
2001.....	4.2	*	*	*	*	0.1	0.3	0.8	2.8	11.3	32.1	61.9
2000.....	4.3	*	*	*	*	0.1	0.4	0.9	3.0	12.7	32.0	59.6
1999.....	4.4	*	*	*	*	*	0.4	0.8	2.8	12.7	32.9	62.2
White, both sexes												
2004.....	5.1	*	*	*	0.1	0.3	0.6	1.5	4.3	16.5	40.6	66.8
2003.....	5.6	*	*	*	0.1	0.2	0.6	1.7	5.2	18.6	44.0	72.4
2002.....	5.7	*	*	*	0.1	0.2	0.6	1.5	5.4	19.9	45.0	71.2
2001.....	5.9	*	*	*	0.1	0.3	0.6	1.5	5.4	19.9	46.9	79.7
2000.....	6.2	*	*	*	0.1	0.2	0.6	1.6	6.0	22.7	48.5	80.8
1999.....	6.3	*	*	*	0.1	0.2	0.6	1.5	6.1	22.6	50.1	82.2
White, male												
2004.....	6.1	*	*	*	0.2	0.4	0.9	2.4	6.5	24.1	56.6	98.7
2003.....	6.9	*	*	*	0.1	0.3	1.0	2.7	8.2	27.3	64.3	110.0
2002.....	7.0	*	*	*	0.1	0.3	0.9	2.3	8.4	29.8	65.5	111.8
2001.....	7.2	*	*	*	0.1	0.5	1.0	2.3	8.4	29.9	68.5	120.3
2000.....	7.8	*	*	*	*	0.3	0.8	2.5	9.5	34.5	73.0	132.8
1999.....	7.8	*	*	*	*	0.4	0.9	2.4	9.8	34.3	75.8	129.8

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Aortic aneurysm and dissection (I71)												
White, female												
2004.....	4.1	*	*	*	*	0.2	0.3	0.7	2.1	10.1	29.8	52.4
2003.....	4.3	*	*	*	*	0.2	0.2	0.8	2.4	11.1	30.3	55.7
2002.....	4.3	*	*	*	*	*	0.3	0.7	2.6	11.4	31.2	53.5
2001.....	4.6	*	*	*	*	*	0.2	0.6	2.6	11.5	32.5	62.6
2000.....	4.6	*	*	*	*	0.1	0.3	0.7	2.8	12.8	32.3	59.6
1999.....	4.8	*	*	*	*	*	0.4	0.7	2.7	12.8	33.3	63.1
Black, both sexes												
2004.....	3.2	*	*	*	*	0.6	1.6	3.6	7.6	13.0	32.0	43.7
2003.....	3.2	*	*	*	*	0.5	1.6	4.2	7.1	14.5	30.0	46.4
2002.....	3.2	*	*	*	*	0.6	1.4	4.2	7.5	14.6	29.9	50.5
2001.....	3.4	*	*	*	*	0.4	1.6	3.5	7.6	14.6	37.6	55.3
2000.....	3.5	*	*	*	*	0.5	1.6	3.2	8.2	17.0	36.3	62.3
1999.....	3.3	*	*	*	*	0.4	1.4	3.7	6.5	16.7	33.8	60.3
Black, male												
2004.....	3.8	*	*	*	*	1.0	2.5	5.5	11.8	18.2	36.7	56.6
2003.....	3.8	*	*	*	*	1.0	2.4	6.4	10.8	18.0	39.2	51.1
2002.....	3.8	*	*	*	*	1.0	2.2	6.6	12.1	19.0	35.0	49.3
2001.....	3.8	*	*	*	*	*	2.6	5.3	11.2	20.0	47.4	48.8
2000.....	3.9	*	*	*	*	*	2.4	4.7	11.5	22.1	46.9	69.2
1999.....	3.6	*	*	*	*	*	1.9	5.8	9.1	21.9	38.4	76.2
Black, female												
2004.....	2.6	*	*	*	*	*	0.8	2.0	4.2	9.4	29.3	38.6
2003.....	2.7	*	*	*	*	*	1.0	2.4	4.2	11.9	24.7	44.6
2002.....	2.7	*	*	*	*	*	*	2.2	3.8	11.5	27.0	50.9
2001.....	3.0	*	*	*	*	*	0.7	1.9	4.6	10.8	32.1	57.7
2000.....	3.2	*	*	*	*	*	0.9	2.0	5.5	13.4	30.3	59.7
1999.....	3.0	*	*	*	*	*	0.9	2.0	4.4	13.0	31.2	54.3

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other diseases of arteries, arterioles and capillaries (I72-I78)												
All races 3/, both sexes												
2004.....	3.4	*	*	*	*	0.1	0.3	1.0	2.7	9.2	24.7	74.5
2003.....	3.4	*	*	*	0.1	0.1	0.3	0.9	2.8	9.1	25.0	76.0
2002.....	3.5	*	*	*	0.1	0.1	0.3	0.9	3.1	10.0	25.3	79.0
2001.....	3.5	*	*	*	*	0.1	0.4	0.9	3.2	10.0	26.1	80.7
2000.....	3.6	*	*	*	0.1	0.1	0.3	1.0	3.1	10.4	27.4	84.0
1999.....	3.6	*	*	*	*	0.1	0.3	0.9	3.3	9.8	27.2	84.9
All races 3/, male												
2004.....	2.9	*	*	*	*	0.1	0.3	1.1	3.0	10.3	28.0	71.6
2003.....	3.0	*	*	*	*	0.1	0.3	1.0	3.1	11.0	29.9	74.2
2002.....	3.0	*	*	*	*	*	0.3	1.0	3.4	11.3	29.6	80.8
2001.....	3.1	*	*	*	*	0.1	0.3	0.9	3.5	11.6	30.6	82.5
2000.....	3.2	*	*	*	*	0.1	0.3	1.1	3.6	12.5	32.3	87.6
1999.....	3.1	*	*	*	*	0.1	0.3	1.0	3.8	11.7	31.5	82.2
All races 3/, female												
2004.....	3.9	*	*	*	*	0.1	0.4	0.8	2.4	8.3	22.5	75.8
2003.....	3.8	*	*	*	*	*	0.3	0.9	2.6	7.5	21.8	76.9
2002.....	4.0	*	*	*	*	0.2	0.3	0.8	2.8	8.9	22.5	78.2
2001.....	4.0	*	*	*	*	0.1	0.4	0.9	2.9	8.6	23.1	79.9
2000.....	4.0	*	*	*	*	0.1	0.4	0.9	2.6	8.7	24.2	82.6
1999.....	4.1	*	*	*	*	0.1	0.4	0.9	2.8	8.3	24.5	86.0
White, both sexes												
2004.....	3.5	*	*	*	*	0.1	0.3	0.9	2.5	8.8	24.0	72.5
2003.....	3.6	*	*	*	*	0.1	0.3	0.8	2.7	8.7	24.2	74.6
2002.....	3.7	*	*	*	*	0.1	0.3	0.8	2.8	9.6	24.7	77.8
2001.....	3.7	*	*	*	*	0.1	0.3	0.9	2.9	9.5	25.2	78.5
2000.....	3.8	*	*	*	*	0.1	0.3	0.8	2.8	10.0	26.6	82.3
1999.....	3.8	*	*	*	*	0.1	0.3	0.9	3.0	9.5	26.3	82.9
White, male												
2004.....	3.0	*	*	*	*	*	0.3	1.0	2.7	10.1	27.7	70.7
2003.....	3.1	*	*	*	*	*	0.2	0.9	2.9	10.6	28.8	74.1
2002.....	3.2	*	*	*	*	*	0.3	0.9	3.1	10.8	29.1	79.1
2001.....	3.2	*	*	*	*	*	0.2	0.8	3.2	11.2	29.8	80.9
2000.....	3.4	*	*	*	*	0.1	0.2	0.9	3.3	12.2	31.6	86.9
1999.....	3.2	*	*	*	*	*	0.3	0.9	3.6	11.4	30.4	81.3

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other diseases of arteries, arterioles and capillaries (I72-I78)

White, female

2004.....	4.0	*	*	*	*	*	0.3	0.7	2.3	7.8	21.6	73.3
2003.....	4.0	*	*	*	*	*	0.3	0.7	2.5	7.2	21.2	74.8
2002.....	4.2	*	*	*	*	0.1	0.3	0.7	2.6	8.5	21.7	77.2
2001.....	4.2	*	*	*	*	0.1	0.4	0.9	2.6	8.2	22.1	77.5
2000.....	4.2	*	*	*	*	*	0.3	0.7	2.3	8.2	23.3	80.4
1999.....	4.3	*	*	*	*	*	0.3	0.8	2.5	7.9	23.6	83.5

Black, both sexes

2004.....	3.4	*	*	*	*	*	0.6	1.7	5.2	14.9	38.4	111.4
2003.....	3.4	*	*	*	*	*	0.7	2.0	4.7	14.8	40.1	103.6
2002.....	3.5	*	*	*	*	*	0.6	1.7	6.1	16.5	38.5	106.3
2001.....	3.7	*	*	*	*	*	0.9	1.7	6.4	17.1	42.2	119.4
2000.....	3.6	*	*	*	*	*	0.6	2.3	6.2	16.0	41.7	115.4
1999.....	3.5	*	*	*	*	*	0.6	1.5	5.8	15.0	42.8	118.6

Black, male

2004.....	2.9	*	*	*	*	*	*	2.0	6.3	17.0	43.2	101.3
2003.....	2.9	*	*	*	*	*	*	1.8	5.6	18.3	51.1	87.6
2002.....	3.0	*	*	*	*	*	*	1.5	7.2	19.4	43.6	116.9
2001.....	3.2	*	*	*	*	*	0.9	2.0	6.5	20.0	49.4	116.9
2000.....	3.1	*	*	*	*	*	*	2.8	7.0	19.0	48.7	108.4
1999.....	2.9	*	*	*	*	*	*	1.4	6.6	17.5	51.3	107.8

Black, female

2004.....	3.9	*	*	*	*	*	0.8	1.4	4.2	13.5	35.7	115.4
2003.....	3.8	*	*	*	*	*	0.7	2.1	4.0	12.3	34.0	109.8
2002.....	3.9	*	*	*	*	*	0.8	1.9	5.1	14.4	35.6	102.3
2001.....	4.2	*	*	*	*	*	0.8	1.4	6.2	15.1	38.2	120.4
2000.....	4.1	*	*	*	*	*	0.7	1.8	5.5	13.9	37.7	118.0
1999.....	4.0	*	*	*	*	*	0.8	1.6	5.2	13.3	38.0	122.7

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other disorders of circulatory system (I80-I99)												
All races 3/, both sexes												
2004.....	1.6	0.7	*	*	0.1	0.3	0.7	1.3	2.1	3.6	9.0	23.3
2003.....	1.6	0.9	*	*	0.1	0.3	0.8	1.4	2.0	4.1	9.0	24.0
2002.....	1.6	0.7	*	*	0.1	0.3	0.7	1.3	2.2	3.8	9.8	24.4
2001.....	1.6	0.5	*	*	0.1	0.3	0.6	1.2	2.2	4.6	9.4	26.1
2000.....	1.6	0.7	*	*	0.1	0.3	0.7	1.4	2.1	4.3	9.4	25.6
1999.....	1.4	0.6	*	*	0.1	0.2	0.6	1.0	1.9	3.8	9.1	23.4
All races 3/, male												
2004.....	1.4	*	*	*	*	0.4	0.7	1.6	2.5	3.9	9.7	19.6
2003.....	1.4	1.0	*	*	0.1	0.3	0.8	1.6	2.4	4.2	8.5	19.1
2002.....	1.4	*	*	*	0.1	0.3	0.7	1.4	2.4	3.9	9.5	20.5
2001.....	1.4	*	*	*	*	0.3	0.7	1.4	2.5	4.9	9.8	22.0
2000.....	1.4	*	*	*	*	0.3	0.8	1.6	2.3	4.7	9.7	22.2
1999.....	1.3	*	*	*	*	0.2	0.7	1.1	2.0	4.2	9.6	20.8
All races 3/, female												
2004.....	1.7	*	*	*	*	0.3	0.7	1.1	1.8	3.4	8.5	25.0
2003.....	1.9	*	*	*	0.1	0.4	0.8	1.2	1.7	4.0	9.4	26.2
2002.....	1.9	*	*	*	*	0.4	0.7	1.2	2.1	3.8	10.1	26.2
2001.....	1.9	*	*	*	0.1	0.4	0.6	1.0	2.0	4.4	9.2	27.8
2000.....	1.8	*	*	*	0.1	0.4	0.6	1.1	2.0	4.0	9.2	26.9
1999.....	1.6	*	*	*	*	0.2	0.5	1.0	1.8	3.4	8.8	24.5
White, both sexes												
2004.....	1.6	0.7	*	*	0.1	0.3	0.6	1.2	2.0	3.4	8.8	23.2
2003.....	1.6	0.9	*	*	0.1	0.3	0.6	1.2	1.9	3.8	8.6	23.6
2002.....	1.6	0.6	*	*	0.1	0.3	0.6	1.2	2.0	3.6	9.5	24.3
2001.....	1.7	*	*	*	0.1	0.3	0.5	1.0	2.0	4.3	9.1	26.0
2000.....	1.6	0.7	*	*	0.1	0.3	0.6	1.2	1.9	4.0	9.1	25.5
1999.....	1.5	*	*	*	*	0.2	0.5	0.9	1.8	3.5	8.9	23.2
White, male												
2004.....	1.4	*	*	*	*	0.3	0.6	1.4	2.4	3.7	9.4	19.7
2003.....	1.3	*	*	*	*	0.2	0.7	1.4	2.2	3.9	7.9	18.5
2002.....	1.3	*	*	*	*	0.3	0.6	1.3	2.2	3.6	9.0	19.9
2001.....	1.4	*	*	*	*	0.2	0.5	1.3	2.2	4.6	9.5	21.7
2000.....	1.4	*	*	*	*	0.2	0.8	1.5	2.0	4.4	9.5	22.0
1999.....	1.2	*	*	*	*	0.2	0.6	0.9	1.9	3.9	9.2	20.7

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other disorders of circulatory system (I80-I99)												
White, female												
2004.....	1.8	*	*	*	*	0.3	0.5	0.9	1.6	3.2	8.4	24.8
2003.....	1.9	*	*	*	*	0.3	0.5	1.1	1.5	3.8	9.1	25.8
2002.....	1.9	*	*	*	*	0.3	0.6	1.1	1.9	3.6	9.8	26.3
2001.....	1.9	*	*	*	*	0.3	0.5	0.8	1.8	4.1	8.9	27.9
2000.....	1.9	*	*	*	*	0.3	0.5	1.0	1.8	3.6	8.9	26.9
1999.....	1.7	*	*	*	*	0.2	0.4	0.9	1.7	3.2	8.7	24.2
Black, both sexes												
2004.....	2.1	*	*	*	*	0.9	1.9	2.9	4.2	6.6	13.6	28.8
2003.....	2.2	*	*	*	*	0.9	1.9	2.6	4.3	7.5	15.4	33.6
2002.....	2.1	*	*	*	*	1.0	1.6	2.6	4.9	6.8	16.6	30.0
2001.....	2.0	*	*	*	*	0.8	1.5	2.2	4.9	7.8	14.4	30.1
2000.....	2.1	*	*	*	*	0.9	1.5	2.8	4.5	8.9	14.4	30.0
1999.....	1.7	*	*	*	*	0.5	1.4	2.4	2.9	6.6	13.8	28.4
Black, male												
2004.....	1.9	*	*	*	*	1.1	2.0	3.4	4.6	7.1	15.0	24.8
2003.....	2.0	*	*	*	*	0.9	1.3	3.4	5.2	8.7	17.5	32.3
2002.....	2.0	*	*	*	*	1.0	1.8	3.0	5.6	7.7	18.4	32.2
2001.....	1.9	*	*	*	*	*	1.7	2.4	6.4	9.0	15.9	29.5
2000.....	1.9	*	*	*	*	0.9	1.8	3.4	5.1	9.1	13.3	30.0
1999.....	1.6	*	*	*	*	*	1.8	2.8	2.6	8.2	16.9	*
Black, female												
2004.....	2.2	*	*	*	*	0.8	1.9	2.5	3.9	6.3	12.7	30.3
2003.....	2.3	*	*	*	*	0.9	2.5	1.9	3.6	6.7	14.2	34.0
2002.....	2.2	*	*	*	*	0.9	1.5	2.3	4.4	6.1	15.6	29.2
2001.....	2.1	*	*	*	*	0.9	1.4	2.1	3.6	7.1	13.6	30.3
2000.....	2.2	*	*	*	*	0.9	1.3	2.3	3.9	8.8	15.0	30.0
1999.....	1.8	*	*	*	*	*	1.1	2.0	3.1	5.5	12.0	30.7

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)

All races 3/, both sexes

2004.....	1.2	*	*	*	0.1	0.3	0.6	1.0	1.5	2.6	6.9	17.3
2003.....	1.2	*	*	*	0.1	0.3	0.6	1.0	1.5	3.1	6.8	18.1
2002.....	1.2	*	*	*	0.1	0.3	0.6	1.0	1.7	2.8	7.3	18.6
2001.....	1.2	*	*	*	0.1	0.3	0.5	0.8	1.7	3.4	7.1	19.6
2000.....	1.2	*	*	*	0.1	0.3	0.6	1.0	1.5	3.2	7.0	18.7
1999.....	0.9	*	*	*	0.1	0.2	0.5	0.7	1.2	2.4	5.8	14.6

All races 3/, male

2004.....	1.0	*	*	*	*	0.3	0.6	1.0	1.7	2.7	7.0	14.3
2003.....	1.0	*	*	*	*	0.3	0.6	1.1	1.7	3.1	6.0	13.8
2002.....	1.0	*	*	*	0.1	0.3	0.5	1.0	1.7	2.8	6.6	14.8
2001.....	1.0	*	*	*	*	0.2	0.5	0.9	1.8	3.5	6.8	15.2
2000.....	1.0	*	*	*	*	0.3	0.6	1.0	1.4	3.3	6.9	15.0
1999.....	0.8	*	*	*	*	0.2	0.5	0.6	1.1	2.7	5.7	11.2

All races 3/, female

2004.....	1.4	*	*	*	*	0.3	0.6	0.9	1.4	2.5	6.7	18.6
2003.....	1.5	*	*	*	0.1	0.4	0.7	1.0	1.3	3.2	7.3	20.1
2002.....	1.5	*	*	*	*	0.3	0.6	1.0	1.7	2.9	7.8	20.2
2001.....	1.5	*	*	*	*	0.3	0.5	0.8	1.6	3.3	7.3	21.4
2000.....	1.4	*	*	*	0.1	0.3	0.5	1.0	1.6	3.1	7.1	20.2
1999.....	1.1	*	*	*	*	0.2	0.4	0.7	1.3	2.3	5.8	15.9

White, both sexes

2004.....	1.2	*	*	*	0.1	0.2	0.5	0.8	1.4	2.4	6.7	17.1
2003.....	1.2	*	*	*	0.1	0.3	0.5	0.9	1.4	3.0	6.5	17.8
2002.....	1.2	*	*	*	0.1	0.2	0.5	0.9	1.6	2.7	7.0	18.4
2001.....	1.2	*	*	*	*	0.2	0.4	0.7	1.5	3.2	6.9	19.6
2000.....	1.2	*	*	*	0.1	0.2	0.5	0.9	1.3	2.9	6.8	18.8
1999.....	0.9	*	*	*	*	0.2	0.4	0.6	1.2	2.3	5.6	14.3

White, male

2004.....	1.0	*	*	*	*	0.2	0.5	0.9	1.5	2.5	6.8	14.3
2003.....	1.0	*	*	*	*	0.2	0.6	0.9	1.6	2.8	5.6	13.3
2002.....	0.9	*	*	*	*	0.2	0.4	0.9	1.5	2.6	6.3	14.4
2001.....	1.0	*	*	*	*	0.2	0.4	0.8	1.5	3.3	6.7	15.0
2000.....	1.0	*	*	*	*	0.2	0.5	0.9	1.3	3.1	6.9	14.9
1999.....	0.7	*	*	*	*	0.2	0.4	0.6	1.1	2.5	5.5	11.4

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)

White, female

2004.....	1.4	*	*	*	*	0.2	0.4	0.7	1.2	2.4	6.6	18.4
2003.....	1.5	*	*	*	*	0.3	0.5	0.9	1.2	3.0	7.1	19.8
2002.....	1.5	*	*	*	*	0.3	0.5	1.0	1.6	2.8	7.5	20.2
2001.....	1.5	*	*	*	*	0.3	0.4	0.7	1.5	3.1	7.1	21.5
2000.....	1.4	*	*	*	*	0.3	0.4	0.8	1.4	2.8	6.8	20.4
1999.....	1.1	*	*	*	*	0.2	0.4	0.6	1.3	2.2	5.7	15.5

Black, both sexes

2004.....	1.7	*	*	*	*	0.9	1.7	2.3	3.6	4.8	10.6	22.8
2003.....	1.8	*	*	*	*	0.9	1.6	2.0	3.6	6.0	11.8	26.3
2002.....	1.7	*	*	*	*	0.9	1.5	2.1	3.7	5.2	12.9	23.8
2001.....	1.6	*	*	*	*	0.8	1.3	1.8	3.9	6.4	10.9	23.3
2000.....	1.6	*	*	*	*	0.9	1.3	2.2	3.4	6.9	10.8	21.9
1999.....	1.2	*	*	*	*	0.5	1.2	1.6	2.0	4.6	9.3	18.8

Black, male

2004.....	1.5	*	*	*	*	1.1	1.7	2.5	3.6	5.0	10.9	19.9
2003.....	1.6	*	*	*	*	0.8	1.2	2.6	4.0	6.6	12.7	25.0
2002.....	1.5	*	*	*	*	1.0	1.5	2.2	4.2	5.6	12.3	22.5
2001.....	1.5	*	*	*	*	*	1.4	1.8	4.9	6.7	10.9	*
2000.....	1.4	*	*	*	*	0.8	1.5	2.3	3.5	6.7	8.2	*
1999.....	1.1	*	*	*	*	*	1.3	1.5	1.9	5.7	10.1	*

Black, female

2004.....	1.8	*	*	*	*	0.7	1.8	2.2	3.5	4.6	10.4	24.0
2003.....	1.9	*	*	*	*	0.9	2.1	1.6	3.3	5.5	11.3	26.7
2002.....	1.8	*	*	*	*	0.9	1.4	2.0	3.3	5.0	13.2	24.2
2001.....	1.8	*	*	*	*	0.8	1.2	1.7	3.2	6.2	10.9	24.0
2000.....	1.8	*	*	*	*	0.9	1.0	2.1	3.3	7.1	12.3	22.3
1999.....	1.3	*	*	*	*	*	1.0	1.6	2.0	3.8	8.8	21.9

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

All other and unspecified disorders of circulatory system (I83-I99)

All races 3/, both sexes

2004.....	0.4	0.6	*	*	*	*	0.1	0.4	0.6	1.0	2.1	6.0
2003.....	0.4	0.7	*	*	*	*	0.1	0.3	0.5	0.9	2.2	5.9
2002.....	0.4	0.5	*	*	*	*	0.1	0.3	0.5	1.0	2.5	5.9
2001.....	0.4	*	*	*	*	*	0.1	0.3	0.5	1.2	2.3	6.5
2000.....	0.4	0.5	*	*	*	*	0.2	0.4	0.6	1.1	2.4	6.8
1999.....	0.5	*	*	*	*	*	0.1	0.3	0.7	1.3	3.4	8.8

All races 3/, male

2004.....	0.4	*	*	*	*	*	0.1	0.6	0.8	1.2	2.7	5.4
2003.....	0.4	*	*	*	*	*	0.1	0.5	0.6	1.1	2.5	5.3
2002.....	0.4	*	*	*	*	*	0.2	0.4	0.7	1.1	2.9	5.7
2001.....	0.4	*	*	*	*	*	0.2	0.5	0.7	1.4	2.9	6.8
2000.....	0.4	*	*	*	*	*	0.2	0.6	0.8	1.4	2.8	7.3
1999.....	0.5	*	*	*	*	*	0.2	0.5	0.9	1.6	3.9	9.6

All races 3/, female

2004.....	0.4	*	*	*	*	*	*	0.2	0.4	0.9	1.8	6.4
2003.....	0.4	*	*	*	*	*	0.1	0.2	0.4	0.8	2.1	6.1
2002.....	0.4	*	*	*	*	*	*	0.2	0.4	0.9	2.2	5.9
2001.....	0.4	*	*	*	*	*	*	0.2	0.4	1.0	1.8	6.4
2000.....	0.4	*	*	*	*	*	0.1	0.2	0.4	0.9	2.1	6.7
1999.....	0.5	*	*	*	*	*	*	0.2	0.5	1.1	3.0	8.5

White, both sexes

2004.....	0.4	*	*	*	*	*	0.1	0.4	0.6	1.0	2.1	6.0
2003.....	0.4	0.7	*	*	*	*	0.1	0.3	0.5	0.9	2.1	5.8
2002.....	0.4	*	*	*	*	*	0.1	0.3	0.5	0.9	2.4	5.9
2001.....	0.4	*	*	*	*	*	0.1	0.3	0.5	1.1	2.2	6.5
2000.....	0.4	*	*	*	*	*	0.1	0.4	0.6	1.0	2.3	6.7
1999.....	0.5	*	*	*	*	*	0.1	0.3	0.6	1.2	3.3	8.8

White, male

2004.....	0.4	*	*	*	*	*	0.1	0.6	0.8	1.2	2.6	5.4
2003.....	0.4	*	*	*	*	*	0.1	0.4	0.6	1.0	2.3	5.2
2002.....	0.4	*	*	*	*	*	0.1	0.4	0.6	1.0	2.7	5.5
2001.....	0.4	*	*	*	*	*	0.2	0.5	0.7	1.2	2.8	6.6
2000.....	0.4	*	*	*	*	*	0.2	0.5	0.7	1.3	2.6	7.1
1999.....	0.5	*	*	*	*	*	0.2	0.4	0.9	1.4	3.7	9.3

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
X. Diseases of the respiratory system (J00-J98)												
All races 3/, both sexes												
2004.....	77.3	15.8	1.9	0.7	1.3	2.0	5.6	17.2	63.6	229.1	627.0	1,497.1
2003.....	81.1	17.5	2.2	0.8	1.5	2.3	5.9	18.2	66.4	239.5	651.9	1,625.4
2002.....	81.4	15.8	1.9	0.8	1.3	2.3	6.0	17.6	65.7	240.3	662.2	1,663.7
2001.....	80.8	17.3	1.8	0.7	1.3	2.3	6.0	17.6	67.3	245.9	651.9	1,671.8
2000.....	82.1	17.1	1.7	0.8	1.4	2.5	6.2	17.9	68.8	250.2	667.5	1,734.0
1999.....	82.4	18.1	2.1	0.8	1.4	2.5	6.0	17.3	70.8	255.0	671.7	1,733.0
All races 3/, male												
2004.....	75.2	17.6	1.8	0.8	1.5	2.3	5.9	19.3	70.6	267.2	779.3	1,915.3
2003.....	78.2	20.6	2.5	0.9	1.6	2.7	6.1	20.3	74.0	279.9	808.5	2,057.2
2002.....	78.9	18.8	2.2	0.8	1.5	2.4	6.5	19.5	73.0	282.2	832.8	2,150.0
2001.....	78.2	19.7	1.9	0.8	1.5	2.4	6.4	19.7	74.6	289.4	824.1	2,187.2
2000.....	80.2	18.9	1.7	0.8	1.6	2.7	6.8	19.9	77.1	297.1	851.4	2,344.4
1999.....	81.3	19.6	2.2	0.9	1.5	2.6	6.4	19.1	79.5	310.4	870.5	2,356.4
All races 3/, female												
2004.....	79.4	14.1	1.9	0.6	1.1	1.7	5.4	15.3	57.2	197.1	524.4	1,309.0
2003.....	84.0	14.2	1.8	0.8	1.3	2.0	5.8	16.2	59.3	205.8	547.3	1,434.5
2002.....	83.7	12.6	1.5	0.8	1.1	2.1	5.6	15.8	58.9	205.5	548.9	1,452.7
2001.....	83.2	14.7	1.8	0.7	1.2	2.3	5.6	15.5	60.6	209.9	538.8	1,456.2
2000.....	84.0	15.2	1.7	0.7	1.2	2.3	5.7	16.0	61.1	211.7	547.6	1,485.4
1999.....	83.4	16.6	1.9	0.7	1.2	2.4	5.7	15.6	62.9	209.7	542.6	1,481.7
White, both sexes												
2004.....	86.4	12.2	1.5	0.5	1.1	1.7	5.0	16.2	64.4	239.6	649.6	1,541.7
2003.....	90.6	14.4	1.8	0.7	1.2	2.0	5.2	17.1	66.9	249.6	674.7	1,676.3
2002.....	90.8	12.0	1.4	0.6	1.1	2.0	5.3	16.6	66.2	250.2	683.4	1,713.3
2001.....	89.7	12.3	1.5	0.6	1.1	1.9	5.1	16.3	68.1	255.1	671.9	1,713.5
2000.....	91.2	13.3	1.5	0.6	1.2	2.1	5.4	16.4	69.0	257.5	687.3	1,783.5
1999.....	91.2	13.3	1.6	0.7	1.2	2.0	5.0	15.9	71.2	262.7	690.3	1,778.5
White, male												
2004.....	82.8	13.5	1.5	0.6	1.3	2.0	5.2	18.0	69.9	274.0	797.2	1,960.9
2003.....	85.8	16.9	2.0	0.7	1.3	2.2	5.3	19.1	73.0	286.1	822.4	2,106.7
2002.....	86.7	14.6	1.7	0.6	1.2	2.1	5.6	18.1	72.0	289.1	848.3	2,201.7
2001.....	85.7	14.3	1.5	0.6	1.2	1.9	5.6	18.1	74.0	295.1	838.0	2,230.5
2000.....	87.7	15.1	1.3	0.7	1.3	2.1	5.9	17.9	76.1	299.6	866.8	2,402.2
1999.....	88.8	14.7	1.5	0.8	1.3	2.1	5.2	17.2	78.4	314.6	884.0	2,409.4

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
X. Diseases of the respiratory system (J00-J98)												
White, female												
2004.....	89.9	10.9	1.4	0.5	0.9	1.5	4.9	14.4	59.2	210.1	548.8	1,352.8
2003.....	95.2	11.8	1.6	0.7	1.1	1.8	5.0	15.2	61.2	218.4	574.7	1,485.6
2002.....	94.7	9.2	1.2	0.7	1.0	1.9	5.0	15.1	60.7	217.2	572.8	1,501.2
2001.....	93.7	10.2	1.6	0.5	0.9	1.9	4.7	14.5	62.5	221.3	561.5	1,497.0
2000.....	94.6	11.3	1.6	0.6	1.0	2.0	4.9	15.0	62.4	222.2	569.1	1,532.2
1999.....	93.6	11.8	1.7	0.6	1.1	1.8	4.8	14.6	64.4	219.4	563.4	1,525.0
Black, both sexes												
2004.....	44.2	33.4	3.7	1.5	2.4	4.2	11.1	29.3	75.5	194.3	482.2	1,082.3
2003.....	45.9	32.7	3.5	1.5	2.9	5.0	11.8	30.6	78.5	206.2	499.7	1,135.5
2002.....	46.7	34.1	4.1	1.8	2.7	4.6	12.5	29.2	79.6	206.0	521.7	1,204.5
2001.....	47.2	39.6	3.2	1.3	2.8	5.6	13.0	31.4	77.5	212.7	515.2	1,258.0
2000.....	48.0	37.7	3.1	1.5	2.9	5.2	12.7	33.0	83.6	231.0	518.4	1,239.4
1999.....	49.2	41.6	4.0	1.6	2.7	6.3	13.7	32.5	84.8	231.0	532.5	1,298.0
Black, male												
2004.....	47.9	37.1	3.5	2.0	2.6	5.5	12.6	34.2	97.8	270.6	697.1	1,501.1
2003.....	50.0	38.7	4.5	1.9	3.5	6.5	12.1	34.5	101.9	285.5	753.9	1,577.5
2002.....	50.2	39.2	4.7	1.8	3.3	5.1	14.3	34.3	102.7	278.9	766.8	1,685.3
2001.....	50.5	43.7	3.8	1.5	3.0	6.3	13.7	36.4	97.4	292.5	773.9	1,775.7
2000.....	52.4	39.3	3.9	1.8	3.4	6.8	14.5	39.5	105.0	326.7	758.8	1,808.3
1999.....	54.0	44.6	5.1	1.9	3.2	6.4	15.2	39.4	109.3	324.6	806.8	1,900.6
Black, female												
2004.....	40.9	29.6	3.8	1.1	2.2	2.9	9.8	25.1	57.7	139.7	360.8	916.3
2003.....	42.1	26.6	2.4	1.2	2.3	3.7	11.6	27.3	59.6	149.7	355.9	963.7
2002.....	43.5	28.8	3.5	1.8	2.1	4.1	10.8	24.9	61.0	154.3	383.0	1,020.3
2001.....	44.1	35.3	2.5	1.2	2.5	5.0	12.4	27.0	61.5	156.4	370.0	1,066.0
2000.....	44.1	36.0	2.2	1.2	2.5	3.8	11.1	27.4	66.3	163.6	383.0	1,027.6
1999.....	44.8	38.6	2.9	1.2	2.3	6.1	12.5	26.5	65.2	165.0	379.0	1,072.8

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other diseases of the respiratory system (J10-J98)

All races 3/, both sexes

2004.....	77.3	15.7	1.8	0.7	1.3	2.0	5.6	17.2	63.6	229.0	626.8	1,496.3
2003.....	81.1	17.1	2.1	0.8	1.4	2.3	5.9	18.2	66.3	239.5	651.8	1,623.9
2002.....	81.3	15.5	1.8	0.8	1.3	2.3	6.0	17.6	65.6	240.2	662.0	1,662.6
2001.....	80.7	17.0	1.7	0.7	1.3	2.3	6.0	17.5	67.3	245.9	651.7	1,670.6
2000.....	82.0	16.9	1.7	0.8	1.4	2.4	6.2	17.9	68.7	250.1	667.2	1,732.2
1999.....	82.3	17.8	2.0	0.8	1.4	2.5	6.0	17.3	70.8	255.0	671.3	1,731.2

All races 3/, male

2004.....	75.1	17.4	1.8	0.8	1.4	2.3	5.9	19.2	70.6	267.1	779.2	1,914.6
2003.....	78.1	20.1	2.4	0.9	1.6	2.7	6.0	20.3	73.9	279.9	808.3	2,055.5
2002.....	78.9	18.6	2.1	0.8	1.5	2.4	6.5	19.4	72.9	282.1	832.7	2,149.3
2001.....	78.1	19.4	1.8	0.7	1.5	2.4	6.4	19.7	74.5	289.4	824.0	2,186.6
2000.....	80.1	18.7	1.7	0.8	1.6	2.6	6.8	19.9	77.1	296.9	851.2	2,343.3
1999.....	81.3	19.2	2.1	0.9	1.5	2.6	6.3	19.1	79.5	310.3	870.1	2,355.3

All races 3/, female

2004.....	79.4	14.0	1.8	0.5	1.1	1.7	5.4	15.3	57.1	197.0	524.2	1,308.1
2003.....	83.9	13.9	1.8	0.8	1.3	2.0	5.7	16.2	59.2	205.7	547.2	1,433.1
2002.....	83.7	12.4	1.4	0.8	1.1	2.1	5.6	15.8	58.9	205.4	548.8	1,451.5
2001.....	83.2	14.6	1.7	0.6	1.1	2.3	5.6	15.5	60.6	209.8	538.5	1,454.6
2000.....	83.9	15.1	1.6	0.7	1.2	2.2	5.6	16.0	61.1	211.5	547.3	1,483.4
1999.....	83.3	16.4	1.9	0.7	1.2	2.4	5.7	15.6	62.8	209.6	542.2	1,479.7

White, both sexes

2004.....	86.3	12.1	1.4	0.5	1.1	1.7	5.0	16.1	64.3	239.5	649.3	1,540.9
2003.....	90.5	14.1	1.7	0.7	1.2	2.0	5.1	17.1	66.9	249.5	674.5	1,674.8
2002.....	90.7	11.9	1.3	0.6	1.1	2.0	5.3	16.6	66.1	250.1	683.3	1,712.2
2001.....	89.7	12.1	1.4	0.6	1.1	1.9	5.1	16.2	68.1	255.0	671.6	1,712.1
2000.....	91.1	13.1	1.4	0.6	1.1	2.1	5.4	16.4	69.0	257.4	687.0	1,781.7
1999.....	91.2	13.1	1.5	0.7	1.2	2.0	5.0	15.8	71.1	262.6	689.9	1,776.7

White, male

2004.....	82.7	13.2	1.4	0.5	1.3	2.0	5.1	18.0	69.8	274.0	797.0	1,960.3
2003.....	85.8	16.4	1.9	0.7	1.3	2.2	5.3	19.1	72.9	286.0	822.2	2,105.2
2002.....	86.7	14.5	1.6	0.6	1.2	2.1	5.6	18.1	71.9	289.0	848.2	2,201.0
2001.....	85.6	14.1	1.4	0.6	1.2	1.9	5.6	18.1	74.0	295.1	837.9	2,229.8
2000.....	87.6	14.8	1.3	0.7	1.3	2.1	5.8	17.9	76.0	299.5	866.6	2,401.1
1999.....	88.7	14.5	1.4	0.7	1.3	2.1	5.1	17.2	78.4	314.5	883.6	2,408.6

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other diseases of the respiratory system (J10-J98)												
White, female												
2004.....	89.9	10.8	1.3	0.4	0.9	1.5	4.9	14.3	59.1	210.0	548.6	1,351.9
2003.....	95.2	11.6	1.5	0.7	1.1	1.7	5.0	15.2	61.1	218.4	574.6	1,484.2
2002.....	94.7	9.0	1.1	0.7	0.9	1.9	5.0	15.1	60.6	217.1	572.6	1,499.8
2001.....	93.6	10.0	1.5	0.5	0.9	1.9	4.7	14.5	62.4	221.2	561.3	1,495.4
2000.....	94.5	11.3	1.6	0.6	0.9	2.0	4.9	15.0	62.4	222.0	568.8	1,530.1
1999.....	93.5	11.7	1.6	0.6	1.1	1.8	4.8	14.5	64.4	219.4	563.0	1,522.9
Black, both sexes												
2004.....	44.2	33.4	3.7	1.5	2.4	4.2	11.0	29.3	75.5	194.1	482.1	1,081.5
2003.....	45.8	32.3	3.4	1.5	2.9	5.0	11.7	30.5	78.4	206.1	499.7	1,134.0
2002.....	46.6	33.5	4.0	1.8	2.7	4.6	12.4	29.1	79.6	205.9	521.5	1,204.5
2001.....	47.1	39.1	3.0	1.3	2.7	5.6	13.0	31.3	77.4	212.7	515.1	1,257.1
2000.....	48.0	37.6	3.0	1.5	2.9	5.2	12.7	32.9	83.6	230.9	518.3	1,238.8
1999.....	49.1	41.0	3.9	1.5	2.7	6.2	13.7	32.4	84.7	230.8	532.4	1,296.4
Black, male												
2004.....	47.8	37.1	3.5	2.0	2.6	5.5	12.5	34.1	97.7	270.5	696.8	1,500.1
2003.....	49.9	38.4	4.4	1.9	3.4	6.5	12.0	34.3	101.7	285.3	753.9	1,574.4
2002.....	50.2	38.3	4.7	1.8	3.3	5.1	14.2	34.2	102.7	278.7	766.5	1,685.3
2001.....	50.4	42.8	3.6	1.4	3.0	6.2	13.6	36.4	97.3	292.5	773.9	1,775.7
2000.....	52.4	39.3	3.9	1.8	3.3	6.7	14.5	39.5	105.0	326.4	758.5	1,806.0
1999.....	53.9	43.9	4.8	1.9	3.2	6.3	15.2	39.4	109.1	324.4	806.8	1,895.9
Black, female												
2004.....	40.9	29.6	3.8	1.1	2.2	2.9	9.8	25.1	57.7	139.6	360.8	915.5
2003.....	42.1	26.0	2.3	1.1	2.3	3.7	11.5	27.2	59.6	149.7	355.9	962.9
2002.....	43.4	28.5	3.4	1.8	2.0	4.1	10.8	24.8	60.9	154.2	382.8	1,020.3
2001.....	44.1	35.3	2.5	1.2	2.5	5.0	12.4	26.9	61.5	156.3	369.8	1,064.7
2000.....	44.0	35.7	2.2	1.2	2.5	3.8	11.1	27.3	66.3	163.6	383.0	1,027.6
1999.....	44.7	37.9	2.9	1.2	2.3	6.1	12.4	26.5	65.1	164.8	378.8	1,072.3

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Influenza (J10-J11)												
All races 3/, both sexes												
2004.....	0.4	*	*	*	*	*	*	0.1	0.2	0.5	2.2	11.4
2003.....	0.6	0.8	0.4	0.1	0.1	*	0.1	0.1	0.3	1.0	3.6	17.0
2002.....	0.3	*	*	*	*	*	*	*	0.1	0.3	1.4	8.8
2001.....	0.1	*	*	*	*	*	*	*	0.1	0.1	0.4	2.2
2000.....	0.6	*	*	*	*	*	0.0	0.1	0.4	1.1	4.0	20.2
1999.....	0.6	*	*	*	*	*	*	0.1	0.3	0.9	3.7	21.2
All races 3/, male												
2004.....	0.3	*	*	*	*	*	*	*	0.2	0.7	2.4	11.1
2003.....	0.5	*	0.4	*	*	*	*	0.2	0.3	1.1	3.7	17.0
2002.....	0.2	*	*	*	*	*	*	*	0.2	0.4	1.6	8.9
2001.....	0.1	*	*	*	*	*	*	*	*	*	0.4	2.2
2000.....	0.5	*	*	*	*	*	*	0.1	0.4	1.3	4.0	20.7
1999.....	0.4	*	*	*	*	*	*	*	0.3	0.9	4.2	19.9
All races 3/, female												
2004.....	0.5	*	*	*	*	*	*	*	0.2	0.4	2.1	11.5
2003.....	0.7	*	0.3	0.1	*	*	*	0.1	0.3	0.9	3.5	17.1
2002.....	0.3	*	*	*	*	*	*	*	*	0.2	1.3	8.7
2001.....	0.1	*	*	*	*	*	*	*	*	*	0.5	2.2
2000.....	0.8	*	*	*	*	*	*	0.1	0.4	0.9	4.0	19.9
1999.....	0.7	*	*	*	*	*	*	*	0.3	0.9	3.4	21.7
White, both sexes												
2004.....	0.4	*	*	*	*	*	*	0.1	0.2	0.6	2.3	12.3
2003.....	0.7	0.7	0.3	0.1	0.1	*	0.1	0.1	0.3	1.1	3.8	18.2
2002.....	0.3	*	*	*	*	*	*	*	0.1	0.3	1.5	9.4
2001.....	0.1	*	*	*	*	*	*	*	*	*	0.5	2.3
2000.....	0.7	*	*	*	*	*	*	0.1	0.4	1.1	4.2	21.4
1999.....	0.7	*	*	*	*	*	*	0.1	0.3	0.9	3.9	22.8
White, male												
2004.....	0.3	*	*	*	*	*	*	*	0.2	0.7	2.4	12.1
2003.....	0.5	*	0.4	*	*	*	*	0.2	0.3	1.1	3.9	18.3
2002.....	0.2	*	*	*	*	*	*	*	*	0.4	1.7	9.7
2001.....	0.1	*	*	*	*	*	*	*	*	*	0.4	2.3
2000.....	0.5	*	*	*	*	*	*	*	0.4	1.3	4.2	21.6
1999.....	0.5	*	*	*	*	*	*	*	0.4	0.9	4.4	21.7

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pneumonia (J12-J18)												
All races 3/, both sexes												
2004.....	19.9	6.3	0.6	0.2	0.4	0.7	2.0	4.5	10.7	34.0	137.0	571.2
2003.....	21.8	7.3	0.7	0.2	0.5	0.9	2.2	5.1	10.9	36.3	147.5	649.1
2002.....	22.5	6.3	0.7	0.2	0.4	0.9	2.1	4.7	11.1	37.2	155.5	687.8
2001.....	21.7	7.2	0.7	0.2	0.4	0.8	2.2	4.6	10.6	36.2	148.1	683.4
2000.....	22.6	7.4	0.6	0.2	0.4	0.9	2.3	4.6	11.5	38.0	156.4	723.9
1999.....	22.2	8.1	0.8	0.2	0.4	0.8	2.3	4.6	10.7	36.4	153.3	730.6
All races 3/, male												
2004.....	18.3	7.5	0.5	0.2	0.5	0.9	2.2	5.5	12.8	42.1	170.3	666.1
2003.....	19.6	8.5	0.7	0.2	0.5	1.1	2.5	6.2	12.9	45.4	183.2	740.9
2002.....	20.2	7.6	0.8	0.2	0.4	0.9	2.4	5.7	13.4	46.0	193.1	796.9
2001.....	19.5	7.8	0.7	0.2	0.5	0.9	2.5	5.7	13.0	44.9	186.0	806.2
2000.....	20.3	8.1	0.6	0.2	0.5	1.0	2.7	5.6	14.0	47.1	195.8	871.3
1999.....	19.8	8.5	0.7	0.2	0.4	0.9	2.6	5.8	12.5	46.0	192.3	874.7
All races 3/, female												
2004.....	21.5	5.0	0.8	0.2	0.3	0.5	1.8	3.5	8.6	27.3	114.6	528.5
2003.....	23.9	6.0	0.6	0.3	0.4	0.7	1.9	4.0	9.1	28.6	123.7	608.5
2002.....	24.8	5.0	0.6	0.2	0.3	0.8	1.9	3.8	9.0	29.8	130.6	640.5
2001.....	23.8	6.7	0.7	0.2	0.4	0.7	1.9	3.5	8.4	29.0	123.2	632.0
2000.....	24.8	6.6	0.6	0.2	0.4	0.8	1.9	3.6	9.1	30.6	130.6	663.9
1999.....	24.6	7.7	0.8	0.2	0.4	0.7	2.0	3.4	9.1	28.4	128.0	672.5
White, both sexes												
2004.....	21.6	4.7	0.5	0.1	0.4	0.7	1.7	4.0	9.9	32.8	136.2	578.4
2003.....	23.7	6.0	0.6	0.2	0.5	0.8	1.9	4.5	10.0	34.8	147.7	661.1
2002.....	24.6	5.0	0.5	0.2	0.4	0.8	1.9	4.3	10.2	36.4	155.4	699.6
2001.....	23.5	5.1	0.6	0.2	0.4	0.7	1.8	4.0	9.8	35.2	146.8	692.7
2000.....	24.4	5.6	0.5	0.2	0.4	0.8	2.0	4.0	10.4	36.0	155.8	737.0
1999.....	24.1	5.9	0.6	0.2	0.4	0.7	1.9	3.9	9.8	35.2	153.9	741.6
White, male												
2004.....	19.5	5.8	0.4	0.2	0.6	0.9	1.9	4.9	11.9	39.8	167.8	673.4
2003.....	20.9	7.1	0.7	0.2	0.5	1.0	2.2	5.5	11.3	42.8	181.0	748.8
2002.....	21.7	6.4	0.5	0.1	0.4	0.8	2.1	5.1	12.1	44.8	191.4	806.2
2001.....	20.6	5.8	0.6	0.2	0.5	0.8	2.1	4.8	11.7	42.9	182.2	812.1
2000.....	21.5	6.1	0.4	0.2	0.5	0.8	2.4	4.7	12.5	43.8	193.6	884.4
1999.....	21.0	6.7	0.4	0.2	0.4	0.8	2.1	4.8	11.1	43.8	191.7	884.0

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pneumonia (J12-J18)												
White, female												
2004.....	23.6	3.5	0.6	0.1	0.3	0.5	1.6	3.1	8.1	26.8	114.6	535.6
2003.....	26.5	4.8	0.5	0.2	0.4	0.6	1.7	3.6	8.7	28.0	125.2	622.3
2002.....	27.4	3.6	0.5	0.2	0.3	0.7	1.7	3.5	8.5	29.3	131.3	653.3
2001.....	26.2	4.4	0.6	0.2	0.3	0.7	1.6	3.1	8.0	28.7	123.3	642.8
2000.....	27.3	5.0	0.6	0.2	0.3	0.7	1.7	3.2	8.5	29.4	130.8	677.1
1999.....	27.1	5.2	0.7	0.2	0.3	0.5	1.7	3.0	8.5	27.9	129.1	684.4
Black, both sexes												
2004.....	14.6	13.5	1.2	*	0.6	1.2	4.0	9.0	18.9	49.4	155.6	502.5
2003.....	15.1	13.0	0.9	*	0.6	1.8	4.4	10.1	20.3	53.7	154.5	519.6
2002.....	15.5	12.7	1.5	*	0.6	1.5	4.3	8.7	21.1	49.9	165.8	572.8
2001.....	15.5	16.3	1.0	*	0.7	1.6	4.7	9.7	18.8	48.2	168.6	579.7
2000.....	16.1	16.4	1.2	*	0.7	1.7	4.4	9.9	21.8	59.3	169.7	591.5
1999.....	16.1	18.3	1.7	*	0.8	1.9	5.4	10.5	20.7	53.3	155.7	629.1
Black, male												
2004.....	14.9	15.1	*	*	*	1.6	4.8	11.7	24.8	71.9	216.6	597.1
2003.....	15.8	15.2	*	*	0.7	2.2	5.2	12.9	28.7	77.7	217.5	638.1
2002.....	15.3	13.7	1.6	*	*	1.7	5.1	11.2	28.7	65.5	228.4	687.2
2001.....	15.8	15.9	*	*	0.7	2.0	5.7	13.3	26.1	69.1	242.2	712.6
2000.....	16.5	17.9	1.7	*	0.7	2.2	5.5	13.6	29.7	82.8	232.5	742.7
1999.....	16.3	17.9	1.9	*	0.8	2.2	6.6	15.0	27.1	76.0	211.8	800.3
Black, female												
2004.....	14.4	11.7	1.7	*	0.6	0.8	3.2	6.7	14.1	33.3	121.2	465.0
2003.....	14.6	10.8	*	*	*	1.4	3.6	7.7	13.6	36.6	119.0	473.5
2002.....	15.6	11.8	*	*	*	1.3	3.7	6.5	14.9	38.8	130.4	529.0
2001.....	15.1	16.7	*	*	0.7	1.3	3.9	6.7	13.0	33.4	127.2	530.5
2000.....	15.8	14.9	*	*	0.8	1.2	3.4	6.6	15.4	42.7	134.3	535.3
1999.....	15.9	18.8	*	*	0.9	1.7	4.3	6.8	15.5	37.2	124.3	565.1

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Bacterial pneumonia (J13-J15)												
All races 3/, both sexes												
2004.....	1.5	0.8	*	*	*	0.1	0.3	0.7	1.4	3.4	10.5	30.1
2003.....	1.6	1.0	0.2	0.1	0.1	0.1	0.3	0.8	1.4	3.7	11.1	34.9
2002.....	1.6	1.1	*	*	*	0.1	0.3	0.7	1.3	3.6	11.3	35.7
2001.....	1.6	0.8	*	*	*	0.1	0.3	0.6	1.5	3.5	11.0	34.8
2000.....	1.6	0.8	*	*	*	0.1	0.4	0.7	1.6	3.7	11.0	35.7
1999.....	1.4	1.3	0.1	*	*	0.1	0.4	0.5	1.3	3.3	9.8	30.1
All races 3/, male												
2004.....	1.5	*	*	*	*	0.1	0.3	0.8	1.5	4.2	14.1	41.1
2003.....	1.7	1.2	*	*	*	0.1	0.4	0.9	1.7	4.9	14.5	45.7
2002.....	1.6	1.2	*	*	*	0.1	0.3	0.7	1.6	4.4	15.1	46.8
2001.....	1.6	*	*	*	*	0.1	0.3	0.7	1.8	4.5	15.1	46.8
2000.....	1.7	1.1	*	*	*	0.1	0.4	0.8	1.8	4.8	15.6	52.3
1999.....	1.4	1.2	*	*	*	*	0.4	0.7	1.4	4.2	12.5	40.4
All races 3/, female												
2004.....	1.5	*	*	*	*	*	0.3	0.5	1.3	2.7	8.0	25.2
2003.....	1.6	*	*	*	*	*	0.3	0.6	1.1	2.7	8.8	30.2
2002.....	1.6	*	*	*	*	0.1	0.3	0.6	1.1	2.9	8.8	31.0
2001.....	1.5	*	*	*	*	*	0.2	0.5	1.2	2.6	8.3	29.8
2000.....	1.5	*	*	*	*	0.1	0.3	0.6	1.3	2.9	8.0	29.0
1999.....	1.4	1.5	*	*	*	*	0.4	0.4	1.2	2.6	7.9	26.0
White, both sexes												
2004.....	1.6	0.7	*	*	*	0.1	0.3	0.6	1.3	3.3	10.2	29.9
2003.....	1.7	0.8	*	0.1	*	0.1	0.3	0.7	1.2	3.7	11.1	34.9
2002.....	1.7	0.9	*	*	*	0.1	0.3	0.6	1.2	3.6	11.2	36.3
2001.....	1.6	0.8	*	*	*	0.1	0.2	0.6	1.4	3.4	10.9	35.0
2000.....	1.7	*	*	*	*	0.1	0.4	0.6	1.5	3.6	10.8	35.6
1999.....	1.5	0.9	*	*	*	0.1	0.3	0.5	1.2	3.2	9.7	30.3
White, male												
2004.....	1.6	*	*	*	*	0.1	0.2	0.7	1.3	3.9	13.7	41.7
2003.....	1.8	*	*	*	*	0.1	0.3	0.8	1.4	4.8	14.5	45.9
2002.....	1.7	*	*	*	*	0.1	0.3	0.6	1.4	4.3	14.7	47.9
2001.....	1.7	*	*	*	*	*	0.3	0.6	1.7	4.3	14.7	46.6
2000.....	1.7	*	*	*	*	*	0.4	0.7	1.7	4.5	15.3	51.1
1999.....	1.4	*	*	*	*	*	0.3	0.6	1.3	4.0	12.3	40.1

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Bacterial pneumonia (J13-J15)												
White, female												
2004.....	1.5	*	*	*	*	*	0.3	0.5	1.3	2.7	7.8	24.7
2003.....	1.7	*	*	*	*	*	0.3	0.6	1.1	2.8	8.8	30.1
2002.....	1.7	*	*	*	*	*	0.3	0.6	1.1	3.0	8.9	31.2
2001.....	1.6	*	*	*	*	*	0.2	0.6	1.2	2.6	8.3	30.2
2000.....	1.6	*	*	*	*	0.1	0.3	0.5	1.3	2.8	7.8	29.2
1999.....	1.5	*	*	*	*	*	0.4	0.4	1.1	2.5	8.0	26.3
Black, both sexes												
2004.....	1.3	*	*	*	*	*	0.5	1.3	2.4	4.8	13.4	35.0
2003.....	1.3	*	*	*	*	*	0.6	1.3	2.7	4.0	11.6	35.0
2002.....	1.2	*	*	*	*	*	0.6	1.0	2.2	4.0	11.4	28.5
2001.....	1.2	*	*	*	*	*	0.6	0.9	2.2	4.6	13.0	32.5
2000.....	1.3	*	*	*	*	*	0.5	1.1	2.4	5.7	12.2	34.7
1999.....	1.2	*	*	*	*	*	0.8	1.0	2.3	4.6	9.7	25.2
Black, male												
2004.....	1.4	*	*	*	*	*	*	1.5	3.6	7.5	18.7	41.7
2003.....	1.5	*	*	*	*	*	0.9	1.7	4.1	5.8	15.2	41.7
2002.....	1.3	*	*	*	*	*	*	1.2	3.2	5.0	18.1	32.2
2001.....	1.5	*	*	*	*	*	0.8	1.5	2.8	7.2	20.3	49.9
2000.....	1.5	*	*	*	*	*	*	1.2	3.4	7.8	17.9	60.0
1999.....	1.3	*	*	*	*	*	0.9	1.5	2.6	6.5	14.5	38.7
Black, female												
2004.....	1.3	*	*	*	*	*	*	1.0	1.5	2.9	10.4	32.3
2003.....	1.2	*	*	*	*	*	*	1.0	1.5	2.8	9.6	32.4
2002.....	1.2	*	*	*	*	*	*	0.9	1.4	3.2	7.6	27.1
2001.....	1.0	*	*	*	*	*	*	*	1.8	2.7	8.9	26.1
2000.....	1.2	*	*	*	*	*	*	1.1	1.6	4.3	9.0	25.3
1999.....	1.1	*	*	*	*	*	0.8	*	2.1	3.3	7.1	20.2

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pneumonia due to other or unspecified organisms (J16,J18)												
All races 3/, both sexes												
2004.....	18.4	4.1	0.4	0.1	0.4	0.6	1.7	3.8	9.2	30.6	126.1	539.8
2003.....	20.0	4.6	0.4	0.2	0.4	0.8	1.8	4.3	9.5	32.4	135.9	612.1
2002.....	20.8	3.9	0.5	0.2	0.3	0.7	1.8	4.1	9.7	33.4	143.7	650.3
2001.....	20.0	4.6	0.5	0.2	0.4	0.7	1.9	3.9	9.1	32.6	136.6	647.1
2000.....	20.8	5.1	0.4	0.1	0.4	0.7	1.9	3.9	9.8	34.0	144.6	685.3
1999.....	20.7	4.7	0.5	0.1	0.4	0.7	1.9	4.0	9.4	32.8	142.8	698.1
All races 3/, male												
2004.....	16.7	4.9	0.3	0.1	0.5	0.8	1.9	4.7	11.3	37.8	155.8	623.9
2003.....	17.8	5.5	0.4	0.2	0.5	0.9	2.1	5.3	11.1	40.3	168.0	693.0
2002.....	18.5	4.6	0.5	0.1	0.4	0.8	2.1	5.0	11.8	41.4	177.4	748.2
2001.....	17.8	4.9	0.5	0.2	0.4	0.8	2.1	4.9	11.2	40.2	170.0	757.4
2000.....	18.5	5.3	0.4	0.2	0.4	0.9	2.3	4.8	12.1	42.0	179.5	816.5
1999.....	18.3	5.0	0.5	0.2	0.4	0.8	2.2	5.1	11.0	41.7	178.5	832.5
All races 3/, female												
2004.....	20.0	3.3	0.5	0.1	0.3	0.5	1.4	2.9	7.3	24.5	106.2	502.0
2003.....	22.2	3.8	0.5	0.1	0.3	0.6	1.6	3.4	7.9	25.7	114.5	576.3
2002.....	23.0	3.1	0.4	0.2	0.3	0.6	1.5	3.2	7.8	26.7	121.3	607.9
2001.....	22.2	4.2	0.5	0.2	0.3	0.6	1.6	2.9	7.1	26.3	114.6	601.0
2000.....	23.1	4.9	0.4	0.1	0.4	0.6	1.6	3.0	7.7	27.4	121.9	631.9
1999.....	23.0	4.3	0.5	0.1	0.4	0.6	1.6	2.9	7.9	25.6	119.6	643.9
White, both sexes												
2004.....	19.9	2.9	0.3	0.1	0.4	0.6	1.5	3.4	8.6	29.5	125.5	547.1
2003.....	21.8	3.8	0.4	0.2	0.4	0.7	1.6	3.8	8.7	31.0	136.2	624.0
2002.....	22.7	2.9	0.4	0.2	0.3	0.6	1.6	3.7	8.9	32.7	143.6	661.6
2001.....	21.7	3.0	0.5	0.2	0.4	0.6	1.6	3.3	8.3	31.7	135.4	656.2
2000.....	22.6	3.9	0.3	0.1	0.4	0.7	1.7	3.3	8.8	32.1	144.2	698.3
1999.....	22.5	3.3	0.4	0.1	0.3	0.6	1.5	3.4	8.5	31.7	143.4	708.8
White, male												
2004.....	17.8	3.4	*	*	0.5	0.7	1.6	4.2	10.5	35.8	153.6	630.7
2003.....	19.0	4.5	0.4	0.2	0.4	0.8	1.8	4.7	9.9	37.8	165.9	700.5
2002.....	19.9	3.7	0.4	0.1	0.4	0.7	1.8	4.4	10.7	40.2	176.0	756.4
2001.....	18.9	3.5	0.5	0.2	0.4	0.6	1.8	4.1	10.0	38.4	166.7	763.5
2000.....	19.6	4.3	*	0.2	0.4	0.7	1.9	4.0	10.7	39.0	177.4	830.6
1999.....	19.5	3.8	*	0.2	0.3	0.7	1.7	4.2	9.7	39.6	178.2	841.9

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pneumonia due to other or unspecified organisms (J16,J18)												
White, female												
2004.....	22.0	2.5	0.3	*	0.2	0.5	1.3	2.6	6.8	24.0	106.3	509.4
2003.....	24.6	3.0	0.4	0.1	0.3	0.5	1.4	3.0	7.5	25.1	116.0	590.1
2002.....	25.5	2.0	0.3	0.2	0.3	0.6	1.4	2.9	7.3	26.2	121.9	620.4
2001.....	24.5	2.5	0.4	0.2	0.3	0.6	1.4	2.6	6.7	26.1	114.7	611.3
2000.....	25.5	3.5	0.4	0.1	0.3	0.6	1.4	2.7	7.1	26.3	122.3	644.6
1999.....	25.4	2.8	0.4	*	0.3	0.5	1.3	2.6	7.3	25.2	120.6	655.4
Black, both sexes												
2004.....	13.2	10.3	1.0	*	0.6	1.0	3.5	7.7	16.4	44.6	142.2	467.3
2003.....	13.7	8.8	*	*	0.5	1.6	3.8	8.8	17.6	49.6	142.7	484.0
2002.....	14.2	8.2	1.0	*	0.5	1.3	3.7	7.6	18.8	45.9	154.2	543.4
2001.....	14.1	11.8	0.8	*	0.7	1.4	4.2	8.8	16.6	43.4	155.5	546.9
2000.....	14.7	11.2	0.8	*	0.6	1.6	3.9	8.7	19.3	53.4	157.4	556.5
1999.....	14.7	11.4	1.1	*	0.7	1.7	4.5	9.5	18.3	48.5	145.4	602.6
Black, male												
2004.....	13.4	12.5	*	*	*	1.3	4.4	10.1	21.2	64.4	197.9	554.3
2003.....	14.2	10.1	*	*	0.7	1.9	4.4	11.2	24.4	71.9	201.7	595.3
2002.....	13.9	8.1	*	*	*	1.6	4.4	10.0	25.5	60.5	209.7	652.9
2001.....	14.3	11.1	*	*	0.7	1.8	4.9	11.8	23.3	61.5	221.9	661.5
2000.....	14.9	10.6	*	*	*	2.0	4.9	12.4	26.1	74.9	214.7	682.7
1999.....	14.8	12.0	*	*	0.7	2.0	5.6	13.4	24.5	69.6	195.8	760.5
Black, female												
2004.....	13.0	8.0	*	*	*	*	2.6	5.7	12.6	30.4	110.8	432.8
2003.....	13.3	7.4	*	*	*	1.3	3.2	6.7	12.1	33.7	109.4	440.7
2002.....	14.4	8.2	*	*	*	1.1	3.1	5.6	13.5	35.5	122.8	501.5
2001.....	14.0	12.6	*	*	0.7	1.0	3.6	6.2	11.2	30.7	118.2	504.4
2000.....	14.6	11.9	*	*	*	1.1	2.9	5.6	13.7	38.2	125.1	509.5
1999.....	14.6	10.7	*	*	0.7	1.4	3.5	6.2	13.4	33.7	117.2	543.6

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other acute lower respiratory infections (J20-J22)												
All races 3/, both sexes												
2004.....	0.1	1.5	*	*	*	*	*	*	*	0.1	0.4	2.6
2003.....	0.1	1.3	0.2	*	*	*	*	*	0.1	0.2	0.6	3.5
2002.....	0.1	1.3	0.1	*	*	*	*	*	*	0.2	0.5	3.3
2001.....	0.1	1.2	*	*	*	*	*	*	*	0.2	0.5	3.3
2000.....	0.2	1.1	*	*	*	*	*	*	*	0.2	0.7	4.5
1999.....	0.2	1.7	0.2	*	*	*	*	*	0.1	0.2	0.9	6.0
All races 3/, male												
2004.....	0.1	1.6	*	*	*	*	*	*	*	*	*	2.5
2003.....	0.1	1.7	*	*	*	*	*	*	*	*	0.7	2.4
2002.....	0.1	1.9	*	*	*	*	*	*	*	*	0.5	2.8
2001.....	0.1	1.3	*	*	*	*	*	*	*	*	0.4	2.3
2000.....	0.1	1.3	*	*	*	*	*	*	*	0.3	0.7	3.8
1999.....	0.1	1.9	*	*	*	*	*	*	*	*	1.1	4.0
All races 3/, female												
2004.....	0.1	1.3	*	*	*	*	*	*	*	*	0.5	2.7
2003.....	0.2	*	*	*	*	*	*	*	*	*	0.6	3.9
2002.....	0.2	*	*	*	*	*	*	*	*	0.2	0.5	3.5
2001.....	0.2	1.2	*	*	*	*	*	*	*	*	0.5	3.7
2000.....	0.2	*	*	*	*	*	*	*	*	0.2	0.6	4.8
1999.....	0.2	1.5	*	*	*	*	*	*	*	*	0.8	6.8
White, both sexes												
2004.....	0.1	1.4	*	*	*	*	*	*	*	*	0.4	2.8
2003.....	0.2	1.3	*	*	*	*	*	*	*	0.2	0.7	3.7
2002.....	0.1	1.1	0.2	*	*	*	*	*	*	0.2	0.5	3.5
2001.....	0.1	1.0	*	*	*	*	*	*	*	0.1	0.5	3.5
2000.....	0.2	0.8	*	*	*	*	*	*	*	0.3	0.7	4.7
1999.....	0.2	1.2	*	*	*	*	*	*	*	0.2	0.9	6.3
White, male												
2004.....	0.1	1.5	*	*	*	*	*	*	*	*	*	2.6
2003.....	0.1	1.7	*	*	*	*	*	*	*	*	0.7	2.7
2002.....	0.1	1.6	*	*	*	*	*	*	*	*	0.5	2.9
2001.....	0.1	*	*	*	*	*	*	*	*	*	*	2.4
2000.....	0.1	*	*	*	*	*	*	*	*	0.3	0.7	3.8
1999.....	0.1	*	*	*	*	*	*	*	*	*	1.1	4.0

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute bronchitis and bronchiolitis (J20-J21)												
All races 3/, both sexes												
2004.....	0.1	1.4	*	*	*	*	*	*	*	*	0.2	1.4
2003.....	0.1	1.3	0.2	*	*	*	*	*	*	0.1	0.3	2.1
2002.....	0.1	1.2	0.1	*	*	*	*	*	*	0.1	0.3	2.0
2001.....	0.1	1.2	*	*	*	*	*	*	*	0.1	0.3	2.1
2000.....	0.1	1.1	*	*	*	*	*	*	*	0.2	0.4	2.6
1999.....	0.1	1.6	0.2	*	*	*	*	*	*	0.2	0.6	3.6
All races 3/, male												
2004.....	0.1	1.6	*	*	*	*	*	*	*	*	*	1.8
2003.....	0.1	1.7	*	*	*	*	*	*	*	*	*	*
2002.....	0.1	1.8	*	*	*	*	*	*	*	*	*	1.7
2001.....	0.1	1.2	*	*	*	*	*	*	*	*	*	*
2000.....	0.1	1.2	*	*	*	*	*	*	*	*	0.6	2.4
1999.....	0.1	1.9	*	*	*	*	*	*	*	*	0.7	2.6
All races 3/, female												
2004.....	0.1	1.3	*	*	*	*	*	*	*	*	0.3	1.3
2003.....	0.1	*	*	*	*	*	*	*	*	*	0.4	2.5
2002.....	0.1	*	*	*	*	*	*	*	*	*	0.3	2.2
2001.....	0.1	1.2	*	*	*	*	*	*	*	*	0.4	2.4
2000.....	0.1	*	*	*	*	*	*	*	*	*	0.3	2.7
1999.....	0.2	1.4	*	*	*	*	*	*	*	*	0.5	4.0
White, both sexes												
2004.....	0.1	1.3	*	*	*	*	*	*	*	*	0.2	1.5
2003.....	0.1	1.3	*	*	*	*	*	*	*	*	0.4	2.3
2002.....	0.1	1.0	0.2	*	*	*	*	*	*	0.1	0.3	2.1
2001.....	0.1	1.0	*	*	*	*	*	*	*	*	0.3	2.1
2000.....	0.1	0.8	*	*	*	*	*	*	*	0.2	0.4	2.7
1999.....	0.1	1.2	*	*	*	*	*	*	*	0.1	0.6	3.8
White, male												
2004.....	0.1	1.4	*	*	*	*	*	*	*	*	*	1.8
2003.....	0.1	1.6	*	*	*	*	*	*	*	*	*	*
2002.....	0.1	1.5	*	*	*	*	*	*	*	*	*	1.7
2001.....	0.1	*	*	*	*	*	*	*	*	*	*	*
2000.....	0.1	*	*	*	*	*	*	*	*	*	0.6	2.4
1999.....	0.1	*	*	*	*	*	*	*	*	*	0.7	2.6

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other diseases of upper respiratory tract (J30-J39)

All races 3/, both sexes

2004.....	0.1	*	*	*	*	*	0.0	0.1	0.2	0.2	0.8	2.1
2003.....	0.2	*	*	*	*	*	*	0.1	0.2	0.3	0.8	2.3
2002.....	0.1	*	*	*	*	*	*	0.1	0.2	0.3	0.6	2.1
2001.....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.3	0.9	2.4
2000.....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.4	0.9	2.1
1999.....	0.2	*	*	*	*	*	0.0	0.1	0.2	0.4	1.0	2.4

All races 3/, male

2004.....	0.2	*	*	*	*	*	*	0.1	0.2	0.3	1.1	3.1
2003.....	0.2	*	*	*	*	*	*	*	0.2	0.5	1.2	3.0
2002.....	0.1	*	*	*	*	*	*	0.1	0.2	0.3	0.7	2.5
2001.....	0.2	*	*	*	*	*	*	0.1	0.2	0.3	1.2	3.4
2000.....	0.2	*	*	*	*	*	*	0.1	*	0.5	1.1	3.1
1999.....	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.2	3.5

All races 3/, female

2004.....	0.1	*	*	*	*	*	*	0.1	*	0.2	0.7	1.6
2003.....	0.1	*	*	*	*	*	*	*	0.2	0.3	0.6	2.0
2002.....	0.1	*	*	*	*	*	*	*	0.2	0.3	0.6	1.9
2001.....	0.2	*	*	*	*	*	*	0.1	*	0.3	0.8	1.9
2000.....	0.2	*	*	*	*	*	*	0.1	0.2	0.4	0.7	1.7
1999.....	0.2	*	*	*	*	*	*	*	0.2	0.3	0.8	2.0

White, both sexes

2004.....	0.1	*	*	*	*	*	*	0.1	0.1	0.2	0.8	2.1
2003.....	0.2	*	*	*	*	*	*	0.1	0.2	0.3	0.8	2.3
2002.....	0.1	*	*	*	*	*	*	0.1	0.2	0.3	0.6	2.2
2001.....	0.2	*	*	*	*	*	*	0.1	0.1	0.3	0.9	2.3
2000.....	0.2	*	*	*	*	*	*	0.1	0.2	0.4	0.9	2.2
1999.....	0.2	*	*	*	*	*	*	0.1	0.2	0.4	0.9	2.4

White, male

2004.....	0.2	*	*	*	*	*	*	*	0.2	*	1.0	3.2
2003.....	0.2	*	*	*	*	*	*	*	0.2	0.4	1.2	3.2
2002.....	0.1	*	*	*	*	*	*	*	*	0.4	0.7	2.7
2001.....	0.1	*	*	*	*	*	*	*	*	0.3	1.1	3.2
2000.....	0.2	*	*	*	*	*	*	0.1	*	0.5	1.2	3.0
1999.....	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.2	3.2

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Chronic lower respiratory diseases (J40-J47)												
All races 3/, both sexes												
2004.....	41.5	0.9	0.3	0.3	0.4	0.6	2.0	8.4	40.4	153.8	366.7	601.7
2003.....	43.5	0.8	0.3	0.3	0.5	0.7	2.1	8.7	43.3	163.2	383.0	635.1
2002.....	43.3	1.0	0.4	0.3	0.5	0.8	2.2	8.7	42.4	163.0	386.7	637.6
2001.....	43.2	1.0	0.3	0.3	0.4	0.7	2.2	8.5	44.1	167.9	379.8	644.7
2000.....	43.4	0.9	0.3	0.3	0.5	0.7	2.1	8.6	44.2	169.4	386.1	648.6
1999.....	44.5	0.9	0.4	0.3	0.5	0.8	2.0	8.5	47.5	177.2	397.8	646.0
All races 3/, male												
2004.....	40.6	1.2	0.4	0.4	0.5	0.7	2.0	8.8	43.1	172.1	445.6	811.1
2003.....	42.4	1.1	0.5	0.4	0.5	0.8	1.9	9.1	46.5	183.6	464.9	865.9
2002.....	42.9	1.1	0.6	0.4	0.6	0.8	2.2	9.1	45.2	184.8	480.8	894.8
2001.....	42.7	1.1	0.4	0.3	0.5	0.7	2.0	8.8	46.9	191.3	475.1	916.9
2000.....	43.5	1.2	0.4	0.4	0.6	0.8	1.9	9.0	47.8	195.2	488.5	967.9
1999.....	45.6	*	0.4	0.4	0.6	0.8	1.8	8.6	52.3	210.7	513.2	996.7
All races 3/, female												
2004.....	42.5	*	*	0.2	0.3	0.5	2.1	8.1	38.0	138.4	313.6	507.4
2003.....	44.4	*	*	0.2	0.4	0.6	2.3	8.2	40.3	146.0	328.3	533.0
2002.....	43.7	*	0.3	0.3	0.4	0.7	2.3	8.2	39.8	144.9	324.1	526.0
2001.....	43.7	*	*	0.2	0.4	0.7	2.3	8.1	41.5	148.5	317.3	530.8
2000.....	43.2	*	0.3	0.3	0.4	0.7	2.2	8.3	41.0	148.2	319.2	518.5
1999.....	43.4	*	0.3	0.2	0.5	0.9	2.1	8.4	43.1	149.8	322.9	504.6
White, both sexes												
2004.....	47.4	0.7	0.2	0.2	0.3	0.5	1.8	8.2	42.3	164.7	387.0	630.1
2003.....	49.5	0.7	0.2	0.2	0.3	0.5	1.9	8.5	45.2	173.8	403.0	664.5
2002.....	49.2	0.8	0.2	0.2	0.3	0.6	1.9	8.4	44.2	173.2	405.6	666.2
2001.....	49.0	0.7	0.2	0.2	0.3	0.5	1.8	8.3	46.0	177.7	398.6	671.8
2000.....	49.0	0.8	0.2	0.2	0.3	0.5	1.7	8.2	45.8	178.7	404.4	675.6
1999.....	50.2	*	0.3	0.2	0.4	0.5	1.6	8.1	49.3	186.2	414.3	672.9
White, male												
2004.....	45.5	*	*	0.2	0.3	0.5	1.7	8.5	43.8	180.6	462.4	837.7
2003.....	47.4	1.3	*	0.2	0.4	0.5	1.7	8.9	47.6	191.6	478.5	894.4
2002.....	47.8	*	0.3	0.2	0.4	0.6	1.8	8.8	46.0	192.3	495.2	923.4
2001.....	47.6	*	*	0.2	0.3	0.5	1.7	8.6	48.0	198.3	489.4	943.6
2000.....	48.3	*	*	0.2	0.4	0.5	1.6	8.4	48.6	201.4	503.6	997.4
1999.....	50.6	*	*	0.3	0.4	0.5	1.5	8.1	53.1	217.3	525.4	1,029.4

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Chronic lower respiratory diseases (J40-J47)												
White, female												
2004.....	49.3	*	*	0.1	0.2	0.4	1.9	7.9	40.8	151.1	335.5	536.5
2003.....	51.5	*	*	0.1	0.3	0.6	2.1	8.1	42.9	158.6	352.0	562.8
2002.....	50.5	*	*	0.2	0.3	0.6	2.0	8.1	42.4	157.0	345.4	554.5
2001.....	50.3	*	*	0.1	0.3	0.5	1.9	8.0	44.1	160.4	338.3	557.9
2000.....	49.7	*	*	0.2	0.3	0.5	1.8	7.9	43.2	159.6	339.1	544.8
1999.....	49.8	*	*	0.2	0.4	0.6	1.8	8.1	45.6	160.3	341.5	529.7
Black, both sexes												
2004.....	19.2	*	0.8	1.0	1.2	1.8	3.9	12.6	38.1	102.1	222.8	333.9
2003.....	20.2	*	1.0	0.9	1.3	1.9	4.3	12.5	40.5	113.2	235.9	353.0
2002.....	20.7	*	1.2	1.1	1.3	1.9	5.1	12.9	40.7	113.5	248.7	368.2
2001.....	20.4	*	0.9	0.8	1.3	2.4	4.8	12.5	40.7	118.9	234.1	375.8
2000.....	20.8	*	0.8	1.0	1.4	2.1	4.8	14.2	43.9	121.7	234.7	373.8
1999.....	21.9	*	*	1.0	1.3	2.6	4.6	13.8	45.8	128.5	262.6	376.2
Black, male												
2004.....	22.0	*	*	1.3	1.5	2.5	4.0	13.9	50.1	140.4	336.3	566.3
2003.....	23.3	*	1.6	1.2	1.5	2.8	4.0	13.3	50.5	155.1	382.2	601.6
2002.....	24.1	*	1.6	1.2	1.9	2.2	5.7	14.4	52.3	158.0	392.2	645.4
2001.....	23.6	*	*	0.9	1.6	2.7	4.7	13.3	49.8	168.0	380.8	671.7
2000.....	24.3	*	*	1.2	1.7	2.8	4.8	15.0	54.6	176.9	370.3	693.1
1999.....	26.2	*	*	1.4	1.8	2.7	4.7	15.3	59.3	184.6	434.4	701.9
Black, female												
2004.....	16.6	*	*	0.6	1.0	1.2	3.9	11.5	28.5	74.9	158.6	241.8
2003.....	17.3	*	*	0.6	1.1	1.1	4.6	11.9	32.4	83.3	153.2	256.4
2002.....	17.7	*	*	1.0	0.6	1.6	4.6	11.6	31.5	82.0	167.4	262.0
2001.....	17.5	*	*	0.7	1.0	2.1	4.9	11.7	33.3	84.3	151.7	266.1
2000.....	17.6	*	*	0.8	1.0	1.5	4.7	13.4	35.3	82.9	158.4	255.0
1999.....	18.0	*	*	0.6	0.9	2.6	4.6	12.6	34.9	88.9	166.4	254.5

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)												
All races 3/, both sexes												
2004.....	41.2	0.9	0.3	0.3	0.4	0.6	2.0	8.4	40.2	153.0	363.6	595.3
2003.....	43.1	0.8	0.3	0.3	0.5	0.7	2.1	8.6	43.1	162.4	379.9	629.1
2002.....	42.9	1.0	0.4	0.3	0.5	0.7	2.2	8.6	42.1	162.1	383.3	631.8
2001.....	42.9	1.0	0.3	0.3	0.4	0.7	2.1	8.4	43.9	167.0	376.9	637.7
2000.....	43.0	0.9	0.3	0.3	0.5	0.7	2.0	8.6	44.0	168.5	382.8	642.4
1999.....	44.2	0.9	0.3	0.3	0.5	0.8	2.0	8.4	47.2	176.3	394.4	639.8
All races 3/, male												
2004.....	40.4	1.2	0.4	0.4	0.5	0.7	1.9	8.8	42.9	171.6	443.6	806.6
2003.....	42.3	1.1	0.5	0.4	0.5	0.8	1.9	9.1	46.3	183.2	462.7	861.8
2002.....	42.7	1.1	0.6	0.4	0.6	0.8	2.2	9.1	45.0	184.2	478.6	890.1
2001.....	42.5	1.1	0.4	0.3	0.5	0.7	2.0	8.8	46.6	190.6	473.2	912.1
2000.....	43.3	1.2	0.4	0.4	0.6	0.8	1.9	8.9	47.5	194.5	486.4	963.6
1999.....	45.4	*	0.4	0.4	0.6	0.7	1.8	8.6	52.0	210.1	510.1	990.1
All races 3/, female												
2004.....	42.0	*	*	0.2	0.3	0.5	2.0	8.0	37.8	137.4	309.8	500.3
2003.....	44.0	*	*	0.2	0.4	0.6	2.3	8.2	40.0	145.0	324.6	526.2
2002.....	43.2	*	0.3	0.3	0.4	0.7	2.2	8.2	39.4	143.7	320.0	519.7
2001.....	43.2	*	*	0.2	0.4	0.7	2.2	8.1	41.3	147.3	313.6	522.9
2000.....	42.8	*	0.3	0.3	0.4	0.6	2.1	8.2	40.6	147.1	315.2	511.6
1999.....	43.0	*	0.3	0.2	0.5	0.9	2.1	8.2	42.8	148.6	319.3	498.6
White, both sexes												
2004.....	47.0	0.7	0.2	0.2	0.3	0.5	1.8	8.1	42.1	163.8	383.6	623.4
2003.....	49.1	0.7	0.2	0.2	0.3	0.5	1.9	8.4	44.9	173.0	399.8	658.1
2002.....	48.8	0.8	0.2	0.2	0.3	0.6	1.9	8.4	43.9	172.3	402.0	660.2
2001.....	48.6	0.7	0.2	0.2	0.3	0.5	1.8	8.2	45.8	176.8	395.5	664.3
2000.....	48.7	0.8	0.2	0.2	0.3	0.5	1.7	8.1	45.6	177.7	400.9	669.0
1999.....	49.8	*	0.3	0.2	0.4	0.5	1.6	8.0	48.9	185.3	410.8	666.2
White, male												
2004.....	45.3	*	*	0.2	0.3	0.5	1.7	8.5	43.6	180.0	460.3	832.8
2003.....	47.2	1.3	*	0.2	0.3	0.5	1.7	8.8	47.4	191.1	476.2	889.8
2002.....	47.6	*	0.3	0.2	0.4	0.6	1.8	8.7	45.8	191.7	492.9	918.7
2001.....	47.4	*	*	0.2	0.3	0.4	1.7	8.5	47.8	197.7	487.6	938.8
2000.....	48.1	*	*	0.2	0.4	0.5	1.5	8.4	48.4	200.7	501.3	992.8
1999.....	50.3	*	*	0.3	0.4	0.5	1.4	8.1	52.9	216.6	522.3	1,022.5

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)												
White, female												
2004.....	48.7	*	*	0.1	0.2	0.4	1.8	7.8	40.6	150.0	331.3	529.0
2003.....	51.0	*	*	0.1	0.3	0.6	2.0	8.0	42.6	157.5	348.1	555.5
2002.....	49.9	*	*	0.2	0.3	0.6	2.0	8.0	42.1	155.8	341.0	547.9
2001.....	49.8	*	*	0.1	0.3	0.5	1.9	7.9	43.9	159.1	334.3	549.4
2000.....	49.2	*	*	0.2	0.2	0.5	1.8	7.9	42.9	158.5	334.9	537.5
1999.....	49.3	*	*	0.2	0.4	0.6	1.8	8.0	45.2	159.2	337.7	523.1
Black, both sexes												
2004.....	19.1	*	0.8	0.9	1.2	1.8	3.9	12.6	38.0	101.7	222.1	332.5
2003.....	20.1	*	1.0	0.9	1.3	1.9	4.2	12.4	40.2	112.9	235.2	352.7
2002.....	20.7	*	1.2	1.1	1.3	1.9	5.1	12.7	40.6	113.0	247.7	367.0
2001.....	20.3	*	0.9	0.8	1.3	2.3	4.8	12.4	40.5	118.6	233.0	374.6
2000.....	20.7	*	0.8	1.0	1.4	2.1	4.7	14.1	43.6	120.9	234.0	373.2
1999.....	21.8	*	*	1.0	1.3	2.6	4.6	13.7	45.5	127.9	261.1	375.3
Black, male												
2004.....	21.9	*	*	1.3	1.5	2.5	4.0	13.9	49.9	140.2	336.3	564.3
2003.....	23.3	*	1.6	1.2	1.5	2.7	3.9	13.2	50.3	154.7	381.3	601.6
2002.....	24.1	*	1.6	1.2	1.9	2.2	5.7	14.2	52.1	157.1	391.0	645.4
2001.....	23.6	*	*	0.9	1.6	2.7	4.7	13.3	49.6	167.4	378.7	669.4
2000.....	24.3	*	*	1.2	1.7	2.8	4.8	15.0	54.4	175.6	369.4	693.1
1999.....	26.1	*	*	1.4	1.8	2.7	4.7	15.2	58.9	184.3	432.9	700.7
Black, female												
2004.....	16.5	*	*	*	1.0	1.2	3.9	11.5	28.4	74.3	157.5	240.6
2003.....	17.3	*	*	0.6	1.1	1.1	4.5	11.7	32.1	83.2	152.6	256.0
2002.....	17.6	*	*	1.0	0.6	1.6	4.6	11.4	31.4	81.7	166.6	260.4
2001.....	17.4	*	*	0.7	1.0	2.0	4.9	11.6	33.2	84.2	151.2	265.2
2000.....	17.5	*	*	0.8	1.0	1.5	4.7	13.4	34.9	82.5	157.8	254.1
1999.....	17.9	*	*	0.6	0.9	2.5	4.6	12.5	34.8	88.2	164.9	253.6

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Bronchitis, not specified as acute or chronic (J40)

All races 3/, both sexes

2004.....	0.1	*	*	*	*	*	*	0.1	0.1	0.2	0.7	3.8
2003.....	0.2	*	*	*	*	*	*	*	0.1	0.2	0.9	4.6
2002.....	0.2	*	*	*	*	*	*	*	0.1	0.3	1.0	5.2
2001.....	0.2	*	*	*	*	*	*	0.1	0.1	0.3	1.1	5.2
2000.....	0.2	*	*	*	*	*	*	0.1	0.2	0.4	1.5	6.8
1999.....	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.4	6.2

All races 3/, male

2004.....	0.1	*	*	*	*	*	*	*	*	0.4	0.7	3.9
2003.....	0.1	*	*	*	*	*	*	*	*	0.3	0.9	3.9
2002.....	0.1	*	*	*	*	*	*	*	*	0.3	1.0	4.7
2001.....	0.1	*	*	*	*	*	*	*	*	0.4	1.1	5.1
2000.....	0.2	*	*	*	*	*	*	*	0.2	0.4	2.0	6.4
1999.....	0.2	*	*	*	*	*	*	*	0.2	0.5	1.6	5.9

All races 3/, female

2004.....	0.2	*	*	*	*	*	*	*	*	*	0.7	3.7
2003.....	0.2	*	*	*	*	*	*	*	*	0.2	0.8	4.9
2002.....	0.2	*	*	*	*	*	*	*	*	0.3	1.0	5.4
2001.....	0.2	*	*	*	*	*	*	*	*	0.3	1.1	5.3
2000.....	0.3	*	*	*	*	*	*	*	*	0.3	1.1	7.0
1999.....	0.3	*	*	*	*	*	*	*	0.2	0.5	1.3	6.3

White, both sexes

2004.....	0.2	*	*	*	*	*	*	0.1	0.1	0.3	0.7	3.9
2003.....	0.2	*	*	*	*	*	*	*	*	0.3	0.9	4.7
2002.....	0.2	*	*	*	*	*	*	*	0.1	0.3	1.1	5.5
2001.....	0.2	*	*	*	*	*	*	0.1	0.1	0.3	1.1	5.5
2000.....	0.3	*	*	*	*	*	*	*	0.2	0.3	1.5	7.3
1999.....	0.2	*	*	*	*	*	*	*	0.2	0.5	1.4	6.4

White, male

2004.....	0.1	*	*	*	*	*	*	*	*	0.4	0.7	4.1
2003.....	0.1	*	*	*	*	*	*	*	*	0.3	1.0	4.0
2002.....	0.1	*	*	*	*	*	*	*	*	0.3	1.0	5.1
2001.....	0.2	*	*	*	*	*	*	*	*	0.3	1.1	5.3
2000.....	0.2	*	*	*	*	*	*	*	*	0.4	2.1	6.9
1999.....	0.2	*	*	*	*	*	*	*	*	0.5	1.5	6.3

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Chronic bronchitis (J41-J42)												
All races 3/, both sexes												
2004.....	0.1	*	*	*	*	*	*	*	0.1	0.3	0.8	2.8
2003.....	0.1	*	*	*	*	*	*	0.1	0.1	0.3	1.1	2.7
2002.....	0.2	*	*	*	*	*	*	0.1	0.1	0.4	1.2	2.9
2001.....	0.2	*	*	*	*	*	*	*	0.1	0.4	1.0	3.9
2000.....	0.2	*	*	*	*	*	*	*	0.1	0.5	1.2	5.0
1999.....	0.2	*	*	*	*	*	*	*	0.2	0.4	1.3	4.5
All races 3/, male												
2004.....	0.1	*	*	*	*	*	*	*	0.1	0.3	0.9	3.1
2003.....	0.1	*	*	*	*	*	*	*	*	0.3	1.3	3.0
2002.....	0.2	*	*	*	*	*	*	*	*	0.6	1.5	4.0
2001.....	0.1	*	*	*	*	*	*	*	*	0.6	1.2	4.1
2000.....	0.2	*	*	*	*	*	*	*	*	0.6	1.6	6.4
1999.....	0.2	*	*	*	*	*	*	*	0.3	0.6	1.7	5.9
All races 3/, female												
2004.....	0.1	*	*	*	*	*	*	*	*	0.3	0.8	2.6
2003.....	0.1	*	*	*	*	*	*	*	*	0.4	0.9	2.6
2002.....	0.1	*	*	*	*	*	*	*	*	0.3	1.0	2.4
2001.....	0.2	*	*	*	*	*	*	*	*	0.3	0.8	3.8
2000.....	0.2	*	*	*	*	*	*	*	*	0.4	1.0	4.4
1999.....	0.2	*	*	*	*	*	*	*	0.2	0.3	1.1	4.0
White, both sexes												
2004.....	0.1	*	*	*	*	*	*	*	0.1	0.3	0.9	2.8
2003.....	0.1	*	*	*	*	*	*	*	*	0.3	1.1	2.8
2002.....	0.2	*	*	*	*	*	*	*	0.1	0.5	1.3	3.1
2001.....	0.2	*	*	*	*	*	*	*	0.2	0.4	1.0	4.1
2000.....	0.2	*	*	*	*	*	*	*	0.1	0.5	1.2	5.3
1999.....	0.2	*	*	*	*	*	*	*	0.3	0.5	1.4	4.8
White, male												
2004.....	0.1	*	*	*	*	*	*	*	*	0.3	1.0	3.0
2003.....	0.1	*	*	*	*	*	*	*	*	0.3	1.2	3.3
2002.....	0.2	*	*	*	*	*	*	*	*	0.6	1.6	4.2
2001.....	0.2	*	*	*	*	*	*	*	*	0.6	1.2	4.2
2000.....	0.2	*	*	*	*	*	*	*	*	0.6	1.6	6.9
1999.....	0.2	*	*	*	*	*	*	*	0.3	0.6	1.7	6.1

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Emphysema (J43)												
All races 3/, both sexes												
2004.....	4.6	*	*	*	*	*	0.2	1.0	5.4	20.5	41.0	49.9
2003.....	5.1	*	*	*	*	*	0.2	1.1	6.3	22.2	46.0	54.6
2002.....	5.4	*	*	*	*	*	0.2	1.1	6.4	24.0	48.0	59.3
2001.....	5.7	*	*	*	*	*	0.2	1.1	7.2	26.2	50.4	62.1
2000.....	5.9	*	*	*	*	*	0.2	1.3	7.4	27.8	52.7	63.3
1999.....	6.4	*	*	*	*	*	0.2	1.3	8.5	30.7	56.9	62.1
All races 3/, male												
2004.....	4.9	*	*	*	*	*	0.2	1.2	6.2	24.3	52.2	74.4
2003.....	5.3	*	*	*	*	*	0.2	1.4	6.9	25.6	58.1	79.2
2002.....	5.7	*	*	*	*	*	0.3	1.4	7.2	28.1	63.5	90.0
2001.....	6.0	*	*	*	*	*	0.2	1.4	7.9	30.8	65.3	101.2
2000.....	6.3	*	*	*	*	*	0.2	1.5	8.1	33.0	69.1	104.2
1999.....	6.9	*	*	*	*	*	0.2	1.5	9.5	37.8	76.4	105.4
All races 3/, female												
2004.....	4.4	*	*	*	*	*	0.1	0.8	4.7	17.4	33.5	38.9
2003.....	4.9	*	*	*	*	*	0.2	0.8	5.7	19.2	37.9	43.7
2002.....	5.0	*	*	*	*	*	0.1	0.9	5.6	20.6	37.6	46.0
2001.....	5.4	*	*	*	*	*	0.1	0.9	6.6	22.5	40.6	45.8
2000.....	5.6	*	*	*	*	*	0.2	1.0	6.8	23.6	41.9	46.7
1999.....	5.8	*	*	*	*	*	0.2	1.1	7.5	24.9	44.2	44.7
White, both sexes												
2004.....	5.4	*	*	*	*	*	0.2	1.0	5.8	22.3	43.7	52.5
2003.....	5.9	*	*	*	*	*	0.2	1.1	6.7	23.9	48.5	57.0
2002.....	6.2	*	*	*	*	*	0.2	1.2	6.9	25.7	50.7	62.3
2001.....	6.6	*	*	*	*	*	0.2	1.2	7.7	28.2	53.5	65.1
2000.....	6.8	*	*	*	*	*	0.2	1.2	7.9	29.7	55.6	65.8
1999.....	7.3	*	*	*	*	*	0.2	1.3	9.1	32.9	59.8	64.7
White, male												
2004.....	5.5	*	*	*	*	*	0.2	1.1	6.4	25.8	54.5	77.1
2003.....	5.9	*	*	*	*	*	0.2	1.4	7.2	27.1	59.8	80.4
2002.....	6.4	*	*	*	*	*	0.3	1.3	7.5	29.3	65.8	93.0
2001.....	6.8	*	*	*	*	*	0.2	1.4	8.2	32.3	68.1	105.1
2000.....	7.0	*	*	*	*	*	0.2	1.4	8.4	34.3	72.1	105.5
1999.....	7.8	*	*	*	*	*	0.2	1.4	9.9	39.4	79.0	109.2

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Emphysema (J43)												
White, female												
2004.....	5.2	*	*	*	*	*	0.1	0.8	5.2	19.3	36.3	41.4
2003.....	5.8	*	*	*	*	*	0.2	0.9	6.2	21.2	40.8	46.6
2002.....	5.9	*	*	*	*	*	*	1.0	6.2	22.7	40.5	48.9
2001.....	6.3	*	*	*	*	*	0.1	0.9	7.2	24.7	43.7	48.4
2000.....	6.5	*	*	*	*	*	0.2	1.0	7.4	25.9	44.7	49.7
1999.....	6.8	*	*	*	*	*	0.2	1.1	8.3	27.4	47.2	46.9
Black, both sexes												
2004.....	1.9	*	*	*	*	*	*	1.3	4.5	10.7	22.1	27.6
2003.....	2.1	*	*	*	*	*	*	1.2	5.1	12.9	27.3	30.9
2002.....	2.3	*	*	*	*	*	0.4	1.2	4.4	15.9	28.0	32.1
2001.....	2.2	*	*	*	*	*	*	1.3	5.6	15.3	23.6	35.3
2000.....	2.5	*	*	*	*	*	*	2.0	6.0	17.0	26.9	37.2
1999.....	2.6	*	*	*	*	*	0.4	1.8	6.1	17.1	31.3	37.6
Black, male												
2004.....	2.5	*	*	*	*	*	*	2.0	6.9	17.0	36.5	56.6
2003.....	2.7	*	*	*	*	*	*	1.7	7.3	18.2	46.0	70.9
2002.....	3.0	*	*	*	*	*	*	1.8	6.5	24.8	48.8	64.3
2001.....	2.7	*	*	*	*	*	*	1.6	8.0	22.6	39.7	68.1
2000.....	3.2	*	*	*	*	*	*	2.9	8.1	27.4	41.5	87.6
1999.....	3.5	*	*	*	*	*	*	2.9	8.9	29.0	54.7	72.6
Black, female												
2004.....	1.3	*	*	*	*	*	*	*	2.6	6.2	14.0	16.1
2003.....	1.6	*	*	*	*	*	*	*	3.4	9.2	16.7	15.4
2002.....	1.6	*	*	*	*	*	*	*	2.7	9.5	16.2	19.7
2001.....	1.7	*	*	*	*	*	*	1.1	3.6	10.2	14.5	23.2
2000.....	1.8	*	*	*	*	*	*	1.3	4.4	9.7	18.8	18.5
1999.....	1.7	*	*	*	*	*	*	1.0	3.9	8.7	18.2	24.5

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other chronic obstructive pulmonary disease (J44)												
All races 3/, both sexes												
2004.....	35.0	*	*	*	*	0.1	0.9	5.9	32.8	129.2	315.8	524.8
2003.....	36.3	*	*	*	*	0.1	0.9	5.9	34.6	136.7	326.2	552.9
2002.....	35.8	*	*	*	*	0.1	0.9	5.9	33.5	134.2	326.7	549.7
2001.....	35.3	*	*	*	0.1	0.1	0.9	5.6	34.2	136.5	318.0	550.3
2000.....	35.1	*	*	*	*	0.1	0.8	5.6	34.0	135.8	320.6	550.6
1999.....	35.7	*	*	*	*	0.1	0.7	5.4	35.9	140.5	327.2	549.6
All races 3/, male												
2004.....	34.3	*	*	*	*	0.1	1.0	6.4	35.1	144.5	386.3	714.1
2003.....	35.7	*	*	*	*	0.1	0.9	6.4	37.6	154.9	397.8	764.9
2002.....	35.5	*	*	*	*	0.1	1.0	6.4	36.0	152.8	407.6	780.4
2001.....	35.1	*	*	*	*	0.1	0.9	6.2	36.9	156.3	401.1	789.6
2000.....	35.4	*	*	*	*	*	0.8	6.0	37.5	157.5	408.7	833.7
1999.....	36.9	*	*	*	*	*	0.7	5.9	40.3	167.9	424.8	860.0
All races 3/, female												
2004.....	35.7	*	*	*	*	*	0.9	5.5	30.7	116.3	268.4	439.6
2003.....	36.9	*	*	*	*	*	1.0	5.4	31.8	121.6	278.3	459.2
2002.....	36.0	*	*	*	*	0.1	0.9	5.4	31.2	118.7	273.0	449.5
2001.....	35.5	*	*	*	*	*	0.9	5.1	31.7	120.2	263.4	450.2
2000.....	34.7	*	*	*	*	*	0.8	5.2	30.8	117.9	263.1	435.3
1999.....	34.5	*	*	*	*	0.1	0.7	4.9	31.8	118.0	263.8	424.5
White, both sexes												
2004.....	40.3	*	*	*	*	0.1	1.0	6.1	34.7	138.5	333.6	550.5
2003.....	41.7	*	*	*	*	0.1	1.0	6.1	36.5	146.1	343.9	579.4
2002.....	40.9	*	*	*	*	0.1	0.9	6.1	35.3	143.1	342.8	574.7
2001.....	40.4	*	*	*	*	0.1	0.9	5.8	36.2	144.9	333.9	573.9
2000.....	40.0	*	*	*	*	*	0.8	5.7	35.7	143.8	336.0	574.7
1999.....	40.6	*	*	*	*	0.1	0.7	5.5	37.5	147.9	341.0	573.7
White, male												
2004.....	38.8	*	*	*	*	0.1	1.0	6.6	36.1	151.8	401.2	738.3
2003.....	40.2	*	*	*	*	*	0.9	6.5	38.8	161.8	410.2	791.5
2002.....	40.0	*	*	*	*	*	1.0	6.4	36.9	159.6	419.8	805.8
2001.....	39.5	*	*	*	*	*	0.9	6.2	38.2	162.3	413.3	812.9
2000.....	39.7	*	*	*	*	*	0.8	5.9	38.5	162.9	420.8	862.1
1999.....	41.2	*	*	*	*	*	0.7	5.8	41.2	173.5	434.9	890.0

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other chronic obstructive pulmonary disease (J44)

White, female

2004.....	41.7	*	*	*	*	*	0.9	5.7	33.4	127.2	287.5	465.8
2003.....	43.2	*	*	*	*	*	1.0	5.7	34.3	132.7	299.1	485.4
2002.....	41.9	*	*	*	*	*	0.9	5.7	33.7	129.1	291.1	474.3
2001.....	41.3	*	*	*	*	*	0.9	5.3	34.3	130.2	281.3	473.8
2000.....	40.4	*	*	*	*	*	0.8	5.4	33.0	127.7	280.2	457.9
1999.....	39.9	*	*	*	*	*	0.7	5.1	34.1	126.5	279.4	446.6

Black, both sexes

2004.....	14.5	*	*	*	*	*	1.1	6.6	27.5	85.1	190.0	284.5
2003.....	15.1	*	*	*	*	*	1.2	6.2	29.7	92.6	197.6	304.3
2002.....	15.3	*	*	*	*	*	1.2	6.6	29.6	88.9	210.4	318.9
2001.....	15.0	*	*	*	*	*	1.3	6.4	28.0	94.7	197.7	319.7
2000.....	14.8	*	*	*	*	*	1.2	6.6	30.5	93.7	194.6	310.0
1999.....	15.8	*	*	*	*	*	1.2	6.3	32.3	100.3	216.5	312.8

Black, male

2004.....	16.8	*	*	*	*	*	1.2	7.6	37.6	117.2	290.6	490.8
2003.....	17.9	*	*	*	*	*	1.2	7.3	38.6	130.9	324.7	520.3
2002.....	18.1	*	*	*	*	*	1.4	8.2	39.7	124.5	333.9	565.0
2001.....	18.0	*	*	*	*	*	1.4	7.4	36.8	136.8	328.4	580.9
2000.....	18.1	*	*	*	*	*	1.3	8.3	39.9	139.4	319.4	576.6
1999.....	19.6	*	*	*	*	*	1.4	8.1	44.2	145.2	364.9	605.8

Black, female

2004.....	12.3	*	*	*	*	*	1.0	5.7	19.4	62.1	133.1	202.8
2003.....	12.5	*	*	*	*	*	1.1	5.2	22.5	65.4	125.8	220.4
2002.....	12.8	*	*	*	*	*	1.1	5.2	21.4	63.7	140.5	224.7
2001.....	12.2	*	*	*	*	*	1.2	5.5	20.9	65.0	124.3	222.7
2000.....	11.9	*	*	*	*	*	1.2	5.1	23.0	61.6	124.3	210.8
1999.....	12.4	*	*	*	*	*	1.0	4.8	22.7	68.7	133.4	203.3

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Asthma (J45-J46)												
All races 3/, both sexes												
2004.....	1.3	*	0.2	0.3	0.4	0.5	0.9	1.4	1.8	2.7	5.2	14.1
2003.....	1.4	*	0.2	0.3	0.4	0.6	0.9	1.5	2.0	2.9	5.8	14.2
2002.....	1.5	*	0.3	0.3	0.4	0.6	1.1	1.5	2.0	3.2	6.4	14.7
2001.....	1.5	*	0.2	0.2	0.4	0.6	1.0	1.5	2.2	3.5	6.4	16.1
2000.....	1.6	*	0.2	0.3	0.4	0.6	1.0	1.6	2.2	4.0	6.9	16.7
1999.....	1.7	*	0.2	0.3	0.5	0.7	1.0	1.7	2.5	4.2	7.6	17.3
All races 3/, male												
2004.....	1.0	*	*	0.3	0.5	0.6	0.7	1.0	1.3	2.1	3.5	11.1
2003.....	1.0	*	0.4	0.3	0.4	0.6	0.7	1.2	1.6	2.1	4.5	10.7
2002.....	1.1	*	0.4	0.3	0.5	0.6	0.9	1.2	1.6	2.4	5.0	10.9
2001.....	1.1	*	0.3	0.3	0.4	0.6	0.9	1.1	1.5	2.7	4.5	12.1
2000.....	1.2	*	0.3	0.4	0.5	0.7	0.9	1.3	1.7	3.0	5.0	12.9
1999.....	1.2	*	*	0.4	0.5	0.6	0.8	1.1	1.8	3.2	5.7	12.7
All races 3/, female												
2004.....	1.6	*	*	0.2	0.3	0.4	1.0	1.6	2.2	3.2	6.3	15.4
2003.....	1.8	*	*	0.2	0.3	0.5	1.1	1.9	2.4	3.6	6.7	15.8
2002.....	1.8	*	*	0.3	0.3	0.5	1.2	1.8	2.4	3.9	7.3	16.3
2001.....	1.9	*	*	0.2	0.3	0.6	1.1	1.9	2.8	4.1	7.6	17.8
2000.....	2.0	*	*	0.3	0.3	0.5	1.2	2.0	2.7	4.8	8.1	18.2
1999.....	2.1	*	*	0.2	0.4	0.7	1.2	2.2	3.0	5.0	8.8	19.2
White, both sexes												
2004.....	1.1	*	*	0.1	0.2	0.3	0.6	0.9	1.4	2.4	4.7	13.7
2003.....	1.2	*	*	0.2	0.2	0.4	0.7	1.1	1.6	2.4	5.4	14.3
2002.....	1.3	*	*	0.1	0.3	0.5	0.7	1.1	1.5	2.7	6.1	14.7
2001.....	1.3	*	*	0.1	0.2	0.4	0.7	1.1	1.6	3.0	6.0	15.7
2000.....	1.4	*	*	0.2	0.3	0.4	0.7	1.1	1.7	3.4	6.5	15.9
1999.....	1.5	*	*	0.2	0.3	0.4	0.7	1.2	1.9	3.5	7.2	16.6
White, male												
2004.....	0.7	*	*	0.1	0.3	0.3	0.5	0.7	0.9	1.8	2.8	10.3
2003.....	0.8	*	*	0.2	0.3	0.4	0.5	0.8	1.2	1.7	4.0	10.6
2002.....	0.9	*	*	0.2	0.3	0.5	0.5	0.8	1.2	1.9	4.6	10.7
2001.....	0.8	*	*	0.2	0.2	0.3	0.6	0.8	1.1	2.2	4.0	11.3
2000.....	0.9	*	*	0.2	0.3	0.4	0.6	1.0	1.2	2.5	4.7	11.4
1999.....	0.9	*	*	0.2	0.4	0.4	0.5	0.8	1.4	2.5	5.3	10.8

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Asthma (J45-J46)												
White, female												
2004.....	1.5	*	*	0.1	0.2	0.3	0.8	1.2	1.8	3.0	6.0	15.2
2003.....	1.6	*	*	0.1	0.2	0.5	0.8	1.4	2.0	3.1	6.3	15.9
2002.....	1.7	*	*	*	0.2	0.5	0.9	1.3	1.9	3.3	7.2	16.4
2001.....	1.7	*	*	*	0.2	0.4	0.8	1.5	2.1	3.7	7.3	17.6
2000.....	1.8	*	*	0.2	0.2	0.4	0.8	1.3	2.1	4.1	7.7	17.7
1999.....	2.0	*	*	0.1	0.3	0.4	0.9	1.7	2.4	4.4	8.5	18.9
Black, both sexes												
2004.....	2.6	*	*	0.8	1.1	1.6	2.5	4.5	5.6	5.7	9.1	15.5
2003.....	2.7	*	0.9	0.9	1.2	1.6	2.7	4.8	5.1	7.0	9.1	12.0
2002.....	2.9	*	0.9	1.0	1.2	1.6	3.4	4.8	6.4	7.8	8.0	11.9
2001.....	3.0	*	0.8	0.8	1.1	2.1	3.3	4.6	6.8	7.7	10.6	16.3
2000.....	3.2	*	*	1.0	1.3	1.9	3.1	5.4	6.9	9.2	10.4	22.2
1999.....	3.2	*	*	1.0	1.2	2.4	2.9	5.2	6.8	9.6	10.4	20.4
Black, male												
2004.....	2.5	*	*	1.1	1.4	2.3	2.3	3.9	5.2	5.5	8.6	*
2003.....	2.4	*	1.5	1.2	1.4	2.4	2.3	3.9	4.2	5.3	8.7	*
2002.....	2.8	*	*	1.1	1.8	2.0	3.6	3.9	5.7	7.3	6.9	*
2001.....	2.6	*	*	0.9	1.4	2.4	3.0	4.1	4.6	6.8	8.8	*
2000.....	2.8	*	*	1.2	1.5	2.5	2.9	3.8	6.1	7.9	6.4	25.4
1999.....	2.8	*	*	1.4	1.6	2.6	2.8	3.8	5.4	9.1	8.3	*
Black, female												
2004.....	2.7	*	*	*	0.9	1.1	2.7	5.0	6.0	5.7	9.4	16.9
2003.....	2.9	*	*	*	1.0	0.9	3.1	5.6	5.8	8.2	9.2	13.8
2002.....	3.0	*	*	1.0	*	1.2	3.3	5.6	7.0	8.2	8.6	12.3
2001.....	3.3	*	*	0.7	0.9	1.9	3.5	5.0	8.5	8.3	11.6	16.8
2000.....	3.5	*	*	0.8	1.0	1.4	3.3	6.8	7.5	10.2	12.6	21.0
1999.....	3.5	*	*	0.6	0.9	2.1	3.1	6.4	7.9	10.0	11.5	21.0

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Bronchiectasis (J47)												
All races 3/, both sexes												
2004.....	0.3	*	*	*	*	*	*	0.1	0.2	0.8	3.1	6.3
2003.....	0.3	*	*	*	*	*	*	0.1	0.2	0.8	3.1	6.0
2002.....	0.3	*	*	*	*	*	*	0.1	0.3	0.9	3.4	5.8
2001.....	0.3	*	*	*	*	*	*	0.1	0.3	1.0	3.0	7.0
2000.....	0.3	*	*	*	*	*	*	*	0.3	0.9	3.3	6.1
1999.....	0.3	*	*	*	*	*	*	0.1	0.3	0.9	3.4	6.2
All races 3/, male												
2004.....	0.2	*	*	*	*	*	*	*	0.2	0.5	2.1	4.5
2003.....	0.2	*	*	*	*	*	*	*	0.2	0.5	2.3	4.2
2002.....	0.2	*	*	*	*	*	*	*	0.2	0.6	2.2	4.7
2001.....	0.2	*	*	*	*	*	*	*	0.3	0.7	1.8	4.8
2000.....	0.2	*	*	*	*	*	*	*	0.2	0.7	2.2	4.3
1999.....	0.2	*	*	*	*	*	*	*	0.2	0.7	3.1	6.6
All races 3/, female												
2004.....	0.5	*	*	*	*	*	*	*	0.2	1.0	3.8	7.2
2003.....	0.5	*	*	*	*	*	*	*	0.3	1.1	3.6	6.8
2002.....	0.5	*	*	*	*	*	*	*	0.3	1.1	4.1	6.3
2001.....	0.5	*	*	*	*	*	*	*	0.3	1.2	3.7	7.9
2000.....	0.5	*	*	*	*	*	*	*	0.3	1.1	4.0	6.9
1999.....	0.5	*	*	*	*	*	*	0.1	0.4	1.2	3.6	6.0
White, both sexes												
2004.....	0.4	*	*	*	*	*	*	0.1	0.2	0.8	3.3	6.7
2003.....	0.4	*	*	*	*	*	*	*	0.2	0.8	3.2	6.4
2002.....	0.4	*	*	*	*	*	*	*	0.3	0.9	3.6	6.1
2001.....	0.4	*	*	*	*	*	*	0.1	0.3	1.0	3.1	7.4
2000.....	0.4	*	*	*	*	*	*	*	0.3	0.9	3.4	6.6
1999.....	0.4	*	*	*	*	*	*	0.1	0.3	1.0	3.5	6.7
White, male												
2004.....	0.2	*	*	*	*	*	*	*	0.2	0.6	2.2	4.8
2003.....	0.2	*	*	*	*	*	*	*	*	0.5	2.3	4.5
2002.....	0.2	*	*	*	*	*	*	*	0.2	0.6	2.3	4.7
2001.....	0.2	*	*	*	*	*	*	*	0.3	0.6	1.7	4.8
2000.....	0.2	*	*	*	*	*	*	*	0.2	0.7	2.3	4.6
1999.....	0.3	*	*	*	*	*	*	*	0.2	0.7	3.1	6.9

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Lung diseases due to external agents (J60-J70)												
All races 3/, both sexes												
2004.....	6.1	*	*	*	0.1	0.2	0.4	1.0	2.8	10.9	46.6	171.1
2003.....	6.4	*	*	*	0.1	0.2	0.4	1.0	3.1	11.0	49.0	182.8
2002.....	6.5	*	*	*	0.1	0.2	0.4	0.9	2.9	11.2	50.3	192.9
2001.....	6.5	*	*	*	0.1	0.2	0.4	1.0	2.9	11.6	50.1	195.9
2000.....	6.3	*	*	*	0.1	0.2	0.4	1.0	2.7	12.0	48.6	195.3
1999.....	5.9	*	*	*	0.1	0.2	0.4	0.8	2.8	11.1	45.5	181.8
All races 3/, male												
2004.....	6.7	*	*	*	0.1	0.2	0.5	1.3	3.8	15.5	67.1	256.1
2003.....	6.9	*	*	*	0.2	0.2	0.4	1.3	4.0	15.6	71.2	266.2
2002.....	7.0	*	*	*	0.1	0.2	0.5	1.2	3.7	15.8	73.1	290.1
2001.....	7.0	*	*	*	0.1	0.2	0.6	1.3	3.7	16.6	73.3	291.3
2000.....	6.9	*	*	*	0.1	0.2	0.5	1.3	3.5	17.7	72.1	301.5
1999.....	6.5	*	*	*	0.1	0.2	0.5	1.0	3.8	16.4	70.2	279.1
All races 3/, female												
2004.....	5.5	*	*	*	*	0.1	0.3	0.7	2.0	7.0	32.7	132.9
2003.....	5.9	*	*	*	*	0.1	0.4	0.8	2.2	7.0	34.2	145.9
2002.....	6.0	*	*	*	*	0.2	0.3	0.6	2.1	7.4	35.2	150.7
2001.....	6.1	*	*	*	*	0.2	0.3	0.8	2.1	7.4	34.9	156.0
2000.....	5.8	*	*	*	*	0.1	0.3	0.8	2.0	7.2	33.3	152.1
1999.....	5.3	*	*	*	*	0.2	0.3	0.7	1.8	6.8	29.5	142.6
White, both sexes												
2004.....	6.7	*	*	*	0.1	0.2	0.4	1.0	2.7	10.8	47.1	174.3
2003.....	7.0	*	*	*	0.1	0.2	0.4	1.0	2.9	10.8	48.9	185.7
2002.....	7.2	*	*	*	0.1	0.2	0.4	0.9	2.8	11.0	50.7	196.8
2001.....	7.1	*	*	*	0.1	0.2	0.4	0.9	2.8	11.4	50.4	197.6
2000.....	7.0	*	*	*	0.1	0.2	0.4	0.9	2.6	11.6	48.7	199.2
1999.....	6.4	*	*	*	0.1	0.2	0.3	0.8	2.6	10.7	45.4	184.4
White, male												
2004.....	7.3	*	*	*	*	0.2	0.5	1.3	3.5	15.2	67.9	261.3
2003.....	7.5	*	*	*	0.1	0.2	0.4	1.3	3.8	15.5	71.0	272.3
2002.....	7.8	*	*	*	0.1	0.2	0.5	1.2	3.6	15.5	73.6	297.5
2001.....	7.7	*	*	*	*	0.2	0.5	1.2	3.6	16.5	74.1	297.5
2000.....	7.6	*	*	*	*	0.2	0.5	1.2	3.4	16.9	72.4	311.0
1999.....	7.0	*	*	*	*	0.2	0.4	0.9	3.5	15.7	69.7	285.0

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Lung diseases due to external agents (J60-J70)												
White, female												
2004.....	6.2	*	*	*	*	0.1	0.3	0.7	1.9	7.0	33.0	135.0
2003.....	6.5	*	*	*	*	0.1	0.3	0.8	2.0	6.9	34.0	147.3
2002.....	6.7	*	*	*	*	0.2	0.3	0.6	2.0	7.2	35.3	153.0
2001.....	6.6	*	*	*	*	0.2	0.3	0.7	2.1	7.1	34.7	155.8
2000.....	6.4	*	*	*	*	0.2	0.3	0.7	1.9	7.1	33.1	153.8
1999.....	5.9	*	*	*	*	0.2	0.3	0.6	1.8	6.6	29.4	144.0
Black, both sexes												
2004.....	4.0	*	*	*	*	*	0.6	1.4	5.0	14.8	48.7	149.7
2003.....	4.4	*	*	*	*	*	0.7	1.3	5.3	14.4	56.9	166.0
2002.....	4.2	*	*	*	*	*	0.7	1.5	4.0	15.5	54.5	169.3
2001.....	4.5	*	*	*	*	*	0.6	1.7	4.3	15.9	55.5	189.2
2000.....	4.3	*	*	*	*	*	0.7	1.9	4.3	18.6	54.2	163.6
1999.....	4.4	*	*	*	*	*	0.8	1.6	4.8	17.2	54.4	172.2
Black, male												
2004.....	4.5	*	*	*	*	*	0.7	1.8	7.5	23.1	72.1	220.5
2003.....	4.6	*	*	*	*	*	*	1.7	7.1	21.1	84.6	221.0
2002.....	4.5	*	*	*	*	*	0.9	2.1	5.3	23.1	79.5	247.7
2001.....	4.5	*	*	*	*	*	0.8	2.0	6.0	22.2	81.5	257.6
2000.....	4.7	*	*	*	*	*	0.9	2.8	5.2	30.3	81.5	223.7
1999.....	4.9	*	*	*	*	*	0.9	2.0	7.4	27.2	86.1	246.1
Black, female												
2004.....	3.6	*	*	*	*	*	*	1.0	3.0	8.8	35.4	121.7
2003.....	4.2	*	*	*	*	*	0.8	1.0	3.8	9.6	41.3	144.6
2002.....	4.0	*	*	*	*	*	*	0.9	3.0	10.0	40.4	139.2
2001.....	4.4	*	*	*	*	*	*	1.5	2.9	11.4	40.8	163.8
2000.....	4.0	*	*	*	*	*	*	1.2	3.5	10.4	38.9	141.2
1999.....	3.9	*	*	*	*	*	*	1.2	2.8	10.1	36.7	144.6

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pneumoconioses and chemical effects (J60-J66,J68)												
All races 3/, both sexes												
2004.....	0.4	*	*	*	*	*	*	*	0.2	1.1	3.4	6.0
2003.....	0.4	*	*	*	*	*	*	*	0.3	1.3	3.8	6.4
2002.....	0.4	*	*	*	*	*	*	*	0.3	1.2	3.8	6.8
2001.....	0.4	*	*	*	*	*	*	*	0.3	1.5	4.1	6.2
2000.....	0.4	*	*	*	*	*	*	*	0.2	1.5	3.9	7.5
1999.....	0.4	*	*	*	*	*	*	*	0.2	1.5	4.1	6.1
All races 3/, male												
2004.....	0.7	*	*	*	*	*	*	*	0.5	2.4	8.1	18.4
2003.....	0.7	*	*	*	*	*	*	*	0.5	2.7	9.1	19.8
2002.....	0.7	*	*	*	*	*	*	*	0.5	2.6	9.0	21.6
2001.....	0.8	*	*	*	*	*	*	*	0.5	3.1	9.9	20.2
2000.....	0.8	*	*	*	*	*	*	*	0.5	3.2	9.6	24.9
1999.....	0.8	*	*	*	*	*	*	*	0.5	3.2	10.1	20.4
All races 3/, female												
2004.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2003.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*	*	*	*	0.3	*
2001.....	0.0	*	*	*	*	*	*	*	*	*	0.3	*
2000.....	0.0	*	*	*	*	*	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	*	*	*	*	*	*
White, both sexes												
2004.....	0.4	*	*	*	*	*	*	*	0.3	1.2	3.7	6.2
2003.....	0.4	*	*	*	*	*	*	*	0.3	1.4	4.0	6.9
2002.....	0.4	*	*	*	*	*	*	*	0.3	1.3	4.0	7.2
2001.....	0.5	*	*	*	*	*	*	*	0.3	1.6	4.4	6.5
2000.....	0.5	*	*	*	*	*	*	*	0.3	1.7	4.1	7.9
1999.....	0.5	*	*	*	*	*	*	*	0.2	1.6	4.4	6.3
White, male												
2004.....	0.8	*	*	*	*	*	*	*	0.5	2.6	8.7	19.0
2003.....	0.9	*	*	*	*	*	*	*	0.6	2.9	9.6	21.3
2002.....	0.9	*	*	*	*	*	*	*	0.6	2.8	9.4	22.8
2001.....	0.9	*	*	*	*	*	*	*	0.5	3.4	10.5	21.2
2000.....	0.9	*	*	*	*	*	*	*	0.5	3.4	10.0	26.4
1999.....	0.9	*	*	*	*	*	*	*	0.5	3.4	10.7	21.0

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Pneumonitis due to solids and liquids (J69)

All races 3/, both sexes

2004.....	5.7	*	*	*	0.1	0.2	0.4	0.9	2.6	9.6	42.9	164.9
2003.....	6.0	*	*	*	0.1	0.2	0.4	1.0	2.8	9.6	45.0	176.2
2002.....	6.1	*	*	*	0.1	0.2	0.4	0.9	2.5	9.8	46.3	186.0
2001.....	6.1	*	*	*	0.1	0.2	0.4	1.0	2.6	10.0	45.8	189.4
2000.....	5.9	*	*	*	0.1	0.2	0.4	1.0	2.5	10.3	44.5	187.6
1999.....	5.5	*	*	*	0.1	0.2	0.4	0.8	2.5	9.5	41.1	175.6

All races 3/, male

2004.....	5.9	*	*	*	0.1	0.2	0.4	1.2	3.2	12.9	58.6	237.2
2003.....	6.1	*	*	*	0.1	0.2	0.4	1.2	3.5	12.8	61.7	246.2
2002.....	6.3	*	*	*	0.1	0.2	0.5	1.1	3.1	13.0	63.8	268.5
2001.....	6.1	*	*	*	0.1	0.2	0.5	1.2	3.2	13.3	63.1	270.3
2000.....	6.1	*	*	*	0.1	0.2	0.5	1.2	3.0	14.4	62.3	276.0
1999.....	5.7	*	*	*	0.1	0.2	0.4	0.9	3.2	13.1	59.7	258.4

All races 3/, female

2004.....	5.5	*	*	*	*	0.1	0.3	0.7	1.9	6.8	32.3	132.4
2003.....	5.8	*	*	*	*	0.1	0.3	0.7	2.1	6.9	33.8	145.3
2002.....	5.9	*	*	*	*	0.2	0.3	0.6	2.0	7.1	34.7	150.2
2001.....	6.0	*	*	*	*	0.2	0.3	0.7	2.1	7.2	34.5	155.6
2000.....	5.8	*	*	*	*	0.1	0.3	0.7	1.9	7.0	33.0	151.6
1999.....	5.3	*	*	*	*	0.2	0.3	0.6	1.8	6.6	29.1	142.2

White, both sexes

2004.....	6.3	*	*	*	0.1	0.2	0.3	0.9	2.4	9.4	43.2	167.8
2003.....	6.5	*	*	*	0.1	0.2	0.3	1.0	2.6	9.3	44.7	178.7
2002.....	6.7	*	*	*	0.1	0.2	0.4	0.8	2.4	9.5	46.5	189.5
2001.....	6.6	*	*	*	0.1	0.2	0.4	0.9	2.5	9.6	45.8	190.8
2000.....	6.5	*	*	*	0.1	0.1	0.4	0.9	2.3	9.8	44.4	191.0
1999.....	5.9	*	*	*	0.1	0.2	0.3	0.7	2.3	9.0	40.7	178.0

White, male

2004.....	6.5	*	*	*	*	0.2	0.4	1.2	2.9	12.4	58.8	241.7
2003.....	6.6	*	*	*	0.1	0.2	0.4	1.2	3.2	12.4	61.1	250.8
2002.....	6.9	*	*	*	0.1	0.2	0.5	1.1	3.0	12.5	64.0	274.6
2001.....	6.7	*	*	*	*	0.2	0.5	1.2	3.1	12.9	63.3	275.4
2000.....	6.6	*	*	*	*	0.1	0.5	1.1	2.9	13.4	62.1	284.0
1999.....	6.1	*	*	*	*	0.2	0.4	0.8	2.9	12.2	58.7	263.6

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pneumonitis due to solids and liquids (J69)												
White, female												
2004.....	6.1	*	*	*	*	0.1	0.3	0.6	1.9	6.8	32.6	134.5
2003.....	6.4	*	*	*	*	0.1	0.3	0.7	2.0	6.7	33.6	146.7
2002.....	6.6	*	*	*	*	0.2	0.3	0.6	1.9	6.9	34.8	152.5
2001.....	6.6	*	*	*	*	0.2	0.3	0.7	2.1	6.9	34.2	155.3
2000.....	6.3	*	*	*	*	0.2	0.3	0.7	1.8	6.8	32.8	153.2
1999.....	5.8	*	*	*	*	0.2	0.3	0.6	1.7	6.4	29.0	143.6
Black, both sexes												
2004.....	3.9	*	*	*	*	*	0.6	1.3	4.8	13.9	46.9	144.7
2003.....	4.2	*	*	*	*	*	0.7	1.3	5.1	13.7	54.4	163.4
2002.....	4.1	*	*	*	*	*	0.7	1.4	3.7	14.8	52.2	165.1
2001.....	4.3	*	*	*	*	*	0.6	1.6	4.0	15.1	53.2	185.5
2000.....	4.2	*	*	*	*	*	0.7	1.9	4.2	17.5	51.5	159.8
1999.....	4.2	*	*	*	*	*	0.7	1.5	4.4	16.2	52.2	167.1
Black, male												
2004.....	4.2	*	*	*	*	*	0.7	1.7	7.1	21.2	67.7	202.7
2003.....	4.3	*	*	*	*	*	*	1.6	7.1	19.6	77.8	212.7
2002.....	4.2	*	*	*	*	*	0.8	2.0	4.8	21.7	73.4	233.7
2001.....	4.2	*	*	*	*	*	0.7	1.9	5.3	20.4	75.3	243.9
2000.....	4.4	*	*	*	*	*	0.8	2.7	5.0	27.8	74.5	209.9
1999.....	4.5	*	*	*	*	*	0.9	1.8	6.4	25.0	80.9	228.5
Black, female												
2004.....	3.6	*	*	*	*	*	*	1.0	3.0	8.7	35.2	121.7
2003.....	4.1	*	*	*	*	*	0.8	1.0	3.6	9.6	41.1	144.2
2002.....	3.9	*	*	*	*	*	*	0.9	2.9	9.9	40.1	138.8
2001.....	4.4	*	*	*	*	*	*	1.4	2.9	11.3	40.8	163.8
2000.....	3.9	*	*	*	*	*	*	1.2	3.5	10.3	38.5	141.2
1999.....	3.8	*	*	*	*	*	*	1.2	2.8	10.0	36.1	144.1

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pneumonitis due to food and vomit (J69.0)												
All races 3/, both sexes												
2004.....	5.7	*	*	*	0.1	0.2	0.4	0.9	2.6	9.6	42.9	164.9
2003.....	6.0	*	*	*	0.1	0.2	0.4	1.0	2.8	9.6	45.0	176.2
2002.....	6.1	*	*	*	0.1	0.2	0.4	0.9	2.5	9.8	46.3	186.0
2001.....	6.1	*	*	*	0.1	0.2	0.4	1.0	2.6	10.0	45.8	189.4
2000.....	5.9	*	*	*	0.1	0.2	0.4	1.0	2.5	10.3	44.5	187.6
1999.....	5.5	*	*	*	0.1	0.2	0.4	0.8	2.5	9.5	41.1	175.6
All races 3/, male												
2004.....	5.9	*	*	*	0.1	0.2	0.4	1.2	3.2	12.9	58.6	237.2
2003.....	6.1	*	*	*	0.1	0.2	0.4	1.2	3.5	12.8	61.7	246.2
2002.....	6.3	*	*	*	0.1	0.2	0.5	1.1	3.1	13.0	63.8	268.5
2001.....	6.1	*	*	*	0.1	0.2	0.5	1.2	3.2	13.3	63.0	270.2
2000.....	6.1	*	*	*	0.1	0.2	0.5	1.2	3.0	14.4	62.3	276.0
1999.....	5.7	*	*	*	0.1	0.2	0.4	0.9	3.2	13.0	59.7	258.4
All races 3/, female												
2004.....	5.5	*	*	*	*	0.1	0.3	0.7	1.9	6.8	32.3	132.4
2003.....	5.8	*	*	*	*	0.1	0.3	0.7	2.1	6.9	33.8	145.3
2002.....	5.9	*	*	*	*	0.2	0.3	0.6	2.0	7.1	34.7	150.2
2001.....	6.0	*	*	*	*	0.2	0.3	0.7	2.1	7.2	34.5	155.5
2000.....	5.8	*	*	*	*	0.1	0.3	0.7	1.9	7.0	33.0	151.6
1999.....	5.3	*	*	*	*	0.2	0.3	0.6	1.8	6.6	29.0	142.2
White, both sexes												
2004.....	6.3	*	*	*	0.1	0.2	0.3	0.9	2.4	9.4	43.2	167.8
2003.....	6.5	*	*	*	0.1	0.2	0.3	1.0	2.6	9.3	44.6	178.7
2002.....	6.7	*	*	*	0.1	0.2	0.4	0.8	2.4	9.5	46.5	189.5
2001.....	6.6	*	*	*	0.1	0.2	0.4	0.9	2.5	9.6	45.8	190.7
2000.....	6.5	*	*	*	0.1	0.1	0.4	0.9	2.3	9.8	44.4	191.0
1999.....	5.9	*	*	*	0.1	0.2	0.3	0.7	2.3	9.0	40.7	178.0
White, male												
2004.....	6.5	*	*	*	*	0.2	0.4	1.2	2.9	12.4	58.7	241.7
2003.....	6.6	*	*	*	0.1	0.2	0.4	1.2	3.2	12.4	61.0	250.8
2002.....	6.9	*	*	*	0.1	0.2	0.5	1.1	3.0	12.5	64.0	274.6
2001.....	6.7	*	*	*	*	0.2	0.5	1.2	3.1	12.9	63.2	275.3
2000.....	6.6	*	*	*	*	0.1	0.5	1.1	2.9	13.4	62.1	284.0
1999.....	6.1	*	*	*	*	0.2	0.4	0.8	2.9	12.1	58.7	263.6

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pneumonitis due to food and vomit (J69.0)												
White, female												
2004.....	6.1	*	*	*	*	0.1	0.3	0.6	1.9	6.8	32.6	134.5
2003.....	6.4	*	*	*	*	0.1	0.3	0.7	2.0	6.7	33.5	146.7
2002.....	6.6	*	*	*	*	0.2	0.3	0.6	1.9	6.9	34.8	152.5
2001.....	6.6	*	*	*	*	0.2	0.3	0.7	2.1	6.9	34.2	155.3
2000.....	6.3	*	*	*	*	0.2	0.3	0.7	1.8	6.8	32.8	153.2
1999.....	5.8	*	*	*	*	0.2	0.3	0.6	1.7	6.4	28.9	143.6
Black, both sexes												
2004.....	3.9	*	*	*	*	*	0.6	1.3	4.8	13.9	46.8	144.7
2003.....	4.2	*	*	*	*	*	0.7	1.3	5.1	13.7	54.4	163.1
2002.....	4.1	*	*	*	*	*	0.7	1.4	3.7	14.8	52.1	165.1
2001.....	4.3	*	*	*	*	*	0.6	1.6	4.0	15.1	53.2	185.5
2000.....	4.2	*	*	*	*	*	0.7	1.9	4.2	17.5	51.5	159.8
1999.....	4.2	*	*	*	*	*	0.7	1.5	4.4	16.1	52.1	167.1
Black, male												
2004.....	4.2	*	*	*	*	*	0.7	1.7	7.1	21.2	67.4	202.7
2003.....	4.3	*	*	*	*	*	*	1.6	7.1	19.6	77.8	212.7
2002.....	4.2	*	*	*	*	*	0.8	2.0	4.8	21.7	73.4	233.7
2001.....	4.2	*	*	*	*	*	0.7	1.9	5.3	20.4	75.3	243.9
2000.....	4.4	*	*	*	*	*	0.8	2.7	5.0	27.8	74.5	209.9
1999.....	4.5	*	*	*	*	*	0.9	1.8	6.4	24.9	80.9	228.5
Black, female												
2004.....	3.6	*	*	*	*	*	*	1.0	3.0	8.7	35.2	121.7
2003.....	4.1	*	*	*	*	*	0.8	1.0	3.6	9.6	41.1	143.8
2002.....	3.9	*	*	*	*	*	*	0.9	2.9	9.9	40.0	138.8
2001.....	4.4	*	*	*	*	*	*	1.4	2.9	11.3	40.8	163.8
2000.....	3.9	*	*	*	*	*	*	1.2	3.5	10.3	38.5	141.2
1999.....	3.8	*	*	*	*	*	*	1.2	2.8	10.0	36.0	144.1

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Suppurative and necrotic conditions of lower respiratory tract (J85-J86)

All races 3/, both sexes

2004.....	0.2	*	*	*	*	*	0.1	0.2	0.4	0.6	1.2	2.2
2003.....	0.2	*	*	*	*	*	0.1	0.3	0.4	0.8	1.3	2.1
2002.....	0.2	*	*	*	*	*	0.1	0.2	0.4	0.8	1.5	2.3
2001.....	0.3	*	*	*	*	*	0.1	0.3	0.4	0.8	1.7	3.0
2000.....	0.3	*	*	*	*	*	0.1	0.2	0.4	0.8	1.5	2.9
1999.....	0.3	*	*	*	*	0.1	0.1	0.3	0.4	1.0	1.5	2.8

All races 3/, male

2004.....	0.3	*	*	*	*	*	0.1	0.3	0.5	1.0	1.7	3.8
2003.....	0.3	*	*	*	*	*	0.1	0.4	0.5	1.1	1.6	3.3
2002.....	0.3	*	*	*	*	*	0.1	0.4	0.6	1.1	2.3	3.4
2001.....	0.3	*	*	*	*	*	0.2	0.4	0.6	1.0	2.3	4.7
2000.....	0.3	*	*	*	*	*	0.1	0.3	0.5	1.2	2.2	5.8
1999.....	0.4	*	*	*	*	*	0.2	0.4	0.6	1.5	2.1	4.0

All races 3/, female

2004.....	0.2	*	*	*	*	*	*	0.2	0.2	0.4	0.9	1.5
2003.....	0.2	*	*	*	*	*	*	0.2	0.3	0.5	1.2	1.5
2002.....	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.0	1.8
2001.....	0.2	*	*	*	*	*	0.1	0.2	0.3	0.6	1.2	2.3
2000.....	0.2	*	*	*	*	*	*	0.1	0.2	0.6	1.1	1.7
1999.....	0.2	*	*	*	*	*	*	0.2	0.3	0.5	1.1	2.3

White, both sexes

2004.....	0.2	*	*	*	*	*	0.1	0.2	0.3	0.6	1.2	2.2
2003.....	0.3	*	*	*	*	*	0.1	0.3	0.4	0.8	1.3	2.1
2002.....	0.2	*	*	*	*	*	0.1	0.2	0.3	0.8	1.4	2.2
2001.....	0.3	*	*	*	*	*	0.1	0.2	0.4	0.8	1.6	3.0
2000.....	0.3	*	*	*	*	*	0.1	0.2	0.3	0.8	1.5	2.7
1999.....	0.3	*	*	*	*	*	0.1	0.3	0.4	0.9	1.5	2.5

White, male

2004.....	0.3	*	*	*	*	*	0.1	0.3	0.5	1.0	1.7	3.8
2003.....	0.3	*	*	*	*	*	0.1	0.4	0.5	1.0	1.5	3.2
2002.....	0.3	*	*	*	*	*	0.1	0.3	0.5	1.1	2.1	3.2
2001.....	0.3	*	*	*	*	*	0.1	0.3	0.6	0.9	2.2	4.7
2000.....	0.3	*	*	*	*	*	0.1	0.3	0.5	1.1	2.1	5.5
1999.....	0.3	*	*	*	*	*	0.2	0.4	0.5	1.5	2.1	3.3

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pleural effusion and plaque (J90,J92)												
All races 3/, both sexes												
2004.....	0.3	*	*	*	*	*	*	0.0	0.2	0.7	2.5	8.8
2003.....	0.3	*	*	*	*	*	*	*	0.2	0.7	2.4	8.8
2002.....	0.3	*	*	*	*	*	*	0.1	0.2	0.7	2.5	9.9
2001.....	0.3	*	*	*	*	*	*	0.1	0.2	0.7	2.5	10.0
2000.....	0.3	*	*	*	*	*	*	0.1	0.2	0.6	2.4	8.9
1999.....	0.3	*	*	*	*	*	*	0.1	0.2	0.6	2.4	9.0
All races 3/, male												
2004.....	0.3	*	*	*	*	*	*	*	0.2	0.8	3.0	11.1
2003.....	0.3	*	*	*	*	*	*	*	0.2	0.7	2.8	11.3
2002.....	0.3	*	*	*	*	*	*	*	0.2	0.6	2.9	11.0
2001.....	0.3	*	*	*	*	*	*	*	0.2	0.8	2.8	12.2
2000.....	0.3	*	*	*	*	*	*	*	0.2	0.8	2.8	10.8
1999.....	0.3	*	*	*	*	*	*	*	0.2	0.8	3.0	11.5
All races 3/, female												
2004.....	0.4	*	*	*	*	*	*	*	0.2	0.6	2.1	7.8
2003.....	0.4	*	*	*	*	*	*	*	0.2	0.6	2.2	7.7
2002.....	0.4	*	*	*	*	*	*	*	*	0.7	2.2	9.4
2001.....	0.4	*	*	*	*	*	*	*	0.2	0.5	2.4	9.1
2000.....	0.3	*	*	*	*	*	*	*	*	0.5	2.1	8.1
1999.....	0.3	*	*	*	*	*	*	*	0.2	0.5	2.0	8.0
White, both sexes												
2004.....	0.4	*	*	*	*	*	*	*	0.2	0.7	2.5	9.0
2003.....	0.4	*	*	*	*	*	*	*	0.2	0.7	2.4	9.2
2002.....	0.4	*	*	*	*	*	*	0.1	0.2	0.6	2.5	10.2
2001.....	0.4	*	*	*	*	*	*	*	0.2	0.6	2.6	10.0
2000.....	0.3	*	*	*	*	*	*	*	0.2	0.6	2.4	9.0
1999.....	0.3	*	*	*	*	*	*	*	0.2	0.7	2.4	9.0
White, male												
2004.....	0.3	*	*	*	*	*	*	*	0.2	0.8	3.1	11.5
2003.....	0.3	*	*	*	*	*	*	*	*	0.7	2.7	11.9
2002.....	0.3	*	*	*	*	*	*	*	0.2	0.6	3.0	11.6
2001.....	0.3	*	*	*	*	*	*	*	*	0.8	2.8	12.5
2000.....	0.3	*	*	*	*	*	*	*	*	0.7	2.9	10.9
1999.....	0.3	*	*	*	*	*	*	*	*	0.9	3.0	11.6

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other diseases of respiratory system (J80-J84,J93-J98)												
All races 3/, both sexes												
2004.....	8.5	5.9	0.5	0.1	0.2	0.4	1.0	2.8	8.7	28.1	69.2	125.2
2003.....	8.0	6.2	0.4	0.1	0.2	0.4	0.9	2.8	7.8	26.2	63.5	123.3
2002.....	7.9	6.1	0.4	0.1	0.2	0.4	1.0	2.7	8.3	26.6	63.0	118.1
2001.....	8.3	6.5	0.4	0.1	0.3	0.4	1.0	3.0	8.8	28.1	67.6	125.7
2000.....	8.3	6.4	0.4	0.2	0.3	0.5	1.1	3.1	9.2	27.5	66.8	125.8
1999.....	8.1	5.8	0.4	0.1	0.2	0.5	1.0	2.8	8.6	27.1	65.3	131.6
All races 3/, male												
2004.....	8.5	5.9	0.6	0.1	0.2	0.3	1.0	3.0	9.7	34.6	87.5	149.7
2003.....	7.8	7.1	0.4	0.1	0.3	0.4	0.9	2.9	9.1	31.7	79.1	145.5
2002.....	7.7	7.3	0.5	0.1	0.2	0.4	1.0	2.7	9.4	32.9	77.5	138.9
2001.....	8.1	8.1	0.4	0.2	0.3	0.4	1.1	3.2	9.9	34.1	82.5	147.4
2000.....	8.1	7.1	0.4	0.2	0.3	0.5	1.2	3.2	10.4	33.0	83.8	158.3
1999.....	7.9	6.6	0.4	0.1	0.3	0.5	1.1	2.9	9.5	33.1	82.9	161.8
All races 3/, female												
2004.....	8.6	5.8	0.3	*	0.3	0.4	1.0	2.6	7.8	22.7	56.9	114.2
2003.....	8.1	5.3	0.4	*	0.2	0.4	1.0	2.7	6.7	21.6	53.0	113.5
2002.....	8.0	4.9	0.3	0.2	0.3	0.4	1.0	2.8	7.3	21.4	53.4	109.0
2001.....	8.6	4.9	0.4	0.1	0.3	0.5	1.0	2.7	7.7	23.2	57.9	116.6
2000.....	8.4	5.7	0.3	0.1	0.3	0.5	1.1	2.9	8.0	23.0	55.7	112.6
1999.....	8.3	4.9	0.3	0.2	0.2	0.6	1.0	2.8	7.7	22.2	53.8	119.4
White, both sexes												
2004.....	9.3	4.2	0.4	0.1	0.2	0.3	0.9	2.5	8.5	29.0	71.8	129.8
2003.....	8.7	4.6	0.3	0.1	0.2	0.4	0.8	2.5	7.7	27.0	65.9	128.0
2002.....	8.6	4.2	0.3	0.1	0.2	0.4	0.9	2.5	8.1	27.3	65.1	122.1
2001.....	9.1	4.5	0.4	0.2	0.2	0.4	0.9	2.6	8.5	28.7	69.8	129.0
2000.....	9.0	4.9	0.4	0.2	0.2	0.5	1.0	2.7	9.0	28.1	68.5	130.0
1999.....	8.8	4.1	0.3	0.1	0.2	0.4	0.9	2.5	8.4	27.4	66.8	134.9
White, male												
2004.....	9.3	4.2	0.5	*	0.2	0.3	0.8	2.7	9.6	35.5	90.4	154.6
2003.....	8.6	4.9	0.3	0.1	0.2	0.4	0.8	2.7	8.9	32.6	81.7	150.4
2002.....	8.4	5.0	0.5	*	0.2	0.3	0.9	2.5	9.1	33.8	79.9	143.8
2001.....	8.8	5.3	0.3	0.2	0.2	0.3	1.0	2.9	9.6	35.1	85.3	151.5
2000.....	8.8	5.8	0.4	0.2	0.3	0.4	1.0	2.9	10.3	33.6	85.9	163.4
1999.....	8.6	4.8	0.4	0.1	0.3	0.5	0.9	2.6	9.3	33.8	84.9	166.3

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

All other diseases of respiratory system (J80-J84,J93-J98)

White, female

2004.....	9.4	4.2	*	*	0.2	0.3	0.9	2.2	7.6	23.4	59.1	118.7
2003.....	8.9	4.4	*	*	0.2	0.4	0.7	2.3	6.6	22.2	55.2	118.0
2002.....	8.8	3.5	*	*	0.2	0.4	0.9	2.5	7.1	21.7	55.1	112.7
2001.....	9.3	3.5	0.4	0.1	0.2	0.4	0.8	2.3	7.4	23.4	59.5	119.6
2000.....	9.2	4.1	*	0.1	0.2	0.5	1.0	2.6	7.8	23.4	57.0	116.4
1999.....	9.0	3.4	*	0.1	0.2	0.4	0.8	2.5	7.5	22.2	54.9	122.2

Black, both sexes

2004.....	5.5	14.1	0.8	*	*	0.7	2.3	5.6	12.2	25.2	48.0	79.0
2003.....	5.1	14.1	*	*	0.5	0.9	2.1	5.4	10.5	22.0	44.4	77.3
2002.....	5.3	14.8	0.9	0.3	0.5	0.8	1.9	5.2	12.1	23.9	45.0	76.9
2001.....	5.8	16.6	*	*	0.5	1.1	2.4	6.3	12.6	26.6	50.3	90.0
2000.....	5.7	15.1	*	*	0.5	0.8	2.3	6.1	12.2	26.7	52.6	82.3
1999.....	5.6	14.5	*	*	*	1.1	2.4	5.6	11.0	27.7	52.3	94.1

Black, male

2004.....	5.6	14.8	*	*	*	0.8	2.6	5.9	13.5	31.9	61.5	100.3
2003.....	5.2	17.6	*	*	0.8	0.9	1.8	4.9	13.1	27.6	59.5	98.0
2002.....	5.2	17.7	*	*	*	0.8	2.1	5.2	13.9	28.8	55.7	86.8
2001.....	5.4	21.3	*	*	*	1.0	1.9	6.4	13.8	29.5	58.6	106.7
2000.....	5.6	15.0	*	*	*	0.8	2.6	6.8	13.5	30.7	66.0	104.9
1999.....	5.3	16.9	*	*	*	0.8	2.2	6.0	12.0	31.2	66.1	114.8

Black, female

2004.....	5.5	13.3	*	*	*	0.7	2.0	5.3	11.2	20.4	40.3	70.5
2003.....	5.0	10.5	*	*	*	0.9	2.4	5.8	8.4	18.0	35.9	69.3
2002.....	5.4	11.8	*	*	*	0.8	1.8	5.3	10.6	20.4	39.0	73.1
2001.....	6.2	11.7	*	*	*	1.1	2.8	6.2	11.6	24.5	45.6	83.8
2000.....	5.7	15.2	*	*	*	0.9	2.0	5.4	11.2	23.8	45.0	73.8
1999.....	5.9	12.1	*	*	*	1.4	2.5	5.3	10.3	25.3	44.6	86.3

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
XI. Diseases of the digestive system (K00-K92)												
All races 3/, both sexes												
2004.....	29.4	14.2	0.8	0.2	0.7	2.1	10.9	29.5	44.5	78.4	162.5	387.5
2003.....	30.2	13.8	0.8	0.3	0.6	2.4	11.5	30.2	45.4	82.5	167.1	403.8
2002.....	30.1	12.6	0.8	0.3	0.6	2.3	11.7	29.8	44.7	81.7	170.0	415.3
2001.....	30.1	11.9	0.8	0.3	0.5	2.4	11.7	29.7	45.2	83.8	168.8	431.4
2000.....	29.9	13.1	0.7	0.3	0.7	2.4	11.9	28.6	45.6	82.1	170.4	440.7
1999.....	29.7	13.2	0.7	0.3	0.6	2.4	11.8	28.0	45.8	85.2	169.0	436.4
All races 3/, male												
2004.....	30.0	15.6	0.9	0.2	0.7	2.8	14.1	41.3	58.6	91.6	176.1	366.9
2003.....	30.9	16.6	0.9	0.3	0.7	2.9	14.6	42.6	60.5	97.6	181.6	388.9
2002.....	30.7	15.1	0.9	0.3	0.7	2.9	15.0	42.5	59.0	96.9	184.1	404.4
2001.....	30.6	13.4	0.9	0.3	0.6	2.9	15.2	41.9	59.8	99.5	181.8	428.2
2000.....	30.3	14.8	0.9	0.3	0.7	3.0	15.8	40.7	60.1	97.7	186.4	441.9
1999.....	30.4	16.5	0.7	0.3	0.6	2.9	15.8	40.4	60.7	102.3	185.8	438.5
All races 3/, female												
2004.....	28.8	12.8	0.6	0.3	0.7	1.5	7.7	18.2	31.4	67.3	153.3	396.8
2003.....	29.4	10.8	0.7	0.3	0.5	1.8	8.4	18.2	31.4	69.9	157.4	410.4
2002.....	29.5	10.0	0.7	0.3	0.6	1.8	8.5	17.7	31.5	69.0	160.6	420.0
2001.....	29.7	10.3	0.6	0.3	0.5	1.9	8.3	18.0	31.8	70.8	160.2	432.7
2000.....	29.4	11.4	0.5	0.3	0.5	1.8	8.1	16.9	32.3	69.3	159.9	440.2
1999.....	29.1	9.7	0.8	0.4	0.6	1.9	7.8	15.9	32.2	71.1	158.1	435.6
White, both sexes												
2004.....	31.5	10.7	0.6	0.2	0.5	2.0	10.8	29.2	43.5	78.8	164.8	396.3
2003.....	32.3	11.0	0.7	0.3	0.5	2.3	11.4	29.5	44.3	83.0	169.9	412.6
2002.....	32.2	10.1	0.7	0.3	0.6	2.1	11.7	29.0	43.8	82.2	172.3	426.3
2001.....	32.1	8.7	0.7	0.3	0.4	2.2	11.4	28.6	43.7	84.0	170.3	441.6
2000.....	31.7	10.0	0.5	0.3	0.6	2.1	11.5	27.2	44.1	81.8	171.7	449.3
1999.....	31.5	10.1	0.7	0.3	0.5	2.3	11.3	26.6	44.2	84.6	170.1	445.2
White, male												
2004.....	31.6	11.4	0.7	0.2	0.6	2.6	14.1	40.9	56.9	91.1	177.0	375.1
2003.....	32.6	13.9	0.9	0.3	0.7	2.8	14.7	41.8	58.6	97.2	182.9	395.9
2002.....	32.4	12.1	0.7	0.3	0.6	2.7	15.2	41.6	57.6	96.8	185.5	412.5
2001.....	31.9	9.3	0.8	0.3	0.5	2.6	14.7	40.4	57.4	98.8	182.5	434.7
2000.....	31.7	11.2	0.6	0.3	0.7	2.8	15.3	39.1	58.2	96.9	186.8	448.1
1999.....	31.7	12.5	0.5	0.3	0.6	2.8	15.4	38.5	58.6	101.7	186.8	443.1

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
XI. Diseases of the digestive system (K00-K92)												
White, female												
2004.....	31.3	9.9	0.5	0.2	0.5	1.4	7.4	17.6	30.8	68.3	156.5	405.9
2003.....	32.0	8.0	0.6	0.3	0.4	1.7	8.1	17.3	30.7	70.8	161.1	420.0
2002.....	32.1	7.9	0.7	0.2	0.5	1.5	8.2	16.7	30.8	69.8	163.4	432.3
2001.....	32.2	8.1	0.5	0.3	0.4	1.7	8.0	16.9	30.8	71.4	162.3	444.5
2000.....	31.6	8.8	0.4	0.3	0.5	1.4	7.5	15.6	30.9	69.2	161.8	449.8
1999.....	31.2	7.5	0.8	0.3	0.5	1.8	7.2	14.8	30.7	70.3	159.2	446.0
Black, both sexes												
2004.....	23.7	31.6	1.4	0.4	1.2	2.9	12.6	37.7	60.9	88.5	158.5	329.7
2003.....	24.2	29.0	1.2	0.4	1.1	3.4	13.2	40.2	63.2	91.6	157.5	346.0
2002.....	23.7	25.9	1.4	0.5	1.0	3.4	12.8	41.0	59.0	89.5	165.2	333.2
2001.....	25.1	28.6	1.2	0.5	1.0	3.5	15.0	43.1	65.0	96.7	171.0	351.0
2000.....	25.2	29.6	1.5	0.4	1.1	3.8	16.0	43.3	64.8	96.6	170.5	382.0
1999.....	25.1	29.6	1.1	0.4	1.2	3.4	16.1	42.9	65.9	100.4	171.9	364.4
Black, male												
2004.....	26.4	36.2	1.7	*	1.1	3.8	15.5	52.7	86.9	116.0	188.8	322.9
2003.....	27.1	32.4	*	*	1.1	4.0	15.7	56.7	90.0	119.6	191.2	345.1
2002.....	26.0	31.4	1.7	*	0.8	3.9	15.0	57.4	82.2	113.9	189.3	355.9
2001.....	28.4	33.5	*	0.6	0.9	4.3	19.8	62.1	90.9	127.0	196.3	386.9
2000.....	27.7	33.9	2.0	*	1.3	4.1	20.6	60.3	88.2	122.0	202.9	392.1
1999.....	27.8	36.4	*	*	1.1	4.0	20.8	63.2	89.2	122.2	195.2	400.7
Black, female												
2004.....	21.1	26.9	*	*	1.3	2.1	10.0	24.9	40.1	68.8	141.5	332.3
2003.....	21.6	25.4	*	*	1.1	2.8	10.9	26.0	41.5	71.7	138.4	346.4
2002.....	21.5	20.3	*	*	1.3	3.0	10.8	27.0	40.3	72.1	151.5	324.5
2001.....	22.1	23.3	*	*	1.2	2.7	10.8	26.9	44.2	75.4	156.8	337.6
2000.....	23.0	25.1	*	*	0.8	3.6	11.9	28.6	45.9	78.7	152.2	378.2
1999.....	22.7	22.5	*	*	1.3	2.8	12.0	25.5	47.2	84.9	158.9	350.9

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Diseases of other parts of digestive system (K20-K92)

All races 3/, both sexes

2004.....	29.3	14.1	0.8	0.2	0.7	2.1	10.9	29.5	44.4	78.3	162.0	385.7
2003.....	30.1	13.7	0.8	0.3	0.6	2.4	11.5	30.2	45.4	82.4	166.9	402.6
2002.....	30.0	12.5	0.8	0.3	0.6	2.3	11.7	29.8	44.6	81.6	169.7	413.8
2001.....	30.1	11.8	0.8	0.3	0.5	2.4	11.7	29.7	45.2	83.8	168.3	430.2
2000.....	29.8	13.0	0.7	0.3	0.7	2.4	11.9	28.6	45.6	82.1	170.1	439.5
1999.....	29.7	13.1	0.7	0.3	0.6	2.4	11.8	27.9	45.8	85.0	168.8	435.4

All races 3/, male

2004.....	29.9	15.5	0.9	0.2	0.7	2.7	14.1	41.3	58.5	91.5	175.6	365.5
2003.....	30.9	16.5	0.9	0.3	0.7	2.9	14.6	42.6	60.5	97.5	181.4	388.0
2002.....	30.6	15.0	0.9	0.3	0.7	2.9	15.0	42.4	58.9	96.7	183.8	403.1
2001.....	30.5	13.3	0.9	0.3	0.6	2.8	15.2	41.8	59.7	99.5	181.4	427.0
2000.....	30.3	14.8	0.9	0.3	0.7	3.0	15.8	40.6	60.1	97.6	186.2	440.8
1999.....	30.3	16.4	0.7	0.3	0.6	2.9	15.8	40.4	60.6	102.2	185.6	437.4

All races 3/, female

2004.....	28.7	12.6	0.6	0.3	0.7	1.5	7.7	18.1	31.3	67.2	152.9	394.7
2003.....	29.4	10.8	0.7	0.3	0.5	1.8	8.3	18.2	31.4	69.8	157.2	409.1
2002.....	29.4	9.9	0.7	0.3	0.6	1.8	8.5	17.6	31.4	68.9	160.3	418.4
2001.....	29.7	10.2	0.6	0.3	0.5	1.8	8.3	18.0	31.8	70.8	159.8	431.5
2000.....	29.4	11.2	0.5	0.3	0.5	1.8	8.1	16.9	32.3	69.3	159.7	439.0
1999.....	29.0	9.7	0.8	0.4	0.6	1.9	7.8	15.9	32.2	71.0	157.8	434.6

White, both sexes

2004.....	31.4	10.5	0.6	0.2	0.5	2.0	10.8	29.1	43.4	78.7	164.4	394.3
2003.....	32.2	11.0	0.7	0.3	0.5	2.2	11.4	29.4	44.2	82.9	169.6	411.3
2002.....	32.2	10.0	0.7	0.3	0.6	2.1	11.7	29.0	43.7	82.1	172.0	424.8
2001.....	32.0	8.6	0.7	0.3	0.4	2.2	11.4	28.6	43.7	83.9	170.0	440.4
2000.....	31.6	9.9	0.5	0.3	0.6	2.1	11.4	27.2	44.1	81.8	171.5	448.1
1999.....	31.4	10.0	0.7	0.3	0.5	2.3	11.3	26.5	44.2	84.5	169.9	444.1

White, male

2004.....	31.6	11.3	0.7	0.2	0.6	2.6	14.1	40.9	56.8	91.0	176.5	373.5
2003.....	32.6	13.8	0.9	0.3	0.6	2.8	14.7	41.8	58.6	97.2	182.7	394.9
2002.....	32.4	12.0	0.7	0.3	0.6	2.7	15.2	41.5	57.5	96.7	185.2	411.2
2001.....	31.9	9.3	0.8	0.3	0.5	2.6	14.7	40.4	57.4	98.8	182.2	433.4
2000.....	31.7	11.2	0.6	0.3	0.7	2.8	15.3	39.1	58.2	96.8	186.6	446.9
1999.....	31.7	12.4	0.5	0.3	0.5	2.8	15.3	38.4	58.6	101.5	186.6	441.9

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diseases of other parts of digestive system (K20-K92)												
White, female												
2004.....	31.2	9.7	0.5	0.2	0.5	1.4	7.4	17.5	30.7	68.1	156.1	403.7
2003.....	31.9	8.0	0.6	0.3	0.4	1.7	8.1	17.3	30.6	70.7	160.8	418.6
2002.....	32.0	7.9	0.7	0.2	0.5	1.5	8.2	16.7	30.8	69.7	163.1	430.6
2001.....	32.1	8.0	0.5	0.3	0.4	1.7	8.0	16.9	30.8	71.4	161.9	443.3
2000.....	31.6	8.6	0.4	0.3	0.5	1.4	7.5	15.6	30.9	69.2	161.6	448.6
1999.....	31.2	7.5	0.8	0.3	0.5	1.8	7.2	14.8	30.7	70.2	158.9	445.0
Black, both sexes												
2004.....	23.6	31.6	1.4	0.4	1.2	2.8	12.6	37.6	60.9	88.2	158.3	329.4
2003.....	24.2	28.8	1.2	0.4	1.1	3.3	13.1	40.2	63.0	91.5	157.5	346.0
2002.....	23.6	25.9	1.4	0.5	1.0	3.4	12.8	41.0	58.9	89.4	165.0	332.3
2001.....	25.0	28.6	1.2	0.5	1.0	3.5	15.0	43.1	65.0	96.7	170.2	350.1
2000.....	25.2	29.4	1.5	0.4	1.1	3.8	15.9	43.3	64.8	96.4	170.1	381.6
1999.....	25.1	29.6	1.1	0.4	1.2	3.4	16.1	42.9	65.9	100.3	171.5	363.8
Black, male												
2004.....	26.4	36.2	1.7	*	1.1	3.7	15.5	52.5	86.9	115.7	188.5	322.9
2003.....	27.1	32.4	*	*	1.1	4.0	15.7	56.7	90.0	119.5	191.2	345.1
2002.....	25.9	31.4	1.7	*	0.8	3.9	15.0	57.2	82.2	113.8	189.1	355.9
2001.....	28.3	33.5	*	0.6	0.9	4.3	19.8	62.1	90.9	127.0	195.1	385.8
2000.....	27.7	33.9	2.0	*	1.3	4.1	20.6	60.3	88.2	121.9	202.9	392.1
1999.....	27.7	36.4	*	*	1.1	4.0	20.7	63.2	89.1	122.2	194.9	400.7
Black, female												
2004.....	21.1	26.9	*	*	1.2	2.0	10.0	24.9	40.0	68.5	141.3	332.0
2003.....	21.5	25.0	*	*	1.1	2.8	10.9	26.0	41.4	71.6	138.4	346.4
2002.....	21.5	20.3	*	*	1.2	2.9	10.8	27.0	40.2	72.1	151.4	323.2
2001.....	22.0	23.3	*	*	1.2	2.7	10.8	26.9	44.2	75.4	156.2	336.8
2000.....	22.9	24.8	*	*	0.8	3.6	11.8	28.6	45.9	78.6	151.7	377.7
1999.....	22.6	22.5	*	*	1.3	2.8	12.0	25.5	47.2	84.8	158.4	350.0

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diseases of esophagus (K20-K22)												
All races 3/, both sexes												
2004.....	0.7	*	*	*	*	*	0.1	0.2	0.4	1.2	4.1	17.9
2003.....	0.7	*	*	*	*	*	0.1	0.2	0.4	1.2	4.4	19.3
2002.....	0.7	*	*	*	*	0.1	0.1	0.3	0.4	1.3	4.9	20.7
2001.....	0.7	*	*	*	*	*	0.1	0.2	0.5	1.4	5.1	21.2
2000.....	0.8	0.5	*	*	*	*	0.1	0.2	0.5	1.4	5.2	23.3
1999.....	0.8	*	*	*	*	0.1	0.1	0.2	0.5	1.5	5.0	23.5
All races 3/, male												
2004.....	0.6	*	*	*	*	*	0.1	0.3	0.6	1.4	5.0	19.1
2003.....	0.7	*	*	*	*	*	0.2	0.3	0.6	1.6	4.9	23.6
2002.....	0.7	*	*	*	*	*	0.2	0.4	0.4	1.6	5.8	25.4
2001.....	0.7	*	*	*	*	*	0.1	0.3	0.7	1.7	5.9	23.6
2000.....	0.7	*	*	*	*	*	0.1	0.3	0.7	1.8	6.4	28.6
1999.....	0.7	*	*	*	*	*	0.1	0.3	0.6	2.0	6.5	28.3
All races 3/, female												
2004.....	0.7	*	*	*	*	*	*	0.1	0.3	1.1	3.5	17.3
2003.....	0.7	*	*	*	*	*	*	0.1	0.3	0.9	4.0	17.4
2002.....	0.8	*	*	*	*	*	*	0.2	0.3	1.1	4.3	18.7
2001.....	0.8	*	*	*	*	*	*	*	0.3	1.1	4.6	20.2
2000.....	0.8	*	*	*	*	*	*	0.1	0.3	1.1	4.5	21.1
1999.....	0.8	*	*	*	*	*	*	0.2	0.4	1.1	4.1	21.5
White, both sexes												
2004.....	0.7	*	*	*	*	*	0.1	0.2	0.4	1.2	4.2	18.8
2003.....	0.8	*	*	*	*	*	0.1	0.2	0.4	1.3	4.6	20.2
2002.....	0.8	*	*	*	*	*	0.1	0.3	0.4	1.3	4.9	21.7
2001.....	0.8	*	*	*	*	*	0.1	0.2	0.5	1.4	5.2	22.2
2000.....	0.9	*	*	*	*	*	0.1	0.2	0.5	1.4	5.4	24.4
1999.....	0.9	*	*	*	*	*	0.1	0.2	0.5	1.5	5.2	24.5
White, male												
2004.....	0.6	*	*	*	*	*	0.1	0.3	0.5	1.3	5.1	20.1
2003.....	0.7	*	*	*	*	*	0.2	0.3	0.6	1.7	5.1	24.9
2002.....	0.8	*	*	*	*	*	0.2	0.4	0.4	1.6	5.8	26.7
2001.....	0.7	*	*	*	*	*	0.1	0.3	0.6	1.6	5.9	24.7
2000.....	0.8	*	*	*	*	*	0.1	0.3	0.7	1.8	6.5	30.0
1999.....	0.8	*	*	*	*	*	0.1	0.2	0.6	2.0	6.6	29.8

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Peptic ulcer (K25-K28)												
All races 3/, both sexes												
2004.....	1.3	*	*	*	*	*	0.2	0.6	1.3	3.2	8.9	24.4
2003.....	1.3	*	*	*	*	0.1	0.2	0.7	1.4	3.2	9.9	26.4
2002.....	1.4	*	*	*	*	0.1	0.3	0.7	1.5	3.6	10.1	28.3
2001.....	1.6	*	*	*	*	0.1	0.3	0.7	1.7	3.9	11.4	33.4
2000.....	1.6	*	*	*	*	*	0.3	0.8	1.8	4.1	11.4	35.6
1999.....	1.6	*	*	*	*	0.1	0.3	0.7	1.8	4.5	12.3	34.6
All races 3/, male												
2004.....	1.2	*	*	*	*	*	0.3	0.9	1.6	3.6	10.2	24.6
2003.....	1.3	*	*	*	*	*	0.3	1.0	1.8	4.1	11.8	28.9
2002.....	1.4	*	*	*	*	*	0.3	0.9	1.9	4.6	12.2	31.6
2001.....	1.5	*	*	*	*	0.1	0.4	1.0	2.2	4.8	13.0	34.9
2000.....	1.6	*	*	*	*	*	0.4	1.2	2.4	5.3	13.7	40.7
1999.....	1.6	*	*	*	*	*	0.3	0.9	2.3	6.1	14.8	37.5
All races 3/, female												
2004.....	1.3	*	*	*	*	*	0.1	0.4	1.0	2.8	8.0	24.3
2003.....	1.4	*	*	*	*	*	0.2	0.4	1.0	2.4	8.7	25.3
2002.....	1.5	*	*	*	*	*	0.2	0.5	1.2	2.9	8.7	26.9
2001.....	1.7	*	*	*	*	*	0.1	0.5	1.1	3.2	10.4	32.7
2000.....	1.6	*	*	*	*	*	0.1	0.5	1.3	3.0	10.0	33.5
1999.....	1.7	*	*	*	*	*	0.2	0.5	1.3	3.3	10.6	33.5
White, both sexes												
2004.....	1.3	*	*	*	*	*	0.2	0.6	1.2	3.1	8.9	25.1
2003.....	1.4	*	*	*	*	0.1	0.2	0.6	1.3	3.2	9.8	26.9
2002.....	1.5	*	*	*	*	*	0.2	0.6	1.4	3.6	10.3	29.1
2001.....	1.7	*	*	*	*	*	0.2	0.7	1.5	3.9	11.5	34.4
2000.....	1.7	*	*	*	*	*	0.2	0.7	1.8	4.0	11.5	36.3
1999.....	1.8	*	*	*	*	*	0.2	0.6	1.7	4.4	12.3	35.3
White, male												
2004.....	1.2	*	*	*	*	*	0.2	0.8	1.6	3.6	10.0	25.5
2003.....	1.4	*	*	*	*	*	0.2	0.8	1.6	4.1	11.4	29.0
2002.....	1.5	*	*	*	*	*	0.3	0.8	1.8	4.4	12.3	32.5
2001.....	1.5	*	*	*	*	*	0.3	0.8	2.0	4.5	12.9	35.7
2000.....	1.7	*	*	*	*	*	0.3	1.0	2.4	5.2	13.5	40.7
1999.....	1.7	*	*	*	*	*	0.2	0.8	2.2	5.9	14.9	38.4

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Peptic ulcer (K25-K28)												
White, female												
2004.....	1.5	*	*	*	*	*	0.1	0.4	1.0	2.7	8.2	25.0
2003.....	1.5	*	*	*	*	*	0.2	0.4	1.0	2.5	8.8	26.0
2002.....	1.6	*	*	*	*	*	0.1	0.5	1.1	2.9	9.0	27.6
2001.....	1.9	*	*	*	*	*	0.1	0.5	1.1	3.3	10.7	33.9
2000.....	1.8	*	*	*	*	*	0.1	0.4	1.2	3.0	10.2	34.5
1999.....	1.9	*	*	*	*	*	0.2	0.4	1.3	3.2	10.7	34.1
Black, both sexes												
2004.....	0.9	*	*	*	*	*	0.5	1.3	2.0	4.2	7.7	16.1
2003.....	1.0	*	*	*	*	*	0.4	1.5	2.2	3.2	11.0	16.9
2002.....	1.0	*	*	*	*	*	0.6	1.2	2.6	4.1	8.0	19.0
2001.....	1.1	*	*	*	*	*	0.6	1.3	2.7	4.4	9.1	20.9
2000.....	1.2	*	*	*	*	*	0.5	1.7	2.7	4.8	10.0	26.6
1999.....	1.2	*	*	*	*	*	0.8	1.5	2.8	5.7	10.9	25.5
Black, male												
2004.....	1.0	*	*	*	*	*	*	1.8	2.9	4.6	11.1	*
2003.....	1.2	*	*	*	*	*	*	2.3	3.4	5.2	16.1	*
2002.....	1.2	*	*	*	*	*	0.8	1.9	3.8	5.9	11.5	*
2001.....	1.4	*	*	*	*	*	1.0	2.2	4.2	7.0	12.4	23.8
2000.....	1.3	*	*	*	*	*	0.8	2.3	3.7	7.0	13.9	31.1
1999.....	1.4	*	*	*	*	*	1.2	2.3	4.4	8.1	12.3	28.1
Black, female												
2004.....	0.9	*	*	*	*	*	*	0.8	1.3	4.0	5.8	16.5
2003.....	0.8	*	*	*	*	*	*	0.8	*	*	8.1	15.8
2002.....	0.8	*	*	*	*	*	*	*	1.6	2.8	6.0	18.5
2001.....	0.9	*	*	*	*	*	*	*	1.5	2.7	7.3	19.8
2000.....	1.0	*	*	*	*	*	*	1.1	1.8	3.3	7.8	24.9
1999.....	1.1	*	*	*	*	*	*	*	*	3.9	10.2	24.5

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Gastric ulcer (K25)												
All races 3/, both sexes												
2004.....	0.4	*	*	*	*	*	0.1	0.3	0.4	0.9	2.7	6.7
2003.....	0.4	*	*	*	*	*	0.1	0.3	0.5	0.9	2.8	7.0
2002.....	0.4	*	*	*	*	*	0.1	0.2	0.5	1.1	2.8	7.7
2001.....	0.4	*	*	*	*	*	0.1	0.3	0.5	1.3	3.3	8.0
2000.....	0.5	*	*	*	*	*	0.1	0.3	0.6	1.3	3.0	9.5
1999.....	0.5	*	*	*	*	*	0.1	0.3	0.6	1.3	3.5	8.8
All races 3/, male												
2004.....	0.3	*	*	*	*	*	0.1	0.3	0.5	1.0	2.9	6.3
2003.....	0.3	*	*	*	*	*	*	0.4	0.6	1.0	2.8	6.4
2002.....	0.4	*	*	*	*	*	0.1	0.3	0.7	1.4	2.9	8.1
2001.....	0.4	*	*	*	*	*	0.1	0.3	0.6	1.4	3.4	6.0
2000.....	0.4	*	*	*	*	*	0.1	0.4	0.7	1.6	3.3	9.9
1999.....	0.4	*	*	*	*	*	0.1	0.4	0.7	1.5	3.7	8.7
All races 3/, female												
2004.....	0.4	*	*	*	*	*	*	0.2	0.3	0.9	2.5	6.9
2003.....	0.4	*	*	*	*	*	*	0.2	0.4	0.8	2.8	7.3
2002.....	0.4	*	*	*	*	*	*	0.2	0.4	0.9	2.8	7.5
2001.....	0.5	*	*	*	*	*	*	0.2	0.4	1.1	3.2	8.8
2000.....	0.5	*	*	*	*	*	*	0.2	0.5	1.0	2.9	9.4
1999.....	0.5	*	*	*	*	*	0.1	0.2	0.5	1.1	3.4	8.8
White, both sexes												
2004.....	0.4	*	*	*	*	*	0.1	0.2	0.4	0.9	2.6	6.8
2003.....	0.4	*	*	*	*	*	0.1	0.2	0.5	0.9	2.7	7.0
2002.....	0.4	*	*	*	*	*	0.1	0.2	0.5	1.1	2.7	7.8
2001.....	0.5	*	*	*	*	*	0.1	0.2	0.4	1.2	3.3	8.0
2000.....	0.5	*	*	*	*	*	0.1	0.3	0.5	1.2	3.0	9.4
1999.....	0.5	*	*	*	*	*	0.1	0.2	0.5	1.2	3.5	8.6
White, male												
2004.....	0.3	*	*	*	*	*	*	0.3	0.4	0.9	2.7	6.7
2003.....	0.3	*	*	*	*	*	*	0.3	0.6	0.9	2.7	6.1
2002.....	0.4	*	*	*	*	*	*	0.2	0.6	1.3	2.8	8.3
2001.....	0.4	*	*	*	*	*	*	0.3	0.5	1.2	3.4	5.9
2000.....	0.4	*	*	*	*	*	*	0.4	0.5	1.4	3.1	9.3
1999.....	0.4	*	*	*	*	*	*	0.3	0.6	1.4	3.7	8.1

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Duodenal ulcer (K26)												
All races 3/, both sexes												
2004.....	0.4	*	*	*	*	*	0.1	0.2	0.5	1.2	3.1	7.7
2003.....	0.5	*	*	*	*	*	0.1	0.3	0.4	1.1	3.3	8.5
2002.....	0.5	*	*	*	*	*	0.1	0.2	0.6	1.3	3.6	9.0
2001.....	0.5	*	*	*	*	*	0.1	0.3	0.6	1.3	3.9	11.0
2000.....	0.5	*	*	*	*	*	0.1	0.3	0.7	1.4	4.0	10.1
1999.....	0.5	*	*	*	*	*	0.1	0.3	0.7	1.5	3.7	10.0
All races 3/, male												
2004.....	0.4	*	*	*	*	*	0.1	0.3	0.7	1.5	3.6	9.2
2003.....	0.5	*	*	*	*	*	0.1	0.4	0.6	1.6	4.4	10.2
2002.....	0.5	*	*	*	*	*	0.1	0.3	0.8	1.7	4.5	10.1
2001.....	0.6	*	*	*	*	*	0.2	0.4	0.8	1.7	4.7	13.8
2000.....	0.6	*	*	*	*	*	0.2	0.5	1.0	1.8	4.8	12.3
1999.....	0.6	*	*	*	*	*	0.1	0.4	0.9	2.4	5.1	11.9
All races 3/, female												
2004.....	0.4	*	*	*	*	*	*	0.1	0.3	0.9	2.7	7.1
2003.....	0.4	*	*	*	*	*	*	0.2	0.3	0.7	2.6	7.8
2002.....	0.5	*	*	*	*	*	*	0.2	0.4	1.0	3.0	8.5
2001.....	0.5	*	*	*	*	*	*	0.2	0.4	1.0	3.4	9.8
2000.....	0.5	*	*	*	*	*	*	0.1	0.5	1.0	3.6	9.2
1999.....	0.5	*	*	*	*	*	*	0.2	0.4	0.9	2.9	9.3
White, both sexes												
2004.....	0.5	*	*	*	*	*	0.1	0.2	0.5	1.2	3.2	8.1
2003.....	0.5	*	*	*	*	*	0.1	0.3	0.4	1.2	3.3	8.8
2002.....	0.5	*	*	*	*	*	0.1	0.2	0.6	1.3	3.8	9.5
2001.....	0.6	*	*	*	*	*	0.1	0.3	0.6	1.4	4.0	11.6
2000.....	0.6	*	*	*	*	*	0.1	0.3	0.7	1.4	4.2	10.5
1999.....	0.6	*	*	*	*	*	0.1	0.2	0.7	1.6	3.8	10.2
White, male												
2004.....	0.5	*	*	*	*	*	*	0.3	0.7	1.5	3.7	9.4
2003.....	0.5	*	*	*	*	*	*	0.4	0.5	1.6	4.4	10.2
2002.....	0.6	*	*	*	*	*	0.1	0.3	0.8	1.7	4.7	10.9
2001.....	0.6	*	*	*	*	*	0.2	0.4	0.8	1.7	4.7	14.5
2000.....	0.6	*	*	*	*	*	0.1	0.4	1.0	1.8	4.9	12.7
1999.....	0.6	*	*	*	*	*	*	0.3	0.9	2.4	5.2	12.5

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Peptic ulcer, site unspecified (K27)												
All races 3/, both sexes												
2004.....	0.4	*	*	*	*	*	0.0	0.2	0.4	1.0	3.0	9.7
2003.....	0.5	*	*	*	*	*	0.1	0.2	0.4	1.1	3.6	10.5
2002.....	0.5	*	*	*	*	*	*	0.2	0.4	1.2	3.6	11.3
2001.....	0.6	*	*	*	*	*	0.1	0.2	0.6	1.3	4.1	14.2
2000.....	0.6	*	*	*	*	*	0.1	0.2	0.5	1.4	4.2	15.5
1999.....	0.6	*	*	*	*	*	0.1	0.1	0.5	1.6	4.9	15.6
All races 3/, male												
2004.....	0.4	*	*	*	*	*	*	0.2	0.5	1.1	3.5	9.0
2003.....	0.5	*	*	*	*	*	*	0.2	0.6	1.4	4.3	11.9
2002.....	0.5	*	*	*	*	*	*	0.3	0.4	1.4	4.6	12.8
2001.....	0.5	*	*	*	*	*	*	0.2	0.8	1.6	4.8	14.9
2000.....	0.6	*	*	*	*	*	*	0.3	0.8	1.9	5.4	18.2
1999.....	0.6	*	*	*	*	*	*	0.2	0.7	2.1	5.8	16.4
All races 3/, female												
2004.....	0.5	*	*	*	*	*	*	0.1	0.3	0.9	2.6	10.0
2003.....	0.5	*	*	*	*	*	*	0.1	0.3	0.8	3.2	9.9
2002.....	0.5	*	*	*	*	*	*	0.1	0.4	0.9	2.9	10.7
2001.....	0.6	*	*	*	*	*	*	0.1	0.4	1.0	3.7	13.9
2000.....	0.6	*	*	*	*	*	*	0.1	0.3	1.0	3.5	14.5
1999.....	0.7	*	*	*	*	*	*	0.1	0.4	1.2	4.3	15.2
White, both sexes												
2004.....	0.5	*	*	*	*	*	*	0.1	0.4	1.0	3.0	9.9
2003.....	0.5	*	*	*	*	*	*	0.1	0.4	1.1	3.6	10.7
2002.....	0.5	*	*	*	*	*	*	0.2	0.4	1.1	3.6	11.5
2001.....	0.6	*	*	*	*	*	*	0.2	0.5	1.3	4.1	14.6
2000.....	0.7	*	*	*	*	*	0.1	0.2	0.5	1.4	4.2	15.9
1999.....	0.7	*	*	*	*	*	*	0.1	0.5	1.6	4.9	16.2
White, male												
2004.....	0.4	*	*	*	*	*	*	0.2	0.4	1.1	3.4	9.2
2003.....	0.5	*	*	*	*	*	*	0.2	0.6	1.4	4.1	12.2
2002.....	0.5	*	*	*	*	*	*	0.3	0.4	1.4	4.6	12.9
2001.....	0.5	*	*	*	*	*	*	0.2	0.7	1.6	4.7	15.0
2000.....	0.6	*	*	*	*	*	*	0.2	0.9	1.9	5.4	18.4
1999.....	0.6	*	*	*	*	*	*	0.2	0.7	2.1	5.8	17.4

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Peptic ulcer, site unspecified (K27)												
White, female												
2004.....	0.5	*	*	*	*	*	*	*	0.3	0.9	2.7	10.3
2003.....	0.5	*	*	*	*	*	*	*	0.3	0.9	3.3	10.1
2002.....	0.6	*	*	*	*	*	*	0.2	0.3	0.9	3.0	10.9
2001.....	0.7	*	*	*	*	*	*	0.1	0.3	1.1	3.7	14.4
2000.....	0.7	*	*	*	*	*	*	*	0.2	1.0	3.5	15.0
1999.....	0.8	*	*	*	*	*	*	*	0.4	1.2	4.3	15.7
Black, both sexes												
2004.....	0.3	*	*	*	*	*	*	*	0.7	1.2	2.7	8.2
2003.....	0.3	*	*	*	*	*	*	*	*	*	3.7	5.8
2002.....	0.3	*	*	*	*	*	*	*	0.8	1.5	2.6	8.9
2001.....	0.4	*	*	*	*	*	*	*	1.0	1.4	4.0	9.5
2000.....	0.4	*	*	*	*	*	*	*	*	1.2	3.7	11.9
1999.....	0.4	*	*	*	*	*	*	*	0.9	2.2	4.4	9.6
Black, male												
2004.....	0.3	*	*	*	*	*	*	*	*	*	*	*
2003.....	0.4	*	*	*	*	*	*	*	*	*	7.3	*
2002.....	0.4	*	*	*	*	*	*	*	*	*	*	*
2001.....	0.5	*	*	*	*	*	*	*	1.9	*	5.9	*
2000.....	0.3	*	*	*	*	*	*	*	*	*	*	*
1999.....	0.4	*	*	*	*	*	*	*	*	3.2	*	*
Black, female												
2004.....	0.3	*	*	*	*	*	*	*	*	*	*	8.3
2003.....	0.2	*	*	*	*	*	*	*	*	*	*	*
2002.....	0.3	*	*	*	*	*	*	*	*	*	*	8.2
2001.....	0.3	*	*	*	*	*	*	*	*	*	*	8.4
2000.....	0.4	*	*	*	*	*	*	*	*	*	3.6	11.2
1999.....	0.4	*	*	*	*	*	*	*	*	*	4.1	10.5

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Gastritis and duodenitis (K29)												
All races 3/, both sexes												
2004.....	0.1	*	*	*	*	*	*	0.1	0.2	0.2	0.8	2.6
2003.....	0.1	*	*	*	*	*	*	0.1	0.1	0.3	0.7	2.9
2002.....	0.1	*	*	*	*	*	*	0.1	0.1	0.2	0.8	2.9
2001.....	0.1	*	*	*	*	*	*	*	0.1	0.3	0.7	3.8
2000.....	0.1	*	*	*	*	*	*	0.1	0.2	0.3	0.8	3.5
1999.....	0.1	*	*	*	*	*	0.0	0.1	0.1	0.3	0.9	3.8
All races 3/, male												
2004.....	0.1	*	*	*	*	*	*	*	0.3	0.3	0.9	3.0
2003.....	0.1	*	*	*	*	*	*	0.2	0.2	0.3	0.9	2.8
2002.....	0.1	*	*	*	*	*	*	*	0.2	0.3	0.9	2.9
2001.....	0.1	*	*	*	*	*	*	*	*	0.3	0.7	4.8
2000.....	0.1	*	*	*	*	*	*	*	0.3	0.4	0.8	3.3
1999.....	0.1	*	*	*	*	*	*	0.2	*	0.3	1.1	3.7
All races 3/, female												
2004.....	0.1	*	*	*	*	*	*	*	*	*	0.7	2.4
2003.....	0.1	*	*	*	*	*	*	*	*	0.2	0.6	3.0
2002.....	0.1	*	*	*	*	*	*	*	*	*	0.7	2.9
2001.....	0.1	*	*	*	*	*	*	*	*	0.3	0.7	3.3
2000.....	0.2	*	*	*	*	*	*	*	*	0.2	0.9	3.6
1999.....	0.2	*	*	*	*	*	*	*	*	0.3	0.8	3.9
White, both sexes												
2004.....	0.1	*	*	*	*	*	*	*	0.2	0.2	0.7	2.6
2003.....	0.1	*	*	*	*	*	*	0.1	0.1	0.3	0.7	2.9
2002.....	0.1	*	*	*	*	*	*	0.1	0.1	0.2	0.8	2.9
2001.....	0.1	*	*	*	*	*	*	*	0.1	0.3	0.7	4.0
2000.....	0.2	*	*	*	*	*	*	*	0.2	0.3	0.9	3.5
1999.....	0.1	*	*	*	*	*	*	0.1	0.1	0.2	0.8	3.9
White, male												
2004.....	0.1	*	*	*	*	*	*	*	0.2	0.3	0.8	3.1
2003.....	0.1	*	*	*	*	*	*	0.2	0.2	*	0.8	2.8
2002.....	0.1	*	*	*	*	*	*	*	0.2	0.3	1.0	2.9
2001.....	0.1	*	*	*	*	*	*	*	*	*	0.7	5.0
2000.....	0.1	*	*	*	*	*	*	*	0.2	0.4	0.8	3.3
1999.....	0.1	*	*	*	*	*	*	0.2	*	*	1.0	3.8

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Dyspepsia and other diseases of stomach and duodenum (K30-K31)

All races 3/, both sexes

2004.....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.5	1.4	3.0
2003.....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.6	1.4	3.5
2002.....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.5	1.4	3.2
2001.....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.5	1.3	3.6
2000.....	0.2	*	*	*	*	*	*	0.1	0.2	0.6	1.3	4.0
1999.....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.5	1.4	4.2

All races 3/, male

2004.....	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.5	3.6
2003.....	0.2	*	*	*	*	*	*	*	0.1	0.6	1.4	2.7
2002.....	0.2	*	*	*	*	*	*	*	0.2	0.6	1.4	2.9
2001.....	0.2	*	*	*	*	*	*	*	0.2	0.5	1.4	3.3
2000.....	0.2	*	*	*	*	*	*	*	0.2	0.8	1.4	4.3
1999.....	0.2	*	*	*	*	*	*	*	0.2	0.5	1.4	4.8

All races 3/, female

2004.....	0.2	*	*	*	*	*	0.1	0.2	0.3	0.4	1.3	2.8
2003.....	0.3	*	*	*	*	*	*	0.1	0.2	0.5	1.3	3.9
2002.....	0.2	*	*	*	*	*	0.1	0.1	0.3	0.4	1.4	3.4
2001.....	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.2	3.7
2000.....	0.2	*	*	*	*	*	*	0.1	0.3	0.4	1.3	3.9
1999.....	0.2	*	*	*	*	*	*	*	0.2	0.4	1.4	4.0

White, both sexes

2004.....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.4	1.4	3.0
2003.....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.6	1.3	3.6
2002.....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.5	1.4	3.3
2001.....	0.2	*	*	*	*	*	*	0.1	0.1	0.5	1.2	3.7
2000.....	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.3	4.1
1999.....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.4	1.4	4.2

White, male

2004.....	0.2	*	*	*	*	*	*	*	0.2	0.4	1.6	3.5
2003.....	0.2	*	*	*	*	*	*	*	*	0.6	1.4	2.9
2002.....	0.2	*	*	*	*	*	*	*	0.2	0.6	1.4	2.8
2001.....	0.2	*	*	*	*	*	*	*	*	0.5	1.4	3.5
2000.....	0.2	*	*	*	*	*	*	*	*	0.7	1.4	4.4
1999.....	0.2	*	*	*	*	*	*	*	0.2	0.4	1.4	4.9

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diseases of appendix (K35-K38)												
All races 3/, both sexes												
2004.....	0.2	*	*	*	*	*	*	0.1	0.2	0.4	1.1	1.7
2003.....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.3	1.0	1.7
2002.....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.5	0.9	2.0
2001.....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.5	0.7	2.1
2000.....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.4	1.0	1.6
1999.....	0.1	*	*	0.1	*	*	0.1	0.1	0.2	0.4	0.8	1.5
All races 3/, male												
2004.....	0.2	*	*	*	*	*	*	0.2	0.2	0.5	1.3	2.5
2003.....	0.2	*	*	*	*	*	*	0.1	0.2	0.6	1.5	2.4
2002.....	0.2	*	*	*	*	*	*	0.2	0.4	0.7	1.3	2.2
2001.....	0.2	*	*	*	*	*	*	0.1	0.2	0.7	1.0	2.8
2000.....	0.2	*	*	*	*	*	*	0.2	0.4	0.5	1.2	2.0
1999.....	0.2	*	*	*	*	*	*	0.2	0.3	0.5	1.0	2.9
All races 3/, female												
2004.....	0.1	*	*	*	*	*	*	*	0.2	0.4	0.9	1.4
2003.....	0.1	*	*	*	*	*	*	0.1	*	*	0.7	1.3
2002.....	0.1	*	*	*	*	*	*	*	*	0.3	0.7	2.0
2001.....	0.1	*	*	*	*	*	*	*	*	0.3	0.5	1.8
2000.....	0.1	*	*	*	*	*	*	*	*	0.3	0.8	1.5
1999.....	0.1	*	*	*	*	*	*	*	*	0.3	0.7	0.9
White, both sexes												
2004.....	0.2	*	*	*	*	*	*	0.1	0.2	0.4	1.0	1.8
2003.....	0.2	*	*	*	*	*	*	0.1	0.1	0.3	1.0	1.7
2002.....	0.2	*	*	*	*	*	*	0.1	0.2	0.4	1.0	2.1
2001.....	0.1	*	*	*	*	*	*	0.1	0.1	0.5	0.7	2.1
2000.....	0.2	*	*	*	*	*	*	0.1	0.2	0.3	0.9	1.7
1999.....	0.1	*	*	*	*	*	*	0.1	0.1	0.4	0.8	1.5
White, male												
2004.....	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.3	2.3
2003.....	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.4	2.5
2002.....	0.2	*	*	*	*	*	*	0.2	0.4	0.6	1.4	2.0
2001.....	0.2	*	*	*	*	*	*	0.1	0.2	0.7	1.0	2.5
2000.....	0.2	*	*	*	*	*	*	*	0.3	0.4	1.2	2.0
1999.....	0.2	*	*	*	*	*	*	0.2	0.2	0.4	0.9	3.1

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hernia (K40-K46)												
All races 3/, both sexes												
2004.....	0.5	1.0	*	*	*	*	0.1	0.2	0.4	1.3	3.9	11.0
2003.....	0.6	0.9	*	*	*	*	0.1	0.3	0.6	1.3	3.6	11.9
2002.....	0.6	0.7	*	*	*	*	0.1	0.2	0.5	1.2	3.8	12.8
2001.....	0.5	1.1	*	*	*	*	0.1	0.2	0.5	1.2	3.5	12.6
2000.....	0.5	1.2	*	*	*	*	0.1	0.2	0.5	1.1	3.8	13.1
1999.....	0.5	1.2	*	*	*	*	0.1	0.2	0.4	1.2	3.7	12.7
All races 3/, male												
2004.....	0.5	1.0	*	*	*	*	0.1	0.2	0.4	1.4	3.8	11.3
2003.....	0.5	1.0	*	*	*	*	*	0.3	0.5	1.3	3.9	12.3
2002.....	0.4	*	*	*	*	*	*	0.2	0.4	1.2	3.7	11.7
2001.....	0.4	1.4	*	*	*	*	*	0.2	0.5	1.1	3.5	12.0
2000.....	0.4	1.4	*	*	*	*	*	0.2	0.4	1.1	4.0	12.6
1999.....	0.4	1.3	*	*	*	*	0.1	0.2	0.5	1.1	4.0	13.2
All races 3/, female												
2004.....	0.6	1.0	*	*	*	*	*	0.2	0.5	1.2	4.0	10.8
2003.....	0.6	*	*	*	*	*	*	0.2	0.6	1.3	3.4	11.8
2002.....	0.7	*	*	*	*	*	*	0.2	0.5	1.2	3.9	13.3
2001.....	0.6	*	*	*	*	*	*	0.2	0.5	1.2	3.6	12.8
2000.....	0.7	*	*	*	*	*	*	0.2	0.5	1.2	3.6	13.3
1999.....	0.6	*	*	*	*	*	*	0.1	0.4	1.2	3.5	12.6
White, both sexes												
2004.....	0.6	1.0	*	*	*	*	0.1	0.2	0.4	1.3	4.1	11.2
2003.....	0.6	0.9	*	*	*	*	0.1	0.3	0.5	1.3	3.6	12.3
2002.....	0.6	0.8	*	*	*	*	0.1	0.2	0.4	1.2	4.0	13.4
2001.....	0.6	1.0	*	*	*	*	0.1	0.2	0.5	1.2	3.5	12.8
2000.....	0.6	1.2	*	*	*	*	0.1	0.2	0.5	1.1	3.8	13.3
1999.....	0.6	1.1	*	*	*	*	0.1	0.1	0.5	1.1	3.7	13.0
White, male												
2004.....	0.5	*	*	*	*	*	*	0.2	0.4	1.4	3.9	11.4
2003.....	0.5	*	*	*	*	*	*	0.3	0.5	1.3	3.9	12.5
2002.....	0.5	*	*	*	*	*	*	0.2	0.4	1.2	3.9	11.9
2001.....	0.4	*	*	*	*	*	*	0.2	0.5	1.2	3.4	11.8
2000.....	0.5	1.5	*	*	*	*	*	0.2	0.4	1.1	4.1	12.7
1999.....	0.5	1.3	*	*	*	*	0.1	0.2	0.6	1.2	4.1	12.9

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Crohn's disease and ulcerative colitis (K50-K51)

All races 3/, both sexes

2004.....	0.3	*	*	*	*	0.1	0.1	0.3	0.5	0.9	2.1	3.3
2003.....	0.3	*	*	*	*	0.1	0.1	0.2	0.4	0.9	1.9	3.1
2002.....	0.3	*	*	*	*	*	0.2	0.2	0.3	0.9	2.0	3.1
2001.....	0.3	*	*	*	*	0.1	0.1	0.2	0.5	0.9	1.9	3.8
2000.....	0.3	*	*	*	*	0.1	0.1	0.2	0.4	0.9	1.9	3.3
1999.....	0.3	*	*	*	*	0.1	0.1	0.3	0.4	0.8	2.0	3.3

All races 3/, male

2004.....	0.3	*	*	*	*	*	0.1	0.2	0.5	0.9	2.0	2.7
2003.....	0.3	*	*	*	*	*	0.1	0.2	0.5	0.9	2.0	3.5
2002.....	0.2	*	*	*	*	*	0.1	0.2	0.3	0.9	1.9	2.7
2001.....	0.3	*	*	*	*	*	*	0.2	0.5	1.0	1.9	3.5
2000.....	0.2	*	*	*	*	*	*	0.2	0.3	0.8	1.6	2.9
1999.....	0.2	*	*	*	*	*	*	0.2	0.3	0.9	1.9	2.7

All races 3/, female

2004.....	0.4	*	*	*	*	*	0.1	0.3	0.5	0.9	2.1	3.6
2003.....	0.3	*	*	*	*	*	*	0.3	0.4	1.0	1.8	2.8
2002.....	0.3	*	*	*	*	*	0.2	0.3	0.4	0.8	2.0	3.2
2001.....	0.3	*	*	*	*	*	0.1	0.2	0.4	0.8	1.9	3.9
2000.....	0.3	*	*	*	*	*	0.1	0.2	0.4	0.9	2.0	3.4
1999.....	0.3	*	*	*	*	*	0.1	0.3	0.4	0.8	2.0	3.5

White, both sexes

2004.....	0.4	*	*	*	*	0.1	0.1	0.3	0.5	0.9	2.3	3.6
2003.....	0.3	*	*	*	*	0.1	0.1	0.3	0.5	1.0	2.0	3.3
2002.....	0.3	*	*	*	*	*	0.1	0.3	0.4	0.9	2.1	3.4
2001.....	0.3	*	*	*	*	*	0.1	0.2	0.5	1.0	2.0	3.9
2000.....	0.3	*	*	*	*	*	0.1	0.2	0.4	1.0	2.0	3.5
1999.....	0.3	*	*	*	*	*	0.1	0.3	0.4	0.9	2.1	3.5

White, male

2004.....	0.3	*	*	*	*	*	0.1	0.3	0.5	0.9	2.1	2.9
2003.....	0.3	*	*	*	*	*	*	0.2	0.5	0.9	2.1	3.7
2002.....	0.3	*	*	*	*	*	0.1	0.2	0.3	0.9	2.0	2.9
2001.....	0.3	*	*	*	*	*	*	0.2	0.5	1.1	2.0	3.5
2000.....	0.2	*	*	*	*	*	*	0.2	0.4	0.9	1.7	3.2
1999.....	0.3	*	*	*	*	*	*	0.2	0.4	1.0	2.0	3.0

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Vascular disorders and obstruction of intestine without hernia (K55-K56)

All races 3/, both sexes

2004.....	4.9	6.9	0.2	0.0	0.1	0.1	0.4	1.1	3.5	12.1	38.2	106.9
2003.....	5.1	6.8	0.1	0.1	0.1	0.1	0.4	1.2	3.7	13.3	39.2	111.5
2002.....	5.0	5.8	0.1	0.1	0.1	0.1	0.4	1.1	3.7	12.8	39.4	113.3
2001.....	5.0	5.8	0.1	0.1	*	0.1	0.3	1.2	3.9	13.8	38.9	113.5
2000.....	5.1	6.0	0.1	0.1	0.1	0.1	0.4	1.3	3.9	13.4	40.1	116.2
1999.....	5.0	6.1	0.1	0.1	0.1	0.1	0.4	1.1	4.0	13.5	38.9	116.2

All races 3/, male

2004.....	3.5	7.7	0.2	*	*	0.1	0.5	1.1	3.3	10.9	35.0	88.8
2003.....	3.7	8.2	*	0.1	0.1	0.1	0.4	1.2	3.7	12.2	35.6	92.6
2002.....	3.6	6.5	*	*	0.1	0.1	0.4	1.1	3.4	11.6	35.8	96.7
2001.....	3.6	6.5	*	*	*	0.1	0.3	1.1	3.5	13.1	35.2	98.5
2000.....	3.6	6.3	*	0.1	0.1	0.1	0.4	1.2	3.8	12.5	36.8	98.5
1999.....	3.5	7.6	*	*	*	0.1	0.4	1.1	3.8	12.5	35.6	95.6

All races 3/, female

2004.....	6.3	6.0	*	*	*	0.2	0.4	1.1	3.7	13.1	40.5	115.1
2003.....	6.5	5.3	*	*	*	0.1	0.4	1.3	3.7	14.2	41.6	119.8
2002.....	6.4	5.0	*	*	*	0.1	0.3	1.1	3.8	13.8	41.7	120.5
2001.....	6.4	5.1	*	*	*	0.1	0.3	1.2	4.1	14.4	41.3	119.8
2000.....	6.5	5.8	*	*	*	0.1	0.4	1.3	4.0	14.0	42.1	123.4
1999.....	6.4	4.4	*	*	*	0.2	0.4	1.1	4.2	14.3	41.1	124.5

White, both sexes

2004.....	5.3	4.6	0.2	*	*	0.1	0.4	1.0	3.4	12.1	38.2	108.5
2003.....	5.5	4.8	*	*	0.1	0.1	0.3	1.1	3.4	13.2	39.4	113.7
2002.....	5.4	4.2	*	*	0.1	0.1	0.4	1.0	3.5	12.4	39.5	115.9
2001.....	5.4	4.1	*	*	*	0.1	0.3	1.0	3.5	13.6	38.9	116.2
2000.....	5.4	3.8	*	0.1	0.1	0.1	0.3	1.0	3.6	13.1	39.9	118.0
1999.....	5.4	4.2	*	*	0.1	0.1	0.3	1.0	3.9	13.0	38.7	118.0

White, male

2004.....	3.7	5.1	*	*	*	*	0.4	1.0	3.0	10.4	34.0	88.6
2003.....	3.8	6.1	*	*	0.1	0.1	0.3	1.0	3.2	11.6	34.8	92.9
2002.....	3.8	4.6	*	*	*	*	0.4	1.0	3.1	11.2	35.5	97.5
2001.....	3.8	4.4	*	*	*	*	0.3	1.0	3.2	12.6	34.8	98.9
2000.....	3.7	3.8	*	*	*	0.1	0.3	1.0	3.3	12.0	35.8	98.9
1999.....	3.7	5.5	*	*	*	*	0.3	0.9	3.6	11.9	35.1	94.8

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Vascular disorders and obstruction of intestine without hernia (K55-K56)

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
White, female												
2004.....	6.9	4.0	*	*	*	*	0.3	1.0	3.6	13.5	41.2	117.5
2003.....	7.2	3.4	*	*	*	*	0.3	1.2	3.6	14.5	42.4	122.9
2002.....	7.1	3.8	*	*	*	*	0.3	1.0	3.8	13.5	42.3	123.9
2001.....	7.0	3.8	*	*	*	*	0.3	1.1	3.8	14.4	41.6	123.4
2000.....	7.1	3.9	*	*	*	*	0.3	1.1	3.8	14.0	42.5	125.8
1999.....	7.0	2.9	*	*	*	0.2	0.3	1.0	4.2	13.9	41.2	127.4
Black, both sexes												
2004.....	4.1	18.8	*	*	*	0.4	0.8	2.0	6.0	15.1	45.4	106.0
2003.....	4.1	18.0	*	*	*	*	0.8	2.4	6.6	17.1	44.3	102.1
2002.....	4.0	14.1	*	*	*	*	0.7	2.4	5.8	18.6	44.6	99.8
2001.....	4.1	15.8	*	*	*	*	0.6	2.8	7.4	19.2	45.4	96.7
2000.....	4.3	16.7	*	*	*	*	0.8	3.1	7.7	17.6	47.8	112.0
1999.....	4.2	15.5	*	*	*	*	0.9	2.4	5.7	20.5	47.1	108.7
Black, male												
2004.....	3.7	21.4	*	*	*	*	*	2.1	6.8	17.7	52.9	107.3
2003.....	3.8	20.5	*	*	*	*	*	2.9	8.2	21.0	51.3	101.1
2002.....	3.5	16.6	*	*	*	*	0.8	2.6	6.8	18.8	45.9	105.1
2001.....	3.7	18.6	*	*	*	*	0.8	2.8	7.1	22.3	48.0	120.3
2000.....	4.0	18.6	*	*	*	*	0.9	3.5	9.6	19.5	54.8	110.7
1999.....	3.5	17.9	*	*	*	*	1.0	2.7	6.2	19.6	49.8	110.1
Black, female												
2004.....	4.5	16.1	*	*	*	*	1.0	2.0	5.4	13.2	41.2	105.5
2003.....	4.4	15.5	*	*	*	*	1.0	2.0	5.3	14.3	40.3	102.5
2002.....	4.5	11.5	*	*	*	*	*	2.2	4.9	18.4	43.9	97.8
2001.....	4.5	12.9	*	*	*	*	*	2.7	7.6	17.1	44.0	88.0
2000.....	4.7	14.9	*	*	*	*	0.7	2.8	6.1	16.4	43.8	112.5
1999.....	4.8	13.1	*	*	*	*	0.8	2.1	5.3	21.2	45.6	108.2

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diverticular disease of intestine (K57)												
All races 3/, both sexes												
2004.....	1.1	*	*	*	*	*	0.1	0.2	0.8	2.8	9.3	27.0
2003.....	1.2	*	*	*	*	*	0.1	0.2	0.8	2.7	9.8	28.9
2002.....	1.2	*	*	*	*	*	0.1	0.2	0.7	2.7	9.9	27.7
2001.....	1.2	*	*	*	*	*	0.0	0.2	0.9	2.9	9.4	31.4
2000.....	1.2	*	*	*	*	*	0.1	0.2	0.7	3.0	9.2	30.9
1999.....	1.2	*	*	*	*	*	0.1	0.2	0.9	3.1	9.3	33.4
All races 3/, male												
2004.....	0.7	*	*	*	*	*	*	0.2	0.8	2.4	7.8	18.8
2003.....	0.8	*	*	*	*	*	*	0.2	0.8	2.4	7.6	22.3
2002.....	0.8	*	*	*	*	*	0.1	0.3	0.7	2.4	8.0	19.9
2001.....	0.8	*	*	*	*	*	*	0.3	0.8	2.4	8.2	23.4
2000.....	0.7	*	*	*	*	*	*	0.3	0.7	2.7	7.3	22.3
1999.....	0.8	*	*	*	*	*	*	0.2	0.8	2.7	7.4	25.8
All races 3/, female												
2004.....	1.5	*	*	*	*	*	*	0.2	0.8	3.1	10.3	30.6
2003.....	1.6	*	*	*	*	*	*	0.2	0.9	3.0	11.3	31.9
2002.....	1.5	*	*	*	*	*	*	0.2	0.7	2.9	11.1	31.0
2001.....	1.6	*	*	*	*	*	*	0.2	1.0	3.3	10.2	34.7
2000.....	1.6	*	*	*	*	*	*	0.2	0.7	3.1	10.5	34.4
1999.....	1.7	*	*	*	*	*	*	0.2	0.9	3.4	10.5	36.4
White, both sexes												
2004.....	1.3	*	*	*	*	*	0.1	0.2	0.8	2.8	9.7	27.9
2003.....	1.4	*	*	*	*	*	0.1	0.2	0.8	2.8	10.2	30.3
2002.....	1.3	*	*	*	*	*	0.1	0.3	0.7	2.7	10.1	28.5
2001.....	1.4	*	*	*	*	*	0.1	0.2	0.9	2.8	9.8	32.4
2000.....	1.3	*	*	*	*	*	0.1	0.2	0.7	3.0	9.5	32.3
1999.....	1.4	*	*	*	*	*	0.1	0.2	0.8	3.2	9.6	34.6
White, male												
2004.....	0.8	*	*	*	*	*	*	0.2	0.8	2.4	7.9	19.3
2003.....	0.9	*	*	*	*	*	*	0.2	0.8	2.4	7.8	23.4
2002.....	0.8	*	*	*	*	*	*	0.4	0.7	2.5	7.9	20.5
2001.....	0.9	*	*	*	*	*	*	0.3	0.9	2.4	8.4	24.3
2000.....	0.8	*	*	*	*	*	*	0.3	0.7	2.7	7.6	23.7
1999.....	0.9	*	*	*	*	*	*	0.2	0.8	2.8	7.6	27.0

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diverticular disease of intestine (K57)												
White, female												
2004.....	1.8	*	*	*	*	*	*	0.1	0.8	3.2	10.9	31.8
2003.....	1.8	*	*	*	*	*	*	0.2	0.8	3.1	11.7	33.3
2002.....	1.7	*	*	*	*	*	*	0.2	0.7	2.9	11.5	32.0
2001.....	1.8	*	*	*	*	*	*	0.2	0.9	3.2	10.8	35.8
2000.....	1.8	*	*	*	*	*	*	0.2	0.6	3.2	10.8	35.7
1999.....	1.9	*	*	*	*	*	*	0.3	0.9	3.4	10.9	37.7
Black, both sexes												
2004.....	0.6	*	*	*	*	*	*	*	*	3.0	7.2	21.7
2003.....	0.6	*	*	*	*	*	*	*	1.5	2.6	7.9	18.4
2002.....	0.7	*	*	*	*	*	*	*	0.9	2.5	9.7	21.7
2001.....	0.7	*	*	*	*	*	*	*	1.0	4.0	6.8	24.3
2000.....	0.7	*	*	*	*	*	*	*	1.3	3.7	7.2	20.0
1999.....	0.7	*	*	*	*	*	*	*	1.5	3.5	7.3	23.6
Black, male												
2004.....	0.5	*	*	*	*	*	*	*	*	*	7.8	*
2003.....	0.4	*	*	*	*	*	*	*	*	*	6.2	*
2002.....	0.5	*	*	*	*	*	*	*	*	*	11.2	*
2001.....	0.4	*	*	*	*	*	*	*	*	2.9	7.1	*
2000.....	0.4	*	*	*	*	*	*	*	*	3.7	*	*
1999.....	0.4	*	*	*	*	*	*	*	*	*	6.5	*
Black, female												
2004.....	0.8	*	*	*	*	*	*	*	*	3.3	6.9	23.2
2003.....	0.9	*	*	*	*	*	*	*	1.8	2.9	8.9	20.3
2002.....	0.8	*	*	*	*	*	*	*	*	3.0	8.9	24.6
2001.....	0.9	*	*	*	*	*	*	*	*	4.7	6.6	27.8
2000.....	0.9	*	*	*	*	*	*	*	*	3.7	8.5	24.5
1999.....	0.9	*	*	*	*	*	*	*	1.5	4.1	7.7	27.2

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other diseases of intestines and peritoneum (K52,K58-K63)												
All races 3/, both sexes												
2004.....	1.7	2.0	0.2	*	0.1	0.1	0.2	0.5	1.4	4.4	12.8	33.3
2003.....	1.7	2.4	0.1	*	0.1	0.1	0.2	0.6	1.6	4.3	12.6	32.8
2002.....	1.7	2.0	0.2	0.0	0.0	0.1	0.2	0.6	1.6	4.3	12.5	32.8
2001.....	1.7	1.4	0.2	*	*	0.1	0.2	0.6	1.4	4.5	12.8	34.1
2000.....	1.6	1.7	*	*	*	0.1	0.2	0.5	1.6	4.1	12.4	33.3
1999.....	1.6	1.5	0.2	*	*	0.1	0.2	0.5	1.5	4.4	12.1	32.0
All races 3/, male												
2004.....	1.3	2.3	*	*	*	*	0.1	0.6	1.2	4.1	12.5	26.4
2003.....	1.3	2.9	*	*	*	0.1	0.2	0.6	1.6	4.2	12.3	27.1
2002.....	1.3	2.4	*	*	*	*	0.2	0.6	1.5	4.4	11.8	29.4
2001.....	1.2	1.6	*	*	*	*	0.2	0.5	1.4	4.2	12.1	29.9
2000.....	1.2	1.8	*	*	*	*	0.2	0.4	1.7	4.0	11.4	28.7
1999.....	1.2	2.0	*	*	*	*	0.2	0.6	1.5	4.2	11.9	28.9
All races 3/, female												
2004.....	2.1	1.7	*	*	*	*	0.2	0.5	1.6	4.6	12.9	36.4
2003.....	2.1	1.8	*	*	*	*	0.2	0.5	1.5	4.5	12.8	35.2
2002.....	2.0	1.5	*	*	*	*	0.2	0.5	1.7	4.3	13.0	34.3
2001.....	2.1	1.1	*	*	*	*	0.2	0.6	1.5	4.7	13.3	35.8
2000.....	2.0	1.6	*	*	*	*	0.2	0.6	1.4	4.2	13.1	35.2
1999.....	1.9	*	*	*	*	*	0.2	0.5	1.6	4.6	12.2	33.2
White, both sexes												
2004.....	1.9	1.6	*	*	*	*	0.2	0.5	1.4	4.4	13.1	34.4
2003.....	1.8	2.1	*	*	*	0.1	0.2	0.5	1.5	4.3	12.9	33.8
2002.....	1.8	1.6	0.2	*	*	*	0.2	0.6	1.6	4.3	12.9	34.1
2001.....	1.8	0.8	*	*	*	*	0.2	0.5	1.3	4.6	12.8	35.2
2000.....	1.7	1.4	*	*	*	0.1	0.2	0.5	1.5	4.1	12.7	33.7
1999.....	1.7	1.2	0.2	*	*	0.1	0.2	0.5	1.5	4.5	12.2	33.0
White, male												
2004.....	1.3	1.8	*	*	*	*	0.1	0.5	1.2	3.9	12.6	26.9
2003.....	1.4	2.7	*	*	*	*	0.2	0.5	1.5	4.1	12.5	27.5
2002.....	1.4	2.2	*	*	*	*	0.2	0.6	1.5	4.2	11.9	30.3
2001.....	1.3	*	*	*	*	*	0.2	0.5	1.2	4.0	12.1	31.2
2000.....	1.3	1.4	*	*	*	*	0.2	0.4	1.5	4.1	11.4	28.1
1999.....	1.3	1.6	*	*	*	*	0.2	0.6	1.4	4.3	11.9	28.3

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other diseases of intestines and peritoneum (K52,K58-K63)

White, female

2004.....	2.4	1.4	*	*	*	*	0.2	0.5	1.6	4.8	13.5	37.8
2003.....	2.3	1.4	*	*	*	*	0.2	0.5	1.5	4.4	13.3	36.5
2002.....	2.3	*	*	*	*	*	0.2	0.5	1.7	4.4	13.5	35.8
2001.....	2.3	*	*	*	*	*	0.2	0.6	1.4	5.0	13.3	36.8
2000.....	2.2	1.5	*	*	*	*	0.2	0.6	1.4	4.2	13.5	36.1
1999.....	2.1	*	*	*	*	*	0.2	0.4	1.6	4.6	12.5	34.9

Black, both sexes

2004.....	1.2	3.6	*	*	*	*	*	0.7	2.2	5.3	11.5	24.0
2003.....	1.3	3.8	*	*	*	*	0.4	1.0	2.5	6.0	12.1	25.7
2002.....	1.2	3.9	*	*	*	*	0.4	0.7	2.4	5.3	10.9	23.2
2001.....	1.3	3.8	*	*	*	*	0.4	0.9	2.4	5.0	15.7	27.0
2000.....	1.2	3.7	*	*	*	*	0.4	0.8	2.5	4.7	10.7	29.7
1999.....	1.1	*	*	*	*	*	0.4	1.0	1.9	5.2	11.7	23.6

Black, male

2004.....	1.1	*	*	*	*	*	*	1.0	2.1	6.7	13.4	24.8
2003.....	1.2	*	*	*	*	*	*	1.1	3.0	6.2	13.5	25.0
2002.....	1.1	*	*	*	*	*	*	*	2.5	7.4	12.6	24.7
2001.....	1.2	*	*	*	*	*	*	*	3.0	7.1	16.2	*
2000.....	1.0	*	*	*	*	*	*	*	3.1	4.0	13.3	30.0
1999.....	1.1	*	*	*	*	*	*	1.2	1.9	4.7	12.9	39.8

Black, female

2004.....	1.2	*	*	*	*	*	*	*	2.2	4.4	10.4	23.6
2003.....	1.4	*	*	*	*	*	*	1.0	2.2	5.8	11.3	25.9
2002.....	1.2	*	*	*	*	*	*	*	2.3	3.7	9.9	22.6
2001.....	1.4	*	*	*	*	*	*	1.1	2.0	3.5	15.4	29.5
2000.....	1.3	*	*	*	*	*	*	0.9	2.1	5.1	9.2	29.6
1999.....	1.2	*	*	*	*	*	*	*	1.9	5.6	11.0	17.5

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diseases of peritoneum (K65-K66)												
All races 3/, both sexes												
2004.....	0.6	*	*	*	*	0.0	0.2	0.4	0.7	1.7	4.1	8.6
2003.....	0.6	*	*	*	*	0.1	0.2	0.5	0.8	1.9	4.1	9.2
2002.....	0.7	*	*	*	*	0.1	0.2	0.5	0.8	1.8	4.8	10.3
2001.....	0.7	*	*	*	*	*	0.2	0.5	0.8	1.8	4.4	10.4
2000.....	0.7	*	*	*	*	0.1	0.2	0.4	1.0	2.0	4.4	10.6
1999.....	0.7	0.5	*	*	*	0.1	0.2	0.5	0.9	2.2	4.4	9.7
All races 3/, male												
2004.....	0.5	*	*	*	*	*	0.2	0.5	0.8	2.0	4.4	7.7
2003.....	0.6	*	*	*	*	*	0.2	0.5	0.9	2.0	4.5	8.7
2002.....	0.5	*	*	*	*	*	0.2	0.5	0.9	1.6	4.3	9.1
2001.....	0.6	*	*	*	*	*	0.2	0.5	1.0	2.0	4.6	10.1
2000.....	0.6	*	*	*	*	*	0.2	0.5	1.1	2.0	4.6	11.0
1999.....	0.6	*	*	*	*	*	0.2	0.6	1.0	2.4	4.9	10.1
All races 3/, female												
2004.....	0.7	*	*	*	*	*	0.2	0.3	0.7	1.5	3.9	9.0
2003.....	0.7	*	*	*	*	*	0.1	0.4	0.7	1.9	3.9	9.4
2002.....	0.8	*	*	*	*	*	0.2	0.4	0.8	1.9	5.1	10.8
2001.....	0.7	*	*	*	*	*	0.1	0.4	0.7	1.7	4.3	10.5
2000.....	0.8	*	*	*	*	*	0.2	0.4	0.9	2.0	4.3	10.5
1999.....	0.7	*	*	*	*	*	0.2	0.3	0.8	2.0	4.1	9.6
White, both sexes												
2004.....	0.6	*	*	*	*	*	0.1	0.4	0.7	1.6	4.1	8.8
2003.....	0.7	*	*	*	*	0.1	0.2	0.4	0.7	1.9	4.2	9.1
2002.....	0.7	*	*	*	*	*	0.2	0.4	0.8	1.7	4.8	10.5
2001.....	0.7	*	*	*	*	*	0.1	0.4	0.7	1.7	4.3	10.4
2000.....	0.7	*	*	*	*	*	0.2	0.4	0.9	1.9	4.3	10.6
1999.....	0.7	*	*	*	*	0.1	0.2	0.4	0.8	2.0	4.4	9.6
White, male												
2004.....	0.6	*	*	*	*	*	0.2	0.4	0.7	1.9	4.4	7.9
2003.....	0.6	*	*	*	*	*	0.2	0.4	0.8	1.9	4.6	8.8
2002.....	0.5	*	*	*	*	*	0.1	0.5	0.8	1.6	4.1	9.3
2001.....	0.6	*	*	*	*	*	0.2	0.5	0.8	1.9	4.6	9.7
2000.....	0.6	*	*	*	*	*	0.2	0.4	1.0	1.9	4.5	10.8
1999.....	0.6	*	*	*	*	*	0.2	0.5	0.9	2.2	4.8	10.2

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diseases of peritoneum (K65-K66)												
White, female												
2004.....	0.7	*	*	*	*	*	0.1	0.3	0.7	1.5	4.0	9.2
2003.....	0.7	*	*	*	*	*	0.1	0.3	0.6	1.8	3.9	9.3
2002.....	0.8	*	*	*	*	*	0.2	0.3	0.7	1.7	5.2	11.0
2001.....	0.7	*	*	*	*	*	*	0.3	0.6	1.6	4.2	10.7
2000.....	0.8	*	*	*	*	*	0.2	0.3	0.8	2.0	4.2	10.5
1999.....	0.7	*	*	*	*	*	0.2	0.3	0.7	1.8	4.1	9.3
Black, both sexes												
2004.....	0.6	*	*	*	*	*	0.4	0.7	1.2	2.7	5.1	7.9
2003.....	0.7	*	*	*	*	*	*	1.1	1.9	2.7	4.4	12.0
2002.....	0.7	*	*	*	*	*	*	1.1	1.5	2.8	5.4	10.4
2001.....	0.8	*	*	*	*	*	0.5	1.1	2.0	2.9	6.0	11.7
2000.....	0.8	*	*	*	*	*	0.5	1.1	2.1	3.1	6.3	11.6
1999.....	0.8	*	*	*	*	*	*	1.0	2.1	3.8	6.0	12.1
Black, male												
2004.....	0.5	*	*	*	*	*	*	*	*	3.0	*	*
2003.....	0.6	*	*	*	*	*	*	0.9	1.9	2.8	*	*
2002.....	0.6	*	*	*	*	*	*	1.2	*	*	6.0	*
2001.....	0.7	*	*	*	*	*	*	1.1	2.4	3.3	*	*
2000.....	0.8	*	*	*	*	*	*	1.3	2.4	3.8	7.6	*
1999.....	0.7	*	*	*	*	*	*	1.3	2.5	4.1	7.1	*
Black, female												
2004.....	0.7	*	*	*	*	*	*	*	*	2.5	5.2	7.9
2003.....	0.8	*	*	*	*	*	*	1.2	1.9	2.6	3.8	12.6
2002.....	0.8	*	*	*	*	*	*	1.0	1.4	3.2	5.0	11.1
2001.....	0.8	*	*	*	*	*	*	1.1	1.7	2.6	6.3	10.5
2000.....	0.8	*	*	*	*	*	*	0.9	1.9	2.6	5.6	10.7
1999.....	0.8	*	*	*	*	*	*	*	1.7	3.6	5.3	12.7

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diseases of liver (K70-K76)												
All races 3/, both sexes												
2004.....	12.3	2.5	0.2	*	0.2	1.2	8.1	23.0	29.4	36.9	40.0	32.8
2003.....	12.6	1.5	0.2	*	0.2	1.4	8.7	23.4	30.0	38.7	41.0	33.6
2002.....	12.4	2.2	0.2	0.1	0.2	1.3	8.9	23.0	29.1	38.1	41.9	35.7
2001.....	12.2	1.9	0.2	0.1	0.2	1.3	9.0	22.9	28.8	38.1	41.4	36.1
2000.....	12.0	2.0	0.2	0.1	0.2	1.4	9.2	21.8	29.3	37.9	41.2	36.7
1999.....	12.0	2.3	0.1	0.1	0.2	1.4	9.0	21.4	29.4	39.4	42.2	37.2
All races 3/, male												
2004.....	15.7	2.5	*	*	0.2	1.6	10.8	33.3	41.7	47.5	49.5	40.7
2003.....	16.1	2.2	0.2	*	0.2	1.7	11.4	34.6	42.9	50.3	49.6	44.4
2002.....	15.8	3.1	*	*	0.2	1.6	11.6	33.9	41.2	49.5	51.7	47.3
2001.....	15.7	2.3	*	*	0.2	1.6	11.9	33.7	40.9	50.2	50.9	49.4
2000.....	15.5	2.7	*	*	0.2	1.8	12.5	32.5	41.3	49.7	51.0	52.2
1999.....	15.6	2.9	*	*	0.2	1.8	12.5	32.4	42.1	52.5	52.0	52.3
All races 3/, female												
2004.....	9.0	2.4	*	*	0.2	0.8	5.5	13.0	18.1	27.9	33.7	29.3
2003.....	9.1	*	*	*	0.2	1.0	6.1	12.7	18.1	29.0	35.2	28.9
2002.....	9.0	1.3	*	*	0.2	1.1	6.2	12.5	17.9	28.5	35.4	30.7
2001.....	8.9	1.6	*	*	0.2	1.0	6.2	12.5	17.5	28.1	35.1	30.5
2000.....	8.7	1.3	*	*	0.2	1.0	5.9	11.4	18.3	28.3	34.7	30.4
1999.....	8.6	1.6	*	*	0.2	1.1	5.6	10.7	17.7	28.8	35.8	31.1
White, both sexes												
2004.....	13.0	1.9	0.2	*	0.2	1.2	8.2	23.1	29.2	37.9	41.5	33.4
2003.....	13.3	1.2	0.2	*	0.1	1.3	9.0	23.4	29.9	39.9	42.7	34.3
2002.....	13.1	1.9	*	*	0.2	1.3	9.1	22.8	29.0	39.3	43.3	36.6
2001.....	12.9	1.6	0.2	0.1	0.1	1.3	9.0	22.5	28.5	39.0	42.6	37.0
2000.....	12.6	1.6	*	*	0.2	1.3	9.0	21.2	28.9	38.7	42.4	37.5
1999.....	12.5	1.7	*	0.1	0.2	1.4	8.8	20.6	28.6	40.0	43.3	37.7
White, male												
2004.....	16.6	1.8	*	*	0.2	1.6	11.0	33.6	41.2	48.6	51.2	42.6
2003.....	17.1	1.8	*	*	0.1	1.7	11.9	34.5	42.4	51.5	51.5	45.1
2002.....	16.8	2.7	*	*	0.2	1.6	12.0	33.8	41.0	50.9	53.3	48.5
2001.....	16.4	1.9	*	*	0.1	1.6	11.8	33.2	40.2	51.1	52.1	50.8
2000.....	16.2	2.0	*	*	0.2	1.7	12.4	31.9	40.9	50.5	52.6	53.3
1999.....	16.3	2.0	*	*	0.2	1.7	12.3	31.4	41.2	53.0	53.6	53.8

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diseases of liver (K70-K76)												
White, female												
2004.....	9.5	1.9	*	*	0.1	0.8	5.4	12.8	17.9	28.8	34.9	29.3
2003.....	9.7	*	*	*	0.2	0.9	6.1	12.3	18.0	30.0	36.8	29.5
2002.....	9.5	*	*	*	0.2	0.9	6.2	11.9	17.7	29.5	36.5	31.4
2001.....	9.4	1.3	*	*	*	1.0	6.2	11.9	17.5	28.7	36.3	31.2
2000.....	9.1	*	*	*	0.1	0.9	5.6	10.7	17.7	28.7	35.6	31.1
1999.....	8.9	1.4	*	*	0.2	0.9	5.2	10.0	16.8	29.1	36.6	31.2
Black, both sexes												
2004.....	9.8	5.4	*	*	0.4	1.0	7.9	25.8	35.4	32.4	25.2	24.3
2003.....	10.0	3.0	*	*	0.5	1.6	7.9	27.4	35.7	34.8	24.1	27.1
2002.....	9.9	3.7	*	*	0.4	1.5	7.9	28.5	33.9	31.6	29.6	27.6
2001.....	10.4	3.4	*	*	0.4	1.4	9.7	29.5	35.5	35.1	31.4	21.8
2000.....	10.4	4.2	*	*	0.4	1.8	10.5	29.4	35.2	36.2	28.8	26.0
1999.....	10.8	5.0	*	*	0.4	1.7	10.8	29.8	39.0	37.4	32.0	31.2
Black, male												
2004.....	12.9	*	*	*	*	1.3	10.2	37.3	54.1	47.1	33.4	19.9
2003.....	13.3	*	*	*	*	1.8	9.1	40.9	55.4	49.8	32.4	40.7
2002.....	12.6	*	*	*	*	1.6	9.1	40.5	50.8	44.8	37.6	36.5
2001.....	14.0	*	*	*	*	1.7	13.2	44.3	54.1	51.1	41.8	27.2
2000.....	13.5	6.4	*	*	*	1.9	13.6	42.9	50.6	50.8	36.9	35.8
1999.....	14.1	7.2	*	*	*	1.7	14.2	45.3	56.1	53.5	39.0	37.5
Black, female												
2004.....	7.0	*	*	*	*	0.8	5.8	15.9	20.4	21.9	20.6	26.0
2003.....	7.1	*	*	*	0.6	1.5	6.8	15.8	19.9	24.1	19.5	21.9
2002.....	7.4	*	*	*	*	1.4	6.8	18.1	20.4	22.3	25.0	24.2
2001.....	7.2	*	*	*	*	1.2	6.6	16.8	20.6	23.8	25.6	19.8
2000.....	7.7	*	*	*	*	1.7	7.7	17.8	22.7	25.9	24.2	22.3
1999.....	7.9	*	*	*	*	1.7	7.8	16.6	25.3	26.0	28.1	28.9

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Alcoholic liver disease (K70)												
All races 3/, both sexes												
2004.....	4.3	*	*	*	*	0.6	4.2	10.6	11.7	10.0	5.5	2.1
2003.....	4.3	*	*	*	*	0.6	4.4	10.5	11.6	9.8	5.7	1.8
2002.....	4.2	*	*	*	*	0.6	4.4	10.3	11.6	10.1	5.7	2.1
2001.....	4.3	*	*	*	*	0.7	4.8	10.7	11.5	10.1	5.7	2.4
2000.....	4.3	*	*	*	*	0.7	4.9	10.3	12.2	10.3	6.1	2.2
1999.....	4.3	*	*	*	*	0.6	4.8	10.4	12.0	10.5	6.5	2.8
All races 3/, male												
2004.....	6.3	*	*	*	*	0.8	5.8	15.5	18.2	16.5	10.1	4.6
2003.....	6.4	*	*	*	*	0.9	6.0	16.1	18.3	16.1	10.1	3.6
2002.....	6.2	*	*	*	*	0.8	5.8	15.6	18.3	16.5	9.7	4.7
2001.....	6.4	*	*	*	*	0.9	6.5	16.1	18.4	16.6	10.1	5.5
2000.....	6.5	*	*	*	*	1.0	6.8	16.0	19.0	17.3	11.5	5.3
1999.....	6.5	*	*	*	*	0.8	6.8	16.3	19.2	18.1	11.2	6.1
All races 3/, female												
2004.....	2.3	*	*	*	*	0.3	2.7	5.7	5.6	4.5	2.3	1.0
2003.....	2.2	*	*	*	*	0.3	2.9	5.1	5.4	4.5	2.7	1.0
2002.....	2.3	*	*	*	*	0.4	3.0	5.2	5.4	4.8	3.1	1.0
2001.....	2.3	*	*	*	*	0.4	3.1	5.5	5.1	4.7	2.8	1.0
2000.....	2.2	*	*	*	*	0.4	3.1	4.8	6.0	4.5	2.5	1.0
1999.....	2.1	*	*	*	*	0.5	2.9	4.7	5.4	4.3	3.4	1.5
White, both sexes												
2004.....	4.6	*	*	*	*	0.6	4.4	11.0	11.7	10.3	5.7	2.2
2003.....	4.5	*	*	*	*	0.6	4.6	10.7	11.7	10.0	5.9	1.9
2002.....	4.4	*	*	*	*	0.6	4.6	10.5	11.7	10.4	5.8	2.1
2001.....	4.5	*	*	*	*	0.7	4.8	10.8	11.5	10.2	5.8	2.3
2000.....	4.5	*	*	*	*	0.7	4.9	10.3	12.2	10.5	6.1	2.3
1999.....	4.4	*	*	*	*	0.6	4.8	10.2	11.8	10.5	6.4	2.7
White, male												
2004.....	6.8	*	*	*	*	0.9	6.0	16.2	18.3	16.8	10.4	4.8
2003.....	6.8	*	*	*	*	0.9	6.2	16.4	18.5	16.4	10.4	3.7
2002.....	6.6	*	*	*	*	0.8	6.1	16.0	18.5	16.8	9.9	4.5
2001.....	6.7	*	*	*	*	0.9	6.5	16.3	18.4	16.6	10.4	5.4
2000.....	6.8	*	*	*	*	1.0	6.8	16.2	19.2	17.5	11.7	5.4
1999.....	6.7	*	*	*	*	0.8	6.7	16.2	19.0	17.9	11.1	5.8

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Alcoholic liver disease (K70)												
White, female												
2004.....	2.4	*	*	*	*	0.3	2.8	5.8	5.6	4.7	2.4	1.0
2003.....	2.3	*	*	*	*	0.3	3.0	5.1	5.3	4.5	2.9	1.1
2002.....	2.3	*	*	*	*	0.3	3.0	5.1	5.2	5.0	3.1	1.1
2001.....	2.3	*	*	*	*	0.4	3.1	5.3	5.1	4.8	2.8	1.0
2000.....	2.1	*	*	*	*	0.3	3.0	4.4	5.6	4.6	2.4	1.0
1999.....	2.1	*	*	*	*	0.4	2.8	4.4	5.0	4.3	3.3	1.5
Black, both sexes												
2004.....	3.0	*	*	*	*	*	2.9	9.0	12.5	8.9	3.1	*
2003.....	3.2	*	*	*	*	*	3.3	9.9	12.5	9.3	3.4	*
2002.....	3.2	*	*	*	*	0.5	2.9	10.0	12.3	8.6	5.3	*
2001.....	3.6	*	*	*	*	0.5	4.2	11.5	12.9	10.7	4.7	*
2000.....	3.8	*	*	*	*	0.6	5.0	12.0	13.3	10.3	5.5	*
1999.....	4.1	*	*	*	*	0.5	5.0	12.8	15.1	11.8	7.1	*
Black, male												
2004.....	4.4	*	*	*	*	*	3.9	12.9	20.3	16.9	*	*
2003.....	4.8	*	*	*	*	*	4.1	15.8	20.1	16.0	6.8	*
2002.....	4.4	*	*	*	*	*	3.8	14.7	18.7	15.0	9.2	*
2001.....	5.3	*	*	*	*	*	5.9	17.1	21.4	19.5	7.7	*
2000.....	5.3	*	*	*	*	*	6.9	17.4	19.4	18.1	10.0	*
1999.....	5.9	*	*	*	*	*	6.7	19.7	24.2	21.8	12.6	*
Black, female												
2004.....	1.8	*	*	*	*	*	1.9	5.6	6.2	3.3	*	*
2003.....	1.8	*	*	*	*	*	2.5	4.7	6.4	4.5	*	*
2002.....	2.0	*	*	*	*	*	2.2	5.9	7.1	4.1	*	*
2001.....	2.1	*	*	*	*	*	2.7	6.7	6.1	4.4	*	*
2000.....	2.4	*	*	*	*	*	3.3	7.3	8.4	4.7	*	*
1999.....	2.4	*	*	*	*	*	3.4	6.9	7.9	4.8	4.0	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Chronic hepatitis, not elsewhere classified (K73)

All races 3/, both sexes

2004.....	0.0	*	*	*	*	*	*	0.1	*	0.1	0.2	*
2003.....	0.0	*	*	*	*	*	*	0.1	0.1	0.2	0.2	*
2002.....	0.2	*	*	*	*	*	0.1	0.2	0.3	0.7	1.0	0.8
2001.....	0.2	*	*	*	*	*	0.1	0.2	0.3	0.7	0.8	0.7
2000.....	0.2	*	*	*	*	*	0.1	0.1	0.3	0.7	0.8	0.8
1999.....	0.2	*	*	*	*	*	0.1	0.2	0.3	0.6	0.9	0.6

All races 3/, male

2004.....	0.0	*	*	*	*	*	*	0.1	*	*	*	*
2003.....	0.1	*	*	*	*	*	*	0.1	*	*	*	*
2002.....	0.1	*	*	*	*	*	*	0.2	*	0.4	0.6	*
2001.....	0.1	*	*	*	*	*	*	0.1	*	0.3	0.4	*
2000.....	0.1	*	*	*	*	*	*	0.1	*	*	*	*
1999.....	0.1	*	*	*	*	*	*	0.2	0.2	0.3	0.5	*

All races 3/, female

2004.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2003.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2002.....	0.3	*	*	*	*	*	0.1	0.2	0.5	1.0	1.3	0.9
2001.....	0.2	*	*	*	*	*	*	0.2	0.5	0.9	1.1	0.7
2000.....	0.2	*	*	*	*	*	*	0.2	0.5	1.0	1.1	0.9
1999.....	0.2	*	*	*	*	*	*	0.2	0.5	0.8	1.1	0.7

White, both sexes

2004.....	0.0	*	*	*	*	*	*	0.1	*	0.1	0.2	*
2003.....	0.1	*	*	*	*	*	*	0.1	0.1	0.2	0.2	*
2002.....	0.2	*	*	*	*	*	0.1	0.2	0.3	0.8	1.1	0.8
2001.....	0.2	*	*	*	*	*	0.1	0.2	0.3	0.7	0.8	0.8
2000.....	0.2	*	*	*	*	*	0.1	0.1	0.3	0.6	0.9	0.9
1999.....	0.2	*	*	*	*	*	0.1	0.2	0.3	0.5	0.8	0.6

White, male

2004.....	0.0	*	*	*	*	*	*	0.1	*	*	*	*
2003.....	0.1	*	*	*	*	*	*	0.1	*	*	*	*
2002.....	0.1	*	*	*	*	*	*	0.2	*	0.4	0.6	*
2001.....	0.1	*	*	*	*	*	*	0.1	*	0.4	0.5	*
2000.....	0.1	*	*	*	*	*	*	*	*	*	*	*
1999.....	0.1	*	*	*	*	*	*	0.2	0.2	0.3	0.5	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Fibrosis and cirrhosis of liver (K74)												
All races 3/, both sexes												
2004.....	4.9	*	*	*	*	0.2	2.1	7.4	10.9	17.6	23.1	17.4
2003.....	5.2	*	*	*	*	0.3	2.3	7.7	11.3	19.6	24.1	18.0
2002.....	5.1	*	*	*	*	0.3	2.5	7.5	11.0	18.6	24.7	18.5
2001.....	5.0	*	*	*	*	0.3	2.6	7.6	10.9	19.2	23.7	19.1
2000.....	5.0	*	*	*	0.1	0.3	2.5	7.2	11.3	18.9	24.1	20.0
1999.....	5.0	*	*	*	*	0.3	2.4	6.8	11.3	19.6	24.6	19.8
All races 3/, male												
2004.....	5.8	*	*	*	*	0.3	2.9	10.6	14.5	20.4	26.5	20.9
2003.....	6.1	*	*	*	*	0.3	3.2	11.3	15.4	22.8	26.7	23.7
2002.....	6.0	*	*	*	*	0.3	3.4	11.1	14.7	21.7	28.1	23.2
2001.....	6.0	*	*	*	*	0.4	3.3	11.3	14.6	23.2	26.2	25.7
2000.....	5.9	*	*	*	*	0.4	3.4	10.7	15.3	22.5	26.5	28.4
1999.....	5.9	*	*	*	*	0.4	3.5	10.4	15.2	23.3	27.9	26.6
All races 3/, female												
2004.....	4.0	*	*	*	*	0.1	1.3	4.2	7.5	15.3	20.9	15.8
2003.....	4.2	*	*	*	*	0.2	1.5	4.2	7.5	16.8	22.4	15.5
2002.....	4.2	*	*	*	*	0.2	1.7	4.1	7.6	16.0	22.4	16.5
2001.....	4.2	*	*	*	*	0.2	1.8	4.1	7.5	15.9	22.1	16.4
2000.....	4.1	*	*	*	*	0.2	1.5	3.8	7.6	15.9	22.5	16.6
1999.....	4.1	*	*	*	*	0.2	1.3	3.4	7.8	16.5	22.5	17.0
White, both sexes												
2004.....	5.3	*	*	*	*	0.2	2.1	7.4	10.9	18.5	24.4	17.9
2003.....	5.6	*	*	*	*	0.3	2.4	7.7	11.3	20.6	25.5	18.8
2002.....	5.5	*	*	*	*	0.3	2.7	7.4	11.1	19.5	25.9	19.4
2001.....	5.4	*	*	*	*	0.3	2.6	7.4	10.9	20.1	24.9	20.1
2000.....	5.4	*	*	*	*	0.3	2.5	7.1	11.3	19.5	25.3	20.8
1999.....	5.3	*	*	*	*	0.3	2.4	6.6	11.2	20.3	25.7	20.4
White, male												
2004.....	6.2	*	*	*	*	0.3	2.9	10.6	14.4	21.1	27.9	21.9
2003.....	6.6	*	*	*	*	0.3	3.3	11.3	15.2	24.0	27.9	24.5
2002.....	6.4	*	*	*	*	0.3	3.6	11.0	14.6	22.6	29.7	24.4
2001.....	6.3	*	*	*	*	0.3	3.3	10.8	14.3	24.1	27.5	27.3
2000.....	6.3	*	*	*	*	0.3	3.5	10.5	15.2	23.2	27.8	29.4
1999.....	6.3	*	*	*	*	0.5	3.5	10.0	14.9	24.2	29.2	27.9

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Fibrosis and cirrhosis of liver (K74)												
White, female												
2004.....	4.4	*	*	*	*	0.1	1.2	4.2	7.5	16.3	22.0	16.1
2003.....	4.7	*	*	*	*	0.2	1.5	4.1	7.7	17.7	23.8	16.3
2002.....	4.6	*	*	*	*	0.2	1.7	3.8	7.9	17.0	23.4	17.2
2001.....	4.6	*	*	*	*	0.2	1.9	3.9	7.7	16.7	23.1	17.1
2000.....	4.5	*	*	*	*	0.2	1.4	3.7	7.6	16.3	23.6	17.3
1999.....	4.4	*	*	*	*	0.2	1.3	3.2	7.7	17.1	23.5	17.5
Black, both sexes												
2004.....	3.5	*	*	*	*	*	2.5	9.3	12.8	12.6	11.3	11.3
2003.....	3.6	*	*	*	*	0.4	2.4	9.8	13.3	14.2	11.0	10.8
2002.....	3.5	*	*	*	*	0.4	2.3	10.4	12.5	12.5	12.8	9.2
2001.....	3.7	*	*	*	*	*	2.9	10.8	12.8	13.8	13.0	7.4
2000.....	3.6	*	*	*	*	*	2.7	9.9	12.4	15.3	12.2	11.6
1999.....	3.6	*	*	*	*	0.4	2.9	9.7	13.7	14.1	13.8	12.1
Black, male												
2004.....	4.6	*	*	*	*	*	3.4	14.0	18.7	16.9	15.3	*
2003.....	4.7	*	*	*	*	*	2.9	14.5	20.7	17.0	13.8	*
2002.....	4.5	*	*	*	*	*	2.9	14.9	19.1	17.3	12.9	*
2001.....	5.1	*	*	*	*	*	4.0	17.2	20.0	19.3	14.7	*
2000.....	4.6	*	*	*	*	*	3.4	15.0	18.5	19.5	14.5	*
1999.....	4.7	*	*	*	*	*	4.3	15.5	19.1	18.7	16.3	*
Black, female												
2004.....	2.5	*	*	*	*	*	1.7	5.1	8.0	9.5	9.0	11.0
2003.....	2.7	*	*	*	*	*	1.9	5.7	7.3	12.1	9.4	8.1
2002.....	2.7	*	*	*	*	*	1.9	6.6	7.1	9.1	12.7	9.0
2001.....	2.5	*	*	*	*	*	1.9	5.4	7.0	9.9	12.1	*
2000.....	2.7	*	*	*	*	*	2.1	5.5	7.5	12.4	10.9	10.3
1999.....	2.6	*	*	*	*	*	1.6	4.7	9.4	10.9	12.4	12.3

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other diseases of liver (K71-K72,K75-K76)												
All races 3/, both sexes												
2004.....	3.1	2.3	0.2	*	0.2	0.4	1.8	5.0	6.9	9.1	11.3	13.1
2003.....	3.1	1.5	0.2	*	0.2	0.5	1.9	5.1	7.0	9.2	11.0	13.5
2002.....	2.9	2.1	0.2	0.1	0.1	0.4	1.9	5.0	6.2	8.6	10.5	14.3
2001.....	2.7	1.9	0.2	0.1	0.1	0.4	1.6	4.4	6.0	8.1	11.1	13.8
2000.....	2.6	1.9	0.2	0.1	0.1	0.4	1.7	4.1	5.5	8.1	10.2	13.6
1999.....	2.6	2.1	0.1	0.1	0.1	0.4	1.7	4.0	5.7	8.8	10.3	14.0
All races 3/, male												
2004.....	3.5	2.3	*	*	0.2	0.5	2.1	6.9	8.9	10.5	12.8	15.1
2003.....	3.6	2.1	0.2	*	0.1	0.5	2.2	7.1	9.1	11.2	12.7	16.9
2002.....	3.5	2.8	*	*	0.1	0.5	2.3	7.1	8.1	11.0	13.3	18.9
2001.....	3.2	2.2	*	*	0.1	0.3	2.1	6.2	7.8	10.0	14.1	17.4
2000.....	3.0	2.5	*	*	0.1	0.4	2.2	5.6	6.9	9.7	12.6	17.8
1999.....	3.1	2.7	*	*	0.1	0.5	2.1	5.5	7.5	10.8	12.5	19.2
All races 3/, female												
2004.....	2.6	2.3	*	*	0.2	0.4	1.5	3.0	4.9	8.0	10.2	12.3
2003.....	2.6	*	*	*	0.2	0.5	1.7	3.3	5.0	7.6	9.8	12.1
2002.....	2.3	1.3	*	*	0.1	0.4	1.5	3.0	4.4	6.6	8.6	12.3
2001.....	2.3	1.5	*	*	0.1	0.4	1.2	2.7	4.4	6.6	9.2	12.3
2000.....	2.2	1.3	*	*	*	0.3	1.2	2.7	4.3	6.8	8.6	11.9
1999.....	2.2	1.5	*	*	0.2	0.3	1.3	2.5	4.0	7.2	8.8	11.9
White, both sexes												
2004.....	3.1	1.6	0.2	*	0.1	0.4	1.7	4.7	6.5	9.0	11.3	13.1
2003.....	3.2	1.2	0.2	*	0.1	0.4	1.9	4.9	6.7	9.1	11.1	13.4
2002.....	3.0	1.8	*	*	0.1	0.4	1.8	4.7	5.9	8.6	10.5	14.3
2001.....	2.8	1.5	0.2	*	0.1	0.3	1.5	4.2	5.8	8.1	11.0	13.8
2000.....	2.6	1.6	*	*	0.1	0.3	1.6	3.8	5.1	8.1	10.1	13.6
1999.....	2.6	1.5	*	*	0.1	0.3	1.5	3.6	5.3	8.6	10.3	13.9
White, male												
2004.....	3.6	1.6	*	*	0.1	0.5	2.0	6.7	8.4	10.4	12.8	15.9
2003.....	3.7	1.7	*	*	*	0.5	2.2	6.8	8.6	10.9	13.0	16.7
2002.....	3.6	2.4	*	*	*	0.4	2.3	6.7	7.7	11.1	13.1	18.9
2001.....	3.3	1.9	*	*	*	0.3	1.9	5.9	7.4	10.0	13.7	17.3
2000.....	3.0	2.0	*	*	*	0.3	2.1	5.1	6.4	9.6	12.7	17.8
1999.....	3.1	1.7	*	*	*	0.4	2.0	5.1	7.1	10.7	12.8	19.6

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other diseases of liver (K71-K72,K75-K76)

White, female

2004.....	2.6	1.7	*	*	*	0.3	1.4	2.8	4.8	7.8	10.3	11.9
2003.....	2.7	*	*	*	0.1	0.4	1.6	3.1	5.0	7.6	9.8	11.9
2002.....	2.4	*	*	*	*	0.3	1.4	2.8	4.2	6.5	8.7	12.3
2001.....	2.3	*	*	*	*	0.4	1.1	2.5	4.2	6.4	9.2	12.4
2000.....	2.2	*	*	*	*	0.3	1.1	2.4	4.0	6.8	8.4	11.9
1999.....	2.2	1.4	*	*	*	0.2	1.1	2.2	3.7	6.9	8.7	11.6

Black, both sexes

2004.....	3.2	5.1	*	*	0.3	0.7	2.4	7.5	10.2	10.8	10.9	12.1
2003.....	3.2	*	*	*	0.5	0.9	2.2	7.6	9.9	11.1	9.8	14.9
2002.....	3.0	3.6	*	*	*	0.6	2.4	7.7	8.9	10.0	11.1	15.4
2001.....	3.0	3.4	*	*	*	0.6	2.4	6.9	9.6	10.1	12.9	11.4
2000.....	3.0	3.9	*	*	*	0.7	2.6	7.3	9.3	10.0	10.8	12.2
1999.....	3.0	5.0	*	*	*	0.7	2.8	7.1	9.7	10.8	10.5	14.0

Black, male

2004.....	3.8	*	*	*	*	0.8	2.8	10.2	15.1	13.3	12.8	*
2003.....	3.8	*	*	*	*	0.8	2.1	10.5	14.5	16.5	11.8	*
2002.....	3.6	*	*	*	*	*	2.4	10.5	13.0	12.4	15.2	*
2001.....	3.6	*	*	*	*	*	3.1	9.8	12.6	12.0	19.4	*
2000.....	3.6	*	*	*	*	0.8	3.1	10.2	12.7	13.0	12.4	*
1999.....	3.4	7.2	*	*	*	*	3.1	9.9	12.7	12.8	9.5	*

Black, female

2004.....	2.7	*	*	*	*	*	2.1	5.2	6.2	9.0	9.8	14.6
2003.....	2.6	*	*	*	*	0.9	2.4	5.2	6.2	7.2	8.6	13.4
2002.....	2.5	*	*	*	*	*	2.4	5.2	5.7	8.3	8.8	13.6
2001.....	2.4	*	*	*	*	*	1.8	4.4	7.2	8.8	9.3	10.1
2000.....	2.4	*	*	*	*	*	2.2	4.7	6.5	7.8	9.9	10.7
1999.....	2.7	*	*	*	*	0.8	2.5	4.7	7.3	9.3	11.0	13.1

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Cholelithiasis and other disorders of gallbladder (K80-K82)												
All races 3/, both sexes												
2004.....	1.1	*	*	*	*	*	0.1	0.3	0.8	2.5	7.8	23.8
2003.....	1.0	*	*	*	*	0.1	0.1	0.3	0.7	2.7	8.2	21.8
2002.....	1.0	*	*	*	*	0.1	0.1	0.3	0.8	2.8	7.8	22.9
2001.....	1.0	*	*	*	*	*	0.1	0.3	1.0	2.9	7.5	23.7
2000.....	1.0	*	*	*	*	*	0.1	0.3	0.8	2.4	7.5	24.8
1999.....	1.0	*	*	*	*	*	0.1	0.3	0.9	2.6	7.8	23.9
All races 3/, male												
2004.....	0.9	*	*	*	*	*	0.1	0.4	1.0	2.9	9.3	25.5
2003.....	0.9	*	*	*	*	*	0.1	0.3	0.8	3.2	10.1	22.7
2002.....	0.9	*	*	*	*	*	0.1	0.4	1.0	3.1	9.8	24.1
2001.....	1.0	*	*	*	*	*	0.1	0.4	1.2	3.2	9.3	26.9
2000.....	0.9	*	*	*	*	*	0.1	0.3	0.9	2.9	9.4	27.6
1999.....	0.9	*	*	*	*	*	0.1	0.3	1.1	3.0	9.3	27.7
All races 3/, female												
2004.....	1.2	*	*	*	*	*	0.1	0.3	0.7	2.1	6.8	23.0
2003.....	1.1	*	*	*	*	*	*	0.3	0.5	2.2	6.9	21.4
2002.....	1.1	*	*	*	*	*	0.1	0.2	0.7	2.5	6.4	22.4
2001.....	1.1	*	*	*	*	*	*	0.3	0.8	2.6	6.3	22.4
2000.....	1.1	*	*	*	*	*	0.1	0.3	0.6	2.1	6.2	23.7
1999.....	1.1	*	*	*	*	*	0.1	0.4	0.7	2.2	6.8	22.4
White, both sexes												
2004.....	1.1	*	*	*	*	*	0.1	0.3	0.8	2.3	7.7	24.2
2003.....	1.1	*	*	*	*	*	0.1	0.3	0.6	2.7	8.2	21.9
2002.....	1.1	*	*	*	*	*	0.1	0.3	0.8	2.7	7.6	23.5
2001.....	1.1	*	*	*	*	*	0.1	0.3	0.9	2.7	7.4	23.9
2000.....	1.1	*	*	*	*	*	0.1	0.2	0.7	2.3	7.4	25.6
1999.....	1.1	*	*	*	*	*	0.1	0.3	0.7	2.5	7.8	24.4
White, male												
2004.....	1.0	*	*	*	*	*	0.1	0.3	0.9	2.9	9.3	26.7
2003.....	1.0	*	*	*	*	*	*	0.3	0.7	3.2	10.0	23.0
2002.....	1.0	*	*	*	*	*	0.1	0.4	0.9	3.1	9.8	24.7
2001.....	1.0	*	*	*	*	*	0.1	0.4	1.1	3.1	9.3	26.7
2000.....	1.0	*	*	*	*	*	0.1	0.2	0.9	2.8	9.6	28.9
1999.....	1.0	*	*	*	*	*	0.1	0.3	0.9	3.0	9.6	28.1

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Cholelithiasis and other disorders of gallbladder (K80-K82)

White, female

2004.....	1.2	*	*	*	*	*	0.1	0.3	0.6	1.9	6.6	23.1
2003.....	1.2	*	*	*	*	*	*	0.2	0.5	2.2	6.9	21.4
2002.....	1.2	*	*	*	*	*	*	0.2	0.6	2.4	6.2	23.0
2001.....	1.2	*	*	*	*	*	*	0.3	0.7	2.4	6.1	22.8
2000.....	1.2	*	*	*	*	*	*	0.2	0.5	1.8	6.0	24.3
1999.....	1.2	*	*	*	*	*	*	0.3	0.6	2.1	6.7	22.9

Black, both sexes

2004.....	0.8	*	*	*	*	*	*	0.8	1.6	3.5	9.1	19.5
2003.....	0.7	*	*	*	*	*	*	0.7	1.2	2.9	7.9	20.7
2002.....	0.7	*	*	*	*	*	*	0.6	1.1	3.3	9.7	14.8
2001.....	0.8	*	*	*	*	*	*	0.7	1.7	4.7	8.3	19.7
2000.....	0.8	*	*	*	*	*	*	0.7	1.2	3.9	8.3	17.8
1999.....	0.7	*	*	*	*	*	0.4	0.5	1.5	3.6	7.2	15.3

Black, male

2004.....	0.7	*	*	*	*	*	*	1.0	1.6	3.6	8.9	*
2003.....	0.7	*	*	*	*	*	*	*	*	3.2	10.4	20.9
2002.....	0.6	*	*	*	*	*	*	*	*	*	10.9	*
2001.....	0.7	*	*	*	*	*	*	*	2.1	4.9	6.2	27.2
2000.....	0.6	*	*	*	*	*	*	*	*	3.7	8.2	*
1999.....	0.6	*	*	*	*	*	*	*	*	3.7	7.1	*

Black, female

2004.....	1.0	*	*	*	*	*	*	*	1.6	3.5	9.3	21.7
2003.....	0.8	*	*	*	*	*	*	0.8	*	2.7	6.5	20.7
2002.....	0.8	*	*	*	*	*	*	*	*	3.7	9.1	16.0
2001.....	0.9	*	*	*	*	*	*	*	*	4.5	9.4	16.8
2000.....	1.0	*	*	*	*	*	*	*	*	4.1	8.4	19.3
1999.....	0.8	*	*	*	*	*	*	*	*	3.5	7.2	15.3

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Disorders of biliary tract and pancreas (K83-K86)												
All races 3/, both sexes												
2004.....	1.7	*	*	*	0.1	0.3	0.8	1.4	2.2	4.7	9.4	20.8
2003.....	1.7	*	*	*	0.1	0.3	0.8	1.4	2.3	4.8	10.0	21.9
2002.....	1.7	*	*	*	0.1	0.3	0.8	1.4	2.5	4.8	9.7	21.7
2001.....	1.7	*	*	*	0.1	0.4	0.9	1.6	2.5	5.1	9.4	22.9
2000.....	1.7	*	*	*	0.1	0.3	0.8	1.4	2.5	4.6	9.4	22.0
1999.....	1.6	*	*	*	0.1	0.4	0.8	1.4	2.4	4.5	8.5	22.2
All races 3/, male												
2004.....	1.8	*	*	*	0.1	0.5	1.1	2.0	3.0	5.7	10.6	21.1
2003.....	1.8	*	*	*	*	0.4	1.0	1.9	3.1	6.1	11.8	20.7
2002.....	1.9	*	*	*	0.2	0.5	1.1	2.0	3.4	6.4	11.9	21.2
2001.....	1.8	*	*	*	0.1	0.5	1.1	2.0	3.3	6.5	10.2	22.4
2000.....	1.8	*	*	*	0.1	0.5	1.2	1.9	3.0	5.9	11.4	22.9
1999.....	1.7	*	*	*	0.1	0.5	1.1	1.9	3.1	5.6	10.1	23.1
All races 3/, female												
2004.....	1.6	*	*	*	*	0.1	0.5	0.9	1.5	3.8	8.6	20.7
2003.....	1.6	*	*	*	*	0.2	0.6	0.9	1.5	3.7	8.8	22.4
2002.....	1.5	*	*	*	*	0.2	0.5	0.8	1.6	3.5	8.3	21.9
2001.....	1.7	*	*	*	*	0.3	0.6	1.1	1.8	3.9	8.8	23.2
2000.....	1.6	*	*	*	*	0.2	0.5	1.0	2.0	3.5	8.0	21.6
1999.....	1.5	*	*	*	0.1	0.2	0.5	0.9	1.7	3.6	7.4	21.8
White, both sexes												
2004.....	1.7	*	*	*	0.1	0.3	0.7	1.3	2.0	4.6	9.4	21.0
2003.....	1.7	*	*	*	0.1	0.3	0.7	1.2	2.1	4.6	10.2	22.0
2002.....	1.8	*	*	*	0.1	0.3	0.7	1.2	2.3	4.8	9.9	22.0
2001.....	1.8	*	*	*	*	0.3	0.8	1.3	2.4	5.0	9.4	23.0
2000.....	1.7	*	*	*	0.1	0.3	0.8	1.3	2.3	4.4	9.3	21.9
1999.....	1.6	*	*	*	0.1	0.3	0.7	1.2	2.2	4.5	8.3	22.0
White, male												
2004.....	1.8	*	*	*	*	0.4	1.1	1.7	2.7	5.6	10.6	21.4
2003.....	1.8	*	*	*	*	0.4	0.8	1.7	2.8	5.9	11.9	20.5
2002.....	1.9	*	*	*	0.1	0.4	1.0	1.8	3.2	6.3	12.1	21.2
2001.....	1.8	*	*	*	*	0.4	1.0	1.7	3.1	6.4	10.3	22.2
2000.....	1.8	*	*	*	*	0.4	1.1	1.7	2.8	5.6	11.3	22.0
1999.....	1.7	*	*	*	*	0.4	1.0	1.6	2.8	5.6	10.0	22.0

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Disorders of biliary tract and pancreas (K83-K86)												
White, female												
2004.....	1.6	*	*	*	*	*	0.4	0.8	1.3	3.7	8.5	20.8
2003.....	1.7	*	*	*	*	0.1	0.5	0.7	1.5	3.5	9.0	22.6
2002.....	1.6	*	*	*	*	0.2	0.5	0.7	1.5	3.5	8.3	22.3
2001.....	1.8	*	*	*	*	0.2	0.5	0.9	1.8	3.8	8.8	23.4
2000.....	1.6	*	*	*	*	0.2	0.5	0.9	1.8	3.3	8.0	21.9
1999.....	1.6	*	*	*	*	0.2	0.5	0.8	1.6	3.5	7.2	22.0
Black, both sexes												
2004.....	1.8	*	*	*	*	0.5	1.5	2.9	4.8	6.3	10.6	20.3
2003.....	1.9	*	*	*	*	0.5	1.8	3.1	4.5	7.4	8.9	23.3
2002.....	1.7	*	*	*	*	0.6	1.4	3.0	4.4	5.9	8.6	21.4
2001.....	1.9	*	*	*	*	0.8	1.9	3.6	4.1	6.7	9.4	22.1
2000.....	1.8	*	*	*	*	0.8	1.5	3.0	4.5	7.2	10.0	23.5
1999.....	1.8	*	*	*	*	0.6	1.5	3.3	4.7	5.5	9.9	23.0
Black, male												
2004.....	2.1	*	*	*	*	0.9	1.9	4.3	6.9	8.5	12.0	*
2003.....	2.2	*	*	*	*	*	2.7	4.1	6.8	9.1	11.8	26.1
2002.....	2.1	*	*	*	*	0.9	1.9	4.6	6.4	8.4	9.8	25.7
2001.....	2.2	*	*	*	*	1.0	2.3	5.1	5.9	8.7	10.0	*
2000.....	2.2	*	*	*	*	1.0	2.2	4.1	6.0	10.1	12.1	32.3
1999.....	2.2	*	*	*	*	1.1	2.2	5.2	6.5	6.2	10.8	32.8
Black, female												
2004.....	1.5	*	*	*	*	*	1.2	1.8	3.2	4.7	9.8	20.9
2003.....	1.5	*	*	*	*	*	1.0	2.2	2.7	6.1	7.2	22.3
2002.....	1.4	*	*	*	*	*	1.0	1.7	2.9	4.2	8.0	19.7
2001.....	1.7	*	*	*	*	*	1.5	2.4	2.6	5.3	9.1	22.3
2000.....	1.5	*	*	*	*	*	0.8	2.0	3.4	5.1	8.9	20.2
1999.....	1.4	*	*	*	*	*	0.9	1.6	3.3	5.0	9.5	19.3

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pancreatitis (K85,K86.0-K86.1)												
All races 3/, both sexes												
2004.....	1.2	*	*	*	0.1	0.3	0.7	1.3	1.8	3.4	5.9	11.4
2003.....	1.2	*	*	*	0.1	0.3	0.7	1.2	1.8	3.4	6.7	11.8
2002.....	1.2	*	*	*	0.1	0.3	0.7	1.2	2.0	3.4	6.5	11.9
2001.....	1.3	*	*	*	0.1	0.3	0.8	1.3	2.1	3.8	6.4	13.2
2000.....	1.2	*	*	*	0.1	0.3	0.8	1.2	2.0	3.4	6.3	12.1
1999.....	1.2	*	*	*	0.1	0.3	0.7	1.2	1.9	3.5	5.8	12.8
All races 3/, male												
2004.....	1.3	*	*	*	*	0.5	1.0	1.8	2.5	4.3	6.6	10.3
2003.....	1.3	*	*	*	*	0.4	1.0	1.6	2.5	4.5	7.7	10.6
2002.....	1.4	*	*	*	0.1	0.4	1.0	1.8	2.8	4.6	7.8	11.2
2001.....	1.4	*	*	*	*	0.4	1.0	1.8	2.8	4.9	6.8	12.9
2000.....	1.3	*	*	*	0.1	0.4	1.1	1.6	2.5	4.5	7.4	11.9
1999.....	1.3	*	*	*	*	0.5	1.0	1.7	2.5	4.4	6.8	12.7
All races 3/, female												
2004.....	1.0	*	*	*	*	0.1	0.4	0.7	1.1	2.7	5.4	11.8
2003.....	1.1	*	*	*	*	0.1	0.5	0.7	1.1	2.6	6.0	12.3
2002.....	1.1	*	*	*	*	0.1	0.5	0.7	1.3	2.4	5.7	12.2
2001.....	1.2	*	*	*	*	0.2	0.5	0.9	1.5	3.0	6.1	13.4
2000.....	1.1	*	*	*	*	0.2	0.4	0.8	1.5	2.5	5.6	12.2
1999.....	1.1	*	*	*	0.1	0.2	0.5	0.8	1.3	2.7	5.2	12.8
White, both sexes												
2004.....	1.2	*	*	*	0.1	0.3	0.6	1.1	1.6	3.3	5.9	11.4
2003.....	1.2	*	*	*	*	0.2	0.6	1.0	1.7	3.3	6.8	11.6
2002.....	1.3	*	*	*	0.1	0.3	0.7	1.1	1.9	3.5	6.7	12.1
2001.....	1.3	*	*	*	*	0.3	0.7	1.1	2.0	3.8	6.4	13.3
2000.....	1.2	*	*	*	0.1	0.3	0.7	1.1	1.8	3.2	6.3	12.2
1999.....	1.2	*	*	*	0.1	0.3	0.7	1.0	1.7	3.4	5.7	12.8
White, male												
2004.....	1.3	*	*	*	*	0.4	1.0	1.6	2.2	4.2	6.5	10.4
2003.....	1.3	*	*	*	*	0.4	0.8	1.4	2.2	4.2	7.7	10.3
2002.....	1.4	*	*	*	0.1	0.4	0.9	1.6	2.6	4.7	7.9	11.0
2001.....	1.4	*	*	*	*	0.3	0.9	1.5	2.6	5.0	6.8	12.9
2000.....	1.3	*	*	*	*	0.4	1.0	1.4	2.2	4.3	7.3	11.5
1999.....	1.3	*	*	*	*	0.4	0.9	1.3	2.3	4.4	6.8	12.0

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pancreatitis (K85,K86.0-K86.1)												
White, female												
2004.....	1.1	*	*	*	*	*	0.3	0.6	1.0	2.6	5.5	11.8
2003.....	1.1	*	*	*	*	*	0.4	0.6	1.1	2.4	6.1	12.2
2002.....	1.1	*	*	*	*	0.1	0.4	0.6	1.3	2.5	5.9	12.6
2001.....	1.2	*	*	*	*	0.2	0.4	0.7	1.5	2.9	6.2	13.4
2000.....	1.1	*	*	*	*	0.1	0.4	0.7	1.4	2.4	5.6	12.5
1999.....	1.1	*	*	*	*	0.2	0.4	0.7	1.1	2.6	5.0	13.1
Black, both sexes												
2004.....	1.4	*	*	*	*	0.5	1.3	2.7	4.0	4.7	7.1	13.3
2003.....	1.4	*	*	*	*	0.4	1.6	2.7	3.7	5.3	5.8	14.0
2002.....	1.3	*	*	*	*	0.5	1.3	2.7	3.7	3.7	5.2	11.3
2001.....	1.5	*	*	*	*	0.6	1.7	3.3	3.4	4.6	6.5	13.2
2000.....	1.4	*	*	*	*	0.7	1.4	2.6	3.9	5.4	6.2	11.9
1999.....	1.4	*	*	*	*	0.6	1.4	3.0	3.7	4.4	7.1	12.8
Black, male												
2004.....	1.8	*	*	*	*	0.8	1.7	3.9	5.8	6.8	8.6	*
2003.....	1.9	*	*	*	*	*	2.4	3.7	6.0	7.3	7.9	*
2002.....	1.7	*	*	*	*	0.8	1.8	4.0	5.5	5.1	6.6	*
2001.....	1.8	*	*	*	*	0.9	2.1	4.5	5.2	5.6	7.4	*
2000.....	1.8	*	*	*	*	0.9	2.2	3.5	5.1	8.0	8.2	*
1999.....	1.8	*	*	*	*	1.0	2.1	4.8	5.2	5.1	7.7	*
Black, female												
2004.....	1.1	*	*	*	*	*	0.9	1.6	2.6	3.2	6.3	13.4
2003.....	1.1	*	*	*	*	*	0.9	1.7	1.8	3.9	4.6	13.4
2002.....	0.9	*	*	*	*	*	0.8	1.5	2.3	2.6	4.4	9.4
2001.....	1.2	*	*	*	*	*	1.3	2.2	1.9	3.9	5.9	13.9
2000.....	1.1	*	*	*	*	*	0.7	1.8	2.9	3.5	5.1	9.9
1999.....	1.1	*	*	*	*	*	0.7	1.6	2.6	3.8	6.7	10.5

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other disorders of biliary tract and pancreas (K83,K86.2-K86.9)

All races 3/, both sexes

2004.....	0.5	*	*	*	*	*	0.1	0.2	0.4	1.3	3.5	9.4
2003.....	0.5	*	*	*	*	*	0.1	0.2	0.5	1.4	3.3	10.1
2002.....	0.5	*	*	*	*	*	0.1	0.2	0.5	1.4	3.2	9.8
2001.....	0.5	*	*	*	*	0.1	0.1	0.3	0.4	1.2	3.0	9.7
2000.....	0.5	*	*	*	*	*	0.1	0.2	0.5	1.2	3.1	9.9
1999.....	0.4	*	*	*	*	*	0.1	0.2	0.5	1.0	2.6	9.4

All races 3/, male

2004.....	0.4	*	*	*	*	*	*	0.2	0.5	1.4	4.1	10.7
2003.....	0.5	*	*	*	*	*	*	0.3	0.6	1.6	4.1	10.1
2002.....	0.5	*	*	*	*	*	0.1	0.2	0.6	1.8	4.2	10.0
2001.....	0.4	*	*	*	*	*	0.1	0.3	0.5	1.6	3.4	9.5
2000.....	0.4	*	*	*	*	*	*	0.3	0.5	1.3	4.0	11.0
1999.....	0.4	*	*	*	*	*	*	0.3	0.6	1.2	3.2	10.5

All races 3/, female

2004.....	0.5	*	*	*	*	*	*	0.2	0.3	1.1	3.1	8.9
2003.....	0.5	*	*	*	*	*	0.1	0.1	0.4	1.2	2.8	10.2
2002.....	0.5	*	*	*	*	*	*	0.1	0.3	1.1	2.5	9.8
2001.....	0.5	*	*	*	*	*	*	0.2	0.4	0.9	2.7	9.8
2000.....	0.5	*	*	*	*	*	0.1	0.1	0.4	1.0	2.5	9.4
1999.....	0.4	*	*	*	*	*	*	0.1	0.5	0.9	2.2	9.0

White, both sexes

2004.....	0.5	*	*	*	*	*	0.1	0.2	0.4	1.2	3.5	9.6
2003.....	0.5	*	*	*	*	*	0.1	0.2	0.5	1.3	3.4	10.3
2002.....	0.5	*	*	*	*	*	0.1	0.2	0.4	1.3	3.2	9.9
2001.....	0.5	*	*	*	*	*	0.1	0.2	0.4	1.2	3.0	9.7
2000.....	0.5	*	*	*	*	*	0.1	0.2	0.5	1.1	3.0	9.7
1999.....	0.4	*	*	*	*	*	0.1	0.2	0.5	1.0	2.6	9.2

White, male

2004.....	0.5	*	*	*	*	*	*	0.2	0.5	1.5	4.2	10.9
2003.....	0.5	*	*	*	*	*	*	0.3	0.5	1.6	4.2	10.2
2002.....	0.5	*	*	*	*	*	*	0.2	0.6	1.6	4.2	10.3
2001.....	0.4	*	*	*	*	*	*	0.3	0.5	1.5	3.6	9.3
2000.....	0.5	*	*	*	*	*	*	0.3	0.5	1.3	4.1	10.6
1999.....	0.4	*	*	*	*	*	*	0.3	0.6	1.2	3.2	10.0

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

All other diseases of digestive system (K90,K92)

All races 3/, both sexes

2004.....	2.7	0.7	*	*	*	0.1	0.3	0.9	2.2	5.6	18.2	68.5
2003.....	2.8	0.7	*	*	*	0.1	0.3	0.8	2.2	6.1	19.1	74.2
2002.....	2.9	0.6	0.1	*	*	0.1	0.3	1.0	2.1	6.1	19.7	76.2
2001.....	2.9	0.6	*	*	*	0.1	0.3	0.8	2.4	6.2	19.9	77.8
2000.....	2.9	0.6	*	*	*	0.1	0.3	0.9	2.1	5.9	20.6	80.6
1999.....	2.8	0.7	0.1	*	*	0.1	0.3	0.8	2.2	6.2	19.6	77.2

All races 3/, male

2004.....	2.5	*	*	*	*	*	0.3	1.3	2.9	7.4	21.8	69.8
2003.....	2.6	*	*	*	*	*	0.5	1.2	2.8	7.7	23.6	73.3
2002.....	2.6	*	*	*	*	0.1	0.4	1.4	2.9	7.8	23.0	76.2
2001.....	2.6	*	*	*	*	0.1	0.4	1.3	3.1	7.9	23.5	81.4
2000.....	2.6	*	*	*	*	0.1	0.4	1.3	2.8	7.1	25.1	83.1
1999.....	2.5	*	*	*	*	*	0.4	1.2	2.9	7.9	23.7	80.8

All races 3/, female

2004.....	2.9	*	*	*	*	*	0.2	0.5	1.5	4.2	15.7	67.9
2003.....	3.1	*	*	*	*	*	0.2	0.5	1.6	4.7	16.0	74.6
2002.....	3.1	*	*	*	*	*	0.1	0.5	1.4	4.7	17.6	76.2
2001.....	3.1	*	*	*	*	*	0.2	0.4	1.7	4.8	17.5	76.3
2000.....	3.2	*	*	*	*	*	0.2	0.5	1.4	4.9	17.7	79.6
1999.....	3.0	*	*	*	*	*	0.2	0.5	1.6	4.8	16.9	75.8

White, both sexes

2004.....	2.9	*	*	*	*	*	0.3	0.9	2.0	5.3	18.0	69.8
2003.....	3.1	0.7	*	*	*	*	0.3	0.8	2.0	5.7	18.9	75.5
2002.....	3.1	*	*	*	*	*	0.2	0.9	1.9	5.8	19.5	77.8
2001.....	3.1	*	*	*	*	0.1	0.3	0.8	2.1	5.9	19.7	79.1
2000.....	3.1	*	*	*	*	*	0.2	0.7	1.8	5.6	20.2	81.7
1999.....	3.0	*	*	*	*	*	0.2	0.7	2.0	5.9	19.2	78.9

White, male

2004.....	2.6	*	*	*	*	*	0.3	1.3	2.6	6.9	21.6	71.6
2003.....	2.7	*	*	*	*	*	0.4	1.1	2.6	7.2	23.4	75.2
2002.....	2.7	*	*	*	*	*	0.3	1.3	2.7	7.3	22.8	77.3
2001.....	2.7	*	*	*	*	0.1	0.4	1.1	2.7	7.5	23.2	82.8
2000.....	2.7	*	*	*	*	*	0.3	1.1	2.5	6.7	24.5	84.9
1999.....	2.6	*	*	*	*	*	0.3	1.0	2.6	7.5	23.3	82.1

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

All other diseases of digestive system (K90,K92)

White, female

2004.....	3.2	*	*	*	*	*	0.2	0.5	1.5	4.0	15.6	69.0
2003.....	3.4	*	*	*	*	*	0.2	0.5	1.4	4.4	15.9	75.6
2002.....	3.4	*	*	*	*	*	0.1	0.5	1.2	4.5	17.3	78.1
2001.....	3.4	*	*	*	*	*	0.1	0.4	1.5	4.7	17.4	77.5
2000.....	3.4	*	*	*	*	*	0.1	0.4	1.1	4.6	17.3	80.5
1999.....	3.3	*	*	*	*	*	0.1	0.4	1.4	4.6	16.5	77.6

Black, both sexes

2004.....	2.3	*	*	*	*	*	*	1.6	4.1	9.8	24.4	62.0
2003.....	2.3	*	*	*	*	*	0.5	1.4	4.2	10.5	24.4	70.3
2002.....	2.4	*	*	*	*	*	0.5	1.8	4.3	9.7	26.3	65.3
2001.....	2.4	*	*	*	*	*	0.6	1.7	5.2	9.7	25.7	73.1
2000.....	2.6	*	*	*	*	*	0.7	2.0	5.0	9.8	28.0	77.9
1999.....	2.3	*	*	*	*	*	0.4	1.7	4.2	9.8	26.3	66.0

Black, male

2004.....	2.3	*	*	*	*	*	*	2.2	6.4	13.7	29.2	58.6
2003.....	2.3	*	*	*	*	*	0.9	2.0	5.6	13.8	31.3	59.4
2002.....	2.4	*	*	*	*	*	0.8	2.6	5.7	13.7	30.1	70.8
2001.....	2.5	*	*	*	*	*	0.8	2.9	7.0	13.6	32.4	71.5
2000.....	2.5	*	*	*	*	*	0.9	2.8	6.6	12.0	35.7	78.4
1999.....	2.2	*	*	*	*	*	*	2.5	5.7	12.8	32.3	71.5

Black, female

2004.....	2.2	*	*	*	*	*	*	1.1	2.2	6.9	21.7	63.4
2003.....	2.4	*	*	*	*	*	*	0.9	3.0	8.1	20.6	74.5
2002.....	2.3	*	*	*	*	*	*	1.1	3.2	6.9	24.2	63.3
2001.....	2.4	*	*	*	*	*	*	*	3.7	6.9	22.0	73.7
2000.....	2.6	*	*	*	*	*	*	1.3	3.6	8.2	23.7	77.7
1999.....	2.3	*	*	*	*	*	*	1.0	3.0	7.7	22.9	64.0

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
XII. Diseases of the skin and subcutaneous tissue (L00-L98)												
All races 3/, both sexes												
2004.....	1.4	*	*	*	0.1	0.1	0.3	0.7	1.4	3.4	9.5	28.8
2003.....	1.4	*	*	*	*	0.2	0.3	0.7	1.4	3.3	9.2	30.5
2002.....	1.4	*	*	*	0.1	0.1	0.3	0.6	1.2	3.1	8.7	32.8
2001.....	1.3	*	*	*	*	0.1	0.3	0.6	1.2	2.8	8.7	32.2
2000.....	1.3	*	*	*	*	0.1	0.2	0.5	1.2	3.0	8.9	34.7
1999.....	1.3	*	*	*	*	0.1	0.2	0.5	1.0	2.9	9.0	35.0
All races 3/, male												
2004.....	1.1	*	*	*	*	0.1	0.3	0.7	1.3	3.0	9.0	25.3
2003.....	1.0	*	*	*	*	*	0.3	0.7	1.4	3.3	8.4	26.6
2002.....	1.0	*	*	*	*	*	0.2	0.6	1.1	3.2	8.6	28.6
2001.....	0.9	*	*	*	*	*	0.2	0.6	1.2	3.0	8.3	24.9
2000.....	0.9	*	*	*	*	*	0.2	0.4	1.3	2.8	8.4	29.7
1999.....	0.9	*	*	*	*	*	0.2	0.5	1.0	2.8	8.4	31.7
All races 3/, female												
2004.....	1.8	*	*	*	*	0.2	0.3	0.6	1.5	3.7	9.8	30.4
2003.....	1.8	*	*	*	*	0.3	0.3	0.7	1.4	3.4	9.7	32.2
2002.....	1.7	*	*	*	*	0.2	0.3	0.6	1.3	3.0	8.8	34.7
2001.....	1.7	*	*	*	*	0.2	0.3	0.6	1.2	2.6	9.1	35.2
2000.....	1.7	*	*	*	*	0.2	0.3	0.5	1.1	3.1	9.2	36.7
1999.....	1.7	*	*	*	*	0.1	0.3	0.5	1.1	2.9	9.4	36.3
White, both sexes												
2004.....	1.3	*	*	*	*	*	0.2	0.6	1.2	2.8	8.5	25.7
2003.....	1.3	*	*	*	*	0.1	0.2	0.5	1.2	2.7	8.1	27.2
2002.....	1.2	*	*	*	*	0.1	0.2	0.5	1.0	2.5	7.5	28.3
2001.....	1.2	*	*	*	*	0.1	0.2	0.5	1.0	2.4	7.6	28.1
2000.....	1.2	*	*	*	*	0.1	0.2	0.3	1.0	2.4	7.6	29.7
1999.....	1.2	*	*	*	*	0.1	0.2	0.4	0.8	2.3	7.5	29.3
White, male												
2004.....	1.0	*	*	*	*	*	0.3	0.6	1.2	2.5	8.2	23.1
2003.....	1.0	*	*	*	*	*	0.2	0.5	1.2	2.5	7.5	24.0
2002.....	1.0	*	*	*	*	*	0.2	0.5	1.0	2.5	7.7	26.1
2001.....	0.9	*	*	*	*	*	0.2	0.5	1.0	2.7	7.0	21.8
2000.....	0.8	*	*	*	*	*	0.2	0.4	1.1	2.1	7.0	25.8
1999.....	0.8	*	*	*	*	*	0.1	0.4	0.7	2.3	6.9	27.8

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
XII. Diseases of the skin and subcutaneous tissue (L00-L98)												
White, female												
2004.....	1.7	*	*	*	*	*	0.2	0.6	1.2	3.1	8.7	27.0
2003.....	1.7	*	*	*	*	0.2	0.3	0.5	1.2	3.0	8.6	28.7
2002.....	1.5	*	*	*	*	*	0.2	0.4	1.0	2.5	7.3	29.3
2001.....	1.5	*	*	*	*	*	0.2	0.5	0.9	2.1	7.9	30.8
2000.....	1.6	*	*	*	*	*	0.2	0.3	0.9	2.6	7.9	31.2
1999.....	1.5	*	*	*	*	*	0.2	0.4	0.8	2.3	7.9	30.0
Black, both sexes												
2004.....	2.4	*	*	*	*	0.6	0.8	1.5	3.6	9.6	23.4	70.8
2003.....	2.4	*	*	*	*	0.7	0.8	1.8	3.5	9.5	23.7	73.2
2002.....	2.5	*	*	*	*	0.5	0.9	1.4	3.7	8.4	25.5	92.9
2001.....	2.3	*	*	*	*	0.4	0.8	1.4	3.1	7.3	24.9	88.4
2000.....	2.4	*	*	*	*	0.4	0.7	1.3	3.0	8.9	26.2	97.3
1999.....	2.7	*	*	*	*	*	0.8	1.8	3.5	8.9	29.1	105.2
Black, male												
2004.....	1.7	*	*	*	*	*	*	1.5	2.9	9.0	22.0	57.6
2003.....	1.9	*	*	*	*	*	0.7	1.5	3.8	11.4	23.4	65.7
2002.....	1.7	*	*	*	*	*	*	1.3	3.0	10.0	23.8	63.3
2001.....	1.7	*	*	*	*	*	*	1.6	2.6	7.2	28.3	71.5
2000.....	1.8	*	*	*	*	*	*	*	2.9	10.1	27.9	81.9
1999.....	1.9	*	*	*	*	*	*	2.1	3.2	9.1	29.5	85.5
Black, female												
2004.....	3.0	*	*	*	*	0.8	0.9	1.4	4.1	10.0	24.2	76.0
2003.....	2.9	*	*	*	*	1.0	0.9	1.9	3.3	8.1	23.9	76.2
2002.....	3.3	*	*	*	*	0.7	1.2	1.6	4.3	7.2	26.5	104.3
2001.....	2.9	*	*	*	*	*	1.0	1.2	3.5	7.4	23.0	94.7
2000.....	3.1	*	*	*	*	*	0.7	1.7	3.1	8.1	25.2	103.0
1999.....	3.3	*	*	*	*	*	0.9	1.6	3.7	8.7	28.9	112.6

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Infections of skin and subcutaneous tissue (I00-L08)												
All races 3/, both sexes												
2004.....	0.6	*	*	*	*	*	0.1	0.3	0.7	1.4	3.8	10.4
2003.....	0.5	*	*	*	*	*	0.1	0.3	0.7	1.3	3.5	10.6
2002.....	0.4	*	*	*	*	*	0.1	0.3	0.6	1.1	2.8	9.3
2001.....	0.4	*	*	*	*	*	0.1	0.3	0.5	1.0	2.6	9.4
2000.....	0.4	*	*	*	*	*	0.1	0.2	0.5	1.0	2.5	9.3
1999.....	0.4	*	*	*	*	*	0.1	0.2	0.4	0.9	2.1	8.3
All races 3/, male												
2004.....	0.5	*	*	*	*	*	0.2	0.4	0.7	1.4	4.0	10.3
2003.....	0.5	*	*	*	*	*	0.1	0.4	0.8	1.4	3.5	10.4
2002.....	0.4	*	*	*	*	*	0.1	0.4	0.6	1.1	3.0	9.1
2001.....	0.4	*	*	*	*	*	0.1	0.4	0.6	1.2	2.4	8.1
2000.....	0.3	*	*	*	*	*	*	0.2	0.7	1.1	2.8	8.9
1999.....	0.3	*	*	*	*	*	*	0.3	0.4	0.9	2.1	9.1
All races 3/, female												
2004.....	0.7	*	*	*	*	*	0.1	0.3	0.7	1.5	3.7	10.4
2003.....	0.6	*	*	*	*	*	0.1	0.2	0.6	1.3	3.4	10.7
2002.....	0.5	*	*	*	*	*	*	0.2	0.5	1.0	2.6	9.4
2001.....	0.5	*	*	*	*	*	*	0.2	0.4	0.9	2.7	10.0
2000.....	0.5	*	*	*	*	*	0.1	0.1	0.4	0.9	2.4	9.5
1999.....	0.4	*	*	*	*	*	*	0.2	0.4	0.9	2.1	8.0
White, both sexes												
2004.....	0.6	*	*	*	*	*	0.1	0.3	0.6	1.4	3.9	10.4
2003.....	0.6	*	*	*	*	*	0.1	0.3	0.7	1.3	3.5	10.8
2002.....	0.5	*	*	*	*	*	0.1	0.2	0.5	1.0	2.7	9.7
2001.....	0.5	*	*	*	*	*	0.1	0.3	0.5	1.0	2.6	9.6
2000.....	0.4	*	*	*	*	*	0.1	0.2	0.5	1.0	2.5	9.3
1999.....	0.4	*	*	*	*	*	*	0.2	0.3	0.9	2.1	8.3
White, male												
2004.....	0.5	*	*	*	*	*	0.2	0.4	0.6	1.3	4.1	10.8
2003.....	0.5	*	*	*	*	*	0.1	0.4	0.8	1.3	3.5	10.6
2002.....	0.4	*	*	*	*	*	*	0.3	0.6	1.0	3.0	9.4
2001.....	0.4	*	*	*	*	*	*	0.3	0.6	1.3	2.4	8.4
2000.....	0.4	*	*	*	*	*	*	0.2	0.7	1.1	2.8	8.7
1999.....	0.3	*	*	*	*	*	*	0.2	0.4	0.9	2.0	8.9

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified diseases of skin and subcutaneous tissue (L10-L98)												
All races 3/, both sexes												
2004.....	0.8	*	*	*	0.1	0.1	0.2	0.3	0.7	2.0	5.7	18.4
2003.....	0.9	*	*	*	*	0.1	0.2	0.3	0.7	2.0	5.7	19.9
2002.....	0.9	*	*	*	*	0.1	0.2	0.3	0.7	2.0	5.9	23.5
2001.....	0.9	*	*	*	*	0.1	0.1	0.3	0.7	1.8	6.1	22.7
2000.....	0.9	*	*	*	*	0.1	0.2	0.3	0.6	1.9	6.4	25.4
1999.....	1.0	*	*	*	*	0.1	0.2	0.3	0.6	2.0	7.0	26.6
All races 3/, male												
2004.....	0.6	*	*	*	*	*	0.1	0.3	0.7	1.6	5.0	14.9
2003.....	0.6	*	*	*	*	*	0.2	0.2	0.6	1.9	4.9	16.2
2002.....	0.6	*	*	*	*	*	0.1	0.2	0.5	2.0	5.7	19.4
2001.....	0.6	*	*	*	*	*	0.1	0.2	0.5	1.8	5.8	16.8
2000.....	0.6	*	*	*	*	*	0.1	0.2	0.6	1.6	5.6	20.8
1999.....	0.6	*	*	*	*	*	0.1	0.2	0.5	2.0	6.3	22.6
All races 3/, female												
2004.....	1.1	*	*	*	*	0.1	0.2	0.4	0.8	2.2	6.1	20.0
2003.....	1.2	*	*	*	*	0.2	0.2	0.4	0.8	2.1	6.2	21.5
2002.....	1.2	*	*	*	*	0.2	0.3	0.4	0.8	2.0	6.1	25.3
2001.....	1.2	*	*	*	*	0.1	0.2	0.3	0.8	1.8	6.4	25.2
2000.....	1.2	*	*	*	*	0.1	0.2	0.4	0.7	2.2	6.9	27.3
1999.....	1.3	*	*	*	*	*	0.2	0.4	0.7	2.0	7.4	28.3
White, both sexes												
2004.....	0.7	*	*	*	*	*	0.1	0.2	0.6	1.4	4.6	15.3
2003.....	0.7	*	*	*	*	*	0.1	0.2	0.5	1.5	4.6	16.4
2002.....	0.8	*	*	*	*	*	0.1	0.2	0.4	1.5	4.7	18.7
2001.....	0.8	*	*	*	*	*	0.1	0.2	0.5	1.4	5.0	18.5
2000.....	0.8	*	*	*	*	*	0.1	0.2	0.5	1.4	5.1	20.4
1999.....	0.8	*	*	*	*	*	0.1	0.2	0.4	1.5	5.4	21.0
White, male												
2004.....	0.5	*	*	*	*	*	*	0.2	0.5	1.2	4.2	12.3
2003.....	0.5	*	*	*	*	*	0.1	0.2	0.5	1.2	4.0	13.5
2002.....	0.5	*	*	*	*	*	*	0.2	0.3	1.5	4.7	16.7
2001.....	0.5	*	*	*	*	*	*	0.2	0.4	1.4	4.5	13.3
2000.....	0.5	*	*	*	*	*	*	0.2	0.4	1.0	4.3	17.0
1999.....	0.5	*	*	*	*	*	*	0.2	0.4	1.4	4.9	18.9

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other and unspecified diseases of skin and subcutaneous tissue (L10-L98)

White, female

2004.....	1.0	*	*	*	*	*	0.1	0.3	0.6	1.6	4.9	16.7
2003.....	1.0	*	*	*	*	*	0.1	0.3	0.6	1.7	5.1	17.8
2002.....	1.0	*	*	*	*	*	0.2	0.3	0.5	1.6	4.8	19.6
2001.....	1.0	*	*	*	*	*	*	0.3	0.6	1.3	5.2	20.6
2000.....	1.1	*	*	*	*	*	0.1	0.2	0.5	1.7	5.6	21.7
1999.....	1.1	*	*	*	*	*	0.2	0.3	0.5	1.5	5.7	21.9

Black, both sexes

2004.....	1.9	*	*	*	*	0.5	0.6	0.9	2.5	7.4	19.1	58.1
2003.....	2.0	*	*	*	*	0.7	0.7	1.2	2.6	7.2	19.9	63.6
2002.....	2.1	*	*	*	*	0.4	0.7	1.1	2.8	6.4	21.8	85.8
2001.....	1.9	*	*	*	*	*	0.6	0.9	2.1	6.2	21.7	79.2
2000.....	2.1	*	*	*	*	*	0.5	1.0	2.3	7.5	23.0	87.3
1999.....	2.3	*	*	*	*	*	0.6	1.2	2.4	7.8	27.5	96.3

Black, male

2004.....	1.2	*	*	*	*	*	*	*	1.9	6.8	17.5	48.7
2003.....	1.4	*	*	*	*	*	*	*	2.7	9.1	19.5	55.3
2002.....	1.3	*	*	*	*	*	*	*	2.2	7.7	20.7	56.8
2001.....	1.3	*	*	*	*	*	*	*	1.8	6.2	25.3	65.8
2000.....	1.5	*	*	*	*	*	*	*	2.3	8.0	24.5	72.7
1999.....	1.6	*	*	*	*	*	*	1.1	2.2	8.7	26.4	73.8

Black, female

2004.....	2.4	*	*	*	*	*	0.7	1.1	3.0	7.8	20.0	61.8
2003.....	2.5	*	*	*	*	1.0	0.8	1.6	2.5	5.8	20.1	66.8
2002.....	2.9	*	*	*	*	*	1.0	1.3	3.3	5.5	22.4	96.9
2001.....	2.4	*	*	*	*	*	0.8	0.9	2.3	6.1	19.7	84.2
2000.....	2.7	*	*	*	*	*	*	1.5	2.3	7.1	22.2	92.7
1999.....	3.0	*	*	*	*	*	0.8	1.2	2.6	7.2	28.1	104.7

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

XIII. Diseases of the musculoskeletal system and connective tissue (M00-M99)

All races 3/, both sexes

2004.....	4.9	*	*	0.1	0.4	0.7	1.3	2.7	5.4	12.1	31.4	85.9
2003.....	5.0	*	*	0.1	0.4	0.8	1.4	2.7	5.6	12.7	31.7	90.4
2002.....	5.0	*	*	0.1	0.4	0.7	1.4	2.7	5.5	12.9	31.7	94.9
2001.....	4.9	*	*	0.1	0.5	0.7	1.5	2.6	5.3	12.8	32.2	93.4
2000.....	4.9	*	*	0.1	0.4	0.8	1.4	2.6	5.4	12.6	32.2	94.0
1999.....	4.8	*	*	0.1	0.4	0.7	1.4	2.7	5.4	12.7	31.1	91.2

All races 3/, male

2004.....	2.9	*	*	*	0.3	0.3	0.8	2.1	4.1	9.0	24.2	59.1
2003.....	3.0	*	*	*	0.2	0.4	1.0	1.9	4.1	10.3	24.5	59.9
2002.....	3.0	*	*	*	0.2	0.4	0.9	1.9	4.0	10.2	24.1	63.5
2001.....	2.9	*	*	*	0.2	0.4	1.0	1.9	3.9	9.6	24.9	63.2
2000.....	2.9	*	*	*	0.3	0.5	0.9	1.7	4.0	9.8	25.2	67.5
1999.....	2.8	*	*	*	0.2	0.4	0.9	1.8	3.8	9.5	24.2	63.7

All races 3/, female

2004.....	6.7	*	*	0.1	0.5	1.1	1.9	3.4	6.7	14.6	36.3	98.0
2003.....	6.9	*	*	*	0.6	1.2	1.8	3.5	6.9	14.6	36.5	103.8
2002.....	7.0	*	*	0.1	0.6	1.0	1.8	3.3	6.9	15.1	36.8	108.5
2001.....	6.9	*	*	*	0.7	1.1	2.0	3.2	6.6	15.6	37.0	106.0
2000.....	6.8	*	*	0.1	0.5	1.1	1.9	3.5	6.6	15.0	36.8	104.8
1999.....	6.7	*	*	0.1	0.6	1.1	1.9	3.7	6.9	15.3	35.5	102.3

White, both sexes

2004.....	5.0	*	*	0.1	0.2	0.4	1.0	2.1	5.0	11.9	31.9	88.8
2003.....	5.2	*	*	0.1	0.3	0.5	1.1	2.2	5.1	12.4	32.0	92.9
2002.....	5.2	*	*	0.1	0.2	0.4	0.9	2.2	5.1	12.8	32.1	98.2
2001.....	5.1	*	*	0.1	0.3	0.5	1.1	2.1	4.8	12.6	32.9	95.8
2000.....	5.1	*	*	*	0.2	0.6	1.0	2.1	4.9	12.5	32.6	96.9
1999.....	5.0	*	*	0.1	0.2	0.5	0.9	2.2	5.0	12.6	31.7	94.4

White, male

2004.....	3.0	*	*	*	0.2	0.2	0.6	1.6	3.8	8.7	24.2	59.8
2003.....	3.1	*	*	*	0.2	0.3	0.8	1.6	3.7	9.8	24.0	60.1
2002.....	3.0	*	*	*	*	0.3	0.7	1.7	3.7	9.8	24.0	64.3
2001.....	3.0	*	*	*	0.2	0.3	0.8	1.6	3.5	9.2	25.1	64.3
2000.....	3.0	*	*	*	0.1	0.4	0.7	1.4	3.7	9.4	25.0	67.3
1999.....	2.9	*	*	*	0.1	0.3	0.6	1.5	3.5	9.4	24.1	64.4

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

XIII. Diseases of the musculoskeletal system and connective tissue (M00-M99)

White, female

2004.....	7.0	*	*	*	0.3	0.6	1.3	2.6	6.0	14.6	37.1	101.8
2003.....	7.3	*	*	*	0.4	0.7	1.4	2.7	6.5	14.7	37.4	107.4
2002.....	7.4	*	*	0.1	0.3	0.6	1.2	2.7	6.4	15.2	37.6	112.9
2001.....	7.2	*	*	*	0.4	0.6	1.4	2.6	6.1	15.5	38.0	109.0
2000.....	7.1	*	*	*	0.3	0.8	1.3	2.7	6.0	15.2	37.7	108.9
1999.....	7.0	*	*	*	0.3	0.7	1.2	2.8	6.3	15.2	36.7	106.5

Black, both sexes

2004.....	5.0	*	*	*	1.1	2.5	3.8	7.3	10.2	15.9	30.9	64.9
2003.....	4.9	*	*	*	1.2	2.5	3.4	7.1	9.9	16.9	31.9	68.0
2002.....	4.8	*	*	*	1.2	2.5	4.2	6.2	10.0	15.8	30.9	67.4
2001.....	4.8	*	*	*	1.1	2.4	4.2	6.4	9.4	16.6	29.3	75.5
2000.....	4.9	*	*	*	1.3	2.4	4.6	7.1	9.9	15.2	31.0	68.2
1999.....	4.7	*	*	*	1.2	2.1	4.7	7.0	9.6	16.2	27.0	60.3

Black, male

2004.....	3.3	*	*	*	0.6	1.5	1.9	5.4	7.2	14.5	29.0	60.6
2003.....	3.3	*	*	*	*	1.0	2.3	4.1	8.1	18.0	33.6	58.4
2002.....	3.1	*	*	*	*	1.4	2.2	4.0	7.7	15.1	28.4	61.1
2001.....	3.1	*	*	*	*	1.2	2.8	4.9	7.7	14.0	28.0	59.0
2000.....	3.4	*	*	*	1.0	1.4	3.0	4.0	7.2	16.2	30.6	78.4
1999.....	2.9	*	*	*	*	0.8	2.9	4.1	6.6	13.1	29.2	63.3

Black, female

2004.....	6.5	*	*	*	1.5	3.5	5.4	8.9	12.6	16.8	31.9	66.5
2003.....	6.4	*	*	*	2.0	3.8	4.4	9.7	11.4	16.0	30.9	71.7
2002.....	6.3	*	*	*	1.9	3.4	5.9	8.1	11.7	16.3	32.3	69.8
2001.....	6.3	*	*	*	1.9	3.6	5.6	7.7	10.8	18.4	30.1	81.7
2000.....	6.2	*	*	*	1.6	3.4	6.0	9.7	12.0	14.5	31.2	64.4
1999.....	6.2	*	*	*	1.8	3.2	6.4	9.6	12.1	18.5	25.8	59.1

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Rheumatoid arthritis and related inflammatory polyarthropathies (M05-M08)												
All races 3/, both sexes												
2004.....	0.9	*	*	*	*	*	0.1	0.2	0.9	3.1	8.0	11.9
2003.....	0.9	*	*	*	*	*	0.1	0.2	0.9	3.1	8.7	12.8
2002.....	1.0	*	*	*	*	*	0.0	0.2	1.0	3.3	9.1	14.2
2001.....	0.9	*	*	*	*	*	0.1	0.2	1.0	3.3	8.8	12.7
2000.....	1.0	*	*	*	*	*	0.1	0.2	1.0	3.5	9.2	12.9
1999.....	1.0	*	*	*	*	*	0.0	0.2	1.1	3.5	9.5	13.9
All races 3/, male												
2004.....	0.4	*	*	*	*	*	*	0.1	0.6	1.9	4.3	6.5
2003.....	0.4	*	*	*	*	*	*	0.1	0.7	1.9	5.0	6.5
2002.....	0.5	*	*	*	*	*	*	0.2	0.7	2.1	5.3	7.6
2001.....	0.4	*	*	*	*	*	*	0.1	0.7	2.0	5.1	5.5
2000.....	0.5	*	*	*	*	*	*	0.1	0.6	2.3	5.2	6.9
1999.....	0.5	*	*	*	*	*	*	0.2	0.8	2.3	5.2	7.2
All races 3/, female												
2004.....	1.3	*	*	*	*	*	*	0.3	1.1	4.1	10.5	14.3
2003.....	1.4	*	*	*	*	*	*	0.3	1.1	4.1	11.1	15.6
2002.....	1.4	*	*	*	*	*	*	0.3	1.3	4.3	11.6	17.0
2001.....	1.4	*	*	*	*	*	0.1	0.3	1.3	4.4	11.2	15.7
2000.....	1.4	*	*	*	*	*	*	0.4	1.3	4.5	11.7	15.3
1999.....	1.5	*	*	*	*	*	*	0.3	1.5	4.4	12.2	16.7
White, both sexes												
2004.....	1.0	*	*	*	*	*	*	0.2	0.9	3.2	8.2	12.1
2003.....	1.0	*	*	*	*	*	0.1	0.2	1.0	3.3	8.9	13.2
2002.....	1.1	*	*	*	*	*	*	0.3	1.0	3.5	9.4	14.8
2001.....	1.0	*	*	*	*	*	0.1	0.2	1.0	3.5	9.2	12.8
2000.....	1.1	*	*	*	*	*	0.1	0.2	1.0	3.7	9.5	13.4
1999.....	1.1	*	*	*	*	*	*	0.2	1.2	3.7	9.8	14.9
White, male												
2004.....	0.5	*	*	*	*	*	*	0.1	0.6	1.9	4.5	6.6
2003.....	0.5	*	*	*	*	*	*	0.1	0.7	2.0	5.0	6.7
2002.....	0.5	*	*	*	*	*	*	0.2	0.7	2.2	5.5	8.0
2001.....	0.5	*	*	*	*	*	*	0.1	0.8	2.1	5.2	5.5
2000.....	0.5	*	*	*	*	*	*	*	0.7	2.5	5.4	7.3
1999.....	0.5	*	*	*	*	*	*	0.2	0.9	2.4	5.4	7.8

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Rheumatoid arthritis and related inflammatory polyarthropathies (M05-M08)

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
White, female												
2004.....	1.5	*	*	*	*	*	*	0.3	1.2	4.4	10.7	14.6
2003.....	1.6	*	*	*	*	*	*	0.3	1.2	4.4	11.5	16.0
2002.....	1.6	*	*	*	*	*	*	0.3	1.3	4.5	12.0	17.7
2001.....	1.6	*	*	*	*	*	0.1	0.3	1.3	4.7	11.8	15.9
2000.....	1.6	*	*	*	*	*	*	0.4	1.3	4.8	12.3	15.8
1999.....	1.7	*	*	*	*	*	*	0.3	1.5	4.8	12.8	17.7
Black, both sexes												
2004.....	0.5	*	*	*	*	*	*	*	0.7	2.5	6.3	9.3
2003.....	0.4	*	*	*	*	*	*	*	*	2.2	6.8	9.6
2002.....	0.5	*	*	*	*	*	*	*	0.8	2.4	6.5	8.0
2001.....	0.4	*	*	*	*	*	*	*	0.9	2.2	5.0	12.0
2000.....	0.4	*	*	*	*	*	*	*	0.9	2.0	5.1	7.8
1999.....	0.4	*	*	*	*	*	*	*	0.8	1.9	6.1	*
Black, male												
2004.....	0.2	*	*	*	*	*	*	*	*	*	*	*
2003.....	0.2	*	*	*	*	*	*	*	*	*	*	*
2002.....	0.2	*	*	*	*	*	*	*	*	*	*	*
2001.....	0.2	*	*	*	*	*	*	*	*	*	*	*
2000.....	0.2	*	*	*	*	*	*	*	*	*	*	*
1999.....	0.2	*	*	*	*	*	*	*	*	*	*	*
Black, female												
2004.....	0.7	*	*	*	*	*	*	*	*	2.8	8.2	11.0
2003.....	0.6	*	*	*	*	*	*	*	*	2.8	8.0	11.3
2002.....	0.7	*	*	*	*	*	*	*	*	2.6	8.0	9.9
2001.....	0.6	*	*	*	*	*	*	*	*	2.8	5.3	14.7
2000.....	0.6	*	*	*	*	*	*	*	*	*	6.3	9.4
1999.....	0.5	*	*	*	*	*	*	*	*	2.2	6.7	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Systemic lupus erythematosus (M32)												
All races 3/, both sexes												
2004.....	0.4	*	*	*	0.2	0.4	0.5	0.6	0.8	0.8	1.3	1.1
2003.....	0.5	*	*	*	0.2	0.4	0.5	0.6	0.8	0.9	1.4	1.1
2002.....	0.4	*	*	*	0.2	0.3	0.5	0.6	0.8	1.0	1.3	1.3
2001.....	0.5	*	*	*	0.3	0.4	0.5	0.6	0.8	1.0	1.5	1.2
2000.....	0.5	*	*	*	0.2	0.3	0.6	0.7	0.9	0.9	1.5	1.2
1999.....	0.5	*	*	*	0.2	0.3	0.5	0.7	0.8	1.1	1.3	1.4
All races 3/, male												
2004.....	0.1	*	*	*	*	*	0.2	0.2	0.2	0.3	0.5	*
2003.....	0.1	*	*	*	*	*	0.2	0.2	0.2	0.4	*	*
2002.....	0.1	*	*	*	*	*	0.1	0.2	0.2	0.4	0.6	*
2001.....	0.1	*	*	*	*	0.1	0.1	0.2	0.2	0.3	0.6	*
2000.....	0.1	*	*	*	*	*	0.2	0.1	0.3	0.4	0.7	*
1999.....	0.1	*	*	*	*	*	0.1	0.2	0.2	0.3	0.6	*
All races 3/, female												
2004.....	0.7	*	*	*	0.3	0.7	0.8	1.0	1.3	1.3	1.8	1.3
2003.....	0.8	*	*	*	0.4	0.6	0.8	1.1	1.4	1.3	2.1	1.3
2002.....	0.8	*	*	*	0.4	0.5	0.9	1.0	1.4	1.5	1.8	1.7
2001.....	0.8	*	*	*	0.5	0.7	0.8	1.1	1.3	1.6	2.1	1.3
2000.....	0.8	*	*	*	0.3	0.6	1.0	1.1	1.4	1.3	2.0	1.4
1999.....	0.8	*	*	*	0.4	0.6	0.9	1.2	1.3	1.8	1.8	1.6
White, both sexes												
2004.....	0.3	*	*	*	0.1	0.2	0.3	0.4	0.6	0.7	1.2	1.2
2003.....	0.4	*	*	*	0.1	0.2	0.3	0.4	0.7	0.8	1.4	1.1
2002.....	0.3	*	*	*	0.1	0.1	0.2	0.4	0.6	0.8	1.3	1.4
2001.....	0.4	*	*	*	0.1	0.2	0.2	0.5	0.6	1.0	1.5	1.2
2000.....	0.4	*	*	*	0.1	0.2	0.3	0.4	0.7	0.8	1.5	1.3
1999.....	0.4	*	*	*	0.1	0.2	0.3	0.5	0.6	1.0	1.3	1.5
White, male												
2004.....	0.1	*	*	*	*	*	*	0.1	0.2	0.3	0.5	*
2003.....	0.1	*	*	*	*	*	0.1	0.1	0.2	0.4	*	*
2002.....	0.1	*	*	*	*	*	*	0.2	0.2	0.3	0.6	*
2001.....	0.1	*	*	*	*	*	*	0.1	*	0.3	0.6	*
2000.....	0.1	*	*	*	*	*	*	*	0.3	0.4	0.7	*
1999.....	0.1	*	*	*	*	*	*	0.2	*	0.3	0.7	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Systemic lupus erythematosus (M32)												
White, female												
2004.....	0.5	*	*	*	0.2	0.3	0.4	0.6	0.9	1.1	1.7	1.4
2003.....	0.6	*	*	*	0.2	0.4	0.5	0.7	1.1	1.1	2.1	1.3
2002.....	0.6	*	*	*	0.2	0.2	0.4	0.7	1.1	1.2	1.8	1.8
2001.....	0.6	*	*	*	0.3	0.3	0.4	0.8	1.1	1.5	2.1	1.1
2000.....	0.6	*	*	*	0.2	0.3	0.6	0.8	1.1	1.2	2.0	1.5
1999.....	0.6	*	*	*	0.2	0.3	0.5	0.8	1.1	1.5	1.8	1.7
Black, both sexes												
2004.....	1.2	*	*	*	0.7	1.5	1.7	2.3	2.7	2.0	2.3	*
2003.....	1.1	*	*	*	0.8	1.2	1.4	2.3	2.1	2.0	*	*
2002.....	1.2	*	*	*	0.8	1.3	1.9	2.0	2.4	2.5	*	*
2001.....	1.2	*	*	*	0.8	1.5	2.0	2.0	2.1	1.7	*	*
2000.....	1.2	*	*	*	0.8	1.3	2.0	2.3	2.4	1.8	*	*
1999.....	1.2	*	*	*	0.8	1.2	2.2	2.2	1.8	2.9	*	*
Black, male												
2004.....	0.4	*	*	*	*	*	*	*	*	*	*	*
2003.....	0.2	*	*	*	*	*	*	*	*	*	*	*
2002.....	0.2	*	*	*	*	*	*	*	*	*	*	*
2001.....	0.3	*	*	*	*	*	*	*	*	*	*	*
2000.....	0.4	*	*	*	*	*	0.7	*	*	*	*	*
1999.....	0.3	*	*	*	*	*	*	*	*	*	*	*
Black, female												
2004.....	2.0	*	*	*	1.1	2.4	2.5	3.6	4.4	3.0	*	*
2003.....	1.9	*	*	*	1.5	2.1	2.4	3.8	3.3	3.3	*	*
2002.....	2.1	*	*	*	1.3	2.2	3.5	3.4	4.2	3.7	*	*
2001.....	2.0	*	*	*	1.4	2.4	3.3	3.3	3.1	2.6	*	*
2000.....	2.0	*	*	*	1.2	2.2	3.2	4.0	3.7	2.8	*	*
1999.....	2.0	*	*	*	1.3	2.0	3.6	3.7	2.9	4.4	*	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other arthropathies and related disorders (M00,M02,M10-M31,M33-M35)

All races 3/, both sexes

2004.....	1.6	*	*	*	0.1	0.2	0.5	1.1	2.2	4.3	9.5	24.5
2003.....	1.6	*	*	*	0.1	0.3	0.6	1.1	2.2	4.6	9.4	25.9
2002.....	1.7	*	*	*	0.1	0.3	0.6	1.2	2.2	4.7	9.7	27.2
2001.....	1.7	*	*	*	0.1	0.2	0.6	1.1	2.1	5.0	10.1	27.7
2000.....	1.7	*	*	*	0.1	0.3	0.6	1.1	2.2	5.0	10.5	28.0
1999.....	1.7	*	*	*	0.1	0.2	0.6	1.3	2.3	5.0	9.5	27.6

All races 3/, male

2004.....	1.0	*	*	*	*	0.2	0.3	0.8	1.5	2.9	7.2	17.5
2003.....	1.0	*	*	*	*	0.2	0.4	0.7	1.5	3.5	7.7	19.4
2002.....	1.0	*	*	*	*	0.2	0.4	0.8	1.6	3.3	7.6	18.6
2001.....	1.0	*	*	*	*	0.2	0.4	0.9	1.5	3.4	8.1	20.5
2000.....	1.1	*	*	*	*	0.3	0.4	0.8	1.6	3.4	8.8	21.1
1999.....	1.0	*	*	*	*	0.2	0.5	0.8	1.6	3.5	7.8	22.5

All races 3/, female

2004.....	2.2	*	*	*	0.1	0.3	0.8	1.4	2.8	5.6	11.1	27.7
2003.....	2.2	*	*	*	0.2	0.4	0.7	1.5	2.9	5.6	10.5	28.8
2002.....	2.3	*	*	*	0.1	0.3	0.7	1.5	2.8	5.9	11.1	31.0
2001.....	2.3	*	*	*	0.2	0.3	0.8	1.3	2.7	6.3	11.5	30.7
2000.....	2.3	*	*	*	0.1	0.4	0.7	1.5	2.8	6.3	11.6	30.8
1999.....	2.3	*	*	*	0.2	0.3	0.7	1.7	3.0	6.3	10.6	29.7

White, both sexes

2004.....	1.7	*	*	*	0.1	0.1	0.4	0.8	2.0	4.3	9.8	25.5
2003.....	1.7	*	*	*	0.1	0.1	0.4	0.8	2.0	4.5	9.5	26.4
2002.....	1.7	*	*	*	*	0.2	0.4	1.0	2.1	4.7	9.9	28.0
2001.....	1.8	*	*	*	0.1	0.2	0.5	0.8	2.0	4.9	10.5	28.6
2000.....	1.7	*	*	*	0.1	0.2	0.4	0.9	2.0	4.9	10.7	28.8
1999.....	1.7	*	*	*	0.1	0.2	0.4	0.9	2.0	5.0	9.7	28.4

White, male

2004.....	1.0	*	*	*	*	*	0.3	0.6	1.4	2.8	7.4	18.0
2003.....	1.0	*	*	*	*	*	0.3	0.6	1.3	3.3	7.5	19.9
2002.....	1.0	*	*	*	*	*	0.3	0.6	1.4	3.2	7.8	19.1
2001.....	1.1	*	*	*	*	0.1	0.3	0.7	1.4	3.3	8.3	21.1
2000.....	1.0	*	*	*	*	0.2	0.2	0.6	1.5	3.2	8.7	20.6
1999.....	1.0	*	*	*	*	*	0.3	0.7	1.4	3.5	7.9	23.3

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other arthropathies and related disorders (M00,M02,M10-M31,M33-M35)												
White, female												
2004.....	2.3	*	*	*	*	0.2	0.5	1.1	2.5	5.6	11.4	28.9
2003.....	2.3	*	*	*	*	0.2	0.5	1.1	2.7	5.6	10.8	29.2
2002.....	2.4	*	*	*	*	0.2	0.5	1.3	2.8	6.0	11.3	31.9
2001.....	2.4	*	*	*	0.1	0.2	0.6	1.0	2.5	6.2	12.0	31.7
2000.....	2.4	*	*	*	*	0.3	0.5	1.1	2.6	6.4	11.9	32.1
1999.....	2.3	*	*	*	*	0.2	0.5	1.2	2.7	6.2	10.9	30.4
Black, both sexes												
2004.....	1.7	*	*	*	*	0.8	1.6	3.2	3.9	5.3	7.7	16.6
2003.....	1.8	*	*	*	*	1.0	1.5	3.2	4.2	6.0	9.2	20.4
2002.....	1.7	*	*	*	*	0.8	1.8	2.8	3.8	5.5	8.7	21.7
2001.....	1.7	*	*	*	*	0.8	1.8	3.1	3.2	6.1	8.5	22.4
2000.....	1.8	*	*	*	*	0.8	1.9	3.1	3.9	5.7	9.3	21.9
1999.....	1.9	*	*	*	*	0.7	2.0	3.7	4.7	6.1	7.8	20.4
Black, male												
2004.....	1.1	*	*	*	*	*	0.7	2.4	2.9	3.9	5.8	*
2003.....	1.2	*	*	*	*	*	1.3	1.8	3.3	5.9	10.4	*
2002.....	1.2	*	*	*	*	*	1.3	2.1	3.8	4.9	6.3	*
2001.....	1.2	*	*	*	*	*	1.4	2.8	2.3	4.3	7.4	*
2000.....	1.3	*	*	*	*	*	1.4	1.7	3.3	5.0	10.6	27.7
1999.....	1.3	*	*	*	*	*	1.8	2.3	3.5	4.6	8.0	*
Black, female												
2004.....	2.2	*	*	*	*	0.8	2.3	3.9	4.8	6.3	8.8	17.7
2003.....	2.3	*	*	*	*	1.3	1.7	4.3	5.0	6.0	8.5	23.5
2002.....	2.2	*	*	*	*	1.0	2.2	3.4	3.8	6.0	10.1	24.6
2001.....	2.2	*	*	*	*	1.1	2.1	3.5	3.8	7.3	9.1	24.4
2000.....	2.2	*	*	*	*	0.9	2.4	4.3	4.4	6.3	8.5	19.7
1999.....	2.4	*	*	*	*	1.0	2.1	4.8	5.7	7.2	7.7	22.3

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Dorsopathies (M40-M54)												
All races 3/, both sexes												
2004.....	0.4	*	*	*	*	*	0.1	0.1	0.4	0.9	2.6	7.1
2003.....	0.3	*	*	*	*	*	0.1	0.2	0.4	0.9	2.7	5.7
2002.....	0.3	*	*	*	*	*	0.1	0.1	0.4	0.9	2.2	6.0
2001.....	0.3	*	*	*	*	*	0.1	0.1	0.3	0.8	2.4	7.4
2000.....	0.3	*	*	*	*	*	0.0	0.1	0.3	0.8	2.4	6.1
1999.....	0.3	*	*	*	*	*	0.0	0.2	0.3	0.8	1.9	5.4
All races 3/, male												
2004.....	0.3	*	*	*	*	*	*	0.2	0.4	0.9	2.7	6.6
2003.....	0.3	*	*	*	*	*	*	0.2	0.4	1.0	2.8	4.8
2002.....	0.3	*	*	*	*	*	*	0.2	0.4	0.9	2.4	5.5
2001.....	0.3	*	*	*	*	*	0.1	0.1	0.3	0.7	2.5	6.4
2000.....	0.3	*	*	*	*	*	*	0.2	0.3	1.0	2.4	6.5
1999.....	0.2	*	*	*	*	*	*	0.1	0.3	0.8	2.2	5.9
All races 3/, female												
2004.....	0.4	*	*	*	*	*	*	0.1	0.3	0.8	2.5	7.3
2003.....	0.4	*	*	*	*	*	*	0.1	0.4	0.8	2.6	6.1
2002.....	0.4	*	*	*	*	*	*	*	0.3	0.9	2.1	6.2
2001.....	0.4	*	*	*	*	*	*	0.1	0.3	0.8	2.3	7.8
2000.....	0.3	*	*	*	*	*	*	*	0.3	0.6	2.4	6.0
1999.....	0.3	*	*	*	*	*	*	0.2	0.2	0.7	1.7	5.2
White, both sexes												
2004.....	0.4	*	*	*	*	*	*	0.1	0.4	0.8	2.7	7.4
2003.....	0.4	*	*	*	*	*	*	0.2	0.3	0.8	2.7	5.9
2002.....	0.4	*	*	*	*	*	*	0.1	0.3	0.9	2.3	6.2
2001.....	0.4	*	*	*	*	*	0.1	0.1	0.3	0.7	2.3	7.8
2000.....	0.3	*	*	*	*	*	*	0.1	0.3	0.8	2.4	6.4
1999.....	0.3	*	*	*	*	*	*	0.2	0.2	0.7	2.0	5.5
White, male												
2004.....	0.3	*	*	*	*	*	*	0.1	0.4	0.8	2.7	6.8
2003.....	0.3	*	*	*	*	*	*	0.2	0.3	0.9	2.8	4.8
2002.....	0.3	*	*	*	*	*	*	0.2	0.3	0.9	2.4	5.3
2001.....	0.3	*	*	*	*	*	*	0.1	0.3	0.7	2.5	6.6
2000.....	0.3	*	*	*	*	*	*	0.2	0.3	0.9	2.3	6.9
1999.....	0.3	*	*	*	*	*	*	0.2	0.3	0.8	2.2	5.8

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Soft tissue disorders (M60-M79)												
All races 3/, both sexes												
2004.....	0.4	*	*	*	*	0.1	0.1	0.4	0.6	1.2	2.7	5.1
2003.....	0.4	*	*	*	*	0.1	0.2	0.4	0.6	1.3	2.6	5.4
2002.....	0.4	*	*	*	*	0.1	0.2	0.3	0.5	1.1	2.4	5.3
2001.....	0.4	*	*	*	*	0.1	0.2	0.3	0.6	1.0	2.3	4.3
2000.....	0.4	*	*	*	*	0.1	0.1	0.4	0.5	1.0	2.1	4.2
1999.....	0.3	*	*	*	*	0.1	0.2	0.3	0.5	1.0	2.1	4.3
All races 3/, male												
2004.....	0.5	*	*	*	*	*	0.2	0.5	0.7	1.2	3.4	7.8
2003.....	0.4	*	*	*	*	0.1	0.2	0.4	0.6	1.5	3.4	7.1
2002.....	0.4	*	*	*	*	*	0.2	0.4	0.6	1.4	2.9	7.8
2001.....	0.4	*	*	*	*	*	0.2	0.4	0.6	1.4	3.2	6.2
2000.....	0.4	*	*	*	*	*	0.2	0.4	0.6	1.2	2.9	6.7
1999.....	0.4	*	*	*	*	*	0.2	0.3	0.6	1.3	3.1	6.8
All races 3/, female												
2004.....	0.4	*	*	*	*	*	0.1	0.3	0.5	1.2	2.2	3.9
2003.....	0.4	*	*	*	*	*	0.1	0.4	0.7	1.1	2.0	4.6
2002.....	0.4	*	*	*	*	*	0.1	0.3	0.5	0.9	2.1	4.2
2001.....	0.3	*	*	*	*	*	0.1	0.3	0.6	0.8	1.7	3.5
2000.....	0.3	*	*	*	*	*	0.1	0.3	0.5	0.9	1.6	3.2
1999.....	0.3	*	*	*	*	*	0.1	0.2	0.5	0.8	1.5	3.3
White, both sexes												
2004.....	0.4	*	*	*	*	*	0.1	0.4	0.5	1.1	2.6	4.8
2003.....	0.4	*	*	*	*	0.1	0.2	0.3	0.6	1.2	2.5	5.4
2002.....	0.4	*	*	*	*	*	0.1	0.3	0.5	1.0	2.4	5.0
2001.....	0.4	*	*	*	*	*	0.2	0.3	0.5	0.9	2.2	4.2
2000.....	0.3	*	*	*	*	0.1	0.1	0.3	0.5	0.9	1.9	4.0
1999.....	0.3	*	*	*	*	0.1	0.1	0.2	0.4	0.9	2.0	4.0
White, male												
2004.....	0.4	*	*	*	*	*	0.1	0.4	0.6	1.1	3.3	7.2
2003.....	0.5	*	*	*	*	*	0.2	0.3	0.5	1.4	3.3	7.1
2002.....	0.4	*	*	*	*	*	0.2	0.4	0.5	1.3	2.8	7.6
2001.....	0.4	*	*	*	*	*	0.2	0.4	0.4	1.3	3.1	6.0
2000.....	0.4	*	*	*	*	*	0.1	0.3	0.6	1.1	2.8	6.3
1999.....	0.4	*	*	*	*	*	0.2	0.2	0.5	1.2	2.8	6.3

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Osteopathies, chondropathies and other disorders of musculoskeletal system and connective tissue (M80-M99)

All races 3/, both sexes

2004.....	1.1	*	*	*	*	*	0.1	0.2	0.6	1.7	7.4	36.2
2003.....	1.2	*	*	*	*	*	0.1	0.2	0.6	1.9	6.9	39.5
2002.....	1.2	*	*	*	*	*	0.1	0.2	0.6	1.9	7.0	40.8
2001.....	1.1	*	*	*	*	*	0.1	0.2	0.5	1.7	7.1	40.1
2000.....	1.1	*	*	*	*	*	0.1	0.1	0.4	1.4	6.6	41.6
1999.....	1.0	*	*	*	*	*	*	0.2	0.4	1.4	6.7	38.6

All races 3/, male

2004.....	0.7	*	*	*	*	*	*	0.3	0.6	1.9	6.0	20.0
2003.....	0.7	*	*	*	*	*	0.1	0.3	0.7	2.1	5.2	21.5
2002.....	0.6	*	*	*	*	*	*	0.2	0.6	2.1	5.3	23.6
2001.....	0.6	*	*	*	*	*	0.1	0.2	0.6	1.7	5.4	23.6
2000.....	0.6	*	*	*	*	*	0.1	0.2	0.5	1.5	5.3	25.6
1999.....	0.5	*	*	*	*	*	*	0.2	0.4	1.3	5.3	20.4

All races 3/, female

2004.....	1.6	*	*	*	*	*	*	0.2	0.6	1.6	8.3	43.5
2003.....	1.7	*	*	*	*	*	*	0.2	0.5	1.7	8.1	47.5
2002.....	1.7	*	*	*	*	*	*	0.1	0.5	1.7	8.1	48.3
2001.....	1.6	*	*	*	*	*	*	0.1	0.4	1.7	8.2	47.0
2000.....	1.6	*	*	*	*	*	*	0.1	0.4	1.4	7.5	48.1
1999.....	1.5	*	*	*	*	*	*	0.1	0.5	1.5	7.7	45.9

White, both sexes

2004.....	1.3	*	*	*	*	*	0.1	0.2	0.6	1.7	7.4	37.7
2003.....	1.3	*	*	*	*	*	0.1	0.2	0.6	1.8	7.0	41.0
2002.....	1.3	*	*	*	*	*	0.1	0.2	0.5	1.8	6.9	42.8
2001.....	1.2	*	*	*	*	*	0.1	0.2	0.5	1.6	7.2	41.3
2000.....	1.2	*	*	*	*	*	0.1	0.1	0.4	1.4	6.7	43.0
1999.....	1.1	*	*	*	*	*	*	0.2	0.4	1.3	6.8	40.2

White, male

2004.....	0.7	*	*	*	*	*	*	0.3	0.6	1.7	5.7	20.4
2003.....	0.7	*	*	*	*	*	*	0.2	0.6	1.8	5.0	20.9
2002.....	0.7	*	*	*	*	*	*	0.2	0.6	2.0	5.0	23.8
2001.....	0.6	*	*	*	*	*	*	0.2	0.5	1.5	5.3	23.8
2000.....	0.6	*	*	*	*	*	*	0.1	0.4	1.3	5.1	25.4
1999.....	0.5	*	*	*	*	*	*	0.2	0.4	1.2	5.1	20.2

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Osteopathies, chondropathies and other disorders of musculoskeletal system and connective tissue (M80-M99)

White, female

2004.....	1.8	*	*	*	*	*	*	0.1	0.5	1.7	8.5	45.5
2003.....	1.9	*	*	*	*	*	*	0.2	0.5	1.8	8.4	49.9
2002.....	1.9	*	*	*	*	*	*	0.1	0.5	1.7	8.2	51.0
2001.....	1.8	*	*	*	*	*	*	0.1	0.5	1.7	8.4	48.6
2000.....	1.8	*	*	*	*	*	*	*	0.3	1.4	7.7	50.2
1999.....	1.7	*	*	*	*	*	*	0.2	0.5	1.4	7.9	48.2

Black, both sexes

2004.....	0.7	*	*	*	*	*	*	0.5	0.9	2.6	9.1	24.8
2003.....	0.7	*	*	*	*	*	*	0.5	1.2	3.0	7.3	26.5
2002.....	0.7	*	*	*	*	*	*	*	1.2	2.8	8.7	24.1
2001.....	0.7	*	*	*	*	*	*	*	1.0	3.3	6.8	29.8
2000.....	0.7	*	*	*	*	*	*	*	1.1	2.5	7.5	27.5
1999.....	0.6	*	*	*	*	*	*	*	*	2.3	7.0	21.4

Black, male

2004.....	0.7	*	*	*	*	*	*	*	*	3.3	11.1	21.9
2003.....	0.8	*	*	*	*	*	*	*	1.7	5.2	9.9	27.1
2002.....	0.6	*	*	*	*	*	*	*	*	4.0	10.0	25.7
2001.....	0.6	*	*	*	*	*	*	*	*	3.8	7.7	26.1
2000.....	0.7	*	*	*	*	*	*	*	*	4.0	8.5	32.3
1999.....	0.4	*	*	*	*	*	*	*	*	*	7.1	24.6

Black, female

2004.....	0.8	*	*	*	*	*	*	*	*	2.1	8.0	26.0
2003.....	0.7	*	*	*	*	*	*	*	*	*	5.9	26.3
2002.....	0.8	*	*	*	*	*	*	*	*	*	8.0	23.4
2001.....	0.8	*	*	*	*	*	*	*	*	3.0	6.3	31.2
2000.....	0.7	*	*	*	*	*	*	*	*	*	7.0	25.8
1999.....	0.6	*	*	*	*	*	*	*	*	2.1	6.9	20.2

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Osteoporosis (M80-M81)												
All races 3/, both sexes												
2004.....	0.6	*	*	*	*	*	*	*	0.1	0.3	3.2	24.4
2003.....	0.6	*	*	*	*	*	*	*	*	0.4	3.2	26.9
2002.....	0.6	*	*	*	*	*	*	*	*	0.4	3.4	29.0
2001.....	0.6	*	*	*	*	*	*	*	*	0.4	3.3	26.8
2000.....	0.6	*	*	*	*	*	*	*	*	0.4	3.4	28.5
1999.....	0.6	*	*	*	*	*	*	*	*	0.4	3.5	27.1
All races 3/, male												
2004.....	0.1	*	*	*	*	*	*	*	*	*	1.2	7.2
2003.....	0.1	*	*	*	*	*	*	*	*	*	1.0	7.9
2002.....	0.2	*	*	*	*	*	*	*	*	*	1.1	9.9
2001.....	0.1	*	*	*	*	*	*	*	*	*	1.0	9.3
2000.....	0.1	*	*	*	*	*	*	*	*	*	1.3	9.9
1999.....	0.1	*	*	*	*	*	*	*	*	*	1.3	7.0
All races 3/, female												
2004.....	1.0	*	*	*	*	*	*	*	*	0.5	4.6	32.2
2003.....	1.1	*	*	*	*	*	*	*	*	0.6	4.7	35.2
2002.....	1.1	*	*	*	*	*	*	*	*	0.6	4.9	37.2
2001.....	1.0	*	*	*	*	*	*	*	*	0.5	4.8	34.1
2000.....	1.1	*	*	*	*	*	*	*	*	0.5	4.9	36.0
1999.....	1.0	*	*	*	*	*	*	*	*	0.6	5.0	35.1
White, both sexes												
2004.....	0.7	*	*	*	*	*	*	*	*	0.4	3.5	26.3
2003.....	0.7	*	*	*	*	*	*	*	*	0.5	3.5	29.1
2002.....	0.8	*	*	*	*	*	*	*	*	0.5	3.6	31.3
2001.....	0.7	*	*	*	*	*	*	*	*	0.4	3.5	28.8
2000.....	0.7	*	*	*	*	*	*	*	*	0.4	3.7	30.7
1999.....	0.7	*	*	*	*	*	*	*	*	0.4	3.8	29.1
White, male												
2004.....	0.2	*	*	*	*	*	*	*	*	*	1.2	7.8
2003.....	0.2	*	*	*	*	*	*	*	*	*	1.1	8.2
2002.....	0.2	*	*	*	*	*	*	*	*	*	1.1	10.6
2001.....	0.2	*	*	*	*	*	*	*	*	*	1.1	10.0
2000.....	0.2	*	*	*	*	*	*	*	*	*	1.3	10.4
1999.....	0.1	*	*	*	*	*	*	*	*	*	1.4	7.4

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other osteopathies, chondropathies and disorders of musculoskeletal system and connective tissue (M83-M99)

All races 3/, both sexes

2004.....	0.6	*	*	*	*	*	0.1	0.2	0.5	1.4	4.1	11.8
2003.....	0.6	*	*	*	*	*	0.1	0.2	0.6	1.5	3.7	12.6
2002.....	0.5	*	*	*	*	*	0.0	0.2	0.5	1.5	3.6	11.8
2001.....	0.5	*	*	*	*	*	0.1	0.2	0.4	1.3	3.8	13.3
2000.....	0.5	*	*	*	*	*	0.1	0.1	0.4	1.1	3.2	13.1
1999.....	0.4	*	*	*	*	*	*	0.1	0.4	1.0	3.2	11.5

All races 3/, male

2004.....	0.5	*	*	*	*	*	*	0.3	0.5	1.7	4.8	12.9
2003.....	0.5	*	*	*	*	*	0.1	0.2	0.7	1.9	4.2	13.6
2002.....	0.5	*	*	*	*	*	*	0.2	0.6	1.9	4.2	13.7
2001.....	0.5	*	*	*	*	*	*	0.2	0.5	1.5	4.4	14.2
2000.....	0.5	*	*	*	*	*	0.1	0.1	0.5	1.4	4.1	15.6
1999.....	0.4	*	*	*	*	*	*	0.2	0.4	1.1	4.0	13.3

All races 3/, female

2004.....	0.6	*	*	*	*	*	*	0.1	0.5	1.2	3.7	11.3
2003.....	0.6	*	*	*	*	*	*	0.2	0.4	1.1	3.3	12.2
2002.....	0.6	*	*	*	*	*	*	0.1	0.5	1.1	3.2	11.1
2001.....	0.6	*	*	*	*	*	*	0.1	0.3	1.2	3.4	12.9
2000.....	0.5	*	*	*	*	*	*	*	0.3	0.9	2.6	12.0
1999.....	0.5	*	*	*	*	*	*	0.1	0.4	0.9	2.8	10.8

White, both sexes

2004.....	0.6	*	*	*	*	*	0.1	0.2	0.5	1.3	3.9	11.4
2003.....	0.6	*	*	*	*	*	0.1	0.2	0.5	1.3	3.5	11.9
2002.....	0.5	*	*	*	*	*	0.1	0.1	0.5	1.4	3.3	11.5
2001.....	0.5	*	*	*	*	*	0.1	0.2	0.4	1.2	3.6	12.5
2000.....	0.5	*	*	*	*	*	0.1	0.1	0.3	1.0	2.9	12.4
1999.....	0.4	*	*	*	*	*	*	0.1	0.4	0.9	3.0	11.1

White, male

2004.....	0.5	*	*	*	*	*	*	0.2	0.5	1.5	4.5	12.6
2003.....	0.5	*	*	*	*	*	*	0.2	0.6	1.6	3.9	12.7
2002.....	0.5	*	*	*	*	*	*	0.2	0.6	1.8	3.8	13.2
2001.....	0.5	*	*	*	*	*	*	0.2	0.4	1.3	4.2	13.8
2000.....	0.4	*	*	*	*	*	*	*	0.4	1.2	3.8	15.1
1999.....	0.4	*	*	*	*	*	*	0.1	0.4	1.0	3.8	12.8

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other osteopathies, chondropathies and disorders of musculoskeletal system and connective tissue (M83-M99)

White, female

2004.....	0.6	*	*	*	*	*	*	0.1	0.5	1.1	3.4	10.8
2003.....	0.6	*	*	*	*	*	*	0.2	0.4	1.1	3.2	11.5
2002.....	0.6	*	*	*	*	*	*	*	0.4	1.0	2.9	10.7
2001.....	0.6	*	*	*	*	*	*	0.1	0.3	1.1	3.2	12.0
2000.....	0.5	*	*	*	*	*	*	*	0.3	0.8	2.4	11.2
1999.....	0.5	*	*	*	*	*	*	0.1	0.4	0.7	2.5	10.4

Black, both sexes

2004.....	0.7	*	*	*	*	*	*	0.5	0.9	2.5	8.6	20.0
2003.....	0.7	*	*	*	*	*	*	0.4	1.2	3.0	6.8	23.3
2002.....	0.6	*	*	*	*	*	*	*	1.2	2.7	7.7	19.6
2001.....	0.6	*	*	*	*	*	*	*	0.9	3.2	6.3	24.3
2000.....	0.6	*	*	*	*	*	*	*	1.0	2.4	7.0	23.8
1999.....	0.5	*	*	*	*	*	*	*	*	2.2	6.2	17.5

Black, male

2004.....	0.7	*	*	*	*	*	*	*	*	3.3	10.6	20.9
2003.....	0.7	*	*	*	*	*	*	*	1.7	5.2	9.3	24.0
2002.....	0.6	*	*	*	*	*	*	*	*	4.0	9.8	24.7
2001.....	0.6	*	*	*	*	*	*	*	*	3.8	7.7	23.8
2000.....	0.6	*	*	*	*	*	*	*	*	3.8	7.9	27.7
1999.....	0.4	*	*	*	*	*	*	*	*	*	6.8	*

Black, female

2004.....	0.7	*	*	*	*	*	*	*	*	2.0	7.6	19.7
2003.....	0.7	*	*	*	*	*	*	*	*	*	5.4	23.1
2002.....	0.6	*	*	*	*	*	*	*	*	*	6.5	17.7
2001.....	0.7	*	*	*	*	*	*	*	*	2.8	5.6	24.4
2000.....	0.6	*	*	*	*	*	*	*	*	*	6.5	22.3
1999.....	0.6	*	*	*	*	*	*	*	*	2.1	5.9	15.8

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
XIV. Diseases of the genitourinary system (N00-N98)												
All races 3/, both sexes												
2004.....	20.5	4.9	0.2	0.1	0.3	0.8	2.2	6.2	16.5	49.2	153.4	458.5
2003.....	20.7	5.1	0.1	0.1	0.3	0.9	2.2	6.0	16.7	50.7	153.8	469.5
2002.....	20.2	4.6	*	0.1	0.3	0.9	2.1	5.7	15.8	49.7	153.5	469.4
2001.....	20.0	3.9	0.2	*	0.3	0.8	2.1	5.7	15.8	51.2	149.8	476.4
2000.....	19.4	5.0	0.1	0.1	0.3	0.8	2.0	5.4	15.6	48.0	148.2	471.4
1999.....	19.0	5.1	0.2	0.1	0.2	0.8	2.0	5.0	15.1	48.1	144.7	471.1
All races 3/, male												
2004.....	18.5	5.8	*	*	0.3	0.7	2.4	7.0	18.2	53.8	179.0	533.9
2003.....	18.6	6.0	*	*	0.3	0.9	2.4	6.8	18.3	55.3	181.1	547.9
2002.....	18.1	5.6	*	*	0.2	0.8	2.1	6.5	17.0	55.9	178.6	553.0
2001.....	17.7	4.2	*	*	0.3	0.7	2.2	6.4	16.7	56.9	173.5	564.2
2000.....	17.2	6.1	*	*	0.3	0.8	2.2	6.3	16.8	52.6	173.2	567.5
1999.....	16.9	5.0	*	0.1	0.2	0.8	2.1	5.6	16.5	53.3	172.1	558.3
All races 3/, female												
2004.....	22.5	4.0	*	*	0.3	0.9	2.1	5.4	15.0	45.4	136.3	424.5
2003.....	22.6	4.3	*	*	0.3	0.9	2.1	5.2	15.3	46.7	135.6	434.9
2002.....	22.3	3.6	*	*	0.3	0.9	2.0	5.0	14.8	44.5	136.8	433.2
2001.....	22.2	3.6	*	*	0.3	0.8	2.0	5.1	15.0	46.5	134.3	439.6
2000.....	21.5	3.9	*	*	0.3	0.8	1.8	4.6	14.5	44.3	132.0	432.3
1999.....	21.1	5.1	*	0.1	0.3	0.8	1.8	4.5	13.8	43.9	127.0	436.0
White, both sexes												
2004.....	20.6	3.6	*	0.1	0.2	0.6	1.7	4.5	12.9	43.0	145.5	449.9
2003.....	20.7	4.1	*	0.1	0.3	0.7	1.5	4.3	13.2	43.9	145.8	461.3
2002.....	20.3	4.0	*	0.1	0.2	0.6	1.5	4.1	12.4	43.3	145.5	458.5
2001.....	20.0	3.0	*	*	0.2	0.6	1.6	4.1	12.3	44.5	141.1	465.1
2000.....	19.4	3.6	*	0.1	0.3	0.6	1.4	3.9	12.2	41.2	139.9	460.5
1999.....	18.9	4.0	*	0.1	0.2	0.6	1.4	3.5	11.9	41.4	135.9	456.0
White, male												
2004.....	18.5	4.2	*	*	0.2	0.4	1.9	5.0	14.4	47.4	172.1	527.3
2003.....	18.6	4.8	*	*	0.3	0.7	1.7	4.8	14.4	48.3	173.0	547.4
2002.....	18.1	4.7	*	*	0.2	0.6	1.6	4.6	13.1	49.4	171.1	543.2
2001.....	17.7	3.4	*	*	0.2	0.5	1.8	4.5	13.2	50.2	165.9	556.8
2000.....	17.2	4.6	*	*	0.3	0.6	1.6	4.4	13.2	46.4	165.2	561.2
1999.....	16.6	4.5	*	0.1	0.2	0.6	1.4	3.9	13.3	46.3	162.2	542.8

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
XIV. Diseases of the genitourinary system (N00-N98)												
White, female												
2004.....	22.7	2.8	*	*	0.2	0.7	1.5	3.9	11.5	39.3	127.4	415.0
2003.....	22.8	3.4	*	*	0.3	0.7	1.4	3.8	12.1	40.1	127.4	423.2
2002.....	22.5	3.2	*	*	0.2	0.6	1.4	3.6	11.8	38.2	128.3	421.7
2001.....	22.3	2.6	*	*	0.3	0.7	1.4	3.6	11.4	39.6	124.7	426.7
2000.....	21.5	2.6	*	*	0.3	0.6	1.3	3.4	11.2	36.8	123.3	419.6
1999.....	21.2	3.4	*	*	0.2	0.6	1.3	3.2	10.6	37.2	118.7	421.1
Black, both sexes												
2004.....	25.7	11.6	*	*	0.6	2.2	6.3	19.3	50.7	113.8	266.5	634.5
2003.....	26.0	11.1	*	*	0.7	2.5	7.1	18.8	50.2	121.5	270.3	645.3
2002.....	25.2	8.3	*	*	0.7	2.5	6.2	18.3	47.0	117.6	267.8	662.5
2001.....	25.1	9.2	*	*	0.5	2.1	6.1	18.3	47.2	122.0	272.0	664.5
2000.....	24.4	12.7	*	*	0.5	2.1	6.2	17.8	46.3	118.9	267.2	637.5
1999.....	24.3	11.1	*	*	0.6	2.2	6.2	17.5	43.3	118.2	264.7	679.1
Black, male												
2004.....	23.4	13.0	*	*	0.6	2.2	7.0	23.4	56.8	128.0	301.5	718.3
2003.....	24.0	12.5	*	*	0.7	2.7	7.6	23.4	58.6	139.5	321.3	681.9
2002.....	23.2	10.5	*	*	*	2.2	6.5	22.1	55.4	134.0	307.5	794.4
2001.....	22.2	9.3	*	*	*	2.6	5.8	21.7	51.7	135.7	301.3	755.7
2000.....	21.9	14.1	*	*	*	2.1	7.2	22.5	51.8	125.7	309.5	716.2
1999.....	22.4	8.1	*	*	*	1.9	7.2	21.2	47.5	132.5	328.1	802.7
Black, female												
2004.....	27.7	10.2	*	*	*	2.2	5.8	15.7	45.7	103.6	246.7	601.3
2003.....	27.7	9.6	*	*	0.7	2.4	6.6	14.8	43.5	108.7	241.4	631.1
2002.....	27.0	6.1	*	*	0.9	2.7	6.0	15.1	40.2	106.0	245.2	612.0
2001.....	27.8	9.1	*	*	*	1.6	6.4	15.4	43.6	112.3	255.5	630.6
2000.....	26.6	11.2	*	*	*	2.0	5.3	13.8	41.8	114.2	243.4	608.3
1999.....	26.1	14.1	*	*	0.7	2.4	5.3	14.3	39.9	108.2	229.3	633.0

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diseases of urinary system (N00-N39)												
All races 3/, both sexes												
2004.....	20.2	4.8	0.2	0.1	0.3	0.8	2.2	6.0	16.3	48.5	150.8	448.9
2003.....	20.3	5.0	0.1	0.1	0.3	0.9	2.1	5.9	16.5	50.0	151.5	460.5
2002.....	19.9	4.6	*	0.1	0.3	0.8	2.0	5.6	15.7	49.1	151.2	460.8
2001.....	19.7	3.7	0.2	*	0.3	0.7	2.1	5.6	15.6	50.5	147.6	467.4
2000.....	19.1	4.8	0.1	0.1	0.3	0.7	2.0	5.3	15.4	47.4	145.9	462.5
1999.....	18.7	4.7	0.1	0.1	0.2	0.8	1.9	4.9	14.8	47.5	142.3	462.6
All races 3/, male												
2004.....	18.0	5.8	*	*	0.3	0.7	2.4	6.9	17.8	52.8	174.3	513.6
2003.....	18.2	5.9	*	*	0.3	0.9	2.4	6.8	18.1	54.4	177.2	528.3
2002.....	17.7	5.6	*	*	0.2	0.8	2.1	6.4	16.8	55.0	174.6	533.8
2001.....	17.3	4.2	*	*	0.3	0.7	2.2	6.3	16.5	56.0	169.8	542.9
2000.....	16.8	6.0	*	*	0.3	0.8	2.2	6.2	16.5	51.7	169.2	546.9
1999.....	16.4	5.0	*	0.1	0.2	0.8	2.1	5.5	16.2	52.4	167.7	537.0
All races 3/, female												
2004.....	22.2	3.8	*	*	0.2	0.9	2.0	5.2	14.8	44.9	135.1	419.9
2003.....	22.4	4.1	*	*	0.3	0.9	1.9	5.0	15.1	46.2	134.3	430.5
2002.....	22.0	3.5	*	*	0.3	0.9	1.9	4.9	14.6	44.2	135.7	429.1
2001.....	22.0	3.2	*	*	0.3	0.8	1.9	4.9	14.8	46.0	133.0	435.8
2000.....	21.2	3.6	*	*	0.3	0.7	1.7	4.4	14.3	43.8	130.6	428.1
1999.....	20.9	4.3	*	0.1	0.3	0.8	1.7	4.4	13.5	43.4	125.7	432.6
White, both sexes												
2004.....	20.2	3.5	*	0.1	0.2	0.5	1.6	4.4	12.6	42.3	143.0	440.3
2003.....	20.4	4.1	*	0.1	0.2	0.7	1.5	4.2	13.0	43.2	143.4	451.9
2002.....	20.0	3.9	*	0.1	0.2	0.6	1.4	4.0	12.3	42.8	143.3	449.9
2001.....	19.7	2.8	*	*	0.2	0.6	1.5	4.0	12.1	43.7	138.9	456.0
2000.....	19.0	3.5	*	0.1	0.3	0.6	1.4	3.8	12.0	40.5	137.6	451.6
1999.....	18.6	3.7	*	0.1	0.2	0.6	1.3	3.5	11.7	40.8	133.5	447.4
White, male												
2004.....	18.0	4.2	*	*	0.2	0.4	1.8	4.9	14.1	46.4	167.5	506.5
2003.....	18.1	4.8	*	*	0.3	0.7	1.7	4.8	14.2	47.4	169.1	527.0
2002.....	17.7	4.7	*	*	0.2	0.6	1.6	4.6	12.9	48.6	167.3	523.8
2001.....	17.3	3.3	*	*	0.2	0.5	1.8	4.4	13.0	49.3	162.2	535.0
2000.....	16.7	4.5	*	*	0.3	0.6	1.6	4.4	13.0	45.6	161.4	540.8
1999.....	16.2	4.5	*	0.1	0.2	0.6	1.4	3.8	13.1	45.4	157.8	521.3

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diseases of urinary system (N00-N39)												
White, female												
2004.....	22.4	2.8	*	*	0.2	0.7	1.4	3.8	11.3	38.9	126.2	410.4
2003.....	22.5	3.3	*	*	0.2	0.7	1.3	3.7	11.9	39.6	126.0	418.6
2002.....	22.2	3.0	*	*	0.2	0.6	1.3	3.5	11.6	37.9	127.2	417.8
2001.....	22.0	2.3	*	*	0.2	0.6	1.3	3.5	11.2	39.1	123.4	423.0
2000.....	21.3	2.4	*	*	0.3	0.5	1.2	3.2	11.1	36.3	121.9	415.4
1999.....	20.9	2.8	*	*	0.2	0.6	1.2	3.1	10.4	36.9	117.5	417.7
Black, both sexes												
2004.....	25.3	11.2	*	*	0.5	2.2	6.2	18.8	50.2	112.1	263.3	625.5
2003.....	25.7	10.6	*	*	0.6	2.4	6.9	18.5	49.7	120.4	268.1	638.9
2002.....	24.8	8.3	*	*	0.7	2.4	6.0	18.1	46.6	116.2	264.9	652.1
2001.....	24.8	8.9	*	*	0.5	2.1	5.8	17.9	46.5	121.1	268.5	655.3
2000.....	24.1	12.2	*	*	0.4	2.0	5.9	17.7	45.8	117.7	264.2	628.5
1999.....	23.9	9.9	*	*	0.6	2.1	6.0	17.1	42.7	116.5	260.9	669.3
Black, male												
2004.....	23.0	13.0	*	*	0.6	2.2	6.9	23.0	56.2	126.1	294.8	701.4
2003.....	23.7	12.2	*	*	0.7	2.7	7.6	23.2	57.9	138.3	316.5	666.2
2002.....	22.8	10.5	*	*	*	2.2	6.4	21.9	54.9	131.7	300.6	776.2
2001.....	21.8	9.3	*	*	*	2.6	5.6	21.4	50.9	134.5	295.7	736.4
2000.....	21.6	14.1	*	*	*	2.1	7.1	22.4	50.9	124.1	303.7	695.4
1999.....	22.0	8.1	*	*	*	1.9	7.2	20.9	46.6	130.2	322.2	779.2
Black, female												
2004.....	27.4	9.3	*	*	*	2.2	5.6	15.2	45.5	102.2	245.5	595.4
2003.....	27.4	9.0	*	*	*	2.2	6.2	14.4	43.0	107.7	240.8	628.3
2002.....	26.6	6.1	*	*	0.9	2.6	5.6	14.7	39.9	105.2	244.6	604.6
2001.....	27.5	8.5	*	*	*	1.6	6.1	14.9	43.0	111.6	253.2	625.2
2000.....	26.3	10.3	*	*	*	1.9	4.9	13.6	41.6	113.3	241.9	603.5
1999.....	25.7	11.7	*	*	*	2.2	5.0	13.9	39.5	106.9	226.5	628.2

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Glomerular and renal tubulo-interstitial diseases (N00-N15)

All races 3/, both sexes

2004.....	0.7	*	*	*	*	0.1	0.2	0.3	0.5	1.4	4.7	12.6
2003.....	0.7	*	*	*	0.1	0.1	0.1	0.3	0.5	1.5	4.8	13.3
2002.....	0.7	*	*	*	*	0.1	0.1	0.3	0.5	1.5	4.9	13.9
2001.....	0.6	*	*	*	*	0.1	0.2	0.3	0.5	1.6	4.6	13.7
2000.....	0.7	0.5	*	*	*	0.1	0.1	0.3	0.6	1.5	4.8	15.7
1999.....	0.7	*	*	*	*	0.1	0.1	0.3	0.7	1.7	5.4	16.4

All races 3/, male

2004.....	0.6	*	*	*	*	*	0.1	0.2	0.7	1.5	5.2	16.9
2003.....	0.6	*	*	*	*	*	0.1	0.3	0.5	1.4	5.2	17.8
2002.....	0.6	*	*	*	*	*	0.1	0.3	0.6	1.7	5.5	18.1
2001.....	0.6	*	*	*	*	*	0.2	0.2	0.5	1.8	5.3	17.2
2000.....	0.6	*	*	*	*	*	0.1	0.3	0.6	1.4	5.4	20.0
1999.....	0.7	*	*	*	*	*	0.1	0.3	0.7	1.6	6.7	22.6

All races 3/, female

2004.....	0.7	*	*	*	*	0.1	0.2	0.3	0.4	1.4	4.5	10.7
2003.....	0.7	*	*	*	*	0.1	0.1	0.3	0.6	1.6	4.5	11.3
2002.....	0.7	*	*	*	*	*	0.1	0.3	0.5	1.4	4.5	12.0
2001.....	0.7	*	*	*	*	*	0.2	0.3	0.5	1.4	4.1	12.3
2000.....	0.8	*	*	*	*	*	0.1	0.3	0.6	1.7	4.5	13.9
1999.....	0.8	*	*	*	*	*	0.2	0.4	0.6	1.9	4.6	13.9

White, both sexes

2004.....	0.7	*	*	*	*	0.1	0.1	0.3	0.5	1.4	4.7	12.4
2003.....	0.7	*	*	*	*	0.1	0.1	0.3	0.5	1.5	4.6	13.2
2002.....	0.7	*	*	*	*	0.1	0.1	0.2	0.5	1.6	4.8	13.9
2001.....	0.7	*	*	*	*	0.1	0.2	0.2	0.4	1.5	4.5	13.9
2000.....	0.7	*	*	*	*	0.1	0.1	0.2	0.5	1.4	4.6	15.2
1999.....	0.8	*	*	*	*	0.1	0.1	0.3	0.6	1.6	5.3	16.3

White, male

2004.....	0.6	*	*	*	*	*	0.1	0.2	0.6	1.4	5.1	16.5
2003.....	0.6	*	*	*	*	*	0.1	0.3	0.4	1.4	4.9	18.0
2002.....	0.6	*	*	*	*	*	0.1	0.2	0.5	1.7	5.3	18.4
2001.....	0.6	*	*	*	*	*	0.2	0.2	0.4	1.7	5.2	17.5
2000.....	0.6	*	*	*	*	*	*	0.2	0.5	1.3	5.3	19.5
1999.....	0.7	*	*	*	*	*	0.1	0.3	0.6	1.3	6.6	22.1

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Glomerular and renal tubulo-interstitial diseases (N00-N15)												
White, female												
2004.....	0.8	*	*	*	*	*	0.2	0.3	0.4	1.4	4.4	10.6
2003.....	0.8	*	*	*	*	*	0.1	0.2	0.5	1.6	4.4	11.0
2002.....	0.8	*	*	*	*	*	0.1	0.2	0.5	1.5	4.5	11.9
2001.....	0.8	*	*	*	*	*	0.1	0.3	0.5	1.4	4.1	12.4
2000.....	0.8	*	*	*	*	*	*	0.2	0.6	1.6	4.2	13.5
1999.....	0.9	*	*	*	*	*	0.1	0.4	0.6	1.7	4.5	14.0
Black, both sexes												
2004.....	0.6	*	*	*	*	*	*	0.5	1.1	2.0	5.7	14.7
2003.....	0.6	*	*	*	*	*	*	0.6	1.0	1.9	6.8	14.9
2002.....	0.6	*	*	*	*	*	*	0.6	0.9	1.4	5.6	15.4
2001.....	0.6	*	*	*	*	*	*	0.6	0.9	2.5	5.8	12.3
2000.....	0.7	*	*	*	*	*	0.4	0.9	1.0	2.6	7.2	18.1
1999.....	0.7	*	*	*	*	*	*	0.6	1.3	3.4	6.0	19.8
Black, male												
2004.....	0.6	*	*	*	*	*	*	*	1.5	*	6.1	21.9
2003.....	0.6	*	*	*	*	*	*	*	*	*	10.2	*
2002.....	0.5	*	*	*	*	*	*	*	*	*	7.7	*
2001.....	0.6	*	*	*	*	*	*	*	*	3.8	6.5	*
2000.....	0.7	*	*	*	*	*	*	1.0	*	2.9	7.3	24.2
1999.....	0.7	*	*	*	*	*	*	*	*	4.1	7.4	35.2
Black, female												
2004.....	0.7	*	*	*	*	*	*	*	*	*	5.5	11.8
2003.....	0.7	*	*	*	*	*	*	*	*	*	4.9	13.8
2002.....	0.6	*	*	*	*	*	*	*	*	*	4.4	14.4
2001.....	0.6	*	*	*	*	*	*	*	*	*	5.5	11.8
2000.....	0.8	*	*	*	*	*	*	*	*	2.4	7.2	15.9
1999.....	0.7	*	*	*	*	*	*	*	*	2.9	5.2	14.0

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Glomerular diseases (N00-N07)												
All races 3/, both sexes												
2004.....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.6	1.6	3.3
2003.....	0.2	*	*	*	*	*	0.0	0.1	0.2	0.7	1.5	3.0
2002.....	0.2	*	*	*	*	0.1	0.1	0.1	0.2	0.6	1.7	3.3
2001.....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.7	1.5	3.3
2000.....	0.2	*	*	*	*	*	0.1	0.1	0.3	0.7	1.7	3.6
1999.....	0.3	*	*	*	*	*	0.1	0.1	0.3	0.7	2.0	3.7
All races 3/, male												
2004.....	0.2	*	*	*	*	*	*	0.1	0.3	0.7	1.9	4.2
2003.....	0.2	*	*	*	*	*	*	0.2	0.2	0.7	1.8	3.9
2002.....	0.2	*	*	*	*	*	*	0.1	0.3	0.7	1.9	4.3
2001.....	0.2	*	*	*	*	*	*	0.1	0.3	0.9	2.0	4.6
2000.....	0.3	*	*	*	*	*	*	0.2	0.3	0.8	2.0	5.6
1999.....	0.3	*	*	*	*	*	*	0.2	0.3	0.7	2.6	5.6
All races 3/, female												
2004.....	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.3	2.9
2003.....	0.2	*	*	*	*	*	*	*	0.2	0.6	1.4	2.6
2002.....	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.5	2.8
2001.....	0.2	*	*	*	*	*	*	*	*	0.6	1.1	2.8
2000.....	0.2	*	*	*	*	*	*	*	0.2	0.6	1.4	2.8
1999.....	0.3	*	*	*	*	*	*	0.1	0.3	0.7	1.5	2.9
White, both sexes												
2004.....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.6	1.5	3.2
2003.....	0.2	*	*	*	*	*	*	0.1	0.2	0.6	1.4	2.8
2002.....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.6	1.6	3.0
2001.....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.7	1.4	3.3
2000.....	0.2	*	*	*	*	*	*	0.1	0.2	0.6	1.6	3.3
1999.....	0.2	*	*	*	*	*	*	0.1	0.2	0.6	1.9	3.4
White, male												
2004.....	0.2	*	*	*	*	*	*	0.1	0.2	0.6	1.8	3.9
2003.....	0.2	*	*	*	*	*	*	0.1	0.2	0.6	1.5	3.7
2002.....	0.2	*	*	*	*	*	*	*	0.3	0.7	1.8	4.1
2001.....	0.2	*	*	*	*	*	*	*	0.2	0.8	2.0	4.8
2000.....	0.2	*	*	*	*	*	*	0.1	0.3	0.7	2.0	4.9
1999.....	0.3	*	*	*	*	*	*	0.1	0.3	0.6	2.5	5.1

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other and unspecified glomerular diseases (N02-N03,N05-N07)

All races 3/, both sexes

2004.....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.4	1.2	2.5
2003.....	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.2	2.5
2002.....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.4	1.3	2.4
2001.....	0.2	*	*	*	*	*	0.1	0.1	0.1	0.6	1.1	2.5
2000.....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.5	1.3	2.5
1999.....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.5	1.5	2.9

All races 3/, male

2004.....	0.2	*	*	*	*	*	*	*	0.3	0.5	1.4	3.5
2003.....	0.2	*	*	*	*	*	*	0.2	0.2	0.5	1.4	3.6
2002.....	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.6	3.3
2001.....	0.2	*	*	*	*	*	*	0.1	0.2	0.8	1.6	3.5
2000.....	0.2	*	*	*	*	*	*	0.1	0.3	0.6	1.5	3.8
1999.....	0.2	*	*	*	*	*	*	0.1	0.3	0.4	2.0	4.4

All races 3/, female

2004.....	0.2	*	*	*	*	*	*	*	0.2	0.4	1.0	2.1
2003.....	0.2	*	*	*	*	*	*	*	0.2	0.5	1.0	2.0
2002.....	0.2	*	*	*	*	*	*	0.1	0.2	0.3	1.1	1.9
2001.....	0.2	*	*	*	*	*	*	*	*	0.5	0.8	2.1
2000.....	0.2	*	*	*	*	*	*	*	0.2	0.5	1.1	1.9
1999.....	0.2	*	*	*	*	*	*	*	0.2	0.5	1.1	2.3

White, both sexes

2004.....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.4	1.1	2.4
2003.....	0.2	*	*	*	*	*	*	0.1	0.2	0.4	1.1	2.3
2002.....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.4	1.2	2.2
2001.....	0.2	*	*	*	*	*	*	0.1	0.1	0.6	1.1	2.5
2000.....	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.2	2.1
1999.....	0.2	*	*	*	*	*	*	0.1	0.2	0.4	1.4	2.7

White, male

2004.....	0.2	*	*	*	*	*	*	*	0.2	0.4	1.4	3.2
2003.....	0.2	*	*	*	*	*	*	0.1	*	0.5	1.2	3.3
2002.....	0.2	*	*	*	*	*	*	*	0.2	0.5	1.5	3.3
2001.....	0.2	*	*	*	*	*	*	*	*	0.7	1.6	3.7
2000.....	0.2	*	*	*	*	*	*	*	0.2	0.6	1.5	3.2
1999.....	0.2	*	*	*	*	*	*	*	0.2	0.4	1.9	4.2

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Renal tubulo-interstitial diseases (N10-N15)												
All races 3/, both sexes												
2004.....	0.4	*	*	*	*	*	0.1	0.2	0.3	0.9	3.2	9.3
2003.....	0.5	*	*	*	*	0.1	0.1	0.2	0.3	0.9	3.2	10.3
2002.....	0.4	*	*	*	*	*	0.1	0.1	0.3	1.0	3.2	10.6
2001.....	0.4	*	*	*	*	*	0.1	0.1	0.3	0.8	3.1	10.4
2000.....	0.4	*	*	*	*	*	0.1	0.2	0.3	0.9	3.2	12.0
1999.....	0.5	*	*	*	*	*	0.1	0.2	0.4	1.0	3.4	12.7
All races 3/, male												
2004.....	0.4	*	*	*	*	*	*	0.1	0.4	0.9	3.3	12.7
2003.....	0.4	*	*	*	*	*	*	0.2	0.3	0.7	3.4	13.8
2002.....	0.4	*	*	*	*	*	*	0.1	0.3	1.1	3.5	13.8
2001.....	0.3	*	*	*	*	*	*	*	0.2	0.9	3.3	12.5
2000.....	0.3	*	*	*	*	*	*	0.1	0.2	0.7	3.3	14.4
1999.....	0.4	*	*	*	*	*	*	0.2	0.4	0.9	4.0	17.0
All races 3/, female												
2004.....	0.5	*	*	*	*	*	0.1	0.2	0.3	0.8	3.1	7.8
2003.....	0.5	*	*	*	*	*	0.1	0.2	0.4	1.0	3.1	8.7
2002.....	0.5	*	*	*	*	*	*	0.2	0.3	0.9	3.0	9.2
2001.....	0.5	*	*	*	*	*	0.1	0.2	0.4	0.7	3.0	9.5
2000.....	0.5	*	*	*	*	*	*	0.2	0.4	1.0	3.0	11.1
1999.....	0.6	*	*	*	*	*	0.1	0.2	0.3	1.1	3.0	11.0
White, both sexes												
2004.....	0.5	*	*	*	*	*	0.1	0.1	0.3	0.9	3.2	9.3
2003.....	0.5	*	*	*	*	*	0.1	0.2	0.3	0.9	3.2	10.4
2002.....	0.5	*	*	*	*	*	0.1	0.1	0.3	1.0	3.2	10.9
2001.....	0.5	*	*	*	*	*	0.1	0.1	0.3	0.8	3.1	10.6
2000.....	0.5	*	*	*	*	*	*	0.1	0.3	0.8	3.0	12.0
1999.....	0.5	*	*	*	*	*	0.1	0.2	0.3	0.9	3.5	12.9
White, male												
2004.....	0.4	*	*	*	*	*	*	*	0.3	0.8	3.3	12.6
2003.....	0.4	*	*	*	*	*	*	0.2	0.2	0.8	3.3	14.3
2002.....	0.4	*	*	*	*	*	*	0.1	0.3	1.1	3.4	14.3
2001.....	0.4	*	*	*	*	*	*	*	*	0.8	3.2	12.8
2000.....	0.4	*	*	*	*	*	*	*	0.2	0.6	3.2	14.6
1999.....	0.4	*	*	*	*	*	*	0.1	0.3	0.7	4.1	17.0

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Renal failure (N17-N19)												
All races 3/, both sexes												
2004.....	14.2	4.0	*	*	0.2	0.6	1.7	4.8	13.4	37.9	106.6	282.9
2003.....	14.4	4.2	*	0.1	0.2	0.7	1.7	4.8	13.4	39.3	107.8	289.7
2002.....	13.9	4.1	*	*	0.2	0.6	1.6	4.6	12.7	38.6	107.2	284.8
2001.....	13.6	3.1	*	*	0.2	0.5	1.6	4.5	12.7	39.4	102.6	284.0
2000.....	13.0	4.0	*	*	0.2	0.6	1.6	4.2	12.5	37.2	98.9	273.7
1999.....	12.4	4.3	*	0.1	0.2	0.6	1.5	3.9	11.7	36.4	95.3	264.8
All races 3/, male												
2004.....	13.8	5.1	*	*	0.2	0.5	2.0	5.7	14.8	42.9	132.1	367.4
2003.....	14.1	4.8	*	*	0.2	0.7	2.0	5.7	15.2	44.5	134.8	383.1
2002.....	13.7	5.1	*	*	0.2	0.6	1.8	5.4	14.0	45.0	134.0	380.7
2001.....	13.2	3.5	*	*	0.2	0.5	1.8	5.3	13.8	45.2	128.0	388.7
2000.....	12.6	5.1	*	*	0.2	0.6	1.8	5.1	13.8	42.1	125.1	374.2
1999.....	12.1	4.5	*	*	0.1	0.5	1.7	4.5	13.1	41.5	122.3	364.6
All races 3/, female												
2004.....	14.6	3.0	*	*	0.2	0.6	1.5	3.9	12.0	33.7	89.5	244.9
2003.....	14.6	3.7	*	*	0.2	0.6	1.5	3.8	11.8	34.9	89.7	248.4
2002.....	14.2	3.0	*	*	0.2	0.7	1.4	3.8	11.6	33.2	89.5	243.2
2001.....	14.0	2.7	*	*	0.2	0.5	1.5	3.7	11.8	34.5	85.9	240.1
2000.....	13.3	2.8	*	*	0.2	0.5	1.3	3.4	11.2	33.3	81.8	232.8
1999.....	12.7	4.1	*	*	0.2	0.6	1.3	3.3	10.4	32.2	77.8	224.5
White, both sexes												
2004.....	13.9	2.9	*	*	0.1	0.4	1.2	3.3	10.0	32.3	99.8	274.7
2003.....	14.0	3.5	*	*	0.2	0.5	1.1	3.3	10.3	33.2	100.7	281.5
2002.....	13.6	3.4	*	*	0.1	0.4	1.1	3.1	9.6	32.7	100.5	275.3
2001.....	13.2	2.2	*	*	0.2	0.4	1.1	3.1	9.5	33.4	95.3	274.6
2000.....	12.6	2.7	*	*	0.2	0.4	1.1	2.9	9.4	31.2	92.2	265.0
1999.....	12.0	3.3	*	*	0.1	0.4	1.0	2.5	8.9	30.8	87.9	251.9
White, male												
2004.....	13.6	3.6	*	*	0.2	0.3	1.4	3.9	11.4	37.3	126.5	362.3
2003.....	13.8	4.1	*	*	0.2	0.5	1.3	3.9	11.7	38.3	128.3	379.3
2002.....	13.4	4.2	*	*	*	0.4	1.2	3.6	10.4	39.2	128.3	372.7
2001.....	13.0	2.6	*	*	0.2	0.3	1.3	3.7	10.7	39.5	121.5	383.9
2000.....	12.4	3.6	*	*	0.2	0.4	1.2	3.4	10.6	36.9	118.9	370.4
1999.....	11.8	3.9	*	*	*	0.4	1.1	3.0	10.3	36.2	114.4	354.9

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Renal failure (N17-N19)												
White, female												
2004.....	14.1	2.1	*	*	0.1	0.4	1.0	2.8	8.7	28.1	81.6	235.2
2003.....	14.2	2.8	*	*	*	0.5	0.9	2.7	9.0	28.8	82.1	238.2
2002.....	13.8	2.6	*	*	*	0.5	1.0	2.6	8.9	27.2	81.8	232.9
2001.....	13.4	1.8	*	*	0.2	0.4	0.9	2.5	8.4	28.3	78.0	228.8
2000.....	12.8	1.7	*	*	0.2	0.4	0.9	2.3	8.2	26.5	74.6	222.1
1999.....	12.2	2.8	*	*	0.1	0.4	0.8	2.2	7.6	26.2	70.6	210.6
Black, both sexes												
2004.....	20.0	9.8	*	*	0.4	1.7	5.3	16.3	44.0	94.2	200.3	426.4
2003.....	20.3	8.9	*	*	0.5	2.1	6.1	16.1	42.5	102.0	204.4	436.4
2002.....	19.5	7.7	*	*	0.5	1.9	5.3	15.7	40.6	99.6	200.2	434.2
2001.....	19.3	8.1	*	*	0.4	1.7	5.2	15.4	41.1	101.3	200.0	425.6
2000.....	18.5	11.1	*	*	0.4	1.7	5.1	15.2	40.1	98.6	192.4	400.4
1999.....	18.2	9.6	*	*	0.4	1.8	5.2	14.8	36.4	94.1	191.8	425.3
Black, male												
2004.....	18.9	11.9	*	*	*	1.9	6.0	20.6	49.0	106.8	230.0	506.7
2003.....	19.7	9.2	*	*	*	2.4	7.0	20.8	51.2	118.8	241.7	513.0
2002.....	18.8	9.6	*	*	*	1.7	5.9	19.6	48.8	114.2	231.5	555.3
2001.....	17.7	8.7	*	*	*	2.1	5.0	18.8	44.7	112.4	231.6	508.3
2000.....	17.3	13.4	*	*	*	1.8	6.3	19.5	44.2	103.3	229.2	480.9
1999.....	17.1	8.1	*	*	*	1.7	6.2	18.5	39.7	101.5	245.7	513.2
Black, female												
2004.....	21.0	7.7	*	*	*	1.6	4.7	12.6	39.9	85.3	183.5	394.6
2003.....	20.9	8.7	*	*	*	1.8	5.3	12.1	35.5	90.0	183.2	406.7
2002.....	20.2	*	*	*	*	2.0	4.7	12.3	34.1	89.2	182.6	387.7
2001.....	20.7	7.6	*	*	*	1.3	5.4	12.6	38.2	93.5	182.3	394.9
2000.....	19.6	8.6	*	*	*	1.6	4.0	11.5	36.7	95.3	171.7	370.4
1999.....	19.1	11.1	*	*	*	1.9	4.3	11.7	33.8	88.9	161.6	392.5

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Urinary tract infection, site not specified (N39.0)

All races 3/, both sexes

2004.....	4.7	*	*	*	*	0.1	0.2	0.6	1.7	7.7	35.6	143.5
2003.....	4.8	*	*	*	*	0.1	0.2	0.5	1.9	7.9	35.5	147.7
2002.....	4.8	*	*	*	*	0.1	0.2	0.5	1.9	7.7	35.6	152.2
2001.....	4.9	*	*	*	*	0.1	0.2	0.5	1.8	8.2	37.0	160.4
2000.....	4.9	*	*	*	*	0.1	0.1	0.5	1.7	7.3	38.4	163.4
1999.....	5.0	*	*	*	*	0.1	0.2	0.5	1.8	8.0	38.1	172.0

All races 3/, male

2004.....	3.1	*	*	*	*	*	0.2	0.6	1.6	6.8	32.1	117.7
2003.....	3.0	*	*	*	*	*	0.1	0.4	1.7	6.9	33.1	114.6
2002.....	3.0	*	*	*	*	*	0.2	0.4	1.6	6.7	31.0	123.5
2001.....	3.0	*	*	*	*	*	0.1	0.4	1.6	7.4	32.6	125.8
2000.....	3.1	*	*	*	*	*	0.1	0.5	1.5	6.7	34.3	139.8
1999.....	3.1	*	*	*	*	*	0.1	0.4	1.8	7.5	34.7	137.5

All races 3/, female

2004.....	6.4	*	*	*	*	0.1	0.2	0.7	1.9	8.4	38.0	155.1
2003.....	6.5	*	*	*	*	*	0.2	0.7	2.1	8.7	37.0	162.3
2002.....	6.5	*	*	*	*	*	0.2	0.6	2.1	8.5	38.7	164.7
2001.....	6.7	*	*	*	*	*	0.2	0.6	2.0	8.9	39.8	174.9
2000.....	6.6	*	*	*	*	*	0.1	0.6	1.9	7.8	41.1	173.0
1999.....	6.9	*	*	*	*	*	0.2	0.6	1.9	8.4	40.3	185.9

White, both sexes

2004.....	5.1	*	*	*	*	0.1	0.2	0.6	1.5	7.2	34.7	143.2
2003.....	5.1	*	*	*	*	0.1	0.1	0.5	1.7	7.3	34.5	147.2
2002.....	5.1	*	*	*	*	*	0.1	0.5	1.7	7.2	34.6	150.8
2001.....	5.2	*	*	*	*	0.1	0.2	0.4	1.6	7.6	35.6	158.2
2000.....	5.2	*	*	*	*	0.1	0.1	0.5	1.6	6.7	37.1	161.9
1999.....	5.3	*	*	*	*	0.1	0.1	0.4	1.6	7.2	37.0	169.8

White, male

2004.....	3.2	*	*	*	*	*	0.2	0.5	1.4	6.2	31.0	115.9
2003.....	3.2	*	*	*	*	*	0.1	0.4	1.5	6.3	31.9	116.5
2002.....	3.1	*	*	*	*	*	0.1	0.4	1.5	6.1	29.7	121.1
2001.....	3.1	*	*	*	*	*	0.2	0.3	1.4	6.7	31.6	122.1
2000.....	3.2	*	*	*	*	*	*	0.4	1.3	6.0	32.7	137.8
1999.....	3.2	*	*	*	*	*	0.1	0.4	1.6	6.4	32.9	132.1

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Urinary tract infection, site not specified (N39.0)

White, female

2004.....	6.9	*	*	*	*	*	0.2	0.6	1.7	8.1	37.2	155.5
2003.....	7.0	*	*	*	*	*	0.1	0.6	1.9	8.2	36.3	160.8
2002.....	7.1	*	*	*	*	*	0.1	0.5	1.9	8.1	37.9	163.8
2001.....	7.2	*	*	*	*	*	0.2	0.5	1.9	8.4	38.3	173.3
2000.....	7.1	*	*	*	*	*	0.1	0.5	1.8	7.3	40.0	171.7
1999.....	7.4	*	*	*	*	*	0.2	0.5	1.7	8.0	39.7	185.0

Black, both sexes

2004.....	4.1	*	*	*	*	*	0.4	1.4	3.9	13.0	51.9	172.9
2003.....	4.2	*	*	*	*	*	*	1.2	4.3	14.5	53.6	178.5
2002.....	4.1	*	*	*	*	*	0.3	1.0	3.7	13.3	53.8	191.5
2001.....	4.4	*	*	*	*	*	*	1.2	3.4	14.4	58.1	208.5
2000.....	4.3	*	*	*	*	*	*	1.1	3.7	13.7	59.8	196.8
1999.....	4.5	*	*	*	*	*	0.4	1.1	3.7	16.3	56.6	213.0

Black, male

2004.....	3.0	*	*	*	*	*	*	1.2	4.1	13.4	52.9	162.9
2003.....	2.8	*	*	*	*	*	*	*	3.9	15.1	58.1	125.1
2002.....	2.9	*	*	*	*	*	*	*	3.5	13.4	54.8	188.7
2001.....	3.0	*	*	*	*	*	*	1.0	3.7	14.8	52.7	204.2
2000.....	3.1	*	*	*	*	*	*	1.3	4.2	14.3	61.8	175.3
1999.....	3.5	*	*	*	*	*	*	*	4.0	20.3	61.8	215.6

Black, female

2004.....	5.1	*	*	*	*	*	*	1.5	3.7	12.7	51.3	176.8
2003.....	5.4	*	*	*	*	*	*	1.5	4.6	14.1	51.0	199.3
2002.....	5.3	*	*	*	*	*	*	1.2	3.9	13.2	53.3	192.6
2001.....	5.6	*	*	*	*	*	*	1.3	3.1	14.1	61.1	210.1
2000.....	5.3	*	*	*	*	*	*	*	3.3	13.3	58.6	204.8
1999.....	5.4	*	*	*	*	*	*	1.4	3.5	13.5	53.7	212.0

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other diseases of urinary system (N28-N36,N39.1-N39.9)

All races 3/, both sexes

2004.....	0.4	*	*	*	*	*	0.1	0.2	0.5	1.1	2.9	7.8
2003.....	0.4	*	*	*	*	*	0.1	0.2	0.5	0.9	2.6	7.1
2002.....	0.4	*	*	*	*	0.1	0.1	0.2	0.4	1.0	2.6	7.5
2001.....	0.4	*	*	*	*	*	0.1	0.2	0.5	1.0	2.7	7.1
2000.....	0.4	*	*	*	*	*	0.1	0.2	0.5	1.0	2.8	7.5
1999.....	0.4	*	*	*	*	*	0.1	0.2	0.5	1.1	2.6	7.1

All races 3/, male

2004.....	0.4	*	*	*	*	*	0.1	0.3	0.6	1.3	3.8	9.2
2003.....	0.4	*	*	*	*	*	*	0.2	0.6	1.1	3.2	9.8
2002.....	0.4	*	*	*	*	*	0.1	0.3	0.5	1.2	3.2	9.0
2001.....	0.4	*	*	*	*	*	0.1	0.2	0.5	1.2	3.2	8.9
2000.....	0.4	*	*	*	*	*	0.1	0.2	0.5	1.3	3.5	10.4
1999.....	0.4	*	*	*	*	*	0.1	0.2	0.5	1.5	3.2	9.9

All races 3/, female

2004.....	0.4	*	*	*	*	*	*	0.2	0.3	0.9	2.2	7.1
2003.....	0.4	*	*	*	*	*	0.1	0.2	0.4	0.7	2.1	6.0
2002.....	0.4	*	*	*	*	*	*	0.1	0.3	0.9	2.2	6.9
2001.....	0.4	*	*	*	*	*	*	0.2	0.4	0.9	2.3	6.3
2000.....	0.4	*	*	*	*	*	0.1	0.2	0.4	0.8	2.2	6.3
1999.....	0.4	*	*	*	*	*	*	0.1	0.4	0.7	2.2	6.0

White, both sexes

2004.....	0.4	*	*	*	*	*	0.1	0.2	0.4	1.0	2.8	7.8
2003.....	0.4	*	*	*	*	*	0.1	0.2	0.4	0.9	2.6	7.2
2002.....	0.4	*	*	*	*	*	0.1	0.2	0.3	1.0	2.6	7.5
2001.....	0.4	*	*	*	*	*	0.1	0.2	0.4	0.9	2.6	7.2
2000.....	0.4	*	*	*	*	*	0.1	0.2	0.4	0.9	2.7	7.3
1999.....	0.4	*	*	*	*	*	0.1	0.1	0.4	0.9	2.4	7.1

White, male

2004.....	0.4	*	*	*	*	*	*	0.2	0.6	1.2	3.8	9.3
2003.....	0.4	*	*	*	*	*	*	0.1	0.5	1.1	3.1	9.7
2002.....	0.4	*	*	*	*	*	*	0.2	0.4	1.2	3.1	8.8
2001.....	0.4	*	*	*	*	*	0.1	0.2	0.4	1.1	3.2	9.1
2000.....	0.4	*	*	*	*	*	0.1	0.2	0.4	1.1	3.5	10.2
1999.....	0.4	*	*	*	*	*	*	0.2	0.5	1.3	3.0	10.1

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Diseases of male genital organs (N40-N50)

All races 3/, both sexes

2004.....	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.9	6.3
2003.....	0.2	*	*	*	*	*	*	*	0.1	0.4	1.6	6.0
2002.....	0.2	*	*	*	*	*	*	*	0.1	0.4	1.6	5.8
2001.....	0.2	*	*	*	*	*	*	*	0.1	0.4	1.5	6.3
2000.....	0.2	*	*	*	*	*	*	*	0.1	0.4	1.6	5.9
1999.....	0.2	*	*	*	*	*	*	*	0.1	0.4	1.7	6.1

All races 3/, male

2004.....	0.5	*	*	*	*	*	*	0.1	0.4	1.0	4.7	20.3
2003.....	0.4	*	*	*	*	*	*	*	0.3	0.9	3.9	19.6
2002.....	0.4	*	*	*	*	*	*	*	0.2	0.9	4.0	19.2
2001.....	0.4	*	*	*	*	*	*	*	0.2	0.9	3.7	21.2
2000.....	0.4	*	*	*	*	*	*	*	0.3	0.9	3.9	20.5
1999.....	0.4	*	*	*	*	*	*	*	0.3	1.0	4.3	21.2

All races 3/, female

2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

White, both sexes

2004.....	0.3	*	*	*	*	*	*	*	0.2	0.4	1.8	6.5
2003.....	0.2	*	*	*	*	*	*	*	0.1	0.4	1.6	6.2
2002.....	0.2	*	*	*	*	*	*	*	*	0.4	1.5	5.9
2001.....	0.2	*	*	*	*	*	*	*	*	0.4	1.5	6.4
2000.....	0.2	*	*	*	*	*	*	*	0.1	0.4	1.5	5.9
1999.....	0.2	*	*	*	*	*	*	*	0.1	0.4	1.7	6.1

White, male

2004.....	0.5	*	*	*	*	*	*	*	0.3	1.0	4.5	20.8
2003.....	0.5	*	*	*	*	*	*	*	0.2	0.9	3.9	20.4
2002.....	0.4	*	*	*	*	*	*	*	*	0.8	3.8	19.5
2001.....	0.5	*	*	*	*	*	*	*	*	0.9	3.7	21.8
2000.....	0.4	*	*	*	*	*	*	*	0.2	0.8	3.8	20.5
1999.....	0.5	*	*	*	*	*	*	*	0.2	0.9	4.4	21.4

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Hyperplasia of prostate (N40)

All races 3/, both sexes

2004.....	0.2	*	*	*	*	*	*	*	*	0.2	1.4	5.5
2003.....	0.2	*	*	*	*	*	*	*	*	0.3	1.3	5.3
2002.....	0.2	*	*	*	*	*	*	*	*	0.2	1.3	4.8
2001.....	0.2	*	*	*	*	*	*	*	*	0.2	1.1	5.5
2000.....	0.2	*	*	*	*	*	*	*	*	0.2	1.3	5.2
1999.....	0.2	*	*	*	*	*	*	*	*	0.3	1.3	5.0

All races 3/, male

2004.....	0.4	*	*	*	*	*	*	*	*	0.5	3.4	17.8
2003.....	0.3	*	*	*	*	*	*	*	*	0.6	3.2	17.2
2002.....	0.3	*	*	*	*	*	*	*	*	0.5	3.2	15.8
2001.....	0.3	*	*	*	*	*	*	*	*	0.5	2.8	18.6
2000.....	0.3	*	*	*	*	*	*	*	*	0.5	3.3	18.1
1999.....	0.3	*	*	*	*	*	*	*	*	0.6	3.3	17.5

All races 3/, female

2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

White, both sexes

2004.....	0.2	*	*	*	*	*	*	*	*	0.2	1.3	5.7
2003.....	0.2	*	*	*	*	*	*	*	*	0.3	1.3	5.5
2002.....	0.2	*	*	*	*	*	*	*	*	0.2	1.3	4.9
2001.....	0.2	*	*	*	*	*	*	*	*	0.3	1.1	5.7
2000.....	0.2	*	*	*	*	*	*	*	*	0.2	1.3	5.2
1999.....	0.2	*	*	*	*	*	*	*	*	0.2	1.3	5.1

White, male

2004.....	0.4	*	*	*	*	*	*	*	*	0.5	3.3	18.3
2003.....	0.4	*	*	*	*	*	*	*	*	0.6	3.2	18.0
2002.....	0.3	*	*	*	*	*	*	*	*	0.5	3.2	16.1
2001.....	0.4	*	*	*	*	*	*	*	*	0.5	2.9	19.4
2000.....	0.3	*	*	*	*	*	*	*	*	0.5	3.3	18.1
1999.....	0.3	*	*	*	*	*	*	*	*	0.5	3.4	17.9

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other diseases of male genital organs (N41-N50)

All races 3/, both sexes

2004.....	0.1	*	*	*	*	*	*	0.1	0.1	0.2	0.5	0.8
2003.....	0.0	*	*	*	*	*	*	*	0.1	0.1	0.3	0.7
2002.....	0.0	*	*	*	*	*	*	*	*	0.2	0.3	1.0
2001.....	0.1	*	*	*	*	*	*	*	*	0.2	0.4	0.8
2000.....	0.0	*	*	*	*	*	*	*	0.1	0.2	0.3	0.7
1999.....	0.1	*	*	*	*	*	*	*	*	0.2	0.4	1.1

All races 3/, male

2004.....	0.2	*	*	*	*	*	*	0.1	0.3	0.5	1.3	2.5
2003.....	0.1	*	*	*	*	*	*	*	0.2	0.3	0.7	2.4
2002.....	0.1	*	*	*	*	*	*	*	*	0.4	0.7	3.4
2001.....	0.1	*	*	*	*	*	*	*	*	0.4	0.9	2.6
2000.....	0.1	*	*	*	*	*	*	*	0.2	0.4	0.7	2.4
1999.....	0.1	*	*	*	*	*	*	*	*	0.4	1.0	3.7

All races 3/, female

2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

White, both sexes

2004.....	0.1	*	*	*	*	*	*	*	0.1	0.2	0.5	0.8
2003.....	0.0	*	*	*	*	*	*	*	*	0.1	0.3	0.7
2002.....	0.0	*	*	*	*	*	*	*	*	0.1	0.2	1.0
2001.....	0.1	*	*	*	*	*	*	*	*	0.2	0.3	0.7
2000.....	0.0	*	*	*	*	*	*	*	*	0.2	0.2	0.7
1999.....	0.1	*	*	*	*	*	*	*	*	0.2	0.4	1.0

White, male

2004.....	0.1	*	*	*	*	*	*	*	0.2	0.4	1.2	2.5
2003.....	0.1	*	*	*	*	*	*	*	*	0.3	0.6	2.4
2002.....	0.1	*	*	*	*	*	*	*	*	0.3	0.6	3.3
2001.....	0.1	*	*	*	*	*	*	*	*	0.4	0.9	2.4
2000.....	0.1	*	*	*	*	*	*	*	*	0.3	0.6	2.3
1999.....	0.1	*	*	*	*	*	*	*	*	0.3	1.0	3.5

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

XV. Pregnancy, childbirth and the puerperium (000-099)

White, female

2004.....	0.3	*	0.6	1.1	0.6	0.3	*	*	*	*
2003.....	0.3	*	0.4	0.9	0.5	0.1	*	*	*	*
2002.....	0.2	*	0.3	0.6	0.3	*	*	*	*	*
2001.....	0.2	*	0.4	0.7	0.3	*	*	*	*	*
2000.....	0.2	*	0.4	0.7	0.4	*	*	*	*	*
1999.....	0.2	*	0.3	0.6	0.4	*	*	*	*	*

Black, both sexes

2004.....	0.7	*	1.1	2.0	0.8	0.5	*	*	*	*
2003.....	0.5	*	0.7	1.5	1.0	*	*	*	*	*
2002.....	0.4	*	0.7	1.2	0.7	*	*	*	*	*
2001.....	0.4	*	0.7	1.2	0.7	*	*	*	*	*
2000.....	0.4	*	0.6	1.2	0.7	*	*	*	*	*
1999.....	0.4	*	0.8	1.3	0.7	*	*	*	*	*

Black, male

2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

Black, female

2004.....	1.2	*	2.3	3.8	1.5	0.9	*	*	*	*
2003.....	1.0	*	1.4	2.9	1.9	*	*	*	*	*
2002.....	0.8	*	1.5	2.3	1.4	*	*	*	*	*
2001.....	0.8	*	1.5	2.3	1.4	*	*	*	*	*
2000.....	0.7	*	1.2	2.3	1.2	*	*	*	*	*
1999.....	0.9	*	1.6	2.5	1.4	*	*	*	*	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other direct obstetric deaths (O10-O92)

White, female

2004.....	0.2	*	0.2	0.6	0.3	*	*	*	*	*
2003.....	0.2	*	0.3	0.5	0.3	*	*	*	*	*
2002.....	0.1	*	0.2	0.5	0.3	*	*	*	*	*
2001.....	0.2	*	0.3	0.6	0.2	*	*	*	*	*
2000.....	0.2	*	0.3	0.5	0.3	*	*	*	*	*
1999.....	0.1	*	0.2	0.5	0.3	*	*	*	*	*

Black, both sexes

2004.....	0.4	*	0.6	1.3	0.3	*	*	*	*	*
2003.....	0.4	*	0.4	1.1	0.8	*	*	*	*	*
2002.....	0.3	*	0.6	0.9	0.5	*	*	*	*	*
2001.....	0.3	*	0.5	0.9	0.5	*	*	*	*	*
2000.....	0.3	*	0.5	0.8	0.4	*	*	*	*	*
1999.....	0.3	*	0.6	1.0	0.6	*	*	*	*	*

Black, male

2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

Black, female

2004.....	0.7	*	1.3	2.4	0.7	*	*	*	*	*
2003.....	0.7	*	0.8	2.1	1.5	*	*	*	*	*
2002.....	0.6	*	1.2	1.8	0.9	*	*	*	*	*
2001.....	0.6	*	1.0	1.8	1.0	*	*	*	*	*
2000.....	0.5	*	0.9	1.6	0.8	*	*	*	*	*
1999.....	0.6	*	1.1	1.9	1.1	*	*	*	*	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

XVII. Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)

All races 3/, both sexes

2004.....	3.6	137.9	3.6	1.0	1.2	1.0	1.1	1.7	2.2	2.3	3.6	6.1
2003.....	3.6	140.4	3.4	0.9	1.1	1.1	1.3	1.6	2.4	2.3	3.6	6.8
2002.....	3.7	139.4	3.4	1.0	1.2	1.2	1.3	1.8	2.4	2.4	3.8	6.6
2001.....	3.7	136.7	3.6	0.9	1.3	1.2	1.3	1.6	2.3	2.4	3.7	7.2
2000.....	3.8	150.9	3.2	1.0	1.1	1.2	1.3	1.6	2.3	2.5	3.9	7.5
1999.....	3.7	144.2	3.6	1.0	1.1	1.2	1.2	1.8	2.3	2.5	4.0	7.8

All races 3/, male

2004.....	3.8	139.0	3.9	1.0	1.3	1.2	1.3	1.8	2.3	2.5	3.7	6.8
2003.....	3.8	147.6	3.4	0.9	1.3	1.2	1.4	1.7	2.5	2.4	3.5	6.7
2002.....	3.9	142.6	3.5	1.0	1.4	1.4	1.3	1.9	2.5	2.6	4.0	7.0
2001.....	3.9	143.1	3.6	1.0	1.5	1.2	1.4	1.7	2.3	2.7	4.1	7.7
2000.....	4.0	155.4	3.3	1.0	1.3	1.4	1.4	1.9	2.3	2.7	4.0	10.6
1999.....	3.9	147.9	4.0	1.0	1.3	1.2	1.3	1.9	2.5	2.5	4.2	8.5

All races 3/, female

2004.....	3.4	136.8	3.2	0.9	1.0	0.9	1.0	1.5	2.1	2.2	3.6	5.8
2003.....	3.4	132.8	3.5	0.9	0.9	1.0	1.2	1.5	2.3	2.2	3.7	6.8
2002.....	3.5	136.1	3.3	1.0	1.1	1.0	1.2	1.6	2.3	2.3	3.8	6.5
2001.....	3.4	129.9	3.7	0.8	1.0	1.1	1.2	1.6	2.3	2.2	3.5	7.0
2000.....	3.5	146.2	3.1	1.0	0.9	1.0	1.1	1.4	2.4	2.4	3.9	6.3
1999.....	3.5	140.3	3.2	1.1	0.9	1.1	1.1	1.6	2.2	2.6	3.9	7.6

White, both sexes

2004.....	3.5	134.5	3.2	0.9	1.1	1.1	1.1	1.7	2.3	2.5	3.7	6.3
2003.....	3.5	137.6	3.2	0.9	1.1	1.1	1.3	1.7	2.5	2.3	3.7	6.9
2002.....	3.6	138.3	3.3	1.0	1.2	1.2	1.3	1.8	2.4	2.5	3.9	6.9
2001.....	3.6	135.0	3.2	0.9	1.2	1.2	1.3	1.7	2.4	2.5	3.8	7.4
2000.....	3.6	147.1	2.9	0.9	1.1	1.2	1.2	1.7	2.4	2.5	4.1	7.8
1999.....	3.6	141.3	3.2	0.9	1.1	1.2	1.2	1.8	2.4	2.6	4.0	8.0

White, male

2004.....	3.6	134.3	3.5	1.0	1.2	1.2	1.3	1.9	2.4	2.6	3.7	7.2
2003.....	3.7	143.2	3.1	0.9	1.2	1.2	1.4	1.7	2.6	2.3	3.4	6.9
2002.....	3.8	143.3	3.4	1.0	1.3	1.4	1.3	1.9	2.5	2.6	3.9	7.3
2001.....	3.8	143.0	3.2	1.0	1.4	1.3	1.4	1.7	2.4	2.9	4.2	7.5
2000.....	3.9	151.9	2.9	0.9	1.3	1.4	1.4	1.9	2.4	2.6	4.3	11.1
1999.....	3.8	146.5	3.5	0.9	1.3	1.2	1.3	1.9	2.6	2.5	4.1	8.6

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

XVII. Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)

White, female

2004.....	3.4	134.7	2.9	0.8	1.0	1.0	1.0	1.6	2.2	2.3	3.6	6.0
2003.....	3.4	131.8	3.2	0.9	0.9	1.0	1.2	1.6	2.4	2.2	3.9	6.9
2002.....	3.5	133.1	3.1	1.0	1.1	1.0	1.2	1.7	2.3	2.4	3.9	6.7
2001.....	3.4	126.7	3.3	0.7	1.0	1.1	1.2	1.6	2.5	2.2	3.6	7.3
2000.....	3.5	142.0	2.9	0.9	0.9	1.0	1.1	1.5	2.4	2.5	4.0	6.5
1999.....	3.5	135.7	2.9	1.0	0.9	1.2	1.2	1.7	2.3	2.6	4.0	7.7

Black, both sexes

2004.....	4.5	162.9	5.1	1.3	1.7	1.1	1.2	1.7	1.6	1.6	3.9	*
2003.....	4.5	157.5	4.7	1.1	1.5	1.4	1.2	1.6	2.2	2.3	3.1	5.8
2002.....	4.7	155.2	4.5	1.2	1.6	1.4	1.6	1.9	2.6	2.4	3.3	*
2001.....	4.6	158.4	5.3	1.1	1.9	1.3	1.6	1.7	1.9	2.2	3.0	6.1
2000.....	4.9	179.2	5.0	1.3	1.5	1.4	1.6	1.6	2.4	2.6	2.7	*
1999.....	4.8	169.3	5.5	1.6	1.4	1.1	1.3	1.9	1.9	2.8	4.2	7.7

Black, male

2004.....	5.1	169.9	5.8	1.1	2.2	1.4	1.6	1.8	1.9	*	*	*
2003.....	5.1	170.7	4.8	1.1	1.7	1.5	1.2	1.9	2.7	3.5	*	*
2002.....	5.1	151.7	4.6	1.4	1.9	1.7	1.7	2.3	3.4	3.0	*	*
2001.....	5.0	160.8	5.5	1.0	2.4	1.2	1.7	1.7	2.0	*	*	*
2000.....	5.4	181.4	5.5	1.5	2.0	1.8	1.9	2.1	*	3.8	*	*
1999.....	5.2	172.7	6.2	1.6	1.8	1.2	1.6	2.3	2.1	*	*	*

Black, female

2004.....	4.0	155.6	4.3	1.4	1.2	0.8	0.9	1.6	1.4	*	3.3	*
2003.....	3.9	143.8	4.6	1.0	1.2	1.3	1.2	1.4	1.8	*	*	*
2002.....	4.3	158.9	4.4	1.0	1.3	1.1	1.5	1.5	2.0	*	*	*
2001.....	4.3	155.9	5.1	1.2	1.4	1.3	1.4	1.7	1.8	2.5	*	*
2000.....	4.4	176.9	4.6	1.2	1.1	1.1	1.4	1.1	3.0	*	*	*
1999.....	4.3	165.8	4.8	1.6	1.1	1.0	1.2	1.5	1.8	3.1	*	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Congenital malformations of heart (Q20-Q24)												
All races 3/, both sexes												
2004.....	1.0	34.1	1.2	0.3	0.5	0.5	0.4	0.5	0.5	0.5	1.0	1.5
2003.....	1.1	36.1	1.4	0.4	0.4	0.5	0.6	0.5	0.5	0.7	1.1	2.1
2002.....	1.1	37.4	1.2	0.4	0.5	0.6	0.5	0.5	0.5	0.7	1.2	2.3
2001.....	1.2	38.6	1.3	0.3	0.6	0.5	0.5	0.6	0.5	0.7	1.2	2.6
2000.....	1.2	42.9	1.2	0.3	0.6	0.6	0.5	0.5	0.6	0.7	1.4	2.5
1999.....	1.3	42.1	1.3	0.4	0.6	0.6	0.6	0.6	0.7	0.9	1.4	3.3
All races 3/, male												
2004.....	1.1	36.7	1.4	0.4	0.6	0.6	0.5	0.5	0.5	0.5	0.9	1.5
2003.....	1.2	40.1	1.5	0.4	0.6	0.6	0.6	0.5	0.5	0.6	0.9	*
2002.....	1.2	40.4	1.5	0.4	0.7	0.7	0.6	0.5	0.5	0.6	1.1	1.4
2001.....	1.3	41.8	1.4	0.3	0.8	0.6	0.6	0.6	0.5	0.8	1.2	1.7
2000.....	1.3	44.5	1.3	0.4	0.7	0.7	0.6	0.6	0.6	0.8	1.5	2.7
1999.....	1.4	45.7	1.5	0.4	0.7	0.6	0.6	0.7	0.7	0.8	1.3	3.1
All races 3/, female												
2004.....	0.9	31.4	1.0	0.3	0.4	0.4	0.4	0.4	0.4	0.5	1.1	1.6
2003.....	1.0	31.9	1.4	0.3	0.3	0.4	0.5	0.5	0.5	0.7	1.3	2.4
2002.....	1.0	34.4	1.0	0.3	0.4	0.5	0.5	0.6	0.5	0.7	1.3	2.6
2001.....	1.1	35.2	1.2	0.3	0.4	0.4	0.5	0.6	0.5	0.5	1.2	3.0
2000.....	1.1	41.3	1.2	0.3	0.4	0.5	0.5	0.5	0.6	0.7	1.4	2.4
1999.....	1.2	38.3	1.1	0.3	0.4	0.6	0.6	0.5	0.6	0.9	1.5	3.4
White, both sexes												
2004.....	1.0	33.2	1.1	0.3	0.5	0.5	0.4	0.5	0.5	0.5	1.1	1.6
2003.....	1.0	35.0	1.3	0.3	0.3	0.5	0.6	0.5	0.5	0.7	1.2	2.2
2002.....	1.1	36.4	1.3	0.3	0.5	0.6	0.5	0.6	0.5	0.7	1.3	2.4
2001.....	1.1	37.4	1.2	0.3	0.5	0.5	0.6	0.6	0.5	0.7	1.3	2.6
2000.....	1.2	41.5	1.1	0.3	0.5	0.6	0.5	0.6	0.6	0.7	1.5	2.7
1999.....	1.3	42.0	1.1	0.3	0.5	0.6	0.6	0.6	0.7	0.9	1.5	3.3
White, male												
2004.....	1.1	35.9	1.3	0.4	0.5	0.6	0.5	0.5	0.6	0.5	1.0	1.6
2003.....	1.1	39.3	1.3	0.4	0.5	0.6	0.7	0.5	0.5	0.7	0.9	*
2002.....	1.2	40.0	1.5	0.4	0.7	0.7	0.6	0.5	0.5	0.6	1.1	1.6
2001.....	1.2	41.3	1.3	0.3	0.7	0.7	0.6	0.6	0.5	0.8	1.3	*
2000.....	1.3	43.6	1.2	0.3	0.6	0.8	0.6	0.6	0.6	0.7	1.6	2.9
1999.....	1.3	46.2	1.3	0.4	0.7	0.7	0.6	0.7	0.7	0.8	1.3	3.0

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Congenital malformations of heart (Q20-Q24)												
White, female												
2004.....	0.9	30.5	0.9	0.3	0.4	0.4	0.4	0.4	0.5	0.5	1.2	1.6
2003.....	1.0	30.5	1.3	0.3	0.2	0.4	0.5	0.5	0.5	0.7	1.4	2.6
2002.....	1.0	32.7	1.0	0.3	0.3	0.5	0.5	0.6	0.5	0.7	1.3	2.7
2001.....	1.0	33.2	1.0	0.2	0.4	0.4	0.5	0.6	0.4	0.6	1.3	3.1
2000.....	1.1	39.3	1.1	0.3	0.4	0.5	0.5	0.5	0.6	0.7	1.4	2.6
1999.....	1.2	37.6	1.0	0.3	0.4	0.6	0.6	0.6	0.6	0.9	1.6	3.5
Black, both sexes												
2004.....	1.3	42.2	1.7	0.5	0.8	0.6	0.5	*	*	*	*	*
2003.....	1.4	41.7	2.1	0.4	0.8	0.7	0.5	0.6	*	*	*	*
2002.....	1.4	44.3	1.3	0.4	0.7	0.7	0.6	*	*	*	*	*
2001.....	1.6	47.7	1.9	0.4	1.1	0.6	0.7	0.6	*	*	*	*
2000.....	1.6	51.5	1.8	0.5	0.9	0.6	0.7	0.5	*	*	*	*
1999.....	1.6	47.9	2.2	0.6	0.8	0.5	0.7	0.5	*	*	*	*
Black, male												
2004.....	1.5	44.8	2.0	*	1.1	0.8	*	*	*	*	*	*
2003.....	1.6	44.9	2.2	*	1.0	0.8	*	*	*	*	*	*
2002.....	1.6	43.0	*	0.6	1.0	0.9	*	*	*	*	*	*
2001.....	1.7	48.8	1.7	*	1.6	*	*	*	*	*	*	*
2000.....	1.8	49.9	2.1	*	1.2	0.9	0.9	*	*	*	*	*
1999.....	1.8	50.1	2.5	0.6	1.1	*	0.8	*	*	*	*	*
Black, female												
2004.....	1.1	39.5	*	*	*	*	*	*	*	*	*	*
2003.....	1.2	38.3	2.1	*	*	*	0.7	*	*	*	*	*
2002.....	1.3	45.7	*	*	*	*	*	*	*	*	*	*
2001.....	1.4	46.6	2.0	*	0.7	*	*	*	*	*	*	*
2000.....	1.4	53.2	*	*	*	*	*	*	*	*	*	*
1999.....	1.4	45.6	2.0	0.6	*	*	0.7	*	*	*	*	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other congenital malformations of circulatory system (Q25-Q28)

All races 3/, both sexes

2004.....	0.3	6.3	0.2	0.1	0.1	0.1	0.1	0.2	0.3	0.4	0.9	1.6
2003.....	0.3	5.9	0.2	0.1	0.1	0.1	0.1	0.2	0.3	0.4	0.8	1.5
2002.....	0.3	5.5	0.2	0.1	0.1	0.1	0.1	0.2	0.3	0.5	1.0	1.5
2001.....	0.3	4.8	*	0.1	0.1	0.1	0.1	0.2	0.3	0.6	0.7	1.7
2000.....	0.3	5.5	0.2	0.1	0.1	0.1	0.2	0.2	0.3	0.4	0.9	2.1
1999.....	0.3	5.8	0.2	0.1	0.1	0.1	0.2	0.3	0.3	0.5	0.9	1.7

All races 3/, male

2004.....	0.3	6.7	*	0.1	0.1	*	0.2	0.3	0.3	0.4	1.1	1.9
2003.....	0.3	6.5	*	*	*	0.1	0.2	0.2	0.3	0.5	1.0	1.7
2002.....	0.3	5.8	*	*	*	0.2	0.1	0.3	0.4	0.6	1.0	1.8
2001.....	0.3	4.9	*	0.1	0.1	*	0.1	0.2	0.3	0.6	1.0	2.1
2000.....	0.3	5.5	*	*	0.1	0.1	0.2	0.3	0.3	0.5	1.0	3.0
1999.....	0.3	5.9	*	*	0.1	0.1	0.2	0.3	0.4	0.5	1.1	2.2

All races 3/, female

2004.....	0.3	5.8	*	0.1	0.1	*	0.1	0.2	0.2	0.3	0.8	1.5
2003.....	0.3	5.4	*	0.1	*	*	0.1	0.2	0.3	0.4	0.6	1.4
2002.....	0.3	5.2	*	*	*	*	0.1	0.2	0.2	0.4	1.1	1.4
2001.....	0.3	4.6	*	*	*	*	0.1	0.2	0.2	0.6	0.6	1.6
2000.....	0.3	5.4	*	0.1	*	*	0.2	0.2	0.3	0.4	0.9	1.7
1999.....	0.3	5.7	*	*	*	*	0.1	0.2	0.3	0.6	0.8	1.5

White, both sexes

2004.....	0.3	5.6	0.2	0.1	0.1	0.1	0.1	0.2	0.2	0.4	0.9	1.7
2003.....	0.3	5.9	0.2	0.1	0.1	0.1	0.1	0.2	0.3	0.4	0.8	1.5
2002.....	0.3	5.0	*	0.1	0.1	0.1	0.1	0.2	0.3	0.5	1.0	1.6
2001.....	0.3	4.6	*	0.1	0.1	0.1	0.1	0.2	0.3	0.6	0.8	1.7
2000.....	0.3	5.1	*	0.1	0.1	0.1	0.2	0.2	0.3	0.4	1.0	2.2
1999.....	0.3	5.2	*	*	0.1	0.1	0.2	0.3	0.3	0.5	0.9	1.7

White, male

2004.....	0.3	5.8	*	*	*	*	0.1	0.3	0.2	0.4	1.0	2.0
2003.....	0.3	6.8	*	*	*	0.1	0.2	0.2	0.3	0.4	1.0	1.7
2002.....	0.3	4.9	*	*	*	0.2	*	0.2	0.3	0.5	0.9	1.9
2001.....	0.3	5.0	*	0.1	*	*	0.1	0.2	0.3	0.6	1.1	2.0
2000.....	0.3	5.2	*	*	*	*	0.2	0.3	0.3	0.5	1.1	3.2
1999.....	0.3	5.5	*	*	*	*	0.2	0.3	0.3	0.5	1.0	2.2

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other congenital malformations and deformations (Q10-Q18,Q35-Q89)

All races 3/, both sexes

2004.....	0.9	36.4	0.8	0.2	0.2	0.2	0.2	0.3	0.4	0.8	1.4	2.6
2003.....	0.9	37.5	0.7	0.2	0.2	0.2	0.2	0.3	0.5	0.7	1.5	2.8
2002.....	0.9	36.2	0.7	0.2	0.3	0.2	0.2	0.4	0.5	0.8	1.3	2.7
2001.....	0.9	36.3	0.8	0.2	0.2	0.2	0.3	0.3	0.5	0.8	1.5	2.5
2000.....	0.9	37.9	0.7	0.2	0.2	0.2	0.3	0.3	0.5	0.8	1.4	2.8
1999.....	0.9	38.3	0.9	0.2	0.2	0.2	0.2	0.3	0.5	0.8	1.5	2.6

All races 3/, male

2004.....	1.0	38.7	0.9	0.2	0.3	0.2	0.3	0.4	0.5	0.9	1.4	3.2
2003.....	1.0	42.8	0.7	0.2	0.3	0.2	0.2	0.4	0.5	0.8	1.4	3.3
2002.....	1.0	39.2	0.6	0.2	0.3	0.2	0.3	0.5	0.5	0.9	1.5	3.6
2001.....	1.0	41.1	0.8	0.2	0.3	0.2	0.3	0.3	0.5	0.9	1.5	3.7
2000.....	1.0	42.1	0.7	0.2	0.2	0.3	0.3	0.4	0.5	0.8	1.4	4.6
1999.....	1.0	42.5	1.1	0.2	0.2	0.2	0.3	0.4	0.5	0.9	1.6	3.0

All races 3/, female

2004.....	0.9	34.0	0.7	0.1	0.2	0.2	0.2	0.3	0.4	0.8	1.4	2.4
2003.....	0.8	31.9	0.7	0.2	0.2	0.2	0.2	0.2	0.5	0.6	1.6	2.6
2002.....	0.8	33.1	0.8	0.2	0.2	0.2	0.2	0.3	0.5	0.7	1.2	2.3
2001.....	0.8	31.3	0.8	0.2	0.2	0.2	0.3	0.3	0.5	0.7	1.5	2.0
2000.....	0.8	33.6	0.6	0.2	0.2	0.2	0.2	0.2	0.6	0.8	1.4	2.1
1999.....	0.8	33.8	0.8	0.2	0.1	0.2	0.2	0.3	0.5	0.8	1.4	2.5

White, both sexes

2004.....	0.9	35.6	0.8	0.2	0.2	0.2	0.2	0.3	0.4	0.9	1.4	2.7
2003.....	0.9	36.4	0.7	0.1	0.2	0.2	0.2	0.3	0.5	0.7	1.5	2.8
2002.....	0.9	37.0	0.7	0.2	0.3	0.2	0.2	0.4	0.5	0.8	1.4	2.7
2001.....	0.9	36.0	0.7	0.2	0.2	0.2	0.3	0.3	0.5	0.8	1.5	2.6
2000.....	0.9	37.4	0.6	0.2	0.2	0.2	0.3	0.3	0.6	0.8	1.4	2.8
1999.....	0.9	36.8	0.9	0.2	0.2	0.2	0.2	0.3	0.5	0.9	1.5	2.7

White, male

2004.....	0.9	37.5	0.8	0.2	0.3	0.2	0.3	0.4	0.4	1.0	1.3	3.3
2003.....	1.0	41.0	0.7	0.1	0.3	0.3	0.3	0.4	0.5	0.8	1.4	3.4
2002.....	1.0	40.4	0.6	0.2	0.3	0.2	0.3	0.5	0.5	0.9	1.5	3.7
2001.....	1.0	40.6	0.6	0.2	0.3	0.2	0.3	0.3	0.5	0.9	1.5	3.8
2000.....	1.0	41.3	0.7	0.2	0.2	0.2	0.3	0.4	0.5	0.8	1.4	4.6
1999.....	1.0	42.1	1.0	0.1	0.2	0.2	0.3	0.4	0.5	1.0	1.6	3.2

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Down's syndrome (Q90)												
All races 3/, both sexes												
2004.....	0.3	2.8	0.1	*	0.1	0.1	0.2	0.5	0.9	0.4	*	*
2003.....	0.3	2.9	*	*	0.1	0.1	0.2	0.5	1.0	0.3	*	*
2002.....	0.3	2.9	0.1	*	0.1	0.1	0.2	0.5	0.9	0.4	*	*
2001.....	0.3	3.0	0.1	*	0.1	0.2	0.2	0.5	0.9	0.3	*	*
2000.....	0.3	2.4	*	*	0.1	0.1	0.1	0.5	0.8	0.4	*	*
1999.....	0.2	2.5	*	*	0.1	0.1	0.1	0.4	0.8	0.2	*	*
All races 3/, male												
2004.....	0.3	2.9	*	*	*	0.1	0.2	0.6	1.0	0.5	*	*
2003.....	0.3	2.6	*	*	*	*	0.2	0.5	1.1	0.3	*	*
2002.....	0.3	3.1	*	*	*	0.1	0.2	0.5	1.0	0.3	*	*
2001.....	0.3	3.4	*	*	*	0.2	0.2	0.5	0.9	0.4	*	*
2000.....	0.3	2.8	*	*	*	0.1	0.2	0.5	0.8	0.3	*	*
1999.....	0.2	2.4	*	*	*	0.1	0.1	0.4	0.8	*	*	*
All races 3/, female												
2004.....	0.3	2.7	*	*	*	*	0.1	0.5	0.9	0.4	*	*
2003.....	0.3	3.2	*	*	*	*	0.2	0.5	0.9	0.3	*	*
2002.....	0.3	2.7	*	*	*	*	0.2	0.4	0.9	0.4	*	*
2001.....	0.3	2.4	*	*	*	0.2	0.2	0.5	1.0	0.3	*	*
2000.....	0.3	2.1	*	*	*	0.2	0.1	0.5	0.8	0.4	*	*
1999.....	0.2	2.5	*	*	*	*	0.1	0.4	0.7	0.3	*	*
White, both sexes												
2004.....	0.3	2.6	*	*	*	0.1	0.2	0.6	1.0	0.5	*	*
2003.....	0.3	2.9	*	*	*	0.1	0.2	0.6	1.1	0.3	*	*
2002.....	0.3	3.1	*	*	0.1	0.1	0.2	0.5	1.0	0.4	*	*
2001.....	0.3	3.1	*	*	0.1	0.2	0.2	0.5	1.0	0.3	*	*
2000.....	0.3	2.4	*	*	*	0.1	0.2	0.5	0.8	0.4	*	*
1999.....	0.2	2.7	*	*	*	0.1	0.1	0.5	0.8	0.2	*	*
White, male												
2004.....	0.3	2.7	*	*	*	*	0.2	0.6	1.1	0.6	*	*
2003.....	0.3	2.4	*	*	*	*	0.2	0.6	1.1	0.3	*	*
2002.....	0.3	3.3	*	*	*	*	0.2	0.5	1.0	0.4	*	*
2001.....	0.3	3.7	*	*	*	0.2	0.2	0.5	0.9	0.4	*	*
2000.....	0.3	3.0	*	*	*	0.1	0.2	0.5	0.8	0.3	*	*
1999.....	0.2	2.6	*	*	*	*	0.1	0.5	0.9	*	*	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

XVIII. Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

All races 3/, both sexes

2004.....	10.3	82.4	1.2	0.3	1.6	2.7	4.3	5.8	7.0	12.6	39.8	221.5
2003.....	10.8	82.9	1.6	0.4	2.0	3.3	5.4	7.0	7.4	13.8	38.9	224.1
2002.....	10.4	85.7	1.6	0.3	1.8	3.3	5.3	6.7	7.5	13.1	38.8	208.0
2001.....	11.4	86.5	1.8	0.4	2.4	3.9	6.9	8.9	9.1	15.1	39.8	206.4
2000.....	11.3	95.6	1.6	0.4	2.2	3.9	6.6	8.3	9.3	16.5	40.7	208.1
1999.....	9.6	93.2	1.5	0.3	1.6	2.7	4.6	5.9	7.8	14.6	36.1	187.7

All races 3/, male

2004.....	9.5	96.9	1.2	0.3	2.1	3.6	5.6	7.6	9.1	15.0	41.8	183.6
2003.....	10.2	96.5	1.8	0.4	2.7	4.5	6.9	9.2	9.6	17.1	41.9	177.9
2002.....	10.1	98.5	1.8	0.4	2.4	4.5	6.9	8.8	10.0	15.9	43.8	175.5
2001.....	11.3	98.9	2.1	0.5	3.2	5.2	9.0	11.9	12.0	18.0	43.8	175.0
2000.....	11.5	110.6	1.7	0.5	2.9	5.3	8.9	11.5	12.3	20.6	46.0	182.7
1999.....	9.4	106.6	1.7	0.4	2.1	3.6	6.0	8.1	10.4	18.2	40.1	166.3

All races 3/, female

2004.....	11.0	67.3	1.1	0.3	1.0	1.8	3.0	4.0	5.1	10.6	38.4	238.5
2003.....	11.4	68.6	1.5	0.4	1.4	2.1	3.8	5.0	5.3	10.9	36.9	244.6
2002.....	10.7	72.3	1.4	0.3	1.2	2.1	3.8	4.7	5.3	10.8	35.4	222.1
2001.....	11.4	73.6	1.5	0.4	1.5	2.5	4.9	6.1	6.3	12.7	37.2	219.5
2000.....	11.2	79.9	1.5	0.4	1.5	2.5	4.4	5.3	6.5	13.2	37.2	218.5
1999.....	9.7	79.2	1.3	0.3	1.0	1.9	3.2	3.8	5.5	11.7	33.5	196.4

White, both sexes

2004.....	10.5	65.8	1.0	0.2	1.6	2.7	4.2	5.5	6.5	12.1	39.5	224.4
2003.....	11.1	68.1	1.4	0.4	2.0	3.2	5.1	6.4	6.9	13.2	39.0	228.3
2002.....	10.5	71.0	1.3	0.3	1.8	3.2	5.1	5.9	7.0	12.7	38.6	211.6
2001.....	11.6	70.9	1.5	0.4	2.4	3.8	6.8	8.6	8.5	14.4	39.6	210.0
2000.....	11.4	77.2	1.5	0.4	2.1	3.7	6.5	7.8	8.6	15.6	40.2	212.2
1999.....	9.4	74.7	1.3	0.3	1.4	2.5	4.1	5.1	7.1	13.9	35.1	190.2

White, male

2004.....	9.4	76.9	1.1	0.3	2.1	3.6	5.5	7.1	8.3	14.4	41.4	186.0
2003.....	10.1	82.4	1.4	0.4	2.7	4.4	6.6	8.4	8.8	16.3	41.7	180.0
2002.....	9.8	82.5	1.4	0.3	2.4	4.3	6.5	7.7	9.2	15.3	42.9	178.2
2001.....	11.2	82.0	1.7	0.4	3.2	5.2	8.9	11.4	11.0	17.2	43.9	177.0
2000.....	11.3	89.4	1.7	0.4	2.9	5.1	8.9	10.7	11.2	19.2	45.5	186.8
1999.....	9.0	86.4	1.6	0.3	2.0	3.2	5.4	6.9	9.3	17.3	38.7	167.3

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

XVIII. Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

White, female

2004.....	11.6	54.2	1.0	0.2	1.0	1.7	2.8	3.9	4.8	10.1	38.2	241.7
2003.....	12.0	53.3	1.3	0.4	1.3	2.0	3.7	4.4	5.1	10.5	37.2	249.7
2002.....	11.2	59.0	1.2	0.2	1.1	2.0	3.6	4.2	5.0	10.5	35.7	226.1
2001.....	11.9	59.3	1.4	0.4	1.5	2.4	4.8	5.8	6.2	12.1	36.7	223.8
2000.....	11.5	64.3	1.3	0.3	1.3	2.3	4.1	4.9	6.1	12.7	36.7	222.5
1999.....	9.9	62.4	1.1	0.3	0.8	1.7	2.7	3.4	5.0	11.2	32.8	199.4

Black, both sexes

2004.....	11.6	171.7	1.9	0.5	1.8	4.0	6.5	9.7	13.3	20.4	53.0	220.8
2003.....	12.3	159.3	3.1	0.8	2.4	5.0	8.5	13.4	14.2	23.4	45.3	210.9
2002.....	12.3	162.3	2.6	0.7	2.5	5.1	8.5	13.7	13.7	20.1	49.6	196.0
2001.....	12.6	169.5	2.8	0.6	2.7	5.0	9.0	13.9	14.8	24.4	49.5	191.6
2000.....	13.7	197.1	2.3	0.7	3.2	5.8	9.2	14.7	17.4	28.9	55.5	186.4
1999.....	12.7	192.8	2.7	0.5	2.6	4.8	9.0	13.2	15.9	24.5	54.1	182.4

Black, male

2004.....	12.6	204.3	2.1	*	2.3	5.2	8.0	13.8	19.0	24.8	59.0	181.8
2003.....	13.5	170.7	3.4	0.8	3.0	6.7	11.1	17.6	20.3	31.4	52.5	193.9
2002.....	14.1	183.0	3.5	0.8	3.0	6.8	11.3	19.4	19.7	26.4	68.3	181.2
2001.....	13.9	187.4	3.5	0.8	3.7	6.0	11.1	18.0	23.3	30.4	51.8	184.9
2000.....	15.5	227.1	2.3	0.8	3.7	7.7	10.8	20.6	25.2	39.9	65.1	161.5
1999.....	14.6	217.6	2.8	0.6	3.5	6.8	11.5	19.4	23.1	32.5	66.1	184.0

Black, female

2004.....	10.8	137.7	1.7	0.6	1.3	3.0	5.1	6.2	8.8	17.3	49.6	236.3
2003.....	11.3	147.5	2.9	0.7	1.9	3.3	6.1	9.8	9.2	17.7	41.3	217.5
2002.....	10.6	140.8	1.8	0.7	2.0	3.6	6.0	8.8	8.9	15.7	39.0	201.7
2001.....	11.4	150.6	2.0	*	1.7	4.1	7.2	10.3	7.9	20.2	48.3	194.1
2000.....	12.0	166.0	2.3	*	2.6	4.1	7.7	9.7	11.0	21.2	50.1	195.7
1999.....	10.9	167.1	2.7	*	1.8	3.1	6.8	7.8	10.1	18.9	47.3	181.8

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Senility (R54)												
All races 3/, both sexes												
2004.....	1.6	*	*	*	*	*	*	*	0.1	0.8	7.1	76.3
2003.....	1.6	*	*	*	*	*	*	*	0.1	0.9	6.8	76.7
2002.....	1.5	*	*	*	*	*	*	*	*	0.8	6.3	72.0
2001.....	1.5	*	*	*	*	*	*	*	*	0.8	6.4	73.0
2000.....	1.3	*	*	*	*	*	*	*	*	0.8	6.3	67.2
1999.....	1.1	*	*	*	*	*	*	*	*	0.7	5.3	56.5
All races 3/, male												
2004.....	0.9	*	*	*	*	*	*	*	*	0.9	7.3	57.2
2003.....	0.9	*	*	*	*	*	*	*	*	1.0	6.6	55.6
2002.....	0.8	*	*	*	*	*	*	*	*	0.8	6.3	56.5
2001.....	0.8	*	*	*	*	*	*	*	*	0.9	6.4	57.2
2000.....	0.7	*	*	*	*	*	*	*	*	0.8	6.4	51.7
1999.....	0.7	*	*	*	*	*	*	*	*	0.9	5.2	45.8
All races 3/, female												
2004.....	2.3	*	*	*	*	*	*	*	*	0.7	7.1	84.9
2003.....	2.3	*	*	*	*	*	*	*	*	0.8	6.9	86.0
2002.....	2.1	*	*	*	*	*	*	*	*	0.8	6.2	78.8
2001.....	2.1	*	*	*	*	*	*	*	*	0.8	6.3	79.6
2000.....	1.9	*	*	*	*	*	*	*	*	0.8	6.2	73.6
1999.....	1.6	*	*	*	*	*	*	*	*	0.5	5.4	60.8
White, both sexes												
2004.....	1.9	*	*	*	*	*	*	*	*	0.8	7.2	79.0
2003.....	1.8	*	*	*	*	*	*	*	0.1	0.9	7.1	79.3
2002.....	1.7	*	*	*	*	*	*	*	*	0.8	6.4	74.2
2001.....	1.7	*	*	*	*	*	*	*	*	0.8	6.5	75.1
2000.....	1.5	*	*	*	*	*	*	*	*	0.8	6.2	69.0
1999.....	1.3	*	*	*	*	*	*	*	*	0.7	5.2	58.8
White, male												
2004.....	1.0	*	*	*	*	*	*	*	*	0.9	7.3	59.9
2003.....	1.0	*	*	*	*	*	*	*	*	1.0	6.8	57.6
2002.....	0.9	*	*	*	*	*	*	*	*	0.8	6.5	58.0
2001.....	0.9	*	*	*	*	*	*	*	*	1.0	6.6	58.3
2000.....	0.8	*	*	*	*	*	*	*	*	0.8	6.3	53.9
1999.....	0.7	*	*	*	*	*	*	*	*	1.0	5.1	47.4

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Senility (R54)												
White, female												
2004.....	2.7	*	*	*	*	*	*	*	*	0.7	7.2	87.6
2003.....	2.7	*	*	*	*	*	*	*	*	0.8	7.3	88.9
2002.....	2.4	*	*	*	*	*	*	*	*	0.8	6.4	81.2
2001.....	2.4	*	*	*	*	*	*	*	*	0.7	6.4	82.2
2000.....	2.2	*	*	*	*	*	*	*	*	0.7	6.1	75.1
1999.....	1.8	*	*	*	*	*	*	*	*	0.5	5.4	63.3
Black, both sexes												
2004.....	0.8	*	*	*	*	*	*	*	*	*	7.9	56.1
2003.....	0.7	*	*	*	*	*	*	*	*	*	4.6	55.7
2002.....	0.7	*	*	*	*	*	*	*	*	*	5.2	55.5
2001.....	0.7	*	*	*	*	*	*	*	*	*	6.7	56.8
2000.....	0.8	*	*	*	*	*	*	*	*	1.5	8.5	55.7
1999.....	0.6	*	*	*	*	*	*	*	*	*	7.8	39.9
Black, male												
2004.....	0.5	*	*	*	*	*	*	*	*	*	10.0	32.8
2003.....	0.4	*	*	*	*	*	*	*	*	*	*	37.5
2002.....	0.4	*	*	*	*	*	*	*	*	*	*	49.3
2001.....	0.5	*	*	*	*	*	*	*	*	*	6.2	57.9
2000.....	0.4	*	*	*	*	*	*	*	*	*	8.8	34.6
1999.....	0.4	*	*	*	*	*	*	*	*	*	9.2	36.3
Black, female												
2004.....	1.1	*	*	*	*	*	*	*	*	*	6.8	65.4
2003.....	1.0	*	*	*	*	*	*	*	*	*	4.1	62.8
2002.....	0.9	*	*	*	*	*	*	*	*	*	5.4	57.9
2001.....	1.0	*	*	*	*	*	*	*	*	*	6.9	56.4
2000.....	1.1	*	*	*	*	*	*	*	*	*	8.4	63.5
1999.....	0.7	*	*	*	*	*	*	*	*	*	7.1	41.2

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Ill-defined and unknown causes of mortality (R95-R99)												
All races 3/, both sexes												
2004.....	5.3	80.7	1.0	0.2	1.2	2.1	3.5	4.6	4.9	6.8	14.3	52.4
2003.....	6.0	81.2	1.3	0.3	1.6	2.6	4.5	5.7	5.4	7.8	15.1	59.5
2002.....	6.0	83.8	1.3	0.2	1.5	2.6	4.4	5.4	5.6	7.6	16.5	58.3
2001.....	7.1	84.8	1.5	0.3	2.0	3.3	6.0	7.8	7.2	9.6	18.1	62.1
2000.....	7.3	93.9	1.2	0.3	1.8	3.2	5.7	7.1	7.2	11.0	20.5	73.0
1999.....	5.9	91.3	1.3	0.3	1.3	2.2	3.7	4.8	5.7	9.2	17.3	66.7
All races 3/, male												
2004.....	6.0	94.9	1.1	0.2	1.7	2.9	4.7	6.1	6.6	8.7	16.7	50.7
2003.....	6.8	94.3	1.4	0.3	2.1	3.6	5.8	7.6	7.3	10.7	18.4	53.6
2002.....	6.8	96.5	1.5	0.3	2.0	3.5	5.8	7.3	7.7	9.8	20.0	55.0
2001.....	8.2	96.5	1.8	0.3	2.7	4.5	7.8	10.4	9.6	12.2	21.8	58.1
2000.....	8.5	108.9	1.4	0.4	2.4	4.4	7.6	10.0	9.8	14.0	24.3	72.1
1999.....	6.7	104.3	1.4	0.3	1.8	2.9	4.9	6.6	7.8	12.0	20.5	64.4
All races 3/, female												
2004.....	4.6	65.8	0.9	0.2	0.8	1.4	2.4	3.1	3.3	5.2	12.8	53.2
2003.....	5.2	67.6	1.2	0.3	1.0	1.6	3.1	4.0	3.6	5.4	12.9	62.1
2002.....	5.1	70.4	1.1	0.2	0.9	1.6	3.0	3.5	3.8	5.8	14.1	59.8
2001.....	6.1	72.4	1.2	0.3	1.2	2.0	4.3	5.3	5.0	7.4	15.7	63.8
2000.....	6.2	78.0	1.1	0.3	1.2	2.0	3.8	4.4	4.9	8.4	17.9	73.3
1999.....	5.2	77.7	1.1	0.3	0.7	1.4	2.6	3.0	3.8	6.8	15.2	67.7
White, both sexes												
2004.....	5.2	64.4	0.8	0.2	1.3	2.2	3.5	4.4	4.7	6.5	14.4	53.1
2003.....	5.9	66.4	1.1	0.3	1.6	2.6	4.4	5.4	5.1	7.6	15.3	60.9
2002.....	5.8	69.5	1.1	0.2	1.5	2.5	4.3	4.8	5.3	7.4	16.5	59.3
2001.....	7.1	69.3	1.3	0.3	2.0	3.3	6.1	7.6	6.9	9.3	18.0	63.2
2000.....	7.2	75.6	1.2	0.3	1.8	3.2	5.6	6.8	6.9	10.5	20.4	74.5
1999.....	5.6	73.1	1.1	0.2	1.2	2.0	3.4	4.2	5.2	8.8	16.9	67.0
White, male												
2004.....	5.8	75.7	0.9	0.1	1.7	3.0	4.8	5.8	6.2	8.5	16.7	51.4
2003.....	6.6	80.2	1.1	0.3	2.2	3.6	5.7	7.2	6.7	10.3	18.4	54.7
2002.....	6.5	81.1	1.1	0.2	2.0	3.5	5.6	6.5	7.2	9.5	19.5	56.2
2001.....	8.1	79.9	1.5	0.3	2.8	4.6	7.9	10.1	9.1	11.8	21.6	59.3
2000.....	8.2	87.8	1.4	0.3	2.4	4.4	7.7	9.5	9.2	13.2	24.2	73.2
1999.....	6.2	84.5	1.3	0.2	1.7	2.7	4.5	5.7	7.0	11.5	19.8	63.1

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Ill-defined and unknown causes of mortality (R95-R99)												
White, female												
2004.....	4.6	52.7	0.7	0.2	0.8	1.3	2.3	3.0	3.2	4.9	12.8	53.8
2003.....	5.2	51.9	1.1	0.3	1.0	1.6	3.1	3.7	3.6	5.4	13.1	63.7
2002.....	5.1	57.4	1.0	0.1	0.9	1.5	3.0	3.2	3.6	5.7	14.4	60.6
2001.....	6.1	58.2	1.1	0.3	1.3	1.9	4.2	5.0	4.9	7.3	15.6	64.9
2000.....	6.2	62.7	0.9	0.2	1.1	1.9	3.5	4.1	4.7	8.3	17.9	75.0
1999.....	5.1	61.2	0.8	0.2	0.6	1.3	2.3	2.8	3.5	6.6	15.0	68.5
Black, both sexes												
2004.....	7.3	168.5	1.7	0.3	1.1	2.5	4.7	7.2	8.1	10.6	16.9	55.0
2003.....	8.2	158.1	2.5	0.5	1.9	3.6	6.6	9.8	9.6	11.8	16.6	54.6
2002.....	8.4	158.0	2.3	0.5	1.6	3.5	6.5	10.6	9.6	11.0	20.4	56.1
2001.....	8.8	167.3	2.3	0.4	1.8	3.9	6.9	11.0	10.3	12.8	22.3	55.6
2000.....	9.9	194.0	1.8	0.5	2.4	4.2	7.5	11.7	12.0	17.0	25.0	66.0
1999.....	9.1	189.1	2.3	0.4	2.0	3.4	7.0	10.1	11.2	13.7	24.3	71.4
Black, male												
2004.....	8.8	199.0	1.7	*	1.4	3.2	5.5	10.5	12.1	12.7	20.3	52.7
2003.....	9.9	168.7	2.8	*	2.3	4.9	8.7	12.9	14.9	18.9	22.0	59.4
2002.....	10.4	178.1	2.9	*	1.9	4.6	9.0	15.4	14.4	15.7	31.3	53.6
2001.....	10.4	184.7	2.9	*	2.5	4.8	8.1	14.2	15.9	17.7	27.4	49.9
2000.....	12.0	224.2	1.9	0.7	2.9	5.7	8.8	16.5	17.5	25.5	32.7	70.3
1999.....	11.2	212.7	2.3	*	2.6	5.0	9.0	15.2	16.9	20.1	31.4	89.1
Black, female												
2004.....	6.0	136.8	1.6	*	0.8	1.9	4.0	4.3	4.9	9.0	14.9	55.9
2003.....	6.7	147.2	2.1	*	1.5	2.3	4.8	7.2	5.4	6.7	13.6	52.7
2002.....	6.5	137.1	1.7	*	1.2	2.5	4.3	6.4	5.7	7.6	14.3	57.1
2001.....	7.4	149.0	1.6	*	1.2	3.1	5.7	8.4	5.8	9.4	19.5	57.7
2000.....	7.9	162.7	1.6	*	2.0	2.8	6.3	7.6	7.6	11.1	20.6	64.4
1999.....	7.2	164.8	2.4	*	1.3	2.1	5.2	5.8	6.7	9.2	20.3	64.8

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other ill-defined and unknown causes of mortality (R96-R99)

All races 3/, both sexes

2004.....	4.5	25.6	1.0	0.2	1.2	2.1	3.5	4.6	4.9	6.8	14.3	52.4
2003.....	5.2	27.2	1.3	0.3	1.6	2.6	4.5	5.7	5.4	7.8	15.1	59.5
2002.....	5.2	26.9	1.3	0.2	1.5	2.6	4.4	5.4	5.6	7.6	16.5	58.3
2001.....	6.3	29.4	1.5	0.3	2.0	3.3	6.0	7.8	7.2	9.6	18.1	62.1
2000.....	6.4	27.6	1.2	0.3	1.8	3.2	5.7	7.1	7.2	11.0	20.5	73.0
1999.....	5.0	21.6	1.3	0.3	1.3	2.2	3.7	4.8	5.7	9.2	17.3	66.7

All races 3/, male

2004.....	5.1	31.2	1.1	0.2	1.7	2.9	4.7	6.1	6.6	8.7	16.7	50.7
2003.....	5.9	31.7	1.4	0.3	2.1	3.6	5.8	7.6	7.3	10.7	18.4	53.6
2002.....	5.9	31.1	1.5	0.3	2.0	3.5	5.8	7.3	7.7	9.8	20.0	55.0
2001.....	7.3	32.7	1.8	0.3	2.7	4.5	7.8	10.4	9.6	12.2	21.8	58.1
2000.....	7.4	31.5	1.4	0.4	2.4	4.4	7.6	10.0	9.8	14.0	24.3	72.1
1999.....	5.5	24.4	1.4	0.3	1.8	2.9	4.9	6.6	7.8	12.0	20.5	64.4

All races 3/, female

2004.....	4.0	19.7	0.9	0.2	0.8	1.4	2.4	3.1	3.3	5.2	12.8	53.2
2003.....	4.6	22.5	1.2	0.3	1.0	1.6	3.1	4.0	3.6	5.4	12.9	62.1
2002.....	4.5	22.5	1.1	0.2	0.9	1.6	3.0	3.5	3.8	5.8	14.1	59.8
2001.....	5.5	25.8	1.2	0.3	1.2	2.0	4.3	5.3	5.0	7.4	15.7	63.8
2000.....	5.5	23.4	1.1	0.3	1.2	2.0	3.8	4.4	4.9	8.4	17.9	73.3
1999.....	4.5	18.6	1.1	0.3	0.7	1.4	2.6	3.0	3.8	6.8	15.2	67.7

White, both sexes

2004.....	4.6	19.1	0.8	0.2	1.3	2.2	3.5	4.4	4.7	6.5	14.4	53.1
2003.....	5.3	22.5	1.1	0.3	1.6	2.6	4.4	5.4	5.1	7.6	15.3	60.9
2002.....	5.2	21.8	1.1	0.2	1.5	2.5	4.3	4.8	5.3	7.4	16.5	59.3
2001.....	6.5	24.6	1.3	0.3	2.0	3.3	6.1	7.6	6.9	9.3	18.0	63.2
2000.....	6.5	21.5	1.2	0.3	1.8	3.2	5.6	6.8	6.9	10.5	20.4	74.5
1999.....	4.9	15.7	1.1	0.2	1.2	2.0	3.4	4.2	5.2	8.8	16.9	67.0

White, male

2004.....	5.1	23.5	0.9	0.1	1.7	3.0	4.8	5.8	6.2	8.5	16.7	51.4
2003.....	5.9	27.3	1.1	0.3	2.2	3.6	5.7	7.2	6.7	10.3	18.4	54.7
2002.....	5.8	25.6	1.1	0.2	2.0	3.5	5.6	6.5	7.2	9.5	19.5	56.2
2001.....	7.3	27.6	1.5	0.3	2.8	4.6	7.9	10.1	9.1	11.8	21.6	59.3
2000.....	7.4	24.4	1.4	0.3	2.4	4.4	7.7	9.5	9.2	13.2	24.2	73.2
1999.....	5.3	18.2	1.3	0.2	1.7	2.7	4.5	5.7	7.0	11.5	19.8	63.1

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other ill-defined and unknown causes of mortality (R96-R99)												
White, female												
2004.....	4.1	14.4	0.7	0.2	0.8	1.3	2.3	3.0	3.2	4.9	12.8	53.8
2003.....	4.7	17.6	1.1	0.3	1.0	1.6	3.1	3.7	3.6	5.4	13.1	63.7
2002.....	4.6	17.9	1.0	0.1	0.9	1.5	3.0	3.2	3.6	5.7	14.4	60.6
2001.....	5.6	21.4	1.1	0.3	1.3	1.9	4.2	5.0	4.9	7.3	15.6	64.9
2000.....	5.7	18.4	0.9	0.2	1.1	1.9	3.5	4.1	4.7	8.3	17.9	75.0
1999.....	4.5	13.1	0.8	0.2	0.6	1.3	2.3	2.8	3.5	6.6	15.0	68.5
Black, both sexes												
2004.....	5.4	58.7	1.7	0.3	1.1	2.5	4.7	7.2	8.1	10.6	16.9	55.0
2003.....	6.4	53.4	2.5	0.5	1.9	3.6	6.6	9.8	9.6	11.8	16.6	54.6
2002.....	6.5	53.8	2.3	0.5	1.6	3.5	6.5	10.6	9.6	11.0	20.4	56.1
2001.....	6.8	53.0	2.3	0.4	1.8	3.9	6.9	11.0	10.3	12.8	22.3	55.6
2000.....	7.6	59.7	1.8	0.5	2.4	4.2	7.5	11.7	12.0	17.0	25.0	66.0
1999.....	6.8	50.5	2.3	0.4	2.0	3.4	7.0	10.1	11.2	13.7	24.3	71.4
Black, male												
2004.....	6.4	69.7	1.7	*	1.4	3.2	5.5	10.5	12.1	12.7	20.3	52.7
2003.....	7.9	59.5	2.8	*	2.3	4.9	8.7	12.9	14.9	18.9	22.0	59.4
2002.....	8.2	60.7	2.9	*	1.9	4.6	9.0	15.4	14.4	15.7	31.3	53.6
2001.....	8.0	58.4	2.9	*	2.5	4.8	8.1	14.2	15.9	17.7	27.4	49.9
2000.....	9.3	70.0	1.9	0.7	2.9	5.7	8.8	16.5	17.5	25.5	32.7	70.3
1999.....	8.5	57.6	2.3	*	2.6	5.0	9.0	15.2	16.9	20.1	31.4	89.1
Black, female												
2004.....	4.6	47.2	1.6	*	0.8	1.9	4.0	4.3	4.9	9.0	14.9	55.9
2003.....	5.1	47.0	2.1	*	1.5	2.3	4.8	7.2	5.4	6.7	13.6	52.7
2002.....	5.0	46.6	1.7	*	1.2	2.5	4.3	6.4	5.7	7.6	14.3	57.1
2001.....	5.7	47.3	1.6	*	1.2	3.1	5.7	8.4	5.8	9.4	19.5	57.7
2000.....	6.1	48.9	1.6	*	2.0	2.8	6.3	7.6	7.6	11.1	20.6	64.4
1999.....	5.3	43.3	2.4	*	1.3	2.1	5.2	5.8	6.7	9.2	20.3	64.8

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

All other symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R53,R55-R94)

All races 3/, both sexes

2004.....	3.3	1.7	0.2	0.1	0.3	0.6	0.8	1.2	2.1	5.1	18.3	92.7
2003.....	3.2	1.6	0.3	0.1	0.4	0.7	0.9	1.3	1.9	5.1	17.0	87.9
2002.....	2.9	1.9	0.3	0.1	0.4	0.7	0.9	1.3	1.8	4.7	16.0	77.7
2001.....	2.7	1.8	0.3	0.1	0.4	0.6	0.9	1.2	1.8	4.7	15.3	71.3
2000.....	2.6	1.8	0.4	0.1	0.4	0.7	0.9	1.2	2.0	4.8	13.9	67.9
1999.....	2.5	1.9	0.3	0.1	0.3	0.6	0.8	1.1	2.1	4.8	13.5	64.5

All races 3/, male

2004.....	2.5	2.0	*	0.1	0.5	0.7	0.9	1.4	2.4	5.5	17.9	75.7
2003.....	2.5	2.2	0.4	0.1	0.5	0.9	1.1	1.6	2.3	5.4	17.0	68.6
2002.....	2.4	1.9	0.4	0.1	0.5	0.9	1.1	1.5	2.2	5.3	17.6	64.0
2001.....	2.3	2.3	0.3	0.2	0.5	0.7	1.2	1.5	2.3	4.9	15.6	59.7
2000.....	2.3	1.7	0.3	0.1	0.5	0.8	1.3	1.5	2.4	5.7	15.3	58.9
1999.....	2.1	2.3	0.3	0.1	0.4	0.7	1.1	1.4	2.5	5.3	14.4	56.1

All races 3/, female

2004.....	4.1	1.5	*	0.1	0.2	0.5	0.6	1.0	1.8	4.7	18.6	100.4
2003.....	3.9	1.1	0.3	0.1	0.3	0.5	0.7	1.0	1.6	4.8	17.1	96.4
2002.....	3.5	1.9	*	*	0.3	0.5	0.8	1.1	1.4	4.2	15.0	83.6
2001.....	3.2	1.2	0.3	0.1	0.3	0.5	0.7	0.8	1.3	4.6	15.1	76.2
2000.....	3.0	1.8	0.4	0.1	0.3	0.5	0.6	0.9	1.6	4.0	13.1	71.6
1999.....	2.9	1.5	*	*	0.3	0.5	0.6	0.7	1.7	4.4	12.9	67.9

White, both sexes

2004.....	3.5	1.4	0.2	0.1	0.3	0.5	0.6	1.1	1.8	4.8	17.9	92.3
2003.....	3.3	1.8	0.3	0.1	0.4	0.6	0.8	1.0	1.7	4.6	16.7	88.1
2002.....	3.0	1.5	0.3	0.1	0.3	0.6	0.8	1.1	1.6	4.5	15.7	78.2
2001.....	2.8	1.7	0.2	0.1	0.3	0.5	0.8	1.0	1.5	4.3	15.1	71.6
2000.....	2.7	1.6	0.4	0.1	0.4	0.6	0.8	1.0	1.7	4.3	13.5	68.8
1999.....	2.5	1.6	0.3	0.1	0.3	0.5	0.7	0.9	1.8	4.4	13.0	64.5

White, male

2004.....	2.5	1.2	*	*	0.4	0.5	0.7	1.3	2.0	5.1	17.4	74.7
2003.....	2.5	2.2	0.4	*	0.5	0.8	0.9	1.3	2.0	5.1	16.5	67.6
2002.....	2.4	1.4	*	*	0.3	0.8	0.9	1.2	1.9	5.0	16.9	64.0
2001.....	2.2	2.2	*	0.1	0.4	0.6	1.0	1.3	1.9	4.4	15.7	59.4
2000.....	2.2	1.6	0.3	*	0.5	0.7	1.2	1.2	1.9	5.1	14.9	59.8
1999.....	2.1	1.9	*	0.1	0.3	0.5	0.9	1.1	2.2	4.9	13.8	56.8

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

All other symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R53,R55-R94)

White, female

2004.....	4.4	1.5	*	*	0.2	0.4	0.6	0.9	1.6	4.5	18.2	100.3
2003.....	4.2	1.3	*	*	0.3	0.4	0.7	0.8	1.4	4.3	16.8	97.1
2002.....	3.7	1.6	*	*	0.2	0.4	0.7	1.0	1.3	4.1	14.9	84.3
2001.....	3.4	*	*	*	0.3	0.4	0.6	0.7	1.2	4.1	14.7	76.7
2000.....	3.2	1.6	0.4	*	0.3	0.4	0.5	0.8	1.4	3.7	12.6	72.4
1999.....	3.0	*	*	*	0.2	0.4	0.5	0.6	1.5	4.0	12.4	67.6

Black, both sexes

2004.....	3.5	3.2	*	*	0.7	1.5	1.7	2.5	5.0	8.9	28.1	109.7
2003.....	3.4	*	*	*	0.6	1.4	1.8	3.6	4.4	10.7	24.1	100.7
2002.....	3.2	4.3	*	*	0.9	1.6	2.0	3.1	4.1	8.1	24.0	84.3
2001.....	3.1	*	*	*	0.9	1.1	2.2	2.8	4.3	10.5	20.5	79.2
2000.....	3.0	*	*	*	0.7	1.6	1.7	3.0	5.2	10.4	22.0	64.8
1999.....	3.0	3.6	*	*	0.7	1.4	2.0	3.0	4.5	10.0	22.0	71.1

Black, male

2004.....	3.3	*	*	*	0.9	2.0	2.5	3.2	6.6	10.8	28.7	96.4
2003.....	3.2	*	*	*	0.7	1.8	2.4	4.7	5.3	11.0	25.1	97.0
2002.....	3.3	*	*	*	1.1	2.3	2.4	3.9	5.4	9.4	32.1	78.3
2001.....	3.1	*	*	*	1.2	1.2	3.0	3.7	7.2	12.0	18.2	77.2
2000.....	3.0	*	*	*	0.8	1.9	2.0	4.1	7.6	13.3	23.6	56.5
1999.....	3.0	*	*	*	0.9	1.8	2.5	4.2	5.9	11.3	25.5	58.6

Black, female

2004.....	3.7	*	*	*	*	1.1	1.1	1.9	3.7	7.5	27.9	115.0
2003.....	3.6	*	*	*	*	1.0	1.3	2.6	3.7	10.5	23.6	102.1
2002.....	3.2	*	*	*	0.7	1.1	1.8	2.4	3.1	7.2	19.3	86.7
2001.....	3.0	*	*	*	*	1.0	1.4	1.9	2.0	9.4	21.8	80.0
2000.....	3.0	*	*	*	*	1.3	1.4	2.1	3.3	8.3	21.1	67.8
1999.....	3.0	*	*	*	*	1.0	1.6	2.0	3.3	9.1	20.0	75.8

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
XX. External causes of mortality (*U01-*U03,V01-Y89)												
All races 3/, both sexes												
2004.....	57.9	36.7	13.0	8.3	61.1	58.9	62.9	66.1	52.8	62.6	129.5	307.5
2003.....	57.4	34.4	13.6	8.0	61.5	58.1	63.9	63.5	52.2	62.8	127.7	310.4
2002.....	56.9	33.8	13.7	8.4	62.2	57.9	63.7	60.8	50.7	63.9	128.4	308.3
2001 4/.....	56.2	35.1	14.3	8.5	60.6	58.1	63.4	58.9	49.9	62.7	128.6	310.1
2000.....	53.8	34.5	14.6	9.1	60.0	54.1	59.2	54.8	48.6	60.9	122.6	310.5
1999.....	54.2	33.9	15.3	9.4	59.7	55.2	58.7	53.2	48.2	64.4	128.6	318.2
All races 3/, male												
2004.....	80.3	40.4	14.7	10.1	92.8	90.8	90.8	95.7	76.8	89.9	180.9	407.1
2003.....	80.2	39.2	15.9	10.1	93.6	89.9	93.4	92.7	76.9	89.8	179.3	417.7
2002.....	79.8	37.6	16.3	10.4	95.1	89.9	93.3	89.2	75.3	92.9	180.3	414.8
2001 4/.....	79.3	38.9	16.8	10.7	93.7	90.5	93.9	88.0	73.0	89.9	183.3	429.1
2000.....	75.8	38.1	17.0	11.4	91.9	84.5	88.8	82.5	71.1	86.9	175.5	438.2
1999.....	76.2	38.3	17.3	11.8	90.8	85.7	87.7	80.1	71.5	92.9	183.6	454.8
All races 3/, female												
2004.....	36.2	32.9	11.3	6.3	27.6	26.0	35.1	37.5	30.5	39.7	94.8	262.7
2003.....	35.3	29.3	11.2	5.7	27.5	25.4	34.6	35.4	29.4	40.3	93.3	262.9
2002.....	34.8	29.8	11.1	6.3	27.5	25.1	34.3	33.3	28.0	39.8	94.0	262.0
2001 4/.....	34.0	31.2	11.8	6.2	25.8	25.1	33.1	30.8	28.5	40.1	92.7	260.3
2000.....	32.5	30.8	12.1	6.7	26.5	23.3	29.9	28.1	27.9	39.4	88.1	258.5
1999.....	33.0	29.2	13.3	6.9	27.0	24.3	30.1	27.3	26.8	41.0	92.8	263.2
White, both sexes												
2004.....	59.0	31.3	11.9	7.6	59.4	56.5	64.1	66.7	52.8	63.0	132.7	321.4
2003.....	58.2	28.9	12.5	7.4	58.9	55.5	64.6	63.7	52.3	63.0	130.6	323.4
2002.....	57.4	26.1	12.3	7.7	59.7	54.8	63.9	60.6	50.7	64.2	131.4	319.7
2001 4/.....	56.5	28.8	13.3	8.0	57.4	55.4	63.2	58.2	49.5	62.6	131.1	319.5
2000.....	53.6	27.6	13.1	8.7	56.6	50.6	58.3	53.6	48.2	60.3	124.5	320.4
1999.....	53.7	27.9	13.6	8.8	55.8	51.9	57.7	51.7	47.0	63.8	129.5	326.8
White, male												
2004.....	80.3	35.3	13.7	9.3	88.3	85.0	91.5	95.3	75.6	90.2	184.9	423.8
2003.....	79.7	32.8	14.7	9.3	87.5	84.0	93.2	91.9	76.2	90.1	182.8	434.4
2002.....	79.0	30.7	14.5	9.4	89.2	83.2	92.8	88.1	74.4	92.5	183.4	427.4
2001 4/.....	78.3	31.8	15.8	9.8	86.7	84.7	93.1	86.1	71.6	89.5	186.2	441.2
2000.....	74.1	31.8	15.4	10.8	84.6	77.5	86.8	79.6	69.4	85.4	177.4	451.7
1999.....	74.2	32.6	15.4	10.9	82.7	79.2	85.5	76.9	68.9	91.5	184.6	465.8

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
XX. External causes of mortality (*U01-*U03,V01-Y89)												
White, female												
2004.....	38.1	27.0	10.2	5.8	28.5	26.3	36.1	38.5	31.2	39.8	97.0	275.2
2003.....	37.1	24.8	10.3	5.4	28.3	25.4	35.4	36.0	29.6	39.9	95.4	274.2
2002.....	36.4	21.3	10.1	5.8	28.3	25.0	34.6	33.6	28.4	40.2	96.5	272.8
2001 4/.....	35.2	25.7	10.5	6.0	26.3	24.7	32.8	30.8	28.7	39.8	94.5	268.6
2000.....	33.6	23.2	10.7	6.5	26.7	22.6	29.5	28.0	28.2	39.3	89.6	267.0
1999.....	33.9	23.1	11.7	6.5	27.2	23.4	29.6	27.0	26.5	40.6	93.3	270.9
Black, both sexes												
2004.....	62.3	66.5	19.4	11.6	77.9	87.6	70.4	76.3	62.7	64.9	109.8	176.0
2003.....	62.8	63.5	20.2	10.8	81.7	87.4	73.9	75.0	60.0	66.9	103.4	184.4
2002.....	63.7	72.8	21.0	12.1	82.8	89.2	75.9	74.3	58.7	65.6	106.5	199.6
2001 4/.....	64.2	68.9	21.2	11.6	84.3	85.6	77.5	75.5	60.7	70.7	110.6	216.8
2000.....	63.9	70.6	22.8	12.1	86.1	84.9	76.4	73.7	58.7	72.3	108.7	212.4
1999.....	65.2	67.1	24.8	13.3	87.1	84.0	76.3	73.1	61.8	76.2	127.6	239.5
Black, male												
2004.....	96.9	68.2	21.4	14.1	129.8	152.3	109.1	119.7	102.6	101.8	168.7	247.4
2003.....	98.2	72.0	23.9	13.9	137.7	151.0	116.8	118.4	97.6	99.4	154.6	256.5
2002.....	99.9	73.5	26.0	15.6	138.8	155.0	117.8	117.8	97.8	105.0	166.7	307.7
2001 4/.....	100.2	77.2	24.1	15.9	144.0	146.6	118.7	122.6	98.8	107.6	168.9	306.4
2000.....	100.2	72.3	25.5	15.7	145.1	146.1	121.3	120.3	97.1	114.3	168.0	318.3
1999.....	101.7	71.2	27.7	17.3	147.0	142.9	120.8	121.1	101.2	119.3	189.4	367.9
Black, female												
2004.....	30.8	64.8	17.4	9.0	25.2	28.2	36.1	38.9	30.6	38.6	76.5	147.7
2003.....	30.5	54.7	16.4	7.7	24.9	29.3	36.0	37.7	29.8	43.9	74.5	156.4
2002.....	30.9	72.0	15.8	8.5	26.4	29.5	38.9	37.0	27.3	37.6	72.4	158.1
2001 4/.....	31.5	60.2	18.3	7.2	24.8	30.4	40.9	35.0	30.1	44.8	77.8	183.6
2000.....	30.9	68.8	19.9	8.4	27.8	29.7	36.7	33.6	27.6	42.7	75.3	173.0
1999.....	32.2	62.8	21.9	9.1	27.9	30.9	37.0	32.1	30.2	45.7	92.9	191.4

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All	Under 1	1-4	5-14	15-24	25-34	35-44	45-54	55-64	65-74	75-84	85 years
	ages 1/ year 1/	year 2/ year 2/	years	years	years	years	years	years	years	years	years	and over

Accidents (unintentional injuries) (V01-X59,Y85-Y86)

All races 3/, both sexes

2004.....	38.1	25.8	10.3	6.5	37.0	32.6	37.3	40.7	33.2	44.0	103.7	276.7
2003.....	37.6	23.6	10.9	6.4	37.1	31.5	37.8	38.8	32.9	44.1	101.9	278.9
2002.....	37.0	23.5	10.5	6.6	38.0	31.5	37.2	36.6	31.4	44.2	101.3	275.4
2001.....	35.7	24.2	11.2	6.9	36.1	29.9	35.4	34.1	30.3	42.8	100.9	276.4
2000.....	34.8	23.1	11.9	7.3	36.0	29.5	34.1	32.6	30.9	41.9	95.1	273.5
1999.....	35.1	22.3	12.4	7.6	35.3	29.6	33.8	31.8	30.6	44.6	100.5	282.4

All races 3/, male

2004.....	49.8	29.0	11.8	8.0	53.3	48.5	53.0	58.6	46.9	59.2	134.7	345.5
2003.....	49.3	26.4	13.0	7.9	53.3	47.1	54.2	55.9	47.1	58.9	132.9	352.5
2002.....	48.9	26.8	12.9	8.2	54.9	47.7	53.7	53.3	45.2	60.6	131.6	346.6
2001.....	47.2	26.2	13.3	8.6	52.9	45.4	51.8	50.6	43.0	57.3	133.4	358.5
2000.....	46.2	25.7	14.0	8.8	52.1	45.0	51.0	49.0	44.1	55.9	124.3	359.7
1999.....	46.4	25.4	14.4	9.4	50.7	45.0	50.2	47.9	44.0	60.2	132.4	375.4

All races 3/, female

2004.....	26.8	22.5	8.7	5.1	19.9	16.0	21.7	23.4	20.5	31.1	82.9	245.8
2003.....	26.2	20.7	8.7	4.8	19.9	15.3	21.4	22.4	19.7	31.6	81.1	246.4
2002.....	25.6	20.0	8.1	5.0	20.1	14.9	20.8	20.5	18.6	30.6	81.2	244.6
2001.....	24.5	22.1	8.9	5.2	18.4	14.0	19.2	18.0	18.5	30.8	79.6	242.0
2000.....	23.8	20.5	9.7	5.6	19.1	13.7	17.5	16.8	18.8	30.3	76.1	238.4
1999.....	24.1	19.0	10.3	5.6	19.2	14.0	17.6	16.3	18.3	31.8	79.7	244.8

White, both sexes

2004.....	40.2	22.2	10.0	6.2	40.3	34.2	38.9	40.7	32.6	43.9	106.3	289.9
2003.....	39.5	20.0	10.6	6.1	40.1	33.0	39.0	38.5	32.3	43.8	104.0	291.1
2002.....	38.7	18.6	9.9	6.2	40.8	32.8	38.1	36.2	30.7	43.9	103.6	286.1
2001.....	37.0	20.6	11.0	6.4	38.2	30.9	35.8	33.1	29.4	42.4	102.7	285.5
2000.....	35.9	19.1	11.1	6.9	38.3	29.9	34.3	31.3	30.2	41.1	96.5	282.8
1999.....	36.0	18.9	11.5	7.1	37.2	30.0	33.8	30.5	29.2	43.5	100.9	290.4

White, male

2004.....	52.0	25.5	11.5	7.5	57.8	50.5	54.8	58.0	45.1	58.8	137.0	359.4
2003.....	51.3	22.5	12.8	7.6	57.4	49.3	55.6	54.9	45.8	58.3	134.4	365.7
2002.....	50.4	22.3	11.8	7.6	58.7	49.3	54.7	52.3	43.5	59.2	133.0	356.6
2001.....	48.4	22.1	13.3	7.8	55.8	46.7	52.3	48.6	41.1	56.3	134.3	368.1
2000.....	47.0	21.7	13.2	8.4	55.0	45.4	51.0	46.7	42.1	54.2	124.3	369.8
1999.....	46.9	21.8	13.2	8.8	52.8	45.4	50.0	45.4	41.5	58.0	131.8	383.6

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidents (unintentional injuries) (V01-X59,Y85-Y86)												
White, female												
2004.....	28.8	18.8	8.4	4.7	21.5	17.0	22.6	23.6	20.7	31.2	85.4	258.6
2003.....	28.0	17.4	8.3	4.7	21.7	15.9	22.1	22.3	19.5	31.4	83.5	258.0
2002.....	27.3	14.7	7.8	4.7	21.8	15.5	21.3	20.3	18.6	30.9	83.9	255.4
2001.....	25.9	18.9	8.5	5.0	19.5	14.4	19.2	17.8	18.3	30.7	81.6	251.0
2000.....	25.1	16.4	9.0	5.4	20.4	13.8	17.4	16.2	18.9	30.1	78.1	247.4
1999.....	25.3	15.8	9.6	5.4	20.5	14.1	17.4	15.8	17.7	31.3	80.6	253.0
Black, both sexes												
2004.....	32.8	45.8	12.5	8.5	25.8	31.4	36.7	49.8	43.7	48.2	87.1	152.3
2003.....	32.4	43.5	13.4	7.7	25.6	29.9	38.6	48.4	42.3	50.4	82.7	160.2
2002.....	33.1	48.9	14.1	9.1	27.8	31.4	39.1	46.8	41.4	49.3	83.6	173.4
2001.....	33.5	43.1	13.6	9.4	28.7	30.2	40.0	47.7	42.0	51.0	88.5	183.6
2000.....	33.5	44.5	16.2	9.3	28.7	32.5	39.7	47.0	40.8	52.5	84.9	181.1
1999.....	35.2	41.6	17.7	10.5	29.5	32.2	40.0	47.8	44.7	57.8	103.3	207.6
Black, male												
2004.....	46.7	48.3	14.2	10.3	37.8	50.3	54.1	76.7	70.5	73.7	132.0	213.6
2003.....	46.1	47.6	15.7	9.7	38.6	46.7	57.8	74.5	67.9	74.3	123.3	225.2
2002.....	47.9	51.1	18.5	11.6	41.8	50.6	58.4	72.2	68.7	76.9	129.7	260.5
2001.....	48.2	48.5	15.4	12.8	43.2	47.4	59.9	77.2	67.6	74.9	136.0	255.3
2000.....	49.0	47.7	18.8	12.0	43.0	51.8	62.0	75.4	68.8	79.4	132.6	275.6
1999.....	51.0	45.9	21.0	13.8	44.9	51.4	62.0	78.6	71.1	88.2	150.0	316.4
Black, female												
2004.....	20.1	43.2	10.6	6.6	13.6	14.1	21.3	26.7	22.2	30.0	61.7	128.0
2003.....	19.9	39.3	11.0	5.7	12.5	14.5	21.6	26.1	21.7	33.4	59.8	134.9
2002.....	19.7	46.6	9.6	6.6	13.8	14.1	22.0	25.1	19.5	29.7	57.5	140.1
2001.....	20.1	37.5	11.9	6.0	14.2	14.7	22.4	22.3	21.4	34.2	61.8	157.0
2000.....	19.5	41.3	13.5	6.5	14.6	15.0	19.9	22.6	18.3	33.5	58.0	145.9
1999.....	20.8	37.3	14.3	7.1	14.3	14.9	20.6	21.4	23.5	36.4	77.1	166.9

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Transport accidents (V01-V99,Y85)												
All races 3/, both sexes												
2004.....	16.4	3.6	4.1	4.3	27.3	18.8	16.4	16.7	15.5	17.5	25.5	27.2
2003.....	16.5	3.7	4.1	4.3	27.7	18.3	17.1	16.6	15.9	17.5	26.2	29.7
2002.....	16.8	3.1	4.1	4.2	29.1	18.7	17.1	16.4	15.7	18.3	27.0	28.8
2001.....	16.6	3.7	4.4	4.4	27.9	18.8	17.4	16.0	15.3	17.9	27.9	28.9
2000.....	16.6	4.6	4.5	4.7	28.1	18.6	17.0	15.8	16.0	18.0	27.0	31.5
1999.....	16.6	5.0	4.6	4.8	27.6	18.5	16.8	15.4	16.0	19.4	28.4	32.0
All races 3/, male												
2004.....	23.0	3.9	4.4	5.0	37.8	27.7	24.0	24.7	22.2	23.7	36.5	46.5
2003.....	23.3	4.0	4.5	5.2	38.6	27.3	24.8	24.3	23.0	23.7	36.6	52.0
2002.....	23.7	3.5	4.8	4.9	40.7	28.1	24.6	24.0	22.5	25.3	37.7	53.0
2001.....	23.5	3.3	5.0	5.3	39.6	28.3	25.1	23.3	21.6	24.3	39.8	52.7
2000.....	23.3	4.8	5.0	5.4	39.3	27.7	24.8	23.0	22.5	24.3	38.1	59.8
1999.....	23.1	5.2	5.1	5.8	38.1	27.2	24.0	22.4	22.5	26.1	40.7	63.5
All races 3/, female												
2004.....	9.9	3.2	3.8	3.6	16.1	9.5	8.8	9.0	9.3	12.3	18.1	18.5
2003.....	10.0	3.4	3.7	3.4	16.3	9.0	9.4	9.1	9.4	12.4	19.3	19.9
2002.....	10.1	2.8	3.4	3.5	16.9	9.2	9.7	9.0	9.3	12.5	19.8	18.3
2001.....	10.0	4.2	3.8	3.5	15.6	9.2	9.8	8.9	9.4	12.7	20.1	18.9
2000.....	10.2	4.3	4.0	3.9	16.4	9.3	9.3	8.9	10.0	12.8	19.8	19.9
1999.....	10.4	4.8	4.2	3.9	16.6	9.6	9.6	8.7	10.1	14.0	20.4	19.3
White, both sexes												
2004.....	16.9	3.3	4.0	4.4	29.5	19.1	16.7	16.7	15.5	17.6	25.9	28.4
2003.....	17.0	3.6	4.0	4.4	29.8	18.7	17.4	16.6	15.7	17.5	26.5	30.5
2002.....	17.3	2.7	3.9	4.2	31.3	19.1	17.3	16.4	15.6	18.4	27.5	29.6
2001.....	17.0	3.7	4.3	4.3	29.5	19.2	17.6	15.8	15.1	17.9	28.5	29.9
2000.....	17.0	4.1	4.3	4.7	30.0	18.5	17.1	15.6	16.0	17.9	27.5	32.2
1999.....	17.0	4.6	4.1	4.8	29.3	18.7	16.8	15.2	15.7	19.4	28.6	32.3
White, male												
2004.....	23.7	3.3	4.2	5.1	40.8	27.8	24.4	24.4	22.0	23.6	36.7	48.6
2003.....	23.9	4.0	4.5	5.2	41.1	27.8	25.0	24.1	22.7	23.7	36.7	52.8
2002.....	24.2	3.1	4.5	5.0	43.4	28.3	24.7	24.0	22.3	24.9	37.7	53.8
2001.....	23.9	3.2	4.9	5.1	41.7	28.7	25.1	22.9	21.2	24.0	40.3	54.1
2000.....	23.6	4.5	4.7	5.5	41.6	27.4	24.7	22.6	22.0	23.7	38.1	61.6
1999.....	23.4	5.2	4.4	5.7	40.1	27.3	23.8	21.8	21.9	25.8	40.6	63.7

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Transport accidents (V01-V99,Y85)												
White, female												
2004.....	10.3	3.2	3.8	3.6	17.4	9.9	8.9	9.0	9.5	12.4	18.5	19.3
2003.....	10.3	3.1	3.4	3.5	17.7	9.1	9.6	9.1	9.1	12.2	19.7	20.6
2002.....	10.5	2.2	3.4	3.4	18.3	9.4	9.8	8.9	9.4	12.8	20.6	19.1
2001.....	10.3	4.1	3.6	3.5	16.5	9.3	10.0	8.8	9.4	12.7	20.7	19.7
2000.....	10.5	3.7	3.8	4.0	17.7	9.3	9.4	8.7	10.3	12.9	20.5	20.3
1999.....	10.7	3.9	3.8	3.9	17.8	9.8	9.6	8.6	10.0	14.0	20.7	19.7
Black, both sexes												
2004.....	15.2	4.8	5.1	4.4	19.6	20.8	17.1	19.5	17.3	16.4	21.6	14.7
2003.....	15.1	4.2	5.0	4.1	19.7	19.3	18.3	18.1	18.5	18.2	22.4	21.0
2002.....	15.4	5.3	5.0	4.6	21.2	20.4	18.2	17.8	17.1	17.1	21.7	18.4
2001.....	15.9	*	5.2	5.3	22.2	19.9	18.7	18.8	18.1	18.8	21.2	17.5
2000.....	16.0	6.5	5.6	4.7	21.7	21.4	18.6	19.3	17.4	19.6	20.0	21.6
1999.....	16.3	7.3	7.1	5.4	22.0	19.9	19.0	19.0	18.3	19.3	26.0	27.7
Black, male												
2004.....	22.8	5.9	5.4	4.8	27.9	33.8	26.8	30.9	27.7	26.7	38.4	28.8
2003.....	22.6	*	5.1	5.1	29.3	30.6	28.4	29.0	29.1	26.6	37.5	42.7
2002.....	23.1	*	6.4	5.3	30.5	33.1	27.5	27.8	28.3	28.5	40.4	46.1
2001.....	23.8	*	5.7	6.9	32.5	31.6	28.9	30.2	29.6	27.4	36.2	35.2
2000.....	24.1	6.7	6.2	5.7	31.6	34.4	29.7	29.6	29.4	32.2	36.3	45.0
1999.....	24.2	*	8.0	6.8	32.0	31.8	29.0	30.3	28.9	29.1	43.4	68.0
Black, female												
2004.....	8.2	*	4.8	3.9	11.1	8.9	8.6	9.6	9.0	9.1	12.1	9.1
2003.....	8.3	*	5.0	3.1	9.8	9.0	9.4	8.8	9.9	12.2	13.9	12.6
2002.....	8.3	*	3.4	3.9	11.8	9.0	9.9	9.2	8.1	9.0	11.0	*
2001.....	8.7	*	4.7	3.7	11.9	9.4	9.7	9.1	8.9	12.7	12.7	10.9
2000.....	8.5	*	5.0	3.6	11.9	9.7	8.7	10.5	7.6	10.8	10.7	12.9
1999.....	9.2	8.7	6.2	4.0	12.2	9.0	10.1	9.4	9.7	12.4	16.4	12.7

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Railway accidents (V05,V15,V80.6,V81.2-V81.9)												
All races 3/, both sexes												
2004.....	0.2	*	*	*	0.2	0.2	0.2	0.2	0.1	0.1	*	*
2003.....	0.1	*	*	*	0.2	0.2	0.2	0.2	0.1	*	*	*
2002.....	0.2	*	*	*	0.2	0.2	0.3	0.2	0.1	*	*	*
2001.....	0.2	*	*	*	0.2	0.2	0.3	0.2	0.2	*	*	*
2000.....	0.2	*	*	*	0.2	0.2	0.3	0.2	0.1	*	*	*
1999.....	0.2	*	*	0.1	0.2	0.2	0.3	0.2	0.1	*	*	*
All races 3/, male												
2004.....	0.3	*	*	*	0.3	0.4	0.4	0.4	0.2	*	*	*
2003.....	0.3	*	*	*	0.3	0.3	0.4	0.4	0.2	*	*	*
2002.....	0.3	*	*	*	0.4	0.4	0.5	0.4	0.3	*	*	*
2001.....	0.3	*	*	*	0.3	0.4	0.5	0.4	0.3	*	*	*
2000.....	0.3	*	*	*	0.4	0.3	0.5	0.4	0.2	*	*	*
1999.....	0.3	*	*	0.1	0.4	0.4	0.5	0.3	0.2	*	*	*
All races 3/, female												
2004.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2003.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	*	*	*	*	*	*
White, both sexes												
2004.....	0.1	*	*	*	0.1	0.2	0.2	0.2	0.1	*	*	*
2003.....	0.1	*	*	*	0.2	0.2	0.2	0.2	0.1	*	*	*
2002.....	0.2	*	*	*	0.2	0.2	0.3	0.2	0.1	*	*	*
2001.....	0.2	*	*	*	0.2	0.2	0.3	0.2	0.1	*	*	*
2000.....	0.2	*	*	*	0.2	0.2	0.3	0.2	0.1	*	*	*
1999.....	0.2	*	*	*	0.2	0.2	0.3	0.2	0.1	*	*	*
White, male												
2004.....	0.3	*	*	*	0.3	0.4	0.4	0.4	0.2	*	*	*
2003.....	0.3	*	*	*	0.3	0.4	0.4	0.4	0.2	*	*	*
2002.....	0.3	*	*	*	0.4	0.4	0.4	0.3	0.2	*	*	*
2001.....	0.3	*	*	*	0.3	0.4	0.4	0.4	0.2	*	*	*
2000.....	0.3	*	*	*	0.4	0.3	0.5	0.3	0.2	*	*	*
1999.....	0.3	*	*	*	0.4	0.4	0.5	0.3	0.2	*	*	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)

All races 3/, both sexes

2004.....	15.3	3.5	4.0	4.1	26.3	17.6	15.1	15.1	14.1	16.1	24.5	26.1
2003.....	15.4	3.6	3.9	4.0	26.6	17.1	15.7	14.9	14.2	16.2	24.9	28.8
2002.....	15.7	3.0	3.9	3.9	28.2	17.8	15.8	14.8	14.1	17.0	25.7	28.0
2001.....	15.4	3.6	4.1	4.1	26.8	17.5	15.7	14.2	13.7	16.3	26.3	27.8
2000.....	15.4	4.4	4.2	4.3	26.9	17.3	15.3	14.2	14.4	16.5	25.7	30.4
1999.....	15.2	4.8	4.2	4.3	26.2	16.9	14.9	13.6	14.2	17.8	26.5	30.4

All races 3/, male

2004.....	21.3	3.8	4.2	4.7	36.4	25.8	21.9	21.9	19.8	21.3	34.7	44.2
2003.....	21.4	3.9	4.4	4.8	36.9	25.5	22.5	21.5	20.0	21.3	34.3	50.0
2002.....	21.9	3.3	4.5	4.6	39.3	26.5	22.3	21.3	19.9	22.7	35.3	51.7
2001.....	21.5	3.2	4.5	4.9	38.0	26.2	22.4	20.5	19.0	21.5	37.3	51.1
2000.....	21.3	4.6	4.7	5.0	37.4	25.5	22.0	20.4	19.8	21.7	35.6	57.5
1999.....	20.9	5.0	4.7	5.1	36.0	24.7	21.1	19.4	19.3	23.3	38.0	59.8

All races 3/, female

2004.....	9.5	3.2	3.7	3.4	15.7	9.1	8.3	8.5	8.8	11.7	17.7	17.9
2003.....	9.5	3.3	3.5	3.2	15.8	8.5	8.9	8.5	8.8	11.9	18.7	19.5
2002.....	9.8	2.8	3.3	3.3	16.6	8.8	9.3	8.6	8.9	12.3	19.3	17.7
2001.....	9.5	4.0	3.6	3.3	15.1	8.7	9.1	8.1	8.8	12.0	19.2	18.0
2000.....	9.7	4.2	3.8	3.6	15.9	8.8	8.8	8.2	9.5	12.3	19.2	19.3
1999.....	9.7	4.7	3.8	3.6	15.9	8.9	8.8	8.0	9.5	13.3	19.1	18.5

White, both sexes

2004.....	15.8	3.2	3.9	4.1	28.5	17.8	15.3	15.0	14.1	16.2	24.9	27.3
2003.....	15.8	3.5	3.8	4.1	28.6	17.5	15.9	14.9	13.9	16.1	25.2	29.5
2002.....	16.2	2.6	3.7	3.9	30.3	18.0	15.9	14.8	14.1	17.1	26.3	28.8
2001.....	15.7	3.5	3.9	4.0	28.4	17.8	15.9	14.1	13.6	16.3	27.0	28.8
2000.....	15.7	3.9	4.0	4.3	28.7	17.1	15.4	14.0	14.3	16.4	26.1	31.1
1999.....	15.5	4.5	3.8	4.3	27.8	17.0	14.9	13.3	13.9	17.7	26.8	30.7

White, male

2004.....	21.9	3.3	4.0	4.8	39.4	25.8	22.2	21.7	19.5	21.1	34.9	46.3
2003.....	22.0	3.9	4.3	4.9	39.2	25.9	22.6	21.3	19.6	21.3	34.4	50.7
2002.....	22.4	2.9	4.2	4.6	41.9	26.6	22.3	21.2	19.6	22.4	35.5	52.4
2001.....	21.9	3.1	4.4	4.7	40.0	26.4	22.4	20.1	18.6	21.3	37.8	52.6
2000.....	21.6	4.2	4.5	5.0	39.6	25.1	21.8	20.0	19.2	21.1	35.6	59.2
1999.....	21.1	5.0	4.1	5.0	37.8	24.6	20.9	18.8	18.7	23.0	37.9	60.3

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)

White, female

2004.....	9.8	3.2	3.7	3.4	16.8	9.4	8.4	8.4	9.0	11.9	18.1	18.7
2003.....	9.8	3.0	3.2	3.2	17.2	8.6	9.0	8.5	8.6	11.7	19.0	20.1
2002.....	10.1	2.2	3.2	3.2	17.9	9.0	9.4	8.5	8.9	12.6	20.1	18.5
2001.....	9.8	3.8	3.4	3.2	16.0	8.8	9.3	8.1	8.8	12.0	19.8	18.9
2000.....	10.0	3.5	3.6	3.7	17.1	8.9	8.9	8.0	9.8	12.4	19.8	19.7
1999.....	10.0	3.9	3.5	3.6	17.1	9.1	8.8	7.9	9.3	13.3	19.6	18.8

Black, both sexes

2004.....	14.3	4.7	4.9	4.2	18.7	19.8	16.0	17.8	15.7	15.5	20.2	14.7
2003.....	14.2	4.2	4.9	3.9	18.8	18.4	17.1	16.5	17.0	17.0	21.0	20.7
2002.....	14.5	5.2	4.8	4.2	20.6	19.7	16.9	16.1	15.6	15.9	20.0	18.1
2001.....	14.8	*	5.0	4.9	21.5	18.9	17.2	16.9	16.4	17.3	19.7	16.6
2000.....	15.0	6.5	5.4	4.4	20.9	20.4	17.1	17.9	16.1	18.2	18.9	20.3
1999.....	15.1	7.1	6.4	5.0	21.1	18.6	17.4	17.1	16.5	17.9	24.1	25.5

Black, male

2004.....	21.2	*	5.2	4.6	26.4	31.8	24.7	28.0	24.7	24.9	35.9	28.8
2003.....	21.1	*	5.0	4.7	27.9	29.0	26.4	25.9	26.3	24.2	35.0	41.7
2002.....	21.5	*	6.3	4.9	29.6	31.7	25.3	24.7	25.2	25.5	36.1	45.0
2001.....	22.0	*	5.3	6.3	31.4	29.9	26.1	26.9	26.3	24.2	34.1	35.2
2000.....	22.5	6.7	5.8	5.4	30.2	32.6	27.2	27.1	27.1	29.4	34.5	43.8
1999.....	22.2	*	7.0	6.0	30.5	29.7	26.3	27.0	25.5	26.5	41.2	60.9

Black, female

2004.....	7.9	*	4.7	3.8	10.9	8.7	8.3	9.0	8.5	8.7	11.3	9.1
2003.....	8.0	*	4.8	3.1	9.6	8.7	8.9	8.4	9.6	11.9	13.1	12.6
2002.....	8.0	*	3.4	3.5	11.6	8.8	9.4	8.8	7.9	9.0	10.9	*
2001.....	8.2	*	4.6	3.3	11.7	8.9	9.2	8.3	8.3	12.5	11.6	9.7
2000.....	8.2	*	4.9	3.5	11.7	9.4	8.2	10.0	7.3	10.3	10.1	11.6
1999.....	8.7	8.7	5.8	3.9	11.8	8.6	9.6	8.7	9.2	11.9	14.5	12.3

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pedestrian involved in collision with motor vehicle (V02-V04)												
All races 3/, both sexes												
2004.....	1.1	*	1.0	0.5	0.9	0.9	1.2	1.4	1.4	1.5	2.5	2.7
2003.....	1.2	*	1.1	0.5	1.1	1.0	1.3	1.4	1.5	1.6	2.6	2.9
2002.....	1.3	*	1.0	0.6	1.1	1.0	1.4	1.4	1.4	2.0	2.6	3.3
2001.....	1.2	*	1.1	0.7	1.0	1.0	1.4	1.4	1.4	1.6	3.0	3.0
2000.....	1.2	*	1.2	0.6	0.9	1.0	1.3	1.3	1.5	1.6	2.8	3.6
1999.....	1.2	*	1.1	0.7	0.9	1.0	1.3	1.3	1.5	1.7	2.5	3.7
All races 3/, male												
2004.....	1.6	*	1.2	0.6	1.3	1.4	1.7	2.1	2.1	2.0	3.4	4.9
2003.....	1.7	*	1.4	0.7	1.5	1.5	1.9	2.1	2.2	2.3	3.3	5.7
2002.....	1.7	*	1.3	0.7	1.6	1.6	1.9	2.1	2.0	2.7	3.4	5.8
2001.....	1.8	*	1.4	0.9	1.5	1.5	2.0	2.0	2.0	2.1	4.1	5.9
2000.....	1.7	*	1.4	0.8	1.3	1.5	1.9	2.0	2.2	2.2	3.8	7.0
1999.....	1.7	*	1.2	0.8	1.4	1.5	1.9	2.0	2.2	2.3	3.7	6.5
All races 3/, female												
2004.....	0.7	*	0.8	0.4	0.4	0.5	0.6	0.7	0.9	1.2	1.8	1.7
2003.....	0.7	*	0.8	0.4	0.6	0.5	0.7	0.8	0.8	1.0	2.2	1.7
2002.....	0.8	*	0.6	0.5	0.5	0.5	0.8	0.7	0.9	1.4	2.0	2.2
2001.....	0.8	*	0.8	0.4	0.5	0.4	0.8	0.8	0.7	1.1	2.2	1.8
2000.....	0.8	*	0.9	0.5	0.5	0.4	0.7	0.7	0.8	1.1	2.1	2.3
1999.....	0.7	*	1.0	0.5	0.5	0.5	0.6	0.6	0.9	1.2	1.7	2.6
White, both sexes												
2004.....	1.1	*	1.0	0.5	0.9	0.9	1.1	1.2	1.3	1.5	2.4	2.6
2003.....	1.2	*	1.1	0.5	1.1	1.0	1.2	1.3	1.4	1.4	2.5	2.9
2002.....	1.2	*	0.9	0.5	1.0	1.0	1.2	1.3	1.3	1.8	2.4	3.2
2001.....	1.2	*	1.0	0.6	1.0	1.0	1.2	1.2	1.3	1.4	2.8	2.9
2000.....	1.1	*	1.0	0.6	0.9	0.9	1.1	1.2	1.4	1.5	2.7	3.4
1999.....	1.1	*	1.0	0.6	0.9	0.9	1.0	1.1	1.4	1.5	2.3	3.6
White, male												
2004.....	1.5	*	1.2	0.6	1.3	1.4	1.7	1.9	1.9	1.8	3.2	4.6
2003.....	1.6	*	1.4	0.6	1.5	1.5	1.7	1.9	2.2	2.0	3.0	5.4
2002.....	1.6	*	1.2	0.6	1.5	1.5	1.7	1.9	1.9	2.5	3.1	5.4
2001.....	1.7	*	1.4	0.8	1.5	1.5	1.8	1.8	1.8	1.9	3.9	5.5
2000.....	1.6	*	1.2	0.7	1.3	1.4	1.6	1.8	2.0	1.9	3.4	6.2
1999.....	1.5	*	1.1	0.7	1.4	1.3	1.5	1.7	2.0	2.0	3.4	6.0

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pedestrian involved in collision with motor vehicle (V02-V04)												
White, female												
2004.....	0.7	*	0.8	0.3	0.4	0.4	0.6	0.6	0.9	1.1	1.8	1.7
2003.....	0.7	*	0.8	0.3	0.6	0.4	0.6	0.7	0.7	0.9	2.1	1.8
2002.....	0.7	*	0.6	0.4	0.5	0.4	0.7	0.6	0.8	1.3	1.9	2.2
2001.....	0.7	*	0.7	0.3	0.5	0.4	0.6	0.7	0.7	1.0	2.0	1.9
2000.....	0.7	*	0.8	0.5	0.6	0.4	0.6	0.6	0.8	1.1	2.2	2.2
1999.....	0.7	*	1.0	0.5	0.5	0.4	0.5	0.6	0.8	1.1	1.7	2.6
Black, both sexes												
2004.....	1.4	*	1.1	0.7	0.8	1.1	1.5	2.6	2.3	1.6	2.5	*
2003.....	1.5	*	1.5	0.9	1.0	1.2	2.0	2.3	1.9	2.4	2.8	*
2002.....	1.6	*	1.3	0.8	1.1	1.4	2.2	2.5	2.0	2.4	3.0	*
2001.....	1.7	*	1.5	1.1	1.0	1.1	2.5	2.6	2.2	2.0	2.9	*
2000.....	1.7	*	1.9	1.0	0.8	1.2	2.4	2.5	2.5	2.2	2.7	*
1999.....	1.8	*	1.6	1.2	1.0	1.6	2.7	2.3	2.7	2.3	2.6	*
Black, male												
2004.....	2.1	*	*	0.9	1.2	1.7	2.2	4.3	4.1	2.8	*	*
2003.....	2.3	*	1.9	1.0	1.3	1.7	3.1	3.7	3.1	4.9	*	*
2002.....	2.4	*	1.7	0.9	1.6	2.1	3.3	4.1	3.5	3.9	6.3	*
2001.....	2.4	*	1.7	1.6	1.5	1.6	3.6	3.8	4.2	3.2	*	*
2000.....	2.5	*	2.3	1.2	1.3	1.9	3.7	3.9	4.2	4.2	6.1	*
1999.....	2.8	*	2.0	1.4	1.4	2.6	4.5	4.1	4.7	3.8	*	*
Black, female												
2004.....	0.8	*	*	*	*	*	1.0	1.1	*	*	*	*
2003.....	0.8	*	*	0.7	0.7	0.7	1.0	1.0	*	*	*	*
2002.....	0.9	*	*	0.7	0.7	0.7	1.3	1.0	*	*	*	*
2001.....	1.0	*	*	*	*	*	1.5	1.5	*	*	*	*
2000.....	0.8	*	*	0.7	*	*	1.3	1.2	*	*	*	*
1999.....	0.9	*	*	0.9	*	*	1.2	*	*	*	*	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Pedalcyclist involved in collision with motor vehicle (V12-V14,V19.0-V19.2,V19.4-V19.6)

All races 3/, both sexes

2004.....	0.2	*	*	0.3	0.2	0.1	0.3	0.3	0.2	0.3	0.2	*
2003.....	0.2	*	*	0.3	0.2	0.1	0.3	0.3	0.2	0.2	0.2	*
2002.....	0.2	*	*	0.3	0.2	0.2	0.2	0.3	0.2	0.2	0.2	*
2001.....	0.2	*	*	0.3	0.2	0.2	0.3	0.3	0.2	0.2	*	*
2000.....	0.2	*	*	0.3	0.2	0.1	0.2	0.3	0.2	0.2	0.2	*
1999.....	0.2	*	*	0.4	0.3	0.2	0.3	0.2	0.1	0.2	*	*

All races 3/, male

2004.....	0.4	*	*	0.4	0.3	0.2	0.5	0.5	0.4	0.6	0.5	*
2003.....	0.4	*	*	0.4	0.3	0.2	0.5	0.5	0.4	0.4	*	*
2002.....	0.4	*	*	0.4	0.3	0.3	0.4	0.5	0.3	0.3	*	*
2001.....	0.4	*	*	0.4	0.4	0.3	0.5	0.5	0.4	0.4	*	*
2000.....	0.4	*	*	0.6	0.3	0.2	0.4	0.5	0.4	0.4	0.4	*
1999.....	0.4	*	*	0.5	0.5	0.4	0.5	0.4	0.2	0.4	*	*

All races 3/, female

2004.....	0.1	*	*	0.1	*	*	*	*	*	*	*	*
2003.....	0.0	*	*	0.1	*	*	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2001.....	0.0	*	*	0.1	*	*	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*	*	*	*	*	*
1999.....	0.1	*	*	0.2	*	*	*	*	*	*	*	*

White, both sexes

2004.....	0.2	*	*	0.3	0.2	0.1	0.3	0.3	0.2	0.3	0.2	*
2003.....	0.2	*	*	0.3	0.2	0.1	0.2	0.3	0.2	0.2	*	*
2002.....	0.2	*	*	0.2	0.2	0.2	0.2	0.3	0.2	0.1	*	*
2001.....	0.2	*	*	0.2	0.2	0.2	0.3	0.3	0.2	0.2	*	*
2000.....	0.2	*	*	0.3	0.2	0.1	0.2	0.2	0.2	0.2	*	*
1999.....	0.2	*	*	0.3	0.3	0.2	0.3	0.2	0.1	0.2	*	*

White, male

2004.....	0.4	*	*	0.4	0.4	0.3	0.5	0.5	0.4	0.5	0.6	*
2003.....	0.3	*	*	0.4	0.3	0.2	0.4	0.4	0.4	0.4	*	*
2002.....	0.4	*	*	0.4	0.4	0.4	0.4	0.5	0.3	0.3	*	*
2001.....	0.4	*	*	0.4	0.4	0.4	0.5	0.5	0.4	0.3	*	*
2000.....	0.4	*	*	0.5	0.4	0.2	0.4	0.4	0.4	0.3	*	*
1999.....	0.4	*	*	0.5	0.5	0.4	0.5	0.4	0.2	0.4	*	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Motorcyclist involved in any accident except collision with railway train (V20-V24,V26-V34,V36-V39)

All races 3/, both sexes

2004.....	1.4	*	*	0.1	1.9	2.2	2.1	2.1	1.3	0.6	0.3	*
2003.....	1.3	*	*	0.1	1.7	2.1	2.0	1.9	1.2	0.5	0.2	*
2002.....	1.1	*	*	0.1	1.5	1.9	1.7	1.6	1.0	0.4	0.2	*
2001.....	1.1	*	*	0.1	1.6	2.0	1.7	1.4	0.7	0.3	0.2	*
2000.....	1.0	*	*	0.1	1.4	1.8	1.6	1.3	0.8	0.4	*	*
1999.....	0.8	*	*	0.1	1.3	1.5	1.3	1.0	0.7	0.3	*	*

All races 3/, male

2004.....	2.5	*	*	0.2	3.5	4.0	3.6	3.7	2.4	1.2	0.5	*
2003.....	2.3	*	*	0.2	3.1	3.7	3.5	3.3	2.2	1.1	0.4	*
2002.....	2.1	*	*	0.1	2.8	3.5	3.1	3.0	1.9	0.8	0.4	*
2001.....	2.0	*	*	0.2	2.8	3.7	2.9	2.5	1.4	0.7	0.4	*
2000.....	1.8	*	*	0.1	2.6	3.4	2.8	2.3	1.4	0.8	*	*
1999.....	1.6	*	*	0.1	2.3	2.9	2.4	1.8	1.4	0.6	*	*

All races 3/, female

2004.....	0.3	*	*	*	0.3	0.3	0.5	0.5	0.2	*	*	*
2003.....	0.3	*	*	*	0.3	0.4	0.5	0.5	0.2	*	*	*
2002.....	0.2	*	*	*	0.2	0.3	0.4	0.4	0.2	*	*	*
2001.....	0.2	*	*	*	0.2	0.3	0.4	0.3	*	*	*	*
2000.....	0.2	*	*	*	0.2	0.2	0.4	0.3	0.2	*	*	*
1999.....	0.2	*	*	*	0.2	0.2	0.3	0.3	*	*	*	*

White, both sexes

2004.....	1.5	*	*	0.2	2.1	2.2	2.2	2.4	1.4	0.6	0.3	*
2003.....	1.4	*	*	0.1	1.9	2.0	2.1	2.2	1.3	0.6	0.2	*
2002.....	1.2	*	*	0.1	1.6	1.9	1.9	1.9	1.1	0.4	0.2	*
2001.....	1.1	*	*	0.1	1.7	2.0	1.9	1.6	0.8	0.4	0.2	*
2000.....	1.0	*	*	0.1	1.5	1.7	1.7	1.4	0.9	0.4	*	*
1999.....	0.9	*	*	0.1	1.3	1.5	1.4	1.2	0.8	0.3	*	*

White, male

2004.....	2.7	*	*	0.3	3.8	3.9	3.8	4.1	2.6	1.3	0.5	*
2003.....	2.4	*	*	0.2	3.4	3.5	3.6	3.8	2.5	1.2	0.5	*
2002.....	2.2	*	*	*	3.0	3.4	3.3	3.4	2.1	0.9	*	*
2001.....	2.1	*	*	0.2	3.0	3.6	3.2	2.8	1.5	0.7	*	*
2000.....	1.9	*	*	0.1	2.8	3.1	3.0	2.6	1.5	0.8	*	*
1999.....	1.6	*	*	0.1	2.4	2.6	2.5	2.1	1.5	0.6	*	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Motor vehicle accident involving collision with railway train (V25,V35,V45,V55,V65,V75,V81.0-V81.1,V87.6,V88.6)

All races 3/, both sexes

2004.....	0.1	*	*	*	0.1	0.1	0.0	0.1	*	*	*	*
2003.....	0.1	*	*	*	0.1	0.1	0.1	0.1	0.1	*	0.2	*
2002.....	0.1	*	*	*	0.1	0.1	0.1	0.1	0.1	*	*	*
2001.....	0.1	*	*	*	0.1	0.1	0.1	0.1	*	*	*	*
2000.....	0.1	*	*	*	0.2	0.1	0.1	0.1	0.1	0.1	0.2	*
1999.....	0.1	*	*	*	0.2	0.1	0.1	0.1	0.1	*	0.2	*

All races 3/, male

2004.....	0.1	*	*	*	0.2	0.2	*	0.1	*	*	*	*
2003.....	0.1	*	*	*	0.2	0.1	0.1	0.1	0.2	*	*	*
2002.....	0.1	*	*	*	0.2	0.1	0.1	0.1	*	*	*	*
2001.....	0.1	*	*	*	0.2	0.1	0.2	0.1	*	*	*	*
2000.....	0.1	*	*	*	0.2	0.2	0.1	0.2	*	*	*	*
1999.....	0.1	*	*	*	0.2	0.1	0.1	0.2	*	*	*	*

All races 3/, female

2004.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2003.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2002.....	0.1	*	*	*	*	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*	*	*	*	*	*
2000.....	0.1	*	*	*	*	*	*	*	*	*	*	*
1999.....	0.1	*	*	*	0.1	*	*	*	*	*	*	*

White, both sexes

2004.....	0.1	*	*	*	0.1	0.1	*	0.1	*	*	*	*
2003.....	0.1	*	*	*	0.1	0.1	0.1	0.1	0.1	*	*	*
2002.....	0.1	*	*	*	0.2	0.1	0.1	0.1	0.1	*	*	*
2001.....	0.1	*	*	*	0.1	0.1	0.1	0.1	*	*	*	*
2000.....	0.1	*	*	*	0.2	0.1	0.1	0.1	*	*	*	*
1999.....	0.1	*	*	*	0.2	0.1	0.1	0.1	0.1	*	0.2	*

White, male

2004.....	0.1	*	*	*	0.2	0.2	*	*	*	*	*	*
2003.....	0.1	*	*	*	0.2	*	*	*	0.2	*	*	*
2002.....	0.1	*	*	*	0.2	*	0.1	0.1	*	*	*	*
2001.....	0.1	*	*	*	0.2	0.2	0.2	*	*	*	*	*
2000.....	0.2	*	*	*	0.3	0.2	0.2	0.2	*	*	*	*
1999.....	0.1	*	*	*	0.2	0.1	0.1	0.2	*	*	*	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other motor vehicle accident involving collision with railway train (V45,V55,V65,V75,V81.0-V81.1,V87.6,V88.6)

All races 3/, both sexes

2004.....	0.1	*	*	*	0.1	0.1	0.0	0.1	*	*	*	*
2003.....	0.1	*	*	*	0.1	0.1	0.1	0.1	0.1	*	0.2	*
2002.....	0.1	*	*	*	0.1	0.1	0.1	0.1	0.1	*	*	*
2001.....	0.1	*	*	*	0.1	0.1	0.1	0.1	*	*	*	*
2000.....	0.1	*	*	*	0.2	0.1	0.1	0.1	0.1	0.1	0.2	*
1999.....	0.1	*	*	*	0.2	0.1	0.1	0.1	0.1	*	0.2	*

All races 3/, male

2004.....	0.1	*	*	*	0.2	0.2	*	0.1	*	*	*	*
2003.....	0.1	*	*	*	0.2	0.1	0.1	*	0.2	*	*	*
2002.....	0.1	*	*	*	0.2	0.1	0.1	0.1	*	*	*	*
2001.....	0.1	*	*	*	0.2	0.1	0.2	0.1	*	*	*	*
2000.....	0.1	*	*	*	0.2	0.2	0.1	0.1	*	*	*	*
1999.....	0.1	*	*	*	0.2	0.1	0.1	0.1	*	*	*	*

All races 3/, female

2004.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2003.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2002.....	0.1	*	*	*	*	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*	*	*	*	*	*
2000.....	0.1	*	*	*	*	*	*	*	*	*	*	*
1999.....	0.1	*	*	*	0.1	*	*	*	*	*	*	*

White, both sexes

2004.....	0.1	*	*	*	0.1	0.1	*	0.1	*	*	*	*
2003.....	0.1	*	*	*	0.1	0.1	0.1	0.1	0.1	*	*	*
2002.....	0.1	*	*	*	0.1	0.1	0.1	0.1	0.1	*	*	*
2001.....	0.1	*	*	*	0.1	0.1	0.1	0.1	*	*	*	*
2000.....	0.1	*	*	*	0.2	0.1	0.1	0.1	*	*	*	*
1999.....	0.1	*	*	*	0.2	0.1	0.1	0.1	0.1	*	0.2	*

White, male

2004.....	0.1	*	*	*	0.2	0.1	*	*	*	*	*	*
2003.....	0.1	*	*	*	0.2	*	*	*	0.2	*	*	*
2002.....	0.1	*	*	*	0.2	*	0.1	0.1	*	*	*	*
2001.....	0.1	*	*	*	0.2	0.2	0.2	*	*	*	*	*
2000.....	0.2	*	*	*	0.2	0.2	0.2	0.2	*	*	*	*
1999.....	0.1	*	*	*	0.2	0.1	0.1	0.1	*	*	*	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Occupant of motor vehicle involved in collision with other (non-motorized) road vehicle, streetcar, animal or pedestrian (V40-V41,V46-V47,V50-V51,V56-V57,V60-V61,V66-V67, V70-V71,V76-V77)

All races 3/, both sexes

2004.....	1.6	*	0.2	0.2	4.1	2.1	1.5	1.2	1.1	1.2	1.6	1.2
2003.....	1.7	*	0.2	0.3	4.2	2.3	1.6	1.4	1.1	1.3	1.6	1.9
2002.....	1.8	*	0.2	0.3	4.5	2.3	1.7	1.4	1.3	1.2	1.6	1.5
2001.....	1.5	*	0.1	0.2	3.8	2.0	1.4	1.2	0.9	1.1	1.6	1.7
2000.....	1.4	*	0.2	0.2	3.5	1.7	1.4	1.0	1.0	1.2	1.4	1.7
1999.....	1.4	*	0.1	0.2	3.3	1.8	1.3	1.0	1.0	1.2	1.5	1.2

All races 3/, male

2004.....	2.4	*	*	0.3	6.0	3.1	2.3	1.8	1.7	1.8	2.5	1.9
2003.....	2.6	*	*	0.4	6.3	3.6	2.4	2.1	1.7	1.9	2.3	3.7
2002.....	2.6	*	*	0.3	6.7	3.6	2.5	1.9	2.0	1.7	2.2	3.2
2001.....	2.2	*	*	0.2	5.6	3.0	2.2	1.8	1.5	1.5	2.5	3.1
2000.....	2.0	*	*	0.2	5.2	2.7	2.0	1.5	1.4	1.6	2.1	3.0
1999.....	2.0	*	*	0.3	4.9	2.8	1.9	1.6	1.4	1.8	2.3	2.5

All races 3/, female

2004.....	0.8	*	*	0.2	2.1	1.0	0.7	0.6	0.6	0.8	1.0	0.9
2003.....	0.9	*	0.3	0.2	2.0	1.1	0.8	0.8	0.5	0.9	1.1	1.1
2002.....	0.9	*	*	0.3	2.3	0.9	0.9	0.8	0.6	0.8	1.2	0.8
2001.....	0.8	*	*	0.2	1.9	0.9	0.7	0.6	0.4	0.7	1.1	1.2
2000.....	0.8	*	*	0.2	1.6	0.8	0.8	0.6	0.6	0.8	0.9	1.2
1999.....	0.7	*	*	0.2	1.7	0.7	0.7	0.5	0.6	0.8	1.0	0.7

White, both sexes

2004.....	1.6	*	*	0.2	4.5	2.1	1.5	1.2	1.1	1.2	1.7	1.2
2003.....	1.8	*	0.2	0.3	4.6	2.4	1.7	1.4	1.1	1.3	1.6	2.0
2002.....	1.8	*	0.2	0.2	4.8	2.3	1.8	1.3	1.2	1.2	1.7	1.6
2001.....	1.5	*	*	0.2	3.9	2.0	1.4	1.2	0.9	1.1	1.6	1.9
2000.....	1.4	*	*	0.2	3.6	1.7	1.4	1.0	0.9	1.1	1.3	1.8
1999.....	1.3	*	*	0.2	3.4	1.8	1.3	1.0	1.0	1.2	1.5	1.2

White, male

2004.....	2.4	*	*	0.2	6.5	3.1	2.3	1.8	1.6	1.7	2.6	2.1
2003.....	2.6	*	*	0.4	6.8	3.6	2.5	2.0	1.6	1.8	2.4	3.9
2002.....	2.6	*	*	0.3	7.0	3.6	2.6	1.9	1.8	1.6	2.2	3.3
2001.....	2.2	*	*	0.3	5.7	3.0	2.1	1.8	1.5	1.5	2.5	3.2
2000.....	2.0	*	*	0.2	5.3	2.6	2.0	1.4	1.3	1.5	1.9	3.2
1999.....	2.0	*	*	0.3	5.0	2.8	1.9	1.5	1.4	1.7	2.3	2.5

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Occupant of motor vehicle involved in collision with other (non-motorized) road vehicle, streetcar, animal or pedestrian (V40-V41,V46-V47,V50-V51,V56-V57,V60-V61,V66-V67, V70-V71,V76-V77)

White, female

2004.....	0.9	*	*	0.2	2.2	1.0	0.7	0.6	0.6	0.8	1.0	0.9
2003.....	0.9	*	*	0.2	2.2	1.2	0.9	0.8	0.6	0.8	1.2	1.2
2002.....	1.0	*	*	0.2	2.5	1.0	1.0	0.8	0.6	0.8	1.3	0.9
2001.....	0.8	*	*	0.2	1.9	0.9	0.7	0.6	0.4	0.8	1.1	1.3
2000.....	0.8	*	*	0.3	1.7	0.8	0.9	0.6	0.6	0.8	0.9	1.2
1999.....	0.7	*	*	0.2	1.8	0.8	0.6	0.5	0.6	0.8	0.9	*

Black, both sexes

2004.....	1.7	*	*	0.3	3.0	2.7	1.7	1.5	2.0	1.4	*	*
2003.....	1.6	*	*	*	3.1	2.4	1.3	2.0	1.5	2.0	*	*
2002.....	1.9	*	*	0.5	3.8	2.6	1.8	1.8	2.3	1.7	*	*
2001.....	1.8	*	*	*	4.1	2.4	1.9	1.6	1.1	*	2.1	*
2000.....	1.6	*	*	*	3.1	2.0	1.7	1.7	1.4	2.0	*	*
1999.....	1.6	*	*	*	3.4	2.3	1.7	1.5	1.6	1.4	2.3	*

Black, male

2004.....	2.5	*	*	*	4.1	4.5	2.7	2.2	3.4	*	*	*
2003.....	2.6	*	*	*	4.9	4.2	2.2	3.1	2.8	*	*	*
2002.....	3.0	*	*	*	5.9	4.5	2.7	2.9	4.0	3.6	*	*
2001.....	2.8	*	*	*	6.3	4.0	3.0	2.7	1.9	*	*	*
2000.....	2.5	*	*	*	4.8	3.7	2.7	2.5	2.6	3.4	*	*
1999.....	2.6	*	*	*	5.1	4.0	2.7	2.4	2.5	*	*	*

Black, female

2004.....	0.9	*	*	*	1.8	1.0	0.9	0.9	*	*	*	*
2003.....	0.7	*	*	*	1.2	0.8	*	1.0	*	*	*	*
2002.....	0.9	*	*	*	1.7	0.8	1.1	0.8	*	*	*	*
2001.....	0.8	*	*	*	1.9	1.1	0.9	*	*	*	*	*
2000.....	0.7	*	*	*	1.4	*	0.8	1.0	*	*	*	*
1999.....	0.8	*	*	*	1.7	0.7	0.9	*	*	*	*	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Occupant of car, pickup truck or van involved in collision with other motor vehicle (V42-V44,V49,V52-V54,V59)

All races 3/, both sexes

2004.....	3.6	1.5	0.8	0.9	6.0	3.6	3.0	3.1	3.5	5.0	8.3	9.6
2003.....	3.7	1.5	0.9	0.8	6.1	3.7	3.2	3.2	3.5	4.9	8.7	10.2
2002.....	3.9	1.1	1.0	0.9	6.8	3.8	3.4	3.3	3.8	5.2	9.2	9.9
2001.....	3.7	1.6	1.0	0.9	6.1	3.7	3.3	3.1	3.5	5.1	8.4	10.1
2000.....	3.7	1.5	0.9	0.9	6.1	3.7	3.1	3.2	3.8	5.0	9.0	10.1
1999.....	3.7	1.6	0.9	0.9	5.8	3.7	3.1	3.2	3.8	5.3	8.8	9.2

All races 3/, male

2004.....	4.4	1.9	0.8	1.0	7.4	4.7	3.8	3.8	4.0	5.8	10.9	15.9
2003.....	4.5	1.4	0.9	0.8	7.4	4.9	4.1	4.0	4.2	5.4	11.2	16.6
2002.....	4.8	1.5	1.0	0.9	8.7	5.1	4.2	4.1	4.7	6.0	12.2	16.9
2001.....	4.5	1.6	0.9	0.9	7.8	5.0	4.2	3.8	4.1	6.1	10.9	18.1
2000.....	4.5	1.5	0.9	0.9	7.5	4.8	3.9	4.0	4.5	5.8	11.6	18.2
1999.....	4.4	1.7	1.0	0.9	7.3	4.9	3.8	3.9	4.3	6.2	11.6	17.8

All races 3/, female

2004.....	2.9	1.1	0.9	0.9	4.4	2.4	2.1	2.5	3.0	4.3	6.5	6.7
2003.....	2.9	1.5	0.9	0.8	4.7	2.5	2.3	2.4	2.9	4.4	7.0	7.4
2002.....	3.0	*	0.9	0.8	4.9	2.4	2.6	2.5	3.0	4.4	7.2	6.8
2001.....	2.9	1.6	1.1	0.9	4.3	2.3	2.5	2.4	3.0	4.3	6.7	6.8
2000.....	2.9	1.5	0.9	0.8	4.5	2.6	2.3	2.4	3.2	4.3	7.3	6.8
1999.....	2.9	1.4	0.7	1.0	4.3	2.5	2.4	2.5	3.3	4.6	7.0	5.7

White, both sexes

2004.....	3.8	1.5	0.8	1.0	6.5	3.6	3.0	3.1	3.5	5.1	8.5	10.1
2003.....	3.9	1.4	0.9	0.8	6.6	3.9	3.2	3.3	3.4	4.9	9.0	10.6
2002.....	4.1	1.0	0.9	0.9	7.4	3.8	3.4	3.3	3.8	5.3	9.6	10.4
2001.....	3.8	1.5	1.0	0.9	6.5	3.8	3.3	3.0	3.6	5.1	8.7	10.6
2000.....	3.8	1.5	0.9	0.9	6.5	3.7	3.1	3.1	3.8	5.1	9.3	10.6
1999.....	3.8	1.5	0.8	1.0	6.2	3.8	3.0	3.2	3.8	5.4	9.1	9.4

White, male

2004.....	4.5	1.9	0.8	1.0	8.0	4.7	3.9	3.7	4.0	5.9	11.0	16.6
2003.....	4.6	1.5	0.9	0.8	8.0	5.2	4.1	4.1	4.0	5.5	11.4	17.3
2002.....	5.0	1.4	1.0	1.0	9.3	5.1	4.2	4.0	4.6	6.0	12.5	17.9
2001.....	4.7	1.6	0.9	0.9	8.3	5.1	4.1	3.6	4.2	6.1	11.3	18.9
2000.....	4.6	1.5	0.9	0.9	7.9	4.7	3.9	3.9	4.3	5.9	11.9	19.3
1999.....	4.5	1.8	0.9	0.9	7.7	5.0	3.6	3.8	4.2	6.1	11.9	18.1

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Occupant of car, pickup truck or van involved in collision with other motor vehicle (V42-V44,V49,V52-V54,V59)												
White, female												
2004.....	3.0	*	0.8	0.9	4.9	2.5	2.1	2.5	3.1	4.5	6.8	7.2
2003.....	3.1	1.4	0.9	0.8	5.2	2.6	2.3	2.5	2.8	4.4	7.4	7.7
2002.....	3.2	*	0.9	0.8	5.3	2.4	2.6	2.5	3.1	4.7	7.7	7.1
2001.....	3.0	1.5	1.1	0.9	4.6	2.4	2.5	2.5	3.0	4.3	7.0	7.2
2000.....	3.1	1.5	0.9	0.9	4.9	2.6	2.3	2.3	3.3	4.5	7.6	7.1
1999.....	3.1	*	0.7	1.0	4.7	2.6	2.5	2.5	3.4	4.7	7.3	5.9
Black, both sexes												
2004.....	3.1	*	0.9	0.8	3.9	4.2	2.9	3.5	3.7	4.4	6.7	5.6
2003.....	3.2	*	0.9	0.8	4.3	3.6	3.6	3.4	4.7	5.0	6.5	6.4
2002.....	3.3	*	1.2	0.8	4.8	4.8	3.6	3.5	3.7	4.0	5.7	*
2001.....	3.3	*	1.1	0.9	4.7	3.9	3.7	3.9	3.8	5.9	5.1	*
2000.....	3.3	*	1.0	0.7	4.6	4.2	3.4	4.2	4.0	4.4	5.6	*
1999.....	3.4	*	1.1	1.0	4.3	4.1	4.0	3.6	4.0	5.0	6.2	7.7
Black, male												
2004.....	4.0	*	*	0.6	5.0	6.3	3.7	4.5	5.0	5.7	12.0	*
2003.....	4.1	*	*	0.6	5.8	5.0	4.5	4.5	6.3	6.1	11.0	*
2002.....	4.4	*	1.6	0.6	6.3	6.7	4.7	4.7	5.3	5.6	9.8	*
2001.....	4.4	*	*	0.9	6.2	5.5	5.2	5.9	4.8	7.1	8.5	*
2000.....	4.4	*	*	0.9	6.1	5.9	4.7	5.7	6.7	5.6	8.5	*
1999.....	4.5	*	*	1.1	5.9	5.6	5.6	4.9	5.4	7.6	10.8	*
Black, female												
2004.....	2.2	*	*	1.0	2.8	2.3	2.2	2.6	2.7	3.5	3.8	*
2003.....	2.4	*	*	0.9	2.7	2.4	2.7	2.4	3.5	4.3	4.0	*
2002.....	2.4	*	*	0.9	3.3	3.0	2.6	2.5	2.5	2.9	3.4	*
2001.....	2.3	*	*	0.8	3.1	2.4	2.4	2.2	3.1	5.0	*	*
2000.....	2.2	*	*	*	3.0	2.7	2.3	2.9	1.8	3.5	3.9	*
1999.....	2.3	*	*	0.8	2.8	2.8	2.5	2.5	2.9	3.2	3.6	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Occupant of heavy transport vehicle or bus involved in collision with other motor vehicle (V62-V64,V69,V72-V74,V79)

All races 3/, both sexes

2004.....	0.1	*	*	*	*	0.1	0.1	0.2	0.1	*	*	*
2003.....	0.1	*	*	*	*	0.1	0.1	0.1	0.2	0.1	*	*
2002.....	0.1	*	*	*	*	0.1	0.1	0.1	0.1	*	*	*
2001.....	0.1	*	*	*	*	0.1	0.1	0.1	0.1	*	*	*
2000.....	0.1	*	*	*	*	0.1	0.1	0.1	*	*	*	*
1999.....	0.1	*	*	*	*	0.1	0.1	0.1	0.1	*	*	*

All races 3/, male

2004.....	0.1	*	*	*	*	0.1	0.2	0.3	0.2	*	*	*
2003.....	0.1	*	*	*	*	0.1	0.2	0.2	0.3	*	*	*
2002.....	0.1	*	*	*	*	0.2	0.2	0.3	0.2	*	*	*
2001.....	0.1	*	*	*	*	0.1	0.1	0.2	0.2	*	*	*
2000.....	0.1	*	*	*	*	0.1	0.2	0.2	*	*	*	*
1999.....	0.1	*	*	*	*	0.1	0.2	0.2	0.2	*	*	*

All races 3/, female

2004.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2003.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	*	*	*	*	*	*

White, both sexes

2004.....	0.1	*	*	*	*	0.1	0.1	0.2	0.1	*	*	*
2003.....	0.1	*	*	*	*	0.1	0.1	0.1	0.2	*	*	*
2002.....	0.1	*	*	*	*	0.1	0.1	0.1	0.1	*	*	*
2001.....	0.1	*	*	*	*	0.1	0.1	0.1	0.1	*	*	*
2000.....	0.1	*	*	*	*	0.1	0.1	0.1	*	*	*	*
1999.....	0.1	*	*	*	*	0.1	0.1	0.1	0.1	*	*	*

White, male

2004.....	0.1	*	*	*	*	0.1	0.2	0.3	0.2	*	*	*
2003.....	0.1	*	*	*	*	0.1	0.2	0.2	0.3	*	*	*
2002.....	0.1	*	*	*	*	0.2	0.2	0.3	*	*	*	*
2001.....	0.1	*	*	*	*	*	0.1	0.2	0.2	*	*	*
2000.....	0.1	*	*	*	*	*	0.2	0.2	*	*	*	*
1999.....	0.1	*	*	*	*	*	0.2	0.2	0.2	*	*	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Occupant of motor vehicle involved in noncollision accident (V48,V58,V68,V78)

All races 3/, both sexes

2004.....	1.4	*	0.3	0.4	3.1	1.7	1.5	1.2	1.1	1.0	0.9	1.0
2003.....	1.5	*	0.3	0.4	3.3	1.9	1.6	1.4	1.2	1.0	1.0	1.3
2002.....	1.5	*	0.3	0.4	3.5	2.0	1.6	1.3	1.2	1.0	1.0	0.9
2001.....	1.4	*	0.2	0.4	3.0	1.8	1.5	1.3	1.1	1.0	1.0	0.7
2000.....	1.3	0.6	0.2	0.4	3.1	1.8	1.4	1.2	0.9	0.8	0.9	1.0
1999.....	1.3	*	0.3	0.3	3.1	1.7	1.5	1.1	1.1	0.9	0.8	1.0

All races 3/, male

2004.....	1.9	*	0.3	0.4	4.2	2.5	2.2	1.8	1.6	1.4	1.2	1.7
2003.....	2.1	*	0.3	0.4	4.5	2.8	2.2	1.9	1.6	1.5	1.6	2.0
2002.....	2.2	*	0.4	0.4	4.8	3.0	2.3	2.0	1.6	1.4	1.5	1.8
2001.....	2.0	*	*	0.4	4.2	2.7	2.1	1.9	1.6	1.4	1.5	*
2000.....	1.9	*	*	0.4	4.3	2.8	2.0	1.8	1.3	1.2	1.4	2.2
1999.....	1.9	*	0.3	0.3	4.2	2.6	2.1	1.6	1.6	1.3	1.2	2.0

All races 3/, female

2004.....	0.8	*	0.3	0.4	1.9	0.9	0.9	0.7	0.6	0.6	0.6	0.7
2003.....	0.9	*	0.4	0.4	2.1	1.0	1.0	0.8	0.8	0.6	0.6	1.0
2002.....	0.9	*	*	0.4	2.2	1.0	1.0	0.7	0.7	0.6	0.7	*
2001.....	0.8	*	*	0.4	1.8	0.9	0.9	0.6	0.6	0.6	0.8	*
2000.....	0.7	*	*	0.3	1.7	0.8	0.8	0.6	0.6	0.5	0.6	*
1999.....	0.8	*	0.3	0.3	1.9	0.9	0.8	0.6	0.6	0.7	0.5	*

White, both sexes

2004.....	1.4	*	0.2	0.4	3.3	1.8	1.6	1.2	1.1	0.9	0.8	1.1
2003.....	1.5	*	0.3	0.4	3.6	2.0	1.7	1.4	1.1	1.0	1.0	1.3
2002.....	1.6	*	0.3	0.4	3.8	2.2	1.6	1.3	1.2	1.0	1.0	0.8
2001.....	1.4	*	0.2	0.4	3.3	1.8	1.6	1.3	1.0	1.0	1.1	0.7
2000.....	1.4	*	0.2	0.3	3.3	1.9	1.4	1.2	0.9	0.8	0.9	1.0
1999.....	1.3	*	0.3	0.3	3.3	1.8	1.4	1.1	1.0	0.9	0.8	1.0

White, male

2004.....	2.0	*	*	0.4	4.5	2.6	2.2	1.7	1.7	1.3	1.2	1.9
2003.....	2.1	*	*	0.4	4.9	3.0	2.2	1.9	1.5	1.5	1.6	2.1
2002.....	2.2	*	0.3	0.4	5.2	3.1	2.2	1.9	1.6	1.5	1.6	1.7
2001.....	2.0	*	*	0.4	4.6	2.7	2.2	1.9	1.4	1.5	1.6	*
2000.....	2.0	*	*	0.4	4.7	2.9	2.1	1.8	1.2	1.2	1.4	2.2
1999.....	1.9	*	*	0.3	4.5	2.6	2.1	1.6	1.5	1.2	1.2	1.9

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Occupant of motor vehicle involved in noncollision accident (V48,V58,V68,V78)

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
White, female												
2004.....	0.8	*	*	0.4	2.0	0.9	0.9	0.7	0.6	0.6	0.6	0.7
2003.....	0.9	*	0.3	0.4	2.3	0.9	1.1	0.8	0.8	0.6	0.6	0.9
2002.....	0.9	*	*	0.4	2.3	1.2	1.0	0.7	0.7	0.6	0.7	*
2001.....	0.8	*	*	0.4	1.9	0.9	0.9	0.7	0.6	0.6	0.8	*
2000.....	0.7	*	*	0.3	1.9	0.9	0.8	0.6	0.5	0.5	0.6	*
1999.....	0.8	*	*	0.3	2.0	0.9	0.8	0.6	0.6	0.7	0.5	*
Black, both sexes												
2004.....	1.1	*	*	*	1.8	1.6	1.4	1.4	1.0	1.3	*	*
2003.....	1.2	*	*	0.4	1.7	1.7	1.4	1.2	1.4	*	*	*
2002.....	1.2	*	*	0.4	2.0	1.4	1.5	1.2	1.1	*	*	*
2001.....	1.1	*	*	0.4	1.6	1.8	1.1	1.2	1.7	*	*	*
2000.....	1.1	*	*	0.4	1.9	1.4	1.3	1.0	1.6	*	*	*
1999.....	1.2	*	*	*	1.9	1.5	1.6	1.3	1.0	*	*	*
Black, male												
2004.....	1.7	*	*	*	2.5	2.2	2.1	2.4	*	*	*	*
2003.....	1.7	*	*	*	2.4	2.2	2.1	2.0	2.4	*	*	*
2002.....	1.8	*	*	*	2.7	2.6	2.2	2.3	1.9	*	*	*
2001.....	1.6	*	*	*	2.1	2.9	1.7	2.2	3.1	*	*	*
2000.....	1.6	*	*	*	2.6	2.3	2.1	1.7	2.5	*	*	*
1999.....	1.7	*	*	*	2.6	2.4	2.2	2.1	*	*	*	*
Black, female												
2004.....	0.6	*	*	*	1.1	1.0	0.8	*	*	*	*	*
2003.....	0.7	*	*	*	0.9	1.3	0.7	*	*	*	*	*
2002.....	0.6	*	*	*	1.3	*	0.9	*	*	*	*	*
2001.....	0.6	*	*	*	1.2	0.7	*	*	*	*	*	*
2000.....	0.6	*	*	*	1.2	*	*	*	*	*	*	*
1999.....	0.7	*	*	*	1.3	0.7	1.0	*	*	*	*	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Occupant of special-use motor vehicle involved in any accident (V83-V86)

All races 3/, both sexes

2004.....	0.4	*	*	0.4	0.6	0.4	0.4	0.3	0.3	0.3	0.3	*
2003.....	0.4	*	*	0.4	0.6	0.4	0.4	0.4	0.3	0.4	0.4	*
2002.....	0.3	*	*	0.3	0.5	0.4	0.3	0.3	0.3	0.4	0.5	*
2001.....	0.4	*	*	0.3	0.6	0.4	0.3	0.2	0.4	0.4	0.6	*
2000.....	0.4	*	*	0.3	0.6	0.4	0.3	0.3	0.5	0.4	0.4	*
1999.....	0.4	*	*	0.3	0.4	0.4	0.3	0.3	0.4	0.6	0.6	0.7

All races 3/, male

2004.....	0.7	*	*	0.6	1.0	0.7	0.7	0.6	0.6	0.6	0.7	*
2003.....	0.7	*	*	0.5	1.0	0.7	0.7	0.7	0.7	0.7	1.0	*
2002.....	0.6	*	*	0.5	0.9	0.6	0.6	0.5	0.5	0.8	1.0	*
2001.....	0.7	*	*	0.5	1.0	0.7	0.6	0.4	0.7	0.9	1.4	*
2000.....	0.7	*	*	0.4	1.0	0.7	0.6	0.5	0.9	0.9	1.1	*
1999.....	0.7	*	*	0.5	0.8	0.7	0.6	0.6	0.8	1.3	1.4	2.2

All races 3/, female

2004.....	0.1	*	*	0.2	0.2	0.1	*	*	*	*	*	*
2003.....	0.1	*	*	0.2	0.1	*	*	0.1	*	*	*	*
2002.....	0.1	*	*	0.2	0.1	*	*	*	*	*	*	*
2001.....	0.1	*	*	0.2	0.1	*	*	*	*	*	*	*
2000.....	0.1	*	*	0.2	0.2	*	*	*	*	*	*	*
1999.....	0.1	*	*	*	*	*	*	*	*	*	*	*

White, both sexes

2004.....	0.4	*	*	0.5	0.7	0.5	0.4	0.4	0.3	0.3	0.3	*
2003.....	0.4	*	*	0.4	0.6	0.5	0.4	0.4	0.4	0.4	0.5	*
2002.....	0.4	*	*	0.4	0.6	0.4	0.4	0.3	0.3	0.4	0.5	*
2001.....	0.4	*	*	0.4	0.7	0.5	0.4	0.2	0.4	0.5	0.6	*
2000.....	0.4	*	*	0.4	0.7	0.4	0.4	0.3	0.5	0.5	0.5	*
1999.....	0.4	*	*	0.3	0.5	0.5	0.3	0.4	0.4	0.6	0.6	0.7

White, male

2004.....	0.7	*	*	0.7	1.2	0.9	0.8	0.6	0.6	0.7	0.7	*
2003.....	0.8	*	*	0.6	1.1	0.9	0.8	0.7	0.7	0.8	1.1	*
2002.....	0.7	*	*	0.6	1.1	0.8	0.7	0.6	0.6	0.8	1.1	*
2001.....	0.7	*	*	0.5	1.1	0.8	0.7	0.4	0.8	1.0	1.5	*
2000.....	0.8	*	*	0.5	1.2	0.8	0.6	0.6	1.0	1.0	1.2	*
1999.....	0.8	*	*	0.5	0.9	0.8	0.6	0.7	0.8	1.4	1.5	2.3

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other and unspecified motor vehicle accidents (V09.0,V09.2,V80.3-V80.5,V82.0-V82.1,V87.0-V87.5,V87.7-V87.8,V88.0-V88.5,V88.7-V88.8,V89.0,V89.2)

All races 3/, both sexes

2004.....	5.5	1.2	1.6	1.2	9.4	6.3	5.1	5.2	4.9	6.1	10.4	10.8
2003.....	5.3	1.3	1.3	1.2	9.3	5.4	5.2	4.8	5.0	6.1	9.9	11.9
2002.....	5.5	1.1	1.3	1.1	9.9	6.0	5.2	5.0	4.8	6.6	10.3	11.4
2001.....	5.8	1.3	1.5	1.2	10.4	6.3	5.6	5.1	5.3	6.5	11.3	11.7
2000.....	6.0	1.8	1.6	1.4	11.1	6.5	5.8	5.5	5.7	6.7	10.7	13.1
1999.....	6.0	2.3	1.6	1.5	10.8	6.4	5.7	5.2	5.3	7.4	11.8	13.9

All races 3/, male

2004.....	7.3	1.2	1.6	1.2	12.6	8.7	7.0	7.2	6.6	7.9	14.5	17.9
2003.....	7.0	1.6	1.5	1.3	12.5	7.8	7.0	6.6	6.6	7.7	13.7	20.3
2002.....	7.3	1.1	1.4	1.3	13.3	8.5	7.0	6.8	6.5	8.7	13.9	21.7
2001.....	7.7	1.1	1.6	1.4	14.4	8.9	7.6	7.1	6.9	8.1	15.8	21.2
2000.....	8.0	1.9	1.8	1.5	14.8	9.2	8.0	7.4	7.5	8.5	14.7	24.7
1999.....	7.9	2.1	1.8	1.6	14.4	8.8	7.6	7.0	7.1	9.3	16.7	27.5

All races 3/, female

2004.....	3.7	1.3	1.5	1.2	6.1	3.8	3.2	3.2	3.4	4.7	7.6	7.7
2003.....	3.6	*	1.1	1.1	5.9	2.9	3.4	3.0	3.5	4.7	7.4	8.1
2002.....	3.7	1.1	1.3	1.0	6.3	3.4	3.4	3.3	3.3	4.8	8.0	6.9
2001.....	3.9	1.5	1.3	1.1	6.2	3.6	3.7	3.2	3.8	5.1	8.3	7.7
2000.....	4.1	1.7	1.4	1.4	7.1	3.8	3.5	3.6	4.0	5.3	8.1	8.4
1999.....	4.2	2.5	1.5	1.3	7.0	3.9	3.8	3.5	3.7	5.9	8.7	8.5

White, both sexes

2004.....	5.6	1.1	1.5	1.1	10.0	6.4	5.1	5.0	4.9	6.0	10.6	11.5
2003.....	5.3	1.2	1.2	1.2	9.7	5.5	5.1	4.5	4.8	6.1	10.0	12.1
2002.....	5.6	0.8	1.3	1.1	10.6	6.0	5.2	4.9	4.8	6.6	10.5	11.8
2001.....	5.9	1.1	1.4	1.1	11.0	6.4	5.6	5.1	5.1	6.4	11.6	12.2
2000.....	6.2	1.5	1.6	1.4	11.8	6.4	5.8	5.3	5.7	6.6	10.9	13.5
1999.....	6.2	2.2	1.4	1.5	11.5	6.5	5.8	5.0	5.1	7.5	11.9	14.3

White, male

2004.....	7.3	*	1.5	1.1	13.4	8.7	6.8	6.8	6.4	7.6	14.8	19.0
2003.....	7.1	1.6	1.4	1.4	12.9	7.9	6.9	6.2	6.3	7.8	13.8	20.4
2002.....	7.4	*	1.3	1.2	14.1	8.5	6.9	6.6	6.3	8.7	13.9	21.9
2001.....	7.8	*	1.5	1.2	15.0	9.1	7.4	7.0	6.6	8.0	16.0	22.3
2000.....	8.2	1.6	1.8	1.5	15.8	9.1	7.9	7.1	7.3	8.1	14.9	25.8
1999.....	8.0	2.2	1.4	1.6	15.2	8.9	7.7	6.6	6.8	9.4	16.6	28.3

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other and unspecified motor vehicle accidents (V09.0,V09.2,V80.3-V80.5,V82.0-V82.1,V87.0-V87.5,V87.7-V87.8,V88.0-V88.5,V88.7-V88.8,V89.0,V89.2)

White, female

2004.....	3.9	1.4	1.5	1.1	6.4	4.0	3.2	3.2	3.4	4.7	7.7	8.0
2003.....	3.6	*	1.0	1.1	6.3	2.9	3.3	2.9	3.3	4.7	7.5	8.4
2002.....	3.9	*	1.3	1.0	6.8	3.5	3.4	3.3	3.3	4.8	8.3	7.4
2001.....	4.0	1.3	1.3	1.1	6.6	3.6	3.8	3.2	3.8	5.1	8.6	7.9
2000.....	4.2	1.4	1.4	1.4	7.5	3.7	3.6	3.5	4.1	5.3	8.3	8.5
1999.....	4.3	2.3	1.4	1.3	7.6	4.0	3.8	3.3	3.6	5.9	8.9	8.7

Black, both sexes

2004.....	5.4	*	2.1	1.7	7.4	6.7	5.9	7.2	5.7	6.2	8.1	*
2003.....	5.2	*	1.6	1.2	7.3	5.8	6.3	6.3	6.5	6.3	8.0	10.8
2002.....	5.1	*	1.5	1.3	7.5	6.6	5.7	6.0	5.6	6.2	8.9	7.1
2001.....	5.6	*	1.7	1.7	8.3	6.6	6.4	6.2	6.7	6.9	9.1	7.1
2000.....	5.9	3.6	1.9	1.4	8.8	8.1	6.3	7.4	5.7	8.0	7.7	8.8
1999.....	5.8	*	2.6	1.7	8.6	6.4	5.9	7.4	6.4	7.7	12.0	10.2

Black, male

2004.....	7.8	*	2.3	1.8	10.3	10.0	8.9	11.1	8.7	10.5	13.1	*
2003.....	7.4	*	1.6	1.2	10.8	8.7	8.9	9.9	9.7	8.7	12.1	20.9
2002.....	7.3	*	1.8	1.6	10.4	10.0	8.1	8.4	8.9	9.7	15.5	21.4
2001.....	8.0	*	1.8	2.2	11.9	9.7	9.4	9.7	10.6	9.1	15.9	*
2000.....	8.5	*	2.0	1.5	12.1	11.6	10.0	10.9	9.3	12.9	13.3	*
1999.....	8.2	*	3.1	2.0	12.1	9.6	8.2	11.2	9.7	9.6	21.5	23.4

Black, female

2004.....	3.2	*	1.9	1.5	4.5	3.6	3.3	3.8	3.4	3.2	5.3	*
2003.....	3.2	*	*	1.1	3.8	3.1	3.9	3.3	4.0	4.6	5.7	*
2002.....	3.1	*	*	0.9	4.5	3.6	3.5	4.0	3.0	3.7	5.2	*
2001.....	3.4	*	*	1.3	4.6	3.8	3.8	3.2	3.6	5.3	5.3	*
2000.....	3.6	*	1.8	1.3	5.5	4.9	3.1	4.4	2.8	4.5	4.6	*
1999.....	3.7	*	2.1	1.4	5.1	3.5	4.0	4.1	3.7	6.3	6.7	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other and unspecified land transport accidents (V01,V06,V09.1,V09.3,V09.9,V10-V11,V16-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.7-V80.9,V87.9,V88.9,V89.1,V89.3,V89.9)

All races 3/, both sexes

2004.....	0.3	*	*	0.1	0.3	0.2	0.3	0.3	0.3	0.4	0.4	*
2003.....	0.3	*	*	0.1	0.5	0.3	0.3	0.3	0.4	0.4	0.5	0.5
2002.....	0.2	*	*	0.1	0.2	0.1	0.2	0.3	0.3	0.3	0.4	*
2001.....	0.3	*	0.2	0.1	0.3	0.2	0.4	0.4	0.4	0.5	0.6	0.6
2000.....	0.4	*	0.2	0.2	0.4	0.3	0.4	0.4	0.4	0.5	0.6	0.7
1999.....	0.6	*	0.3	0.3	0.7	0.6	0.6	0.6	0.6	0.7	1.2	1.2

All races 3/, male

2004.....	0.4	*	*	0.1	0.4	0.3	0.4	0.4	0.5	0.5	0.6	*
2003.....	0.5	*	*	0.1	0.7	0.5	0.4	0.6	0.6	0.6	0.8	*
2002.....	0.3	*	*	0.2	0.3	0.2	0.4	0.5	0.4	0.5	0.9	*
2001.....	0.5	*	*	0.2	0.5	0.3	0.6	0.6	0.6	0.8	0.9	*
2000.....	0.5	*	*	0.2	0.6	0.5	0.6	0.5	0.6	0.7	1.0	*
1999.....	0.8	*	0.3	0.3	1.0	0.8	0.9	1.0	1.0	1.1	1.7	2.7

All races 3/, female

2004.....	0.1	*	*	*	0.2	0.1	0.1	0.1	0.2	0.3	*	*
2003.....	0.2	*	*	*	0.3	0.1	0.2	0.1	0.2	*	0.3	*
2002.....	0.1	*	*	*	*	0.1	0.1	*	*	*	*	*
2001.....	0.2	*	*	0.1	0.1	*	0.2	0.2	0.2	0.3	0.5	*
2000.....	0.2	*	*	0.2	0.2	0.2	0.2	0.3	0.2	0.2	0.4	*
1999.....	0.3	*	0.3	0.2	0.4	0.3	0.3	0.3	0.3	0.4	0.9	*

White, both sexes

2004.....	0.3	*	*	0.1	0.3	0.2	0.3	0.3	0.3	0.4	0.4	*
2003.....	0.3	*	*	0.1	0.5	0.3	0.3	0.3	0.4	0.4	0.5	0.5
2002.....	0.2	*	*	0.1	0.2	0.1	0.2	0.3	0.3	0.3	0.4	*
2001.....	0.3	*	0.2	0.1	0.3	0.2	0.3	0.4	0.4	0.5	0.6	0.5
2000.....	0.4	*	*	0.2	0.4	0.3	0.4	0.4	0.4	0.4	0.6	0.7
1999.....	0.6	*	0.3	0.3	0.7	0.6	0.6	0.6	0.7	0.7	1.2	1.2

White, male

2004.....	0.4	*	*	0.1	0.4	0.3	0.4	0.4	0.5	0.5	0.6	*
2003.....	0.5	*	*	*	0.8	0.5	0.4	0.5	0.6	0.6	0.8	*
2002.....	0.3	*	*	0.2	0.3	0.2	0.3	0.5	0.4	0.5	0.8	*
2001.....	0.5	*	*	0.2	0.5	0.3	0.5	0.6	0.6	0.8	0.9	*
2000.....	0.5	*	*	0.3	0.6	0.5	0.6	0.5	0.6	0.7	1.0	*
1999.....	0.8	*	*	0.3	1.0	0.8	0.9	1.0	1.0	1.0	1.6	2.6

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Water transport accidents (V90-V94)												
All races 3/, both sexes												
2004.....	0.2	*	*	0.1	0.2	0.3	0.2	0.3	0.2	0.2	0.2	*
2003.....	0.2	*	*	0.1	0.3	0.2	0.3	0.3	0.2	0.2	0.2	*
2002.....	0.2	*	*	0.1	0.2	0.3	0.3	0.2	0.2	0.3	0.2	*
2001.....	0.2	*	*	0.1	0.3	0.2	0.2	0.3	0.3	0.2	*	*
2000.....	0.2	*	*	0.1	0.3	0.3	0.3	0.3	0.2	0.3	*	*
1999.....	0.2	*	*	0.1	0.3	0.3	0.3	0.3	0.3	0.2	*	*
All races 3/, male												
2004.....	0.3	*	*	*	0.4	0.4	0.4	0.5	0.4	0.4	0.4	*
2003.....	0.4	*	*	0.1	0.4	0.4	0.5	0.5	0.4	0.3	0.4	*
2002.....	0.4	*	*	*	0.4	0.5	0.6	0.5	0.5	0.7	0.4	*
2001.....	0.4	*	*	0.1	0.5	0.4	0.4	0.5	0.5	0.5	*	*
2000.....	0.4	*	*	0.1	0.5	0.5	0.6	0.5	0.5	0.5	*	*
1999.....	0.4	*	*	0.2	0.5	0.5	0.5	0.6	0.6	0.5	*	*
All races 3/, female												
2004.....	0.1	*	*	*	*	0.1	*	*	*	*	*	*
2003.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*	*	*	*	*	*
1999.....	0.1	*	*	*	0.1	0.1	0.1	*	*	*	*	*
White, both sexes												
2004.....	0.2	*	*	0.1	0.2	0.3	0.2	0.3	0.2	0.2	*	*
2003.....	0.2	*	*	0.1	0.3	0.2	0.3	0.3	0.2	0.1	*	*
2002.....	0.2	*	*	0.1	0.3	0.3	0.3	0.2	0.2	0.3	0.2	*
2001.....	0.2	*	*	0.1	0.3	0.3	0.3	0.3	0.3	0.2	*	*
2000.....	0.2	*	*	0.1	0.3	0.3	0.3	0.3	0.2	0.3	*	*
1999.....	0.2	*	*	0.1	0.3	0.3	0.3	0.3	0.3	0.2	*	*
White, male												
2004.....	0.4	*	*	*	0.4	0.5	0.4	0.5	0.4	0.5	*	*
2003.....	0.4	*	*	*	0.4	0.4	0.5	0.5	0.4	0.3	*	*
2002.....	0.4	*	*	*	0.4	0.5	0.6	0.5	0.5	0.6	*	*
2001.....	0.4	*	*	0.1	0.5	0.5	0.4	0.5	0.5	0.5	*	*
2000.....	0.4	*	*	*	0.5	0.5	0.6	0.5	0.5	0.6	*	*
1999.....	0.4	*	*	0.2	0.5	0.6	0.5	0.6	0.5	0.5	*	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Air and space transport accidents (V95-V97)

All races 3/, both sexes

2004.....	0.2	*	*	*	0.1	0.3	0.3	0.4	0.4	0.4	*	*
2003.....	0.3	*	*	0.1	0.1	0.2	0.3	0.5	0.6	0.4	0.2	*
2002.....	0.2	*	*	*	0.1	0.2	0.3	0.4	0.5	0.4	0.2	*
2001.....	0.3	*	*	*	0.2	0.4	0.5	0.5	0.5	0.5	0.2	*
2000.....	0.3	*	*	*	0.2	0.3	0.3	0.5	0.6	0.4	0.2	*
1999.....	0.3	*	*	0.0	0.1	0.3	0.3	0.4	0.5	0.4	0.2	*

All races 3/, male

2004.....	0.4	*	*	*	0.1	0.4	0.5	0.7	0.8	0.7	*	*
2003.....	0.4	*	*	*	0.2	0.3	0.5	0.8	1.1	0.7	0.4	*
2002.....	0.4	*	*	*	0.2	0.3	0.5	0.7	0.9	0.8	*	*
2001.....	0.5	*	*	*	0.2	0.6	0.7	0.8	0.8	0.8	0.5	*
2000.....	0.5	*	*	*	0.3	0.5	0.6	0.8	1.0	0.7	0.5	*
1999.....	0.4	*	*	*	0.2	0.5	0.5	0.7	1.0	0.7	*	*

All races 3/, female

2004.....	0.1	*	*	*	0.1	*	*	0.1	*	*	*	*
2003.....	0.1	*	*	*	*	*	0.1	0.1	*	*	*	*
2002.....	0.1	*	*	*	*	*	*	0.1	0.1	*	*	*
2001.....	0.2	*	*	*	0.1	0.2	0.2	0.3	0.2	*	*	*
2000.....	0.1	*	*	*	*	0.1	*	0.1	0.2	*	*	*
1999.....	0.1	*	*	*	*	0.1	0.1	0.1	*	*	*	*

White, both sexes

2004.....	0.3	*	*	*	0.2	0.3	0.3	0.5	0.5	0.4	*	*
2003.....	0.3	*	*	0.1	0.1	0.2	0.4	0.6	0.7	0.4	0.2	*
2002.....	0.3	*	*	*	0.2	0.2	0.3	0.5	0.6	0.4	0.2	*
2001.....	0.4	*	*	*	0.2	0.5	0.5	0.6	0.6	0.5	0.2	*
2000.....	0.3	*	*	*	0.2	0.4	0.4	0.5	0.6	0.4	0.2	*
1999.....	0.3	*	*	*	0.1	0.4	0.4	0.5	0.6	0.4	0.2	*

White, male

2004.....	0.5	*	*	*	0.2	0.5	0.6	0.9	0.9	0.8	*	*
2003.....	0.5	*	*	*	0.2	0.4	0.6	1.0	1.2	0.8	0.5	*
2002.....	0.5	*	*	*	0.2	0.4	0.6	0.8	1.0	0.9	*	*
2001.....	0.6	*	*	*	0.2	0.7	0.9	0.9	0.9	0.9	0.5	*
2000.....	0.6	*	*	*	0.3	0.6	0.7	0.9	1.1	0.7	0.5	*
1999.....	0.5	*	*	*	0.2	0.6	0.6	0.8	1.1	0.8	*	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other and unspecified transport accidents and their sequelae (V98-V99,Y85)

All races 3/, both sexes

2004.....	0.2	*	*	*	0.1	0.2	0.3	0.4	0.3	0.3	0.2	*
2003.....	0.2	*	*	*	0.1	0.2	0.3	0.4	0.4	0.4	0.3	*
2002.....	0.2	*	*	*	0.1	0.2	0.3	0.4	0.3	0.2	0.4	*
2001.....	0.2	*	*	*	0.1	0.2	0.3	0.3	0.3	0.3	0.4	*
2000.....	0.2	*	*	*	0.1	0.2	0.3	0.2	0.3	0.3	0.3	*
1999.....	0.2	*	*	*	0.1	0.2	0.3	0.3	0.3	0.2	0.3	*

All races 3/, male

2004.....	0.3	*	*	*	0.2	0.4	0.5	0.6	0.6	0.5	*	*
2003.....	0.3	*	*	*	0.2	0.3	0.5	0.5	0.6	0.6	0.4	*
2002.....	0.3	*	*	*	0.2	0.3	0.4	0.6	0.5	0.4	0.7	*
2001.....	0.3	*	*	*	0.2	0.3	0.5	0.6	0.4	0.4	0.6	*
2000.....	0.3	*	*	*	0.1	0.3	0.4	0.4	0.4	0.5	0.5	*
1999.....	0.3	*	*	*	0.1	0.3	0.4	0.4	0.4	0.4	*	*

All races 3/, female

2004.....	0.1	*	*	*	*	*	0.2	0.2	*	*	*	*
2003.....	0.1	*	*	*	*	0.1	0.1	0.2	0.2	*	*	*
2002.....	0.1	*	*	*	*	*	0.2	0.2	0.2	*	*	*
2001.....	0.1	*	*	*	*	0.1	0.2	0.1	*	*	0.3	*
2000.....	0.1	*	*	*	*	*	0.1	0.1	*	*	*	*
1999.....	0.1	*	*	*	*	0.1	0.1	0.1	0.2	*	*	*

White, both sexes

2004.....	0.2	*	*	*	0.2	0.2	0.3	0.4	0.3	0.3	0.2	*
2003.....	0.2	*	*	*	0.1	0.2	0.3	0.3	0.4	0.4	0.3	*
2002.....	0.2	*	*	*	0.1	0.2	0.3	0.4	0.3	0.2	0.4	*
2001.....	0.2	*	*	*	0.1	0.2	0.3	0.3	0.2	0.3	0.4	*
2000.....	0.2	*	*	*	0.1	0.2	0.3	0.2	0.2	0.3	0.3	*
1999.....	0.2	*	*	*	0.1	0.2	0.3	0.3	0.2	0.2	0.2	*

White, male

2004.....	0.4	*	*	*	0.2	0.3	0.5	0.6	0.5	0.5	*	*
2003.....	0.3	*	*	*	0.1	0.3	0.4	0.5	0.6	0.6	*	*
2002.....	0.3	*	*	*	0.2	0.3	0.4	0.6	0.5	0.4	0.6	*
2001.....	0.3	*	*	*	0.2	0.3	0.5	0.5	0.4	0.4	0.6	*
2000.....	0.3	*	*	*	0.1	0.3	0.4	0.4	0.4	0.5	0.5	*
1999.....	0.3	*	*	*	0.1	0.3	0.4	0.4	0.3	0.4	*	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Falls (W00-W19)												
All races 3/, both sexes												
2004.....	6.4	0.6	0.3	0.1	0.6	0.8	1.5	2.8	4.8	12.2	43.8	143.3
2003.....	5.9	*	0.3	0.1	0.6	0.7	1.4	2.6	4.4	11.2	40.8	135.9
2002.....	5.6	*	0.2	0.1	0.6	0.8	1.5	2.5	4.1	10.8	38.3	130.4
2001.....	5.3	0.6	0.2	0.2	0.6	0.9	1.4	2.6	4.0	10.0	35.3	121.5
2000.....	4.7	*	0.2	0.1	0.6	0.8	1.3	2.3	3.9	9.0	31.1	112.6
1999.....	4.7	*	0.4	0.1	0.6	0.9	1.4	2.3	3.7	9.0	31.5	110.2
All races 3/, male												
2004.....	6.8	*	0.4	0.1	1.0	1.3	2.4	4.5	7.1	16.5	54.8	174.4
2003.....	6.2	*	0.5	0.1	0.9	1.2	2.4	4.0	6.2	14.5	51.6	165.1
2002.....	6.0	*	0.4	0.1	1.0	1.3	2.3	3.9	6.0	14.5	48.1	158.9
2001.....	5.8	*	0.3	0.2	1.1	1.5	2.4	4.3	5.9	13.1	45.7	155.7
2000.....	5.2	*	0.3	0.1	1.0	1.4	2.2	3.7	5.8	11.9	39.9	145.4
1999.....	5.2	*	0.5	0.2	1.1	1.5	2.4	3.7	5.7	12.4	40.5	142.5
All races 3/, female												
2004.....	6.0	*	*	*	0.2	0.2	0.6	1.3	2.7	8.6	36.4	129.2
2003.....	5.6	*	*	*	0.2	0.2	0.5	1.2	2.7	8.4	33.5	122.9
2002.....	5.3	*	*	*	0.2	0.2	0.6	1.2	2.3	7.7	31.8	118.1
2001.....	4.8	*	*	0.1	0.1	0.2	0.5	1.0	2.2	7.4	28.5	107.2
2000.....	4.3	*	*	*	0.2	0.2	0.5	0.9	2.1	6.6	25.3	99.2
1999.....	4.3	*	*	*	0.2	0.2	0.5	0.9	1.9	6.3	25.7	97.2
White, both sexes												
2004.....	7.3	*	0.3	0.1	0.7	0.9	1.6	2.9	4.8	12.7	46.4	152.6
2003.....	6.7	*	0.3	0.1	0.6	0.8	1.5	2.5	4.4	11.5	42.8	144.4
2002.....	6.4	*	0.2	0.1	0.7	0.9	1.5	2.6	4.1	11.1	40.4	138.0
2001.....	6.0	*	0.2	0.2	0.8	0.9	1.5	2.6	4.0	10.4	37.0	128.0
2000.....	5.3	*	0.2	0.1	0.7	0.8	1.4	2.3	3.9	9.2	32.6	119.1
1999.....	5.2	*	0.4	0.1	0.7	0.9	1.5	2.2	3.5	9.1	32.9	115.9
White, male												
2004.....	7.6	*	0.4	*	1.1	1.4	2.5	4.5	6.9	17.1	57.6	183.7
2003.....	6.9	*	0.5	0.2	1.0	1.3	2.4	3.9	6.2	14.7	53.8	175.1
2002.....	6.6	*	*	0.1	1.1	1.5	2.4	3.9	6.0	14.8	50.2	166.4
2001.....	6.4	*	0.3	0.2	1.3	1.6	2.5	4.2	5.8	13.5	47.4	162.3
2000.....	5.7	*	*	0.2	1.2	1.5	2.2	3.7	5.7	12.1	41.4	152.8
1999.....	5.6	*	0.5	0.2	1.2	1.5	2.4	3.5	5.4	12.3	41.9	149.1

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Falls (W00-W19)												
White, female												
2004.....	7.0	*	*	*	0.2	0.2	0.6	1.3	2.8	8.9	38.7	138.6
2003.....	6.5	*	*	*	0.2	0.3	0.5	1.1	2.8	8.8	35.4	130.8
2002.....	6.2	*	*	*	0.2	0.2	0.7	1.2	2.4	8.0	33.9	125.6
2001.....	5.5	*	*	*	0.1	0.2	0.4	1.1	2.2	7.7	30.2	113.6
2000.....	5.0	*	*	*	0.2	0.2	0.5	0.9	2.1	6.7	26.8	105.4
1999.....	4.9	*	*	*	0.2	0.2	0.6	0.8	1.8	6.4	27.1	102.6
Black, both sexes												
2004.....	2.6	*	*	*	*	0.7	1.4	3.0	5.6	10.1	20.3	46.5
2003.....	2.4	*	*	*	*	0.5	1.3	3.3	4.8	8.9	19.4	44.1
2002.....	2.2	*	*	*	*	0.4	1.3	2.9	4.2	8.2	16.1	49.3
2001.....	2.3	*	*	*	*	0.5	1.6	2.9	4.4	8.7	18.5	44.5
2000.....	2.1	*	*	*	*	0.4	1.3	2.7	4.7	8.3	14.9	38.8
1999.....	2.4	*	*	*	0.4	0.8	1.3	3.4	5.5	9.2	17.1	45.3
Black, male												
2004.....	3.4	*	*	*	*	1.1	2.1	4.9	10.1	14.4	29.2	71.5
2003.....	3.0	*	*	*	*	0.8	2.2	5.2	7.1	12.5	24.8	56.3
2002.....	3.0	*	*	*	*	0.8	2.2	4.8	6.7	11.8	24.4	77.2
2001.....	3.1	*	*	*	*	0.8	2.6	5.4	7.0	12.6	28.5	68.1
2000.....	2.8	*	*	*	*	*	2.1	4.7	8.2	11.4	21.2	61.1
1999.....	3.3	*	*	*	*	1.3	2.5	5.8	8.8	14.7	24.3	58.6
Black, female												
2004.....	1.8	*	*	*	*	*	0.8	1.3	2.0	7.0	15.3	36.6
2003.....	1.9	*	*	*	*	*	*	1.6	2.9	6.3	16.3	39.3
2002.....	1.5	*	*	*	*	*	*	1.2	2.2	5.6	11.4	38.6
2001.....	1.6	*	*	*	*	*	*	*	2.3	5.9	12.9	35.8
2000.....	1.4	*	*	*	*	*	*	1.0	2.0	6.1	11.4	30.5
1999.....	1.7	*	*	*	*	*	*	1.3	2.9	5.3	13.1	40.3

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Fall on same level (W00-W09,W18)												
All races 3/, both sexes												
2004.....	2.1	*	*	*	0.1	0.2	0.3	0.7	1.3	3.8	15.1	53.6
2003.....	1.9	*	*	*	0.1	0.1	0.3	0.6	1.2	3.3	14.2	50.3
2002.....	1.8	*	*	*	0.1	0.1	0.3	0.6	1.1	3.1	12.6	50.8
2001.....	1.4	*	*	0.1	0.1	0.1	0.2	0.6	1.0	2.6	9.9	37.3
2000.....	1.2	*	*	*	0.1	0.1	0.2	0.4	0.8	2.0	8.0	33.0
1999.....	0.8	*	*	*	0.1	0.1	0.1	0.3	0.4	1.5	5.8	21.4
All races 3/, male												
2004.....	2.1	*	*	*	0.2	0.3	0.4	1.0	1.8	4.8	18.6	64.1
2003.....	1.9	*	*	*	*	0.2	0.4	0.8	1.6	4.0	18.0	60.1
2002.....	1.8	*	*	*	0.2	0.2	0.4	0.9	1.4	4.2	15.2	60.8
2001.....	1.4	*	*	*	0.2	0.2	0.3	0.8	1.3	3.0	12.7	45.3
2000.....	1.1	*	*	*	0.1	0.1	0.3	0.6	1.0	2.5	9.8	39.6
1999.....	0.8	*	*	*	0.1	*	0.2	0.5	0.5	2.0	7.3	26.6
All races 3/, female												
2004.....	2.1	*	*	*	*	*	0.2	0.3	0.9	2.9	12.7	48.9
2003.....	2.0	*	*	*	*	*	0.2	0.4	0.8	2.7	11.6	45.9
2002.....	1.9	*	*	*	*	*	0.1	0.3	0.8	2.2	10.9	46.4
2001.....	1.4	*	*	*	*	*	0.1	0.3	0.6	2.2	8.0	33.9
2000.....	1.2	*	*	*	*	*	*	0.2	0.6	1.6	6.9	30.4
1999.....	0.8	*	*	*	*	*	*	0.2	0.3	1.1	4.9	19.4
White, both sexes												
2004.....	2.4	*	*	*	0.1	0.2	0.3	0.7	1.3	3.9	15.9	57.5
2003.....	2.2	*	*	*	0.1	0.1	0.3	0.6	1.2	3.5	15.0	53.5
2002.....	2.1	*	*	*	0.1	0.1	0.3	0.6	1.1	3.3	13.3	53.7
2001.....	1.6	*	*	*	0.1	0.1	0.2	0.6	1.0	2.7	10.3	39.3
2000.....	1.3	*	*	*	0.1	0.1	0.2	0.4	0.8	2.1	8.5	35.0
1999.....	0.9	*	*	*	0.1	0.1	0.1	0.3	0.4	1.5	6.0	22.7
White, male												
2004.....	2.3	*	*	*	0.2	0.3	0.4	1.0	1.7	5.0	19.3	68.2
2003.....	2.1	*	*	*	*	0.2	0.4	0.8	1.5	4.1	18.9	63.5
2002.....	2.0	*	*	*	0.2	0.2	0.4	0.9	1.5	4.3	15.7	63.3
2001.....	1.6	*	*	*	0.2	0.2	0.3	0.8	1.3	3.2	13.1	47.7
2000.....	1.2	*	*	*	*	0.1	0.3	0.6	0.9	2.6	10.2	41.9
1999.....	0.9	*	*	*	0.1	*	0.2	0.4	0.5	2.0	7.4	28.0

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Fall from one level to another (W10-W17)												
All races 3/, both sexes												
2004.....	1.1	*	0.1	*	0.3	0.4	0.7	1.1	1.6	2.8	5.8	10.6
2003.....	1.1	*	0.2	0.0	0.4	0.5	0.7	1.0	1.4	2.7	5.9	11.4
2002.....	1.2	*	0.1	*	0.4	0.5	0.8	0.9	1.4	2.8	6.1	10.9
2001.....	1.2	*	0.1	0.1	0.5	0.6	0.7	1.1	1.4	2.7	6.1	11.3
2000.....	1.0	*	0.2	*	0.4	0.5	0.7	1.0	1.3	2.4	5.1	8.8
1999.....	1.1	*	0.2	0.1	0.4	0.6	0.7	0.9	1.5	2.6	5.8	10.4
All races 3/, male												
2004.....	1.7	*	*	*	0.6	0.8	1.3	1.9	2.7	4.4	8.5	16.4
2003.....	1.6	*	0.3	*	0.7	0.8	1.2	1.8	2.2	4.3	8.9	18.1
2002.....	1.7	*	*	*	0.7	0.9	1.3	1.6	2.4	4.3	9.7	17.4
2001.....	1.7	*	*	0.1	0.8	1.0	1.3	2.0	2.4	4.2	9.5	18.2
2000.....	1.6	*	*	*	0.8	1.0	1.3	1.8	2.2	3.8	8.4	16.4
1999.....	1.7	*	0.3	0.1	0.7	1.0	1.3	1.6	2.6	4.1	9.0	17.8
All races 3/, female												
2004.....	0.6	*	*	*	*	*	0.2	0.3	0.6	1.3	4.0	7.9
2003.....	0.7	*	*	*	0.1	0.1	0.1	0.3	0.6	1.4	3.8	8.5
2002.....	0.6	*	*	*	0.1	0.1	0.2	0.3	0.5	1.5	3.7	8.1
2001.....	0.6	*	*	*	*	0.1	0.1	0.3	0.5	1.4	3.9	8.3
2000.....	0.5	*	*	*	0.1	*	0.2	0.3	0.5	1.2	2.9	5.7
1999.....	0.6	*	*	*	0.1	0.1	0.2	0.2	0.6	1.4	3.7	7.4
White, both sexes												
2004.....	1.3	*	*	*	0.4	0.5	0.8	1.1	1.6	2.9	6.1	11.2
2003.....	1.2	*	0.2	*	0.5	0.5	0.7	1.0	1.5	2.8	6.1	12.1
2002.....	1.3	*	*	*	0.5	0.6	0.8	1.0	1.5	3.0	6.4	11.2
2001.....	1.3	*	*	0.1	0.5	0.6	0.8	1.1	1.4	2.8	6.4	11.7
2000.....	1.1	*	*	*	0.5	0.6	0.8	1.0	1.4	2.5	5.4	9.2
1999.....	1.2	*	0.2	0.1	0.5	0.6	0.8	0.9	1.6	2.7	6.1	10.9
White, male												
2004.....	1.8	*	*	*	0.6	0.8	1.3	1.9	2.7	4.7	9.1	17.5
2003.....	1.8	*	*	*	0.8	0.9	1.2	1.7	2.3	4.2	9.3	19.4
2002.....	1.9	*	*	*	0.8	1.1	1.4	1.7	2.5	4.6	10.2	18.1
2001.....	1.9	*	*	*	1.0	1.1	1.4	1.9	2.3	4.3	9.9	19.0
2000.....	1.7	*	*	*	0.9	1.1	1.3	1.8	2.3	3.9	8.8	17.0
1999.....	1.8	*	0.3	*	0.8	1.1	1.4	1.6	2.6	4.1	9.3	19.2

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Unspecified fall (W19)												
All races 3/, both sexes												
2004.....	3.2	*	*	*	0.1	0.2	0.5	1.1	1.9	5.7	22.9	79.1
2003.....	2.9	*	*	*	0.1	0.1	0.5	0.9	1.8	5.1	20.8	74.2
2002.....	2.7	*	*	*	0.1	0.1	0.5	1.0	1.6	4.8	19.6	68.8
2001.....	2.7	*	*	*	0.1	0.2	0.5	1.0	1.6	4.8	19.3	73.0
2000.....	2.5	*	*	*	0.1	0.1	0.4	0.9	1.8	4.6	17.9	70.7
1999.....	2.8	*	*	*	0.1	0.2	0.6	1.0	1.8	4.9	19.9	78.4
All races 3/, male												
2004.....	3.1	*	*	*	0.2	0.3	0.8	1.6	2.6	7.3	27.7	93.9
2003.....	2.7	*	*	*	0.1	0.2	0.7	1.4	2.3	6.2	24.7	86.9
2002.....	2.5	*	*	*	0.1	0.2	0.7	1.4	2.2	5.9	23.2	80.7
2001.....	2.6	*	*	*	0.1	0.3	0.8	1.5	2.2	5.9	23.4	92.1
2000.....	2.5	*	*	*	0.2	0.2	0.7	1.4	2.6	5.6	21.7	89.4
1999.....	2.8	*	*	*	0.2	0.3	0.9	1.6	2.6	6.3	24.2	98.1
All races 3/, female												
2004.....	3.2	*	*	*	*	*	0.2	0.7	1.2	4.3	19.7	72.4
2003.....	3.0	*	*	*	*	*	0.2	0.5	1.3	4.2	18.1	68.5
2002.....	2.8	*	*	*	*	*	0.3	0.6	1.0	3.9	17.2	63.6
2001.....	2.7	*	*	*	*	*	0.2	0.4	1.1	3.9	16.6	64.9
2000.....	2.6	*	*	*	*	*	0.2	0.5	1.0	3.8	15.5	63.1
1999.....	2.8	*	*	*	*	0.1	0.3	0.5	1.0	3.8	17.1	70.4
White, both sexes												
2004.....	3.6	*	*	*	0.2	0.2	0.5	1.1	1.9	5.8	24.4	83.9
2003.....	3.2	*	*	*	0.1	0.2	0.5	0.9	1.8	5.3	21.6	78.8
2002.....	3.0	*	*	*	0.1	0.1	0.5	1.0	1.5	4.9	20.8	73.0
2001.....	3.1	*	*	*	0.1	0.2	0.5	1.0	1.6	4.9	20.3	77.0
2000.....	2.9	*	*	*	0.1	0.1	0.4	0.9	1.7	4.7	18.7	74.9
1999.....	3.1	*	*	*	0.2	0.2	0.6	0.9	1.6	4.9	20.9	82.3
White, male												
2004.....	3.5	*	*	*	0.3	0.3	0.8	1.5	2.5	7.4	29.3	98.0
2003.....	3.0	*	*	*	0.1	0.2	0.8	1.3	2.3	6.3	25.6	92.2
2002.....	2.8	*	*	*	*	0.2	0.6	1.3	2.0	5.9	24.4	85.1
2001.....	2.9	*	*	*	0.2	0.3	0.8	1.5	2.2	6.0	24.3	95.7
2000.....	2.7	*	*	*	0.2	0.2	0.6	1.4	2.5	5.7	22.4	94.0
1999.....	3.0	*	*	*	0.3	0.3	0.8	1.5	2.3	6.2	25.1	101.9

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Unspecified fall (W19)												
White, female												
2004.....	3.8	*	*	*	*	*	0.2	0.7	1.3	4.5	21.1	77.6
2003.....	3.4	*	*	*	*	*	0.2	0.5	1.3	4.4	19.0	72.8
2002.....	3.2	*	*	*	*	*	0.3	0.6	1.0	4.0	18.3	67.8
2001.....	3.2	*	*	*	*	*	0.2	0.5	1.1	4.0	17.6	69.2
2000.....	3.0	*	*	*	*	*	0.2	0.5	1.0	3.8	16.2	67.2
1999.....	3.3	*	*	*	*	*	0.3	0.4	0.9	3.8	18.1	74.4
Black, both sexes												
2004.....	1.2	*	*	*	*	*	0.5	1.3	2.4	5.5	9.6	30.2
2003.....	1.2	*	*	*	*	*	0.5	1.4	2.2	5.0	12.0	28.6
2002.....	1.1	*	*	*	*	*	0.5	1.5	2.3	4.5	8.7	25.2
2001.....	1.1	*	*	*	*	*	0.7	1.1	2.1	4.7	10.8	26.4
2000.....	1.2	*	*	*	*	*	0.7	1.2	2.4	4.9	10.0	25.7
1999.....	1.4	*	*	*	*	*	0.6	1.8	3.4	5.5	11.0	33.5
Black, male												
2004.....	1.6	*	*	*	*	*	*	2.0	4.5	8.0	13.9	48.7
2003.....	1.4	*	*	*	*	*	*	2.3	3.2	6.3	13.8	36.5
2002.....	1.4	*	*	*	*	*	0.9	2.7	3.9	6.4	10.6	38.6
2001.....	1.4	*	*	*	*	*	1.0	2.0	3.1	6.2	15.6	46.5
2000.....	1.4	*	*	*	*	*	1.1	1.9	4.3	6.1	14.2	38.1
1999.....	1.8	*	*	*	*	*	1.0	3.1	5.5	8.2	15.1	46.9
Black, female												
2004.....	0.9	*	*	*	*	*	*	*	*	3.8	7.1	22.8
2003.....	1.1	*	*	*	*	*	*	*	1.5	4.1	11.0	25.5
2002.....	0.8	*	*	*	*	*	*	*	*	3.1	7.6	20.1
2001.....	0.9	*	*	*	*	*	*	*	*	3.7	8.1	18.9
2000.....	0.9	*	*	*	*	*	*	*	*	4.0	7.7	21.0
1999.....	1.1	*	*	*	*	*	*	*	1.8	3.6	8.8	28.5

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Accidental discharge of firearms (W32-W34)

All races 3/, both sexes

2004.....	0.2	*	*	0.1	0.4	0.2	0.2	0.3	0.1	0.2	0.2	*
2003.....	0.3	*	*	0.1	0.5	0.3	0.3	0.2	0.2	0.2	0.3	*
2002.....	0.3	*	*	0.1	0.5	0.4	0.3	0.2	0.2	0.2	0.3	*
2001.....	0.3	*	*	0.1	0.5	0.3	0.3	0.3	0.3	0.2	0.3	*
2000.....	0.3	*	*	0.2	0.5	0.3	0.3	0.2	0.2	0.2	0.2	*
1999.....	0.3	*	*	0.2	0.6	0.4	0.3	0.3	0.2	0.2	0.2	*

All races 3/, male

2004.....	0.4	*	*	0.2	0.7	0.5	0.4	0.4	0.2	0.4	0.4	*
2003.....	0.5	*	*	0.2	0.9	0.5	0.5	0.4	0.4	0.4	0.6	*
2002.....	0.5	*	*	0.2	0.9	0.6	0.4	0.4	0.4	0.4	0.5	*
2001.....	0.5	*	*	0.2	0.9	0.5	0.5	0.5	0.5	0.3	0.7	*
2000.....	0.5	*	*	0.3	0.9	0.5	0.6	0.4	0.4	0.4	0.5	*
1999.....	0.5	*	*	0.3	1.2	0.6	0.5	0.4	0.4	0.4	0.4	*

All races 3/, female

2004.....	0.1	*	*	*	*	*	*	*	*	*	*	*
2003.....	0.1	*	*	*	*	*	*	*	*	*	*	*
2002.....	0.1	*	*	*	*	*	0.1	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	0.1	*	*	*	*	*
2000.....	0.1	*	*	*	*	0.1	0.1	*	*	*	0.1	*
1999.....	0.1	*	*	*	0.1	0.1	0.1	*	*	*	*	*

White, both sexes

2004.....	0.2	*	*	0.1	0.4	0.2	0.2	0.3	0.1	0.2	0.2	*
2003.....	0.2	*	*	0.1	0.5	0.3	0.3	0.2	0.2	0.2	0.3	*
2002.....	0.3	*	*	0.1	0.4	0.3	0.3	0.2	0.2	0.2	0.2	*
2001.....	0.3	*	*	0.1	0.5	0.3	0.3	0.3	0.3	0.2	0.3	*
2000.....	0.3	*	*	0.2	0.5	0.3	0.3	0.2	0.2	0.2	0.2	*
1999.....	0.3	*	*	0.2	0.6	0.4	0.3	0.3	0.2	0.2	0.2	*

White, male

2004.....	0.4	*	*	0.2	0.6	0.4	0.4	0.5	0.2	0.4	*	*
2003.....	0.4	*	*	0.2	0.8	0.5	0.5	0.4	0.5	0.4	0.6	*
2002.....	0.4	*	*	0.2	0.8	0.5	0.4	0.4	0.4	0.4	0.5	*
2001.....	0.5	*	*	0.3	0.9	0.5	0.6	0.5	0.5	0.3	0.7	*
2000.....	0.5	*	*	0.3	0.8	0.5	0.6	0.4	0.4	0.4	0.5	*
1999.....	0.5	*	*	0.3	1.0	0.6	0.5	0.4	0.4	0.4	*	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Accidental explosion (W39-W40)

All races 3/, both sexes

2004.....	0.1	*	*	*	*	*	0.1	0.1	*	*	*	*
2003.....	0.1	*	*	*	0.1	0.1	0.1	0.1	*	*	*	*
2002.....	0.0	*	*	*	*	*	0.1	0.1	*	*	*	*
2001.....	0.1	*	*	*	*	0.1	0.1	0.1	*	*	*	*
2000.....	0.1	*	*	*	*	0.1	0.1	0.1	*	0.1	*	*
1999.....	0.1	*	*	*	*	0.1	0.1	0.1	*	*	*	*

All races 3/, male

2004.....	0.1	*	*	*	*	*	0.2	0.1	*	*	*	*
2003.....	0.1	*	*	*	*	0.1	0.1	0.1	*	*	*	*
2002.....	0.1	*	*	*	*	*	0.2	0.1	*	*	*	*
2001.....	0.1	*	*	*	*	*	0.1	0.1	*	*	*	*
2000.....	0.1	*	*	*	*	0.1	0.1	0.1	*	*	*	*
1999.....	0.1	*	*	*	*	0.1	0.2	0.2	*	*	*	*

All races 3/, female

2004.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2003.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	*	*	*	*	*	*

White, both sexes

2004.....	0.1	*	*	*	*	*	0.1	0.1	*	*	*	*
2003.....	0.1	*	*	*	*	0.1	0.1	0.1	*	*	*	*
2002.....	0.1	*	*	*	*	*	0.1	0.1	*	*	*	*
2001.....	0.1	*	*	*	*	0.1	0.1	0.1	*	*	0.1	*
2000.....	0.1	*	*	*	*	0.1	0.1	0.1	*	*	*	*
1999.....	0.1	*	*	*	*	0.1	0.1	0.1	*	*	*	*

White, male

2004.....	0.1	*	*	*	*	*	0.2	0.1	*	*	*	*
2003.....	0.1	*	*	*	*	0.1	0.2	0.1	*	*	*	*
2002.....	0.1	*	*	*	*	*	0.2	0.1	*	*	*	*
2001.....	0.1	*	*	*	*	*	0.1	*	*	*	*	*
2000.....	0.1	*	*	*	*	0.2	0.2	0.1	*	*	*	*
1999.....	0.1	*	*	*	*	0.1	0.2	0.2	*	*	*	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidental drowning and submersion (W65-W74)												
All races 3/, both sexes												
2004.....	1.1	1.5	2.7	0.7	1.4	1.0	1.0	1.1	0.9	1.0	1.2	1.6
2003.....	1.1	1.4	2.9	0.7	1.4	0.9	1.0	1.0	0.9	1.1	1.3	1.8
2002.....	1.2	1.6	2.9	0.8	1.5	1.1	1.2	0.9	0.8	0.9	1.3	1.7
2001.....	1.2	1.7	3.0	0.8	1.5	0.9	1.0	0.9	0.8	0.9	1.2	1.7
2000.....	1.2	2.0	3.2	0.9	1.6	1.1	1.1	0.9	0.9	1.0	1.3	1.5
1999.....	1.3	1.8	3.2	0.9	1.7	1.1	1.1	1.0	0.9	1.0	1.3	1.6
All races 3/, male												
2004.....	1.8	1.8	3.2	0.9	2.4	1.6	1.7	1.8	1.5	1.7	2.2	2.9
2003.....	1.8	1.7	3.7	0.9	2.5	1.6	1.8	1.6	1.6	1.7	2.0	3.3
2002.....	1.9	1.9	3.9	1.2	2.8	1.9	2.0	1.5	1.3	1.6	2.1	3.6
2001.....	1.8	1.5	3.7	1.1	2.7	1.6	1.7	1.6	1.3	1.5	2.0	3.5
2000.....	2.0	2.2	4.0	1.3	2.9	1.8	1.8	1.6	1.5	1.6	1.8	2.8
1999.....	2.0	2.0	4.0	1.3	3.0	2.0	1.8	1.7	1.5	1.6	2.3	2.3
All races 3/, female												
2004.....	0.5	1.2	2.1	0.4	0.3	0.3	0.3	0.4	0.4	0.5	0.6	1.0
2003.....	0.5	1.2	2.0	0.3	0.2	0.2	0.3	0.4	0.4	0.6	0.8	1.1
2002.....	0.5	1.2	1.9	0.4	0.3	0.3	0.4	0.4	0.4	0.4	0.8	0.9
2001.....	0.5	1.9	2.2	0.5	0.3	0.2	0.4	0.3	0.4	0.3	0.7	1.0
2000.....	0.5	1.7	2.4	0.5	0.3	0.3	0.3	0.3	0.3	0.5	0.9	1.0
1999.....	0.5	1.6	2.4	0.5	0.3	0.2	0.3	0.3	0.4	0.6	0.6	1.4
White, both sexes												
2004.....	1.1	1.6	3.0	0.4	1.2	0.9	1.0	1.1	0.9	1.0	1.2	1.5
2003.....	1.1	1.4	3.1	0.5	1.3	0.9	1.0	1.0	0.9	1.1	1.3	1.6
2002.....	1.1	1.6	3.0	0.5	1.4	1.0	1.1	0.9	0.8	0.9	1.3	1.6
2001.....	1.1	1.7	3.3	0.5	1.3	0.9	1.0	0.9	0.8	0.9	1.2	1.7
2000.....	1.2	1.6	3.5	0.7	1.5	1.0	1.0	0.9	0.9	1.0	1.2	1.5
1999.....	1.2	1.5	3.5	0.6	1.5	1.0	1.1	1.0	0.9	1.0	1.3	1.6
White, male												
2004.....	1.7	1.8	3.5	0.6	2.2	1.6	1.6	1.7	1.4	1.6	2.1	2.8
2003.....	1.8	1.5	4.0	0.7	2.3	1.5	1.7	1.5	1.5	1.8	2.1	3.3
2002.....	1.8	1.9	4.0	0.7	2.5	1.8	1.9	1.4	1.2	1.5	2.1	3.7
2001.....	1.7	1.6	4.2	0.7	2.4	1.4	1.6	1.5	1.2	1.5	2.0	3.5
2000.....	1.9	1.8	4.4	0.9	2.6	1.7	1.7	1.5	1.6	1.6	1.8	2.7
1999.....	1.9	1.7	4.3	0.8	2.7	1.8	1.8	1.7	1.4	1.4	2.2	2.4

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Accidental inhalation and ingestion of food or other objects causing obstruction of respiratory tract (W78-W80)

All races 3/, both sexes

2004.....	1.5	1.4	0.4	0.1	0.2	0.2	0.4	0.8	1.3	2.9	9.8	29.9
2003.....	1.5	1.5	0.6	0.1	0.1	0.2	0.4	0.8	1.4	3.1	9.3	27.4
2002.....	1.4	1.5	0.5	0.1	0.1	0.2	0.4	0.6	1.2	3.1	9.5	28.4
2001.....	1.5	1.5	0.5	0.1	0.1	0.1	0.4	0.9	1.3	2.9	9.5	30.0
2000.....	1.5	2.0	0.6	0.1	0.1	0.2	0.4	0.8	1.4	3.1	10.0	32.6
1999.....	1.4	1.7	0.6	0.1	0.1	0.2	0.4	0.7	1.1	3.1	8.8	30.3

All races 3/, male

2004.....	1.6	2.3	0.6	*	0.2	0.3	0.4	0.9	1.6	3.6	12.5	39.0
2003.....	1.6	1.6	0.7	*	0.2	0.3	0.5	0.9	1.8	3.7	12.5	37.5
2002.....	1.5	2.0	0.6	0.1	0.2	0.3	0.5	0.8	1.6	3.9	12.1	37.0
2001.....	1.5	2.0	0.6	*	0.2	0.2	0.5	1.1	1.6	3.5	12.4	39.8
2000.....	1.6	2.3	0.7	*	0.2	0.3	0.5	1.1	1.6	3.6	13.2	41.0
1999.....	1.4	1.9	0.8	0.1	0.1	0.3	0.5	0.8	1.3	3.8	10.9	39.2

All races 3/, female

2004.....	1.5	*	0.3	*	0.1	0.2	0.3	0.7	1.0	2.4	8.0	25.8
2003.....	1.4	1.5	0.4	*	0.1	0.1	0.3	0.7	1.0	2.6	7.2	23.0
2002.....	1.4	1.0	0.3	*	*	0.1	0.3	0.5	0.9	2.4	7.8	24.6
2001.....	1.4	1.0	0.4	*	0.1	0.1	0.3	0.7	1.0	2.4	7.6	26.0
2000.....	1.5	1.8	0.5	*	*	0.2	0.3	0.5	1.1	2.7	7.9	29.1
1999.....	1.4	1.4	0.4	0.1	*	0.1	0.2	0.5	1.0	2.5	7.4	26.7

White, both sexes

2004.....	1.6	1.4	0.3	0.1	0.2	0.2	0.4	0.8	1.3	2.7	9.7	30.1
2003.....	1.6	1.3	0.5	*	0.1	0.2	0.4	0.8	1.3	3.0	9.3	27.9
2002.....	1.5	1.4	0.4	0.1	0.1	0.2	0.4	0.6	1.1	2.8	9.4	28.3
2001.....	1.5	1.3	0.5	0.1	0.1	0.1	0.4	0.9	1.2	2.8	9.1	29.7
2000.....	1.6	1.7	0.5	*	0.1	0.2	0.4	0.7	1.3	2.9	9.9	32.5
1999.....	1.4	1.4	0.5	0.1	0.1	0.2	0.4	0.6	1.0	2.9	8.4	30.4

White, male

2004.....	1.7	2.3	0.4	*	0.2	0.2	0.5	1.0	1.6	3.3	12.5	39.2
2003.....	1.6	1.3	0.7	*	0.2	0.3	0.5	0.9	1.7	3.6	12.4	38.2
2002.....	1.5	1.9	0.5	*	0.2	0.3	0.5	0.8	1.5	3.5	11.8	37.2
2001.....	1.5	1.8	0.5	*	0.2	0.2	0.5	1.0	1.4	3.4	11.6	39.3
2000.....	1.6	1.9	0.7	*	0.1	0.3	0.5	1.0	1.5	3.3	12.9	40.2
1999.....	1.4	1.7	0.6	*	0.1	0.2	0.5	0.8	1.1	3.6	10.4	39.7

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Accidental inhalation and ingestion of food or other objects causing obstruction of respiratory tract (W78-W80)

White, female

2004.....	1.6	*	*	*	0.1	0.2	0.4	0.7	1.0	2.2	7.8	26.0
2003.....	1.5	1.4	0.4	*	*	*	0.3	0.7	1.0	2.5	7.2	23.3
2002.....	1.5	*	*	*	*	0.1	0.3	0.5	0.8	2.2	7.8	24.5
2001.....	1.5	*	0.4	*	*	*	0.3	0.7	1.0	2.3	7.4	25.7
2000.....	1.6	1.5	*	*	*	0.2	0.3	0.5	1.1	2.5	7.9	29.4
1999.....	1.5	*	*	*	*	*	0.3	0.5	0.9	2.3	7.1	26.7

Black, both sexes

2004.....	1.3	*	0.8	*	*	*	0.4	0.9	1.9	5.2	13.2	30.5
2003.....	1.3	*	*	*	*	0.4	0.6	1.0	2.1	4.9	10.6	24.5
2002.....	1.4	*	0.8	*	*	*	0.4	1.0	2.0	6.4	13.0	33.0
2001.....	1.5	3.1	*	*	*	*	0.5	1.2	2.5	5.3	14.3	36.8
2000.....	1.5	3.7	1.0	*	*	0.4	0.4	1.1	2.2	6.0	13.1	35.3
1999.....	1.5	*	1.1	*	*	0.4	0.4	1.1	2.5	5.5	14.6	33.2

Black, male

2004.....	1.4	*	*	*	*	*	*	1.1	2.2	6.9	17.0	42.7
2003.....	1.4	*	*	*	*	*	0.8	1.3	2.8	5.5	16.4	30.2
2002.....	1.6	*	*	*	*	*	*	1.3	2.6	8.4	19.2	41.8
2001.....	1.7	*	*	*	*	*	*	1.5	3.5	6.8	23.2	51.1
2000.....	1.6	*	*	*	*	*	*	1.7	2.8	7.3	20.0	51.9
1999.....	1.7	*	*	*	*	*	*	1.3	3.3	6.9	20.3	43.4

Black, female

2004.....	1.3	*	*	*	*	*	*	0.8	1.6	4.0	11.0	25.6
2003.....	1.1	*	*	*	*	*	*	*	1.4	4.5	7.3	22.3
2002.....	1.3	*	*	*	*	*	*	*	1.6	5.0	9.4	29.6
2001.....	1.3	*	*	*	*	*	*	0.8	1.6	4.3	9.3	31.6
2000.....	1.4	*	*	*	*	*	*	*	1.6	5.0	9.2	29.2
1999.....	1.3	*	*	*	*	*	*	*	1.8	4.5	11.4	29.3

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other accidental and unspecified threats to breathing (W75-W77,W81-W84)

All races 3/, both sexes

2004.....	0.5	16.4	0.3	0.2	0.2	0.2	0.3	0.3	0.2	0.2	0.3	0.6
2003.....	0.4	13.9	0.5	0.1	0.2	0.3	0.3	0.3	0.2	0.2	0.3	0.8
2002.....	0.5	14.3	0.4	0.2	0.3	0.3	0.3	0.3	0.2	0.3	0.4	1.0
2001.....	0.5	13.7	0.4	0.2	0.2	0.2	0.4	0.3	0.2	0.3	0.4	0.9
2000.....	0.5	11.8	0.4	0.2	0.3	0.3	0.4	0.3	0.3	0.3	0.5	1.2
1999.....	0.6	10.8	0.5	0.2	0.3	0.2	0.3	0.4	0.3	0.6	1.5	4.7

All races 3/, male

2004.....	0.7	17.7	0.4	0.3	0.4	0.3	0.6	0.5	0.3	0.3	0.4	*
2003.....	0.6	16.3	0.5	0.2	0.4	0.4	0.6	0.5	0.4	0.3	*	*
2002.....	0.7	15.7	0.5	0.3	0.4	0.5	0.5	0.5	0.4	0.3	0.5	*
2001.....	0.7	15.3	0.5	0.3	0.4	0.4	0.6	0.5	0.3	0.4	0.5	*
2000.....	0.7	13.2	0.5	0.3	0.5	0.5	0.6	0.5	0.5	0.4	0.6	*
1999.....	0.8	12.6	0.5	0.4	0.4	0.4	0.5	0.6	0.5	0.9	1.8	5.4

All races 3/, female

2004.....	0.3	14.9	*	*	*	0.1	0.1	0.1	*	*	*	0.6
2003.....	0.3	11.4	0.4	*	*	*	0.1	*	*	*	0.3	0.7
2002.....	0.3	12.8	0.4	*	*	*	0.1	*	*	*	*	1.1
2001.....	0.3	12.0	0.3	*	*	*	0.1	*	*	0.2	0.4	0.9
2000.....	0.3	10.3	0.3	0.1	*	*	0.1	0.1	*	0.2	0.4	1.1
1999.....	0.4	8.9	0.5	*	*	*	0.1	0.1	0.2	0.4	1.3	4.5

White, both sexes

2004.....	0.5	13.2	0.4	0.2	0.3	0.3	0.4	0.4	0.2	0.2	0.3	0.6
2003.....	0.4	11.1	0.5	0.1	0.2	0.3	0.4	0.3	0.2	0.2	0.3	0.8
2002.....	0.4	10.7	0.4	0.2	0.3	0.3	0.3	0.3	0.3	0.3	0.4	1.0
2001.....	0.5	10.8	0.4	0.2	0.3	0.3	0.4	0.3	0.2	0.3	0.4	0.9
2000.....	0.5	9.4	0.4	0.2	0.3	0.3	0.4	0.3	0.4	0.3	0.5	1.2
1999.....	0.6	8.7	0.5	0.2	0.3	0.3	0.3	0.4	0.3	0.6	1.5	4.9

White, male

2004.....	0.7	15.0	0.5	0.3	0.5	0.4	0.6	0.6	0.3	0.3	*	*
2003.....	0.6	13.2	0.6	0.2	0.4	0.5	0.6	0.5	0.4	0.3	*	*
2002.....	0.6	12.3	0.5	0.3	0.5	0.5	0.5	0.6	0.4	0.4	0.5	*
2001.....	0.6	11.9	0.4	0.3	0.4	0.5	0.7	0.6	0.3	0.5	0.5	*
2000.....	0.7	10.8	0.5	0.4	0.5	0.5	0.7	0.5	0.6	0.4	0.6	*
1999.....	0.8	9.7	0.5	0.4	0.5	0.5	0.6	0.6	0.4	0.8	1.8	5.7

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Accidental exposure to electric current (W85-W87)

All races 3/, both sexes

2004.....	0.1	*	*	*	0.2	0.2	0.2	0.2	0.1	*	*	*
2003.....	0.1	*	*	*	0.2	0.2	0.2	0.1	0.1	*	*	*
2002.....	0.1	*	*	*	0.2	0.2	0.3	0.2	0.1	*	*	*
2001.....	0.1	*	*	*	0.2	0.2	0.3	0.1	0.1	*	*	*
2000.....	0.1	*	*	*	0.2	0.2	0.2	0.2	0.1	*	*	*
1999.....	0.2	*	*	*	0.2	0.2	0.3	0.2	0.2	0.1	*	*

All races 3/, male

2004.....	0.2	*	*	*	0.3	0.4	0.3	0.4	0.3	*	*	*
2003.....	0.3	*	*	*	0.3	0.4	0.4	0.3	0.2	*	*	*
2002.....	0.3	*	*	*	0.4	0.5	0.5	0.4	0.2	*	*	*
2001.....	0.3	*	*	*	0.3	0.4	0.5	0.3	0.2	*	*	*
2000.....	0.3	*	*	*	0.3	0.4	0.4	0.4	0.3	*	*	*
1999.....	0.3	*	*	*	0.4	0.4	0.5	0.3	0.3	0.2	*	*

All races 3/, female

2004.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	*	*	*	*	*	*

White, both sexes

2004.....	0.1	*	*	*	0.2	0.3	0.2	0.2	0.1	*	*	*
2003.....	0.1	*	*	*	0.2	0.2	0.2	0.2	0.1	*	*	*
2002.....	0.2	*	*	*	0.2	0.3	0.3	0.2	0.1	*	*	*
2001.....	0.2	*	*	*	0.2	0.3	0.3	0.1	0.1	*	*	*
2000.....	0.2	*	*	*	0.2	0.3	0.3	0.2	0.1	*	*	*
1999.....	0.2	*	*	*	0.2	0.2	0.3	0.2	0.2	*	*	*

White, male

2004.....	0.3	*	*	*	0.3	0.5	0.4	0.4	0.3	*	*	*
2003.....	0.3	*	*	*	0.4	0.5	0.4	0.3	0.2	*	*	*
2002.....	0.3	*	*	*	0.4	0.5	0.6	0.4	*	*	*	*
2001.....	0.3	*	*	*	0.3	0.5	0.6	0.3	0.2	*	*	*
2000.....	0.3	*	*	*	0.4	0.5	0.5	0.4	0.3	*	*	*
1999.....	0.3	*	*	*	0.4	0.5	0.6	0.3	0.4	*	*	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidental exposure to smoke, fire and flames (X00-X09)												
All races 3/, both sexes												
2004.....	1.1	0.6	1.4	0.6	0.4	0.6	0.7	1.2	1.4	1.9	3.5	5.6
2003.....	1.2	0.7	1.4	0.5	0.5	0.6	0.9	1.3	1.4	2.3	3.6	5.4
2002.....	1.1	0.9	1.4	0.6	0.5	0.7	0.8	1.1	1.3	2.1	3.4	4.8
2001.....	1.2	1.2	1.5	0.6	0.5	0.6	1.0	1.1	1.5	2.2	3.4	5.7
2000.....	1.2	1.0	1.9	0.6	0.5	0.6	0.9	1.2	1.5	2.2	3.8	6.2
1999.....	1.2	1.1	2.0	0.6	0.5	0.7	0.9	1.0	1.3	2.3	4.0	5.9
All races 3/, male												
2004.....	1.3	*	1.5	0.7	0.5	0.8	0.9	1.6	1.8	2.5	4.5	7.2
2003.....	1.4	*	1.6	0.5	0.6	0.8	1.2	1.6	1.9	2.9	4.4	7.4
2002.....	1.4	1.0	1.6	0.7	0.6	0.8	1.1	1.5	1.7	2.9	4.5	7.0
2001.....	1.4	1.5	1.8	0.7	0.7	0.8	1.4	1.5	1.9	2.8	4.1	8.3
2000.....	1.5	1.0	2.2	0.7	0.6	0.8	1.2	1.6	2.1	2.8	5.0	9.4
1999.....	1.5	1.1	2.2	0.7	0.7	1.0	1.3	1.4	1.7	2.9	5.1	9.3
All races 3/, female												
2004.....	0.9	*	1.3	0.5	0.4	0.4	0.5	0.8	1.1	1.4	2.9	4.9
2003.....	1.0	1.1	1.2	0.5	0.4	0.5	0.6	0.9	1.0	1.8	3.0	4.5
2002.....	0.8	*	1.2	0.5	0.3	0.5	0.5	0.8	0.9	1.4	2.6	3.9
2001.....	0.9	1.0	1.1	0.5	0.4	0.4	0.6	0.7	1.1	1.8	2.9	4.6
2000.....	0.9	*	1.5	0.6	0.4	0.4	0.6	0.7	1.0	1.7	3.0	4.9
1999.....	0.9	1.1	1.7	0.6	0.4	0.4	0.5	0.6	1.0	1.7	3.2	4.5
White, both sexes												
2004.....	1.0	*	1.1	0.4	0.4	0.6	0.7	1.0	1.2	1.6	3.1	4.9
2003.....	1.1	0.7	1.2	0.4	0.5	0.6	0.9	1.2	1.3	2.0	3.1	4.7
2002.....	1.0	*	1.0	0.4	0.4	0.6	0.8	1.0	1.1	1.8	3.0	4.0
2001.....	1.0	1.1	1.2	0.5	0.5	0.6	0.9	0.9	1.3	1.9	2.8	4.8
2000.....	1.0	0.8	1.3	0.5	0.5	0.5	0.8	1.0	1.3	1.8	3.1	5.1
1999.....	1.0	0.9	1.5	0.5	0.5	0.7	0.8	0.9	1.2	1.8	3.3	4.7
White, male												
2004.....	1.2	*	1.2	0.5	0.5	0.8	0.9	1.4	1.6	2.1	3.9	6.5
2003.....	1.3	*	1.4	0.4	0.6	0.7	1.2	1.5	1.7	2.6	3.6	6.7
2002.....	1.2	*	1.1	0.6	0.6	0.8	1.1	1.3	1.5	2.5	4.1	5.8
2001.....	1.3	*	1.6	0.6	0.7	0.7	1.2	1.2	1.6	2.2	3.4	7.0
2000.....	1.3	*	1.6	0.5	0.6	0.7	1.1	1.5	1.7	2.3	4.1	8.2
1999.....	1.3	*	1.7	0.5	0.7	1.0	1.2	1.2	1.5	2.3	4.2	8.0

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidental exposure to smoke, fire and flames (X00-X09)												
White, female												
2004.....	0.8	*	1.0	0.4	0.4	0.4	0.5	0.7	0.9	1.2	2.5	4.2
2003.....	0.9	*	1.0	0.4	0.4	0.5	0.6	0.8	0.9	1.5	2.7	3.8
2002.....	0.7	*	1.0	0.3	0.3	0.4	0.5	0.7	0.7	1.3	2.2	3.2
2001.....	0.8	*	0.8	0.4	0.3	0.4	0.5	0.6	0.9	1.6	2.5	3.8
2000.....	0.8	*	1.1	0.5	0.3	0.4	0.5	0.6	0.8	1.5	2.4	3.9
1999.....	0.8	*	1.3	0.5	0.3	0.4	0.5	0.5	0.8	1.4	2.6	3.3
Black, both sexes												
2004.....	2.1	*	2.8	1.6	0.6	1.0	1.2	2.4	3.4	4.8	9.7	14.9
2003.....	2.0	*	3.0	1.4	0.6	0.9	1.2	2.6	2.9	5.3	10.2	14.9
2002.....	2.1	*	3.5	1.4	0.8	1.1	1.0	2.1	3.3	5.0	8.6	15.7
2001.....	2.3	*	3.3	1.3	0.7	1.2	1.8	2.4	3.5	6.1	9.7	17.8
2000.....	2.5	*	4.8	1.5	0.7	1.2	1.7	2.4	4.2	5.9	13.2	20.3
1999.....	2.5	*	4.5	1.5	0.7	1.0	1.8	2.1	3.4	6.9	13.2	21.7
Black, male												
2004.....	2.6	*	2.9	1.9	0.7	1.3	1.7	3.8	4.4	7.5	13.4	*
2003.....	2.4	*	3.2	1.5	*	1.1	1.4	3.3	4.3	7.5	16.1	*
2002.....	2.6	*	4.1	1.6	1.0	1.4	1.7	3.0	4.7	7.8	10.9	25.7
2001.....	2.9	*	3.6	1.6	0.8	1.4	2.7	3.9	5.0	9.0	12.9	27.2
2000.....	3.2	*	5.6	1.7	0.9	1.5	2.4	3.5	6.3	8.8	18.5	27.7
1999.....	3.0	*	5.0	1.7	0.8	1.2	2.7	3.2	4.1	10.1	18.1	29.3
Black, female												
2004.....	1.6	*	2.7	1.2	*	*	0.7	1.2	2.6	3.0	7.6	13.4
2003.....	1.7	*	2.7	1.3	0.6	*	1.0	2.0	1.7	3.7	6.9	13.4
2002.....	1.6	*	2.8	1.3	*	0.9	*	1.4	2.2	3.0	7.3	11.9
2001.....	1.7	*	2.9	1.0	*	1.1	1.0	1.1	2.4	4.1	7.9	14.3
2000.....	2.0	*	4.1	1.3	*	0.9	1.1	1.4	2.4	3.9	10.2	17.6
1999.....	2.0	*	3.9	1.3	*	0.8	1.0	1.3	2.7	4.6	10.5	18.8

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Accidental poisoning and exposure to noxious substances (X40-X49)

All races 3/, both sexes

2004.....	7.1	*	*	0.1	5.4	9.1	14.6	14.5	5.4	2.3	2.4	3.4
2003.....	6.7	0.5	0.3	0.1	4.9	8.6	14.0	13.3	4.9	2.2	2.2	3.5
2002.....	6.1	0.6	0.2	0.1	4.1	7.8	13.4	11.7	4.1	2.1	2.3	3.9
2001.....	4.9	*	0.2	0.1	3.4	6.3	11.2	9.1	3.2	1.6	2.2	3.5
2000.....	4.5	*	0.2	0.1	3.0	6.0	10.3	8.1	2.8	1.5	2.0	4.3
1999.....	4.4	*	0.2	0.1	2.5	5.9	10.1	7.8	2.8	1.6	2.2	3.4

All races 3/, male

2004.....	9.6	*	*	0.1	8.2	13.1	19.0	19.1	6.6	2.6	2.7	4.2
2003.....	9.2	*	0.3	0.1	7.4	12.4	18.8	18.0	6.3	2.7	2.8	4.0
2002.....	8.5	1.0	*	0.1	6.3	11.5	18.4	15.9	5.4	2.5	2.7	5.1
2001.....	7.1	*	0.3	0.1	5.3	9.3	15.7	13.0	4.1	2.0	2.7	3.8
2000.....	6.6	*	0.3	0.1	4.6	9.0	15.1	12.0	3.4	1.8	2.1	6.2
1999.....	6.5	*	*	0.1	3.8	9.0	14.9	11.7	3.9	2.1	2.4	4.4

All races 3/, female

2004.....	4.7	*	*	0.1	2.4	4.9	10.3	10.0	4.3	2.0	2.2	3.0
2003.....	4.3	*	0.3	0.2	2.2	4.7	9.3	8.8	3.7	1.8	1.8	3.3
2002.....	3.7	*	*	*	1.9	4.0	8.4	7.6	3.0	1.7	2.1	3.4
2001.....	2.9	*	*	0.1	1.4	3.3	6.7	5.2	2.3	1.2	1.9	3.3
2000.....	2.5	*	*	*	1.3	2.8	5.6	4.4	2.3	1.3	1.9	3.6
1999.....	2.3	*	*	*	1.1	2.6	5.3	4.0	1.8	1.3	2.0	3.0

White, both sexes

2004.....	7.6	*	*	0.1	6.4	10.3	15.7	14.6	5.0	2.2	2.4	3.4
2003.....	7.1	*	0.3	0.1	5.8	9.7	14.8	13.3	4.8	2.0	2.3	3.5
2002.....	6.3	*	*	0.1	4.9	8.7	14.0	11.3	3.8	2.1	2.3	3.9
2001.....	5.0	*	0.2	0.1	3.9	6.9	11.4	8.4	2.8	1.5	2.1	3.3
2000.....	4.6	*	*	0.1	3.4	6.4	10.4	7.5	2.6	1.4	2.1	4.1
1999.....	4.3	*	0.2	0.1	2.8	6.1	10.0	7.1	2.5	1.5	2.2	3.3

White, male

2004.....	10.2	*	*	0.2	9.7	14.9	20.3	18.9	5.7	2.4	2.5	4.3
2003.....	9.7	*	*	0.1	8.8	13.9	19.8	17.6	5.8	2.2	2.8	3.7
2002.....	8.8	*	*	0.2	7.4	12.7	19.2	15.2	4.5	2.3	2.6	5.2
2001.....	7.2	*	*	*	6.0	10.2	16.2	11.9	3.4	1.9	2.5	3.7
2000.....	6.7	*	*	*	5.3	9.7	15.2	10.9	2.9	1.6	2.1	5.9
1999.....	6.4	*	*	*	4.2	9.5	14.9	10.5	3.2	1.9	2.5	4.5

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Accidental poisoning and exposure to noxious substances (X40-X49)

White, female

2004.....	5.0	*	*	*	2.8	5.5	11.0	10.3	4.4	2.0	2.2	3.0
2003.....	4.5	*	*	*	2.6	5.2	9.8	9.0	3.8	1.8	1.9	3.4
2002.....	3.9	*	*	*	2.2	4.5	8.7	7.5	3.0	1.8	2.1	3.4
2001.....	2.9	*	*	0.1	1.6	3.5	6.6	5.0	2.3	1.2	1.8	3.1
2000.....	2.5	*	*	*	1.3	3.0	5.5	4.1	2.3	1.3	2.0	3.4
1999.....	2.3	*	*	*	1.2	2.6	5.1	3.8	1.8	1.2	2.1	2.8

Black, both sexes

2004.....	6.5	*	*	*	1.8	5.3	12.8	18.8	9.8	3.7	3.5	*
2003.....	6.3	*	*	*	1.4	5.7	13.5	17.9	7.6	4.1	2.0	*
2002.....	6.3	*	*	*	1.4	5.6	13.8	17.9	8.4	2.7	2.8	*
2001.....	5.8	*	*	*	1.6	5.2	13.1	16.7	6.5	2.5	3.6	6.4
2000.....	5.5	*	*	*	1.3	5.6	13.0	15.5	5.5	2.6	*	8.1
1999.....	5.7	*	*	*	1.6	6.1	13.1	15.5	6.0	3.0	*	*

Black, male

2004.....	8.9	*	*	*	2.7	7.6	16.4	27.1	16.2	6.2	5.6	*
2003.....	8.8	*	*	*	2.0	8.0	18.6	26.2	12.8	7.6	*	*
2002.....	9.1	*	*	*	2.2	8.5	19.3	26.5	15.0	5.0	*	*
2001.....	8.4	*	*	*	2.3	7.5	17.9	26.6	11.3	3.9	5.9	*
2000.....	8.2	*	*	*	1.7	8.4	18.7	24.9	9.5	4.5	*	*
1999.....	8.7	*	*	*	2.7	9.1	19.4	25.9	11.0	4.4	*	*

Black, female

2004.....	4.3	*	*	*	1.0	3.3	9.7	11.8	4.7	*	*	*
2003.....	3.9	*	*	*	0.7	3.6	8.9	10.8	3.3	*	*	*
2002.....	3.7	*	*	*	0.7	3.0	8.9	10.5	3.2	*	*	*
2001.....	3.5	*	*	*	0.9	3.1	9.0	8.2	2.6	*	*	*
2000.....	3.1	*	*	*	1.0	3.0	7.8	7.4	2.3	*	*	*
1999.....	3.0	*	*	*	*	3.4	7.6	6.5	2.0	2.1	*	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidental poisoning by and exposure to drugs and other biological substances (X40-X44)												
All races 3/, both sexes												
2004.....	6.8	*	*	0.1	5.2	8.8	14.1	13.9	5.0	1.8	1.8	2.5
2003.....	6.3	*	0.2	0.1	4.6	8.3	13.4	12.7	4.4	1.7	1.7	2.8
2002.....	5.7	*	*	*	3.9	7.4	12.8	11.1	3.6	1.6	1.7	3.1
2001.....	4.6	*	*	0.1	3.1	6.0	10.6	8.5	2.7	1.3	1.6	2.5
2000.....	4.2	*	*	*	2.7	5.6	9.8	7.6	2.3	1.1	1.4	3.3
1999.....	4.0	*	*	*	2.2	5.6	9.5	7.2	2.3	1.2	1.6	2.8
All races 3/, male												
2004.....	9.1	*	*	*	7.8	12.6	18.1	18.1	6.0	1.9	1.8	2.6
2003.....	8.6	*	*	*	6.9	11.9	17.9	16.9	5.4	2.0	2.1	2.7
2002.....	7.9	*	*	*	5.9	10.9	17.5	15.0	4.5	1.7	1.8	3.7
2001.....	6.5	*	*	*	4.8	8.9	14.8	12.2	3.5	1.5	1.7	1.8
2000.....	6.1	*	*	*	4.1	8.5	14.3	11.2	2.7	1.3	1.3	4.4
1999.....	5.9	*	*	*	3.3	8.5	14.0	10.9	3.2	1.5	1.5	3.3
All races 3/, female												
2004.....	4.5	*	*	*	2.3	4.8	10.1	9.7	4.1	1.7	1.8	2.5
2003.....	4.1	*	*	*	2.1	4.6	9.0	8.6	3.4	1.5	1.5	2.9
2002.....	3.5	*	*	*	1.8	3.9	8.1	7.3	2.7	1.4	1.6	2.8
2001.....	2.7	*	*	*	1.3	3.1	6.4	5.0	2.0	1.1	1.4	2.7
2000.....	2.3	*	*	*	1.2	2.7	5.5	4.1	1.9	1.0	1.5	2.8
1999.....	2.1	*	*	*	1.0	2.5	5.1	3.7	1.5	1.0	1.6	2.6
White, both sexes												
2004.....	7.2	*	*	0.1	6.1	10.0	15.1	13.9	4.6	1.7	1.8	2.6
2003.....	6.7	*	*	*	5.5	9.3	14.2	12.7	4.2	1.5	1.8	2.8
2002.....	5.9	*	*	*	4.6	8.3	13.4	10.8	3.3	1.5	1.7	3.2
2001.....	4.7	*	*	0.1	3.6	6.6	10.9	7.9	2.4	1.2	1.6	2.3
2000.....	4.2	*	*	*	3.1	6.0	9.9	7.0	2.1	1.1	1.5	3.1
1999.....	4.0	*	*	*	2.5	5.8	9.5	6.6	2.1	1.1	1.6	2.6
White, male												
2004.....	9.6	*	*	*	9.3	14.3	19.4	17.9	5.1	1.7	1.7	2.7
2003.....	9.1	*	*	*	8.3	13.3	18.8	16.6	5.0	1.6	2.1	2.4
2002.....	8.2	*	*	*	6.9	12.0	18.3	14.4	3.8	1.6	1.7	3.8
2001.....	6.6	*	*	*	5.5	9.7	15.3	11.0	2.9	1.4	1.7	1.7
2000.....	6.1	*	*	*	4.8	9.0	14.4	10.1	2.2	1.1	1.3	4.1
1999.....	5.9	*	*	*	3.7	9.0	14.0	9.7	2.6	1.3	1.5	3.2

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Accidental poisoning by and exposure to drugs and other biological substances (X40-X44)

White, female

2004.....	4.8	*	*	*	2.7	5.4	10.8	9.9	4.2	1.8	1.8	2.5
2003.....	4.3	*	*	*	2.5	5.0	9.5	8.7	3.5	1.5	1.5	3.0
2002.....	3.7	*	*	*	2.1	4.4	8.5	7.3	2.8	1.5	1.7	2.9
2001.....	2.7	*	*	*	1.5	3.3	6.4	4.8	2.1	1.1	1.5	2.6
2000.....	2.3	*	*	*	1.2	2.9	5.4	3.9	2.0	1.0	1.6	2.6
1999.....	2.1	*	*	*	1.1	2.5	4.9	3.6	1.6	0.9	1.7	2.3

Black, both sexes

2004.....	6.1	*	*	*	1.7	5.1	12.5	18.3	9.2	2.9	2.2	*
2003.....	5.9	*	*	*	1.2	5.4	13.1	17.3	6.9	3.2	*	*
2002.....	5.8	*	*	*	1.3	5.3	13.1	17.1	7.5	2.0	*	*
2001.....	5.4	*	*	*	1.3	5.0	12.5	15.9	6.0	2.0	*	*
2000.....	5.2	*	*	*	1.3	5.3	12.4	14.8	4.7	2.0	*	6.3
1999.....	5.3	*	*	*	1.4	5.9	12.6	14.6	5.2	2.3	*	*

Black, male

2004.....	8.4	*	*	*	2.5	7.4	15.8	26.1	15.2	5.0	*	*
2003.....	8.2	*	*	*	1.8	7.4	17.9	25.0	11.6	5.8	*	*
2002.....	8.4	*	*	*	2.0	8.0	18.2	25.2	13.3	3.4	*	*
2001.....	7.8	*	*	*	2.0	7.2	16.8	25.3	10.4	3.2	*	*
2000.....	7.7	*	*	*	1.6	8.0	17.8	23.9	8.3	3.7	*	*
1999.....	8.1	*	*	*	2.3	8.8	18.5	24.5	9.5	3.4	*	*

Black, female

2004.....	4.1	*	*	*	0.9	3.1	9.5	11.6	4.4	*	*	*
2003.....	3.8	*	*	*	0.7	3.6	8.8	10.7	3.2	*	*	*
2002.....	3.5	*	*	*	*	2.9	8.6	10.1	2.9	*	*	*
2001.....	3.2	*	*	*	*	3.0	8.6	7.7	2.4	*	*	*
2000.....	2.9	*	*	*	0.9	2.9	7.7	7.0	1.9	*	*	*
1999.....	2.8	*	*	*	*	3.3	7.4	6.2	1.7	*	*	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Accidental poisoning by and exposure to other and unspecified solid or liquids substances (X45-X46,X48-X49)

All races 3/, both sexes

2004.....	0.2	*	*	*	0.1	0.1	0.3	0.4	0.2	0.1	0.2	*
2003.....	0.2	*	*	*	0.1	0.1	0.4	0.4	0.2	0.1	*	*
2002.....	0.2	*	*	*	0.1	0.2	0.3	0.3	0.3	0.1	0.2	*
2001.....	0.2	*	*	*	0.1	0.1	0.3	0.3	0.2	0.1	*	*
2000.....	0.2	*	*	*	0.1	0.1	0.3	0.3	0.2	0.1	*	*
1999.....	0.2	*	*	*	0.1	0.2	0.3	0.3	0.2	0.2	0.2	*

All races 3/, male

2004.....	0.3	*	*	*	0.2	0.2	0.5	0.6	0.3	*	*	*
2003.....	0.3	*	*	*	0.2	0.2	0.6	0.6	0.4	*	*	*
2002.....	0.3	*	*	*	0.2	0.3	0.5	0.5	0.5	*	*	*
2001.....	0.2	*	*	*	0.2	0.2	0.5	0.4	0.3	*	*	*
2000.....	0.2	*	*	*	0.2	0.2	0.4	0.4	0.3	*	*	*
1999.....	0.3	*	*	*	0.2	0.2	0.5	0.4	0.3	0.3	*	*

All races 3/, female

2004.....	0.1	*	*	*	*	*	0.1	0.2	0.1	*	*	*
2003.....	0.1	*	*	*	*	*	0.2	0.1	*	*	*	*
2002.....	0.1	*	*	*	*	*	0.1	0.1	*	*	*	*
2001.....	0.1	*	*	*	*	*	0.2	0.2	*	*	*	*
2000.....	0.1	*	*	*	*	*	0.1	0.1	0.2	*	*	*
1999.....	0.1	*	*	*	*	*	0.1	0.1	*	*	*	*

White, both sexes

2004.....	0.2	*	*	*	0.2	0.2	0.3	0.4	0.2	0.1	0.2	*
2003.....	0.2	*	*	*	0.1	0.2	0.4	0.4	0.2	*	*	*
2002.....	0.2	*	*	*	0.1	0.2	0.3	0.3	0.2	0.1	*	*
2001.....	0.2	*	*	*	0.1	0.1	0.3	0.3	0.2	*	*	*
2000.....	0.2	*	*	*	0.1	0.1	0.3	0.3	0.2	*	*	*
1999.....	0.2	*	*	*	0.1	0.1	0.3	0.3	0.2	0.1	0.2	*

White, male

2004.....	0.3	*	*	*	0.3	0.3	0.6	0.6	0.2	*	*	*
2003.....	0.3	*	*	*	0.2	0.3	0.6	0.6	0.4	*	*	*
2002.....	0.3	*	*	*	0.2	0.3	0.5	0.4	0.4	*	*	*
2001.....	0.2	*	*	*	0.2	0.2	0.5	0.4	0.3	*	*	*
2000.....	0.2	*	*	*	0.2	0.2	0.4	0.4	0.3	*	*	*
1999.....	0.3	*	*	*	0.2	0.2	0.5	0.4	0.3	*	*	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Accidental poisoning by and exposure to other gases and vapors (X47)

All races 3/, both sexes

2004.....	0.2	*	*	0.1	0.1	0.2	0.2	0.3	0.2	0.3	0.4	0.7
2003.....	0.2	*	0.1	0.1	0.2	0.2	0.2	0.2	0.3	0.4	0.4	0.6
2002.....	0.2	*	*	0.1	0.2	0.2	0.3	0.3	0.3	0.4	0.5	0.6
2001.....	0.2	*	*	0.1	0.2	0.2	0.2	0.2	0.2	0.2	0.6	0.9
2000.....	0.2	*	*	0.1	0.2	0.2	0.2	0.2	0.3	0.3	0.5	0.8
1999.....	0.2	*	*	*	0.2	0.2	0.2	0.2	0.3	0.2	0.4	0.5

All races 3/, male

2004.....	0.3	*	*	*	0.2	0.3	0.3	0.4	0.4	0.5	0.6	1.3
2003.....	0.3	*	*	*	0.2	0.3	0.4	0.4	0.4	0.6	0.6	*
2002.....	0.3	*	*	*	0.2	0.3	0.5	0.4	0.4	0.6	0.7	*
2001.....	0.3	*	*	*	0.3	0.2	0.4	0.4	0.3	0.3	0.8	1.7
2000.....	0.3	*	*	*	0.3	0.3	0.4	0.4	0.4	0.3	0.7	*
1999.....	0.3	*	*	*	0.3	0.3	0.4	0.4	0.4	0.4	0.7	*

All races 3/, female

2004.....	0.1	*	*	*	*	*	*	0.1	*	0.2	0.3	*
2003.....	0.1	*	*	*	*	*	0.1	*	*	0.2	0.3	*
2002.....	0.1	*	*	*	*	*	*	0.1	0.2	0.2	0.3	*
2001.....	0.1	*	*	*	*	*	0.1	*	*	*	0.4	*
2000.....	0.1	*	*	*	*	*	*	0.1	0.2	0.2	0.3	*
1999.....	0.1	*	*	*	*	*	*	*	*	*	*	*

White, both sexes

2004.....	0.2	*	*	*	0.1	0.2	0.2	0.3	0.2	0.3	0.4	0.7
2003.....	0.2	*	*	*	0.2	0.2	0.3	0.3	0.3	0.3	0.4	0.5
2002.....	0.2	*	*	0.1	0.2	0.2	0.3	0.3	0.3	0.4	0.5	0.7
2001.....	0.2	*	*	*	0.2	0.2	0.2	0.2	0.2	0.2	0.4	0.8
2000.....	0.2	*	*	*	0.2	0.2	0.2	0.2	0.3	0.3	0.5	0.7
1999.....	0.2	*	*	*	0.2	0.2	0.2	0.2	0.2	0.2	0.4	*

White, male

2004.....	0.3	*	*	*	0.2	0.3	0.4	0.4	0.4	0.4	0.6	*
2003.....	0.3	*	*	*	0.3	0.3	0.4	0.4	0.4	0.5	0.6	*
2002.....	0.3	*	*	*	0.3	0.3	0.4	0.4	0.4	0.6	0.7	*
2001.....	0.3	*	*	*	0.3	0.2	0.4	0.4	0.3	0.4	0.6	1.7
2000.....	0.3	*	*	*	0.3	0.4	0.4	0.4	0.4	0.3	0.7	*
1999.....	0.3	*	*	*	0.3	0.3	0.4	0.4	0.4	0.3	0.7	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

All other and unspecified accidents and their sequelae (W20-W31,W35-W38,W41-W64,W92-W99,X20-X32,X39,X50-X59,Y86)

All races 3/, both sexes

2004.....	3.5	1.3	0.7	0.3	0.9	1.3	1.8	2.7	3.2	5.5	16.6	64.5
2003.....	3.7	1.1	0.7	0.3	1.0	1.2	1.9	2.5	3.2	6.0	17.6	73.5
2002.....	3.8	0.9	0.6	0.3	0.9	1.2	1.9	2.4	3.5	6.2	18.5	75.8
2001.....	4.0	1.3	0.8	0.3	1.0	1.2	1.9	2.7	3.5	6.4	20.1	83.4
2000.....	3.9	1.1	0.7	0.3	1.0	1.3	2.1	2.6	3.6	6.2	18.9	82.6
1999.....	4.3	1.2	0.8	0.4	1.0	1.4	2.0	2.7	3.7	7.0	22.1	92.8

All races 3/, male

2004.....	4.2	1.6	0.9	0.4	1.5	2.2	3.0	4.4	5.0	7.6	20.2	69.6
2003.....	4.3	1.1	0.8	0.4	1.6	2.0	3.1	4.1	5.2	8.6	21.6	80.4
2002.....	4.3	1.2	0.7	0.4	1.5	2.1	3.0	4.0	5.5	8.8	22.8	79.9
2001.....	4.5	1.6	0.9	0.4	1.6	2.1	3.1	4.4	5.3	8.7	24.8	92.4
2000.....	4.5	1.2	0.8	0.4	1.7	2.3	3.6	4.5	5.6	8.5	22.6	91.7
1999.....	4.8	1.4	1.0	0.5	1.7	2.3	3.4	4.4	5.8	9.6	27.3	107.0

All races 3/, female

2004.....	2.9	*	0.4	0.2	0.3	0.3	0.7	1.0	1.5	3.6	14.2	62.2
2003.....	3.1	1.1	0.5	0.1	0.3	0.4	0.6	1.0	1.4	3.8	14.9	70.5
2002.....	3.2	*	0.5	0.2	0.3	0.3	0.7	0.9	1.6	4.0	15.7	74.0
2001.....	3.5	1.1	0.7	0.2	0.3	0.3	0.7	1.1	1.8	4.4	17.1	79.6
2000.....	3.3	*	0.5	0.2	0.3	0.3	0.6	0.8	1.7	4.4	16.5	78.9
1999.....	3.7	*	0.5	0.3	0.3	0.4	0.7	1.0	1.8	4.8	18.7	87.1

White, both sexes

2004.....	3.8	1.2	0.7	0.3	1.0	1.3	1.9	2.7	3.1	5.5	16.9	67.6
2003.....	4.0	1.0	0.6	0.3	1.0	1.2	1.9	2.4	3.1	5.9	17.9	77.0
2002.....	4.1	0.8	0.5	0.3	1.0	1.3	1.9	2.4	3.4	6.0	18.7	79.1
2001.....	4.3	1.3	0.8	0.3	1.1	1.3	1.9	2.6	3.4	6.3	20.6	86.4
2000.....	4.2	1.0	0.6	0.3	1.0	1.4	2.0	2.5	3.4	6.1	19.1	86.2
1999.....	4.5	1.0	0.8	0.4	1.1	1.4	2.0	2.5	3.5	6.7	22.1	96.1

White, male

2004.....	4.4	1.5	1.0	0.4	1.6	2.3	3.1	4.5	4.8	7.7	20.3	72.6
2003.....	4.5	*	0.8	0.4	1.7	2.1	3.1	3.9	4.9	8.5	21.8	83.8
2002.....	4.5	*	0.6	0.4	1.6	2.2	3.0	4.0	5.3	8.5	22.8	82.5
2001.....	4.8	1.5	0.9	0.4	1.7	2.1	3.1	4.2	5.2	8.6	25.2	95.8
2000.....	4.6	*	0.7	0.4	1.8	2.4	3.5	4.1	5.2	8.3	22.4	95.0
1999.....	5.0	*	1.0	0.5	1.7	2.3	3.3	4.1	5.5	9.0	27.1	108.9

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other and unspecified accidents and their sequelae (W20-W31,W35-W38,W41-W64,W92-W99,X20-X32,X39,X50-X59,Y86)												
White, female												
2004.....	3.3	*	0.4	0.2	0.3	0.4	0.7	0.9	1.4	3.6	14.5	65.4
2003.....	3.5	*	0.5	0.1	0.3	0.4	0.6	0.9	1.4	3.8	15.3	74.0
2002.....	3.6	*	0.4	0.3	0.3	0.4	0.6	0.8	1.6	3.9	15.9	77.6
2001.....	3.9	*	0.6	0.2	0.3	0.3	0.6	1.0	1.7	4.4	17.6	82.4
2000.....	3.7	*	0.5	0.2	0.3	0.3	0.6	0.8	1.7	4.3	17.0	82.6
1999.....	4.1	*	0.5	0.3	0.4	0.4	0.7	0.9	1.6	4.7	18.8	91.0
Black, both sexes												
2004.....	2.5	*	*	0.3	0.7	1.4	2.0	3.2	4.3	6.1	16.6	38.4
2003.....	2.8	*	0.9	*	1.0	1.1	2.2	4.0	4.6	7.5	16.1	45.8
2002.....	2.8	*	1.0	0.4	0.7	1.2	2.4	3.1	4.7	8.0	18.5	46.9
2001.....	2.9	*	0.9	0.3	0.7	1.3	2.3	3.8	4.6	7.8	18.2	58.3
2000.....	3.0	*	0.9	0.3	0.9	1.3	2.7	3.9	5.4	8.0	19.1	51.0
1999.....	3.6	*	0.8	0.3	0.9	1.5	2.5	4.6	6.3	11.1	25.3	67.0
Black, male												
2004.....	3.5	*	*	*	1.1	2.4	3.4	5.4	7.3	9.0	23.9	42.7
2003.....	3.7	*	*	*	1.4	1.8	3.5	6.5	8.0	12.2	21.7	54.2
2002.....	3.7	*	*	0.7	1.1	2.0	3.6	5.0	8.3	11.6	25.2	57.9
2001.....	3.7	*	*	*	1.0	2.2	3.7	6.2	7.2	11.6	23.8	62.4
2000.....	4.4	*	*	*	1.6	2.4	4.9	7.5	9.9	11.3	29.7	69.2
1999.....	4.9	*	*	*	1.7	2.6	4.3	8.1	10.3	17.4	32.6	106.6
Black, female												
2004.....	1.7	*	*	*	*	*	0.8	1.3	1.9	4.1	12.4	36.6
2003.....	1.9	*	*	*	*	*	1.0	1.9	1.9	4.1	12.9	42.5
2002.....	2.0	*	*	*	*	*	1.3	1.5	1.8	5.5	14.6	42.7
2001.....	2.2	*	*	*	*	*	1.0	1.7	2.5	5.1	15.0	56.8
2000.....	1.8	*	*	*	*	*	0.9	0.9	1.7	5.7	13.1	44.2
1999.....	2.4	*	*	*	*	*	0.9	1.6	3.2	6.6	21.2	52.1

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)												
All races 3/, both sexes												
2004.....	11.0	0.7	10.3	12.7	15.0	16.6	13.8	12.3	16.3	16.4
2003.....	10.8	0.6	9.7	12.7	14.9	15.9	13.8	12.7	16.4	16.9
2002.....	11.0	0.6	9.9	12.6	15.3	15.7	13.6	13.5	17.7	18.0
2001 4/.....	10.8	0.7	9.9	12.8	14.7	15.2	13.1	13.3	17.4	17.5
2000.....	10.4	0.7	10.2	12.0	14.5	14.4	12.1	12.5	17.6	19.6
1999.....	10.5	0.6	10.1	12.7	14.3	13.9	12.2	13.4	18.1	19.3
All races 3/, male												
2004.....	17.7	0.9	16.8	20.4	23.0	24.8	22.1	22.6	34.8	45.0
2003.....	17.6	0.9	16.0	20.6	23.2	24.4	22.3	23.4	35.1	47.8
2002.....	17.9	0.9	16.5	20.5	23.7	24.4	22.2	24.7	38.1	50.7
2001 4/.....	17.6	1.0	16.6	21.0	23.1	23.4	21.1	24.6	37.8	51.1
2000.....	17.1	1.2	17.1	19.6	22.8	22.4	19.4	22.7	38.6	57.5
1999.....	17.1	0.9	16.8	20.7	22.4	21.5	19.8	24.7	38.8	57.1
All races 3/, female												
2004.....	4.6	0.5	3.6	4.7	7.1	8.6	6.1	3.8	3.9	3.6
2003.....	4.3	0.3	3.0	4.6	6.6	7.7	5.9	3.8	4.0	3.3
2002.....	4.3	0.3	2.9	4.6	6.9	7.4	5.7	4.1	4.2	3.8
2001 4/.....	4.1	0.3	2.9	4.4	6.4	7.2	5.7	3.9	4.0	3.4
2000.....	4.0	0.3	3.0	4.3	6.4	6.7	5.4	4.0	4.0	4.2
1999.....	4.0	0.3	3.0	4.6	6.4	6.6	5.2	4.1	4.7	4.1
White, both sexes												
2004.....	12.3	0.7	11.0	13.6	16.9	18.6	15.1	13.4	17.5	17.5
2003.....	12.1	0.6	10.2	13.7	16.6	17.9	15.2	13.8	17.6	18.1
2002.....	12.2	0.7	10.7	13.6	17.0	17.6	15.0	14.7	19.0	19.1
2001 4/.....	11.9	0.7	10.6	13.8	16.4	16.9	14.3	14.3	18.6	18.6
2000.....	11.5	0.8	10.7	12.8	16.1	15.9	13.3	13.5	18.8	21.0
1999.....	11.5	0.6	10.5	13.5	15.8	15.3	13.2	14.4	19.3	20.7
White, male												
2004.....	19.6	0.9	17.9	21.8	25.6	27.6	23.9	24.2	37.1	48.4
2003.....	19.5	0.9	16.9	21.8	25.7	27.3	24.3	25.2	37.5	51.4
2002.....	19.9	1.0	17.7	21.6	26.2	27.0	24.3	26.8	40.6	53.9
2001 4/.....	19.5	1.0	17.6	22.4	25.4	26.0	22.8	26.3	40.2	55.0
2000.....	18.8	1.2	17.9	20.7	24.9	24.5	21.2	24.3	41.1	61.6
1999.....	18.7	1.0	17.5	21.8	24.4	23.4	21.2	26.4	41.2	61.5

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)

White, female

2004.....	5.1	0.5	3.8	5.0	8.1	9.8	6.8	4.1	4.1	3.6
2003.....	4.7	0.3	3.1	5.1	7.5	8.7	6.5	4.1	4.2	3.3
2002.....	4.8	0.4	3.1	5.2	7.8	8.4	6.2	4.4	4.4	4.0
2001 4/.....	4.6	0.3	3.1	4.9	7.3	8.0	6.4	4.2	4.3	3.4
2000.....	4.4	0.3	3.1	4.6	7.2	7.5	6.0	4.4	4.1	4.5
1999.....	4.5	0.3	3.2	4.9	7.2	7.4	5.7	4.4	4.9	4.3

Black, both sexes

2004.....	5.2	0.7	7.2	9.4	6.6	5.8	5.7	4.4	6.0	*
2003.....	5.1	0.6	7.1	9.3	7.2	5.6	5.0	4.5	4.5	*
2002.....	5.1	0.5	6.5	9.3	7.6	5.9	4.8	4.9	5.5	5.9
2001 4/.....	5.3	0.6	7.1	9.1	7.3	6.2	5.2	5.5	5.7	*
2000.....	5.4	0.6	8.1	8.7	7.6	6.5	4.2	5.1	5.8	*
1999.....	5.4	0.4	8.1	9.2	7.6	5.7	4.9	5.8	5.7	*

Black, male

2004.....	9.0	0.9	12.2	16.2	11.3	9.7	10.8	9.8	15.0	*
2003.....	8.8	1.0	12.1	16.5	12.2	9.1	8.7	8.3	11.3	*
2002.....	9.1	0.8	11.3	17.3	13.0	10.2	8.4	9.7	13.8	*
2001 4/.....	9.2	0.9	13.0	16.5	12.3	9.9	9.4	10.7	13.5	*
2000.....	9.4	1.0	14.2	15.5	13.2	11.2	7.5	11.1	12.1	*
1999.....	9.6	0.6	14.3	16.8	13.3	10.0	9.1	11.8	13.8	*

Black, female

2004.....	1.8	*	2.2	3.2	2.5	2.5	1.6	*	*	*
2003.....	1.8	*	2.0	2.7	2.9	2.6	2.0	*	*	*
2002.....	1.5	*	1.7	2.0	2.9	2.2	1.9	*	*	*
2001 4/.....	1.7	*	1.3	2.4	2.9	3.0	1.8	*	*	*
2000.....	1.7	*	2.2	2.6	2.7	2.5	1.6	*	*	*
1999.....	1.6	*	1.9	2.3	2.6	2.0	1.6	*	*	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Intentional self-poisoning (suicide) by and exposure to drugs and other biological substances (X60-X64)

All races 3/, both sexes

2004.....	1.4	*	0.7	1.4	2.6	3.1	2.0	0.9	0.9	1.1
2003.....	1.3	*	0.5	1.4	2.5	2.9	1.8	1.0	0.9	0.9
2002.....	1.3	*	0.6	1.4	2.6	2.8	1.7	1.0	1.0	0.8
2001.....	1.2	*	0.6	1.3	2.4	2.7	1.5	1.0	0.8	0.9
2000.....	1.2	*	0.5	1.2	2.2	2.4	1.5	0.8	0.9	1.0
1999.....	1.1	*	0.5	1.2	2.2	2.3	1.4	1.0	1.1	0.8

All races 3/, male

2004.....	1.4	*	0.7	1.5	2.4	2.8	1.9	0.7	0.9	1.6
2003.....	1.3	*	0.6	1.4	2.5	2.7	1.6	0.8	0.7	1.7
2002.....	1.3	*	0.6	1.5	2.5	2.7	1.5	0.9	1.2	*
2001.....	1.2	*	0.5	1.3	2.6	2.4	1.2	1.0	0.7	*
2000.....	1.1	*	0.5	1.2	2.2	2.3	1.2	0.7	1.0	*
1999.....	1.1	*	0.5	1.1	2.3	2.2	1.2	1.0	0.9	*

All races 3/, female

2004.....	1.5	*	0.6	1.4	2.7	3.4	2.2	1.0	0.9	0.8
2003.....	1.4	*	0.5	1.4	2.4	3.1	2.0	1.1	1.0	*
2002.....	1.4	*	0.6	1.2	2.6	3.0	2.0	1.1	0.9	0.7
2001.....	1.3	*	0.6	1.2	2.3	2.9	1.7	0.9	0.8	0.6
2000.....	1.2	*	0.5	1.1	2.3	2.5	1.7	0.9	0.7	0.9
1999.....	1.2	*	0.5	1.2	2.2	2.4	1.6	0.9	1.2	*

White, both sexes

2004.....	1.6	*	0.7	1.6	3.0	3.6	2.2	1.0	1.0	1.1
2003.....	1.5	*	0.6	1.6	2.8	3.3	2.0	1.1	1.0	1.0
2002.....	1.5	*	0.7	1.5	2.9	3.2	2.0	1.1	1.1	0.9
2001.....	1.4	*	0.6	1.4	2.8	3.0	1.6	1.0	0.8	0.9
2000.....	1.3	*	0.6	1.3	2.5	2.7	1.7	0.9	0.9	1.1
1999.....	1.3	*	0.6	1.3	2.5	2.6	1.5	1.1	1.2	0.8

White, male

2004.....	1.6	*	0.8	1.7	2.8	3.2	2.0	0.8	1.0	1.7
2003.....	1.5	*	0.7	1.6	2.9	3.1	1.8	0.9	0.8	1.8
2002.....	1.5	*	0.6	1.6	2.8	3.0	1.7	0.9	1.2	*
2001.....	1.4	*	0.6	1.5	3.0	2.7	1.3	1.1	0.8	*
2000.....	1.2	*	0.6	1.4	2.5	2.5	1.4	0.7	1.1	*
1999.....	1.3	*	0.5	1.2	2.6	2.5	1.3	1.1	1.0	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Intentional self-poisoning (suicide) by and exposure to drugs and other biological substances (X60-X64)

White, female

2004.....	1.7	*	0.7	1.5	3.1	3.9	2.4	1.1	1.0	0.8
2003.....	1.6	*	0.5	1.6	2.7	3.5	2.2	1.2	1.2	*
2002.....	1.6	*	0.7	1.3	3.0	3.4	2.2	1.2	1.0	0.8
2001.....	1.5	*	0.7	1.3	2.6	3.3	1.9	1.0	0.9	*
2000.....	1.4	*	0.6	1.3	2.6	2.9	1.9	1.0	0.8	1.0
1999.....	1.3	*	0.6	1.4	2.5	2.7	1.7	1.0	1.3	*

Black, both sexes

2004.....	0.4	*	*	0.7	0.9	0.7	*	*	*	*
2003.....	0.4	*	*	0.6	1.1	1.0	*	*	*	*
2002.....	0.4	*	*	0.8	0.9	0.8	*	*	*	*
2001.....	0.4	*	*	0.5	1.0	0.9	*	*	*	*
2000.....	0.4	*	*	0.6	1.0	1.1	*	*	*	*
1999.....	0.4	*	*	0.7	0.9	0.7	*	*	*	*

Black, male

2004.....	0.4	*	*	*	*	*	*	*	*	*
2003.....	0.3	*	*	*	1.1	*	*	*	*	*
2002.....	0.4	*	*	1.0	1.0	*	*	*	*	*
2001.....	0.4	*	*	*	1.0	*	*	*	*	*
2000.....	0.5	*	*	*	0.9	1.3	*	*	*	*
1999.....	0.4	*	*	0.9	0.9	*	*	*	*	*

Black, female

2004.....	0.5	*	*	0.8	1.0	0.8	*	*	*	*
2003.....	0.5	*	*	0.7	1.1	1.4	*	*	*	*
2002.....	0.4	*	*	*	0.9	1.0	*	*	*	*
2001.....	0.4	*	*	*	1.0	1.0	*	*	*	*
2000.....	0.4	*	*	*	1.0	0.9	*	*	*	*
1999.....	0.4	*	*	*	0.9	*	*	*	*	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Intentional self-poisoning (suicide) by and exposure to other and unspecified solid or liquid substances and their vapors (X65-X66,X68-X69)

All races 3/, both sexes

2004.....	0.1	*	*	0.1	0.1	0.1	0.2	*	*	*
2003.....	0.1	*	*	0.1	0.1	0.1	0.1	*	*	*
2002.....	0.1	*	*	0.1	0.1	0.1	*	*	*	*
2001.....	0.1	*	*	0.1	0.1	0.1	0.1	*	*	*
2000.....	0.1	*	*	0.1	0.1	0.1	*	*	*	*
1999.....	0.1	*	*	0.1	0.1	0.1	0.1	*	*	*

All races 3/, male

2004.....	0.1	*	*	0.2	0.2	0.2	0.2	*	*	*
2003.....	0.1	*	*	0.1	0.2	0.2	*	*	*	*
2002.....	0.1	*	*	0.1	0.2	0.2	*	*	*	*
2001.....	0.1	*	*	0.1	0.2	0.1	*	*	*	*
2000.....	0.1	*	*	0.1	0.2	0.2	*	*	*	*
1999.....	0.1	*	*	0.1	0.2	0.2	0.2	*	*	*

All races 3/, female

2004.....	0.1	*	*	*	*	0.1	*	*	*	*
2003.....	0.0	*	*	*	*	0.1	*	*	*	*
2002.....	0.0	*	*	*	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	*	*	*	*

White, both sexes

2004.....	0.1	*	*	0.1	0.1	0.2	0.2	*	*	*
2003.....	0.1	*	*	0.1	0.1	0.1	0.1	*	*	*
2002.....	0.1	*	*	0.1	0.1	0.1	*	*	*	*
2001.....	0.1	*	*	0.1	0.1	0.1	*	*	*	*
2000.....	0.1	*	*	0.1	0.1	0.1	*	*	*	*
1999.....	0.1	*	*	0.1	0.1	0.1	0.1	*	*	*

White, male

2004.....	0.1	*	*	0.2	0.2	0.2	0.2	*	*	*
2003.....	0.1	*	*	0.2	0.2	0.2	*	*	*	*
2002.....	0.1	*	*	*	0.2	0.2	*	*	*	*
2001.....	0.1	*	*	*	0.2	0.1	*	*	*	*
2000.....	0.1	*	*	0.1	0.2	0.2	*	*	*	*
1999.....	0.1	*	*	0.1	0.2	0.2	*	*	*	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Intentional self-poisoning (suicide) by and exposure to other gases and vapors (X67)

All races 3/, both sexes

2004.....	0.5	*	0.2	0.5	0.8	0.9	0.6	0.3	0.5	0.6
2003.....	0.5	*	0.2	0.5	0.9	0.8	0.7	0.4	0.5	0.7
2002.....	0.5	*	0.2	0.5	0.8	1.0	0.6	0.4	0.7	0.8
2001.....	0.5	*	0.2	0.6	0.9	0.9	0.7	0.4	0.6	0.8
2000.....	0.5	*	0.2	0.6	0.9	0.9	0.7	0.4	0.6	0.8
1999.....	0.6	*	0.3	0.7	0.9	0.9	0.6	0.4	0.8	1.2

All races 3/, male

2004.....	0.7	*	0.3	0.8	1.3	1.5	1.0	0.5	0.9	*
2003.....	0.7	*	0.3	0.9	1.4	1.3	1.1	0.7	0.8	1.6
2002.....	0.8	*	0.3	0.9	1.4	1.5	1.0	0.6	1.2	2.2
2001.....	0.8	*	0.4	0.9	1.3	1.4	1.1	0.7	1.2	1.9
2000.....	0.8	*	0.4	1.0	1.4	1.5	1.0	0.6	1.2	2.3
1999.....	0.9	*	0.6	1.1	1.5	1.4	0.9	0.6	1.5	3.9

All races 3/, female

2004.....	0.2	*	*	0.2	0.3	0.4	0.2	*	0.3	*
2003.....	0.2	*	*	0.2	0.4	0.4	0.3	*	0.3	*
2002.....	0.2	*	*	0.2	0.3	0.4	0.2	0.2	0.3	*
2001.....	0.2	*	*	0.2	0.4	0.4	0.4	0.2	*	*
2000.....	0.2	*	*	0.2	0.4	0.4	0.3	0.2	*	*
1999.....	0.2	*	*	0.3	0.4	0.5	0.4	*	0.4	*

White, both sexes

2004.....	0.5	*	0.2	0.5	0.9	1.1	0.7	0.4	0.6	0.7
2003.....	0.6	*	0.2	0.6	1.0	1.0	0.7	0.5	0.6	0.8
2002.....	0.6	*	0.2	0.7	1.0	1.1	0.7	0.5	0.7	0.9
2001.....	0.6	*	0.3	0.7	1.0	1.1	0.8	0.5	0.7	0.9
2000.....	0.6	*	0.3	0.7	1.0	1.1	0.8	0.4	0.6	0.9
1999.....	0.6	*	0.4	0.8	1.1	1.1	0.7	0.4	0.9	1.3

White, male

2004.....	0.8	*	0.4	0.9	1.5	1.7	1.1	0.6	1.0	*
2003.....	0.9	*	0.3	1.1	1.6	1.5	1.2	0.8	0.9	1.8
2002.....	0.9	*	0.4	1.0	1.6	1.7	1.1	0.7	1.3	2.3
2001.....	0.9	*	0.5	1.1	1.5	1.6	1.2	0.8	1.3	2.1
2000.....	1.0	*	0.5	1.1	1.6	1.7	1.2	0.6	1.3	2.5
1999.....	1.0	*	0.7	1.3	1.7	1.6	1.1	0.7	1.6	4.4

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Intentional self-harm (suicide) by hanging, strangulation and suffocation (X70)

All races 3/, both sexes

2004.....	2.5	0.5	3.6	4.0	3.8	3.0	2.0	1.2	1.6	2.2
2003.....	2.3	0.4	3.3	3.7	3.5	2.7	1.8	1.2	1.6	2.1
2002.....	2.2	0.4	3.2	3.5	3.4	2.7	1.7	1.4	1.6	2.3
2001.....	2.2	0.4	3.1	3.5	3.4	2.4	1.5	1.3	1.7	2.1
2000.....	2.0	0.4	2.9	3.1	3.1	2.2	1.4	1.1	1.8	2.9
1999.....	1.9	0.3	2.5	3.3	3.0	2.0	1.4	1.4	1.9	2.8

All races 3/, male

2004.....	4.1	0.7	5.7	6.7	6.5	4.8	3.3	2.1	2.9	4.8
2003.....	3.8	0.6	5.3	6.2	5.9	4.5	3.0	2.2	3.1	5.0
2002.....	3.8	0.6	5.2	6.0	5.7	4.4	2.9	2.3	2.7	5.2
2001.....	3.7	0.6	5.1	5.9	5.8	4.2	2.5	2.3	3.1	5.2
2000.....	3.4	0.7	4.8	5.4	5.3	3.8	2.2	1.7	3.3	7.2
1999.....	3.3	0.5	4.1	5.6	5.2	3.3	2.3	2.2	3.1	6.6

All races 3/, female

2004.....	0.9	0.4	1.5	1.2	1.1	1.2	0.8	0.5	0.7	1.1
2003.....	0.8	0.2	1.2	1.1	1.0	0.9	0.7	0.4	0.7	0.9
2002.....	0.8	0.2	1.0	1.0	1.1	1.0	0.6	0.6	0.8	1.0
2001.....	0.7	0.2	0.9	0.9	1.0	0.7	0.6	0.5	0.8	0.8
2000.....	0.7	0.2	0.9	0.8	0.8	0.7	0.7	0.7	0.9	1.2
1999.....	0.7	0.1	0.9	0.9	0.8	0.7	0.5	0.7	1.0	1.3

White, both sexes

2004.....	2.6	0.5	3.8	4.2	4.2	3.2	2.0	1.2	1.6	2.3
2003.....	2.4	0.3	3.5	4.0	3.7	2.9	1.8	1.2	1.7	2.1
2002.....	2.4	0.4	3.5	3.7	3.7	2.9	1.8	1.3	1.6	2.3
2001.....	2.3	0.4	3.4	3.6	3.7	2.6	1.6	1.3	1.7	2.1
2000.....	2.1	0.4	3.1	3.3	3.3	2.4	1.5	1.1	1.8	2.9
1999.....	2.0	0.3	2.6	3.4	3.2	2.1	1.4	1.3	1.8	2.9

White, male

2004.....	4.4	0.6	6.0	7.1	7.1	5.1	3.3	2.1	2.9	4.9
2003.....	4.1	0.6	5.6	6.6	6.3	4.8	2.9	2.0	3.1	5.1
2002.....	4.0	0.6	5.8	6.3	6.1	4.7	3.1	2.3	2.8	5.2
2001.....	4.0	0.6	5.6	6.2	6.3	4.6	2.6	2.2	3.1	5.3
2000.....	3.6	0.6	5.1	5.7	5.6	4.0	2.3	1.7	3.2	7.1
1999.....	3.4	0.5	4.2	5.9	5.5	3.5	2.4	2.2	3.1	6.9

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Intentional self-harm (suicide) by discharge of firearms (X72-X74)

All races 3/, both sexes

2004.....	5.7	0.1	5.0	5.7	6.5	8.0	8.0	8.8	12.2	11.1
2003.....	5.8	0.2	5.0	6.0	6.6	8.0	8.3	9.3	12.4	11.9
2002.....	5.9	0.2	5.1	6.0	7.0	7.8	8.4	9.7	13.2	12.0
2001.....	5.9	0.2	5.3	6.5	6.7	7.7	8.2	9.6	13.1	12.2
2000.....	5.9	0.3	5.8	6.1	7.1	7.6	7.6	9.2	13.0	13.2
1999.....	5.9	0.3	6.0	6.5	6.9	7.3	7.8	9.7	13.1	12.6

All races 3/, male

2004.....	10.0	0.2	8.8	9.7	10.8	13.5	14.2	17.5	28.3	34.4
2003.....	10.4	0.3	8.9	10.4	11.2	13.8	14.9	18.4	28.7	36.6
2002.....	10.6	0.3	9.1	10.3	11.9	13.6	15.1	19.5	31.0	37.8
2001.....	10.6	0.3	9.4	11.2	11.4	13.3	14.6	19.2	30.7	38.9
2000.....	10.5	0.4	10.2	10.6	11.8	13.0	13.5	18.4	30.8	42.9
1999.....	10.6	0.4	10.5	11.2	11.5	12.5	14.0	19.5	31.1	41.6

All races 3/, female

2004.....	1.5	*	1.1	1.5	2.2	2.8	2.2	1.6	1.4	*
2003.....	1.4	*	1.0	1.4	2.1	2.5	2.2	1.6	1.5	0.9
2002.....	1.4	*	1.0	1.6	2.2	2.2	2.2	1.6	1.4	0.8
2001.....	1.5	0.1	1.0	1.6	2.1	2.4	2.4	1.7	1.5	1.0
2000.....	1.5	0.1	1.2	1.7	2.4	2.3	2.1	1.7	1.4	1.1
1999.....	1.5	0.1	1.2	1.8	2.3	2.3	2.1	1.7	1.4	0.9

White, both sexes

2004.....	6.5	0.2	5.4	6.1	7.4	9.2	9.0	9.7	13.3	12.1
2003.....	6.6	0.2	5.3	6.4	7.6	9.2	9.3	10.3	13.4	12.9
2002.....	6.7	0.2	5.5	6.5	7.9	8.9	9.4	10.8	14.2	12.9
2001.....	6.7	0.2	5.6	7.1	7.5	8.7	9.2	10.5	14.1	13.2
2000.....	6.6	0.3	6.1	6.6	7.9	8.5	8.3	10.1	14.1	14.3
1999.....	6.6	0.3	6.2	7.0	7.7	8.2	8.6	10.6	14.1	13.6

White, male

2004.....	11.3	0.3	9.4	10.3	12.1	15.3	15.8	19.1	30.4	37.5
2003.....	11.7	0.3	9.4	10.9	12.6	15.6	16.5	20.2	30.9	39.9
2002.....	11.9	0.4	9.6	10.8	13.3	15.2	16.7	21.4	33.3	40.5
2001.....	11.8	0.4	9.7	12.0	12.6	14.9	16.1	20.9	32.9	42.1
2000.....	11.6	0.5	10.6	11.2	13.1	14.4	14.7	19.8	33.3	46.5
1999.....	11.7	0.4	10.8	11.8	12.6	13.9	15.2	21.0	33.3	44.9

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Intentional self-harm (suicide) by discharge of firearms (X72-X74)												
White, female												
2004.....	1.7	*	1.1	1.7	2.6	3.2	2.5	1.8	1.6	*
2003.....	1.6	*	1.0	1.7	2.4	2.9	2.5	1.8	1.6	0.9
2002.....	1.6	*	1.0	1.9	2.5	2.6	2.5	1.7	1.5	0.9
2001.....	1.7	*	1.1	1.8	2.4	2.7	2.7	1.8	1.6	1.1
2000.....	1.7	*	1.2	1.8	2.7	2.6	2.4	1.9	1.5	1.2
1999.....	1.7	0.1	1.3	2.0	2.6	2.6	2.4	1.9	1.5	1.0
Black, both sexes												
2004.....	2.6	*	4.0	4.8	2.7	2.4	3.0	3.5	4.8	*
2003.....	2.6	*	4.3	5.1	2.6	2.5	2.7	3.3	4.0	*
2002.....	2.8	*	4.1	5.0	3.5	2.9	2.8	3.3	4.4	*
2001.....	2.9	*	4.9	4.9	3.4	2.8	2.8	3.7	5.2	*
2000.....	3.0	*	5.2	5.0	3.4	3.3	2.9	3.7	3.5	*
1999.....	3.1	*	5.6	5.0	3.5	3.0	3.4	4.0	4.7	*
Black, male												
2004.....	4.9	*	7.1	9.0	5.2	4.4	5.8	8.3	12.8	*
2003.....	5.0	*	7.8	10.0	4.8	4.7	5.1	6.6	9.9	*
2002.....	5.4	*	7.5	10.0	6.3	5.5	5.5	7.1	11.5	*
2001.....	5.5	*	9.2	9.4	6.3	4.8	5.7	7.7	12.4	*
2000.....	5.5	*	9.6	9.3	6.2	5.8	5.7	8.8	8.2	*
1999.....	5.9	*	10.3	9.6	6.7	5.6	7.0	8.7	12.0	*
Black, female												
2004.....	0.5	*	0.9	0.9	*	*	*	*	*	*
2003.....	0.5	*	0.7	*	0.7	*	*	*	*	*
2002.....	0.5	*	0.7	*	0.9	*	*	*	*	*
2001.....	0.6	*	*	0.9	0.9	1.0	*	*	*	*
2000.....	0.7	*	0.9	1.1	0.8	1.1	*	*	*	*
1999.....	0.6	*	1.0	0.8	0.7	*	*	*	*	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Intentional self-harm (suicide) by jumping from a high place (X80)

All races 3/, both sexes

2004.....	0.2	*	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.5
2003.....	0.2	*	0.2	0.3	0.4	0.3	0.3	0.2	0.3	0.6
2002.....	0.3	*	0.3	0.4	0.3	0.3	0.3	0.2	0.3	0.6
2001.....	0.2	*	0.2	0.3	0.3	0.3	0.3	0.3	0.4	*
2000.....	0.2	*	0.3	0.3	0.3	0.3	0.2	0.2	0.3	0.6
1999.....	0.2	*	0.3	0.3	0.4	0.3	0.2	0.3	0.4	0.5

All races 3/, male

2004.....	0.3	*	0.4	0.5	0.5	0.4	0.4	0.4	0.4	*
2003.....	0.4	*	0.4	0.5	0.5	0.4	0.4	0.3	0.4	*
2002.....	0.4	*	0.4	0.6	0.5	0.4	0.4	0.3	0.5	*
2001.....	0.3	*	0.3	0.4	0.4	0.4	0.4	0.3	0.6	*
2000.....	0.3	*	0.4	0.4	0.4	0.5	0.2	0.3	0.6	*
1999.....	0.4	*	0.4	0.5	0.6	0.5	0.2	0.4	0.7	*

All races 3/, female

2004.....	0.1	*	0.1	0.1	0.2	0.2	*	*	*	*
2003.....	0.1	*	*	0.2	0.2	0.2	0.2	*	*	*
2002.....	0.2	*	0.1	0.2	0.2	0.2	0.2	*	*	*
2001.....	0.1	*	0.1	0.1	0.2	0.2	0.2	0.2	*	*
2000.....	0.1	*	0.1	0.1	0.1	0.1	0.2	*	*	*
1999.....	0.1	*	0.1	0.1	0.2	0.2	0.2	*	*	*

White, both sexes

2004.....	0.2	*	0.3	0.3	0.3	0.3	0.3	0.3	0.3	*
2003.....	0.3	*	0.2	0.3	0.4	0.3	0.3	0.2	0.3	0.5
2002.....	0.3	*	0.3	0.4	0.4	0.3	0.3	0.2	0.3	0.6
2001.....	0.2	*	0.2	0.3	0.3	0.3	0.3	0.3	0.4	*
2000.....	0.2	*	0.3	0.2	0.3	0.3	0.2	0.2	0.3	0.5
1999.....	0.2	*	0.3	0.3	0.4	0.3	0.2	0.3	0.4	0.6

White, male

2004.....	0.3	*	0.4	0.4	0.5	0.4	0.4	0.4	0.4	*
2003.....	0.4	*	0.4	0.4	0.5	0.5	0.4	0.3	0.5	*
2002.....	0.4	*	0.4	0.5	0.5	0.4	0.4	0.3	0.5	*
2001.....	0.3	*	0.3	0.4	0.4	0.4	0.4	0.4	0.6	*
2000.....	0.3	*	0.4	0.3	0.4	0.5	0.3	0.3	0.5	*
1999.....	0.4	*	0.4	0.4	0.6	0.5	0.2	0.5	0.7	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Intentional self-harm (suicide) by all other and unspecified means and their sequelae (*U03,X71,X75-X79,X81-X84,X87.0)

All races 3/, both sexes

2004.....	0.6	*	0.5	0.6	0.9	1.1	0.8	0.7	0.7	0.9
2003.....	0.6	*	0.4	0.7	1.0	1.0	0.8	0.5	0.6	0.7
2002.....	0.6	*	0.5	0.7	1.0	1.0	0.8	0.7	0.8	1.4
2001 4/.....	0.6	*	0.5	0.7	0.9	1.0	0.8	0.6	0.8	1.0
2000.....	0.6	*	0.5	0.6	0.9	0.8	0.8	0.6	0.9	1.2
1999.....	0.6	*	0.5	0.7	0.9	0.9	0.6	0.7	0.8	1.3

All races 3/, male

2004.....	1.0	*	0.8	1.0	1.4	1.7	1.2	1.1	1.1	2.2
2003.....	0.9	*	0.6	1.1	1.5	1.5	1.2	0.9	1.2	1.7
2002.....	1.0	*	0.8	1.1	1.5	1.6	1.2	1.0	1.4	3.1
2001 4/.....	0.9	*	0.7	1.1	1.3	1.5	1.2	0.9	1.4	2.5
2000.....	0.9	*	0.7	0.9	1.4	1.2	1.2	1.0	1.6	2.4
1999.....	0.8	*	0.7	1.1	1.2	1.3	0.9	1.0	1.4	2.5

All races 3/, female

2004.....	0.3	*	0.2	0.3	0.5	0.6	0.5	0.4	0.4	*
2003.....	0.3	*	0.2	0.3	0.5	0.5	0.5	0.3	*	*
2002.....	0.3	*	0.2	0.3	0.4	0.4	0.5	0.4	0.5	0.7
2001 4/.....	0.3	*	0.2	0.3	0.4	0.5	0.4	0.4	0.4	*
2000.....	0.3	*	0.2	0.2	0.4	0.5	0.4	0.4	0.5	0.7
1999.....	0.3	*	0.2	0.2	0.5	0.5	0.3	0.5	0.4	0.8

White, both sexes

2004.....	0.7	*	0.5	0.7	1.0	1.2	0.8	0.8	0.6	0.9
2003.....	0.6	*	0.4	0.7	1.0	1.1	0.9	0.6	0.6	0.7
2002.....	0.7	*	0.5	0.7	1.0	1.1	0.9	0.7	0.9	1.5
2001 4/.....	0.6	*	0.5	0.7	0.9	1.0	0.8	0.6	0.9	1.1
2000.....	0.6	*	0.4	0.6	0.9	0.9	0.8	0.7	0.9	1.2
1999.....	0.6	*	0.5	0.6	0.9	0.9	0.6	0.7	0.8	1.4

White, male

2004.....	1.0	*	0.8	1.0	1.4	1.7	1.1	1.2	1.0	2.1
2003.....	1.0	*	0.5	1.1	1.5	1.6	1.3	0.9	1.3	1.8
2002.....	1.0	*	0.8	1.1	1.6	1.7	1.2	1.0	1.4	3.3
2001 4/.....	1.0	*	0.7	1.1	1.4	1.6	1.2	0.9	1.5	2.8
2000.....	0.9	*	0.7	0.9	1.4	1.2	1.2	1.0	1.6	2.7
1999.....	0.9	*	0.7	1.0	1.2	1.4	0.9	0.9	1.5	2.8

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)												
All races 3/, both sexes												
2004.....	5.9	8.0	2.4	0.8	12.2	11.2	6.8	4.8	3.0	2.4	2.2	2.1
2003.....	6.1	8.5	2.4	0.8	13.0	11.3	7.0	4.9	2.8	2.4	2.5	2.2
2002.....	6.1	7.5	2.7	0.9	12.9	11.2	7.2	4.8	3.2	2.3	2.3	2.1
2001 4/.....	7.1	8.2	2.7	0.8	13.3	13.1	9.5	6.3	4.0	2.9	2.5	2.4
2000.....	6.0	9.2	2.3	0.9	12.6	10.4	7.1	4.7	3.0	2.4	2.4	2.4
1999.....	6.1	8.7	2.5	1.1	12.9	10.5	7.1	4.6	3.0	2.6	2.5	2.4
All races 3/, male												
2004.....	9.4	8.0	2.5	1.0	20.4	18.5	10.1	7.3	4.6	3.6	2.6	2.7
2003.....	9.7	10.0	2.5	1.0	21.8	18.5	10.4	7.4	4.2	3.4	3.3	3.2
2002.....	9.6	7.9	2.9	0.9	21.5	18.2	10.7	7.1	4.8	3.3	3.1	3.0
2001 4/.....	11.1	9.5	3.0	0.9	22.2	20.8	13.9	9.5	5.9	4.0	3.1	3.2
2000.....	9.3	10.4	2.5	1.1	20.9	16.7	10.3	6.9	4.6	3.4	3.2	3.3
1999.....	9.3	9.6	2.5	1.2	21.1	16.5	10.3	6.8	4.7	3.7	3.4	3.4
All races 3/, female												
2004.....	2.5	7.9	2.2	0.6	3.5	3.7	3.4	2.4	1.6	1.4	1.9	1.9
2003.....	2.6	6.9	2.3	0.5	3.7	3.9	3.6	2.5	1.5	1.5	1.9	1.8
2002.....	2.7	7.1	2.5	0.8	3.8	4.2	3.8	2.6	1.6	1.4	1.8	1.7
2001 4/.....	3.3	6.9	2.4	0.7	3.9	5.3	5.1	3.2	2.2	2.0	2.1	2.0
2000.....	2.8	7.9	2.1	0.7	3.9	4.1	4.0	2.5	1.6	1.6	2.0	2.0
1999.....	2.9	7.8	2.4	0.9	4.3	4.4	4.0	2.4	1.5	1.7	1.9	2.0
White, both sexes												
2004.....	3.6	6.5	1.6	0.6	6.5	6.1	4.5	3.4	2.3	2.0	1.9	1.9
2003.....	3.7	6.9	1.7	0.5	6.7	6.1	4.6	3.5	2.2	1.9	2.1	2.0
2002.....	3.7	5.4	2.0	0.6	6.7	5.9	4.8	3.3	2.6	1.9	1.9	1.8
2001 4/.....	4.9	6.2	1.9	0.7	7.2	8.3	7.3	5.0	3.4	2.5	2.1	2.2
2000.....	3.6	6.6	1.7	0.7	6.4	5.7	4.7	3.3	2.4	1.9	2.0	2.2
1999.....	3.8	6.7	1.7	0.8	6.8	6.0	4.8	3.3	2.4	2.2	2.0	2.1
White, male												
2004.....	5.3	6.9	1.8	0.7	10.2	9.3	6.3	4.9	3.4	2.8	2.1	2.3
2003.....	5.4	8.1	1.5	0.6	10.6	9.2	6.3	5.0	3.1	2.7	2.8	3.0
2002.....	5.4	6.2	2.1	0.6	10.6	8.9	6.8	4.6	3.7	2.6	2.4	2.7
2001 4/.....	7.2	7.3	2.2	0.7	11.2	12.3	10.7	7.4	4.9	3.2	2.6	2.8
2000.....	5.2	8.2	1.9	0.9	9.9	8.4	6.5	4.6	3.5	2.4	2.5	2.8
1999.....	5.5	7.9	1.8	0.9	10.3	8.8	6.7	4.8	3.6	2.9	2.7	2.7

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)												
White, female												
2004.....	1.9	6.2	1.4	0.5	2.5	2.7	2.6	1.9	1.4	1.4	1.7	1.8
2003.....	2.0	5.7	1.8	0.3	2.5	2.8	2.8	2.1	1.4	1.3	1.7	1.6
2002.....	2.0	4.6	1.8	0.6	2.5	2.8	2.8	2.1	1.5	1.3	1.6	1.5
2001 4/.....	2.6	5.1	1.6	0.6	3.0	4.1	3.8	2.7	2.0	1.8	1.8	2.0
2000.....	2.1	5.0	1.5	0.6	2.7	2.9	2.8	2.1	1.4	1.5	1.8	1.9
1999.....	2.1	5.4	1.7	0.7	3.0	3.1	2.9	1.8	1.3	1.6	1.6	1.9
Black, both sexes												
2004.....	21.1	15.6	6.3	1.9	43.5	44.2	22.6	15.3	8.8	6.0	6.3	*
2003.....	22.0	16.1	6.1	2.2	47.6	45.2	23.4	15.4	8.1	6.2	6.6	*
2002.....	22.0	17.3	6.3	2.0	46.8	45.3	23.7	15.5	8.3	5.7	7.4	*
2001 4/.....	22.1	18.9	6.8	1.4	47.3	43.7	25.0	16.1	8.9	7.1	6.7	*
2000.....	21.5	22.8	5.8	1.8	47.8	40.8	23.9	14.8	8.3	7.6	7.0	*
1999.....	21.1	19.5	6.1	2.2	47.7	39.3	23.2	14.0	8.2	6.3	7.4	*
Black, male												
2004.....	37.1	14.2	6.3	2.4	77.6	81.6	37.9	26.0	15.1	11.5	10.3	*
2003.....	38.9	17.8	7.4	2.9	84.6	82.5	40.2	26.3	14.8	11.1	11.6	*
2002.....	38.4	16.3	7.1	2.5	83.1	82.2	38.8	27.1	15.4	10.6	13.8	*
2001 4/.....	38.3	21.0	8.0	2.0	85.7	78.8	38.5	26.9	15.0	12.6	10.0	*
2000.....	37.2	23.3	5.9	2.1	85.3	73.9	38.5	26.0	14.7	13.9	11.8	*
1999.....	36.1	19.5	5.9	2.7	84.7	69.2	37.2	23.4	15.1	11.8	12.0	*
Black, female												
2004.....	6.4	17.0	6.3	1.5	8.8	9.8	9.1	6.0	3.7	2.1	4.1	*
2003.....	6.6	14.2	4.8	1.5	10.1	11.1	8.5	6.0	2.7	2.7	3.8	*
2002.....	7.0	18.5	5.5	1.5	10.3	11.9	10.4	5.6	2.5	2.3	3.7	*
2001 4/.....	7.4	16.7	5.6	0.8	8.9	12.1	13.0	6.8	3.9	3.3	4.8	*
2000.....	7.2	22.2	5.7	1.6	10.7	10.9	11.0	5.3	3.3	3.1	4.3	*
1999.....	7.6	19.5	6.3	1.7	11.2	12.4	10.8	5.9	2.6	2.5	4.8	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Assault (homicide) by hanging, strangulation and suffocation (X91)

All races 3/, both sexes

2004.....	0.2	1.0	0.2	0.1	0.3	0.3	0.3	0.2	0.1	0.2	0.2	*
2003.....	0.2	1.0	*	0.1	0.3	0.3	0.3	0.2	0.1	0.2	0.3	*
2002.....	0.2	0.8	0.2	0.1	0.3	0.3	0.3	0.3	0.1	0.2	*	*
2001.....	0.2	1.0	*	*	0.3	0.3	0.3	0.2	0.2	0.3	0.2	*
2000.....	0.2	0.7	0.2	0.1	0.3	0.3	0.3	0.2	0.2	0.2	0.2	*
1999.....	0.3	0.9	0.1	0.1	0.3	0.4	0.3	0.2	0.1	0.2	0.2	*

All races 3/, male

2004.....	0.1	*	*	*	0.1	0.2	0.2	0.2	*	*	*	*
2003.....	0.2	1.0	*	*	0.1	0.1	0.2	0.1	*	*	*	*
2002.....	0.2	*	*	*	0.2	0.2	0.2	0.2	0.2	*	*	*
2001.....	0.2	1.2	*	*	0.2	0.2	0.2	0.2	0.2	0.3	*	*
2000.....	0.2	*	*	*	0.1	0.2	0.2	0.1	0.2	*	*	*
1999.....	0.2	1.1	*	*	0.1	0.2	0.2	0.1	*	0.3	*	*

All races 3/, female

2004.....	0.3	1.3	*	*	0.4	0.5	0.5	0.3	0.1	0.2	*	*
2003.....	0.3	*	*	*	0.4	0.5	0.4	0.3	0.2	0.2	*	*
2002.....	0.3	*	*	0.1	0.4	0.4	0.5	0.3	*	*	*	*
2001.....	0.3	*	*	*	0.4	0.5	0.5	0.3	0.2	0.2	0.3	*
2000.....	0.3	*	*	*	0.5	0.5	0.4	0.3	*	*	*	*
1999.....	0.3	*	*	0.1	0.5	0.6	0.5	0.2	*	0.2	0.3	*

White, both sexes

2004.....	0.2	0.9	*	*	0.2	0.2	0.2	0.2	0.1	0.2	0.2	*
2003.....	0.2	0.9	*	*	0.2	0.3	0.3	0.1	0.1	0.1	0.2	*
2002.....	0.2	0.6	*	0.1	0.2	0.2	0.2	0.2	0.1	0.1	*	*
2001.....	0.2	0.7	*	*	0.2	0.3	0.2	0.2	0.1	0.2	*	*
2000.....	0.2	*	0.2	0.1	0.2	0.3	0.2	0.2	0.1	0.1	*	*
1999.....	0.2	0.7	*	0.1	0.2	0.3	0.2	0.1	*	0.2	0.2	*

White, male

2004.....	0.1	*	*	*	*	0.1	0.1	0.1	*	*	*	*
2003.....	0.1	*	*	*	*	0.1	0.2	*	*	*	*	*
2002.....	0.1	*	*	*	0.1	0.1	0.1	0.2	*	*	0.1	*
2001.....	0.1	*	*	*	*	0.1	0.1	0.2	*	*	*	*
2000.....	0.1	*	*	*	*	0.2	0.2	0.1	*	*	*	*
1999.....	0.1	*	*	*	*	0.2	0.2	*	*	*	*	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Assault (homicide) by hanging, strangulation and suffocation (X91)

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
White, female												
2004.....	0.2	1.4	*	*	0.3	0.4	0.3	0.2	*	*	*	*
2003.....	0.2	*	*	*	0.3	0.4	0.3	0.2	*	*	*	*
2002.....	0.2	*	*	*	0.3	0.3	0.3	0.3	*	*	*	*
2001.....	0.2	*	*	*	0.4	0.4	0.3	0.2	*	*	*	*
2000.....	0.2	*	*	*	0.4	0.3	0.3	0.2	*	*	*	*
1999.....	0.2	*	*	*	0.4	0.4	0.3	0.2	*	*	*	*
Black, both sexes												
2004.....	0.5	*	*	*	0.5	0.7	1.1	0.5	*	*	*	*
2003.....	0.5	*	*	*	0.7	0.7	0.7	0.7	*	*	*	*
2002.....	0.6	*	*	*	0.6	0.9	1.0	0.7	*	*	*	*
2001.....	0.6	*	*	*	0.6	0.7	0.9	0.5	*	*	*	*
2000.....	0.6	*	*	*	0.7	0.9	0.9	*	*	*	*	*
1999.....	0.6	*	*	*	0.7	1.1	1.0	*	*	*	*	*
Black, male												
2004.....	0.3	*	*	*	*	*	*	*	*	*	*	*
2003.....	0.3	*	*	*	*	*	*	*	*	*	*	*
2002.....	0.4	*	*	*	*	*	*	*	*	*	*	*
2001.....	0.4	*	*	*	*	*	*	*	*	*	*	*
2000.....	0.3	*	*	*	*	*	*	*	*	*	*	*
1999.....	0.4	*	*	*	*	*	*	*	*	*	*	*
Black, female												
2004.....	0.7	*	*	*	0.8	1.1	1.5	*	*	*	*	*
2003.....	0.7	*	*	*	1.1	1.0	0.9	0.9	*	*	*	*
2002.....	0.7	*	*	*	0.8	1.3	1.5	*	*	*	*	*
2001.....	0.7	*	*	*	0.9	1.0	1.3	*	*	*	*	*
2000.....	0.8	*	*	*	1.1	1.4	1.3	*	*	*	*	*
1999.....	0.8	*	*	*	1.0	1.6	1.4	*	*	*	*	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Assault (homicide) by discharge of firearms (*U01.4,X93-X95)

All races 3/, both sexes

2004.....	4.0	*	0.2	0.5	9.9	8.8	4.3	2.6	1.6	1.0	0.8	0.6
2003.....	4.1	*	0.3	0.5	10.7	8.9	4.4	2.7	1.4	0.9	0.7	0.5
2002.....	4.1	*	0.3	0.5	10.6	8.7	4.5	2.6	1.5	0.9	0.9	*
2001.....	4.0	*	0.4	0.4	10.5	8.4	4.4	2.4	1.4	1.0	0.8	*
2000.....	3.8	*	0.2	0.5	10.1	7.7	4.2	2.5	1.5	1.0	0.7	0.7
1999.....	3.9	*	0.3	0.5	10.4	7.6	4.1	2.4	1.6	1.0	0.8	0.6

All races 3/, male

2004.....	6.9	*	*	0.6	17.3	15.3	7.1	4.0	2.5	1.6	0.8	*
2003.....	7.1	*	0.2	0.6	18.8	15.5	6.9	4.3	2.2	1.4	0.7	*
2002.....	7.0	*	0.4	0.6	18.5	14.9	7.3	3.9	2.4	1.2	1.3	*
2001.....	6.8	*	0.3	0.6	18.5	14.3	6.8	3.8	2.2	1.5	1.1	*
2000.....	6.5	*	*	0.7	17.7	13.2	6.6	3.9	2.3	1.3	0.8	*
1999.....	6.5	*	0.3	0.7	18.0	12.9	6.3	3.8	2.6	1.5	1.0	*

All races 3/, female

2004.....	1.1	*	*	0.3	2.0	2.0	1.5	1.1	0.8	0.5	0.8	*
2003.....	1.2	*	0.3	0.3	2.2	2.1	1.8	1.2	0.7	0.5	0.7	*
2002.....	1.3	*	*	0.4	2.4	2.3	1.9	1.3	0.8	0.6	0.7	*
2001.....	1.3	*	0.4	0.3	2.1	2.3	2.0	1.0	0.8	0.7	0.6	*
2000.....	1.3	*	*	0.2	2.2	2.1	1.8	1.2	0.8	0.8	0.6	*
1999.....	1.3	*	0.3	0.3	2.5	2.2	2.0	1.2	0.6	0.7	0.6	*

White, both sexes

2004.....	2.1	*	*	0.3	4.7	4.3	2.7	1.8	1.3	0.9	0.7	0.5
2003.....	2.2	*	0.2	0.3	5.0	4.2	2.7	1.9	1.1	0.7	0.6	0.5
2002.....	2.2	*	0.3	0.3	5.0	4.1	2.8	1.8	1.2	0.7	0.8	*
2001.....	2.2	*	0.3	0.4	5.0	4.2	2.9	1.7	1.2	0.9	0.7	*
2000.....	2.1	*	*	0.4	4.6	3.7	2.7	1.8	1.3	0.8	0.6	0.6
1999.....	2.2	*	0.2	0.4	4.9	3.8	2.7	1.7	1.2	0.9	0.7	0.6

White, male

2004.....	3.5	*	*	0.4	7.8	7.0	4.1	2.6	1.8	1.2	0.7	*
2003.....	3.5	*	*	0.4	8.5	6.9	3.9	2.8	1.6	1.1	0.5	*
2002.....	3.5	*	0.3	0.4	8.4	6.6	4.2	2.5	1.8	0.9	1.0	*
2001.....	3.5	*	*	0.5	8.3	6.7	4.3	2.6	1.6	1.1	0.8	*
2000.....	3.2	*	*	0.5	7.7	5.9	3.9	2.6	1.8	1.0	0.6	*
1999.....	3.4	*	*	0.5	8.0	6.1	4.0	2.6	1.9	1.2	0.7	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Assault (homicide) by discharge of firearms (*U01.4,X93-X95)

White, female

2004.....	0.9	*	*	0.2	1.3	1.3	1.2	0.9	0.7	0.6	0.7	*
2003.....	0.9	*	*	0.1	1.3	1.4	1.5	1.0	0.7	0.4	0.6	*
2002.....	1.0	*	*	0.3	1.4	1.5	1.4	1.1	0.7	0.6	0.6	*
2001.....	1.0	*	*	0.3	1.5	1.6	1.5	0.9	0.8	0.7	0.6	*
2000.....	1.0	*	*	0.2	1.4	1.5	1.4	1.0	0.8	0.7	0.6	*
1999.....	0.9	*	*	0.3	1.5	1.5	1.5	0.9	0.6	0.7	0.6	*

Black, both sexes

2004.....	16.1	*	0.8	1.2	38.8	37.7	15.6	8.4	4.8	2.4	*	*
2003.....	16.8	*	*	1.4	41.9	38.8	15.9	8.7	3.9	2.5	*	*
2002.....	16.7	*	*	1.2	41.8	38.2	16.3	8.5	4.1	2.4	2.2	*
2001.....	15.8	*	*	0.8	41.8	35.1	15.0	7.5	3.7	2.7	*	*
2000.....	15.6	*	*	1.0	41.2	33.2	15.0	8.4	3.5	2.7	*	*
1999.....	15.4	*	*	1.2	42.0	31.4	14.1	8.0	4.4	2.4	2.3	*

Black, male

2004.....	30.3	*	*	1.8	71.3	72.2	28.9	15.2	8.6	5.4	*	*
2003.....	31.6	*	*	1.9	76.8	74.1	29.1	15.9	7.5	4.6	*	*
2002.....	31.0	*	*	1.6	76.1	72.8	29.2	15.1	7.6	4.7	*	*
2001.....	29.8	*	*	1.3	78.2	66.9	26.0	13.7	7.1	5.6	*	*
2000.....	29.2	*	*	1.5	76.5	63.3	26.7	15.1	6.3	5.4	*	*
1999.....	28.4	*	*	1.8	76.9	59.1	24.0	13.9	8.9	4.6	*	*

Black, female

2004.....	3.0	*	*	*	5.8	6.1	3.8	2.5	1.7	*	*	*
2003.....	3.3	*	*	0.8	6.5	6.6	4.2	2.5	*	*	*	*
2002.....	3.6	*	*	0.8	7.3	6.9	4.8	2.8	1.4	*	*	*
2001.....	3.1	*	*	*	5.4	6.3	5.4	2.2	*	*	*	*
2000.....	3.2	*	*	*	6.4	6.1	4.6	2.6	*	*	*	*
1999.....	3.7	*	*	0.6	7.6	6.5	5.3	3.0	*	*	*	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Assault (homicide) by sharp object (X99)												
All races 3/, both sexes												
2004.....	0.7	*	*	0.1	1.2	1.2	1.0	0.8	0.4	0.4	0.4	*
2003.....	0.7	*	*	0.1	1.1	1.2	1.1	0.8	0.4	0.4	0.4	*
2002.....	0.7	*	*	0.1	1.0	1.2	1.1	0.8	0.5	0.4	0.4	*
2001.....	0.7	*	*	0.1	1.2	1.2	1.0	0.7	0.4	0.4	0.4	*
2000.....	0.6	*	*	0.1	1.0	1.1	1.0	0.6	0.4	0.3	0.5	*
1999.....	0.7	*	*	0.1	1.1	1.2	1.0	0.7	0.4	0.3	0.3	*
All races 3/, male												
2004.....	1.0	*	*	*	1.8	1.7	1.4	1.1	0.7	0.6	0.6	*
2003.....	1.0	*	*	*	1.7	1.7	1.5	1.2	0.6	0.5	0.6	*
2002.....	1.0	*	*	0.1	1.6	1.7	1.5	1.2	0.8	0.7	0.4	*
2001.....	1.0	*	*	*	1.8	1.7	1.3	1.0	0.6	0.5	0.4	*
2000.....	0.9	*	*	*	1.6	1.5	1.4	0.8	0.7	0.5	0.7	*
1999.....	1.0	*	*	*	1.6	1.7	1.6	1.0	0.7	0.5	0.5	*
All races 3/, female												
2004.....	0.4	*	*	*	0.5	0.7	0.6	0.5	0.3	0.2	0.3	*
2003.....	0.4	*	*	*	0.5	0.7	0.6	0.4	0.3	0.3	0.3	*
2002.....	0.4	*	*	*	0.4	0.8	0.7	0.4	0.3	0.3	0.3	*
2001.....	0.4	*	*	*	0.5	0.7	0.7	0.3	0.3	0.4	0.4	*
2000.....	0.4	*	*	*	0.4	0.6	0.7	0.4	0.2	*	0.3	*
1999.....	0.4	*	*	*	0.5	0.6	0.5	0.4	0.3	0.2	*	*
White, both sexes												
2004.....	0.5	*	*	0.1	0.9	0.8	0.7	0.5	0.3	0.3	0.3	*
2003.....	0.5	*	*	*	0.7	0.8	0.7	0.5	0.3	0.3	0.3	*
2002.....	0.5	*	*	0.1	0.7	0.8	0.8	0.5	0.4	0.4	0.3	*
2001.....	0.5	*	*	*	0.9	0.8	0.6	0.4	0.3	0.3	0.3	*
2000.....	0.4	*	*	0.1	0.8	0.8	0.7	0.4	0.3	0.2	0.4	*
1999.....	0.5	*	*	0.1	0.7	0.8	0.6	0.5	0.3	0.3	0.2	*
White, male												
2004.....	0.7	*	*	*	1.3	1.1	0.9	0.7	0.4	0.4	*	*
2003.....	0.7	*	*	*	1.1	1.2	0.9	0.7	0.4	0.4	0.4	*
2002.....	0.7	*	*	*	1.2	1.1	1.1	0.7	0.5	0.5	*	*
2001.....	0.7	*	*	*	1.3	1.2	0.8	0.6	0.4	0.4	*	*
2000.....	0.6	*	*	*	1.2	1.1	0.9	0.5	0.5	0.4	0.6	*
1999.....	0.7	*	*	*	1.1	1.2	0.9	0.7	0.4	0.3	*	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Assault (homicide) by sharp object (X99)												
White, female												
2004.....	0.3	*	*	*	0.4	0.6	0.4	0.3	0.2	*	*	*
2003.....	0.3	*	*	*	0.3	0.5	0.4	0.3	0.2	0.3	*	*
2002.....	0.3	*	*	*	0.3	0.5	0.4	0.3	0.3	0.2	*	*
2001.....	0.3	*	*	*	0.3	0.4	0.4	0.3	0.2	0.3	0.3	*
2000.....	0.3	*	*	*	0.3	0.4	0.4	0.3	*	*	*	*
1999.....	0.3	*	*	*	0.4	0.4	0.4	0.2	0.3	*	*	*
Black, both sexes												
2004.....	2.1	*	*	*	2.7	3.4	3.3	3.0	1.6	1.6	2.4	*
2003.....	2.2	*	*	*	3.3	3.3	3.9	2.8	1.4	1.2	*	*
2002.....	2.2	*	*	*	2.6	3.8	3.5	2.9	1.8	1.2	*	*
2001.....	2.0	*	*	*	2.9	3.5	3.5	2.4	1.3	1.2	*	*
2000.....	1.9	*	*	*	2.5	2.8	3.6	2.3	1.6	1.5	*	*
1999.....	2.0	*	*	*	2.8	3.1	3.6	2.4	1.4	*	*	*
Black, male												
2004.....	3.2	*	*	*	4.0	5.4	4.7	4.7	2.8	*	*	*
2003.....	3.4	*	*	*	5.0	5.0	6.1	4.6	2.4	*	*	*
2002.....	3.2	*	*	*	4.2	5.6	4.7	5.0	3.2	*	*	*
2001.....	3.0	*	*	*	4.3	5.2	5.0	4.4	1.9	*	*	*
2000.....	2.8	*	*	*	3.8	4.4	4.8	3.7	2.5	*	*	*
1999.....	3.1	*	*	*	4.1	4.6	6.2	3.7	2.6	*	*	*
Black, female												
2004.....	1.1	*	*	*	1.3	1.5	2.1	1.6	*	*	*	*
2003.....	1.1	*	*	*	1.6	1.8	2.0	1.1	*	*	*	*
2002.....	1.2	*	*	*	1.1	2.1	2.4	1.0	*	*	*	*
2001.....	1.1	*	*	*	1.5	1.9	2.2	*	*	*	*	*
2000.....	1.1	*	*	*	1.3	1.3	2.5	1.1	*	*	*	*
1999.....	1.0	*	*	*	1.5	1.8	1.4	1.3	*	*	*	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Assault (homicide) by blunt object (Y00)

All races 3/, both sexes

2004.....	0.0	*	*	*	*	*	*	0.0	*	*	*	*
2003.....	0.0	*	*	*	*	*	0.0	0.1	*	*	*	*
2002.....	0.1	*	*	*	0.1	0.1	0.1	0.1	*	*	*	*
2001.....	0.1	*	*	*	*	0.1	0.1	0.1	0.1	*	*	*
2000.....	0.1	*	*	*	0.1	0.1	0.1	0.1	*	*	*	*
1999.....	0.1	*	*	*	0.1	0.1	0.1	0.1	0.1	0.1	*	*

All races 3/, male

2004.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2003.....	0.1	*	*	*	*	*	*	0.1	*	*	*	*
2002.....	0.1	*	*	*	0.1	0.1	0.1	0.1	*	*	*	*
2001.....	0.1	*	*	*	*	*	0.1	0.2	*	*	*	*
2000.....	0.1	*	*	*	0.1	*	0.2	0.2	*	*	*	*
1999.....	0.1	*	*	*	0.1	0.1	0.2	0.1	*	*	*	*

All races 3/, female

2004.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2003.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*	*	*	*	*	*
1999.....	0.1	*	*	*	*	0.1	*	*	*	*	*	*

White, both sexes

2004.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2003.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2002.....	0.1	*	*	*	*	*	0.1	0.1	*	*	*	*
2001.....	0.1	*	*	*	*	*	0.1	0.1	*	*	*	*
2000.....	0.1	*	*	*	*	*	0.1	0.1	*	*	*	*
1999.....	0.1	*	*	*	0.1	0.1	0.1	0.1	0.1	*	*	*

White, male

2004.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2003.....	0.1	*	*	*	*	*	*	*	*	*	*	*
2002.....	0.1	*	*	*	*	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	0.1	0.1	*	*	*	*
2000.....	0.1	*	*	*	*	*	*	0.2	*	*	*	*
1999.....	0.1	*	*	*	*	*	0.1	*	*	*	*	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Assault (homicide) by bodily force (Y04)

All races 3/, both sexes

2004.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2003.....	0.0	*	*	*	*	*	*	0.1	*	*	*	*
2002.....	0.0	*	*	*	0.0	0.1	*	0.1	*	*	*	*
2001.....	0.0	*	*	*	0.1	0.1	0.1	0.1	*	*	*	*
2000.....	0.0	*	*	*	0.1	0.1	0.1	0.1	*	*	*	*
1999.....	0.1	*	*	*	0.1	0.1	0.1	0.1	*	*	*	*

All races 3/, male

2004.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2003.....	0.1	*	*	*	*	*	*	0.1	*	*	*	*
2002.....	0.1	*	*	*	*	*	*	0.1	*	*	*	*
2001.....	0.1	*	*	*	0.1	*	0.1	0.1	*	*	*	*
2000.....	0.1	*	*	*	*	0.2	0.2	0.1	*	*	*	*
1999.....	0.1	*	*	*	0.1	0.1	0.2	0.1	*	*	*	*

All races 3/, female

2004.....	*	*	*	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	*	*	*	*	*	*

White, both sexes

2004.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2003.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	0.1	*	*	*	*	*
2000.....	0.0	*	*	*	*	0.1	0.1	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	0.1	0.1	*	*	*	*

White, male

2004.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2003.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*	*	*	*	*	*
2000.....	0.1	*	*	*	*	0.1	0.1	*	*	*	*	*
1999.....	0.1	*	*	*	*	*	0.1	*	*	*	*	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Assault (homicide) by all other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5,*U01.8-*U01.9,*U02,X92,X96-X98,Y01-Y03,Y08-Y09,Y87.1)

All races 3/, both sexes

2004.....	0.9	3.8	1.3	0.1	0.8	0.9	1.0	1.2	0.8	0.7	0.7	0.9
2003.....	0.9	4.4	1.3	0.2	0.8	0.8	1.1	1.1	0.7	0.7	0.9	1.1
2002.....	0.9	3.7	1.4	0.1	0.7	0.9	1.0	1.0	0.8	0.7	0.7	0.9
2001 4/.....	2.0	3.5	1.6	0.2	1.1	3.1	3.5	2.8	1.8	1.0	0.9	1.1
2000.....	1.0	4.3	1.3	0.2	1.0	1.1	1.3	1.1	0.8	0.7	1.0	0.9
1999.....	1.0	3.8	1.3	0.3	0.9	1.1	1.3	1.1	0.8	0.7	1.0	1.1

All races 3/, male

2004.....	1.2	3.8	1.3	0.1	1.0	1.3	1.3	1.8	1.2	1.1	1.0	*
2003.....	1.2	4.7	1.5	0.2	1.1	1.1	1.5	1.6	1.2	1.1	1.5	2.0
2002.....	1.2	4.0	1.6	0.1	1.0	1.1	1.4	1.5	1.4	1.2	1.1	*
2001 4/.....	2.9	4.0	1.7	0.2	1.5	4.4	5.3	4.2	2.8	1.6	1.2	1.6
2000.....	1.4	4.7	1.5	0.2	1.3	1.5	1.8	1.7	1.3	1.2	1.4	*
1999.....	1.3	4.0	1.4	0.3	1.2	1.4	1.8	1.6	1.2	1.1	1.6	*

All races 3/, female

2004.....	0.6	3.8	1.3	0.2	0.5	0.5	0.7	0.5	0.4	0.4	0.5	0.8
2003.....	0.6	4.0	1.2	0.1	0.5	0.6	0.7	0.6	0.3	0.4	0.6	0.7
2002.....	0.5	3.3	1.2	0.2	0.5	0.6	0.6	0.5	0.3	0.4	0.5	0.8
2001 4/.....	1.1	3.1	1.4	0.2	0.8	1.8	1.7	1.4	1.0	0.5	0.7	0.9
2000.....	0.7	3.9	1.1	0.3	0.7	0.8	0.9	0.6	0.4	0.3	0.8	0.7
1999.....	0.7	3.6	1.2	0.3	0.7	0.8	0.9	0.5	0.4	0.4	0.6	0.9

White, both sexes

2004.....	0.7	3.1	0.9	0.1	0.6	0.7	0.8	0.9	0.6	0.6	0.6	0.9
2003.....	0.7	3.5	0.9	0.1	0.7	0.6	0.9	0.8	0.6	0.6	0.9	1.1
2002.....	0.7	2.5	0.9	0.1	0.6	0.6	0.8	0.7	0.7	0.6	0.6	0.7
2001 4/.....	1.8	2.7	1.1	0.2	1.0	2.9	3.3	2.5	1.7	0.8	0.8	1.0
2000.....	0.7	3.0	0.8	0.2	0.7	0.8	0.9	0.8	0.6	0.6	0.8	0.8
1999.....	0.8	2.9	1.0	0.2	0.8	0.8	1.0	0.8	0.6	0.6	0.8	0.9

White, male

2004.....	0.9	3.5	1.0	*	0.8	0.9	1.0	1.4	0.9	0.9	0.8	*
2003.....	0.9	3.8	0.9	*	0.9	0.8	1.2	1.2	1.0	0.9	1.3	1.9
2002.....	0.9	2.9	1.1	*	0.7	0.9	1.1	1.0	1.1	0.9	0.8	*
2001 4/.....	2.6	3.1	1.2	0.2	1.2	4.2	5.2	3.8	2.5	1.3	1.0	*
2000.....	0.9	3.5	1.0	0.2	0.8	1.0	1.3	1.1	0.9	0.9	1.1	*
1999.....	1.0	3.1	1.1	0.2	0.9	1.0	1.3	1.2	1.0	0.9	1.4	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Assault (homicide) by all other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5,*U01.8-*U01.9,*U02,X92,X96-X98,Y01-Y03,Y08-Y09,Y87.1)

White, female

2004.....	0.5	2.6	0.8	*	0.5	0.4	0.5	0.4	0.3	0.4	0.5	0.8
2003.....	0.5	3.1	0.9	*	0.4	0.5	0.6	0.5	0.3	0.3	0.5	0.7
2002.....	0.4	2.2	0.8	0.1	0.4	0.4	0.5	0.5	0.3	0.3	0.5	0.7
2001 4/.....	1.0	2.3	0.9	0.2	0.7	1.6	1.4	1.2	0.9	0.4	0.6	0.9
2000.....	0.5	2.4	0.7	0.2	0.5	0.6	0.6	0.5	0.3	0.3	0.7	*
1999.....	0.5	2.5	0.9	0.2	0.6	0.6	0.6	0.5	0.3	0.4	0.5	0.9

Black, both sexes

2004.....	2.0	7.7	3.7	0.4	1.4	2.2	2.4	3.0	1.9	1.4	2.0	*
2003.....	2.1	8.8	3.6	0.4	1.5	2.1	2.7	2.9	2.1	1.8	2.1	*
2002.....	2.1	9.2	3.8	0.4	1.5	2.1	2.5	3.1	2.0	1.6	2.6	*
2001 4/.....	3.1	8.0	4.1	0.3	1.7	4.1	5.0	5.1	3.0	2.4	2.2	*
2000.....	2.9	11.2	3.8	0.6	3.0	3.3	3.8	3.5	2.5	2.2	3.4	*
1999.....	2.5	8.9	3.2	0.6	1.9	3.0	3.9	2.9	1.6	2.1	2.3	*

Black, male

2004.....	2.8	5.9	3.4	*	1.8	3.5	3.3	5.2	3.2	2.8	*	*
2003.....	3.1	8.6	4.5	0.6	2.0	2.9	4.1	4.7	4.1	3.5	*	*
2002.....	3.1	9.3	4.3	*	2.2	2.9	3.7	5.4	4.0	3.0	6.0	*
2001 4/.....	4.3	8.4	4.8	*	2.5	5.8	6.5	7.4	5.0	4.2	*	*
2000.....	4.2	11.2	3.9	*	4.3	5.0	11.2	6.3	4.8	4.4	6.1	*
1999.....	3.4	9.1	3.6	0.6	2.9	4.0	5.4	4.9	2.5	4.1	*	*

Black, female

2004.....	1.3	9.6	4.0	*	0.9	0.9	1.5	1.2	*	*	*	*
2003.....	1.2	9.0	2.6	*	1.0	1.5	1.4	1.3	*	*	*	*
2002.....	1.2	9.1	3.3	*	0.9	1.5	1.5	1.1	*	*	*	*
2001 4/.....	2.1	7.6	3.3	*	1.0	2.6	3.8	3.0	1.4	*	*	*
2000.....	1.7	11.2	3.7	0.8	1.8	1.8	2.4	1.0	*	*	*	*
1999.....	1.6	8.7	2.9	0.6	0.9	2.1	2.5	1.1	*	*	*	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Event of undetermined intent (Y10-Y34,Y87.2,Y89.9)												
All races 3/, both sexes												
2004.....	1.7	2.6	0.3	0.2	1.2	2.0	3.1	3.1	1.4	0.8	0.9	1.6
2003.....	1.7	1.9	0.2	0.1	1.4	2.1	3.6	3.1	1.3	0.7	0.7	1.1
2002.....	1.7	2.5	0.4	0.2	1.2	2.1	3.4	2.8	1.2	0.8	0.7	1.4
2001.....	1.5	2.1	0.3	0.1	1.0	1.8	3.1	2.5	0.9	0.5	0.9	1.0
2000.....	1.4	1.7	0.3	0.1	0.9	1.7	2.8	2.2	0.9	0.6	0.6	1.3
1999.....	1.4	1.4	0.3	0.1	1.1	1.8	2.9	2.2	0.9	0.6	0.8	1.4
All races 3/, male												
2004.....	2.2	3.1	0.3	0.2	1.8	2.7	4.0	4.0	1.7	1.1	1.0	1.9
2003.....	2.3	2.4	*	0.2	1.9	2.9	4.6	4.0	1.7	0.8	1.0	1.5
2002.....	2.2	2.5	0.4	0.2	1.7	2.9	4.4	3.5	1.6	1.0	0.9	2.2
2001.....	2.0	2.4	*	0.2	1.5	2.6	4.2	3.4	1.2	0.8	1.1	1.5
2000.....	1.9	1.4	0.3	0.2	1.4	2.4	3.9	3.1	1.3	0.8	1.0	1.7
1999.....	2.0	1.7	0.3	0.2	1.7	2.6	4.0	3.0	1.3	0.9	1.1	2.1
All races 3/, female												
2004.....	1.2	2.0	0.3	0.1	0.6	1.3	2.3	2.3	1.1	0.6	0.9	1.6
2003.....	1.2	1.3	*	*	0.8	1.3	2.6	2.2	1.0	0.7	0.6	0.9
2002.....	1.2	2.5	0.4	0.1	0.7	1.2	2.4	2.2	0.8	0.6	0.6	1.1
2001.....	0.9	1.7	0.3	*	0.5	1.1	2.1	1.6	0.6	0.3	0.7	0.8
2000.....	0.8	2.0	*	*	0.4	1.0	1.8	1.4	0.6	0.3	0.4	1.2
1999.....	0.9	1.2	0.3	*	0.5	1.1	1.8	1.4	0.6	0.4	0.6	1.1
White, both sexes												
2004.....	1.7	2.1	0.2	0.1	1.3	2.1	3.3	3.2	1.4	0.8	0.9	1.6
2003.....	1.8	1.7	0.2	0.1	1.5	2.3	3.7	3.1	1.3	0.6	0.7	1.2
2002.....	1.7	1.8	0.4	0.2	1.3	2.1	3.4	2.8	1.2	0.7	0.7	1.4
2001.....	1.5	1.4	0.3	0.1	1.1	1.9	3.1	2.5	0.9	0.5	0.9	1.0
2000.....	1.3	1.4	0.2	0.1	1.0	1.7	2.8	2.1	0.9	0.5	0.6	1.3
1999.....	1.4	1.2	0.2	0.1	1.1	1.9	2.8	2.0	0.9	0.6	0.8	1.4
White, male												
2004.....	2.2	2.7	*	0.1	1.9	2.9	4.0	3.9	1.6	1.0	1.0	1.9
2003.....	2.3	1.9	*	0.2	2.1	3.0	4.7	3.9	1.5	0.7	1.0	1.5
2002.....	2.2	1.8	0.4	0.2	1.9	2.9	4.4	3.3	1.5	0.9	0.9	2.3
2001.....	2.0	1.7	*	0.2	1.6	2.7	4.0	3.2	1.1	0.7	1.2	*
2000.....	1.8	*	*	0.2	1.4	2.4	3.8	2.8	1.1	0.7	1.0	*
1999.....	1.9	1.4	*	0.2	1.8	2.5	3.8	2.6	1.1	0.8	1.1	2.1

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Event of undetermined intent (Y10-Y34,Y87.2,Y89.9)												
White, female												
2004.....	1.3	1.5	*	*	0.7	1.3	2.5	2.5	1.2	0.6	0.9	1.5
2003.....	1.3	1.4	*	*	0.9	1.5	2.7	2.4	1.0	0.6	0.6	1.0
2002.....	1.2	1.8	0.4	0.1	0.7	1.3	2.5	2.2	0.9	0.6	0.6	1.1
2001.....	1.0	*	*	*	0.6	1.2	2.2	1.7	0.6	0.4	0.6	0.8
2000.....	0.8	1.6	*	*	0.5	1.0	1.7	1.4	0.6	0.4	0.4	1.1
1999.....	0.9	*	*	*	0.4	1.2	1.8	1.4	0.7	0.4	0.6	1.1
Black, both sexes												
2004.....	1.8	4.8	*	0.4	0.9	1.8	3.3	3.9	1.7	1.3	*	*
2003.....	1.8	3.2	*	*	0.8	1.8	3.8	4.0	1.9	1.2	*	*
2002.....	2.0	5.5	*	*	0.9	2.3	4.2	4.2	1.5	*	*	*
2001.....	1.7	5.5	*	*	0.7	1.7	3.9	3.8	1.3	*	*	*
2000.....	1.8	*	*	0.3	0.9	1.9	4.1	3.7	1.8	*	*	*
1999.....	1.9	*	*	*	1.2	2.2	4.2	4.1	1.2	*	*	*
Black, male												
2004.....	2.6	*	*	*	1.3	2.8	4.6	5.8	2.9	*	*	*
2003.....	2.7	*	*	*	1.4	2.9	5.3	6.4	3.2	*	*	*
2002.....	2.9	*	*	*	1.4	3.5	6.0	6.4	2.5	*	*	*
2001.....	2.7	6.0	*	*	1.3	2.6	6.2	6.3	2.4	*	*	*
2000.....	2.8	*	*	*	1.6	3.1	6.0	5.9	3.0	*	*	*
1999.....	3.0	*	*	*	1.9	3.6	6.4	7.2	2.5	*	*	*
Black, female												
2004.....	1.1	*	*	*	*	0.9	2.2	2.2	*	*	*	*
2003.....	0.9	*	*	*	*	*	2.4	1.9	*	*	*	*
2002.....	1.2	*	*	*	*	1.2	2.6	2.4	*	*	*	*
2001.....	0.9	*	*	*	*	0.9	1.9	1.6	*	*	*	*
2000.....	0.9	*	*	*	*	0.9	2.4	1.8	*	*	*	*
1999.....	0.9	*	*	*	*	1.0	2.3	1.6	*	*	*	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Poisoning by and exposure to drugs and biological substances, undetermined intent (Y10-Y14)

All races 3/, both sexes

2004.....	1.1	*	*	*	0.7	1.5	2.5	2.3	0.8	0.3	*	*
2003.....	1.2	*	*	*	0.9	1.6	2.9	2.4	0.8	0.2	*	*
2002.....	1.1	*	*	*	0.7	1.5	2.6	2.2	0.7	0.2	*	*
2001.....	1.0	*	*	*	0.6	1.3	2.4	1.8	0.5	0.2	0.2	*
2000.....	0.9	*	*	*	0.5	1.1	2.2	1.6	0.4	*	*	*
1999.....	0.9	*	*	*	0.5	1.3	2.2	1.5	0.4	0.2	*	*

All races 3/, male

2004.....	1.4	*	*	*	1.1	1.9	3.0	2.8	0.8	0.3	*	*
2003.....	1.6	*	*	*	1.3	2.1	3.6	2.9	0.9	*	*	*
2002.....	1.4	*	*	*	0.9	2.0	3.2	2.5	0.8	*	*	*
2001.....	1.3	*	*	*	0.8	1.7	3.1	2.5	0.5	*	*	*
2000.....	1.2	*	*	*	0.7	1.6	3.1	2.2	0.5	*	*	*
1999.....	1.2	*	*	*	0.7	1.8	3.0	2.0	0.5	*	*	*

All races 3/, female

2004.....	0.9	*	*	*	0.4	1.0	1.9	1.9	0.8	0.3	*	*
2003.....	0.9	*	*	*	0.5	1.1	2.2	1.8	0.7	0.3	*	*
2002.....	0.8	*	*	*	0.4	1.0	2.0	1.8	0.6	0.3	*	*
2001.....	0.7	*	*	*	0.3	0.9	1.7	1.2	0.4	*	*	*
2000.....	0.5	*	*	*	0.3	0.7	1.4	1.1	0.3	*	*	*
1999.....	0.6	*	*	*	0.3	0.8	1.5	1.1	0.4	0.2	*	*

White, both sexes

2004.....	1.2	*	*	*	0.9	1.6	2.6	2.4	0.8	0.3	*	*
2003.....	1.3	*	*	*	1.1	1.8	3.1	2.4	0.8	0.2	*	*
2002.....	1.2	*	*	*	0.8	1.6	2.7	2.1	0.7	0.2	*	*
2001.....	1.0	*	*	*	0.7	1.4	2.5	1.8	0.5	0.1	*	*
2000.....	0.9	*	*	*	0.6	1.2	2.2	1.6	0.4	*	*	*
1999.....	0.9	*	*	*	0.6	1.4	2.2	1.4	0.4	0.2	*	*

White, male

2004.....	1.5	*	*	*	1.3	2.2	3.1	2.7	0.8	*	*	*
2003.....	1.6	*	*	*	1.5	2.3	3.8	2.8	0.8	*	*	*
2002.....	1.4	*	*	*	1.1	2.2	3.3	2.4	0.7	*	*	*
2001.....	1.3	*	*	*	1.0	1.9	3.0	2.3	0.5	*	*	*
2000.....	1.2	*	*	*	0.8	1.6	3.0	2.1	0.5	*	*	*
1999.....	1.1	*	*	*	0.8	1.8	2.8	1.7	0.4	*	*	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Poisoning by and exposure to drugs and biological substances, undetermined intent (Y10-Y14)

White, female

2004.....	1.0	*	*	*	0.5	1.1	2.1	2.1	0.9	0.3	*	*
2003.....	1.0	*	*	*	0.6	1.3	2.4	1.9	0.8	0.3	*	*
2002.....	0.9	*	*	*	0.5	1.1	2.0	1.9	0.6	0.3	*	*
2001.....	0.7	*	*	*	0.4	1.0	1.9	1.3	0.5	*	*	*
2000.....	0.6	*	*	*	0.3	0.7	1.4	1.1	0.4	*	*	*
1999.....	0.6	*	*	*	0.3	0.9	1.5	1.1	0.4	0.2	*	*

Black, both sexes

2004.....	1.0	*	*	*	*	1.0	2.5	2.6	1.0	*	*	*
2003.....	1.1	*	*	*	*	1.0	2.9	3.1	1.0	*	*	*
2002.....	1.2	*	*	*	0.3	1.4	3.2	3.2	0.9	*	*	*
2001.....	1.1	*	*	*	*	1.2	3.0	2.7	*	*	*	*
2000.....	1.0	*	*	*	*	1.1	3.1	2.6	*	*	*	*
1999.....	1.2	*	*	*	*	1.3	3.2	3.0	*	*	*	*

Black, male

2004.....	1.4	*	*	*	*	1.4	3.5	3.6	1.7	*	*	*
2003.....	1.7	*	*	*	*	1.6	4.0	4.9	1.6	*	*	*
2002.....	1.6	*	*	*	*	1.9	4.2	4.5	*	*	*	*
2001.....	1.7	*	*	*	*	1.6	4.8	4.5	*	*	*	*
2000.....	1.5	*	*	*	*	1.7	4.4	4.0	*	*	*	*
1999.....	1.8	*	*	*	*	2.2	4.7	5.0	*	*	*	*

Black, female

2004.....	0.7	*	*	*	*	0.7	1.7	1.7	*	*	*	*
2003.....	0.7	*	*	*	*	*	2.0	1.6	*	*	*	*
2002.....	0.9	*	*	*	*	0.9	2.3	2.1	*	*	*	*
2001.....	0.5	*	*	*	*	0.8	1.5	1.2	*	*	*	*
2000.....	0.6	*	*	*	*	*	1.9	1.5	*	*	*	*
1999.....	0.6	*	*	*	*	*	1.9	1.3	*	*	*	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Discharge of firearms, undetermined intent (Y22-Y24)

All races 3/, both sexes

2004.....	0.1	*	*	*	0.1	0.1	0.1	0.1	0.1	*	*	*
2003.....	0.1	*	*	*	0.2	0.1	0.1	0.1	0.1	*	*	*
2002.....	0.1	*	*	*	0.2	0.1	0.1	0.1	*	*	*	*
2001.....	0.1	*	*	*	0.2	0.1	0.1	0.1	*	*	*	*
2000.....	0.1	*	*	*	0.2	0.1	0.1	0.1	*	*	*	*
1999.....	0.1	*	*	*	0.3	0.1	0.1	0.1	*	*	*	*

All races 3/, male

2004.....	0.1	*	*	*	0.2	0.2	0.1	0.1	*	*	*	*
2003.....	0.1	*	*	*	0.3	0.2	0.1	0.1	*	*	*	*
2002.....	0.1	*	*	*	0.3	0.2	0.2	0.1	*	*	*	*
2001.....	0.1	*	*	*	0.3	0.2	0.1	0.1	*	*	*	*
2000.....	0.1	*	*	*	0.3	0.2	0.1	*	*	*	*	*
1999.....	0.2	*	*	*	0.6	0.2	0.2	0.1	*	*	*	*

All races 3/, female

2004.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2003.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	*	*	*	*	*	*

White, both sexes

2004.....	0.1	*	*	*	0.1	0.1	0.1	0.1	*	*	*	*
2003.....	0.1	*	*	*	0.1	0.1	0.1	0.1	*	*	*	*
2002.....	0.1	*	*	*	0.2	0.1	0.1	0.1	*	*	*	*
2001.....	0.1	*	*	*	0.1	0.1	0.1	0.1	*	*	*	*
2000.....	0.1	*	*	*	0.2	0.1	0.1	0.1	*	*	*	*
1999.....	0.1	*	*	*	0.3	0.1	0.1	0.1	*	*	*	*

White, male

2004.....	0.1	*	*	*	0.2	0.1	0.1	*	*	*	*	*
2003.....	0.1	*	*	*	0.2	0.1	0.1	0.1	*	*	*	*
2002.....	0.1	*	*	*	0.3	0.2	0.2	0.1	*	*	*	*
2001.....	0.1	*	*	*	0.2	0.2	0.2	*	*	*	*	*
2000.....	0.1	*	*	*	0.3	0.2	0.1	*	*	*	*	*
1999.....	0.2	*	*	*	0.5	0.2	0.2	*	*	*	*	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

All other and unspecified events of undetermined intent and their sequelae (Y20-Y21,Y25-Y29,Y31-Y34,Y87.2,Y89.9)

All races 3/, both sexes

2004.....	0.4	2.3	0.2	0.1	0.2	0.4	0.5	0.6	0.4	0.4	0.7	1.3
2003.....	0.4	1.7	0.1	0.1	0.2	0.3	0.5	0.5	0.4	0.4	0.5	0.8
2002.....	0.4	2.1	0.3	0.1	0.3	0.3	0.5	0.5	0.4	0.4	0.5	1.1
2001.....	0.3	1.7	0.2	0.1	0.2	0.3	0.5	0.5	0.3	0.3	0.5	0.7
2000.....	0.3	1.6	0.2	0.1	0.2	0.4	0.4	0.5	0.4	0.3	0.4	0.9
1999.....	0.3	1.2	0.2	0.1	0.2	0.3	0.4	0.5	0.4	0.3	0.4	1.0

All races 3/, male

2004.....	0.6	2.9	0.3	0.1	0.4	0.5	0.7	0.9	0.6	0.5	0.8	1.4
2003.....	0.5	2.2	*	0.1	0.3	0.5	0.7	0.8	0.6	0.4	0.6	*
2002.....	0.5	2.0	0.3	0.1	0.3	0.5	0.8	0.7	0.6	0.6	0.6	1.7
2001.....	0.5	2.0	*	0.1	0.4	0.5	0.7	0.7	0.5	0.4	0.7	*
2000.....	0.5	1.2	0.3	0.1	0.4	0.5	0.6	0.7	0.6	0.5	0.7	*
1999.....	0.5	1.4	*	0.1	0.4	0.5	0.6	0.7	0.6	0.4	0.6	*

All races 3/, female

2004.....	0.3	1.7	*	*	0.1	0.2	0.3	0.3	0.2	0.3	0.6	1.3
2003.....	0.2	1.2	*	*	0.1	0.2	0.3	0.3	0.2	0.3	0.4	0.7
2002.....	0.3	2.1	0.3	0.1	0.2	0.1	0.3	0.2	0.2	0.2	0.4	0.8
2001.....	0.2	1.5	0.3	*	0.1	0.1	0.2	0.3	*	*	0.4	*
2000.....	0.2	1.9	*	*	0.1	0.2	0.3	0.2	0.2	0.2	0.3	0.9
1999.....	0.2	*	*	*	*	0.2	0.2	0.2	*	*	0.3	0.8

White, both sexes

2004.....	0.4	2.0	0.2	0.1	0.2	0.3	0.5	0.5	0.4	0.3	0.7	1.3
2003.....	0.3	1.5	*	0.1	0.2	0.3	0.4	0.5	0.3	0.3	0.4	0.9
2002.....	0.4	1.6	0.3	0.1	0.3	0.3	0.5	0.4	0.4	0.3	0.5	1.1
2001.....	0.3	1.1	0.2	0.1	0.2	0.3	0.4	0.4	0.3	0.3	0.5	0.7
2000.....	0.3	1.2	0.2	0.1	0.2	0.3	0.4	0.4	0.3	0.3	0.4	0.8
1999.....	0.3	1.1	0.2	0.1	0.2	0.3	0.4	0.4	0.4	0.3	0.4	0.9

White, male

2004.....	0.5	2.6	*	*	0.3	0.5	0.7	0.8	0.5	0.4	0.7	*
2003.....	0.5	1.7	*	*	0.3	0.4	0.6	0.8	0.5	0.4	0.6	*
2002.....	0.5	1.6	0.4	*	0.3	0.5	0.7	0.6	0.6	0.5	0.6	1.8
2001.....	0.4	1.4	*	*	0.3	0.4	0.7	0.6	0.4	0.4	0.7	*
2000.....	0.4	*	*	0.1	0.3	0.4	0.5	0.6	0.5	0.4	0.6	*
1999.....	0.4	1.4	*	0.1	0.4	0.4	0.5	0.6	0.5	0.4	0.6	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Legal intervention (Y35,Y89.0)												
All races 3/, both sexes												
2004.....	0.1	*	*	*	0.2	0.3	0.2	0.1	*	*	*	*
2003.....	0.1	*	*	*	0.2	0.3	0.3	0.1	*	*	*	*
2002.....	0.1	*	*	*	0.2	0.3	0.2	0.1	*	*	*	*
2001.....	0.1	*	*	*	0.2	0.3	0.3	0.1	0.1	*	*	*
2000.....	0.1	*	*	*	0.2	0.3	0.2	0.1	*	*	*	*
1999.....	0.1	*	*	*	0.2	0.3	0.3	0.1	0.1	*	*	*
All races 3/, male												
2004.....	0.3	*	*	*	0.4	0.5	0.5	0.3	*	*	*	*
2003.....	0.3	*	*	*	0.4	0.6	0.5	0.2	*	*	*	*
2002.....	0.3	*	*	*	0.4	0.5	0.5	0.3	*	*	*	*
2001.....	0.3	*	*	*	0.4	0.5	0.5	0.3	0.2	*	*	*
2000.....	0.3	*	*	*	0.4	0.5	0.4	0.3	*	*	*	*
1999.....	0.3	*	*	*	0.3	0.7	0.5	0.3	0.2	*	*	*
All races 3/, female												
2004.....	*	*	*	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*	*	*
White, both sexes												
2004.....	0.1	*	*	*	0.2	0.2	0.2	0.1	*	*	*	*
2003.....	0.1	*	*	*	0.2	0.2	0.3	0.1	*	*	*	*
2002.....	0.1	*	*	*	0.2	0.2	0.2	0.1	*	*	*	*
2001.....	0.1	*	*	*	0.2	0.2	0.2	0.1	*	*	*	*
2000.....	0.1	*	*	*	0.1	0.2	0.2	0.1	*	*	*	*
1999.....	0.1	*	*	*	0.1	0.3	0.2	0.1	*	*	*	*
White, male												
2004.....	0.2	*	*	*	0.3	0.4	0.4	0.2	*	*	*	*
2003.....	0.2	*	*	*	0.3	0.5	0.5	0.2	*	*	*	*
2002.....	0.2	*	*	*	0.3	0.4	0.4	0.2	*	*	*	*
2001.....	0.2	*	*	*	0.4	0.5	0.4	0.2	*	*	*	*
2000.....	0.2	*	*	*	0.3	0.4	0.3	0.3	*	*	*	*
1999.....	0.2	*	*	*	0.2	0.5	0.4	0.3	*	*	*	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Legal execution (Y35.5)

All races 3/, both sexes

2004.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2003.....	0.0	*	*	*	*	*	0.0	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	0.1	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	0.1	*	*	*	*	*
2000.....	0.0	*	*	*	*	0.1	0.1	0.1	*	*	*	*
1999.....	0.0	*	*	*	*	*	0.1	*	*	*	*	*

All races 3/, male

2004.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2003.....	0.0	*	*	*	*	*	0.1	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	0.1	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	0.1	*	*	*	*	*
2000.....	0.1	*	*	*	*	0.1	0.1	0.1	*	*	*	*
1999.....	0.1	*	*	*	*	*	0.2	*	*	*	*	*

All races 3/, female

2004.....	*	*	*	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*	*	*

White, both sexes

2004.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2003.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	0.1	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	0.1	*	*	*	*	*

White, male

2004.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2003.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	0.1	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	0.1	*	*	*	*	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other legal intervention and their sequelae (Y35.0-Y35.4,Y35.6-Y35.7,Y89.0)

All races 3/, both sexes

2004.....	0.1	*	*	*	0.2	0.2	0.2	0.1	*	*	*	*
2003.....	0.1	*	*	*	0.2	0.3	0.2	0.1	*	*	*	*
2002.....	0.1	*	*	*	0.2	0.2	0.2	0.1	*	*	*	*
2001.....	0.1	*	*	*	0.2	0.2	0.2	0.1	*	*	*	*
2000.....	0.1	*	*	*	0.2	0.2	0.2	0.1	*	*	*	*
1999.....	0.1	*	*	*	0.2	0.3	0.2	0.1	*	*	*	*

All races 3/, male

2004.....	0.2	*	*	*	0.4	0.4	0.4	0.2	*	*	*	*
2003.....	0.2	*	*	*	0.4	0.6	0.4	0.2	*	*	*	*
2002.....	0.2	*	*	*	0.4	0.4	0.3	0.2	*	*	*	*
2001.....	0.2	*	*	*	0.4	0.5	0.4	0.2	*	*	*	*
2000.....	0.2	*	*	*	0.3	0.4	0.3	0.2	*	*	*	*
1999.....	0.2	*	*	*	0.3	0.6	0.3	0.2	*	*	*	*

All races 3/, female

2004.....	*	*	*	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*	*	*

White, both sexes

2004.....	0.1	*	*	*	0.2	0.2	0.2	0.1	*	*	*	*
2003.....	0.1	*	*	*	0.2	0.2	0.2	0.1	*	*	*	*
2002.....	0.1	*	*	*	0.2	0.2	0.2	0.1	*	*	*	*
2001.....	0.1	*	*	*	0.2	0.2	0.2	0.1	*	*	*	*
2000.....	0.1	*	*	*	0.1	0.2	0.1	0.1	*	*	*	*
1999.....	0.1	*	*	*	0.1	0.2	0.2	0.1	*	*	*	*

White, male

2004.....	0.2	*	*	*	0.3	0.4	0.4	0.2	*	*	*	*
2003.....	0.2	*	*	*	0.3	0.4	0.4	0.2	*	*	*	*
2002.....	0.2	*	*	*	0.3	0.4	0.3	0.2	*	*	*	*
2001.....	0.2	*	*	*	0.4	0.4	0.3	0.2	*	*	*	*
2000.....	0.2	*	*	*	0.3	0.4	0.3	0.2	*	*	*	*
1999.....	0.2	*	*	*	0.2	0.5	0.3	0.2	*	*	*	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Complications of medical and surgical care (Y40-Y84,Y88)

All races 3/, both sexes

2004.....	1.0	*	*	*	0.1	0.2	0.4	0.7	1.4	3.0	6.1	10.6
2003.....	1.0	*	0.2	*	0.1	0.2	0.4	0.7	1.4	2.9	6.2	11.2
2002.....	1.0	*	*	0.1	0.1	0.2	0.4	0.7	1.3	3.1	6.2	11.3
2001.....	1.1	0.6	0.2	*	0.1	0.2	0.4	0.7	1.5	3.1	6.9	12.8
2000.....	1.1	*	0.1	0.1	0.1	0.2	0.3	0.8	1.5	3.5	6.7	13.6
1999.....	1.0	1.4	0.2	*	0.1	0.2	0.3	0.6	1.3	3.2	6.6	12.7

All races 3/, male

2004.....	0.9	*	*	*	0.1	0.2	0.3	0.7	1.6	3.3	7.4	11.9
2003.....	0.9	*	*	*	0.1	0.2	0.4	0.7	1.5	3.2	6.9	12.9
2002.....	0.9	*	*	0.1	*	0.2	0.4	0.6	1.3	3.1	6.3	12.3
2001.....	1.0	*	*	*	*	0.2	0.4	0.7	1.6	3.2	7.7	14.5
2000.....	1.0	*	*	*	*	0.2	0.3	0.8	1.6	4.0	8.2	16.0
1999.....	1.0	1.6	*	*	0.1	0.2	0.3	0.5	1.4	3.4	7.8	16.5

All races 3/, female

2004.....	1.0	*	*	*	0.1	0.2	0.4	0.7	1.2	2.8	5.2	9.9
2003.....	1.0	*	*	*	*	0.2	0.4	0.6	1.3	2.7	5.7	10.5
2002.....	1.1	*	*	*	*	0.2	0.4	0.7	1.3	3.0	6.2	10.9
2001.....	1.1	*	*	*	0.1	0.2	0.4	0.7	1.4	3.0	6.3	12.1
2000.....	1.1	*	*	*	*	0.2	0.3	0.8	1.5	3.1	5.7	12.7
1999.....	1.1	1.2	*	*	*	0.2	0.3	0.6	1.2	3.0	5.9	11.2

White, both sexes

2004.....	1.0	*	*	*	0.1	0.2	0.3	0.7	1.3	2.9	5.9	10.4
2003.....	1.0	*	*	*	0.1	0.2	0.4	0.6	1.2	2.8	6.1	11.0
2002.....	1.0	*	*	0.1	0.1	0.2	0.3	0.6	1.2	3.0	6.1	11.2
2001.....	1.1	*	*	*	0.1	0.2	0.3	0.6	1.4	2.8	6.8	12.1
2000.....	1.1	*	*	0.1	0.1	0.2	0.3	0.7	1.4	3.3	6.4	13.1
1999.....	1.0	1.2	0.2	*	0.1	0.2	0.3	0.5	1.2	3.0	6.3	12.1

White, male

2004.....	1.0	*	*	*	*	0.2	0.3	0.7	1.5	3.3	7.3	11.8
2003.....	1.0	*	*	*	0.1	0.2	0.4	0.6	1.4	3.1	7.0	12.8
2002.....	0.9	*	*	*	*	0.2	0.4	0.6	1.2	2.9	6.3	11.8
2001.....	0.9	*	*	*	*	0.2	0.3	0.6	1.4	2.8	7.7	13.6
2000.....	1.1	*	*	*	*	0.2	0.3	0.7	1.4	3.7	8.2	15.8
1999.....	1.0	1.4	*	*	*	0.2	0.3	0.5	1.3	3.2	7.5	15.6

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Complications of medical and surgical care (Y40-Y84,Y88)

White, female

2004.....	1.0	*	*	*	*	0.2	0.3	0.6	1.1	2.5	4.9	9.8
2003.....	1.0	*	*	*	*	0.2	0.3	0.5	1.1	2.5	5.4	10.2
2002.....	1.1	*	*	*	*	0.1	0.3	0.5	1.2	3.0	6.0	10.9
2001.....	1.2	*	*	*	*	0.2	0.3	0.6	1.3	2.7	6.2	11.5
2000.....	1.1	*	*	*	*	0.2	0.3	0.7	1.3	3.0	5.3	12.0
1999.....	1.1	*	*	*	*	0.1	0.3	0.6	1.1	2.9	5.6	10.7

Black, both sexes

2004.....	1.1	*	*	*	*	*	0.8	1.2	2.6	5.0	9.3	13.8
2003.....	1.1	*	*	*	*	*	0.6	1.4	2.7	4.6	8.8	15.2
2002.....	1.2	*	*	*	*	*	0.8	1.6	2.7	4.6	8.3	14.0
2001.....	1.4	*	*	*	*	0.4	0.9	1.6	3.4	6.3	9.2	23.3
2000.....	1.3	*	*	*	*	0.4	0.5	1.4	3.5	6.1	10.5	21.0
1999.....	1.2	*	*	*	*	*	0.6	1.3	2.6	5.2	10.6	20.1

Black, male

2004.....	0.9	*	*	*	*	*	*	1.0	2.9	4.6	10.0	*
2003.....	1.0	*	*	*	*	*	*	1.8	2.8	3.9	7.6	*
2002.....	1.0	*	*	*	*	*	*	1.3	2.5	5.3	7.2	*
2001.....	1.4	*	*	*	*	*	1.0	1.8	4.2	7.8	9.1	29.5
2000.....	1.1	*	*	*	*	*	*	1.3	3.1	7.6	10.3	*
1999.....	1.1	*	*	*	*	*	*	1.4	2.8	5.9	12.3	23.4

Black, female

2004.....	1.3	*	*	*	*	*	1.0	1.4	2.3	5.2	9.0	13.4
2003.....	1.3	*	*	*	*	*	0.7	1.1	2.5	5.0	9.4	15.4
2002.....	1.3	*	*	*	*	*	1.1	1.8	2.7	4.1	8.9	11.5
2001.....	1.4	*	*	*	*	*	0.7	1.3	2.7	5.3	9.3	21.0
2000.....	1.5	*	*	*	*	*	*	1.5	3.8	5.0	10.6	21.9
1999.....	1.3	*	*	*	*	*	*	1.2	2.4	4.8	9.6	18.8

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other complications of medical and surgical care and their sequelae (Y70-Y84,Y88.2-Y88.3)

All races 3/, both sexes

2004.....	0.8	*	*	*	0.1	0.1	0.2	0.5	1.1	2.6	5.0	9.3
2003.....	0.8	*	0.1	*	0.1	0.1	0.3	0.5	1.1	2.5	5.5	9.9
2002.....	0.8	*	*	0.1	0.1	0.1	0.3	0.5	1.1	2.6	5.5	10.4
2001.....	0.9	*	*	*	0.1	0.1	0.3	0.6	1.4	2.7	6.1	11.4
2000.....	0.9	*	*	0.1	0.1	0.1	0.2	0.6	1.3	3.1	5.8	12.2
1999.....	0.9	1.2	0.1	*	0.1	0.1	0.2	0.5	1.1	2.7	5.9	11.8

All races 3/, male

2004.....	0.8	*	*	*	*	*	0.2	0.6	1.3	2.9	6.2	10.7
2003.....	0.8	*	*	*	*	*	0.3	0.6	1.2	2.7	6.2	11.5
2002.....	0.7	*	*	*	*	0.1	0.3	0.5	1.1	2.7	5.7	11.7
2001.....	0.8	*	*	*	*	0.1	0.2	0.6	1.4	2.8	6.8	13.2
2000.....	0.9	*	*	*	*	0.1	0.2	0.6	1.3	3.6	7.1	14.3
1999.....	0.8	1.4	*	*	*	0.1	0.2	0.4	1.2	2.9	7.0	15.4

All races 3/, female

2004.....	0.8	*	*	*	*	0.1	0.3	0.5	0.9	2.3	4.2	8.7
2003.....	0.9	*	*	*	*	0.1	0.3	0.4	1.1	2.3	5.0	9.2
2002.....	0.9	*	*	*	*	0.1	0.3	0.6	1.1	2.6	5.4	9.9
2001.....	1.0	*	*	*	0.1	0.1	0.3	0.5	1.3	2.5	5.6	10.6
2000.....	1.0	*	*	*	*	0.1	0.2	0.6	1.3	2.7	5.0	11.4
1999.....	0.9	*	*	*	*	0.1	0.2	0.5	1.0	2.6	5.2	10.4

White, both sexes

2004.....	0.8	*	*	*	0.1	0.1	0.2	0.5	1.0	2.5	4.9	9.1
2003.....	0.8	*	*	*	*	0.1	0.3	0.4	1.0	2.4	5.3	9.6
2002.....	0.9	*	*	*	*	0.1	0.2	0.5	1.0	2.5	5.4	10.2
2001.....	0.9	*	*	*	0.1	0.1	0.2	0.5	1.2	2.4	6.0	10.7
2000.....	0.9	*	*	0.1	*	0.1	0.2	0.6	1.2	2.9	5.6	11.8
1999.....	0.9	1.0	*	*	0.1	0.1	0.2	0.4	1.0	2.7	5.7	11.2

White, male

2004.....	0.8	*	*	*	*	*	0.2	0.6	1.2	2.9	6.2	10.4
2003.....	0.8	*	*	*	*	*	0.3	0.4	1.1	2.7	6.2	11.5
2002.....	0.8	*	*	*	*	0.2	0.3	0.5	1.0	2.5	5.7	11.2
2001.....	0.8	*	*	*	*	*	0.2	0.5	1.2	2.5	6.8	12.3
2000.....	0.9	*	*	*	*	0.1	0.2	0.6	1.2	3.4	7.0	14.0
1999.....	0.8	1.3	*	*	*	0.1	0.2	0.4	1.1	2.8	6.8	14.5

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other complications of medical and surgical care and their sequelae (Y70-Y84,Y88.2-Y88.3)

White, female

2004.....	0.8	*	*	*	*	0.1	0.2	0.4	0.8	2.1	4.0	8.4
2003.....	0.9	*	*	*	*	*	0.3	0.4	0.9	2.1	4.8	8.8
2002.....	0.9	*	*	*	*	*	0.2	0.5	1.0	2.6	5.2	9.8
2001.....	1.0	*	*	*	*	0.1	0.3	0.5	1.2	2.3	5.5	10.0
2000.....	1.0	*	*	*	*	0.1	0.2	0.6	1.1	2.6	4.7	10.8
1999.....	0.9	*	*	*	*	*	0.2	0.4	0.9	2.5	5.0	9.9

Black, both sexes

2004.....	0.9	*	*	*	*	*	0.5	0.9	2.0	4.3	7.9	13.0
2003.....	0.9	*	*	*	*	*	0.4	1.2	2.1	3.9	7.8	14.0
2002.....	1.0	*	*	*	*	*	0.7	1.3	2.0	4.0	7.7	13.4
2001.....	1.2	*	*	*	*	*	0.7	1.1	3.0	5.3	7.9	22.1
2000.....	1.1	*	*	*	*	*	0.5	1.0	2.9	5.2	9.5	19.1
1999.....	1.0	*	*	*	*	*	0.4	1.0	2.3	4.3	9.4	19.1

Black, male

2004.....	0.7	*	*	*	*	*	*	*	2.3	3.9	8.4	*
2003.....	0.8	*	*	*	*	*	*	1.6	2.1	3.2	7.1	*
2002.....	0.9	*	*	*	*	*	*	1.2	2.1	4.7	6.9	*
2001.....	1.2	*	*	*	*	*	0.7	1.4	3.9	6.8	7.9	28.4
2000.....	0.9	*	*	*	*	*	*	1.1	2.7	6.3	9.7	*
1999.....	1.0	*	*	*	*	*	*	1.1	2.5	5.1	11.1	*

Worktable 12. Death rates for 358 selected causes, by 10-year age groups, race, and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other complications of medical and surgical care and their sequelae (Y70-Y84,Y88.2-Y88.3)

Black, female

2004.....	1.1	*	*	*	*	*	0.7	1.1	1.7	4.6	7.7	12.6
2003.....	1.1	*	*	*	*	*	*	0.8	2.1	4.4	8.3	15.0
2002.....	1.1	*	*	*	*	*	0.9	1.3	1.9	3.4	8.1	11.1
2001.....	1.2	*	*	*	*	*	0.7	0.9	2.3	4.3	7.9	19.8
2000.....	1.3	*	*	*	*	*	*	0.9	3.1	4.4	9.4	19.3
1999.....	1.1	*	*	*	*	*	*	0.9	2.2	3.6	8.4	18.0

* Figure does not meet standards of reliability or precision.

0.0 Quantity more than zero but less than 0.05.

... Category not applicable.

1/ Figures for age not stated included in "All ages" but not distributed among age groups.

2/ Death rates for "Under 1 year" (based on population estimates) differ from infant mortality rates (based on live births).

3/ Includes races other than white and black.

4/ Figures include September 11, 2001 related deaths for which death certificates were filed as of 10/24/02; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001."

Source: CDC/NCHS, National Vital Statistics System, Mortality.