

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
All causes										
All races 3/, both sexes										
2005.....	825.9	692.5	29.4	14.5	18.1	65.1	97.6	97.5	111.4	149.6
2004.....	816.5	685.2	29.9	14.7	18.7	66.1	94.0	96.0	107.9	151.7
2003.....	841.9	700.0	31.5	14.7	19.1	66.4	96.4	95.2	111.3	159.1
2002.....	847.3	695.0	31.2	15.2	19.5	67.8	95.2	94.7	111.6	161.3
2001.....	848.5	683.4	33.3	15.3	19.2	66.9	95.0	96.2	113.5	165.9
2000.....	854.0	736.7	32.4	15.8	20.3	67.1	93.6	91.2	111.0	159.2
1999.....	857.0	736.0	34.2	16.9	20.4	68.6	90.8	90.4	113.4	160.8
All races 3/, male										
2005.....	827.2	762.3	33.4	15.6	21.5	91.6	143.9	138.7	148.1	189.4
2004.....	817.6	753.7	32.4	16.4	21.7	91.0	138.0	135.6	143.2	193.8
2003.....	840.3	777.4	35.1	16.2	23.1	92.3	140.3	133.5	148.9	202.7
2002.....	846.6	761.5	35.2	16.7	23.1	94.0	140.8	134.4	149.4	206.6
2001.....	846.4	749.8	37.0	16.7	22.8	93.7	141.1	135.6	151.2	213.7
2000.....	853.0	806.5	35.9	17.6	24.2	93.3	138.0	128.8	147.9	205.4
1999.....	859.2	805.0	37.9	18.6	24.4	94.7	133.1	127.0	152.0	207.6
All races 3/, female										
2005.....	824.6	619.4	25.1	13.4	14.4	37.2	48.2	54.4	73.7	109.3
2004.....	815.4	613.4	27.3	13.0	15.4	39.9	47.3	54.5	72.0	109.3
2003.....	843.4	619.1	27.8	13.0	15.0	39.0	49.8	55.4	73.0	115.3
2002.....	848.0	625.3	27.0	13.5	15.9	40.1	47.3	53.6	73.3	116.1
2001.....	850.4	613.9	29.5	13.9	15.3	38.5	46.8	55.8	75.2	118.2
2000.....	855.0	663.4	28.7	14.0	16.1	39.3	47.1	52.8	73.6	113.2
1999.....	854.9	663.6	30.3	15.0	16.2	41.0	46.6	53.2	74.4	114.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
All causes										
White, both sexes										
2005 4/.....	873.7	579.1	27.0	13.3	16.7	62.5	91.5	89.3	101.7	138.3
2004 4/.....	863.2	574.0	27.0	13.4	17.4	64.4	88.3	87.3	98.3	140.6
2003 4/.....	890.1	591.3	28.5	13.7	17.8	64.7	89.4	86.1	101.9	145.5
2002.....	895.7	586.7	28.1	14.0	18.1	66.3	88.0	85.0	101.1	146.8
2001.....	895.1	570.9	30.7	14.4	17.9	64.2	87.1	87.0	102.7	151.2
2000.....	900.2	610.6	29.2	14.8	19.2	64.5	85.1	81.0	99.0	144.0
1999.....	901.4	606.1	30.8	15.3	19.2	65.3	81.6	80.8	101.4	144.5
White, male										
2005 4/.....	864.5	640.0	30.9	14.2	19.8	86.5	133.8	126.5	135.1	176.1
2004 4/.....	854.2	631.6	29.4	15.1	20.6	87.1	129.1	123.2	130.8	180.9
2003 4/.....	877.6	658.7	31.5	15.3	21.4	88.3	129.1	120.0	137.0	187.6
2002.....	884.0	650.9	31.5	15.4	21.2	90.5	128.9	119.7	136.0	188.7
2001.....	881.9	627.6	34.2	15.3	21.3	88.3	128.2	122.4	137.4	196.7
2000.....	887.8	667.6	32.6	16.5	23.1	88.1	124.6	113.9	133.6	188.2
1999.....	892.1	667.0	33.9	16.8	22.7	88.3	118.5	113.5	136.5	188.8
White, female										
2005 4/.....	882.8	515.3	22.9	12.3	13.4	37.2	45.8	49.4	66.6	99.0
2004 4/.....	871.9	513.6	24.4	11.6	14.2	40.4	44.3	48.8	64.3	98.9
2003 4/.....	902.3	520.7	25.5	12.1	14.0	39.6	46.7	50.0	65.2	102.2
2002.....	907.0	519.4	24.5	12.6	14.8	40.4	44.4	48.1	64.8	103.9
2001.....	907.9	511.5	27.1	13.3	14.4	38.5	43.2	49.6	66.6	104.9
2000.....	912.3	550.5	25.5	13.0	15.2	39.3	43.0	46.5	63.0	99.2
1999.....	910.4	542.0	27.5	13.8	15.4	40.8	42.3	46.7	65.0	99.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
All causes										
Black, both sexes										
2005 4/.....	749.4	1,311.2	41.8	21.1	25.2	83.6	143.1	162.0	197.3	254.1
2004 4/.....	744.3	1,284.7	44.8	21.6	25.3	81.4	138.8	167.2	191.3	254.4
2003 4/.....	763.6	1,273.8	46.8	19.7	25.7	80.0	147.3	168.6	194.1	275.2
2002.....	768.4	1,263.6	47.1	21.3	27.5	82.0	146.8	172.1	200.7	284.0
2001.....	773.5	1,304.5	47.5	20.7	25.8	86.3	151.3	168.8	200.5	286.0
2000.....	781.1	1,426.1	49.9	21.8	26.6	87.4	155.0	168.4	206.9	283.1
1999.....	788.1	1,457.3	52.6	25.2	28.0	91.9	154.4	163.5	204.9	293.5
Black, male										
2005 4/.....	799.2	1,437.2	46.7	22.6	30.9	127.2	220.4	239.1	270.7	323.9
2004 4/.....	792.6	1,414.2	48.6	23.1	28.6	121.0	210.1	244.7	259.9	323.6
2003 4/.....	813.7	1,410.0	53.7	21.6	31.4	120.9	224.5	249.6	262.5	344.8
2002.....	816.7	1,351.5	54.4	24.1	33.4	121.7	227.6	255.7	272.9	366.5
2001.....	823.9	1,426.4	53.2	23.8	31.6	130.4	236.3	247.0	271.0	364.1
2000.....	834.1	1,567.6	54.5	24.5	32.1	130.1	240.6	246.8	275.0	361.6
1999.....	847.4	1,592.8	59.0	29.3	35.0	137.7	239.9	238.0	278.9	372.8
Black, female										
2005 4/.....	703.9	1,179.7	36.7	19.6	19.3	38.8	64.5	89.1	131.0	191.5
2004 4/.....	700.3	1,149.9	40.9	20.1	21.9	40.7	67.0	94.7	129.5	192.6
2003 4/.....	717.9	1,132.2	39.7	17.8	19.8	37.8	70.6	93.6	132.5	213.1
2002.....	724.4	1,172.0	39.5	18.5	21.3	41.0	68.0	95.7	135.7	210.6
2001.....	727.7	1,176.3	41.7	17.5	19.8	40.8	69.4	97.5	137.3	216.6
2000.....	733.0	1,279.8	45.3	19.0	21.0	43.7	73.9	97.3	145.6	213.4
1999.....	734.3	1,317.4	46.1	21.0	20.7	45.2	73.2	96.1	138.6	223.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Septicemia (A40-A41)

All races 3/, both sexes

2005.....	11.5	7.4	0.5	0.2	0.2	0.3	0.4	0.6	0.9	1.5
2004.....	11.4	6.6	0.5	0.2	0.1	0.2	0.4	0.6	1.0	1.5
2003.....	11.7	6.9	0.5	0.2	0.2	0.3	0.5	0.6	0.9	1.5
2002.....	11.7	7.3	0.5	0.2	0.3	0.2	0.3	0.6	0.9	1.5
2001.....	11.3	7.7	0.7	0.1	0.2	0.3	0.4	0.5	0.9	1.4
2000.....	11.1	7.2	0.6	0.2	0.1	0.2	0.3	0.5	0.9	1.4
1999.....	11.0	7.5	0.6	0.2	0.1	0.2	0.5	0.5	0.8	1.4

All races 3/, male

2005.....	10.5	8.1	0.6	0.2	0.2	0.3	0.5	0.7	0.8	1.4
2004.....	10.4	8.0	0.5	*	*	0.2	0.4	0.5	1.1	1.5
2003.....	10.5	7.4	0.7	*	*	0.3	0.4	0.6	1.0	1.6
2002.....	10.6	7.7	0.5	0.3	0.2	0.2	0.4	0.7	1.0	1.5
2001.....	10.2	8.4	0.8	*	0.2	0.3	0.4	0.5	0.9	1.5
2000.....	9.8	7.1	0.7	0.2	*	0.2	0.4	0.6	0.9	1.4
1999.....	9.8	9.0	0.6	0.2	*	*	0.5	0.5	0.9	1.5

All races 3/, female

2005.....	12.5	6.5	0.5	*	0.2	0.3	0.4	0.6	1.0	1.6
2004.....	12.3	5.2	0.6	0.2	*	0.2	0.4	0.7	1.0	1.4
2003.....	12.9	6.5	0.4	0.2	*	0.2	0.5	0.7	0.8	1.4
2002.....	12.9	7.0	0.5	*	0.3	0.3	0.3	0.6	0.9	1.5
2001.....	12.4	7.0	0.6	*	*	0.3	0.3	0.5	0.9	1.2
2000.....	12.3	7.3	0.6	*	*	*	0.2	0.4	0.8	1.4
1999.....	12.2	5.8	0.5	0.2	*	*	0.4	0.5	0.7	1.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Septicemia (A40-A41)

White, both sexes

2005 4/.....	11.3	5.0	0.5	0.2	0.2	0.2	0.4	0.5	0.7	1.2
2004 4/.....	11.2	5.2	0.5	0.2	*	0.2	0.3	0.5	0.8	1.3
2003 4/.....	11.5	4.6	0.4	0.2	0.2	0.2	0.4	0.5	0.7	1.2
2002.....	11.5	6.1	0.5	0.2	0.2	0.2	0.3	0.5	0.8	1.2
2001.....	11.1	5.4	0.7	0.2	0.2	0.2	0.3	0.4	0.7	1.1
2000.....	10.8	5.7	0.6	0.2	0.1	0.2	0.2	0.4	0.6	1.1
1999.....	10.6	5.5	0.5	0.2	0.1	0.2	0.4	0.4	0.7	1.0

White, male

2005 4/.....	10.2	5.9	0.5	*	*	*	0.4	0.6	0.6	1.1
2004 4/.....	10.2	6.3	0.5	*	*	*	0.3	0.4	0.8	1.4
2003 4/.....	10.2	4.6	0.6	*	*	0.3	0.4	0.4	0.8	1.3
2002.....	10.3	6.2	0.5	0.3	0.2	*	0.4	0.6	0.8	1.2
2001.....	9.9	6.2	0.8	*	*	0.3	0.3	0.4	0.6	1.3
2000.....	9.4	5.8	0.6	*	*	*	0.3	0.4	0.7	1.2
1999.....	9.3	6.9	0.5	*	*	*	0.5	0.5	0.8	1.2

White, female

2005 4/.....	12.4	4.1	0.4	*	*	*	0.3	0.4	0.8	1.3
2004 4/.....	12.2	4.0	0.5	*	*	*	0.3	0.6	0.8	1.2
2003 4/.....	12.8	4.5	*	*	*	*	0.4	0.6	0.7	1.1
2002.....	12.8	6.0	0.4	*	*	0.3	*	0.4	0.8	1.2
2001.....	12.3	4.6	0.6	*	*	*	0.3	0.4	0.7	0.9
2000.....	12.1	5.6	0.5	*	*	*	*	0.4	0.5	1.0
1999.....	11.9	4.0	0.5	*	*	*	*	0.4	0.6	0.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Septicemia (A40-A41)

Black, both sexes

2005 4/.....	15.9	19.8	0.8	*	*	0.6	0.8	1.5	2.3	3.6
2004 4/.....	15.6	14.5	*	*	*	*	1.0	1.3	2.5	2.7
2003 4/.....	16.3	18.6	0.9	*	*	*	0.7	1.6	2.2	3.6
2002.....	16.3	13.5	*	*	*	*	*	1.6	2.2	3.3
2001.....	15.8	19.3	0.9	*	*	*	0.7	1.1	2.5	3.7
2000.....	16.2	16.4	1.0	*	*	*	0.8	1.5	2.5	3.7
1999.....	16.1	17.7	*	*	*	*	1.0	1.4	1.7	3.7

Black, male

2005 4/.....	14.9	20.4	*	*	*	*	*	1.6	2.1	3.6
2004 4/.....	14.6	16.3	*	*	*	*	*	*	2.8	2.5
2003 4/.....	15.3	21.1	*	*	*	*	*	1.6	2.7	3.6
2002.....	15.0	14.8	*	*	*	*	*	*	2.5	3.7
2001.....	14.9	20.1	*	*	*	*	*	*	2.5	3.7
2000.....	14.9	16.0	*	*	*	*	*	2.1	2.5	3.5
1999.....	15.3	19.8	*	*	*	*	*	*	2.1	4.0

Black, female

2005 4/.....	16.8	19.1	*	*	*	*	*	1.4	2.6	3.7
2004 4/.....	16.4	12.7	*	*	*	*	*	1.4	2.1	2.9
2003 4/.....	17.2	16.1	*	*	*	*	*	1.6	1.6	3.5
2002.....	17.4	12.1	*	*	*	*	*	1.9	2.0	3.1
2001.....	16.7	18.6	*	*	*	*	*	*	2.5	3.7
2000.....	17.4	16.9	*	*	*	*	*	*	2.5	3.9
1999.....	16.9	15.4	*	*	*	*	*	1.6	1.4	3.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Viral hepatitis (B15-B19)										
All races 3/, both sexes										
2005.....	1.9	*	*	*	*	*	*	0.1	0.2	0.6
2004.....	1.8	*	*	*	*	*	*	0.1	0.2	0.7
2003.....	1.9	*	*	*	*	*	*	0.1	0.2	0.8
2002.....	2.0	*	*	*	*	*	*	0.2	0.3	1.0
2001.....	2.0	*	*	*	*	*	*	*	0.3	0.9
2000.....	1.9	*	*	*	*	*	*	0.1	0.3	1.0
1999.....	1.7	*	*	*	*	*	*	*	0.3	0.9
All races 3/, male										
2005.....	2.5	*	*	*	*	*	*	*	*	0.8
2004.....	2.4	*	*	*	*	*	*	*	0.2	0.8
2003.....	2.5	*	*	*	*	*	*	*	0.3	1.2
2002.....	2.7	*	*	*	*	*	*	0.2	0.4	1.4
2001.....	2.5	*	*	*	*	*	*	*	0.3	1.2
2000.....	2.4	*	*	*	*	*	*	*	0.3	1.3
1999.....	2.2	*	*	*	*	*	*	*	0.4	1.2
All races 3/, female										
2005.....	1.3	*	*	*	*	*	*	*	*	0.4
2004.....	1.3	*	*	*	*	*	*	*	*	0.5
2003.....	1.3	*	*	*	*	*	*	*	*	0.4
2002.....	1.4	*	*	*	*	*	*	*	*	0.6
2001.....	1.4	*	*	*	*	*	*	*	0.2	0.6
2000.....	1.4	*	*	*	*	*	*	*	0.2	0.6
1999.....	1.3	*	*	*	*	*	*	*	0.2	0.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Viral hepatitis (B15-B19)

White, both sexes

2005 4/.....	1.8	*	*	*	*	*	*	*	0.1	0.6
2004 4/.....	1.8	*	*	*	*	*	*	*	0.2	0.6
2003 4/.....	1.8	*	*	*	*	*	*	*	0.2	0.8
2002.....	2.0	*	*	*	*	*	*	0.1	0.3	1.0
2001.....	1.9	*	*	*	*	*	*	*	0.2	0.9
2000.....	1.8	*	*	*	*	*	*	*	0.2	1.0
1999.....	1.7	*	*	*	*	*	*	*	0.3	0.9

White, male

2005 4/.....	2.5	*	*	*	*	*	*	*	*	0.9
2004 4/.....	2.4	*	*	*	*	*	*	*	*	0.8
2003 4/.....	2.4	*	*	*	*	*	*	*	0.2	1.2
2002.....	2.6	*	*	*	*	*	*	*	0.4	1.5
2001.....	2.5	*	*	*	*	*	*	*	0.3	1.2
2000.....	2.4	*	*	*	*	*	*	*	0.3	1.3
1999.....	2.2	*	*	*	*	*	*	*	0.4	1.3

White, female

2005 4/.....	1.2	*	*	*	*	*	*	*	*	0.3
2004 4/.....	1.2	*	*	*	*	*	*	*	*	0.5
2003 4/.....	1.2	*	*	*	*	*	*	*	*	0.5
2002.....	1.3	*	*	*	*	*	*	*	*	0.5
2001.....	1.3	*	*	*	*	*	*	*	*	0.6
2000.....	1.3	*	*	*	*	*	*	*	*	0.6
1999.....	1.2	*	*	*	*	*	*	*	0.2	0.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Human immunodeficiency virus (HIV) disease (B20-B24)										
All races 3/, both sexes										
2005.....	4.2	*	*	*	*	0.1	0.6	2.1	4.5	8.1
2004.....	4.4	*	*	*	*	0.1	0.8	2.2	5.1	9.5
2003.....	4.7	*	*	*	0.1	0.2	0.7	2.3	5.5	11.2
2002.....	4.9	*	*	*	0.1	0.2	0.7	2.6	6.4	11.9
2001.....	5.0	*	*	*	0.1	0.2	0.9	3.1	7.3	12.3
2000.....	5.1	*	*	*	0.1	0.2	0.8	3.3	8.7	12.8
1999.....	5.3	*	0.2	0.2	0.1	0.2	0.9	3.9	9.5	13.0
All races 3/, male										
2005.....	6.3	*	*	*	*	*	0.7	2.4	5.5	11.2
2004.....	6.6	*	*	*	*	*	0.9	2.6	6.4	13.3
2003.....	7.1	*	*	*	*	*	0.8	2.8	7.3	15.9
2002.....	7.4	*	*	*	*	*	0.7	3.0	8.5	17.7
2001.....	7.6	*	*	*	*	0.2	0.8	3.7	10.2	18.4
2000.....	7.9	*	*	*	*	*	0.8	3.8	11.9	19.2
1999.....	8.2	*	*	0.2	*	*	0.8	5.1	13.6	19.5
All races 3/, female										
2005.....	2.2	*	*	*	*	*	0.5	1.7	3.5	4.9
2004.....	2.4	*	*	*	*	*	0.7	1.8	3.7	5.6
2003.....	2.4	*	*	*	*	*	0.7	1.8	3.7	6.5
2002.....	2.5	*	*	*	*	0.2	0.6	2.3	4.2	6.2
2001.....	2.5	*	*	*	*	0.2	1.0	2.5	4.4	6.3
2000.....	2.5	*	*	*	*	*	0.7	2.9	5.5	6.4
1999.....	2.5	*	*	*	*	*	1.0	2.7	5.4	6.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Human immunodeficiency virus (HIV) disease (B20-B24)

White, both sexes

2005 4/.....	2.2	*	*	*	*	*	0.3	0.8	2.1	4.2
2004 4/.....	2.4	*	*	*	*	*	0.2	0.8	2.3	5.4
2003 4/.....	2.5	*	*	*	*	*	0.3	0.9	2.9	6.1
2002.....	2.6	*	*	*	*	*	0.3	1.2	3.0	6.8
2001.....	2.7	*	*	*	*	*	0.3	1.3	3.8	7.1
2000.....	2.8	*	*	*	*	*	0.3	1.5	4.5	7.6
1999.....	3.0	*	*	*	*	*	0.4	2.0	5.4	7.9

White, male

2005 4/.....	3.7	*	*	*	*	*	0.4	1.1	3.0	6.8
2004 4/.....	3.9	*	*	*	*	*	0.4	1.2	3.6	8.4
2003 4/.....	4.2	*	*	*	*	*	0.3	1.4	4.5	9.7
2002.....	4.3	*	*	*	*	*	0.3	1.8	4.7	11.1
2001.....	4.5	*	*	*	*	*	0.4	1.8	5.8	11.6
2000.....	4.7	*	*	*	*	*	0.4	2.0	7.1	12.5
1999.....	5.0	*	*	*	*	*	0.4	2.8	8.6	13.0

White, female

2005 4/.....	0.8	*	*	*	*	*	*	0.5	1.2	1.6
2004 4/.....	0.8	*	*	*	*	*	*	0.4	1.0	2.4
2003 4/.....	0.9	*	*	*	*	*	*	0.5	1.3	2.4
2002.....	0.9	*	*	*	*	*	*	0.7	1.3	2.3
2001.....	0.9	*	*	*	*	*	0.3	0.8	1.6	2.4
2000.....	1.0	*	*	*	*	*	*	1.0	1.9	2.6
1999.....	1.0	*	*	*	*	*	0.3	1.0	2.1	2.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Human immunodeficiency virus (HIV) disease (B20-B24)										
Black, both sexes										
2005 4/.....	18.0	*	*	*	*	0.6	2.7	9.7	20.4	34.3
2004 4/.....	18.8	*	*	*	*	0.7	3.7	10.8	23.2	37.4
2003 4/.....	19.6	*	*	*	*	0.6	3.2	11.2	23.2	46.0
2002.....	20.8	*	*	*	*	0.8	3.3	11.7	29.1	48.2
2001.....	21.1	*	*	*	*	1.1	4.4	14.5	31.5	48.8
2000.....	21.4	*	*	*	*	0.7	3.7	15.2	37.1	49.8
1999.....	21.8	*	*	0.7	*	0.7	4.1	16.6	37.8	49.9
Black, male										
2005 4/.....	25.1	*	*	*	*	*	2.9	11.0	23.5	44.5
2004 4/.....	25.9	*	*	*	*	*	3.8	11.9	26.5	50.1
2003 4/.....	27.7	*	*	*	*	*	3.2	12.7	28.6	61.9
2002.....	29.5	*	*	*	*	*	3.3	11.7	36.6	68.4
2001.....	30.1	*	*	*	*	*	3.5	16.8	42.3	69.2
2000.....	31.0	*	*	*	*	*	3.6	16.4	47.9	71.2
1999.....	31.9	*	*	*	*	*	3.5	21.1	50.6	69.9
Black, female										
2005 4/.....	11.5	*	*	*	*	*	2.5	8.5	17.6	25.1
2004 4/.....	12.4	*	*	*	*	*	3.7	9.8	20.3	25.9
2003 4/.....	12.3	*	*	*	*	*	3.1	9.8	18.3	31.9
2002.....	12.8	*	*	*	*	*	3.3	11.7	22.2	30.3
2001.....	12.9	*	*	*	*	*	5.2	12.4	21.8	30.7
2000.....	12.8	*	*	*	*	*	3.8	14.1	27.4	30.8
1999.....	12.6	*	*	*	*	*	4.7	12.5	26.3	32.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1, A85.8,A86-B04,B06-B09,B25-B49,B55-B99)

All races 3/, both sexes

2005.....	2.6	4.1	0.4	0.2	0.1	0.2	0.2	0.2	0.4	0.7
2004.....	2.6	4.1	0.4	0.1	0.1	0.2	0.3	0.3	0.4	0.7
2003.....	2.6	4.3	0.5	0.1	0.1	0.2	0.2	0.3	0.4	0.7
2002.....	2.3	5.6	0.4	0.2	0.2	0.2	0.3	0.4	0.5	0.7
2001.....	2.0	4.7	0.4	0.2	0.1	0.2	0.2	0.3	0.4	0.7
2000.....	2.0	4.4	0.5	0.1	0.2	0.2	0.3	0.3	0.4	0.7
1999.....	2.6	5.3	0.5	0.2	0.2	0.2	0.3	0.5	0.7	1.0

All races 3/, male

2005.....	2.9	4.8	0.5	*	*	0.2	0.2	0.3	0.5	0.9
2004.....	2.8	4.8	0.4	*	*	0.2	0.4	0.2	0.6	0.8
2003.....	2.8	4.2	0.5	*	*	0.3	0.2	0.3	0.5	0.9
2002.....	2.6	6.1	0.5	*	0.2	0.2	0.3	0.4	0.5	0.9
2001.....	2.1	4.9	0.5	*	*	0.2	0.3	0.3	0.5	0.8
2000.....	2.1	4.8	0.5	*	*	0.2	0.3	0.3	0.4	0.9
1999.....	2.7	6.1	0.6	0.2	0.2	0.2	0.3	0.5	0.8	1.3

All races 3/, female

2005.....	2.3	3.4	0.3	0.2	*	0.2	0.2	0.2	0.3	0.5
2004.....	2.3	3.4	0.4	*	*	0.2	*	0.4	0.3	0.6
2003.....	2.3	4.5	0.5	*	*	*	*	0.3	0.3	0.6
2002.....	2.1	5.0	0.3	0.3	*	0.2	0.2	0.3	0.4	0.6
2001.....	1.9	4.6	0.4	0.2	*	*	*	0.3	0.4	0.6
2000.....	1.9	3.9	0.4	*	*	*	0.2	0.3	0.4	0.5
1999.....	2.5	4.5	0.4	0.2	*	0.2	0.2	0.5	0.6	0.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1, A85.8,A86-B04,B06-B09,B25-B49,B55-B99)

White, both sexes

2005 4/.....	2.7	3.7	0.4	0.2	0.2	0.2	0.2	0.2	0.3	0.6
2004 4/.....	2.6	3.7	0.4	*	*	0.2	0.3	0.2	0.3	0.7
2003 4/.....	2.6	3.5	0.4	0.1	*	0.2	0.2	0.2	0.4	0.6
2002.....	2.4	5.0	0.4	0.2	0.2	0.2	0.2	0.3	0.4	0.6
2001.....	2.0	3.8	0.4	0.2	0.1	0.2	0.2	0.3	0.3	0.6
2000.....	2.0	3.4	0.4	0.2	0.2	0.2	0.3	0.2	0.3	0.6
1999.....	2.5	4.5	0.5	0.2	0.2	0.2	0.2	0.4	0.5	0.8

White, male

2005 4/.....	3.0	4.2	0.4	*	*	*	*	0.2	0.4	0.8
2004 4/.....	2.9	4.2	0.4	*	*	*	0.3	*	0.4	0.7
2003 4/.....	2.8	3.2	0.4	*	*	0.3	*	0.3	0.4	0.8
2002.....	2.6	5.3	0.5	*	0.2	*	*	0.3	0.5	0.7
2001.....	2.1	3.9	0.4	*	*	*	*	0.3	0.4	0.7
2000.....	2.1	3.9	0.4	*	*	*	0.3	*	0.3	0.7
1999.....	2.6	5.6	0.6	*	*	*	*	0.3	0.6	1.2

White, female

2005 4/.....	2.4	3.1	*	*	*	*	*	*	*	0.4
2004 4/.....	2.4	3.0	0.3	*	*	*	*	0.3	*	0.6
2003 4/.....	2.4	3.9	0.5	*	*	*	*	*	0.3	0.5
2002.....	2.2	4.6	*	*	*	*	*	0.3	0.4	0.4
2001.....	1.9	3.8	*	*	*	*	*	*	0.2	0.5
2000.....	1.9	2.9	0.4	*	*	*	*	*	0.3	0.4
1999.....	2.5	3.4	*	*	*	*	*	0.4	0.5	0.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1, A85.8,A86-B04,B06-B09,B25-B49,B55-B99)

Black, both sexes

2005 4/.....	2.5	6.5	*	*	*	*	*	*	1.1	1.2
2004 4/.....	2.5	6.7	*	*	*	*	*	*	0.9	1.1
2003 4/.....	2.5	8.0	1.0	*	*	*	*	*	0.7	1.4
2002.....	2.5	8.7	*	*	*	*	*	*	1.0	1.8
2001.....	2.2	9.5	*	*	*	*	*	*	1.0	1.4
2000.....	2.3	9.9	*	*	*	*	*	*	1.2	1.6
1999.....	3.5	9.9	0.9	*	*	*	*	1.1	2.1	2.5

Black, male

2005 4/.....	2.8	8.0	*	*	*	*	*	*	*	1.5
2004 4/.....	2.9	7.4	*	*	*	*	*	*	*	*
2003 4/.....	2.9	8.9	*	*	*	*	*	*	*	1.6
2002.....	2.9	9.6	*	*	*	*	*	*	*	2.1
2001.....	2.4	10.8	*	*	*	*	*	*	*	1.7
2000.....	2.5	9.9	*	*	*	*	*	*	*	2.2
1999.....	3.6	10.1	*	*	*	*	*	*	2.6	2.7

Black, female

2005 4/.....	2.2	*	*	*	*	*	*	*	*	*
2004 4/.....	2.2	*	*	*	*	*	*	*	*	*
2003 4/.....	2.2	7.1	*	*	*	*	*	*	*	*
2002.....	2.1	7.9	*	*	*	*	*	*	*	1.6
2001.....	2.0	8.2	*	*	*	*	*	*	*	*
2000.....	2.1	9.9	*	*	*	*	*	*	*	*
1999.....	3.4	9.7	*	*	*	*	*	*	1.6	2.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasms (C00-C97)										
All races 3/, both sexes										
2005.....	188.7	1.8	2.3	2.5	2.5	3.5	4.7	6.7	11.3	20.8
2004.....	188.6	1.8	2.5	2.7	2.3	3.5	4.7	6.6	11.5	21.5
2003.....	191.5	1.9	2.5	2.6	2.6	3.4	4.6	6.8	11.8	23.0
2002.....	193.2	1.8	2.6	2.7	2.5	3.5	5.0	6.6	12.5	23.5
2001.....	194.4	1.6	2.7	2.4	2.5	3.6	4.9	7.3	12.6	24.6
2000.....	196.5	2.4	2.7	2.4	2.6	3.7	5.1	7.0	12.5	24.5
1999.....	197.0	1.8	2.7	2.5	2.5	3.7	5.3	7.3	12.5	25.2
All races 3/, male										
2005.....	198.9	2.1	2.6	2.6	2.8	4.1	5.5	7.3	10.4	17.4
2004.....	198.4	1.8	2.6	3.0	2.5	4.1	5.5	6.8	10.4	18.1
2003.....	201.3	1.7	2.8	2.9	2.7	4.2	5.1	6.8	10.9	19.3
2002.....	203.8	2.0	2.7	2.9	2.8	4.0	5.7	7.0	11.3	19.8
2001.....	205.3	1.5	2.9	2.4	2.6	4.3	5.7	7.2	11.2	21.0
2000.....	207.2	2.6	3.0	2.6	2.9	4.3	5.9	7.1	11.3	21.2
1999.....	208.9	1.9	2.9	2.6	2.7	4.3	6.1	7.6	11.6	21.6
All races 3/, female										
2005.....	178.8	1.5	2.0	2.3	2.2	2.8	3.9	6.0	12.2	24.3
2004.....	179.1	1.9	2.4	2.3	2.2	2.9	3.8	6.3	12.6	24.9
2003.....	182.0	2.1	2.1	2.3	2.5	2.5	4.2	6.8	12.6	26.7
2002.....	183.0	1.6	2.4	2.5	2.2	3.0	4.2	6.2	13.7	27.2
2001.....	183.9	1.8	2.5	2.4	2.3	2.9	4.1	7.4	14.0	28.1
2000.....	186.2	2.3	2.5	2.2	2.2	3.0	4.2	7.0	13.7	27.8
1999.....	185.6	1.7	2.5	2.3	2.3	3.0	4.4	7.0	13.4	28.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasms (C00-C97)										
White, both sexes										
2005 4/.....	200.8	2.0	2.3	2.5	2.5	3.6	4.7	6.3	10.8	20.1
2004 4/.....	200.7	2.0	2.5	2.7	2.3	3.6	4.5	6.4	11.1	20.8
2003 4/.....	203.7	2.1	2.6	2.8	2.6	3.5	4.7	6.7	11.3	22.2
2002.....	205.5	1.8	2.5	2.8	2.5	3.6	5.2	6.5	12.1	22.7
2001.....	206.4	1.7	3.0	2.5	2.4	3.7	5.0	7.2	12.2	23.7
2000.....	208.6	2.5	2.8	2.5	2.6	3.6	5.1	7.0	12.1	23.3
1999.....	208.7	1.8	2.9	2.5	2.4	3.6	5.2	7.3	12.1	24.0
White, male										
2005 4/.....	210.6	2.3	2.6	2.6	2.8	4.2	5.4	7.0	10.0	17.1
2004 4/.....	209.9	1.9	2.5	3.1	2.5	4.2	5.3	6.8	10.4	18.0
2003 4/.....	213.1	1.8	2.9	3.1	2.7	4.3	5.1	7.0	10.6	19.3
2002.....	215.5	2.2	2.7	2.9	2.7	4.2	5.9	6.9	11.1	19.2
2001.....	216.4	1.5	3.1	2.4	2.4	4.4	5.8	7.2	11.2	20.5
2000.....	218.1	2.8	3.1	2.6	2.9	4.2	5.9	7.1	11.2	20.5
1999.....	219.5	2.0	3.2	2.7	2.6	4.1	5.9	7.6	11.3	21.1
White, female										
2005 4/.....	191.1	1.7	2.0	2.4	2.3	2.9	3.9	5.5	11.7	23.3
2004 4/.....	191.7	2.1	2.5	2.3	2.1	2.9	3.7	6.0	11.9	23.7
2003 4/.....	194.6	2.3	2.3	2.4	2.5	2.6	4.2	6.5	12.0	25.2
2002.....	195.8	1.4	2.3	2.6	2.4	3.0	4.4	6.1	13.2	26.3
2001.....	196.7	2.0	2.8	2.5	2.4	2.9	4.1	7.2	13.3	27.1
2000.....	199.4	2.3	2.4	2.3	2.3	3.0	4.2	7.0	13.0	26.3
1999.....	198.2	1.5	2.7	2.2	2.2	3.0	4.3	7.0	12.9	27.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasms (C00-C97)										
Black, both sexes										
2005 4/.....	161.7	*	2.4	2.6	2.6	3.0	5.1	9.2	15.6	28.1
2004 4/.....	161.9	*	2.5	2.5	2.6	3.8	5.9	8.9	15.1	28.4
2003 4/.....	164.3	*	2.2	2.3	2.8	2.9	4.9	8.5	15.7	31.0
2002.....	165.9	*	2.9	2.6	2.5	3.6	4.5	8.0	16.3	31.7
2001.....	167.1	*	1.9	2.4	2.8	3.6	5.1	9.1	16.4	32.7
2000.....	169.3	*	2.8	2.1	2.3	4.4	5.9	8.0	15.8	34.5
1999.....	171.3	*	2.0	2.4	3.0	4.5	6.6	8.6	15.8	35.8
Black, male										
2005 4/.....	175.4	*	2.8	2.9	3.2	3.4	6.1	9.3	14.7	21.7
2004 4/.....	176.2	*	2.8	2.5	2.4	4.1	6.8	7.9	12.2	19.7
2003 4/.....	178.3	*	2.4	2.4	3.1	3.5	5.6	7.0	13.5	21.7
2002.....	181.5	*	2.6	3.3	2.9	4.1	5.0	8.2	14.1	25.6
2001.....	184.5	*	2.4	2.5	3.4	4.3	5.9	8.5	12.4	26.8
2000.....	188.5	*	2.7	2.4	2.7	5.2	7.1	7.3	12.9	27.6
1999.....	191.0	*	2.1	2.3	3.3	5.6	7.3	8.8	14.0	27.8
Black, female										
2005 4/.....	149.1	*	2.1	2.3	2.0	2.7	4.1	9.0	16.3	33.8
2004 4/.....	148.9	*	2.2	2.5	2.7	3.5	5.0	9.9	17.8	36.2
2003 4/.....	151.4	*	2.1	2.1	2.5	2.3	4.3	10.0	17.7	39.2
2002.....	151.7	*	3.2	1.9	2.1	3.1	4.0	7.9	18.3	37.1
2001.....	151.3	*	*	2.2	2.2	2.9	4.3	9.7	20.0	37.9
2000.....	151.8	*	2.9	1.8	2.0	3.5	4.7	8.5	18.3	40.5
1999.....	153.3	*	1.9	2.6	2.7	3.4	5.9	8.3	17.5	43.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Malignant neoplasm of stomach (C16)

All races 3/, both sexes

2005.....	3.9	*	*	*	*	*	*	0.1	0.4	0.7
2004.....	4.0	*	*	*	*	*	*	0.1	0.5	0.7
2003.....	4.2	*	*	*	*	*	*	0.2	0.4	0.8
2002.....	4.2	*	*	*	*	*	0.1	0.3	0.4	0.7
2001.....	4.3	*	*	*	*	*	*	0.2	0.4	0.8
2000.....	4.5	*	*	*	*	*	*	0.2	0.4	0.7
1999.....	4.6	*	*	*	*	*	*	0.2	0.4	0.8

All races 3/, male

2005.....	4.6	*	*	*	*	*	*	*	0.4	0.8
2004.....	4.8	*	*	*	*	*	*	*	0.4	0.7
2003.....	4.9	*	*	*	*	*	*	*	0.4	0.8
2002.....	5.0	*	*	*	*	*	*	0.3	0.5	0.7
2001.....	5.1	*	*	*	*	*	*	0.2	0.5	0.9
2000.....	5.3	*	*	*	*	*	*	0.2	0.5	0.9
1999.....	5.5	*	*	*	*	*	*	0.2	0.5	0.9

All races 3/, female

2005.....	3.2	*	*	*	*	*	*	*	0.4	0.5
2004.....	3.3	*	*	*	*	*	*	*	0.5	0.6
2003.....	3.4	*	*	*	*	*	*	*	0.4	0.8
2002.....	3.5	*	*	*	*	*	*	*	0.4	0.7
2001.....	3.5	*	*	*	*	*	*	*	0.4	0.7
2000.....	3.7	*	*	*	*	*	*	*	0.4	0.6
1999.....	3.7	*	*	*	*	*	*	0.3	0.3	0.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Malignant neoplasms of colon, rectum and anus (C18-C21)

All races 3/, both sexes

2005.....	18.0	*	*	*	*	*	0.2	0.4	1.1	1.9
2004.....	18.3	*	*	*	*	*	0.2	0.4	0.9	2.0
2003.....	19.2	*	*	*	*	*	0.2	0.5	0.9	2.1
2002.....	19.7	*	*	*	*	*	0.2	0.5	1.0	2.0
2001.....	20.0	*	*	*	*	*	0.2	0.4	1.1	2.3
2000.....	20.4	*	*	*	*	*	0.1	0.4	0.9	2.1
1999.....	20.5	*	*	*	*	*	0.2	0.4	1.0	1.9

All races 3/, male

2005.....	18.4	*	*	*	*	*	0.2	0.5	1.1	2.1
2004.....	18.6	*	*	*	*	*	0.2	0.5	1.0	2.1
2003.....	19.6	*	*	*	*	*	*	0.6	0.9	2.2
2002.....	20.1	*	*	*	*	*	0.3	0.5	1.1	2.1
2001.....	20.2	*	*	*	*	*	*	0.4	1.1	2.6
2000.....	20.6	*	*	*	*	*	*	0.4	1.1	2.4
1999.....	20.6	*	*	*	*	*	0.2	0.6	1.0	2.0

All races 3/, female

2005.....	17.6	*	*	*	*	*	*	0.4	1.0	1.7
2004.....	18.0	*	*	*	*	*	*	0.4	0.9	2.0
2003.....	18.9	*	*	*	*	*	*	0.4	0.9	2.0
2002.....	19.2	*	*	*	*	*	*	0.4	0.9	1.9
2001.....	19.8	*	*	*	*	*	*	0.5	1.0	1.9
2000.....	20.2	*	*	*	*	*	*	0.4	0.8	1.8
1999.....	20.3	*	*	*	*	*	*	0.3	0.9	1.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Malignant neoplasms of colon, rectum and anus (C18-C21)

White, both sexes

2005 4/.....	18.7	*	*	*	*	*	0.2	0.4	1.0	1.8
2004 4/.....	19.2	*	*	*	*	*	0.2	0.4	0.9	1.9
2003 4/.....	20.2	*	*	*	*	*	0.1	0.5	0.9	1.9
2002.....	20.7	*	*	*	*	*	0.2	0.5	1.0	1.9
2001.....	21.0	*	*	*	*	*	0.2	0.4	1.0	2.1
2000.....	21.5	*	*	*	*	*	*	0.4	0.9	1.9
1999.....	21.6	*	*	*	*	*	0.2	0.4	0.9	1.8

White, male

2005 4/.....	19.2	*	*	*	*	*	*	0.5	1.0	2.1
2004 4/.....	19.5	*	*	*	*	*	*	0.5	0.9	2.1
2003 4/.....	20.5	*	*	*	*	*	*	0.6	0.9	2.0
2002.....	21.1	*	*	*	*	*	0.2	0.5	1.0	2.0
2001.....	21.2	*	*	*	*	*	*	0.4	1.1	2.5
2000.....	21.6	*	*	*	*	*	*	0.4	1.0	2.2
1999.....	21.8	*	*	*	*	*	*	0.5	0.9	1.9

White, female

2005 4/.....	18.3	*	*	*	*	*	*	0.3	1.1	1.5
2004 4/.....	18.9	*	*	*	*	*	*	0.4	0.8	1.7
2003 4/.....	19.8	*	*	*	*	*	*	0.4	0.9	1.8
2002.....	20.3	*	*	*	*	*	*	0.4	0.9	1.9
2001.....	20.7	*	*	*	*	*	*	0.4	0.9	1.8
2000.....	21.4	*	*	*	*	*	*	0.3	0.7	1.6
1999.....	21.4	*	*	*	*	*	*	0.3	0.9	1.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasm of pancreas (C25)										
All races 3/, both sexes										
2005.....	11.1	*	*	*	*	*	*	0.1	0.2	0.6
2004.....	10.8	*	*	*	*	*	*	*	0.2	0.6
2003.....	10.6	*	*	*	*	*	*	0.1	0.2	0.7
2002.....	10.5	*	*	*	*	*	*	*	0.2	0.7
2001.....	10.5	*	*	*	*	*	*	*	0.2	0.8
2000.....	10.4	*	*	*	*	*	*	*	0.2	0.6
1999.....	10.4	*	*	*	*	*	*	*	0.2	0.8
All races 3/, male										
2005.....	11.1	*	*	*	*	*	*	*	0.2	0.8
2004.....	10.9	*	*	*	*	*	*	*	0.3	0.7
2003.....	10.5	*	*	*	*	*	*	*	0.2	0.9
2002.....	10.5	*	*	*	*	*	*	*	0.2	0.8
2001.....	10.3	*	*	*	*	*	*	*	0.2	0.9
2000.....	10.3	*	*	*	*	*	*	*	0.2	0.8
1999.....	10.4	*	*	*	*	*	*	*	0.3	0.9
All races 3/, female										
2005.....	11.0	*	*	*	*	*	*	*	*	0.5
2004.....	10.7	*	*	*	*	*	*	*	*	0.5
2003.....	10.6	*	*	*	*	*	*	*	*	0.5
2002.....	10.5	*	*	*	*	*	*	*	*	0.6
2001.....	10.6	*	*	*	*	*	*	*	*	0.6
2000.....	10.5	*	*	*	*	*	*	*	*	0.5
1999.....	10.5	*	*	*	*	*	*	*	0.2	0.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasms of trachea, bronchus and lung (C33-C34)										
All races 3/, both sexes										
2005.....	53.7	*	*	*	*	*	*	0.2	0.5	2.1
2004.....	53.8	*	*	*	*	*	*	0.2	0.5	2.2
2003.....	54.4	*	*	*	*	*	*	0.2	0.6	2.4
2002.....	54.7	*	*	*	*	*	*	0.2	0.6	2.7
2001.....	54.8	*	*	*	*	*	*	0.2	0.6	2.9
2000.....	55.3	*	*	*	*	*	*	0.2	0.8	2.9
1999.....	54.5	*	*	*	*	*	*	0.2	0.7	2.9
All races 3/, male										
2005.....	61.8	*	*	*	*	*	*	*	0.6	2.2
2004.....	62.0	*	*	*	*	*	*	0.3	0.6	2.2
2003.....	62.9	*	*	*	*	*	*	*	0.7	2.4
2002.....	63.7	*	*	*	*	*	*	*	0.7	2.6
2001.....	64.7	*	*	*	*	*	*	*	0.6	2.8
2000.....	65.5	*	*	*	*	*	*	0.3	0.7	3.3
1999.....	65.4	*	*	*	*	*	*	0.2	0.7	2.8
All races 3/, female										
2005.....	45.9	*	*	*	*	*	*	0.2	0.4	2.0
2004.....	45.9	*	*	*	*	*	*	*	0.5	2.2
2003.....	46.1	*	*	*	*	*	*	0.2	0.5	2.5
2002.....	46.0	*	*	*	*	*	*	*	0.6	2.7
2001.....	45.3	*	*	*	*	*	*	*	0.6	2.9
2000.....	45.4	*	*	*	*	*	*	*	0.8	2.6
1999.....	44.1	*	*	*	*	*	*	*	0.6	3.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasms of trachea, bronchus and lung (C33-C34)										
White, both sexes										
2005 4/.....	58.1	*	*	*	*	*	*	0.2	0.4	2.1
2004 4/.....	58.1	*	*	*	*	*	*	0.2	0.5	2.2
2003 4/.....	58.7	*	*	*	*	*	*	0.2	0.6	2.4
2002.....	59.1	*	*	*	*	*	*	0.2	0.6	2.6
2001.....	59.1	*	*	*	*	*	*	0.2	0.6	2.9
2000.....	59.5	*	*	*	*	*	*	0.2	0.8	2.7
1999.....	58.5	*	*	*	*	*	*	0.2	0.7	2.7
White, male										
2005 4/.....	65.9	*	*	*	*	*	*	*	0.6	2.2
2004 4/.....	65.9	*	*	*	*	*	*	0.3	0.5	2.1
2003 4/.....	66.9	*	*	*	*	*	*	*	0.6	2.3
2002.....	67.7	*	*	*	*	*	*	*	0.7	2.5
2001.....	68.6	*	*	*	*	*	*	*	0.6	2.8
2000.....	69.4	*	*	*	*	*	*	0.3	0.7	3.0
1999.....	69.1	*	*	*	*	*	*	*	0.8	2.6
White, female										
2005 4/.....	50.4	*	*	*	*	*	*	*	0.3	2.0
2004 4/.....	50.4	*	*	*	*	*	*	*	0.4	2.3
2003 4/.....	50.7	*	*	*	*	*	*	*	0.6	2.4
2002.....	50.7	*	*	*	*	*	*	*	0.5	2.8
2001.....	49.8	*	*	*	*	*	*	*	0.6	3.0
2000.....	49.9	*	*	*	*	*	*	*	0.8	2.4
1999.....	48.2	*	*	*	*	*	*	*	0.6	2.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant melanoma of skin (C43)										
All races 3/, both sexes										
2005.....	2.8	*	*	*	*	*	0.2	0.5	0.6	0.9
2004.....	2.7	*	*	*	*	*	0.1	0.3	0.6	0.9
2003.....	2.7	*	*	*	*	*	0.1	0.3	0.6	0.9
2002.....	2.6	*	*	*	*	*	0.2	0.4	0.6	1.0
2001.....	2.6	*	*	*	*	*	0.2	0.4	0.6	1.0
2000.....	2.6	*	*	*	*	*	0.2	0.3	0.6	1.0
1999.....	2.6	*	*	*	*	*	0.2	0.3	0.6	1.0
All races 3/, male										
2005.....	3.6	*	*	*	*	*	*	0.6	0.8	0.9
2004.....	3.5	*	*	*	*	*	*	0.4	0.7	1.0
2003.....	3.5	*	*	*	*	*	*	0.4	0.8	1.1
2002.....	3.3	*	*	*	*	*	*	0.4	0.8	1.1
2001.....	3.4	*	*	*	*	*	*	0.4	0.6	1.2
2000.....	3.3	*	*	*	*	*	*	0.3	0.7	1.0
1999.....	3.3	*	*	*	*	*	*	0.4	0.7	1.3
All races 3/, female										
2005.....	2.0	*	*	*	*	*	*	0.3	0.5	0.8
2004.....	1.9	*	*	*	*	*	*	0.3	0.5	0.8
2003.....	1.9	*	*	*	*	*	*	0.2	0.5	0.7
2002.....	1.9	*	*	*	*	*	*	0.3	0.4	0.9
2001.....	1.9	*	*	*	*	*	*	0.3	0.5	0.7
2000.....	2.0	*	*	*	*	*	*	0.3	0.5	1.0
1999.....	1.9	*	*	*	*	*	*	0.2	0.6	0.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Malignant melanoma of skin (C43)

White, both sexes

2005 4/.....	3.4	*	*	*	*	*	0.2	0.6	0.8	1.0
2004 4/.....	3.3	*	*	*	*	*	0.2	0.4	0.8	1.1
2003 4/.....	3.2	*	*	*	*	*	0.2	0.4	0.8	1.1
2002.....	3.1	*	*	*	*	*	0.2	0.5	0.7	1.2
2001.....	3.2	*	*	*	*	*	0.2	0.5	0.7	1.2
2000.....	3.1	*	*	*	*	*	0.2	0.4	0.8	1.2
1999.....	3.1	*	*	*	*	*	0.2	0.4	0.8	1.3

White, male

2005 4/.....	4.4	*	*	*	*	*	*	0.7	0.9	1.1
2004 4/.....	4.3	*	*	*	*	*	*	0.5	0.9	1.3
2003 4/.....	4.2	*	*	*	*	*	*	0.5	0.9	1.3
2002.....	4.0	*	*	*	*	*	*	0.5	1.0	1.3
2001.....	4.1	*	*	*	*	*	*	0.5	0.7	1.4
2000.....	4.0	*	*	*	*	*	*	0.4	0.9	1.2
1999.....	4.0	*	*	*	*	*	*	0.5	0.8	1.6

White, female

2005 4/.....	2.4	*	*	*	*	*	*	0.4	0.6	1.0
2004 4/.....	2.3	*	*	*	*	*	*	0.4	0.6	1.0
2003 4/.....	2.3	*	*	*	*	*	*	0.3	0.6	0.9
2002.....	2.3	*	*	*	*	*	*	0.4	0.5	1.1
2001.....	2.3	*	*	*	*	*	*	0.4	0.6	0.9
2000.....	2.3	*	*	*	*	*	*	0.3	0.6	1.2
1999.....	2.3	*	*	*	*	*	*	0.3	0.7	0.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Malignant neoplasm of breast (C50)

All races 3/, both sexes

2005.....	14.0	*	*	*	*	*	*	0.4	1.4	3.8
2004.....	14.1	*	*	*	*	*	*	0.4	1.6	3.9
2003.....	14.4	*	*	*	*	*	*	0.3	1.7	4.3
2002.....	14.5	*	*	*	*	*	*	0.3	1.7	4.1
2001.....	14.7	*	*	*	*	*	*	0.5	1.8	4.3
2000.....	15.0	*	*	*	*	*	*	0.4	1.9	4.3
1999.....	14.9	*	*	*	*	*	*	0.4	1.7	4.3

All races 3/, male

2005.....	0.3	*	*	*	*	*	*	*	*	*
2004.....	0.3	*	*	*	*	*	*	*	*	*
2003.....	0.3	*	*	*	*	*	*	*	*	*
2002.....	0.3	*	*	*	*	*	*	*	*	*
2001.....	0.3	*	*	*	*	*	*	*	*	*
2000.....	0.3	*	*	*	*	*	*	*	*	*
1999.....	0.3	*	*	*	*	*	*	*	*	*

All races 3/, female

2005.....	27.3	*	*	*	*	*	*	0.8	2.8	7.7
2004.....	27.5	*	*	*	*	*	*	0.8	3.2	7.8
2003.....	28.2	*	*	*	*	*	*	0.6	3.4	8.5
2002.....	28.3	*	*	*	*	*	*	0.7	3.5	8.2
2001.....	28.6	*	*	*	*	*	*	0.9	3.6	8.6
2000.....	29.2	*	*	*	*	*	*	0.8	3.8	8.5
1999.....	28.9	*	*	*	*	*	*	0.9	3.4	8.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Malignant neoplasm of breast (C50)

White, both sexes

2005 4/.....	14.4	*	*	*	*	*	*	0.3	1.2	3.3
2004 4/.....	14.6	*	*	*	*	*	*	0.3	1.3	3.4
2003 4/.....	15.0	*	*	*	*	*	*	0.2	1.5	3.8
2002.....	15.1	*	*	*	*	*	*	0.3	1.5	3.6
2001.....	15.2	*	*	*	*	*	*	0.4	1.5	3.9
2000.....	15.7	*	*	*	*	*	*	0.4	1.6	3.9
1999.....	15.5	*	*	*	*	*	*	0.4	1.5	3.7

White, male

2005 4/.....	0.3	*	*	*	*	*	*	*	*	*
2004 4/.....	0.3	*	*	*	*	*	*	*	*	*
2003 4/.....	0.3	*	*	*	*	*	*	*	*	*
2002.....	0.3	*	*	*	*	*	*	*	*	*
2001.....	0.3	*	*	*	*	*	*	*	*	*
2000.....	0.3	*	*	*	*	*	*	*	*	*
1999.....	0.3	*	*	*	*	*	*	*	*	*

White, female

2005 4/.....	28.3	*	*	*	*	*	*	0.6	2.4	6.7
2004 4/.....	28.6	*	*	*	*	*	*	0.6	2.7	6.9
2003 4/.....	29.3	*	*	*	*	*	*	0.4	3.0	7.7
2002.....	29.5	*	*	*	*	*	*	0.5	3.0	7.2
2001.....	29.8	*	*	*	*	*	*	0.8	3.0	7.8
2000.....	30.7	*	*	*	*	*	*	0.8	3.3	7.7
1999.....	30.2	*	*	*	*	*	*	0.8	3.0	7.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Malignant neoplasm of breast (C50)

Black, both sexes

2005 4/.....	15.0	*	*	*	*	*	*	0.8	2.9	7.8
2004 4/.....	14.5	*	*	*	*	*	*	0.9	3.2	8.1
2003 4/.....	15.1	*	*	*	*	*	*	0.9	3.4	8.0
2002.....	14.9	*	*	*	*	*	*	0.8	3.7	8.1
2001.....	15.0	*	*	*	*	*	*	1.1	4.0	7.5
2000.....	14.8	*	*	*	*	*	*	0.7	3.7	7.7
1999.....	15.0	*	*	*	*	*	*	0.9	3.7	8.8

Black, male

2005 4/.....	0.3	*	*	*	*	*	*	*	*	*
2004 4/.....	0.3	*	*	*	*	*	*	*	*	*
2003 4/.....	0.3	*	*	*	*	*	*	*	*	*
2002.....	0.4	*	*	*	*	*	*	*	*	*
2001.....	0.4	*	*	*	*	*	*	*	*	*
2000.....	0.4	*	*	*	*	*	*	*	*	*
1999.....	0.4	*	*	*	*	*	*	*	*	*

Black, female

2005 4/.....	28.4	*	*	*	*	*	*	1.6	5.6	14.7
2004 4/.....	27.5	*	*	*	*	*	*	1.8	6.2	15.1
2003 4/.....	28.5	*	*	*	*	*	*	1.6	6.5	15.1
2002.....	28.2	*	*	*	*	*	*	1.6	6.9	15.1
2001.....	28.3	*	*	*	*	*	*	2.2	7.6	14.2
2000.....	27.9	*	*	*	*	*	*	*	7.1	14.3
1999.....	28.2	*	*	*	*	*	*	1.8	6.9	16.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Malignant neoplasm of cervix uteri (C53)

All races 3/, both sexes

2005.....	1.3	*	*	*	*	*	*	0.2	0.6	1.1
2004.....	1.3	*	*	*	*	*	*	0.4	0.7	1.1
2003.....	1.3	*	*	*	*	*	*	0.3	0.7	1.2
2002.....	1.4	*	*	*	*	*	*	0.2	0.8	1.3
2001.....	1.4	*	*	*	*	*	*	0.3	0.8	1.3
2000.....	1.5	*	*	*	*	*	*	0.4	0.7	1.3
1999.....	1.5	*	*	*	*	*	*	0.3	0.8	1.3

All races 3/, male

2005.....
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

All races 3/, female

2005.....	2.6	*	*	*	*	*	*	0.4	1.3	2.3
2004.....	2.6	*	*	*	*	*	*	0.7	1.4	2.2
2003.....	2.7	*	*	*	*	*	*	0.6	1.4	2.3
2002.....	2.7	*	*	*	*	*	*	0.5	1.5	2.5
2001.....	2.8	*	*	*	*	*	*	0.5	1.5	2.6
2000.....	2.9	*	*	*	*	*	*	0.8	1.5	2.6
1999.....	3.0	*	*	*	*	*	*	0.6	1.5	2.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Malignant neoplasm of cervix uteri (C53)

White, both sexes

2005 4/.....	1.2	*	*	*	*	*	*	0.2	0.7	1.1
2004 4/.....	1.2	*	*	*	*	*	*	0.3	0.7	1.1
2003 4/.....	1.2	*	*	*	*	*	*	0.3	0.6	1.1
2002.....	1.3	*	*	*	*	*	*	0.2	0.8	1.2
2001.....	1.3	*	*	*	*	*	*	0.3	0.7	1.3
2000.....	1.4	*	*	*	*	*	*	0.4	0.7	1.2
1999.....	1.4	*	*	*	*	*	*	0.3	0.7	1.2

White, male

2005 4/.....
2004 4/.....
2003 4/.....
2002.....
2001.....
2000.....
1999.....

White, female

2005 4/.....	2.5	*	*	*	*	*	*	0.4	1.4	2.2
2004 4/.....	2.4	*	*	*	*	*	*	0.6	1.4	2.2
2003 4/.....	2.5	*	*	*	*	*	*	0.5	1.2	2.1
2002.....	2.5	*	*	*	*	*	*	0.5	1.5	2.4
2001.....	2.7	*	*	*	*	*	*	0.6	1.5	2.5
2000.....	2.7	*	*	*	*	*	*	0.8	1.4	2.4
1999.....	2.8	*	*	*	*	*	*	0.7	1.5	2.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasm of cervix uteri (C53)

Black, both sexes

2005 4/.....	2.0	*	*	*	*	*	*	*	0.8	1.6
2004 4/.....	2.0	*	*	*	*	*	*	0.7	0.9	1.6
2003 4/.....	2.1	*	*	*	*	*	*	*	1.2	2.0
2002.....	2.2	*	*	*	*	*	*	*	1.1	2.1
2001.....	2.1	*	*	*	*	*	*	*	1.1	1.8
2000.....	2.4	*	*	*	*	*	*	*	1.4	2.1
1999.....	2.3	*	*	*	*	*	*	*	1.0	2.1

Black, male

2005 4/.....
2004 4/.....
2003 4/.....
2002.....
2001.....
2000.....
1999.....

Black, female

2005 4/.....	3.8	*	*	*	*	*	*	*	1.5	3.0
2004 4/.....	3.9	*	*	*	*	*	*	1.4	1.7	3.0
2003 4/.....	4.0	*	*	*	*	*	*	*	2.2	3.8
2002.....	4.2	*	*	*	*	*	*	*	2.0	3.9
2001.....	4.0	*	*	*	*	*	*	*	2.1	3.5
2000.....	4.5	*	*	*	*	*	*	*	2.7	4.0
1999.....	4.4	*	*	*	*	*	*	*	1.9	3.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Malignant neoplasm of ovary (C56)

All races 3/, both sexes

2005.....	5.0	*	*	*	*	*	*	0.2	0.3	0.5
2004.....	5.0	*	*	*	*	*	0.1	0.2	0.3	0.6
2003.....	5.0	*	*	*	*	*	*	0.2	0.3	0.6
2002.....	5.1	*	*	*	*	*	0.1	0.2	0.3	0.7
2001.....	5.1	*	*	*	*	*	*	0.2	0.3	0.7
2000.....	5.0	*	*	*	*	*	*	0.2	0.3	0.6
1999.....	4.9	*	*	*	*	*	*	0.1	0.2	0.7

All races 3/, male

2005.....
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

All races 3/, female

2005.....	9.8	*	*	*	*	*	*	0.3	0.5	1.0
2004.....	9.9	*	*	*	*	*	0.2	0.3	0.6	1.2
2003.....	9.9	*	*	*	*	*	*	0.3	0.5	1.3
2002.....	10.0	*	*	*	*	*	0.3	0.4	0.6	1.4
2001.....	9.9	*	*	*	*	*	*	0.4	0.6	1.4
2000.....	9.8	*	*	*	*	*	*	0.4	0.6	1.3
1999.....	9.6	*	*	*	*	*	*	0.3	0.5	1.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Malignant neoplasm of ovary (C56)

White, both sexes

2005 4/.....	5.5	*	*	*	*	*	*	0.2	0.3	0.5
2004 4/.....	5.5	*	*	*	*	*	*	0.1	0.3	0.6
2003 4/.....	5.6	*	*	*	*	*	*	0.2	0.3	0.6
2002.....	5.6	*	*	*	*	*	0.1	0.2	0.3	0.7
2001.....	5.6	*	*	*	*	*	*	0.2	0.3	0.7
2000.....	5.5	*	*	*	*	*	*	0.2	0.3	0.6
1999.....	5.4	*	*	*	*	*	*	0.1	0.2	0.7

White, male

2005 4/.....
2004 4/.....
2003 4/.....
2002.....
2001.....
2000.....
1999.....

White, female

2005 4/.....	10.9	*	*	*	*	*	*	0.3	0.5	1.1
2004 4/.....	10.9	*	*	*	*	*	*	0.3	0.6	1.2
2003 4/.....	11.0	*	*	*	*	*	*	0.3	0.6	1.2
2002.....	11.1	*	*	*	*	*	0.3	0.4	0.6	1.4
2001.....	11.0	*	*	*	*	*	*	0.4	0.6	1.3
2000.....	10.9	*	*	*	*	*	*	0.4	0.6	1.3
1999.....	10.6	*	*	*	*	*	*	0.3	0.5	1.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)

All races 3/, both sexes

2005.....	4.4	*	0.5	1.0	0.7	0.6	0.6	0.8	1.1	1.6
2004.....	4.4	*	0.6	0.9	0.7	0.5	0.5	0.7	1.1	1.7
2003.....	4.4	*	0.7	0.9	0.8	0.6	0.6	0.9	1.2	1.9
2002.....	4.4	*	0.7	1.0	0.6	0.5	0.5	0.7	1.2	1.7
2001.....	4.4	*	0.7	0.8	0.6	0.5	0.6	0.7	1.4	1.8
2000.....	4.5	0.5	0.7	0.8	0.7	0.5	0.6	0.8	1.2	2.0
1999.....	4.6	*	0.7	0.9	0.6	0.6	0.6	0.9	1.3	1.9

All races 3/, male

2005.....	5.0	*	0.5	0.9	0.7	0.7	0.6	1.1	1.3	2.0
2004.....	4.9	*	0.7	1.0	0.7	0.6	0.7	0.8	1.3	2.1
2003.....	5.0	*	0.7	0.9	0.8	0.6	0.6	0.9	1.4	2.3
2002.....	5.0	*	0.8	1.1	0.7	0.6	0.7	0.7	1.5	2.0
2001.....	5.0	*	0.7	0.8	0.6	0.6	0.7	0.9	1.7	2.2
2000.....	5.1	*	0.7	0.8	0.8	0.6	0.7	1.0	1.6	2.6
1999.....	5.1	*	0.7	0.9	0.6	0.7	0.7	1.1	1.7	2.3

All races 3/, female

2005.....	3.9	*	0.6	1.0	0.6	0.4	0.5	0.5	0.8	1.2
2004.....	3.9	*	0.6	0.9	0.7	0.5	0.4	0.6	0.8	1.2
2003.....	3.9	*	0.6	0.9	0.7	0.5	0.5	0.9	0.9	1.5
2002.....	3.9	*	0.7	0.9	0.5	0.4	0.3	0.7	1.0	1.4
2001.....	3.9	*	0.6	0.9	0.6	0.4	0.6	0.5	1.1	1.4
2000.....	3.9	*	0.6	0.8	0.6	0.4	0.5	0.7	0.8	1.4
1999.....	4.1	*	0.8	0.9	0.7	0.5	0.5	0.8	0.9	1.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)

White, both sexes

2005 4/.....	5.0	*	0.5	0.9	0.7	0.6	0.6	0.9	1.2	1.8
2004 4/.....	4.9	*	0.7	1.0	0.6	0.6	0.6	0.7	1.2	1.9
2003 4/.....	5.0	*	0.7	0.9	0.8	0.6	0.6	1.0	1.3	2.1
2002.....	5.0	*	0.7	1.0	0.6	0.5	0.5	0.7	1.4	1.9
2001.....	5.0	*	0.7	0.9	0.6	0.6	0.6	0.8	1.5	2.1
2000.....	5.1	*	0.7	0.8	0.7	0.5	0.7	0.9	1.3	2.2
1999.....	5.2	*	0.8	0.9	0.6	0.5	0.6	1.1	1.4	2.2

White, male

2005 4/.....	5.6	*	0.4	0.9	0.8	0.8	0.7	1.2	1.5	2.2
2004 4/.....	5.5	*	0.7	1.0	0.7	0.6	0.7	0.8	1.5	2.4
2003 4/.....	5.6	*	0.8	0.9	0.8	0.7	0.7	1.0	1.6	2.5
2002.....	5.7	*	0.7	1.1	0.7	0.7	0.8	0.7	1.6	2.3
2001.....	5.7	*	0.8	0.8	0.6	0.7	0.7	1.1	1.9	2.6
2000.....	5.7	*	0.7	0.9	0.8	0.6	0.7	1.0	1.7	2.8
1999.....	5.8	*	0.8	0.9	0.6	0.6	0.7	1.2	1.8	2.6

White, female

2005 4/.....	4.4	*	0.6	1.0	0.6	0.4	0.5	0.6	0.9	1.3
2004 4/.....	4.4	*	0.6	0.9	0.6	0.5	0.4	0.7	0.9	1.4
2003 4/.....	4.4	*	0.6	0.9	0.8	0.6	0.5	0.9	1.0	1.8
2002.....	4.4	*	0.6	1.0	0.6	0.4	0.3	0.7	1.2	1.6
2001.....	4.4	*	0.7	1.0	0.7	0.4	0.6	0.5	1.2	1.6
2000.....	4.4	*	0.6	0.8	0.7	0.4	0.6	0.8	0.9	1.5
1999.....	4.6	*	0.8	0.8	0.7	0.5	0.5	0.9	1.0	1.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)

All races 3/, both sexes

2005.....	18.6	0.6	0.8	0.8	0.9	1.3	1.8	1.8	2.2	2.7
2004.....	18.6	0.6	0.8	0.9	0.9	1.4	1.7	1.8	2.3	2.7
2003.....	19.1	0.8	0.9	0.9	1.0	1.2	1.7	1.9	2.4	3.1
2002.....	19.5	0.7	0.8	0.8	1.0	1.5	1.9	1.9	2.6	3.2
2001.....	19.8	0.7	1.0	0.8	1.0	1.5	2.0	2.3	2.7	3.5
2000.....	20.1	1.0	0.9	0.8	1.0	1.6	2.1	2.1	2.5	3.4
1999.....	20.2	0.8	0.9	0.7	1.0	1.5	2.2	2.2	2.6	3.7

All races 3/, male

2005.....	20.6	*	1.0	0.9	1.0	1.6	2.2	2.1	2.4	3.0
2004.....	20.4	*	0.9	1.1	1.0	1.6	2.1	2.1	2.7	3.2
2003.....	21.0	*	1.0	1.0	1.1	1.6	2.0	2.3	3.0	3.5
2002.....	21.4	*	0.8	0.9	1.2	1.8	2.2	2.3	3.0	3.7
2001.....	21.5	*	1.0	0.9	1.1	1.8	2.3	2.6	3.1	4.2
2000.....	21.6	1.1	1.0	0.9	1.2	1.9	2.6	2.3	3.0	3.9
1999.....	21.7	*	1.0	0.8	1.2	1.7	2.7	2.4	3.1	4.4

All races 3/, female

2005.....	16.6	*	0.6	0.6	0.7	1.0	1.3	1.5	2.0	2.3
2004.....	16.8	*	0.6	0.7	0.7	1.1	1.3	1.5	1.9	2.2
2003.....	17.3	1.1	0.8	0.7	0.9	0.9	1.4	1.6	1.7	2.6
2002.....	17.7	*	0.7	0.8	0.8	1.2	1.7	1.5	2.2	2.6
2001.....	18.1	*	1.0	0.7	0.8	1.2	1.7	2.0	2.2	2.7
2000.....	18.6	*	0.9	0.7	0.8	1.2	1.5	1.9	1.9	3.0
1999.....	18.7	*	0.8	0.6	0.9	1.3	1.8	1.9	2.2	3.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)										
White, both sexes										
2005 4/.....	20.3	0.6	0.7	0.8	0.9	1.4	1.8	1.6	2.1	2.7
2004 4/.....	20.2	0.6	0.8	1.0	0.9	1.4	1.7	1.7	2.2	2.6
2003 4/.....	20.9	0.9	0.9	0.9	1.0	1.3	1.7	1.9	2.3	3.0
2002.....	21.3	*	0.8	0.9	1.0	1.5	2.1	1.9	2.5	3.1
2001.....	21.5	0.7	1.1	0.8	1.0	1.6	2.0	2.1	2.6	3.4
2000.....	21.9	1.2	1.0	0.9	1.1	1.6	2.0	2.1	2.4	3.3
1999.....	21.9	0.8	1.0	0.7	1.0	1.5	2.3	2.1	2.5	3.6
White, male										
2005 4/.....	22.5	*	0.9	0.9	1.1	1.7	2.2	1.9	2.2	3.1
2004 4/.....	22.2	*	0.8	1.2	1.0	1.7	2.1	2.1	2.6	3.1
2003 4/.....	23.1	*	1.1	1.2	1.1	1.6	2.0	2.2	2.9	3.4
2002.....	23.4	*	0.9	0.9	1.2	1.9	2.4	2.3	2.9	3.5
2001.....	23.3	*	1.1	0.9	1.0	1.9	2.3	2.4	3.2	4.0
2000.....	23.6	*	1.0	0.9	1.2	1.9	2.5	2.3	2.9	3.8
1999.....	23.6	*	1.0	0.9	1.1	1.7	2.8	2.3	2.9	4.3
White, female										
2005 4/.....	18.1	*	0.5	0.7	0.7	1.0	1.4	1.3	1.9	2.3
2004 4/.....	18.3	*	0.7	0.7	0.8	1.1	1.3	1.3	1.8	2.1
2003 4/.....	18.8	1.4	0.8	0.7	0.9	0.8	1.5	1.6	1.7	2.5
2002.....	19.2	*	0.8	0.9	0.8	1.1	1.7	1.5	2.2	2.6
2001.....	19.7	*	1.1	0.7	0.9	1.2	1.6	1.9	2.0	2.8
2000.....	20.3	*	1.0	0.8	0.9	1.2	1.4	1.8	1.9	2.9
1999.....	20.4	*	0.9	0.6	0.9	1.3	1.7	1.9	2.1	2.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)										
Black, both sexes										
2005 4/.....	12.9	*	1.0	*	0.8	1.3	1.9	2.7	3.8	3.2
2004 4/.....	13.6	*	*	0.7	0.8	1.4	1.9	2.8	3.1	3.7
2003 4/.....	13.4	*	*	0.6	1.0	1.1	1.7	2.4	3.3	4.1
2002.....	13.6	*	*	*	1.0	1.4	1.5	2.3	3.4	3.8
2001.....	14.1	*	*	0.9	1.0	1.5	2.1	3.2	3.5	4.5
2000.....	13.9	*	0.8	0.6	0.7	1.9	2.6	2.8	3.1	4.4
1999.....	14.1	*	*	0.6	1.3	1.7	2.9	2.6	3.2	4.6
Black, male										
2005 4/.....	14.1	*	*	*	*	1.4	2.5	2.8	4.7	3.7
2004 4/.....	14.4	*	*	*	*	1.4	2.5	2.9	3.6	4.3
2003 4/.....	13.9	*	*	*	1.2	*	2.1	2.5	4.2	4.7
2002.....	14.6	*	*	*	1.1	1.6	1.3	2.8	4.4	5.1
2001.....	15.1	*	*	*	1.7	1.8	2.2	3.8	4.0	6.5
2000.....	14.7	*	*	*	*	2.3	3.2	2.6	4.4	4.8
1999.....	15.1	*	*	*	1.7	2.1	3.1	3.3	4.2	5.3
Black, female										
2005 4/.....	11.8	*	*	*	*	1.3	*	2.6	3.0	2.8
2004 4/.....	12.8	*	*	*	*	1.4	1.3	2.7	2.7	3.1
2003 4/.....	12.9	*	*	*	*	*	1.3	2.4	2.4	3.6
2002.....	12.8	*	*	*	*	1.3	1.6	2.0	2.5	2.7
2001.....	13.1	*	*	*	*	*	2.1	2.7	3.1	2.8
2000.....	13.1	*	*	*	*	1.4	2.1	3.0	1.9	4.1
1999.....	13.2	*	*	*	*	1.4	2.8	2.0	2.3	3.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Hodgkin's disease (C81)										
All races 3/, both sexes										
2005.....	0.4	*	*	*	*	*	0.2	0.3	0.3	0.3
2004.....	0.4	*	*	*	*	*	0.2	0.3	0.3	0.3
2003.....	0.5	*	*	*	*	*	0.2	0.4	0.5	0.4
2002.....	0.5	*	*	*	*	0.1	0.3	0.3	0.4	0.4
2001.....	0.5	*	*	*	*	0.1	0.3	0.4	0.4	0.4
2000.....	0.5	*	*	*	*	0.1	0.3	0.3	0.4	0.3
1999.....	0.5	*	*	*	*	0.2	0.3	0.4	0.3	0.4
All races 3/, male										
2005.....	0.5	*	*	*	*	*	0.2	0.3	0.3	0.3
2004.....	0.5	*	*	*	*	*	0.3	0.3	0.4	0.3
2003.....	0.5	*	*	*	*	*	0.2	0.4	0.5	0.4
2002.....	0.5	*	*	*	*	*	0.3	0.4	0.4	0.4
2001.....	0.5	*	*	*	*	*	0.3	0.4	0.5	0.5
2000.....	0.5	*	*	*	*	*	0.3	0.3	0.5	0.4
1999.....	0.6	*	*	*	*	0.2	0.4	0.4	0.4	0.5
All races 3/, female										
2005.....	0.4	*	*	*	*	*	0.3	0.2	0.3	0.2
2004.....	0.4	*	*	*	*	*	*	0.3	0.3	0.2
2003.....	0.4	*	*	*	*	*	0.2	0.4	0.4	0.3
2002.....	0.4	*	*	*	*	*	0.3	0.3	0.4	0.3
2001.....	0.4	*	*	*	*	*	0.3	0.4	0.4	0.3
2000.....	0.4	*	*	*	*	*	0.2	0.3	0.4	0.3
1999.....	0.4	*	*	*	*	*	0.3	0.4	0.3	0.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Hodgkin's disease (C81)										
White, both sexes										
2005 4/.....	0.5	*	*	*	*	*	0.2	0.2	0.2	0.3
2004 4/.....	0.5	*	*	*	*	*	0.2	0.3	0.3	0.3
2003 4/.....	0.5	*	*	*	*	*	0.2	0.4	0.5	0.4
2002.....	0.5	*	*	*	*	0.1	0.3	0.3	0.4	0.4
2001.....	0.5	*	*	*	*	0.1	0.3	0.4	0.4	0.4
2000.....	0.5	*	*	*	*	*	0.3	0.4	0.5	0.4
1999.....	0.6	*	*	*	*	0.2	0.3	0.4	0.4	0.4
White, male										
2005 4/.....	0.5	*	*	*	*	*	*	0.3	*	0.3
2004 4/.....	0.5	*	*	*	*	*	0.3	0.3	0.4	0.3
2003 4/.....	0.5	*	*	*	*	*	*	0.4	0.5	0.4
2002.....	0.6	*	*	*	*	*	0.3	0.4	0.4	0.4
2001.....	0.6	*	*	*	*	*	0.3	0.4	0.5	0.5
2000.....	0.5	*	*	*	*	*	0.3	0.4	0.5	0.4
1999.....	0.6	*	*	*	*	*	0.4	0.4	0.5	0.5
White, female										
2005 4/.....	0.4	*	*	*	*	*	0.3	*	0.3	*
2004 4/.....	0.4	*	*	*	*	*	*	0.3	0.3	*
2003 4/.....	0.5	*	*	*	*	*	*	0.4	0.4	0.3
2002.....	0.5	*	*	*	*	*	0.4	0.3	0.5	0.3
2001.....	0.4	*	*	*	*	*	*	0.4	0.3	0.3
2000.....	0.5	*	*	*	*	*	*	0.3	0.4	0.3
1999.....	0.5	*	*	*	*	*	0.3	0.4	0.3	0.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Non-Hodgkin's lymphoma (C82-C85)										
All races 3/, both sexes										
2005.....	7.0	*	*	*	0.1	0.2	0.4	0.4	0.7	0.9
2004.....	7.1	*	*	0.1	*	0.3	0.4	0.6	0.7	0.9
2003.....	7.4	*	*	*	0.1	0.3	0.4	0.5	0.8	1.1
2002.....	7.6	*	*	*	0.2	0.2	0.5	0.6	0.8	1.2
2001.....	7.8	*	*	*	0.1	0.3	0.4	0.7	0.8	1.4
2000.....	8.1	*	*	0.1	0.1	0.3	0.6	0.6	0.8	1.3
1999.....	8.2	*	*	*	0.1	0.3	0.5	0.6	0.9	1.4
All races 3/, male										
2005.....	7.6	*	*	*	0.2	0.3	0.5	0.5	0.8	1.1
2004.....	7.6	*	*	*	*	0.3	0.5	0.8	0.8	1.2
2003.....	8.0	*	*	*	*	0.3	0.5	0.6	1.0	1.4
2002.....	8.1	*	*	*	0.3	0.3	0.6	0.7	1.0	1.5
2001.....	8.4	*	*	*	0.2	0.4	0.5	0.8	1.0	1.8
2000.....	8.6	*	*	*	*	0.4	0.8	0.7	1.0	1.7
1999.....	8.6	*	*	*	0.2	0.3	0.6	0.7	1.1	1.8
All races 3/, female										
2005.....	6.5	*	*	*	*	*	0.3	0.4	0.5	0.7
2004.....	6.6	*	*	*	*	*	0.3	0.4	0.6	0.6
2003.....	6.8	*	*	*	*	*	0.3	0.4	0.5	0.7
2002.....	7.1	*	*	*	*	*	0.4	0.4	0.6	0.8
2001.....	7.3	*	*	*	*	*	0.4	0.6	0.6	1.0
2000.....	7.6	*	*	*	*	*	0.4	0.6	0.6	0.9
1999.....	7.7	*	*	*	*	0.2	0.4	0.4	0.7	1.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Non-Hodgkin's lymphoma (C82-C85)

White, both sexes

2005 4/.....	7.9	*	*	*	0.2	0.2	0.3	0.4	0.6	0.8
2004 4/.....	8.0	*	*	*	*	0.3	0.4	0.5	0.6	0.9
2003 4/.....	8.3	*	*	*	*	0.2	0.4	0.4	0.7	1.1
2002.....	8.5	*	*	*	0.2	0.2	0.5	0.6	0.7	1.1
2001.....	8.8	*	*	*	0.1	0.3	0.4	0.6	0.8	1.3
2000.....	9.1	*	*	*	0.1	0.3	0.6	0.6	0.8	1.2
1999.....	9.2	*	*	*	*	0.2	0.5	0.5	0.8	1.4

White, male

2005 4/.....	8.5	*	*	*	0.3	0.4	0.5	0.4	0.7	1.1
2004 4/.....	8.5	*	*	*	*	0.4	0.5	0.8	0.7	1.2
2003 4/.....	8.9	*	*	*	*	0.3	0.5	0.6	0.9	1.4
2002.....	9.1	*	*	*	0.3	0.3	0.6	0.8	0.9	1.4
2001.....	9.3	*	*	*	*	0.4	0.5	0.7	1.0	1.6
2000.....	9.6	*	*	*	*	0.5	0.8	0.6	1.0	1.6
1999.....	9.6	*	*	*	*	0.3	0.6	0.6	1.0	1.7

White, female

2005 4/.....	7.3	*	*	*	*	*	*	0.3	0.5	0.6
2004 4/.....	7.5	*	*	*	*	*	0.3	0.3	0.5	0.6
2003 4/.....	7.7	*	*	*	*	*	0.4	0.3	0.4	0.7
2002.....	8.0	*	*	*	*	*	0.3	0.4	0.6	0.8
2001.....	8.2	*	*	*	*	*	0.4	0.5	0.5	1.0
2000.....	8.6	*	*	*	*	*	0.4	0.5	0.6	0.8
1999.....	8.8	*	*	*	*	*	0.4	0.4	0.6	1.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Leukemia (C91-C95)										
All races 3/, both sexes										
2005.....	7.3	0.5	0.7	0.6	0.7	1.1	1.2	1.1	1.2	1.4
2004.....	7.3	0.6	0.8	0.8	0.7	1.0	1.1	0.9	1.2	1.4
2003.....	7.4	0.7	0.8	0.8	0.9	0.9	1.1	1.1	1.1	1.5
2002.....	7.5	0.5	0.7	0.7	0.8	1.2	1.2	1.0	1.3	1.5
2001.....	7.5	0.6	1.0	0.8	0.8	1.1	1.3	1.2	1.4	1.6
2000.....	7.6	1.0	0.9	0.7	0.8	1.2	1.2	1.2	1.2	1.6
1999.....	7.5	0.7	0.8	0.6	0.9	1.1	1.4	1.2	1.3	1.6
All races 3/, male										
2005.....	8.4	*	1.0	0.8	0.8	1.2	1.5	1.3	1.2	1.4
2004.....	8.3	*	0.9	1.0	0.8	1.2	1.3	1.0	1.4	1.5
2003.....	8.4	*	0.9	0.9	1.0	1.2	1.2	1.3	1.4	1.6
2002.....	8.5	*	0.7	0.7	0.9	1.4	1.4	1.2	1.5	1.6
2001.....	8.5	*	1.0	0.8	0.9	1.3	1.5	1.4	1.6	1.8
2000.....	8.5	1.1	0.9	0.8	1.0	1.4	1.5	1.3	1.5	1.6
1999.....	8.4	*	0.8	0.7	0.9	1.3	1.7	1.3	1.4	1.8
All races 3/, female										
2005.....	6.2	*	0.5	0.5	0.7	0.9	0.8	0.9	1.2	1.3
2004.....	6.3	*	0.6	0.6	0.7	0.8	0.8	0.8	0.9	1.2
2003.....	6.4	1.1	0.8	0.7	0.8	0.6	0.9	0.9	0.8	1.4
2002.....	6.5	*	0.6	0.8	0.7	1.0	1.0	0.8	1.1	1.3
2001.....	6.6	*	1.0	0.7	0.7	0.9	1.0	1.0	1.2	1.3
2000.....	6.7	*	0.9	0.6	0.6	0.9	0.9	1.0	0.9	1.6
1999.....	6.7	*	0.7	0.5	0.8	0.9	1.1	1.1	1.2	1.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Leukemia (C91-C95)										
White, both sexes										
2005 4/.....	8.1	*	0.7	0.7	0.7	1.1	1.2	1.0	1.2	1.5
2004 4/.....	8.0	*	0.7	0.8	0.8	1.0	1.1	0.9	1.2	1.3
2003 4/.....	8.2	0.9	0.9	0.9	0.9	0.9	1.1	1.1	1.1	1.4
2002.....	8.2	*	0.7	0.8	0.8	1.2	1.3	1.0	1.3	1.5
2001.....	8.3	0.6	1.1	0.7	0.8	1.2	1.3	1.1	1.4	1.5
2000.....	8.3	1.2	1.0	0.7	0.9	1.2	1.2	1.1	1.1	1.6
1999.....	8.3	0.8	0.9	0.7	0.8	1.1	1.4	1.2	1.3	1.6
White, male										
2005 4/.....	9.3	*	0.9	0.8	0.8	1.3	1.5	1.2	1.3	1.5
2004 4/.....	9.1	*	0.8	1.0	0.8	1.2	1.4	1.0	1.4	1.4
2003 4/.....	9.4	*	1.0	1.0	1.0	1.2	1.2	1.3	1.4	1.4
2002.....	9.3	*	0.8	0.7	0.9	1.4	1.5	1.2	1.5	1.6
2001.....	9.3	*	1.1	0.8	0.8	1.4	1.6	1.3	1.6	1.7
2000.....	9.4	*	0.9	0.8	1.1	1.4	1.5	1.3	1.4	1.7
1999.....	9.2	*	0.9	0.8	0.9	1.2	1.8	1.3	1.4	1.8
White, female										
2005 4/.....	6.9	*	0.5	0.6	0.7	0.9	0.9	0.8	1.2	1.4
2004 4/.....	6.9	*	0.7	0.7	0.7	0.8	0.9	0.8	1.0	1.2
2003 4/.....	7.0	1.4	0.8	0.7	0.8	0.6	0.9	0.9	0.8	1.4
2002.....	7.1	*	0.6	0.8	0.7	1.0	1.1	0.8	1.1	1.4
2001.....	7.3	*	1.1	0.7	0.8	0.9	1.0	0.9	1.2	1.4
2000.....	7.4	*	1.0	0.7	0.7	0.9	0.9	0.9	0.9	1.6
1999.....	7.3	*	0.8	0.6	0.8	1.0	1.0	1.1	1.2	1.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Leukemia (C91-C95)										
Black, both sexes										
2005 4/.....	4.4	*	1.0	*	0.7	1.0	0.9	1.5	1.5	1.1
2004 4/.....	4.8	*	*	*	0.6	1.1	0.9	1.1	1.5	1.8
2003 4/.....	4.6	*	*	*	0.8	0.7	1.0	1.0	1.4	2.0
2002.....	4.7	*	*	*	0.7	0.9	0.8	1.1	1.4	1.5
2001.....	4.7	*	*	0.8	0.9	0.9	1.2	1.4	1.4	1.8
2000.....	4.8	*	*	*	0.6	1.5	1.3	1.4	1.8	1.7
1999.....	4.7	*	*	*	1.1	1.2	1.7	1.1	1.4	1.8
Black, male										
2005 4/.....	4.9	*	*	*	*	*	1.6	1.5	*	*
2004 4/.....	5.3	*	*	*	*	*	*	*	1.8	2.1
2003 4/.....	5.0	*	*	*	*	*	1.3	*	1.8	2.5
2002.....	5.1	*	*	*	*	*	*	*	1.6	1.9
2001.....	5.2	*	*	*	1.5	*	*	1.7	1.8	2.3
2000.....	5.3	*	*	*	*	2.0	1.8	*	2.5	1.4
1999.....	5.2	*	*	*	1.5	1.5	1.7	*	1.7	1.8
Black, female										
2005 4/.....	4.0	*	*	*	*	*	*	1.4	1.6	*
2004 4/.....	4.4	*	*	*	*	*	*	*	*	1.5
2003 4/.....	4.2	*	*	*	*	*	*	*	*	1.7
2002.....	4.3	*	*	*	*	*	*	*	*	*
2001.....	4.3	*	*	*	*	*	*	*	*	1.4
2000.....	4.2	*	*	*	*	*	*	1.5	*	1.9
1999.....	4.3	*	*	*	*	*	1.8	*	*	1.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)

All races 3/, both sexes

2005.....	21.2	*	0.8	0.6	0.8	1.3	1.5	1.5	2.0	2.9
2004.....	21.1	0.6	0.8	0.6	0.7	1.2	1.4	1.5	2.0	3.1
2003.....	21.5	*	0.7	0.6	0.7	1.2	1.3	1.6	1.9	3.1
2002.....	21.8	0.6	0.8	0.7	0.7	1.1	1.5	1.5	2.1	3.3
2001.....	22.0	*	0.8	0.6	0.8	1.2	1.4	1.7	2.0	3.2
2000.....	22.4	0.6	0.8	0.6	0.7	1.2	1.4	1.5	2.0	3.4
1999.....	23.7	*	0.8	0.7	0.7	1.2	1.5	1.6	2.2	3.6

All races 3/, male

2005.....	22.1	*	0.8	0.6	0.8	1.4	1.9	1.9	2.4	3.1
2004.....	21.9	*	0.7	0.8	0.8	1.5	1.8	1.9	2.3	3.6
2003.....	22.3	*	0.8	0.8	0.7	1.5	1.6	1.9	2.4	3.5
2002.....	22.7	*	0.8	0.8	0.7	1.3	1.9	2.0	2.5	3.7
2001.....	22.6	*	0.9	0.6	0.8	1.5	1.9	2.0	2.4	3.5
2000.....	23.0	*	1.0	0.6	0.7	1.4	1.8	1.9	2.4	3.7
1999.....	24.5	*	0.8	0.8	0.8	1.5	1.8	2.1	2.8	4.1

All races 3/, female

2005.....	20.4	*	0.7	0.6	0.7	1.2	1.0	1.0	1.6	2.7
2004.....	20.3	*	0.9	0.5	0.6	1.0	0.9	1.1	1.7	2.5
2003.....	20.8	*	0.5	0.5	0.7	0.8	1.1	1.2	1.4	2.7
2002.....	21.0	*	0.8	0.6	0.8	0.9	1.0	1.0	1.7	2.8
2001.....	21.4	*	0.6	0.5	0.7	0.9	0.8	1.4	1.6	3.0
2000.....	21.9	*	0.7	0.6	0.6	1.0	1.0	1.1	1.6	3.0
1999.....	23.1	*	0.7	0.7	0.6	0.9	1.1	1.1	1.6	3.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)										
White, both sexes										
2005 4/.....	22.7	*	0.8	0.6	0.7	1.4	1.5	1.4	1.9	2.9
2004 4/.....	22.6	*	0.8	0.7	0.7	1.2	1.3	1.6	2.0	3.0
2003 4/.....	23.1	*	0.6	0.7	0.7	1.3	1.4	1.6	1.8	3.1
2002.....	23.3	0.7	0.7	0.7	0.8	1.2	1.5	1.5	2.1	3.2
2001.....	23.5	*	0.8	0.6	0.8	1.2	1.5	1.7	1.9	3.1
2000.....	23.9	*	0.8	0.7	0.7	1.2	1.5	1.6	2.0	3.2
1999.....	25.3	*	0.8	0.7	0.7	1.2	1.4	1.6	2.2	3.5
White, male										
2005 4/.....	23.6	*	0.9	0.6	0.7	1.5	2.0	1.9	2.3	3.2
2004 4/.....	23.4	*	0.7	0.8	0.8	1.5	1.8	2.0	2.4	3.5
2003 4/.....	23.8	*	0.7	0.9	0.6	1.6	1.6	2.0	2.3	3.6
2002.....	24.2	*	0.8	0.7	0.7	1.4	1.9	2.0	2.5	3.6
2001.....	24.0	*	1.0	0.6	0.8	1.5	2.0	2.0	2.3	3.4
2000.....	24.4	*	1.0	0.7	0.7	1.3	1.9	2.1	2.5	3.6
1999.....	25.9	*	0.9	0.8	0.8	1.5	1.7	2.2	2.7	4.0
White, female										
2005 4/.....	21.9	*	0.6	0.5	0.8	1.2	1.0	0.9	1.5	2.6
2004 4/.....	21.8	*	0.8	0.5	0.6	1.0	0.9	1.1	1.5	2.4
2003 4/.....	22.5	*	0.5	0.5	0.7	0.9	1.1	1.1	1.3	2.6
2002.....	22.5	*	0.7	0.6	0.8	1.0	1.0	1.0	1.7	2.8
2001.....	22.9	*	0.7	0.5	0.7	1.0	0.8	1.4	1.5	2.8
2000.....	23.5	*	0.6	0.6	0.6	1.0	1.1	1.1	1.5	2.8
1999.....	24.6	*	0.7	0.7	0.6	1.0	1.1	1.0	1.7	3.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)										
Black, both sexes										
2005 4/.....	17.1	*	0.8	0.7	1.1	0.9	1.5	2.1	2.6	3.6
2004 4/.....	17.2	*	0.8	0.8	1.0	1.4	1.8	1.5	2.1	4.3
2003 4/.....	17.3	*	0.9	0.6	0.9	0.9	1.3	1.9	2.7	3.7
2002.....	18.1	*	1.1	0.8	0.7	1.1	1.6	1.9	2.6	4.5
2001.....	18.3	*	*	*	0.8	1.2	1.3	2.0	2.6	4.3
2000.....	18.3	*	1.0	*	0.7	1.5	1.3	1.3	2.0	5.2
1999.....	19.5	*	*	0.6	0.9	1.3	1.6	1.5	2.4	4.9
Black, male										
2005 4/.....	18.0	*	*	*	1.4	*	1.9	2.4	3.2	3.7
2004 4/.....	18.4	*	*	*	*	1.7	2.1	1.7	2.1	4.6
2003 4/.....	18.3	*	*	*	*	1.2	1.7	1.7	3.0	3.6
2002.....	18.8	*	*	*	*	1.4	2.2	2.5	2.7	5.1
2001.....	19.4	*	*	*	*	1.5	1.9	2.5	3.0	4.4
2000.....	19.4	*	*	*	*	1.7	2.1	1.6	2.0	5.4
1999.....	20.6	*	*	*	*	1.9	2.0	1.5	3.1	5.1
Black, female										
2005 4/.....	16.3	*	*	*	*	*	*	1.7	2.0	3.6
2004 4/.....	16.2	*	*	*	1.1	*	1.5	*	2.2	4.0
2003 4/.....	16.5	*	*	*	*	*	*	2.0	2.3	3.9
2002.....	17.4	*	*	*	*	*	*	1.4	2.6	4.0
2001.....	17.3	*	*	*	*	*	*	1.5	2.3	4.2
2000.....	17.3	*	*	*	*	1.4	*	*	1.9	5.1
1999.....	18.5	*	*	*	*	*	*	1.5	1.8	4.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)										
All races 3/, both sexes										
2005.....	4.6	1.4	0.3	0.2	0.2	0.2	0.2	0.3	0.5	0.7
2004.....	4.6	1.7	0.3	0.2	0.2	0.2	0.3	0.3	0.5	0.6
2003.....	4.7	1.5	0.3	0.2	0.2	0.2	0.3	0.4	0.5	0.7
2002.....	4.6	1.7	0.4	0.2	0.2	0.2	0.2	0.3	0.5	0.6
2001.....	4.8	1.3	0.4	0.3	0.3	0.2	0.3	0.3	0.4	0.7
2000.....	4.8	1.6	0.3	0.3	0.2	0.2	0.3	0.4	0.5	0.7
1999.....	4.7	1.5	0.4	0.3	0.2	0.3	0.2	0.3	0.6	0.6
All races 3/, male										
2005.....	4.7	1.0	0.4	*	*	0.3	0.2	0.4	0.5	0.7
2004.....	4.7	1.7	0.3	0.2	0.2	0.2	0.4	0.4	0.6	0.6
2003.....	4.7	1.6	0.4	0.2	0.2	0.2	0.3	0.4	0.6	0.7
2002.....	4.6	1.5	0.4	0.2	0.2	0.2	0.3	0.3	0.5	0.7
2001.....	4.8	1.3	0.4	0.3	0.2	0.2	0.3	0.3	0.4	0.7
2000.....	4.8	1.6	0.4	0.3	*	0.3	0.4	0.3	0.5	0.8
1999.....	4.8	1.5	0.4	0.3	*	0.3	0.3	0.3	0.6	0.7
All races 3/, female										
2005.....	4.6	1.8	0.3	0.2	*	0.2	*	0.3	0.4	0.7
2004.....	4.6	1.8	0.3	0.2	*	0.2	0.2	0.3	0.5	0.6
2003.....	4.7	1.5	0.3	*	*	*	0.2	0.3	0.5	0.6
2002.....	4.6	2.0	0.4	0.2	*	0.2	*	0.3	0.5	0.5
2001.....	4.8	1.3	0.4	0.3	0.3	0.3	0.2	0.3	0.5	0.7
2000.....	4.9	1.5	0.3	0.3	0.2	*	0.2	0.4	0.4	0.6
1999.....	4.7	1.5	0.4	0.3	*	0.3	*	0.2	0.5	0.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)

White, both sexes

2005 4/.....	5.1	1.3	0.3	0.2	0.2	0.2	0.2	0.3	0.5	0.6
2004 4/.....	5.1	1.9	0.3	0.2	0.2	0.2	0.3	0.3	0.5	0.6
2003 4/.....	5.2	1.5	0.3	0.2	0.2	0.2	0.3	0.4	0.5	0.6
2002.....	5.1	1.9	0.4	0.2	0.2	0.2	0.2	0.3	0.5	0.6
2001.....	5.3	1.2	0.4	0.2	0.2	0.2	0.3	0.4	0.4	0.7
2000.....	5.3	1.6	0.4	0.3	0.2	0.2	0.3	0.3	0.5	0.7
1999.....	5.2	1.4	0.4	0.3	0.2	0.2	0.2	0.3	0.5	0.6

White, male

2005 4/.....	5.2	*	0.4	*	*	0.3	0.3	0.4	0.5	0.7
2004 4/.....	5.1	1.8	0.3	*	*	0.2	0.4	0.4	0.6	0.6
2003 4/.....	5.2	1.6	0.3	*	*	*	0.3	0.4	0.5	0.7
2002.....	5.1	1.7	0.3	*	0.2	*	0.2	0.3	0.5	0.7
2001.....	5.3	1.2	0.4	*	*	*	0.3	0.4	0.4	0.7
2000.....	5.2	1.6	0.4	0.3	*	*	0.4	0.3	0.5	0.7
1999.....	5.2	1.4	0.4	0.3	*	*	0.3	0.3	0.5	0.7

White, female

2005 4/.....	5.0	1.7	*	*	*	*	*	0.3	0.5	0.6
2004 4/.....	5.0	1.9	0.3	*	*	*	*	0.3	0.4	0.6
2003 4/.....	5.1	1.5	*	*	*	*	0.3	0.3	0.5	0.5
2002.....	5.1	2.1	0.4	0.3	*	*	*	*	0.4	0.5
2001.....	5.2	*	0.4	*	0.3	0.3	*	0.3	0.5	0.7
2000.....	5.4	1.6	*	0.3	*	*	*	0.3	0.4	0.6
1999.....	5.1	1.4	0.4	0.4	*	0.3	*	*	0.5	0.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Anemias (D50-D64)										
All races 3/, both sexes										
2005.....	1.6	*	0.1	*	*	0.2	0.3	0.4	0.4	0.4
2004.....	1.6	*	0.2	0.1	0.1	0.2	0.4	0.4	0.4	0.4
2003.....	1.6	0.5	0.2	*	0.1	0.2	0.3	0.4	0.4	0.4
2002.....	1.6	*	0.1	0.1	0.1	0.2	0.3	0.4	0.5	0.5
2001.....	1.6	*	0.1	0.1	0.1	0.2	0.3	0.3	0.5	0.4
2000.....	1.6	*	0.2	0.1	0.1	0.1	0.3	0.3	0.4	0.4
1999.....	1.6	*	0.2	0.1	0.1	0.2	0.4	0.3	0.5	0.4
All races 3/, male										
2005.....	1.2	*	*	*	*	0.2	0.3	0.4	0.4	0.4
2004.....	1.3	*	*	*	*	0.3	0.4	0.4	0.5	0.5
2003.....	1.3	*	*	*	*	*	0.3	0.4	0.4	0.3
2002.....	1.3	*	*	*	*	0.2	0.4	0.5	0.5	0.5
2001.....	1.3	*	*	*	*	0.2	0.3	0.4	0.5	0.4
2000.....	1.2	*	*	*	*	*	0.3	0.3	0.4	0.3
1999.....	1.3	*	*	*	*	0.2	0.5	0.3	0.5	0.4
All races 3/, female										
2005.....	1.9	*	*	*	*	*	0.3	0.4	0.3	0.4
2004.....	1.8	*	*	*	*	*	0.4	0.5	0.3	0.3
2003.....	1.9	*	*	*	*	*	0.3	0.3	0.4	0.4
2002.....	1.9	*	*	*	*	*	0.3	0.3	0.5	0.4
2001.....	2.0	*	*	*	*	0.2	0.3	0.3	0.4	0.4
2000.....	2.0	*	*	*	*	*	0.3	0.3	0.5	0.4
1999.....	1.9	*	*	*	*	*	0.4	0.3	0.5	0.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Anemias (D50-D64)

Black, both sexes

2005 4/.....	2.6	*	*	*	*	0.9	1.5	2.3	2.4	2.2
2004 4/.....	2.7	*	*	*	*	0.9	2.0	2.2	2.4	2.4
2003 4/.....	2.7	*	*	*	*	*	1.9	1.9	2.2	2.2
2002.....	2.8	*	*	*	*	0.9	1.7	2.5	2.3	2.6
2001.....	2.7	*	*	*	*	0.9	1.5	1.9	2.5	2.1
2000.....	2.6	*	*	*	*	*	1.6	1.7	2.8	2.2
1999.....	2.8	*	0.8	*	*	0.7	2.2	2.1	3.0	2.2

Black, male

2005 4/.....	2.4	*	*	*	*	*	1.7	2.4	2.7	2.7
2004 4/.....	2.6	*	*	*	*	1.2	1.9	2.3	3.0	3.0
2003 4/.....	2.7	*	*	*	*	*	1.8	2.2	2.3	2.2
2002.....	2.7	*	*	*	*	*	2.3	3.2	2.5	2.7
2001.....	2.5	*	*	*	*	*	1.6	2.5	3.3	2.2
2000.....	2.3	*	*	*	*	*	1.7	1.6	2.7	1.9
1999.....	2.7	*	*	*	*	*	2.8	2.3	3.3	2.5

Black, female

2005 4/.....	2.8	*	*	*	*	*	1.3	2.1	2.2	1.8
2004 4/.....	2.8	*	*	*	*	*	2.1	2.2	1.9	1.9
2003 4/.....	2.8	*	*	*	*	*	2.1	1.7	2.2	2.2
2002.....	3.0	*	*	*	*	*	*	1.8	2.2	2.4
2001.....	2.8	*	*	*	*	*	1.5	*	1.8	2.1
2000.....	3.0	*	*	*	*	*	1.6	1.9	2.9	2.5
1999.....	3.0	*	*	*	*	*	1.7	1.8	2.8	2.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Diabetes mellitus (E10-E14)										
All races 3/, both sexes										
2005.....	25.3	*	*	*	0.2	0.3	0.6	1.0	2.0	3.6
2004.....	24.9	*	*	*	0.1	0.2	0.5	0.9	2.0	3.4
2003.....	25.5	*	*	*	0.1	0.2	0.5	1.2	2.1	3.3
2002.....	25.4	*	*	*	0.1	0.3	0.6	1.2	2.0	3.5
2001.....	25.1	*	*	*	0.2	0.2	0.5	1.1	1.8	3.3
2000.....	24.6	*	*	*	*	0.2	0.6	1.1	2.0	3.0
1999.....	24.5	*	*	*	0.1	0.3	0.6	1.0	1.9	3.2
All races 3/, male										
2005.....	25.0	*	*	*	*	0.4	0.7	1.1	2.5	4.3
2004.....	24.4	*	*	*	*	0.3	0.5	1.0	2.3	4.2
2003.....	24.8	*	*	*	*	0.3	0.6	1.2	2.5	3.9
2002.....	24.2	*	*	*	*	0.3	0.7	1.3	2.4	4.1
2001.....	23.5	*	*	*	0.2	0.3	0.6	1.2	2.2	3.9
2000.....	22.9	*	*	*	*	0.3	0.7	1.2	2.5	3.5
1999.....	22.8	*	*	*	*	0.2	0.6	1.0	2.2	3.8
All races 3/, female										
2005.....	25.7	*	*	*	0.2	0.3	0.6	0.9	1.5	2.9
2004.....	25.4	*	*	*	*	*	0.6	0.8	1.8	2.5
2003.....	26.2	*	*	*	*	*	0.5	1.1	1.7	2.8
2002.....	26.5	*	*	*	*	0.3	0.5	1.1	1.5	3.0
2001.....	26.6	*	*	*	*	0.2	0.5	1.1	1.5	2.7
2000.....	26.3	*	*	*	*	*	0.5	0.9	1.6	2.5
1999.....	26.2	*	*	*	*	0.3	0.5	1.0	1.6	2.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Diabetes mellitus (E10-E14)										
White, both sexes										
2005 4/.....	24.9	*	*	*	0.1	0.2	0.5	0.8	1.6	3.0
2004 4/.....	24.4	*	*	*	*	0.2	0.4	0.8	1.7	2.9
2003 4/.....	25.0	*	*	*	*	0.2	0.5	0.9	1.8	2.8
2002.....	24.9	*	*	*	*	0.2	0.5	1.0	1.7	3.1
2001.....	24.6	*	*	*	*	0.2	0.4	0.8	1.6	2.8
2000.....	24.1	*	*	*	*	0.2	0.5	0.8	1.8	2.7
1999.....	23.9	*	*	*	*	0.2	0.5	0.9	1.6	2.8
White, male										
2005 4/.....	24.9	*	*	*	*	0.3	0.6	0.9	1.9	3.5
2004 4/.....	24.3	*	*	*	*	*	0.4	0.8	2.0	3.6
2003 4/.....	24.8	*	*	*	*	*	0.6	0.8	2.2	3.4
2002.....	24.2	*	*	*	*	*	0.5	1.1	2.1	3.7
2001.....	23.5	*	*	*	*	*	0.4	0.8	1.8	3.3
2000.....	22.9	*	*	*	*	0.3	0.5	1.0	2.3	3.1
1999.....	22.7	*	*	*	*	0.3	0.5	0.9	1.8	3.4
White, female										
2005 4/.....	24.9	*	*	*	*	*	0.4	0.7	1.2	2.5
2004 4/.....	24.5	*	*	*	*	*	0.4	0.7	1.5	2.1
2003 4/.....	25.2	*	*	*	*	*	0.4	1.0	1.4	2.3
2002.....	25.6	*	*	*	*	*	0.4	0.9	1.3	2.5
2001.....	25.7	*	*	*	*	*	0.3	0.8	1.4	2.3
2000.....	25.3	*	*	*	*	*	0.4	0.7	1.2	2.3
1999.....	25.0	*	*	*	*	*	0.4	0.9	1.4	2.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Diabetes mellitus (E10-E14)										
Black, both sexes										
2005 4/.....	33.2	*	*	*	*	0.7	1.5	2.8	5.1	8.1
2004 4/.....	33.2	*	*	*	*	0.6	1.3	2.0	4.6	7.5
2003 4/.....	33.8	*	*	*	*	*	1.0	3.2	4.4	7.3
2002.....	33.6	*	*	*	*	*	1.4	2.8	4.3	7.3
2001.....	33.1	*	*	*	*	*	1.4	3.2	4.2	6.9
2000.....	32.8	*	*	*	*	*	1.5	2.8	4.1	6.0
1999.....	33.0	*	*	*	*	*	1.1	1.8	4.1	6.6
Black, male										
2005 4/.....	30.7	*	*	*	*	*	1.4	2.9	6.7	10.8
2004 4/.....	30.2	*	*	*	*	*	*	2.5	5.4	9.5
2003 4/.....	29.8	*	*	*	*	*	*	4.1	5.2	8.4
2002.....	29.0	*	*	*	*	*	1.7	3.0	5.7	8.5
2001.....	28.5	*	*	*	*	*	1.5	3.9	5.6	8.7
2000.....	27.4	*	*	*	*	*	1.5	3.1	4.5	7.1
1999.....	27.7	*	*	*	*	*	*	1.6	5.0	7.7
Black, female										
2005 4/.....	35.5	*	*	*	*	*	1.7	2.6	3.6	5.7
2004 4/.....	36.0	*	*	*	*	*	1.6	1.6	3.8	5.6
2003 4/.....	37.4	*	*	*	*	*	*	2.3	3.7	6.3
2002.....	37.8	*	*	*	*	*	*	2.6	3.1	6.3
2001.....	37.2	*	*	*	*	*	1.3	2.7	2.9	5.3
2000.....	37.8	*	*	*	*	*	1.5	2.5	3.8	5.0
1999.....	37.8	*	*	*	*	*	*	2.0	3.3	5.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Major cardiovascular diseases (I00-I78)										
All races 3/, both sexes										
2005.....	288.8	12.2	1.3	0.8	1.0	2.3	4.3	7.3	12.8	24.1
2004.....	293.3	13.8	1.5	0.6	1.0	2.2	4.2	7.3	12.6	24.3
2003.....	310.3	13.9	1.5	0.7	1.0	2.5	4.5	7.1	13.2	25.1
2002.....	318.6	15.9	1.4	0.6	1.1	2.4	3.9	7.1	12.6	25.3
2001.....	323.9	14.9	1.9	0.7	1.1	2.2	4.3	7.1	13.0	25.0
2000.....	332.9	16.7	1.5	0.7	1.1	2.4	4.3	6.7	12.2	24.2
1999.....	340.6	17.0	1.5	0.7	1.0	2.7	4.1	6.5	12.6	25.9
All races 3/, male										
2005.....	277.9	13.4	1.5	0.8	1.1	3.0	5.5	9.5	16.6	32.0
2004.....	281.2	14.7	1.5	0.5	1.1	2.8	5.3	9.2	16.3	32.1
2003.....	296.3	15.2	1.4	0.7	1.2	3.1	5.3	8.5	17.0	33.4
2002.....	303.3	16.6	1.5	0.6	1.3	3.1	5.0	9.0	16.1	32.9
2001.....	306.2	15.3	1.9	0.6	1.1	2.7	5.3	8.8	16.5	33.1
2000.....	315.8	17.5	1.6	0.7	1.2	2.8	5.0	8.2	15.3	31.3
1999.....	323.0	17.9	1.6	0.8	1.1	3.3	4.8	7.8	16.2	34.2
All races 3/, female										
2005.....	299.3	11.0	1.1	0.9	0.8	1.7	3.0	5.0	9.0	16.1
2004.....	304.9	12.9	1.5	0.7	1.0	1.6	3.0	5.3	8.9	16.4
2003.....	323.9	12.5	1.5	0.6	0.8	2.0	3.6	5.6	9.3	16.8
2002.....	333.3	15.1	1.3	0.7	0.9	1.7	2.8	5.1	9.0	17.7
2001.....	340.9	14.5	1.8	0.8	1.1	1.7	3.2	5.5	9.5	17.0
2000.....	349.5	15.9	1.4	0.6	1.0	2.0	3.5	5.2	9.0	17.1
1999.....	357.5	16.0	1.4	0.6	1.0	2.1	3.4	5.2	9.1	17.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Major cardiovascular diseases (I00-I78)										
White, both sexes										
2005 4/.....	307.0	9.9	1.1	0.7	0.8	2.0	3.6	5.9	10.5	20.3
2004 4/.....	311.7	11.9	1.2	0.5	0.9	1.8	3.6	5.8	10.4	21.1
2003 4/.....	330.0	11.0	1.2	0.6	0.9	2.2	3.7	5.7	11.0	21.4
2002.....	339.4	13.4	1.3	0.6	0.9	2.1	3.3	5.7	10.2	21.6
2001.....	344.8	12.2	1.5	0.6	1.0	1.9	3.4	5.9	10.9	21.4
2000.....	354.7	14.6	1.3	0.6	0.9	2.1	3.5	5.5	10.1	20.6
1999.....	362.7	14.6	1.3	0.7	0.9	2.4	3.4	5.4	10.6	22.0
White, male										
2005 4/.....	293.0	10.8	1.3	0.7	1.0	2.5	4.7	7.9	13.7	27.5
2004 4/.....	296.7	12.7	1.3	0.5	1.0	2.3	4.6	7.6	13.8	28.3
2003 4/.....	312.6	11.5	1.1	0.6	1.1	2.7	4.4	7.0	14.4	29.3
2002.....	320.9	14.3	1.5	0.5	1.1	2.7	4.3	7.5	13.3	28.5
2001.....	323.5	13.2	1.6	0.6	1.0	2.3	4.3	7.3	14.0	28.8
2000.....	334.2	15.3	1.6	0.7	1.1	2.4	4.2	6.9	13.2	27.6
1999.....	341.2	15.3	1.5	0.8	0.9	2.9	3.9	6.5	14.0	29.8
White, female										
2005 4/.....	320.8	9.0	0.9	0.7	0.7	1.5	2.5	3.7	7.1	13.0
2004 4/.....	326.5	11.0	1.1	0.6	0.8	1.3	2.5	3.9	7.0	13.7
2003 4/.....	347.0	10.4	1.4	0.5	0.7	1.7	2.8	4.3	7.4	13.2
2002.....	357.5	12.5	1.1	0.7	0.8	1.5	2.2	3.9	7.1	14.5
2001.....	365.5	11.1	1.5	0.7	0.9	1.5	2.5	4.3	7.6	13.9
2000.....	374.8	13.9	1.1	0.5	0.7	1.7	2.8	4.1	6.9	13.5
1999.....	383.6	13.8	1.2	0.6	0.9	1.9	2.9	4.2	7.2	14.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Major cardiovascular diseases (I00-I78)										
Black, both sexes										
2005 4/.....	256.2	23.1	2.3	1.6	1.5	4.2	8.3	16.6	29.5	52.6
2004 4/.....	260.3	21.9	2.9	1.1	1.8	4.2	7.7	16.8	28.5	49.8
2003 4/.....	273.1	26.2	2.8	1.2	1.6	4.4	9.6	16.1	29.3	52.7
2002.....	276.1	27.6	2.0	0.8	2.0	4.0	7.8	16.6	29.6	52.9
2001.....	280.4	28.6	3.5	1.0	1.6	4.0	9.4	16.4	28.5	51.5
2000.....	285.0	27.8	2.5	1.0	1.9	4.6	9.1	15.1	27.7	51.2
1999.....	289.4	29.2	2.4	1.0	1.8	4.8	9.0	14.4	26.8	55.2
Black, male										
2005 4/.....	255.0	26.3	2.2	1.4	1.7	5.6	10.6	20.6	38.0	69.2
2004 4/.....	257.2	23.7	2.8	*	1.8	5.6	9.8	20.3	35.2	65.4
2003 4/.....	268.6	31.2	3.3	1.4	1.7	5.6	11.6	19.5	36.3	65.7
2002.....	268.5	28.2	1.9	*	2.3	5.1	9.4	20.6	36.9	66.8
2001.....	272.4	28.1	3.6	*	1.5	5.1	11.3	20.3	35.2	65.6
2000.....	275.6	29.1	2.1	*	1.8	5.1	10.2	18.0	32.7	61.6
1999.....	282.5	30.6	2.5	*	2.1	6.3	10.9	17.7	33.0	69.6
Black, female										
2005 4/.....	257.2	19.7	2.3	1.7	1.4	2.6	6.1	12.9	22.0	37.8
2004 4/.....	263.2	20.1	3.0	1.3	1.8	2.7	5.5	13.5	22.5	35.7
2003 4/.....	277.2	21.0	2.4	*	1.5	3.1	7.6	13.0	23.0	41.0
2002.....	283.0	26.9	2.1	*	1.6	2.8	6.3	13.0	23.0	40.6
2001.....	287.8	29.0	3.4	1.3	1.6	2.8	7.5	12.9	22.4	38.9
2000.....	293.4	26.5	2.9	*	2.0	4.2	8.1	12.4	23.3	42.0
1999.....	295.7	27.9	2.4	*	1.5	3.3	7.1	11.5	21.3	42.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Diseases of heart (I00-I09,I11,I13,I20-I51)										
All races 3/, both sexes										
2005.....	220.0	8.7	0.9	0.5	0.7	1.8	3.5	5.9	10.3	19.4
2004.....	222.2	10.3	1.2	0.4	0.8	1.8	3.2	5.7	10.0	19.4
2003.....	235.6	11.0	1.2	0.5	0.8	1.9	3.6	5.6	10.5	20.1
2002.....	241.7	12.4	1.1	0.5	0.8	2.0	3.1	5.7	10.0	20.3
2001.....	245.8	11.9	1.5	0.5	0.8	1.7	3.3	5.6	10.2	19.9
2000.....	252.6	13.0	1.2	0.5	0.8	2.0	3.3	5.2	9.5	19.1
1999.....	259.9	13.8	1.2	0.6	0.8	2.3	3.3	5.1	10.1	20.6
All races 3/, male										
2005.....	221.1	9.4	1.0	0.5	0.8	2.5	4.6	7.9	13.7	26.8
2004.....	222.8	10.9	1.1	0.3	0.8	2.3	4.2	7.4	13.6	26.5
2003.....	235.0	12.1	1.1	0.6	0.9	2.4	4.4	6.9	13.9	27.8
2002.....	240.7	12.9	1.1	0.4	0.9	2.6	4.0	7.4	13.4	27.8
2001.....	242.5	11.8	1.5	0.4	0.9	2.1	4.2	7.0	13.4	27.4
2000.....	249.8	13.3	1.4	0.6	0.9	2.4	4.0	6.5	12.4	26.0
1999.....	257.0	13.9	1.3	0.6	0.9	2.9	3.9	6.3	13.2	28.5
All races 3/, female										
2005.....	218.9	8.0	0.9	0.6	0.6	1.2	2.2	3.7	6.8	12.0
2004.....	221.6	9.7	1.2	0.5	0.7	1.2	2.2	3.9	6.4	12.3
2003.....	236.2	9.8	1.3	0.5	0.6	1.4	2.7	4.2	7.1	12.5
2002.....	242.7	11.8	1.0	0.5	0.6	1.3	2.1	3.9	6.5	12.8
2001.....	249.0	12.0	1.4	0.6	0.8	1.3	2.4	4.1	6.9	12.4
2000.....	255.3	12.5	1.0	0.4	0.7	1.6	2.6	3.9	6.5	12.2
1999.....	262.6	13.7	1.1	0.5	0.7	1.7	2.6	3.9	6.9	12.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Diseases of heart (I00-I09,I11,I13,I20-I51)										
White, both sexes										
2005 4/.....	235.2	7.1	0.7	0.4	0.6	1.6	2.9	4.8	8.4	16.7
2004 4/.....	237.4	9.0	0.9	0.4	0.6	1.4	2.7	4.5	8.4	17.2
2003 4/.....	251.7	8.8	1.0	0.4	0.7	1.6	2.9	4.5	8.8	17.7
2002.....	258.5	10.6	0.9	0.4	0.7	1.7	2.5	4.6	8.1	17.7
2001.....	262.8	9.9	1.2	0.4	0.7	1.5	2.6	4.5	8.5	17.3
2000.....	270.2	11.5	1.1	0.4	0.7	1.7	2.6	4.2	7.9	16.7
1999.....	277.7	11.8	1.0	0.5	0.7	2.0	2.6	4.1	8.5	17.8
White, male										
2005 4/.....	234.9	7.6	0.9	0.4	0.7	2.1	3.9	6.8	11.4	23.4
2004 4/.....	236.5	9.6	1.0	0.3	0.7	1.8	3.6	6.0	11.6	23.8
2003 4/.....	249.5	9.4	0.8	0.4	0.8	2.0	3.5	5.7	12.0	25.1
2002.....	256.0	11.3	1.0	0.4	0.8	2.2	3.3	6.1	11.2	24.6
2001.....	257.8	10.6	1.2	0.4	0.8	1.8	3.4	5.9	11.3	24.4
2000.....	265.8	11.9	1.3	0.5	0.8	2.0	3.3	5.3	10.7	23.5
1999.....	272.9	12.0	1.2	0.6	0.7	2.4	3.0	5.2	11.4	25.2
White, female										
2005 4/.....	235.5	6.6	0.6	0.4	0.5	1.1	1.8	2.7	5.3	9.7
2004 4/.....	238.3	8.4	0.9	0.4	0.6	1.0	1.7	2.9	5.0	10.3
2003 4/.....	253.8	8.1	1.1	0.4	0.5	1.3	2.2	3.2	5.4	10.2
2002.....	261.0	9.9	0.8	0.5	0.5	1.1	1.6	3.0	5.0	10.7
2001.....	267.7	9.2	1.1	0.5	0.7	1.1	1.8	3.1	5.6	10.0
2000.....	274.5	11.1	0.8	0.4	0.5	1.3	1.9	2.9	5.0	9.8
1999.....	282.4	11.6	0.9	0.5	0.6	1.6	2.1	3.0	5.4	10.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Diseases of heart (I00-I09,I11,I13,I20-I51)										
Black, both sexes										
2005 4/.....	189.8	16.1	1.7	1.1	1.2	3.4	6.9	13.1	23.7	41.5
2004 4/.....	192.3	15.3	2.4	0.9	1.5	3.6	6.3	13.3	22.1	38.1
2003 4/.....	202.8	20.2	2.2	1.0	1.2	3.5	7.8	13.0	23.0	39.6
2002.....	205.6	21.3	1.6	*	1.3	3.4	6.5	13.4	23.2	40.8
2001.....	208.8	21.6	3.0	0.8	1.3	3.3	7.5	12.9	22.6	39.8
2000.....	211.8	19.8	2.0	0.9	1.4	4.0	7.4	12.8	21.5	38.2
1999.....	217.2	23.8	2.0	0.9	1.4	4.2	7.7	11.7	21.5	42.7
Black, male										
2005 4/.....	194.8	18.3	*	*	1.2	4.9	9.0	16.5	31.1	56.1
2004 4/.....	196.7	16.3	2.4	*	1.4	4.8	8.0	16.7	28.4	50.9
2003 4/.....	206.0	24.1	2.5	*	1.3	4.5	10.2	15.7	28.6	51.1
2002.....	206.3	22.1	*	*	1.5	4.5	8.0	17.5	30.0	54.4
2001.....	209.0	19.5	3.2	*	1.4	4.4	9.7	16.0	29.3	52.1
2000.....	211.1	19.8	1.9	*	1.4	4.8	8.7	15.7	26.5	48.6
1999.....	218.5	23.4	2.1	*	1.6	5.8	9.3	14.8	27.1	56.3
Black, female										
2005 4/.....	185.2	13.9	2.0	1.4	1.2	1.9	4.7	9.8	17.0	28.4
2004 4/.....	188.3	14.2	2.3	*	1.5	2.3	4.5	10.1	16.4	26.7
2003 4/.....	200.0	16.1	2.0	*	*	2.3	5.5	10.5	18.1	29.3
2002.....	205.0	20.6	1.8	*	*	2.4	5.0	9.7	17.0	28.6
2001.....	208.6	23.9	2.7	*	1.2	2.3	5.5	10.1	16.7	28.9
2000.....	212.6	19.8	2.0	*	1.3	3.3	6.3	10.2	16.9	29.0
1999.....	216.0	24.2	1.9	*	*	2.5	6.1	9.0	16.5	30.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Hypertensive heart disease (I11)										
All races 3/, both sexes										
2005.....	9.9	*	*	*	*	*	0.2	0.6	1.1	2.2
2004.....	9.7	*	*	*	*	*	0.1	0.5	1.0	1.9
2003.....	9.7	*	*	*	*	*	0.2	0.5	1.0	1.9
2002.....	9.2	*	*	*	*	*	0.1	0.4	1.0	1.8
2001.....	8.7	*	*	*	*	*	*	0.3	0.9	1.6
2000.....	8.4	*	*	*	*	*	0.1	0.4	0.7	1.4
1999.....	8.1	*	*	*	*	*	0.1	0.3	0.8	1.5
All races 3/, male										
2005.....	9.1	*	*	*	*	*	0.2	0.7	1.4	3.2
2004.....	8.9	*	*	*	*	*	*	0.7	1.5	2.8
2003.....	8.8	*	*	*	*	*	0.3	0.7	1.4	2.7
2002.....	8.2	*	*	*	*	*	0.2	0.7	1.4	2.6
2001.....	7.5	*	*	*	*	*	*	0.4	1.2	2.1
2000.....	7.2	*	*	*	*	*	0.2	0.4	0.9	1.9
1999.....	6.8	*	*	*	*	*	*	0.4	1.0	2.0
All races 3/, female										
2005.....	10.7	*	*	*	*	*	*	0.4	0.7	1.2
2004.....	10.5	*	*	*	*	*	*	0.3	0.5	1.0
2003.....	10.7	*	*	*	*	*	*	0.3	0.7	1.2
2002.....	10.2	*	*	*	*	*	*	*	0.6	1.0
2001.....	9.8	*	*	*	*	*	*	0.3	0.5	1.1
2000.....	9.7	*	*	*	*	*	*	0.3	0.5	1.0
1999.....	9.4	*	*	*	*	*	*	*	0.5	1.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Hypertensive heart disease (I11)										
Black, both sexes										
2005 4/.....	17.7	*	*	*	*	*	0.7	1.6	3.4	6.7
2004 4/.....	17.6	*	*	*	*	*	*	2.1	3.2	6.1
2003 4/.....	17.9	*	*	*	*	*	0.6	1.5	3.5	6.0
2002.....	17.2	*	*	*	*	*	*	1.5	3.9	5.8
2001.....	16.0	*	*	*	*	*	*	1.1	3.1	5.1
2000.....	16.1	*	*	*	*	*	*	1.4	2.8	4.9
1999.....	15.5	*	*	*	*	*	*	1.0	2.6	5.7
Black, male										
2005 4/.....	18.5	*	*	*	*	*	*	1.7	4.2	9.2
2004 4/.....	18.4	*	*	*	*	*	*	2.5	4.6	8.8
2003 4/.....	18.1	*	*	*	*	*	*	1.7	4.3	7.3
2002.....	17.5	*	*	*	*	*	*	2.4	5.4	8.5
2001.....	15.9	*	*	*	*	*	*	*	4.0	5.9
2000.....	15.6	*	*	*	*	*	*	1.6	3.5	5.8
1999.....	15.2	*	*	*	*	*	*	*	2.8	7.3
Black, female										
2005 4/.....	17.0	*	*	*	*	*	*	1.6	2.7	4.4
2004 4/.....	16.9	*	*	*	*	*	*	1.8	2.1	3.6
2003 4/.....	17.8	*	*	*	*	*	*	*	2.7	4.8
2002.....	16.9	*	*	*	*	*	*	*	2.6	3.4
2001.....	16.2	*	*	*	*	*	*	*	2.3	4.4
2000.....	16.5	*	*	*	*	*	*	*	2.1	4.1
1999.....	15.9	*	*	*	*	*	*	*	2.5	4.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Ischemic heart diseases (I20-I25)										
All races 3/, both sexes										
2005.....	150.4	*	*	*	*	0.1	0.6	1.4	3.6	9.1
2004.....	153.7	*	*	*	*	0.1	0.5	1.5	3.9	9.4
2003.....	165.1	0.5	*	*	*	0.2	0.6	1.4	3.9	10.1
2002.....	171.4	0.6	*	*	*	0.1	0.5	1.3	3.6	10.1
2001.....	176.3	0.6	*	*	*	0.2	0.7	1.5	4.0	9.9
2000.....	183.1	0.8	*	*	*	0.2	0.5	1.2	3.4	9.8
1999.....	189.8	0.7	*	*	*	0.3	0.5	1.3	3.7	10.4
All races 3/, male										
2005.....	159.0	*	*	*	*	0.2	0.9	2.2	5.4	13.6
2004.....	161.6	*	*	*	*	0.2	0.7	2.1	6.0	13.9
2003.....	172.2	*	*	*	*	0.2	0.8	2.0	5.9	15.2
2002.....	178.4	*	*	*	*	*	0.7	2.0	5.2	15.2
2001.....	181.7	*	*	*	*	0.2	0.9	2.2	5.9	15.0
2000.....	188.7	*	*	*	*	0.2	0.6	1.8	5.1	14.7
1999.....	195.4	*	*	*	*	0.3	0.6	1.7	5.5	15.8
All races 3/, female										
2005.....	142.0	*	*	*	*	*	0.2	0.6	1.9	4.6
2004.....	146.1	*	*	*	*	*	0.2	0.9	1.8	4.9
2003.....	158.1	*	*	*	*	*	0.4	0.8	1.9	4.9
2002.....	164.7	*	*	*	*	*	0.3	0.6	1.9	4.9
2001.....	171.2	*	*	*	*	*	0.4	0.7	2.1	4.9
2000.....	177.6	*	*	*	*	*	0.3	0.6	1.7	5.0
1999.....	184.5	*	*	*	*	*	0.3	0.8	1.9	5.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Ischemic heart diseases (I20-I25)										
White, both sexes										
2005 4/.....	162.6	*	*	*	*	0.1	0.5	1.2	3.5	8.5
2004 4/.....	166.1	*	*	*	*	*	0.4	1.3	3.6	9.2
2003 4/.....	178.3	*	*	*	*	0.2	0.5	1.3	3.7	9.8
2002.....	185.4	*	*	*	*	0.1	0.4	1.2	3.3	9.6
2001.....	190.5	*	*	*	*	0.1	0.5	1.3	3.8	9.5
2000.....	197.9	0.8	*	*	*	0.2	0.4	1.0	3.1	9.4
1999.....	205.1	*	*	*	*	0.2	0.3	1.2	3.5	9.9
White, male										
2005 4/.....	171.5	*	*	*	*	*	0.8	1.9	5.0	12.7
2004 4/.....	174.3	*	*	*	*	*	0.5	1.9	5.6	13.6
2003 4/.....	185.5	*	*	*	*	*	0.7	1.8	5.6	15.0
2002.....	192.5	*	*	*	*	*	0.6	1.9	5.0	14.6
2001.....	195.8	*	*	*	*	*	0.8	2.0	5.5	14.5
2000.....	203.6	*	*	*	*	*	0.6	1.6	4.8	14.3
1999.....	210.5	*	*	*	*	0.3	0.5	1.6	5.1	15.1
White, female										
2005 4/.....	153.9	*	*	*	*	*	*	0.5	1.8	4.1
2004 4/.....	158.1	*	*	*	*	*	*	0.7	1.6	4.7
2003 4/.....	171.4	*	*	*	*	*	0.4	0.7	1.7	4.4
2002.....	178.4	*	*	*	*	*	*	0.5	1.6	4.5
2001.....	185.3	*	*	*	*	*	0.3	0.6	1.9	4.5
2000.....	192.4	*	*	*	*	*	0.3	0.5	1.4	4.5
1999.....	200.0	*	*	*	*	*	*	0.7	1.7	4.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Ischemic heart diseases (I20-I25)										
Black, both sexes										
2005 4/.....	117.8	*	*	*	*	*	0.9	2.6	5.3	15.5
2004 4/.....	121.0	*	*	*	*	*	1.2	2.9	6.5	12.7
2003 4/.....	129.2	*	*	*	*	*	1.2	2.4	6.1	14.0
2002.....	132.9	*	*	*	*	*	0.9	2.1	5.6	15.1
2001.....	136.9	*	*	*	*	*	1.5	2.8	6.4	14.9
2000.....	140.1	*	*	*	*	*	0.7	2.4	6.0	14.6
1999.....	143.7	*	*	*	*	*	1.2	1.9	5.8	16.1
Black, male										
2005 4/.....	122.9	*	*	*	*	*	1.3	4.2	7.9	23.0
2004 4/.....	125.2	*	*	*	*	*	1.8	3.4	9.6	18.4
2003 4/.....	133.4	*	*	*	*	*	1.6	3.1	9.2	19.6
2002.....	135.3	*	*	*	*	*	*	2.7	7.5	22.0
2001.....	138.9	*	*	*	*	*	1.9	3.9	9.3	21.6
2000.....	141.5	*	*	*	*	*	*	3.7	8.0	20.3
1999.....	146.0	*	*	*	*	*	*	2.9	8.4	23.3
Black, female										
2005 4/.....	113.1	*	*	*	*	*	*	*	3.0	8.8
2004 4/.....	117.1	*	*	*	*	*	*	2.4	3.7	7.6
2003 4/.....	125.3	*	*	*	*	*	*	1.7	3.3	9.1
2002.....	130.8	*	*	*	*	*	*	1.5	4.0	9.0
2001.....	135.2	*	*	*	*	*	*	1.7	3.8	9.0
2000.....	138.8	*	*	*	*	*	*	*	4.3	9.7
1999.....	141.7	*	*	*	*	*	*	*	3.4	9.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Acute myocardial infarction (I21-I22)										
All races 3/, both sexes										
2005.....	50.9	*	*	*	*	*	0.3	0.6	1.3	3.5
2004.....	53.4	*	*	*	*	*	0.2	0.6	1.6	3.8
2003.....	58.7	*	*	*	*	*	0.3	0.5	1.5	4.1
2002.....	62.3	*	*	*	*	*	0.2	0.5	1.5	4.0
2001.....	64.9	*	*	*	*	*	0.2	0.6	1.8	4.2
2000.....	68.5	0.5	*	*	*	*	0.2	0.6	1.3	4.0
1999.....	71.5	*	*	*	*	0.1	0.2	0.5	1.6	4.1
All races 3/, male										
2005.....	54.8	*	*	*	*	*	0.4	1.0	2.0	5.1
2004.....	57.4	*	*	*	*	*	0.3	0.8	2.3	5.7
2003.....	62.6	*	*	*	*	*	0.3	0.8	2.2	6.2
2002.....	66.2	*	*	*	*	*	0.3	0.7	2.1	6.0
2001.....	68.6	*	*	*	*	*	0.3	0.7	2.5	6.2
2000.....	72.7	*	*	*	*	*	0.2	0.8	1.9	5.8
1999.....	76.1	*	*	*	*	*	0.3	0.7	2.3	6.2
All races 3/, female										
2005.....	47.2	*	*	*	*	*	*	0.3	0.7	2.0
2004.....	49.6	*	*	*	*	*	*	0.4	0.8	1.9
2003.....	54.8	*	*	*	*	*	0.2	0.3	0.8	2.0
2002.....	58.4	*	*	*	*	*	*	0.3	0.8	2.1
2001.....	61.3	*	*	*	*	*	*	0.4	1.0	2.1
2000.....	64.6	*	*	*	*	*	*	0.3	0.7	2.2
1999.....	67.0	*	*	*	*	*	*	0.3	0.9	2.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Acute myocardial infarction (I21-I22)										
White, both sexes										
2005 4/.....	55.1	*	*	*	*	*	0.2	0.5	1.3	3.4
2004 4/.....	57.8	*	*	*	*	*	0.1	0.5	1.6	3.8
2003 4/.....	63.4	*	*	*	*	*	0.2	0.5	1.4	4.1
2002.....	67.3	*	*	*	*	*	0.2	0.4	1.4	4.0
2001.....	70.1	*	*	*	*	*	0.2	0.5	1.7	4.1
2000.....	74.1	*	*	*	*	*	0.2	0.5	1.2	4.0
1999.....	77.4	*	*	*	*	*	0.1	0.5	1.5	4.0
White, male										
2005 4/.....	59.5	*	*	*	*	*	0.4	0.8	1.9	4.9
2004 4/.....	62.3	*	*	*	*	*	*	0.7	2.4	5.6
2003 4/.....	67.7	*	*	*	*	*	0.3	0.7	2.0	6.3
2002.....	71.8	*	*	*	*	*	0.3	0.6	2.1	5.9
2001.....	74.4	*	*	*	*	*	0.3	0.6	2.5	6.1
2000.....	78.8	*	*	*	*	*	*	0.6	1.8	6.0
1999.....	82.7	*	*	*	*	*	*	0.6	2.1	6.2
White, female										
2005 4/.....	50.8	*	*	*	*	*	*	*	0.7	1.8
2004 4/.....	53.3	*	*	*	*	*	*	0.4	0.7	1.9
2003 4/.....	59.2	*	*	*	*	*	*	*	0.8	1.9
2002.....	62.8	*	*	*	*	*	*	*	0.7	1.9
2001.....	66.0	*	*	*	*	*	*	0.3	0.9	2.0
2000.....	69.6	*	*	*	*	*	*	0.3	0.6	2.0
1999.....	72.3	*	*	*	*	*	*	0.3	0.9	1.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Acute myocardial infarction (I21-I22)

Black, both sexes

2005 4/.....	39.8	*	*	*	*	*	*	1.4	2.0	5.4
2004 4/.....	42.0	*	*	*	*	*	*	1.2	2.2	4.9
2003 4/.....	45.7	*	*	*	*	*	*	0.9	2.3	5.6
2002.....	48.5	*	*	*	*	*	*	0.9	2.3	5.5
2001.....	50.0	*	*	*	*	*	*	1.1	2.5	5.8
2000.....	52.2	*	*	*	*	*	*	1.2	2.3	4.9
1999.....	53.3	*	*	*	*	*	*	0.8	2.4	5.5

Black, male

2005 4/.....	40.3	*	*	*	*	*	*	2.1	2.9	7.5
2004 4/.....	42.4	*	*	*	*	*	*	*	2.7	7.7
2003 4/.....	46.6	*	*	*	*	*	*	*	3.5	7.5
2002.....	48.3	*	*	*	*	*	*	*	3.0	7.6
2001.....	49.2	*	*	*	*	*	*	*	3.4	8.0
2000.....	52.0	*	*	*	*	*	*	1.6	2.7	6.5
1999.....	53.4	*	*	*	*	*	*	*	3.4	7.7

Black, female

2005 4/.....	39.2	*	*	*	*	*	*	*	*	3.6
2004 4/.....	41.6	*	*	*	*	*	*	*	1.8	2.4
2003 4/.....	44.9	*	*	*	*	*	*	*	*	3.9
2002.....	48.8	*	*	*	*	*	*	*	1.6	3.6
2001.....	50.6	*	*	*	*	*	*	*	1.7	3.8
2000.....	52.5	*	*	*	*	*	*	*	1.9	3.6
1999.....	53.2	*	*	*	*	*	*	*	1.4	3.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other forms of chronic ischemic heart disease (I20,I25)										
All races 3/, both sexes										
2005.....	98.2	*	*	*	*	*	0.3	0.8	2.3	5.5
2004.....	99.5	*	*	*	*	*	0.3	0.9	2.3	5.5
2003.....	105.3	*	*	*	*	*	0.3	0.9	2.4	5.8
2002.....	108.0	*	*	*	*	*	0.3	0.8	2.0	6.0
2001.....	110.3	*	*	*	*	*	0.4	0.9	2.2	5.6
2000.....	113.3	*	*	*	*	*	0.3	0.6	2.0	5.7
1999.....	117.2	*	*	*	*	0.1	0.3	0.7	2.1	6.2
All races 3/, male										
2005.....	102.9	*	*	*	*	*	0.5	1.2	3.4	8.4
2004.....	103.3	*	*	*	*	*	0.4	1.3	3.6	8.1
2003.....	108.5	*	*	*	*	*	0.4	1.2	3.7	8.8
2002.....	110.9	*	*	*	*	*	0.4	1.3	3.0	9.2
2001.....	111.9	*	*	*	*	*	0.5	1.5	3.3	8.6
2000.....	114.8	*	*	*	*	*	0.4	1.1	3.1	8.7
1999.....	118.0	*	*	*	*	0.2	0.3	1.0	3.2	9.4
All races 3/, female										
2005.....	93.7	*	*	*	*	*	*	0.3	1.2	2.5
2004.....	95.7	*	*	*	*	*	*	0.5	0.9	3.0
2003.....	102.3	*	*	*	*	*	0.2	0.5	1.0	2.8
2002.....	105.2	*	*	*	*	*	*	0.4	1.0	2.8
2001.....	108.8	*	*	*	*	*	0.2	0.3	1.1	2.7
2000.....	111.9	*	*	*	*	*	*	0.2	1.0	2.8
1999.....	116.4	*	*	*	*	*	*	0.4	1.0	3.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other forms of chronic ischemic heart disease (I20,I25)

White, both sexes

2005 4/.....	106.2	*	*	*	*	*	0.3	0.7	2.2	5.0
2004 4/.....	107.5	*	*	*	*	*	0.2	0.8	2.0	5.4
2003 4/.....	113.8	*	*	*	*	*	0.3	0.8	2.2	5.6
2002.....	116.9	*	*	*	*	*	0.2	0.8	1.8	5.6
2001.....	119.1	*	*	*	*	*	0.3	0.8	2.0	5.3
2000.....	122.6	*	*	*	*	*	0.2	0.6	1.9	5.3
1999.....	126.5	*	*	*	*	*	0.2	0.7	1.9	5.8

White, male

2005 4/.....	110.7	*	*	*	*	*	0.4	1.1	3.1	7.6
2004 4/.....	111.1	*	*	*	*	*	0.3	1.2	3.1	7.9
2003 4/.....	116.6	*	*	*	*	*	0.4	1.1	3.5	8.5
2002.....	119.4	*	*	*	*	*	0.4	1.3	2.8	8.6
2001.....	120.1	*	*	*	*	*	0.4	1.4	3.0	8.2
2000.....	123.5	*	*	*	*	*	0.3	0.9	2.9	8.2
1999.....	126.5	*	*	*	*	*	*	0.9	3.0	8.8

White, female

2005 4/.....	101.9	*	*	*	*	*	*	0.3	1.1	2.3
2004 4/.....	103.9	*	*	*	*	*	*	0.3	0.9	2.8
2003 4/.....	111.0	*	*	*	*	*	*	0.4	0.9	2.5
2002.....	114.4	*	*	*	*	*	*	0.3	0.8	2.5
2001.....	118.1	*	*	*	*	*	*	*	0.9	2.4
2000.....	121.6	*	*	*	*	*	*	*	0.8	2.4
1999.....	126.5	*	*	*	*	*	*	0.4	0.9	2.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other forms of chronic ischemic heart disease (I20,I25)

Black, both sexes

2005 4/.....	76.6	*	*	*	*	*	*	1.1	3.3	9.8
2004 4/.....	78.0	*	*	*	*	*	*	1.6	4.1	7.7
2003 4/.....	82.3	*	*	*	*	*	*	1.5	3.7	8.2
2002.....	83.2	*	*	*	*	*	*	1.1	3.2	9.6
2001.....	85.8	*	*	*	*	*	0.8	1.7	3.8	8.9
2000.....	86.6	*	*	*	*	*	*	1.3	3.6	9.5
1999.....	89.3	*	*	*	*	*	0.7	1.1	3.4	10.1

Black, male

2005 4/.....	81.0	*	*	*	*	*	*	2.0	4.9	15.3
2004 4/.....	81.5	*	*	*	*	*	*	1.9	6.7	10.5
2003 4/.....	85.5	*	*	*	*	*	*	2.1	5.7	11.7
2002.....	85.7	*	*	*	*	*	*	*	4.1	14.4
2001.....	88.2	*	*	*	*	*	*	2.4	5.9	13.3
2000.....	88.0	*	*	*	*	*	*	2.1	5.0	13.4
1999.....	91.1	*	*	*	*	*	*	1.7	5.0	14.9

Black, female

2005 4/.....	72.6	*	*	*	*	*	*	*	1.8	4.9
2004 4/.....	74.7	*	*	*	*	*	*	*	1.7	5.2
2003 4/.....	79.4	*	*	*	*	*	*	*	2.0	5.0
2002.....	81.0	*	*	*	*	*	*	*	2.3	5.3
2001.....	83.5	*	*	*	*	*	*	*	1.9	4.9
2000.....	85.3	*	*	*	*	*	*	*	2.4	5.9
1999.....	87.6	*	*	*	*	*	*	*	1.9	5.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Atherosclerotic cardiovascular disease, so described (I25.0)										
All races 3/, both sexes										
2005.....	21.2	*	*	*	*	*	*	0.2	0.7	1.9
2004.....	21.7	*	*	*	*	*	*	0.2	0.8	1.7
2003.....	23.1	*	*	*	*	*	*	0.3	0.8	1.9
2002.....	23.6	*	*	*	*	*	*	0.2	0.7	1.9
2001.....	23.9	*	*	*	*	*	*	0.2	0.7	1.8
2000.....	24.7	*	*	*	*	*	*	0.2	0.5	1.7
1999.....	25.8	*	*	*	*	*	*	0.2	0.6	2.0
All races 3/, male										
2005.....	23.6	*	*	*	*	*	*	0.4	1.1	3.0
2004.....	23.9	*	*	*	*	*	*	0.3	1.2	2.5
2003.....	25.2	*	*	*	*	*	*	0.4	1.3	2.9
2002.....	25.4	*	*	*	*	*	*	0.3	1.1	3.0
2001.....	25.3	*	*	*	*	*	*	0.3	1.0	2.8
2000.....	26.0	*	*	*	*	*	*	0.3	0.9	2.8
1999.....	26.8	*	*	*	*	*	*	0.2	0.9	3.1
All races 3/, female										
2005.....	18.8	*	*	*	*	*	*	*	0.3	0.8
2004.....	19.5	*	*	*	*	*	*	*	0.3	0.8
2003.....	21.1	*	*	*	*	*	*	*	0.2	0.9
2002.....	21.9	*	*	*	*	*	*	*	0.3	0.8
2001.....	22.5	*	*	*	*	*	*	*	0.4	0.8
2000.....	23.5	*	*	*	*	*	*	*	0.2	0.7
1999.....	24.8	*	*	*	*	*	*	*	0.3	0.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)

All races 3/, both sexes

2005.....	77.0	*	*	*	*	*	0.2	0.5	1.6	3.6
2004.....	77.8	*	*	*	*	*	0.2	0.7	1.5	3.9
2003.....	82.2	*	*	*	*	*	0.3	0.6	1.6	3.9
2002.....	84.4	*	*	*	*	*	0.2	0.6	1.3	4.1
2001.....	86.4	*	*	*	*	*	0.3	0.7	1.5	3.8
2000.....	88.6	*	*	*	*	*	0.2	0.5	1.5	4.0
1999.....	91.4	*	*	*	*	0.1	0.2	0.6	1.5	4.3

All races 3/, male

2005.....	79.3	*	*	*	*	*	0.4	0.9	2.3	5.4
2004.....	79.4	*	*	*	*	*	0.3	1.0	2.3	5.6
2003.....	83.3	*	*	*	*	*	0.3	0.8	2.3	5.9
2002.....	85.5	*	*	*	*	*	0.3	0.9	1.9	6.2
2001.....	86.5	*	*	*	*	*	0.4	1.1	2.3	5.8
2000.....	88.8	*	*	*	*	*	0.2	0.8	2.2	5.9
1999.....	91.2	*	*	*	*	0.2	0.2	0.8	2.2	6.3

All races 3/, female

2005.....	74.8	*	*	*	*	*	*	0.2	0.9	1.7
2004.....	76.2	*	*	*	*	*	*	0.4	0.7	2.1
2003.....	81.1	*	*	*	*	*	*	0.4	0.8	1.9
2002.....	83.3	*	*	*	*	*	*	0.3	0.8	2.0
2001.....	86.2	*	*	*	*	*	*	0.2	0.7	1.9
2000.....	88.4	*	*	*	*	*	*	*	0.8	2.1
1999.....	91.6	*	*	*	*	*	*	0.3	0.7	2.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)										
White, both sexes										
2005 4/.....	84.7	*	*	*	*	*	0.2	0.5	1.6	3.3
2004 4/.....	85.4	*	*	*	*	*	0.2	0.6	1.4	3.7
2003 4/.....	90.2	*	*	*	*	*	0.2	0.5	1.5	3.7
2002.....	92.8	*	*	*	*	*	0.2	0.6	1.2	3.9
2001.....	94.8	*	*	*	*	*	0.2	0.6	1.4	3.6
2000.....	97.4	*	*	*	*	*	0.1	0.4	1.4	3.7
1999.....	100.4	*	*	*	*	*	0.1	0.5	1.4	4.0
White, male										
2005 4/.....	87.0	*	*	*	*	*	0.3	0.8	2.3	5.0
2004 4/.....	87.0	*	*	*	*	*	0.2	0.9	2.1	5.5
2003 4/.....	91.3	*	*	*	*	*	0.3	0.7	2.2	5.7
2002.....	93.9	*	*	*	*	*	0.3	0.9	1.8	5.9
2001.....	94.8	*	*	*	*	*	0.4	1.1	2.1	5.5
2000.....	97.5	*	*	*	*	*	*	0.7	2.1	5.6
1999.....	99.9	*	*	*	*	*	*	0.7	2.1	6.0
White, female										
2005 4/.....	82.4	*	*	*	*	*	*	*	0.9	1.6
2004 4/.....	83.8	*	*	*	*	*	*	0.3	0.6	2.0
2003 4/.....	89.1	*	*	*	*	*	*	0.3	0.7	1.6
2002.....	91.7	*	*	*	*	*	*	*	0.6	1.9
2001.....	94.9	*	*	*	*	*	*	*	0.6	1.7
2000.....	97.3	*	*	*	*	*	*	*	0.6	1.8
1999.....	100.8	*	*	*	*	*	*	0.3	0.6	2.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other heart diseases (I26-I51)										
All races 3/, both sexes										
2005.....	57.6	8.3	0.9	0.5	0.6	1.7	2.7	3.8	5.4	7.8
2004.....	56.6	10.0	1.1	0.4	0.7	1.5	2.5	3.6	5.0	7.8
2003.....	58.5	10.4	1.1	0.5	0.7	1.7	2.7	3.6	5.4	7.8
2002.....	58.8	11.7	1.0	0.4	0.7	1.8	2.3	3.8	5.2	8.1
2001.....	58.6	11.1	1.4	0.5	0.7	1.5	2.5	3.6	5.1	8.0
2000.....	58.8	12.0	1.1	0.5	0.7	1.8	2.6	3.5	5.2	7.5
1999.....	59.4	12.9	1.2	0.5	0.7	2.0	2.6	3.5	5.3	8.2
All races 3/, male										
2005.....	51.4	8.9	0.9	0.4	0.7	2.3	3.4	4.9	6.7	9.6
2004.....	50.6	10.5	1.1	0.3	0.8	1.9	3.2	4.5	6.1	9.4
2003.....	52.3	11.4	1.0	0.5	0.8	2.1	3.2	4.1	6.5	9.5
2002.....	52.4	12.1	1.0	0.4	0.8	2.4	3.0	4.6	6.6	9.6
2001.....	51.8	10.8	1.5	0.4	0.8	1.9	3.1	4.3	6.0	9.9
2000.....	52.2	12.2	1.2	0.6	0.9	2.1	3.1	4.2	6.2	9.1
1999.....	53.1	13.1	1.3	0.5	0.8	2.4	3.0	4.0	6.4	10.3
All races 3/, female										
2005.....	63.5	7.7	0.8	0.5	0.6	1.1	1.9	2.7	4.1	6.0
2004.....	62.4	9.4	1.1	0.5	0.6	1.2	1.8	2.6	3.9	6.1
2003.....	64.5	9.3	1.2	0.4	0.5	1.3	2.1	3.1	4.3	6.1
2002.....	65.0	11.4	1.0	0.5	0.5	1.1	1.7	3.0	3.8	6.6
2001.....	65.2	11.5	1.3	0.5	0.7	1.1	1.9	2.9	4.1	6.0
2000.....	65.1	11.8	0.9	0.4	0.6	1.4	2.1	2.9	4.1	5.9
1999.....	65.5	12.7	1.0	0.4	0.7	1.5	2.2	2.9	4.2	6.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other heart diseases (I26-I51)										
White, both sexes										
2005 4/.....	61.4	6.9	0.7	0.4	0.5	1.5	2.3	3.2	4.1	6.4
2004 4/.....	60.4	8.7	0.9	0.3	0.6	1.3	2.2	2.8	4.0	6.5
2003 4/.....	62.2	8.2	0.9	0.4	0.6	1.5	2.2	2.8	4.4	6.4
2002.....	62.6	10.0	0.9	0.4	0.6	1.5	1.9	3.0	4.1	6.6
2001.....	62.3	9.3	1.1	0.4	0.7	1.3	2.0	2.9	4.1	6.5
2000.....	62.5	10.6	1.0	0.4	0.6	1.4	2.1	2.8	4.3	6.1
1999.....	62.9	11.0	1.0	0.4	0.6	1.8	2.1	2.7	4.4	6.7
White, male										
2005 4/.....	54.0	7.2	0.8	0.3	0.6	1.9	2.9	4.2	5.2	8.1
2004 4/.....	53.0	9.1	0.9	*	0.7	1.6	2.9	3.6	5.0	8.0
2003 4/.....	54.7	8.8	0.8	0.4	0.8	1.8	2.5	3.2	5.5	7.9
2002.....	54.9	10.5	1.0	0.4	0.7	2.1	2.5	3.6	5.2	7.9
2001.....	54.1	9.6	1.2	0.4	0.7	1.5	2.4	3.5	4.8	8.1
2000.....	54.6	10.8	1.2	0.5	0.8	1.7	2.5	3.4	5.2	7.5
1999.....	55.1	11.4	1.1	0.5	0.7	2.1	2.4	3.3	5.4	8.5
White, female										
2005 4/.....	68.7	6.5	0.6	0.4	0.5	1.0	1.5	2.1	3.0	4.6
2004 4/.....	67.6	8.3	0.8	0.4	0.5	1.0	1.5	2.0	3.1	4.9
2003 4/.....	69.6	7.6	1.0	0.3	0.4	1.1	1.8	2.4	3.2	5.0
2002.....	70.2	9.6	0.8	0.4	0.5	1.0	1.3	2.3	2.9	5.3
2001.....	70.4	8.9	1.1	0.5	0.6	1.0	1.5	2.3	3.4	4.8
2000.....	70.2	10.4	0.7	0.3	0.4	1.2	1.6	2.2	3.3	4.6
1999.....	70.5	10.7	0.8	0.4	0.6	1.4	1.9	2.2	3.4	4.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other heart diseases (I26-I51)										
Black, both sexes										
2005 4/.....	51.4	15.2	1.7	1.1	1.1	3.2	5.1	8.6	14.0	18.5
2004 4/.....	50.8	14.8	2.3	0.8	1.3	3.0	4.5	8.0	11.9	17.9
2003 4/.....	52.5	19.6	2.0	0.9	1.0	3.0	5.9	8.9	12.8	18.1
2002.....	52.5	20.2	1.5	*	1.2	3.1	5.1	9.5	13.0	18.9
2001.....	52.8	20.1	2.9	0.8	1.1	2.9	5.5	8.7	12.3	18.4
2000.....	52.7	19.0	1.9	0.8	1.4	3.6	6.0	8.8	11.9	17.8
1999.....	54.4	22.3	2.0	0.7	1.2	3.5	5.6	8.6	12.4	19.5
Black, male										
2005 4/.....	50.6	17.7	*	*	1.2	4.6	6.6	10.4	17.8	22.6
2004 4/.....	50.4	15.7	2.3	*	1.3	4.0	5.6	10.4	13.9	21.5
2003 4/.....	51.5	23.2	2.2	*	1.2	3.9	7.5	10.5	14.3	22.5
2002.....	50.8	20.9	*	*	1.3	4.0	6.2	11.9	16.4	22.8
2001.....	51.7	17.7	3.2	*	1.2	3.8	7.0	10.7	14.9	23.1
2000.....	51.5	18.9	1.9	*	1.4	4.3	6.7	10.1	14.3	21.6
1999.....	54.3	21.8	2.1	*	1.5	4.8	6.8	10.2	15.2	24.3
Black, female										
2005 4/.....	52.2	12.6	2.0	1.3	*	1.7	3.7	6.9	10.6	14.8
2004 4/.....	51.1	13.9	2.3	*	1.3	2.0	3.4	5.8	10.1	14.7
2003 4/.....	53.4	15.8	1.8	*	*	2.1	4.2	7.3	11.4	14.3
2002.....	54.0	19.4	1.8	*	*	2.2	4.0	7.4	9.8	15.5
2001.....	53.8	22.7	2.5	*	*	2.0	4.1	6.9	10.0	14.1
2000.....	53.9	19.2	2.0	*	1.3	2.9	5.3	7.6	9.8	14.5
1999.....	54.5	22.8	1.8	*	*	2.2	4.5	7.1	9.9	15.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)										
All races 3/, both sexes										
2005.....	37.0	7.1	0.7	0.4	0.5	1.5	2.4	3.5	4.9	7.0
2004.....	36.5	8.7	0.9	0.3	0.6	1.4	2.3	3.2	4.5	6.9
2003.....	38.0	8.6	0.9	0.4	0.6	1.5	2.4	3.3	4.9	7.0
2002.....	38.5	10.5	0.8	0.3	0.6	1.6	2.1	3.5	4.6	7.3
2001.....	37.9	9.7	1.1	0.4	0.7	1.3	2.3	3.3	4.6	7.0
2000.....	38.3	10.6	0.9	0.4	0.6	1.5	2.4	3.2	4.8	6.8
1999.....	39.0	11.9	0.9	0.4	0.7	1.8	2.4	3.2	4.9	7.4
All races 3/, male										
2005.....	34.9	7.7	0.8	0.4	0.6	2.1	3.1	4.4	6.0	8.7
2004.....	34.5	9.2	0.9	0.2	0.7	1.8	3.0	4.1	5.5	8.3
2003.....	35.8	9.2	0.9	0.4	0.7	1.8	2.9	3.7	5.8	8.6
2002.....	36.3	10.9	0.8	0.3	0.7	2.2	2.7	4.2	5.8	8.8
2001.....	35.6	9.1	1.2	0.3	0.7	1.7	2.8	3.8	5.4	8.8
2000.....	36.1	10.5	1.0	0.5	0.8	1.9	2.8	3.8	5.7	8.3
1999.....	37.1	12.1	1.0	0.4	0.8	2.2	2.7	3.7	5.9	9.3
All races 3/, female										
2005.....	39.0	6.4	0.7	0.4	0.4	1.0	1.7	2.4	3.7	5.3
2004.....	38.5	8.2	0.8	0.4	0.6	1.0	1.6	2.4	3.5	5.5
2003.....	40.1	7.9	1.0	0.3	0.4	1.1	1.8	2.8	3.9	5.5
2002.....	40.6	10.0	0.8	0.4	0.4	1.0	1.5	2.7	3.4	5.9
2001.....	40.2	10.4	1.0	0.4	0.6	1.0	1.7	2.7	3.7	5.3
2000.....	40.4	10.7	0.7	0.3	0.5	1.2	2.0	2.6	3.8	5.3
1999.....	40.9	11.6	0.8	0.4	0.5	1.4	2.0	2.7	3.9	5.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)										
White, both sexes										
2005 4/.....	38.9	6.0	0.6	0.3	0.5	1.3	2.0	2.9	3.8	5.8
2004 4/.....	38.2	7.7	0.7	0.3	0.5	1.2	2.0	2.6	3.7	5.8
2003 4/.....	39.8	7.1	0.7	0.3	0.5	1.3	1.9	2.5	4.0	5.9
2002.....	40.4	9.1	0.7	0.3	0.5	1.4	1.7	2.7	3.6	6.1
2001.....	39.6	8.2	0.9	0.3	0.6	1.1	1.8	2.6	3.7	5.7
2000.....	40.0	9.6	0.8	0.3	0.5	1.3	1.9	2.5	3.9	5.5
1999.....	40.6	10.3	0.8	0.4	0.6	1.6	1.9	2.5	4.0	6.0
White, male										
2005 4/.....	36.0	6.4	0.7	0.3	0.6	1.7	2.6	3.8	4.7	7.4
2004 4/.....	35.5	8.1	0.7	*	0.6	1.4	2.6	3.3	4.6	7.0
2003 4/.....	36.9	7.4	0.7	0.3	0.7	1.6	2.2	2.9	5.0	7.2
2002.....	37.5	9.5	0.8	0.3	0.6	1.9	2.3	3.3	4.6	7.3
2001.....	36.5	8.2	0.9	0.3	0.6	1.3	2.3	3.1	4.4	7.2
2000.....	37.1	9.4	0.9	0.4	0.7	1.5	2.3	3.1	4.8	6.9
1999.....	37.9	10.7	1.0	0.4	0.6	1.9	2.2	3.0	4.9	7.7
White, female										
2005 4/.....	41.6	5.6	0.6	0.3	0.4	0.9	1.4	1.9	2.8	4.2
2004 4/.....	40.9	7.3	0.6	0.4	0.5	0.9	1.4	1.9	2.7	4.5
2003 4/.....	42.6	6.8	0.8	*	0.4	1.0	1.5	2.2	2.9	4.4
2002.....	43.2	8.7	0.6	0.3	0.4	0.9	1.1	2.1	2.6	4.8
2001.....	42.6	8.2	0.8	0.3	0.5	0.9	1.3	2.1	3.0	4.2
2000.....	42.8	9.7	0.7	*	0.3	1.0	1.5	1.9	3.0	4.2
1999.....	43.2	10.0	0.7	0.4	0.5	1.4	1.7	2.0	3.1	4.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)										
Black, both sexes										
2005 4/.....	36.1	12.2	1.1	0.8	0.9	2.9	4.7	7.7	12.5	16.1
2004 4/.....	36.3	12.4	1.8	*	1.2	2.7	4.1	7.3	10.7	15.7
2003 4/.....	37.5	15.0	1.6	0.7	0.8	2.6	5.2	8.0	11.3	15.9
2002.....	37.8	17.2	1.1	*	0.9	2.9	4.8	8.7	11.4	17.0
2001.....	38.0	17.2	2.3	0.6	1.1	2.6	4.9	7.9	11.2	16.3
2000.....	38.2	15.6	1.4	0.7	1.1	3.2	5.4	8.0	10.8	15.9
1999.....	39.6	20.2	1.4	0.6	1.1	3.3	5.2	7.8	11.4	17.5
Black, male										
2005 4/.....	37.3	14.8	*	*	1.1	4.2	6.1	9.2	15.7	19.7
2004 4/.....	37.5	13.3	1.8	*	1.1	3.7	5.1	9.4	12.2	19.3
2003 4/.....	38.6	17.8	1.8	*	*	3.4	6.7	9.6	12.2	19.9
2002.....	38.3	18.9	*	*	1.2	3.7	5.8	10.9	14.5	20.5
2001.....	38.9	15.0	2.6	*	1.1	3.4	6.3	9.3	13.5	20.3
2000.....	39.2	14.7	1.6	*	1.2	3.9	6.1	9.1	13.2	19.3
1999.....	41.5	19.8	*	*	1.5	4.6	6.2	9.2	13.8	21.8
Black, female										
2005 4/.....	34.9	9.6	*	*	*	1.6	3.2	6.2	9.6	12.9
2004 4/.....	35.1	11.4	1.7	*	1.3	1.7	3.1	5.3	9.3	12.4
2003 4/.....	36.5	12.1	*	*	*	1.8	3.6	6.6	10.4	12.4
2002.....	37.4	15.4	*	*	*	2.0	3.8	6.7	8.7	13.9
2001.....	37.2	19.5	2.0	*	*	1.7	3.5	6.6	9.2	12.8
2000.....	37.3	16.5	*	*	*	2.6	4.8	6.9	8.7	12.9
1999.....	37.9	20.5	*	*	*	2.0	4.2	6.5	9.2	13.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Cerebrovascular diseases (I60-I69)										
All races 3/, both sexes										
2005.....	48.4	3.1	0.4	0.3	0.2	0.4	0.6	1.0	1.8	3.3
2004.....	51.1	3.1	0.3	0.2	0.2	0.3	0.7	1.1	1.8	3.4
2003.....	54.2	2.5	0.3	0.1	0.2	0.4	0.6	1.0	1.9	3.6
2002.....	56.4	2.9	0.3	0.2	0.3	0.3	0.6	1.0	1.8	3.6
2001.....	57.4	2.7	0.4	0.2	0.2	0.3	0.7	1.1	1.9	3.8
2000.....	59.6	3.3	0.3	0.1	0.2	0.3	0.7	1.1	1.9	3.9
1999.....	60.0	2.7	0.3	0.1	0.2	0.3	0.6	1.0	1.8	4.0
All races 3/, male										
2005.....	38.8	3.5	0.5	0.3	0.3	0.3	0.6	1.1	1.9	3.4
2004.....	40.7	3.4	0.3	*	0.2	0.4	0.7	1.1	1.7	3.7
2003.....	42.9	2.8	0.3	*	0.2	0.5	0.6	1.0	2.0	3.8
2002.....	44.2	3.2	0.4	*	0.3	0.3	0.7	1.0	1.7	3.6
2001.....	45.2	3.1	0.3	0.2	0.2	0.3	0.7	1.1	2.0	3.9
2000.....	46.9	3.8	*	*	0.2	0.4	0.7	1.1	1.9	3.8
1999.....	47.1	3.3	0.3	*	*	0.3	0.6	1.0	1.9	3.9
All races 3/, female										
2005.....	57.8	2.6	0.3	0.3	*	0.4	0.6	0.8	1.6	3.2
2004.....	61.2	2.8	*	*	0.2	0.3	0.7	1.0	1.8	3.1
2003.....	65.1	2.2	0.3	*	*	0.4	0.7	0.9	1.8	3.3
2002.....	68.2	2.5	0.3	*	0.3	*	0.5	0.9	1.9	3.7
2001.....	69.2	2.3	0.4	*	0.2	0.3	0.6	1.0	1.9	3.6
2000.....	71.8	2.7	0.4	*	0.3	0.3	0.7	1.0	1.9	4.0
1999.....	72.3	2.1	0.3	*	0.2	0.3	0.6	1.0	1.8	4.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Cerebrovascular diseases (I60-I69)										
White, both sexes										
2005 4/.....	50.7	2.5	0.4	0.2	0.2	0.3	0.5	0.7	1.5	2.7
2004 4/.....	53.7	2.5	0.2	0.2	0.2	0.3	0.6	0.9	1.4	2.9
2003 4/.....	57.0	2.0	0.2	0.1	0.2	0.4	0.5	0.8	1.6	2.7
2002.....	59.5	2.3	0.3	0.2	0.2	0.2	0.6	0.8	1.5	2.9
2001.....	60.5	1.9	0.3	0.2	0.2	0.3	0.5	0.9	1.7	3.1
2000.....	62.8	2.5	0.3	0.1	0.2	0.3	0.6	1.0	1.6	3.0
1999.....	63.3	2.3	0.3	0.1	0.2	0.3	0.6	0.9	1.5	3.2
White, male										
2005 4/.....	39.7	2.8	0.4	*	*	0.2	0.5	0.8	1.6	2.8
2004 4/.....	41.8	2.8	*	*	*	0.3	0.6	1.1	1.4	3.1
2003 4/.....	44.2	2.0	*	*	*	0.4	0.6	0.8	1.7	2.8
2002.....	45.7	2.6	0.4	*	*	0.3	0.7	1.0	1.4	2.8
2001.....	46.6	2.2	*	*	*	0.3	0.7	0.9	1.8	3.1
2000.....	48.4	3.0	*	*	*	0.4	0.7	1.1	1.6	3.0
1999.....	48.7	2.8	*	*	*	0.3	0.6	0.9	1.7	3.2
White, female										
2005 4/.....	61.6	2.1	*	*	*	0.4	0.5	0.7	1.4	2.7
2004 4/.....	65.3	2.2	*	*	*	0.3	0.6	0.7	1.5	2.7
2003 4/.....	69.5	2.0	*	*	*	0.3	0.5	0.8	1.5	2.5
2002.....	73.0	2.0	*	*	0.3	*	0.5	0.7	1.6	2.9
2001.....	74.0	1.6	0.4	*	*	0.3	0.4	0.9	1.6	3.0
2000.....	76.9	2.0	*	*	0.3	*	0.6	0.9	1.5	3.0
1999.....	77.6	1.9	*	*	*	*	0.6	0.9	1.4	3.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Cerebrovascular diseases (I60-I69)										
Black, both sexes										
2005 4/.....	44.9	6.0	*	*	*	*	1.0	2.2	3.6	6.8
2004 4/.....	46.9	6.0	*	*	*	*	1.0	2.2	4.1	7.0
2003 4/.....	49.3	4.9	*	*	*	0.7	1.2	2.0	4.1	8.8
2002.....	50.0	5.5	*	*	*	*	0.8	1.9	4.2	8.2
2001.....	51.1	6.6	*	*	*	*	1.3	2.2	3.6	8.1
2000.....	52.5	7.5	*	*	*	*	1.1	1.5	4.3	9.6
1999.....	52.2	4.6	*	*	*	*	0.8	1.8	3.6	9.1
Black, male										
2005 4/.....	40.3	6.8	*	*	*	*	*	2.6	4.1	7.1
2004 4/.....	41.5	6.8	*	*	*	*	*	1.7	4.1	8.5
2003 4/.....	43.2	5.9	*	*	*	*	*	2.6	4.1	9.7
2002.....	43.5	5.8	*	*	*	*	*	1.7	4.0	7.5
2001.....	44.6	8.1	*	*	*	*	*	2.8	3.4	8.8
2000.....	46.1	8.6	*	*	*	*	*	*	4.1	9.2
1999.....	45.9	*	*	*	*	*	*	1.9	3.7	8.9
Black, female										
2005 4/.....	49.1	*	*	*	*	*	*	1.9	3.2	6.5
2004 4/.....	51.9	*	*	*	*	*	*	2.6	4.0	5.7
2003 4/.....	54.8	*	*	*	*	*	1.6	1.5	4.1	8.0
2002.....	55.8	*	*	*	*	*	*	2.2	4.3	8.9
2001.....	56.9	*	*	*	*	*	1.7	1.7	3.8	7.4
2000.....	58.3	*	*	*	*	*	*	1.6	4.6	10.0
1999.....	57.9	*	*	*	*	*	*	1.8	3.6	9.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Other diseases of circulatory system (I71-I78)

All races 3/, both sexes

2005.....	8.0	*	*	*	*	*	0.2	0.3	0.5	0.8
2004.....	8.1	*	*	*	*	*	0.2	0.3	0.5	0.8
2003.....	8.5	*	*	*	*	0.1	0.2	0.3	0.4	0.8
2002.....	8.6	*	*	*	*	0.2	0.2	0.3	0.5	0.7
2001.....	8.9	*	*	*	*	0.1	0.2	0.4	0.5	0.9
2000.....	9.3	*	*	*	*	*	0.2	0.3	0.5	0.7
1999.....	9.2	*	*	*	*	*	0.2	0.3	0.4	0.8

All races 3/, male

2005.....	8.5	*	*	*	*	*	0.2	0.3	0.7	1.1
2004.....	8.5	*	*	*	*	*	0.3	0.5	0.6	1.1
2003.....	9.2	*	*	*	*	*	0.3	0.4	0.6	1.1
2002.....	9.4	*	*	*	*	*	0.2	0.3	0.7	0.7
2001.....	9.6	*	*	*	*	*	0.2	0.4	0.7	1.2
2000.....	10.2	*	*	*	*	*	0.2	0.4	0.6	0.8
1999.....	10.1	*	*	*	*	*	0.2	0.4	0.6	1.0

All races 3/, female

2005.....	7.4	*	*	*	*	*	*	0.3	0.3	0.4
2004.....	7.6	*	*	*	*	*	*	*	0.4	0.6
2003.....	7.8	*	*	*	*	*	*	*	0.3	0.5
2002.....	7.9	*	*	*	*	*	*	0.2	0.4	0.7
2001.....	8.2	*	*	*	*	*	*	0.3	0.3	0.6
2000.....	8.3	*	*	*	*	*	*	*	0.4	0.6
1999.....	8.4	*	*	*	*	*	*	*	0.2	0.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other disorders of circulatory system (I80-I99)										
All races 3/, both sexes										
2005.....	1.6	0.6	*	*	*	*	0.1	0.2	0.4	0.6
2004.....	1.6	0.7	*	*	*	*	0.1	0.3	0.4	0.5
2003.....	1.6	0.9	*	*	*	*	0.2	0.3	0.4	0.6
2002.....	1.6	0.7	*	*	*	*	0.1	0.2	0.5	0.6
2001.....	1.6	0.5	*	*	*	*	0.1	0.2	0.4	0.4
2000.....	1.6	0.7	*	*	*	*	0.1	0.3	0.4	0.6
1999.....	1.4	0.6	*	*	*	*	*	0.2	0.3	0.4
All races 3/, male										
2005.....	1.5	*	*	*	*	*	*	0.2	0.4	0.5
2004.....	1.4	*	*	*	*	*	*	0.3	0.4	0.5
2003.....	1.4	1.0	*	*	*	*	*	0.2	0.4	0.5
2002.....	1.4	*	*	*	*	*	*	*	0.5	0.6
2001.....	1.4	*	*	*	*	*	*	*	0.3	0.5
2000.....	1.4	*	*	*	*	*	*	*	0.4	0.6
1999.....	1.3	*	*	*	*	*	*	*	0.3	0.5
All races 3/, female										
2005.....	1.8	*	*	*	*	*	*	0.3	0.4	0.7
2004.....	1.7	*	*	*	*	*	*	0.3	0.4	0.5
2003.....	1.9	*	*	*	*	*	0.2	0.3	0.5	0.6
2002.....	1.9	*	*	*	*	*	*	0.3	0.4	0.6
2001.....	1.9	*	*	*	*	*	*	0.3	0.4	0.4
2000.....	1.8	*	*	*	*	*	*	0.4	0.4	0.5
1999.....	1.6	*	*	*	*	*	*	*	0.3	0.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Other disorders of circulatory system (I80-I99)

White, both sexes

2005 4/.....	1.6	0.7	*	*	*	*	*	0.2	0.3	0.5
2004 4/.....	1.6	0.7	*	*	*	*	*	0.2	0.3	0.4
2003 4/.....	1.6	0.9	*	*	*	*	0.1	0.2	0.3	0.5
2002.....	1.6	0.6	*	*	*	*	*	0.2	0.4	0.5
2001.....	1.7	*	*	*	*	*	*	0.2	0.3	0.3
2000.....	1.6	0.7	*	*	*	*	*	0.2	0.3	0.5
1999.....	1.5	*	*	*	*	*	*	0.1	0.3	0.4

White, male

2005 4/.....	1.4	*	*	*	*	*	*	*	0.3	0.5
2004 4/.....	1.4	*	*	*	*	*	*	*	0.3	0.4
2003 4/.....	1.3	*	*	*	*	*	*	*	0.3	0.5
2002.....	1.3	*	*	*	*	*	*	*	0.4	0.5
2001.....	1.4	*	*	*	*	*	*	*	0.3	0.4
2000.....	1.4	*	*	*	*	*	*	*	0.3	0.5
1999.....	1.2	*	*	*	*	*	*	*	0.2	0.4

White, female

2005 4/.....	1.8	*	*	*	*	*	*	0.3	0.3	0.6
2004 4/.....	1.8	*	*	*	*	*	*	*	0.3	0.4
2003 4/.....	1.9	*	*	*	*	*	*	0.3	0.3	0.4
2002.....	1.9	*	*	*	*	*	*	*	0.3	0.4
2001.....	1.9	*	*	*	*	*	*	*	0.3	0.3
2000.....	1.9	*	*	*	*	*	*	0.3	0.3	0.4
1999.....	1.7	*	*	*	*	*	*	*	0.3	0.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Influenza and pneumonia (J10-J18)

All races 3/, both sexes

2005.....	21.3	6.5	0.7	0.3	0.3	0.3	0.5	0.7	1.0	1.5
2004.....	20.3	6.7	0.7	0.2	0.2	0.3	0.6	0.5	1.0	1.5
2003.....	22.4	8.0	1.0	0.4	0.3	0.4	0.7	0.8	1.1	1.8
2002.....	22.8	6.5	0.7	0.2	0.3	0.4	0.5	0.7	1.0	1.7
2001.....	21.8	7.4	0.7	0.2	0.2	0.3	0.6	0.7	1.0	1.8
2000.....	23.2	7.6	0.7	0.2	0.2	0.3	0.7	0.7	1.1	1.8
1999.....	22.8	8.4	0.8	0.2	0.2	0.4	0.6	0.7	1.0	2.0

All races 3/, male

2005.....	19.2	7.0	0.7	0.3	0.3	0.5	0.6	0.8	1.2	1.8
2004.....	18.6	8.0	0.6	*	0.3	0.4	0.7	0.7	1.2	1.7
2003.....	20.1	9.4	1.1	0.4	0.3	0.5	0.7	0.9	1.3	1.9
2002.....	20.4	7.8	0.8	*	0.2	0.4	0.5	0.7	1.2	1.8
2001.....	19.6	8.1	0.7	0.2	0.2	0.3	0.7	0.7	1.2	2.0
2000.....	20.8	8.4	0.7	0.3	0.2	0.3	0.7	0.8	1.2	1.9
1999.....	20.3	8.8	0.8	0.3	0.3	0.3	0.6	0.8	1.2	2.2

All races 3/, female

2005.....	23.2	5.8	0.7	0.3	0.3	*	0.4	0.7	0.9	1.2
2004.....	22.0	5.4	0.9	*	0.2	0.2	0.4	0.4	0.7	1.2
2003.....	24.6	6.6	0.9	0.4	0.4	0.3	0.7	0.6	0.8	1.6
2002.....	25.1	5.1	0.6	0.2	0.3	0.3	0.4	0.7	0.8	1.6
2001.....	23.9	6.7	0.7	0.2	0.2	0.3	0.5	0.6	0.9	1.6
2000.....	25.6	6.8	0.7	0.2	*	0.3	0.6	0.6	1.0	1.7
1999.....	25.3	8.0	0.9	*	0.2	0.4	0.5	0.6	0.8	1.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Influenza and pneumonia (J10-J18)

White, both sexes

2005 4/.....	23.1	4.9	0.6	0.2	0.2	0.3	0.5	0.7	1.0	1.4
2004 4/.....	22.0	5.1	0.6	*	0.2	0.3	0.5	0.5	0.9	1.3
2003 4/.....	24.4	6.7	0.9	0.3	0.3	0.4	0.7	0.7	1.0	1.6
2002.....	24.9	5.2	0.5	0.2	0.2	0.3	0.4	0.7	0.9	1.5
2001.....	23.6	5.3	0.7	0.2	0.2	0.3	0.5	0.6	0.9	1.5
2000.....	25.2	5.8	0.6	0.2	0.2	0.3	0.6	0.6	1.0	1.6
1999.....	24.8	6.2	0.6	0.2	0.2	0.3	0.5	0.5	0.8	1.7

White, male

2005 4/.....	20.5	5.9	0.6	*	0.2	0.4	0.6	0.7	1.0	1.6
2004 4/.....	19.8	6.2	0.5	*	0.2	0.4	0.7	0.6	1.1	1.4
2003 4/.....	21.4	8.0	1.1	0.3	0.3	0.4	0.7	0.9	1.2	1.8
2002.....	21.9	6.6	0.6	*	*	0.4	0.5	0.7	1.0	1.5
2001.....	20.7	6.2	0.7	*	*	0.3	0.7	0.5	1.0	1.7
2000.....	22.0	6.4	0.5	0.3	*	0.3	0.7	0.6	1.0	1.7
1999.....	21.5	6.7	0.5	0.3	0.3	0.3	0.5	0.6	1.0	1.9

White, female

2005 4/.....	25.7	4.0	0.6	0.3	0.3	*	0.4	0.7	0.9	1.1
2004 4/.....	24.2	3.9	0.7	*	*	*	0.3	0.4	0.7	1.1
2003 4/.....	27.3	5.4	0.8	0.4	0.4	0.3	0.7	0.5	0.7	1.4
2002.....	27.8	3.7	0.5	*	0.3	0.3	0.4	0.7	0.7	1.4
2001.....	26.4	4.4	0.6	*	*	0.3	0.3	0.6	0.8	1.4
2000.....	28.2	5.1	0.7	*	*	0.3	0.5	0.6	0.9	1.6
1999.....	27.9	5.6	0.7	*	*	0.4	0.4	0.5	0.6	1.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Influenza and pneumonia (J10-J18)										
Black, both sexes										
2005 4/.....	14.8	14.8	1.1	*	*	*	*	1.0	1.8	2.6
2004 4/.....	14.8	14.1	1.3	*	*	*	0.8	0.9	1.5	3.0
2003 4/.....	15.4	14.1	1.1	0.6	*	*	0.8	1.6	2.1	3.3
2002.....	15.6	12.9	1.6	*	*	*	0.7	1.0	2.0	3.5
2001.....	15.5	16.3	1.0	*	*	*	1.0	1.2	2.1	3.7
2000.....	16.4	16.7	1.2	*	*	*	1.1	1.4	2.0	2.9
1999.....	16.2	19.0	1.7	*	*	*	1.3	1.7	2.2	4.1
Black, male										
2005 4/.....	14.6	13.9	*	*	*	*	*	*	2.2	3.4
2004 4/.....	15.1	16.3	*	*	*	*	*	*	2.0	3.7
2003 4/.....	16.0	16.1	*	*	*	*	*	1.7	2.8	3.5
2002.....	15.4	13.9	1.7	*	*	*	*	*	2.4	4.0
2001.....	15.9	15.9	*	*	*	*	*	*	2.7	4.5
2000.....	16.7	17.9	1.7	*	*	*	*	2.1	2.5	3.0
1999.....	16.4	18.9	2.0	*	*	*	*	1.9	2.5	4.5
Black, female										
2005 4/.....	14.9	15.7	*	*	*	*	*	*	1.5	1.8
2004 4/.....	14.5	11.7	1.8	*	*	*	*	*	*	2.3
2003 4/.....	14.9	12.1	*	*	*	*	*	1.6	1.4	3.1
2002.....	15.7	11.8	*	*	*	*	*	*	1.6	3.1
2001.....	15.2	16.7	*	*	*	*	*	*	1.4	3.0
2000.....	16.0	15.5	*	*	*	*	*	*	1.6	2.7
1999.....	16.1	19.1	*	*	*	*	*	1.5	1.9	3.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Pneumonia (J12-J18)

All races 3/, both sexes

2005.....	20.6	6.0	0.6	0.2	0.2	0.3	0.5	0.7	1.0	1.5
2004.....	19.9	6.3	0.6	0.2	0.2	0.3	0.5	0.5	0.9	1.4
2003.....	21.8	7.3	0.7	0.2	0.3	0.3	0.6	0.8	1.0	1.7
2002.....	22.5	6.3	0.7	0.2	0.2	0.3	0.4	0.7	1.0	1.7
2001.....	21.7	7.2	0.7	0.2	0.2	0.3	0.6	0.6	1.0	1.8
2000.....	22.6	7.4	0.6	0.2	0.2	0.3	0.6	0.7	1.1	1.7
1999.....	22.2	8.1	0.8	0.2	0.2	0.3	0.5	0.7	1.0	2.0

All races 3/, male

2005.....	18.8	6.6	0.6	0.2	0.2	0.4	0.6	0.8	1.1	1.8
2004.....	18.3	7.5	0.5	*	0.2	0.4	0.7	0.7	1.2	1.7
2003.....	19.6	8.5	0.7	0.2	0.2	0.4	0.7	0.9	1.3	1.9
2002.....	20.2	7.6	0.8	*	0.2	0.4	0.5	0.7	1.2	1.8
2001.....	19.5	7.8	0.7	0.2	0.2	0.3	0.7	0.6	1.2	2.0
2000.....	20.3	8.1	0.6	0.2	*	0.3	0.7	0.8	1.2	1.8
1999.....	19.8	8.5	0.7	0.2	0.2	0.3	0.6	0.8	1.1	2.2

All races 3/, female

2005.....	22.5	5.3	0.6	0.2	0.2	*	0.3	0.7	0.9	1.2
2004.....	21.5	5.0	0.8	*	*	0.2	0.4	0.4	0.7	1.2
2003.....	23.9	6.0	0.6	0.2	0.3	0.2	0.6	0.6	0.8	1.5
2002.....	24.8	5.0	0.6	*	0.2	0.3	0.4	0.7	0.8	1.5
2001.....	23.8	6.7	0.7	*	0.2	0.3	0.5	0.6	0.8	1.6
2000.....	24.8	6.6	0.6	*	*	0.3	0.6	0.6	0.9	1.7
1999.....	24.6	7.7	0.8	*	*	0.3	0.5	0.6	0.8	1.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Pneumonia (J12-J18)

White, both sexes

2005 4/.....	22.4	4.5	0.5	0.2	0.2	0.3	0.5	0.7	0.9	1.4
2004 4/.....	21.6	4.7	0.5	*	0.2	0.3	0.5	0.5	0.9	1.2
2003 4/.....	23.7	6.0	0.6	0.2	0.3	0.3	0.7	0.6	0.9	1.5
2002.....	24.6	5.0	0.5	0.2	0.2	0.3	0.4	0.7	0.9	1.4
2001.....	23.5	5.1	0.6	0.2	0.2	0.3	0.5	0.5	0.9	1.5
2000.....	24.4	5.6	0.5	0.2	0.2	0.3	0.6	0.6	1.0	1.6
1999.....	24.1	5.9	0.6	0.2	0.2	0.3	0.4	0.5	0.8	1.7

White, male

2005 4/.....	20.0	5.3	0.4	*	*	0.4	0.6	0.7	1.0	1.6
2004 4/.....	19.5	5.8	0.4	*	*	0.4	0.7	0.6	1.1	1.4
2003 4/.....	20.9	7.1	0.7	*	0.3	0.4	0.7	0.8	1.1	1.7
2002.....	21.7	6.4	0.5	*	*	0.4	0.5	0.6	1.0	1.5
2001.....	20.6	5.8	0.6	*	*	0.3	0.7	0.5	1.0	1.6
2000.....	21.5	6.1	0.4	0.3	*	0.3	0.6	0.6	1.0	1.7
1999.....	21.0	6.7	0.4	*	0.2	0.3	0.5	0.6	0.9	1.9

White, female

2005 4/.....	24.8	3.7	0.5	*	*	*	0.3	0.7	0.8	1.1
2004 4/.....	23.6	3.5	0.6	*	*	*	0.3	0.4	0.7	1.0
2003 4/.....	26.5	4.8	0.5	*	0.3	*	0.6	0.5	0.7	1.3
2002.....	27.4	3.6	0.5	*	*	*	0.4	0.7	0.7	1.4
2001.....	26.2	4.4	0.6	*	*	0.3	0.3	0.6	0.7	1.4
2000.....	27.3	5.0	0.6	*	*	*	0.5	0.6	0.9	1.5
1999.....	27.1	5.2	0.7	*	*	0.3	0.4	0.4	0.6	1.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Pneumonia (J12-J18)										
Black, both sexes										
2005 4/.....	14.6	14.0	1.0	*	*	*	*	0.9	1.8	2.5
2004 4/.....	14.6	13.5	1.2	*	*	*	0.8	0.9	1.5	3.0
2003 4/.....	15.1	13.0	0.9	*	*	*	0.7	1.6	2.0	3.3
2002.....	15.5	12.7	1.5	*	*	*	0.7	1.0	2.0	3.5
2001.....	15.5	16.3	1.0	*	*	*	1.0	1.2	2.0	3.7
2000.....	16.1	16.4	1.2	*	*	*	1.0	1.4	2.0	2.9
1999.....	16.1	18.3	1.7	*	*	*	1.3	1.7	2.2	4.0
Black, male										
2005 4/.....	14.5	13.9	*	*	*	*	*	*	2.2	3.2
2004 4/.....	14.9	15.1	*	*	*	*	*	*	1.9	3.7
2003 4/.....	15.8	15.2	*	*	*	*	*	1.6	2.8	3.5
2002.....	15.3	13.7	1.6	*	*	*	*	*	2.4	4.0
2001.....	15.8	15.9	*	*	*	*	*	*	2.7	4.5
2000.....	16.5	17.9	1.7	*	*	*	*	2.0	2.5	3.0
1999.....	16.3	17.9	1.9	*	*	*	*	1.9	2.5	4.4
Black, female										
2005 4/.....	14.8	14.2	*	*	*	*	*	*	1.5	1.8
2004 4/.....	14.4	11.7	1.7	*	*	*	*	*	*	2.3
2003 4/.....	14.6	10.8	*	*	*	*	*	1.5	1.4	3.1
2002.....	15.6	11.8	*	*	*	*	*	*	1.6	3.1
2001.....	15.1	16.7	*	*	*	*	*	*	1.4	3.0
2000.....	15.8	14.9	*	*	*	*	*	*	1.6	2.7
1999.....	15.9	18.8	*	*	*	*	*	1.5	1.9	3.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Chronic lower respiratory diseases (J40-J47)										
All races 3/, both sexes										
2005.....	44.2	0.8	0.3	0.3	0.3	0.3	0.4	0.5	0.8	1.2
2004.....	41.5	0.9	0.3	0.2	0.3	0.4	0.4	0.6	0.7	1.3
2003.....	43.5	0.8	0.3	0.2	0.4	0.4	0.5	0.6	0.8	1.4
2002.....	43.3	1.0	0.4	0.2	0.4	0.5	0.5	0.6	0.9	1.6
2001.....	43.2	1.0	0.3	0.2	0.3	0.4	0.5	0.5	0.9	1.6
2000.....	43.4	0.9	0.3	0.2	0.4	0.4	0.5	0.6	0.8	1.4
1999.....	44.5	0.9	0.4	0.2	0.4	0.5	0.5	0.7	0.9	1.3
All races 3/, male										
2005.....	42.8	*	0.4	0.3	0.3	0.4	0.5	0.6	0.7	1.2
2004.....	40.6	1.2	0.4	0.3	0.4	0.5	0.5	0.6	0.9	1.2
2003.....	42.4	1.1	0.5	0.2	0.5	0.5	0.6	0.7	0.9	1.3
2002.....	42.9	1.1	0.6	0.2	0.5	0.6	0.5	0.7	0.9	1.5
2001.....	42.7	1.1	0.4	0.2	0.3	0.4	0.6	0.6	0.9	1.5
2000.....	43.5	1.2	0.4	0.2	0.5	0.6	0.6	0.7	0.9	1.3
1999.....	45.6	*	0.4	0.3	0.6	0.6	0.7	0.7	0.8	1.2
All races 3/, female										
2005.....	45.5	*	0.3	0.2	0.2	*	0.4	0.4	0.9	1.3
2004.....	42.5	*	*	*	0.3	0.3	0.4	0.5	0.5	1.4
2003.....	44.4	*	*	*	0.3	0.3	0.4	0.5	0.7	1.5
2002.....	43.7	*	0.3	*	0.4	0.3	0.5	0.5	0.9	1.7
2001.....	43.7	*	*	*	0.2	0.3	0.4	0.5	0.9	1.7
2000.....	43.2	*	0.3	0.2	0.4	0.3	0.5	0.6	0.8	1.4
1999.....	43.4	*	0.3	0.2	0.3	0.5	0.4	0.8	1.0	1.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Chronic lower respiratory diseases (J40-J47)

White, both sexes

2005 4/.....	50.3	0.6	0.2	0.2	0.1	0.2	0.2	0.4	0.6	1.1
2004 4/.....	47.4	0.7	0.2	*	0.2	0.2	0.3	0.4	0.5	1.1
2003 4/.....	49.5	0.7	0.2	*	0.2	0.2	0.4	0.5	0.6	1.2
2002.....	49.2	0.8	0.2	*	0.2	0.3	0.4	0.5	0.7	1.3
2001.....	49.0	0.7	0.2	*	0.2	0.3	0.3	0.4	0.6	1.2
2000.....	49.0	0.8	0.2	0.1	0.2	0.3	0.4	0.4	0.6	1.0
1999.....	50.2	*	0.3	0.2	0.3	0.4	0.4	0.5	0.6	1.1

White, male

2005 4/.....	47.9	*	*	*	*	0.3	0.3	0.4	0.4	1.0
2004 4/.....	45.5	*	*	*	0.3	0.3	0.4	0.4	0.6	1.0
2003 4/.....	47.4	1.3	*	*	0.3	0.2	0.5	0.4	0.5	1.1
2002.....	47.8	*	0.3	*	*	0.4	0.4	0.5	0.8	1.1
2001.....	47.6	*	*	*	*	0.3	0.3	0.4	0.5	1.2
2000.....	48.3	*	*	*	0.3	0.3	0.5	0.5	0.6	0.9
1999.....	50.6	*	*	*	0.4	0.4	0.4	0.4	0.5	1.0

White, female

2005 4/.....	52.8	*	*	*	*	*	*	0.4	0.7	1.2
2004 4/.....	49.3	*	*	*	*	*	0.3	0.4	0.4	1.2
2003 4/.....	51.5	*	*	*	*	*	0.3	0.5	0.7	1.3
2002.....	50.5	*	*	*	*	*	0.4	0.4	0.7	1.5
2001.....	50.3	*	*	*	*	0.3	*	0.3	0.7	1.2
2000.....	49.7	*	*	*	*	*	0.3	0.4	0.6	1.2
1999.....	49.8	*	*	*	*	0.4	0.4	0.5	0.7	1.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Chronic lower respiratory diseases (J40-J47)										
Black, both sexes										
2005 4/.....	21.1	*	1.2	0.7	0.9	0.7	1.6	1.2	2.3	2.6
2004 4/.....	19.2	*	0.8	0.9	1.0	1.3	1.2	1.7	1.9	3.0
2003 4/.....	20.2	*	1.0	0.7	1.1	1.3	1.3	1.7	2.1	2.8
2002.....	20.7	*	1.2	0.6	1.6	1.4	1.2	1.6	2.2	3.8
2001.....	20.4	*	0.9	0.7	0.9	1.0	1.6	1.8	3.0	4.5
2000.....	20.8	*	0.8	0.7	1.4	1.4	1.3	1.9	2.4	3.5
1999.....	21.9	*	*	0.6	1.5	1.2	1.5	2.5	2.7	3.2
Black, male										
2005 4/.....	23.9	*	1.6	*	1.2	1.2	1.8	1.5	2.5	2.7
2004 4/.....	22.0	*	*	1.3	1.3	1.6	1.3	2.4	2.6	3.1
2003 4/.....	23.3	*	1.6	*	1.5	1.6	1.5	2.4	3.1	2.8
2002.....	24.1	*	1.6	*	1.7	2.2	1.5	2.2	2.3	4.5
2001.....	23.6	*	*	*	*	1.4	1.8	2.1	3.4	4.3
2000.....	24.3	*	*	*	1.7	2.0	1.5	2.4	3.1	3.7
1999.....	26.2	*	*	*	2.1	1.5	2.2	2.6	2.7	3.3
Black, female										
2005 4/.....	18.4	*	*	*	*	*	1.4	*	2.2	2.5
2004 4/.....	16.6	*	*	*	*	*	*	*	1.4	2.9
2003 4/.....	17.3	*	*	*	*	*	*	*	*	2.9
2002.....	17.7	*	*	*	1.4	*	*	*	2.1	3.2
2001.....	17.5	*	*	*	*	*	1.4	1.5	2.6	4.7
2000.....	17.6	*	*	*	*	*	*	1.4	1.7	3.3
1999.....	18.0	*	*	*	*	*	*	2.4	2.8	3.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Asthma (J45-J46)										
All races 3/, both sexes										
2005.....	1.3	*	0.2	0.2	0.3	0.2	0.4	0.5	0.6	0.7
2004.....	1.3	*	0.2	0.2	0.3	0.4	0.4	0.5	0.5	0.7
2003.....	1.4	*	0.2	0.2	0.4	0.3	0.4	0.5	0.6	0.7
2002.....	1.5	*	0.3	0.2	0.4	0.4	0.4	0.5	0.6	0.9
2001.....	1.5	*	0.2	0.2	0.3	0.3	0.4	0.5	0.7	0.9
2000.....	1.6	*	0.2	0.2	0.4	0.4	0.5	0.5	0.7	0.8
1999.....	1.7	*	0.2	0.2	0.4	0.5	0.5	0.6	0.7	0.9
All races 3/, male										
2005.....	0.9	*	0.3	0.3	0.3	0.3	0.4	0.5	0.5	0.5
2004.....	1.0	*	*	0.2	0.4	0.5	0.5	0.5	0.6	0.6
2003.....	1.0	*	0.4	0.2	0.4	0.4	0.5	0.6	0.7	0.6
2002.....	1.1	*	0.4	0.2	0.5	0.6	0.5	0.7	0.6	0.8
2001.....	1.1	*	0.3	0.2	0.3	0.3	0.4	0.5	0.6	0.8
2000.....	1.2	*	0.3	0.2	0.5	0.5	0.5	0.7	0.8	0.7
1999.....	1.2	*	*	0.2	0.6	0.5	0.6	0.6	0.6	0.8
All races 3/, female										
2005.....	1.7	*	*	*	*	*	0.3	0.4	0.7	0.8
2004.....	1.6	*	*	*	0.2	0.3	0.3	0.5	0.4	0.9
2003.....	1.8	*	*	*	0.3	0.3	0.4	0.4	0.6	0.9
2002.....	1.8	*	*	*	0.4	0.2	0.4	0.4	0.6	1.0
2001.....	1.9	*	*	*	0.2	0.3	0.4	0.4	0.8	1.0
2000.....	2.0	*	*	0.2	0.3	0.3	0.4	0.4	0.6	0.9
1999.....	2.1	*	*	*	0.3	0.4	0.4	0.6	0.8	1.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Asthma (J45-J46)										
White, both sexes										
2005 4/.....	1.1	*	*	0.1	*	0.2	0.2	0.4	0.4	0.5
2004 4/.....	1.1	*	*	*	0.2	0.2	0.3	0.3	0.3	0.6
2003 4/.....	1.2	*	*	*	0.2	0.2	0.3	0.4	0.4	0.5
2002.....	1.3	*	*	*	0.2	0.3	0.3	0.4	0.5	0.7
2001.....	1.3	*	*	*	0.2	0.2	0.2	0.3	0.4	0.5
2000.....	1.4	*	*	*	0.2	0.2	0.4	0.3	0.5	0.5
1999.....	1.5	*	*	0.1	0.2	0.4	0.3	0.4	0.4	0.7
White, male										
2005 4/.....	0.7	*	*	*	*	*	0.2	0.4	0.3	0.4
2004 4/.....	0.7	*	*	*	*	0.3	0.3	0.3	0.4	0.4
2003 4/.....	0.8	*	*	*	0.3	*	0.3	0.4	0.4	0.4
2002.....	0.9	*	*	*	*	0.3	0.3	0.4	0.5	0.5
2001.....	0.8	*	*	*	*	*	0.2	0.3	0.3	0.5
2000.....	0.9	*	*	*	0.2	0.3	0.4	0.4	0.5	0.4
1999.....	0.9	*	*	*	0.3	0.4	0.4	0.4	0.4	0.6
White, female										
2005 4/.....	1.5	*	*	*	*	*	*	0.4	0.5	0.6
2004 4/.....	1.5	*	*	*	*	*	*	0.4	0.3	0.7
2003 4/.....	1.6	*	*	*	*	*	*	0.4	0.5	0.7
2002.....	1.7	*	*	*	*	*	0.3	0.4	0.6	0.9
2001.....	1.7	*	*	*	*	*	*	*	0.6	0.6
2000.....	1.8	*	*	*	*	*	0.3	0.3	0.5	0.6
1999.....	2.0	*	*	*	*	0.4	*	0.4	0.5	0.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Asthma (J45-J46)										
Black, both sexes										
2005 4/.....	2.6	*	1.1	0.7	0.9	0.7	1.5	1.2	1.9	1.8
2004 4/.....	2.6	*	*	0.7	1.0	1.2	1.1	1.6	1.7	2.0
2003 4/.....	2.7	*	0.9	0.7	1.0	1.1	1.3	1.4	1.9	2.0
2002.....	2.9	*	0.9	*	1.5	1.3	1.2	1.6	1.6	2.9
2001.....	3.0	*	0.8	0.7	0.8	0.9	1.4	1.7	2.6	3.4
2000.....	3.2	*	*	0.7	1.4	1.3	1.2	1.8	2.0	2.7
1999.....	3.2	*	*	0.6	1.4	1.1	1.4	2.3	2.4	2.6
Black, male										
2005 4/.....	2.2	*	1.6	*	1.2	*	1.6	1.5	1.9	1.7
2004 4/.....	2.5	*	*	*	1.2	1.5	1.3	2.2	2.3	1.8
2003 4/.....	2.4	*	1.5	*	1.4	1.3	1.4	2.1	2.7	1.9
2002.....	2.8	*	*	*	1.7	2.1	1.5	2.2	1.7	3.4
2001.....	2.6	*	*	*	*	*	1.5	1.9	3.0	3.0
2000.....	2.8	*	*	*	1.7	1.7	*	2.3	2.6	2.7
1999.....	2.8	*	*	*	2.0	1.3	2.0	2.6	2.6	2.6
Black, female										
2005 4/.....	3.0	*	*	*	*	*	1.3	*	1.9	1.9
2004 4/.....	2.7	*	*	*	*	*	*	*	*	2.2
2003 4/.....	2.9	*	*	*	*	*	*	*	*	2.1
2002.....	3.0	*	*	*	1.4	*	*	*	1.4	2.3
2001.....	3.3	*	*	*	*	*	*	1.4	2.3	3.7
2000.....	3.5	*	*	*	*	*	*	*	1.4	2.6
1999.....	3.5	*	*	*	*	*	*	2.0	2.3	2.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)

All races 3/, both sexes

2005.....	9.1	7.4	0.6	0.2	0.1	0.2	0.3	0.3	0.6	0.7
2004.....	9.3	6.5	0.7	0.2	0.1	0.3	0.3	0.4	0.5	0.9
2003.....	8.8	7.0	0.5	0.1	0.2	0.3	0.4	0.4	0.6	0.9
2002.....	8.7	6.9	0.5	0.2	0.2	0.3	0.4	0.4	0.5	0.9
2001.....	9.2	7.3	0.6	0.1	0.2	0.3	0.4	0.5	0.6	1.0
2000.....	9.1	7.2	0.6	0.2	0.2	0.3	0.4	0.5	0.8	1.0
1999.....	9.0	6.6	0.6	0.2	0.2	0.3	0.3	0.5	0.8	1.0

All races 3/, male

2005.....	9.0	8.9	0.7	*	*	0.3	0.3	0.4	0.5	0.7
2004.....	9.3	6.6	0.7	*	*	0.2	0.3	0.4	0.5	0.9
2003.....	8.7	8.0	0.6	*	0.2	0.3	0.4	0.5	0.6	0.9
2002.....	8.5	7.9	0.7	0.2	*	0.3	0.4	0.3	0.5	0.9
2001.....	8.9	9.0	0.6	*	0.2	0.3	0.5	0.5	0.6	0.9
2000.....	9.0	7.9	0.6	0.2	*	0.3	0.4	0.5	0.8	1.2
1999.....	8.9	7.5	0.7	0.2	*	0.4	0.3	0.5	0.8	1.1

All races 3/, female

2005.....	9.2	5.8	0.5	*	*	*	0.3	0.3	0.6	0.8
2004.....	9.3	6.4	0.6	*	*	0.3	0.3	0.4	0.5	0.9
2003.....	8.9	5.9	0.5	*	*	0.3	0.3	0.3	0.6	0.9
2002.....	8.8	5.8	0.4	0.2	0.2	0.2	0.4	0.4	0.6	0.8
2001.....	9.4	5.5	0.6	*	0.2	0.3	0.4	0.6	0.6	1.0
2000.....	9.2	6.4	0.6	*	0.2	0.3	0.3	0.5	0.7	0.9
1999.....	9.1	5.8	0.5	*	0.2	0.2	0.3	0.5	0.8	0.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)										
White, both sexes										
2005 4/.....	9.9	6.0	0.5	0.2	0.1	0.2	0.3	0.3	0.5	0.6
2004 4/.....	10.2	4.9	0.5	*	0.1	0.3	0.3	0.3	0.4	0.7
2003 4/.....	9.6	5.4	0.4	0.1	0.2	0.2	0.3	0.4	0.6	0.7
2002.....	9.5	4.8	0.5	0.2	0.2	0.2	0.3	0.3	0.5	0.7
2001.....	10.0	5.0	0.5	0.1	0.2	0.2	0.4	0.4	0.5	0.8
2000.....	9.9	5.6	0.5	0.2	0.2	0.2	0.3	0.5	0.7	0.8
1999.....	9.7	4.9	0.5	0.2	0.2	0.3	0.3	0.4	0.7	0.8
White, male										
2005 4/.....	9.7	7.0	0.6	*	*	0.3	*	0.4	0.5	0.6
2004 4/.....	10.1	4.9	0.6	*	*	*	0.3	0.3	0.4	0.7
2003 4/.....	9.4	5.8	0.4	*	*	*	0.3	0.4	0.6	0.8
2002.....	9.2	5.4	0.6	*	*	*	0.4	0.3	0.5	0.8
2001.....	9.7	5.9	0.5	*	0.2	0.3	0.4	0.3	0.5	0.8
2000.....	9.7	6.6	0.5	*	*	*	0.4	0.4	0.7	0.9
1999.....	9.6	5.6	0.6	*	*	0.4	0.4	0.5	0.7	0.9
White, female										
2005 4/.....	10.1	4.9	0.5	*	*	*	0.3	*	0.5	0.7
2004 4/.....	10.2	4.9	0.4	*	*	0.3	0.3	0.3	0.4	0.8
2003 4/.....	9.8	5.0	0.4	*	*	0.3	*	0.3	0.5	0.7
2002.....	9.7	4.2	*	*	*	*	*	0.4	0.5	0.7
2001.....	10.2	4.0	0.6	*	*	*	0.4	0.5	0.5	0.7
2000.....	10.0	4.6	0.6	*	*	0.3	*	0.5	0.6	0.7
1999.....	9.8	4.3	0.5	*	*	*	*	0.3	0.7	0.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)										
Black, both sexes										
2005 4/.....	6.3	15.4	0.9	*	*	*	*	*	1.2	1.4
2004 4/.....	6.2	14.8	1.2	*	*	*	*	*	1.1	2.2
2003 4/.....	5.9	15.0	1.1	*	*	*	0.8	0.8	1.2	1.8
2002.....	6.0	16.6	1.1	*	*	*	0.9	*	1.0	1.8
2001.....	6.7	18.0	1.0	*	*	*	0.8	1.1	1.4	2.5
2000.....	6.5	16.7	0.8	*	*	*	0.8	0.8	1.3	2.4
1999.....	6.5	15.4	1.0	*	*	*	*	1.1	1.6	2.3
Black, male										
2005 4/.....	6.5	20.1	*	*	*	*	*	*	*	*
2004 4/.....	6.3	15.7	*	*	*	*	*	*	*	2.8
2003 4/.....	6.0	18.4	*	*	*	*	*	*	*	1.9
2002.....	6.0	19.5	*	*	*	*	*	*	*	1.9
2001.....	6.4	23.1	*	*	*	*	*	*	1.5	2.0
2000.....	6.5	16.3	*	*	*	*	*	*	*	3.0
1999.....	6.4	17.6	*	*	*	*	*	*	1.7	2.5
Black, female										
2005 4/.....	6.2	10.5	*	*	*	*	*	*	*	1.7
2004 4/.....	6.1	13.9	*	*	*	*	*	*	*	1.7
2003 4/.....	5.7	11.4	*	*	*	*	*	*	*	1.7
2002.....	6.0	13.6	*	*	*	*	*	*	*	1.6
2001.....	7.0	12.6	*	*	*	*	*	*	*	2.9
2000.....	6.4	17.2	*	*	*	*	*	*	*	1.9
1999.....	6.7	13.1	*	*	*	*	*	1.7	1.6	2.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Chronic liver disease and cirrhosis (K70,K73-K74)										
All races 3/, both sexes										
2005.....	9.3	*	*	*	*	*	*	0.4	1.2	3.4
2004.....	9.2	*	*	*	*	*	*	0.3	1.3	3.7
2003.....	9.5	*	*	*	*	*	*	0.4	1.4	4.1
2002.....	9.5	*	*	*	*	*	0.1	0.4	1.4	4.2
2001.....	9.5	*	*	*	*	*	0.1	0.4	1.5	5.0
2000.....	9.4	*	*	*	*	*	0.2	0.5	1.6	4.8
1999.....	9.4	*	*	*	*	*	*	0.4	1.6	4.8
All races 3/, male										
2005.....	12.3	*	*	*	*	*	*	0.5	1.4	4.5
2004.....	12.1	*	*	*	*	*	*	0.3	1.8	4.9
2003.....	12.5	*	*	*	*	*	*	0.6	1.9	5.4
2002.....	12.3	*	*	*	*	*	*	0.5	1.8	5.5
2001.....	12.4	*	*	*	*	*	*	0.6	2.0	6.4
2000.....	12.5	*	*	*	*	*	*	0.6	2.1	6.3
1999.....	12.5	*	*	*	*	*	*	0.4	2.1	6.3
All races 3/, female										
2005.....	6.4	*	*	*	*	*	*	0.3	0.9	2.3
2004.....	6.3	*	*	*	*	*	*	*	0.7	2.5
2003.....	6.5	*	*	*	*	*	*	0.3	0.8	2.9
2002.....	6.7	*	*	*	*	*	*	0.3	1.0	3.0
2001.....	6.7	*	*	*	*	*	*	0.3	1.0	3.5
2000.....	6.5	*	*	*	*	*	*	0.3	1.1	3.2
1999.....	6.4	*	*	*	*	*	*	0.3	1.2	3.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Chronic liver disease and cirrhosis (K70,K73-K74)										
White, both sexes										
2005 4/.....	10.0	*	*	*	*	*	*	0.4	1.1	3.6
2004 4/.....	9.9	*	*	*	*	*	*	0.3	1.3	3.8
2003 4/.....	10.2	*	*	*	*	*	*	0.4	1.4	4.2
2002.....	10.1	*	*	*	*	*	*	0.4	1.4	4.3
2001.....	10.1	*	*	*	*	*	*	0.3	1.5	5.1
2000.....	10.0	*	*	*	*	*	*	0.4	1.5	4.7
1999.....	9.9	*	*	*	*	*	*	0.3	1.6	4.8
White, male										
2005 4/.....	13.2	*	*	*	*	*	*	0.5	1.4	4.8
2004 4/.....	13.0	*	*	*	*	*	*	0.3	1.9	5.0
2003 4/.....	13.4	*	*	*	*	*	*	0.5	2.0	5.6
2002.....	13.2	*	*	*	*	*	*	0.4	1.9	5.6
2001.....	13.1	*	*	*	*	*	*	0.5	2.1	6.5
2000.....	13.2	*	*	*	*	*	*	0.6	2.1	6.3
1999.....	13.1	*	*	*	*	*	*	0.4	2.1	6.4
White, female										
2005 4/.....	6.9	*	*	*	*	*	*	*	0.8	2.4
2004 4/.....	6.8	*	*	*	*	*	*	*	0.7	2.6
2003 4/.....	7.0	*	*	*	*	*	*	*	0.8	2.9
2002.....	7.2	*	*	*	*	*	*	0.3	0.9	3.0
2001.....	7.1	*	*	*	*	*	*	*	1.0	3.7
2000.....	6.8	*	*	*	*	*	*	*	1.0	3.1
1999.....	6.8	*	*	*	*	*	*	*	1.1	3.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Alcoholic liver disease (K70)										
All races 3/, both sexes										
2005.....	4.4	*	*	*	*	*	*	0.3	0.8	2.4
2004.....	4.3	*	*	*	*	*	*	0.2	0.9	2.5
2003.....	4.3	*	*	*	*	*	*	0.3	0.9	2.9
2002.....	4.2	*	*	*	*	*	*	0.2	0.9	2.8
2001.....	4.3	*	*	*	*	*	*	0.2	1.1	3.3
2000.....	4.3	*	*	*	*	*	*	0.3	1.1	3.3
1999.....	4.3	*	*	*	*	*	*	0.2	1.1	3.3
All races 3/, male										
2005.....	6.5	*	*	*	*	*	*	0.4	1.0	3.2
2004.....	6.3	*	*	*	*	*	*	0.3	1.3	3.3
2003.....	6.4	*	*	*	*	*	*	0.4	1.4	3.7
2002.....	6.2	*	*	*	*	*	*	0.3	1.2	3.6
2001.....	6.4	*	*	*	*	*	*	0.3	1.4	4.4
2000.....	6.5	*	*	*	*	*	*	0.5	1.5	4.3
1999.....	6.5	*	*	*	*	*	*	0.2	1.4	4.3
All races 3/, female										
2005.....	2.3	*	*	*	*	*	*	*	0.6	1.6
2004.....	2.3	*	*	*	*	*	*	*	0.5	1.8
2003.....	2.2	*	*	*	*	*	*	*	0.5	2.0
2002.....	2.3	*	*	*	*	*	*	*	0.7	2.0
2001.....	2.3	*	*	*	*	*	*	*	0.7	2.3
2000.....	2.2	*	*	*	*	*	*	*	0.6	2.2
1999.....	2.1	*	*	*	*	*	*	*	0.8	2.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Alcoholic liver disease (K70)										
White, both sexes										
2005 4/.....	4.7	*	*	*	*	*	*	0.3	0.8	2.6
2004 4/.....	4.6	*	*	*	*	*	*	0.2	1.0	2.6
2003 4/.....	4.5	*	*	*	*	*	*	0.2	1.0	2.9
2002.....	4.4	*	*	*	*	*	*	0.2	0.9	2.8
2001.....	4.5	*	*	*	*	*	*	0.2	1.1	3.4
2000.....	4.5	*	*	*	*	*	*	0.3	1.1	3.2
1999.....	4.4	*	*	*	*	*	*	0.2	1.0	3.3
White, male										
2005 4/.....	6.9	*	*	*	*	*	*	0.4	1.0	3.4
2004 4/.....	6.8	*	*	*	*	*	*	0.3	1.5	3.3
2003 4/.....	6.8	*	*	*	*	*	*	0.4	1.4	3.9
2002.....	6.6	*	*	*	*	*	*	0.3	1.3	3.7
2001.....	6.7	*	*	*	*	*	*	0.3	1.5	4.5
2000.....	6.8	*	*	*	*	*	*	0.5	1.6	4.3
1999.....	6.7	*	*	*	*	*	*	0.3	1.4	4.3
White, female										
2005 4/.....	2.4	*	*	*	*	*	*	*	0.5	1.7
2004 4/.....	2.4	*	*	*	*	*	*	*	0.5	1.9
2003 4/.....	2.3	*	*	*	*	*	*	*	0.5	1.9
2002.....	2.3	*	*	*	*	*	*	*	0.6	2.0
2001.....	2.3	*	*	*	*	*	*	*	0.7	2.3
2000.....	2.1	*	*	*	*	*	*	*	0.5	2.2
1999.....	2.1	*	*	*	*	*	*	*	0.7	2.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other chronic liver disease and cirrhosis (K73-K74)										
All races 3/, both sexes										
2005.....	4.9	*	*	*	*	*	*	*	0.4	1.0
2004.....	4.9	*	*	*	*	*	*	*	0.3	1.2
2003.....	5.2	*	*	*	*	*	*	0.2	0.4	1.3
2002.....	5.2	*	*	*	*	*	*	0.2	0.5	1.5
2001.....	5.2	*	*	*	*	*	*	0.2	0.4	1.6
2000.....	5.1	*	*	*	*	*	0.1	0.2	0.5	1.5
1999.....	5.1	*	*	*	*	*	*	0.2	0.6	1.5
All races 3/, male										
2005.....	5.8	*	*	*	*	*	*	*	0.4	1.3
2004.....	5.8	*	*	*	*	*	*	*	0.5	1.7
2003.....	6.2	*	*	*	*	*	*	*	0.5	1.6
2002.....	6.1	*	*	*	*	*	*	*	0.6	1.9
2001.....	6.1	*	*	*	*	*	*	0.2	0.6	2.0
2000.....	6.0	*	*	*	*	*	*	*	0.6	2.0
1999.....	6.0	*	*	*	*	*	*	*	0.7	2.0
All races 3/, female										
2005.....	4.0	*	*	*	*	*	*	*	0.4	0.7
2004.....	4.1	*	*	*	*	*	*	*	0.2	0.7
2003.....	4.3	*	*	*	*	*	*	*	0.3	0.9
2002.....	4.5	*	*	*	*	*	*	0.2	0.4	1.0
2001.....	4.4	*	*	*	*	*	*	*	0.3	1.3
2000.....	4.3	*	*	*	*	*	*	*	0.4	1.0
1999.....	4.3	*	*	*	*	*	*	*	0.4	1.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Other chronic liver disease and cirrhosis (K73-K74)

White, both sexes

2005 4/.....	5.3	*	*	*	*	*	*	*	0.3	1.1
2004 4/.....	5.3	*	*	*	*	*	*	*	0.3	1.2
2003 4/.....	5.7	*	*	*	*	*	*	*	0.4	1.3
2002.....	5.7	*	*	*	*	*	*	0.2	0.4	1.5
2001.....	5.6	*	*	*	*	*	*	*	0.5	1.7
2000.....	5.5	*	*	*	*	*	*	*	0.5	1.5
1999.....	5.5	*	*	*	*	*	*	0.1	0.6	1.5

White, male

2005 4/.....	6.2	*	*	*	*	*	*	*	0.4	1.4
2004 4/.....	6.2	*	*	*	*	*	*	*	0.5	1.7
2003 4/.....	6.6	*	*	*	*	*	*	*	0.6	1.7
2002.....	6.5	*	*	*	*	*	*	*	0.6	2.0
2001.....	6.4	*	*	*	*	*	*	*	0.6	2.0
2000.....	6.4	*	*	*	*	*	*	*	0.5	2.1
1999.....	6.4	*	*	*	*	*	*	*	0.7	2.0

White, female

2005 4/.....	4.4	*	*	*	*	*	*	*	0.3	0.7
2004 4/.....	4.5	*	*	*	*	*	*	*	*	0.7
2003 4/.....	4.7	*	*	*	*	*	*	*	0.3	1.0
2002.....	4.9	*	*	*	*	*	*	*	0.3	1.0
2001.....	4.8	*	*	*	*	*	*	*	0.4	1.4
2000.....	4.7	*	*	*	*	*	*	*	0.4	0.9
1999.....	4.6	*	*	*	*	*	*	*	0.4	1.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)										
All races 3/, both sexes										
2005.....	14.8	3.9	*	*	*	0.2	0.3	0.6	0.9	1.2
2004.....	14.5	4.3	*	*	*	0.1	0.3	0.5	0.7	1.4
2003.....	14.6	4.5	*	*	*	0.1	0.3	0.5	0.9	1.3
2002.....	14.2	4.3	*	*	*	0.1	0.3	0.6	0.8	1.3
2001.....	13.9	3.3	*	*	*	0.1	0.4	0.4	0.7	1.3
2000.....	13.2	4.3	*	*	*	0.2	0.3	0.4	0.8	1.2
1999.....	12.7	4.4	*	*	*	0.1	0.3	0.5	0.8	1.2
All races 3/, male										
2005.....	14.6	4.2	*	*	*	*	0.3	0.5	1.0	1.3
2004.....	14.1	5.3	*	*	*	*	0.3	0.5	0.6	1.5
2003.....	14.3	5.2	*	*	*	*	0.4	0.5	0.9	1.4
2002.....	13.9	5.3	*	*	*	*	0.3	0.5	0.8	1.4
2001.....	13.5	3.7	*	*	*	*	0.3	0.5	0.6	1.4
2000.....	12.9	5.3	*	*	*	*	0.3	0.4	0.8	1.3
1999.....	12.4	4.6	*	*	*	*	0.2	0.4	0.8	1.3
All races 3/, female										
2005.....	15.0	3.5	*	*	*	0.2	0.3	0.6	0.8	1.1
2004.....	14.8	3.2	*	*	*	*	0.2	0.5	0.8	1.3
2003.....	14.9	3.8	*	*	*	*	0.3	0.5	0.8	1.1
2002.....	14.5	3.2	*	*	*	*	0.3	0.7	0.7	1.1
2001.....	14.2	2.9	*	*	*	*	0.4	0.4	0.7	1.3
2000.....	13.6	3.2	*	*	*	*	0.3	0.4	0.8	1.1
1999.....	13.0	4.1	*	*	*	*	0.3	0.6	0.7	1.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)

White, both sexes

2005 4/.....	14.5	3.0	*	*	*	0.1	0.2	0.4	0.6	0.9
2004 4/.....	14.1	3.1	*	*	*	*	0.2	0.3	0.5	1.0
2003 4/.....	14.3	3.8	*	*	*	*	0.2	0.4	0.6	0.8
2002.....	13.9	3.6	*	*	*	*	0.2	0.4	0.6	0.8
2001.....	13.5	2.4	*	*	*	*	0.3	0.3	0.5	0.9
2000.....	12.9	2.9	*	*	*	0.1	0.3	0.3	0.5	0.8
1999.....	12.3	3.4	*	*	*	*	0.2	0.3	0.5	0.8

White, male

2005 4/.....	14.4	3.1	*	*	*	*	0.3	0.3	0.7	1.0
2004 4/.....	13.9	3.8	*	*	*	*	0.2	0.4	0.3	1.0
2003 4/.....	14.0	4.5	*	*	*	*	0.3	0.4	0.6	0.8
2002.....	13.7	4.4	*	*	*	*	*	0.3	0.6	0.9
2001.....	13.3	2.8	*	*	*	*	*	0.3	0.4	1.1
2000.....	12.7	3.8	*	*	*	*	0.3	0.4	0.5	0.9
1999.....	12.1	4.1	*	*	*	*	*	0.3	0.5	0.9

White, female

2005 4/.....	14.6	2.9	*	*	*	*	*	0.4	0.5	0.8
2004 4/.....	14.4	2.3	*	*	*	*	*	0.3	0.6	1.0
2003 4/.....	14.5	3.0	*	*	*	*	*	0.4	0.6	0.8
2002.....	14.1	2.7	*	*	*	*	*	0.5	0.5	0.8
2001.....	13.7	2.1	*	*	*	*	0.4	0.3	0.6	0.7
2000.....	13.0	2.0	*	*	*	*	0.3	0.3	0.5	0.7
1999.....	12.4	2.8	*	*	*	*	*	0.4	0.5	0.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)										
Black, both sexes										
2005 4/.....	20.7	8.7	*	*	*	*	*	1.6	2.7	3.5
2004 4/.....	20.3	10.1	*	*	*	*	0.8	1.3	2.4	4.1
2003 4/.....	20.6	9.1	*	*	*	*	0.7	1.4	2.9	4.4
2002.....	19.8	8.2	*	*	*	*	0.9	1.9	2.1	4.0
2001.....	19.6	8.1	*	*	*	*	0.7	1.3	2.2	4.0
2000.....	18.9	11.5	*	*	*	*	*	1.1	2.6	4.4
1999.....	18.6	9.6	*	*	*	*	0.8	1.5	2.3	4.0
Black, male										
2005 4/.....	19.5	10.6	*	*	*	*	*	1.7	2.9	3.7
2004 4/.....	19.2	12.2	*	*	*	*	*	*	2.8	5.2
2003 4/.....	20.0	9.5	*	*	*	*	*	*	3.5	5.4
2002.....	19.1	10.2	*	*	*	*	*	1.6	2.2	4.4
2001.....	18.0	8.7	*	*	*	*	*	1.7	2.6	3.4
2000.....	17.7	13.4	*	*	*	*	*	*	2.8	5.1
1999.....	17.5	8.1	*	*	*	*	*	*	2.3	4.8
Black, female										
2005 4/.....	21.7	6.8	*	*	*	*	*	1.6	2.6	3.4
2004 4/.....	21.3	8.0	*	*	*	*	*	1.5	2.1	3.1
2003 4/.....	21.1	8.7	*	*	*	*	*	*	2.5	3.5
2002.....	20.5	6.1	*	*	*	*	*	2.1	2.1	3.6
2001.....	21.0	7.6	*	*	*	*	*	*	1.8	4.5
2000.....	20.0	9.6	*	*	*	*	*	*	2.3	3.8
1999.....	19.5	11.1	*	*	*	*	*	1.6	2.3	3.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Renal failure (N17-N19)										
All races 3/, both sexes										
2005.....	14.5	3.7	*	*	*	0.1	0.2	0.5	0.8	1.2
2004.....	14.2	4.0	*	*	*	0.1	0.3	0.4	0.7	1.3
2003.....	14.4	4.2	*	*	*	0.1	0.3	0.5	0.8	1.2
2002.....	13.9	4.1	*	*	*	*	0.3	0.5	0.7	1.2
2001.....	13.6	3.1	*	*	*	*	0.3	0.4	0.6	1.2
2000.....	13.0	4.0	*	*	*	0.1	0.3	0.4	0.7	1.2
1999.....	12.4	4.3	*	*	*	0.1	0.2	0.4	0.7	1.1
All races 3/, male										
2005.....	14.2	4.1	*	*	*	*	0.3	0.5	0.9	1.2
2004.....	13.8	5.1	*	*	*	*	0.3	0.4	0.6	1.4
2003.....	14.1	4.8	*	*	*	*	0.3	0.5	0.9	1.3
2002.....	13.7	5.1	*	*	*	*	0.3	0.4	0.7	1.3
2001.....	13.2	3.5	*	*	*	*	0.3	0.4	0.6	1.3
2000.....	12.6	5.1	*	*	*	*	0.2	0.4	0.7	1.3
1999.....	12.1	4.5	*	*	*	*	*	0.3	0.7	1.3
All races 3/, female										
2005.....	14.7	3.2	*	*	*	*	0.2	0.5	0.7	1.1
2004.....	14.6	3.0	*	*	*	*	0.2	0.4	0.8	1.2
2003.....	14.6	3.7	*	*	*	*	0.2	0.5	0.8	1.1
2002.....	14.2	3.0	*	*	*	*	0.3	0.6	0.7	1.1
2001.....	14.0	2.7	*	*	*	*	0.4	0.4	0.7	1.2
2000.....	13.3	2.8	*	*	*	*	0.3	0.3	0.7	1.0
1999.....	12.7	4.1	*	*	*	*	0.3	0.5	0.7	0.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Renal failure (N17-N19)

White, both sexes

2005 4/.....	14.2	2.8	*	*	*	*	0.2	0.4	0.5	0.8
2004 4/.....	13.9	2.9	*	*	*	*	0.2	0.3	0.4	0.9
2003 4/.....	14.0	3.5	*	*	*	*	0.2	0.4	0.5	0.8
2002.....	13.6	3.4	*	*	*	*	0.2	0.3	0.5	0.8
2001.....	13.2	2.2	*	*	*	*	0.3	0.3	0.4	0.8
2000.....	12.6	2.7	*	*	*	*	0.2	0.3	0.5	0.7
1999.....	12.0	3.3	*	*	*	*	0.2	0.3	0.5	0.7

White, male

2005 4/.....	14.1	3.1	*	*	*	*	0.2	0.3	0.6	0.9
2004 4/.....	13.6	3.6	*	*	*	*	*	0.3	0.3	0.9
2003 4/.....	13.8	4.1	*	*	*	*	*	0.4	0.5	0.8
2002.....	13.4	4.2	*	*	*	*	*	*	0.5	0.9
2001.....	13.0	2.6	*	*	*	*	*	0.3	0.4	1.0
2000.....	12.4	3.6	*	*	*	*	0.3	0.3	0.5	0.8
1999.....	11.8	3.9	*	*	*	*	*	*	0.5	0.8

White, female

2005 4/.....	14.2	2.6	*	*	*	*	*	0.4	0.4	0.7
2004 4/.....	14.1	2.1	*	*	*	*	*	0.3	0.5	1.0
2003 4/.....	14.2	2.8	*	*	*	*	*	0.4	0.5	0.7
2002.....	13.8	2.6	*	*	*	*	*	0.4	0.5	0.7
2001.....	13.4	1.8	*	*	*	*	0.3	0.3	0.5	0.7
2000.....	12.8	1.7	*	*	*	*	*	*	0.5	0.7
1999.....	12.2	2.8	*	*	*	*	*	0.3	0.5	0.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Renal failure (N17-N19)										
Black, both sexes										
2005 4/.....	20.2	8.4	*	*	*	*	*	1.4	2.6	3.5
2004 4/.....	20.0	9.8	*	*	*	*	0.7	1.2	2.3	4.0
2003 4/.....	20.3	8.9	*	*	*	*	0.6	1.3	2.8	4.4
2002.....	19.5	7.7	*	*	*	*	0.8	1.7	2.0	3.9
2001.....	19.3	8.1	*	*	*	*	0.7	1.2	2.1	3.9
2000.....	18.5	11.1	*	*	*	*	*	1.0	2.5	4.1
1999.....	18.2	9.6	*	*	*	*	*	1.4	2.2	3.8
Black, male										
2005 4/.....	19.1	10.6	*	*	*	*	*	1.4	2.7	3.5
2004 4/.....	18.9	11.9	*	*	*	*	*	*	2.7	5.1
2003 4/.....	19.7	9.2	*	*	*	*	*	*	3.4	5.4
2002.....	18.8	9.6	*	*	*	*	*	*	2.0	4.4
2001.....	17.7	8.7	*	*	*	*	*	1.6	2.6	3.3
2000.....	17.3	13.4	*	*	*	*	*	*	2.7	4.9
1999.....	17.1	8.1	*	*	*	*	*	*	2.2	4.5
Black, female										
2005 4/.....	21.2	6.2	*	*	*	*	*	1.4	2.5	3.4
2004 4/.....	21.0	7.7	*	*	*	*	*	*	1.9	3.0
2003 4/.....	20.9	8.7	*	*	*	*	*	*	2.3	3.5
2002.....	20.2	*	*	*	*	*	*	1.9	2.0	3.5
2001.....	20.7	7.6	*	*	*	*	*	*	1.7	4.4
2000.....	19.6	8.6	*	*	*	*	*	*	2.3	3.4
1999.....	19.1	11.1	*	*	*	*	*	1.6	2.2	3.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Pregnancy, childbirth and the puerperium (000-099)

All races 3/, both sexes

2005.....	0.3	*	0.2	0.7	0.8	0.8	0.6
2004.....	0.2	*	0.2	0.6	0.7	0.7	0.5
2003.....	0.2	*	0.1	0.4	0.6	0.6	0.5
2002.....	0.1	*	0.1	0.3	0.4	0.4	0.4
2001.....	0.1	*	0.2	0.4	0.5	0.5	0.3
2000.....	0.1	*	*	0.4	0.4	0.5	0.4
1999.....	0.1	*	0.2	0.3	0.5	0.4	0.4

All races 3/, male

2005.....
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

All races 3/, female

2005.....	0.5	*	0.4	1.4	1.6	1.6	1.3
2004.....	0.5	*	0.4	1.3	1.5	1.4	0.9
2003.....	0.4	*	0.3	0.9	1.2	1.2	1.0
2002.....	0.3	*	0.3	0.6	0.9	0.9	0.7
2001.....	0.3	*	0.4	0.8	1.0	0.9	0.7
2000.....	0.3	*	*	0.8	0.9	0.9	0.8
1999.....	0.3	*	0.3	0.7	0.9	0.9	0.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Pregnancy, childbirth and the puerperium (000-099)

White, both sexes

2005 4/.....	0.2	*	0.1	0.5	0.5	0.6	0.5
2004 4/.....	0.2	*	0.1	0.4	0.5	0.6	0.4
2003 4/.....	0.1	*	*	0.3	0.5	0.4	0.3
2002.....	0.1	*	*	0.2	0.3	0.3	0.2
2001.....	0.1	*	0.1	0.3	0.3	0.4	0.2
2000.....	0.1	*	*	0.3	0.3	0.3	0.3
1999.....	0.1	*	*	0.2	0.3	0.3	0.3

White, male

2005 4/.....
2004 4/.....
2003 4/.....
2002.....
2001.....
2000.....
1999.....

White, female

2005 4/.....	0.4	*	0.3	1.1	1.0	1.3	1.0
2004 4/.....	0.3	*	0.3	0.8	1.0	1.2	0.7
2003 4/.....	0.3	*	*	0.6	1.0	0.7	0.7
2002.....	0.2	*	*	0.4	0.6	0.7	0.5
2001.....	0.2	*	0.3	0.6	0.7	0.8	0.4
2000.....	0.2	*	*	0.6	0.7	0.7	0.6
1999.....	0.2	*	*	0.4	0.6	0.6	0.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Pregnancy, childbirth and the puerperium (000-099)

Black, both sexes

2005 4/.....	0.7	*	0.6	1.5	2.3	1.9	1.5
2004 4/.....	0.7	*	*	2.0	2.2	1.7	1.1
2003 4/.....	0.5	*	*	1.0	1.2	1.8	1.4
2002.....	0.4	*	*	1.0	1.4	1.0	1.2
2001.....	0.4	*	*	1.1	1.4	1.1	1.0
2000.....	0.4	*	*	1.1	1.1	1.2	0.9
1999.....	0.4	*	*	1.1	1.4	1.2	0.9

Black, male

2005 4/.....
2004 4/.....
2003 4/.....
2002.....
2001.....
2000.....
1999.....

Black, female

2005 4/.....	1.3	*	1.2	3.1	4.5	3.6	2.8
2004 4/.....	1.2	*	*	4.0	4.3	3.2	2.0
2003 4/.....	1.0	*	*	2.1	2.3	3.5	2.7
2002.....	0.8	*	*	2.0	2.8	1.9	2.3
2001.....	0.8	*	*	2.1	2.6	2.1	1.9
2000.....	0.7	*	*	2.2	2.2	2.3	1.7
1999.....	0.9	*	*	2.1	2.7	2.3	1.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Pregnancy with abortive outcome (000-007)

All races 3/, both sexes

2005.....	0.0	*	*	*	*	*	*
2004.....	0.0	*	*	*	*	*	*
2003.....	0.0	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	*

All races 3/, male

2005.....
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

All races 3/, female

2005.....	0.0	*	*	*	*	*	*
2004.....	0.0	*	*	*	*	*	*
2003.....	0.0	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Pregnancy with abortive outcome (000-007)

White, both sexes

2005 4/.....	*	*	*	*	*	*	*
2004 4/.....	*	*	*	*	*	*	*
2003 4/.....	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

White, male

2005 4/.....
2004 4/.....
2003 4/.....
2002.....
2001.....
2000.....
1999.....

White, female

2005 4/.....	*	*	*	*	*	*	*
2004 4/.....	*	*	*	*	*	*	*
2003 4/.....	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Pregnancy with abortive outcome (000-007)

Black, both sexes

2005 4/.....	*	*	*	*	*	*	*
2004 4/.....	*	*	*	*	*	*	*
2003 4/.....	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

Black, male

2005 4/.....
2004 4/.....
2003 4/.....
2002.....
2001.....
2000.....
1999.....

Black, female

2005 4/.....	*	*	*	*	*	*	*
2004 4/.....	*	*	*	*	*	*	*
2003 4/.....	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Other complications of pregnancy, childbirth and the puerperium (010-099)

All races 3/, both sexes

2005.....	0.2	*	0.2	0.7	0.7	0.8	0.6
2004.....	0.2	*	0.2	0.6	0.7	0.7	0.4
2003.....	0.2	*	0.1	0.4	0.5	0.5	0.5
2002.....	0.1	*	0.1	0.3	0.4	0.4	0.3
2001.....	0.1	*	0.2	0.3	0.5	0.4	0.3
2000.....	0.1	*	*	0.4	0.4	0.4	0.3
1999.....	0.1	*	0.1	0.3	0.4	0.4	0.4

All races 3/, male

2005.....
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

All races 3/, female

2005.....	0.5	*	0.4	1.4	1.5	1.5	1.2
2004.....	0.4	*	0.4	1.2	1.4	1.4	0.9
2003.....	0.3	*	0.2	0.8	1.1	1.1	0.9
2002.....	0.2	*	0.3	0.6	0.8	0.8	0.7
2001.....	0.3	*	0.3	0.7	0.9	0.9	0.6
2000.....	0.3	*	*	0.8	0.8	0.9	0.7
1999.....	0.3	*	0.3	0.6	0.9	0.8	0.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other complications of pregnancy, childbirth and the puerperium (010-099)										
White, both sexes										
2005 4/.....	0.2	*	*	0.5	0.5	0.6	0.4
2004 4/.....	0.2	*	0.1	0.4	0.5	0.6	0.4
2003 4/.....	0.1	*	*	0.3	0.4	0.3	0.3
2002.....	0.1	*	*	0.2	0.3	0.3	0.2
2001.....	0.1	*	*	0.2	0.3	0.4	0.2
2000.....	0.1	*	*	0.3	0.3	0.3	0.3
1999.....	0.1	*	*	0.2	0.3	0.3	0.3
White, male										
2005 4/.....
2004 4/.....
2003 4/.....
2002.....
2001.....
2000.....
1999.....
White, female										
2005 4/.....	0.4	*	*	1.1	1.0	1.3	0.9
2004 4/.....	0.3	*	0.3	0.7	1.0	1.1	0.7
2003 4/.....	0.3	*	*	0.6	0.9	0.7	0.6
2002.....	0.2	*	*	0.4	0.5	0.6	0.5
2001.....	0.2	*	*	0.5	0.6	0.7	0.3
2000.....	0.2	*	*	0.6	0.6	0.7	0.5
1999.....	0.2	*	*	0.4	0.6	0.6	0.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Other complications of pregnancy, childbirth and the puerperium (010-099)

Black, both sexes

2005 4/.....	0.7	*	*	1.5	2.1	1.8	1.3
2004 4/.....	0.6	*	*	1.7	2.1	1.6	0.9
2003 4/.....	0.5	*	*	0.9	1.2	1.6	1.3
2002.....	0.4	*	*	0.9	1.4	0.9	1.0
2001.....	0.4	*	*	1.0	1.3	1.0	0.9
2000.....	0.3	*	*	1.0	1.0	1.1	0.7
1999.....	0.4	*	*	1.0	1.3	1.0	0.8

Black, male

2005 4/.....
2004 4/.....
2003 4/.....
2002.....
2001.....
2000.....
1999.....

Black, female

2005 4/.....	1.2	*	*	3.1	4.1	3.3	2.5
2004 4/.....	1.1	*	*	3.5	4.1	3.0	1.8
2003 4/.....	0.9	*	*	1.9	2.3	3.1	2.5
2002.....	0.7	*	*	1.8	2.6	1.8	2.0
2001.....	0.7	*	*	1.9	2.5	1.8	1.7
2000.....	0.6	*	*	1.9	1.8	2.1	1.4
1999.....	0.8	*	*	1.9	2.4	2.0	1.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Certain conditions originating in the perinatal period (P00-P96)

All races 3/, both sexes

2005.....	4.9	351.2	0.4	*	*	*	*	*	*	*
2004.....	4.8	345.3	0.4	0.1	*	*	*	*	*	*
2003.....	4.9	356.0	0.5	*	*	*	*	*	*	*
2002.....	4.9	349.7	0.4	*	*	*	*	*	*	*
2001.....	4.9	340.5	0.5	0.1	*	*	*	*	*	*
2000.....	5.0	365.7	0.5	0.1	*	*	*	*	*	*
1999.....	5.1	371.0	0.6	0.1	*	*	*	*	*	*

All races 3/, male

2005.....	5.6	386.1	0.4	*	*	*	*	*	*	*
2004.....	5.6	384.2	0.5	*	*	*	*	*	*	*
2003.....	5.7	397.4	0.5	*	*	*	*	*	*	*
2002.....	5.7	385.7	0.5	*	*	*	*	*	*	*
2001.....	5.6	376.6	0.6	*	*	*	*	*	*	*
2000.....	5.8	403.9	0.7	*	*	*	*	*	*	*
1999.....	5.9	408.6	0.7	*	*	*	*	*	*	*

All races 3/, female

2005.....	4.2	314.6	0.3	*	*	*	*	*	*	*
2004.....	4.1	304.6	0.3	*	*	*	*	*	*	*
2003.....	4.2	312.8	0.5	*	*	*	*	*	*	*
2002.....	4.2	312.0	0.3	*	*	*	*	*	*	*
2001.....	4.2	302.6	0.4	*	*	*	*	*	*	*
2000.....	4.3	325.6	0.4	*	*	*	*	*	*	*
1999.....	4.4	331.7	0.5	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Certain conditions originating in the perinatal period (P00-P96)

White, both sexes

2005 4/.....	3.7	275.7	0.3	*	*	*	*	*	*	*
2004 4/.....	3.7	272.1	0.3	*	*	*	*	*	*	*
2003 4/.....	3.8	285.1	0.4	*	*	*	*	*	*	*
2002.....	3.7	277.2	0.3	*	*	*	*	*	*	*
2001.....	3.7	268.5	0.4	*	*	*	*	*	*	*
2000.....	3.7	283.7	0.4	*	*	*	*	*	*	*
1999.....	3.8	285.9	0.5	*	*	*	*	*	*	*

White, male

2005 4/.....	4.2	303.8	0.3	*	*	*	*	*	*	*
2004 4/.....	4.3	305.9	0.3	*	*	*	*	*	*	*
2003 4/.....	4.4	319.1	0.4	*	*	*	*	*	*	*
2002.....	4.3	309.4	*	*	*	*	*	*	*	*
2001.....	4.2	297.1	0.5	*	*	*	*	*	*	*
2000.....	4.3	312.4	0.5	*	*	*	*	*	*	*
1999.....	4.3	315.0	0.5	*	*	*	*	*	*	*

White, female

2005 4/.....	3.2	246.3	*	*	*	*	*	*	*	*
2004 4/.....	3.1	236.8	*	*	*	*	*	*	*	*
2003 4/.....	3.2	249.5	0.3	*	*	*	*	*	*	*
2002.....	3.2	243.5	*	*	*	*	*	*	*	*
2001.....	3.1	238.6	*	*	*	*	*	*	*	*
2000.....	3.2	253.5	*	*	*	*	*	*	*	*
1999.....	3.2	255.3	0.4	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)

All races 3/, both sexes

2005.....	3.5	135.2	3.2	1.0	1.0	1.2	1.2	1.2	1.0	1.1
2004.....	3.6	137.9	3.6	1.0	0.9	1.2	1.1	1.1	1.0	1.0
2003.....	3.6	140.4	3.4	0.9	1.0	1.1	1.1	1.0	1.1	1.1
2002.....	3.7	139.4	3.4	1.0	1.0	1.2	1.2	1.1	1.3	1.2
2001.....	3.7	136.7	3.6	0.9	0.9	1.3	1.3	1.2	1.1	1.2
2000.....	3.8	150.9	3.2	1.0	1.0	1.1	1.1	1.1	1.3	1.2
1999.....	3.7	144.2	3.6	1.0	1.1	1.1	1.1	1.1	1.2	1.2

All races 3/, male

2005.....	3.7	141.1	3.2	0.9	1.0	1.6	1.4	1.3	1.1	1.2
2004.....	3.8	139.0	3.9	1.1	0.9	1.4	1.3	1.2	1.1	1.2
2003.....	3.8	147.6	3.4	0.9	1.0	1.4	1.2	1.2	1.1	1.2
2002.....	3.9	142.6	3.5	1.0	1.1	1.4	1.3	1.4	1.4	1.2
2001.....	3.9	143.1	3.6	1.0	1.0	1.5	1.4	1.3	1.2	1.2
2000.....	4.0	155.4	3.3	1.0	1.0	1.3	1.4	1.2	1.5	1.4
1999.....	3.9	147.9	4.0	1.0	1.1	1.3	1.3	1.1	1.3	1.3

All races 3/, female

2005.....	3.3	129.0	3.3	1.2	0.9	0.8	1.0	1.0	0.9	1.0
2004.....	3.4	136.8	3.2	1.0	0.8	1.1	0.9	0.9	1.0	0.8
2003.....	3.4	132.8	3.5	0.9	1.0	0.8	1.0	0.8	1.1	1.1
2002.....	3.5	136.1	3.3	1.0	1.0	1.0	1.1	0.9	1.1	1.2
2001.....	3.4	129.9	3.7	0.8	0.8	1.0	1.1	1.1	1.0	1.1
2000.....	3.5	146.2	3.1	0.9	1.0	0.9	0.9	0.9	1.1	1.1
1999.....	3.5	140.3	3.2	1.0	1.1	0.9	1.0	1.1	1.2	1.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)

White, both sexes

2005 4/.....	3.4	131.0	3.1	0.9	0.9	1.1	1.2	1.1	1.0	1.1
2004 4/.....	3.5	134.5	3.2	1.0	0.9	1.1	1.0	1.1	1.1	1.0
2003 4/.....	3.5	137.6	3.2	0.9	1.0	1.1	1.0	1.1	1.1	1.2
2002.....	3.6	138.3	3.3	1.0	1.0	1.2	1.2	1.1	1.3	1.2
2001.....	3.6	135.0	3.2	0.9	0.9	1.2	1.2	1.2	1.2	1.1
2000.....	3.6	147.1	2.9	0.9	0.9	1.0	1.2	1.1	1.4	1.2
1999.....	3.6	141.3	3.2	0.9	1.0	1.1	1.1	1.1	1.3	1.2

White, male

2005 4/.....	3.6	136.9	3.1	0.8	1.0	1.5	1.4	1.2	1.1	1.2
2004 4/.....	3.6	134.3	3.5	1.1	0.9	1.2	1.1	1.3	1.1	1.1
2003 4/.....	3.7	143.2	3.1	0.8	0.9	1.3	1.1	1.2	1.1	1.2
2002.....	3.8	143.3	3.4	0.9	1.0	1.3	1.3	1.4	1.5	1.2
2001.....	3.8	143.0	3.2	1.0	1.1	1.4	1.3	1.4	1.3	1.2
2000.....	3.9	151.9	2.9	0.9	0.9	1.1	1.4	1.2	1.6	1.4
1999.....	3.8	146.5	3.5	0.8	1.0	1.3	1.3	1.2	1.3	1.4

White, female

2005 4/.....	3.2	124.8	3.0	1.0	0.8	0.7	0.9	1.0	0.9	1.1
2004 4/.....	3.4	134.7	2.9	0.8	0.8	1.0	0.9	1.0	1.1	0.8
2003 4/.....	3.4	131.8	3.2	0.9	1.0	0.8	1.0	0.9	1.1	1.1
2002.....	3.5	133.1	3.1	1.1	1.0	1.0	1.1	0.9	1.1	1.2
2001.....	3.4	126.7	3.3	0.8	0.7	0.9	1.1	1.0	1.1	1.0
2000.....	3.5	142.0	2.9	1.0	0.9	0.9	0.9	0.9	1.1	1.0
1999.....	3.5	135.7	2.9	1.0	1.0	0.8	0.9	1.1	1.2	1.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)

Black, both sexes

2005 4/.....	4.4	162.9	4.0	1.3	1.3	1.6	1.4	1.6	1.3	1.3
2004 4/.....	4.5	162.9	5.1	1.5	1.0	1.8	1.7	1.1	1.1	1.2
2003 4/.....	4.5	157.5	4.7	1.1	1.1	1.2	1.7	1.2	1.5	1.2
2002.....	4.7	155.2	4.5	1.2	1.2	1.7	1.5	1.3	1.4	1.4
2001.....	4.6	158.4	5.3	1.0	1.2	1.9	1.8	1.4	1.2	1.5
2000.....	4.9	179.2	5.0	1.3	1.4	1.8	1.3	1.4	1.4	1.6
1999.....	4.8	169.3	5.5	1.4	1.7	1.4	1.5	1.0	1.2	1.4

Black, male

2005 4/.....	4.8	166.5	3.6	*	1.3	2.1	1.8	2.1	*	1.8
2004 4/.....	5.1	169.9	5.8	1.2	*	2.0	2.5	*	1.6	1.7
2003 4/.....	5.1	170.7	4.8	1.2	*	1.7	1.8	*	*	*
2002.....	5.1	151.7	4.6	1.4	1.4	2.1	1.7	1.7	1.6	*
2001.....	5.0	160.8	5.5	*	*	2.6	2.2	*	*	*
2000.....	5.4	181.4	5.5	1.6	1.4	2.5	*	1.9	1.7	1.8
1999.....	5.2	172.7	6.2	1.4	1.8	1.7	1.8	*	*	*

Black, female

2005 4/.....	4.1	159.0	4.5	1.5	1.4	*	*	*	1.5	*
2004 4/.....	4.0	155.6	4.3	1.8	*	1.6	*	*	*	*
2003 4/.....	3.9	143.8	4.6	*	*	*	1.5	*	1.6	*
2002.....	4.3	158.9	4.4	*	*	1.3	1.3	*	*	1.4
2001.....	4.3	155.9	5.1	*	1.4	1.3	1.5	1.7	*	1.7
2000.....	4.4	176.9	4.6	*	1.5	*	*	*	*	1.5
1999.....	4.3	165.8	4.8	1.5	1.7	*	*	*	*	1.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

All races 3/, both sexes

2005.....	10.8	87.4	1.4	0.3	0.5	1.2	2.5	2.8	3.1	4.1
2004.....	10.3	82.4	1.2	0.3	0.3	1.0	2.1	2.7	2.8	3.6
2003.....	10.8	82.9	1.6	0.4	0.5	1.5	2.6	3.2	3.4	4.8
2002.....	10.4	85.7	1.6	0.3	0.4	1.3	2.4	3.0	3.6	4.6
2001.....	11.4	86.5	1.8	0.4	0.5	1.8	3.0	3.5	4.2	6.2
2000.....	11.3	95.6	1.6	0.4	0.4	1.6	2.9	3.4	4.3	5.7
1999.....	9.6	93.2	1.5	0.3	0.4	1.1	2.0	2.3	3.1	4.1

All races 3/, male

2005.....	10.0	101.8	1.6	0.3	0.6	1.6	3.4	3.9	4.0	5.1
2004.....	9.5	96.9	1.2	0.3	0.2	1.3	3.0	3.6	3.6	4.6
2003.....	10.2	96.5	1.8	0.3	0.6	1.9	3.4	4.4	4.5	6.4
2002.....	10.1	98.5	1.8	0.4	0.4	1.6	3.2	4.1	4.9	6.1
2001.....	11.3	98.9	2.1	0.4	0.5	2.4	4.0	4.8	5.6	8.0
2000.....	11.5	110.6	1.7	0.5	0.5	1.9	4.0	4.7	5.9	7.6
1999.....	9.4	106.6	1.7	0.3	0.4	1.5	2.8	2.9	4.2	5.3

All races 3/, female

2005.....	11.5	72.3	1.3	0.3	0.3	0.8	1.4	1.7	2.1	3.1
2004.....	11.0	67.3	1.1	0.2	0.4	0.7	1.3	1.7	2.0	2.7
2003.....	11.4	68.6	1.5	0.5	0.4	1.0	1.7	1.9	2.3	3.3
2002.....	10.7	72.3	1.4	0.2	0.3	0.8	1.6	1.8	2.3	3.1
2001.....	11.4	73.6	1.5	0.4	0.5	1.1	1.9	2.2	2.8	4.4
2000.....	11.2	79.9	1.5	0.4	0.4	1.2	1.8	2.2	2.8	3.7
1999.....	9.7	79.2	1.3	0.3	0.3	0.8	1.2	1.6	2.1	3.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

White, both sexes

2005 4/.....	11.1	72.9	1.2	0.2	0.4	1.1	2.4	2.7	3.1	4.0
2004 4/.....	10.5	65.8	1.0	0.2	0.2	1.0	2.2	2.6	2.7	3.6
2003 4/.....	11.1	68.1	1.4	0.4	0.4	1.5	2.5	3.1	3.4	4.7
2002.....	10.5	71.0	1.3	0.2	0.3	1.2	2.3	2.8	3.5	4.4
2001.....	11.6	70.9	1.5	0.3	0.4	1.8	3.0	3.5	4.1	6.0
2000.....	11.4	77.2	1.5	0.4	0.4	1.5	2.8	3.2	4.2	5.6
1999.....	9.4	74.7	1.3	0.3	0.3	1.0	1.8	2.0	2.9	3.7

White, male

2005 4/.....	10.0	87.7	1.4	*	0.5	1.6	3.4	3.8	4.1	4.9
2004 4/.....	9.4	76.9	1.1	0.3	*	1.2	3.0	3.6	3.5	4.6
2003 4/.....	10.1	82.4	1.4	0.3	0.4	2.0	3.4	4.2	4.5	6.1
2002.....	9.8	82.5	1.4	0.3	0.4	1.6	3.2	3.9	4.7	5.8
2001.....	11.2	82.0	1.7	0.3	0.5	2.4	4.0	4.9	5.5	7.7
2000.....	11.3	89.4	1.7	0.4	0.4	1.9	3.9	4.4	5.7	7.7
1999.....	9.0	86.4	1.6	0.3	0.3	1.3	2.6	2.7	3.8	4.9

White, female

2005 4/.....	12.2	57.4	1.1	*	*	0.7	1.4	1.5	2.0	3.0
2004 4/.....	11.6	54.2	1.0	*	0.3	0.7	1.3	1.5	1.9	2.5
2003 4/.....	12.0	53.3	1.3	0.4	0.3	1.0	1.6	1.9	2.1	3.2
2002.....	11.2	59.0	1.2	*	*	0.8	1.4	1.6	2.2	3.0
2001.....	11.9	59.3	1.4	0.4	0.4	1.2	1.9	1.9	2.7	4.3
2000.....	11.5	64.3	1.3	0.3	0.4	1.1	1.6	1.9	2.6	3.5
1999.....	9.9	62.4	1.1	0.3	*	0.7	1.0	1.3	2.1	2.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)										
Black, both sexes										
2005 4/.....	12.3	169.0	2.5	0.8	0.9	1.9	3.1	4.2	4.2	6.0
2004 4/.....	11.6	171.7	1.9	*	0.6	1.3	2.3	3.9	4.2	5.2
2003 4/.....	12.3	159.3	3.1	*	1.0	1.5	3.4	4.9	5.0	7.1
2002.....	12.3	162.3	2.6	0.7	0.7	1.8	3.2	4.6	5.6	7.5
2001.....	12.6	169.5	2.8	0.6	0.6	1.8	3.8	4.6	5.3	8.2
2000.....	13.7	197.1	2.3	0.6	0.8	2.2	4.3	5.3	6.3	7.3
1999.....	12.7	192.8	2.7	*	*	1.7	3.7	4.3	5.4	7.8
Black, male										
2005 4/.....	13.3	186.6	2.7	*	*	2.4	4.5	5.7	5.2	7.1
2004 4/.....	12.6	204.3	2.1	*	*	1.6	3.1	4.8	5.5	6.3
2003 4/.....	13.5	170.7	3.4	*	1.3	1.8	4.2	7.3	6.2	9.6
2002.....	14.1	183.0	3.5	*	*	2.4	3.6	5.8	7.8	10.3
2001.....	13.9	187.4	3.5	*	*	2.4	5.2	5.5	6.5	10.5
2000.....	15.5	227.1	2.3	*	*	2.0	5.6	7.1	8.2	8.3
1999.....	14.6	217.6	2.8	*	*	2.4	4.7	5.2	8.3	9.4
Black, female										
2005 4/.....	11.4	150.7	2.2	*	*	1.3	1.7	2.8	3.3	5.0
2004 4/.....	10.8	137.7	1.7	*	*	*	1.5	3.0	2.9	4.3
2003 4/.....	11.3	147.5	2.9	*	*	*	2.7	2.7	3.9	4.9
2002.....	10.6	140.8	1.8	*	*	*	2.7	3.5	3.7	5.1
2001.....	11.4	150.6	2.0	*	*	*	2.4	3.9	4.2	6.2
2000.....	12.0	166.0	2.3	*	*	2.3	3.0	3.6	4.6	6.3
1999.....	10.9	167.1	2.7	*	*	*	2.6	3.4	2.8	6.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)

All races 3/, both sexes

2005.....	73.4	29.2	3.8	1.9	2.4	4.0	5.8	7.0	8.7	13.9
2004.....	68.8	29.2	3.8	1.9	2.4	3.8	5.6	7.1	9.0	14.3
2003.....	69.3	29.7	4.0	2.1	2.4	4.1	5.8	7.4	9.3	14.5
2002.....	67.5	28.4	4.2	1.9	2.3	3.5	5.6	6.6	9.0	14.1
2001.....	65.5	28.0	4.2	1.9	2.3	3.8	5.2	6.5	8.9	14.1
2000.....	63.4	29.0	4.0	1.8	2.5	3.6	5.4	6.5	9.0	13.9
1999.....	61.5	30.2	4.2	2.1	2.4	3.9	5.3	6.3	8.9	14.0

All races 3/, male

2005.....	60.9	34.8	4.3	2.0	2.5	5.0	6.5	7.9	9.9	16.1
2004.....	57.2	33.8	4.0	2.0	2.6	4.4	6.4	8.2	10.1	16.6
2003.....	57.3	34.8	4.2	2.3	2.8	4.6	6.6	8.2	10.4	16.5
2002.....	55.8	33.4	4.6	2.2	2.5	4.0	6.3	7.4	10.6	16.4
2001.....	53.8	31.4	4.6	1.8	2.5	4.4	6.0	7.3	9.9	16.8
2000.....	52.3	32.3	4.3	1.9	2.6	4.2	6.5	7.5	10.4	16.5
1999.....	51.0	35.0	4.5	2.2	2.5	4.7	5.9	7.3	10.4	16.8

All races 3/, female

2005.....	85.6	23.2	3.3	1.7	2.2	2.9	5.0	6.1	7.5	11.7
2004.....	79.9	24.5	3.6	1.8	2.2	3.1	4.8	6.0	7.8	12.0
2003.....	81.0	24.5	3.8	1.8	2.1	3.4	4.9	6.5	8.1	12.5
2002.....	78.8	23.0	3.9	1.7	2.1	3.0	4.9	5.8	7.3	11.9
2001.....	76.7	24.5	3.8	2.1	2.0	3.1	4.4	5.6	8.0	11.5
2000.....	74.0	25.6	3.6	1.8	2.3	3.1	4.4	5.4	7.5	11.3
1999.....	71.6	25.3	3.9	2.1	2.3	3.1	4.7	5.3	7.4	11.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)

White, both sexes

2005 4/.....	78.5	25.1	3.5	1.7	2.1	3.9	5.6	6.5	7.9	12.6
2004 4/.....	73.3	25.4	3.5	1.8	2.4	3.5	5.2	6.4	7.9	12.9
2003 4/.....	73.9	25.5	3.7	2.0	2.3	4.0	5.5	6.7	8.3	13.0
2002.....	71.8	25.2	3.9	1.9	2.2	3.4	5.0	6.0	8.1	12.8
2001.....	69.4	23.7	3.9	2.0	2.2	3.6	4.9	5.8	7.9	12.6
2000.....	67.0	24.9	3.5	1.7	2.3	3.5	5.1	5.7	8.0	12.3
1999.....	65.0	25.2	4.0	2.0	2.3	3.8	4.8	5.9	8.1	12.1

White, male

2005 4/.....	64.1	30.0	4.0	1.9	2.3	5.0	6.5	7.6	9.6	15.1
2004 4/.....	59.9	29.6	3.7	2.0	2.6	4.2	6.2	7.8	9.0	15.6
2003 4/.....	59.9	31.5	3.8	2.4	2.6	4.7	6.6	7.8	9.6	15.2
2002.....	58.2	30.8	4.1	2.2	2.3	4.1	5.9	6.8	9.9	15.5
2001.....	55.8	25.9	4.4	1.9	2.4	4.4	5.8	6.6	8.9	15.5
2000.....	54.1	26.8	3.9	1.9	2.6	4.1	6.2	7.0	9.6	15.0
1999.....	52.7	29.2	4.3	2.0	2.2	4.8	5.5	7.0	9.8	15.1

White, female

2005 4/.....	92.6	19.9	3.1	1.5	1.9	2.7	4.6	5.5	6.2	10.0
2004 4/.....	86.4	21.0	3.3	1.6	2.1	2.8	4.2	4.9	6.7	10.2
2003 4/.....	87.6	19.3	3.7	1.6	2.0	3.1	4.2	5.4	6.8	10.7
2002.....	85.2	19.3	3.6	1.6	2.1	2.8	4.1	5.0	6.1	9.9
2001.....	82.7	21.4	3.4	2.0	2.0	2.8	3.9	4.8	6.8	9.7
2000.....	79.6	22.9	3.2	1.6	2.0	2.8	3.9	4.4	6.4	9.5
1999.....	77.0	20.9	3.7	1.9	2.3	2.8	4.1	4.7	6.3	9.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)

Black, both sexes

2005 4/.....	63.6	50.5	5.0	2.8	3.8	4.9	8.1	11.3	15.5	24.7
2004 4/.....	60.9	49.3	5.3	2.6	2.9	5.1	8.5	12.8	17.7	26.1
2003 4/.....	60.9	51.8	5.3	2.2	3.2	4.9	8.1	13.3	16.9	27.2
2002.....	59.7	45.2	5.9	2.2	2.9	4.3	9.4	12.2	16.6	25.7
2001.....	59.1	51.3	5.4	2.0	2.9	5.3	7.7	11.5	16.6	26.6
2000.....	58.1	49.6	5.9	2.3	3.3	4.8	8.3	12.8	17.1	26.9
1999.....	56.2	56.7	5.3	2.9	3.1	4.7	8.9	10.4	15.4	28.7

Black, male

2005 4/.....	56.8	62.6	5.8	2.9	4.1	5.7	7.9	11.7	15.3	25.1
2004 4/.....	56.1	55.7	5.7	2.1	3.0	5.8	8.0	12.4	19.7	26.6
2003 4/.....	55.7	55.3	6.1	1.8	3.9	4.7	7.2	12.9	17.1	28.7
2002.....	54.8	48.8	6.4	2.3	3.4	4.2	9.4	12.7	17.9	26.4
2001.....	54.6	58.4	5.1	1.5	3.3	5.6	7.7	12.3	17.5	29.7
2000.....	54.3	59.8	6.1	2.0	3.1	5.1	9.1	12.9	18.3	29.9
1999.....	52.8	64.7	5.4	3.2	3.8	4.8	9.7	10.5	16.2	31.5

Black, female

2005 4/.....	69.7	37.9	4.1	2.7	3.4	4.2	8.2	10.9	15.6	24.3
2004 4/.....	65.4	42.6	4.9	3.1	2.9	4.3	9.0	13.2	15.9	25.6
2003 4/.....	65.6	48.2	4.5	2.7	2.5	5.2	9.0	13.6	16.7	25.9
2002.....	64.1	41.5	5.4	2.2	2.5	4.3	9.5	11.8	15.5	25.0
2001.....	63.1	43.8	5.6	2.5	2.5	5.1	7.6	10.8	15.7	24.0
2000.....	61.6	39.0	5.7	2.6	3.5	4.6	7.6	12.6	16.1	24.2
1999.....	59.3	48.3	5.1	2.6	2.4	4.5	8.1	10.3	14.6	26.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Accidents (unintentional injuries) (V01-X59,Y85-Y86)

All races 3/, both sexes

2005.....	39.7	26.4	10.3	5.5	6.4	31.4	43.4	36.0	33.8	35.3
2004.....	38.1	25.8	10.3	5.7	7.3	32.9	41.1	34.6	30.6	34.8
2003.....	37.6	23.6	10.9	5.5	7.2	33.0	41.1	32.8	30.2	35.0
2002.....	37.0	23.5	10.5	5.9	7.3	35.0	40.9	32.3	30.8	34.9
2001.....	35.7	24.2	11.2	6.4	7.4	32.8	39.5	31.4	28.5	34.1
2000.....	34.8	23.1	11.9	6.8	7.7	33.4	38.8	30.4	28.6	32.9
1999.....	35.1	22.3	12.4	7.1	8.1	33.3	37.5	30.0	29.2	32.2

All races 3/, male

2005.....	52.3	27.5	12.4	6.3	8.1	43.0	66.2	54.8	49.7	50.6
2004.....	49.8	29.0	11.8	6.8	9.0	44.6	61.8	52.2	45.0	49.7
2003.....	49.3	26.4	13.0	6.4	9.3	44.7	61.8	49.4	45.1	50.2
2002.....	48.9	26.8	12.9	7.1	9.2	47.6	62.4	49.5	46.1	50.5
2001.....	47.2	26.2	13.3	7.6	9.5	45.3	60.7	48.0	43.1	49.5
2000.....	46.2	25.7	14.0	7.8	9.9	45.8	58.9	47.0	43.1	48.9
1999.....	46.4	25.4	14.4	8.5	10.4	44.8	57.1	46.1	43.9	47.4

All races 3/, female

2005.....	27.5	25.2	8.0	4.7	4.7	19.2	19.2	16.3	17.5	19.8
2004.....	26.8	22.5	8.7	4.6	5.4	20.6	19.1	16.2	15.9	19.7
2003.....	26.2	20.7	8.7	4.6	4.9	20.6	19.2	15.5	15.1	19.7
2002.....	25.6	20.0	8.1	4.7	5.3	21.8	18.4	14.5	15.2	19.2
2001.....	24.5	22.1	8.9	5.0	5.3	19.5	17.2	14.3	13.8	18.7
2000.....	23.8	20.5	9.7	5.7	5.5	20.3	17.8	13.5	13.9	17.0
1999.....	24.1	19.0	10.3	5.6	5.6	21.1	17.1	13.7	14.3	17.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Accidents (unintentional injuries) (V01-X59,Y85-Y86)

White, both sexes

2005 4/.....	41.8	23.0	10.0	5.1	6.3	33.9	46.9	38.5	35.3	36.8
2004 4/.....	40.2	22.2	10.0	5.2	7.1	36.0	44.4	36.5	32.1	36.1
2003 4/.....	39.5	20.0	10.6	5.2	7.0	35.9	44.3	34.6	31.6	36.2
2002.....	38.7	18.6	9.9	5.2	7.0	37.9	43.8	33.5	32.2	35.5
2001.....	37.0	20.6	11.0	5.8	7.1	34.9	41.7	32.7	29.3	34.8
2000.....	35.9	19.1	11.1	6.2	7.6	36.0	40.7	30.9	29.1	33.4
1999.....	36.0	18.9	11.5	6.3	7.9	35.6	38.9	30.8	29.3	32.4

White, male

2005 4/.....	54.3	23.7	12.3	5.9	7.8	46.1	71.6	58.1	51.2	52.1
2004 4/.....	52.0	25.5	11.5	6.1	8.9	48.5	67.0	54.6	46.5	51.4
2003 4/.....	51.3	22.5	12.8	6.0	9.0	48.4	66.2	51.9	46.8	51.4
2002.....	50.4	22.3	11.8	6.3	8.8	51.0	66.4	50.7	47.9	50.9
2001.....	48.4	22.1	13.3	6.7	9.0	47.8	64.2	49.6	44.0	50.3
2000.....	47.0	21.7	13.2	7.2	9.6	48.8	61.7	47.2	43.7	49.5
1999.....	46.9	21.8	13.2	7.4	10.1	47.2	58.9	47.1	43.8	47.5

White, female

2005 4/.....	29.5	22.2	7.6	4.3	4.7	21.0	20.3	17.4	18.6	20.9
2004 4/.....	28.8	18.8	8.4	4.3	5.2	22.9	20.1	17.2	16.9	20.3
2003 4/.....	28.0	17.4	8.3	4.4	4.8	22.7	20.7	16.1	15.7	20.6
2002.....	27.3	14.7	7.8	4.2	5.2	23.9	19.6	15.1	15.9	19.7
2001.....	25.9	18.9	8.5	4.8	5.1	21.1	17.8	14.8	14.0	19.0
2000.....	25.1	16.4	9.0	5.3	5.5	22.3	18.3	13.8	13.8	17.0
1999.....	25.3	15.8	9.6	5.1	5.6	23.3	17.6	13.8	14.3	17.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Accidents (unintentional injuries) (V01-X59,Y85-Y86)										
Black, both sexes										
2005 4/.....	34.9	47.4	12.5	7.5	7.6	22.8	31.8	29.4	34.9	36.1
2004 4/.....	32.8	45.8	12.5	8.5	8.5	21.6	30.1	32.2	30.6	34.9
2003 4/.....	32.4	43.5	13.4	7.3	8.1	20.7	30.6	29.8	29.9	35.5
2002.....	33.1	48.9	14.1	9.3	9.0	23.9	32.0	32.8	30.1	38.1
2001.....	33.5	43.1	13.6	9.4	9.4	24.7	33.0	30.5	30.0	35.8
2000.....	33.5	44.5	16.2	9.7	8.8	23.7	34.3	33.1	31.9	35.9
1999.....	35.2	41.6	17.7	11.3	9.6	25.0	34.6	31.5	32.9	36.6
Black, male										
2005 4/.....	50.0	50.8	13.8	8.3	10.3	32.5	48.5	47.2	54.8	55.7
2004 4/.....	46.7	48.3	14.2	10.4	10.1	31.2	44.8	51.7	48.9	51.5
2003 4/.....	46.1	47.6	15.7	8.7	10.6	30.0	47.6	46.6	46.9	54.9
2002.....	47.9	51.1	18.5	11.1	12.1	34.6	49.5	53.6	47.7	58.9
2001.....	48.2	48.5	15.4	12.6	12.9	36.1	51.0	48.3	46.6	53.3
2000.....	49.0	47.7	18.8	11.9	12.1	34.5	52.9	54.1	49.6	55.7
1999.....	51.0	45.9	21.0	14.5	13.0	36.6	54.7	51.0	51.7	55.4
Black, female										
2005 4/.....	21.2	43.8	11.1	6.7	4.8	12.7	14.9	12.6	16.8	18.5
2004 4/.....	20.1	43.2	10.6	6.5	6.7	11.8	15.4	14.1	14.1	20.0
2003 4/.....	19.9	39.3	11.0	5.9	5.5	11.2	13.7	14.3	14.7	18.3
2002.....	19.7	46.6	9.6	7.5	5.8	12.8	14.9	13.8	14.4	19.7
2001.....	20.1	37.5	11.9	6.2	5.8	12.9	15.6	14.2	15.1	20.2
2000.....	19.5	41.3	13.5	7.6	5.4	12.7	16.7	14.1	15.9	18.3
1999.....	20.8	37.3	14.3	8.0	6.1	13.1	15.6	13.8	16.0	19.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Transport accidents (V01-V99,Y85)

All races 3/, both sexes

2005.....	16.3	3.6	4.0	3.3	4.3	24.4	29.3	20.5	17.6	16.3
2004.....	16.4	3.6	4.1	3.5	5.1	26.0	28.5	20.9	16.7	16.4
2003.....	16.5	3.7	4.1	3.5	5.1	26.7	28.8	20.1	16.5	16.7
2002.....	16.8	3.1	4.1	3.5	4.9	28.4	29.9	20.5	17.1	17.1
2001.....	16.6	3.7	4.4	3.8	5.0	26.6	29.2	20.9	16.9	17.4
2000.....	16.6	4.6	4.5	4.1	5.3	27.1	29.3	20.4	16.8	17.1
1999.....	16.6	5.0	4.6	4.4	5.3	27.1	28.2	20.0	17.0	16.9

All races 3/, male

2005.....	23.3	3.5	4.5	3.7	5.1	31.8	44.0	31.2	26.0	23.5
2004.....	23.0	3.9	4.4	3.9	6.0	33.6	42.0	31.2	24.4	24.0
2003.....	23.3	4.0	4.5	4.0	6.3	34.7	42.4	29.8	24.8	24.1
2002.....	23.7	3.5	4.8	4.0	5.9	37.0	44.4	31.0	25.4	24.7
2001.....	23.5	3.3	5.0	4.3	6.2	35.4	43.9	31.4	25.4	24.7
2000.....	23.3	4.8	5.0	4.4	6.4	35.5	43.3	30.8	24.7	24.8
1999.....	23.1	5.2	5.1	5.0	6.5	34.8	41.8	29.9	24.7	24.0

All races 3/, female

2005.....	9.5	3.6	3.5	3.0	3.5	16.5	13.6	9.3	8.9	8.9
2004.....	9.9	3.2	3.8	3.0	4.1	18.0	14.2	10.2	8.9	8.8
2003.....	10.0	3.4	3.7	3.0	3.8	18.3	14.3	10.0	8.1	9.3
2002.....	10.1	2.8	3.4	3.1	3.9	19.3	14.6	9.7	8.7	9.6
2001.....	10.0	4.2	3.8	3.3	3.8	17.3	13.8	10.1	8.3	10.2
2000.....	10.2	4.3	4.0	3.8	4.1	18.1	14.6	9.8	8.7	9.5
1999.....	10.4	4.8	4.2	3.6	4.1	19.0	14.0	10.0	9.3	9.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Transport accidents (V01-V99,Y85)										
White, both sexes										
2005 4/.....	16.9	3.1	3.9	3.3	4.3	26.4	31.1	21.2	17.9	16.6
2004 4/.....	16.9	3.3	4.0	3.4	5.2	28.6	30.3	21.4	17.0	16.7
2003 4/.....	17.0	3.6	4.0	3.5	5.2	29.2	30.4	20.7	16.8	17.0
2002.....	17.3	2.7	3.9	3.4	5.0	31.0	31.5	20.7	17.6	17.2
2001.....	17.0	3.7	4.3	3.6	5.0	28.6	30.4	21.6	17.1	17.6
2000.....	17.0	4.1	4.3	4.1	5.4	29.6	30.5	20.3	16.8	17.3
1999.....	17.0	4.6	4.1	4.1	5.5	29.4	29.2	20.4	17.2	16.8
White, male										
2005 4/.....	24.0	3.3	4.3	3.7	5.1	34.2	46.7	32.0	26.0	23.5
2004 4/.....	23.7	3.3	4.2	3.8	6.3	36.7	44.8	31.5	24.3	24.3
2003 4/.....	23.9	4.0	4.5	3.9	6.5	37.6	44.4	30.6	25.1	24.2
2002.....	24.2	3.1	4.5	3.8	6.0	40.2	46.7	31.0	25.9	24.5
2001.....	23.9	3.2	4.9	4.0	6.2	37.8	45.8	32.3	25.4	24.7
2000.....	23.6	4.5	4.7	4.4	6.5	38.4	45.0	30.3	24.6	24.8
1999.....	23.4	5.2	4.4	4.7	6.7	37.3	43.1	30.2	24.6	23.7
White, female										
2005 4/.....	10.0	2.9	3.5	3.0	3.6	18.2	14.2	9.7	9.5	9.4
2004 4/.....	10.3	3.2	3.8	2.9	4.1	20.0	14.7	10.5	9.3	8.9
2003 4/.....	10.3	3.1	3.4	3.1	3.9	20.2	15.3	10.2	8.2	9.6
2002.....	10.5	2.2	3.4	2.9	3.9	21.3	15.3	9.8	9.0	9.7
2001.....	10.3	4.1	3.6	3.1	3.8	18.9	14.0	10.4	8.4	10.4
2000.....	10.5	3.7	3.8	3.7	4.3	20.2	15.1	9.9	8.7	9.8
1999.....	10.7	3.9	3.8	3.5	4.3	21.0	14.4	10.1	9.6	9.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Transport accidents (V01-V99,Y85)										
Black, both sexes										
2005 4/.....	15.0	5.7	4.5	3.6	4.3	17.1	23.6	19.2	20.0	17.9
2004 4/.....	15.2	4.8	5.1	4.0	4.8	16.6	22.7	22.4	19.3	17.6
2003 4/.....	15.1	4.2	5.0	3.6	4.5	16.2	23.2	20.3	18.3	17.9
2002.....	15.4	5.3	5.0	4.6	4.6	17.9	24.6	23.0	18.0	19.2
2001.....	15.9	*	5.2	5.3	5.3	18.8	25.8	21.0	18.9	18.1
2000.....	16.0	6.5	5.6	4.5	4.8	17.3	26.6	23.5	19.4	17.9
1999.....	16.3	7.3	7.1	5.8	5.0	18.7	25.8	21.3	18.5	19.9
Black, male										
2005 4/.....	22.7	*	4.9	3.6	5.5	23.5	35.8	31.2	33.6	29.4
2004 4/.....	22.8	5.9	5.4	4.4	5.3	22.9	33.2	36.1	31.5	27.9
2003 4/.....	22.6	*	5.1	4.5	5.7	22.7	36.3	31.8	29.4	28.4
2002.....	23.1	*	6.4	5.0	5.6	24.5	37.1	37.5	28.9	29.8
2001.....	23.8	*	5.7	6.6	7.3	26.5	39.1	33.2	30.0	27.2
2000.....	24.1	6.7	6.2	4.9	6.5	24.2	40.3	38.2	30.7	28.6
1999.....	24.2	*	8.0	7.2	6.4	25.8	39.2	34.3	29.4	29.7
Black, female										
2005 4/.....	7.9	6.8	4.2	3.5	3.0	10.6	11.2	7.9	7.8	7.7
2004 4/.....	8.2	*	4.8	3.6	4.2	10.0	12.1	9.6	8.3	8.5
2003 4/.....	8.3	*	5.0	2.8	3.4	9.5	10.2	9.8	8.2	8.4
2002.....	8.3	*	3.4	4.2	3.5	11.2	12.5	9.8	8.2	9.8
2001.....	8.7	*	4.7	4.0	3.4	10.8	13.0	9.9	8.9	10.0
2000.....	8.5	*	5.0	4.1	3.0	10.3	13.7	10.2	9.2	8.4
1999.....	9.2	8.7	6.2	4.4	3.6	11.4	13.1	9.5	8.6	11.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)

All races 3/, both sexes

2005.....	15.3	3.6	3.8	3.2	4.0	23.6	28.2	19.6	16.5	15.1
2004.....	15.3	3.5	4.0	3.3	4.8	25.2	27.5	19.7	15.6	15.2
2003.....	15.4	3.6	3.9	3.3	4.7	25.7	27.5	19.0	15.4	15.4
2002.....	15.7	3.0	3.9	3.3	4.5	27.6	28.8	19.7	16.0	15.8
2001.....	15.4	3.6	4.1	3.6	4.6	25.7	28.0	19.6	15.6	15.9
2000.....	15.4	4.4	4.2	3.8	4.8	26.0	28.0	19.2	15.5	15.6
1999.....	15.2	4.8	4.2	3.9	4.8	25.9	26.5	18.4	15.4	15.1

All races 3/, male

2005.....	21.7	3.5	4.2	3.5	4.7	30.7	42.3	29.6	24.2	21.8
2004.....	21.3	3.8	4.2	3.7	5.6	32.5	40.4	29.3	22.4	21.9
2003.....	21.4	3.9	4.4	3.7	5.9	33.3	40.4	28.2	23.0	21.9
2002.....	21.9	3.3	4.5	3.8	5.3	36.0	42.6	29.6	23.6	22.4
2001.....	21.5	3.2	4.5	4.0	5.7	34.0	42.1	29.5	23.1	22.3
2000.....	21.3	4.6	4.7	4.1	5.8	33.9	41.2	28.6	22.6	22.1
1999.....	20.9	5.0	4.7	4.4	5.7	33.1	39.1	27.2	22.2	21.2

All races 3/, female

2005.....	9.1	3.6	3.4	2.9	3.2	16.2	13.2	9.1	8.6	8.4
2004.....	9.5	3.2	3.7	2.8	3.9	17.5	13.8	9.7	8.5	8.3
2003.....	9.5	3.3	3.5	2.8	3.5	17.8	13.9	9.5	7.7	8.8
2002.....	9.8	2.8	3.3	2.9	3.6	18.9	14.3	9.4	8.3	9.2
2001.....	9.5	4.0	3.6	3.0	3.5	16.9	13.4	9.5	8.0	9.6
2000.....	9.7	4.2	3.8	3.5	3.8	17.5	14.2	9.5	8.2	9.1
1999.....	9.7	4.7	3.8	3.3	3.8	18.3	13.4	9.4	8.5	9.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)

White, both sexes

2005 4/.....	15.8	3.1	3.7	3.2	4.0	25.6	30.0	20.3	16.8	15.4
2004 4/.....	15.8	3.2	3.9	3.2	4.9	27.8	29.2	20.0	15.7	15.3
2003 4/.....	15.8	3.5	3.8	3.3	4.8	28.1	29.1	19.5	15.7	15.6
2002.....	16.2	2.6	3.7	3.2	4.6	30.2	30.4	19.8	16.4	15.8
2001.....	15.7	3.5	3.9	3.3	4.6	27.6	29.1	20.2	15.7	16.1
2000.....	15.7	3.9	4.0	3.7	5.0	28.4	29.1	19.0	15.4	15.8
1999.....	15.5	4.5	3.8	3.7	5.0	28.1	27.4	18.6	15.5	14.9

White, male

2005 4/.....	22.3	3.3	4.1	3.5	4.6	33.0	45.1	30.4	24.1	21.7
2004 4/.....	21.9	3.3	4.0	3.6	5.9	35.6	43.2	29.5	22.2	22.1
2003 4/.....	22.0	3.9	4.3	3.6	6.0	36.1	42.3	28.8	23.3	21.9
2002.....	22.4	2.9	4.2	3.6	5.5	39.0	44.8	29.5	23.9	22.2
2001.....	21.9	3.1	4.4	3.7	5.6	36.2	43.8	30.1	23.1	22.3
2000.....	21.6	4.2	4.5	4.0	5.9	36.8	42.6	27.9	22.4	22.1
1999.....	21.1	5.0	4.1	4.2	5.9	35.5	40.4	27.4	22.1	20.8

White, female

2005 4/.....	9.5	2.9	3.4	2.8	3.2	17.7	13.8	9.4	9.1	8.8
2004 4/.....	9.8	3.2	3.7	2.7	3.9	19.5	14.1	9.9	8.9	8.3
2003 4/.....	9.8	3.0	3.2	2.9	3.6	19.6	14.8	9.6	7.7	9.0
2002.....	10.1	2.2	3.2	2.7	3.7	20.9	15.0	9.5	8.6	9.3
2001.....	9.8	3.8	3.4	2.9	3.5	18.5	13.5	9.7	8.0	9.8
2000.....	10.0	3.5	3.6	3.4	4.0	19.5	14.6	9.6	8.2	9.3
1999.....	10.0	3.9	3.5	3.2	4.0	20.2	13.7	9.5	8.7	8.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)

Black, both sexes

2005 4/.....	14.1	5.7	4.3	3.4	4.0	16.7	22.4	18.4	19.0	17.0
2004 4/.....	14.3	4.7	4.9	3.8	4.5	15.8	21.8	21.2	18.4	16.6
2003 4/.....	14.2	4.2	4.9	3.4	4.3	15.5	22.3	19.5	17.3	16.7
2002.....	14.5	5.2	4.8	4.3	4.2	17.4	24.0	22.4	17.2	17.9
2001.....	14.8	*	5.0	4.9	4.8	18.1	25.2	20.1	17.7	16.7
2000.....	15.0	6.5	5.4	4.3	4.6	16.5	25.8	22.8	18.0	16.4
1999.....	15.1	7.1	6.4	5.3	4.6	17.8	24.8	20.1	17.0	18.2

Black, male

2005 4/.....	21.2	*	4.7	3.4	5.1	22.7	33.8	29.9	31.9	27.6
2004 4/.....	21.2	*	5.2	4.2	5.0	21.6	31.5	33.8	29.8	26.0
2003 4/.....	21.1	*	5.0	4.1	5.2	21.6	34.6	30.5	27.7	26.3
2002.....	21.5	*	6.3	4.8	5.0	23.7	36.1	36.2	27.4	27.6
2001.....	22.0	*	5.3	6.2	6.5	25.5	38.0	31.9	28.1	24.9
2000.....	22.5	6.7	5.8	4.5	6.2	22.8	38.8	36.7	28.5	26.0
1999.....	22.2	*	7.0	6.3	5.7	24.5	37.7	32.4	27.0	27.0

Black, female

2005 4/.....	7.5	6.8	3.9	3.4	3.0	10.6	10.8	7.6	7.5	7.5
2004 4/.....	7.9	*	4.7	3.5	4.0	9.8	12.0	9.4	8.1	8.2
2003 4/.....	8.0	*	4.8	2.7	3.4	9.2	10.1	9.4	8.0	8.2
2002.....	8.0	*	3.4	3.8	3.3	11.0	12.2	9.7	7.9	9.3
2001.....	8.2	*	4.6	3.6	3.1	10.5	12.9	9.5	8.4	9.5
2000.....	8.2	*	4.9	4.0	2.9	10.1	13.5	10.2	8.6	7.9
1999.....	8.7	8.7	5.8	4.2	3.5	11.0	12.6	9.0	8.1	10.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9,V88.9,V89.1,V89.3,V89.9)

All races 3/, both sexes

2005.....	0.4	*	0.2	*	0.2	0.4	0.5	0.4	0.4	0.5
2004.....	0.4	*	*	0.1	0.2	0.5	0.4	0.5	0.4	0.4
2003.....	0.5	*	*	0.1	0.2	0.6	0.7	0.5	0.5	0.5
2002.....	0.4	*	*	0.1	0.2	0.3	0.5	0.3	0.4	0.5
2001.....	0.5	*	0.2	0.1	0.3	0.5	0.5	0.4	0.4	0.6
2000.....	0.5	*	0.2	0.2	0.3	0.6	0.6	0.5	0.6	0.7
1999.....	0.8	*	0.3	0.3	0.3	0.9	1.0	0.8	0.8	0.9

All races 3/, male

2005.....	0.7	*	*	*	0.3	0.7	0.7	0.7	0.6	0.7
2004.....	0.6	*	*	*	0.2	0.7	0.7	0.8	0.7	0.7
2003.....	0.7	*	*	*	0.2	0.9	1.0	0.8	0.8	0.8
2002.....	0.6	*	*	*	0.3	0.5	0.8	0.5	0.6	0.8
2001.....	0.8	*	0.3	*	0.3	0.7	0.9	0.7	0.8	1.1
2000.....	0.8	*	*	0.2	0.4	1.0	1.0	0.8	0.9	1.1
1999.....	1.1	*	0.3	0.4	0.5	1.2	1.7	1.3	1.1	1.4

All races 3/, female

2005.....	0.2	*	*	*	*	*	*	*	*	0.2
2004.....	0.2	*	*	*	*	0.2	*	*	*	*
2003.....	0.2	*	*	*	*	0.3	0.2	0.2	*	0.2
2002.....	0.1	*	*	*	*	*	*	*	*	*
2001.....	0.2	*	*	*	*	0.2	*	*	*	0.2
2000.....	0.3	*	*	*	*	0.3	0.2	*	0.3	0.2
1999.....	0.4	*	0.3	*	*	0.5	0.3	0.3	0.5	0.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9,V88.9,V89.1,V89.3,V89.9)

White, both sexes

2005 4/.....	0.4	*	0.2	*	0.2	0.5	0.4	0.4	0.4	0.5
2004 4/.....	0.4	*	*	*	0.2	0.4	0.4	0.5	0.4	0.5
2003 4/.....	0.5	*	*	*	0.2	0.7	0.7	0.6	0.5	0.5
2002.....	0.4	*	*	*	0.2	0.3	0.5	0.3	0.4	0.5
2001.....	0.5	*	0.2	*	0.2	0.5	0.6	0.5	0.4	0.5
2000.....	0.5	*	0.2	0.2	0.3	0.7	0.6	0.5	0.6	0.6
1999.....	0.8	*	0.3	0.3	0.3	0.9	1.0	0.8	0.8	0.9

White, male

2005 4/.....	0.6	*	*	*	0.3	0.7	0.7	0.6	0.6	0.7
2004 4/.....	0.6	*	*	*	0.2	0.7	0.6	0.7	0.7	0.7
2003 4/.....	0.7	*	*	*	0.3	1.0	1.1	0.9	0.8	0.8
2002.....	0.6	*	*	*	0.3	0.5	0.9	0.5	0.6	0.8
2001.....	0.7	*	*	*	0.3	0.7	0.9	0.8	0.7	0.9
2000.....	0.8	*	*	*	0.4	1.0	1.0	0.9	0.9	1.1
1999.....	1.1	*	*	0.4	0.5	1.2	1.7	1.3	1.1	1.4

White, female

2005 4/.....	0.2	*	*	*	*	*	*	*	*	0.2
2004 4/.....	0.2	*	*	*	*	*	*	*	*	*
2003 4/.....	0.2	*	*	*	*	0.4	0.3	*	*	*
2002.....	0.1	*	*	*	*	*	*	*	*	*
2001.....	0.2	*	*	*	*	*	*	*	*	*
2000.....	0.3	*	*	*	*	0.3	*	*	0.3	0.2
1999.....	0.4	*	*	*	*	0.6	0.4	0.3	0.5	0.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)

All races 3/, both sexes

2005.....	0.6	*	*	*	0.2	0.3	0.6	0.5	0.7	0.7
2004.....	0.7	*	*	*	0.1	0.4	0.6	0.8	0.7	0.9
2003.....	0.7	*	*	0.1	0.2	0.4	0.6	0.6	0.6	0.9
2002.....	0.7	*	*	0.1	0.2	0.4	0.6	0.5	0.7	0.9
2001.....	0.7	*	0.2	0.1	0.1	0.4	0.6	0.8	0.9	0.9
2000.....	0.7	*	*	0.1	0.1	0.5	0.7	0.8	0.8	0.9
1999.....	0.7	*	*	0.2	0.2	0.4	0.6	0.8	0.8	0.9

All races 3/, male

2005.....	1.0	*	*	*	*	0.4	0.9	0.9	1.1	1.1
2004.....	1.1	*	*	*	*	0.5	0.9	1.2	1.2	1.4
2003.....	1.1	*	*	*	0.2	0.6	1.0	0.8	1.0	1.4
2002.....	1.1	*	*	*	0.2	0.6	0.9	0.9	1.2	1.5
2001.....	1.2	*	*	*	0.2	0.7	1.0	1.2	1.5	1.4
2000.....	1.2	*	*	*	0.2	0.6	1.2	1.3	1.3	1.5
1999.....	1.1	*	*	0.2	0.3	0.6	0.9	1.4	1.3	1.4

All races 3/, female

2005.....	0.2	*	*	*	*	0.2	*	*	0.2	0.3
2004.....	0.3	*	*	*	*	0.2	0.3	0.3	0.2	0.3
2003.....	0.2	*	*	*	*	*	0.2	0.3	0.2	0.3
2002.....	0.2	*	*	*	*	0.2	*	*	0.2	0.3
2001.....	0.3	*	*	*	*	*	0.3	0.4	0.3	0.4
2000.....	0.2	*	*	*	*	0.3	*	*	0.2	0.2
1999.....	0.3	*	*	*	*	0.2	0.3	0.3	0.3	0.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)										
White, both sexes										
2005 4/.....	0.7	*	*	*	0.2	0.4	0.6	0.6	0.7	0.7
2004 4/.....	0.7	*	*	*	0.2	0.4	0.7	0.9	0.8	0.9
2003 4/.....	0.7	*	*	0.1	0.2	0.4	0.6	0.6	0.7	0.9
2002.....	0.7	*	*	*	0.2	0.5	0.6	0.6	0.8	0.9
2001.....	0.8	*	0.2	0.1	0.2	0.5	0.7	0.9	1.0	1.0
2000.....	0.7	*	*	*	0.2	0.5	0.8	0.9	0.8	0.9
1999.....	0.7	*	*	0.2	0.2	0.4	0.7	1.0	0.9	1.0
White, male										
2005 4/.....	1.1	*	*	*	*	0.5	1.0	1.0	1.2	1.1
2004 4/.....	1.2	*	*	*	*	0.5	1.0	1.3	1.4	1.5
2003 4/.....	1.2	*	*	*	*	0.6	1.0	1.0	1.1	1.5
2002.....	1.2	*	*	*	*	0.6	1.1	1.0	1.4	1.6
2001.....	1.3	*	*	*	*	0.8	1.1	1.4	1.6	1.5
2000.....	1.3	*	*	*	*	0.6	1.4	1.5	1.4	1.6
1999.....	1.2	*	*	*	0.3	0.6	1.1	1.5	1.4	1.5
White, female										
2005 4/.....	0.3	*	*	*	*	0.3	*	*	*	0.4
2004 4/.....	0.3	*	*	*	*	0.3	0.4	0.4	0.3	0.3
2003 4/.....	0.3	*	*	*	*	*	*	0.3	0.3	0.4
2002.....	0.2	*	*	*	*	0.3	*	*	*	0.3
2001.....	0.3	*	*	*	*	*	0.4	0.5	0.3	0.4
2000.....	0.2	*	*	*	*	0.3	*	*	*	*
1999.....	0.3	*	*	*	*	*	0.3	0.3	0.4	0.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Nontransport accidents (W00-X59,Y86)

All races 3/, both sexes

2005.....	23.4	22.8	6.3	2.2	2.1	7.1	14.2	15.5	16.2	19.1
2004.....	21.8	22.2	6.1	2.3	2.2	6.9	12.6	13.7	13.8	18.3
2003.....	21.0	19.9	6.8	2.0	2.1	6.3	12.3	12.7	13.7	18.3
2002.....	20.2	20.3	6.4	2.4	2.4	6.6	11.1	11.8	13.6	17.8
2001.....	19.0	20.5	6.8	2.5	2.4	6.2	10.3	10.5	11.6	16.7
2000.....	18.2	18.6	7.4	2.7	2.5	6.3	9.5	10.0	11.8	15.8
1999.....	18.4	17.3	7.7	2.7	2.7	6.2	9.3	10.0	12.1	15.3

All races 3/, male

2005.....	29.0	24.0	7.9	2.6	3.0	11.2	22.2	23.6	23.7	27.1
2004.....	26.8	25.1	7.4	2.9	3.1	11.0	19.8	21.1	20.5	25.7
2003.....	26.0	22.4	8.5	2.5	3.0	10.0	19.4	19.6	20.2	26.2
2002.....	25.2	23.3	8.1	3.1	3.3	10.6	18.0	18.4	20.6	25.9
2001.....	23.8	22.9	8.3	3.3	3.3	9.9	16.8	16.6	17.7	24.8
2000.....	22.9	20.9	9.0	3.4	3.5	10.2	15.5	16.2	18.4	24.2
1999.....	23.3	20.2	9.3	3.5	3.9	10.0	15.3	16.2	19.2	23.4

All races 3/, female

2005.....	18.0	21.5	4.5	1.7	1.2	2.7	5.6	7.0	8.6	10.9
2004.....	16.9	19.3	4.8	1.6	1.3	2.6	4.9	6.0	7.0	10.9
2003.....	16.2	17.3	5.0	1.6	1.2	2.3	4.9	5.5	7.0	10.4
2002.....	15.4	17.2	4.7	1.6	1.5	2.5	3.9	4.9	6.5	9.7
2001.....	14.5	17.9	5.1	1.8	1.5	2.3	3.4	4.2	5.5	8.5
2000.....	13.6	16.2	5.7	1.9	1.4	2.2	3.2	3.7	5.2	7.4
1999.....	13.7	14.2	6.1	2.0	1.5	2.1	3.1	3.7	5.0	7.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Nontransport accidents (W00-X59,Y86)

White, both sexes

2005 4/.....	24.9	19.8	6.1	1.8	1.9	7.5	15.8	17.3	17.4	20.2
2004 4/.....	23.3	19.0	6.0	1.8	1.9	7.4	14.1	15.2	15.1	19.4
2003 4/.....	22.5	16.4	6.6	1.7	1.8	6.8	13.9	13.9	14.8	19.2
2002.....	21.4	15.9	5.9	1.8	2.0	6.9	12.3	12.7	14.6	18.3
2001.....	20.0	16.9	6.7	2.2	2.1	6.3	11.3	11.0	12.2	17.2
2000.....	18.9	15.1	6.9	2.2	2.2	6.4	10.2	10.5	12.2	16.0
1999.....	19.0	14.3	7.3	2.2	2.4	6.2	9.7	10.4	12.1	15.6

White, male

2005 4/.....	30.3	20.4	8.0	2.2	2.7	11.9	24.8	26.1	25.3	28.6
2004 4/.....	28.3	22.1	7.3	2.2	2.6	11.7	22.1	23.1	22.2	27.1
2003 4/.....	27.4	18.6	8.3	2.1	2.5	10.8	21.7	21.3	21.7	27.2
2002.....	26.2	19.2	7.4	2.4	2.8	10.9	19.7	19.7	22.0	26.4
2001.....	24.5	18.9	8.4	2.7	2.8	10.0	18.4	17.3	18.6	25.5
2000.....	23.4	17.3	8.4	2.8	3.1	10.3	16.7	16.9	19.1	24.6
1999.....	23.5	16.5	8.8	2.7	3.4	10.0	15.8	16.9	19.2	23.8

White, female

2005 4/.....	19.6	19.2	4.1	1.3	1.1	2.8	6.1	7.8	9.1	11.5
2004 4/.....	18.5	15.6	4.6	1.3	1.0	2.9	5.4	6.7	7.6	11.5
2003 4/.....	17.7	14.2	4.9	1.3	0.9	2.5	5.4	5.9	7.6	11.0
2002.....	16.8	12.5	4.4	1.2	1.2	2.7	4.3	5.3	6.9	10.1
2001.....	15.6	14.8	4.8	1.7	1.3	2.3	3.8	4.4	5.6	8.6
2000.....	14.6	12.8	5.2	1.6	1.2	2.2	3.3	3.8	5.1	7.3
1999.....	14.6	11.8	5.8	1.6	1.3	2.2	3.2	3.7	4.8	7.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Nontransport accidents (W00-X59,Y86)

Black, both sexes

2005 4/.....	20.0	41.6	7.9	3.9	3.3	5.6	8.2	10.2	14.8	18.2
2004 4/.....	17.6	41.0	7.4	4.5	3.7	5.0	7.5	9.9	11.3	17.2
2003 4/.....	17.2	39.3	8.4	3.7	3.6	4.5	7.4	9.5	11.6	17.7
2002.....	17.8	43.6	9.2	4.7	4.4	6.0	7.4	9.8	12.1	18.9
2001.....	17.6	40.2	8.4	4.1	4.1	5.9	7.2	9.4	11.1	17.7
2000.....	17.6	38.0	10.6	5.2	4.0	6.4	7.7	9.6	12.5	18.0
1999.....	18.9	34.4	10.6	5.6	4.6	6.3	8.8	10.2	14.4	16.7

Black, male

2005 4/.....	27.3	46.1	8.9	4.6	4.8	9.1	12.7	16.0	21.3	26.3
2004 4/.....	23.9	42.4	8.9	6.1	4.9	8.2	11.7	15.6	17.4	23.7
2003 4/.....	23.5	44.0	10.6	4.2	5.0	7.3	11.3	14.8	17.5	26.4
2002.....	24.8	45.9	12.1	6.1	6.5	10.1	12.4	16.2	18.8	29.1
2001.....	24.4	46.4	9.7	6.0	5.7	9.6	11.8	15.1	16.5	26.1
2000.....	24.9	40.9	12.6	6.9	5.6	10.3	12.6	15.9	18.9	27.1
1999.....	26.8	40.0	13.0	7.4	6.6	10.8	15.5	16.7	22.3	25.7

Black, female

2005 4/.....	13.3	37.0	6.9	3.2	1.8	2.1	3.7	4.7	9.0	10.9
2004 4/.....	11.9	39.5	5.9	2.9	2.5	1.8	3.3	4.5	5.8	11.5
2003 4/.....	11.6	34.3	6.0	3.1	2.1	1.7	3.5	4.5	6.4	9.8
2002.....	11.4	41.2	6.2	3.3	2.3	1.7	2.4	4.0	6.2	9.8
2001.....	11.5	33.7	7.1	2.2	2.5	2.1	2.7	4.3	6.2	10.2
2000.....	11.0	35.1	8.5	3.5	2.3	2.3	3.0	3.8	6.7	9.9
1999.....	11.7	28.5	8.1	3.7	2.5	1.6	2.5	4.3	7.3	8.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Falls (W00-W19)

All races 3/, both sexes

2005.....	6.6	*	0.2	*	*	0.4	0.8	0.7	0.8	1.1
2004.....	6.4	0.6	0.3	*	0.1	0.4	0.7	0.8	0.8	1.1
2003.....	5.9	*	0.3	*	0.1	0.4	0.7	0.7	0.7	1.2
2002.....	5.6	*	0.2	*	0.1	0.4	0.8	0.7	0.8	1.2
2001.....	5.3	0.6	0.2	0.2	0.2	0.4	0.9	0.8	0.9	1.2
2000.....	4.7	*	0.2	*	0.1	0.5	0.7	0.7	0.8	1.0
1999.....	4.7	*	0.4	0.1	0.1	0.6	0.7	0.8	0.9	1.2

All races 3/, male

2005.....	7.0	*	0.2	*	*	0.6	1.2	1.2	1.3	1.8
2004.....	6.8	*	0.4	*	*	0.7	1.3	1.3	1.4	1.8
2003.....	6.2	*	0.5	*	0.2	0.7	1.1	1.3	1.1	2.0
2002.....	6.0	*	0.4	*	*	0.6	1.4	1.2	1.4	1.7
2001.....	5.8	*	0.3	0.2	0.2	0.8	1.5	1.5	1.5	2.1
2000.....	5.2	*	0.3	*	*	0.8	1.2	1.2	1.5	1.8
1999.....	5.2	*	0.5	*	0.2	0.9	1.2	1.4	1.5	2.0

All races 3/, female

2005.....	6.3	*	*	*	*	*	0.3	*	0.3	0.4
2004.....	6.0	*	*	*	*	*	*	0.2	0.2	0.5
2003.....	5.6	*	*	*	*	*	0.2	*	0.3	0.3
2002.....	5.3	*	*	*	*	*	*	*	0.3	0.6
2001.....	4.8	*	*	*	*	*	*	*	0.3	0.4
2000.....	4.3	*	*	*	*	*	0.2	*	*	0.3
1999.....	4.3	*	*	*	*	*	*	*	0.3	0.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Falls (W00-W19)

White, both sexes

2005 4/.....	7.5	*	*	*	*	0.4	0.9	0.8	0.8	1.1
2004 4/.....	7.3	*	0.3	*	0.1	0.5	0.8	0.8	0.9	1.2
2003 4/.....	6.7	*	0.3	*	0.2	0.5	0.8	0.8	0.8	1.2
2002.....	6.4	*	0.2	*	*	0.4	0.9	0.8	0.9	1.2
2001.....	6.0	*	0.2	0.2	0.2	0.5	1.0	0.9	0.9	1.3
2000.....	5.3	*	0.2	*	*	0.6	0.8	0.8	0.9	1.0
1999.....	5.2	*	0.4	*	0.2	0.6	0.8	0.9	0.9	1.3

White, male

2005 4/.....	7.8	*	*	*	*	0.7	1.4	1.4	1.3	1.9
2004 4/.....	7.6	*	0.4	*	*	0.8	1.4	1.4	1.5	2.0
2003 4/.....	6.9	*	0.5	*	*	0.8	1.3	1.4	1.2	2.1
2002.....	6.6	*	*	*	*	0.7	1.6	1.4	1.5	1.8
2001.....	6.4	*	0.3	0.2	0.2	0.9	1.8	1.6	1.6	2.1
2000.....	5.7	*	*	*	*	1.0	1.4	1.4	1.5	1.7
1999.....	5.6	*	0.5	*	*	1.1	1.3	1.5	1.5	2.0

White, female

2005 4/.....	7.3	*	*	*	*	*	0.3	*	*	0.4
2004 4/.....	7.0	*	*	*	*	*	*	*	0.3	0.4
2003 4/.....	6.5	*	*	*	*	*	0.3	*	0.3	0.3
2002.....	6.2	*	*	*	*	*	*	*	0.3	0.7
2001.....	5.5	*	*	*	*	*	*	*	0.3	0.4
2000.....	5.0	*	*	*	*	*	0.3	*	*	0.3
1999.....	4.9	*	*	*	*	*	*	*	0.3	0.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Accidental drowning and submersion (W65-W74)										
All races 3/, both sexes										
2005.....	1.2	1.6	3.0	0.6	0.6	1.5	1.6	1.0	0.9	1.0
2004.....	1.1	1.5	2.7	0.7	0.7	1.5	1.3	1.1	0.8	0.9
2003.....	1.1	1.4	2.9	0.6	0.7	1.4	1.4	1.0	0.8	1.0
2002.....	1.2	1.6	2.9	0.8	0.8	1.6	1.5	1.2	1.0	1.2
2001.....	1.2	1.7	3.0	0.8	0.8	1.6	1.4	1.1	0.8	1.0
2000.....	1.2	2.0	3.2	1.0	0.8	1.8	1.5	1.1	1.0	1.1
1999.....	1.3	1.8	3.2	0.9	0.9	1.8	1.5	1.1	1.1	1.1
All races 3/, male										
2005.....	1.9	1.3	4.1	0.8	1.0	2.6	2.8	1.7	1.5	1.7
2004.....	1.8	1.8	3.2	0.9	1.0	2.6	2.3	1.9	1.4	1.6
2003.....	1.8	1.7	3.7	0.9	1.0	2.5	2.5	1.8	1.4	1.7
2002.....	1.9	1.9	3.9	1.2	1.1	2.8	2.8	2.1	1.7	2.0
2001.....	1.8	1.5	3.7	1.2	1.1	2.8	2.5	1.9	1.4	1.6
2000.....	2.0	2.2	4.0	1.3	1.2	3.3	2.5	1.9	1.7	1.9
1999.....	2.0	2.0	4.0	1.3	1.3	3.2	2.8	2.0	2.0	1.9
All races 3/, female										
2005.....	0.5	1.8	2.0	0.4	0.3	0.3	0.3	0.3	0.3	0.3
2004.....	0.5	1.2	2.1	0.4	0.3	0.3	0.2	0.3	0.2	0.2
2003.....	0.5	1.2	2.0	0.4	0.3	*	0.2	*	*	0.3
2002.....	0.5	1.2	1.9	0.3	0.4	0.3	*	0.3	0.3	0.4
2001.....	0.5	1.9	2.2	0.5	0.5	0.3	0.3	0.3	0.2	0.4
2000.....	0.5	1.7	2.4	0.6	0.4	0.3	0.3	0.3	0.3	0.3
1999.....	0.5	1.6	2.4	0.5	0.4	0.3	0.3	*	0.3	0.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Accidental drowning and submersion (W65-W74)										
White, both sexes										
2005 4/.....	1.2	1.8	3.3	0.5	0.5	1.3	1.6	1.0	0.9	0.9
2004 4/.....	1.1	1.6	3.0	0.5	0.4	1.2	1.3	1.1	0.8	0.9
2003 4/.....	1.1	1.4	3.1	0.6	0.5	1.3	1.3	1.0	0.8	1.0
2002.....	1.1	1.6	3.0	0.6	0.5	1.4	1.5	1.1	1.0	1.1
2001.....	1.1	1.7	3.3	0.6	0.5	1.3	1.4	1.0	0.7	1.0
2000.....	1.2	1.6	3.5	0.8	0.6	1.6	1.4	1.1	1.0	1.1
1999.....	1.2	1.5	3.5	0.7	0.5	1.6	1.5	1.0	1.1	1.1
White, male										
2005 4/.....	1.8	1.3	4.5	0.6	0.6	2.2	2.8	1.8	1.4	1.5
2004 4/.....	1.7	1.8	3.5	0.7	0.6	2.1	2.2	1.8	1.3	1.5
2003 4/.....	1.8	1.5	4.0	0.8	0.7	2.3	2.4	1.7	1.3	1.7
2002.....	1.8	1.9	4.0	0.9	0.6	2.4	2.7	1.9	1.7	1.9
2001.....	1.7	1.6	4.2	0.8	0.6	2.3	2.4	1.7	1.2	1.5
2000.....	1.9	1.8	4.4	1.0	0.8	2.8	2.4	1.8	1.7	1.8
1999.....	1.9	1.7	4.3	0.9	0.7	2.8	2.5	1.7	1.9	1.8
White, female										
2005 4/.....	0.5	2.2	2.1	0.4	0.3	0.3	0.3	0.3	0.3	0.3
2004 4/.....	0.5	1.4	2.4	0.3	*	*	0.3	0.3	0.3	0.3
2003 4/.....	0.5	1.3	2.2	0.3	*	*	*	*	*	0.3
2002.....	0.4	*	2.0	0.3	0.3	0.3	*	0.3	*	0.4
2001.....	0.5	1.8	2.4	0.5	0.3	*	0.3	0.3	*	0.4
2000.....	0.5	*	2.5	0.5	0.3	0.3	0.3	0.3	*	0.4
1999.....	0.5	1.4	2.6	0.4	*	0.3	0.3	*	0.3	0.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Accidental drowning and submersion (W65-W74)										
Black, both sexes										
2005 4/.....	1.4	*	2.2	1.2	1.4	2.2	1.9	1.1	1.1	1.5
2004 4/.....	1.4	*	1.9	1.3	1.8	2.4	1.2	1.3	0.9	1.1
2003 4/.....	1.3	*	2.1	1.0	1.6	1.8	1.4	1.1	1.1	0.8
2002.....	1.6	*	2.5	1.8	2.2	2.3	1.9	1.6	1.2	1.5
2001.....	1.6	*	1.9	1.7	2.3	2.9	1.5	1.2	1.1	1.2
2000.....	1.6	4.2	2.2	1.9	2.1	2.8	1.7	1.3	0.9	1.2
1999.....	1.8	3.5	2.4	2.2	2.6	2.7	1.9	1.8	1.4	1.1
Black, male										
2005 4/.....	2.5	*	2.7	1.6	2.6	4.2	3.3	1.9	2.1	2.7
2004 4/.....	2.3	*	2.8	1.6	2.7	4.3	2.3	2.2	1.6	2.2
2003 4/.....	2.2	*	3.0	1.4	2.4	3.4	2.6	2.1	2.2	1.7
2002.....	2.8	*	3.4	2.8	3.5	4.3	3.7	3.1	2.0	2.8
2001.....	2.7	*	2.2	2.7	3.2	5.2	3.1	2.6	2.2	2.2
2000.....	2.7	*	2.8	2.7	3.2	5.2	3.3	2.3	1.8	2.3
1999.....	3.1	*	3.2	3.4	3.9	5.0	3.8	3.5	2.7	2.2
Black, female										
2005 4/.....	0.4	*	1.7	*	*	*	*	*	*	*
2004 4/.....	0.5	*	*	*	*	*	*	*	*	*
2003 4/.....	0.4	*	*	*	*	*	*	*	*	*
2002.....	0.5	*	*	*	*	*	*	*	*	*
2001.....	0.5	*	1.6	*	1.3	*	*	*	*	*
2000.....	0.6	*	1.7	*	*	*	*	*	*	*
1999.....	0.5	*	1.6	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Accidental exposure to smoke, fire and flames (X00-X09)										
All races 3/, both sexes										
2005.....	1.1	0.8	1.3	0.7	0.4	0.3	0.5	0.6	0.5	0.7
2004.....	1.1	0.6	1.4	0.9	0.4	0.3	0.6	0.7	0.5	0.7
2003.....	1.2	0.7	1.4	0.7	0.4	0.4	0.6	0.6	0.6	0.8
2002.....	1.1	0.9	1.4	0.8	0.5	0.4	0.5	0.6	0.7	0.7
2001.....	1.2	1.2	1.5	0.8	0.4	0.4	0.7	0.6	0.6	0.9
2000.....	1.2	1.0	1.9	0.9	0.4	0.4	0.6	0.6	0.6	0.8
1999.....	1.2	1.1	2.0	0.8	0.5	0.4	0.6	0.7	0.7	0.9
All races 3/, male										
2005.....	1.3	*	1.5	0.8	0.4	0.4	0.6	0.7	0.6	0.9
2004.....	1.3	*	1.5	1.1	0.4	0.4	0.6	0.9	0.7	0.9
2003.....	1.4	*	1.6	0.7	0.4	0.4	0.7	0.7	0.8	1.1
2002.....	1.4	1.0	1.6	0.9	0.6	0.6	0.7	0.8	0.9	0.9
2001.....	1.4	1.5	1.8	1.0	0.4	0.4	0.9	0.8	0.8	1.2
2000.....	1.5	1.0	2.2	1.0	0.3	0.4	0.8	0.8	0.8	1.2
1999.....	1.5	1.1	2.2	0.9	0.5	0.6	0.8	0.9	1.0	1.2
All races 3/, female										
2005.....	0.9	*	1.0	0.6	0.4	0.2	0.4	0.4	0.5	0.4
2004.....	0.9	*	1.3	0.6	0.4	0.2	0.5	0.4	0.3	0.4
2003.....	1.0	1.1	1.2	0.7	0.4	0.3	0.5	0.5	0.5	0.5
2002.....	0.8	*	1.2	0.6	0.4	0.3	0.4	0.4	0.6	0.4
2001.....	0.9	1.0	1.1	0.6	0.4	0.3	0.4	0.4	0.4	0.5
2000.....	0.9	*	1.5	0.7	0.5	0.4	0.4	0.4	0.4	0.5
1999.....	0.9	1.1	1.7	0.7	0.4	0.3	0.4	0.5	0.4	0.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Accidental exposure to smoke, fire and flames (X00-X09)										
White, both sexes										
2005 4/.....	1.0	0.7	1.0	0.5	0.3	0.2	0.5	0.5	0.5	0.6
2004 4/.....	1.0	*	1.1	0.6	0.3	0.3	0.5	0.6	0.5	0.7
2003 4/.....	1.1	0.7	1.2	0.5	0.2	0.3	0.6	0.6	0.7	0.8
2002.....	1.0	*	1.0	0.5	0.4	0.4	0.5	0.5	0.7	0.6
2001.....	1.0	1.1	1.2	0.7	0.4	0.3	0.7	0.5	0.6	0.8
2000.....	1.0	0.8	1.3	0.7	0.3	0.4	0.5	0.5	0.5	0.7
1999.....	1.0	0.9	1.5	0.6	0.4	0.4	0.6	0.7	0.7	0.8
White, male										
2005 4/.....	1.2	*	1.3	0.6	0.3	0.3	0.6	0.7	0.6	0.8
2004 4/.....	1.2	*	1.2	0.7	0.3	0.4	0.6	0.8	0.7	0.8
2003 4/.....	1.3	*	1.4	0.5	*	0.4	0.8	0.7	0.8	1.1
2002.....	1.2	*	1.1	0.6	0.5	0.5	0.6	0.7	0.9	0.8
2001.....	1.3	*	1.6	0.8	0.4	0.4	0.9	0.7	0.8	1.0
2000.....	1.3	*	1.6	0.7	0.3	0.4	0.8	0.7	0.7	1.0
1999.....	1.3	*	1.7	0.6	0.4	0.6	0.7	0.9	1.0	1.0
White, female										
2005 4/.....	0.8	*	0.7	0.4	0.3	*	0.4	0.4	0.3	0.4
2004 4/.....	0.8	*	1.0	0.5	0.3	0.3	0.5	0.4	0.4	0.5
2003 4/.....	0.9	*	1.0	0.5	0.3	0.3	0.5	0.5	0.5	0.5
2002.....	0.7	*	1.0	0.4	*	*	0.3	0.3	0.5	0.4
2001.....	0.8	*	0.8	0.5	0.3	0.3	0.4	*	0.4	0.5
2000.....	0.8	*	1.1	0.6	0.4	0.4	0.3	0.4	0.3	0.4
1999.....	0.8	*	1.3	0.6	0.3	0.3	0.4	0.4	0.3	0.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Accidental exposure to smoke, fire and flames (X00-X09)										
Black, both sexes										
2005 4/.....	1.9	*	2.8	1.6	0.9	*	*	0.9	1.0	1.1
2004 4/.....	2.1	*	2.8	2.3	0.9	*	0.8	1.2	0.8	0.9
2003 4/.....	2.0	*	3.0	1.9	1.0	0.6	*	0.9	0.8	1.0
2002.....	2.1	*	3.5	1.9	0.9	0.7	0.9	1.2	1.1	1.0
2001.....	2.3	*	3.3	1.8	0.8	*	1.0	1.4	1.0	1.5
2000.....	2.5	*	4.8	2.1	0.8	*	1.0	1.1	1.2	1.5
1999.....	2.5	*	4.5	2.1	0.9	*	0.9	1.1	0.9	1.9
Black, male										
2005 4/.....	2.4	*	2.8	1.7	*	*	*	*	*	*
2004 4/.....	2.6	*	2.9	3.1	*	*	*	1.6	*	*
2003 4/.....	2.4	*	3.2	1.9	*	*	*	*	*	*
2002.....	2.6	*	4.1	2.2	*	*	*	*	*	1.5
2001.....	2.9	*	3.6	2.3	*	*	*	*	*	2.5
2000.....	3.2	*	5.6	2.6	*	*	*	*	1.5	2.2
1999.....	3.0	*	5.0	2.4	*	*	*	*	*	2.8
Black, female										
2005 4/.....	1.5	*	2.7	1.6	*	*	*	*	*	*
2004 4/.....	1.6	*	2.7	1.4	*	*	*	*	*	*
2003 4/.....	1.7	*	2.7	1.8	*	*	*	*	*	*
2002.....	1.6	*	2.8	1.7	*	*	*	*	*	*
2001.....	1.7	*	2.9	1.2	*	*	*	1.5	*	*
2000.....	2.0	*	4.1	1.6	*	*	*	*	*	*
1999.....	2.0	*	3.9	1.8	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Accidental poisoning and exposure to noxious substances (X40-X49)

All races 3/, both sexes

2005.....	8.0	0.5	0.1	*	0.2	3.0	8.8	10.6	11.3	13.5
2004.....	7.1	*	*	*	0.2	3.1	7.7	8.9	9.3	12.8
2003.....	6.7	0.5	0.3	*	0.2	2.5	7.1	7.9	9.3	12.5
2002.....	6.1	0.6	0.2	*	0.1	2.4	5.9	6.9	8.6	11.9
2001.....	4.9	*	0.2	*	0.2	2.0	4.9	5.7	6.9	10.4
2000.....	4.5	*	0.2	*	0.1	1.7	4.3	5.1	6.8	9.5
1999.....	4.4	*	0.2	*	0.1	1.3	3.8	4.9	6.8	9.1

All races 3/, male

2005.....	10.9	*	*	*	0.2	4.5	13.4	15.5	15.7	18.0
2004.....	9.6	*	*	*	0.2	4.6	11.8	13.3	13.0	16.8
2003.....	9.2	*	0.3	*	0.2	3.8	10.9	11.7	13.1	16.8
2002.....	8.5	1.0	*	*	*	3.5	9.1	10.4	12.5	16.5
2001.....	7.1	*	0.3	*	*	3.0	7.6	8.7	9.9	14.6
2000.....	6.6	*	0.3	*	*	2.6	6.7	7.9	10.1	13.7
1999.....	6.5	*	*	*	*	1.9	5.9	7.7	10.3	13.3

All races 3/, female

2005.....	5.1	*	*	*	*	1.5	3.9	5.4	6.7	8.9
2004.....	4.7	*	*	*	0.2	1.5	3.4	4.4	5.4	8.7
2003.....	4.3	*	0.3	*	*	1.2	3.1	4.0	5.4	8.1
2002.....	3.7	*	*	*	*	1.2	2.6	3.3	4.7	7.2
2001.....	2.9	*	*	*	*	1.0	1.9	2.6	3.9	6.1
2000.....	2.5	*	*	*	*	0.8	1.7	2.1	3.5	5.3
1999.....	2.3	*	*	*	*	0.7	1.6	2.0	3.2	4.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Accidental poisoning and exposure to noxious substances (X40-X49)

White, both sexes

2005 4/.....	8.4	*	*	*	0.2	3.6	10.4	12.1	12.5	14.6
2004 4/.....	7.6	*	*	*	0.2	3.7	9.1	10.4	10.3	13.8
2003 4/.....	7.1	*	0.3	*	0.1	3.0	8.5	9.1	10.2	13.3
2002.....	6.3	*	*	*	0.2	2.8	6.9	7.8	9.5	12.5
2001.....	5.0	*	0.2	*	0.2	2.3	5.6	6.4	7.4	10.8
2000.....	4.6	*	*	*	0.1	2.0	4.9	5.5	7.2	9.8
1999.....	4.3	*	0.2	*	0.1	1.5	4.2	5.2	6.9	9.3

White, male

2005 4/.....	11.4	*	*	*	*	5.5	15.8	17.7	17.4	19.5
2004 4/.....	10.2	*	*	*	0.3	5.6	13.8	15.3	14.5	18.2
2003 4/.....	9.7	*	*	*	*	4.6	12.9	13.4	14.4	17.7
2002.....	8.8	*	*	*	*	4.1	10.6	11.6	13.6	17.3
2001.....	7.2	*	*	*	*	3.5	8.7	9.5	10.7	15.2
2000.....	6.7	*	*	*	*	3.1	7.8	8.6	10.7	14.3
1999.....	6.4	*	*	*	*	2.1	6.5	8.3	10.6	13.7

White, female

2005 4/.....	5.4	*	*	*	*	1.7	4.4	6.2	7.3	9.5
2004 4/.....	5.0	*	*	*	*	1.7	4.0	5.1	6.0	9.3
2003 4/.....	4.5	*	*	*	*	1.4	3.8	4.5	5.8	8.7
2002.....	3.9	*	*	*	*	1.4	3.0	3.8	5.1	7.5
2001.....	2.9	*	*	*	*	1.0	2.3	3.0	3.9	6.2
2000.....	2.5	*	*	*	*	0.9	1.8	2.3	3.6	5.2
1999.....	2.3	*	*	*	*	0.8	1.7	2.1	3.1	4.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Accidental poisoning and exposure to noxious substances (X40-X49)										
Black, both sexes										
2005 4/.....	7.6	*	*	*	*	0.7	3.0	5.6	8.8	11.6
2004 4/.....	6.5	*	*	*	*	0.9	2.9	4.5	6.2	11.0
2003 4/.....	6.3	*	*	*	*	*	2.3	4.0	7.3	11.8
2002.....	6.3	*	*	*	*	*	2.3	4.4	6.8	11.9
2001.....	5.8	*	*	*	*	0.9	2.3	4.0	6.3	11.1
2000.....	5.5	*	*	*	*	0.7	2.1	4.3	6.8	10.7
1999.....	5.7	*	*	*	*	*	2.8	4.2	8.0	10.3
Black, male										
2005 4/.....	10.8	*	*	*	*	*	4.2	8.6	12.0	15.5
2004 4/.....	8.9	*	*	*	*	*	4.4	6.5	8.7	13.5
2003 4/.....	8.8	*	*	*	*	*	3.5	5.7	10.2	16.7
2002.....	9.1	*	*	*	*	*	3.5	7.0	9.9	17.0
2001.....	8.4	*	*	*	*	*	3.8	6.7	8.2	15.4
2000.....	8.2	*	*	*	*	*	2.7	7.1	9.6	14.9
1999.....	8.7	*	*	*	*	*	4.7	6.7	11.5	14.9
Black, female										
2005 4/.....	4.8	*	*	*	*	*	1.7	2.8	5.9	8.1
2004 4/.....	4.3	*	*	*	*	*	1.4	2.5	4.0	8.8
2003 4/.....	3.9	*	*	*	*	*	*	2.4	4.8	7.4
2002.....	3.7	*	*	*	*	*	*	2.0	4.0	7.4
2001.....	3.5	*	*	*	*	*	*	1.6	4.6	7.3
2000.....	3.1	*	*	*	*	*	1.5	1.7	4.3	7.1
1999.....	3.0	*	*	*	*	*	*	2.0	4.9	6.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)

All races 3/, both sexes

2005.....	6.3	19.5	1.5	0.6	0.7	1.4	2.0	2.3	2.4	2.6
2004.....	5.8	19.2	1.6	0.6	0.6	1.2	1.9	1.9	2.2	2.6
2003.....	5.9	16.9	1.8	0.5	0.6	1.1	2.0	2.0	2.0	2.6
2002.....	6.0	16.8	1.6	0.6	0.8	1.3	1.8	2.0	2.1	2.6
2001.....	6.2	16.6	1.8	0.6	0.7	1.2	2.0	1.8	2.2	2.9
2000.....	6.2	15.0	1.7	0.6	0.7	1.4	2.0	2.1	2.3	3.0
1999.....	6.6	13.8	1.9	0.7	0.9	1.4	2.0	2.2	2.3	2.7

All races 3/, male

2005.....	7.4	21.1	1.8	0.8	1.0	2.3	3.3	3.8	4.0	4.2
2004.....	6.8	21.7	2.0	0.8	1.0	2.0	3.1	3.1	3.7	4.1
2003.....	6.9	19.5	2.2	0.6	0.9	1.8	3.3	3.3	3.4	4.0
2002.....	6.9	18.8	2.0	0.7	1.1	2.1	3.1	3.2	3.6	4.3
2001.....	7.2	18.9	2.1	0.7	1.0	1.9	3.4	3.1	3.7	4.8
2000.....	7.2	16.8	2.1	0.7	1.1	2.1	3.5	3.7	3.9	5.1
1999.....	7.6	16.2	2.3	0.9	1.2	2.3	3.4	3.7	3.7	4.5

All races 3/, female

2005.....	5.1	17.9	1.2	0.5	0.3	0.6	0.7	0.7	0.8	0.9
2004.....	4.8	16.5	1.1	0.4	0.3	0.4	0.5	0.6	0.7	1.0
2003.....	4.9	14.1	1.3	0.3	0.2	0.3	0.7	0.7	0.6	1.1
2002.....	5.0	14.6	1.2	0.4	0.4	0.5	0.5	0.7	0.6	1.0
2001.....	5.3	14.2	1.5	0.4	0.3	0.6	0.6	0.6	0.6	1.1
2000.....	5.2	13.2	1.4	0.4	0.3	0.5	0.4	0.6	0.7	0.8
1999.....	5.6	11.3	1.5	0.5	0.4	0.5	0.5	0.6	0.8	0.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)

White, both sexes

2005 4/.....	6.5	16.6	1.5	0.6	0.7	1.5	2.1	2.5	2.5	2.6
2004 4/.....	6.2	16.0	1.5	0.6	0.6	1.4	2.0	2.1	2.3	2.6
2003 4/.....	6.3	13.6	1.7	0.5	0.6	1.2	2.1	2.1	2.2	2.6
2002.....	6.3	12.9	1.4	0.5	0.8	1.4	2.0	2.2	2.3	2.7
2001.....	6.6	13.4	1.7	0.6	0.7	1.4	2.2	1.9	2.2	3.0
2000.....	6.6	12.2	1.6	0.6	0.8	1.4	2.1	2.3	2.4	3.0
1999.....	6.9	11.2	1.8	0.7	0.9	1.6	2.2	2.2	2.3	2.9

White, male

2005 4/.....	7.7	17.8	1.9	0.8	1.0	2.4	3.4	4.1	4.2	4.4
2004 4/.....	7.2	19.0	2.0	0.7	1.0	2.2	3.4	3.4	3.8	4.2
2003 4/.....	7.3	15.7	2.2	0.6	0.9	2.0	3.6	3.6	3.6	4.1
2002.....	7.3	15.3	1.8	0.7	1.0	2.4	3.4	3.6	3.8	4.3
2001.....	7.5	15.2	2.0	0.7	1.0	2.1	3.7	3.2	3.7	5.0
2000.....	7.4	14.0	2.0	0.7	1.2	2.2	3.7	3.8	4.1	5.2
1999.....	7.8	12.9	2.1	1.0	1.3	2.5	3.7	3.9	3.8	4.8

White, female

2005 4/.....	5.4	15.3	1.1	0.5	0.3	0.5	0.6	0.7	0.8	0.8
2004 4/.....	5.2	12.8	0.9	0.4	0.3	0.4	0.5	0.7	0.8	1.0
2003 4/.....	5.3	11.4	1.2	0.3	*	0.4	0.6	0.6	0.7	1.1
2002.....	5.4	10.5	1.1	0.4	0.5	0.5	0.5	0.7	0.6	1.0
2001.....	5.8	11.5	1.4	0.5	0.3	0.6	0.6	0.6	0.6	1.0
2000.....	5.7	10.3	1.2	0.4	0.3	0.5	0.4	0.6	0.7	0.8
1999.....	6.1	9.5	1.4	0.4	0.5	0.6	0.5	0.6	0.7	0.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)

Black, both sexes

2005 4/.....	6.0	37.4	1.8	*	0.7	1.3	1.8	1.8	2.3	2.7
2004 4/.....	4.8	37.1	1.8	*	0.6	0.6	1.4	1.6	2.2	2.8
2003 4/.....	4.9	34.9	2.0	*	*	0.7	1.9	2.0	1.6	2.7
2002.....	5.2	37.5	2.4	0.7	0.9	1.0	1.2	1.5	1.9	3.0
2001.....	5.4	34.2	2.5	*	0.7	0.8	1.3	1.8	1.9	2.6
2000.....	5.4	30.6	2.3	0.7	0.7	1.3	1.8	2.0	2.4	3.1
1999.....	6.1	27.9	2.5	0.7	0.7	1.0	1.6	2.2	2.6	2.2

Black, male

2005 4/.....	7.6	41.0	1.9	*	*	2.0	2.8	2.9	3.3	4.3
2004 4/.....	6.1	38.3	1.7	*	*	*	2.2	2.8	3.8	4.6
2003 4/.....	6.4	40.5	2.8	*	*	*	2.6	3.0	2.9	4.6
2002.....	6.6	38.9	3.3	*	1.5	1.5	1.9	2.3	3.4	5.2
2001.....	6.8	40.4	2.8	*	*	*	2.0	2.8	3.6	3.7
2000.....	7.3	32.0	2.5	*	1.2	1.8	3.2	3.5	3.8	5.5
1999.....	8.0	33.2	3.4	*	1.2	1.8	3.0	3.5	4.4	3.5

Black, female

2005 4/.....	4.5	33.6	1.6	*	*	*	*	*	*	1.4
2004 4/.....	3.6	35.8	2.0	*	*	*	*	*	*	*
2003 4/.....	3.6	29.1	*	*	*	*	1.3	*	*	*
2002.....	4.0	36.0	*	*	*	*	*	*	*	*
2001.....	4.1	27.7	2.2	*	*	*	*	*	*	1.5
2000.....	3.8	29.1	2.0	*	*	*	*	*	*	*
1999.....	4.4	22.5	1.6	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)

All races 3/, both sexes

2005.....	11.0	*	1.3	7.7	12.4	11.8	13.0	13.8
2004.....	11.0	*	1.3	8.2	12.5	12.3	13.0	14.0
2003.....	10.8	*	1.2	7.3	12.1	11.9	13.5	13.8
2002.....	11.0	*	1.2	7.4	12.4	12.8	12.5	14.3
2001 5/.....	10.8	*	1.3	7.9	12.0	12.6	13.0	14.3
2000.....	10.4	*	1.5	8.0	12.5	12.1	12.0	13.7
1999.....	10.5	*	1.2	8.0	12.3	12.3	13.1	14.4

All races 3/, male

2005.....	17.7	*	1.9	12.1	20.2	19.2	20.6	21.4
2004.....	17.7	*	1.7	12.6	20.8	20.2	20.6	21.7
2003.....	17.6	*	1.7	11.6	20.2	19.5	21.6	21.9
2002.....	17.9	*	1.8	12.2	20.8	20.8	20.2	22.7
2001 5/.....	17.6	*	1.9	12.9	20.5	21.0	21.0	22.7
2000.....	17.1	*	2.3	13.0	21.4	20.0	19.2	21.7
1999.....	17.1	*	1.9	13.0	20.8	20.3	21.1	22.7

All races 3/, female

2005.....	4.5	*	0.7	3.0	4.0	4.1	5.3	6.1
2004.....	4.6	*	1.0	3.5	3.6	4.2	5.3	6.3
2003.....	4.3	*	0.5	2.7	3.4	4.0	5.2	5.7
2002.....	4.3	*	0.6	2.4	3.5	4.5	4.7	6.0
2001 5/.....	4.1	*	0.6	2.7	3.1	4.0	4.8	5.9
2000.....	4.0	*	0.6	2.7	3.2	4.0	4.6	5.8
1999.....	4.0	*	0.5	2.7	3.4	4.1	5.1	6.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)										
White, both sexes										
2005 4/.....	12.3	*	1.3	8.3	13.1	12.5	14.3	15.4
2004 4/.....	12.3	*	1.3	8.8	13.2	13.0	14.2	15.6
2003 4/.....	12.1	*	1.1	7.9	12.5	12.6	14.7	15.3
2002.....	12.2	*	1.3	8.2	13.2	13.5	13.7	15.9
2001 5/.....	11.9	*	1.3	8.6	12.6	13.5	14.2	15.8
2000.....	11.5	*	1.5	8.5	13.1	12.7	13.0	15.1
1999.....	11.5	*	1.3	8.5	12.7	12.8	14.2	15.7
White, male										
2005 4/.....	19.7	*	1.9	13.3	21.3	20.3	22.2	23.7
2004 4/.....	19.6	*	1.7	13.6	22.0	21.2	22.3	24.0
2003 4/.....	19.5	*	1.7	12.7	21.1	20.3	23.3	23.9
2002.....	19.9	*	1.9	13.4	22.0	21.4	21.7	24.7
2001 5/.....	19.5	*	2.0	14.0	21.3	22.2	22.6	24.7
2000.....	18.8	*	2.4	13.7	22.3	20.8	20.7	23.6
1999.....	18.7	*	2.0	13.7	21.5	21.0	22.5	24.3
White, female										
2005 4/.....	5.0	*	0.6	3.1	4.3	4.2	6.0	6.8
2004 4/.....	5.1	*	0.9	3.7	3.8	4.3	5.6	7.0
2003 4/.....	4.7	*	0.5	2.9	3.3	4.3	5.7	6.5
2002.....	4.8	*	0.7	2.6	3.7	5.1	5.3	6.8
2001 5/.....	4.6	*	0.6	2.9	3.3	4.3	5.4	6.7
2000.....	4.4	*	0.7	2.9	3.3	4.1	5.1	6.5
1999.....	4.5	*	0.5	2.9	3.5	4.3	5.5	6.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)										
Black, both sexes										
2005 4/.....	5.1	*	1.1	4.4	9.1	8.8	8.5	7.2
2004 4/.....	5.2	*	1.4	4.7	9.8	10.1	8.7	7.0
2003 4/.....	5.1	*	1.2	3.8	10.5	9.5	9.1	7.5
2002.....	5.1	*	1.0	4.0	9.1	10.3	8.2	8.1
2001 5/.....	5.3	*	1.1	4.3	10.1	9.6	8.6	7.8
2000.....	5.4	*	1.2	5.5	11.0	9.9	7.6	7.3
1999.....	5.4	*	0.9	5.8	10.7	9.6	8.7	8.4
Black, male										
2005 4/.....	8.7	*	1.5	7.2	16.2	14.9	15.0	12.0
2004 4/.....	9.0	*	1.6	7.4	17.2	18.0	14.4	11.7
2003 4/.....	8.8	*	1.9	6.6	18.0	17.4	15.7	12.9
2002.....	9.1	*	1.6	6.9	16.1	19.5	15.2	14.5
2001 5/.....	9.2	*	1.7	7.3	19.4	17.7	15.4	13.8
2000.....	9.4	*	2.1	9.5	19.5	17.5	13.6	13.0
1999.....	9.6	*	1.3	9.9	19.5	17.9	15.7	15.6
Black, female										
2005 4/.....	1.8	*	*	1.4	2.0	3.0	2.7	2.8
2004 4/.....	1.8	*	*	1.9	2.4	2.8	3.6	2.7
2003 4/.....	1.8	*	*	*	3.1	2.2	3.1	2.6
2002.....	1.5	*	*	*	2.3	2.0	2.0	2.5
2001 5/.....	1.7	*	*	1.3	*	2.2	2.5	2.5
2000.....	1.7	*	*	1.4	3.0	3.0	2.2	2.2
1999.....	1.6	*	*	1.6	2.3	2.1	2.5	2.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Intentional self-harm (suicide) by discharge of firearms (X72-X74)										
All races 3/, both sexes										
2005.....	5.7	*	0.4	3.5	5.8	5.6	5.7	5.9
2004.....	5.7	*	0.3	3.8	6.3	5.7	5.7	5.9
2003.....	5.8	*	0.3	3.6	6.5	5.8	6.1	6.0
2002.....	5.9	*	0.4	3.6	6.7	6.4	5.6	6.6
2001.....	5.9	*	0.4	4.1	6.6	6.7	6.3	6.4
2000.....	5.9	*	0.5	4.4	7.2	6.3	6.0	6.7
1999.....	5.9	*	0.5	4.9	7.2	6.6	6.5	7.1
All races 3/, male										
2005.....	10.2	*	0.7	6.1	10.3	9.7	9.7	10.1
2004.....	10.0	*	0.5	6.5	11.1	9.9	9.6	9.8
2003.....	10.4	*	0.6	6.3	11.4	10.1	10.6	10.2
2002.....	10.6	*	0.6	6.4	11.9	11.1	9.6	11.3
2001.....	10.6	*	0.6	7.1	11.8	11.7	10.8	10.8
2000.....	10.5	*	0.9	7.6	12.9	11.1	10.1	11.1
1999.....	10.6	*	0.8	8.4	12.8	11.5	10.9	11.8
All races 3/, female										
2005.....	1.4	*	*	0.8	1.1	1.3	1.6	1.7
2004.....	1.5	*	*	1.0	1.1	1.3	1.7	1.9
2003.....	1.4	*	*	0.8	1.2	1.3	1.5	1.8
2002.....	1.4	*	*	0.7	1.2	1.6	1.7	2.0
2001.....	1.5	*	0.2	1.0	1.0	1.5	1.7	2.1
2000.....	1.5	*	0.2	1.1	1.3	1.5	1.9	2.3
1999.....	1.5	*	0.2	1.1	1.4	1.6	1.9	2.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Intentional self-harm (suicide) by discharge of firearms (X72-X74)										
White, both sexes										
2005 4/.....	6.5	*	0.5	3.8	6.1	6.0	6.3	6.7
2004 4/.....	6.5	*	0.3	4.2	6.6	6.0	6.3	6.6
2003 4/.....	6.6	*	0.4	4.0	6.7	6.1	6.7	6.8
2002.....	6.7	*	0.4	3.9	7.0	6.8	6.2	7.4
2001.....	6.7	*	0.4	4.5	6.7	7.2	7.0	7.2
2000.....	6.6	*	0.6	4.6	7.6	6.6	6.6	7.5
1999.....	6.6	*	0.6	5.2	7.4	7.0	7.1	7.8
White, male										
2005 4/.....	11.5	*	0.8	6.6	10.7	10.3	10.5	11.2
2004 4/.....	11.3	*	0.5	7.1	11.7	10.2	10.4	11.0
2003 4/.....	11.7	*	0.7	6.9	11.8	10.5	11.3	11.3
2002.....	11.9	*	0.7	6.9	12.3	11.3	10.4	12.4
2001.....	11.8	*	0.7	7.6	11.9	12.3	11.7	11.9
2000.....	11.6	*	1.0	7.9	13.5	11.4	11.0	12.2
1999.....	11.7	*	0.9	8.9	13.0	11.9	11.8	12.7
White, female										
2005 4/.....	1.6	*	*	0.9	1.2	1.4	1.9	2.0
2004 4/.....	1.7	*	*	1.1	1.2	1.5	2.0	2.1
2003 4/.....	1.6	*	*	0.9	1.2	1.5	1.8	2.1
2002.....	1.6	*	*	0.8	1.3	2.0	1.9	2.3
2001.....	1.7	*	*	1.1	1.1	1.7	2.0	2.4
2000.....	1.7	*	*	1.1	1.3	1.6	2.0	2.7
1999.....	1.7	*	0.3	1.2	1.4	1.8	2.2	2.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Intentional self-harm (suicide) by discharge of firearms (X72-X74)										
Black, both sexes										
2005 4/.....	2.6	*	*	2.4	5.3	4.6	4.2	3.5
2004 4/.....	2.6	*	*	2.2	5.9	5.4	4.2	2.9
2003 4/.....	2.6	*	*	2.1	6.5	5.5	4.8	2.8
2002.....	2.8	*	*	2.5	5.8	6.1	4.0	4.0
2001.....	2.9	*	*	2.8	7.1	5.8	4.1	3.7
2000.....	3.0	*	*	3.8	6.8	6.0	4.0	3.4
1999.....	3.1	*	*	3.9	7.6	5.7	4.2	4.2
Black, male										
2005 4/.....	4.9	*	*	4.3	9.8	8.4	8.3	6.7
2004 4/.....	4.9	*	*	3.7	10.7	10.2	7.8	5.5
2003 4/.....	5.0	*	*	3.9	11.9	11.0	9.1	5.5
2002.....	5.4	*	*	4.3	11.0	12.4	7.7	7.4
2001.....	5.5	*	*	5.0	13.9	10.9	8.0	7.1
2000.....	5.5	*	*	6.9	12.6	11.3	7.3	6.4
1999.....	5.9	*	*	7.0	14.2	11.2	8.1	8.1
Black, female										
2005 4/.....	0.5	*	*	*	*	*	*	*
2004 4/.....	0.5	*	*	*	*	*	*	*
2003 4/.....	0.5	*	*	*	*	*	*	*
2002.....	0.5	*	*	*	*	*	*	*
2001.....	0.6	*	*	*	*	*	*	*
2000.....	0.7	*	*	*	*	*	*	*
1999.....	0.6	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)

All races 3/, both sexes

2005.....	5.3	*	0.9	4.2	6.5	6.2	7.3	7.8
2004.....	5.3	*	1.1	4.4	6.2	6.6	7.3	8.1
2003.....	5.0	*	0.8	3.7	5.6	6.1	7.4	7.8
2002.....	5.0	*	0.8	3.8	5.7	6.4	6.9	7.7
2001 5/.....	4.8	*	0.9	3.8	5.4	5.9	6.7	7.9
2000.....	4.5	*	0.9	3.6	5.3	5.8	6.0	7.0
1999.....	4.5	*	0.7	3.2	5.1	5.7	6.7	7.3

All races 3/, male

2005.....	7.5	*	1.2	6.0	10.0	9.5	10.9	11.3
2004.....	7.6	*	1.2	6.2	9.7	10.3	11.0	11.9
2003.....	7.3	*	1.2	5.4	8.8	9.3	11.0	11.7
2002.....	7.3	*	1.2	5.8	8.9	9.7	10.6	11.4
2001 5/.....	7.1	*	1.3	5.8	8.7	9.3	10.2	11.9
2000.....	6.6	*	1.4	5.4	8.5	8.9	9.1	10.6
1999.....	6.6	*	1.1	4.6	8.0	8.8	10.2	10.9

All races 3/, female

2005.....	3.1	*	0.6	2.2	2.9	2.8	3.6	4.4
2004.....	3.1	*	0.9	2.5	2.4	2.8	3.5	4.4
2003.....	2.8	*	0.4	1.9	2.2	2.7	3.7	4.0
2002.....	2.9	*	0.4	1.6	2.3	2.9	3.1	4.0
2001 5/.....	2.6	*	0.4	1.7	2.0	2.5	3.1	3.9
2000.....	2.5	*	0.4	1.7	1.9	2.6	2.7	3.4
1999.....	2.5	*	0.3	1.6	2.0	2.5	3.2	3.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)										
White, both sexes										
2005 4/.....	5.8	*	0.8	4.5	7.0	6.5	8.0	8.8
2004 4/.....	5.8	*	1.0	4.6	6.6	7.0	7.9	9.0
2003 4/.....	5.5	*	0.8	4.0	5.8	6.4	8.0	8.5
2002.....	5.5	*	0.8	4.2	6.2	6.8	7.4	8.5
2001 5/.....	5.3	*	0.9	4.2	5.9	6.3	7.2	8.6
2000.....	4.9	*	0.9	3.8	5.5	6.1	6.4	7.6
1999.....	4.9	*	0.7	3.3	5.4	5.9	7.1	7.9
White, male										
2005 4/.....	8.2	*	1.1	6.7	10.6	9.9	11.8	12.5
2004 4/.....	8.3	*	1.2	6.5	10.4	10.9	11.9	13.0
2003 4/.....	7.9	*	1.1	5.8	9.3	9.8	11.9	12.6
2002.....	8.0	*	1.2	6.5	9.7	10.2	11.2	12.3
2001 5/.....	7.7	*	1.3	6.4	9.4	9.8	10.8	12.8
2000.....	7.1	*	1.3	5.8	8.8	9.4	9.7	11.4
1999.....	7.0	*	1.1	4.9	8.5	9.1	10.7	11.6
White, female										
2005 4/.....	3.4	*	0.5	2.2	3.2	2.8	4.0	4.8
2004 4/.....	3.4	*	0.8	2.7	2.6	2.8	3.6	4.9
2003 4/.....	3.1	*	0.4	2.0	2.2	2.8	4.0	4.3
2002.....	3.2	*	0.5	1.8	2.5	3.1	3.4	4.5
2001 5/.....	2.9	*	0.4	1.8	2.2	2.6	3.4	4.3
2000.....	2.8	*	0.5	1.8	1.9	2.5	3.1	3.8
1999.....	2.8	*	0.3	1.7	2.1	2.5	3.3	4.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)										
Black, both sexes										
2005 4/.....	2.5	*	1.0	2.0	3.9	4.1	4.3	3.7
2004 4/.....	2.6	*	1.1	2.5	3.9	4.8	4.5	4.0
2003 4/.....	2.5	*	1.0	1.7	4.0	4.0	4.3	4.6
2002.....	2.3	*	0.7	1.6	3.2	4.2	4.2	4.1
2001 5/.....	2.3	*	0.8	1.5	3.0	3.8	4.5	4.1
2000.....	2.4	*	1.1	1.8	4.2	3.9	3.6	3.8
1999.....	2.3	*	0.7	1.9	3.1	3.9	4.6	4.2
Black, male										
2005 4/.....	3.8	*	1.2	2.9	6.4	6.5	6.7	5.3
2004 4/.....	4.1	*	1.2	3.7	6.5	7.8	6.5	6.2
2003 4/.....	3.8	*	1.6	2.6	6.1	6.4	6.7	7.4
2002.....	3.7	*	1.1	2.6	5.0	7.0	7.5	7.0
2001 5/.....	3.7	*	1.1	2.3	5.5	6.8	7.4	6.7
2000.....	3.9	*	1.8	2.6	6.8	6.2	6.4	6.6
1999.....	3.7	*	*	2.9	5.3	6.7	7.6	7.5
Black, female										
2005 4/.....	1.3	*	*	*	1.3	1.9	2.1	2.3
2004 4/.....	1.3	*	*	*	1.3	2.0	2.6	2.0
2003 4/.....	1.3	*	*	*	1.9	1.8	2.2	2.2
2002.....	1.1	*	*	*	1.5	1.6	1.4	1.6
2001 5/.....	1.1	*	*	*	*	*	1.8	1.8
2000.....	1.0	*	*	*	1.7	1.8	*	1.4
1999.....	1.0	*	*	*	*	*	1.9	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)

All races 3/, both sexes

2005.....	6.1	7.5	2.3	0.6	1.1	9.9	16.1	13.6	10.0	7.5
2004.....	5.9	8.0	2.4	0.6	1.0	9.3	15.0	13.3	9.3	7.3
2003.....	6.1	8.5	2.4	0.6	1.0	9.5	16.5	13.4	9.4	7.6
2002.....	6.1	7.5	2.7	0.7	1.0	9.3	16.5	13.3	9.4	7.9
2001 5/.....	7.1	8.2	2.7	0.7	0.9	9.4	17.3	14.9	11.5	10.4
2000.....	6.0	9.2	2.3	0.7	1.1	9.5	16.0	11.9	9.0	7.6
1999.....	6.1	8.7	2.5	0.9	1.2	10.4	15.6	12.1	9.1	7.7

All races 3/, male

2005.....	9.8	8.2	2.6	0.7	1.3	16.8	27.2	22.8	16.4	11.2
2004.....	9.4	8.0	2.5	0.6	1.3	15.3	25.4	22.3	14.9	11.2
2003.....	9.7	10.0	2.5	0.7	1.3	15.9	27.6	22.4	14.8	11.4
2002.....	9.6	7.9	2.9	0.6	1.2	15.3	27.7	22.0	14.7	11.8
2001 5/.....	11.1	9.5	3.0	0.7	1.1	15.7	28.9	24.1	17.7	15.3
2000.....	9.3	10.4	2.5	0.7	1.4	15.5	26.7	19.6	14.0	10.9
1999.....	9.3	9.6	2.5	0.9	1.5	16.9	25.7	19.5	13.8	11.0

All races 3/, female

2005.....	2.5	6.6	2.0	0.6	0.8	2.5	4.3	4.0	3.5	3.9
2004.....	2.5	7.9	2.2	0.6	0.7	3.0	4.0	3.9	3.6	3.3
2003.....	2.6	6.9	2.3	0.5	0.6	2.6	4.8	4.0	3.9	3.9
2002.....	2.7	7.1	2.5	0.8	0.8	2.9	4.6	4.2	4.1	4.1
2001 5/.....	3.3	6.9	2.4	0.6	0.7	2.7	5.1	5.5	5.2	5.5
2000.....	2.8	7.9	2.1	0.6	0.8	3.1	4.7	4.1	4.0	4.4
1999.....	2.9	7.8	2.4	0.9	0.9	3.5	5.1	4.5	4.3	4.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)										
White, both sexes										
2005 4/.....	3.7	6.1	1.7	0.4	0.6	5.3	7.9	7.2	5.6	4.7
2004 4/.....	3.6	6.5	1.6	0.5	0.7	5.1	7.9	7.0	5.2	4.6
2003 4/.....	3.7	6.9	1.7	0.4	0.6	5.0	8.4	6.9	5.3	5.0
2002.....	3.7	5.4	2.0	0.5	0.7	5.2	8.2	6.8	5.1	5.0
2001 5/.....	4.9	6.2	1.9	0.7	0.7	5.1	9.4	8.9	7.8	7.8
2000.....	3.6	6.6	1.7	0.6	0.9	5.2	7.7	6.2	5.3	4.9
1999.....	3.8	6.7	1.7	0.7	1.0	5.6	8.0	6.5	5.5	5.1
White, male										
2005 4/.....	5.4	6.7	1.8	0.5	0.7	8.6	12.5	11.1	8.4	6.5
2004 4/.....	5.3	6.9	1.8	0.5	0.9	7.9	12.5	10.9	7.7	6.8
2003 4/.....	5.4	8.1	1.5	0.4	0.9	8.0	13.3	10.7	7.8	6.9
2002.....	5.4	6.2	2.1	0.4	0.7	8.2	12.9	10.5	7.3	7.0
2001 5/.....	7.2	7.3	2.2	0.6	0.8	7.9	14.5	13.5	11.3	11.6
2000.....	5.2	8.2	1.9	0.7	1.1	8.0	12.0	9.2	7.6	6.6
1999.....	5.5	7.9	1.8	0.7	1.1	8.6	12.1	9.9	7.8	7.0
White, female										
2005 4/.....	1.9	5.5	1.6	0.4	0.6	1.8	2.9	3.0	2.7	2.9
2004 4/.....	1.9	6.2	1.4	0.5	0.5	2.1	3.0	2.9	2.6	2.4
2003 4/.....	2.0	5.7	1.8	0.4	0.3	1.8	3.2	2.9	2.6	3.0
2002.....	2.0	4.6	1.8	0.6	0.6	1.9	3.1	2.8	2.8	3.1
2001 5/.....	2.6	5.1	1.6	0.7	0.6	2.0	3.9	4.1	4.2	3.9
2000.....	2.1	5.0	1.5	0.5	0.7	2.1	3.2	3.0	2.9	3.2
1999.....	2.1	5.4	1.7	0.7	0.8	2.3	3.7	2.9	3.2	3.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)										
Black, both sexes										
2005 4/.....	22.2	14.3	5.3	1.4	3.0	33.5	61.2	53.0	39.2	26.5
2004 4/.....	21.1	15.6	6.3	1.4	2.4	31.6	55.9	52.8	35.6	25.2
2003 4/.....	22.0	16.1	6.1	1.7	2.7	33.2	62.4	54.5	36.3	25.4
2002.....	22.0	17.3	6.3	1.4	2.6	31.0	63.4	54.2	37.0	26.9
2001 5/.....	22.1	18.9	6.8	0.9	1.9	32.7	62.8	52.6	35.3	28.2
2000.....	21.5	22.8	5.8	1.2	2.5	33.1	64.1	48.2	33.5	25.9
1999.....	21.1	19.5	6.1	1.9	2.5	36.5	60.4	47.0	31.8	25.1
Black, male										
2005 4/.....	39.7	15.9	6.6	1.6	4.1	60.0	110.1	98.6	72.8	44.7
2004 4/.....	37.1	14.2	6.3	1.4	3.3	54.8	101.8	98.4	64.7	43.1
2003 4/.....	38.9	17.8	7.4	2.0	3.7	58.9	111.6	101.6	63.9	43.7
2002.....	38.4	16.3	7.1	1.4	3.5	53.3	115.3	99.9	65.4	45.2
2001 5/.....	38.3	21.0	8.0	1.2	2.7	58.2	116.3	96.0	62.3	42.8
2000.....	37.2	23.3	5.9	*	3.3	57.2	117.8	89.4	58.6	42.5
1999.....	36.1	19.5	5.9	2.0	3.3	62.4	110.9	84.0	54.7	40.1
Black, female										
2005 4/.....	6.2	12.6	4.0	1.3	1.9	6.3	11.5	9.9	8.8	10.1
2004 4/.....	6.4	17.0	6.3	1.4	1.5	7.8	9.8	10.2	9.3	9.2
2003 4/.....	6.6	14.2	4.8	1.3	1.7	6.7	13.6	10.9	11.4	9.0
2002.....	7.0	18.5	5.5	1.3	1.6	8.0	12.7	12.3	11.5	10.6
2001 5/.....	7.4	16.7	5.6	*	*	6.4	11.4	13.1	11.1	15.3
2000.....	7.2	22.2	5.7	1.4	1.7	8.4	13.2	10.9	10.9	11.1
1999.....	7.6	19.5	6.3	1.8	1.6	10.0	12.5	13.6	11.2	11.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)										
All races 3/, both sexes										
2005.....	4.2	*	0.2	0.2	0.7	8.3	13.1	11.0	7.8	5.1
2004.....	4.0	*	0.2	0.2	0.7	7.6	12.2	10.5	7.0	4.9
2003.....	4.1	*	0.3	0.2	0.7	7.7	13.6	10.8	7.1	5.0
2002.....	4.1	*	0.3	0.3	0.7	7.7	13.6	10.6	6.9	5.2
2001.....	4.0	*	0.4	0.3	0.6	7.5	13.6	10.3	6.6	5.1
2000.....	3.8	*	0.2	0.2	0.7	7.7	12.7	9.2	6.3	4.7
1999.....	3.9	*	0.3	0.3	0.8	8.5	12.5	9.2	6.1	4.6
All races 3/, male										
2005.....	7.2	*	0.3	0.3	1.0	14.7	23.1	19.3	13.5	8.1
2004.....	6.9	*	*	0.2	1.0	13.1	21.5	18.8	11.9	8.3
2003.....	7.1	*	0.2	0.3	1.0	13.7	23.8	19.1	12.0	8.0
2002.....	7.0	*	0.4	0.3	0.9	13.2	23.8	18.6	11.6	8.4
2001.....	6.8	*	0.3	0.3	0.8	13.2	24.0	17.8	11.1	8.0
2000.....	6.5	*	*	0.3	1.0	13.3	22.4	16.1	10.4	7.4
1999.....	6.5	*	0.3	0.4	1.1	14.5	21.8	15.9	10.1	7.1
All races 3/, female										
2005.....	1.2	*	*	*	0.4	1.5	2.4	2.3	2.1	2.1
2004.....	1.1	*	*	0.2	0.3	1.8	2.2	2.0	2.0	1.4
2003.....	1.2	*	0.3	0.2	0.3	1.5	2.8	2.2	2.1	1.9
2002.....	1.3	*	*	0.3	0.5	1.8	2.9	2.4	2.1	2.0
2001.....	1.3	*	0.4	0.2	0.3	1.5	2.7	2.5	2.1	2.2
2000.....	1.3	*	*	0.2	0.3	1.7	2.7	2.2	2.1	1.9
1999.....	1.3	*	0.3	0.2	0.5	2.2	2.8	2.3	2.1	2.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Assault (homicide) by discharge of firearms (*U01.4,X93-X95)

White, both sexes

2005 4/.....	2.2	*	0.2	0.2	0.4	4.1	5.5	5.1	4.0	3.0
2004 4/.....	2.1	*	*	0.2	0.4	3.8	5.5	4.9	3.7	2.9
2003 4/.....	2.2	*	0.2	0.1	0.4	3.8	6.3	4.9	3.6	3.0
2002.....	2.2	*	0.3	0.2	0.4	4.1	6.0	4.9	3.4	3.0
2001.....	2.2	*	0.3	0.3	0.4	3.7	6.3	5.0	3.5	3.2
2000.....	2.1	*	*	0.2	0.5	3.9	5.5	4.2	3.3	2.8
1999.....	2.2	*	0.2	0.2	0.6	4.2	5.6	4.3	3.4	3.0

White, male

2005 4/.....	3.5	*	*	*	0.5	7.0	9.3	8.4	6.3	4.2
2004 4/.....	3.5	*	*	*	0.6	6.3	9.3	8.3	5.8	4.7
2003 4/.....	3.5	*	*	*	0.6	6.4	10.6	8.2	5.7	4.4
2002.....	3.5	*	0.3	*	0.5	6.7	10.0	8.1	5.3	4.5
2001.....	3.5	*	*	0.3	0.6	6.2	10.5	8.2	5.4	4.9
2000.....	3.2	*	*	0.3	0.7	6.5	9.0	6.7	5.1	4.1
1999.....	3.4	*	*	0.2	0.8	6.9	9.1	7.1	5.2	4.3

White, female

2005 4/.....	0.9	*	*	*	*	1.0	1.4	1.6	1.5	1.7
2004 4/.....	0.9	*	*	*	*	1.2	1.5	1.2	1.4	1.0
2003 4/.....	0.9	*	*	*	*	1.0	1.7	1.4	1.4	1.6
2002.....	1.0	*	*	*	0.3	1.2	1.6	1.6	1.4	1.5
2001.....	1.0	*	*	*	0.3	1.1	1.9	1.6	1.6	1.6
2000.....	1.0	*	*	*	*	1.1	1.7	1.6	1.5	1.5
1999.....	0.9	*	*	*	0.4	1.2	1.9	1.4	1.5	1.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Assault (homicide) by discharge of firearms (*U01.4,X93-X95)

Black, both sexes

2005 4/.....	17.2	*	*	*	2.1	30.0	55.1	46.9	33.5	19.4
2004 4/.....	16.1	*	0.8	*	1.8	27.8	50.4	46.3	29.3	18.5
2003 4/.....	16.8	*	*	0.7	2.0	29.0	55.3	47.9	30.1	18.1
2002.....	16.7	*	*	*	1.9	27.3	57.0	46.9	30.1	19.5
2001.....	15.8	*	*	*	1.3	28.7	55.7	43.6	26.9	18.4
2000.....	15.6	*	*	*	1.7	28.6	55.3	40.7	26.0	17.3
1999.....	15.4	*	*	0.7	1.8	32.0	53.4	39.3	23.7	15.9

Black, male

2005 4/.....	32.5	*	*	*	3.1	54.9	101.6	89.5	64.2	35.7
2004 4/.....	30.3	*	*	*	2.9	49.8	93.9	88.9	55.2	34.6
2003 4/.....	31.6	*	*	*	2.9	53.2	101.7	92.4	56.4	33.9
2002.....	31.0	*	*	*	2.7	48.4	105.9	90.2	56.3	35.7
2001.....	29.8	*	*	*	2.1	52.8	106.5	83.2	51.3	31.5
2000.....	29.2	*	*	*	2.8	51.7	105.2	78.7	48.1	31.7
1999.....	28.4	*	*	*	2.8	56.3	101.0	74.4	44.0	27.3

Black, female

2005 4/.....	3.1	*	*	*	*	4.4	7.6	6.6	5.8	4.7
2004 4/.....	3.0	*	*	*	*	5.1	6.6	6.4	5.9	4.0
2003 4/.....	3.3	*	*	*	1.1	4.0	9.1	6.8	6.3	4.1
2002.....	3.6	*	*	*	1.2	5.4	9.2	7.3	6.5	5.0
2001.....	3.1	*	*	*	*	3.9	7.0	7.6	5.0	6.8
2000.....	3.2	*	*	*	*	4.9	8.0	6.2	6.0	4.5
1999.....	3.7	*	*	*	*	7.0	8.2	7.5	5.5	5.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)										
All races 3/, both sexes										
2005.....	1.9	7.3	2.1	0.4	0.4	1.6	3.0	2.6	2.2	2.5
2004.....	2.0	7.8	2.1	0.4	0.3	1.7	2.9	2.7	2.2	2.4
2003.....	2.0	8.3	2.1	0.4	0.3	1.7	2.9	2.6	2.3	2.7
2002.....	2.0	7.3	2.4	0.4	0.3	1.6	2.9	2.6	2.5	2.7
2001 5/.....	3.1	8.0	2.3	0.4	0.3	1.8	3.7	4.6	4.9	5.2
2000.....	2.1	8.9	2.1	0.4	0.5	1.8	3.2	2.7	2.8	2.9
1999.....	2.2	8.5	2.1	0.6	0.4	1.9	3.1	2.9	3.0	3.0
All races 3/, male										
2005.....	2.6	7.9	2.3	0.4	0.3	2.1	4.0	3.5	3.0	3.1
2004.....	2.5	7.9	2.3	0.4	0.3	2.2	4.0	3.5	2.9	2.9
2003.....	2.6	9.7	2.3	0.4	0.4	2.3	3.8	3.3	2.8	3.4
2002.....	2.6	7.8	2.6	0.4	0.3	2.1	3.9	3.4	3.0	3.4
2001 5/.....	4.3	9.2	2.7	0.4	0.3	2.4	4.9	6.3	6.7	7.3
2000.....	2.8	10.0	2.3	0.5	0.4	2.2	4.4	3.5	3.6	3.5
1999.....	2.8	9.4	2.1	0.5	0.4	2.4	3.9	3.6	3.7	3.9
All races 3/, female										
2005.....	1.3	6.6	1.8	0.4	0.4	1.0	1.9	1.7	1.5	1.8
2004.....	1.4	7.7	2.0	0.4	0.4	1.2	1.7	1.9	1.5	1.9
2003.....	1.4	6.9	2.0	0.3	0.2	1.1	2.0	1.8	1.8	1.9
2002.....	1.4	6.8	2.2	0.5	0.3	1.1	1.8	1.8	2.0	2.1
2001 5/.....	2.0	6.7	2.0	0.4	0.4	1.2	2.4	3.0	3.2	3.2
2000.....	1.5	7.7	2.0	0.4	0.5	1.4	2.0	1.9	2.0	2.4
1999.....	1.6	7.6	2.1	0.7	0.5	1.4	2.2	2.2	2.3	2.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)

White, both sexes

2005 4/.....	1.5	6.0	1.5	0.3	0.2	1.2	2.4	2.1	1.7	1.7
2004 4/.....	1.5	6.5	1.5	0.3	0.3	1.3	2.4	2.1	1.5	1.7
2003 4/.....	1.5	6.9	1.5	0.3	0.2	1.2	2.1	2.0	1.7	2.0
2002.....	1.5	5.3	1.7	0.3	0.2	1.1	2.2	1.9	1.7	2.0
2001 5/.....	2.6	6.0	1.6	0.4	0.3	1.4	3.0	3.9	4.3	4.6
2000.....	1.5	6.5	1.5	0.4	0.4	1.3	2.3	2.0	2.0	2.1
1999.....	1.6	6.6	1.5	0.5	0.4	1.4	2.4	2.1	2.2	2.1

White, male

2005 4/.....	1.9	6.4	1.6	0.3	*	1.6	3.2	2.7	2.2	2.3
2004 4/.....	1.9	6.8	1.7	0.3	0.2	1.6	3.2	2.6	1.9	2.1
2003 4/.....	1.9	8.0	1.4	0.3	0.3	1.6	2.6	2.4	2.2	2.5
2002.....	1.9	6.2	1.8	*	*	1.5	2.9	2.5	2.0	2.5
2001 5/.....	3.7	7.0	1.9	0.3	*	1.8	3.9	5.3	5.9	6.8
2000.....	2.0	8.1	1.7	0.5	0.4	1.6	3.0	2.5	2.5	2.5
1999.....	2.1	7.7	1.5	0.4	0.3	1.7	3.0	2.7	2.6	2.7

White, female

2005 4/.....	1.0	5.5	1.4	*	0.4	0.8	1.5	1.3	1.2	1.2
2004 4/.....	1.1	6.1	1.3	0.3	0.3	0.9	1.5	1.7	1.2	1.3
2003 4/.....	1.1	5.6	1.6	*	*	0.8	1.5	1.5	1.2	1.4
2002.....	1.1	4.5	1.6	0.4	0.3	0.7	1.5	1.2	1.4	1.5
2001 5/.....	1.6	4.9	1.4	0.4	0.3	0.9	2.0	2.5	2.6	2.3
2000.....	1.1	4.8	1.3	0.3	0.5	1.0	1.5	1.4	1.4	1.7
1999.....	1.2	5.4	1.5	0.5	0.4	1.1	1.8	1.5	1.7	1.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)

Black, both sexes

2005 4/.....	5.0	14.0	4.8	0.9	0.9	3.5	6.2	6.1	5.7	7.1
2004 4/.....	5.0	15.1	5.4	0.9	0.6	3.9	5.6	6.6	6.3	6.7
2003 4/.....	5.2	15.2	5.3	1.0	0.6	4.2	7.2	6.6	6.2	7.2
2002.....	5.3	16.5	5.8	1.0	0.6	3.7	6.4	7.3	7.0	7.4
2001 5/.....	6.3	18.6	6.1	*	0.6	4.0	7.1	9.0	8.4	9.8
2000.....	5.9	21.6	5.5	0.8	0.8	4.5	8.8	7.6	7.6	8.6
1999.....	5.7	19.2	5.4	1.3	0.6	4.5	7.0	7.7	8.1	9.2

Black, male

2005 4/.....	7.2	15.4	5.9	*	*	5.1	8.5	9.1	8.6	9.0
2004 4/.....	6.8	13.9	5.5	*	*	5.0	7.9	9.4	9.5	8.5
2003 4/.....	7.4	16.1	6.5	*	*	5.7	9.9	9.2	7.6	9.8
2002.....	7.3	15.7	6.4	*	*	4.8	9.4	9.7	9.1	9.5
2001 5/.....	8.5	20.7	7.3	*	*	5.4	9.8	12.8	11.0	11.3
2000.....	8.0	21.1	5.5	*	*	5.5	12.6	10.7	10.5	10.9
1999.....	7.7	19.2	5.4	1.2	*	6.1	9.8	9.5	10.7	12.8

Black, female

2005 4/.....	3.0	12.6	3.7	*	*	1.9	3.9	3.3	3.0	5.4
2004 4/.....	3.4	16.4	5.4	*	*	2.8	3.3	3.9	3.4	5.2
2003 4/.....	3.3	14.2	4.1	*	*	2.7	4.5	4.1	5.0	5.0
2002.....	3.4	17.3	5.1	*	*	2.6	3.5	5.0	5.0	5.6
2001 5/.....	4.3	16.4	4.8	*	*	2.5	4.5	5.5	6.1	8.4
2000.....	4.0	22.2	5.5	*	*	3.4	5.2	4.7	4.9	6.7
1999.....	3.9	19.1	5.4	1.3	*	2.9	4.3	6.1	5.7	6.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)										
All races 3/, both sexes										
2005.....	1.6	2.5	0.4	0.1	0.2	0.7	1.6	1.8	1.9	2.3
2004.....	1.7	2.6	0.3	0.1	0.2	0.7	1.7	1.9	2.1	2.9
2003.....	1.7	1.9	0.2	*	0.2	0.8	1.9	1.8	2.3	3.2
2002.....	1.7	2.5	0.4	0.1	0.2	0.7	1.7	1.9	2.3	3.3
2001.....	1.5	2.1	0.3	*	0.2	0.7	1.4	1.6	2.0	3.0
2000.....	1.4	1.7	0.3	*	0.2	0.6	1.3	1.6	1.8	2.6
1999.....	1.4	1.4	0.3	*	0.2	0.8	1.5	1.6	2.1	2.8
All races 3/, male										
2005.....	2.0	3.2	0.5	*	0.3	1.0	2.3	2.4	2.3	2.7
2004.....	2.2	3.1	0.3	*	0.2	1.1	2.4	2.6	2.9	3.8
2003.....	2.3	2.4	*	*	0.3	1.2	2.7	2.5	3.2	4.2
2002.....	2.2	2.5	0.4	*	0.2	1.0	2.4	2.7	3.1	4.5
2001.....	2.0	2.4	*	*	0.3	1.0	2.1	2.4	2.7	3.9
2000.....	1.9	1.4	0.3	*	0.4	0.9	2.0	2.3	2.6	3.7
1999.....	2.0	1.7	0.3	*	0.3	1.3	2.2	2.3	2.8	3.9
All races 3/, female										
2005.....	1.3	1.8	0.3	*	*	0.4	0.9	1.2	1.5	1.9
2004.....	1.2	2.0	0.3	*	*	0.4	0.8	1.1	1.4	1.9
2003.....	1.2	1.3	*	*	*	0.5	1.1	1.1	1.5	2.2
2002.....	1.2	2.5	0.4	*	*	0.4	1.0	1.0	1.4	2.1
2001.....	0.9	1.7	0.3	*	*	0.3	0.7	0.9	1.3	2.1
2000.....	0.8	2.0	*	*	*	0.3	0.6	0.9	1.1	1.6
1999.....	0.9	1.2	0.3	*	*	0.3	0.6	0.9	1.3	1.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)										
White, both sexes										
2005 4/.....	1.6	2.1	0.3	*	0.2	0.7	1.7	2.0	2.0	2.4
2004 4/.....	1.7	2.1	0.2	*	0.2	0.8	1.9	2.1	2.2	3.0
2003 4/.....	1.8	1.7	0.2	*	0.2	0.9	2.1	2.0	2.6	3.4
2002.....	1.7	1.8	0.4	*	0.2	0.7	1.9	1.9	2.3	3.4
2001.....	1.5	1.4	0.3	*	0.2	0.7	1.5	1.7	2.1	3.0
2000.....	1.3	1.4	0.2	*	0.2	0.6	1.3	1.6	1.8	2.7
1999.....	1.4	1.2	0.2	*	0.2	0.8	1.5	1.6	2.1	2.8
White, male										
2005 4/.....	2.0	2.8	0.4	*	0.3	1.0	2.5	2.5	2.4	2.8
2004 4/.....	2.2	2.7	*	*	*	1.1	2.7	2.9	2.9	3.9
2003 4/.....	2.3	1.9	*	*	0.3	1.3	2.9	2.7	3.3	4.4
2002.....	2.2	1.8	0.4	*	0.2	1.1	2.6	2.7	3.1	4.5
2001.....	2.0	1.7	*	*	0.3	1.1	2.2	2.5	2.9	3.8
2000.....	1.8	*	*	*	0.3	1.0	1.9	2.3	2.5	3.7
1999.....	1.9	1.4	*	*	0.3	1.3	2.3	2.3	2.8	3.7
White, female										
2005 4/.....	1.3	1.5	*	*	*	0.4	0.9	1.4	1.6	2.0
2004 4/.....	1.3	1.5	*	*	*	0.4	0.9	1.2	1.5	2.1
2003 4/.....	1.3	1.4	*	*	*	0.5	1.2	1.2	1.7	2.3
2002.....	1.2	1.8	0.4	*	*	0.4	1.1	1.1	1.5	2.3
2001.....	1.0	*	*	*	*	0.3	0.8	0.9	1.4	2.3
2000.....	0.8	1.6	*	*	*	0.3	0.7	0.9	1.1	1.6
1999.....	0.9	*	*	*	*	*	0.6	1.0	1.4	1.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)										
Black, both sexes										
2005 4/.....	1.8	4.7	0.9	*	*	0.6	1.1	1.4	1.9	2.5
2004 4/.....	1.8	4.8	*	*	*	0.7	1.1	1.0	2.6	3.1
2003 4/.....	1.8	3.2	*	*	*	*	1.1	1.4	2.1	3.2
2002.....	2.0	5.5	*	*	*	*	1.3	2.0	2.5	4.1
2001.....	1.7	5.5	*	*	*	*	1.1	1.5	1.9	3.8
2000.....	1.8	*	*	*	*	*	1.4	1.7	2.2	3.3
1999.....	1.9	*	*	*	*	0.8	1.7	1.9	2.5	3.9
Black, male										
2005 4/.....	2.4	*	*	*	*	*	1.6	2.3	2.3	3.1
2004 4/.....	2.6	*	*	*	*	*	1.5	1.5	4.0	4.6
2003 4/.....	2.7	*	*	*	*	*	1.9	2.5	3.4	4.4
2002.....	2.9	*	*	*	*	*	1.9	3.2	3.7	6.5
2001.....	2.7	6.0	*	*	*	*	1.9	2.5	2.7	5.9
2000.....	2.8	*	*	*	*	*	2.5	2.7	3.4	4.8
1999.....	3.0	*	*	*	*	1.3	2.7	3.4	3.7	6.3
Black, female										
2005 4/.....	1.1	*	*	*	*	*	*	*	1.5	2.0
2004 4/.....	1.1	*	*	*	*	*	*	*	1.4	1.8
2003 4/.....	0.9	*	*	*	*	*	*	*	*	2.2
2002.....	1.2	*	*	*	*	*	*	*	1.5	1.9
2001.....	0.9	*	*	*	*	*	*	*	*	2.0
2000.....	0.9	*	*	*	*	*	*	*	*	2.0
1999.....	0.9	*	*	*	*	*	*	*	1.4	1.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)										
All races 3/, both sexes										
2005.....	1.5	2.5	0.4	*	0.2	0.6	1.3	1.7	1.8	2.3
2004.....	1.6	2.6	0.3	0.1	0.2	0.6	1.5	1.8	2.0	2.8
2003.....	1.7	1.9	0.2	*	0.1	0.7	1.7	1.7	2.3	3.1
2002.....	1.6	2.5	0.4	0.1	0.2	0.5	1.5	1.7	2.2	3.2
2001.....	1.4	2.1	0.3	*	0.1	0.5	1.2	1.5	1.9	2.9
2000.....	1.3	1.7	0.3	*	0.2	0.5	1.1	1.4	1.7	2.6
1999.....	1.3	1.4	0.3	*	0.1	0.5	1.1	1.4	2.0	2.7
All races 3/, male										
2005.....	1.8	3.2	0.5	*	0.2	0.8	1.9	2.2	2.2	2.6
2004.....	2.1	3.1	0.3	*	0.2	0.9	2.1	2.4	2.7	3.7
2003.....	2.2	2.4	*	*	0.2	0.9	2.4	2.4	3.0	4.1
2002.....	2.1	2.5	0.3	*	0.2	0.7	2.1	2.4	3.0	4.3
2001.....	1.9	2.4	*	*	0.2	0.8	1.8	2.1	2.6	3.8
2000.....	1.8	1.4	0.3	*	0.3	0.7	1.6	2.0	2.4	3.6
1999.....	1.8	1.7	*	*	0.2	0.7	1.7	2.0	2.7	3.7
All races 3/, female										
2005.....	1.2	1.8	0.3	*	*	0.4	0.8	1.1	1.5	1.9
2004.....	1.2	2.0	0.3	*	*	0.3	0.8	1.1	1.3	1.9
2003.....	1.2	1.3	*	*	*	0.4	1.0	1.1	1.5	2.1
2002.....	1.1	2.5	0.4	*	*	0.3	0.9	1.0	1.4	2.0
2001.....	0.9	1.7	0.3	*	*	0.3	0.6	0.8	1.3	2.0
2000.....	0.8	2.0	*	*	*	0.3	0.5	0.8	1.0	1.6
1999.....	0.8	1.2	0.3	*	*	0.2	0.5	0.8	1.3	1.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)										
White, both sexes										
2005 4/.....	1.6	2.1	0.3	*	0.1	0.6	1.5	1.9	1.9	2.4
2004 4/.....	1.7	2.1	0.2	*	0.1	0.7	1.7	2.0	2.1	2.9
2003 4/.....	1.7	1.7	0.2	*	0.1	0.8	1.9	1.9	2.5	3.3
2002.....	1.6	1.8	0.4	*	0.2	0.6	1.7	1.8	2.2	3.3
2001.....	1.4	1.4	0.3	*	0.2	0.6	1.3	1.6	2.0	2.9
2000.....	1.3	1.4	0.2	*	0.1	0.5	1.1	1.5	1.8	2.6
1999.....	1.3	1.2	0.2	*	0.1	0.5	1.2	1.5	2.0	2.6
White, male										
2005 4/.....	1.8	2.8	0.4	*	*	0.8	2.1	2.4	2.3	2.7
2004 4/.....	2.1	2.7	*	*	*	1.0	2.4	2.8	2.8	3.8
2003 4/.....	2.2	1.9	*	*	*	1.1	2.7	2.5	3.2	4.2
2002.....	2.0	1.8	0.4	*	*	0.8	2.3	2.5	3.0	4.3
2001.....	1.9	1.7	*	*	0.3	0.9	1.9	2.2	2.7	3.6
2000.....	1.7	*	*	*	0.3	0.7	1.6	2.1	2.4	3.6
1999.....	1.7	1.4	*	*	*	0.8	1.8	2.0	2.7	3.5
White, female										
2005 4/.....	1.3	1.5	*	*	*	0.4	0.8	1.3	1.6	2.0
2004 4/.....	1.2	1.5	*	*	*	0.3	0.9	1.2	1.4	2.0
2003 4/.....	1.3	1.4	*	*	*	0.5	1.2	1.2	1.7	2.2
2002.....	1.2	1.8	0.4	*	*	0.4	1.0	1.1	1.5	2.2
2001.....	1.0	*	*	*	*	0.3	0.8	0.9	1.4	2.1
2000.....	0.8	1.6	*	*	*	0.3	0.6	0.9	1.1	1.6
1999.....	0.8	*	*	*	*	*	0.5	0.9	1.4	1.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)

Black, both sexes

2005 4/.....	1.7	4.7	0.8	*	*	*	0.9	1.3	1.8	2.5
2004 4/.....	1.7	4.8	*	*	*	*	0.8	0.8	2.5	3.1
2003 4/.....	1.7	3.2	*	*	*	*	0.8	1.3	2.0	3.2
2002.....	1.9	5.5	*	*	*	*	1.1	1.8	2.4	3.9
2001.....	1.7	5.5	*	*	*	*	0.8	1.3	1.9	3.8
2000.....	1.7	*	*	*	*	*	0.9	1.4	1.9	3.2
1999.....	1.7	*	*	*	*	*	1.1	1.5	2.4	3.8

Black, male

2005 4/.....	2.3	*	*	*	*	*	*	1.9	2.2	3.0
2004 4/.....	2.4	*	*	*	*	*	*	*	3.7	4.5
2003 4/.....	2.6	*	*	*	*	*	1.4	2.2	3.2	4.3
2002.....	2.7	*	*	*	*	*	1.4	2.7	3.4	6.3
2001.....	2.5	6.0	*	*	*	*	1.5	2.2	2.7	5.9
2000.....	2.5	*	*	*	*	*	1.5	2.2	3.1	4.5
1999.....	2.7	*	*	*	*	*	1.8	2.7	3.5	6.1

Black, female

2005 4/.....	1.1	*	*	*	*	*	*	*	1.5	2.0
2004 4/.....	1.1	*	*	*	*	*	*	*	*	1.8
2003 4/.....	0.9	*	*	*	*	*	*	*	*	2.1
2002.....	1.2	*	*	*	*	*	*	*	1.5	1.8
2001.....	0.8	*	*	*	*	*	*	*	*	2.0
2000.....	0.9	*	*	*	*	*	*	*	*	2.0
1999.....	0.9	*	*	*	*	*	*	*	1.4	1.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Complications of medical and surgical care (Y40-Y84,Y88)										
All races 3/, both sexes										
2005.....	0.9	*	*	*	*	*	*	0.2	0.2	0.3
2004.....	1.0	*	*	*	*	0.1	0.1	0.2	0.2	0.3
2003.....	1.0	*	0.2	*	*	0.1	0.1	0.1	0.2	0.4
2002.....	1.0	*	*	*	0.1	*	*	0.1	0.2	0.3
2001.....	1.1	0.6	0.2	*	*	*	0.1	0.1	0.3	0.3
2000.....	1.1	*	0.1	*	*	*	0.1	0.2	0.2	0.3
1999.....	1.0	1.4	0.2	*	*	*	*	0.1	0.2	0.3
All races 3/, male										
2005.....	0.8	*	*	*	*	*	*	*	*	0.2
2004.....	0.9	*	*	*	*	*	*	*	*	0.2
2003.....	0.9	*	*	*	*	*	*	*	0.2	0.3
2002.....	0.9	*	*	*	*	*	*	*	0.3	0.3
2001.....	1.0	*	*	*	*	*	*	*	0.3	0.4
2000.....	1.0	*	*	*	*	*	*	0.2	0.2	0.3
1999.....	1.0	1.6	*	*	*	*	*	*	0.3	0.3
All races 3/, female										
2005.....	1.0	*	*	*	*	*	*	*	0.3	0.3
2004.....	1.0	*	*	*	*	*	*	*	0.3	0.3
2003.....	1.0	*	*	*	*	*	*	*	0.2	0.4
2002.....	1.1	*	*	*	*	*	*	*	0.2	0.3
2001.....	1.1	*	*	*	*	*	*	*	0.2	0.3
2000.....	1.1	*	*	*	*	*	*	*	0.2	0.3
1999.....	1.1	1.2	*	*	*	*	*	0.2	*	0.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K86.0,R78.0,X45,X65,Y15) 6/										
All races 3/, both sexes										
2005.....	7.3	*	*	*	*	0.2	0.5	1.0	1.7	4.7
2004.....	7.2	*	*	*	*	0.2	0.5	0.9	2.1	5.0
2003.....	7.1	*	*	*	*	0.2	0.4	0.9	2.1	5.3
2002.....	7.0	*	*	*	*	0.2	0.4	0.8	2.1	5.4
2001.....	7.1	*	*	*	*	0.2	0.3	0.9	2.3	6.0
2000.....	7.0	*	*	*	*	0.2	0.3	0.9	2.3	6.0
1999.....	7.0	*	*	*	*	0.1	0.4	0.8	2.4	6.2
All races 3/, male										
2005.....	11.1	*	*	*	*	0.3	0.9	1.5	2.5	6.6
2004.....	11.0	*	*	*	*	0.2	0.8	1.4	3.2	7.1
2003.....	10.9	*	*	*	*	0.4	0.6	1.4	3.2	7.4
2002.....	10.8	*	*	*	*	0.3	0.6	1.4	3.1	7.6
2001.....	10.8	*	*	*	*	0.3	0.6	1.3	3.3	8.5
2000.....	10.9	*	*	*	*	0.3	0.5	1.5	3.5	8.7
1999.....	10.9	*	*	*	*	0.2	0.6	1.2	3.3	8.7
All races 3/, female										
2005.....	3.6	*	*	*	*	*	*	0.4	1.0	2.7
2004.....	3.5	*	*	*	*	*	*	0.4	1.0	2.9
2003.....	3.4	*	*	*	*	*	*	0.3	0.9	3.2
2002.....	3.4	*	*	*	*	*	*	0.3	1.1	3.3
2001.....	3.4	*	*	*	*	*	*	0.5	1.3	3.5
2000.....	3.2	*	*	*	*	*	*	0.3	1.1	3.2
1999.....	3.2	*	*	*	*	*	*	0.3	1.5	3.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K86.0,R78.0,X45,X65,Y15) 6/										
White, both sexes										
2005 4/.....	7.7	*	*	*	*	0.2	0.5	1.0	1.7	4.9
2004 4/.....	7.5	*	*	*	*	0.2	0.5	0.9	2.2	5.2
2003 4/.....	7.4	*	*	*	*	0.2	0.4	0.9	2.1	5.4
2002.....	7.2	*	*	*	*	0.2	0.3	0.8	2.1	5.5
2001.....	7.2	*	*	*	*	0.2	0.3	0.9	2.3	6.0
2000.....	7.1	*	*	*	*	0.2	0.3	0.9	2.2	5.9
1999.....	7.0	*	*	*	*	*	0.4	0.8	2.4	6.0
White, male										
2005 4/.....	11.7	*	*	*	*	0.3	0.9	1.5	2.5	6.9
2004 4/.....	11.5	*	*	*	*	*	0.8	1.3	3.3	7.2
2003 4/.....	11.3	*	*	*	*	0.4	0.6	1.5	3.2	7.6
2002.....	11.1	*	*	*	*	0.3	0.6	1.4	3.1	7.6
2001.....	11.0	*	*	*	*	0.3	0.5	1.3	3.3	8.4
2000.....	11.0	*	*	*	*	0.3	0.5	1.5	3.4	8.7
1999.....	10.9	*	*	*	*	*	0.7	1.3	3.2	8.5
White, female										
2005 4/.....	3.7	*	*	*	*	*	*	0.4	0.9	2.7
2004 4/.....	3.6	*	*	*	*	*	*	0.4	1.0	3.1
2003 4/.....	3.5	*	*	*	*	*	*	0.3	1.0	3.2
2002.....	3.4	*	*	*	*	*	*	*	1.0	3.3
2001.....	3.4	*	*	*	*	*	*	0.4	1.2	3.6
2000.....	3.2	*	*	*	*	*	*	*	1.1	3.1
1999.....	3.1	*	*	*	*	*	*	*	1.4	3.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K86.0,R78.0,X45,X65,Y15) 6/										
Black, both sexes										
2005 4/.....	5.9	*	*	*	*	*	*	*	1.2	3.0
2004 4/.....	6.1	*	*	*	*	*	*	*	1.3	3.7
2003 4/.....	6.3	*	*	*	*	*	*	0.9	1.3	4.7
2002.....	6.4	*	*	*	*	*	*	*	1.9	4.4
2001.....	7.3	*	*	*	*	*	*	0.8	1.9	5.2
2000.....	7.4	*	*	*	*	*	*	0.9	2.5	5.5
1999.....	7.8	*	*	*	*	*	*	*	2.2	7.1
Black, male										
2005 4/.....	9.1	*	*	*	*	*	*	*	1.5	4.1
2004 4/.....	9.7	*	*	*	*	*	*	*	2.2	5.9
2003 4/.....	10.0	*	*	*	*	*	*	*	2.2	6.5
2002.....	10.0	*	*	*	*	*	*	*	2.9	6.2
2001.....	11.6	*	*	*	*	*	*	*	2.5	8.4
2000.....	11.4	*	*	*	*	*	*	1.6	3.4	8.1
1999.....	12.2	*	*	*	*	*	*	*	2.9	10.2
Black, female										
2005 4/.....	3.0	*	*	*	*	*	*	*	*	2.1
2004 4/.....	2.8	*	*	*	*	*	*	*	*	1.7
2003 4/.....	2.9	*	*	*	*	*	*	*	*	3.2
2002.....	3.2	*	*	*	*	*	*	*	*	2.9
2001.....	3.5	*	*	*	*	*	*	*	*	2.4
2000.....	3.7	*	*	*	*	*	*	*	1.7	3.2
1999.....	3.9	*	*	*	*	*	*	*	1.5	4.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7-F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 6/

All races 3/, both sexes

2005.....	11.3	0.9	0.2	*	0.2	3.7	11.0	13.5	15.1	18.3
2004.....	10.5	0.7	0.2	*	0.2	3.9	10.0	12.2	13.1	18.4
2003.....	9.9	0.6	0.2	*	0.2	3.3	9.4	10.9	13.3	18.3
2002.....	9.0	0.7	0.2	*	0.2	3.1	7.9	9.8	12.4	17.5
2001.....	7.6	0.5	0.2	*	0.2	2.7	6.5	8.2	10.6	15.5
2000.....	7.0	*	*	*	0.2	2.2	5.9	7.5	10.1	14.4
1999.....	6.9	0.6	0.2	*	0.1	1.8	5.3	7.4	10.3	14.2

All races 3/, male

2005.....	14.5	*	0.3	*	*	5.4	16.0	18.9	20.0	23.0
2004.....	13.4	*	*	*	0.2	5.6	14.7	17.3	17.6	23.6
2003.....	12.9	*	0.3	*	*	4.5	13.9	15.4	17.7	23.8
2002.....	11.8	1.0	*	*	*	4.3	11.4	14.0	17.0	23.1
2001.....	10.2	*	0.3	*	*	3.7	9.6	11.9	14.4	20.7
2000.....	9.5	*	*	*	*	3.1	8.8	10.8	14.1	19.8
1999.....	9.4	*	*	*	*	2.4	7.7	10.8	14.6	19.6

All races 3/, female

2005.....	8.2	*	*	*	0.2	2.0	5.6	7.7	10.1	13.5
2004.....	7.6	*	*	*	0.3	2.1	5.0	6.7	8.5	13.2
2003.....	7.0	*	*	*	*	1.9	4.7	6.2	8.8	12.8
2002.....	6.3	*	0.3	*	*	1.8	4.2	5.4	7.8	11.9
2001.....	5.1	*	*	*	*	1.6	3.2	4.4	6.7	10.3
2000.....	4.6	*	*	*	*	1.4	2.9	4.1	6.1	9.0
1999.....	4.4	*	*	*	*	1.2	2.8	4.0	6.0	8.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7-F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 6/

White, both sexes

2005 4/.....	12.0	0.8	*	*	0.2	4.5	13.0	15.3	16.8	19.8
2004 4/.....	11.1	*	*	*	0.2	4.7	11.8	14.0	14.5	19.7
2003 4/.....	10.4	*	0.2	*	0.1	3.9	11.2	12.4	14.7	19.4
2002.....	9.4	*	*	*	0.2	3.6	9.2	10.9	13.5	18.5
2001.....	7.8	*	*	*	0.2	3.1	7.5	9.1	11.4	16.1
2000.....	7.1	*	*	*	0.2	2.6	6.7	8.2	10.7	14.8
1999.....	6.9	*	*	*	0.1	2.1	5.9	7.9	10.7	14.4

White, male

2005 4/.....	15.3	*	*	*	*	6.5	19.0	21.5	22.2	24.8
2004 4/.....	14.1	*	*	*	*	6.7	17.3	19.9	19.5	25.3
2003 4/.....	13.5	*	*	*	*	5.5	16.4	17.6	19.5	25.1
2002.....	12.2	*	*	*	*	5.0	13.2	15.5	18.5	24.2
2001.....	10.4	*	*	*	*	4.4	10.9	13.1	15.6	21.6
2000.....	9.6	*	*	*	*	3.6	10.0	11.6	14.9	20.4
1999.....	9.3	*	*	*	*	2.8	8.5	11.3	15.2	19.9

White, female

2005 4/.....	8.8	*	*	*	*	2.3	6.5	8.8	11.1	14.6
2004 4/.....	8.2	*	*	*	0.3	2.5	5.8	7.7	9.4	14.1
2003 4/.....	7.4	*	*	*	*	2.2	5.5	6.8	9.7	13.6
2002.....	6.7	*	*	*	*	2.1	4.9	6.1	8.3	12.6
2001.....	5.4	*	*	*	*	1.6	3.8	5.0	7.0	10.6
2000.....	4.7	*	*	*	*	1.5	3.2	4.6	6.3	9.1
1999.....	4.5	*	*	*	*	1.4	3.0	4.3	6.1	8.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7-F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 6/

Black, both sexes

2005 4/.....	10.5	*	*	*	*	1.0	3.4	7.5	11.5	16.1
2004 4/.....	9.4	*	*	*	*	1.0	3.5	6.2	9.6	16.3
2003 4/.....	9.2	*	*	*	*	0.7	2.9	5.8	10.4	17.9
2002.....	9.2	*	*	*	*	1.0	3.2	6.6	10.2	17.3
2001.....	8.5	*	*	*	*	0.9	3.0	5.8	9.5	16.7
2000.....	8.3	*	*	*	*	0.9	3.2	6.2	10.3	16.5
1999.....	8.6	*	*	*	*	0.7	3.7	6.8	11.2	16.5

Black, male

2005 4/.....	14.3	*	*	*	*	*	4.8	11.1	14.8	20.1
2004 4/.....	12.8	*	*	*	*	1.4	5.0	9.0	13.2	20.5
2003 4/.....	12.7	*	*	*	*	*	4.4	7.7	13.6	24.3
2002.....	12.8	*	*	*	*	1.4	4.6	9.9	13.9	24.2
2001.....	12.2	*	*	*	*	*	5.1	9.3	12.3	22.4
2000.....	12.0	*	*	*	*	*	4.6	9.8	14.3	22.8
1999.....	12.7	*	*	*	*	*	5.6	11.0	15.7	23.9

Black, female

2005 4/.....	7.0	*	*	*	*	*	2.0	4.0	8.6	12.5
2004 4/.....	6.3	*	*	*	*	*	2.0	3.6	6.4	12.5
2003 4/.....	6.1	*	*	*	*	*	1.5	4.0	7.6	12.2
2002.....	5.8	*	*	*	*	*	1.9	3.6	6.8	11.2
2001.....	5.1	*	*	*	*	*	*	2.6	7.0	11.6
2000.....	4.9	*	*	*	*	*	1.8	2.9	6.8	11.0
1999.....	4.8	*	*	*	*	*	1.9	3.1	7.3	10.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 6/										
All races 3/, both sexes										
2005.....	10.4	*	0.4	0.3	1.3	12.5	20.0	17.2	14.2	11.6
2004.....	10.1	*	0.3	0.3	1.1	12.0	19.3	16.9	13.3	11.2
2003.....	10.4	*	0.3	0.3	1.2	12.1	21.1	17.4	13.8	11.6
2002.....	10.5	*	0.4	0.4	1.3	12.1	21.3	17.8	13.2	12.4
2001.....	10.4	*	0.5	0.4	1.2	12.4	21.2	17.7	13.5	12.2
2000.....	10.2	*	0.3	0.3	1.5	12.9	20.9	16.3	12.8	12.0
1999.....	10.3	*	0.4	0.4	1.7	14.4	21.0	16.6	13.2	12.3
All races 3/, male										
2005.....	18.3	*	0.5	0.4	2.0	22.0	35.3	30.2	24.3	19.3
2004.....	17.6	*	0.4	0.3	1.8	20.7	34.2	29.8	22.5	19.0
2003.....	18.3	*	0.3	0.4	2.0	21.2	37.1	30.7	23.7	19.3
2002.....	18.4	*	0.5	0.4	1.9	21.1	37.6	31.1	22.3	20.6
2001.....	18.2	*	0.5	0.5	1.9	21.8	37.7	30.9	22.9	20.0
2000.....	17.8	*	0.4	0.4	2.4	22.4	37.0	28.5	21.4	19.5
1999.....	18.1	*	0.5	0.5	2.5	24.8	36.8	28.8	22.3	20.0
All races 3/, female										
2005.....	2.7	*	0.3	0.3	0.6	2.4	3.6	3.7	3.8	3.9
2004.....	2.7	*	0.3	0.3	0.4	2.9	3.5	3.4	3.9	3.3
2003.....	2.7	*	0.3	0.2	0.4	2.4	4.2	3.6	3.7	3.8
2002.....	2.8	*	0.3	0.3	0.7	2.7	4.2	4.1	3.9	4.2
2001.....	2.8	*	0.4	0.3	0.5	2.6	3.8	4.1	3.9	4.6
2000.....	2.8	*	*	0.2	0.5	2.9	4.2	3.9	4.1	4.5
1999.....	2.9	*	0.4	0.3	0.8	3.4	4.4	4.1	4.1	4.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 6/										
White, both sexes										
2005 4/.....	9.1	*	0.3	0.2	1.1	8.5	12.6	11.7	10.7	10.2
2004 4/.....	9.0	*	*	0.2	0.9	8.5	12.9	11.4	10.4	10.0
2003 4/.....	9.2	*	0.2	0.2	1.0	8.4	13.9	11.7	10.8	10.4
2002.....	9.3	*	0.3	0.3	1.1	8.7	13.9	12.3	10.2	10.9
2001.....	9.4	*	0.4	0.4	1.1	8.9	14.0	12.9	11.1	11.2
2000.....	9.1	*	0.2	0.3	1.4	9.2	13.9	11.5	10.4	10.9
1999.....	9.2	*	0.3	0.3	1.6	10.2	14.0	12.1	11.1	11.3
White, male										
2005 4/.....	15.8	*	0.3	0.3	1.7	14.6	21.8	19.7	17.6	16.3
2004 4/.....	15.5	*	*	*	1.5	14.4	22.3	19.5	17.0	16.6
2003 4/.....	16.0	*	*	*	1.6	14.4	23.9	19.8	17.9	16.8
2002.....	16.1	*	0.4	0.3	1.5	14.9	23.9	20.5	16.7	17.7
2001.....	16.2	*	0.4	0.5	1.7	15.0	24.1	21.8	18.2	17.9
2000.....	15.6	*	*	0.4	2.3	15.6	23.9	19.3	16.9	17.2
1999.....	15.9	*	*	0.3	2.3	17.3	23.8	20.3	18.1	18.1
White, female										
2005 4/.....	2.6	*	*	*	0.4	1.9	2.6	3.2	3.6	3.9
2004 4/.....	2.7	*	*	*	0.3	2.3	2.8	2.8	3.5	3.2
2003 4/.....	2.6	*	*	*	0.3	2.0	3.1	3.0	3.3	3.9
2002.....	2.7	*	*	0.3	0.6	2.1	3.1	3.7	3.4	4.0
2001.....	2.7	*	*	0.3	0.5	2.3	3.1	3.4	3.7	4.3
2000.....	2.7	*	*	*	0.5	2.3	3.2	3.3	3.7	4.5
1999.....	2.8	*	*	*	0.7	2.6	3.5	3.4	3.8	4.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 6/

Black, both sexes

2005 4/.....	20.4	*	1.0	0.8	2.4	33.6	61.8	52.6	39.4	23.8
2004 4/.....	19.3	*	1.2	0.8	2.2	30.9	58.0	53.1	34.5	22.0
2003 4/.....	20.1	*	0.9	0.8	2.5	32.2	63.8	55.5	35.8	21.5
2002.....	20.2	*	*	*	2.4	31.4	64.7	54.8	35.1	24.4
2001.....	19.3	*	0.9	*	1.9	32.9	64.4	50.4	31.5	22.6
2000.....	19.3	*	*	*	2.2	33.9	64.0	48.1	30.9	21.7
1999.....	19.4	*	1.1	0.9	2.3	38.0	63.5	46.5	29.0	20.9

Black, male

2005 4/.....	38.7	*	*	*	3.7	61.5	114.1	100.1	75.9	44.4
2004 4/.....	36.4	*	*	*	3.5	55.2	107.7	102.1	64.9	41.3
2003 4/.....	37.8	*	*	*	3.7	59.3	117.4	107.6	67.4	40.3
2002.....	37.8	*	*	*	3.5	56.0	120.6	106.3	65.9	44.8
2001.....	36.4	*	*	*	3.0	60.5	123.4	96.0	60.1	39.5
2000.....	36.1	*	*	*	3.7	61.5	121.3	92.9	57.0	39.9
1999.....	36.0	*	*	1.3	3.4	67.1	119.9	88.6	54.3	37.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 6/

Black, female

2005 4/.....	3.7	*	*	*	1.2	5.0	8.5	7.7	6.4	5.3
2004 4/.....	3.7	*	*	*	*	5.9	7.9	7.3	7.0	4.7
2003 4/.....	3.9	*	*	*	1.3	4.2	10.5	7.4	7.2	4.7
2002.....	4.2	*	*	*	1.3	6.0	10.2	7.7	7.4	6.3
2001.....	3.8	*	*	*	*	4.5	7.8	8.9	5.8	7.5
2000.....	4.0	*	*	*	*	5.7	9.6	7.5	7.4	5.5
1999.....	4.4	*	*	*	*	8.2	9.9	8.4	6.4	6.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
All causes										
All races 3/, both sexes										
2005.....	233.4	353.1	520.8	734.6	1,136.9	1,700.0	2,657.6	4,154.0	6,712.9	13,798.6
2004.....	231.7	352.3	511.7	734.8	1,140.1	1,727.4	2,676.4	4,193.2	6,717.2	13,823.5
2003.....	241.3	359.2	517.8	755.6	1,182.7	1,793.7	2,778.3	4,363.4	6,976.2	14,593.3
2002.....	242.6	357.1	512.8	770.1	1,187.7	1,832.7	2,845.9	4,449.1	7,103.8	14,828.3
2001.....	240.5	354.8	512.4	771.8	1,210.7	1,869.7	2,878.3	4,494.0	7,151.9	15,112.8
2000.....	239.1	352.5	509.0	792.6	1,241.0	1,906.3	2,929.4	4,553.8	7,335.0	15,524.4
1999.....	236.3	346.6	501.4	801.6	1,254.0	1,956.7	2,995.3	4,608.0	7,378.5	15,554.6
All races 3/, male										
2005.....	292.7	443.7	666.0	925.7	1,410.0	2,084.2	3,267.0	5,103.4	8,147.4	14,889.4
2004.....	289.5	444.4	657.0	917.3	1,411.0	2,125.1	3,279.2	5,173.5	8,179.0	15,031.1
2003.....	304.1	457.9	660.8	942.3	1,462.2	2,216.6	3,428.7	5,400.3	8,513.7	15,794.0
2002.....	306.4	454.7	653.5	961.8	1,476.0	2,265.9	3,533.0	5,522.7	8,652.3	16,254.5
2001.....	304.8	455.4	646.8	958.4	1,496.8	2,315.6	3,587.4	5,603.2	8,781.9	16,744.8
2000.....	305.9	456.3	642.2	989.8	1,535.9	2,362.0	3,676.1	5,693.2	9,095.4	17,501.4
1999.....	304.0	448.0	632.4	1,003.8	1,560.8	2,444.1	3,785.1	5,806.4	9,216.8	17,597.2
All races 3/, female										
2005.....	174.7	264.6	381.6	554.3	887.9	1,364.7	2,164.0	3,464.7	5,822.0	13,297.7
2004.....	174.5	262.5	372.7	562.7	893.6	1,381.2	2,191.0	3,488.7	5,816.6	13,280.3
2003.....	179.2	263.2	380.9	579.7	928.8	1,427.1	2,256.6	3,625.3	6,036.0	14,062.5
2002.....	179.7	262.4	378.2	590.0	926.0	1,458.7	2,296.7	3,690.9	6,162.2	14,209.6
2001.....	177.2	257.2	383.7	596.9	951.3	1,485.9	2,314.8	3,717.6	6,175.1	14,429.9
2000.....	173.4	252.0	381.3	608.1	973.7	1,515.7	2,341.2	3,760.2	6,296.6	14,719.2
1999.....	169.8	248.5	375.9	612.6	976.4	1,539.6	2,378.2	3,774.8	6,302.8	14,731.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
All causes										
White, both sexes										
2005 4/.....	217.8	326.6	480.7	689.7	1,085.7	1,649.5	2,618.1	4,128.9	6,731.9	14,021.1
2004 4/.....	215.0	325.0	471.5	688.5	1,090.5	1,675.5	2,634.1	4,160.5	6,736.0	14,010.1
2003 4/.....	222.6	329.4	474.5	710.6	1,127.8	1,739.2	2,728.6	4,329.5	6,992.1	14,792.3
2002.....	224.0	327.4	471.3	723.5	1,133.0	1,777.6	2,796.3	4,407.8	7,111.3	15,015.9
2001.....	218.8	322.8	471.4	723.2	1,156.1	1,805.8	2,823.2	4,446.8	7,149.4	15,287.5
2000.....	215.9	317.9	467.4	743.9	1,184.9	1,844.5	2,866.8	4,496.6	7,333.2	15,707.0
1999.....	210.8	310.5	460.0	747.2	1,193.5	1,895.5	2,923.9	4,542.8	7,370.9	15,718.0
White, male										
2005 4/.....	276.4	413.4	616.6	864.0	1,341.8	2,016.7	3,208.8	5,065.8	8,169.0	15,156.5
2004 4/.....	272.8	412.3	607.8	855.8	1,343.7	2,055.9	3,219.3	5,123.2	8,214.4	15,250.7
2003 4/.....	284.0	423.5	606.9	884.2	1,390.6	2,146.2	3,359.5	5,349.4	8,522.6	16,037.9
2002.....	287.2	420.4	601.2	902.0	1,402.7	2,196.1	3,469.7	5,469.3	8,660.8	16,473.2
2001.....	281.4	417.1	595.5	896.0	1,427.9	2,237.6	3,513.9	5,543.5	8,771.1	16,935.4
2000.....	279.4	416.1	588.3	925.1	1,465.1	2,285.1	3,593.1	5,620.3	9,092.5	17,716.4
1999.....	274.0	402.1	579.3	934.6	1,484.0	2,367.8	3,698.3	5,727.2	9,211.2	17,752.9
White, female										
2005 4/.....	158.6	239.9	347.5	521.9	847.6	1,322.9	2,129.8	3,435.6	5,831.3	13,498.3
2004 4/.....	156.6	238.0	338.1	527.6	855.6	1,338.1	2,153.2	3,456.3	5,817.7	13,450.9
2003 4/.....	160.7	235.8	344.9	543.9	884.5	1,379.6	2,212.3	3,592.1	6,048.6	14,240.6
2002.....	160.5	235.1	344.3	552.6	883.7	1,409.1	2,247.4	3,648.0	6,161.6	14,382.8
2001.....	155.9	229.4	350.1	558.4	905.3	1,426.7	2,263.9	3,669.9	6,169.3	14,597.6
2000.....	152.2	220.7	349.3	571.5	926.5	1,459.3	2,284.0	3,706.2	6,286.6	14,890.7
1999.....	147.4	219.9	343.6	569.0	925.9	1,483.5	2,308.5	3,711.2	6,284.5	14,900.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
All causes										
Black, both sexes										
2005 4/.....	379.3	602.7	914.9	1,254.6	1,812.2	2,466.3	3,514.3	5,161.7	7,547.7	13,083.1
2004 4/.....	386.6	608.3	910.1	1,271.1	1,813.5	2,509.7	3,582.0	5,266.0	7,493.0	13,338.5
2003 4/.....	410.4	632.9	940.3	1,295.1	1,889.9	2,595.8	3,754.9	5,421.2	7,745.3	13,976.7
2002.....	412.9	634.1	925.9	1,321.6	1,880.7	2,632.0	3,824.9	5,571.3	7,962.2	14,260.8
2001.....	428.4	644.1	928.5	1,328.9	1,891.7	2,738.8	3,904.2	5,660.2	8,028.9	14,550.8
2000.....	434.5	666.6	938.7	1,347.0	1,929.0	2,769.0	4,019.7	5,782.6	8,161.0	14,714.3
1999.....	448.6	668.3	929.9	1,399.3	1,969.2	2,813.7	4,148.0	5,878.3	8,169.4	14,757.9
Black, male										
2005 4/.....	463.8	755.9	1,184.8	1,667.1	2,380.7	3,174.7	4,509.2	6,579.2	9,376.2	13,809.8
2004 4/.....	467.1	766.8	1,186.6	1,668.7	2,374.4	3,242.7	4,587.3	6,747.0	9,229.2	14,452.5
2003 4/.....	507.1	800.8	1,227.5	1,688.7	2,454.5	3,334.8	4,843.9	6,978.2	9,827.8	14,903.4
2002.....	502.9	801.4	1,206.6	1,719.3	2,464.7	3,357.1	4,918.1	7,088.6	9,968.4	15,635.5
2001.....	527.8	830.2	1,201.4	1,718.2	2,424.6	3,497.3	5,053.2	7,255.7	10,256.3	16,439.9
2000.....	549.1	853.0	1,227.4	1,776.2	2,462.1	3,513.8	5,203.9	7,410.0	10,405.7	16,791.0
1999.....	572.7	885.6	1,220.1	1,822.6	2,549.7	3,599.1	5,336.3	7,538.1	10,433.8	17,304.5
Black, female										
2005 4/.....	304.8	469.0	686.6	914.1	1,369.2	1,940.8	2,833.5	4,316.0	6,581.9	12,789.9
2004 4/.....	315.7	470.1	676.4	943.8	1,376.3	1,965.5	2,899.8	4,388.5	6,578.2	12,896.9
2003 4/.....	325.1	486.7	697.2	971.3	1,449.4	2,047.3	3,020.5	4,495.8	6,654.1	13,616.7
2002.....	333.6	489.0	688.0	995.2	1,423.7	2,093.9	3,091.2	4,664.6	6,917.8	13,734.2
2001.....	340.6	482.8	696.2	1,009.5	1,472.5	2,178.4	3,138.8	4,707.8	6,894.3	13,849.9
2000.....	333.3	505.0	692.6	995.2	1,508.3	2,219.3	3,233.3	4,801.6	7,030.5	13,941.3
1999.....	339.3	480.9	683.9	1,053.5	1,514.0	2,231.6	3,359.6	4,878.6	7,047.2	13,805.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Certain other intestinal infections (A04,A07-A09)

All races 3/, both sexes

2005.....	0.1	0.1	0.4	0.5	1.2	2.8	6.5	11.6	21.5	45.8
2004.....	0.1	0.2	0.3	0.6	1.3	2.2	4.6	9.3	17.5	34.3
2003.....	*	0.1	0.2	0.4	0.9	1.7	3.3	7.0	12.3	24.8
2002.....	*	0.1	0.2	0.4	0.7	1.4	2.6	5.0	9.7	21.0
2001.....	*	*	0.2	0.2	0.5	0.8	1.8	3.2	6.4	12.8
2000.....	*	*	*	0.2	0.3	0.9	1.5	2.7	5.3	11.8
1999.....	*	*	*	*	0.3	0.7	1.2	2.3	4.5	10.3

All races 3/, male

2005.....	*	*	0.4	0.5	1.4	2.9	6.3	12.4	22.1	46.1
2004.....	*	*	0.3	0.6	1.3	2.2	4.3	9.3	17.8	32.0
2003.....	*	*	0.2	0.5	0.8	1.5	3.1	6.9	12.9	21.4
2002.....	*	*	*	0.4	0.8	1.5	3.1	5.1	10.1	20.4
2001.....	*	*	*	*	0.4	0.7	2.0	3.5	6.2	12.1
2000.....	*	*	*	0.3	*	1.0	1.6	2.9	5.8	11.4
1999.....	*	*	*	*	*	0.6	1.2	2.7	4.9	9.5

All races 3/, female

2005.....	*	*	0.4	0.6	1.0	2.6	6.6	11.0	21.1	45.6
2004.....	*	*	0.3	0.6	1.2	2.2	4.9	9.3	17.3	35.4
2003.....	*	*	0.2	0.4	0.9	1.8	3.5	7.1	12.0	26.3
2002.....	*	*	*	0.4	0.6	1.3	2.3	5.0	9.5	21.3
2001.....	*	*	0.2	*	0.5	0.8	1.8	3.0	6.5	13.1
2000.....	*	*	*	*	*	0.8	1.5	2.6	4.9	11.9
1999.....	*	*	*	*	*	0.8	1.3	2.0	4.3	10.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Certain other intestinal infections (A04,A07-A09)

White, both sexes

2005 4/.....	*	0.2	0.3	0.5	1.2	2.9	6.7	12.2	22.6	48.6
2004 4/.....	*	0.1	0.3	0.6	1.3	2.2	4.9	9.8	18.4	36.6
2003 4/.....	*	*	0.2	0.5	0.9	1.8	3.5	7.2	13.2	26.1
2002.....	*	0.1	0.2	0.4	0.6	1.5	2.7	5.3	10.3	22.0
2001.....	*	*	0.2	0.2	0.4	0.7	2.0	3.3	6.6	13.5
2000.....	*	*	*	0.3	0.3	0.9	1.6	2.9	5.6	12.2
1999.....	*	*	*	*	0.3	0.7	1.3	2.3	4.7	10.9

White, male

2005 4/.....	*	*	0.3	0.4	1.5	3.2	6.5	13.0	22.7	49.3
2004 4/.....	*	*	0.3	0.5	1.4	2.2	4.5	9.7	18.8	33.9
2003 4/.....	*	*	*	0.4	0.8	1.6	3.2	7.2	13.7	22.7
2002.....	*	*	*	0.4	0.7	1.6	3.0	5.3	10.7	21.4
2001.....	*	*	*	*	*	0.7	2.0	3.6	6.2	12.7
2000.....	*	*	*	*	*	1.0	1.6	3.0	6.0	11.9
1999.....	*	*	*	*	*	0.7	1.3	2.8	5.2	9.9

White, female

2005 4/.....	*	*	0.4	0.5	1.0	2.7	6.9	11.7	22.6	48.3
2004 4/.....	*	*	0.3	0.6	1.2	2.3	5.2	9.9	18.1	37.8
2003 4/.....	*	*	*	0.5	0.9	1.9	3.8	7.2	12.8	27.6
2002.....	*	*	*	0.4	0.6	1.4	2.4	5.3	10.1	22.3
2001.....	*	*	0.3	*	0.5	0.8	1.9	3.2	6.8	13.9
2000.....	*	*	*	*	*	0.8	1.6	2.7	5.3	12.3
1999.....	*	*	*	*	*	0.8	1.3	2.1	4.4	11.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Tuberculosis (A16-A19)										
All races 3/, both sexes										
2005.....	*	0.2	0.2	0.2	0.4	0.5	0.7	1.3	1.3	2.2
2004.....	0.1	0.1	0.2	0.2	0.3	0.6	1.0	1.1	1.5	2.4
2003.....	0.1	0.2	0.3	0.4	0.3	0.5	0.8	1.4	2.0	2.3
2002.....	0.1	0.2	0.2	0.3	0.5	0.9	0.9	1.5	2.3	2.6
2001.....	0.1	0.2	0.3	0.4	0.5	0.6	1.0	1.4	2.0	2.8
2000.....	0.1	0.2	0.3	0.2	0.5	0.7	1.0	1.8	2.4	2.7
1999.....	0.1	0.3	0.3	0.5	0.6	0.7	1.2	1.8	2.4	4.2
All races 3/, male										
2005.....	*	0.2	0.3	0.3	0.6	0.7	1.1	1.7	1.7	3.3
2004.....	0.2	0.2	0.3	0.3	0.6	0.8	1.4	1.2	2.2	3.8
2003.....	*	0.4	0.4	0.6	0.4	0.6	1.2	2.0	2.7	3.1
2002.....	0.2	0.2	0.4	0.5	0.7	1.2	1.2	1.8	3.2	4.0
2001.....	*	0.3	0.4	0.7	0.7	0.9	1.5	1.7	2.4	4.4
2000.....	*	0.4	0.5	0.4	0.7	1.0	1.1	2.3	3.8	4.6
1999.....	0.2	0.4	0.5	0.8	0.9	1.1	1.7	2.4	3.7	7.1
All races 3/, female										
2005.....	*	*	*	*	*	0.4	0.4	0.9	1.0	1.7
2004.....	*	*	*	*	*	*	0.6	1.0	1.1	1.7
2003.....	*	*	*	*	*	0.4	0.5	1.0	1.7	1.9
2002.....	*	*	*	*	0.4	0.5	0.6	1.3	1.7	2.1
2001.....	*	*	*	*	*	0.4	0.7	1.1	1.8	2.2
2000.....	*	*	*	*	*	0.5	0.9	1.4	1.6	2.0
1999.....	*	*	*	*	*	0.4	0.7	1.5	1.7	3.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Tuberculosis (A16-A19)										
White, both sexes										
2005 4/.....	*	0.1	0.1	*	0.2	0.4	0.4	0.8	0.9	1.7
2004 4/.....	*	*	0.1	*	0.2	0.3	0.7	0.8	1.2	1.7
2003 4/.....	*	0.1	0.1	0.3	0.2	0.3	0.7	1.0	1.7	1.8
2002.....	*	*	0.1	0.2	0.4	0.5	0.6	1.2	1.8	2.1
2001.....	*	0.1	0.1	0.3	0.3	0.5	0.7	1.1	1.4	2.2
2000.....	*	*	0.2	*	0.3	0.5	0.6	1.2	1.7	2.2
1999.....	*	0.2	0.2	0.3	0.4	0.6	0.8	1.3	2.0	3.2
White, male										
2005 4/.....	*	*	*	*	0.4	0.5	*	1.0	1.0	2.4
2004 4/.....	*	*	*	*	*	*	1.0	1.0	1.6	2.8
2003 4/.....	*	*	*	0.4	*	*	1.0	1.3	2.2	2.4
2002.....	*	*	*	*	0.5	0.8	1.0	1.3	2.5	3.2
2001.....	*	*	*	0.5	0.5	0.6	0.8	1.4	1.8	3.1
2000.....	*	*	0.3	*	*	0.7	0.8	1.5	2.3	3.8
1999.....	*	0.3	0.4	0.5	0.6	0.9	1.1	1.5	3.0	5.4
White, female										
2005 4/.....	*	*	*	*	*	*	*	0.6	0.8	1.4
2004 4/.....	*	*	*	*	*	*	*	0.7	0.9	1.3
2003 4/.....	*	*	*	*	*	*	*	0.9	1.3	1.6
2002.....	*	*	*	*	*	*	*	1.1	1.3	1.6
2001.....	*	*	*	*	*	*	0.5	0.9	1.2	1.8
2000.....	*	*	*	*	*	*	0.5	1.0	1.4	1.6
1999.....	*	*	*	*	*	*	0.5	1.1	1.4	2.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Respiratory tuberculosis (A16)										
All races 3/, both sexes										
2005.....	*	0.1	0.2	0.2	0.3	0.3	0.6	1.0	1.0	1.9
2004.....	0.1	0.1	0.2	0.2	0.3	0.4	0.8	0.8	1.1	2.2
2003.....	0.1	0.2	0.2	0.3	0.2	0.4	0.6	1.2	1.6	1.8
2002.....	0.1	0.1	0.2	0.2	0.4	0.6	0.7	1.2	2.0	2.3
2001.....	*	0.1	0.2	0.4	0.4	0.5	0.8	1.1	1.6	2.4
2000.....	0.1	0.1	0.2	0.2	0.3	0.5	0.8	1.4	2.1	2.3
1999.....	0.1	0.2	0.2	0.4	0.5	0.6	0.9	1.5	2.1	3.7
All races 3/, male										
2005.....	*	*	0.3	0.2	0.4	0.5	0.9	1.2	1.4	2.7
2004.....	*	*	0.3	*	0.4	0.6	1.2	0.9	1.7	3.7
2003.....	*	0.3	0.3	0.5	*	0.5	1.0	1.7	2.2	2.5
2002.....	0.2	*	0.3	0.4	0.5	1.0	0.9	1.3	2.8	3.5
2001.....	*	0.2	0.3	0.6	0.7	0.7	1.2	1.5	2.0	3.8
2000.....	*	0.3	0.4	0.3	0.6	0.9	0.9	1.9	3.2	4.1
1999.....	*	0.3	0.4	0.6	0.9	0.9	1.4	2.0	3.3	6.4
All races 3/, female										
2005.....	*	*	*	*	*	*	*	0.8	0.8	1.5
2004.....	*	*	*	*	*	*	0.5	0.7	0.8	1.5
2003.....	*	*	*	*	*	*	*	0.8	1.3	1.5
2002.....	*	*	*	*	*	*	0.5	1.1	1.5	1.8
2001.....	*	*	*	*	*	*	0.5	0.8	1.4	1.8
2000.....	*	*	*	*	*	*	0.6	1.1	1.4	1.6
1999.....	*	*	*	*	*	*	0.5	1.2	1.4	2.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Respiratory tuberculosis (A16)										
White, both sexes										
2005 4/.....	*	*	*	*	0.2	0.3	0.3	0.6	0.7	1.3
2004 4/.....	*	*	*	*	*	0.3	0.5	0.6	0.9	1.6
2003 4/.....	*	*	*	0.2	*	*	0.5	0.9	1.4	1.4
2002.....	*	*	*	*	0.3	0.4	0.5	0.9	1.5	1.8
2001.....	*	*	*	0.2	0.3	0.4	0.5	0.9	1.2	1.8
2000.....	*	*	0.2	*	*	0.3	0.5	1.0	1.5	1.9
1999.....	*	0.1	0.1	0.2	0.4	0.5	0.6	1.1	1.7	3.0
White, male										
2005 4/.....	*	*	*	*	*	*	*	*	*	1.9
2004 4/.....	*	*	*	*	*	*	0.8	0.8	1.2	2.6
2003 4/.....	*	*	*	0.3	*	*	0.9	1.1	1.8	1.8
2002.....	*	*	*	*	*	0.7	0.8	0.9	2.0	2.9
2001.....	*	*	*	0.4	0.5	*	0.7	1.2	1.4	2.6
2000.....	*	*	0.3	*	*	0.5	0.7	1.3	1.9	3.4
1999.....	*	*	*	0.4	0.5	0.7	1.0	1.3	2.6	5.2
White, female										
2005 4/.....	*	*	*	*	*	*	*	*	0.7	1.1
2004 4/.....	*	*	*	*	*	*	*	*	0.7	1.1
2003 4/.....	*	*	*	*	*	*	*	0.8	1.1	1.2
2002.....	*	*	*	*	*	*	*	0.9	1.1	1.3
2001.....	*	*	*	*	*	*	*	0.6	1.0	1.4
2000.....	*	*	*	*	*	*	*	0.7	1.2	1.2
1999.....	*	*	*	*	*	*	*	0.9	1.1	2.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Septicemia (A40-A41)										
All races 3/, both sexes										
2005.....	2.3	4.0	6.6	10.5	16.1	25.7	40.8	63.6	104.8	187.3
2004.....	2.3	4.1	6.9	10.3	16.2	24.9	41.2	65.1	103.6	186.7
2003.....	2.6	4.2	6.5	10.3	16.7	25.2	40.9	68.4	107.8	202.5
2002.....	2.3	4.2	6.2	10.0	16.0	27.0	43.1	70.0	109.4	203.0
2001.....	2.3	4.0	6.2	9.8	15.5	25.3	40.9	67.3	103.9	205.9
2000.....	2.5	3.9	6.1	9.4	15.2	24.2	38.4	64.0	105.0	215.7
1999.....	2.2	3.7	5.6	9.0	14.4	24.7	38.3	63.8	102.9	220.7
All races 3/, male										
2005.....	2.4	4.4	7.1	11.5	17.8	28.6	45.9	71.7	119.8	201.9
2004.....	2.4	4.5	7.6	11.4	17.1	26.9	46.6	74.5	120.9	203.9
2003.....	2.5	4.6	7.1	11.3	17.7	28.7	45.1	76.7	122.5	213.8
2002.....	2.5	4.7	6.8	10.6	17.8	29.1	48.2	80.6	127.0	213.9
2001.....	2.5	4.3	6.7	10.6	16.6	27.8	46.7	78.6	122.3	224.3
2000.....	2.6	4.4	7.0	10.1	16.7	26.9	43.2	73.9	121.7	223.0
1999.....	2.6	4.2	6.2	9.7	15.7	27.7	43.7	74.8	120.6	233.5
All races 3/, female										
2005.....	2.3	3.6	6.1	9.5	14.5	23.2	36.7	57.8	95.4	180.6
2004.....	2.3	3.7	6.4	9.3	15.5	23.2	36.8	58.4	92.9	179.0
2003.....	2.7	3.8	6.0	9.4	15.7	22.2	37.5	62.5	98.8	197.5
2002.....	2.2	3.8	5.6	9.5	14.5	25.2	39.0	62.5	98.7	198.3
2001.....	2.1	3.8	5.7	9.1	14.4	23.1	36.2	59.5	92.9	198.3
2000.....	2.3	3.4	5.2	8.7	13.8	21.8	34.6	57.1	95.2	212.7
1999.....	1.9	3.3	5.1	8.4	13.1	22.1	34.1	56.2	92.5	215.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Septicemia (A40-A41)										
White, both sexes										
2005 4/.....	1.9	3.3	5.4	9.0	14.0	22.9	37.5	58.9	98.3	177.3
2004 4/.....	1.8	3.2	5.7	8.8	13.8	21.9	37.6	60.7	97.5	178.1
2003 4/.....	2.1	3.4	5.2	8.8	14.3	21.7	36.9	63.7	102.5	191.1
2002.....	1.9	3.6	5.0	8.4	13.5	23.6	39.2	64.4	103.1	191.9
2001.....	1.8	3.2	4.9	8.1	13.2	22.2	36.8	62.1	98.5	195.3
2000.....	2.0	3.0	4.9	7.8	12.6	20.7	34.5	58.0	97.4	203.0
1999.....	1.7	2.8	4.5	7.2	12.1	21.3	33.6	57.6	95.5	205.9
White, male										
2005 4/.....	2.0	3.6	5.7	9.7	15.8	25.2	41.8	66.7	113.0	190.7
2004 4/.....	1.8	3.5	6.4	9.7	14.4	23.7	42.5	69.8	113.4	196.5
2003 4/.....	2.0	3.8	5.7	9.3	15.3	24.6	40.8	71.3	116.7	203.3
2002.....	2.0	4.0	5.4	8.9	15.3	25.2	44.0	73.4	119.0	202.9
2001.....	2.1	3.3	5.3	8.5	14.4	24.4	41.6	72.5	116.0	211.8
2000.....	2.2	3.5	5.5	8.4	14.0	23.1	39.4	66.7	114.0	209.8
1999.....	2.0	3.1	4.9	7.6	12.8	23.7	38.2	68.3	111.6	217.4
White, female										
2005 4/.....	1.8	2.9	5.1	8.3	12.4	20.8	33.9	53.1	89.2	171.0
2004 4/.....	1.8	2.9	5.1	8.0	13.3	20.2	33.5	54.1	87.7	169.9
2003 4/.....	2.2	3.0	4.7	8.2	13.4	19.2	33.7	58.2	93.7	185.8
2002.....	1.9	3.1	4.5	7.8	11.8	22.3	35.2	58.0	93.4	187.1
2001.....	1.6	3.0	4.5	7.7	12.2	20.4	32.9	54.8	87.9	188.3
2000.....	1.9	2.5	4.3	7.2	11.3	18.7	30.6	51.9	87.6	200.2
1999.....	1.4	2.4	4.1	6.9	11.5	19.2	29.9	50.1	86.1	201.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Septicemia (A40-A41)										
Black, both sexes										
2005 4/.....	5.5	10.0	16.6	25.5	38.4	55.8	81.2	125.4	203.7	336.8
2004 4/.....	6.2	10.3	17.3	25.5	39.9	57.9	85.1	128.0	196.1	320.3
2003 4/.....	6.3	10.5	17.5	25.3	41.5	61.0	88.0	133.4	193.4	372.0
2002.....	5.5	9.5	17.0	27.2	40.9	62.0	90.6	146.5	208.8	367.0
2001.....	5.6	10.6	16.9	27.1	38.4	57.4	89.6	139.1	193.1	364.8
2000.....	5.8	11.0	16.9	24.3	40.2	59.7	86.6	146.4	216.3	392.6
1999.....	6.2	11.0	15.5	25.4	36.3	57.9	94.6	142.8	209.5	412.3
Black, male										
2005 4/.....	5.5	10.9	19.5	30.5	43.4	68.2	97.1	143.5	238.8	374.6
2004 4/.....	6.7	12.0	19.0	30.4	44.5	65.2	102.3	153.8	244.6	341.7
2003 4/.....	6.4	11.8	19.7	31.8	46.1	74.9	100.2	160.2	235.5	404.5
2002.....	6.1	10.3	20.0	29.0	45.1	71.6	105.5	183.7	259.8	393.5
2001.....	5.7	11.6	19.1	33.3	41.5	66.3	111.1	171.0	236.8	424.4
2000.....	5.8	12.4	21.3	27.8	45.1	68.9	97.0	182.9	250.4	427.8
1999.....	7.7	12.8	18.1	29.7	46.5	71.4	115.0	166.2	266.8	453.5
Black, female										
2005 4/.....	5.5	9.2	14.1	21.3	34.4	46.5	70.3	114.7	185.1	321.5
2004 4/.....	5.7	8.9	15.8	21.5	36.4	52.5	73.4	112.7	170.5	311.9
2003 4/.....	6.3	9.4	15.6	20.0	37.9	50.8	79.8	117.5	171.4	359.3
2002.....	5.0	8.9	14.4	25.7	37.6	54.9	80.7	124.3	182.2	356.9
2001.....	5.4	9.7	15.0	22.1	36.0	50.9	75.3	120.1	170.9	342.7
2000.....	5.8	9.7	13.2	21.5	36.3	52.8	79.7	124.5	199.1	379.5
1999.....	5.0	9.4	13.3	21.9	28.3	47.9	81.0	128.7	181.1	396.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Viral hepatitis (B15-B19)										
All races 3/, both sexes										
2005.....	1.7	4.2	6.8	5.3	3.5	3.2	3.8	3.6	3.8	2.4
2004.....	2.0	4.3	6.5	4.9	3.5	3.3	3.6	3.6	3.6	2.7
2003.....	2.1	4.8	6.3	4.5	3.1	3.4	4.3	4.3	4.1	2.4
2002.....	2.3	5.6	6.1	4.3	3.7	3.9	4.5	4.7	4.4	3.9
2001.....	2.6	5.3	5.2	4.1	3.8	4.3	4.9	5.1	4.6	3.8
2000.....	2.5	5.2	4.8	3.7	3.8	4.3	4.9	5.4	5.2	4.0
1999.....	2.3	4.4	3.8	3.3	3.3	4.5	5.1	5.3	5.4	4.7
All races 3/, male										
2005.....	2.3	6.3	10.2	8.2	4.7	3.5	4.0	3.3	4.2	2.6
2004.....	2.8	6.4	9.9	7.3	4.5	4.1	3.7	3.8	3.3	3.0
2003.....	2.9	7.1	9.7	6.6	3.9	3.7	4.8	4.7	5.0	2.8
2002.....	3.2	8.5	9.5	6.2	4.6	4.3	4.7	5.7	4.8	5.0
2001.....	3.7	8.1	8.2	5.9	4.5	4.7	4.9	5.1	5.5	4.4
2000.....	3.5	8.0	7.4	5.2	4.7	4.6	5.0	6.5	6.2	5.7
1999.....	3.5	6.7	5.9	4.7	3.8	4.9	5.4	6.1	7.3	6.9
All races 3/, female										
2005.....	1.1	2.2	3.6	2.7	2.3	2.9	3.7	3.8	3.6	2.3
2004.....	1.3	2.3	3.3	2.5	2.6	2.6	3.5	3.4	3.7	2.5
2003.....	1.2	2.6	3.1	2.5	2.4	3.1	3.9	4.0	3.5	2.3
2002.....	1.4	2.8	2.8	2.5	2.9	3.4	4.4	4.1	4.2	3.3
2001.....	1.5	2.5	2.4	2.5	3.3	3.9	4.9	5.0	4.1	3.5
2000.....	1.5	2.5	2.4	2.2	3.1	4.1	4.9	4.6	4.7	3.3
1999.....	1.2	2.2	1.7	2.1	2.8	4.1	4.9	4.7	4.2	3.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Viral hepatitis (B15-B19)										
White, both sexes										
2005 4/.....	1.7	4.3	6.7	4.9	3.0	2.7	3.2	3.3	3.6	2.2
2004 4/.....	2.1	4.3	6.2	4.2	3.1	3.0	3.2	3.3	3.1	2.5
2003 4/.....	2.1	4.9	6.0	3.8	2.7	2.9	3.8	4.0	3.8	2.3
2002.....	2.2	5.5	5.8	3.8	3.2	3.4	4.1	4.5	4.3	3.7
2001.....	2.6	5.2	4.7	3.7	3.5	3.8	4.4	4.9	4.4	3.6
2000.....	2.4	4.9	4.3	3.0	3.4	3.9	4.6	5.3	5.0	3.9
1999.....	2.2	4.2	3.3	2.9	2.9	4.0	4.6	4.9	5.3	4.5
White, male										
2005 4/.....	2.4	6.4	10.0	7.6	4.0	3.1	3.4	3.3	3.9	2.4
2004 4/.....	2.9	6.4	9.4	6.4	4.0	3.7	3.4	3.6	2.8	2.7
2003 4/.....	2.9	7.2	9.2	5.7	3.4	3.3	4.3	4.2	4.8	2.8
2002.....	3.1	8.3	9.1	5.5	3.9	4.0	4.3	5.5	4.8	4.9
2001.....	3.7	8.1	7.6	5.4	4.1	4.4	4.6	5.2	5.6	4.5
2000.....	3.4	7.6	6.6	4.4	4.2	4.0	4.6	6.4	5.9	5.7
1999.....	3.3	6.1	5.2	4.1	3.6	4.6	5.1	5.8	7.7	6.5
White, female										
2005 4/.....	1.1	2.2	3.4	2.4	2.0	2.4	3.0	3.2	3.4	2.2
2004 4/.....	1.3	2.3	3.0	2.2	2.4	2.4	3.0	3.1	3.3	2.4
2003 4/.....	1.2	2.6	2.9	2.0	2.0	2.6	3.5	3.8	3.2	2.1
2002.....	1.3	2.6	2.7	2.2	2.6	2.8	3.9	3.8	4.0	3.2
2001.....	1.4	2.4	2.0	2.1	2.9	3.3	4.2	4.6	3.7	3.2
2000.....	1.4	2.3	2.0	1.8	2.7	3.7	4.5	4.5	4.4	3.2
1999.....	1.1	2.2	1.5	1.7	2.3	3.5	4.3	4.3	3.9	3.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Viral hepatitis (B15-B19)

Black, both sexes

2005 4/.....	1.7	4.7	9.7	9.2	7.2	5.8	6.4	5.5	4.9	*
2004 4/.....	2.0	5.2	10.3	10.2	6.5	4.2	5.1	3.9	6.7	*
2003 4/.....	2.7	5.2	10.1	9.4	6.3	5.1	5.7	6.5	5.3	*
2002.....	2.8	7.2	9.0	7.7	6.3	6.1	6.2	6.0	*	*
2001.....	3.3	6.8	9.4	7.3	6.0	5.1	6.2	6.1	*	*
2000.....	3.1	7.3	8.8	6.9	4.6	5.2	6.3	5.7	7.1	*
1999.....	3.5	7.0	6.7	5.7	4.2	6.6	7.2	6.8	*	6.4

Black, male

2005 4/.....	2.2	6.7	14.5	14.3	10.5	6.7	7.9	*	*	*
2004 4/.....	2.2	8.3	16.0	16.6	9.9	6.8	6.5	*	*	*
2003 4/.....	3.6	7.6	16.2	14.9	7.5	5.4	6.9	*	*	*
2002.....	3.8	10.5	14.6	12.4	8.1	8.0	*	*	*	*
2001.....	4.7	10.1	13.9	10.2	7.3	*	*	*	*	*
2000.....	4.2	11.0	13.9	10.4	5.2	6.8	7.4	*	*	*
1999.....	5.1	11.9	11.2	7.6	4.5	5.5	*	*	*	*

Black, female

2005 4/.....	*	3.0	5.7	5.0	4.7	5.1	5.4	7.5	*	*
2004 4/.....	1.8	2.5	5.5	4.9	3.9	*	*	*	*	*
2003 4/.....	1.9	3.2	5.0	4.8	5.3	4.9	4.9	5.7	*	*
2002.....	1.9	4.4	4.2	3.9	4.8	4.6	6.3	5.8	*	*
2001.....	2.0	3.9	5.5	4.9	5.0	5.4	6.7	7.2	*	*
2000.....	2.1	4.0	4.4	4.0	4.1	4.0	5.5	*	*	*
1999.....	2.0	2.8	2.8	4.2	4.0	7.5	7.7	6.6	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Human immunodeficiency virus (HIV) disease (B20-B24)

All races 3/, both sexes

2005.....	11.7	11.3	9.9	6.3	4.0	2.8	1.8	1.0	0.5	*
2004.....	12.3	11.5	9.6	6.3	4.2	3.1	1.5	1.1	0.5	*
2003.....	12.8	12.1	9.5	6.4	4.1	2.8	1.9	1.0	0.4	*
2002.....	13.4	12.6	9.6	5.9	4.0	2.9	1.4	1.0	0.5	*
2001.....	13.7	12.1	8.7	5.8	4.5	2.7	1.4	0.9	0.4	*
2000.....	13.5	12.7	9.0	5.9	4.1	2.6	1.7	0.9	*	*
1999.....	14.6	12.4	8.7	5.9	3.5	2.7	1.5	0.7	*	*

All races 3/, male

2005.....	17.2	17.2	15.4	10.5	6.6	4.8	3.1	1.8	*	*
2004.....	17.9	17.5	14.9	10.5	7.0	5.1	2.6	1.7	*	*
2003.....	19.1	18.8	15.3	10.7	6.9	4.6	3.3	1.8	1.0	*
2002.....	19.8	19.4	15.7	9.8	6.8	5.0	2.6	1.7	*	*
2001.....	20.5	19.0	14.4	9.8	7.1	4.2	2.7	1.8	*	*
2000.....	20.4	20.3	15.0	10.1	7.0	4.5	2.9	1.8	*	*
1999.....	22.6	20.0	14.5	10.2	5.9	4.8	2.7	1.3	*	*

All races 3/, female

2005.....	6.2	5.6	4.6	2.4	1.6	1.1	0.7	*	*	*
2004.....	6.7	5.7	4.5	2.3	1.5	1.4	0.6	0.6	*	*
2003.....	6.6	5.6	3.9	2.5	1.6	1.3	0.7	*	*	*
2002.....	7.2	5.9	3.6	2.3	1.5	1.0	0.5	*	*	*
2001.....	7.0	5.4	3.3	2.0	2.0	1.4	0.4	*	*	*
2000.....	6.7	5.3	3.4	2.0	1.5	1.0	0.7	*	*	*
1999.....	6.9	5.0	3.1	1.9	1.2	1.0	0.6	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Human immunodeficiency virus (HIV) disease (B20-B24)

White, both sexes

2005 4/.....	6.7	5.9	5.1	3.2	1.9	1.4	0.9	0.6	*	*
2004 4/.....	6.8	6.1	4.8	3.3	2.0	1.5	0.8	0.6	*	*
2003 4/.....	7.4	6.7	4.9	3.5	2.1	1.3	1.0	0.6	*	*
2002.....	7.3	6.8	4.6	3.3	2.2	1.3	0.6	0.5	*	*
2001.....	7.5	6.4	4.5	3.3	2.0	1.2	0.8	0.5	*	*
2000.....	7.5	6.9	4.7	3.4	2.1	1.3	0.7	0.5	*	*
1999.....	8.1	6.5	4.9	3.1	1.9	1.4	0.7	0.4	*	*

White, male

2005 4/.....	11.0	10.0	8.4	5.6	3.4	2.5	1.7	1.1	*	*
2004 4/.....	11.2	10.1	8.2	5.9	3.7	2.7	1.5	1.2	*	*
2003 4/.....	12.1	11.3	8.4	6.0	3.6	2.2	2.0	1.3	*	*
2002.....	12.0	11.4	8.0	5.8	3.8	2.4	1.2	0.8	*	*
2001.....	12.6	10.8	8.0	5.9	3.4	2.1	1.7	1.0	*	*
2000.....	12.3	11.7	8.3	6.0	3.7	2.5	1.3	1.0	*	*
1999.....	13.4	11.0	8.6	5.6	3.5	2.7	1.2	0.8	*	*

White, female

2005 4/.....	2.4	1.9	1.9	0.9	0.5	*	*	*	*	*
2004 4/.....	2.4	2.1	1.5	0.8	0.4	0.5	*	*	*	*
2003 4/.....	2.6	2.1	1.3	1.1	0.6	0.5	*	*	*	*
2002.....	2.6	2.2	1.3	0.9	0.7	*	*	*	*	*
2001.....	2.4	2.0	1.0	0.9	0.7	0.5	*	*	*	*
2000.....	2.7	2.1	1.2	0.9	0.7	*	*	*	*	*
1999.....	2.8	2.0	1.2	0.7	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Human immunodeficiency virus (HIV) disease (B20-B24)

Black, both sexes

2005 4/.....	47.2	51.2	47.8	33.8	24.0	16.6	10.5	6.0	*	*
2004 4/.....	51.4	52.1	47.4	33.4	24.6	17.6	8.5	6.3	*	*
2003 4/.....	52.3	53.4	47.2	33.9	23.8	17.2	11.0	4.9	*	*
2002.....	57.3	57.4	49.7	30.2	20.7	17.5	10.5	7.0	*	*
2001.....	58.8	56.5	44.7	28.9	26.2	16.3	8.5	6.4	*	*
2000.....	57.3	58.7	46.6	29.3	22.0	15.1	11.6	6.2	*	*
1999.....	62.8	59.0	42.6	32.7	18.3	15.4	11.2	4.8	*	*

Black, male

2005 4/.....	65.8	74.8	75.4	57.8	40.7	29.7	18.7	11.6	*	*
2004 4/.....	69.9	76.9	72.4	56.1	42.0	29.6	15.2	9.6	*	*
2003 4/.....	74.0	79.5	75.7	57.6	41.3	30.0	20.0	*	*	*
2002.....	79.6	86.3	83.5	50.0	37.0	31.4	19.4	14.7	*	*
2001.....	82.8	87.5	72.2	49.9	43.5	26.4	15.5	13.0	*	*
2000.....	84.1	92.7	77.4	51.0	39.5	26.3	22.4	11.8	*	*
1999.....	94.9	96.1	70.6	57.5	30.6	27.3	21.4	*	*	*

Black, female

2005 4/.....	30.8	30.6	24.6	14.0	10.9	6.9	5.0	*	*	*
2004 4/.....	35.0	30.5	26.3	14.7	10.9	8.6	*	*	*	*
2003 4/.....	33.1	30.7	23.1	14.3	10.2	7.7	4.9	*	*	*
2002.....	37.6	32.4	21.0	13.9	7.9	7.2	4.5	*	*	*
2001.....	37.5	29.7	21.2	11.7	12.6	8.8	*	*	*	*
2000.....	33.7	29.2	20.4	11.5	8.2	6.9	4.4	*	*	*
1999.....	34.6	26.9	18.8	12.5	8.7	6.5	4.5	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1, A85.8,A86-B04,B06-B09,B25-B49,B55-B99)

All races 3/, both sexes

2005.....	1.1	2.4	4.2	3.9	4.5	5.9	8.3	12.0	15.8	21.2
2004.....	1.2	2.5	3.5	4.2	4.4	6.0	8.3	11.2	15.9	22.5
2003.....	1.3	2.4	3.5	3.7	4.5	5.9	8.9	12.3	14.8	22.5
2002.....	1.3	1.9	2.7	3.0	3.8	5.4	8.1	11.6	13.8	21.7
2001.....	0.9	1.5	1.9	2.4	3.9	5.6	6.9	9.4	12.3	20.1
2000.....	0.8	1.4	1.8	2.4	3.6	4.7	7.0	10.0	12.9	21.2
1999.....	1.4	2.0	2.6	3.3	4.4	6.8	8.9	12.6	16.6	27.0

All races 3/, male

2005.....	1.5	3.3	5.8	5.2	5.3	7.0	10.0	14.1	19.6	23.6
2004.....	1.5	3.3	4.9	5.6	5.2	7.2	9.9	12.9	18.4	29.6
2003.....	1.6	3.3	4.7	4.9	5.4	7.0	10.3	13.9	18.8	26.5
2002.....	1.7	2.6	3.8	3.8	4.7	6.5	9.7	14.0	17.7	26.8
2001.....	1.2	2.1	2.4	3.0	4.5	6.2	7.5	11.4	15.9	23.6
2000.....	1.0	1.7	2.3	2.8	4.0	5.7	8.8	12.1	16.2	27.6
1999.....	1.7	2.4	3.0	3.5	5.2	8.0	10.5	15.3	20.4	31.9

All races 3/, female

2005.....	0.7	1.5	2.5	2.7	3.8	5.0	7.0	10.5	13.4	20.1
2004.....	0.9	1.6	2.2	2.8	3.6	5.0	7.0	10.0	14.4	19.3
2003.....	0.9	1.6	2.2	2.6	3.7	5.0	7.8	11.1	12.4	20.8
2002.....	0.8	1.2	1.7	2.2	3.0	4.5	6.8	9.9	11.5	19.5
2001.....	0.6	0.9	1.5	1.8	3.3	5.1	6.5	8.0	10.1	18.7
2000.....	0.6	1.1	1.3	2.1	3.2	3.9	5.6	8.5	10.9	18.6
1999.....	1.1	1.6	2.3	3.1	3.7	5.9	7.7	10.8	14.4	25.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1, A85.8,A86-B04,B06-B09,B25-B49,B55-B99)

White, both sexes										
2005 4/.....	1.1	2.2	4.1	3.8	4.3	5.9	8.0	11.7	15.6	21.2
2004 4/.....	1.1	2.5	3.5	4.0	4.1	5.7	8.0	10.9	15.7	22.5
2003 4/.....	1.2	2.3	3.3	3.5	4.2	5.9	8.9	11.8	14.7	23.0
2002.....	1.1	1.8	2.5	2.6	3.6	5.4	7.9	11.3	13.8	22.0
2001.....	0.8	1.4	1.7	2.2	3.6	5.4	6.7	9.3	12.1	20.1
2000.....	0.7	1.3	1.6	2.3	3.2	4.6	6.8	9.8	12.7	21.0
1999.....	1.1	1.6	2.2	2.9	3.9	6.4	8.5	12.1	16.1	26.5
White, male										
2005 4/.....	1.4	3.1	5.9	5.0	5.0	7.0	9.8	13.8	19.4	23.7
2004 4/.....	1.4	3.3	4.9	5.2	4.9	7.1	9.4	12.6	18.1	30.1
2003 4/.....	1.6	3.2	4.5	4.6	5.0	6.8	10.0	13.5	18.6	27.2
2002.....	1.5	2.5	3.5	3.2	4.3	6.6	9.4	13.6	17.6	27.2
2001.....	1.0	1.9	2.2	2.8	4.3	6.0	7.2	11.3	15.9	23.8
2000.....	0.9	1.5	2.0	2.7	3.6	5.5	8.6	11.6	16.1	26.9
1999.....	1.4	2.0	2.6	3.3	4.5	7.5	10.2	14.8	18.6	30.6
White, female										
2005 4/.....	0.7	1.4	2.4	2.7	3.6	5.0	6.6	10.2	13.3	20.0
2004 4/.....	0.9	1.6	2.2	2.8	3.4	4.6	6.9	9.7	14.1	19.0
2003 4/.....	0.8	1.5	2.1	2.4	3.5	5.0	8.0	10.6	12.3	21.2
2002.....	0.7	1.1	1.5	2.0	2.9	4.3	6.7	9.7	11.5	19.8
2001.....	0.6	0.8	1.3	1.6	3.1	5.0	6.2	7.8	9.9	18.5
2000.....	0.6	1.0	1.1	1.9	2.8	3.8	5.4	8.6	10.8	18.7
1999.....	0.7	1.2	1.9	2.6	3.4	5.6	7.2	10.2	14.6	24.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1, A85.8,A86-B04,B06-B09,B25-B49,B55-B99)

Black, both sexes

2005 4/.....	1.7	3.4	5.0	5.5	6.6	6.3	9.6	13.4	17.2	22.7
2004 4/.....	2.1	2.9	3.9	5.8	6.7	7.5	10.2	14.7	19.1	24.3
2003 4/.....	2.1	3.6	5.3	6.0	6.2	5.9	8.3	16.2	16.2	19.5
2002.....	2.5	3.2	4.8	6.0	5.4	5.3	9.7	15.3	11.5	21.7
2001.....	1.8	2.5	3.4	4.4	6.0	7.1	9.8	9.7	13.3	20.6
2000.....	1.7	2.6	3.5	4.0	5.9	6.1	9.9	11.6	16.1	24.1
1999.....	3.6	5.1	5.9	6.3	8.2	10.7	13.1	18.7	23.4	34.1

Black, male

2005 4/.....	2.5	4.3	6.1	7.8	8.4	8.3	10.8	18.8	20.4	23.2
2004 4/.....	2.4	4.1	5.9	8.7	9.2	7.5	14.6	18.7	21.5	29.8
2003 4/.....	2.6	4.6	7.1	7.7	8.3	8.9	13.1	16.4	22.1	22.9
2002.....	3.2	4.5	6.7	9.2	7.3	6.5	12.0	18.4	*	28.9
2001.....	2.7	3.6	4.3	5.5	7.3	9.7	10.4	10.2	16.1	*
2000.....	2.1	3.2	4.8	4.4	7.2	8.6	12.4	17.0	20.2	32.3
1999.....	3.6	5.8	6.5	5.8	10.4	12.5	15.9	21.8	41.9	45.7

Black, female

2005 4/.....	*	2.6	4.0	3.5	5.2	4.8	8.9	10.1	15.6	22.5
2004 4/.....	1.8	1.9	2.3	3.4	4.8	7.6	7.2	12.4	17.8	22.1
2003 4/.....	1.6	2.6	3.8	4.6	4.5	3.7	5.1	16.0	13.2	18.2
2002.....	1.9	2.0	3.1	3.4	3.9	4.4	8.1	13.4	9.9	18.9
2001.....	*	1.6	2.7	3.6	5.0	5.3	9.4	9.4	11.9	20.2
2000.....	*	2.1	2.4	3.6	4.9	4.2	8.2	8.3	14.0	21.0
1999.....	3.5	4.5	5.3	6.8	6.5	9.4	11.3	16.9	14.3	29.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms (C00-C97)										
All races 3/, both sexes										
2005.....	44.6	86.5	154.8	255.6	422.1	622.2	886.3	1,158.4	1,427.7	1,637.7
2004.....	44.3	86.7	155.6	261.5	427.6	637.8	892.4	1,172.1	1,424.8	1,653.3
2003.....	46.1	89.8	159.1	267.9	440.9	647.9	909.2	1,195.4	1,450.0	1,698.2
2002.....	47.5	90.6	161.5	276.8	447.0	663.8	933.6	1,207.0	1,458.3	1,723.9
2001.....	48.7	92.4	164.8	280.7	453.3	673.2	943.5	1,211.7	1,465.9	1,765.6
2000.....	48.8	93.6	166.2	289.0	463.4	690.6	951.6	1,222.5	1,505.3	1,819.4
1999.....	49.4	93.7	167.0	295.3	471.6	700.2	963.4	1,224.0	1,493.1	1,805.8
All races 3/, male										
2005.....	39.5	83.4	164.9	284.8	484.5	743.6	1,091.8	1,475.1	1,901.6	2,319.3
2004.....	39.2	86.1	168.2	289.9	492.5	756.9	1,091.6	1,503.3	1,894.3	2,349.5
2003.....	41.7	90.6	169.8	295.9	507.6	774.4	1,117.9	1,533.3	1,939.7	2,413.8
2002.....	42.8	89.8	171.7	308.6	519.7	796.8	1,157.8	1,554.1	1,951.5	2,491.1
2001.....	44.0	91.7	174.2	311.6	526.7	814.1	1,178.0	1,568.4	1,978.7	2,613.6
2000.....	44.4	93.5	173.9	321.3	535.7	834.7	1,190.4	1,586.0	2,050.4	2,710.7
1999.....	45.5	94.7	174.9	330.6	550.1	848.2	1,212.9	1,600.1	2,046.0	2,722.2
All races 3/, female										
2005.....	49.6	89.6	145.1	228.1	365.2	516.2	719.8	928.5	1,133.4	1,324.6
2004.....	49.2	87.4	143.5	234.8	368.5	534.2	732.0	934.2	1,135.5	1,340.1
2003.....	50.5	89.1	148.8	241.5	380.3	538.1	741.8	954.8	1,150.5	1,381.9
2002.....	52.2	91.4	151.8	246.8	381.1	548.9	754.4	961.9	1,158.4	1,391.1
2001.....	53.5	93.2	155.9	251.7	386.7	551.9	757.2	962.0	1,158.6	1,410.7
2000.....	53.2	93.7	158.8	258.9	398.0	567.2	763.5	969.3	1,183.8	1,456.4
1999.....	53.3	92.7	159.5	262.3	400.6	573.7	768.4	962.5	1,169.5	1,436.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms (C00-C97)										
White, both sexes										
2005 4/.....	43.0	82.0	147.8	247.2	414.2	618.5	888.9	1,164.9	1,432.7	1,645.4
2004 4/.....	42.0	82.2	148.2	253.8	421.6	633.6	891.5	1,174.1	1,429.3	1,659.7
2003 4/.....	44.0	84.8	151.3	261.2	434.5	644.8	908.5	1,195.1	1,450.7	1,701.4
2002.....	45.5	86.2	154.7	268.7	439.5	659.2	931.4	1,205.1	1,456.9	1,725.7
2001.....	46.4	87.8	158.4	272.2	447.1	665.6	938.4	1,207.5	1,458.8	1,765.2
2000.....	45.8	88.2	159.5	281.1	457.1	685.0	944.1	1,214.2	1,503.0	1,815.7
1999.....	46.8	88.5	159.9	285.8	462.2	693.1	955.3	1,213.0	1,488.4	1,798.8
White, male										
2005 4/.....	38.7	79.2	156.6	271.7	470.2	731.9	1,083.4	1,472.2	1,894.9	2,322.7
2004 4/.....	37.4	81.1	158.7	278.1	479.7	745.4	1,077.9	1,490.7	1,887.6	2,348.9
2003 4/.....	39.8	85.1	159.7	285.4	494.7	763.2	1,108.1	1,518.9	1,930.1	2,412.1
2002.....	41.2	85.4	162.8	296.5	503.4	785.6	1,145.2	1,537.3	1,934.6	2,486.8
2001.....	41.9	86.5	165.1	298.2	513.9	797.6	1,160.4	1,547.8	1,950.7	2,597.6
2000.....	41.4	87.7	164.0	307.0	522.0	819.7	1,166.6	1,559.2	2,026.9	2,693.7
1999.....	43.1	88.0	164.3	316.0	532.7	831.3	1,191.9	1,571.1	2,024.1	2,696.3
White, female										
2005 4/.....	47.3	84.8	139.2	223.6	362.1	517.7	728.2	937.5	1,143.0	1,333.6
2004 4/.....	46.6	83.4	137.9	230.5	367.7	534.4	738.3	942.5	1,144.7	1,348.9
2003 4/.....	48.3	84.6	143.1	237.8	378.9	540.3	745.2	961.0	1,155.2	1,386.5
2002.....	49.9	86.9	146.8	242.0	380.4	548.0	757.1	967.4	1,164.2	1,395.1
2001.....	51.0	89.0	152.0	247.4	385.4	549.7	758.6	966.4	1,161.4	1,416.7
2000.....	50.1	88.7	155.2	256.5	397.2	567.2	765.6	971.5	1,191.3	1,459.1
1999.....	50.5	89.1	155.5	257.1	397.2	572.5	767.3	961.5	1,172.1	1,438.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms (C00-C97)										
Black, both sexes										
2005 4/.....	60.6	129.1	231.4	374.3	573.3	782.6	1,026.2	1,296.6	1,585.5	1,744.7
2004 4/.....	64.2	129.6	235.7	377.1	570.2	803.0	1,059.1	1,355.6	1,578.2	1,774.0
2003 4/.....	66.5	138.3	243.2	382.6	589.1	802.7	1,080.1	1,385.5	1,640.8	1,853.1
2002.....	68.3	135.1	238.2	399.6	597.1	833.2	1,126.1	1,425.8	1,665.1	1,885.7
2001.....	70.4	137.8	241.5	404.7	589.4	858.4	1,156.8	1,433.0	1,716.9	1,931.7
2000.....	74.4	145.6	251.9	407.9	604.6	861.4	1,179.2	1,481.7	1,700.6	1,998.9
1999.....	73.3	144.5	255.6	429.5	633.9	879.1	1,191.7	1,502.1	1,704.2	2,008.4
Black, male										
2005 4/.....	50.0	127.0	258.6	459.7	729.5	1,024.8	1,395.3	1,803.3	2,354.0	2,683.7
2004 4/.....	56.2	134.0	274.7	455.9	729.5	1,035.5	1,442.5	1,926.3	2,340.8	2,748.8
2003 4/.....	61.2	145.6	283.0	464.8	747.3	1,045.9	1,455.8	1,975.5	2,417.1	2,825.5
2002.....	60.4	137.1	271.5	484.4	777.8	1,079.3	1,536.2	2,047.9	2,514.3	2,976.1
2001.....	62.8	143.6	280.7	494.9	746.2	1,135.3	1,600.3	2,084.3	2,645.3	3,209.9
2000.....	70.3	151.8	293.7	512.8	769.2	1,137.3	1,654.9	2,166.0	2,683.2	3,264.8
1999.....	68.3	160.5	300.8	533.7	815.7	1,159.8	1,644.1	2,205.4	2,642.0	3,340.7
Black, female										
2005 4/.....	70.0	131.0	208.4	303.9	451.6	603.0	773.5	994.4	1,179.6	1,365.8
2004 4/.....	71.2	125.9	202.6	312.3	446.1	630.3	798.9	1,017.5	1,176.3	1,387.7
2003 4/.....	71.1	132.0	209.5	314.9	465.7	622.1	826.8	1,034.8	1,234.0	1,475.3
2002.....	75.2	133.5	210.0	330.0	455.6	650.5	850.9	1,054.1	1,223.1	1,468.0
2001.....	77.1	132.8	208.1	330.6	466.1	653.9	861.4	1,044.2	1,244.0	1,457.5
2000.....	78.1	140.2	216.2	321.9	474.8	657.8	863.3	1,069.2	1,205.8	1,527.7
1999.....	77.7	130.6	217.2	344.4	491.4	671.1	891.6	1,078.5	1,239.4	1,510.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)										
All races 3/, both sexes										
2005.....	0.8	1.9	3.5	5.6	7.5	9.2	10.6	11.8	14.2	17.0
2004.....	0.9	2.0	3.8	5.1	7.5	9.4	11.0	12.7	14.9	18.0
2003.....	0.9	2.0	3.6	5.5	7.9	9.7	10.8	12.8	13.6	18.6
2002.....	0.8	2.0	3.6	5.8	8.1	8.8	12.1	13.1	14.1	18.7
2001.....	0.8	2.1	3.5	6.1	7.9	9.4	11.6	13.5	15.2	18.5
2000.....	0.9	2.0	3.8	5.7	8.1	9.9	11.5	12.6	14.0	19.4
1999.....	0.9	2.3	3.6	5.8	8.1	9.3	11.3	13.5	15.4	19.3
All races 3/, male										
2005.....	1.2	2.8	5.7	9.1	11.9	14.8	16.0	17.9	20.4	22.0
2004.....	1.2	3.2	6.2	8.6	11.9	14.9	16.4	19.3	21.7	24.9
2003.....	1.4	3.2	6.2	9.1	12.6	15.0	16.4	19.0	19.9	26.9
2002.....	1.2	3.2	5.8	9.4	12.9	13.9	18.1	19.2	20.3	26.9
2001.....	1.2	3.2	5.8	10.0	13.0	14.3	17.9	19.6	22.1	25.8
2000.....	1.4	3.1	6.0	9.5	12.7	15.2	16.9	18.7	20.2	27.1
1999.....	1.3	3.6	5.7	9.0	13.1	14.3	17.4	20.1	22.3	27.1
All races 3/, female										
2005.....	0.4	1.0	1.4	2.3	3.4	4.3	6.2	7.3	10.4	14.7
2004.....	0.5	0.9	1.5	1.7	3.6	4.7	6.7	8.0	10.8	14.9
2003.....	0.5	0.9	1.1	2.1	3.6	5.2	6.3	8.4	9.7	15.0
2002.....	0.4	0.9	1.4	2.5	3.7	4.4	7.3	8.8	10.3	15.1
2001.....	0.4	1.0	1.4	2.4	3.3	5.1	6.6	9.3	11.1	15.4
2000.....	0.4	1.0	1.6	2.2	3.9	5.4	7.2	8.3	10.4	16.3
1999.....	0.4	1.0	1.6	2.7	3.6	5.1	6.6	9.0	11.3	16.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)										
White, both sexes										
2005 4/.....	0.7	1.7	3.1	5.1	7.1	8.7	10.5	11.7	14.2	17.3
2004 4/.....	0.7	1.7	3.4	4.6	7.0	9.0	11.0	12.7	15.0	18.6
2003 4/.....	0.8	1.8	3.2	5.2	7.3	9.3	10.4	12.6	13.4	19.0
2002.....	0.7	1.7	3.1	5.3	7.5	8.4	12.3	13.3	14.2	18.9
2001.....	0.7	1.7	3.2	5.5	7.3	8.7	11.4	13.6	15.5	19.0
2000.....	0.7	1.7	3.2	5.0	7.3	9.4	11.1	12.5	14.0	19.7
1999.....	0.7	1.8	3.0	5.0	7.5	8.9	11.0	13.2	15.5	19.8
White, male										
2005 4/.....	1.0	2.4	5.1	8.1	11.2	13.8	15.3	17.3	20.0	22.1
2004 4/.....	0.9	2.6	5.4	7.8	10.9	14.0	15.8	18.8	21.5	24.8
2003 4/.....	1.3	2.7	5.5	8.5	11.3	14.1	15.2	18.2	19.2	27.2
2002.....	1.0	2.7	5.0	8.6	11.7	12.9	18.0	19.4	20.2	27.0
2001.....	1.0	2.6	5.2	8.9	11.8	13.2	17.1	19.2	22.3	26.4
2000.....	1.1	2.7	4.9	8.3	11.2	14.0	15.9	18.0	19.9	27.6
1999.....	1.1	2.7	4.7	7.6	12.0	13.5	16.8	19.3	22.0	27.3
White, female										
2005 4/.....	0.5	0.9	1.3	2.2	3.3	4.2	6.5	7.6	10.6	15.1
2004 4/.....	0.4	0.8	1.4	1.5	3.4	4.6	7.1	8.2	10.9	15.7
2003 4/.....	0.3	0.9	0.9	2.1	3.7	5.1	6.4	8.5	9.8	15.4
2002.....	0.4	0.7	1.3	2.2	3.6	4.3	7.6	8.8	10.5	15.3
2001.....	0.3	0.8	1.2	2.2	3.2	4.8	6.7	9.6	11.3	15.9
2000.....	0.4	0.8	1.6	1.9	3.6	5.3	7.2	8.6	10.6	16.5
1999.....	0.4	0.8	1.3	2.5	3.3	4.9	6.4	9.0	11.7	16.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)

Black, both sexes

2005 4/.....	1.1	3.6	6.6	9.2	12.6	14.1	13.7	13.0	14.3	12.6
2004 4/.....	1.9	4.6	7.8	9.5	13.3	13.8	13.7	14.9	15.8	12.4
2003 4/.....	1.7	3.9	6.5	8.8	14.4	15.5	15.3	14.8	16.0	15.5
2002.....	1.5	4.2	7.1	10.1	14.7	13.3	12.5	11.3	12.6	16.0
2001.....	1.7	4.3	6.2	12.2	13.1	15.3	15.7	14.2	13.3	12.9
2000.....	1.6	4.1	8.3	12.2	15.2	14.9	16.1	14.6	15.8	15.3
1999.....	1.5	5.9	8.8	12.6	14.6	14.4	16.2	17.7	15.6	14.3

Black, male

2005 4/.....	2.1	6.1	11.7	17.5	22.4	26.6	25.9	26.8	28.8	21.4
2004 4/.....	2.9	7.6	13.8	17.0	23.7	26.2	27.2	25.9	27.3	26.8
2003 4/.....	1.9	7.0	11.4	16.3	27.8	27.6	30.2	26.5	30.3	25.0
2002.....	2.4	6.7	12.9	18.2	26.9	24.2	21.4	16.5	21.4	26.8
2001.....	2.4	7.0	10.8	22.2	24.8	25.4	29.3	25.1	20.9	*
2000.....	2.7	6.3	15.9	21.9	27.3	27.6	29.1	28.8	28.7	*
1999.....	2.3	10.3	14.5	21.5	25.2	24.1	26.5	33.2	28.8	24.6

Black, female

2005 4/.....	*	1.5	2.2	2.4	5.0	4.8	5.4	*	*	9.0
2004 4/.....	*	1.9	2.7	3.4	5.2	4.7	4.6	8.4	9.8	*
2003 4/.....	1.5	*	2.3	2.7	3.9	6.6	5.3	7.9	8.5	11.7
2002.....	*	1.9	2.2	3.4	5.1	5.2	6.5	8.2	8.0	11.9
2001.....	*	2.0	2.2	4.0	3.8	7.9	6.7	7.8	9.4	10.5
2000.....	*	2.1	*	4.2	5.6	5.6	7.5	6.0	9.3	14.2
1999.....	*	2.1	3.9	5.3	6.3	7.1	9.5	8.3	9.1	10.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasm of esophagus (C15)										
All races 3/, both sexes										
2005.....	1.1	2.3	4.9	8.4	12.6	17.5	22.4	27.6	27.3	24.9
2004.....	1.0	2.3	4.7	8.1	13.2	17.5	21.3	26.0	27.7	25.8
2003.....	0.9	2.4	4.6	8.8	12.8	17.6	22.2	25.4	27.6	25.0
2002.....	1.0	2.4	4.7	8.6	12.7	18.1	22.5	25.7	27.8	24.6
2001.....	1.1	2.4	5.0	7.9	13.0	18.5	22.9	26.1	26.5	25.8
2000.....	1.0	2.4	4.9	8.8	13.2	17.9	22.1	24.5	27.2	25.9
1999.....	1.0	2.4	4.7	8.7	13.0	18.2	22.0	24.3	24.6	26.8
All races 3/, male										
2005.....	1.7	3.9	8.8	14.8	22.2	31.1	39.6	49.6	52.4	46.0
2004.....	1.6	4.0	8.1	14.1	22.9	30.5	37.0	46.8	51.7	48.5
2003.....	1.5	4.1	8.1	15.3	22.7	30.6	38.3	45.7	51.0	45.3
2002.....	1.6	4.1	8.3	15.1	22.1	31.4	40.3	46.8	50.6	43.4
2001.....	1.8	4.1	8.7	13.7	22.6	32.6	39.9	46.5	48.8	48.1
2000.....	1.8	4.2	8.7	15.2	23.0	30.7	38.0	43.9	48.8	47.8
1999.....	1.7	4.2	8.3	15.3	21.9	31.7	39.3	44.1	44.8	48.0
All races 3/, female										
2005.....	0.4	0.7	1.2	2.5	3.9	5.6	8.5	11.6	11.6	15.3
2004.....	0.3	0.7	1.4	2.4	4.3	6.1	8.7	11.1	12.9	15.7
2003.....	0.3	0.7	1.3	2.6	3.8	6.3	9.2	11.0	13.3	16.1
2002.....	0.4	0.7	1.3	2.4	4.2	6.7	8.2	10.9	14.0	16.5
2001.....	0.3	0.7	1.4	2.4	4.2	6.4	9.4	11.9	13.2	16.5
2000.....	0.2	0.6	1.4	2.8	4.4	7.1	9.6	11.0	14.5	17.1
1999.....	0.3	0.6	1.3	2.5	5.0	6.7	8.5	10.5	12.8	18.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasm of esophagus (C15)										
White, both sexes										
2005 4/.....	1.1	2.3	4.9	8.3	12.5	17.6	22.9	28.0	28.0	25.9
2004 4/.....	1.0	2.2	4.6	7.9	12.8	17.4	21.2	26.5	28.0	26.2
2003 4/.....	0.9	2.3	4.3	8.5	12.3	17.0	22.5	25.8	27.8	25.8
2002.....	0.9	2.3	4.6	8.1	12.2	17.5	22.4	25.7	28.2	25.1
2001.....	1.0	2.3	4.6	7.1	12.3	17.7	22.1	26.2	26.9	26.3
2000.....	0.9	2.3	4.5	8.1	12.4	17.4	21.8	24.0	27.2	26.6
1999.....	0.9	2.2	4.3	8.0	12.1	17.5	21.5	23.7	24.9	27.2
White, male										
2005 4/.....	1.8	3.9	9.0	14.7	22.0	31.2	40.4	50.6	53.7	48.5
2004 4/.....	1.7	3.9	8.0	13.9	22.5	30.5	37.2	47.6	52.8	49.2
2003 4/.....	1.6	3.9	7.6	15.0	22.0	29.5	39.0	46.5	52.3	46.7
2002.....	1.6	4.0	8.1	14.5	21.5	30.7	40.2	47.2	51.3	44.6
2001.....	1.8	3.9	8.1	12.5	21.6	31.2	39.0	46.9	49.5	48.8
2000.....	1.8	4.1	8.0	14.1	22.0	30.0	37.5	43.2	49.7	49.3
1999.....	1.6	4.0	7.6	14.1	20.5	30.8	38.4	43.0	45.6	48.0
White, female										
2005 4/.....	0.3	0.6	1.0	2.2	3.6	5.5	8.4	11.3	11.8	15.5
2004 4/.....	0.3	0.5	1.1	2.1	3.8	5.9	8.1	11.1	12.6	15.9
2003 4/.....	0.3	0.6	1.0	2.3	3.4	6.0	9.0	10.9	12.7	16.5
2002.....	0.3	0.6	1.2	2.1	3.7	5.9	7.8	10.3	14.1	16.7
2001.....	0.3	0.6	1.2	1.9	3.7	5.8	8.5	11.6	13.2	16.8
2000.....	*	0.4	1.1	2.4	3.6	6.4	9.1	10.6	13.8	17.3
1999.....	*	0.5	1.0	2.1	4.4	5.8	8.0	10.1	12.6	18.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasm of esophagus (C15)										
Black, both sexes										
2005 4/.....	1.3	3.2	6.8	11.8	17.4	21.4	23.9	30.2	22.6	17.4
2004 4/.....	1.2	4.0	6.6	12.0	20.4	23.0	26.8	27.4	27.0	21.4
2003 4/.....	1.3	4.1	8.4	14.1	20.7	26.1	24.9	26.9	29.4	21.3
2002.....	1.6	3.6	6.9	14.9	20.4	27.5	29.8	31.8	26.2	20.8
2001.....	1.6	3.7	8.6	15.9	21.9	29.6	35.8	30.7	27.6	23.6
2000.....	1.5	4.1	9.8	16.9	23.3	27.0	29.6	36.1	29.9	22.2
1999.....	1.7	4.3	9.4	17.1	23.5	28.7	31.3	35.7	23.7	23.9
Black, male										
2005 4/.....	2.1	5.1	11.4	19.8	31.3	39.7	41.4	52.7	48.5	27.0
2004 4/.....	1.7	6.3	10.8	19.9	33.4	40.0	43.1	49.6	45.2	38.7
2003 4/.....	2.1	7.1	14.0	23.3	36.2	46.5	41.4	47.5	43.6	38.6
2002.....	2.5	5.7	11.5	26.8	34.4	45.4	54.4	54.2	48.1	37.5
2001.....	2.6	6.0	15.3	27.5	38.8	51.8	59.9	55.3	53.8	46.5
2000.....	2.4	6.3	17.4	30.5	38.5	44.8	49.8	65.7	45.5	38.1
1999.....	3.2	7.4	16.4	30.3	40.3	46.5	56.0	67.9	36.6	51.6
Black, female										
2005 4/.....	*	1.6	2.9	5.2	6.5	7.8	11.9	16.8	8.9	13.5
2004 4/.....	*	1.9	3.1	5.5	10.2	10.4	15.8	14.3	17.4	14.6
2003 4/.....	*	1.6	3.6	6.5	8.6	11.0	13.7	14.7	22.0	14.6
2002.....	*	1.8	3.0	5.1	9.4	14.2	13.2	18.4	14.7	14.4
2001.....	*	1.7	2.9	6.3	8.6	13.1	19.7	16.1	14.3	15.2
2000.....	*	2.1	3.4	5.8	11.3	13.8	16.2	18.2	22.1	16.3
1999.....	*	1.6	3.4	6.4	10.3	15.6	14.9	16.3	17.3	13.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasm of stomach (C16)										
All races 3/, both sexes										
2005.....	1.3	2.1	3.1	4.8	7.3	11.8	16.7	23.5	30.6	38.5
2004.....	1.4	2.1	3.5	5.3	8.0	12.1	18.2	23.4	30.3	42.2
2003.....	1.3	2.3	3.8	5.3	8.6	12.4	18.4	24.2	32.2	43.6
2002.....	1.2	2.1	3.5	5.6	8.5	12.3	18.8	25.6	33.0	47.6
2001.....	1.3	2.2	3.7	5.3	9.2	13.7	19.5	27.0	34.7	45.4
2000.....	1.4	2.3	3.6	5.6	9.7	13.8	19.6	27.1	36.9	52.7
1999.....	1.4	2.3	3.8	6.2	9.8	14.2	20.2	28.1	36.8	50.1
All races 3/, male										
2005.....	1.6	2.5	4.3	6.6	10.4	16.4	23.8	32.5	44.3	54.2
2004.....	1.7	2.6	4.6	7.5	11.2	17.0	24.9	33.6	42.5	62.9
2003.....	1.5	2.8	5.0	7.5	12.3	17.9	26.6	34.2	44.7	60.0
2002.....	1.4	2.7	4.8	7.7	11.9	18.1	27.5	36.4	47.1	67.2
2001.....	1.4	2.8	4.9	7.5	13.7	19.5	27.5	39.1	49.4	62.6
2000.....	1.7	2.9	4.8	7.8	14.1	19.5	28.3	39.4	51.4	75.6
1999.....	1.7	3.1	5.4	8.8	13.7	20.3	28.6	41.5	54.9	71.3
All races 3/, female										
2005.....	1.1	1.8	2.0	3.0	4.5	7.7	11.0	17.0	22.1	31.2
2004.....	1.2	1.6	2.4	3.3	5.1	7.8	12.8	16.0	22.7	32.9
2003.....	1.1	1.7	2.6	3.1	5.3	7.7	11.9	17.1	24.5	36.4
2002.....	1.0	1.5	2.3	3.7	5.4	7.3	11.9	18.1	24.4	39.1
2001.....	1.1	1.6	2.5	3.2	5.1	8.8	13.1	18.5	25.9	38.3
2000.....	1.2	1.7	2.4	3.5	5.8	8.9	12.7	18.5	28.3	43.4
1999.....	1.2	1.5	2.4	3.7	6.4	9.1	13.7	18.8	26.2	41.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasm of stomach (C16)										
White, both sexes										
2005 4/.....	1.1	1.7	2.7	4.2	6.2	10.0	14.7	20.5	27.0	34.8
2004 4/.....	1.2	1.7	2.9	4.5	6.9	10.2	15.9	20.9	27.3	38.8
2003 4/.....	1.1	1.8	3.1	4.6	7.6	11.0	16.3	21.4	28.8	38.7
2002.....	1.0	1.8	2.9	4.9	7.3	10.5	16.5	22.6	29.3	42.2
2001.....	1.1	1.9	3.1	4.7	7.9	11.7	17.4	24.0	31.0	41.1
2000.....	1.1	1.9	3.1	4.9	8.5	11.9	17.2	23.6	32.7	47.3
1999.....	1.1	2.0	3.1	5.2	8.3	12.8	17.7	24.8	33.0	44.8
White, male										
2005 4/.....	1.3	2.0	3.7	5.8	8.6	13.7	20.7	28.7	39.1	49.0
2004 4/.....	1.4	2.1	3.9	6.5	9.5	14.4	21.3	29.9	38.0	58.4
2003 4/.....	1.2	2.3	4.1	6.6	10.6	15.5	23.5	30.3	40.3	53.5
2002.....	1.1	2.4	3.9	6.8	10.3	15.7	24.3	32.8	42.4	60.5
2001.....	1.2	2.4	4.1	6.5	11.7	16.3	24.5	35.4	44.0	57.7
2000.....	1.2	2.3	4.2	6.7	12.5	17.1	25.1	34.6	46.1	68.9
1999.....	1.2	2.7	4.4	7.6	11.5	18.4	25.2	36.9	49.0	63.4
White, female										
2005 4/.....	0.9	1.5	1.8	2.6	3.9	6.7	9.7	14.5	19.5	28.2
2004 4/.....	1.0	1.3	2.0	2.7	4.4	6.5	11.5	14.2	20.7	29.9
2003 4/.....	0.9	1.4	2.2	2.7	4.8	7.0	10.5	15.0	21.7	32.1
2002.....	0.8	1.2	1.8	3.0	4.6	5.9	10.1	15.4	21.3	34.2
2001.....	1.0	1.3	2.1	2.9	4.3	7.7	11.7	16.0	23.1	34.1
2000.....	1.0	1.5	2.0	3.1	4.9	7.4	10.9	15.8	24.7	38.6
1999.....	0.9	1.2	1.9	2.9	5.4	8.0	11.7	16.3	23.5	37.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasm of stomach (C16)										
Black, both sexes										
2005 4/.....	2.3	4.0	5.2	9.0	14.4	23.5	30.1	48.6	57.3	74.3
2004 4/.....	2.7	3.9	6.9	10.6	15.8	27.3	34.0	43.8	59.9	71.3
2003 4/.....	2.4	4.3	7.4	9.6	16.7	22.3	33.1	48.8	62.4	85.8
2002.....	2.2	3.6	7.5	10.4	16.1	24.6	38.9	52.7	63.1	100.4
2001.....	2.5	3.7	7.5	9.2	17.7	27.1	34.7	51.1	69.9	88.1
2000.....	2.9	4.7	7.0	11.1	17.3	26.8	38.2	56.9	75.1	101.7
1999.....	3.2	3.9	8.5	13.2	19.1	23.6	40.3	58.6	74.3	102.4
Black, male										
2005 4/.....	3.2	5.4	7.7	13.5	24.2	34.9	48.4	67.5	79.4	114.3
2004 4/.....	3.1	5.3	9.5	15.3	25.2	40.7	56.0	63.7	93.3	100.3
2003 4/.....	3.3	6.2	10.9	14.9	26.0	36.4	54.2	74.9	87.1	106.3
2002.....	2.4	5.1	11.0	15.2	24.0	36.9	59.8	73.5	90.2	137.2
2001.....	2.8	5.0	11.5	13.8	27.5	43.7	54.2	75.7	106.7	104.4
2000.....	3.9	6.6	9.6	16.5	25.9	38.5	57.2	83.2	105.4	136.1
1999.....	4.0	5.7	12.2	19.0	28.2	35.1	60.4	89.3	121.2	147.6
Black, female										
2005 4/.....	1.5	2.8	3.1	5.3	6.8	15.0	17.5	37.3	45.6	58.1
2004 4/.....	2.3	2.7	4.7	6.7	8.4	17.3	19.1	32.1	42.3	59.9
2003 4/.....	1.5	2.7	4.5	5.3	9.5	11.9	18.8	33.4	49.5	77.8
2002.....	2.1	2.3	4.6	6.5	9.9	15.5	24.9	40.3	48.9	86.3
2001.....	2.2	2.5	4.1	5.4	10.1	14.8	21.7	36.3	51.1	82.1
2000.....	1.9	3.0	4.7	6.6	10.5	18.2	25.5	41.0	59.9	88.9
1999.....	2.4	2.4	5.4	8.4	12.0	15.2	27.0	40.0	51.0	85.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of colon, rectum and anus (C18-C21)										
All races 3/, both sexes										
2005.....	4.0	7.4	13.4	21.4	34.2	52.7	75.8	105.7	149.2	216.9
2004.....	3.9	7.4	13.5	22.4	35.9	53.2	78.3	111.3	152.3	225.6
2003.....	3.8	7.9	14.3	23.0	38.6	59.3	83.3	117.0	159.0	239.7
2002.....	4.3	8.2	15.0	24.6	40.7	61.6	86.2	118.9	163.2	241.4
2001.....	4.1	8.4	15.2	25.4	39.3	62.9	88.8	121.1	165.8	255.7
2000.....	4.0	8.2	14.7	25.2	42.0	62.0	92.4	126.9	174.8	270.4
1999.....	4.0	8.3	14.4	25.3	43.6	63.9	91.9	129.0	173.2	271.7
All races 3/, male										
2005.....	4.2	8.2	15.2	25.9	42.4	67.7	94.4	131.8	180.2	245.9
2004.....	4.2	8.3	16.1	27.0	43.7	66.4	98.0	139.0	183.1	254.7
2003.....	4.2	9.0	16.6	27.6	47.1	75.0	103.4	143.0	196.5	273.3
2002.....	4.9	9.1	17.5	30.1	51.6	77.8	106.8	147.7	197.4	275.1
2001.....	4.3	9.5	17.1	30.8	48.0	78.9	110.4	149.4	201.3	298.3
2000.....	4.4	9.0	17.0	30.6	51.0	77.4	115.7	159.4	213.0	315.6
1999.....	4.1	9.1	16.6	30.6	52.9	78.0	114.3	161.3	216.2	324.0
All races 3/, female										
2005.....	3.7	6.6	11.6	17.1	26.7	39.7	60.8	86.8	130.0	203.6
2004.....	3.6	6.5	11.1	18.0	28.7	41.8	62.4	91.3	133.3	212.5
2003.....	3.4	6.9	12.1	18.6	30.9	45.6	67.3	98.5	136.1	224.8
2002.....	3.6	7.4	12.6	19.5	30.8	47.6	69.7	98.6	142.3	226.8
2001.....	3.8	7.4	13.4	20.4	31.5	49.2	71.5	101.2	144.6	237.8
2000.....	3.5	7.5	12.4	20.2	33.9	48.8	74.1	104.3	152.2	251.9
1999.....	3.9	7.5	12.3	20.4	35.1	51.8	74.3	106.6	148.0	250.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of colon, rectum and anus (C18-C21)										
White, both sexes										
2005 4/.....	3.7	6.8	12.2	19.9	32.1	50.3	74.4	103.1	147.3	215.7
2004 4/.....	3.6	6.8	12.6	21.3	34.2	51.2	76.2	109.5	150.9	225.1
2003 4/.....	3.4	7.3	13.0	21.5	36.9	58.0	80.7	115.1	158.2	238.5
2002.....	3.9	7.5	14.2	23.3	38.9	59.9	84.6	116.9	160.3	240.7
2001.....	3.8	7.7	14.3	23.9	37.9	60.8	86.8	118.0	163.7	255.3
2000.....	3.6	7.5	13.8	23.7	40.5	60.3	90.6	125.0	172.1	269.2
1999.....	3.7	7.7	13.5	23.8	42.0	62.0	90.0	126.9	170.8	270.9
White, male										
2005 4/.....	4.1	7.6	13.9	24.1	40.0	64.6	92.9	128.0	179.5	245.3
2004 4/.....	3.9	7.8	15.1	26.0	41.9	64.0	95.3	135.9	181.7	255.4
2003 4/.....	3.8	8.3	15.2	26.2	45.0	73.6	100.8	140.7	196.0	272.9
2002.....	4.5	8.4	16.5	28.7	49.4	76.0	105.4	144.9	193.3	273.6
2001.....	4.0	8.7	16.4	29.3	46.7	76.7	108.7	146.7	198.9	297.3
2000.....	3.9	8.3	16.1	28.6	49.1	75.3	113.5	157.6	208.6	313.3
1999.....	3.9	8.5	15.8	28.8	51.1	76.1	113.3	158.1	214.4	324.5
White, female										
2005 4/.....	3.4	6.0	10.5	15.9	24.8	37.6	59.2	84.7	127.1	202.0
2004 4/.....	3.4	5.9	10.3	16.8	27.1	39.8	60.5	90.3	131.8	211.4
2003 4/.....	2.9	6.3	10.8	17.0	29.5	44.1	64.2	96.6	134.8	223.2
2002.....	3.3	6.5	12.0	18.2	29.3	45.7	67.6	96.9	140.1	226.4
2001.....	3.6	6.7	12.2	18.8	29.8	46.9	69.1	97.7	142.5	237.7
2000.....	3.3	6.6	11.5	19.1	32.6	47.2	72.3	102.1	150.4	251.3
1999.....	3.5	6.9	11.2	19.2	33.6	49.7	71.5	104.9	145.1	249.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of colon, rectum and anus (C18-C21)										
Black, both sexes										
2005 4/.....	5.7	12.6	23.8	36.5	57.5	82.9	101.3	148.3	197.6	261.7
2004 4/.....	5.9	12.2	21.9	36.1	56.7	79.9	113.2	148.6	189.1	260.8
2003 4/.....	6.6	13.2	25.5	39.4	59.7	81.2	124.6	154.7	197.7	281.5
2002.....	7.2	14.5	23.5	40.0	61.9	86.8	114.9	160.5	218.7	274.7
2001.....	6.5	14.5	24.6	43.1	58.3	90.9	120.3	172.6	211.0	286.5
2000.....	6.6	15.2	24.8	39.9	62.7	86.9	126.1	167.4	226.2	310.0
1999.....	6.6	13.3	23.4	42.0	65.7	90.5	125.4	172.3	226.8	306.1
Black, male										
2005 4/.....	5.7	13.7	27.3	46.0	74.2	106.7	124.9	197.1	231.8	295.6
2004 4/.....	6.6	13.2	26.3	42.0	69.7	100.5	147.0	198.8	239.6	278.2
2003 4/.....	7.1	15.2	30.3	46.1	76.4	102.7	156.7	194.9	242.2	317.0
2002.....	8.4	15.7	27.9	47.5	79.4	109.2	138.2	209.0	285.1	328.1
2001.....	7.6	16.4	26.4	51.3	67.7	114.1	146.8	208.2	263.2	347.2
2000.....	8.0	16.4	29.2	51.4	76.4	109.3	160.1	208.9	298.4	377.1
1999.....	6.3	15.0	26.9	54.1	80.4	110.6	147.6	228.0	274.7	350.4
Black, female										
2005 4/.....	5.6	11.6	20.9	28.7	44.5	65.2	85.1	119.2	179.5	248.0
2004 4/.....	5.3	11.3	18.2	31.3	46.6	64.6	90.3	118.9	162.5	254.0
2003 4/.....	6.1	11.4	21.4	33.8	46.8	65.2	103.0	130.8	174.5	267.8
2002.....	6.1	13.5	19.7	33.9	48.2	70.1	99.2	131.5	184.2	254.2
2001.....	5.6	12.8	23.1	36.3	50.9	73.8	102.7	151.4	184.4	264.0
2000.....	5.3	14.2	21.1	30.5	52.0	70.5	103.4	142.4	189.8	285.0
1999.....	6.7	11.9	20.5	32.1	54.2	75.6	110.7	138.7	203.1	289.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of liver and intrahepatic bile ducts (C22)										
All races 3/, both sexes										
2005.....	1.2	3.7	8.5	10.2	12.9	16.6	24.3	30.2	34.4	32.5
2004.....	1.2	3.9	7.6	9.4	12.5	16.6	23.7	28.8	34.6	32.4
2003.....	1.2	3.7	7.4	8.5	11.7	16.5	23.5	29.6	32.1	33.4
2002.....	1.3	3.4	6.6	8.1	11.6	16.9	22.6	27.6	31.6	35.0
2001.....	1.2	3.8	5.8	7.5	11.6	16.3	21.5	27.0	32.0	34.1
2000.....	1.4	3.7	5.4	7.7	11.3	16.0	21.6	27.1	31.3	32.7
1999.....	1.4	3.4	4.9	7.1	10.9	16.3	21.8	26.1	30.9	30.6
All races 3/, male										
2005.....	1.6	5.9	13.8	16.7	19.5	24.2	34.8	42.8	48.2	44.8
2004.....	1.8	6.2	12.8	15.1	18.6	23.6	34.0	38.8	48.4	41.5
2003.....	1.8	6.0	12.2	13.0	17.3	23.3	33.4	43.7	46.1	47.5
2002.....	2.0	5.7	11.1	12.4	17.4	24.8	32.8	39.5	44.8	46.2
2001.....	1.8	6.0	9.7	11.4	17.2	23.5	30.6	39.2	43.9	50.3
2000.....	2.0	6.0	8.5	11.7	16.5	23.6	30.9	39.3	46.7	44.7
1999.....	2.1	5.5	7.6	10.6	15.6	23.7	31.9	36.6	42.9	40.1
All races 3/, female										
2005.....	0.8	1.5	3.4	4.1	6.8	9.9	15.8	21.0	25.9	26.8
2004.....	0.6	1.8	2.6	4.1	6.9	10.6	15.4	21.7	26.1	28.3
2003.....	0.7	1.6	2.8	4.3	6.6	10.7	15.6	19.6	23.6	27.2
2002.....	0.6	1.2	2.4	4.0	6.4	10.1	14.4	19.2	23.6	30.2
2001.....	0.5	1.7	2.1	3.9	6.5	10.2	14.2	18.4	24.8	27.2
2000.....	0.8	1.4	2.4	4.0	6.5	9.5	14.2	18.6	22.2	27.8
1999.....	0.8	1.3	2.3	3.8	6.6	10.1	13.8	18.7	23.9	26.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of liver and intrahepatic bile ducts (C22)										
White, both sexes										
2005 4/.....	1.0	3.3	7.4	8.9	11.3	15.3	22.5	29.1	32.8	31.7
2004 4/.....	1.0	3.3	6.6	8.0	11.2	15.1	21.9	27.3	33.7	31.2
2003 4/.....	1.0	3.1	6.3	7.5	10.3	15.2	21.7	28.3	31.0	32.1
2002.....	1.1	3.0	5.5	6.8	10.6	15.8	21.1	26.7	30.7	33.8
2001.....	0.8	3.2	4.9	6.4	10.5	15.1	20.2	26.1	30.4	33.3
2000.....	1.1	3.1	4.4	6.6	10.0	14.6	20.2	25.8	30.7	32.0
1999.....	1.1	2.7	4.2	5.9	9.7	15.2	20.5	25.2	30.2	29.9
White, male										
2005 4/.....	1.3	5.2	11.6	14.2	17.1	22.0	32.2	41.5	45.8	45.0
2004 4/.....	1.5	5.0	11.1	12.4	16.5	21.5	31.1	37.4	47.2	39.9
2003 4/.....	1.4	4.9	10.2	11.4	15.2	21.4	31.4	41.8	44.8	45.6
2002.....	1.6	4.9	9.0	10.4	15.6	23.0	30.4	38.0	42.9	45.1
2001.....	1.3	5.0	8.1	9.5	15.7	22.1	29.1	37.7	42.6	49.7
2000.....	1.5	4.8	6.8	9.6	14.4	21.4	28.9	38.1	45.8	43.1
1999.....	1.7	4.3	6.3	8.8	13.9	21.9	30.2	35.3	42.4	40.6
White, female										
2005 4/.....	0.7	1.3	3.2	3.7	6.0	9.3	14.5	19.9	24.7	25.6
2004 4/.....	0.5	1.5	2.2	3.7	6.3	9.5	14.3	19.8	25.3	27.3
2003 4/.....	0.6	1.3	2.5	3.8	5.7	9.7	13.9	18.6	22.5	26.2
2002.....	0.5	1.1	2.0	3.4	5.9	9.3	13.5	18.6	23.2	29.0
2001.....	0.4	1.4	1.9	3.4	5.7	8.9	13.0	17.8	23.1	26.5
2000.....	0.6	1.3	2.0	3.6	5.9	8.5	13.2	17.1	21.7	27.4
1999.....	0.6	1.2	2.1	3.1	5.8	9.3	12.8	18.1	23.0	25.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of liver and intrahepatic bile ducts (C22)										
Black, both sexes										
2005 4/.....	1.8	5.6	14.5	18.9	22.1	21.4	29.4	29.1	39.3	29.1
2004 4/.....	1.8	7.1	12.7	17.9	16.6	23.2	28.5	33.7	35.4	36.1
2003 4/.....	1.9	6.2	14.2	14.7	18.4	18.9	27.9	30.3	32.7	41.7
2002.....	2.1	5.8	12.3	15.5	14.0	19.5	27.3	30.2	31.9	38.0
2001.....	2.3	7.1	10.7	15.2	14.3	20.0	24.0	26.9	40.9	35.0
2000.....	2.7	7.0	10.6	12.7	15.8	20.9	26.9	28.8	28.5	31.6
1999.....	2.2	6.6	8.1	12.0	14.7	21.1	28.0	26.6	33.8	34.4
Black, male										
2005 4/.....	2.5	8.9	26.8	35.0	36.7	35.9	44.6	45.1	59.7	31.6
2004 4/.....	2.9	11.7	22.9	31.9	27.1	36.8	45.0	39.6	50.9	46.7
2003 4/.....	3.0	10.3	25.5	24.0	28.5	26.8	42.4	43.8	45.0	63.6
2002.....	3.3	9.8	21.7	25.3	22.8	32.2	45.1	44.6	51.2	41.8
2001.....	3.6	11.6	18.6	25.3	21.4	28.0	32.7	39.5	49.0	42.0
2000.....	3.8	12.6	17.7	22.1	25.3	34.1	41.5	36.4	45.5	48.4
1999.....	3.4	11.8	14.2	18.6	21.6	32.2	42.1	40.4	42.7	30.5
Black, female										
2005 4/.....	*	2.6	4.2	5.6	10.8	10.6	19.0	19.5	28.6	28.1
2004 4/.....	*	3.0	4.1	6.4	8.4	13.1	17.4	30.2	27.2	31.9
2003 4/.....	*	2.7	4.5	7.0	10.5	13.0	18.2	22.2	26.3	33.2
2002.....	*	2.3	4.3	7.5	7.1	10.2	15.5	21.7	21.9	36.6
2001.....	*	3.2	3.9	6.8	8.8	14.1	18.2	19.4	36.8	32.4
2000.....	1.8	2.2	4.7	5.0	8.3	11.1	17.3	24.2	20.0	25.3
1999.....	*	2.0	2.9	6.5	9.3	12.9	18.7	18.3	29.4	35.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasm of pancreas (C25)										
All races 3/, both sexes										
2005.....	1.9	4.5	8.7	15.7	25.1	37.6	52.6	69.6	88.4	97.4
2004.....	1.7	4.2	8.8	15.2	25.8	37.3	52.4	68.9	85.4	97.7
2003.....	1.7	4.1	8.7	14.7	24.7	36.4	53.0	67.6	86.2	95.2
2002.....	1.9	4.2	8.2	15.5	24.0	37.2	53.2	70.2	81.8	94.5
2001.....	1.8	4.0	8.6	14.9	24.8	37.0	52.8	67.2	84.2	100.1
2000.....	1.8	4.2	8.7	15.1	24.8	37.2	51.4	70.4	85.7	96.4
1999.....	1.9	4.5	8.7	15.6	25.0	38.1	52.5	67.2	82.0	99.7
All races 3/, male										
2005.....	2.2	5.2	10.6	19.6	30.6	43.9	61.6	78.7	96.5	102.8
2004.....	2.2	5.5	11.0	18.6	31.5	45.2	61.6	77.7	94.8	102.0
2003.....	2.0	5.2	10.7	18.2	29.8	42.5	60.9	76.1	99.5	95.6
2002.....	2.4	5.3	10.6	19.2	29.3	43.3	60.8	78.9	91.9	100.9
2001.....	2.0	4.9	10.7	18.5	30.3	43.6	59.0	77.2	94.9	105.1
2000.....	2.1	5.4	11.1	18.5	30.0	44.7	57.0	80.4	94.9	102.9
1999.....	2.4	5.9	11.0	19.1	29.9	44.9	60.9	75.7	92.0	107.5
All races 3/, female										
2005.....	1.6	3.8	6.9	12.0	20.0	32.1	45.3	63.0	83.4	94.9
2004.....	1.3	2.9	6.7	12.0	20.6	30.5	45.1	62.6	79.6	95.8
2003.....	1.4	3.0	6.8	11.5	20.2	31.1	46.7	61.6	78.1	95.0
2002.....	1.4	3.1	6.0	12.0	19.2	31.9	47.1	64.1	75.6	91.7
2001.....	1.6	3.1	6.5	11.5	19.8	31.3	47.8	60.1	77.8	98.1
2000.....	1.5	3.0	6.4	11.8	20.0	30.8	47.1	63.5	80.2	93.8
1999.....	1.4	3.1	6.5	12.3	20.5	32.2	46.0	61.2	76.1	96.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasm of pancreas (C25)										
White, both sexes										
2005 4/.....	1.9	4.4	8.2	15.2	24.6	37.1	52.2	68.8	87.7	96.8
2004 4/.....	1.6	4.0	8.4	14.6	25.1	36.8	51.3	68.3	85.1	97.1
2003 4/.....	1.6	3.9	8.1	14.3	23.8	36.0	52.2	67.2	85.2	94.4
2002.....	1.9	4.1	7.8	15.0	23.3	36.1	52.0	69.2	81.4	94.3
2001.....	1.7	3.8	8.3	14.4	24.7	36.3	51.7	66.3	83.3	99.0
2000.....	1.7	4.1	8.4	14.5	24.3	36.2	50.4	69.2	84.5	94.7
1999.....	1.8	4.4	8.2	14.8	24.3	37.2	51.5	65.8	80.6	98.6
White, male										
2005 4/.....	2.2	5.1	10.0	18.9	30.4	43.4	62.0	78.1	96.2	102.9
2004 4/.....	2.0	5.3	10.5	17.7	30.8	44.5	60.0	77.7	95.8	102.9
2003 4/.....	1.8	4.8	10.0	17.7	28.9	42.7	60.2	76.1	99.3	96.0
2002.....	2.4	5.1	10.1	18.6	28.8	42.6	59.7	78.9	92.2	101.6
2001.....	1.9	4.7	10.5	18.2	30.3	42.8	58.0	77.0	94.5	106.5
2000.....	2.0	5.3	10.7	17.8	29.6	44.3	55.9	79.9	94.0	102.4
1999.....	2.3	5.8	10.3	17.9	29.2	44.0	60.1	74.6	90.9	107.4
White, female										
2005 4/.....	1.5	3.6	6.4	11.7	19.3	31.5	44.0	61.9	82.4	94.0
2004 4/.....	1.3	2.8	6.2	11.5	19.7	30.0	44.1	61.5	78.5	94.5
2003 4/.....	1.4	2.9	6.3	11.0	19.1	30.0	45.8	60.7	76.6	93.8
2002.....	1.4	3.0	5.5	11.6	18.2	30.3	45.7	62.2	74.9	91.1
2001.....	1.6	2.9	6.3	10.8	19.6	30.6	46.5	58.7	76.5	95.9
2000.....	1.4	2.9	6.1	11.4	19.4	29.2	46.0	61.6	78.9	91.5
1999.....	1.3	3.1	6.3	11.9	19.9	31.3	44.6	59.5	74.5	95.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasm of pancreas (C25)										
Black, both sexes										
2005 4/.....	2.2	6.2	13.7	22.3	33.6	49.1	66.4	86.9	105.1	111.4
2004 4/.....	2.6	6.0	13.3	22.6	36.3	48.0	70.5	84.3	93.3	113.6
2003 4/.....	2.7	6.4	14.5	20.7	36.1	47.1	68.0	83.6	107.4	112.9
2002.....	2.5	5.6	12.3	22.4	34.4	50.4	73.7	88.6	91.8	106.6
2001.....	2.2	6.4	12.1	20.7	31.6	48.5	70.8	84.9	104.8	121.3
2000.....	2.6	5.8	13.2	22.8	33.3	50.3	69.8	93.8	104.2	119.8
1999.....	2.8	6.2	14.1	25.0	32.2	50.9	70.1	87.5	104.6	118.6
Black, male										
2005 4/.....	2.3	7.2	17.3	30.1	39.6	59.2	71.5	96.1	116.6	100.4
2004 4/.....	3.4	7.6	16.1	30.0	44.8	60.6	88.1	88.7	91.1	100.3
2003 4/.....	3.5	9.0	18.6	26.2	42.8	50.0	80.8	89.0	112.2	104.3
2002.....	2.9	7.2	16.5	29.0	40.8	56.6	84.1	90.5	105.5	100.8
2001.....	2.3	7.9	15.2	24.2	38.2	60.0	78.8	95.2	109.2	99.8
2000.....	3.0	7.3	17.5	29.3	40.4	55.7	78.9	101.6	116.3	109.6
1999.....	3.4	8.3	19.5	34.1	39.5	60.0	78.4	100.7	117.7	114.8
Black, female										
2005 4/.....	2.2	5.3	10.7	15.8	28.9	41.6	63.0	81.3	99.0	115.9
2004 4/.....	1.9	4.6	11.0	16.6	29.7	38.7	58.6	81.7	94.5	118.9
2003 4/.....	2.1	4.1	11.0	16.2	30.8	45.0	59.4	80.3	104.8	116.3
2002.....	2.1	4.3	8.8	17.0	29.4	45.9	66.8	87.5	84.7	108.8
2001.....	2.2	5.1	9.6	17.8	26.4	40.0	65.5	78.8	102.6	129.2
2000.....	2.3	4.4	9.4	17.4	27.8	46.3	63.7	89.1	98.1	123.6
1999.....	2.2	4.3	9.5	17.6	26.5	44.3	64.6	79.5	98.1	120.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasm of larynx (C32)										
All races 3/, both sexes										
2005.....	0.2	0.7	1.3	2.4	3.9	5.8	6.7	6.8	7.3	6.3
2004.....	0.2	0.7	1.4	2.6	4.1	5.3	6.2	7.1	6.0	6.5
2003.....	0.3	0.7	1.4	2.6	4.6	5.8	6.6	6.6	7.4	6.5
2002.....	0.2	0.7	1.4	2.7	4.2	5.3	6.6	7.9	7.2	5.5
2001.....	0.3	0.7	1.5	2.9	4.5	5.6	7.3	7.4	6.9	6.7
2000.....	0.2	0.8	1.5	2.9	4.5	6.0	7.3	8.0	8.1	6.5
1999.....	0.3	0.8	1.5	3.1	4.6	6.1	7.4	7.2	7.4	7.0
All races 3/, male										
2005.....	0.4	1.0	2.1	4.0	6.7	10.2	11.4	12.2	14.5	15.7
2004.....	0.4	1.2	2.4	4.4	6.8	9.4	11.0	13.3	12.4	15.6
2003.....	0.5	1.2	2.4	4.2	7.9	10.3	11.4	12.6	15.4	15.3
2002.....	0.4	1.2	2.4	4.7	7.2	8.9	11.3	14.8	14.9	13.2
2001.....	0.5	1.2	2.5	4.9	7.5	9.5	12.9	14.5	13.9	18.3
2000.....	0.4	1.2	2.5	4.9	7.9	10.0	12.9	14.8	17.1	17.4
1999.....	0.4	1.3	2.6	5.4	7.8	10.4	12.6	13.2	15.8	17.8
All races 3/, female										
2005.....	*	0.4	0.6	0.8	1.3	1.9	3.0	2.8	2.8	1.9
2004.....	*	0.2	0.5	0.9	1.6	1.7	2.2	2.6	2.1	2.4
2003.....	*	0.3	0.4	1.0	1.6	2.0	2.8	2.3	2.5	2.7
2002.....	*	0.2	0.5	0.9	1.5	2.2	2.9	3.1	2.5	2.2
2001.....	*	0.2	0.5	1.0	1.7	2.2	2.8	2.5	2.8	1.9
2000.....	*	0.3	0.6	0.9	1.4	2.6	2.8	3.3	2.8	2.1
1999.....	*	0.2	0.4	1.0	1.7	2.5	3.3	3.0	2.6	2.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasm of larynx (C32)										
White, both sexes										
2005 4/.....	0.2	0.6	1.1	2.1	3.4	5.4	6.3	6.8	7.2	6.0
2004 4/.....	0.2	0.7	1.2	2.3	3.8	5.0	5.9	6.9	5.7	6.3
2003 4/.....	0.3	0.6	1.1	2.3	4.2	5.5	6.2	6.3	7.1	6.2
2002.....	0.2	0.6	1.2	2.5	3.9	5.0	6.4	7.5	6.9	5.6
2001.....	0.3	0.6	1.2	2.5	4.1	5.2	6.7	7.0	6.8	6.6
2000.....	0.2	0.6	1.3	2.4	3.9	5.6	6.7	7.5	8.0	6.6
1999.....	0.2	0.6	1.3	2.7	4.2	5.7	7.1	7.1	7.3	6.7
White, male										
2005 4/.....	0.4	0.9	1.8	3.5	5.7	9.5	10.5	12.1	14.0	14.9
2004 4/.....	0.3	1.2	1.9	3.9	6.2	8.8	10.4	12.8	11.7	15.0
2003 4/.....	0.5	1.1	2.0	3.7	7.2	9.3	10.4	11.9	14.4	14.5
2002.....	0.3	1.0	2.0	4.1	6.3	8.3	10.7	13.8	14.6	13.3
2001.....	0.4	0.9	2.0	4.2	6.9	8.8	11.9	13.3	13.2	18.3
2000.....	0.3	1.0	2.1	4.1	6.8	9.2	11.7	13.9	16.8	17.6
1999.....	0.3	1.1	2.2	4.6	7.0	9.6	12.0	12.9	15.1	16.8
White, female										
2005 4/.....	*	0.3	0.5	0.8	1.3	1.7	2.9	2.9	2.9	1.9
2004 4/.....	*	0.2	0.5	0.9	1.6	1.7	2.1	2.6	2.0	2.5
2003 4/.....	*	*	0.3	0.9	1.4	2.1	2.7	2.3	2.5	2.5
2002.....	*	*	0.4	0.9	1.5	2.1	2.8	2.9	2.3	2.3
2001.....	*	*	0.4	0.9	1.5	2.1	2.6	2.5	2.9	1.7
2000.....	*	0.3	0.5	0.9	1.3	2.5	2.6	3.0	2.7	2.1
1999.....	*	*	0.3	1.0	1.5	2.3	3.3	3.0	2.7	2.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasm of larynx (C32)										
Black, both sexes										
2005 4/.....	*	1.3	3.3	5.5	9.1	11.5	11.4	9.2	10.9	10.2
2004 4/.....	*	1.2	3.5	5.5	8.0	9.3	11.6	11.2	11.1	8.2
2003 4/.....	*	1.7	3.6	5.8	9.7	10.6	12.7	11.1	12.7	11.1
2002.....	*	1.9	3.6	6.1	9.3	9.1	11.8	15.3	11.3	*
2001.....	*	1.9	3.8	7.1	8.9	9.5	13.9	13.4	9.2	8.3
2000.....	*	2.0	4.1	7.5	10.8	10.6	15.3	15.6	11.6	6.9
1999.....	0.9	2.1	3.9	7.5	10.0	11.4	11.8	10.2	9.8	8.3
Black, male										
2005 4/.....	*	2.1	5.0	10.3	18.5	22.1	22.1	18.3	26.0	28.8
2004 4/.....	*	2.1	6.7	11.0	14.9	18.7	22.7	24.1	25.8	23.8
2003 4/.....	*	2.7	6.8	10.7	17.5	22.7	25.3	25.6	31.7	27.1
2002.....	*	3.2	6.0	11.5	18.8	17.0	22.4	33.1	22.9	*
2001.....	*	3.6	7.0	13.0	15.9	17.5	26.3	30.7	22.5	*
2000.....	*	3.2	7.6	14.3	20.7	19.8	30.1	29.3	27.0	*
1999.....	1.6	3.9	6.9	14.4	17.8	21.0	22.7	19.9	27.9	24.6
Black, female										
2005 4/.....	*	*	1.8	*	*	3.7	*	*	*	*
2004 4/.....	*	*	*	*	*	*	*	*	*	*
2003 4/.....	*	*	*	*	3.6	*	*	*	*	*
2002.....	*	*	*	*	*	*	4.7	*	*	*
2001.....	*	*	*	*	3.4	*	5.6	*	*	*
2000.....	*	*	*	*	*	3.8	5.5	7.4	*	*
1999.....	*	*	*	*	3.8	4.2	4.5	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of trachea, bronchus and lung (C33-C34)										
All races 3/, both sexes										
2005.....	8.2	20.4	40.1	73.8	142.7	218.7	308.3	372.3	380.0	302.3
2004.....	8.5	20.6	40.4	77.9	144.9	226.5	311.2	372.5	375.1	297.5
2003.....	8.5	21.2	40.7	81.8	149.0	230.0	314.0	376.1	380.1	298.9
2002.....	9.2	20.4	41.6	86.3	152.9	233.4	320.9	376.9	378.5	297.2
2001.....	9.5	20.5	42.3	88.8	154.5	239.7	323.2	373.4	368.6	302.7
2000.....	9.3	21.0	43.8	92.5	159.8	245.8	325.4	368.2	374.7	302.1
1999.....	9.3	19.9	44.4	93.0	160.7	243.6	322.3	361.7	362.6	297.0
All races 3/, male										
2005.....	8.5	22.8	49.0	91.0	177.3	275.5	399.2	497.5	540.5	468.0
2004.....	9.1	23.5	49.8	95.8	179.7	285.8	400.8	508.4	530.2	466.8
2003.....	9.5	24.8	50.0	101.3	183.7	292.1	411.0	514.2	541.5	475.1
2002.....	9.6	23.6	51.4	107.0	192.7	297.5	423.0	515.5	546.8	482.2
2001.....	10.5	24.3	52.2	109.6	195.7	312.9	434.7	522.3	541.0	512.4
2000.....	10.5	25.7	53.2	115.9	202.3	322.5	440.5	516.5	558.1	521.2
1999.....	10.5	24.3	54.4	118.1	207.7	322.4	439.2	513.4	551.3	529.2
All races 3/, female										
2005.....	8.0	18.1	31.7	57.6	111.2	169.1	234.7	281.3	280.3	226.2
2004.....	7.9	17.8	31.5	61.1	113.2	174.9	239.1	274.9	279.5	221.3
2003.....	7.5	17.8	31.9	63.5	117.4	176.2	236.1	277.9	281.4	221.0
2002.....	8.8	17.3	32.2	66.8	116.8	177.9	239.3	279.1	276.1	217.0
2001.....	8.4	16.8	32.8	69.2	117.2	176.7	234.5	269.1	265.2	215.0
2000.....	8.1	16.5	34.7	70.5	121.3	180.2	234.7	264.9	266.6	212.8
1999.....	8.1	15.7	34.9	69.5	118.1	176.1	230.9	256.2	252.2	203.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of trachea, bronchus and lung (C33-C34)										
White, both sexes										
2005 4/.....	8.1	19.6	38.7	72.0	142.7	222.2	315.0	380.7	385.6	305.6
2004 4/.....	8.1	19.7	38.8	76.1	145.5	229.2	318.0	378.3	378.9	302.0
2003 4/.....	8.2	19.9	39.1	80.7	149.9	233.5	319.7	381.3	384.5	302.4
2002.....	8.8	19.4	40.0	84.9	152.6	236.4	327.0	382.1	384.1	300.9
2001.....	9.1	19.3	40.9	87.9	155.9	242.0	327.4	377.2	371.2	305.4
2000.....	8.6	19.5	42.2	91.5	161.2	248.2	328.0	370.7	380.7	304.5
1999.....	8.8	18.4	42.6	92.0	160.6	244.5	324.6	362.9	365.2	298.8
White, male										
2005 4/.....	8.1	21.5	46.6	86.3	173.3	275.0	401.3	503.1	543.8	469.9
2004 4/.....	8.5	21.8	46.4	91.4	175.5	284.9	401.9	508.8	530.1	472.2
2003 4/.....	8.9	22.8	46.9	97.3	180.4	291.1	412.1	513.0	541.9	478.5
2002.....	8.9	21.8	48.1	102.8	187.3	295.9	424.5	514.5	547.8	486.6
2001.....	9.7	22.2	49.0	105.6	193.4	309.8	432.8	519.4	537.4	513.9
2000.....	9.3	23.5	49.5	111.5	200.5	319.1	436.8	511.6	560.0	522.4
1999.....	9.7	21.9	50.5	114.3	203.1	317.9	435.0	508.8	548.0	531.5
White, female										
2005 4/.....	8.0	17.7	30.9	58.3	114.2	175.2	243.6	290.1	286.4	229.9
2004 4/.....	7.7	17.7	31.3	61.3	117.7	179.7	249.0	282.8	285.0	225.3
2003 4/.....	7.5	17.0	31.5	64.8	121.6	182.6	244.0	286.0	287.4	224.4
2002.....	8.7	17.1	32.0	67.7	120.6	184.0	247.5	287.3	283.8	220.2
2001.....	8.4	16.4	33.1	71.0	121.3	182.5	242.1	276.4	270.7	218.1
2000.....	8.0	15.6	35.0	72.5	125.0	186.2	240.7	271.6	274.1	215.9
1999.....	7.9	15.0	35.0	70.8	121.6	180.5	236.9	260.4	257.3	205.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of trachea, bronchus and lung (C33-C34)										
Black, both sexes										
2005 4/.....	10.9	30.9	60.6	107.8	180.3	248.0	316.4	361.1	374.0	289.2
2004 4/.....	12.7	31.0	62.7	114.0	179.5	256.4	309.0	382.2	387.7	277.5
2003 4/.....	12.4	35.4	62.0	114.8	182.9	256.6	328.9	387.1	377.2	280.9
2002.....	14.1	32.4	64.4	120.6	193.7	267.2	330.7	391.2	366.3	272.3
2001.....	14.5	32.6	63.5	119.3	182.2	273.6	347.6	395.1	370.3	280.7
2000.....	15.7	35.5	69.3	123.1	188.7	279.9	359.1	394.9	343.3	288.1
1999.....	14.5	35.1	70.8	125.2	198.4	282.1	351.1	402.9	363.5	290.5
Black, male										
2005 4/.....	12.1	38.4	79.5	156.2	262.7	363.2	469.6	544.5	597.6	495.5
2004 4/.....	14.5	40.8	89.1	161.1	271.7	367.8	467.4	600.5	627.7	481.8
2003 4/.....	15.6	45.1	86.6	167.9	267.8	376.2	495.0	618.5	628.3	502.5
2002.....	17.0	42.6	91.6	172.1	292.0	398.4	506.4	630.6	619.0	484.6
2001.....	18.3	45.0	92.4	174.1	268.7	418.8	552.7	659.8	659.7	549.2
2000.....	21.4	47.4	100.4	185.2	271.7	432.2	561.6	656.8	617.1	562.8
1999.....	18.6	48.8	103.6	186.3	301.7	429.2	556.6	658.7	665.3	574.2
Black, female										
2005 4/.....	9.9	24.4	44.6	67.7	116.2	162.5	211.5	251.7	255.9	205.9
2004 4/.....	11.1	22.5	40.4	75.2	107.7	173.8	201.4	252.8	261.2	196.5
2003 4/.....	9.6	26.9	41.2	71.2	116.7	167.8	216.9	249.6	245.6	194.8
2002.....	11.5	23.6	41.5	78.3	116.8	169.8	212.8	248.1	234.7	191.0
2001.....	11.1	21.9	38.8	74.3	114.1	166.3	211.0	237.1	222.8	181.0
2000.....	10.7	25.1	42.7	72.2	123.2	167.6	224.7	237.0	205.5	185.9
1999.....	10.8	23.3	42.9	75.3	117.3	173.2	214.7	248.9	213.9	184.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant melanoma of skin (C43)										
All races 3/, both sexes										
2005.....	1.5	2.2	3.5	4.6	5.9	8.1	10.9	15.1	16.2	19.8
2004.....	1.4	2.3	3.3	4.7	6.1	8.1	10.6	13.6	16.3	17.7
2003.....	1.6	2.5	3.4	4.9	6.4	7.3	10.3	13.9	15.5	17.6
2002.....	1.7	2.3	3.4	4.4	6.0	7.8	10.0	12.9	14.9	17.9
2001.....	1.8	2.6	3.6	4.6	6.1	7.8	10.7	12.4	14.7	17.8
2000.....	1.7	2.7	3.8	4.7	6.3	7.9	10.0	12.0	14.9	18.9
1999.....	1.7	2.6	3.6	4.7	6.1	8.3	9.7	12.9	14.2	16.4
All races 3/, male										
2005.....	1.9	2.7	4.6	6.4	7.9	11.9	16.2	23.5	26.0	34.9
2004.....	1.8	2.9	4.5	6.6	8.9	11.8	16.1	21.8	26.7	31.4
2003.....	1.9	3.1	4.6	6.9	8.9	10.4	15.1	22.3	24.9	30.2
2002.....	2.1	3.3	4.4	6.1	8.5	11.3	14.9	19.8	23.3	29.2
2001.....	2.2	3.2	4.7	6.3	8.8	11.9	16.3	19.6	24.0	29.9
2000.....	2.1	3.2	5.2	6.5	8.7	11.2	14.7	18.4	24.7	31.8
1999.....	2.2	3.5	4.6	6.7	8.4	12.0	14.6	19.8	22.7	27.5
All races 3/, female										
2005.....	1.1	1.8	2.4	2.9	4.2	4.8	6.6	9.0	10.1	12.9
2004.....	1.1	1.7	2.2	3.0	3.6	4.9	6.2	7.7	9.9	11.5
2003.....	1.3	1.9	2.2	3.1	4.0	4.6	6.4	7.8	9.8	12.1
2002.....	1.2	1.4	2.5	2.8	3.6	4.8	6.1	7.9	9.8	12.9
2001.....	1.4	1.9	2.6	3.0	3.7	4.3	6.4	7.3	9.2	12.8
2000.....	1.4	2.1	2.4	3.0	4.1	5.0	6.3	7.5	9.1	13.7
1999.....	1.3	1.8	2.6	2.8	4.0	5.2	5.8	8.1	9.2	11.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant melanoma of skin (C43)										
White, both sexes										
2005 4/.....	1.8	2.6	4.1	5.3	6.8	9.3	12.3	16.8	17.6	21.4
2004 4/.....	1.7	2.8	3.9	5.5	7.0	9.2	12.0	15.1	17.6	19.2
2003 4/.....	1.9	2.9	3.9	5.6	7.3	8.3	11.6	15.2	17.0	19.1
2002.....	2.0	2.8	4.0	5.0	6.8	8.9	11.0	14.2	16.3	19.2
2001.....	2.2	3.1	4.3	5.3	7.0	8.8	11.9	13.6	15.9	19.3
2000.....	2.0	3.1	4.4	5.3	7.2	8.9	11.0	13.1	16.0	20.3
1999.....	2.1	3.1	4.2	5.4	6.9	9.4	10.7	14.0	15.3	17.6
White, male										
2005 4/.....	2.2	3.1	5.4	7.3	9.0	13.6	18.3	26.0	28.3	38.1
2004 4/.....	2.1	3.5	5.3	7.5	10.1	13.4	18.1	24.0	28.9	34.2
2003 4/.....	2.3	3.6	5.3	7.9	10.1	11.7	17.1	24.4	27.1	32.9
2002.....	2.5	3.8	5.1	6.9	9.7	12.7	16.4	21.8	25.6	31.7
2001.....	2.6	3.8	5.5	7.2	10.0	13.3	18.0	21.5	26.1	32.7
2000.....	2.5	3.8	6.0	7.3	9.9	12.5	16.2	20.1	26.6	34.7
1999.....	2.7	4.1	5.3	7.7	9.5	13.4	16.2	21.5	24.5	29.7
White, female										
2005 4/.....	1.3	2.1	2.8	3.4	4.7	5.5	7.4	10.0	10.9	13.8
2004 4/.....	1.4	2.1	2.6	3.5	4.1	5.4	7.0	8.6	10.6	12.4
2003 4/.....	1.5	2.3	2.6	3.5	4.6	5.2	7.1	8.5	10.8	12.9
2002.....	1.5	1.7	2.9	3.2	4.2	5.5	6.7	8.8	10.7	13.8
2001.....	1.7	2.3	3.1	3.5	4.2	4.8	7.0	8.0	9.8	13.7
2000.....	1.6	2.5	2.8	3.4	4.6	5.7	6.9	8.2	9.7	14.5
1999.....	1.5	2.2	3.0	3.3	4.6	5.8	6.4	8.8	9.9	12.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasm of breast (C50)										
All races 3/, both sexes										
2005.....	7.4	11.8	17.9	25.0	33.1	38.7	49.3	63.5	82.9	122.7
2004.....	7.3	11.7	18.7	26.1	33.1	40.8	49.6	64.7	82.4	124.2
2003.....	7.9	12.6	19.1	26.6	33.5	39.9	51.8	66.0	87.3	132.5
2002.....	7.9	12.9	19.7	26.2	33.7	41.3	52.4	67.1	89.8	134.6
2001.....	8.2	13.4	20.6	27.1	34.1	41.7	54.0	69.1	88.0	134.2
2000.....	8.3	13.6	20.8	28.1	35.0	43.4	55.2	70.9	90.9	147.4
1999.....	7.9	13.3	21.0	28.1	34.8	43.5	55.9	70.0	92.4	144.4
All races 3/, male										
2005.....	*	*	0.3	0.4	0.7	1.0	1.7	1.8	2.5	2.4
2004.....	*	*	0.2	0.4	0.6	0.9	1.3	2.1	2.1	3.3
2003.....	*	*	0.4	0.4	0.7	0.9	1.2	1.5	2.8	3.9
2002.....	*	*	0.3	0.4	0.7	0.8	1.3	1.9	2.7	3.4
2001.....	*	*	0.4	0.5	0.8	1.2	1.7	2.3	2.4	3.5
2000.....	*	*	0.3	0.7	0.8	1.1	1.7	2.1	2.4	4.2
1999.....	*	*	0.4	0.6	0.7	1.1	1.4	1.9	2.3	4.5
All races 3/, female										
2005.....	14.7	23.2	34.9	48.3	62.6	71.6	87.9	108.3	132.8	177.9
2004.....	14.5	23.0	36.3	50.4	62.6	75.6	88.5	109.6	131.8	178.6
2003.....	15.6	24.7	37.0	51.3	63.4	73.7	92.4	111.9	139.0	189.4
2002.....	15.6	25.3	38.3	50.4	63.7	76.2	93.2	113.2	142.7	191.5
2001.....	16.2	26.3	40.1	52.0	64.3	76.5	95.6	115.9	139.4	188.9
2000.....	16.4	26.6	40.4	53.8	66.0	79.6	97.3	118.8	143.1	205.7
1999.....	15.6	26.1	40.8	53.9	65.6	79.7	98.4	117.4	145.1	200.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasm of breast (C50)										
White, both sexes										
2005 4/.....	6.6	10.4	16.4	23.4	32.0	38.1	49.2	63.6	83.6	123.4
2004 4/.....	6.5	10.5	17.1	25.1	32.0	40.6	49.5	65.0	83.4	124.8
2003 4/.....	7.1	11.4	17.8	25.3	32.6	39.2	51.7	66.0	87.5	132.5
2002.....	7.0	11.7	18.5	25.0	33.1	40.8	52.0	67.5	89.0	135.3
2001.....	7.3	12.4	19.4	25.8	33.3	40.9	53.6	69.4	88.2	134.8
2000.....	7.5	12.4	19.7	26.9	34.4	43.3	55.5	71.1	91.7	147.3
1999.....	7.1	12.1	19.8	26.8	33.8	42.8	55.5	69.7	92.8	146.0
White, male										
2005 4/.....	*	*	0.3	0.4	0.6	0.9	1.6	1.6	2.7	2.4
2004 4/.....	*	*	*	0.3	0.6	0.9	1.2	2.2	2.0	3.0
2003 4/.....	*	*	0.4	0.4	0.7	0.9	1.2	1.5	2.9	3.9
2002.....	*	*	0.3	0.4	0.6	0.6	1.3	1.7	2.7	3.3
2001.....	*	*	0.3	0.5	0.7	1.2	1.6	2.1	2.3	3.3
2000.....	*	*	0.3	0.6	0.7	1.1	1.7	2.2	2.3	3.9
1999.....	*	*	0.3	0.5	0.6	1.0	1.3	1.8	2.2	4.4
White, female										
2005 4/.....	13.3	20.7	32.3	45.7	61.1	71.1	88.6	109.5	134.4	179.1
2004 4/.....	12.9	20.9	33.7	48.9	61.1	75.9	89.2	111.0	134.0	179.7
2003 4/.....	14.2	22.7	34.8	49.2	62.0	72.9	93.0	112.7	139.7	189.4
2002.....	13.9	23.2	36.3	48.6	63.1	76.2	93.3	114.7	142.0	192.6
2001.....	14.7	24.5	38.1	50.1	63.4	75.7	95.8	117.0	140.0	189.9
2000.....	15.0	24.4	38.7	51.8	65.4	80.2	98.6	119.7	144.9	205.5
1999.....	14.2	23.9	38.9	51.7	64.4	79.2	98.6	117.4	146.3	202.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasm of breast (C50)										
Black, both sexes										
2005 4/.....	13.6	23.1	31.8	43.6	47.7	53.8	61.5	75.5	91.5	144.3
2004 4/.....	13.7	21.2	31.8	40.0	48.9	51.2	62.8	77.0	90.1	143.5
2003 4/.....	14.1	22.6	31.2	43.4	48.1	54.6	65.1	80.3	106.4	161.6
2002.....	15.3	22.9	32.1	40.5	46.3	54.5	67.2	78.5	117.5	152.6
2001.....	14.8	22.6	33.1	42.4	47.4	58.0	70.4	83.5	105.9	152.0
2000.....	14.8	23.9	32.8	43.9	47.3	52.7	65.6	84.2	100.5	176.4
1999.....	14.6	24.2	34.7	44.4	48.6	57.8	69.9	88.0	104.9	147.3
Black, male										
2005 4/.....	*	*	*	*	*	*	*	*	*	*
2004 4/.....	*	*	*	*	*	*	*	*	*	*
2003 4/.....	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*
Black, female										
2005 4/.....	25.4	43.0	58.4	79.1	83.9	91.8	101.5	117.3	139.5	201.8
2004 4/.....	25.6	39.5	58.1	72.4	85.8	88.1	103.7	121.6	135.7	196.9
2003 4/.....	26.3	42.0	57.3	78.3	84.7	94.1	107.4	127.2	160.5	222.4
2002.....	28.7	42.4	58.7	73.1	81.2	92.9	111.1	123.2	176.7	209.1
2001.....	27.5	41.9	60.5	76.7	83.7	99.5	115.0	130.9	157.8	205.9
2000.....	27.7	44.3	60.4	78.8	83.3	91.0	107.4	133.6	149.0	238.7
1999.....	27.5	44.9	63.3	79.6	85.2	99.5	114.1	139.3	154.7	199.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasm of cervix uteri (C53)										
All races 3/, both sexes										
2005.....	1.7	2.2	2.3	2.6	2.6	3.0	3.1	3.5	4.6	5.4
2004.....	1.7	2.1	2.0	2.7	2.9	3.2	3.3	3.3	4.3	5.1
2003.....	1.7	2.0	2.4	2.8	2.9	3.2	3.0	3.7	5.1	5.7
2002.....	1.7	2.1	2.4	2.7	2.8	3.3	3.6	4.1	4.2	6.2
2001.....	1.8	2.3	2.6	2.6	3.2	2.9	3.5	4.0	5.5	7.2
2000.....	1.8	2.2	2.7	3.1	3.1	3.6	3.9	4.3	5.6	7.3
1999.....	1.8	2.4	2.7	3.0	3.5	3.7	3.9	4.7	4.6	7.9
All races 3/, male										
2005.....
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....
All races 3/, female										
2005.....	3.4	4.3	4.6	5.1	5.0	5.7	5.7	6.0	7.5	7.8
2004.....	3.4	4.1	4.0	5.2	5.6	6.0	6.0	5.6	7.0	7.3
2003.....	3.3	4.0	4.6	5.5	5.5	5.9	5.4	6.3	8.3	8.3
2002.....	3.4	4.1	4.7	5.2	5.3	6.1	6.5	7.0	6.7	8.9
2001.....	3.7	4.5	5.1	5.1	6.0	5.4	6.3	6.8	8.9	10.1
2000.....	3.5	4.4	5.2	6.0	5.9	6.7	6.9	7.3	9.0	10.3
1999.....	3.6	4.6	5.2	5.8	6.7	6.8	6.9	8.0	7.2	11.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasm of cervix uteri (C53)										
White, both sexes										
2005 4/.....	1.6	1.9	2.2	2.3	2.3	2.7	2.8	3.0	4.0	4.8
2004 4/.....	1.5	1.9	1.8	2.5	2.6	2.7	2.8	2.9	3.8	4.1
2003 4/.....	1.6	1.9	2.1	2.6	2.6	2.8	2.4	2.8	4.6	4.9
2002.....	1.6	1.8	2.2	2.5	2.4	2.7	3.0	3.6	3.5	5.1
2001.....	1.7	2.2	2.3	2.3	2.9	2.4	3.0	3.4	4.7	6.2
2000.....	1.6	2.0	2.4	2.7	2.6	2.9	3.3	3.8	5.0	6.3
1999.....	1.7	2.3	2.5	2.7	3.0	3.1	3.1	4.0	4.1	6.8
White, male										
2005 4/.....
2004 4/.....
2003 4/.....
2002.....
2001.....
2000.....
1999.....
White, female										
2005 4/.....	3.1	3.8	4.3	4.5	4.5	5.1	5.2	5.2	6.6	6.9
2004 4/.....	3.1	3.8	3.5	4.9	5.1	5.1	5.1	5.0	6.1	5.9
2003 4/.....	3.2	3.7	4.2	5.1	4.9	5.2	4.4	4.9	7.4	7.1
2002.....	3.1	3.7	4.3	4.9	4.5	5.2	5.4	6.2	5.7	7.3
2001.....	3.5	4.3	4.6	4.6	5.5	4.4	5.5	5.8	7.6	8.8
2000.....	3.1	4.0	4.7	5.3	5.0	5.5	5.9	6.5	8.0	8.8
1999.....	3.4	4.5	5.0	5.2	5.7	5.9	5.5	6.8	6.5	9.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasm of cervix uteri (C53)										
Black, both sexes										
2005 4/.....	3.0	4.0	3.3	5.1	5.3	6.5	6.2	9.4	12.4	13.1
2004 4/.....	2.9	3.5	3.9	4.7	6.1	7.1	8.1	8.0	10.1	16.1
2003 4/.....	2.4	3.5	3.8	5.1	6.1	7.1	7.7	11.2	11.4	14.6
2002.....	3.0	3.7	4.6	4.2	6.0	8.2	10.1	9.6	12.6	17.5
2001.....	3.0	3.2	4.4	4.6	5.9	7.9	7.7	9.7	15.2	19.0
2000.....	3.1	3.7	4.9	6.2	6.4	9.4	9.7	10.8	13.6	19.7
1999.....	3.0	3.0	4.1	6.1	8.3	8.7	13.1	12.3	9.5	21.7
Black, male										
2005 4/.....
2004 4/.....
2003 4/.....
2002.....
2001.....
2000.....
1999.....
Black, female										
2005 4/.....	5.7	7.6	6.2	9.3	9.5	11.3	10.4	14.9	18.9	18.4
2004 4/.....	5.4	6.6	7.2	8.6	10.9	12.4	13.6	12.7	15.5	22.4
2003 4/.....	4.5	6.5	7.1	9.3	10.8	12.4	12.8	17.9	17.4	20.3
2002.....	5.6	7.0	8.4	7.6	10.7	14.2	16.8	15.4	19.1	24.2
2001.....	5.6	6.1	8.2	8.3	10.6	13.7	12.8	15.5	22.9	26.1
2000.....	5.9	7.0	9.1	11.2	11.4	16.3	16.2	17.4	20.4	27.0
1999.....	5.7	5.5	7.5	11.1	14.8	15.2	21.8	19.7	14.3	29.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)										
All races 3/, both sexes										
2005.....	0.4	0.9	1.8	3.7	6.4	8.7	10.8	13.1	17.4	22.1
2004.....	0.5	0.9	1.8	3.4	5.8	8.8	11.5	13.8	17.7	22.3
2003.....	0.5	0.8	1.9	3.5	6.1	9.0	11.3	13.8	17.1	22.8
2002.....	0.4	0.9	2.0	3.5	5.8	8.4	11.8	14.2	17.0	24.6
2001.....	0.4	0.9	2.0	3.2	5.8	8.2	11.4	14.5	19.2	24.7
2000.....	0.4	0.9	1.8	3.5	6.0	8.4	11.1	14.0	18.6	24.7
1999.....	0.5	0.9	1.7	3.3	5.5	8.8	10.9	14.4	19.4	24.0
All races 3/, male										
2005.....
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....
All races 3/, female										
2005.....	0.8	1.8	3.5	7.2	12.1	16.4	19.6	22.6	28.2	32.2
2004.....	1.0	1.8	3.5	6.7	11.0	16.5	20.8	23.8	28.7	32.3
2003.....	1.0	1.6	3.7	6.8	11.7	16.8	20.4	23.5	27.6	33.0
2002.....	0.9	1.7	3.9	6.9	11.0	15.7	21.2	24.2	27.4	35.2
2001.....	0.8	1.8	4.0	6.2	11.1	15.3	20.4	24.6	30.7	35.0
2000.....	0.9	1.9	3.5	6.9	11.4	15.6	19.8	23.8	29.5	34.8
1999.....	0.9	1.8	3.3	6.3	10.5	16.2	19.5	24.5	30.7	33.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)										
White, both sexes										
2005 4/.....	0.4	0.9	1.7	3.4	5.5	7.9	9.8	12.0	16.9	21.8
2004 4/.....	0.5	0.9	1.6	3.2	5.2	7.7	10.2	13.0	17.4	22.4
2003 4/.....	0.4	0.7	1.9	3.4	5.4	7.9	10.5	13.0	16.2	22.7
2002.....	0.4	0.8	1.9	3.2	5.2	7.2	10.8	13.1	16.3	23.8
2001.....	0.4	0.9	1.9	3.1	5.2	7.2	10.5	13.7	18.6	24.5
2000.....	0.4	0.9	1.8	3.4	5.4	7.3	9.9	13.0	17.5	24.4
1999.....	0.4	0.9	1.6	3.0	5.2	7.6	10.2	13.7	18.8	23.4
White, male										
2005 4/.....
2004 4/.....
2003 4/.....
2002.....
2001.....
2000.....
1999.....
White, female										
2005 4/.....	0.8	1.8	3.4	6.7	10.6	15.0	18.0	20.8	27.5	31.8
2004 4/.....	1.0	1.8	3.2	6.3	10.0	14.5	18.6	22.6	28.2	32.6
2003 4/.....	0.8	1.5	3.7	6.6	10.4	14.9	19.1	22.4	26.2	32.8
2002.....	0.9	1.7	3.7	6.3	10.1	13.6	19.6	22.4	26.3	34.2
2001.....	0.8	1.7	3.8	6.1	9.9	13.6	18.9	23.5	29.8	34.7
2000.....	0.8	1.9	3.5	6.7	10.3	13.6	17.8	22.2	28.0	34.3
1999.....	0.9	1.8	3.2	5.9	9.9	14.2	18.4	23.3	30.0	32.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)										
Black, both sexes										
2005 4/.....	*	1.3	2.5	6.1	15.1	18.0	22.7	27.6	27.4	31.3
2004 4/.....	0.7	1.0	3.0	5.2	11.9	20.2	26.4	26.6	25.5	26.2
2003 4/.....	0.9	1.1	2.5	5.3	13.9	19.9	22.4	25.5	31.2	30.0
2002.....	*	1.1	2.9	6.6	10.8	20.0	25.2	27.8	29.0	38.9
2001.....	*	1.4	2.9	4.6	11.6	18.8	24.2	26.6	31.1	32.2
2000.....	*	1.0	2.4	5.3	12.0	19.6	26.4	27.7	35.0	33.2
1999.....	0.8	*	2.2	5.8	10.5	20.9	21.8	25.7	29.8	35.4
Black, male										
2005 4/.....
2004 4/.....
2003 4/.....
2002.....
2001.....
2000.....
1999.....
Black, female										
2005 4/.....	*	2.5	4.7	11.2	26.9	31.4	38.3	44.0	41.9	43.9
2004 4/.....	1.4	1.9	5.5	9.4	21.1	35.3	44.4	42.3	38.9	36.6
2003 4/.....	1.6	2.1	4.7	9.6	24.8	34.7	37.4	40.7	47.6	41.7
2002.....	*	2.0	5.4	12.1	19.2	34.8	42.1	44.5	44.2	53.8
2001.....	*	2.6	5.4	8.3	20.8	32.7	40.4	42.4	47.0	44.2
2000.....	*	1.9	4.4	9.6	21.4	34.0	44.0	44.4	52.6	45.5
1999.....	1.6	*	4.1	10.5	18.8	36.4	36.2	41.2	44.5	48.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasm of ovary (C56)										
All races 3/, both sexes										
2005.....	1.3	2.8	5.3	8.4	12.2	16.5	22.3	28.0	34.7	38.1
2004.....	1.4	2.9	5.2	7.9	11.9	17.8	22.5	28.9	36.4	37.7
2003.....	1.4	2.9	5.2	8.3	13.0	17.3	21.9	29.4	36.9	38.4
2002.....	1.4	3.2	5.5	8.3	12.9	17.3	22.3	30.3	37.8	38.1
2001.....	1.4	2.9	5.5	8.3	13.0	17.6	22.6	30.2	37.4	39.6
2000.....	1.5	3.1	5.5	8.3	12.2	17.5	21.8	30.9	36.5	40.9
1999.....	1.5	3.1	5.5	8.3	12.6	17.0	22.4	29.2	35.5	37.6
All races 3/, male										
2005.....
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....
All races 3/, female										
2005.....	2.7	5.5	10.3	16.3	23.4	30.9	40.4	48.3	56.3	55.6
2004.....	2.8	5.8	10.1	15.3	22.7	33.3	40.7	49.7	58.8	54.6
2003.....	2.8	5.7	10.2	16.2	24.8	32.3	39.5	50.4	59.5	55.4
2002.....	2.8	6.4	10.7	16.2	24.6	32.3	40.1	51.7	60.8	54.6
2001.....	2.8	5.8	10.8	16.0	24.7	32.8	40.5	51.4	59.8	56.1
2000.....	2.9	6.0	10.8	16.1	23.3	32.5	39.0	52.5	58.0	57.6
1999.....	3.0	6.2	10.8	16.0	24.0	31.6	39.9	49.5	56.2	52.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasm of ovary (C56)										
White, both sexes										
2005 4/.....	1.3	2.9	5.5	8.8	12.7	16.9	22.8	28.8	35.5	39.4
2004 4/.....	1.5	3.0	5.3	8.0	12.4	18.2	23.1	29.8	37.8	38.6
2003 4/.....	1.4	3.0	5.4	8.7	13.5	17.9	22.4	30.3	38.1	39.1
2002.....	1.4	3.3	5.8	8.6	13.5	18.0	22.9	31.1	38.8	39.0
2001.....	1.4	3.1	5.7	8.5	13.2	18.1	23.3	31.1	38.6	40.4
2000.....	1.5	3.2	5.6	8.7	12.7	18.0	22.4	32.0	37.6	42.0
1999.....	1.6	3.3	5.7	8.6	13.0	17.5	22.9	29.6	36.4	38.7
White, male										
2005 4/.....
2004 4/.....
2003 4/.....
2002.....
2001.....
2000.....
1999.....
White, female										
2005 4/.....	2.7	5.7	10.9	17.2	24.4	32.0	41.6	50.1	57.8	57.5
2004 4/.....	2.9	5.9	10.4	15.8	23.8	34.4	42.1	51.6	61.3	56.0
2003 4/.....	2.9	6.0	10.6	17.1	25.9	33.7	40.7	52.1	61.5	56.4
2002.....	2.9	6.6	11.4	16.8	25.9	33.9	41.6	53.4	62.5	55.9
2001.....	2.9	6.2	11.4	16.6	25.3	33.9	42.2	53.1	61.9	57.3
2000.....	3.0	6.4	11.1	17.0	24.5	33.7	40.3	54.5	60.0	59.0
1999.....	3.2	6.5	11.2	16.8	25.1	32.8	41.0	50.5	57.9	54.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasm of ovary (C56)										
Black, both sexes										
2005 4/.....	1.4	2.3	4.2	6.9	11.1	15.6	22.4	25.2	31.1	31.3
2004 4/.....	1.1	3.0	4.9	7.5	10.2	18.3	22.5	26.2	29.0	32.7
2003 4/.....	1.2	2.8	5.1	6.8	11.0	15.3	20.6	26.2	29.4	36.8
2002.....	1.2	2.9	4.1	7.7	10.9	14.9	20.0	26.5	32.4	34.7
2001.....	1.4	1.9	4.4	7.9	12.6	16.3	20.7	26.2	27.9	36.5
2000.....	1.5	2.2	4.8	6.4	10.9	16.8	19.6	26.1	27.4	34.4
1999.....	1.1	2.4	5.0	7.2	11.4	15.0	22.7	28.9	28.3	28.7
Black, male										
2005 4/.....
2004 4/.....
2003 4/.....
2002.....
2001.....
2000.....
1999.....
Black, female										
2005 4/.....	2.7	4.3	7.8	12.5	19.7	27.1	37.7	40.3	47.5	43.9
2004 4/.....	2.0	5.7	9.1	13.7	18.1	31.8	37.8	41.8	44.2	45.7
2003 4/.....	2.3	5.3	9.4	12.5	19.6	26.7	34.6	41.8	44.9	51.0
2002.....	2.3	5.3	7.7	13.9	19.5	25.9	33.4	42.3	49.3	48.1
2001.....	2.7	3.6	8.1	14.4	22.6	28.3	34.5	41.9	42.1	50.1
2000.....	2.9	4.0	8.8	11.6	19.4	29.2	32.6	41.9	41.2	47.2
1999.....	2.1	4.5	9.3	13.2	20.3	26.2	37.8	46.3	42.3	39.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of kidney and renal pelvis (C64-C65)										
All races 3/, both sexes										
2005.....	0.9	2.0	4.1	6.4	10.8	14.4	19.9	24.3	30.3	32.6
2004.....	0.8	2.1	4.0	6.8	11.3	14.6	19.1	24.5	29.8	32.7
2003.....	1.0	2.0	4.0	7.1	11.4	14.5	19.6	24.8	30.0	33.2
2002.....	0.9	2.4	4.3	7.2	10.9	16.0	19.6	24.6	28.7	32.5
2001.....	1.2	2.4	4.4	7.1	11.9	14.6	20.8	24.4	29.1	32.3
2000.....	1.1	2.2	4.5	7.6	10.8	15.0	19.4	24.5	29.6	33.8
1999.....	1.1	2.2	4.3	7.3	10.8	14.4	19.2	23.1	26.9	31.4
All races 3/, male										
2005.....	1.2	2.9	5.8	9.3	15.7	20.7	28.9	34.1	43.3	50.6
2004.....	1.2	2.9	5.9	9.8	16.6	20.7	27.8	34.2	44.0	47.0
2003.....	1.5	3.0	5.7	10.4	17.2	20.2	27.7	35.9	44.9	48.7
2002.....	1.3	3.4	6.4	10.6	16.1	23.0	27.6	36.6	39.9	50.1
2001.....	1.6	3.5	6.5	10.5	17.2	21.0	30.0	34.4	42.2	52.9
2000.....	1.6	3.1	6.6	11.2	15.6	21.4	27.4	34.6	44.2	55.1
1999.....	1.5	3.2	6.1	10.6	15.8	20.3	27.3	33.7	38.9	49.4
All races 3/, female										
2005.....	0.7	1.2	2.4	3.6	6.4	8.8	12.5	17.2	22.2	24.3
2004.....	0.5	1.3	2.2	3.8	6.4	9.2	12.1	17.6	21.0	26.3
2003.....	0.6	1.0	2.3	4.0	6.1	9.4	13.0	16.9	20.9	26.4
2002.....	0.5	1.5	2.3	3.9	6.2	10.0	13.2	16.2	21.9	24.9
2001.....	0.9	1.4	2.4	3.9	7.0	9.1	13.5	17.4	21.2	23.7
2000.....	0.7	1.4	2.4	4.3	6.5	9.5	13.0	17.5	21.0	25.1
1999.....	0.6	1.2	2.5	4.3	6.2	9.3	12.8	15.7	19.9	24.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of kidney and renal pelvis (C64-C65)										
White, both sexes										
2005 4/.....	0.9	2.0	4.2	6.5	11.0	14.8	20.6	24.9	31.1	33.1
2004 4/.....	0.8	2.2	4.2	7.0	11.7	15.1	19.4	24.9	30.6	33.7
2003 4/.....	1.0	2.0	4.2	7.2	11.6	15.0	20.2	25.4	30.8	33.8
2002.....	1.0	2.4	4.4	7.4	11.2	16.6	20.2	25.0	29.2	33.2
2001.....	1.2	2.5	4.4	7.2	12.1	15.3	21.3	24.7	29.5	32.8
2000.....	1.1	2.3	4.6	7.8	11.1	15.4	19.7	24.9	30.0	34.5
1999.....	1.1	2.1	4.3	7.5	11.2	14.8	19.4	23.4	27.3	31.5
White, male										
2005 4/.....	1.2	2.8	5.9	9.5	15.7	21.3	29.6	34.8	44.2	52.0
2004 4/.....	1.1	3.0	6.2	10.2	17.2	21.3	28.2	34.5	45.4	48.6
2003 4/.....	1.5	3.1	5.9	10.7	17.5	20.8	28.2	36.5	46.2	50.0
2002.....	1.4	3.5	6.6	10.9	16.4	23.8	28.3	37.5	40.2	51.9
2001.....	1.6	3.6	6.5	10.8	17.2	21.9	30.6	34.4	42.8	54.4
2000.....	1.6	3.2	6.7	11.3	16.1	21.5	27.7	35.1	44.5	56.6
1999.....	1.5	3.1	6.0	10.7	16.4	20.6	27.6	34.1	40.1	49.8
White, female										
2005 4/.....	0.7	1.2	2.6	3.7	6.7	9.1	13.1	17.6	22.9	24.3
2004 4/.....	0.5	1.3	2.3	4.0	6.7	9.6	12.2	17.8	21.3	27.0
2003 4/.....	0.6	0.9	2.5	3.8	6.2	9.7	13.7	17.4	21.3	26.6
2002.....	0.5	1.4	2.2	4.1	6.4	10.3	13.6	16.1	22.5	25.1
2001.....	0.8	1.3	2.4	3.8	7.4	9.5	13.8	17.8	21.4	23.8
2000.....	0.7	1.4	2.5	4.5	6.5	10.0	13.3	17.7	21.4	25.5
1999.....	0.6	1.1	2.6	4.4	6.3	9.7	13.0	15.8	19.8	24.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of kidney and renal pelvis (C64-C65)										
Black, both sexes										
2005 4/.....	1.1	2.4	4.4	6.6	12.1	13.5	17.5	22.2	26.2	30.5
2004 4/.....	1.2	2.3	4.0	6.2	10.5	13.1	19.2	26.2	25.3	28.2
2003 4/.....	1.5	2.5	4.1	8.2	11.2	13.4	16.1	22.3	26.1	33.8
2002.....	0.9	2.8	4.4	7.0	10.6	14.3	16.8	25.8	26.7	28.2
2001.....	1.8	2.6	5.0	7.4	12.2	11.7	19.2	24.0	28.2	29.8
2000.....	1.1	2.5	4.7	7.4	10.4	14.8	18.5	23.5	29.1	29.7
1999.....	1.2	2.9	4.9	7.8	9.7	13.5	19.3	23.0	24.3	31.2
Black, male										
2005 4/.....	1.9	3.6	7.1	10.0	19.6	20.0	29.1	32.2	39.3	39.0
2004 4/.....	1.6	3.2	6.1	9.2	16.1	19.6	29.8	38.2	35.9	38.7
2003 4/.....	2.5	3.6	6.5	11.5	17.0	19.5	25.9	35.6	39.1	43.8
2002.....	*	3.5	6.4	11.8	16.8	19.9	23.7	34.4	39.0	39.7
2001.....	1.9	3.5	7.8	10.5	20.5	17.0	29.3	39.0	42.5	46.5
2000.....	*	3.2	7.1	11.4	15.3	25.0	28.1	32.1	48.9	49.6
1999.....	1.8	4.1	8.0	12.5	14.0	20.0	28.2	34.2	27.9	49.2
Black, female										
2005 4/.....	*	*	2.1	3.8	6.2	8.6	9.5	16.3	19.3	27.0
2004 4/.....	*	1.5	2.2	3.7	6.1	8.3	12.1	19.1	19.7	24.0
2003 4/.....	*	1.5	2.1	5.5	6.6	9.0	9.5	14.4	19.3	30.0
2002.....	*	2.3	2.6	3.1	5.7	10.2	12.1	20.6	20.3	23.8
2001.....	1.7	1.8	2.7	4.9	5.6	7.7	12.6	15.0	20.8	23.6
2000.....	*	1.8	2.6	4.2	6.5	7.3	12.2	18.2	19.1	22.3
1999.....	*	1.9	2.3	3.9	6.3	8.7	13.5	16.3	22.5	24.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasm of bladder (C67)										
All races 3/, both sexes										
2005.....	0.4	0.9	1.6	3.5	6.8	11.5	18.9	30.0	46.7	68.3
2004.....	0.4	0.9	1.8	3.6	6.8	11.5	18.9	31.1	45.7	67.9
2003.....	0.3	0.8	1.8	3.5	6.9	11.8	18.5	30.0	45.1	64.8
2002.....	0.4	0.9	1.7	3.8	6.4	12.0	20.2	31.9	44.1	66.6
2001.....	0.4	0.8	1.7	3.4	7.0	12.0	19.0	29.3	44.8	68.9
2000.....	0.3	0.8	1.6	3.6	6.8	11.7	19.6	29.1	45.5	69.6
1999.....	0.3	0.7	1.9	3.8	7.0	11.8	20.2	29.7	43.8	68.5
All races 3/, male										
2005.....	0.5	1.2	2.2	5.3	10.8	18.7	31.2	50.4	86.3	132.1
2004.....	0.5	1.3	2.8	5.7	10.8	18.5	30.6	53.1	81.6	133.0
2003.....	0.4	1.4	2.7	5.5	10.9	19.1	29.6	51.7	80.3	125.1
2002.....	0.5	1.2	2.4	5.9	10.1	19.0	32.2	54.0	76.6	128.9
2001.....	0.5	1.2	2.6	5.2	11.3	19.5	31.6	50.8	81.0	134.8
2000.....	0.4	1.1	2.5	5.7	10.6	19.0	32.2	50.6	82.4	139.4
1999.....	0.4	1.1	2.9	5.9	11.0	19.3	32.9	52.3	79.5	134.8
All races 3/, female										
2005.....	0.3	0.6	1.0	1.7	3.1	5.2	8.9	15.2	22.1	39.0
2004.....	0.3	0.5	0.8	1.7	3.1	5.3	9.5	15.3	23.5	38.6
2003.....	0.2	0.3	0.9	1.7	3.3	5.4	9.5	14.6	23.5	38.1
2002.....	0.3	0.5	1.0	1.9	2.9	6.0	10.6	16.3	24.2	39.6
2001.....	0.3	0.4	0.9	1.7	3.2	5.6	9.0	14.3	23.2	41.4
2000.....	0.3	0.5	0.8	1.7	3.4	5.5	9.7	14.0	23.8	41.2
1999.....	0.2	0.3	0.9	1.7	3.4	5.4	10.3	13.9	22.9	41.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasm of bladder (C67)										
White, both sexes										
2005 4/.....	0.4	0.9	1.6	3.6	7.1	12.0	19.8	31.1	48.7	70.0
2004 4/.....	0.3	0.9	1.8	3.8	7.3	12.3	19.8	32.6	47.6	69.6
2003 4/.....	0.3	0.8	1.8	3.7	7.3	12.2	19.6	31.1	46.2	66.6
2002.....	0.4	0.9	1.8	3.9	6.7	12.5	20.9	32.9	45.2	67.7
2001.....	0.4	0.8	1.8	3.5	7.3	12.6	19.8	30.3	45.7	70.8
2000.....	0.3	0.8	1.7	3.8	7.2	12.3	20.2	29.7	46.4	72.0
1999.....	0.3	0.7	2.0	3.8	7.2	12.1	20.8	30.5	45.0	69.8
White, male										
2005 4/.....	0.5	1.4	2.3	5.6	11.3	19.9	32.8	52.5	90.5	137.4
2004 4/.....	0.4	1.3	2.9	6.0	11.6	20.2	32.3	55.9	86.2	139.3
2003 4/.....	0.4	1.4	2.8	5.7	11.7	19.8	31.6	53.8	83.5	131.0
2002.....	0.5	1.3	2.6	6.2	10.8	20.1	33.7	56.0	79.4	134.1
2001.....	0.5	1.2	2.7	5.4	12.0	20.7	33.1	53.1	83.7	141.1
2000.....	0.4	1.2	2.7	6.0	11.4	20.2	33.6	52.7	84.6	145.7
1999.....	0.4	1.1	3.0	6.2	11.4	20.1	34.4	54.3	82.1	140.7
White, female										
2005 4/.....	0.3	0.5	0.9	1.6	3.2	5.1	9.1	15.3	22.4	39.0
2004 4/.....	0.3	0.5	0.7	1.7	3.3	5.3	9.5	15.6	23.7	38.1
2003 4/.....	*	0.2	0.8	1.7	3.3	5.5	9.8	14.6	23.3	38.1
2002.....	0.3	0.5	1.0	1.7	2.9	5.9	10.4	16.4	24.3	38.9
2001.....	0.3	0.3	0.9	1.7	3.0	5.5	9.0	14.2	22.7	41.3
2000.....	0.3	0.5	0.8	1.8	3.4	5.5	9.6	13.6	23.7	42.1
1999.....	0.2	0.3	1.0	1.5	3.4	5.1	10.0	13.8	23.2	41.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasm of bladder (C67)										
Black, both sexes										
2005 4/.....	*	1.0	2.0	3.7	5.9	9.7	15.7	25.1	36.2	57.5
2004 4/.....	*	1.2	2.0	3.3	5.1	7.9	16.5	23.0	33.4	60.1
2003 4/.....	*	1.2	2.2	3.2	5.0	11.4	12.7	26.5	40.9	53.4
2002.....	*	*	1.6	4.0	5.2	10.9	18.6	27.0	37.2	63.3
2001.....	*	0.9	1.7	3.4	6.9	10.0	14.1	24.8	41.2	55.0
2000.....	*	0.8	1.3	3.0	5.4	9.5	17.9	28.4	44.6	49.7
1999.....	*	1.0	2.2	4.5	6.9	11.8	18.9	25.4	35.3	60.3
Black, male										
2005 4/.....	*	*	2.7	4.9	9.1	13.1	25.0	38.0	56.9	101.3
2004 4/.....	*	1.7	3.0	4.4	8.0	8.2	23.0	35.0	46.6	81.5
2003 4/.....	*	1.7	2.8	4.5	6.6	18.2	16.4	41.1	61.3	80.3
2002.....	*	*	*	4.5	7.1	14.2	23.7	41.8	57.3	91.1
2001.....	*	*	2.8	4.2	9.2	12.7	20.5	34.4	57.8	78.3
2000.....	*	*	*	4.6	7.2	12.5	24.1	41.1	73.3	88.8
1999.....	*	*	3.9	5.4	9.8	15.6	23.7	38.9	59.3	86.7
Black, female										
2005 4/.....	*	*	*	2.7	3.4	7.2	9.3	17.3	25.2	39.8
2004 4/.....	*	*	*	2.5	*	7.7	12.1	15.9	26.5	51.6
2003 4/.....	*	*	1.7	*	3.8	6.4	10.2	17.9	30.2	42.9
2002.....	*	*	*	3.7	3.7	8.5	15.2	18.1	26.7	52.6
2001.....	*	*	*	2.7	5.1	8.1	9.9	19.1	32.7	46.3
2000.....	*	*	*	*	3.9	7.3	13.8	20.8	30.1	35.2
1999.....	*	*	*	3.7	4.7	9.0	15.8	17.2	23.3	50.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)										
All races 3/, both sexes										
2005.....	2.7	3.7	5.6	8.3	10.7	14.0	18.0	20.0	20.4	15.8
2004.....	2.4	3.8	5.5	8.4	10.9	14.0	17.7	19.0	20.4	16.6
2003.....	2.9	3.9	5.8	8.5	11.2	13.8	17.3	19.4	19.6	15.2
2002.....	2.8	4.2	6.2	8.5	11.1	14.7	17.8	19.6	18.5	16.4
2001.....	2.8	4.0	6.1	8.3	11.6	14.2	16.7	20.3	20.6	16.4
2000.....	2.9	4.1	6.3	8.2	11.4	14.8	18.3	19.4	21.1	16.9
1999.....	2.8	4.3	6.1	9.2	11.7	15.7	18.3	21.0	19.4	16.0
All races 3/, male										
2005.....	3.5	4.3	7.0	10.0	13.1	17.7	21.2	25.3	25.3	19.3
2004.....	2.9	4.7	7.1	10.4	13.7	16.0	21.3	23.1	24.5	19.4
2003.....	3.4	5.1	7.4	9.7	13.7	16.4	21.3	24.0	25.4	20.0
2002.....	3.5	5.3	7.2	10.6	13.1	18.4	22.2	24.4	22.9	20.1
2001.....	3.5	4.9	7.5	10.7	13.9	17.3	21.2	25.4	26.5	20.1
2000.....	3.7	4.9	7.7	10.1	14.0	17.9	22.8	23.2	26.3	22.4
1999.....	3.4	5.2	7.4	10.6	14.7	19.5	22.3	26.1	24.5	20.7
All races 3/, female										
2005.....	1.9	3.0	4.3	6.8	8.6	10.7	15.4	16.1	17.4	14.3
2004.....	1.9	2.8	4.0	6.4	8.4	12.2	14.7	16.0	17.9	15.4
2003.....	2.4	2.8	4.3	7.3	9.0	11.6	14.1	16.2	16.0	13.1
2002.....	2.1	3.1	5.1	6.5	9.2	11.4	14.2	16.2	15.8	14.8
2001.....	2.1	3.1	4.8	6.0	9.5	11.5	13.1	16.7	17.1	14.8
2000.....	2.0	3.3	4.9	6.5	9.0	12.1	14.8	16.7	18.0	14.7
1999.....	2.3	3.4	4.9	7.9	8.9	12.4	15.1	17.5	16.5	14.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)										
White, both sexes										
2005 4/.....	3.0	4.0	6.2	9.1	11.6	15.2	19.4	21.1	21.5	16.5
2004 4/.....	2.6	4.1	6.0	9.2	11.9	15.2	19.1	20.2	21.7	17.3
2003 4/.....	3.3	4.3	6.3	9.1	12.2	14.9	18.5	20.6	20.7	15.7
2002.....	3.1	4.7	6.7	9.1	12.0	15.9	19.0	20.7	19.3	17.1
2001.....	3.0	4.4	6.6	8.8	12.6	15.2	17.8	21.6	21.6	17.0
2000.....	3.1	4.5	6.9	8.8	12.3	16.0	19.7	20.5	21.8	17.5
1999.....	3.0	4.7	6.6	10.0	12.5	16.9	19.6	22.2	20.3	16.6
White, male										
2005 4/.....	3.9	4.6	7.6	10.7	14.0	19.3	22.5	26.3	26.5	20.2
2004 4/.....	3.1	5.1	7.6	11.3	14.8	17.4	23.1	24.2	25.8	20.6
2003 4/.....	3.8	5.5	8.0	10.3	14.8	17.4	22.6	25.3	26.6	20.7
2002.....	3.9	5.9	7.9	11.2	14.0	19.9	23.6	25.6	23.7	21.2
2001.....	3.8	5.4	8.1	11.4	14.9	18.6	22.6	26.7	27.8	20.5
2000.....	4.0	5.4	8.3	10.9	15.2	19.4	24.3	24.3	27.5	23.5
1999.....	3.6	5.7	7.9	11.4	15.7	20.9	23.9	27.4	25.8	21.8
White, female										
2005 4/.....	2.1	3.3	4.8	7.4	9.4	11.5	16.7	17.3	18.4	14.8
2004 4/.....	2.1	3.1	4.4	7.2	9.1	13.2	15.7	17.2	19.2	15.8
2003 4/.....	2.7	3.0	4.8	7.9	9.8	12.7	15.2	17.3	17.0	13.4
2002.....	2.3	3.5	5.5	7.1	10.2	12.4	15.3	17.2	16.7	15.4
2001.....	2.3	3.4	5.2	6.4	10.5	12.2	13.9	18.0	17.9	15.6
2000.....	2.1	3.5	5.5	6.9	9.7	13.0	16.0	17.7	18.5	15.0
1999.....	2.5	3.8	5.4	8.7	9.7	13.4	16.2	18.5	17.1	14.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)										
Black, both sexes										
2005 4/.....	1.3	2.4	3.1	5.2	5.6	7.2	9.8	12.9	10.9	9.4
2004 4/.....	1.8	2.2	3.5	4.1	6.0	7.1	9.2	10.0	9.9	10.4
2003 4/.....	1.6	2.7	3.6	5.5	5.6	8.3	9.7	10.6	10.4	10.5
2002.....	1.3	1.9	3.7	5.7	5.6	8.0	9.5	12.2	11.3	9.5
2001.....	1.6	2.6	3.4	5.4	6.4	9.0	9.4	9.4	10.8	11.7
2000.....	2.0	2.2	3.7	5.4	6.0	7.5	8.8	10.3	14.4	10.6
1999.....	1.8	2.6	4.0	5.3	6.7	8.5	7.8	12.0	11.6	9.9
Black, male										
2005 4/.....	1.5	2.9	4.5	6.6	7.5	8.1	13.3	19.2	14.7	*
2004 4/.....	2.8	2.9	5.0	5.9	7.8	7.8	8.1	14.1	14.3	*
2003 4/.....	2.4	3.1	5.1	6.6	6.7	11.8	12.2	16.0	*	*
2002.....	*	2.1	3.3	8.2	7.9	9.5	11.7	15.6	17.6	*
2001.....	1.9	3.0	4.4	6.9	9.2	9.1	11.4	12.5	*	*
2000.....	2.7	2.2	5.7	6.6	7.7	7.8	12.4	11.8	*	*
1999.....	2.6	3.7	5.3	5.9	8.7	10.9	8.8	14.2	*	*
Black, female										
2005 4/.....	*	2.0	1.9	4.0	4.1	6.5	7.4	9.1	8.9	9.8
2004 4/.....	*	1.6	2.2	2.6	4.5	6.7	9.9	7.5	7.6	11.4
2003 4/.....	*	2.3	2.2	4.6	4.7	5.7	8.0	7.3	8.5	11.7
2002.....	*	1.8	4.0	3.7	3.7	6.8	8.1	10.2	8.0	10.3
2001.....	1.3	2.3	2.6	4.1	4.3	8.8	8.1	7.5	10.2	9.7
2000.....	1.5	2.2	2.1	4.5	4.7	7.3	6.4	9.4	14.0	10.7
1999.....	*	1.6	2.8	4.8	5.2	6.7	7.2	10.6	12.5	9.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)										
All races 3/, both sexes										
2005.....	4.1	6.6	11.0	19.0	33.2	52.9	83.8	121.4	161.5	189.8
2004.....	4.1	6.7	11.6	20.2	33.1	54.5	84.2	122.7	162.6	189.4
2003.....	4.7	7.2	11.7	20.1	35.5	55.2	87.3	127.4	165.2	198.0
2002.....	4.6	7.5	12.6	21.5	36.9	59.2	90.1	125.6	168.8	197.5
2001.....	4.8	8.1	13.0	22.7	38.1	58.8	91.8	129.0	168.5	199.9
2000.....	4.8	8.3	13.4	23.4	38.2	62.2	91.6	131.4	176.0	204.6
1999.....	4.8	8.4	13.6	25.6	39.4	62.8	93.9	130.7	174.0	199.4
All races 3/, male										
2005.....	4.9	7.8	13.3	23.9	40.7	68.1	109.5	161.0	214.7	262.4
2004.....	5.0	8.1	14.4	24.9	41.7	70.0	108.7	158.6	214.5	254.9
2003.....	5.8	8.8	14.5	25.2	45.6	71.9	112.3	163.5	216.8	268.7
2002.....	5.5	9.1	15.1	26.8	45.7	76.4	116.8	164.3	224.2	269.1
2001.....	5.9	9.8	16.0	27.7	47.5	75.2	116.9	165.7	220.9	275.3
2000.....	5.7	10.3	16.1	28.4	45.8	78.3	117.7	169.8	230.6	281.8
1999.....	5.9	10.2	16.7	31.0	48.6	78.1	118.9	169.4	221.8	279.0
All races 3/, female										
2005.....	3.3	5.5	8.7	14.4	26.4	39.6	63.0	92.7	128.4	156.4
2004.....	3.2	5.3	9.0	15.7	25.2	40.9	64.5	96.8	130.6	160.0
2003.....	3.5	5.5	9.1	15.3	26.4	40.8	67.3	101.7	133.6	166.8
2002.....	3.7	6.0	10.1	16.4	28.9	44.4	68.8	98.2	135.1	166.5
2001.....	3.7	6.4	10.1	18.1	29.5	44.6	71.8	103.4	137.1	168.4
2000.....	3.9	6.5	10.8	18.8	31.3	48.4	71.0	104.7	143.8	173.1
1999.....	3.8	6.6	10.6	20.5	31.0	49.8	74.3	103.8	145.9	167.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)										
White, both sexes										
2005 4/.....	3.9	6.3	10.5	18.7	33.3	53.6	85.6	125.4	167.1	197.3
2004 4/.....	4.0	6.5	11.2	19.9	33.1	55.0	85.0	125.4	167.2	194.9
2003 4/.....	4.6	7.0	11.3	20.1	35.8	56.1	88.7	130.3	169.5	204.5
2002.....	4.6	7.2	12.1	21.0	36.9	60.3	91.7	128.7	173.6	204.0
2001.....	4.6	7.9	12.8	22.3	38.0	59.0	93.7	131.9	172.8	205.9
2000.....	4.7	8.0	13.2	23.3	38.5	63.5	93.4	134.4	181.5	208.7
1999.....	4.7	8.3	13.4	25.3	39.2	63.8	95.6	133.1	178.4	204.1
White, male										
2005 4/.....	4.7	7.4	12.7	23.5	41.1	69.4	111.8	165.6	221.9	275.3
2004 4/.....	5.0	7.8	13.9	24.8	41.9	70.4	109.7	162.3	221.1	264.0
2003 4/.....	5.7	8.7	13.9	25.1	46.3	73.6	115.2	168.7	224.3	279.5
2002.....	5.5	8.8	14.8	26.5	45.7	77.9	119.2	169.0	231.0	279.4
2001.....	5.7	9.7	15.8	27.1	47.3	75.6	119.2	169.8	226.4	284.8
2000.....	5.8	9.9	15.6	28.3	46.1	80.4	119.0	174.1	238.0	290.2
1999.....	5.6	10.0	16.5	30.6	48.6	79.4	121.0	172.1	228.5	285.8
White, female										
2005 4/.....	3.0	5.3	8.3	14.1	26.1	39.5	64.0	95.7	132.7	161.4
2004 4/.....	3.0	5.2	8.6	15.2	25.0	41.4	64.6	98.3	133.7	163.8
2003 4/.....	3.4	5.4	8.8	15.3	26.1	40.7	67.0	102.6	135.8	171.3
2002.....	3.7	5.7	9.6	15.8	28.8	44.8	69.4	99.9	138.5	171.2
2001.....	3.5	6.1	9.8	17.7	29.3	44.4	73.0	105.0	140.4	172.8
2000.....	3.6	6.1	10.7	18.6	31.4	48.7	72.8	106.5	147.9	175.5
1999.....	3.8	6.6	10.3	20.2	30.5	50.3	75.5	105.7	148.8	171.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)										
Black, both sexes										
2005 4/.....	5.8	9.2	16.0	24.9	38.7	58.5	81.2	103.1	127.4	136.8
2004 4/.....	5.2	9.1	16.8	26.3	40.8	60.7	93.3	115.8	134.7	154.0
2003 4/.....	6.3	9.5	17.3	23.5	41.1	59.4	90.5	117.4	139.1	150.5
2002.....	5.7	10.5	17.3	29.8	43.9	62.3	93.2	111.8	138.2	152.3
2001.....	6.6	10.4	16.5	30.2	44.9	67.8	90.7	116.7	142.7	155.7
2000.....	6.3	12.5	17.5	29.1	44.0	62.2	90.5	114.3	131.6	175.5
1999.....	6.5	10.4	18.2	33.2	45.9	64.6	90.4	119.1	139.3	162.0
Black, male										
2005 4/.....	6.8	11.9	19.9	33.1	46.7	73.4	108.2	139.0	172.1	164.5
2004 4/.....	5.9	11.3	21.4	31.5	51.2	80.4	122.7	139.2	168.6	193.7
2003 4/.....	8.0	11.9	22.5	32.0	51.1	73.4	105.1	137.8	166.8	184.5
2002.....	7.2	13.1	19.7	36.0	55.6	78.8	123.2	137.8	177.3	202.6
2001.....	8.2	11.9	21.0	37.5	56.9	87.9	114.1	141.2	191.0	206.5
2000.....	6.6	14.9	21.8	36.7	54.2	73.1	125.4	147.9	167.8	224.9
1999.....	8.4	13.3	22.5	41.2	55.2	80.7	114.6	159.1	164.8	220.3
Black, female										
2005 4/.....	5.0	6.7	12.8	18.2	32.4	47.4	62.8	81.6	103.9	125.7
2004 4/.....	4.6	7.2	12.9	21.9	32.8	46.1	73.4	101.9	116.8	138.2
2003 4/.....	4.8	7.4	12.9	16.6	33.2	49.0	80.6	105.3	124.6	137.3
2002.....	4.3	8.3	15.2	24.7	34.7	50.1	73.0	96.3	117.8	133.1
2001.....	5.3	9.2	12.7	24.2	35.4	52.9	75.1	102.1	118.1	136.8
2000.....	6.0	10.4	13.9	22.9	36.0	54.1	67.3	94.0	113.4	157.1
1999.....	4.8	8.0	14.5	26.6	38.6	52.7	74.3	95.0	126.6	140.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Hodgkin's disease (C81)										
All races 3/, both sexes										
2005.....	0.3	0.4	0.4	0.5	0.7	1.0	1.2	2.2	2.5	2.1
2004.....	0.3	0.4	0.4	0.7	0.8	1.0	1.4	1.8	2.1	2.2
2003.....	0.3	0.3	0.4	0.6	0.8	0.9	1.5	2.1	2.3	2.4
2002.....	0.3	0.4	0.5	0.5	0.8	1.0	1.6	1.7	2.3	2.7
2001.....	0.4	0.4	0.5	0.4	0.7	1.1	1.2	2.0	2.5	2.2
2000.....	0.3	0.4	0.5	0.5	0.9	1.0	1.6	1.8	2.4	2.5
1999.....	0.3	0.4	0.5	0.7	0.9	1.3	1.8	2.0	2.3	2.4
All races 3/, male										
2005.....	0.3	0.5	0.4	0.8	0.9	1.2	1.6	2.3	2.7	3.1
2004.....	0.3	0.5	0.5	0.9	1.0	1.4	1.9	2.0	3.2	3.2
2003.....	0.5	0.4	0.5	0.7	1.1	1.1	2.0	2.4	2.9	2.6
2002.....	0.4	0.5	0.6	0.7	1.0	1.4	2.2	2.0	3.2	3.6
2001.....	0.5	0.5	0.6	0.6	0.9	1.5	1.6	2.5	2.9	2.6
2000.....	0.4	0.5	0.6	0.5	1.1	1.3	1.9	2.3	2.7	3.7
1999.....	0.4	0.5	0.6	0.9	1.2	1.6	2.3	2.3	2.8	3.8
All races 3/, female										
2005.....	0.2	0.4	0.3	0.3	0.5	0.8	1.0	2.1	2.3	1.6
2004.....	0.3	0.3	0.3	0.5	0.5	0.8	1.0	1.6	1.5	1.7
2003.....	0.2	0.2	0.4	0.5	0.6	0.6	1.2	1.8	1.9	2.3
2002.....	0.3	0.3	0.4	0.3	0.7	0.6	1.1	1.5	1.7	2.3
2001.....	0.3	0.4	0.4	0.3	0.6	0.7	0.9	1.7	2.2	2.0
2000.....	0.3	0.3	0.4	0.5	0.7	0.7	1.3	1.4	2.3	2.0
1999.....	0.3	0.3	0.4	0.5	0.5	0.9	1.5	1.8	2.0	1.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Hodgkin's disease (C81)										
White, both sexes										
2005 4/.....	0.2	0.4	0.4	0.5	0.8	1.1	1.3	2.4	2.6	2.1
2004 4/.....	0.3	0.4	0.4	0.7	0.8	1.1	1.5	1.9	2.2	2.3
2003 4/.....	0.4	0.4	0.5	0.6	0.9	0.8	1.7	2.2	2.4	2.5
2002.....	0.4	0.4	0.5	0.5	0.8	1.0	1.7	1.8	2.4	2.9
2001.....	0.4	0.4	0.5	0.4	0.8	1.2	1.3	2.1	2.6	2.4
2000.....	0.3	0.4	0.5	0.5	0.9	1.0	1.6	2.0	2.5	2.6
1999.....	0.4	0.4	0.5	0.7	0.9	1.3	2.0	2.2	2.5	2.4
White, male										
2005 4/.....	0.3	0.5	0.4	0.7	1.0	1.3	1.6	2.4	2.9	3.4
2004 4/.....	0.4	0.5	0.6	0.9	1.0	1.3	2.0	2.1	3.4	3.5
2003 4/.....	0.5	0.4	0.5	0.7	1.1	1.1	2.0	2.4	3.0	2.7
2002.....	0.4	0.5	0.5	0.7	0.9	1.5	2.3	2.1	3.5	3.9
2001.....	0.5	0.5	0.7	0.6	0.9	1.6	1.8	2.5	3.0	2.7
2000.....	0.4	0.5	0.6	0.5	1.1	1.4	1.9	2.6	2.8	3.9
1999.....	0.5	0.5	0.6	0.9	1.3	1.7	2.4	2.5	3.0	3.8
White, female										
2005 4/.....	*	0.4	0.3	0.4	0.6	0.9	1.0	2.3	2.5	1.5
2004 4/.....	0.2	0.4	0.3	0.5	0.5	0.8	1.0	1.7	1.5	1.7
2003 4/.....	0.2	0.3	0.4	0.5	0.6	0.6	1.4	2.0	2.0	2.4
2002.....	0.3	0.3	0.5	*	0.7	0.6	1.2	1.6	1.7	2.5
2001.....	0.3	0.4	0.4	*	0.7	0.8	0.9	1.8	2.4	2.2
2000.....	0.3	0.3	0.4	0.5	0.7	0.7	1.3	1.6	2.4	2.1
1999.....	0.3	0.4	0.4	0.5	0.6	0.9	1.6	2.0	2.2	1.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Non-Hodgkin's lymphoma (C82-C85)										
All races 3/, both sexes										
2005.....	1.6	2.4	4.3	7.6	12.3	19.2	31.5	46.9	64.4	75.2
2004.....	1.5	2.8	4.8	7.7	12.6	21.1	32.0	46.9	64.8	75.0
2003.....	1.7	3.0	4.7	8.1	14.3	21.6	32.8	49.3	65.4	78.8
2002.....	1.9	3.0	5.0	8.8	14.9	23.2	35.0	50.2	67.0	77.3
2001.....	2.0	3.4	5.5	9.6	15.6	23.0	36.0	50.9	69.8	79.6
2000.....	2.1	3.6	5.8	10.6	16.1	25.0	36.3	53.7	71.8	80.2
1999.....	2.1	3.5	6.2	10.8	16.3	25.6	38.4	54.6	72.6	77.7
All races 3/, male										
2005.....	2.2	3.1	5.7	9.7	15.0	24.5	39.9	59.0	79.4	100.5
2004.....	2.1	3.7	6.1	9.8	15.7	26.7	39.6	57.5	81.6	95.8
2003.....	2.2	3.9	6.0	10.1	18.7	27.8	41.0	60.7	81.8	103.6
2002.....	2.4	3.9	6.2	11.3	18.4	30.0	42.8	61.8	84.4	99.7
2001.....	2.7	4.4	7.4	11.9	19.1	29.2	44.7	61.3	87.3	106.6
2000.....	2.6	4.7	7.5	13.3	19.2	30.5	45.2	66.3	88.1	105.9
1999.....	2.8	4.7	7.9	13.5	19.6	30.8	46.8	68.1	85.7	104.4
All races 3/, female										
2005.....	1.1	1.7	3.0	5.7	9.9	14.5	24.7	38.1	55.2	63.5
2004.....	0.9	1.8	3.5	5.8	9.8	16.2	25.8	39.3	54.5	65.7
2003.....	1.2	2.1	3.5	6.1	10.4	16.2	26.1	41.2	55.4	67.8
2002.....	1.3	2.2	3.8	6.4	11.7	17.3	28.7	41.9	56.4	67.6
2001.....	1.3	2.5	3.8	7.5	12.4	17.7	29.1	43.6	59.3	68.3
2000.....	1.5	2.6	4.1	8.2	13.3	20.3	29.2	45.0	62.2	69.8
1999.....	1.4	2.4	4.5	8.3	13.3	21.1	31.8	45.2	65.0	66.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Non-Hodgkin's lymphoma (C82-C85)										
White, both sexes										
2005 4/.....	1.6	2.4	4.2	7.8	12.6	20.2	33.2	50.0	67.9	79.7
2004 4/.....	1.5	2.7	4.7	7.9	13.1	22.2	33.5	49.2	68.6	78.8
2003 4/.....	1.7	2.9	4.7	8.4	14.9	22.8	34.2	51.8	69.0	83.3
2002.....	1.8	3.0	4.9	9.0	15.6	24.5	36.5	52.9	70.4	82.1
2001.....	2.0	3.4	5.7	9.8	16.2	23.9	37.9	53.6	73.0	83.4
2000.....	2.1	3.6	5.9	11.2	16.9	26.3	38.1	56.5	75.8	84.3
1999.....	2.0	3.6	6.4	11.1	17.0	27.1	40.2	57.0	76.7	81.2
White, male										
2005 4/.....	2.2	3.0	5.5	9.8	15.3	25.7	41.9	62.8	83.6	107.0
2004 4/.....	2.1	3.6	6.1	9.9	16.2	27.7	41.3	60.1	86.2	101.2
2003 4/.....	2.1	3.7	6.0	10.6	19.4	29.6	43.1	63.9	86.5	109.7
2002.....	2.3	4.0	6.2	11.6	19.1	31.6	44.5	65.0	88.6	105.6
2001.....	2.6	4.4	7.4	12.0	19.7	30.3	46.6	64.2	91.2	111.5
2000.....	2.7	4.7	7.6	13.7	20.0	32.2	46.8	69.8	92.6	111.8
1999.....	2.7	4.7	8.1	13.7	20.2	32.3	48.9	70.2	90.8	108.0
White, female										
2005 4/.....	1.0	1.7	3.0	5.8	10.2	15.3	26.0	40.4	58.1	67.1
2004 4/.....	0.9	1.8	3.4	6.0	10.2	17.3	27.1	41.3	57.7	68.7
2003 4/.....	1.2	2.1	3.5	6.4	10.7	16.7	27.0	43.0	58.2	71.6
2002.....	1.3	2.1	3.7	6.5	12.3	18.2	29.9	44.2	59.3	71.8
2001.....	1.3	2.4	3.9	7.7	13.0	18.3	30.9	46.0	62.0	71.6
2000.....	1.4	2.6	4.3	8.8	14.1	21.2	31.1	47.2	65.8	73.1
1999.....	1.4	2.4	4.8	8.7	14.1	22.6	33.3	47.7	68.3	70.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Non-Hodgkin's lymphoma (C82-C85)										
Black, both sexes										
2005 4/.....	2.4	3.2	5.5	7.5	11.5	13.9	20.6	22.4	32.0	33.7
2004 4/.....	1.7	3.9	5.9	7.6	11.0	15.9	21.7	28.9	26.5	38.6
2003 4/.....	2.3	3.9	5.9	6.1	12.7	14.6	22.8	26.5	31.0	31.2
2002.....	2.7	3.5	6.1	9.3	12.8	17.6	25.7	26.8	35.3	30.6
2001.....	2.4	4.3	5.9	9.8	12.3	19.1	23.4	25.7	37.9	39.3
2000.....	2.3	4.5	5.5	8.8	12.6	16.6	22.7	25.9	28.2	36.3
1999.....	2.8	4.1	5.6	10.5	11.7	17.0	23.9	32.0	31.5	40.5
Black, male										
2005 4/.....	2.8	4.4	8.2	10.4	14.7	17.8	29.1	25.9	37.2	40.0
2004 4/.....	2.2	5.4	7.8	10.0	14.4	19.6	28.5	31.8	34.4	41.7
2003 4/.....	3.0	5.7	7.5	7.7	16.0	17.2	24.6	28.8	34.7	34.4
2002.....	3.8	4.0	7.4	11.5	16.6	22.9	34.4	31.7	41.3	42.9
2001.....	3.4	5.1	8.7	12.2	15.5	24.6	31.6	30.7	47.4	54.5
2000.....	2.6	6.1	7.9	13.5	15.7	19.3	32.1	30.7	32.0	43.8
1999.....	3.9	5.6	7.9	13.7	15.7	23.4	28.5	47.0	28.8	59.8
Black, female										
2005 4/.....	2.0	2.1	3.2	5.2	8.9	10.9	14.7	20.3	29.3	31.1
2004 4/.....	1.3	2.5	4.2	5.6	8.3	13.1	17.1	27.2	22.3	37.4
2003 4/.....	1.6	2.3	4.5	4.7	10.1	12.6	21.5	25.2	29.0	30.0
2002.....	1.7	3.1	5.0	7.6	9.8	13.7	19.9	23.9	32.2	25.9
2001.....	1.6	3.6	3.6	7.8	9.8	15.0	17.9	22.7	33.1	33.7
2000.....	2.1	3.2	3.5	5.0	10.1	14.6	16.4	23.1	26.3	33.5
1999.....	1.7	2.8	3.7	7.9	8.7	12.3	20.9	22.9	32.8	33.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Leukemia (C91-C95)										
All races 3/, both sexes										
2005.....	1.7	2.7	4.1	6.5	11.8	19.8	31.3	45.7	61.0	78.7
2004.....	1.8	2.5	4.2	7.1	12.0	19.9	31.8	45.6	61.5	78.8
2003.....	2.1	2.7	4.0	7.0	12.4	19.6	32.5	47.4	62.9	80.3
2002.....	1.9	2.8	4.5	7.4	12.9	20.9	32.4	45.7	63.4	80.5
2001.....	1.9	3.0	4.6	7.6	13.1	20.6	33.3	47.7	61.3	81.4
2000.....	1.9	3.1	4.5	7.7	12.7	21.7	33.2	47.4	65.1	83.6
1999.....	1.9	3.2	4.4	8.5	13.1	21.8	33.1	46.3	62.7	82.4
All races 3/, male										
2005.....	1.8	2.9	4.4	8.3	15.4	26.1	43.9	66.0	87.9	112.4
2004.....	2.1	2.8	5.0	9.0	15.7	26.7	43.9	64.1	86.6	110.7
2003.....	2.4	3.2	4.9	8.9	15.8	26.8	44.9	66.3	88.8	111.8
2002.....	2.2	3.2	5.3	9.2	16.1	28.0	45.8	65.0	90.0	114.5
2001.....	2.1	3.4	5.2	9.2	17.4	27.6	44.9	66.2	86.9	117.5
2000.....	2.1	3.6	5.1	9.7	16.0	29.2	46.2	65.1	93.4	119.8
1999.....	2.1	3.6	5.2	10.1	16.8	29.0	45.7	64.6	87.6	119.2
All races 3/, female										
2005.....	1.6	2.5	3.7	4.9	8.5	14.4	21.1	31.0	44.3	63.3
2004.....	1.6	2.2	3.3	5.4	8.6	14.1	22.1	32.3	46.0	64.5
2003.....	1.7	2.2	3.2	5.2	9.4	13.3	22.7	33.9	47.0	66.4
2002.....	1.6	2.5	3.8	5.7	10.0	14.7	21.8	32.1	47.3	65.7
2001.....	1.7	2.7	4.0	6.0	9.3	14.6	24.1	34.8	45.9	66.3
2000.....	1.7	2.6	3.9	5.9	9.7	15.3	22.9	35.1	48.4	68.8
1999.....	1.7	2.8	3.6	7.0	9.8	15.6	23.2	33.6	48.1	67.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Leukemia (C91-C95)										
White, both sexes										
2005 4/.....	1.7	2.7	4.0	6.7	12.1	20.5	32.7	47.5	63.7	82.4
2004 4/.....	1.8	2.5	4.2	7.3	12.2	20.5	32.5	47.1	63.6	81.6
2003 4/.....	2.1	2.8	4.0	7.1	12.7	20.3	33.6	49.3	65.0	83.4
2002.....	2.0	2.7	4.5	7.4	13.1	21.8	33.6	47.0	65.7	83.0
2001.....	1.9	3.1	4.5	7.6	13.5	20.9	34.5	49.1	63.8	84.7
2000.....	1.9	3.0	4.5	7.8	12.8	22.6	34.4	49.2	67.6	86.0
1999.....	1.9	3.2	4.3	8.7	13.3	22.2	34.2	48.0	64.1	85.4
White, male										
2005 4/.....	1.7	2.9	4.5	8.5	15.8	27.0	45.6	67.9	91.4	118.7
2004 4/.....	2.1	2.8	4.9	9.2	16.0	27.2	44.4	66.2	89.8	115.3
2003 4/.....	2.4	3.4	4.8	9.0	16.4	27.7	46.4	69.2	92.8	116.9
2002.....	2.3	3.0	5.3	9.3	16.3	29.0	47.2	67.0	93.3	119.1
2001.....	2.0	3.5	5.1	9.4	17.8	27.8	46.5	68.3	90.1	123.1
2000.....	2.2	3.5	5.0	9.8	16.1	30.5	47.4	67.5	96.7	123.6
1999.....	2.1	3.5	5.1	10.3	17.0	29.5	47.1	66.7	90.2	124.1
White, female										
2005 4/.....	1.6	2.5	3.6	4.9	8.7	14.7	22.0	32.4	46.4	65.7
2004 4/.....	1.6	2.3	3.4	5.4	8.6	14.5	22.6	33.2	47.3	66.4
2003 4/.....	1.7	2.3	3.2	5.4	9.3	13.8	23.1	34.9	47.9	68.6
2002.....	1.7	2.5	3.8	5.6	10.1	15.4	22.6	32.8	48.9	67.3
2001.....	1.7	2.7	3.9	6.0	9.5	14.8	24.7	35.4	47.9	68.6
2000.....	1.6	2.5	4.0	5.9	9.8	15.7	23.9	36.3	50.2	70.8
1999.....	1.8	2.9	3.6	7.2	9.8	15.8	23.9	34.9	48.7	69.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Leukemia (C91-C95)										
Black, both sexes										
2005 4/.....	2.0	3.1	5.0	6.8	10.7	18.7	24.3	35.1	43.2	52.1
2004 4/.....	2.2	2.9	5.3	7.7	13.8	19.4	32.5	37.1	47.0	60.1
2003 4/.....	2.5	2.5	5.1	7.2	12.8	17.9	29.6	37.3	48.0	59.2
2002.....	1.7	3.6	5.1	9.1	13.8	17.3	28.6	39.2	46.6	64.7
2001.....	2.3	3.0	5.6	7.9	12.3	22.3	28.0	41.3	42.0	57.1
2000.....	2.4	4.5	5.2	7.9	14.1	17.9	28.4	34.7	43.8	66.0
1999.....	2.0	3.5	5.4	8.1	13.6	22.2	27.9	35.0	51.1	55.8
Black, male										
2005 4/.....	2.2	3.7	5.1	8.3	14.0	24.5	34.2	52.7	66.7	62.3
2004 4/.....	2.3	3.1	7.1	9.4	17.6	28.3	48.6	48.2	58.8	79.5
2003 4/.....	3.1	2.7	6.8	10.7	14.5	24.9	39.4	50.2	58.3	77.2
2002.....	1.7	4.8	5.7	11.0	17.2	23.4	42.4	52.4	64.2	86.8
2001.....	2.7	2.8	6.6	9.9	17.1	32.3	36.4	51.6	61.0	77.2
2000.....	2.5	4.9	6.2	10.0	18.6	22.4	43.5	49.6	64.1	98.0
1999.....	2.2	4.4	6.9	10.3	17.2	29.1	38.0	51.3	61.0	77.3
Black, female										
2005 4/.....	1.8	2.6	4.9	5.5	8.2	14.5	17.5	24.5	30.8	48.0
2004 4/.....	2.0	2.6	3.8	6.3	10.9	12.8	21.5	30.5	40.8	52.4
2003 4/.....	2.1	2.4	3.8	4.3	11.4	12.8	23.0	29.6	42.6	52.3
2002.....	1.6	2.6	4.7	7.6	11.2	12.8	19.3	31.3	37.4	56.3
2001.....	2.0	3.3	4.7	6.3	8.5	15.0	22.4	35.2	32.3	49.7
2000.....	2.3	4.2	4.4	6.2	10.5	14.6	18.4	25.6	33.5	54.1
1999.....	1.7	2.6	4.2	6.2	10.8	17.1	21.2	25.2	46.2	47.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Multiple myeloma and immunoproliferative neoplasms (C88,C90)										
All races 3/, both sexes										
2005.....	0.4	1.1	2.2	4.3	8.4	12.8	19.6	26.5	33.4	33.6
2004.....	0.5	1.0	2.3	4.6	7.7	12.3	18.9	28.3	33.9	33.1
2003.....	0.5	1.1	2.5	4.4	7.9	13.1	20.4	28.5	34.5	36.3
2002.....	0.4	1.3	2.5	4.7	8.3	14.1	21.0	27.9	35.8	36.7
2001.....	0.5	1.2	2.3	5.1	8.5	14.0	21.1	28.3	34.7	36.3
2000.....	0.5	1.2	2.6	4.5	8.5	14.4	20.4	28.3	36.4	37.8
1999.....	0.5	1.2	2.5	5.6	9.0	14.1	20.6	27.6	36.2	36.6
All races 3/, male										
2005.....	0.6	1.3	2.7	5.2	9.4	16.3	23.9	33.5	44.5	46.1
2004.....	0.5	1.1	2.7	5.2	9.3	15.3	23.2	35.0	42.8	44.6
2003.....	0.7	1.3	3.1	5.4	10.1	16.0	24.3	33.9	43.3	50.4
2002.....	0.5	1.5	3.0	5.6	10.2	16.8	25.8	35.5	46.2	50.8
2001.....	0.7	1.5	2.8	6.0	10.1	16.8	25.3	35.6	43.4	48.0
2000.....	0.6	1.4	2.9	4.9	9.5	17.1	24.2	35.7	46.1	51.8
1999.....	0.6	1.4	3.0	6.5	11.0	16.6	24.1	34.1	45.5	51.3
All races 3/, female										
2005.....	0.3	0.9	1.7	3.5	7.4	9.9	16.1	21.4	26.5	27.9
2004.....	0.4	0.9	1.8	4.0	6.2	9.8	15.5	23.5	28.4	28.0
2003.....	0.4	0.9	1.9	3.5	6.0	10.6	17.2	24.7	29.2	30.1
2002.....	0.4	1.0	2.0	4.0	6.5	11.7	17.2	22.5	29.5	30.6
2001.....	0.4	0.9	1.9	4.2	7.1	11.5	17.7	23.2	29.4	31.3
2000.....	0.4	1.0	2.3	4.2	7.6	12.1	17.3	23.1	30.7	32.2
1999.....	0.4	1.1	2.1	4.7	7.3	12.0	17.8	23.0	30.7	30.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Multiple myeloma and immunoproliferative neoplasms (C88,C90)										
White, both sexes										
2005 4/.....	0.4	0.9	1.9	3.7	7.7	11.7	18.3	25.5	32.7	32.9
2004 4/.....	0.4	0.9	1.9	4.0	7.1	11.3	17.4	27.0	32.5	32.0
2003 4/.....	0.5	0.9	2.1	3.9	7.3	12.1	19.1	26.9	33.1	35.0
2002.....	0.4	1.1	2.1	4.2	7.5	13.0	19.9	26.9	34.8	35.7
2001.....	0.4	1.0	2.1	4.4	7.5	12.9	19.9	27.1	33.0	35.0
2000.....	0.4	1.0	2.2	3.8	7.8	13.4	19.1	26.5	35.2	35.2
1999.....	0.4	1.1	2.1	4.7	7.9	13.2	19.2	25.7	34.9	34.8
White, male										
2005 4/.....	0.5	1.0	2.3	4.4	8.9	15.2	22.5	32.2	43.9	45.8
2004 4/.....	0.5	1.0	2.3	4.6	8.7	14.2	21.8	33.8	41.2	43.5
2003 4/.....	0.6	1.1	2.6	4.8	9.4	15.1	23.4	32.9	41.9	49.7
2002.....	0.4	1.3	2.7	4.9	9.4	15.8	24.9	34.8	45.3	50.3
2001.....	0.6	1.3	2.5	5.2	9.0	15.7	24.1	34.6	41.7	47.1
2000.....	0.5	1.2	2.5	4.3	8.8	16.1	22.8	33.9	45.4	50.2
1999.....	0.4	1.3	2.6	5.6	10.0	15.8	22.5	32.4	44.3	49.7
White, female										
2005 4/.....	0.3	0.7	1.4	3.1	6.6	8.6	14.9	20.5	25.6	26.9
2004 4/.....	0.3	0.8	1.5	3.3	5.6	8.7	13.8	22.0	27.0	26.8
2003 4/.....	0.3	0.7	1.6	3.1	5.4	9.5	15.6	22.6	27.7	28.5
2002.....	0.3	0.8	1.6	3.4	5.7	10.5	15.8	21.2	28.4	29.4
2001.....	*	0.7	1.6	3.7	6.1	10.4	16.5	21.7	27.8	29.9
2000.....	0.2	0.8	2.0	3.4	6.8	11.1	16.2	21.3	29.2	29.1
1999.....	0.3	1.0	1.5	3.8	5.9	10.9	16.6	21.0	29.4	28.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Multiple myeloma and immunoproliferative neoplasms (C88,C90)										
Black, both sexes										
2005 4/.....	0.9	2.3	5.0	9.9	16.2	25.4	35.1	44.3	51.2	49.4
2004 4/.....	1.0	2.0	5.2	10.2	15.3	24.4	38.1	48.6	59.4	53.9
2003 4/.....	1.2	2.8	5.9	9.4	15.1	25.6	37.3	52.6	59.4	59.8
2002.....	1.0	2.9	5.6	10.7	16.3	26.7	37.7	44.7	54.9	55.5
2001.....	1.5	2.6	4.8	11.8	19.4	25.8	38.2	47.9	61.2	58.3
2000.....	1.1	3.1	6.3	11.7	16.5	27.3	37.9	53.3	57.9	71.6
1999.....	1.4	2.4	6.6	13.8	20.0	24.2	37.6	51.1	56.3	63.1
Black, male										
2005 4/.....	*	3.0	6.2	12.7	17.5	30.4	42.7	58.1	67.4	61.4
2004 4/.....	*	2.1	6.1	11.3	18.1	30.8	44.3	57.8	73.9	71.5
2003 4/.....	1.5	3.2	7.7	12.2	19.7	29.3	39.4	57.5	73.1	73.0
2002.....	*	3.5	5.9	12.4	20.2	31.4	44.7	52.4	70.3	71.8
2001.....	1.6	3.4	5.3	14.5	23.4	30.2	45.1	57.1	80.3	71.5
2000.....	*	3.3	7.2	12.3	18.4	31.0	48.1	67.1	69.1	79.6
1999.....	1.9	2.6	6.7	16.1	21.6	26.2	46.8	59.4	74.1	80.9
Black, female										
2005 4/.....	*	1.7	4.1	7.5	15.2	21.7	29.9	36.0	42.7	44.6
2004 4/.....	*	1.9	4.4	9.3	13.1	19.6	33.8	43.1	51.8	46.9
2003 4/.....	*	2.4	4.4	7.0	11.6	22.8	35.9	49.6	52.2	54.7
2002.....	*	2.4	5.3	9.3	13.3	23.1	32.9	40.1	46.9	49.3
2001.....	1.5	2.0	4.3	9.5	16.3	22.5	33.6	42.4	51.5	53.5
2000.....	*	2.8	5.5	11.2	15.0	24.6	31.1	45.0	52.2	68.7
1999.....	*	2.3	6.6	11.9	18.8	22.7	31.5	46.1	47.5	56.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)

All races 3/, both sexes

2005.....	5.3	9.9	16.6	27.4	43.5	62.7	90.1	121.9	167.4	212.9
2004.....	5.3	9.6	16.2	27.6	43.3	64.6	91.1	126.4	165.2	213.1
2003.....	5.3	10.3	17.7	28.1	44.4	64.6	92.9	131.5	167.7	224.0
2002.....	5.8	10.4	17.5	28.8	46.5	66.0	97.0	131.0	168.7	230.1
2001.....	5.9	10.5	18.0	30.1	46.4	67.2	98.1	133.4	170.9	234.5
2000.....	5.9	10.5	18.2	30.4	48.2	71.5	100.4	136.0	177.4	239.8
1999.....	6.7	11.4	19.1	32.5	52.5	76.1	107.8	142.3	188.5	250.4

All races 3/, male

2005.....	5.9	11.2	19.1	32.8	52.3	74.7	109.2	148.5	208.5	263.5
2004.....	5.5	10.6	18.8	32.5	51.4	78.8	109.3	157.8	207.2	260.6
2003.....	5.9	11.9	20.1	33.0	52.9	78.3	111.3	162.7	208.2	275.0
2002.....	6.2	11.7	20.8	33.0	56.5	79.9	118.9	161.8	207.8	290.8
2001.....	6.4	12.0	21.5	34.9	55.4	79.4	120.3	162.3	211.0	295.2
2000.....	6.4	12.0	20.6	35.1	57.2	86.2	122.5	167.5	221.4	300.7
1999.....	7.6	13.2	22.0	38.2	62.6	90.6	132.8	174.0	233.0	318.9

All races 3/, female

2005.....	4.7	8.7	14.2	22.4	35.6	52.1	74.7	102.5	141.9	189.6
2004.....	5.1	8.6	13.7	23.0	35.9	52.3	76.5	103.8	139.3	191.8
2003.....	4.8	8.7	15.5	23.6	36.7	52.8	78.1	109.2	142.9	201.4
2002.....	5.4	9.1	14.3	24.9	37.5	53.9	79.5	109.2	144.9	203.8
2001.....	5.3	9.1	14.7	25.7	38.3	56.7	80.5	113.1	146.8	209.1
2000.....	5.3	9.1	15.8	25.9	40.0	59.0	83.1	114.1	151.4	215.0
1999.....	5.8	9.7	16.3	27.1	43.3	63.8	88.2	120.3	162.5	222.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)

White, both sexes

2005 4/.....	5.2	9.4	15.8	26.7	42.9	62.4	90.9	123.8	170.0	217.2
2004 4/.....	5.1	9.0	15.4	26.7	42.6	63.9	92.0	127.8	167.1	217.6
2003 4/.....	5.0	9.7	17.0	27.3	44.0	64.7	93.5	132.7	170.1	227.9
2002.....	5.5	9.9	16.8	28.1	45.7	65.6	96.9	131.7	170.0	233.0
2001.....	5.6	9.9	17.2	29.0	45.3	66.5	97.4	134.1	171.4	238.1
2000.....	5.4	9.9	17.2	29.7	47.1	71.4	99.9	135.6	179.5	243.7
1999.....	6.3	10.8	18.3	31.1	51.2	76.3	107.1	142.3	189.4	252.4

White, male

2005 4/.....	5.8	10.4	18.3	31.7	51.4	73.8	109.4	151.6	211.5	270.3
2004 4/.....	5.2	9.9	17.5	31.0	50.2	77.5	109.7	159.9	210.9	266.8
2003 4/.....	5.4	11.0	19.2	31.8	52.2	78.0	111.6	164.1	212.2	280.9
2002.....	5.7	11.1	20.0	32.0	55.1	80.2	118.8	162.2	210.9	296.5
2001.....	6.2	11.3	20.1	33.1	53.5	78.0	119.7	163.1	212.0	300.2
2000.....	5.8	11.2	19.5	34.1	54.7	86.2	121.8	165.7	225.2	307.0
1999.....	7.2	12.2	20.8	36.6	60.9	90.1	131.8	174.6	235.3	323.6

White, female

2005 4/.....	4.5	8.4	13.4	21.9	35.0	52.1	75.7	103.2	144.0	192.8
2004 4/.....	5.0	8.2	13.2	22.5	35.5	51.9	77.5	104.4	139.9	195.4
2003 4/.....	4.6	8.4	14.9	23.1	36.3	52.9	78.8	110.0	144.1	204.4
2002.....	5.4	8.7	13.7	24.3	36.9	52.7	79.1	109.9	144.9	205.5
2001.....	5.1	8.5	14.3	25.1	37.7	56.3	79.3	113.7	146.8	212.1
2000.....	5.0	8.6	14.8	25.6	40.1	58.4	82.4	114.5	152.3	218.0
1999.....	5.3	9.5	15.8	26.0	42.3	64.3	87.5	119.6	162.3	223.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)

Black, both sexes

2005 4/.....	7.2	14.9	25.1	39.4	56.1	76.0	97.9	127.4	168.9	183.9
2004 4/.....	7.5	14.7	25.0	40.6	57.1	84.2	99.1	133.0	167.8	186.7
2003 4/.....	7.8	16.0	26.5	41.1	57.7	77.0	102.4	139.4	166.0	206.3
2002.....	8.0	15.8	25.3	41.5	63.3	82.5	113.5	146.5	174.5	222.1
2001.....	7.7	16.7	27.8	44.1	64.1	85.2	120.3	140.7	182.3	217.1
2000.....	10.0	16.8	28.8	41.3	64.8	85.6	117.7	155.7	172.0	210.8
1999.....	10.1	18.3	29.0	48.3	73.5	85.4	128.3	157.3	193.0	242.6

Black, male

2005 4/.....	7.7	18.1	28.3	50.5	71.6	98.6	127.5	144.4	217.0	221.2
2004 4/.....	8.3	17.1	32.7	52.1	73.3	107.1	128.8	169.7	203.0	222.5
2003 4/.....	9.4	19.8	31.4	53.0	72.3	98.5	128.4	176.2	206.0	241.9
2002.....	9.6	18.4	30.9	50.2	82.0	97.2	142.9	187.4	209.4	267.0
2001.....	8.2	19.0	35.9	57.1	80.9	110.0	151.5	178.9	219.1	283.7
2000.....	11.8	20.8	33.1	51.4	89.0	102.0	149.1	211.7	208.2	253.7
1999.....	11.7	23.2	36.5	61.9	94.0	109.8	163.8	191.4	233.7	292.9

Black, female

2005 4/.....	6.7	12.1	22.4	30.3	44.1	59.2	77.7	117.3	143.6	168.8
2004 4/.....	6.8	12.5	18.5	31.1	44.4	67.3	78.9	111.3	149.3	172.5
2003 4/.....	6.4	12.6	22.3	31.3	46.3	61.0	84.8	117.5	145.1	192.4
2002.....	6.5	13.5	20.5	34.5	48.6	71.6	93.9	122.1	156.4	205.0
2001.....	7.3	14.8	20.9	33.5	50.9	66.8	99.5	117.9	163.5	192.4
2000.....	8.3	13.4	25.1	33.1	45.8	73.5	96.8	121.9	153.7	194.9
1999.....	8.8	14.0	22.7	37.2	57.4	67.3	104.6	136.7	172.9	223.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)

All races 3/, both sexes

2005.....	1.0	1.4	2.1	3.3	5.7	9.5	16.3	27.9	44.8	76.7
2004.....	0.8	1.4	2.0	3.5	5.8	9.3	16.7	27.3	47.2	77.7
2003.....	0.9	1.4	2.0	3.2	6.0	9.8	16.3	28.5	47.4	80.5
2002.....	1.0	1.5	2.1	3.7	5.4	9.7	16.3	28.8	46.3	78.7
2001.....	0.9	1.5	2.3	3.9	6.3	10.5	17.9	29.5	47.8	82.7
2000.....	0.9	1.5	2.3	3.8	6.6	10.6	17.4	28.9	50.6	84.9
1999.....	0.9	1.4	2.3	3.5	6.3	10.8	18.6	29.3	47.5	82.9

All races 3/, male

2005.....	1.0	1.5	2.4	3.6	7.2	12.1	20.3	36.3	59.5	92.7
2004.....	0.9	1.4	2.3	4.0	7.1	11.7	20.5	34.8	61.3	98.9
2003.....	1.0	1.4	2.3	3.6	7.0	11.6	21.2	35.9	61.5	101.7
2002.....	1.0	1.6	2.1	4.5	6.3	12.2	20.0	37.3	59.1	101.1
2001.....	1.0	1.7	2.4	4.1	7.5	12.8	22.5	39.2	62.4	107.6
2000.....	0.9	1.6	2.6	4.1	7.5	13.3	21.2	36.8	65.3	108.9
1999.....	1.1	1.5	2.7	4.0	7.9	13.4	23.9	37.2	61.0	109.2

All races 3/, female

2005.....	1.0	1.3	1.8	3.1	4.3	7.3	13.0	21.8	35.6	69.4
2004.....	0.7	1.3	1.7	3.0	4.6	7.1	13.7	21.9	38.5	68.1
2003.....	0.7	1.3	1.7	2.8	5.1	8.3	12.4	23.2	38.7	71.1
2002.....	0.9	1.5	2.0	3.0	4.6	7.5	13.4	22.8	38.5	68.9
2001.....	0.8	1.2	2.1	3.7	5.2	8.5	14.3	22.7	39.1	72.3
2000.....	1.0	1.3	1.9	3.5	5.7	8.2	14.4	23.4	41.9	75.2
1999.....	0.8	1.3	1.9	3.0	4.8	8.5	14.5	23.8	39.7	72.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)										
White, both sexes										
2005 4/.....	0.9	1.3	2.1	3.3	5.6	9.7	16.6	28.7	46.3	79.6
2004 4/.....	0.7	1.4	1.9	3.4	5.7	9.4	17.1	27.7	48.5	80.2
2003 4/.....	0.9	1.3	2.0	3.1	6.1	9.8	16.8	29.5	48.7	83.0
2002.....	0.9	1.6	2.0	3.7	5.6	9.9	16.7	29.7	47.2	80.7
2001.....	0.9	1.5	2.3	3.9	6.3	10.7	18.1	30.4	48.9	83.9
2000.....	0.9	1.3	2.2	3.9	6.6	10.7	17.3	29.5	52.2	87.3
1999.....	0.9	1.3	2.2	3.5	6.2	10.7	19.0	29.7	48.7	85.3
White, male										
2005 4/.....	0.9	1.4	2.4	3.7	7.1	12.4	21.0	37.5	61.9	97.0
2004 4/.....	0.9	1.4	2.3	3.9	7.0	12.0	21.0	35.4	63.2	103.2
2003 4/.....	1.0	1.3	2.3	3.5	7.1	11.8	21.8	37.4	63.3	105.5
2002.....	1.1	1.7	2.1	4.4	6.4	12.4	20.7	38.4	60.1	104.2
2001.....	1.0	1.7	2.5	4.1	7.4	13.2	23.0	40.4	63.7	109.9
2000.....	0.9	1.5	2.6	4.3	7.3	13.4	21.0	37.6	68.0	112.6
1999.....	1.1	1.4	2.6	3.9	8.0	13.3	24.6	37.7	62.9	114.3
White, female										
2005 4/.....	0.9	1.1	1.8	3.0	4.1	7.3	13.0	22.1	36.5	71.6
2004 4/.....	0.6	1.3	1.5	2.9	4.6	7.0	13.8	22.1	39.3	69.9
2003 4/.....	0.7	1.2	1.7	2.8	5.3	8.0	12.7	23.9	39.8	73.0
2002.....	0.8	1.4	1.8	3.0	4.8	7.6	13.5	23.4	39.2	70.5
2001.....	0.8	1.2	2.0	3.8	5.2	8.6	14.2	23.3	40.0	73.0
2000.....	1.0	1.2	1.9	3.5	6.0	8.4	14.3	23.8	42.9	77.1
1999.....	0.7	1.2	1.8	3.0	4.6	8.3	14.7	24.0	40.4	73.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)										
Black, both sexes										
2005 4/.....	1.6	2.4	2.5	3.9	7.5	9.9	16.4	24.1	33.5	54.5
2004 4/.....	1.3	1.6	3.1	5.5	7.1	10.2	16.1	26.1	39.4	57.0
2003 4/.....	1.0	2.2	2.7	4.2	6.6	11.2	15.3	22.6	39.9	57.8
2002.....	1.7	2.0	3.0	5.2	5.8	10.6	15.0	24.4	41.9	63.6
2001.....	1.2	1.4	2.7	4.5	7.6	10.9	18.6	24.5	42.5	73.4
2000.....	1.4	2.4	3.1	4.2	7.1	10.6	20.3	26.7	37.8	64.1
1999.....	1.3	2.2	3.2	5.0	7.8	13.7	18.1	29.5	38.7	61.9
Black, male										
2005 4/.....	1.7	2.6	2.8	4.3	9.1	10.9	18.0	28.2	42.8	54.8
2004 4/.....	*	1.7	3.5	6.6	9.3	12.4	19.4	34.6	46.6	61.6
2003 4/.....	*	2.1	2.9	5.5	8.5	12.3	21.4	26.5	53.2	66.7
2002.....	*	1.7	2.8	7.4	6.5	12.7	16.0	29.9	53.5	79.3
2001.....	*	*	2.4	4.5	9.2	13.0	22.9	31.6	58.6	88.5
2000.....	*	2.2	3.6	4.4	10.1	15.6	23.7	34.5	38.8	76.1
1999.....	*	*	3.3	6.1	8.3	16.9	21.7	36.1	42.7	63.3
Black, female										
2005 4/.....	1.4	2.2	2.3	3.6	6.2	9.2	15.4	21.6	28.6	54.4
2004 4/.....	1.4	1.6	2.6	4.7	5.4	8.6	13.8	21.0	35.5	55.1
2003 4/.....	*	2.3	2.6	3.1	5.1	10.4	11.3	20.3	32.9	54.3
2002.....	2.1	2.2	3.3	3.3	5.3	9.1	14.3	21.1	35.8	57.5
2001.....	*	*	3.0	4.5	6.4	9.4	15.7	20.2	34.3	67.8
2000.....	1.3	2.5	2.7	4.0	4.7	6.9	18.0	21.9	37.4	59.7
1999.....	1.4	2.5	3.0	4.0	7.5	11.4	15.8	25.5	36.7	61.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Anemias (D50-D64)										
All races 3/, both sexes										
2005.....	0.5	0.6	0.6	0.8	1.2	2.0	3.5	6.3	12.8	36.6
2004.....	0.5	0.5	0.6	0.8	1.3	1.8	3.4	5.9	12.2	38.1
2003.....	0.4	0.6	0.5	0.8	1.3	1.9	3.5	6.0	12.7	40.7
2002.....	0.4	0.6	0.6	0.8	1.2	1.9	3.7	6.4	12.9	40.3
2001.....	0.6	0.5	0.7	0.8	1.2	2.0	3.5	7.1	13.6	42.0
2000.....	0.5	0.5	0.7	0.9	1.2	2.2	3.4	6.5	13.9	42.4
1999.....	0.5	0.7	0.6	0.8	1.2	1.8	3.5	6.8	13.7	43.8
All races 3/, male										
2005.....	0.5	0.5	0.5	0.7	1.3	1.9	3.5	6.4	14.1	33.5
2004.....	0.5	0.5	0.6	1.0	1.4	1.9	3.6	6.3	12.8	36.5
2003.....	0.5	0.7	0.5	0.9	1.5	2.3	3.8	6.6	13.3	38.0
2002.....	0.4	0.5	0.6	0.8	1.3	1.7	4.3	6.6	12.4	38.3
2001.....	0.4	0.6	0.6	0.8	1.1	2.1	3.8	7.2	14.3	40.0
2000.....	0.4	0.5	0.7	1.0	1.2	2.2	3.6	6.5	14.6	40.1
1999.....	0.5	0.7	0.6	0.7	1.2	2.1	3.4	7.2	14.8	43.4
All races 3/, female										
2005.....	0.5	0.7	0.7	0.9	1.2	2.0	3.5	6.3	12.1	38.0
2004.....	0.5	0.6	0.6	0.7	1.3	1.7	3.3	5.7	11.8	38.8
2003.....	0.4	0.5	0.6	0.7	1.0	1.5	3.2	5.6	12.3	41.9
2002.....	0.5	0.7	0.7	0.8	1.1	2.1	3.2	6.2	13.2	41.2
2001.....	0.7	0.5	0.7	0.9	1.2	1.9	3.3	7.0	13.2	42.9
2000.....	0.5	0.6	0.7	0.9	1.2	2.2	3.3	6.5	13.6	43.3
1999.....	0.4	0.7	0.6	0.9	1.1	1.6	3.6	6.5	13.0	44.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Anemias (D50-D64)										
White, both sexes										
2005 4/.....	0.1	0.3	0.3	0.6	1.0	1.6	3.1	6.0	12.4	36.6
2004 4/.....	0.2	0.3	0.3	0.5	1.0	1.7	3.2	5.6	12.2	37.7
2003 4/.....	0.1	0.3	0.3	0.4	0.9	1.6	3.2	5.7	12.4	40.6
2002.....	0.1	0.3	0.3	0.6	0.9	1.7	3.4	5.9	12.6	39.7
2001.....	0.2	0.2	0.4	0.6	1.0	1.8	3.2	6.7	13.2	41.7
2000.....	0.1	0.2	0.4	0.7	0.9	1.9	3.2	6.2	13.8	42.0
1999.....	0.1	0.3	0.3	0.4	0.8	1.6	3.2	6.5	13.6	43.0
White, male										
2005 4/.....	*	0.3	0.3	0.6	1.0	1.5	3.3	5.9	13.5	33.7
2004 4/.....	0.2	0.3	0.3	0.5	1.2	1.7	3.5	5.7	12.7	36.7
2003 4/.....	*	0.3	0.3	0.5	1.1	2.0	3.3	6.3	13.4	37.9
2002.....	*	0.3	0.3	0.6	1.0	1.6	3.7	6.3	11.9	37.2
2001.....	*	*	0.4	0.5	1.0	1.9	3.3	7.0	14.1	40.4
2000.....	*	*	0.4	0.8	0.9	2.1	3.4	6.1	14.2	39.1
1999.....	*	0.4	0.3	*	0.7	1.9	3.1	7.0	14.8	43.4
White, female										
2005 4/.....	*	0.4	0.3	0.7	1.0	1.7	3.0	6.1	11.7	37.9
2004 4/.....	*	0.2	0.3	0.5	0.9	1.6	3.0	5.6	11.9	38.2
2003 4/.....	*	*	0.3	0.3	0.8	1.3	3.1	5.3	11.8	41.8
2002.....	*	0.3	0.3	0.5	0.8	1.8	3.1	5.6	13.0	40.8
2001.....	0.3	*	0.4	0.6	1.1	1.6	3.1	6.5	12.6	42.3
2000.....	*	0.3	0.4	0.7	0.9	1.8	3.0	6.3	13.5	43.2
1999.....	*	0.3	0.3	0.5	0.9	1.4	3.4	6.1	12.9	42.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Anemias (D50-D64)										
Black, both sexes										
2005 4/.....	2.9	2.5	2.9	2.5	3.7	5.3	7.3	11.0	20.1	42.5
2004 4/.....	2.4	2.5	3.0	3.9	4.2	3.7	6.2	10.7	13.4	48.2
2003 4/.....	2.5	3.3	2.7	4.1	4.4	4.3	7.1	10.9	19.8	49.3
2002.....	2.4	3.1	3.1	3.2	4.1	4.4	7.9	13.7	19.4	52.3
2001.....	3.2	3.2	2.9	3.2	2.6	3.9	7.4	12.5	21.4	48.5
2000.....	2.7	2.9	3.2	3.0	3.9	4.8	6.1	10.3	18.1	51.0
1999.....	2.8	3.2	2.9	4.2	4.3	3.6	6.6	11.4	16.5	52.9
Black, male										
2005 4/.....	3.1	2.4	2.6	*	*	6.7	*	13.4	23.2	37.2
2004 4/.....	2.5	2.1	2.9	5.3	4.5	*	*	13.6	14.3	42.7
2003 4/.....	2.9	3.9	2.6	4.9	6.2	5.7	10.8	12.8	18.5	49.0
2002.....	2.3	2.6	2.6	3.0	5.1	*	11.7	12.9	23.7	56.8
2001.....	2.7	3.9	3.2	3.3	*	*	10.4	12.1	20.1	42.0
2000.....	2.2	3.2	3.4	*	*	*	*	10.9	19.4	55.4
1999.....	2.9	3.3	3.2	4.2	6.2	*	7.8	10.9	*	44.5
Black, female										
2005 4/.....	2.7	2.6	3.2	2.9	4.0	4.2	8.2	9.6	18.5	44.6
2004 4/.....	2.4	2.8	3.1	2.7	3.9	*	6.2	8.9	12.9	50.4
2003 4/.....	2.1	2.8	2.7	3.3	3.0	*	4.7	9.8	20.5	49.4
2002.....	2.6	3.6	3.5	3.3	3.3	4.6	5.4	14.3	17.1	50.5
2001.....	3.7	2.6	2.6	3.2	*	3.9	5.4	12.8	22.1	50.9
2000.....	3.2	2.7	3.0	3.0	3.9	5.8	6.2	10.0	17.4	49.4
1999.....	2.6	3.2	2.7	4.2	*	*	5.9	11.7	16.4	56.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Diabetes mellitus (E10-E14)										
All races 3/, both sexes										
2005.....	5.6	9.9	17.3	29.7	47.3	71.7	104.9	151.4	211.2	312.1
2004.....	5.7	10.0	17.2	29.4	47.2	71.3	105.7	149.8	213.1	307.0
2003.....	5.8	10.3	17.9	30.4	48.9	75.0	108.8	155.3	216.5	317.5
2002.....	6.0	9.9	18.0	30.1	47.5	75.7	108.8	157.7	217.9	320.6
2001.....	5.4	9.9	17.8	29.5	48.4	75.5	108.6	155.5	218.7	321.8
2000.....	5.5	9.5	17.3	29.6	48.1	74.0	108.6	156.5	213.9	319.7
1999.....	5.5	9.9	16.5	29.5	48.9	75.6	109.2	153.5	214.8	317.2
All races 3/, male										
2005.....	7.2	12.7	21.0	36.3	56.6	82.4	121.2	174.0	244.1	344.4
2004.....	7.1	12.4	21.2	34.5	56.6	82.7	121.2	172.2	246.9	333.9
2003.....	7.4	12.6	21.3	35.4	57.8	86.8	124.4	179.0	249.7	340.7
2002.....	7.7	11.9	21.5	36.0	55.5	84.3	121.1	175.9	243.5	348.5
2001.....	6.6	12.1	21.3	34.4	54.4	84.7	121.0	171.5	242.3	343.1
2000.....	6.8	11.4	19.7	33.4	53.9	80.8	119.7	173.7	242.5	345.1
1999.....	6.4	12.3	19.4	33.5	53.9	82.5	119.4	172.3	241.3	345.7
All races 3/, female										
2005.....	4.1	7.2	13.8	23.4	38.9	62.3	91.6	134.9	190.8	297.2
2004.....	4.3	7.7	13.4	24.6	38.6	61.4	93.2	133.6	192.2	294.9
2003.....	4.2	8.1	14.7	25.7	40.9	64.8	96.2	138.4	196.2	307.2
2002.....	4.4	7.9	14.7	24.5	40.2	68.2	99.0	144.9	202.3	308.4
2001.....	4.2	7.9	14.5	24.9	42.9	67.6	98.8	144.3	204.5	312.9
2000.....	4.3	7.8	14.9	26.1	42.8	68.1	99.9	144.6	197.0	309.4
1999.....	4.6	7.5	13.8	25.8	44.5	69.7	101.2	140.4	199.3	305.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Diabetes mellitus (E10-E14)										
White, both sexes										
2005 4/.....	4.9	8.6	15.0	25.9	41.8	63.3	94.6	139.9	198.7	296.3
2004 4/.....	5.0	8.9	14.6	25.5	41.3	63.6	94.8	137.4	199.5	289.4
2003 4/.....	5.1	8.8	15.2	26.4	42.7	66.5	98.1	143.1	202.7	300.8
2002.....	5.3	8.5	15.4	26.0	41.3	66.0	98.2	145.4	204.9	303.7
2001.....	4.8	8.6	15.3	25.3	42.3	66.8	98.0	143.0	205.9	306.0
2000.....	5.0	8.0	14.6	25.7	42.3	65.2	98.1	143.7	199.1	303.5
1999.....	4.7	8.3	13.8	24.5	41.9	66.9	97.6	141.2	200.8	298.9
White, male										
2005 4/.....	6.5	11.3	18.4	32.0	51.0	74.8	111.5	164.6	235.4	336.2
2004 4/.....	6.3	11.1	18.7	30.3	51.0	75.0	111.6	162.1	238.5	323.0
2003 4/.....	6.6	11.1	18.7	31.6	51.5	79.1	115.3	170.2	241.1	333.4
2002.....	6.9	10.3	18.8	32.0	49.5	75.9	112.9	167.9	235.5	344.3
2001.....	6.1	10.5	18.6	30.4	48.8	77.6	113.5	161.8	232.1	337.3
2000.....	6.3	9.7	17.3	29.5	48.7	73.8	111.6	164.0	234.1	337.4
1999.....	5.6	10.6	16.8	28.9	47.7	75.2	111.3	163.7	233.2	333.4
White, female										
2005 4/.....	3.3	5.9	11.6	20.0	33.2	53.1	80.7	121.7	175.7	278.0
2004 4/.....	3.6	6.7	10.6	20.8	32.4	53.4	81.1	119.4	175.3	274.3
2003 4/.....	3.5	6.5	11.8	21.4	34.5	55.4	84.0	123.5	179.1	286.3
2002.....	3.8	6.6	12.0	20.2	33.7	57.3	86.3	129.3	186.1	286.1
2001.....	3.4	6.7	12.0	20.4	36.3	57.4	85.4	129.7	190.1	293.0
2000.....	3.7	6.4	12.0	22.1	36.3	57.6	87.3	129.3	178.3	289.7
1999.....	3.9	6.1	11.0	20.4	36.6	59.6	86.7	125.4	181.7	285.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Diabetes mellitus (E10-E14)										
Black, both sexes										
2005 4/.....	10.8	20.7	36.3	63.6	102.9	150.2	211.2	283.6	377.0	506.5
2004 4/.....	11.5	19.2	38.8	64.8	102.0	147.7	219.3	297.6	384.2	532.4
2003 4/.....	11.7	22.1	39.9	69.8	106.9	156.1	221.2	300.0	389.9	527.5
2002.....	12.0	21.4	40.7	67.8	105.5	161.6	221.6	300.3	392.7	537.8
2001.....	10.7	21.0	40.2	69.6	104.4	157.6	226.8	305.2	382.2	529.7
2000.....	10.1	21.9	40.7	66.3	102.4	159.2	223.7	316.5	404.9	524.9
1999.....	11.3	22.5	40.5	72.7	111.6	159.6	229.0	294.9	395.8	529.0
Black, male										
2005 4/.....	12.3	25.9	42.2	78.0	118.2	161.8	236.9	310.7	391.9	464.8
2004 4/.....	13.9	24.2	44.1	74.6	115.7	168.6	238.3	309.8	389.5	505.7
2003 4/.....	14.4	25.4	44.0	77.3	121.7	170.6	232.6	306.3	387.6	453.5
2002.....	14.9	25.4	46.4	76.5	118.8	165.3	223.3	295.8	386.7	417.0
2001.....	11.7	25.5	45.6	76.7	112.5	162.1	221.5	308.1	400.5	450.5
2000.....	11.8	25.5	43.5	72.1	108.5	154.3	224.3	311.4	382.7	452.1
1999.....	13.2	27.5	44.1	78.2	113.3	163.8	216.1	282.6	374.1	506.2
Black, female										
2005 4/.....	9.5	16.2	31.3	51.7	91.0	141.5	193.7	267.5	369.1	523.3
2004 4/.....	9.4	14.9	34.4	56.8	91.3	132.2	206.5	290.3	381.4	543.0
2003 4/.....	9.3	19.2	36.4	63.6	95.4	145.3	213.6	296.3	391.1	556.2
2002.....	9.4	17.9	35.8	60.6	95.1	158.9	220.4	303.0	395.9	584.1
2001.....	9.8	17.0	35.6	63.9	98.1	154.3	230.3	303.4	372.8	559.1
2000.....	8.7	18.7	38.4	61.5	97.6	162.8	223.3	319.6	416.1	552.0
1999.....	9.7	18.1	37.4	68.3	110.3	156.4	237.6	302.4	406.6	537.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Nutritional deficiencies (E40-E64)										
All races 3/, both sexes										
2005.....	0.1	0.2	0.4	0.5	0.6	1.2	2.7	4.7	10.0	30.8
2004.....	0.1	0.2	0.4	0.4	0.7	1.2	2.1	4.6	10.0	30.3
2003.....	0.1	0.2	0.3	0.5	0.8	1.3	2.5	5.7	10.4	35.5
2002.....	0.1	0.2	0.3	0.5	0.8	1.6	2.7	6.1	12.8	41.5
2001.....	0.1	0.2	0.3	0.5	0.8	1.4	2.8	5.9	12.8	42.9
2000.....	0.1	0.2	0.3	0.5	1.0	2.0	3.0	6.8	15.8	50.0
1999.....	0.2	0.2	0.4	0.6	0.9	1.6	3.2	6.4	15.9	53.9
All races 3/, male										
2005.....	*	0.2	0.3	0.5	0.7	1.2	2.6	4.5	9.3	27.9
2004.....	*	*	0.4	0.5	0.8	1.3	2.2	4.9	9.6	27.4
2003.....	*	*	0.3	0.5	0.7	1.4	2.7	5.6	10.9	30.6
2002.....	*	0.2	0.3	0.6	0.9	1.6	3.0	6.3	12.0	37.0
2001.....	*	0.3	0.3	0.5	0.9	1.4	2.9	5.6	13.2	35.2
2000.....	*	0.3	0.2	0.7	1.0	2.0	3.4	6.8	14.4	43.7
1999.....	0.2	0.2	0.4	0.7	1.1	1.6	3.8	6.8	16.0	48.4
All races 3/, female										
2005.....	*	0.2	0.4	0.4	0.5	1.2	2.8	5.0	10.4	32.1
2004.....	*	0.2	0.4	0.4	0.7	1.1	2.0	4.3	10.3	31.7
2003.....	*	0.2	0.3	0.5	0.8	1.2	2.3	5.8	10.1	37.6
2002.....	*	*	0.4	0.4	0.7	1.6	2.4	6.0	13.3	43.5
2001.....	*	0.2	0.4	0.4	0.7	1.3	2.7	6.1	12.6	46.1
2000.....	*	0.2	0.4	0.4	0.9	2.0	2.7	6.8	16.6	52.6
1999.....	0.2	0.2	0.4	0.5	0.7	1.5	2.6	6.2	15.8	56.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Nutritional deficiencies (E40-E64)										
White, both sexes										
2005 4/.....	*	0.2	0.3	0.4	0.5	1.1	2.6	4.4	9.4	30.0
2004 4/.....	*	0.1	0.3	0.4	0.6	1.1	1.9	4.2	9.7	29.2
2003 4/.....	*	0.2	0.2	0.4	0.6	1.2	2.3	5.6	9.8	34.2
2002.....	0.1	0.2	0.3	0.4	0.7	1.4	2.5	5.9	12.3	40.2
2001.....	*	0.2	0.3	0.4	0.8	1.2	2.6	5.5	12.0	42.2
2000.....	*	0.2	0.2	0.5	0.8	1.9	2.7	6.4	15.3	48.8
1999.....	0.2	0.2	0.3	0.4	0.8	1.4	2.7	5.9	14.8	52.8
White, male										
2005 4/.....	*	*	0.3	0.5	0.6	1.1	2.5	4.1	8.6	27.2
2004 4/.....	*	*	0.3	0.5	0.6	1.0	2.1	4.5	9.1	26.3
2003 4/.....	*	*	*	0.4	0.6	1.3	2.3	5.2	10.1	29.6
2002.....	*	*	0.3	0.4	0.7	1.4	2.8	6.1	11.0	36.3
2001.....	*	*	*	0.4	0.7	1.1	2.6	5.3	12.2	33.8
2000.....	*	0.2	*	0.5	0.8	1.8	2.9	6.2	13.9	42.8
1999.....	*	*	0.3	0.4	1.0	1.3	3.0	6.1	14.7	46.5
White, female										
2005 4/.....	*	*	0.4	0.4	0.5	1.0	2.6	4.6	9.9	31.4
2004 4/.....	*	*	0.4	0.4	0.6	1.1	1.8	4.1	10.1	30.4
2003 4/.....	*	*	*	0.5	0.7	1.1	2.3	5.9	9.7	36.3
2002.....	*	*	0.4	0.4	0.7	1.3	2.2	5.7	13.1	41.9
2001.....	*	0.2	0.3	0.4	0.8	1.3	2.7	5.7	11.8	45.7
2000.....	*	*	0.3	0.4	0.9	2.0	2.5	6.5	16.2	51.3
1999.....	*	*	0.3	0.4	0.6	1.4	2.4	5.9	14.9	55.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Nutritional deficiencies (E40-E64)

Black, both sexes

2005 4/.....	*	*	*	*	1.6	2.7	4.4	8.7	18.0	42.5
2004 4/.....	*	*	0.9	*	1.7	2.7	4.2	8.1	14.1	46.8
2003 4/.....	*	*	1.0	*	1.9	*	4.6	9.0	19.5	55.1
2002.....	*	*	*	1.5	2.0	3.5	5.1	10.5	20.7	60.6
2001.....	*	*	1.0	*	*	3.0	4.4	10.8	23.8	54.7
2000.....	*	*	*	*	2.1	3.2	6.5	12.1	22.6	66.9
1999.....	*	*	*	2.0	2.0	3.2	8.3	12.3	30.9	70.8

Black, male

2005 4/.....	*	*	*	*	*	*	*	*	19.0	37.2
2004 4/.....	*	*	*	*	*	*	*	10.5	15.1	45.7
2003 4/.....	*	*	*	*	*	*	7.2	13.7	24.4	48.0
2002.....	*	*	*	*	*	*	*	9.6	27.5	47.2
2001.....	*	*	*	*	*	*	6.7	11.6	24.9	57.9
2000.....	*	*	*	*	*	*	9.7	15.1	22.8	63.4
1999.....	*	*	*	*	*	*	12.9	16.1	34.9	75.0

Black, female

2005 4/.....	*	*	*	*	*	*	5.0	9.1	17.4	44.6
2004 4/.....	*	*	*	*	*	*	*	6.7	13.6	47.3
2003 4/.....	*	*	*	*	*	*	*	6.2	17.0	57.9
2002.....	*	*	*	*	*	*	4.7	11.0	17.1	65.7
2001.....	*	*	*	*	*	*	*	10.3	23.3	53.5
2000.....	*	*	*	*	*	*	4.4	10.3	22.5	68.3
1999.....	*	*	*	*	*	*	5.2	10.0	29.0	69.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malnutrition (E40-E46)										
All races 3/, both sexes										
2005.....	0.1	0.2	0.3	0.4	0.6	1.2	2.5	4.6	9.3	28.9
2004.....	0.1	0.2	0.4	0.4	0.7	1.2	2.0	4.4	9.6	28.3
2003.....	0.1	0.2	0.3	0.5	0.7	1.2	2.4	5.5	10.0	33.4
2002.....	0.1	0.2	0.3	0.5	0.7	1.4	2.5	5.6	12.2	38.4
2001.....	0.1	0.2	0.3	0.4	0.8	1.3	2.6	5.5	12.0	40.2
2000.....	0.1	0.2	0.3	0.5	0.9	1.9	2.8	6.4	14.6	46.3
1999.....	0.2	0.2	0.3	0.5	0.8	1.5	3.0	6.0	14.7	50.0
All races 3/, male										
2005.....	*	0.2	0.3	0.5	0.6	1.2	2.4	4.4	8.8	26.3
2004.....	*	*	0.3	0.5	0.7	1.3	2.2	4.8	9.2	25.3
2003.....	*	*	0.3	0.5	0.7	1.3	2.5	5.4	10.4	28.2
2002.....	*	*	0.3	0.5	0.8	1.4	2.8	5.7	11.4	35.1
2001.....	*	0.2	0.3	0.5	0.8	1.3	2.6	5.1	12.6	33.6
2000.....	*	0.2	*	0.6	0.9	1.8	3.3	6.4	13.1	40.7
1999.....	*	0.2	0.3	0.6	1.1	1.5	3.6	6.2	14.8	44.4
All races 3/, female										
2005.....	*	0.2	0.4	0.3	0.5	1.2	2.6	4.7	9.7	30.2
2004.....	*	0.2	0.4	0.3	0.7	1.1	1.9	4.2	9.9	29.6
2003.....	*	*	0.3	0.5	0.8	1.1	2.2	5.7	9.7	35.6
2002.....	*	*	0.3	0.4	0.6	1.4	2.2	5.6	12.6	39.8
2001.....	*	0.2	0.3	0.4	0.7	1.3	2.5	5.7	11.6	43.0
2000.....	*	0.2	0.3	0.3	0.9	1.9	2.4	6.5	15.5	48.6
1999.....	0.2	0.2	0.3	0.4	0.6	1.4	2.5	5.8	14.7	52.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malnutrition (E40-E46)										
White, both sexes										
2005 4/.....	*	0.2	0.3	0.4	0.5	1.0	2.4	4.2	8.8	28.2
2004 4/.....	*	0.1	0.3	0.4	0.6	1.0	1.9	4.1	9.3	26.9
2003 4/.....	*	0.1	0.2	0.4	0.6	1.1	2.2	5.4	9.3	32.1
2002.....	0.1	0.2	0.3	0.4	0.6	1.3	2.4	5.4	11.7	37.0
2001.....	*	0.2	0.2	0.4	0.7	1.1	2.4	5.1	11.2	39.5
2000.....	*	0.2	0.2	0.4	0.8	1.8	2.4	6.1	14.2	45.2
1999.....	0.2	0.2	0.3	0.4	0.7	1.3	2.5	5.5	13.8	48.9
White, male										
2005 4/.....	*	*	0.3	0.5	0.5	1.0	2.3	4.0	8.1	25.7
2004 4/.....	*	*	0.3	0.4	0.6	1.0	2.0	4.3	8.7	24.1
2003 4/.....	*	*	*	0.4	0.5	1.2	2.2	4.9	9.5	27.2
2002.....	*	*	*	0.4	0.6	1.3	2.7	5.5	10.5	34.3
2001.....	*	*	*	0.4	0.7	1.0	2.3	4.7	11.7	32.2
2000.....	*	*	*	0.5	0.7	1.6	2.8	5.9	12.5	39.7
1999.....	*	*	0.3	0.4	0.9	1.2	2.8	5.6	13.4	42.2
White, female										
2005 4/.....	*	*	0.3	0.4	0.5	1.0	2.4	4.4	9.2	29.4
2004 4/.....	*	*	0.3	0.4	0.6	1.1	1.7	3.9	9.7	28.2
2003 4/.....	*	*	*	0.5	0.7	1.0	2.2	5.7	9.3	34.2
2002.....	*	*	0.3	0.4	0.6	1.2	2.1	5.3	12.4	38.2
2001.....	*	*	0.3	0.4	0.8	1.2	2.5	5.4	10.9	42.6
2000.....	*	*	0.3	*	0.9	1.9	2.1	6.2	15.1	47.4
1999.....	*	*	0.3	0.4	0.5	1.3	2.3	5.5	14.0	51.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malnutrition (E40-E46)										
Black, both sexes										
2005 4/.....	*	*	*	*	*	2.7	4.2	8.7	16.7	40.4
2004 4/.....	*	*	0.9	*	*	2.5	4.2	8.1	13.6	46.0
2003 4/.....	*	*	1.0	*	1.8	*	4.4	8.8	19.3	53.4
2002.....	*	*	*	1.5	1.9	3.0	4.4	10.1	19.9	57.6
2001.....	*	*	*	*	*	2.8	4.0	10.1	22.2	51.6
2000.....	*	*	*	*	1.9	3.1	6.5	11.2	20.9	63.8
1999.....	*	*	*	1.6	2.0	3.2	8.0	11.1	28.9	66.0
Black, male										
2005 4/.....	*	*	*	*	*	*	*	*	17.6	34.4
2004 4/.....	*	*	*	*	*	*	*	10.5	14.3	43.7
2003 4/.....	*	*	*	*	*	*	6.9	13.2	23.6	45.9
2002.....	*	*	*	*	*	*	*	9.2	25.2	46.1
2001.....	*	*	*	*	*	*	*	11.2	24.1	55.6
2000.....	*	*	*	*	*	*	9.7	14.2	22.8	61.1
1999.....	*	*	*	*	*	*	12.2	13.3	33.1	73.8
Black, female										
2005 4/.....	*	*	*	*	*	*	4.8	9.1	16.3	42.8
2004 4/.....	*	*	*	*	*	*	*	6.7	13.2	46.9
2003 4/.....	*	*	*	*	*	*	*	6.2	17.0	56.3
2002.....	*	*	*	*	*	*	*	10.7	17.1	62.0
2001.....	*	*	*	*	*	*	*	9.4	21.3	50.1
2000.....	*	*	*	*	*	*	4.4	9.4	20.0	64.8
1999.....	*	*	*	*	*	*	5.2	9.7	26.8	63.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other nutritional deficiencies (E50-E64)

All races 3/, both sexes

2005.....	*	*	*	*	*	*	*	*	0.7	1.8
2004.....	*	*	*	*	*	*	*	*	0.4	2.1
2003.....	*	*	*	*	*	*	*	*	0.5	2.1
2002.....	*	*	*	*	*	*	*	0.4	0.6	3.1
2001.....	*	*	*	*	*	*	*	0.4	0.8	2.7
2000.....	*	*	*	*	*	*	0.3	0.4	1.2	3.7
1999.....	*	*	*	*	*	*	*	0.4	1.1	3.9

All races 3/, male

2005.....	*	*	*	*	*	*	*	*	*	1.6
2004.....	*	*	*	*	*	*	*	*	*	2.1
2003.....	*	*	*	*	*	*	*	*	*	2.4
2002.....	*	*	*	*	*	*	*	*	*	1.9
2001.....	*	*	*	*	*	*	*	*	*	1.6
2000.....	*	*	*	*	*	*	*	*	1.3	3.0
1999.....	*	*	*	*	*	*	*	*	1.3	3.9

All races 3/, female

2005.....	*	*	*	*	*	*	*	*	0.7	1.9
2004.....	*	*	*	*	*	*	*	*	*	2.1
2003.....	*	*	*	*	*	*	*	*	*	2.0
2002.....	*	*	*	*	*	*	*	*	0.6	3.6
2001.....	*	*	*	*	*	*	*	*	1.0	3.1
2000.....	*	*	*	*	*	*	*	*	1.1	4.0
1999.....	*	*	*	*	*	*	*	*	1.1	3.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other nutritional deficiencies (E50-E64)

White, both sexes

2005 4/.....	*	*	*	*	*	*	*	*	0.6	1.8
2004 4/.....	*	*	*	*	*	*	*	*	0.4	2.2
2003 4/.....	*	*	*	*	*	*	*	*	0.5	2.2
2002.....	*	*	*	*	*	*	*	0.5	0.6	3.1
2001.....	*	*	*	*	*	*	*	0.4	0.8	2.7
2000.....	*	*	*	*	*	*	0.3	0.3	1.2	3.7
1999.....	*	*	*	*	*	*	*	0.4	1.1	3.9

White, male

2005 4/.....	*	*	*	*	*	*	*	*	*	1.5
2004 4/.....	*	*	*	*	*	*	*	*	*	2.2
2003 4/.....	*	*	*	*	*	*	*	*	*	2.4
2002.....	*	*	*	*	*	*	*	*	*	2.0
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	1.4	3.2
1999.....	*	*	*	*	*	*	*	*	1.3	4.3

White, female

2005 4/.....	*	*	*	*	*	*	*	*	0.7	1.9
2004 4/.....	*	*	*	*	*	*	*	*	*	2.2
2003 4/.....	*	*	*	*	*	*	*	*	*	2.1
2002.....	*	*	*	*	*	*	*	*	0.7	3.6
2001.....	*	*	*	*	*	*	*	*	0.9	3.1
2000.....	*	*	*	*	*	*	*	*	1.0	3.9
1999.....	*	*	*	*	*	*	*	*	1.0	3.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Meningitis (G00,G03)										
All races 3/, both sexes										
2005.....	0.2	0.3	0.3	0.4	0.3	0.4	0.4	0.5	0.6	0.9
2004.....	0.2	0.3	0.3	0.3	0.3	0.5	0.5	0.6	0.7	0.9
2003.....	0.2	0.3	0.3	0.3	0.4	0.6	0.5	0.7	0.6	1.1
2002.....	0.2	0.3	0.3	0.3	0.3	0.5	0.7	0.6	0.7	1.2
2001.....	0.2	0.3	0.3	0.4	0.5	0.5	0.5	0.6	1.0	0.9
2000.....	0.2	0.2	0.3	0.3	0.4	0.5	0.7	0.9	1.1	1.2
1999.....	0.2	0.3	0.3	0.3	0.5	0.5	0.7	0.9	1.2	1.5
All races 3/, male										
2005.....	0.2	0.3	0.3	0.4	0.3	*	*	*	*	*
2004.....	0.3	0.3	0.4	0.3	0.3	*	*	*	*	*
2003.....	*	0.3	0.4	0.4	0.5	0.6	*	0.8	*	*
2002.....	0.2	0.4	0.3	0.4	*	0.6	0.9	0.7	1.0	*
2001.....	*	0.3	0.3	0.4	0.6	0.5	0.7	0.9	1.1	*
2000.....	0.2	0.3	0.4	0.4	0.5	0.6	0.9	1.3	1.1	*
1999.....	0.2	0.4	0.4	0.4	0.4	0.6	0.8	0.9	1.3	1.7
All races 3/, female										
2005.....	0.2	0.2	0.3	0.4	0.3	*	*	0.5	0.6	0.9
2004.....	*	*	0.3	0.3	0.3	0.6	0.6	0.6	0.6	0.7
2003.....	*	0.2	0.2	0.3	0.3	0.6	0.4	0.7	0.6	1.0
2002.....	*	0.2	0.2	*	0.3	0.4	0.6	0.6	*	1.1
2001.....	*	0.2	0.2	0.4	0.4	0.4	0.4	0.5	0.9	0.8
2000.....	*	0.2	0.3	*	*	0.4	0.5	0.6	1.1	1.0
1999.....	*	0.2	*	*	0.5	0.4	0.7	0.9	1.1	1.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Meningitis (G00,G03)										
White, both sexes										
2005 4/.....	0.2	0.3	0.2	0.4	0.3	0.3	0.4	0.5	0.6	0.9
2004 4/.....	0.2	0.2	0.2	0.2	0.3	0.4	0.4	0.6	0.6	0.8
2003 4/.....	0.1	0.2	0.2	0.3	0.3	0.5	0.4	0.7	0.6	1.0
2002.....	0.1	0.2	0.2	0.3	0.3	0.5	0.7	0.6	0.6	1.2
2001.....	0.1	0.2	0.2	0.3	0.5	0.4	0.6	0.6	0.9	0.8
2000.....	0.1	0.2	0.2	0.3	0.4	0.4	0.6	0.8	1.1	1.0
1999.....	0.1	0.2	0.2	0.3	0.4	0.5	0.6	0.7	1.1	1.3
White, male										
2005 4/.....	*	0.3	0.3	0.4	*	*	*	*	*	*
2004 4/.....	0.2	0.3	0.3	*	*	*	*	*	*	*
2003 4/.....	*	0.3	0.3	0.3	0.4	0.6	*	0.9	*	*
2002.....	*	0.3	*	0.3	*	0.6	0.8	*	*	*
2001.....	*	0.2	0.3	0.4	0.5	*	0.7	0.8	*	*
2000.....	*	0.3	0.3	0.4	0.5	*	0.8	1.2	*	*
1999.....	*	0.3	*	0.4	*	0.5	0.7	*	1.3	*
White, female										
2005 4/.....	*	*	*	0.3	*	*	*	*	*	0.9
2004 4/.....	*	*	*	*	*	0.5	0.5	0.6	*	*
2003 4/.....	*	*	*	*	*	0.5	*	0.6	*	0.9
2002.....	*	*	*	*	*	*	0.5	0.5	*	1.2
2001.....	*	*	*	*	0.4	*	*	*	0.8	0.8
2000.....	*	*	*	*	*	*	*	0.6	1.0	1.0
1999.....	*	*	*	*	0.5	*	0.6	0.8	1.0	1.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Parkinson's disease (G20-G21)										
All races 3/, both sexes										
2005.....	*	0.1	0.2	0.8	2.3	6.8	20.3	50.2	98.7	143.7
2004.....	*	*	0.2	0.6	2.0	6.4	18.5	47.6	94.0	135.8
2003.....	*	*	0.2	0.7	2.0	6.4	19.9	49.3	93.3	138.2
2002.....	*	*	0.2	0.6	1.9	6.5	18.4	45.4	89.8	135.2
2001.....	*	*	0.2	0.7	1.8	5.8	18.1	46.4	90.9	134.2
2000.....	*	*	0.2	0.6	1.7	6.6	16.7	44.8	87.6	131.9
1999.....	*	*	0.2	0.6	1.6	5.6	16.7	41.8	82.8	124.4
All races 3/, male										
2005.....	*	*	0.3	0.9	3.2	9.6	30.0	75.0	153.6	228.6
2004.....	*	*	0.4	0.9	2.6	9.1	27.3	70.6	147.1	220.0
2003.....	*	*	0.2	0.8	2.7	8.9	29.2	74.2	140.7	222.4
2002.....	*	*	0.3	0.8	2.5	9.8	27.3	67.2	136.6	220.1
2001.....	*	*	0.3	1.0	2.7	8.2	26.5	68.8	141.7	224.4
2000.....	*	*	*	0.8	2.2	9.0	24.1	67.4	135.0	222.3
1999.....	*	*	0.3	0.7	2.3	7.7	24.8	64.4	131.3	205.6
All races 3/, female										
2005.....	*	*	*	0.6	1.5	4.3	12.5	32.3	64.6	104.7
2004.....	*	*	*	0.4	1.4	4.1	11.4	31.0	61.3	97.9
2003.....	*	*	0.2	0.6	1.4	4.3	12.4	31.5	64.3	101.0
2002.....	*	*	*	0.5	1.4	3.8	11.3	30.0	61.3	98.4
2001.....	*	*	*	0.4	1.0	3.7	11.5	30.7	60.4	96.4
2000.....	*	*	*	0.4	1.3	4.4	10.9	29.1	59.5	95.0
1999.....	*	*	*	0.4	0.9	3.9	10.4	26.0	54.4	91.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Parkinson's disease (G20-G21)										
White, both sexes										
2005 4/.....	*	0.1	0.2	0.8	2.4	7.1	21.8	53.7	104.4	152.9
2004 4/.....	*	*	0.2	0.7	2.1	6.8	20.2	50.6	99.9	143.1
2003 4/.....	*	*	0.2	0.7	2.0	6.8	21.2	52.4	99.0	146.0
2002.....	*	*	0.2	0.6	2.1	6.9	19.6	48.3	95.7	143.3
2001.....	*	*	0.2	0.7	1.9	6.2	19.4	49.6	96.4	141.8
2000.....	*	*	0.2	0.6	1.9	7.1	17.8	47.5	92.5	139.5
1999.....	*	*	0.2	0.6	1.6	6.0	17.7	44.1	87.4	131.2
White, male										
2005 4/.....	*	*	0.3	1.0	3.3	10.0	31.9	79.5	162.2	242.7
2004 4/.....	*	*	0.4	1.0	2.7	9.6	29.4	74.7	156.1	231.3
2003 4/.....	*	*	*	0.8	2.7	9.4	30.9	78.4	148.9	233.3
2002.....	*	*	0.3	0.8	2.7	10.2	28.8	71.2	145.0	232.7
2001.....	*	*	0.3	1.0	3.0	8.8	28.1	73.4	149.6	237.6
2000.....	*	*	*	0.9	2.5	9.6	25.5	70.7	141.5	235.3
1999.....	*	*	0.3	0.7	2.4	8.1	26.4	67.7	138.6	216.7
White, female										
2005 4/.....	*	*	*	0.7	1.6	4.6	13.4	34.6	68.2	111.5
2004 4/.....	*	*	*	0.4	1.5	4.3	12.6	33.0	65.0	103.3
2003 4/.....	*	*	*	0.6	1.4	4.5	13.3	33.6	68.2	107.4
2002.....	*	*	*	0.5	1.5	3.9	12.0	31.9	65.5	104.4
2001.....	*	*	*	0.5	0.9	4.0	12.3	32.7	64.2	101.7
2000.....	*	*	*	0.4	1.4	4.8	11.7	31.2	63.3	100.6
1999.....	*	*	*	0.4	0.9	4.2	10.8	27.6	57.3	96.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Parkinson's disease (G20-G21)

Black, both sexes

2005 4/.....	*	*	*	*	*	4.3	10.9	22.7	50.0	56.7
2004 4/.....	*	*	*	*	*	3.9	7.2	22.3	39.1	61.2
2003 4/.....	*	*	*	*	1.9	4.1	10.7	22.8	38.3	52.8
2002.....	*	*	*	*	*	4.6	9.5	20.6	32.7	51.7
2001.....	*	*	*	*	*	2.7	8.9	18.9	34.9	59.0
2000.....	*	*	*	*	*	3.1	8.8	21.0	37.6	51.6
1999.....	*	*	*	*	*	3.6	8.7	19.5	32.4	52.0

Black, male

2005 4/.....	*	*	*	*	*	7.4	18.7	36.2	82.2	97.6
2004 4/.....	*	*	*	*	*	5.3	12.6	34.1	58.1	106.3
2003 4/.....	*	*	*	*	*	5.4	16.8	37.0	59.1	97.0
2002.....	*	*	*	*	*	7.5	16.0	30.3	55.0	89.0
2001.....	*	*	*	*	*	*	14.8	25.6	57.8	99.8
2000.....	*	*	*	*	*	*	13.7	37.8	65.8	86.5
1999.....	*	*	*	*	*	5.7	12.6	31.8	48.8	86.7

Black, female

2005 4/.....	*	*	*	*	*	*	5.6	14.7	33.0	40.1
2004 4/.....	*	*	*	*	*	*	*	15.4	29.1	43.3
2003 4/.....	*	*	*	*	*	*	6.6	14.4	27.5	35.6
2002.....	*	*	*	*	*	*	5.2	14.8	21.1	37.4
2001.....	*	*	*	*	*	*	4.9	15.0	23.3	43.8
2000.....	*	*	*	*	*	*	5.5	10.8	23.4	38.6
1999.....	*	*	*	*	*	*	6.1	12.0	24.2	39.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Alzheimer's disease (G30)										
All races 3/, both sexes										
2005.....	*	0.1	0.3	1.1	3.5	9.7	33.3	102.5	275.5	861.6
2004.....	*	0.2	0.3	0.9	3.1	10.1	31.0	99.4	261.0	818.8
2003.....	*	0.1	0.3	1.0	3.3	10.0	33.3	98.6	255.0	802.4
2002.....	*	*	0.2	0.9	3.2	9.8	30.6	95.4	245.7	752.3
2001.....	*	*	0.3	1.2	3.3	9.0	29.2	88.5	232.5	710.3
2000.....	*	*	0.3	1.1	3.2	9.1	28.9	85.2	221.1	667.7
1999.....	*	0.1	0.2	1.0	3.0	8.7	26.7	78.6	206.2	601.3
All races 3/, male										
2005.....	*	*	0.2	1.0	3.2	9.6	31.4	95.9	246.9	639.9
2004.....	*	*	0.3	0.8	3.1	9.4	29.6	89.3	236.4	619.6
2003.....	*	*	0.3	1.0	2.9	10.1	32.1	92.1	231.2	606.5
2002.....	*	*	0.2	0.8	3.1	9.2	29.5	90.9	220.6	572.0
2001.....	*	*	0.3	0.9	3.2	9.0	27.7	85.6	216.6	556.1
2000.....	*	*	0.3	0.9	2.9	9.5	29.1	79.9	206.8	522.1
1999.....	*	*	*	1.1	3.0	8.8	26.9	77.4	198.2	485.7
All races 3/, female										
2005.....	*	*	0.4	1.2	3.7	9.8	34.8	107.3	293.2	963.4
2004.....	*	0.2	0.4	1.0	3.1	10.7	32.1	106.7	276.2	908.4
2003.....	*	*	0.3	0.9	3.7	10.0	34.2	103.2	269.5	889.0
2002.....	*	*	*	1.0	3.3	10.3	31.5	98.5	260.9	830.5
2001.....	*	*	0.3	1.4	3.4	9.0	30.4	90.5	242.1	774.9
2000.....	*	*	0.3	1.2	3.5	8.7	28.8	88.9	229.5	727.0
1999.....	*	*	0.3	0.8	3.0	8.6	26.6	79.4	210.9	648.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Alzheimer's disease (G30)										
White, both sexes										
2005 4/.....	*	*	0.3	1.2	3.7	10.0	34.1	105.3	284.3	891.6
2004 4/.....	*	0.2	0.4	1.0	3.3	10.4	32.0	102.3	268.5	846.8
2003 4/.....	*	*	0.3	1.1	3.6	10.3	34.5	101.8	264.4	834.0
2002.....	*	*	0.2	1.0	3.4	10.3	31.9	99.1	255.3	781.8
2001.....	*	*	0.3	1.3	3.6	9.3	30.6	90.9	240.5	738.1
2000.....	*	*	0.3	1.1	3.4	9.6	29.9	88.1	229.1	695.3
1999.....	*	*	0.2	1.0	3.3	9.1	27.6	81.3	213.6	627.9
White, male										
2005 4/.....	*	*	0.2	1.1	3.5	9.8	31.9	97.5	254.3	661.6
2004 4/.....	*	*	0.4	0.9	3.3	9.7	30.2	91.6	243.9	640.9
2003 4/.....	*	*	0.3	1.1	3.0	10.3	33.1	94.9	238.5	630.3
2002.....	*	*	*	0.8	3.3	9.4	30.5	94.3	227.6	594.8
2001.....	*	*	0.3	0.9	3.4	9.2	28.8	88.1	222.7	577.2
2000.....	*	*	0.3	1.0	3.1	10.1	29.9	82.6	214.4	542.1
1999.....	*	*	*	1.1	3.3	9.2	27.6	80.1	205.6	506.9
White, female										
2005 4/.....	*	*	0.4	1.4	3.9	10.2	35.8	111.1	303.1	997.5
2004 4/.....	*	0.3	0.4	1.1	3.3	11.1	33.5	110.2	283.8	939.6
2003 4/.....	*	*	0.4	1.0	4.2	10.3	35.7	106.8	280.3	924.3
2002.....	*	*	*	1.2	3.5	11.0	33.1	102.6	272.2	863.0
2001.....	*	*	0.4	1.6	3.8	9.4	32.1	92.9	251.3	805.4
2000.....	*	*	0.3	1.3	3.7	9.2	29.9	92.0	237.7	757.6
1999.....	*	*	*	0.9	3.3	9.1	27.5	82.1	218.3	676.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Alzheimer's disease (G30)										
Black, both sexes										
2005 4/.....	*	*	*	*	2.8	10.3	37.0	100.4	241.5	689.5
2004 4/.....	*	*	*	*	2.2	10.3	30.6	96.0	236.7	657.9
2003 4/.....	*	*	*	*	1.8	10.5	30.2	91.4	207.9	583.8
2002.....	*	*	*	*	2.5	8.5	27.2	81.3	186.8	543.7
2001.....	*	*	*	*	1.9	8.7	23.3	86.8	183.6	506.7
2000.....	*	*	*	*	2.1	7.4	25.9	71.1	171.7	445.1
1999.....	*	*	*	*	*	6.9	25.3	65.0	152.0	372.4
Black, male										
2005 4/.....	*	*	*	*	*	11.6	36.1	103.7	222.6	518.7
2004 4/.....	*	*	*	*	*	9.4	32.0	89.6	210.9	497.7
2003 4/.....	*	*	*	*	*	11.1	32.5	93.1	206.0	459.8
2002.....	*	*	*	*	*	9.5	29.4	82.2	189.5	412.8
2001.....	*	*	*	*	*	9.9	24.2	84.6	185.4	414.1
2000.....	*	*	*	*	*	7.8	27.7	64.7	156.0	371.3
1999.....	*	*	*	*	*	8.0	27.5	62.7	139.5	317.5
Black, female										
2005 4/.....	*	*	*	*	3.5	9.3	37.7	98.4	251.5	758.5
2004 4/.....	*	*	*	*	*	11.0	29.7	99.7	250.2	721.4
2003 4/.....	*	*	*	*	*	10.1	28.6	90.3	208.9	631.9
2002.....	*	*	*	*	*	7.8	25.8	80.7	185.4	593.9
2001.....	*	*	*	*	*	7.9	22.6	88.2	182.7	541.0
2000.....	*	*	*	*	*	7.1	24.6	74.9	179.6	472.6
1999.....	*	*	*	*	*	6.0	23.9	66.4	158.2	392.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Major cardiovascular diseases (I00-I78)

All races 3/, both sexes

2005.....	47.2	84.2	140.0	208.2	336.6	515.2	848.2	1,452.7	2,635.2	6,398.5
2004.....	47.8	86.0	139.2	211.1	343.0	537.5	885.1	1,514.3	2,730.4	6,616.6
2003.....	50.3	87.7	142.8	224.5	363.7	574.0	946.1	1,627.3	2,918.8	7,138.0
2002.....	49.7	87.9	144.2	233.5	374.0	602.6	993.2	1,700.2	3,034.4	7,389.6
2001.....	48.8	85.6	144.9	234.9	385.7	619.6	1,017.5	1,759.4	3,133.9	7,673.7
2000.....	49.6	88.5	146.7	248.4	407.5	647.9	1,066.0	1,824.4	3,253.7	8,030.3
1999.....	50.0	88.3	148.7	254.8	416.0	678.9	1,111.3	1,888.7	3,356.4	8,197.1

All races 3/, male

2005.....	64.1	116.8	199.9	294.8	462.0	679.9	1,093.3	1,820.2	3,159.0	6,615.9
2004.....	64.6	120.2	198.5	296.2	472.0	714.7	1,134.9	1,900.8	3,283.3	6,818.1
2003.....	67.7	123.1	204.5	314.7	497.9	765.9	1,225.3	2,051.8	3,517.1	7,329.3
2002.....	67.9	123.9	206.1	326.7	512.6	802.0	1,287.4	2,140.7	3,633.4	7,638.8
2001.....	66.5	121.4	204.7	327.6	526.3	821.4	1,325.6	2,223.2	3,774.2	7,936.0
2000.....	68.0	126.8	209.0	350.6	556.9	857.2	1,403.8	2,305.7	3,956.7	8,448.1
1999.....	68.9	125.7	212.5	358.4	568.4	908.7	1,461.9	2,392.7	4,078.9	8,603.2

All races 3/, female

2005.....	30.5	52.4	82.5	126.6	222.2	371.4	649.7	1,185.8	2,309.9	6,298.7
2004.....	31.2	52.6	82.4	130.8	225.5	383.3	684.0	1,236.7	2,389.8	6,526.0
2003.....	33.2	53.3	83.7	139.7	241.8	407.6	722.2	1,325.1	2,552.9	7,053.5
2002.....	31.6	52.9	85.0	146.0	248.2	430.4	758.0	1,389.0	2,670.1	7,281.5
2001.....	31.5	50.7	87.6	148.1	258.3	445.9	772.7	1,434.7	2,750.2	7,563.9
2000.....	31.5	51.4	87.0	152.8	272.0	468.5	799.9	1,489.2	2,839.0	7,860.1
1999.....	31.4	52.1	87.5	158.0	278.2	482.2	837.3	1,538.3	2,933.7	8,033.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Major cardiovascular diseases (I00-I78)

White, both sexes

2005 4/.....	41.4	73.7	122.4	187.2	309.5	483.0	814.8	1,420.7	2,623.8	6,507.8
2004 4/.....	41.9	74.9	122.1	189.5	316.2	503.9	851.4	1,479.8	2,721.9	6,704.8
2003 4/.....	43.4	76.1	124.8	203.3	333.1	537.6	907.0	1,593.5	2,909.9	7,233.1
2002.....	43.6	76.6	127.4	211.8	344.7	567.9	957.1	1,663.0	3,023.3	7,482.7
2001.....	41.6	74.2	127.7	212.4	355.6	580.3	980.2	1,721.1	3,122.8	7,770.6
2000.....	42.6	76.5	130.6	225.6	376.3	612.7	1,027.2	1,783.3	3,243.6	8,133.9
1999.....	42.4	76.6	132.4	230.2	385.3	644.7	1,068.3	1,848.2	3,345.3	8,299.2

White, male

2005 4/.....	58.0	105.2	179.2	268.7	430.2	643.2	1,057.0	1,793.2	3,164.9	6,772.6
2004 4/.....	59.0	107.8	179.1	270.3	439.8	677.7	1,101.5	1,869.3	3,298.4	6,940.8
2003 4/.....	60.6	110.4	183.6	289.9	463.8	727.3	1,183.5	2,022.8	3,520.9	7,466.8
2002.....	62.0	111.9	186.7	301.4	479.1	765.2	1,253.1	2,112.9	3,642.0	7,766.6
2001.....	59.1	109.0	185.6	301.7	493.8	781.8	1,286.2	2,192.9	3,777.7	8,060.7
2000.....	60.7	114.5	190.5	323.5	523.7	822.7	1,366.4	2,273.2	3,963.6	8,585.0
1999.....	60.8	113.0	193.5	329.9	533.4	874.1	1,420.2	2,361.9	4,089.2	8,717.5

White, female

2005 4/.....	24.6	42.4	66.8	108.8	197.3	340.5	614.5	1,145.0	2,284.8	6,385.8
2004 4/.....	24.6	42.1	66.4	111.9	201.6	349.8	645.9	1,194.8	2,363.8	6,598.4
2003 4/.....	26.2	42.1	67.2	120.1	212.2	370.0	680.7	1,283.1	2,533.3	7,129.6
2002.....	25.2	41.6	69.3	126.1	220.5	394.1	715.8	1,341.0	2,644.0	7,359.3
2001.....	24.0	39.8	71.0	127.1	228.1	403.3	732.3	1,386.9	2,726.9	7,649.2
2000.....	24.4	39.0	72.0	132.4	240.4	429.2	755.0	1,438.7	2,815.1	7,950.6
1999.....	24.0	40.6	72.8	135.3	248.8	444.5	788.7	1,487.4	2,906.2	8,131.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Major cardiovascular diseases (I00-I78)

Black, both sexes

2005 4/.....	93.7	172.2	295.3	422.3	640.6	882.2	1,301.4	2,041.3	3,168.4	6,016.9
2004 4/.....	97.5	179.4	293.5	434.4	650.7	923.3	1,374.7	2,132.1	3,217.2	6,368.6
2003 4/.....	104.2	183.8	305.8	451.7	695.7	983.6	1,489.0	2,249.7	3,405.2	6,862.0
2002.....	99.1	184.1	302.3	468.4	697.7	1,001.3	1,510.2	2,343.9	3,555.3	7,081.7
2001.....	105.7	181.9	307.2	471.1	713.7	1,057.5	1,548.8	2,422.8	3,623.2	7,316.7
2000.....	104.9	192.0	303.5	489.3	736.9	1,061.9	1,623.2	2,496.2	3,719.8	7,518.9
1999.....	109.8	188.5	311.0	509.4	743.3	1,093.2	1,704.6	2,553.3	3,787.7	7,582.5

Black, male

2005 4/.....	117.6	224.6	402.0	585.8	873.4	1,156.7	1,667.3	2,508.7	3,664.5	5,707.6
2004 4/.....	118.5	234.5	393.3	594.3	890.6	1,210.4	1,729.0	2,625.1	3,677.3	6,198.0
2003 4/.....	129.8	239.7	412.9	610.0	924.3	1,273.7	1,900.2	2,789.8	4,035.4	6,547.7
2002.....	122.7	237.2	406.5	633.2	934.4	1,292.7	1,908.8	2,803.5	4,093.1	6,929.0
2001.....	129.8	237.2	405.7	627.8	934.4	1,342.7	1,988.0	2,943.9	4,284.2	7,363.9
2000.....	131.7	247.2	407.0	670.8	955.1	1,328.0	2,040.7	2,999.5	4,397.8	7,687.4
1999.....	138.3	246.5	423.5	686.4	993.1	1,400.0	2,146.5	3,046.5	4,407.7	7,921.1

Black, female

2005 4/.....	72.7	126.5	205.1	287.3	459.1	678.7	1,051.0	1,762.4	2,906.3	6,141.8
2004 4/.....	79.0	131.5	209.1	302.8	463.7	710.0	1,134.3	1,839.9	2,974.8	6,436.3
2003 4/.....	81.7	135.2	215.1	321.5	517.3	768.3	1,211.6	1,928.7	3,075.0	6,984.1
2002.....	78.2	138.0	213.9	333.0	512.5	785.0	1,242.7	2,069.3	3,275.3	7,140.2
2001.....	84.3	134.0	223.4	342.6	540.2	846.9	1,256.2	2,111.8	3,286.5	7,299.2
2000.....	81.2	144.0	215.2	340.5	564.6	865.5	1,346.0	2,192.8	3,378.3	7,456.2
1999.....	84.7	138.5	215.6	364.7	547.4	865.8	1,411.4	2,256.2	3,480.4	7,456.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Diseases of heart (I00-I09,I11,I13,I20-I51)										
All races 3/, both sexes										
2005.....	37.7	68.0	114.1	171.0	273.2	405.5	654.0	1,090.7	1,946.8	4,778.4
2004.....	38.3	69.6	113.7	173.4	278.1	423.7	679.6	1,130.0	2,007.8	4,895.9
2003.....	40.5	71.3	116.6	185.0	296.1	456.3	730.9	1,218.3	2,151.6	5,278.4
2002.....	40.2	71.3	119.2	192.8	304.4	479.6	766.1	1,278.2	2,234.3	5,446.8
2001.....	39.1	69.3	119.5	194.2	314.1	494.4	787.7	1,321.2	2,309.0	5,664.2
2000.....	39.4	71.2	120.4	205.5	330.6	515.8	826.7	1,372.3	2,392.1	5,926.1
1999.....	40.1	71.7	123.5	211.0	341.9	547.1	867.9	1,427.8	2,484.7	6,063.0
All races 3/, male										
2005.....	53.6	98.7	168.8	249.4	385.1	549.7	869.1	1,414.1	2,428.3	5,143.4
2004.....	54.2	101.5	167.5	252.3	393.5	580.0	899.4	1,472.2	2,509.6	5,239.3
2003.....	56.9	104.0	173.3	268.1	416.1	624.7	975.4	1,589.0	2,695.8	5,621.5
2002.....	57.7	104.7	177.0	278.2	429.1	654.3	1,025.8	1,667.3	2,786.9	5,823.5
2001.....	55.8	102.6	175.2	279.7	440.9	672.2	1,054.4	1,732.2	2,882.7	6,040.5
2000.....	57.0	107.1	178.1	298.9	463.9	702.0	1,119.7	1,787.5	3,012.3	6,430.0
1999.....	58.1	107.0	182.9	306.5	478.2	752.8	1,174.1	1,865.5	3,123.3	6,561.6
All races 3/, female										
2005.....	21.8	38.0	61.6	97.1	171.2	279.6	479.8	856.0	1,647.8	4,610.8
2004.....	22.7	38.5	62.2	99.1	173.2	287.7	502.6	884.1	1,698.7	4,741.5
2003.....	24.3	39.5	62.4	106.7	187.0	310.3	534.8	954.5	1,818.9	5,126.7
2002.....	22.9	38.8	63.9	112.5	191.2	328.8	558.5	1,003.3	1,898.3	5,283.3
2001.....	22.6	37.0	66.1	114.2	199.0	341.4	575.8	1,033.5	1,965.3	5,506.8
2000.....	22.2	36.4	65.0	118.1	209.8	356.3	596.0	1,083.0	2,026.2	5,720.9
1999.....	22.4	37.5	66.6	121.7	218.6	371.2	628.7	1,123.5	2,111.0	5,862.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Diseases of heart (I00-I09,I11,I13,I20-I51)										
White, both sexes										
2005 4/.....	34.0	61.3	102.5	156.6	255.0	385.2	634.7	1,073.2	1,946.5	4,879.3
2004 4/.....	34.4	62.4	102.3	158.8	261.2	402.0	658.9	1,112.1	2,010.3	4,977.6
2003 4/.....	35.9	63.6	104.7	170.9	275.7	432.1	706.5	1,199.6	2,153.0	5,363.5
2002.....	36.3	64.0	108.1	177.9	284.5	457.0	744.1	1,255.7	2,234.0	5,529.1
2001.....	34.2	61.8	108.6	178.5	294.7	468.4	765.0	1,297.7	2,308.3	5,752.1
2000.....	34.9	63.7	109.9	190.0	309.5	492.6	801.0	1,347.0	2,393.0	6,020.8
1999.....	34.9	64.1	112.7	193.7	321.3	524.0	839.7	1,403.0	2,484.1	6,152.7
White, male										
2005 4/.....	49.8	91.0	155.3	231.6	364.2	527.6	848.6	1,402.3	2,445.7	5,288.4
2004 4/.....	50.5	93.4	154.7	234.6	373.5	556.2	880.1	1,458.9	2,532.6	5,348.4
2003 4/.....	52.1	95.4	159.4	251.9	394.1	599.0	949.5	1,575.7	2,710.0	5,747.2
2002.....	53.9	96.8	164.4	261.0	406.6	631.3	1,006.4	1,654.3	2,805.0	5,939.8
2001.....	50.9	94.4	163.4	261.7	420.5	647.2	1,031.2	1,716.7	2,896.4	6,157.2
2000.....	52.3	99.4	166.1	280.7	442.0	680.1	1,096.8	1,771.0	3,030.0	6,560.8
1999.....	52.4	98.8	170.3	286.1	454.8	730.0	1,148.4	1,851.7	3,144.4	6,669.0
White, female										
2005 4/.....	18.0	31.8	50.8	84.4	153.5	258.6	457.9	829.6	1,633.6	4,691.0
2004 4/.....	18.2	31.5	51.0	85.9	157.2	265.2	477.1	858.3	1,685.8	4,810.4
2003 4/.....	19.6	32.1	51.1	93.1	166.1	284.6	507.6	927.7	1,809.7	5,193.6
2002.....	18.6	31.4	53.1	98.3	171.5	303.6	530.3	970.4	1,884.1	5,350.6
2001.....	17.5	29.4	54.9	99.1	178.6	311.5	549.5	1,001.0	1,952.9	5,582.5
2000.....	17.5	28.3	54.9	103.7	187.4	328.6	563.7	1,048.7	2,014.1	5,801.4
1999.....	17.4	29.7	56.4	105.8	198.3	344.3	594.5	1,088.0	2,094.3	5,945.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Diseases of heart (I00-I09,I11,I13,I20-I51)										
Black, both sexes										
2005 4/.....	69.4	128.5	224.1	326.8	494.7	662.2	958.8	1,494.7	2,282.2	4,326.7
2004 4/.....	73.8	134.9	226.0	337.4	492.5	697.0	1,020.9	1,534.2	2,297.3	4,563.8
2003 4/.....	79.0	139.6	232.4	348.4	533.8	752.7	1,115.2	1,638.3	2,445.2	4,941.2
2002.....	74.8	138.7	232.1	365.8	542.7	764.2	1,126.8	1,715.1	2,557.1	5,115.2
2001.....	79.5	137.2	232.2	369.7	546.5	809.7	1,160.0	1,780.6	2,619.7	5,275.7
2000.....	77.5	142.2	231.2	380.7	571.0	813.9	1,230.4	1,836.2	2,673.7	5,389.5
1999.....	82.9	143.0	240.2	400.6	579.9	852.1	1,289.1	1,878.6	2,749.2	5,487.4
Black, male										
2005 4/.....	91.1	175.9	312.9	461.0	679.8	877.7	1,259.7	1,881.5	2,710.8	4,230.5
2004 4/.....	93.6	182.7	310.5	471.5	685.8	932.8	1,315.6	1,939.0	2,703.1	4,637.3
2003 4/.....	102.7	188.1	322.4	481.2	718.0	997.2	1,460.0	2,083.2	2,981.9	4,850.3
2002.....	96.7	186.2	319.6	503.6	740.6	1,001.5	1,448.8	2,095.3	3,039.3	5,125.7
2001.....	101.9	184.4	314.7	503.1	731.0	1,039.8	1,518.8	2,214.9	3,181.5	5,459.9
2000.....	101.3	190.8	319.1	533.0	754.7	1,033.7	1,570.9	2,245.4	3,226.1	5,633.5
1999.....	109.3	193.3	335.5	556.7	789.2	1,111.8	1,652.9	2,277.2	3,259.4	5,858.8
Black, female										
2005 4/.....	50.2	87.1	149.0	216.0	350.4	502.4	753.0	1,264.0	2,055.7	4,365.6
2004 4/.....	56.4	93.3	154.5	227.0	341.8	521.9	820.9	1,294.4	2,083.5	4,534.7
2003 4/.....	58.2	97.4	156.3	239.2	390.1	571.3	882.6	1,373.9	2,164.0	4,976.5
2002.....	55.5	97.5	157.9	252.6	388.0	588.1	910.7	1,488.0	2,306.1	5,111.2
2001.....	59.7	96.3	162.0	260.3	401.4	639.8	921.0	1,521.3	2,333.5	5,207.3
2000.....	56.5	100.1	156.3	255.9	426.1	651.7	1,004.3	1,589.5	2,395.4	5,298.7
1999.....	59.6	99.6	159.4	273.1	415.8	659.7	1,047.8	1,638.6	2,496.4	5,348.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)										
All races 3/, both sexes										
2005.....	0.2	0.4	0.5	0.9	1.5	2.5	4.2	6.7	10.9	19.0
2004.....	0.3	0.4	0.6	0.9	1.4	2.6	4.2	7.3	10.9	17.1
2003.....	0.3	0.5	0.6	1.0	1.8	3.0	4.6	8.1	11.6	20.5
2002.....	0.2	0.5	0.7	1.3	2.1	2.8	5.0	8.4	11.4	17.8
2001.....	0.4	0.5	0.7	1.2	1.7	3.2	5.0	8.1	11.4	17.7
2000.....	0.4	0.6	0.8	1.4	2.2	3.4	5.6	8.1	11.6	16.8
1999.....	0.3	0.5	0.9	1.4	2.1	3.5	5.8	8.4	12.6	18.0
All races 3/, male										
2005.....	0.2	0.3	0.5	0.9	1.1	1.9	3.1	4.9	8.0	14.0
2004.....	0.2	0.4	0.5	0.7	0.9	2.1	2.8	5.7	9.5	11.5
2003.....	0.2	0.4	0.4	0.8	1.5	2.3	3.6	6.4	9.0	16.2
2002.....	0.2	0.4	0.6	1.0	1.5	1.9	4.2	5.8	8.2	11.8
2001.....	0.3	0.4	0.5	1.1	1.2	2.4	3.2	6.0	7.7	12.8
2000.....	0.3	0.5	0.6	1.1	1.8	2.2	3.6	5.8	7.8	12.2
1999.....	0.3	0.4	0.6	1.0	1.4	2.4	4.7	5.4	8.9	10.3
All races 3/, female										
2005.....	0.3	0.5	0.6	1.0	1.9	3.0	5.1	8.0	12.7	21.3
2004.....	0.3	0.4	0.7	1.1	1.9	3.0	5.3	8.3	11.8	19.5
2003.....	0.3	0.6	0.8	1.1	2.1	3.6	5.5	9.3	13.2	22.3
2002.....	0.3	0.5	0.9	1.5	2.6	3.7	5.6	10.2	13.4	20.4
2001.....	0.4	0.6	0.9	1.3	2.2	3.9	6.4	9.5	13.7	19.8
2000.....	0.5	0.7	0.9	1.6	2.5	4.5	7.1	9.7	13.9	18.7
1999.....	0.4	0.6	1.1	1.8	2.7	4.5	6.7	10.4	14.7	21.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)										
White, both sexes										
2005 4/.....	0.2	0.3	0.4	0.8	1.4	2.5	4.4	7.0	11.5	20.4
2004 4/.....	0.2	0.3	0.6	0.7	1.4	2.6	4.4	7.6	11.6	18.2
2003 4/.....	0.2	0.4	0.5	0.8	1.6	3.1	4.7	8.6	12.4	21.8
2002.....	0.2	0.4	0.7	1.2	2.0	2.8	5.2	8.6	12.1	18.9
2001.....	0.3	0.4	0.6	1.0	1.7	3.2	5.0	8.4	12.2	18.7
2000.....	0.3	0.5	0.7	1.3	2.2	3.5	5.6	8.6	12.4	17.6
1999.....	0.3	0.5	0.8	1.4	2.0	3.6	6.0	8.8	13.1	18.9
White, male										
2005 4/.....	*	0.3	0.4	0.8	1.1	1.9	3.3	5.2	8.3	14.7
2004 4/.....	*	0.3	0.4	0.6	0.9	2.0	3.1	6.1	10.1	12.4
2003 4/.....	*	0.3	0.3	0.7	1.3	2.3	3.7	6.9	9.8	17.3
2002.....	*	0.3	0.5	0.9	1.5	1.8	4.4	5.9	8.8	12.6
2001.....	0.3	0.4	0.4	1.0	1.2	2.3	3.3	6.3	8.0	13.3
2000.....	0.3	0.5	0.7	1.1	1.9	2.3	3.5	6.1	8.4	12.7
1999.....	0.2	0.4	0.6	1.0	1.3	2.6	4.9	5.5	9.4	11.1
White, female										
2005 4/.....	*	0.4	0.5	0.9	1.8	3.1	5.3	8.4	13.6	23.1
2004 4/.....	0.2	0.3	0.7	0.9	1.9	3.0	5.4	8.6	12.5	20.8
2003 4/.....	0.2	0.5	0.6	1.0	1.9	3.8	5.5	9.8	14.0	23.8
2002.....	0.2	0.5	0.9	1.4	2.5	3.7	5.9	10.6	14.2	21.6
2001.....	0.3	0.4	0.7	1.1	2.3	3.9	6.4	9.9	14.7	21.0
2000.....	0.4	0.6	0.8	1.5	2.4	4.6	7.3	10.4	14.7	19.6
1999.....	0.3	0.6	0.9	1.8	2.7	4.6	6.9	11.0	15.4	22.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)

Black, both sexes

2005 4/.....	*	*	1.3	1.2	1.8	2.5	*	4.2	*	6.1
2004 4/.....	0.7	1.0	1.0	2.0	*	2.6	2.7	4.1	*	5.9
2003 4/.....	*	1.2	1.4	2.2	3.2	3.1	4.8	3.9	*	7.6
2002.....	*	0.8	1.1	2.0	2.9	3.2	*	4.8	*	*
2001.....	0.9	1.1	1.6	2.2	2.0	3.3	5.1	4.9	*	6.8
2000.....	0.9	1.0	1.1	1.9	2.2	3.4	3.6	3.7	*	8.4
1999.....	0.7	0.9	1.6	2.2	2.1	2.8	4.2	4.3	*	8.3

Black, male

2005 4/.....	*	*	*	*	*	*	*	*	*	*
2004 4/.....	*	*	*	*	*	*	*	*	*	*
2003 4/.....	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*

Black, female

2005 4/.....	*	*	*	*	*	*	*	5.3	*	*
2004 4/.....	*	*	*	*	*	*	*	5.4	*	*
2003 4/.....	*	1.6	1.7	*	3.3	*	6.2	*	*	8.1
2002.....	*	*	*	*	3.6	*	*	5.5	*	*
2001.....	*	*	2.2	2.7	*	3.8	7.2	6.4	*	*
2000.....	*	*	*	*	*	4.4	*	*	*	*
1999.....	*	*	*	2.8	*	*	4.7	*	*	11.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Hypertensive heart disease (I11)										
All races 3/, both sexes										
2005.....	4.0	6.5	9.2	11.7	14.7	17.6	23.8	35.8	65.9	198.6
2004.....	4.0	6.8	9.0	12.2	14.9	18.8	25.1	36.7	66.1	192.8
2003.....	3.9	6.6	9.4	11.7	14.9	19.4	25.1	39.5	66.8	195.0
2002.....	3.5	5.9	8.6	11.1	14.3	18.9	24.0	37.9	66.0	187.0
2001.....	3.2	5.1	7.7	9.7	12.5	16.3	23.5	36.6	66.0	189.5
2000.....	2.9	4.7	6.6	9.3	13.2	16.5	23.0	36.3	66.1	193.1
1999.....	2.8	4.5	6.2	9.6	11.9	16.3	22.1	36.4	64.7	184.9
All races 3/, male										
2005.....	5.4	9.1	13.0	16.5	20.3	22.5	28.5	39.3	62.2	147.5
2004.....	5.2	9.2	12.4	17.0	20.4	24.7	30.5	39.1	60.9	145.4
2003.....	5.2	9.0	12.8	16.0	18.8	25.2	29.4	42.8	63.3	142.2
2002.....	4.9	8.0	11.9	15.2	19.0	23.6	27.0	41.0	62.2	134.6
2001.....	4.5	7.0	10.3	12.8	16.5	20.5	27.1	38.9	63.5	134.4
2000.....	3.9	6.3	9.4	12.1	17.4	20.7	27.2	37.3	61.8	139.1
1999.....	3.9	6.1	8.1	12.7	15.0	19.5	24.6	39.4	59.2	133.6
All races 3/, female										
2005.....	2.5	3.9	5.5	7.2	9.7	13.3	20.0	33.2	68.2	222.0
2004.....	2.7	4.4	5.8	7.6	9.9	13.5	20.7	35.0	69.3	214.1
2003.....	2.6	4.4	6.1	7.7	11.2	14.4	21.6	37.2	69.0	218.3
2002.....	2.2	3.9	5.4	7.2	10.0	14.8	21.6	35.8	68.2	209.7
2001.....	2.0	3.2	5.1	6.8	8.9	12.7	20.6	35.1	67.5	212.6
2000.....	2.0	3.2	3.8	6.6	9.3	12.9	19.6	35.7	68.6	215.1
1999.....	1.7	3.0	4.3	6.8	9.2	13.5	20.2	34.3	67.9	205.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Hypertensive heart disease (I11)										
White, both sexes										
2005 4/.....	2.8	4.6	6.5	8.9	11.7	13.5	19.4	30.4	58.0	188.9
2004 4/.....	2.8	4.8	6.4	9.0	11.2	15.1	20.4	31.0	59.1	183.7
2003 4/.....	2.7	4.4	6.6	8.5	11.4	14.7	20.2	33.2	61.2	185.3
2002.....	2.4	4.3	6.0	8.1	10.5	14.4	19.3	31.9	58.1	177.3
2001.....	2.1	3.3	5.2	6.9	9.1	12.1	18.5	30.9	59.2	179.9
2000.....	2.0	3.1	4.3	6.2	9.4	12.5	17.6	30.2	58.4	181.8
1999.....	1.8	2.9	4.1	6.5	8.7	12.0	16.9	30.0	57.3	175.3
White, male										
2005 4/.....	4.0	6.9	9.6	12.8	16.4	17.7	22.9	33.6	54.7	136.9
2004 4/.....	4.1	7.0	9.3	12.8	15.3	20.0	25.0	33.4	54.3	136.4
2003 4/.....	4.0	6.4	9.7	12.2	15.0	19.4	23.6	36.1	57.9	131.9
2002.....	3.7	6.1	8.9	11.6	14.0	18.1	22.0	34.5	54.8	124.8
2001.....	3.1	4.8	7.5	9.6	12.5	15.9	21.9	32.9	56.1	123.7
2000.....	2.8	4.6	6.5	8.6	12.7	16.0	21.5	31.1	53.4	129.3
1999.....	2.7	4.2	5.8	8.9	11.3	14.4	19.1	32.4	51.8	123.7
White, female										
2005 4/.....	1.5	2.3	3.6	5.2	7.3	9.7	16.5	28.0	60.1	212.9
2004 4/.....	1.6	2.7	3.5	5.3	7.4	10.8	16.6	29.3	62.1	205.0
2003 4/.....	1.3	2.4	3.5	5.0	8.1	10.7	17.4	31.1	63.3	209.0
2002.....	1.2	2.4	3.3	4.7	7.1	11.1	17.1	30.0	60.1	200.1
2001.....	1.0	1.8	3.0	4.4	6.0	8.8	15.8	29.5	61.1	203.5
2000.....	1.2	1.6	2.2	4.0	6.3	9.4	14.5	29.6	61.3	203.1
1999.....	0.8	1.6	2.5	4.2	6.3	10.0	15.1	28.2	60.6	196.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Hypertensive heart disease (I11)										
Black, both sexes										
2005 4/.....	12.2	20.8	30.3	37.8	46.0	57.8	69.9	100.9	168.5	338.1
2004 4/.....	12.3	22.0	31.3	41.6	49.9	55.4	76.2	103.1	161.4	314.7
2003 4/.....	12.8	23.9	32.8	41.9	48.3	63.0	77.1	113.3	145.7	332.3
2002.....	11.4	19.3	30.3	39.7	50.5	61.2	76.5	110.3	167.7	320.4
2001.....	11.5	18.6	28.4	35.7	43.6	55.8	77.8	105.1	154.7	326.1
2000.....	9.9	17.8	25.4	37.6	47.9	54.6	81.6	111.1	164.9	340.0
1999.....	10.2	17.9	24.8	38.0	41.6	57.6	79.0	113.7	161.2	305.1
Black, male										
2005 4/.....	16.3	27.1	42.3	53.5	64.9	74.2	91.1	118.9	169.2	301.2
2004 4/.....	14.5	27.8	40.8	59.1	72.3	78.3	97.8	114.6	158.5	267.2
2003 4/.....	14.9	29.8	41.5	55.1	59.8	86.7	99.9	129.6	149.1	287.8
2002.....	14.6	24.4	39.9	53.1	71.1	79.0	89.8	125.4	167.4	267.0
2001.....	14.8	25.1	36.6	45.1	55.9	69.6	91.6	117.1	171.8	284.8
2000.....	12.7	21.4	35.8	47.9	63.8	70.0	95.9	117.2	178.7	273.3
1999.....	13.1	22.5	31.3	49.5	52.0	73.0	90.2	129.2	166.5	263.6
Black, female										
2005 4/.....	8.6	15.3	20.2	24.9	31.3	45.7	55.4	90.1	168.0	353.0
2004 4/.....	10.3	17.0	23.2	27.2	32.5	38.5	61.5	96.2	162.9	333.5
2003 4/.....	11.0	18.7	25.5	31.0	39.4	45.5	61.8	103.6	143.9	349.6
2002.....	8.6	14.9	22.3	28.7	34.4	47.9	67.7	101.3	167.9	340.9
2001.....	8.6	13.0	21.5	28.0	33.9	45.6	68.6	97.9	145.9	341.4
2000.....	7.5	14.6	16.5	29.2	35.3	43.2	72.1	107.4	157.9	364.9
1999.....	7.6	13.8	19.3	28.7	33.4	46.2	71.6	104.4	158.6	320.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Hypertensive heart and renal disease (I13)										
All races 3/, both sexes										
2005.....	0.2	0.3	0.6	0.6	1.0	1.8	2.9	5.0	9.5	25.3
2004.....	0.3	0.5	0.5	0.7	1.3	1.6	2.6	4.5	9.4	24.8
2003.....	0.3	0.4	0.6	0.9	1.1	1.6	2.9	5.2	9.5	25.6
2002.....	0.2	0.4	0.5	0.9	1.2	1.6	2.8	4.9	9.5	23.5
2001.....	0.3	0.3	0.4	0.8	1.2	1.8	2.8	5.0	9.3	23.3
2000.....	0.3	0.3	0.4	0.8	1.0	1.6	3.0	5.1	9.8	24.3
1999.....	0.3	0.4	0.4	0.9	1.4	2.1	3.0	6.6	12.3	29.1
All races 3/, male										
2005.....	0.3	0.5	0.7	0.8	1.2	2.1	2.9	5.3	9.8	27.4
2004.....	0.3	0.5	0.7	0.7	1.7	1.7	2.8	5.1	10.4	24.3
2003.....	0.3	0.6	0.7	1.1	1.3	1.7	3.0	5.6	10.1	26.7
2002.....	0.2	0.5	0.6	1.1	1.4	1.5	3.4	5.1	9.9	23.1
2001.....	0.4	0.4	0.6	0.9	1.2	2.1	2.9	5.2	9.4	24.2
2000.....	0.3	0.4	0.5	0.9	1.0	1.7	3.4	5.0	10.8	27.8
1999.....	0.4	0.5	0.6	1.0	1.6	2.3	3.2	7.3	13.3	31.3
All races 3/, female										
2005.....	0.2	*	0.4	0.5	0.8	1.6	2.8	4.8	9.3	24.3
2004.....	*	0.4	0.4	0.7	0.9	1.5	2.3	4.0	8.8	25.0
2003.....	0.2	0.2	0.5	0.6	0.9	1.5	2.8	4.9	9.2	25.2
2002.....	0.2	0.2	0.4	0.7	1.0	1.7	2.4	4.7	9.2	23.7
2001.....	*	*	0.3	0.7	1.3	1.5	2.8	4.9	9.3	22.9
2000.....	0.2	0.2	0.4	0.7	1.0	1.5	2.7	5.1	9.3	22.9
1999.....	0.3	0.3	0.3	0.7	1.2	2.0	2.9	6.1	11.7	28.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Hypertensive heart and renal disease (I13)										
White, both sexes										
2005 4/.....	*	0.1	0.2	0.3	0.5	1.1	2.0	3.9	7.9	23.9
2004 4/.....	0.1	0.1	0.2	0.3	0.7	0.9	1.8	3.5	8.1	23.0
2003 4/.....	*	0.1	0.2	0.3	0.6	0.9	2.1	4.2	8.3	23.4
2002.....	*	*	0.1	0.3	0.6	0.9	1.9	3.8	8.4	21.0
2001.....	*	*	0.2	0.4	0.7	1.1	2.0	3.9	8.3	21.0
2000.....	0.2	0.1	0.2	0.4	0.5	0.9	2.2	4.0	8.1	22.0
1999.....	*	0.1	0.2	0.4	0.7	1.2	1.9	5.2	10.4	26.1
White, male										
2005 4/.....	*	*	0.3	0.4	0.5	1.1	2.1	4.2	8.2	25.5
2004 4/.....	*	*	*	0.3	1.0	1.0	2.1	3.9	9.2	22.9
2003 4/.....	*	*	*	0.5	0.8	1.1	2.2	4.9	9.3	24.3
2002.....	*	*	*	0.5	0.7	0.8	2.4	4.2	8.8	20.6
2001.....	*	*	0.3	0.4	0.7	1.3	2.0	4.1	8.7	22.7
2000.....	*	*	*	0.4	0.5	1.0	2.5	4.4	9.2	24.4
1999.....	*	*	*	0.5	0.8	1.4	2.0	5.8	12.1	28.8
White, female										
2005 4/.....	*	*	*	*	0.4	1.0	1.9	3.7	7.8	23.2
2004 4/.....	*	*	*	0.3	0.4	0.8	1.5	3.2	7.4	23.0
2003 4/.....	*	*	*	*	0.5	0.7	1.9	3.7	7.8	22.9
2002.....	*	*	*	*	0.5	0.9	1.5	3.6	8.1	21.1
2001.....	*	*	*	0.3	0.8	0.8	2.0	3.7	8.0	20.3
2000.....	*	*	*	0.4	0.5	0.8	2.0	3.7	7.4	21.1
1999.....	*	*	*	0.4	0.6	1.1	1.8	4.7	9.5	25.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Hypertensive heart and renal disease (I13)										
Black, both sexes										
2005 4/.....	1.2	1.7	3.4	3.7	5.7	8.7	10.7	16.5	28.4	42.8
2004 4/.....	1.3	2.8	2.9	3.9	6.4	7.5	9.7	15.2	26.0	48.2
2003 4/.....	1.5	2.3	3.8	5.3	5.5	7.4	11.0	16.5	24.1	53.1
2002.....	1.1	2.2	3.3	6.0	6.2	8.0	12.3	17.4	23.0	56.1
2001.....	1.4	1.9	2.3	4.4	5.7	8.4	11.8	18.4	22.8	50.1
2000.....	1.2	1.9	2.5	4.1	5.3	8.3	11.9	17.6	31.1	50.4
1999.....	2.1	2.5	2.3	5.0	7.3	10.1	15.4	21.8	35.3	65.4
Black, male										
2005 4/.....	*	2.8	4.0	4.7	8.0	11.6	10.4	17.0	33.0	50.2
2004 4/.....	2.0	3.3	3.8	4.4	8.6	8.2	9.4	20.5	27.3	41.7
2003 4/.....	2.1	3.4	4.6	7.6	6.4	6.6	11.8	15.5	24.4	50.0
2002.....	*	3.0	4.7	7.6	7.3	7.2	14.0	17.0	22.9	61.1
2001.....	2.1	3.0	3.5	5.6	6.3	10.7	13.5	18.1	19.3	44.3
2000.....	1.5	2.7	2.9	5.1	5.6	9.4	13.7	13.7	33.7	63.4
1999.....	2.6	3.5	3.4	6.3	9.3	10.6	17.3	23.7	29.6	59.8
Black, female										
2005 4/.....	*	*	2.9	2.9	3.8	6.5	10.8	16.3	26.0	39.8
2004 4/.....	*	2.4	2.2	3.5	4.7	7.0	9.9	12.1	25.3	50.8
2003 4/.....	*	*	3.1	3.5	4.8	8.0	10.4	17.1	24.0	54.3
2002.....	*	1.5	2.1	4.6	5.3	8.5	11.2	17.6	23.1	54.2
2001.....	*	*	*	3.3	5.3	6.8	10.8	18.6	24.5	52.2
2000.....	*	*	2.1	3.2	5.1	7.5	10.7	19.9	29.7	45.5
1999.....	1.6	1.6	*	4.0	5.7	9.6	14.2	20.6	38.0	67.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Ischemic heart diseases (I20-I25)										
All races 3/, both sexes										
2005.....	21.6	42.9	78.3	120.9	198.6	295.4	472.3	778.6	1,359.8	3,158.1
2004.....	22.4	44.0	77.9	123.9	203.1	310.5	495.9	813.5	1,418.0	3,287.4
2003.....	23.5	45.9	80.7	132.4	218.7	335.0	538.1	886.4	1,538.2	3,610.2
2002.....	23.8	46.4	83.5	140.1	226.2	358.4	569.8	940.2	1,611.2	3,775.0
2001.....	23.4	45.6	84.8	142.8	236.6	372.4	591.8	984.3	1,685.2	3,958.2
2000.....	23.6	48.0	86.7	153.9	250.0	390.8	628.0	1,032.5	1,761.7	4,182.5
1999.....	24.2	48.5	90.1	157.2	260.0	420.1	661.9	1,081.1	1,841.7	4,318.1
All races 3/, male										
2005.....	33.0	66.2	121.3	183.8	291.3	415.2	653.7	1,054.3	1,780.2	3,576.8
2004.....	34.0	68.2	121.1	187.5	297.8	440.2	682.0	1,106.0	1,855.0	3,680.3
2003.....	35.5	71.4	126.9	200.3	319.7	476.1	744.1	1,200.1	2,013.5	4,005.2
2002.....	36.5	72.3	130.3	211.4	331.4	508.5	791.5	1,274.8	2,094.2	4,205.3
2001.....	35.8	71.6	131.1	215.5	344.4	525.7	820.3	1,337.5	2,180.0	4,395.8
2000.....	37.0	76.9	134.2	232.8	362.9	550.5	879.6	1,394.4	2,305.0	4,701.7
1999.....	37.9	77.0	140.3	237.7	376.1	595.2	924.6	1,462.5	2,400.4	4,837.5
All races 3/, female										
2005.....	10.4	20.2	37.0	61.5	114.1	190.9	325.4	578.5	1,098.7	2,965.8
2004.....	10.8	20.4	36.6	64.0	116.8	197.7	346.0	603.2	1,148.8	3,110.6
2003.....	11.6	21.1	36.5	68.4	126.9	212.6	373.0	663.1	1,247.5	3,435.5
2002.....	11.2	21.4	38.7	73.1	130.7	228.8	392.6	704.0	1,317.5	3,588.3
2001.....	11.1	20.3	40.3	74.8	138.8	240.5	410.2	737.0	1,388.7	3,775.1
2000.....	10.3	20.0	41.3	80.0	147.7	253.9	429.9	780.5	1,441.2	3,971.0
1999.....	10.7	20.8	42.1	82.1	155.0	270.2	456.7	815.9	1,514.7	4,108.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Ischemic heart diseases (I20-I25)										
White, both sexes										
2005 4/.....	21.0	41.1	73.8	114.2	189.8	286.3	463.7	771.1	1,364.5	3,227.0
2004 4/.....	21.6	42.0	73.4	117.5	195.2	298.7	485.8	807.0	1,423.8	3,343.9
2003 4/.....	22.4	43.5	76.1	126.4	208.2	322.6	526.3	877.2	1,542.2	3,671.7
2002.....	22.9	44.2	79.1	133.3	216.1	346.6	559.2	928.9	1,616.4	3,836.4
2001.....	22.1	43.1	80.3	135.4	225.8	357.9	580.5	972.9	1,689.9	4,022.7
2000.....	22.2	45.3	82.4	146.2	238.5	378.4	615.0	1,019.7	1,768.5	4,257.7
1999.....	22.6	46.0	85.4	149.1	249.5	408.5	647.3	1,070.6	1,849.5	4,389.2
White, male										
2005 4/.....	32.7	64.1	116.2	175.7	281.5	406.5	646.8	1,053.1	1,798.9	3,687.1
2004 4/.....	33.7	66.3	116.1	180.0	289.4	428.5	673.9	1,105.8	1,880.8	3,768.8
2003 4/.....	34.4	68.8	121.6	194.2	308.7	464.4	731.5	1,195.8	2,028.8	4,106.1
2002.....	35.9	70.3	125.6	203.9	320.5	498.1	784.6	1,273.1	2,115.1	4,299.7
2001.....	34.6	69.5	126.4	207.2	333.9	512.8	810.3	1,335.1	2,201.7	4,494.0
2000.....	35.9	74.5	130.0	223.7	352.1	541.2	870.1	1,390.0	2,328.3	4,813.9
1999.....	36.2	74.7	135.0	228.2	364.4	586.0	913.4	1,463.9	2,429.7	4,933.9
White, female										
2005 4/.....	9.2	18.1	32.1	54.9	104.5	179.4	312.3	562.4	1,092.2	3,015.2
2004 4/.....	9.4	17.9	31.5	57.3	107.8	183.6	331.3	588.5	1,140.0	3,152.3
2003 4/.....	10.4	18.4	31.5	61.3	115.2	197.3	358.4	646.9	1,242.2	3,479.3
2002.....	9.8	18.2	33.6	65.7	119.6	213.2	375.5	682.5	1,310.6	3,635.2
2001.....	9.5	17.0	35.2	66.8	126.1	222.0	394.4	716.3	1,380.6	3,825.5
2000.....	8.5	16.4	36.0	72.5	133.8	236.0	410.4	759.2	1,435.5	4,031.8
1999.....	8.9	17.6	37.0	73.8	143.6	253.7	435.9	794.5	1,507.0	4,170.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Ischemic heart diseases (I20-I25)										
Black, both sexes										
2005 4/.....	30.2	65.0	129.0	202.1	320.9	431.0	638.8	1,004.8	1,517.3	2,804.4
2004 4/.....	33.2	67.4	130.6	207.3	320.6	471.5	691.7	1,030.8	1,554.7	3,014.3
2003 4/.....	35.2	71.5	133.3	213.8	352.2	503.0	756.8	1,139.8	1,696.1	3,310.1
2002.....	35.1	72.8	136.5	230.9	362.0	522.5	779.6	1,196.0	1,768.0	3,462.9
2001.....	36.5	72.5	138.8	236.1	376.4	562.5	810.8	1,249.3	1,837.2	3,615.1
2000.....	37.1	78.2	140.3	250.9	392.9	566.1	866.5	1,296.9	1,882.8	3,684.7
1999.....	39.7	76.9	149.3	258.5	395.4	596.2	906.3	1,321.5	1,926.9	3,805.8
Black, male										
2005 4/.....	41.2	95.4	187.9	294.0	454.6	584.9	857.4	1,296.8	1,887.0	2,785.0
2004 4/.....	44.9	96.6	189.8	297.2	454.8	641.3	922.9	1,332.6	1,860.9	3,100.5
2003 4/.....	50.2	103.2	195.6	302.7	490.5	677.1	1,027.8	1,501.3	2,138.8	3,285.3
2002.....	48.4	101.6	196.9	330.3	509.6	703.3	1,024.8	1,495.0	2,171.1	3,556.2
2001.....	50.8	101.9	199.9	333.7	515.8	743.8	1,093.0	1,582.5	2,244.8	3,775.0
2000.....	51.8	111.6	197.6	366.7	530.3	727.5	1,139.7	1,630.6	2,316.5	3,885.2
1999.....	56.8	110.4	217.6	372.5	553.5	788.8	1,188.9	1,630.3	2,312.4	4,080.1
Black, female										
2005 4/.....	20.5	38.6	79.1	126.3	216.6	316.9	489.1	830.7	1,322.0	2,812.2
2004 4/.....	22.9	42.0	80.5	133.4	215.9	345.4	534.9	852.0	1,393.3	2,980.1
2003 4/.....	21.9	43.9	80.5	140.7	244.3	373.7	574.0	924.9	1,464.2	3,319.7
2002.....	23.5	47.9	85.4	149.3	246.6	388.3	615.0	1,017.3	1,558.1	3,427.2
2001.....	23.9	47.1	86.8	156.0	266.7	428.6	622.8	1,050.3	1,629.6	3,555.7
2000.....	24.1	49.2	91.4	156.0	284.5	447.0	685.1	1,095.7	1,664.3	3,610.0
1999.....	24.7	48.0	91.4	165.5	271.4	453.5	718.8	1,135.4	1,735.9	3,703.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Acute myocardial infarction (I21-I22)										
All races 3/, both sexes										
2005.....	8.7	17.3	30.9	47.8	78.9	115.0	176.6	272.1	448.1	923.1
2004.....	8.9	17.7	31.6	50.6	82.7	124.1	187.7	291.1	476.0	992.9
2003.....	9.6	18.8	33.2	55.5	90.8	134.5	207.4	325.6	532.5	1,118.8
2002.....	9.9	19.0	34.9	58.4	96.2	148.5	227.5	352.5	572.7	1,191.3
2001.....	9.9	19.7	36.5	61.3	100.1	154.1	235.6	374.0	611.5	1,265.4
2000.....	9.9	21.1	37.8	67.0	110.9	166.8	254.7	405.3	650.1	1,338.5
1999.....	10.4	21.5	39.0	69.7	114.9	181.7	271.6	422.9	689.4	1,379.3
All races 3/, male										
2005.....	13.1	26.5	47.2	72.2	114.0	159.3	240.5	361.8	567.1	1,045.0
2004.....	13.2	27.4	48.7	76.3	120.6	173.0	254.1	386.8	608.9	1,120.5
2003.....	14.4	29.2	52.1	83.4	131.6	188.5	281.4	430.0	682.4	1,263.5
2002.....	14.8	29.2	53.7	86.6	139.0	207.6	312.0	466.1	726.6	1,366.4
2001.....	15.0	30.8	55.6	91.3	143.5	214.2	322.0	495.6	776.1	1,461.3
2000.....	15.5	33.6	57.5	99.7	158.6	231.3	349.6	535.2	835.8	1,568.3
1999.....	16.0	33.7	60.9	105.3	164.9	253.5	373.8	564.2	885.1	1,622.1
All races 3/, female										
2005.....	4.4	8.4	15.2	24.7	47.0	76.4	124.8	206.9	374.1	867.1
2004.....	4.5	8.3	15.3	26.4	48.2	81.5	134.3	222.3	394.1	935.5
2003.....	5.0	8.7	15.1	29.3	53.8	87.7	148.0	251.3	440.9	1,054.8
2002.....	5.1	9.1	17.0	32.0	57.4	97.6	159.8	272.2	479.1	1,115.3
2001.....	4.9	9.0	18.2	33.2	60.7	102.4	167.0	288.8	512.9	1,183.5
2000.....	4.4	8.9	18.9	36.5	67.7	111.5	179.9	314.9	540.6	1,244.8
1999.....	4.9	9.7	18.0	36.5	69.6	120.2	191.7	324.7	574.9	1,281.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Acute myocardial infarction (I21-I22)										
White, both sexes										
2005 4/.....	8.6	17.0	30.3	46.3	78.0	113.8	175.8	270.1	449.4	936.8
2004 4/.....	8.7	17.4	30.6	49.4	81.8	121.7	185.3	291.1	478.6	1,004.1
2003 4/.....	9.3	18.2	32.1	54.4	89.1	132.3	205.3	324.2	532.9	1,132.6
2002.....	9.7	18.4	34.2	57.3	94.3	146.4	225.0	349.5	572.7	1,201.2
2001.....	9.5	19.3	35.4	59.8	98.4	150.8	233.3	371.9	612.7	1,277.8
2000.....	9.5	20.6	36.7	65.6	108.9	164.1	251.8	401.7	652.6	1,353.5
1999.....	10.0	21.1	38.3	68.2	113.4	180.6	268.3	421.1	692.5	1,393.2
White, male										
2005 4/.....	13.1	26.3	47.2	70.7	114.2	160.0	241.8	363.3	572.8	1,070.1
2004 4/.....	13.4	27.2	48.0	75.4	121.2	171.7	252.9	391.6	618.6	1,143.7
2003 4/.....	14.1	28.7	51.2	83.0	130.8	188.3	280.0	430.3	685.7	1,289.3
2002.....	14.7	29.0	53.3	86.2	138.7	207.7	313.1	467.1	730.8	1,386.1
2001.....	14.9	30.9	55.1	90.6	143.7	213.3	321.9	497.6	784.0	1,481.1
2000.....	15.3	33.6	57.2	98.8	158.3	231.1	349.3	536.3	843.5	1,592.9
1999.....	15.8	33.8	60.8	104.0	165.1	254.3	374.0	568.8	896.6	1,644.0
White, female										
2005 4/.....	3.9	7.7	13.7	22.9	44.3	72.6	121.3	201.2	372.1	875.4
2004 4/.....	3.9	7.5	13.5	24.3	45.2	77.4	129.7	217.6	391.6	941.1
2003 4/.....	4.5	7.7	13.4	26.8	50.5	82.8	144.1	247.5	438.7	1,063.2
2002.....	4.6	8.0	15.4	29.7	53.3	92.5	153.2	265.3	475.8	1,120.8
2001.....	4.2	7.7	16.1	30.3	56.7	96.0	161.5	282.8	509.1	1,192.7
2000.....	3.7	7.7	16.6	34.1	63.5	105.5	173.6	307.0	539.1	1,256.2
1999.....	4.3	8.4	16.2	34.2	65.8	116.3	184.3	317.3	572.1	1,292.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Acute myocardial infarction (I21-I22)										
Black, both sexes										
2005 4/.....	11.8	24.2	43.3	69.9	107.8	150.2	217.0	345.4	499.1	892.1
2004 4/.....	12.4	25.0	47.9	75.3	112.7	169.2	249.7	351.1	520.9	979.9
2003 4/.....	14.4	27.5	49.5	79.4	124.3	180.3	270.2	401.4	597.0	1,094.9
2002.....	13.9	27.7	50.4	84.2	136.3	193.6	295.9	434.1	643.9	1,201.2
2001.....	14.3	27.8	53.2	89.2	135.9	211.1	304.3	452.3	671.2	1,239.9
2000.....	15.1	29.6	55.5	94.7	151.4	219.0	328.8	490.6	698.8	1,275.7
1999.....	15.4	30.0	54.9	96.9	150.2	224.0	347.5	495.1	729.6	1,318.8
Black, male										
2005 4/.....	15.8	34.4	59.5	100.2	145.2	190.8	278.3	425.6	598.3	885.9
2004 4/.....	15.4	36.3	67.6	104.8	150.1	223.1	322.7	419.5	591.1	1,009.3
2003 4/.....	20.0	38.6	70.9	108.1	167.9	230.2	353.4	516.2	736.1	1,113.5
2002.....	19.0	37.1	70.4	114.5	177.6	247.8	370.2	531.4	776.4	1,280.1
2001.....	18.5	36.4	71.2	120.1	174.5	264.8	386.4	548.7	792.2	1,382.0
2000.....	20.6	41.0	72.7	131.4	196.7	270.9	419.2	592.5	856.5	1,418.5
1999.....	21.1	41.1	76.9	138.8	198.8	290.8	435.5	591.7	858.0	1,498.7
Black, female										
2005 4/.....	8.4	15.3	29.6	44.9	78.6	120.0	175.1	297.6	446.6	894.6
2004 4/.....	9.7	15.2	31.2	51.0	83.6	129.2	200.1	310.5	483.8	968.3
2003 4/.....	9.5	18.0	31.5	55.7	90.3	143.3	214.0	333.2	524.2	1,087.7
2002.....	9.3	19.5	33.4	59.4	104.1	153.4	246.0	376.0	574.9	1,171.0
2001.....	10.5	20.3	37.8	63.8	105.6	171.4	249.5	394.7	609.5	1,187.2
2000.....	10.1	19.8	40.8	64.6	115.7	180.7	268.8	429.2	619.4	1,222.5
1999.....	10.4	20.5	36.2	62.6	112.1	174.5	289.2	436.8	665.9	1,251.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other acute ischemic heart diseases (I24)										
All races 3/, both sexes										
2005.....	0.3	0.6	1.1	1.4	2.1	2.4	4.2	5.4	9.5	20.8
2004.....	0.3	0.4	0.7	1.1	1.4	1.8	2.6	4.2	6.5	14.8
2003.....	0.3	0.5	0.9	1.3	1.7	2.7	3.6	5.2	8.3	20.0
2002.....	0.3	0.6	1.2	1.4	2.2	3.0	3.9	5.7	9.0	21.1
2001.....	0.3	0.6	1.0	1.4	2.4	2.9	4.1	5.3	9.2	21.5
2000.....	0.3	0.5	1.1	1.6	2.5	3.1	4.0	5.9	9.4	21.9
1999.....	0.3	0.5	0.9	1.6	2.2	3.1	4.3	5.5	9.2	21.9
All races 3/, male										
2005.....	0.4	0.8	1.6	2.1	3.2	3.4	5.3	7.1	10.6	20.2
2004.....	0.4	0.5	1.0	1.6	2.0	2.7	3.4	5.0	8.3	16.3
2003.....	0.5	0.7	1.3	1.8	2.4	3.8	4.9	7.1	9.4	19.5
2002.....	0.4	0.8	1.8	2.1	3.0	4.3	5.1	7.0	11.3	21.4
2001.....	0.4	0.9	1.6	2.0	3.5	4.1	5.5	6.7	11.3	20.7
2000.....	0.4	0.8	1.6	2.5	3.5	4.4	5.2	7.6	11.7	20.7
1999.....	0.4	0.7	1.3	2.5	3.1	4.3	5.6	7.6	10.8	22.3
All races 3/, female										
2005.....	*	0.4	0.6	0.7	1.2	1.6	3.3	4.2	8.8	21.1
2004.....	*	0.2	0.4	0.6	0.8	1.0	1.9	3.6	5.4	14.1
2003.....	*	0.3	0.5	0.8	1.0	1.8	2.6	3.9	7.6	20.2
2002.....	*	0.3	0.7	0.7	1.4	1.9	3.0	4.7	7.6	21.0
2001.....	*	0.3	0.5	0.9	1.3	2.0	3.1	4.4	8.0	21.8
2000.....	*	*	0.5	0.8	1.5	1.9	2.9	4.8	8.1	22.4
1999.....	*	0.2	0.6	0.7	1.5	2.0	3.2	4.1	8.3	21.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other acute ischemic heart diseases (I24)										
White, both sexes										
2005 4/.....	0.2	0.5	0.9	1.2	1.9	2.3	3.9	5.1	9.2	20.5
2004 4/.....	0.3	0.3	0.6	0.9	1.3	1.6	2.4	3.9	6.3	15.0
2003 4/.....	0.3	0.4	0.8	1.1	1.6	2.6	3.4	5.0	8.1	20.2
2002.....	0.2	0.5	1.1	1.3	2.0	2.9	3.8	5.6	8.9	21.4
2001.....	0.2	0.5	0.9	1.3	2.1	2.7	4.0	5.2	9.2	21.5
2000.....	0.2	0.4	1.0	1.4	2.2	2.9	3.8	5.6	9.3	22.1
1999.....	0.2	0.3	0.8	1.4	2.1	3.0	4.0	5.4	8.9	22.4
White, male										
2005 4/.....	0.3	0.7	1.4	1.8	2.8	3.2	4.9	6.6	10.3	19.7
2004 4/.....	0.4	0.4	0.8	1.4	1.8	2.3	3.1	4.6	8.3	16.2
2003 4/.....	0.4	0.6	1.2	1.5	2.2	3.6	4.8	6.7	8.9	19.7
2002.....	0.4	0.7	1.6	2.0	2.8	4.1	4.8	7.0	11.1	21.9
2001.....	0.3	0.8	1.3	1.8	3.1	3.5	5.4	6.5	11.6	20.7
2000.....	0.4	0.7	1.5	2.2	3.2	4.0	4.9	7.2	11.6	21.0
1999.....	0.3	0.5	1.2	2.2	2.8	4.1	5.2	7.6	10.3	22.9
White, female										
2005 4/.....	*	0.2	0.4	0.6	1.0	1.6	3.1	4.0	8.5	20.8
2004 4/.....	*	*	0.3	0.5	0.8	1.0	1.9	3.4	5.1	14.4
2003 4/.....	*	0.2	0.4	0.7	0.9	1.6	2.4	3.8	7.7	20.5
2002.....	*	0.3	0.6	0.7	1.2	1.8	2.9	4.6	7.5	21.1
2001.....	*	0.2	0.4	0.8	1.2	1.9	2.9	4.3	7.7	21.7
2000.....	*	*	0.5	0.6	1.3	1.9	2.8	4.5	8.0	22.6
1999.....	*	*	0.5	0.7	1.5	2.0	3.0	3.8	8.1	22.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other acute ischemic heart diseases (I24)										
Black, both sexes										
2005 4/.....	*	1.3	2.8	3.2	4.5	4.1	7.3	9.7	14.8	27.3
2004 4/.....	*	1.0	1.9	2.9	3.0	3.9	4.3	8.1	9.9	15.2
2003 4/.....	0.7	1.1	1.5	2.7	3.3	4.9	6.2	8.7	11.7	19.8
2002.....	*	1.1	2.4	2.1	4.1	5.0	7.1	7.7	11.8	21.1
2001.....	0.7	1.1	2.1	2.8	5.2	5.3	6.2	6.9	10.8	24.0
2000.....	*	1.0	2.1	3.8	4.7	5.2	6.1	10.0	12.1	21.9
1999.....	*	1.5	2.2	3.3	3.9	4.3	7.3	8.7	13.9	17.2
Black, male										
2005 4/.....	*	*	3.6	5.0	6.9	6.4	9.8	14.8	17.6	27.9
2004 4/.....	*	*	2.7	4.5	4.7	7.0	7.1	11.8	*	22.8
2003 4/.....	*	1.7	2.2	4.6	5.0	7.4	7.9	14.2	19.2	24.0
2002.....	*	1.9	3.6	3.1	5.7	8.0	10.7	9.2	16.0	21.4
2001.....	*	*	3.6	3.4	8.1	9.1	7.7	11.2	*	23.8
2000.....	*	*	3.6	5.8	6.6	8.3	9.0	13.2	*	*
1999.....	*	2.5	2.8	5.8	6.6	6.8	10.2	10.0	17.4	*
Black, female										
2005 4/.....	*	*	2.2	*	*	*	5.6	6.7	13.4	27.0
2004 4/.....	*	*	*	*	*	*	*	5.9	9.1	12.2
2003 4/.....	*	*	*	*	*	*	5.1	5.4	7.7	18.2
2002.....	*	*	*	*	*	*	4.7	6.9	9.5	20.9
2001.....	*	*	*	*	*	*	5.2	*	11.9	24.0
2000.....	*	*	*	*	3.3	*	*	8.0	10.6	22.3
1999.....	*	*	*	*	*	*	5.4	8.0	12.1	16.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other forms of chronic ischemic heart disease (I20,I25)										
All races 3/, both sexes										
2005.....	12.7	25.1	46.3	71.7	117.5	178.0	291.6	501.1	902.3	2,214.2
2004.....	13.2	25.9	45.6	72.2	119.0	184.6	305.6	518.2	935.6	2,279.7
2003.....	13.5	26.6	46.6	75.6	126.1	197.8	327.1	555.6	997.3	2,471.4
2002.....	13.6	26.8	47.3	80.2	127.8	206.8	338.4	582.1	1,029.5	2,562.5
2001.....	13.2	25.2	47.3	80.1	134.1	215.4	352.0	605.0	1,064.5	2,671.3
2000.....	13.4	26.5	47.9	85.2	136.6	221.0	369.4	621.2	1,102.2	2,822.1
1999.....	13.5	26.5	50.2	85.9	142.9	235.3	386.1	652.6	1,143.1	2,917.0
All races 3/, male										
2005.....	19.5	39.0	72.5	109.5	174.1	252.6	407.9	685.4	1,202.5	2,511.6
2004.....	20.4	40.3	71.4	109.5	175.2	264.4	424.5	714.2	1,237.9	2,543.6
2003.....	20.7	41.5	73.5	115.1	185.6	283.8	457.8	763.1	1,321.6	2,722.2
2002.....	21.4	42.2	74.9	122.7	189.4	296.6	474.3	801.6	1,356.2	2,817.4
2001.....	20.4	40.0	74.0	122.2	197.4	307.4	492.9	835.3	1,392.6	2,913.8
2000.....	21.1	42.5	75.1	130.6	200.8	314.8	524.7	851.5	1,457.6	3,112.7
1999.....	21.5	42.6	78.1	129.9	208.1	337.3	545.2	890.7	1,504.6	3,193.1
All races 3/, female										
2005.....	5.9	11.5	21.2	36.1	65.9	112.9	197.3	367.3	715.8	2,077.6
2004.....	6.1	11.9	20.9	37.0	67.8	115.1	209.8	377.3	749.3	2,161.0
2003.....	6.5	12.1	20.9	38.4	72.0	123.2	222.4	408.0	799.1	2,360.5
2002.....	6.0	11.9	21.0	40.3	72.0	129.4	229.7	427.1	830.8	2,451.9
2001.....	6.1	11.0	21.7	40.6	76.8	136.2	240.1	443.8	867.9	2,569.8
2000.....	5.8	10.9	21.8	42.7	78.5	140.5	247.0	460.8	892.5	2,703.8
1999.....	5.6	10.9	23.4	44.9	83.9	148.1	261.7	487.1	931.6	2,805.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other forms of chronic ischemic heart disease (I20,I25)										
White, both sexes										
2005 4/.....	12.2	23.6	42.6	66.7	109.9	170.2	283.9	495.8	905.8	2,269.8
2004 4/.....	12.6	24.4	42.2	67.2	112.1	175.4	298.1	512.0	939.0	2,324.8
2003 4/.....	12.9	24.9	43.2	71.0	117.6	187.7	317.6	548.0	1,001.1	2,518.8
2002.....	13.0	25.2	43.9	74.7	119.8	197.3	330.5	573.8	1,034.7	2,613.9
2001.....	12.3	23.3	44.0	74.3	125.3	204.5	343.2	595.8	1,068.1	2,723.5
2000.....	12.5	24.3	44.8	79.2	127.3	211.4	359.4	612.4	1,106.6	2,882.1
1999.....	12.3	24.6	46.3	79.4	134.0	225.0	375.0	644.2	1,148.1	2,973.6
White, male										
2005 4/.....	19.2	37.1	67.6	103.3	164.5	243.3	400.0	683.2	1,215.8	2,597.3
2004 4/.....	19.9	38.7	67.2	103.3	166.4	254.5	417.9	709.7	1,254.0	2,608.9
2003 4/.....	19.9	39.5	69.2	109.6	175.8	272.6	446.7	758.8	1,334.2	2,797.1
2002.....	20.8	40.6	70.7	115.8	179.1	286.2	466.8	799.0	1,373.2	2,891.6
2001.....	19.4	37.7	69.9	114.8	187.1	296.0	483.0	831.0	1,406.1	2,992.2
2000.....	20.3	40.1	71.3	122.7	190.6	306.1	515.9	846.5	1,473.2	3,200.0
1999.....	20.1	40.4	72.9	122.0	196.5	327.6	534.1	887.5	1,522.9	3,267.1
White, female										
2005 4/.....	5.1	10.2	18.0	31.4	59.2	105.2	187.9	357.2	711.6	2,119.0
2004 4/.....	5.3	10.2	17.7	32.5	61.8	105.2	199.7	367.4	743.3	2,196.8
2003 4/.....	5.8	10.5	17.7	33.8	63.8	112.8	211.9	395.6	795.8	2,395.5
2002.....	5.1	10.0	17.6	35.3	65.1	118.9	219.3	412.6	827.2	2,493.2
2001.....	5.2	9.0	18.7	35.8	68.2	124.1	230.0	429.2	863.8	2,611.0
2000.....	4.7	8.6	18.8	37.8	69.0	128.7	233.9	447.7	888.4	2,753.0
1999.....	4.5	9.0	20.3	38.9	76.3	135.5	248.5	473.4	926.8	2,855.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other forms of chronic ischemic heart disease (I20,I25)										
Black, both sexes										
2005 4/.....	17.9	39.5	82.8	129.0	208.5	276.7	414.4	649.7	1,003.4	1,885.0
2004 4/.....	20.3	41.4	80.8	129.1	204.8	298.4	437.7	671.6	1,023.9	2,019.1
2003 4/.....	20.1	42.8	82.2	131.7	224.6	317.7	480.4	729.7	1,087.4	2,195.3
2002.....	20.7	44.0	83.7	144.6	221.6	323.9	476.6	754.2	1,112.3	2,240.6
2001.....	21.5	43.7	83.6	144.1	235.3	346.1	500.3	790.1	1,155.2	2,351.2
2000.....	21.5	47.6	82.7	152.4	236.7	341.9	531.5	796.3	1,171.8	2,387.1
1999.....	23.8	45.4	92.2	158.4	241.2	367.9	551.4	817.7	1,183.5	2,469.8
Black, male										
2005 4/.....	24.7	59.5	124.8	188.8	302.5	387.6	569.3	856.5	1,271.1	1,871.2
2004 4/.....	28.8	59.1	119.5	187.9	300.1	411.1	593.0	901.3	1,258.3	2,068.3
2003 4/.....	29.4	62.9	122.5	189.9	317.6	439.5	666.5	970.9	1,383.5	2,147.8
2002.....	29.0	62.5	122.8	212.7	326.3	447.5	643.9	954.4	1,378.6	2,254.6
2001.....	31.2	64.0	125.1	210.1	333.1	469.8	698.8	1,022.7	1,443.9	2,369.1
2000.....	30.3	69.2	121.3	229.5	326.9	448.3	711.4	1,024.9	1,444.9	2,446.0
1999.....	34.9	66.9	138.0	227.8	348.1	491.2	743.2	1,028.6	1,437.0	2,561.5
Black, female										
2005 4/.....	11.9	22.1	47.3	79.7	135.3	194.4	308.4	526.4	862.1	1,890.6
2004 4/.....	12.8	26.0	48.2	80.7	130.5	214.6	332.3	535.6	900.4	1,999.6
2003 4/.....	11.9	25.3	48.2	83.8	152.1	227.3	354.9	586.3	932.3	2,213.8
2002.....	13.3	28.0	50.5	88.6	139.7	232.2	364.3	634.5	973.6	2,235.2
2001.....	13.0	26.1	48.2	89.9	158.2	254.8	368.1	651.2	1,008.1	2,344.5
2000.....	13.7	28.8	49.7	89.2	165.6	263.4	412.0	658.5	1,034.3	2,365.2
1999.....	14.1	26.8	53.5	101.6	157.4	276.5	424.2	690.6	1,057.9	2,435.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Atherosclerotic cardiovascular disease, so described (I25.0)										
All races 3/, both sexes										
2005.....	4.4	9.6	18.6	26.9	37.7	46.8	64.6	96.5	161.9	376.4
2004.....	4.8	9.5	18.2	26.6	39.1	49.8	69.0	100.8	166.6	400.4
2003.....	4.7	9.9	18.6	26.6	40.5	53.3	73.3	110.7	186.2	442.9
2002.....	4.8	9.8	17.5	27.9	39.7	53.9	76.5	116.6	187.6	474.5
2001.....	4.6	8.8	17.4	27.7	40.6	55.3	76.7	121.8	197.2	493.6
2000.....	4.4	9.4	17.5	28.5	40.7	55.7	81.9	125.0	207.8	534.6
1999.....	4.7	9.1	17.4	28.6	43.4	59.6	86.8	131.6	218.2	563.4
All races 3/, male										
2005.....	7.0	15.3	30.0	42.5	57.7	68.3	91.5	127.0	204.6	390.7
2004.....	7.7	15.3	29.6	42.0	58.4	73.1	96.4	133.8	210.5	414.0
2003.....	7.4	16.0	30.5	41.8	60.6	77.1	102.8	146.6	233.1	462.1
2002.....	7.8	15.6	28.4	44.8	59.7	78.8	106.7	154.5	232.4	491.8
2001.....	7.2	14.6	28.3	43.4	61.8	79.0	107.6	160.2	247.4	502.0
2000.....	7.1	15.5	28.4	45.8	61.3	80.8	115.6	167.1	252.8	550.8
1999.....	7.6	14.9	28.0	45.0	64.6	87.3	120.8	172.7	264.3	578.6
All races 3/, female										
2005.....	1.9	4.0	7.7	12.2	19.4	28.1	42.9	74.3	135.4	369.9
2004.....	2.0	3.9	7.4	12.0	21.5	29.5	47.0	77.0	139.6	394.3
2003.....	2.0	4.0	7.2	12.4	22.2	32.6	49.7	85.2	157.6	434.4
2002.....	1.8	4.1	7.1	12.1	21.4	32.4	52.5	89.7	160.4	467.1
2001.....	2.1	3.3	7.0	12.9	21.4	34.9	52.1	94.9	167.1	490.1
2000.....	1.8	3.5	7.2	12.4	22.1	34.1	55.4	95.7	181.2	528.0
1999.....	1.8	3.5	7.3	13.4	24.1	36.0	60.2	103.1	191.3	557.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Atherosclerotic cardiovascular disease, so described (I25.0)										
White, both sexes										
2005 4/.....	4.3	9.0	16.6	23.8	33.3	42.5	58.7	89.9	154.9	373.8
2004 4/.....	4.6	8.7	16.4	23.7	35.1	44.4	63.4	93.3	160.3	395.6
2003 4/.....	4.4	9.0	16.7	24.2	35.4	47.5	66.6	103.0	179.6	439.3
2002.....	4.4	9.1	15.6	24.7	35.3	47.3	69.7	107.8	180.5	470.6
2001.....	4.2	7.9	15.6	23.9	35.0	47.9	69.4	112.2	190.6	489.7
2000.....	4.0	8.3	15.7	24.8	35.2	49.5	73.7	116.0	198.5	529.5
1999.....	4.2	8.0	15.2	24.6	37.5	52.3	77.9	121.5	209.8	557.5
White, male										
2005 4/.....	7.0	14.6	27.2	38.2	51.2	62.2	82.8	118.8	194.8	392.6
2004 4/.....	7.5	14.4	27.3	37.8	52.4	66.1	89.3	123.9	203.3	408.8
2003 4/.....	6.9	14.8	27.7	38.3	53.4	69.2	92.5	136.9	224.4	458.7
2002.....	7.4	14.7	25.7	40.0	53.6	69.4	98.0	143.3	222.2	488.0
2001.....	6.6	13.3	25.7	38.3	54.0	69.3	97.6	148.7	239.9	498.6
2000.....	6.6	14.0	25.9	40.0	54.2	72.8	104.4	154.6	241.6	545.6
1999.....	7.0	13.2	24.8	39.4	56.2	77.6	109.0	159.8	253.9	571.2
White, female										
2005 4/.....	1.5	3.4	6.2	9.9	16.6	25.1	38.9	68.5	129.9	365.1
2004 4/.....	1.7	3.0	5.8	10.2	19.1	25.1	42.1	70.9	133.6	389.7
2003 4/.....	1.8	3.2	5.9	10.6	18.7	28.3	45.4	78.5	152.0	430.7
2002.....	1.5	3.5	5.6	10.2	18.4	27.8	46.6	82.5	154.9	463.0
2001.....	1.8	2.6	5.7	10.1	17.6	29.1	46.7	86.4	160.8	486.0
2000.....	1.4	2.7	5.8	10.2	17.6	29.0	49.0	88.8	172.9	522.9
1999.....	1.3	2.8	5.8	10.6	20.3	30.3	53.1	94.6	183.8	552.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Atherosclerotic cardiovascular disease, so described (I25.0)										
Black, both sexes										
2005 4/.....	6.5	15.5	37.3	56.6	83.4	92.4	130.4	183.1	265.1	469.6
2004 4/.....	7.5	17.5	35.7	55.9	81.5	105.1	139.0	198.5	268.4	505.1
2003 4/.....	7.3	17.4	35.9	53.2	91.8	111.4	149.2	214.1	282.8	543.2
2002.....	8.1	17.0	35.9	60.5	85.5	119.5	157.5	231.9	300.4	575.5
2001.....	8.2	16.8	35.7	64.2	94.4	128.0	162.6	244.0	301.5	594.5
2000.....	7.4	19.4	36.4	65.4	94.0	119.6	176.8	239.7	340.0	646.6
1999.....	9.1	19.1	38.4	67.8	98.2	133.9	188.9	261.7	338.9	668.6
Black, male										
2005 4/.....	8.9	23.1	58.1	87.6	130.3	140.2	200.8	251.7	361.7	450.8
2004 4/.....	10.7	25.1	54.0	88.3	127.8	153.1	197.1	278.0	350.8	531.5
2003 4/.....	11.4	26.4	57.8	81.3	140.0	161.3	228.9	301.3	380.2	583.9
2002.....	11.8	25.5	55.0	97.1	130.5	180.0	220.3	321.0	417.3	599.3
2001.....	12.4	25.7	56.0	97.0	144.0	183.5	241.0	330.4	388.5	609.3
2000.....	11.0	30.3	56.8	106.4	137.1	168.6	257.1	338.3	425.7	701.2
1999.....	13.7	29.9	59.7	103.8	147.9	197.3	271.0	350.0	435.1	715.9
Black, female										
2005 4/.....	4.3	8.9	19.7	31.1	46.8	56.9	82.2	142.1	214.0	477.1
2004 4/.....	4.7	10.9	20.2	29.3	45.3	69.4	99.5	151.5	224.9	494.6
2003 4/.....	3.7	9.6	17.3	30.0	54.2	74.4	95.5	162.2	231.7	527.4
2002.....	4.8	9.7	19.7	30.4	50.3	74.6	115.4	178.7	239.5	566.4
2001.....	4.5	9.0	18.4	37.3	55.4	87.1	110.3	192.5	257.1	589.0
2000.....	4.2	10.0	19.0	31.8	60.0	83.5	123.4	180.3	296.8	626.3
1999.....	5.0	9.8	20.4	38.4	59.2	87.0	134.3	208.5	291.3	650.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)										
All races 3/, both sexes										
2005.....	8.2	15.5	27.7	44.9	79.9	131.2	226.9	404.6	740.4	1,837.8
2004.....	8.4	16.4	27.4	45.7	79.9	134.8	236.6	417.4	769.0	1,879.3
2003.....	8.9	16.7	28.0	49.0	85.6	144.5	253.8	444.9	811.1	2,028.5
2002.....	8.8	17.1	29.9	52.3	88.2	152.9	261.8	465.6	841.9	2,088.0
2001.....	8.6	16.4	29.8	52.4	93.5	160.1	275.3	483.2	867.3	2,177.7
2000.....	9.0	17.0	30.3	56.7	95.9	165.3	287.4	496.2	894.4	2,287.5
1999.....	8.8	17.4	32.8	57.3	99.5	175.7	299.3	521.0	924.9	2,353.6
All races 3/, male										
2005.....	12.5	23.7	42.4	67.0	116.4	184.3	316.4	558.3	997.9	2,120.9
2004.....	12.7	25.0	41.8	67.5	116.8	191.3	328.2	580.4	1,027.4	2,129.6
2003.....	13.3	25.5	43.0	73.3	125.0	206.7	355.0	616.5	1,088.5	2,260.1
2002.....	13.6	26.6	46.5	78.0	129.6	217.8	367.6	647.1	1,123.8	2,325.6
2001.....	13.2	25.4	45.7	78.8	135.6	228.4	385.2	675.1	1,145.1	2,411.8
2000.....	14.0	27.0	46.7	84.9	139.5	234.0	409.1	684.5	1,204.8	2,561.9
1999.....	13.9	27.7	50.2	84.9	143.5	250.0	424.4	718.0	1,240.3	2,614.4
All races 3/, female										
2005.....	4.0	7.4	13.5	23.9	46.5	84.8	154.5	293.0	580.4	1,707.7
2004.....	4.1	8.0	13.6	25.0	46.3	85.6	162.8	300.3	609.8	1,766.7
2003.....	4.5	8.2	13.7	26.0	49.8	90.5	172.7	322.7	641.5	1,926.2
2002.....	4.2	7.8	13.9	28.2	50.5	97.0	177.3	337.3	670.4	1,984.9
2001.....	4.0	7.6	14.7	27.7	55.4	101.2	188.0	348.9	700.7	2,079.8
2000.....	4.0	7.3	14.6	30.3	56.4	106.3	191.6	365.1	711.3	2,175.8
1999.....	3.9	7.4	16.1	31.5	59.7	112.1	201.5	384.0	740.3	2,248.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)										
White, both sexes										
2005 4/.....	8.0	14.6	26.0	42.9	76.7	127.6	225.2	405.9	750.9	1,896.0
2004 4/.....	8.1	15.7	25.8	43.5	77.0	131.0	234.7	418.8	778.7	1,929.2
2003 4/.....	8.5	16.0	26.5	46.8	82.2	140.3	251.0	445.0	821.5	2,079.5
2002.....	8.6	16.1	28.3	49.9	84.5	150.0	260.7	466.0	854.2	2,143.3
2001.....	8.1	15.4	28.4	50.5	90.2	156.6	273.8	483.6	877.5	2,233.8
2000.....	8.5	16.0	29.1	54.4	92.2	162.0	285.8	496.4	908.0	2,352.7
1999.....	8.2	16.6	31.1	54.8	96.5	172.6	297.1	522.7	938.2	2,416.1
White, male										
2005 4/.....	12.3	22.5	40.4	65.0	113.3	181.1	317.3	564.5	1,021.0	2,204.7
2004 4/.....	12.4	24.3	40.0	65.5	114.0	188.4	328.6	585.8	1,050.7	2,200.2
2003 4/.....	12.9	24.7	41.5	71.3	122.4	203.4	354.1	622.0	1,109.8	2,338.4
2002.....	13.4	25.9	45.0	75.8	125.4	216.8	368.7	655.7	1,151.0	2,403.6
2001.....	12.8	24.4	44.2	76.5	133.1	226.8	385.4	682.4	1,166.2	2,493.6
2000.....	13.7	26.1	45.5	82.7	136.4	233.3	411.5	691.9	1,231.6	2,654.4
1999.....	13.1	27.2	48.1	82.6	140.3	249.9	425.1	727.7	1,269.0	2,695.9
White, female										
2005 4/.....	3.6	6.8	11.8	21.5	42.6	80.1	149.1	288.6	581.7	1,753.8
2004 4/.....	3.7	7.2	11.9	22.3	42.8	80.1	157.6	296.5	609.7	1,807.0
2003 4/.....	4.0	7.3	11.8	23.3	45.1	84.5	166.6	317.1	643.8	1,964.9
2002.....	3.6	6.5	12.0	25.2	46.7	91.1	172.7	330.1	672.3	2,030.2
2001.....	3.5	6.4	13.0	25.7	50.7	95.0	183.3	342.8	703.0	2,125.0
2000.....	3.3	5.9	13.0	27.5	51.4	99.6	184.9	358.9	715.6	2,230.1
1999.....	3.2	6.2	14.5	28.3	56.1	105.2	195.4	378.8	743.0	2,303.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)										
Black, both sexes										
2005 4/.....	11.4	24.0	45.5	72.4	125.2	184.3	284.0	466.7	738.4	1,415.4
2004 4/.....	12.8	23.9	45.2	73.2	123.4	193.3	298.8	473.1	755.5	1,514.0
2003 4/.....	12.8	25.4	46.4	78.5	132.8	206.3	331.2	515.6	804.7	1,652.1
2002.....	12.6	27.0	47.8	84.1	136.1	204.4	319.1	522.2	811.9	1,665.1
2001.....	13.3	26.9	47.9	79.9	140.9	218.1	337.8	546.0	853.7	1,756.7
2000.....	14.1	28.1	46.3	87.0	142.8	222.3	354.7	556.6	831.8	1,740.5
1999.....	14.7	26.3	53.8	90.6	143.1	233.9	362.6	555.9	844.6	1,801.2
Black, male										
2005 4/.....	15.8	36.4	66.7	101.3	172.1	247.4	368.4	604.8	909.5	1,420.4
2004 4/.....	18.1	34.0	65.5	99.6	172.3	258.0	395.9	623.3	907.5	1,536.8
2003 4/.....	18.0	36.5	64.7	108.6	177.6	278.2	437.5	669.6	1,003.3	1,563.9
2002.....	17.2	37.0	67.8	115.7	195.8	267.5	423.6	633.4	961.4	1,655.3
2001.....	18.8	38.3	69.2	113.1	189.2	286.4	457.8	692.3	1,055.4	1,759.8
2000.....	19.3	39.0	64.5	123.2	189.9	279.7	454.3	686.6	1,019.2	1,744.8
1999.....	21.2	37.0	78.2	124.1	200.3	293.9	472.2	678.6	1,001.9	1,845.5
Black, female										
2005 4/.....	7.6	13.2	27.6	48.5	88.6	137.5	226.2	384.3	648.0	1,413.4
2004 4/.....	8.1	15.1	28.0	51.4	85.2	145.2	232.8	384.1	675.5	1,505.0
2003 4/.....	8.3	15.7	30.9	53.8	97.9	153.0	259.4	424.0	700.6	1,686.4
2002.....	8.5	18.4	30.8	58.2	89.4	157.6	248.9	455.8	734.1	1,668.8
2001.....	8.5	17.0	29.8	52.6	102.9	167.6	257.8	458.7	751.0	1,755.5
2000.....	9.5	18.7	30.7	57.4	105.6	179.9	288.6	478.2	737.5	1,738.9
1999.....	9.1	17.1	33.1	63.3	98.2	189.5	289.9	482.0	766.6	1,784.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other heart diseases (I26-I51)										
All races 3/, both sexes										
2005.....	11.6	17.9	25.5	36.8	57.4	88.2	150.8	264.6	500.7	1,377.4
2004.....	11.5	17.9	25.6	35.7	57.5	90.3	151.8	268.1	503.3	1,373.9
2003.....	12.6	17.9	25.3	39.1	59.7	97.3	160.2	279.1	525.5	1,427.2
2002.....	12.4	18.1	25.9	39.4	60.6	97.9	164.5	286.7	536.3	1,443.5
2001.....	11.9	17.9	25.9	39.7	62.0	100.7	164.7	287.2	537.0	1,475.5
2000.....	12.3	17.6	25.9	40.2	64.3	103.6	167.1	290.2	542.8	1,509.4
1999.....	12.5	17.8	25.9	41.8	66.5	105.2	175.0	295.4	553.5	1,512.9
All races 3/, male										
2005.....	14.8	22.6	33.3	47.4	71.3	108.0	180.8	310.3	568.1	1,377.8
2004.....	14.4	23.2	32.8	46.4	72.7	111.3	181.2	316.2	573.7	1,377.7
2003.....	15.6	22.7	32.4	49.9	74.9	119.4	195.4	334.0	599.9	1,431.3
2002.....	15.9	23.5	33.6	49.4	75.7	118.8	199.8	340.7	612.4	1,448.6
2001.....	14.9	23.1	32.7	49.4	77.7	121.5	200.9	344.7	622.1	1,473.4
2000.....	15.4	23.0	33.4	52.0	80.8	126.9	205.9	345.1	626.8	1,549.1
1999.....	15.6	23.0	33.3	54.1	84.3	133.4	217.0	350.9	641.4	1,548.9
All races 3/, female										
2005.....	8.5	13.3	18.0	26.9	44.8	70.9	126.4	231.4	458.9	1,377.3
2004.....	8.6	12.8	18.6	25.7	43.8	72.0	128.2	233.5	459.9	1,372.1
2003.....	9.5	13.2	18.6	28.8	45.9	78.1	131.9	240.0	480.0	1,425.4
2002.....	9.0	12.8	18.5	30.1	46.8	79.8	136.3	248.6	490.0	1,441.3
2001.....	8.9	12.8	19.5	30.6	47.8	82.8	135.9	247.0	486.0	1,476.4
2000.....	9.2	12.3	18.7	29.2	49.3	83.6	136.6	252.1	493.2	1,493.2
1999.....	9.4	12.8	18.9	30.2	50.5	81.0	142.2	256.8	502.0	1,498.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other heart diseases (I26-I51)										
White, both sexes										
2005 4/.....	10.0	15.2	21.6	32.4	51.6	81.9	145.2	260.8	504.5	1,419.0
2004 4/.....	9.7	15.0	21.8	31.3	52.7	84.7	146.5	262.9	507.6	1,408.8
2003 4/.....	10.5	15.2	21.4	34.8	53.8	90.7	153.2	276.4	528.9	1,461.4
2002.....	10.7	15.0	22.1	35.0	55.3	92.4	158.5	282.4	539.1	1,475.5
2001.....	9.7	14.9	22.3	34.8	57.2	94.1	159.0	281.6	538.8	1,509.6
2000.....	10.2	14.6	22.1	35.9	59.0	97.3	160.6	284.5	545.7	1,541.6
1999.....	10.2	14.6	22.2	36.3	60.4	98.6	167.6	288.5	553.7	1,543.3
White, male										
2005 4/.....	12.8	19.6	28.7	41.9	64.7	100.4	173.5	306.2	575.6	1,424.2
2004 4/.....	12.4	19.6	28.7	40.9	66.8	104.6	176.0	309.6	578.2	1,407.9
2003 4/.....	13.5	19.7	27.5	44.5	68.3	111.8	188.4	332.0	604.3	1,467.6
2002.....	14.1	19.9	29.2	44.1	69.9	112.5	193.0	336.5	617.5	1,482.1
2001.....	12.7	19.7	28.9	43.5	72.1	114.9	193.7	338.2	621.9	1,503.4
2000.....	13.0	19.7	28.6	46.9	74.8	119.7	199.2	339.4	630.8	1,580.5
1999.....	13.3	19.4	28.7	47.5	76.9	125.6	209.0	344.0	641.4	1,571.5
White, female										
2005 4/.....	7.1	10.8	14.5	23.3	39.5	65.4	121.9	227.1	460.0	1,416.6
2004 4/.....	6.9	10.5	15.1	22.0	39.7	66.9	122.2	228.7	463.8	1,409.2
2003 4/.....	7.5	10.8	15.4	25.5	40.4	72.1	124.3	236.1	482.4	1,458.6
2002.....	7.3	10.2	15.2	26.2	41.8	74.7	130.3	243.7	491.0	1,472.6
2001.....	6.6	10.1	15.8	26.5	43.5	75.9	130.9	241.5	488.5	1,512.2
2000.....	7.4	9.6	15.8	25.4	44.3	77.7	129.5	245.9	495.2	1,525.8
1999.....	7.2	9.8	15.8	25.7	45.1	75.0	134.8	249.5	501.9	1,531.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other heart diseases (I26-I51)										
Black, both sexes										
2005 4/.....	25.3	40.3	60.2	82.0	120.4	162.1	237.3	368.3	563.4	1,135.3
2004 4/.....	26.3	41.7	60.2	82.6	114.5	159.9	240.5	381.0	551.3	1,180.7
2003 4/.....	28.9	40.8	61.1	85.3	124.6	176.1	265.5	364.8	574.9	1,238.1
2002.....	26.6	43.6	60.8	87.1	121.2	169.4	255.8	386.7	594.7	1,270.4
2001.....	29.2	43.1	60.9	91.4	118.8	179.7	254.5	402.9	601.8	1,277.7
2000.....	28.5	43.3	62.0	86.2	122.8	181.5	266.8	406.9	590.7	1,306.0
1999.....	30.2	44.9	62.2	96.8	133.5	185.5	284.2	417.4	621.2	1,302.8
Black, male										
2005 4/.....	31.8	50.1	77.6	107.6	151.1	204.4	300.1	446.6	614.5	1,088.5
2004 4/.....	31.6	54.3	75.0	109.0	149.1	202.5	284.5	469.5	653.5	1,224.9
2003 4/.....	34.7	51.0	79.5	113.5	158.3	223.6	318.0	434.1	668.1	1,220.9
2002.....	32.0	56.7	77.1	110.7	150.7	208.7	317.5	454.2	675.6	1,238.3
2001.....	33.7	53.5	73.7	117.2	151.5	212.9	318.8	494.8	743.2	1,349.1
2000.....	34.8	54.3	82.1	112.1	153.6	224.8	318.9	480.5	696.3	1,402.3
1999.....	36.2	56.1	81.9	127.0	172.7	237.8	353.1	489.1	747.3	1,454.1
Black, female										
2005 4/.....	19.6	31.7	45.4	60.8	96.5	130.8	194.4	321.6	536.4	1,154.2
2004 4/.....	21.6	30.6	47.7	60.9	87.6	128.3	210.7	328.6	497.4	1,163.2
2003 4/.....	23.8	31.9	45.4	62.1	98.2	140.9	230.2	323.7	526.1	1,244.8
2002.....	21.8	32.3	47.0	67.7	98.2	140.2	214.4	346.3	552.6	1,282.7
2001.....	25.3	34.0	50.1	70.3	93.1	155.1	211.6	348.1	529.8	1,251.2
2000.....	22.9	33.9	44.8	64.9	98.4	149.6	232.2	362.6	537.5	1,270.2
1999.....	24.9	35.2	45.4	72.1	102.8	146.8	238.5	374.2	558.7	1,246.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Acute and subacute endocarditis (I33)										
All races 3/, both sexes										
2005.....	0.2	0.4	0.5	0.6	0.7	1.1	1.4	2.4	2.7	3.0
2004.....	0.2	0.4	0.5	0.5	0.7	1.0	1.5	2.1	2.9	3.2
2003.....	0.3	0.4	0.5	0.6	0.9	1.0	1.5	2.3	2.7	3.0
2002.....	0.3	0.4	0.5	0.5	0.8	1.1	1.5	2.2	2.7	2.6
2001.....	0.3	0.4	0.5	0.5	0.8	1.1	1.7	2.2	3.0	2.5
2000.....	0.3	0.4	0.6	0.7	0.7	1.1	1.3	2.2	2.8	2.8
1999.....	0.3	0.3	0.4	0.7	0.7	1.1	1.5	2.1	2.7	2.9
All races 3/, male										
2005.....	0.2	0.4	0.7	0.8	0.9	1.2	1.8	3.1	3.5	5.1
2004.....	0.3	0.5	0.7	0.6	0.8	1.4	1.4	2.3	4.2	4.8
2003.....	0.4	0.6	0.7	0.8	1.1	1.3	1.6	2.9	3.7	4.1
2002.....	0.4	0.5	0.7	0.6	0.9	1.5	1.7	2.5	3.7	3.5
2001.....	0.3	0.7	0.7	0.6	1.1	1.1	2.1	3.1	4.3	3.8
2000.....	0.4	0.6	0.7	0.8	0.9	1.3	1.6	3.1	4.0	4.2
1999.....	0.4	0.5	0.7	1.0	0.8	1.2	1.7	2.8	3.9	4.7
All races 3/, female										
2005.....	*	0.3	0.3	0.4	0.6	1.0	1.1	1.9	2.2	2.1
2004.....	*	0.3	0.4	0.3	0.6	0.7	1.7	2.0	2.1	2.5
2003.....	0.2	0.3	0.3	0.4	0.8	0.9	1.4	1.9	2.1	2.5
2002.....	0.2	0.3	0.3	0.4	0.8	0.8	1.3	1.9	2.1	2.2
2001.....	0.3	0.2	0.3	0.5	0.5	1.0	1.3	1.6	2.2	2.0
2000.....	0.2	0.3	0.4	0.5	0.6	0.9	1.1	1.6	2.0	2.3
1999.....	*	*	0.2	0.5	0.7	1.0	1.4	1.7	2.0	2.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Acute and subacute endocarditis (I33)										
White, both sexes										
2005 4/.....	0.1	0.3	0.3	0.4	0.7	1.0	1.5	2.4	2.6	3.1
2004 4/.....	0.2	0.3	0.4	0.4	0.6	1.0	1.5	2.0	2.9	3.2
2003 4/.....	0.3	0.3	0.4	0.5	0.8	1.0	1.4	2.3	2.8	3.1
2002.....	0.2	0.3	0.4	0.4	0.8	1.0	1.4	2.2	2.9	2.7
2001.....	0.2	0.4	0.4	0.4	0.7	0.9	1.5	2.0	3.0	2.6
2000.....	0.3	0.3	0.4	0.5	0.7	1.1	1.3	2.2	2.7	2.7
1999.....	0.2	0.3	0.3	0.5	0.6	1.0	1.4	2.2	2.8	3.0
White, male										
2005 4/.....	*	0.3	0.5	0.6	0.8	1.2	1.8	3.2	3.5	5.3
2004 4/.....	0.3	0.4	0.5	0.6	0.7	1.3	1.4	2.2	4.3	4.9
2003 4/.....	0.3	0.4	0.5	0.6	1.0	1.2	1.5	2.8	3.9	4.4
2002.....	0.3	0.4	0.6	0.4	0.9	1.3	1.7	2.7	3.9	3.6
2001.....	0.3	0.5	0.5	0.5	0.9	0.9	1.8	3.0	4.3	3.9
2000.....	0.4	0.5	0.6	0.7	0.9	1.2	1.6	3.2	4.0	4.2
1999.....	0.3	0.4	0.5	0.8	0.7	1.2	1.6	3.0	4.0	5.1
White, female										
2005 4/.....	*	0.2	*	0.3	0.5	0.9	1.2	1.7	2.0	2.1
2004 4/.....	*	*	0.2	*	0.5	0.6	1.6	1.8	2.1	2.4
2003 4/.....	*	0.2	0.2	0.4	0.6	0.8	1.3	1.8	2.1	2.6
2002.....	*	*	*	0.3	0.7	0.7	1.2	1.8	2.3	2.3
2001.....	*	*	*	0.4	0.4	0.9	1.3	1.3	2.2	2.1
2000.....	*	*	0.3	0.4	0.5	1.0	1.0	1.6	2.0	2.1
1999.....	*	*	*	*	0.6	0.8	1.3	1.6	2.0	2.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Diseases of pericardium and acute myocarditis (I30-I31,I40)										
All races 3/, both sexes										
2005.....	0.2	0.3	0.3	0.3	0.4	0.7	0.8	1.1	1.3	1.9
2004.....	0.2	0.2	0.3	0.3	0.5	0.6	0.6	1.1	1.3	1.8
2003.....	0.2	0.3	0.3	0.4	0.4	0.8	0.8	0.9	1.5	1.6
2002.....	0.2	0.3	0.3	0.4	0.4	0.6	1.0	1.1	1.4	2.1
2001.....	0.2	0.3	0.3	0.4	0.5	0.5	0.8	1.2	1.3	1.9
2000.....	0.2	0.2	0.2	0.3	0.4	0.5	0.9	1.1	1.3	1.8
1999.....	0.3	0.3	0.3	0.5	0.6	0.7	0.9	1.2	1.5	2.0
All races 3/, male										
2005.....	0.3	0.4	0.4	0.3	0.5	0.7	1.0	1.4	1.5	1.9
2004.....	0.2	0.2	0.4	0.4	0.6	0.8	0.7	1.2	1.4	2.3
2003.....	0.3	0.4	0.3	0.4	0.4	1.0	1.0	1.1	1.5	1.5
2002.....	0.3	0.4	0.3	0.5	0.6	0.7	1.1	1.1	1.3	2.4
2001.....	*	0.3	0.3	0.4	0.6	0.6	0.8	1.3	1.1	1.8
2000.....	0.3	0.3	*	0.5	0.4	0.5	1.1	1.4	1.7	2.1
1999.....	0.3	0.3	0.3	0.6	0.6	0.8	0.9	1.3	1.5	2.5
All races 3/, female										
2005.....	*	0.3	0.3	0.3	0.4	0.6	0.6	0.9	1.1	1.9
2004.....	*	0.2	0.2	0.3	0.4	0.5	0.6	1.1	1.3	1.6
2003.....	0.2	0.2	*	0.3	0.3	0.7	0.7	0.8	1.5	1.7
2002.....	*	*	0.2	0.4	*	0.5	1.0	1.1	1.4	2.0
2001.....	0.3	0.3	0.3	0.4	*	*	0.9	1.1	1.4	2.0
2000.....	*	*	0.2	*	0.4	0.5	0.7	0.9	1.0	1.7
1999.....	0.3	0.3	0.2	0.3	0.6	0.7	0.8	1.2	1.5	1.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Diseases of pericardium and acute myocarditis (I30-I31,I40)										
White, both sexes										
2005 4/.....	0.2	0.3	0.3	0.3	0.4	0.7	0.8	1.1	1.2	1.9
2004 4/.....	0.1	0.2	0.3	0.3	0.4	0.6	0.6	1.0	1.3	1.8
2003 4/.....	0.2	0.3	0.2	0.4	0.4	0.8	0.8	1.0	1.5	1.7
2002.....	0.2	0.2	0.2	0.4	0.4	0.6	1.0	1.0	1.3	2.2
2001.....	0.2	0.2	0.2	0.3	0.5	0.4	0.9	1.2	1.3	1.9
2000.....	0.2	0.2	0.2	0.3	0.4	0.5	0.9	1.0	1.4	1.8
1999.....	0.2	0.3	0.2	0.5	0.6	0.7	0.8	1.2	1.5	2.0
White, male										
2005 4/.....	0.2	0.3	0.4	0.3	0.4	0.7	1.0	1.3	1.5	1.9
2004 4/.....	*	*	0.3	0.4	0.6	0.6	0.7	1.2	1.4	2.2
2003 4/.....	0.2	0.3	0.3	0.4	0.5	1.0	1.1	1.1	1.6	1.7
2002.....	0.3	0.3	0.3	0.4	0.6	0.6	1.1	1.1	1.2	2.5
2001.....	*	0.2	0.3	0.4	0.6	0.5	0.9	1.3	1.1	2.0
2000.....	0.2	*	*	0.5	*	*	1.1	1.2	1.9	2.1
1999.....	0.3	0.3	*	0.6	0.6	0.7	0.9	1.3	1.5	2.5
White, female										
2005 4/.....	*	*	0.3	*	0.4	0.6	0.6	0.9	1.1	1.8
2004 4/.....	*	*	*	0.3	*	0.5	0.6	0.9	1.3	1.6
2003 4/.....	*	*	*	0.4	*	0.7	0.6	0.8	1.4	1.7
2002.....	*	*	*	0.4	*	0.5	0.9	1.0	1.4	2.0
2001.....	*	0.3	*	0.3	*	*	0.9	1.2	1.4	1.9
2000.....	*	*	*	*	*	0.6	0.7	0.9	1.1	1.6
1999.....	*	0.3	*	*	0.5	0.7	0.8	1.1	1.4	1.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Heart failure (I50)										
All races 3/, both sexes										
2005.....	1.0	1.7	3.5	6.2	11.8	21.5	43.6	88.8	190.5	618.6
2004.....	0.9	1.7	3.3	6.1	11.4	22.4	43.8	88.7	187.7	621.2
2003.....	1.0	1.6	3.3	6.4	12.1	23.6	44.7	89.9	192.7	641.1
2002.....	0.8	1.7	3.4	6.0	12.4	22.9	45.7	89.0	194.0	645.0
2001.....	0.8	1.7	3.3	6.5	13.0	23.8	45.6	92.6	199.2	674.6
2000.....	0.8	1.4	3.0	6.1	12.8	24.0	46.5	93.5	199.9	684.9
1999.....	0.8	1.5	2.9	6.5	11.7	23.5	48.3	94.3	201.6	681.9
All races 3/, male										
2005.....	1.3	2.1	4.3	7.6	14.0	24.6	50.1	102.1	213.7	603.1
2004.....	1.0	2.2	4.0	7.5	13.7	26.4	51.1	103.2	211.1	605.5
2003.....	1.3	2.1	3.9	7.9	14.7	28.3	53.1	105.3	213.9	629.8
2002.....	1.0	2.0	4.4	7.2	14.8	27.0	53.2	103.9	218.0	621.4
2001.....	0.9	2.2	4.0	8.1	15.0	28.0	54.2	107.9	226.6	646.4
2000.....	1.1	1.9	3.5	7.4	15.3	28.2	55.6	107.9	229.5	672.2
1999.....	1.0	1.9	3.6	8.0	14.2	28.0	58.9	108.4	224.9	667.0
All races 3/, female										
2005.....	0.6	1.4	2.7	4.9	9.8	18.7	38.4	79.1	176.1	625.6
2004.....	0.7	1.3	2.8	4.8	9.4	18.8	37.9	78.3	173.3	628.2
2003.....	0.7	1.2	2.7	4.9	9.7	19.6	37.8	78.9	179.8	646.1
2002.....	0.5	1.3	2.4	4.9	10.1	19.3	39.8	78.4	179.3	655.3
2001.....	0.7	1.2	2.6	5.1	11.3	20.2	38.7	81.9	182.7	686.4
2000.....	0.6	0.9	2.5	5.0	10.5	20.3	39.3	83.5	182.4	690.0
1999.....	0.6	1.1	2.1	5.1	9.5	19.6	39.9	84.5	188.0	687.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Heart failure (I50)										
White, both sexes										
2005 4/.....	0.7	1.2	2.7	5.3	10.5	19.8	42.3	87.6	192.4	640.6
2004 4/.....	0.7	1.4	2.5	5.1	10.3	21.5	42.9	87.9	191.0	641.5
2003 4/.....	0.7	1.3	2.6	5.6	10.7	22.4	43.2	89.5	195.3	660.4
2002.....	0.6	1.2	2.7	5.3	11.4	21.7	44.1	88.1	195.6	664.5
2001.....	0.6	1.4	2.7	5.7	12.1	22.5	44.3	92.1	201.1	695.1
2000.....	0.6	1.1	2.5	5.4	11.8	22.6	44.9	92.8	201.8	703.7
1999.....	0.6	1.1	2.3	5.6	10.9	22.0	46.8	92.5	202.5	699.2
White, male										
2005 4/.....	0.8	1.5	3.4	6.5	12.5	22.7	48.4	100.6	217.9	629.3
2004 4/.....	0.8	1.6	3.1	6.2	12.2	25.2	50.7	101.7	215.6	625.5
2003 4/.....	0.9	1.7	3.0	7.0	13.1	27.0	52.1	105.9	217.1	650.8
2002.....	0.8	1.5	3.6	6.4	13.6	25.5	51.4	103.4	221.4	641.9
2001.....	0.7	1.9	3.4	7.2	14.2	26.5	52.8	107.6	228.2	666.5
2000.....	0.9	1.5	3.0	6.8	14.1	27.2	54.3	107.7	231.7	690.0
1999.....	0.8	1.4	2.9	7.1	13.3	26.2	58.1	107.2	226.3	680.0
White, female										
2005 4/.....	0.5	1.0	2.1	4.2	8.6	17.3	37.2	78.0	176.5	645.8
2004 4/.....	0.6	1.1	2.0	4.0	8.6	18.1	36.5	77.8	175.7	648.7
2003 4/.....	0.4	0.9	2.1	4.2	8.5	18.3	35.9	77.6	181.9	664.7
2002.....	0.4	0.9	1.9	4.2	9.3	18.4	38.2	77.1	179.8	674.3
2001.....	0.5	0.8	2.0	4.2	10.1	18.9	37.4	81.0	184.7	707.1
2000.....	0.4	0.6	2.0	4.1	9.6	18.6	37.3	82.4	183.9	709.3
1999.....	0.4	0.8	1.7	4.2	8.8	18.4	37.8	82.1	188.5	707.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Heart failure (I50)										
Black, both sexes										
2005 4/.....	3.0	5.2	10.0	15.6	26.8	41.3	69.1	123.8	216.0	489.6
2004 4/.....	2.2	5.0	10.5	16.3	23.8	35.7	64.0	120.2	194.6	506.5
2003 4/.....	3.3	4.7	9.4	15.3	27.7	41.1	70.9	114.7	203.6	531.8
2002.....	2.2	5.5	9.3	13.9	24.6	38.6	71.7	118.5	215.3	528.0
2001.....	2.7	4.5	8.7	15.7	24.6	41.2	71.1	121.1	213.7	543.8
2000.....	2.2	4.2	7.6	14.0	24.3	41.2	72.6	119.8	209.5	558.4
1999.....	2.4	4.8	8.4	16.4	22.0	42.0	74.0	130.5	216.7	552.9
Black, male										
2005 4/.....	4.6	6.4	12.6	20.1	32.7	48.5	82.5	152.0	224.0	423.0
2004 4/.....	2.6	6.8	12.2	21.0	30.6	45.5	69.3	148.3	220.2	503.7
2003 4/.....	3.9	5.5	11.6	19.4	35.9	51.0	79.5	124.6	227.4	513.0
2002.....	2.7	6.6	12.0	16.5	31.3	47.6	87.5	136.9	236.9	479.2
2001.....	2.6	5.0	10.4	18.9	26.4	49.6	86.2	140.8	256.8	541.2
2000.....	2.5	5.5	9.1	14.8	29.6	45.5	82.9	134.7	242.8	581.2
1999.....	2.7	6.4	11.0	19.3	27.4	51.4	83.1	146.7	238.9	603.5
Black, female										
2005 4/.....	1.6	4.0	7.7	11.9	22.1	36.0	60.0	106.9	211.8	516.5
2004 4/.....	1.9	3.4	8.9	12.4	18.5	28.4	60.4	103.5	181.1	507.6
2003 4/.....	2.8	4.0	7.6	11.9	21.4	33.8	65.1	108.8	191.1	539.2
2002.....	1.7	4.5	7.0	11.7	19.4	31.8	61.2	107.6	204.1	546.7
2001.....	2.8	4.1	7.2	13.1	23.2	35.1	61.0	109.3	191.7	544.8
2000.....	1.9	3.0	6.4	13.4	20.1	38.0	65.7	110.8	192.8	549.9
1999.....	2.1	3.5	6.3	14.0	17.8	35.0	68.0	120.7	205.7	534.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)										
All races 3/, both sexes										
2005.....	10.3	15.5	21.2	29.7	44.5	65.0	104.9	172.3	306.2	754.0
2004.....	10.2	15.5	21.4	28.8	45.0	66.3	105.9	176.1	311.3	747.7
2003.....	11.0	15.5	21.3	31.7	46.3	71.8	113.2	186.0	328.5	781.5
2002.....	11.2	15.7	21.7	32.5	46.9	73.3	116.3	194.4	338.2	793.7
2001.....	10.5	15.4	21.9	32.3	47.8	75.3	116.6	191.2	333.6	796.5
2000.....	10.9	15.5	22.1	33.1	50.4	78.0	118.4	193.4	338.9	820.0
1999.....	11.1	15.7	22.3	34.1	53.5	79.9	124.4	197.7	347.6	826.1
All races 3/, male										
2005.....	12.9	19.8	27.9	38.7	55.8	81.5	128.0	203.6	349.3	767.7
2004.....	12.9	20.2	27.8	37.9	57.6	82.7	128.0	209.5	357.0	765.2
2003.....	13.8	19.7	27.5	40.8	58.6	88.8	139.6	224.8	380.7	795.9
2002.....	14.2	20.6	28.1	41.1	59.4	89.7	143.8	233.1	389.4	821.3
2001.....	13.4	19.9	27.7	40.3	61.1	91.7	143.7	232.4	390.0	821.3
2000.....	13.6	20.2	28.9	43.4	64.2	96.8	147.6	232.6	391.5	870.7
1999.....	13.9	20.2	28.7	44.5	68.7	103.5	155.5	238.5	411.1	874.7
All races 3/, female										
2005.....	7.6	11.4	14.8	21.3	34.1	50.5	86.2	149.6	279.5	747.6
2004.....	7.6	11.0	15.3	20.3	33.4	52.0	88.0	152.2	283.2	739.9
2003.....	8.4	11.5	15.4	23.1	35.1	57.0	92.0	158.4	296.6	775.1
2002.....	8.2	11.0	15.6	24.4	35.6	59.2	94.3	167.1	307.1	781.7
2001.....	7.6	11.1	16.3	24.8	35.7	61.2	95.0	162.4	299.7	786.1
2000.....	8.2	11.0	15.6	23.4	37.8	61.9	95.4	166.0	307.9	799.3
1999.....	8.3	11.2	16.2	24.3	39.8	59.7	100.1	169.4	310.5	806.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)										
White, both sexes										
2005 4/.....	8.9	13.4	18.2	26.4	40.1	60.3	100.7	169.7	308.2	773.4
2004 4/.....	8.6	13.2	18.6	25.4	41.3	61.7	101.4	172.0	312.3	762.4
2003 4/.....	9.4	13.4	18.2	28.4	42.0	66.5	107.8	183.7	329.3	796.1
2002.....	9.7	13.3	18.8	28.9	42.7	69.1	111.9	191.2	339.3	806.1
2001.....	8.7	12.9	18.9	28.3	44.0	70.3	112.3	186.3	333.4	809.9
2000.....	9.1	13.0	18.9	29.6	46.2	73.1	113.5	188.4	339.9	833.4
1999.....	9.2	12.9	19.4	29.8	48.2	74.9	118.6	192.7	346.9	839.0
White, male										
2005 4/.....	11.5	17.3	24.4	34.5	51.0	75.8	122.3	201.1	352.7	787.7
2004 4/.....	11.2	17.4	24.7	33.7	53.3	77.5	123.2	204.5	356.9	775.3
2003 4/.....	12.0	17.3	23.6	36.5	53.7	82.6	133.8	222.2	381.7	810.7
2002.....	12.6	17.7	24.8	36.9	54.7	85.0	138.8	229.4	391.0	834.2
2001.....	11.5	17.0	24.7	35.4	56.5	86.9	138.2	226.4	388.3	831.0
2000.....	11.5	17.5	24.8	38.9	59.5	90.9	142.2	227.3	393.2	884.2
1999.....	11.9	17.3	25.1	39.0	62.4	97.5	148.4	232.5	409.5	883.9
White, female										
2005 4/.....	6.3	9.4	12.0	18.6	30.0	46.6	82.9	146.5	280.4	766.9
2004 4/.....	6.0	9.0	12.7	17.5	30.3	47.7	83.5	148.2	284.6	756.5
2003 4/.....	6.8	9.5	12.9	20.5	31.1	52.3	86.5	155.9	297.0	789.7
2002.....	6.7	9.0	13.0	21.4	31.5	55.1	90.0	163.8	307.6	793.9
2001.....	5.8	8.8	13.4	21.5	32.5	55.8	91.3	158.0	300.1	801.1
2000.....	6.7	8.6	13.2	20.7	33.9	57.6	90.4	161.1	308.1	812.7
1999.....	6.5	8.6	13.7	21.0	35.2	55.1	94.9	164.7	310.0	821.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)

Black, both sexes										
2005 4/.....	21.7	33.7	48.4	63.6	91.0	118.1	165.9	239.4	341.0	640.9
2004 4/.....	23.3	35.0	47.6	64.9	88.5	121.2	173.4	255.4	352.5	668.3
2003 4/.....	24.5	34.4	49.8	67.9	94.2	132.5	190.7	246.1	366.8	703.7
2002.....	23.6	36.5	49.5	70.6	94.5	127.7	180.3	263.4	376.5	738.5
2001.....	25.1	36.9	50.0	73.6	91.5	135.2	179.9	276.1	383.5	730.2
2000.....	25.2	37.3	52.5	69.7	96.5	137.5	191.0	282.9	377.8	742.0
1999.....	26.3	38.5	51.8	77.4	109.1	140.1	206.0	283.3	400.8	746.1
Black, male										
2005 4/.....	26.2	42.0	62.5	84.2	114.3	153.5	215.1	288.8	384.1	661.9
2004 4/.....	27.9	45.0	59.8	86.2	116.2	152.2	212.4	315.7	428.3	715.3
2003 4/.....	29.7	43.0	65.2	90.9	120.3	170.1	234.2	304.9	436.3	707.9
2002.....	28.2	47.7	62.8	90.7	117.2	157.1	226.3	314.2	434.1	753.7
2001.....	30.0	45.9	60.5	95.7	121.2	158.8	227.2	347.1	479.1	805.6
2000.....	30.9	46.3	71.0	95.0	122.0	174.3	233.4	340.7	449.3	816.5
1999.....	31.8	47.7	67.7	103.6	142.8	183.3	265.3	341.0	503.1	849.5
Black, female										
2005 4/.....	17.6	26.4	36.4	46.5	72.8	91.8	132.2	209.9	318.3	632.4
2004 4/.....	19.2	26.2	37.2	47.5	66.9	98.2	147.0	219.7	312.6	649.7
2003 4/.....	19.9	26.8	36.8	48.9	73.9	104.5	161.3	211.1	330.4	702.0
2002.....	19.5	26.8	38.3	54.1	76.8	105.8	149.4	233.0	346.5	732.8
2001.....	20.8	29.1	41.0	55.4	68.2	117.7	148.4	233.8	334.8	702.2
2000.....	20.3	29.5	36.8	49.0	76.5	110.4	162.9	248.1	341.8	714.3
1999.....	21.5	30.6	38.3	56.1	82.7	108.1	166.8	248.6	350.0	707.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Essential (primary) hypertension and hypertensive renal disease (I10,I12)										
All races 3/, both sexes										
2005.....	1.3	2.2	3.3	5.0	8.3	13.9	22.2	40.5	75.4	210.0
2004.....	1.0	2.2	3.4	4.9	8.1	13.0	21.9	38.5	71.3	198.5
2003.....	1.1	1.8	3.3	5.0	8.0	13.1	21.3	38.4	69.9	188.9
2002.....	1.0	1.8	2.9	4.5	7.3	11.8	20.7	34.9	66.7	180.4
2001.....	1.0	1.9	3.0	4.4	7.5	12.2	19.0	35.5	65.2	171.9
2000.....	1.1	1.8	3.0	4.4	7.8	11.5	19.1	33.8	63.0	162.9
1999.....	0.9	1.7	2.8	4.1	7.3	12.0	18.6	32.7	59.9	152.1
All races 3/, male										
2005.....	1.5	2.8	4.4	6.4	9.9	15.9	23.3	41.9	74.5	169.3
2004.....	1.2	2.6	4.5	5.6	9.8	14.9	22.2	40.0	72.7	166.0
2003.....	1.3	2.5	4.2	6.0	9.2	14.9	22.5	37.5	66.1	153.5
2002.....	1.3	2.3	3.5	5.5	8.2	12.7	22.2	35.9	64.8	153.4
2001.....	1.3	2.3	3.7	5.2	8.1	13.7	20.0	36.0	61.3	150.8
2000.....	1.4	2.3	3.8	5.4	9.1	11.5	20.6	34.5	62.9	142.4
1999.....	1.1	2.0	3.5	5.0	8.2	13.4	20.1	33.1	59.4	132.4
All races 3/, female										
2005.....	1.0	1.7	2.3	3.8	6.8	12.2	21.4	39.4	76.0	228.7
2004.....	0.7	1.7	2.3	4.3	6.6	11.3	21.6	37.4	70.4	213.2
2003.....	0.9	1.2	2.4	4.0	6.9	11.4	20.4	39.1	72.2	204.6
2002.....	0.7	1.3	2.3	3.7	6.5	11.0	19.5	34.2	67.9	192.1
2001.....	0.7	1.4	2.3	3.8	6.9	11.0	18.2	35.2	67.5	180.8
2000.....	0.8	1.3	2.2	3.5	6.7	11.5	17.9	33.4	63.0	171.3
1999.....	0.7	1.3	2.1	3.3	6.5	10.8	17.5	32.4	60.2	160.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Essential (primary) hypertension and hypertensive renal disease (I10,I12)

White, both sexes										
2005 4/.....	0.8	1.4	2.1	3.5	6.2	10.6	17.6	35.0	69.5	200.3
2004 4/.....	0.6	1.3	2.2	3.2	5.8	9.8	17.6	32.9	64.9	188.1
2003 4/.....	0.7	1.2	2.3	3.3	5.8	9.8	16.6	33.0	64.1	181.0
2002.....	0.6	1.0	1.9	3.0	5.4	9.0	16.8	29.8	60.4	170.4
2001.....	0.5	1.1	1.8	2.9	5.3	9.2	15.3	30.9	58.6	163.3
2000.....	0.6	1.0	1.8	3.0	5.5	8.5	15.3	28.9	57.4	153.2
1999.....	0.5	1.0	1.6	2.7	5.0	9.1	15.2	27.6	53.3	142.0
White, male										
2005 4/.....	1.0	1.8	2.8	4.5	7.5	12.2	18.1	35.3	68.9	160.2
2004 4/.....	0.8	1.7	3.0	3.8	7.3	11.5	18.2	33.7	66.6	159.4
2003 4/.....	0.8	1.6	2.9	4.1	6.6	11.6	18.0	32.7	60.3	145.3
2002.....	0.8	1.4	2.4	3.8	6.3	9.8	18.3	30.8	57.7	145.2
2001.....	0.7	1.4	2.2	3.5	5.9	10.3	16.4	31.4	54.5	142.2
2000.....	0.9	1.4	2.5	3.8	6.6	8.8	16.8	29.9	57.5	132.5
1999.....	0.7	1.3	2.1	3.5	5.8	10.4	16.6	28.1	52.5	121.0
White, female										
2005 4/.....	0.5	0.9	1.4	2.6	5.1	9.2	17.3	34.7	69.9	218.8
2004 4/.....	0.4	0.9	1.4	2.7	4.4	8.3	17.1	32.3	63.9	201.1
2003 4/.....	0.5	0.7	1.6	2.4	5.0	8.2	15.5	33.2	66.5	196.8
2002.....	0.4	0.7	1.4	2.3	4.5	8.3	15.6	29.0	62.0	181.3
2001.....	0.3	0.7	1.4	2.4	4.8	8.2	14.5	30.5	61.1	172.2
2000.....	0.4	0.6	1.1	2.2	4.5	8.3	14.1	28.2	57.4	161.6
1999.....	0.3	0.8	1.2	1.9	4.3	7.9	14.0	27.2	53.8	150.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Essential (primary) hypertension and hypertensive renal disease (I10,I12)

Black, both sexes

2005 4/.....	4.6	8.5	12.8	18.4	28.6	44.6	66.4	100.7	153.2	341.0
2004 4/.....	3.6	8.3	12.4	19.7	29.6	41.8	64.6	103.9	150.0	335.6
2003 4/.....	3.9	7.1	12.0	20.5	29.0	42.9	68.9	102.1	145.4	301.1
2002.....	3.8	7.4	10.9	18.4	24.3	36.5	61.5	96.4	151.0	308.8
2001.....	4.4	7.8	12.5	18.0	27.0	40.4	56.3	89.8	148.2	288.0
2000.....	4.2	7.2	12.7	17.9	29.0	40.3	58.8	92.4	130.4	283.4
1999.....	4.0	6.3	12.8	17.0	27.0	38.0	57.2	95.0	138.7	269.4

Black, male

2005 4/.....	5.2	9.8	17.4	24.4	36.5	52.8	79.1	123.8	154.5	289.1
2004 4/.....	4.2	9.8	16.6	22.7	36.3	49.9	66.0	123.7	157.8	263.3
2003 4/.....	4.5	9.8	16.4	24.3	36.2	50.0	74.6	102.2	147.7	274.2
2002.....	4.7	9.2	13.3	21.7	25.7	40.6	67.1	105.2	164.3	274.5
2001.....	5.5	10.1	16.3	21.7	30.3	48.8	61.9	95.2	158.1	275.7
2000.....	5.3	9.0	15.6	22.6	34.4	40.1	63.9	93.1	137.4	269.9
1999.....	4.6	7.8	16.6	19.5	31.0	43.1	62.1	98.8	155.2	266.0

Black, female

2005 4/.....	4.0	7.4	9.0	13.4	22.4	38.6	57.8	86.9	152.5	362.0
2004 4/.....	3.0	7.1	8.8	17.3	24.5	35.8	63.7	92.2	145.9	364.2
2003 4/.....	3.3	4.7	8.4	17.4	23.3	37.6	65.1	102.0	144.3	311.5
2002.....	3.0	5.9	8.8	15.6	23.2	33.5	57.8	91.1	144.0	322.0
2001.....	3.4	5.8	9.3	15.0	24.5	34.2	52.5	86.5	143.1	292.6
2000.....	3.2	5.7	10.3	14.2	24.7	40.5	55.5	92.0	126.9	288.5
1999.....	3.5	5.0	9.5	15.0	24.0	34.2	54.0	92.7	130.5	270.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Cerebrovascular diseases (I60-I69)										
All races 3/, both sexes										
2005.....	6.9	11.7	18.8	25.8	42.7	73.8	133.6	253.6	497.3	1,141.8
2004.....	7.1	11.9	18.3	26.5	44.5	78.4	142.2	276.7	532.1	1,245.9
2003.....	7.4	11.8	18.6	27.8	45.9	79.7	150.7	294.1	571.1	1,370.1
2002.....	7.1	12.3	18.3	28.6	48.3	83.8	160.7	308.8	601.7	1,445.9
2001.....	7.2	12.1	18.5	28.8	49.6	85.7	164.3	320.2	622.4	1,500.2
2000.....	7.7	13.1	19.3	30.8	53.8	91.0	169.1	332.8	654.0	1,589.2
1999.....	7.5	12.5	18.4	31.7	51.4	91.1	173.5	340.6	664.0	1,614.8
All races 3/, male										
2005.....	7.0	12.1	21.5	30.1	50.0	84.6	149.5	275.8	513.0	1,023.3
2004.....	7.4	12.8	21.1	29.9	52.2	90.2	158.8	301.2	551.6	1,118.1
2003.....	7.6	12.9	21.1	31.3	53.5	91.4	170.8	322.5	595.6	1,236.0
2002.....	6.9	13.4	20.4	32.7	55.8	97.0	178.7	334.8	615.6	1,317.9
2001.....	7.4	13.2	20.5	32.8	57.2	98.5	187.9	349.0	654.6	1,380.2
2000.....	7.8	14.2	21.3	35.2	62.3	102.6	192.9	367.6	695.1	1,484.3
1999.....	7.8	13.3	20.7	35.7	60.0	102.8	197.7	376.9	706.8	1,512.8
All races 3/, female										
2005.....	6.7	11.3	16.2	21.8	36.0	64.3	120.7	237.6	487.5	1,196.1
2004.....	6.8	11.0	15.5	23.2	37.5	68.1	128.8	259.1	520.2	1,303.4
2003.....	7.2	10.8	16.3	24.6	38.9	69.5	134.5	273.9	556.1	1,429.4
2002.....	7.2	11.2	16.3	24.8	41.4	72.4	146.2	290.4	593.3	1,501.5
2001.....	7.1	10.9	16.5	25.1	42.7	74.7	145.6	300.0	603.2	1,550.4
2000.....	7.5	12.0	17.3	26.6	46.0	81.0	150.3	308.5	629.7	1,632.0
1999.....	7.1	11.7	16.1	28.0	43.7	81.1	154.5	315.3	639.0	1,656.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Cerebrovascular diseases (I60-I69)										
White, both sexes										
2005 4/.....	5.3	9.1	14.5	21.1	36.2	65.6	123.8	244.2	491.2	1,153.3
2004 4/.....	5.7	9.1	14.1	21.8	37.5	70.0	132.6	264.8	526.2	1,257.8
2003 4/.....	5.6	9.0	14.2	22.6	38.3	70.8	140.3	283.2	565.9	1,382.2
2002.....	5.5	9.5	14.0	23.8	41.2	74.2	150.1	298.6	595.7	1,459.5
2001.....	5.6	9.3	13.8	23.9	41.7	75.2	153.2	309.7	617.1	1,511.2
2000.....	5.9	9.7	15.4	25.4	46.5	81.9	159.0	321.3	647.0	1,601.6
1999.....	5.7	9.4	14.4	26.2	43.5	82.9	161.3	329.3	658.4	1,631.6
White, male										
2005 4/.....	5.5	9.5	16.5	24.2	41.8	74.3	138.0	266.4	505.3	1,033.7
2004 4/.....	6.1	9.7	16.3	24.1	43.3	80.6	147.0	288.0	546.9	1,129.3
2003 4/.....	5.9	10.1	16.0	24.7	44.5	81.8	158.5	310.0	590.4	1,247.0
2002.....	5.5	10.6	15.5	26.9	47.2	85.1	167.3	323.4	610.1	1,327.4
2001.....	5.9	10.2	15.4	26.9	48.1	86.7	174.8	337.7	648.8	1,386.2
2000.....	6.1	10.8	16.9	28.6	53.8	91.7	180.5	354.6	686.1	1,490.7
1999.....	6.1	9.8	16.2	29.5	50.6	93.2	183.6	363.2	700.5	1,522.7
White, female										
2005 4/.....	5.2	8.6	12.5	18.2	31.1	57.8	112.1	227.7	482.4	1,208.5
2004 4/.....	5.2	8.5	11.9	19.5	32.2	60.5	120.7	247.9	513.3	1,315.7
2003 4/.....	5.3	7.9	12.3	20.5	32.5	61.1	125.4	263.8	550.7	1,442.1
2002.....	5.5	8.4	12.6	20.9	35.6	64.5	136.0	280.8	586.8	1,516.9
2001.....	5.4	8.3	12.2	21.1	35.8	65.1	135.7	289.8	598.0	1,563.5
2000.....	5.7	8.7	13.9	22.4	39.8	73.3	141.9	297.9	623.8	1,646.7
1999.....	5.3	8.9	12.7	23.0	37.0	73.9	143.7	305.5	633.6	1,675.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Cerebrovascular diseases (I60-I69)

Black, both sexes

2005 4/.....	17.1	30.5	50.3	65.4	99.5	145.1	231.4	372.3	611.0	1,106.7
2004 4/.....	17.2	31.5	48.0	66.3	108.1	154.9	246.5	418.6	647.9	1,211.2
2003 4/.....	18.6	31.4	52.7	72.4	113.1	157.4	256.3	433.2	677.0	1,348.7
2002.....	17.4	32.1	51.2	71.1	109.9	169.6	272.5	447.2	712.5	1,383.9
2001.....	18.4	32.6	55.0	70.6	117.1	178.3	277.6	460.2	716.4	1,452.7
2000.....	19.7	37.2	51.2	77.4	114.0	174.8	281.1	475.2	769.4	1,530.0
1999.....	19.9	34.4	50.6	78.4	118.2	170.0	307.8	487.9	756.4	1,492.5

Black, male

2005 4/.....	17.4	32.0	60.5	83.6	133.4	184.9	276.7	407.2	655.2	983.5
2004 4/.....	16.9	35.4	56.4	84.7	139.4	188.3	297.8	470.9	677.2	1,061.0
2003 4/.....	18.8	34.2	62.6	90.6	141.7	184.9	307.4	503.0	727.9	1,180.3
2002.....	16.9	34.3	61.6	88.9	138.8	213.7	328.8	502.5	729.8	1,252.2
2001.....	17.8	36.5	63.9	87.6	143.4	218.5	336.3	524.6	768.1	1,349.1
2000.....	20.5	41.0	60.4	97.8	137.5	212.8	340.0	544.8	863.2	1,458.8
1999.....	20.8	38.8	61.3	92.7	149.4	204.3	366.7	559.0	818.8	1,448.3

Black, female

2005 4/.....	16.8	29.3	41.8	50.3	73.0	115.6	200.4	351.5	587.6	1,156.5
2004 4/.....	17.5	28.1	40.8	51.2	83.6	130.1	211.8	387.6	632.4	1,270.7
2003 4/.....	18.5	29.0	44.4	57.3	90.7	137.1	221.8	391.8	650.3	1,414.2
2002.....	17.9	30.1	42.4	56.5	87.3	136.9	234.8	414.1	703.4	1,434.3
2001.....	19.0	29.1	47.4	56.7	96.5	148.5	238.6	421.9	690.1	1,491.2
2000.....	19.0	33.9	43.4	60.6	95.4	146.7	242.0	433.2	722.2	1,556.5
1999.....	19.0	30.5	41.6	66.6	93.7	144.5	268.7	445.1	725.6	1,509.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Atherosclerosis (I70)										
All races 3/, both sexes										
2005.....	0.1	0.3	0.7	1.0	2.1	3.9	7.2	16.9	37.8	132.9
2004.....	0.1	0.3	0.5	1.0	2.3	4.1	8.5	17.5	38.0	137.0
2003.....	0.2	0.3	0.6	1.1	2.4	4.5	9.0	19.9	43.5	154.6
2002.....	0.1	0.3	0.6	1.4	2.4	5.2	9.3	20.5	47.8	168.1
2001.....	0.1	0.3	0.7	1.4	2.5	4.9	9.8	21.9	49.2	179.1
2000.....	0.1	0.4	0.6	1.3	2.6	5.2	10.5	22.7	52.4	188.8
1999.....	0.1	0.3	0.8	1.3	2.9	5.1	11.2	24.1	53.6	202.0
All races 3/, male										
2005.....	*	0.4	1.0	1.2	2.7	4.8	8.7	20.0	42.0	121.7
2004.....	*	0.4	0.7	1.2	3.0	4.9	10.3	20.2	43.5	128.2
2003.....	0.2	0.5	0.8	1.5	3.2	5.4	10.6	23.9	48.3	138.6
2002.....	0.2	0.4	0.8	1.7	3.2	6.4	11.5	24.4	52.5	156.6
2001.....	*	0.4	0.8	1.7	3.2	6.2	12.1	25.0	53.2	167.2
2000.....	0.2	0.5	0.8	1.7	3.6	6.6	13.1	27.5	59.8	176.5
1999.....	0.2	0.4	1.0	1.8	3.8	6.3	13.3	28.1	60.0	189.0
All races 3/, female										
2005.....	*	0.2	0.4	0.8	1.5	3.1	5.9	14.7	35.2	138.1
2004.....	*	*	0.3	0.8	1.7	3.4	7.1	15.5	34.6	141.0
2003.....	*	*	0.4	0.7	1.7	3.7	7.8	17.0	40.6	161.7
2002.....	*	*	0.5	1.0	1.7	4.1	7.5	17.8	45.0	173.1
2001.....	*	0.2	0.5	1.1	1.8	3.9	8.1	19.7	46.8	184.1
2000.....	*	0.2	0.4	0.9	1.7	4.0	8.4	19.4	48.1	193.8
1999.....	*	*	0.5	0.9	2.2	4.0	9.5	21.4	49.8	207.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Atherosclerosis (I70)										
White, both sexes										
2005 4/.....	0.1	0.3	0.6	0.9	2.0	3.8	7.1	17.2	38.8	138.7
2004 4/.....	0.1	0.3	0.4	1.0	2.3	4.0	8.5	17.8	39.3	142.0
2003 4/.....	0.2	0.3	0.6	1.0	2.4	4.4	8.9	20.2	44.1	159.5
2002.....	0.1	0.2	0.5	1.3	2.3	5.3	9.1	20.4	48.4	174.7
2001.....	0.1	0.3	0.6	1.4	2.3	4.9	9.8	22.2	50.3	185.9
2000.....	0.1	0.3	0.5	1.2	2.4	5.2	10.4	23.1	53.0	195.2
1999.....	*	0.3	0.7	1.2	2.9	5.0	11.3	24.2	54.7	207.7
White, male										
2005 4/.....	*	0.4	0.9	1.2	2.6	4.8	8.5	20.6	42.9	128.3
2004 4/.....	*	0.4	0.7	1.2	2.9	4.9	10.4	20.5	44.8	134.4
2003 4/.....	0.2	0.5	0.8	1.4	3.1	5.2	10.1	23.8	48.4	143.3
2002.....	*	0.4	0.6	1.7	3.0	6.5	11.3	24.0	53.1	163.2
2001.....	*	0.4	0.8	1.7	3.0	6.2	12.1	25.2	54.0	173.9
2000.....	*	0.4	0.7	1.7	3.5	6.5	13.2	28.2	60.3	181.4
1999.....	*	0.4	1.0	1.6	3.7	6.2	13.3	27.7	60.8	193.7
White, female										
2005 4/.....	*	*	0.3	0.7	1.5	3.0	5.9	14.7	36.2	143.5
2004 4/.....	*	*	0.2	0.8	1.7	3.2	7.0	15.9	35.9	145.4
2003 4/.....	*	*	0.4	0.7	1.7	3.6	7.8	17.6	41.4	166.7
2002.....	*	*	0.4	1.0	1.6	4.2	7.3	17.7	45.5	179.7
2001.....	*	*	0.5	1.0	1.6	3.9	8.0	20.2	48.0	190.9
2000.....	*	*	0.3	0.8	1.5	4.1	8.2	19.5	48.7	200.8
1999.....	*	*	0.5	0.9	2.0	3.9	9.7	21.7	51.0	213.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Atherosclerosis (I70)										
Black, both sexes										
2005 4/.....	*	*	1.3	1.3	3.1	5.8	9.5	16.5	33.7	97.8
2004 4/.....	*	*	*	1.7	3.2	5.4	9.9	18.4	31.9	102.9
2003 4/.....	*	*	1.3	1.9	3.5	6.5	12.7	20.8	45.4	121.1
2002.....	*	*	1.5	1.9	4.2	5.2	11.5	26.8	51.3	117.0
2001.....	*	*	1.0	2.0	5.0	6.3	12.1	22.9	42.8	125.6
2000.....	*	*	1.5	2.1	4.7	5.6	12.9	23.5	53.9	138.3
1999.....	*	*	1.3	2.1	4.7	7.0	11.9	27.5	46.8	154.3
Black, male										
2005 4/.....	*	*	*	*	4.4	7.6	12.3	17.0	40.0	75.3
2004 4/.....	*	*	*	*	4.9	6.1	12.0	23.2	40.9	78.5
2003 4/.....	*	*	*	*	4.8	8.1	18.1	28.8	59.1	104.3
2002.....	*	*	2.3	*	6.3	6.2	15.4	34.9	59.6	110.4
2001.....	*	*	*	*	5.9	7.6	15.1	30.2	49.0	113.5
2000.....	*	*	*	*	6.0	8.6	14.0	25.5	66.6	147.6
1999.....	*	*	*	3.4	5.7	8.8	15.6	37.0	56.7	164.0
Black, female										
2005 4/.....	*	*	*	*	*	4.4	7.6	16.3	30.4	106.9
2004 4/.....	*	*	*	*	*	4.9	8.6	15.6	27.2	112.6
2003 4/.....	*	*	*	*	*	5.3	9.1	16.0	38.3	127.6
2002.....	*	*	*	*	*	4.4	9.0	22.0	46.9	119.5
2001.....	*	*	*	*	4.3	5.3	10.1	18.6	39.7	130.1
2000.....	*	*	*	*	3.6	*	12.2	22.2	47.6	134.8
1999.....	*	*	*	*	3.8	5.6	9.5	21.7	41.9	150.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other diseases of circulatory system (I71-I78)										
All races 3/, both sexes										
2005.....	1.3	2.0	3.2	5.4	10.2	18.0	31.2	50.9	77.9	135.5
2004.....	1.2	2.1	3.4	5.2	9.9	18.3	33.0	51.7	81.1	139.3
2003.....	1.2	2.4	3.5	5.7	11.3	20.5	34.2	56.6	82.7	146.0
2002.....	1.3	2.2	3.1	6.2	11.7	22.3	36.5	57.8	83.8	148.4
2001.....	1.3	2.0	3.3	6.0	12.1	22.3	36.6	60.6	88.1	158.2
2000.....	1.3	2.1	3.5	6.4	12.7	24.5	40.6	62.9	92.2	163.2
1999.....	1.4	2.1	3.3	6.7	12.5	23.5	40.1	63.5	94.3	165.2
All races 3/, male										
2005.....	1.8	2.9	4.4	7.8	14.2	24.8	42.7	68.4	101.2	158.2
2004.....	1.7	3.0	4.8	7.2	13.6	24.7	44.1	67.1	105.8	166.5
2003.....	1.7	3.2	5.1	7.9	15.9	29.4	46.1	78.9	111.3	179.7
2002.....	1.8	3.2	4.4	8.6	16.3	31.7	49.2	78.3	113.7	187.4
2001.....	1.8	2.8	4.4	8.2	16.8	30.8	51.2	81.0	122.5	197.3
2000.....	1.7	2.8	4.9	9.3	18.0	34.6	57.6	88.6	126.6	214.9
1999.....	1.6	3.0	4.4	9.4	18.2	33.5	56.7	89.1	129.3	207.5
All races 3/, female										
2005.....	0.9	1.1	2.0	3.2	6.6	12.1	21.9	38.2	63.4	125.0
2004.....	0.8	1.2	2.1	3.3	6.6	12.8	24.0	40.6	65.9	127.0
2003.....	0.8	1.6	2.1	3.7	7.2	12.8	24.6	40.7	65.2	131.1
2002.....	0.7	1.4	2.0	3.9	7.5	14.2	26.4	43.4	65.6	131.5
2001.....	0.9	1.2	2.2	3.9	7.8	15.0	25.1	46.3	67.4	141.8
2000.....	0.9	1.5	2.1	3.7	7.9	15.7	27.2	45.0	71.9	142.2
1999.....	1.1	1.3	2.1	4.1	7.3	15.0	27.2	45.7	73.7	148.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other diseases of circulatory system (I71-I78)

White, both sexes

2005 4/.....	1.2	1.7	2.8	5.0	10.0	17.7	31.6	51.1	77.9	136.1
2004 4/.....	1.1	1.8	3.0	4.8	9.3	18.1	33.8	52.1	81.2	139.3
2003 4/.....	1.1	2.0	3.1	5.5	11.0	20.6	34.7	57.5	82.9	146.9
2002.....	1.1	1.9	2.7	5.8	11.4	22.4	37.0	58.7	84.8	148.9
2001.....	1.1	1.8	2.9	5.6	11.7	22.6	36.8	60.6	88.4	158.2
2000.....	1.0	1.8	3.0	6.0	12.3	24.5	41.4	63.0	93.0	163.0
1999.....	1.2	1.9	3.0	6.3	12.6	23.7	40.8	64.1	94.8	165.1

White, male

2005 4/.....	1.5	2.4	3.8	7.2	14.1	24.3	43.9	68.5	102.1	162.0
2004 4/.....	1.5	2.6	4.3	6.6	12.8	24.5	45.7	68.2	107.5	169.3
2003 4/.....	1.5	2.8	4.6	7.7	15.4	29.7	47.4	80.5	111.9	184.1
2002.....	1.6	2.7	3.8	7.9	16.0	32.5	49.8	80.4	116.1	190.9
2001.....	1.4	2.5	3.8	7.9	16.4	31.5	51.7	82.0	124.0	201.3
2000.....	1.3	2.5	4.4	8.9	17.8	35.5	59.1	89.4	129.7	219.6
1999.....	1.5	2.7	4.0	9.2	18.6	34.4	58.4	91.1	131.1	211.1

White, female

2005 4/.....	0.8	0.9	1.8	2.9	6.2	11.8	21.3	38.3	62.7	124.2
2004 4/.....	0.7	1.1	1.8	3.1	6.1	12.4	23.9	40.4	64.8	125.7
2003 4/.....	0.7	1.3	1.7	3.4	6.9	12.6	24.4	40.9	65.0	130.5
2002.....	0.6	1.0	1.7	3.7	7.2	13.5	26.5	43.1	65.6	130.7
2001.....	0.7	1.1	2.0	3.4	7.4	14.7	24.7	45.5	67.0	140.1
2000.....	0.7	1.1	1.8	3.3	7.2	15.0	27.1	44.5	71.2	140.1
1999.....	0.9	1.0	1.9	3.6	7.1	14.5	26.9	45.1	73.4	146.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other diseases of circulatory system (I71-I78)

Black, both sexes

2005 4/.....	2.6	4.3	6.7	10.4	14.7	24.5	35.2	57.0	88.4	144.6
2004 4/.....	2.6	4.4	6.4	9.4	17.4	24.3	32.7	56.9	90.1	155.1
2003 4/.....	2.6	5.3	7.3	8.5	16.3	24.0	35.9	55.3	92.1	150.0
2002.....	2.8	5.3	6.7	11.2	16.5	25.7	37.8	58.4	83.5	156.8
2001.....	3.2	4.0	6.5	10.7	18.0	22.9	42.8	69.3	96.2	174.7
2000.....	3.2	4.5	6.8	11.2	18.3	27.3	39.9	68.9	92.3	177.7
1999.....	2.8	4.5	6.1	11.4	13.5	26.1	38.6	64.3	96.5	178.9

Black, male

2005 4/.....	3.7	6.7	9.5	15.1	19.3	33.7	39.5	79.1	103.9	129.2
2004 4/.....	3.6	6.3	8.9	13.7	24.3	33.4	37.6	68.2	98.3	158.0
2003 4/.....	3.5	6.8	9.9	11.2	23.5	33.5	40.1	72.6	118.9	138.7
2002.....	4.1	6.7	9.8	16.5	23.0	30.7	48.7	65.7	100.1	166.2
2001.....	4.5	5.5	9.4	13.1	23.8	28.0	55.9	79.0	127.6	165.7
2000.....	4.2	5.5	10.0	15.1	22.6	32.8	51.8	90.7	104.5	177.6
1999.....	3.1	6.1	8.6	14.1	17.8	31.9	49.2	74.6	117.7	184.0

Black, female

2005 4/.....	1.5	2.3	4.4	6.5	11.2	17.6	32.2	43.7	80.1	150.8
2004 4/.....	1.8	2.7	4.2	5.8	11.9	17.4	29.4	50.1	85.8	154.0
2003 4/.....	1.8	4.0	5.1	6.2	10.7	17.0	33.0	45.0	78.1	154.3
2002.....	1.7	4.0	4.0	6.8	11.5	22.0	30.5	54.1	74.8	153.2
2001.....	2.0	2.7	4.1	8.7	13.4	19.1	34.1	63.5	80.1	178.1
2000.....	2.3	3.7	4.1	8.0	14.9	23.2	32.0	55.8	86.2	177.7
1999.....	2.5	3.2	4.0	9.1	10.1	21.7	31.5	58.1	86.0	177.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Aortic aneurysm and dissection (I71)										
All races 3/, both sexes										
2005.....	0.9	1.4	2.2	3.3	6.8	11.7	19.8	31.6	46.6	64.9
2004.....	0.8	1.4	2.2	3.1	6.4	11.4	21.1	32.6	48.8	64.8
2003.....	0.8	1.7	2.4	3.7	7.3	13.5	22.7	36.6	50.7	70.0
2002.....	0.9	1.5	2.0	4.0	7.4	14.7	23.9	37.5	51.5	69.4
2001.....	0.9	1.4	2.0	3.7	7.8	15.0	23.8	39.0	55.4	77.5
2000.....	0.9	1.3	2.2	4.3	8.4	16.3	27.8	41.0	56.5	79.2
1999.....	0.9	1.4	2.1	4.4	8.1	16.5	27.3	42.2	58.2	80.3
All races 3/, male										
2005.....	1.5	2.3	3.3	5.3	10.0	17.1	29.0	43.8	65.4	91.0
2004.....	1.3	2.1	3.3	4.9	9.7	17.0	30.6	44.2	70.3	95.0
2003.....	1.3	2.5	3.7	5.8	11.4	20.2	33.0	53.8	74.2	105.5
2002.....	1.4	2.4	3.2	6.3	11.5	23.0	34.9	55.0	74.5	106.6
2001.....	1.4	2.1	3.2	5.8	12.0	22.2	36.0	56.3	82.5	114.8
2000.....	1.3	1.9	3.5	6.7	13.1	24.9	41.8	61.9	84.9	127.3
1999.....	1.2	2.2	3.3	6.8	13.0	25.3	41.0	63.3	88.6	125.3
All races 3/, female										
2005.....	0.4	0.6	1.1	1.5	3.8	7.0	12.4	22.7	35.0	52.9
2004.....	0.4	0.7	1.0	1.5	3.3	6.6	13.5	24.3	35.6	51.2
2003.....	0.4	0.9	1.1	1.7	3.6	7.8	14.5	24.3	36.3	54.2
2002.....	0.3	0.7	1.0	1.8	3.7	7.5	15.1	25.1	37.6	53.3
2001.....	0.3	0.6	0.9	1.8	3.9	8.8	14.0	26.9	39.2	61.9
2000.....	0.5	0.8	1.0	2.1	4.1	8.9	16.7	26.5	39.7	59.6
1999.....	0.5	0.7	1.0	2.1	3.7	9.0	16.6	27.5	40.4	62.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Aortic aneurysm and dissection (I71)										
White, both sexes										
2005 4/.....	0.8	1.2	1.8	3.2	6.7	11.7	20.6	32.5	47.6	66.2
2004 4/.....	0.7	1.2	1.9	2.8	6.1	11.7	22.1	33.7	49.8	66.8
2003 4/.....	0.7	1.4	2.1	3.6	7.3	14.0	23.6	38.1	52.2	72.4
2002.....	0.7	1.2	1.7	3.8	7.5	15.3	24.8	39.0	53.1	71.2
2001.....	0.7	1.2	1.7	3.6	7.7	15.5	24.6	39.8	57.0	79.7
2000.....	0.7	1.2	2.0	4.1	8.4	16.7	29.0	42.0	58.1	80.8
1999.....	0.7	1.2	1.9	4.2	8.5	17.1	28.4	43.5	60.0	82.2
White, male										
2005 4/.....	1.2	1.9	2.8	5.1	10.0	17.2	30.6	45.1	67.1	94.2
2004 4/.....	1.1	1.8	3.0	4.5	9.3	17.4	32.1	45.8	72.1	98.7
2003 4/.....	1.2	2.1	3.4	5.7	11.4	21.1	34.6	56.2	76.5	110.0
2002.....	1.2	2.0	2.7	5.9	11.6	24.3	36.1	57.5	77.7	111.8
2001.....	1.1	1.9	2.8	5.6	11.9	23.2	37.3	57.7	85.5	120.3
2000.....	1.0	1.8	3.2	6.5	13.3	26.1	43.8	63.6	88.6	132.8
1999.....	1.0	1.9	2.9	6.7	13.6	26.5	43.0	66.3	91.5	129.8
White, female										
2005 4/.....	0.3	0.4	0.9	1.4	3.5	6.9	12.3	23.1	35.3	53.3
2004 4/.....	0.3	0.5	0.9	1.2	3.2	6.7	13.9	24.8	35.9	52.4
2003 4/.....	0.3	0.7	0.8	1.5	3.5	7.9	14.6	25.0	37.2	55.7
2002.....	0.3	0.5	0.8	1.7	3.7	7.4	15.5	25.8	38.1	53.5
2001.....	0.3	0.5	0.7	1.6	3.9	8.8	14.3	27.1	39.8	62.6
2000.....	0.3	0.5	0.9	1.9	3.9	8.5	17.1	26.8	40.0	59.6
1999.....	0.4	0.5	0.9	1.8	3.7	8.9	16.7	27.5	41.3	63.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Aortic aneurysm and dissection (I71)										
Black, both sexes										
2005 4/.....	1.8	3.3	4.9	5.6	8.8	13.3	15.9	25.6	38.6	50.2
2004 4/.....	1.8	3.2	4.0	5.8	10.0	11.6	14.9	26.1	40.6	43.7
2003 4/.....	1.8	3.8	4.7	5.7	9.0	11.7	17.9	25.9	36.0	46.4
2002.....	2.1	3.8	4.7	6.8	8.3	11.6	18.4	24.9	37.4	50.5
2001.....	2.1	2.7	4.4	6.2	9.3	11.6	18.4	34.4	42.5	55.3
2000.....	2.3	2.9	3.7	7.3	9.3	14.3	20.3	33.8	40.4	62.3
1999.....	2.0	3.5	4.0	6.5	6.5	14.1	19.8	29.6	40.5	60.3
Black, male										
2005 4/.....	3.1	5.3	7.1	8.8	12.0	18.8	17.1	34.0	47.1	51.1
2004 4/.....	2.8	5.1	6.0	8.8	15.9	17.9	18.5	29.6	48.1	56.6
2003 4/.....	2.8	5.8	7.1	8.0	14.7	16.3	20.4	32.9	49.5	51.1
2002.....	3.5	5.7	7.7	11.3	13.1	14.7	24.7	31.2	41.3	49.3
2001.....	3.4	3.9	6.9	8.6	14.6	15.0	26.6	43.2	54.6	48.8
2000.....	3.5	3.6	6.0	10.4	12.8	17.7	27.7	47.7	45.5	69.2
1999.....	2.7	5.4	6.4	9.0	9.3	19.2	25.4	29.9	54.1	76.2
Black, female										
2005 4/.....	*	1.6	3.1	3.0	6.4	9.2	15.2	20.5	34.1	49.9
2004 4/.....	*	1.6	2.4	3.3	5.4	6.8	12.5	24.0	36.7	38.6
2003 4/.....	*	2.1	2.7	3.8	4.7	8.4	16.2	21.7	29.0	44.6
2002.....	*	2.1	2.2	3.2	4.5	9.3	14.1	21.1	35.4	50.9
2001.....	*	1.7	2.2	4.2	5.1	9.0	13.0	29.1	36.4	57.7
2000.....	*	2.3	*	4.7	6.5	11.7	15.3	25.3	37.8	59.7
1999.....	1.4	2.0	*	4.4	4.3	10.4	16.0	29.5	33.7	54.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other diseases of arteries, arterioles and capillaries (I72-I78)										
All races 3/, both sexes										
2005.....	0.4	0.5	1.0	2.1	3.5	6.3	11.4	19.3	31.3	70.6
2004.....	0.4	0.7	1.2	2.1	3.5	6.9	11.9	19.1	32.3	74.5
2003.....	0.4	0.7	1.2	2.0	4.0	7.0	11.4	20.0	32.0	76.0
2002.....	0.4	0.7	1.1	2.2	4.3	7.7	12.6	20.4	32.3	79.0
2001.....	0.5	0.6	1.2	2.2	4.3	7.3	12.9	21.5	32.7	80.7
2000.....	0.4	0.8	1.3	2.1	4.3	8.2	12.8	21.8	35.8	84.0
1999.....	0.5	0.7	1.2	2.3	4.4	7.0	12.9	21.3	36.1	84.9
All races 3/, male										
2005.....	0.3	0.6	1.1	2.5	4.2	7.7	13.7	24.6	35.8	67.2
2004.....	0.4	0.8	1.4	2.3	3.9	7.7	13.6	22.9	35.5	71.6
2003.....	0.3	0.7	1.3	2.0	4.4	9.3	13.0	25.1	37.1	74.2
2002.....	0.4	0.8	1.2	2.3	4.9	8.7	14.2	23.4	39.2	80.8
2001.....	0.4	0.7	1.2	2.4	4.8	8.5	15.2	24.7	40.0	82.5
2000.....	0.3	0.8	1.4	2.6	4.9	9.7	15.7	26.6	41.7	87.6
1999.....	0.4	0.8	1.2	2.6	5.2	8.1	15.7	25.9	40.8	82.2
All races 3/, female										
2005.....	0.5	0.5	0.9	1.7	2.8	5.2	9.6	15.5	28.4	72.2
2004.....	0.5	0.6	1.0	1.8	3.2	6.2	10.5	16.3	30.3	75.8
2003.....	0.5	0.7	1.0	1.9	3.6	5.0	10.1	16.3	28.9	76.9
2002.....	0.4	0.6	1.0	2.1	3.7	6.7	11.3	18.3	28.1	78.2
2001.....	0.5	0.6	1.3	2.0	3.9	6.3	11.1	19.3	28.3	79.9
2000.....	0.5	0.7	1.1	1.6	3.7	6.9	10.5	18.5	32.2	82.6
1999.....	0.5	0.6	1.2	2.1	3.6	6.0	10.6	18.2	33.3	86.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other diseases of arteries, arterioles and capillaries (I72-I78)

White, both sexes

2005 4/.....	0.4	0.5	1.0	1.8	3.3	6.0	11.0	18.6	30.3	69.9
2004 4/.....	0.4	0.7	1.1	1.9	3.2	6.4	11.7	18.4	31.4	72.5
2003 4/.....	0.3	0.6	1.0	1.9	3.7	6.6	11.1	19.5	30.7	74.6
2002.....	0.4	0.6	1.0	2.0	3.9	7.1	12.2	19.7	31.7	77.8
2001.....	0.4	0.6	1.2	2.1	4.0	7.1	12.2	20.8	31.4	78.5
2000.....	0.3	0.7	1.0	1.9	3.9	7.8	12.4	21.0	34.9	82.3
1999.....	0.5	0.7	1.1	2.1	4.1	6.6	12.5	20.6	34.8	82.9

White, male

2005 4/.....	0.3	0.5	1.0	2.1	4.1	7.1	13.3	23.4	35.0	67.8
2004 4/.....	0.4	0.8	1.3	2.1	3.6	7.1	13.6	22.3	35.4	70.7
2003 4/.....	0.3	0.7	1.2	2.0	4.0	8.7	12.8	24.4	35.4	74.1
2002.....	0.4	0.8	1.1	2.0	4.4	8.2	13.7	23.0	38.4	79.1
2001.....	0.3	0.6	1.1	2.3	4.4	8.3	14.4	24.2	38.5	80.9
2000.....	0.3	0.7	1.1	2.4	4.5	9.4	15.3	25.9	41.1	86.9
1999.....	0.4	0.8	1.1	2.4	4.9	7.9	15.3	24.8	39.6	81.3

White, female

2005 4/.....	0.5	0.5	0.9	1.5	2.6	4.9	9.0	15.1	27.4	70.8
2004 4/.....	0.4	0.5	0.9	1.8	2.8	5.7	10.1	15.6	28.9	73.3
2003 4/.....	0.4	0.6	0.9	1.9	3.4	4.8	9.8	15.9	27.8	74.8
2002.....	0.3	0.5	0.9	2.0	3.4	6.1	11.0	17.3	27.5	77.2
2001.....	0.5	0.6	1.2	1.8	3.5	6.0	10.4	18.4	27.1	77.5
2000.....	0.4	0.6	0.9	1.4	3.3	6.5	10.0	17.6	31.2	80.4
1999.....	0.5	0.5	1.0	1.8	3.4	5.6	10.2	17.6	32.0	83.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other diseases of arteries, arterioles and capillaries (I72-I78)

Black, both sexes

2005 4/.....	0.7	1.0	1.8	4.7	5.9	11.2	19.3	31.4	49.8	94.3
2004 4/.....	0.8	1.1	2.3	3.6	7.4	12.7	17.8	30.8	49.5	111.4
2003 4/.....	0.8	1.5	2.6	2.8	7.3	12.3	18.0	29.4	56.1	103.6
2002.....	0.7	1.5	2.0	4.4	8.3	14.1	19.4	33.5	46.1	106.3
2001.....	1.1	1.3	2.1	4.5	8.7	11.3	24.4	34.9	53.6	119.4
2000.....	0.9	1.7	3.1	3.9	9.0	13.0	19.6	35.2	52.0	115.4
1999.....	0.8	1.0	2.1	4.9	7.0	11.9	18.8	34.6	56.1	118.6

Black, male

2005 4/.....	*	*	2.4	6.2	7.3	15.0	22.5	45.1	56.9	78.1
2004 4/.....	*	*	2.9	4.9	8.4	15.5	19.1	38.7	50.2	101.3
2003 4/.....	*	*	2.8	3.2	8.9	17.2	19.7	39.7	69.4	87.6
2002.....	*	*	2.1	5.2	9.9	16.0	24.0	34.4	58.8	116.9
2001.....	*	*	2.5	4.5	9.2	13.0	29.3	35.8	73.0	116.9
2000.....	*	1.9	4.0	4.8	9.7	15.1	24.1	43.0	59.0	108.4
1999.....	*	*	*	5.1	8.5	12.7	23.7	44.6	63.7	107.8

Black, female

2005 4/.....	*	*	*	3.4	4.8	8.5	17.1	23.2	46.0	100.9
2004 4/.....	*	*	1.8	2.5	6.6	10.6	16.9	26.1	49.1	115.4
2003 4/.....	*	1.8	2.4	2.4	6.0	8.6	16.8	23.3	49.1	109.8
2002.....	*	1.9	1.8	3.7	7.0	12.8	16.4	32.9	39.4	102.3
2001.....	*	*	1.8	4.5	8.3	10.1	21.1	34.4	43.7	120.4
2000.....	*	*	2.4	3.2	8.3	11.5	16.6	30.5	48.4	118.0
1999.....	*	*	*	4.7	5.8	11.4	15.5	28.6	52.3	122.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other disorders of circulatory system (I80-I99)										
All races 3/, both sexes										
2005.....	0.9	1.3	1.6	1.9	2.4	3.3	4.4	7.0	10.8	23.0
2004.....	0.9	1.3	1.4	1.9	2.5	3.1	4.3	7.1	11.5	23.3
2003.....	0.9	1.3	1.5	1.7	2.5	3.3	4.9	7.0	11.9	24.0
2002.....	0.8	1.2	1.5	2.0	2.5	3.4	4.3	7.4	13.2	24.4
2001.....	0.8	1.1	1.3	1.9	2.6	3.8	5.4	7.7	11.9	26.1
2000.....	0.9	1.2	1.5	1.7	2.6	3.4	5.3	7.7	11.9	25.6
1999.....	0.8	1.0	1.0	1.6	2.2	3.3	4.3	6.8	12.7	23.4
All races 3/, male										
2005.....	1.0	1.6	2.1	2.3	3.0	4.0	5.0	6.6	11.1	19.2
2004.....	1.0	1.4	1.8	2.2	3.0	3.3	4.6	7.9	12.2	19.6
2003.....	1.0	1.4	1.8	2.2	2.7	3.5	4.9	6.9	10.9	19.1
2002.....	0.8	1.3	1.6	2.3	2.6	3.6	4.2	7.0	13.3	20.5
2001.....	0.9	1.2	1.5	2.2	2.9	3.9	5.9	8.3	12.1	22.0
2000.....	1.1	1.5	1.7	1.9	2.7	3.9	5.6	8.0	12.4	22.2
1999.....	0.9	1.1	1.2	1.7	2.3	3.7	4.8	7.5	13.2	20.8
All races 3/, female										
2005.....	0.9	1.0	1.2	1.5	1.8	2.7	4.0	7.2	10.7	24.8
2004.....	0.8	1.1	1.1	1.6	2.0	2.9	4.0	6.5	11.0	25.0
2003.....	0.9	1.1	1.2	1.3	2.3	3.0	5.0	7.0	12.5	26.2
2002.....	0.8	1.1	1.4	1.7	2.5	3.2	4.4	7.7	13.1	26.2
2001.....	0.8	0.9	1.0	1.7	2.3	3.7	5.0	7.3	11.7	27.8
2000.....	0.7	1.0	1.3	1.5	2.6	3.0	5.0	7.5	11.6	26.9
1999.....	0.6	1.0	0.9	1.5	2.1	2.9	3.9	6.3	12.4	24.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other disorders of circulatory system (I80-I99)										
White, both sexes										
2005 4/.....	0.7	1.2	1.4	1.7	2.1	3.2	4.2	6.7	10.5	23.5
2004 4/.....	0.7	1.1	1.2	1.8	2.3	3.0	3.9	6.8	11.3	23.2
2003 4/.....	0.8	1.1	1.4	1.6	2.2	3.0	4.7	6.5	11.6	23.6
2002.....	0.6	1.0	1.4	1.9	2.2	3.2	4.1	7.1	12.7	24.3
2001.....	0.7	0.9	1.1	1.8	2.3	3.6	5.1	7.2	11.8	26.0
2000.....	0.8	1.1	1.4	1.5	2.4	2.9	5.0	7.3	11.8	25.5
1999.....	0.6	0.9	0.9	1.6	2.1	2.9	4.1	6.5	12.4	23.2
White, male										
2005 4/.....	0.8	1.4	1.9	2.1	2.6	3.7	4.6	6.4	10.5	19.1
2004 4/.....	0.8	1.4	1.5	2.1	2.7	3.1	4.3	7.6	11.9	19.7
2003 4/.....	0.9	1.2	1.6	2.1	2.3	3.2	4.6	6.4	10.2	18.5
2002.....	0.6	1.1	1.5	2.1	2.3	3.3	4.0	6.7	12.4	19.9
2001.....	0.7	1.2	1.3	2.0	2.5	3.6	5.6	7.8	12.0	21.7
2000.....	1.0	1.3	1.6	1.7	2.4	3.4	5.5	7.8	12.3	22.0
1999.....	0.8	0.9	1.0	1.7	2.3	3.2	4.7	7.0	12.8	20.7
White, female										
2005 4/.....	0.7	1.0	1.0	1.4	1.6	2.7	3.9	6.9	10.5	25.5
2004 4/.....	0.6	0.9	0.9	1.4	1.8	2.8	3.6	6.2	11.0	24.8
2003 4/.....	0.7	1.1	1.2	1.1	2.2	2.9	4.7	6.5	12.4	25.8
2002.....	0.7	1.0	1.2	1.6	2.1	3.0	4.2	7.4	12.9	26.3
2001.....	0.7	0.7	1.0	1.5	2.1	3.5	4.8	6.8	11.7	27.9
2000.....	0.5	0.9	1.1	1.4	2.4	2.5	4.7	7.0	11.5	26.9
1999.....	0.5	0.9	0.9	1.5	1.9	2.7	3.7	6.1	12.2	24.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other disorders of circulatory system (I80-I99)										
Black, both sexes										
2005 4/.....	2.3	2.6	3.5	4.0	5.2	5.5	7.6	11.5	17.5	24.3
2004 4/.....	2.5	2.4	3.6	3.6	5.0	4.9	8.9	12.9	14.6	28.8
2003 4/.....	2.3	2.6	2.5	3.6	5.2	6.2	9.3	13.8	17.8	33.6
2002.....	1.8	2.4	2.9	4.2	5.9	6.4	7.2	12.9	22.2	30.0
2001.....	1.8	2.1	2.4	3.9	6.1	6.9	9.0	14.1	14.9	30.1
2000.....	1.8	2.5	3.2	3.9	5.1	8.7	9.1	14.6	14.1	30.0
1999.....	1.9	2.4	2.3	2.1	3.9	6.9	6.4	11.4	17.6	28.4
Black, male										
2005 4/.....	2.4	3.6	4.1	4.8	6.9	7.8	10.1	12.1	22.5	25.1
2004 4/.....	2.5	2.1	5.0	3.6	6.2	5.8	8.7	14.1	16.5	24.8
2003 4/.....	1.8	3.4	3.4	3.6	7.3	7.6	10.2	15.5	20.7	32.3
2002.....	2.1	3.0	2.9	5.4	5.9	7.7	7.7	12.4	28.3	32.2
2001.....	2.2	1.7	3.1	4.7	8.5	7.9	10.4	16.3	*	29.5
2000.....	2.0	3.1	3.7	4.8	5.6	9.4	8.7	13.2	*	30.0
1999.....	2.6	2.7	3.0	*	*	9.3	6.8	14.7	20.9	*
Black, female										
2005 4/.....	2.2	1.8	3.0	3.3	3.8	3.7	5.8	11.2	14.8	24.0
2004 4/.....	2.6	2.6	2.4	3.7	4.1	4.1	9.0	12.1	13.6	30.3
2003 4/.....	2.7	1.9	1.8	3.6	3.6	5.1	8.6	12.8	16.2	34.0
2002.....	1.6	1.8	2.9	3.2	5.9	5.4	6.9	13.2	19.1	29.2
2001.....	1.5	2.4	*	3.2	4.2	6.2	8.1	12.8	14.7	30.3
2000.....	1.7	1.9	2.8	3.2	4.7	8.3	9.3	15.4	14.4	30.0
1999.....	*	2.1	*	*	4.2	5.0	6.1	9.4	16.0	30.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Influenza and pneumonia (J10-J18)										
All races 3/, both sexes										
2005.....	2.7	4.2	6.1	8.8	14.6	24.5	48.6	96.1	202.7	593.9
2004.....	2.5	3.6	5.6	8.3	14.2	23.8	47.2	94.7	198.7	582.6
2003.....	2.7	4.2	6.4	8.2	15.2	26.6	49.4	103.3	216.8	666.1
2002.....	2.6	4.2	5.5	8.5	14.8	25.3	50.9	106.5	227.3	696.6
2001.....	2.6	3.9	5.4	8.1	13.9	24.4	49.3	101.1	216.9	685.6
2000.....	3.0	3.9	5.6	8.9	15.6	26.6	52.6	106.7	240.8	744.1
1999.....	2.7	4.1	5.3	8.4	14.3	25.0	50.5	105.9	233.8	751.8
All races 3/, male										
2005.....	3.1	5.1	7.5	10.7	17.5	29.6	59.0	119.3	253.7	673.5
2004.....	2.7	4.3	7.1	9.9	17.2	29.8	58.6	118.3	252.3	677.2
2003.....	3.1	4.9	8.1	9.6	17.9	33.3	62.1	130.1	272.3	757.8
2002.....	3.1	4.9	6.7	10.2	18.0	31.2	63.9	133.9	287.7	805.8
2001.....	3.0	4.9	6.7	10.0	17.0	30.3	61.6	130.7	274.7	808.4
2000.....	3.8	4.7	7.0	10.9	18.9	32.9	65.7	134.1	308.8	892.0
1999.....	3.1	5.2	6.8	9.7	16.8	31.7	64.2	133.5	301.7	894.7
All races 3/, female										
2005.....	2.3	3.4	4.9	6.9	12.0	20.1	40.2	79.3	171.1	557.4
2004.....	2.3	3.0	4.2	6.8	11.4	18.6	37.9	77.7	165.7	540.0
2003.....	2.3	3.5	4.8	6.8	12.7	20.7	39.2	84.2	182.9	625.6
2002.....	2.2	3.4	4.3	6.8	11.8	20.1	40.6	87.2	190.6	649.2
2001.....	2.2	3.0	4.1	6.4	11.1	19.2	39.5	80.5	182.3	634.2
2000.....	2.2	3.1	4.3	7.0	12.5	21.2	42.2	87.6	200.7	683.9
1999.....	2.3	3.0	3.9	7.2	12.0	19.2	39.7	86.7	194.1	694.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Influenza and pneumonia (J10-J18)										
White, both sexes										
2005 4/.....	2.4	3.8	5.5	8.0	13.7	23.3	47.0	95.1	204.0	606.4
2004 4/.....	2.2	3.2	5.0	7.7	13.3	22.3	46.2	93.1	198.3	590.7
2003 4/.....	2.3	3.7	5.7	7.4	14.0	25.2	47.9	102.4	218.4	679.3
2002.....	2.3	3.8	5.0	7.7	13.8	24.5	49.9	105.4	228.0	709.1
2001.....	2.2	3.3	4.8	7.5	13.0	23.3	48.1	98.6	216.6	695.0
2000.....	2.5	3.4	4.9	8.1	14.3	24.5	50.3	105.6	240.7	758.3
1999.....	2.1	3.4	4.6	7.7	13.1	24.0	48.8	105.5	235.5	764.3
White, male										
2005 4/.....	2.7	4.5	6.6	9.4	16.1	27.6	56.5	117.7	253.9	684.4
2004 4/.....	2.4	3.8	6.2	9.1	15.9	27.1	56.7	114.6	250.8	685.5
2003 4/.....	2.6	4.3	7.2	8.5	15.7	30.8	59.2	127.2	271.4	767.1
2002.....	2.7	4.4	6.0	9.1	16.5	29.8	62.4	131.4	286.5	815.9
2001.....	2.5	4.2	5.7	9.0	15.4	28.3	59.3	125.5	272.3	814.4
2000.....	3.2	3.9	5.9	9.5	17.2	29.7	62.2	131.8	306.4	906.1
1999.....	2.3	4.2	5.7	8.6	15.1	30.1	60.9	132.2	301.9	905.7
White, female										
2005 4/.....	2.1	3.2	4.4	6.6	11.4	19.5	39.2	78.4	172.8	570.4
2004 4/.....	2.1	2.7	3.8	6.2	10.9	18.1	37.5	77.3	165.6	548.0
2003 4/.....	2.1	3.1	4.3	6.3	12.4	20.2	38.6	84.5	185.7	640.4
2002.....	1.9	3.2	3.9	6.3	11.3	19.8	39.7	86.8	192.1	662.6
2001.....	1.8	2.5	4.0	6.0	10.7	19.0	39.0	79.6	182.9	645.0
2000.....	1.8	2.8	4.0	6.7	11.6	20.0	40.8	87.1	201.7	698.3
1999.....	1.9	2.6	3.6	6.7	11.2	18.7	39.1	86.8	196.4	707.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Influenza and pneumonia (J10-J18)										
Black, both sexes										
2005 4/.....	4.9	7.9	12.1	16.8	24.5	39.2	68.2	115.8	199.3	471.2
2004 4/.....	5.0	7.0	11.6	15.5	23.8	39.8	62.4	117.6	215.1	505.1
2003 4/.....	5.5	8.5	12.6	15.7	27.1	41.6	70.1	122.5	205.6	525.4
2002.....	5.2	7.5	10.1	17.2	26.3	35.5	68.9	123.3	231.6	575.2
2001.....	5.8	9.0	10.7	15.5	23.3	34.3	65.7	133.7	224.0	581.3
2000.....	6.1	8.3	12.2	17.8	27.2	46.0	77.5	124.4	245.4	600.9
1999.....	6.9	9.7	11.8	16.2	26.6	36.8	75.2	118.7	219.9	634.8
Black, male										
2005 4/.....	6.1	10.5	15.9	23.8	33.3	53.7	90.5	144.4	259.1	531.7
2004 4/.....	5.9	8.5	15.8	19.1	33.1	61.3	87.4	169.7	297.7	598.0
2003 4/.....	7.0	10.4	16.7	21.3	39.7	61.6	100.5	177.6	285.0	641.2
2002.....	6.2	9.9	12.9	23.7	36.0	48.1	89.8	173.2	321.7	689.4
2001.....	7.0	11.4	15.7	20.8	33.6	50.1	94.9	203.5	309.0	714.8
2000.....	8.3	11.3	17.2	25.9	35.2	64.8	107.6	172.5	343.9	755.4
1999.....	9.0	13.4	17.2	21.7	34.2	51.9	109.6	160.0	312.2	803.8
Black, female										
2005 4/.....	3.9	5.7	9.0	11.0	17.7	28.4	53.0	98.7	167.7	446.8
2004 4/.....	4.2	5.7	8.0	12.6	16.6	23.9	45.5	86.8	171.6	468.2
2003 4/.....	4.1	7.0	9.1	11.1	17.3	26.9	49.6	89.8	164.0	480.4
2002.....	4.3	5.5	7.7	12.0	18.7	26.1	54.9	93.6	184.6	531.5
2001.....	4.8	6.9	6.5	11.2	15.2	22.7	46.2	92.1	180.7	531.7
2000.....	4.2	5.8	7.9	11.2	20.9	32.1	57.5	95.4	195.7	543.4
1999.....	5.1	6.6	7.1	11.8	20.6	25.6	52.4	93.8	174.1	571.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Influenza (J10-J11)										
All races 3/, both sexes										
2005.....	*	*	0.1	0.1	0.3	0.4	1.1	2.4	5.4	19.3
2004.....	*	*	*	0.1	0.3	0.3	0.8	1.6	3.2	11.4
2003.....	*	0.1	0.2	0.3	0.3	0.7	1.3	2.7	4.7	17.0
2002.....	*	*	*	*	*	*	0.4	0.8	2.2	8.8
2001.....	*	*	*	*	*	*	*	*	0.7	2.2
2000.....	*	0.1	0.1	0.3	0.5	0.8	1.4	2.5	6.1	20.2
1999.....	*	*	*	0.2	0.4	0.6	1.2	2.1	6.1	21.2
All races 3/, male										
2005.....	*	*	*	*	0.4	*	1.1	2.5	5.3	18.2
2004.....	*	*	*	*	*	*	1.1	1.8	3.3	11.1
2003.....	*	*	*	0.3	*	0.8	1.3	2.9	4.9	17.0
2002.....	*	*	*	*	*	*	*	0.9	2.6	8.9
2001.....	*	*	*	*	*	*	*	*	*	2.2
2000.....	*	*	*	0.3	0.5	1.0	1.6	2.7	6.3	20.7
1999.....	*	*	*	*	0.5	0.6	1.2	2.4	7.3	19.9
All races 3/, female										
2005.....	*	*	*	*	*	0.5	1.1	2.3	5.5	19.8
2004.....	*	*	*	*	*	*	0.6	1.4	3.1	11.5
2003.....	*	*	*	0.3	0.3	0.7	1.3	2.5	4.7	17.1
2002.....	*	*	*	*	*	*	*	0.7	2.0	8.7
2001.....	*	*	*	*	*	*	*	*	0.7	2.2
2000.....	*	*	*	*	0.5	0.6	1.2	2.5	6.0	19.9
1999.....	*	*	*	*	*	0.6	1.1	1.9	5.4	21.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Influenza (J10-J11)										
White, both sexes										
2005 4/.....	*	*	0.1	0.2	0.3	0.5	1.2	2.5	5.8	20.8
2004 4/.....	*	*	*	0.2	0.3	0.3	0.9	1.5	3.4	12.3
2003 4/.....	*	0.1	0.2	0.3	0.3	0.8	1.4	2.9	5.1	18.2
2002.....	*	*	*	*	*	*	0.4	0.9	2.4	9.4
2001.....	*	*	*	*	*	*	*	*	0.7	2.3
2000.....	*	*	0.1	0.3	0.5	0.8	1.4	2.7	6.5	21.4
1999.....	*	*	*	0.2	0.4	0.7	1.2	2.2	6.4	22.8
White, male										
2005 4/.....	*	*	*	*	*	*	1.2	2.7	5.6	19.2
2004 4/.....	*	*	*	*	*	*	1.1	1.7	3.6	12.1
2003 4/.....	*	*	*	0.3	*	0.9	1.4	3.1	5.2	18.3
2002.....	*	*	*	*	*	*	*	1.0	2.8	9.7
2001.....	*	*	*	*	*	*	*	*	*	2.3
2000.....	*	*	*	*	0.5	1.0	1.7	2.7	6.7	21.6
1999.....	*	*	*	*	0.5	0.6	1.2	2.6	7.5	21.7
White, female										
2005 4/.....	*	*	*	*	*	0.5	1.2	2.4	5.9	21.5
2004 4/.....	*	*	*	*	*	*	0.7	1.4	3.3	12.4
2003 4/.....	*	*	*	*	0.4	0.7	1.4	2.8	5.0	18.1
2002.....	*	*	*	*	*	*	*	0.8	2.2	9.3
2001.....	*	*	*	*	*	*	*	*	0.7	2.3
2000.....	*	*	*	*	0.6	0.7	1.2	2.7	6.3	21.3
1999.....	*	*	*	*	*	0.7	1.1	2.0	5.7	23.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Pneumonia (J12-J18)										
All races 3/, both sexes										
2005.....	2.6	4.2	6.0	8.6	14.3	24.1	47.5	93.7	197.3	574.7
2004.....	2.5	3.6	5.6	8.2	13.9	23.5	46.4	93.1	195.6	571.2
2003.....	2.6	4.1	6.2	7.9	14.9	25.8	48.1	100.6	212.1	649.1
2002.....	2.6	4.1	5.4	8.4	14.6	25.1	50.5	105.7	225.1	687.8
2001.....	2.6	3.9	5.3	8.1	13.8	24.3	49.2	100.9	216.1	683.4
2000.....	2.9	3.8	5.5	8.6	15.1	25.8	51.2	104.1	234.7	723.9
1999.....	2.7	4.0	5.2	8.2	13.9	24.3	49.3	103.8	227.7	730.6
All races 3/, male										
2005.....	3.0	5.0	7.4	10.5	17.1	29.2	57.9	116.8	248.4	655.3
2004.....	2.7	4.2	7.0	9.7	17.0	29.4	57.6	116.5	249.1	666.1
2003.....	3.0	4.8	7.9	9.3	17.6	32.4	60.8	127.2	267.5	740.9
2002.....	3.0	4.9	6.7	10.1	17.8	30.9	63.4	133.0	285.0	796.9
2001.....	3.0	4.8	6.6	9.9	16.9	30.3	61.4	130.5	273.9	806.2
2000.....	3.7	4.6	6.8	10.6	18.4	31.9	64.1	131.5	302.6	871.3
1999.....	3.1	5.1	6.7	9.5	16.3	31.1	63.0	131.1	294.5	874.7
All races 3/, female										
2005.....	2.3	3.4	4.7	6.8	11.7	19.6	39.1	77.0	165.6	537.6
2004.....	2.3	2.9	4.1	6.7	11.1	18.4	37.3	76.4	162.6	528.5
2003.....	2.2	3.4	4.7	6.5	12.4	20.1	37.9	81.7	178.2	608.5
2002.....	2.2	3.4	4.3	6.8	11.8	20.0	40.2	86.5	188.6	640.5
2001.....	2.2	2.9	4.1	6.3	11.0	19.1	39.4	80.2	181.5	632.0
2000.....	2.1	3.0	4.2	6.7	12.0	20.5	41.0	85.1	194.6	663.9
1999.....	2.3	2.9	3.9	7.0	11.7	18.6	38.6	84.8	188.7	672.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Pneumonia (J12-J18)										
White, both sexes										
2005 4/.....	2.4	3.8	5.4	7.8	13.4	22.8	45.8	92.6	198.2	585.6
2004 4/.....	2.2	3.2	4.9	7.5	13.1	22.0	45.3	91.5	194.9	578.4
2003 4/.....	2.3	3.6	5.6	7.1	13.7	24.4	46.5	99.5	213.3	661.1
2002.....	2.3	3.8	4.9	7.6	13.7	24.3	49.5	104.5	225.6	699.6
2001.....	2.1	3.3	4.8	7.4	12.9	23.2	48.0	98.3	215.8	692.7
2000.....	2.4	3.2	4.8	7.8	13.8	23.7	48.9	102.9	234.3	737.0
1999.....	2.1	3.3	4.6	7.4	12.6	23.4	47.6	103.3	229.2	741.6
White, male										
2005 4/.....	2.7	4.4	6.5	9.2	15.8	27.2	55.3	115.1	248.3	665.2
2004 4/.....	2.3	3.8	6.2	9.0	15.7	26.7	55.6	112.9	247.3	673.4
2003 4/.....	2.5	4.2	7.0	8.2	15.4	29.9	57.8	124.1	266.2	748.8
2002.....	2.7	4.3	6.0	9.0	16.3	29.6	61.9	130.4	283.7	806.2
2001.....	2.5	4.1	5.6	9.0	15.3	28.3	59.2	125.3	271.5	812.1
2000.....	3.1	3.8	5.7	9.2	16.7	28.6	60.6	129.1	299.7	884.4
1999.....	2.2	4.2	5.6	8.4	14.6	29.5	59.7	129.6	294.3	884.0
White, female										
2005 4/.....	2.1	3.1	4.3	6.5	11.1	19.0	38.0	75.9	166.9	548.9
2004 4/.....	2.1	2.6	3.7	6.1	10.6	17.8	36.9	75.9	162.3	535.6
2003 4/.....	2.0	3.0	4.2	6.0	12.1	19.5	37.2	81.7	180.7	622.3
2002.....	1.9	3.2	3.9	6.2	11.3	19.7	39.4	86.0	190.0	653.3
2001.....	1.8	2.4	3.9	5.9	10.6	18.8	38.9	79.3	182.2	642.8
2000.....	1.8	2.7	3.9	6.4	11.1	19.4	39.6	84.5	195.4	677.1
1999.....	1.9	2.5	3.5	6.5	10.8	18.0	37.9	84.8	190.7	684.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Pneumonia (J12-J18)										
Black, both sexes										
2005 4/.....	4.9	7.8	12.0	16.7	24.1	39.0	67.8	114.9	197.1	467.2
2004 4/.....	4.9	6.9	11.5	15.5	23.5	39.5	61.9	115.6	214.1	502.5
2003 4/.....	5.4	8.3	12.4	15.4	27.0	41.0	69.6	121.7	203.6	519.6
2002.....	5.2	7.5	10.1	17.2	26.1	35.0	68.6	123.2	230.8	572.8
2001.....	5.8	9.0	10.7	15.4	23.1	34.1	65.7	133.7	222.9	579.7
2000.....	6.0	8.1	12.1	17.7	26.9	45.4	76.0	123.5	243.1	591.5
1999.....	6.8	9.7	11.6	16.0	26.4	36.5	73.9	117.8	217.0	629.1
Black, male										
2005 4/.....	6.1	10.2	15.7	23.6	32.5	53.7	89.8	143.9	257.7	524.3
2004 4/.....	5.9	8.4	15.8	19.1	32.9	61.1	86.4	165.6	297.0	597.1
2003 4/.....	7.0	10.0	16.5	20.8	39.5	60.8	100.2	176.6	283.5	638.1
2002.....	6.2	9.9	12.9	23.5	35.6	47.4	89.8	173.2	320.2	687.2
2001.....	7.0	11.4	15.7	20.6	33.2	49.6	94.9	203.5	309.0	712.6
2000.....	8.1	11.0	17.0	25.5	35.0	64.3	106.6	170.6	343.1	742.7
1999.....	8.9	13.3	17.1	21.7	33.9	51.4	108.2	159.6	307.8	800.3
Black, female										
2005 4/.....	3.8	5.7	8.8	10.9	17.6	28.0	52.8	97.6	165.1	444.1
2004 4/.....	4.1	5.7	7.9	12.6	16.2	23.6	45.3	86.0	170.5	465.0
2003 4/.....	4.1	6.7	8.9	11.0	17.1	26.3	49.0	89.0	161.7	473.5
2002.....	4.3	5.5	7.7	12.0	18.6	25.9	54.4	93.3	184.2	529.0
2001.....	4.7	6.9	6.5	11.2	15.2	22.7	46.2	92.1	179.1	530.5
2000.....	4.2	5.6	7.9	11.2	20.4	31.5	55.7	95.1	192.8	535.3
1999.....	5.0	6.6	7.0	11.4	20.5	25.4	51.1	92.7	172.0	565.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other acute lower respiratory infections (J20-J22)

All races 3/, both sexes

2005.....	*	*	*	*	*	*	*	0.4	0.9	3.3
2004.....	*	*	*	*	*	*	*	0.3	0.6	2.6
2003.....	*	*	*	*	*	*	*	0.5	0.8	3.5
2002.....	*	*	*	*	*	*	0.3	0.3	0.8	3.3
2001.....	*	*	*	*	*	*	*	0.3	0.7	3.3
2000.....	*	*	*	*	*	*	0.3	0.6	0.8	4.5
1999.....	*	*	*	*	*	*	0.2	0.6	1.3	6.0

All races 3/, male

2005.....	*	*	*	*	*	*	*	*	1.3	3.1
2004.....	*	*	*	*	*	*	*	*	*	2.5
2003.....	*	*	*	*	*	*	*	*	*	2.4
2002.....	*	*	*	*	*	*	*	*	*	2.8
2001.....	*	*	*	*	*	*	*	*	*	2.3
2000.....	*	*	*	*	*	*	*	0.7	*	3.8
1999.....	*	*	*	*	*	*	*	0.7	1.6	4.0

All races 3/, female

2005.....	*	*	*	*	*	*	*	*	0.6	3.5
2004.....	*	*	*	*	*	*	*	*	0.7	2.7
2003.....	*	*	*	*	*	*	*	*	0.8	3.9
2002.....	*	*	*	*	*	*	*	*	0.9	3.5
2001.....	*	*	*	*	*	*	*	*	0.7	3.7
2000.....	*	*	*	*	*	*	*	0.5	0.9	4.8
1999.....	*	*	*	*	*	*	*	0.5	1.1	6.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other acute lower respiratory infections (J20-J22)

White, both sexes

2005 4/.....	*	*	*	*	*	*	*	0.4	0.9	3.5
2004 4/.....	*	*	*	*	*	*	*	*	0.7	2.8
2003 4/.....	*	*	*	*	*	*	*	0.5	0.9	3.7
2002.....	*	*	*	*	*	*	*	*	0.8	3.5
2001.....	*	*	*	*	*	*	*	0.3	0.7	3.5
2000.....	*	*	*	*	*	*	0.3	0.6	0.9	4.7
1999.....	*	*	*	*	*	*	*	0.6	1.3	6.3

White, male

2005 4/.....	*	*	*	*	*	*	*	*	1.4	3.3
2004 4/.....	*	*	*	*	*	*	*	*	*	2.6
2003 4/.....	*	*	*	*	*	*	*	*	*	2.7
2002.....	*	*	*	*	*	*	*	*	*	2.9
2001.....	*	*	*	*	*	*	*	*	*	2.4
2000.....	*	*	*	*	*	*	*	0.8	*	3.8
1999.....	*	*	*	*	*	*	*	0.8	1.6	4.0

White, female

2005 4/.....	*	*	*	*	*	*	*	*	*	3.6
2004 4/.....	*	*	*	*	*	*	*	*	0.7	2.8
2003 4/.....	*	*	*	*	*	*	*	*	0.8	4.2
2002.....	*	*	*	*	*	*	*	*	0.9	3.7
2001.....	*	*	*	*	*	*	*	*	0.8	3.9
2000.....	*	*	*	*	*	*	*	*	1.0	5.1
1999.....	*	*	*	*	*	*	*	0.5	1.1	7.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Acute bronchitis and bronchiolitis (J20-J21)

All races 3/, both sexes

2005.....	*	*	*	*	*	*	*	0.3	0.5	1.9
2004.....	*	*	*	*	*	*	*	*	*	1.4
2003.....	*	*	*	*	*	*	*	*	0.5	2.1
2002.....	*	*	*	*	*	*	*	*	0.5	2.0
2001.....	*	*	*	*	*	*	*	*	0.4	2.1
2000.....	*	*	*	*	*	*	*	0.4	0.5	2.6
1999.....	*	*	*	*	*	*	*	0.4	0.8	3.6

All races 3/, male

2005.....	*	*	*	*	*	*	*	*	*	1.7
2004.....	*	*	*	*	*	*	*	*	*	1.8
2003.....	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	1.7
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	2.4
1999.....	*	*	*	*	*	*	*	*	*	2.6

All races 3/, female

2005.....	*	*	*	*	*	*	*	*	*	2.0
2004.....	*	*	*	*	*	*	*	*	*	1.3
2003.....	*	*	*	*	*	*	*	*	*	2.5
2002.....	*	*	*	*	*	*	*	*	*	2.2
2001.....	*	*	*	*	*	*	*	*	*	2.4
2000.....	*	*	*	*	*	*	*	*	*	2.7
1999.....	*	*	*	*	*	*	*	*	0.7	4.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Acute bronchitis and bronchiolitis (J20-J21)

White, both sexes

2005 4/.....	*	*	*	*	*	*	*	0.3	0.6	2.0
2004 4/.....	*	*	*	*	*	*	*	*	*	1.5
2003 4/.....	*	*	*	*	*	*	*	*	0.5	2.3
2002.....	*	*	*	*	*	*	*	*	0.5	2.1
2001.....	*	*	*	*	*	*	*	*	0.5	2.1
2000.....	*	*	*	*	*	*	*	0.4	0.5	2.7
1999.....	*	*	*	*	*	*	*	0.4	0.9	3.8

White, male

2005 4/.....	*	*	*	*	*	*	*	*	*	1.7
2004 4/.....	*	*	*	*	*	*	*	*	*	1.8
2003 4/.....	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	1.7
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	2.4
1999.....	*	*	*	*	*	*	*	*	*	2.6

White, female

2005 4/.....	*	*	*	*	*	*	*	*	*	2.1
2004 4/.....	*	*	*	*	*	*	*	*	*	1.3
2003 4/.....	*	*	*	*	*	*	*	*	*	2.7
2002.....	*	*	*	*	*	*	*	*	*	2.3
2001.....	*	*	*	*	*	*	*	*	*	2.5
2000.....	*	*	*	*	*	*	*	*	*	2.9
1999.....	*	*	*	*	*	*	*	*	0.7	4.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Chronic lower respiratory diseases (J40-J47)										
All races 3/, both sexes										
2005.....	2.8	6.5	12.6	26.9	62.1	118.6	210.3	330.6	457.8	637.2
2004.....	2.6	5.6	11.7	25.5	60.0	113.7	200.6	315.7	434.7	601.7
2003.....	2.8	5.9	11.9	27.5	63.9	121.1	210.8	330.0	455.9	635.1
2002.....	2.9	5.8	12.0	28.2	60.8	119.0	211.5	330.1	465.7	637.6
2001.....	2.7	5.4	11.9	29.4	62.9	124.2	215.4	329.2	452.9	644.7
2000.....	2.8	5.8	11.8	29.6	62.5	124.1	218.2	327.8	473.4	648.6
1999.....	2.6	5.5	12.0	31.9	66.6	130.4	227.6	340.2	484.5	646.0
All races 3/, male										
2005.....	2.7	6.4	13.6	28.5	67.6	132.9	240.0	393.6	564.7	819.9
2004.....	2.6	5.6	12.5	26.9	64.6	126.1	228.2	378.8	543.4	811.1
2003.....	2.5	6.3	12.3	29.4	69.2	132.6	244.0	395.8	569.1	865.9
2002.....	3.0	5.9	12.8	29.5	65.9	131.5	246.0	405.6	595.9	894.8
2001.....	2.5	5.8	12.3	31.4	67.0	137.1	252.9	406.7	583.5	916.9
2000.....	2.6	6.2	12.1	31.8	68.0	139.7	257.8	404.2	628.5	967.9
1999.....	2.4	5.7	12.1	34.3	74.6	151.0	278.3	432.4	648.3	996.7
All races 3/, female										
2005.....	2.8	6.6	11.7	25.5	57.1	106.1	186.2	284.8	391.5	553.3
2004.....	2.7	5.5	10.9	24.2	55.8	102.9	178.4	270.4	367.7	507.4
2003.....	3.1	5.4	11.5	25.7	59.0	111.2	184.2	283.1	386.7	533.0
2002.....	2.8	5.6	11.2	26.9	56.1	108.3	183.8	276.8	386.5	526.0
2001.....	2.9	5.1	11.5	27.5	59.1	113.1	185.6	275.0	374.6	530.8
2000.....	2.9	5.5	11.5	27.5	57.5	110.7	187.0	274.6	381.9	518.5
1999.....	2.9	5.3	11.9	29.7	59.3	112.8	188.0	276.1	388.7	504.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Chronic lower respiratory diseases (J40-J47)										
White, both sexes										
2005 4/.....	2.5	6.2	12.4	27.7	65.6	126.2	224.5	349.4	481.5	667.6
2004 4/.....	2.4	5.2	11.6	26.1	63.2	121.7	214.3	333.3	457.6	630.1
2003 4/.....	2.5	5.6	11.7	28.0	67.3	129.5	223.2	347.2	478.9	664.5
2002.....	2.5	5.5	11.7	28.9	63.8	126.8	223.3	346.9	486.5	666.2
2001.....	2.4	5.1	11.7	30.2	66.2	131.3	227.1	346.1	473.5	671.8
2000.....	2.3	5.3	11.4	30.1	65.5	130.5	229.6	343.1	495.5	675.6
1999.....	2.2	4.9	11.8	32.6	69.6	137.2	237.9	353.4	504.9	672.9
White, male										
2005 4/.....	2.4	6.0	13.2	28.5	70.2	138.6	251.2	406.6	583.5	846.1
2004 4/.....	2.4	5.3	12.3	26.6	66.5	132.4	238.5	392.8	563.4	837.7
2003 4/.....	2.2	6.1	12.0	29.4	71.5	139.3	252.4	406.9	585.6	894.4
2002.....	2.5	5.6	12.4	29.6	67.5	137.1	254.3	417.4	612.9	923.4
2001.....	2.2	5.6	11.9	31.8	69.1	142.0	260.8	418.9	600.0	943.6
2000.....	2.2	5.7	11.5	31.8	69.8	143.1	265.8	415.5	648.5	997.4
1999.....	2.0	5.0	11.6	34.5	76.3	155.7	285.6	441.3	664.4	1,029.4
White, female										
2005 4/.....	2.6	6.3	11.7	26.9	61.4	115.1	202.4	307.0	417.6	585.4
2004 4/.....	2.5	5.1	11.0	25.6	60.1	112.2	194.4	289.7	391.9	536.5
2003 4/.....	2.8	5.1	11.4	26.7	63.5	120.8	199.3	304.0	413.2	562.8
2002.....	2.5	5.4	11.0	28.3	60.4	117.6	198.0	296.4	408.9	554.5
2001.....	2.6	4.6	11.6	28.6	63.5	122.0	199.7	294.5	397.1	557.9
2000.....	2.4	4.9	11.3	28.4	61.5	119.4	200.5	292.1	404.4	544.8
1999.....	2.5	4.7	12.0	30.8	63.5	121.1	200.0	291.6	410.7	529.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Chronic lower respiratory diseases (J40-J47)

Black, both sexes

2005 4/.....	5.0	10.5	17.9	30.4	52.4	92.0	142.8	220.8	271.1	366.7
2004 4/.....	4.8	9.6	16.3	27.9	52.1	81.7	128.1	204.6	249.3	333.9
2003 4/.....	5.7	9.1	16.7	31.3	52.7	86.5	146.7	211.9	271.9	353.0
2002.....	6.5	9.1	17.4	30.9	53.4	86.1	148.1	215.9	298.5	368.2
2001.....	5.1	9.1	16.6	31.6	52.2	94.3	149.6	206.9	276.5	375.8
2000.....	6.1	11.0	18.2	33.0	57.4	97.3	151.2	211.6	271.3	373.8
1999.....	6.1	11.2	17.2	36.1	57.6	100.0	163.3	242.4	295.3	376.2

Black, male

2005 4/.....	4.9	11.0	20.1	37.5	66.0	126.0	198.9	345.5	422.8	594.9
2004 4/.....	4.9	9.9	18.9	37.7	67.6	109.0	182.3	300.7	392.4	566.3
2003 4/.....	5.1	9.8	17.7	38.0	67.7	112.0	212.5	326.8	471.8	601.6
2002.....	6.9	9.9	19.9	38.1	71.1	116.2	214.0	340.8	477.6	645.4
2001.....	5.1	8.9	18.8	37.1	66.3	127.1	222.2	329.9	468.7	671.7
2000.....	5.9	11.4	19.6	39.5	73.5	138.4	226.3	326.5	448.5	693.1
1999.....	6.2	11.9	19.7	43.2	79.3	139.4	243.5	387.0	521.4	701.9

Black, female

2005 4/.....	5.1	10.1	16.1	24.6	41.8	66.8	104.3	146.4	191.0	274.6
2004 4/.....	4.8	9.3	14.2	19.9	40.1	61.3	91.4	147.7	173.9	241.8
2003 4/.....	6.2	8.6	15.8	25.7	41.1	67.6	102.4	143.5	167.1	256.4
2002.....	6.1	8.5	15.3	25.1	39.6	63.8	104.0	141.3	205.3	262.0
2001.....	5.1	9.2	14.7	27.1	41.1	70.0	101.3	133.4	178.7	266.1
2000.....	6.2	10.6	16.9	27.7	44.6	67.0	101.2	142.4	182.1	255.0
1999.....	6.1	10.7	15.0	30.2	40.6	70.8	110.0	155.3	183.2	254.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Bronchitis, chronic and unspecified (J40-J42)										
All races 3/, both sexes										
2005.....	*	*	*	0.2	0.3	0.5	0.9	1.1	2.2	7.4
2004.....	*	*	0.1	0.2	0.3	0.4	0.7	1.2	2.0	6.5
2003.....	*	*	*	0.1	0.2	0.4	0.7	1.4	2.6	7.3
2002.....	*	*	0.1	0.1	0.3	0.5	1.0	1.8	2.9	8.1
2001.....	*	0.1	0.1	0.2	0.3	0.6	0.9	1.7	2.6	9.1
2000.....	*	*	*	0.1	0.4	0.7	1.0	2.1	3.6	11.8
1999.....	*	*	0.1	0.3	0.6	0.7	1.2	1.9	4.1	10.7
All races 3/, male										
2005.....	*	*	*	*	0.4	*	0.9	1.4	2.5	7.3
2004.....	*	*	*	*	*	0.5	0.8	1.2	2.2	7.0
2003.....	*	*	*	*	*	*	0.8	1.9	2.6	7.0
2002.....	*	*	*	*	*	0.7	1.1	2.2	2.9	8.8
2001.....	*	*	*	*	0.4	0.8	1.1	2.2	2.5	9.2
2000.....	*	*	*	*	0.4	0.9	1.0	2.8	4.9	12.9
1999.....	*	*	*	0.4	0.5	0.8	1.5	2.4	4.7	11.9
All races 3/, female										
2005.....	*	*	*	0.2	*	0.5	0.8	0.8	2.0	7.4
2004.....	*	*	*	*	*	*	0.6	1.2	1.9	6.3
2003.....	*	*	*	*	*	0.4	0.7	1.1	2.5	7.5
2002.....	*	*	*	*	*	*	0.9	1.4	2.9	7.8
2001.....	*	*	*	*	*	0.4	0.7	1.4	2.6	9.1
2000.....	*	*	*	*	0.4	0.5	1.1	1.6	2.8	11.4
1999.....	*	*	*	*	0.6	0.5	1.0	1.5	3.8	10.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Bronchitis, chronic and unspecified (J40-J42)										
White, both sexes										
2005 4/.....	*	*	*	0.1	0.3	0.5	0.9	1.1	2.2	7.9
2004 4/.....	*	*	*	0.2	0.3	0.4	0.7	1.2	2.1	6.7
2003 4/.....	*	*	*	*	0.2	0.5	0.7	1.5	2.6	7.5
2002.....	*	*	*	*	0.3	0.6	1.1	1.9	3.0	8.6
2001.....	*	*	0.1	0.2	0.3	0.6	0.9	1.8	2.7	9.6
2000.....	*	*	*	*	0.5	0.7	1.0	2.2	3.7	12.7
1999.....	*	*	*	0.3	0.6	0.7	1.2	1.9	4.1	11.2
White, male										
2005 4/.....	*	*	*	*	0.4	*	0.9	1.4	2.5	7.4
2004 4/.....	*	*	*	*	*	0.6	0.8	1.2	2.4	7.1
2003 4/.....	*	*	*	*	*	*	0.7	1.9	2.6	7.3
2002.....	*	*	*	*	*	0.7	1.2	2.4	2.9	9.3
2001.....	*	*	*	*	*	0.8	1.0	2.2	2.5	9.4
2000.....	*	*	*	*	0.5	1.0	1.0	2.8	5.2	13.8
1999.....	*	*	*	0.4	0.5	0.9	1.5	2.4	4.4	12.4
White, female										
2005 4/.....	*	*	*	*	*	0.6	0.9	0.9	2.1	8.1
2004 4/.....	*	*	*	*	*	*	0.6	1.2	2.0	6.5
2003 4/.....	*	*	*	*	*	0.5	0.8	1.2	2.6	7.6
2002.....	*	*	*	*	*	*	1.0	1.6	3.0	8.3
2001.....	*	*	*	*	*	*	0.7	1.5	2.8	9.7
2000.....	*	*	*	*	0.5	0.5	1.1	1.7	2.8	12.2
1999.....	*	*	*	*	0.7	0.5	1.0	1.5	3.9	10.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Emphysema (J43)										
All races 3/, both sexes										
2005.....	0.2	0.8	1.4	3.7	8.0	15.3	25.8	37.8	46.0	51.9
2004.....	0.2	0.6	1.5	3.4	8.1	15.8	26.1	37.5	45.7	49.9
2003.....	0.3	0.7	1.5	4.0	9.3	16.6	28.5	41.9	51.6	54.6
2002.....	0.3	0.7	1.6	4.3	9.1	17.4	31.3	43.2	54.7	59.3
2001.....	0.2	0.7	1.7	5.0	10.1	19.8	33.2	46.9	55.5	62.1
2000.....	0.3	0.8	1.8	5.0	10.4	20.7	35.5	47.5	60.4	63.3
1999.....	0.4	0.7	2.0	5.5	12.1	23.5	38.4	51.3	65.3	62.1
All races 3/, male										
2005.....	0.3	1.1	1.6	4.4	9.7	18.4	30.7	47.5	59.5	71.5
2004.....	0.3	0.8	1.6	3.8	9.5	18.4	31.6	46.5	60.5	74.4
2003.....	0.3	1.0	1.9	4.4	10.3	18.3	34.4	51.9	67.5	79.2
2002.....	0.5	0.9	1.9	4.6	10.7	19.7	37.8	56.1	74.9	90.0
2001.....	0.3	0.9	2.0	5.8	10.8	22.4	40.3	59.2	74.9	101.2
2000.....	0.3	1.0	2.1	5.6	11.2	24.5	42.6	60.4	83.6	104.2
1999.....	0.4	0.9	2.2	6.2	13.6	28.0	48.9	67.3	91.5	105.4
All races 3/, female										
2005.....	*	0.5	1.1	3.0	6.5	12.6	21.9	30.8	37.6	42.8
2004.....	0.2	0.4	1.3	3.1	6.8	13.5	21.7	31.1	36.6	38.9
2003.....	0.3	0.5	1.2	3.6	8.4	15.2	23.7	34.8	41.9	43.7
2002.....	0.2	0.6	1.3	4.0	7.7	15.4	26.1	34.0	42.4	46.0
2001.....	*	0.5	1.4	4.2	9.5	17.6	27.5	38.3	43.8	45.8
2000.....	0.2	0.6	1.5	4.5	9.7	17.5	29.9	38.5	46.8	46.7
1999.....	0.3	0.5	1.7	4.9	10.8	19.8	30.2	40.2	49.9	44.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Emphysema (J43)										
White, both sexes										
2005 4/.....	0.2	0.8	1.3	3.9	8.6	16.4	27.9	39.9	48.2	54.5
2004 4/.....	0.2	0.6	1.4	3.6	8.6	17.2	28.2	40.1	48.4	52.5
2003 4/.....	0.3	0.7	1.6	4.2	9.9	18.0	30.5	44.3	54.2	57.0
2002.....	0.3	0.7	1.7	4.6	9.8	18.6	33.3	45.8	57.4	62.3
2001.....	0.2	0.7	1.7	5.1	10.9	21.5	35.2	49.8	58.7	65.1
2000.....	0.2	0.7	1.8	5.3	11.2	22.0	37.9	50.2	63.6	65.8
1999.....	0.3	0.6	2.0	5.8	13.0	25.4	40.9	53.8	68.8	64.7
White, male										
2005 4/.....	0.3	1.1	1.5	4.5	10.1	19.2	32.4	49.3	60.6	73.6
2004 4/.....	0.3	0.8	1.6	3.8	9.8	19.4	33.4	48.5	63.2	77.1
2003 4/.....	0.3	1.0	1.9	4.5	10.6	19.5	35.9	53.2	69.6	80.4
2002.....	0.4	0.9	1.9	4.8	11.1	20.2	39.5	58.2	77.4	93.0
2001.....	0.3	1.0	1.9	5.8	11.3	23.9	41.6	61.7	78.0	105.1
2000.....	0.2	0.9	2.0	5.8	11.8	25.0	44.6	63.1	87.0	105.5
1999.....	0.3	0.8	2.1	6.4	14.2	29.4	50.6	69.6	94.4	109.2
White, female										
2005 4/.....	*	0.5	1.2	3.3	7.2	13.8	24.2	33.0	40.5	45.7
2004 4/.....	*	0.4	1.3	3.4	7.5	15.2	24.0	33.9	39.2	41.4
2003 4/.....	0.3	0.5	1.4	3.9	9.2	16.7	26.0	37.8	44.7	46.6
2002.....	*	0.6	1.5	4.4	8.6	17.3	28.3	37.0	45.1	48.9
2001.....	*	0.5	1.4	4.5	10.6	19.4	30.0	41.3	47.0	48.4
2000.....	0.2	0.6	1.5	4.8	10.7	19.3	32.5	41.1	49.8	49.7
1999.....	0.3	0.5	1.9	5.3	11.9	21.9	33.1	42.6	53.7	46.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Emphysema (J43)										
Black, both sexes										
2005 4/.....	*	0.9	1.9	3.5	6.0	11.6	13.6	26.2	26.2	27.3
2004 4/.....	*	*	2.1	3.1	6.5	8.4	13.7	20.7	24.3	27.6
2003 4/.....	*	1.0	1.3	3.6	7.2	10.2	16.4	25.2	30.5	30.9
2002.....	0.8	1.0	1.5	3.3	5.9	11.9	20.9	23.5	34.8	32.1
2001.....	*	*	2.3	4.9	6.4	11.6	20.1	21.9	26.3	35.3
2000.....	*	1.6	2.5	4.7	7.7	14.9	19.5	24.5	30.8	37.2
1999.....	0.8	1.4	2.4	4.6	8.0	14.0	20.8	31.4	31.2	37.6
Black, male										
2005 4/.....	*	*	2.8	5.6	9.8	17.8	20.2	39.8	45.6	51.1
2004 4/.....	*	*	3.0	4.9	9.7	13.8	21.4	34.6	39.5	56.6
2003 4/.....	*	*	2.0	5.2	10.2	13.3	24.6	43.4	50.2	70.9
2002.....	*	*	2.4	4.5	9.3	20.4	30.7	41.3	61.1	64.3
2001.....	*	*	2.9	6.9	9.6	15.8	31.6	34.4	49.0	68.1
2000.....	*	2.4	3.5	5.9	10.8	25.2	30.1	35.0	53.1	87.6
1999.....	*	2.0	4.1	6.4	11.9	22.8	37.0	48.9	65.4	72.6
Black, female										
2005 4/.....	*	*	*	*	3.0	7.1	9.1	18.1	16.0	17.6
2004 4/.....	*	*	*	*	3.9	4.3	8.6	12.4	16.3	16.1
2003 4/.....	*	*	*	2.3	4.8	7.9	10.9	14.4	20.1	15.4
2002.....	*	*	*	*	3.3	5.6	14.3	12.9	21.1	19.7
2001.....	*	*	*	3.3	4.0	8.4	12.3	14.4	14.7	23.2
2000.....	*	*	*	3.6	5.2	7.3	12.4	18.2	19.5	18.5
1999.....	*	*	*	3.0	5.0	7.5	10.1	20.9	14.3	24.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Asthma (J45-J46)										
All races 3/, both sexes										
2005.....	1.0	1.4	1.4	1.5	1.9	2.3	2.9	4.1	7.1	14.5
2004.....	0.9	1.3	1.4	1.7	1.9	2.5	3.0	4.2	6.6	14.1
2003.....	1.1	1.5	1.7	1.9	2.2	2.7	3.1	4.9	7.2	14.2
2002.....	1.2	1.4	1.6	1.9	2.1	2.7	3.8	5.4	7.7	14.7
2001.....	1.1	1.3	1.8	1.9	2.6	3.0	3.9	5.2	8.1	16.1
2000.....	1.2	1.6	1.7	2.0	2.5	3.4	4.6	5.7	8.7	16.7
1999.....	1.1	1.6	1.7	2.6	2.2	3.7	4.7	6.2	9.6	17.3
All races 3/, male										
2005.....	0.8	0.9	1.2	1.1	1.2	1.7	1.9	3.7	4.8	8.7
2004.....	0.8	1.1	1.0	1.4	1.3	2.0	2.3	2.8	4.4	11.1
2003.....	0.9	1.2	1.2	1.5	1.7	1.8	2.4	3.7	5.7	10.7
2002.....	0.9	1.1	1.3	1.5	1.7	2.0	2.8	4.3	6.2	10.9
2001.....	0.9	1.0	1.3	1.2	1.9	2.4	3.0	3.8	5.6	12.1
2000.....	1.0	1.3	1.3	1.6	1.8	2.7	3.4	3.9	6.7	12.9
1999.....	0.8	1.0	1.2	1.9	1.7	2.6	3.9	4.8	7.2	12.7
All races 3/, female										
2005.....	1.2	1.8	1.7	1.9	2.6	2.8	3.7	4.4	8.6	17.2
2004.....	1.1	1.5	1.8	2.0	2.5	2.9	3.6	5.1	7.9	15.4
2003.....	1.4	1.7	2.1	2.2	2.6	3.5	3.7	5.7	8.0	15.8
2002.....	1.4	1.8	1.9	2.3	2.5	3.3	4.5	6.2	8.7	16.3
2001.....	1.3	1.7	2.2	2.5	3.3	3.6	4.7	6.1	9.6	17.8
2000.....	1.4	1.9	2.1	2.4	3.1	4.0	5.5	6.9	9.8	18.2
1999.....	1.3	2.2	2.2	3.3	2.7	4.6	5.4	7.2	11.1	19.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Asthma (J45-J46)										
White, both sexes										
2005 4/.....	0.8	1.0	1.1	1.1	1.6	1.8	2.3	3.8	6.4	14.5
2004 4/.....	0.7	0.8	1.0	1.3	1.4	2.1	2.8	3.6	6.2	13.7
2003 4/.....	0.8	1.1	1.1	1.5	1.8	2.3	2.6	4.4	6.7	14.3
2002.....	0.8	1.1	1.1	1.4	1.8	2.2	3.2	5.1	7.5	14.7
2001.....	0.8	1.0	1.4	1.4	1.9	2.6	3.4	4.7	7.8	15.7
2000.....	0.9	1.1	1.2	1.4	2.0	2.8	4.0	5.2	8.4	15.9
1999.....	0.8	1.2	1.3	2.0	1.8	3.1	4.0	5.8	9.4	16.6
White, male										
2005 4/.....	0.6	0.6	1.0	0.8	0.9	1.3	1.4	3.1	4.1	8.6
2004 4/.....	0.6	0.6	0.7	0.9	0.8	1.6	1.9	2.2	3.8	10.3
2003 4/.....	0.6	0.9	0.8	1.2	1.2	1.5	2.0	3.0	5.4	10.6
2002.....	0.6	0.8	0.9	1.0	1.4	1.5	2.3	3.8	5.8	10.7
2001.....	0.6	0.7	0.9	1.0	1.2	2.0	2.4	3.3	5.1	11.3
2000.....	0.7	1.0	1.0	1.0	1.4	2.0	3.0	3.7	6.3	11.4
1999.....	0.5	0.7	0.9	1.5	1.2	2.0	3.0	4.2	7.0	10.8
White, female										
2005 4/.....	0.9	1.3	1.2	1.4	2.2	2.2	3.0	4.3	7.9	17.2
2004 4/.....	0.8	1.0	1.4	1.6	2.1	2.6	3.5	4.7	7.7	15.2
2003 4/.....	1.0	1.3	1.5	1.7	2.3	3.0	3.1	5.4	7.5	15.9
2002.....	1.0	1.3	1.3	1.7	2.2	2.8	3.9	6.1	8.6	16.4
2001.....	1.0	1.2	1.8	1.9	2.5	3.2	4.2	5.7	9.4	17.6
2000.....	1.0	1.2	1.5	1.8	2.6	3.4	4.8	6.3	9.7	17.7
1999.....	1.1	1.7	1.7	2.6	2.2	4.0	4.8	6.9	10.8	18.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Asthma (J45-J46)										
Black, both sexes										
2005 4/.....	2.8	4.4	3.9	5.4	5.3	6.0	8.7	7.9	12.6	11.8
2004 4/.....	2.9	4.8	4.1	5.1	6.3	5.7	5.6	9.0	9.4	15.5
2003 4/.....	3.4	4.1	5.7	5.2	5.0	6.6	7.5	8.2	10.4	12.0
2002.....	4.0	4.1	5.7	7.1	5.6	6.1	10.1	7.4	8.9	11.9
2001.....	3.1	4.3	4.9	5.9	7.9	6.3	9.4	10.2	11.1	16.3
2000.....	3.6	5.4	5.3	6.6	7.2	8.8	9.7	10.7	9.9	22.2
1999.....	3.3	4.8	5.8	7.3	6.2	8.4	11.1	9.8	11.3	20.4
Black, male										
2005 4/.....	2.5	3.3	2.5	4.3	*	4.8	6.3	9.8	*	*
2004 4/.....	2.7	4.7	2.9	4.8	5.8	4.8	6.5	*	*	*
2003 4/.....	2.7	3.5	4.4	3.8	4.8	4.9	*	9.1	*	*
2002.....	3.7	3.4	4.6	6.0	5.3	6.2	8.7	*	*	*
2001.....	2.9	3.8	4.5	3.6	5.9	*	9.8	*	*	*
2000.....	3.1	4.0	3.5	6.3	6.0	8.6	7.0	*	*	25.4
1999.....	2.9	3.6	4.2	5.1	5.7	6.8	12.2	9.5	*	*
Black, female										
2005 4/.....	3.1	5.3	5.1	6.4	6.7	6.9	10.4	6.7	13.0	13.5
2004 4/.....	3.1	4.9	5.1	5.4	6.7	6.3	5.1	9.7	9.1	16.9
2003 4/.....	4.1	4.7	6.7	6.3	5.1	7.9	8.6	7.6	11.6	13.8
2002.....	4.2	4.7	6.6	8.1	5.7	5.9	11.0	8.0	9.5	12.3
2001.....	3.3	4.8	5.3	7.8	9.4	7.5	9.2	11.4	11.9	16.8
2000.....	4.0	6.6	6.9	6.9	8.2	9.0	11.5	14.2	10.2	21.0
1999.....	3.6	5.9	7.1	9.1	6.5	9.6	10.4	10.0	13.8	21.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other chronic lower respiratory diseases (J44,J47)										
All races 3/, both sexes										
2005.....	1.5	4.3	9.7	21.5	51.8	100.6	180.7	287.5	402.5	563.5
2004.....	1.4	3.6	8.8	20.2	49.7	95.1	170.8	272.9	380.4	531.1
2003.....	1.4	3.6	8.6	21.5	52.2	101.4	178.5	281.8	394.6	558.9
2002.....	1.4	3.5	8.7	21.8	49.3	98.4	175.4	279.8	400.3	555.5
2001.....	1.3	3.3	8.4	22.4	49.8	100.7	177.4	275.4	386.8	557.4
2000.....	1.2	3.4	8.2	22.4	49.1	99.2	177.1	272.5	400.7	556.7
1999.....	1.2	3.1	8.2	23.5	51.6	102.5	183.2	280.8	405.5	555.8
All races 3/, male										
2005.....	1.4	4.3	10.7	22.8	56.3	112.4	206.5	341.0	497.9	732.4
2004.....	1.5	3.6	9.8	21.5	53.5	105.2	193.5	328.2	476.3	718.6
2003.....	1.3	4.1	9.1	23.3	57.0	112.2	206.5	338.3	493.3	769.0
2002.....	1.6	3.8	9.5	23.3	53.2	109.1	204.3	343.0	511.9	785.1
2001.....	1.3	3.8	9.0	24.2	54.0	111.6	208.6	341.5	500.4	794.5
2000.....	1.2	3.8	8.7	24.4	54.5	111.6	210.9	337.1	533.3	838.0
1999.....	1.2	3.7	8.6	25.8	58.7	119.6	223.9	357.9	544.9	866.7
All races 3/, female										
2005.....	1.5	4.3	8.8	20.4	47.7	90.2	159.8	248.7	343.2	485.9
2004.....	1.4	3.6	7.8	18.9	46.2	86.2	152.5	233.1	321.3	446.8
2003.....	1.4	3.2	8.0	19.8	47.8	92.1	156.0	241.5	334.2	466.0
2002.....	1.3	3.3	7.9	20.4	45.7	89.3	152.3	235.2	332.4	455.8
2001.....	1.4	2.8	7.8	20.7	46.1	91.4	152.7	229.1	318.6	458.2
2000.....	1.2	2.9	7.8	20.5	44.3	88.7	150.4	227.6	322.5	442.2
1999.....	1.1	2.6	7.8	21.4	45.2	87.9	151.4	227.2	324.0	430.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other chronic lower respiratory diseases (J44,J47)										
White, both sexes										
2005 4/.....	1.5	4.4	9.9	22.5	55.1	107.6	193.4	304.5	424.6	590.7
2004 4/.....	1.5	3.7	9.1	21.1	52.9	102.0	182.6	288.3	400.9	557.2
2003 4/.....	1.3	3.8	8.9	22.3	55.5	108.7	189.4	297.0	415.4	585.8
2002.....	1.4	3.6	8.8	22.9	51.9	105.4	185.7	294.0	418.6	580.7
2001.....	1.4	3.3	8.6	23.4	53.1	106.6	187.6	289.8	404.4	581.3
2000.....	1.2	3.4	8.4	23.2	51.8	105.0	186.7	285.5	419.7	581.2
1999.....	1.1	3.0	8.4	24.4	54.2	108.1	191.8	292.0	422.6	580.3
White, male										
2005 4/.....	1.4	4.3	10.6	23.1	58.7	117.8	216.4	352.8	516.3	756.5
2004 4/.....	1.5	3.7	9.9	21.7	55.6	110.7	202.4	340.9	494.0	743.2
2003 4/.....	1.2	4.2	9.3	23.6	59.4	117.8	213.8	348.8	507.9	796.0
2002.....	1.6	3.8	9.5	23.7	54.7	114.7	211.4	353.0	526.8	810.4
2001.....	1.3	3.9	9.0	24.8	56.2	115.4	215.7	351.7	514.4	817.8
2000.....	1.2	3.8	8.5	24.9	56.2	115.1	217.3	345.8	550.1	866.7
1999.....	1.1	3.5	8.6	26.2	60.3	123.4	230.5	365.1	558.5	897.0
White, female										
2005 4/.....	1.5	4.5	9.2	22.0	51.8	98.5	174.4	268.8	367.2	514.4
2004 4/.....	1.4	3.6	8.2	20.5	50.3	94.2	166.3	249.9	343.0	473.4
2003 4/.....	1.5	3.3	8.5	21.0	51.8	100.6	169.5	259.5	358.4	492.7
2002.....	1.3	3.5	8.2	22.1	49.3	97.1	164.8	251.8	352.3	480.9
2001.....	1.4	2.8	8.2	22.1	50.2	98.9	164.8	246.0	337.9	482.3
2000.....	1.1	3.0	8.2	21.6	47.7	96.1	162.1	243.0	342.1	465.2
1999.....	1.1	2.6	8.3	22.7	48.7	94.7	161.1	240.6	342.4	453.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other chronic lower respiratory diseases (J44,J47)										
Black, both sexes										
2005 4/.....	1.8	5.1	12.0	21.2	40.8	73.9	119.9	186.1	230.8	326.1
2004 4/.....	1.5	4.1	9.8	19.5	39.0	67.5	108.2	174.2	214.9	285.9
2003 4/.....	1.8	3.8	9.4	22.1	40.4	69.5	122.2	178.0	228.7	304.6
2002.....	1.7	3.9	10.1	20.3	41.8	67.8	116.9	184.3	252.5	320.1
2001.....	1.6	4.1	9.4	20.7	37.6	75.7	119.1	173.3	238.4	320.9
2000.....	1.9	3.9	10.1	21.7	42.2	72.8	120.7	174.7	227.9	310.6
1999.....	2.0	4.8	8.5	23.9	43.1	77.1	129.9	199.1	248.8	313.7
Black, male										
2005 4/.....	1.7	6.3	14.7	27.3	52.4	102.7	171.4	294.1	363.1	532.6
2004 4/.....	1.5	3.9	12.3	28.1	51.6	90.1	153.8	257.5	342.9	492.7
2003 4/.....	1.9	4.5	11.0	28.8	52.4	93.8	181.3	273.0	410.5	520.3
2002.....	2.0	4.8	12.7	27.5	56.2	88.8	174.3	291.6	407.3	565.0
2001.....	1.6	4.4	11.3	26.6	50.4	105.7	179.4	285.3	408.5	583.2
2000.....	1.7	5.0	12.5	27.3	56.1	104.1	187.9	284.9	383.6	576.6
1999.....	2.5	6.0	11.0	31.2	61.3	109.6	192.3	325.8	441.2	607.0
Black, female										
2005 4/.....	1.8	4.1	9.8	16.1	31.7	52.5	84.6	121.6	161.0	242.7
2004 4/.....	1.4	4.2	7.6	12.4	29.1	50.7	77.3	124.8	147.4	204.0
2003 4/.....	1.7	3.1	8.0	16.6	31.0	51.5	82.4	121.5	133.5	220.8
2002.....	1.4	3.1	8.0	14.4	30.5	52.2	78.4	120.2	171.9	226.3
2001.....	1.5	3.8	7.7	15.8	27.5	53.5	78.9	106.5	151.7	223.5
2000.....	1.9	2.8	8.0	17.0	31.2	49.7	76.1	108.2	149.4	211.6
1999.....	1.5	3.7	6.5	17.9	28.8	53.1	88.4	122.7	153.4	204.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Pneumonitis due to solids and liquids (J69)										
All races 3/, both sexes										
2005.....	0.4	0.8	1.5	2.0	3.7	6.4	12.9	28.5	61.1	163.9
2004.....	0.5	0.8	1.2	1.9	3.4	6.0	13.7	28.4	62.2	164.9
2003.....	0.4	0.9	1.1	2.0	3.8	6.3	13.3	30.1	65.4	176.2
2002.....	0.5	0.7	1.1	1.8	3.4	6.5	13.4	31.3	67.3	186.0
2001.....	0.5	0.7	1.3	1.8	3.6	6.2	14.1	30.4	68.0	189.4
2000.....	0.5	0.8	1.2	1.9	3.2	6.7	14.2	30.4	65.7	187.6
1999.....	0.5	0.7	0.9	2.0	3.1	6.1	13.2	28.2	60.7	175.6
All races 3/, male										
2005.....	0.4	0.9	1.8	2.4	4.9	8.4	17.5	38.3	86.2	229.3
2004.....	0.6	1.0	1.5	2.3	4.5	8.1	18.7	39.0	87.4	237.2
2003.....	0.4	1.0	1.4	2.6	4.8	8.6	17.8	41.5	92.3	246.2
2002.....	0.7	0.9	1.5	2.2	4.2	8.6	18.2	43.4	95.1	268.5
2001.....	0.7	0.8	1.7	2.3	4.2	8.2	19.1	41.0	98.1	270.3
2000.....	0.6	1.0	1.5	2.4	3.9	9.2	20.2	42.4	95.2	276.0
1999.....	0.6	0.9	1.0	2.4	4.2	8.3	18.4	41.6	90.1	258.4
All races 3/, female										
2005.....	0.4	0.7	1.1	1.6	2.7	4.6	9.1	21.3	45.5	133.9
2004.....	0.3	0.5	0.8	1.5	2.5	4.2	9.8	20.9	46.6	132.4
2003.....	0.4	0.7	0.8	1.5	2.9	4.3	9.7	22.1	49.0	145.3
2002.....	0.4	0.5	0.8	1.4	2.8	4.8	9.6	22.8	50.4	150.2
2001.....	0.4	0.6	0.9	1.4	3.0	4.4	10.2	23.0	49.9	155.6
2000.....	0.4	0.6	0.9	1.5	2.5	4.6	9.4	22.1	48.2	151.6
1999.....	0.4	0.5	0.8	1.6	2.0	4.2	9.2	18.8	43.5	142.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Pneumonitis due to solids and liquids (J69)										
White, both sexes										
2005 4/.....	0.5	0.8	1.4	2.0	3.5	6.0	12.7	27.9	61.4	166.5
2004 4/.....	0.4	0.7	1.1	1.8	3.1	5.9	13.4	28.2	62.9	167.8
2003 4/.....	0.4	0.8	1.1	2.0	3.3	5.9	13.1	29.7	65.0	178.7
2002.....	0.5	0.7	1.0	1.8	3.3	6.1	13.1	30.9	68.1	189.5
2001.....	0.5	0.7	1.2	1.8	3.5	6.0	13.6	30.2	68.1	190.8
2000.....	0.5	0.7	1.1	1.9	2.9	6.4	13.4	30.1	65.7	191.0
1999.....	0.4	0.6	0.8	1.8	2.9	5.8	12.5	27.6	60.3	178.0
White, male										
2005 4/.....	0.5	0.9	1.7	2.4	4.5	7.8	17.1	37.1	86.1	234.2
2004 4/.....	0.5	1.0	1.4	2.2	3.9	7.8	17.9	38.3	88.4	241.7
2003 4/.....	0.4	1.0	1.4	2.5	4.2	8.2	17.2	40.6	91.7	250.8
2002.....	0.6	0.8	1.3	2.1	4.1	8.0	17.6	42.5	96.5	274.6
2001.....	0.7	0.8	1.5	2.3	4.0	8.0	18.4	40.6	98.7	275.4
2000.....	0.7	0.9	1.3	2.3	3.6	8.5	18.7	41.4	96.2	284.0
1999.....	0.5	0.8	0.9	2.2	3.9	7.7	17.1	40.4	88.9	263.6
White, female										
2005 4/.....	0.4	0.7	1.1	1.6	2.6	4.5	9.0	21.1	45.9	135.3
2004 4/.....	0.3	0.5	0.8	1.5	2.3	4.2	9.7	20.8	47.0	134.5
2003 4/.....	0.4	0.7	0.8	1.5	2.5	3.9	9.7	21.8	48.6	146.7
2002.....	0.4	0.5	0.7	1.5	2.6	4.5	9.5	22.6	50.7	152.5
2001.....	0.4	0.5	0.8	1.3	3.0	4.2	9.7	22.8	49.6	155.3
2000.....	0.4	0.5	0.9	1.4	2.3	4.6	9.1	22.2	47.5	153.2
1999.....	0.4	0.4	0.8	1.5	2.0	4.1	8.8	18.6	43.4	143.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Pneumonitis due to solids and liquids (J69)

Black, both sexes

2005 4/.....	*	1.2	2.4	3.1	6.4	10.4	17.0	40.9	69.2	153.1
2004 4/.....	0.9	1.0	1.7	3.0	7.3	8.8	20.4	34.7	64.9	144.7
2003 4/.....	0.9	1.3	1.2	2.7	8.5	11.0	17.2	39.8	76.2	163.4
2002.....	0.9	0.8	2.1	2.4	5.5	11.7	18.8	41.9	67.8	165.1
2001.....	0.7	1.0	2.4	2.4	6.0	9.3	22.3	38.6	76.1	185.5
2000.....	*	1.7	2.2	2.9	5.8	11.3	25.1	38.9	71.4	159.8
1999.....	1.0	1.2	1.8	3.9	5.0	10.1	23.7	40.5	71.1	167.1

Black, male

2005 4/.....	*	1.5	3.4	3.6	9.8	15.4	25.6	62.1	106.0	201.7
2004 4/.....	*	*	2.2	4.4	11.0	13.6	31.4	51.4	93.3	202.7
2003 4/.....	*	1.7	*	3.8	11.6	14.5	26.3	59.3	107.8	212.7
2002.....	*	*	3.1	3.7	6.3	16.7	28.4	62.5	91.7	233.7
2001.....	*	*	3.4	3.1	8.1	12.5	31.0	56.2	108.3	243.9
2000.....	*	2.4	3.1	3.3	7.2	17.4	41.1	62.4	96.1	209.9
1999.....	*	*	*	5.1	8.1	15.6	37.3	65.1	109.9	228.5

Black, female

2005 4/.....	*	*	*	2.7	3.7	6.7	11.0	28.3	49.7	133.5
2004 4/.....	*	*	*	*	4.4	5.2	13.0	24.8	49.9	121.7
2003 4/.....	*	*	*	*	6.0	8.4	11.1	28.2	59.6	144.2
2002.....	*	*	*	*	4.8	8.0	12.3	29.6	55.3	138.8
2001.....	*	*	*	*	4.3	6.9	16.6	28.0	59.7	163.8
2000.....	*	*	*	*	4.6	6.7	14.4	24.8	59.0	141.2
1999.....	*	*	*	2.9	*	6.0	14.6	25.7	51.9	144.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)										
All races 3/, both sexes										
2005.....	1.4	2.4	4.1	6.6	11.8	21.5	37.5	59.4	89.7	136.5
2004.....	1.5	2.6	4.0	7.3	12.4	21.7	39.5	61.4	91.3	139.4
2003.....	1.3	2.6	4.0	6.6	11.6	21.0	36.2	57.3	83.8	138.1
2002.....	1.5	2.5	4.1	7.0	11.9	20.6	37.6	56.8	83.7	133.5
2001.....	1.5	2.6	4.4	7.3	12.6	22.8	38.0	61.8	89.6	142.7
2000.....	1.7	2.6	4.6	7.8	12.7	22.9	36.9	60.5	89.2	141.7
1999.....	1.6	2.7	4.1	6.9	12.7	22.5	36.8	59.5	87.6	147.6
All races 3/, male										
2005.....	1.4	2.7	4.4	7.3	14.1	25.9	46.0	74.5	113.9	157.7
2004.....	1.5	2.7	4.5	8.1	14.3	26.6	49.4	77.3	118.1	168.8
2003.....	1.4	2.8	4.3	7.5	13.8	25.9	43.8	71.8	105.5	165.0
2002.....	1.6	2.5	4.3	7.9	13.9	24.4	47.8	70.5	104.2	156.6
2001.....	1.8	2.9	4.9	7.7	15.0	27.1	47.0	76.2	110.0	169.1
2000.....	1.8	2.8	5.0	8.8	14.6	26.8	45.7	77.3	112.4	179.7
1999.....	1.7	2.8	4.4	7.4	14.7	27.3	46.4	74.9	115.0	182.3
All races 3/, female										
2005.....	1.4	2.1	3.8	5.9	9.8	17.7	30.6	48.5	74.6	126.7
2004.....	1.5	2.6	3.4	6.6	10.7	17.4	31.6	50.0	74.8	126.1
2003.....	1.3	2.5	3.7	5.7	9.6	16.8	30.0	47.0	70.5	126.2
2002.....	1.4	2.4	3.9	6.2	10.2	17.3	29.4	47.0	71.1	123.4
2001.....	1.3	2.3	4.0	6.8	10.5	19.1	30.8	51.7	77.3	131.6
2000.....	1.6	2.4	4.2	6.8	10.8	19.6	29.9	48.8	75.6	126.2
1999.....	1.4	2.6	3.8	6.5	10.9	18.4	29.3	48.8	71.6	133.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)										
White, both sexes										
2005 4/.....	1.3	2.1	3.7	6.2	11.6	21.6	38.4	60.7	92.8	140.2
2004 4/.....	1.3	2.2	3.6	7.1	12.2	22.0	40.9	63.6	94.1	144.1
2003 4/.....	1.1	2.2	3.7	6.3	11.5	21.5	37.4	58.5	87.4	143.2
2002.....	1.3	2.1	3.8	6.7	11.6	21.0	38.2	58.7	85.5	137.9
2001.....	1.3	2.3	3.9	7.0	12.4	23.0	38.7	63.5	92.3	146.0
2000.....	1.6	2.2	4.3	7.5	12.4	22.8	37.7	62.0	91.2	145.9
1999.....	1.3	2.3	3.8	6.4	12.6	22.6	37.0	60.6	89.5	150.7
White, male										
2005 4/.....	1.2	2.4	4.0	6.8	13.6	26.3	47.2	75.5	117.0	160.0
2004 4/.....	1.3	2.2	4.3	7.8	14.2	27.0	50.8	79.7	121.5	174.3
2003 4/.....	1.2	2.5	4.1	7.1	13.7	26.4	45.2	72.7	110.0	170.5
2002.....	1.4	2.1	4.0	7.6	13.4	25.3	48.3	72.5	106.8	162.0
2001.....	1.7	2.5	4.4	7.6	14.5	27.5	48.3	78.4	113.3	173.5
2000.....	1.7	2.4	4.6	8.5	14.4	26.8	46.3	78.9	115.0	184.4
1999.....	1.5	2.4	4.2	6.7	14.7	27.7	47.0	76.5	117.6	185.7
White, female										
2005 4/.....	1.3	1.8	3.3	5.7	9.7	17.4	31.1	49.8	77.5	131.1
2004 4/.....	1.4	2.2	3.0	6.3	10.4	17.5	32.7	51.8	77.2	130.5
2003 4/.....	1.0	2.0	3.3	5.4	9.6	17.2	30.9	48.2	73.6	131.0
2002.....	1.2	2.0	3.6	5.9	10.0	17.2	30.0	48.7	72.5	127.4
2001.....	1.0	2.0	3.4	6.4	10.4	19.1	31.0	52.9	79.6	134.4
2000.....	1.5	2.1	3.9	6.6	10.6	19.3	30.7	50.1	77.1	130.3
1999.....	1.1	2.2	3.5	6.2	10.7	18.2	29.1	49.5	72.9	136.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)

Black, both sexes										
2005 4/.....	2.8	4.8	7.7	9.9	15.3	24.7	34.4	52.6	60.9	102.6
2004 4/.....	3.0	5.5	7.1	11.1	16.5	23.2	32.8	46.2	63.4	92.8
2003 4/.....	3.1	5.7	7.3	10.5	14.4	20.2	29.1	51.4	50.0	89.6
2002.....	2.9	5.5	6.9	11.1	16.8	20.7	33.9	40.5	68.3	90.3
2001.....	3.3	5.7	9.3	10.9	17.0	24.2	36.1	50.2	66.9	109.6
2000.....	3.0	6.0	7.8	11.6	15.5	28.1	33.4	49.8	71.2	98.2
1999.....	3.4	5.8	7.1	11.7	15.4	25.0	38.0	52.8	66.2	111.6
Black, male										
2005 4/.....	2.8	5.3	8.4	12.1	20.0	28.3	40.8	70.2	82.2	132.0
2004 4/.....	3.3	6.3	7.2	12.8	18.7	29.3	41.4	61.4	81.1	115.2
2003 4/.....	2.5	5.5	7.3	12.6	18.5	25.6	37.8	70.8	65.0	113.6
2002.....	3.4	5.7	7.6	12.4	21.2	22.2	44.4	53.3	86.4	100.8
2001.....	3.2	6.2	9.4	10.5	21.6	28.0	39.4	60.4	85.1	129.3
2000.....	3.3	7.2	8.6	13.3	17.6	32.5	40.5	67.6	82.6	132.6
1999.....	3.3	6.6	7.8	13.7	17.8	28.3	44.1	66.5	83.7	145.3
Black, female										
2005 4/.....	2.8	4.4	7.2	8.1	11.6	22.0	30.1	42.1	49.7	90.8
2004 4/.....	2.7	4.9	7.0	9.7	14.9	18.7	27.0	37.2	54.1	83.9
2003 4/.....	3.6	5.8	7.3	8.8	11.1	16.3	23.3	39.9	42.2	80.2
2002.....	2.3	5.3	6.2	10.0	13.3	19.6	26.9	32.9	58.9	86.3
2001.....	3.3	5.3	9.1	11.3	13.4	21.4	33.9	44.1	57.6	102.3
2000.....	2.6	5.0	7.1	10.3	13.9	24.8	28.6	39.0	65.4	85.4
1999.....	3.5	5.1	6.6	10.1	13.5	22.5	34.0	44.6	57.5	99.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Peptic ulcer (K25-K28)										
All races 3/, both sexes										
2005.....	0.3	0.5	0.8	0.9	1.6	2.2	3.8	5.8	10.8	21.4
2004.....	0.3	0.5	0.8	1.0	1.7	2.5	3.9	6.8	11.6	24.4
2003.....	0.3	0.6	0.8	1.2	1.7	2.3	4.2	7.3	13.6	26.4
2002.....	0.3	0.6	0.8	1.4	1.7	2.7	4.7	8.0	13.2	28.3
2001.....	0.4	0.7	0.8	1.3	2.1	3.1	4.8	8.3	16.0	33.4
2000.....	0.3	0.6	1.0	1.5	2.2	3.1	5.1	8.7	15.6	35.6
1999.....	0.4	0.6	0.8	1.4	2.2	3.3	5.9	9.1	17.1	34.6
All races 3/, male										
2005.....	0.5	0.8	1.2	1.3	2.2	2.6	4.8	7.3	13.1	22.4
2004.....	0.3	0.7	1.0	1.2	2.2	2.9	4.5	8.3	12.9	24.6
2003.....	0.4	0.8	1.2	1.4	2.3	3.1	5.4	8.8	16.2	28.9
2002.....	0.4	0.7	1.2	1.8	2.1	3.2	6.1	9.3	16.7	31.6
2001.....	0.5	0.9	1.1	1.8	2.8	4.1	5.6	10.3	17.3	34.9
2000.....	0.4	0.9	1.5	2.1	2.9	3.9	6.9	11.2	17.8	40.7
1999.....	0.4	0.7	1.2	1.8	3.0	4.5	7.9	11.0	21.1	37.5
All races 3/, female										
2005.....	0.2	0.3	0.5	0.6	1.1	1.8	3.0	4.7	9.4	21.0
2004.....	0.2	0.4	0.5	0.7	1.3	2.2	3.4	5.8	10.8	24.3
2003.....	0.2	0.4	0.5	1.0	1.1	1.7	3.2	6.2	11.9	25.3
2002.....	0.3	0.4	0.5	1.0	1.4	2.2	3.6	7.0	11.0	26.9
2001.....	0.2	0.4	0.6	0.8	1.5	2.2	4.2	6.9	15.2	32.7
2000.....	0.2	0.4	0.5	0.9	1.6	2.5	3.6	6.9	14.4	33.5
1999.....	0.3	0.5	0.5	1.1	1.5	2.3	4.3	7.7	14.7	33.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Peptic ulcer (K25-K28)										
White, both sexes										
2005 4/.....	0.3	0.4	0.8	0.9	1.6	2.2	3.7	5.7	10.9	21.6
2004 4/.....	0.2	0.4	0.7	0.9	1.7	2.5	3.8	6.8	11.7	25.1
2003 4/.....	0.3	0.5	0.7	1.1	1.6	2.4	4.2	7.2	13.4	26.9
2002.....	0.2	0.5	0.8	1.3	1.6	2.6	4.8	7.9	13.6	29.1
2001.....	0.3	0.6	0.7	1.1	2.0	3.0	4.8	8.3	16.2	34.4
2000.....	0.3	0.6	0.9	1.4	2.2	3.0	5.1	8.5	16.0	36.3
1999.....	0.3	0.5	0.7	1.4	2.2	3.1	5.8	9.1	17.1	35.3
White, male										
2005 4/.....	0.4	0.6	1.1	1.2	2.1	2.5	4.5	7.0	13.5	22.5
2004 4/.....	0.3	0.6	1.0	1.2	2.1	2.9	4.3	8.1	12.6	25.5
2003 4/.....	0.3	0.7	1.0	1.3	2.2	3.1	5.3	8.6	15.6	29.0
2002.....	0.3	0.6	1.0	1.6	2.0	3.1	6.0	9.2	17.1	32.5
2001.....	0.5	0.8	0.9	1.5	2.7	3.8	5.3	10.1	17.3	35.7
2000.....	0.4	0.8	1.3	2.0	2.9	3.8	6.7	10.9	17.8	40.7
1999.....	0.3	0.6	1.1	1.6	2.9	4.1	7.8	11.1	21.1	38.4
White, female										
2005 4/.....	*	0.2	0.5	0.7	1.1	1.9	3.0	4.7	9.3	21.2
2004 4/.....	*	0.3	0.5	0.7	1.3	2.1	3.3	5.8	11.1	25.0
2003 4/.....	0.2	0.3	0.5	0.9	1.0	1.7	3.3	6.2	12.1	26.0
2002.....	*	0.4	0.5	1.0	1.3	2.1	3.8	7.0	11.5	27.6
2001.....	*	0.4	0.6	0.8	1.4	2.2	4.4	7.0	15.6	33.9
2000.....	*	0.3	0.5	0.9	1.6	2.4	3.8	6.9	14.9	34.5
1999.....	0.3	0.4	0.4	1.1	1.5	2.2	4.2	7.7	14.8	34.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Peptic ulcer (K25-K28)										
Black, both sexes										
2005 4/.....	0.7	1.1	1.4	1.2	2.2	2.7	5.4	5.8	9.7	17.4
2004 4/.....	*	1.3	1.3	1.6	2.6	3.5	5.1	6.6	9.4	16.1
2003 4/.....	*	1.2	1.8	2.1	2.4	2.1	4.5	8.7	14.5	16.9
2002.....	0.9	0.9	1.6	2.3	2.9	3.2	5.2	8.8	6.8	19.0
2001.....	*	1.1	1.6	2.5	3.0	3.6	5.5	7.8	11.1	20.9
2000.....	*	1.4	2.0	2.6	2.7	4.1	5.7	9.4	11.0	26.6
1999.....	1.0	1.2	1.8	2.4	3.2	5.0	6.5	7.7	16.2	25.5
Black, male										
2005 4/.....	*	1.9	2.5	*	*	*	7.9	9.4	*	20.5
2004 4/.....	*	2.1	*	*	3.9	*	*	10.0	*	*
2003 4/.....	*	1.6	3.1	3.1	3.9	*	7.2	11.4	23.6	*
2002.....	*	*	2.8	3.6	4.2	*	8.0	11.9	*	*
2001.....	*	1.8	2.7	4.1	4.5	5.3	9.1	10.7	*	23.8
2000.....	*	*	3.3	3.5	*	*	10.0	12.8	*	31.1
1999.....	1.5	1.9	2.8	3.7	5.3	7.5	8.8	*	18.3	28.1
Black, female										
2005 4/.....	*	*	*	*	*	*	*	*	10.4	16.1
2004 4/.....	*	*	*	*	*	3.6	4.4	*	7.6	16.5
2003 4/.....	*	*	*	*	*	*	*	7.1	9.7	15.8
2002.....	*	*	*	*	*	*	*	6.9	*	18.5
2001.....	*	*	*	*	*	*	*	6.1	9.0	19.8
2000.....	*	*	*	*	*	*	*	7.4	8.5	24.9
1999.....	*	*	*	*	*	*	5.0	6.9	15.1	24.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Diseases of appendix (K35-K38)

All races 3/, both sexes

2005.....	*	*	0.1	0.2	0.2	0.4	0.4	0.8	0.9	2.1
2004.....	*	*	0.2	0.1	0.2	0.3	0.6	0.8	1.4	1.7
2003.....	*	0.1	0.1	*	0.2	0.3	0.4	0.8	1.4	1.7
2002.....	*	0.1	0.2	0.2	0.3	0.3	0.7	0.8	1.1	2.0
2001.....	*	0.1	*	0.2	0.2	0.4	0.6	0.5	1.0	2.1
2000.....	*	*	0.1	0.2	0.3	0.3	0.5	0.7	1.4	1.6
1999.....	*	0.1	0.1	0.2	0.2	0.3	0.4	0.6	1.1	1.5

All races 3/, male

2005.....	*	*	*	0.3	*	0.6	0.7	0.9	1.3	2.8
2004.....	*	*	0.2	*	*	*	0.7	1.1	1.7	2.5
2003.....	*	*	*	*	*	0.4	0.7	0.9	2.3	2.4
2002.....	*	*	0.2	*	0.5	0.5	0.8	1.1	1.6	2.2
2001.....	*	*	*	*	*	0.5	0.9	0.7	1.6	2.8
2000.....	*	*	*	0.3	0.4	*	0.6	1.0	1.6	2.0
1999.....	*	*	*	*	*	*	0.5	0.9	*	2.9

All races 3/, female

2005.....	*	*	*	*	*	*	*	0.7	0.7	1.8
2004.....	*	*	*	*	*	*	0.5	0.6	1.2	1.4
2003.....	*	*	*	*	*	*	*	0.6	0.9	1.3
2002.....	*	*	*	*	*	*	0.5	0.6	0.8	2.0
2001.....	*	*	*	*	*	*	0.4	*	0.7	1.8
2000.....	*	*	*	*	*	*	0.4	0.5	1.2	1.5
1999.....	*	*	*	*	*	*	*	*	1.1	0.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Diseases of appendix (K35-K38)

White, both sexes

2005 4/.....	*	*	*	0.2	0.2	0.4	0.4	0.7	1.0	2.2
2004 4/.....	*	*	0.1	0.2	0.2	0.3	0.6	0.8	1.3	1.8
2003 4/.....	*	*	*	*	*	0.3	0.4	0.7	1.5	1.7
2002.....	*	*	0.1	0.2	0.3	0.3	0.6	0.8	1.2	2.1
2001.....	*	*	*	*	*	0.3	0.7	0.5	0.9	2.1
2000.....	*	*	*	0.2	0.3	0.2	0.4	0.6	1.3	1.7
1999.....	*	*	0.2	*	*	0.3	0.4	0.6	1.0	1.5

White, male

2005 4/.....	*	*	*	*	*	0.6	0.7	0.9	1.4	2.9
2004 4/.....	*	*	*	*	*	*	0.7	1.1	1.7	2.3
2003 4/.....	*	*	*	*	*	*	0.7	0.8	2.4	2.5
2002.....	*	*	*	*	0.5	*	0.7	1.2	1.6	2.0
2001.....	*	*	*	*	*	*	1.0	0.7	1.5	2.5
2000.....	*	*	*	*	0.5	*	*	0.9	1.5	2.0
1999.....	*	*	*	*	*	*	*	0.9	*	3.1

White, female

2005 4/.....	*	*	*	*	*	*	*	0.6	0.7	1.9
2004 4/.....	*	*	*	*	*	*	0.5	0.6	1.0	1.6
2003 4/.....	*	*	*	*	*	*	*	0.6	0.9	1.3
2002.....	*	*	*	*	*	*	0.5	0.6	0.9	2.1
2001.....	*	*	*	*	*	*	*	*	*	1.9
2000.....	*	*	*	*	*	*	*	*	1.2	1.6
1999.....	*	*	*	*	*	*	*	*	1.1	0.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Hernia (K40-K46)										
All races 3/, both sexes										
2005.....	0.1	0.1	0.3	0.4	0.9	0.9	1.4	2.5	4.8	12.1
2004.....	0.1	0.2	0.2	0.4	0.5	0.9	1.8	2.9	5.3	11.0
2003.....	0.1	0.2	0.3	0.5	0.7	1.0	1.6	2.7	4.8	11.9
2002.....	0.1	0.2	0.2	0.4	0.6	0.9	1.6	2.8	5.3	12.8
2001.....	0.1	0.1	0.3	0.4	0.7	0.8	1.5	2.5	5.0	12.6
2000.....	*	0.1	0.3	0.4	0.6	0.9	1.4	3.0	4.9	13.1
1999.....	0.1	*	0.2	0.3	0.7	0.8	1.6	2.5	5.4	12.7
All races 3/, male										
2005.....	*	*	0.3	0.4	0.9	0.7	1.3	2.4	4.6	12.7
2004.....	*	0.2	0.3	0.4	0.5	0.8	2.2	3.2	4.7	11.3
2003.....	*	0.2	0.4	0.4	0.7	1.1	1.6	3.0	5.2	12.3
2002.....	*	*	*	0.4	0.5	0.9	1.6	2.7	5.2	11.7
2001.....	*	*	0.3	0.3	0.8	0.9	1.4	2.3	5.3	12.0
2000.....	*	*	0.4	0.3	0.6	0.9	1.4	3.0	5.7	12.6
1999.....	*	*	0.2	0.3	0.8	0.8	1.5	2.5	6.5	13.2
All races 3/, female										
2005.....	*	*	0.2	0.4	0.8	1.1	1.4	2.5	5.0	11.8
2004.....	*	0.2	0.2	0.4	0.5	1.0	1.4	2.7	5.6	10.8
2003.....	*	0.2	0.3	0.6	0.7	1.0	1.5	2.5	4.6	11.8
2002.....	*	*	0.3	0.4	0.7	1.0	1.5	2.9	5.3	13.3
2001.....	*	*	0.3	0.5	0.6	0.8	1.6	2.6	4.8	12.8
2000.....	*	*	0.3	0.4	0.6	0.9	1.5	3.0	4.4	13.3
1999.....	*	*	*	*	0.6	0.8	1.6	2.6	4.8	12.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Hernia (K40-K46)										
White, both sexes										
2005 4/.....	*	0.1	0.3	0.4	0.9	0.9	1.4	2.5	5.0	12.5
2004 4/.....	*	0.2	0.2	0.4	0.5	0.9	1.8	3.1	5.4	11.2
2003 4/.....	*	0.2	0.3	0.4	0.7	1.1	1.6	2.8	4.7	12.3
2002.....	*	0.1	0.2	0.4	0.6	0.9	1.6	3.0	5.4	13.4
2001.....	*	0.1	0.3	0.4	0.7	0.8	1.5	2.5	5.0	12.8
2000.....	*	0.1	0.3	0.4	0.6	0.8	1.5	3.0	5.0	13.3
1999.....	*	*	0.2	0.3	0.7	0.7	1.6	2.5	5.5	13.0
White, male										
2005 4/.....	*	*	0.3	0.4	0.9	0.7	1.4	2.4	4.6	12.8
2004 4/.....	*	*	0.3	0.4	0.5	0.7	2.2	3.3	4.7	11.4
2003 4/.....	*	*	0.4	0.4	0.7	1.1	1.6	3.1	5.1	12.5
2002.....	*	*	*	0.4	0.5	0.8	1.6	2.9	5.4	11.9
2001.....	*	*	0.3	*	0.7	0.9	1.4	2.3	5.2	11.8
2000.....	*	*	0.3	*	0.6	0.8	1.5	3.1	5.7	12.7
1999.....	*	*	*	0.4	0.8	0.8	1.6	2.5	6.6	12.9
White, female										
2005 4/.....	*	*	*	0.4	0.8	1.1	1.4	2.6	5.2	12.3
2004 4/.....	*	*	*	0.4	0.5	1.1	1.4	2.9	5.8	11.2
2003 4/.....	*	*	*	0.5	0.7	1.0	1.6	2.6	4.5	12.2
2002.....	*	*	*	0.4	0.6	0.9	1.6	3.0	5.4	14.0
2001.....	*	*	0.3	0.5	0.6	0.8	1.6	2.6	5.0	13.2
2000.....	*	*	0.3	0.5	0.6	0.8	1.5	2.9	4.6	13.6
1999.....	*	*	*	*	0.5	0.7	1.5	2.5	4.9	13.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Chronic liver disease and cirrhosis (K70,K73-K74)										
All races 3/, both sexes										
2005.....	8.6	15.0	20.7	23.0	24.1	25.4	29.3	29.8	27.9	19.7
2004.....	8.7	15.1	21.3	21.7	23.7	26.2	29.5	30.1	27.0	19.7
2003.....	9.3	15.9	21.1	22.0	24.4	28.0	31.2	31.0	28.5	20.1
2002.....	9.7	16.0	20.3	21.7	24.5	27.9	31.2	32.5	29.9	21.4
2001.....	9.8	17.1	20.1	21.1	24.7	28.9	31.1	31.3	28.7	22.2
2000.....	10.2	16.8	18.7	21.9	26.2	28.9	30.8	31.9	29.6	23.1
1999.....	9.9	16.4	18.6	21.4	26.5	29.7	31.6	32.6	30.8	23.2
All races 3/, male										
2005.....	11.5	21.3	31.3	34.5	35.3	33.9	38.1	37.2	35.1	27.6
2004.....	12.2	21.5	31.9	32.3	33.4	35.7	38.7	38.6	34.0	25.5
2003.....	12.8	23.4	32.3	32.4	35.6	38.4	40.0	38.1	35.0	27.5
2002.....	12.9	23.3	31.0	32.0	34.5	37.2	40.0	39.4	36.9	28.4
2001.....	13.2	25.2	30.1	31.5	35.3	39.7	40.8	37.6	35.3	32.0
2000.....	14.4	25.4	28.6	31.7	37.7	39.3	40.7	40.7	34.6	34.3
1999.....	14.5	25.3	28.8	32.5	37.2	41.8	41.5	40.1	38.7	33.1
All races 3/, female										
2005.....	5.7	8.9	10.6	12.2	13.9	17.9	22.2	24.4	23.4	16.0
2004.....	5.4	9.0	11.1	11.8	14.9	18.0	22.1	24.0	22.7	17.1
2003.....	5.9	8.6	10.3	12.2	14.2	19.0	24.1	26.0	24.5	16.8
2002.....	6.5	8.9	10.1	12.1	15.3	19.8	24.1	27.6	25.6	18.4
2001.....	6.4	9.3	10.5	11.4	15.2	19.6	23.5	26.9	24.7	18.1
2000.....	6.1	8.4	9.1	12.7	15.7	19.9	23.1	25.8	26.6	18.5
1999.....	5.4	7.8	8.8	11.1	16.8	19.4	23.9	27.4	26.2	19.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Chronic liver disease and cirrhosis (K70,K73-K74)										
White, both sexes										
2005 4/.....	9.0	15.4	21.0	23.5	24.2	26.5	30.9	30.8	28.8	20.2
2004 4/.....	9.0	15.6	21.6	21.6	24.0	27.1	31.1	31.8	28.1	20.3
2003 4/.....	9.7	16.2	21.0	22.0	24.6	28.9	32.8	32.9	29.8	20.9
2002.....	10.0	16.1	20.2	21.8	24.8	28.9	32.7	34.2	30.9	22.3
2001.....	9.8	17.2	19.6	20.7	25.3	29.7	32.3	32.6	30.0	23.2
2000.....	10.1	16.9	18.1	21.6	26.5	29.4	31.8	33.4	30.5	23.9
1999.....	9.7	15.8	18.3	20.8	26.3	30.2	32.6	33.7	32.0	23.8
White, male										
2005 4/.....	12.1	21.8	31.9	35.0	35.2	35.0	39.8	38.1	35.9	28.5
2004 4/.....	12.6	22.0	32.5	32.0	33.7	36.4	40.3	40.5	35.4	26.7
2003 4/.....	13.4	24.0	32.0	32.2	35.9	39.5	41.8	40.1	36.2	28.4
2002.....	13.5	23.8	30.9	32.1	34.7	38.2	41.6	41.6	38.1	29.6
2001.....	13.2	25.5	29.3	30.6	35.6	40.2	42.1	39.1	37.2	33.5
2000.....	14.4	26.0	27.8	31.5	38.3	40.2	41.7	42.3	35.9	35.5
1999.....	14.2	24.6	28.4	31.5	37.4	42.2	42.5	41.1	40.3	34.2
White, female										
2005 4/.....	5.9	9.1	10.3	12.4	13.9	19.0	23.5	25.5	24.3	16.4
2004 4/.....	5.4	9.2	11.0	11.6	15.0	18.8	23.5	25.5	23.6	17.4
2003 4/.....	5.9	8.4	10.2	12.2	14.2	19.6	25.4	27.8	25.9	17.5
2002.....	6.5	8.5	9.8	11.9	15.6	20.7	25.4	28.9	26.5	19.1
2001.....	6.3	8.9	10.0	11.3	15.8	20.4	24.3	28.0	25.7	18.9
2000.....	5.9	7.8	8.7	12.2	15.6	20.0	23.9	27.2	27.3	19.2
1999.....	5.1	7.2	8.5	10.7	16.2	19.8	24.7	28.4	27.2	19.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Chronic liver disease and cirrhosis (K70,K73-K74)										
Black, both sexes										
2005 4/.....	6.2	14.2	21.6	24.6	25.4	20.0	20.4	19.9	18.0	12.3
2004 4/.....	7.7	14.4	23.0	25.7	24.7	22.6	20.3	13.9	15.1	12.1
2003 4/.....	7.7	15.0	25.6	26.4	25.0	25.4	21.7	14.1	14.7	12.3
2002.....	7.9	17.4	25.0	24.6	25.5	22.9	20.0	20.1	16.0	12.2
2001.....	10.6	19.2	26.8	27.7	23.6	26.9	22.6	21.0	14.6	10.4
2000.....	10.9	19.2	25.9	25.9	25.9	27.5	24.7	18.3	17.5	13.8
1999.....	11.9	22.5	22.9	28.2	30.7	27.7	25.3	23.2	18.8	17.2
Black, male										
2005 4/.....	8.1	20.8	31.7	38.3	40.5	30.9	31.6	32.6	23.2	*
2004 4/.....	10.5	21.3	34.3	40.9	36.4	35.1	32.0	20.0	21.5	*
2003 4/.....	9.7	21.4	41.5	42.2	38.9	36.7	28.9	19.6	22.1	21.9
2002.....	9.8	23.4	38.2	36.9	39.0	34.9	29.0	23.0	21.4	*
2001.....	14.5	28.2	42.2	43.8	38.4	43.2	33.7	26.9	*	*
2000.....	14.6	26.4	40.7	38.5	37.3	38.0	37.4	27.4	19.4	*
1999.....	17.1	34.5	36.6	44.1	42.4	43.4	37.3	31.8	25.3	*
Black, female										
2005 4/.....	4.5	8.5	13.1	13.2	13.6	12.0	12.8	12.3	15.2	10.9
2004 4/.....	5.2	8.5	13.5	13.1	15.6	13.3	12.3	10.2	11.7	11.4
2003 4/.....	5.9	9.3	12.2	13.4	14.1	17.0	16.8	10.9	10.8	8.5
2002.....	6.3	12.2	13.7	14.5	14.9	14.1	13.9	18.4	13.1	10.7
2001.....	7.1	11.3	13.7	14.5	12.0	14.8	15.2	17.5	14.7	9.7
2000.....	7.7	13.0	13.3	15.5	17.0	19.8	16.2	12.8	16.6	11.6
1999.....	7.3	12.2	11.4	15.2	21.5	16.2	17.3	18.0	15.6	15.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Alcoholic liver disease (K70)										
All races 3/, both sexes										
2005.....	5.7	9.3	12.0	12.4	12.0	10.3	9.4	6.4	4.8	1.8
2004.....	5.8	9.1	12.2	11.8	11.5	10.8	9.0	6.5	4.1	2.1
2003.....	5.9	9.4	11.8	11.6	11.6	11.0	8.3	6.6	4.4	1.8
2002.....	5.9	9.4	11.4	11.4	11.8	10.7	9.5	6.7	4.4	2.1
2001.....	6.1	10.2	11.3	11.1	12.0	11.1	9.0	6.9	3.9	2.4
2000.....	6.6	10.0	10.7	11.7	12.9	11.4	9.1	7.2	4.4	2.2
1999.....	6.4	10.0	10.9	11.5	12.6	11.6	9.4	7.3	5.2	2.8
All races 3/, male										
2005.....	7.5	13.2	18.5	19.5	19.4	16.5	15.0	11.2	8.7	4.2
2004.....	8.0	12.9	18.6	18.2	18.3	17.8	15.0	11.5	8.0	4.6
2003.....	8.0	14.2	18.3	18.1	18.6	17.7	14.2	11.3	8.2	3.6
2002.....	8.0	13.8	17.7	17.6	19.2	17.4	15.4	10.9	8.0	4.7
2001.....	8.4	15.3	17.0	18.0	19.0	17.9	15.2	12.1	7.0	5.5
2000.....	9.4	15.3	16.9	18.0	20.3	18.7	15.7	13.8	7.7	5.3
1999.....	9.4	15.5	17.2	18.7	20.0	19.7	16.2	12.1	9.7	6.1
All races 3/, female										
2005.....	3.9	5.4	5.8	5.7	5.3	4.9	4.8	3.0	2.4	0.7
2004.....	3.6	5.5	6.0	5.7	5.4	4.8	4.1	2.9	1.7	1.0
2003.....	3.8	4.8	5.5	5.5	5.3	5.3	3.6	3.3	2.1	1.0
2002.....	3.9	5.1	5.4	5.6	5.1	4.9	4.8	3.7	2.2	1.0
2001.....	3.9	5.3	5.8	4.7	5.6	5.3	4.0	3.3	2.1	1.0
2000.....	3.9	4.8	4.8	5.8	6.2	5.1	3.9	2.6	2.4	1.0
1999.....	3.5	4.6	4.8	4.9	5.9	4.6	4.0	3.9	2.7	1.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Alcoholic liver disease (K70)										
White, both sexes										
2005 4/.....	5.9	9.6	12.5	12.9	12.1	10.7	9.7	6.6	4.9	1.9
2004 4/.....	6.1	9.5	12.6	11.8	11.7	11.1	9.3	6.8	4.1	2.2
2003 4/.....	6.2	9.7	11.8	11.8	11.6	11.2	8.6	7.0	4.5	1.9
2002.....	6.2	9.6	11.5	11.4	12.0	11.0	9.8	6.8	4.5	2.1
2001.....	6.2	10.5	11.0	11.0	12.2	11.2	9.2	7.1	4.0	2.3
2000.....	6.6	10.0	10.6	11.5	13.0	11.6	9.3	7.3	4.3	2.3
1999.....	6.2	9.8	10.8	11.2	12.4	11.5	9.4	7.2	5.3	2.7
White, male										
2005 4/.....	7.8	13.7	19.2	20.3	19.5	17.0	15.4	11.3	8.9	4.4
2004 4/.....	8.5	13.3	19.4	18.1	18.4	18.1	15.3	12.0	8.0	4.8
2003 4/.....	8.4	14.6	18.4	18.3	18.7	17.9	14.6	11.8	8.2	3.7
2002.....	8.3	14.3	17.9	17.9	19.4	17.7	15.8	11.1	8.2	4.5
2001.....	8.5	15.9	16.8	17.6	19.4	17.8	15.4	12.5	7.2	5.4
2000.....	9.4	15.6	16.8	18.0	20.6	19.0	15.9	14.1	7.8	5.4
1999.....	9.2	15.4	17.1	18.2	20.0	19.5	16.2	11.9	9.8	5.8
White, female										
2005 4/.....	3.9	5.6	6.0	5.8	5.2	5.1	5.0	3.1	2.4	0.8
2004 4/.....	3.7	5.7	5.9	5.7	5.4	4.9	4.4	3.0	1.7	1.0
2003 4/.....	3.9	4.9	5.4	5.5	5.1	5.3	3.7	3.5	2.2	1.1
2002.....	4.0	5.0	5.2	5.3	5.1	5.0	4.9	3.7	2.2	1.1
2001.....	3.9	5.2	5.4	4.6	5.7	5.4	4.2	3.3	2.1	1.0
2000.....	3.8	4.4	4.5	5.3	6.0	5.1	4.0	2.6	2.3	1.0
1999.....	3.3	4.3	4.5	4.7	5.5	4.6	4.0	3.9	2.6	1.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Alcoholic liver disease (K70)

Black, both sexes

2005 4/.....	3.9	7.8	10.1	11.0	11.9	9.0	8.4	5.3	*	*
2004 4/.....	4.1	7.5	10.8	12.9	12.0	10.8	6.5	*	*	*
2003 4/.....	4.2	7.6	12.7	12.2	12.9	11.2	6.9	*	*	*
2002.....	4.2	8.2	12.1	12.5	12.0	9.6	7.4	6.2	*	*
2001.....	5.9	9.0	14.7	14.1	11.5	13.0	7.8	5.7	*	*
2000.....	6.9	11.0	13.3	13.9	12.6	11.8	8.4	6.2	*	*
1999.....	7.2	12.8	12.8	14.8	15.6	12.9	10.4	8.0	*	*

Black, male

2005 4/.....	4.6	11.7	15.0	16.8	20.5	15.4	15.2	12.1	*	*
2004 4/.....	5.6	10.7	15.6	20.8	19.4	19.4	13.6	*	*	*
2003 4/.....	5.4	12.0	20.6	19.5	20.8	18.7	12.5	*	*	*
2002.....	5.6	11.5	18.7	17.4	20.4	16.5	13.0	10.6	*	*
2001.....	8.2	12.9	22.3	23.5	18.7	22.9	15.1	*	*	*
2000.....	9.6	15.0	20.6	20.4	18.0	19.3	16.7	11.3	*	*
1999.....	10.1	19.2	20.2	24.9	23.3	24.1	18.7	14.2	*	*

Black, female

2005 4/.....	3.3	4.3	5.9	6.2	5.1	4.2	*	*	*	*
2004 4/.....	2.8	4.6	6.8	6.3	6.1	4.5	*	*	*	*
2003 4/.....	3.0	3.7	6.0	6.1	6.8	5.7	*	*	*	*
2002.....	3.0	5.3	6.6	8.4	5.4	4.4	*	*	*	*
2001.....	3.9	5.5	8.2	6.4	5.8	5.6	*	*	*	*
2000.....	4.5	7.5	7.1	8.5	8.3	6.3	*	*	*	*
1999.....	4.7	7.2	6.6	6.5	9.5	4.6	5.0	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other chronic liver disease and cirrhosis (K73-K74)										
All races 3/, both sexes										
2005.....	2.9	5.7	8.7	10.6	12.0	15.0	20.0	23.3	23.1	17.8
2004.....	2.9	6.0	9.1	9.9	12.2	15.4	20.5	23.6	22.9	17.6
2003.....	3.4	6.5	9.3	10.4	12.7	17.0	22.9	24.5	24.1	18.2
2002.....	3.7	6.6	8.9	10.3	12.6	17.2	21.7	25.8	25.5	19.3
2001.....	3.6	6.9	8.8	10.0	12.8	17.7	22.2	24.4	24.7	19.9
2000.....	3.6	6.8	7.9	10.2	13.3	17.5	21.7	24.7	25.2	20.9
1999.....	3.5	6.4	7.7	9.9	13.9	18.1	22.3	25.4	25.6	20.4
All races 3/, male										
2005.....	4.0	8.1	12.8	14.9	15.9	17.4	23.1	26.0	26.5	23.3
2004.....	4.1	8.6	13.3	14.1	15.2	17.9	23.8	27.1	26.0	21.0
2003.....	4.8	9.2	14.0	14.3	17.0	20.7	25.8	26.8	26.8	23.9
2002.....	4.9	9.5	13.3	14.4	15.3	19.8	24.6	28.6	28.9	23.7
2001.....	4.8	10.0	13.1	13.5	16.3	21.8	25.6	25.5	28.4	26.5
2000.....	5.0	10.2	11.6	13.7	17.4	20.6	25.0	27.0	26.8	29.0
1999.....	5.2	9.8	11.6	13.8	17.3	22.0	25.3	28.0	29.0	27.0
All races 3/, female										
2005.....	1.9	3.5	4.7	6.5	8.5	13.0	17.4	21.4	21.0	15.3
2004.....	1.8	3.5	5.1	6.0	9.5	13.2	18.0	21.1	21.0	16.1
2003.....	2.0	3.8	4.8	6.6	8.9	13.8	20.5	22.8	22.5	15.8
2002.....	2.6	3.8	4.7	6.5	10.2	14.9	19.4	23.9	23.4	17.4
2001.....	2.5	4.0	4.7	6.7	9.6	14.2	19.5	23.6	22.6	17.1
2000.....	2.2	3.6	4.3	6.9	9.5	14.8	19.2	23.2	24.2	17.5
1999.....	1.8	3.2	4.0	6.2	10.8	14.8	19.9	23.5	23.6	17.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other chronic liver disease and cirrhosis (K73-K74)										
White, both sexes										
2005 4/.....	3.1	5.8	8.5	10.6	12.0	15.8	21.2	24.2	23.8	18.3
2004 4/.....	2.9	6.1	9.0	9.8	12.3	16.0	21.7	25.0	24.0	18.1
2003 4/.....	3.5	6.5	9.2	10.2	13.0	17.7	24.2	26.0	25.3	19.0
2002.....	3.9	6.5	8.7	10.4	12.8	18.0	22.8	27.4	26.4	20.2
2001.....	3.6	6.6	8.5	9.7	13.1	18.5	23.1	25.5	26.0	20.9
2000.....	3.6	6.9	7.5	10.1	13.5	17.8	22.5	26.1	26.2	21.6
1999.....	3.4	6.0	7.6	9.6	13.9	18.7	23.2	26.5	26.8	21.1
White, male										
2005 4/.....	4.2	8.1	12.7	14.7	15.7	18.0	24.5	26.7	26.9	24.2
2004 4/.....	4.1	8.6	13.1	13.9	15.3	18.3	24.9	28.5	27.3	22.0
2003 4/.....	5.0	9.4	13.6	13.8	17.2	21.5	27.2	28.2	27.9	24.7
2002.....	5.2	9.5	13.0	14.3	15.3	20.5	25.8	30.5	30.0	25.1
2001.....	4.7	9.6	12.5	13.0	16.2	22.4	26.7	26.6	30.0	28.1
2000.....	5.0	10.3	11.0	13.5	17.6	21.2	25.8	28.2	28.1	30.1
1999.....	5.1	9.2	11.3	13.4	17.3	22.8	26.3	29.1	30.5	28.4
White, female										
2005 4/.....	2.0	3.5	4.4	6.6	8.7	13.9	18.5	22.3	21.9	15.6
2004 4/.....	1.7	3.5	5.0	5.9	9.6	13.9	19.1	22.4	21.9	16.4
2003 4/.....	1.9	3.6	4.8	6.7	9.2	14.3	21.7	24.3	23.7	16.5
2002.....	2.5	3.5	4.6	6.6	10.5	15.7	20.4	25.1	24.3	18.1
2001.....	2.4	3.8	4.6	6.6	10.2	15.1	20.2	24.7	23.5	17.9
2000.....	2.1	3.4	4.2	6.9	9.6	14.8	19.9	24.6	25.1	18.2
1999.....	1.8	2.9	3.9	6.0	10.7	15.1	20.7	24.6	24.6	18.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other chronic liver disease and cirrhosis (K73-K74)										
Black, both sexes										
2005 4/.....	2.3	6.5	11.5	13.6	13.5	11.0	12.1	14.5	14.6	10.7
2004 4/.....	3.6	7.0	12.2	12.8	12.8	11.8	13.7	11.2	11.4	11.3
2003 4/.....	3.6	7.4	13.0	14.2	12.1	14.2	14.8	11.1	10.9	10.8
2002.....	3.7	9.2	12.8	12.1	13.5	13.4	12.6	13.9	12.0	9.8
2001.....	4.7	10.2	12.1	13.6	12.2	13.9	14.8	15.3	11.6	7.7
2000.....	4.1	8.2	12.6	12.0	13.3	15.7	16.3	12.1	13.3	12.2
1999.....	4.7	9.8	10.1	13.4	15.1	14.8	14.9	15.2	13.3	12.8
Black, male										
2005 4/.....	3.5	9.1	16.7	21.6	20.0	15.4	16.4	20.6	19.0	*
2004 4/.....	4.9	10.6	18.7	20.1	17.0	15.7	18.5	15.9	14.3	*
2003 4/.....	4.3	9.5	20.9	22.7	18.1	18.0	16.4	14.6	*	*
2002.....	4.2	11.9	19.5	19.5	18.6	18.4	16.0	12.4	*	*
2001.....	6.4	15.3	19.9	20.3	19.7	20.3	18.5	18.1	*	*
2000.....	5.0	11.4	20.1	18.1	19.3	18.7	20.7	16.1	*	*
1999.....	7.0	15.3	16.4	19.2	19.1	19.2	18.7	17.6	*	*
Black, female										
2005 4/.....	*	4.2	7.1	7.1	8.5	7.8	9.1	10.9	12.2	10.1
2004 4/.....	2.4	3.8	6.7	6.8	9.5	8.8	10.5	8.4	9.8	11.0
2003 4/.....	2.9	5.6	6.2	7.3	7.4	11.3	13.7	9.0	10.1	8.1
2002.....	3.3	6.9	7.1	6.1	9.4	9.6	10.3	14.8	10.7	9.9
2001.....	3.2	5.8	5.5	8.1	6.2	9.2	12.3	13.6	13.1	*
2000.....	3.2	5.5	6.1	7.0	8.7	13.4	13.3	9.7	14.0	11.2
1999.....	2.6	5.0	4.8	8.7	12.0	11.5	12.4	13.7	12.1	13.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Cholelithiasis and other disorders of gallbladder (K80-K82)

All races 3/, both sexes

2005.....	0.1	0.2	0.4	0.5	0.9	1.7	3.3	5.8	11.1	22.9
2004.....	0.2	0.3	0.4	0.6	1.2	1.8	3.2	5.8	10.4	23.8
2003.....	0.1	0.2	0.4	0.5	0.9	2.0	3.4	5.9	11.3	21.8
2002.....	0.2	0.2	0.4	0.6	1.1	2.0	3.6	5.7	10.8	22.9
2001.....	0.1	0.3	0.4	0.7	1.3	2.0	3.8	5.8	10.0	23.7
2000.....	0.1	0.2	0.4	0.6	1.0	1.9	3.0	5.7	10.3	24.8
1999.....	0.2	0.3	0.4	0.7	1.0	1.8	3.4	5.7	10.9	23.9

All races 3/, male

2005.....	*	*	0.5	0.5	1.2	2.3	4.3	7.1	12.7	24.3
2004.....	*	0.3	0.4	0.7	1.4	2.2	3.8	7.0	12.5	25.5
2003.....	*	0.2	0.5	0.6	1.1	2.2	4.3	7.5	13.9	22.7
2002.....	0.2	0.3	0.5	0.8	1.1	2.3	4.0	7.5	13.5	24.1
2001.....	*	0.4	0.4	0.9	1.5	2.2	4.4	7.6	12.1	26.9
2000.....	*	0.2	0.3	0.7	1.2	2.3	3.6	7.4	12.9	27.6
1999.....	*	0.3	0.4	0.9	1.3	2.0	4.2	6.9	13.3	27.7

All races 3/, female

2005.....	*	0.2	0.3	0.6	0.7	1.2	2.4	4.8	10.1	22.3
2004.....	*	0.3	0.4	0.5	1.0	1.5	2.8	5.0	9.0	23.0
2003.....	*	0.2	0.4	0.4	0.7	1.8	2.8	4.8	9.8	21.4
2002.....	*	*	0.3	0.4	1.0	1.8	3.2	4.4	9.1	22.4
2001.....	*	0.3	0.3	0.5	1.1	1.9	3.3	4.6	8.7	22.4
2000.....	*	0.2	0.4	0.5	0.8	1.6	2.5	4.5	8.7	23.7
1999.....	*	0.3	0.5	0.6	0.7	1.6	2.8	4.8	9.5	22.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Cholelithiasis and other disorders of gallbladder (K80-K82)

White, both sexes

2005 4/.....	*	0.1	0.3	0.5	0.9	1.5	3.1	5.7	11.1	23.1
2004 4/.....	0.2	0.2	0.3	0.5	1.1	1.8	3.0	5.7	10.3	24.2
2003 4/.....	*	0.2	0.3	0.4	0.8	1.8	3.6	5.9	11.3	21.9
2002.....	0.2	0.2	0.3	0.6	1.0	2.0	3.5	5.5	10.6	23.5
2001.....	*	0.3	0.3	0.6	1.3	1.8	3.7	5.7	9.8	23.9
2000.....	0.1	0.2	0.3	0.6	0.9	1.8	2.7	5.6	10.1	25.6
1999.....	0.1	0.3	0.4	0.6	0.9	1.6	3.4	5.7	11.0	24.4

White, male

2005 4/.....	*	*	0.4	0.5	1.1	2.1	4.1	7.1	13.1	24.8
2004 4/.....	*	0.2	0.3	0.6	1.4	2.2	3.7	6.9	12.7	26.7
2003 4/.....	*	*	0.4	0.5	0.9	2.1	4.4	7.5	13.9	23.0
2002.....	*	0.3	0.5	0.8	1.1	2.3	4.0	7.4	13.3	24.7
2001.....	*	0.3	0.4	0.8	1.4	1.9	4.4	7.6	12.0	26.7
2000.....	*	*	0.3	0.7	1.1	2.3	3.4	7.4	13.2	28.9
1999.....	*	0.3	0.3	0.7	1.2	1.9	4.2	7.0	13.8	28.1

White, female

2005 4/.....	*	*	0.2	0.5	0.7	0.9	2.3	4.7	9.8	22.4
2004 4/.....	*	0.2	0.3	0.5	0.8	1.4	2.4	4.8	8.8	23.1
2003 4/.....	*	*	0.3	0.3	0.7	1.6	2.8	4.7	9.8	21.4
2002.....	*	*	0.3	0.4	0.9	1.7	3.1	4.1	9.0	23.0
2001.....	*	0.3	*	0.4	1.1	1.8	3.1	4.4	8.4	22.8
2000.....	*	*	0.3	0.4	0.7	1.4	2.2	4.4	8.3	24.3
1999.....	*	0.3	0.4	0.6	0.6	1.4	2.7	4.8	9.3	22.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Cholelithiasis and other disorders of gallbladder (K80-K82)

Black, both sexes

2005 4/.....	*	*	*	*	*	3.6	4.6	7.0	12.4	20.6
2004 4/.....	*	*	1.0	1.3	2.0	2.3	5.1	7.1	12.1	19.5
2003 4/.....	*	*	1.0	*	*	3.0	2.8	6.0	10.9	20.7
2002.....	*	*	*	*	*	2.3	4.6	7.7	12.8	14.8
2001.....	*	*	*	1.4	2.0	3.5	6.2	6.1	11.6	19.7
2000.....	*	*	*	*	*	3.1	4.9	6.6	11.0	17.8
1999.....	*	*	*	*	2.0	3.2	4.1	4.6	11.3	15.3

Black, male

2005 4/.....	*	*	*	*	*	*	6.3	9.8	*	19.5
2004 4/.....	*	*	*	*	*	*	*	*	*	*
2003 4/.....	*	*	*	*	*	*	*	*	*	20.9
2002.....	*	*	*	*	*	*	*	*	16.0	*
2001.....	*	*	*	*	*	*	*	*	*	27.2
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*

Black, female

2005 4/.....	*	*	*	*	*	*	*	5.3	13.4	21.0
2004 4/.....	*	*	*	*	*	*	5.5	6.7	12.9	21.7
2003 4/.....	*	*	*	*	*	*	*	*	9.3	20.7
2002.....	*	*	*	*	*	*	5.4	7.7	11.1	16.0
2001.....	*	*	*	*	*	*	6.3	6.9	13.1	16.8
2000.....	*	*	*	*	*	*	4.9	6.3	11.5	19.3
1999.....	*	*	*	*	*	*	*	*	12.1	15.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)										
All races 3/, both sexes										
2005.....	2.1	3.8	5.8	10.6	17.7	29.2	51.3	85.6	142.8	288.3
2004.....	2.2	3.7	6.4	10.6	17.6	29.6	49.0	85.4	139.0	286.6
2003.....	2.3	3.8	6.1	10.4	17.9	30.7	50.6	84.5	143.9	293.1
2002.....	2.1	3.7	5.9	10.0	16.8	29.4	50.0	85.4	142.2	288.6
2001.....	2.1	3.6	5.8	9.9	16.9	31.7	49.3	82.2	135.8	287.7
2000.....	2.1	3.4	5.5	9.4	16.9	29.5	47.2	79.5	132.8	277.8
1999.....	2.0	3.1	5.2	9.0	15.8	28.2	46.7	77.7	127.4	268.9
All races 3/, male										
2005.....	2.5	4.8	7.2	12.4	21.1	31.4	59.1	102.7	183.8	376.5
2004.....	2.6	4.4	7.7	11.8	19.6	33.3	56.2	102.2	181.2	372.2
2003.....	2.7	4.5	7.6	12.2	19.6	34.6	58.1	102.9	187.9	387.6
2002.....	2.3	4.3	6.9	11.0	18.6	33.6	59.8	105.7	182.7	385.5
2001.....	2.3	4.4	6.7	10.9	18.2	36.5	57.0	100.5	176.8	393.7
2000.....	2.5	4.4	6.4	10.7	18.6	32.8	54.3	99.8	173.0	380.7
1999.....	2.3	3.8	5.8	9.9	17.8	31.0	54.9	99.2	168.9	371.1
All races 3/, female										
2005.....	1.7	2.9	4.5	8.8	14.7	27.3	44.9	73.2	117.2	247.8
2004.....	1.8	3.1	5.2	9.4	15.9	26.4	43.2	73.3	113.0	248.1
2003.....	1.9	3.2	4.7	8.6	16.4	27.4	44.7	71.4	117.1	251.4
2002.....	1.9	3.1	4.9	9.2	15.2	25.8	42.1	71.1	117.6	246.6
2001.....	1.9	2.9	4.9	8.9	15.7	27.6	43.2	69.4	111.2	243.3
2000.....	1.6	2.5	4.6	8.2	15.5	26.6	41.6	65.3	109.0	235.9
1999.....	1.7	2.4	4.5	8.1	14.0	25.8	40.3	62.8	103.1	227.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)										
White, both sexes										
2005 4/.....	1.4	2.7	4.3	7.7	13.8	23.9	44.3	78.7	135.3	280.3
2004 4/.....	1.6	2.5	4.5	7.8	13.6	24.3	42.9	78.2	132.3	278.2
2003 4/.....	1.5	2.6	4.3	7.8	14.1	25.2	43.5	77.6	136.0	284.8
2002.....	1.4	2.6	3.9	7.4	13.1	24.4	43.2	78.1	135.5	278.8
2001.....	1.4	2.4	4.1	7.3	12.8	25.8	43.1	75.1	127.9	278.2
2000.....	1.5	2.2	3.8	6.8	13.2	23.8	40.6	73.0	125.3	268.7
1999.....	1.3	2.0	3.5	6.6	12.2	23.0	40.4	70.9	118.5	255.7
White, male										
2005 4/.....	1.7	3.4	5.5	8.9	16.8	27.1	52.7	95.9	178.1	373.7
2004 4/.....	2.0	2.9	5.3	9.0	15.2	27.6	50.4	95.6	176.4	366.7
2003 4/.....	1.9	3.1	5.1	9.2	15.5	29.1	50.5	96.9	179.7	383.6
2002.....	1.6	3.0	4.6	8.0	14.3	28.4	53.0	99.0	177.9	377.4
2001.....	1.6	2.8	4.9	8.2	14.4	30.5	51.3	94.6	169.0	388.9
2000.....	1.8	2.9	4.4	7.9	14.9	27.8	48.7	93.4	166.7	376.2
1999.....	1.5	2.4	4.0	7.8	14.0	26.0	48.9	91.6	159.6	360.8
White, female										
2005 4/.....	1.1	2.0	3.2	6.5	11.0	21.0	37.4	66.0	108.5	237.3
2004 4/.....	1.1	2.1	3.7	6.5	12.0	21.5	36.7	65.4	104.9	238.3
2003 4/.....	1.1	2.1	3.4	6.5	12.8	21.8	37.7	63.6	109.1	241.0
2002.....	1.3	2.3	3.3	6.7	12.0	20.8	35.1	63.2	109.4	235.9
2001.....	1.2	1.9	3.3	6.3	11.4	21.7	36.4	61.3	103.1	231.9
2000.....	1.1	1.7	3.2	5.8	11.6	20.3	34.1	58.6	100.7	225.0
1999.....	1.1	1.6	3.1	5.5	10.6	20.4	33.6	56.4	94.3	213.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)										
Black, both sexes										
2005 4/.....	7.3	12.1	18.2	35.8	55.7	79.4	124.8	175.9	254.9	424.4
2004 4/.....	6.9	12.6	21.5	35.6	56.7	79.2	115.9	175.4	242.9	430.9
2003 4/.....	7.9	12.6	21.0	34.3	54.2	83.8	127.8	171.0	260.7	440.5
2002.....	6.7	11.7	21.6	33.7	50.4	79.2	126.6	178.0	241.5	441.6
2001.....	6.7	12.5	19.8	32.5	53.1	88.6	119.8	172.3	248.6	429.6
2000.....	6.3	12.7	19.7	32.3	50.7	83.5	119.7	165.4	242.3	406.4
1999.....	7.0	11.8	19.6	29.1	47.1	77.8	117.0	163.7	246.2	433.3
Black, male										
2005 4/.....	8.5	15.9	22.6	45.8	68.3	79.4	133.5	208.8	303.4	482.4
2004 4/.....	7.1	15.4	27.8	39.7	64.1	92.1	129.5	199.3	285.5	512.6
2003 4/.....	8.8	14.8	29.3	43.3	62.7	94.1	155.7	191.7	333.7	518.2
2002.....	7.6	14.7	26.8	40.0	61.2	89.3	149.6	207.6	281.2	561.8
2001.....	7.0	16.5	22.8	37.7	55.5	101.1	130.9	189.1	309.8	511.7
2000.....	7.9	17.4	24.1	37.4	54.2	86.9	127.7	198.9	290.0	490.1
1999.....	8.2	15.5	23.4	29.2	54.5	81.8	130.6	208.0	327.0	526.1
Black, female										
2005 4/.....	6.3	8.8	14.5	27.5	45.9	79.5	118.8	156.3	229.2	401.0
2004 4/.....	6.6	10.1	16.2	32.3	51.0	69.6	106.8	161.2	220.4	398.5
2003 4/.....	7.2	10.7	13.9	26.9	47.5	76.2	109.0	158.7	222.4	410.4
2002.....	5.9	9.1	17.2	28.5	42.0	71.8	111.1	160.3	220.8	395.5
2001.....	6.4	9.1	17.1	28.2	51.2	79.4	112.3	162.3	217.5	399.1
2000.....	4.8	8.6	16.1	28.2	47.9	81.0	114.3	145.3	218.2	375.2
1999.....	5.9	8.6	16.4	29.1	41.3	74.8	108.0	137.0	206.1	398.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01,N04)

All races 3/, both sexes

2005.....	*	*	*	*	*	*	*	0.3	*	0.8
2004.....	*	*	*	*	*	*	*	0.3	0.4	0.8
2003.....	*	*	*	*	*	*	*	*	0.6	0.6
2002.....	*	*	*	*	*	*	*	*	0.6	0.9
2001.....	*	*	*	*	*	*	*	0.4	0.4	0.9
2000.....	*	*	*	*	*	*	*	*	0.7	1.2
1999.....	*	*	*	*	*	*	0.3	0.5	0.5	0.8

All races 3/, male

2005.....	*	*	*	*	*	*	*	*	*	*
2004.....	*	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	1.8
1999.....	*	*	*	*	*	*	*	*	*	*

All races 3/, female

2005.....	*	*	*	*	*	*	*	*	*	0.7
2004.....	*	*	*	*	*	*	*	*	*	0.8
2003.....	*	*	*	*	*	*	*	*	*	0.6
2002.....	*	*	*	*	*	*	*	*	0.6	0.9
2001.....	*	*	*	*	*	*	*	*	*	0.7
2000.....	*	*	*	*	*	*	*	*	0.7	0.9
1999.....	*	*	*	*	*	*	*	0.5	*	0.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02-N03,N05-N07,N26)

All races 3/, both sexes

2005.....	*	*	0.2	0.3	0.4	0.5	0.9	1.6	2.4	5.4
2004.....	*	*	0.1	0.2	0.3	0.4	0.6	1.0	1.6	2.8
2003.....	*	*	0.2	0.1	0.2	0.4	0.7	1.0	1.6	2.8
2002.....	*	0.1	0.1	0.2	*	0.3	0.6	1.2	1.7	2.8
2001.....	*	*	0.1	*	0.2	0.5	0.7	0.9	1.5	2.8
2000.....	0.1	*	0.1	0.2	0.3	0.5	0.7	1.1	1.9	2.8
1999.....	*	*	0.2	0.2	0.3	0.4	0.7	1.4	2.0	3.2

All races 3/, male

2005.....	*	*	0.2	0.3	0.4	0.6	1.1	1.9	3.2	7.5
2004.....	*	*	*	0.2	0.3	0.4	0.7	1.3	2.0	4.0
2003.....	*	*	0.2	*	*	0.5	0.6	1.3	2.0	4.1
2002.....	*	*	*	0.3	*	*	0.9	1.6	2.1	3.8
2001.....	*	*	*	*	*	0.5	1.0	1.3	2.3	3.7
2000.....	*	*	*	*	*	0.5	0.8	1.5	2.1	4.5
1999.....	*	*	*	*	*	0.5	0.6	1.9	2.7	5.0

All races 3/, female

2005.....	*	*	*	*	0.3	0.5	0.8	1.4	1.9	4.4
2004.....	*	*	*	*	*	*	0.5	0.8	1.4	2.2
2003.....	*	*	*	*	*	*	0.8	0.9	1.4	2.2
2002.....	*	*	*	*	*	*	0.4	0.8	1.5	2.3
2001.....	*	*	*	*	*	0.5	0.5	0.7	1.0	2.4
2000.....	*	*	*	*	*	*	0.6	0.8	1.8	2.1
1999.....	*	*	*	*	*	0.4	0.7	1.1	1.5	2.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02-N03,N05-N07,N26)

White, both sexes

2005 4/.....	*	*	*	0.2	0.3	0.5	0.8	1.5	2.3	5.2
2004 4/.....	*	*	0.1	0.2	0.2	0.4	0.6	1.0	1.6	2.7
2003 4/.....	*	*	*	0.2	0.2	0.4	0.6	0.9	1.5	2.6
2002.....	*	*	*	0.2	*	0.3	0.7	1.0	1.7	2.6
2001.....	*	*	*	*	*	0.4	0.7	0.9	1.5	2.9
2000.....	*	*	*	0.2	*	0.4	0.7	1.0	1.9	2.4
1999.....	*	*	*	*	0.3	0.3	0.6	1.4	1.8	2.9

White, male

2005 4/.....	*	*	*	*	*	0.5	1.0	1.7	3.1	7.3
2004 4/.....	*	*	*	*	*	*	0.6	1.2	2.0	3.7
2003 4/.....	*	*	*	*	*	0.5	*	1.1	1.9	3.8
2002.....	*	*	*	*	*	*	0.9	1.5	2.1	3.8
2001.....	*	*	*	*	*	*	1.1	1.2	2.3	4.0
2000.....	*	*	*	*	*	*	0.8	1.4	2.2	4.0
1999.....	*	*	*	*	*	*	0.6	1.9	2.6	4.6

White, female

2005 4/.....	*	*	*	*	*	*	0.6	1.3	1.8	4.2
2004 4/.....	*	*	*	*	*	*	0.5	0.8	1.3	2.3
2003 4/.....	*	*	*	*	*	*	0.6	0.8	1.3	2.1
2002.....	*	*	*	*	*	*	0.5	0.7	1.4	2.0
2001.....	*	*	*	*	*	*	0.5	0.7	0.9	2.4
2000.....	*	*	*	*	*	*	0.6	0.7	1.7	1.8
1999.....	*	*	*	*	*	*	0.7	1.0	1.4	2.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Renal failure (N17-N19)										
All races 3/, both sexes										
2005.....	2.0	3.8	5.6	10.3	17.3	28.6	50.2	83.7	140.0	282.1
2004.....	2.1	3.6	6.2	10.3	17.3	29.1	48.2	84.0	136.8	282.9
2003.....	2.2	3.7	5.9	10.2	17.6	30.2	49.7	83.2	141.7	289.7
2002.....	2.0	3.6	5.7	9.8	16.5	29.0	49.1	83.9	139.8	284.8
2001.....	2.0	3.5	5.7	9.7	16.6	31.1	48.4	80.9	133.8	284.0
2000.....	2.0	3.3	5.3	9.2	16.6	28.9	46.2	78.1	130.0	273.7
1999.....	1.9	3.0	4.9	8.7	15.3	27.7	45.7	75.7	124.8	264.8
All races 3/, male										
2005.....	2.3	4.7	6.9	12.0	20.5	30.7	57.8	100.4	180.3	368.0
2004.....	2.5	4.3	7.4	11.6	19.1	32.8	55.3	100.5	178.4	367.4
2003.....	2.6	4.3	7.4	12.0	19.3	33.9	57.2	101.3	185.2	383.1
2002.....	2.2	4.2	6.7	10.6	18.4	33.1	58.8	103.9	180.0	380.7
2001.....	2.2	4.2	6.5	10.7	17.9	35.9	55.8	98.9	173.9	388.7
2000.....	2.4	4.2	6.1	10.4	18.1	32.1	53.3	97.9	170.1	374.2
1999.....	2.2	3.6	5.6	9.7	17.3	30.5	53.9	96.7	165.1	364.6
All races 3/, female										
2005.....	1.7	2.8	4.4	8.6	14.3	26.7	44.0	71.6	115.0	242.6
2004.....	1.8	3.0	5.0	9.2	15.6	25.8	42.5	72.1	111.2	244.9
2003.....	1.8	3.2	4.6	8.5	16.0	27.0	43.6	70.2	115.0	248.4
2002.....	1.8	2.9	4.8	9.0	14.9	25.4	41.4	69.9	115.3	243.2
2001.....	1.8	2.8	4.8	8.8	15.5	27.0	42.5	68.3	109.8	240.1
2000.....	1.6	2.4	4.5	8.0	15.2	26.1	40.6	64.4	106.4	232.8
1999.....	1.6	2.3	4.3	7.9	13.5	25.3	39.3	61.1	101.2	224.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Renal failure (N17-N19)										
White, both sexes										
2005 4/.....	1.3	2.6	4.2	7.4	13.3	23.3	43.3	76.8	132.7	274.3
2004 4/.....	1.5	2.5	4.3	7.6	13.3	23.8	42.1	76.8	130.1	274.7
2003 4/.....	1.4	2.5	4.1	7.6	13.8	24.7	42.7	76.4	133.9	281.5
2002.....	1.4	2.5	3.8	7.1	12.9	23.9	42.3	76.8	133.2	275.3
2001.....	1.4	2.3	4.0	7.1	12.6	25.2	42.1	73.8	126.0	274.6
2000.....	1.4	2.1	3.7	6.6	12.8	23.3	39.6	71.7	122.6	265.0
1999.....	1.2	1.8	3.4	6.4	11.9	22.5	39.4	69.0	116.2	251.9
White, male										
2005 4/.....	1.6	3.4	5.3	8.6	16.3	26.5	51.4	93.7	174.6	365.4
2004 4/.....	1.9	2.9	5.1	8.8	14.9	27.1	49.5	94.0	173.7	362.3
2003 4/.....	1.8	3.0	4.9	9.0	15.3	28.4	49.7	95.6	177.3	379.3
2002.....	1.5	2.9	4.5	7.7	14.0	27.9	52.0	97.3	175.3	372.7
2001.....	1.6	2.7	4.7	8.0	14.1	30.0	50.1	93.0	166.1	383.9
2000.....	1.7	2.7	4.2	7.6	14.5	27.2	47.7	91.6	163.8	370.4
1999.....	1.4	2.2	3.8	7.6	13.6	25.5	48.0	89.1	156.2	354.9
White, female										
2005 4/.....	1.0	1.9	3.2	6.3	10.5	20.6	36.7	64.3	106.4	232.4
2004 4/.....	1.1	2.1	3.5	6.4	11.8	21.0	36.0	64.2	103.1	235.2
2003 4/.....	1.0	2.1	3.3	6.4	12.4	21.4	36.9	62.4	107.1	238.2
2002.....	1.2	2.1	3.2	6.6	11.8	20.3	34.4	62.1	107.4	232.9
2001.....	1.1	1.9	3.3	6.2	11.2	21.1	35.6	60.2	101.8	228.8
2000.....	1.1	1.6	3.1	5.6	11.3	19.9	33.2	57.8	98.1	222.1
1999.....	1.1	1.5	2.9	5.3	10.3	19.9	32.5	54.8	92.5	210.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Renal failure (N17-N19)										
Black, both sexes										
2005 4/.....	7.0	11.9	17.6	34.9	54.9	78.4	122.1	172.7	250.0	413.7
2004 4/.....	6.6	12.2	21.3	35.2	56.1	78.0	114.8	173.3	239.6	426.4
2003 4/.....	7.8	12.4	20.5	34.2	53.6	82.9	126.1	168.8	257.4	436.4
2002.....	6.6	11.2	21.1	33.3	50.1	78.8	125.7	175.4	238.1	434.2
2001.....	6.5	12.3	19.3	32.2	52.5	87.1	119.0	170.6	245.9	425.6
2000.....	6.1	12.3	18.9	32.0	50.1	82.1	118.6	163.1	238.9	400.4
1999.....	6.7	11.7	18.9	28.5	46.1	76.5	115.7	161.0	241.6	425.3
Black, male										
2005 4/.....	8.0	15.6	21.9	44.7	67.1	78.7	131.2	203.4	297.1	469.4
2004 4/.....	6.8	15.0	27.6	39.1	63.0	91.1	127.9	196.5	282.7	506.7
2003 4/.....	8.6	14.4	28.7	43.3	61.9	92.6	153.7	188.1	328.5	513.0
2002.....	7.4	14.4	26.1	39.4	61.2	88.8	148.2	203.5	278.2	555.3
2001.....	6.8	16.1	22.1	37.2	54.5	99.1	129.9	187.7	307.4	508.3
2000.....	7.7	17.1	22.7	37.0	53.2	84.8	127.0	196.6	287.5	480.9
1999.....	8.0	15.3	22.7	28.7	53.5	80.5	128.9	204.7	320.9	513.2
Black, female										
2005 4/.....	6.1	8.6	13.9	26.8	45.3	78.3	115.8	154.4	225.2	391.2
2004 4/.....	6.3	9.8	16.0	31.9	50.7	68.4	105.9	159.6	217.0	394.6
2003 4/.....	7.0	10.7	13.7	26.7	47.1	75.7	107.4	157.4	220.1	406.7
2002.....	5.9	8.5	16.8	28.2	41.5	71.4	110.7	158.6	217.2	387.7
2001.....	6.4	9.0	16.9	28.1	50.9	78.3	111.7	160.3	214.6	394.9
2000.....	4.7	8.2	15.7	27.8	47.6	80.1	113.0	143.0	214.4	370.4
1999.....	5.6	8.5	15.8	28.4	40.3	73.5	106.9	134.7	202.2	392.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Infections of kidney (N10-N12,N13.6,N15.1)

All races 3/, both sexes

2005.....	*	0.1	0.1	0.1	0.2	0.4	0.7	1.3	2.5	5.2
2004.....	0.1	0.1	0.1	0.2	0.2	0.4	0.7	1.3	2.6	5.2
2003.....	0.1	0.1	0.2	0.2	0.2	0.6	0.6	1.3	2.8	5.9
2002.....	*	*	0.1	0.2	0.3	0.5	0.7	1.6	2.3	6.3
2001.....	*	0.1	0.1	0.2	0.2	0.4	0.6	1.7	2.4	6.0
2000.....	*	0.1	0.1	0.2	0.3	0.4	0.7	1.5	2.7	7.5
1999.....	*	0.1	0.2	0.2	0.3	0.5	0.9	1.5	2.7	7.5

All races 3/, male

2005.....	*	*	*	*	*	*	0.7	0.9	2.5	3.6
2004.....	*	*	*	0.3	*	*	*	1.0	2.2	4.5
2003.....	*	*	*	*	*	*	*	0.8	2.4	5.4
2002.....	*	*	*	*	*	*	0.7	1.4	1.4	5.3
2001.....	*	*	*	*	*	*	0.5	1.2	2.1	4.3
2000.....	*	*	*	*	*	*	*	1.1	2.2	5.6
1999.....	*	*	*	*	*	*	0.6	1.2	2.6	6.8

All races 3/, female

2005.....	*	*	*	*	*	0.5	0.8	1.6	2.6	5.9
2004.....	*	*	*	*	*	0.5	0.9	1.5	2.8	5.5
2003.....	*	*	*	0.3	*	0.8	0.7	1.6	3.1	6.1
2002.....	*	*	*	*	0.4	0.6	0.7	1.7	2.8	6.7
2001.....	*	*	0.2	*	*	*	0.6	2.1	2.6	6.7
2000.....	*	*	*	0.3	0.4	0.5	1.1	1.8	3.0	8.3
1999.....	*	*	0.3	*	*	0.6	1.1	1.7	2.8	7.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Infections of kidney (N10-N12,N13.6,N15.1)

White, both sexes

2005 4/.....	*	0.1	*	*	0.2	0.4	0.7	1.2	2.6	5.2
2004 4/.....	0.1	*	*	0.2	0.2	0.4	0.7	1.2	2.5	5.2
2003 4/.....	0.1	0.1	0.2	0.2	0.2	0.6	0.6	1.3	2.7	6.0
2002.....	*	*	*	0.2	0.2	0.5	0.8	1.6	2.3	6.4
2001.....	*	*	*	*	*	0.3	0.6	1.7	2.2	6.1
2000.....	*	*	*	0.2	0.3	0.4	0.7	1.4	2.5	7.6
1999.....	*	*	0.2	0.2	0.2	0.5	0.9	1.5	2.8	7.7

White, male

2005 4/.....	*	*	*	*	*	*	0.6	0.9	2.6	3.5
2004 4/.....	*	*	*	*	*	*	*	0.9	2.4	4.7
2003 4/.....	*	*	*	*	*	*	*	0.8	2.2	5.8
2002.....	*	*	*	*	*	*	0.8	1.4	1.4	5.4
2001.....	*	*	*	*	*	*	*	1.1	2.0	4.3
2000.....	*	*	*	*	*	*	*	1.1	2.1	5.9
1999.....	*	*	*	*	*	*	*	1.3	2.6	7.2

White, female

2005 4/.....	*	*	*	*	*	0.5	0.7	1.5	2.7	6.0
2004 4/.....	*	*	*	*	*	0.5	1.0	1.4	2.7	5.5
2003 4/.....	*	*	*	*	*	0.8	0.7	1.6	3.1	6.0
2002.....	*	*	*	*	*	0.6	0.8	1.7	2.9	6.8
2001.....	*	*	*	*	*	*	0.7	2.2	2.4	6.9
2000.....	*	*	*	*	*	0.5	1.0	1.7	2.8	8.3
1999.....	*	*	*	*	*	0.6	1.1	1.7	2.9	7.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)										
All races 3/, both sexes										
2005.....	1.2	1.5	1.8	2.3	2.4	1.7	2.0	2.4	4.0	6.4
2004.....	1.2	1.5	1.8	2.2	2.2	2.3	2.3	3.1	4.3	6.1
2003.....	1.4	1.5	1.8	2.4	2.4	2.3	2.2	3.0	4.4	6.8
2002.....	1.3	1.6	1.9	2.3	2.4	2.3	2.5	3.1	4.8	6.6
2001.....	1.4	1.5	1.8	2.1	2.5	2.2	2.7	3.3	4.3	7.2
2000.....	1.3	1.4	1.8	2.2	2.5	2.6	2.4	3.3	5.0	7.5
1999.....	1.2	1.7	1.8	2.3	2.4	2.6	2.5	3.4	5.0	7.8
All races 3/, male										
2005.....	1.3	1.5	1.9	2.5	2.6	1.6	2.0	2.8	4.9	6.9
2004.....	1.3	1.7	2.0	2.3	2.2	2.3	2.7	3.7	3.8	6.8
2003.....	1.5	1.6	1.8	2.5	2.6	2.4	2.4	2.8	4.5	6.7
2002.....	1.4	1.9	2.0	2.3	2.7	2.4	2.7	3.4	4.8	7.0
2001.....	1.6	1.5	1.9	2.1	2.5	2.5	3.1	3.6	4.8	7.7
2000.....	1.5	1.6	2.2	2.0	2.6	2.7	2.6	3.3	5.3	10.6
1999.....	1.3	1.9	2.0	2.5	2.5	2.4	2.6	3.4	5.7	8.5
All races 3/, female										
2005.....	1.2	1.4	1.8	2.1	2.3	1.7	2.0	2.2	3.5	6.2
2004.....	1.2	1.3	1.7	2.0	2.1	2.3	2.0	2.7	4.6	5.8
2003.....	1.3	1.3	1.8	2.3	2.3	2.2	2.1	3.2	4.4	6.8
2002.....	1.2	1.4	1.8	2.3	2.2	2.3	2.3	2.9	4.9	6.5
2001.....	1.3	1.4	1.8	2.1	2.6	2.0	2.4	3.1	4.0	7.0
2000.....	1.1	1.3	1.5	2.4	2.5	2.5	2.3	3.3	4.8	6.3
1999.....	1.2	1.6	1.7	2.2	2.3	2.8	2.3	3.4	4.5	7.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)

White, both sexes

2005 4/.....	1.3	1.5	2.0	2.4	2.6	1.8	2.2	2.5	4.3	6.5
2004 4/.....	1.3	1.6	1.9	2.3	2.3	2.4	2.5	3.1	4.4	6.3
2003 4/.....	1.5	1.5	1.9	2.5	2.5	2.3	2.3	3.1	4.5	6.9
2002.....	1.3	1.7	2.0	2.4	2.5	2.5	2.5	3.2	4.8	6.9
2001.....	1.5	1.4	1.9	2.2	2.7	2.3	2.7	3.4	4.4	7.4
2000.....	1.3	1.5	2.0	2.2	2.6	2.6	2.4	3.4	5.2	7.8
1999.....	1.3	1.8	1.8	2.4	2.4	2.7	2.4	3.3	5.1	8.0

White, male

2005 4/.....	1.3	1.6	2.0	2.6	2.7	1.8	2.2	2.8	5.2	6.9
2004 4/.....	1.4	1.8	2.0	2.4	2.3	2.4	2.9	3.6	3.9	7.2
2003 4/.....	1.6	1.5	2.0	2.6	2.6	2.3	2.3	2.8	4.3	6.9
2002.....	1.4	1.8	2.0	2.3	2.8	2.6	2.7	3.4	4.6	7.3
2001.....	1.6	1.5	1.9	2.2	2.6	2.7	3.2	3.9	4.6	7.5
2000.....	1.4	1.6	2.3	2.0	2.8	2.6	2.6	3.4	5.6	11.1
1999.....	1.3	1.9	2.0	2.6	2.6	2.5	2.6	3.0	5.8	8.6

White, female

2005 4/.....	1.3	1.4	1.9	2.2	2.5	1.8	2.2	2.2	3.7	6.3
2004 4/.....	1.2	1.3	1.9	2.2	2.3	2.5	2.2	2.8	4.7	6.0
2003 4/.....	1.3	1.4	1.9	2.4	2.4	2.2	2.2	3.3	4.7	6.9
2002.....	1.2	1.5	2.0	2.4	2.2	2.4	2.4	3.0	5.0	6.7
2001.....	1.3	1.4	1.9	2.2	2.8	2.1	2.3	3.1	4.2	7.3
2000.....	1.1	1.3	1.7	2.4	2.4	2.7	2.3	3.4	4.9	6.5
1999.....	1.2	1.7	1.7	2.3	2.3	2.9	2.3	3.6	4.6	7.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)										
All races 3/, both sexes										
2005.....	5.2	6.4	7.0	6.9	7.7	10.4	15.7	27.6	54.0	224.4
2004.....	4.9	5.7	5.8	6.3	7.9	10.2	15.4	28.0	55.5	221.5
2003.....	5.9	7.3	6.8	6.8	8.2	10.6	17.3	26.7	55.7	224.1
2002.....	6.0	6.4	6.9	7.0	8.3	10.4	16.1	28.3	53.3	208.0
2001.....	7.7	9.1	8.8	8.4	9.9	12.7	17.8	29.8	54.3	206.4
2000.....	7.6	8.6	8.0	8.6	10.2	14.2	19.0	30.1	56.5	208.1
1999.....	5.1	5.9	5.8	6.9	8.9	11.9	17.6	26.5	50.5	187.7
All races 3/, male										
2005.....	6.4	8.2	9.3	9.2	9.6	12.8	18.8	31.0	54.8	183.9
2004.....	6.5	7.4	7.7	8.3	10.1	12.6	17.9	30.6	58.1	183.6
2003.....	7.4	9.4	8.9	8.9	10.6	13.7	21.3	31.2	58.0	177.9
2002.....	7.6	8.5	9.0	9.0	11.2	12.7	19.7	32.9	60.6	175.5
2001.....	9.9	11.9	11.8	11.0	13.2	15.5	20.9	33.2	60.6	175.0
2000.....	10.2	11.7	11.2	11.4	13.4	18.9	22.4	36.1	62.3	182.7
1999.....	6.7	8.2	7.9	9.1	11.9	15.1	21.8	31.5	54.4	166.3
All races 3/, female										
2005.....	4.1	4.6	4.8	4.7	6.0	8.3	13.2	25.1	53.5	243.0
2004.....	3.3	4.1	4.0	4.5	5.9	8.1	13.4	26.1	53.8	238.5
2003.....	4.4	5.1	4.8	4.9	6.0	7.9	14.2	23.5	54.3	244.6
2002.....	4.5	4.4	5.0	5.0	5.6	8.4	13.3	25.1	48.9	222.1
2001.....	5.5	6.3	5.9	5.9	6.9	10.2	15.4	27.4	50.4	219.5
2000.....	5.0	5.6	5.0	5.9	7.3	10.1	16.4	26.0	53.0	218.5
1999.....	3.4	3.7	3.9	4.9	6.2	9.2	14.2	23.0	48.3	196.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

White, both sexes										
2005 4/.....	5.0	5.9	6.6	6.3	7.3	9.7	15.0	27.2	53.7	228.9
2004 4/.....	4.7	5.6	5.4	5.8	7.4	9.8	14.7	27.6	55.1	224.4
2003 4/.....	5.6	6.6	6.2	6.3	7.6	10.1	16.6	26.5	56.1	228.3
2002.....	5.7	5.8	6.1	6.4	7.9	9.8	15.9	27.9	53.3	211.6
2001.....	7.6	8.8	8.3	7.9	9.4	11.8	17.2	29.5	53.9	210.0
2000.....	7.4	8.2	7.3	7.9	9.5	13.0	18.5	29.5	56.0	212.2
1999.....	4.4	5.2	5.0	6.2	8.2	11.1	17.0	25.5	49.5	190.2
White, male										
2005 4/.....	6.3	7.8	8.7	8.3	8.9	11.8	17.9	30.4	53.6	185.6
2004 4/.....	6.3	7.2	6.9	7.5	9.3	11.9	17.4	30.1	57.8	186.0
2003 4/.....	7.0	8.6	8.2	8.1	9.7	12.8	20.4	31.0	57.8	180.0
2002.....	7.2	7.6	7.7	8.1	10.7	11.9	19.2	31.6	60.0	178.2
2001.....	10.0	11.6	11.2	10.1	12.2	14.3	20.3	33.2	60.6	177.0
2000.....	10.1	11.2	10.1	10.2	12.4	17.1	21.5	35.4	62.0	186.8
1999.....	5.9	7.0	6.7	8.0	10.9	13.9	21.0	30.3	52.6	167.3
White, female										
2005 4/.....	3.8	4.1	4.5	4.3	5.8	7.9	12.6	24.9	53.7	248.8
2004 4/.....	3.1	4.0	3.8	4.2	5.6	7.9	12.5	25.8	53.5	241.7
2003 4/.....	4.2	4.6	4.2	4.7	5.6	7.7	13.5	23.3	55.0	249.7
2002.....	4.2	4.0	4.4	4.7	5.3	7.9	13.1	25.3	49.2	226.1
2001.....	5.3	6.0	5.5	5.7	6.7	9.7	14.6	26.8	49.9	223.8
2000.....	4.6	5.2	4.6	5.6	6.7	9.4	16.0	25.4	52.4	222.5
1999.....	2.9	3.4	3.4	4.5	5.7	8.6	13.8	22.1	47.6	199.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

Black, both sexes										
2005 4/.....	8.0	11.2	12.4	13.9	14.2	19.2	26.2	38.8	70.9	215.1
2004 4/.....	7.6	8.6	11.0	12.5	14.6	16.1	25.9	39.9	72.0	220.8
2003 4/.....	9.7	13.4	13.4	13.0	15.7	17.9	30.3	34.9	60.9	210.9
2002.....	9.5	12.1	15.7	13.8	13.6	17.8	23.1	39.5	64.9	196.0
2001.....	9.8	13.2	14.6	14.3	15.4	21.7	27.9	38.6	66.6	191.6
2000.....	11.2	13.8	15.9	16.0	19.0	28.3	29.7	44.2	73.4	186.4
1999.....	10.3	13.1	13.3	14.9	17.0	22.3	27.2	42.7	72.5	182.4
Black, male										
2005 4/.....	9.2	13.8	17.2	19.5	20.0	26.4	33.5	49.2	85.7	199.9
2004 4/.....	9.7	11.8	16.3	18.2	20.2	22.3	28.2	45.0	81.1	181.8
2003 4/.....	12.5	17.5	17.8	18.5	22.8	25.9	38.8	41.5	70.1	193.9
2002.....	12.4	17.1	22.1	19.5	20.0	22.9	31.0	57.0	87.1	181.2
2001.....	11.7	17.3	18.9	21.7	25.4	29.5	31.6	40.9	70.6	184.9
2000.....	13.5	18.2	23.6	25.0	25.5	41.4	38.1	54.8	83.5	161.5
1999.....	13.6	19.7	19.1	22.4	24.0	30.1	35.6	54.1	88.1	184.0
Black, female										
2005 4/.....	6.9	8.9	8.4	9.3	9.6	13.9	21.2	32.5	63.1	221.3
2004 4/.....	5.9	5.9	6.6	7.7	10.2	11.5	24.4	36.9	67.3	236.3
2003 4/.....	7.3	9.9	9.7	8.4	10.2	12.1	24.6	30.9	56.1	217.5
2002.....	7.0	7.7	10.2	9.0	8.7	14.1	17.7	29.1	53.3	201.7
2001.....	8.1	9.7	11.0	8.2	7.5	16.0	25.3	37.2	64.6	194.1
2000.....	9.1	9.9	9.4	8.6	13.9	18.6	24.2	37.9	68.4	195.7
1999.....	7.3	7.3	8.5	8.9	11.5	16.5	21.6	35.8	64.8	181.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)

All races 3/, both sexes

2005.....	22.8	33.6	47.8	60.3	81.9	115.6	182.4	313.2	593.5	1,559.1
2004.....	21.9	33.6	45.0	58.1	78.7	111.5	176.4	300.6	554.8	1,461.6
2003.....	23.1	33.3	45.3	56.6	80.9	114.5	178.3	303.7	559.8	1,508.4
2002.....	23.5	33.6	42.6	56.2	77.1	110.4	173.8	297.8	555.5	1,493.0
2001.....	21.7	32.1	40.9	53.2	76.4	112.0	172.3	295.5	539.3	1,476.7
2000.....	22.1	31.0	40.4	53.1	74.1	106.0	168.2	288.9	537.8	1,445.7
1999.....	21.6	29.9	38.4	51.0	72.1	107.7	167.8	281.3	510.8	1,410.7

All races 3/, male

2005.....	27.6	40.7	58.5	71.9	93.1	125.3	194.2	327.1	595.6	1,305.3
2004.....	26.0	40.6	56.0	68.3	88.8	123.1	187.2	315.5	549.4	1,236.6
2003.....	27.8	40.3	55.2	65.6	91.0	125.7	191.0	316.5	563.7	1,258.1
2002.....	28.2	41.0	52.4	64.4	87.9	122.8	186.8	309.6	555.2	1,262.9
2001.....	25.7	38.8	49.1	59.9	85.8	124.3	186.5	307.6	539.6	1,271.6
2000.....	26.3	38.5	47.6	60.9	83.1	117.6	183.1	303.4	537.6	1,266.2
1999.....	26.6	36.3	45.4	59.5	80.9	117.1	184.3	298.1	528.1	1,225.4

All races 3/, female

2005.....	17.9	26.7	37.6	49.4	71.7	107.2	172.9	303.1	592.2	1,675.6
2004.....	17.9	26.7	34.6	48.4	69.5	101.4	167.6	289.9	558.0	1,562.8
2003.....	18.5	26.5	36.0	48.1	71.7	104.8	168.0	294.7	557.4	1,619.1
2002.....	18.9	26.4	33.2	48.6	67.3	99.6	163.5	289.5	555.6	1,592.8
2001.....	17.7	25.6	33.0	46.9	67.9	101.4	161.0	287.0	539.1	1,562.6
2000.....	17.9	23.7	33.4	45.8	65.9	96.1	156.5	278.8	537.8	1,518.8
1999.....	16.6	23.7	31.7	43.1	64.2	99.5	154.8	269.7	500.7	1,485.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)

White, both sexes										
2005 4/.....	21.4	31.1	44.7	57.4	78.8	112.6	179.6	311.2	596.4	1,586.5
2004 4/.....	20.4	31.0	42.0	54.8	75.6	108.1	173.5	298.9	558.1	1,486.9
2003 4/.....	21.4	30.8	41.6	53.5	77.1	111.9	175.4	301.8	562.0	1,534.2
2002.....	21.6	30.9	39.2	53.1	73.4	107.7	170.2	295.6	556.2	1,517.6
2001.....	19.3	28.7	37.3	49.5	72.6	108.9	169.2	292.6	541.2	1,498.0
2000.....	19.8	27.4	37.1	49.3	70.1	102.5	164.5	285.7	537.2	1,464.6
1999.....	19.1	26.3	34.8	47.4	68.6	103.2	163.6	277.1	512.1	1,434.0

White, male										
2005 4/.....	26.8	38.6	55.5	68.5	89.5	120.8	190.8	322.7	594.7	1,328.2
2004 4/.....	25.2	38.4	52.7	64.4	85.2	118.1	183.1	309.7	550.1	1,251.8
2003 4/.....	26.4	38.0	50.8	62.0	86.2	121.6	186.0	311.4	561.0	1,279.1
2002.....	26.8	38.2	48.7	60.6	83.4	118.6	181.2	304.6	553.0	1,274.7
2001.....	23.4	35.3	45.1	55.3	81.8	119.9	182.1	302.0	540.6	1,282.9
2000.....	24.0	35.0	44.1	56.8	77.5	112.7	177.5	298.2	533.4	1,274.8
1999.....	24.3	32.5	41.6	55.4	76.8	111.8	179.1	290.9	525.5	1,231.6

White, female										
2005 4/.....	15.9	23.7	34.2	46.7	68.9	105.3	170.4	302.7	597.5	1,705.4
2004 4/.....	15.7	23.6	31.6	45.5	66.7	99.1	165.6	291.0	563.1	1,592.9
2003 4/.....	16.3	23.6	32.6	45.3	68.6	103.4	166.8	294.9	562.7	1,647.2
2002.....	16.3	23.6	29.9	46.0	64.2	98.1	161.2	289.2	558.2	1,623.1
2001.....	15.3	22.3	29.7	44.0	64.2	99.2	158.8	285.9	541.6	1,588.0
2000.....	15.5	19.8	30.3	42.2	63.2	93.6	154.2	277.0	539.4	1,541.7
1999.....	13.8	20.2	28.3	39.9	61.0	95.7	151.2	267.4	504.2	1,515.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)

Black, both sexes

2005 4/.....	36.6	58.1	80.1	98.8	128.6	164.8	247.6	401.0	680.4	1,501.2
2004 4/.....	35.5	59.5	78.0	99.9	125.7	165.8	244.3	385.2	624.6	1,405.8
2003 4/.....	38.7	58.2	83.5	96.1	131.7	161.7	241.2	382.2	626.5	1,435.7
2002.....	41.6	60.3	78.4	96.1	123.7	155.7	242.8	369.7	642.1	1,425.4
2001.....	41.2	63.0	79.4	95.6	122.1	163.6	237.0	380.4	612.1	1,418.0
2000.....	41.9	64.3	75.6	95.5	122.2	157.2	233.9	372.5	618.9	1,388.9
1999.....	42.5	61.8	75.4	92.1	117.2	167.2	239.7	369.5	556.5	1,286.2

Black, male

2005 4/.....	39.8	65.7	94.2	119.4	151.1	194.4	274.5	447.9	743.7	1,241.0
2004 4/.....	37.6	66.9	95.4	121.7	147.6	200.1	280.0	464.5	659.3	1,291.5
2003 4/.....	42.0	65.9	102.4	113.0	158.7	191.6	289.7	441.8	710.2	1,224.1
2002.....	44.2	70.9	94.6	116.3	146.1	185.2	288.4	419.3	693.1	1,319.8
2001.....	47.0	74.8	95.9	114.8	141.6	196.9	273.3	445.6	659.7	1,313.9
2000.....	48.0	74.7	89.1	114.4	150.1	188.4	279.8	428.6	692.9	1,325.1
1999.....	47.6	73.3	89.5	111.0	134.9	189.5	274.1	434.1	635.7	1,291.3

Black, female

2005 4/.....	33.8	51.4	68.1	81.8	111.1	142.8	229.2	373.1	646.9	1,606.2
2004 4/.....	33.7	53.1	63.3	82.0	108.5	140.3	220.1	338.3	606.3	1,451.1
2003 4/.....	35.7	51.5	67.6	82.2	110.7	139.6	208.5	346.7	582.6	1,517.9
2002.....	39.3	51.0	64.6	79.6	106.1	133.8	212.2	340.0	615.5	1,465.9
2001.....	36.2	52.9	65.3	79.8	106.7	138.9	212.8	341.4	587.9	1,456.6
2000.....	36.5	55.2	64.0	79.9	100.2	134.2	203.3	338.6	581.7	1,412.7
1999.....	37.9	52.0	63.5	76.6	103.3	150.6	216.9	330.7	517.3	1,284.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Accidents (unintentional injuries) (V01-X59,Y85-Y86)										
All races 3/, both sexes										
2005.....	41.6	44.6	41.6	35.5	36.0	40.1	53.7	82.7	137.0	279.5
2004.....	39.7	42.6	38.6	32.8	33.7	38.4	50.5	79.6	135.9	276.7
2003.....	40.4	40.7	36.6	32.8	32.9	38.1	50.9	78.8	133.5	278.9
2002.....	39.4	38.7	34.3	30.4	32.6	38.3	50.8	80.0	131.1	275.4
2001.....	36.7	36.0	31.9	29.3	31.5	36.8	49.3	80.4	130.4	276.4
2000.....	35.4	34.2	30.8	29.9	32.2	35.4	48.8	75.5	124.6	273.5
1999.....	35.4	34.1	29.2	30.1	31.2	38.7	50.9	78.6	133.3	282.4
All races 3/, male										
2005.....	58.7	63.5	60.7	51.3	51.4	54.6	72.6	110.0	180.8	344.2
2004.....	56.0	60.5	56.5	46.9	46.8	52.8	67.0	104.0	179.7	345.5
2003.....	58.0	58.9	52.3	47.2	47.0	52.3	66.8	104.9	175.2	352.5
2002.....	56.8	56.6	49.6	44.8	45.7	53.4	68.9	106.4	170.1	346.6
2001.....	54.1	54.0	46.9	41.8	44.5	49.9	65.6	108.3	173.2	358.5
2000.....	53.1	51.6	46.0	43.1	45.3	48.1	64.6	99.8	164.8	359.7
1999.....	53.1	51.7	43.6	43.6	44.5	53.1	68.2	102.8	181.8	375.4
All races 3/, female										
2005.....	24.6	26.0	23.3	20.7	22.0	27.4	38.5	62.8	109.7	249.8
2004.....	23.6	25.2	21.4	19.5	21.7	25.8	37.2	62.1	109.0	245.8
2003.....	23.0	23.0	21.6	19.3	20.1	25.8	38.1	60.3	108.1	246.4
2002.....	22.4	21.3	19.6	16.8	20.7	25.3	36.3	61.4	107.3	244.6
2001.....	19.6	18.5	17.5	17.6	19.6	25.5	36.3	60.9	104.8	242.0
2000.....	18.0	17.3	16.1	17.5	20.4	24.5	36.4	58.5	100.9	238.4
1999.....	18.1	17.0	15.5	17.6	19.2	26.4	37.3	61.8	105.0	244.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Accidents (unintentional injuries) (V01-X59,Y85-Y86)										
White, both sexes										
2005 4/.....	43.2	45.4	40.7	34.7	35.3	39.8	53.8	83.1	140.0	290.9
2004 4/.....	41.3	43.2	37.9	32.0	33.3	38.2	50.5	80.6	140.2	289.9
2003 4/.....	41.5	41.0	35.7	32.1	32.5	37.3	50.9	80.2	136.5	291.1
2002.....	40.5	38.5	33.6	29.6	32.1	37.3	51.0	81.1	134.6	286.1
2001.....	36.8	34.9	31.1	28.5	30.5	36.4	48.9	81.2	133.3	285.5
2000.....	35.2	33.2	29.2	28.9	31.8	34.5	48.1	75.7	127.3	282.8
1999.....	35.2	32.8	27.8	28.9	29.5	37.7	49.5	78.0	135.0	290.4
White, male										
2005 4/.....	60.6	63.8	58.6	49.2	49.2	53.7	71.7	110.0	182.9	356.6
2004 4/.....	57.9	61.0	54.7	44.8	45.4	52.4	66.5	104.6	183.9	359.4
2003 4/.....	59.6	58.6	50.6	45.6	46.1	50.8	66.9	105.8	177.3	365.7
2002.....	58.2	56.4	47.8	42.8	44.5	50.9	68.6	106.5	173.0	356.6
2001.....	54.2	51.8	45.1	40.0	42.6	48.9	64.6	108.2	175.2	368.1
2000.....	52.5	49.8	43.2	40.8	43.8	45.9	63.3	98.7	166.4	369.8
1999.....	52.5	49.1	41.1	41.1	42.0	51.1	65.7	100.7	183.4	383.6
White, female										
2005 4/.....	25.6	26.9	23.0	20.8	22.4	27.5	39.0	63.2	113.1	260.7
2004 4/.....	24.5	25.5	21.5	19.7	22.0	25.6	37.3	63.0	113.0	258.6
2003 4/.....	23.4	23.4	21.1	19.2	19.9	25.4	37.9	61.7	111.3	258.0
2002.....	22.8	20.7	19.7	16.9	20.7	25.3	36.7	62.9	111.2	255.4
2001.....	19.3	18.1	17.4	17.6	19.3	25.4	36.1	62.0	107.9	251.0
2000.....	17.8	16.8	15.5	17.5	20.7	24.6	35.8	59.6	104.0	247.4
1999.....	17.9	16.6	14.9	17.4	18.0	26.1	36.6	62.0	106.5	253.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Accidents (unintentional injuries) (V01-X59,Y85-Y86)										
Black, both sexes										
2005 4/.....	41.1	49.5	57.2	49.7	45.6	47.5	58.9	87.4	110.9	174.8
2004 4/.....	38.4	49.0	50.7	45.2	41.7	43.8	53.8	77.9	100.5	152.3
2003 4/.....	41.7	47.9	49.1	43.7	40.4	47.7	53.8	67.7	105.1	160.2
2002.....	40.0	48.3	45.1	42.0	40.6	47.3	51.9	74.9	96.8	173.4
2001.....	44.3	50.1	44.8	40.6	43.7	45.5	58.0	78.2	104.5	183.6
2000.....	43.8	47.0	47.0	42.0	39.4	46.1	60.2	75.3	100.0	181.1
1999.....	43.6	50.4	44.4	43.4	46.3	50.7	66.6	90.3	124.2	207.6
Black, male										
2005 4/.....	60.1	75.7	90.9	80.3	79.6	73.2	90.5	130.5	167.1	225.0
2004 4/.....	56.5	72.2	82.2	73.2	66.7	67.4	82.2	116.0	157.1	213.6
2003 4/.....	60.8	73.9	75.2	70.1	64.8	74.4	74.2	105.0	152.8	225.2
2002.....	57.9	72.0	72.6	71.2	65.3	75.8	78.4	117.6	149.8	260.5
2001.....	66.6	81.7	71.8	65.9	69.8	69.1	82.5	119.4	164.5	255.3
2000.....	68.8	74.5	76.5	71.5	65.4	74.2	86.3	119.5	156.0	275.6
1999.....	69.0	82.7	73.2	71.5	70.6	77.9	101.8	135.3	177.0	316.4
Black, female										
2005 4/.....	24.3	26.7	28.7	24.5	19.1	28.4	37.2	61.6	81.2	154.5
2004 4/.....	22.6	28.8	24.2	22.0	22.3	26.3	34.5	55.3	70.7	128.0
2003 4/.....	24.8	25.3	27.0	22.0	21.4	28.0	40.1	45.6	80.1	134.9
2002.....	24.2	27.7	21.8	18.1	21.2	26.1	34.1	49.4	69.2	140.1
2001.....	24.5	22.7	21.8	19.9	23.2	28.0	41.7	53.5	74.0	157.0
2000.....	21.7	23.1	21.9	17.8	18.8	25.3	42.8	48.7	71.8	145.9
1999.....	21.3	22.5	20.1	20.4	27.3	30.6	43.2	63.2	98.1	166.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Transport accidents (V01-V99,Y85)										
All races 3/, both sexes										
2005.....	17.0	16.8	16.6	16.2	16.3	17.2	18.9	23.0	26.0	26.1
2004.....	16.4	17.1	16.3	15.2	15.9	17.0	18.1	23.6	28.1	27.2
2003.....	17.5	16.8	16.3	15.9	16.0	16.2	19.0	23.9	29.4	29.7
2002.....	17.1	16.9	15.8	14.9	16.7	16.7	20.2	25.0	29.7	28.8
2001.....	17.3	16.3	15.6	15.2	15.4	16.4	19.5	26.2	30.4	28.9
2000.....	16.8	16.2	15.5	15.7	16.4	16.8	19.4	25.1	30.0	31.5
1999.....	16.6	15.9	14.9	16.3	15.7	18.1	20.9	25.9	32.1	32.0
All races 3/, male										
2005.....	24.8	24.9	25.0	23.7	23.2	23.9	27.0	32.5	38.0	46.0
2004.....	24.1	24.9	24.4	22.2	22.4	23.2	24.3	32.3	42.6	46.5
2003.....	25.5	24.9	23.6	23.2	22.7	22.5	25.0	32.9	42.1	52.0
2002.....	24.5	25.1	22.7	22.2	22.8	23.4	27.5	34.5	42.6	53.0
2001.....	25.4	24.0	22.5	21.4	22.0	22.1	26.8	36.2	45.5	52.7
2000.....	24.8	23.6	22.3	22.4	22.6	22.6	26.3	34.5	44.0	59.8
1999.....	24.0	23.1	21.5	23.0	21.8	24.8	27.5	36.2	48.3	63.5
All races 3/, female										
2005.....	9.2	8.9	8.6	9.0	10.0	11.3	12.4	16.1	18.5	17.0
2004.....	8.9	9.4	8.6	8.7	10.0	11.5	13.1	17.3	19.1	18.5
2003.....	9.5	9.0	9.2	9.0	9.9	10.7	14.2	17.5	21.7	19.9
2002.....	9.9	8.8	9.2	7.9	11.1	10.9	14.3	18.3	21.8	18.3
2001.....	9.4	8.8	8.9	9.4	9.3	11.6	13.8	19.2	21.3	18.9
2000.....	9.0	8.9	8.9	9.5	10.7	11.7	13.9	18.5	21.7	19.9
1999.....	9.4	8.9	8.6	10.0	10.2	12.4	15.7	18.8	22.6	19.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Transport accidents (V01-V99,Y85)										
White, both sexes										
2005 4/.....	17.5	17.2	16.6	16.4	16.2	17.4	19.0	23.0	26.8	27.1
2004 4/.....	16.8	17.2	16.0	15.2	15.9	17.1	18.2	23.6	29.0	28.4
2003 4/.....	17.7	16.9	16.1	15.7	15.8	16.0	19.2	23.9	30.0	30.5
2002.....	17.5	16.9	15.9	14.8	16.7	16.5	20.5	25.5	30.3	29.6
2001.....	17.5	15.9	15.7	15.1	15.2	16.4	19.5	26.6	31.2	29.9
2000.....	16.8	16.0	15.1	15.6	16.4	16.6	19.2	25.3	30.7	32.2
1999.....	16.7	15.6	14.6	16.1	15.4	18.0	20.8	26.0	32.4	32.3
White, male										
2005 4/.....	25.3	25.3	24.8	23.8	23.0	23.7	26.9	32.1	39.0	47.5
2004 4/.....	24.4	25.0	23.7	21.9	22.0	23.2	24.1	32.2	43.2	48.6
2003 4/.....	25.8	24.9	23.3	22.8	22.6	22.2	25.4	32.7	42.6	52.8
2002.....	24.9	25.3	22.5	21.9	22.8	22.7	27.5	34.4	42.8	53.8
2001.....	25.4	23.2	22.6	21.0	21.5	21.7	26.6	36.2	46.6	54.1
2000.....	24.6	23.4	21.7	21.8	22.2	21.9	25.8	34.3	44.5	61.6
1999.....	24.0	22.5	21.0	22.5	21.1	24.6	27.1	35.9	48.4	63.7
White, female										
2005 4/.....	9.5	9.1	8.5	9.2	10.0	11.7	12.5	16.2	19.2	17.7
2004 4/.....	9.0	9.5	8.5	8.8	10.3	11.6	13.4	17.3	20.1	19.3
2003 4/.....	9.6	9.1	9.1	8.8	9.6	10.5	14.0	17.6	22.2	20.6
2002.....	10.0	8.6	9.3	8.1	11.0	11.0	14.8	19.1	22.6	19.1
2001.....	9.5	8.7	8.9	9.5	9.3	11.6	13.8	19.8	21.8	19.7
2000.....	9.0	8.7	8.6	9.6	11.1	12.0	13.9	19.0	22.6	20.3
1999.....	9.4	8.8	8.4	9.9	10.1	12.3	15.8	19.1	23.0	19.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Transport accidents (V01-V99,Y85)										
Black, both sexes										
2005 4/.....	16.8	16.9	18.9	16.9	17.3	17.2	18.8	22.4	16.3	16.6
2004 4/.....	16.7	19.1	19.9	17.2	17.4	16.6	16.2	23.0	19.6	14.7
2003 4/.....	18.7	18.1	18.2	18.9	17.8	18.5	17.9	22.8	21.8	21.0
2002.....	17.1	18.9	16.5	16.8	17.4	17.3	16.9	20.1	24.1	18.4
2001.....	19.4	20.5	16.8	17.8	18.6	18.6	19.0	22.1	19.8	17.5
2000.....	19.3	19.4	19.3	18.1	16.4	18.2	21.3	19.4	20.9	21.6
1999.....	18.0	19.6	18.3	18.7	17.7	18.0	21.0	24.6	28.3	27.7
Black, male										
2005 4/.....	26.2	25.5	30.8	27.4	27.8	28.8	30.4	38.9	27.4	33.5
2004 4/.....	25.8	29.6	32.6	27.1	28.6	27.1	26.2	36.9	40.9	28.8
2003 4/.....	28.3	29.1	28.9	30.3	27.6	28.6	24.0	37.4	37.7	42.7
2002.....	25.2	29.1	26.3	29.0	27.3	28.7	28.4	38.1	44.3	46.1
2001.....	30.6	33.8	25.7	28.6	30.9	28.2	26.3	39.0	31.3	35.2
2000.....	31.0	29.3	30.0	31.0	27.5	31.2	33.4	34.5	39.6	45.0
1999.....	28.3	31.2	29.1	29.3	28.4	27.3	31.5	40.4	48.8	68.0
Black, female										
2005 4/.....	8.5	9.4	8.8	8.3	9.1	8.6	10.8	12.5	10.4	9.8
2004 4/.....	8.6	10.0	9.1	9.1	8.7	8.8	9.4	14.8	8.3	9.1
2003 4/.....	10.3	8.4	9.2	9.6	10.2	11.0	13.7	14.1	13.5	12.6
2002.....	10.0	10.0	8.2	6.8	9.6	8.9	9.2	9.3	13.5	*
2001.....	9.4	8.9	9.3	8.9	9.0	11.5	14.1	11.9	13.9	10.9
2000.....	9.0	10.8	10.2	7.6	7.7	8.6	13.3	10.3	11.5	12.9
1999.....	9.0	9.6	9.1	10.1	9.3	11.2	14.0	15.2	18.1	12.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)

All races 3/, both sexes

2005.....	15.6	15.3	14.8	14.6	14.8	15.8	17.8	21.6	24.6	25.1
2004.....	15.1	15.5	14.6	13.9	14.3	15.5	16.8	22.7	26.9	26.1
2003.....	16.0	15.1	14.6	14.1	14.3	14.8	17.7	22.5	28.2	28.8
2002.....	15.7	15.2	14.4	13.4	15.2	15.3	18.9	23.8	28.4	28.0
2001.....	15.5	14.5	13.9	13.7	13.7	14.8	18.0	24.6	28.9	27.8
2000.....	15.1	14.6	13.8	14.2	14.7	15.1	18.1	23.8	28.5	30.4
1999.....	14.8	14.0	13.1	14.4	13.9	16.4	19.3	24.1	30.1	30.4

All races 3/, male

2005.....	22.6	22.2	22.1	21.0	20.7	21.5	25.2	30.1	35.7	44.1
2004.....	21.9	22.3	21.6	19.9	19.6	20.8	22.0	30.8	40.4	44.2
2003.....	23.1	22.0	20.8	20.3	19.6	20.1	22.8	30.5	40.0	50.0
2002.....	22.1	22.2	20.3	19.6	20.2	20.8	25.0	32.3	39.9	51.7
2001.....	22.5	21.2	19.7	18.9	19.2	19.2	24.2	33.6	43.1	51.1
2000.....	21.9	21.1	19.6	19.9	19.7	19.8	23.9	32.2	41.4	57.5
1999.....	21.0	20.0	18.6	19.8	18.7	22.0	24.7	33.6	45.2	59.8

All races 3/, female

2005.....	8.7	8.5	7.9	8.5	9.4	10.7	11.8	15.5	17.8	16.4
2004.....	8.3	8.8	8.0	8.2	9.5	10.9	12.6	16.9	18.6	17.9
2003.....	9.0	8.3	8.7	8.3	9.4	10.3	13.7	16.9	21.0	19.5
2002.....	9.4	8.4	8.7	7.5	10.6	10.6	14.1	17.7	21.4	17.7
2001.....	8.7	8.0	8.3	8.8	8.8	11.0	13.1	18.2	20.4	18.0
2000.....	8.4	8.3	8.2	9.0	10.2	11.0	13.5	17.9	20.9	19.3
1999.....	8.6	8.1	8.0	9.4	9.5	11.6	15.0	17.5	21.3	18.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)

White, both sexes

2005 4/.....	16.1	15.7	14.9	14.7	14.7	15.9	17.8	21.6	25.5	26.1
2004 4/.....	15.4	15.6	14.3	13.9	14.3	15.6	16.8	22.8	27.7	27.3
2003 4/.....	16.2	15.2	14.4	13.8	14.0	14.6	17.8	22.6	28.8	29.5
2002.....	16.0	15.2	14.3	13.3	15.1	15.1	19.2	24.3	29.0	28.8
2001.....	15.6	14.2	13.9	13.6	13.5	14.7	17.9	25.0	29.7	28.8
2000.....	15.0	14.5	13.4	14.1	14.7	14.9	17.9	24.0	29.2	31.1
1999.....	14.8	13.7	12.9	14.1	13.5	16.3	19.2	24.3	30.6	30.7

White, male

2005 4/.....	23.1	22.7	22.1	21.0	20.5	21.4	25.0	29.7	36.7	45.6
2004 4/.....	22.2	22.3	20.9	19.6	19.2	20.7	21.7	30.8	40.8	46.3
2003 4/.....	23.2	22.1	20.5	19.7	19.4	19.9	23.0	30.3	40.6	50.7
2002.....	22.5	22.3	20.0	19.3	20.0	20.1	25.0	32.4	40.1	52.4
2001.....	22.5	20.5	19.7	18.7	18.5	19.0	23.9	33.6	44.3	52.6
2000.....	21.6	20.9	18.9	19.3	19.2	19.1	23.4	32.0	41.7	59.2
1999.....	20.9	19.4	18.1	19.2	18.1	21.8	24.4	33.3	45.5	60.3

White, female

2005 4/.....	9.0	8.7	7.8	8.7	9.3	11.0	11.9	15.5	18.5	17.1
2004 4/.....	8.4	8.9	7.9	8.3	9.8	11.1	12.8	16.9	19.6	18.7
2003 4/.....	9.0	8.4	8.6	8.1	9.1	10.0	13.5	17.1	21.5	20.1
2002.....	9.5	8.2	8.8	7.6	10.5	10.7	14.5	18.5	22.2	18.5
2001.....	8.8	8.0	8.2	8.8	8.8	11.0	13.0	19.0	20.9	18.9
2000.....	8.4	8.1	7.9	9.1	10.6	11.3	13.5	18.4	21.8	19.7
1999.....	8.6	8.0	7.7	9.3	9.3	11.6	15.1	18.0	21.8	18.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)

Black, both sexes

2005 4/.....	15.6	15.0	16.7	15.9	16.2	15.9	18.0	21.4	15.0	15.8
2004 4/.....	15.4	17.3	18.3	15.9	15.5	15.6	15.3	20.8	19.3	14.7
2003 4/.....	17.5	16.2	16.8	17.5	16.4	16.9	17.2	21.3	20.6	20.7
2002.....	15.9	16.9	15.1	15.0	16.2	15.9	15.8	18.4	22.5	18.1
2001.....	17.6	18.1	15.3	15.9	16.9	16.5	18.3	20.1	19.0	16.6
2000.....	17.9	17.6	18.3	16.7	15.4	16.6	20.1	18.1	20.0	20.3
1999.....	16.6	17.8	16.3	17.1	15.8	16.6	19.6	22.9	26.0	25.5

Black, male

2005 4/.....	24.3	22.4	26.8	25.5	26.0	25.7	29.4	36.7	26.0	32.5
2004 4/.....	23.5	26.7	29.7	24.5	25.0	25.4	24.3	33.2	40.2	28.8
2003 4/.....	26.4	25.8	26.0	27.6	24.5	25.1	23.0	35.1	34.7	41.7
2002.....	23.0	25.6	23.6	25.3	24.9	25.4	25.7	33.5	40.5	45.0
2001.....	27.5	29.9	23.2	24.7	28.5	23.9	24.6	36.2	30.5	35.2
2000.....	28.4	26.3	28.1	28.2	25.7	27.8	31.4	32.1	38.8	43.8
1999.....	25.6	28.2	25.5	26.3	24.6	24.9	28.5	38.9	45.3	60.9

Black, female

2005 4/.....	7.9	8.5	8.1	8.0	8.6	8.6	10.2	12.3	9.3	9.0
2004 4/.....	8.3	9.1	8.8	8.8	8.0	8.3	9.2	13.5	8.3	9.1
2003 4/.....	9.7	7.9	9.1	9.2	10.1	10.8	13.3	13.0	13.2	12.6
2002.....	9.6	9.4	8.0	6.6	9.4	8.9	9.2	9.3	13.1	*
2001.....	8.9	8.0	8.7	8.7	7.8	11.1	14.1	10.5	13.1	9.7
2000.....	8.6	10.0	9.9	7.3	7.4	8.3	12.7	9.7	10.6	11.6
1999.....	8.7	8.9	8.5	9.6	8.8	10.4	13.7	13.2	16.4	12.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9,V88.9,V89.1,V89.3,V89.9)

All races 3/, both sexes

2005.....	0.5	0.5	0.7	0.5	0.5	0.4	0.4	0.6	0.6	0.5
2004.....	0.5	0.6	0.5	0.4	0.5	0.4	0.6	0.4	0.6	0.6
2003.....	0.6	0.6	0.5	0.6	0.5	0.4	0.5	0.6	0.7	0.5
2002.....	0.5	0.6	0.5	0.4	0.4	0.3	0.4	0.5	0.6	*
2001.....	0.6	0.7	0.6	0.5	0.6	0.7	0.6	0.8	0.8	0.7
2000.....	0.7	0.6	0.6	0.5	0.5	0.5	0.5	0.7	0.8	0.8
1999.....	0.9	0.9	0.7	0.8	0.8	0.8	0.7	1.3	1.4	1.3

All races 3/, male

2005.....	0.8	0.9	1.2	0.7	0.8	0.7	0.6	1.1	1.1	*
2004.....	0.8	0.9	0.8	0.7	0.7	0.6	0.8	0.6	1.2	*
2003.....	0.9	1.1	0.8	0.9	0.8	0.8	0.7	0.8	1.3	*
2002.....	0.9	1.0	0.8	0.7	0.7	0.6	0.8	0.8	1.2	*
2001.....	1.0	1.0	0.9	0.8	0.9	1.1	0.9	1.2	*	*
2000.....	1.2	0.9	0.9	0.7	1.0	0.9	0.9	1.1	1.3	*
1999.....	1.5	1.4	1.2	1.2	1.1	1.2	1.2	1.7	2.1	2.9

All races 3/, female

2005.....	0.2	*	0.4	*	*	*	*	*	*	*
2004.....	0.2	0.2	*	*	*	*	*	*	*	*
2003.....	0.3	0.2	*	*	*	*	*	*	*	*
2002.....	0.2	*	*	*	*	*	*	*	*	*
2001.....	0.2	0.4	0.2	*	*	*	*	0.5	0.7	*
2000.....	0.3	0.3	0.4	*	*	*	*	*	*	*
1999.....	0.4	0.4	0.3	0.4	0.4	0.5	0.4	0.9	1.0	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9,V81.2-V81.9, V82.2-V82.9,V87.9,V88.9,V89.1,V89.3,V89.9)

White, both sexes

2005 4/.....	0.5	0.5	0.6	0.5	0.4	0.4	0.3	0.6	0.7	0.5
2004 4/.....	0.5	0.5	0.4	0.4	0.4	0.4	0.6	0.3	0.7	0.6
2003 4/.....	0.6	0.6	0.5	0.5	0.5	0.4	0.4	0.6	0.7	0.5
2002.....	0.5	0.5	0.5	0.4	0.4	0.3	0.4	0.4	0.6	*
2001.....	0.6	0.6	0.6	0.5	0.5	0.6	0.6	0.8	0.7	0.6
2000.....	0.7	0.6	0.6	0.4	0.6	0.5	0.5	0.7	0.8	0.8
1999.....	0.9	0.9	0.7	0.8	0.8	0.8	0.7	1.2	1.4	1.3

White, male

2005 4/.....	0.7	0.8	0.9	0.8	0.6	0.5	*	1.0	1.1	*
2004 4/.....	0.8	0.8	0.7	0.7	0.7	0.6	0.8	*	1.4	*
2003 4/.....	0.9	0.9	0.8	0.9	0.8	0.7	0.7	0.8	1.2	*
2002.....	0.8	0.9	0.8	0.6	0.7	0.5	0.7	*	1.3	*
2001.....	0.9	0.9	0.9	0.8	0.9	1.0	0.9	1.3	*	*
2000.....	1.2	0.8	0.8	0.6	1.0	0.8	0.9	1.1	1.3	*
1999.....	1.5	1.3	1.2	1.3	1.1	1.2	1.1	1.7	2.0	2.8

White, female

2005 4/.....	0.2	*	0.3	*	*	*	*	*	*	*
2004 4/.....	0.2	*	*	*	*	*	*	*	*	*
2003 4/.....	0.2	0.2	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	0.2	0.3	*	*	*	*	*	*	*	*
2000.....	0.3	0.3	0.4	*	*	*	*	*	*	*
1999.....	0.4	0.5	0.3	0.4	0.4	0.5	*	0.9	1.0	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)

All races 3/, both sexes

2005.....	0.8	1.0	1.0	1.1	1.0	1.0	0.7	0.7	0.7	0.5
2004.....	0.8	1.0	1.2	0.9	1.1	1.0	0.8	0.5	0.5	0.5
2003.....	0.9	1.1	1.1	1.2	1.2	0.9	0.8	0.8	0.5	*
2002.....	0.9	1.1	1.0	1.1	1.1	1.0	0.8	0.8	0.7	0.4
2001.....	1.2	1.1	1.2	1.0	1.1	1.0	0.9	0.8	0.7	0.5
2000.....	1.0	0.9	1.0	1.0	1.1	1.2	0.7	0.6	0.7	*
1999.....	0.9	1.0	1.0	1.1	1.1	0.9	0.8	0.5	0.6	*

All races 3/, male

2005.....	1.4	1.7	1.8	1.9	1.7	1.7	1.3	1.3	1.2	*
2004.....	1.3	1.7	2.0	1.5	2.1	1.8	1.5	0.9	1.0	*
2003.....	1.6	1.7	2.0	2.0	2.3	1.7	1.5	1.6	*	*
2002.....	1.5	1.9	1.6	1.9	1.9	2.0	1.7	1.4	1.5	*
2001.....	1.8	1.8	1.9	1.7	1.9	1.8	1.7	1.3	1.4	*
2000.....	1.7	1.6	1.8	1.8	1.9	2.0	1.4	1.2	1.4	*
1999.....	1.5	1.7	1.8	2.0	1.9	1.6	1.6	0.9	*	*

All races 3/, female

2005.....	0.3	0.3	0.3	0.3	0.4	0.4	*	*	*	*
2004.....	0.3	0.4	0.4	0.3	*	*	*	*	*	*
2003.....	0.3	0.4	0.3	0.4	*	*	*	*	*	*
2002.....	0.3	0.3	0.3	0.3	0.4	*	*	*	*	*
2001.....	0.5	0.4	0.4	0.4	*	*	*	0.5	*	*
2000.....	0.3	0.3	0.3	*	0.4	0.4	*	*	*	*
1999.....	0.4	0.3	0.3	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)

White, both sexes										
2005 4/.....	0.9	1.0	1.1	1.2	1.1	1.1	0.8	0.8	0.7	0.5
2004 4/.....	0.9	1.1	1.3	0.9	1.2	1.1	0.8	0.5	0.6	0.6
2003 4/.....	1.0	1.1	1.2	1.3	1.3	0.9	0.9	0.8	0.5	*
2002.....	1.0	1.2	1.0	1.1	1.2	1.0	0.9	0.8	0.7	*
2001.....	1.3	1.1	1.2	1.0	1.2	1.0	1.0	0.8	0.7	*
2000.....	1.1	1.0	1.1	1.1	1.2	1.2	0.8	0.6	0.7	*
1999.....	1.0	1.1	1.1	1.1	1.1	0.9	0.9	0.5	0.5	*
White, male										
2005 4/.....	1.6	1.8	1.8	2.1	1.9	1.8	1.4	1.4	1.2	*
2004 4/.....	1.4	1.9	2.1	1.6	2.1	1.9	1.6	0.9	1.0	*
2003 4/.....	1.7	1.9	2.1	2.2	2.5	1.6	1.7	1.6	*	*
2002.....	1.6	2.1	1.7	1.9	2.1	2.0	1.8	1.3	1.5	*
2001.....	2.0	1.8	2.0	1.6	2.1	1.8	1.8	1.3	1.5	*
2000.....	1.9	1.6	2.0	1.9	2.0	2.0	1.5	1.2	1.5	*
1999.....	1.5	1.8	1.8	2.0	1.9	1.6	1.6	0.9	*	*
White, female										
2005 4/.....	0.3	0.3	0.4	0.3	0.4	0.4	*	*	*	*
2004 4/.....	0.4	0.4	0.4	0.3	*	*	*	*	*	*
2003 4/.....	0.4	0.4	0.4	0.4	*	*	*	*	*	*
2002.....	0.3	0.2	0.4	0.4	0.4	*	*	*	*	*
2001.....	0.6	0.4	0.5	0.5	*	*	*	*	*	*
2000.....	0.4	0.3	0.3	*	0.4	*	*	*	*	*
1999.....	0.4	0.3	0.3	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Nontransport accidents (W00-X59,Y86)										
All races 3/, both sexes										
2005.....	24.6	27.8	25.0	19.4	19.8	22.9	34.8	59.7	111.0	253.4
2004.....	23.3	25.5	22.3	17.6	17.8	21.4	32.4	56.0	107.8	249.5
2003.....	22.9	23.9	20.3	17.0	16.9	21.9	31.9	54.9	104.1	249.2
2002.....	22.3	21.8	18.4	15.5	16.0	21.7	30.6	55.0	101.4	246.7
2001.....	19.4	19.7	16.3	14.1	16.1	20.3	29.8	54.3	100.1	247.5
2000.....	18.6	18.0	15.3	14.2	15.9	18.6	29.4	50.4	94.6	242.0
1999.....	18.8	18.2	14.3	13.9	15.5	20.6	30.0	52.7	101.3	250.3
All races 3/, male										
2005.....	33.9	38.7	35.7	27.5	28.3	30.7	45.6	77.5	142.8	298.3
2004.....	31.9	35.6	32.1	24.7	24.5	29.6	42.8	71.7	137.0	299.0
2003.....	32.4	34.1	28.7	24.0	24.3	29.7	41.8	72.0	133.0	300.5
2002.....	32.3	31.5	26.8	22.6	22.9	30.0	41.4	71.9	127.5	293.6
2001.....	28.7	30.0	24.3	20.5	22.5	27.8	38.8	72.1	127.8	305.8
2000.....	28.3	28.0	23.7	20.7	22.7	25.5	38.4	65.3	120.8	299.8
1999.....	29.1	28.6	22.0	20.6	22.7	28.3	40.7	66.6	133.5	311.9
All races 3/, female										
2005.....	15.3	17.1	14.7	11.7	12.0	16.0	26.1	46.7	91.3	232.8
2004.....	14.7	15.8	12.8	10.8	11.7	14.3	24.1	44.8	89.9	227.3
2003.....	13.5	14.1	12.3	10.4	10.2	15.1	23.9	42.7	86.4	226.5
2002.....	12.5	12.5	10.4	8.9	9.7	14.5	21.9	43.1	85.5	226.3
2001.....	10.2	9.7	8.6	8.2	10.3	13.9	22.6	41.8	83.5	223.1
2000.....	9.0	8.4	7.3	8.0	9.7	12.8	22.4	40.0	79.2	218.4
1999.....	8.7	8.1	6.9	7.6	9.0	14.0	21.7	43.0	82.4	225.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Nontransport accidents (W00-X59,Y86)										
White, both sexes										
2005 4/.....	25.7	28.1	24.1	18.3	19.1	22.4	34.8	60.2	113.2	263.9
2004 4/.....	24.5	26.0	21.9	16.8	17.4	21.2	32.3	57.0	111.2	261.5
2003 4/.....	23.8	24.0	19.5	16.5	16.7	21.4	31.8	56.2	106.5	260.6
2002.....	23.1	21.6	17.7	14.7	15.5	20.8	30.5	55.6	104.3	256.5
2001.....	19.3	19.0	15.4	13.4	15.3	20.0	29.4	54.6	102.1	255.7
2000.....	18.3	17.2	14.1	13.3	15.3	17.9	28.9	50.4	96.5	250.6
1999.....	18.5	17.1	13.2	12.9	14.2	19.7	28.7	51.9	102.6	258.1
White, male										
2005 4/.....	35.2	38.5	33.8	25.4	26.2	29.9	44.8	77.9	143.9	309.1
2004 4/.....	33.5	36.0	30.9	22.9	23.4	29.2	42.4	72.5	140.7	310.8
2003 4/.....	33.8	33.8	27.2	22.8	23.4	28.6	41.5	73.1	134.7	312.9
2002.....	33.3	31.1	25.2	20.9	21.7	28.2	41.1	72.1	130.1	302.8
2001.....	28.7	28.6	22.4	19.0	21.1	27.1	38.1	72.0	128.6	314.0
2000.....	27.9	26.4	21.5	19.0	21.6	24.0	37.5	64.4	122.0	308.2
1999.....	28.5	26.6	20.0	18.6	20.9	26.5	38.6	64.8	135.0	319.8
White, female										
2005 4/.....	16.1	17.8	14.5	11.6	12.5	15.8	26.5	47.1	94.0	243.0
2004 4/.....	15.5	16.0	13.1	10.9	11.7	14.1	24.0	45.7	92.9	239.3
2003 4/.....	13.8	14.3	12.0	10.4	10.4	14.9	23.9	44.0	89.1	237.4
2002.....	12.8	12.1	10.4	8.8	9.7	14.3	21.9	43.8	88.5	236.3
2001.....	9.7	9.4	8.5	8.1	10.0	13.8	22.3	42.2	86.1	231.2
2000.....	8.7	8.1	6.9	7.9	9.5	12.6	21.9	40.6	81.4	227.2
1999.....	8.5	7.8	6.5	7.4	8.0	13.7	20.8	42.9	83.5	233.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Nontransport accidents (W00-X59,Y86)										
Black, both sexes										
2005 4/.....	24.3	32.6	38.3	32.8	28.3	30.3	40.1	65.0	94.7	158.2
2004 4/.....	21.8	29.9	30.9	27.9	24.3	27.2	37.6	54.8	81.0	137.6
2003 4/.....	22.9	29.9	30.9	24.8	22.6	29.3	36.0	44.9	83.3	139.2
2002.....	22.9	29.4	28.6	25.2	23.2	29.9	35.0	54.8	72.7	155.0
2001.....	24.9	29.6	28.0	22.8	25.1	26.9	39.0	56.1	84.8	166.1
2000.....	24.5	27.6	27.7	23.9	22.9	27.8	38.8	56.0	79.1	159.5
1999.....	25.6	30.7	26.2	24.6	28.6	32.7	45.6	65.7	95.9	179.8
Black, male										
2005 4/.....	34.0	50.2	60.1	52.9	51.8	44.4	60.1	91.6	139.8	191.5
2004 4/.....	30.7	42.6	49.6	46.2	38.1	40.2	56.0	79.2	116.2	184.8
2003 4/.....	32.4	44.8	46.3	39.8	37.2	45.8	50.3	67.6	115.2	182.5
2002.....	32.7	42.9	46.3	42.1	38.0	47.1	50.1	79.5	105.5	214.4
2001.....	36.0	47.8	46.1	37.2	38.8	40.9	56.2	80.4	133.2	220.1
2000.....	37.8	45.3	46.5	40.5	37.9	42.9	52.8	85.1	116.3	230.6
1999.....	40.7	51.5	44.1	42.2	42.2	50.6	70.2	95.0	128.2	248.4
Black, female										
2005 4/.....	15.8	17.3	19.8	16.2	10.1	19.7	26.4	49.1	70.8	144.8
2004 4/.....	13.9	18.9	15.1	12.9	13.5	17.4	25.0	40.4	62.4	118.9
2003 4/.....	14.5	16.8	17.8	12.5	11.1	17.0	26.4	31.5	66.5	122.3
2002.....	14.2	17.7	13.6	11.2	11.6	17.2	24.9	40.1	55.7	132.3
2001.....	15.2	13.8	12.5	11.0	14.2	16.5	27.6	41.6	60.1	146.1
2000.....	12.7	12.3	11.7	10.3	11.1	16.7	29.5	38.4	60.3	133.1
1999.....	12.3	12.9	11.0	10.2	18.0	19.4	29.3	48.1	79.9	154.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Falls (W00-W19)										
All races 3/, both sexes										
2005.....	1.7	2.4	3.2	3.9	5.9	8.8	16.7	31.9	63.6	147.7
2004.....	1.8	2.6	3.2	4.1	5.8	8.9	16.1	30.3	61.8	143.3
2003.....	1.7	2.3	2.8	3.7	5.3	8.1	14.7	28.7	57.5	135.9
2002.....	1.8	2.3	2.8	3.4	4.9	7.9	13.9	27.1	54.0	130.4
2001.....	1.6	2.5	2.7	3.3	4.9	7.8	12.4	25.2	49.9	121.5
2000.....	1.6	2.0	2.6	3.3	4.6	6.2	12.0	22.1	44.5	112.6
1999.....	1.7	2.3	2.2	3.0	4.6	6.9	11.3	22.3	45.5	110.2
All races 3/, male										
2005.....	2.6	3.7	5.0	5.9	8.5	12.1	21.4	41.0	82.1	174.8
2004.....	2.9	4.1	4.9	6.2	8.1	12.4	21.6	38.4	78.7	174.4
2003.....	2.7	3.6	4.4	5.3	7.3	10.7	18.9	36.9	73.8	165.1
2002.....	2.9	3.6	4.3	5.2	7.1	10.7	18.9	35.1	67.9	158.9
2001.....	2.8	4.2	4.4	5.0	7.1	10.6	16.0	34.0	64.1	155.7
2000.....	2.7	3.4	4.2	5.0	6.9	8.6	15.7	29.4	57.3	145.4
1999.....	2.8	3.8	3.6	4.7	7.0	9.8	15.3	27.9	61.6	142.5
All races 3/, female										
2005.....	0.7	1.2	1.4	2.1	3.6	6.0	13.0	25.4	52.2	135.2
2004.....	0.7	1.1	1.5	2.0	3.6	5.8	11.7	24.5	51.4	129.2
2003.....	0.6	1.1	1.3	2.1	3.5	5.8	11.3	22.8	47.5	122.9
2002.....	0.7	1.0	1.4	1.8	3.0	5.5	9.9	21.4	45.5	118.1
2001.....	0.6	0.9	1.2	1.6	2.9	5.4	9.5	19.0	41.3	107.2
2000.....	0.6	0.8	1.2	1.8	2.6	4.2	9.1	17.1	36.9	99.2
1999.....	0.6	0.8	0.9	1.5	2.4	4.5	8.1	18.4	36.0	97.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Falls (W00-W19)										
White, both sexes										
2005 4/.....	1.8	2.5	3.2	4.0	6.0	9.1	17.6	33.4	66.6	156.7
2004 4/.....	1.9	2.6	3.2	4.0	5.8	9.2	16.7	32.0	65.3	152.6
2003 4/.....	1.7	2.3	2.7	3.7	5.4	8.2	15.2	30.1	60.0	144.4
2002.....	1.8	2.3	2.8	3.4	5.1	8.1	14.4	28.3	57.1	138.0
2001.....	1.7	2.5	2.8	3.2	4.9	8.0	12.8	26.3	52.4	128.0
2000.....	1.7	2.1	2.5	3.2	4.6	6.2	12.3	23.1	46.6	119.1
1999.....	1.7	2.2	2.1	2.9	4.3	6.9	11.4	23.0	47.7	115.9
White, male										
2005 4/.....	2.8	3.7	5.0	5.9	8.4	12.4	22.4	43.1	84.6	183.8
2004 4/.....	3.1	4.2	4.8	6.1	7.9	13.0	22.0	40.4	82.7	183.7
2003 4/.....	2.8	3.6	4.1	5.3	7.3	10.6	19.5	38.6	76.4	175.1
2002.....	3.0	3.7	4.2	5.1	7.2	10.8	19.3	36.4	71.2	166.4
2001.....	2.8	4.1	4.4	4.9	7.0	10.8	16.5	35.1	66.6	162.3
2000.....	2.7	3.5	3.9	4.8	6.9	8.4	16.2	30.3	59.7	152.8
1999.....	2.7	3.7	3.3	4.4	6.6	9.6	15.3	28.3	64.2	149.1
White, female										
2005 4/.....	0.8	1.3	1.4	2.1	3.7	6.2	13.6	26.3	55.3	144.2
2004 4/.....	0.7	1.1	1.6	2.1	3.8	5.8	12.3	25.9	54.6	138.6
2003 4/.....	0.6	1.0	1.3	2.1	3.6	6.1	11.6	24.0	49.9	130.8
2002.....	0.7	1.0	1.5	1.8	3.1	5.8	10.3	22.6	48.5	125.6
2001.....	0.5	0.9	1.2	1.6	2.9	5.6	9.9	20.0	43.9	113.6
2000.....	0.6	0.7	1.2	1.8	2.6	4.3	9.3	18.1	38.8	105.4
1999.....	0.6	0.8	0.9	1.5	2.2	4.6	8.3	19.3	37.9	102.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Falls (W00-W19)										
Black, both sexes										
2005 4/.....	1.0	2.4	3.9	4.1	5.7	7.2	10.9	20.5	32.5	52.7
2004 4/.....	1.7	2.6	3.5	5.1	6.3	7.9	12.8	16.8	25.5	46.5
2003 4/.....	1.5	3.0	3.6	4.3	5.5	7.6	10.6	13.4	28.2	44.1
2002.....	1.6	2.5	3.2	3.8	4.7	6.1	10.9	14.8	18.1	49.3
2001.....	2.0	2.6	3.3	3.9	4.9	7.9	9.7	16.3	21.9	44.5
2000.....	1.6	2.0	3.6	4.5	5.1	7.1	9.7	11.6	20.3	38.8
1999.....	1.7	3.2	3.6	4.5	6.7	9.0	9.5	13.4	23.1	45.3
Black, male										
2005 4/.....	1.7	4.4	6.4	6.5	9.8	10.9	14.5	23.7	52.7	63.2
2004 4/.....	2.6	4.1	6.0	8.8	11.8	9.7	20.7	23.7	38.0	71.5
2003 4/.....	2.6	4.8	5.8	6.5	8.1	12.8	12.2	18.3	35.4	56.3
2002.....	2.6	4.1	5.6	6.3	7.3	9.2	15.4	22.0	28.3	77.2
2001.....	3.3	4.9	6.0	6.4	7.7	11.9	13.5	24.2	36.1	68.1
2000.....	2.6	3.4	6.3	7.7	8.7	10.7	12.4	18.4	26.1	61.1
1999.....	3.3	5.6	6.0	7.6	10.2	14.8	14.6	20.4	31.4	58.6
Black, female										
2005 4/.....	*	*	1.8	2.1	*	4.4	8.4	18.7	21.9	48.4
2004 4/.....	*	*	*	*	*	6.7	7.5	12.7	18.9	36.6
2003 4/.....	*	1.4	1.7	2.5	3.5	3.7	9.5	10.6	24.4	39.3
2002.....	*	*	*	*	*	3.7	7.8	10.4	12.7	38.6
2001.....	*	*	*	*	*	4.9	7.2	11.6	14.7	35.8
2000.....	*	*	*	*	*	4.4	8.0	7.4	17.4	30.5
1999.....	*	*	*	*	4.0	4.6	6.1	9.2	19.0	40.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Accidental drowning and submersion (W65-W74)										
All races 3/, both sexes										
2005.....	1.3	1.2	1.1	0.9	0.9	0.9	1.2	1.4	1.4	1.8
2004.....	1.0	1.1	1.0	0.9	0.9	1.1	1.0	1.0	1.5	1.6
2003.....	1.1	1.0	1.0	1.0	0.9	1.0	1.2	1.3	1.3	1.8
2002.....	1.1	1.0	0.9	0.8	0.8	0.8	1.0	1.2	1.5	1.7
2001.....	1.1	1.0	0.8	0.8	0.9	0.8	0.9	1.1	1.4	1.7
2000.....	1.0	0.9	1.0	0.9	0.9	0.8	1.1	1.0	1.7	1.5
1999.....	1.0	1.0	1.0	0.9	0.9	0.9	1.1	1.2	1.4	1.6
All races 3/, male										
2005.....	2.0	2.0	1.7	1.5	1.5	1.4	1.8	2.1	2.5	3.9
2004.....	1.7	1.8	1.7	1.5	1.5	1.7	1.6	1.7	2.8	2.9
2003.....	1.8	1.6	1.6	1.6	1.5	1.6	1.9	1.8	2.3	3.3
2002.....	1.9	1.6	1.5	1.3	1.2	1.4	1.8	1.8	2.5	3.6
2001.....	1.7	1.8	1.3	1.2	1.4	1.4	1.6	1.8	2.4	3.5
2000.....	1.7	1.6	1.7	1.6	1.5	1.4	1.8	1.2	2.7	2.8
1999.....	1.7	1.7	1.8	1.5	1.5	1.5	1.7	2.1	2.7	2.3
All races 3/, female										
2005.....	0.5	0.5	0.4	0.3	0.3	0.5	0.7	0.8	0.7	0.9
2004.....	0.4	0.4	0.5	0.4	0.5	0.5	0.5	0.5	0.7	1.0
2003.....	0.3	0.3	0.4	0.3	0.4	0.6	0.6	0.8	0.6	1.1
2002.....	0.4	0.4	0.3	0.3	0.4	*	0.4	0.7	0.9	0.9
2001.....	0.4	0.3	0.3	0.4	0.4	*	*	0.7	0.8	1.0
2000.....	0.3	0.3	0.3	*	0.4	*	0.6	0.8	1.1	1.0
1999.....	0.3	0.3	0.3	0.4	0.4	0.4	0.7	0.6	0.7	1.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Accidental drowning and submersion (W65-W74)										
White, both sexes										
2005 4/.....	1.3	1.2	1.0	0.9	0.9	0.9	1.2	1.3	1.2	1.7
2004 4/.....	1.0	1.1	1.0	0.9	0.9	1.1	0.9	1.0	1.4	1.5
2003 4/.....	1.0	1.0	1.0	1.0	0.9	1.1	1.1	1.2	1.3	1.6
2002.....	1.2	0.9	0.8	0.8	0.8	0.8	0.9	1.1	1.4	1.6
2001.....	1.0	1.0	0.8	0.7	0.9	0.8	0.9	1.1	1.3	1.7
2000.....	1.0	0.9	0.9	0.9	0.9	0.8	1.1	0.9	1.8	1.5
1999.....	1.0	1.1	1.0	0.9	0.9	0.9	1.0	1.1	1.4	1.6
White, male										
2005 4/.....	2.0	2.0	1.6	1.5	1.4	1.3	1.7	2.1	2.3	3.9
2004 4/.....	1.7	1.8	1.6	1.4	1.4	1.8	1.4	1.6	2.8	2.8
2003 4/.....	1.8	1.5	1.5	1.6	1.4	1.7	1.9	1.8	2.4	3.3
2002.....	1.9	1.4	1.4	1.3	1.2	1.4	1.6	1.8	2.4	3.7
2001.....	1.6	1.7	1.3	1.1	1.3	1.4	1.6	1.8	2.3	3.5
2000.....	1.6	1.5	1.5	1.6	1.6	1.4	1.7	1.1	3.0	2.7
1999.....	1.7	1.8	1.7	1.4	1.4	1.4	1.5	1.9	2.7	2.4
White, female										
2005 4/.....	0.6	0.5	0.5	0.3	0.4	0.4	0.7	0.7	*	0.8
2004 4/.....	0.3	0.4	0.4	0.4	0.5	*	*	*	*	0.9
2003 4/.....	0.3	0.4	0.4	0.4	0.4	0.6	0.6	0.8	0.7	0.9
2002.....	0.4	0.4	0.3	0.3	0.4	*	*	0.6	0.8	0.7
2001.....	0.4	0.3	0.3	0.4	0.5	*	*	0.7	0.7	1.0
2000.....	0.3	0.3	0.3	*	*	*	0.6	0.7	1.0	1.0
1999.....	0.4	0.4	0.3	0.4	*	*	0.7	0.5	0.7	1.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Accidental exposure to smoke, fire and flames (X00-X09)										
All races 3/, both sexes										
2005.....	0.9	1.2	1.1	1.1	1.6	1.9	2.3	3.1	4.2	5.0
2004.....	0.7	1.1	1.2	1.2	1.6	1.8	2.0	2.9	4.4	5.6
2003.....	1.0	1.2	1.3	1.3	1.6	2.0	2.6	3.0	4.3	5.4
2002.....	0.9	1.0	1.3	1.2	1.4	1.8	2.4	2.8	4.2	4.8
2001.....	1.1	1.0	1.1	1.5	1.5	2.0	2.5	3.0	3.8	5.7
2000.....	1.0	1.1	1.3	1.4	1.7	2.0	2.4	3.5	4.3	6.2
1999.....	1.0	1.0	1.0	1.2	1.5	2.0	2.5	3.5	4.7	5.9
All races 3/, male										
2005.....	1.2	1.5	1.6	1.5	2.0	2.5	2.8	3.9	5.6	7.0
2004.....	1.0	1.5	1.8	1.6	2.1	2.6	2.3	3.5	5.9	7.2
2003.....	1.2	1.7	1.6	1.8	2.0	2.5	3.4	4.0	5.1	7.4
2002.....	1.3	1.4	1.6	1.5	2.0	2.6	3.2	3.5	6.1	7.0
2001.....	1.6	1.5	1.5	2.1	1.8	2.5	3.0	3.5	5.1	8.3
2000.....	1.2	1.5	1.8	1.9	2.2	2.4	3.2	4.5	5.9	9.4
1999.....	1.5	1.5	1.3	1.6	1.9	2.5	3.4	4.3	6.6	9.3
All races 3/, female										
2005.....	0.5	0.9	0.7	0.8	1.2	1.4	1.9	2.5	3.4	4.0
2004.....	0.5	0.8	0.7	0.9	1.3	1.2	1.6	2.5	3.5	4.9
2003.....	0.7	0.8	1.1	0.8	1.1	1.6	1.9	2.3	3.9	4.5
2002.....	0.6	0.6	0.9	0.9	0.8	1.1	1.8	2.4	3.0	3.9
2001.....	0.7	0.6	0.7	1.0	1.2	1.5	2.1	2.7	3.1	4.6
2000.....	0.7	0.7	0.7	0.9	1.2	1.6	1.8	2.8	3.4	4.9
1999.....	0.5	0.5	0.7	0.9	1.2	1.6	1.8	3.0	3.6	4.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Accidental exposure to smoke, fire and flames (X00-X09)										
White, both sexes										
2005 4/.....	0.8	1.2	1.0	0.8	1.4	1.7	2.1	2.7	3.8	4.2
2004 4/.....	0.7	1.0	1.1	1.1	1.4	1.6	1.6	2.5	3.9	4.9
2003 4/.....	1.0	1.2	1.2	1.2	1.4	1.8	2.2	2.6	3.7	4.7
2002.....	0.9	0.9	1.1	1.1	1.1	1.6	2.1	2.5	3.7	4.0
2001.....	1.0	0.9	1.0	1.3	1.3	1.7	2.1	2.6	3.2	4.8
2000.....	0.9	0.9	1.2	1.1	1.4	1.7	2.0	2.8	3.5	5.1
1999.....	0.9	0.9	0.9	1.1	1.2	1.6	2.0	2.8	3.9	4.7
White, male										
2005 4/.....	1.1	1.5	1.4	1.1	1.7	2.1	2.6	3.4	5.1	6.0
2004 4/.....	0.9	1.2	1.6	1.3	1.9	2.2	1.9	2.9	5.4	6.5
2003 4/.....	1.3	1.5	1.5	1.6	1.9	2.2	3.0	3.2	4.2	6.7
2002.....	1.3	1.3	1.4	1.4	1.6	2.2	2.7	3.1	5.8	5.8
2001.....	1.4	1.2	1.2	1.8	1.5	2.1	2.4	2.8	4.4	7.0
2000.....	1.1	1.3	1.7	1.6	1.8	2.0	2.7	3.6	4.8	8.2
1999.....	1.4	1.3	1.2	1.4	1.7	2.0	2.6	3.3	5.8	8.0
White, female										
2005 4/.....	0.6	0.9	0.7	0.6	1.2	1.3	1.8	2.2	2.9	3.4
2004 4/.....	0.4	0.7	0.7	0.9	1.0	1.1	1.3	2.2	2.9	4.2
2003 4/.....	0.7	0.8	0.9	0.8	1.0	1.5	1.6	2.2	3.4	3.8
2002.....	0.6	0.6	0.9	0.8	0.6	1.0	1.6	2.1	2.5	3.2
2001.....	0.6	0.5	0.7	0.8	1.1	1.3	1.9	2.4	2.5	3.8
2000.....	0.6	0.6	0.7	0.7	1.1	1.4	1.5	2.3	2.7	3.9
1999.....	0.5	0.5	0.6	0.9	0.8	1.3	1.5	2.5	2.8	3.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Accidental exposure to smoke, fire and flames (X00-X09)										
Black, both sexes										
2005 4/.....	1.2	1.7	2.2	3.7	3.2	4.4	4.2	8.0	11.2	14.4
2004 4/.....	1.4	2.6	2.2	2.8	4.2	4.0	5.9	8.0	12.1	14.9
2003 4/.....	1.4	2.3	2.9	2.8	2.9	4.4	6.4	8.7	12.4	14.9
2002.....	1.1	1.6	2.8	2.6	4.2	4.0	6.3	7.2	10.7	15.7
2001.....	2.1	2.3	2.5	3.7	3.3	5.2	7.3	8.5	11.6	17.8
2000.....	2.0	2.2	2.6	3.9	4.5	5.0	7.1	11.4	16.1	20.3
1999.....	1.7	2.0	2.3	2.6	4.3	5.7	8.3	12.0	15.3	21.7
Black, male										
2005 4/.....	1.9	2.3	3.8	5.4	5.3	6.7	*	12.1	15.5	22.3
2004 4/.....	2.0	3.9	3.7	4.2	4.7	7.3	7.8	12.3	15.1	*
2003 4/.....	1.5	3.3	3.2	4.6	3.9	6.4	8.9	15.5	17.0	*
2002.....	1.8	2.1	4.0	3.4	6.3	6.7	9.3	9.2	*	25.7
2001.....	2.9	3.7	4.2	5.0	4.9	7.4	11.1	12.1	*	27.2
2000.....	2.7	3.2	4.0	5.8	7.0	7.5	10.4	16.5	21.9	27.7
1999.....	2.5	3.0	3.3	4.1	4.2	7.8	13.2	17.1	20.1	29.3
Black, female										
2005 4/.....	*	*	*	2.3	*	*	*	5.6	8.9	11.3
2004 4/.....	*	1.5	*	*	3.8	*	4.6	5.4	10.6	13.4
2003 4/.....	*	1.4	2.6	*	*	*	4.7	*	10.1	13.4
2002.....	*	*	*	*	*	*	*	6.0	9.2	11.9
2001.....	1.3	*	*	2.7	*	*	4.7	6.4	10.2	14.3
2000.....	1.3	*	*	*	*	*	4.9	8.3	13.2	17.6
1999.....	*	*	*	*	4.3	4.2	5.0	8.9	13.0	18.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Accidental poisoning and exposure to noxious substances (X40-X49)										
All races 3/, both sexes										
2005.....	17.1	18.2	14.4	8.3	4.4	2.6	2.0	2.3	2.6	3.5
2004.....	16.3	16.7	12.0	6.7	3.7	2.4	2.1	2.4	2.5	3.4
2003.....	15.5	15.4	10.9	6.1	3.4	2.3	2.0	2.0	2.5	3.5
2002.....	14.8	13.8	9.2	5.0	3.0	2.3	1.9	2.1	2.7	3.9
2001.....	12.0	10.9	7.0	3.7	2.4	1.7	1.5	2.1	2.4	3.5
2000.....	11.2	9.9	6.1	3.3	2.2	1.5	1.5	1.6	2.6	4.3
1999.....	11.1	9.8	5.4	3.4	2.0	1.7	1.6	1.9	2.6	3.4
All races 3/, male										
2005.....	22.1	24.0	19.1	10.9	5.7	3.0	2.3	3.2	2.8	4.6
2004.....	20.9	21.6	16.3	8.4	4.2	2.7	2.5	2.6	2.9	4.2
2003.....	20.7	20.8	14.7	7.6	4.5	3.0	2.2	2.3	3.5	4.0
2002.....	20.3	18.7	12.8	6.8	3.6	2.8	2.1	2.3	3.2	5.1
2001.....	16.8	15.7	9.9	5.0	3.0	2.0	2.1	2.7	2.8	3.8
2000.....	16.4	14.7	8.9	4.0	2.7	1.8	1.7	1.6	2.9	6.2
1999.....	16.6	15.1	7.9	4.7	3.0	2.2	1.9	2.1	2.9	4.4
All races 3/, female										
2005.....	12.0	12.6	10.0	5.8	3.2	2.3	1.7	1.7	2.4	3.0
2004.....	11.7	11.9	7.9	5.1	3.2	2.1	1.8	2.2	2.3	3.0
2003.....	10.3	10.2	7.3	4.6	2.5	1.8	1.8	1.7	2.0	3.3
2002.....	9.4	9.1	5.8	3.4	2.4	1.8	1.7	1.9	2.3	3.4
2001.....	7.3	6.2	4.1	2.6	1.9	1.4	1.0	1.6	2.2	3.3
2000.....	6.0	5.3	3.4	2.7	1.8	1.3	1.3	1.6	2.4	3.6
1999.....	5.7	4.7	3.1	2.2	1.2	1.2	1.4	1.7	2.4	3.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Accidental poisoning and exposure to noxious substances (X40-X49)										
White, both sexes										
2005 4/.....	18.0	18.7	13.8	7.6	4.1	2.4	1.9	2.3	2.5	3.6
2004 4/.....	17.3	17.1	11.7	6.2	3.5	2.3	2.1	2.3	2.4	3.4
2003 4/.....	16.3	15.7	10.6	5.8	3.4	2.0	2.0	2.0	2.7	3.5
2002.....	15.4	13.7	8.7	4.6	2.7	2.2	1.9	2.1	2.6	3.9
2001.....	12.1	10.5	6.1	3.4	2.1	1.6	1.4	2.0	2.3	3.3
2000.....	11.0	9.4	5.3	3.0	2.1	1.5	1.4	1.7	2.6	4.1
1999.....	10.8	9.1	4.8	3.1	1.7	1.5	1.4	1.9	2.6	3.3
White, male										
2005 4/.....	23.3	24.2	17.9	9.4	4.9	2.3	2.0	3.1	2.6	4.8
2004 4/.....	22.2	22.0	15.4	7.3	3.6	2.3	2.4	2.4	2.7	4.3
2003 4/.....	21.8	20.9	13.8	7.0	4.2	2.3	2.2	2.3	3.6	3.7
2002.....	21.0	18.3	11.7	5.8	2.9	2.6	2.0	2.3	3.0	5.2
2001.....	17.1	14.9	8.5	4.3	2.3	1.9	2.0	2.4	2.7	3.7
2000.....	16.2	13.7	7.6	3.3	2.4	1.6	1.6	1.7	2.9	5.9
1999.....	16.2	13.8	6.6	4.0	2.3	2.0	1.7	2.1	3.0	4.5
White, female										
2005 4/.....	12.6	13.3	9.9	5.9	3.4	2.4	1.8	1.7	2.5	3.0
2004 4/.....	12.5	12.2	8.2	5.2	3.4	2.2	1.8	2.3	2.2	3.0
2003 4/.....	10.8	10.5	7.3	4.7	2.6	1.7	1.8	1.8	2.1	3.4
2002.....	9.8	9.0	5.8	3.4	2.5	1.9	1.8	1.9	2.5	3.4
2001.....	7.1	6.1	3.8	2.6	1.9	1.4	1.0	1.7	2.0	3.1
2000.....	5.8	5.1	3.1	2.7	1.8	1.4	1.3	1.7	2.5	3.4
1999.....	5.4	4.5	3.0	2.3	1.2	1.1	1.2	1.8	2.4	2.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Accidental poisoning and exposure to noxious substances (X40-X49)

Black, both sexes

2005 4/.....	17.0	20.7	23.1	16.3	8.1	5.0	3.7	*	*	*
2004 4/.....	14.6	19.6	17.9	12.6	6.0	4.1	3.1	*	*	*
2003 4/.....	15.2	18.8	16.9	9.6	4.8	5.5	*	*	*	*
2002.....	15.7	19.4	16.1	10.3	6.0	3.0	*	*	*	*
2001.....	15.2	17.7	15.4	7.4	5.4	2.6	*	*	*	6.4
2000.....	15.3	16.5	14.2	6.8	3.9	2.7	*	*	*	8.1
1999.....	16.1	17.9	12.3	6.5	5.5	2.9	3.2	*	*	*

Black, male

2005 4/.....	22.0	30.3	34.6	26.9	15.1	9.3	6.6	*	*	*
2004 4/.....	19.1	25.8	28.7	20.5	10.1	7.3	*	*	*	*
2003 4/.....	20.5	26.7	25.7	15.8	8.7	10.3	*	*	*	*
2002.....	21.6	27.2	25.6	17.7	11.3	5.5	*	*	*	*
2001.....	20.3	28.0	24.8	12.7	9.6	*	*	*	*	*
2000.....	22.8	26.3	23.1	11.5	7.0	*	*	*	*	*
1999.....	24.2	29.7	20.9	11.9	10.0	*	*	*	*	*

Black, female

2005 4/.....	12.6	12.4	13.4	7.6	*	*	*	*	*	*
2004 4/.....	10.6	14.2	8.9	6.1	*	*	*	*	*	*
2003 4/.....	10.5	11.9	9.5	4.5	*	*	*	*	*	*
2002.....	10.5	12.5	8.1	4.3	*	*	*	*	*	*
2001.....	10.7	8.9	7.5	3.1	*	*	*	*	*	*
2000.....	8.6	8.0	6.6	3.0	*	*	*	*	*	*
1999.....	9.1	7.8	4.9	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)										
All races 3/, both sexes										
2005.....	3.5	4.4	4.9	5.0	6.8	8.4	12.3	20.7	38.9	95.1
2004.....	3.2	3.9	4.5	4.5	5.7	7.0	11.0	19.2	37.4	95.5
2003.....	3.4	3.8	4.0	4.7	5.5	8.1	11.2	19.8	38.1	102.5
2002.....	3.3	3.5	4.0	4.8	5.7	8.6	11.1	21.6	38.8	105.6
2001.....	3.2	3.9	4.5	4.6	6.3	7.8	12.3	22.5	42.2	115.0
2000.....	3.4	3.9	4.1	5.0	6.3	7.8	12.2	22.0	41.4	117.1
1999.....	3.7	3.8	4.4	5.1	6.3	8.9	13.3	23.6	46.9	128.9
All races 3/, male										
2005.....	5.6	7.1	7.8	7.4	10.2	11.5	16.6	26.8	49.0	107.1
2004.....	5.0	6.3	7.0	6.7	8.4	9.8	14.3	25.1	46.1	109.8
2003.....	5.5	6.0	6.1	7.1	8.7	11.4	15.1	26.5	47.6	120.2
2002.....	5.4	5.8	6.2	7.4	8.7	12.0	15.0	28.6	47.2	118.6
2001.....	5.3	6.3	6.8	6.7	8.9	10.9	15.9	29.3	52.7	134.0
2000.....	5.6	6.5	6.7	7.7	9.1	10.8	15.6	28.1	51.6	135.2
1999.....	6.1	6.0	7.2	7.8	9.1	12.0	18.0	29.9	59.3	152.7
All races 3/, female										
2005.....	1.5	1.8	2.1	2.7	3.6	5.7	8.7	16.3	32.6	89.6
2004.....	1.4	1.5	2.1	2.3	3.2	4.6	8.4	15.0	32.1	89.1
2003.....	1.4	1.6	2.1	2.5	2.7	5.3	8.2	15.0	32.3	94.7
2002.....	1.3	1.3	1.9	2.4	3.0	5.7	8.1	16.6	33.8	100.0
2001.....	1.2	1.6	2.2	2.5	3.8	5.2	9.5	17.7	35.9	107.0
2000.....	1.3	1.4	1.7	2.4	3.8	5.2	9.6	17.8	35.4	109.8
1999.....	1.4	1.7	1.7	2.5	3.7	6.3	9.6	19.3	39.6	119.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)										
White, both sexes										
2005 4/.....	3.5	4.3	4.7	4.8	6.5	8.2	11.7	20.3	38.8	97.4
2004 4/.....	3.4	4.0	4.5	4.4	5.7	6.8	10.8	19.1	38.0	98.8
2003 4/.....	3.5	3.7	3.9	4.6	5.4	8.0	11.1	20.0	38.4	106.3
2002.....	3.4	3.5	4.0	4.6	5.6	7.9	11.0	21.4	39.1	108.8
2001.....	3.2	3.8	4.4	4.5	5.9	7.7	11.9	22.3	42.5	117.7
2000.....	3.4	3.7	4.0	4.7	6.2	7.4	11.8	21.7	41.9	120.5
1999.....	3.7	3.6	4.2	4.7	5.8	8.6	12.7	22.8	46.8	132.4
White, male										
2005 4/.....	5.6	6.8	7.4	7.0	9.5	11.4	15.5	25.9	48.6	110.0
2004 4/.....	5.2	6.4	7.0	6.5	8.5	9.5	14.2	24.8	46.6	112.9
2003 4/.....	5.7	5.8	5.9	6.9	8.2	11.4	14.7	26.6	47.3	123.9
2002.....	5.6	5.9	6.1	6.9	8.5	10.9	14.9	28.1	47.2	121.2
2001.....	5.2	6.2	6.6	6.5	8.5	10.7	15.3	29.1	51.9	137.0
2000.....	5.6	6.1	6.4	7.3	8.7	10.1	15.1	27.1	51.2	137.6
1999.....	6.0	5.5	6.8	7.1	8.5	11.3	17.0	28.7	59.0	155.2
White, female										
2005 4/.....	1.4	1.7	1.9	2.6	3.7	5.3	8.6	16.1	32.6	91.6
2004 4/.....	1.5	1.6	2.2	2.4	3.1	4.4	8.1	14.9	32.6	92.5
2003 4/.....	1.3	1.6	1.9	2.4	2.7	5.0	8.1	15.3	32.9	98.5
2002.....	1.2	1.1	1.9	2.4	2.9	5.3	7.8	16.5	34.2	103.4
2001.....	1.1	1.5	2.3	2.6	3.5	5.1	9.1	17.4	36.8	109.7
2000.....	1.3	1.4	1.6	2.3	3.8	5.1	9.1	17.9	36.4	113.6
1999.....	1.5	1.6	1.7	2.4	3.3	6.2	9.3	18.7	39.6	123.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)

Black, both sexes

2005 4/.....	3.9	6.4	7.6	7.6	10.1	12.4	19.3	31.9	46.6	86.1
2004 4/.....	2.9	4.0	5.5	6.3	6.9	10.1	13.7	25.9	37.4	70.2
2003 4/.....	3.7	4.7	6.2	6.7	8.0	11.2	14.7	19.6	39.6	73.2
2002.....	3.3	4.3	5.1	7.3	7.3	15.9	14.3	27.8	39.0	83.1
2001.....	3.9	5.4	5.8	5.9	10.4	10.4	18.7	27.3	43.9	96.1
2000.....	3.9	5.7	5.6	7.9	8.6	12.3	18.1	30.2	39.2	90.1
1999.....	4.7	6.6	6.6	9.3	10.9	13.9	22.9	36.2	53.7	105.5

Black, male

2005 4/.....	6.0	10.8	12.1	12.1	18.9	15.4	29.1	48.3	63.2	97.6
2004 4/.....	4.9	6.6	8.7	10.6	9.9	14.5	18.8	35.0	53.8	87.4
2003 4/.....	5.6	7.9	9.2	10.4	13.9	15.3	22.3	29.2	56.8	91.8
2002.....	4.3	6.4	8.6	12.7	11.3	23.7	18.7	40.9	53.5	106.1
2001.....	6.6	8.6	9.5	9.9	14.6	15.8	25.2	35.3	72.2	114.6
2000.....	6.5	10.1	10.0	13.8	13.9	18.2	23.1	46.8	64.1	124.5
1999.....	8.1	11.1	11.0	15.4	16.1	20.8	34.9	50.3	69.8	153.5

Black, female

2005 4/.....	2.0	2.6	3.8	3.9	3.3	10.2	12.6	22.1	37.8	81.4
2004 4/.....	*	1.7	2.7	2.7	4.5	6.8	10.3	20.5	28.7	63.4
2003 4/.....	2.0	2.0	3.7	3.7	3.5	8.2	9.5	13.8	30.6	66.0
2002.....	2.4	2.5	2.1	2.9	4.2	10.2	11.4	20.0	31.4	74.3
2001.....	1.6	2.7	2.7	2.7	7.0	6.4	14.3	22.5	29.4	89.3
2000.....	1.5	1.8	*	3.0	4.4	7.9	14.9	20.2	26.7	77.3
1999.....	1.6	2.7	2.8	4.3	6.8	8.9	14.9	27.7	45.8	87.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)										
All races 3/, both sexes										
2005.....	16.0	16.7	16.1	14.3	13.3	11.8	13.5	15.4	18.7	16.9
2004.....	16.0	16.9	16.2	14.2	13.2	11.4	13.5	16.0	16.7	16.4
2003.....	15.9	16.0	15.7	14.7	12.6	12.0	13.6	15.6	17.7	16.9
2002.....	16.1	16.3	15.1	14.6	12.3	12.5	14.6	16.6	19.3	18.0
2001 5/.....	15.2	15.7	14.6	14.0	12.0	12.7	13.9	16.4	18.9	17.5
2000.....	15.4	14.9	13.9	12.7	11.4	11.3	13.8	16.5	19.3	19.6
1999.....	14.3	14.3	13.4	12.8	11.4	12.4	14.5	17.4	19.2	19.3
All races 3/, male										
2005.....	24.6	25.4	25.0	22.4	22.0	20.6	25.4	31.9	41.4	45.0
2004.....	24.2	25.2	24.4	22.3	21.7	20.2	25.4	32.4	38.3	45.0
2003.....	24.5	24.4	24.4	23.1	21.1	21.2	26.0	31.8	40.1	47.8
2002.....	24.7	24.9	23.8	23.5	20.4	22.0	27.9	34.3	44.0	50.7
2001 5/.....	23.6	24.3	22.4	22.3	19.5	22.7	26.7	33.8	44.2	51.1
2000.....	23.9	23.3	21.4	20.1	18.7	19.8	26.1	34.9	44.8	57.5
1999.....	22.1	22.3	20.6	20.3	19.1	22.1	27.7	35.6	44.1	57.1
All races 3/, female										
2005.....	7.4	8.2	7.7	6.6	5.5	4.2	3.8	3.5	4.7	4.0
2004.....	7.9	8.9	8.4	6.6	5.5	3.7	3.9	4.3	3.5	3.6
2003.....	7.4	7.9	7.5	6.7	4.9	4.0	3.6	4.0	3.9	3.3
2002.....	7.7	8.0	6.7	6.2	5.0	4.3	3.9	4.1	4.4	3.8
2001 5/.....	6.9	7.4	7.0	6.2	5.2	4.1	3.7	4.3	3.8	3.4
2000.....	7.0	6.7	6.8	5.8	4.9	4.0	4.1	3.8	4.2	4.2
1999.....	6.7	6.7	6.5	5.8	4.4	4.2	4.1	4.8	4.6	4.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)										
White, both sexes										
2005 4/.....	18.0	18.9	18.1	15.9	14.7	13.0	14.7	16.7	20.0	17.9
2004 4/.....	18.1	19.0	18.3	15.7	14.4	12.4	14.5	17.3	17.7	17.5
2003 4/.....	17.9	18.1	17.7	16.2	13.8	13.0	14.7	16.7	18.9	18.1
2002.....	18.2	18.4	16.8	16.1	13.6	13.7	15.7	17.6	20.9	19.1
2001 5/.....	16.9	17.6	16.2	15.3	13.1	13.8	14.9	17.6	20.0	18.6
2000.....	17.0	16.5	15.3	13.9	12.6	12.3	14.7	17.6	20.4	21.0
1999.....	16.0	15.9	14.8	14.0	12.3	13.3	15.6	18.6	20.4	20.7
White, male										
2005 4/.....	27.3	28.5	27.7	24.8	24.1	22.6	27.6	34.2	44.3	48.2
2004 4/.....	27.0	27.8	27.3	24.1	23.6	21.9	27.1	34.8	40.4	48.4
2003 4/.....	27.3	27.3	27.2	25.4	22.9	22.9	27.9	34.0	42.9	51.4
2002.....	27.6	27.6	26.4	25.7	22.5	23.8	30.1	36.3	47.2	53.9
2001 5/.....	26.1	27.1	24.7	24.3	21.0	24.5	28.3	36.0	46.8	55.0
2000.....	26.2	25.6	23.2	21.8	20.4	21.3	27.6	37.1	47.7	61.6
1999.....	24.5	24.4	22.3	21.9	20.5	23.6	29.5	37.9	46.8	61.5
White, female										
2005 4/.....	8.5	9.4	8.7	7.4	6.0	4.4	4.1	3.7	4.8	3.9
2004 4/.....	9.1	10.2	9.5	7.6	5.8	4.1	4.1	4.5	3.6	3.6
2003 4/.....	8.4	9.0	8.3	7.4	5.5	4.3	3.9	4.2	4.1	3.3
2002.....	8.7	9.3	7.4	6.9	5.3	4.8	3.9	4.2	4.7	4.0
2001 5/.....	7.8	8.1	7.9	6.8	5.8	4.3	4.0	4.6	3.8	3.4
2000.....	7.9	7.5	7.5	6.4	5.5	4.4	4.4	3.9	4.2	4.5
1999.....	7.5	7.4	7.4	6.4	4.8	4.4	4.5	5.0	4.8	4.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Intentional self-harm (suicide) by discharge of firearms (X72-X74)										
All races 3/, both sexes										
2005.....	7.0	8.0	8.3	8.1	8.2	8.2	9.9	11.9	13.6	11.2
2004.....	7.1	8.1	8.0	7.8	8.3	8.0	9.8	12.2	12.3	11.1
2003.....	7.2	7.8	8.3	8.4	8.2	8.5	10.1	11.9	13.1	11.9
2002.....	7.4	7.8	7.9	8.6	8.1	8.8	10.7	12.4	14.3	12.0
2001.....	7.0	7.5	8.0	8.4	8.0	9.0	10.3	12.4	14.1	12.2
2000.....	7.4	7.4	7.8	7.4	7.7	8.3	10.3	12.4	14.0	13.2
1999.....	6.7	7.4	7.2	8.1	7.5	8.7	10.9	12.9	13.5	12.6
All races 3/, male										
2005.....	11.8	13.7	14.6	14.1	15.3	15.5	20.2	26.6	33.3	33.8
2004.....	11.7	13.4	13.6	13.8	14.9	15.5	20.0	26.7	30.7	34.4
2003.....	12.1	13.2	14.5	14.6	15.3	16.5	20.7	26.3	32.4	36.6
2002.....	12.6	13.3	14.0	15.4	14.8	17.0	22.5	27.8	35.9	37.8
2001.....	12.0	12.8	13.8	14.9	14.2	17.3	21.4	27.7	35.5	38.9
2000.....	12.6	12.8	13.3	13.1	13.9	15.9	21.3	28.1	35.4	42.9
1999.....	11.2	12.6	12.4	14.0	14.0	16.9	22.4	29.1	34.6	41.6
All races 3/, female										
2005.....	2.3	2.5	2.4	2.3	1.8	1.7	1.6	1.3	1.3	0.8
2004.....	2.6	2.8	2.7	2.1	2.3	1.4	1.7	1.7	1.0	*
2003.....	2.3	2.6	2.3	2.5	1.7	1.6	1.6	1.6	1.3	0.9
2002.....	2.3	2.4	2.1	2.3	2.0	1.8	1.3	1.5	1.2	0.8
2001.....	2.2	2.3	2.4	2.4	2.3	1.9	1.4	1.7	1.3	1.0
2000.....	2.4	2.2	2.5	2.1	2.1	1.8	1.6	1.4	1.4	1.1
1999.....	2.2	2.4	2.1	2.6	1.6	1.7	1.8	1.6	1.2	0.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Intentional self-harm (suicide) by discharge of firearms (X72-X74)										
White, both sexes										
2005 4/.....	8.0	9.2	9.5	9.2	9.2	9.2	10.9	13.1	14.7	12.2
2004 4/.....	8.1	9.2	9.2	8.7	9.3	8.8	10.8	13.3	13.3	12.1
2003 4/.....	8.3	9.1	9.4	9.5	9.1	9.5	11.1	12.9	14.1	12.9
2002.....	8.5	8.9	8.9	9.7	9.0	9.9	11.7	13.3	15.6	12.9
2001.....	7.9	8.5	9.0	9.4	8.9	10.0	11.1	13.4	15.0	13.2
2000.....	8.4	8.3	8.7	8.2	8.5	9.2	11.1	13.4	15.2	14.3
1999.....	7.5	8.3	8.0	8.9	8.2	9.5	11.8	13.9	14.4	13.6
White, male										
2005 4/.....	13.3	15.5	16.3	15.9	17.0	17.4	22.0	28.7	36.0	36.8
2004 4/.....	13.1	15.1	15.5	15.1	16.6	16.9	21.6	28.7	32.9	37.5
2003 4/.....	13.8	15.1	16.2	16.4	16.8	18.3	22.5	28.3	34.8	39.9
2002.....	14.3	15.0	15.5	16.9	16.3	18.8	24.4	29.6	38.8	40.5
2001.....	13.3	14.5	15.4	16.3	15.7	19.0	22.9	29.7	37.8	42.1
2000.....	14.0	14.2	14.7	14.2	15.3	17.3	22.7	30.4	38.1	46.5
1999.....	12.5	14.0	13.7	15.1	15.2	18.3	24.0	31.2	36.8	44.9
White, female										
2005 4/.....	2.7	2.9	2.8	2.7	2.0	1.9	1.7	1.5	1.4	0.9
2004 4/.....	3.0	3.3	3.1	2.5	2.6	1.7	1.9	2.0	1.1	*
2003 4/.....	2.7	3.1	2.6	2.9	2.0	1.8	1.8	1.8	1.4	0.9
2002.....	2.7	2.8	2.4	2.7	2.3	2.1	1.4	1.6	1.3	0.9
2001.....	2.4	2.6	2.8	2.7	2.7	2.0	1.5	1.8	1.3	1.1
2000.....	2.7	2.5	2.8	2.4	2.3	2.1	1.8	1.5	1.5	1.2
1999.....	2.5	2.7	2.5	3.0	1.8	1.8	2.1	1.8	1.3	1.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)										
All races 3/, both sexes										
2005.....	9.0	8.7	7.8	6.2	5.1	3.7	3.6	3.5	5.2	5.7
2004.....	8.9	8.9	8.2	6.5	4.9	3.4	3.6	3.9	4.4	5.4
2003.....	8.7	8.2	7.5	6.2	4.5	3.4	3.5	3.7	4.6	5.1
2002.....	8.7	8.5	7.2	5.9	4.2	3.7	3.9	4.2	5.0	6.0
2001 5/.....	8.2	8.2	6.6	5.6	4.0	3.7	3.6	4.0	4.8	5.3
2000.....	7.9	7.4	6.2	5.2	3.7	3.0	3.5	4.2	5.3	6.5
1999.....	7.7	6.9	6.2	4.7	3.9	3.8	3.6	4.5	5.7	6.7
All races 3/, male										
2005.....	12.8	11.8	10.4	8.3	6.7	5.1	5.2	5.3	8.1	11.2
2004.....	12.5	11.8	10.8	8.6	6.8	4.7	5.4	5.7	7.6	10.7
2003.....	12.4	11.2	9.9	8.5	5.8	4.7	5.4	5.5	7.7	11.1
2002.....	12.1	11.6	9.9	8.2	5.7	5.0	5.5	6.5	8.1	12.9
2001 5/.....	11.6	11.5	8.6	7.5	5.3	5.4	5.4	6.1	8.7	12.2
2000.....	11.3	10.6	8.1	6.9	4.8	3.9	4.8	6.8	9.4	14.6
1999.....	10.8	9.6	8.1	6.4	5.1	5.2	5.3	6.6	9.5	15.5
All races 3/, female										
2005.....	5.1	5.7	5.3	4.3	3.6	2.4	2.2	2.2	3.4	3.1
2004.....	5.3	6.0	5.7	4.5	3.2	2.2	2.2	2.5	2.5	3.0
2003.....	5.1	5.3	5.1	4.1	3.2	2.4	2.0	2.4	2.6	2.4
2002.....	5.4	5.6	4.6	3.9	2.9	2.5	2.6	2.6	3.1	3.0
2001 5/.....	4.8	5.1	4.6	3.8	2.8	2.2	2.3	2.6	2.4	2.4
2000.....	4.6	4.4	4.3	3.7	2.8	2.1	2.5	2.4	2.8	3.2
1999.....	4.5	4.2	4.4	3.2	2.8	2.5	2.3	3.1	3.4	3.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)										
White, both sexes										
2005 4/.....	9.9	9.7	8.6	6.8	5.5	3.8	3.8	3.6	5.3	5.6
2004 4/.....	10.0	9.7	9.1	7.0	5.1	3.6	3.7	4.0	4.4	5.5
2003 4/.....	9.6	9.1	8.3	6.7	4.7	3.5	3.6	3.8	4.8	5.2
2002.....	9.7	9.6	7.9	6.4	4.6	3.8	4.0	4.3	5.3	6.2
2001 5/.....	9.1	9.1	7.2	5.9	4.2	3.8	3.7	4.2	5.0	5.5
2000.....	8.7	8.2	6.6	5.7	4.1	3.1	3.6	4.2	5.3	6.7
1999.....	8.4	7.6	6.7	5.1	4.1	3.8	3.8	4.7	5.9	7.1
White, male										
2005 4/.....	14.0	13.0	11.4	8.9	7.2	5.3	5.6	5.5	8.4	11.4
2004 4/.....	13.9	12.6	11.8	9.0	7.1	4.9	5.4	6.1	7.5	10.9
2003 4/.....	13.5	12.2	11.0	9.0	6.1	4.7	5.4	5.6	8.1	11.5
2002.....	13.4	12.6	10.9	8.7	6.2	5.0	5.8	6.7	8.4	13.4
2001 5/.....	12.7	12.6	9.3	7.9	5.3	5.5	5.3	6.3	9.0	12.9
2000.....	12.2	11.4	8.5	7.6	5.2	4.0	4.9	6.7	9.6	15.2
1999.....	11.9	10.4	8.6	6.7	5.2	5.3	5.5	6.7	10.0	16.6
White, female										
2005 4/.....	5.8	6.5	5.9	4.8	3.9	2.5	2.3	2.2	3.4	3.0
2004 4/.....	6.0	6.8	6.4	5.1	3.3	2.5	2.2	2.5	2.5	3.0
2003 4/.....	5.7	5.9	5.7	4.5	3.5	2.5	2.2	2.4	2.8	2.4
2002.....	6.0	6.5	5.0	4.2	3.0	2.7	2.5	2.6	3.4	3.1
2001 5/.....	5.4	5.6	5.1	4.1	3.1	2.3	2.5	2.8	2.5	2.3
2000.....	5.1	5.0	4.8	4.0	3.1	2.3	2.6	2.5	2.7	3.3
1999.....	4.9	4.7	4.9	3.5	3.0	2.6	2.4	3.3	3.5	3.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)										
All races 3/, both sexes										
2005.....	6.7	5.4	4.2	2.9	2.7	2.3	2.4	2.2	2.1	2.1
2004.....	6.3	5.3	4.2	3.2	2.8	2.6	2.2	2.2	2.2	2.1
2003.....	6.4	5.6	4.2	3.1	2.4	2.3	2.4	2.5	2.4	2.2
2002.....	6.5	5.4	4.0	3.5	2.7	2.4	2.2	2.3	2.4	2.1
2001 5/.....	8.6	7.0	5.5	4.4	3.5	3.3	2.5	2.5	2.5	2.4
2000.....	6.6	5.3	4.0	3.4	2.6	2.4	2.4	2.3	2.6	2.4
1999.....	6.6	5.1	4.0	3.2	2.8	2.7	2.4	2.5	2.4	2.4
All races 3/, male										
2005.....	10.0	8.3	6.7	4.5	4.2	3.1	3.5	2.8	2.4	2.7
2004.....	9.0	7.8	6.7	4.9	4.1	3.8	3.3	2.6	2.7	2.7
2003.....	9.5	8.3	6.4	4.6	3.7	3.4	3.4	3.5	3.0	3.2
2002.....	9.6	8.0	6.0	5.3	4.2	3.4	3.2	3.3	2.9	3.0
2001 5/.....	12.6	10.6	8.3	6.7	5.0	4.6	3.3	3.2	3.0	3.2
2000.....	9.7	7.8	5.9	5.2	3.9	3.3	3.4	2.8	3.8	3.3
1999.....	9.5	7.7	5.8	5.1	4.3	4.2	3.1	3.4	3.4	3.4
All races 3/, female										
2005.....	3.4	2.6	1.8	1.5	1.4	1.6	1.5	1.8	2.0	1.9
2004.....	3.6	3.0	1.9	1.6	1.6	1.6	1.3	1.9	2.0	1.9
2003.....	3.4	3.0	2.1	1.7	1.3	1.4	1.6	1.8	2.1	1.8
2002.....	3.5	2.9	2.2	1.8	1.4	1.5	1.4	1.6	2.1	1.7
2001 5/.....	4.7	3.6	2.7	2.3	2.2	2.1	1.9	1.9	2.2	2.0
2000.....	3.6	2.8	2.1	1.7	1.5	1.7	1.6	2.0	1.9	2.0
1999.....	3.6	2.5	2.2	1.4	1.5	1.5	1.9	1.9	1.9	2.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)										
White, both sexes										
2005 4/.....	4.4	3.8	2.9	2.3	2.1	1.8	1.9	2.0	1.9	1.9
2004 4/.....	4.3	3.8	3.0	2.4	2.3	2.2	1.8	1.8	2.0	1.9
2003 4/.....	4.2	3.9	3.1	2.4	2.0	1.9	1.9	2.2	2.1	2.0
2002.....	4.5	3.8	2.9	2.9	2.2	1.9	1.9	1.8	2.1	1.8
2001 5/.....	6.8	5.6	4.4	3.8	3.0	2.7	2.2	2.1	2.2	2.2
2000.....	4.4	3.7	2.9	2.7	2.0	1.8	2.0	2.1	2.0	2.2
1999.....	4.5	3.6	3.0	2.6	2.3	2.3	2.0	2.0	2.1	2.1
White, male										
2005 4/.....	6.2	5.6	4.5	3.3	3.3	2.2	2.7	2.2	2.0	1.9
2004 4/.....	5.8	5.2	4.6	3.6	3.2	3.0	2.6	2.0	2.2	2.3
2003 4/.....	5.8	5.4	4.5	3.3	2.9	2.7	2.6	2.9	2.5	3.0
2002.....	6.5	5.1	4.0	4.1	3.2	2.7	2.6	2.4	2.3	2.7
2001 5/.....	9.8	8.2	6.5	5.5	4.2	3.7	2.6	2.7	2.6	2.8
2000.....	6.4	5.1	4.0	4.0	2.8	2.2	2.6	2.4	2.5	2.8
1999.....	6.3	5.3	4.3	3.8	3.3	3.4	2.5	2.6	2.9	2.7
White, female										
2005 4/.....	2.6	2.0	1.4	1.3	1.1	1.4	1.3	1.8	1.8	1.9
2004 4/.....	2.9	2.3	1.4	1.3	1.4	1.6	1.1	1.6	1.9	1.8
2003 4/.....	2.7	2.4	1.7	1.6	1.0	1.2	1.3	1.6	1.8	1.6
2002.....	2.5	2.5	1.8	1.7	1.3	1.3	1.4	1.4	1.9	1.5
2001 5/.....	3.8	3.0	2.4	2.1	1.9	1.9	1.8	1.7	1.9	2.0
2000.....	2.5	2.3	1.8	1.4	1.3	1.4	1.5	1.8	1.7	1.9
1999.....	2.6	1.9	1.7	1.3	1.3	1.4	1.7	1.6	1.6	1.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)

Black, both sexes

2005 4/.....	21.6	17.2	13.9	8.3	7.8	7.1	7.1	5.8	5.6	*
2004 4/.....	20.2	16.8	13.4	9.4	7.9	5.7	6.4	7.1	5.2	*
2003 4/.....	21.4	17.7	12.7	9.1	6.8	5.7	6.9	6.6	6.6	*
2002.....	20.6	17.6	13.0	8.9	7.5	5.8	5.6	8.2	6.0	*
2001 5/.....	21.7	18.1	13.6	9.9	7.5	7.7	6.5	6.9	6.2	*
2000.....	21.8	16.8	12.4	9.1	7.4	8.2	6.8	5.3	9.6	*
1999.....	21.2	16.2	11.2	8.8	7.4	6.6	6.0	8.2	6.1	*

Black, male

2005 4/.....	36.7	29.3	24.1	15.3	12.9	12.6	12.7	11.2	*	*
2004 4/.....	32.9	27.8	23.8	16.5	13.1	11.6	11.3	10.5	*	*
2003 4/.....	36.8	30.0	21.8	17.3	11.4	10.8	11.5	12.3	*	*
2002.....	32.4	30.8	22.5	16.8	13.5	10.0	11.3	15.2	*	*
2001 5/.....	34.2	30.0	23.1	17.4	12.0	13.2	11.8	11.6	*	*
2000.....	34.1	28.7	22.4	16.0	13.0	15.1	12.4	*	19.4	*
1999.....	34.1	27.3	18.5	17.0	12.7	13.0	10.2	13.8	*	*

Black, female

2005 4/.....	8.3	6.7	5.2	2.5	3.8	*	*	*	*	*
2004 4/.....	8.9	7.2	4.6	3.6	3.8	*	*	*	*	*
2003 4/.....	7.9	7.0	5.0	2.3	3.2	*	*	*	*	*
2002.....	10.1	6.1	4.9	*	*	*	*	*	*	*
2001 5/.....	10.6	7.7	5.6	3.8	4.0	*	*	*	*	*
2000.....	10.9	6.5	3.8	3.5	*	*	*	*	*	*
1999.....	9.8	6.7	5.0	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)										
All races 3/, both sexes										
2005.....	4.1	3.0	2.1	1.4	1.3	1.1	1.1	0.8	0.6	0.5
2004.....	3.8	2.9	2.2	1.7	1.5	1.2	0.8	0.8	0.7	0.6
2003.....	3.8	3.1	2.2	1.6	1.2	0.9	0.9	0.7	0.6	0.5
2002.....	3.9	2.8	2.2	1.8	1.3	1.0	0.8	1.0	0.8	*
2001.....	3.7	2.7	2.1	1.5	1.4	1.2	0.9	0.8	0.7	*
2000.....	3.8	2.9	2.1	1.7	1.2	1.0	1.0	0.7	0.5	0.7
1999.....	3.7	2.8	2.0	1.6	1.6	1.1	1.0	0.8	0.7	0.6
All races 3/, male										
2005.....	6.6	4.7	3.4	2.2	2.0	1.4	1.7	1.0	*	*
2004.....	5.9	4.4	3.5	2.7	2.2	1.8	1.4	1.0	*	*
2003.....	5.9	5.0	3.5	2.4	1.8	1.4	1.4	0.9	*	*
2002.....	6.2	4.2	3.4	2.7	1.9	1.4	1.0	1.4	1.0	*
2001.....	5.6	4.2	3.2	2.3	2.0	1.8	1.1	1.2	*	*
2000.....	5.8	4.5	3.3	2.6	1.8	1.3	1.3	0.8	*	*
1999.....	5.5	4.4	3.0	2.6	2.5	1.5	1.4	1.1	*	*
All races 3/, female										
2005.....	1.7	1.4	0.8	0.6	0.6	0.8	0.6	0.7	0.6	*
2004.....	1.6	1.4	0.9	0.7	0.9	0.7	*	0.7	0.8	*
2003.....	1.7	1.4	1.0	0.8	0.6	0.4	0.5	0.6	0.7	*
2002.....	1.7	1.5	1.1	0.9	0.7	0.6	0.6	0.6	0.7	*
2001.....	1.7	1.1	1.0	0.7	0.8	0.7	0.7	0.5	0.7	*
2000.....	1.7	1.3	1.0	0.9	0.7	0.8	0.7	0.7	*	*
1999.....	1.8	1.3	1.1	0.6	0.7	0.7	0.7	0.6	0.6	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Assault (homicide) by discharge of firearms (*U01.4,X93-X95)

White, both sexes

2005 4/.....	2.6	2.1	1.5	1.1	1.0	0.9	0.9	0.8	0.5	*
2004 4/.....	2.5	2.0	1.5	1.3	1.2	1.1	0.7	0.7	0.7	0.5
2003 4/.....	2.4	2.2	1.7	1.2	1.0	0.7	0.8	0.6	0.6	0.5
2002.....	2.6	1.9	1.6	1.5	0.9	0.8	0.7	0.8	0.8	*
2001.....	2.6	1.9	1.5	1.2	1.2	1.0	0.7	0.7	0.7	*
2000.....	2.5	2.0	1.5	1.4	1.0	0.8	0.8	0.7	*	0.6
1999.....	2.5	1.9	1.5	1.2	1.2	1.0	0.9	0.7	0.7	0.6

White, male

2005 4/.....	3.9	3.0	2.3	1.6	1.6	1.1	1.2	0.9	*	*
2004 4/.....	3.5	2.9	2.3	1.9	1.7	1.4	1.0	0.9	*	*
2003 4/.....	3.4	3.1	2.5	1.8	1.4	1.1	1.1	0.7	*	*
2002.....	4.0	2.6	2.4	2.1	1.4	1.0	0.8	1.2	*	*
2001.....	3.7	2.8	2.3	1.6	1.7	1.3	0.8	1.0	*	*
2000.....	3.7	3.0	2.2	2.1	1.4	1.0	0.9	0.7	*	*
1999.....	3.6	2.8	2.3	1.9	1.8	1.3	1.1	0.7	*	*

White, female

2005 4/.....	1.4	1.1	0.7	0.6	0.5	0.7	0.6	0.8	*	*
2004 4/.....	1.5	1.1	0.8	0.7	0.8	0.8	*	0.6	0.9	*
2003 4/.....	1.3	1.2	0.9	0.8	0.5	*	0.5	0.5	0.7	*
2002.....	1.3	1.3	0.9	0.9	0.5	0.5	0.6	0.6	0.7	*
2001.....	1.4	0.9	0.8	0.8	0.7	0.8	0.7	0.5	0.7	*
2000.....	1.3	1.1	0.9	0.8	0.7	0.7	0.7	0.7	*	*
1999.....	1.4	1.0	0.7	0.6	0.7	0.7	0.7	0.6	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Assault (homicide) by discharge of firearms (*U01.4,X93-X95)

Black, both sexes

2005 4/.....	14.4	9.9	6.9	3.7	3.3	2.8	3.3	*	*	*
2004 4/.....	12.8	9.4	7.2	5.2	4.2	2.6	*	*	*	*
2003 4/.....	13.7	10.2	6.8	4.6	3.1	2.4	*	*	*	*
2002.....	13.1	9.6	7.1	4.2	4.0	2.8	*	*	*	*
2001.....	11.6	8.4	6.5	4.3	3.0	2.9	*	*	*	*
2000.....	12.6	9.4	7.0	4.2	2.6	2.8	2.7	*	*	*
1999.....	12.2	9.7	5.9	4.5	4.2	2.2	*	*	*	*

Black, male

2005 4/.....	26.1	17.9	12.6	7.3	5.6	5.2	7.0	*	*	*
2004 4/.....	23.4	16.5	13.7	9.6	7.1	5.6	*	*	*	*
2003 4/.....	24.3	19.1	11.9	9.1	5.2	*	*	*	*	*
2002.....	22.7	17.4	12.3	8.3	6.5	5.2	*	*	*	*
2001.....	20.3	15.3	11.8	8.6	5.1	6.4	*	*	*	*
2000.....	21.6	16.7	13.1	7.1	5.4	5.2	*	*	*	*
1999.....	20.5	17.3	9.5	9.3	8.3	*	*	*	*	*

Black, female

2005 4/.....	4.2	3.0	2.1	*	*	*	*	*	*	*
2004 4/.....	3.5	3.2	1.7	*	*	*	*	*	*	*
2003 4/.....	4.4	2.5	2.5	*	*	*	*	*	*	*
2002.....	4.6	2.8	2.8	*	*	*	*	*	*	*
2001.....	3.9	2.4	1.9	*	*	*	*	*	*	*
2000.....	4.7	3.2	*	*	*	*	*	*	*	*
1999.....	4.8	3.2	2.8	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)

All races 3/, both sexes

2005.....	2.5	2.4	2.1	1.6	1.4	1.2	1.4	1.4	1.5	1.7
2004.....	2.5	2.5	2.1	1.5	1.3	1.4	1.4	1.4	1.6	1.5
2003.....	2.6	2.5	1.9	1.5	1.3	1.5	1.5	1.8	1.8	1.7
2002.....	2.6	2.6	1.8	1.7	1.5	1.4	1.4	1.3	1.6	1.7
2001 5/.....	4.9	4.4	3.4	2.9	2.1	2.1	1.6	1.6	1.8	2.0
2000.....	2.9	2.4	1.8	1.6	1.4	1.4	1.4	1.6	2.1	1.7
1999.....	2.9	2.3	1.9	1.6	1.3	1.6	1.4	1.7	1.7	1.8

All races 3/, male

2005.....	3.4	3.6	3.2	2.3	2.1	1.7	1.9	1.8	1.8	2.1
2004.....	3.1	3.4	3.2	2.2	2.0	2.0	1.8	1.6	2.2	1.7
2003.....	3.6	3.3	2.9	2.2	1.9	2.1	2.0	2.7	2.6	2.5
2002.....	3.4	3.8	2.6	2.6	2.3	2.0	2.2	1.8	1.9	2.3
2001 5/.....	7.0	6.4	5.0	4.4	3.0	2.8	2.2	2.0	2.1	2.7
2000.....	3.9	3.3	2.6	2.6	2.0	2.0	2.1	2.0	3.0	2.5
1999.....	4.0	3.3	2.8	2.5	1.8	2.6	1.7	2.2	2.6	2.6

All races 3/, female

2005.....	1.7	1.2	1.0	0.9	0.8	0.8	1.0	1.1	1.3	1.5
2004.....	1.9	1.6	1.0	0.8	0.8	0.8	1.0	1.2	1.2	1.5
2003.....	1.6	1.6	1.0	0.9	0.7	1.0	1.0	1.1	1.3	1.4
2002.....	1.8	1.4	1.1	1.0	0.7	0.9	0.8	0.9	1.4	1.4
2001 5/.....	2.9	2.5	1.8	1.5	1.4	1.4	1.2	1.4	1.6	1.7
2000.....	1.8	1.5	1.0	0.8	0.8	0.9	0.9	1.3	1.5	1.4
1999.....	1.8	1.3	1.1	0.9	0.8	0.8	1.2	1.3	1.2	1.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)

White, both sexes										
2005 4/.....	1.8	1.7	1.4	1.2	1.1	0.9	1.1	1.1	1.3	1.5
2004 4/.....	1.8	1.8	1.5	1.2	1.0	1.2	1.1	1.0	1.3	1.4
2003 4/.....	1.9	1.7	1.4	1.2	1.0	1.2	1.1	1.5	1.5	1.6
2002.....	1.9	1.8	1.3	1.4	1.3	1.2	1.2	1.0	1.3	1.4
2001 5/.....	4.2	3.7	2.9	2.6	1.8	1.7	1.4	1.4	1.5	1.8
2000.....	1.9	1.7	1.4	1.2	1.0	1.0	1.1	1.4	1.6	1.5
1999.....	2.0	1.7	1.5	1.3	1.0	1.4	1.2	1.4	1.4	1.5
White, male										
2005 4/.....	2.4	2.5	2.2	1.7	1.6	1.1	1.5	1.3	1.6	1.6
2004 4/.....	2.3	2.3	2.3	1.7	1.4	1.6	1.5	1.1	1.8	1.5
2003 4/.....	2.4	2.3	2.0	1.5	1.5	1.6	1.5	2.2	2.2	2.3
2002.....	2.6	2.5	1.6	1.9	1.9	1.7	1.8	1.2	1.5	2.0
2001 5/.....	6.0	5.3	4.2	3.9	2.5	2.4	1.8	1.7	2.0	2.3
2000.....	2.7	2.1	1.8	1.9	1.5	1.3	1.6	1.7	2.2	2.1
1999.....	2.8	2.4	2.1	1.9	1.5	2.1	1.3	1.8	2.2	1.9
White, female										
2005 4/.....	1.2	0.9	0.7	0.8	0.6	0.7	0.8	1.0	1.2	1.5
2004 4/.....	1.4	1.2	0.7	0.7	0.6	0.8	0.7	1.0	1.0	1.3
2003 4/.....	1.3	1.2	0.9	0.8	0.5	0.9	0.8	1.0	1.1	1.2
2002.....	1.2	1.2	0.9	0.9	0.7	0.7	0.7	0.8	1.2	1.2
2001 5/.....	2.3	2.0	1.6	1.4	1.1	1.1	1.1	1.2	1.2	1.6
2000.....	1.2	1.2	0.9	0.6	0.6	0.7	0.7	1.1	1.2	1.3
1999.....	1.2	0.9	1.0	0.8	0.6	0.8	1.0	1.0	1.0	1.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)

Black, both sexes

2005 4/.....	7.2	7.3	7.0	4.6	4.5	4.3	3.7	4.5	*	*
2004 4/.....	7.3	7.4	6.2	4.2	3.7	3.1	4.3	5.2	*	*
2003 4/.....	7.7	7.5	5.9	4.5	3.6	3.3	4.4	4.8	5.1	*
2002.....	7.5	8.0	5.9	4.6	3.5	3.1	3.8	5.7	*	*
2001 5/.....	10.1	9.7	7.2	5.6	4.6	4.8	4.0	4.7	*	*
2000.....	9.2	7.4	5.3	5.0	4.7	5.4	4.1	4.1	6.8	*
1999.....	9.0	6.5	5.3	4.3	3.2	4.4	3.4	5.2	*	*

Black, male

2005 4/.....	10.6	11.5	11.5	8.0	7.3	7.4	*	*	*	*
2004 4/.....	9.5	11.3	10.1	6.9	6.0	6.1	*	*	*	*
2003 4/.....	12.5	10.9	9.9	8.1	6.2	6.2	6.9	9.1	*	*
2002.....	9.7	13.4	10.3	8.5	6.9	*	7.3	10.1	*	*
2001 5/.....	13.9	14.7	11.3	8.8	6.9	6.9	7.1	*	*	*
2000.....	12.6	12.0	9.3	8.9	7.7	9.9	6.7	*	*	*
1999.....	13.6	10.0	9.0	7.6	4.5	8.6	*	*	*	*

Black, female

2005 4/.....	4.2	3.7	3.1	*	*	*	*	*	*	*
2004 4/.....	5.4	4.0	2.9	*	*	*	*	*	*	*
2003 4/.....	3.5	4.5	2.5	*	*	*	*	*	*	*
2002.....	5.5	3.4	2.1	*	*	*	*	*	*	*
2001 5/.....	6.8	5.3	3.7	3.1	*	*	*	*	*	*
2000.....	6.2	3.3	2.0	*	*	*	*	*	*	*
1999.....	5.0	3.5	2.2	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)										
All races 3/, both sexes										
2005.....	3.1	3.2	2.9	1.6	1.1	0.9	0.7	0.8	0.9	1.5
2004.....	3.4	3.6	2.6	1.7	1.0	0.9	0.8	0.7	1.3	1.6
2003.....	3.9	3.6	2.6	1.6	1.0	0.8	0.7	0.6	0.9	1.1
2002.....	3.4	3.3	2.3	1.4	1.0	0.8	0.7	0.6	1.0	1.4
2001.....	3.2	3.0	2.0	1.1	0.7	0.6	0.5	0.7	1.1	1.0
2000.....	3.1	2.7	1.7	1.1	0.7	0.6	0.6	0.6	0.7	1.3
1999.....	2.9	2.7	1.6	1.0	0.8	0.8	0.5	0.7	0.9	1.4
All races 3/, male										
2005.....	3.6	3.8	3.5	2.1	1.1	1.1	1.0	1.0	*	1.6
2004.....	4.1	4.5	3.3	2.0	1.1	1.2	0.9	0.8	1.3	1.9
2003.....	4.9	4.6	3.4	2.0	1.2	0.8	0.7	0.8	1.1	1.5
2002.....	4.3	4.1	2.9	1.8	1.3	1.0	1.0	0.8	1.1	2.2
2001.....	4.4	4.2	2.5	1.5	0.8	0.9	0.6	0.9	1.5	1.5
2000.....	4.2	3.8	2.3	1.6	0.9	0.9	0.8	1.0	1.1	1.7
1999.....	4.0	3.7	2.1	1.5	1.1	1.1	0.6	1.0	1.3	2.1
All races 3/, female										
2005.....	2.6	2.7	2.4	1.2	1.1	0.8	0.5	0.6	0.9	1.4
2004.....	2.7	2.6	2.0	1.3	0.8	0.7	0.6	0.6	1.2	1.6
2003.....	2.9	2.6	1.8	1.1	0.9	0.7	0.6	*	0.8	0.9
2002.....	2.6	2.6	1.7	1.0	0.7	0.7	0.5	0.5	0.9	1.1
2001.....	2.0	1.8	1.5	0.6	0.5	*	*	0.6	0.8	0.8
2000.....	1.9	1.7	1.1	0.7	0.5	*	*	*	*	1.2
1999.....	1.9	1.7	1.0	0.6	0.6	0.5	*	0.5	0.7	1.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)										
White, both sexes										
2005 4/.....	3.1	3.3	2.9	1.6	1.1	0.9	0.7	0.7	0.9	1.5
2004 4/.....	3.5	3.7	2.6	1.7	1.0	0.8	0.8	0.6	1.3	1.6
2003 4/.....	4.0	3.6	2.5	1.5	0.9	0.7	0.6	0.6	0.9	1.2
2002.....	3.5	3.3	2.2	1.4	0.9	0.8	0.7	0.6	0.9	1.4
2001.....	3.2	2.9	1.9	1.1	0.6	0.6	0.5	0.7	1.0	1.0
2000.....	2.9	2.6	1.6	1.0	0.7	0.5	0.5	0.6	0.8	1.3
1999.....	2.8	2.5	1.4	1.0	0.8	0.8	0.4	0.7	0.9	1.4
White, male										
2005 4/.....	3.5	3.7	3.3	2.0	0.9	1.1	1.0	0.8	*	1.7
2004 4/.....	4.1	4.5	3.1	2.0	1.0	1.0	0.9	*	1.4	1.9
2003 4/.....	5.0	4.5	3.2	1.8	1.0	0.7	0.7	0.9	1.1	1.5
2002.....	4.3	4.0	2.5	1.8	1.2	0.9	0.8	*	1.1	2.3
2001.....	4.2	4.0	2.3	1.4	0.7	0.8	0.6	0.9	1.5	*
2000.....	4.0	3.5	2.1	1.3	0.9	0.8	0.7	1.0	*	*
1999.....	3.8	3.2	1.8	1.3	1.0	1.1	*	1.0	1.3	2.1
White, female										
2005 4/.....	2.6	2.9	2.5	1.3	1.2	0.8	0.5	0.6	0.8	1.3
2004 4/.....	2.9	2.9	2.0	1.4	0.9	0.6	0.6	0.6	1.2	1.5
2003 4/.....	3.1	2.8	1.9	1.2	0.8	0.7	0.5	*	0.8	1.0
2002.....	2.6	2.6	1.9	1.1	0.7	0.7	0.5	*	0.7	1.1
2001.....	2.1	1.9	1.6	0.7	0.6	*	*	0.6	0.7	0.8
2000.....	1.9	1.7	1.1	0.7	0.5	*	*	*	*	1.1
1999.....	1.7	1.8	1.0	0.7	0.7	0.5	*	*	0.8	1.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)										
All races 3/, both sexes										
2005.....	3.0	3.2	2.8	1.6	1.0	0.9	0.7	0.7	0.9	1.4
2004.....	3.3	3.5	2.5	1.6	0.9	0.8	0.7	0.6	1.2	1.6
2003.....	3.8	3.5	2.5	1.5	1.0	0.7	0.6	0.6	0.8	1.1
2002.....	3.3	3.2	2.2	1.3	0.9	0.8	0.6	0.6	0.9	1.4
2001.....	3.1	2.9	1.9	1.0	0.6	0.5	0.4	0.6	1.0	1.0
2000.....	3.0	2.7	1.6	1.1	0.7	0.5	0.5	0.5	0.7	1.3
1999.....	2.8	2.6	1.5	1.0	0.7	0.7	0.4	0.6	0.8	1.3
All races 3/, male										
2005.....	3.5	3.7	3.3	2.0	1.0	1.1	0.9	0.9	*	1.5
2004.....	3.9	4.5	3.1	1.9	1.0	1.0	0.8	0.7	1.1	1.8
2003.....	4.8	4.5	3.3	1.8	1.1	0.8	0.6	0.7	*	1.4
2002.....	4.1	4.0	2.8	1.7	1.3	0.9	0.8	0.7	1.0	2.2
2001.....	4.2	4.1	2.4	1.4	0.7	0.8	0.6	0.7	1.3	*
2000.....	4.1	3.7	2.1	1.5	0.9	0.7	0.7	0.8	*	*
1999.....	3.8	3.5	2.0	1.4	0.9	0.9	*	0.9	*	1.8
All races 3/, female										
2005.....	2.6	2.7	2.4	1.1	1.0	0.7	0.5	0.6	0.9	1.4
2004.....	2.6	2.6	2.0	1.3	0.8	0.6	0.6	0.6	1.2	1.5
2003.....	2.8	2.5	1.8	1.1	0.8	0.7	0.6	*	0.8	0.9
2002.....	2.6	2.5	1.7	0.9	0.7	0.6	0.4	0.5	0.8	1.1
2001.....	2.0	1.7	1.5	0.6	0.5	*	*	0.6	0.7	0.8
2000.....	1.9	1.6	1.1	0.7	0.4	*	*	*	*	1.2
1999.....	1.8	1.7	1.0	0.6	0.5	0.5	*	*	0.7	1.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)										
White, both sexes										
2005 4/.....	3.0	3.3	2.8	1.6	1.0	0.9	0.6	0.7	0.8	1.4
2004 4/.....	3.4	3.6	2.5	1.6	0.9	0.7	0.7	0.6	1.2	1.6
2003 4/.....	3.9	3.5	2.5	1.4	0.9	0.7	0.5	0.5	0.8	1.2
2002.....	3.4	3.2	2.1	1.3	0.9	0.7	0.6	0.5	0.8	1.4
2001.....	3.1	2.9	1.8	1.0	0.6	0.5	0.4	0.6	0.9	1.0
2000.....	2.8	2.5	1.5	0.9	0.6	0.5	0.5	0.5	0.7	1.2
1999.....	2.6	2.4	1.3	0.9	0.7	0.7	0.4	0.6	0.8	1.3
White, male										
2005 4/.....	3.4	3.7	3.1	1.9	0.8	1.0	0.8	0.7	*	1.6
2004 4/.....	4.0	4.4	3.0	1.9	0.9	0.8	0.8	*	1.1	1.8
2003 4/.....	4.8	4.4	3.1	1.7	0.9	0.6	*	0.7	*	*
2002.....	4.2	3.9	2.4	1.7	1.2	0.8	0.7	*	*	2.3
2001.....	4.1	3.9	2.2	1.3	0.6	0.7	*	0.7	1.4	*
2000.....	3.8	3.5	2.0	1.2	0.8	0.6	0.6	0.8	*	*
1999.....	3.6	3.1	1.7	1.2	0.9	0.9	*	0.8	*	*
White, female										
2005 4/.....	2.6	2.9	2.5	1.2	1.1	0.8	0.5	0.6	0.8	1.3
2004 4/.....	2.8	2.8	2.0	1.4	0.8	0.5	0.6	0.6	1.2	1.5
2003 4/.....	3.0	2.7	1.9	1.2	0.8	0.7	0.5	*	0.8	1.0
2002.....	2.6	2.5	1.9	1.0	0.7	0.6	0.5	*	0.7	1.1
2001.....	2.1	1.8	1.5	0.7	0.6	*	*	0.6	*	0.8
2000.....	1.8	1.6	1.1	0.7	0.4	*	*	*	*	1.1
1999.....	1.7	1.7	1.0	0.7	0.6	0.5	*	*	0.8	1.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Complications of medical and surgical care (Y40-Y84,Y88)										
All races 3/, both sexes										
2005.....	0.4	0.5	0.8	1.0	1.4	2.0	3.3	4.7	7.2	9.1
2004.....	0.4	0.6	0.8	1.2	1.6	2.6	3.5	5.0	7.6	10.6
2003.....	0.5	0.7	0.7	1.1	1.7	2.2	3.7	5.3	7.3	11.2
2002.....	0.4	0.5	0.8	1.1	1.6	2.3	3.9	5.1	7.7	11.3
2001.....	0.4	0.7	0.8	1.3	1.9	2.6	3.6	5.9	8.3	12.8
2000.....	0.4	0.6	0.9	1.2	2.0	2.8	4.3	5.8	8.0	13.6
1999.....	0.4	0.5	0.7	1.1	1.7	2.4	4.0	5.5	8.4	12.7
All races 3/, male										
2005.....	0.4	0.5	0.8	1.1	1.7	2.3	3.7	4.6	8.9	10.8
2004.....	0.4	0.6	0.8	1.5	1.7	2.5	4.3	5.8	9.6	11.9
2003.....	0.5	0.7	0.7	1.3	1.6	2.3	4.2	5.6	8.8	12.9
2002.....	0.4	0.6	0.7	1.2	1.5	2.5	3.8	5.4	7.7	12.3
2001.....	0.4	0.7	0.8	1.3	2.0	2.6	3.8	6.3	9.9	14.5
2000.....	0.4	0.6	1.0	1.2	2.0	2.8	5.3	7.0	10.4	16.0
1999.....	0.4	0.4	0.7	1.2	1.7	2.3	4.5	6.4	10.1	16.5
All races 3/, female										
2005.....	0.4	0.6	0.9	1.0	1.2	1.8	3.0	4.8	6.2	8.3
2004.....	0.5	0.6	0.8	1.0	1.5	2.7	2.9	4.3	6.3	9.9
2003.....	0.4	0.6	0.6	0.9	1.7	2.1	3.3	5.1	6.4	10.5
2002.....	0.4	0.5	0.9	1.0	1.7	2.2	3.9	5.0	7.7	10.9
2001.....	0.5	0.6	0.8	1.2	1.7	2.6	3.5	5.7	7.3	12.1
2000.....	0.4	0.7	0.9	1.1	2.0	2.7	3.6	4.9	6.7	12.7
1999.....	0.4	0.5	0.7	0.9	1.6	2.5	3.6	4.8	7.4	11.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Complications of medical and surgical care (Y40-Y84,Y88)										
White, both sexes										
2005 4/.....	0.4	0.5	0.7	0.9	1.3	1.8	3.0	4.4	7.3	9.2
2004 4/.....	0.4	0.6	0.8	1.1	1.5	2.4	3.4	4.8	7.3	10.4
2003 4/.....	0.4	0.6	0.6	1.0	1.6	2.1	3.6	5.1	7.4	11.0
2002.....	0.4	0.5	0.7	1.0	1.5	2.2	3.8	5.0	7.6	11.2
2001.....	0.4	0.6	0.7	1.1	1.6	2.4	3.2	5.8	8.2	12.1
2000.....	0.3	0.6	0.9	1.0	1.8	2.6	4.1	5.4	8.0	13.1
1999.....	0.3	0.4	0.6	0.9	1.5	2.3	3.8	5.2	8.0	12.1
White, male										
2005 4/.....	0.4	0.4	0.7	0.9	1.5	2.3	3.4	4.3	8.8	11.2
2004 4/.....	0.3	0.6	0.8	1.3	1.7	2.5	4.2	5.8	9.5	11.8
2003 4/.....	0.6	0.6	0.5	1.2	1.6	2.3	4.1	5.5	9.1	12.8
2002.....	0.4	0.5	0.6	1.0	1.5	2.2	3.7	5.3	7.8	11.8
2001.....	0.3	0.6	0.6	1.1	1.7	2.5	3.2	6.2	10.0	13.6
2000.....	0.4	0.5	1.0	1.1	1.8	2.5	5.0	6.8	10.6	15.8
1999.....	0.3	0.3	0.6	1.0	1.6	2.3	4.2	6.2	9.7	15.6
White, female										
2005 4/.....	0.4	0.5	0.7	1.0	1.1	1.5	2.6	4.5	6.3	8.3
2004 4/.....	0.4	0.5	0.7	0.9	1.4	2.4	2.7	4.0	6.0	9.8
2003 4/.....	0.3	0.5	0.6	0.8	1.6	2.0	3.1	4.7	6.3	10.2
2002.....	0.3	0.4	0.7	0.9	1.5	2.1	3.9	4.8	7.5	10.9
2001.....	0.4	0.5	0.7	1.1	1.6	2.4	3.2	5.6	7.0	11.5
2000.....	0.3	0.6	0.9	0.9	1.8	2.7	3.3	4.5	6.4	12.0
1999.....	0.4	0.5	0.6	0.7	1.5	2.4	3.4	4.5	7.1	10.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Complications of medical and surgical care (Y40-Y84,Y88)

Black, both sexes

2005 4/.....	*	1.0	2.1	2.2	2.9	4.5	6.4	9.4	8.5	9.1
2004 4/.....	0.9	1.0	1.4	2.7	2.5	4.5	5.5	7.6	11.9	13.8
2003 4/.....	*	1.3	1.6	2.5	3.0	3.6	5.8	9.5	7.6	15.2
2002.....	1.0	1.2	2.1	2.6	2.8	4.0	5.2	7.6	9.4	14.0
2001.....	1.0	1.6	1.5	2.7	4.2	4.6	8.5	8.0	11.1	23.3
2000.....	*	1.1	1.7	2.6	4.7	4.6	7.9	10.5	10.4	21.0
1999.....	0.8	0.9	1.8	2.6	2.6	3.5	7.3	9.5	12.4	20.1

Black, male

2005 4/.....	*	*	2.2	2.6	3.8	*	*	9.4	*	*
2004 4/.....	*	*	*	3.4	*	*	*	*	*	*
2003 4/.....	*	*	2.1	2.8	*	*	*	*	*	*
2002.....	*	*	*	*	*	5.0	*	*	*	*
2001.....	*	1.7	*	3.4	5.1	5.3	11.1	*	*	29.5
2000.....	*	*	*	*	4.6	6.2	9.4	10.4	*	*
1999.....	*	*	*	*	*	*	9.2	10.4	*	23.4

Black, female

2005 4/.....	*	*	2.1	*	*	4.8	6.7	9.3	7.4	9.0
2004 4/.....	*	*	1.7	*	*	5.4	5.1	7.5	11.0	13.4
2003 4/.....	*	*	*	*	3.0	3.8	6.4	10.6	7.7	15.4
2002.....	*	*	2.5	2.4	3.1	*	4.9	8.2	9.9	11.5
2001.....	*	*	*	*	3.5	4.1	6.7	8.0	11.0	21.0
2000.....	*	*	*	3.1	4.7	*	6.9	10.5	10.6	21.9
1999.....	*	*	*	*	*	*	6.1	8.9	10.8	18.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K86.0,R78.0,X45,X65,Y15) 6/										
All races 3/, both sexes										
2005.....	10.1	15.7	19.7	19.7	19.3	16.3	13.7	10.1	8.1	4.6
2004.....	10.0	15.5	19.5	18.9	18.5	17.0	14.1	10.7	7.3	4.3
2003.....	10.6	16.0	18.9	18.3	18.8	16.8	13.2	10.3	7.6	4.1
2002.....	10.6	15.6	18.2	18.2	18.6	16.5	14.4	10.9	7.2	4.4
2001.....	10.6	16.5	17.9	17.6	18.9	17.3	13.7	11.0	7.5	5.0
2000.....	11.1	15.9	16.7	18.2	19.3	17.2	14.3	11.2	7.9	5.4
1999.....	10.9	16.0	16.9	18.1	19.3	17.6	14.1	11.9	8.5	5.5
All races 3/, male										
2005.....	14.4	22.9	30.9	31.7	31.3	27.0	22.6	18.4	14.9	9.3
2004.....	14.6	22.7	30.6	30.0	30.4	28.1	24.5	19.4	13.8	9.0
2003.....	15.2	24.5	29.9	29.1	30.6	27.6	22.8	18.0	15.0	7.8
2002.....	15.3	23.7	29.2	29.2	31.0	27.9	23.8	18.7	13.4	9.7
2001.....	15.3	25.4	28.0	28.7	31.0	28.5	23.8	19.4	13.8	11.6
2000.....	16.4	25.0	26.9	28.8	31.4	28.9	25.0	21.1	14.9	11.9
1999.....	16.6	25.2	27.4	29.7	31.5	29.9	24.5	21.2	16.6	11.8
All races 3/, female										
2005.....	5.8	8.6	8.9	8.4	8.4	7.1	6.5	4.0	3.8	2.5
2004.....	5.3	8.4	8.8	8.3	7.8	7.3	5.8	4.4	3.3	2.3
2003.....	6.0	7.7	8.3	8.1	8.1	7.5	5.4	4.9	3.1	2.5
2002.....	6.1	7.7	7.7	7.9	7.2	6.7	6.9	5.3	3.4	2.1
2001.....	6.0	7.9	8.3	7.2	8.0	7.6	5.7	5.1	3.8	2.3
2000.....	5.8	7.1	7.0	8.3	8.3	7.2	5.9	4.3	3.8	2.7
1999.....	5.3	7.1	6.8	7.3	8.3	7.0	6.0	5.5	3.8	3.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K86.0,R78.0,X45,X65,Y15) 6/

White, both sexes										
2005 4/.....	10.4	16.1	20.0	20.2	19.1	16.6	13.8	10.1	8.0	4.6
2004 4/.....	10.4	16.0	19.8	18.7	18.2	16.6	14.0	10.9	7.5	4.5
2003 4/.....	10.8	16.2	18.7	18.2	18.2	17.0	13.2	10.6	7.6	4.2
2002.....	10.9	15.8	18.2	17.7	17.9	16.5	14.2	10.9	7.1	4.3
2001.....	10.4	16.5	17.1	16.8	18.6	16.7	13.8	10.9	7.7	4.8
2000.....	10.8	15.5	16.1	17.4	18.7	16.9	14.2	11.1	7.9	5.3
1999.....	10.5	15.3	16.2	17.1	18.5	17.0	13.6	11.6	8.3	5.3
White, male										
2005 4/.....	14.9	23.4	31.2	32.3	30.8	27.0	22.4	18.2	14.8	9.4
2004 4/.....	15.2	23.3	31.1	29.5	29.5	27.1	23.8	19.5	14.2	9.3
2003 4/.....	15.6	24.7	29.3	28.9	29.2	27.6	22.7	18.2	14.6	7.8
2002.....	15.8	24.0	29.1	28.4	29.8	27.5	23.0	18.7	13.2	9.3
2001.....	14.9	25.4	26.8	27.0	30.2	27.1	23.5	19.2	14.0	11.1
2000.....	16.0	24.6	26.0	27.6	30.3	28.1	24.5	20.8	14.9	11.6
1999.....	16.1	24.2	26.3	27.7	30.4	28.4	23.4	20.4	16.4	11.4
White, female										
2005 4/.....	5.8	8.9	9.0	8.5	8.2	7.3	6.6	4.1	3.8	2.5
2004 4/.....	5.5	8.7	8.8	8.3	7.7	7.4	6.0	4.6	3.3	2.3
2003 4/.....	6.0	7.8	8.3	8.0	8.0	7.5	5.4	5.1	3.3	2.5
2002.....	6.1	7.7	7.4	7.5	6.9	6.8	7.1	5.3	3.4	2.2
2001.....	5.9	7.6	7.6	7.0	8.0	7.6	5.8	5.1	3.9	2.2
2000.....	5.6	6.5	6.5	7.7	8.0	7.1	5.9	4.4	3.7	2.7
1999.....	4.8	6.5	6.3	7.0	7.6	7.0	5.9	5.4	3.6	2.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K86.0,R78.0,X45,X65,Y15) 6/

Black, both sexes										
2005 4/.....	8.1	14.6	19.6	20.3	23.5	17.0	15.9	10.7	8.5	5.6
2004 4/.....	7.5	14.2	20.6	23.2	23.9	22.4	17.1	8.6	6.2	*
2003 4/.....	8.7	15.1	22.6	22.4	26.7	19.0	14.3	9.0	8.6	*
2002.....	9.0	15.8	21.8	25.3	25.3	18.7	17.7	12.5	9.2	*
2001.....	12.2	18.4	27.4	26.9	24.6	25.3	14.7	13.5	7.3	*
2000.....	13.0	21.6	24.1	26.6	26.1	23.9	17.6	13.5	8.2	6.3
1999.....	12.9	22.8	24.1	29.3	28.9	24.9	21.2	16.4	11.6	7.7
Black, male										
2005 4/.....	11.1	23.1	31.7	33.6	41.3	31.1	30.4	24.1	15.5	*
2004 4/.....	11.5	22.2	33.0	40.6	42.6	42.9	35.0	17.3	*	*
2003 4/.....	12.6	24.7	39.4	37.2	49.2	33.0	27.6	17.3	21.4	*
2002.....	12.5	24.4	35.3	42.0	45.5	36.4	35.1	23.4	19.1	*
2001.....	18.2	28.9	43.9	47.9	44.3	47.8	29.6	26.0	*	*
2000.....	19.3	32.9	39.9	43.6	44.1	42.9	36.1	28.4	16.9	*
1999.....	18.9	35.9	39.9	52.9	48.6	48.3	41.4	31.3	22.7	*
Black, female										
2005 4/.....	5.4	7.3	9.4	9.3	9.6	6.5	6.1	*	*	*
2004 4/.....	4.0	7.3	10.1	8.8	9.3	7.2	5.1	*	*	*
2003 4/.....	5.3	6.7	8.4	10.3	9.2	8.6	5.3	*	*	*
2002.....	5.8	8.4	10.3	11.6	9.4	5.6	6.0	6.0	*	*
2001.....	6.9	9.2	13.4	9.6	9.1	8.6	4.7	6.1	*	*
2000.....	7.4	11.7	10.7	12.7	11.9	9.8	5.3	*	*	*
1999.....	7.6	11.6	10.8	10.1	13.5	7.5	7.9	7.4	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5, F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7-F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5, F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R78.1-R78.5,X40-X44, X60-X64,X85,Y10-Y14) 6/

All races 3/, both sexes

2005.....	24.0	25.8	21.2	12.8	7.8	5.0	4.4	4.9	6.1	7.6
2004.....	23.2	24.4	18.5	11.0	6.6	4.4	4.1	4.7	4.9	6.3
2003.....	22.8	22.7	17.0	9.8	5.7	4.5	3.7	3.9	4.6	6.0
2002.....	21.8	20.9	14.5	8.2	5.1	3.8	3.3	3.6	4.1	5.7
2001.....	18.5	17.6	11.6	6.3	4.1	3.1	2.9	3.4	3.6	5.1
2000.....	17.5	15.8	10.3	5.9	3.6	2.6	2.7	3.2	3.9	5.7
1999.....	17.4	15.5	9.3	5.8	3.8	3.2	2.9	3.6	4.2	4.8

All races 3/, male

2005.....	29.9	31.9	26.6	16.0	9.0	5.4	5.2	6.6	6.9	9.9
2004.....	28.3	30.1	23.6	13.0	7.6	4.8	4.5	5.2	5.6	8.6
2003.....	29.1	29.2	21.7	11.5	6.6	5.2	3.9	4.7	5.3	7.8
2002.....	28.0	26.4	19.1	10.0	5.7	4.1	3.4	3.9	4.7	7.3
2001.....	24.6	23.9	15.0	7.7	4.3	3.7	3.3	3.8	3.9	5.7
2000.....	23.8	21.9	13.8	6.7	3.9	2.9	2.8	3.6	4.3	7.5
1999.....	24.3	21.9	12.2	7.1	4.7	3.7	3.1	3.5	4.4	6.2

All races 3/, female

2005.....	18.2	19.8	16.1	9.8	6.6	4.8	3.8	3.7	5.5	6.6
2004.....	18.2	19.0	13.7	9.1	5.7	4.1	3.7	4.4	4.4	5.2
2003.....	16.6	16.5	12.4	8.2	4.9	4.0	3.4	3.3	4.2	5.2
2002.....	15.8	15.6	10.2	6.6	4.6	3.6	3.3	3.3	3.8	4.9
2001.....	12.6	11.6	8.3	5.1	3.9	2.6	2.6	3.2	3.4	4.9
2000.....	11.4	9.8	7.0	5.2	3.4	2.3	2.7	2.9	3.8	4.9
1999.....	10.6	9.2	6.6	4.6	2.9	2.7	2.7	3.7	4.0	4.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7-F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 6/

White, both sexes										
2005 4/.....	25.4	26.5	20.5	12.2	7.8	5.1	4.6	5.0	6.3	8.0
2004 4/.....	24.8	24.9	18.2	10.6	6.4	4.3	4.3	4.9	5.0	6.4
2003 4/.....	23.9	23.1	16.5	9.5	5.6	4.4	3.8	4.2	4.9	6.3
2002.....	22.6	20.8	13.8	7.9	4.9	3.9	3.5	3.7	4.3	5.8
2001.....	18.7	16.9	10.8	6.1	3.9	3.1	3.0	3.5	3.7	5.1
2000.....	17.3	15.0	9.5	5.7	3.6	2.5	2.8	3.4	4.1	5.6
1999.....	16.9	14.5	8.6	5.5	3.5	3.1	2.9	3.8	4.4	4.8

White, male										
2005 4/.....	31.4	32.0	24.8	14.4	8.4	5.0	5.2	6.7	6.9	10.6
2004 4/.....	29.9	30.4	22.1	11.6	6.8	4.2	4.6	5.0	5.7	8.7
2003 4/.....	30.3	29.2	20.4	10.7	6.2	4.7	3.9	4.9	5.5	7.8
2002.....	29.0	25.9	17.3	9.0	4.9	3.8	3.4	4.0	4.6	7.6
2001.....	24.8	22.3	13.3	6.9	3.6	3.6	3.2	3.6	4.1	5.6
2000.....	23.4	20.4	12.1	5.9	3.6	2.7	2.7	3.8	4.4	7.2
1999.....	23.5	20.0	10.6	6.2	4.0	3.5	3.1	3.5	4.6	6.4

White, female										
2005 4/.....	19.2	21.0	16.2	10.2	7.2	5.1	4.1	3.8	5.9	6.8
2004 4/.....	19.6	19.5	14.3	9.6	6.1	4.5	4.1	4.8	4.5	5.3
2003 4/.....	17.3	17.0	12.7	8.5	5.2	4.2	3.6	3.6	4.4	5.6
2002.....	16.3	15.8	10.4	7.0	4.9	3.9	3.5	3.5	4.2	4.9
2001.....	12.7	11.6	8.4	5.3	4.2	2.6	2.8	3.5	3.5	4.9
2000.....	11.3	9.7	6.9	5.5	3.7	2.4	3.0	3.0	3.9	4.9
1999.....	10.1	9.1	6.6	4.8	3.2	2.8	2.7	4.0	4.2	4.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5, F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7-F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5, F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R78.1-R78.5,X40-X44, X60-X64,X85,Y10-Y14) 6/

Black, both sexes

2005 4/.....	23.8	29.3	32.8	22.4	10.4	5.6	4.6	4.3	*	*
2004 4/.....	21.0	29.0	27.0	18.0	9.4	5.9	3.0	3.7	*	*
2003 4/.....	23.6	28.2	25.4	14.6	7.8	6.4	3.2	*	*	*
2002.....	24.1	28.2	25.0	13.6	8.9	3.7	*	*	*	*
2001.....	23.3	28.0	21.7	10.2	7.1	3.5	3.0	*	*	6.1
2000.....	23.9	26.0	21.0	9.6	4.8	3.6	*	*	*	7.5
1999.....	25.7	27.8	18.4	9.8	6.5	3.9	3.0	*	*	*

Black, male

2005 4/.....	30.2	41.3	48.4	36.0	19.1	10.0	7.6	*	*	*
2004 4/.....	26.9	37.1	41.6	29.7	16.1	10.9	*	*	*	*
2003 4/.....	31.1	39.3	38.5	23.3	13.1	11.6	*	*	*	*
2002.....	31.1	38.7	39.9	22.8	16.0	7.0	*	*	*	*
2001.....	31.7	43.3	34.4	17.1	12.6	5.3	*	*	*	*
2000.....	33.8	40.7	33.6	16.0	7.9	6.0	*	*	*	*
1999.....	36.8	44.3	30.2	17.6	12.1	6.5	*	*	*	*

Black, female

2005 4/.....	18.1	18.9	19.6	11.2	3.5	*	*	*	*	*
2004 4/.....	15.9	22.0	14.6	8.4	4.2	*	*	*	*	*
2003 4/.....	17.0	18.5	14.3	7.5	3.6	*	*	*	*	*
2002.....	17.9	19.1	12.4	6.1	3.3	*	*	*	*	*
2001.....	15.9	14.8	10.8	4.6	*	*	*	*	*	*
2000.....	15.1	13.3	10.2	4.5	*	*	*	*	*	*
1999.....	15.9	13.5	8.5	3.3	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 6/

All races 3/, both sexes

2005.....	11.6	11.4	10.9	9.8	9.9	9.5	11.3	13.0	14.5	12.0
2004.....	11.4	11.3	10.7	9.7	10.0	9.5	11.0	13.3	13.4	11.9
2003.....	11.5	11.4	10.9	10.4	9.6	9.7	11.3	12.9	14.2	12.5
2002.....	11.9	11.1	10.5	10.7	9.6	10.1	11.8	13.6	15.5	12.5
2001.....	11.3	10.6	10.4	10.4	9.7	10.5	11.4	13.7	15.3	12.8
2000.....	11.8	10.7	10.3	9.6	9.1	9.7	11.5	13.4	14.7	14.2
1999.....	10.9	10.7	9.6	9.9	9.4	10.0	12.2	14.0	14.5	13.5

All races 3/, male

2005.....	19.2	19.1	18.7	17.0	17.9	17.4	22.6	28.1	34.6	35.4
2004.....	18.5	18.4	18.0	16.9	17.4	17.8	22.1	28.2	32.0	35.9
2003.....	18.9	18.9	18.6	17.8	17.6	18.5	22.5	27.8	33.8	37.8
2002.....	19.7	18.4	18.0	18.6	17.1	18.8	24.1	29.8	37.6	38.9
2001.....	18.6	17.9	17.6	17.9	16.7	19.6	22.9	29.9	37.3	40.2
2000.....	19.4	17.9	17.3	16.5	16.1	17.8	23.1	29.6	36.6	44.7
1999.....	17.7	17.8	16.1	17.1	17.1	18.9	24.5	30.8	36.2	43.3

All races 3/, female

2005.....	4.1	4.0	3.3	3.0	2.6	2.7	2.3	2.1	2.0	1.3
2004.....	4.4	4.3	3.7	3.0	3.2	2.3	2.1	2.6	1.9	1.0
2003.....	4.2	4.1	3.5	3.4	2.4	2.1	2.3	2.3	2.2	1.3
2002.....	4.2	4.0	3.2	3.3	2.8	2.5	2.0	2.2	2.0	1.1
2001.....	4.1	3.6	3.5	3.3	3.2	2.6	2.2	2.3	2.2	1.3
2000.....	4.3	3.7	3.5	3.1	2.8	2.7	2.4	2.2	1.9	1.7
1999.....	4.2	3.7	3.4	3.2	2.4	2.4	2.5	2.3	1.9	1.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 6/										
White, both sexes										
2005 4/.....	11.1	11.7	11.4	10.7	10.7	10.3	12.2	14.2	15.6	13.0
2004 4/.....	11.1	11.6	11.3	10.2	10.7	10.3	11.8	14.3	14.4	12.8
2003 4/.....	11.2	11.6	11.5	11.1	10.4	10.5	12.1	13.9	15.1	13.5
2002.....	11.6	11.3	10.8	11.5	10.2	11.0	12.8	14.4	16.7	13.4
2001.....	11.0	10.9	10.9	10.9	10.5	11.3	12.1	14.6	16.1	13.8
2000.....	11.5	10.7	10.6	10.0	9.8	10.4	12.1	14.5	15.7	15.2
1999.....	10.6	10.6	10.0	10.4	9.8	10.7	12.9	14.9	15.4	14.5
White, male										
2005 4/.....	18.0	19.2	19.4	18.2	19.2	18.9	24.0	30.0	37.2	38.0
2004 4/.....	17.6	18.6	18.7	17.4	18.6	18.9	23.4	30.1	34.2	38.9
2003 4/.....	18.1	18.9	19.4	18.9	18.9	20.0	24.1	29.7	36.0	41.0
2002.....	19.1	18.4	18.5	19.7	18.1	20.3	25.8	31.2	40.3	41.7
2001.....	18.0	18.2	18.3	18.6	18.0	20.9	24.1	31.8	39.2	43.3
2000.....	18.7	17.8	17.6	17.0	17.0	18.9	24.1	31.9	38.8	48.4
1999.....	17.1	17.6	16.7	17.5	17.7	20.1	25.8	32.6	38.2	46.8
White, female										
2005 4/.....	4.2	4.2	3.6	3.4	2.7	2.8	2.4	2.4	2.1	1.4
2004 4/.....	4.6	4.5	4.0	3.2	3.4	2.6	2.3	2.7	2.0	1.1
2003 4/.....	4.2	4.4	3.6	3.7	2.5	2.2	2.3	2.4	2.2	1.3
2002.....	4.1	4.2	3.3	3.6	2.9	2.8	2.1	2.3	2.2	1.2
2001.....	4.0	3.7	3.7	3.6	3.6	2.9	2.3	2.4	2.2	1.4
2000.....	4.1	3.7	3.7	3.3	3.1	2.9	2.6	2.2	1.9	1.8
1999.....	4.1	3.8	3.4	3.6	2.5	2.5	2.7	2.4	1.9	1.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 6/

Black, both sexes

2005 4/.....	18.1	12.9	10.1	6.0	6.2	5.7	7.6	5.3	6.3	*
2004 4/.....	15.8	12.1	9.9	8.8	7.0	6.0	6.5	7.1	5.9	*
2003 4/.....	16.6	12.4	10.2	7.3	6.3	5.9	6.2	6.0	6.3	*
2002.....	16.6	13.0	10.3	7.3	7.0	5.1	6.7	8.8	*	*
2001.....	15.4	11.3	9.6	8.1	5.5	5.9	7.4	6.9	8.4	*
2000.....	16.5	13.2	10.6	7.4	5.8	6.2	7.7	4.6	6.8	*
1999.....	15.6	13.6	8.6	8.5	7.9	6.4	6.9	8.4	6.1	*

Black, male

2005 4/.....	33.3	23.5	18.8	11.8	11.1	10.2	16.1	14.3	14.7	*
2004 4/.....	29.2	21.6	18.6	16.5	12.3	13.3	15.9	15.9	15.8	*
2003 4/.....	29.4	23.4	18.2	13.7	11.8	11.3	12.2	13.2	14.8	*
2002.....	29.1	24.1	18.1	14.4	12.3	10.2	14.7	20.7	*	*
2001.....	26.8	20.4	17.4	16.3	10.4	12.2	15.8	16.3	20.9	*
2000.....	28.5	23.4	19.4	13.7	11.2	12.7	18.1	9.5	16.9	*
1999.....	26.9	24.4	15.2	17.3	15.5	12.7	15.6	19.5	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2005

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 6/

Black, female

2005 4/.....	4.8	3.7	2.7	*	*	*	*	*	*	*
2004 4/.....	4.0	4.0	2.5	2.5	*	*	*	*	*	*
2003 4/.....	5.4	2.8	3.4	*	*	*	*	*	*	*
2002.....	5.6	3.3	3.7	*	*	*	*	*	*	*
2001.....	5.4	3.5	2.9	*	*	*	*	*	*	*
2000.....	5.9	4.4	3.1	*	*	*	*	*	*	*
1999.....	5.6	4.3	3.1	*	*	*	*	*	*	*

* Figure does not meet standards of reliability or precision.

0.0 Quantity more than zero but less than 0.05.

... Category not applicable.

1/ Figures for age not stated included in "All ages" but not distributed among age groups.

2/ Death rates for "Under 1 year" (based on population estimates) differ from infant mortality rates (based on live births).

3/ Includes races other than white and black.

4/ Multiple-race data were reported for California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming in 2005, for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2003. The multiple-race data for these areas were bridged to the single-race and age categories in selected categories; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001."

5/ Appendix from Vital Statistics of the United States: Mortality, 2001.

6/ Tables in selected categories.

Source: CDC/NCHS, National Vital Statistics System, Mortality.